

A&A Ref: 154781

PUBLICATION PARTICULARS AND ABSTRACT
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

2006/03685 9 May 2006

26-07-2007

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

C12N, A61K

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

Powderject Vaccines, Inc.

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

FULLER, James

EARLIEST PRIORITY CLAIMED	COUNTRY		NUMBER	DATE	
	33	US	31	60/509,936	32 10 October 2003

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------

Nucleic acid constructs

57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	139
------------------	-----

The sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

WO 2005/035771 A3

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
21 April 2005 (21.04.2005)

PCT

(10) International Publication Number
WO 2005/035771 A3

(51) International Patent Classification⁷: C12N 15/85,
15/38, 15/36, 15/17, 15/12, A61K 48/00, 9/16

(74) Agent: WOODS, Geoffrey, Corlett; J.A. Kemp & Co., 14
South Square, Gray's Inn, London WC1R 5JJ (GB).

(21) International Application Number:
PCT/GB2004/004279

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, IT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(22) International Filing Date: 11 October 2004 (11.10.2004)

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/509,936 10 October 2003 (10.10.2003) US

(71) Applicant (for all designated States except US): POW-
DERJECT VACCINES, INC. [US/US]; 585 Science
Drive, Madison, WI 53711 (US).


(71) Applicant (for MG only): POWDERMED LIMITED
[GB/GB]; 2nd Floor Park Gate, 25 Milton Park, Abingdon,
Oxfordshire OX14 4SH (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): FULLER, James
[US/US]; 8551 Research Way Blvd., Middleton, WI 63562
(US).

(54) Title: NUCLEIC ACID CONSTRUCTS

[Continued on next page]



(57) Abstract: A nucleic acid construct comprising a chimeric promoter sequence and a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises: (a) a hCMV immediate early promoter sequence; (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

NUCLEIC ACID CONSTRUCTS

FIELD OF THE INVENTION

The invention relates to the fields of molecular biology and immunology and generally to reagents useful in nucleic acid immunisation techniques. More specifically, the invention relates to nucleic acid constructs for the expression of antigenic polypeptides, and to nucleic acid immunisation strategies using such reagents.

BACKGROUND OF THE INVENTION

Gene therapy and nucleic acid immunisation are promising approaches for the treatment and prevention of both acquired and inherited diseases. These techniques provide for the transfer of a desired nucleic acid into a subject with subsequent *in vivo* expression. Transfer can be accomplished by transfecting the subject's cells or tissues *ex vivo* and reintroducing the transformed material into the host. Alternatively, the nucleic acid can be administered *in vivo* directly to the recipient.

Each of these techniques requires efficient expression of the nucleic acid in the transfected cell, to provide a sufficient amount of the therapeutic or antigenic gene product. Several factors are known to affect the levels of expression obtained, including transfection efficiency, and the efficiency with which the gene or sequence of interest is transcribed and the mRNA translated.

A number of expression systems have been described in the art, each of which typically consists of a vector containing a gene or nucleotide sequence of interest operably linked to expression control sequences. These control sequences include transcriptional promoter sequences and transcriptional start and termination sequences. Commonly used promoters for mammalian cell expression systems include the SV40 early promoter, a cytomegalovirus (CMV) promoter

such as the CMV immediate early promoter (Chapman *et al* (1991) *Nucl. Acids Res.* 19:3979-3986), the mouse mammary tumour virus long terminal repeat (LTR) promoter, the adenovirus major late promoter (Ad MLP) and the herpes simplex virus (HSV) promoter, among others.

5 Nonviral promoters, such as a promoter derived from the murine metallothionein gene are also commonly used.

Expression systems often include transcriptional modulator elements, referred to as "enhancers". Enhancers are broadly defined as a

cis-acting agent, which when operably linked to a promoter/gene

10 sequence, will increase transcription of that gene sequence. Enhancers can function from positions that are much further away from a sequence of interest than other expression control elements (e.g. promoters), and can operate when positioned in either orientation relative to the sequence of interest (Banerji *et al.* (1981) *Cell* 27:299-308, deVilleirs *et al.* (1981)

15 *Nucl. Acids Res* 9: 6251-6264). Enhancers have been identified from a number of viral sources, including polyoma virus, BK virus, cytomegalovirus (CMV), adenovirus, simian virus 40 (SV40), Moloney sarcoma virus, bovine papilloma virus and Rous sarcoma virus (deVilleirs *et al supra*, Rosenthal *et al.* (1983) *Science* 222:749-755, Hearing *et al.* (1983) *Cell* 33:695-703, Weeks *et al.* (1983) *Mol. Cell. Biol.* 3:1222-1234, Levinson *et al.* (1982) *Nature* 295: 568-572, and Luciw *et al.* (1983) *Cell* 33: 705-716).

A number of expression systems for nucleic acid immunisation and gene therapy make use of the hCMV immediate early promoter. See eg

25 US Patent Nos. 5168062 and 5385839 to Stinski, and EP Patent Specification 0323997 B1. Expression vectors using the hCMV immediate early promoter include for example, pWRG7128 (Roy *et al*, Vaccine 19, 764-778, 2001), and pBC12/CMV and pJW4303 which are mentioned in WO 95/20660. Chapman *et al* (1991) report reduced

levels of expression from the hCMV immediate early promoter in the absence of hCMV Intron A.

SUMMARY OF THE INVENTION

A nucleic acid construct has been developed using manipulated 5 viral promoter/expression sequences, that provides enhanced expression of heterologous coding sequences in host cells. The construct is suitable for efficient expression of antigen-encoding genes, and can therefore be used in nucleic acid immunisation. In particular, the construct can be provided on carrier particles, for use in particle-mediated nucleic acid 10 immunisation.

Accordingly, the present invention provides a nucleic acid construct comprising a chimeric promoter sequence and a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises:

- 15 (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

20 The invention also provides:

- a nucleic acid construct comprising:
 - (i) a chimeric promoter sequence which comprises:
 - (a) a hCMV immediate early promoter sequence;
 - (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene; and

(ii) a cloning site for insertion of a coding sequence in 5 operable linkage with the chimeric promoter; and

(iii) (a) a non-translated leader sequence which is derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2gD antigen sequence and which is in operable linkage with the chimeric promoter; and/or

10 (b) an enhancer sequence which is derived from a 3' untranslated region (UTR) of a HBsAg sequence, or a 3'UTR of a simian CMV immediate early gene sequence and which is in operable linkage with the chimeric promoter, wherein the enhancer sequence is downstream of the 15 cloning site;

- a nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen 20 sequence; and

(iii) a coding sequence operably linked to (i) and (ii) wherein the coding sequence is heterologous to the non-translated leader sequence;

- a nucleic acid construct comprising:

25 (i) a promoter sequence;

- (ii) a coding sequence operably linked to the promoter sequence (i); and
 - (iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

5 wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence, and the coding sequence (ii) is heterologous to the 3' enhancer sequence;

- a method of obtaining expression in mammalian cells of a 10 polypeptide of interest, which method comprises transferring into said cells a nucleic acid construct of the invention, the construct including a coding sequence encoding the polypeptide;
- coated particles, suitable for delivery from a particle-mediated 15 delivery device, which particles comprise carrier particles coated with a nucleic acid construct of the invention, the construct including a coding sequence encoding the polypeptide;
- a dosage receptacle for a particle mediated delivery device comprising the coated particles;
- a particle mediated delivery device loaded with the coated 20 particles;
- a method of nucleic acid immunisation comprising administering to a subject an effective amount of the coated particles in which particles the coding sequence encodes an antigen;
- a purified isolated chimeric promoter sequence which comprises:

25 (a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

5 These and other objects, aspects, embodiments and advantages of the present invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the levels of expression of hepatitis B virus 10 surface antigen (HBsAg) obtained using various plasmid expression vectors.

Figure 2 shows the effect of intron inclusion on expression of HBsAg in SCC15 cells and of beta-gal in B16 cells (average of three expts).

15 Figure 3 shows the effect of rat insulin intron A and HBV3'UTR on expression of beta-gal in SCC15 cells (average of three expts).

Figure 4 shows the effect of rat insulin intron A and HBV3'UTR on expression of HSVgD in SCC15 cells (average of three expts).

20 Figure 5 shows the effect of rat insulin intron A and HBV3'UTR on expression of SEAP in SCC15 and B16 cells (three repetitions per cell line).

Figure 6 shows the ability of heterologous signal peptides to direct secretion of SEAP or hFc fragment in B16 cells.

Figure 7 illustrates levels of antibodies detected in the sera of mice immunised with antigen-encoding nucleic acids contained in a variety of plasmid expression vectors.

Figure 8 is a diagrammatic representation of pJV expression
5 vector.

Figure 9 is a diagrammatic representation of pJV7389.

Figure 10 is a diagrammatic representation of pJV7400.

Figure 11 is a diagrammatic representation of pJV7468.

Figure 12 is a diagrammatic representation of pJV7563.

10 Figure 13 sets out the base composition for pJV7563.

Brief Description of the Sequences

SEQ ID NO:1 is hCMV immediate early promoter sequence
(GenBank #M60321, X17403)

15 SEQ ID NO:2 is sequence from exons 1 and 2 of the hCMV major
immediate early gene (GenBank #M60321, X17403)

SEQ ID NO:3 is rat insulin intron A sequence (GenBank #J00748)

SEQ ID NO:4 is the sequence of a chimeric promoter according to
the present invention

20 SEQ ID NO:5 is a leader sequence from HBV preS2 antigen 5' UTR
sequence (GenBank #M54923)

SEQ ID NO:6 is a leader sequence from HSV type 2 gD 5'UTR
sequence (GenBank #Z86099)

SEQ ID NO:7 is a leader sequence from HBV e antigen 5'UTR sequence (GenBank #M54923)

SEQ ID NO:8 is HBVenh 3'UTR sequence (GenBank #AF143308)

SEQ ID NO:9 is simian immediate early gene 3'UTR sequence
5 (GenBank #M16019)

SEQ ID NO:10 is rabbit β globin poly A sequence (GenBank #K03256)

SEQ ID NO:11 is simian sCMV immediate early gene poly A sequence (GenBank #M16019)

10 SEQ ID NO:12 is HSV2 gB gene poly A sequence (GenBank #Z86099)

SEQ ID NO:13 is HPV16 early gene poly A sequence (GenBank #K02718)

SEQ ID NO:14 is the sequence of the pJV expression vector

15 SEQ ID NO:15 is PCR primer JF93

SEQ ID NO:16 is PCR primer F110

SEQ ID NO:17 is PCR primer GW1

SEQ ID NO:18 is PCR primer JF254

SEQ ID NO:19 is PCR primer GW150

20 SEQ ID NO:20 is PCR primer JF255

SEQ ID NO:21 is PCR primer DS1

SEQ ID NO:22 is PCR primer DA1

SEQ ID NO:23 is PCR primer JF301

SEQ ID NO:24 is PCR primer JF302

SEQ ID NO:25 is PCR primer JF84

SEQ ID NO:26 is PCR primer JF225

5 SEQ ID NO:27 is PCR primer JF335

SEQ ID NO:28 is PCR primer JF336

SEQ ID NO:29 is PCR primer JF357

SEQ ID NO:30 is PCR primer JF365

SEQ ID NO:31 is PCR primer JF393

10 SEQ ID NO:32 is PCR primer JF406

SEQ ID NO:33 is PCR primer JF256

SEQ ID NO:34 is PCR primer JF257

SEQ ID NO:35 is PCR primer JF320

SEQ ID NO:36 is PCR primer JF321

15 SEQ ID NO:37 is PCR primer JF386

SEQ ID NO:38 is PCR primer FcAS

SEQ ID NO:39 is oligonucleotide JF354

SEQ ID NO:40 is PCR primer JF355

SEQ ID NO:41 is PCR primer JF356

20 SEQ ID NO:42 is oligonucleotide JF348

SEQ ID NO:43 is PCR primer JF349

SEQ ID NO:44 is PCR primer JF350

SEQ ID NO:45 is oligonucleotide JF351

SEQ ID NO:46 is PCR primer JF352

5 SEQ ID NO:47 is PCR primer JF353

SEQ ID NO:48 is PCR primer JF430

SEQ ID NO:49 is PCR primer JF442

SEQ ID NO:50 is PCR primer JF421

SEQ ID NO:51 is PCR primer JF444

10 SEQ ID NO:52 is Pseudorabies virus (PRV) promoter sequence.

SEQ ID NO:53 is Rous sarcoma virus (RSV) promoter sequence.

DETAILED DESCRIPTION

Before describing the present invention in detail, it is to be understood that this invention is not limited to particularly exemplified molecules or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting. In addition, the practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology, recombinant DNA techniques and immunology all of which are within the ordinary skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.);

Oligonucleotide Synthesis (N. Gait, ed., 1984); *A Practical Guide to Molecular Cloning* (1984); and *Fundamental Virology*, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.).

All publications, patents and patent applications cited herein,
5 whether *supra* or *infra*, are hereby incorporated by reference in their entirety.

It must be noted that, as used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the content clearly dictates otherwise.

10 A. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although a number of methods and materials similar or equivalent to those described 15 herein can be used in the practice of the present invention, the preferred materials and methods are described herein.

In describing the present invention, the following terms will be employed, and are intended to be defined as indicated below.

The term "nucleic acid immunization" is used herein to refer to the
20 introduction of a nucleic acid molecule encoding one or more selected antigens into a host cell for the *in vivo* expression of the antigen or antigens. The nucleic acid molecule can be introduced directly into the recipient subject, such as by standard intramuscular or intradermal injection; transdermal particle delivery; inhalation; topically, or by oral, 25 intranasal or mucosal modes of administration. The molecule alternatively can be introduced *ex vivo* into cells which have been removed from a subject. In this latter case, cells containing the nucleic acid molecule of interest are re-introduced into the subject such that an

immune response can be mounted against the antigen encoded by the nucleic acid molecule. The nucleic acid molecules used in such immunization are generally referred to herein as "nucleic acid vaccines."

The term "adjuvant" intends an material or composition capable of 5 specifically or non-specifically altering, enhancing, directing, redirecting, potentiating or initiating an antigen specific immune response. Thus, coadministration of an adjuvant with an antigen may result in a lower dose or fewer dose of antigen being necessary to achieve a desired immune response in the subject to which the antigen is administered, or 10 coadministration may result in a qualitatively and/or quantitatively different immune response in the subject. In particular, the administration of the adjuvant may result in an enhanced immune response such as one of greater magnitude and/or duration. The effectiveness of an adjuvant can be determined by administering the adjuvant with a vaccine composition 15 in parallel with a vaccine composition alone to animals and comparing antibody and/or cellular mediated immunity in the two groups using standard assays such as radioimmunoassay, ELISAs, and CTL assays.

By "core carrier" is meant a carrier on which a guest nucleic acid (e.g., DNA, RNA) is coated in order to impart a defined particle size as 20 well as a sufficiently high density to achieve the momentum required for cell membrane penetration, such that the guest molecule can be delivered using particle-mediated techniques (see, e.g., U.S. Patent No. 5,100,792). Core carriers typically include materials such as tungsten, gold, platinum, ferrite, polystyrene and latex. See e.g., *Particle 25 Bombardment Technology for Gene Transfer*, (1994) Yang, N. ed., Oxford University Press, New York, NY pages 10-11.

By "needleless syringe" is meant an instrument which delivers a particulate composition transdermally without the aid of a conventional

needle to pierce the skin. Needleless syringes for use with the present invention are discussed herein.

The term "transdermal" delivery intends intradermal (e.g., into the dermis or epidermis), transdermal (e.g., "percutaneous") and 5 transmucosal administration, i.e., delivery by passage of an agent into or through skin or mucosal tissue. *See, e.g., Transdermal Drug Delivery: Developmental Issues and Research Initiatives*, Hadgraft and Guy (eds.), Marcel Dekker, Inc., (1989); *Controlled Drug Delivery: Fundamentals and Applications*, Robinson and Lee (eds.), Marcel Dekker Inc., (1987); and 10 *Transdermal Delivery of Drugs*, Vols. 1-3, Kydonieus and Berner (eds.), CRC Press, (1987). Thus, the term encompasses delivery from a needleless syringe as described in U.S. Patent No. 5,630,796, as well as particle-mediated delivery as described in U.S. Patent No. 5,865,796.

A "polypeptide" is used in its broadest sense to refer to a 15 compound of two or more subunit amino acids, amino acid analogs, or other peptidomimetics. The subunits may be linked by peptide bonds or by other bonds, for example ester, ether, etc. As used herein, the term "amino acid" refers to either natural and/or unnatural or synthetic amino acids, including glycine and both the D or L optical isomers, and amino 20 acid analogs and peptidomimetics. A peptide of three or more amino acids is commonly called an oligopeptide if the peptide chain is short. If the peptide chain is long, the peptide is typically called a polypeptide or a protein.

An "antigen" refers to any agent, generally a macromolecule, 25 which can elicit an immunological response in an individual. The term may be used to refer to an individual macromolecule or to a homogeneous or heterogeneous population of antigenic macromolecules. As used herein, "antigen" is generally used to refer to a protein molecule or portion thereof which contains one or more epitopes. For the

purposes of the present invention, antigens can be obtained or derived from any appropriate source. Furthermore, for the purposes of the present invention, an "antigen" includes a protein having modifications, such as deletions, additions and substitutions (generally conservative in nature) to the native sequence, so long as the protein maintains sufficient immunogenicity. These modifications may be deliberate, for example through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the antigens.

An "immune response" against an antigen of interest is the development in an individual of a humoral and/or a cellular immune response to that antigen. For purposes of the present invention, a "humoral immune response" refers to an immune response mediated by antibody molecules, while a "cellular immune response" is one mediated by T-lymphocytes and/or other white blood cells.

The terms "nucleic acid molecule" and "polynucleotide" are used interchangeably herein and refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. Polynucleotides may have any three-dimensional structure, and may perform any function, known or unknown. Non-limiting examples of polynucleotides include a gene, a gene fragment, exons, introns, messenger RNA (mRNA), transfer RNA, ribosomal RNA, ribozymes, cDNA, recombinant polynucleotides, branched polynucleotides, plasmids, vectors, isolated DNA of any sequence, isolated RNA of any sequence, nucleic acid probes, and primers.

A polynucleotide is typically composed of a specific sequence of four nucleotide bases: adenine (A); cytosine (C); guanine (G); and thymine (T) (uracil (U) for thymine (T) when the polynucleotide is RNA). Thus, the term nucleic acid sequence is the alphabetical representation of a polynucleotide molecule. This alphabetical representation can be input

into databases in a computer having a central processing unit and used for bioinformatics applications such as functional genomics and homology searching.

A "vector" is capable of transferring nucleic acid sequences to target cells (e.g., viral vectors, non-viral vectors, particulate carriers, and liposomes). Typically, "vector construct," "expression vector," and "gene transfer vector," mean any nucleic acid construct capable of directing the expression of a gene of interest and which can transfer gene sequences to target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors. A "plasmid" is a vector in the form of an extrachromosomal genetic element.

A nucleic acid sequence which "encodes" a selected antigen is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. For the purposes of the invention, such nucleic acid sequences can include, but are not limited to, cDNA from viral, prokaryotic or eukaryotic mRNA, genomic sequences from viral or prokaryotic DNA or RNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence.

A "promoter" is a nucleotide sequence which initiates and regulates transcription of a polypeptide-encoding polynucleotide. Promoters can include inducible promoters (where expression of a polynucleotide sequence operably linked to the promoter is induced by an analyte, cofactor, regulatory protein, etc.), repressible promoters (where expression of a polynucleotide sequence operably linked to the promoter is repressed by an analyte, cofactor, regulatory protein, etc.), and

constitutive promoters. It is intended that the term "promoter" or "control element" includes full-length promoter regions and functional (e.g., controls transcription or translation) segments of these regions.

"Operably linked" refers to an arrangement of elements wherein 5 the components so described are configured so as to perform their usual function. Thus, a given promoter operably linked to a nucleic acid sequence is capable of effecting the expression of that sequence when the proper enzymes are present. The promoter need not be contiguous with the sequence, so long as it functions to direct the expression 10 thereof. Thus, for example, intervening untranslated yet transcribed sequences can be present between the promoter sequence and the nucleic acid sequence and the promoter sequence can still be considered "operably linked" to the coding sequence.

"Recombinant" is used herein to describe a nucleic acid molecule 15 (polynucleotide) of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation is not associated with all or a portion of the polynucleotide with which it is associated in nature and/or is linked to a polynucleotide other than that to which it is linked in nature. Two nucleic acid sequences which are contained within a single 20 recombinant nucleic acid molecule are "heterologous" relative to each other when they are not normally associated with each other in nature.

Homologues of polynucleotides are referred to herein. Typically a polynucleotide which is homologous to another polynucleotide is at least 70% homologous to the polynucleotide, preferably at least 80 or 90%. 25 and more preferably at least 95%, 97% or 99% homologous thereto. Methods of measuring homology are well known in the art and it will be understood by those of skill in the art that in the present context, homology is calculated on the basis of nucleic acid identity. Such

homology may exist over a region of at least 15, preferably at least 30, for instance at least 40, 60 or 100 or more contiguous nucleotides.

Methods of measuring polynucleotide homology or identity are known in the art. For example the UWGCG Package provides the 5 BESTFIT program which can be used to calculate homology (e.g. used on its default settings) (Devereux *et al* (1984) *Nucleic Acids Research* 12, p387-395).

The PILEUP and BLAST algorithms can also be used to calculate homology or line up sequences (typically on their default settings), for 10 example as described in Altschul S.F. (1993) *J Mol Evol* 36:290-300; Altschul, S, F *et al* (1990) *J Mol Biol* 215:403-10.

Software for performing BLAST analysis is publicly available through the National Centre for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying 15 high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, *supra*). These initial neighbourhood word hits 20 act as seeds for initiating searches to find HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score goes to zero or below, due to the accumulation of one or more 25 negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89:10915-

10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) 5 *Proc. Natl. Acad. Sci. USA* 90:5873-5787. One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a sequence is considered similar to another 10 sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

The homologues typically hybridize with the relevant 15 polynucleotide at a level significantly above background. The signal level generated by the interaction between the homologue and the polynucleotide is typically at least 10 fold, preferably at least 100 fold, as intense as "background hybridisation". The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P . 20 Selective hybridisation is typically achieved using conditions of medium to high stringency, (for example, 0.03M sodium chloride and 0.003M sodium citrate at from about 50 °C to about 60 °C.

Stringent hybridization conditions can include 50% formamide, 5x 25 Denhardt's Solution, 5x SSC, 0.1% SDS and 100 $\mu\text{g}/\text{ml}$ denatured salmon sperm DNA and the washing conditions can include 2x SSC, 0.1% SDS at 37 °C followed by 1x SSC, 0.1% SDS at 68 °C. Defining appropriate hybridization conditions is within the skill of the art. See, e.g., Sambrook *et al.*, *supra*.

The homologue may differ from a sequence in the relevant polynucleotide by less than 3, 5, 10, 15, 20 or more mutations (each of which may be a substitution, deletion or insertion). These mutations may be measured over a region of at least 30, for instance at least 40, 5 60 or 100 or more contiguous nucleotides of the homologue. Where a polynucleotide encodes a polypeptide, substitutions preferably create "conservative" changes in the amino acid encoded. These are defined according to the following table. Amino acids in the same block in the second column and preferably in the same line in the third column may 10 be substituted for each other in conservative changes.

ALIPHATIC	Non-Polar	G A P
		I L V
	Polar-uncharged	C S T M
		N Q
AROMATIC	Polar-charged	D E
		K R
AROMATIC		H F W Y

The terms "individual" and "subject" are used interchangeably herein to refer to any member of the subphylum cordata, including, 15 without limitation, humans and other primates, including non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and guinea pigs; birds, including domestic, wild and 20 game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The terms do not denote a particular age. Thus, both adult and newborn individuals are intended to be covered. The methods described herein are intended for use in any of the above

vertebrate species, since the immune systems of all of these vertebrates operate similarly.

B. General Overview

The invention is concerned with nucleic acid constructs which 5 allow efficient expression of heterologous coding sequences, and in particular antigen-encoding genes, in host cells. More specifically, the invention provides nucleic acid constructs comprising, or in some embodiments, consisting essentially of a chimeric promoter sequence and a cloning site, such that when a coding sequence is inserted in the 10 cloning site, the coding sequence is in operable linkage with the chimeric promoter.

The chimeric promoter comprises, or in some embodiments consists essentially of:

- (a) a hCMV immediate early promoter sequence;
- 15 (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

The hCMV immediate early promoter sequence (a) may comprise:

- 20 (i) a native hCMV immediate early promoter sequence;
- (ii) a functional homologous variant thereof; or
- (iii) a functional fragment of (i) or (ii).

In general sequence (a) comprises about 100 to 600, preferably 200 to 600, for instance 400 to 600 nucleotides. Typically sequence (a) 25 comprises the sequences present in (i) which bind transcription factors or

the RNA polymerase, or instead of any of these sequences, homologues of these sequences able to bind the same transcription factors and RNA polymerase. Typically such sequences or their homologues are present in promoter sequence (a) in the same order and/or substantially the same 5 relative spacing as in (i).

Generally, (i) comprises at least from nucleotides -100 to -1, typically -150 to -1, for example -500 to -1 or -600 to -1 of the hCMV major immediate early gene. Sequence (i) typically comprises the hCMV core promoter sequence and may also include one or more enhancer 10 elements present in hCMV immediate early promoter. For example, (i) may comprise from nucleotides -118 to -1, or -524 to -1 as in US 6218140, or from nucleotides -62 to 1 or -465 to -1 as in US 538 5839.

Generally (i) includes a TATA box or CAAT box commonly found in promoter sequences. Preferably the sequence includes one or more of 15 the repeat sequences in the hCMV immediate early promoter.

In a preferred embodiment, (i) comprises SEQ ID NO: 1.

A hCMV immediate early promoter sequence can be obtained using known methods. A native hCMV immediate early promoter can be isolated directly from a sample of the virus, using standard techniques. 20 US 5385839, for example, describes the cloning of a hCMV promoter region. The sequence of a hCMV immediate early promoter is available at Genbank #M60321 (hCMV Towne strain) and X17403 (hCMV Ad169 strain). A native sequence could therefore be isolated by PCR using PCR primers based on the known sequence. See e.g Sambrook *et al*, *supra*, 25 for a description of techniques used to obtain and isolate DNA. A suitable hCMV promoter sequence could also be isolated from an existing plasmid vector. Promoter sequences can also be produced synthetically.

A functional variant (ii) or fragment (iii) is generally one which retains and/or complements the activity of the native promoter (i). Typically this activity is the ability to cause (including initiating and regulating) transcription of an operably linked polynucleotide, in particular 5 the hCMV major immediate early gene. In one embodiment, the variant or fragment would be able to complement the activity of the native promoter in a hCMV virus, for example allowing the virus to retain the ability to infect and/or replicate in cells.

A homologous variant (ii) or fragment (iii) can be assayed for the 10 ability to retain and/or complement the activity of (i). For example, a variant or fragment may be assayed for ability to restore functionality (such as infection and/or replication ability) to mutant hCMV in which the native hCMV immediate early promoter is defective.

A homologous variant (ii) or fragment (iii) may be tested for utility 15 using the Comparative Expression Assay below. The test promoter sequence is swapped into the base vector in place of the native hCMV immediate early promoter. Typically, a functional variant or fragment allows at least 50%, for instance, 60, 70, 80, 90% or more of the expression provided using the base vector. Generally expression is 20 provided in at least one but preferably two reference cell types. Typically, the reference cells are mammalian HEK 293T, CHO, HeLa, BHK, 3T3 or COS cells.

Additionally or alternatively, promoter sequence may be tested in 25 the Comparative Immunogenicity Assay below. Test promoter sequence is swapped into the base vector in place of the native hCMV immediate early promoter. A functional promoter sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the

base vector. Preferred antigens are HBsAg, HSV 2gD and flu-M2 antigens. According to this assay, a functional homologous variant (ii) or functional fragment (iii) of native promoter sequence (i) is one which allows the highest antibody titres achieved by the native sequence.

5 As mentioned above, the construct may comprise exon sequence (b) which comprises sequence derived from exon 1 and exon 2 of the hCMV major immediate early gene. Exons are coding sequences, which in nature are generally separated by introns. In the native hCMV major immediate early gene, exons 1 and 2 are usually separated by the native 10 intron A. In the present chimeric construct exon 2 sequence is generally positioned 3' of exon 1 sequence, without intervening intron sequence, so that the exon 1 and exon 2 sequences are contiguous.

Exon sequence (b) may comprise:

- (i) native exon sequence, typically exon 1 and (whole or 15 partial) exon 2;
- (ii) a homologous variant of (i) which is functional; or
- (iii) a functional fragment of (i) or (ii).

Sequence (i) may comprise from about 50 to 100% of the native hCMV major immediate early gene exon 1 sequence, for example, 60 to 20 90% or 70 to 80%. Typically at least 50% of the natural exon 1 sequence is present, such as 60%, 70%, 80%, 90% or more. Exon sequence (b) also comprises at least a part of exon 2 sequence. In sequence (i), typically 2 or more bases of native exon 2, for example 2 to 9, 2 to 7 or 3 to 5 bases are present. Up to and including 100% of 25 natural exon 2 sequence, for example 5 to 95%, 20 to 80% or 40 to 60% of natural exon 2 sequence may be present. Typically the homologous variant has any of the above lengths mentioned for the native sequence.

Preferably (i) comprises SEQ ID No. 2.

Suitable exon sequence (b) can be obtained using known methods.

See e.g Sambrook *et al.*, *supra*, for a description of techniques used to obtain and isolate DNA. Native hCMV major immediate early gene

5 sequence can be isolated directly from a sample of the virus, using standard techniques (see for example, MacLean, A (1998) "Preparation of HSV-DNA and Production of Infectious Virus" in *Herpes Simplex Virus Protocols* S. Brown, A Maclean, editors, Humana Press, Totowa, NJ, pp.19-26). The sequence of a hCMV major immediate early gene, 10 including the location of exon 1 and exon 2, is available at Genbank #M60321, X17403. Native exon 1 and 2 sequences could therefore be isolated by cutting the native major gene sequence at appropriate restriction sites or by PCR using PCR primers based on the known sequence. Suitable exon sequences could alternatively be isolated from 15 an existing plasmid vector, such as pWRG7128. Exon sequences can also be produced synthetically, rather than cloned. Variant sequences can readily be constructed by routine methodologies such as site-directed mutagenesis.

Generally the exon sequence will, when present in the construct of 20 the invention, enhance expression, typically causing comparable enhancement to the native exon 1 and exon 2 sequence (i) mentioned above.

Exon sequence may be assayed for functionality using the Comparative Expression Assay below. Test exon sequence is swapped 25 into the base vector in place of the exon sequence already present. Generally exon sequence is functional if the sequence does not abrogate expression but preferably increases expression in at least one but preferably two reference cell types when compared to the base vector. Typically the reference cells are mammalian HEK293T, CHO, HeLa, BHK,

3T3 or COS cells. Preferably expression increases by at least 5%, 10%, 20%, 30% or 40%. According to this assay, a functional homologous variant (ii) or functional fragment (iii) of natural exon sequence (i) is one which allows at least 50% of the expression improvement provided by 5 the natural sequence.

Additionally or alternatively, exon sequence may be tested in the Comparative Immunogenicity Assay below. Test exon sequence is swapped into the base vector in place of the exon sequence already present. Functional exon sequence provides antibody titres that are at 10 least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBsAg, HSV 2gD and flu-M2 antigens. According to this assay, a functional homologous variant (ii) or functional fragment 15 (iii) of natural exon sequence (i) is one which allows the highest antibody titres achieved by the natural sequence.

The chimeric promoter construct comprises heterologous intron (c) in place of the native intron A region of the hCMV major immediate early gene. An intron is a non-coding sequence which is spliced from the 20 hnRNA transcribed from a gene. A heterologous intron is one which is not present in the coding sequence in nature.

The heterologous intron (c) replaces wholly or partially, native intron A of the hCMV major immediate early gene. Typically the native intron A is absent.

25 In general the heterologous intron (c) is 3' of exon sequence (b).

Typically the heterologous intron (c) comprises:

(i) a natural intron;

- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

Heterologous intron (c) is in general a viral or eukaryotic intron. Preferably the intron is a mammalian intron, in particular a non-human 5 intron, for example a rat or chicken intron. Preferably the intron is an intron A, for example, rat insulin intron A, chicken keratin intron A or chicken cardiac action intron A.

Typically intron (c) has a length of from about 50 nucleotides to about 1000 nucleotides, for instance from about 100 to about 500 10 nucleotides. The intron (c) may for example, comprise 50 to 500 nucleotides, such as up to 100, 200, 300 or 400 nucleotides. Preferably the intron comprises sequence found at about nucleotides 50 to 133 of native rat insulin intron A, or a homologue of this sequence.

Preferably heterologous intron (c) is capable of being spliced from 15 an RNA transcript in a eukaryotic host cell. In general the intron comprises one or more of a donor sequence (such as GT), an acceptor sequence (such as AG), a 3' pyrimidine rich region and a branch point sequence. The pyrimidine rich region, if present, may include, for example at least 5, 8, 10 or more pyrimidines. Preferably the intron 20 comprises at least a donor sequence, acceptor sequence and a branch point sequence. Typically in the chimeric construct, intron (c) comprises non-intron flanking sequences which are derived from exon sequences found on the intron/exon boundaries of the natural intron (i). The flanking exon sequence may be native exon sequence or a homologue of 25 this sequences which retains substantially the same activity as the native sequence, for example retains splicing function. Typically from 5 to 10, preferably from 7 to 10 bases of exon sequence are included at each end of the intron.

Intron (c) may be an artificial intron, provided that the intron is functional. For example, a recombinant or chimeric intron may be used. Such an intron may comprise sequence from more than one natural intron.

5 Typically intron (c) comprises sequences present in hCMV intron A which bind transcription factors or regulatory proteins or instead of any of these sequences, homologues of these sequences able to bind the same factors or proteins. Typically such sequences or their homologues are present in the intron (c) in the same order and/or substantially the
10 same relative spacing as in hCMV intron A.

Intron (c) may comprise an homologous variant (ii) in which the sequence of the natural intron (i) has been modified to remove an internal restriction site. For example, an homologous variant of rat insulin intron A may be used in which an internal *N*hel site has been destroyed.

15 Preferably, intron (c) comprises:

- (i) SEQ ID No. 3;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

Intron sequence (c) may be obtained using standard cloning
20 techniques. For example, rat insulin intron A sequence is available at GenBank J00748, chicken keratin intron A sequence at GenBank J00847 and chicken cardiac intron A sequence at GenBank X02212. Intron sequence can be isolated from natural sources using primers based on known sequence. Sequence may be prepared synthetically. Variant
25 sequencer may be obtained by mutagenesis.

Typically a functional intron sequence, for example a functional variant (ii) or a functional fragment (iii) is one which has substantially the

same activity as, and/or complements the activity of a natural intron (i). In one embodiment the activity is splicing activity.

Intron (c) sequences may be tested for splicing activity using a routine splicing assay. In general a functional homologue (ii) or 5 functional fragment (iii) will show at least 50%, for example 60%, 70%, 80%, 90% and up to 100% or more of the splicing efficiency of the natural intron (i) in the assay.

In general the heterologous intron sequence will, when present in the construct of the invention, enhance expression. Typically, a variant 10 (ii) or fragment (iii) intron will cause comparable enhancement to a natural intron (i).

Functionality of potential intron sequence (c) can be tested using the Comparative Expression Assay below. The heterologous intron is swapped into the base vector. Generally heterologous intron sequence is 15 functional if the addition of the sequence increases expression in at least one, but preferably two reference cell types by 25% or more, compared to the base vector. Typically the reference cells are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. The increase in expression may be at least 35%, 45%, 55% or more. According to this 20 assay, a functional variant (i) or functional fragment (ii) of a natural intron sequence (i) is one which allows greater than 50% of the expression improvement achieved by the natural sequence.

A heterologous intron (c) sequence may, additionally or alternatively be tested for functionality using the Comparative 25 Immunogenicity Assay below. Intron (c) sequence is added to the base vector. A functional intron (c) sequence provides antibody titres that are higher than those achieved by the base vector with at least one, preferably two antigens. Preferably, the antibody titres are at least 5 or 10%, for instance 20%, 30% or 40% higher than with the base vector.

Preferred antigens are HBsAg, HSV2gD and Flu-M2 antigens. According to this assay, a functional variant (ii) or functional fragment (iii) of a natural intron sequence (i) is one which allows the highest antibody titres achieved by the natural sequence.

5 Suitable heterologous intron sequence can be obtained using standard cloning techniques. For example, rat insulin intron A sequence is available at GenBank J00748, chicken keratin intron A at GenBank J00847, and chicken cardiac actin intron A at X02212. Intron sequence can be isolated from native sources using primers based on the 10 known sequence data. Suitable sequence may also be prepared synthetically.

The component sequences (a), (b) and (c) may be provided suitably linked together to form a chimeric promoter using standard cloning or molecular biology techniques. Preferably intron sequence (c) is 15 provided 3' of exon sequence (b). The chimeric promoter construct is linked to a cloning site, in such a way that the promoter will effect the expression of a coding sequence inserted in the site, when the proper enzymes are present. Suitable cloning sites, including multi-cloning sites are known in the art, e.g the pUC19, pBC SK, pBluescript II KS, 20 cDNA3.1, pSP72, pGEM 7Z multicloning site.

Typically, a nucleic acid for insertion (or inserted) in the cloning site encodes a therapeutically relevant polypeptide. It is preferred that the coding sequence is suitable for use in nucleic acid immunisation or gene therapy. The nucleic acid insert may thus comprise a sequence 25 capable of providing immunity, for example an immunogenic sequence that elicits a humoral and/or cellular immune response when delivered to a subject. Alternatively, the nucleic acid may comprise one or more genes encoding a therapeutic polypeptide e.g a protein defective or missing from a target cell genome or a non-native protein having a

desired biological or therapeutic effect (e.g., an antiviral function). For the treatment of genetic disorders, functional genes corresponding to genes known to be deficient in the particular disorder can be administered to a subject. Preferably the nucleic acid is DNA.

5 Suitable nucleic acids for insertion include those used for the treatment of inflammatory diseases, autoimmune, chronic and infectious diseases, including such disorders as AIDS, cancer, neurological diseases, cardiovascular disease, hypercholesterolemia; various blood disorders including various anemias, thalassemia and hemophilia; genetic 10 defects such as cystic fibrosis, Gaucher's Disease, adenosine deaminase (ADA) deficiency, emphysema, etc.

For example, in methods for the treatment of solid tumors, genes encoding toxic peptides (i.e., chemotherapeutic agents such as ricin, diphtheria toxin and cobra venom factor), tumor suppressor genes such as 15 p53, genes coding for mRNA sequences which are antisense to transforming oncogenes, antineoplastic peptides such as tumor necrosis factor (TNF) and other cytokines, or transdominant negative mutants of transforming oncogenes, can be inserted for expression at or near the tumor site.

20 Similarly, nucleic acids coding for polypeptides known to display antiviral and/or antibacterial activity, or stimulate the host's immune system, can also be included. The nucleic acid may encode one of the various cytokines (or functional fragments thereof), such as the interleukins, interferons and colony stimulating factors. The nucleic acid 25 may encode an antigen for the treatment or prevention of a number of conditions including but not limited to cancer, allergies, toxicity and infection by a pathogen such as, but not limited to, fungus, viruses including Human Papilloma Viruses (HPV), HIV, HSV2/HSV1, influenza virus (types A, B and C), Polio virus, RSV virus, Rhinoviruses,

Rotaviruses, Hepatitis A virus, Norwalk Virus Group, Enteroviruses, Astroviruses, Measles virus, Par Influenza virus, Mumps virus, Varicella-Zoster virus, Cytomegalovirus, Epstein-Barr virus, Adenoviruses, Rubella virus, Human T-cell Lymphoma type I virus (HTLV-I), Hepatitis B virus (HBV), Hepatitis C virus (HCV), Hepatitis D virus, Pox virus, Marburg and Ebola; bacteria including *M. tuberculosis*, *Chlamydia*, *N. gonorrhoeae*, *Shigella*, *Salmonella*, *Vibrio Cholera*, *Treponema pallidua*, *Pseudomonas*, *Bordetella pertussis*, *Brucella*, *Franciscella tularensis*, *Helicobacter pylori*, *Leptospira interrogans*, *Legionella pneumophila*, *Yersinia pestis*, *Streptococcus* (types A and B), *Pneumococcus*, *Meningococcus*, *Hemophilus influenza* (type b), *Toxoplasma gondii*, *Complbybacteriosis*, *Moraxella catarrhalis*, *Donovanosis*, and *Actinomycosis*; fungal pathogens including Candidiasis and Aspergillosis; parasitic pathogens including Taenia, Flukes, Roundworms, Amebiasis, Giardiasis, Cryptosporidium, *Schitosoma*, *Pneumocystis carinii*, Trichomoniasis and Trichinosis. The nucleic acid may also be used to provide a suitable immune response against numerous veterinary diseases, such as Foot and Mouth diseases, Coronavirus, *Pasteurella multocida*, *Helicobacter*, *Strongylus vulgaris*, *Actinobacillus pleuropneumonia*, Bovine viral diarrhea virus (BVDV), *Klebsiella pneumoniae*, *E. Coli*, *Bordetella pertussis*, *Bordetella parapertussis* and *Bordetella bronchiseptica*. Thus in one aspect, the nucleic acid constructs of the present invention may find use in a vaccine.

In some embodiments, the nucleic acid construct will encode an adjuvant. Thus, the sequences inserted into the cloning site for insertion of a coding sequence may encode a polypeptide that can act as an adjuvant. In a preferred instance, the encoded adjuvant may be an ADP-ribosylating bacterial toxin. These include diphtheria toxin (DT), pertussis toxin (PT), cholera toxin (CT), the *E. coli* heat labile toxins (LT1 and LT2), *Pseudomonas* endotoxin A, *Pseudomonas* exotoxin S, *B. cereus*

exoenzyme, *B. sphaericus* toxin, *C. botulinum* C2 and C3 toxins, *C. limosum* exoenzyme, as well as toxins from *C. perfringens*, *C. spiriforma* and *C. difficile* and *Staphylococcus aureus* EDIN. Most ADP-ribosylating bacterial toxins contain A and B subunits. The construct may express 5 the A subunit, the B subunit and/or both subunits.

In a preferred instance, the nucleic acid construct may encode the *E. coli* heat labile toxin and/or cholera toxin and in particular may express *E. coli* heat labile toxin. A GenBank entry for the complete sequences of the CT subunit A and B genes can be found at Locus VIBCTXABB 10 (Accession No. D30053), while a GenBank entry for the complete sequences of the LT subunit A and B genes can be found at locus AB0116677 (Accession No. AB011677).

The construct may express an active variants or fragment of a particular adjuvant. The variant or fragment will be said to be active if it 15 retains at least some of the adjuvant activity of the polypeptide it is derived from. Thus, the variant and/or fragment will still be able to enhance an immune response against a particular antigen in comparison to the immune response seen when no adjuvant is administered with the antigen. The encoded sequence may be active fragments or variants of 20 CT A and/or B subunits and in particular may be active fragments of LT A and/or B subunits.

The toxin subunit may have had its naturally occurring signal sequence deleted. A naturally occurring exotoxin subunit may have been modified to detoxify the toxin. The A subunit may have been modified to 25 disrupt or inactivate ADP-ribosyl transferase activity.

Thus, the adjuvant constructs of the invention may be used to enhance an immune response against a particular antigen. The enhanced immune response may involve an immune response of greater magnitude or duration. It may mean that when the antigen is re-encountered the

immune response then is greater than if no adjuvant was administered. The enhance immune response may result in higher antibody titres. In the case of some constructs, and in particular those expressing exotoxin subunits the adjuvant may result in an augmented cellular response and a 5 T helper 1-like immune response against the antigen in question.

The nucleic acid for insertion into the cloning site may comprise a polyadenylation (polyA) sequence. Such a polyA sequence is generally native to the coding sequence. Alternatively, a heterologous polyA sequence may be provided in the nucleic acid construct of the invention.

10 Typically the polyA sequence will be provided downstream of the cloning site, such that it is operatively linked to a coding sequence inserted in the cloning site. Any suitable polyA sequence may be included in the construct using standard cloning techniques. Such polyA sequences are known in the art.

15 The poly A sequence may be:

- (i) a natural poly A sequence;
- (ii) a functional homologous variant of (i); or
- (iii) a functional fragment of (i) or (ii).

20 The natural poly A sequence (i) may be, for example a rabbit β globin gene poly A, Human Papilloma Virus (HPV) early or late gene poly A, HSV-2gB gene poly A, a simian CMV immediate early gene poly A or HSV gD late gene poly A.

25 Preferably the natural poly A sequence (i) is selected from the group consisting of SEQ ID No. 10 (GenBank K03256), SEQ ID No. 11 (GenBank M16019), SEQ ID No. 12 (GenBank Z80699) and SEQ ID No. 13 (GenBank K02718).

In general, a functional polyA sequence is one which retains polyadenylation activity.

A poly A sequence may be tested for the ability to bring about polyadenylation of an RNA transcript using a routine expression assay. A 5 functional homologous variant (ii) or functional fragment (iii) typically shows at least 50%, for example 60%, 70% 80% or more of the poly A activity of the natural poly A sequence in the assay.

Generally the poly A sequence will, when present in the construct of the invention, enhance expression, typically causing comparable 10 enhancement to the natural poly A sequence (i) mentioned above.

A poly A sequence may also be assayed for functionality using the Comparative Expression Assay below. A test poly A region is swapped into the base vector in place of the RBGpA. A test poly A is considered functional if the poly A does not abrogate expression but preferably 15 increases expression in at least one but preferably two reference cell types, compared to the base vector. Preferably there is an increase in expression of at least 5%, 10%, 20%, 30%, 40% or 50% or more. Preferred cell types are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. According to the assay, an homologous variant (ii) or 20 fragment (iii) is functional if it allows greater than 50% of the expression improvement achieved by the natural poly A sequence (i).

Alternatively, or additionally, poly A sequences may be tested for activity in the Comparative Immunogenicity Assay below. Poly A sequence is swapped into the base vector in place of RBGpA. A 25 functional poly A sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector with at least one, preferably two antigens. Preferably the antibody titres are at least 5%, 10%, for instance 20%, 30% or 40% higher than those achieved with the base vector. Preferred antigens are HBsAg, HSV2gD and Flu-

M2 antigens. An homologous variant (ii) or fragment (iii) is functional if it allows the highest antibody titres achieved by the natural poly A sequence (i).

The nucleic acid construct may comprise additional control sequences which influence expression of a coding sequence inserted in the cloning site. The construct may include a non-translated leader sequence. The sequence is provided in the construct in operable linkage with the chimeric promoter, and therefore also with a coding sequence inserted at the cloning site. The leader provides a translational start site for expression of an inserted coding sequence and typically includes a Kozak sequence.

Typically the untranslated leader sequence comprises:

- (i) a natural untranslated leader sequence;
- (ii) a functional homologous variant of (i); or
- 15 (iii) a functional fragment of (i) or (ii).

In general the natural sequence (i) is a eukaryotic sequence or a viral sequence, in particular, of a virus which infects a eukaryote. Preferably the natural sequence (i) is HBV or HSV sequence, for example HBV preS2 antigen sequence, HBV e-antigen sequence, or HSV type 2gD 20 antigen sequence. Particularly preferably, (i) is selected from the group consisting of SEQ ID No. 5, SEQ ID No. 6 and SEQ ID No. 7.

Typically the leader sequence comprises sequences present in (i) which bind transcription components or regulatory proteins, or homologues of these sequences which are able to bind the same 25 components or proteins. Typically such sequences or their homologues are present in the leader sequence in the same order and/or substantially the same relative spacing as in (i). In general the leader sequence

comprises a translational start site for expression of an inserted coding sequence. Typically the leader sequence includes a Kozak sequence.

In general the untranslated leader sequence has a length of from about 10 to about 200 nucleotides, for example from about 15 to 150 5 nucleotides, preferably 15 to about 130 nucleotides. Leader sequences comprising, for example, 15, 50, 75 or 100 nucleotides may be used.

Generally a functional untranslated leader sequence is one which is able to provide a translational start site for expression of a coding sequence in operable linkage with the leader sequence. Typically a 10 functional variant (ii) or fragment (iii) has substantially the same activity as and/or complements the activity of the natural sequence (i), usually in facilitating or enhancing expression of a coding sequence in operable linkage with the sequence.

A variant (ii) or fragment (iii) may be tested for activity as an 15 untranslated leader sequence relative to natural leader sequence using standard protocols. For example, expression vectors may be prepared comprising a natural leader sequence (i) operably linked to its native coding sequence, and expression monitored in suitable host cells e.g. mammalian HEK 293T, CHO, HeLa, BHK, 3T3 or COS cells. Test 20 constructs may be prepared in which the natural leader sequence is replaced by an homologous variant or fragment and expression is monitored again in the same host cells. In general, a variant (ii) or fragment (iii) provides at least 50%, such as 60%, 70%, 80%, 90% or 100% or more of the expression provided by the natural sequence.

25 A potential leader sequence can also be tested for utility in the Comparative Expression Assay below. A test leader sequence is swapped into the base vector in place of the HBV preS2 5'UTR. A functional leader sequence does not abrogate expression but preferably increases expression in at least one but preferably two reference cell

types, compared to the base vector. In general expression is increased by at least 5%, 10%, 20%, 30%, 40% or 50%. Preferred cell types are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. According to the assay, an homologous variant (ii) or fragment (iii) is functional if it 5 allows greater than 50% of the expression improvement achieved by the natural leader sequence.

Alternatively, or additionally, a leader sequence may be tested for activity in the Comparative Immunogenicity Assay below. A leader sequence is swapped into the base vector in place of HBV preS2 5'UTR. 10 A functional leader sequence provides antibody titres that are at least as high as or higher than those achieved by the base vector, with at least one preferably two antigens. Preferably the antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBsAg, HSV 2gD and Flu-M2 antigens. An 15 homologous variant (ii) or fragment (iii) is functional if it allows the highest antibody titres allowed by the natural leader sequence (i).

Suitable leader sequence can be obtained using standard protocols. For example, HBV preS2 antigen sequence, HBV e-antigen sequence and HSV type 2gD antigen sequence is available at GenBank M54923, 20 M54923 and Z86099 respectively. Primers can be designed based on this known sequence and used to isolate homologous sequences. Leader sequences may be synthesised based on known sequences.

The nucleic acid construct may comprise an enhancer sequence. An enhancer sequence is typically provided 3' of the cloning site, in 25 operable linkage with both the chimeric promoter and an inserted coding sequence, and acts to increase transcription of the inserted sequence.

In general the enhancer comprises:

(i) a natural enhancer;

(ii) a functional homologous variant of (i); or

(iii) a functional fragment of (i) or (ii).

The enhancer sequence generally comprises from about 50 to about 850 nucleotides, for example from about 75 to about 500 5 nucleotides. Enhancers of about 100, 200, 300 or 400 nucleotides may be used.

Typically (i) is a eukaryotic or viral enhancer, in particular, of a virus which infects eukaryotes. Usually such enhancers occur in the 3' untranslated region (3'UTR) of a gene. Preferably (i) is an HBV or a CMV 10 enhancer, for example an HBs Ag 3' UTR or a simian CMV immediate early gene 3' UTR. Preferably (i) comprises SEQ ID No. 8 or SEQ ID No. 9.

In general, the enhancer in the construct comprises sequences found in (i) which bind transcription components or regulatory proteins, 15 for example transcription factors, or homologues of these sequences which bind the same components or proteins. Preferably these sequences are present in the enhancer in the same order and/or substantially the same relative spacing as in (i).

Generally a functional enhancer is one which enhances or 20 increases expression of a polynucleotide, for example, a coding sequence, which is operably linked to the enhancer sequence. Typically a functional homologous variant (ii) or fragment (iii) has substantially the same activity (for example, enhancement of expression) as and/or complements the activity of the natural enhancer (i).

25 Enhancer activity may be assayed using an enhancer trap assay. Protocols are known in the art. A functional homologous variant (ii) or fragment (iii) preferably provides at least 50% of the enhancer activity

shown by the natural enhancer in such as assay. Typically the activity is at least 60%, 70%, 80%, 90, 100% or more of the activity of the natural enhancer. In general, a functional variant (ii) or fragment (iii) is able to complement the activity of the natural enhancer (i) in the assay.

5 Enhancer utility may also be tested using the Comparative Expression Assay set out below. A test 3'UTR sequence is swapped into the base vector. A 3'UTR has utility if it does not abrogate expression but preferably increases expression in at least one but preferably two reference cell types compared to the base vectors in the assay.

10 Preferably expression is increased by at least 5%, 10%, 20%, 30%, 40% or 50%. Preferred cell types are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. According to this assay, an homologous variant (ii) or fragment (iii) is functional if it allows greater than 50% of the expression improvement achieved by the natural enhancer sequence

15 (i).

Additionally or alternatively, enhancer sequences may be tested for activity in the Comparative Immunogenicity Assay below. A 3'UTR is swapped into the base vector. A functional enhancer sequence provides antibody titres that are at least as high as or higher than those achieved

20 by the base vector with at least one, preferably two antigens. Preferably the antibody titres are at least 5%, 10%, 20%, 30% or 40% higher than with the base vector. Preferred antigens are HBs Ag, HSV2gD and Flu-M2 antigens. An homologous variant (ii) or fragment (iii) is functional if it allows the highest antibody titre allowed by the natural enhancer

25 sequence (i).

Suitable enhancer sequence can be obtained using standard cloning methods. For example, HBsAg 3'UTR sequence, or simian CMV immediate early gene 3'UTR sequence can be accessed at GenBank

AF143308 and M16019. Primers can be designed based on this known sequence and used to isolate homologous sequences.

In a preferred embodiment, the nucleic acid construct comprises a heterologous polyA sequence, a heterologous leader sequence and a 5 heterologous enhancer, all in operable linkage with the chimeric promoter, for efficient expression of an inserted coding sequence.

In a further aspect, the present invention also provides a nucleic acid construct comprising, or sometimes consisting essentially of:

- (i) a promoter sequence
- 10 (ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and
- (iii) a coding sequence operably linked to (i) and (ii)

wherein the coding sequence is heterologous to the non-translated 15 leader sequence.

Typically the promoter sequence (i) is derived from a viral or eukaryotic promoter sequence. The promoter sequence may be a natural promoter sequence, a functional homologue of the natural sequence or a functional fragment of either. Suitable natural promoters include, for 20 example, the hCMV immediate early promoter, Pseudorabies virus (PRV) promoter or Rous sarcoma virus (RSV) promoter. Preferably the natural promoter comprises SEQ ID NO: 52 or SEQ ID NO: 53.

An artificial promoter construct, such as the chimeric promoter described above, may be used, provided that the promoter is functional.

A functional promoter sequence is generally one which is able to cause (including initiate and regulate) transcription of an operably linked coding sequence in a suitable host cell.

A promoter sequence may be tested for promoter activity using a 5 routine expression assay. A functional homologue or fragment of a natural promoter sequence typically provides at least 50%, for example, or least 60, 70, 80 or 90% of the expression provided by the natural sequence in such an assay.

The non-translated leader sequence (ii) is as described above.

10 Suitable coding sequences (iii) include those already described in relation to the chimeric promoter construct. However, in the present aspect of the invention, the coding sequence is heterologous to the non-translated leader sequence. The present construct typically includes a poly A sequence, which as already described, may be native to the coding 15 sequence, or provided as a heterologous poly A sequence in the construct. Suitable poly A sequences have already been described. The construct may additionally include an enhancer sequence 3' of the coding sequence. Suitable enhancer sequences are described above in relation to the chimeric promoter construct.

20 In another aspect, the invention provides a nucleic acid construct comprising, or in some embodiments consisting essentially of:

- (i) a promoter sequence;
- (ii) a coding sequence operably linked to the promoter sequence (i) and;
- 25 (iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence, and the coding sequence (ii) is heterologous to the enhancer sequence.

5 The construct may include a non-translated leader sequence such as the ones already described in relation to the chimeric promoter construct.

Typically the promoter sequence (i) is derived from a viral or eukaryotic promoter sequence. The promoter sequence may be a natural 10 promoter sequence, a functional homologue of the natural sequence or a functional fragment of either. Suitable natural promoters include, for example, the hCMV immediate early promoter, Pseudorabies virus (PRV) promoter or Rous sarcoma virus (RSV) promoter. Preferably the natural promoter comprises SEQ ID NO: 52 or SEQ ID NO: 53.

15 An artificial promoter construct, such as the chimeric promoter described above, may be used, provided that the promoter is functional.

A functional promoter sequence is generally one which is able to cause (including initiate and regulate) transcription of an operably linked coding sequence in a suitable host cell.

20 A promoter sequence may be tested for promoter activity using a routine expression assay. Functional homologues or fragments of a natural promoter sequence typically provide at least 50%, for example, or least 60, 70, 80 or 90% or the expression provided by the natural sequence in such an assay.

25 Suitable coding sequences (ii) include those already mentioned in relation to the chimeric promoter construct. However, in the present aspect, the coding sequence is heterologous to the 3'enhancer sequence. The enhancer sequence (iii) of the construct is described above. The

present construct also typically includes a poly A sequence. As in the case of the chimeric promoter construct, this poly A region may be native to the coding sequence (ii) or may be provided as a heterologous poly A component in the construct.

5 A construct according to any aspect of the present invention may comprise a signal peptide sequence. The signal peptide sequence is inserted in operable linkage with the promoter such that the signal peptide is expressed and facilitates secretion of a polypeptide encoded by coding sequence also in operable linkage with the promoter.

10 Typically a signal peptide sequence encodes a peptide of 10 to 30 amino acids for example 15 to 20 amino acids. Often the amino acids are predominantly hydrophobic. In a typical situation, a signal peptide targets a growing polypeptide chain bearing the signal peptide to the endoplasmic reticulum of the expressing cell. The signal peptide is 15 cleaved off in the endoplasmic reticulum, allow for secretion of the polypeptide via the Golgi apparatus.

A signal peptide for use in the invention may comprise:

- (i) a natural signal peptide sequence;
- (ii) a homologous variant of (i) which retains signal peptide activity; or
- (iii) a fragment of (i) or (ii) which retains signal peptide activity.

Sequence (i) may be for example human tissue plasminogen activator signal peptide (hTPAAsp) (GenBank L00141), the aprotinin signal peptide (GenBank AAD13685), tobacco extensin signal peptide (GenBank JU0465), or chicken lysosyme signal peptide (GenBank AF410481).

A signal peptide, suitable for use in the present invention, is one which will enable the secretion of heterologous proteins. A functional signal peptide can be identified in an assay which compares the effect of a test signal peptide with the effect of a known signal peptide - e.g.

5 human tissue plasminogen activator signal peptide (hTPAsP) - and with the effect of having no signal peptide. The Comparative Expression Assay set out below may be used but with the following modification. Secretion expression vectors are constructed containing the base vector with either the test signal peptide, hTPAsP or no signal peptide. Coding 10 sequences for polypeptides devoid of their naturally occurring signal peptides are inserted into the vectors and the vectors transformed into reference host cells. Preferably cells are mammalian HEK293T, CHO, HeLa, BHK, 3T3 or COS cells. The cell media is analysed for polypeptide expression levels. A functional signal peptide enables polypeptide 15 secretion at a higher level than a vector lacking a signal peptide with at least one, preferably two polypeptides. Typically, secretion is 5% higher, or more preferably 10% higher or more, for example 20 or 50% higher or more. Typically, secretion levels are comparable to those obtained using hTPAsP.

20 Allowing secretion of encoded protein outside of an expressing cell may have a number of advantages, in particular where the protein is an antigen. For example, increased antigen secretion could allow greater antigen uptake and response by immune cells (macrophages, Langhans' cells, B-cells, T-cells etc), enable the ability of antigen to reach the 25 bloodstream and signal cells (cytokines), enable an antigen to find cellular ligands and effect a function (antibodies, toxins such as cholera toxin, *E.coli* LT) and participate in normal cellular biochemical processes (cellular receptors).

A nucleic acid construct of the invention may be in the form of a 30 plasmid expression vector. The vector may then include additional

elements, such as an origin of replication, or selector genes. Such elements are known in the art and can be included using standard techniques. In one embodiment, the plasmid vector has the sequence in SEQ ID NO:14. Alternatively, the construct may be included in a viral vector construct.

In some embodiments, the nucleic acid construct of the invention may comprise two or more of the chimeric promoters defined herein. Thus, the construct may comprise a plurality of chimeric promoters and in particular the construct may have two, three, four, five or more chimeric promoters. The chimeric promoters will preferably be each separately operably linked to a cloning site for insertion of a coding sequence. Thus, the construct may express two, three, four, five or more more coding sequences. The coding sequences expressed may be any of those specified herein. In a preferred instance, the construct has two chimeric promoters with each having a coding sequence operably linked to them. In particular, the two promoters may be transcribed away from each other.

In particular, the constructs with two promoters may express the A and B subunits of an ADP-ribosylating bacterial toxin, including any of those mentioned herein and preferably an LTA and B subunit.

In cases where the construct has multiple chimeric promoters each may comprise, or be operably linked to, any of the sequences mentioned herein. In a particularly preferred instance, the heterologous intron of one or more of the promoters may be the rat insulin gene intron A sequence. One or more of the chimeric promoters may also preferably comprise the 5' UTR of HSV-2gB pre-S2. One or more of the promoters may comprise the poly adenylation sequence of the rat beta globin gene.

In a preferred case, the nucleic acid construct of the invention may comprise two chimeric promoter sequences, with each promoter

sequence being operably linked to a cloning site which has a coding sequence inserted into it, where each chimeric promoter comprises

- (a) a hCMV immediate early promoter sequence;
- (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
- 5 (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

with the coding sequence operably linked to one chimeric promoter encoding a LTA subunit and the coding sequence linked to the other 10 encoding an LTB subunit. The construct can therefore express both subunits. Preferably:

- the heterologous intron of each promoter is the rat insulin gene intron A sequence;
- the sequence encoding each LT subunit is operably linked to the 5' UTR 15 of HBV pre-S2; and/or
- each LT encoding sequence is operably linked to the rat beta globin gene polyadenylation sequence.

A polynucleotide construct of the invention may be substantially free of or associated with cells or with cellular material. It may be in 20 substantially isolated form, or it may be in substantially purified form, in which case it will generally comprise at least 90% e.g. at least 95%, 98% or 99% of the polynucleotide or dry mass in the preparation.

The present nucleic acid molecules may be delivered to suitable host cells, for expression of a polynucleotide in operable linkage with the 25 promoter. Preferably the host cells are mammalian cells, in particular human cells. Suitable methods for delivery of nucleic acids to such cells are known in the art and include, for example, dextran mediated

transfection, calcium phosphate precipitation, electroporation and direct microinjection into nuclei.

As described above, a nucleic acid coding sequence in a construct may encode a therapeutically relevant polypeptide. The present constructs may therefore be used for nucleic acid immunisation or gene therapy using standard gene delivery protocols. Suitable methods for gene delivery are known in the art, as discussed below. The nucleic acid molecules can be delivered either directly to a subject, or alternatively, delivered *ex vivo* to cells derived from the subject whereafter the cells are reimplanted in the subject.

For use in nucleic acid immunisation or gene therapy, the nucleic acid constructs may be formulated as conventional pharmaceutical preparations. This can be done using standard pharmaceutical formulation chemistries and methodologies, which are available to those skilled in the art. For example, compositions containing one or more nucleic acid sequences (e.g., present in a suitable vector form such as a DNA plasmid) can be combined with one or more pharmaceutically acceptable excipients or vehicles to provide a liquid preparation.

Auxiliary substances, such as wetting or emulsifying agents, pH buffering substances and the like, may be present in the excipient or vehicle. These excipients, vehicles and auxiliary substances are generally pharmaceutical agents which may be administered without undue toxicity and which, in the case of vaccine compositions will not induce an immune response in the individual receiving the composition.

Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, polyethyleneglycol, hyaluronic acid, glycerol and ethanol. Pharmaceutically acceptable salts can also be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of

organic acids such as acetates, propionates, malonates, benzoates, and the like. It is also preferred, although not required, that the preparation will contain a pharmaceutically acceptable excipient that serves as a stabilizer, particularly for peptide, protein or other like molecules if they are to be included in the composition. Examples of suitable carriers that also act as stabilizers for peptides include, without limitation, pharmaceutical grades of dextrose, sucrose, lactose, trehalose, mannitol, sorbitol, inositol, dextran, and the like. Other suitable carriers include, again without limitation, starch, cellulose, sodium or calcium phosphates, citric acid, tartaric acid, glycine, high molecular weight polyethylene glycols (PEGs), and combination thereof. A thorough discussion of pharmaceutically acceptable excipients, vehicles and auxiliary substances is available in REMINGTON'S PHARMACEUTICAL SCIENCES (Mack Pub. Co., N.J. 1991), incorporated herein by reference.

15 Certain facilitators of nucleic acid uptake and/or expression ("transfection facilitating agents") can also be included in the compositions, for example, facilitators such as bupivacaine, cardiotoxin and sucrose, and transfection facilitating vehicles such as liposomal or lipid preparations that are routinely used to deliver nucleic acid molecules. Anionic and neutral liposomes are widely available and well known for delivering nucleic acid molecules (see, e.g., *Liposomes: A Practical Approach*, (1990) RPC New Ed., IRL Press). Cationic lipid preparations are also well known vehicles for use in delivery of nucleic acid molecules. Suitable lipid preparations include DOTMA (N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride), available under the tradename Lipofectin™, and DOTAP (1,2-bis(oleyloxy)-3-(trimethylammonio)propane), see, e.g., Felgner *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7416; Malone *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:6077-6081; US Patent Nos 5,283,185 and 5,527,928, and 20 International Publication Nos WO 90/11092, WO 91/15501 and WO 25 25 30

95/26356. These cationic lipids may preferably be used in association with a neutral lipid, for example DOPE (dioleyl phosphatidylethanolamine). Still further transfection-facilitating compositions that can be added to the above lipid or liposome 5 preparations include spermine derivatives (see, e.g., International Publication No. WO 93/18759) and membrane-permeabilizing compounds such as GALA, Gramicidine S and cationic bile salts (see, e.g., International Publication No. WO 93/19768).

Alternatively, the nucleic acid molecules of the present invention 10 may be encapsulated, adsorbed to, or associated with, particulate carriers. Suitable particulate carriers include those derived from polymethyl methacrylate polymers, as well as PLG microparticles derived from poly(lactides) and poly(lactide-co-glycolides). See, e.g., Jeffery *et al.* (1993) *Pharm. Res.* 10:362-368. Other particulate systems and 15 polymers can also be used, for example, polymers such as polylysine, polyarginine, polyornithine, spermine, spermidine, as well as conjugates of these molecules.

Once formulated the compositions can be delivered to a subject *in vivo* using a variety of known routes and techniques. For example, the 20 liquid preparations can be provided as an injectable solution, suspension or emulsion and administered via parenteral, subcutaneous, intradermal, intramuscular, intravenous injection using a conventional needle and syringe, or using a liquid jet injection system. Liquid preparations can also be administered topically to skin or mucosal tissue, or provided as a 25 finely divided spray suitable for respiratory or pulmonary administration. Other modes of administration include oral administration, suppositories, and active or passive transdermal delivery techniques.

Alternatively, the compositions can be administered *ex vivo*, for example delivery and reimplantation of transformed cells into a subject

are known (e.g., dextran-mediated transfection, calcium phosphate precipitation, electroporation, and direct microinjection of into nuclei).

The compositions are administered to a subject in an amount that is compatible with the dosage formulation and that will be prophylactically and/or therapeutically effective. An appropriate effective amount will fall in a relatively broad range but can be readily determined by one of skill in the art by routine trials. The "Physicians Desk Reference" and "Goodman and Gilman's The Pharmacological Basis of Therapeutics" are useful for the purpose of determining the amount needed. For example, it is generally expected that an effective dose of the polynucleotide will fall within a range of about 0.001 to 1000 μ g, more preferably 0.01 to 10.0 μ g.

In one instance, a nucleic acid construct of the invention may be used in conjunction with another nucleic acid construct. In one case, the nucleic acid construct may be one of those described herein for the expression of an adjuvant and the other construct may be a construct encoding one or more antigens. In a preferred case, both constructs may employ the chimeric promoters of the invention.

In the case where one construct expresses an adjuvant and the other an antigen or antigens, the antigens may in particular be from HSV or Hepatitis virus (particularly Hepatitis B virus). The antigens may in particular be the HSV ICP0, ICP4, ICP 22 and/ICP 27 antigens and preferably all four. In cases where such antigens are expressed, the adjuvant construct will in particular express LTA and/or LTB and in particular both.

The two constructs may be administered separately, simultaneously or sequentially. The two may be administered in the same or different compositions. In particular, where one construct has an adjuvant effect the two will be delivered so that an adjuvant effect is

seen, that is the immune response generated will be greater and/or for a longer period than if the adjuvant had not been administered with the antigen. In a preferred instance, the two constructs may be delivered in the same composition, preferably on the same carrier particles.

5 In a preferred embodiment, the nucleic acid constructs of the invention are delivered to target cells using a particle-mediated delivery technique. Particle mediated methods for delivering nucleic acid preparations are known in the art.

Particles for particle mediated delivery may be formed by coating 10 the present nucleic acid molecules onto carrier particles (e.g., core carriers) using a variety of techniques known in the art. Carrier particles are selected from materials which have a suitable density in the range of particle sizes typically used for intracellular delivery from a particle-mediated delivery device. Typically carrier particles have a diameter of 15 from 0.1 to 5 μ m, for example 0.5 to 3 μ m, preferably 1 to 2 μ m. The optimum carrier particle size will, of course, depend on the diameter of the target cells. Alternatively, colloidal gold particles can be used wherein the coated colloidal gold is administered (e.g., injected) into tissue (e.g.; skin or muscle) and subsequently taken-up by immune- 20 competent cells.

Usually carrier particles are selected from inert metals. The metals are inert in that they are not physiologically active. For the purposes of the invention, tungsten, gold, platinum and iridium carrier particles can be used. Tungsten and gold particles are preferred. Tungsten particles are readily available in average sizes of 0.5 to 2.0 μ m in diameter. 25 Although such particles have optimal density for use in particle acceleration delivery methods, and allow highly efficient coating with DNA, tungsten may potentially be toxic to certain cell types. Gold particles or microcrystalline gold (e.g., gold powder A1570, available

from Engelhard Corp., East Newark, NJ) will also find use with the present methods. Gold particles provide uniformity in size (available from Alpha Chemicals in particle sizes of 1-3 μm , or available from Degussa, South Plainfield, NJ in a range of particle sizes including 0.95 μm) and 5 reduced toxicity. Microcrystalline gold provides a diverse particle size distribution, typically in the range of 0.1-5 μm . However, the irregular surface area of microcrystalline gold provides for highly efficient coating with nucleic acids.

A number of methods are known and have been described for 10 coating or precipitating DNA or RNA onto gold or tungsten particles. Most such methods generally combine a predetermined amount of gold or tungsten with plasmid DNA, CaCl_2 and spermidine. The resulting solution is vortexed continually during the coating procedure to ensure uniformity of the reaction mixture. After precipitation of the nucleic acid, 15 the coated particles can be transferred to suitable membranes and allowed to dry prior to use, coated onto surfaces of a sample module or cassette, or loaded into a delivery cassette for use in particular particle-mediated delivery instruments.

As an alternative, the polynucleotides of the invention can be 20 formulated as a particulate composition. Formulation can be carried out using the above-described standard pharmaceutical formulation chemistries. For example, the polynucleotides can be combined with one or more pharmaceutically acceptable excipients or vehicles to provide a suitable composition. The formulated compositions are then prepared as 25 particles using standard techniques such as by simple evaporation (air drying), vacuum drying, spray drying, freeze drying (lyophilisation), spray-freeze drying, spray coating, precipitation, supercritical fluid particle formulation, and the like. If desired, the resultant particles can be densified using the techniques described in commonly owned

International Publication No. WO 97/48485, incorporated herein by reference.

These methods can be used to obtain nucleic acid particles having a size ranging from about 0.01 to about 250 μm , preferably about 10 to 5 about 150 μm , and most preferably about 20 to about 60 μm ; and a particle density ranging from about 0.1 to about 25 g/cm³, and a bulk density of about 0.5 to about 3.0 g/cm³, or greater.

Once formed, the particles comprising the nucleic acid molecules may be packaged in single unit dosages or multidose containers. Such 10 containers may comprise an hermetically sealed container enclosing a suitable amount of the particles. The particles can be packaged as a sterile formulation, and the hermetically sealed container can thus be designed to preserve the sterility of the formulation until use in delivery to a subject. The containers are preferably adapted for direct use in a 15 particle mediated delivery device. Typically such containers take the form of capsules, foil pouches, sachets, cassettes and the like. The particle delivery devices can also be provided in a preloaded condition containing a suitable dosage of the particles. The preloaded device may then also be prepackaged in a hermetically sealed container.

20 The container in which the particles are packaged can further be labelled to identify the composition and provide relevant dosage information. In addition, the container can be labelled with a notice in the form prescribed by a governmental agency, for example, the Food 25 and Drug Administration, wherein the notice indicates approval by the agency under Federal Law of the manufacture, use or sale of the nucleic acid preparation contained therein for human administration.

Particle acceleration devices, suitable for particle-mediated delivery are known in the art. For example, current gene gun devices employ an explosive, electric or gaseous discharge to propel coated carrier particles

towards target cells. The coated carrier particles can be releasably attached to a movable carrier sheet, or removably attached to a surface along which a gas stream passes, lifting the particles from the surface and accelerating them toward the target. An example of a gaseous 5 discharge device is described in U.S.Patent No. 5,204,253. An explosive-type device is described in U.S.Patent No. 4,945,050. One example of an electric discharge apparatus suitable for use herein is described in U.S.Patent No. 5,120,657. Another electric discharge apparatus is described in US Patent No 5,149,655. The disclosure of all 10 of these patents is incorporated herein by reference in their entireties.

Particles may also be administered using a needleless syringe device, such as those described in U.S.Patent No. 5,630,796 to Bellhouse *et al* ("the PowderJect® needle less syringe device") and in International Publication Nos. WO 94/24263, WO 96/04947, WO 15 96/12513 and WO 96/20022, all of which are incorporated herein by reference.

Devices such as the one described in US Patent No.5,630,796 may be provided as a pen-shaped instrument containing, in linear order moving from top to bottom, a gas cylinder, a particle cassette or 20 package, and a supersonic nozzle with an associated silencer medium. The particles are provided within a suitable container, e.g. a cassette formed by two rupturable polymer membranes that are heat-sealed to a washer-shaped spacer to form a self-contained sealed unit. Membrane materials can be selected to achieve a specific mode of opening and 25 burst pressure that dictate the conditions at which the supersonic flow is initiated.

In operation, the device is actuated to release the compressed gas from the cylinder into an expansion chamber within the device. The released gas contacts the particle cassette and, when sufficient pressure

is built up, suddenly breaches the cassette membranes sweeping the particles into the supersonic nozzle for subsequent delivery. The nozzle is designed to achieve a specific gas velocity and flow pattern to deliver a quantity of particles to a target surface of predefined area. The 5 silencer is used to attenuate the noise produced by the supersonic gas flow.

The delivery system described in International Publication No. WO 96/20022 also uses the energy of a compressed gas source to accelerate and deliver powdered compositions. However, it is

10 distinguished from the system of US Patent No. 5,630,796 in its use of a shock wave instead of gas flow to accelerate the particles. More particularly, an instantaneous pressure rise provided by a shock wave generated behind a flexible dome strikes the back of the dome, causing a sudden eversion of the flexible dome in the direction of a target surface.

15 This sudden eversion catapults a powdered composition (which is located on the outside of the dome) at a sufficient velocity, thus momentum, to penetrate target tissue, e.g., oral mucosal tissue. The powdered composition is released at the point of full dome eversion. The dome also serves to completely contain the high-pressure gas flow which

20 therefore does not come into contact with the tissue. Because the gas is not released during this delivery operation, the system is inherently quiet. This design can be used in other enclosed or other wise sensitive applications for example, to deliver particles to minimally invasive surgical sites.

25 Particles may be delivered *in vivo* directly to a subject, or *ex vivo* to cells taken from a subject, the transformed cells then being reimplanted in the subject. For *in vivo* delivery, particle injection is typically subcutaneously, epidermally, intradermally, intramucosally (e.g. nasally, rectally and/or vaginally), intraperitoneally, intravenously, orally

30 or intramuscularly. Preferably, delivery is to terminally differentiated

cells; however, the particles can also be delivered to non-differentiated, or partially differentiated cells such as stem cells of blood and skin fibroblasts. Most preferably, delivery is to skin epidermal cells.

The particles are administered to a subject in a manner compatible with the dosage formulation and in an amount that will be prophylactically and/or therapeutically effective. A "therapeutically effective amount" of the present particulate compositions will be sufficient to bring about treatment or prevention of disease or condition symptoms, and will fall in a relatively broad range that can be determined by routine trials. Generally the particles are delivered in an amount of from 0.001 to 1000 μ g, more preferably 0.01 to 10.0 μ g of nucleic acid per dose. However, the exact amount necessary will vary depending on the age and general condition of the individual being treated and the particular nucleotide sequence selected, as well as other factors. An appropriate effective amount can be readily determined through clinical testing. The "Physicians Desk Reference" and "Goodman and Gilman's The Pharmacological Basis of Therapeutics" are useful for the purpose of determining the amount needed.

AssaysComparative Expression Assay

A suitable test for element utility determines the effect the element has on expression of a polypeptide. The basis of comparison for testing utility of the elements is a 'base vector', generally (unless otherwise noted) a plasmid with a hCMV promoter, hCMV exon 1, 9 bases of hCMV exon 2, the 5'UTR from HBV preS2 and the rabbit- β globin_polyadenylation region, positioned to drive expression of a coding sequence. Typically, the base vector is pJV7384, pJV 7401, pJV 7450 or pJV 7533. Heterologous introns and 3'UTRs are added, or promoter sequences, exons, 5'UTRs and polyA sites are swapped into the base vectors to create test expression vectors. Thus functional variants or fragments can be tested.

The base vectors and test vectors are transformed into suitable host cells and the cells analysed for polypeptide expression levels. Preferably mammalian host cells are used. Suitable cells include mammalian HEK 293T, CHO, HeLa, BHK, 3T3 or COS cells.

Typically, a functional element causes expression which is comparable to the base vector, for example at least the same as or greater. Preferably expression is tested in more than one cell type and with more than one coding sequence.

Suitable experimental protocols are provided, for example, in Examples 1 to 13 below.

Comparative Immunogenicity Assay

Where the polypeptide to be expressed is an antigen, a further test may be carried out to identify functional or particularly preferred construct elements. In the assay, the effect of an element on immune response is determined after delivery of an expression vector to a test organism. Antibody levels against the antigen are the easiest way to judge immune response. Groups of mice are vaccinated with the base vectors or test vectors constructed as above. Sera is collected after an appropriate amount of time and analyzed for antibody levels.

10 This experiment is performed twice, and the antibody levels from all the groups in both experiments are plotted. Functional elements will give rise to at least as high as or higher antibody titres in both experiments for a particular antigen than the base vector. Preferably, the result will be seen with more than one antigen to demonstrate the 15 breadth of utility of the element(s) in that expression panel.

Suitable experimental protocols are provided for example, in Example 14 below.

C. Experimental

Below are examples of specific embodiments for carrying 20 out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some 25 experimental error and deviation should, of course, be allowed for.

MethodsStandard PCR Conditions

The standard PCR conditions used for the construction of vectors were as follows: 1x PCR core buffer w/ 1.5mM MgCl₂ (Promega Corporation, Madison, WI), 0.400 μ M each of each primer, 200 μ M of each dNTP (USB, Inc, Cleveland, OH) , 2.5 μ Taq polymerase (Promega Corporation, Madison, WI), 1.0 ng template DNA, water to 100 μ l, and a mineral oil (Aldrich Chemical, Inc, Milwaukee, WI) overlay. The PTC-200 thermocycler (MJ Research, Inc, Waltham, MA) was programmed to run the following routine: 4'@95°C, 30 cycles of (1'@95°C/ 1'15" @55°C/ 1'@72°C), 10'@72°C, 4°C hold). The amplification products were removed from the PCR reaction by using the QIAquick \ddagger PCR Purification Kit (Qiagen Inc, Valencia, CA) prior to cutting with restriction enzymes (New England Biolabs, Beverly, MA).

All PCR products were sequenced after cloning to ensure fidelity of the amplification.

Example 1. Construction of hepatitis B virus surface antigen (HBsAg) vector panels

A number of plasmid expression vectors were constructed for expression of HBsAg.

Starting materials

(i) pWRG7128 (Roy, M, et.al. *Vaccine* (2001) 19: 764-778), which contains the hCMV immediate early promoter sequence, the first exon, first intron, and a partial second exon of the hCMV major immediate early gene, the HBsAg coding sequence with flanking regions (HBV preS2 5'UTR derived sequence and 3' posttranscriptional response

element) and the bovine growth hormone polyadenylation region (BGHpA)

(ii) pJV7284, a derivative of pWRG7128 that exchanges the rabbit β globin polyadenylation region (RBGpA) for BGHpA.

5 (a) pJV7384 (CMV(no intron), HBV preS2 5'UTR and RBGpA)

pWRG7128 was PCR'd with JF93 (SEQ ID NO:15) and F110 (SEQ ID NO:16) using standard conditions and cut with *Sa*/1 and *Bam*H1 to isolate an insert fragment containing the CMV promoter, exon 1 and part of the exon 2 sequence. pAM6 (ATCC, Mannassas, VA) was cut with 10 *Bam*H1 and *Bst*X1 to isolate an insert fragment that contained the 5'-UTR of HBsAg, and roughly 70% of the HBsAg coding region. pJV7284 was cut with *Sa*/1 and *Bst*X1 to generate a vector fragment into which the two insert fragments were ligated, resulting in pJV7293.

pWRG7128 was PCR'd with primers GW1 (SEQ ID NO:17) and 15 JF254 (SEQ ID NO:18) and cut with *Bst*X1 and *Bg*/2 to isolate an insert fragment that contained the 3'-end of the HBsAg coding region. pJV7293 was cut with *Bst*X1 and *Bg*/2 to generate a vector fragment into which the insert fragment was ligated, resulting in vector pJV7384.

20 (b) pJV7382 (CMV(no intron), HBsAg 3'UTR, HBV preS2 5'UTR and RBGpA)

pJV7293 was cut with *Xho*1 and *Xba*1 to generate an insert fragment containing the CMV promoter/exons and the 5'-UTR with 5'-end of the HBsAg coding sequence. pWRG7128 was cut with *Xba*1 and 25 *Bc*/1 to generate an insert fragment containing the majority of the HBsAg coding sequence and the 3'-UTR. pJV7284 was cut with *Xho*1 and *Bg*/2 to generate a vector fragment into which the two insert fragments were ligated, resulting in pJV7382.

(c) pJV7389 (CMV (RIA), HBsAg 3'UTR, HBV preS2 5'UTR and RBGpA)

The rat insulin intron A (RIA) was PCR'd out of plasmid p5'rIns (unknown origin) with primers GW150 (SEQ ID NO:19) and JF255 (SEQ ID NO:20). The PCR product was cut with *Bam*H1 and inserted into 5 *Bam*H1 linearized pJV7382, resulting in pJV7389.

(d) pJV7387 (CMV(RIA), HBV preS2 5'UTR and RBGpA)

pJV7384 was cut with *Bst*X1 and *Eco*R1 to generate an insert fragment containing the 3' end of the HBsAg coding region and RBGpA. 10 pJV7389 was cut with *Bst*X1 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7387.

Example 2. Construction of herpes simplex virus glycoprotein D antigen (HSVgD) vector panels

A number of plasmid expression vectors were constructed for 15 expression of HSVgD.

Starting materials

(a) pJV7334, a derivative of pWRG7284 (pJV 7284) that replaces the HBsAg coding sequence with an in-frame *Nhe*1 directly downstream of the ATG start codon, followed by a stuffer fragment with a *Bam*H1 20 immediately 5' of the HBV Enh

(b) pWRG7202 , a derivative of pGem3Z (Promega) with a stuffer fragment that allows the fusion of a coding sequence to the human tissue plasminogen activator (TPA) signal peptide downstream of a *Nhe*1 site.

(a) pJV7392(CMV(native intron), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

The coding region for HSV2 gD was PCR'd out of a viral DNA stock (Advanced Biotech, Inc, Columbia, MD) using primers DS1 (SEQ ID NO:21) and DA1 (SEQ ID NO:22) and was cut with *Nhe*1 and *Eco*R1 to generate an insert fragment. pWRG7202 was cut with *Nhe*1 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7391.

pJV7391 was cut with *Nhe*1 and *Bgl*2 to generate an insert fragment containing the HSV2 gD coding sequence. pJV7334 was cut with *Nhe*1 and *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7392. This vector consists of the following expression elements: the hCMV immediate early promoter sequence, the first exon, first intron, and a partial second exon of the hCMV major immediate early gene, the 5'-UTR from HBsAg, the coding sequence for HSV2 gD gene, the 3'-UTR from HBsAg, and RBGpA.

(b) pJV7399 (CMV(no intron), HBsAg 3'UTR HBsAg 5'UTR and RBGpA)

An intronless version of pJV7392 was constructed as follows. pJV7384 was cut with *Hind*3 and *Nde*1 to isolate an insert fragment containing the 5' ends of the kanamycin resistance gene and the CMV promoter. pJV7384 was cut with *Nde*1 and *Ssp*1 to isolate an insert fragment containing the 3' end of the CMV promoter, the CMV exon1/2 and 5' end of the 5'-UTR from HBsAg. These insert fragments were inserted into pJV7392 from which the *Hind*3-*Ssp*1 fragment was removed, resulting in pJV7399.

(c) pJV7400 (CMV(RIA), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

A RIA version of pJV7392 was constructed as follows. pJV7384 was cut with *Hind*3 and *Nde*1 to isolate an insert fragment containing 5 the 5' ends of the kanamycin resistance gene and CMV promoter. pJV7387 was cut with *Nde*1 and *Ssp*1 to isolate an insert fragment containing the 3' end of the CMV promoter, the CMV exon1/2(partial), RIA, and 5' end of the 5'-UTR from HBsAg. These insert fragments were inserted into pJV7392 from which the *Hind*3-*Ssp*1 fragment was 10 removed, resulting in pJV7400.

(d) pJV7401 (CMV(no intron), HBsAg 5'UTR and RBGpA)

A 3'-UTR-less version of pJV7399 was constructed as follows. pJV7391 was cut with *Bsp*120I and *Bgl*2 to isolate an insert fragment containing the 3' end of the HSV2 gD gene. pJV7284 was cut with *Bgl*2 15 and *Eco*R1 to isolate the RBGpA signal. These insert fragments were inserted into pJV7399 from which the *Bsp*120I-*Eco*R1 fragment was removed, resulting in pJV7401.

(e) pJV7402 (CMV(RIA), HBsAg 5'UTR and RBGpA)

A 3'-UTR-less version of pJV7400 was constructed as follows. 20 pJV7391 was cut with *Bsp*120I and *Bgl*2 to isolate an insert fragment containing the 3' end of the HSV2 gD gene. pJV7284 was cut with *Bgl*2 and *Eco*R1 to isolate the RBGpA signal. These insert fragments were ligated into pJV7400 from which the *Bsp*120I-*Eco*R1 fragment was removed, resulting in pJV7402.

Example 3. Construction of Flu M2 antigen vector panels(a) pJV7450 (CMV(no intron), HBsAg 5'UTR and RBGpA)

A coding region for Flu M2 was PCR'd out of plasmid pFL-M2 (Joel Haynes, PJV) using primers JF301 (SEQ ID NO:23) and JF302 (SEQ ID NO:24) and was cut with *Nhe*1 and *Bgl*2 to generate an insert fragment. pJV7401 was cut with *Nhe*1 and *Bgl*2 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7450.

(b) pJV7452 (CMV(no intron), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

A 3'UTR fragment was PCR'd out of pJV7389 with primers JF84 (SEQ ID NO:25) and JF225 (SEQ ID NO:26) was cut with *Bsp*120I, filled with T4 DNA polymerase, and linkerered with *Bgl*2 linkers (cat# 1036, New England Biolabs). The fragment was then cut with *Bgl*2 and *Eco*R1 to isolate an insert fragment containing the 3'UTR of HBsAg and the RBGpA region. pJV7450 was cut with *Bgl*2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7452.

(c) pJV7458 (CMV(RIA) HBsAg 5' UTR and RBGpA)

A version of pJV7450 containing the RIA was constructed as follows: pJV7389 was cut with *Bam*H1 to isolate a RIA containing insert fragment. pJV7450 was cut with *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7458.

(d) pJV7468 (CMV(RIA), HBsAg 3'UTR ,HBsAg 5'UTR and RBGpA)

A version of pJV7458 containing the 3'UTR of HBsAg was constructed as follows: pJV7452 was cut with *Bgl*2 and *Eco*R1 to 5 produce an insert fragment containing the HBsAg 3'UTR and RBGpA. pJV7458 was cut with *Bgl*2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7468.

Example 4. Construction of Beta-gal vector panels

(a) pJV7488 (CMV(no intron), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

CMV-beta (Clontech) was PCRed with primers JF335 (SEQ ID NO:27) and JF336 (SEQ ID NO:28) and cut with *Nhe*1 and *Bgl*2 to isolate an insert fragment coding for beta-galactosidase. pJV7452 was cut with *Nhe*1 and *Bgl*2 to generate a vector fragment into which the 15 insert fragment was ligated, resulting in pJV7488.

(b) pJV7533 (CMV(no intron), HBsAg 5'UTR and RBGpA)

pJV7450 was cut with *Bgl*2 and *Eco*R1 to isolate an insert fragment containing the RBGpA. pJV7488 was cut with *Bgl*2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, 20 resulting in pJV7533.

(c) pJV7551(CMV(RIA/*Nhe*1), HBsAg 3'UTR, HBsAg 5'UTR and RBGpA)

pJV7530 (see Example 5) was cut with *Xho*1 and *Bam*H1 to isolate an insert fragment containing the CMV promoter through RIA. 25 pJV7488 was cut with *Xho*1 and *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7551.

(d) pJV7552(CMV(RIA/Nhe1), HBsAg 5'UTR and RBGpA)

pJV7530 was cut with *Xho*1 and *Bam*H1 to isolate an insert fragment containing the CMV promoter through RIA. pJV7533 was cut with *Xho*1 and *Bam*H1 to generate a vector fragment into which the 5 insert fragment was ligated, resulting in pJV7552.

Example 5. Construction of pJV Expression (pJV7563)(a) pJV7496

pJV7389 was PCR'd with primers JF357 (SEQ ID NO:29) and JF365 (SEQ ID NO: 30), treated with T4 DNA polymerase to blunt the 10 ends, and cut with *Sa*1 to isolate an insert fragment coding for kanamycin resistance. pJV7389 was cut with *Ava*1, treated with T4 DNA polymerase to blunt the ends, and cut with *Sa*1 to isolate a vector fragment into which the insert fragment was ligated, resulting in pJV7496.

15 (b) pJV7530

pJV7389 was PCR'd with primers JF393 (SEQ.ID NO:31) and JF406 (SEQ ID NO:32) and cut with *Bg*/2 and *Bam*H1 to isolate an insert fragment containing the RIA devoid of an internal *Nhe*1 site. pJV7496 was cut with *Bam*H1 to prepare a vector fragment into which the insert 20 fragment was ligated, resulting in pJV7530.

(c) pJV7549

pJV7468 was cut with *Bam*H1 and *Eco*R5 to isolate an insert fragment containing M2 and part of the HBV 3'ENH. pJV7530 was cut with *Bam*H1 and *Eco*R5 to prepare a vector fragment into which the 25 insert fragment was ligated, resulting in pJV7549.

(d) pJV7563

Primers JF256 (SEQ ID NO:33) and JF257 (SEQ ID NO:34) were annealed to prepare an insert fragment consisting of a multiple cloning site. pJV7549 was cut Nhe1 and Bgl2 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7563. A pJV7563 plasmid map is provided in Figure 12. The base composition for the pJV7563 plasmid is provided in Figure 13. The components and their position in the plasmid pJV7563 are as follows:

1-44 Transposon 903 sequences

10 45-860 Kanamycin resistance coding sequence from
Transposon 903

861-896 Transposon 903 sequences

897-902 Sal1 site

903-1587 CMV promoter

15 1588-1718 untranslated leader sequence from the immediate-early
gene of CMV

1719-1724 Fusion of BamH1 and BgIII restriction enzymes

1725-1857 Rat insulin intron A

1858-1863 BamH1 site

20 1864-1984 HBV surface antigen 5'- untranslated leader

1985-1993 Synthetic start codon/ Nhe1 cloning site

1994-2011 Synthetic cloning sites

2012-2544 HBV enhancer

2545-2555 Old vector sequence. No hits against NCBI databases

2556-2686 Rabbit beta-globin polyadenylation region

2687-3759 pUC19 vector sequence

Example 6. Construction of Signal Peptide Expression Panels using

5 Human Secreted Alkaline Phosphatase (SEAP) and Human IgG Fc
Fragment (hFc) as Model Antigens

(i) pJV7507 (hTPAAsp and SEAP)

pSEAP-Basic (Clontech) was PCR'd with primers JF320 (SEQ ID NO:35) and JF321 (SEQ ID NO:36) then cut with *Nhe*1 and *Bg*II to 10 isolate an insert fragment consisting of the human SEAP fragment. pJV7079 (Macklin, et.al.) was cut with *Nhe*1 and *Bg*II to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7507.

(ii) pJV7508 (hTPAAsp and hFc)

15 Human DNA was PCR'd with primers JF386 (SEQ ID NO:37) and FcAS (SEQ ID NO:38) then cut with *Nhe*1 and *Bg*II to isolate an insert fragment consisting of the human IgG Fc fragment. pJV7079 was cut with *Nhe*1 and *Bg*II to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7508.

20 (iii) Preparation of Aprotinin Signal Peptide Coding Sequence

Synthetic oligo JF354 (SEQ ID NO:39) was PCR'd with primers JF355 (SEQ ID NO:40) and JF356 (SEQ ID NO:41) to generate the coding sequence for the aprotinin signal peptide.

(iv) Preparation of Tobacco Extensin Signal Peptide CodingSequence

Synthetic oligo JF348 (SEQ ID NO:42) was PCR'd with primers JF349 (SEQ ID NO:43) and JF350 (SEQ ID NO:44) to generate the 5 coding sequence for the tobacco extensin signal peptide.

(v) Preparation of Chicken Lysozyme Signal Peptide CodingSequence

Synthetic oligo JF351 (SEQ ID NO:45) was PCR'd with primers JF352 (SEQ ID NO:46) and JF353 (SEQ ID NO:47) to generate the 10 coding sequence for the chicken lysozyme signal peptide.

(a) Flu M2 antigen signal peptide panels

pJV7499 (CMV(no intron), HBsAg5'UTR, RBGpA, aprotinin s.p.)

pJV7497 (CMV(no intron), HBsAg5'UTR, RBGpA, tobacco extensin s.p.)

15 PJV7500 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken lysozyme s.p.)

Coding sequences for the signal peptides were cut with *Spe*1 and *Nhe*1 to isolate insert fragments. pJV7450 was cut with *Nhe*1 to prepare a vector fragment into which the insert fragments were ligated, resulting 20 in pJV7499 (aprotinin), pJV7497 (tobacco extensin), and pJV7500 (chicken lysozyme).

(b) SEAP signal peptide panels

pJV7513 (CMV(no intron), HBsAg5'UTR, RBGpA, aprotinin s.p.)

pJV7512 (CMV(no intron), HBsAg5'UTR, RBGpA, tobacco 25 extensin s.p.)

pJV7510 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken
lysosome s.p.)

pJV7499, 7497, and 7500 were cut with *Xho*1 and *Nhe*1 to
isolate an insert fragment consisting of the CMV promoter through the
5 signal peptide coding sequence of the plasmids. pJV7507 was cut with
*Xho*1 and *Nhe*1 to prepare a vector fragment into which the insert
fragments were ligated, resulting in pJV7513 (aprotinin), pJV7512
(tobacco extensin), and pJV7510 (chicken lysozyme).

(c) hFc signal peptide panels

10 pJV7524 (CMV(no intron), HBsAg5'UTR, RBGpA, aprotinin s.p.)

pJV7525 (CMV(no intron), HBsAg5'UTR, RBGpA, tobacco
extensin s.p.)

pJV7526 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken
lysosome signal peptide)

15 pJV7499, 7497, and 7500 were cut with *Xho*1 and *Nhe*1 to
isolate an insert fragment consisting of the CMV promoter through the
signal peptide coding sequence of the plasmids. pJV7508 was cut with
*Xho*1 and *Nhe*1 to prepare a vector fragment into which the insert
fragments were ligated, resulting in pJV7524 (aprotinin), pJV7525
20 (tobacco extensin), and pJV7526 (chicken lysozyme).

Example 7. Construction of Human Secreted Alkaline Phosphatase
(SEAP) panels

(a) pJV7531 (CMV(no intron), HBsAg5'UTR, RBGpA, chicken
lysosome s.p.)

25 pJV7510 was cut with *Sa*1 and *Bgl*2 to isolate an insert fragment
containing the CMV promoter through lysozyme signal peptide. pJV7450

was cut with *Sa*/1 and *Bg*/2 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7531.

(b) pJV7554 (CMV(RIA/*Nhe*l), HBsAg5'UTR, RBGpA, chicken lysosome s.p.)

5 pJV7530 was cut with *Xho*1 and *Bam*H1 to isolate an insert fragment containing the CMV promoter through RIA. pJV7531 was cut with *Xho*1 and *Bam*H1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7554.

10 (c) pJV7568 (CMV(no intron), HBsAg 3'UTR, HBsAg5'UTR, RBGpA, chicken lysosome s.p.)

pJV7563 was cut with *Bg*/2 and *Eco*R1 to isolate an insert fragment containing the HBV 3'-UTR and RBGpA. pJV7531 was cut with *Bg*/2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7568.

15 (d) pJV7572 (CMV(RIA/*Nhe*l), HBsAg 3'UTR, HBsAg5'UTR, RBGpA, chicken lysosome s.p.)

20 pJV7563 was cut with *Bg*/2 and *Eco*R1 to isolate an insert fragment containing the HBV 3'UTR and RBGpA. pJV7554 was cut with *Bg*/2 and *Eco*R1 to generate a vector fragment into which the insert fragment was ligated, resulting in pJV7572.

Example 8. Construction of Beta-gal and HBsAg Vectors Using the Chicken Keratin and Chicken Cardiac Actin Introns

(a) pJV7557 (Beta-gal, CMV(cA intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

26 Chicken DNA was PCR'd with primers JF430 (SEQ ID NO:48) and JF442 (SEQ ID NO:49) and cut with *Bg*/2 and *Bam*H1 to isolate an

insert fragment consisting of the intron and flanking exon sequences from chicken cardiac actin. pJV7488 was cut with *Bam*H1 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7557.

5 (b) pJV7558 (Beta-gal, CMV(cK intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

Chicken DNA was PCR'd with primers JF421 (SEQ ID NO:50) and JF444 (SEQ ID NO:51) and cut with *Bgl*II and *Bam*H1 to isolate an insert fragment consisting of the intron and flanking exon sequences from the 10 chicken keratin gene. pJV7488 was cut with *Bam*H1 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7558.

(c) pJV7578 (HBsAg, CMV(cA intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

15 pJV7557 was cut with *Sa*/1 and *Bam*H1 to isolate an insert fragment consisting of the CMV promoter through intron regions. pJV7496 was cut with *Sa*/1 and *Bam*H1 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7558.

(d) pJV7579 (HBsAg, CMV(cK intron), HBsAg3'UTR, HBsAg5'UTR and RBGpA)

pJV7558 was cut with *Sa*/1 and *Bam*H1 to isolate an insert fragment consisting of the CMV promoter through intron regions. pJV7496 was cut with *Sa*/1 and *Bam*H1 to prepare a vector fragment into which the insert fragment was ligated, resulting in pJV7579.

Example 9. In-Vitro Analysis of Antigen Expression by HBsAgVector Panels

On day one, SCC15 (ATCC) or B16 (origin unknown, versions available at ATCC) cells were plated on 6 well tissue culture plates at 20-
5 40% confluency, and allowed to grow overnight in an incubator. The host cells were propagated in media recommended by ATCC.

On day two, the transfection reaction was performed. For each vector to be tested, 20 μ l of Lipofectin® re agent (Life Technologies Inc, Grand Island, NY) was added to 180 μ l of Optimem® media (Life 10 Technologies, Grand Island, NY), and allowed to incubate at room temperature for 45 minutes. For each vector to be tested, 2 μ g of vector was mixed into 200 μ l of Optimem® at 40 minutes. At 45 minutes, the vector and Lipofectin® solutions were mixed together and allowed to sit at room temperature for an additional 10 minutes. During this final 15 incubation, the plated host cells were removed from the incubator and washed twice with Optimem® media. At 10 minutes, 1.6ml of Optimem® was added to the Lipofectin® / vector mix, and 1ml of the resultant mix was added to each of two cell wells. The host cells were returned to the incubator and allowed to sit undisturbed for 5 hours, at 20 which point the Lipofectin® / vector mix was removed and replaced by standard cell maintenance media.

At 18 to 24 hours after the media change, from 50 to 100 μ l of cell maintenance media was removed from the tissue culture plates and analyzed for antigen expression by placing the samples into reaction 25 vessels provided in the AUSZYME® Monoclonal Diagnostic Kit (Abbott Laboratories, Abbott Park, IL). The volume of the test samples was brought to a volume of 200 μ l with PBS, then 50 μ l of conjugate and a reaction bead were added to each sample. The vessel is incubated for 80 minutes at 40°C, after which the wells were washed clean of all liquid

reaction components. The beads were transferred to new tubes after which 300 μ l of color development buffer was added. At 30 minutes, the color development reaction was stopped by the addition of 1M sulfuric acid, and the absorbance of the reaction was measured at 490nm. The 5 data shown in Figure 1 is the average absorbance readings of the duplicate wells from two experiments.

As shown in Figure 1, the addition of RIA, the HBV 3'UTR or both elements to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of HBsAg in SCC15 cells. As shown in 10 Figure 2, the addition of either the chicken keratin or the chicken cardiac actin intron to a base vector (CMV promoter, exon, HBV 3'UTR and polyadenylation region) increased expression of HBsAg in SCC15 cells.

Example 10. *In-Vitro* Analysis of Antigen Expression by Beta-gal Vector Panels

15 SSC-15 or B16 host cells were transfected as described in Example 9.

At eighteen to forty hours after the media change, the media supernatants were removed and the cells were washed with PBS. After removal of the wash, the cells were lysed by incubating the cells in 20 500 μ l lysis buffer (50mM NaPO₄, 0.1% Triton X-100, pH 7) for 5 minutes, followed by physically scraping the cells off the plastic dish. The lysates were microfuged for two minutes to remove cell debris, and 10 to 25 μ l of the cleared lysate were added to 500 μ l of reaction buffer (80ug/ml α -nitrophenyl galactopyranoside, 50mM NaPO₄, pH 7) and 25 incubated at 37°C for 10 to 20 minutes. The reaction was stopped by the addition of 500 μ l of 1M Na₂CO₃ and read at 405 nm. Data is presented as the ratio of the expression of enhanced (containing an intron, HBVenh, or both) vector to a base vector.

The addition of RIA, the HBV 3'UTR or both elements to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of beta-gal in both cell lines. The results for SCC15 cells are shown in Figure 3. Addition of either the chicken keratin or the chicken 5 cardiac actin intron to a base vector (CMV promoter, exon, the HBV 3'UTR and polyadenylation region) increased the expression of beta-gal in both cell lines. Results for B16 cells are shown in Figure 2.

Example 11. In-Vitro Analysis of Antigen Expression by HSV gD
Vector Panel

10 SCC15 or B16 host cells were transfected as described in Example 9. Eighteen hours post transfection, plates were placed on ice for 15 minutes. Each well was then washed with 2 ml of PBS (Biowhittaker, Walkerville, MD). Cells were fixed with 0.05% gluteraldehyde (Polysciences Inc, Warrington, PA) diluted in PBS and incubated for 30 15 minutes at room temperature. All subsequent incubations lasted 1 hour at room temperature and washes between each incubation were as stated above. The plates were blocked with 2 ml of 5% dry milk (Bio Rad Laboratories, Melville, NY) in PBS. Incubations with 1 ml of a 1:1000 dilution of anti-gD monoclonal (ABI, Columbia, MD) in 2% dry milk / PBS 20 / 0.05% Tween-20 (Sigma, St. Louis, MO) and 1 ml of a 1:2500 dilution of goat anti-mouse HRP (KPL, Gaithersburg, MD) in PBS / 0.1% Tween-20 followed. Color was developed using 1 ml of TMB microwell substrate (BioFX, Owings Mills, MD). The reactions were stopped with 1M H₂SO₄, the liquid was transferred to plastic cuvettes and the optical 25 density read at 450nm. Data is presented as the ratio of the expression of enhanced (containing an intron, HBVenh, or both) vector to a base vector.

Addition of RIA with or without the HBV 3'UTR to a base vector (CMV promoter, exon and polyadenylation region) increased the

expression of HSV gD in both cell lines. Results for SC15 cells are shown in Figure 4.

Example 12. In-Vitro Analysis of Antigen Expression by SEAP

Vector Panels

5 SCC15 or B16 host cells were transfected as described in Example 9. At eighteen to fourty hours after the media change, the media supernatants were removed and heated at 70°C for 30 minutes. 10 to 25 μ l of the heat-inactivated supernatants were incubated for 5 minutes with 1/10th volume of 100mM L-homoarginine. 500 μ l of alkaline phosphatase reaction buffer (cat # 172-1063, Bio-Rad, prepared according to instructions) were added to the lysates and incubated at 37°C for 10 to 20 minutes. The reaction was stopped by the addition of 500 μ l of 1M NaOH and read at 405 nm. Data is presented as the ratio of the expression of enhanced (containing an intron, HBVen, or both) 10 vector to a base vector, or the ratio of the expression of the experimental signal peptides to the human TPA signal peptide vector.

15

As shown in Figure 5, the addition of RIA, the HBV 3'UTR or both elements to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of SEAP in B16 cells. Unexpectedly, 20 only the addition of the HBV 3'UTR to a base vector (CMV promoter, exon and polyadenylation region) increased the expression of SEAP in SCC15 cells.

Addition of signal peptides from either bovine aprotinin, chicken lysozyme, or tobacco extensin to the N-terminus of mature SEAP allowed 25 for efficient secretion of SEAP into cell media supernatants of both cell lines. Results for B16 cells are shown in Figure 6.

Example 13. In-Vitro Analysis of Antigen Expression by Human IgG
Fc Fragment Signal Peptide Panel

SCC15 or B16 host cells were transfected as described in Example 9. The media supernatants were removed from eighteen to forty hours 5 after the media change.

ELISA plates (Costar) were incubated overnight at 4°C with 100 μ l of goat anti-human IgG (Sigma #I3382, 1/1000 dilution in carbonate coating buffer) per well. All subsequent incubations lasted 1 hour at room temperature with washes (10mM Tris, 150mM NaCl, 0.1% Brij-35, 10 pH8.0) between each incubation. The wells were then blocked with 100 μ l of 5 % dry in PBS, followed by incubation with serially diluted media supernatants in dilution buffer (2% dry milk, PBS, 0.05% Tween-20). This was followed by incubation with 100 μ l of goat anti-human IgG/HRP (Sigma #A6029, 1/5000 dilution in dilution buffer) per 15 well, followed by color development using 100 μ l of TMB microwell substrate. The reactions were stopped with 100 μ l of 1M H₂SO₄, and read at 450 nm. Data is presented as the ratio of the expression of the experimental signal peptides to the human TPA signal peptide vector.

Addition of signal peptides from either bovine aprotinin, chicken 20 lysozyme, or tobacco extensin to the N-terminus of the human Fc fragment allowed for efficient secretion of hFc into cell media supernatants of both cell lines. Results for B16 cells are shown in Figure 6.

Example 14. Use of the HBsAg, HSVgD and Flu-M2 plasmid expression vectors for immunisation of mice

a. Preparation of cartridges of immunisation

For each plasmid to be tested, 25mg of 2 micron gold powder was weighed into a microfuge tube. After the addition of a 250 μ l aliquot of 50 mM spermidine (Aldrich Chemical, Inc, Milwaukee, WI), the tube was vortexed and briefly sonicated. The gold was microfuged out, and the spermidine replaced by a fresh 100 μ l aliquot. The gold was resuspended by vortexing, after which 25 μ g of DNA was added to the tube and mixed. While the tube was lightly vortexed, 100 μ l of 10% CaCl (Fujisawa USA, Inc, Deerfield, IL) was added to precipitate the DNA onto the gold beads. The precipitation reaction was allowed to proceed for 10 minutes on the benchtop, after which the gold was collected by a brief microfuge spin and washed three times with absolute ethanol (Spectrum Quality Products, Inc, Gardena, CA) to remove excess precipitation reagents. The washed gold/DNA complex was then resuspended in 3.6 ml of 0.05 mg/ml polyvinylpyrrolidone (360KD, Spectrum Quality Products, Inc, Gardena, CA) in absolute ethanol. This slurry was then injected into a Tefzel $\text{^{\text{a}}}$ tube (McMaster-Carr, Chicago, IL) located in a tube turner (PowderJect Vaccines) which coats the inside of the Tefzel $\text{^{\text{a}}}$ tube with the gold/DNA complex. After the tube turning procedure was completed, the tube was cut into 0.5" "shots" of vaccine which were loaded into the XR1 device (PowderJect Vaccines) for delivery to the mice.

b. Vaccination Procedure

Four to six week old mice were anesthetized with a mixture of Ketasetâ (Fort Dodge) and Rompunâ (Bayer). The bellies were shaved with a pair of electric clippers to remove hair, and two non-overlapping 5 "shots" of vaccine were delivered via the XR1 device (450psi) to the shaved area. The animals were returned to their cages and bled at six weeks post vaccination. Balb/c mice were used to evaluate the HBsAg expression vectors, and Swiss Webster mice were used to evaluate the HSV-gD and Flu M2 expression vectors.

10 Analysis of Sera for Anti-HBsAg Antibodies

At six weeks, blood samples were harvested from vaccinated animals. A volume of serum isolated from these samples was placed into wells of a reaction vessel supplied with the AUSAB® EIA Diagnostic Kit (Abbott Laboratories, Abbott Park, IL). The volume of sera added 15 depended upon the antibody titer of the sample, and the sample was diluted with sample dilution buffer to fall within values obtainable with a quantification panel panel. 200µl from each vial of the AUSAB® Quantification Panel (Abbott Laboratories, Abbott Park, IL) was added to wells of the reaction vessel. To each well a bead was added, after which 20 the vessel was sealed and incubated for two hours at 40°C. The wells were then washed of all liquid reaction components. To each washed well was added 200µl of conjugate mix, after which the vessel was sealed and incubated for two hours at 40°C. The wells were then washed of all liquid reaction components. The beads were transferred to 25 new tubes after which 300µl of color development buffer was added. At 30 minutes, the color development reaction was stopped by the addition of 1M sulfuric acid, and the absorbance of the reactions was measured at 490nm in a Quantum II® spectrophotometer (Abbott Laboratories, Abbott Park, IL) . This spectrophotometer calculates the antibody levels

of a sample by comparing the absorbance of the sample with a standard curve generated with the quantification panel. These antibody levels were then corrected for dilution factors. The data shown in Figure 7 are the geometric mean titers of all animals vaccinated with a particular 5 vector.

Analysis of Sera for Anti-Flu M2 Antigen Antibodies

96-well Costar medium-binding ELISA plates (Fisher Scientific, Pittsburgh, PA) were coated with a synthetic Flu M2 peptide (QCB / Biosource, Hopkinton, MA) at a concentration of 1ug/ml in PBS 10 (Biowhittaker, Walkerville, MD) and incubated overnight at 4°C. The plates were washed three times with 10mM Tris (Sigma, St. Louis, MO) / 150mM NaCl (Fisher Scientific) / 0.1% Brij-35 (Sigma), then blocked with 5% dry milk (Bio Rad Laboratories, Melville, NY) in PBS for 1 hour at room temperature. All subsequent incubations were at room 15 temperature for one hour and washes between each incubation were as stated above. Sample mouse sera, a standard (high titer, anti-M2 mouse sera) and a negative control (anti-HBsAg mouse sera) were diluted in 2% dry milk / PBS / 0.05% Tween-20 (Sigma) and incubated in the ELISA plates. Goat anti-mouse IgG (H + L) biotin conjugated antibody (Southern 20 Biotechnology Associate, Birmingham, AL) diluted 1:8000 in 2% dry milk / PBS / 0.05% Tween-20 and streptavidin-horseradish peroxidase conjugate (Southern Biotechnology) diluted 1:8000 in PBS / 0.1% Tween-20 followed. Color was developed using TMB substrate (BioFX, Owings Mills, MD). The reactions were stopped with 1M H₂SO₄ and the 25 plates read at 450nm with an Emax precision microplate reader (Molecular Devices, Sunnyvale, CA). SoftMax Pro 4.1 software (Molecular Devices) was used to calculate endpoint titers using a four-parameter analysis. Titers were normalized to the standard serum, which had a pre-determined titer, to minimize assay-to-assay and plate-to-plate 30 variation. Results are shown in Figure 7.

Analysis of Sera for Anti-HSV gD Antigen Antibodies

96-well Costar medium-binding ELISA plates (Fisher Scientific, Pittsburgh, PA) were coated with HSV gD (Viral Therapeutics, Ithaca, NY) protein at a concentration of 1ug/ml in PBS (Biowhittaker, Walkerville, MD) and incubated overnight at 4°C. The plates were washed three times with 10mM Tris (Sigma, St. Louis, MO) / 150mM NaCl (Fisher Scientific) / 0.1% Brij-35 (Sigma), then blocked with 5% dry milk (Bio Rad Laboratories, Melville, NY) in PBS for 1 hour at room temperature. All subsequent incubations were at room temperature for one hour and washes between each incubation were as stated above.

Sample mouse sera, a standard (high titer, anti-gD mouse sera) and a negative control (anti HBsAg mouse sera) were diluted in 2% dry milk / PBS / 0.05% Tween-20 (Sigma) and incubated in the ELISA plates. Goat anti-mouse IgG (H+L) biotin conjugated antibody (Southern Biotechnology Associate, Birmingham, AL) diluted 1:8000 in 2% dry milk / PBS / 0.05% Tween-20 and streptavidin-horseradish peroxidase conjugate (Southern Biotechnology) diluted 1:8000 in PBS / 0.1% Tween-20 followed. Color was developed using TMB substrate (BioFX, Owings Mills, MD). The reactions were stopped with 1M H₂SO₄ and the plates read at 450nm with an Emax precision microplate reader (Molecular Devices, Sunnyvale, CA). SoftMax Pro 4.1 software (Molecular Devices) was used to calculate endpoint titers using a four-parameter analysis. Titers were normalized to the standard serum, which had a pre-determined titer, to minimize assay-to-assay and plate-to-plate variation. Results are shown in Figure 7.

Accordingly, novel nucleic acid constructs, compositions comprising these constructs, and nucleic acid immunization techniques using these constructs have been described. Although preferred embodiments of the subject invention have been described in some detail, it is understood that obvious variations can be made without

departing from the spirit and the scope of the invention as defined by the appended claims.

CLAIMS

1. A nucleic acid construct comprising:

- (i) a chimeric promoter sequence which comprises:
 - (a) a hCMV immediate early promoter sequence;
 - (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
 - (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene;
- (ii) a coding sequence in operable linkage with the chimeric promoter;
- (iii) a non-translated leader sequence which is selected from the HBVpreS2 antigen sequence, HBV e-antigen sequence and HSV type 2gD antigen sequence and which is in operable linkage with the chimeric promoter; and
- (iv) an enhancer sequence which is derived from a 3' untranslated region (UTR) of a HBsAg sequence or of a simian CMV immediate early gene sequence, which is in operable linkage with the chimeric promoter and which is downstream of the coding sequence.

2. A nucleic acid construct according to claim 1 wherein:

- (i) the hCMV immediate early promoter sequence (a) comprises SEQ ID NO:1, a functional variant thereof with at least 80% homology or a functional fragment of either; and/ or

(ii) exon sequence (b) comprises SEQ ID NO:2, a functional variant thereof with at least 80% homology or a functional fragment of either.

5 3. A nucleic acid construct according to claim 1 or 2 wherein the heterologous intron (c) comprises a sequence selected from rat insulin gene intron A sequence, chicken keratin gene intron A sequence, chicken cardiac actin gene intron A sequence, a functional variant with at least 80% homology to a said intron A sequence and a functional fragment of 10 a said intron A sequence or of a said functional variant thereof.

4. A nucleic acid construct according to claim 3 wherein the rat insulin gene intron A sequence comprises SEQ ID NO: 3, a functional variant thereof with at least 80% homology, or a functional fragment of 15 either.

5. A nucleic acid construct according to any one of the preceding claims wherein:

(i) the non-translated leader sequence is selected from SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO:7, a functional variant with at least 80% homology to any one of SEQ ID NOS: 5 to 20 7 or a functional fragment of any one of SEQ ID NOS: 5 to 7 or of a said variant thereof; and /or

(ii) the enhancer sequence is selected from SEQ ID NO: 8, SEQ ID NO: 9, a functional variant with at least 80% homology to SEQ ID NO: 8 or SEQ ID NO: 9 or a functional fragment 25

of SEQ ID NO: 8 or SEQ ID NO: 9 or of a said variant thereof.

6. A nucleic acid construct according to any one of the preceding 5 claims which comprises:

- (i) a polyadenylation sequence; and/or
- (ii) a sequence encoding a signal peptide.

7. A nucleic acid construct according to claim 6 wherein:

10 (i) the polyadenylation sequence is a polyadenylation sequence of a gene selected from the rabbit β -globin gene, human papilloma virus (HPV) early or late gene, the HSV-2gB gene, a simian CMV immediate early gene and HSVgD late gene, a functional variant with at least 80% homology to any one of said sequences or a functional fragment of any one of said

15 polyadenylation sequences or said function variants thereof; and/or

(ii) the signal peptide is selected from human tissue plasminogen activator signal peptide (hTPAAsp), aprotinin signal peptide, tobacco extensin signal peptide, chicken lysozyme signal peptide, a functional variant with at least 80% homology to any one of said sequences or a functional fragment of any of said sequence encoding the signal peptide or said functional variant thereof.

25

8. A nucleic acid construct according to claim 7 wherein the

polyadenylation sequence is selected from SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, a functional variant with at least 80% homology to any one of SEQ ID NOS: 10 to 13 or a functional fragment of any one of SEQ ID NOS: 10 to 13 or of a said variant 5 thereof.

9. A nucleic acid construct according to any one of the preceding claims wherein the coding sequence encodes an antigen.

10 10. A nucleic acid construct according to claim 9 wherein the antigen is an antigen of a viral, bacterial, parasitic or fungal pathogen, an allergy antigen or a cancer antigen.

11. A nucleic acid construct according to claim 10 wherein the viral 15 antigen is a HPV, HIV, HSV1, HSV2, influenza virus, Hepatitis A virus or Hepatitis B virus antigen.

12. A nucleic acid construct according to claim 11 wherein the antigen is HBsAg.

20 13. A nucleic acid construct according to any one of claims 1 to 8, wherein the coding sequence encodes an ADP ribosylating bacterial subunit A and/or B subunit, an active homolog of one or both said subunits wherein each homolog has at least 80% homology to the 25 respective said subunit or an active fragment of a said subunit or of a said active homolog thereof.

14. A nucleic acid construct according to claim 13 wherein the bacterial subunit is selected from Cholera toxin subunit A, Cholera toxin subunit B, *E.coli* heat labile toxin subunit A, *E.coli* heat labile toxin subunit B, an active homolog having at least 80% homology to any one of said subunits or an active fragment of any one of said subunits or said active homologs.

15. A nucleic acid construct according to any one of the preceding claims which is a DNA plasmid.

10

16. Coated particles, suitable for delivery from a particle-mediated delivery device, which particles comprise carrier particles coated with a nucleic acid construct according to any one of the preceding claims.

15

17. Coated particles according to claim 16 wherein the carrier particles are gold or tungsten.

18. A dosage receptacle for a particle mediated delivery device comprising coated particles according to claim 16 or 17.

19. A particle mediated delivery device loaded with coated particles according to claim 16 or 17.

20. A particle mediated delivery device according to claim 19 which is a needleless syringe.

21. A pharmaceutical preparation comprising a nucleic acid construct according to any one of claims 1 to 15 and a pharmaceutically acceptable excipient.

5 22. A vaccine comprising a nucleic acid construct according to any one of claims 9 to 12.

23. A vaccine according to claim 22, further comprising an adjuvant construct according to claim 13 or 14.

10

24. An *in vitro* method of obtaining expression in mammalian cells of a polypeptide of interest, which method comprises transferring into said cells a nucleic acid construct according to any one of claim 1 to 15 or coated particles according to claim 16 or 17.

15

25. Use of a nucleic acid construct according to any one of claims 9 to 12 for the manufacture of a medicament for immunization.

20 26. Use according to claim 25, wherein the medicament is for immunization against infection by a pathogen, allergy or cancer.

27. Use according to claim 25 or 26 wherein the medicament is to be delivered by injection, transdermal particle delivery, inhalation, topically, orally, intranasally or transmucosally.

25

28. Use according to claim 27 wherein the construct is delivered by needleless injection.

29. A nucleic acid construct comprising a chimeric promoter sequence and a coding sequence in operable linkage with the chimeric promoter, wherein the chimeric promoter sequence comprises:

5 (a) a hCMV immediate early promoter sequence;

(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and

(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

10

30. A nucleic acid construct comprising:

(i) a promoter sequence;

(ii) a coding sequence operably linked to the promoter sequence (i); and

15 (iii) an enhancer sequence 3' of and operably linked to the coding sequence (ii);

wherein the enhancer sequence (iii) is derived from a 3'UTR of an HBsAg sequence or a 3'UTR of a simian CMV immediate early gene sequence, and the coding sequence (ii) is heterologous to the enhancer sequence.

20

31. A nucleic acid construct according to claim 30 wherein the promoter sequence (i) is selected from the hCMV immediate early promoter sequence, Pseudorabies virus promoter sequence and Rous sarcoma virus promoter sequence.

25

32. A nucleic acid construct comprising:

(i) a promoter sequence;

- (ii) a non-translated leader sequence derived from HBV preS2 antigen sequence, HBV e-antigen sequence or HSV type 2 gD antigen sequence; and
- 5 (iii) a coding sequence operably linked to (i) and (ii) wherein the coding sequence is heterologous to the non-translated leader sequence.

33. A nucleic acid construct according to any one of claims 29 to 32
10 wherein the coding sequences encodes an antigen as defined in any one of claims 9 to 12 or an ADP ribosylating bacterial subunit, active homolog, or active fragment as defined in claim 13 or 14.

34. A purified isolated chimeric promoter sequence which comprises:
15 (a) a hCMV immediate early promoter sequence;
(b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
(c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene.

20 35. Coated particles, suitable for delivery from a particle-mediated delivery device, which particles comprise carrier particles coated with a nucleic acid construct according to any one of claims 29 to 33.

25 36. Coated particles according to claim 35 wherein the carrier particles are gold or tungsten.

37. A dosage receptacle for a particle mediated delivery device comprising coated particles according to claim 35 or 36.

5 38. A particle mediated delivery device loaded with coated particles according to claim 35 or 36.

39. A particle mediated delivery device according to claim 38 which is a needleless syringe.

10

40. A pharmaceutical preparation comprising a nucleic acid construct according to any one of claims 29 to 33 and a pharmaceutically acceptable excipient.

15

41. A vaccine comprising a nucleic acid construct according to claim 33 which encodes an antigen.

20

42. A vaccine according to claim 41, further comprising an additional construct according to claim 34 which encodes a ADP ribosylating bacterial subunit, active homolog thereof or an active fragment thereof.

25

43. An *in vitro* method of obtaining expression in mammalian cells of a polypeptide of interest, which method comprises transferring into said cells a nucleic acid construct according to any one of claim 29 to 33 or coated particles according to claim 35 or 36.

44. Use of a nucleic acid construct according to claim 33 which encodes an antigen for the manufacture of a medicament for immunization.

5 45. Use according to claim 44, wherein the medicament is for immunization against infection by a pathogen, allergy or cancer.

10 46. Use according to claim 44 or 45 wherein the medicament is to be delivered by injection, transdermal particle delivery, inhalation, topically, orally, intranasally or transmucosally.

47. Use according to claim 46 wherein the construct is delivered by needleless injection.

15 48. A nucleic acid construct comprising:

- (i) a chimeric promoter sequence which comprises:
 - (a) a hCMV immediate early promoter sequence;
 - (b) exon 1 and at least a part of exon 2 of the hCMV major immediate early gene; and
 - (c) a heterologous intron provided in place of the intron A region of the hCMV major immediate early gene;
- (ii) a cloning site for insertion of a coding sequence in operable linkage with the chimeric promoter;
- (iii) a non-translated leader sequence which is from the HBVpreS2 antigen sequence, HBV e-antigen sequence and HSV type 2gD antigen sequence and

which is in operable linkage with the chimeric promoter; and

(v) an enhancer sequence which is derived from a 3' untranslated region (UTR) of a HBsAg sequence or of a simian CMV immediate early gene sequence, which is in operable linkage with the chimeric promoter and which is downstream of the cloning site.

49. A nucleic acid construct according to any one of claims 1 to 15, 10 29 to 33 or 48, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

50. Coated particles according to any one of claims 16, 17, 35 or 36, substantially as herein described with reference to and as illustrated in 15 any of the examples and accompanying drawings.

51. A dosage receptacle according to claim 18 or claim 37, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

20 52. A device according to any one of claims 19, 20, 38 or 39, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

25 53. A preparation according to claim 21 or claim 40, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

54. A vaccine according to any one of claims 22, 23, 41 or 42, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

5

55. A method according to claim 24 or claim 43, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

10 56. Use according to any one of claims 25 to 28, or 44 to 47, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.

15 57. A promoter sequence according to claim 34, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.




Figure 1:

2/13






Figure 2

3/13

**Figure 3**

4/13

**Figure 4**

5/13

Effect of RIA, ENH on SEAP Expression (Two Lines ,Three Reps per Line)






Figure 5

6/13

**Figure 6**

7/13

**Figure 7**





Figure 8

9/13

**Figure 9**




Figure 10




Figure 11

12/13




Figure 12

Figure 13

SEQUENCE LISTING

<110> JAMES FULLER
<120> NUCLEIC ACID CONSTRUCTS
<130> APF 33 (X.13149 GCW)

<160> 53

<170> PatentIn version 3.1

<210> 1
<211> 685
<212> DNA
<213> Human cytomegalovirus

<400> 1

aatattggot	attggccatt	gcatacgtt	tatctata	ataatatgt	a	catttatatt	60
ggctcatgtc	caatatgacc	gcacgttga	cattgattat	tgactagtt	a	ttaatagtaa	120
tcaattacgg	ggtoattagt	tcata	ccca	tatatggagt	tccgcgttac	ataacttacg	180
gtaaatggcc	cgcotggctg	accgc	ccaaac	gaccccccgc	cattgaogtc	aataatgaog	240
tatgttccca	tagtaacgccc	aatagg	gact	ttccattgac	gtcaatgggt	ggagtattt	300
cggtaaactg	cccacttggc	agta	catca	gtgtatcata	tgccaagtcc	gccccotatt	360
gacgtcaatg	acggtaaatg	gc	ccgcctgg	cattatgc	agtacatgac	cttacgggac	420
tttco	actt	ggcag	tacat	ctacgtt	gtcatcgct	ttaccatgg	480
tggcagtaca	ccaaatggcg	tggatagogg	tttgactc	ac	ggggatttcc	aagtctccac	540
cccattgacg	tcaatgggag	ttt	gttttgg	cacccaaatc	aacgggactt	tccaaaatgt	600
cgtataacc	ccgccccgtt	gacgcaa	atg	ggcggttaggc	gtgtacgg	tg ggaggttat	660
ataagcagag	ctcg	tttagt	gaa	cc			685

<210> 2
<211> 131
<212> DNA
<213> Human cytomegalovirus

<400> 2

gtcagatcgc	ctggagacgc	cattccacgt	gttttgac	ccatagaaga	cacccggacc	60	
gateccag	ccgcggccgg	gaa	cgggt	gca	tttggaaacgcg	gattccccgt	120
actcaccgtc	c						131

<210> 3
<211> 135

<212> DNA

<213> Rattus rattus

<400> 3

atcagcaagc aggtatgtac tctccagggt gggcctggct tccccagtca agactacagg	60
gatttgaggg acgctgtggg ctatcttctt acatgtacot tttgctagcc tcaaccctga	120
ctatcttcca ggtca	135

<210> 4

<211> 955

<212> DNA

<213> Artificial sequence

<220>

<223> Chimeric promoter sequence

<400> 4

aatattggct attggccatt gcatacgttg tatctatato ataataatgtt catttatattt	60
ggctcatgtc caaatatgacc gccatgttga cattgattat tgacttagtta ttaatagtaa	120
tcaattaaagg ggtoatttagt tcatacgccca tatatggagt tccgcgttac ataacttacg	180
gttaaatggcc cgcctggctg accgcaccaac gaccccccggc cattgaoctc aataatgacg	240
tatgttccca tagtaacgcc aataggact ttccattgac gtcaatgggt ggagtattta	300
cggtaaatgt cccacttggc agtacatcaa gtgtatcata tgccaaatgc gccccctatt	360
gacgtcaatg acggtaaatgt gcccgcgttgc cattatgccc agtacatgac ttacgggac	420
tttcctactt ggcagtagat ctacgttata gtcatcgctt ttaccatgggt gatgcgggtt	480
tggcagtaca ccaatggcgg tggatagcgg tttgacttac ggggatttcc aagtctccac	540
cccatgtacg tcaatggag tttgttttgg caccaaaatc aacgggactt tccaaaatgt	600
cgtataaaacc cccggccgtt gacgcaaatg ggoggttaggc gtgtacgggt ggaggttotat	660
ataaggcagag ctgttttagt gaaccgttag atgccttggaa gaogcoatcc acgtgtttt	720
gacccatata gaagacacccg ggacogatcc agccctccggc gcccggaaacg gtgcatttgg	780
acgcggatcc ccogtgccaa gagtactca cctgtccggat ctcagcaagc aggtatgtac	840
tctccagggt gggcctggct tccccagtca agactccagg gatttgaggg acgtgtggg	900
ctcttcttcc acatgtaccc ttgttgcctcaaccctga ctatcttcca ggtca	955

<210> 5

<211> 121

<212> DNA

<213> Hepatitis B virus

<400> 5

cagagtcaagg ggttgttatt ttccctgtgg tggcttcaagt taaaccctgc	60
tccgaatatt gccttcaca tctgtcaat ctccgcagg actggggacc ctgtgacgaa	120

<210> 6
 <211> 57
 <212> DNA
 <213> Herpes simplex virus
 <400> 6
 ataaagctgca ttgcgaacca ctatgcgccc ttttttgtgtt gcatcggtt tcaaggc

57

<210> 7
 <211> 48
 <212> DNA
 <213> Hepatitis B virus
 <400> 7
 ctttgtacta ggaggctgttta ggcataaaatt ggtatgttca ccaggcacc

48

<210> 8
 <211> 533
 <212> DNA
 <213> Hepatitis B virus
 <400> 8
 taacaaaaca aaaagatggg gttattccct aaacttcatg gttacgtaa ttggaaagttg
 ggggacattt ccacaagatc atattgtaca aaagatcaaa cactgttttta gaaaacttcc
 tgtaaacagg cttattgatt gggaaagtatg tcaaaggatt gttttttttt tggctttgc
 tgctccatcc acadaatgtt gatattctgc cttaatgcct ttgtatgtat gtatacaagc
 taaaacaggct ttcaatttttcc cggcaacttta caaggcctttt ctaagtaaaac agtacatgaa
 cttttacccc gttgtcgcc aacggcctgg totgtgcctt gttttgtcg acgcaaccc
 cactggatgg ggcttgcata taggcccattca ggcattgcgtt ggaacccccc tggotccctt
 ggcgatccat actggggaaac tcttagccgc ttgtttgtt cgcagccgggtt ctggagcaaa
 gctcatagga actgacaatt ctgtcgctt ctgcggaaa tatacatatgt ttc

60

120

180

240

300

360

420

480

533

<210> 9
 <211> 158
 <212> DNA
 <213> Simian cytomegalovirus

<400> 9
 gtcagacaga cagacagtta tatggcgttccctataac tctgcatttgc taacccatca

60

tagccagaca gttagcattg catctattga ttagttaota atgtattgtt accccccata 120
tgccatttgtc taactgtact aatgtatgtt attataacc 158

<210> 10
<211> 131
<212> DNA
<213> *Oryctolagus cuniculus*

<400> 10
gatcttttc cctctgcaa aaattatggg gacatcatga agcccttga gcatctgact 60
totggctaat aaaggaaatt tatttcatt gcaatagtgt gttgaaattt. tttgtgtctc 120
tcactcgaa g 131

<210> 11
<211> 204
<212> DNA
<213> Simian cytomegalovirus

<400> 11
atataatactatgttatac totatgatata acaaataata ctcataaaca ctatgtactt 60
ggtgttatgactcattattgt ctgggacttg gttgggactt ggttgggttgg gaagaatgtt 120
gtgcctgtac ttgtgtctgtc ctgtggatct caataaaatgt gactatgttc aaaacactaa 180
gtgccccccgt gtcttcttta acta 204

<210> 12
<211> 163
<212> DNA
<213> Herpes simplex virus 2

<400> 12
gaagacgagc tctaagggag gggaggggag ctgggattgt gtataaataa aaagacacccg 60
atgttcaaaa atacacatga cttctggtat tgttttgact tggtttttat ttgggggggg 120
aaaaacgtgt gactagaaaa acaaatgcag acatgtgcata aog 163

<210> 13
<211> 191
<212> DNA
<213> Human papillomavirus type 16

<400> 13

aattgttaca tataattgtt gtataccata acttactatt ttttctttt tattttcata	60
tataattttt tttttgttt gttgtttgt ttttaataa actgttatta cttacaatg	12
cgacacaaac gttctgaaa acgcacaaaaa cgtgcatogg ctacccaact ttataaaaca	180
tgcaaacagg c	191

<210> 14
<211> 3759
<212> DNA
<213> Artificial sequence

<220>
<223> pJV expression vector

<220>
<221> Intron
<222> (1725) .. (1857)
<223> Rat Ins IntA

<220>
<221> misc_feature
<222> (1) .. (44)
<223> Tn903, pUC4K Remnants

<220>
<221> misc_feature
<222> (861) .. (896)
<223> Tn903, pUC4K Remnants

<220>
<221> misc_feature
<222> (897) .. (902)
<223> pUC19 MCS

<220>
<221> polyA_signal
<222> (2556) .. (2686)
<223> rGLOB pA

<220>
<221> polyA_site
<222> (2647) .. (2647)
<223> PolyA_Site_1

<220>
<221> promoter
<222> (903) .. (1587)
<223> CMV Pro

<220>
<221> 3'UTR
<222> (2012) .. (2544)
<223> HBVenh

<220>
<221> 5'UTR
<222> (1864) .. (1984)
<223> 5'-UTR of HBV pre-S2

<220>
<221> misc_feature
<222> (1719) .. (1724)
<223> Bam/Bgl fusion
<220>
<221> misc_feature
<222> (1985) .. (1987)
<223> ATG-Nhe

<220>
<221> misc_feature
<222> (1988) .. (2011)
<223> CDS insertion site

<220>
<221> misc_feature
<222> (2545) .. (2555)
<223> unknown

<220>
<221> exon
<222> (1588) .. (1718)
<223> CMV Exon 1/2

<220>
<221> misc_feature
<222> (2693) .. (3759)

<223> pUC19

<220>

<221> misc_feature

<222> (45)..(860)

<223> KanR (Tn903) complement

<400> 14

ggcgttaatgc tctgcagggt ttacaaccaa ttaaccaatt ctgatttagaa aaactcatcg 60
 agcatcaaat gaaactgca tttattcata tcaggattat caataccata tttttgaaaa 120
 agccgtttct gtaatgaagg agaaaactca cccaggcagt tccataggat ggcaagatcc 180
 tggtatecggt ctgcgattcc gactcgccaa acatcaatac aacctattaa ttcccccctcg 240
 tcaaaaataa ggttatcaag tgagaaatca ccatgagtga cgactgaatc cggtgagaat 300
 ggccaaaagot tatgcatttc ttccagact tgttcaacag gccagccatt acgtcgctca 360
 tcaaaaatcac tcgcacatcaac caaaccgtta ttccattcggtt attgcgcctg agcgagacga 420
 aatacgcgtt cgtgttaaa aggacaatta caaacaggaa tccaaatgca cccggcgagg 480
 aacactgcac ggcacatcaac aatattttca cctgaatcgtt gatatttttc taataactgg 540
 aatgcgtttt tcccccggat cgcgtgggtt agttaaccatg catcatcagg agtacggata 600
 aatgcgttga tggtcggaa aggccataat tccgtcagcc agttttagtctt gaccatctca 660
 tctgttaacat cattggcaac gtcaccccttgc ccatgtttca gaaacaactc tggccatcg 720
 gggttcccat acaatcgata gattgtcgca cctgattgcc cgcacattato ggcggccat 780
 ttatacccat ataaatcgcg atccatgttg gaatttaato gggccctcgca gcaagacgtt 840
 tcccggttgaat tggcgtcat aacaccccttgc gtattactgtt ttatgtaaagc agacaggctt 900
 acaatattgg ctattggcca ttgcatacgatgtt tttatctata tcataatatgt tacattata 960
 ttggcgtcatg tccaaatatga cccgcattgtt gacattgtt attgactgtt tattatagt 1020
 aatcaattac ggggtcatta gttcatagcc catatatgtt gttcccggtt acataactta 1080
 cggtaaatgg cccgcctggc tgaccggca acgaccccccggcc cccattgttgc tcaataatgt 1140
 cgtatgttcc catagtaacg ccaataggaa ctttccatgtt acgtcaatgg gtggagtatt 1200
 tacggtaaaatggttgc gcaatgttgc aagtgtatca tatgtcaatggt ccggcccccata 1260
 ttgacgttcaatggcgttgc ggcattatgc ccagtacatgtt accttacggg 1320
 actttccatgttgc tggcgttgc atctacatgtt tagtcatgtt tattaccatgtt gtgtatgttgc 1380
 tttggcgttgc caccatggg cgtggatagc ggtttgactc acggggattt ccaatgttgc 1440
 accccatgttgc cgtcaatggg agtttgcgtt ggcacccaaa tcaacgggac tttccaaaat 1500
 gtcgttgcataatggcgttgc ggcacccaaa tggccgttgc ggtgttgc gttggaggctt 1560
 atataagcgttgc gtcgttgc gtc aga tgg cct gga gac gcc atc cac 1614
 Val Arg Ser Pro Gly Asp Ala Ile His

1

5

gct gtt ttg acc tcc ata gaa gac acc ggg acc gat cca gcc tcc ggg 1662
 Ala Val Leu Thr Ser Ile Glu Asp Thr Gly Thr Asp Pro Ala Ser Ala
 10 15 20 25

gcc ggg aac ggt gca ttg gaa cgc gga ttc ccc gtg cca aga gtg act 1710
 Ala Gly Asn Gly Ala Leu Glu Arg Gly Phe Pro Val Pro Arg Val Thr

30

35

40

cac cgt cc ggatctcago aagcaggtat gtactctoca ggggtgggcct ggcttcacca 1768
His Arg

His Arg

gtcaagactc cagggatttg agggacgctg tgggtcttc tcttacatgt accttttgat	1828
tgccctcaacc ctgactatct tcacggctcg gatcccagag tcaggggtot gtattttct	1888
gctggtgct ccagttcagg aacagtaaac cctgctccga atattgcctc tcacatctcg	1948
tcaatctccg cgaggactgg ggaccctgtg acoaacatgg ctagcccccc cagatctggg	2008
ccctaacaaa acaaaaagat ggggttatto cctaaacttc atgggttacg taatttggaaag	2068
ttgggggaca ttgcacaag atoatattgt acaaaaagatc aaacactgtt ttagaaaact	2128
tcctgtaaac aggcttattt attggaaagt atgtcaaaagg attgtgggtc tttgggctt	2188
tgctgctcca tttacacaat gtggatatec tgocttaatg cctttgtatg catgtataca	2248
agctaaacag gcttcactt totgcacaac ttacaaggcc ttcttaagta aacagtacat	2308
gaacotttaac ccogttgtc ggcacaaoggcc tggctgtgc caagtgttgc ctgacgcaac	2368
ccccactggc tggggcttgg ccataggccca tcagcgcacg cgtggaaacct ttgtggctcc	2428
tctgcgcgtc catactgggg aactcttagc cgcttggttt gctcgacgccc ggtctggagc	2488
aaagctcata ggaactgaca attctgtcgt cctctcgccg aaatatacat ogtttcgatc	2548
tacgatgtatc cttttccct ctgcacaaaaa ttatggggac atcatgaagc cccttgagca	2608
tctgacttct ggctaaataaa gggaaattttat ttctattgtca atagtgtttt ggaatttttt	2668
gtgtctctca ctgcgaagga attctgcatt aatgaatogg ccaacgcggc gggagaggcg	2728
gtttgogtat tgggcgtct tcgcgttctc ogctoactga ctgcgtgcgc tggctgttcc	2788
ggctgaggcgc agcgttatca gctoactcaa aggccgttaat acggttatcc acagaatcc	2848
gggataaaacg aggaaagaac atgtgagccaa aaggccagca aaaggccagg aacogtaaaa	2908
aggccgcgtt gctggcgttt ttccataggc tcggcccccc tgaogagcat cacaacaaatc	2968
gacgctcaag tcagagggtgg cggaaaccccgaa caggactata aagataccag gctttcccc	3028
ctggaaagctc cctctgtgcgc tctctgttcc cgaccctgcc gcttaccggaa tacatgtccg	3088
cctttctccaa ttctggaaagc gtggcgcttt ctcatagctc acgctgttagg tatctcagtt	3148
cggtgttaggt cggtcgctcc aacgtgggtt gtgtgcacgaa acccccccgtt cagcccgacc	3208
gctgcgcgtt atccggtaac tategttggc agtccaaaccc ggtaaagacac gacttacgc	3268
cactggcagc agccactggt aacaggatttgcagagcgag gtatgttaggc ggtgotacag	3328
agttcttggaa gtgggtggcct aactacggctt acactagaag aacagtattt ggtatctcg	3388
ctctgtcgaa gccagttacc ttctggaaaaa gagttggtag ctcttgcattcc ggccaaacaaa	3448
ccacccgttgg tagccgtgggtt tttttgtttt gcaaggcagca gattacgccc agaaaaaaaag	3508
gatctcaaga agatcccttg atcttttcta cggggcttgc cgtctcgtgg aacggaaaact	3568
cacgttaagg gattttggtc atgagattt caaaaaggat cttcacccatgc atctttttaa	3628
ataaaaaaatg aagttttaaa tcaatctaaa gtatataatgc gtaaaacttgg tctgacagtt	3688
accaatgtttt aatcagtgtgag gacccatctc cagcgatctg tctatttgc toatccatag	3748
ttgoctgact c	3759

<210> 15

<211> 42

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 15

ggaggatccg gacggtgagt cactcttggc acggggaaatc cg

42

<210> 16

<211> 21

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 16

ggtaatatg gtcataaca c

21

<210> 17

<211> 23

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 17

ccgcgaaca tggagaacat cgc

23

<210> 18

<211> 33

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 18

cacagatctt ttgttagggt ttaaatgtat acc

33

<210> 19
<211> 29
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 19
ggaggatctt gaccttggaaat atagtcacc

29

<210> 20
<211> 26
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 20
ggaggatcca toagcaagca ggtatg

26

<210> 21
<211> 33
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 21
ggagctatcg ggcttttgcac ctccggcgta ggg

33

<210> 22
<211> 37
<212> DNA
<213> Artificial sequence

<220>

<223> Primer

<400> 22

ggagaattca gatctcttot agtaaaaacaa tggctgg

37

<210> 23

<211> 25

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 23

ggagctagcc ttcttaaccga ggttag

25

<210> 24

<211> 30

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 24

ggaagatctc cttactccag ctctatgctg.

30

<210> 25

<211> 27

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 25

ggcgaattcc ttccgagtga gagacac

27

<210> 26

<211> 43

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 26

ggagtataca tttaaaggc cotaacaaaa caaaaagatg ggg

43

<210> 27

<211> 31

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 27

ggagctagct cgtttacttt gaccaagaac g

31

<210> 28

<211> 36

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 28

ggaagatctc cttatttttg acaccagacc aactgg

36

<210> 29

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 29

ggagtcgacc tgtctgctta cataaacag

29

<210> 30
<211> 26
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 30
cgtaatgctc tgccagtgtt acaacc

26

<210> 31
<211> 20
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 31
gaaaagatctc agcaaggcagg

20

<210> 32
<211> 47
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 32
ggaggaggatccct gacatggaaag atagtcaggg ttgaggcaag caaaaagg

47

<210> 33
<211> 12
<212> DNA
<213> Artificial sequence

<220>

<223> Primer

<400> 33

ctagcgggco ca

12

<210> 34

<211> 12

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 34

gatctgggcc cg

12

<210> 35

<211> 28

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 35

ggagctagca tcatccagt tgaggagg

28

<210> 36

<211> 28

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 36

ggtagatctc ctcatgtctg ctogaagc

28

<210> 37

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 37

ccaagctagc gacaaaactc acacatgcc

29

<210> 38

<211> 45

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 38

ggaagatctc gtttaccact gtcatttacc cggagacagg gagag

45

<210> 39

<211> 57

<212> DNA

<213> Artificial sequence

<220>

<223> Oligonucleotide

<400> 39

aagatgtcca gactctgtct ctccgtggcc ctactctgtgc tcctatggac actcgcc

57

<210> 40

<211> 24

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 40

gaaactagta agatgtccag actc

24

<210> 41
<211> 25
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 41
ggaagctagc ggccgagtgtc cccag

25

<210> 42
<211> 75
<212> DNA
<213> Artificial sequence

<220>
<223> Oligonucleotide

<400> 42
ggaaagatgg ccagcccttt tgccacatcc ctcgtgggtcc tcgtgacccct cccgcctggcc
agccaaagca ggcgc

60

75

<210> 43
<211> 24
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 43
ggaactatgtc gaaagatggc cccgc

24

<210> 44
<211> 26
<212> DNA
<213> Artificial sequence

<220>

<223> Primer

<400> 44

ggaagctagc ggccgtcgctt tcgtcg

26

<210> 45

<211> 51

<212> DNA

<213> Artificial sequence

<220>

<223> Oligonucleotide

<400> 45

aggctttgc taatcttggc gctttgttc atgcacactgg ctgtatctggg g

51

<210> 46

<211> 24

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 46

ggaacttagta ggtctttgct aatc

24

<210> 47

<211> 25

<212> DNA

<213> Artificial sequence

<220>

<223> Primer

<400> 47

ggaagctagc ccccaagagca gccag

25

<210> 48

<211> 31
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 48

ggagctagct cgtttacttt gaccaagaac g

31

<210> 49
<211> 25
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 49

ggaagatctc cggtgagtgg tgctg

25

<210> 50
<211> 32
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 50

gcaggatcca gtagaactgg agagaggaca ag

32

<210> 51
<211> 29
<212> DNA
<213> Artificial sequence

<220>
<223> Primer

<400> 51

ggaagatcta caaggtgago tgctgtggc

29

<210> 52
<211> 490
<212> DNA
<213> Pseudo rabies virus

<400> 1

tggccgcaga	gccccccggg	catgcaaatac	agaggcgccg	gggagacgcc	tccgcgcgc	60
cattggcccg	ggcgagccga	gatggccgc	gccccggccg	gacatgcaaa	gtagacgoga	120
gaggaagtag	ggagagaaat	ccattggcc	gtcgaggggc	caagatggcg	ccctcggggc	180
cggacatgca	aatgtagacgc	gagaggaagt	gggcgagaga	aatccatgt	gcccgtcgatg	240
gggcaagatg	gccccccggg	ggccgggca	tgcaaattgt	cctcgcgagg	aagtccctcg	300
cgaaatccca	ttggccggcg	gcgcgcatac	tggccggggc	atgcaaagca	gacggcagag	360
gaagggggcg	agaaaaatcc	cattggccgg	ccgtcgggga	atcccgccgc	gaaaatcgcc	420
cattggtcgc	cttacotggg	ggcggtctct	catacgggcg	cttataagcg	cggctctccat	480
cgtagcaactt						490

<210> 53
<211> 495
<212> DNA
<213> Rous sarcoma virus

<400> 2

ctgtccctcg	tttgtgtgtt	ggaggcgct	gagtagtgcg	cgagcaaaat	ttaagctaca	60
acaaggcaag	gtttgacoga	caattgcatg	aagaatotgc	ttagggttag	gogttttgag	120
ctgttcgcgc	atgtacgggc	cagatatacg	cgtatctgag	gggacttaggg	tgtttttagg	180
cggaaagccg	ggcttcgggtt	gtaogcggtt	aggagttccc	tcaggatata	gtagtttcgc	240
ttttgcatac	ggagggggaa	atgtagtcctt	atgcaatacaca	cttgcgtct	tgcaacatgg	300
taacgatgag	ttagcaacat	gccttacaag	gagagaaaaa	gcaccgtgca	tgccgattgg	360
tggaaagtaag	gtggtacgat	cgtgccttat	taggaaggca	acagacaggt	otgacatgga	420
ttggacgaaac	cactgaattc	cgcattgcag	agataattgt	atthaagtgc	ctagctcgat	480
acaataaaacg	ccatt					495