



(86) Date de dépôt PCT/PCT Filing Date: 2009/12/04

(87) Date publication PCT/PCT Publication Date: 2010/06/10

(85) Entrée phase nationale/National Entry: 2011/05/31

(86) N° demande PCT/PCT Application No.: US 2009/066815

(87) N° publication PCT/PCT Publication No.: 2010/065882

(30) Priorités/Priorities: 2008/12/04 (US61/200,877);
2009/04/07 (US61/212,071)

(51) Cl.Int./Int.Cl. *C07K 16/46* (2006.01),
A61K 39/395 (2006.01), *C07K 16/00* (2006.01),
C12N 15/00 (2006.01), *C12N 15/13* (2006.01),
G01N 33/53 (2006.01), *G01N 33/68* (2006.01)

(71) Demandeur/Applicant:
ABBOTT LABORATORIES, US


(72) Inventeurs/Inventors:
JAKOB, CLARISSA G., US;
WU, CHENGBIN, US;
WALTER, KARL A., US;
GHAYUR, TARIQ, US

(74) Agent: TORYS LLP

(54) Titre : IMMUNOGLOBULINES A DOUBLE DOMAINE VARIABLE ET LEURS UTILISATIONS

(54) Title: DUAL VARIABLE DOMAIN IMMUNOGLOBULINS AND USES THEREOF

Figure 1



(57) Abrégé/Abstract:

The present invention relates to engineered multivalent and multispecific binding proteins, methods of making, and specifically to their uses in the prevention, diagnosis, and/or treatment of disease.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
10 June 2010 (10.06.2010)

PCT

(10) International Publication Number
WO 2010/065882 A1(51) International Patent Classification:
C07K 16/46 (2006.01) *A61K 39/00* (2006.01)(21) International Application Number:
PCT/US2009/066815(22) International Filing Date:
4 December 2009 (04.12.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/200,877 4 December 2008 (04.12.2008)
61/212,071 7 April 2009 (07.04.2009)US
US(71) Applicant (for all designated States except US): **ABBOTT LABORATORIES** [US/US]; 100 Abbott Park Road, Abbott Park, IL 60064 (US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): **JAKOB, Clarissa, G.** [US/US]; 820 Amelia Court, Grayslake, IL 60030 (US). **WU, Chengbin** [CN/US]; 386 Prospect Street, Shrewsbury, MA 01545 (US). **WALTER, Karl, A.** [US/US]; 4458 W. Forest Avenue, Waukegan, IL 60085 (US). **GHAYUR, Tariq** [US/US]; 1014 Washington Street, Holliston, MA 01746 (US).(74) Agents: **STEEL, Diana, M.** et al.; Abbott Bioresearch Center, 100 Research Drive, Worcester, MA 01605 (US).

[Continued on next page]

(54) Title: DUAL VARIABLE DOMAIN IMMUNOGLOBULINS AND USES THEREOF

Figure 1



(57) Abstract: The present invention relates to engineered multivalent and multispecific binding proteins, methods of making, and specifically to their uses in the prevention, diagnosis, and/or treatment of disease.

WO 2010/065882 A1

WO 2010/065882 A1



(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *with international search report (Art. 21(3))*

DUAL VARIABLE DOMAIN IMMUNOGLOBULINS AND USES THEREOF**Reference to Related Applications**

This application is a non-provisional application claiming priority to U.S. Provisional
5 Application Serial No. 61/200,877, filed December 4, 2008 and U.S. Provisional Application
Serial No. 61/212,071, filed April 7, 2009, the contents of which are hereby incorporated by
reference.

Field of the Invention

The present invention relates to multivalent and multispecific binding proteins, methods
10 of making, and specifically to their uses in the, diagnosis, prevention and/or treatment of acute
and chronic inflammatory diseases, cancer, and other diseases.

Background of the Invention

Engineered proteins, such as multispecific antibodies capable of binding two or more
antigens are known in the art. Such multispecific binding proteins can be generated using cell
15 fusion, chemical conjugation, or recombinant DNA techniques.

Bispecific antibodies have been produced using quadroma technology (see Milstein, C.
and A.C. Cuello (1983) Nature 305(5934):537-40) based on the somatic fusion of two different
hybridoma cell lines expressing murine monoclonal antibodies (mAbs) with the desired
specificities of the bispecific antibody. Because of the random pairing of two different
20 immunoglobulin (Ig) heavy and light chains within the resulting hybrid-hybridoma (or quadroma)
cell line, up to ten different Ig species are generated, of which only one is the functional bispecific
antibody. The presence of mis-paired by-products, and significantly reduced production yields,
means sophisticated purification procedures are required.

Bispecific antibodies can also be produced by chemical conjugation of two different
25 mAbs (see Staerz, U.D., et al. (1985) Nature 314(6012): 628-31). This approach does not yield
homogeneous preparation. Other approaches have used chemical conjugation of two different
mAbs or smaller antibody fragments (see Brennan, M., et al. (1985) Science 229(4708): 81-3).

Another method used to produce bispecific antibodies is the coupling of two parental
antibodies with a hetero-bifunctional crosslinker, but the resulting bispecific antibodies suffer
30 from significant molecular heterogeneity because reaction of the crosslinker with the parental
antibodies is not site-directed. To obtain more homogeneous preparations of bispecific antibodies
two different Fab fragments have been chemically crosslinked at their hinge cysteine residues in a

site-directed manner (see Glennie, M.J., et al. (1987) J. Immunol. 139(7): 2367-75). But this method results in Fab'2 fragments, not full IgG molecule.

A wide variety of other recombinant bispecific antibody formats have been developed (see Kriangkum, J., et al. (2001) Biomol. Eng. 18(2): 31-40). Amongst them tandem single-chain Fv molecules and diabodies, and various derivatives thereof, are the most widely used. Routinely, construction of these molecules starts from two single-chain Fv (scFv) fragments that recognize different antigens (see Economides, A.N., et al. (2003) Nat. Med. 9(1): 47-52). Tandem scFv molecules (taFv) represent a straightforward format simply connecting the two scFv molecules with an additional peptide linker. The two scFv fragments present in these tandem scFv molecules form separate folding entities. Various linkers can be used to connect the two scFv fragments and linkers with a length of up to 63 residues (see Nakanishi, K., et al. (2001) Ann. Rev. Immunol. 19: 423-74). Although the parental scFv fragments can normally be expressed in soluble form in bacteria, it is, however, often observed that tandem scFv molecules form insoluble aggregates in bacteria. Hence, refolding protocols or the use of mammalian expression systems are routinely applied to produce soluble tandem scFv molecules. In a recent study, *in vivo* expression by transgenic rabbits and cattle of a tandem scFv directed against CD28 and a melanoma-associated proteoglycan was reported (see Gracie, J.A., et al. (1999) J. Clin. Invest. 104(10): 1393-401). In this construct, the two scFv molecules were connected by a CH1 linker and serum concentrations of up to 100 mg/L of the bispecific antibody were found. Various strategies including variations of the domain order or using middle linkers with varying length or flexibility were employed to allow soluble expression in bacteria. A few studies have now reported expression of soluble tandem scFv molecules in bacteria (see Leung, B.P., et al. (2000) J. Immunol. 164(12): 6495-502; Ito, A., et al. (2003) J. Immunol. 170(9): 4802-9; Karni, A., et al. (2002) J. Neuroimmunol. 125(1-2): 134-40) using either a very short Ala3 linker or long glycine/serine-rich linkers. In another study, phage display of a tandem scFv repertoire containing randomized middle linkers with a length of 3 or 6 residues was employed to enrich for those molecules that are produced in soluble and active form in bacteria. This approach resulted in the isolation of a tandem scFv molecule with a 6 amino acid residue linker (see Arndt, M. and J. Krauss (2003) Methods Mol. Biol. 207: 305-21). It is unclear whether this linker sequence represents a general solution to the soluble expression of tandem scFv molecules. Nevertheless, this study demonstrated that phage display of tandem scFv molecules in combination with directed mutagenesis is a powerful tool to enrich for these molecules, which can be expressed in bacteria in an active form.

Bispecific diabodies (Db) utilize the diabody format for expression. Diabodies are produced from scFv fragments by reducing the length of the linker connecting the VH and VL domain to approximately 5 residues (see Peipp, M. and T. Valerius (2002) Biochem. Soc. Trans.

30(4): 507-11). This reduction of linker size facilitates dimerization of two polypeptide chains by crossover pairing of the VH and VL domains. Bispecific diabodies are produced by expressing, two polypeptide chains with, either the structure VHA-VLB and VHB-VLA (VH-VL configuration), or VLA-VHB and VLB-VHA (VL-VH configuration) within the same cell. A large variety of different bispecific diabodies have been produced in the past and most of them are expressed in soluble form in bacteria. However, a recent comparative study demonstrates that the orientation of the variable domains can influence expression and formation of active binding sites (see Mack, M. et al. (1995) Proc. Natl. Acad. Sci. U S A 92(15): 7021-5). Nevertheless, soluble expression in bacteria represents an important advantage over tandem scFv molecules. However, since two different polypeptide chains are expressed within a single cell inactive homodimers can be produced together with active heterodimers. This necessitates the implementation of additional purification steps in order to obtain homogenous preparations of bispecific diabodies. One approach to force the generation of bispecific diabodies is the production of knob-into-hole diabodies (see Holliger, P., T. Prospero, and G. Winter (1993) Proc. Natl. Acad. Sci. U S A 90(14): 6444-8.18). This approach was demonstrated for a bispecific diabody directed against HER2 and CD3. A large knob was introduced in the VH domain by exchanging Val37 with Phe and Leu45 with Trp and a complementary hole was produced in the VL domain by mutating Phe98 to Met and Tyr87 to Ala, either in the anti- HER2 or the anti-CD3 variable domains. By using this approach the production of bispecific diabodies could be increased from 72% by the parental diabody to over 90% by the knob-into-hole diabody. Importantly, production yields did only slightly decrease as a result of these mutations. However, a reduction in antigen-binding activity was observed for several analyzed constructs. Thus, this rather elaborate approach requires the analysis of various constructs in order to identify those mutations that produce heterodimeric molecule with unaltered binding activity. In addition, such approach requires mutational modification of the immunoglobulin sequence at the constant region, thus creating non-native and non-natural form of the antibody sequence, which may result in increased immunogenicity, poor *in vivo* stability, as well as undesirable pharmacokinetics.

Single-chain diabodies (scDb) represent an alternative strategy for improving the formation of bispecific diabody-like molecules (see Holliger, P. and G. Winter (1997) Cancer Immunol. Immunother. 45(3-4): 128-30; Wu, A.M., et al. (1996) Immunotechnology 2(1): p. 21-36). Bispecific single-chain diabodies are produced by connecting the two diabody-forming polypeptide chains with an additional middle linker with a length of approximately 15 amino acid residues. Consequently, all molecules with a molecular weight corresponding to monomeric single-chain diabodies (50-60 kDa) are bispecific. Several studies have demonstrated that bispecific single chain diabodies are expressed in bacteria in soluble and active form with the majority of purified molecules present as monomers (see Holliger, P. and G. Winter (1997)

Cancer Immunol. Immunother. 45(3-4): 128-30; Wu, A.M., et al. (1996) Immunotechnol. 2(1): 21-36; Pluckthun, A. and P. Pack (1997) Immunotechnol. 3(2): 83-105; Ridgway, J.B., et al. (1996) Protein Engin. 9(7): 617-21). Thus, single-chain diabodies combine the advantages of tandem scFvs (all monomers are bispecific) and diabodies (soluble expression in bacteria).

5 More recently diabodies have been fused to Fc to generate more Ig-like molecules, named di-diabodies (see Lu, D., et al. (2004) J. Biol. Chem. 279(4): 2856-65). In addition, multivalent antibody construct comprising two Fab repeats in the heavy chain of an IgG and capable of binding four antigen molecules has been described (see WO 0177342A1, and Miller, K., et al. (2003) J. Immunol. 170(9): 4854-61).

10 There is a need in the art for improved multivalent binding proteins capable of binding two or more antigens. U.S. Patent Application Serial No. 11/507,050 provides a novel family of binding proteins capable of binding two or more antigens with high affinity, which are called dual variable domain immunoglobulins (DVD-IgTM). The present invention provides further novel binding proteins capable of binding two or more antigens.

15 Summary of the Invention

This invention pertains to multivalent binding proteins capable of binding two or more antigens. The present invention provides a novel family of binding proteins capable of binding two or more antigens with high affinity.

In one embodiment the invention provides a binding protein comprising a polypeptide
20 chain, wherein the polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first variable domain, VD2 is a second variable domain, C is a constant domain, X1 represents an amino acid or polypeptide, X2 represents an Fc region and n is 0 or 1. In an embodiment the VD1 and VD2 in the binding protein are heavy chain variable domains. In another embodiment, the heavy chain variable domain is selected from the group consisting of a murine heavy chain
25 variable domain, a human heavy chain variable domain, a CDR grafted heavy chain variable domain, and a humanized heavy chain variable domain. In yet another, embodiment VD1 and VD2 are capable of binding the same antigen. In another embodiment VD1 and VD2 are capable of binding different antigens. In still another embodiment, C is a heavy chain constant domain. For example, X1 is a linker with the proviso that X1 is not CH1.

30 For example, X1 is a linker selected from the group consisting of
AKTTPKLEEGEFSEAR (SEQ ID NO: 1); AKTTPKLEEGEFSEARV (SEQ ID NO: 2);
AKTTPKLGG (SEQ ID NO: 3); SAKTTPKLGG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5);
RADAAP (SEQ ID NO: 6); RADAAPTVS (SEQ ID NO: 7); RADAAAAGGPGS (SEQ ID NO:
8); RADAAA(G₄S)₄ (SEQ ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP
35 (SEQ ID NO: 11); ADAAPTVSIFPP (SEQ ID NO: 12); TVAAP (SEQ ID NO: 13);

TVAAPSVFIFPP (SEQ ID NO: 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID NO: 16); AKTTPP (SEQ ID NO: 17); AKTTPPSVTPLAP (SEQ ID NO: 18); AKTTAP (SEQ ID NO: 19); AKTTAPSVYPLAP (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21); ASTKGPSVFPLAP (SEQ ID NO: 22); GGGGSGGGGSGGGGS (SEQ ID NO: 23);

5 GENKVEYAPALMALS (SEQ ID NO: 24); GPAKELTPLKEAKVS (SEQ ID NO: 25); GHEAAAVMQVQYPAS (SEQ ID NO: 26); GGGGGGGGP (SEQ ID NO: 27); GGGGGGGGP (SEQ ID NO: 28); PAPNLLGGP (SEQ ID NO: 29); PNLLGGP (SEQ ID NO: 30); GGGGGGGP (SEQ ID NO: 31); PAPELLGGP (SEQ ID NO: 32); PTISPAPNLLGGP (SEQ ID NO: 33); TVAADDDDKSVFIVPP (SEQ ID NO: 34); TVDDDDKAAAP (SEQ ID NO: 35); LVPRGSAAP (SEQ ID NO: 36); ASDDDDK GGP (SEQ ID NO: 37); ALVPR GSGP (SEQ ID NO: 38); ASTDDDDK SVFPLAP (SEQ ID NO: 39); TVALVPR GSVFIFPP (SEQ ID NO: 40); ASTLVPR GSVFPLAP (SEQ ID NO: 41); TVAADDDK SVFIVPP (SEQ ID NO: 42); ASTDDDK SVFPLAP (SEQ ID NO: 43); LEVLFFQ GP (SEQ ID NO: 44); TVAALEVLFFQ GPAP (SEQ ID NO: 45); ASTLEVLFFQ GPLAP (SEQ ID NO: 46); PAPLEVLFFQ GP (SEQ ID NO: 47); TAENLYFFQ GAP (SEQ ID NO: 48); AENLYFFQ GA (SEQ ID NO: 49); PGPFGR SAGGP (SEQ ID NO: 50); PGPFGR SAGG (SEQ ID NO: 51); PQRGR SAG (SEQ ID NO: 52); PHYGR SGG (SEQ ID NO: 53); GPFGR SAGP (SEQ ID NO: 54); GDDDDDK GGP (SEQ ID NO: 55); AGDDDDDK GGP (SEQ ID NO: 56); GGDDDDDK GGP (SEQ ID NO: 57); AS; TVA; ASTK (SEQ ID NO: 58); ASTKGPSV (SEQ ID NO: 59); ASTKGPSVFP (SEQ ID NO: 60);

15 NO: 47); TAENLYFFQ GAP (SEQ ID NO: 48); AENLYFFQ GA (SEQ ID NO: 49); PGPFGR SAGGP (SEQ ID NO: 50); PGPFGR SAGG (SEQ ID NO: 51); PQRGR SAG (SEQ ID NO: 52); PHYGR SGG (SEQ ID NO: 53); GPFGR SAGP (SEQ ID NO: 54); GDDDDDK GGP (SEQ ID NO: 55); AGDDDDDK GGP (SEQ ID NO: 56); GGDDDDDK GGP (SEQ ID NO: 57); AS; TVA; ASTK (SEQ ID NO: 58); ASTKGPSV (SEQ ID NO: 59); ASTKGPSVFP (SEQ ID NO: 60);

20 TVAAPSV (SEQ ID NO: 61), and TVAAPSVFI (SEQ ID NO: 62). In an embodiment, X2 is an Fc region. In another embodiment, X2 is a variant Fc region.

In an embodiment the binding protein disclosed herein comprises a polypeptide chain, wherein the polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region.

In an embodiment, VD1 and VD2 in the binding protein are light chain variable domains. In an embodiment, the light chain variable domain is selected from the group consisting of a murine light chain variable domain, a human light chain variable domain, a CDR grafted light chain variable domain, and a humanized light chain variable domain. In one embodiment VD1 and VD2 are capable of binding the same antigen. In another embodiment VD1 and VD2 are capable of binding different antigens. In an embodiment, C is a light chain constant domain. In another embodiment, X1 is a linker with the proviso that X1 is not CL1.

In an embodiment, X1 is a linker selected from the group consisting of AKTTPKLEEGEFSEAR (SEQ ID NO: 1); AKTTPKLEEGEFSEARV (SEQ ID NO: 2); AKTTPKLGG (SEQ ID NO: 3);

35 SAKTTPKLGG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5); RADAAP (SEQ ID NO: 6); RADAAPTVS (SEQ ID NO: 7); RADAAAAGGPGS (SEQ ID NO: 8); RADAAA(G₄S)₄ (SEQ

ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP (SEQ ID NO: 11);
 ADAAPTVSIFPP (SEQ ID NO: 12); TVAAP (SEQ ID NO: 13); TVAAPSVFIFPP (SEQ ID NO:
 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID NO: 16); AKTTTPP (SEQ ID NO:
 17); AKTTPPSVTPLAP (SEQ ID NO: 18); AKTTAP (SEQ ID NO: 19); AKTTAPSVYPLAP
 5 (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21); ASTKGPSVFPLAP (SEQ ID NO: 22),
 GGGGSGGGGSGGGGS (SEQ ID NO: 23); GENKVEYAPALMALS (SEQ ID NO: 24);
 GPAKELTPLKEAKVS (SEQ ID NO: 25); GHEAAVMQVQYPAS (SEQ ID NO: 26);
 GGGGGGGGP (SEQ ID NO: 27); GGGGGGGGGP (SEQ ID NO: 28); PAPNLLGGP (SEQ ID NO:
 29); PNLLGGP (SEQ ID NO: 30); GGGGGGGP (SEQ ID NO: 31); PAPELLGGP (SEQ ID NO:
 10 32); PTISPAPNLLGGP (SEQ ID NO: 33); TVAADDDDKSVFIVPP (SEQ ID NO: 34);
 TVDDDDDKAAP (SEQ ID NO: 35); LVPRGSAAP (SEQ ID NO: 36); ASDDDDK GGP (SEQ
 ID NO: 37); ALVPR GSGP (SEQ ID NO: 38); ASTDDDDK SVFPLAP (SEQ ID NO: 39);
 TVALVPR GSVFIFPP (SEQ ID NO: 40); ASTLVPR GSVFPLAP (SEQ ID NO: 41);
 TVAADDDK SVFIVPP (SEQ ID NO: 42); ASTDDDK SVFPLAP (SEQ ID NO: 43); LEVLQ
 15 GP (SEQ ID NO: 44); TVAALEVLQ GPAP (SEQ ID NO: 45); ASTLEVLQ GPLAP (SEQ ID
 NO: 46); PAPLEVLQ GP (SEQ ID NO: 47); TAENLYFQ GAP (SEQ ID NO: 48); AENLYFQ
 GA (SEQ ID NO: 49); PGPFGR SAGGP (SEQ ID NO: 50); PGPFGR SAGG (SEQ ID NO: 51);
 PQRGR SAG (SEQ ID NO: 52); PHYGR SGG (SEQ ID NO: 53); GPFGR SAGP (SEQ ID NO:
 54); GDDDDK GGP (SEQ ID NO: 55); AGDDDDK GGP (SEQ ID NO: 56); GGDDDDK GGP
 20 (SEQ ID NO: 57); AS; TVA; ASTK (SEQ ID NO: 58); ASTKGPSV (SEQ ID NO: 59);
 ASTKGPSVFP (SEQ ID NO: 60); TVAAPSV (SEQ ID NO: 61), and TVAAPSVFI (SEQ ID
 NO: 62).

In an embodiment, the binding protein does not comprise X2.

In an embodiment, both the variable heavy and variable light chain comprise the same
 25 linker. In another embodiment, the variable heavy and variable light chain comprise different
 linkers. In another embodiment, both the variable heavy and variable light chain comprise a short
 (about 6 amino acids) linker. In another embodiment, both the variable heavy and variable light
 chain comprise a long (greater than 6 amino acids) linker. In another embodiment, the variable
 heavy chain comprises a short linker and the variable light chain comprises a long linker. In
 30 another embodiment, the variable heavy chain comprises a long linker and the variable light chain
 comprises a short linker.

In an embodiment the binding protein disclosed herein comprises a polypeptide chain,
 wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first light
 chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant
 35 domain, X1 is a linker with the proviso that it is not CH1, and X2 does not comprise an Fc region.

In another embodiment the invention provides a binding protein comprising two polypeptide chains, wherein said first polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and said second polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 does not comprise an Fc region. In a particular embodiment, the Dual Variable Domain (DVD) binding protein comprises four polypeptide chains wherein the first two polypeptide chains comprises VD1-(X1)_n-VD2-C-(X2)_n, respectively wherein VD1 is a first heavy chain variable domain, VD2 is a second heavy chain variable domain, C is a heavy chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 is an Fc region; and the second two polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n respectively, wherein VD1 is a first light chain variable domain, VD2 is a second light chain variable domain, C is a light chain constant domain, X1 is a linker with the proviso that it is not CH1, and X2 does not comprise an Fc region. Such a Dual Variable Domain (DVD) protein has four antigen binding sites.

In another embodiment the binding proteins disclosed herein are capable of binding one or more targets. In an embodiment, the target is selected from the group consisting of cytokines, cell surface proteins, enzymes and receptors. In another embodiment, the binding protein is capable of modulating a biological function of one or more targets. In another embodiment, the binding protein is capable of neutralizing one or more targets. The binding protein of the invention is capable of binding cytokines selected from the group consisting of lymphokines, monokines, polypeptide hormones, receptors, or tumor markers. For example, the DVD-Ig of the invention is capable of binding two or more of the following: VEGF, NRP1, SOST, and TNF (see also Table 4). In a specific embodiment the binding protein is capable of binding pairs of targets selected from the group consisting of VEGF and NRP1; and TNF and SOST.

In an embodiment, the binding protein capable of binding NRP1 (seq. 1) and VEGF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 84 and a DVD light chain amino acid sequence of SEQ ID NO. 85.

In a second embodiment, the binding protein capable of binding NRP1 (seq. 1) and VEGF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 86 and a DVD light chain amino acid sequence of SEQ ID NO. 87.

In a third embodiment, the binding protein capable of binding NRP1 (seq. 1) and VEGF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 88 and a DVD light chain amino acid sequence of SEQ ID NO. 89.

In a fourth embodiment, the binding protein capable of binding NRP1 (seq. 1) and VEGF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 90 and a DVD light chain amino acid sequence of SEQ ID NO. 91.

5 In a fifth embodiment, the binding protein capable of binding NRP1 (seq. 1) and VEGF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 92 and a DVD light chain amino acid sequence of SEQ ID NO. 93.

In an embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 94 and a DVD light chain amino acid sequence of SEQ ID NO. 95.

10 In a second embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 96 and a DVD light chain amino acid sequence of SEQ ID NO. 97.

In a third embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 98 and a DVD light chain
15 amino acid sequence of SEQ ID NO. 99.

In a fourth embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 100 and a DVD light chain amino acid sequence of SEQ ID NO. 101.

In a fifth embodiment, the binding protein capable of binding SOST and TNF (seq. 1)
20 comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 102 and a DVD light chain amino acid sequence of SEQ ID NO. 103.

In a sixth embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 104 and a DVD light chain amino acid sequence of SEQ ID NO. 105.

25 In a seventh embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 106 and a DVD light chain amino acid sequence of SEQ ID NO. 107.

In an eighth embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 108 and a DVD light chain
30 amino acid sequence of SEQ ID NO. 109.

In a ninth embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 110 and a DVD light chain amino acid sequence of SEQ ID NO. 111.

In a tenth embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 112 and a DVD light chain amino acid sequence of SEQ ID NO. 113.

In a eleventh embodiment, the binding protein capable of binding SOST and TNF (seq. 1) comprises a DVD heavy chain amino acid sequence of SEQ ID NO. 114 and a DVD light chain amino acid sequence of SEQ ID NO. 115.

In an embodiment, the binding protein capable of binding TNF (seq. 3) and IL-13 (seq. 1) comprises a DVD heavy chain amino acid sequence of any one of SEQ ID NOs: 116-122. In an embodiment, the binding protein capable of binding TNF (seq. 3) and IL-13 (seq. 1) comprises a DVD light chain amino acid sequence of any one of SEQ ID NOs: 123-128. Any of the heavy chains of SEQ ID NOs: 116-122 can be combined with any of the light chain of SEQ ID NOs: 123-128 to make a DVD-Ig of the invention.

In an embodiment, the binding protein capable of binding TNF (seq. 2) and IL-13 (seq. 2) comprises a DVD heavy chain amino acid sequence of any one of SEQ ID NOs: 129-130. In an embodiment, the binding protein capable of binding TNF (seq. 3) and IL-13 (seq. 1) comprises a DVD light chain amino acid sequence of any one of SEQ ID NOs: 131-132. Any of the heavy chains of SEQ ID NOs: 129-130 can be combined with any of the light chain of SEQ ID NOs: 131-132 to make a DVD-Ig of the invention.

In another embodiment the invention provides a binding protein comprising a polypeptide chain, wherein said polypeptide chain comprises $VD1-(X1)_n-VD2-C-(X2)_n$, wherein; VD1 is a first heavy chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; $(X1)_n$ is a linker with the proviso that it is not CH1, wherein said $(X1)_n$ is either present or absent; and $(X2)_n$ is an Fc region, wherein said $(X2)_n$ is either present or absent. In an embodiment, the Fc region is absent from the binding protein.

In another embodiment, the invention provides a binding protein comprising a polypeptide chain, wherein said polypeptide chain comprises $VD1-(X1)_n-VD2-C-(X2)_n$, wherein, VD1 is a first light chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a light chain constant domain; $(X1)_n$ is a linker with the proviso that it is not CH1, wherein said $(X1)_n$ is either present or absent; and $(X2)_n$ does not comprise an Fc region, wherein said $(X2)_n$ is either present or absent. In an embodiment, $(X2)_n$ is absent from the binding protein.

In another embodiment the binding protein of the invention comprises first and second polypeptide chains, wherein said first polypeptide chain comprises a first VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first heavy chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region, wherein said (X2)_n is either present or absent; and wherein said second polypeptide chain comprises a second VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first light chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a light chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent. In another embodiment, the binding protein comprises two first polypeptide chains and two second polypeptide chains. In yet another embodiment, (X2)_n is absent from the second polypeptide. In still another embodiment, the Fc region, if present in the first polypeptide is selected from the group consisting of native sequence Fc region and a variant sequence Fc region. In still another embodiment, the Fc region is selected from the group consisting of an Fc region from an IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, and IgD.

In another embodiment the binding protein of the invention is a DVD-Ig capable of binding two antigens comprising four polypeptide chains, wherein, first and third polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein, VD1 is a first heavy chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second heavy chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region, wherein said (X2)_n is either present or absent; and wherein second and fourth polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first light chain variable domain obtained from a first parent antibody or antigen binding portion thereof; VD2 is a second light chain variable domain obtained from a second parent antibody or antigen binding portion thereof; C is a light chain constant domain; (X1)_n is a linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein said (X2)_n is either present or absent.

The invention provides a method of making a DVD-Ig binding protein by preselecting the parent antibodies. In an embodiment, the method of making a Dual Variable Domain Immunoglobulin capable of binding two antigens comprising the steps of a) obtaining a first parent antibody or antigen binding portion thereof, capable of binding a first antigen; b) obtaining

a second parent antibody or antigen binding portion thereof, capable of binding a second antigen;
 c) constructing first and third polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein,
 VD1 is a first heavy chain variable domain obtained from said first parent antibody or antigen
 binding portion thereof; VD2 is a second heavy chain variable domain obtained from said second
 5 parent antibody or antigen binding portion thereof; C is a heavy chain constant domain; (X1)_n is a
 linker with the proviso that it is not CH1, wherein said (X1)_n is either present or absent; and
 (X2)_n is an Fc region, wherein said (X2)_n is either present or absent; d) constructing second and
 fourth polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein, VD1 is a first light
 chain variable domain obtained from said first parent antibody or antigen binding portion thereof;
 10 VD2 is a second light chain variable domain obtained from said second parent antibody or antigen
 binding thereof; C is a light chain constant domain; (X1)_n is a linker with the proviso that it is not
 CH1, wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region,
 wherein said (X2)_n is either present or absent; e) expressing said first, second, third and fourth
 polypeptide chains; such that a Dual Variable Domain Immunoglobulin capable of binding said
 15 first and said second antigen is generated.

In still another embodiment, the invention provides a method of generating a Dual
 Variable Domain Immunoglobulin capable of binding two antigens with desired properties
 comprising the steps of a) obtaining a first parent antibody or antigen binding portion thereof,
 capable of binding a first antigen and possessing at least one desired property exhibited by the
 20 Dual Variable Domain Immunoglobulin; b) obtaining a second parent antibody or antigen binding
 portion thereof, capable of binding a second antigen and possessing at least one desired property
 exhibited by the Dual Variable Domain Immunoglobulin; c) constructing first and third
 polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein; VD1 is a first heavy chain
 variable domain obtained from said first parent antibody or antigen binding portion thereof; VD2
 25 is a second heavy chain variable domain obtained from said second parent antibody or antigen
 binding portion thereof; C is a heavy chain constant domain; (X1)_n is a linker with the proviso
 that it is not CH1, wherein said (X1)_n is either present or absent; and (X2)_n is an Fc region,
 wherein said (X2)_n is either present or absent; d) constructing second and fourth polypeptide
 chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein; VD1 is a first light chain variable domain
 30 obtained from said first parent antibody or antigen binding portion thereof; VD2 is a second light
 chain variable domain obtained from said second parent antibody or antigen binding portion
 thereof; C is a light chain constant domain; (X1)_n is a linker with the proviso that it is not CH1,
 wherein said (X1)_n is either present or absent; and (X2)_n does not comprise an Fc region, wherein
 said (X2)_n is either present or absent; e) expressing said first, second, third and fourth polypeptide
 35 chains; such that a Dual Variable Domain Immunoglobulin capable of binding said first and said
 second antigen with desired properties is generated.

In one embodiment, the VDI of the first and second polypeptide chains disclosed herein are obtained from the same parent antibody or antigen binding portion thereof. In another embodiment, the VDI of the first and second polypeptide chains disclosed herein are obtained from different parent antibodies or antigen binding portions thereof. In another embodiment, the
5 VD2 of the first and second polypeptide chains disclosed herein are obtained from the same parent antibody or antigen binding portion thereof. In another embodiment, the VD2 of the first and second polypeptide chains disclosed herein are obtained from different parent antibodies or antigen binding portions thereof.

In one embodiment the first parent antibody or antigen binding portion thereof, and the
10 second parent antibody or antigen binding portion thereof, are the same antibody. In another embodiment the first parent antibody or antigen binding portion thereof, and the second parent antibody or antigen binding portion thereof, are different antibodies.

In one embodiment the first parent antibody or antigen binding portion thereof, binds a first antigen and the second parent antibody or antigen binding portion thereof, binds a second
15 antigen. In a particular embodiment, the first and second antigens are the same antigen. In another embodiment, the parent antibodies bind different epitopes on the same antigen. In another embodiment the first and second antigens are different antigens. In another embodiment, the first parent antibody or antigen binding portion thereof, binds the first antigen with a potency different from the potency with which the second parent antibody or antigen binding portion
20 thereof, binds the second antigen. In yet another embodiment, the first parent antibody or antigen binding portion thereof, binds the first antigen with an affinity different from the affinity with which the second parent antibody or antigen binding portion thereof, binds the second antigen.

In another embodiment the first parent antibody or antigen binding portion thereof, and the second parent antibody or antigen binding portion thereof, are selected from the group
25 consisting of, human antibody, CDR grafted antibody, and humanized antibody. In an embodiment, the antigen binding portions are selected from the group consisting of a Fab fragment, a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consisting of the VH and CH1 domains; a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, a dAb fragment,
30 an isolated complementarity determining region (CDR), a single chain antibody, and diabodies.

In another embodiment the binding protein of the invention possesses at least one desired property exhibited by the first parent antibody or antigen binding portion thereof, or the second parent antibody or antigen binding portion thereof. Alternatively, the first parent antibody or antigen binding portion thereof and the second parent antibody or antigen binding portion thereof
35 possess at least one desired property exhibited by the Dual Variable Domain Immunoglobulin. In

an embodiment, the desired property is selected from one or more antibody parameters. In another embodiment, the antibody parameters are selected from the group consisting of antigen specificity, affinity to antigen, potency, biological function, epitope recognition, stability, solubility, production efficiency, immunogenicity, pharmacokinetics, bioavailability, tissue cross reactivity, and orthologous antigen binding. In an embodiment the binding protein is multivalent. In another embodiment, the binding protein is multispecific. The multivalent and or multispecific binding proteins described herein have desirable properties particularly from a therapeutic standpoint. For instance, the multivalent and or multispecific binding protein may (1) be internalized (and/or catabolized) faster than a bivalent antibody by a cell expressing an antigen to which the antibodies bind; (2) be an agonist antibody; and/or (3) induce cell death and/or apoptosis of a cell expressing an antigen which the multivalent antibody is capable of binding to. The "parent antibody" which provides at least one antigen binding specificity of the multivalent and or multispecific binding proteins may be one which is internalized (and/or catabolized) by a cell expressing an antigen to which the antibody binds; and/or may be an agonist, cell death-inducing, and/or apoptosis-inducing antibody, and the multivalent and or multispecific binding protein as described herein may display improvement(s) in one or more of these properties. Moreover, the parent antibody may lack any one or more of these properties, but may be endowed with them when constructed as a multivalent binding protein as described herein.

In another embodiment the binding protein of the invention has an on rate constant (K_{on}) to one or more targets selected from the group consisting of: at least about $10^2 M^{-1}s^{-1}$; at least about $10^3 M^{-1}s^{-1}$; at least about $10^4 M^{-1}s^{-1}$; at least about $10^5 M^{-1}s^{-1}$; and at least about $10^6 M^{-1}s^{-1}$, as measured by surface plasmon resonance. In an embodiment, the binding protein of the invention has an on rate constant (K_{on}) to one or more targets between $10^2 M^{-1}s^{-1}$ and $10^3 M^{-1}s^{-1}$; between $10^3 M^{-1}s^{-1}$ and $10^4 M^{-1}s^{-1}$; between $10^4 M^{-1}s^{-1}$ and $10^5 M^{-1}s^{-1}$; or between $10^5 M^{-1}s^{-1}$ and $10^6 M^{-1}s^{-1}$, as measured by surface plasmon resonance.

In another embodiment the binding protein has an off rate constant (K_{off}) for one or more targets selected from the group consisting of: at most about $10^{-3} s^{-1}$; at most about $10^{-4} s^{-1}$; at most about $10^{-5} s^{-1}$; and at most about $10^{-6} s^{-1}$, as measured by surface plasmon resonance. In an embodiment, the binding protein of the invention has an off rate constant (K_{off}) to one or more targets of $10^{-3} s^{-1}$ to $10^{-4} s^{-1}$; of $10^{-4} s^{-1}$ to $10^{-5} s^{-1}$; or of $10^{-5} s^{-1}$ to $10^{-6} s^{-1}$, as measured by surface plasmon resonance.

In another embodiment the binding protein has a dissociation constant (K_D) to one or more targets selected from the group consisting of: at most about $10^{-7} M$; at most about $10^{-8} M$; at most about $10^{-9} M$; at most about $10^{-10} M$; at most about $10^{-11} M$; at most about $10^{-12} M$; and at most $10^{-13} M$. In an embodiment, the binding protein of the invention has a dissociation constant

(K_D) to its targets of 10⁻⁷ M to 10⁻⁸ M; of 10⁻⁸ M to 10⁻⁹ M; of 10⁻⁹ M to 10⁻¹⁰ M; of 10⁻¹⁰ to 10⁻¹¹ M; of 10⁻¹¹ M to 10⁻¹² M; or of 10⁻¹² to M 10⁻¹³ M.

In another embodiment, the binding protein described herein is a conjugate further comprising an agent selected from the group consisting of an immunoadhesion molecule, an
 5 imaging agent, a therapeutic agent, and a cytotoxic agent. In an embodiment, the imaging agent is selected from the group consisting of a radiolabel, an enzyme, a fluorescent label, a luminescent label, a bioluminescent label, a magnetic label, and biotin. In another embodiment, the imaging agent is a radiolabel selected from the group consisting of: ³H, ¹⁴C, ³⁵S, ⁹⁰Y, ⁹⁹Tc, ¹¹¹In, ¹²⁵I, ¹³¹I, ¹⁷⁷Lu, ¹⁶⁶Ho, and ¹⁵³Sm. In yet another embodiment, the therapeutic or cytotoxic agent is selected
 10 from the group consisting of an anti-metabolite, an alkylating agent, an antibiotic, a growth factor, a cytokine, an anti-angiogenic agent, an anti-mitotic agent, an anthracycline, toxin, and an apoptotic agent.

In another embodiment, the binding protein described herein is a crystallized binding protein and exists as a crystal. In an embodiment, the crystal is a carrier-free pharmaceutical
 15 controlled release crystal. In yet another embodiment, the crystallized binding protein has a greater half life in vivo than the soluble counterpart of said binding protein. In still another embodiment, the crystallized binding protein retains biological activity.

In another embodiment, the binding protein described herein is glycosylated. For example, the glycosylation is a human glycosylation pattern.

20 One aspect of the invention pertains to an isolated nucleic acid encoding any one of the binding proteins disclosed herein. A further embodiment provides a vector comprising the isolated nucleic acid disclosed herein wherein said vector is selected from the group consisting of pcDNA; pTT (Durocher et al., *Nucleic Acids Research* 2002, Vol 30, No.2); pTT3 (pTT with additional multiple cloning site; pEFBOS (Mizushima, S. and Nagata, S., (1990) *Nucleic acids*
 25 *Research* Vol 18, No. 17); pBV; pJV; pcDNA3.1 TOPO, pEF6 TOPO and pBJ. In an embodiment, the vector is a vector disclosed in US Patent Application Serial No. 61/021,282.

In another aspect a host cell is transformed with the vector disclosed herein. In an embodiment, the host cell is a prokaryotic cell. In another embodiment, the host cell is E.Coli. In a related embodiment the host cell is a eukaryotic cell. In another embodiment, the eukaryotic
 30 cell is selected from the group consisting of protist cell, animal cell, plant cell and fungal cell. In yet another embodiment, the host cell is a mammalian cell including, but not limited to, CHO, COS; NS0, SP2, PER.C6 or a fungal cell such as *Saccharomyces cerevisiae*; or an insect cell such as Sf9.

In an embodiment, two or more DVD-Igs, e.g., with different specificities, are produced in a single recombinant host cell. For example, the expression of a mixture of antibodies has been called Oligoclonics™, (Merus B.V., The Netherlands) U.S. Patent Nos. 7,262,028; 7,429,486.

Another aspect of the invention provides a method of producing a binding protein disclosed herein comprising culturing any one of the host cells also disclosed herein in a culture medium under conditions sufficient to produce the binding protein. In an embodiment, 50%-75% of the binding protein produced by this method is a dual specific tetravalent binding protein. In a particular embodiment, 75%-90% of the binding protein produced by this method is a dual specific tetravalent binding protein. In a particular embodiment, 90%-95% of the binding protein produced is a dual specific tetravalent binding protein.

One embodiment provides a composition for the release of a binding protein wherein the composition comprises a formulation that in turn comprises a crystallized binding protein, as disclosed herein, and an ingredient, and at least one polymeric carrier. For example, the polymeric carrier is a polymer selected from one or more of the group consisting of: poly (acrylic acid), poly (cyanoacrylates), poly (amino acids), poly (anhydrides), poly (depsipeptide), poly (esters), poly (lactic acid), poly (lactic-co-glycolic acid) or PLGA, poly (b-hydroxybutyrate), poly (caprolactone), poly (dioxanone); poly (ethylene glycol), poly ((hydroxypropyl) methacrylamide, poly [(organo)phosphazene], poly (ortho esters), poly (vinyl alcohol), poly (vinylpyrrolidone), maleic anhydride- alkyl vinyl ether copolymers, pluronic polyols, albumin, alginate, cellulose and cellulose derivatives, collagen, fibrin, gelatin, hyaluronic acid, oligosaccharides, glycaminoglycans, sulfated polyeaccharides, blends and copolymers thereof. For example, the ingredient is selected from the group consisting of albumin, sucrose, trehalose, lactitol, gelatin, hydroxypropyl-β- cyclodextrin, methoxypolyethylene glycol and polyethylene glycol. Another embodiment provides a method for treating a mammal comprising the step of administering to the mammal an effective amount of the composition disclosed herein.

The invention also provides a pharmaceutical composition comprising a binding protein, as disclosed herein and a pharmaceutically acceptable carrier. In a further embodiment the pharmaceutical composition comprises at least one additional therapeutic agent for treating a disorder. For example, the additional agent is selected from the group consisting of: a therapeutic agent, an imaging agent, a cytotoxic agent, an angiogenesis inhibitor (including but not limited to an anti-VEGF antibody or a VEGF-trap), a kinase inhibitor (including but not limited to a KDR and a TIE-2 inhibitor), a co-stimulation molecule blocker (including but not limited to anti-B7.1, anti-B7.2, CTLA4-Ig, anti-CD20), an adhesion molecule blocker (including but not limited to an anti-LFA-1 antibody, an anti-E/L selectin antibody, a small molecule inhibitor), an anti-cytokine antibody or functional fragment thereof (including but not limited to an anti-IL-18, an anti-TNF, and an anti-IL-6/cytokine receptor antibody), methotrexate, cyclosporin, rapamycin, FK506, a

detectable label or reporter, a TNF antagonist, an antirheumatic, a muscle relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive,
 5 a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an antipsychotic, a stimulant, an asthma medication, a beta agonist, an inhaled steroid, an epinephrine or analog, a cytokine, and a cytokine antagonist.

In another aspect, the invention provides a method for treating a human subject suffering from a disorder in which the target, or targets, capable of being bound by the binding protein
 10 disclosed herein is detrimental, comprising administering to the human subject a binding protein disclosed herein such that the activity of the target, or targets in the human subject is inhibited and one of more symptoms is alleviated or treatment is achieved. For example, the disorder is selected from the group comprising arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus
 15 erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's
 20 granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison's disease,
 25 sporadic polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, spondyloarthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous
 30 disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases,
 35 Hepatitis B, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature

ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory
 interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial
 lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis
 associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease,
 5 systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated
 lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung
 disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced
 interstitial lung disease, fibrosis, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic
 pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty
 10 arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid
 hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated
 hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute
 immune disease associated with organ transplantation, chronic immune disease associated with
 organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis
 15 type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS,
 glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus
 erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all
 subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue
 disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute
 20 rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjörgren's syndrome,
 Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia,
 autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism
 (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic
 uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis,
 25 alcohol-induced liver injury, choleostasis, idiosyncratic liver disease, Drug-Induced hepatitis,
 Non-alcoholic Steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental
 disorders (*e.g.*, depression and schizophrenia), Th2 Type and Th1 Type mediated diseases, acute
 and chronic pain (different forms of pain), and cancers such as lung, breast, stomach, bladder,
 colon, pancreas, ovarian, prostate and rectal cancer and hematopoietic malignancies (leukemia
 30 and lymphoma), Abetalipoproteinemia, Acrocyanosis, acute and chronic parasitic or infectious
 processes, acute leukemia, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML),
 acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, arial
 ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic
 contact dermatitis, allergic rhinitis, allograft rejection, alpha-1- antitrypsin deficiency,
 35 amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3
 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral
 aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia,

atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, Burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors, cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical
5 degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob
10 disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, Dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, Diffuse Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's Syndrome in middle age, drug- induced movement disorders induced by drugs which block CNS dopamine
15 receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottitis, epstein-barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hematophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer, glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive sepsis,
20 granulomas due to intracellular organisms, hairy cell leukemia, Hallerorden-Spatz disease, hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis, hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, hepatitis (A), His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease, hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hypertension,
25 hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity, Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing radiation exposure, iridocyclitis/uveitis/optic neuritis, ischemia- reperfusion injury, ischemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney
30 transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system, lipedema, liver transplant rejection, lymphoderma, malaria, malignant Lymphoma, malignant histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic diseases, migraine headache, mitochondrial multi.system disorder, mixed connective tissue disease, monoclonal gammopathy, multiple myeloma, multiple systems degenerations (Mencel Dejerine-
35 Thomas Shi-Drager and Machado-Joseph), myasthenia gravis, mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease,

nephritis, nephrosis, neurodegenerative diseases, neurogenic I muscular atrophies, neutropenic fever, non- hodgkins lymphoma, occlusion of the abdominal aorta and its branches, occlusive arterial disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy reversal procedures, organomegaly, osteoporosis, pancreas transplant rejection, pancreatic carcinoma, paraneoplastic syndrome/hypercalcemia of malignancy, parathyroid transplant rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral vascular disorders, peritonitis, pernicious anemia, pneumocystis carinii pneumonia, pneumonia, POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, preeclampsia, Progressive supranucleo Palsy, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon and disease, Raynaud's disease, Refsum's disease, regular narrow QRS tachycardia, renovascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcomas, scleroderma, senile chorea, Senile Dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural lesions of the cerebellum, Subacute sclerosing panencephalitis, Syncope, syphilis of the cardiovascular system, systemic anaphalaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, Telangiectasia, thromboangitis obliterans, thrombocytopenia, toxicity, transplants, trauma/hemorrhage, type III hypersensitivity reactions, type IV hypersensitivity, unstable angina, uremia, urosepsis, urticaria, valvular heart diseases, varicose veins, ,vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, vital encephalitis/aseptic meningitis, vital-associated hemaphagocytic syndrome, Wernicke- Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue , acute coronary syndromes, acute idiopathic polyneuritis, acute inflammatory demyelinating polyradiculoneuropathy, acute ischemia, adult Still's disease, alopecia areata, anaphylaxis, anti-phospholipid antibody syndrome, aplastic anemia, arteriosclerosis, atopic eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic antiphospholipid syndrome, celiac disease, cervical spondylosis, chronic ischemia, cicatricial pemphigoid, clinically isolated syndrome (cis) with risk for multiple sclerosis, conjunctivitis, childhood onset psychiatric disorder, chronic obstructive pulmonary disease (COPD), dacryocystitis, dermatomyositis, diabetic retinopathy, diabetes mellitus, disk herniation, disk prolaps, drug induced immune hemolytic anemia, endocarditis, endometriosis, endophthalmitis, episcleritis, erythema multiforme, erythema multiforme major, gestational pemphigoid, Guillain-Barré

syndrome (GBS), hay fever, Hughes syndrome, idiopathic Parkinson's disease, idiopathic
 interstitial pneumonia, IgE-mediated allergy, immune hemolytic anemia, inclusion body myositis,
 infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart
 disease, inflammatory kidney disease, IPF/UIP, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul
 5 disease or Kussmaul-Meier disease, Landry's paralysis, Langerhan's cell histiocytosis, livedo
 reticularis, macular degeneration, microscopic polyangiitis, morbus bechterev, motor neuron
 disorders, mucous membrane pemphigoid, multiple organ failure, myasthenia gravis,
 myelodysplastic syndrome, myocarditis, nerve root disorders, neuropathy, non-A non-B hepatitis,
 optic neuritis, osteolysis, ovarian cancer, pauciarticular JRA, peripheral artery occlusive disease
 10 (PAOD), peripheral vascular disease (PVD), peripheral artery, disease (PAD), phlebitis,
 polyarteritis nodosa (or periarteritis nodosa), polychondritis, polymyalgia rheumatica, poliosis,
 polyarticular JRA, polyendocrine deficiency syndrome, polymyositis, polymyalgia rheumatica
 (PMR), post-pump syndrome, primary Parkinsonism, prostate and rectal cancer and
 hematopoietic malignancies (leukemia and lymphoma), prostatitis, pure red cell aplasia, primary
 15 adrenal insufficiency, recurrent neuromyelitis optica, restenosis, rheumatic heart disease, sapho
 (synovitis, acne, pustulosis, hyperostosis, and osteitis), scleroderma, secondary amyloidosis,
 shock lung, scleritis, sciatica, secondary adrenal insufficiency, silicone associated connective
 tissue disease, sneddon-wilkinson dermatosis, spondylitis ankylosans, Stevens-Johnson syndrome
 (SJS), systemic inflammatory response syndrome, temporal arteritis, toxoplasmic retinitis, toxic
 20 epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor receptor, type 1 allergic
 reaction, type II diabetes, urticaria, usual interstitial pneumonia (UIP), vasculitis, vernal
 conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular
 degeneration, wound healing, yersinia and salmonella associated arthropathy.

In an embodiment, diseases that can be treated or diagnosed with the compositions and
 25 methods of the invention include, but are not limited to, primary and metastatic cancers, including
 carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach,
 pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney, bladder
 and urothelium), female genital tract (including cervix, uterus, and ovaries as well as
 choriocarcinoma and gestational trophoblastic disease), male genital tract (including prostate,
 30 seminal vesicles, testes and germ cell tumors), endocrine glands (including the thyroid, adrenal,
 and pituitary glands), and skin, as well as hemangiomas, melanomas, sarcomas (including those
 arising from bone and soft tissues as well as Kaposi's sarcoma), tumors of the brain, nerves, eyes,
 and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas,
 neuroblastomas, Schwannomas, and meningiomas), solid tumors arising from hematopoietic
 35 malignancies such as leukemias, and lymphomas (both Hodgkin's and non-Hodgkin's
 lymphomas).

In an embodiment, the antibodies of the invention or antigen-binding portions thereof, are used to treat cancer or in the prevention of metastases from the tumors described herein either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

In another aspect the invention provides a method of treating a patient suffering from a disorder comprising the step of administering any one of the binding proteins disclosed herein before, concurrent, or after the administration of a second agent, as discussed herein. In a particular embodiment the second agent is selected from the group consisting of budenoside, epidermal growth factor, corticosteroids, cyclosporin, sulfasalazine, aminosalicylates, 6-mercaptopurine, azathioprine, metronidazole, lipoxigenase inhibitors, mesalamine, olsalazine, balsalazide, antioxidants, thromboxane inhibitors, IL-1 receptor antagonists, anti-IL-1 β mAbs, anti-IL-6 or IL-6 receptor mAbs, growth factors, elastase inhibitors, pyridinyl-imidazole compounds, antibodies or agonists of TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-12, IL-13, IL-15, IL-16, IL-18, IL-23, EMAP-II, GM-CSF, FGF, and PDGF, antibodies of CD2, CD3, CD4, CD8, CD-19, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands, methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, ibuprofen, corticosteroids, prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, IRAK, NIK, IKK, p38, MAP kinase inhibitors, IL-1 β converting enzyme inhibitors, TNF α converting enzyme inhibitors, T-cell signalling inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors, soluble p55 TNF receptor, soluble p75 TNF receptor, sIL-1RI, sIL-1RII, sIL-6R, antiinflammatory cytokines, IL-4, IL-10, IL-11, IL-13 and TGF β .

In a particular embodiment the pharmaceutical compositions disclosed herein are administered to the patient by at least one mode selected from parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelial, intracerebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, and transdermal.

One aspect of the invention provides at least one anti-idiotypic antibody to at least one binding protein of the present invention. The anti-idiotypic antibody includes any protein or peptide containing molecule that comprises at least a portion of an immunoglobulin molecule such as, but not limited to, at least one complementarily determining region (CDR) of a heavy or light chain or a ligand binding portion thereof, a heavy chain or light chain variable region, a

heavy chain or light chain constant region, a framework region, or any portion thereof, that can be incorporated into a binding protein of the present invention.

In another aspect, the invention provides a method for improving a characteristic of a binding protein of the invention comprising the steps of (a) determining the characteristic of the binding protein prior to alteration; (a) altering the length and/or sequence of $(X1)_1$ of the heavy and/or light chain thereby providing an altered heavy and/ or light chain; and (b) determining the improved characteristic of the altered binding protein comprising the altered heavy and light chains. In another embodiment, the invention provides a method for improving a characteristic of the binding protein of the invention comprising the steps of (a) determining the characteristic of the binding protein prior to alteration; (b) altering the first and second polypeptide chains such that $VD1-(X1)_n-VD2-C-(X2)_n$ is changed to $VD2-(X1)_n-VD1-C-(X2)_n$, thereby providing altered heavy and light chains; and (c) determining the improved characteristic of the altered binding protein comprising the altered heavy and light chains. In another embodiment, the invention provides a method for improving a characteristic of the binding protein of the invention comprising the steps of (a) determining the characteristic of the binding protein prior to alteration; (b) altering the first and/or second polypeptide chains such that the sequence of only one of the VD1 or VD2 of the heavy and/or light chain is changed; and (c) determining the characteristic of the altered binding protein comprising the altered heavy and light chains. The characteristic is selected from the group consisting of binding to target antigen, expression yield from host cell, in vitro halflife, in vivo halflife, stability, solubility, affinity, avidity, and improved effector function.

In an embodiment, the length of the linker of the altered heavy chain is increased. In another embodiment, the length of the linker of the altered heavy chain is decreased.

In an embodiment, the length of the linker of the altered light chain is increased. In another embodiment, the length of the linker of the altered light chain is decreased.

In yet another embodiment of the methods and compositions of the invention, the altered heavy and/ or light chain comprises a cleavage site. In an embodiment, the cleavage site is between at least one VD1 and VD2. In another embodiment, the cleavage site is in at least one linker. In an embodiment, the heavy and/ or light chain is cleaved by an enzyme or agent selected from the group consisting of enterokinase, thrombin, PreScission, Tobacco Etch Virus protease (TEV), and tissue plasminogen activator (tPA) + Proline. In another embodiment, the binding protein is cleaved by an enzyme or agent selected from the group consisting of a zinc-dependent endopeptidase, Matrix Metalloproteinase (MMP), a serralyisin, an astacin, an adamalysin, MMP-1; MMP-2; MMP-3; MMP-7; MMP-8; MMP-9; MMP-10; MMP-11; MMP-12; MMP-13; MMP-14; MMP-15; MMP-16; MMP-17; MMP-18; MMP-19; MMP-20; MMP-21; MMP-22; MMP-

23A; MMP-23B; MMP-24; MMP-25; MMP-26; MMP-27; MMP-28; a Disintegrin and Metalloproteinase (ADAM); ADAM17; ADAMTS1; ADAM1; ADAM10; ADAM8; ADAMTS4; ADAMTS13; ADAM12; ADAM15; ADAM9; ADAMTS5; ADAM33; ADAM11; ADAM2; ADAMTS2; ADAMTS9; ADAMTS3; ADAMTS7; ADAM22; ADAM28; ADAMTS12; ADAM19; ADAMTS8; ADAM29; ADAM23; ADAM3A; ADAM18; ADAMTS6; ADAM7; ADAMDES1; ADAM20; ADAM6; ADAM21; ADAM3B; ADAMTSL3; ADAMTSL4; ADAM30; ADAMTS20; ADAMTSL2; a Caspase; Caspases 1-12, Caspase 14; a Cathepsin; Cathepsin G; Cathepsin B; Cathepsin D; Cathepsin L1; Cathepsin C; Cathepsin K; Cathepsin S; Cathepsin H; Cathepsin A; Cathepsin E; Cathepsin L; Cathepsin Z; Cathepsin F; Cathepsin G-like 2; Cathepsin L-like 1; Cathepsin W; Cathepsin L-like 2; Cathepsin L-like 3; Cathepsin L-like 4; Cathepsin L-like 5; Cathepsin L-like 6; Cathepsin L-like 7; Cathepsin O; a Calpain; Calpain 3; Calpain 10; Calpain 1 (mu/l) large subunit; Calpain, small subunit 1; Calpain 2, (mu/l); large subunit; Calpain 9; Calpain 11; Calpain 5; Calpain 6; Calpain 13; Calpain 8; Calpain, small subunit 2; Calpain 15; Calpain 12; Calpain 7; and Calpain 8.

In an embodiment, at least one of the VD1 or VD2 does not bind to its target until a cleavage between the VD1 and VD2 occurs. In another embodiment, the linker of the binding protein has been selectively cleaved by an enzyme. In another embodiment, the linker of the binding protein has been selectively cleaved by an enzyme during the manufacturing process. In another embodiment, the linker of the binding protein has been selectively cleaved by an enzyme when the DVD-Ig is adjacent to at least one target. In another embodiment, the binding protein is selectively cleaved by an enzyme when the DVD-Ig is bound to at least one target.

In another aspect, the invention provides a method for treating a subject for a disease or a disorder by administering to the subject the binding protein of the invention, such that treatment is achieved. In an embodiment, at least one of a VD1 or VD2 does not bind its target until the binding protein is cleaved. In another embodiment, the VD2 does not bind its target until the binding protein is cleaved. In another embodiment, VD1 is released when the binding protein is cleaved. In another embodiment, the VD1 is released when the VD2 binds to its target.

Brief Description of the Drawings

Figure 1A is a schematic representation of Dual Variable Domain (DVD)-Ig constructs and shows the strategy for generation of a DVD-Ig from two parent antibodies;

Figure 1B, is a schematic representation of constructs DVD1-Ig, DVD2-Ig, and two chimeric mono-specific antibodies.

Detailed Description of the Invention

This invention pertains to multivalent and/or multispecific binding proteins capable of binding two or more antigens. Specifically, the invention relates to dual variable domain immunoglobulins (DVD-Ig), and pharmaceutical compositions thereof, as well as nucleic acids, recombinant expression vectors and host cells for making such DVD-Igs. Methods of using the DVD-Igs of the invention to detect specific antigens, either in vitro or in vivo are also encompassed by the invention.

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. The meaning and scope of the terms should be clear, however, in the event of any latent ambiguity, definitions provided herein take precedent over any dictionary or extrinsic definition. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. In this application, the use of "or" means "and/or" unless stated otherwise. Furthermore, the use of the term "including", as well as other forms, such as "includes" and "included", is not limiting. Also, terms such as "element" or "component" encompass both elements and components comprising one unit and elements and components that comprise more than one subunit unless specifically stated otherwise.

Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclatures used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques are used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

That the present invention may be more readily understood, select terms are defined below.

The term "polypeptide" as used herein, refers to any polymeric chain of amino acids. The terms "peptide" and "protein" are used interchangeably with the term polypeptide and also refer to a polymeric chain of amino acids. The term "polypeptide" encompasses native or artificial

proteins, protein fragments and polypeptide analogs of a protein sequence. A polypeptide may be monomeric or polymeric. Use of "polypeptide" herein is intended to encompass polypeptide and fragments and variants (including fragments of variants) thereof, unless otherwise contradicted by context. For an antigenic polypeptide, a fragment of polypeptide optionally contains at least one
5 contiguous or nonlinear epitope of polypeptide. The precise boundaries of the at least one epitope fragment can be confirmed using ordinary skill in the art. The fragment comprises at least about 5 contiguous amino acids, such as at least about 10 contiguous amino acids, at least about 15 contiguous amino acids, or at least about 20 contiguous amino acids. A variant of polypeptide is as described herein.

10 The term "isolated protein" or "isolated polypeptide" is a protein or polypeptide that by virtue of its origin or source of derivation is not associated with naturally associated components that accompany it in its native state; is substantially free of other proteins from the same species; is expressed by a cell from a different species; or does not occur in nature. Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from
15 which it naturally originates will be "isolated" from its naturally associated components. A protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art.

The term "recovering" as used herein, refers to the process of rendering a chemical species such as a polypeptide substantially free of naturally associated components by isolation,
20 e.g., using protein purification techniques well known in the art.

"Biological activity" as used herein, refers to any one or more inherent biological properties of a molecule (whether present naturally as found in vivo, or provided or enabled by recombinant means). Biological properties include but are not limited to binding receptor; induction of cell proliferation, inhibiting cell growth, inductions of other cytokines, induction of
25 apoptosis, and enzymatic activity. Biological activity also includes activity of an Ig molecule.

The terms "specific binding" or "specifically binding", as used herein, in reference to the interaction of an antibody, a protein, or a peptide with a second chemical species, mean that the interaction is dependent upon the presence of a particular structure (e.g., an antigenic determinant or epitope) on the chemical species; for example, an antibody recognizes and binds to a specific
30 protein structure rather than to proteins generally. If an antibody is specific for epitope "A", the presence of a molecule containing epitope A (or free, unlabeled A), in a reaction containing labeled "A" and the antibody, will reduce the amount of labeled A bound to the antibody.

The term "antibody", as used herein, broadly refers to any immunoglobulin (Ig) molecule comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains, or any
35 functional fragment, mutant, variant, or derivation thereof, which retains the essential epitope

binding features of an Ig molecule. Such mutant, variant, or derivative antibody formats are known in the art. Nonlimiting embodiments of which are discussed below.

In a full-length antibody, each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain
5 constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework
10 regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. Immunoglobulin molecules can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG 1, IgG2, IgG 3, IgG4, IgA1 and IgA2) or subclass.

The term "Fc region" is used to define the C-terminal region of an immunoglobulin heavy
15 chain, which may be generated by papain digestion of an intact antibody. The Fc region may be a native sequence Fc region or a variant Fc region. The Fc region of an immunoglobulin generally comprises two constant domains, a CH2 domain and a CH3 domain, and optionally comprises a CH4 domain. Replacements of amino acid residues in the Fc portion to alter antibody effector function are known in the art (Winter, *et al.* US Patent Nos 5,648,260 and 5,624,821). The Fc
20 portion of an antibody mediates several important effector functions e.g., cytokine induction, ADCC, phagocytosis, complement dependent cytotoxicity (CDC) and half-life/ clearance rate of antibody and antigen-antibody complexes. In some cases these effector functions are desirable for therapeutic antibody but in other cases might be unnecessary or even deleterious, depending on the therapeutic objectives. Certain human IgG isotypes, particularly IgG1 and IgG3, mediate
25 ADCC and CDC via binding to FcγRs and complement C1q, respectively. Neonatal Fc receptors (FcRn) are the critical components determining the circulating half-life of antibodies. In still another embodiment at least one amino acid residue is replaced in the constant region of the antibody, for example the Fc region of the antibody, such that effector functions of the antibody are altered. The dimerization of two identical heavy chains of an immunoglobulin is mediated by
30 the dimerization of CH3 domains and is stabilized by the disulfide bonds within the hinge region (Huber et al. Nature; 264: 415-20; Thies et al 1999 J Mol Biol; 293: 67-79.). Mutation of cysteine residues within the hinge regions to prevent heavy chain-heavy chain disulfide bonds will destabilize dimeration of CH3 domains. Residues responsible for CH3 dimerization have been identified (Dall'Acqua 1998 Biochemistry 37: 9266-73.). Therefore, it is possible to generate a
35 monovalent half-Ig. Interestingly, these monovalent half Ig molecules have been found in nature for both IgG and IgA subclasses (Seligman 1978 Ann Immunol 129: 855-70; Biewenga et al 1983

Clin Exp Immunol 51: 395-400). The stoichiometry of FcRn: Ig Fc region has been determined to be 2:1 (West et al .2000 Biochemistry 39: 9698-708), and half Fc is sufficient for mediating FcRn binding (Kim et al 1994 Eur J Immunol; 24: 542-548.). Mutations to disrupt the dimerization of CH3 domain may not have greater adverse effect on its FcRn binding as the residues important for CH3 dimerization are located on the inner interface of CH3 b sheet structure, whereas the region responsible for FcRn binding is located on the outside interface of CH2-CH3 domains. However the half Ig molecule may have certain advantage in tissue penetration due to its smaller size than that of a regular antibody. In one embodiment at least one amino acid residue is replaced in the constant region of the binding protein of the invention, for example the Fc region, such that the dimerization of the heavy chains is disrupted, resulting in half DVD Ig molecules. The anti-inflammatory activity of IgG is completely dependent on sialylation of the N-linked glycan of the IgG Fc fragment. The precise glycan requirements for anti-inflammatory activity has been determined, such that an appropriate IgG1 Fc fragment can be created, thereby generating a fully recombinant, sialylated IgG1 Fc with greatly enhanced potency (Anthony, R.M., et al. (2008) Science 320:373-376).

The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen. It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Such antibody embodiments may also be bispecific, dual specific, or multi-specific formats; specifically binding to two or more different antigens. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546, Winter et al., PCT publication WO 90/05144 A1 herein incorporated by reference), which comprises a single variable domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too

short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see *e.g.*, Holliger, P., *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994) *Structure* 2:1121-1123). Such antibody binding portions are known in the art
5 (Kontermann and Dubel eds., Antibody Engineering (2001) Springer-Verlag. New York. 790 pp. (ISBN 3-540-41354-5). In addition single chain antibodies also include "linear antibodies" comprising a pair of tandem Fv segments (VH-CH1-VH-CH1) which, together with complementary light chain polypeptides, form a pair of antigen binding regions (Zapata et al. Protein Eng. 8(10):1057-1062 (1995); and US Patent No. 5,641,870).

10 The term "multivalent binding protein" is used throughout this specification to denote a binding protein comprising two or more antigen binding sites. In an embodiment, the multivalent binding protein is engineered to have the three or more antigen binding sites, and is generally not a naturally occurring antibody. The term "multispecific binding protein" refers to a binding protein capable of binding two or more related or unrelated targets. Dual variable domain (DVD)
15 binding proteins of the invention comprise two or more antigen binding sites and are tetravalent or multivalent binding proteins. DVDs may be monospecific, i.e., capable of binding one antigen or multispecific, i.e. capable of binding two or more antigens. DVD binding proteins comprising two heavy chain DVD polypeptides and two light chain DVD polypeptides are referred to as DVD-Ig. Each half of a DVD-Ig comprises a heavy chain DVD polypeptide, and a light chain
20 DVD polypeptide, and two antigen binding sites. Each binding site comprises a heavy chain variable domain and a light chain variable domain with a total of 6 CDRs involved in antigen binding per antigen binding site.

The term "bispecific antibody", as used herein, refers to full-length antibodies that are generated by quadroma technology (see Milstein, C. and A.C. Cuello, *Nature*, 1983. 305(5934): p.
25 537-40), by chemical conjugation of two different monoclonal antibodies (see Staerz, U.D., et al., *Nature*, 1985. 314(6012): p. 628-31), or by knob-into-hole or similar approaches which introduces mutations in the Fc region (see Holliger, P., T. Prospero, and G. Winter, *Proc Natl Acad Sci U S A*, 1993. 90(14): p. 6444-8.18), resulting in multiple different immunoglobulin species of which only one is the functional bispecific antibody. By molecular function, a bispecific antibody binds
30 one antigen (or epitope) on one of its two binding arms (one pair of HC/LC), and binds a different antigen (or epitope) on its second arm (a different pair of HC/LC). By this definition, a bispecific antibody has two distinct antigen binding arms (in both specificity and CDR sequences), and is monovalent for each antigen it binds to.

The term "dual-specific antibody", as used herein, refers to full-length antibodies that can
35 bind two different antigens (or epitopes) in each of its two binding arms (a pair of HC/LC) (see PCT publication WO 02/02773). Accordingly a dual-specific binding protein has two identical

antigen binding arms, with identical specificity and identical CDR sequences, and is bivalent for each antigen it binds to.

A "functional antigen binding site" of a binding protein is one that is capable of binding a target antigen. The antigen binding affinity of the antigen binding site is not necessarily as strong as the parent antibody from which the antigen binding site is derived, but the ability to bind antigen must be measurable using any one of a variety of methods known for evaluating antibody binding to an antigen. Moreover, the antigen binding affinity of each of the antigen binding sites of a multivalent antibody herein need not be quantitatively the same.

The term "cytokine" is a generic term for proteins released by one cell population, which act on another cell population as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental lactogen; tumor necrosis factor-alpha and -beta; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors such as NGF-alpha; platelet-growth factor; placental growth factor, transforming growth factors (TGFs) such as TGF- alpha and TGF-beta; insulin-like growth factor-1 and -11; erythropoietin (EPO); osteoinductive factors; interferons such as interferon-alpha, -beta and -gamma colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-18, IL-21, IL-22, IL-23, IL-33; a tumor necrosis factor such as TNF-alpha or TNF-beta; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

The term "linker" is used to denote polypeptides comprising two or more amino acid residues joined by peptide bonds and are used to link one or more antigen binding portions. Such linker polypeptides are well known in the art (see *e.g.*, Holliger, P., *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994) *Structure* 2:1121-1123). Exemplary linkers include, but are not limited to, AKTTPKLEEGEFSEAR (SEQ ID NO: 1); AKTTPKLEEGEFSEARV (SEQ ID NO: 2); AKTTPKLGG (SEQ ID NO: 3); SAKTTPKLGG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5); RADAAP (SEQ ID NO: 6); RADAAPTVS (SEQ ID NO: 7); RADAAAAGGPGS (SEQ ID NO: 8); RADAAAA(G₄S)₄ (SEQ ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP (SEQ ID NO: 11); ADAAPTVSIFPP

(SEQ ID NO: 12); TVAAP (SEQ ID NO: 13); TVAAPSVFIFPP (SEQ ID NO: 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID NO: 16); AKTTPP (SEQ ID NO: 17); AKTTPPSVTPLAP (SEQ ID NO: 18); AKTTAP (SEQ ID NO: 19); AKTTAPSVYPLAP (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21); ASTKGPSVFPLAP (SEQ ID NO: 22),
5 GGGGSGGGGSGGGGS (SEQ ID NO: 23); GENKVEYAPALMALS (SEQ ID NO: 24); GPAKELTPLKEAKVS (SEQ ID NO: 25); GHEAAVMQVQYPAS (SEQ ID NO: 26); GGGGGGGP (SEQ ID NO: 27); GGGGGGGGP (SEQ ID NO: 28); PAPNLLGGP (SEQ ID NO: 29); PNLLGGP (SEQ ID NO: 30); GGGGGGP (SEQ ID NO: 31); PAPELLGGP (SEQ ID NO: 32); PTISPAPNLLGGP (SEQ ID NO: 33); TVAADDDDKSVFIVPP (SEQ ID NO: 34);
10 TVDDDDKAAP (SEQ ID NO: 35); LVPRGSAAP (SEQ ID NO: 36); ASDDDDK GGP (SEQ ID NO: 37); ALVPR GSGP (SEQ ID NO: 38); ASTDDDDK SVFPLAP (SEQ ID NO: 39); TVALVPR GSVFIFPP (SEQ ID NO: 40); ASTLVPR GSVFPLAP (SEQ ID NO: 41); TVAADDDK SVFIVPP (SEQ ID NO: 42); ASTDDDK SVFPLAP (SEQ ID NO: 43); LEVLFGP (SEQ ID NO: 44); TAAALEVLFG GPAP (SEQ ID NO: 45); ASTLEVLFG GPLAP (SEQ ID NO: 46); PAPLEVLFG GP (SEQ ID NO: 47); TAENLYFQ GAP (SEQ ID NO: 48); AENLYFQ GA (SEQ ID NO: 49); PGPFGR SAGGP (SEQ ID NO: 50); PGPFGR SAGG (SEQ ID NO: 51); PQRGR SAG (SEQ ID NO: 52); PHYGR SGG (SEQ ID NO: 53); GPFGR SAGP (SEQ ID NO: 54); GDDDDK GGP (SEQ ID NO: 55); AGDDDDK GGP (SEQ ID NO: 56); GGDDDDK GGP (SEQ ID NO: 57); AS; TVA; ASTK (SEQ ID NO: 58); ASTKGPSV (SEQ ID NO: 59);
20 ASTKGPSVFP (SEQ ID NO: 60); TVAAPSV (SEQ ID NO: 61), TVAAPSVFI (SEQ ID NO: 62).

In some instances, the DVD-Ig architecture, in particular the linkers connecting the VL1 – VL2 and VH1 – VH2, may impose constraints on ligand binding to the inner (C terminal) domain. In an embodiment, the DVD-Ig is cleavable by an enzyme so that such constraints are ameliorated or removed. In an embodiment, the DVD-Ig is cleavable by an enzyme between the
25 VD1 (VH1, VL1) and VD2 (VH2, VL2) domains of at least one of a heavy chain and a light chain. In an embodiment, a cleavable linker joins the VD1 and VD2 domain of at least one of a heavy chain and a light chain. In an embodiment, the DVD-Ig is cleaved by enterokinase, thrombin, PreScission, Tobacco Etch Virus protease (TEV), and tissue plasminogen activator
30 (tPA) + Proline.

In an embodiment, cleavable linkers between the outer variable domain (N terminal) and the inner variable domain (C terminal) of the DVD-Ig, or a single chain (VH/VL no linkers) attachment of the outer variable domain to the inner variable domain of the DVD-Ig via the VH1 – VH2 or VL1 – VL2 linkage were tested to attempt to overcome such potential constraints.
35 Inner domain binding can also be modulated by linker length. The VH1 – VH2 and VL1 – VL2 linkers can consist of, but are not limited to, sequences derived from CH1/CL, the hinge region of

IgG, and sequences such as glycine – serine repeats. Providing such alternative linkers may (1) return full antigen affinity to the inner antigen binding site of a DVD-Ig; (2) modulate the affinity of the inner antigen binding site of the DVD-Ig; (3) activate of the inner antigen binding site of a DVD-Ig at the site of action; (4) increase avidity of a DVD-Ig for an antigen; and/or (5) release the outer variable domain as an Fv unit.

The attachment of the outer domain Fv is accomplished in a number of ways. First, a protease cleavage site is engineered into the sequence of the linker. This is performed on one or both of the heavy or light chain linkers connecting the inner or outer variable domains of the DVD-Ig. The protease cleavage sequence is selected from any number of protease cleavage sequences (see Tables 1, 2, 5, 6, and Example 2.3, for example). Ideally, the protease is selected to cleave at a specific sequence, and does not readily cleave other sites in the DVD-Ig. The DVD-Ig construct containing a protease cleavage sequence is proteolytically cleaved in the manufacturing process itself or, if the selected protease is expressed by the targeted cell type, the DVD-Ig is cleaved at the targeted tissue or by another endogenous protease. The DVD-Ig is thereby considered a pro-drug that is activated at the intended therapeutic site. Additionally, such a mechanism allows the DVD-Ig to mask a particular antigen binding affinity at the inner binding site until it is activated at the targeted therapeutic site. Another advantage to the compositions and methods of the invention is the ability to increase the affinity for a particular targeted antigen in a site – specific manner. This is accomplished via avidity in the following manner. If the DVD-Ig contains antigen binding domains that bind two different epitopes on the same antigen, it is possible with the increased flexibility of the outer variable domain for it to bind twice - but at different locations - to the same antigen, thereby increasing the apparent affinity for that particular antigen via avidity. A DVD – Ig with cleavable linkers can also be designed that targets two distinct antigens (on the same cells or two different cells) to enhance affinity (by avidity) in a site – specific manner. For instance the two antigens of interest may be expressed on a tumor simultaneously, but individually on normal cells. Thus, binding of two DVD-Ig domains on the tumor cell with increased avidity will improve tumor binding specificity and reduce DVD-Ig side effects (toxicity) on normal cells. It is also foreseen that a disulfide linkage between the heavy and light chains, or specific mutations in the VH and VL, of the outer variable domain could be engineered to maintain their connectivity in vivo. It is also advantageous, in some cases, to cleave the entire outer Fv from the DVD – Ig using the same cleavage technology resulting in free Fv's and IgG's. The Fv cleavage could occur before or following binding of the target to the outer domain.

Table 1: Examples of Linkers with Protease Cleavage Sites (indicated by gap) in VL1 – VL2 or VH1 – VH2 Linkers

Heavy Chain linker (VH1 – VH2)	SEQ ID NO	Light Chain Linker (VL1 – VL2)	SEQ ID NO	Description
ASTKGP	21	TVDDDDDK AAP	35	Short HC linker; EK cleavable LC linker
ASDDDDDK GGP	37	TVAAP	13	EK cleavable HC linker; Short LC linker
ASTKGP	21	LVPR GSAAP	36	Short HC linker; Thrombin cleavable LC linker
ALVPR GSGP	38	TVAAP	13	Thrombin cleavable HC linker; Short LC linker
ASTKGP	21	TVAADDDDK SVFIVPP	34	Short HC linker; EK cleavable longer LC linker
ASTDDDDDK SVFPLAP	39	TVAAP	13	EK cleavable longer HC linker; Short LC linker
ASTKGP	21	TVALVPR GSVFIFPP	40	Short HC linker; Thrombin cleavable longer LC linker
ASTLVPR GSVFPLAP	41	TVAAP	3	Thrombin cleavable longer HC linker; Short LC linker
ASTKGPSVFPLAP	22	TVDDDDDK AAP	35	Long HC linker; EK cleavable LC linker
ASDDDDDK GGP	37	TVAAPSVFIFPP	14	EK cleavable HC linker; Long LC linker
ASTKGPSVFPLAP	22	LVPR GSAAP	36	Long HC linker; Thrombin cleavable LC linker
ALVPR GSGP	38	TVAAPSVFIFPP	14	Thrombin cleavable HC linker; Long LC linker
ASTKGPSVFPLAP	22	TVAADDDDK SVFIVPP	34	Long HC linker; EK cleavable longer LC linker
ASTDDDK SVFPLAP	43	TVAAPSVFIFPP	14	EK cleavable longer HC linker; Long LC linker
ASTKGPSVFPLAP	22	TVALVPR GSVFIFPP	40	Long HC linker; Thrombin cleavable longer LC linker
ASTLVPR GSVFPLAP	41	TVAAPSVFIFPP	14	Thrombin cleavable Longer HC linker; Long LC linker
ASTKGP	21	LEVLFQ GP	44	Short HC linker; PreScission cleavable LC linker
LEVLFQ GP	44	TVAAP	13	PreScission cleavable HC linker; Short LC linker
ASTKGP	21	TVAALEVLFQ GPAP	45	Short HC linker; PreScission cleavable Longer LC
ASTLEVLFQ GPLAP	46	TVAAP	3	PreScission cleavable longer HC linker; Short LC linker
ASTKGPSVFPLAP	22	LEVLFQ GP	44	Long HC linker; PreScission cleavable LC linker
LEVLFQ GP	44	TVAAPSVFIFPP	14	PreScission cleavable HC linker; Long LC linker
ASTKGPSVFPLAP	22	TVAALEVLFQ GPAP	45	Long HC linker; PreScission cleavable longer LC linker
ASTLEVLFQ GPLAP	46	TVAAPSVFIFPP	14	PreScission cleavable longer HC linker; Long LC linker
PAPLEVLFQ GP	47			PreScission cleavable longer HC linker based on Hinge
		PAPLEVLFQ GP	47	PreScission cleavable longer LC linker based on Hinge
ASTKGP	21	TAENLYFQ GAP	48	Short HC linker; TEV cleavable LC linker
AENLYFQ GA	49	TVAAPSVFIFPP	14	Shorter TEV cleavable HC linker; Long LC linker
AENLYFQ GA	49	PAPNLLGGP	29	Shorter TEV cleavable HC linker; Long LC linker based on Hinge (9)

Table 2: Examples of Linkers with Protease Cleavage Sites (indicated by gap) in Both the VH1 – VH2 and the VL1 – VL2 Chain Linkers

Heavy Chain (VH1 – VH2)	SEQ ID NO	Light Chain (VL1 – VL2)	SEQ ID NO	Comments
ASDDDDK GGP	37	TVALVPR GSVFIFPP	40	Shorter EK cleavable HC linker; Longer Thrombin cleavable LC linker
LEVLFQ GP		TVAADDDDK SVFIVPP	34	Shorter PreScission cleavable HC linker; Longer EK cleavable LC linker
ALVPR GSGP	38	TVAALEVLFQ GPAP	45	Shorter Thrombin cleavable HC linker; Longer PreScission cleavable LC linker
LEVLFQ GP	44	TAENLYFQ GAP	48	Shorter PreScission cleavable HC linker; Longer TEV cleavable LC linker
AENLYFQ GA	49	TVAADDDDK SVFIVPP	34	Shorter TEV cleavable HC linker; Longer EK cleavable LC linker
PGPFGR SAGGP	50			Tissue plasminogen activator (tPA) + Proline closest to hinge-like sequence
		PGPFGR SAGGP	50	tPA + Proline closest to hinge-like sequence
PGPFGR SAGG	51			tPA + hinge - can be used for either HC or LC linker
PQRGR SAG	52			tPA + hinge - can be used for either HC or LC linker
PHYGR SGG	53			tPA + hinge - can be used for either HC or LC linker
GPFGR SAGP	54			tPA + hinge - can be used for either HC or LC linker
GDDDDK GGP	55			EK – short linker can be used for either HC or LC linker
AGDDDDK GGP	56			EK – short linker can be used for either HC or LC linker
GGDDDDK GGP	57			EK – short linker can be used for either HC or LC linker

5

Although the introduction of a cleavage site may be achieved by the insertion of a linker, other methods for insertion of a cleavage site into a protein are well known in the art. Any of a number of proteases target sites may be introduced into a DVD-Ig according to the compositions and methods of the invention such that the DVD-Ig is cleaved by an enzyme of choice and/or where and when it is desired (e.g., during the manufacturing process of at a certain location or time of choice in the body).

10

Enzymes useful in the practice of the invention are extensive and well known in the art (Barrett, D., Rawlings, N.D., Woessner, J.F. (2004) Handbook of Proteolytic Enzymes, Academic Press; Rawlings, N., Morton, F., Barrett, A. (2006) Nucleic Acid Research 34, database issue,

D270-D272; International Proteolysis Society (IPS) websites <http://www.protease.org/index.html> and <http://www.protease.org/blog.html>; Merops (database of all proteases) website <http://merops.sanger.ac.uk/>; <http://www.ihop-net.org/UniPub/iHOP/>).

In an embodiment, the DVD-Ig is cleaved by a protease of the metzincin superfamily, which is a family of zinc-dependent endopeptidases, such a Matrix Metalloproteinase (MMP), serralsins, astacins, adamalysins. MMPs can degrade the extracellular matrix surrounding cells and tissues and therefore play a role in tissue degradation and repair in a number of disease states. In cancer, MMPs degrade the extracellular matrix surrounding the cancer cells enabling cancer cell progression and metastasis. Other specific MMP's are expressed in different tissue types or pathological processes and therefore are considered as tissue specific proteases for diseases related to those tissues or disease states such as arthritis, tissue repair, cirrhosis, angiogenesis, metastasis, and morphogenesis. The cleavage motifs for these enzymes are known in the art. See for example Tables 1, 4A, 4B, and 4D in Turk, B.E. et al. (2001) Nature Biotechnology 19:661-667 for the motifs/sequences cleaved by MMPs. Exemplary MMPs include, for example, MMP-1; MMP-2; MMP-3; MMP-7; MMP-8; MMP-9; MMP-10; MMP-11; MMP-12; MMP-13; MMP-14; MMP-15; MMP-16; MMP-17; MMP-18; MMP-19; MMP-20; MMP-21; MMP-22; MMP-23A; MMP-23B; MMP-24; MMP-25; MMP-26; MMP-27; and MMP-28.

In an embodiment, the DVD-Ig is cleaved by a Disintegrin and Metalloproteinase (ADAM). For example ADAM-17 (or TACE) is a TNF-alpha converting enzyme. ADAMs proteinases are important in cell-cell interactions, and protein ectodomain shedding (e.g., cleaving off parts of other important cell surface proteins to make them active). For example, HER-2 is activated when cleaved by ADAM-10, which leads to cell proliferation. ADAM-10 is also just now being recognized as playing a role in glioblastoma cell migration via cleavage of N-cadherin. See, for example, J Neuroscience (2009); 29 (14): 4605-15 (2009) and Cancer Biol Ther. (2006) Jun; 5(6):657-64. For example, HER-2 is cleaved by ADAM-10, which plays a role in glioblastoma cell migration. See, for example, Caescu, C.I. et al. (2009) Biochem J. 424:79-88 for the motifs/sequences cleaved by these two enzymes. Exemplary ADAMs include, for example, ADAM17; ADAMTS1; ADAM1; ADAM10; ADAM8; ADAMTS4; ADAMTS13; ADAM12; ADAM15; ADAM9; ADAMTS5; ADAM33; ADAM11; ADAM2; ADAMTS2; ADAMTS9; ADAMTS3; ADAMTS7; ADAM22; ADAM28; ADAMTS12; ADAM19; ADAMTS8; ADAM29; ADAM23; ADAM3A; ADAM18; ADAMTS6; ADAM7; ADAMDES1; ADAM20; ADAM6; ADAM21; ADAM3B; ADAMTSL3; ADAMTSL4; ADAM30; ADAMTS20; and ADAMTSL2.

In an embodiment, DVD-Ig is cleaved by a Caspase (*cysteine-aspartic proteases*) which is a family of cysteine proteases that play essential roles in apoptosis, necrosis, and inflammation).

Some caspases are also required for the maturation of cytokines. Exemplary Caspases include, for example, Caspases 1-12, and 14.

In an embodiment, the DVD-Ig is cleaved by a Cathepsin. Cathepsins have a role in mammalian cellular turnover, e.g., neoplasia, metastasis, bone resorption; inflammation,
5 degenerative joint diseases. Exemplary Cathepsins include, for example, Cathepsin G; Cathepsin B; Cathepsin D; Cathepsin L1; Cathepsin C; Cathepsin K; Cathepsin S; Cathepsin H; Cathepsin A; Cathepsin E; Cathepsin L; Cathepsin Z; Cathepsin F; Cathepsin G-like 2; Cathepsin L-like 1; Cathepsin W; Cathepsin L-like 2; Cathepsin L-like 3; Cathepsin L-like 4; Cathepsin L-like 5; Cathepsin L-like 6; Cathepsin L-like 7; or Cathepsin O.

10 In an embodiment, the DVD-Ig is cleaved by a Calpain. Calpains are calcium activated cysteine proteases that play roles in cell cycle, neuronal function and memory, and blood vessels and clotting. Exemplary Calpains include, for example, Calpain 3; Calpain 10; Calpain 1 (mu/l) large subunit; Calpain, small subunit 1; Calpain 2, (mu/l); large subunit; Calpain 9; Calpain 11; Calpain 5; Calpain 6; Calpain 13; Calpain 8; Calpain, small subunit 2; Calpain 15; Calpain 12;
15 Calpain 7; or Calpain 8.

The term "immunoglobulin constant domain" refers to a heavy or light chain constant domain. Human IgG heavy chain and light chain constant domain amino acid sequences are known in the art.

The term "monoclonal antibody" or "mAb" as used herein refers to an antibody obtained
20 from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigen. Furthermore, in contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each mAb is directed
25 against a single determinant on the antigen. The modifier "monoclonal" is not to be construed as requiring production of the antibody by any particular method.

The term "human antibody", as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the invention may include amino acid residues not encoded by human
30 germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs and in particular CDR3. However, the term "human antibody", as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies expressed using a recombinant expression vector transfected into a host cell (described further in Section II C, below), antibodies isolated from a recombinant, combinatorial human antibody library (Hoogenboom H.R. (1997) TIB Tech. 15:62-70; Azzazy H., and Highsmith W.E. (2002) Clin. Biochem. 35:425-445; Gavilondo J.V., and Larrick J.W. (2002) BioTechniques 29:128-145; Hoogenboom H., and Chames P. (2000) Immunology Today 21:371-378), antibodies isolated from an animal (e.g., a mouse) that is transgenic for human immunoglobulin genes (see, Taylor, L. D., et al. (1992) Nucl. Acids Res. 20:6287-6295; Kellermann S-A. and Green L.L. (2002) Current Opinion in Biotechnology 13:593-597; Little M. et al. (2000) Immunology Today 21:364-370) or antibodies prepared, expressed, created or isolated by any other means that involves splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable and constant regions derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies are subjected to in vitro mutagenesis (or, when an animal transgenic for human Ig sequences is used, in vivo somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire in vivo.

An "affinity matured" antibody is an antibody with one or more alterations in one or more CDRs thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Exemplary affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks et al. *BidlTechnology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas et al. *Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier et al. *Gene* 169:147- 155 (1995); Yelton et al. *J. Immunol.* 155:1994-2004 (1995); Jackson et al., *J. Immunol.* 154(7):3310-9 (1995); Hawkins et al, *J. Mol. Biol* 226:889-896 (1992) and selective mutation at selective mutagenesis positions, contact or hypermutation positions with an activity enhancing amino acid residue as described in US patent US 6914128B1.

The term "chimeric antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species and constant region sequences from another species, such as antibodies having murine heavy and light chain variable regions linked to human constant regions.

The term "CDR-grafted antibody" refers to antibodies which comprise heavy and light chain variable region sequences from one species but in which the sequences of one or more of the CDR regions of VH and/or VL are replaced with CDR sequences of another species, such as antibodies having murine heavy and light chain variable regions in which one or more of the murine CDRs (e.g., CDR3) has been replaced with human CDR sequences.

The term "humanized antibody" refers to antibodies which comprise heavy and light chain variable region sequences from a non-human species (e.g., a mouse) but in which at least a portion of the VH and/or VL sequence has been altered to be more "human-like", i.e., more similar to human germline variable sequences. One type of humanized antibody is a CDR-grafted antibody, in which human CDR sequences are introduced into non-human VH and VL sequences to replace the corresponding nonhuman CDR sequences. Also "humanized antibody" is an antibody or a variant, derivative, analog or fragment thereof which immunospecifically binds to an antigen of interest and which comprises a framework (FR) region having substantially the amino acid sequence of a human antibody and a complementary determining region (CDR) having substantially the amino acid sequence of a non-human antibody. As used herein, the term "substantially" in the context of a CDR refers to a CDR having an amino acid sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the amino acid sequence of a non-human antibody CDR. A humanized antibody comprises substantially all of at least one, and typically two, variable domains (Fab, Fab', F(ab')₂, FabC, Fv) in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin (i.e., donor antibody) and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. In an embodiment, a humanized antibody also comprises at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. In some embodiments, a humanized antibody contains both the light chain as well as at least the variable domain of a heavy chain. The antibody also may include the CH1, hinge, CH2, CH3, and CH4 regions of the heavy chain. In some embodiments, a humanized antibody only contains a humanized light chain. In some embodiments, a humanized antibody only contains a humanized heavy chain. In specific embodiments, a humanized antibody only contains a humanized variable domain of a light chain and/or humanized heavy chain.

The terms "Kabat numbering", "Kabat definitions" and "Kabat labeling" are used interchangeably herein. These terms, which are recognized in the art, refer to a system of numbering amino acid residues which are more variable (*i.e.* hypervariable) than other amino acid residues in the heavy and light chain variable regions of an antibody, or an antigen binding portion thereof (Kabat *et al.* (1971) *Ann. NY Acad. Sci.* 190:382-391 and, Kabat, E.A., *et al.* (1991) *Sequences of Proteins of Immunological Interest, Fifth Edition*, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). For the heavy chain variable region, the

hypervariable region ranges from amino acid positions 31 to 35 for CDR1, amino acid positions 50 to 65 for CDR2, and amino acid positions 95 to 102 for CDR3. For the light chain variable region, the hypervariable region ranges from amino acid positions 24 to 34 for CDR1, amino acid positions 50 to 56 for CDR2, and amino acid positions 89 to 97 for CDR3.

5 As used herein, the term "CDR" refers to the complementarity determining region within antibody variable sequences. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The term "CDR set" as used herein refers to a group of three CDRs that occur in a single variable region capable of binding the antigen. The exact boundaries of these CDRs have been
10 defined differently according to different systems. The system described by Kabat (Kabat et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987) and (1991)) not only provides an unambiguous residue numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the three CDRs. These CDRs may be referred to as Kabat CDRs. Chothia and coworkers (Chothia & Lesk,
15 J. Mol. Biol. 196:901-917 (1987) and Chothia et al., Nature 342:877-883 (1989)) found that certain sub- portions within Kabat CDRs adopt nearly identical peptide backbone conformations, despite having great diversity at the level of amino acid sequence. These sub-portions were designated as L1, L2 and L3 or H1, H2 and H3 where the "L" and the "H" designates the light chain and the heavy chains regions, respectively. These regions may be referred to as Chothia
20 CDRs, which have boundaries that overlap with Kabat CDRs. Other boundaries defining CDRs overlapping with the Kabat CDRs have been described by Padlan (FASEB J. 9:133-139 (1995)) and MacCallum (J Mol Biol 262(5):732-45 (1996)). Still other CDR boundary definitions may not strictly follow one of the herein systems, but will nonetheless overlap with the Kabat CDRs, although they may be shortened or lengthened in light of prediction or experimental findings that
25 particular residues or groups of residues or even entire CDRs do not significantly impact antigen binding. The methods used herein may utilize CDRs defined according to any of these systems, although certain embodiments use Kabat or Chothia defined CDRs.

As used herein, the term "framework" or "framework sequence" refers to the remaining sequences of a variable region minus the CDRs. Because the exact definition of a CDR sequence
30 can be determined by different systems, the meaning of a framework sequence is subject to correspondingly different interpretations. The six CDRs (CDR-L1, -L2, and -L3 of light chain and CDR-H1, -H2, and -H3 of heavy chain) also divide the framework regions on the light chain and the heavy chain into four sub-regions (FR1, FR2, FR3 and FR4) on each chain, in which CDR1 is positioned between FR1 and FR2, CDR2 between FR2 and FR3, and CDR3 between FR3 and
35 FR4. Without specifying the particular sub-regions as FR1, FR2, FR3 or FR4, a framework region, as referred by others, represents the combined FR's within the variable region of a single,

naturally occurring immunoglobulin chain. As used herein, a FR represents one of the four sub-regions, and FRs represents two or more of the four sub- regions constituting a framework region.

As used herein, the term "germline antibody gene" or "gene fragment" refers to an immunoglobulin sequence encoded by non- lymphoid cells that have not undergone the maturation process that leads to genetic rearrangement and mutation for expression of a particular immunoglobulin. (See, e.g., Shapiro et al., Crit. Rev. Immunol. 22(3): 183-200 (2002); Marchalonis et al., Adv Exp Med Biol. 484:13-30 (2001)). One of the advantages provided by various embodiments of the present invention stems from the recognition that germline antibody genes are more likely than mature antibody genes to conserve essential amino acid sequence structures characteristic of individuals in the species, hence less likely to be recognized as from a foreign source when used therapeutically in that species.

As used herein, the term "neutralizing" refers to counteracting the biological activity of an antigen when a binding protein specifically binds the antigen. In an embodiment, the neutralizing binding protein binds the cytokine and reduces its biological activity by at least about 20%, 40%, 60%, 80%, 85% or more.

The term "activity" includes activities such as the binding specificity and affinity of a DVD-Ig for two or more antigens.

The term "epitope" includes any polypeptide determinant capable of specific binding to an immunoglobulin or T-cell receptor. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and/or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody. In certain embodiments, an antibody is said to specifically bind an antigen when it recognizes its target antigen in a complex mixture of proteins and/or macromolecules. Antibodies are said to "bind to the same epitope" if the antibodies cross-compete (one prevents the binding or modulating effect of the other). In addition structural definitions of epitopes (overlapping, similar, identical) are informative, but functional definitions are often more relevant as they encompass structural (binding) and functional (modulation, competition) parameters.

The term "surface plasmon resonance", as used herein, refers to an optical phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore® system (BIAcore International AB, a GE Healthcare company, Uppsala, Sweden and Piscataway, NJ). For further descriptions, see Jönsson, U., *et al.* (1993) *Ann. Biol. Clin.* 51:19-26; Jönsson, U., *et*

al. (1991) *Biotechniques* 11:620-627; Johnsson, B., *et al.* (1995) *J. Mol. Recognit.* 8:125-131; and Johnsson, B., *et al.* (1991) *Anal. Biochem.* 198:268-277.

The term " K_{on} ", as used herein, is intended to refer to the on rate constant for association of a binding protein (e.g., an antibody) to the antigen to form the, e.g., antibody/antigen complex as is known in the art. The " K_{on} " also is known by the terms "association rate constant", or " k_a ", as used interchangeably herein. This value indicating the binding rate of an antibody to its target antigen or the rate of complex formation between an antibody and antigen also is shown by the equation below:



The term " K_{off} ", as used herein, is intended to refer to the off rate constant for dissociation, or "dissociation rate constant", of a binding protein (e.g., an antibody) from the, e.g., antibody/antigen complex as is known in the art. This value indicates the dissociation rate of an antibody from its target antigen or separation of Ab-Ag complex over time into free antibody and antigen as shown by the equation below:

$\text{Ab} + \text{Ag} \leftarrow \text{Ab-Ag}$. The term " K_D ", as used herein, is intended to refer to the "equilibrium dissociation constant", and refers to the value obtained in a titration measurement at equilibrium, or by dividing the dissociation rate constant (k_{off}) by the association rate constant (k_{on}). The association rate constant, the dissociation rate constant and the equilibrium dissociation constant are used to represent the binding affinity of an antibody to an antigen.

Methods for determining association and dissociation rate constants are well known in the art. Using fluorescence-based techniques offers high sensitivity and the ability to examine samples in physiological buffers at equilibrium. Other experimental approaches and instruments such as a BIAcore® (biomolecular interaction analysis) assay can be used (e.g., instrument available from BIAcore International AB, a GE Healthcare company, Uppsala, Sweden). Additionally, a KinExA® (Kinetic Exclusion Assay) assay, available from Sapidyne Instruments (Boise, Idaho) can also be used.

"Label" and "detectable label" mean a moiety attached to a specific binding partner, such as an antibody or an analyte, e.g., to render the reaction between members of a specific binding pair, such as an antibody and an analyte, detectable, and the specific binding partner, e.g., antibody or analyte, so labeled is referred to as "detectably labeled." Thus, the term "labeled binding protein" as used herein, refers to a protein with a label incorporated that provides for the identification of the binding protein. In an embodiment, the label is a detectable marker that can produce a signal that is detectable by visual or instrumental means, e.g., incorporation of a radiolabeled amino acid or attachment to a polypeptide of biotinyl moieties that can be detected

by marked avidin (e.g., streptavidin containing a fluorescent marker or enzymatic activity that can be detected by optical or colorimetric methods). Examples of labels for polypeptides include, but are not limited to, the following: radioisotopes or radionuclides (e.g., ^3H , ^{14}C , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , or ^{153}Sm); chromogens, fluorescent labels (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic labels (e.g., horseradish peroxidase, luciferase, alkaline phosphatase); chemiluminescent markers; biotinyl groups; predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags); and magnetic agents, such as gadolinium chelates. Representative examples of labels commonly employed for immunassays include moieties that produce light, e.g., acridinium compounds, and moieties that produce fluorescence, e.g., fluorescein. Other labels are described herein. In this regard, the moiety itself may not be detectably labeled but may become detectable upon reaction with yet another moiety. Use of "detectably labeled" is intended to encompass the latter type of detectable labeling.

The term "conjugate" refers to a binding protein, such as an antibody, chemically linked to a second chemical moiety, such as a therapeutic or cytotoxic agent. The term "agent" is used herein to denote a chemical compound, a mixture of chemical compounds, a biological macromolecule, or an extract made from biological materials. In an embodiment, the therapeutic or cytotoxic agents include, but are not limited to, pertussis toxin, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. When employed in the context of an immunoassay, the conjugate antibody may be a detectably labeled antibody used as the detection antibody.

The terms "crystal" and "crystallized" as used herein, refer to a binding protein (e.g., an antibody), or antigen binding portion thereof, that exists in the form of a crystal. Crystals are one form of the solid state of matter, which is distinct from other forms such as the amorphous solid state or the liquid crystalline state. Crystals are composed of regular, repeating, three-dimensional arrays of atoms, ions, molecules (e.g., proteins such as antibodies), or molecular assemblies (e.g., antigen/antibody complexes). These three-dimensional arrays are arranged according to specific mathematical relationships that are well-understood in the field. The fundamental unit, or building block, that is repeated in a crystal is called the asymmetric unit. Repetition of the asymmetric unit in an arrangement that conforms to a given, well-defined crystallographic symmetry provides the "unit cell" of the crystal. Repetition of the unit cell by regular translations in all three dimensions provides the crystal. See Giege, R. and Ducruix, A.

Barrett, Crystallization of Nucleic Acids and Proteins, a Practical Approach, 2nd ea., pp. 20 1-16, Oxford University Press, New York, New York, (1999)."

The term "polynucleotide" means a polymeric form of two or more nucleotides, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term
5 includes single and double stranded forms of DNA.

The term "isolated polynucleotide" shall mean a polynucleotide (e.g., of genomic, cDNA, or synthetic origin, or some combination thereof) that, by virtue of its origin, the "isolated polynucleotide" is not associated with all or a portion of a polynucleotide with which the "isolated polynucleotide" is found in nature; is operably linked to a polynucleotide that it is not linked to in
10 nature; or does not occur in nature as part of a larger sequence.

The term "vector", is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the
15 viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes
20 to which they are operatively linked. Such vectors are referred to herein as "recombinant expression vectors" (or simply, "expression vectors"). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression
25 vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding
30 sequence is achieved under conditions compatible with the control sequences. "Operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in *trans* or at a distance to control the gene of interest. The term "expression control sequence" as used herein refers to polynucleotide sequences which are necessary to effect the expression and processing of coding sequences to which they are
35 ligated. Expression control sequences include appropriate transcription initiation, termination,

promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences
5 differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include components whose presence is essential for expression and processing, and can also include additional components whose presence is
10 advantageous, for example, leader sequences and fusion partner sequences.

"Transformation", refers to any process by which exogenous DNA enters a host cell. Transformation may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on
15 the host cell being transformed and may include, but is not limited to, viral infection, electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

The term "recombinant host cell" (or simply "host cell"), is intended to refer to a cell into which exogenous DNA has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell, but, to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included
25 within the scope of the term "host cell" as used herein. In an embodiment, host cells include prokaryotic and eukaryotic cells selected from any of the Kingdoms of life. In another embodiment, eukaryotic cells include protist, fungal, plant and animal cells. In another embodiment, host cells include but are not limited to the prokaryotic cell line E.Coli; mammalian cell lines CHO, HEK 293, COS, NS0, SP2 and PER.C6; the insect cell line Sf9; and the fungal
30 cell *Saccharomyces cerevisiae*.

Standard techniques may be used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Enzymatic reactions and purification techniques may be performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. The foregoing techniques and
35 procedures may be generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed

throughout the present specification. See e.g., Sambrook et al. *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)), which is incorporated herein by reference for any purpose.

“Transgenic organism”, as known in the art, refers to an organism having cells that contain a transgene, wherein the transgene introduced into the organism (or an ancestor of the organism) expresses a polypeptide not naturally expressed in the organism. A “transgene” is a DNA construct, which is stably and operably integrated into the genome of a cell from which a transgenic organism develops, directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic organism.

The term “regulate” and “modulate” are used interchangeably, and, as used herein, refers to a change or an alteration in the activity of a molecule of interest (e.g., the biological activity of a cytokine). Modulation may be an increase or a decrease in the magnitude of a certain activity or function of the molecule of interest. Exemplary activities and functions of a molecule include, but are not limited to, binding characteristics, enzymatic activity, cell receptor activation, and signal transduction.

Correspondingly, the term “modulator” is a compound capable of changing or altering an activity or function of a molecule of interest (e.g., the biological activity of a cytokine). For example, a modulator may cause an increase or decrease in the magnitude of a certain activity or function of a molecule compared to the magnitude of the activity or function observed in the absence of the modulator. In certain embodiments, a modulator is an inhibitor, which decreases the magnitude of at least one activity or function of a molecule. Exemplary inhibitors include, but are not limited to, proteins, peptides, antibodies, peptibodies, carbohydrates or small organic molecules. Peptibodies are described, e.g., in WO01/83525.

The term “agonist”, refers to a modulator that, when contacted with a molecule of interest, causes an increase in the magnitude of a certain activity or function of the molecule compared to the magnitude of the activity or function observed in the absence of the agonist. Particular agonists of interest may include, but are not limited to, polypeptides, nucleic acids, carbohydrates, or any other molecules that bind to the antigen.

The term “antagonist” or “inhibitor”, refer to a modulator that, when contacted with a molecule of interest causes a decrease in the magnitude of a certain activity or function of the molecule compared to the magnitude of the activity or function observed in the absence of the antagonist. Particular antagonists of interest include those that block or modulate the biological or immunological activity of the antigen. Antagonists and inhibitors of antigens may include, but are not limited to, proteins, nucleic acids, carbohydrates, or any other molecules, which bind to the antigen.

As used herein, the term "effective amount" refers to the amount of a therapy which is sufficient to reduce or ameliorate the severity and/or duration of a disorder or one or more symptoms thereof, prevent the advancement of a disorder, cause regression of a disorder, prevent the recurrence, development, onset or progression of one or more symptoms associated with a disorder, detect a disorder, or enhance or improve the prophylactic or therapeutic effect(s) of another therapy (e.g., prophylactic or therapeutic agent).

"Patient" and "subject" may be used interchangeably herein to refer to an animal, such as a mammal, including a primate (for example, a human, a monkey, and a chimpanzee), a non-primate (for example, a cow, a pig, a camel, a llama, a horse, a goat, a rabbit, a sheep, a hamster, a guinea pig, a cat, a dog, a rat, a mouse, a whale), a bird (e.g., a duck or a goose), and a shark. Preferably, the patient or subject is a human, such as a human being treated or assessed for a disease, disorder or condition, a human at risk for a disease, disorder or condition, a human having a disease, disorder or condition, and/or human being treated for a disease, disorder or condition.

The term "sample", as used herein, is used in its broadest sense. A "biological sample", as used herein, includes, but is not limited to, any quantity of a substance from a living thing or formerly living thing. Such living things include, but are not limited to, humans, mice, rats, monkeys, dogs, rabbits and other animals. Such substances include, but are not limited to, blood, (e.g., whole blood), plasma, serum, urine, amniotic fluid, synovial fluid, endothelial cells, leukocytes, monocytes, other cells, organs, tissues, bone marrow, lymph nodes and spleen.

"Component," "components," and "at least one component," refer generally to a capture antibody, a detection or conjugate antibody, a control, a calibrator, a series of calibrators, a sensitivity panel, a container, a buffer, a diluent, a salt, an enzyme, a co-factor for an enzyme, a detection reagent, a pretreatment reagent/solution, a substrate (e.g., as a solution), a stop solution, and the like that can be included in a kit for assay of a test sample, such as a patient urine, serum or plasma sample, in accordance with the methods described herein and other methods known in the art. Thus, in the context of the present disclosure, "at least one component," "component," and "components" can include a polypeptide or other analyte as above, such as a composition comprising an analyte such as polypeptide, which is optionally immobilized on a solid support, such as by binding to an anti-analyte (e.g., anti-polypeptide) antibody. Some components can be in solution or lyophilized for reconstitution for use in an assay.

"Control" refers to a composition known to not analyte ("negative control") or to contain analyte ("positive control"). A positive control can comprise a known concentration of analyte. "Control," "positive control," and "calibrator" may be used interchangeably herein to refer to a composition comprising a known concentration of analyte. A "positive control" can be used to

establish assay performance characteristics and is a useful indicator of the integrity of reagents (e.g., analytes).

“Predetermined cutoff” and “predetermined level” refer generally to an assay cutoff value that is used to assess diagnostic/prognostic/therapeutic efficacy results by comparing the assay results against the predetermined cutoff/level, where the predetermined cutoff/level already has been linked or associated with various clinical parameters (e.g., severity of disease, progression/nonprogression/improvement, etc.). While the present disclosure may provide exemplary predetermined levels, it is well-known that cutoff values may vary depending on the nature of the immunoassay (e.g., antibodies employed, etc.). It further is well within the ordinary skill of one in the art to adapt the disclosure herein for other immunoassays to obtain immunoassay-specific cutoff values for those other immunoassays based on this disclosure. Whereas the precise value of the predetermined cutoff/level may vary between assays, correlations as described herein (if any) should be generally applicable.

“Pretreatment reagent,” e.g., lysis, precipitation and/or solubilization reagent, as used in a diagnostic assay as described herein is one that lyses any cells and/or solubilizes any analyte that is/are present in a test sample. Pretreatment is not necessary for all samples, as described further herein. Among other things, solubilizing the analyte (e.g., polypeptide of interest) may entail release of the analyte from any endogenous binding proteins present in the sample. A pretreatment reagent may be homogeneous (not requiring a separation step) or heterogeneous (requiring a separation step). With use of a heterogeneous pretreatment reagent there is removal of any precipitated analyte binding proteins from the test sample prior to proceeding to the next step of the assay.

“Quality control reagents” in the context of immunoassays and kits described herein, include, but are not limited to, calibrators, controls, and sensitivity panels. A “calibrator” or “standard” typically is used (e.g., one or more, such as a plurality) in order to establish calibration (standard) curves for interpolation of the concentration of an analyte, such as an antibody or an analyte. Alternatively, a single calibrator, which is near a predetermined positive/negative cutoff, can be used. Multiple calibrators (i.e., more than one calibrator or a varying amount of calibrator(s)) can be used in conjunction so as to comprise a “sensitivity panel.”

“Risk” refers to the possibility or probability of a particular event occurring either presently or at some point in the future. “Risk stratification” refers to an array of known clinical risk factors that allows physicians to classify patients into a low, moderate, high or highest risk of developing a particular disease, disorder or condition.

“Specific” and “specificity” in the context of an interaction between members of a specific binding pair (e.g., an antigen (or fragment thereof) and an antibody (or antigenically

reactive fragment thereof)) refer to the selective reactivity of the interaction. The phrase “specifically binds to” and analogous phrases refer to the ability of antibodies (or antigenically reactive fragments thereof) to bind specifically to analyte (or a fragment thereof) and not bind specifically to other entities.

5 “Specific binding partner” is a member of a specific binding pair. A specific binding pair comprises two different molecules, which specifically bind to each other through chemical or physical means. Therefore, in addition to antigen and antibody specific binding pairs of common immunoassays, other specific binding pairs can include biotin and avidin (or streptavidin), carbohydrates and lectins, complementary nucleotide sequences, effector and receptor molecules, 10 cofactors and enzymes, enzyme inhibitors and enzymes, and the like. Furthermore, specific binding pairs can include members that are analogs of the original specific binding members, for example, an analyte-analog. Immunoreactive specific binding members include antigens, antigen fragments, and antibodies, including monoclonal and polyclonal antibodies as well as complexes, fragments, and variants (including fragments of variants) thereof, whether isolated or 15 recombinantly produced.

“Variant” as used herein means a polypeptide that differs from a given polypeptide (e.g., IL-18, BNP, NGAL or HIV polypeptide or anti-polypeptide antibody) in amino acid sequence by the addition (e.g., insertion), deletion, or conservative substitution of amino acids, but that retains the biological activity of the given polypeptide (e.g., a variant IL-18 can compete with anti-IL-18 20 antibody for binding to IL-18). A conservative substitution of an amino acid, i.e., replacing an amino acid with a different amino acid of similar properties (e.g., hydrophilicity and degree and distribution of charged regions) is recognized in the art as typically involving a minor change. These minor changes can be identified, in part, by considering the hydropathic index of amino acids, as understood in the art (see, e.g., Kyte et al., J. Mol. Biol. 157: 105-132 (1982)). The 25 hydropathic index of an amino acid is based on a consideration of its hydrophobicity and charge. It is known in the art that amino acids of similar hydropathic indexes can be substituted and still retain protein function. In one aspect, amino acids having hydropathic indexes of ± 2 are substituted. The hydrophilicity of amino acids also can be used to reveal substitutions that would result in proteins retaining biological function. A consideration of the hydrophilicity of amino 30 acids in the context of a peptide permits calculation of the greatest local average hydrophilicity of that peptide, a useful measure that has been reported to correlate well with antigenicity and immunogenicity (see, e.g., U.S. Pat. No. 4,554,101, which is incorporated herein by reference). Substitution of amino acids having similar hydrophilicity values can result in peptides retaining biological activity, for example immunogenicity, as is understood in the art. In one aspect, 35 substitutions are performed with amino acids having hydrophilicity values within ± 2 of each other. Both the hydrophobicity index and the hydrophilicity value of amino acids are influenced

by the particular side chain of that amino acid. Consistent with that observation, amino acid substitutions that are compatible with biological function are understood to depend on the relative similarity of the amino acids, and particularly the side chains of those amino acids, as revealed by the hydrophobicity, hydrophilicity, charge, size, and other properties. "Variant" also can be used to describe a polypeptide or fragment thereof that has been differentially processed, such as by proteolysis, phosphorylation, or other post-translational modification, yet retains its biological activity or antigen reactivity, e.g., the ability to bind to IL-18. Use of "variant" herein is intended to encompass fragments of a variant unless otherwise contradicted by context.

I. Generation of DVD binding protein

The invention pertains to Dual Variable Domain binding proteins capable of binding one or more targets and methods of making the same. In an embodiment, the binding protein comprises a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein VD1 is a first variable domain, VD2 is a second variable domain, C is a constant domain, X1 represents an amino acid or polypeptide, X2 represents an Fc region and n is 0 or 1.

The binding protein of the invention can be generated using various techniques. The invention provides expression vectors, host cell and methods of generating the binding protein.

A. Generation of parent monoclonal antibodies

The variable domains of the DVD binding protein can be obtained from parent antibodies, including polyclonal and mAbs capable of binding antigens of interest. These antibodies may be naturally occurring or may be generated by recombinant technology.

MAbs can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, mAbs can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced. Hybridomas are selected, cloned and further screened for desirable characteristics, including robust hybridoma growth, high antibody production and desirable antibody characteristics, as discussed in Example 1 below. Hybridomas may be cultured and expanded *in vivo* in syngeneic animals, in animals that lack an immune system, e.g., nude mice, or in cell culture *in vitro*. Methods of selecting, cloning and expanding hybridomas are well known to those of ordinary skill in the art. In a particular embodiment, the hybridomas are mouse

hybridomas. In another embodiment, the hybridomas are produced in a non-human, non-mouse species such as rats, sheep, pigs, goats, cattle or horses. In another embodiment, the hybridomas are human hybridomas, in which a human non-secretory myeloma is fused with a human cell expressing an antibody capable of binding a specific antigen.

5 Recombinant mAbs are also generated from single, isolated lymphocytes using a procedure referred to in the art as the selected lymphocyte antibody method (SLAM), as described in U.S. Patent No. 5,627,052, PCT Publication WO 92/02551 and Babcock, J.S. *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93:7843-7848. In this method, single cells secreting antibodies of interest, *e.g.*, lymphocytes derived from an immunized animal, are identified, and, heavy- and
10 light-chain variable region cDNAs are rescued from the cells by reverse transcriptase-PCR and these variable regions can then be expressed, in the context of appropriate immunoglobulin constant regions (*e.g.*, human constant regions), in mammalian host cells, such as COS or CHO cells. The host cells transfected with the amplified immunoglobulin sequences, derived from *in vivo* selected lymphocytes, can then undergo further analysis and selection *in vitro*, for example
15 by panning the transfected cells to isolate cells expressing antibodies to the antigen of interest. The amplified immunoglobulin sequences further can be manipulated *in vitro*, such as by *in vitro* affinity maturation methods such as those described in PCT Publication WO 97/29131 and PCT Publication WO 00/56772.

 Monoclonal antibodies are also produced by immunizing a non-human animal
20 comprising some, or all, of the human immunoglobulin locus with an antigen of interest. In an embodiment, the non-human animal is a XENOMOUSE transgenic mouse, an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production. *See, e.g.*, Green et al. *Nature Genetics* 7:13-21 (1994) and United States Patents Nos. 5,916,771, 5,939,598, 5,985,615, 5,998,209, 6,075,181, 6,091,001,
25 6,114,598 and 6,130,364. See also WO 91/10741, published July 25, 1991, WO 94/02602, published February 3, 1994, WO 96/34096 and WO 96/33735, both published October 31, 1996, WO 98/16654, published April 23, 1998, WO 98/24893, published June 11, 1998, WO 98/50433, published November 12, 1998, WO 99/45031, published September 10, 1999, WO 99/53049, published October 21, 1999, WO 00 09560, published February 24, 2000 and WO
30 00/037504, published June 29, 2000. The XENOMOUSE transgenic mouse produces an adult-like human repertoire of fully human antibodies, and generates antigen-specific human monoclonal antibodies. The XENOMOUSE transgenic mouse contains approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and x light chain loci. See Mendez et al., *Nature Genetics* 15:146-156 (1997), Green and Jakobovits *J. Exp. Med.* 188:483-495 (1998), the
35 disclosures of which are hereby incorporated by reference.

In vitro methods also can be used to make the parent antibodies, wherein an antibody library is screened to identify an antibody having the desired binding specificity. Methods for such screening of recombinant antibody libraries are well known in the art and include methods described in, for example, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* PCT Publication No. WO 92/18619; Dower *et al.* PCT Publication No. WO 91/17271; Winter *et al.* PCT Publication No. WO 92/20791; Markland *et al.* PCT Publication No. WO 92/15679; Breitling *et al.* PCT Publication No. WO 93/01288; McCafferty *et al.* PCT Publication No. WO 92/01047; Garrard *et al.* PCT Publication No. WO 92/09690; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; McCafferty *et al.*, *Nature* (1990) 348:552-554; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrard *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982, US patent application publication 20030186374, and PCT Publication No. WO 97/29131, the contents of each of which are incorporated herein by reference.

Parent antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular, such phage can be utilized to display antigen-binding domains expressed from a repertoire or combinatorial antibody library (e. g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman *et al.*, *J. Immunol. Methods* 182:41-50 (1995); Ames *et al.*, *J. Immunol. Methods* 184:177-186 (1995); Kettleborough *et al.*, *Eur. J. Immunol.* 24:952-958 (1994); Persic *et al.*, *Gene* 187 9-18 (1997); Burton *et al.*, *Advances in Immunology* 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Pat. Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the herein references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies including human antibodies

or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., *BioTechniques* 12(6):864-869 (1992); and Sawai et al., *AJRI* 34:26-34 (1995); and Better et al., *Science* 240:1041-1043 (1988) (said references incorporated by reference in their entireties). Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Pat. 4,946,778 and 5,258,498; Huston et al., *Methods in Enzymology* 203:46-88 (1991); Shu et al., *PNAS* 90:7995-7999 (1993); and Skerra et al., *Science* 240:1038-1040 (1988).

Alternative to screening of recombinant antibody libraries by phage display, other methodologies known in the art for screening large combinatorial libraries can be applied to the identification of parent antibodies. One type of alternative expression system is one in which the recombinant antibody library is expressed as RNA-protein fusions, as described in PCT Publication No. WO 98/31700 by Szostak and Roberts, and in Roberts, R.W. and Szostak, J.W. (1997) *Proc. Natl. Acad. Sci. USA* 94:12297-12302. In this system, a covalent fusion is created between an mRNA and the peptide or protein that it encodes by *in vitro* translation of synthetic mRNAs that carry puromycin, a peptidyl acceptor antibiotic, at their 3' end. Thus, a specific mRNA can be enriched from a complex mixture of mRNAs (e.g., a combinatorial library) based on the properties of the encoded peptide or protein, e.g., antibody, or portion thereof, such as binding of the antibody, or portion thereof, to the dual specificity antigen. Nucleic acid sequences encoding antibodies, or portions thereof, recovered from screening of such libraries can be expressed by recombinant means as described herein (e.g., in mammalian host cells) and, moreover, can be subjected to further affinity maturation by either additional rounds of screening of mRNA-peptide fusions in which mutations have been introduced into the originally selected sequence(s), or by other methods for affinity maturation *in vitro* of recombinant antibodies, as described herein.

In another approach the parent antibodies can also be generated using yeast display methods known in the art. In yeast display methods, genetic methods are used to tether antibody domains to the yeast cell wall and display them on the surface of yeast. In particular, such yeast can be utilized to display antigen-binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Examples of yeast display methods that can be used to make the parent antibodies include those disclosed in Wittrup, et al. U.S. Patent No. 6,699,658 incorporated herein by reference.

The antibodies described herein can be further modified to generate CDR grafted and humanized parent antibodies. CDR-grafted parent antibodies comprise heavy and light chain

variable region sequences from a human antibody wherein one or more of the CDR regions of V_H and/or V_L are replaced with CDR sequences of murine antibodies capable of binding antigen of interest. A framework sequence from any human antibody may serve as the template for CDR grafting. However, straight chain replacement onto such a framework often leads to some loss of binding affinity to the antigen. The more homologous a human antibody is to the original murine antibody, the less likely the possibility that combining the murine CDRs with the human framework will introduce distortions in the CDRs that could reduce affinity. Therefore, in an embodiment, the human variable framework that is chosen to replace the murine variable framework apart from the CDRs have at least a 65% sequence identity with the murine antibody variable region framework. In an embodiment, the human and murine variable regions apart from the CDRs have at least 70% sequence identity. In a particular embodiment, that the human and murine variable regions apart from the CDRs have at least 75% sequence identity. In another embodiment, the human and murine variable regions apart from the CDRs have at least 80% sequence identity. Methods for producing such antibodies are known in the art (see EP 239,400; PCT publication WO 91/09967; U.S. Pat. Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Pat. No. 5,565,352); and anti-idiotypic antibodies.

Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and framework regions from a human immunoglobulin molecule. Known human Ig sequences are disclosed, e.g., www.ncbi.nlm.nih.gov/entrez-/query.fcgi; www.atcc.org/phage/hdb.html; www.sciquest.com/; www.abcam.com/; www.antibodyresource.com/onlinecomp.html; www.public.iastate.edu/.about.pedro/research_tools.html; www.mgen.uni-heidelberg.de/SD/IT/IT.html; www.whfreeman.com/immunology/CH-05/kuby05.htm; www.library.thinkquest.org/12429/Immune/Antibody.html; www.hhmi.org/grants/lectures/1996/vlab/; www.path.cam.ac.uk/.about.mrc7/m-ikeimages.html; www.antibodyresource.com/; mcb.harvard.edu/BioLinks/Immunology.html; www.immunologylink.com/; pathbox.wustl.edu/.about.hcenter/index.-html; www.biotech.ufl.edu/.about.hcl/; www.pebio.com/pa/340913/340913.html; www.nal.usda.gov/awic/pubs/antibody/; www.m.ehime-u.ac.jp/.about.yasuhito-/Elisa.html; www.biodesign.com/table.asp; www.icnet.uk/axp/facs/davies/links.html; www.biotech.ufl.edu/.about.fccl/protocol.html; www.isac-net.org/sites_geo.html; aximtl.imt.uni-marburg.de/.about.rek/AEP-Start.html; baserv.uci.kun.nl/.about.jraats/links1.html; www.recab.uni-hd.de/immuno.bme.nwu.edu/; www.mrc-cpe.cam.ac.uk/imt-doc/pu-

blic/INTRO.html; www.ibt.unam.mx/vir/V_mice.html; imgt.cnusc.fr:8104/;
 www.biochem.ucl.ac.uk/.about.martin/abs/index.html; antibody.bath.ac.uk/;
 abgen.cvm.tamu.edu/lab/wwwabgen.html; www.unizh.ch/.about.honegger/AHOsem-
 inar/Slide01.html; www.cryst.bbk.ac.uk/.about.ubcg07s/;
 5 www.nimr.mrc.ac.uk/CC/ccaewg/ccaewg.htm; www.path.cam.ac.uk/.about.mrc7/h-
 umanisation/TAHHP.html; www.ibt.unam.mx/vir/structure/stat_aim.html;
 www.biosci.missouri.edu/smithgp/index.html; www.cryst.bioc.cam.ac.uk/.abo- ut.fmolina/Web-
 pages/Pept/spottech.html; www.jerini.de/fr oducts.htm; www.patents.ibm.com/ibm.html.Kabat et
 al., Sequences of Proteins of Immunological Interest, U.S. Dept. Health (1983), each entirely
 10 incorporated herein by reference. Such imported sequences can be used to reduce immunogenicity
 or reduce, enhance or modify binding, affinity, on-rate, off-rate, avidity, specificity, half-life, or
 any other suitable characteristic, as known in the art.

Framework residues in the human framework regions may be substituted with the
 corresponding residue from the CDR donor antibody to alter, e.g., improve, antigen binding.
 15 These framework substitutions are identified by methods well known in the art, e.g., by modeling
 of the interactions of the CDR and framework residues to identify framework residues important
 for antigen binding and sequence comparison to identify unusual framework residues at particular
 positions. (See, e.g., Queen et al., U.S. Pat. No. 5,585,089; Riechmann et al., Nature 332:323
 (1988), which are incorporated herein by reference in their entireties.) Three-dimensional
 20 immunoglobulin models are commonly available and are familiar to those skilled in the art.
 Computer programs are available which illustrate and display probable three-dimensional
 conformational structures of selected candidate immunoglobulin sequences. Inspection of these
 displays permits analysis of the likely role of the residues in the functioning of the candidate
 immunoglobulin sequence, i.e., the analysis of residues that influence the ability of the candidate
 25 immunoglobulin to bind its antigen. In this way, FR residues can be selected and combined from
 the consensus and import sequences so that the desired antibody characteristic, such as increased
 affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most
 substantially involved in influencing antigen binding. Antibodies can be humanized using a
 variety of techniques known in the art, such as but not limited to those described in Jones et al.,
 30 Nature 321:522 (1986); Verhoeyen et al., Science 239:1534 (1988)), Sims et al., J. Immunol. 151:
 2296 (1993); Chothia and Lesk, J. Mol. Biol. 196:901 (1987), Carter et al., Proc. Natl. Acad. Sci.
 U.S.A. 89:4285 (1992); Presta et al., J. Immunol. 151:2623 (1993), Padlan, Molecular
 Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994);
 Roguska. et al. , PNAS 91:969-973 (1994); PCT publication WO 91/09967, PCT/: US98/16280,
 35 US96/18978, US91/09630, US91/05939, US94/01234, GB89/01334, GB91/01134, GB92/01755;
 WO90/14443, WO90/14424, WO90/14430, EP 229246, EP 592,106; EP 519,596, EP 239,400,

U.S. Pat. Nos. 5,565,332, 5,723,323, 5,976,862, 5,824,514, 5,817,483, 5,814,476, 5,763,192, 5,723,323, 5,766,886, 5,714,352, 6,204,023, 6,180,370, 5,693,762, 5,530,101, 5,585,089, 5,225,539; 4,816,567, each entirely incorporated herein by reference, included references cited therein.

5 **B. Criteria for selecting parent monoclonal antibodies**

An embodiment of the invention pertains to selecting parent antibodies with at least one or more properties desired in the DVD-Ig molecule. In an embodiment, the desired property is selected from one or more antibody parameters. In another embodiment, the antibody parameters are selected from the group consisting of antigen specificity, affinity to antigen, potency,
10 biological function, epitope recognition, stability, solubility, production efficiency, immunogenicity, pharmacokinetics, bioavailability, tissue cross reactivity, and orthologous antigen binding.

B1. Affinity to Antigen

The desired affinity of a therapeutic mAb may depend upon the nature of the antigen, and
15 the desired therapeutic end-point. In an embodiment, monoclonal antibodies have higher affinities ($K_d = 0.01 - 0.50$ pM) when blocking a cytokine-cytokine receptor interaction as such interaction are usually high affinity interactions (e.g., <pM – <nM ranges). In such instances, the mAb affinity for its target should be equal to or better than the affinity of the cytokine (ligand) for its receptor. On the other hand, mAb with lesser affinity (> nM range) could be therapeutically
20 effective e.g., in clearing circulating potentially pathogenic proteins e.g., monoclonal antibodies that bind to, sequester, and clear circulating species of A- β amyloid. In other instances, reducing the affinity of an existing high affinity mAb by site-directed mutagenesis or using a mAb with lower affinity for its target could be used to avoid potential side-effects e.g., a high affinity mAb may sequester/neutralize all of its intended target, thereby completely depleting/eliminating the
25 function(s) of the targeted protein. In this scenario, a low affinity mAb may sequester/neutralize a fraction of the target that may be responsible for the disease symptoms (the pathological or over-produced levels), thus allowing a fraction of the target to continue to perform its normal physiological function(s). Therefore, it may be possible to reduce the K_d to adjust dose and/or reduce side-effects. The affinity of the parental mAb might play a role in appropriately targeting
30 cell surface molecules to achieve desired therapeutic out-come. For example, if a target is expressed on cancer cells with high density and on normal cells with low density, a lower affinity mAb will bind a greater number of targets on tumor cells than normal cells, resulting in tumor cell elimination via ADCC or CDC, and therefore might have therapeutically desirable effects. Thus selecting a mAb with desired affinity may be relevant for both soluble and surface targets.

Signaling through a receptor upon interaction with its ligand may depend upon the affinity of the receptor-ligand interaction. Similarly, it is conceivable that the affinity of a mAb for a surface receptor could determine the nature of intracellular signaling and whether the mAb may deliver an agonist or an antagonist signal. The affinity-based nature of mAb-mediated
 5 signaling may have an impact of its side-effect profile. Therefore, the desired affinity and desired functions of therapeutic monoclonal antibodies need to be determined carefully by in vitro and in vivo experimentation.

The desired K_d of a binding protein (e.g., an antibody) may be determined experimentally depending on the desired therapeutic outcome. In an embodiment parent antibodies with affinity
 10 (K_d) for a particular antigen equal to, or better than, the desired affinity of the DVD-Ig for the same antigen are selected. The antigen binding affinity and kinetics are assessed by Biacore or another similar technique. In one embodiment, each parent antibody has a dissociation constant (K_d) to its antigen selected from the group consisting of: at most about 10^{-7} M; at most about 10^{-8} M; at most about 10^{-9} M; at most about 10^{-10} M; at most about 10^{-11} M; at most about 10^{-12} M; and
 15 at most 10^{-13} M. First parent antibody from which VD1 is obtained and second parent antibody from which VD2 is obtained may have similar or different affinity (K_D) for the respective antigen. Each parent antibody has an on rate constant (K_{on}) to the antigen selected from the group consisting of: at least about $10^2 M^{-1} s^{-1}$; at least about $10^3 M^{-1} s^{-1}$; at least about $10^4 M^{-1} s^{-1}$; at least
 20 about $10^5 M^{-1} s^{-1}$; and at least about $10^6 M^{-1} s^{-1}$, as measured by surface plasmon resonance. The first parent antibody from which VD1 is obtained and the second parent antibody from which VD2 is obtained may have similar or different on rate constant (K_{on}) for the respective antigen. In one embodiment, each parent antibody has an off rate constant (K_{off}) to the antigen selected from the group consisting of: at most about $10^{-3} s^{-1}$; at most about $10^{-4} s^{-1}$; at most about $10^{-5} s^{-1}$; and at most
 25 about $10^{-6} s^{-1}$, as measured by surface plasmon resonance. The first parent antibody from which VD1 is obtained and the second parent antibody from which VD2 is obtained may have similar or different off rate constants (K_{off}) for the respective antigen.

B2. Potency

The desired affinity/potency of parental monoclonal antibodies will depend on the desired therapeutic outcome. For example, for receptor-ligand (R-L) interactions the affinity (k_d) is equal
 30 to or better than the R-L k_d (pM range). For simple clearance of a pathologic circulating protein, the k_d could be in low nM range e.g., clearance of various species of circulating A- β peptide. In addition, the k_d will also depend on whether the target expresses multiple copies of the same epitope e.g a mAb targeting conformational epitope in A β oligomers.

Where VDI and VD2 bind the same antigen, but distinct epitopes, the DVD-Ig will contain
 35 4 binding sites for the same antigen, thus increasing avidity and thereby the apparent k_d of the

DVD-Ig. In an embodiment, parent antibodies with equal or lower k_d than that desired in the DVD-Ig are chosen. The affinity considerations of a parental mAb may also depend upon whether the DVD-Ig contains four or more identical antigen binding sites (i.e; a DVD-Ig from a single mAb). In this case, the apparent k_d would be greater than the mAb due to avidity. Such
 5 DVD-Igs can be employed for cross-linking surface receptor, increase neutralization potency, enhance clearance of pathological proteins etc.

In an embodiment parent antibodies with neutralization potency for specific antigen equal to or better than the desired neutralization potential of the DVD-Ig for the same antigen are selected. The neutralization potency can be assessed by a target-dependent bioassay where cells
 10 of appropriate type produce a measurable signal (i.e. proliferation or cytokine production) in response to target stimulation, and target neutralization by the mAb can reduce the signal in a dose-dependent manner.

B3. Biological functions

Monoclonal antibodies can perform potentially several functions. Some of these functions
 15 are listed in Table 3. These functions can be assessed by both in vitro assays (e.g., cell-based and biochemical assays) and in vivo animal models.

Table 3: Some Potential Applications For Therapeutic Antibodies

Target (Class)	Mechanism of Action (target)
Soluble (cytokines, other)	Neutralization of activity (e.g., a cytokine) Enhance clearance (e.g., A β oligomers) Increase half-life (e.g., GLP 1)
Cell Surface (Receptors, other)	Agonist (e.g., GLP1 R; EPO R; etc.) Antagonist (e.g., integrins; etc.) Cytotoxic (CD 20; etc.)
Protein deposits	Enhance clearance/degradation (e.g., A β plaques, amyloid deposits)

MAbs with distinct functions described in the examples herein in Table 3 can be selected to achieve desired therapeutic outcomes. Two or more selected parent monoclonal antibodies can
 20 then be used in DVD-Ig format to achieve two distinct functions in a single DVD-Ig molecule. For example, a DVD-Ig can be generated by selecting a parent mAb that neutralizes function of a specific cytokine, and selecting a parent mAb that enhances clearance of a pathological protein. Similarly, we can select two parent monoclonal antibodies that recognize two different cell surface receptors, one mAb with an agonist function on one receptor and the other mAb with an
 25 antagonist function on a different receptor. These two selected monoclonal antibodies each with a distinct function can be used to construct a single DVD-Ig molecule that will possess the two

distinct functions (agonist and antagonist) of the selected monoclonal antibodies in a single molecule. Similarly, two antagonistic monoclonal antibodies to cell surface receptors each blocking binding of respective receptor ligands (e.g., EGF and IGF) can be used in a DVD-Ig format. Conversely, an antagonistic anti-receptor mAb (e.g., anti-EGFR) and a neutralizing anti-soluble mediator (e.g., anti-IGF1/2) mAb can be selected to make a DVD-Ig.

B4. Epitope Recognition:

Different regions of proteins may perform different functions. For example specific regions of a cytokine interact with the cytokine receptor to bring about receptor activation whereas other regions of the protein may be required for stabilizing the cytokine. In this instance one may select a mAb that binds specifically to the receptor interacting region(s) on the cytokine and thereby block cytokine-receptor interaction. In some cases, for example certain chemokine receptors that bind multiple ligands, a mAb that binds to the epitope (region on chemokine receptor) that interacts with only one ligand can be selected. In other instances, monoclonal antibodies can bind to epitopes on a target that are not directly responsible for physiological functions of the protein, but binding of a mAb to these regions could either interfere with physiological functions (steric hindrance) or alter the conformation of the protein such that the protein cannot function (mAb to receptors with multiple ligand which alter the receptor conformation such that none of the ligand can bind). Anti-cytokine monoclonal antibodies that do not block binding of the cytokine to its receptor, but block signal transduction have also been identified (e.g., 125-2H, an anti-IL-18 mAb).

Examples of epitopes and mAb functions include, but are not limited to, blocking Receptor-Ligand (R-L) interaction (neutralizing mAb that binds R-interacting site); steric hindrance resulting in diminished or no R-binding. An Ab can bind the target at a site other than a receptor binding site, but still interferes with receptor binding and functions of the target by inducing conformational change and eliminate function (e.g., Xolair), binding to R but block signaling (125-2H).

In an embodiment, the parental mAb needs to target the appropriate epitope for maximum efficacy. Such epitope should be conserved in the DVD-Ig. The binding epitope of a mAb can be determined by several approaches, including co-crystallography, limited proteolysis of mAb-antigen complex plus mass spectrometric peptide mapping (Legros V. et al 2000 Protein Sci. 9:1002-10), phage displayed peptide libraries (O'Connor KH et al 2005 J Immunol Methods. 299:21-35), as well as mutagenesis (Wu C. et al . 2003 J Immunol 170:5571-7).

B5. Physicochemical and pharmaceutical properties:

Therapeutic treatment with antibodies often requires administration of high doses, often several mg/kg (due to a low potency on a mass basis as a consequence of a typically large

molecular weight). In order to accommodate patient compliance and to adequately address chronic disease therapies and outpatient treatment, subcutaneous (s.c.) or intramuscular (i.m.) administration of therapeutic mAbs is desirable. For example, the maximum desirable volume for s.c. administration is ~1.0 mL, and therefore, concentrations of >100 mg/mL are desirable to limit the number of injections per dose. In an embodiment, the therapeutic antibody is administered in one dose. The development of such formulations is constrained, however, by protein-protein interactions (e.g., aggregation, which potentially increases immunogenicity risks) and by limitations during processing and delivery (e.g., viscosity). Consequently, the large quantities required for clinical efficacy and the associated development constraints limit full exploitation of the potential of antibody formulation and s.c. administration in high-dose regimens. It is apparent that the physicochemical and pharmaceutical properties of a protein molecule and the protein solution are of utmost importance, e.g., stability, solubility and viscosity features.

B5.1. Stability:

A "stable" antibody formulation is one in which the antibody therein essentially retains its physical stability and/or chemical stability and/or biological activity upon storage. Stability can be measured at a selected temperature for a selected time period. In an embodiment, the antibody in the formulation is stable at room temperature (about 30°C) or at 40°C for at least 1 month and/or stable at about 2-8°C. for at least 1 year for at least 2 years. Furthermore, in an embodiment, the formulation is stable following freezing (to, e.g., -70°C) and thawing of the formulation, hereinafter referred to as a "freeze/thaw cycle." In another example, a "stable" formulation may be one wherein less than about 10% and less than about 5% of the protein is present as an aggregate in the formulation.

A DVD-Ig stable in vitro at various temperatures for an extended time period is desirable. One can achieve this by rapid screening of parental mAbs stable in vitro at elevated temperature, e.g., at 40°C for 2-4 weeks, and then assess stability. During storage at 2-8°C, the protein reveals stability for at least 12 months, e.g., at least 24 months. Stability (% of monomeric, intact molecule) can be assessed using various techniques such as cation exchange chromatography, size exclusion chromatography, SDS-PAGE, as well as bioactivity testing. For a more comprehensive list of analytical techniques that may be employed to analyze covalent and conformational modifications please see Jones, A. J. S. (1993) Analytical methods for the assessment of protein formulations and delivery systems. In: Cleland, J. L.; Langer, R., editors. Formulation and delivery of peptides and proteins, 1st edition, Washington, ACS, pg. 22-45; and Pearlman, R.; Nguyen, T. H. (1990) Analysis of protein drugs. In: Lee, V. H., editor. Peptide and protein drug delivery, 1st edition, New York, Marcel Dekker, Inc., pg. 247-301.

Heterogeneity and aggregate formation: stability of the antibody may be such that the formulation may reveal less than about 10%, and, in an embodiment, less than about 5%, in another embodiment, less than about 2%, or, in an embodiment, within the range of 0.5% to 1.5% or less in the GMP antibody material that is present as aggregate. Size exclusion chromatography is a method that is sensitive, reproducible, and very robust in the detection of protein aggregates.

In addition to low aggregate levels, the antibody must, in an embodiment, be chemically stable. Chemical stability may be determined by ion exchange chromatography (e.g., cation or anion exchange chromatography), hydrophobic interaction chromatography, or other methods such as isoelectric focusing or capillary electrophoresis. For instance, chemical stability of the antibody may be such that after storage of at least 12 months at 2-8°C the peak representing unmodified antibody in a cation exchange chromatography may increase not more than 20%, in an embodiment, not more than 10%, or, in another embodiment, not more than 5% as compared to the antibody solution prior to storage testing.

In an embodiment, the parent antibodies display structural integrity; correct disulfide bond formation, and correct folding: Chemical instability due to changes in secondary or tertiary structure of an antibody may impact antibody activity. For instance, stability as indicated by activity of the antibody may be such that after storage of at least 12 months at 2-8°C the activity of the antibody may decrease not more than 50%, in an embodiment not more than 30%, or even not more than 10%, or in an embodiment not more than 5% or 1% as compared to the antibody solution prior to storage testing. Suitable antigen-binding assays can be employed to determine antibody activity.

B5.2. Solubility:

The "solubility" of a mAb correlates with the production of correctly folded, monomeric IgG. The solubility of the IgG may therefore be assessed by HPLC. For example, soluble (monomeric) IgG will give rise to a single peak on the HPLC chromatograph, whereas insoluble (e.g., multimeric and aggregated) will give rise to a plurality of peaks. A person skilled in the art will therefore be able to detect an increase or decrease in solubility of an IgG using routine HPLC techniques. For a more comprehensive list of analytical techniques that may be employed to analyze solubility (see Jones, A. G. *Dep. Chem. Biochem. Eng., Univ. Coll. London, London, UK*. Editor(s): Shamlou, P. Ayazi. *Process. Solid-Liq. Suspensions* (1993), 93-117. Publisher: Butterworth-Heinemann, Oxford, UK and Pearlman, Rodney; Nguyen, Tue H, *Advances in Parenteral Sciences* (1990), 4 (Pept. Protein Drug Delivery), 247-301). Solubility of a therapeutic mAb is critical for formulating to high concentration often required for adequate dosing. As outlined herein, solubilities of >100 mg/mL may be required to accommodate efficient antibody dosing. For instance, antibody solubility may be not less than about 5 mg/mL in early research

phase, in an embodiment not less than about 25 mg/mL in advanced process science stages, or in an embodiment not less than about 100 mg/mL, or in an embodiment not less than about 150 mg/mL. It is obvious to a person skilled in the art that the intrinsic properties of a protein molecule are important the physico-chemical properties of the protein solution, e.g., stability, solubility, viscosity. However, a person skilled in the art will appreciate that a broad variety of excipients exist that may be used as additives to beneficially impact the characteristics of the final protein formulation. These excipients may include: (i) liquid solvents, cosolvents (e.g., alcohols such as ethanol); (ii) buffering agents (e.g., phosphate, acetate, citrate, amino acid buffers); (iii) sugars or sugar alcohols (e.g., sucrose, trehalose, fructose, raffinose, mannitol, sorbitol, dextrans); (iv) surfactants (e.g., polysorbate 20, 40, 60, 80, poloxamers); (v) isotonicity modifiers (e.g., salts such as NaCl, sugars, sugar alcohols); and (vi) others (e.g., preservatives, chelating agents, antioxidants, chelating substances (e.g., EDTA), biodegradable polymers, carrier molecules (e.g., HSA, PEGs)

Viscosity is a parameter of high importance with regard to antibody manufacture and antibody processing (e.g., diafiltration/ultrafiltration), fill-finish processes (pumping aspects, filtration aspects) and delivery aspects (syringeability, sophisticated device delivery). Low viscosities enable the liquid solution of the antibody having a higher concentration. This enables the same dose may be administered in smaller volumes. Small injection volumes inhere the advantage of lower pain on injection sensations, and the solutions not necessarily have to be isotonic to reduce pain on injection in the patient. The viscosity of the antibody solution may be such that at shear rates of 100 (1/s) antibody solution viscosity is below 200 mPa s, in an embodiment below 125 mPa s, in another embodiment below 70 mPa s, and in yet another embodiment below 25 mPa s or even below 10 mPa s.

B 5.3. Production efficiency

The generation of a DVD-Ig that is efficiently expressed in mammalian cells, such as Chinese hamster ovary cells (CHO), will in an embodiment require two parental monoclonal antibodies which are themselves expressed efficiently in mammalian cells. The production yield from a stable mammalian line (i.e., CHO) should be above about 0.5g/L, in an embodiment above about 1 g/L, and in another embodiment in the range of from about 2-5 g/L or more (Kipriyanov SM, Little M. 1999 Mol Biotechnol. 12:173-201; Carroll S, Al-Rubeai M. 2004 Expert Opin Biol Ther. 4:1821-9).

Production of antibodies and Ig fusion proteins in mammalian cells is influenced by several factors. Engineering of the expression vector via incorporation of strong promoters, enhancers and selection markers can maximize transcription of the gene of interest from an integrated vector copy. The identification of vector integration sites that are permissive for high

levels of gene transcription can augment protein expression from a vector (Wurm et al, 2004, Nature Biotechnology , 2004, Vol/Iss/Pg. 22/11 (1393-1398)). Furthermore, levels of production are affected by the ratio of antibody heavy and light chains and various steps in the process of protein assembly and secretion (Jiang et al. 2006, Biotechnology Progress, Jan-Feb 2006, vol. 22, no. 1, p. 313-8).

B 6. Immunogenicity

Administration of a therapeutic mAb may results in certain incidence of an immune response (ie, the formation of endogenous antibodies directed against the therapeutic mAb). Potential elements that might induce immunogenicity should be analyzed during selection of the parental monoclonal antibodies, and steps to reduce such risk can be taken to optimize the parental monoclonal antibodies prior to DVD-Ig construction. Mouse-derived antibodies have been found to be highly immunogenic in patients. The generation of chimeric antibodies comprised of mouse variable and human constant regions presents a logical next step to reduce the immunogenicity of therapeutic antibodies (Morrison and Schlom, 1990). Alternatively, immunogenicity can be reduced by transferring murine CDR sequences into a human antibody framework (reshaping/CDR grafting/humanization), as described for a therapeutic antibody by Riechmann et al., 1988. Another method is referred to as “resurfacing” or “veneering”, starting with the rodent variable light and heavy domains, only surface-accessible framework amino acids are altered to human ones, while the CDR and buried amino acids remain from the parental rodent antibody (Roguska et al., 1996). In another type of humanization, instead of grafting the entire CDRs, one technique grafts only the “specificity-determining regions” (SDRs), defined as the subset of CDR residues that are involved in binding of the antibody to its target (Kashmiri et al., 2005). This necessitates identification of the SDRs either through analysis of available three-dimensional structures of antibody-target complexes or mutational analysis of the antibody CDR residues to determine which interact with the target. Alternatively, fully human antibodies may have reduced immunogenicity compared to murine, chimeric or humanized antibodies.

Another approach to reduce the immunogenicity of therapeutic antibodies is the elimination of certain specific sequences that are predicted to be immunogenic. In one approach, after a first generation biologic has been tested in humans and found to be unacceptably immunogenic, the B-cell epitopes can be mapped and then altered to avoid immune detection. Another approach uses methods to predict and remove potential T-cell epitopes. Computational methods have been developed to scan and to identify the peptide sequences of biologic therapeutics with the potential to bind to MHC proteins (Desmet et al., 2005). Alternatively a human dendritic cell-based method can be used to identify CD4⁺ T-cell epitopes in potential protein allergens (Stickler et al., 2005; S.L. Morrison and J. Schlom, *Important Adv. Oncol.* (1990), pp. 3–18; Riechmann, L., Clark, M., Waldmann, H. and Winter, G. "Reshaping human

antibodies for therapy." Nature (1988) 332: 323-327; Roguska-M-A, Pedersen-J-T, Henry-A-H, Searle-S-M, Roja-C-M, Avery-B, Hoffee-M, Cook-S, Lambert-J-M, Blättler-W-A, Rees-A-R, Guild-B-C. A comparison of two murine mAbs humanized by CDR-grafting and variable domain resurfacing. Protein engineering, {Protein-Eng}, 1996, vol. 9, p. 895-904; Kashmiri-Syed-V-S, De-Pascalis-Roberto, Gonzales-Noreen-R, Schlom-Jeffrey. SDR grafting--a new approach to antibody humanization. Methods (San Diego Calif.), {Methods}, May 2005, vol. 36, no. 1, p. 25-34; Desmet-Johan, Meersseman-Geert, Boutonnet-Nathalie, Pletinckx-Jurgen, De-Clercq-Krista, Debulpaep-Maja, Braeckman-Tessa, Lasters-Ignace. Anchor profiles of HLA-specific peptides: analysis by a novel affinity scoring method and experimental validation. Proteins, 2005, vol. 58, p. 53-69; Stickler-M-M, Estell-D-A, Harding-F-A. CD4+ T-cell epitope determination using unexposed human donor peripheral blood mononuclear cells. Journal of immunotherapy 2000, vol. 23, p. 654-60.)

B 7. In vivo efficacy

To generate a DVD-Ig molecule with desired in vivo efficacy, it is important to generate and select mAbs with similarly desired in vivo efficacy when given in combination. However, in some instances the DVD-Ig may exhibit in vivo efficacy that cannot be achieved with the combination of two separate mAbs. For instance, a DVD-Ig may bring two targets in close proximity leading to an activity that cannot be achieved with the combination of two separate mAbs. Additional desirable biological functions are described herein in section **B 3**. Parent antibodies with characteristics desirable in the DVD-Ig molecule may be selected based on factors such as pharmacokinetic $t_{1/2}$; tissue distribution; soluble versus cell surface targets; and target concentration- soluble/density –surface.

B 8. In vivo tissue distribution

To generate a DVD-Ig molecule with desired in vivo tissue distribution, in an embodiment parent mAbs with similar desired in vivo tissue distribution profile must be selected. Alternatively, based on the mechanism of the dual-specific targeting strategy, it may at other times not be required to select parent mAbs with the similarly desired in vivo tissue distribution when given in combination. For instance, in the case of a DVD-Ig in which one binding component targets the DVD-Ig to a specific site thereby bringing the second binding component to the same target site. For example, one binding specificity of a DVD-Ig could target pancreas (islet cells) and the other specificity could bring GLP1 to the pancreas to induce insulin.

B 9. Isotype:

To generate a DVD-Ig molecule with desired properties including, but not limited to, Isotype, Effector functions and the circulating half-life, in an embodiment parent mAbs with

appropriate Fc-effector functions depending on the therapeutic utility and the desired therapeutic end-point are selected. There are five main heavy-chain classes or isotypes some of which have several sub-types and these determine the effector functions of an antibody molecule. These effector functions reside in the hinge region, CH2 and CH3 domains of the antibody molecule.

5 However, residues in other parts of an antibody molecule may have effects on effector functions as well. The hinge region Fc-effector functions include: (i) antibody-dependent cellular cytotoxicity, (ii) complement (C1q) binding, activation and complement-dependent cytotoxicity (CDC), (iii) phagocytosis/clearance of antigen-antibody complexes, and (iv) cytokine release in some instances. These Fc-effector functions of an antibody molecule are mediated through the
10 interaction of the Fc-region with a set of class-specific cell surface receptors. Antibodies of the IgG1 isotype are most active while IgG2 and IgG4 having minimal or no effector functions. The effector functions of the IgG antibodies are mediated through interactions with three structurally homologous cellular Fc receptor types (and sub-types) (FcγR1, FcγR2 and FcγR3). These effector functions of an IgG1 can be eliminated by mutating specific amino acid residues in the
15 lower hinge region (e.g., L234A, L235A) that are required for FcγR and C1q binding. Amino acid residues in the Fc region, in particular the CH2-CH3 domains, also determine the circulating half-life of the antibody molecule. This Fc function is mediated through the binding of the Fc-region to the neonatal Fc receptor (FcRn) which is responsible for recycling of antibody molecules from the acidic lysosomes back to the general circulation.

20 Whether a mAb should have an active or an inactive isotype will depend on the desired therapeutic end-point for an antibody. Some examples of usage of isotypes and desired therapeutic outcome are listed below:

- a) If the desired end-point is functional neutralization of a soluble cytokine then an inactive isotype may be used;
- 25 b) If the desired out-come is clearance of a pathological protein an active isotype may be used;
- c) If the desired out-come is clearance of protein aggregates an active isotype may be used;
- d) If the desired outcome is to antagonize a surface receptor an inactive isotype is used (Tysabri, IgG4; OKT3, mutated IgG1);
- 30 e) If the desired outcome is to eliminate target cells an active isotype is used (Herceptin, IgG1 (and with enhanced effector functions); and
- f) If the desired outcome is to clear proteins from circulation without entering the CNS an IgM isotype may be used (e.g., clearing circulating Ab peptide species).

The Fc effector functions of a parental mAb can be determined by various in vitro methods well known in the art.

As discussed, the selection of isotype, and thereby the effector functions will depend upon the desired therapeutic end-point. In cases where simple neutralization of a circulating target is desired, for example blocking receptor-ligand interactions, the effector functions may not be required. In such instances isotypes or mutations in the Fc-region of an antibody that eliminate effector functions are desirable. In other instances where elimination of target cells is the therapeutic end-point, for example elimination of tumor cells, isotypes or mutations or defucosylation in the Fc-region that enhance effector functions are desirable (Presta GL, Adv. Drug Delivery Rev. 58:640-656, 2006; Satoh M., Iida S., Shitara K. Expert Opinion Biol. Ther. 6:1161-1173, 2006). Similarly, depending up on the therapeutic utility, the circulating half-life of an antibody molecule can be reduced/prolonged by modulating antibody-FcRn interactions by introducing specific mutations in the Fc region (Dall'Acqua WF, Kiener PA, Wu H. J. Biol. Chem. 281:23514-23524, 2006; Petkova SB., Akilesh S., Sproule TJ. et al. Internat. Immunol. 18:1759-1769, 2006; Vaccaro C., Bawdon R., Wanjie S et al. PNAS 103:18709-18714, 2007).

The published information on the various residues that influence the different effector functions of a normal therapeutic mAb may need to be confirmed for DVD-Ig. It may be possible that in a DVD-Ig format additional (different) Fc-region residues, other than those identified for the modulation of monoclonal antibody effector functions, may be important.

Overall, the decision as to which Fc-effector functions (isotype) will be critical in the final DVD-Ig format will depend up on the disease indication, therapeutic target, desired therapeutic end-point and safety considerations. Listed below are exemplary appropriate heavy chain and light chain constant regions including, but not limited to:

- IgG1 – allotype: G1mz
- IgG1 mutant – A234, A235
- IgG2 – allotype: G2m(n-)
- Kappa – Km3
- Lambda

Fc Receptor and C1q Studies: The possibility of unwanted antibody-dependent cell-mediated cytotoxicity (ADCC) and complement-dependent cytotoxicity (CDC) by antibody complexing to any overexpressed target on cell membranes can be abrogated by the (for example, L234A, L235A) hinge-region mutations. These substituted amino acids, present in the IgG1 hinge region of mAb, are expected to result in diminished binding of mAb to human Fc receptors (but not FcRn), as FcγR binding is thought to occur within overlapping sites on the IgG1 hinge

region. This feature of mAb may lead to an improved safety profile over antibodies containing a wild-type IgG. Binding of mAb to human Fc receptors can be determined by flow cytometry experiments using cell lines (e.g., THP-1, K562) and an engineered CHO cell line that expresses FcγRIIb (or other FcγRs). Compared to IgG1 control monoclonal antibodies, mAb show reduced binding to FcγRI and FcγRIIa whereas binding to FcγRIIb is unaffected. The binding and activation of C1q by antigen/IgG immune complexes triggers the classical complement cascade with consequent inflammatory and/or immunoregulatory responses. The C1q binding site on IgGs has been localized to residues within the IgG hinge region. C1q binding to increasing concentrations of mAb was assessed by C1q ELISA. The results demonstrate that mAb is unable to bind to C1q, as expected when compared to the binding of a wildtype control IgG1. Overall, the L234A, L235A hinge region mutation abolishes binding of mAb to FcγRI, FcγRIIa and C1q but does not impact the interaction of mAb with FcγRIIb. This data suggests that in vivo, mAb with mutant Fc will interact normally with the inhibitory FcγRIIb but will likely fail to interact with the activating FcγRI and FcγRIIa receptors or C1q.

Human FcRn binding: The neonatal receptor (FcRn) is responsible for transport of IgG across the placenta and to control the catabolic half-life of the IgG molecules. It might be desirable to increase the terminal half-life of an antibody to improve efficacy, to reduce the dose or frequency of administration, or to improve localization to the target. Alternatively, it might be advantageous to do the converse that is, to decrease the terminal half-life of an antibody to reduce whole body exposure or to improve the target-to-non-target binding ratios. Tailoring the interaction between IgG and its salvage receptor, FcRn, offers a way to increase or decrease the terminal half-life of IgG. Proteins in the circulation, including IgG, are taken up in the fluid phase through micropinocytosis by certain cells, such as those of the vascular endothelia. IgG can bind FcRn in endosomes under slightly acidic conditions (pH 6.0–6.5) and can recycle to the cell surface, where it is released under almost neutral conditions (pH 7.0–7.4). Mapping of the Fc-region-binding site on FcRn80, 16, 17 showed that two histidine residues that are conserved across species, His310 and His435, are responsible for the pH dependence of this interaction. Using phage-display technology, a mouse Fc-region mutation that increases binding to FcRn and extends the half-life of mouse IgG was identified (see Victor, G. et al.; *Nature Biotechnology* (1997), 15(7), 637-640). Fc-region mutations that increase the binding affinity of human IgG for FcRn at pH 6.0, but not at pH 7.4, have also been identified (see Dall'Acqua William F, et al., *Journal of Immunology* (2002), 169(9), 5171-80). Moreover, in one case, a similar pH-dependent increase in binding (up to 27-fold) was also observed for rhesus FcRn, and this resulted in a twofold increase in serum half-life in rhesus monkeys compared with the parent IgG (see Hinton, Paul R. et al., *Journal of Biological Chemistry* (2004), 279(8), 6213-6216). These findings indicate that it is feasible to extend the plasma half-life of antibody therapeutics by tailoring the

interaction of the Fc region with FcRn. Conversely, Fc-region mutations that attenuate interaction with FcRn can reduce antibody half-life.

B.10 Pharmacokinetics (PK):

To generate a DVD-Ig molecule with desired pharmacokinetic profile, in an embodiment
5 parent mAbs with the similarly desired pharmacokinetic profile are selected. One consideration is that immunogenic response to monoclonal antibodies (ie, HAAA, human anti-human antibody response; HACA, human anti-chimeric antibody response) further complicates the pharmacokinetics of these therapeutic agents. In an embodiment, monoclonal antibodies with minimal or no immunogenicity are used for constructing DVD-Ig molecules such that the
10 resulting DVD-Igs will also have minimal or no immunogenicity. Some of the factors that determine the PK of a mAb include, but are not limited to, Intrinsic properties of the mAb (VH amino acid sequence); immunogenicity; FcRn binding and Fc functions.

The PK profile of selected parental monoclonal antibodies can be easily determined in rodents as the PK profile in rodents correlates well with (or closely predicts) the PK profile of
15 monoclonal antibodies in cynomolgus monkey and humans. The PK profile is determined as described in Example section 1.2.2.3.A.

After the parental monoclonal antibodies with desired PK characteristics (and other desired functional properties as discussed herein) are selected, the DVD-Ig is constructed. As the DVD-Ig molecules contain two antigen-binding domains from two parental monoclonal antibodies, the PK
20 properties of the DVD-Ig are assessed as well. Therefore, while determining the PK properties of the DVD-Ig, PK assays may be employed that determine the PK profile based on functionality of both antigen-binding domains derived from the 2 parent monoclonal antibodies. The PK profile of a DVD-Ig can be determined as described in Example 1.2.2.3.A. Additional factors that may impact the PK profile of DVD-Ig include the antigen-binding domain (CDR) orientation; Linker
25 size; and Fc / FcRn interactions. PK characteristics of parent antibodies can be evaluated by assessing the following parameters: absorption, distribution, metabolism and excretion.

Absorption: To date, administration of therapeutic monoclonal antibodies is via parenteral routes (e.g., intravenous [IV], subcutaneous [SC], or intramuscular [IM]). Absorption of a mAb into the systemic circulation following either SC or IM administration from the
30 interstitial space is primarily through the lymphatic pathway. Saturable, presystemic, proteolytic degradation may result in variable absolute bioavailability following extravascular administration. Usually, increases in absolute bioavailability with increasing doses of monoclonal antibodies may be observed due to saturated proteolytic capacity at higher doses. The absorption process for a mAb is usually quite slow as the lymph fluid drains slowly into the vascular system, and the

duration of absorption may occur over hours to several days. The absolute bioavailability of monoclonal antibodies following SC administration generally ranges from 50% to 100%.

Distribution: Following IV administration, monoclonal antibodies usually follow a biphasic serum (or plasma) concentration-time profile, beginning with a rapid distribution phase, followed by a slow elimination phase. In general, a biexponential pharmacokinetic model best describes this kind of pharmacokinetic profile. The volume of distribution in the central compartment (V_c) for a mAb is usually equal to or slightly larger than the plasma volume (2-3 liters). A distinct biphasic pattern in serum (plasma) concentration versus time profile may not be apparent with other parenteral routes of administration, such as IM or SC, because the distribution phase of the serum (plasma) concentration-time curve is masked by the long absorption portion. Many factors, including physicochemical properties, site-specific and target-oriented receptor mediated uptake, binding capacity of tissue, and mAb dose can influence biodistribution of a mAb. Some of these factors can contribute to nonlinearity in biodistribution for a mAb.

Metabolism and Excretion: Due to the molecular size, intact monoclonal antibodies are not excreted into the urine via kidney. They are primarily inactivated by metabolism (e.g., catabolism). For IgG-based therapeutic monoclonal antibodies, half-lives typically ranges from hours or 1-2 days to over 20 days. The elimination of a mAb can be affected by many factors, including, but not limited to, affinity for the FcRn receptor, immunogenicity of the mAb, the degree of glycosylation of the mAb, the susceptibility for the mAb to proteolysis, and receptor-mediated elimination.

B.11 Tissue cross-reactivity pattern on human and tox species:

Identical staining pattern suggests that potential human toxicity can be evaluated in tox species. Tox species are those animal in which unrelated toxicity is studied.

The individual antibodies are selected to meet two criteria. (1) Tissue staining appropriate for the known expression of the antibody target. (2) Similar staining pattern between human and tox species tissues from the same organ.

Criterion 1: Immunizations and/or antibody selections typically employ recombinant or synthesized antigens (proteins, carbohydrates or other molecules). Binding to the natural counterpart and counterscreen against unrelated antigens are often part of the screening funnel for therapeutic antibodies. However, screening against a multitude of antigens is often impractical. Therefore tissue cross-reactivity studies with human tissues from all major organs serve to rule out unwanted binding of the antibody to any unrelated antigens.

Criterion 2: Comparative tissue cross reactivity studies with human and tox species tissues (cynomolgus monkey, dog, possibly rodents and others, the same 36 or 37 tissues are

being tested as in the human study) help to validate the selection of a tox species. In the typical tissue cross-reactivity studies on frozen tissues sections therapeutic antibodies may demonstrate the expected binding to the known antigen and/or to a lesser degree binding to tissues based either on low level interactions (unspecific binding, low level binding to similar antigens, low level charge based interactions etc.). In any case the most relevant toxicology animal species is the one with the highest degree of coincidence of binding to human and animal tissue.

Tissue cross reactivity studies follow the appropriate regulatory guidelines including EC CPMP Guideline III/5271/94 “Production and quality control of mAbs” and the 1997 US FDA/CBER “Points to Consider in the Manufacture and Testing of Monoclonal Antibody Products for Human Use”. . Cryosections (5 μ m) of human tissues obtained at autopsy or biopsy were fixed and dried on object glass. The peroxidase staining of tissue sections was performed, using the avidin-biotin system. FDA’s Guidance “*Points to Consider in the Manufacture and Testing of Monoclonal Antibody Products for Human Use*”. Relevant references include Clarke J 2004, Boon L. 2002a, Boon L 2002b, Ryan A 1999.

Tissue cross reactivity studies are often done in two stages, with the first stage including cryosections of 32 tissues (typically: Adrenal Gland, Gastrointestinal Tract, Prostate, Bladder, Heart, Skeletal Muscle, Blood Cells, Kidney, Skin, Bone Marrow, Liver, Spinal Cord, Breast, Lung, Spleen, Cerebellum, Lymph Node, Testes, Cerebral Cortex, Ovary, Thymus, Colon, Pancreas, Thyroid, Endothelium, Parathyroid, Ureter, Eye, Pituitary, Uterus, Fallopian Tube and Placenta) from one human donor. In the second phase a full cross reactivity study is performed with up to 38 tissues (including adrenal, blood, blood vessel, bone marrow, cerebellum, cerebrum, cervix, esophagus, eye, heart, kidney, large intestine, liver, lung, lymph node, breast mammary gland, ovary, oviduct, pancreas, parathyroid, peripheral nerve, pituitary, placenta, prostate, salivary gland, skin, small intestine, spinal cord, spleen, stomach, striated muscle, testis, thymus, thyroid, tonsil, ureter, urinary bladder, and uterus) from 3 unrelated adults. Studies are done typically at minimally two dose levels.

The therapeutic antibody (i.e. test article) and isotype matched control antibody may be biotinylated for avidin-biotin complex (ABC) detection; other detection methods may include tertiary antibody detection for a FITC (or otherwise) labeled test article, or precomplexing with a labeled anti-human IgG for an unlabeled test article.

Briefly, cryosections (about 5 μ m) of human tissues obtained at autopsy or biopsy are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system. First (in case of a precomplexing detection system), the test article is incubated with the secondary biotinylated anti-human IgG and developed into immune complex. The immune complex at the final concentrations of 2 and 10 μ g/mL of test article is added onto

tissue sections on object glass and then the tissue sections were reacted for 30 minutes with a avidin-biotin-peroxidase kit. Subsequently, DAB (3,3'-diaminobenzidine), a substrate for the peroxidase reaction, was applied for 4 minutes for tissue staining. Antigen-Sepharose beads are used as positive control tissue sections.

5 Any specific staining is judged to be either an expected (e.g., consistent with antigen expression) or unexpected reactivity based upon known expression of the target antigen in question. Any staining judged specific is scored for intensity and frequency. Antigen or serum competition or blocking studies can assist further in determining whether observed staining is specific or nonspecific.

10 If two selected antibodies are found to meet the selection criteria – appropriate tissue staining, matching staining between human and toxicology animal specific tissue – they can be selected for DVD-Ig generation.

 The tissue cross reactivity study has to be repeated with the final DVD-Ig construct, but while these studies follow the same protocol as outline herein, they are more complex to evaluate
15 because any binding can come from any of the two parent antibodies, and any unexplained binding needs to be confirmed with complex antigen competition studies.

 It is readily apparent that the complex undertaking of tissue crossreactivity studies with a multispecific molecule like a DVD-Ig is greatly simplified if the two parental antibodies are selected for (1) lack of unexpected tissue cross reactivity findings and (2) for appropriate
20 similarity of tissue cross reactivity findings between the corresponding human and toxicology animal species tissues.

B.12 Specificity and selectivity:

 To generate a DVD-Ig molecule with desired specificity and selectivity, one needs to generate and select parent mAbs with the similarly desired specificity and selectivity profile.

25 Binding studies for specificity and selectivity with a DVD-Ig can be complex due to the four or more binding sites, two each for each antigen. Briefly, binding studies using ELISA, BIAcore. KinExA or other interaction studies with a DVD-Ig need to monitor the binding of one, two or more antigens to the DVD-Ig molecule. While BIAcore technology can resolve the sequential, independent binding of multiple antigens, more traditional methods including ELISA
30 or more modern techniques like KinExA cannot. Therefore careful characterization of each parent antibody is critical. After each individual antibody has been characterized for specificity, confirmation of specificity retention of the individual binding sites in the DVD-Ig molecule is greatly simplified.

It is readily apparent that the complex undertaking of determining the specificity of a DVD-Ig is greatly simplified if the two parental antibodies are selected for specificity prior to being combined into a DVD-Ig.

Antigen-antibody interaction studies can take many forms, including many classical protein protein interaction studies, including ELISA (Enzyme linked immunosorbent assay), Mass spectrometry, chemical cross linking, SEC with light scattering, equilibrium dialysis, gel permeation, ultrafiltration, gel chromatography, large-zone analytical SEC, micropreparative ultracentrifugation (sedimentation equilibrium), spectroscopic methods, titration microcalorimetry, sedimentation equilibrium (in analytical ultracentrifuge), sedimentation velocity (in analytical centrifuge), surface plasmon resonance (including BIAcore). Relevant references include "Current Protocols in Protein Science", John E. Coligan, Ben M. Dunn, David W. Speicher, Paul T. Wingfield (eds.) Volume 3, chapters 19 and 20, published by John Wiley & Sons Inc., and references included therein and "Current Protocols in Immunology", John E. Coligan, Barbara E. Bierer, David H. Margulies, Ethan M. Shevach, Warren Strober (eds.) published by John Wiley & Sons Inc and relevant references included therein.

Cytokine Release in Whole Blood: The interaction of mAb with human blood cells can be investigated by a cytokine release assay (Wing, M. G. Therapeutic Immunology (1995), 2(4), 183-190; "Current Protocols in Pharmacology", S.J. Enna, Michael Williams, John W. Ferkany, Terry Kenakin, Paul Moser, (eds.) published by John Wiley & Sons Inc; Madhusudan, S. Clinical Cancer Research (2004), 10(19), 6528-6534; Cox, J. Methods (2006), 38(4), 274-282; Choi, I. European Journal of Immunology (2001), 31(1), 94-106). Briefly, various concentrations of mAb are incubated with human whole blood for 24 hours. The concentration tested should cover a wide range including final concentrations mimicking typical blood levels in patients (including but not limited to 100 ng/ml – 100µg/ml). Following the incubation, supernatants and cell lysates were analyzed for the presence of IL-1R α , TNF- α , IL-1b, IL-6 and IL-8. Cytokine concentration profiles generated for mAb were compared to profiles produced by a negative human IgG control and a positive LPS or PHA control. The cytokine profile displayed by mAb from both cell supernatants and cell lysates was comparable to control human IgG. In an embodiment, the monoclonal antibody does not interact with human blood cells to spontaneously release inflammatory cytokines.

Cytokine release studies for a DVD-Ig are complex due to the four or more binding sites, two each for each antigen. Briefly, cytokine release studies as described herein measure the effect of the whole DVD-Ig molecule on whole blood or other cell systems, but can resolve which portion of the molecule causes cytokine release. Once cytokine release has been detected, the purity of the DVD-Ig preparation has to be ascertained, because some co-purifying cellular components can cause cytokine release on their own. If purity is not the issue, fragmentation of

DVD-Ig (including but not limited to removal of Fc portion, separation of binding sites etc.), binding site mutagenesis or other methods may need to be employed to deconvolute any observations. It is readily apparent that this complex undertaking is greatly simplified if the two parental antibodies are selected for lack of cytokine release prior to being combined into a DVD-Ig.

B.13 Cross reactivity to other species for toxicological studies:

In an embodiment, the individual antibodies selected with sufficient cross-reactivity to appropriate tox species, for example, cynomolgus monkey. Parental antibodies need to bind to orthologous species target (i.e. cynomolgus monkey) and elicit appropriate response (modulation, neutralization, activation). In an embodiment, the cross-reactivity (affinity/potency) to orthologous species target should be within 10-fold of the human target. In practice, the parental antibodies are evaluated for multiple species, including mouse, rat, dog, monkey (and other non-human primates), as well as disease model species (i.e. sheep for asthma model). The acceptable cross-reactivity to tox species from the parental monoclonal antibodies allows future toxicology studies of DVD-Ig-Ig in the same species. For that reason, the two parental monoclonal antibodies should have acceptable cross-reactivity for a common tox species therefore allowing toxicology studies of DVD-Ig in the same species.

Parent mAbs may be selected from various mAbs capable of binding specific targets and well known in the art. These include, but are not limited to anti-TNF antibody (US Patent No. 6,258,562), anti-IL-12 and/or anti-IL-12p40 antibody (US Patent No. 6,914,128); anti-IL-18 antibody (US 2005/0147610 A1), anti-C5, anti-CBL, anti-CD147, anti-gp120, anti-VLA-4, anti-CD11a, anti-CD18, anti-VEGF, anti-CD40L, anti CD-40 (e.g., see WO2007124299) anti-Id, anti-ICAM-1, anti-CXCL13, anti-CD2, anti-EGFR, anti-TGF-beta 2, anti-HGF, anti-cMet, anti DLL-4, anti-NPR1, anti-PLGF, anti-ErbB3, anti-E-selectin, anti-Fact VII, anti-Her2/neu, anti-F gp, anti-CD11/18, anti-CD14, anti-ICAM-3, anti-RON, anti-SOST, anti CD-19, anti-CD80 (e.g., see WO2003039486, anti-CD4, anti-CD3, anti-CD23, anti-beta2-integrin, anti-alpha4beta7, anti-CD52, anti-HLA DR, anti-CD22 (e.g., see US Patent NO: 5,789,554), anti-CD20, anti-MIF, anti-CD64 (FcR), anti-TCR alpha beta, anti-CD2, anti-Hep B, anti-CA 125, anti-EpCAM, anti-gp120, anti-CMV, anti-gpIIbIIIa, anti-IgE, anti-CD25, anti-CD33, anti-HLA, anti-IGF1,2, anti IGFR, anti-VNRintegrin, anti-IL-1alpha, anti-IL-1beta, anti-IL-1 receptor, anti-IL-2 receptor, anti-IL-4, anti-IL-4 receptor, anti-IL5, anti-IL-5 receptor, anti-IL-6, anti-IL-8, anti-IL-9, anti-IL-13, anti-IL-13 receptor, anti-IL-17, and anti-IL-23 (see Presta LG. 2005 Selection, design, and engineering of therapeutic antibodies J Allergy Clin Immunol. 116:731-6 and <http://www.path.cam.ac.uk/~mrc7/humanisation/antibodies.html>).

Parent mAbs may also be selected from various therapeutic antibodies approved for use, in clinical trials, or in development for clinical use. Such therapeutic antibodies include, but are not limited to, rituximab (Rituxan®, IDEC/Genentech/Roche) (see for example U. S. Pat. No. 5,736,137), a chimeric anti-CD20 antibody approved to treat Non-Hodgkin's lymphoma;

5 HuMax-CD20, an anti-CD20 currently being developed by Genmab, an anti-CD20 antibody described in U.S. Pat. No. 5, 500,362, AME-133 (Applied Molecular Evolution), hA20 (Immunomedics, Inc.), HumaLYM (Intracel), and PRO70769 (PCT/US2003/040426, entitled "Immunoglobulin Variants and Uses Thereof"), trastuzumab (Herceptin®, Genentech) (see for example U.S. Pat. No. 5,677,171), a humanized anti- Her2/neu antibody approved to treat breast

10 cancer; pertuzumab (rhuMab-2C4, Omnitarg®), currently being developed by Genentech; an anti-Her2 antibody described in U.S. Pat. No. 4,753,894; cetuximab (Erbix®, Imclone) (U.S. Pat. No. 4,943,533; PCT WO 96/40210), a chimeric anti-EGFR antibody in clinical trials for a variety of cancers; ABX-EGF (U.S. Pat. No. 6,235,883), currently being developed by Abgenix-Immunex-Amgen; HuMax- EGFr (U.S. Ser. No. 10/172,317), currently being developed by

15 Genmab; 425, EMD55900, EMD62000, and EMD72000 (Merck KGaA) (U.S. Pat. No. 5,558,864; Murthy et al. 1987, Arch Biochem Biophys. 252(2):549-60; Rodeck et al., 1987, J Cell Biochem. 35(4):315-20; Kettleborough et al., 1991, Protein Eng. 4(7):773-83); ICR62 (Institute of Cancer Research) (PCT WO 95/20045; Modjtahedi et al., 1993, J. Cell Biophys. 1993, 22(1-3):129-46; Modjtahedi et al., 1993, Br J Cancer. 1993, 67(2):247-53; Modjtahedi et

20 al, 1996, Br J Cancer, 73(2):228-35; Modjtahedi et al, 2003, Int J Cancer, 105(2):273-80); TheraCIM hR3 (YM Biosciences, Canada and Centro de Immunologia Molecular, Cuba (U.S. Pat. No. 5,891,996; U.S. Pat. No. 6,506, 883; Mateo et al, 1997, Immunotechnology, 3(1):71-81); mAb-806 (Ludwig Institue for Cancer Research, Memorial Sloan-Kettering) (Jungbluth et al. 2003, Proc Natl Acad Sci USA. 100(2):639-44); KSB-102 (KS Biomedix); MR1-1 (IVAX, National Cancer Institute) (PCT WO 0162931A2); and SC100 (Scancell) (PCT WO 01/88138);

25 alemtuzumab (Campath®, Millenium), a humanized mAb currently approved for treatment of B-cell chronic lymphocytic leukemia; muromonab-CD3 (Orthoclone OKT3®), an anti-CD3 antibody developed by Ortho Biotech/Johnson & Johnson, ibritumomab tiuxetan (Zevalin®), an anti-CD20 antibody developed by IDEC/Schering AG, gemtuzumab ozogamicin (Mylotarg®),

30 an anti-CD33 (p67 protein) antibody developed by Celltech/Wyeth, alefacept (Amevive®), an anti-LFA-3 Fc fusion developed by Biogen), abciximab (ReoPro®), developed by Centocor/Lilly, basiliximab (Simulect®), developed by Novartis, palivizumab (Synagis®), developed by Medimmune, infliximab (Remicade®), an anti-TNFalpha antibody developed by Centocor, adalimumab (Humira®), an anti-TNFalpha antibody developed by Abbott,

35 Humicade®, an anti-TNFalpha antibody developed by Celltech, golimumab (CNTO-148), a fully human TNF antibody developed by Centocor, etanercept (Enbrel®), an p75 TNF receptor Fc fusion developed by Immunex/Amgen, lenercept, an p55TNF receptor Fc fusion previously

developed by Roche, ABX-CBL, an anti-CD147 antibody being developed by Abgenix, ABX-IL8, an anti-IL8 antibody being developed by Abgenix, ABX-MA1, an anti-MUC18 antibody being developed by Abgenix, Pemtumomab (R1549, 90Y-muHMFG1), an anti-MUC1 in development by Antisoma, Therex (R1550), an anti-MUC1 antibody being developed by
 5 Antisoma, AngioMab (AS1405), being developed by Antisoma, HuBC-1, being developed by Antisoma, Thioplatin (AS1407) being developed by Antisoma, Antegren® (natalizumab), an anti-alpha-4-beta-1 (VLA-4) and alpha-4-beta-7 antibody being developed by Biogen, VLA-1 mAb, an anti-VLA-1 integrin antibody being developed by Biogen, LTBR mAb, an anti-lymphotoxin beta receptor (LTBR) antibody being developed by Biogen, CAT-152, an anti-
 10 TGF-β2 antibody being developed by Cambridge Antibody Technology, ABT 874 (J695), an anti-IL-12 p40 antibody being developed by Abbott, CAT-192, an anti-TGFβ1 antibody being developed by Cambridge Antibody Technology and Genzyme, CAT-213, an anti-Eotaxin1 antibody being developed by Cambridge Antibody Technology, LymphoStat-B® an anti-Blys antibody being developed by Cambridge Antibody Technology and Human Genome Sciences
 15 Inc., TRAIL-R1mAb, an anti-TRAIL-R1 antibody being developed by Cambridge Antibody Technology and Human Genome Sciences, Inc., Avastin® bevacizumab, rhuMAb-VEGF), an anti-VEGF antibody being developed by Genentech, an anti-HER receptor family antibody being developed by Genentech, Anti-Tissue Factor (ATF), an anti-Tissue Factor antibody being developed by Genentech, Xolair® (Omalizumab), an anti-IgE antibody being developed by
 20 Genentech, Raptiva® (Efalizumab), an anti-CD11a antibody being developed by Genentech and Xoma, MLN-02 Antibody (formerly LDP-02), being developed by Genentech and Millenium Pharmaceuticals, HuMax CD4, an anti-CD4 antibody being developed by Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Genmab and Amgen, HuMax-Inflam, being developed by Genmab and Medarex, HuMax-Cancer, an anti-Heparanase I antibody being
 25 developed by Genmab and Medarex and Oxford GcoSciences, HuMax-Lymphoma, being developed by Genmab and Amgen, HuMax-TAC, being developed by Genmab, IDEC-131, and anti-CD40L antibody being developed by IDEC Pharmaceuticals, IDEC-151 (Clenoliximab), an anti-CD4 antibody being developed by IDEC Pharmaceuticals, IDEC-114, an anti-CD80 antibody being developed by IDEC Pharmaceuticals, IDEC-152, an anti-CD23 being developed
 30 by IDEC Pharmaceuticals, anti-macrophage migration factor (MIF) antibodies being developed by IDEC Pharmaceuticals, BEC2, an anti-idiotypic antibody being developed by Imclone, IMC-1C11, an anti-KDR antibody being developed by Imclone, DC101, an anti-flk-1 antibody being developed by Imclone, anti-VE cadherin antibodies being developed by Imclone, CEA-Cide® (labetuzumab), an anti-carcinoembryonic antigen (CEA) antibody being developed by
 35 Immunomedics, LymphoCide® (Epratuzumab), an anti-CD22 antibody being developed by Immunomedics, AFP-Cide, being developed by Immunomedics, MyelomaCide, being developed by Immunomedics, LkoCide, being developed by Immunomedics, ProstaCide, being developed

by Immunomedics, MDX-010, an anti-CTLA4 antibody being developed by Medarex, MDX-060, an anti-CD30 antibody being developed by Medarex, MDX-070 being developed by Medarex, MDX-018 being developed by Medarex, Osidem® (IDM-1), and anti-Her2 antibody being developed by Medarex and Immuno-Designed Molecules, HuMax®-CD4, an anti-CD4 antibody being developed by Medarex and Genmab, HuMax-IL15, an anti-IL15 antibody being developed by Medarex and Genmab, CNTO 148, an anti-TNF α antibody being developed by Medarex and Centocor/J&J, CNTO 1275, an anti-cytokine antibody being developed by Centocor/J&J, MOR101 and MOR102, anti-intercellular adhesion molecule-1 (ICAM-1) (CD54) antibodies being developed by MorphoSys, MOR201, an anti-fibroblast growth factor receptor 3 (FGFR-3) antibody being developed by MorphoSys, Nuvion® (visilizumab), an anti-CD3 antibody being developed by Protein Design Labs, HuZAF®, an anti-gamma interferon antibody being developed by Protein Design Labs, Anti- α 5 β 1 Integrin, being developed by Protein Design Labs, anti-IL-12, being developed by Protein Design Labs, ING-1, an anti-Ep-CAM antibody being developed by Xoma, Xolair® (Omalizumab) a humanized anti-IgE antibody developed by Genentech and Novartis, and MLN01, an anti-Beta2 integrin antibody being developed by Xoma, all of the herein-cited references in this paragraph are expressly incorporated herein by reference. In another embodiment, the therapeutics include KRN330 (Kirin); huA33 antibody (A33, Ludwig Institute for Cancer Research); CNTO 95 (alpha V integrins, Centocor); MEDI-522 (alpha V β 3 integrin, Medimmune); volociximab (alpha V β 1 integrin, Biogen/PDL); Human mAb 216 (B cell glycosylated epitope, NCI); BiTE MT103 (bispecific CD19 x CD3, Medimmune); 4G7xH22 (Bispecific BcellxFcgammaR1, Medarex/Merck KGaA); rM28 (Bispecific CD28 x MAPG, US Patent No. EP1444268); MDX447 (EMD 82633) (Bispecific CD64 x EGFR, Medarex); Catumaxomab (removab) (Bispecific EpCAM x anti-CD3, Trion/Fres); Ertumaxomab (bispecific HER2/CD3, Fresenius Biotech); oregovomab (OvaRex) (CA-125, ViRexx); Rencarex® (WX G250) (carbonic anhydrase IX, Willex); CNTO 888 (CCL2, Centocor); TRC105 (CD105 (endoglin), Tracon); BMS-663513 (CD137 agonist, Bristol Myers Squibb); MDX-1342 (CD19, Medarex); Siplizumab (MEDI-507) (CD2, Medimmune); Ofatumumab (Humax-CD20) (CD20, Genmab); Rituximab (Rituxan) (CD20, Genentech); velutuzumab (hA20) (CD20, Immunomedics); Epratuzumab (CD22, Amgen); lumiliximab (IDEC 152) (CD23, Biogen); muromonab-CD3 (CD3, Ortho); HuM291 (CD3 fc receptor, PDL Biopharma); HeFi-1, CD30, NCI); MDX-060 (CD30, Medarex); MDX-1401 (CD30, Medarex); SGN-30 (CD30, Seattle Genentics); SGN-33 (Lintuzumab) (CD33, Seattle Genentics); Zanolimumab (HuMax-CD4) (CD4, Genmab); HCD122 (CD40, Novartis); SGN-40 (CD40, Seattle Genentics); Campath1h (Alemtuzumab) (CD52, Genzyme); MDX-1411 (CD70, Medarex); hLL1 (EPB-1) (CD74.38, Immunomedics); Galiximab (IDEC-144) (CD80, Biogen); MT293 (TRC093/D93) (cleaved collagen, Tracon); HuLuc63 (CS1, PDL Pharma); ipilimumab (MDX-010) (CTLA4, Bristol Myers Squibb); Tremelimumab (Ticilimumab, CP-

675,2) (CTLA4, Pfizer); HGS-ETR1 (Mapatumumab) (DR4 TRAIL-R1 agonist, Human Genome Science /Glaxo Smith Kline); AMG-655 (DR5, Amgen); Apomab (DR5, Genentech); CS-1008 (DR5, Daiichi Sankyo); HGS-ETR2 (lexatumumab) (DR5 TRAIL-R2 agonist, HGS); Cetuximab (Erbix) (EGFR, Imclone); IMC-11F8, (EGFR, Imclone); Nimotuzumab (EGFR, YM Bio); Panitumumab (Vectabix) (EGFR, Amgen); Zalutumumab (HuMaxEGFr) (EGFR, Genmab); CDX-110 (EGFRvIII, AVANT Immunotherapeutics); adecatumumab (MT201) (Epcam, Merck); edrecolomab (Panorex, 17-1A) (Epcam, Glaxo/Centocor); MORAb-003 (folate receptor α , Morphotech); KW-2871 (ganglioside GD3, Kyowa); MORAb-009 (GP-9, Morphotech); CDX-1307 (MDX-1307) (hCgb, Celldex); Trastuzumab (Herceptin) (HER2, Celldex); Pertuzumab (rhuMAb 2C4) (HER2 (DI), Genentech); apolizumab (HLA-DR beta chain, PDL Pharma); AMG-479 (IGF-1R, Amgen); anti-IGF-1R R1507 (IGF1-R, Roche); CP 751871 (IGF1-R, Pfizer); IMC-A12 (IGF1-R, Imclone); BIIB022 (IGF-1R, Biogen); Mik-beta-1 (IL-2Rb (CD122), Hoffman LaRoche); CNTO 328 (IL6, Centocor); Anti-KIR (1-7F9) (Killer cell Ig-like Receptor (KIR), Novo); Hu3S193 (Lewis (y), Wyeth, Ludwig Institute of Cancer Research); hCBE-11 (LT β R, Biogen); HuHMFG1 (MUC1, Antisoma/NCI); RAV12 (N-linked carbohydrate epitope, Raven); CAL (parathyroid hormone-related protein (PTH-rP), University of California); CT-011 (PD1, CureTech); MDX-1106 (ono-4538) (PD1, Medarex/Ono); MAb CT-011 (PD1, Curetech); IMC-3G3 (PDGFR α , Imclone); bavituximab (phosphatidylserine, Peregrine); huJ591 (PSMA, Cornell Research Foundation); muJ591 (PSMA, Cornell Research Foundation); GC1008 (TGF β (pan) inhibitor (IgG4), Genzyme); Infliximab (Remicade) (TNF α , Centocor); A27.15 (transferrin receptor, Salk Institute, INSERN WO 2005/111082); E2.3 (transferrin receptor, Salk Institute); Bevacizumab (Avastin) (VEGF, Genentech); HuMV833 (VEGF, Tsukuba Research Lab-WO/2000/034337, University of Texas); IMC-18F1 (VEGFR1, Imclone); IMC-1121 (VEGFR2, Imclone).

25 **B. Construction of DVD molecules:**

The dual variable domain immunoglobulin (DVD-Ig) molecule is designed such that two different light chain variable domains (VL) from the two different parent monoclonal antibodies are linked in tandem directly or via a short linker by recombinant DNA techniques, followed by the light chain constant domain. Similarly, the heavy chain comprises two different heavy chain variable domains (VH) linked in tandem, followed by the constant domain CH1 and Fc region (Fig.1A).

The variable domains can be obtained using recombinant DNA techniques from a parent antibody generated by any one of the methods described herein. In an embodiment, the variable domain is a murine heavy or light chain variable domain. In another embodiment, the variable domain is a CDR grafted or a humanized variable heavy or light chain domain. In an embodiment, the variable domain is a human heavy or light chain variable domain.

In one embodiment the first and second variable domains are linked directly to each other using recombinant DNA techniques. In another embodiment the variable domains are linked via a linker sequence. In an embodiment, two variable domains are linked. Three or more variable domains may also be linked directly or via a linker sequence. The variable domains may bind the same antigen or may bind different antigens. DVD molecules of the invention may include one immunoglobulin variable domain and one non- immunoglobulin variable domain such as ligand binding domain of a receptor, active domain of an enzyme. DVD molecules may also comprise 2 or more non-Ig domains.

The linker sequence may be a single amino acid or a polypeptide sequence. In an embodiment, the linker sequences are selected from the group consisting of

AKTTPKLEEGEFSEAR (SEQ ID NO: 1); AKTTPKLEEGEFSEARV (SEQ ID NO: 2);
 AKTTPKLGG (SEQ ID NO: 3); SAKTTPKLGG (SEQ ID NO: 4); SAKTTP (SEQ ID NO: 5);
 RADAAP (SEQ ID NO: 6); RADAAPTVS (SEQ ID NO: 7); RADAAAAGGPGS (SEQ ID NO: 8);
 RADAAA(G₄S)₄ (SEQ ID NO: 9); SAKTTPKLEEGEFSEARV (SEQ ID NO: 10); ADAAP
 (SEQ ID NO: 11); ADAAPTVSIFPP (SEQ ID NO: 12); TVAAP (SEQ ID NO: 13);
 TVAAPSVFIFPP (SEQ ID NO: 14); QPKAAP (SEQ ID NO: 15); QPKAAPSVTLFPP (SEQ ID
 NO: 16); AKTTPP (SEQ ID NO: 17); AKTTPPSVTPLAP (SEQ ID NO: 18); AKTTAP (SEQ ID
 NO: 19); AKTTAPSVYPLAP (SEQ ID NO: 20); ASTKGP (SEQ ID NO: 21);
 ASTKGPSVFPLAP (SEQ ID NO: 22); GGGGSGGGGSGGGGS (SEQ ID NO: 23);
 GENKVEYAPALMALS (SEQ ID NO: 24); GPAKELTPLKEAKVS (SEQ ID NO: 25);
 GHEAAVMQVQYPAS (SEQ ID NO: 26); GGGGGGGGP (SEQ ID NO: 27); GGGGGGGGGP
 (SEQ ID NO: 28); PAPNLLGGP (SEQ ID NO: 29); PNLLGGP (SEQ ID NO: 30); GGGGGGGP
 (SEQ ID NO: 31); PAPELLGGP (SEQ ID NO: 32); PTISPAPNLLGGP (SEQ ID NO: 33);
 TVAADDDDKSVFIVPP (SEQ ID NO: 34); TVDDDDKAAP (SEQ ID NO: 35); LVPRGSAAP
 (SEQ ID NO: 36); ASDDDDK GGP (SEQ ID NO: 37); ALVPR GSGP (SEQ ID NO: 38);
 ASTDDDDK SVFPLAP (SEQ ID NO: 39); TVALVPR GSVFIFPP (SEQ ID NO: 40);
 ASTLVPR GSVFPLAP (SEQ ID NO: 41); TVAADDDK SVFIVPP (SEQ ID NO: 42);
 ASTDDDK SVFPLAP (SEQ ID NO: 43); LEVLFFQ GP (SEQ ID NO: 44); TVAALEVLFFQ
 GPAP (SEQ ID NO: 45); ASTLEVLFFQ GPLAP (SEQ ID NO: 46); PAPLEVLFFQ GP (SEQ ID
 NO: 47); TAENLYFQ GAP (SEQ ID NO: 48); AENLYFQ GA (SEQ ID NO: 49); PGPFGR
 SAGGP (SEQ ID NO: 50); PGPFGR SAGG (SEQ ID NO: 51); PQRGR SAG (SEQ ID NO: 52);
 PHYGR SGG (SEQ ID NO: 53); GPFGR SAGP (SEQ ID NO: 54); GDDDDDK GGP (SEQ ID
 NO: 55); AGDDDDDK GGP (SEQ ID NO: 56); GGDDDDDK GGP (SEQ ID NO: 57); AS; TVA;
 ASTK (SEQ ID NO: 58); ASTKGPSV (SEQ ID NO: 59); ASTKGPSVFP (SEQ ID NO: 60);
 TVAAPSV (SEQ ID NO: 61), TVAAPSVFI (SEQ ID NO: 62). The choice of linker sequences is
 based on crystal structure analysis of several Fab molecules. There is a natural flexible linkage

between the variable domain and the CH1/CL constant domain in Fab or antibody molecular structure. This natural linkage comprises approximately 10-12 amino acid residues, contributed by 4-6 residues from C-terminus of V domain and 4-6 residues from the N-terminus of CL/CH1 domain. DVD Igs of the invention were generated using N-terminal 5-6 amino acid residues, or
5 11-12 amino acid residues, of CL or CH1 as linker in light chain and heavy chain of DVD-Ig, respectively. The N-terminal residues of CL or CH1 domains, particularly the first 5-6 amino acid residues, adopt a loop conformation without strong secondary structures, therefore can act as flexible linkers between the two variable domains. The N-terminal residues of CL or CH1 domains are natural extension of the variable domains, as they are part of the Ig sequences,
10 therefore minimize to a large extent any immunogenicity potentially arising from the linkers and junctions.

Other linker sequences may include any sequence of any length of CL/CH1 domain but not all residues of CL/CH1 domain; for example the first 5-12 amino acid residues of the CL/CH1 domains; the light chain linkers can be from C κ or C λ ; and the heavy chain linkers can be derived
15 from CH1 of any isotypes, including C γ 1, C γ 2, C γ 3, C γ 4, C α 1, C α 2, C δ , C ϵ , and C μ . Linker sequences may also be derived from other proteins such as Ig-like proteins, (e.g. TCR, FcR, KIR); G/S based sequences (e.g. G4S repeats; SEQ ID NO: 63); hinge region-derived sequences; and other natural sequences from other proteins.

In an embodiment a constant domain is linked to the two linked variable domains using
20 recombinant DNA techniques. In an embodiment, sequence comprising linked heavy chain variable domains is linked to a heavy chain constant domain and sequence comprising linked light chain variable domains is linked to a light chain constant domain. In an embodiment, the constant domains are human heavy chain constant domain and human light chain constant domain respectively. In an embodiment, the DVD heavy chain is further linked to an Fc region. The Fc
25 region may be a native sequence Fc region, or a variant Fc region. In another embodiment, the Fc region is a human Fc region. In another embodiment the Fc region includes Fc region from IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, or IgD.

In another embodiment two heavy chain DVD polypeptides and two light chain DVD polypeptides are combined to form a DVD-Ig molecule. Table 4 lists amino acid sequences of
30 VH and VL regions of exemplary antibodies for targets useful for treating disease, e.g., for treating cancer. In an embodiment, the invention provides a DVD comprising at least two of the VH and/or VL regions listed in Table 4, in any orientation.

Table 4: List of Amino Acid Sequences of VH and VL regions of Antibodies for Generating DVD-Igs

SEQ ID No.	ABT Unique ID	Protein region	Sequence 1234567890123456789012345678901234567890
64	AB014VH	VH VEGF (seq. 1)	EVQLVESGGGLVQPGGSLRLSCAASGYTFTNYGMNWVRQA PGKGLEWVGWINTYTGEPTYAADFKRRFTFSLDTSKSTAY LQMNSLRAEDTAVYYCAKYPHYYGSSHWYFDVWGQGTLVTVSS
65	AB014VL	VL VEGF (seq. 1)	DIQMTQSPSSLSASVGDRVTITCSASQDISNYLNWYQQKP GKAPKVLIIYFTSSLHSGVPSRFSGSGSGTDFTLTISLQPE EDFATYYCQQYSTVPWTFGQGTKVEIKR
66	AB016VH	VH NRP1 (seq. 1)	EVQLVESGGGLVQPGGSLRLSCAASGFSFSSEPISWVRQA PGKGLEWVSSITGKNGYTTYADSVKGRFTISADTSKNTAY LQMNSLRAEDTAVYYCARWGKKVYGMDVWGQGTLVTVSS
67	AB016VL	VL NRP1 (seq. 1)	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLAWYQQKP GKAPKLLIYGASSRASGVPSRFSGSGSGTDFTLTISLQPE EDFATYYCQQYMSVPITFGQGTKVEIKR
68	AB017VH	VH TNF (seq. 1)	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQA PGKGLEWVSAITWNSGHIDYADSVTEGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTLVTVSS
69	AB017VL	VL TNF (seq. 1)	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKP GKAPKLLIYAASTLQSGVPSRFSGSGSGTDFTLTISLQPE EDVATYYCQRYNRAPYTFGQGTKVEIKR
70	AB050VH	VH SOST	EVQLQQSGPELMKPGASVKMSCASGYTFTDYNMHWMKQN QGKSLEWIGEINPNSGGSGYNQKFKGKATLTVDKSSSTAY MELRSLTSEDSAVYYCARLGYYGNYEDWYFDVWGAGTTVT VSS
71	AB050VL	VL SOST	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNWYQQKP DGTVKLLIFYTSTLQSGVPSRFSGSGSGTNYSLTITNLEQ DDAATYFCQQGDTLPYTFGGGTKLEIKR
72	AB229VH	VH TNF (D2E7) (seq. 2)	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQA PGKGLEWVSAITWNSGHIDYADSVTEGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGTLVTVSS

SEQ ID No.	ABT Unique ID	Protein region	Sequence
			1234567890123456789012345678901234567890
73	AB229VL	VL TNF (D2E7) (seq. 2)	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKP GKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISSLQP EDVATYYCARYNRAPYTFGQGTKVEIKR
74	AB230VH	VH TNF (D2E7.1) (seq. 3)	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQA PGKGLEWVSAITWNSGHIDYADSVTEGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCAKVAYLSTASSLDYWGQGTLLVTVS S
75	AB230VL	VL TNF (D2E7.1) (seq. 3)	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWYQQKP GKAPKLLIYAASLTQSGVPSRFSGSGSGTDFTLTISSLQP EDVATYYCARYNRAPYTFGQGTKVEIKR
76	AB231VH	VH IL-13 (13C5.5) (seq. 1)	EVTLRSGPGLVKPTQTLTLTCTLYGFSLSLSDMGVDWIR QPPGKGLEWLAHIWDDVKRYNPALKSRLTISKDTSKNQV VLKLTSVDPVDTATYYCARTVSSGYIYYAMDYWGQGTLLV VSS
77	AB231VL	VL IL-13 (13C5.5) (seq. 1)	DIQMTQSPSSLSASVGDRVTISCRASQDIRNYLNWYQQKP GKAPKLLIFYTSKLHSGVPSRFSGSGSGTDYTLTISSLQP EDIATYYCQQGNTLPLTFGGGTKVEIKR
78	AB232VH	VH IL-13 (13C5.5L3F) (seq. 2)	EVTLRSGPGLVKPTQTLTLTCTLYGFSLSLSDMGVDWIR QPPGKGLEWLAHIWDDVKRYNPALKSRLTISKDTSKNQV VLKLTSVDPVDTATYYCARTVSSGYIYYAMDYWGQGTLLV VSS
79	AB232VL	VL IL-13 (13C5.5L3F) (seq. 2)	DIQMTQSPSSLSASVGDRVTISCRASQDIRNYLNWYQQKP GKAPKLLIFYTSKLHSGVPSRFSGSGSGTDYTLTISSLQP EDIATYYCQQGLTPPLTFGGGTKVEIKR

Detailed description of specific DVD-Ig molecules capable of binding specific targets, and methods of making the same, is provided in the Examples section below.

C. Production of DVD proteins

Binding proteins of the present invention may be produced by any of a number of techniques known in the art. For example, expression from host cells, wherein expression vector(s) encoding the DVD heavy and DVD light chains is (are) transfected into a host cell by standard techniques. The various forms of the term "transfection" are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, *e.g.*, electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like. Although it is possible to express the DVD proteins of the invention in either prokaryotic or eukaryotic host cells, DVD proteins are expressed in eukaryotic cells, for example, mammalian host cells, because such eukaryotic cells (and in particular

mammalian cells) are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active DVD protein.

Exemplary mammalian host cells for expressing the recombinant antibodies of the invention include Chinese Hamster Ovary (CHO cells) (including dhfr- CHO cells, described in 5 Urlaub and Chasin, (1980) *Proc. Natl. Acad. Sci. USA* 77:4216-4220, used with a DHFR selectable marker, *e.g.*, as described in R.J. Kaufman and P.A. Sharp (1982) *Mol. Biol.* 159:601-621), NS0 myeloma cells, COS cells, SP2 and PER.C6 cells. When recombinant expression vectors encoding DVD proteins are introduced into mammalian host cells, the DVD proteins are produced by culturing the host cells for a period of time sufficient to allow for expression of the 10 DVD proteins in the host cells or secretion of the DVD proteins into the culture medium in which the host cells are grown. DVD proteins can be recovered from the culture medium using standard protein purification methods.

In an exemplary system for recombinant expression of DVD proteins of the invention, a recombinant expression vector encoding both the DVD heavy chain and the DVD light chain is 15 introduced into dhfr- CHO cells by calcium phosphate-mediated transfection. Within the recombinant expression vector, the DVD heavy and light chain genes are each operatively linked to CMV enhancer/AdMLP promoter regulatory elements to drive high levels of transcription of the genes. The recombinant expression vector also carries a DHFR gene, which allows for selection of CHO cells that have been transfected with the vector using methotrexate 20 selection/amplification. The selected transformant host cells are cultured to allow for expression of the DVD heavy and light chains and intact DVD protein is recovered from the culture medium. Standard molecular biology techniques are used to prepare the recombinant expression vector, transfect the host cells, select for transformants, culture the host cells and recover the DVD protein from the culture medium. Still further the invention provides a method of synthesizing a 25 DVD protein of the invention by culturing a host cell of the invention in a suitable culture medium until a DVD protein of the invention is synthesized. The method can further comprise isolating the DVD protein from the culture medium.

An important feature of DVD-Ig is that it can be produced and purified in a similar way as a conventional antibody. The production of DVD-Ig results in a homogeneous, single major 30 product with desired dual-specific activity, without any sequence modification of the constant region or chemical modifications of any kind. Other previously described methods to generate “bi-specific”, “multi-specific”, and “multi-specific multivalent” full length binding proteins do not lead to a single primary product but instead lead to the intracellular or secreted production of a mixture of assembled inactive, mono-specific, multi-specific, multivalent, full length binding 35 proteins, and multivalent full length binding proteins with combination of different binding sites. As an example, based on the design described by Miller and Presta (PCT publication

WO2001/077342(A1), there are 16 possible combinations of heavy and light chains.

Consequently only 6.25% of protein is likely to be in the desired active form, and not as a single major product or single primary product compared to the other 15 possible combinations.

Separation of the desired, fully active forms of the protein from inactive and partially active forms of the protein using standard chromatography techniques, typically used in large scale manufacturing, is yet to be demonstrated.

Surprisingly the design of the “dual-specific multivalent full length binding proteins” of the present invention leads to a dual variable domain light chain and a dual variable domain heavy chain which assemble primarily to the desired “dual-specific multivalent full length binding proteins”.

At least 50%, at least 75% and at least 90% of the assembled, and expressed dual variable domain immunoglobulin molecules are the desired dual-specific tetravalent protein. This aspect of the invention particularly enhances the commercial utility of the invention. Therefore, the present invention includes a method to express a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a single primary product of a “dual-specific tetravalent full length binding protein”.

The present invention provides a methods of expressing a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a “primary product” of a “dual-specific tetravalent full length binding protein”, where the “primary product” is more than 50% of all assembled protein, comprising a dual variable domain light chain and a dual variable domain heavy chain.

The present invention provides methods of expressing a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a single “primary product” of a “dual-specific tetravalent full length binding protein”, where the “primary product” is more than 75% of all assembled protein, comprising a dual variable domain light chain and a dual variable domain heavy chain.

The present invention provides methods of expressing a dual variable domain light chain and a dual variable domain heavy chain in a single cell leading to a single “primary product” of a “dual-specific tetravalent full length binding protein”, where the “primary product” is more than 90% of all assembled protein, comprising a dual variable domain light chain and a dual variable domain heavy chain.

II. Derivatized DVD binding proteins:

One embodiment provides a labeled binding protein wherein the binding protein of the invention is derivatized or linked to another functional molecule (e.g., another peptide or protein).

For example, a labeled binding protein of the invention can be derived by functionally linking an binding protein of the invention (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (e.g., a bispecific antibody or a diabody), a detectable agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the binding protein with another molecule (such as a streptavidin core region or a polyhistidine tag).

Useful detectable agents with which a binding protein of the invention may be derivatized include fluorescent compounds. Exemplary fluorescent detectable agents include fluorescein, fluorescein isothiocyanate, rhodamine, 5-dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin and the like. A binding protein may also be derivatized with detectable enzymes, such as alkaline phosphatase, horseradish peroxidase, glucose oxidase and the like. When a binding protein is derivatized with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a detectable reaction product. For example, when the detectable agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. a binding protein may also be derivatized with biotin, and detected through indirect measurement of avidin or streptavidin binding.

Another embodiment of the invention provides a crystallized binding protein and formulations and compositions comprising such crystals. In one embodiment the crystallized binding protein has a greater half-life *in vivo* than the soluble counterpart of the binding protein. In another embodiment the binding protein retains biological activity after crystallization.

Crystallized binding protein of the invention may be produced according to methods known in the art and as disclosed in WO 02072636, incorporated herein by reference.

Another embodiment of the invention provides a glycosylated binding protein wherein the antibody or antigen-binding portion thereof comprises one or more carbohydrate residues. Nascent *in vivo* protein production may undergo further processing, known as post-translational modification. In particular, sugar (glycosyl) residues may be added enzymatically, a process known as glycosylation. The resulting proteins bearing covalently linked oligosaccharide side chains are known as glycosylated proteins or glycoproteins. Antibodies are glycoproteins with one or more carbohydrate residues in the Fc domain, as well as the variable domain. Carbohydrate residues in the Fc domain have important effect on the effector function of the Fc domain, with minimal effect on antigen binding or half-life of the antibody (R. Jefferis, *Biotechnol. Prog.* **21** (2005), pp. 11–16). In contrast, glycosylation of the variable domain may have an effect on the antigen binding activity of the antibody. Glycosylation in the variable domain may have a negative effect on antibody binding affinity, likely due to steric hindrance

(Co, M.S., et al., Mol. Immunol. (1993) 30:1361- 1367), or result in increased affinity for the antigen (Wallick, S.C., et al., Exp. Med. (1988) 168:1099-1109; Wright, A., et al., EMBO J. (1991) 10:2717 2723).

One aspect of the present invention is directed to generating glycosylation site mutants in which the O- or N-linked glycosylation site of the binding protein has been mutated. One skilled in the art can generate such mutants using standard well-known technologies. Glycosylation site mutants that retain the biological activity but have increased or decreased binding activity are another object of the present invention.

In still another embodiment, the glycosylation of the antibody or antigen-binding portion of the invention is modified. For example, an aglycosylated antibody can be made (i.e., the antibody lacks glycosylation). Glycosylation can be altered to, for example, increase the affinity of the antibody for antigen. Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the antibody sequence. For example, one or more amino acid substitutions can be made that result in elimination of one or more variable region glycosylation sites to thereby eliminate glycosylation at that site. Such aglycosylation may increase the affinity of the antibody for antigen. Such an approach is described in further detail in PCT Publication WO2003016466A2, and U.S. Pat. Nos. 5,714,350 and 6,350,861, each of which is incorporated herein by reference in its entirety.

Additionally or alternatively, a modified binding protein of the invention can be made that has an altered type of glycosylation, such as a hypofucosylated antibody having reduced amounts of fucosyl residues (see Kanda, Yutaka et al., Journal of Biotechnology (2007), 130(3), 300-310.) or an antibody having increased bisecting GlcNAc structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Such carbohydrate modifications can be accomplished by, for example, expressing the antibody in a host cell with altered glycosylation machinery. Cells with altered glycosylation machinery have been described in the art and can be used as host cells in which to express recombinant antibodies of the invention to thereby produce an antibody with altered glycosylation. See, for example, Shields, R. L. et al. (2002) J. Biol. Chem. 277:26733-26740; Umana et al. (1999) Nat. Biotech. 17:176-1, as well as, European Patent No: EP 1,176,195; PCT Publications WO 03/035835; WO 99/54342 80, each of which is incorporated herein by reference in its entirety.

Protein glycosylation depends on the amino acid sequence of the protein of interest, as well as the host cell in which the protein is expressed. Different organisms may produce different glycosylation enzymes (eg., glycosyltransferases and glycosidases), and have different substrates (nucleotide sugars) available. Due to such factors, protein glycosylation pattern, and composition of glycosyl residues, may differ depending on the host system in which the particular protein is

expressed. Glycosyl residues useful in the invention may include, but are not limited to, glucose, galactose, mannose, fucose, n-acetylglucosamine and sialic acid. In an embodiment, the glycosylated binding protein comprises glycosyl residues such that the glycosylation pattern is human.

5 It is known to those skilled in the art that differing protein glycosylation may result in differing protein characteristics. For instance, the efficacy of a therapeutic protein produced in a microorganism host, such as yeast, and glycosylated utilizing the yeast endogenous pathway may be reduced compared to that of the same protein expressed in a mammalian cell, such as a CHO cell line. Such glycoproteins may also be immunogenic in humans and show reduced half-life *in*
10 *vivo* after administration. Specific receptors in humans and other animals may recognize specific glycosyl residues and promote the rapid clearance of the protein from the bloodstream. Other adverse effects may include changes in protein folding, solubility, susceptibility to proteases, trafficking, transport, compartmentalization, secretion, recognition by other proteins or factors, antigenicity, or allergenicity. Accordingly, a practitioner may choose a therapeutic protein with a
15 specific composition and pattern of glycosylation, for example glycosylation composition and pattern identical, or at least similar, to that produced in human cells or in the species-specific cells of the intended subject animal.

Expressing glycosylated proteins different from that of a host cell may be achieved by genetically modifying the host cell to express heterologous glycosylation enzymes. Using
20 techniques known in the art a practitioner may generate antibodies or antigen-binding portions thereof exhibiting human protein glycosylation. For example, yeast strains have been genetically modified to express non-naturally occurring glycosylation enzymes such that glycosylated proteins (glycoproteins) produced in these yeast strains exhibit protein glycosylation identical to that of animal cells, especially human cells (U.S patent applications 20040018590 and
25 20020137134 and PCT publication WO2005100584 A2).

In addition to the binding proteins, the present invention is also directed to anti-idiotypic (anti-Id) antibodies specific for such binding proteins of the invention. An anti-Id antibody is an antibody, which recognizes unique determinants generally associated with the antigen-binding region of another antibody. The anti-Id can be prepared by immunizing an animal with the
30 binding protein or a CDR containing region thereof. The immunized animal will recognize, and respond to the idiotypic determinants of the immunizing antibody and produce an anti-Id antibody. It is readily apparent that it may be easier to generate anti-idiotypic antibodies to the two or more parent antibodies incorporated into a DVD-Ig molecule; and confirm binding studies by methods well recognized in the art (e.g., BIAcore, ELISA) to verify that anti-idiotypic
35 antibodies specific for the idio type of each parent antibody also recognize the idio type (e.g., antigen binding site) in the context of the DVD-Ig. The anti-idiotypic antibodies specific for

each of the two or more antigen binding sites of a DVD-Ig provide ideal reagents to measure DVD-Ig concentrations of a human DVD-Ig in patient serum; DVD-Ig concentration assays can be established using a "sandwich assay ELISA format" with an antibody to a first antigen binding regions coated on the solid phase (e.g., BIAcore chip, ELISA plate etc.), rinsed with rinsing
 5 buffer, incubation with the serum sample, another rinsing step and ultimately incubation with another anti-idiotypic antibody to the another antigen binding site, itself labeled with an enzyme for quantitation of the binding reaction. In an embodiment, for a DVD-Ig with more than two different binding sites, anti-idiotypic antibodies to the two outermost binding sites (most distal and proximal from the constant region) will not only help in determining the DVD-Ig
 10 concentration in human serum but also document the integrity of the molecule *in vivo*. Each anti-Id antibody may also be used as an "immunogen" to induce an immune response in yet another animal, producing a so-called anti-anti-Id antibody.

Further, it will be appreciated by one skilled in the art that a protein of interest may be expressed using a library of host cells genetically engineered to express various glycosylation
 15 enzymes, such that member host cells of the library produce the protein of interest with variant glycosylation patterns. A practitioner may then select and isolate the protein of interest with particular novel glycosylation patterns. In an embodiment, the protein having a particularly selected novel glycosylation pattern exhibits improved or altered biological properties.

III. Uses of DVD-Ig

20 Given their ability to bind to two or more antigens the binding proteins of the invention can be used to detect the antigens (e.g., in a biological sample, such as serum or plasma), using a conventional immunoassay, such as an enzyme linked immunosorbent assays (ELISA), an radioimmunoassay (RIA) or tissue immunohistochemistry. The DVD-Ig is directly or indirectly labeled with a detectable substance to facilitate detection of the bound or unbound antibody.
 25 Suitable detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a
 30 luminescent material includes luminol; and examples of suitable radioactive material include ^3H , ^{14}C , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , or ^{153}Sm .

In an embodiment, the binding proteins of the invention are capable of neutralizing the activity of the antigens both *in vitro* and *in vivo*. Accordingly, such DVD-Igs can be used to
 35 inhibit antigen activity, e.g., in a cell culture containing the antigens, in human subjects or in other

mammalian subjects having the antigens with which a binding protein of the invention cross-reacts. In another embodiment, the invention provides a method for reducing antigen activity in a subject suffering from a disease or disorder in which the antigen activity is detrimental. A binding protein of the invention can be administered to a human subject for therapeutic purposes.

5 As used herein, the term "a disorder in which antigen activity is detrimental" is intended to include diseases and other disorders in which the presence of the antigen in a subject suffering from the disorder has been shown to be or is suspected of being either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which antigen activity is detrimental is a disorder in which reduction of
 10 antigen activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of the antigen in a biological fluid of a subject suffering from the disorder (*e.g.*, an increase in the concentration of antigen in serum, plasma, synovial fluid, *etc.* of the subject). Non-limiting examples of disorders that can be treated with the binding proteins of the invention include those disorders discussed
 15 below and in the section pertaining to pharmaceutical compositions of the antibodies of the invention.

The DVD-Igs of the invention may bind one antigen or multiple antigens. Such antigens include, but are not limited to, the targets listed in the following databases, which databases are incorporated herein by reference. These target databases include those listings:

- 20 Therapeutic targets (<http://xin.cz3.nus.edu.sg/group/cjttd/ttd.asp>);
- Cytokines and cytokine receptors (<http://www.cytokinewebfacts.com/>,
<http://www.copewithcytokines.de/cope.cgi>, and
http://cmbi.bjmu.edu.cn/cmbidata/cgf/CGF_Database/cytokine.medic.kumamoto-u.ac.jp/CFC/indexR.html);
- 25 Chemokines (<http://cytokine.medic.kumamoto-u.ac.jp/CFC/CK/Chemokine.html>);
- Chemokine receptors and GPCRs (<http://csp.medic.kumamoto-u.ac.jp/CSP/Receptor.html>,
<http://www.gpcr.org/7tm/>);
- Olfactory Receptors (<http://senselab.med.yale.edu/senselab/ORDB/default.asp>);
- Receptors (<http://www.iuphar-db.org/iuphar-rd/list/index.htm>);
- 30 Cancer targets (<http://cged.hgc.jp/cgi-bin/input.cgi>);
- Secreted proteins as potential antibody targets (<http://spd.cbi.pku.edu.cn/>);
- Protein kinases (<http://spd.cbi.pku.edu.cn/>), and

Human CD markers (http://content.labvelocity.com/tools/6/1226/CD_table_final_locked.pdf) and (Zola H, 2005 CD molecules 2005: human cell differentiation molecules Blood, 106:3123-6).

DVD-Igs are useful as therapeutic agents to simultaneously block two different targets to enhance efficacy/safety and/or increase patient coverage. Such targets may include soluble
5 targets (TNF) and cell surface receptor targets (VEGFR and EGFR). It can also be used to induce redirected cytotoxicity between tumor cells and T cells (Her2 and CD3) for cancer therapy, or between autoreactive cell and effector cells for autoimmune disease or transplantation, or between any target cell and effector cell to eliminate disease-causing cells in any given disease.

In addition, DVD-Ig can be used to trigger receptor clustering and activation when it is
10 designed to target two different epitopes on the same receptor. This may have benefit in making agonistic and antagonistic anti-GPCR therapeutics. In this case, DVD-Ig can be used to target two different epitopes (including epitopes on both the loop regions and the extracellular domain) on one cell for clustering/signaling (two cell surface molecules) or signaling (on one molecule). Similarly, a DVD-Ig molecule can be designed to trigger CTLA-4 ligation, and a negative signal
15 by targeting two different epitopes (or 2 copies of the same epitope) of CTLA-4 extracellular domain, leading to down regulation of the immune response. CTLA-4 is a clinically validated target for therapeutic treatment of a number of immunological disorders. CTLA-4/B7 interactions negatively regulate T cell activation by attenuating cell cycle progression, IL-2 production, and proliferation of T cells following activation, and CTLA-4 (CD152) engagement can down-
20 regulate T cell activation and promote the induction of immune tolerance. However, the strategy of attenuating T cell activation by agonistic antibody engagement of CTLA-4 has been unsuccessful since CTLA-4 activation requires ligation. The molecular interaction of CTLA-4/B7 is in “skewed zipper” arrays, as demonstrated by crystal structural analysis (Stamper 2001 Nature 410:608). However none of the currently available CTLA-4 binding reagents have ligation
25 properties, including anti-CTLA-4 mAbs. There have been several attempts to address this issue. In one case, a cell member-bound single chain antibody was generated, and significantly inhibited allogeneic rejection in mice (Hwang 2002 JI 169:633). In a separate case, artificial APC surface-linked single-chain antibody to CTLA-4 was generated and demonstrated to attenuate T cell responses (Griffin 2000 JI 164:4433). In both cases, CTLA-4 ligation was achieved by closely
30 localized member-bound antibodies in artificial systems. While these experiments provide proof-of-concept for immune down-regulation by triggering CTLA-4 negative signaling, the reagents used in these reports are not suitable for therapeutic use. To this end, CTLA-4 ligation may be achieved by using a DVD-Ig molecule, which target two different epitopes (or 2 copies of the same epitope) of CTLA-4 extracellular domain. The rationale is that the distance spanning two
35 binding sites of an IgG, approximately 150-170Å, is too large for active ligation of CTLA-4 (30-50 Å between 2 CTLA-4 homodimer). However the distance between the two binding sites on

DVD-Ig (one arm) is much shorter, also in the range of 30-50 Å, allowing proper ligation of CTLA-4.

Similarly, DVD-Ig can target two different members of a cell surface receptor complex (e.g., IL-12R alpha and beta). Furthermore, DVD-Ig can target CR1 and a soluble
5 protein/pathogen to drive rapid clearance of the target soluble protein/pathogen.

Additionally, DVD-Igs of the invention can be employed for tissue-specific delivery (target a tissue marker and a disease mediator for enhanced local PK thus higher efficacy and/or lower toxicity), including intracellular delivery (targeting an internalizing receptor and a intracellular molecule), delivering to inside brain (targeting transferrin receptor and a CNS disease
10 mediator for crossing the blood-brain barrier). DVD-Ig can also serve as a carrier protein to deliver an antigen to a specific location via binding to a non-neutralizing epitope of that antigen and also to increase the half-life of the antigen. Furthermore, DVD-Ig can be designed to either be physically linked to medical devices implanted into patients or target these medical devices (see Burke, Sandra E.; Kuntz, Richard E.; Schwartz, Lewis B., Zotarolimus eluting stents.

15 Advanced Drug Delivery Reviews (2006), 58(3), 437-446; Surface coatings for biological activation and functionalization of medical devices, Hildebrand, H. F.; Blanchemain, N.; Mayer, G.; Chai, F.; Lefebvre, M.; Boschini, F., Surface and Coatings Technology (2006), 200(22-23), 6318-6324; Drug/ device combinations for local drug therapies and infection prophylaxis, Wu, Peng; Grainger, David W., Biomaterials (2006), 27(11), 2450-2467; Mediation of the cytokine
20 network in the implantation of orthopedic devices., Marques, A. P.; Hunt, J. A.; Reis, Rui L., Biodegradable Systems in Tissue Engineering and Regenerative Medicine (2005), 377-397).

Briefly, directing appropriate types of cell to the site of medical implant may promote healing and restoring normal tissue function. Alternatively, inhibition of mediators (including but not limited to cytokines), released upon device implantation by a DVD coupled to or target to a device is also

25 provided. For example, Stents have been used for years in interventional cardiology to clear blocked arteries and to improve the flow of blood to the heart muscle. However, traditional bare metal stents have been known to cause restenosis (re-narrowing of the artery in a treated area) in some patients and can lead to blood clots. Recently, an anti-CD34 antibody coated stent has been described which reduced restenosis and prevents blood clots from occurring by capturing
30 endothelial progenitor cells (EPC) circulating throughout the blood. Endothelial cells are cells that line blood vessels, allowing blood to flow smoothly. The EPCs adhere to the hard surface of the stent forming a smooth layer that not only promotes healing but prevents restenosis and blood clots, complications previously associated with the use of stents (Aoji et al. 2005 J Am Coll Cardiol. 45(10):1574-9). In addition to improving outcomes for patients requiring stents, there
35 are also implications for patients requiring cardiovascular bypass surgery. For example, a prosthetic vascular conduit (artificial artery) coated with anti-EPC antibodies would eliminate the

need to use arteries from patients legs or arms for bypass surgery grafts. This would reduce surgery and anesthesia times, which in turn will reduce coronary surgery deaths. DVD-Ig are designed in such a way that it binds to a cell surface marker (such as CD34) as well as a protein (or an epitope of any kind, including but not limited to proteins, lipids and polysaccharides) that has been coated on the implanted device to facilitate the cell recruitment. Such approaches can also be applied to other medical implants in general. Alternatively, DVD-Igs can be coated on medical devices and upon implantation and releasing all DVDs from the device (or any other need which may require additional fresh DVD-Ig, including aging and denaturation of the already loaded DVD-Ig) the device could be reloaded by systemic administration of fresh DVD-Ig to the patient, where the DVD-Ig is designed to binds to a target of interest (a cytokine, a cell surface marker (such as CD34) etc.) with one set of binding sites and to a target coated on the device (including a protein, an epitope of any kind, including but not limited to lipids, polysaccharides and polymers) with the other. This technology has the advantage of extending the usefulness of coated implants.

15 **A. Use of DVD-Igs in various diseases**

DVD-Ig molecules of the invention are also useful as therapeutic molecules to treat various diseases. Such DVD molecules may bind one or more targets involved in a specific disease. Examples of such targets in various diseases are described below.

1. **Human Autoimmune and Inflammatory Response**

Many proteins have been implicated in general autoimmune and inflammatory responses, including C5, CCL1 (I-309), CCL11 (eotaxin), CCL13 (mcp-4), CCL15 (MIP-1d), CCL16 (HCC-4), CCL17 (TARC), CCL18 (PARC), CCL19, CCL2 (mcp-1), CCL20 (MIP-3a), CCL21 (MIP-2), CCL23 (MPIF-1), CCL24 (MPIF-2 / eotaxin-2), CCL25 (TECK), CCL26, CCL3 (MIP-1a), CCL4 (MIP-1b), CCL5 (RANTES), CCL7 (mcp-3), CCL8 (mcp-2), CXCL1, CXCL10 (IP-10), CXCL11 (I-TAC / IP-9), CXCL12 (SDF1), CXCL13, CXCL14, CXCL2, CXCL3, CXCL5 (ENA-78 / LIX), CXCL6 (GCP-2), CXCL9, IL13, IL8, CCL13 (mcp-4), CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CX3CR1, IL8RA, XCR1 (CCXCR1), IFNA2, IL10, IL13, IL17C, IL1A, IL1B, IL1F10, IL1F5, IL1F6, IL1F7, IL1F8, IL1F9, IL22, IL5, IL8, IL9, LTA, LTB, MIF, SCYE1 (endothelial Monocyte-activating cytokine), SPP1, TNF, TNFSF5, IFNA2, IL10RA, IL10RB, IL13, IL13RA1, IL5RA, IL9, IL9R, ABCF1, BCL6, C3, C4A, CEBPB, CRP, ICEBERG, IL1R1, IL1RN, IL8RB, LTB4R, TOLLIP, FADD, IRAK1, IRAK2, MYD88, NCK2, TNFAIP3, TRADD, TRAF1, TRAF2, TRAF3, TRAF4, TRAF5, TRAF6, ACVR1, ACVR1B, ACVR2, ACVR2B, ACVRL1, CD28, CD3E, CD3G, CD3Z, CD69, CD80, CD86, CNR1, CTLA4, CYSLTR1, FCER1A, FCER2, FCGR3A, GPR44, HAVCR2, OPRD1, P2RX7, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, BLR1, CCL1, CCL2,

CCL3, CCL4, CCL5, CCL7, CCL8, CCL11, CCL13, CCL15, CCL16, CCL17, CCL18, CCL19, CCL20, CCL21, CCL22, CCL23, CCL24, CCL25, CCR1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CX3CL1, CX3CR1, CXCL1, CXCL2, CXCL3, CXCL5, CXCL6, CXCL10, CXCL11, CXCL12, CXCL13, CXCR4, GPR2, SCYE1, SDF2, XCL1, XCL2, XCR1, AMH, AMHR2, BMPR1A, BMPR1B, BMPR2, C19orf10 (IL27w), CER1, CSF1, CSF2, CSF3, DKFZp451J0118, FGF2, GFI1, IFNA1, IFNB1, IFNG, IGF1, IL1A, IL1B, IL1R1, IL1R2, IL2, IL2RA, IL2RB, IL2RG, IL3, IL4, IL4R, IL5, IL5RA, IL6, IL6R, IL6ST, IL7, IL8, IL8RA, IL8RB, IL9, IL9R, IL10, IL10RA, IL10RB, IL11, IL11RA, IL12A, IL12B, IL12RB1, IL12RB2, IL13, IL13RA1, IL13RA2, IL15, IL15RA, IL16, IL17, IL17R, IL18, IL18R1, IL19, IL20, KITLG, LEP, LTA, LTB, LTB4R, LTB4R2, LTBR, MIF, NPPB, PDGFB, TBX21, TDGF1, TGFA, TGFB1, TGFB1I1, TGFB2, TGFB3, TGFBI, TGFBR1, TGFBR2, TGFBR3, TH1L, TNF, TNFRSF1A, TNFRSF1B, TNFRSF7, TNFRSF8, TNFRSF9, TNFRSF11A, TNFRSF21, TNFSF4, TNFSF5, TNFSF6, TNFSF11, VEGF, ZFPM2, and RNF110 (ZNF144). In one aspect, DVD-Igs capable of binding one or more of the targets listed herein are provided.

2. Asthma

Allergic asthma is characterized by the presence of eosinophilia, goblet cell metaplasia, epithelial cell alterations, airway hyperreactivity (AHR), and Th2 and Th1 cytokine expression, as well as elevated serum IgE levels. It is now widely accepted that airway inflammation is the key factor underlying the pathogenesis of asthma, involving a complex interplay of inflammatory cells such as T cells, B cells, eosinophils, mast cells and macrophages, and of their secreted mediators including cytokines and chemokines. Corticosteroids are the most important anti-inflammatory treatment for asthma today, however their mechanism of action is non-specific and safety concerns exist, especially in the juvenile patient population. The development of more specific and targeted therapies is therefore warranted. There is increasing evidence that IL-13 in mice mimics many of the features of asthma, including AHR, mucus hypersecretion and airway fibrosis, independently of eosinophilic inflammation (Finotto et al., *International Immunology* (2005), 17(8), 993-1007; Padilla et al., *Journal of Immunology* (2005), 174(12), 8097-8105).

IL-13 has been implicated as having a pivotal role in causing pathological responses associated with asthma. The development of anti-IL-13 mAb therapy to reduce the effects of IL-13 in the lung is an exciting new approach that offers considerable promise as a novel treatment for asthma. However other mediators of differential immunological pathways are also involved in asthma pathogenesis, and blocking these mediators, in addition to IL-13, may offer additional therapeutic benefit. Such target pairs include, but are not limited to, IL-13 and a pro-inflammatory cytokine, such as tumor necrosis factor- α (TNF- α). TNF- α may amplify the inflammatory response in asthma and may be linked to disease severity (McDonnell, et al., *Progress in Respiratory Research* (2001), 31(New Drugs for Asthma, Allergy and COPD), 247-

250.). This suggests that blocking both IL-13 and TNF- α may have beneficial effects, particularly in severe airway disease. In another embodiment the DVD-Ig of the invention binds the targets IL-13 and TNF α and is used for treating asthma.

Animal models such as OVA-induced asthma mouse model, where both inflammation
 5 and AHR can be assessed, are known in the art and may be used to determine the ability of various DVD-Ig molecules to treat asthma. Animal models for studying asthma are disclosed in Coffman, et al., *Journal of Experimental Medicine* (2005), 201(12), 1875-1879; Lloyd, et al., *Advances in Immunology* (2001), 77, 263-295; Boyce et al., *Journal of Experimental Medicine* (2005), 201(12), 1869-1873; and Snibson, et al., *Journal of the British Society for Allergy and*
 10 *Clinical Immunology* (2005), 35(2), 146-52. In addition to routine safety assessments of these target pairs specific tests for the degree of immunosuppression may be warranted and helpful in selecting the best target pairs (see Luster et al., *Toxicology* (1994), 92(1-3), 229-43; Descotes, et al., *Developments in biological standardization* (1992), 77 99-102; Hart et al., *Journal of Allergy and Clinical Immunology* (2001), 108(2), 250-257).

15 Based on the rationale disclosed herein and using the same evaluation model for efficacy and safety other pairs of targets that DVD-Ig molecules can bind and be useful to treat asthma may be determined. In an embodiment, such targets include, but are not limited to, IL-13 and IL-1beta, since IL-1beta is also implicated in inflammatory response in asthma; IL-13 and cytokines and chemokines that are involved in inflammation, such as IL-13 and IL-9; IL-13 and IL-4; IL-13
 20 and IL-5; IL-13 and IL-25; IL-13 and TARC; IL-13 and MDC; IL-13 and MIF; IL-13 and TGF- β ; IL-13 and LHR agonist; IL-13 and CL25; IL-13 and SPRR2a; IL-13 and SPRR2b; and IL-13 and ADAM8. The present invention also provides DVD-Igs capable of binding one or more targets involved in asthma selected from the group consisting of CSF1 (MCSF), CSF2 (GM-CSF), CSF3 (GCSF), FGF2, IFNA1, IFNB1, IFNG, histamine and histamine receptors, IL1A, IL1B, IL2, IL3,
 25 IL4, IL5, IL6, IL7, IL8, IL9, IL10, IL11, IL12A, IL12B, IL13, IL14, IL15, IL16, IL17, IL18, IL19, KITLG, PDGFB, IL2RA, IL4R, IL5RA, IL8RA, IL8RB, IL12RB1, IL12RB2, IL13RA1, IL13RA2, IL18R1, TSLP, CCL1, CCL2, CCL3, CCL4, CCL5, CCL7, CCL8, CCL13, CCL17, CCL18, CCL19, CCL20, CCL22, CCL24, CX3CL1, CXCL1, CXCL2, CXCL3, XCL1, CCR2, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CX3CR1, GPR2, XCR1, FOS, GATA3, JAK1,
 30 JAK3, STAT6, TBX21, TGFB1, TNF, TNFSF6, YY1, CYSLTR1, FCER1A, FCER2, LTB4R, TB4R2, LTBR, and Chitinase.

3. Rheumatoid arthritis

Rheumatoid arthritis (RA), a systemic disease, is characterized by a chronic inflammatory
 reaction in the synovium of joints and is associated with degeneration of cartilage and erosion of
 35 juxta-articular bone. Many pro-inflammatory cytokines including TNF, chemokines, and growth

factors are expressed in diseased joints. Systemic administration of anti-TNF antibody or sTNFR fusion protein to mouse models of RA was shown to be anti-inflammatory and joint protective. Clinical investigations in which the activity of TNF in RA patients was blocked with intravenously administered infliximab (Harriman G, Harper LK, Schaible TF. 1999 Summary of clinical trials in rheumatoid arthritis using infliximab, an anti-TNF α treatment. *Ann Rheum Dis* 58 Suppl 1:I61-4), a chimeric anti-TNF mAb, has provided evidence that TNF regulates IL-6, IL-8, MCP-1, and VEGF production, recruitment of immune and inflammatory cells into joints, angiogenesis, and reduction of blood levels of matrix metalloproteinases-1 and -3. A better understanding of the inflammatory pathway in rheumatoid arthritis has led to identification of other therapeutic targets involved in rheumatoid arthritis. Promising treatments such as interleukin-6 antagonists (IL-6 receptor antibody MRA, developed by Chugai, Roche (see Nishimoto, Norihiro et al., *Arthritis & Rheumatism* (2004), 50(6), 1761-1769), CTLA4Ig (abatacept, Genovese Mc et al 2005 Abatacept for rheumatoid arthritis refractory to tumor necrosis factor alpha inhibition. *N Engl J Med*. 353:1114-23.), and anti-B cell therapy (rituximab, Okamoto H, Kamatani N. 2004 Rituximab for rheumatoid arthritis. *N Engl J Med*. 351:1909) have already been tested in randomized controlled trials over the past year. Other cytokines have been identified and have been shown to be of benefit in animal models, including interleukin-15 (therapeutic antibody HuMax-IL_15, AMG 714 see Baslund, Bo et al., *Arthritis & Rheumatism* (2005), 52(9), 2686-2692), interleukin-17, and interleukin-18, and clinical trials of these agents are currently under way. Dual-specific antibody therapy, combining anti-TNF and another mediator, has great potential in enhancing clinical efficacy and/or patient coverage. For example, blocking both TNF and VEGF can potentially eradicate inflammation and angiogenesis, both of which are involved in pathophysiology of RA. Blocking other pairs of targets involved in RA including, but not limited to, TNF and IL-18; TNF and IL-12; TNF and IL-23; TNF and IL-1 β ; TNF and MIF; TNF and IL-17; TNF and IL-15, TNF and SOST with specific DVD Igs is also contemplated. In addition to routine safety assessments of these target pairs, specific tests for the degree of immunosuppression may be warranted and helpful in selecting the best target pairs (see Luster et al., *Toxicology* (1994), 92(1-3), 229-43; Descotes, et al., *Developments in biological standardization* (1992), 77 99-102; Hart et al., *Journal of Allergy and Clinical Immunology* (2001), 108(2), 250-257). Whether a DVD Ig molecule will be useful for the treatment of rheumatoid arthritis can be assessed using pre-clinical animal RA models such as the collagen-induced arthritis mouse model. Other useful models are also well known in the art (see Brand DD., *Comp Med*. (2005) 55(2):114-22). Based on the cross-reactivity of the parental antibodies for human and mouse orthologues (e.g., reactivity for human and mouse TNF, human and mouse IL-15 etc.) validation studies in the mouse CIA model may be conducted with "matched surrogate antibody" derived DVD-Ig molecules; briefly, a DVD-Ig based on two (or more) mouse target specific antibodies may be matched to the extent possible to the characteristics of the parental

human or humanized antibodies used for human DVD-Ig construction (similar affinity, similar neutralization potency, similar half-life etc.).

4. SLE

The immunopathogenic hallmark of SLE is the polyclonal B cell activation, which leads to hyperglobulinemia, autoantibody production and immune complex formation. The fundamental abnormality appears to be the failure of T cells to suppress the forbidden B cell clones due to generalized T cell dysregulation. In addition, B and T-cell interaction is facilitated by several cytokines such as IL-10 as well as co-stimulatory molecules such as CD40 and CD40L, B7 and CD28 and CTLA-4, which initiate the second signal. These interactions together with impaired phagocytic clearance of immune complexes and apoptotic material, perpetuate the immune response with resultant tissue injury. The following targets may be involved in SLE and can potentially be used for DVD-Ig approach for therapeutic intervention: B cell targeted therapies: CD-20, CD-22, CD-19, CD28, CD4, CD80, HLA-DRA, IL10, IL2, IL4, TNFRSF5, TNFRSF6, TNFSF5, TNFSF6, BLR1, HDAC4, HDAC5, HDAC7A, HDAC9, ICOSL, IGBP1, MS4A1, RGS1, SLA2, CD81, IFNB1, IL10, TNFRSF5, TNFRSF7, TNFSF5, AICDA, BLNK, GALNAC4S-6ST, HDAC4, HDAC5, HDAC7A, HDAC9, IL10, IL11, IL4, INHA, INHBA, KLF6, TNFRSF7, CD28, CD38, CD69, CD80, CD83, CD86, DPP4, FCER2, IL2RA, TNFRSF8, TNFSF7, CD24, CD37, CD40, CD72, CD74, CD79A, CD79B, CR2, IL1R2, ITGA2, ITGA3, MS4A1, ST6GAL1, CD1C, CHST10, HLA-A, HLA-DRA, and NT5E.; co-stimulatory signals: CTLA4 or B7.1/B7.2; inhibition of B cell survival: BlyS, BAFF; Complement inactivation: C5; Cytokine modulation: the key principle is that the net biologic response in any tissue is the result of a balance between local levels of proinflammatory or anti-inflammatory cytokines (see Sfikakis PP et al 2005 Curr Opin Rheumatol 17:550-7). SLE is considered to be a Th-2 driven disease with documented elevations in serum IL-4, IL-6, IL-10. DVD Igs capable of binding one or more targets selected from the group consisting of IL-4, IL-6, IL-10, IFN- α , and TNF- α are also contemplated. Combination of targets discussed herein will enhance therapeutic efficacy for SLE which can be tested in a number of lupus preclinical models (see Peng SL (2004) Methods Mol Med.;102:227-72). Based on the cross-reactivity of the parental antibodies for human and mouse orthologues (e.g., reactivity for human and mouse CD20, human and mouse Interferon alpha etc.) validation studies in a mouse lupus model may be conducted with "matched surrogate antibody" derived DVD-Ig molecules; briefly, a DVD-Ig based two (or more) mouse target specific antibodies may be matched to the extent possible to the characteristics of the parental human or humanized antibodies used for human DVD-Ig construction (similar affinity, similar neutralization potency, similar half-life etc.).

5. Multiple sclerosis

Multiple sclerosis (MS) is a complex human autoimmune-type disease with a predominantly unknown etiology. Immunologic destruction of myelin basic protein (MBP) throughout the nervous system is the major pathology of multiple sclerosis. MS is a disease of complex pathologies, which involves infiltration by CD4+ and CD8+ T cells and of response within the central nervous system. Expression in the CNS of cytokines, reactive nitrogen species and costimulator molecules have all been described in MS. Of major consideration are immunological mechanisms that contribute to the development of autoimmunity. In particular, antigen expression, cytokine and leukocyte interactions, and regulatory T-cells, which help balance/modulate other T-cells such as Th1 and Th2 cells, are important areas for therapeutic target identification.

IL-12 is a proinflammatory cytokine that is produced by APC and promotes differentiation of Th1 effector cells. IL-12 is produced in the developing lesions of patients with MS as well as in EAE-affected animals. Previously it was shown that interference in IL-12 pathways effectively prevents EAE in rodents, and that in vivo neutralization of IL-12p40 using a anti-IL-12 mAb has beneficial effects in the myelin-induced EAE model in common marmosets.

TWEAK is a member of the TNF family, constitutively expressed in the central nervous system (CNS), with pro-inflammatory, proliferative or apoptotic effects depending upon cell types. Its receptor, Fn14, is expressed in CNS by endothelial cells, reactive astrocytes and neurons. TWEAK and Fn14 mRNA expression increased in spinal cord during experimental autoimmune encephalomyelitis (EAE). Anti-TWEAK antibody treatment in myelin oligodendrocyte glycoprotein (MOG) induced EAE in C57BL/6 mice resulted in a reduction of disease severity and leukocyte infiltration when mice were treated after the priming phase.

One aspect of the invention pertains to DVD Ig molecules capable of binding one or more, for example two, targets selected from the group consisting of IL-12, TWEAK, IL-23, CXCL13, CD40, CD40L, IL-18, VEGF, VLA-4, TNF, CD45RB, CD200, IFN γ , GM-CSF, FGF, C5, CD52, and CCR2. An embodiment includes a dual-specific anti-IL-12/TWEAK DVD Ig as a therapeutic agent beneficial for the treatment of MS.

Several animal models for assessing the usefulness of the DVD molecules to treat MS are known in the art (see Steinman L, et al., (2005) Trends Immunol. 26(11):565-71; Lublin FD., et al., (1985) Springer Semin Immunopathol.8(3):197-208; Genain CP, et al., (1997) J Mol Med. 75(3):187-97; Tuohy VK, et al., (1999) J Exp Med. 189(7):1033-42; Owens T, et al., (1995) Neurol Clin.13(1):51-73; and 't Hart BA, et al., (2005) J Immunol 175(7):4761-8. Based on the cross-reactivity of the parental antibodies for human and animal species orthologues (e.g., reactivity for human and mouse IL-12, human and mouse TWEAK etc.) validation studies in

the mouse EAE model may be conducted with “matched surrogate antibody” derived DVD-Ig molecules; briefly, a DVD-Ig based on to (or more) mouse target specific antibodies may be matched to the extent possible to the characteristics of the parental human or humanized antibodies used for human DVD-Ig construction (similar affinity, similar neutralization potency, similar half-life etc.). The same concept applies to animal models in other non-rodent species, where a “matched surrogate antibody” derived DVD-Ig would be selected for the anticipated pharmacology and possibly safety studies. In addition to routine safety assessments of these target pairs specific tests for the degree of immunosuppression may be warranted and helpful in selecting the best target pairs (see Luster et al., Toxicology (1994), 92(1-3), 229-43; Descotes, et al., Developments in biological standardization (1992), 77 99-102; Jones R. 2000 Rovelizumab (ICOS Corp). IDrugs.3(4):442-6).

6. Sepsis

The pathophysiology of sepsis is initiated by the outer membrane components of both gram-negative organisms (lipopolysaccharide [LPS], lipid A, endotoxin) and gram-positive organisms (lipoteichoic acid, peptidoglycan). These outer membrane components are able to bind to the CD14 receptor on the surface of monocytes. By virtue of the recently described toll-like receptors, a signal is then transmitted to the cell, leading to the eventual production of the proinflammatory cytokines tumor necrosis factor-alpha (TNF-alpha) and interleukin-1 (IL-1). Overwhelming inflammatory and immune responses are essential features of septic shock and play a central part in the pathogenesis of tissue damage, multiple organ failure, and death induced by sepsis. Cytokines, especially tumor necrosis factor (TNF) and interleukin (IL-1), have been shown to be critical mediators of septic shock. These cytokines have a direct toxic effect on tissues; they also activate phospholipase A2. These and other effects lead to increased concentrations of platelet-activating factor, promotion of nitric oxide synthase activity, promotion of tissue infiltration by neutrophils, and promotion of neutrophil activity.

The treatment of sepsis and septic shock remains a clinical conundrum, and recent prospective trials with biological response modifiers (i.e. anti-TNF, anti-MIF) aimed at the inflammatory response have shown only modest clinical benefit. Recently, interest has shifted toward therapies aimed at reversing the accompanying periods of immune suppression. Studies in experimental animals and critically ill patients have demonstrated that increased apoptosis of lymphoid organs and some parenchymal tissues contribute to this immune suppression, anergy, and organ system dysfunction. During sepsis syndromes, lymphocyte apoptosis can be triggered by the absence of IL-2 or by the release of glucocorticoids, granzymes, or the so-called 'death' cytokines: tumor necrosis factor alpha or Fas ligand. Apoptosis proceeds via auto-activation of cytosolic and/or mitochondrial caspases, which can be influenced by the pro- and anti-apoptotic members of the Bcl-2 family. In experimental animals, not only can treatment with inhibitors of

apoptosis prevent lymphoid cell apoptosis; it may also improve outcome. Although clinical trials with anti-apoptotic agents remain distant due in large part to technical difficulties associated with their administration and tissue targeting, inhibition of lymphocyte apoptosis represents an attractive therapeutic target for the septic patient. Likewise, a dual-specific agent targeting both inflammatory mediator and a apoptotic mediator, may have added benefit. One aspect of the invention pertains to DVD Igs capable of binding one or more targets involved in sepsis, in an embodiment two targets, selected from the group consisting TNF, IL-1, MIF, IL-6, IL-8, IL-18, IL-12, IL-23, FasL, LPS, Toll-like receptors, TLR-4, tissue factor, MIP-2, ADORA2A, CASP1, CASP4, IL-10, IL-1B, NFkB1, PROC, TNFRSF1A, CSF3, CCR3, IL1RN, MIF, NFkB1, PTAFR, TLR2, TLR4, GPR44, HMOX1, midkine, IRAK1, NFkB2, SERPINA1, SERPINE1, and TREM1. The efficacy of such DVD Igs for sepsis can be assessed in preclinical animal models known in the art (see Buras JA, et al.,(2005) Nat Rev Drug Discov. 4(10):854-65 and Calandra T, et al., (2000) Nat Med. 6(2):164-70).

7. Neurological disorders

7.1. Neurodegenerative Diseases

Chronic neurodegenerative diseases are usually age-dependent diseases characterized by progressive loss of neuronal functions (neuronal cell death, demyelination), loss of mobility and loss of memory. Emerging knowledge of the mechanisms underlying chronic neurodegenerative diseases (e.g., Alzheimer's disease) show a complex etiology and a variety of factors have been recognized to contribute to their development and progression e.g., age, glycemic status, amyloid production and multimerization, accumulation of advanced glycation-end products (AGE) which bind to their receptor RAGE (receptor for AGE), increased brain oxidative stress, decreased cerebral blood flow, neuroinflammation including release of inflammatory cytokines and chemokines, neuronal dysfunction and microglial activation. Thus these chronic neurodegenerative diseases represent a complex interaction between multiple cell types and mediators. Treatment strategies for such diseases are limited and mostly constitute either blocking inflammatory processes with non-specific anti-inflammatory agents (e.g., corticosteroids, COX inhibitors) or agents to prevent neuron loss and/or synaptic functions. These treatments fail to stop disease progression. Recent studies suggest that more targeted therapies such as antibodies to soluble A-b peptide (including the A-b oligomeric forms) can not only help stop disease progression but may help maintain memory as well. These preliminary observations suggest that specific therapies targeting more than one disease mediator (e.g., A-b and a pro-inflammatory cytokine such as TNF) may provide even better therapeutic efficacy for chronic neurodegenerative diseases than observed with targeting a single disease mechanism (e.g., soluble A-b alone) (see C.E. Shepherd, et al, Neurobiol Aging. 2005 Oct 24; Nelson RB., Curr Pharm Des. 2005;11:3335; William L. Klein.; Neurochem Int. 2002 ;41:345; Michelle C

Janelins, et al., J Neuroinflammation. 2005 ;2:23; Soloman B., Curr Alzheimer Res. 2004;1:149; Igor Klyubin, et al., Nat Med. 2005;11:556-61; Arancio O, et al., EMBO Journal (2004) 1-10; Bornemann KD, et al., Am J Pathol. 2001;158:63; Deane R, et al., Nat Med. 2003;9:907-13; and Eliezer Masliah, et al., Neuron. 2005;46:857).

5 The DVD-Ig molecules of the invention can bind one or more targets involved in Chronic neurodegenerative diseases such as Alzheimers. Such targets include, but are not limited to, any mediator, soluble or cell surface, implicated in AD pathogenesis e.g AGE (S100 A, amphoterin), pro-inflammatory cytokines (e.g.,IL-1), chemokines (e.g.,MCP 1), molecules that inhibit nerve regeneration (e.g.,Nogo, RGM A), molecules that enhance neurite growth (neurotrophins). The efficacy of DVD-Ig molecules can be validated in pre-clinical animal models such as the transgenic mice that over-express amyloid precursor protein or RAGE and develop Alzheimer's disease-like symptoms. In addition, DVD-Ig molecules can be constructed and tested for efficacy in the animal models and the best therapeutic DVD-Ig can be selected for testing in human patients. DVD-Ig molecules can also be employed for treatment of other

10

15 neurodegenerative diseases such as Parkinson's disease. Alpha-Synuclein is involved in Parkinson's pathology. A DVD-Ig capable of targeting alpha-synuclein and inflammatory mediators such as TNF, IL-1, MCP-1 can prove effective therapy for Parkinson's disease and are contemplated in the invention.

7.2 Neuronal Regeneration and Spinal Cord Injury

20 Despite an increase in knowledge of the pathologic mechanisms, spinal cord injury (SCI) is still a devastating condition and represents a medical indication characterized by a high medical need. Most spinal cord injuries are contusion or compression injuries and the primary injury is usually followed by secondary injury mechanisms (inflammatory mediators e.g.,cytokines and chemokines) that worsen the initial injury and result in significant enlargement of the lesion area, sometimes more than 10-fold. These primary and secondary mechanisms in SCI are very similar to those in brain injury caused by other means e.g.,stroke. No satisfying treatment exists and high dose bolus injection of methylprednisolone (MP) is the only used therapy within a narrow time window of 8 h post injury. This treatment, however, is only intended to prevent secondary injury without causing any significant functional recovery. It is heavily criticized for the lack of

25

30 unequivocal efficacy and severe adverse effects, like immunosuppression with subsequent infections and severe histopathological muscle alterations. No other drugs, biologics or small molecules, stimulating the endogenous regenerative potential are approved, but promising treatment principles and drug candidates have shown efficacy in animal models of SCI in recent years. To a large extent the lack of functional recovery in human SCI is caused by factors

35

inhibiting neurite growth, at lesion sites, in scar tissue, in myelin as well as on injury-associated cells. Such factors are the myelin-associated proteins NogoA, OMgp and MAG, RGM A, the

scar-associated CSPG (Chondroitin Sulfate Proteoglycans) and inhibitory factors on reactive astrocytes (some semaphorins and ephrins). However, at the lesion site not only growth inhibitory molecules are found but also neurite growth stimulating factors like neurotrophins, laminin, L1 and others. This ensemble of neurite growth inhibitory and growth promoting molecules may explain that blocking single factors, like NogoA or RGM A, resulted in significant functional recovery in rodent SCI models, because a reduction of the inhibitory influences could shift the balance from growth inhibition to growth promotion. However, recoveries observed with blocking a single neurite outgrowth inhibitory molecule were not complete. To achieve faster and more pronounced recoveries either blocking two neurite outgrowth inhibitory molecules e.g Nogo and RGM A, or blocking an neurite outgrowth inhibitory molecule and enhancing functions of a neurite outgrowth enhancing molecule e.g Nogo and neurotrophins, or blocking a neurite outgrowth inhibitory molecule e.g., Nogo and a pro-inflammatory molecule e.g., TNF, may be desirable (see McGee AW, et al., Trends Neurosci. 2003;26:193; Marco Domeniconi, et al., J Neurol Sci. 2005;233:43; Milan Makwana¹, et al., FEBS J. 2005;272:2628; Barry J. Dickson, Science. 2002;298:1959; Felicia Yu Hsuan Teng, et al., J Neurosci Res. 2005;79:273; Tara Karnezis, et al., Nature Neuroscience 2004; 7, 736; Gang Xu, et al., J. Neurochem. 2004; 91; 1018).

In one aspect, DVD-Igs capable of binding target pairs such as NgR and RGM A; NogoA and RGM A; MAG and RGM A; OMgp and RGM A; RGM A and RGM B; CSPGs and RGM A; aggrecan, midkine, neurocan, versican, phosphacan, Te38 and TNF- α ; A β globulomer-specific antibodies combined with antibodies promoting dendrite & axon sprouting are provided. Dendrite pathology is a very early sign of AD and it is known that Nogo A restricts dendrite growth. One can combine such type of ab with any of the SCI-candidate (myelin-proteins) Ab. Other DVD-Ig targets may include any combination of NgR-p75, NgR-Troy, NgR-Nogo66 (Nogo), NgR-Lingo, Lingo-Troy, Lingo-p75, MAG or Omgp. Additionally, targets may also include any mediator, soluble or cell surface, implicated in inhibition of neurite e.g Nogo, Omgp, MAG, RGM A, semaphorins, ephrins, soluble A-b, pro-inflammatory cytokines (e.g., IL-1), chemokines (e.g., MIP 1a), molecules that inhibit nerve regeneration. The efficacy of anti-nogo / anti-RGM A or similar DVD-Ig molecules can be validated in pre-clinical animal models of spinal cord injury. In addition, these DVD-Ig molecules can be constructed and tested for efficacy in the animal models and the best therapeutic DVD-Ig can be selected for testing in human patients. In addition, DVD-Ig molecules can be constructed that target two distinct ligand binding sites on a single receptor e.g., Nogo receptor which binds three ligand Nogo, Omgp, and MAG and RAGE that binds A-b and S100 A. Furthermore, neurite outgrowth inhibitors e.g., nogo and nogo receptor, also play a role in preventing nerve regeneration in immunological diseases like multiple sclerosis. Inhibition of nogo-nogo receptor interaction has been shown to

enhance recovery in animal models of multiple sclerosis. Therefore, DVD-Ig molecules that can block the function of one immune mediator eg a cytokine like IL-12 and a neurite outgrowth inhibitor molecule eg nogo or RGM may offer faster and greater efficacy than blocking either an immune or an neurite outgrowth inhibitor molecule alone.

5 8. Oncological disorders

Monoclonal antibody therapy has emerged as an important therapeutic modality for cancer (von Mehren M, et al 2003 Monoclonal antibody therapy for cancer. Annu Rev Med.;54:343-69). Antibodies may exert antitumor effects by inducing apoptosis, redirected cytotoxicity, interfering with ligand-receptor interactions, or preventing the expression of proteins that are critical to the neoplastic phenotype. In addition, antibodies can target components of the tumor microenvironment, perturbing vital structures such as the formation of tumor-associated vasculature. Antibodies can also target receptors whose ligands are growth factors, such as the epidermal growth factor receptor. The antibody thus inhibits natural ligands that stimulate cell growth from binding to targeted tumor cells. Alternatively, antibodies may induce an anti-idiotypic network, complement-mediated cytotoxicity, or antibody-dependent cellular cytotoxicity (ADCC). The use of dual-specific antibody that targets two separate tumor mediators will likely give additional benefit compared to a mono-specific therapy. DVD Igs capable of binding the following pairs of targets to treat oncological disease are also contemplated: IGF1 and IGF2; IGF1/2 and HER-2; VEGFR and EGFR; CD20 and CD3; CD138 and CD20; CD38 and CD20; CD38 and CD138; CD40 and CD20; CD138 and CD40; CD38 and CD40; CD-20 and CD-19; CD-20 and EGFR; CD-20 and CD-80; CD-20 and CD-22; CD-3 and HER-2; CD-3 and CD-19; EGFR and HER-2; EGFR and CD-3; EGFR and IGF1,2; EGFR and IGF1R; EGFR and RON; EGFR and HGF; EGFR and c-MET; HER-2 and IGF1,2; HER-2 and IGF1R; RON and HGF; VEGF and EGFR; VEGF and HER-2; VEGF and CD-20; VEGF and IGF1,2; VEGF and DLL4; VEGF and HGF; VEGF and RON; VEGF and NRP1; CD20 and CD3; VEGF and PLGF; DLL4 and PLGF; ErbB3 and EGFR; HGF and ErbB3, HER-2 and ErbB3; c-Met and ErbB3; HER-2 and PLGF; HER-2 and HER-2; and TNF and SOST.

In another embodiment, a DVD of the invention is capable of binding VEGF and phosphatidylserine; VEGF and ErbB3; VEGF and PLGF; VEGF and ROBO4; VEGF and BSG2; VEGF and CDCP1; VEGF and ANPEP; VEGF and c-MET; HER-2 and ERB3; HER-2 and BSG2; HER-2 and CDCP1; HER-2 and ANPEP; EGFR and CD64; EGFR and BSG2; EGFR and CDCP1; EGFR and ANPEP; IGF1R and PDGFR; IGF1R and VEGF; IGF1R and CD20; CD20 and CD74; CD20 and CD30; CD20 and DR4; CD20 and VEGFR2; CD20 and CD52; CD20 and CD4; HGF and c-MET; HGF and NRP1; HGF and phosphatidylserine; ErbB3 and IGF1R; ErbB3 and IGF1,2; c-Met and Her-2; c-Met and NRP1; c-Met and IGF1R; IGF1,2 and PDGFR; IGF1,2 and CD20; IGF1,2 and IGF1R; IGF2 and EGFR; IGF2 and HER2; IGF2 and CD20; IGF2 and

VEGF; IGF2 and IGF1R; IGF1 and IGF2; PDGFRa and VEGFR2; PDGFRa and PLGF; PDGFRa
 and VEGF; PDGFRa and c-Met; PDGFRa and EGFR; PDGFRb and VEGFR2; PDGFRb and c-
 Met; PDGFRb and EGFR; RON and c-Met; RON and MTSP1; RON and MSP; RON and
 CDCP1; VGFR1 and PLGF; VGFR1 and RON; VGFR1 and EGFR; VEGFR2 and PLGF;
 5 VEGFR2 and NRP1; VEGFR2 and RON; VEGFR2 and DLL4; VEGFR2 and EGFR; VEGFR2
 and ROBO4; VEGFR2 and CD55; LPA and S1P; EPHB2 and RON; CTLA4 and VEGF; CD3
 and EPCAM; CD40 and IL6; CD40 and IGF; CD40 and CD56; CD40 and CD70; CD40 and
 VEGFR1; CD40 and DR5; CD40 and DR4; CD40 and APRIL; CD40 and BCMA; CD40 and
 RANKL; CD28 and MAPG; CD80 and CD40; CD80 and CD30; CD80 and CD33; CD80 and
 10 CD74; CD80 and CD2; CD80 and CD3; CD80 and CD19; CD80 and CD4; CD80 and CD52;
 CD80 and VEGF; CD80 and DR5; CD80 and VEGFR2; CD22 and CD20; CD22 and CD80;
 CD22 and CD40; CD22 and CD23; CD22 and CD33; CD22 and CD74; CD22 and CD19; CD22
 and DR5; CD22 and DR4; CD22 and VEGF; CD22 and CD52; CD30 and CD20; CD30 and
 CD22; CD30 and CD23; CD30 and CD40; CD30 and VEGF; CD30 and CD74; CD30 and CD19;
 15 CD30 and DR5; CD30 and DR4; CD30 and VEGFR2; CD30 and CD52; CD30 and CD4; CD138
 and RANKL; CD33 and FTL3; CD33 and VEGF; CD33 and VEGFR2; CD33 and CD44; CD33
 and DR4; CD33 and DR5; DR4 and CD137; DR4 and IGF1,2; DR4 and IGF1R; DR4 and DR5;
 DR5 and CD40; DR5 and CD137; DR5 and CD20; DR5 and EGFR; DR5 and IGF1,2; DR5 and
 IGFR, DR5 and HER-2, EGFR and DLL4; and TNF and SOST. Other target combinations
 20 include one or more members of the EGF/erb-2/erb-3 family. Other targets (one or more)
 involved in oncological diseases that DVD Igs may bind include, but are not limited to those
 selected from the group consisting of: CD52, CD20, CD19, CD3, CD4, CD8, BMP6, IL12A,
 IL1A, IL1B, IL2, IL24, INHA, TNF, TNFSF10, BMP6, EGF, FGF1, FGF10, FGF11, FGF12,
 FGF13, FGF14, FGF16, FGF17, FGF18, FGF19, FGF2, FGF20, FGF21, FGF22, FGF23, FGF3,
 25 FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, GRP, IGF1, IGF2, IL12A, IL1A, IL1B, IL2, INHA,
 TGFA, TGFB1, TGFB2, TGFB3, VEGF, CDK2, FGF10, FGF18, FGF2, FGF4, FGF7, IGF1R,
 IL2, BCL2, CD164, CDKN1A, CDKN1B, CDKN1C, CDKN2A, CDKN2B, CDKN2C, CDKN3,
 GNRH1, IGFBP6, IL1A, IL1B, ODZ1, PAWR, PLG, TGFB1I1, AR, BRCA1, CDK3, CDK4,
 CDK5, CDK6, CDK7, CDK9, E2F1, EGFR, ENO1, ERBB2, ESR1, ESR2, IGFBP3, IGFBP6,
 30 IL2, INSL4, MYC, NOX5, NR6A1, PAP, PCNA, PRKCQ, PRKD1, PRL, TP53, FGF22, FGF23,
 FGF9, IGFBP3, IL2, INHA, KLK6, TP53, CHGB, GNRH1, IGF1, IGF2, INHA, INSL3, INSL4,
 PRL, KLK6, SHBG, NR1D1, NR1H3, NR1I3, NR2F6, NR4A3, ESR1, ESR2, NR0B1, NR0B2,
 NR1D2, NR1H2, NR1H4, NR1I2, NR2C1, NR2C2, NR2E1, NR2E3, NR2F1, NR2F2, NR3C1,
 NR3C2, NR4A1, NR4A2, NR5A1, NR5A2, NR6A1, PGR, RARB, FGF1, FGF2, FGF6, KLK3,
 35 KRT1, APOC1, BRCA1, CHGA, CHGB, CLU, COL1A1, COL6A1, EGF, ERBB2, ERK8,
 FGF1, FGF10, FGF11, FGF13, FGF14, FGF16, FGF17, FGF18, FGF2, FGF20, FGF21, FGF22,
 FGF23, FGF3, FGF4, FGF5, FGF6, FGF7, FGF8, FGF9, GNRH1, IGF1, IGF2, IGFBP3,

IGFBP6, IL12A, IL1A, IL1B, IL2, IL24, INHA, INSL3, INSL4, KLK10, KLK12, KLK13, KLK14, KLK15, KLK3, KLK4, KLK5, KLK6, KLK9, MMP2, MMP9, MSMB, NTN4, ODZ1, PAP, PLAUI, PRL, PSAP, SERPINA3, SHBG, TGFA, TIMP3, CD44, CDH1, CDH10, CDH19, CDH20, CDH7, CDH9, CDH1, CDH10, CDH13, CDH18, CDH19, CDH20, CDH7, CDH8, 5 CDH9, ROBO2, CD44, ILK, ITGA1, APC, CD164, COL6A1, MTSS1, PAP, TGFB1I1, AGR2, AIG1, AKAP1, AKAP2, CANT1, CAV1, CDH12, CLDN3, CLN3, CYB5, CYC1, DAB2IP, DES, DNCL1, ELAC2, ENO2, ENO3, FASN, FLJ12584, FLJ25530, GAGEB1, GAGEC1, GGT1, GSTP1, HIP1, HUMCYT2A, IL29, K6HF, KAI1, KRT2A, MIB1, PART1, PATE, PCA3, PIAS2, PIK3CG, PPID, PR1, PSCA, SLC2A2, SLC33A1, SLC43A1, STEAP, STEAP2, TPM1, 10 TPM2, TRPC6, ANGPT1, ANGPT2, ANPEP, ECGF1, EREG, FGF1, FGF2, FIGF, FLT1, JAG1, KDR, LAMA5, NRP1, NRP2, PGF, PLXDC1, STAB1, VEGF, VEGFC, ANGPTL3, BAI1, COL4A3, IL8, LAMA5, NRP1, NRP2, STAB1, ANGPTL4, PECAM1, PF4, PROK2, SERPINF1, TNFAIP2, CCL11, CCL2, CXCL1, CXCL10, CXCL3, CXCL5, CXCL6, CXCL9, IFNA1, IFNB1, IFNG, IL1B, IL6, MDK, EDG1, EFNA1, EFNA3, EFNB2, EGF, EPHB4, 15 FGFR3, HGF, IGF1, ITGB3, PDGFA, TEK, TGFA, TGFB1, TGFB2, TGFB1R1, CCL2, CDH5, COL18A1, EDG1, ENG, ITGAV, ITGB3, THBS1, THBS2, BAD, BAG1, BCL2, CCNA1, CCNA2, CCND1, CCNE1, CCNE2, CDH1 (E-cadherin), CDKN1B (p27Kip1), CDKN2A (p16INK4a), COL6A1, CTNNB1 (b-catenin), CTSB (cathepsin B), ERBB2 (Her-2), ESR1, ESR2, F3 (TF), FOSL1 (FRA-1), GATA3, GSN (Gelsolin), IGFBP2, IL2RA, IL6, IL6R, IL6ST 20 (glycoprotein 130), ITGA6 (a6 integrin), JUN, KLK5, KRT19, MAP2K7 (c-Jun), MKI67 (Ki-67), NGFB (NGF), NGFR, NME1 (NM23A), PGR, PLAUI (uPA), PTEN, SERPINB5 (maspin), SERPINE1 (PAI-1), TGFA, THBS1 (thrombospondin-1), TIE (Tie-1), TNFRSF6 (Fas), TNFSF6 (FasL), TOP2A (topoisomerase Iia), TP53, AZGP1 (zinc-a-glycoprotein), BPAG1 (plectin), CDKN1A (p21Wap1/Cip1), CLDN7 (claudin-7), CLU (clusterin), ERBB2 (Her-2), FGF1, 25 FLRT1 (fibronectin), GABRP (GABAa), GNAS1, ID2, ITGA6 (a6 integrin), ITGB4 (b 4 integrin), KLF5 (GC Box BP), KRT19 (Keratin 19), KRTHB6 (hair-specific type II keratin), MACMARCKS, MT3 (metallothionein-III), MUC1 (mucin), PTGS2 (COX-2), RAC2 (p21Rac2), S100A2, SCGB1D2 (lipophilin B), SCGB2A1 (mammaglobin 2), SCGB2A2 (mammaglobin 1), SPRR1B (Spr1), THBS1, THBS2, THBS4, and TNFAIP2 (B94), RON, c-Met, 30 CD64, DLL4, PLGF, CTLA4, phosphatidylserine, ROBO4, CD80, CD22, CD40, CD23, CD28, CD80, CD55, CD38, CD70, CD74, CD30, CD138, CD56, CD33, CD2, CD137, DR4, DR5, RANKL, VEGFR2, PDGFR, VEGFR1, MTSP1, MSP, EPHB2, EPHA1, EPHA2, EpCAM, PGE2, NKG2D, LPA, SIP, APRIL, BCMA, MAPG, FLT3, PDGFR alpha, PDGFR beta, ROR1, PSMA, PSCA, SCD1, and CD59.

IV. Pharmaceutical Composition

The invention also provides pharmaceutical compositions comprising a binding protein, of the invention and a pharmaceutically acceptable carrier. The pharmaceutical compositions comprising binding proteins of the invention are for use in, but not limited to, diagnosing, 5 detecting, or monitoring a disorder, in preventing, treating, managing, or ameliorating of a disorder or one or more symptoms thereof, and/or in research. In a specific embodiment, a composition comprises one or more binding proteins of the invention. In another embodiment, the pharmaceutical composition comprises one or more binding proteins of the invention and one or more prophylactic or therapeutic agents other than binding proteins of the invention for treating a 10 disorder. In an embodiment, the prophylactic or therapeutic agents known to be useful for or having been or currently being used in the prevention, treatment, management, or amelioration of a disorder or one or more symptoms thereof. In accordance with these embodiments, the composition may further comprise of a carrier, diluent or excipient.

The binding proteins of the invention can be incorporated into pharmaceutical 15 compositions suitable for administration to a subject. Typically, the pharmaceutical composition comprises a binding protein of the invention and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Examples of pharmaceutically acceptable carriers include 20 one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In some embodiments, isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride, are included in the composition. Pharmaceutically acceptable carriers may further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or 25 effectiveness of the antibody or antibody portion.

Various delivery systems are known and can be used to administer one or more antibodies of the invention or the combination of one or more antibodies of the invention and a prophylactic agent or therapeutic agent useful for preventing, managing, treating, or ameliorating a disorder or one or more symptoms thereof, e.g., encapsulation in liposomes, microparticles, microcapsules, 30 recombinant cells capable of expressing the antibody or antibody fragment, receptor-mediated endocytosis (see, e. g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of administering a prophylactic or therapeutic agent of the invention include, but are not limited to, parenteral administration (e.g., intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous) , epidurala 35 administration, intratumoral administration, and mucosal administration (e.g., intranasal and oral routes). In addition, pulmonary administration can be employed, e.g., by use of an inhaler or

nebulizer, and formulation with an aerosolizing agent. See, e.g., U.S. Pat. Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013; WO 98/31346; and WO 99/66903, each of which is incorporated herein by reference their entireties. In one embodiment, a binding protein of the invention, combination therapy, or a composition of the invention is administered using Alkermes AIR® pulmonary drug delivery technology (Alkermes, Inc., Cambridge, Mass.). In a specific embodiment, prophylactic or therapeutic agents of the invention are administered intramuscularly, intravenously, intratumorally, orally, intranasally, pulmonary, or subcutaneously. The prophylactic or therapeutic agents may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local.

In an embodiment, specific binding of antibody-coupled carbon nanotubes (CNTs) to tumor cells in vitro, followed by their highly specific ablation with near-infrared (NIR) light can be used to target tumor cells. For example, biotinylated polar lipids can be used to prepare stable, biocompatible, noncytotoxic CNT dispersions that are then attached to one or two different neutralite avidin-derivatized DVD-Igs directed against one or more tumor antigens (e.g., CD22) (Chakravarty, P. et al. (2008) Proc. Natl. Acad. Sci. USA 105:8697-8702).

In a specific embodiment, it may be desirable to administer the prophylactic or therapeutic agents of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by injection, or by means of an implant, said implant being of a porous or non-porous material, including membranes and matrices, such as sialastic membranes, polymers, fibrous matrices (e.g., Tissuel®), or collagen matrices. In one embodiment, an effective amount of one or more antibodies of the invention antagonists is administered locally to the affected area to a subject to prevent, treat, manage, and/or ameliorate a disorder or a symptom thereof. In another embodiment, an effective amount of one or more antibodies of the invention is administered locally to the affected area in combination with an effective amount of one or more therapies (e. g., one or more prophylactic or therapeutic agents) other than a binding protein of the invention of a subject to prevent, treat, manage, and/or ameliorate a disorder or one or more symptoms thereof.

In another embodiment, the prophylactic or therapeutic agent can be delivered in a controlled release or sustained release system. In one embodiment, a pump may be used to achieve controlled or sustained release (see Langer, supra; Sefton, 1987, CRC Crit. Ref. Biomed. Eng. 14:20; Buchwald et al., 1980, Surgery 88:507; Saudek et al., 1989, N. Engl. J. Med. 321:574). In another embodiment, polymeric materials can be used to achieve controlled or

sustained release of the therapies of the invention (see e.g., Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, 1983, J., Macromol. Sci. Rev. Macromol. Chem. 23:61; see also Levy et al., 1985, Science 228:190; During et al., 1989, Ann. Neurol. 25:351; Howard et al., 1989, J. Neurosurg. 71:105); U.S. Pat. No. 5,679,377; U.S. Pat. No. 5,916,597; U. S. Pat. No. 5,912,015; U.S. Pat. No. 5,989,463; U.S. Pat. No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO 99/20253. Examples of polymers used in sustained release formulations include, but are not limited to, poly(2-hydroxy ethyl methacrylate), poly(methyl methacrylate), poly(acrylic acid), poly(ethylene-co-vinyl acetate), poly(methacrylic acid), polyglycolides (PLG), polyanhydrides, poly(N-vinyl pyrrolidone), poly(vinyl alcohol), polyacrylamide, poly(ethylene glycol), polylactides (PLA), poly(lactide-co-glycolides) (PLGA), and polyorthoesters. In an embodiment, the polymer used in a sustained release formulation is inert, free of leachable impurities, stable on storage, sterile, and biodegradable. In yet another embodiment, a controlled or sustained release system can be placed in proximity of the prophylactic or therapeutic target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138 (1984)).

Controlled release systems are discussed in the review by Langer (1990, Science 249:1527-1533). Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more therapeutic agents of the invention. See, e.g., U. S. Pat. No. 4,526,938, PCT publication WO 91/05548, PCT publication WO 96/20698, Ning et al., 1996, "Intratumoral Radioimmunotherapy of a Human Colon Cancer Xenograft Using a Sustained-Release Gel," Radiotherapy & Oncology 39:179-189, Song et al., 1995, "Antibody Mediated Lung Targeting of Long-Circulating Emulsions," PDA Journal of Pharmaceutical Science & Technology 50:372-397, Cleek et al., 1997, "Biodegradable Polymeric Carriers for a bFGF Antibody for Cardiovascular Application," Pro. Int'l. Symp. Control. Rel. Bioact. Mater. 24:853-854, and Lam et al., 1997, "Microencapsulation of Recombinant Humanized Monoclonal Antibody for Local Delivery," Proc. Int'l. Symp. Control Rel. Bioact. Mater. 24:759-760, each of which is incorporated herein by reference in their entireties.

In a specific embodiment, where the composition of the invention is a nucleic acid encoding a prophylactic or therapeutic agent, the nucleic acid can be administered in vivo to promote expression of its encoded prophylactic or therapeutic agent, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U. S. Pat. No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like

peptide which is known to enter the nucleus (see, e.g., Joliot et al., 1991, Proc. Natl. Acad. Sci. USA 88:1864-1868). Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression by homologous recombination.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include, but are not limited to, parenteral, e.g., intravenous, intradermal, subcutaneous, oral, intranasal (e.g., inhalation), transdermal (e.g., topical), transmucosal, and rectal administration. In a specific embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous, subcutaneous, intramuscular, oral, intranasal, or topical administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection.

If the compositions of the invention are to be administered topically, the compositions can be formulated in the form of an ointment, cream, transdermal patch, lotion, gel, shampoo, spray, aerosol, solution, emulsion, or other form well-known to one of skill in the art. See, e.g., Remington's Pharmaceutical Sciences and Introduction to Pharmaceutical Dosage Forms, 19th ed., Mack Pub. Co., Easton, Pa. (1995). In an embodiment, for non-sprayable topical dosage forms, viscous to semi-solid or solid forms comprising a carrier or one or more excipients compatible with topical application and having a dynamic viscosity greater than water are employed. Suitable formulations include, without limitation, solutions, suspensions, emulsions, creams, ointments, powders, liniments, salves, and the like, which are, if desired, sterilized or mixed with auxiliary agents (e.g., preservatives, stabilizers, wetting agents, buffers, or salts) for influencing various properties, such as, for example, osmotic pressure. Other suitable topical dosage forms include sprayable aerosol preparations wherein the active ingredient, in an embodiment, in combination with a solid or liquid inert carrier, is packaged in a mixture with a pressurized volatile (e.g., a gaseous propellant, such as freon) or in a squeeze bottle. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well-known in the art.

If the method of the invention comprises intranasal administration of a composition, the composition can be formulated in an aerosol form, spray, mist or in the form of drops. In particular, prophylactic or therapeutic agents for use according to the present invention can be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant (e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas). In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges (composed of, e.g., gelatin) for use in an inhaler or

insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

If the method of the invention comprises oral administration, compositions can be formulated orally in the form of tablets, capsules, cachets, gelcaps, solutions, suspensions, and the like. Tablets or capsules can be prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinised maize starch, polyvinylpyrrolidone, or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose, or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc, or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulphate). The tablets may be coated by methods well-known in the art. Liquid preparations for oral administration may take the form of, but not limited to, solutions, syrups or suspensions, or they may be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives, or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters, ethyl alcohol, or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations may also contain buffer salts, flavoring, coloring, and sweetening agents as appropriate. Preparations for oral administration may be suitably formulated for slow release, controlled release, or sustained release of a prophylactic or therapeutic agent(s).

The method of the invention may comprise pulmonary administration, e.g., by use of an inhaler or nebulizer, of a composition formulated with an aerosolizing agent. See, e.g., U.S. Pat. Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013; WO 98/31346; and WO 99/66903, each of which is incorporated herein by reference their entireties. In a specific embodiment, a binding protein of the invention, combination therapy, and/or composition of the invention is administered using Alkermes AIR® pulmonary drug delivery technology (Alkermes, Inc., Cambridge, Mass.).

The method of the invention may comprise administration of a composition formulated for parenteral administration by injection (e.g., by bolus injection or continuous infusion). Formulations for injection may be presented in unit dosage form (e.g., in ampoules or in multi-dose containers) with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle (e.g., sterile pyrogen-free water) before use.

The methods of the invention may additionally comprise of administration of compositions formulated as depot preparations. Such long acting formulations may be administered by implantation (e.g., subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compositions may be formulated with suitable polymeric or hydrophobic materials (e.g., as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives (e.g., as a sparingly soluble salt).

The methods of the invention encompass administration of compositions formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

Generally, the ingredients of compositions are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the mode of administration is infusion, composition can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the mode of administration is by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

In particular, the invention also provides that one or more of the prophylactic or therapeutic agents, or pharmaceutical compositions of the invention is packaged in a hermetically sealed container such as an ampoule or sachette indicating the quantity of the agent. In one embodiment, one or more of the prophylactic or therapeutic agents, or pharmaceutical compositions of the invention is supplied as a dry sterilized lyophilized powder or water free concentrate in a hermetically sealed container and can be reconstituted (e.g., with water or saline) to the appropriate concentration for administration to a subject. In an embodiment, one or more of the prophylactic or therapeutic agents or pharmaceutical compositions of the invention is supplied as a dry sterile lyophilized powder in a hermetically sealed container at a unit dosage of at least 5 mg, at least 10 mg, at least 15 mg, at least 25 mg, at least 35 mg, at least 45 mg, at least 50 mg, at least 75 mg, or at least 100 mg. The lyophilized prophylactic or therapeutic agents or pharmaceutical compositions of the invention should be stored at between 2° C. and 8° C. in its original container and the prophylactic or therapeutic agents, or pharmaceutical compositions of the invention should be administered within 1 week, e.g., within 5 days, within 72 hours, within 48 hours, within 24 hours, within 12 hours, within 6 hours, within 5 hours, within 3 hours, or within 1 hour after being reconstituted. In an alternative embodiment, one or more of the prophylactic or therapeutic agents or pharmaceutical compositions of the invention is supplied in liquid form in a hermetically sealed container indicating the quantity and concentration of the

agent. In an embodiment, the liquid form of the administered composition is supplied in a hermetically sealed container at least 0.25 mg/ml, at least 0.5 mg/ml, at least 1 mg/ml, at least 2.5 mg/ml, at least 5 mg/ml, at least 8 mg/ml, at least 10 mg/ml, at least 15 mg/kg, at least 25 mg/ml, at least 50 mg/ml, at least 75 mg/ml or at least 100 mg/ml. The liquid form should be stored at
 5 between 2° C. and 8° C. in its original container.

The binding proteins of the invention can be incorporated into a pharmaceutical composition suitable for parenteral administration. In an embodiment, the antibody or antibody-
 portions will be prepared as an injectable solution containing 0.1-250 mg/ml binding protein. The injectable solution can be composed of either a liquid or lyophilized dosage form in a flint or
 10 amber vial, ampule or pre-filled syringe. The buffer can be L-histidine (1-50 mM), optimally 5-10mM, at pH 5.0 to 7.0 (optimally pH 6.0). Other suitable buffers include but are not limited to, sodium succinate, sodium citrate, sodium phosphate or potassium phosphate. Sodium chloride can be used to modify the toxicity of the solution at a concentration of 0-300 mM (optimally 150 mM for a liquid dosage form). Cryoprotectants can be included for a lyophilized dosage form,
 15 principally 0-10% sucrose (optimally 0.5-1.0%). Other suitable cryoprotectants include trehalose and lactose. Bulking agents can be included for a lyophilized dosage form, principally 1-10% mannitol (optimally 2-4%). Stabilizers can be used in both liquid and lyophilized dosage forms, principally 1-50 mM L-Methionine (optimally 5-10 mM). Other suitable bulking agents include glycine, arginine, can be included as 0-0.05% polysorbate-80 (optimally 0.005-0.01%).
 20 Additional surfactants include but are not limited to polysorbate 20 and BRIJ surfactants. The pharmaceutical composition comprising the binding proteins of the invention prepared as an injectable solution for parenteral administration, can further comprise an agent useful as an adjuvant, such as those used to increase the absorption, or dispersion of a therapeutic protein (e.g., antibody). A particularly useful adjuvant is hyaluronidase, such as Hylenex® (recombinant
 25 human hyaluronidase). Addition of hyaluronidase in the injectable solution improves human bioavailability following parenteral administration, particularly subcutaneous administration. It also allows for greater injection site volumes (i.e. greater than 1 ml) with less pain and discomfort, and minimum incidence of injection site reactions. (see WO2004078140, and US2006104968 incorporated herein by reference).

30 The compositions of this invention may be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The form chosen depends on the intended mode of administration and therapeutic application. Typical compositions are in the form of injectable or infusible solutions, such as
 35 compositions similar to those used for passive immunization of humans with other antibodies. The chosen mode of administration is parenteral (e.g., intravenous, subcutaneous, intraperitoneal,

intramuscular). In an embodiment, the antibody is administered by intravenous infusion or injection. In another embodiment, the antibody is administered by intramuscular or subcutaneous injection.

Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the active compound (*i.e.*, antibody or antibody portion) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated herein, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated herein. In the case of sterile, lyophilized powders for the preparation of sterile injectable solutions, the methods of preparation are vacuum drying and spray-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including, in the composition, an agent that delays absorption, for example, monostearate salts and gelatin.

The binding proteins of the present invention can be administered by a variety of methods known in the art, although for many therapeutic applications, in an embodiment, the route/mode of administration is subcutaneous injection, intravenous injection or infusion. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, *e.g.*, *Sustained and Controlled Release Drug Delivery Systems*, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

In certain embodiments, a binding protein of the invention may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer

a compound of the invention by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

Supplementary active compounds can also be incorporated into the compositions. In certain embodiments, a binding protein of the invention is coformulated with and/or
5 coadministered with one or more additional therapeutic agents that are useful for treating disorders with binding protein of the invention. For example, a binding protein of the invention may be coformulated and/or coadministered with one or more additional antibodies that bind other targets (*e.g.*, antibodies that bind other cytokines or that bind cell surface molecules). Furthermore, one or more antibodies of the invention may be used in combination with two or
10 more of the foregoing therapeutic agents. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

In certain embodiments, a binding protein is linked to a half-life extending vehicle known in the art. Such vehicles include, but are not limited to, the Fc domain, polyethylene
15 glycol, and dextran. Such vehicles are described, *e.g.*, in U.S. Application Serial No. 09/428,082 and published PCT Application No. WO 99/25044, which are hereby incorporated by reference for any purpose.

In a specific embodiment, nucleic acid sequences encoding a binding protein of the invention or another prophylactic or therapeutic agent of the invention are administered to treat,
20 prevent, manage, or ameliorate a disorder or one or more symptoms thereof by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded antibody or prophylactic or therapeutic agent of the invention that mediates a prophylactic or therapeutic effect.

Any of the methods for gene therapy available in the art can be used according to the
25 present invention. For general reviews of the methods of gene therapy, see Goldspiel et al., 1993, Clinical Pharmacy 12:488-505; Wu and Wu, 1991, Biotherapy 3:87-95; Tolstoshev, 1993, Ann. Rev. Pharmacol. Toxicol. 32:573-596; Mulligan, Science 260:926- 932 (1993); and Morgan and Anderson, 1993, Ann. Rev. Biochem. 62:191-217; May, 1993, TIBTECH 11(5):155-215.
30 Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990). Detailed description of various methods of gene therapy are disclosed in US20050042664 A1 which is incorporated herein by reference.

The binding proteins of the invention are useful in treating various diseases wherein the targets that are recognized by the binding proteins are detrimental. Such diseases include, but are not limited to, rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's

granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, spondyloarthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases, Hepatitis B, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, fibrosis, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated

hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS,

5 glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjögren's syndrome,

10 Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis, alcohol-induced liver injury, choleosatatis, idiosyncratic liver disease, Drug-Induced hepatitis,

15 Non-alcoholic Steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental disorders (*e.g.*, depression and schizophrenia), Th2 Type and Th1 Type mediated diseases, acute and chronic pain (different forms of pain), and cancers such as lung, breast, stomach, bladder, colon, pancreas, ovarian, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), Abetalipoproteinemia, Acrocyanosis, acute and chronic parasitic or infectious

20 processes, acute leukemia, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, arial ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-1- antitrypsin deficiency, amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3

25 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, Burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors,

30 cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive

35 heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, Dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis,

dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, Diffuse
 Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's
 Syndrome in middle age, drug- induced movement disorders induced by drugs which block CNS
 dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy,
 5 epiglottitis, epstein-barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders,
 familial hemophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's
 ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer,
 glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive
 sepsis, granulomas due to intracellular organisms, hairy cell leukemia, Hallerorden-Spatz
 10 disease, hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis,
 hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage,
 hepatitis (A), His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease,
 hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis,
 hypertension, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation,
 15 idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity,
 Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing
 radiation exposure, iridocyclitis/uveitis/optic neuritis, ischemia- reperfusion injury, ischemic
 stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney
 transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system,
 20 lipedema, liver transplant rejection, lymphoderma, malaria, malignant Lymphoma, malignant
 histiocytosis, malignant melanoma, meningitis, meningococemia, metabolic/idiopathic, migraine
 headache, mitochondrial multi.system disorder, mixed connective tissue disease, monoclonal
 gammopathy, multiple myeloma, multiple systems degenerations (Mencel Dejerine- Thomas Shi-
 Drager and Machado-Joseph), myasthenia gravis, mycobacterium avium intracellulare,
 25 mycobacterium tuberculosis, myelodysplastic syndrome, myocardial infarction, myocardial
 ischemic disorders, nasopharyngeal carcinoma, neonatal chronic lung disease, nephritis,
 nephrosis, neurodegenerative diseases, neurogenic I muscular atrophies , neutropenic fever, non-
 hodgkins lymphoma, occlusion of the abdominal aorta and its branches, occlusive arterial
 disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy reversal procedures,
 30 organomegaly, osteoporosis, pancreas transplant rejection, pancreatic carcinoma, paraneoplastic
 syndrome/hypercalcemia of malignancy, parathyroid transplant rejection, pelvic inflammatory
 disease, perennial rhinitis, pericardial disease, peripheral atherosclerotic disease, peripheral
 vascular disorders, peritonitis, pernicious anemia, pneumocystis carinii pneumonia, pneumonia,
 POEMS syndrome (polyneuropathy, organomegaly, endocrinopathy, monoclonal gammopathy,
 35 and skin changes syndrome), post perfusion syndrome, post pump syndrome, post-MI cardiomyopathy
 syndrome, preeclampsia, Progressive supranuclear Palsy, primary pulmonary hypertension,
 radiation therapy, Raynaud's phenomenon and disease, Raynaud's disease, Refsum's disease,

regular narrow QRS tachycardia, renovascular hypertension, reperfusion injury, restrictive cardiomyopathy, sarcomas, scleroderma, senile chorea, Senile Dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia, skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors, specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural lesions of the cerebellum, Subacute sclerosing panencephalitis, Syncope, syphilis of the cardiovascular system, systemic anaphalaxis, systemic inflammatory response syndrome, systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, Telangiectasia, thromboangitis obliterans, thrombocytopenia, toxicity, transplants, trauma/hemorrhage, type III hypersensitivity reactions, type IV hypersensitivity, unstable angina, uremia, urosepsis, urticaria, valvular heart diseases, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation, viral and fungal infections, vital encephalitis/aseptic meningitis, vital-associated hemaphagocytic syndrome, Wernicke- Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or tissue. (see Peritt et al. PCT publication No. WO2002097048A2, Leonard et al., PCT publication No. WO9524918 A1, and Salfeld et al., PCT publication No. WO00/56772A1).

The binding proteins of the invention can be used to treat humans suffering from autoimmune diseases, in particular those associated with inflammation, including, rheumatoid arthritis, spondylitis, allergy, autoimmune diabetes, autoimmune uveitis. In an embodiment, the binding proteins of the invention or antigen-binding portions thereof, are used to treat rheumatoid arthritis, Crohn's disease, multiple sclerosis, insulin dependent diabetes mellitus and psoriasis.

In an embodiment, diseases that can be treated or diagnosed with the compositions and methods of the invention include, but are not limited to, primary and metastatic cancers, including carcinomas of breast, colon, rectum, lung, oropharynx, hypopharynx, esophagus, stomach, pancreas, liver, gallbladder and bile ducts, small intestine, urinary tract (including kidney, bladder and urothelium), female genital tract (including cervix, uterus, and ovaries as well as choriocarcinoma and gestational trophoblastic disease), male genital tract (including prostate, seminal vesicles, testes and germ cell tumors), endocrine glands (including the thyroid, adrenal, and pituitary glands), and skin, as well as hemangiomas, melanomas, sarcomas (including those arising from bone and soft tissues as well as Kaposi's sarcoma), tumors of the brain, nerves, eyes, and meninges (including astrocytomas, gliomas, glioblastomas, retinoblastomas, neuromas, neuroblastomas, Schwannomas, and meningiomas), solid tumors arising from hematopoietic malignancies such as leukemias, and lymphomas (both Hodgkin's and non-Hodgkin's lymphomas).

In an embodiment, the antibodies of the invention or antigen-binding portions thereof, are used to treat cancer or in the prevention of metastases from the tumors described herein either when used alone or in combination with radiotherapy and/or other chemotherapeutic agents.

The antibodies of the invention, or antigen binding portions thereof, may be combined with agents that include but are not limited to, antineoplastic agents, radiotherapy, chemotherapy such as DNA alkylating agents, cisplatin, carboplatin, anti-tubulin agents, paclitaxel, docetaxel, taxol, doxorubicin, gemcitabine, gemzar, anthracyclines, adriamycin, topoisomerase I inhibitors, 5 topoisomerase II inhibitors, 5-fluorouracil (5-FU), leucovorin, irinotecan, receptor tyrosine kinase inhibitors (e.g., erlotinib, gefitinib), COX-2 inhibitors (e.g., celecoxib), kinase inhibitors, and siRNAs.

A binding protein of the invention also can be administered with one or more additional therapeutic agents useful in the treatment of various diseases.

10 A binding protein of the invention can be used alone or in combination to treat such diseases. It should be understood that the binding proteins can be used alone or in combination with an additional agent, e.g., a therapeutic agent, said additional agent being selected by the skilled artisan for its intended purpose. For example, the additional agent can be a therapeutic agent art-recognized as being useful to treat the disease or condition being treated by the antibody 15 of the present invention. The additional agent also can be an agent that imparts a beneficial attribute to the therapeutic composition e.g., an agent which effects the viscosity of the composition.

It should further be understood that the combinations which are to be included within this invention are those combinations useful for their intended purpose. The agents set forth below are 20 illustrative for purposes and not intended to be limited. The combinations, which are part of this invention, can be the antibodies of the present invention and at least one additional agent selected from the lists below. The combination can also include more than one additional agent, e.g., two or three additional agents if the combination is such that the formed composition can perform its intended function.

25 Combinations to treat autoimmune and inflammatory diseases are non-steroidal anti-inflammatory drug(s) also referred to as NSAIDS which include drugs like ibuprofen. Other combinations are corticosteroids including prednisolone; the well known side-effects of steroid use can be reduced or even eliminated by tapering the steroid dose required when treating patients in combination with the DVD Igs of this invention. Non-limiting examples of therapeutic agents 30 for rheumatoid arthritis with which an antibody, or antibody portion, of the invention can be combined include the following: cytokine suppressive anti-inflammatory drug(s) (CSAIDs); antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-15, IL-16, IL-18, IL-21, IL-23, interferons, EMAP-II, GM-CSF, FGF, and PDGF. Binding proteins of the invention, or antigen binding 35 portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3,

CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD80 (B7.1), CD86 (B7.2), CD90, CTLA or their ligands including CD154 (gp39 or CD40L).

Combinations of therapeutic agents may interfere at different points in the autoimmune and subsequent inflammatory cascade; examples include TNF antagonists like chimeric, humanized or human TNF antibodies, ADALIMUMAB, (PCT Publication No. WO 97/29131), CA2 (RemicadeTM), CDP 571, and soluble p55 or p75 TNF receptors, derivatives, thereof, (p75TNFR1gG (EnbrelTM) or p55TNFR1gG (Lenercept)), and also TNF α converting enzyme (TACE) inhibitors; similarly IL-1 inhibitors (Interleukin-1-converting enzyme inhibitors, IL-1RA etc.) may be effective for the same reason. Other combinations include Interleukin 11. Yet another combination include key players of the autoimmune response which may act parallel to, dependent on or in concert with IL-12 function; especially are IL-18 antagonists including IL-18 antibodies or soluble IL-18 receptors, or IL-18 binding proteins. It has been shown that IL-12 and IL-18 have overlapping but distinct functions and a combination of antagonists to both may be most effective. Yet another combination are non-depleting anti-CD4 inhibitors. Yet other combinations include antagonists of the co-stimulatory pathway CD80 (B7.1) or CD86 (B7.2) including antibodies, soluble receptors or antagonistic ligands.

The binding proteins of the invention may also be combined with agents, such as methotrexate, 6-MP, azathioprine sulphasalazine, mesalazine, olsalazine chloroquine/hydroxychloroquine, pencillamine, aurothiomalate (intramuscular and oral), azathioprine, cochlincine, corticosteroids (oral, inhaled and local injection), beta-2 adrenoreceptor agonists (salbutamol, terbutaline, salmeteral), xanthines (theophylline, aminophylline), cromoglycate, nedocromil, ketotifen, ipratropium and oxitropium, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF- α or IL-1 (e.g., IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme (TACE) inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g., soluble p55 or p75 TNF receptors and the derivatives p75TNFR1gG (EnbrelTM and p55TNFR1gG (Lenercept)), sIL-1RI, sIL-1RII, sIL-6R), antiinflammatory cytokines (e.g., IL-4, IL-10, IL-11, IL-13 and TGF β), celecoxib, folic acid, hydroxychloroquine sulfate, rofecoxib, etanercept, infliximab, naproxen, valdecoxib, sulfasalazine, methylprednisolone, meloxicam, methylprednisolone acetate, gold sodium thiomalate, aspirin, triamcinolone acetonide, propoxyphene napsylate/apap, folate, nabumetone, diclofenac, piroxicam, etodolac, diclofenac sodium, oxaprozin, oxycodone hcl, hydrocodone

bitartrate/apap, diclofenac sodium/misoprostol, fentanyl, anakinra, human recombinant, tramadol hcl, salsalate, sulindac, cyanocobalamin/fa/pyridoxine, acetaminophen, alendronate sodium, prednisolone, morphine sulfate, lidocaine hydrochloride, indomethacin, glucosamine sulf/chondroitin, amitriptyline hcl, sulfadiazine, oxycodone hcl/acetaminophen, olopatadine hcl, misoprostol, naproxen sodium, omeprazole, cyclophosphamide, rituximab, IL-1 TRAP, MRA, CTLA4-IG, IL-18 BP, anti-IL-18, Anti-IL15, BIRB-796, SCIO-469, VX-702, AMG-548, VX-740, Roflumilast, IC-485, CDC-801, and Mesopram. Combinations include methotrexate or leflunomide and in moderate or severe rheumatoid arthritis cases, cyclosporine.

Nonlimiting additional agents which can also be used in combination with a binding protein to treat rheumatoid arthritis include, but are not limited to, the following: non-steroidal anti-inflammatory drug(s) (NSAIDs); cytokine suppressive anti-inflammatory drug(s) (CSAIDs); CDP-571/BAY-10-3356 (humanized anti-TNF α antibody; Celltech/Bayer); cA2/infliximab (chimeric anti-TNF α antibody; Centocor); 75 kDTNFR-IgG/etanercept (75 kD TNF receptor-IgG fusion protein; Immunex; see *e.g.*, *Arthritis & Rheumatism* (1994) Vol. 37, S295; *J. Invest. Med.* (1996) Vol. 44, 235A); 55 kDTNF-IgG (55 kD TNF receptor-IgG fusion protein; Hoffmann-LaRoche); IDEC-CE9.1/SB 210396 (non-depleting primatized anti-CD4 antibody; IDEC/SmithKline; see *e.g.*, *Arthritis & Rheumatism* (1995) Vol. 38, S185); DAB 486-IL-2 and/or DAB 389-IL-2 (IL-2 fusion proteins; Seragen; see *e.g.*, *Arthritis & Rheumatism* (1993) Vol. 36, 1223); Anti-Tac (humanized anti-IL-2R α ; Protein Design Labs/Roche); IL-4 (anti-inflammatory cytokine; DNAX/Schering); IL-10 (SCH 52000; recombinant IL-10, anti-inflammatory cytokine; DNAX/Schering); IL-4; IL-10 and/or IL-4 agonists (*e.g.*, agonist antibodies); IL-1RA (IL-1 receptor antagonist; Synergen/Amgen); anakinra (Kineret[®]/Amgen); TNF-bp/s-TNF (soluble TNF binding protein; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S284; *Amer. J. Physiol. - Heart and Circulatory Physiology* (1995) Vol. 268, pp. 37-42); R973401 (phosphodiesterase Type IV inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282); MK-966 (COX-2 Inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S81); Iloprost (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S82); methotrexate; thalidomide (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282) and thalidomide-related drugs (*e.g.*, Celgen); leflunomide (anti-inflammatory and cytokine inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S131; *Inflammation Research* (1996) Vol. 45, pp. 103-107); tranexamic acid (inhibitor of plasminogen activation; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S284); T-614 (cytokine inhibitor; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282); prostaglandin E1 (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S282); Tenidap (non-steroidal anti-inflammatory drug; see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S280); Naproxen (non-steroidal anti-

inflammatory drug; see *e.g.*, *Neuro Report* (1996) Vol. 7, pp. 1209-1213); Meloxicam (non-steroidal anti-inflammatory drug); Ibuprofen (non-steroidal anti-inflammatory drug); Piroxicam (non-steroidal anti-inflammatory drug); Diclofenac (non-steroidal anti-inflammatory drug); Indomethacin (non-steroidal anti-inflammatory drug); Sulfasalazine (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S281); Azathioprine (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S281); ICE inhibitor (inhibitor of the enzyme interleukin-1 β converting enzyme); zap-70 and/or lck inhibitor (inhibitor of the tyrosine kinase zap-70 or lck); VEGF inhibitor and/or VEGF-R inhibitor (inhibitors of vascular endothelial cell growth factor or vascular endothelial cell growth factor receptor; inhibitors of angiogenesis); corticosteroid anti-inflammatory drugs (*e.g.*, SB203580); TNF-convertase inhibitors; anti-IL-12 antibodies; anti-IL-18 antibodies; interleukin-11 (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S296); interleukin-13 (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S308); interleukin -17 inhibitors (see *e.g.*, *Arthritis & Rheumatism* (1996) Vol. 39, No. 9 (supplement), S120); gold; penicillamine; chloroquine; chlorambucil; hydroxychloroquine; cyclosporine; cyclophosphamide; total lymphoid irradiation; anti-thymocyte globulin; anti-CD4 antibodies; CD5-toxins; orally-administered peptides and collagen; lobenzarit disodium; Cytokine Regulating Agents (CRAs) HP228 and HP466 (Houghten Pharmaceuticals, Inc.); ICAM-1 antisense phosphorothioate oligo-deoxynucleotides (ISIS 2302; Isis Pharmaceuticals, Inc.); soluble complement receptor 1 (TP10; T Cell Sciences, Inc.); prednisone; orgotein; glycosaminoglycan polysulphate; minocycline; anti-IL2R antibodies; marine and botanical lipids (fish and plant seed fatty acids; see *e.g.*, DeLuca *et al.* (1995) *Rheum. Dis. Clin. North Am.* 21:759-777); auranofin; phenylbutazone; meclofenamic acid; flufenamic acid; intravenous immune globulin; zileuton; azaribine; mycophenolic acid (RS-61443); tacrolimus (FK-506); sirolimus (rapamycin); amiprilose (therafectin); cladribine (2-chlorodeoxyadenosine); methotrexate; bcl-2 inhibitors (see Bruncko, Milan *et al.*, *Journal of Medicinal Chemistry* (2007), 50(4), 641-662); antivirals and immune modulating agents.

In one embodiment, the binding protein or antigen-binding portion thereof, is administered in combination with one of the following agents for the treatment of rheumatoid arthritis: small molecule inhibitor of KDR, small molecule inhibitor of Tie-2; methotrexate; prednisone; celecoxib; folic acid; hydroxychloroquine sulfate; rofecoxib; etanercept; infliximab; leflunomide; naproxen; valdecoxib; sulfasalazine; methylprednisolone; ibuprofen; meloxicam; methylprednisolone acetate; gold sodium thiomalate; aspirin; azathioprine; triamcinolone acetonide; propxyphene napsylate/apap; folate; nabumetone; diclofenac; piroxicam; etodolac; diclofenac sodium; oxaprozin; oxycodone hcl; hydrocodone bitartrate/apap; diclofenac sodium/misoprostol; fentanyl; anakinra, human recombinant; tramadol hcl; salsalate; sulindac; cyanocobalamin/fa/pyridoxine; acetaminophen; alendronate sodium; prednisolone; morphine

sulfate; lidocaine hydrochloride; indomethacin; glucosamine sulfate/chondroitin; cyclosporine; amitriptyline hcl; sulfadiazine; oxycodone hcl/acetaminophen; olopatadine hcl; misoprostol; naproxen sodium; omeprazole; mycophenolate mofetil; cyclophosphamide; rituximab; IL-1 TRAP; MRA; CTLA4-IG; IL-18 BP; IL-12/23; anti-IL 18; anti-IL 15; BIRB-796; SCIO-469; 5 VX-702; AMG-548; VX-740; Roflumilast; IC-485; CDC-801; and mesopram.

Non-limiting examples of therapeutic agents for inflammatory bowel disease with which a binding protein of the invention can be combined include the following: budenoside; epidermal growth factor; corticosteroids; cyclosporin, sulfasalazine; aminosalicylates; 6-mercaptopurine; azathioprine; metronidazole; lipoxygenase inhibitors; mesalamine; olsalazine; balsalazide; 10 antioxidants; thromboxane inhibitors; IL-1 receptor antagonists; anti-IL-1 β mAbs; anti-IL-6 mAbs; growth factors; elastase inhibitors; pyridinyl-imidazole compounds; antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-15, IL-16, IL-17, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface 15 molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands. The antibodies of the invention, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, 20 adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (e.g., IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof 25 (e.g., soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R) and antiinflammatory cytokines (e.g., IL-4, IL-10, IL-11, IL-13 and TGF β) and bcl-2 inhibitors.

Examples of therapeutic agents for Crohn's disease in which a binding protein can be combined include the following: TNF antagonists, for example, anti-TNF antibodies, ADALIMUMAB (PCT Publication No. WO 97/29131; HUMIRA), CA2 (REMICADE), CDP 30 571, TNFR-Ig constructs, (p75TNFR IgG (ENBREL) and p55TNFR IgG (LENERCEPT)) inhibitors and PDE4 inhibitors. Antibodies of the invention, or antigen binding portions thereof, can be combined with corticosteroids, for example, budenoside and dexamethasone. Binding proteins of the invention or antigen binding portions thereof, may also be combined with agents such as sulfasalazine, 5-aminosalicylic acid and olsalazine, and agents which interfere with 35 synthesis or action of proinflammatory cytokines such as IL-1, for example, IL-1 β converting enzyme inhibitors and IL-1ra. Antibodies of the invention or antigen binding portion thereof may

also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors 6-
 mercaptopurines. Binding proteins of the invention, or antigen binding portions thereof, can be
 combined with IL-11. Binding proteins of the invention, or antigen binding portions thereof, can
 be combined with mesalamine, prednisone, azathioprine, mercaptopurine, infliximab,
 5 methylprednisolone sodium succinate, diphenoxylate/atrop sulfate, loperamide hydrochloride,
 methotrexate, omeprazole, folate, ciprofloxacin/dextrose-water, hydrocodone bitartrate/apap,
 tetracycline hydrochloride, fluocinonide, metronidazole, thimerosal/boric acid,
 cholestyramine/sucrose, ciprofloxacin hydrochloride, hyoscyamine sulfate, meperidine
 hydrochloride, midazolam hydrochloride, oxycodone hcl/acetaminophen, promethazine
 10 hydrochloride, sodium phosphate, sulfamethoxazole/trimethoprim, celecoxib, polycarbophil,
 propoxyphene napsylate, hydrocortisone, multivitamins, balsalazide disodium, codeine
 phosphate/apap, colesevelam hcl, cyanocobalamin, folic acid, levofloxacin, methylprednisolone,
 natalizumab and interferon-gamma

Non-limiting examples of therapeutic agents for multiple sclerosis with which binding
 15 proteins of the invention can be combined include the following: corticosteroids; prednisolone;
 methylprednisolone; azathioprine; cyclophosphamide; cyclosporine; methotrexate; 4-
 aminopyridine; tizanidine; interferon- β 1a (AVONEX; Biogen); interferon- β 1b (BETASERON;
 Chiron/Berlex); interferon α -n3) (Interferon Sciences/Fujimoto), interferon- α (Alfa
 Wassermann/J&J), interferon β 1A-IF (Serono/Inhale Therapeutics), Peginterferon α 2b
 20 (Enzon/Schering-Plough), Copolymer 1 (Cop-1; COPAXONE; Teva Pharmaceutical Industries,
 Inc.); hyperbaric oxygen; intravenous immunoglobulin; clabribine; antibodies to or antagonists of
 other human cytokines or growth factors and their receptors, for example, TNF, LT, IL-1, IL-2,
 IL-6, IL-7, IL-8, IL-23, IL-15, IL-16, IL-18, EMAP-II, GM-CSF, FGF, and PDGF. Binding
 proteins of the invention can be combined with antibodies to cell surface molecules such as CD2,
 25 CD3, CD4, CD8, CD19, CD20, CD25, CD28, CD30, CD40, CD45, CD69, CD80, CD86, CD90
 or their ligands. Binding proteins of the invention, may also be combined with agents, such as
 methotrexate, cyclosporine, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs,
 for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors,
 adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents
 30 which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (e.g., IRAK,
 NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TACE inhibitors,
 T-cell signaling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine,
 azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine
 receptors and derivatives thereof (e.g., soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-
 35 6R), antiinflammatory cytokines (e.g., IL-4, IL-10, IL-13 and TGF β) and bcl-2 inhibitors.

Examples of therapeutic agents for multiple sclerosis in which binding proteins of the invention can be combined include interferon- β , for example, IFN β 1a and IFN β 1b; copaxone, corticosteroids, caspase inhibitors, for example inhibitors of caspase-1, IL-1 inhibitors, TNF inhibitors, and antibodies to CD40 ligand and CD80.

5 The binding proteins of the invention, may also be combined with agents, such as alemtuzumab, dronabinol, Unimed, daclizumab, mitoxantrone, xaliproden hydrochloride, fampridine, glatiramer acetate, natalizumab, sinnabidol, a-immunokine NNSO3, ABR-215062, AnergiX.MS, chemokine receptor antagonists, BBR-2778, calagualine, CPI-1189, LEM (liposome encapsulated mitoxantrone), THC.CBD (cannabinoid agonist) MBP-8298, mesopram
10 (PDE4 inhibitor), MNA-715, anti-IL-6 receptor antibody, neurovax, pirfenidone allotrap 1258 (RDP-1258), sTNF-R1, talampanel, teriflunomide, TGF-beta2, tiplimotide, VLA-4 antagonists (for example, TR-14035, VLA4 Ultrahaler, Antegran-ELAN/Biogen), interferon gamma antagonists, IL-4 agonists.

Non-limiting examples of therapeutic agents for Angina with which binding proteins of
15 the invention can be combined include the following: aspirin, nitroglycerin, isosorbide mononitrate, metoprolol succinate, atenolol, metoprolol tartrate, amlodipine besylate, diltiazem hydrochloride, isosorbide dinitrate, clopidogrel bisulfate, nifedipine, atorvastatin calcium, potassium chloride, furosemide, simvastatin, verapamil hcl, digoxin, propranolol hydrochloride, carvedilol, lisinopril, spironolactone, hydrochlorothiazide, enalapril maleate, nadolol, ramipril,
20 enoxaparin sodium, heparin sodium, valsartan, sotalol hydrochloride, fenofibrate, ezetimibe, bumetanide, losartan potassium, lisinopril/hydrochlorothiazide, felodipine, captopril, bisoprolol fumarate.

Non-limiting examples of therapeutic agents for Ankylosing Spondylitis with which binding proteins of the invention can be combined include the following: ibuprofen, diclofenac
25 and misoprostol, naproxen, meloxicam, indomethacin, diclofenac, celecoxib, rofecoxib, Sulfasalazine, Methotrexate, azathioprine, minocyclin, prednisone, etanercept, infliximab.

Non-limiting examples of therapeutic agents for Asthma with which binding proteins of the invention can be combined include the following: albuterol, salmeterol/fluticasone, montelukast sodium, fluticasone propionate, budesonide, prednisone, salmeterol xinafoate,
30 levalbuterol hcl, albuterol sulfate/ipratropium, prednisolone sodium phosphate, triamcinolone acetonide, beclomethasone dipropionate, ipratropium bromide, azithromycin, pirbuterol acetate, prednisolone, theophylline anhydrous, methylprednisolone sodium succinate, clarithromycin, zafirlukast, formoterol fumarate, influenza virus vaccine, methylprednisolone, amoxicillin trihydrate, flunisolide, allergy injection, cromolyn sodium, fexofenadine hydrochloride,
35 flunisolide/menthol, amoxicillin/clavulanate, levofloxacin, inhaler assist device, guaifenesin,

dexamethasone sodium phosphate, moxifloxacin hcl, doxycycline hyclate, guaifenesin/d-methorphan, p-ephedrine/cod/chlorphenir, gatifloxacin, cetirizine hydrochloride, mometasone furoate, salmeterol xinafoate, benzonatate, cephalixin, pe/hydrocodone/chlorphenir, cetirizine hcl/pseudoephed, phenylephrine/cod/promethazine, codeine/promethazine, cefprozil,
 5 dexamethasone, guaifenesin/pseudoephedrine, chlorpheniramine/hydrocodone, nedocromil sodium, terbutaline sulfate, epinephrine, methylprednisolone, metaproterenol sulfate.

Non-limiting examples of therapeutic agents for COPD with which binding proteins of the invention can be combined include the following: albuterol sulfate/ipratropium, ipratropium bromide, salmeterol/fluticasone, albuterol, salmeterol xinafoate, fluticasone propionate,
 10 prednisone, theophylline anhydrous, methylprednisolone sodium succinate, montelukast sodium, budesonide, formoterol fumarate, triamcinolone acetonide, levofloxacin, guaifenesin, azithromycin, beclomethasone dipropionate, levalbuterol hcl, flunisolide, ceftriaxone sodium, amoxicillin trihydrate, gatifloxacin, zafirlukast, amoxicillin/clavulanate, flunisolide/menthol, chlorpheniramine/hydrocodone, metaproterenol sulfate, methylprednisolone, mometasone furoate,
 15 p-ephedrine/cod/chlorphenir, pirbuterol acetate, p-ephedrine/loratadine, terbutaline sulfate, tiotropium bromide, (R,R)-formoterol, TgAAT, Cilomilast, Roflumilast.

Non-limiting examples of therapeutic agents for HCV with which binding proteins of the invention can be combined include the following: Interferon-alpha-2a, Interferon-alpha-2b, Interferon-alpha con1, Interferon-alpha-n1, Pegylated interferon-alpha-2a, Pegylated interferon-alpha-2b, ribavirin, Peginterferon alfa-2b + ribavirin, Ursodeoxycholic Acid, Glycyrrhizic Acid,
 20 Thymalfasin, Maxamine, VX-497 and any compounds that are used to treat HCV through intervention with the following targets: HCV polymerase, HCV protease, HCV helicase, HCV IRES (internal ribosome entry site).

Non-limiting examples of therapeutic agents for Idiopathic Pulmonary Fibrosis with
 25 which binding proteins of the invention can be combined include the following: prednisone, azathioprine, albuterol, colchicine, albuterol sulfate, digoxin, gamma interferon, methylprednisolone sod succ, lorazepam, furosemide, lisinopril, nitroglycerin, spironolactone, cyclophosphamide, ipratropium bromide, actinomycin d, alteplase, fluticasone propionate, levofloxacin, metaproterenol sulfate, morphine sulfate, oxycodone hcl, potassium chloride,
 30 triamcinolone acetonide, tacrolimus anhydrous, calcium, interferon-alpha, methotrexate, mycophenolate mofetil, Interferon-gamma-1 β .

Non-limiting examples of therapeutic agents for Myocardial Infarction with which binding proteins of the invention can be combined include the following: aspirin, nitroglycerin, metoprolol tartrate, enoxaparin sodium, heparin sodium, clopidogrel bisulfate, carvedilol,
 35 atenolol, morphine sulfate, metoprolol succinate, warfarin sodium, lisinopril, isosorbide

mononitrate, digoxin, furosemide, simvastatin, ramipril, tenecteplase, enalapril maleate, torsemide, retavase, losartan potassium, quinapril hcl/mag carb, bumetanide, alteplase, enalaprilat, amiodarone hydrochloride, tirofiban hcl m-hydrate, diltiazem hydrochloride, captopril, irbesartan, valsartan, propranolol hydrochloride, fosinopril sodium, lidocaine hydrochloride, eptifibatide, cefazolin sodium, atropine sulfate, aminocaproic acid, spironolactone, interferon, sotalol hydrochloride, potassium chloride, docusate sodium, dobutamine hcl, alprazolam, pravastatin sodium, atorvastatin calcium, midazolam hydrochloride, meperidine hydrochloride, isosorbide dinitrate, epinephrine, dopamine hydrochloride, bivalirudin, rosuvastatin, ezetimibe/simvastatin, avasimibe, cariporide.

10 Non-limiting examples of therapeutic agents for Psoriasis with which binding proteins of the invention can be combined include the following: small molecule inhibitor of KDR, small molecule inhibitor of Tie-2, calcipotriene, clobetasol propionate, triamcinolone acetonide, halobetasol propionate, tazarotene, methotrexate, fluocinonide, betamethasone diprop augmented, fluocinolone acetonide, acitretin, tar shampoo, betamethasone valerate, mometasone furoate, ketoconazole, pramoxine/fluocinolone, hydrocortisone valerate, flurandrenolide, urea, betamethasone, clobetasol propionate/emoll, fluticasone propionate, azithromycin, hydrocortisone, moisturizing formula, folic acid, desonide, pimecrolimus, coal tar, diflorasone diacetate, etanercept folate, lactic acid, methoxsalen, hc/bismuth subgal/znox/resor, methylprednisolone acetate, prednisone, sunscreen, halcinonide, salicylic acid, anthralin, clocortolone pivalate, coal extract, coal tar/salicylic acid, coal tar/salicylic acid/sulfur, desoximetasone, diazepam, emollient, fluocinonide/emollient, mineral oil/castor oil/na lact, mineral oil/peanut oil, petroleum/isopropyl myristate, psoralen, salicylic acid, soap/tribromsalan, thimerosal/boric acid, celecoxib, infliximab, cyclosporine, alefacept, efalizumab, tacrolimus, pimecrolimus, PUVA, UVB, sulfasalazine.

25 Non-limiting examples of therapeutic agents for Psoriatic Arthritis with which binding proteins of the invention can be combined include the following: methotrexate, etanercept, rofecoxib, celecoxib, folic acid, sulfasalazine, naproxen, leflunomide, methylprednisolone acetate, indomethacin, hydroxychloroquine sulfate, prednisone, sulindac, betamethasone diprop augmented, infliximab, methotrexate, folate, triamcinolone acetonide, diclofenac, dimethylsulfoxide, piroxicam, diclofenac sodium, ketoprofen, meloxicam, methylprednisolone, nabumetone, tolmetin sodium, calcipotriene, cyclosporine, diclofenac sodium/misoprostol, fluocinonide, glucosamine sulfate, gold sodium thiomalate, hydrocodone bitartrate/apap, ibuprofen, risedronate sodium, sulfadiazine, thioguanine, valdecoxib, alefacept, efalizumab and bcl-2 inhibitors.

Non-limiting examples of therapeutic agents for Restenosis with which binding proteins of the invention can be combined include the following: sirolimus, paclitaxel, everolimus, tacrolimus, Zotarolimus, acetaminophen.

Non-limiting examples of therapeutic agents for Sciatica with which binding proteins of the invention can be combined include the following: hydrocodone bitartrate/apap, rofecoxib, cyclobenzaprine hcl, methylprednisolone, naproxen, ibuprofen, oxycodone hcl/acetaminophen, celecoxib, valdecoxib, methylprednisolone acetate, prednisone, codeine phosphate/apap, tramadol hcl/acetaminophen, metaxalone, meloxicam, methocarbamol, lidocaine hydrochloride, diclofenac sodium, gabapentin, dexamethasone, carisoprodol, ketorolac tromethamine, indomethacin, acetaminophen, diazepam, nabumetone, oxycodone hcl, tizanidine hcl, diclofenac sodium/misoprostol, propoxyphene napsylate/apap, asa/oxycod/oxycodone ter, ibuprofen/hydrocodone bit, tramadol hcl, etodolac, propoxyphene hcl, amitriptyline hcl, carisoprodol/codeine phos/asa, morphine sulfate, multivitamins, naproxen sodium, orphenadrine citrate, temazepam.

Examples of therapeutic agents for SLE (Lupus) in which binding proteins of the invention can be combined include the following: NSAIDS, for example, diclofenac, naproxen, ibuprofen, piroxicam, indomethacin; COX2 inhibitors, for example, Celecoxib, rofecoxib, valdecoxib; anti-malarials, for example, hydroxychloroquine; Steroids, for example, prednisone, prednisolone, budenoside, dexamethasone; Cytotoxics, for example, azathioprine, cyclophosphamide, mycophenolate mofetil, methotrexate; inhibitors of PDE4 or purine synthesis inhibitor, for example Cellcept. Binding proteins of the invention, may also be combined with agents such as sulfasalazine, 5-aminosalicylic acid, olsalazine, Imuran and agents which interfere with synthesis, production or action of proinflammatory cytokines such as IL-1, for example, caspase inhibitors like IL-1 β converting enzyme inhibitors and IL-1ra. Binding proteins of the invention may also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors; or molecules that target T cell activation molecules, for example, CTLA-4-IgG or anti-B7 family antibodies, anti-PD-1 family antibodies. Binding proteins of the invention, can be combined with IL-11 or anti-cytokine antibodies, for example, fonotolizumab (anti-IFN γ antibody), or anti-receptor antibodies, for example, anti-IL-6 receptor antibody and antibodies to B-cell surface molecules. Antibodies of the invention or antigen binding portion thereof may also be used with LJP 394 (abetimus), agents that deplete or inactivate B-cells, for example, Rituximab (anti-CD20 antibody), lymphostat-B (anti-BlyS antibody), TNF antagonists, for example, anti-TNF antibodies, Adalimumab (PCT Publication No. WO 97/29131; HUMIRA), CA2 (REMICADE), CDP 571, TNFR-Ig constructs, (p75TNFR-IgG (ENBREL)¹ and p55TNFR-IgG (LENERCEPT)) and bcl-2 inhibitors, because bcl-2 overexpression in transgenic mice has been demonstrated to cause a lupus like phenotype (see Marquina, Regina et al., Journal

of Immunology (2004), 172(11), 7177-7185), therefore inhibition is expected to have therapeutic effects.

The pharmaceutical compositions of the invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of a binding protein of the invention.

5 A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the binding protein may be determined by a person skilled in the art and may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the binding protein to elicit a desired response in the individual. A therapeutically effective amount is

10 also one in which any toxic or detrimental effects of the antibody, or antibody portion, are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically

15 effective amount.

Dosage regimens may be adjusted to provide the optimum desired response (*e.g.*, a therapeutic or prophylactic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous

20 to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the

25 invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

An exemplary, non-limiting range for a therapeutically or prophylactically effective

30 amount of a binding protein of the invention is 0.1-20 mg/kg, for example, 1-10 mg/kg. It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set

35 forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

It will be readily apparent to those skilled in the art that other suitable modifications and adaptations of the methods of the invention described herein are obvious and may be made using suitable equivalents without departing from the scope of the invention or the embodiments disclosed herein. Having now described the present invention in detail, the same will be more
5 clearly understood by reference to the following examples, which are included for purposes of illustration only and are not intended to be limiting of the invention.

V. Diagnostics

The disclosure herein also provides diagnostic applications. This is further elucidated below.

10 I. Method of Assay

The present disclosure also provides a method for determining the presence, amount or concentration of an analyte (or a fragment thereof) in a test sample using at least one DVD-Ig as described herein. Any suitable assay as is known in the art can be used in the method. Examples include, but are not limited to, immunoassay, such as sandwich immunoassay (e.g., monoclonal,
15 polyclonal and/or DVD-Ig sandwich immunoassays or any variation thereof (e.g., monoclonal/DVD-Ig, DVD-Ig/polyclonal, etc.), including radioisotope detection (radioimmunoassay (RIA)) and enzyme detection (enzyme immunoassay (EIA) or enzyme-linked immunosorbent assay (ELISA) (e.g., Quantikine ELISA assays, R&D Systems, Minneapolis, MN))), competitive inhibition immunoassay (e.g., forward and reverse), fluorescence polarization
20 immunoassay (FPIA), enzyme multiplied immunoassay technique (EMIT), bioluminescence resonance energy transfer (BRET), and homogeneous chemiluminescent assay, etc. In a SELDI-based immunoassay, a capture reagent that specifically binds an analyte (or a fragment thereof) of interest is attached to the surface of a mass spectrometry probe, such as a pre-activated protein chip array. The analyte (or a fragment thereof) is then specifically captured on the biochip, and
25 the captured analyte (or a fragment thereof) is detected by mass spectrometry. Alternatively, the analyte (or a fragment thereof) can be eluted from the capture reagent and detected by traditional MALDI (matrix-assisted laser desorption/ionization) or by SELDI. A chemiluminescent microparticle immunoassay, in particular one employing the ARCHITECT® automated analyzer (Abbott Laboratories, Abbott Park, IL), is an example of a preferred immunoassay.

30 Methods well-known in the art for collecting, handling and processing urine, blood, serum and plasma, and other body fluids, are used in the practice of the present disclosure, for instance, when a DVD-Ig as described herein is employed as an immunodiagnostic reagent and/or in an analyte immunoassay kit. The test sample can comprise further moieties in addition to the analyte of interest, such as antibodies, antigens, haptens, hormones, drugs, enzymes, receptors,
35 proteins, peptides, polypeptides, oligonucleotides and/or polynucleotides. For example, the sample can be a whole blood sample obtained from a subject. It can be necessary or desired that a

test sample, particularly whole blood, be treated prior to immunoassay as described herein, e.g., with a pretreatment reagent. Even in cases where pretreatment is not necessary (e.g., most urine samples), pretreatment optionally can be done (e.g., as part of a regimen on a commercial platform).

5 The pretreatment reagent can be any reagent appropriate for use with the immunoassay and kits of the invention. The pretreatment optionally comprises: (a) one or more solvents (e.g., methanol and ethylene glycol) and optionally, salt, (b) one or more solvents and salt, and optionally, detergent, (c) detergent, or (d) detergent and salt. Pretreatment reagents are known in the art, and such pretreatment can be employed, e.g., as used for assays on Abbott TDx, AxSYM®, and
10 ARCHITECT® analyzers (Abbott Laboratories, Abbott Park, IL), as described in the literature (see, e.g., Yatscoff et al., Abbott TDx Monoclonal Antibody Assay Evaluated for Measuring Cyclosporine in Whole Blood, Clin. Chem. 36: 1969-1973 (1990), and Wallemacq et al., Evaluation of the New AxSYM Cyclosporine Assay: Comparison with TDx Monoclonal Whole Blood and EMIT Cyclosporine Assays, Clin. Chem. 45: 432-435 (1999)), and/or as commercially
15 available. Additionally, pretreatment can be done as described in Abbott's U.S. Pat. No. 5,135,875, European Pat. Pub. No. 0 471 293, U.S. Provisional Pat. App. 60/878,017, filed December 29, 2006, and U.S. Pat. App. Pub. No. 2008/0020401 (incorporated by reference in its entirety for its teachings regarding pretreatment). The pretreatment reagent can be a heterogeneous agent or a homogeneous agent.

20 With use of a heterogeneous pretreatment reagent, the pretreatment reagent precipitates analyte binding protein (e.g., protein that can bind to an analyte or a fragment thereof) present in the sample. Such a pretreatment step comprises removing any analyte binding protein by separating from the precipitated analyte binding protein the supernatant of the mixture formed by addition of the pretreatment agent to sample. In such an assay, the supernatant of the mixture
25 absent any binding protein is used in the assay, proceeding directly to the antibody capture step.

With use of a homogeneous pretreatment reagent there is no such separation step. The entire mixture of test sample and pretreatment reagent are contacted with a labeled specific binding partner for analyte (or a fragment thereof), such as a labeled anti-analyte antibody (or an antigenically reactive fragment thereof). The pretreatment reagent employed for such an assay
30 typically is diluted in the pretreated test sample mixture, either before or during capture by the first specific binding partner. Despite such dilution, a certain amount of the pretreatment reagent is still present (or remains) in the test sample mixture during capture. According to the invention, the labeled specific binding partner can be a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof).

35 In a heterogeneous format, after the test sample is obtained from a subject, a first mixture is prepared. The mixture contains the test sample being assessed for an analyte (or a fragment thereof) and a first specific binding partner, wherein the first specific binding partner and any

analyte contained in the test sample form a first specific binding partner-analyte complex.

Preferably, the first specific binding partner is an anti-analyte antibody or a fragment thereof. The first specific binding partner can be a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein. The order in which the test sample and the first specific binding partner are added to form the mixture is not critical. Preferably, the first specific binding partner is immobilized on a solid phase. The solid phase used in the immunoassay (for the first specific binding partner and, optionally, the second specific binding partner) can be any solid phase known in the art, such as, but not limited to, a magnetic particle, a bead, a test tube, a microtiter plate, a cuvette, a membrane, a scaffolding molecule, a film, a filter paper, a disc and a chip.

After the mixture containing the first specific binding partner-analyte complex is formed, any unbound analyte is removed from the complex using any technique known in the art. For example, the unbound analyte can be removed by washing. Desirably, however, the first specific binding partner is present in excess of any analyte present in the test sample, such that all analyte that is present in the test sample is bound by the first specific binding partner.

After any unbound analyte is removed, a second specific binding partner is added to the mixture to form a first specific binding partner-analyte-second specific binding partner complex. The second specific binding partner is preferably an anti-analyte antibody that binds to an epitope on analyte that differs from the epitope on analyte bound by the first specific binding partner. Moreover, also preferably, the second specific binding partner is labeled with or contains a detectable label as described above. The second specific binding partner can be a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein.

Any suitable detectable label as is known in the art can be used. For example, the detectable label can be a radioactive label (such as ^3H , ^{125}I , ^{35}S , ^{14}C , ^{32}P , and ^{33}P), an enzymatic label (such as horseradish peroxidase, alkaline peroxidase, glucose 6-phosphate dehydrogenase, and the like), a chemiluminescent label (such as acridinium esters, thioesters, or sulfonamides; luminol, isoluminol, phenanthridinium esters, and the like), a fluorescent label (such as fluorescein (e.g., 5-fluorescein, 6-carboxyfluorescein, 3'-carboxyfluorescein, 5(6)-carboxyfluorescein, 6-hexachloro-fluorescein, 6-tetrachlorofluorescein, fluorescein isothiocyanate, and the like)), rhodamine, phycobiliproteins, R-phycoerythrin, quantum dots (e.g., zinc sulfide-capped cadmium selenide), a thermometric label, or an immuno-polymerase chain reaction label. An introduction to labels, labeling procedures and detection of labels is found in Polak and Van Noorden, *Introduction to Immunocytochemistry*, 2nd ed., Springer Verlag, N.Y. (1997), and in Haugland, *Handbook of Fluorescent Probes and Research Chemicals* (1996), which is a combined handbook and catalogue published by Molecular Probes, Inc., Eugene, Oregon. A fluorescent label can be used in FPIA (see, e.g., U.S. Patent Nos. 5,593,896, 5,573,904, 5,496,925, 5,359,093, and 5,352,803, which are hereby incorporated by reference in their entireties). An acridinium compound can be used as a detectable label in a homogeneous or

heterogeneous chemiluminescent assay (see, e.g., Adamczyk et al., *Bioorg. Med. Chem. Lett.* 16: 1324-1328 (2006); Adamczyk et al., *Bioorg. Med. Chem. Lett.* 4: 2313-2317 (2004); Adamczyk et al., *Bioorg. Med. Chem. Lett.* 14: 3917-3921 (2004); and Adamczyk et al., *Org. Lett.* 5: 3779-3782 (2003)).

5 A preferred acridinium compound is an acridinium-9-carboxamide. Methods for preparing acridinium 9-carboxamides are described in Mattingly, J. *Biolumin. Chemilumin.* 6: 107-114 (1991); Adamczyk et al., *J. Org. Chem.* 63: 5636-5639 (1998); Adamczyk et al., *Tetrahedron* 55: 10899-10914 (1999); Adamczyk et al., *Org. Lett.* 1: 779-781 (1999); Adamczyk et al., *Bioconjugate Chem.* 11: 714-724 (2000); Mattingly et al., In *Luminescence Biotechnology: Instruments and Applications*; Dyke, K. V. Ed.; CRC Press: Boca Raton, pp. 77-105 (2002); Adamczyk et al., *Org. Lett.* 5: 3779-3782 (2003); and U.S. Pat. Nos. 5,468,646, 5,543,524 and 5,783,699 (each of which is incorporated herein by reference in its entirety for its teachings regarding same). Another preferred acridinium compound is an acridinium-9-carboxylate aryl ester. An example of an acridinium-9-carboxylate aryl ester is 10-methyl-9-
15 (phenoxy carbonyl)acridinium fluorosulfonate (available from Cayman Chemical, Ann Arbor, MI). Methods for preparing acridinium 9-carboxylate aryl esters are described in McCapra et al., *Photochem. Photobiol.* 4: 1111-21 (1965); Razavi et al., *Luminescence* 15: 245-249 (2000); Razavi et al., *Luminescence* 15: 239-244 (2000); and U.S. Patent No. 5,241,070 (each of which is incorporated herein by reference in its entirety for its teachings regarding same). Further details
20 regarding acridinium-9-carboxylate aryl ester and its use are set forth in US 2008-0248493.

Chemiluminescent assays (e.g., using acridinium as described above or other chemiluminescent agents) can be performed in accordance with the methods described in Adamczyk et al., *Anal. Chim. Acta* 579(1): 61-67 (2006). While any suitable assay format can be used, a microplate chemiluminometer (Mithras LB-940, Berthold Technologies U.S.A., LLC, Oak
25 Ridge, TN) enables the assay of multiple samples of small volumes rapidly.

The order in which the test sample and the specific binding partner(s) are added to form the mixture for chemiluminescent assay is not critical. If the first specific binding partner is detectably labeled with a chemiluminescent agent such as an acridinium compound, detectably labeled first specific binding partner-analyte complexes form. Alternatively, if a second specific
30 binding partner is used and the second specific binding partner is detectably labeled with a chemiluminescent agent such as an acridinium compound, detectably labeled first specific binding partner-analyte-second specific binding partner complexes form. Any unbound specific binding partner, whether labeled or unlabeled, can be removed from the mixture using any technique known in the art, such as washing.

35 Hydrogen peroxide can be generated in situ in the mixture or provided or supplied to the mixture (e.g., the source of the hydrogen peroxide being one or more buffers or other solutions that are known to contain hydrogen peroxide) before, simultaneously with, or after the addition of

an above-described acridinium compound. Hydrogen peroxide can be generated in situ in a number of ways such as would be apparent to one skilled in the art.

Upon the simultaneous or subsequent addition of at least one basic solution to the sample, a detectable signal, namely, a chemiluminescent signal, indicative of the presence of analyte is generated. The basic solution contains at least one base and has a pH greater than or equal to 10, preferably, greater than or equal to 12. Examples of basic solutions include, but are not limited to, sodium hydroxide, potassium hydroxide, calcium hydroxide, ammonium hydroxide, magnesium hydroxide, sodium carbonate, sodium bicarbonate, calcium hydroxide, calcium carbonate, and calcium bicarbonate. The amount of basic solution added to the sample depends on the concentration of the basic solution. Based on the concentration of the basic solution used, one skilled in the art can easily determine the amount of basic solution to add to the sample.

The chemiluminescent signal that is generated can be detected using routine techniques known to those skilled in the art. Based on the intensity of the signal generated, the amount of analyte in the sample can be quantified. Specifically, the amount of analyte in the sample is proportional to the intensity of the signal generated. The amount of analyte present can be quantified by comparing the amount of light generated to a standard curve for analyte or by comparison to a reference standard. The standard curve can be generated using serial dilutions or solutions of known concentrations of analyte by mass spectroscopy, gravimetric methods, and other techniques known in the art. While the above is described with emphasis on use of an acridinium compound as the chemiluminescent agent, one of ordinary skill in the art can readily adapt this description for use of other chemiluminescent agents.

Analyte immunoassays generally can be conducted using any format known in the art, such as, but not limited to, a sandwich format. Specifically, in one immunoassay format, at least two antibodies are employed to separate and quantify analyte, such as human analyte, or a fragment thereof in a sample. More specifically, the at least two antibodies bind to different epitopes on an analyte (or a fragment thereof) forming an immune complex, which is referred to as a “sandwich.” Generally, in the immunoassays one or more antibodies can be used to capture the analyte (or a fragment thereof) in the test sample (these antibodies are frequently referred to as a “capture” antibody or “capture” antibodies) and one or more antibodies can be used to bind a detectable (namely, quantifiable) label to the sandwich (these antibodies are frequently referred to as the “detection antibody,” the “detection antibodies,” the “conjugate,” or the “conjugates”). Thus, in the context of a sandwich immunoassay format, a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein can be used as a capture antibody, a detection antibody, or both. For example, one DVD-Ig having a domain that can bind a first epitope on an analyte (or a fragment thereof) can be used as a capture antibody and/or another DVD-Ig having a domain that can bind a second epitope on an analyte (or a fragment thereof) can be used as a detection antibody. In this regard, a DVD-Ig having a first domain that can bind a first epitope on

an analyte (or a fragment thereof) and a second domain that can bind a second epitope on an analyte (or a fragment thereof) can be used as a capture antibody and/or a detection antibody. Alternatively, one DVD-Ig having a first domain that can bind an epitope on a first analyte (or a fragment thereof) and a second domain that can bind an epitope on a second analyte (or a fragment thereof) can be used as a capture antibody and/or a detection antibody to detect, and optionally quantify, two or more analytes. In the event that an analyte can be present in a sample in more than one form, such as a monomeric form and a dimeric/multimeric form, which can be homomeric or heteromeric, one DVD-Ig having a domain that can bind an epitope that is only exposed on the monomeric form and another DVD-Ig having a domain that can bind an epitope on a different part of a dimeric/multimeric form can be used as capture antibodies and/or detection antibodies, thereby enabling the detection, and optional quantification, of different forms of a given analyte. Furthermore, employing DVD-Igs with differential affinities within a single DVD-Ig and/or between DVD-Igs can provide an avidity advantage. In the context of immunoassays as described herein, it generally may be helpful or desired to incorporate one or more linkers within the structure of a DVD-Ig. When present, optimally the linker should be of sufficient length and structural flexibility to enable binding of an epitope by the inner domains as well as binding of another epitope by the outer domains. In this regard, if a DVD-Ig can bind two different analytes and one analyte is larger than the other, desirably the larger analyte is bound by the outer domains.

Generally speaking, a sample being tested for (for example, suspected of containing) analyte (or a fragment thereof) can be contacted with at least one capture antibody (or antibodies) and at least one detection antibody (which can be a second detection antibody or a third detection antibody or even a successively numbered antibody, e.g., as where the capture and/or detection antibody comprise multiple antibodies) either simultaneously or sequentially and in any order. For example, the test sample can be first contacted with at least one capture antibody and then (sequentially) with at least one detection antibody. Alternatively, the test sample can be first contacted with at least one detection antibody and then (sequentially) with at least one capture antibody. In yet another alternative, the test sample can be contacted simultaneously with a capture antibody and a detection antibody.

In the sandwich assay format, a sample suspected of containing analyte (or a fragment thereof) is first brought into contact with at least one first capture antibody under conditions that allow the formation of a first antibody/analyte complex. If more than one capture antibody is used, a first capture antibody/analyte complex comprising two or more capture antibodies is formed. In a sandwich assay, the antibodies, i.e., preferably, the at least one capture antibody, are used in molar excess amounts of the maximum amount of analyte (or a fragment thereof) expected in the test sample. For example, from about 5 μ g to about 1 mg of antibody per mL of buffer (e.g., microparticle coating buffer) can be used.

Competitive inhibition immunoassays, which are often used to measure small analytes because binding by only one antibody is required, comprise sequential and classic formats. In a sequential competitive inhibition immunoassay a capture antibody to an analyte of interest is coated onto a well of a microtiter plate or other solid support. When the sample containing the analyte of interest is added to the well, the analyte of interest binds to the capture antibody. After washing, a known amount of labeled (e.g., biotin or horseradish peroxidase (HRP)) analyte is added to the well. A substrate for an enzymatic label is necessary to generate a signal. An example of a suitable substrate for HRP is 3,3',5,5'-tetramethylbenzidine (TMB). After washing, the signal generated by the labeled analyte is measured and is inversely proportional to the amount of analyte in the sample. In a classic competitive inhibition immunoassay an antibody to an analyte of interest is coated onto a solid support (e.g., a well of a microtiter plate). However, unlike the sequential competitive inhibition immunoassay, the sample and the labeled analyte are added to the well at the same time. Any analyte in the sample competes with labeled analyte for binding to the capture antibody. After washing, the signal generated by the labeled analyte is measured and is inversely proportional to the amount of analyte in the sample.

Optionally, prior to contacting the test sample with the at least one capture antibody (for example, the first capture antibody), the at least one capture antibody can be bound to a solid support, which facilitates the separation of the first antibody/analyte (or a fragment thereof) complex from the test sample. The substrate to which the capture antibody is bound can be any suitable solid support or solid phase that facilitates separation of the capture antibody-analyte complex from the sample.

Examples include a well of a plate, such as a microtiter plate, a test tube, a porous gel (e.g., silica gel, agarose, dextran, or gelatin), a polymeric film (e.g., polyacrylamide), beads (e.g., polystyrene beads or magnetic beads), a strip of a filter/membrane (e.g., nitrocellulose or nylon), microparticles (e.g., latex particles, magnetizable microparticles (e.g., microparticles having ferric oxide or chromium oxide cores and homo- or hetero-polymeric coats and radii of about 1-10 microns). The substrate can comprise a suitable porous material with a suitable surface affinity to bind antigens and sufficient porosity to allow access by detection antibodies. A microporous material is generally preferred, although a gelatinous material in a hydrated state can be used. Such porous substrates are preferably in the form of sheets having a thickness of about 0.01 to about 0.5 mm, preferably about 0.1 mm. While the pore size may vary quite a bit, preferably the pore size is from about 0.025 to about 15 microns, more preferably from about 0.15 to about 15 microns. The surface of such substrates can be activated by chemical processes that cause covalent linkage of an antibody to the substrate. Irreversible binding, generally by adsorption through hydrophobic forces, of the antigen or the antibody to the substrate results; alternatively, a chemical coupling agent or other means can be used to bind covalently the antibody to the substrate, provided that such binding does not interfere with the ability of the antibody to bind to

analyte. Alternatively, the antibody can be bound with microparticles, which have been previously coated with streptavidin (e.g., DYNAL® Magnetic Beads, Invitrogen, Carlsbad, CA) or biotin (e.g., using Power-Bind™-SA-MP streptavidin-coated microparticles (Seradyn, Indianapolis, IN)) or anti-species-specific monoclonal antibodies. If necessary, the substrate can be derivatized to allow reactivity with various functional groups on the antibody. Such derivatization requires the use of certain coupling agents, examples of which include, but are not limited to, maleic anhydride, N-hydroxysuccinimide, and 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide. If desired, one or more capture reagents, such as antibodies (or fragments thereof), each of which is specific for analyte(s) can be attached to solid phases in different physical or addressable locations (e.g., such as in a biochip configuration (see, e.g., U.S. Pat. No. 6,225,047; Int'l Pat. App. Pub. No. WO 99/51773; U.S. Pat. No. 6,329,209; Int'l Pat. App. Pub. No. WO 00/56934, and U.S. Pat. No. 5,242,828). If the capture reagent is attached to a mass spectrometry probe as the solid support, the amount of analyte bound to the probe can be detected by laser desorption ionization mass spectrometry. Alternatively, a single column can be packed with different beads, which are derivatized with the one or more capture reagents, thereby capturing the analyte in a single place (see, antibody-derivatized, bead-based technologies, e.g., the xMAP technology of Luminex (Austin, TX)).

After the test sample being assayed for analyte (or a fragment thereof) is brought into contact with the at least one capture antibody (for example, the first capture antibody), the mixture is incubated in order to allow for the formation of a first antibody (or multiple antibody)-analyte (or a fragment thereof) complex. The incubation can be carried out at a pH of from about 4.5 to about 10.0, at a temperature of from about 2°C to about 45°C, and for a period from at least about one (1) minute to about eighteen (18) hours, preferably from about 1 to about 24 minutes, most preferably for about 4 to about 18 minutes. The immunoassay described herein can be conducted in one step (meaning the test sample, at least one capture antibody and at least one detection antibody are all added sequentially or simultaneously to a reaction vessel) or in more than one step, such as two steps, three steps, etc.

After formation of the (first or multiple) capture antibody/analyte (or a fragment thereof) complex, the complex is then contacted with at least one detection antibody under conditions which allow for the formation of a (first or multiple) capture antibody/analyte (or a fragment thereof)/second detection antibody complex). While captioned for clarity as the "second" antibody (e.g., second detection antibody), in fact, where multiple antibodies are used for capture and/or detection, the at least one detection antibody can be the second, third, fourth, etc. antibodies used in the immunoassay. If the capture antibody/analyte (or a fragment thereof) complex is contacted with more than one detection antibody, then a (first or multiple) capture antibody/analyte (or a fragment thereof)/(multiple) detection antibody complex is formed. As with the capture antibody (e.g., the first capture antibody), when the at least one (e.g., second and

any subsequent) detection antibody is brought into contact with the capture antibody/analyte (or a fragment thereof) complex, a period of incubation under conditions similar to those described above is required for the formation of the (first or multiple) capture antibody/analyte (or a fragment thereof)/(second or multiple) detection antibody complex. Preferably, at least one
5 detection antibody contains a detectable label. The detectable label can be bound to the at least one detection antibody (e.g., the second detection antibody) prior to, simultaneously with, or after the formation of the (first or multiple) capture antibody/analyte (or a fragment thereof)/(second or multiple) detection antibody complex. Any detectable label known in the art can be used (see discussion above, including of the Polak and Van Noorden (1997) and Haugland (1996)
10 references).

The detectable label can be bound to the antibodies either directly or through a coupling agent. An example of a coupling agent that can be used is EDAC (1-ethyl-3-(3-dimethylaminopropyl) carbodiimide, hydrochloride), which is commercially available from Sigma-Aldrich, St. Louis, MO. Other coupling agents that can be used are known in the art. Methods for binding a
15 detectable label to an antibody are known in the art. Additionally, many detectable labels can be purchased or synthesized that already contain end groups that facilitate the coupling of the detectable label to the antibody, such as CPSP-Acrininium Ester (i.e., 9-[N-tosyl-N-(3-carboxypropyl)]-10-(3-sulfopropyl)acridinium carboxamide) or SPSP-Acrininium Ester (i.e., N10-(3-sulfopropyl)-N-(3-sulfopropyl)-acridinium-9-carboxamide).

20 The (first or multiple) capture antibody/analyte/(second or multiple) detection antibody complex can be, but does not have to be, separated from the remainder of the test sample prior to quantification of the label. For example, if the at least one capture antibody (e.g., the first capture antibody) is bound to a solid support, such as a well or a bead, separation can be accomplished by removing the fluid (of the test sample) from contact with the solid support. Alternatively, if the at
25 least first capture antibody is bound to a solid support, it can be simultaneously contacted with the analyte-containing sample and the at least one second detection antibody to form a first (multiple) antibody/analyte/second (multiple) antibody complex, followed by removal of the fluid (test sample) from contact with the solid support. If the at least one first capture antibody is not bound to a solid support, then the (first or multiple) capture antibody/analyte/(second or multiple)
30 detection antibody complex does not have to be removed from the test sample for quantification of the amount of the label.

After formation of the labeled capture antibody/analyte/detection antibody complex (e.g., the first capture antibody/analyte/second detection antibody complex), the amount of label in the complex is quantified using techniques known in the art. For example, if an enzymatic label is
35 used, the labeled complex is reacted with a substrate for the label that gives a quantifiable reaction such as the development of color. If the label is a radioactive label, the label is quantified using appropriate means, such as a scintillation counter. If the label is a fluorescent label, the label is

quantified by stimulating the label with a light of one color (which is known as the “excitation wavelength”) and detecting another color (which is known as the “emission wavelength”) that is emitted by the label in response to the stimulation. If the label is a chemiluminescent label, the label is quantified by detecting the light emitted either visually or by using luminometers, x-ray
5 film, high speed photographic film, a CCD camera, etc. Once the amount of the label in the complex has been quantified, the concentration of analyte or a fragment thereof in the test sample is determined by appropriate means, such as by use of a standard curve that has been generated using serial dilutions of analyte or a fragment thereof of known concentration. Other than using serial dilutions of analyte or a fragment thereof, the standard curve can be generated
10 gravimetrically, by mass spectroscopy and by other techniques known in the art.

In a chemiluminescent microparticle assay employing the ARCHITECT® analyzer, the conjugate diluent pH should be about 6.0 +/- 0.2, the microparticle coating buffer should be maintained at about room temperature (i.e., at from about 17 to about 27 °C), the microparticle coating buffer pH should be about 6.5 +/- 0.2, and the microparticle diluent pH should be about
15 7.8 +/- 0.2. Solids preferably are less than about 0.2%, such as less than about 0.15%, less than about 0.14%, less than about 0.13%, less than about 0.12%, or less than about 0.11%, such as about 0.10%.

FPIAs are based on competitive binding immunoassay principles. A fluorescently labeled compound, when excited by a linearly polarized light, will emit fluorescence having a degree of
20 polarization inversely proportional to its rate of rotation. When a fluorescently labeled tracer-antibody complex is excited by a linearly polarized light, the emitted light remains highly polarized because the fluorophore is constrained from rotating between the time light is absorbed and the time light is emitted. When a “free” tracer compound (i.e., a compound that is not bound to an antibody) is excited by linearly polarized light, its rotation is much faster than the
25 corresponding tracer-antibody conjugate produced in a competitive binding immunoassay. FPIAs are advantageous over RIAs inasmuch as there are no radioactive substances requiring special handling and disposal. In addition, FPIAs are homogeneous assays that can be easily and rapidly performed.

In view of the above, a method of determining the presence, amount, or concentration of
30 analyte (or a fragment thereof) in a test sample is provided. The method comprises assaying the test sample for an analyte (or a fragment thereof) by an assay (i) employing (i') at least one of an antibody, a fragment of an antibody that can bind to an analyte, a variant of an antibody that can bind to an analyte, a fragment of a variant of an antibody that can bind to an analyte, and a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) that can bind to an analyte, and (ii')
35 at least one detectable label and (ii) comprising comparing a signal generated by the detectable label as a direct or indirect indication of the presence, amount or concentration of analyte (or a fragment thereof) in the test sample to a signal generated as a direct or indirect indication of the

presence, amount or concentration of analyte (or a fragment thereof) in a control or calibrator. The calibrator is optionally part of a series of calibrators, in which each of the calibrators differs from the other calibrators by the concentration of analyte.

The method can comprise (i) contacting the test sample with at least one first specific
5 binding partner for analyte (or a fragment thereof) selected from the group consisting of an antibody, a fragment of an antibody that can bind to an analyte, a variant of an antibody that can bind to an analyte, a fragment of a variant of an antibody that can bind to an analyte, and a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) that can bind to an analyte so as to form a first specific binding partner/analyte (or fragment thereof) complex, (ii) contacting the first
10 specific binding partner/analyte (or fragment thereof) complex with at least one second specific binding partner for analyte (or fragment thereof) selected from the group consisting of a detectably labeled anti-analyte antibody, a detectably labeled fragment of an anti-analyte antibody that can bind to analyte, a detectably labeled variant of an anti-analyte antibody that can bind to analyte, a detectably labeled fragment of a variant of an anti-analyte antibody that can bind to
15 analyte, and a detectably labeled DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) so as to form a first specific binding partner/analyte (or fragment thereof)/second specific binding partner complex, and (iii) determining the presence, amount or concentration of analyte in the test sample by detecting or measuring the signal generated by the detectable label in the first specific binding partner/analyte (or fragment thereof)/second specific binding partner complex
20 formed in (ii). A method in which at least one first specific binding partner for analyte (or a fragment thereof) and/or at least one second specific binding partner for analyte (or a fragment thereof) is a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) as described herein can be preferred.

Alternatively, the method can comprise contacting the test sample with at least one first
25 specific binding partner for analyte (or a fragment thereof) selected from the group consisting of an antibody, a fragment of an antibody that can bind to an analyte, a variant of an antibody that can bind to an analyte, a fragment of a variant of an antibody that can bind to an analyte, and a DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof) and simultaneously or sequentially, in either order, contacting the test sample with at least one second specific binding
30 partner, which can compete with analyte (or a fragment thereof) for binding to the at least one first specific binding partner and which is selected from the group consisting of a detectably labeled analyte, a detectably labeled fragment of analyte that can bind to the first specific binding partner, a detectably labeled variant of analyte that can bind to the first specific binding partner, and a detectably labeled fragment of a variant of analyte that can bind to the first specific binding
35 partner. Any analyte (or a fragment thereof) present in the test sample and the at least one second specific binding partner compete with each other to form a first specific binding partner/analyte (or fragment thereof) complex and a first specific binding partner/second specific binding partner

complex, respectively. The method further comprises determining the presence, amount or concentration of analyte in the test sample by detecting or measuring the signal generated by the detectable label in the first specific binding partner/second specific binding partner complex formed in (ii), wherein the signal generated by the detectable label in the first specific binding partner/second specific binding partner complex is inversely proportional to the amount or concentration of analyte in the test sample.

The above methods can further comprise diagnosing, prognosticating, or assessing the efficacy of a therapeutic/prophylactic treatment of a patient from whom the test sample was obtained. If the method further comprises assessing the efficacy of a therapeutic/prophylactic treatment of the patient from whom the test sample was obtained, the method optionally further comprises modifying the therapeutic/prophylactic treatment of the patient as needed to improve efficacy. The method can be adapted for use in an automated system or a semi-automated system.

With regard to the methods of assay (and kit therefor), it may be possible to employ commercially available anti-analyte antibodies or methods for production of anti-analyte as described in the literature. Commercial supplies of various antibodies include, but are not limited to, Santa Cruz Biotechnology Inc. (Santa Cruz, CA), GenWay Biotech, Inc. (San Diego, CA), and R&D Systems (RDS; Minneapolis, MN).

Generally, a predetermined level can be employed as a benchmark against which to assess results obtained upon assaying a test sample for analyte or a fragment thereof, e.g., for detecting disease or risk of disease. Generally, in making such a comparison, the predetermined level is obtained by running a particular assay a sufficient number of times and under appropriate conditions such that a linkage or association of analyte presence, amount or concentration with a particular stage or endpoint of a disease, disorder or condition or with particular clinical indicia can be made. Typically, the predetermined level is obtained with assays of reference subjects (or populations of subjects). The analyte measured can include fragments thereof, degradation products thereof, and/or enzymatic cleavage products thereof.

In particular, with respect to a predetermined level as employed for monitoring disease progression and/or treatment, the amount or concentration of analyte or a fragment thereof may be “unchanged,” “favorable” (or “favorably altered”), or “unfavorable” (or “unfavorably altered”). “Elevated” or “increased” refers to an amount or a concentration in a test sample that is higher than a typical or normal level or range (e.g., predetermined level), or is higher than another reference level or range (e.g., earlier or baseline sample). The term “lowered” or “reduced” refers to an amount or a concentration in a test sample that is lower than a typical or normal level or range (e.g., predetermined level), or is lower than another reference level or range (e.g., earlier or baseline sample). The term “altered” refers to an amount or a concentration in a sample that is altered (increased or decreased) over a typical or normal level or range (e.g., predetermined level), or over another reference level or range (e.g., earlier or baseline sample).

The typical or normal level or range for analyte is defined in accordance with standard practice. Because the levels of analyte in some instances will be very low, a so-called altered level or alteration can be considered to have occurred when there is any net change as compared to the typical or normal level or range, or reference level or range, that cannot be explained by experimental error or sample variation. Thus, the level measured in a particular sample will be compared with the level or range of levels determined in similar samples from a so-called normal subject. In this context, a “normal subject” is an individual with no detectable disease, for example, and a “normal” (sometimes termed “control”) patient or population is/are one(s) that exhibit(s) no detectable disease, respectively, for example. Furthermore, given that analyte is not routinely found at a high level in the majority of the human population, a “normal subject” can be considered an individual with no substantial detectable increased or elevated amount or concentration of analyte, and a “normal” (sometimes termed “control”) patient or population is/are one(s) that exhibit(s) no substantial detectable increased or elevated amount or concentration of analyte. An “apparently normal subject” is one in which analyte has not yet been or currently is being assessed. The level of an analyte is said to be “elevated” when the analyte is normally undetectable (e.g., the normal level is zero, or within a range of from about 25 to about 75 percentiles of normal populations), but is detected in a test sample, as well as when the analyte is present in the test sample at a higher than normal level. Thus, inter alia, the disclosure provides a method of screening for a subject having, or at risk of having, a particular disease, disorder, or condition. The method of assay can also involve the assay of other markers and the like.

Accordingly, the methods described herein also can be used to determine whether or not a subject has or is at risk of developing a given disease, disorder or condition. Specifically, such a method can comprise the steps of:

- (a) determining the concentration or amount in a test sample from a subject of analyte (or a fragment thereof) (e.g., using the methods described herein, or methods known in the art); and
- (b) comparing the concentration or amount of analyte (or a fragment thereof) determined in step (a) with a predetermined level, wherein, if the concentration or amount of analyte determined in step (a) is favorable with respect to a predetermined level, then the subject is determined not to have or be at risk for a given disease, disorder or condition. However, if the concentration or amount of analyte determined in step (a) is unfavorable with respect to the predetermined level, then the subject is determined to have or be at risk for a given disease, disorder or condition.

Additionally, provided herein is method of monitoring the progression of disease in a subject. Optimally the method comprising the steps of:

- (a) determining the concentration or amount in a test sample from a subject of analyte;
- (b) determining the concentration or amount in a later test sample from the subject of analyte; and
- (c) comparing the concentration or amount of analyte as determined in step (b) with the concentration or amount of analyte determined in step (a), wherein if the concentration or amount

determined in step (b) is unchanged or is unfavorable when compared to the concentration or amount of analyte determined in step (a), then the disease in the subject is determined to have continued, progressed or worsened. By comparison, if the concentration or amount of analyte as determined in step (b) is favorable when compared to the concentration or amount of analyte as determined in step (a), then the disease in the subject is determined to have discontinued, regressed or improved.

Optionally, the method further comprises comparing the concentration or amount of analyte as determined in step (b), for example, with a predetermined level. Further, optionally the method comprises treating the subject with one or more pharmaceutical compositions for a period of time if the comparison shows that the concentration or amount of analyte as determined in step (b), for example, is unfavorably altered with respect to the predetermined level.

Still further, the methods can be used to monitor treatment in a subject receiving treatment with one or more pharmaceutical compositions. Specifically, such methods involve providing a first test sample from a subject before the subject has been administered one or more pharmaceutical compositions. Next, the concentration or amount in a first test sample from a subject of analyte is determined (e.g., using the methods described herein or as known in the art). After the concentration or amount of analyte is determined, optionally the concentration or amount of analyte is then compared with a predetermined level. If the concentration or amount of analyte as determined in the first test sample is lower than the predetermined level, then the subject is not treated with one or more pharmaceutical compositions. However, if the concentration or amount of analyte as determined in the first test sample is higher than the predetermined level, then the subject is treated with one or more pharmaceutical compositions for a period of time. The period of time that the subject is treated with the one or more pharmaceutical compositions can be determined by one skilled in the art (for example, the period of time can be from about seven (7) days to about two years, preferably from about fourteen (14) days to about one (1) year).

During the course of treatment with the one or more pharmaceutical compositions, second and subsequent test samples are then obtained from the subject. The number of test samples and the time in which said test samples are obtained from the subject are not critical. For example, a second test sample could be obtained seven (7) days after the subject is first administered the one or more pharmaceutical compositions, a third test sample could be obtained two (2) weeks after the subject is first administered the one or more pharmaceutical compositions, a fourth test sample could be obtained three (3) weeks after the subject is first administered the one or more pharmaceutical compositions, a fifth test sample could be obtained four (4) weeks after the subject is first administered the one or more pharmaceutical compositions, etc.

After each second or subsequent test sample is obtained from the subject, the concentration or amount of analyte is determined in the second or subsequent test sample is

determined (e.g., using the methods described herein or as known in the art). The concentration or amount of analyte as determined in each of the second and subsequent test samples is then compared with the concentration or amount of analyte as determined in the first test sample (e.g., the test sample that was originally optionally compared to the predetermined level). If the concentration or amount of analyte as determined in step (c) is favorable when compared to the concentration or amount of analyte as determined in step (a), then the disease in the subject is determined to have discontinued, regressed or improved, and the subject should continue to be administered the one or pharmaceutical compositions of step (b). However, if the concentration or amount determined in step (c) is unchanged or is unfavorable when compared to the concentration or amount of analyte as determined in step (a), then the disease in the subject is determined to have continued, progressed or worsened, and the subject should be treated with a higher concentration of the one or more pharmaceutical compositions administered to the subject in step (b) or the subject should be treated with one or more pharmaceutical compositions that are different from the one or more pharmaceutical compositions administered to the subject in step (b). Specifically, the subject can be treated with one or more pharmaceutical compositions that are different from the one or more pharmaceutical compositions that the subject had previously received to decrease or lower said subject's analyte level.

Generally, for assays in which repeat testing may be done (e.g., monitoring disease progression and/or response to treatment), a second or subsequent test sample is obtained at a period in time after the first test sample has been obtained from the subject. Specifically, a second test sample from the subject can be obtained minutes, hours, days, weeks or years after the first test sample has been obtained from the subject. For example, the second test sample can be obtained from the subject at a time period of about 1 minute, about 5 minutes, about 10 minutes, about 15 minutes, about 30 minutes, about 45 minutes, about 60 minutes, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, about 21 hours, about 22 hours, about 23 hours, about 24 hours, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days, about 7 days, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, about 7 weeks, about 8 weeks, about 9 weeks, about 10 weeks, about 11 weeks, about 12 weeks, about 13 weeks, about 14 weeks, about 15 weeks, about 16 weeks, about 17 weeks, about 18 weeks, about 19 weeks, about 20 weeks, about 21 weeks, about 22 weeks, about 23 weeks, about 24 weeks, about 25 weeks, about 26 weeks, about 27 weeks, about 28 weeks, about 29 weeks, about 30 weeks, about 31 weeks, about 32 weeks, about 33 weeks, about 34 weeks, about 35 weeks, about 36 weeks, about 37 weeks, about 38 weeks, about 39 weeks, about 40 weeks, about 41 weeks, about 42 weeks, about 43 weeks, about 44 weeks, about 45 weeks, about 46 weeks, about 47 weeks, about 48 weeks, about 49 weeks, about 50 weeks, about 51

weeks, about 52 weeks, about 1.5 years, about 2 years, about 2.5 years, about 3.0 years, about 3.5 years, about 4.0 years, about 4.5 years, about 5.0 years, about 5.5 years, about 6.0 years, about 6.5 years, about 7.0 years, about 7.5 years, about 8.0 years, about 8.5 years, about 9.0 years, about 9.5 years or about 10.0 years after the first test sample from the subject is obtained.

5 When used to monitor disease progression, the above assay can be used to monitor the progression of disease in subjects suffering from acute conditions. Acute conditions, also known as critical care conditions, refer to acute, life-threatening diseases or other critical medical conditions involving, for example, the cardiovascular system or excretory system. Typically, critical care conditions refer to those conditions requiring acute medical intervention in a hospital-
 10 based setting (including, but not limited to, the emergency room, intensive care unit, trauma center, or other emergent care setting) or administration by a paramedic or other field-based medical personnel. For critical care conditions, repeat monitoring is generally done within a shorter time frame, namely, minutes, hours or days (e.g., about 1 minute, about 5 minutes, about 10 minutes, about 15 minutes, about 30 minutes, about 45 minutes, about 60 minutes, about 2
 15 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours, about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, about 21 hours, about 22 hours, about 23 hours, about 24 hours, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days or about 7 days), and the initial assay likewise is generally
 20 done within a shorter timeframe, e.g., about minutes, hours or days of the onset of the disease or condition.

The assays also can be used to monitor the progression of disease in subjects suffering from chronic or non-acute conditions. Non-critical care or, non-acute conditions, refers to conditions other than acute, life-threatening disease or other critical medical conditions involving,
 25 for example, the cardiovascular system and/or excretory system. Typically, non-acute conditions include those of longer-term or chronic duration. For non-acute conditions, repeat monitoring generally is done with a longer timeframe, e.g., hours, days, weeks, months or years (e.g., about 1 hour, about 2 hours, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 7 hours, about 8 hours, about 9 hours, about 10 hours, about 11 hours, about 12 hours, about 13 hours,
 30 about 14 hours, about 15 hours, about 16 hours, about 17 hours, about 18 hours, about 19 hours, about 20 hours, about 21 hours, about 22 hours, about 23 hours, about 24 hours, about 2 days, about 3 days, about 4 days, about 5 days, about 6 days, about 7 days, about 2 weeks, about 3 weeks, about 4 weeks, about 5 weeks, about 6 weeks, about 7 weeks, about 8 weeks, about 9 weeks, about 10 weeks, about 11 weeks, about 12 weeks, about 13 weeks, about 14 weeks, about
 35 15 weeks, about 16 weeks, about 17 weeks, about 18 weeks, about 19 weeks, about 20 weeks, about 21 weeks, about 22 weeks, about 23 weeks, about 24 weeks, about 25 weeks, about 26 weeks, about 27 weeks, about 28 weeks, about 29 weeks, about 30 weeks, about 31 weeks, about

32 weeks, about 33 weeks, about 34 weeks, about 35 weeks, about 36 weeks, about 37 weeks, about 38 weeks, about 39 weeks, about 40 weeks, about 41 weeks, about 42 weeks, about 43 weeks, about 44 weeks, about 45 weeks, about 46 weeks, about 47 weeks, about 48 weeks, about 49 weeks, about 50 weeks, about 51 weeks, about 52 weeks, about 1.5 years, about 2 years, about 2.5 years, about 3.0 years, about 3.5 years, about 4.0 years, about 4.5 years, about 5.0 years, about 5.5 years, about 6.0 years, about 6.5 years, about 7.0 years, about 7.5 years, about 8.0 years, about 8.5 years, about 9.0 years, about 9.5 years or about 10.0 years), and the initial assay likewise generally is done within a longer time frame, e.g., about hours, days, months or years of the onset of the disease or condition.

Furthermore, the above assays can be performed using a first test sample obtained from a subject where the first test sample is obtained from one source, such as urine, serum or plasma. Optionally, the above assays can then be repeated using a second test sample obtained from the subject where the second test sample is obtained from another source. For example, if the first test sample was obtained from urine, the second test sample can be obtained from serum or plasma. The results obtained from the assays using the first test sample and the second test sample can be compared. The comparison can be used to assess the status of a disease or condition in the subject.

Moreover, the present disclosure also relates to methods of determining whether a subject predisposed to or suffering from a given disease, disorder or condition will benefit from treatment. In particular, the disclosure relates to analyte companion diagnostic methods and products. Thus, the method of “monitoring the treatment of disease in a subject” as described herein further optimally also can encompass selecting or identifying candidates for therapy.

Thus, in particular embodiments, the disclosure also provides a method of determining whether a subject having, or at risk for, a given disease, disorder or condition is a candidate for therapy. Generally, the subject is one who has experienced some symptom of a given disease, disorder or condition or who has actually been diagnosed as having, or being at risk for, a given disease, disorder or condition, and/or who demonstrates an unfavorable concentration or amount of analyte or a fragment thereof, as described herein.

The method optionally comprises an assay as described herein, where analyte is assessed before and following treatment of a subject with one or more pharmaceutical compositions (e.g., particularly with a pharmaceutical related to a mechanism of action involving analyte), with immunosuppressive therapy, or by immunoabsorption therapy, or where analyte is assessed following such treatment and the concentration or the amount of analyte is compared against a predetermined level. An unfavorable concentration or amount of analyte observed following treatment confirms that the subject will not benefit from receiving further or continued treatment, whereas a favorable concentration or amount of analyte observed following treatment confirms

that the subject will benefit from receiving further or continued treatment. This confirmation assists with management of clinical studies, and provision of improved patient care.

It goes without saying that, while certain embodiments herein are advantageous when employed to assess a given disease, disorder or condition as discussed herein, the assays and kits
5 can be employed to assess analyte in other diseases, disorders and conditions. The method of assay can also involve the assay of other markers and the like.

The method of assay also can be used to identify a compound that ameliorates a given disease, disorder or condition. For example, a cell that expresses analyte can be contacted with a candidate compound. The level of expression of analyte in the cell contacted with the compound
10 can be compared to that in a control cell using the method of assay described herein.

II. Kit

A kit for assaying a test sample for the presence, amount or concentration of an analyte (or a fragment thereof) in a test sample is also provided. The kit comprises at least one
15 component for assaying the test sample for the analyte (or a fragment thereof) and instructions for assaying the test sample for the analyte (or a fragment thereof). The at least one component for assaying the test sample for the analyte (or a fragment thereof) can include a composition comprising an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof), which is optionally immobilized on a solid phase.

20 The kit can comprise at least one component for assaying the test sample for an analyte by immunoassay, e.g., chemiluminescent microparticle immunoassay, and instructions for assaying the test sample for an analyte by immunoassay, e.g., chemiluminescent microparticle immunoassay. For example, the kit can comprise at least one specific binding partner for an analyte, such as an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof that can
25 bind to the analyte, a variant thereof that can bind to the analyte, or a fragment of a variant that can bind to the analyte) or an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof), either of which can be detectably labeled. Alternatively or additionally, the kit can comprise detectably labeled analyte (or a fragment thereof that can bind to an anti-analyte, monoclonal/polyclonal antibody or an anti-analyte DVD-Ig (or a fragment, a variant, or a
30 fragment of a variant thereof)), which can compete with any analyte in a test sample for binding to an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof that can bind to the analyte, a variant thereof that can bind to the analyte, or a fragment of a variant that can bind to the analyte) or an anti-analyte DVD-Ig (or a fragment, a variant, or a fragment of a variant thereof), either of which can be immobilized on a solid support. The kit can comprise a calibrator
35 or control, e.g., isolated or purified analyte. The kit can comprise at least one container (e.g., tube, microtiter plates or strips, which can be already coated with a first specific binding partner, for example) for conducting the assay, and/or a buffer, such as an assay buffer or a wash buffer,

either one of which can be provided as a concentrated solution, a substrate solution for the detectable label (e.g., an enzymatic label), or a stop solution. Preferably, the kit comprises all components, i.e., reagents, standards, buffers, diluents, etc., which are necessary to perform the assay. The instructions can be in paper form or computer-readable form, such as a disk, CD,
5 DVD, or the like.

Any antibodies, such as an anti-analyte antibody or an anti-analyte DVD-Ig, or tracer can incorporate a detectable label as described herein, such as a fluorophore, a radioactive moiety, an enzyme, a biotin/avidin label, a chromophore, a chemiluminescent label, or the like, or the kit can include reagents for carrying out detectable labeling. The antibodies, calibrators and/or controls
10 can be provided in separate containers or pre-dispensed into an appropriate assay format, for example, into microtiter plates.

Optionally, the kit includes quality control components (for example, sensitivity panels, calibrators, and positive controls). Preparation of quality control reagents is well-known in the art and is described on insert sheets for a variety of immunodiagnostic products. Sensitivity panel
15 members optionally are used to establish assay performance characteristics, and further optionally are useful indicators of the integrity of the immunoassay kit reagents, and the standardization of assays.

The kit can also optionally include other reagents required to conduct a diagnostic assay or facilitate quality control evaluations, such as buffers, salts, enzymes, enzyme co-factors,
20 enzyme substrates, detection reagents, and the like. Other components, such as buffers and solutions for the isolation and/or treatment of a test sample (e.g., pretreatment reagents), also can be included in the kit. The kit can additionally include one or more other controls. One or more of the components of the kit can be lyophilized, in which case the kit can further comprise reagents suitable for the reconstitution of the lyophilized components.

The various components of the kit optionally are provided in suitable containers as necessary, e.g., a microtiter plate. The kit can further include containers for holding or storing a sample (e.g., a container or cartridge for a urine sample). Where appropriate, the kit optionally also can contain reaction vessels, mixing vessels, and other components that facilitate the preparation of reagents or the test sample. The kit can also include one or more instruments for
30 assisting with obtaining a test sample, such as a syringe, pipette, forceps, measured spoon, or the like.

If the detectable label is at least one acridinium compound, the kit can comprise at least one acridinium-9-carboxamide, at least one acridinium-9-carboxylate aryl ester, or any combination thereof. If the detectable label is at least one acridinium compound, the kit also can
35 comprise a source of hydrogen peroxide, such as a buffer, a solution, and/or at least one basic solution. If desired, the kit can contain a solid phase, such as a magnetic particle, bead, test tube, microtiter plate, cuvette, membrane, scaffolding molecule, film, filter paper, disc or chip.

III. Adaptation of Kit and Method

The kit (or components thereof), as well as the method of determining the presence, amount or concentration of an analyte in a test sample by an assay, such as an immunoassay as described herein, can be adapted for use in a variety of automated and semi-automated systems (including those wherein the solid phase comprises a microparticle), as described, e.g., in U.S. Patent Nos. 5,089,424 and 5,006,309, and as commercially marketed, e.g., by Abbott Laboratories (Abbott Park, IL) as ARCHITECT®.

Some of the differences between an automated or semi-automated system as compared to a non-automated system (e.g., ELISA) include the substrate to which the first specific binding partner (e.g., an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof, a variant thereof, or a fragment of a variant thereof) or an anti-analyte DVD-Ig (or a fragment thereof, a variant thereof, or a fragment of a variant thereof) is attached; either way, sandwich formation and analyte reactivity can be impacted), and the length and timing of the capture, detection and/or any optional wash steps. Whereas a non-automated format, such as an ELISA, may require a relatively longer incubation time with sample and capture reagent (e.g., about 2 hours), an automated or semi-automated format (e.g., ARCHITECT®, Abbott Laboratories) may have a relatively shorter incubation time (e.g., approximately 18 minutes for ARCHITECT®). Similarly, whereas a non-automated format, such as an ELISA, may incubate a detection antibody, such as the conjugate reagent, for a relatively longer incubation time (e.g., about 2 hours), an automated or semi-automated format (e.g., ARCHITECT®) may have a relatively shorter incubation time (e.g., approximately 4 minutes for the ARCHITECT®).

Other platforms available from Abbott Laboratories include, but are not limited to, AxSYM®, IMx® (see, e.g., U.S. Pat. No. 5,294,404, which is hereby incorporated by reference in its entirety), PRISM®, EIA (bead), and Quantum™ II, as well as other platforms. Additionally, the assays, kits and kit components can be employed in other formats, for example, on electrochemical or other hand-held or point-of-care assay systems. The present disclosure is, for example, applicable to the commercial Abbott Point of Care (i-STAT®, Abbott Laboratories) electrochemical immunoassay system that performs sandwich immunoassays. Immunosensors and their methods of manufacture and operation in single-use test devices are described, for example in, U.S. Patent No. 5,063,081, U.S. Pat. App. Pub. No. 2003/0170881, U.S. Pat. App. Pub. No. 2004/0018577, U.S. Pat. App. Pub. No. 2005/0054078, and U.S. Pat. App. Pub. No. 2006/0160164, which are incorporated in their entireties by reference for their teachings regarding same.

In particular, with regard to the adaptation of an analyte assay to the I-STAT® system, the following configuration is preferred. A microfabricated silicon chip is manufactured with a pair of gold amperometric working electrodes and a silver-silver chloride reference electrode. On

one of the working electrodes, polystyrene beads (0.2 mm diameter) with immobilized anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof, a variant thereof, or a fragment of a variant thereof) or anti-analyte DVD-Ig (or a fragment thereof, a variant thereof, or a fragment of a variant thereof), are adhered to a polymer coating of patterned polyvinyl alcohol
5 over the electrode. This chip is assembled into an I-STAT® cartridge with a fluidics format suitable for immunoassay. On a portion of the wall of the sample-holding chamber of the cartridge there is a layer comprising a specific binding partner for an analyte, such as an anti-analyte, monoclonal/polyclonal antibody (or a fragment thereof, a variant thereof, or a fragment of a variant thereof that can bind the analyte) or an anti-analyte DVD-Ig (or a fragment thereof, a
10 variant thereof, or a fragment of a variant thereof that can bind the analyte), either of which can be detectably labeled. Within the fluid pouch of the cartridge is an aqueous reagent that includes p-aminophenol phosphate.

In operation, a sample suspected of containing an analyte is added to the holding chamber of the test cartridge, and the cartridge is inserted into the I-STAT® reader. After the specific
15 binding partner for an analyte has dissolved into the sample, a pump element within the cartridge forces the sample into a conduit containing the chip. Here it is oscillated to promote formation of the sandwich. In the penultimate step of the assay, fluid is forced out of the pouch and into the conduit to wash the sample off the chip and into a waste chamber. In the final step of the assay, the alkaline phosphatase label reacts with p-aminophenol phosphate to cleave the phosphate group
20 and permit the liberated p-aminophenol to be electrochemically oxidized at the working electrode. Based on the measured current, the reader is able to calculate the amount of analyte in the sample by means of an embedded algorithm and factory-determined calibration curve.

It further goes without saying that the methods and kits as described herein necessarily encompass other reagents and methods for carrying out the immunoassay. For instance,
25 encompassed are various buffers such as are known in the art and/or which can be readily prepared or optimized to be employed, e.g., for washing, as a conjugate diluent, microparticle diluent, and/or as a calibrator diluent. An exemplary conjugate diluent is ARCHITECT® conjugate diluent employed in certain kits (Abbott Laboratories, Abbott Park, IL) and containing 2-(N-morpholino)ethanesulfonic acid (MES), a salt, a protein blocker, an antimicrobial agent, and
30 a detergent. An exemplary calibrator diluent is ARCHITECT® human calibrator diluent employed in certain kits (Abbott Laboratories, Abbott Park, IL), which comprises a buffer containing MES, other salt, a protein blocker, and an antimicrobial agent. Additionally, as described in U.S. Patent Application No. 61/142,048 filed December 31, 2008, improved signal generation may be obtained, e.g., in an I-Stat cartridge format, using a nucleic acid sequence
35 linked to the signal antibody as a signal amplifier.

EXEMPLIFICATION

Example 1: Design, Construction, and Analysis of a DVD-Ig

Example 1.1: Proteolytic Cleavage of DVD-Ig with Cleavable Linkers

Protein samples of enterokinase-cleaved DVD699 (denoted DVD699-C) and DVD701 (denoted DVD701-C) were generated as follows. 180µl of purified DVD-Ig at 1.1 mg/ml was combined with 20µl of 10X enterokinase-cleavage buffer [500mM Tris-HCl (pH 8.0), 10mM CaCl₂, 1% Tween-20]. To this mixture, 5 U/mg of EKMax (Invitrogen, Cat# E180-01) was added, and the mixture was incubated for 2 days at room-temperature. To verify that all the light-chain had been processed to completion at the engineered site, samples (reducing and non-reducing) were run on SDS-PAGE before and after cleavage, and the reaction was deemed to have gone to completion when there was no longer a ~36 kDa band (i.e., intact light-chain) present in the reducing samples and instead there were ~24 kDa and ~12 kDa bands present, corresponding to the expected cleaved LC-fragments. To verify the 12 kDa fragment still remained associated with the DVD-Ig molecule post-cleavage, the mixture was additionally purified over a Protein-A column, and the presence of the 12 kDa (and 24 kDa) protein fragments was confirmed by SDS-PAGE (of reduced samples).

Protein samples of thrombin-cleaved DVD700 (denoted DVD700-C) were generated in a similar manner except that 5 U/mg of thrombin (GE Healthcare, Cat# 27-0846-01) was utilized.

Table 5: Linkers Used to Construct NRP1 – VEGF DVD – Igs

DVD-Ig ID	N-term Variable Domain (VD)	C-term Variable Domain (VD)	Heavy Chain Linker Sequence	SEQ ID NO	Light Chain Linker Sequence	SEQ ID NO
DVD050	NRP1	VEGF	ASTKGP	21	TVAAP	13
DVD695	NRP1	VEGF	ASTKGP	21	TVDDDDKAAP	35
DVD695-C	NRP1	VEGF	ASTKGP	21	TVDDDDK / AAP	35
DVD696	NRP1	VEGF	ASTKGP	21	LVPRGSAAP	36
DVD696-C	NRP1	VEGF	ASTKGP	21	LVPR / GSAAP	36
DVD697	NRP1	VEGF	GGGGGGGGP	27	GGGGGGGP	31
DVD698	NRP1	VEGF	GGGGGGGGGP	28	GGGGGGGP	27

“-C” denotes that the engineered light-chain linker has been treated with either Thrombin (DVD700-C) or Enterokinase (DVD699-C, DVD701-C).

Table 6: Linkers Used Construct SOST - TNF DVD-Igs

DVD-Ig ID	N-term Variable Domain (VD)	C-term Variable Domain (VD)	Heavy Chain Linker Seq	SEQ ID NO	Light Chain Linker Seq	SEQ ID NO
DVD278	SOST	TNF	ASTKGP	21	TVAAP	13
DVD699	SOST	TNF	ASTKGP	21	TVDDDDKAAP	35
DVD699-C	SOST	TNF	ASTKGP	21	TVDDDDK / AAP	35
DVD700	SOST	TNF	ASTKGP	21	LVPRGSAAP	36
DVD700-C	SOST	TNF	ASTKGP	21	LVPR / GSAAP	36
DVD701	SOST	TNF	ASTKGPSVFPL AP	22	TVAADDDDKSV FIVPP	34
DVD701-C	SOST	TNF	ASTKGPSVFPL AP	22	TVAADDDDK / SVFIVPP	4
DVD702	SOST	TNF	GGGGGGGP	27	GGGGGGGP	31
DVD703	SOST	TNF	GGGGGGGGGP	28	GGGGGGGGGP	27
DVD704	SOST	TNF	PAPNLLGGP	29	PAPNLLGGP	29
DVD705	SOST	TNF	PAPNLLGGP	29	PAPELLGGP	32
DVD706	SOST	TNF	PNLLGGP	30	PAPNLLGGP	29
DVD707	SOST	TNF	PAPNLLGGP	29	PTISPAPNLLGGP	33
DVD708	SOST	TNF	PNLLGGP	30	PTISPAPNLLGGP	33

5 “-C” denotes that the engineered light-chain linker has been treated with either Thrombin (DVD700-C) or Enterokinase (DVD699-C, DVD701-C).

Example 1.2: Assays Used to Identify and Characterize Parent Antibodies and DVD-Ig

The following assays are used throughout the Examples to identify and characterize parent antibodies and DVD-Ig unless otherwise stated.

10 **Example 1.2.1: Assays Used To Determine Binding and Affinity of Parent Antibodies and DVD-Ig for Their Target Antigen(s)**

Example 1.2.1.A: Direct Bind ELISA

Enzyme Linked Immunosorbent Assays to screen for antibodies that bind a desired target antigen were performed as follows. High bind ELISA plates (Corning Costar # 3369, Acton, MA) were coated with 100mL/well of 10mg/ml of desired target antigen (R&D Systems, Minneapolis, MN) or desired target antigen extra-cellular domain / FC fusion protein (R&D Systems, Minneapolis, MN) or monoclonal mouse anti-polyHistidine antibody (R&D Systems # MAB050, Minneapolis, MN) in phosphate buffered saline (10X PBS, Abbott Bioresearch Center, Media Prep# MPS-073, Worcester, MA) overnight at 4°C. Plates were washed four times with PBS containing 0.02% Tween 20. Plates were blocked by the addition of 300 mL/well blocking solution (non-fat dry milk powder, various retail suppliers, diluted to 2% in PBS) for 1/2 hour at

15

20

room temperature. Plates were washed four times after blocking with PBS containing 0.02% Tween 20.

Alternatively, one hundred microliters per well of 10 mg/ml of Histidine (His) tagged desired target antigen (R&D Systems, Minneapolis, MN) was added to ELISA plates coated with monoclonal mouse anti-polyHistidine antibody as described above and incubated for 1 hour at room temperature. Wells were washed four times with PBS containing 0.02% Tween 20.

One hundred microliters of antibody or DVD-Ig preparations diluted in blocking solution as described above was added to the desired target antigen plate or desired target antigen / FC fusion plate or the anti-polyHistidine antibody / His tagged desired target antigen plate prepared as described above and incubated for 1 hour at room temperature. Wells are washed four times with PBS containing 0.02% Tween 20.

One hundred microliters of 10ng/mL goat anti-human IgG –FC specific HRP conjugated antibody (Southern Biotech # 2040-05, Birmingham, AL) was added to each well of the desired target antigen plate or anti-polyHistidine antibody / Histidine tagged desired target antigen plate. Alternatively, one hundred microliters of 10 ng/mL goat anti-human IgG –kappa light chain specific HRP conjugated antibody (Southern Biotech # 2060-05 Birmingham, AL) was added to each well of the desired target antigen / FC fusion plate and incubated for 1 hour at room temperature. Plates were washed 4 times with PBS containing 0.02% Tween 20.

One hundred microliters of enhanced TMB solution (Neogen Corp. #308177, K Blue, Lexington, KY) was added to each well and incubated for 10 minutes at room temperature. The reaction was stopped by the addition of 50 mL 1N sulphuric acid. Plates were read spectrophotometrically at a wavelength of 450 nm.

Table 7 contains a list of the antigens used in the NRP1/VEGF Direct Bind Assay.

Table 8 contains the binding data expressed as EC50 in nM for those antibodies and DVD-Ig constructs tested in the NRP1/VEGF Direct Bind ELISA assay.

In the Direct Bind ELISA, binding was sometimes not observed, probably because the antibody binding site on the target antigen was either “masked” or the antigen is “distorted” when coated to the plastic surface. The inability of a DVD-Ig to bind its target may also be due to steric limitation imposed on DVD-Ig by the Direct Bind ELISA format. The parent antibodies and DVD-Igs that did not bind in the Direct Bind ELISA format bound to target antigen in other ELISA formats, such as FACS, Biacore or bioassay. Non-binding of a DVD-Ig was also restored by adjusting the linker length between the two variable domains of the DVD-Ig, as shown previously.

Table 7: Antigens Used in Direct Bind ELISA

Assay	Antigen	Vendor Designation	Vendor	Catalog #
NRP1	NRP1	Neuropilin-1 Npn-1-His tag	R&D	3870-N1-025
VEGF	VEGF	VEGF 1-165 aa	R&D	293-VE-010

Table 8: NRP1 And VEGF Direct Bind ELISA Of 4 DVD Constructs With Various Sequences, Orientations And Linker Combinations

<u>DVD-Ig</u> <u>ID</u>	<u>N-Term.</u> <u>VD</u> <u>Seq.</u> <u>ID</u>	<u>N-Term.</u> <u>VD</u> <u>EC50</u> <u>(nM)</u>	<u>N-Term.</u> <u>Ref.</u> <u>Ab. ID</u>	<u>N-Term.</u> <u>Ref.</u> <u>Ab.E</u> <u>C50</u> <u>(nM)</u>	<u>HC</u> <u>linker</u>	<u>LC</u> <u>linker</u>	<u>C-Term.</u> <u>VD</u> <u>Seq.</u> <u>ID</u>	<u>C-Term.</u> <u>VD</u> <u>EC50</u> <u>(nM)</u>	<u>C-Term.</u> <u>Ref.</u> <u>Ab. ID</u>	<u>C-Term.</u> <u>Ref.</u> <u>Ab.E</u> <u>C50</u> <u>(nM)</u>
DVD695	NRP1 (seq 1)	0.32	AB016	0.28	HG-short	LK-(EK)-short	VEGF (seq1)	0.37	AB014	0.45
DVD696	NRP1 (seq 1)	0.28	AB016	0.28	HG-short	LK-(THR)-short	VEGF (seq1)	0.33	AB014	0.45
DVD697	NRP1 (seq 1)	0.27	AB016	0.28	H-7GP	L-6GP	VEGF (seq1)	0.42	AB014	0.45
DVD698	NRP1 (seq 1)	0.30	AB016	0.28	H-8GP	L-7GP	VEGF (seq1)	0.35	AB014	0.45

Binding of all DVD-Ig constructs was maintained and comparable to that of parent antibodies. All N-terminal and C-terminal variable domains of DVD-Ig constructs DVD695, DVD696, DVD697, and DVD698 bound their target antigens with a similar high affinity as the parent antibody.

Table 9 contains a list of the antigens used in the SOST/TNF Direct Bind Assay.

Table 10 contains the binding data expressed as EC50 in nM for those antibodies and DVD-Ig constructs tested in the SOST/TNF Direct Bind ELISA assay.

Table 9: Antigens Used in Direct Bind ELISA

Assay	Antigen	Vendor Designation	Vendor	Catalog #
SOST	SOST	SOST/His	R&D	210-TA
TNF α	TNF α	TNF α /TNFSF1A	R&D	293-VE-010

Table 10: SOST & TNF α Direct Bind ELISA Of 9 DVD Constructs With Various Sequences, Orientations And Linker Combinations

<u>DVD-Ig ID</u>	<u>N-Term. VD Seq. ID</u>	<u>N-Term. VD EC50 (nM)</u>	<u>N-Term. Ref. Ab. ID</u>	<u>N-Term. Ref. Ab.E C50 (nM)</u>	<u>HC linker</u>	<u>LC linker</u>	<u>C-Term. VD Seq. ID</u>	<u>C-Term. VD EC50 (nM)</u>	<u>C-Term. Ref. Ab. ID</u>	<u>C-Term. Ref. Ab.E C50 (nM)</u>
DVD699	SOST (seq2)	0.33	AB050	0.30	HG-short	LK-(EK)-short	TNF α	17.76	AB017	0.14
DVD700	SOST (seq2)	0.26	AB050	0.30	HG-short	LK-(THR)-short	TNF α	7.76	AB017	0.14
DVD702	SOST (seq2)	2.44	AB050	0.30	H-7GP	L-6GP	TNF α	12.43	AB017	0.14
DVD703	SOST (seq2)	0.32	AB050	0.30	H-8GP	L-7GP	TNF α	6.91	AB017	0.14
DVD704	SOST (seq2)	0.44	AB050	0.30	HH-long	LH-N-short	TNF α	2.92	AB017	0.14
DVD705	SOST (seq2)	0.18	AB050	0.30	HH-long	LH-E-short	TNF α	3.33	AB017	0.14
DVD706	SOST (seq2)	0.42	AB050	0.37	HH-short	LH-N-short	TNF α	6.50	AB017	0.14
DVD707	SOST (seq2)	0.4	AB050	0.37	HH-long	LH-long	TNF α	3.09	AB017	0.14
DVD708	SOST (seq2)	0.57	AB050	0.37	HH-short	LH-long	TNF α	3.60	AB017	0.14

5 Binding of all DVD-Ig constructs was maintained. All N-terminal and C-terminal variable domains of DVD-Ig constructs DVD699, DVD700, DVD702, DVD703, DVD704, DVD705, DVD706, DVD707, DVD708 bound their target antigens with a similar high affinity as the parent antibody.

Example 1.2.1.B: Affinity Determination using BIACORE Technology

10 The BIACORE assay (Biacore, Inc, Piscataway, NJ) determines the affinity of antibodies or DVD-Ig with kinetic measurements of on-rate and off-rate constants. Binding of antibodies or DVD-Ig to a target antigen (for example, a purified recombinant target antigen) is determined by surface plasmon resonance-based measurements with a Biacore® 1000 or 3000 instrument (Biacore® AB, Uppsala, Sweden) using running HBS-EP (10 mM HEPES [pH 7.4], 150 mM

15 NaCl, 3 mM EDTA, and 0.005% surfactant P20) at 25° C. All chemicals are obtained from Biacore® AB (Uppsala, Sweden) or otherwise from a different source as described in the text. For example, approximately 5000 RU of goat anti-mouse IgG, (Fc γ), fragment specific polyclonal antibody (Pierce Biotechnology Inc, Rockford, IL) diluted in 10 mM sodium acetate (pH 4.5) is directly immobilized across a CM5 research grade biosensor chip using a standard amine

20 coupling kit according to manufacturer's instructions and procedures at 25 μ g/ml. Unreacted moieties on the biosensor surface are blocked with ethanolamine. Modified carboxymethyl

dextran surface in flowcell 2 and 4 is used as a reaction surface. Unmodified carboxymethyl dextran without goat anti-mouse IgG in flow cell 1 and 3 is used as the reference surface. For kinetic analysis, rate equations derived from the 1:1 Langmuir binding model are fitted simultaneously to association and dissociation phases of all eight injections (using global fit analysis) with the use of Biaevaluation 4.0.1 software. Purified antibodies or DVD-Ig are diluted in HEPES-buffered saline for capture across goat anti-mouse IgG specific reaction surfaces. Antibodies or DVD-Ig to be captured as a ligand (25 µg/ml) are injected over reaction matrices at a flow rate of 5 µl/min. The association and dissociation rate constants, k_{on} ($M^{-1}s^{-1}$) and k_{off} (s^{-1}) are determined under a continuous flow rate of 25 µl/min. Rate constants are derived by making kinetic binding measurements at different antigen concentrations ranging from 10 – 200 nM. The equilibrium dissociation constant (M) of the reaction between antibodies or DVD-Igs and the target antigen is then calculated from the kinetic rate constants by the following formula: $K_D = k_{off}/k_{on}$. Binding is recorded as a function of time and kinetic rate constants are calculated. In this assay, on-rates as fast as $10^6 M^{-1}s^{-1}$ and off-rates as slow as $10^{-6} s^{-1}$ can be measured.

Table 11: BIACORE Analysis of SOST and TNF DVD-Igs

Parent Antibody or DVD-Ig ID	N-Terminal Variable Domain (VD)	C-Terminal Variable Domain (VD)	k_{on} ($M^{-1}s^{-1}$)	k_{off} (s^{-1})	K_D (M)
AB017	TNF		1.28E+06	1.58E-04	1.23E-10
DVD278	SOST	TNF	1.74E+04	2.26E-04	1.30E-08
DVD699	SOST	TNF	4.70E+04	2.39E-04	5.09E-09
DVD699-C	SOST	TNF	3.71E+05	1.95E-04	5.26E-10
DVD700	SOST	TNF	5.62E+04	2.00E-04	3.56E-09
DVD700-C	SOST	TNF	3.64E+05	1.90E-04	5.22E-10
DVD701	SOST	TNF	2.41E+05	1.97E-04	8.17E-10
DVD701-C	SOST	TNF	4.45E+05	1.79E-04	4.02E-10
DVD702	SOST	TNF	3.14E+04	2.13E-04	6.78E-09
DVD703	SOST	TNF	5.11E+04	1.93E-04	3.78E-09
DVD704	SOST	TNF	2.11E+05	1.90E-04	9.00E-10
DVD705	SOST	TNF	1.73E+05	2.01E-04	1.16E-09
DVD706	SOST	TNF	8.40E+04	2.05E-04	2.44E-09
DVD707	SOST	TNF	2.57E+05	1.85E-04	7.20E-10
DVD708	SOST	TNF	1.56E+05	1.84E-04	1.18E-09

Note: “-C” denotes that the engineered light-chain linker has been treated with either Thrombin (DVD700-C) or Enterokinase (DVD699-C, DVD701-C).

Cleavage of light chain linker with either thrombin (DVD 700-C) or enterokinase (DVD699-C, DVD701C) improved inner domain binding to TNF compared to the inner domain TNF binding of non-cleaved linker DVD278 and DVD699. Linker derived from the hinge region in DVD – Ig 704 and DVD – Ig 707 improved inner domain binding.

Example 1.2.2: Assays Used To Determine the Functional Activity Of Parent Antibodies And DVD-Ig

Example 1.2.2.A: Cytokine Bioassay

The ability of an anti-cytokine or an anti-growth factor parent antibody or DVD-Ig containing anti-cytokine or an anti-growth factor sequences to inhibit or neutralize a target cytokine or an anti-growth factor bioactivity is analyzed by determining inhibitory potential of the antibody or DVD-Ig. For example, the ability of an anti-IL-4 antibody to inhibit IL-4 mediated IgE production may be used. For example, human naive B cells are isolated from peripheral blood, respectively, buffy coats by Ficoll-paque density centrifugation, followed by magnetic separation with MACS beads (Miltenyi Biotec, Bergisch Gladbach, Germany) specific for human sIgD FITC labeled goat F(ab)2 antibodies followed by anti-FITC MACS beads. Magnetically sorted naive B cells are adjusted to 3×10^5 cells per ml in XV15 and plated out in 100 μ l per well of 96-well plates in a 6 x 6 array in the center of the plate, surrounded by PBS filled wells during the 10 days of culture at 37° C in the presence of 5% CO₂. One plate each is prepared per antibody to be tested, consisting of 3 wells each of un-induced and induced controls and quintuplicate repeats of antibody titrations starting at 7 μ g/ml and running in 3-fold dilution down to 29 ng/ml final concentrations added in 50 μ l four times concentrated pre-dilution. To induce IgE production, rhIL-4 at 20 ng/ml plus anti-CD40 monoclonal antibody (Novartis, Basel, Switzerland) at 0.5 μ g/ml final concentrations in 50 μ l each are added to each well, and IgE concentrations are determined at the end of the culture period by a standard sandwich ELISA method.

Example 1.2.2.B: Cytokine Release Assay

The ability of a parent antibody or DVD-Ig to cause cytokine release is analyzed. Peripheral blood is withdrawn from three healthy donors by venipuncture into heparized vacutainer tubes. Whole blood is diluted 1:5 with RPMI-1640 medium and placed in 24-well tissue culture plates at 0.5 mL per well. The anti-cytokine antibodies (e.g., anti-IL-4) are diluted into RPMI-1640 and placed in the plates at 0.5 mL/well to give final concentrations of 200, 100, 50, 10, and 1 μ g/mL. The final dilution of whole blood in the culture plates is 1:10. LPS and PHA are added to separate wells at 2 μ g/mL and 5 μ g/mL final concentration as a positive control for cytokine release. Polyclonal human IgG is used as negative control antibody. The experiment is performed in duplicate. Plates are incubated at 37°C at 5% CO₂. Twenty-four hours later the contents of the wells are transferred into test tubes and spun for 5 minutes at 1200 rpm. Cell-free supernatants are collected and frozen for cytokine assays. Cells left over on the plates and in the tubes are lysed with 0.5 mL of lysis solution, and placed at -20°C and thawed. 0.5 mL of medium is added (to bring the volume to the same level as the cell-free supernatant samples) and the cell preparations are collected and frozen for cytokine assays. Cell-free supernatants and cell lysates are assayed for cytokine levels by ELISA, for example, for levels of IL-8, IL-6, IL-1 β , IL-1RA, or TNF- α .

Example 1.2.2.C: Cytokine Cross-Reactivity Study

The ability of an anti-cytokine parent antibody or DVD-Ig directed to a cytokine(s) of interest to cross react with other cytokines is analyzed. Parent antibodies or DVD-Ig are immobilized on a Biacore biosensor matrix. An anti-human Fc mAb is covalently linked via free amine groups to the dextran matrix by first activating carboxyl groups on the matrix with 100mM N-hydroxysuccinimide (NHS) and 400mM N-Ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC). Approximately 50μL of each antibody or DVD-Ig preparation at a concentration of 25μg/mL, diluted in sodium acetate, pH4.5, is injected across the activated biosensor and free amines on the protein are bound directly to the activated carboxyl groups. Typically, 5000 Resonance Units (RU's) are immobilized. Unreacted matrix EDC-esters are deactivated by an injection of 1 M ethanolamine. A second flow cell is prepared as a reference standard by immobilizing human IgG1/K using the standard amine coupling kit. SPR measurements are performed using the CM biosensor chip. All antigens to be analyzed on the biosensor surface are diluted in HBS-EP running buffer containing 0.01% P20.

To examine the cytokine binding specificity, excess cytokine of interest (100nM, e.g., soluble recombinant human) is injected across the anti-cytokine parent antibody or DVD-Ig immobilized biosensor surface (5 minute contact time). Before injection of the cytokine of interest and immediately afterward, HBS-EP buffer alone flows through each flow cell. The net difference in the signals between the baseline and the point corresponding to approximately 30 seconds after completion of cytokine injection are taken to represent the final binding value. Again, the response is measured in Resonance Units. Biosensor matrices are regenerated using 10mM HCl before injection of the next sample where a binding event is observed, otherwise running buffer was injected over the matrices. Human cytokines (e.g., IL-1α, IL-1β, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, IL-22, IL-23, IL-27, TNF-α, TNF-β, and IFN-γ, for example) are also simultaneously injected over the immobilized mouse IgG1/K reference surface to record any nonspecific binding background. By preparing a reference and reaction surface, Biacore can automatically subtract the reference surface data from the reaction surface data in order to eliminate the majority of the refractive index change and injection noise. Thus, it is possible to ascertain the true binding response attributed to an anti-cytokine antibody or DVD-Ig binding reaction.

When a cytokine of interest is injected across immobilized anti-cytokine antibody, significant binding is observed. 10mM HCl regeneration completely removes all non-covalently associated proteins. Examination of the sensorgram shows that immobilized anti-cytokine antibody or DVD-Ig binding to soluble cytokine is strong and robust. After confirming the expected result with the cytokine of interest, the panel of remaining recombinant human cytokines

is tested, for each antibody or DVD-Ig separately. The amount of anti-cytokine antibody or DVD-Ig bound or unbound cytokine for each injection cycle is recorded. The results from three independent experiments are used to determine the specificity profile of each antibody or DVD-Ig. Antibodies or DVD-Ig with the expected binding to the cytokine of interest and no binding to any other cytokine are selected.

Example 1.2.2.D: Tissue Cross Reactivity

Tissue cross reactivity studies are done in three stages, with the first stage including cryosections of 32 tissues, second stage including up to 38 tissues, and the 3rd stage including additional tissues from 3 unrelated adults as described below. Studies are done typically at two dose levels.

Stage 1: Cryosections (about 5 μ m) of human tissues (32 tissues (typically: Adrenal Gland, Gastrointestinal Tract, Prostate, Bladder, Heart, Skeletal Muscle, Blood Cells, Kidney, Skin, Bone Marrow, Liver, Spinal Cord, Breast, Lung, Spleen, Cerebellum, Lymph Node, Testes, Cerebral Cortex, Ovary, Thymus, Colon, Pancreas, Thyroid, Endothelium, Parathyroid, Ureter, Eye, Pituitary, Uterus, Fallopian Tube and Placenta) from one human donor obtained at autopsy or biopsy) are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system.

Stage 2: Cryosections (about 5 μ m) of human tissues 38 tissues (including adrenal, blood, blood vessel, bone marrow, cerebellum, cerebrum, cervix, esophagus, eye, heart, kidney, large intestine, liver, lung, lymph node, breast mammary gland, ovary, oviduct, pancreas, parathyroid, peripheral nerve, pituitary, placenta, prostate, salivary gland, skin, small intestine, spinal cord, spleen, stomach, striated muscle, testis, thymus, thyroid, tonsil, ureter, urinary bladder, and uterus) from 3 unrelated adults obtained at autopsy or biopsy) are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system.

Stage 3: Cryosections (about 5 μ m) of cynomolgus monkey tissues (38 tissues (including adrenal, blood, blood vessel, bone marrow, cerebellum, cerebrum, cervix, esophagus, eye, heart, kidney, large intestine, liver, lung, lymph node, breast mammary gland, ovary, oviduct, pancreas, parathyroid, peripheral nerve, pituitary, placenta, prostate, salivary gland, skin, small intestine, spinal cord, spleen, stomach, striated muscle, testis, thymus, thyroid, tonsil, ureter, urinary bladder, and uterus) from 3 unrelated adult monkeys obtained at autopsy or biopsy) are fixed and dried on object glass. The peroxidase staining of tissue sections is performed, using the avidin-biotin system.

The antibody or DVD-Ig is incubated with the secondary biotinylated anti-human IgG and developed into immune complex. The immune complex at the final concentrations of 2 and 10 μ g/mL of antibody or DVD-Ig is added onto tissue sections on object glass and then the tissue

sections are reacted for 30 minutes with a avidin-biotin-peroxidase kit. Subsequently, DAB (3,3'-diaminobenzidine), a substrate for the peroxidase reaction, is applied for 4 minutes for tissue staining. Antigen-Sepharose beads are used as positive control tissue sections. Target antigen and human serum blocking studies serve as additional controls. The immune complex at the final concentrations of 2 and 10 µg/mL of antibody or DVD-Ig is pre-incubated with target antigen (final concentration of 100 µg/ml) or human serum (final concentration 10%) for 30 minutes, and then added onto the tissue sections on object glass and then the tissue sections are reacted for 30 minutes with a avidin-biotin-peroxidase kit. Subsequently, DAB (3,3'-diaminobenzidine), a substrate for the peroxidase reaction, is applied for 4 minutes for tissue staining.

Any specific staining is judged to be either an expected (e.g., consistent with antigen expression) or unexpected reactivity based upon known expression of the target antigen in question. Any staining judged specific is scored for intensity and frequency. The tissue staining between stage 2 (human tissue) and stage 3 (cynomolgus monkey tissue) is either judged to be similar or different.

Example 1.2.2.E: Tumoricidal Effect Of A Parent or DVD-Ig Antibody *In Vitro*

Parent antibodies or DVD-Ig that bind to target antigens on tumor cells may be analyzed for tumoricidal activity. Briefly, parent antibodies or DVD-Ig are diluted in D-PBS-BSA (Dulbecco's phosphate buffered saline with 0.1%BSA) and added to human tumor cells at final concentrations of 0.01 µg/mL to 200 µg/mL. The plates are incubated at 37 °C in a humidified, 5% CO₂ atmosphere for 3 days. The number of live cells in each well is quantified using MTS reagents according to the manufacturer's instructions (Promega, Madison, WI) to determine the percent of tumor growth inhibition. Wells without antibody treatment are used as controls of 0% inhibition whereas wells without cells are considered to show 100% inhibition.

For assessment of apoptosis, caspase-3 activation is determined by the following protocol: antibody-treated cells in 96 well plates are lysed in 120 µl of 1x lysis buffer (1.67mM Hepes, pH 7.4, 7mM KCl, 0.83mM MgCl₂, 0.11mM EDTA, 0.11mM EGTA, 0.57% CHAPS, 1mM DTT, 1x protease inhibitor cocktail tablet; EDTA-free; Roche Pharmaceuticals, Nutley, NJ) at room temperature with shaking for 20 minutes. After cell lysis, 80 µl of a caspase-3 reaction buffer (48mM Hepes, pH 7.5, 252mM sucrose, 0.1% CHAPS, 4mM DTT, and 20 µM Ac-DEVD-AMC substrate; Biomol Research Labs, Inc., Plymouth Meeting, PA) is added and the plates are incubated for 2 hours at 37°C. The plates are read on a 1420 VICTOR Multilabel Counter (Perkin Elmer Life Sciences, Downers Grove, IL) using the following settings: excitation= 360/40, emission= 460/40. An increase of fluorescence units from antibody-treated cells relative to the isotype antibody control-treated cells is indicative of apoptosis.

Example 1.2.2.F: Inhibition Of Receptor Activation By Antibodies or DVD-Ig *In Vitro*

Parent antibodies or DVD-Ig that bind to cell receptors or their ligands may be tested for inhibition of receptor activation. Parent antibodies or DVD-Ig diluted in D-PBS-BSA (Dulbecco's phosphate buffered saline with 0.1%BSA) are added to human carcinoma cells at

5 final concentrations of 0.01 $\mu\text{g/mL}$ to 100 $\mu\text{g/mL}$. The plates are incubated at 37 °C in a humidified, 5% CO₂ atmosphere for 1h. Growth factors (e.g., IGF1 or IGF2) at concentration of 1-100 ng/mL are added to the cells for 5-15 minutes to stimulate receptor (e.g., IGF1R) autophosphorylation. Wells without antibody treatment are used as controls of 0% inhibition whereas wells without growth factor stimulation are considered to show 100% inhibition. Cell

10 lysates are made by incubation with cell extraction buffer (10 mM Tris, pH 7.4, 100 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM NaF, 1 mM sodium orthovanadate, 1% Triton X-100, 10% Glycerol, 0.1% SDS, and protease inhibitor cocktail). Phospho-IGF1R in these cell lysates is determined using specific ELISA kits purchased from R&D System (Minneapolis, MN).

15 **Example 1.2.2.G: Efficacy Of An Anti-Tumor Cell Antigen Antibody or DVD-Ig By Itself Or In Combination With Chemotherapy On The Growth Of Human Carcinoma Xenografts (Subcutaneous Flank, Orthotopic, Or Spontaneous Metastases)**

Human cancer cells are grown *in vitro* to 99% viability, 85% confluence in tissue culture flasks. SCID female or male mice (Charles Rivers Labs, Wilmington, MA) at 19-25 grams are ear tagged and shaved. Mice are then inoculated subcutaneously into the right flank with 1 x 10⁶

20 human tumor cells (1:1 matrigel) on study day 0. Administration (IP, QD, 3x/ week) of human IgG control antibody or DVD-Ig, and/or chemotherapy is initiated after mice are size matched into groups of mice with mean tumor volumes of approximately 200-320 mm³. The tumors are measured by a pair of calipers twice a week starting on approximately day 10 post tumor cell injection. Reduction in tumor volume is seen in animals treated with antibody or DVD-Ig alone

25 or in combination with chemotherapy relative to tumors in animals that received only vehicle or an isotype control mAb.

Example 1.2.2.H: Binding of Monoclonal Antibodies to the Surface of Human Tumor Cell Lines as Assessed by Flow Cytometry

Stable cell lines overexpressing a cell-surface antigen of interest or human tumor cell

30 lines were harvested from tissue culture flasks and resuspended in phosphate buffered saline (PBS) containing 5% fetal calf serum (PBS/FCS). Prior to staining, human tumor cells were incubated on ice with 100 μl human IgG at 5 $\mu\text{g/mL}$ in PBS/FCS. 1-5 x10⁵ cells were incubated with antibody or DVD-Ig (1-2 $\mu\text{g/mL}$) in PBS/FCS for 30-60 minutes on ice. Cells were washed twice and 100 μl of F(ab')₂ goat anti human IgG, Fc γ - phycoerythrin (1:200 dilution in

35 PBS/BSA) (Jackson ImmunoResearch, West Grove, PA, Cat.#109-116-170) was added. After 30

minutes incubation on ice, cells were washed twice and resuspended in PBS/FCS. Fluorescence was measured using a Becton Dickinson FACSCalibur (Becton Dickinson, San Jose, CA).

Example 1.3: Generation Of Parent Monoclonal Antibodies to a Human Antigen of Interest

Parent mouse mAbs able to bind to and neutralize a human antigen of interest and a
5 variant thereof are obtained as follows:

Example 1.3.A: Immunization Of Mice With a Human Antigen of Interest

Twenty micrograms of recombinant purified human antigen (e.g., IGF1,2) mixed with complete Freund's adjuvant or Immuneasy adjuvant (Qiagen, Valencia, CA) is injected subcutaneously into five 6-8 week-old Balb/C, five C57B/6 mice, and five AJ mice on Day 1. On
10 days 24, 38, and 49, twenty micrograms of recombinant purified human antigen variant mixed with incomplete Freund's adjuvant or Immuneasy adjuvant is injected subcutaneously into the same mice. On day 84 or day 112 or day 144, mice are injected intravenously with 1 µg recombinant purified human antigen of interest.

Example 1.3.B: Generation of Hybridomas

15 Splenocytes obtained from the immunized mice described in Example 1.3.A are fused with SP2/O-Ag-14 cells at a ratio of 5:1 according to the established method described in Kohler, G. and Milstein (1975) Nature, 256:495 to generate hybridomas. Fusion products are plated in selection media containing azaserine and hypoxanthine in 96-well plates at a density of 2.5×10^6 spleen cells per well. Seven to ten days post fusion, macroscopic hybridoma colonies are
20 observed. Supernatant from each well containing hybridoma colonies is tested by ELISA for the presence of antibody to the antigen of interest (as described in Example 1.3.A). Supernatants displaying antigen-specific activity are then tested for activity (as described in the assays of Example 1.2.2), for example, the ability to neutralize the antigen of interest in a bioassay such as that described in Example 1.2.2.A).

25 **Example 1.3.C: Identification And Characterization Of Parent Monoclonal Antibodies to a Human Target Antigen of Interest**

Example 1.3.C.1: Analyzing Parent Monoclonal Antibody Neutralizing Activity

Hybridoma supernatants are assayed for the presence of parent antibodies that bind an antigen of interest, generated according to Examples 1.3.A and 1.3.B, and are also capable of
30 binding a variant of the antigen of interest ("antigen variant"). Supernatants with antibodies positive in both assays are then tested for their antigen neutralization potency, for example, in the cytokine bioassay of Example 1.2.2.A. The hybridomas producing antibodies with IC_{50} values in the bioassay less than 1000pM, in an embodiment, less than 100pM are scaled up and cloned by limiting dilution. Hybridoma cells are expanded into media containing 10% low IgG fetal bovine

serum (Hyclone #SH30151, Logan, UT.). On average, 250 mL of each hybridoma supernatant (derived from a clonal population) is harvested, concentrated and purified by protein A affinity chromatography, as described in Harlow, E. and Lane, D. 1988 "Antibodies: A Laboratory Manual". The ability of purified mAbs to inhibit the activity of its target antigen is determined, for example, using the cytokine bioassay as described in Example 1.2.2.A.

Example 1.3.C.2: Analyzing Parent Monoclonal Antibody Cross-Reactivity To Cynomolgus Target Antigen Of Interest

To determine whether the selected mAbs described herein recognize cynomolgus antigen of interest, BIACORE analysis is conducted as described herein (Example 1.2.1.B) using recombinant cynomolgus target antigen. In addition, neutralization potencies of mAbs against recombinant cynomolgus antigen of interest may also be measured in the cytokine bioassay (Example 1.2.2.A). MAbs with good cyno cross-reactivity (in an embodiment, within 5-fold of reactivity for human antigen) are selected for future characterization.

Example 1.3.D: Determination Of The Amino Acid Sequence Of The Variable Region For Each Murine Anti-Human Monoclonal Antibody

Isolation of the cDNAs, expression and characterization of the recombinant anti-human mouse mAbs is conducted as follows. For each amino acid sequence determination, approximately 1×10^6 hybridoma cells are isolated by centrifugation and processed to isolate total RNA with Trizol (Gibco BRL/Invitrogen, Carlsbad, CA.) following manufacturer's instructions. Total RNA is subjected to first strand DNA synthesis using the SuperScript First-Strand Synthesis System (Invitrogen, Carlsbad, CA) per the manufacturer's instructions. Oligo(dT) is used to prime first-strand synthesis to select for poly(A)+ RNA. The first-strand cDNA product is then amplified by PCR with primers designed for amplification of murine immunoglobulin variable regions (Ig-Primer Sets, Novagen, Madison, WI). PCR products are resolved on an agarose gel, excised, purified, and then subcloned with the TOPO Cloning kit into pCR2.1-TOPO vector (Invitrogen, Carlsbad, CA) and transformed into TOP10 chemically competent E. coli (Invitrogen, Carlsbad, CA). Colony PCR is performed on the transformants to identify clones containing insert. Plasmid DNA is isolated from clones containing insert using a QIAprep Miniprep kit (Qiagen, Valencia, CA). Inserts in the plasmids are sequenced on both strands to determine the variable heavy or variable light chain DNA sequences using M13 forward and M13 reverse primers (Fermentas Life Sciences, Hanover MD). Variable heavy and variable light chain sequences of the mAbs are identified. In an embodiment, the selection criteria for a panel of lead mAbs for next step development (humanization) includes the following:

- The antibody does not contain any N-linked glycosylation sites (NXS), except from the standard one in CH2

- The antibody does not contain any extra cysteines in addition to the normal cysteines in every antibody
- The antibody sequence is aligned with the closest human germline sequences for VH and VL and any unusual amino acids should be checked for occurrence in other natural human antibodies
- N-terminal Glutamine (Q) is changed to Glutamic acid (E) if it does not affect the activity of the antibody. This will reduce heterogeneity due to cyclization of Q
- Efficient signal sequence cleavage is confirmed by Mass Spectrophotometry. This can be done with COS cell or 293 cell material
- The protein sequence is checked for the risk of deamidation of Asn that could result in loss of activity
- The antibody has a low level of aggregation
- The antibody has solubility >5-10 mg/ml (in research phase); >25 mg/ml
- The antibody has a normal size (5-6 nm) by Dynamic Light Scattering (DLS)
- The antibody has a low charge heterogeneity
- The antibody lacks cytokine release (see Example 1.2.2.B)
- The antibody has specificity for the intended cytokine (see Example 1.2.2.C)
- The antibody lacks unexpected tissue cross reactivity (see Example 1.2.2.D)
- The antibody has similarity between human and cynomolgus tissue cross reactivity (see Example 1.2.2.D)

Example 1.3.2: Recombinant Humanized Parent Antibodies

Example 1.3.2.1: Construction And Expression Of Recombinant Chimeric Anti Human Parent Antibodies

The DNA encoding the heavy chain constant region of murine anti-human parent mAbs is replaced by a cDNA fragment encoding the human IgG1 constant region containing 2 hinge-region amino acid mutations by homologous recombination in bacteria. These mutations are a leucine to alanine change at position 234 (EU numbering) and a leucine to alanine change at position 235 (Lund et al., 1991, J. Immunol., 147:2657). The light chain constant region of each of these antibodies is replaced by a human kappa constant region. Full-length chimeric antibodies are transiently expressed in COS cells by co-transfection of chimeric heavy and light chain cDNAs ligated into the pBOS expression plasmid (Mizushima and Nagata, Nucleic Acids Research 1990, Vol 18, pg 5322). Cell supernatants containing recombinant chimeric antibody are purified by Protein A Sepharose chromatography and bound antibody is eluted by addition of acid buffer. Antibodies are neutralized and dialyzed into PBS.

The heavy chain cDNA encoding a chimeric mAb is co-transfected with its chimeric light chain cDNA (both ligated in the pBOS vector) into COS cells. Cell supernatant containing recombinant chimeric antibody is purified by Protein A Sepharose chromatography and bound antibody is eluted by addition of acid buffer. Antibodies are neutralized and dialyzed into PBS.

5 The purified chimeric anti-human parent mAbs are then tested for their ability to bind (by Biacore) and for functional activity, e.g., to inhibit the cytokine induced production of IgE as described in Examples 1.2.1.B and 1.2.2.B. Chimeric mAbs that maintain the activity of the parent hybridoma mAbs are selected for future development.

10 **Example 1.3.2.2: Construction And Expression Of Humanized Anti Human Parent Antibodies**

Example 1.3.2.2.A: Selection Of Human Antibody Frameworks

Each murine variable heavy and variable light chain gene sequence is separately aligned against 44 human immunoglobulin germline variable heavy chain or 46 germline variable light chain sequences (derived from NCBI Ig Blast website at
15 <http://www.ncbi.nlm.nih.gov/igblast/retrieveig.html>.) using Vector NTI software.

Humanization is based on amino acid sequence homology, CDR cluster analysis, frequency of use among expressed human antibodies, and available information on the crystal structures of human antibodies. Taking into account possible effects on antibody binding, VH-VL pairing, and other factors, murine residues are mutated to human residues where murine and
20 human framework residues are different, with a few exceptions. Additional humanization strategies are designed based on an analysis of human germline antibody sequences, or a subgroup thereof, that possessed a high degree of homology, i.e., sequence similarity, to the actual amino acid sequence of the murine antibody variable regions.

Homology modeling is used to identify residues unique to the murine antibody sequences
25 that are predicted to be critical to the structure of the antibody combining site, the CDRs. Homology modeling is a computational method whereby approximate three dimensional coordinates are generated for a protein. The source of initial coordinates and guidance for their further refinement is a second protein, the reference protein, for which the three dimensional coordinates are known and the sequence of which is related to the sequence of the first protein.
30 The relationship among the sequences of the two proteins is used to generate a correspondence between the reference protein and the protein for which coordinates are desired, the target protein. The primary sequences of the reference and target proteins are aligned with coordinates of identical portions of the two proteins transferred directly from the reference protein to the target protein. Coordinates for mismatched portions of the two proteins, e.g., from residue mutations,
35 insertions, or deletions, are constructed from generic structural templates and energy refined to

insure consistency with the already transferred model coordinates. This computational protein structure may be further refined or employed directly in modeling studies. The quality of the model structure is determined by the accuracy of the contention that the reference and target proteins are related and the precision with which the sequence alignment is constructed.

5 For the murine mAbs, a combination of BLAST searching and visual inspection is used to identify suitable reference structures. Sequence identity of 25% between the reference and target amino acid sequences is considered the minimum necessary to attempt a homology modeling exercise. Sequence alignments are constructed manually and model coordinates are generated with the program Jackal (see Petrey, D. et al. (2003) Proteins 53 (Supl. 6): 430–435).

10 The primary sequences of the murine and human framework regions of the selected antibodies share significant identity. Residue positions that differ are candidates for inclusion of the murine residue in the humanized sequence in order to retain the observed binding potency of the murine antibody. A list of framework residues that differ between the human and murine sequences is constructed manually. Table 12 shows the framework sequences chosen for this
15 study.

Table 12: Sequence Of Human IgG Heavy Chain Constant Domain And Light Chain Constant Domain

Protein	SEQ ID NO	Sequence
		12345678901234567890123456789012345678901
Wild type hIgG1 constant region	80	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPE <u>LL</u> GGPSV FLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFSVCSVMHEALHNHYTQKSLSLSP GK
Mutant hIgG1 constant region	81	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPE <u>AA</u> GGPSV FLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDG VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQV SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFSVCSVMHEALHNHYTQKSLSLSP GK
Ig kappa constant region	82	TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWK VDNALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEKHK VYACEVTHQGLSSPVTKSFNRGEC
Ig Lambda constant region	83	QPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAW KADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHR SYSCQVTHEGSTVEKTVAPTECS

20 The likelihood that a given framework residue would impact the binding properties of the antibody depends on its proximity to the CDR residues. Therefore, using the model structures,

the residues that differ between the murine and human sequences are ranked according to their distance from any atom in the CDRs. Those residues that fell within 4.5 Å of any CDR atom are identified as most important and are recommended to be candidates for retention of the murine residue in the humanized antibody (i.e., back mutation).

5 *In silico* constructed humanized antibodies are constructed using oligonucleotides. For each variable region cDNA, 6 oligonucleotides of 60-80 nucleotides each are designed to overlap each other by 20 nucleotides at the 5' and/or 3' end of each oligonucleotide. In an annealing reaction, all 6 oligonucleotides are combined, boiled, and annealed in the presence of dNTPs. DNA polymerase I, Large (Klenow) fragment (New England Biolabs #M0210, Beverly, MA.) is added to fill-in the approximately 40bp gaps between the overlapping oligonucleotides. PCR is performed to amplify the entire variable region gene using two outermost primers containing overhanging sequences complementary to the multiple cloning site in a modified pBOS vector (Mizushima, S. and Nagata, S. (1990) Nucleic Acids Res. 18: 17). The PCR products derived from each cDNA assembly are separated on an agarose gel and the band corresponding to the predicted variable region cDNA size is excised and purified. The variable heavy region is inserted in-frame onto a cDNA fragment encoding the human IgG1 constant region containing 2 hinge-region amino acid mutations by homologous recombination in bacteria. These mutations are a leucine to alanine change at position 234 (EU numbering) and a leucine to alanine change at position 235 (Lund et al. (1991) J. Immunol. 147:2657). The variable light chain region is inserted in-frame with the human kappa constant region by homologous recombination. Bacterial colonies are isolated and plasmid DNA extracted. cDNA inserts are sequenced in their entirety. Correct humanized heavy and light chains corresponding to each antibody are co-transfected into COS cells to transiently produce full-length humanized anti-human antibodies. Cell supernatants containing recombinant chimeric antibody are purified by Protein A Sepharose chromatography and bound antibody is eluted by addition of acid buffer. Antibodies are neutralized and dialyzed into PBS.

Example 1.3.2.3: Characterization Of Humanized Antibodies

The ability of purified humanized antibodies to inhibit a functional activity is determined, e.g., using the cytokine bioassay as described in Examples 1.2.2.A. The binding affinities of the humanized antibodies to recombinant human antigen are determined using surface plasmon resonance (Biacore®) measurement as described in Example 1.2.1.B. The IC₅₀ values from the bioassays and the affinity of the humanized antibodies are ranked. The humanized mAbs that fully maintain the activity of the parent hybridoma mAbs are selected as candidates for future development. The top 2-3 most favorable humanized mAbs are further characterized.

Example 1.3.2.3.A: Pharmacokinetic Analysis Of Humanized Antibodies

Pharmacokinetic studies are carried out in Sprague-Dawley rats and cynomolgus monkeys. Male and female rats and cynomolgus monkeys are dosed intravenously or subcutaneously with a single dose of 4mg/kg mAb and samples are analyzed using antigen capture ELISA, and pharmacokinetic parameters are determined by noncompartmental analysis. Briefly, ELISA plates are coated with goat anti-biotin antibody (5 mg/ml, 4°C, overnight), blocked with Superblock (Pierce), and incubated with biotinylated human antigen at 50 ng/ml in 10% Superblock TTBS at room temperature for 2 hours. Serum samples are serially diluted (0.5% serum, 10% Superblock in TTBS) and incubated on the plate for 30 minutes at room temperature. Detection is carried out with HRP-labeled goat anti human antibody and concentrations are determined with the help of standard curves using the four parameter logistic fit. Values for the pharmacokinetic parameters are determined by non-compartmental model using WinNonlin software (Pharsight Corporation, Mountain View, CA). Humanized mAbs with good pharmacokinetics profile (T_{1/2} is 8-13 days or better, with low clearance and excellent bioavailability 50-100%) are selected.

Example 1.3.2.3.B: Physicochemical And In Vitro Stability Analysis Of Humanized Monoclonal Antibodies**Size exclusion chromatography**

Antibodies are diluted to 2.5 mg/mL with water and 20 mL is analyzed on a Shimadzu HPLC system using a TSK gel G3000 SWXL column (Tosoh Bioscience, cat# k5539-05k). Samples are eluted from the column with 211 mM sodium sulfate, 92 mM sodium phosphate, pH 7.0, at a flow rate of 0.3 mL/minutes. The HPLC system operating conditions are the following:

Mobile phase:	211 mM Na ₂ SO ₄ , 92 mM Na ₂ HPO ₄ *7H ₂ O, pH 7.0
Gradient:	Isocratic
Flow rate:	0.3 mL/minute
Detector wavelength:	280 nm
Autosampler cooler temp:	4°C
Column oven temperature:	Ambient
Run time:	50 minutes

Tables 13 and 14 contain purity data of parent antibodies and DVD-Ig constructs expressed as percent monomer (unaggregated protein of the expected molecular weight) as determined by the above protocol.

Table 13: Purity of DVD-Ig Constructs as Determined by Size Exclusion Chromatography (NRP1, VEGF)

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	% Monomer (purity)
DVD695	NRP1	VEGF	96.8
DVD696	NRP1	VEGF	98.6

DVD695 and DVD696 showed excellent SEC profile >90% monomer. This DVD-Ig profile is similar to that observed for parent antibodies.

Table 14: Purity of DVD-Ig Constructs as Determined by Size Exclusion Chromatography (SOST, TNF)

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	% Monomer (purity)
DVD699	SOST	TNF	94.0
DVD700	SOST	TNF	93.2
DVD701	SOST	TNF	91.9

DVD699, DVD700 and DVD701 showed an excellent SEC profile with most DVD-Ig showing >90% monomer. This DVD-Ig profile is similar to that observed for parent antibodies.

10 SDS-PAGE

Antibodies are analyzed by sodium dodecyl sulfate - polyacrylamide gel electrophoresis (SDS-PAGE) under both reducing and non-reducing conditions. Adalimumab lot AFP04C is used as a control. For reducing conditions, the samples are mixed 1:1 with 2X tris glycine SDS-PAGE sample buffer (Invitrogen, cat# LC2676, lot# 1323208) with 100 mM DTT, and heated at 60°C for 30 minutes. For non-reducing conditions, the samples are mixed 1:1 with sample buffer and heated at 100°C for 5 minutes. The reduced samples (10 mg per lane) are loaded on a 12% pre-cast tris-glycine gel (Invitrogen, cat# EC6005box, lot# 6111021), and the non-reduced samples (10 mg per lane) are loaded on an 8%-16% pre-cast tris-glycine gel (Invitrogen, cat# EC6045box, lot# 6111021). SeeBlue Plus 2 (Invitrogen, cat#LC5925, lot# 1351542) is used as a molecular weight marker. The gels are run in a XCell SureLock mini cell gel box (Invitrogen, cat# EI0001) and the proteins are separated by first applying a voltage of 75 to stack the samples in the gel, followed by a constant voltage of 125 until the dye front reached the bottom of the gel. The running buffer used is 1X tris glycine SDS buffer, prepared from a 10X tris glycine SDS buffer (ABC, MPS-79-080106)). The gels are stained overnight with colloidal blue stain (Invitrogen cat# 46-7015, 46-7016) and destained with Milli-Q water until the background is clear. The stained gels are then scanned using an Epson Expression scanner (model 1680, S/N DASX003641).

Sedimentation Velocity Analysis

Antibodies are loaded into the sample chamber of each of three standard two-sector carbon epon centerpieces. These centerpieces have a 1.2 cm optical path length and are built with sapphire windows. PBS is used for a reference buffer and each chamber contained 140 μ L. All
 5 samples are examined simultaneously using a 4-hole (AN-60Ti) rotor in a Beckman ProteomeLab XL-I analytical ultracentrifuge (serial # PL106C01).

Run conditions are programmed and centrifuge control is performed using ProteomeLab (v5.6). The samples and rotor are allowed to thermally equilibrate for one hour prior to analysis (20.0 $\pm$ 0.1 $^{\circ}$ C). Confirmation of proper cell loading is performed at 3000 rpm and a single scan is
 10 recorded for each cell. The sedimentation velocity conditions are the following:

Sample Cell Volume: 420 mL

Reference Cell Volume: 420 mL

Temperature: 20 $^{\circ}$ C

Rotor Speed: 35,000 rpm

15 Time: 8:00 hours

UV Wavelength: 280 nm

Radial Step Size: 0.003 cm

Data Collection: One data point per step without signal averaging.

Total Number of Scans: 100

20 LC-MS molecular weight measurement of intact antibodies

Molecular weight of intact antibodies are analyzed by LC-MS. Each antibody is diluted to approximately 1 mg/mL with water. An 1100 HPLC (Agilent) system with a protein microtrap (Michrom Bioresources, Inc, cat# 004/25109/03) is used to desalt and introduce 5 mg of the sample into an API Qstar pulsar i mass spectrometer (Applied Biosystems). A short gradient is
 25 used to elute the samples. The gradient is run with mobile phase A (0.08% FA, 0.02% TFA in HPLC water) and mobile phase B (0.08% FA and 0.02% TFA in acetonitrile) at a flow rate of 50 mL/minute. The mass spectrometer is operated at 4.5 kvolts spray voltage with a scan range from 2000 to 3500 mass to charge ratio.

LC-MS molecular weight measurement of antibody light and heavy chains

30 Molecular weight measurement of antibody light chain (LC), heavy chain (HC) and deglycosylated HC are analyzed by LC-MS. Antibody is diluted to 1 mg/mL with water and the sample is reduced to LC and HC with a final concentration of 10 mM DTT for 30 minutes at 37 $^{\circ}$ C. To deglycosylate the antibody, 100 mg of the antibody is incubated with 2 mL of PNGase F, 5 mL of 10% N-octylglucoside in a total volume of 100 mL overnight at 37 $^{\circ}$ C. After

deglycosylation the sample is reduced with a final concentration of 10 mM DTT for 30 minutes at 37°C. An Agilent 1100 HPLC system with a C4 column (Vydac, cat# 214TP5115, S/N 060206537204069) is used to desalt and introduce the sample (5 mg) into an API Qstar pulsar i mass spectrometer (Applied Biosystems). A short gradient is used to elute the sample. The
 5 gradient is run with mobile phase A (0.08% FA, 0.02% TFA in HPLC water) and mobile phase B (0.08% FA and 0.02% TFA in acetonitrile) at a flow rate of 50 mL/minute. The mass spectrometer is operated at 4.5 kvolts spray voltage with a scan range from 800 to 3500 mass to charge ratio.

Peptide mapping

10 Antibody is denatured for 15 minutes at room temperature with a final concentration of 6 M guanidine hydrochloride in 75 mM ammonium bicarbonate. The denatured samples are reduced with a final concentration of 10 mM DTT at 37°C for 60 minutes, followed by alkylation with 50 mM iodoacetic acid (IAA) in the dark at 37°C for 30 minutes. Following alkylation, the sample is dialyzed overnight against four liters of 10 mM ammonium bicarbonate at 4°C. The
 15 dialyzed sample is diluted to 1 mg/mL with 10 mM ammonium bicarbonate, pH 7.8 and 100 mg of antibody is either digested with trypsin (Promega, cat# V5111) or Lys-C (Roche, cat# 11 047 825 001) at a 1:20 (w/w) trypsin/Lys-C:antibody ratio at 37°C for 4 hrs. Digests are quenched with 1 mL of 1 N HCl. For peptide mapping with mass spectrometer detection, 40 mL of the digests are separated by reverse phase high performance liquid chromatography (RPHPLC) on a
 20 C18 column (Vydac, cat# 218TP51, S/N NE9606 10.3.5) with an Agilent 1100 HPLC system. The peptide separation is run with a gradient using mobile phase A (0.02% TFA and 0.08% FA in HPLC grade water) and mobile phase B (0.02% TFA and 0.08% FA in acetonitrile) at a flow rate of 50 mL/minutes. The API QSTAR Pulsar i mass spectrometer is operated in positive mode at 4.5 kvolts spray voltage and a scan range from 800 to 2500 mass to charge ratio.

25 Disulfide Bond Mapping

To denature the antibody, 100 mL of the antibody is mixed with 300 mL of 8 M guanidine HCl in 100 mM ammonium bicarbonate. The pH is checked to ensure that it is between 7 and 8 and the samples are denatured for 15 minutes at room temperature in a final concentration of 6 M guanidine HCl. A portion of the denatured sample (100 mL) is diluted to
 30 600 mL with Milli-Q water to give a final guanidine-HCl concentration of 1 M. The sample (220 mg) is digested with either trypsin (Promega, cat # V5111, lot# 22265901) or Lys-C (Roche, cat# 11047825001, lot# 12808000) at a 1:50 trypsin or 1:50 Lys-C: antibody (w/w) ratios (4.4 mg enzyme: 220 mg sample) at 37°C for approximately 16 hours. An additional 5 mg of trypsin or Lys-C is added to the samples and digestion is allowed to proceed for an additional 2 hours at
 35 37°C. Digestions are stopped by adding 1 mL of TFA to each sample. Digested samples are

separated by RPHPLC using a C18 column (Vydac, cat# 218TP51 S/N NE020630-4-1A) on an Agilent HPLC system. The separation is run with the same gradient used for peptide mapping using mobile phase A (0.02% TFA and 0.08% FA in HPLC grade water) and mobile phase B (0.02% TFA and 0.08% FA in acetonitrile) at a flow rate of 50 mL/minute. The HPLC operating conditions are the same as those used for peptide mapping. The API QSTAR Pulsar i mass spectrometer is operated in positive mode at 4.5 kvolts spray voltage and a scan range from 800 to 2500 mass-to-charge ratio. Disulfide bonds are assigned by matching the observed MWs of peptides with the predicted MWs of tryptic or Lys-C peptides linked by disulfide bonds.

Free sulfhydryl determination

The method used to quantify free cysteines in an antibody is based on the reaction of Ellman's reagent, 5,5'-dithio-bis (2-nitrobenzoic acid) (DTNB), with sulfhydryl groups (SH) which gives rise to a characteristic chromophoric product, 5-thio-(2-nitrobenzoic acid) (TNB-). The reaction is illustrated in the formula:



The absorbance of the TNB- is measured at 412 nm using a Cary 50 spectrophotometer. An absorbance curve is plotted using dilutions of 2 mercaptoethanol (b-ME) as the free SH standard and the concentrations of the free sulfhydryl groups in the protein are determined from absorbance at 412 nm of the sample.

The b-ME standard stock is prepared by a serial dilution of 14.2 M b-ME with HPLC grade water to a final concentration of 0.142 mM. Then standards in triplicate for each concentration are prepared. Antibody is concentrated to 10 mg/mL using an amicon ultra 10,000 MWCO centrifugal filter (Millipore, cat# UFC801096, lot# L3KN5251) and the buffer is changed to the formulation buffer used for adalimumab (5.57 mM sodium phosphate monobasic, 8.69 mM sodium phosphate dibasic, 106.69 mM NaCl, 1.07 mM sodium citrate, 6.45 mM citric acid, 66.68 mM mannitol, pH 5.2, 0.1% (w/v) Tween). The samples are mixed on a shaker at room temperature for 20 minutes. Then 180 mL of 100 mM Tris buffer, pH 8.1 is added to each sample and standard followed by the addition of 300 mL of 2 mM DTNB in 10 mM phosphate buffer, pH 8.1. After thorough mixing, the samples and standards are measured for absorption at 412 nm on a Cary 50 spectrophotometer. The standard curve is obtained by plotting the amount of free SH and OD₄₁₂ nm of the b-ME standards. Free SH content of samples are calculated based on this curve after subtraction of the blank.

Weak Cation Exchange Chromatography

Antibody is diluted to 1 mg/mL with 10 mM sodium phosphate, pH 6.0. Charge heterogeneity is analyzed using a Shimadzu HPLC system with a WCX-10 ProPac analytical

column (Dionex, cat# 054993, S/N 02722). The samples are loaded on the column in 80% mobile phase A (10 mM sodium phosphate, pH 6.0) and 20% mobile phase B (10 mM sodium phosphate, 500 mM NaCl, pH 6.0) and eluted at a flow rate of 1.0 mL/minute.

Oligosaccharide Profiling

5 Oligosaccharides released after PNGase F treatment of antibody are derivatized with 2-aminobenzamide (2-AB) labeling reagent. The fluorescent-labeled oligosaccharides are separated by normal phase high performance liquid chromatography (NPHPLC) and the different forms of oligosaccharides are characterized based on retention time comparison with known standards.

10 The antibody is first digested with PNGaseF to cleave N-linked oligosaccharides from the Fc portion of the heavy chain. The antibody (200 mg) is placed in a 500 mL Eppendorf tube along with 2 mL PNGase F and 3 mL of 10% N-octylglucoside. Phosphate buffered saline is added to bring the final volume to 60 mL. The sample is incubated overnight at 37°C in an Eppendorf thermomixer set at 700 RPM. Adalimumab lot AFP04C is also digested with PNGase F as a control.

15 After PNGase F treatment, the samples are incubated at 95°C for 5 minutes in an Eppendorf thermomixer set at 750 RPM to precipitate out the proteins, then the samples are placed in an Eppendorf centrifuge for 2 minutes at 10,000 RPM to spin down the precipitated proteins. The supernatant containing the oligosaccharides are transferred to a 500 mL Eppendorf tube and dried in a speed-vac at 65°C.

20 The oligosaccharides are labeled with 2AB using a 2AB labeling kit purchased from Prozyme (cat# GKK-404, lot# 132026). The labeling reagent is prepared according to the manufacturer's instructions. Acetic acid (150 mL, provided in kit) is added to the DMSO vial (provided in kit) and mixed by pipeting the solution up and down several times. The acetic acid/DMSO mixture (100 mL) is transferred to a vial of 2-AB dye (just prior to use) and mixed
25 until the dye is fully dissolved. The dye solution is then added to a vial of reductant (provided in kit) and mixed well (labeling reagent). The labeling reagent (5 mL) is added to each dried oligosaccharide sample vial, and mixed thoroughly. The reaction vials are placed in an Eppendorf thermomixer set at 65°C and 700-800 RPM for 2 hours of reaction.

30 After the labeling reaction, the excess fluorescent dye is removed using GlycoClean S Cartridges from Prozyme (cat# GKI-4726). Prior to adding the samples, the cartridges are washed with 1 mL of milli-Q water followed with 5 washes of 1 mL 30% acetic acid solution. Just prior to adding the samples, 1 mL of acetonitrile (Burdick and Jackson, cat# AH015-4) is added to the cartridges.

After all of the acetonitrile passed through the cartridge, the sample is spotted onto the center of the freshly washed disc and allowed to adsorb onto the disc for 10 minutes. The disc is washed with 1 mL of acetonitrile followed by five washes of 1 mL of 96% acetonitrile. The cartridges are placed over a 1.5 mL Eppendorf tube and the 2-AB labeled oligosaccharides are eluted with 3 washes (400 µL each wash) of milli Q water.

The oligosaccharides are separated using a Glycosep N HPLC (cat# GKI-4728) column connected to a Shimadzu HPLC system. The Shimadzu HPLC system consisted of a system controller, degasser, binary pumps, autosampler with a sample cooler, and a fluorescent detector.

Stability at Elevated Temperatures

The buffer of antibody is either 5.57 mM sodium phosphate monobasic, 8.69 mM sodium phosphate dibasic, 106.69 mM NaCl, 1.07 mM sodium citrate, 6.45 mM citric acid, 66.68 mM mannitol, 0.1% (w/v) Tween, pH 5.2; or 10 mM histidine, 10 mM methionine, 4% mannitol, pH 5.9 using Amicon ultra centrifugal filters. The final concentration of the antibodies is adjusted to 2 mg/mL with the appropriate buffers. The antibody solutions are then filter sterilized and 0.25 mL aliquots are prepared under sterile conditions. The aliquots are left at either -80°C, 5°C, 25°C, or 40°C for 1, 2 or 3 weeks. At the end of the incubation period, the samples are analyzed by size exclusion chromatography and SDS-PAGE.

The stability samples are analyzed by SDS-PAGE under both reducing and non-reducing conditions. The procedure used is the same as described herein. The gels are stained overnight with colloidal blue stain (Invitrogen cat# 46-7015, 46-7016) and destained with Milli-Q water until the background is clear. The stained gels are then scanned using an Epson Expression scanner (model 1680, S/N DASX003641). To obtain more sensitivity, the same gels are silver stained using silver staining kit (Owl Scientific) and the recommended procedures given by the manufacturer is used.

Example 1.3.2.3.C: Efficacy Of A Humanized Monoclonal Antibody By Itself Or In Combination With Chemotherapy On The Growth Of Human Carcinoma Xenografts

Human cancer cells are grown *in vitro* to 99% viability, 85% confluence in tissue culture flasks. SCID female or male mice (Charles Rivers Labs) at 19-25 grams, are ear tagged and shaved. Mice are then inoculated subcutaneously into the right flank with 0.2 ml of 2×10^6 human tumor cells (1:1 matrigel) on study day 0. Administration (IP, Q3D/ week) of vehicle (PBS), humanized antibody, and/or chemotherapy is initiated after mice are size matched into separate cages of mice with mean tumor volumes of approximately 150 to 200 mm³. The tumors are measured by a pair of calipers twice a week starting on approximately day 10 post inoculation and the tumor volumes calculated according to the formula $V = L \times W^2/2$ (V: volume, mm³; L: length, mm; W: width, mm). Reduction in tumor volume is seen in animals treated with mAb alone or in combination

with chemotherapy relative to tumors in animals that received only vehicle or an isotype control mAb. **Example 1.3.2.3.D: FACS Based Redirected Cytotoxicity (rCTL) Assay**

Human CD3⁺ T cells were isolated from previously frozen isolated peripheral blood mononuclear cells (PBMC) by a negative selection enrichment column (R&D Systems, Minneapolis, MN; Cat.#HTCC-525). T cells were stimulated for 4 days in flasks (vent cap, Corning, Acton, MA) coated with 10mg/mL anti-CD3 (OKT-3, eBioscience, Inc., San Diego, CA) and 2 μ g/mL anti-CD28 (CD28.2, eBioscience, Inc., San Diego, CA) in D-PBS (Invitrogen, Carlsbad, CA) and cultured in 30U/mL IL-2 (Roche) in complete RPMI 1640 media (Invitrogen, Carlsbad, CA) with L-glutamine, 55mM α -ME, Pen/Strep, 10% FBS). T cells were then rested overnight in 30U/mL IL-2 before using in assay. DoHH2 or Raji target cells were labeled with PKH26 (Sigma-Aldrich, St. Louis, MO) according to manufacturer's instructions. RPMI 1640 media (no phenol, Invitrogen, Carlsbad, CA) containing L-glutamine and 10% FBS (Hyclone, Logan, UT) was used throughout the rCTL assay. (See Dreier et al. (2002) Int J Cancer 100:690).

Effector T cells (E) and targets (T) were plated at a final cell concentration of 10⁵ and 10⁴ cells/well in 96-well plates (Costar #3799, Acton, MA), respectively to give an E:T ratio of 10:1. DVD-Ig molecules were diluted to obtain concentration-dependent titration curves. After an overnight incubation cells are pelleted and washed with D-PBS once before resuspending in FACS buffer containing 0.1% BSA (Invitrogen, Carlsbad, CA), 0.1% sodium azide and 0.5 μ g/mL propidium iodide (BD) in D-PBS. FACS data was collected on a FACS Canto II machine (Becton Dickinson, San Jose, CA) and analyzed in Flowjo (Treestar). The percent live targets in the DVD-Ig treated samples divided by the percent total targets (control, no treatment) was calculated to determine percent specific lysis. IC50s were calculated in Prism (Graphpad).

The CD3 / CD 19 DVD-Ig (AB sequence IDs, AB002 + AB006; Example 2.7) was tested in the redirected toxicity assay for tumor cell killing. This DVD-Ig showed in vitro tumor killing with an IC50 = 5.0 pM. A CD3 / CD20 DVD-Ig was also tested for redirected toxicity and showed in vitro tumor killing with an IC50 = 325pM. The sequence of this CD3 / CD20 DVD-Ig was disclosed in US Patent Application Serial No. 20070071675.

Example 1.4: Generation of a DVD-Ig

DVD-Ig molecules capable of binding two antigens are constructed using two parent monoclonal antibodies, one against human antigen A, and the other against human antigen B, selected as described herein.

Example 1.4.1: Generation Of A DVD-Ig Having Two Linker Lengths

A constant region containing γ 1 Fc with mutations at 234, and 235 to eliminate ADCC/CDC effector functions is used. Four different anti-A/B DVD-Ig constructs are generated:

2 with short linker and 2 with long linker, each in two different domain orientations: V_A-V_B-C and V_B-V_A-C (see Table 15). The linker sequences, derived from the N-terminal sequence of human C1/Ck or CH1 domain, are as follows:

For DVDAB constructs:

5 light chain (if anti-A has λ): Short linker: QPKAAP (SEQ ID NO: 15); Long linker: QPKAAPSVTLFPP (SEQ ID NO: 16)

light chain (if anti-A has κ): Short linker: TVAAP (SEQ ID NO: 13); Long linker: TVAAPSVFIFPP (SEQ ID NO: 14)

heavy chain (γ 1): Short linker: ASTKGP (SEQ ID NO: 21); Long linker: 10 ASTKGPSVFPLAP (SEQ ID NO: 22)

For DVDBA constructs:

light chain (if anti-B has λ): Short linker: QPKAAP (SEQ ID NO: 15); Long linker: QPKAAPSVTLFPP (SEQ ID NO: 16)

light chain (if anti-B has κ): Short linker: TVAAP (SEQ ID NO: 13); Long linker: 15 TVAAPSVFIFPP (SEQ ID NO: 14)

heavy chain (γ 1): Short linker: ASTKGP (SEQ ID NO: 21); Long linker: ASTKGPSVFPLAP (SEQ ID NO: 22)

Heavy and light chain constructs are subcloned into the pBOS expression vector, and expressed in COS cells, followed by purification by Protein A chromatography. The purified 20 materials are subjected to SDS-PAGE and SEC analysis.

Table 15 below describes the heavy chain and light chain constructs used to express each anti-A/B DVD-Ig protein.

Table 15: Anti-A/B DVD-Ig Constructs

DVD-Ig protein	Heavy chain construct	Light chain construct
DVDABSL	DVDABHC-SL	DVDABLC-SL
DVDABLL	DVDABHC-LL	DVDABLC-LL
DVDBASL	DVDBAHC-SL	DVDBALC-SL
DVDBALL	DVDBAHC-LL	DVDBALC-LL

Example 1.4.2: Molecular cloning of DNA constructs for DVDABSL and DVDABLL

25 To generate heavy chain constructs DVDABHC-LL and DVDABHC-SL, VH domain of A antibody is PCR amplified using specific primers (3' primers contain short/long linker sequence

for SL/LL constructs, respectively); meanwhile VH domain of B antibody is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template
 5 for the subsequent overlapping PCR reaction. The overlapping PCR products are subcloned into Srf I and Sal I double digested pBOS-hCγ1,z non-a mammalian expression vector (Abbott) by using standard homologous recombination approach.

To generate light chain constructs DVDABLC-LL and DVDABLC-SL, VL domain of A antibody is PCR amplified using specific primers (3' primers contain short/long liner sequence for
 10 SL/LL constructs, respectively); meanwhile VL domain of B antibody is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping PCR products are
 15 subcloned into Srf I and Not I double digested pBOS-hCκ mammalian expression vector (Abbott) by using standard homologous recombination approach. Similar approach has been used to generate DVDBASL and DVDBALL as described below:

Example 1.4.3: Molecular cloning of DNA constructs for DVDBASL and DVDBALL

To generate heavy chain constructs DVDBAHC-LL and DVDBAHC-SL, VH domain of
 20 antibody B is PCR amplified using specific primers (3' primers contain short/long liner sequence for SL/LL constructs, respectively); meanwhile VH domain of antibody A is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template
 25 for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping PCR products are subcloned into Srf I and Sal I double digested pBOS-hCγ1,z non-a mammalian expression vector (Abbott) by using standard homologous recombination approach.

To generate light chain constructs DVDBALC-LL and DVDBALC-SL, VL domain of
 antibody B is PCR amplified using specific primers (3' primers contain short/long liner sequence
 30 for SL/LL constructs, respectively); meanwhile VL domain of antibody A is amplified using specific primers (5' primers contains short/long liner sequence for SL/LL constructs, respectively). Both PCR reactions are performed according to standard PCR techniques and procedures. The two PCR products are gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping

PCR products are subcloned into Srf I and Not I double digested pBOS-hCk mammalian expression vector (Abbott) by using standard homologous recombination approach.

Example 1.4.4: Construction and Expression of Additional DVD-Ig

Example 1.4.4.1: Preparation of DVD-Ig Vector Constructs

5 Parent antibody amino acid sequences for specific antibodies, which recognize specific antigens or epitopes thereof, for incorporation into a DVD-Ig can be obtained by preparation of hybridomas as described above or can be obtained by sequencing known antibody proteins or nucleic acids. In addition, known sequences can be obtained from the literature. The sequences can be used to synthesize nucleic acids using standard DNA synthesis or amplification
10 technologies and assembling the desired antibody fragments into expression vectors, using standard recombinant DNA technology, for expression in cells.

For example, nucleic acid codons were determined from amino acids sequences and oligonucleotide DNA was synthesized by Blue Heron Biotechnology, Inc. (www.blueheronbio.com) Bothell, WA USA. The oligonucleotides were assembled into 300-
15 2,000 base pair double-stranded DNA fragments, cloned into a plasmid vector and sequence-verified. Cloned fragments were assembled using an enzymatic process to yield the complete gene and subcloned into an expression vector. (See 7,306,914; 7,297,541; 7,279,159; 7,150,969; 20080115243; 20080102475; 20080081379; 20080075690; 20080063780; 20080050506; 20080038777; 20080022422; 20070289033; 20070287170; 20070254338; 20070243194;
20 20070225227; 20070207171; 20070150976; 20070135620; 20070128190; 20070104722; 20070092484; 20070037196; 20070028321; 20060172404; 20060162026; 20060153791; 20030215458; 20030157643).

A group of pHybE vectors (US Patent Application Serial No. 61/021,282) were used for parental antibody and DVD-Ig cloning. V1, derived from pJP183; pHybE-hCg1,z,non-a V2, was
25 used for cloning of antibody and DVD heavy chains with a wildtype constant region. V2, derived from pJP191; pHybE-hCk V2, was used for cloning of antibody and DVD light chains with a kappa constant region. V3, derived from pJP192; pHybE-hCl V2, was used for cloning of antibody and DVDs light chains with a lambda constant region. V4, built with a lambda signal peptide and a kappa constant region, was used for cloning of DVD light chains with a lambda-
30 kappa hybrid V domain. V5, built with a kappa signal peptide and a lambda constant region, was used for cloning of DVD light chains with a kappa-lambda hybrid V domain. V7, derived from pJP183; pHybE-hCg1,z,non-a V2, was used for cloning of antibody and DVD heavy chains with a (234,235 AA) mutant constant region.

Referring to Table 16, a number of vectors were used in the cloning of the parent antibodies and
35 DVD-Ig VH and VL chains.

Table 16: Vectors Used to Clone Parent Antibodies and DVD-Igs

ID	Heavy chain vector	Light chain vector
DVD050	V1	V2
DVD695	V1	V2
DVD695-C	V1	V2
DVD696	V1	V2
DVD696-C	V1	V2
DVD697	V1	V2
DVD698	V1	V2
DVD278	V1	V2
DVD699	V1	V2
DVD699-C	V1	V2
DVD700	V1	V2
DVD700-C	V1	V2
DVD701	V1	V2
DVD701-C	V1	V2
DVD702	V1	V2
DVD703	V1	V2
DVD704	V1	V2
DVD705	V1	V2
DVD706	V1	V2
DVD707	V1	V2
DVD708	V1	V2

Example 1.4.4.2: Transfection and Expression in 293 Cells

The DVD-Ig vector constructs are transfected into 293 cells for production of DVD-Ig protein. The 293 transient transfection procedure used is a modification of the methods published in Durocher et al. (2002) Nucleic Acids Res. 30(2):E9 and Pham et al. (2005) Biotech. Bioengineering 90(3):332-44. Reagents that were used in the transfection included:

- HEK 293-6E cells (human embryonic kidney cell line stably expressing EBNA1; obtained from National Research Council Canada) cultured in disposable Erlenmeyer flasks in a humidified incubator set at 130 rpm, 37°C and 5% CO₂.
- Culture medium: FreeStyle 293 Expression Medium (Invitrogen 12338-018) plus 25 µg/mL Geneticin (G418) (Invitrogen 10131-027) and 0.1% Pluronic F-68 (Invitrogen 24040-032).
- Transfection medium: FreeStyle 293 Expression Medium plus 10 mM HEPES (Invitrogen 15630-080).
- Polyethylenimine (PEI) stock: 1 mg/mL sterile stock solution, pH 7.0, prepared with linear 25kDa PEI (Polysciences) and stored at less than -15°C.

- Tryptone Feed Medium: 5% w/v sterile stock of Tryptone N1 (Organotechnie, 19554) in FreeStyle 293 Expression Medium.

Cell preparation for transfection: Approximately 2 - 4 hours prior to transfection, HEK 293-6E cells are harvested by centrifugation and resuspended in culture medium at a cell density of approximately 1 million viable cells per mL. For each transfection, 40 mL of the cell suspension is transferred into a disposable 250-mL Erlenmeyer flask and incubated for 2 - 4 hours.

Transfection: The transfection medium and PEI stock are prewarmed to room temperature (RT). For each transfection, 25µg of plasmid DNA and 50µg of polyethylenimine (PEI) are combined in 5 mL of transfection medium and incubated for 15 – 20 minutes at RT to allow the DNA:PEI complexes to form. For the BR3-Ig transfections, 25µg of BR3-Ig plasmid is used per transfection. Each 5-mL DNA:PEI complex mixture is added to a 40-mL culture prepared previously and returned to the humidified incubator set at 130 rpm, 37°C and 5% CO₂. After 20-28 hours, 5 mL of Tryptone Feed Medium is added to each transfection and the cultures are continued for six days.

Tables 17 and 18 contains the yield data for parent antibodies or DVD-Ig constructs expressed as milligrams per liter in 293 cells.

Table 17: Transient Expression in Yields of Parent Antibodies and DVD-Ig Constructs in 293 Cells (NRP1, VEGF)

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	Expression Yield (mg/L)
AB014	VEGF		52.4
AB016	NRP1		114.6
DVD050	NRP1	VEGF	27.2
DVD695	NRP1	VEGF	19.2
DVD696	NRP1	VEGF	19.6
DVD697	NRP1	VEGF	13.0
DVD698	NRP1	VEGF	13.8

All DVD – Igs with different linkers express well in HEK293 cells

Table 18: Transient Expression in Yields of Parent Antibodies and DVD-Ig Constructs in 293 Cells (SOST, TNF)

Parent Antibody or DVD-Ig ID	N-terminal Variable Domain (VD)	C-terminal Variable Domain (VD)	Expression Yield (mg/L)
AB017	TNF		52.4
AB050	SOST		77.8
DVD278	SOST	TNF	58.8
DVD699	SOST	TNF	56.0
DVD700	SOST	TNF	42.0
DVD701	SOST	TNF	25.0
DVD702	SOST	TNF	35.2
DVD703	SOST	TNF	45.0
DVD704	SOST	TNF	24.4
DVD705	SOST	TNF	55.0
DVD706	SOST	TNF	51.0
DVD707	SOST	TNF	44.2
DVD708	SOST	TNF	43.8

All DVD – Igs with different linkers express well in HEK293 cells. DVD – Ig yields are similar to that of parental antibodies.

5 **Example 1.4.5: Characterization and lead selection of A/B DVD Igs**

The binding affinities of anti-A/B DVD-Igs are analyzed on Biacore against both protein A and protein B. The tetravalent property of the DVD-Ig is examined by multiple binding studies on Biacore. Meanwhile, the neutralization potency of the DVD-Igs for protein A and protein B are assessed by bioassays, respectively, as described herein. The DVD-Ig molecules that best
 10 retain the affinity and potency of the original parent mAbs are selected for in-depth physicochemical and bio-analytical (rat PK) characterizations as described herein for each mAb. Based on the collection of analyses, the final lead DVD-Ig is advanced into CHO stable cell line development, and the CHO-derived material is employed in stability, pharmacokinetic and efficacy studies in cynomolgus monkey, and preformulation activities.

15

Example 2.0: Generation and Characterization of an Anti-TNF/IL-13 DVD-Ig

Example 2.1: Generation of Monoclonal Antibodies (mAbs) Against TNF and IL-13

DVD-Ig molecules capable of binding TNF and IL-13 were constructed using two pairs
 20 of parent mAbs, one with D2E7.1 (anti-TNF) and 13C5.5 (Anti-IL-13), and the second pair being D2E7 (anti-TNF) and 13C5.5L3F (Anti-IL-13). The two anti-TNF antibodies D2E7 and D2E7.1 have been disclosed previously (US Patent No. 7,541,031). The two anti-IL-13 antibodies 13C5.5

and 13C5.5L3F have been disclosed previously (US Patent Publication No. US20080171014). The variable domain amino acid sequences for the four antibodies are shown in Tables 42 and 43.

Example 2.2: Production And Characterization Of Monoclonal Antibodies Against TNF and IL-13

5 The four monoclonal antibodies were produced in mammalian cells by co-expressing the heavy and light chain construct of each mAb, and purified by protein A chromatography. The composition and purity of the purified mAbs were analyzed by SDS-PAGE in both reduced and non-reduced conditions. The purified proteins were characterized by the following assays:

Example 2.2.A: Mass Spectrometry And SEC Analysis Of Anti-TNF And Anti-IL-13 Mabs

10 For measuring molecular weight (MW) of light and heavy chains of mAbs, 10 μ L of MAbs (0.8 μ g/ μ L) was reduced by 1.0 M DTT solution (5 μ L). A PLRP-S, 8 μ , 4000A, and 1x150 mm protein column (Michrom BioResource, Auburn, MA) was used to separate the heavy and light chains of the MAbs. Agilent HP1100 Capillary HPLC (Agilent Technologies Inc., Palo Alto, CA) was used with the mass spectrometer QSTAR (Applied Biosystems, Foster City, CA).

15 The valco valve was set at 10 minutes to switch the flow from waste to MS for desalting sample. Buffer A was 0.02% TFA, 0.08% FA, 0.1% ACN and 99.8% HPLC-H₂O. Buffer B contained 0.02% TFA, 0.08% FA, 0.1% HPLC-H₂O, and 99.8% ACN. The HPLC flow rate was 50 μ L/min, and the sample injection volume was 8.0 μ L. The temperature of the column oven was set at 60°C, and separation gradient was: 5%B for 5 minutes; 5%B to 65%B for 35 minutes;

20 65%B to 95%B for another 5 minutes, and 95%B to 5%B for 5 minutes. TOFMS scan was from 800 to 2500 amu, and cycles were 3600. To determine the MW of full length MAbs, a Protein MicroTrap cartridge (Michrom BioResource, Auburn, MA) was used for desalting the sample. The HPLC gradient was: 5%B for 5 minutes; 5%B to 95%B in 1 minute; and from 95%B to 5%B in another 4 minutes. The QSTAR TOFMS scan was from 2000 to 3500 amu, and cycles were

25 899. All MS raw data were analyzed using the Analyst QS software (Applied Biosystems). For SEC analysis of the MAbs, purified MAbs and chimeric Abs, in PBS, were applied on a Superose 6 10/300 G2, 300 x 10 mm column (Amersham Bioscience, Piscataway, NJ). An HPLC instrument, Model 10A (Shimadzu, Columbia, MD) was used for SEC. All proteins were determined using UV detection at 280 nm and 214 nm. The elution was isocratic at a flow rate of

30 0.5 mL/min. For stability study, samples in the concentration range of 0.2-0.4 mg/ml in PBS underwent 3 freeze-thaw cycles between -80°C and 25°C, or were incubated at 4°C, 25°C, or 40°C, for 4 weeks and 8 weeks, followed by SEC analysis.

Example 2.2.B: Determination Of Antigen Binding Affinity Of Anti-TNF And Anti-IL-13
Mabs

The kinetics of mAb binding to rhTNF α and rhIL-13 was determined by surface plasmon resonance-based measurements with a Biacore 3000 instrument (Biacore AB, Uppsala, Sweden) using HBS-EP (10 mM HEPES, pH 7.4, 150 mM NaCl, 3 mM EDTA, and 0.005% surfactant P20) at 25°C. All chemicals were obtained from Biacore AB (Uppsala, Sweden) or otherwise from a different source as described herein. Approximately, 5000 RU of goat anti-human IgG Fc γ fragment specific polyclonal antibody (Pierce Biotechnology Inc, Rockford, IL) diluted in 10 mM sodium acetate (pH 4.5) was directly immobilized across a CM5 research grade biosensor chip using a standard amine coupling kit according to manufacturer's instructions and procedures at 25 mg/ml. Unreacted moieties on the biosensor surface were blocked with ethanolamine. Modified carboxymethyl dextran surface in flowcell 2 and 4 was used as a reaction surface. Unmodified carboxymethyl dextran without goat anti-human IgG in flow cell 1 and 3 was used as the reference surface. For kinetic analysis, rate equations derived from the 1:1 Langmuir binding model were fitted simultaneously to association and dissociation phases of all ten injections (using global fit analysis) using the Bioevaluation 4.0.1 software. Purified mAb samples were diluted in HEPES-buffered saline for capture across goat anti-human IgG Fc specific reaction surfaces and injected over reaction matrices at a flow rate of 5 ml/min. The association and dissociation rate constants, k_{on} (M⁻¹s⁻¹) and k_{off} (s⁻¹) were determined under a continuous flow rate of 25 ml/min. Rate constants were derived by making kinetic binding measurements at ten different antigen concentrations ranging from 1.25 to 1000 nM. The equilibrium dissociation constant (M) of the reaction between mAb and antigen was then calculated from the kinetic rate constants by the following formula: $KD = k_{off}/k_{on}$. Aliquots of antigen samples were also simultaneously injected over a blank reference and reaction CM surface to record and subtract any nonspecific binding background to eliminate the majority of the refractive index change and injection noise. Surfaces were regenerated with two subsequent 25 ml injections of 10 mM Glycine (pH 1.5) at a flow rate of 5 ml/min. The anti-Fc antibody immobilized surfaces were completely regenerated and retained their full capture capacity over twelve cycles. The apparent stoichiometry of the captured mAb- antigen complex was calculated under saturating binding conditions (steady-state equilibrium) using the following formula:

$$\text{Stoichiometry} = \frac{\text{antigen response (RU)}}{\text{mAb response (RU)}} \times \frac{\text{mAb MW}}{\text{antigen MW}}$$

The binding kinetic parameters of these four antibodies against their antigens are shown in Table 19.

Example 2.2.C: Determination Of Biological Activity Of Anti-TNF And Anti-IL-13 Mabs

The biological activity of anti-TNF/IL-13 DVD-Ig proteins was measured using L929 (for anti-TNF) and A549 (for anti-IL-13) bioassays.

Example 2.2.C.1: Determination Of Biological Activity Of Anti-TNF Mabs By L929**5 Bioassay**

Human recombinant TNF α (rhTNF α) causes cell cytotoxicity to murine L929 cells after an incubation period of 18-24 hours. Human anti-hTNF α antibodies were evaluated in L929 assays by coincubation of antibodies with rhTNF α and the cells as follows. A 96-well microtiter plate containing 100 μ l of anti-hTNF α Abs was serially diluted 1/3 down the plate in duplicates using RPMI medium containing 10% fetal bovine serum (FBS). Fifty microliters of rhTNF α was added for a final concentration of 500 pg/ml in each sample well. The plates were then incubated for 30 minutes at room temperature. 50 μ l of TNF α -sensitive L929 mouse fibroblasts cells were added for a final concentration of 5×10^4 cells per well, including 1 μ g/ml Actinomycin-D. Controls included medium plus cells and rhTNF α plus cells. These controls, and a TNF α standard curve, ranging from 2 ng/ml to 8.2 pg/ml, were used to determine the quality of the assay and provide a window of neutralization. The plates were then incubated overnight (18-24 hours) at 37° C in 5% CO₂. One hundred microliters of medium was removed from each well and 50 μ l of 5 mg/ml 3-(4,4-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT; commercially available from Sigma Chemical Co., St. Louis, MO) in PBS was added. The plates were then incubated for 4 hours at 37° C. Fifty microliters of 20% sodium dodecyl sulfate (SDS) was then added to each well and the plates were incubated overnight at 37° C. The optical density at 570/630 nm was measured, curves were plotted for each sample and IC₅₀s were determined by standard methods.

Example 2.2.C.2: Determination Of Biological Activity Of Anti-IL-13 Mabs By A549**25 Bioassay**

The ability of anti-human IL-13 antibodies to inhibit the human IL-13 induced production of TARC (CCL-17) by A-549 cells was analyzed as follows. A-549 cells were seeded on day one in 96-well plate (2E5 cells/well) in RPMI growth medium (with 10% FBS). On day two, the medium was replaced with fresh RPMI growth medium containing 400ng/ml rhTNF (100ml/well). Meanwhile, various concentrations of immunized mouse serum, murine hybridoma supernatant or purified anti-human IL-13 antibodies were preincubated for one hour at 37°C with 10 ng/ml recombinant purified human IL-13 or IL-13 variant in 100mL RPMI complete medium in a microtiter plate (U-bottom, 96-well, Costar). The antibody plus recombinant purified human IL-13 mixture was then added (100ml/well) to the TNF-treated A-549 cells, with the final volume of 200ml/well (final IL-13 and TNF concentrations were 5ng/ml and 200ng/ml, respectively), and incubated for 18 hours at 37°C. After incubation, 150mL of cell-free supernatant was withdrawn

from each well and the level of human TARC produced was measured using a human TARC ELISA (R&D Systems Cat#DDN00). Neutralization potencies of IL-13 antagonists were calculated based on the average TARC values for each data point and the IC₅₀ values determined by GraphPad Prism software. The neutralization potency of the two anti-IL-13 mAbs are shown in Table 19.

Table 19: Characterization Of Anti-TNF And Anti-IL-13 Monoclonal Antibodies

mAb	Specificity	K _D (nM)	IC ₅₀ (nM)
D2E7	TNF	0.04	0.08
D2E7.1	TNF	0.05	0.05
13C5.5	IL-13	0.15	0.20
13C5.5L3F	IL-13	0.24	0.31

Example 2.3: Construction Of Anti-TNF/IL-13 Dual Variable Domain Immunoglobulins (DVD-Ig)

DVD-Ig molecules capable of binding TNF and IL-13 were constructed using two pairs of parent mAbs, one with D2E7.1 (anti-TNF) and 13C5.5 (Anti-IL-13), and the second pair being D2E7 (anti-TNF) and 13C5.5L3F (Anti-IL-13). A constant region containing gamma Fc with mutations at 234, and 235 to eliminate ADCC/CDC effector functions was used. Several different anti-TNF/IL-13 DVD-Ig constructs were generated with various linkers, all in the orientation of V_{TNF}-V_{IL-13}-C. DVD-Ig heavy and light chains with various linker length, ranging from 0 to 12 in VL and 0-13 in VH were constructed in order to examine the optimal linker length for each of the LC and HC. The linker sequences, derived from the N-terminal sequence of human C1/Ck or CH1 domain, are as follows:

Heavy chain linkers (with identifiers):

AS (HC2),

ASTK (HC4) (SEQ ID NO: 58),

ASTKGP (HC6) (SEQ ID NO: 21),

ASTKGPSV (HC8) (SEQ ID NO: 59),

ASTKGPSVFP (HC10) (SEQ ID NO: 60), and

ASTKGPSVFPLAP (HC13) (SEQ ID NO: 22)

Light chain linkers (with identifiers):

TVA (LC3),

TVAAP (LC5) (SEQ ID NO: 13),

TVAAPSV (LC7) (SEQ ID NO: 61),

TVAAPSVFI (LC9) (SEQ ID NO: 62), and

TVAAPSVFIFPP (LC12) (SEQ ID NO: 14)

To generate heavy chain constructs of DVD-Igs, VH domain of D2E7.1 or D2E7 was PCR amplified using specific primers (3' primers contain an appropriate linker sequence); meanwhile VH domain of 13C5.5 or 13C5.5L3F was amplified using specific primers (5' primers contains an appropriate linker sequence). Both PCR reactions were performed according to standard PCR techniques and procedures. The two PCR products were gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction. The overlapping PCR products were subcloned into Srf I and Sal I double digested pBOS-hCg1,z non-a mammalian expression vector (Abbott) by using standard homologous recombination approach.

To generate light chain constructs of DVD-Igs, VL domain of D2E7.1 or D2E7 was PCR amplified using specific primers (3' primers contain an appropriate linker sequence); meanwhile VL domain of 13C5.5 or 13C5.5L3F was amplified using specific primers (5' primers contains an appropriate linker sequence). Both PCR reactions were performed according to standard PCR techniques and procedures. The two PCR products were gel-purified, and used together as overlapping template for the subsequent overlapping PCR reaction using standard PCR conditions. The overlapping PCR products were subcloned into Srf I and Not I double digested pBOS-hCk mammalian expression vector (Abbott) by using standard homologous recombination approach.

The amino acid sequence of VH and VL of anti-hTNF/IL-13 DVD-Ig proteins generated from D2E7.1 (anti-TNF) and 13C5.5 (anti-IL-13) monoclonal antibodies are shown in Table 42. The amino acid sequence of VH and VL of anti-hTNF/IL-13 DVD-Ig proteins generated from D2E7 (anti-TNF) and 13C5.5L3F (anti-IL-13) monoclonal antibodies are shown in Table 43. All heavy and light chain constructs contained a mutant hIgG1 constant region and Ig kappa constant region (Table 12) and are subcloned into the pBOS expression vector, and expressed in COS cells, followed by purification by Protein A chromatography. The purified materials are subjected to SDS-PAGE and SEC analysis.

In addition, heavy chains and light chains with various linker sizes were paired as shown in the matrix (Tables 20 and 21) in order to identify the optimal combination of linker choices for this mAb pair.

Example 2.4: Expression and Purification of anti-TNF/IL-13 DVD-Igs

The heavy and light chain of each construct was subcloned into mammalian expression vectors, respectively, and sequenced to ensure accuracy. The plasmids encoding the heavy and light chains of each construct were transiently expressed in human embryonic kidney 293 cells (American Type Culture Collection, Manassas, VA). The cell culture media was harvested 72 hr-post transient transfection and antibodies purified using protein A chromatography (Pierce, Rockford, IL) according to manufacturer's instructions. The Abs were analyzed by SDS-PAGE

and quantitated by A280 and BCA (Pierce, Rockford, IL). Table 20 shows that different D2E7.1-13C5.5 DVD-Ig molecules were generated by co-expressing different pairs of heavy chain and light chain constructs, in which various linkers of different lengths were used in both HC and LC and paired in all possible combinations. Similarly, different D2E7-13C5.5L3F DVD-Ig molecules were generated by co-expressing different pairs of heavy chain and light chain constructs, in which 2 linkers of different lengths were used in both HC and LC and paired in all possible combinations (Table 21).

Table 20: Generation Of Different D2E7.1-13C5.5 DVD-Ig Proteins By Co-Expression Of Different Pairs Of Heavy Chain And Light Chain

Construct		D2E7.1-13C5.5 light chain					
	Linker length	LC0	LC3	LC5	LC7	LC9	LC12
D2E7.1-13C5.5 heavy chain	HC0	LC0HC0	LC3HC0	LC5HC0	LC7HC0	LC9HC0	LC12HC0
	HC2	LC0HC2	LC3HC2	LC5HC2	LC7HC2	LC9HC2	LC12HC2
	HC4	LC0HC4	LC3HC4	LC5HC4	LC7HC4	LC9HC4	LC12HC4
	HC6	LC0HC6	LC3HC6	LC5HC6	LC7HC6	LC9HC6	LC12HC6
	HC8	LC0HC8	LC3HC8	LC5HC8	LC7HC8	LC9HC8	LC12HC8
	HC10	LC0HC10	LC3HC10	LC5HC10	LC7HC10	LC9HC10	LC12HC10
	HC13	LC0HC13	LC3HC13	LC5HC13	LC7HC13	LC9HC13	LC12HC13

Table 21: Generation Of Different D2E7-13C5.5L3F DVD-Ig Proteins By Co-Expression Of Different Pairs Of Heavy Chain And Light Chains

Construct		D2E7-13C5.5L3F light chain	
	Linker length	LC5	LC12
D2E7-13C5.5L3F heavy chain	HC6	LC5HC6	LC12HC6
	HC13	LC5HC13	LC12HC13

Example 2.5: Mass Spectrometry And SEC Analysis Of Anti-TNF/IL-13 DVD-Ig

For measuring molecular weight (MW) of light and heavy chains of DVD-Ig, 10 µL of DVD-Ig (0.8 µg/µL) was reduced by 1.0 M DTT solution (5 µL). A PLRP-S, 8u, 4000A, and 1x150 mm protein column (Michrom BioResource, Auburn, MA) was used to separate heavy and light chains of DVD-Ig. Agilent HP1100 Capillary HPLC (Agilent Technologies Inc., Palo Alto, CA) was used with the mass spectrometer QSTAR (Applied Biosystems, Foster City, CA). The valco valve was set at 10 minutes to switch the flow from waste to MS for desalting sample.

Buffer A was 0.02% TFA, 0.08% FA, 0.1% ACN and 99.8% HPLC-H₂O. Buffer B contained 0.02% TFA, 0.08% FA, 0.1% HPLC-H₂O, and 99.8% ACN. The HPLC flow rate was 50 uL/min, and the sample injection volume was 8.0 mL. The temperature of the column oven was set at 60°C, and separation gradient was: 5%B for 5 minutes; 5%B to 65%B for 35 minutes; 5 65%B to 95%B for another 5 minutes, and 95%B to 5%B for 5 minutes. TOFMS scan was from 800 to 2500 amu, and cycles were 3600. To determine the MW of full length DVD-Ig, a Protein MicroTrap cartridge (Michrom BioResource, Auburn, MA) was used for desalting the sample. The HPLC gradient was: 5%B for 5 minutes; 5%B to 95%B in 1 minute; and from 95%B to 5%B in another 4 minutes. The QSTAR TOFMS scan was from 2000 to 3500 amu, and cycles were 10 899. All MS raw data were analyzed using the Analyst QS software (Applied Biosystems). For SEC analysis of the DVD-Ig, purified DVD-Ig and chimeric Abs, in PBS, were applied on a Superose 6 10/300 G2, 300 x 10 mm column (Amersham Bioscience, Piscataway, NJ). An HPLC instrument, Model 10A (Shimadzu, Columbia, MD) was used for SEC. All proteins were determined using UV detection at 280 nm and 214 nm. The elution was isocratic at a flow rate of 15 0.5 mL/min. For stability study, samples in the concentration range of 0.2-0.4 mg/ml in PBS underwent 3 freeze-thaw cycles between -80°C and 25°C, or were incubated at 4°C, 25°C, or 40°C, for 4 weeks and 8 weeks, followed by SEC analysis.

DVD-Ig and chimeric Abs were purified by protein A chromatography. The purification yield (3-5mg/L) was consistent with hIgG quantification of the expression medium for each 20 protein. The composition and purity of the purified DVD-Igs and chimeric Abs were analyzed by SDS-PAGE in both reduced and non-reduced conditions. In non-reduced condition, each of the four proteins migrated as a single band. The DVD-Ig proteins showed larger M.W. than the chimeric Abs, as expected. In non-reducing condition, each of the four proteins yielded two bands, one heavy chain and one light chain. Again, the heavy and light chains of the DVD-Igs 25 were larger in size than that of the chimeric Abs. The SDS-PAGE showed that each DVD-Ig is expressed as a single species, and the heavy and light chains are efficiently paired to form an IgG-like molecule. The sizes of the heavy and light chains as well as the full-length protein of two DVD-Ig molecules are consistent with their calculated molecular mass based on amino acid sequences.

30 In order to determine the precise molecular weight of DVD-Ig, mass spectrometry was employed. The experimentally determined molecular mass of each DVD-Ig, including the light chain, heavy chain, and the full-length protein, is in good agreement with the predicted value. To further study the physical properties of DVD-Ig in solution, size exclusion chromatography (SEC) was used to analyze each protein, which exhibited largely monomeric properties.

35

Example 2.6: Determination Of IL-13 Binding Activity Of Anti-TNF/IL-13 DVD-Igs By ELISA

The binding activity of all DVD-Igs were first characterized or screened by a high through-put ELISA for their anti-IL-13 activity. Biotinylated human IL-13 was immobilized on anti-biotin polyclonal antibody coated ELISA plate. DVD-Igs were titrated to various concentrations and added to the IL-13-captured plate, followed by incubation for 1 hour at 27°C. After washing, bound DVD-Ig proteins were detected by HRP-conjugated goat anti-human Fc polyclonal antibody. The DVD-Ig molecules exhibited better binding properties were further tested for kinetic binding affinity to both TNF and IL-13 by Biacore and neutralization potency in cell-based bioassays.

Example 2.7: Determination Of Antigen Binding Affinity Of Anti-TNF/IL-13 DVD-Igs By Biacore

The kinetics of DVD-Ig binding to rhTNF α and rhIL-13 was determined by surface plasmon resonance-based measurements with a Biacore 3000 instrument (Biacore AB, Uppsala, Sweden) using HBS-EP (10 mM HEPES, pH 7.4, 150 mM NaCl, 3 mM EDTA, and 0.005% surfactant P20) at 25°C. All chemicals were obtained from Biacore AB (Uppsala, Sweden) or otherwise from a different source as described herein. Approximately, 5000 RU of goat anti-human IgG Fc γ fragment specific polyclonal antibody (Pierce Biotechnology Inc, Rockford, IL) diluted in 10 mM sodium acetate (pH 4.5) was directly immobilized across a CM5 research grade biosensor chip using a standard amine coupling kit according to manufacturer's instructions and procedures at 25 mg/ml. Unreacted moieties on the biosensor surface were blocked with ethanolamine. Modified carboxymethyl dextran surface in flowcell 2 and 4 was used as a reaction surface. Unmodified carboxymethyl dextran without goat anti-human IgG in flow cell 1 and 3 was used as the reference surface. For kinetic analysis, rate equations derived from the 1:1 Langmuir binding model were fitted simultaneously to association and dissociation phases of all ten injections (using global fit analysis) using the Bioevaluation 4.0.1 software. Purified DVD-Ig samples were diluted in HEPES-buffered saline for capture across goat anti-human IgG Fc specific reaction surfaces and injected over reaction matrices at a flow rate of 5 ml/minute. The association and dissociation rate constants, k_{on} (M⁻¹s⁻¹) and k_{off} (s⁻¹) were determined under a continuous flow rate of 25 ml/minute. Rate constants were derived by making kinetic binding measurements at ten different antigen concentrations ranging from 1.25 to 1000 nM. The equilibrium dissociation constant (M) of the reaction between DVD-Ig and antigen was then calculated from the kinetic rate constants by the following formula: $KD = k_{off}/k_{on}$. Aliquots of antigen samples were also simultaneously injected over a blank reference and reaction CM surface to record and subtract any nonspecific binding background to eliminate the majority of the refractive index change and injection noise. Surfaces were regenerated with two subsequent 25

ml injections of 10 mM Glycine (pH 1.5) at a flow rate of 5 ml/minute. The anti-Fc antibody immobilized surfaces were completely regenerated and retained their full capture capacity over twelve cycles. The apparent stoichiometry of the captured DVD-Ig- antigen complex was calculated under saturating binding conditions (steady-state equilibrium) using the following

5 formula:

$$\text{Stoichiometry} = \frac{\text{antigen response (RU)}}{\text{DVD response (RU)}} \times \frac{\text{DVD-Ig MW}}{\text{antigen MW}}$$

The Biacore analysis indicated that all DVD-Igs tested exhibit binding to both TNF and
10 IL-13. The binding affinity parameters of all DVD-Igs against TNF and IL-13 are shown in Tables 22 and 23.

Table 22: Binding Affinity Of D2E7.1-13C5.5 DVD-Ig Molecules For IL-13 And TNF

D2E7.1-13C5.5 DVD-Ig	Binding for IL-13 by ELISA (ED₅₀, nM)	K_D for IL-13 by Biacore (nM)	K_D for TNF by Biacore (nM)
LC0HC0			
LC0HC2	0.035		
LC0HC4	0.04		
LC0HC6	0.04	0.351	0.081
LC0HC8	0.22		
LC0HC10	0.68		
LC0HC13	0.04	0.571	0.321
LC3HC0	0.014	0.147	0.279
LC3HC2	0.34		
LC3HC4	5.42		
LC3HC6	2.25		
LC3HC8	0.16		
LC3HC10	0.38		
LC3HC13	0.035		
LC5HC0	0.049	0.640	0.156
LC5HC2	0.93		
LC5HC4	0.68		
LC5HC6	0.31	0.139	0.134
LC5HC8	0.36		
LC5HC10	0.31		

D2E7.1-13C5.5 DVD-Ig	Binding for IL-13 by ELISA (ED₅₀, nM)	K_D for IL-13 by Biacore (nM)	K_D for TNF by Biacore (nM)
LC5HC13	0.043	0.538	0.067
LC7HC0	0.07		
LC7HC2	0.058		
LC7HC4	0.042		
LC7HC6	0.069		
LC7HC8	0.038		
LC7HC10	0.032		
LC7HC13	0.019	0.567	0.051
LC9HC0	0.057		
LC9HC2	0.041		
LC9HC4	0.024	0.615	0.142
LC9HC6	0.03	0.743	0.098
LC9HC8	0.027		
LC9HC10	0.069		
LC9HC13	0.089		
LC12HC0	0.024	0.199	0.197
LC12HC2	0.02		
LC12HC4	0.019		
LC12HC6	0.030	0.235	0.059
LC12HC8	0.057		
LC12HC10	0.044		
LC12HC13	0.016	0.442	0.069

Table 23: Binding Affinity Of D2E7-13C5.5L3F DVD-Ig Molecules For IL-13 And TNF

D2E7-13C5.5L3F DVD-Ig	K_D for IL-13 by Biacore (nM)	K_D for TNF by Biacore (nM)
LC5HC6	1.23	0.030
LC5HC13	0.474	0.014
LC12HC6	0.402	0.008
LC12HC13		

Example 2.7: Determination of biological activity of anti-TNF/IL-13 DVD-Ig

5 The biological activity of anti-TNF/IL-13 DVD-Ig proteins was measured using L929 (for anti-TNF) and A549 (for anti-IL-13) bioassays. .

As shown in Tables 24 and 25, all DVD-Igs tested were able to neutralize TNF and IL-13.

Table 24: Neutralization Activity Of D2E7.1-13C5.5 DVD-Ig Molecules For IL-13 And TNF

D2E7.1-13C5.5 DVD-Ig	Anti-IL-13 activity by A549 bioassay (IC₅₀, nM)	Anti-TNF activity by L929 bioassay (IC₅₀, nM)
LC0HC0		
LC0HC2	5.32	0.362
LC0HC4	7.66	0.575
LC0HC6	4.76	0.529
LC0HC8		
LC0HC10		
LC0HC13	3.82	0.616
LC3HC0	2.51	1.63
LC3HC2		
LC3HC4		
LC3HC6		
LC3HC8		
LC3HC10		
LC3HC13		
LC5HC0	4.26	0.063
LC5HC2		
LC5HC4		
LC5HC6	4.29	0.67
LC5HC8		
LC5HC10		
LC5HC13	0.691	0.372
LC7HC0		
LC7HC2		
LC7HC4		
LC7HC6		
LC7HC8		
LC7HC10		
LC7HC13	0.80	0.052
LC9HC0		
LC9HC2		

D2E7.1-13C5.5 DVD-Ig	Anti-IL-13 activity by A549 bioassay (IC₅₀, nM)	Anti-TNF activity by L929 bioassay (IC₅₀, nM)
LC9HC4	3.10	0.250
LC9HC6	2.38	0.178
LC9HC8		
LC9HC10		
LC9HC13		
LC12HC0	1.62	0.063
LC12HC2	1.03	0.065
LC12HC4	0.99	0.073
LC12HC6	0.666	0.208
LC12HC8		
LC12HC10		
LC12HC13	0.89	0.063

Table 25 Neutralization Activity Of D2E7-13C5.5L3F DVD-Ig Molecules For IL-13 And TNF

D2E7-13C5.5L3F DVD-Ig	Anti-IL-13 activity by A549 bioassay (IC₅₀, nM)	Anti-TNF activity by L929 bioassay (IC₅₀, nM)
LC5HC6	1.84	0.095
LC5HC13	0.34	0.032
LC12HC6	0.52	0.028
LC12HC13		

5 **Example 3: Generation and Characterization of Dual Variable Domain Immunoglobulins (DVD-Ig)**

Dual variable domain immunoglobulins (DVD-Ig) using parent antibodies with known amino acid sequences were generated by synthesizing polynucleotide fragments encoding DVD-Ig variable heavy and DVD-Ig variable light chain sequences and cloning the fragments into a
10 pHybC-D2 vector according to Example 1.4.4.1. The DVD-Ig constructs were cloned into and expressed in 293 cells as described in Example 1,4.4.2. The DVD-Ig protein was purified according to standard methods. Functional characteristics were determined according to the methods described in Example 1.2.1 and 1.2.2 as indicated.

DVD-Ig VH and VL chains for the DVD-Igs of the invention are provided below.

Example 3.1: Generation of NRP1 (seq. 1) and VEGF (seq. 1) DVD-Igs with Linker Set 1

Table 26

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
84	DVD050H	AB016VH	AB014VH	EVQLVESGGGLVQPGGSLRLSCAASGFSFSSEPIS WVRQAPGKGLEWVSSITGKNGYTTYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARWGKKVYG MDVWGQGTLVTVSS ASTKGPE VQLVESGGGLVQPG GSLRLSCAASGYTFTNYGMNWVRQAPGKGLEWVGW INTYTGEPTYAADFKRRFTFSLDTSKSTAYLQMNS LRAEDTAVYYCAKYPHYYGSSHWYFDVWGQGTLVT VSS
85	DVD050L	AB016VL	AB014VL	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLAW YQQKPGKAPKLLIYGASSRASGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQQYMSVPITFGQGTKVE IKR TVAAP DIQMTQSPSSLSASVGDRVTITCSASQ DISNYLNWYQQKPGKAPKVLIIYFTSSLHSGVPSRF SGSGSGTDFTLTISSLQPEDFATYYCQQYSTVPWT FGQGTKVEIKR

5

Example 3.2: Generation of NRP1 (seq. 1) and VEGF (seq. 1) DVD-Igs with Linker Set 2

Table 27

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
86	DVD695H	AB016VH	AB014VH	EVQLVESGGGLVQPGGSLRLSCAASGFSFSSEPIS WVRQAPGKGLEWVSSITGKNGYTTYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARWGKKVYG MDVWGQGTLVTVSS ASTKGPE VQLVESGGGLVQPG GSLRLSCAASGYTFTNYGMNWVRQAPGKGLEWVGW INTYTGEPTYAADFKRRFTFSLDTSKSTAYLQMNS LRAEDTAVYYCAKYPHYYGSSHWYFDVWGQGTLVT VSS
87	DVD695L	AB016VL	AB014VL	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLAW YQQKPGKAPKLLIYGASSRASGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQQYMSVPITFGQGTKVE IKR TVDDDDKAAP DIQMTQSPSSLSASVGDRVTIT CSASQDISNYLNWYQQKPGKAPKVLIIYFTSSLHSG VPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYS TVPWTFGQGTKVEIKR

Example 3.3: Generation of NRP1 (seq. 1) and VEGF (seq. 1) DVD-Igs with Linker Set 3

28

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
88	DVD696H	AB016VH	AB014VH	EVQLVESGGGLVQPGGSLRLSCAASGFSFSSEPISWVRQAPGKGLEWVSSITGKNGYTTYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARWGKKVYGMVWGQGTLVTVSS ASTKGPE VQLVESGGGLVQPGGSLRLSCAASGYTFTNYGMNWVRQAPGKGLEWVGWINTYTGEPTYAADFKRRFTFSLDTSKSTAYLQMNSLRAEDTAVYYCAKYPHYYGSSHWYFDVWGQGTLTVTVSS
89	DVD696L	AB016VL	AB014VL	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLAWYQQKPGKAPKLLIYGASSRASGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYMSVPITFGQGTKVEIKR LVPRGSAAP DIQMTQSPSSLSASVGDRVTITCSASQDISNYLNWYQQKPGKAPKVLIIYFTSSLHSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSTVPWTFGQGTKVEIKR

5 Example 3.4: Generation of NRP1 (seq. 1) and VEGF (seq. 1) DVD-Igs with Linker Set 4

Table 29

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
90	DVD697H	AB016VH	AB014VH	EVQLVESGGGLVQPGGSLRLSCAASGFSFSSEPISWVRQAPGKGLEWVSSITGKNGYTTYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARWGKKVYGMVWGQGTLVTVSS GGGGGGGPE VQLVESGGGLVQPGGSLRLSCAASGYTFTNYGMNWVRQAPGKGLEWVGWINTYTGEPTYAADFKRRFTFSLDTSKSTAYLQMNSLRAEDTAVYYCAKYPHYYGSSHWYFDVWGQGTLVTVSS
91	DVD697L	AB016VL	AB014VL	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLAWYQQKPGKAPKLLIYGASSRASGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYMSVPITFGQGTKVEIKR GGGGGGP DIQMTQSPSSLSASVGDRVTITCSASQDISNYLNWYQQKPGKAPKVLIIYFTSSLHSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSTVPWTFGQGTKVEIKR

Example 3.5: Generation of NRP1 (seq. 1) and VEGF (seq. 1) DVD-Igs with Linker Set 5

Table 30

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
92	DVD698H	AB016VH	AB014VH	EVQLVESGGGLVQPGGSLRLSCAASGFSFSSEPIS WVRQAPGKGLEWVSSI TGKNGY TYYADSVKGRFTI SADTSKNTAYLQMNSLRAEDTAVYYCARWGKKVYG MDVWGQGT LVTVSSGGGGGGGGPEVQLVESGGGLV QPGGSLRLSCAASGYTFTNYGMNWVRQAPGKGLEW VGWINTYTGEPTYAADFKRRFTFSLDTSKSTAYLQ MNSLRAEDTAVYYCAKYPHYYGSSHWFVDVWGQGT LVTVSS
93	DVD698L	AB016VL	AB014VL	DIQMTQSPSSLSASVGDRVITITCRASQSISSYLAW YQQKPGKAPKLLIYGASSRASGVPSRFSGSGSGTD FTLTISSLQPEDFATYYCQQYMSVPITFGQGTKVE IKRGGGGGGGPDIQMTQSPSSLSASVGDRVITITCS ASQDISNYLNWYQQKPGKAPKVLIIYFTSSLHSGVP SRFSGSGSGTDFTLTISSLQPEDFATYYCQQYSTV PWTFGQGTKVEIKR

5 Example 3.6: Generation of SOST and TNF (seq. 1) DVD-Igs with Linker Set 1

Table 31

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
94	DVD278H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNOGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSSASTKGPEVQLVESGGGL VQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLE WVSAITWNSGHI DYADSVEGRFTISRDNAKNSLYL QMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGT L VTVSS
95	DVD278L	AB050VL	AB017VL	DLQMTQTSSLSASLGDRVITISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKRTVAAPDIQMTQSPSSLSASVGDRVITITCRASQ GIRNYLAWYQQKPGKAPKLLIYAASTLQSGVPSRF SGSGSGTDFTLTISSLQPEDVATYYCQRYNRAPYT FGQGTKVEIKR

Example 3.7: Generation of SOST and TNF DVD-Igs with Linker Set 2

Table 32

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
96	DVD699H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS ASTKGPE VQLVESGGGL VQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLE WVSAITWNSGHIDYADSVEGRFTISRDNAKNSLYL QMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGT LTVSS
97	DVD699L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGKLE IKR TVDDDDKAAP DIQMTQSPSSLSASVGDRVTIT CRASQGIRNYLAWYQQKPGKAPKLLIYAAS TLQSGVPSRFSGSGSGTDFTLTIS SLQPEDVATYYCQRYN RAPHYTFGGQGTKVEIKR

5 Example 23.8: Generation of SOST and TNF DVD-Igs with Linker Set 3

Table 33

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
98	DVD700H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS ASTKGPE VQLVESGGGL VQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLE WVSAITWNSGHIDYADSVEGRFTISRDNAKNSLYL QMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGT LTVSS
99	DVD700L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGKLE IKR LVPRGSAAP DIQMTQSPSSLSASVGDRVTITC RASQGIRNYLAWYQQKPGKAPKLLIYAAS TLQSGVPSRFSGSGSGTDFTLTIS SLQPEDVATYYCQRYN RAPHYTFGGQGTKVEIKR

Example 3.9: Generation of SOST and TNF DVD-Igs with Linker Set 4

Table 34

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
100	DVD701H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS ASTKGPSVFPLAPE EVQL VESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQ APGKGLEWVSAITWNSGHIDYADSVEGRFTISRDN AKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSLD YWGQGTTLTVSS
101	DVD701L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKR TVAADDDDKSVFIVPP DIQMTQSPSSLSASVG DRVTITCRASQGIRNYLAWYQQKPGKAPKLLIYAA STLQSGVPSRFSGSGSGTDFTLTISSLQPEDVATY YCQRYNRAPYTFGQGTKVEIKR

5 Example 3.10: Generation of SOST and TNF DVD-Igs with Linker Set 5

Table 35

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
102	DVD702H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS GGGGGGG PEVQLVESGG GLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKG LEWVSAITWNSGHIDYADSVEGRFTISRDN AKNSL YLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQG TLVTVSS
103	DVD702L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKR GGGGGGP DIQMTQSPSSLSASVGDRVTITCRA SQGIRNYLAWYQQKPGKAPKLLIYA ASTLQSGVPS RFSGSGSGTDFTLTISSLQPEDVATYYCQRYNRAP YTFGQGTKVEIKR

Example 3.11: Generation of SOST and TNF DVD-Igs with Linker Set 6

Table 36

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
104	DVD703H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSSGGGGGGGGPEVQLVESG GGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGK GLEWVSAITWNSGHIDYADSVEGRFTISRDNAKNS LYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQ GTLVTVSS
105	DVD703L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKRGGGGGGGPDIQMTQSPSSLSASVGDRVTITCR ASQGIRNYLAWYQQKPGKAPKLLIYAASTLQSGVP SRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNRA PYTFGQGTKVEIKR

5 Example 3.12: Generation of SOST and TNF DVD-Igs with Linker Set 7

Table 37

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
106	DVD704H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSSPAPNLLGGPEVQLVESG GGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGK GLEWVSAITWNSGHIDYADSVEGRFTISRDNAKNS LYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQ GTLVTVSS
107	DVD704L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKRPAPNLLGGPDIQMTQSPSSLSASVGDRVTITC RASQGIRNYLAWYQQKPGKAPKLLIYAASTLQSGV PSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNR APYTFGQGTKVEIKR

Example 3.13: Generation of SOST and TNF DVD-Igs with Linker Set 8

Table 38

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
108	DVD705H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS PAPNLLGGPE EVQLVESG GGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGK GLEWVSAITWNSGHIDYADSVEGRFTISRDNAKNS LYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQ GTLVTVSS
109	DVD705L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKR PAPELLGGP DIQMTQSPSSLSASVGDRVTITC RASQGIRNYLAWYQQKPGKAPKLLIYAASTLQSGV PSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNR APYTFGQGTKVEIKR

5 Example 3.14: Generation of SOST and TNF DVD-Igs with Linker Set 9

Table 39

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
110	DVD706H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS PNLLGGPE EVQLVESGGG LVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGL EWVSAITWNSGHIDYADSVEGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGT LVTVSS
111	DVD706L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKR PAPNLLGGP DIQMTQSPSSLSASVGDRVTITC RASQGIRNYLAWYQQKPGKAPKLLIYAASTLQSGV PSRFSGSGSGTDFTLTISSLQPEDVATYYCQRYNR APYTFGQGTKVEIKR

Example 3.15: Generation of SOST and TNF DVD-Igs with Linker Set 10

Table 40

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
112	DVD707H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSS PAPNLLGGPE VQLVESG GGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGK GLEWVSAITWNSGHIDYADSVEGRFTISRDNAKNS LYLQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQ GTLVTVSS
113	DVD707L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKR PTISPAPNLLGGP DIQMTQSPSSLSASVGDRV TITCRASQGIRNYLAWYQQKPGKAPKLLIYAASTL QSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQ RYNRAPYTFGQGTKVEIKR

5 Example 3.16: Generation of SOST and TNF DVD-Igs with Linker Set 11

Table 41

SEQ ID NO	DVD Variable Domain Name	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
114	DVD708H	AB050VH	AB017VH	EVQLQQSGPELMKPGASVKMSCKASGYTFTDYNMH WMKQNQGKSLEWIGEINPNSGGSGYNQKFKGKATL TVDKSSSTAYMELRSLTSEDSAVYYCARLGYYGNY EDWYFDVWGAGTTVTVSSPNLLGGPEVQLVESGGG LVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGL EWVSAITWNSGHIDYADSVEGRFTISRDNAKNSLY LQMNSLRAEDTAVYYCAKVSYLSTASSLDYWGQGT LVTVSS
115	DVD708L	AB050VL	AB017VL	DLQMTQTTSSLSASLGDRVTISCRASQDISNYLNW YQQKPDGTVKLLIFYTSTLQSGVPSRFSGSGSGTN YSLTITNLEQDDAATYFCQQGDTLPYTFGGGTKLE IKR PTISPAPNLLGGP DIQMTQSPSSLSASVGDRV TITCRASQGIRNYLAWYQQKPGKAPKLLIYAASTL QSGVPSRFSGSGSGTDFTLTISSLQPEDVATYYCQ RYNRAPYTFGQGTKVEIKR

Example 3.17: Generation of TNF (seq. 3 – D2E7.1) and IL-13 (seq. 1 – 13C5.5) DVD-Igs Heavy and Light Chains

Table 42

5

SEQ ID NO	Descriptor	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				123456789012345678901234567890123456
116	D2E7.1-13C5.5 (HC0) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSSSEVTLRESGPGLVKPTQTLTLTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEWLAHIWWDDVK RYNPALKSRLTISKDTSKNQVVLKLTSVDPVDTATY YCARTVSSGYIYYAMDYWGQGTTLTVSS
117	D2E7.1-13C5.5 (HC2) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSS AS EVTLRESGPGLVKPTQTLTLTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEWLAHIWWDD VKRYNPALKSRLTISKDTSKNQVVLKLTSVDPVDTA TYYCARTVSSGYIYYAMDYWGQGTTLTVSS
118	D2E7.1-13C5.5 (HC4) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSS ASTK EVTLRESGPGLVKPTQTLT LTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEWLAHIWW DDVKRYNPALKSRLTISKDTSKNQVVLKLTSVDPVD TATYYCARTVSSGYIYYAMDYWGQGTTLTVSS
119	D2E7.1-13C5.5 (HC6) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSS ASTKGPE VTLRESGPGLVKPTQT LTLTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEWLAHI WWDDVKRYNPALKSRLTISKDTSKNQVVLKLTSDVP VDTATYYCARTVSSGYIYYAMDYWGQGTTLTVSS
120	D2E7.1-13C5.5 (HC8) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSS ASTKGPSV EVTLRESGPGLVKPT QTLTLTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEWLA HIWWDDVKRYNPALKSRLTISKDTSKNQVVLKLT SVDPVDTATYYCARTVSSGYIYYAMDYWGQGTTLTVSS
121	D2E7.1-13C5.5 (HC10) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSS ASTKGPSVF EVTLRESGPGLVK PTQTLTLTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEW LAHIWWDDVKRYNPALKSRLTISKDTSKNQVVLKLT SVDPVDTATYYCARTVSSGYIYYAMDYWGQGTTLTV SS
122	D2E7.1-13C5.5 (HC13) VH	AB230VH	AB231VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVAYLSTASSL DYWGQGTLVTVSS ASTKGPSVFPLA EVTLRESGPG LVKPTQTLTLTCTLYGFSLSLSTSDMGVDWIRQPPGKG LEWLAHIWWDDVKRYNPALKSRLTISKDTSKNQVVL KLTSDVPVDTATYYCARTVSSGYIYYAMDYWGQGT LTVSS
123	D2E7.1-13C5.5 (LC0) VL	AB230VL	AB231VL	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASLTQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGTKVEIKR DIQMTQSPSSLSASVGDRVTISCRASQDIRNYLNWY QQKPGKAPKLLIFYTSKLHSGVPSRFSGSGSGTDYT LTISSLQPEDIATYYCQQGNTLPLTFGGGTKVEIKR

SEQ ID NO	Descriptor	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				123456789012345678901234567890123456
124	D2E7.1-13C5.5 (LC3) VL	AB230VL	AB231VL	DIQMTQSPSSLSASVGDRVTTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGGTKVEIKR TVADIQMTQSPSSLSASVGDRV TISCRASQDIRNYL NWYQQKPGKAPKLLIFYTSKLHSGVPSRFSGSGSGT DYTLTISSSLQPEDIATYYCQQGNTLPLTFGGGTKVE IKR
125	D2E7.1-13C5.5 (LC5) VL	AB230VL	AB231VL	DIQMTQSPSSLSASVGDRVTTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGGTKVEIKR TVAAPDIQMTQSPSSLSASVGDRV TISCRASQDIRN YLNWYQQKPGKAPKLLIFYTSKLHSGVPSRFSGSGS GTDYTLTISSSLQPEDIATYYCQQGNTLPLTFGGGTK VEIKR
126	D2E7.1-13C5.5 (LC7) VL	AB230VL	AB231VL	DIQMTQSPSSLSASVGDRVTTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGGTKVEIKR TVAAPSVDIQMTQSPSSLSASVGDRV TISCRASQDI RNYLNWYQQKPGKAPKLLIFYTSKLHSGVPSRFSGS GSGTDYTLTISSSLQPEDIATYYCQQGNTLPLTFGGG TKVEIKR
127	D2E7.1-13C5.5 (LC9) VL	AB230VL	AB231VL	DIQMTQSPSSLSASVGDRVTTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGGTKVEIKR TVAAPSVFI DIQMTQSPSSLSASVGDRVTTISCRASQ DIRNYLNWYQQKPGKAPKLLIFYTSKLHSGVPSRFS GSGSGTDYTLTISSSLQPEDIATYYCQQGNTLPLTFG GGTKVEIKR
128	D2E7.1-13C5.5 (LC12) VL	AB230VL	AB231VL	DIQMTQSPSSLSASVGDRVTTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGGTKVEIKR TVAAPSVFIFPPDIQMTQSPSSLSASVGDRV TISCR ASQDIRNYLNWYQQKPGKAPKLLIFYTSKLHSGVPS RFSGSGSGTDYTLTISSSLQPEDIATYYCQQGNTLPL TFGGGTKVEIKR

Example 3.18: Generation of TNF (seq. 2 – D2E7) and IL-13 (seq. 2 – 13C5.5L3F) DVD-Ig Heavy and Light Chains

Table 43

5

SEQ ID NO	Descriptor	Outer Variable Domain Name	Inner Variable Domain Name	Sequence
				12345678901234567890123456789012345
129	D2E7-13C5.5L3F (HC6) VH	AB229VH	AB232VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSL DYWGQGTLLVTVSS ASTKGPE VTTLRESGPGLVKPTQT LTLTCTLYGFSLSLSTSDMGVDWIRQPPGKGLEWLAHI WWDDVKRYNPALKSRLTISKDTSKNQVVLKLTSVDP VDTATYYCARTVSSGYIYYAMDYWGQGTLLTVSS
130	D2E7-13C5.5L3F (HC13) VH	AB229VH	AB232VH	EVQLVESGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSAITWNSGHIDYADSVEGRFTISR DNAKNSLYLQMNSLRAEDTAVYYCAKVSYLSTASSL DYWGQGTLLVTVSS ASTKGPSVFPLAP EVTLRESGPG LVKPTQTLTLTCTLYGFSLSLSTSDMGVDWIRQPPGKG LEWLAHIWWDDVKRYNPALKSRLTISKDTSKNQVVL KLTSVDPVDTATYYCARTVSSGYIYYAMDYWGQGTLL VTVSS
131	D2E7-13C5.5L3F (LC5) VL	AB229VL	AB232VL	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCQRYNRAPYTFGQGTKVEIKR TVAAP DIQMTQSPSSLSASVGDRVTISCRASQDIRN YLNWYQQKPGKAPKLLIFYTSKLHSGVPSRFSGSGS GTDYTLTISSLQPEDATYYCQQGLTPPLTFGGGTK VEIKR
132	D2E7-13C5.5L3F (LC12) VL	AB229VL	AB232VL	DIQMTQSPSSLSASVGDRVTITCRASQGIRNYLAWY QQKPGKAPKLLIYAASTLQSGVPSRFSGSGSGTDFT LTISSLQPEDVATYYCARYNRAPYTFGQGTKVEIKR TVAAPSVFIFPP DIQMTQSPSSLSASVGDRVTISCR ASQDIRNYLNWYQQKPGKAPKLLIFYTSKLHSGVPS RFSGSGSGTDYTLTISSLQPEDATYYCQQGLTPPL TFGGTKVEIKR

Example 3.19: Cloning Vector Sequences Used to Clone Parent Antibody and DVD-Ig Sequences

10

Table 44

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901
133	V1	GCGTCGACCAAGGGCCCATCGGTCTTCCCCCTGGCACCTCCTCCAAGAGC ACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCC GAACCGGTGACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCAC ACCTTCCCGGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTG GTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTG AATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCCAAATCT TGTGACAAAACCTCACACATGCCCACCGTGCCCAGCACCTGAACCTCCTGGGG GGACCGTCAGTCTTCTCTTCCCCCCAAAACCCAAGGACACCCTCATGATC TCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGAC CCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCC AAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGC GTCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC AAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAA GCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGC GAGGAGATGACCAAGAACCAGGTACGCTGACCTGCCTGGTCAAAGGCTTC TATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901
		AACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTC TACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC TCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGC CTCTCCCTGTCTCCGGGTAAATGAGCGGCCGCTCGAGGCCGGAAGGCCGG ATCCCCGACCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAA TAGTGTGTTGGAATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGG CAAATCATTTGGTCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCC CCGCCCCGGACGAACATAACCTGACTACGACATCTCTGCCCCTTCTTCGCG GGGCAGTGCATGTAATCCCTTCAGTTGGTTGGTACAACCTTGCCAACCTGGGC CCTGTTCCACATGTGACACGGGGGGGGACCAAACACAAAGGGGTTCTCTGA CTGTAGTTGACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTG GCTTTCATCCTGGAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTG CCTTTATGTGTAACCTTTGGCTGAAGCTCTTACACCAATGCTGGGGGACAT GTACCTCCCAGGGGCCAGGAAGACTACGGGAGGCTACACCAACGTCAATC AGAGGGGCCTGTGTAGCTACCGATAAGCGGACCCTCAAGAGGGGCATTAGCA ATAGTGTTTATAAGGCCCCCTTGTTAACCTAAACGGGTAGCATATGCTTC CCGGGTAGTAGTATATACTATCCAGACTAACCCTAATTCAATAGCATATGT TACCCAACGGGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTA AGGAACAGCGATATCTCCCACCCCATGAGCTGTCACGGTTTTATTTACATG GGGTCAGGATTCCACGAGGGTAGTGAACCATTTTAGTCACAAGGGCAGTGG CTGAAGATCAAGGAGCGGGCAGTGAACCTCTCCTGAATCTTCGCCTGCTTCT TCATTCTCCTTCGTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAA GGTGTATGTGAGGTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAGAATA AAATTTGGACGGGGGGTTCAGTGGTGGCATTGTGCTATGACACCAATATAA CCCTCACAAACCCCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTCT GAATATCTTTAACAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACT GGATGTCCATCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGT GCAATATGATACTGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACA GGTGAACCATGTTGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGAC GCCGACAGCAGCGGACTCCACTGGTTGTCTCTAACACCCCCGAAAATTAAA CGGGGCTCCACGCCAATGGGGCCCATAAACAAAGACAAGTGGCCACTCTTT TTTTTGAAATTGTGGAGTGGGGGCACGCGTCAGCCCCCACACGCCGCCCTG CGGTTTTGGACTGTAAAATAAGGGTGTAATAACTTGGCTGATTGTAACCCC GCTAACCACTGCGGTCAAACCACTTGCCCACAAAACCACTAATGGCACCCC GGGAATACCTGCATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTGC TGCGATCTGGAGGACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAG GGTTGTTGGTCCTCATATTCACGAGGTGCTGAGAGCACGGTGGGCTAATG TTGCCATGGGTAGCATATACTACCCAAATATCTGGATAGCATATGCTATCC TAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCAT ATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATTTTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAA TCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATG CTATCCTAATAGAGATTAGGGTAGTATATGCTATCCTAATTTTATATCTGGG TAGCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCT AATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATA TGCTATCCTAATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT CTGTATCCGGGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGA ATTTTCTTGAAGACGAAAGGGCCTCGTGATACGCCTATTTTATAGGTTAA TGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTTCGGGGAAA TGTGCGCGGAACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTA TCCGCTCATGAGACAATAACCTGATAAATGCTTCAATAATATTGAAAAAG GAAGAGTATGAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGC GGCATTTTGCCTTCCTGTTTTTGTCTACCCAGAAACGCTGGTGAAAGTAAA AGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCT CAACAGCGGTAAGATCCTTGAGAGTTTTTCGCCCCGAAGAACGTTTTTCCAAT GATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTTGA CGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTT

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		GGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGT AAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAAACTGCGGCCAA CTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGGCA CAACATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAA TGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGC AACAACGTTGCGCAAACCTATTAAC TGGCGAACTACTTACTCTAGCTTCCCG GCAACAATTAATAGACTGGATGGAGGCGGATAAAAGTTGCAGGACCACTTCT GCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCGG TGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCC CTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGA ACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAAGCATTTGGTA ACTGTCAGACCAAGTTTACTCATATATACTTTTAGATTGATTTAAAACTTCA TTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGAC CAAAATCCCTTAACGTGAGTTTTCGTTCCACTGAGCGTCAGACCCCGTAGA AAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTG CTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCA AGAGCTACCAACTCTTTTTCCGAAGGTAAC TGGCTTCAGCAGAGCGCAGAT ACCAAATACTGTTCTTCTAGTG TAGCCGTAGTTAGGCCACCACTTCAAGAA CTCTGTAGCACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGC TGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATA GTTACCGGATAAGGCGCAGCGGTTCGGGCTGAACGGGGGGTTTCGTGCACACA GCCCAGCTTGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGA GCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCC GGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGG AAACGCCTGGTATCTTTATAGTCCTGTCTGGGTTTCGCCACCTCTGACTTGA GCGTCGATTTTTTG TGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGC CAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCA CATGTTCTTTCCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGC CTTTGAGTGAGCTGATAACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGA GTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCC CGCGCGTTGGCCGATTCAATTAATGCAGCTGGCACGACAGGTTTCCCGACTG GAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTA GGCACCCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAAT TGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACGC CAAGCTCTAGCTAGAGGTGAGTCCCTCCCCAGCAGGCAGAAGTATGCAAA GCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTA ACTCCGCCCCA TCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGAC TAATTTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTAT TCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGC TTTGCAAAGATGGATAAAAGTTTTTAAACAGAGAGGAATCTTTGCAGCTAATG GACCTTCTAGGTCTTGAAAGGAGTGGAATTGGCTCCGGTGCCCGTCAGTG GGCAGAGCGCACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTCTGG CAATTGAACCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGA TGTCGTGTACTGGCTCCGCCTTTTTTCCCGAGGGTGGGGGAGAACCGTATAT AAGTGCAGTAGTCGCCGTGAACGTTCTTTTTCGCAACGGGTTTGCCGCCAG AACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGG TTATGGCCCTTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGAT TCTTGATCCCGAGCTTCGGGTTGGAAGTGGGTGGGAGAGTTTCGAGGCCTTG CGCTTAAGGAGCCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCG CTGGGGCCGCGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGC TTTCGATAAGTCTCTAGCCATTTAAAATTTTTTGATGACCTGCTGCGACGCT TTTTTTCTGGCAAGATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGG TATTTTCGGTTTTTTGGGGCCGCGGGCGGCGACGGGGCCCGTGCGTCCCAGCG CACATGTTTCGGCGAGGCGGGGCCTGCGAGCGCGGCCACCGAGAATCGGACG GGGGTAGTCTCAAGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCC GTGTATCGCCCCGCCCTGGGCGGCAAGGCTGGCCCGGTCGGCACCAGTTGC GTGAGCGGAAAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATG GAGGACGCGGCGCTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAAGGAA AAGGGCCTTTCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCG

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		GGCGCCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTTCGTC TTTAGGTTGGGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTG GGTGGAGACTGAAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGA ATTTGCCCTTTTTGAGTTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGT GGTTCAAAGTTTTTTTCTTCCATTTTCAGGTGTCGTGAGGAATTCTCTAGAG ATCCCTCGACCTCGAGATCCATTGTGCCCCGGGCGCCACCATGGAGTTTGGG CTGAGCTGGCTTTTTTCTTGTGCGGATTTTAAAAGGTGTCCAGTGC
134	V2	ACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTG AAATCTGGAACCTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGA GAGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGC AGCACCTTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCC TGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAAC AGGGGAGAGTGTTGAGCGGCCGCTCGAGGCCGGCAAGGCCGGATCCCCCGA CCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTT GGAATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATT TGGTCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCCGG ACGAACAAACCTGACTACGACATCTCTGCCCTTCTTCGCGGGGCAGTGC ATGTAATCCCTTCAGTTGGTTGGTACAACCTGCCAACTGGGCCCTGTTCCA CATGTGACACGGGGGGGGACCAAACACAAAGGGGTCTCTGACTGTAGTTG ACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTCATC CTGGAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTGCCTTTATGT GTAACCTTTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCC AGGGGCCCAGGAAGACTACGGGAGGCTACACCAACGTCAATCAGAGGGGCC TGTGTAGCTACCGATAAGCGGACCCTCAAGAGGGCATTAGCAATAGTGTTT ATAAGGCCCCCTTGTTAACCTAAACGGGTAGCATATGCTTCCCGGGTAGT AGTATATACTATCCAGACTAACCCTAATTCAATAGCATATGTTACCCAACG GGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTAAGGAACAGC GATATCTCCCACCCCATGAGCTGTCACGGTTTTATTTACATGGGGTCAGGA TTCCACGAGGGTAGTGAACCATTTTAGTGCACAAGGGCAGTGGCTGAAGATC AAGGAGCGGGCAGTGAACCTCCTGAATCTTCGCCTGCTTCTTCATTCTCC TTCGTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGT GAGGTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAGAATAAAATTTGGA CGGGGGGTTTCAGTGGTGGCATTGTGCTATGACACCAATATAACCTCACAA ACCCCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTCTGAATATCTT TAACAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACTGGATGTCCA TCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGTGCAATATGA TACTGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACAGGTGAACCA TGTTGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACGCCGACAGC AGCGGACTCCACTGGTTGTCTCTAACACCCCCGAAAATTAAACGGGGCTCC ACGCCAATGGGGCCCATAAACAAAGACAAGTGGCCACTCTTTTTTTTGAAA TTGTGGAGTGGGGGCACGCGTCAGCCCCCACACGCCGCCCTGCGGTTTTGG ACTGTAAAATAAGGGTGTAATAACTTGGCTGATTGTAACCCCGCTAACCAC TGCGGTCAAACCACTTGCCCACAAAACCACTAATGGCACCCCGGGGAATAC CTGCATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTGCTGCGATCTG GAGGACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAGGGTTGTTGG TCCTCATATTACGAGGTGCTGAGAGCACGGTGGGCTAATGTTGCCATGG GTAGCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATA TCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCC TAATCTATATCTGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCAT AGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCT GGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATGCTATCCTAA TAGAGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATA CTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGC ATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCT AATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCG GGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAATTTTCTTG

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901
		AAGACGAAAGGGCCTCGTGATACGCCTATTTTTATAGGTTAATGTCATGAT AATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGG AACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCAT GAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTAT GAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTG CCTTCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGA AGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGG TAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCAC TTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTTGACGCCGGGCA AGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTTGAGTA CTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATT ATGCAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCT GACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGACAACATGGG GGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCAT ACCAAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTT GCGCAAAC TATTA ACTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATT AATAGACTGGATGGAGGCGGATAAAAGTTGCAGGACCACTTCTGCGCTCGGC CCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGG GTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTAT CGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAG ACAGATCGCTGAGATAGGTGCCTCACTGATTAAAGCATTGGTAACTGTCAGA CCAAGTTTACTCATATATACTTTAGATTGATTTAAAAC TTCATTTTTAATT TAAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCC TTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAA AGGATCTTCTTGAGATCCTTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAC AAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCAAGAGCTACC AACTCTTTTTCCGAAGGTA ACTGGCTTCAGCAGAGCGCAGATACCAAATAC TGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGC ACCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAG TGCGGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGA TAAGGCGCAGCGGTGCGGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTT GGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGA AAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGG CAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTG GTATCTTTATAGTCCTGTGCGGTTTTCGCCACCTCTGACTTGAGCGTCGATT TTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGC GGCCTTTTTACGGTTCCCTGGCCTTTTGCTGGCCTTTTGCTCACATGTTCTT TCCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTG AGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTGAG CGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTTG GCCGATTCATTAATGCAGCTGGCACGACAGGTTTCCCGACTGGAAAGCGGG CAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCCA GGCTTTACACTTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGG ATAACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCTA GCTAGAGGTCGAGTCCCTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATC TCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCCCC TAACTCCGCCCAGTTCGCCCCATTCTCCGCCCCATGGCTGACTAATTTTTT TTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATTCCAGAAGT AGTGAGGAGGCTTTTTTGAGGCCTAGGCTTTTGCAAAAAGCTTTGCAAAG ATGGATAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATGGACCTTCTA GGTCTTGAAAGGAGTGGAATTGGCTCCGGTGCCCGTCAGTGGGCAGAGCG CACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTGCGCAATTGAAC CGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTA CTGGCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGT AGTCGCCGTGAACGTTCTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGT AAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCC TTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGATTCTTGATCC CGAGCTTCGGGTGGAAGTGGGTGGGAGAGTTCGAGGCCTTGCGCTTAAGG AGCCCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCG

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901
		CCGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTTCGATAA GTCTCTAGCCATTTTAAATTTTTGATGACCTGCTGCGACGCTTTTTTTCTG GCAAGATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGGTATTTTCGGT TTTTGGGGCCGCGGGCGGCGACGGGGCCCCGTGCGTCCCAGCGCACATGTTT GGCGAGGCGGGGCCCTGCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTC TCAAGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCCGTGTATCGC CCCGCCCTGGGCGGCAAGGCTGGCCCCGGTCGGCACCAAGTTGCGTGAGCGGA AAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCG GCGCTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGGCCTT TCCGTCCTCAGCCGTGCTTTCATGTGACTCCACGGAGTACCGGGCGCCGTC CAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCGTCTTTAGGTTG GGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGAC TGAAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGAATTTGCCCT TTTTGAGTTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAG TTTTTTTCTTCCATTTCAGGTGTGCTGAGGAATTCTCTAGAGATCCCTCGA CCTCGAGATCCATTGTGCCCCGGGCGCACCATGGACATGCGCGTGCCCGCCC AGCTGCTGGGCCTGCTGCTGCTGTGGTTCCCCGGCTCGCGATGC
135	V3	CAACCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAG CTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCG GGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGA GTGGAGACCACCACACCCCTCCAAACAAAGCAACAACAAGTACGCGGCCAGC AGCTACCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGC TGCCAGGTCACGCATGAAGGGAGCACCGTGGAAGACAGTGGCCCCCTACA GAATGTTTCATGAGCGGCCGCTCGAGGCCGGCAAGGCCGGATCCCCCGACCT CGACCTCTGGCTAATAAAGGAAATTTATTTTTCATTGCAATAGTGTGTTGGA ATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTTGG TCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCCGGACG AACTAAACCTGACTACGACATCTCTGCCCCCTTCTTCGCGGGGGCAGTGCATG TAATCCCTTCAGTTGGTTGGTACAACCTTGCCAACCTGGGCCCTGTTCCACAT GTGACACGGGGGGGGACCAAACACAAAGGGGTTCTCTGACTGTAGTTGACA TCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTCATCCTG GAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTGCCTTTATGTGTA ACTCTTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCCAGG GGCCCAGGAAGACTACGGGAGGCTACACCAACGTCAATCAGAGGGGCCTGT GTAGCTACCGATAAGCGGACCCCTCAAGAGGGCATTAGCAATAGTGTTTATA AGGCCCCCTTGTTAACCCTAAACGGGTAGCATATGCTTCCCGGGTAGTAGT ATATACTATCCAGACTAACCCTAATTCAATAGCATATGTTACCCAACGGGA AGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTAAGGAACAGCGAT ATCTCCCACCCCATGAGCTGTCACGGTTTTTATTTACATGGGGTCAGGATTC CACGAGGGTAGTGAACCATTTTAGTCACAAGGGCAGTGGCTGAAGATCAAG GAGCGGGCAGTGAACCTCTCCTGAATCTTCGCCTGCTTCTTCATTCTCCTTC GTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGTGAG GTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAGAATAAAATTTGGACGG GGGGTTCAGTGGTGGCATTTGTGCTATGACACCAATATAACCCTCACAAACC CCTTGGGCAATAAAATACTAGTGTAGGAATGAAACATTCTGAATATCTTTAA CAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACTGGATGTCCATCT CACACGAATTTATGGCTATGGGCAACACATAATCCTAGTGCAATATGATAC TGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACAGGTGAACCATGT TGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACGCCGACAGCAGC GGACTCCACTGGTTGTCTCTAACACCCCCGAAAATTAAACGGGGCTCCACG CCAATGGGGCCCATAAACAAAGACAAGTGGCCACTCTTTTTTTTTTGAAATTG TGGAGTGGGGGCACGCGTCAGCCCCCACACGCCGCCCTGCGGTTTTTGGACT GTAAAATAAGGGTGTAATAACTTGGCTGATTGTAACCCCGCTAACCCTGC GGTCAAACCACTTGCCCAAAAACCACTAATGGCACCCCGGGGAATACCTG CATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTGCTGCGATCTGGAG GACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAGGGTTGTTGGTCC TCATATTCACGAGGTGCTGAGAGCACGGTGGGCTAATGTTGCCATGGGTA GCATATACTACCAAATATCTGGATAGCATATGCTATCCTAATCTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAA

206

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901
		GATAAAGTTTTTAAACAGAGAGGAATCTTTGCAGCTAATGGACCTTCTAGGTCTTGAAAGGAGTGGGAATTGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTTCGGCAATTGAACCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAACGTTCTTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCCTTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGATTCTTGATCCCGAGCTTCGGGTGGAAGTGGGTGGGAGAGTTCGAGGCCTTGCGCTTAAGGAGCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCGCCGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAAAATTTTTGATGACCTGCTGCGACGCTTTTTTTTCTGGCAAGATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGGTATTTTCGGTTTTTGGGGCCGCGGGCGGCGACGGGGCCCCGTGCGTCCCAGCGCACATGTTCCGGCAGGGCGGGCCTGCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTCTCAGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCCGTGTATCGCCCCGCCCTGGGCGGCAAGGCTGGCCCCGGTCGGCACCAGTTGCGTGAGCGGAAAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGCGCTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGGCCTTTCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCGGGCGCCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTGCTCTTTAGGTGGGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTGAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGAATTTGCCCTTTTGAGTTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAGTTT TTTTCTTCCATTTCAGGTGTCGTGAGGAATTCTCTAGAGATCCCTCGACCTCGAGATCCATTGTGCCCGGGCGCCACCATGACTTGGACCCCACTCCTCTTCCTCACCTCCTCCTCCACTGCACAGGAAGCTTATCG
136	V4	ACGGTGGCTGCACCATCTGTCTTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGCCAAAGTACAGTGGAAAGGTGGATAACGCCCTCCAATCGGGTAACTCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTGAGCGGCCGCTCGAGGGCCGGCAAGGCCGGATCCCCCGACCTCGACCTCTGGCTAATAAAGGAAATTTATTTTTCATTGCAATAGTGTGTTGGAATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTGGTCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCCGGACGAATAAACCTGACTACGACATCTCTGCCCCCTTCTTCGCGGGGCAGTGCATGTAATCCCTTCAGTTGGTTGGTACAACCTTGCCAACTGGGCCCTGTTCCACATGTGACACGGGGGGGGACCAAACACAAAGGGGTTCCTTGACTGTAGTTGACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTTCATCTGGAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTGCCTTTATGTGTAACCTCTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCCAGGGGCCAGGAAGACTACGGGAGGCTACACCAACGTCAATCAGAGGGGCCGTGTAGCTACCGATAAGCGGACCTCAAGAGGGCATTAGCAATAGTGTTTATAAGGCCCCCTTGTTAACCCCTAAACGGGTAGCATATGCTTCCCGGGTAGTAGTATATACTATCCAGACTAACCCCTAATTCAATAGCATATGTTACCCAACGGGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTAAGGAACAGCGATATCTCCCACCCCATGAGCTGTCACGGTTTTATTTACATGGGGTCAGGATTCCACGAGGGTAGTGAACCATTTTAGTCAACAAGGGCAGTGGCTGAAGATCAAGGAGCGGGCAGTGAACCTCTCCTGAATCTTCGCCTGCTTCTTCATTCTCCTTCGTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGTGAGGTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAGAATAAAATTTGGA CGGGGGGTTTCAGTGGTGGCATTGTGCTATGACACCAATATAACCCTCACAAACCCCTGGGCAATAAATACTAGTGTAGGAATGAAACATTCTGAATATCTTTAACAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACTGGATGTCCATCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGTGCAATATGATACTGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACAGGTGAACCATGTTGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACGCCGACAGC

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		AGCGGACTCCACTGGTTGTCTCTAACACCCCCGAAAATTAAACGGGGCTCC ACGCCAATGGGGCCCATAAACAAAGACAAGTGGCCACTCTTTTTTTTGAAA TTGTGGAGTGGGGGCACGCGTCAGCCCCCACACGCCGCCCTGCGGTTTTGG ACTGTAAAATAAGGGTGTAATAACTTGGCTGATTGTAACCCCGCTAACCAC TGCGGTCAAACCACTTGCCCACAAAACCACTAATGGCACCCCGGGGAATAC CTGCATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTGCTGCGATCTG GAGGACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAGGGTTGTTGG TCCTCATATTCACGAGGTCGCTGAGAGCACGGTGGGCTAATGTTGCCATGG GTAGCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATA TCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCC TAATCTATATCTGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCAT AGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCT GGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATGCTATCCTAA TAGAGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATA CTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGC ATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCT AATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCG GGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAATTTTCTTG AAGACGAAAGGGCCTCGTGATACGCCTATTTTTTATAGGTTAATGTCATGAT AATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGG AACCCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCAT GAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTAT GAGTATTCAACATTTCCGTGTCGCCCTTATTCCCTTTTTTGCGGCATTTTG CCTTCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGA AGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGG TAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTCCAATGATGAGCAC TTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTTGACGCCGGGCA AGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTTGGTGAGTA CTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATT ATGCAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCT GACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCAACAATGGG GGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCAT ACCAAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTT GCGCAAACTATTAAGTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATT AATAGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGC CCTTCCGGCTGGCTGGTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGG GTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTAT CGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAG ACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAAGTGTGAGA CCAAGTTTACTCATATATACTTTAGATTGATTTAAAACTTCATTTTTTAATT TAAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAAATCCC TTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAA AGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAAC AAAAAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTACC AACTCTTTTTTCCGAAGGTAAGTGGCTTCAGCAGAGCGCAGATACCAAATAC TGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGC ACCGCTACATAACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAG TGCGGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGA TAAGGCGCAGCGGTCGGGCTGAACGGGGGGTTCGTGCACACAGCCCAGCTT GGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGA AAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGG CAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTG GTATCTTTATAGTCCTGTGCGGTTTTCGCCACCTCTGACTTGAGCGTCGATT TTTGTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGC GGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCACATGTTCTT TCCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTG AGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTGAG

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		CGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTTG GCCGATTCATTAATGCAGCTGGCACGACAGGTTTCCCGACTGGAAAGCGGG CAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCCA GGCTTTACACTTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGG ATAACAATTTCACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCTA GCTAGAGGTCGAGTCCCTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATC TCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCCCC TAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTT TTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATTCCAGAAGT AGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTTTGCAAAG ATGGATAAAGTTTTAAACAGAGAGGAATCTTTGCAGCTAATGGACCTTCTA GGTCTTGAAAGGAGTGGAATTGGCTCCGGTGCCCGTCAGTGGGCAGAGCG CACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTCGGCAATTGAAC CGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCGTGTA CTGGCTCCGCCTTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGT AGTCGCCGTGAACGTTCTTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGT AAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTATGGCCC TTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGATTCTTGATCC CGAGCTTCGGGTGGAAGTGGGTGGGAGAGTTCGAGGCCTTGCCTTAAGG AGCCCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCG CCGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTTCGATAA GTCTCTAGCCATTTAAAATTTTTGATGACCTGCTGCGACGCTTTTTTTCTG GCAAGATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGGTATTTCCGT TTTTTGGGGCCGCGGGCGGCGACGGGGCCCCGTGCGTCCCAGCGCACATGTT GGCGAGGCGGGGCCGTGCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTC TCAAGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCCGTGTATCGC CCCGCCCTGGGCGGCAAGGCTGGCCCGGTGGGCACCAAGTTGCGTGAGCGGA AAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCG GCGCTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGGCCTT TCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCGGGCGCCGTC CAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCGTCTTTAGGTTG GGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGAC TGAAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGAATTTGCCCT TTTTTGAGTTTGGATCTTGGTTCAATCTCAAGCCTCAGACAGTGGTTCAAAG TTTTTTTCTTCCATTTCAAGTGTCGTGAGGAATTCTCTAGAGATCCCTCGA CCTCGAGATCCATTGTGCCCCGGGCGCACCATGACTTGGACCCCACTCCTCT TCCTCACCCCTCCTCCTCCACTGCACAGGAAGCTTATCG
137	V5	CAACCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAG CTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCG GGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGA GTGGAGACCACCACACCCCTCAAACAAGCAACAACAAGTACGCGGCCAGC AGCTACCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGC TGCCAGGTCACGCATGAAGGGAGCACCGTGGAAGAAGACAGTGGCCCCCTACA GAATGTTTCATGAGCGGCCGCTCGAGGCCGGCAAGGCCGGATCCCCCGACCT CGACCTCTGGCTAATAAAGGAAATTTATTTTTCATTGCAATAGTGTGTTGGA ATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTTGG TCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCCCCGCCCGGACG AACTAAACCTGACTACGACATCTCTGCCCTTCTTCGCGGGGCAGTGCATG TAATCCCTTCAGTTGGTTGGTACAACCTTGCCAACCTGGGCCCTGTTCCACAT GTGACACGGGGGGGGACCAAACACAAAGGGGTCTCTGACTGTAGTTGACA TCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTGGCTTTCATCCTG GAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTGCCTTTATGTGTA ACTCTTGGCTGAAGCTCTTACACCAATGCTGGGGGACATGTACCTCCCAGG GGCCCAGGAAGACTACGGGAGGCTACACCAACGTCAATCAGAGGGGCCCTGT GTAGCTACCGATAAGCGGACCTCAAGAGGGCATTAGCAATAGTGTTTATA AGGCCCCCTTGTTAACCCTAAACGGGTAGCATATGCTTCCCGGGTAGTAGT ATATACTATCCAGACTAACCCTAATTCAATAGCATATGTTACCCAACGGGA AGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTAAGGAACAGCGAT ATCTCCCACCCCATGAGCTGTCACGGTTTTTATTTACATGGGGTCAGGATTC

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		CACGAGGGTAGTGAACCATTTTAGTCACAAGGGCAGTGGCTGAAGATCAAG GAGCGGGCAGTGAACCTCTCCTGAATCTTCGCCTGCTTCTTCATTCTCCTTC GTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAAGGTGTATGTGAG GTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAGAATAAAATTTGGACGG GGGGTTCAGTGGTGGCATTTGTGCTATGACACCAATATAACCCTCACAAACC CCTTGGGCAATAAAATACTAGTGTAGGAATGAAACATTCTGAATATCTTTAA CAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACTGGATGTCCATCT CACACGAATTTATGGCTATGGGCAACACATAATCCTAGTGCAATATGATAC TGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACAGGTGAACCATGT TGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGACGCCGACAGCAGC GGACTCCACTGGTTGTCTCTAACACCCCCGAAAATTAAACGGGGCTCCACG CCAATGGGGCCCCATAAACAAAGACAAGTGGCCACTCTTTTTTTTTTGAAATTG TGGAGTGGGGGCACGCGTCAGCCCCCACACGCCGCCCTGCGGTTTTTGGACT GTAAAAAAGGGTGTAAATAACTTGGCTGATTGTAACCCCGCTAACCCTGC GGTCAAACCACTTGCCCAAAAACCACTAATGGCACCCCGGGGAATACCTG CATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTGCTGCGATCTGGAG GACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAGGGTTGTTGGTCC TCATATTCACGAGGTCGCTGAGAGCACGGTGGGCTAATGTTGCCATGGGTA GCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAA TCTATATCTGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATAGG CTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGG TAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATGCTATCCTAATAG AGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGGTAGCATATACTA CCCAAATATCTGGATAGCATATGCTATCCTAATCTATATCTGGGTAGCATA TGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT TTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGC TATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCGGGT AGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGAATTTTCTTGAAG ACGAAAGGGCCTCGTGATACGCCTATTTTTTATAGGTTAATGTCATGATAAT AATGGTTTCTTAGACGTCAGGTGGCACTTTTTCGGGGAAATGTGCGCGGAAC CCCTATTTGTTTATTTTTCTAAATACATTCAAATATGTATCCGCTCATGAG ACAATAACCTGATAAATGCTTCAATAATATTGAAAAAGGAAGAGTATGAG TATTCAACATTTCCGTGTGCGCCCTTATTCCCTTTTTTTGCGGCATTTTGCCT TCCTGTTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAAAGATGCTGAAGA TCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCTCAACAGCGGTAA GATCCTTGAGAGTTTTTCGCCCCGAAGAACGTTTTTCCAATGATGAGCACTTT TAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTTGACGCCGGGCAAGA GCAACTCGGTGCGCGCATACACTATTCTCAGAATGACTTGGTTGAGTACTC ACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATG CAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCAACTTACTTCTGAC AACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTGCACAACATGGGGGA TCATGTAACCTCGCCTTGATCGTTGGGAACCGGAGCTGAATGAAGCCATACC AAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGCAACAACGTTGCG CAAACATTAACCTGGCGAACTACTTACTCTAGCTTCCCGGCAACAATTAAT AGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCTGCGCTCGGCCCT TCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCGGTGAGCGTGGGTC TCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCCCTCCCGTATCGT AGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGAACGAAATAGACA GATCGCTGAGATAGGTGCCTCACTGATTAAGCATTGGTAACCTGTCAGACCA AGTTTACTCATATATACTTTAGATTGATTTAAAACTTCATTTTTTAATTTAA AAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGACCAAATCCCTTA ACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAAGATCAAAGG ATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCTTGCAAACAAA AAAACCACCGCTACCAGCGGTGGTTTGTGTTGCCGGATCAAGAGCTACCAAC TCTTTTTCCGAAGGTAACGGCTTCAGCAGAGCGCAGATACCAAATACTGT TCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAACTCTGTAGCACC GCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGCTGCTGCCAGTGG

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901
		CGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGGATAA GGCGCAGCGGTTCGGGCTGAACGGGGGGTTTCGTGCACACAGCCCAGCTTGGA GCGAACGACCTACACCGAACTGAGATACCTACAGCGTGAGCTATGAGAAAG CGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCCGGTAAGCGGCAG GGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGGAAACGCCTGGTA TCTTTATAGTCCTGTTCGGGTTTCGCCACCTCTGACTTGAGCGTCGATTTTT GTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGCGGC CTTTTTACGGTTCCCTGGCCTTTTGCTGGCCTTTTGCTCACATGTTCTTCC TGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGCCTTTGAGTGAGC TGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGAGTCAGTGAGCGA GGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCCCGCGCGTGGCC GATTCATTAATGCAGCTGGCACGACAGGTTTCCCGACTGGAAAGCGGGCAG TGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTAGGCACCCCAGGC TTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAATTGTGAGCGGATA ACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCTAGCT AGAGGTCGAGTCCCTCCCCAGCAGGCAGAAAGTATGCAAAGCATGCATCTCA ATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCCCCCTAA CTCCGCCCAGTTCCGCCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTTTA TTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGT GAGGAGGCTTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTTTGCAAAGATG GATAAAGTTTTTAAACAGAGAGGAATCTTTGCAGCTAATGGACCTTCTAGGT CTTGAAAGGAGTGGGAATTGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCAC ATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTTCGGCAATTGAACCGG TGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGATGTCTGTACTG GCTCCGCCTTTTTTCCCGAGGGTGGGGGAGAACCGTATATAAGTGCAGTAGT CGCCGTGAACGTTCTTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGTAAG TGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGGTTATGGCCCTTG CGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGATTCTTGATCCCGA GCTTCGGGTTGGAAGTGGGTGGGAGAGTTTCGAGGCCTTGCGCTTAAGGAGC CCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCGCCG CGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTC TCTAGCCATTTAAAATTTTTGATGACCTGCTGCGACGCTTTTTTTTCTGGCA AGATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGGTATTTTCGGTTTTT TGGGGCCGCGGGCGGCGACGGGGCCCCGTGCGTCCCAGCGCACATGTTCCGC GAGGCGGGGCCTGCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTCTCA AGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCGGTGTATCGCCCC GCCCTGGGCGGCAAGGCTGGCCCCGGTCGGCACCAGTTGCGTGAGCGGAAAG ATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGCG CTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGGCCTTTCC GTCCTCAGCCGTCGTTTCATGTGACTCCACGGAGTACCGGGCGCCGTCCAG GCACCTCGATTAGTTCTCGAGCTTTTGAGTACGTCTTTAGGTTGGGG GGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGACTGA AGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGGAATTTGCCCTTTT TGAGTTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAGTTT TTTTCTTCCATTTCAGGTGTCGTGAGGAATTCTCTAGAGATCCCTCGACCT CGAGATCCATTGTGCCCCGGGCGCCACCATGGACATGCGCGTGCCCCGCCAG CTGCTGGGCCCTGCTGCTGTGGTTCCCCGGCTCGCGATGC
138	V7	GCGTCGACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGC ACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCC GAACCGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGAC ACCTTCCCGGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTG GTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAACGTG AATCACAAGCCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCT TGTGACAAAACCTCACACATGCCACCCGTGCCCAGCACCTGAAGCCGCGGGG GGACCGTCAGTCTTCTCTTCCCCCAAACCCAAGGACACCCTCATGATC TCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAGAC CCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCC AAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGC GTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		AAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAA GCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGC GAGGAGATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTC TATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC AACTACAAGACCACGCCTCCCGTGTCTGGACTCCGACGGCTCCTTCTTCCTC TACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC TCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGC CTCTCCCTGTCTCCGGGTAAATGAGCGGCCGCTCGAGGCCGGCAAGGCCGG ATCCCCCGACCTCGACCTCTGGCTAATAAAGGAAATTTATTTTCATTGCAA TAGTGTGTTGGAATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGG CAAATCATTTGGTCGAGATCCCTCGGAGATCTCTAGCTAGAGGATCGATCC CCGCCCCGGACGAATAAACCTGACTACGACATCTCTGCCCCTTCTTCGCG GGGCAGTGCAATTAATCCCTTCAGTTGGTTGGTACAACCTTGCCAACCTGGGC CCTGTTCCACATGTGACACGGGGGGGGACCAAACACAAAGGGGTTCTCTGA CTGTAGTTGACATCCTTATAAATGGATGTGCACATTTGCCAACACTGAGTG GCTTTCATCCTGGAGCAGACTTTGCAGTCTGTGGACTGCAACACAACATTG CCTTTATGTGTAACCTTTGGCTGAAGCTCTTACACCAATGCTGGGGGACAT GTACCTCCCAGGGGCCCAGGAAGACTACGGGAGGCTACACCAACGTCAATC AGAGGGGCCTGTGTAGCTACCGATAAGCGGACCCTCAAGAGGGCATTAGCA ATAGTGTTTATAAGGCCCCCTTGTTAACCCATAAACGGGTAGCATATGCTTC CCGGGTAGTAGTATATACTATCCAGACTAACCCTAATTCAATAGCATATGT TACCCAACGGGAAGCATATGCTATCGAATTAGGGTTAGTAAAAGGGTCCTA AGGAACAGCGATATCTCCCACCCCATGAGCTGTCACGGTTTTTATTTACATG GGGTCAGGATTCACGAGGGTAGTGAACCATTTTAGTCAACAGGGCAGTGG CTGAAGATCAAGGAGCGGGCAGTGAACCTCTCCTGAATCTTCGCCTGCTTCT TCATTCTCCTTCGTTTTAGCTAATAGAATAACTGCTGAGTTGTGAACAGTAA GGTGTATGTGAGGTGCTCGAAAACAAGGTTTCAGGTGACGCCCCCAGAATA AAATTTGGACGGGGGGTTCAGTGGTGGCATTTGTGCTATGACACCAATATAA CCCTCACAAACCCCTTGGGCAATAAATACTAGTGTAGGAATGAAACATTCT GAATATCTTTAACAATAGAAATCCATGGGGTGGGGACAAGCCGTAAAGACT GGATGTCCATCTCACACGAATTTATGGCTATGGGCAACACATAATCCTAGT GCAATATGATACTGGGGTTATTAAGATGTGTCCCAGGCAGGGACCAAGACA GGTGAACCATGTTGTTACACTCTATTTGTAACAAGGGGAAAGAGAGTGGAC GCCGACAGCAGCGGACTCCACTGGTTGTCTCTAACACCCCCGAAAATTAAA CGGGGCTCCACGCCAATGGGGCCCATAAACAAAGACAAGTGGCCACTCTTT TTTTTGAAATTGTGGAGTGGGGGCACGCGTCAGCCCCCACACGCCGCCCTG CGGTTTTGGACTGTAAATAAGGGTGTAATAACTTGGCTGATTGTAACCCC GCTAACCACTGCGGTCAAACCACTTGCCCACAAAACCACTAATGGCACCCC GGGGAATACCTGCATAAGTAGGTGGGCGGGCCAAGATAGGGGCGCGATTGC TGCGATCTGGAGGACAAATTACACACACTTGCGCCTGAGCGCCAAGCACAG GGTTGTTGGTCCTCATATTCACGAGGTCGCTGAGAGCACGGTGGGCTAATG TTGCCATGGGTAGCATATACTACCCAAATATCTGGATAGCATATGCTATCC TAATCTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCAT ATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAATTTATATCT GGGTAGCATAGGCTATCCTAATCTATATCTGGGTAGCATATGCTATCCTAA TCTATATCTGGGTAGTATATGCTATCCTAATCTGTATCCGGGTAGCATATG CTATCCTAATAGAGATTAGGGTAGTATATGCTATCCTAATTTATATCTGGG TAGCATATACTACCCAAATATCTGGATAGCATATGCTATCCTAATCTATAT CTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGCATAGGCTATCCT AATCTATATCTGGGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATA TGCTATCCTAATTTATATCTGGGTAGCATAGGCTATCCTAATCTATATCTG GGTAGCATATGCTATCCTAATCTATATCTGGGTAGTATATGCTATCCTAAT CTGTATCCGGGTAGCATATGCTATCCTCATGATAAGCTGTCAAACATGAGA ATTTTCTTGAAGACGAAAGGGCCTCGTGATACGCCTATTTTATAGGTTAA TGTCATGATAATAATGGTTTCTTAGACGTCAGGTGGCACTTTTCGGGGAAA TGTGCGCGGAACCCCTATTTGTTTATTTTCTAAATACATTCAAATATGTA TCCGCTCATGAGACAATAACCCTGATAAATGCTTCAATAATATTGAAAAAG GAAGAGTATGAGTATTCAACATTTCCGTGTGCCCCTTATTCCCTTTTTTGC GGCATTTTGCCTTCCTGTTTTTGCTCACCCAGAAACGCTGGTGAAAGTAAA

SEQ ID NO	Vector name	Nucleotide sequences
		1234567890123456789012345678901234567890123456789012345678901
		AGATGCTGAAGATCAGTTGGGTGCACGAGTGGGTACATCGAACTGGATCT CAACAGCGGTAAGATCCTTGAGAGTTTTCGCCCCGAAGAACGTTTTTCCAAT GATGAGCACTTTTAAAGTTCTGCTATGTGGCGCGGTATTATCCCGTGTTGA CGCCGGGCAAGAGCAACTCGGTCGCCGCATACACTATTCTCAGAATGACTT GGTTGAGTACTCACCAGTCACAGAAAAGCATCTTACGGATGGCATGACAGT AAGAGAATTATGCAGTGCTGCCATAACCATGAGTGATAAACAACGCGGCCAA CTTACTTCTGACAACGATCGGAGGACCGAAGGAGCTAACCGCTTTTTTTGCA CAACATGGGGGATCATGTAACTCGCCTTGATCGTTGGGAACCGGAGCTGAA TGAAGCCATACCAAACGACGAGCGTGACACCACGATGCCTGCAGCAATGGC AACAAAGTTGCGCAAACATTAACCTGGCGAACTACTTACTCTAGCTTCCCG GCAACAATTAATAGACTGGATGGAGGCGGATAAAGTTGCAGGACCACTTCT GCGCTCGGCCCTTCCGGCTGGCTGGTTTTATTGCTGATAAATCTGGAGCCGG TGAGCGTGGGTCTCGCGGTATCATTGCAGCACTGGGGCCAGATGGTAAGCC CTCCCGTATCGTAGTTATCTACACGACGGGGAGTCAGGCAACTATGGATGA ACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGATTAAGCATTTGGTA ACTGTCAGACCAAGTTTACTCATATATACTTTAGATTGATTTAAAACCTTCA TTTTTAATTTAAAAGGATCTAGGTGAAGATCCTTTTTTGATAATCTCATGAC CAAAATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGA AAAGATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTG CTTGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGCCGGATCA AGAGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGAT ACCAAATACTGTTCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGAA CTCTGTAGCACCCGCCTACATACCTCGCTCTGCTAATCCTGTTACCAGTGGC TGCTGCCAGTGGCGATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATA GTTACCGGATAAGGCGCAGCGGTGCGGGCTGAACGGGGGGTTTCGTGCACACA GCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGCGTGA GCTATGAGAAAGCGCCACGCTTCCCGAAGGGAGAAAGGCGGACAGGTATCC GGTAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTTCCAGGGGG AAACGCCTGGTATCTTTATAGTCCTGTGCGGTTTTCGCCACCTCTGACTTGA GCGTCGATTTTTTGATGCTCGTCAGGGGGGCGGAGCCTATGGAAAAACGC CAGCAACGCGGCCTTTTTACGGTTCCTGGCCTTTTGCTGGCCTTTTGCTCA CATGTTCTTTCTGCGTTATCCCCTGATTCTGTGGATAACCGTATTACCGC CTTTGAGTGAGCTGATACCGCTCGCCGCAGCCGAACGACCGAGCGCAGCGA GTCAGTGAGCGAGGAAGCGGAAGAGCGCCCAATACGCAAACCGCCTCTCCC CGCGCGTTGGCCGATTCAATTAATGCAGCTGGCACGACAGGTTTCCCGACTG GAAAGCGGGCAGTGAGCGCAACGCAATTAATGTGAGTTAGCTCACTCATTA GGCACCCCAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTGTGGAAT TGTGAGCGGATAACAATTTACACAGGAAACAGCTATGACCATGATTACGC CAAGCTCTAGCTAGAGGTCGAGTCCCTCCCCAGCAGGCAGAAGTATGCAAA GCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACCTCCGCCCA TCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATTTCTCCGCCCATGGCTGAC TAATTTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTAT TCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGC TTTGCAAAGATGGATAAAGTTTTTAAACAGAGAGGAATCTTTGCAGCTAATG GACCTTCTAGGTCTTGAAAGGAGTGGGAATTGGCTCCGGTGCCCGTCAGTG GGCAGAGCGCACATCGCCACAGTCCCCGAGAAGTTGGGGGGAGGGGTGCG CAATTGAACCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAAAGTGA TGTCGTGTACTGGCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCGTATAT AAGTGCAGTAGTCGCCGTGAACGTTCTTTTTCGCAACGGGTTTGCCGCCAG AACACAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACGGG TTATGGCCCTTGCGTGCCTTGAATTACTTCCACCTGGCTGCAGTACGTGAT TCTTGATCCCGAGCTTCGGGTTGGAAGTGGGTGGGAGAGTTTCGAGGCCCTG CGCTTAAGGAGCCCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGGGCG CTGGGGCCCGCGCGTGCGAATCTGGTGGCACCTTCGCGCCTGTCTCGCTGC TTTCGATAAGTCTCTAGCCATTTAAAATTTTTGATGACCTGCTGCGACGCT TTTTTTCTGGCAAGATAGTCTTGTAATATGCGGGCCAAGATCTGCACACTGG TATTTTCGGTTTTTTGGGGCCGCGGGCGGCGACGGGGCCCGTGCGTCCCAGCG CACATGTTTCGGCGAGGCGGGGCCTGCGAGCGCGGCCACCGAGAATCGGACG GGGGTAGTCTCAAGCTGGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCC

SEQ ID NO	Vector name	Nucleotide sequences
		123456789012345678901234567890123456789012345678901 GTGTATCGCCCCGCCCTGGGCGGCAAGGCTGGCCCGGTCGGCACCAGTTGC GTGAGCGGAAAGATGGCCGCTTCCCGGCCCTGCTGCAGGGAGCTCAAAATG GAGGACGCGGCGCTCGGGAGAGCGGGCGGGTGAGTCACCCACACAAAGGAA AAGGGCCTTTCCGTCCTCAGCCGTCGCTTCATGTGACTCCACGGAGTACCG GGCGCCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCGTC TTTAGGTTGGGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTG GGTGGAGACTGAAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTGA ATTTGCCCTTTTTGAGTTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGT GGTTCAAAGTTTTTTTCTTCCATTTTCAGGTGTCGTGAGGAATTCTCTAGAG ATCCCTCGACCTCGAGATCCATTGTGCCCGGGCGCCACCATGGAGTTTGGG CTGAGCTGGCTTTTTTCTTGTGCGGATTTTAAAGGTGTCCAGTGC

The present invention incorporates by reference in their entirety techniques well known in the field of molecular biology and drug delivery. These techniques include, but are not limited to, techniques described in the following publications:

Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993);
Ausubel, F.M. et al. eds., Short Protocols In Molecular Biology (4th Ed. 1999) John Wiley & Sons, NY. (ISBN 0-471-32938-X).
Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984);
Giege, R. and Ducruix, A. Barrett, Crystallization of Nucleic Acids and Proteins, a Practical Approach, 2nd ea., pp. 20 1-16, Oxford University Press, New York, New York, (1999);
Goodson, in Medical Applications of Controlled Release, vol. 2, pp. 115-138 (1984);
Hammerling, et al., in: Monoclonal Antibodies and T-Cell Hybridomas 563-681 (Elsevier, N.Y., 1981);
Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988);
Kabat et al., Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md. (1987) and (1991);
Kabat, E.A., *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242;
Kontermann and Dubel eds., Antibody Engineering (2001) Springer-Verlag. New York. 790 pp. (ISBN 3-540-41354-5).
Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990);
Lu and Weiner eds., Cloning and Expression Vectors for Gene Function Analysis (2001) BioTechniques Press. Westborough, MA. 298 pp. (ISBN 1-881299-21-X).
Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Fla. (1974);

Old, R.W. & S.B. Primrose, Principles of Gene Manipulation: An Introduction To Genetic Engineering (3d Ed. 1985) Blackwell Scientific Publications, Boston. Studies in Microbiology; V.2:409 pp. (ISBN 0-632-01318-4).

Sambrook, J. et al. eds., Molecular Cloning: A Laboratory Manual (2d Ed. 1989) Cold Spring Harbor Laboratory Press, NY. Vols. 1-3. (ISBN 0-87969-309-6).

Sustained and Controlled Release Drug Delivery Systems, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978

Winnacker, E.L. From Genes To Clones: Introduction To Gene Technology (1987) VCH Publishers, NY (translated by Horst Ibelgauf). 634 pp. (ISBN 0-89573-614-4).

10 J Neuroscience 29 (14): 4605-15 (2009) -- for glioblastoma
and Cancer Biol Ther. 2006 Jun; 5(6):657-64. for Her2

Incorporation by Reference

15 The contents of all cited references (including literature references, patents, patent applications, and websites) that maybe cited throughout this application are hereby expressly incorporated by reference in their entirety, as are the references cited therein. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of immunology, molecular biology and cell biology, which are well known in the art.

Equivalents

20 The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting of the invention described herein. Scope of the invention is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are therefore
25 intended to be embraced herein.

We claim:

1. A binding protein comprising a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein;

VD1 is a first heavy chain variable domain;

5 VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

10 (X2)_n is (X2)₀ or (X2)₁;

wherein the heavy chain is comprises a cleavage site.

2. A binding protein comprising a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein;

VD1 is a first light chain variable domain;

15 VD2 is a second light chain variable domain;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

20 (X2)_n is (X2)₀ or (X2)₁;

wherein the light chain comprises a cleavage site .

3. A binding protein comprising first and second polypeptide chains, wherein said first polypeptide chain comprises a first VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first heavy chain variable domain;

25 VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1; and

X2 is an Fc region; and

wherein said second polypeptide chain comprises a second VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first light chain variable domain;

VD2 is a second light chain variable domain;

5 C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

10 (X2)_n is (X2)₀ or (X2)₁, wherein at least one of the heavy chain and light chain comprises a cleavage site .

4. A binding protein capable of binding two antigens comprising four polypeptide chains, wherein two polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first heavy chain variable domain;

VD2 is a second heavy chain variable domain;

15 C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1; and

X2 is an Fc region; and

wherein two polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first light chain variable domain;

20 VD2 is a second light chain variable domain;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

25 (X2)_n is (X2)₀ or (X2)₁; wherein at least one of the heavy chain and light chain comprises a cleavage site.

5. The binding protein of any one of claims 1-4, wherein n is 0.

6. The binding protein of any one of claims 1-4, wherein X1 or X2 is an amino acid sequence selected from the group consisting of SEQ ID NOs 1-62, TVA, and AS.

30 7. The binding protein of any one of claims 1-4, wherein the binding protein is cleavable such that binding to at least one of the VD1 and VD2 is enhanced.

8. The binding protein of any one of claims 1-4, wherein the binding protein is cleaved by an enzyme or agent selected from the group consisting of enterokinase, thrombin, PreScission, Tobacco Etch Virus protease (TEV), and tissue plasminogen activator (tPA) + Proline.
9. The binding protein of any one of claims 1-4, wherein the binding protein is cleaved by an enzyme or agent selected from the group consisting of a zinc-dependent endopeptidase, Matrix Metalloproteinase (MMP), a serralyisin, an astacin, an adamalysin, MMP-1; MMP-2; MMP-3; MMP-7; MMP-8; MMP-9; MMP-10; MMP-11; MMP-12; MMP-13; MMP-14; MMP-15; MMP-16; MMP-17; MMP-18; MMP-19; MMP-20; MMP-21; MMP-22; MMP-23A; MMP-23B; MMP-24; MMP-25; MMP-26; MMP-27; MMP-28; a Disintegrin and Metalloproteinase (ADAM); ADAM17; ADAMTS1; ADAM1; ADAM10; ADAM8; ADAMTS4; ADAMTS13; ADAM12; ADAM15; ADAM9; ADAMTS5; ADAM33; ADAM11; ADAM2; ADAMTS2; ADAMTS9; ADAMTS3; ADAMTS7; ADAM22; ADAM28; ADAMTS12; ADAM19; ADAMTS8; ADAM29; ADAM23; ADAM3A; ADAM18; ADAMTS6; ADAM7; ADAMDES1; ADAM20; ADAM6; ADAM21; ADAM3B; ADAMTSL3; ADAMTSL4; ADAM30; ADAMTS20; ADAMTSL2; a Caspase; Caspases 1-12, Caspase 14; a Cathepsin; Cathepsin G; Cathepsin B; Cathepsin D; Cathepsin L1; Cathepsin C; Cathepsin K; Cathepsin S; Cathepsin H; Cathepsin A; Cathepsin E; Cathepsin L; Cathepsin Z; Cathepsin F; Cathepsin G-like 2; Cathepsin L-like 1; Cathepsin W; Cathepsin L-like 2; Cathepsin L-like 3; Cathepsin L-like 4; Cathepsin L-like 5; Cathepsin L-like 6; Cathepsin L-like 7; Cathepsin O; a Calpain; Calpain 3; Calpain 10; Calpain 1 (mu/l) large subunit; Calpain, small subunit 1; Calpain 2, (mu/l); large subunit; Calpain 9; Calpain 11; Calpain 5; Calpain 6; Calpain 13; Calpain 8; Calpain, small subunit 2; Calpain 15; Calpain 12; Calpain 7; and Calpain 8.
10. The binding protein of any one of claims 1-4, wherein the cleavage site is between at least one VD1 and VD2.
11. The binding protein of any one of claims 1-4, wherein the cleavage site is in at least one linker.
12. The binding protein of any one of claims 1-4, wherein at least one of the VD1 or VD2 does not bind to its target until a cleavage between the VD1 and VD2 occurs.
13. The binding protein of any one of claims 1-4, wherein the linker of the binding protein has been selectively cleaved by an enzyme.
14. The binding protein of any one of claims 1-4, wherein the linker of the binding protein has been selectively cleaved by an enzyme during the manufacturing process.
15. The binding protein of any one of claims 1-4, wherein the linker of the binding protein has been selectively cleaved by an enzyme when the DVD-Ig is adjacent to at least one target.

16. The binding protein of any one of claims 1-4, wherein the binding protein is selectively cleaved by an enzyme when the DVD-Ig is bound to at least one target.
17. The binding protein of any one of claims 1-4, wherein the Fc region is selected from the group consisting of native sequence Fc region and a variant sequence Fc region.
- 5 18. The binding protein of claim 17, wherein the Fc region is selected from the group consisting of an Fc region from an IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, and IgD.
19. The binding protein of any one of claims 1-4, wherein said binding protein has an on rate constant (K_{on}) to said one or more targets selected from the group consisting of: at least about $10^2 M^{-1} s^{-1}$; at least about $10^3 M^{-1} s^{-1}$; at least about $10^4 M^{-1} s^{-1}$; at least about $10^5 M^{-1} s^{-1}$; and at least
10 about $10^6 M^{-1} s^{-1}$, as measured by surface plasmon resonance.
20. The binding protein of any one of claims 1-4, wherein said binding protein has an off rate constant (K_{off}) to said one or more targets selected from the group consisting of: at most about $10^{-3} s^{-1}$; at most about $10^{-4} s^{-1}$; at most about $10^{-5} s^{-1}$; and at most about $10^{-6} s^{-1}$, as measured by surface plasmon resonance.
- 15 21. The binding protein of any one of claims 1-4, wherein said binding protein has a dissociation constant (K_D) to said one or more targets selected from the group consisting of: at most about $10^{-7} M$; at most about $10^{-8} M$; at most about $10^{-9} M$; at most about $10^{-10} M$; at most about $10^{-11} M$; at most about $10^{-12} M$; and at most $10^{-13} M$.
22. A binding protein conjugate comprising a binding protein according to any one of claims 1-4,
20 said binding protein conjugate further comprising an agent selected from the group consisting of; an immunoadhesion molecule, an imaging agent, a therapeutic agent, and a cytotoxic agent.
23. The binding protein conjugate of claim 22, wherein said agent is an imaging agent selected from the group consisting of a radiolabel, an enzyme, a fluorescent label, a luminescent label, a bioluminescent label, a magnetic label, and biotin.
- 25 24. The binding protein conjugate of claim 22, wherein said imaging agent is a radiolabel selected from the group consisting of: ^3H , ^{14}C , ^{35}S , ^{90}Y , ^{99}Tc , ^{111}In , ^{125}I , ^{131}I , ^{177}Lu , ^{166}Ho , and ^{153}Sm .
25. The binding protein conjugate of claim 22, wherein said agent is a therapeutic or cytotoxic agent selected from the group consisting of; an anti-metabolite, an alkylating agent, an antibiotic,
30 a growth factor, a cytokine, an anti-angiogenic agent, an anti-mitotic agent, an anthracycline, toxin, and an apoptotic agent.
26. The binding protein of any one of claims 1-4, wherein said binding protein is a crystallized binding protein.

27. The binding protein of claim 26, wherein said crystal is a carrier-free pharmaceutical controlled release crystal.
28. An isolated nucleic acid encoding a binding protein amino acid sequence according to any one of claims 1-11.
- 5 29. A vector comprising an isolated nucleic acid according to claim 18.
30. The vector of claim 29, wherein said vector is selected from the group consisting of pcDNA, pTT, pTT3, pEFBOS, pBV, pJV, pcDNA3.1 TOPO, pEF6 TOPO, and pBJ.
31. A host cell comprising a vector according to claim 29.
32. The host cell of claim 31, wherein said host cell is a prokaryotic cell.
- 10 33. The host cell of claim 32, wherein said host cell is E.Coli.
34. The host cell of claim 31, wherein said host cell is a eukaryotic cell.
35. The host cell of claim 34, wherein said eukaryotic cell is selected from the group consisting of protist cell, animal cell, plant cell and fungal cell.
36. The host cell of claim 35, wherein said animal cell is selected from the group consisting of; a
15 mammalian cell, an avian cell, and an insect cell.
37. The host cell of claim 34, wherein said host cell is a CHO cell.
38. The host cell of claim 34, wherein said host cell is COS.
39. The host cell of claim 34, wherein said host cell is a yeast cell.
40. The host cell of claim 39, wherein said yeast cell is *Saccharomyces cerevisiae*.
- 20 41. The host cell of claim 36, wherein said host cell is an insect Sf9 cell.
42. A method of producing a binding protein, comprising culturing a host cell described in any one of claims 31-41 in culture medium under conditions sufficient to produce the binding protein
43. The method of claim 42, wherein 50%-75% of the binding protein produced is a dual specific tetravalent binding protein.
- 25 44. The method of claim 42, wherein 75%-90% of the binding protein produced is a dual specific tetravalent binding protein.
45. The method of claim 42, wherein 90%-95% of the binding protein produced is a dual specific tetravalent binding protein.
46. A protein produced according to the method of claim 42.

47. A pharmaceutical composition comprising the binding protein of any one of claims 1-27, and a pharmaceutically acceptable carrier.

48. The pharmaceutical composition of claim 47 further comprising at least one additional therapeutic agent.

5 49. The pharmaceutical composition of claim 48, wherein said additional therapeutic agent is selected from the group consisting of: Therapeutic agent, imaging agent, cytotoxic agent, angiogenesis inhibitors; kinase inhibitors; co-stimulation molecule blockers; adhesion molecule blockers; anti-cytokine antibody or functional fragment thereof; methotrexate; cyclosporin; rapamycin; FK506; detectable label or reporter; a TNF antagonist; an antirheumatic; a muscle
10 relaxant, a narcotic, a non-steroid anti-inflammatory drug (NSAID), an analgesic, an anesthetic, a sedative, a local anesthetic, a neuromuscular blocker, an antimicrobial, an antipsoriatic, a corticosteroid, an anabolic steroid, an erythropoietin, an immunization, an immunoglobulin, an immunosuppressive, a growth hormone, a hormone replacement drug, a radiopharmaceutical, an antidepressant, an antipsychotic, a stimulant, an asthma medication, a beta agonist, an inhaled
15 steroid, an epinephrine or analog, a cytokine, and a cytokine antagonist.

50. A method for treating a subject for a disease or a disorder by administering to the subject the binding protein of any one of claims 1-27 such that treatment is achieved.

51. The method of claim 50, wherein said disorder is selected from the group comprising rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, septic arthritis, Lyme arthritis,
20 psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's
25 disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis, hemolytic anemia, malignancies,
30 heart failure, myocardial infarction, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella associated arthropathy, spondyloarthropathy, atheromatous disease/arteriosclerosis,
35 atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus,

pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis, primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired

5 Immunodeficiency Related Diseases, Hepatitis B, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease,

10 systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, fibrosis, radiation fibrosis,

15 bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ

20 transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS, glomerulonephritides, microscopic vasculitis of the kidneys, lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension

25 secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease, systemic sclerosis, Sjögren's syndrome, Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism,

30 primary myxoedema, phacogenic uveitis, primary vasculitis, vitiligo acute liver disease, chronic liver diseases, alcoholic cirrhosis, alcohol-induced liver injury, choleostasis, idiosyncratic liver disease, Drug-Induced hepatitis, Non-alcoholic Steatohepatitis, allergy and asthma, group B streptococci (GBS) infection, mental disorders (*e.g.*, depression and schizophrenia), Th2 Type and Th1 Type mediated diseases, acute and chronic pain (different forms of pain), and cancers such as

35 lung, breast, stomach, bladder, colon, pancreas, ovarian, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma) Abetalipoproteinemia, Acrocyanosis, acute and chronic parasitic or infectious processes, acute leukemia, acute lymphoblastic leukemia

(ALL), acute myeloid leukemia (AML), acute or chronic bacterial infection, acute pancreatitis, acute renal failure, adenocarcinomas, aerial ectopic beats, AIDS dementia complex, alcohol-induced hepatitis, allergic conjunctivitis, allergic contact dermatitis, allergic rhinitis, allograft rejection, alpha-1- antitrypsin deficiency, amyotrophic lateral sclerosis, anemia, angina pectoris, anterior horn cell degeneration, anti cd3 therapy, antiphospholipid syndrome, anti-receptor hypersensitivity reactions, aortic and peripheral aneurysms, aortic dissection, arterial hypertension, arteriosclerosis, arteriovenous fistula, ataxia, atrial fibrillation (sustained or paroxysmal), atrial flutter, atrioventricular block, B cell lymphoma, bone graft rejection, bone marrow transplant (BMT) rejection, bundle branch block, Burkitt's lymphoma, Burns, cardiac arrhythmias, cardiac stun syndrome, cardiac tumors, cardiomyopathy, cardiopulmonary bypass inflammation response, cartilage transplant rejection, cerebellar cortical degenerations, cerebellar disorders, chaotic or multifocal atrial tachycardia, chemotherapy associated disorders, chronic myelocytic leukemia (CML), chronic alcoholism, chronic inflammatory pathologies, chronic lymphocytic leukemia (CLL), chronic obstructive pulmonary disease (COPD), chronic salicylate intoxication, colorectal carcinoma, congestive heart failure, conjunctivitis, contact dermatitis, cor pulmonale, coronary artery disease, Creutzfeldt-Jakob disease, culture negative sepsis, cystic fibrosis, cytokine therapy associated disorders, Dementia pugilistica, demyelinating diseases, dengue hemorrhagic fever, dermatitis, dermatologic conditions, diabetes, diabetes mellitus, diabetic atherosclerotic disease, Diffuse Lewy body disease, dilated congestive cardiomyopathy, disorders of the basal ganglia, Down's Syndrome in middle age, drug- induced movement disorders induced by drugs which block CNS dopamine receptors, drug sensitivity, eczema, encephalomyelitis, endocarditis, endocrinopathy, epiglottitis, epstein-barr virus infection, erythromelalgia, extrapyramidal and cerebellar disorders, familial hemophagocytic lymphohistiocytosis, fetal thymus implant rejection, Friedreich's ataxia, functional peripheral arterial disorders, fungal sepsis, gas gangrene, gastric ulcer, glomerular nephritis, graft rejection of any organ or tissue, gram negative sepsis, gram positive sepsis, granulomas due to intracellular organisms, hairy cell leukemia, Hallervorden-Spatz disease, hashimoto's thyroiditis, hay fever, heart transplant rejection, hemachromatosis, hemodialysis, hemolytic uremic syndrome/thrombolytic thrombocytopenic purpura, hemorrhage, hepatitis (A), His bundle arrhythmias, HIV infection/HIV neuropathy, Hodgkin's disease, hyperkinetic movement disorders, hypersensitivity reactions, hypersensitivity pneumonitis, hypertension, hypokinetic movement disorders, hypothalamic-pituitary-adrenal axis evaluation, idiopathic Addison's disease, idiopathic pulmonary fibrosis, antibody mediated cytotoxicity, Asthenia, infantile spinal muscular atrophy, inflammation of the aorta, influenza a, ionizing radiation exposure, iridocyclitis/uveitis/optic neuritis, ischemia- reperfusion injury, ischemic stroke, juvenile rheumatoid arthritis, juvenile spinal muscular atrophy, Kaposi's sarcoma, kidney transplant rejection, legionella, leishmaniasis, leprosy, lesions of the corticospinal system, lipedema, liver transplant rejection, lymphedema,

malaria, malignant Lymphoma, malignant histiocytosis, malignant melanoma, meningitis,
 meningococemia, metabolic/idiopathic, migraine headache, mitochondrial multi.system disorder,
 mixed connective tissue disease, monoclonal gammopathy, multiple myeloma, multiple systems
 degenerations (Mencel Dejerine- Thomas Shi-Drager and Machado-Joseph), myasthenia gravis,
 5 mycobacterium avium intracellulare, mycobacterium tuberculosis, myelodysplastic syndrome,
 myocardial infarction, myocardial ischemic disorders, nasopharyngeal carcinoma, neonatal
 chronic lung disease, nephritis, nephrosis, neurodegenerative diseases, neurogenic I muscular
 atrophies , neutropenic fever, non- hodgkins lymphoma, occlusion of the abdominal aorta and its
 branches, occlusive arterial disorders, okt3 therapy, orchitis/epididymitis, orchitis/vasectomy
 10 reversal procedures, organomegaly, osteoporosis, pancreas transplant rejection, pancreatic
 carcinoma, paraneoplastic syndrome/hypercalcemia of malignancy, parathyroid transplant
 rejection, pelvic inflammatory disease, perennial rhinitis, pericardial disease, peripheral
 atherosclerotic disease, peripheral vascular disorders, peritonitis, pernicious anemia,
 pneumocystis carinii pneumonia, pneumonia, POEMS syndrome (polyneuropathy, organomegaly,
 15 endocrinopathy, monoclonal gammopathy, and skin changes syndrome), post perfusion
 syndrome, post pump syndrome, post-MI cardiomyopathy syndrome, preeclampsia, Progressive
 supranuclear Palsy, primary pulmonary hypertension, radiation therapy, Raynaud's phenomenon
 and disease, Raynaud's disease, Refsum's disease, regular narrow QRS tachycardia, renovascular
 hypertension, reperfusion injury, restrictive cardiomyopathy, sarcomas, scleroderma, senile
 20 chorea, Senile Dementia of Lewy body type, seronegative arthropathies, shock, sickle cell anemia,
 skin allograft rejection, skin changes syndrome, small bowel transplant rejection, solid tumors,
 specific arrhythmias, spinal ataxia, spinocerebellar degenerations, streptococcal myositis, structural
 lesions of the cerebellum, Subacute sclerosing panencephalitis, Syncope, syphilis of the
 cardiovascular system, systemic anaphylaxis, systemic inflammatory response syndrome,
 25 systemic onset juvenile rheumatoid arthritis, T-cell or FAB ALL, Telangiectasia, thromboangitis
 obliterans, thrombocytopenia, toxicity, transplants, trauma/hemorrhage, type III hypersensitivity
 reactions, type IV hypersensitivity, unstable angina, uremia, urosepsis, urticaria, valvular heart
 diseases, varicose veins, vasculitis, venous diseases, venous thrombosis, ventricular fibrillation,
 viral and fungal infections, viral encephalitis/aseptic meningitis, viral-associated hemophagocytic
 30 syndrome, Wernicke- Korsakoff syndrome, Wilson's disease, xenograft rejection of any organ or
 tissue, acute coronary syndromes, acute idiopathic polyneuritis, acute inflammatory
 demyelinating polyradiculoneuropathy, acute ischemia, adult Still's disease, alopecia areata,
 anaphylaxis, anti-phospholipid antibody syndrome, aplastic anemia, arteriosclerosis, atopic
 eczema, atopic dermatitis, autoimmune dermatitis, autoimmune disorder associated with
 35 streptococcus infection, autoimmune enteropathy, autoimmune hearing loss, autoimmune
 lymphoproliferative syndrome (ALPS), autoimmune myocarditis, autoimmune premature ovarian
 failure, blepharitis, bronchiectasis, bullous pemphigoid, cardiovascular disease, catastrophic

- antiphospholipid syndrome, celiac disease, cervical spondylosis, chronic ischemia, cicatricial pemphigoid, clinically isolated syndrome (cis) with risk for multiple sclerosis, conjunctivitis, childhood onset psychiatric disorder, chronic obstructive pulmonary disease (COPD), dacryocystitis, dermatomyositis, diabetic retinopathy, diabetes mellitus, disk herniation, disk prolaps, drug induced immune hemolytic anemia, endocarditis, endometriosis, endophthalmitis, episcleritis, erythema multiforme, erythema multiforme major, gestational pemphigoid, Guillain-Barré syndrome (GBS), hay fever, Hughes syndrome, idiopathic Parkinson's disease, idiopathic interstitial pneumonia, IgE-mediated allergy, immune hemolytic anemia, inclusion body myositis, infectious ocular inflammatory disease, inflammatory demyelinating disease, inflammatory heart disease, inflammatory kidney disease, IPF/UIP, iritis, keratitis, keratoconjunctivitis sicca, Kussmaul disease or Kussmaul-Meier disease, Landry's paralysis, Langerhan's cell histiocytosis, livedo reticularis, macular degeneration, microscopic polyangiitis, morbus bechterev, motor neuron disorders, mucous membrane pemphigoid, multiple organ failure, myasthenia gravis, myelodysplastic syndrome, myocarditis, nerve root disorders, neuropathy, non-A non-B hepatitis, optic neuritis, osteolysis, ovarian cancer, pauciarticular JRA, peripheral artery occlusive disease (PAOD), peripheral vascular disease (PVD), peripheral artery disease (PAD), phlebitis, polyarteritis nodosa (or periarteritis nodosa), polychondritis, polymyalgia rheumatica, poliosis, polyarticular JRA, polyendocrine deficiency syndrome, polymyositis, polymyalgia rheumatica (PMR), post-pump syndrome, primary Parkinsonism, prostate and rectal cancer and hematopoietic malignancies (leukemia and lymphoma), prostatitis, pure red cell aplasia, primary adrenal insufficiency, recurrent neuromyelitis optica, restenosis, rheumatic heart disease, sapho (synovitis, acne, pustulosis, hyperostosis, and osteitis), scleroderma, secondary amyloidosis, shock lung, scleritis, sciatica, secondary adrenal insufficiency, silicone associated connective tissue disease, sneddon-wilkinson dermatosis, spondylitis ankylosans, Stevens-Johnson syndrome (SJS), systemic inflammatory response syndrome, temporal arteritis, toxoplasmic retinitis, toxic epidermal necrolysis, transverse myelitis, TRAPS (tumor necrosis factor receptor, type 1 allergic reaction, type II diabetes, urticaria, usual interstitial pneumonia (UIP), vasculitis, vernal conjunctivitis, viral retinitis, Vogt-Koyanagi-Harada syndrome (VKH syndrome), wet macular degeneration, wound healing, yersinia and salmonella associated arthropathy.
52. The method of claim 50, wherein said administering to the subject is by at least one mode selected from parenteral, subcutaneous, intramuscular, intravenous, intrarticular, intrabronchial, intraabdominal, intracapsular, intracartilaginous, intracavitary, intracelical, intracerebellar, intracerebroventricular, intracolic, intracervical, intragastric, intrahepatic, intramyocardial, intraosteal, intrapelvic, intrapericardiac, intraperitoneal, intrapleural, intraprostatic, intrapulmonary, intrarectal, intrarenal, intraretinal, intraspinal, intrasynovial, intrathoracic, intrauterine, intravesical, bolus, vaginal, rectal, buccal, sublingual, intranasal, and transdermal.

53. A method for generating a Dual Variable Domain Immunoglobulin capable of binding two antigens comprising the steps of

- a) obtaining a first parent antibody or antigen binding portion thereof, capable of binding a first antigen;
- 5 b) obtaining a second parent antibody or antigen binding portion thereof, capable of binding a second antigen;
- c) constructing first and third polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein

10 VD1 is a first heavy chain variable domain obtained from said first parent antibody or antigen binding portion thereof;

VD2 is a second heavy chain variable domain obtained from said second parent antibody or antigen binding portion thereof;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1;

15 X2 is an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

(X2)_n is (X2)₀ or (X2)₁; and
- d) constructing second and fourth polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein

20 VD1 is a first light chain variable domain obtained from said first parent antibody or antigen binding portion thereof;

VD2 is a second light chain variable domain obtained from said second parent antibody or antigen binding thereof;

C is a light chain constant domain;

25 X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

(X2)_n is (X2)₀ or (X2)₁; and
- e) expressing said first, second, third and fourth polypeptide chains;

such that a Dual Variable Domain Immunoglobulin capable of binding said first and said second antigen is generated, wherein the at least one of the heavy and light chain comprises a cleavage site.

54. The method of claim 53, wherein the Fc region is selected from the group consisting of a native sequence Fc region and a variant sequence Fc region.

55. The method of claim 53, wherein the Fc region is selected from the group consisting of an Fc region from an IgG1, IgG2, IgG3, IgG4, IgA, IgM, IgE, and IgD.

56. A method for generating a Dual Variable Domain Immunoglobulin capable of binding two antigens with desired properties comprising the steps of

- a) obtaining a first parent antibody or antigen binding portion thereof, capable of binding a first antigen and possessing at least one desired property exhibited by the Dual Variable Domain Immunoglobulin;
- b) obtaining a second parent antibody or antigen binding portion thereof, capable of binding a second antigen and possessing at least one desired property exhibited by the Dual Variable Domain Immunoglobulin;

- c) constructing first and third polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein;

VD1 is a first heavy chain variable domain obtained from said first parent antibody or antigen binding portion thereof;

- VD2 is a second heavy chain variable domain obtained from said second parent antibody or antigen binding portion thereof;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

- (X1)_n is (X1)₀ or (X1)₁; and

(X2)_n is (X2)₀ or (X2)₁;

- d) constructing second and fourth polypeptide chains comprising VD1-(X1)_n-VD2-C-(X2)_n, wherein;

- VD1 is a first light chain variable domain obtained from said first parent antibody or antigen binding portion thereof;

VD2 is a second light chain variable domain obtained from said second parent antibody or antigen binding portion thereof;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

5 (X2)_n is (X2)₀ or (X2)₁;

e) expressing said first, second, third and fourth polypeptide chains;

such that a Dual Variable Domain Immunoglobulin capable of binding said first and said second antigen with desired properties is generated, wherein the at least one of the heavy and light chain comprises a cleavage site.

10 57. A method for improving a characteristic of the binding protein of claim 3 or 4, the method comprising the steps of:

(a) determining the characteristic of the binding protein prior to alteration;

(a) altering the length and/or sequence of (X1)₁ of the heavy and/or light chain thereby providing an altered heavy and/ or light chain;

15 (b) determining the improved characteristic of the altered binding protein comprising the altered heavy and light chains; wherein at least one of the heavy and light chain comprises a cleavage site.

58. A method for improving a characteristic of the binding protein of claim 3 or 4, the method comprising the steps of:

20 (a) determining the characteristic of the binding protein prior to alteration;

(b) altering the first and second polypeptide chains such that VD1-(X1)_n-VD2-C-(X2)_n is changed to VD2-(X1)_n-VD1-C-(X2)_n, thereby providing altered heavy and light chains;

(c) determining the improved characteristic of the altered binding protein comprising the altered heavy and light chains; wherein at least one of the heavy and light chain comprises a
25 cleavage site.

59. A method for improving a characteristic of the binding protein of claim 3 or 4, the method comprising the steps of:

(a) determining the characteristic of the binding protein prior to alteration;

(b) altering the first and/or second polypeptide chains such that the sequence of only one
30 of the VD1 or VD2 of the heavy and/or light chain is changed; and

(c) determining the characteristic of the altered binding protein comprising the altered heavy and light chains; wherein at least one of the heavy and light chain comprises a cleavage site.

60. The method of any one of claims 57-59, wherein the characteristic is selected from the group consisting of binding to target antigen, expression yield from host cell, in vitro halflife, in vivo halflife, stability, solubility, affinity, avidity, and improved effector function.

61. The method of claim 57, wherein the length of the linker of the altered heavy chain is increased.

62. The method of claim 57, wherein the length of the linker of the altered heavy chain is decreased.

63. The method of claim 57, wherein the length of the linker of the altered light chain is increased.

64. The method of claim 57, wherein the length of the linker of the altered light chain is decreased.

65. The method of claim 65, wherein the cleavage site is between at least one VD1 and VD2.

66. The method of claim 65, wherein the cleavage site is in at least one linker.

67. A method for treating a subject for a disease or a disorder by administering to the subject the binding protein of any one of claims 58-66, such that treatment is achieved.

68. The method of claim 67, wherein at least one of a VD1 or VD2 does not bind its target until the binding protein is cleaved.

69. The method of claim 67, wherein the VD2 does not bind its target until the binding protein is cleaved.

70. The method of claim 67, wherein the VD1 is released when the binding protein is cleaved.

71. The method of claim 67, wherein the VD1 is released when the VD2 binds to its target.

72. A binding protein comprising a polypeptide chain, wherein said polypeptide chain comprises VD1-(X1)_n-VD2-C-(X2)_n, wherein;

VD1 is a first heavy chain variable domain;

VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 is an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

(X2)_n is (X2)₀ or (X2)₁;

wherein the VD1 and VD2 heavy chain variable domains comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, and 78.

73. A binding protein comprising a polypeptide chain, wherein said polypeptide chain comprises
5 VD1-(X1)_n-VD2-C-(X2)_n, wherein;

VD1 is a first light chain variable domain;

VD2 is a second light chain variable domain;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

10 X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

(X2)_n is (X2)₀ or (X2)₁;

wherein the VD1 and VD2 light chain variable domains comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 65, 67, 69, 71, 73, 75, 77, and 79.

15 74. A binding protein comprising first and second polypeptide chains, wherein said first polypeptide chain comprises a first VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first heavy chain variable domain;

VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

20 X1 is a linker with the proviso that it is not CH1; and

X2 is an Fc region; and

wherein said second polypeptide chain comprises a second VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first light chain variable domain;

25 VD2 is a second light chain variable domain;

C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

(X2)_n is (X2)₀ or (X2)₁, wherein the VD1 and VD2 heavy chain variable domains comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, and 78 and wherein the VD1 and VD2 light chain variable domains comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 65, 67, 69, 71, 73, 75, 77, and 79.

- 5 75. A binding protein capable of binding two antigens comprising four polypeptide chains, wherein two polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first heavy chain variable domain;

VD2 is a second heavy chain variable domain;

C is a heavy chain constant domain;

10 X1 is a linker with the proviso that it is not CH1; and

X2 is an Fc region; and

wherein two polypeptide chains comprise VD1-(X1)_n-VD2-C-(X2)_n, wherein

VD1 is a first light chain variable domain;

VD2 is a second light chain variable domain;

15 C is a light chain constant domain;

X1 is a linker with the proviso that it is not CH1;

X2 does not comprise an Fc region;

(X1)_n is (X1)₀ or (X1)₁; and

20 (X2)_n is (X2)₀ or (X2)₁; wherein the VD1 and VD2 heavy chain variable domains comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 64, 66, 68, 70, 72, 74, 76, and 78 and wherein the VD1 and VD2 light chain variable domains comprise an amino acid sequence selected from the group consisting of SEQ ID NOs: 65, 67, 69, 71, 73, 75, 77, and 79.

Attorne




Figure 1**A****B**

Figure 1

A



B

