

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2019364336 B2**

(54) Title
TYK2 inhibitors and uses thereof

(51) International Patent Classification(s)
C07D 403/14 (2006.01) **C07D 413/14** (2006.01)
C07D 213/74 (2006.01) **C07D 471/04** (2006.01)
C07D 401/04 (2006.01) **C07D 471/10** (2006.01)
C07D 401/14 (2006.01) **C07D 487/04** (2006.01)
C07D 405/14 (2006.01) **C07D 498/04** (2006.01)

(21) Application No: **2019364336** (22) Date of Filing: **2019.10.22**

(87) WIPO No: **WO20/086616**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/749,003	2018.10.22	US
62/756,942	2018.11.07	US
62/839,459	2019.04.26	US
62/875,449	2019.07.17	US
62/893,721	2019.08.29	US
62/907,354	2019.09.27	US

(43) Publication Date: **2020.04.30**

(44) Accepted Journal Date: **2023.11.16**

(71) Applicant(s)
Alumis Inc.

(72) Inventor(s)
JIN, Bohan;DONG, Qing;HUNG, Gene;KALDOR, Stephen W.

(74) Agent / Attorney
**Pizzeyes Patent and Trade Mark Attorneys Pty Ltd, Brisbane Club Tower L 15 241
Adelaide St, Brisbane City, QLD, 4000, AU**

(56) Related Art
EP 2428508 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

30 April 2020 (30.04.2020)



(10) International Publication Number

WO 2020/086616 A1

(51) International Patent Classification:

C07D 403/14 (2006.01) C07D 413/14 (2006.01)
C07D 213/74 (2006.01) C07D 471/04 (2006.01)
C07D 401/04 (2006.01) C07D 471/10 (2006.01)
C07D 401/14 (2006.01) C07D 487/04 (2006.01)
C07D 405/14 (2006.01) C07D 498/04 (2006.01)

UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2019/057485

Published:

— with international search report (Art. 21(3))

(22) International Filing Date:

22 October 2019 (22.10.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/749,003 22 October 2018 (22.10.2018) US
62/756,942 07 November 2018 (07.11.2018) US
62/839,459 26 April 2019 (26.04.2019) US
62/875,449 17 July 2019 (17.07.2019) US
62/893,721 29 August 2019 (29.08.2019) US
62/907,354 27 September 2019 (27.09.2019) US

(71) Applicant: **FRONTERA U.S. PHARMACEUTICALS LLC** [US/US]; 11526 Sorrento Valley Road, Suite D, San Diego, California 92121 (US).

(72) Inventors: **JIN, Bohan**; 11526 Sorrento Valley Road, Suite D, San Diego, California 92121 (US). **DONG, Qing**; 11526 Sorrento Valley Road, Suite D, San Diego, California 92121 (US). **HUNG, Gene**; 11526 Sorrento Valley Road, Suite D, San Diego, California 92121 (US). **KALDOR, Stephen W.**; 11526 Sorrento Valley Road, Suite D, San Diego, California 92121 (US).

(74) Agent: **BONNEFOUS, Celine**; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, California 94304 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,

(54) Title: TYK2 INHIBITORS AND USES THEREOF

(57) Abstract: Described herein are compounds that are useful in treating a TYK2-mediated disorder. In some embodiments, the TYK2-mediated disorder is an autoimmune disorder, an inflammatory disorder, a proliferative disorder, an endocrine disorder, a neurological disorder, or a disorder associated with transplantation.



WO 2020/086616 A1

TYK2 INHIBITORS AND USES THEREOF**CROSS-REFERENCE**

[0001] This patent application claims the benefit of US Provisional Application No. 62/749,003, filed October 22, 2018; US Provisional Application No. 62/756,942, filed November 7, 2018; US Provisional Application No. 62/839,459, filed April 26, 2019; US Provisional Application No. 62/875,449, filed July 17, 2019; US Provisional Application No. 62/893,721, filed August 29, 2019; and US Provisional Application No. 62/907,354, filed September 27, 2019 each of which is incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

[0002] Described herein are compounds, methods of making such compounds, pharmaceutical compositions and medicaments comprising such compounds, and methods of using such compounds for inhibiting nonreceptor tyrosine-protein kinase 2 ("TYK2"), also known as Tyrosine kinase 2.

BACKGROUND OF THE INVENTION

[0003] TYK2 is a non-receptor tyrosine kinase member of the Janus kinase (JAKs) family of protein kinases. The mammalian JAK family consists of four members, TYK2, JAK1, JAK2, and JAK3. JAK proteins, including TYK2, are integral to cytokine signaling. TYK2 associates with the cytoplasmic domain of type I and type II cytokine receptors, as well as interferon types I and III receptors, and is activated by those receptors upon cytokine binding. Cytokines implicated in TYK2 activation include interferons (e.g. IFN- α , IFN- β , IFN- κ , IFN- δ , IFN- ϵ , IFN- τ , IFN- ω , and IFN- ζ (also known as limitin), and interleukins (e.g. IL-4, IL-6, IL-10, IL-11, IL-12, IL-13, L-22, IL-23, IL-27, IL-31, oncostatin M, ciliary neurotrophic factor, cardiotrophin 1, cardiotrophin-like cytokine, and LIF). The activated TYK2 then goes on to phosphorylate further signaling proteins such as members of the STAT family, including STAT1, STAT2, STAT4, and STAT6.

[0004] TYK2 activation by IL-23, has been linked to inflammatory bowel disease (IBD), Crohn's disease, and ulcerative colitis. A genome-wide association study of 2,622 individuals with psoriasis identified associations between disease susceptibility and TYK2. Knockout or tyrophostin inhibition of TYK2 significantly reduces both IL-23 and IL-22-induced dermatitis.

[0005] TYK2 also plays a role in respiratory diseases such as asthma, chronic obstructive pulmonary disease (COPD), lung cancer, and cystic fibrosis. Goblet cell hyperplasia (GCH) and mucous hypersecretion is mediated by IL-13-induced activation of TYK2, which in turn activates STAT6.

[0006] Decreased TYK2 activity leads to protection of joints from collagen antibody-induced arthritis, a model of human rheumatoid arthritis. Mechanistically, decreased Tyk2 activity reduced the production of Th1/Th17-related cytokines and matrix metalloproteases, and other key markers of inflammation.

[0007] TYK2 knockout mice showed complete resistance in experimental autoimmune encephalomyelitis (EAE, an animal model of multiple sclerosis (MS)), with no infiltration of CD4 T cells in the spinal cord, as compared to controls, suggesting that TYK2 is essential to pathogenic CD4-

mediated disease development in MS. This corroborates earlier studies linking increased TYK2 expression with MS susceptibility. Loss of function mutation in TYK2, leads to decreased demyelination and increased remyelination of neurons, further suggesting a role for TYK2 inhibitors in the treatment of MS and other CNS demyelination disorders.

[0008] TYK2 is the sole signaling messenger common to both IL-12 and IL-23. TYK2 knockout reduced methylated BSA injection-induced footpad thickness, imiquimod-induced psoriasis-like skin inflammation, and dextran sulfate sodium or 2,4,6-trinitrobenzene sulfonic acid-induced colitis in mice.

[0009] Joint linkage and association studies of various type I IFN signaling genes with systemic lupus erythematosus (SLE, an autoimmune disorder), showed a strong, and significant correlation between loss of function mutations to TYK2 and decreased prevalence of SLE in families with affected members. Genome-wide association studies of individuals with SLE versus an unaffected cohort showed highly significant correlation between the TYK2 locus and SLE.

[0010] TYK2 has been shown to play an important role in maintaining tumor surveillance and TYK2 knockout mice showed compromised cytotoxic T cell response, and accelerated tumor development. However, these effects were linked to the efficient suppression of natural killer (NK) and cytotoxic T lymphocytes, suggesting that TYK2 inhibitors would be highly suitable for the treatment of autoimmune disorders or transplant rejection. Although other JAK family members such as JAK3 have similar roles in the immune system, TYK2 has been suggested as a superior target because of its involvement in fewer and more closely related signaling pathways, leading to fewer off-target effects.

[0011] Studies in T-cell acute lymphoblastic leukemia (T-ALL) indicate that T-ALL is highly dependent on IL-10 via TYK2 via STAT1-mediated signal transduction to maintain cancer cell survival through upregulation of anti-apoptotic protein BCL2. Knockdown of TYK2, but not other JAK family members, reduced cell growth. Specific activating mutations to TYK2 that promote cancer cell survival include those to the FERM domain (G36D, S47N, and R425H), the JH2 domain (V731I), and the kinase domain (E957D and R1027H). However, it was also identified that the kinase function of TYK2 is required for increased cancer cell survival, as TYK2 enzymes featuring kinase-dead mutations (M978Y or M978F) in addition to an activating mutation (E957D) resulted in failure to transform.

[0012] Thus, selective inhibition of TYK2 has been suggested as a suitable target for patients with IL-10 and/or BCL2-addicted tumors, such as 70% of adult T-cell leukemia cases. TYK2 mediated STAT3 signaling has also been shown to mediate neuronal cell death caused by amyloid- β (A β) peptide. Decreased TYK2 phosphorylation of STAT3 following A β administration lead to decreased neuronal cell death, and increased phosphorylation of STAT3 has been observed in postmortem brains of Alzheimer's patients.

[0013] Inhibition of JAK-STAT signaling pathways is also implicated in hair growth, and the reversal of the hair loss associated with alopecia areata.


[0014] Accordingly, compounds that inhibit the activity of TYK2 are beneficial, especially those with selectivity over JAK2. Such compounds should deliver a pharmacological response that favorably treats

one or more of the conditions described herein without the side-effects associated with the inhibition of JAK2.

[0015] Accordingly there is a need to provide novel inhibitors having more effective or advantageous pharmaceutically relevant properties, like selectivity over other JAK kinases (especially JAK2).

BRIEF SUMMARY OF THE INVENTION

[0016] Disclosed herein is a compound of Formula (XII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (XII),

wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, heteroaryl;

R¹⁶ is -C(=O)NR¹R², -C(=N-CN)NR¹R², -P(=O)R¹R², or -C(=O)R¹¹;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

or -L-Ring A is absent;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a};

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;


each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0017] Also disclosed herein is a compound of Formula (XIII), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (XIII),

wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

or -L-Ring A is absent;

each X is independently -CR^x or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷,

$-\text{NR}^8\text{C}(=\text{O})\text{R}^7$, $-\text{NR}^8\text{C}(=\text{N}-\text{CN})\text{OR}^8$, $-\text{NR}^8\text{C}(=\text{O})\text{OR}^8$, $-\text{NR}^8\text{S}(=\text{O})(=\text{NR}^8)\text{R}^7$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ aminoalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ aminoalkyl, $\text{C}_1\text{-C}_6$ heteroalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, $\text{C}_1\text{-C}_6$ alkyl(cycloalkyl), $\text{C}_1\text{-C}_6$ alkyl(heterocycloalkyl), $\text{C}_1\text{-C}_6$ alkyl(aryl), or $\text{C}_1\text{-C}_6$ alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl;

R^7 is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ aminoalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{OMe}$, $-\text{NH}_2$, $-\text{C}(=\text{O})\text{Me}$, $-\text{C}(=\text{O})\text{OH}$, $-\text{C}(=\text{O})\text{OMe}$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl;

each R^8 is independently hydrogen, CN , $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ aminoalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{OMe}$, $-\text{NH}_2$, $-\text{C}(=\text{O})\text{Me}$, $-\text{C}(=\text{O})\text{OH}$, $-\text{C}(=\text{O})\text{OMe}$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl;

R^9 and R^{10} are independently hydrogen, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ aminoalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{OMe}$, $-\text{NH}_2$, $-\text{C}(=\text{O})\text{Me}$, $-\text{C}(=\text{O})\text{OH}$, $-\text{C}(=\text{O})\text{OMe}$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R^9 and R^{10} are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{OMe}$, $-\text{NH}_2$, $-\text{C}(=\text{O})\text{Me}$, $-\text{C}(=\text{O})\text{OH}$, $-\text{C}(=\text{O})\text{OMe}$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl;

or R^8 and R^9 are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OH}$, $-\text{OMe}$, $-\text{NH}_2$, $-\text{C}(=\text{O})\text{Me}$, $-\text{C}(=\text{O})\text{OH}$, $-\text{C}(=\text{O})\text{OMe}$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl;

R^{11} is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_1\text{-C}_6$ hydroxyalkyl, $\text{C}_1\text{-C}_6$ aminoalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a} ;

each R^{1a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0018] Also disclosed herein is a pharmaceutical composition comprising a therapeutically effective amount of the compound disclosed herein, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

[0019] Also disclosed herein is a method of inhibiting a TYK2 enzyme in a patient or biological sample comprising contacting said patient or biological sample with a compound disclosed herein, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

[0020] Also disclosed herein is a method of treating a TYK2-mediated disorder comprising administering to a patient in need thereof a compound disclosed herein, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof. In some embodiments, the TYK2-mediated disorder is an autoimmune disorder, an inflammatory disorder, a proliferative disorder, an endocrine disorder, a neurological disorder, or a disorder associated with transplantation. In some embodiments, the disorder is associated with type I interferon, IL-10, IL-12, or IL-23 signaling.

INCORPORATION BY REFERENCE

[0021] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference for the specific purposes identified herein.

DETAILED DESCRIPTION OF THE INVENTION**Definitions**

[0022] As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “an agent” includes a plurality of such agents, and reference to “the cell” includes reference to one or more cells (or to a plurality of cells) and equivalents thereof known to those skilled in the art, and so forth. When ranges are used herein for physical properties, such as molecular weight, or chemical properties, such as chemical formulae, all combinations and subcombinations of ranges and specific embodiments therein are intended to be included. The term “about” when referring to a number or a numerical range means that the number or numerical range referred to is an approximation within experimental variability (or within statistical experimental error), and thus the number or numerical range, in some instances, will vary between 1% and 15% of the stated number or numerical range. The term “comprising” (and related terms such as “comprise” or “comprises” or “having” or “including”) is not intended to exclude that in other certain embodiments, for example, an embodiment of any composition of matter, composition, method, or process, or the like, described herein, “consist of” or “consist essentially of” the described features.

[0023] As used in the specification and appended claims, unless specified to the contrary, the following terms have the meaning indicated below.

[0024] “Aliphatic chain” refers to a linear chemical moiety that is composed of only carbons and hydrogens. In some embodiments, the aliphatic chain is saturated. In some embodiments, the aliphatic chain is unsaturated. In some embodiments, the unsaturated aliphatic chain contains one unsaturation. In some embodiments, the unsaturated aliphatic chain contains more than one unsaturation. In some embodiments, the unsaturated aliphatic chain contains two unsaturations. In some embodiments, the unsaturated aliphatic chain contains one double bond. In some embodiments, the unsaturated aliphatic chain contains two double bonds.

[0025] “Alkyl” refers to an optionally substituted straight-chain, or optionally substituted branched-chain saturated hydrocarbon monoradical having from one to about ten carbon atoms, or from one to six carbon atoms. Examples include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl, tert-amyl and hexyl, and longer alkyl groups, such as heptyl, octyl, and the like. Whenever it appears herein, a numerical range such as “C₁-C₆ alkyl” means that the alkyl group consists of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, 4 carbon atoms, 5 carbon atoms or 6 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated. In some embodiments, the alkyl is a C₁-C₁₀ alkyl, a C₁-C₉ alkyl, a C₁-C₈ alkyl, a C₁-C₇ alkyl, a C₁-C₆ alkyl, a C₁-C₅ alkyl, a C₁-C₄ alkyl, a C₁-C₃ alkyl, a C₁-C₂ alkyl, or a C₁ alkyl. Unless stated otherwise specifically in the specification, an alkyl group is optionally substituted, for

example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, the alkyl is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, the alkyl is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, or -OMe. In some embodiments, the alkyl is optionally substituted with halogen.

[0026] “Alkenyl” refers to an optionally substituted straight-chain, or optionally substituted branched-chain hydrocarbon monoradical having one or more carbon-carbon double-bonds and having from two to about ten carbon atoms, more preferably two to about six carbon atoms. The group may be in either the *cis* or *trans* conformation about the double bond(s), and should be understood to include both isomers. Examples include, but are not limited to, ethenyl (-CH=CH₂), 1-propenyl (-CH₂CH=CH₂), isopropenyl [-C(CH₃)=CH₂], butenyl, 1,3-butadienyl and the like. Whenever it appears herein, a numerical range such as “C₂-C₆ alkenyl” means that the alkenyl group may consist of 2 carbon atoms, 3 carbon atoms, 4 carbon atoms, 5 carbon atoms or 6 carbon atoms, although the present definition also covers the occurrence of the term “alkenyl” where no numerical range is designated. In some embodiments, the alkenyl is a C₂-C₁₀ alkenyl, a C₂-C₉ alkenyl, a C₂-C₈ alkenyl, a C₂-C₇ alkenyl, a C₂-C₆ alkenyl, a C₂-C₅ alkenyl, a C₂-C₄ alkenyl, a C₂-C₃ alkenyl, or a C₂ alkenyl. Unless stated otherwise specifically in the specification, an alkenyl group is optionally substituted, for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, an alkenyl is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, an alkenyl is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, or -OMe. In some embodiments, the alkenyl is optionally substituted with halogen.

[0027] “Alkynyl” refers to an optionally substituted straight-chain or optionally substituted branched-chain hydrocarbon monoradical having one or more carbon-carbon triple-bonds and having from two to about ten carbon atoms, more preferably from two to about six carbon atoms. Examples include, but are not limited to, ethynyl, 2-propynyl, 2-butylnyl, 1,3-butadiynyl and the like. Whenever it appears herein, a numerical range such as “C₂-C₆ alkynyl” means that the alkynyl group may consist of 2 carbon atoms, 3 carbon atoms, 4 carbon atoms, 5 carbon atoms or 6 carbon atoms, although the present definition also covers the occurrence of the term “alkynyl” where no numerical range is designated. In some embodiments, the alkynyl is a C₂-C₁₀ alkynyl, a C₂-C₉ alkynyl, a C₂-C₈ alkynyl, a C₂-C₇ alkynyl, a C₂-C₆ alkynyl, a C₂-C₅ alkynyl, a C₂-C₄ alkynyl, a C₂-C₃ alkynyl, or a C₂ alkynyl. Unless stated otherwise specifically in the specification, an alkynyl group is optionally substituted, for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, an alkynyl is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, an alkynyl is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, or -OMe. In some embodiments, the alkynyl is optionally substituted with halogen.

[0028] “Alkylene” refers to a straight or branched divalent hydrocarbon chain. Unless stated otherwise specifically in the specification, an alkylene group may be optionally substituted, for example,

with oxo, halogen, amino, nitrile, nitro, hydroxyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, an alkylene is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, an alkylene is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, or -OMe. In some embodiments, the alkylene is optionally substituted with halogen.

[0029] “Alkoxy” refers to a radical of the formula -OR_a where R_a is an alkyl radical as defined.

Unless stated otherwise specifically in the specification, an alkoxy group may be optionally substituted, for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, an alkoxy is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, an alkoxy is optionally substituted with oxo, halogen, -CN, -CF₃, -OH, or -OMe. In some embodiments, the alkoxy is optionally substituted with halogen.

[0030] “Aminoalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more amines. In some embodiments, the alkyl is substituted with one amine. In some embodiments, the alkyl is substituted with one, two, or three amines. Hydroxyalkyl include, for example, aminomethyl, aminoethyl, aminopropyl, aminobutyl, or aminopentyl. In some embodiments, the hydroxyalkyl is aminomethyl.

[0031] “Aryl” refers to a radical derived from a hydrocarbon ring system comprising hydrogen, 6 to 30 carbon atoms and at least one aromatic ring. The aryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused (when fused with a cycloalkyl or heterocycloalkyl ring, the aryl is bonded through an aromatic ring atom) or bridged ring systems. In some embodiments, the aryl is a 6- to 10-membered aryl. In some embodiments, the aryl is a 6-membered aryl. Aryl radicals include, but are not limited to, aryl radicals derived from the hydrocarbon ring systems of anthrylene, naphthylene, phenanthrylene, anthracene, azulene, benzene, chrysene, fluoranthene, fluorene, as-indacene, s-indacene, indane, indene, naphthalene, phenalene, phenanthrene, pleiadene, pyrene, and triphenylene. In some embodiments, the aryl is phenyl. Unless stated otherwise specifically in the specification, an aryl may be optionally substituted, for example, with halogen, amino, nitrile, nitro, hydroxyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, an aryl is optionally substituted with halogen, methyl, ethyl, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, an aryl is optionally substituted with halogen, methyl, ethyl, -CN, -CF₃, -OH, or -OMe. In some embodiments, the aryl is optionally substituted with halogen.

[0032] “Cycloalkyl” refers to a stable, partially or fully saturated, monocyclic or polycyclic carbocyclic ring, which may include fused (when fused with an aryl or a heteroaryl ring, the cycloalkyl is bonded through a non-aromatic ring atom) or bridged ring systems. Representative cycloalkyls include, but are not limited to, cycloalkyls having from three to fifteen carbon atoms (C₃-C₁₅ cycloalkyl), from three to ten carbon atoms (C₃-C₁₀ cycloalkyl), from three to eight carbon atoms (C₃-C₈ cycloalkyl), from three to six carbon atoms (C₃-C₆ cycloalkyl), from three to five carbon atoms (C₃-C₅ cycloalkyl), or three to four carbon atoms (C₃-C₄ cycloalkyl). In some embodiments, the cycloalkyl is a 3- to 6-membered

cycloalkyl. In some embodiments, the cycloalkyl is a 5- to 6-membered cycloalkyl. Monocyclic cycloalkyls include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic cycloalkyls or carbocycles include, for example, adamantyl, norbornyl, decalinyl, bicyclo[3.3.0]octane, bicyclo[4.3.0]nonane, cis-decalin, trans-decalin, bicyclo[2.1.1]hexane, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, bicyclo[3.2.2]nonane, and bicyclo[3.3.2]decane, and 7,7-dimethyl-bicyclo[2.2.1]heptanyl. Partially saturated cycloalkyls include, for example cyclopentenyl, cyclohexenyl, cycloheptenyl, and cyclooctenyl. Unless stated otherwise specifically in the specification, a cycloalkyl is optionally substituted, for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, a cycloalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, a cycloalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, or -OMe. In some embodiments, the cycloalkyl is optionally substituted with halogen.

[0033] “Deuteroalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more deuterium atoms. In some embodiments, the alkyl is substituted with one deuterium atom. In some embodiments, the alkyl is substituted with one, two, or three deuterium atoms. In some embodiments, the alkyl is substituted with one, two, three, four, five, or six deuterium atoms. Deuteroalkyl includes, for example, CD₃, CH₂D, CHD₂, CH₂CD₃, CD₂CD₃, CHDCD₃, CH₂CH₂D, or CH₂CHD₂. In some embodiments, the deuteroalkyl is CD₃.

[0034] “Haloalkyl” refers to an alkyl radical, as defined above, that is substituted by one or more halogen atoms. In some embodiments, the alkyl is substituted with one, two, or three halogen atoms. In some embodiments, the alkyl is substituted with one, two, three, four, five, or six halogen atoms. Haloalkyl includes, for example, trifluoromethyl, difluoromethyl, fluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1,2-difluoroethyl, 3-bromo-2-fluoropropyl, 1,2-dibromoethyl, and the like. In some embodiments, the haloalkyl is trifluoromethyl.

[0035] “Halo” or “halogen” refers to bromo, chloro, fluoro or iodo. In some embodiments, halogen is fluoro or chloro. In some embodiments, halogen is fluoro.

[0036] “Heteroalkyl” refers to an alkyl group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, e.g., oxygen, nitrogen (e.g., -NH-, -N(alkyl)-), sulfur, or combinations thereof. A heteroalkyl is attached to the rest of the molecule at a carbon atom of the heteroalkyl. In one aspect, a heteroalkyl is a C₁-C₆ heteroalkyl wherein the heteroalkyl is comprised of 1 to 6 carbon atoms and one or more atoms other than carbon, e.g., oxygen, nitrogen (e.g. -NH-, -N(alkyl)-), sulfur, or combinations thereof wherein the heteroalkyl is attached to the rest of the molecule at a carbon atom of the heteroalkyl. Examples of such heteroalkyl are, for example, -CH₂OCH₃, -CH₂CH₂OCH₃, -CH₂CH₂OCH₂CH₂OCH₃, or -CH(CH₃)OCH₃. Unless stated otherwise specifically in the specification, a heteroalkyl is optionally substituted for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, a heteroalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -

CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, a heteroalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, or -OMe. In some embodiments, the heteroalkyl is optionally substituted with halogen.

[0037] "Hydroxyalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more hydroxyls. In some embodiments, the alkyl is substituted with one hydroxyl. In some embodiments, the alkyl is substituted with one, two, or three hydroxyls. Hydroxyalkyl include, for example, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, or hydroxypentyl. In some embodiments, the hydroxyalkyl is hydroxymethyl.

[0038] "Heterocycloalkyl" refers to a stable 3- to 24-membered partially or fully saturated ring radical comprising 2 to 23 carbon atoms and from one to 8 heteroatoms selected from the group consisting of nitrogen, oxygen, phosphorous and sulfur. In some embodiments, the heterocycloalkyl comprises 1 or 2 heteroatoms selected from nitrogen and oxygen. Unless stated otherwise specifically in the specification, the heterocycloalkyl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused (when fused with an aryl or a heteroaryl ring, the heterocycloalkyl is bonded through a non-aromatic ring atom) or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heterocycloalkyl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized. Representative heterocycloalkyls include, but are not limited to, heterocycloalkyls having from two to fifteen carbon atoms (C₂-C₁₅ heterocycloalkyl), from two to ten carbon atoms (C₂-C₁₀ heterocycloalkyl), from two to eight carbon atoms (C₂-C₈ heterocycloalkyl), from two to six carbon atoms (C₂-C₆ heterocycloalkyl), from two to five carbon atoms (C₂-C₅ heterocycloalkyl), or two to four carbon atoms (C₂-C₄ heterocycloalkyl). In some embodiments, the heterocycloalkyl is a 3- to 6-membered heterocycloalkyl. In some embodiments, the cycloalkyl is a 5- to 6-membered heterocycloalkyl. Examples of such heterocycloalkyl radicals include, but are not limited to, aziridinyl, azetidiny, dioxolanyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazoliny, imidazolidiny, isothiazolidiny, isoxazolidiny, morpholinyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidiny, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidiny, quinuclidiny, thiazolidiny, tetrahydrofuryl, trithianyl, tetrahydropyranly, thiomorpholinyl, thiamorpholinyl, 1-oxo-thiomorpholinyl, 1,1-dioxo-thiomorpholinyl, 1,3-dihydroisobenzofuran-1-yl, 3-oxo-1,3-dihydroisobenzofuran-1-yl, methyl-2-oxo-1,3-dioxol-4-yl, and 2-oxo-1,3-dioxol-4-yl. The term heterocycloalkyl also includes all ring forms of the carbohydrates, including but not limited to, the monosaccharides, the disaccharides and the oligosaccharides. It is understood that when referring to the number of carbon atoms in a heterocycloalkyl, the number of carbon atoms in the heterocycloalkyl is not the same as the total number of atoms (including the heteroatoms) that make up the heterocycloalkyl (i.e. skeletal atoms of the heterocycloalkyl ring). Unless stated otherwise specifically in the specification, a heterocycloalkyl is optionally substituted, for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, a heterocycloalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, a heterocycloalkyl is optionally substituted with

oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, or -OMe. In some embodiments, the heterocycloalkyl is optionally substituted with halogen.

[0039] “Heteroalkyl” refers to an alkyl group in which one or more skeletal atoms of the alkyl are selected from an atom other than carbon, e.g., oxygen, nitrogen (e.g. -NH-, -N(alkyl)-), sulfur, or combinations thereof. A heteroalkyl is attached to the rest of the molecule at a carbon atom of the heteroalkyl. In one aspect, a heteroalkyl is a C₁-C₆ heteroalkyl. Unless stated otherwise specifically in the specification, a Heteroalkyl is optionally substituted, for example, with oxo, halogen, amino, nitrile, nitro, hydroxyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, a heteroalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, a heteroalkyl is optionally substituted with oxo, halogen, methyl, ethyl, -CN, -CF₃, -OH, or -OMe. In some embodiments, the heteroalkyl is optionally substituted with halogen.

[0040] “Heteroaryl” refers to a 5- to 14-membered ring system radical comprising hydrogen atoms, one to thirteen carbon atoms, one to six heteroatoms selected from the group consisting of nitrogen, oxygen, phosphorous and sulfur, and at least one aromatic ring. The heteroaryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused (when fused with a cycloalkyl or heterocycloalkyl ring, the heteroaryl is bonded through an aromatic ring atom) or bridged ring systems; and the nitrogen, carbon or sulfur atoms in the heteroaryl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized. In some embodiments, the heteroaryl is a 5- to 10-membered heteroaryl. In some embodiments, the heteroaryl is a 5- to 6-membered heteroaryl. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzothiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl, benzofuranonyl, benzothienyl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnoliny, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indoliny, isoindoliny, isoquinolyl, indoliziny, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl, 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-1H-pyrrolyl, phenazinyl, phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, pyridazinyl, quinazoliny, quinoxaliny, quinoliny, quinuclidiny, isoquinoliny, tetrahydroquinoliny, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl (i.e., thienyl). Unless stated otherwise specifically in the specification, a heteroaryl is optionally substituted, for example, with halogen, amino, nitrile, nitro, hydroxyl, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, aryl, cycloalkyl, heterocycloalkyl, heteroaryl, and the like. In some embodiments, a heteroaryl is optionally substituted with halogen, methyl, ethyl, -CN, -CF₃, -OH, -OMe, -NH₂, or -NO₂. In some embodiments, a heteroaryl is optionally substituted with halogen, methyl, ethyl, -CN, -CF₃, -OH, or -OMe. In some embodiments, the heteroaryl is optionally substituted with halogen.

[0041] The terms “treat,” “prevent,” “ameliorate,” and “inhibit,” as well as words stemming therefrom, as used herein, do not necessarily imply 100% or complete treatment, prevention, amelioration, or inhibition. Rather, there are varying degrees of treatment, prevention, amelioration, and inhibition of which one of ordinary skill in the art recognizes as having a potential benefit or therapeutic effect. In this respect, the disclosed methods can provide any amount of any level of treatment, prevention, amelioration, or inhibition of the disorder in a mammal. For example, a disorder, including symptoms or conditions thereof, may be reduced by, for example, about 100%, about 90%, about 80%, about 70%, about 60%, about 50%, about 40%, about 30%, about 20%, or about 10%. Furthermore, the treatment, prevention, amelioration, or inhibition provided by the methods disclosed herein can include treatment, prevention, amelioration, or inhibition of one or more conditions or symptoms of the disorder, e.g., cancer or an inflammatory disease. Also, for purposes herein, “treatment,” “prevention,” “amelioration,” or “inhibition” encompass delaying the onset of the disorder, or a symptom or condition thereof.


[0042] The terms “effective amount” or “therapeutically effective amount,” as used herein, refer to a sufficient amount of a compound disclosed herein being administered which will relieve to some extent one or more of the symptoms of the disease or condition being treated, e.g., cancer or an inflammatory disease. In some embodiments, the result is a reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic uses is the amount of the composition comprising a compound disclosed herein required to provide a clinically significant decrease in disease symptoms. In some embodiments, an appropriate “effective” amount in any individual case is determined using techniques, such as a dose escalation study.

[0043] As used herein, the term “TYK2-mediated” disorders, diseases, and/or conditions as used herein means any disease or other deleterious condition in which TYK2 or a mutant thereof is known to play a role. Accordingly, another embodiment relates to treating or lessening the severity of one or more diseases in which TYK2, or a mutant thereof, is known to play a role. Such TYK2-mediated disorders include but are not limited to autoimmune disorders, inflammatory disorders, proliferative disorders, endocrine disorders, neurological disorders and disorders associated with transplantation.

Compounds

[0044] Described herein are compounds that are useful in treating a TYK2-mediated disorder. In some embodiments, the TYK2-mediated disorder is an autoimmune disorder, an inflammatory disorder, a proliferative disorder, an endocrine disorder, a neurological disorder, or a disorder associated with transplantation.

[0045] Disclosed herein is a compound of Formula (I), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (I),

wherein:

Z is a bond, $-\text{CR}^Z_2-$, or $-(\text{CR}^Z_2)_2-$;

each R^Z is independently hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl;

R^1 and R^2 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

R^3 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl;

R^4 is hydrogen, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{SR}^b$, $-\text{S}(=\text{O})\text{R}^a$, $-\text{S}(=\text{O})_2\text{R}^a$, $-\text{NO}_2$, $-\text{NR}^c\text{R}^d$, $-\text{NHS}(=\text{O})_2\text{R}^a$, $-\text{S}(=\text{O})_2\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{OC}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $-\text{OC}(=\text{O})\text{NR}^c\text{R}^d$, $-\text{NR}^b\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $-\text{NR}^b\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^b\text{C}(=\text{O})\text{OR}^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

or R^3 and R^4 are taken together to form an optionally substituted ring;

L is a bond or $-\text{C}(=\text{O})-$;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A ;

each R^A is independently deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{SR}^b$, $-\text{S}(=\text{O})\text{R}^a$, $-\text{S}(=\text{O})_2\text{R}^a$, $-\text{NO}_2$, $-\text{NR}^c\text{R}^d$, $-\text{NHS}(=\text{O})_2\text{R}^a$, $-\text{S}(=\text{O})_2\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{OC}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $-\text{OC}(=\text{O})\text{NR}^c\text{R}^d$, $-\text{NR}^b\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $-\text{NR}^b\text{C}(=\text{O})\text{R}^a$, $-\text{NR}^b\text{C}(=\text{O})\text{OR}^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

R^5 is halogen, $-\text{CN}$, $-\text{OR}^8$, $-\text{SR}^8$, $-\text{S}(=\text{O})\text{R}^7$, $-\text{S}(=\text{O})_2\text{R}^7$, $-\text{NO}_2$, $-\text{NR}^9\text{R}^{10}$, $-\text{NHS}(=\text{O})_2\text{R}^7$, $-\text{S}(=\text{O})_2\text{NR}^9\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^7$, $-\text{OC}(=\text{O})\text{R}^7$, $-\text{C}(=\text{O})\text{OR}^8$, $-\text{OC}(=\text{O})\text{OR}^8$, $-\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{OC}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{R}^7$, $-\text{NR}^8\text{C}(=\text{O})\text{OR}^8$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

R^7 is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl,

cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0046] In some embodiments of a compound of Formula (I), L is a bond. In some embodiments of a compound of Formula (I), L is -C(=O)-.


[0047] In some embodiments of a compound of Formula (I), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (I),

Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (I), Ring A is heteroaryl optionally substituted with one or more R^A.


[0048] In some embodiments of a compound of Formula (I), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (I), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (I), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (I), each R^A is independently C₁-C₆alkyl.

[0049] In some embodiments of a compound of Formula (I), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (I), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (I), R⁴ is -OR^b. In some embodiments of a compound of Formula (I), R⁴ is hydrogen.

[0050] In some embodiments of a compound of Formula (I), the compound is of Formula (Ia):



[0051] In some embodiments of a compound of Formula (I), the compound is of Formula (Ib):



[0052] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R³ is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R³ is hydrogen.

[0053] In some embodiments of a compound of Formula (I), (Ia), or (Ib), Z is a bond or -CH₂-. In some embodiments of a compound of Formula (I), (Ia), or (Ib), Z is -CH₂-. In some embodiments of a compound of Formula (I), (Ia), or (Ib), Z is a bond.

[0054] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R¹ and R² are independently hydrogen or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R¹ is hydrogen. In some embodiments of a

compound of Formula (I), (Ia), or (Ib), R² is C₁-C₆alkyl or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R² is C₁-C₆deuteroalkyl.

[0055] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0056] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0057] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, or aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0058] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁵ is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁵ is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.


[0059] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁷ is unsubstituted cycloalkyl.

[0060] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl,

cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0061] In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (I), (Ia), or (Ib), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0062] Also disclosed herein is a compound of Formula (II) or (II'), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



wherein:

R¹¹ is deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or

more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0063] In some embodiments of a compound of Formula (II) or (II'), L is a bond. In some embodiments of a compound of Formula (II) or (II'), L is -C(=O)-.

[0064] In some embodiments of a compound of Formula (II) or (II'), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (II) or (II'), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each


optionally substituted with one or more R^A . In some embodiments of a compound of Formula (II) or (II'), Ring A is heteroaryl optionally substituted with one or more R^A .

[0065] In some embodiments of a compound of Formula (II) or (II'), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II) or (II'), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II) or (II'), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (II) or (II'), each R^A is independently C₁-C₆alkyl.


[0066] In some embodiments of a compound of Formula (II) or (II'), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II) or (II'), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (II) or (II'), R⁴ is -OR^b. In some embodiments of a compound of Formula (II) or (II'), R⁴ is hydrogen.

[0067] In some embodiments of a compound of Formula (II) or (II'), X is -CH-. In some embodiments of a compound of Formula (II) or (II'), X is -N-.

[0068] In some embodiments of a compound of Formula (II) or (II'), the compound is of Formula (IIa) or (II'a):



[0069] In some embodiments of a compound of Formula (II) or (II'), the compound is of Formula (IIb) or (II'b):



[0070] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R³ is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R³ is hydrogen.

[0071] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹ and R² are independently hydrogen or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹ is hydrogen. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R² is C₁-C₆alkyl or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R² is C₁-C₆deuteroalkyl.

[0072] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0073] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0074] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, or aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0075] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁵ is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁵ is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0076] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe,


C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁷ is unsubstituted cycloalkyl.

[0077] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0078] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.


[0079] In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹¹ is deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹¹ is deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹¹ is halogen. In some embodiments of a compound of Formula (II), (II'), (IIa), (II'a), (IIb), or (II'b), R¹¹ is hydrogen.

[0080] Also disclosed herein is a compound of Formula (III), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:




Formula (III),

wherein:



Ring B is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; provided that



is not

each R^B is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or two R^B on the same carbon are taken together to form an oxo;

or two R^B on adjacent atoms are taken together to form a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each optionally substituted with one or more deuterium, oxo, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆deuteroalkyl, or C₁-C₆haloalkyl;

n is 0-4;

each Y is independently C or N;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl,

heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0081] In some embodiments of a compound of Formula (III), L is a bond. In some embodiments of a compound of Formula (III), L is -C(=O)-.

[0082] In some embodiments of a compound of Formula (III), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (III), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (III), Ring A is heteroaryl optionally substituted with one or more R^A.


[0083] In some embodiments of a compound of Formula (III), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (III), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some

embodiments of a compound of Formula (III), each R^A is independently halogen or C_1 - C_6 alkyl. In some embodiments of a compound of Formula (III), each R^A is independently C_1 - C_6 alkyl.


[0084] In some embodiments of a compound of Formula (III), R^4 is hydrogen, deuterium, halogen, -CN, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (III), R^4 is hydrogen or $-OR^b$. In some embodiments of a compound of Formula (III), R^4 is $-OR^b$. In some embodiments of a compound of Formula (III), R^4 is hydrogen.

[0085] In some embodiments of a compound of Formula (III), X is $-CH-$. In some embodiments of a compound of Formula (III), X is $-N-$.

[0086] In some embodiments of a compound of Formula (III), the compound is of Formula (IIIa):



[0087] In some embodiments of a compound of Formula (III), the compound is of Formula (IIIb):



[0088] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^3 is hydrogen or C_1 - C_6 alkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^3 is hydrogen.

[0089] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^5 is halogen, -CN, $-OR^8$, $-NR^9R^{10}$, $-C(=O)R^7$, $-C(=O)OR^8$, $-C(=O)NR^9R^{10}$, $-NR^8C(=O)NR^9R^{10}$, $-NR^8C(=O)R^7$, $-NR^8C(=O)OR^8$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

[0090] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^5 is $-OR^8$, $-NR^9R^{10}$, $-C(=O)R^7$, $-C(=O)OR^8$, $-C(=O)NR^9R^{10}$, $-NR^8C(=O)NR^9R^{10}$, $-NR^8C(=O)R^7$, $-NR^8C(=O)OR^8$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

[0091] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^5 is $-OR^8$, $-NR^9R^{10}$, $-NR^8C(=O)R^7$, or aryl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[0092] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^5 is $-OR^8$, $-NR^9R^{10}$, $-NR^8C(=O)R^7$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^5 is $-NR^8C(=O)R^7$. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^5 is aryl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[0093] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^7 is C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^7 is C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^7 is unsubstituted cycloalkyl.



[0094] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^8 is hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^8 is hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[0095] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^9 and R^{10} are independently hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), R^9 and R^{10} are independently hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl,

C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[0096] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), Ring B is a heterocycloalkyl or heteroaryl.

[0097] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb),



[0098] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), each R^B is independently hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl; or two R^B on the same carbon are taken together to form an oxo. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), each R^B is independently hydrogen, deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl; two R^B on the same carbon are taken together to form an oxo.


[0099] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), two R^B on adjacent atoms are taken together to form a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more deuterium, oxo, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆deuteroalkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), two R^B on adjacent atoms are taken together to form a heterocycloalkyl or heteroaryl; each optionally substituted with one or more deuterium, oxo, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆deuteroalkyl, or C₁-C₆haloalkyl.

[00100] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is 0. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is 1. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is 2. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is 0-2. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is

0 or 1. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is 1 or 2. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n is 1-3.

[00101] In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n' is 0. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n' is 1. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n' is 2. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n' is 0-2. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n' is 0 or 1. In some embodiments of a compound of Formula (III), (IIIa), or (IIIb), n' is 1 or 2.

[00102] Also disclosed herein is a compound of Formula (IV), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



wherein:

Ring C is a bicyclic ring system;

each R^C is independently hydrogen, oxo, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

m is 0-4;

R¹² is -C(=O)NR¹R² or -L¹-R¹³;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

L¹ is -O-, -NH-, or -N(CH₃)-;

R¹³ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl,

heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00103] In some embodiments of a compound of Formula (IV), L is a bond. In some embodiments of a compound of Formula (IV), L is -C(=O)-.

[00104] In some embodiments of a compound of Formula (IV), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (IV), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (IV), Ring A is heteroaryl optionally substituted with one or more R^A.


[00105] In some embodiments of a compound of Formula (IV), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (IV), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some

embodiments of a compound of Formula (IV), each R^A is independently halogen or C_1 - C_6 alkyl. In some embodiments of a compound of Formula (IV), each R^A is independently C_1 - C_6 alkyl.


[00106] In some embodiments of a compound of Formula (IV), R^4 is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (IV), R^4 is hydrogen or -OR^b. In some embodiments of a compound of Formula (IV), R^4 is -OR^b. In some embodiments of a compound of Formula (IV), R^4 is hydrogen.

[00107] In some embodiments of a compound of Formula (IV), X is -CH-. In some embodiments of a compound of Formula (IV), X is -N-.

[00108] In some embodiments of a compound of Formula (IV), the compound is of Formula (IVa):



[00109] In some embodiments of a compound of Formula (IV), the compound is of Formula (IVb):



[00110] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^{12} is -C(=O)NR¹R².

[00111] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^1 and R^2 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^1 and R^2 are independently hydrogen or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^1 is hydrogen. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^2 is C_1 - C_6 alkyl or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^2 is C_1 - C_6 deuteroalkyl.

[00112] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^{12} is -L¹-R¹³.

[00113] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), L¹ is -NH-. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), L¹ is -O- or -NH-.

[00114] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^{13} is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl.

[00115] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R^5 is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and

heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00116] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00117] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, or aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00118] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁵ is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁵ is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00119] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁷ is unsubstituted cycloalkyl.

[00120] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo,

deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00121] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.


[00122] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), Ring C is indole or benzimidazole.

[00123] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), each R^C is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl.

[00124] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), each R^C is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), each R^C is independently halogen or C₁-C₆alkyl.

[00125] In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 0. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 1. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 2. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 0 or 1. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 0-2. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 1 or 2. In some embodiments of a compound of Formula (IV), (IVa), or (IVb), m is 1-3.

[00126] Also disclosed herein is a compound of Formula (V), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (V)

wherein:

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -

$C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

or two R^D on adjacent atoms are taken together to form a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more deuterium, oxo, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 deuteroalkyl, or C_1 - C_6 haloalkyl;

r is 0-4;

each Y is independently C or N ;

R^1 and R^2 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

R^3 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl;

R^4 is hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

or R^3 and R^4 are taken together to form an optionally substituted ring;

L is a bond or $-C(=O)-$;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A ;

each R^A is independently deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

each X is independently $-CR^x-$ or $-N-$;

each R^x is independently hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00127] In some embodiments of a compound of Formula (V), L is a bond. In some embodiments of a compound of Formula (V), L is -C(=O)-.


[00128] In some embodiments of a compound of Formula (V), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (V), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (V), Ring A is heteroaryl optionally substituted with one or more R^A.

[00129] In some embodiments of a compound of Formula (V), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (V), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (V), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (V), each R^A is independently C₁-C₆alkyl.

[00130] In some embodiments of a compound of Formula (V), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (V), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (V), R⁴ is -OR^b. In some embodiments of a compound of Formula (V), R⁴ is hydrogen.


[00131] In some embodiments of a compound of Formula (V), each X is -N-. In some embodiments of a compound of Formula (V), each X is -CH-. In some embodiments of a compound of Formula (V), one X is -N- and the other is -CH-.

[00132] In some embodiments of a compound of Formula (V), the compound is of Formula (Va):



Formula (Va).

[00133] In some embodiments of a compound of Formula (V), the compound is of Formula (Vb):




Formula (Vb).

[00134] In some embodiments of a compound of Formula (V), (Va), or (Vb), R³ is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (V), (Va), or (Vb), R³ is hydrogen.


[00135] In some embodiments of a compound of Formula (V), (Va), or (Vb), R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (V), (Va), or (Vb), R¹ and R² are independently hydrogen or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (V), (Va), or (Vb), R¹ is hydrogen. In some embodiments of a compound of Formula (V), (Va), or (Vb), R² is C₁-C₆alkyl or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (V), (Va), or (Vb), R² is C₁-C₆deuteroalkyl.

[00136] In some embodiments of a compound of Formula (V), (Va), or (Vb), Ring D is a heterocycloalkyl or heteroaryl.

[00137] In some embodiments of a compound of Formula (V), (Va), or (Vb),



is




and r' is 0-3.

[00138] In some embodiments of a compound of Formula (V), (Va), or (Vb), each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, or heterocycloalkyl. In some embodiments of a compound of Formula (V), (Va), or (Vb), each R^D is independently hydrogen, deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl. In some embodiments of a compound of Formula (V), (Va), or (Vb), each R^D is independently hydrogen or cycloalkyl.

[00139] In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 0. In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 1. In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 2. In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 0-2. In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 0 or 1. In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 1 or 2. In some embodiments of a compound of Formula (V), (Va), or (Vb), r is 1-3.

[00140] In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 0. In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 1. In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 2. In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 0-2. In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 0 or 1. In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 1 or 2. In some embodiments of a compound of Formula (V), (Va), or (Vb), r¹ is 1-3.

[00141] Also disclosed herein is a compound of Formula (VI), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (VI)

wherein:

Ring E is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^E;

each R^E is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^E on the same carbon are taken together to form an oxo;

L² is a bond, -O-, or $\text{---}\equiv\text{---}$;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl,

heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl,

heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo,

deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or

C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -

NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -

NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl,

or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00142] In some embodiments of a compound of Formula (VI), L is a bond. In some embodiments of a compound of Formula (VI), L is -C(=O)-.


[00143] In some embodiments of a compound of Formula (VI), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (VI), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (VI), Ring A is heteroaryl optionally substituted with one or more R^A.

[00144] In some embodiments of a compound of Formula (VI), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VI), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VI), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (VI), each R^A is independently C₁-C₆alkyl.

[00145] In some embodiments of a compound of Formula (VI), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VI), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (VI), R⁴ is -OR^b. In some embodiments of a compound of Formula (VI), R⁴ is hydrogen.


[00146] In some embodiments of a compound of Formula (VI), X is -CH-. In some embodiments of a compound of Formula (VI), X is -N-.

[00147] In some embodiments of a compound of Formula (VI), the compound is of Formula (VIa):



Formula (VIa).

[00148] In some embodiments of a compound of Formula (VI), the compound is of Formula (VIb):



Formula (VIb).

[00149] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^3 is hydrogen or C_1 - C_6 alkyl. In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^3 is hydrogen.

[00150] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^1 and R^2 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^1 and R^2 are independently hydrogen or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^1 is hydrogen. In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^2 is C_1 - C_6 alkyl or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (VI), (VIa), or (VIb), R^2 is C_1 - C_6 deuteroalkyl.

[00151] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), L^2 is a bond.

[00152] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), Ring E is cycloalkyl, heterocycloalkyl, or aryl; each optionally substituted with one or more R^E .


[00153] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), Ring E is aryl optionally substituted with one or more R^E .

[00154] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), each R^E is independently deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl.

[00155] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), each R^E is independently deuterium, halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl.

[00156] In some embodiments of a compound of Formula (VI), (VIa), or (VIb), each R^E is independently halogen.

[00157] Also disclosed herein is a compound of Formula (VII), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (VII)

wherein:

R^{14} is hydrogen, $-S(=O)R^a$, $-S(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 -

C₆alkynyl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl),

C₁-C₆alkyl(heteroaryl), cycloalkyl, or heterocycloalkyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl,

heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl,

heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo,

deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or

C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -

NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -

C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -

NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl,

or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl

is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -

NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00158] In some embodiments of a compound of Formula (VII), L is a bond. In some embodiments of a compound of Formula (VII), L is -C(=O)-.


[00159] In some embodiments of a compound of Formula (VII), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (VII), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (VII), Ring A is heteroaryl optionally substituted with one or more R^A.

[00160] In some embodiments of a compound of Formula (VII), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VII), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VII), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (VII), each R^A is independently C₁-C₆alkyl.

[00161] In some embodiments of a compound of Formula (VII), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VII), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (VII), R⁴ is -OR^b. In some embodiments of a compound of Formula (VII), R⁴ is hydrogen.


[00162] In some embodiments of a compound of Formula (VII), X is -CH-. In some embodiments of a compound of Formula (VII), X is -N-.

[00163] In some embodiments of a compound of Formula (VII), the compound is of Formula (VIIa):



Formula (VIIa).

[00164] In some embodiments of a compound of Formula (VII), the compound is of Formula (VIIb):



Formula (VIIb).


- [00165]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^3 is hydrogen or C_1 - C_6 alkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^3 is hydrogen.
- [00166]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^5 is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.
- [00167]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^5 is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.
- [00168]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^5 is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, or aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.
- [00169]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^5 is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^5 is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^5 is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.
- [00170]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^7 is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^7 is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^7 is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^7 is unsubstituted cycloalkyl.
- [00171]** In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R^8 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein

each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00172] In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00173] In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R¹⁴ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or C₁-C₆alkyl(cycloalkyl). In some embodiments of a compound of Formula (VII), (VIIa), or (VIIb), R¹⁴ is C₁-C₆alkyl or C₁-C₆alkyl(cycloalkyl).

[00174] Also disclosed herein is a compound of Formula (VIII), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (VIII)

wherein:

R¹⁵ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

L³ is -O-, -NH-, or -N(CH₃)-;

L⁴ is -NR³- or -C(=O)-;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;
or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-

C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00175] In some embodiments of a compound of Formula (VIII), L is a bond. In some embodiments of a compound of Formula (VIII), L is -C(=O)-.


[00176] In some embodiments of a compound of Formula (VIII), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (VIII), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (VIII), Ring A is heteroaryl optionally substituted with one or more R^A.

[00177] In some embodiments of a compound of Formula (VIII), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VIII), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VIII), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (VIII), each R^A is independently C₁-C₆alkyl.


[00178] In some embodiments of a compound of Formula (VIII), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (VIII), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (VIII), R⁴ is -OR^b. In some embodiments of a compound of Formula (VIII), R⁴ is hydrogen.

[00179] In some embodiments of a compound of Formula (VIII), X is -CH-. In some embodiments of a compound of Formula (VIII), X is -N-.

[00180] In some embodiments of a compound of Formula (VIII), the compound is of Formula (VIIIa):



[00181] In some embodiments of a compound of Formula (VIII), the compound is of Formula (VIIIb):



[00182] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and

heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00183] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00184] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, or aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00185] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁵ is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁵ is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00186] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁷ is unsubstituted cycloalkyl.

[00187] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo,

deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00188] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00189] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), L³ is -O-. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), L³ is -NH-.


[00190] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R¹⁵ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl.

[00191] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), L⁴ is -NR³-.

[00192] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R³ is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), R³ is hydrogen.

[00193] In some embodiments of a compound of Formula (VIII), (VIIIa), or (VIIIb), L⁴ is -C(=O)-.

[00194] A compound of Formula (IX), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (IX),

wherein:

Ring F is a heterocycloalkyl or heteroaryl;

each R^F is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or two R^F on the same carbon are taken together to form an oxo;

or two R^F on adjacent atoms are taken together to form a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more deuterium, oxo, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆deuteroalkyl, or C₁-C₆haloalkyl;

p is 0-4;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl,

cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d,

C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00195] In some embodiments of a compound of Formula (IX), L is a bond. In some embodiments of a compound of Formula (IX), L is -C(=O)-.


[00196] In some embodiments of a compound of Formula (IX), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (IX), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (IX), Ring A is heteroaryl optionally substituted with one or more R^A.

[00197] In some embodiments of a compound of Formula (IX), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (IX), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (IX), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (IX), each R^A is independently C₁-C₆alkyl.

[00198] In some embodiments of a compound of Formula (IX), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (IX), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (IX), R⁴ is -OR^b. In some embodiments of a compound of Formula (IX), R⁴ is hydrogen.


[00199] In some embodiments of a compound of Formula (IX), X is -CH-. In some embodiments of a compound of Formula (IX), X is -N-.

[00200] In some embodiments of a compound of Formula (IX), the compound is of Formula (IXa):



Formula (IXa).

[00201] In some embodiments of a compound of Formula (IX), the compound is of Formula (IXb):



Formula (IXb).

[00202] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R³ is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R³ is hydrogen.

[00203] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00204] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00205] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, or aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00206] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁵ is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁵ is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.


[00207] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN,

-OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁷ is unsubstituted cycloalkyl.


[00208] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁸ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00209] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00210] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), Ring F is a heterocycloalkyl.



[00211] In some embodiments of a compound of Formula (IX), (IXa), or (IXb),




[00212] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), each R^F is independently hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), each R^F is independently hydrogen, deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl.

[00213] In some embodiments of a compound of Formula (IX), (IXa), or (IXb), p is 0. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), p is 1. In some embodiments of a

compound of Formula (IX), (IXa), or (IXb), p is 2. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), p is 0-2. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), p is 0 or 1. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), p is 1 or 2. In some embodiments of a compound of Formula (IX), (IXa), or (IXb), p is 1-3.

[00214] Also disclosed herein is a compound of Formula (X), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (X),

wherein:

L^5 is a saturated or unsaturated linear aliphatic chain having 1-10 carbon atoms optionally substituted with one or more R^{L5} , wherein 1-5 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, -S(=O)₂-, or -P(=O)-;

each R^{L5} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or two R^{L5} on the same carbon atom are taken together to form an oxo;

R^{12} is -C(=O)NR¹R² or -L¹-R¹³;

R^1 and R^2 are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

L^1 is -O-, -NH-, or -N(CH₃)-;

R^{13} is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R^3 is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R^4 is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R^3 and R^4 are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A ;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00215] In some embodiments of a compound of Formula (X), L is a bond. In some embodiments of a compound of Formula (X), L is -C(=O)-.

[00216] In some embodiments of a compound of Formula (X), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (X), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally


substituted with one or more R^A . In some embodiments of a compound of Formula (X), Ring A is heteroaryl optionally substituted with one or more R^A .

[00217] In some embodiments of a compound of Formula (X), each R^A is independently deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, C_1-C_6 alkyl, C_1-C_6 haloalkyl, or C_1-C_6 deuteroalkyl. In some embodiments of a compound of Formula (X), each R^A is independently deuterium, halogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, or C_1-C_6 deuteroalkyl. In some embodiments of a compound of Formula (X), each R^A is independently halogen or C_1-C_6 alkyl. In some embodiments of a compound of Formula (X), each R^A is independently C_1-C_6 alkyl.


[00218] In some embodiments of a compound of Formula (X), R^4 is hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, C_1-C_6 alkyl, C_1-C_6 haloalkyl, or C_1-C_6 deuteroalkyl. In some embodiments of a compound of Formula (X), R^4 is hydrogen or $-OR^b$. In some embodiments of a compound of Formula (X), R^4 is $-OR^b$. In some embodiments of a compound of Formula (X), R^4 is hydrogen.

[00219] In some embodiments of a compound of Formula (X), X is $-CH-$. In some embodiments of a compound of Formula (X), X is $-N-$.

[00220] In some embodiments of a compound of Formula (X), the compound is of Formula




. In some embodiments of a compound of Formula (X), the compound is of




Formula

[00221] In some embodiments of a compound of Formula (X), the compound is of Formula (Xa):




Formula (Xa).

[00222] In some embodiments of a compound of Formula (X), the compound is of Formula (Xb):




Formula (Xb).

[00223] In some embodiments of a compound of Formula (X), the compound is of Formula (Xa-1):



Formula (Xa-1).

[00224] In some embodiments of a compound of Formula (X), the compound is of Formula (Xb-1):



Formula (Xb-1).

[00225] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R³ is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R³ is hydrogen.

[00226] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R¹² is -C(=O)NR¹R².

[00227] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R¹ and R² are independently hydrogen or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R¹ is hydrogen. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R² is C₁-C₆alkyl or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R² is C₁-C₆deuteroalkyl.

[00228] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R¹² is -L¹-R¹³.

[00229] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L¹ is -NH-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L¹ is -O- or -NH-.

[00230] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), R¹³ is C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl.

[00231] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is a saturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L5}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, -S(=O)₂-, or -P(=O)-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is a saturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L5}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is a saturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L5}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.

[00232] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is a saturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, or -S(=O)₂-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is a saturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is a saturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.

[00233] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is an unsaturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L5}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, or -S(=O)₂-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is an unsaturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L5}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is an unsaturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L5}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.


[00234] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is an unsaturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, -S(=O)₂-, or -P(=O)-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is an unsaturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is an unsaturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.

[00235] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), each R^{L5} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d,


C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), each R^{L5} is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), each R^{L5} is independently deuterium or halogen.

[00236] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), two R^{L5} on the same carbon atom are taken together to form an oxo.

[00237] In some embodiments of a compound of Formula (X), (Xa), (Xa-1), (Xb), or (Xb-1), L⁵ is



[00238] Also disclosed herein is a compound of Formula (XI), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (XI),

wherein:

L⁶ is a saturated or unsaturated linear aliphatic chain having 1-10 carbon atoms optionally substituted with one or more R^{L6}, wherein 1-5 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, -S(=O)₂-, or -P(=O)-;

each R^{L6} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or two R^{L6} on the same carbon atom are taken together to form an oxo;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A,

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl,

heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

X is -CR^x- or -N-;

R^x is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is hydrogen, halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=O)R⁷, -OC(=O)R⁷, -C(=O)OR⁸, -OC(=O)OR⁸, -C(=O)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00239] In some embodiments of a compound of Formula (XI), L is a bond. In some embodiments of a compound of Formula (XI), L is -C(=O)-.

[00240] In some embodiments of a compound of Formula (XI), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (XI), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (XI), Ring A is heteroaryl optionally substituted with one or more R^A.


[00241] In some embodiments of a compound of Formula (XI), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XI), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some

embodiments of a compound of Formula (XI), each R^A is independently halogen or C_1-C_6 alkyl. In some embodiments of a compound of Formula (XI), each R^A is independently C_1-C_6 alkyl.


[00242] In some embodiments of a compound of Formula (XI), R^4 is hydrogen, deuterium, halogen, -CN, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, C_1-C_6 alkyl, C_1-C_6 haloalkyl, or C_1-C_6 deuteroalkyl. In some embodiments of a compound of Formula (XI), R^4 is hydrogen or $-OR^b$. In some embodiments of a compound of Formula (XI), R^4 is $-OR^b$. In some embodiments of a compound of Formula (XI), R^4 is hydrogen.

[00243] In some embodiments of a compound of Formula (XI), X is $-CH-$. In some embodiments of a compound of Formula (XI), X is $-N-$.

[00244] In some embodiments of a compound of Formula (XI), the compound is of Formula (XIa):



[00245] In some embodiments of a compound of Formula (XI), the compound is of Formula (XIb):



[00246] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^3 is hydrogen or C_1-C_6 alkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^3 is hydrogen.

[00247] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^5 is halogen, -CN, $-OR^8$, $-NR^9R^{10}$, $-C(=O)R^7$, $-C(=O)OR^8$, $-C(=O)NR^9R^{10}$, $-NR^8C(=O)NR^9R^{10}$, $-NR^8C(=O)R^7$, $-NR^8C(=O)OR^8$, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[00248] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^5 is $-OR^8$, $-NR^9R^{10}$, $-C(=O)R^7$, $-C(=O)OR^8$, $-C(=O)NR^9R^{10}$, $-NR^8C(=O)NR^9R^{10}$, $-NR^8C(=O)R^7$, $-NR^8C(=O)OR^8$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[00249] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^5 is $-OR^8$, $-NR^9R^{10}$, $-NR^8C(=O)R^7$, or aryl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[00250] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^5 is $-OR^8$, $-NR^9R^{10}$, $-NR^8C(=O)R^7$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^5 is $-NR^8C(=O)R^7$. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^5 is aryl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[00251] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^7 is C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^7 is C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^7 is unsubstituted cycloalkyl.

[00252] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^8 is hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^8 is hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl.

[00253] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^9 and R^{10} are independently hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1-C_6 alkyl, or C_1-C_6 haloalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), R^9 and R^{10} are independently hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl,

C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00254] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is a saturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L6}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, or -S(=O)₂-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is a saturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L6}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is a saturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L6}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.

[00255] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is a saturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, or -S(=O)₂-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is a saturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is a saturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.

[00256] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is an unsaturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L6}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, -S(=O)₂-, or -P(=O)-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is an unsaturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L6}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is an unsaturated linear aliphatic chain having 1-8 carbon atoms optionally substituted with one or more R^{L6}, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.


[00257] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is an unsaturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, -O-, -S-, -S(=O)-, -S(=O)₂-, or -P(=O)-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is an unsaturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH-, -N(CH₃)-, or -O-. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is an unsaturated linear aliphatic chain having 1-8 carbon atoms, wherein 1, 2, or 3 carbon atoms are optionally replaced with -NH- or -O-.

[00258] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), each R^{L6} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), each R^{L6} is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or


C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XI), (XIa), or (XIb), each R^{L6} is independently deuterium or halogen.

[00259] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), two R^{L6} on the same carbon atom are taken together to form an oxo.

[00260] In some embodiments of a compound of Formula (XI), (XIa), or (XIb), L⁶ is



[00261] Also disclosed herein is a compound of Formula (XII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (XII),

wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, heteroaryl;

R¹⁶ is -C(=O)NR¹R², -C(=N-CN)NR¹R², -P(=O)R¹R², or -C(=O)R¹¹;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl,

C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-

C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more

R^A,

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -

NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -

OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl,

heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl,

heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo,

deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or

C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

or -L-Ring A is absent;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl

optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a},

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl,

C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;


wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl

is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;
 or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.
[00262] Also disclosed herein is a compound of Formula (XII), or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof:



Formula (XII),

wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, heteroaryl;

R¹⁶ is -C(=O)NR¹R², -C(=N-CN)NR¹R², -P(=O)R¹R², or -C(=O)R¹¹;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

or -L-Ring A is absent;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a};

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;


each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00263] Also disclosed herein is a compound of Formula (XII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, heteroaryl;

R^{16} is $-C(=O)NR^1R^2$, $-C(=N-CN)NR^1R^2$, $-P(=O)R^1R^2$, or $-C(=O)R^{11}$;

R^1 and R^2 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl,

C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

R^3 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl;

R^4 is hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$,

$-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$,

$-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, $-P(=O)R^bR^b$, C_1 - C_6 alkyl,

C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 -

C_6 alkynyl;

or R^3 and R^4 are taken together to form an optionally substituted ring;

L is a bond or $-C(=O)-$;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A ;

each R^A is independently deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$,

$-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$,

$-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl,

C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl,

heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl,

heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo,

deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or

C_1 - C_6 haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

each X is independently $-CR^x-$ or $-N-$;

each R^x is independently hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$,

$-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$,

$-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl,

C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 -

C_6 alkynyl;

R^5 is halogen, $-CN$, $-OR^8$, $-SR^8$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NO_2$, $-NR^9R^{10}$, $-NR^8S(=O)R^7$, $-NR^8S(=O)_2R^7$,

$-S(=O)_2NR^9R^{10}$, $-C(=N-CN)R^7$, $-C(=O)R^7$, $-OC(=N-CN)R^7$, $-OC(=O)R^7$, $-C(=N-CN)OR^8$, $-C(=O)OR^8$,

-OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe,

-NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a};

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;


each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00264] Also disclosed herein is a compound of Formula (XII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, heteroaryl;

R¹⁶ is -C(=O)NR¹R², -C(=N-CN)NR¹R², -P(=O)R¹R², or -C(=O)R¹¹;

R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl,

alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a};

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;


each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00265] In some embodiments of a compound of Formula (XII), Ring B is aryl or heteroaryl. In some embodiments of a compound of Formula (XII), Ring B is aryl. In some embodiments of a compound of Formula (XII), Ring B is phenyl. In some embodiments of a compound of Formula (XII), Ring B is heteroaryl. In some embodiments of a compound of Formula (XII), Ring B is 5- or 6-membered heteroaryl. In some embodiments of a compound of Formula (XII), Ring B is 6-membered heteroaryl. In some embodiments of a compound of Formula (XII), Ring B is pyridyl.

[00266] Also disclosed herein is a compound of Formula (XII'), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (XII'),

wherein:

R^{16} is $-C(=O)NR^1R^2$, $-C(=N-CN)NR^1R^2$, $-P(=O)R^1R^2$, or $-C(=O)R^{11}$;

R^1 and R^2 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl,

C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

R^3 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl;

R^4 is hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, $-P(=O)R^bR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

or R^3 and R^4 are taken together to form an optionally substituted ring;

L is a bond or $-C(=O)-$;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A ;

each R^A is independently deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

each X is independently $-CR^x-$ or $-N-$;

each R^x is independently hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

R^5 is halogen, $-CN$, $-OR^8$, $-SR^8$, $-S(=O)R^7$, $-S(=O)_2R^7$, $-NO_2$, $-NR^9R^{10}$, $-NR^8S(=O)R^7$, $-NR^8S(=O)_2R^7$, $-S(=O)_2NR^9R^{10}$, $-C(=N-CN)R^7$, $-C(=O)R^7$, $-OC(=N-CN)R^7$, $-OC(=O)R^7$, $-C(=N-CN)OR^8$, $-C(=O)OR^8$, $-OC(=N-CN)OR^8$, $-OC(=O)OR^8$, $-C(=N-CN)NR^9R^{10}$, $-C(=O)NR^9R^{10}$, $-OC(=N-CN)NR^9R^{10}$, $-OC(=O)NR^9R^{10}$, $-NR^8C(=N-CN)NR^9R^{10}$, $-NR^8C(=O)NR^9R^{10}$, $-NR^8C(=N-CN)R^7$, $-NR^8C(=N-OH)R^7$, $-NR^8C(=O)R^7$, $-NR^8C(=N-CN)OR^8$, $-NR^8C(=O)OR^8$, $-NR^8S(=O)(=NR^8)R^7$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl,

C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R^{11} is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a} ,

each R^{11a} is independently deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;

each R^a is independently C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

each R^b is independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

each R^c and R^d is independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

[00267] In some embodiments of a compound of Formula (XII) or (XII'), L is a bond. In some embodiments of a compound of Formula (XII) or (XII'), L is $-C(=O)-$.


[00268] In some embodiments of a compound of Formula (XII) or (XII'), $-L$ -Ring A is absent.

[00269] In some embodiments of a compound of Formula (XII) or (XII'), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A . In some embodiments of a compound of Formula (XII) or (XII'), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A . In some embodiments of a compound of Formula (XII) or (XII'), Ring A is heteroaryl optionally substituted with one or more R^A .

[00270] In some embodiments of a compound of Formula (XII) or (XII'), each R^A is independently deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl. In some embodiments of a compound of Formula (XII) or (XII'), each R^A is independently deuterium, halogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or


C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII) or (XII'), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (XII) or (XII'), each R^A is independently C₁-C₆alkyl.

[00271] In some embodiments of a compound of Formula (XII), the compound is of Formula (XIIa):



Formula (XIIa).




[00272] In some embodiments of a compound of Formula (XII), the compound is of Formula (XIIb):



Formula (XIIb).




[00273] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R⁴ is hydrogen or -OR^b. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R⁴ is -OR^b. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R⁴ is hydrogen. In some embodiments of a compound of Formula (XII), R⁴ is -P(=O)R^bR^b. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R⁴ is -S(=O)₂R^a.

[00274] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each X is -N-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc) each X is -CR^X-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), two X are -N- and the other is -CR^X-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), one X is -N- and the others are -CR^X-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each X is -CH-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), two X are -N- and the other is -CH-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), one X is -N- and the others are -CH-. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc),



[00275] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^X is independently hydrogen, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, C_1-C_6 hydroxyalkyl, or C_1-C_6 aminoalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^X is independently hydrogen, deuterium, halogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, or C_1-C_6 deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^X is independently hydrogen, deuterium, or halogen.

[00276] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc),



[00277] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^3 is hydrogen or C_1-C_6 alkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^3 is hydrogen.

[00278] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^{16} is $-C(=O)NR^1R^2$ or $-C(=O)R^{11}$. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^{16} is $-C(=O)NR^1R^2$. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^{16} is $-C(=N-CN)NR^1R^2$. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^{16} is $-P(=O)R^1R^2$. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R^{16} is $-C(=O)R^{11}$.

[00279] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹ and R² are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹ and R² are independently hydrogen or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹ and R² are independently C₁-C₆alkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹ is hydrogen. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R² is C₁-C₆alkyl or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R² is C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R² is C₁-C₆alkyl.

[00280] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, cycloalkyl, or heterocycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, or heterocycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or cycloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, or cycloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is cycloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), R¹¹ is C₁-C₆alkyl, C₁-C₆deuteroalkyl, or cycloalkyl.

[00281] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, or C₁-C₆aminoalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, or C₁-C₆aminoalkyl. In some embodiments of a compound of Formula (XII), (XII'),

(XIIa), (XIIb), or (XIIc), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, or C₁-C₆alkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^{11a} is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), (XIIb), or (XIIc), each R^{11a} is independently deuterium, halogen, or C₁-C₆alkyl.

[00282] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NHS(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00283] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00284] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00285] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -

$\text{NR}^8\text{C}(=\text{O})\text{OR}^8$, $-\text{NR}^8\text{S}(=\text{O})(=\text{NR}^8)\text{R}^7$, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ haloalkyl, $\text{C}_1\text{-C}_6$ deuteroalkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl.

[00286] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{OR}^8$, $-\text{NR}^9\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^7$, $-\text{C}(=\text{O})\text{OR}^8$, $-\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{N-CN})\text{R}^7$, $-\text{NR}^8\text{C}(=\text{O})\text{R}^7$, $-\text{NR}^8\text{C}(=\text{O})\text{OR}^8$, $-\text{NR}^8\text{S}(=\text{O})(=\text{NR}^8)\text{R}^7$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl.

[00287] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{OR}^8$, $-\text{NR}^9\text{R}^{10}$, $-\text{C}(=\text{O})\text{R}^7$, $-\text{C}(=\text{O})\text{OR}^8$, $-\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{N-CN})\text{R}^7$, $-\text{NR}^8\text{C}(=\text{O})\text{OR}^8$, $-\text{NR}^8\text{S}(=\text{O})(=\text{NR}^8)\text{R}^7$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{OR}^8$, $-\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{R}^7$, $-\text{NR}^8\text{C}(=\text{N-CN})\text{R}^7$, $-\text{NR}^8\text{S}(=\text{O})(=\text{NR}^8)\text{R}^7$, or aryl optionally substituted with one or more deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl.

[00288] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{OR}^8$, $-\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{N-CN})\text{R}^7$, $-\text{NR}^8\text{S}(=\text{O})(=\text{NR}^8)\text{R}^7$, or aryl optionally substituted with one or more deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl.

[00289] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{OR}^8$, $-\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{R}^7$, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is aryl optionally substituted with one or more oxo, deuterium, halogen, $-\text{CN}$, $-\text{OR}^b$, $-\text{NR}^c\text{R}^d$, $-\text{C}(=\text{O})\text{R}^a$, $-\text{C}(=\text{O})\text{OR}^b$, $-\text{C}(=\text{O})\text{NR}^c\text{R}^d$, $\text{C}_1\text{-C}_6$ alkyl, or $\text{C}_1\text{-C}_6$ haloalkyl.

[00290] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{NR}^9\text{R}^{10}$, $-\text{NR}^8\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$, or $-\text{NR}^8\text{C}(=\text{N-CN})\text{R}^7$.

[00291] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{NR}^9\text{R}^{10}$.

[00292] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R^5 is $-\text{NR}^9\text{R}^{10}$ or $-\text{NR}^8\text{C}(=\text{O})\text{NR}^9\text{R}^{10}$.

[00293] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is -NR⁸C(=N-CN)R⁷.

[00294] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is -NR⁸S(=O)(=NR⁸)R⁷.

[00295] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is -NR⁸C(=O)NR⁹R¹⁰ or -NR⁸C(=O)R⁷.

[00296] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is -NR⁸C(=O)NR⁹R¹⁰.

[00297] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is -NR⁸C(=O)R⁷.

[00298] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is not -NR⁸C(=O)R⁷.

[00299] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is -NR⁸C(=O)NR⁹R¹⁰ or heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.


[00300] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00301] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), or C₁-C₆alkyl(heterocycloalkyl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.


[00302] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁵ is heterocycloalkyl optionally substituted with one or more oxo, halogen, -OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl,

C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, C₁-C₆alkyl(cycloalkyl), or C₁-C₆alkyl(heterocycloalkyl); wherein each alkyl, alkynyl, cycloalkyl, heterocycloalkyl, and aryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00303] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), the



[00304] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), the




[00305] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is C₁-C₆alkyl, cycloalkyl, or heterocycloalkyl; wherein each alkyl, cycloalkyl, or heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is C₁-C₆alkyl or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is unsubstituted cycloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁷ is unsubstituted heterocycloalkyl.

[00306] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁸ is hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁸ is hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00307] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or C₁-C₆hydroxyalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or C₁-C₆hydroxyalkyl.

[00308] In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a C₂-C₈heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a bicyclic heterocycloalkyl or a spiro heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XII), (XII'), (XIIa), or (XIIb), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl selected from aziridinyl, azetidyl, pyrrolidinyl, piperidinyl, 2-azabicyclo[1.1.1]pentanyl, or 2-azaspiro[3.3]heptanyl, each optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00309] In some embodiments of a compound of Formula (XII), the compound is of Formula (XIIc):




Formula (XIIc);

wherein R^{5a} is deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00310] In some embodiments of a compound of Formula (XIIc); R^{5a} is deuterium, halogen, -CN, -OR^b, -NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00311] In some embodiments of a compound of Formula (XIIc); R^{5a} is halogen, -OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, C₁-C₆alkyl(cycloalkyl), or C₁-C₆alkyl(heterocycloalkyl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, and aryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00312] Also disclosed herein is a compound of Formula (XIII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (XIII),

wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

or -L-Ring A is absent;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl,

C_1-C_6 alkyl(cycloalkyl), C_1-C_6 alkyl(heterocycloalkyl), C_1-C_6 alkyl(aryl), or C_1-C_6 alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1-C_6 alkyl, or C_1-C_6 haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 aminoalkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C_1-C_6 alkyl, or C_1-C_6 haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 aminoalkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1-C_6 alkyl, or C_1-C_6 haloalkyl;

each R⁸ is independently hydrogen, CN, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 aminoalkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1-C_6 alkyl, or C_1-C_6 haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C_1-C_6 alkyl, C_1-C_6 haloalkyl, C_1-C_6 deuteroalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 aminoalkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1-C_6 alkyl, C_1-C_6 deuteroalkyl, C_1-C_6 haloalkyl, C_1-C_6 hydroxyalkyl, C_1-C_6 hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1-C_6 alkyl, or C_1-C_6 haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C_1-C_6 alkyl, or C_1-C_6 haloalkyl;

R^{11} is C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a} ,

each R^{11a} is independently deuterium, halogen, $-CN$, $-OR^b$, $-SR^b$, $-S(=O)R^a$, $-S(=O)_2R^a$, $-NO_2$, $-NR^cR^d$, $-NHS(=O)_2R^a$, $-S(=O)_2NR^cR^d$, $-C(=O)R^a$, $-OC(=O)R^a$, $-C(=O)OR^b$, $-OC(=O)OR^b$, $-C(=O)NR^cR^d$, $-OC(=O)NR^cR^d$, $-NR^bC(=O)NR^cR^d$, $-NR^bC(=O)R^a$, $-NR^bC(=O)OR^b$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, or C_2 - C_6 alkynyl;


each R^a is independently C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

each R^b is independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

each R^c and R^d is independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OH$, $-OMe$, $-NH_2$, $-C(=O)Me$, $-C(=O)OH$, $-C(=O)OMe$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

[00313] Also disclosed herein is a compound of Formula (XIII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



Formula (XIII),

wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

R^3 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, or C_1 - C_6 deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

L is a bond or -C(=O)-;

Ring A is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; each optionally substituted with one or more R^A;

each R^A is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^A on the same carbon are taken together to form an oxo;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is

independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl,

alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a};

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;


each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00314] Also disclosed herein is a compound of Formula (XIII), or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof:



wherein:

Ring B is cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

R³ is hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl;

R⁴ is hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, -P(=O)R^bR^b, C₁-C₆alkyl,

C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

or R³ and R⁴ are taken together to form an optionally substituted ring;

each X is independently -CR^x- or -N-;

each R^x is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

R⁵ is halogen, -CN, -OR⁸, -SR⁸, -S(=O)R⁷, -S(=O)₂R⁷, -NO₂, -NR⁹R¹⁰, -NR⁸S(=O)R⁷, -NR⁸S(=O)₂R⁷, -S(=O)₂NR⁹R¹⁰, -C(=N-CN)R⁷, -C(=O)R⁷, -OC(=N-CN)R⁷, -OC(=O)R⁷, -C(=N-CN)OR⁸, -C(=O)OR⁸, -OC(=N-CN)OR⁸, -OC(=O)OR⁸, -C(=N-CN)NR⁹R¹⁰, -C(=O)NR⁹R¹⁰, -OC(=N-CN)NR⁹R¹⁰, -OC(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=N-OH)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)OR⁸, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), or C₁-C₆alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^x and R⁵ are taken together to form ring D optionally substituted with one or more R^D;

Ring D is a cycloalkyl, heterocycloalkyl, aryl, or heteroaryl;

each R^D is independently hydrogen, deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or two R^D on the same carbon are taken together to form an oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or

more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, cycloalkyl, or heterocycloalkyl;

or R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R⁸ and R⁹ are taken together with the atoms to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a};

each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -SR^b, -S(=O)R^a, -S(=O)₂R^a, -NO₂, -NR^cR^d, -NHS(=O)₂R^a, -S(=O)₂NR^cR^d, -C(=O)R^a, -OC(=O)R^a, -C(=O)OR^b, -OC(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, -NR^bC(=O)NR^cR^d, -NR^bC(=O)R^a, -NR^bC(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, or C₂-C₆alkynyl;

each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl,

C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl;

or R^c and R^d are taken together with the nitrogen atom to which they are attached to form a

heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00315] In some embodiments of a compound of Formula (XIII), Ring B is aryl or heteroaryl. In some embodiments of a compound of Formula (XIII), Ring B is aryl. In some embodiments of a compound of Formula (XIII), Ring B is phenyl. In some embodiments of a compound of Formula (XIII), Ring B is heteroaryl. In some embodiments of a compound of Formula (XIII), Ring B is 5- or 6-membered heteroaryl. In some embodiments of a compound of Formula (XIII), Ring B is 6-membered heteroaryl. In some embodiments of a compound of Formula (XIII), Ring B is 6-membered pyridyl.


[00316] In some embodiments of a compound of Formula (XIII), L is a bond. In some embodiments of a compound of Formula (XIII), L is -C(=O)-.

[00317] In some embodiments of a compound of Formula (XIII), L-Ring A is absent.

[00318] In some embodiments of a compound of Formula (XIII), Ring A is heterocycloalkyl or heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (XIII), Ring A is a 5-membered heterocycloalkyl or a 5-membered heteroaryl; each optionally substituted with one or more R^A. In some embodiments of a compound of Formula (XIII), Ring A is heteroaryl optionally substituted with one or more R^A.


[00319] In some embodiments of a compound of Formula (XIII), each R^A is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII), each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII), each R^A is independently halogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (XIII), each R^A is independently C₁-C₆alkyl.

[00320] In some embodiments of a compound of Formula (XIII), the compound is of Formula (XIIIa):




Formula (XIIIa).

[00321] In some embodiments of a compound of Formula (XIII), the compound is of Formula (XIIIb):




Formula (XIIIb).

[00322] In some embodiments of a compound of Formula (XIII), the compound is of Formula (XIIIc):



Formula (XIIIc).

[00323] In some embodiments of a compound of Formula (XIII), the compound is of Formula (XIIIId):






Formula (XIIIId).

[00324] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is hydrogen, deuterium, halogen, -CN, -OR^b, -S(=O)₂R^a, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, -OC(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is hydrogen or -OR^b. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is -OR^b. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is hydrogen. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is -P(=O)R^bR^b. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^4 is -S(=O)₂R^a.


[00325] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each X is -N-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId) each X is -CR^X-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), two X are -N- and the other is -CR^X-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), one X is -N- and the others are -CR^X-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each X is -CH-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), two X are -N- and the other is -

CH-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), one X is -N- and the others are -CH-. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId),




[00326] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^X is independently hydrogen, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, or C₁-C₆aminoalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^X is independently hydrogen, deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^X is independently hydrogen, deuterium, or halogen.


[00327] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId),



In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId),



In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId),



[00328] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^3 is hydrogen or C₁-C₆alkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^3 is hydrogen.

[00329] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^{11} is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more R^{11a} . In some embodiments of a compound of Formula (XIII) or

(XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, cycloalkyl, or heterocycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, or heterocycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more R^{11a}. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or cycloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl, C₁-C₆haloalkyl, or cycloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is C₁-C₆alkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R¹¹ is cycloalkyl.

[00330] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, or C₁-C₆aminoalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, or C₁-C₆aminoalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^{11a} is independently deuterium, halogen, -CN, -OR^b, -NR^cR^d, or C₁-C₆alkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^{11a} is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), each R^{11a} is independently deuterium, halogen, or C₁-C₆alkyl.

[00331] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00332] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-

C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00333] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is halogen, -CN, -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00334] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00335] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is -OR⁸, -NR⁹R¹⁰, -C(=O)R⁷, -C(=O)OR⁸, -C(=O)NR⁹R¹⁰, -NR⁸C(=O)NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸C(=O)OR⁸, -NR⁸S(=O)(=NR⁸)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, -NR⁸C(=N-CN)R⁷, -NR⁸S(=O)(=NR⁸)R⁷, or aryl optionally substituted with one or more deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00336] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=N-CN)R⁷, -NR⁸S(=O)(=NR⁸)R⁷, or aryl optionally substituted with one or more deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00337] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is -OR⁸, -NR⁹R¹⁰, -NR⁸C(=O)R⁷, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is -NR⁸C(=O)R⁷. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is aryl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00338] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^9R^{10}$, $-NR^8C(=O)NR^9R^{10}$, or $-NR^8C(=N-CN)R^7$.

[00339] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^9R^{10}$.

[00340] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^9R^{10}$ or $-NR^8C(=O)NR^9R^{10}$.

[00341] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^8C(=N-CN)R^7$.

[00342] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^8S(=O)(=NR^8)R^7$.

[00343] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^8C(=O)NR^9R^{10}$ or $-NR^8C(=O)R^7$.

[00344] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^8C(=O)NR^9R^{10}$.

[00345] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^8C(=O)R^7$.

[00346] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is $-NR^8C(=O)NR^9R^{10}$ or heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 heteroalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C_1 - C_6 alkyl(cycloalkyl), C_1 - C_6 alkyl(heterocycloalkyl), C_1 - C_6 alkyl(aryl), or C_1 - C_6 alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.


[00347] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 heteroalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C_1 - C_6 alkyl(cycloalkyl), C_1 - C_6 alkyl(heterocycloalkyl), C_1 - C_6 alkyl(aryl), C_1 - C_6 alkyl(heteroaryl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, $-C(=O)R^a$, $-C(=O)OR^b$, $-C(=O)NR^cR^d$, C_1 - C_6 alkyl, or C_1 - C_6 haloalkyl.

[00348] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R^5 is heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, $-CN$, $-OR^b$, $-NR^cR^d$, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 heteroalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, C_1 - C_6 alkyl(cycloalkyl), or C_1 - C_6 alkyl(heterocycloalkyl); wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl,


and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00349] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁵ is heterocycloalkyl optionally substituted with one or more oxo, halogen, -OR^b, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, C₁-C₆hydroxyalkyl, C₁-C₆aminoalkyl, C₁-C₆heteroalkyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, C₁-C₆alkyl(cycloalkyl), or C₁-C₆alkyl(heterocycloalkyl); wherein each alkyl, alkynyl, cycloalkyl, heterocycloalkyl, and aryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OR^b, -NR^cR^d, -C(=O)R^a, -C(=O)OR^b, -C(=O)NR^cR^d, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00350] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), the



[00351] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), the



[00352] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is C₁-C₆alkyl, cycloalkyl, or heterocycloalkyl; wherein each alkyl, cycloalkyl, or heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is C₁-C₆alkyl or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is cycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is unsubstituted cycloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -

C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁷ is unsubstituted heterocycloalkyl.

[00353] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁸ is hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁸ is hydrogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00354] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or C₁-C₆hydroxyalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁹ and R¹⁰ are independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl, cycloalkyl, and heterocycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or C₁-C₆hydroxyalkyl.



[00355] In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a C₂-C₈heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a bicyclic heterocycloalkyl or a spiro heterocycloalkyl optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound of Formula (XIII) or (XIIIa)-(XIIIId), R⁹ and R¹⁰ are taken together with the nitrogen atom to which they are attached to form a heterocycloalkyl selected from aziridinyl, azetidiny, pyrrolidinyl, piperidinyl, 2-azabicyclo[1.1.1]pentanyl, or 2-azaspiro[3.3]heptanyl, each optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl.

[00356] In some embodiments of a compound described above, each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound described above, each R^a is independently C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound described above, each R^a is independently C₁-C₆alkyl.











[00357] In some embodiments of a compound described above, each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound described above, each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound described above, each R^b is independently C₁-C₆alkyl or C₁-C₆deuteroalkyl. In some embodiments of a compound described above, each R^b is independently hydrogen or C₁-C₆alkyl. In some embodiments of a compound described above, each R^b is hydrogen. In some embodiments of a compound described above, each R^b is independently C₁-C₆alkyl.

[00358] In some embodiments of a compound described above, each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, C₁-C₆deuteroalkyl, or cycloalkyl; wherein each alkyl and cycloalkyl is independently optionally substituted with one or more oxo, deuterium, halogen, -CN, -OH, -OMe, -NH₂, -C(=O)Me, -C(=O)OH, -C(=O)OMe, C₁-C₆alkyl, or C₁-C₆haloalkyl. In some embodiments of a compound described above, each R^c and R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆haloalkyl, or C₁-C₆deuteroalkyl. In some embodiments of a compound described above, each R^c and R^d is independently hydrogen or C₁-C₆alkyl. In some embodiments of a compound described above, each R^c and R^d is hydrogen. In some embodiments of a compound described above, each R^c and R^d is independently C₁-C₆alkyl.

[00359] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:











Ex.	Structure	Ex.	Structure
1		99	

Ex.	Structure	Ex.	Structure
2		100	
3		101	
4		102	
5		103	
6		104	

Ex.	Structure	Ex.	Structure
7		105	
8		106	
9		107	
10		108	
11		109	

Ex.	Structure	Ex.	Structure
17		115	
18		116	
19		117	
20		118	
21		119	

Ex.	Structure	Ex.	Structure
22		120	
23		121	
24		122	
25		123	
26		124	











Ex.	Structure	Ex.	Structure
27		125	
28		126	
29		127	
30		128	
31		129	

Ex.	Structure	Ex.	Structure
32		130	
33		131	
34		132	
35		133	
36		134	
37		135	

Ex.	Structure	Ex.	Structure
38		136	
39		137	
40		138	
41		139	
42		140	
43		141	

Ex.	Structure	Ex.	Structure
44		142	
45		143	
46		144	
47		145	
48		146	

Ex.	Structure	Ex.	Structure
49		147	
50		148	
51		149	
52		150	
53		151	

Ex.	Structure	Ex.	Structure
54		152	
55		153	
56		154	
57		155	
58		156	

Ex.	Structure	Ex.	Structure
59		157	
60		158	
61		159	
62		160	
63		161	

Ex.	Structure	Ex.	Structure
69		169	
70		171	
71		172	
72		173	
73		174	
74		175	

Ex.	Structure	Ex.	Structure
75		176	
76		177	
77		178	
78		179	
79		180	


Ex.	Structure	Ex.	Structure
80		181	
81		182	
82		183	
83		184	
84		185	

Ex.	Structure	Ex.	Structure
85		188	
86		189	
87		190	
88		191	
89		193	


Ex.	Structure	Ex.	Structure
90		195	
91		196	
92		198	
93		199	
94		200	

Ex.	Structure	Ex.	Structure
95		201	
96		202	
97		203	
98			


[00360] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00361] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00362] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:







[00363] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00364] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00365] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00366] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00367] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00368] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00369] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00370] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00371] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:




[00372] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:





[00373] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:





[00374] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:












[00375] Disclosed herein is a compound, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, selected from the group consisting of:



wherein:

R₁= -Et, -CH₂-CD₃, or cyclopropyl;

R₂=Me, CD₃, or Et

R₃= Isopropyl, cyclobutyl, cyclopropyl, or t-butyl.

Further Forms of Compounds Disclosed Herein

Isomers/Stereoisomers

[00376] In some embodiments, the compounds described herein exist as geometric isomers. In some embodiments, the compounds described herein possess one or more double bonds. The compounds

presented herein include all cis, trans, syn, anti, entgegen (E), and zusammen (Z) isomers as well as the corresponding mixtures thereof. In some situations, the compounds described herein possess one or more chiral centers and each center exists in the R configuration or S configuration. The compounds described herein include all diastereomeric, enantiomeric, and epimeric forms as well as the corresponding mixtures thereof. In additional embodiments of the compounds and methods provided herein, mixtures of enantiomers and/or diastereoisomers, resulting from a single preparative step, combination, or interconversion are useful for the applications described herein. In some embodiments, the compounds described herein are prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds, separating the diastereomers, and recovering the optically pure enantiomers. In some embodiments, dissociable complexes are preferred. In some embodiments, the diastereomers have distinct physical properties (e.g., melting points, boiling points, solubilities, reactivity, etc.) and are separated by taking advantage of these dissimilarities. In some embodiments, the diastereomers are separated by chiral chromatography, or preferably, by separation/resolution techniques based upon differences in solubility. In some embodiments, the optically pure enantiomer is then recovered, along with the resolving agent.

Labeled compounds

[00377] In some embodiments, the compounds described herein exist in their isotopically-labeled forms. In some embodiments, the methods disclosed herein include methods of treating diseases by administering such isotopically-labeled compounds. In some embodiments, the methods disclosed herein include methods of treating diseases by administering such isotopically-labeled compounds as pharmaceutical compositions. Thus, in some embodiments, the compounds disclosed herein include isotopically-labeled compounds, which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds described herein, or a solvate, or stereoisomer thereof, include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, sulfur, fluorine, and chloride, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds described herein, and the pharmaceutically acceptable salts, solvates, or stereoisomers thereof which contain the aforementioned isotopes and/or other isotopes of other atoms are within the scope of this disclosure. Certain isotopically-labeled compounds, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavy isotopes such as deuterium, i.e., ^2H , produces certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements. In some embodiments, the isotopically labeled compound or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof is prepared by any suitable method.

[00378] In some embodiments, the compounds described herein are labeled by other means, including, but not limited to, the use of chromophores or fluorescent moieties, bioluminescent labels, or chemiluminescent labels.

Pharmaceutically acceptable salts

[00379] In some embodiments, the compounds described herein exist as their pharmaceutically acceptable salts. In some embodiments, the methods disclosed herein include methods of treating diseases by administering such pharmaceutically acceptable salts. In some embodiments, the methods disclosed herein include methods of treating diseases by administering such pharmaceutically acceptable salts as pharmaceutical compositions.

[00380] In some embodiments, the compounds described herein possess acidic or basic groups and therefor react with any of a number of inorganic or organic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt. In some embodiments, these salts are prepared *in situ* during the final isolation and purification of the compounds disclosed herein, or by separately reacting a purified compound in its free form with a suitable acid or base, and isolating the salt thus formed.

[00381] Examples of pharmaceutically acceptable salts include those salts prepared by reaction of the compounds described herein with a mineral, organic acid, or inorganic base, such salts including acetate, acrylate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, bisulfite, bromide, butyrate, butyn-1,4-dioate, camphorate, camphorsulfonate, caproate, caprylate, chlorobenzoate, chloride, citrate, cyclopentanepropionate, decanoate, digluconate, dihydrogenphosphate, dinitrobenzoate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hexyne-1,6-dioate, hydroxybenzoate, γ -hydroxybutyrate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, iodide, isobutyrate, lactate, maleate, malonate, methanesulfonate, mandelate metaphosphate, methanesulfonate, methoxybenzoate, methylbenzoate, monohydrogenphosphate, 1-naphthalenesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, pyrosulfate, pyrophosphate, propiolate, phthalate, phenylacetate, phenylbutyrate, propanesulfonate, salicylate, succinate, sulfate, sulfite, succinate, suberate, sebacate, sulfonate, tartrate, thiocyanate, tosylateundeconate, and xylenesulfonate.

[00382] Further, the compounds described herein can be prepared as pharmaceutically acceptable salts formed by reacting the free base form of the compound with a pharmaceutically acceptable inorganic or organic acid, including, but not limited to, inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid metaphosphoric acid, and the like; and organic acids such as acetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, p-toluenesulfonic acid, tartaric acid, trifluoroacetic acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, arylsulfonic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 4-methylbicyclo-[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-

carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, and muconic acid.

[00383] In some embodiments, those compounds described herein which comprise a free acid group react with a suitable base, such as the hydroxide, carbonate, bicarbonate, or sulfate of a pharmaceutically acceptable metal cation, with ammonia, or with a pharmaceutically acceptable organic primary, secondary, tertiary, or quaternary amine. Representative salts include the alkali or alkaline earth salts, like lithium, sodium, potassium, calcium, and magnesium, and aluminum salts and the like. Illustrative examples of bases include sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, $N^+(C_{1-4} \text{ alkyl})_4$, and the like.

[00384] Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine, and the like. It should be understood that the compounds described herein also include the quaternization of any basic nitrogen-containing groups they contain. In some embodiments, water or oil-soluble or dispersible products are obtained by such quaternization.

Solvates

[00385] In some embodiments, the compounds described herein exist as solvates. The disclosure provides for methods of treating diseases by administering such solvates. The disclosure further provides for methods of treating diseases by administering such solvates as pharmaceutical compositions.

[00386] Solvates contain either stoichiometric or non-stoichiometric amounts of a solvent, and, in some embodiments, are formed during the process of crystallization with pharmaceutically acceptable solvents such as water, ethanol, and the like. Hydrates are formed when the solvent is water, or alcoholates are formed when the solvent is alcohol. Solvates of the compounds described herein can be conveniently prepared or formed during the processes described herein. In addition, the compounds provided herein can exist in unsolvated as well as solvated forms. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the compounds and methods provided herein.

Tautomers

[00387] In some situations, compounds exist as tautomers. The compounds described herein include all possible tautomers within the formulas described herein. Tautomers are compounds that are interconvertible by migration of a hydrogen atom, accompanied by a switch of a single bond and adjacent double bond. In bonding arrangements where tautomerization is possible, a chemical equilibrium of the tautomers will exist. All tautomeric forms of the compounds disclosed herein are contemplated. The exact ratio of the tautomers depends on several factors, including temperature, solvent, and pH.

Preparation of the Compounds

[00388] The compounds used in the reactions described herein are made according to organic synthesis techniques known to those skilled in this art, starting from commercially available chemicals and/or from compounds described in the chemical literature. "Commercially available chemicals" are obtained from standard commercial sources including Acros Organics (Pittsburgh, PA), Aldrich Chemical (Milwaukee,

WI, including Sigma Chemical and Fluka), Apin Chemicals Ltd. (Milton Park, UK), Avocado Research (Lancashire, U.K.), BDH, Inc. (Toronto, Canada), Bionet (Cornwall, U.K.), Chem Service Inc. (West Chester, PA), Crescent Chemical Co. (Hauppauge, NY), Eastman Organic Chemicals, Eastman Kodak Company (Rochester, NY), Fisher Scientific Co. (Pittsburgh, PA), Fisons Chemicals (Leicestershire, UK), Frontier Scientific (Logan, UT), ICN Biomedicals, Inc. (Costa Mesa, CA), Key Organics (Cornwall, U.K.), Lancaster Synthesis (Windham, NH), Maybridge Chemical Co. Ltd. (Cornwall, U.K.), Parish Chemical Co. (Orem, UT), Pfaltz & Bauer, Inc. (Waterbury, CN), Polyorganix (Houston, TX), Pierce Chemical Co. (Rockford, IL), Riedel de Haen AG (Hanover, Germany), Spectrum Quality Product, Inc. (New Brunswick, NJ), TCI America (Portland, OR), Trans World Chemicals, Inc. (Rockville, MD), and Wako Chemicals USA, Inc. (Richmond, VA).

[00389] Suitable reference books and treatises that detail the synthesis of reactants useful in the preparation of compounds described herein, or provide references to articles that describe the preparation, include for example, "Synthetic Organic Chemistry", John Wiley & Sons, Inc., New York; S. R. Sandler et al., "Organic Functional Group Preparations," 2nd Ed., Academic Press, New York, 1983; H. O. House, "Modern Synthetic Reactions", 2nd Ed., W. A. Benjamin, Inc. Menlo Park, Calif. 1972; T. L. Gilchrist, "Heterocyclic Chemistry", 2nd Ed., John Wiley & Sons, New York, 1992; J. March, "Advanced Organic Chemistry: Reactions, Mechanisms and Structure", 4th Ed., Wiley-Interscience, New York, 1992. Additional suitable reference books and treatises that detail the synthesis of reactants useful in the preparation of compounds described herein, or provide references to articles that describe the preparation, include for example, Fuhrhop, J. and Penzlin G. "Organic Synthesis: Concepts, Methods, Starting Materials", Second, Revised and Enlarged Edition (1994) John Wiley & Sons ISBN: 3-527-29074-5; Hoffman, R.V. "Organic Chemistry, An Intermediate Text" (1996) Oxford University Press, ISBN 0-19-509618-5; Larock, R. C. "Comprehensive Organic Transformations: A Guide to Functional Group Preparations" 2nd Edition (1999) Wiley-VCH, ISBN: 0-471-19031-4; March, J. "Advanced Organic Chemistry: Reactions, Mechanisms, and Structure" 4th Edition (1992) John Wiley & Sons, ISBN: 0-471-60180-2; Otera, J. (editor) "Modern Carbonyl Chemistry" (2000) Wiley-VCH, ISBN: 3-527-29871-1; Patai, S. "Patai's 1992 Guide to the Chemistry of Functional Groups" (1992) Interscience ISBN: 0-471-93022-9; Solomons, T. W. G. "Organic Chemistry" 7th Edition (2000) John Wiley & Sons, ISBN: 0-471-19095-0; Stowell, J.C., "Intermediate Organic Chemistry" 2nd Edition (1993) Wiley-Interscience, ISBN: 0-471-57456-2; "Industrial Organic Chemicals: Starting Materials and Intermediates: An Ullmann's Encyclopedia" (1999) John Wiley & Sons, ISBN: 3-527-29645-X, in 8 volumes; "Organic Reactions" (1942-2000) John Wiley & Sons, in over 55 volumes; and "Chemistry of Functional Groups" John Wiley & Sons, in 73 volumes.

[00390] Specific and analogous reactants are optionally identified through the indices of known chemicals prepared by the Chemical Abstract Service of the American Chemical Society, which are available in most public and university libraries, as well as through on-line. Chemicals that are known but not commercially available in catalogs are optionally prepared by custom chemical synthesis houses, where many of the standard chemical supply houses (*e.g.*, those listed above) provide custom synthesis services. A reference for

the preparation and selection of pharmaceutical salts of the compounds described herein is P. H. Stahl & C. G. Wermuth "Handbook of Pharmaceutical Salts", Verlag Helvetica Chimica Acta, Zurich, 2002.

Pharmaceutical Compositions

[00391] In certain embodiments, the compound described herein is administered as a pure chemical. In some embodiments, the compound described herein is combined with a pharmaceutically suitable or acceptable carrier (also referred to herein as a pharmaceutically suitable (or acceptable) excipient, physiologically suitable (or acceptable) excipient, or physiologically suitable (or acceptable) carrier) selected on the basis of a chosen route of administration and standard pharmaceutical practice as described, for example, in *Remington: The Science and Practice of Pharmacy* (Gennaro, 21st Ed. Mack Pub. Co., Easton, PA (2005)).

[00392] Accordingly, provided herein is a pharmaceutical composition comprising a compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, and a pharmaceutically acceptable excipient.

[00393] In certain embodiments, the compound provided herein is substantially pure, in that it contains less than about 5%, or less than about 1%, or less than about 0.1%, of other organic small molecules, such as unreacted intermediates or synthesis by-products that are created, for example, in one or more of the steps of a synthesis method.

[00394] Pharmaceutical compositions are administered in a manner appropriate to the disease to be treated (or prevented). An appropriate dose and a suitable duration and frequency of administration will be determined by such factors as the condition of the patient, the type and severity of the patient's disease, the particular form of the active ingredient, and the method of administration. In general, an appropriate dose and treatment regimen provides the composition(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit (e.g., an improved clinical outcome, such as more frequent complete or partial remissions, or longer disease-free and/or overall survival, or a lessening of symptom severity). Optimal doses are generally determined using experimental models and/or clinical trials. The optimal dose depends upon the body mass, weight, or blood volume of the patient.

[00395] In some embodiments, the pharmaceutical composition is formulated for oral, topical (including buccal and sublingual), rectal, vaginal, transdermal, parenteral, intrapulmonary, intradermal, intrathecal and epidural and intranasal administration. Parenteral administration includes intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In some embodiments, the pharmaceutical composition is formulated for intravenous injection, oral administration, inhalation, nasal administration, topical administration, or ophthalmic administration. In some embodiments, the pharmaceutical composition is formulated for oral administration. In some embodiments, the pharmaceutical composition is formulated for intravenous injection. In some embodiments, the pharmaceutical composition is formulated as a tablet, a pill, a capsule, a liquid, an inhalant, a nasal spray solution, a suppository, a suspension, a gel, a colloid, a dispersion, a suspension, a solution, an emulsion, an

ointment, a lotion, an eye drop, or an ear drop. In some embodiments, the pharmaceutical composition is formulated as a tablet.

[00396] Suitable doses and dosage regimens are determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages that are less than the optimum dose of the compound disclosed herein. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. In some embodiments, the present method involve the administration of about 0.1 μg to about 50 mg of at least one compound described herein per kg body weight of the subject. For a 70 kg patient, dosages of from about 10 μg to about 200 mg of the compound disclosed herein would be more commonly used, depending on a subject's physiological response.

[00397] By way of example only, the dose of the compound described herein for methods of treating a disease as described herein is about 0.001 to about 1 mg/kg body weight of the subject per day, for example, about 0.001 mg, about 0.002 mg, about 0.005 mg, about 0.010 mg, 0.015 mg, about 0.020 mg, about 0.025 mg, about 0.050 mg, about 0.075 mg, about 0.1 mg, about 0.15 mg, about 0.2 mg, about 0.25 mg, about 0.5 mg, about 0.75 mg, or about 1 mg/kg body weight per day. In some embodiments, the dose of compound described herein for the described methods is about 1 to about 1000 mg/kg body weight of the subject being treated per day, for example, about 1 mg, about 2 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 50 mg, about 75 mg, about 100 mg, about 150 mg, about 200 mg, about 250 mg, about 500 mg, about 750 mg, or about 1000 mg per day.

Methods of Treatment

[00398] The compounds disclosed herein, or pharmaceutically acceptable salts, solvates, or stereoisomers thereof, are useful for the inhibition of kinase activity of one or more enzymes. In some embodiments the kinase inhibited by the compounds and methods is TYK2.

[00399] Provided herein are compounds that are inhibitors of TYK2 and are therefore useful for treating one or more disorders associated with activity of TYK2 or mutants thereof.

[00400] Provided herein are methods for treating a disease or disorder, wherein the disease or disorder is an autoimmune disorders, inflammatory disorders, proliferative disorders, endocrine disorders, neurological disorders, or disorders associated with transplantation, said method comprising administering to a patient in need thereof, a pharmaceutical composition comprising an effective amount of a compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof.

[00401] In some embodiments, the disease or disorder is an autoimmune disorder. In some embodiments the disease or disorder is selected from type 1 diabetes, systemic lupus erythematosus, multiple sclerosis, psoriasis, Behçet's disease, POEMS syndrome, Crohn's disease, ulcerative colitis, and inflammatory bowel disease.

[00402] In some embodiments, the disease or disorder is an inflammatory disorder. In some embodiments, the inflammatory disorder is rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, psoriasis, hepatomegaly, Crohn's disease, ulcerative colitis, inflammatory bowel disease.

[00403] In some embodiments, the disease or disorder is a proliferative disorder. In some embodiments, the proliferative disorder is a hematological cancer. In some embodiments the proliferative disorder is a leukemia. In some embodiments, the leukemia is a T-cell leukemia. In some embodiments the T-cell leukemia is T-cell acute lymphoblastic leukemia (T-ALL). In some embodiments the proliferative disorder is polycythemia vera, myelofibrosis, essential or thrombocytosis.

[00404] In some embodiments, the disease or disorder is an endocrine disorder. In some embodiments, the endocrine disorder is polycystic ovary syndrome, Crouzon's syndrome, or type 1 diabetes.

[00405] In some embodiments, the disease or disorder is a neurological disorder. In some embodiments, the neurological disorder is Alzheimer's disease.

[00406] In some embodiments the proliferative disorder is associated with one or more activating mutations in TYK2. In some embodiments, the activating mutation in TYK2 is a mutation to the FERM domain, the JH2 domain, or the kinase domain. In some embodiments the activating mutation in TYK2 is selected from G36D, S47N, R425H, V731I, E957D, and R1027H.

[00407] In some embodiments, the disease or disorder is associated with transplantation. In some embodiments the disease or disorder associated with transplantation is transplant rejection, or graft versus host disease.

[00408] In some embodiments the disease or disorder is associated with type I interferon, IL-10, IL-12, or IL-23 signaling. In some embodiments the disease or disorder is associated with type I interferon signaling. In some embodiments the disease or disorder is associated with IL-10 signaling. In some embodiments the disorder is associated with IL-12 signaling. In some embodiments the disease or disorder is associated with IL-23 signaling.

[00409] Provided herein are methods for treating an inflammatory or allergic condition of the skin, for example psoriasis, contact dermatitis, atopic dermatitis, alopecia areata, erythema multiforma, dermatitis herpetiformis, scleroderma, vitiligo, hypersensitivity angiitis, urticaria, bullous pemphigoid, lupus erythematosus, systemic lupus erythematosus, pemphigus vulgaris, pemphigus foliaceus, paraneoplastic pemphigus, epidermolysis bullosa acquisita, acne vulgaris, and other inflammatory or allergic conditions of the skin.

[00410] Provided herein are methods for treating other diseases or conditions, such as diseases or conditions having an inflammatory component, for example, treatment of diseases and conditions of the eye such as ocular allergy, conjunctivitis, keratoconjunctivitis sicca, and vernal conjunctivitis, diseases affecting the nose including allergic rhinitis, and inflammatory disease in which autoimmune reactions are implicated or having an autoimmune component or etiology, including autoimmune hematological disorders (e.g. hemolytic anemia, aplastic anemia, pure red cell anemia and idiopathic thrombocytopenia), systemic lupus erythematosus, rheumatoid arthritis, polycondritis, scleroderma, Wegener granulomatosis, dermatomyositis, chronic active hepatitis, myasthenia gravis, Steven-Johnson syndrome, idiopathic sprue, autoimmune inflammatory bowel disease (e.g. ulcerative colitis and Crohn's disease), irritable bowel syndrome, celiac disease, periodontitis, hyaline membrane disease, kidney disease, glomerular disease, alcoholic liver disease, multiple sclerosis, endocrine ophthalmopathy, Grave's

disease, sarcoidosis, alveolitis, chronic hypersensitivity pneumonitis, multiple sclerosis, primary biliary cirrhosis, uveitis (anterior and posterior), Sjogren's syndrome, keratoconjunctivitis sicca and vernal keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis, systemic juvenile idiopathic arthritis, cryopyrin-associated periodic syndrome, nephritis, vasculitis, diverticulitis, interstitial cystitis, glomerulonephritis (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy), chronic granulomatous disease, endometriosis, leptospirosis renal disease, glaucoma, retinal disease, ageing, headache, pain, complex regional pain syndrome, cardiac hypertrophy, musclewasting, catabolic disorders, obesity, fetal growth retardation, hypercholesterolemia, heart disease, chronic heart failure, mesothelioma, anhidrotic ectodermal dysplasia, Behcet's disease, incontinentia pigmenti, Paget's disease, pancreatitis, hereditary periodic fever syndrome, asthma (allergic and non-allergic, mild, moderate, severe, bronchitic, and exercise-induced), acute lung injury, acute respiratory distress syndrome, eosinophilia, hypersensitivities, anaphylaxis, nasal sinusitis, ocular allergy, silica induced diseases, COPD (reduction of damage, airways inflammation, bronchial hyperreactivity, remodeling or disease progression), pulmonary disease, cystic fibrosis, acid-induced lung injury, pulmonary hypertension, polyneuropathy, cataracts, muscle inflammation in conjunction with systemic sclerosis, inclusion body myositis, myasthenia gravis, thyroiditis, Addison's disease, lichen planus, Type 1 diabetes, or Type 2 diabetes, appendicitis, atopic dermatitis, asthma, allergy, blepharitis, bronchiolitis, bronchitis, bursitis, cervicitis, cholangitis, cholecystitis, chronic graft rejection, colitis, conjunctivitis, Crohn's disease, cystitis, dacryoadenitis, dermatitis, dermatomyositis, encephalitis, endocarditis, endometritis, enteritis, enterocolitis, epicondylitis, epididymitis, fasciitis, fibrositis, gastritis, gastroenteritis, Henoch-Schonlein purpura, hepatitis, hidradenitis suppurativa, immunoglobulin A nephropathy, interstitial lung disease, laryngitis, mastitis, meningitis, myelitis myocarditis, myositis, nephritis, oophoritis, orchitis, osteitis, otitis, pancreatitis, parotitis, pericarditis, peritonitis, pharyngitis, pleuritis, phlebitis, pneumonitis, pneumonia, polymyositis, proctitis, prostatitis, pyelonephritis, rhinitis, salpingitis, sinusitis, stomatitis, synovitis, tendonitis, tonsillitis, ulcerative colitis, uveitis, vaginitis, vasculitis, or vulvitis.

[00411] In some embodiments the inflammatory disease is acute and chronic gout, chronic gouty arthritis, psoriasis, psoriatic arthritis, rheumatoid arthritis, Juvenile rheumatoid arthritis, Systemic juvenile idiopathic arthritis (SJIA), Cryopyrin Associated Periodic Syndrome (CAPS), or osteoarthritis.

[00412] In some embodiments the inflammatory disease is a Th1 or Th17 mediated disease. In some embodiments the Th17 mediated disease is selected from Systemic lupus erythematosus, Multiple sclerosis, and inflammatory bowel disease (including Crohn's disease or ulcerative colitis).

[00413] In some embodiments the inflammatory disease is Sjogren's syndrome, allergic disorders, osteoarthritis, conditions of the eye such as ocular allergy, conjunctivitis, keratoconjunctivitis sicca, vernal conjunctivitis, or diseases affecting the nose such as allergic rhinitis.

Combination Therapy

[00414] In certain instances, the compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, is administered in combination with a second therapeutic agent.

[00415] In some embodiments, the benefit experienced by a patient is increased by administering one of the compounds described herein with a second therapeutic agent (which also includes a therapeutic regimen) that also has therapeutic benefit.

[00416] In one specific embodiment, a compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, is co-administered with a second therapeutic agent, wherein the compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, and the second therapeutic agent modulate different aspects of the disease, disorder or condition being treated, thereby providing a greater overall benefit than administration of either therapeutic agent alone.

[00417] In any case, regardless of the disease, disorder or condition being treated, the overall benefit experienced by the patient is simply additive of the two therapeutic agents or the patient experiences a synergistic benefit.

[00418] In certain embodiments, different therapeutically-effective dosages of the compounds disclosed herein will be utilized in formulating a pharmaceutical composition and/or in treatment regimens when the compounds disclosed herein are administered in combination with a second therapeutic agent. Therapeutically-effective dosages of drugs and other agents for use in combination treatment regimens are optionally determined by means similar to those set forth hereinabove for the actives themselves. Furthermore, the methods of prevention/treatment described herein encompasses the use of metronomic dosing, i.e., providing more frequent, lower doses in order to minimize toxic side effects. In some embodiments, a combination treatment regimen encompasses treatment regimens in which administration of a compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, is initiated prior to, during, or after treatment with a second agent described herein, and continues until any time during treatment with the second agent or after termination of treatment with the second agent. It also includes treatments in which a compound described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, and the second agent being used in combination are administered simultaneously or at different times and/or at decreasing or increasing intervals during the treatment period. Combination treatment further includes periodic treatments that start and stop at various times to assist with the clinical management of the patient.

[00419] It is understood that the dosage regimen to treat, prevent, or ameliorate the condition(s) for which relief is sought, is modified in accordance with a variety of factors (e.g. the disease, disorder or condition from which the subject suffers; the age, weight, sex, diet, and medical condition of the subject). Thus, in some instances, the dosage regimen actually employed varies and, in some embodiments, deviates from the dosage regimens set forth herein.

[00420] For combination therapies described herein, dosages of the co-administered compounds vary depending on the type of co-drug employed, on the specific drug employed, on the disease or condition being treated, and so forth. In additional embodiments, when co-administered with a second therapeutic

agent, the compound provided herein is administered either simultaneously with the second therapeutic agent, or sequentially.


[00421] In combination therapies, the multiple therapeutic agents (one of which is one of the compounds described herein) are administered in any order or even simultaneously. If administration is simultaneous, the multiple therapeutic agents are, by way of example only, provided in a single, unified form, or in multiple forms (e.g., as a single pill or as two separate pills).

[00422] The compounds described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, as well as combination therapies, are administered before, during, or after the occurrence of a disease or condition, and the timing of administering the composition containing a compound varies. Thus, in one embodiment, the compounds described herein are used as a prophylactic and are administered continuously to subjects with a propensity to develop conditions or diseases in order to prevent the occurrence of the disease or condition. In another embodiment, the compounds and compositions are administered to a subject during or as soon as possible after the onset of the symptoms. In specific embodiments, a compound described herein is administered as soon as is practicable after the onset of a disease or condition is detected or suspected, and for a length of time necessary for the treatment of the disease. In some embodiments, the length required for treatment varies, and the treatment length is adjusted to suit the specific needs of each subject. For example, in specific embodiments, a compound described herein or a formulation containing the compound is administered for at least 2 weeks, about 1 month to about 5 years.

[00423] In some embodiments, the compound of described herein, or a pharmaceutically acceptable salt, solvate, or stereoisomer thereof, is administered in combination with an adjuvant. In one embodiment, the therapeutic effectiveness of one of the compounds described herein is enhanced by administration of an adjuvant (i.e., by itself the adjuvant has minimal therapeutic benefit, but in combination with another therapeutic agent, the overall therapeutic benefit to the patient is enhanced).

EXAMPLES

Intermediate A:



Step 1: synthesis of compound Ab

[00424] To a solution of methyl 2-hydroxy-3-nitrobenzoate (10 g, 50.7 mmol) in DMF (100 mL) at room temperature was added potassium carbonate (14.02 g, 101 mmol) followed by addition of methyl iodide

(6.34 mL, 101 mmol) and the resulting orange mixture was heated to 60 °C for 1 h. The reaction was cooled to room temperature and then crushed ice (~100 mL) was added, followed by water to a total volume of ~400 mL causing a yellow solid to crystallize from solution. The slurry was stirred for a few minutes and then collected by vacuum filtration and the resulting initially yellow solid was rinsed with additional water (~100 mL) until all of the yellow color was rinsed into the filtrate giving a near white solid in the funnel. Partially air-dried solid in funnel then transferred to a flask and further dried under vacuum over night to afford **Ab** (10.5 g, 98%) of a yellow solid as the desired product. LCMS $[M+1]^+ = 197.1$.

Step 2: synthesis of compound **Ac**

[00425] Methyl 2-methoxy-3-nitrobenzoate (11 g, 52.1 mmol) was dissolved in a cold solution of ammonia in methanol (7 N, 250 mL) and cone. aqueous ammonium hydroxide (100 mL) was added. The flask was sealed and the resulting solution was allowed to gently stir at room temperature overnight (~17 h). The reaction mixture was concentrated to afford **Ac** (1.67 g, 86%) as a yellow solid. LCMS $[M+1]^+ = 196.1$.

Step 3: synthesis of compound **Ad**

[00426] 2-methoxy-3-nitrobenzamide (6.5 g, 33.1 mmol) was slurried in dimethyl formamide dimethyl acetal (39.5 g, 331 mmol) and the mixture was heated to 95 °C giving a clear, pale yellow solution. After heating for ~30 min at 95 °C, the reaction was cooled and was concentrated on the rotovap and the resulting yellow oil was azeotroped twice with 1,2-dichloroethane (40 mL portions) to ensure complete removal of any residual dimethyl formamide dimethyl acetal. The crude oil thus obtained was immediately dissolved in 35 mL of ethanol and was immediately used in the following step.

[00427] In a separate flask was prepared a mixture of ethanol (150 mL) and acetic acid (AcOH, 35 mL) and the resulting solution was cooled in an ice bath. Once cooled, hydrazine hydrate (16.1 mL, 331 mmol) was added dropwise. At this time, the solution containing the crude dimethyl formamide dimethyl acetal adduct as prepared above was transferred dropwise over ~15 min by cannula into the previously prepared well-stirred ice-cold mixture containing the hydrazine. During the addition, a pale yellow solid formed in the solution. After the addition was complete, the resulting cloudy yellow mixture was allowed to warm to room temperature and stir for ~4 h. The reaction mixture at this time was concentrated on the rotovap to remove some of the ethanol, diluted with additional water and filtered to collect the solid. The solid was washed with additional portions of water, air dried in the funnel then under vacuum to afford **Ad** (4.0 g, 54.8%) of a pale yellow solid as the desired product. LCMS $[M+1]^+ = 221.1$.

Step 4: synthesis of compound **Ae**


[00428] A solution of 3-(2-methoxy-3-nitrophenyl)-4H-1,2,4-triazole (5.00 g, 22.71 mmol) in DMF (20 mL) was treated with potassium carbonate (4.20 g, 30.4 mmol). After cooling the resulting mixture in an ice bath, a solution of iodomethane (4.20 g, 29.59 mmol) in DMF (5 mL) was slowly added dropwise by syringe over 2 min. After the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to rt. After stirring at room temperature for ~4 h, LCMS analysis indicated complete and clean conversion to the region-isomeric mixture of products in ~2:1 ratio, respectively. The

reaction was cooled in an ice bath and was diluted with water (~50 mL) and the solution was extracted with EtOAc (3 x 40 mL) and the combined extracts were washed with 10% aq. LiCl (2 x 20 mL), water (20 mL) then brine (20 mL), concentrated and purified by CC to afford **Ae** (2.0 g, 38%) of the major isomer as a pale yellow solid. LCMS $[M+1]^+ = 235.1$. $^1\text{H NMR}$ (400MHz, methanol- d_6) δ 8.50 (s, 1H), 8.11 (dd, $J = 7.9, 1.8$ Hz, 1H), 7.85 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.38 (t, $J = 8.0$ Hz, 1H), 4.03 (s, 3H), 3.83 (s, 3H).

Step 5: synthesis of Intermediate A

[00429] A solution of 3-(2-methoxy-3-nitrophenyl)-1-methyl-1H-1,2,4-triazole (2.20 g, 10.77 mmol) in EtOH (50 mL) was sparged with nitrogen for a few minutes before adding 10 % Pd-C (0.90 g, 0.43 mmol) followed by sparging with hydrogen from a balloon for a few minutes then allowing the mixture to stir under a balloon of hydrogen for 1.5 h at rt. The mixture was then sparged with nitrogen to deactivate the catalyst and the mixture was filtered through a pad of celite washing with additional amounts of EtOH and the resulting clear, colorless filtrate containing the product was concentrated and purified by CC to afford an off-white solid **Intermediate A** (1.5 g, 68%). LCMS $[M+1]^+ = 205.2$. $^1\text{H NMR}$ (400MHz, chloroform- d) δ 8.09 (s, 1H), 7.35 (dd, $J = 7.8, 1.7$ Hz, 1H), 7.00 (t, $J = 7.8$ Hz, 1H), 6.82 (dd, $J = 7.8, 1.7$ Hz, 1H), 4.00 (s, 3H), 3.94 (s, 2H), 3.78 (s, 3H).

Intermediate 1:



Step 1: synthesis of compound Xb:


[00430] Xa (10 g, 64.1 mmol) was placed in a 1 L flask and triethylamine (8.9 mL, 64.1 mmol) was added, followed by phosphorus oxychloride (50 mL, 546 mmol). the mixture was heated to 110°C and stirred for 120 minutes, then concentrated under reduced pressure. 200 mL of anhydrous 1,2-dichloroethane was added to the residue and the mixture sonicated and then concentrated. the residue was dissolved in THF (200 mL), deuteromethylamine (HCl salt, 2.26 g, 32 mmol) was then added, followed by *N,N*-diisopropylethylamine (18 mL, 103 mmol). After 1 hour the reaction was concentrated and the residue adsorbed onto celite using dichloromethane, The filtrate was re-concentrated and purified by CC to afford a yellow solid **Xb** (5.1 g, 37%).

Step 2: synthesis of Intermediate 1:

[00431] To a solution of **Xb** (5.00 g, 23.92 mmol) and **intermediate A** (4.88 g, 23.92 mmol) in THF (50 mL) under N_2 was added LiHMDS (1 M, 71.80 mL, 71.80 mmol) at 0°C resulting a mild exotherm. The reaction was stirred at r.t. for 3h. The reaction mixture was cooled to 0 °C, quenched by adding satd.

NH₄Cl (aq.), diluted with water (100 mL) and extracted by EtOAc (50 mL×3). The combined organic layers were washed by brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (DCM/EtOAc = 3/1) to give the desired product **Intermediate 1** (6.5 g, yield: 72.2%) as a yellow solid. LCMS [M+1]⁺ = 377.1.

Intermediate 2:



Step 1: Synthesis of compound 2Ae

[00432] A solution of 3-(2-methoxy-3-nitrophenyl)-4H-1,2,4-triazole (**Ad**) (5.00 g, 22.71 mmol) in DMF (20 mL) was treated with potassium carbonate (4.20 g, 30.4 mmol). After cooling the resulting mixture in an ice bath, a solution of iodomethane-*d*₃ (4.20 g, 29.59 mmol) in DMF (5 mL) was slowly added dropwise by syringe over 2 min. After the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to rt. After stirring at room temperature for ~4 h, LCMS analysis indicated complete and clean conversion to the regioisomeric mixture of products in ~2:1 ratio, respectively. The reaction was cooled in an ice bath and was diluted with water (~50 mL) and the solution was extracted with EtOAc (3 x 40 mL) and the combined extracts were washed with 10% aq. LiCl (2 x 20 mL), water (20 mL) then brine (20 mL), concentrated and purified by CC to afford **2Ae** (2.0 g, 38%) of the major isomer as a pale yellow solid. LCMS [M+1]⁺ = 238.1.

Step 2: Synthesis of compound 2A


[00433] A solution of 3-(2-methoxy-3-nitrophenyl)-1-(methyl-*d*₃)-1H-1,2,4-triazole (2.20 g, 10.77 mmol) in EtOH (50 mL) was sparged with nitrogen for a few minutes before adding 10% Pd-C (0.90 g, 0.43 mmol) followed by sparging with hydrogen from a balloon for a few minutes then allowing the mixture to stir under a balloon of hydrogen for 1.5 h at rt. The mixture was then sparged with nitrogen to deactivate the catalyst and the mixture was filtered through a pad of CELITE® washing with additional amounts of EtOH and the resulting clear, colorless filtrate containing the product was concentrated and purified by CC to afford an off-white solid **2A** (1.5 g, 68%). LCMS [M+1]⁺ = 208.2.

Step 3: Synthesis of intermediate 2:

[00434] To a solution of **Xb** (5.00 g, 23.92 mmol) and **Example 2A** (4.88 g, 23.92 mmol) in THF (50 mL) under N₂ was added LiHMDS (1 M, 71.80 mL, 71.80 mmol) at 0°C resulting a mild exotherm. The reaction was stirred at r.t. When completed, The reaction mixture was cooled to 0°C, quenched by adding satd. NH₄Cl (aq.), diluted with water (100 mL) and extracted by EtOAc (50 mL*3). The combined organic layers were washed by brine (50 mL), dried over Na₂SO₄, filtered and concentrated under

reduced pressure to afford the crude product which was purified by column chromatography (DCM/EtOAc=3/1) to give the desired product **intermediate 2** (6.5 g, yield: 72.2%) as a yellow solid. LCMS $[M+1]^+ = 380.1$.

Intermediate 3A:



Step 1: Synthesis of compound 3Ab

[00435] To a solution of methyl 2-hydroxy-3-nitrobenzoate (7 g, 35.5 mmol) in DMF (70 mL) at room temperature was added potassium carbonate (9.8 g, 71.0 mmol) followed by addition of iodomethane- d_3 (4.42 mL, 71.0 mmol) and the resulting orange mixture was heated to 60 °C for 1 h. The reaction was cooled to room temperature and then crushed ice (~100 mL) was added, followed by water to a total volume of ~400 mL causing a yellow solid to crystallize from solution. The slurry was stirred for a few minutes and then collected by vacuum filtration and the resulting initially yellow solid was rinsed with additional water (~100 mL) until all of the yellow color was rinsed into the filtrate giving a near white solid in the funnel. Partially air-dried solid in funnel then transferred to a flask and further dried under vacuum overnight to afford **3Ab** (6.5 g, 86%) of a yellow solid as the desired product. LCMS $[M+1]^+ = 215.1$.

Step 2: Synthesis of compound 3Ac

[00436] methyl 2-(methoxy- d_3)-3-nitrobenzoate **3Ab** (6.5 g, 30.3 mmol) was dissolved in a cold solution of ammonia in methanol (7N, 140 mL) and cone. aqueous ammonium hydroxide (60 mL) was added. The flask was sealed and the resulting solution was allowed to gently stir at room temperature overnight (~17 h). The reaction mixture was concentrated to afford **3Ac** (5.8 g, 96%) as a yellow solid. LCMS $[M+1]^+ = 200.1$

Step 3: Synthesis of compound 3Ad

[00437] 2-(methoxy- d_3)-3-nitrobenzamide **3Ac** (5.8 g, 29.1 mmol) was slurried in dimethyl formamide dimethyl acetal (38.6 mL, 291 mmol) and the mixture was heated to 95°C giving a clear, pale yellow solution. After heating for ~30 min at this temp the reaction was cooled and was concentrated on the rotovap and the resulting yellow oil was azeotroped twice with 1,2-dichloroethane (40 mL portions) to ensure complete removal of any residual dimethyl formamide dimethyl acetal. The crude oil thus obtained was immediately dissolved in 35 mL of ethanol and was immediately used in the following step.


[00438] In a separate flask was prepared a mixture of ethanol (150 mL) and acetic acid (AcOH, 35 mL) and the resulting solution was cooled in an ice bath. Once cooled, hydrazine hydrate (14.1 mL, 291 mmol) was added dropwise. At this time, the solution containing the crude dimethyl formamide dimethyl acetal adduct as prepared above was transferred dropwise over ~15 min by cannula into the previously prepared well-stirred ice-cold mixture containing the hydrazine. During the addition, a pale yellow solid formed in the solution. After the addition was complete, the resulting cloudy yellow mixture was allowed to warm to room temperature and stir for ~4 h. The reaction mixture at this time was concentrated on the rotovap to remove some of the ethanol, diluted with additional water and filtered to collect the solid. The solid was washed with additional portions of water, air dried in the funnel then under vacuum to afford **3Ad** (5.0 g, 77.0%) of a pale yellow solid as the desired product. LCMS $[M+1]^+ = 224.1$

Step 4: Synthesis of compound **3Ae**

[00439] A solution of 3-(2-(methoxy-d3)-3-nitrophenyl)-1H-1,2,4-triazole **3Ad** (5.00 g, 22.4 mmol) in DMF (20 mL) was treated with potassium carbonate (9.28 g, 67.2 mmol). After cooling the resulting mixture in an ice bath, a solution of iodomethane (1.9 mL, 30.2 mmol) in DMF (5 mL) was slowly added dropwise by syringe over 2 min. After the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to rt. After stirring at room temperature for ~4 h, LCMS analysis indicated complete and clean conversion to the regioisomeric mixture of products in ~2:1 ratio, respectively. The reaction was cooled in an ice bath and was diluted with water (~50 mL) and the solution was extracted with EtOAc (3 x 40 mL) and the combined extracts were washed with 10% aq. LiCl (2 x 20 mL), water (20 mL) then brine (20 mL), concentrated and purified by CC to afford **3Ae** (2.1 g, 39.54%) of the major isomer as a pale yellow solid. LCMS $[M+1]^+ = 238.2$.


Step 5: Synthesis of compound **3A**

[00440] A solution of 3-(2-(methoxy-d3)-3-nitrophenyl)-1-methyl-1H-1,2,4-triazole **3Ae** (1.6 g, 6.75 mmol) in EtOH (50 mL) was sparged with nitrogen for a few minutes before adding 10% Pd-C (0.8 g) followed by sparging with hydrogen from a balloon for a few minutes then allowing the mixture to stir under a balloon of hydrogen for 1.5h at rt. The mixture was then sparged with nitrogen to deactivate the catalyst and the mixture was filtered through a pad of CELITE® washing with additional amounts of EtOH and the resulting clear, colorless filtrate containing the product was concentrated and purified by CC to afford an off-white solid **3A** (1.1 g, 68%).
LCMS $[M+1]^+ = 208.2$. $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 8.09 (s, 1H), 7.35 (dd, 1H), 6.99 (t, 1H), 6.83 (dd, 1H), 3.99 (s, 3H), 3.80 – 3.45 (m, 2H).

Intermediate 3:

[00441] To a solution of **Xb** (0.9 g, 4.32 mmol) and **3A** (0.9 g, 4.32 mmol) in THF (15 mL) under N₂ was added LiHMDS (1 M, 13 mL, 13 mmol) at 0°C resulting a mild exotherm. The reaction was stirred at r.t. When completed, The reaction mixture was cooled to 0°C, quenched by adding satd. NH₄Cl (aq.), diluted with water (100 mL) and extracted by EtOAc (50 mL×3). The combined organic layers were washed by brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (DCM/EtOAc=3/1) to give the desired product **intermediate 3** (1.3 g, yield: 79.26%) as a yellow solid. LCMS [M+1]⁺ = 380.2.

Example 1 : 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-6-(2-oxoimidazolidin-1-yl)-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 6-(3-acetyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide


[00442] To a solution of **intermediate 1** (60mg, 0.16mmol) and **1A** (41mg, 0.32mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100mg, 0.32 mmol), Pd₂(dba)₃ (43mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N₂ for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 1** (6 mg, yield:8%) as a yellow solid. LM-MS: m/z = 469.2[M+H]⁺

Step 2: 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-6-(2-oxoimidazolidin-1-yl)-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00443] To a solution of **1B** (6 mg, 0.013mmol) in methanol (3 mL) was added NaOH (1.04 mg, 0.026mmol) The mixture was heated to 40 °C for 2 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 1** (3 mg, yield: 50%) as a yellow solid. LM-MS: m/z = 428.2[M+H]⁺. ¹H NMR (400


MHZ, DMSO) δ 10.91 (s, 1H), 9.14 (s, 1H), 8.56 (s, 1H), 8.29 (s, 1H), 7.63 (dd, 1H), 7.53 (dd, 1H), 7.43 (s, 1H), 7.25 (t, 1H), 4.17 – 4.12 (m, 2H), 3.95 (s, 3H), 3.74 (s, 3H), 3.49 – 3.37 (m, 2H).

Example 2 : 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-6-(3-methyl-2-oxo-imidazolidin-1-yl)-N-(trideuteriomethyl)pyridazine-3-carboxamide




[00444] To a solution of **intermediate 1** (60 mg, 0.16 mmol) and **2A** (32 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), $\text{Pd}_2(\text{dba})_3$ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **2**. (6 mg, yield: 9%) as a yellow solid. LM-MS: $m/z = 441.2$ $[\text{M}+\text{H}]^+$. ^1H NMR (400 MHz, CDCl_3) δ 11.60 (s, 1H), 8.39 (s, 1H), 8.22 (s, 1H), 8.06 – 7.88 (m, 2H), 7.50 (d, 1H), 7.32 (t, 1H), 4.62 – 4.49 (m, 2H), 4.03 (s, 3H), 3.81 (s, 3H), 3.65 – 3.54 (m, 2H), 2.91 (s, 3H).

Example 3: 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-methyl-6-(2-oxoazetidin-1-yl)pyridazine-3-carboxamide




[00445] **Intermediate 1** (0.05 g, 0.133 mmol), azetidin-2-one (0.02 g, 0.282 mmol), cesium carbonate (0.086 g, 0.264 mmol), tris(dibenzylideneacetone)dipalladium (0.012 g, 0.013 mmol) and xantphos (0.012 g, 0.021 mmol) were added to 45 mL sealed tube. 10 mL 1,4-dioxane was added to the mixture. The resulting solution was stirred at 130 °C for 1h under N_2 atmosphere. The mixture solution was evaporated to dryness, residues was purified by flash chromatography to afford **3** (0.0121 g, 20%) as a light yellow solid. ^1H NMR (400 MHz, CD_3OD) δ 8.70 (s, 1H), 7.77 (d, $J = 7.6$ Hz, 1H), 7.64 – 7.55 (m, 2H), 7.34 (t, $J = 8.0$ Hz, 1H), 4.05 (s, 3H), 3.92 (t, $J = 4.9$ Hz, 2H), 3.75 (s, 3H), 3.25 (t, $J = 4.8$ Hz, 2H). LC-MS (ESI): $m/z = 412.2$ $[\text{M}+\text{H}]^+$.

Example 4 : 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-6-(4-oxo-5-azaspiro[2.4]heptan-5-yl)-N-(trideuteriomethyl)pyridazine-3-carboxamide



[00446] To a solution of **intermediate 1** (60 mg, 0.16 mmol) and **4A** (35 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), Pd₂(dba)₃ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N₂ for 3 times and heated to 110 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, The residue was purified by flash Chromatography (PE/EtOAc=50/50 then DCM/MeOH=97/3) to afford the title compound **Example 4** (24 mg, 33.33%) as a white solid. LM-MS: m/z =452.3 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ 11.22 (s, 1H), 8.53 (s, 1H), 8.32 (s, 1H), 8.16 (s, 1H), 7.78 (d, 1H), 7.55 (d, 1H), 7.31 (t, 1H), 4.34 (t, 2H), 4.05 (s, 3H), 3.78 (s, 3H), 2.26 (t, 2H), 1.28 – 1.24 (m, 2H), 0.98 – 0.93 (m, 2H).

Example 5 : 6-(3-cyclopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-cyclopropyl-urea


[00447] To a solution of cyclopropanamine (2.0 g, 35.03 mmol) in Acetonitrile (20 mL) was added 1-chloro-2-isocyanato-ethane (3.70g, 35.03mmol), then it was stirred at room temperature for 2 h. solid precipitation, filtered, give the desired product **5B** (4.05 g, 71.1%) as a white solid, it was used in the next step without further purification. LM-MS: m/z =163.2 [M+H]⁺

Step 2: 1-cyclopropylimidazolidin-2-one

[00448] To a solution of **5B** (0.5 g, 3.07 mmol) in THF(30ml) was added sodium hydride (0.15g, 6.0 mmol), then it was stirred at room temperature for 2h. Quenched with water (20 mL) and extracted by EtOAc (30 mL×2). The combined organic layers were washed by brine(30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc / PE =1/1) to give the desired product **5C** (0.17g, yield: 38.67%) as a white solid. LM-MS: m/z =127.2 [M+H]⁺

Step 3: 6-(3-cyclopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00449] To a solution of **intermediate 1** (70 mg, 0.18 mmol) and **5C** (35 mg, 0.28 mmol) in 1,4-dioxane (8mL) was added Cesium carbonate (100mg, 0.32 mmol), Pd₂(dba)₃ (43mg, 0.048 mmol), and xantphos (46m g, 0.08 mmol).The mixture degassed by N₂ for 3 times and heated to 120°Cfor 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 5** (10 mg, yield: 10%) as a yellow solid. LM-MS: m/z = 467.2[M+H]⁺. ¹H NMR (400 MHz, MeOD) δ 8.46 (s, 1H), 8.39 (s, 1H), 7.66-7.62 (m, 2H), 7.24 (t, 1H), 4.11 (t, 2H), 4.02 (s, 3H), 3.74 (s, 3H), 3.53 (t, 2H),2.56-2.52(m, 1H), 0.77-0.72(m, 4H).

Example 6 : 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide**Step 1: 1-(2-chloroethyl)-3-isopropyl-urea**

[00450] To a solution of propan-2-amine (2.0 g, 33.8 mmol) in Acetonitrile (20 mL) was added 1-chloro-2-isocyanato-ethane (3.57g, 33.8mmol) , then it was stirred at room temperature for 2h. solid precipitation, filtered, give the desired product Example 6B (4.2 g, 75.4%) as a white solid ,it was used in the next step without further purification. LM-MS: m/z =165.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ 5.39 – 4.32 (m, 2H), 3.85 (dt, 1H), 3.65 – 3.58 (m, 2H), 3.58 – 3.49 (m, 2H), 1.15 (d, 6H).

Step 2: 1-isopropylimidazolidin-2-one


[00451] To a solution of 6B (0.5 g, 3.0 mmol) in THF (30ml) was added sodium hydride (0.15g, 6.0mmol), then it was stirred at room temperature for 2h. Quenched with water (20 mL) and extracted by EtOAc (30 mL×2). The combined organic layers were washed by brine(30 mL), dried over Na₂SO₄,

filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc/PE=1/1) to give the desired product Example 6C (0.17g, yield: 44%) as a white solid. LM-MS: $m/z = 129.2 [M+H]^+$. 1H NMR (400 MHz, $CDCl_3$) δ 4.36 (s, 1H), 4.14 (dt, 1H), 3.42 – 3.33 (m, 4H), 1.14 (d, 3H), 1.12 (s, 3H).

Step 3: 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide;2,2,2-trifluoroacetic acid


[00452] To a solution of **Intermediate 1** (60m g, 0.16mmol) and **6C** (41mg, 0.32mmol) in 1,4-dioxane (8mL) was added Cesium carbonate (100mg, 0.32 mmol), $Pd_2(dba)_3$ (43mg, 0.048 mmol), and xantphos (46m g, 0.08 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc=50/50 then DCM/MeOH=97/3) to give the desired product **Example 6** (5mg, yield: 5%) as a yellow solid. LM-MS: $m/z = 469.2 [M+H]^+$. 1H NMR (400 MHz, $CDCl_3$) δ 11.47 (s, 1H), 8.54 (s, 1H), 8.02 (s, 1H), 7.89 (s, 1H), 7.83 (d, 1H), 7.54 (d, 1H), 7.34 (t, 1H), 4.21 (dd, 3H), 4.06 (s, 3H), 3.76 (s, 3H), 3.64 – 3.49 (m, 2H), 1.19 (d, 6H).

Example 7 : 6-(3,3-dimethyl-2-oxo-pyrrolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



[00453] To a solution of **intermediate 1** (60 mg, 0.16 mmol) and **7A** (36 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), $Pd_2(dba)_3$ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50/50 then DCM/MeOH = 97/3) to give the desired product **Example 7** (24 mg, yield:30%) as a yellow solid. LM-MS: $m/z = 455.3 [M+H]^+$. 1H NMR (400 MHz, $CDCl_3$) δ 11.08 (s, 1H), 8.42 (d, 2H), 8.17 (s, 1H), 7.77 (d, 1H), 7.58 (d, 1H), 7.31 (t, 1H), 4.17 (t, 2H), 4.04 (s, 3H), 3.80 (s, 3H), 2.02 (t, 2H), 1.25 (s, 6H).

Example 8 : 6-[3-(2,2-difluoroethyl)-2-oxo-imidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide;2,2,2-trifluoroacetic acid



Step 1: 1-(2-chloroethyl)-3-(2,2-difluoroethyl)urea

[00454] To a solution of 2,2-difluoroethanamine (2.0 g, 24.67 mmol) in Acetonitrile (20 mL) was added 1-chloro-2-isocyanato-ethane (2.60g, 24.67mmol), then it was stirred at room temperature for 2 h. solid precipitation, filtered, give the desired product **8B** (2.80 g, 60.83%) as a white solid ,it was used in the next step without further purification. LM-MS: $m/z = 187.2$ $[M+H]^+$


Step 2: 1-(2,2-difluoroethyl)imidazolidin-2-one

[00455] To a solution of **8B** (0.5 g, 2.68 mmol) in THF(30ml) was added sodium hydride (0.15g, 6.0mmol), then it was stirred at room temperature for 2h. Quenched with water (20 mL) and extracted by EtOAc (30 mL \times 2). The combined organic layers were washed by brine(30 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc / PE =1/1) to give the desired product **8C** (0.15g, yield: 37.29%) as a white solid. LM-MS: $m/z = 151.2$ $[M+H]^+$

Step 3: 6-[3-(2,2-difluoroethyl)-2-oxo-imidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide;2,2,2-trifluoroacetic acid

[00456] To a solution of **intermediate 1** (80 mg, 0.21 mmol) and **8C** (47 mg, 0.32 mmol) in 1,4-dioxane (8mL) was added Cesium carbonate (100 mg, 0.32 mmol), $Pd_2(dba)_3$ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol).The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 8**. (4 mg, yield: 4%) as a yellow solid. LM-MS: $m/z = 491.2$ $[M+H]^+$. 1H NMR (400 MHz, CD_3OD): δ 7.77-7.71 (m, 2H), 7.64 (d, 1H), 7.33 (t, 1H), 6.18-5.91 (m, 1H), 4.09 (t, 1H), 4.04 (s, 3H), 3.72-3.72 (m, 4H), 3.70-3.65 (m, 2H).

Example 9: 6-[3-(2-methoxyethyl)-2-oxo-imidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-(2-methoxyethyl)urea

[00457] To a solution of 2-methoxyethanamine (0.38 g, 5 mmol) in THF (10 mL) was added 1-chloro-2-isocyanato-ethane (0.53g, 5mmol), then it was stirred at room temperature for 2 h give the desired product **9B** (0.81 g, 90%) as a colorless oil ,it was used in the next step without further purification. LC-MS: $m/z = 180.6$ $[M+H]^+$


Step 2: 1-(3-methoxyethyl)imidazolidin-2-one

[00458] To a solution of **9B** (0.81g, 4.5 mmol) in THF(20 ml) was added sodium hydride (0.13 g, 5.4 mmol), then it was stirred at room temperature for 4h. Quenched with water (20 mL) and extracted by EtOAc (40 mL \times 2). The combined organic layers were washed by brine(60 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc / PE = 1/1) to give the desired product **9C** (0.25 g, yield: 40%) as a colorless oil. LC-MS: $m/z = 144.2$ $[M+H]^+$

Step 3: 6-[3-(2-methoxyethyl)-2-oxo-imidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00459] To a solution of **intermediate 1** (60 mg, 0.16 mmol) and **9C** (30 mg, 0.2 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (160 mg, 0.5 mmol), $Pd_2(dba)_3$ (20 mg, 0.02 mmol), and xantphos (24 mg, 0.04 mmol). The mixture degassed by N_2 for 3 times and heated to 130 °C for 2 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 95/5) to give the desired product **Example 9** (8.4 mg, yield: 11%) as a white solid. LM-MS: $m/z = 484.5$ $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$): δ 11.03 (s, 1H), 8.36 (s, 1H), 8.13 – 8.11 (m, 2H), 7.80 – 7.79 (d, 1H), 7.55 – 7.53 (d, 1H), 7.25 – 7.29 (m, 1H), 4.30 – 4.26 (q, 2H), 4.00 (s, 3H), 3.82 (s, 3H), 3.68 – 3.64 (q, 2H), 3.56 – 3.53 (q, 2H), 3.49 – 3.46 (q, 2H), 3.35 (s, 3H).

Example 10 : 6-[3-(3,3-difluorocyclobutyl)-2-oxo-imidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-(3,3-difluorocyclobutyl)urea

[00460] To a solution of 3,3-difluorocyclobutanamine hydrochloride (**10A**) (1.0 g, 6.96 mmol) and TEA (1.94 mL, 13.93 mmol) in THF (15 mL) was added 1-chloro-2-isocyanato-ethane (0.6 mL, 6.96 mmol) via a syringe, then it was stirred at room temperature for 2 h. The mixture solution was evaporated to dryness, redissolved in EtOAc (80 mL). The organic layer was then washed with NaHCO₃ and brine, dried over Na₂SO₄, filtered and concentrated, then the title compound **10B** (550 mg, 37.4%) was obtained as white solid, which was used in the next step without further purification. LM-MS: m/z = 213.1 [M+H]⁺

Step 2: 1-(3,3-difluorocyclobutyl)imidazolidin-2-one


[00461] To a solution of **10B** (0.5 g, 2.35 mmol) in THF (30 mL) was added sodium hydride (0.19 g, 4.7 mmol), then it was stirred at room temperature for 2 h. Quenched with water (20 mL) and extracted by EtOAc (30 mL × 2). The combined organic layers were washed by brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc/PE = 1/1) to give the desired product **10C** (0.23 g, yield: 55.4%) as a white solid.

Step 3: 6-[3-(3,3-difluorocyclobutyl)-2-oxo-imidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00462] To a solution of **intermediate 1** (60 mg, 0.16 mmol) and **10C** (56 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), Pd₂(dba)₃ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N₂ for 3 times and heated to 110 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, The residue was purified by flash Chromatography

[00463] (PE/EtOAc=50/50 then DCM/MeOH=97/3) to afford the title compound **Example 10** (12 mg, yield: 13.95%) as a white solid. LM-MS: m/z = 517.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): δ 10.98 (s, 1H), 8.32 (s, 1H), 8.18 – 8.07 (m, 2H), 7.77 (d, 1H), 7.55 – 7.49 (d, 1H), 7.30 (d, 1H), 4.51 – 4.41 (m, 1H), 4.26 (t, 2H), 4.03 (s, 3H), 3.78 (s, 3H), 3.60 (t, 2H), 2.95 – 2.71 (m, 4H).

Example 11 : 6-(3-cyclobutyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-cyclobutyl-urea

[00464] To a solution of cyclobutanamine (1.0 g, 14.1 mmol) in acetonitrile (20 mL) was added 1-chloro-2-isocyanatoethane (1.48 g, 14.1 mmol), then it was stirred at room temperature for 2h. solid precipitation, filtered, give the desired product Example 11B (1.5 g, 60.4%) as a white solid, it was used in the next step without further purification. LM-MS: $m/z = 177.1$ [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): δ 4.57 (s, 2H), 4.16 – 4.06 (m, 1H), 3.68 – 3.59 (m, 2H), 3.59 – 3.51 (m, 2H), 2.40 – 2.30 (m, 2H), 1.91 – 1.79 (m, 2H), 1.79 – 1.60 (m, 2H).

Step 2: 1-cyclobutylimidazolidin-2-one


[00465] To a solution of 1-(2-chloroethyl)-3-cyclobutyl-urea (1.0 g, 5.7 mmol) in THF (30 ml) was added sodium hydride (0.27 g, 11 mmol), then it was stirred at room temperature for 2h. Quenched with water (20 mL) and extracted by EtOAc (30 mL×2). The combined organic layers were washed by brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc/PE = 1/1) to give the desired product Example 11C (0.70g, yield: 88%) as a white solid. LM-MS: $m/z = 141.2$ [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ 4.43 (t, 1H), 3.50 (dd, 2H), 3.40 (dd, 2H), 2.14 – 2.05 (m, 4H), 1.68 – 1.61 (m, 2H).

Step 3: 6-(3-cyclobutyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00466] To a solution of Intermediate 1 (60 mg, 0.16 mmol) and 11C (45 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), Pd₂(dba)₃ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N₂ for 3 times and heated to 120°C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50/50 then DCM/MeOH = 97/3) to give the desired product 11 (5 mg, yield: 7%) as a yellow solid. LM-MS: $m/z = 481.2$ [M+H]⁺. ¹H NMR (400 MHz, CD₃OD) δ 8.46 (s, 1H), 8.37 (s, 1H), 7.63 (td, 2H), 7.29

(t, 1H), 4.43 (t, 1H), 4.19 – 4.09 (m, 2H), 4.01 (s, 3H), 3.74 (s, 3H), 3.70 – 3.61 (m, 2H), 2.38 – 2.23 (m, 2H), 2.13 (dd, 2H), 1.77 – 1.68 (m, 2H).

Example 12 : 6-(3-ethyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-ethyl-urea

[00467] To a solution of ethylamine hydrochloride (1.0 g, 12.3 mmol) in Acetonitrile (20 mL) was added Triethylamine (2.48 g, 24.6 mmol) and 1-chloro-2-isocyanatoethane (1.24 g, 12.3 mmol), then it was stirred at room temperature for 2 h. solid precipitation, filtered, give the desired product Example **12B** (1.5 g, 81%) as a white solid, it was used in the next step without further purification.


Step 2: 1-ethylimidazolidin-2-one

[00468] To a solution of **12B** (1.0 g, 6.6mmol) in THF (30 ml) was added sodium hydride (0.32 g, 13 mmol), then it was stirred at room temperature for 2 h. Quenched with water (20 mL) and extracted by EtOAc (30 mL×2). The combined organic layers were washed by brine(30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc/PE=1/1) to give the desired product Example **12C** (0.70g, yield: 92%) as a white solid.

Step 3: 6-(3-ethyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00469] To a solution of **Intermediate 1** (60 mg, 0.16 mmol) and **12C** (36 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), Pd₂(dba)₃ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol).The mixture degassed by N₂ for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc=50/50 then DCM/MeOH=97/3) to give the desired product Example **12** (5 mg, yield: 6%) as a yellow solid. LM-MS: m/z =455.3 [M+H]⁺. ¹H NMR (400 MHz, MeOD): δ 8.53 (s, 1H), 7.76 (dd, 1H), 7.63 (dd, 1H), 7.58 (s, 1H), 7.35 (d, 1H), 4.13 – 3.94 (m, 5H), 3.74 (s, 3H), 3.69 – 3.58 (m, 2H), 3.37 (d, 2H), 1.19 (t, 3H).

Example 13 : 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-[2,2,2-trideuterio-1-(trideuteriomethyl)ethyl]urea

[00470] To a solution of 1,1,1,3,3,3-hexadeuterio-2-propanamine (0.20 g, 3.1 mmol) in acetonitrile (20 mL) was added 1-chloro-2-isocyanatoethane (0.32g, 3.1mmol), then it was stirred at room temperature for 2 h. solid precipitation, filtered, give the desired product **13B** (0.4 g, 76%) as a white solid, it was used in the next step without further purification. LM-MS: $m/z = 171.2$ [M+H]⁺


Step 2: 1-[2,2,2-trideuterio-1-(trideuteriomethyl)ethyl]imidazolidin-2-one

[00471] To a solution of **13B** (0.2 g, 1.2mmol) in THF (10ml) was added sodium hydride (0.056 g, 2.4mmol), then it was stirred at room temperature for 2 h. Quenched with water (20 mL) and extracted by EtOAc (30 mL×2). The combined organic layers were washed by brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc /PE =1/1) to give the desired product Example **13C** (0.070g, yield: 40%) as a white solid.

Step 3: 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide


[00472] To a solution of **intermediate 1** (80 mg, 0.21 mmol) and **13C** (57mg, 0.42mmol) in 1,4-dioxane (8mL) was added Cesium carbonate (140 mg, 0.42 mmol), Pd₂(dba)₃ (57 mg, 0.063 mmol), and xantphos (72 mg, 0.126 mmol). The mixture degassed by N₂ for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc=50/50 then DCM/MeOH = 97/3) to give the desired product Example **13** (7 mg, yield: 7%) as a yellow solid. LM-MS: $m/z = 475.3$ [M+H]⁺. ¹H NMR (400 MHz, MeOD): δ 8.46 (s, 1H), 8.40 (d, 1H), 7.66 – 7.61 (m, 2H), 7.29 (t, 1H), 4.14 (dd, 3H), 4.01 (s, 3H), 3.74 (s, 3H), 3.58 – 3.52 (m, 2H).

Example 14: 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[3-(1-methyl-1,2,4-triazol-3-yl)-2-(trideuteriomethoxy)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide




[00473] To a solution of **intermediate 3** (160 mg, 0.42 mmol) and 1-isopropylimidazolidin-2-one (110 mg, 0.84 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (270 mg, 0.84 mmol), $\text{Pd}_2(\text{dba})_3$ (115.3 mg, 0.126 mmol), and xantphos (145.6 mg, 0.252 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc=50/50 then DCM/MeOH=97/3) to give the desired product **Example 14** (60 mg, yield: 24%) as a yellow solid. LM-MS: $m/z = 472.3$ $[\text{M}+\text{H}]^+$. ^1H NMR (400 MHz, CDCl_3) δ 11.10 (s, 1H), 8.38 (s, 1H), 8.10 (d, 2H), 7.81 (d, 1H), 7.55 (d, 1H), 7.29 (d, 1H), 4.31 (s, 2H), 4.27 – 4.21 (m, 1H), 4.00 (s, 3H), 3.53 – 3.46 (m, 2H), 1.18 (d, 6H).

Example 15 : 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-[1-(trideuteriomethyl)-1,2,4-triazol-3-yl]anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



[00474] To a solution of **Intermediate 2** (60 mg, 0.16 mmol) and 1-isopropylimidazolidin-2-one (41 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), $\text{Pd}_2(\text{dba})_3$ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 15**. (6 mg, yield: 8%) as a yellow solid. LM-MS: $m/z = 472.3$ $[\text{M}+\text{H}]^+$. ^1H NMR (400 MHz, CDCl_3) δ 10.92 (s, 1H), 8.35 (s, 1H), 8.16 (s, 1H), 8.10 (s, 1H), 7.77 (dd, 1H), 7.55 (dd, 1H), 7.28 (d, 1H), 4.25 (ddd, 3H), 3.81 (s, 3H), 3.51 – 3.44 (m, 2H), 1.18 (d, 6H).

Example 16. 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-6-[3-(3-methyloxetan-3-yl)-2-oxoimidazolidin-1-yl]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-(3-methyloxetan-3-yl)urea

[00475] To a solution of 3-methyloxetan-3-amine (0.87 g, 10 mmol) in THF (20 mL) was added 1-chloro-2-isocyanatoethane (1.06 g, 10 mmol), then it was stirred at room temperature for 2 h give the desired product **16B** (1.82 g, 94%) as a colorless oil, it was used in the next step without further purification. LC-MS: $m/z = 192.6$ [M+H]⁺


Step 2: 1-(3-methyloxetan-3-yl)imidazolidin-2-one

[00476] To a solution of **16B** (1.82 g, 9.2 mmol) in THF (30 mL) was added sodium hydride (0.29 g, 12 mmol), then it was stirred at room temperature for 4 h. Quenched with water (20 mL) and extracted by EtOAc (40 mL × 2). The combined organic layers were washed by brine (60 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc / PE = 1/1) to give the desired product **16C** (1.0 g, yield: 69%) as a white solid. LC-MS: $m/z = 156.2$ [M+H]⁺

Step 3: 4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-6-[3-(3-methyloxetan-3-yl)-2-oxoimidazolidin-1-yl]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00477] To a solution of **intermediate 1** (110 mg, 0.3 mmol) and **16C** (63 mg, 0.4 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (300 mg, 0.9 mmol), Pd₂(dba)₃ (30 mg, 0.03 mmol), and xantphos (35 mg, 0.06 mmol). The mixture degassed by N₂ for 3 times and heated to 130°C for 2 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 95/5) to give the desired product **Example 16**. (40 mg, yield: 30%) as a white solid. LM-MS: $m/z = 496.6$ [M+H]⁺. ¹H NMR (400 MHz, CDCl₃): δ 11.09 (s, 1H), 8.13 – 8.06 (m, 3H), 7.76 (s, 1H), 7.45 (s, 1H), 7.19 (m, 1H), 4.83 – 4.81 (d, 2H), 4.35 – 4.30 (m, 4H), 3.94 (s, 3H), 3.75 (s, 3H), 3.39 (m, 2H), 1.57 (s, 3H)

Example 17 : 6-(3-isobutyl-2-oxoimidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(2-chloroethyl)-3-isobutyl-urea

[00478] To a solution of 2-methylpropan-1-amine (1.0 g, 14 mmol) in acetonitrile (20 mL) was added 1-chloro-2-isocyanato-ethane (1.4 g, 14 mmol), then it was stirred at room temperature for 2 h. Solid precipitation, filtered, give the desired product Example 17B (1.5g, 61%) as a white solid, it was used in the next step without further purification. LM-MS: $m/z = 171.2$ $[M+H]^+$


Step 2: 1-(2-chloroethyl)-3-isobutyl-urea

[00479] To a solution of 1-(2-chloroethyl)-3-isobutyl-urea (1.2 g, 6.7 mmol) in THF (10 mL) was added sodium hydride (0.32 g, 13 mmol), then it was stirred at room temperature for 2 h. Quenched with water (20 mL) and extracted by EtOAc (30 mL \times 2). The combined organic layers were washed by brine (30 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc /PE = 1/1) to give the desired product Example 17C (0.5 g, yield: 50%) as a white solid.

Step 3: 6-(3-isobutyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00480] To a solution of **intermediate 1** (110 mg, 0.292 mmol) and 1-(2-chloroethyl)-3-isobutyl-urea (83 mg, 0.584 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (190 mg, 0.584 mmol), $Pd_2(dba)_3$ (80 mg, 0.087 mmol), and xantphos (100.7 mg, 0.175 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc=50/50 then DCM/MeOH=97/3) to give the desired product Example 17 (55 mg, yield: 39%) as a yellow solid. LM-MS: $m/z = 483.3$ $[M+H]^+$. 1H NMR (400 MHz, $CDCl_3$) δ 11.01 (s, 1H), 8.37 (s, 1H), 8.11 (d, 2H), 7.78 (d, 1H), 7.56 (d, 1H), 7.28 (d, 1H), 4.34 – 4.21 (m, 2H), 4.00 (s, 3H), 3.82 (s, 3H), 3.58 – 3.48 (m, 2H), 3.09 (d, 2H), 1.95 – 1.90 (m, 1H), 0.91 (dd, 6H).

Example 18 : 6-(3-cyclopentyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



Step 1: 1-(3-bicyclo[1.1.1]pentanyl)-3-(2-chloroethyl)urea

[00483] To a solution of bicyclo[1.1.1]pentan-3-amine (0.5 g, 4.2 mmol) in acetonitrile (20 mL) was added 1-chloro-2-isocyanatoethane (0.44 g, 4.2 mmol), then it was stirred at room temperature for 2 h. Solid precipitation, filtered, give the desired product Example 19B (0.5 g, 63%) as a white solid, it was used in the next step without further purification. LM-MS: $m/z = 189.1 [M+H]^+$


Step 2: 1-(3-bicyclo[1.1.1]pentanyl)imidazolidin-2-one

[00484] To a solution of 19B (0.5 g, 2.7 mmol) in THF (10 mL) was added sodium hydride (0.13 g, 5.3 mmol), then it was stirred at room temperature for 2 h. Quenched with water (20 mL) and extracted by EtOAc (30 mL \times 2). The combined organic layers were washed by brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (EtOAc/PE = 1/1) to give the desired product Example T202C (0.35 g, yield: 87%) as a white solid.

Step 3: 6-[3-(3-bicyclo[1.1.1]pentanyl)-2-oxoimidazolidin-1-yl]-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide


[00485] To a solution of intermediate 1 (170 mg, 0.451 mmol) and 19C (103 mg, 0.677 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (293 mg, 0.902 mmol), Pd₂(dba)₃ (123.8 mg, 0.135 mmol), and xantphos (155 mg, 0.27 mmol). The mixture degassed by N₂ for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50/50 then DCM/MeOH = 97/3) to give the desired product Example 19 (55 mg, yield: 39%) as a yellow solid. LM-MS: $m/z = 493.3 [M+H]^+$. ¹H NMR (400 MHz, CDCl₃): δ 10.94 (s, 1H), 8.30 (s, 1H), 8.12 (d, 2H), 7.78 (dd, 1H), 7.54 (dd, 1H), 7.28 (d, 1H), 4.24 – 4.15 (m, 2H), 4.00 (s, 3H), 3.81 (s, 3H), 3.57 – 3.46 (m, 2H), 2.48 (s, 1H), 2.13 (s, 6H).

Example 20 : 6-(3-cyclobutyl-2-oxoimidazolidin-1-yl)-4-[3-(1-methyl-1,2,4-triazol-3-yl)-2-(trideuteriomethoxy)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide




[00486] To a solution of **intermediate 3** (60 mg, 0.16 mmol) and **20C** (45 mg, 0.32 mmol) in 1,4-dioxane (8mL) was added Cesium carbonate (100 mg, 0.32 mmol), $\text{Pd}_2(\text{dba})_3$ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 20** (6 mg, yield: 8%) as a yellow solid. LM-MS: $m/z = 484.3$ $[\text{M}+\text{H}]^+$. ^1H NMR (400 MHz, CDCl_3) δ 10.85 (s, 1H), 8.34 (s, 1H), 8.11 (d, 2H), 7.75 (dd, 1H), 7.55 (dd, 1H), 7.27 (d, 1H), 4.62 – 4.47 (m, 1H), 4.27 – 4.11 (m, 2H), 3.99 (s, 3H), 3.65 – 3.52 (m, 2H), 2.26 – 2.03 (m, 4H), 1.70 (dt, 2H).

Example 21 : 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-(1-methyl-1,2,4-triazol-3-yl)anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide




[00487] To a solution of **intermediate 2** (200 mg, 0.527 mmol) and **21C** (141 mg, 1.05 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (342 mg, 1.05 mmol), $\text{Pd}_2(\text{dba})_3$ (144.6 mg, 0.158 mmol), and xantphos (182.6 mg, 0.316 mmol). The mixture degassed by N_2 for 3 times and heated to 120°C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc=50/50 then DCM/MeOH=97/3) to give the desired product **Example 21** (60 mg, yield: 24%) as a yellow solid. LM-MS: $m/z = 478.3$ $[\text{M}+\text{H}]^+$. ^1H NMR (400 MHz, CDCl_3): δ 11.04 – 10.91 (m, 1H), 8.38 (t, 1H), 8.15 – 8.03 (m, 2H), 7.82 – 7.75 (m, 1H), 7.60 – 7.52 (m, 1H), 7.31 – 7.26 (m, 1H), 4.32 – 4.18 (m, 3H), 3.86 – 3.76 (m, 3H), 3.53 – 3.44 (m, 2H).

Example 22 : 6-(3-cyclobutyl-2-oxo-imidazolidin-1-yl)-4-[2-methoxy-3-[1-(trideuteriomethyl)-1,2,4-triazol-3-yl]anilino]-N-(trideuteriomethyl)pyridazine-3-carboxamide



[00488] To a solution of **intermediate 2** (60 mg, 0.16 mmol) and **22C** (45 mg, 0.32 mmol) in 1,4-dioxane (8 mL) was added Cesium carbonate (100 mg, 0.32 mmol), $\text{Pd}_2(\text{dba})_3$ (43 mg, 0.048 mmol), and xantphos (46 mg, 0.08 mmol). The mixture degassed by N_2 for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product **Example 22**. (6 mg, yield: 8 %) as a yellow solid. LM-MS: $m/z = 484.3$ [M+H]⁺. ¹H NMR (400 MHz, CDCl_3) δ 10.97 (s, 1H), 8.35 (s, 1H), 8.11 (d, 2H), 7.79 (dd, 1H), 7.54 (dt, 1H), 7.28 (d, 1H), 4.53 (t, 1H), 4.29 – 4.21 (m, 2H), 3.81 (s, 3H), 3.66 – 3.53 (m, 2H), 2.25 – 2.07 (m, 4H), 1.74 – 1.65 (m, 2H).

Example 23:



Step 1: Example 23b

[00489] To a solution of dimethyl 3-oxopentanedioate (5.0 g, 28.7 mmol) and Ni(acac)₂ (738 mg, 2.87 mmol) in dioxane (30 mL) was added NH₂-CN (3.6 g, 86.2 mmol). The mixture was stirred at 110°C for o.n. The reaction was cooled to r.t.. The mixture was filtered and the filter cake was collected, washed with MeOH (20 mL) and concentrated in *vacuo* to give the desired product **Example 23b** (3.0 g, 56.6% yield) as a yellow solid. LCMS [M+1]⁺ = 185.0

Step 2: Example 23c

[00490] To a solution of **Example 23b** (2.5 g, 13.58 mmol) in POCl₃ (15 mL) was added DIEA (2 mL) at 0°C, which was heated to 30°C and stirred for o.n. The reaction was concentrated in *vacuo*. H₂O (15 mL) and MeOH (3 mL) were added at 0°C, which was stirred at r.t. for 1 h. The mixture was filtered and the filtrate cake was collected by filtration to give **Example 23c** (1.5 g, 50.1% yield) as a yellow solid. LCMS [M+1]⁺ = 221.0

Step 3: Example 23e

[00491] To a solution of **Example 23c** (1.2 g, 5.42 mmol) and **Example 23d** (1.21 g, 5.96 mmol) in EtOH (30 mL) was added conc. HCl (5 mL) and the solution was heated to reflux for o.n. The reaction was concentrated in *vacuo*. The residue was dissolved in EtOAc (100 mL) and H₂O (50 mL), and the pH was adjusted to ~8 with sat. NaHCO₃. The organic layer was separated and concentrated. The residue was purified by silica gel chromatography (DCM/MeOH = 30/1) to afford the product **Example 23e** (700 mg, 33.1% yield) as a yellow solid. LCMS [M+1]⁺ = 389.1

Step 4: Example 23f

[00492] To a solution of **Example 23e** (690 mg, 1.78 mmol) in THF (30 mL) and H₂O (10 mL) cooled at 0 °C was added LiOH.H₂O (112 mg, 2.67 mmol) and the solution was stirred at r.t. for o.n.. The reaction was concentrated in *vacuo*. The residue was dissolved in H₂O (50 mL), adjusted pH ~4 with HCl (2 mol/L), and extracted with EtOAc (100 mL). The organic layer was concentrated to afford the crude product **Example 23f** (750 mg, 100% crude yield) as a yellow solid. LCMS [M+1]⁺ = 375.2

Step 5: Example 23g

[00493] To a solution of **Example 23f** (500 mg, 1.33 mmol) in DMF (10 mL) were added DIEA (515 mg, 3.99 mmol), HATU (610 mg, 1.60 mmol) and CD₃-NH₂.HCl (110 mg, 1.59 mmol) and the solution was stirred at r.t. for 2 h. The reaction was diluted with EtOAc (50 mL), washed with brine (10 mL*3), dried over Na₂SO₄, and concentrated in *vacuo*. The residue was purified by silica gel chromatography (DCM/MeOH = 30/1) to afford the product **Example 14g** (510 mg, 98% yield) as a light yellow solid. LCMS [M+1]⁺ = 391.0

Step 6: Example 23h


[00494] To a mixture of **Example 23g** (500 mg, 1.278 mmol), **Example 23h** (33 mg, 1.917 mmol) and Cs₂CO₃ (167 mg, 2.55 mmol) in dioxane (5 mL) was added 3rd t-Bu-Xphos-Pd (22.5 mg, 0.128 mmol). The mixture was degassed with N₂ three times, then heated to 100°C and stirred for overnight. The reaction was concentrated in *vacuo*. The residue was further purified by prep-HPLC to give the desired product **Example 23h** (60 mg, 54.2% yield) as a white solid. LCMS [M+1]⁺ = 440.1. ¹H NMR (300

MHz, DMSO-*d*₆) δ 9.29 (s, 1H), 8.56 (s, 1H), 8.43 (s, 1H), 7.74 (d, *J* = 6.8 Hz, 1H), 7.39 (d, *J* = 7.6 Hz, 1H), 7.29 (t, *J* = 7.8 Hz, 1H), 3.94 (s, 3H), 3.70 (s, 3H), 1.65-1.80 (m, 1H), 0.8-0.94 (m, 4H).

Step 7: Example 23

[00495] To a solution of **Example 23h** (50 mg, 0.11 mmol, 1.0 eq) in pyridine (2.0 mL) was added pyridine hydrofluoride (70% in pyridine, 0.5 mL) followed by NaNO₂ (10 mg, 0.15 mmol, 1.4 eq) at 0 °C very slowly and the resulting mixture was stirred for 20 min. The reaction was concentrated in *vacuo*. The residue was purified by prep-HPLC to give the desired product **Example 23** (13.5 mg, 28.1% yield) as a light yellow solid. LCMS [M+1]⁺ = 443.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.87 (s, 1H), 10.29 (s, 1H), 8.56 (s, 1H), 8.31 (s, 1H), 7.89 (s, 1H), 7.61 (dd, *J* = 7.8 Hz, *J* = 1.8 Hz, 1H), 7.45 (d, *J* = 8.0 Hz, 1H), 7.24 (d, *J* = 7.8 Hz, 1H), 3.95 (s, 3H), 3.72 (s, 3H), 1.95-1.91 (m, 1H), 0.79-0.77 (m, 4H).

Example 24:



Step 1: Example 24c

[00496] To a solution of **Example 24a** (50.00 g, 287 mmol, 1.0 eq) and TEA (31.9 g, 316 mmol, 1.1 eq) in CH₃CN (1 L) was added **Example 24b** (70.3 g, 293 mmol, 1.02 eq) in portions at 0°C. After addition, it was stirred for 1 h at room temperature. The solid was removed by filtration, rinsed with MTBE. The filtrate was concentrated and then diluted with MTBE. The slurry was stirred for 30 min, filtered and the filtrate was concentrated to provide **Example 24c** (52.47 g, crude) as brown oil.

Step 2: Example 24d

[00497] **Example 24c** (52.47 g, 262 mmol, 1.0 eq) was dissolved in MTBE (600 mL) and PPh₃ (68.74 g, 262 mmol, 1.0 eq) was added. The reaction solution was stirred 16 h at room temperature and then concentrated in *vacuo*. To the residual sludge was added AcOH (500 mL) and H₂O (50 mL). The vessel

was equipped with a condenser and the mixture was heated to reflux for 6 h, and then concentrated in *vacuo*. The crude product (115.0 g, crude) was used to next step without purification.

Step 3: Example 24e

[00498] To last step crude product **Example 24d** (115.0 g, 32% purity, 216 mmol, 1.0 eq) dissolved in THF (150 mL)/MeOH (50 mL)/H₂O (50 mL) was added LiOH.H₂O (36.3 g, 864 mmol, 4.0 eq) and the reaction mixture was stirred for 16 at room temperature. After the reaction was completed, MeOH and THF were concentrated. The residue was diluted with H₂O (200 mL), and then extracted with EtOAc (300 mL*3). Then pH of the aqueous phase was adjusted to 3 with conc. HCl. Acidification of the water solution afforded a brown precipitate which was collected by filtration and dried in *vacuo* to afford **Example 24e** (31.5 g, 93.5% yield) as a yellow solid.

Step 4: Example 24f

[00499] The solution of **Example 24e** (15.0 g, 96.15 mmol, 1.0 eq) in POCl₃ (150 mL) was stirred for 4 h at 100°C. After the reaction was completed, it was concentrated in *vacuo* to give **Example 24f** (17.4 g, crude), which was used to next step without further purification.

Step 5: Example 24h

[00500] To a solution of **Example 24g** (2.30 g, 23.70 mmol, 2.0 eq) and DIEA (12.23 g, 94.79 mmol, 8.0 eq) in THF (50 mL) was added the solution of **Example 24f** (2.5 g, 11.85 mmol, 1.0 eq) in DCM (25 mL) dropwise at 0°C. The reaction solution was stirred for 30 min at r.t. The reaction solution was diluted with EtOAc (100 mL), washed with brine (80 mL*3), dried over anhydrous Na₂SO₄ and concentrated in *vacuo*. The residue was purified by silica gel chromatography (Petroleum Ether/EtOAc = 1/2) to afford the product **Example 24h** (1.8 g, 64.4% yield) as a yellow solid. LCMS [M+1]⁺ = 236.2

Step 6: Example 24j

[00501] To a solution of **Example 24h** (700 mg, 2.98 mmol, 1.0 eq) and **Example 24i** (611 mg, 2.98 mmol, 1.0 eq) in dry THF (15 mL) was add LiHMDS (5.96 mL, 1M, 2.0 eq) dropwise at -30°C under N₂. The reaction mixture was stirred for 30 min at r.t. The crude product was concentrated and purified by silica gel chromatography (DCM/MeOH = 10/1) to afford the product **Example 24j** (450 mg, 37.5% yield) as a yellow solid. LCMS [M+1]⁺ = 404.2.


Step 7: Example 24l

[00502] To the solution of **Example 24j** (450 mg, 1.12 mmol, 1.0 eq) in dioxane (5 mL) were added Cs₂CO₃ (728 mg, 2.23 mmol, 2.0 eq), **Example 24k** (285 mg, 3.35 mmol, 3.0 eq) and 3rd *t*-Bu-Xphos catalyst (99 mg, 0.11 mmol, 0.1 eq) and the reaction mixture was stirred for 4 h at 110°C under N₂. The reaction solution was concentrated and purified by silica gel chromatography (DCM/MeOH = 10/1) to afford the product **Example 24l** (320 mg, 63.5% yield) as a yellow solid. LCMS [M+1]⁺ = 453.2.


Step 5: Example 24

[00503] To a solution of **Example 24h** (120 mg, 0.27 mmol, 1.0 eq) in THF (3 mL) was added CH₃MgBr (0.27 mL, 3.0 M, 3.0 eq) dropwise at 0°C under N₂. The reaction solution was stirred for 30 min at r.t. The reaction solution was poured into saturated aqueous NH₄Cl (10 mL), and extracted with DCM (10 mL*3). The combined organic layer was washed with brine (30 mL), dried by Na₂SO₄ and concentrated.

The crude product was purified by prep-HPLC to give the desired product **Example 24** (8.8 mg, 8.0% yield) as an off white solid. LCMS $[M+1]^+ = 408.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.52 (s, 1H), 10.57 (s, 1H), 8.58 (s, 1H), 8.12 (s, 1H), 7.72 (d, $J = 7.6$ Hz, 1H), 7.53 (d, $J = 7.2$ Hz, 1H), 7.30 (t, $J = 7.8$ Hz, 1H), 3.96 (s, 3H), 3.71 (s, 3H), 2.81 (s, 3H), 1.09-1.03 (m, 1H), 0.86-0.83 (m, 4H).

Example 25:**Step 1 Example 25**


[00504] To the solution of **Example 25a** (50 mg, 0.11 mmol, 1.0 eq) in THF (1 mL) was added CH_3MgBr (0.55 mL, 1.0 M, 5.0 eq) dropwise at 0°C under N_2 . The reaction solution was stirred for 30 min at r.t.. The reaction solution was purified by prep-TLC (DCM/MeOH = 12/1) to obtain crude product (27 mg) and further purified by prep-HPLC (Prep-C18, 5 μM XBridge column, 19×150 mm, Waters; gradient elution of 40% MeCN in water to 50% MeCN in water over a 7 min period, where both solvents contain 0.1% formic acid) to give the desired product **Example 25** (6.8 mg, 14.6% yield) as a yellow solid (FA salt). LCMS $[M+1]^+ = 422.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 10.55 (s, 1H), 8.58 (s, 1H), 8.48 (br, 2H), 8.10 (s, 1H), 7.71 (dd, $J = 7.8, 1.6$ Hz, 1H), 7.52 (dd, $J = 7.8, 1.6$ Hz, 1H), 7.29 (t, $J = 7.8$ Hz, 1H), 3.95 (s, 3H), 3.72 (s, 3H), 3.36 (q, $J = 7.2$ Hz, 2H), 2.11-2.04 (m, 1H), 1.17 (t, $J = 7.2$ Hz, 3H), 0.85-0.81 (m, 4H).

Example 26:

[00505] To a solution of **Example 26a** (135 mg, 0.3 mmol) in dry THF (10 mL) was added **Example 26b** (1.2 mL, 1M in THF, 1.2 mmol) dropwise at 0°C under nitrogen. After addition (about 5 min), the reaction mixture was stirred for 1 h at r.t.. Then, the reaction was quenched with MeOH (1 mL) and concentrated in *vacuo*. The residue was purified by prep-TLC (DCM/MeOH = 20/1) to afford the product

Example 26 (19.5 mg, 15% yield) as an off-white solid. LCMS $[M+1]^+ = 434.3$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 11.53 (s, 1H), 10.61 (s, 1H), 8.57 (s, 1H), 8.11 (s, 1H), 7.72 (dd, $J = 7.8$ Hz, 1.6 Hz, 1H), 7.52 (dd, $J = 7.8$ Hz, 1.8 Hz, 1H), 7.29 (t, $J = 7.8$ Hz, 1H), 3.95 (s, 3H), 3.72-3.69 (m, 4H), 2.11-2.05 (m, 1H), 1.24-1.16 (m, 4H), 0.85-0.82 (m, 4H).

Example 27:



Step 1: Example 027c

[00506] To a solution of **Example 057a** (5 g, 21.3 mmol) and **Example 027b** (2.28 g, 23.4 mmol) in THF (20 mL) were added Et_3N (3.28 g, 31.9 mmol) and EDCI (6 g, 31.9 mmol). The mixture was stirred at r.t. overnight. The reaction mixture was washed by brine (60 mL), and extracted by DCM (60 mL*3). The combined organic layer was dried over Na_2SO_4 , concentrated under reduced pressure, and dried to give **Example 027c** (4.4 g, 95% yield) as a yellow solid. LCMS $[M+1]^+ = 234.9/236.9$

Step 2: Example 027e

[00507] To a solution of **Example 027c** (4.4 g, 18.7 mmol), **Example 027d** (CAS: 1609394-10-6, 4.2 g, 20.6 mmol) in THF (20 mL) under N_2 was added LiHMDS (1 mol/L, 28 mL, 28.05 mmol) dropwise. The mixture was stirred at r.t. for 1 h. The reaction mixture was washed by brine (60 mL), extracted by DCM (60 mL*3). The combined organic layer was dried over Na_2SO_4 , concentrated under reduced pressure, purified by flash C-18 column ($\text{H}_2\text{O}/\text{CH}_3\text{CN} = 3/1$) to give **Example 027e** (2.4 g, 32% yield) as yellow oil. LCMS $[M+1]^+ = 403.1$

Step 3: Example 027g


[00508] To a solution of **Example 027e** (500 mg, 1.24 mmol) in THF (10 mL) under N_2 , was added **Example 027f** (1 mol/L, 10 mL, 12.4 mmol) in one portion. The mixture was stirred at r.t. for 1 h. The reaction mixture was washed by brine (40 mL), and extracted by EtOAc (40 mL*3). The combined

organic layer was dried over Na_2SO_4 , concentrated under reduced pressure to get crude product (600 mg, crude) as a yellow solid, which was directly used in next step without further purification. LCMS $[\text{M}+1]^+ = 384.1$

Step 4: Example 027

[00509] To a solution of **Example 027g** (100 mg, 0.26 mmol), **Example 027h** (33.3mg 0.92 mmol) in 1,4-dioxane (2 mL) were added Cs_2CO_3 (127.3 mg 0.392 mmol), $\text{Pd}_2(\text{dba})_3$ (23.9mg 0.026mmol), and Xant-phos (15.1 mg 0.026 mmol). The mixture was degassed by Ar, heated to 90°C and stirred over night. The reaction mixture was diluted by DCM and filtrated. The filtrate was concentrated under reduced pressure to remove solvent, which was further purified by prep-HPLC to give **Example 027** (17.5 mg, 15% yield) as a white solid. LCMS $[\text{M}+1]^+ = 433.2$. ^1H NMR (400 MHz, Chloroform-*d*) δ 11.25 (s, 1H), 8.84 (s, 1H), 8.11 (s, 1H), 8.09 (s, 1H), 7.80 (d, $J = 8.0, 1.6$ Hz, 1H), 7.55 (d, $J = 8.0, 1.6$ Hz, 1H), 7.27-7.29 (m, 1H), 4.00 (s, 3H), 3.76 (s, 3H), 2.54-2.60 (m, 1H), 1.67-1.56 (m, 1H), 1.26-1.29 (m, 2H), 1.04-1.10 (m, 4H), 0.87-0.93 (m, 2H).

Example 28:



Step 1: Example 028b

[00510] To a suspension of **Example 028a** (20 g, 96.6 mmol) in MeCN (100 mL) and H₂O (15 mL) was added LiBr (25.2 g, 289.8 mmol), DIPEA (35.4 g, 289.8 mmol). The mixture was stirred at 25°C for 16 h. The reaction mixture was isolated by filtration. The crude solid was washed with MeCN (25 mL), dried under vacuum at 45°C to give **Example 028b** (17.8 g, 93% yield) as a light yellow solid.

Step 2: Example 028d

[00511] To a suspension of **Example 028b** (17.5 g, 88.2 mmol) and **Example 028c** (15.0 g, 73.5 mmol) in H₂O (90 mL) and 2-propanol (15 mL) was added Zn(OAc)₂ (13.4 g, 73.5 mmol). The mixture was stirred at 65°C for 15 h. The reaction mixture was cooled to room temperature and isolated by filtration. The crude solid was washed with water (45 mL) and THF (45 mL). The solid was dried under vacuum at 70°C to give **Example 028d** (25.2 g, 61% yield) as a light yellow solid.

Step 3: Example 28b

[00512] To a suspension of **Example 28d** (2.0 g, 5.10 mmol), CD₃NH₂HCl (431 mg, 6.12 mmol) and NMI (293 mg, 3.57 mmol) in NMP/ACN (20 mL /20 mL) were added HOBt (344 mg, 2.55 mmol) and EDCI (1.37 g, 7.13 mmol). The mixture was stirred at r.t. for 16 h. After cooling to 0°C, the mixture was aged for 2 h at 0°C. The product was isolated by filtration. The wet cake was washed with H₂O (20 mL), and then ACN (20 mL). The solid was collected and dried under vacuum to give **Example 28b** (1.1 g, 57.3% yield) as a white solid. LCMS [M+1]⁺ = 377.0

Step 4: Example 28d

[00513] To a solution of **Example 28b** (1.1 g, 2.92 mmol) and **Example 28c** (512 mg, 4.38 mmol) in 1,4-dioxane (15 mL) were added Pd₂(dba)₃ (267 mg, 0.29 mmol), XantPhos (337 mg, 0.58 mmol) and Cs₂CO₃ (1.9 g, 5.84 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 120°C for 4 h. The reaction was cooled to r.t. and concentrated. The residue was purified by silica gel chromatography (DCM/MeOH = 85/15) to give the desired crude product **Example 28d** (1.08 g, 81.2% yield, ~35% purity) as a yellow solid, which was used for the next step directly. LCMS [M+1]⁺ = 458.0

Step 5: Example 28e


[00514] To a solution of the crude **Example 28d** (1.08 g, 2.36 mmol) in DCM/MeOH (10 mL) was added 6N HCl/iPrOAc (20 mL). The mixture was stirred at r.t. for 16 h. The resulting mixture was concentrated, and the residue was purified by silica gel chromatography (EtOAc/MeOH = 85/15) to give the desired product **Example 28e** (240 mg, 28.4% yield) as a yellow solid. CMS [M+1]⁺ = 358.0

Step 6: Example 28h

[00515] To a solution of **Example 28g** (200 mg, 3.51 mmol) and TEA (709 mg, 7.02 mmol) in DCM (15 mL) was slowly added a solution of **Example 28f** (707 mg, 3.51 mmol) in DCM (5 mL) at 0°C. The mixture was then stirred at r.t. for 1 h. The solvent was removed in *vacuo*, and the residue was purified by silica gel chromatography (Petroleum Ether/EtOAc = 92/8) to give the desired product **Example 28h** (470 mg, 60.4% yield) as a yellow solid. LCMS [M+1]⁺ = 223.0

Step 7: Example 28

[00516] To an ice-cooled solution of NaH (14.5 mg, 0.36 mmol, 60% w.t.% in mineral oil) in DMF (1 mL) under nitrogen atmosphere was added (dropwise) a solution of **Example 28e** (35.7 mg, 0.1 mmol) in DMF (1 mL). After 15 min, a solution of **Example 28h** (20.2 mg, 0.091 mmol) in DMF (1 mL) was added. After 5 min, the cooling bath was removed and the reaction was warmed to 80°C and stirred for 2 h. The mixture was quenched by adding water, purified by prep-HPLC to give the desired product **Example 28** (1.1 mg, 2.8% yield) as a yellow solid. LCMS $[M+1]^+ = 441.0$. $^1\text{H NMR}$ (400 MHz, DMSO- d_6) δ 10.89 (s, 1H), 9.60 (s, 1H), 9.00 (s, 1H), 8.54 (s, 1H), 8.00 (s, 1H), 7.61 (dd, $J = 7.8, 1.6$ Hz, 1H), 7.49 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.24 (t, $J = 7.8$ Hz, 1H), 3.99 (t, $J = 6.4$ Hz, 4H), 3.93 (s, 3H), 3.71 (s, 3H), 2.13 (p, $J = 7.6$ Hz, 2H).


Example 29:**Step 1: Example 29c**

[00517] To a solution of **Example 29b** (1.0 g, 7.30 mmol) in 1,4-dioxane (15 mL) was treated with **Example 29a** (2 mL, 24.1 mmol) and stirred at r.t. for 16 h. After the reaction was completed, the solvent was concentrated, and the residue was suspended in DCM (5 mL) and sonicated. The resulting solid was collected *via* filtration, and dried to afford the desired product **Example 29c** (665 mg, 80% yield) as a white solid. LCMS $[M+1]^+ = 115.0$. $^1\text{H NMR}$ (400 MHz, DMSO- d_6) δ 5.62 (s, 2H), 3.18-3.12 (m, 4H), 1.80-1.68 (m, 4H).

Step 2: Example 29

[00518] To a solution of **Example 29d** (50 mg, 0.13 mmol) and **Example 29c** (22.7 mg, 0.20 mmol) in 1,4-dioxane (1 mL) were added $\text{Pd}_2(\text{dba})_3$ (12.1 mg, 0.013 mmol), XantPhos (15.3 mg, 0.026 mmol) and Cs_2CO_3 (86.5 mg, 0.27 mmol). The mixture was sealed, degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by prep-HPLC to give the desired product **Example 29** (13 mg, 21.7% yield) as a white solid. LCMS $[M+1]^+ = 455.0$. $^1\text{H NMR}$ (400 MHz, DMSO- d_6) δ 10.88 (s, 1H), 9.21 (s, 1H), 8.99 (s, 1H), 8.54 (s, 1H), 7.96 (s, 1H), 7.62 (dd, $J = 7.8, 1.6$ Hz, 1H), 7.49 (dd, $J = 8.0, 1.6$ Hz, 1H), 7.25 (t, $J = 8.0$ Hz, 1H), 3.93 (s, 3H), 3.71 (s, 3H), 3.42-3.37 (m, 4H), 1.81 (s, 4H).

Example 30:




Step 1: Example 30b

[00519] To a suspension of **Example 30a** (1.0 g, 2.55 mmol), $\text{CH}_3\text{NH}_2\cdot\text{HCl}$ (206 mg, 3.06 mmol) and NMI (146 mg, 1.78 mmol) in NMP/ACN (7 mL / 7 mL) were added HOBt (172 mg, 1.27 mmol) and EDCI (685 mg, 3.57 mmol). The mixture was stirred at r.t. for 16 h. After cooling to 0°C , the mixture was aged for 2 h at 0°C . The product was isolated by filtration. The wet cake was washed with H_2O (10 mL), and then ACN (10 mL). The solid was collected and dried under vacuum to give **Example 30b** (417 mg, 43.9% yield) as a white solid. LCMS $[\text{M}+1]^+ = 374.0$

Step 2: Example 30

[00520] To a solution of **Example 68b** (72 mg, 0.19 mmol) and **Example 30c** (23 mg, 0.23 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (17.6 mg, 0.019 mmol), XantPhos (22.3 mg, 0.038 mmol) and Cs_2CO_3 (125.5 mg, 0.38 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 3.5 h in a sealed tube. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by prep-HPLC to give the desired product **Example 30** (16 mg, 19.0% yield) as a white solid. LCMS $[\text{M}+1]^+ = 438.0$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.88 (s, 1H), 9.59 (s, 1H), 9.02 (d, $J = 5.4$ Hz, 1H), 8.54 (s, 1H), 7.99 (s, 1H), 7.61 (d, $J = 7.8$ Hz, 1H), 7.49 (d, $J = 8.0$ Hz, 1H), 7.24 (t, $J = 8.0$ Hz, 1H), 3.98 (s, 4H), 3.93 (s, 3H), 3.71 (s, 3H), 2.84 (d, $J = 4.4$ Hz, 3H), 2.13 (t, $J = 7.8$ Hz, 2H).

Example 31:



Step 1: Example 31b

[00521] A solution of **Example 31a** (2.0 g, 5.4 mmol), TsNH₂ (1.1 g, 6.52 mmol), K₂CO₃ (1.1 g, 6.52 mmol) in MeOH (5 mL) was stirred at r.t. for 10 min, and then **Example 31b** (1.86 g, 13.6 mmol) in dry DMSO (20 mL) was added, which was stirred at 110°C for o.n.. When completed, the reaction was cooled to r.t., diluted by EtOAc, washed by water, and dried over Na₂SO₄. The organic layer was concentrated and purified by silica gel column (Petroleum ether/EtOAc = 4/1) to give **Example 031b** (600 mg, 52.1% yield) as a white solid. LCMS [M+1]⁺ = 214.1. ¹H NMR (400 MHz, Chloroform-*d*) δ 7.73 (d, *J* = 7.8 Hz, 1H), 7.37 (d, *J* = 7.8 Hz, 1H), 3.93 (d, *J* = 6.6 Hz, 4H), 3.76 (d, *J* = 7.4 Hz, 2H), 2.46 (s, 3H).

Step 2: Example 31c

[00522] A solution of Mg (676 mg, 28 mmol) in MeOH (5 mL) was stirred at r.t. for 10 min, then **Example 31b** (600 mg, 2.8 mmol) in MeOH (5 mL) was added, and the mixture reaction was stirred at r.t. for o.n. in a sealed tube. Another Mg (676 mg, 28 mmol) was added, and the mixture reaction was stirred at r.t. for another 6 h. The mixture was filtered, and the solid was washed by MeOH. The filtrate **Example 31c** was used directly for next step. LCMS [M+1]⁺ = 61.2

Step 3: Example 31d

[00523] To a solution of **Example 31c** (crude in MeOH) was added (Boc)₂O (940 mg, 5.6 mmol), which was stirred at r.t. for o.n. The mixture was filtered, and the solid was washed by MeOH. The organic layer was added additional (Boc)₂O (940 mg, 5.6 mmol), which was stirred at r.t. for another 4 h. After reaction was completed, the mixture was concentrated and purified by silica gel column (Petroleum Ether/EtOAc = 5/1) to give **Example 031d** (500 mg, crude, containing (Boc)₂O) as a white solid. LCMS [M+1-56]⁺ = 104.1. ¹H NMR (400 MHz, Chloroform-*d*) 3.93 (d, *J* = 6.6 Hz, 4H), 1.43 (s, 9H).

Step 4: Example 31c

[00524] A solution of **Example 31d** (500 mg, 3.1 mmol) in con. HCl (1 mL) was stirred at r.t. for 0.5 h. After reaction was completed, the mixture was concentrated to give **Example 31c** (200 mg, 67.9% yield) as a white solid, which was used for next step without further purification. LCMS [M+1]⁺ = 60.2.

Step 5: Example 31f


[00525] To a solution of **Example 31c** (200 mg, 3.3 mmol) and DIEA (3.3 g, 33 mmol) in 1,4-dioxane (5 mL) was added **Example 31e** (230 mg, 1.67 mmol), which was stirred at r.t. for o.n. After reaction was completed, the mixture was concentrated, and directly purified by silica gel column (DCM/MeOH = 10/1) to give the desired product **Example 31f** (230 mg, crude, containing DIEA, 67.6% yield) as colorless oil. LCMS [M]⁺ = 103.2

Step 6: Example 31

[00526] To a solution of **Example 31f** (200 mg, ~50% purity, 0.5 mmol) and **Example 31g** (188 mg, 0.5 mmol) in 1,4-dioxane (5 mL) were added Pd₂(dba)₃ (45 mg, 0.05 mmol), XantPhos (30 mg, 0.05 mmol) and Cs₂CO₃ (489 mg, 1.5 mmol). The mixture was degassed by nitrogen for 3 times, sealed, and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by prep-HPLC to give the desired product **Example 31** (2.2 mg, 1% yield) as a white solid.

LCMS $[M+1]^+ = 442.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 10.90 (s, 1H), 9.60 (s, 1H), 9.00 (s, 1H), 8.54 (s, 1H), 8.00 (s, 1H), 7.61 (d, $J = 7.8$ Hz, 1H), 7.49 (d, $J = 8.0$ Hz, 1H), 7.24 (t, $J = 8.0$ Hz, 1H), 3.97 (s, 4H), 3.93 (s, 3H), 3.71 (s, 3H).

Example 32:



Step 1: Example 32c


[00527] To a solution of **Example 32a** (300 mg, 2.19 mmol) in 1,4-dioxane (5 mL) were treated with **Example 32b** (623 mg, 3.28 mmol) and TEA (664 mg, 6.57 mmol). The mixture was stirred at r.t. for 16 h. After completion, the mixture was concentrated, and the residue was suspended in DCM (5 mL) and sonicated. The resulting solid was collected via filtration, and dried to afford the desired product **Example 32c** (226 mg, 73.9% yield) as a white solid.

LCMS $[M+1]^+ = 141.0$

Step 2: Example 32

[00528] To a solution of **Example 32d** (60 mg, 0.16 mmol) and **Example 32c** (26.7 mg, 0.19 mmol) in 1,4-dioxane (0.6 mL) were added $\text{Pd}_2(\text{dba})_3$ (14.6 mg, 0.016 mmol), XantPhos (18.4 mg, 0.032 mmol) and Cs_2CO_3 (104 mg, 0.32 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 16 h in a sealed tube. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by prep-HPLC to give the desired product **Example 32** (6.6 mg, 8.7% yield) as a white solid. LCMS $[M+1]^+ = 481.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 10.87 (s, 1H), 9.61 (s, 1H), 8.99 (s, 1H), 8.55 (s, 1H), 7.97 (s, 1H), 7.61 (d, $J = 7.8$ Hz, 1H), 7.48 (d, $J = 8.0$ Hz, 1H), 7.24 (t, $J = 8.0$ Hz, 1H), 3.93 (s, 7H), 3.69 (s, 3H), 2.08 (t, $J = 7.5$ Hz, 4H), 1.80 -1.67 (m, 2H).

Example 33:




Step 1: Example 33c

[00529] To a solution of **Example 33a** (500 mg, 3.65 mmol, 1.0 eq) in dioxane (6 ml) was added **Example 33b** (624 mg, 10.95 mmol, 3.0 eq). The reaction solution was stirred for 16 h at r.t. and concentrated. The residue was treated with DCM. The resulting solid was collected by filtration to afford **Example 33c** (190 mg, 52% yield) as a white solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ 5.72 (brs, 2H), 3.76 (t, *J* = 7.6 Hz, 4H), 2.14-2.04 (m, 2H).

Step 2: Example 33

[00530] To a solution of **Example 33d** (100 mg, 0.26 mmol, 1.0 eq) and **Example 33c** (78.0 mg, 0.78 mmol, 3.0 eq) in dioxane (4 mL) were added Xantphos (30.1 mg, 0.05 mmol, 0.2 eq), Cs₂CO₃ (169.5 mg, 0.52 mmol, 2.0 eq) and Pd₂(dba)₃.CHCl₃ (26.9 mg, 0.03 mmol, 0.1 eq). The reaction mixture was stirred at 110°C for 4 h under N₂ protection. The solid was filtered out and filtrate was concentrated, and the residue was purified by prep-TLC (DCM/MeOH = 15/1) to afford the **Example 33** (43.3 mg, 37% yield) as an off-white solid. LCMS [M+1]⁺ = 448.2. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.00 (s, 1H), 9.20 (s, 1H), 9.05 (s, 1H), 8.56 (s, 1H), 7.91 (s, 1H), 7.62 (dd, *J* = 7.8, 1.6 Hz, 1H), 7.53 (dd, *J* = 8.0, 1.6 Hz, 1H), 7.25 (t, *J* = 7.8 Hz, 1H), 4.00-3.70 (m, 7H), 3.70 (s, 3H), 3.01-2.91 (m, 1H), 2.19-2.09 (m, 2H), 1.12-0.99 (m, 4H).

Example 34:

Step 1: Example 34b

[00531] To a solution of **Example 34a** (10.0 g, 42.7 mmol, 1.0 eq) in DCM (100 mL) was added BBr_3 (11.8 g, 46.9 mmol, 1.1 eq) dropwise at 0°C . The mixture was stirred for 2 h at r.t.. The mixture was quenched with MeOH (12 mL), washed with brine (50 mL*3), dried over Na_2SO_4 , and concentrated to afford the product **Example 34b** (9.2 g, 98% yield) as an orange solid. LCMS $[\text{M}+1]^+ = 221.2$.

Step 2: Example 34c

[00532] To a solution of **Example 34b** (9.2 g, 41.8 mmol, 1.0 eq) in ACN (184 mL) were added K_2CO_3 (11.6 g, 83.6 mmol, 2.0 eq) and CD_3I (7.3 g, 50.2 mmol, 1.2 eq). The reaction mixture was stirred for 6 h at 80°C . After the reaction was completed, the reaction mixture was concentrated and purified by silica gel chromatography (Petroleum Ether/EtOAc = 2/1) to afford the product **Example 34c** (3.9 g, 39% yield) as an orange solid. LCMS $[\text{M}+1]^+ = 238.2$.

Step 3: Example 34d

[00533] To a solution of **Example 34c** (3.9 g, 16.4 mmol, 1.0 eq) in MeOH (78 mL) was added Pd/C (780 mg) under N_2 protection. The suspension was degassed under vacuum and purged with H_2 three times. The reaction mixture was stirred at r.t. for 48 h under H_2 balloon. The solid was filtered out, and the filtrate was concentrated. The residue was purified by silica gel chromatography (Petroleum Ether/EtOAc = 1/1) to afford the product **Example 34d** (1.9 g, 56% yield) as an off-white solid. LCMS $[\text{M}+1]^+ = 208.2$.

Step 4: Example 34f

[00534] To a solution of **Example 34e** (450 mg, 1.91 mmol, 1.0 eq) and **Example 34d** (396 mg, 1.91 mmol, 1.0 eq) in dry THF (15 mL) was added LiHMDS (3.8 mL, 1 M in THF, 2.0 eq) dropwise at -20°C under N_2 protection. The reaction mixture was stirred for 0.5 h at r.t., and then the silica was added to the mixture and concentrated. The residue was purified by silica gel chromatography (DCM/MeOH = 20/1) to afford the product **Example 34f** (450 mg, 58% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 406.2$.

Step 5: Example 34h


[00535] To a solution of **Example 34f** (200 mg, 0.49 mmol, 1.0 eq) in THF (4 mL) was added **Example 34g** (3.0 mL, 1.0 M in THF, 6.0 eq) dropwise at 0°C under N_2 protection. The mixture was stirred for 0.5 h at r.t.. The reaction was poured into saturated aqueous NH_4Cl (10 mL) and extracted with EtOAc (15 mL*3). The combined organic layer was washed with brine, dried over Na_2SO_4 , and concentrated. The crude product was purified by silica gel chromatography (DCM/MeOH = 20/1) to afford the product **Example 34h** (160 mg, 84% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 387.1$.

Step 6: Example 34

[00536] To a solution of **Example 34h** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (4 mL) were added Cs_2CO_3 (169 mg, 0.52 mmol, 2.0 eq), **Example 34i** (66 mg, 0.77 mmol, 3.0 eq), Xantphos (30 mg, 0.05 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (27 mg, 0.03 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 34** (30.9 mg, 27% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 436.2$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.01 (s, 1H),

10.93 (s, 1H), 9.13 (s, 1H), 8.56 (s, 1H), 8.04 (s, 1H), 7.64 (d, $J = 7.6$ Hz, 1H), 7.52 (d, $J = 7.8$ Hz, 1H), 7.26 (t, $J = 7.8$ Hz, 1H), 3.95 (s, 3H), 3.05-2.90 (m, 1H), 2.05-2.00 (m, 1H), 1.14-1.03 (m, 4H), 0.81 (d, $J = 6.4$ Hz, 4H).

Example 35:



Step 1: Example 35c

[00537] To a solution of **Example 35a** (10.0 g, 94.79 mmol, 1.0 eq) in CH_3CN (50 mL) was added **propan-2-amine** (5.59 g, 94.79 mmol, 1.0 eq) at 0°C . The reaction solution was stirred for 2 h at room temperature. The resulting solid was collected by filtration to afford **Example 35b** (8.07 g) as a white solid. The solid was dissolved in THF (150 mL), followed by NaH (5.83 g, 60% in mineral oil, 145.8 mmol, 3.0 eq) in portions at 0°C . The mixture was stirred for 16 h at room temperature. The reaction was quenched with H_2O and extracted with EtOAc. The combined organic layer was washed with brine, dried by Na_2SO_4 and concentrated. The crude product was purified by silica gel chromatography (DCM/MeOH = 30/1) to give the desired product **Example 35c** (3.50 g, 29% yield) as a white solid. ^1H NMR (300 MHz, CDCl_3 - d) δ 5.07 (s, 1H), 4.21-4.06 (m, 1H), 3.38 (s, 4H), 1.13 (d, $J = 6.6$ Hz, 6H).

Step 2: Example 35f

[00538] To a solution of **Example 35d** (10.0 g, 52.08 mmol, 1.0 eq) in DCM (200 mL) were added DIEA (33.59 g, 260.4 mmol, 5.0 eq) and HATU (23.75 g, 62.50 mmol, 1.2 eq). After stirred for 30 min, **Example 35e** (6.09 g, 62.50 mmol, 1.2 eq) was added to the solution. The reaction solution was stirred for 1 h at room temperature. After the reaction was completed, the solvent was removed and the crude was purified by silica gel chromatography (Petroleum Ether/EtOAc = 3/1) to give the desired product **Example 35f** (11.5 g, 94.0% yield) as a yellow solid. LCMS $[M+1]^+ = 235.0$.

Step 3: Example 35h

[00539] To a solution of **Example 35f** (5.0 g, 21.28 mmol, 1.0 eq) and **Example 35g** (4.36 g, 21.28 mmol, 1.0 eq) in THF (100 mL) was added LiHMDS (42.55 mL, 1 M in THF, 2.0 eq) dropwise at -15°C under N_2 . The reaction solution was stirred for 1 h at r.t.. And then the silica was added to the mixture and concentrated. The residue was purified by silica gel chromatography (DCM/MeOH = 50/1) to give the desired product **Example 35h** (4.02 g, 46.9% yield) as a yellow solid. LCMS $[M+1]^+ = 403.2$.


Step 4: Example 35j

[00540] To a solution of **Example 35h** (2.0 g, 4.98 mmol, 1.0 eq) in THF (100 mL) was added **Example 35i** (39.80 mL, 39.80 mmol, 1 M, 8.0 eq) dropwise at -10°C under N_2 . The reaction solution was stirred for 1 h at r.t. The reaction solution was quenched with saturated aqueous NH_4Cl and extracted with EtOAc. The combined organic layer was washed with brine, dried by Na_2SO_4 and concentrated. The crude product was purified by silica gel chromatography (DCM/MeOH = 20/1) to give the desired product **Example 35j** (1.90 g, 99.7% yield) as a yellow solid. LCMS $[M+1]^+ = 384.2$.

Step 5: Example 35

[00541] To the solution of **Example 35j** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (3 mL) were added **Example 70c** (100 mg, 0.78 mmol, 3.0 eq), Cs_2CO_3 (170 mg, 0.52 mmol, 2.0 eq), Xantphos (15 mg, 0.026 mmol, 0.1 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (27 mg, 0.026 mmol, 0.1 eq). The reaction solution was stirred for 4 h at 110°C under N_2 . The reaction solution was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 35** (31.3 mg, 25.3% yield) as an off white solid. LCMS $[M+1]^+ = 476.3$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 10.93 (s, 1H), 9.14 (s, 1H), 8.56 (s, 1H), 8.17 (s, 1H), 7.62 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.56 (dd, $J = 8.4$ Hz, 1.6 Hz, 1H), 7.26 (t, $J = 7.8$ Hz, 1H), 4.12-3.99 (m, 3H), 3.95 (s, 3H), 3.70 (s, 3H), 3.41 (t, $J = 8.0$ Hz, 2H), 3.04-2.91 (m, 1H), 1.17-1.06 (m, 8H), 1.06-0.98 (m, 2H).


Example 36:



[00542] To a solution of **Example 36a** (70 mg, 0.18 mmol, 1.0 eq) in dioxane (2 mL) were added **Example 36b** (48 mg, 0.55 mmol, 3.0 eq), Cs_2CO_3 (119 mg, 0.37 mmol, 2.0 eq), Xantphos (11 mg, 0.018 mmol, 0.1 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (19 mg, 0.018 mmol, 0.1 eq). The reaction solution was stirred for 4 h at 110°C under N_2 . The reaction solution was filtered and the filtrate was concentrated. The crude product was purified by prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 36** (30.9 mg, 39.0% yield) as an off-white solid.


LCMS $[\text{M}+1]^+ = 435.2$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.02 (s, 1H), 10.56 (s, 1H), 9.12 (s, 1H), 8.56 (s, 1H), 8.08 (s, 1H), 7.66 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.56 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.29 (t, $J = 7.4$ Hz, 1H), 3.95 (s, 3H), 3.70 (s, 3H), 3.06-2.94 (m, 1H), 2.83-2.69 (m, 1H), 1.15-1.08 (m, 2H), 1.08-0.98 (m, 8H).

Example 37:



[00543] To a solution of **Example 37a** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (3 mL) were added **Example 37b** (49 mg, 0.52 mmol, 3.0 eq), Cs_2CO_3 (170 mg, 0.52 mmol, 2.0 eq), and 3^{rd} *t*-Bu-Xphos Pd (23 mg, 0.026 mmol, 0.1 eq). The reaction solution was stirred for 6 h at 110°C under N_2 . The reaction solution was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 37** (21.0 mg, 18.3% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 442.2$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.08 (s, 1H), 9.98 (s, 1H), 9.10 (s, 1H), 8.56 (s, 1H), 8.19 (d, $J = 4.8$ Hz, 1H), 8.01 (s, 1H), 7.69 (t, $J = 8.0$ Hz, 2H), 7.60 (t, $J = 8.0$ Hz, 2H), 7.31 (t, $J = 8.0$ Hz, 1H), 6.96-6.88 (m, 1H), 3.96 (s, 3H), 3.72 (s, 3H), 3.04-2.91 (m, 1H), 1.14-0.94 (m, 4H).

Example 38:

**Step 1: Example 38c**


[00544] A solution of **Example 38b** (1.13 g, 8.2 mmol) in 1,4-dioxane (10 mL) were treated with **Example 38a** (1.0 g, 8.2 mmol) and TEA (2.5 g, 24.7 mmol). The mixture was stirred at r.t. for 16 h. After the reaction was completed, the solution was concentrated, and the residue was suspended in a mixed solution (Petroleum Ether/EtOAc = 1/1, 10 mL), and sonicated. The resulting solid was collected via filtration, dried to afford the desired product **Example 38c** (1.0 g, 94.3% yield) as a white solid, which contained some TEA.HCl. LCMS $[M+1]^+ = 129.0$

Step 2: Example 38

[00545] To a solution of **Example 38d** (250 mg, 0.66 mmol) and **Example 38c** (127 mg, 0.99 mmol) in 1,4-dioxane (4 mL) were added $\text{Pd}_2(\text{dba})_3$ (60 mg, 0.068 mmol), Xantphos (77.5 mg, 0.13 mmol) and Cs_2CO_3 (433 mg, 1.32 mmol). The mixture was sealed, degassed by nitrogen for 3 times and stirred at 110°C for 16 h. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by prep-HPLC to give the desired product **Example 38** (116 mg, 37.4% yield) as a yellow solid. LCMS $[M+1]^+ = 469.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 10.91 (s, 1H), 9.58 (s, 1H), 8.99 (s, 1H), 8.54 (s, 1H), 8.02 (s, 1H), 7.62 (d, $J = 7.7$ Hz, 1H), 7.49 (d, $J = 7.8$ Hz, 1H), 7.24 (t, $J = 7.9$ Hz, 1H), 3.93 (s, 3H), 3.72 (s, 3H), 3.66 (s, 4H), 1.17 (s, 6H).


Examples 89-104 were synthesized as described for examples 1-88.

Example 105:




[00546] To a solution of **Example 105a** (100 mg, 0.26 mmol, 1.0 eq), **Example 105b** (57.0 mg, 0.78 mmol, 3.0 eq) in dioxane (3 mL) were added Cs_2CO_3 (169.5 mg, 0.52 mmol, 2.0 eq) and $3^{\text{rd}}\text{-t-Bu-xphos-Pd}$ (26.7 mg, 0.03 mmol, 0.1 eq). The reaction mixture was stirred at 110 °C for 4 h under N_2 protection. The solid was filtered out and filtrate was concentrated. The residue was purified by Prep-TLC (DCM/MeOH = 15/1) to afford **Example 105** (36.1 mg, 33% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 421.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$) δ 11.02 (s, 1H), 10.56 (s, 1H), 9.12 (s, 1H), 8.57 (s, 1H), 8.07 (s, 1H), 7.66 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.55 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.28 (t, $J = 7.8$ Hz, 1H), 3.95 (s, 3H), 3.70 (s, 3H), 3.06 – 2.93 (m, 1H), 2.39 (q, $J = 7.5$ Hz, 2H), 1.16 – 0.96 (m, 7H).

Example 106:




[00547] To a solution of **Example 106a** (50 mg, 0.13 mmol, 1.0 eq) and **Example 106b** (13.9 mg, 0.20 mmol, 1.5 eq) in dioxane (3 mL) were added Cs_2CO_3 (84.9 mg, 0.26 mmol, 2.0 eq), $\text{Pd}_2(\text{dba})_3\cdot\text{CHCl}_3$ (13.5 mg, 0.013 mmol, 0.1 eq) and Xantphos (15.1 mg, 0.026 mmol, 0.2 eq). The reaction mixture was stirred for 2 h at 110°C under N_2 protection. The mixture was filtered and the filtrate was concentrated in *vacuo*. The crude product was purified by pre-HPLC (DCM/MeOH = 25/1) to afford the desired product **Example 106** (5.9 mg, 10.8% yield) as a white solid. LCMS $[\text{M}+1]^+ = 419.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$) δ 11.04 (s, 1H), 10.86 (s, 1H), 9.15 (s, 1H), 8.57 (s, 1H), 8.17 (s, 1H), 7.67 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.59-7.56(m, 1H), 7.29 (t, $J = 7.8$ Hz, 1H), 6.61 (dd, $J = 16.8, 10.2$ Hz, 1H), 6.28 (dd, $J = 16.8, 1.8$ Hz, 1H), 5.81(d, $J = 11.7$ Hz, 1H), 3.96 (s, 3H), 3.71 (s, 3H), 3.05 - 2.95 (m, 1H), 1.15 - 0.95 (m, 4H).

Example 107:




[00548] To a solution of **Example 107a** (110 mg, 0.29 mmol) and **Example 107b** (93 mg, 0.86 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (26 mg, 0.029 mmol), Xantphos (33 mg, 0.057 mmol) and Cs_2CO_3 (187 mg, 0.57 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 1 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 107** (24 mg, 18.5% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 456.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 11.04 (s, 1H), 9.88 (s, 1H), 9.07 (s, 1H), 8.55 (s, 1H), 8.03 (d, $J = 5.1$ Hz, 1H), 7.97 (s, 1H), 7.66 (d, $J = 8.1$ Hz, 1H), 7.57 (d, $J = 7.8$ Hz, 1H), 7.39 (s, 1H), 7.27 (t, $J = 8.0$ Hz, 1H), 6.74 (d, $J = 4.9$ Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.94 (s, 1H), 2.24 (s, 3H), 1.05 (s, 2H), 0.99 (d, $J = 7.6$ Hz, 2H).

Example 108:




[00549] To a solution of **Example 108a** (100 mg, 0.26 mmol) and **Example 108b** (44 mg, 0.39 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 108** (18 mg, 15.1% yield) as a white solid. LCMS $[\text{M}+1]^+ = 460.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 11.01 (s, 1H), 10.15 (s, 1H), 9.11

(s, 1H), 8.54 (s, 1H), 8.20 (t, $J = 7.5$ Hz, 1H), 7.69 (s, 1H), 7.63 (q, $J = 8.2, 5.8$ Hz, 3H), 7.28 (t, $J = 8.0$ Hz, 1H), 6.83 (s, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.95 (s, 1H), 1.07 (s, 2H), 1.00 (d, $J = 7.4$ Hz, 2H).


Example 109:**Example 109a****Example 109**

To a solution of **Example 109a** (100 mg, 0.26 mmol) and **Example 109b** (30 mg, 0.31 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 109** (25 mg, 21.7% yield) as a white solid. LCMS $[\text{M}+1]^+ = 443.0$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.04 (s, 1H), 10.34 (s, 1H), 9.10 (s, 1H), 8.77 (d, $J = 4.4$ Hz, 1H), 8.56 (s, 1H), 8.07 (d, $J = 9.1$ Hz, 1H), 7.72 (s, 1H), 7.62 (dd, $J = 12.0, 8.1$ Hz, 2H), 7.55 (dd, $J = 9.2, 4.5$ Hz, 1H), 7.24 (t, $J = 7.9$ Hz, 1H), 3.93 (s, 3H), 3.70 (s, 3H), 2.95 (t, $J = 7.4$ Hz, 1H), 1.06 (d, $J = 4.1$ Hz, 2H), 1.00 (d, $J = 7.4$ Hz, 2H).

Example 110:**Example 110a****Step 1****Example 110c****Step 2****Example 110****Step 1: Example 110c**


[00550] To a stirred solution of **Example 110a** (1.0 g, 5.78 mmol), xphos (551 mg, 1.16 mmol), $\text{Pd}_2(\text{dba})_3$ (530 mg, 0.58 mmol) and **Example 110b** (2.5 g, 28.9 mmol) in THF (12 mL) was added LiHMDS (32 mL, 1.0 M in THF) and the resulting reaction was heated to 65°C and stirring continued for 2 h. The reaction was poured into water (100 mL) and extracted with EtOAc (150 mL*2) and DCM (150

Example 112:




[00553] A solution of **Example 112a** (100 mg, 0.26 mmol), **Example 112b** (80 mg, 0.78 mmol), Pd₂(dba)₃ (24 mg, 0.026 mmol), Xantphos (8 mg, 0.013 mmol) and Cs₂CO₃ (170 mg, 0.52 mmol) in dioxane (3 mL) was heated to 110°C for 2 h in Ar atmosphere. This mixture was filtered and directly purified by prep-HPLC to afford **Example 112** (34 mg, 28.6% yield) as a yellow solid. LCMS [M+1]⁺ = 456.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.12 (s, 1H), 9.91 (s, 1H), 9.06 (s, 1H), 8.53 (s, 1H), 8.25 (s, 1H), 7.68 (d, *J* = 8.0 Hz, 1H), 7.60 (d, *J* = 7.8 Hz, 1H), 7.52 (t, *J* = 7.8 Hz, 1H), 7.26 (t, *J* = 7.9 Hz, 1H), 7.15 (d, *J* = 8.3 Hz, 1H), 6.73 (d, *J* = 7.3 Hz, 1H), 3.92 (s, 3H), 3.69 (s, 3H), 2.98 – 2.89 (m, 1H), 2.28 (s, 3H), 1.09 – 0.94 (m, 4H).

Example 113:




[00554] A solution of **Example 113a** (100 mg, 0.26 mmol), **Example 113b** (30 mg, 0.26 mmol), Pd₂(dba)₃ (24 mg, 0.026 mmol), Xantphos (8 mg, 0.013 mmol) and Cs₂CO₃ (170 mg, 0.52 mmol) in DMA (3 mL) was heated to 130°C for 2 h in Ar atmosphere. This mixture was filtered and directly purified by prep-HPLC to afford **Example 113** (13 mg, 10.9% yield) as a yellow solid. LCMS [M+1]⁺ = 457.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.04 (s, 1H), 10.21 (s, 1H), 9.09 (s, 1H), 8.54 (s, 1H), 7.97 (d, *J* = 9.1 Hz, 1H), 7.62 (dd, *J* = 15.5, 8.4 Hz, 3H), 7.43 (d, *J* = 9.2 Hz, 1H), 7.25 (t, *J* = 8.0 Hz, 1H), 3.93 (s, 3H), 3.70 (s, 3H), 2.94 (t, *J* = 5.6 Hz, 1H), 1.09 – 0.96 (m, 4H).

0.054 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (28 mg, 0.027 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 . The mixture was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 115** (40.3 mg, 32.3% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 464.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 10.96 (s, 1H), 8.89 (s, 1H), 8.56 (s, 1H), 8.15 (s, 1H), 7.62 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.56 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.26 (t, $J = 7.8$ Hz, 1H), 4.09 - 3.96 (m, 3H), 3.95 (s, 3H), 3.72 (s, 3H), 3.39 (t, $J = 8.1$ Hz, 2H), 3.10 (q, $J = 7.2$ Hz, 2H), 1.19 - 1.03 (m, 9H).

Example 116:

[00558] To a solution of **Example 116a** (100 mg, 0.27 mmol, 1.0 eq) in dioxane (5 mL) were added **Example 116b** (69 mg, 0.81 mmol, 3.0 eq), Cs_2CO_3 (175 mg, 0.54 mmol, 2.0 eq), Xantphos (31 mg, 0.054 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (28 mg, 0.027 mmol, 0.1 eq). The reaction solution was stirred for 4 h at 110°C under N_2 . The reaction solution was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 116** (40.7 mg, 36.0% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 421.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.04 (s, 1H), 10.90 (s, 1H), 8.88 (s, 1H), 8.56 (s, 1H), 8.00 (s, 1H), 7.65 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.52 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.26 (t, $J = 7.8$ Hz, 1H), 3.95 (s, 3H), 3.71 (s, 3H), 3.12 (q, $J = 7.2$ Hz, 2H), 2.07-1.96 (m, 1H), 1.13 (t, $J = 7.2$ Hz, 3H), 0.79 (d, $J = 6.0$ Hz, 4H).

Example 117:

Step 1: Example 117b

[00559] To a solution of dimethyl 3-oxopentanedioate (5.0 g, 28.7 mmol) and Ni(acac)₂ (738 mg, 2.87 mmol) in dioxane (30 mL) was added NH₂-CN (3.6 g, 86.2 mmol). The mixture was stirred at 110°C for o.n. The reaction was cooled to r.t.. The mixture was filtered and the filter cake was collected, washed with MeOH (20 mL) and concentrated in *vacuo* to give the desired product **Example 117b** (3.0 g, 56.6% yield) as a yellow solid. LCMS [M+1]⁺ = 185.0

Step 2: Example 117c

[00560] To a solution of **Example 117b** (2.5 g, 13.58 mmol) in POCl₃ (15 mL) was added DIEA (2 mL) at 0°C, which was heated to 30°C and stirred for o.n. The reaction was concentrated in *vacuo*. H₂O (15 mL) and MeOH (3 mL) were added at 0°C, which was stirred at r.t. for 1 h. The mixture was filtered and the filtrate cake was collected by filtration to give **Example 117c** (1.5 g, 50.1% yield) as a yellow solid. LCMS [M+1]⁺ = 221.0

Step 3: Example 117e

[00561] To a solution of **Example 117c** (1.2 g, 5.42 mmol) and **Example 117d** (1.21 g, 5.96 mmol) in EtOH (30 mL) was added conc. HCl (5 mL) and the solution was heated to reflux for o.n. The reaction was concentrated in *vacuo*. The residue was dissolved in EtOAc (100 mL) and H₂O (50 mL), and the pH was adjusted to ~8 with sat. NaHCO₃. The organic layer was separated and concentrated. The residue was purified by silica gel chromatography (DCM/MeOH = 30/1) to afford the product **Example 117e** (700 mg, 33.1% yield) as a yellow solid. LCMS [M+1]⁺ = 389.11.

Step 4: Example 117f

[00562] To a solution of **Example 117e** (690 mg, 1.78 mmol) in THF (30 mL) and H₂O (10 mL) cooled at 0°C was added LiOH.H₂O (112 mg, 2.67 mmol) and the solution was stirred at r.t. for o.n.. The reaction was concentrated in *vacuo*. The residue was dissolved in H₂O (50 mL), adjusted pH ~4 with HCl (2 mol/L), and extracted with EtOAc (100 mL). The organic layer was concentrated to afford the crude product **Example 117f** (750 mg, 100% crude yield) as a yellow solid. LCMS [M+1]⁺ = 375.2

Step 5: Example 117g


[00563] To a solution of **Example 117f** (500 mg, 1.33 mmol) in DMF (10 mL) were added DIEA (515 mg, 3.99 mmol), HATU (610 mg, 1.60 mmol) and CD₃-NH₂.HCl (110 mg, 1.59 mmol) and the solution was stirred at r.t. for 2 h. The reaction was diluted with EtOAc (50 mL), washed with brine (10 mL*3), dried over Na₂SO₄, and concentrated in *vacuo*. The residue was purified by silica gel chromatography (DCM/MeOH = 30/1) to afford the product **Example 117g** (510 mg, 98% yield) as a light yellow solid. LCMS [M+1]⁺ = 391.0

Step 6: Example 117

[00564] To a mixture of **Example 117g** (500 mg, 1.278 mmol), **Example 117h** (33 mg, 1.917 mmol) and Cs₂CO₃ (167 mg, 2.55 mmol) in dioxane (5 mL) was added 3rd t-Bu-Xphos-Pd (22.5 mg, 0.128 mmol). The mixture was degassed with N₂ three times, then heated to 100°C and stirred for overnight. The reaction was concentrated in *vacuo*. The residue was further purified by prep-HPLC to give the desired product **Example 117** (60 mg, 54.2% yield) as a white solid. LCMS [M+1]⁺ = 440.1. ¹H NMR (300


MHz, DMSO-*d*₆) δ 9.29 (s, 1H), 8.56 (s, 1H), 8.43 (s, 1H), 7.74 (d, *J* = 6.8 Hz, 1H), 7.39 (d, *J* = 7.6 Hz, 1H), 7.29 (t, *J* = 7.8 Hz, 1H), 3.94 (s, 3H), 3.70 (s, 3H), 1.65-1.80 (m, 1H), 0.8-0.94 (m, 4H).

Example 118:



[00565] To the solution of **Example 118a** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (5 mL) were added **Example 118b** (95 mg, 0.78 mmol, 3.0 eq), Cs₂CO₃ (170 mg, 0.52 mmol, 2.0 eq), Xantphos (15 mg, 0.026 mmol, 0.1 eq) and Pd₂(dba)₃CHCl₃ (27 mg, 0.026 mmol, 0.1 eq). The reaction solution was stirred for 4 h at 110°C under N₂. The reaction solution was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the desired product **Example 118** (21.9 mg, 17.9% yield) as a light yellow solid. LCMS [M+1]⁺ = 469.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.94 (s, 1H), 9.04 (s, 1H), 8.59 (s, 1H), 7.69 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.52 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.30 (t, *J* = 7.8 Hz, 1H), 6.64 (s, 1H), 3.95 (s, 3H), 3.69 (s, 3H), 3.00 - 2.89 (m, 1H), 1.22 (s, 9H), 1.13 - 1.00 (m, 4H).

Example 119:





Step 1: Example 119

[00566] To a solution of **Example 119a** (100 mg, 0.26 mmol) and **Example 119b** (44 mg, 0.39 mmol) in 1,4-dioxane (2 mL) were added Pd₂(dba)₃ (24 mg, 0.026 mmol), xantphos (30 mg, 0.052 mmol) and Cs₂CO₃ (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 119** (19.7 mg, 16.6% yield) as a yellow solid. LCMS [M+1]⁺ = 460.0. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.05 (s, 1H), 9.40 (s, 1H), 9.06 (s, 1H), 8.54 (s, 1H), 8.05 (d, *J* = 4.6 Hz, 1H), 7.92 (s, 1H), 7.65 (q, *J* = 8.1, 5.8 Hz, 2H), 7.59 (d, *J* = 7.8

Hz, 1H), 7.29 (t, J = 8.0 Hz, 1H), 7.04 (dt, J = 8.0, 3.9 Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.95 (dt, J = 8.1, 3.7 Hz, 1H), 1.07 (t, J = 3.7 Hz, 2H), 0.99 (dd, J = 7.9, 4.5 Hz, 2H).

Example 120:



Step 1: Example120b

[00567] To a solution of **Example 120a** (5.37 g, 47.9 mmol, 1.0 eq) in ACN (250 mL) was added NBS (6.2 g, 52.7 mmol, 1.1 eq) at 0 °C under N₂ protection. The reaction solution was stirred at r.t. for 1 h, and then the mixture was concentrated. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 5/1) to afford the product **Example 120b** (6.8 g, 75% yield) as a white solid. LCMS [M+1]⁺ = 190.2.

Step 2: Example120c


[00568] To a solution of **Example120b** (6.0 g, 31.4 mmol, 1.0 eq) in dry THF (150 mL) was added n-BuLi (44 mL, 2.5 M in THF, 110 mmol, 3.5 eq) dropwise at -78 °C under N₂ protection. The mixture was stirred for 5 min at the same temperature. Acetone (18.2 g, 314 mmol, 10.0 eq) in THF (50 mL) was added dropwise at -78 °C under N₂ protection. The mixture was warmed to r.t. and stirred for 30 min. The reaction was poured into saturated aqueous NH₄Cl (100 mL) and extracted with EtOAc (200 mL*2). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, and concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 5/1) to afford the product **Example 120c** (3.5 g, 65% yield) as a green solid. LCMS [M+1]⁺ = 171.2.

Step 3: Example 120

[00569] To a solution of **Example 120d** (100 mg, 0.26 mmol, 1.0 eq.) in dioxane (3 mL) were added Cs₂CO₃ (169 mg, 0.52 mmol, 2.0 eq), **Example 120c** (132 mg, 0.77 mmol, 3.0 eq), Xantphos(29 mg, 0.05 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃(31 mg, 0.03 mmol, 0.1 eq). The reaction mixture was stirred for

3 h at 110 °C under N₂ protection. After cooled to room temperature, the solvent was removed, the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 120** (48.4 mg, 36% yield) as an off-white solid. LCMS [M+1]⁺ = 518.4. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.11 (s, 1H), 10.14 (s, 1H), 9.11 (s, 1H), 8.56 (s, 1H), 7.97 (dd, *J* = 10.8, 8.4 Hz, 1H), 7.80 (s, 1H), 7.73 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.62 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.49 (dd, *J* = 8.1, 1.8 Hz, 1H), 7.27 (t, *J* = 7.8 Hz, 1H), 5.26 (s, 1H), 3.95 (s, 3H), 3.72 (s, 3H), 3.05 - 2.90 (m, 1H), 1.46 (s, 6H), 1.16 - 0.94 (m, 4H).

Example 121:




Step 1: Example 121c

[00570] To a solution of **Example 121a** (200 mg, 1.5 mmol, 1.0 eq) in dioxane (3 mL) was added **Example 121b** (265.5 mg, 4.5 mmol, 3.0 eq). The reaction solution was stirred for 16 h at r.t. and concentrated to dryness. The residue was suspended in DCM and sonicated. The resulting solid was collected by filtration to afford **Example 121c** (57 mg, 37% yield) as a white solid.

Step 2: Example 121

To a solution of **Example 121d** (60 mg, 0.16 mmol, 1.0 eq) and **Example 121c** (49 mg, 0.48 mmol, 3.0 eq) in dioxane (2 mL) were added Cs₂CO₃ (104.3 mg, 0.32 mmol, 2.0 eq), Xantphos (17.4 mg, 0.03 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (20.7 mg, 0.02 mmol, 0.1 eq). The reaction mixture was stirred at 110 °C for 4 h under N₂ protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 121** (40.1 mg, 56% yield) as a yellow solid. LCMS [M+1]⁺ = 450.4. ¹H NMR (300 MHz, DMSO-*d*₆) δ 10.97 (s, 1H), 9.13 (s, 1H), 9.05 (s, 1H), 8.56 (s, 1H), 7.76 (d, *J* = 7.5 Hz, 1H), 7.63 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.53 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.39 (s, 1H), 7.26 (t, *J* = 7.8 Hz, 1H), 3.95 (s, 3H), 3.86-3.73 (m, 1H), 3.70 (s, 3H), 3.00-2.88 (m, 1H), 1.16 - 0.95 (m, 10H).

Example 122:




Step 1: Example 122b

[00571] To a solution of **Example 122a** (700 mg, 4.61 mmol) in THF (65 mL) was added MeMgCl (15 mL, 45 mmol) at 0°C under argon protection. The reaction mixture was allowed to warm to room temperature and stirred for 16 h. Then the mixture was quenched by aq. NH₄Cl, extracted with EA (50 mL*2). The combined organic phase was washed by water, brine and dried over anhydrous Na₂SO₄. The solvent was removed under reduced pressure to give **Example 122b** (600 mg, 85.7% yield) as brown oil, which was used for the next step directly.

Step 1: Example 122


[00572] To a solution of **Example 122c** (100 mg, 0.26 mmol) and **Example 122b** (59 mg, 0.39 mmol) in 1, 4-dioxane (2.5 mL) were added Pd₂(dba)₃ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs₂CO₃ (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 122** (16 mg, 11.5% yield) as a yellow solid. LCMS [M+1]⁺ = 500.0. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.05 (s, 1H), 9.90 (s, 1H), 9.06 (s, 1H), 8.54 (s, 1H), 8.25 (s, 1H), 7.95 (s, 1H), 7.72 (d, *J* = 8.8 Hz, 1H), 7.67 (d, *J* = 8.0 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.50 (d, *J* = 8.8 Hz, 1H), 7.28 (t, *J* = 8.0 Hz, 1H), 3.93 (s, 3H), 3.70 (s, 3H), 2.94 (s, 1H), 1.40 (s, 6H), 1.05 (s, 2H), 0.98 (d, *J* = 7.7 Hz, 2H).

Example 123:



[00573] To the solution of **Example 123a** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (5 mL) were added **Example 123b** (95 mg, 0.78 mmol, 3.0 eq), Cs_2CO_3 (170 mg, 0.52 mmol, 2.0 eq), Xantphos (15 mg, 0.026 mmol, 0.1 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (27 mg, 0.026 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 . The reaction solution was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the desired product **Example 123** (26.1 mg, 21.4% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 469.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 10.89 (s, 1H), 9.05 (s, 1H), 8.91 (brs, 1H), 8.56 (s, 1H), 7.67 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.52 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.29 (t, $J = 8.1$ Hz, 1H), 6.64 (s, 1H), 3.95 (s, 3H), 3.69 (s, 3H), 3.00 - 2.88 (m, 1H), 1.22 (s, 9H), 1.12 - 0.99 (m, 4H).

Example 124:



Step 1: Example 124c

[00574] To a solution of **Example 124b** (465 mg, 5.28 mmol, 1.5 eq) in DMF (10 mL) was added Cs_2CO_3 (2.3 g, 7.04 mmol, 2.0 eq). The reaction mixture was stirred for 10 min at r.t., followed by addition of **Example 124a** (500 mg, 3.52 mmol, 1.0 eq) in DMF (3 mL). The mixture was stirred for 4 h at 80°C . After cooled to room temperature, the mixture was poured into H_2O (25 mL) and extracted with EtOAc (50 mL*2). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (1/1) to afford the product **Example 124c** (250 mg, 32% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 211.2$.

Step 2: Example 124d


[00575] To a solution of **Example 124c** (250 mg, 1.19 mmol, 1.0 eq) in MeOH (30 mL) was added Pd/C (25 mg) under N_2 protection. The suspension was degassed under vacuum and purged with H_2 three

times, and the reaction mixture was stirred at r.t. for 0.5 h under H₂ balloon. The solid was filtered out, and the filtrate was concentrated to afford the product **Example 124d** (230 mg, crude, yield: quant.) as yellow oil. LCMS [M+1]⁺ = 181.2.

Step 3: Example 124

[00576] To a solution of **Example 124e** (100 mg, 0.26 mmol, 1.0 eq.) in dioxane (3 mL) were added Cs₂CO₃ (169.5 mg, 0.52 mmol, 2.0 eq), **Example 124d** (93.7 mg, 0.52 mmol, 2.0 eq) and 3rd-t-Bu-xphos-Pd (26.7 mg, 0.03 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 124** (37.0 mg, 27% yield) as a yellow solid. LCMS [M+1]⁺ = 528.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.06 (s, 1H), 9.87 (s, 1H), 9.07 (s, 1H), 8.56 (s, 1H), 7.90 (d, *J* = 3.0 Hz, 1H), 7.76 (s, 1H), 7.67 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.63 - 7.55 (m, 2H), 7.40 (dd, *J* = 9.0, 3.0 Hz, 1H), 7.32 (t, *J* = 7.8 Hz, 1H), 5.08 - 4.99 (m, 1H), 3.95 (s, 3H), 3.91 - 3.75 (m, 4H), 3.72 (s, 3H), 3.00 - 2.88 (m, 1H), 2.29 - 2.05 (m, 1H), 2.03 - 1.89 (m, 1H), 1.15 - 0.91 (m, 4H).

Example 125:



Step 1: Example 125b

[00577] To a solution of **Example 125a** (5.0 g, 35.2 mmol, 1.0 eq) in DMF (50 mL) was added NaSMe (18.5 g, 20% in water, 53 mmol, 1.5 eq) dropwise at 0°C. The reaction mixture was stirred for 1 h at r.t.. The reaction was diluted with EtOAc (100mL) and washed with brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (2/1) to afford the product **Example 125b** (4.9 g, 82% yield) as a yellow solid. LCMS [M+1]⁺ = 171.2.

Step 2: Example 125c

[00578] To a solution of **Example 125b** (4.8 g, 28 mmol, 1.0 eq) in MeOH (100 mL) were added Fe powder (7.9 g, 141 mmol, 5.0 eq), NH₄Cl (7.5 g, 141 mmol, 5.0 eq) and AcOH (3 mL). The reaction mixture was stirred for 1 h at 80°C. The reaction mixture was basified with K₂CO₃ (pH = 7~8). The solid was filtered out, and the filtrate was concentrated. The crude product was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (2/1) to afford the product **Example 125c** (3.6 g, 92% yield) as red oil.

LCMS [M+1]⁺ = 141.2.

Step 3: Example 125f

[00579] To a solution of **Example 125d** (5.0 g, 21 mmol, 1.0 eq) in dry THF (50 mL) was added **Example 125e** (105 mL, 1.0 M in THF, 105 mmol, 5.0 eq) dropwise at -15°C under N₂ protection. The mixture was stirred for 0.5 h at r.t.. The mixture was poured into saturated aqueous of NH₄Cl (100 mL) and extracted with EtOAc (150 mL*2). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (10/1) to afford the product **Example 125f** (2.8 g, 62% yield) as yellow oil.

LCMS [M+1]⁺ = 216.2

Step 4: Example 125g

[00580] To a solution of **Example 125c** (500 mg, 3.57 mmol, 1.0 eq) in DMF (10 mL), was added NaH (1.43 g, 60% in mineral oil, 35.7 mmol, 10.0 eq) in portions at 0°C. After addition, the reaction was stirred for 30 min at 0°C, followed by addition of **Example 125f** (768 mg, 3.57 mmol, 1.0 eq) in DMF. The reaction mixture was stirred for 1 h at r.t.. The reaction solution was poured into water, extracted with EtOAc (50 mL*2), and the combined organic layer dried over Na₂SO₄ and concentrated. The crude product was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (3/1) to afford the product **Example 125g** (370 mg, 32% yield) as a yellow solid. LCMS [M+1]⁺ = 320.1

Step 5: Example 125h


[00581] To a solution of **Example 125g** (200 mg, 0.627 mmol, 1.0 eq) in DCM (4 mL) was added *m*-CPBA (434 mg, 2.51 mmol, 4.0 eq) at 0°C. The reaction mixture was stirred for 1 h at r.t.. The reaction was diluted with DCM (50 mL), washed with NaOH aqueous solution (1M), Na₂SO₃ aqueous solution and brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (1/1) to afford the product **Example 125h** (60 mg, 27% yield) as a yellow solid. LCMS [M+1]⁺ = 352.2.

Step 6: Example 125

[00582] To a solution of **Example 125h** (50 mg, 0.142 mmol, 1.0 eq) in dioxane (2 mL) were added Cs₂CO₃ (93 mg, 0.285 mmol, 2.0 eq), **Example 125i** (27 mg, 0.285 mmol, 2.0 eq), Xantphos (17 mg, 0.028 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (15 mg, 0.014 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 125** (17 mg, 29% yield) as a yellow solid. LCMS [M+1]⁺ = 411.2. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.83 (s,

1H), 10.54 (s, 1H), 9.18 (s, 1H), 8.86- 8.78 (m, 2H), 8.62 (dd, $J = 4.8, 2.1$ Hz, 1H), 8.28 (dd, $J = 7.8, 2.1$ Hz, 1H), 8.17 (dd, $J = 9.3, 1.5$ Hz, 1H), 7.61 (dd, $J = 9.3, 4.8$ Hz, 1H), 7.33 (dd, $J = 7.8, 4.8$ Hz, 1H), 3.31 (s, 3H), 3.03-2.90 (m, 1H), 1.16-0.99(m, 4H).

Example 126:



Step 1: Example 126c

[00583] To a solution of **Example 126a** (1.54 g, 7.1 mmol, 1.0 eq) in DMF (20 mL), was added NaH (2.86 g, 60% in mineral oil, 71 mmol, 10.0 eq) in portions at 0°C. **Example 126b** (1.0 g, 7.1 mmol, 1.0 eq) in DMF was added to the solution. The reaction mixture was stirred for 1 h at r.t.. The reaction was poured into water (100 mL), extracted with EtOAc (100 mL*3), and the combined organic layers were dried over Na_2SO_4 and concentrated. The crude product was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (3/1) to afford the product **Example 126c** (570 mg, 25% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 320.1$

Step 2: Example 126d


[00584] To a solution of **Example 126c** (740 mg, 2.32 mmol, 1.0 eq) in DCM (8 mL) was added *m*-CPBA (1.61 g, 9.28 mmol, 4.0 eq) at 0°C. The reaction mixture was stirred for 2 h at r.t.. The reaction was diluted with DCM (20 mL), washed with NaOH aqueous solution (1M), Na_2SO_3 aqueous solution and brine. The organic layer was dried over Na_2SO_4 , and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (1/1) to afford the product **Example 126d** (470 mg, 57% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 352.2$.

Step 3: Example 126

[00585] To a solution of **Example 126d** (100 mg, 0.285 mmol, 1.0 eq) in dioxane (2 mL) were added Cs_2CO_3 (186 mg, 0.57 mmol, 2.0 eq), **Example 126e** (87 mg, 0.57 mmol, 2.0 eq), BINAP (36 mg, 0.057 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (30 mg, 0.028 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed, and the crude

product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 126** (12 mg, 9% yield) as an off-white solid. LCMS $[M+1]^+ = 468.1$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$) δ 11.84 (s, 1H), 10.04 (s, 1H), 9.14 (s, 1H), 8.95 (s, 1H), 8.69 - 8.63 (m, 1H), 8.35 (dd, $J = 2.7, 0.9$ Hz, 1H), 8.26 (dd, $J = 7.8, 2.4$ Hz, 1H), 7.78 (dd, $J = 8.7, 2.4$ Hz, 1H), 7.68 (d, $J = 8.7$ Hz, 1H), 7.32 (dd, $J = 7.8, 4.8$ Hz, 1H), 5.09 (s, 1H), 3.31 (s, 3H), 3.01-2.88 (m, 1H), 1.45 (s, 6H), 1.12-0.97 (m, 4H).

Example 127:



Step 1: Example 127b

[00586] To a solution of **Example 127a** (550 mg, 2.32 mmol, 1.0 eq) in DCM (10 mL) was added BBr_3 (1.15 g, 4.64 mmol, 2.0 eq) at -20°C . The reaction mixture was stirred for 2 h at r.t.. It was quenched with NaHCO_3 aqueous solution (10 mL), extracted with DCM (20 mL*3). The combined organic layer was washed with brine, dried over Na_2SO_4 and concentrated to afford the product **Example 127b** (500 mg, crude, 96% yield) as a brown solid. LCMS $[M+1]^+ = 224.3$.

Step 2: Example 127c

[00587] To a mixture of **Example 127b** (500 mg, 2.24 mmol, 1.0 eq) and K_2CO_3 (927 mg, 6.72 mmol, 3.0 eq) in ACN (10 mL) was added CD_3I (487.2 mg, 3.36 mmol, 1.5 eq). The reaction mixture was stirred for 6 h at 80°C . After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography eluted with PE/EtOAc (1/3) to afford the product **Example 127c** (200 mg, 37% yield) as a yellow solid. LCMS $[M+1]^+ = 241.2$.

Step 3: Example 127d

[00588] To a solution of **Example 127c** (200 mg, 0.84 mmol, 1.0 eq) in MeOH (20 mL) was added Pd/C (20 mg) under N_2 protection, the suspension was degassed under vacuum and purged with H_2 three times, the reaction mixture was stirred at r.t. for 1 h under H_2 balloon. The solid was filtered out, the filtrate was concentrated to afford the product **Example 127d** (160 mg, 90% yield) as a yellow solid. LCMS $[M+1]^+ = 211.2$.

Step 4: Example 127f


[00589] To a solution of **Example 127e** (170 mg, 0.72 mmol, 1.0 eq) and **Example 127d** (151.2 mg, 0.72 mmol, 1.0 eq) in dry THF (6 mL) was added LiHMDS (1.4 mL, 1 M in THF, 1.44 mmol, 2.0 eq) dropwise at -20 °C under N₂ protection. The reaction mixture was stirred for 0.5 h at r.t., and then the silica was added to the mixture and concentrated. The residue was purified by silica gel flash column chromatography eluted with DCM/MeOH (20/1) to afford the product **Example 127f** (45 mg, 15% yield) as a yellow solid. LCMS [M+1]⁺ = 409.4.

Step 5: Example 127h

[00590] To a solution of **Example 127f** (45 mg, 0.11 mmol, 1.0 eq) in THF (3 mL) was added **Example 127g** (0.9 mL, 1.0 M in THF, 0.88 mmol, 8.0 eq) dropwise at 0°C under N₂ protection. The mixture was stirred for 0.5 h at r.t.. The reaction was poured into saturated aqueous of NH₄Cl (10 mL) and extracted with EtOAc (15 mL x 3). The combined organic layer was washed with brine, dried over Na₂SO₄, and concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 127f** (40 mg, 93% yield) as a yellow solid. LCMS [M+1]⁺ = 390.3.

Step 6: Example 127


[00591] To a solution of **Example 127f** (40 mg, 0.10 mmol, 1.0 eq) in dioxane (3 mL) were added Cs₂CO₃ (65.2 mg, 0.20 mmol, 2.0 eq), **Example 127l** (26.2 mg, 0.30 mmol, 3.0 eq), Xantphos (11.6 mg, 0.02 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (10.4 mg, 0.01 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. After cooled to room temperature, the solvent was removed, the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 127** (3.1 mg, 7% yield) as a yellow solid. LCMS [M+1]⁺ = 439.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.01 (s, 1H), 10.93 (s, 1H), 9.13 (s, 1H), 8.56 (s, 1H), 8.04 (s, 1H), 7.64 (dd, *J* = 7.8, 1.5 Hz, 1H), 7.52 (dd, *J* = 8.1, 1.8 Hz, 1H), 7.26 (t, *J* = 7.8 Hz, 1H), 3.08 - 2.90 (m, 1H), 2.11 - 1.93 (m, 1H), 1.18 - 0.98 (m, 4H), 0.81 (d, *J* = 6.0 Hz, 4H).

Example 128:

[00592] To a solution of **Example 128** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (5 mL) were added **Example 128b** (95 mg, 0.78 mmol, 3.0 eq), Cs₂CO₃ (170 mg, 0.52 mmol, 2.0 eq), Xantphos (15 mg, 0.026 mmol, 0.1 eq) and Pd₂(dba)₃.CHCl₃ (27 mg, 0.026 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂. The reaction solution was filtered and the filtrate was concentrated. The crude

product was purified by Prep-TLC (DCM/MeOH = 15/1) to give 32.0 mg crude product (90% purity) and further purified by Prep-TLC (DCM/MeOH = 15/1) to afford the desired product **Example 128** (12.1 mg, 9.9% yield) as a yellow solid. LCMS $[M+1]^+ = 469.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$) δ 10.89 (s, 1H), 9.04 (s, 1H), 8.91 (brs, 1H), 8.56 (s, 1H), 7.67 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.53 (d, $J = 6.6$ Hz, 1H), 7.29 (t, $J = 7.8$ Hz, 1H), 6.64 (s, 1H), 3.95 (s, 3H), 3.69 (s, 3H), 3.00 - 2.88 (m, 1H), 1.22 (s, 9H), 1.12 - 0.99 (m, 4H).

Example 129:



Example 129a

Step 1

Example 129c

Step 2

Example 129


Step 1: Example 129c

[00593] A solution of **Example 129b** (300 mg, 2.19 mmol) in 1,4-dioxane (5 mL) was treated with **Example 129a** (353 mg, 3.28 mmol) and TEA (664 mg, 6.57 mmol). The mixture was stirred at r.t. for 16 hrs. After reaction completed, the solvent was concentrated, the residue was suspended in DCM (5 mL), sonicated and the resulting solid was collected via filtration, dried to afford the desired product **Example 129c** (253 mg, crude) as a white solid. LCMS $[M+1]^+ = 115.0$.

Step 2: Example 129

[00594] To a solution of **Example 129a** (80 mg, 0.21 mmol) and **Example 129c** (29.3 mg, 0.26 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (19.6 mg, 0.021 mmol), XantPhos (24.7 mg, 0.042 mmol) and Cs_2CO_3 (140 mg, 0.43 mmol). The mixture was sealed, degassed by nitrogen for 3 times and stirred at 110°C for 16 h. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 129** (21.8 mg, 22.7% yield) as a white solid. LCMS $[M+1]^+ = 452.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 10.89 (s, 1H), 9.59 (s, 1H), 9.02 (d, $J = 5.3$ Hz, 1H), 8.54 (s, 1H), 8.00 (s, 1H), 7.62 (d, $J = 7.8$ Hz, 1H), 7.49 (d, $J = 7.9$ Hz, 1H), 7.24 (t, $J = 8.0$ Hz, 1H), 4.09 (s, 2H), 3.93 (s, 3H), 3.71 (s, 3H), 3.54 (s, 2H), 2.84 (d, $J = 4.4$ Hz, 3H), 2.59 (d, $J = 8.4$ Hz, 1H), 1.15 (d, $J = 6.7$ Hz, 3H).

Example 130:




Step 1: Example 130c

[00595] To a solution of **Example 130a** (200 mg, 1.92 mmol) in acetone (4 mL) was added TEA (291 mg, 2.88 mmol) at 0-5°C, and then **Example 130b** (311 mg, 2.88 mmol) was added dropwise. The mixture was stirred at room temperature for 1 h under N₂. The white solid was filtered off. To the filtration was added NH₃-H₂O (5 eq.). The reaction mixture was stirred at room temperature overnight. The mixture was concentrated. The residue was cooled and added Petroleum ether/EtOAc = 1/1 (3 mL). The solid was collected by filtered and dried to give **Example 130c** (17 mg, 8% yield) as a white solid.

Step 2: Example 130

[00596] To a solution of **Example 130d** (63 mg, 0.16 mmol) in dioxane (1.5 mL) were added **Example 130c** (17 mg, 0.16 mmol), Pd₂(dba)₃ (15 mg, 0.016 mmol), Xantphos (10 mg, 0.016 mmol) and Cs₂CO₃ (108 mg, 0.33 mmol). The mixture was sealed and heated to 120°C for 2 h. The mixture was filtrated and concentrated under reduced pressure. The residue was purified by Prep-HPLC to give **Example 130** (8.7 mg, 12% yield) as a yellow solid. LCMS [M+1]⁺ = 451.1. ¹H NMR (400 MHz, Chloroform-*d*) δ 11.12 (s, 1H), 8.90 (s, 1H), 8.22-8.03 (m, 3H), 7.75 (d, *J* = 7.8 Hz, 1H), 7.59 (d, *J* = 7.9 Hz, 1H), 7.25-7.22 (m, 1H), 4.78 (d, *J* = 65.0 Hz, 1H), 3.99 (s, 3H), 3.77 (s, 3H), 2.66-2.57 (m, 1H), 1.94-1.76 (m, 2H), 1.27-1.17 (m, 3H), 1.07-1.0 (m, 2H).


Example 131:



[00597] To a solution of **Example 131a** (40 mg, 0.104 mmol, 1.0 eq) and **Example 131b** (35.4 mg, 0.208 mmol, 2.0 eq) in dioxane (3 mL) were added Xantphos (12.0 mg, 0.02 mmol, 0.2 eq), Cs₂CO₃ (67.8 mg, 0.208 mmol, 2.0 eq) and Pd₂(dba)₃.CHCl₃ (10.4 mg, 0.01 mmol, 0.1 eq). The reaction mixture was stirred at 110°C for 4 h under N₂ protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 131** (7.4 mg, 14% yield) as a yellow solid. LCMS [M+1]⁺ = 519.4.


¹H NMR (300 MHz, DMSO-*d*₆) δ 12.33 (s, 1H), 10.27 (s, 1H), 9.54 (s, 1H), 9.18 (s, 1H), 8.66 (s, 1H), 8.17 (d, *J* = 5.1 Hz, 1H), 8.05 - 7.95 (m, 1H), 7.58 (d, *J* = 8.1 Hz, 1H), 7.50 (d, *J* = 5.1 Hz, 1H), 5.30 (s, 1H), 4.00 (s, 3H), 3.88 (s, 3H), 3.08 - 2.93 (m, 1H), 1.50 (s, 6H), 1.16 - 0.98 (m, 4H).

Example 132:



[00598] To a solution of **Example 132a** (100 mg, 0.26 mmol) and **Example 132b** (43 mg, 0.39 mmol) in DMA (2.5 mL) were added Pd₂(dba)₃ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs₂CO₃ (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 132** (1.3 mg, 3.5% yield) as a white solid. LCMS [M+1]⁺ = 459.0. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.07 (s, 1H), 9.49 (s, 1H), 8.97 (s, 1H), 8.53 (s, 1H), 8.30 (s, 0H), 7.65 (d, *J* = 8.0 Hz, 1H), 7.56 (d, *J* = 8.0 Hz, 1H), 7.27 (t, *J* = 7.9 Hz, 1H), 5.90 (s, 1H), 3.92 (s, 3H), 3.68 (s, 3H), 3.57 (s, 3H), 2.88 (s, 1H), 2.17 (s, 3H), 1.03 (s, 2H), 0.95 (d, *J* = 7.7 Hz, 2H).

Example 133:




Example 133a

Example 133

[00599] To a solution of **Example 133a** (180 mg, 0.47 mmol) and **Example 133b** (180 mg, 1.41 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (43 mg, 0.047 mmol), xantphos (54 mg, 0.094 mmol) and Cs_2CO_3 (306 mg, 0.94 mmol). The mixture was degassed by nitrogen for 3 times, sealed and stirred at 120°C for 16 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 133** (67.3 mg, 30.3% yield) as a white solid. LCMS $[\text{M}+1]^+ = 476.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 10.98 (s, 1H), 9.19 (s, 1H), 9.02 (s, 1H), 8.54 (s, 1H), 7.90 (s, 1H), 7.60 (d, $J = 7.8$ Hz, 1H), 7.51 (d, $J = 8.0$ Hz, 1H), 7.23 (t, $J = 8.0$ Hz, 1H), 3.93 (s, 3H), 3.67 (s, 3H), 3.64 (s, 4H), 2.94 (s, 1H), 1.18 (s, 6H), 1.06 (d, $J = 4.5$ Hz, 2H), 1.00 (d, $J = 7.8$ Hz, 2H).

Example 134:




Example 134a

Example 134

[00600] To a solution of **Example 134a** (130 mg, 0.35 mmol) and **Example 134b** (67 mg, 0.52 mmol) in 1,4-dioxane (4 mL) were added $\text{Pd}_2(\text{dba})_3$ (32 mg, 0.035 mmol), XantPhos (40 mg, 0.070 mmol) and Cs_2CO_3 (227 mg, 0.70 mmol). The mixture was sealed, degassed by nitrogen for 3 times and stirred at 110°C for 16 h. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 134** (34 mg, 21.0% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 466.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 10.86 (s, 1H), 9.17 (d, $J = 5.3$ Hz, 1H), 8.55 (s, 1H), 8.27 (s, 1H), 7.62 (d, $J = 7.8$ Hz, 1H), 7.51 (d, $J = 7.9$ Hz, 1H), 7.25 (t, $J = 7.9$ Hz, 1H), 4.06 (t, $J = 8.2$ Hz, 2H), 4.02 (s, 1H), 3.93 (s, 3H), 3.71 (s, 3H), 3.44 (t, $J = 8.1$ Hz, 2H), 2.83 (d, $J = 4.6$ Hz, 3H), 1.09 (d, $J = 6.7$ Hz, 6H).

Example 135:




Example 135a

Example 135

[00601] To a solution of **Example 135a** (100 mg, 0.26 mmol,) and **Example 135b** (43 mg, 0.39 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 120°C for 16 h. When completed, the reaction was cooled to r.t, diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 135** (41.8 mg, 35.1% yield) as a white solid (contain 0.43 FA salt by $^1\text{H NMR}$). LCMS $[\text{M}+1]^+ = 459.0$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 13.32 (s, 1H), 10.96 (s, 1H), 9.20 (s, 1H), 8.54 (s, 1H), 7.72 (d, $J = 6.6$ Hz, 1H), 7.66 (d, $J = 7.8$ Hz, 1H), 7.53 (d, $J = 7.9$ Hz, 1H), 7.27 (t, $J = 8.0$ Hz, 1H), 6.99 (s, 1H), 5.84 (d, $J = 6.6$ Hz, 1H), 3.93 (s, 3H), 3.68 (s, 3H), 2.98 (d, $J = 7.1$ Hz, 1H), 1.09 (d, $J = 4.3$ Hz, 2H), 1.03 (dd, $J = 9.6, 5.4$ Hz, 2H).

Example 136:




Example 136a

Example 136


[00602] To a solution of **Example 136a** (100 mg, 0.26 mmol,) and **Example 136b** (37 mg, 0.31 mmol) in 1,4-dioxane (1 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), XantPhos (30 mg, 0.052 mmol) and Cs_2CO_3 (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 136** (4.5 mg, 3.7% yield) as a white solid. LCMS $[\text{M}+1]^+ = 467.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 11.00 (s, 1H), 10.34 (s, 1H), 9.16 (s, 1H), 8.54 (s, 1H), 8.40 (d, $J = 8.0$ Hz, 1H), 8.21 (s, 1H), 7.63 - 7.60 (m, 3H), 7.31 - 7.27 (m, 2H), 3.93 (s, 3H), 3.69 (s, 3H), 2.97 (br, 1H), 1.07 - 1.00 (m, 4H).

Example 137:




[00603] To a solution of **Example 137a** (60 mg, 0.171 mmol, 1.0 eq) in dioxane (2 mL) were added Cs_2CO_3 (111 mg, 0.342 mmol, 2.0 eq), **Example 137b** (32 mg, 0.342 mmol, 2.0 eq), BINAP (21.3 mg, 0.034 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (17.7 mg, 0.017 mmol, 0.1eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 30/1) to afford the product **Example 137** (8 mg, 11% yield) as an off-white solid. LCMS $[\text{M}+1]^+ = 410.2$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.85 (s, 1H), 10.11 (s, 1H), 9.16 (s, 1H), 9.03 (s, 1H), 8.66 (dd, $J = 4.8, 1.8$ Hz, 1H), 8.30-8.22 (m, 2H), 7.76-7.68 (m, 2H), 7.31 (dd, $J = 7.8, 4.8$ Hz, 1H), 6.99 - 6.91 (m, 1H), 3.12 (m, 3H), 3.01-2.89 (m, 1H), 1.12 - 0.97 (m, 4H).

Example 138:




[00604] To a solution of **Example 138a** (277 mg, 0.72 mmol) and **Example 138b** (84 mg, 0.87 mmol) in 1,4-dioxane (4 mL) were added $\text{Pd}_2(\text{dba})_3$ (66 mg, 0.072 mmol), Xantphos (83 mg, 0.14 mmol) and Cs_2CO_3 (470 mg, 1.44 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 16 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 138** (21 mg, 6.6% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 445.0$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.06 (s, 1H), 9.60 (s, 1H), 8.99 (s, 1H), 8.54 (s, 1H), 7.65 (d, $J = 8.0$ Hz, 1H), 7.56 (d, $J = 8.1$ Hz, 1H), 7.51 (s, 2H), 7.27 (t, $J = 7.9$ Hz, 1H), 6.08 (s, 1H), 3.92 (s, 3H), 3.69 (d, $J = 4.7$ Hz, 6H), 2.89 (s, 1H), 1.03 (s, 2H), 0.95 (s, 2H).

Example 139:



[00605] To a solution of **Example 139a** (100 mg, 0.26 mmol) and **Example 139b** (37 mg, 0.31 mmol) in 1,4-dioxane (1 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), XantPhos (30 mg, 0.052 mmol) and Cs_2CO_3 (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-TLC (MeOH/DCM = 1/15) to give the desired product **Example 139** (14.6 mg, 12.1% yield) as a white solid. LCMS $[\text{M}+1]^+ = 467.2$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.04 (s, 1H), 10.53 (s, 1H), 9.13 (s, 1H), 8.61 (s, 1H), 8.54 (s, 1H), 8.07 (d, $J = 12.0$ Hz, 1H), 7.83 - 7.79 (m, 2H), 7.62 (t, $J = 8.0$ Hz, 2H), 7.33 (t, $J = 8.0$ Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.97 (br, 1H), 1.21 - 1.01 (m, 4H).


Example 140:



[00606] To a solution of **Example 140a** (100 mg, 0.26 mmol) and **Example 140b** (37 mg, 0.31 mmol) in 1,4-dioxane (1 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 140** (12.2 mg, 10.1% yield) as a white solid. LCMS $[\text{M}+1]^+ = 467.2$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.24 (s, 1H), 10.42 (s, 1H), 9.12 (s, 1H), 8.54 (s, 1H), 7.98 (s, 1H), 7.87 (t, $J = 8.0$ Hz, 1H), 7.78 (d, $J = 8.0$ Hz,

1H), 7.72 (d, $J = 8.0$ Hz, 1H), 7.58 (d, $J = 8.0$ Hz, 1H), 7.52 (d, $J = 8.0$ Hz, 1H), 7.35 (t, $J = 8.0$ Hz, 1H), 3.93 (s, 3H), 3.70 (s, 3H), 2.96 (br, 1H), 1.21 - 1.00 (m, 4H).

Example 141:



Step 1: Example 141c

[00607] To a solution of **Example 141a** (182 mg, 1.94 mmol, 1.0 eq) in dioxane (10 mL) were added Cs_2CO_3 (1.25 g, 3.87 mmol, 2.0 eq), **Example 141b** (500 mg, 2.32 mmol, 1.2 eq), BINAP (240 mg, 0.387 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (199 mg, 0.194 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. The solvent was concentrated, and the crude product was purified by silica gel flash column chromatography, eluted with (DCM/MeOH = 30/1) to afford the product **Example 141c** (490 mg, 72% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 293.1$.

Step 2: Example 141e


To a solution of **Example 141c** (490 mg, 1.678 mmol, 1.0 eq) in THF (10 mL) was added **Example 141d** (25.17 mL, 1.0 M in THF, 25.17 mmol, 15.0 eq) dropwise at 0°C under N_2 protection. The mixture was stirred for 1.5 h at r.t.. After the reaction was completed, the mixture was poured into saturated aqueous of NH_4Cl (20 mL) and extracted with EtOAc (50 mL*2). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by silica gel flash column chromatography, eluted with DCM/MeOH = (30/1) to afford the product **Example 141e** (170 mg, 37% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 274.1$.

Step 3: Example 141

[00608] To a solution of **Example 141e** (60 mg, 0.22 mmol, 1.0 eq) in dioxane (2 mL) were added Cs_2CO_3 (143 mg, 0.44 mmol, 2.0 eq), **Example 141f** (90 mg, 0.44 mmol, 2.0 eq), 3rd-t-Bu-xphos-Pd (39 mg, 0.044 mmol, 0.2 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed, the crude product was purified by Prep-TLC (DCM/MeOH = 20/1) to afford the product **Example 141** (29 mg, 29% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 443.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 12.31 (s, 1H), 10.10 (s, 1H), 9.51 (s, 1H), 9.17 (s,

1H), 8.66 (s, 1H), 8.35 - 8.32 (m, 1H), 8.19 (d, $J = 5.1$ Hz, 1H), 7.82 - 7.64 (m, 2H), 7.48 (d, $J = 5.1$ Hz, 1H), 7.02 - 6.92 (m, 1H), 4.00 (s, 3H), 3.88 (s, 3H), 3.06 - 2.95 (m, 1H), 1.67 - 0.97 (m, 4H).

Example 142:




Step 1: Example 142c

[00609] A mixture of **Example 142a** (250 mg, 2.0 mmol), **Example 142b** (276 mg, 2.0 mmol) and TEA (610 mg, 6.0 mmol) in 1,4-dioxane (2 mL) was stirred at room temperature overnight. The reaction was filtered to give the desired product **Example 142c** (350 mg, 51.4% yield) as a white solid. LCMS $[M+1]^+ = 133.1$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 5.97 (s, 2H), 3.88 - 3.75 (m, 4H), 3.69 (s, 3H), 1.51 (d, $J = 24.0$ Hz, 3H).

Step 2: Example 142

[00610] To a solution of **Example 142c** (40 mg, 0.31 mmol) and **Example 142d** (96 mg, 0.25 mmol) in 1,4-dioxane (1 mL) were added $\text{Pd}_2(\text{dba})_3$ (23 mg, 0.025 mmol), Xantphos (29 mg, 0.051 mmol) and Cs_2CO_3 (166 mg, 0.51 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 142** (46.7 mg, 38.9% yield) as a yellow solid. LCMS $[M+1]^+ = 473.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 10.90 (s, 1H), 9.89 (s, 1H), 9.01 (s, 1H), 8.54 (s, 1H), 7.99 (s, 1H), 7.62 (d, $J = 8.0$ Hz, 1H), 7.49 (d, $J = 8.0$ Hz, 1H), 7.25 (t, $J = 8.0$ Hz, 1H), 4.13 - 4.01 (m, 4H), 3.93 (s, 3H), 3.71 (s, 3H), 1.54 (d, $J = 24.0$ Hz, 3H).

Example 143:


**Step 1: Example 143b**

[00611] A solution of **Example 143a** (2.0 g, 15.5 mmol) in DMSO/H₂O (20 mL/ 20 mL) were treated with NaCN (1.52 g, 31.0 mmol) and DABCO (1.74 g, 15.5 mmol). The mixture was stirred at 60°C for 6 h. After reaction completed, the solvent was extracted by DCM (50 mL) and concentrated to give crude product, which was purified directly by Prep-HPLC to give the desired product **Example 143b** (500 mg, 26.8% yield) as a white solid. LCMS [M+1]⁺ = 121.0

Step 2: Example 143

[00612] To a solution of **Example 143b** (60 mg, 0.5 mmol) and **Example 143c** (191 mg, 0.5 mmol,) in dioxane (5 mL) were added Pd₂(dba)₃ (50 mg, 0.05 mmol), Xantphos (30 mg, 0.05 mmol) and Cs₂CO₃ (224.5 mg, 0.75 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for overnight. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 143** (7.4 mg, 3.2% yield) as a off white solid. LCMS [M+1]⁺ = 468.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.14 (s, 1H), 10.90 (s, 1H), 9.16 (s, 1H), 8.60 – 8.51 (m, 2H), 7.85 (s, 1H), 7.76 (s, 1H), 7.67 (d, *J* = 7.9 Hz, 1H), 7.62 (d, *J* = 7.9 Hz, 1H), 7.34 (d, *J* = 8.0 Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.98 (s, 1H), 1.21 (s, 1H), 1.09 (s, 2H), 1.04 (d, *J* = 7.0 Hz, 2H).

Example 144:




Step 1: Example 144c

[00613] A mixture of **Example 144a** (250 mg, 2.1 mmol), **Example 144b** (288 mg, 2.1 mmol) and TEA (637 mg, 6.3 mmol) in 1,4-dioxane (2 mL) was stirred at room temperature overnight. The reaction was filtered to give the desired product **Example 144c** (200 mg, 76.3% yield) as a white solid. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 6.04 (s, 2H), 4.03 (t, $J = 8.0$ Hz, 2H), 3.86 (t, $J = 8.0$ Hz, 2H), 3.67 - 3.62 (m, 1H).

Step 2: Example 144


[00614] To a solution of **Example 144c** (66 mg, 0.53 mmol) and **Example 144d** (100 mg, 0.27 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.027 mmol), Xantphos (31 mg, 0.053 mmol) and Cs_2CO_3 (173 mg, 0.53 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C overnight. When completed, the reaction was cooled to r.t., diluted with CH_3CN (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 144** (61.3 mg, 49.7% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 466.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 10.91 (s, 1H), 9.96 (s, 1H), 9.02 (s, 1H), 8.54 (s, 1H), 7.96 (s, 1H), 7.62 (d, $J = 8.0$ Hz, 1H), 7.48 (d, $J = 8.0$ Hz, 1H), 7.25 (t, $J = 8.0$ Hz, 1H), 4.27 (t, $J = 8.0$ Hz, 2H), 4.13 (t, $J = 8.0$ Hz, 1H), 3.93 (s, 3H), 3.75 - 3.67 (m, 4H).

Example 145:



[00615] To a solution of **Example 145a** (100 mg, 0.26 mmol) and **Example 145b** (98 mg, 0.78 mmol) in 1,4-dioxane (2 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and

Cs_2CO_3 (170 mg, 0.52 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 145** (32 mg, 26.0% yield) as a white solid. LCMS $[\text{M}+1]^+ = 474.0$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.03 (s, 1H), 9.92 (s, 1H), 9.08 (s, 1H), 8.54 (s, 1H), 8.05 (s, 1H), 7.71 (s, 1H), 7.63 (d, $J = 8.0$ Hz, 1H), 7.58 (d, $J = 8.3$ Hz, 2H), 7.29 (t, $J = 7.9$ Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.94 (s, 1H), 2.23 (s, 3H), 1.05 (s, 2H), 0.99 (d, $J = 7.6$ Hz, 2H).

Example 146:**Step 1: Example 146b**

[00616] The solution of **Example 146a** (3.0 g, 19.2 mmol, 1.0 eq) in POCl_3 (30 mL) was stirred for 4 h at 100°C . After the reaction completed, it was concentrated under vacuo to give crude product (3.0 g, crude) which was used to next step without further purification.

Step 2: Example 146d

[00617] To a solution of **Example 146c** (1.96 g, 28.4 mmol, 2.0 eq) and DIEA (14.7 g, 113.6 mmol, 8.0 eq) in THF (30 mL) was added a solution of **Example 146b** (3.0 g, crude) in DCM (20 mL) dropwise at 0°C . The reaction solution was stirred for 30 min at r.t.. The reaction solution was diluted with EtOAc (100 mL), washed with brine (50 mL*3), dried over Na_2SO_4 and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc = (3/1) to afford the product **Example 146d** (400 mg, 14% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 206.2$.

Step 3: Example 146f


[00618] To a solution of **Example 146d** (400 mg, 1.94 mmol, 1.0 eq) and **Example 146e** (402 mg, 1.94 mmol, 1.0 eq) in dry THF (15 mL) was added LiHMDS (3.88 mL, 1M in THF, 3.88 mmol, 2.0 eq) dropwise at 0°C under N_2 protection. The reaction mixture was stirred for 0.5 h at r.t.. Then the silica was added to the mixture and concentrated. The residue was purified by silica gel flash column

chromatography, eluted with DCM/MeOH = (20/1) to afford the product **Example 146f** (310 mg, 42% yield) as a yellow solid. LCMS $[M+1]^+ = 377.3$.

Step 4: Example 146

[00619] To a solution of **Example 146f** (290 mg, 0.77 mmol, 1.0 eq) in dioxane (5 mL) were added Cs_2CO_3 (502.0 mg, 1.54 mmol, 2.0 eq), **Example 146g** (231 mg, 2.31 mmol, 3.0 eq) and BINAP (95.9 mg, 0.15 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (82.8 mg, 0.08 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 146** (80.5 mg, 24% yield) as an off-white solid. LCMS $[M+1]^+ = 441.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 10.92 (s, 1H), 9.61 (s, 1H), 9.05 (d, $J = 4.8$ Hz, 1H), 8.57 (s, 1H), 8.03 (s, 1H), 7.64 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.52 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.27 (t, $J = 7.8$ Hz, 1H), 4.01 (t, $J = 7.8$ Hz, 4H), 3.74 (s, 3H), 2.87 (d, $J = 4.8$ Hz, 3H), 2.23-2.08 (m, 2H).

Example 147:



Step 1 : 2-chloro-3-methoxyisonicotinic acid

[00620] To a solution of 2-chloro-3-methoxypyridine (50 g, 0.348 mol) in THF (500 mL) at -78°C was added LDA (1.0 M in THF, 418 mL, 0.418 mmol) dropwise. After addition, the mixture was stirred at -78°C for 30 minutes, then dry ice was added to the reaction during 30 minutes. The reaction was quenched with 5% w/v aqueous NaOH (200 mL) and the aqueous layer was washed with EtOAc (200 mL \times 2). The organic fractions were discarded and the pH of the aqueous layer was adjusted to 2 with a 6 M aqueous HCl solution. The aqueous layer was extracted with EtOAc (30 mL \times 3) and the combined organic fractions dried by Na_2SO_4 , filtered and concentrated to give the desired compound as a yellow solid (35 g, 53.8%).

Step 2 : 2-chloro-3-methoxyisonicotinamide

[00621] To a solution of 2-chloro-3-methoxyisonicotinic acid (37 g, 0.197 mol), Boc_2O (49.84 mL, 0.217 mol) and pyridine (19.14 mL, 0.239 mol) in DCM (600 mL) at 0°C was added NH_4HCO_3 (78.0

g, 0.985 mol). After the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to rt. After stirring at room temperature for overnight, the reaction mixture was concentrated on the rotovap to remove some of the DCM, and filtered to collect the liquid. The filtrate washed with H₂O (100 mL×2). The combined organic layers were washed by brine (100 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (DCM/MeOH=20/1) to give the title compound as a brown solid (24.0 g, Yield : 65.3 %). LM-MS: m/z =187.6[M+H]⁺

Step 3: 2-chloro-3-methoxy-4-(1H-1,2,4-triazol-3-yl)pyridine

[00622] 2-chloro-3-methoxyisonicotinamide (24 g, 0.129 mol) was slurried in dimethyl formamide dimethyl acetal (173 mL, 1.29 mol) and the mixture was heated to 95 °C giving a clear, pale yellow solution. After heating for 1 h, the reaction was cooled and was concentrated on the rotovap and the resulting yellow oil was azeotroped twice with 1,2-dichloroethane (40 mL portions) to ensure complete removal of any residual dimethyl formamide dimethyl acetal. The crude oil thus obtained was immediately dissolved in 50 mL of ethanol and was immediately used in the following step.

[00623] In a separate flask was prepared a mixture of ethanol (500 mL) and acetic acid (AcOH, 130 mL) and the resulting solution was cooled in an ice bath. Once cooled, hydrazine hydrate 64 mL, 1.29 mol) was added dropwise. At this time, the solution containing the crude dimethyl formamide dimethyl acetal adduct as prepared above was transferred dropwise over 15 min by cannula into the previously prepared well-stirred ice-cold mixture containing the hydrazine. During the addition, a pale yellow solid formed in the solution. After the addition was completed, the resulting cloudy yellow mixture was allowed to warm to room temperature and stirred for overnight. The reaction mixture at this time was concentrated on the rotovap to remove some of the ethanol, diluted with additional water and filtered to collect the solid. The solid was washed with additional portions of water, air dried in the funnel then under vacuum to afford 24 g (88 %) of a pale yellow solid as the desired product. LM-MS: m/z =211.6 [M+H]⁺

Step 4: 2-chloro-3-methoxy-4-(1-methyl-1H-1, 2, 4-triazol-3-yl) pyridine

[00624] To a solution of 2-chloro-3-methoxy-4-(1H-1, 2, 4-triazol-3-yl) pyridine (9.00 g, 42.7 mmol) in DMF (80 mL) was treated with potassium carbonate (17.7 g, 128.1 mmol). After cooling the resulting mixture in an ice bath, a solution of iodomethane (8.1 g, 57.6 mmol) in DMF (5 mL) was slowly added dropwise by syringe over 2 min. After the addition was complete, the ice bath was removed and the reaction mixture was allowed to warm to rt. After stirring at room temperature for overnight, LCMS analysis indicated complete and clean conversion to the regioisomeric mixture of products in ~3:1 ratio, respectively. The reaction was cooled in an ice bath and was diluted with water (~50 mL) and the solution was extracted with EtOAc (3 x 40 mL) and the combined extracts were washed with 10% aq. LiCl (2 x 20 mL), water (20 mL) then brine (20 mL), concentrated and purified by column chromatography (PE/EA =5/1) to afford the title compound (5.0 g, 52%) of the major isomer as a pale yellow solid. LM-MS: m/z =225.6[M+H]⁺

Step 5: 3-methoxy-4-(1-methyl-1H-1,2,4-triazol-3-yl)pyridin-2-amine

[00625] To a solution of 2-chloro-3-methoxy-4-(1-methyl-1H-1,2,4-triazol-3-yl) pyridine (5 g, 22.3 mmol), diphenylmethanimine (6.01 g, 33.45 mmol), Sodium tert-butoxide (3.2 g, 33.45 mmol) and DPEphos (2.4 g, 4.46 mmol) in 1,4-dioxane (100 mL) was added Pd₂(dba)₃ (2.0 g, 2.23 mmol). The mixture degassed by N₂ for 3 times and heated to 100°C for 1 hrs. When reaction completed, filtered, filtrate was removed in vacuo, added DCM 50 mL, 2M HCl 50ml stirring at room temperature for 20 min, the aqueous layer was washed with DCM (20 mL x 2). the pH of the aqueous layer was adjusted to 9 with 5% w/v aqueous NaOH, The aqueous layer was extracted with DCM (50 mL×6) and the combined organic fractions dried (Na₂SO₄), filtered and concentrated and chromatography (DCM/MeOH = 20/1) to give the title compound as a white solid. (2.6 g, 56.8 %)


Step 6 : 6-chloro-4-((3-methoxy-4-(1-methyl-1H-1,2,4-triazol-3-yl)pyridin-2-yl)amino)-N-(methyl-d₃)pyridazine-3-carboxamide

[00626] To a solution of 4,6-dichloro-N-(trideuteriomethyl)pyridazine-3-carboxamide (3.97 g, 19.0 mmol) and 3-methoxy-4-(1-methyl-1,2,4-triazol-3-yl)pyridin-2-amine (3.00 g, 14.6 mmol) in THF (50 mL) under N₂ was added LiHMDS (1 M, 43.80 mL, 43.80 mmol) at 0°C resulting a mild exotherm. The reaction was stirred at r.t. When completed, the reaction mixture was cooled to 0°C, quenched by adding satd. NH₄Cl (aq.), diluted with water (100 mL) and extracted by EtOAc (50 mL×3). The combined organic layers were washed by brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (DCM/EtOAc = 3/1) to give the title compound as a yellow solid (2.70 g, 48.9%).

Step 7 : 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[[3-methoxy-4-(1-methyl-1,2,4-triazol-3-yl)-2-pyridyl]amino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00627] To a solution of 6-chloro-4-((3-methoxy-4-(1-methyl-1H-1,2,4-triazol-3-yl)pyridin-2-yl)amino)-N-(methyl-d₃)pyridazine-3-carboxamide (3.6g, 9.5 mmol) and 1-isopropylimidazolidin-2-one (2.4g, 19 mmol) in 1,4-dioxane (30mL) was added cesium carbonate (6.2g, 19 mmol), Pd₂(dba)₃ (2.6 g, 2.9 mmol), and Xantphos (3.3 g, 5.7 mmol).The mixture degassed by N₂ for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product T241 as a yellow solid (0.55 g, 12%). ¹H NMR (400 MHz, CDCl₃): δ 12.20 (s, 1H), 10.16 (s, 1H), 8.28 (d, 1H), 8.22 (s, 1H), 8.14 (s, 1H), 7.51 (d, 1H), 4.43 – 4.32 (m, 1H), 4.23 (t, 2H), 4.03 (s, 3H), 4.00 (s, 3H), 3.51 (t, 2H), 1.23 (d, 6H). LM-MS: m/z =470.3 [M+H]⁺

Example 148:

**Step 1: Example 148b**

[00628] To a solution of **Example 148a** (1.0 g, 5.21 mmol, 1.0 eq) in DCM (15 mL) were added $\text{CD}_3\text{NH}_2\cdot\text{HCl}$ (438 mg, 6.25 mmol, 1.2 eq), DIEA (3.36 g, 26.04 mmol, 5.0 eq) and HATU (2.37 g, 6.25 mmol, 1.2 eq). The reaction mixture was stirred for 4 h at r.t.. The solvent was removed, and the residue was purified by silica gel flash column chromatography, eluted with PE/EtOAc (1/1) to afford the product **Example 148b** (670 mg, 61.8% yield) as an off white solid. LCMS $[\text{M}+1]^+ = 208.2$.

Step 2: Example 148d


[00629] To a solution of **Example 148b** (200 mg, 0.96 mmol, 1.0 eq) and **Example 148c** (196 mg, 0.96 mmol, 1.0 eq) in dry THF (5 mL) was added LiHMDS (1.92 mL, 1 M in THF, 1.92 mmol, 2.0 eq) dropwise at -15°C . The reaction solution was stirred at r.t. for 30 min. After the reaction was completed, the solvent was removed, the residue was purified by silica gel flash column chromatography, eluted with DCM/MeOH (20/1) to afford the product **Example 148d** (212 mg, 58.7% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 376.3$

Step 3: Example 148

[00630] To a solution of **Example 148d** (100 mg, 0.27 mmol, 1.0 eq) in dioxane (3 mL) were added Cs_2CO_3 (260 mg, 0.80 mmol, 3.0 eq), **Example 148e** (102 mg, 0.80 mmol, 3.0 eq) and 3rd Brettphos catalyst (48 mg, 0.05 mmol, 0.2 eq). The reaction mixture was stirred for 6 h at 120°C under N_2 . The mixture was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 148** (26.1 mg, 21.0% yield) as a light yellow solid.

LCMS $[M+1]^+ = 468.4$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO-}d_6$) δ 10.66 (s, 1H), 8.96 (s, 1H), 8.55 (s, 1H), 8.52 (s, 1H), 8.46 (s, 1H), 7.94 (s, 1H), 7.56 - 7.48 (m, 2H), 7.22 (t, $J = 7.8$ Hz, 1H), 3.95 (s, 3H), 3.74 (s, 3H), 3.65 (s, 4H), 1.21 (s, 6H).

Example 149:




Step 1: Example 149c

[00631] A solution of **Example 149a** (200 mg, 0.90 mmol), **Example 149b** (95 mg, 0.91 mmol), **Example 149c** (26 mg, 0.18 mmol), CuI (18 mg, 0.09 mmol) and K₃PO₄ (380 mg, 1.82 mmol) in 1,4-dioxane (5 mL) was stirred at 110°C overnight. The reaction was diluted with CH₃CN (5 mL), filtered and concentrated to give the crude product **Example 149d** (210 mg, quant.) as black oil. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 7.85 (br, 1H), 7.33 (d, $J = 8.0$ Hz, 1H), 6.43 (d, $J = 8.0$ Hz, 1H), 5.98 (s, 2H), 4.13 (s, 2H), 3.91 (t, $J = 4.0$ Hz, 2H), 3.61 (t, $J = 4.0$ Hz, 2H).

Step 2: Example 149

[00632] To a solution of **Example 149d** (210 mg, 0.90 mmol) and **Example 149e** (200 mg, 0.52 mmol) in 1,4-dioxane (5 mL) were added Pd₂(dba)₃ (48 mg, 0.05 mmol), XantPhos (60 mg, 0.10 mmol) and Cs₂CO₃ (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C overnight. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 149** (11.5 mg, 4.1% yield) as an off-white solid. LCMS $[M+1]^+ = 541.3$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO-}d_6$) δ 11.05 (s, 1H), 10.08 (s, 1H), 9.09 (s, 1H), 8.54 (s, 1H), 8.20 (s, 1H), 7.91 (s, 1H), 7.72 - 7.58 (m, 4H), 7.29 (t, $J = 8.0$ Hz, 1H), 4.19 (s, 2H), 3.96 (t, $J = 4.0$ Hz, 2H), 3.93 (s, 3H), 3.72 - 3.70 (m, 5H), 2.95 (br, 1H), 1.06 - 0.97 (m, 4H).

Example 150:




Step 1: Example 150c

[00633] To a solution of **Example 150a** (240 mg, 2.0 mmol) in dioxane (4 mL) were added **Example 150b** (411 mg, 3.0 mmol) and TEA (606 mg, 6.0 mmol). The reaction mixture was stirred at room temperature overnight. The mixture was concentrated, and the residue was triturated with Petroleum ether/EtOAc = 1/1 (5 mL). The solid was collected by filtered and dried to give **Example 150c** (120 mg, 47.5% yield) as a white solid. LCMS $[M+1]^+ = 127.1$

Step 2: Example 150

[00634] To a solution of **Example 150d** (100 mg, 0.27 mmol) in dioxane (3 mL) were added **Example 150c** (50 mg, 0.4 mmol), $Pd_2(dba)_3$ (25 mg, 0.027 mmol), Xantphos (16 mg, 0.027 mmol) and Cs_2CO_3 (176 mg, 0.26 mmol). The mixture was sealed and heated to 120°C for overnight. The mixture was filtrated and concentrated under reduced pressure. The residue was purified by Prep-HPLC to give **Example 150** (47 mg, 37.3% yield) as a yellow solid. LCMS $[M+1]^+ = 467.2$. 1H NMR (400 MHz, Chloroform-*d*) δ 10.94 (s, 1H), 8.14-8.02 (m, 3H), 7.76 (d, $J = 7.7$ Hz, 1H), 7.53 (d, $J = 7.2$ Hz, 1H), 7.26 (s, 1H), 7.14 (s, 1H), 4.15 (s, 3H), 3.99 (s, 3H), 3.81 (s, 3H), 0.69 (br, 4H).

Example 151:




Step 1: Example 151c

[00635] A solution of **Example 151a** (200 mg, 1.1 mmol) in 1,4-dioxane (35 mL) were treated with **Example 151b** (274 mg, 2.0 mmol) and TEA (300 mg, 3.0 mmol). The mixture was stirred at r.t. for 16 h. After reaction completed, the solvent was concentrated, and the residue was suspended in DCM (5 mL), and sonicated. The resulting solid was collected via filtration, and dried to afford the desired crude product **Example 151c** (300 mg, crude, 190.7% yield) as a white solid.


Step 2: Example 151

[00636] To a solution of **Example 151d** (94 mg, 0.25 mmol) and **Example 151c** (150 mg crude, 1.0 mmol) in dioxane (5 mL) were added $\text{Pd}_2(\text{dba})_3$ (91 mg, 0.1 mmol), Xantphos (59 mg, 0.1 mmol) and Cs_2CO_3 (652 mg, 2.0 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for overnight. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 151** (3.7 mg, 0.7% yield) as an off white solid. LCMS $[\text{M}+1]^+ = 484.2$. $^1\text{H NMR}$ (400 MHz, Chloroform-*d*) δ 11.12 (s, 1H), 8.11 (s, 1H), 7.97-7.83 (m, 3H), 7.51-7.44 (m, 1H), 4.71-4.61 (m, 2H), 4.52 (dd, $J = 21.4, 10.8$ Hz, 2H), 4.01 (s, 3H), 3.80 (s, 3H).

Example 152:


[00637] To a solution of **Example 152a** (100 mg, 0.26 mmol,) and **Example 152b** (45 mg, 0.41 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 152** (10.0 mg, 8.4% yield) as a white solid. LCMS $[\text{M}+1]^+ = 457.2$. $^1\text{H NMR}$ (400 MHz, DMSO-*d*₆) δ 11.03 (s, 1H), 10.12 (s, 1H), 9.10 (s, 1H), 8.88 (s, 1H), 8.54 (d, $J = 3.4$ Hz, 1H), 8.08 (s, 1H), 7.69 (s, 1H), 7.62 (d, $J = 9.3$ Hz, 2H), 7.29 (t, $J = 7.7$ Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.96 (s, 1H), 2.37 (d, $J = 3.4$ Hz, 3H), 1.06 (s, 2H), 0.99 (s, 2H).

Example 153:




[00638] To a solution of **Example 153a** (100 mg, 0.26 mmol,) and **Example 153b** (45 mg, 0.41 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 153** (20.0 mg, 17.1% yield) as a white solid. LCMS $[\text{M}+1]^+ = 457.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 11.07 (s, 1H), 10.31 (s, 1H), 9.11 (s, 1H), 8.54 (s, 1H), 8.30 (s, 1H), 8.15 (s, 1H), 7.65 (t, $J = 9.6$ Hz, 2H), 7.29 (d, $J = 7.9$ Hz, 1H), 7.23 (s, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.97 (s, 1H), 2.37 (s, 3H), 1.08 (s, 2H), 1.02 (s, 2H).

Example 154:



[00639] To a solution of **Example 154a** (100 mg, 0.26 mmol,) and **Example 154b** (45 mg, 0.39 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 154** (40.0 mg, 34.2% yield) as a white solid. LCMS $[\text{M}+1]^+ = 457.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 11.03 (s, 1H), 10.27 (s, 1H), 9.14 (s, 1H), 8.55 (s, 2H), 7.78 (s, 1H), 7.65-7.57 (m, 3H), 7.30 (d, $J = 7.9$ Hz, 1H), 3.93 (s, 3H), 3.69 (s, 3H), 2.97 (s, 1H), 2.34 (s, 3H), 1.07 (s, 2H), 1.01 (s, 2H).

Example 155:




Step 1: Example 155c

[00640] To a solution of **Example 155a** (120 mg, 0.585 mmol, 1.0 eq) in DMF (10 mL), was added NaH (234 mg, 60% in mineral oil, 5.85 mmol, 10.0 eq) in portions at 0°C. After **Example 155b** (188.8 mg, 0.878 mmol, 1.5 eq) in DMF was added to the mixture, and the reaction mixture was stirred for 1.5 h at r.t.. The reaction solution was poured into water (50 mL), extracted with EtOAc (50 mL*3), and the combined organic layers were dried over Na₂SO₄ and concentrated. The crude product was purified by silica gel flash column chromatography, eluted with DCM/MeOH (20/1) to afford the product **Example 155c** (24 mg, 11% yield) as a yellow solid. LCMS [M+1]⁺=385.1.

Step 2: Example 155

[00641] To a solution of **Example 155c** (18 mg, 0.047 mmol, 1.0 eq) in dioxane (2 mL) were added Cs₂CO₃ (30.1 mg, 0.094 mmol, 2.0 eq), **Example 155d** (10.2 mg, 0.094 mmol, 2.0 eq), BINAP (5.8 mg, 0.009 mmol, 0.2 eq) and Pd₂(dba)₃·CHCl₃ (4.9 mg, 0.005 mmol, 0.1eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by Prep-TLC (DCM/MeOH = 20/1) to afford the product **Example 155** (8.6 mg, 41% yield) as a yellow solid. LCMS [M+1]⁺=458.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.28 (s, 1H), 10.46 (s, 1H), 9.16 (d, *J* = 7.2 Hz, 2H), 8.66 (s, 1H), 8.20 - 8.11 (m, 2H), 7.54 - 7.46 (m, 2H), 4.00 (s, 3H), 3.88 (s, 3H), 3.05 - 2.93 (m, 1H), 2.56 (s, 3H), 1.17 - 0.99 (m, 4H).

Example 156:

**Step 1: Example 156b**

[00642] To a solution of **Example 156a** (5.0 g, 22.73 mmol, 1.0 eq) in DMF (100 mL) were added K_2CO_3 (9.41 g, 68.18 mmol, 3.0 eq) and CD_3I (6.60 g, 45.45 mmol, 2.0 eq) at $0^\circ C$. The reaction mixture was stirred for 16 h at r.t.. The reaction was diluted with EtOAc and washed with brine. The organic layer dried over Na_2SO_4 and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (1/1) to afford the product **Example 156b** (1.58 g, 29.3% yield) as a yellow solid. LCMS $[M+1]^+ = 238.2$.

Step 2: Example 156c


[00643] To a solution of **Example 156b** (500 mg, 2.11 mmol, 1.0 eq) in MeOH (10 mL) was added Pd/C (200 mg) in portions under N_2 protection, the suspension was degassed under vacuum and purged with H_2 three times, the reaction mixture was stirred for 30 min at r.t. under H_2 balloon. The solid was filtered out, the filtrate was concentrated to afford the product **Example 156c** (425 mg, crude, 97.3% yield) as a gray solid. LCMS $[M+1]^+ = 208.2$.

Step 3: Example 156e

[00644] To a solution of **Example 156c** (425 mg, 2.05 mmol, 1.0 eq) and **Example 156d** (429 mg, 2.05 mmol, 1.0 eq) in dry THF (15 mL) was added LiHMDS (4.11 mL, 1 M in THF, 4.11 mmol, 2.0 eq) dropwise at $-15^\circ C$. The reaction solution was stirred at $-15^\circ C$ for 30 min. After the reaction was completed, the reaction solution was concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (1/2) to afford the product **Example 156e** (170 mg, 27.8% yield) as a yellow solid. LCMS $[M+1]^+ = 380.3$.

Step 4: Example 156

[00645] To a solution of **Example 156e** (100 mg, 0.26 mmol, 1.0 eq) in dioxane (3 mL) were added **Example 156f** (102 mg, 0.79 mmol, 3.0 eq), Cs_2CO_3 (257 mg, 0.79 mmol, 3.0 eq), BINAP (66 mg, 0.11 mmol, 0.4 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (54 mg, 0.05 mmol, 0.2 eq). The reaction solution was stirred for 4 h at 120°C under N_2 . The mixture was filtered and the filtrate was concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to give the desired product **Example 156** (18.0 mg, 28.2% yield) as a white solid. LCMS $[\text{M}+1]^+ = 472.3$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.89 (s, 1H), 9.60 (s, 1H), 8.99 (s, 1H), 8.54 (s, 1H), 8.01 (s, 1H), 7.62 (d, $J = 8.1$ Hz, 1H), 7.49 (d, $J = 7.8$ Hz, 1H), 7.24 (t, $J = 8.1$ Hz, 1H), 3.71 (s, 2H), 3.67 (s, 2H), 1.19 (s, 6H).


Example 157:**Step 1: Example 157c**

[00646] A solution of **Example 157a** (200 mg, 1.4 mmol) in 1,4-dioxane (3 mL) were treated with **Example 157b** (374 mg, 2.8 mmol) and TEA (420 mg, 4.2 mmol). The mixture was stirred at r.t. for 16 h. After reaction completed, the solvent was concentrated, and the residue was suspended in DCM (5 mL), and sonicated. The resulting solid was collected via filtration, and dried to afford the desired product **Example 157c** (130 mg, 62% yield) as a white solid. LCMS $[\text{M}+1]^+ = 151.0$

Step 2: Example 157

[00647] To a solution of **Example 157d** (157 mg, 0.42 mmol) and **Example 157c** (300 mg, 2.1 mmol) in dioxane (5 mL) were added $\text{Pd}_2(\text{dba})_3$ (91 mg, 0.1 mmol), Xantphos (59 mg, 0.1 mmol) and Cs_2CO_3 (325.8 mg, 1.0 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for overnight. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 157** (14.8 mg, 7.2% yield) as a off white solid. LCMS $[\text{M}+1]^+ = 491.2$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.90 (s, 1H), 9.84 (s, 1H), 9.01 (s, 1H), 8.54 (s, 1H), 7.98 (s, 1H), 7.62 (d, $J = 7.7$ Hz, 1H), 7.48 (d, $J = 7.8$ Hz, 1H), 7.26 (d, $J = 8.1$ Hz, 1H), 6.16-6.44 (t, 1H), 4.09 (s, 2H), 3.93 (s, 5H), 3.71 (s, 3H), 3.06 (s, 1H).

Example 158:




Step 1: Example 158c

[00648] To a solution of **Example 158a** (200 mg, 2.29 mmol) in dioxane (5 mL) were added **Example 158b** (400 mg, 2.91 mmol), and TEA (795 mg, 7.87 mmol). The reaction mixture was stirred at room temperature overnight. Then the mixture was diluted with EtOAc, washed by water, brine, and dried over anhydrous Na_2SO_4 . The solution was concentrated to afford crude **Example 158c** (170 mg, 57.2% yield) and used directly for next step.

Step 2: Example 158

[00649] To a solution of **Example 158d** (100 mg, 0.26 mmol) and **Example 158c** (120 mg, 0.92 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 158** (14.0 mg, 13.1% yield) as a white solid. LCMS $[\text{M}+1]^+ = 471.3$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.90 (s, 1H), 9.74 (s, 1H), 9.01 (s, 1H), 8.54 (s, 1H), 7.99 (s, 1H), 7.62 (d, $J = 7.6$ Hz, 1H), 7.49 (d, $J = 7.9$ Hz, 1H), 7.25 (d, $J = 8.1$ Hz, 1H), 4.16 (d, $J = 11.1$ Hz, 4H), 3.93 (s, 3H), 3.79 (s, 2H), 3.71 (s, 3H), 3.18 (s, 3H).

Example 159:

**Step 1: Example 159c**

[00650] To a solution of **Example 159a** (1.0 g, 4.2 mmol, 1.0 eq) in dioxane (15 mL) were added Cs_2CO_3 (2.7g, 8.4 mmol, 2.0 eq), **Example 159b** (861.8 mg, 5.04 mmol, 1.2 eq), BINAP (523.3 mg, 0.84 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (434.7 mg, 0.42 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 159c** (1.0 g, 67% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 369.3$.

Step 2: Example 159e


[00651] To a solution of **Example 159c** (1.0 g, 2.7 mmol, 1.0 eq) in THF (10 mL) was added **Example 159d** (40.5 mL, 1.0 M in THF, 40.5 mmol, 15.0 eq) dropwise at 0°C under N_2 protection. The mixture was stirred for 0.5 h at r.t.. The reaction was poured into saturated aqueous of NH_4Cl (70 mL) and extracted with EtOAc (100 mL*3). The combined organic layer was washed with brine, dried over Na_2SO_4 , and concentrated. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 159e** (640 mg, 68% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 350.2$.

Step 3: Example 159

[00652] To a solution of **Example 159e** (300 mg, 0.86 mmol, 1.0 eq) in dioxane (5 mL) were added Cs_2CO_3 (560 mg, 1.72 mmol, 2.0 eq), **Example 159f** (176 mg, 1.0 mmol, 1.2 eq), BINAP (107.2 mg, 0.172 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (89 mg, 0.086 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to give 270 mg crude product (90% purity) and further purified by Prep-TLC (DCM/MeOH = 20/1) to afford the product **Example 159** (120 mg, 32% yield) as an off-white solid. LCMS $[\text{M}+1]^+ = 486.4$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.89 (s, 1H), 10.28 (s, 1H), 9.17 (s, 1H), 9.02 (s, 1H), 8.64 (dd, $J = 4.8, 1.8$ Hz, 1H),

8.27 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.99 (dd, $J = 10.8, 8.4$ Hz, 1H), 7.58 (dd, $J = 8.1, 2.1$ Hz, 1H), 7.34 (dd, $J = 7.8, 4.8$ Hz, 1H), 5.28 (s, 1H), 3.30 (s, 3H), 3.01-2.86 (m, 1H), 1.49 (s, 6H), 1.14 - 0.97 (m, 4H).

Example 160:



Step 1: 2-chloro-3-methoxy-4-[1-(trideuteriomethyl)-1,2,4-triazol-3-yl]pyridine

[00653] To a solution of 2-chloro-3-methoxy-4-(1H-1,2,4-triazol-3-yl)pyridine (10.00 g, 47.5 mmol) in DMF (50 mL) was treated with potassium carbonate (13.1 g, 95.0 mmol). a solution of trideuterioiodomethane (8.26 g, 57.0 mmol) in DMF (5 mL) was slowly added dropwise by syringe over 2 min. The reaction was stirring at room temperature for 4 h, The reaction was cooled in an ice bath and was diluted with water (50 mL) and the solution was extracted with EtOAc (3×50 mL) and the combined extracts were washed water (50 mL) then brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (PE/EtOAc = 3/1) to give the title compound as a yellow solid (8.0 g, 74%).

Step 2: 3-methoxy-4-[1-(trideuteriomethyl)-1,2,4-triazol-3-yl]pyridin-2-amine

[00654] To a solution of 2-chloro-3-methoxy-4-(1-methyl-1H-1,2,4-triazol-3-yl)pyridine (8.0 g, 35.1 mmol), diphenylmethanimine (7.64 g, 42.2 mmol), Sodium tert-butoxide (6.75 g, 70.3 mmol) and DPEphos (1.88 g, 3.51 mmol) in 1,4-dioxane (100 mL) was added Pd₂(dba)₃ (3.22 g, 3.51 mmol). The mixture degassed by N₂ for 3 times and heated to 100°C for 1 hr. When the reaction completed, filtered, filtrate was removed in vacuo, Added DCM 50 mL, 2N HCl 50 mL stirring at room temperature for 20 min, the aqueous layer was washed with DCM (20 mL×2). the pH of the aqueous layer was adjusted to 9 with 5% w/v aqueous NaOH, The aqueous layer was extracted with DCM (50 mL×6) and the combined organic fractions dried (Na₂SO₄), filtered and concentrated and chromatography (DCM/MeOH = 20/1) to give the title compound as a white solid. (3.2 g, 43.7 %)


Step 3: 6-chloro-4-[[3-methoxy-4-(1-methyl-1,2,4-triazol-3-yl)-2-pyridyl]amino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00655] To a solution of 4,6-dichloro-N-(trideuteriomethyl)pyridazine-3-carboxamide (4.18 g, 20.0 mmol) and 3-methoxy-4-[1-(trideuteriomethyl)-1,2,4-triazol-3-yl]pyridin-2-amine (3.20 g, 15.4 mmol) in THF (50 mL) under N₂ was added LiHMDS (1 M, 46.2 mL, 46.2 mmol) at 0°C resulting a mild exotherm. The reaction was stirred at r.t. When completed, The reaction mixture was cooled to 0°C, quenched by adding satd. NH₄Cl (aq.), diluted with water (100 mL) and extracted by EtOAc (50 mL×3). The combined organic layers were washed by brine (50 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude product which was purified by column chromatography (DCM/EtOAc=3/1) to give the title compound as a yellow solid (2.70g, 46.1%).

Step 4 : 6-(3-isopropyl-2-oxo-imidazolidin-1-yl)-4-[[3-methoxy-4-[1-(trideuteriomethyl)-1,2,4-triazol-3-yl]-2-pyridyl]amino]-N-(trideuteriomethyl)pyridazine-3-carboxamide

[00656] To a solution of 6-chloro-4-[[3-methoxy-4-(1-methyl-1,2,4-triazol-3-yl)-2-pyridyl]amino]-N-(trideuteriomethyl)pyridazine-3-carboxamide (2.8 g, 7.4 mmol) and 1-isopropylimidazolidin-2-one (1.9 g, 15 mmol) in 1,4-dioxane (30 mL) was added cesium carbonate (4.8 g, 15 mmol), Pd₂(dba)₃ (2.0 g, 2.2 mmol), and Xantphos (2.6 g, 4.4 mmol). The mixture degassed by N₂ for 3 times and heated to 120 °C for 4 hrs. When reaction completed, filtered, filtrate was removed in vacuo, chromatography (PE/EtOAc = 50 / 50 then DCM/MeOH = 97/3) to give the desired product T 311 as a yellow solid (0.58 g, 17%). ¹H NMR (400 MHz, CDCl₃) δ 12.20 (s, 1H), 10.16 (s, 1H), 8.28 (d, 1H), 8.22 (s, 1H), 8.14 (s, 1H), 7.51 (d, 1H), 4.43 – 4.32 (m, 1H), 4.23 (t, 2H), 4.00 (s, 3H), 3.51 (t, 2H), 1.23 (d, 6H). LM-MS: m/z =473.3[M+H]⁺

Example 161:

**Step 1: Example 161b**

[00657] To a solution of **Example 161a** (383 mg, 1.0 mmol) in dioxane (10 mL) were added NH_2Boc (176 mg, 1.5 mmol), $\text{Pd}_2(\text{dba})_3$ (91 mg, 0.1 mmol), Xantphos (60 mg, 0.1 mmol) and Cs_2CO_3 (652 mg, 2.0 mmol). The mixture was sealed and heated to 100°C for 8 h. The mixture was cooled to room temperature, diluted by EtOAc, washed by water, dried over anhydrous Na_2SO_4 , and then concentrated under reduced pressure. The residue was purified by silica gel flash column chromatography, eluted with DCM/MeOH (10/1) to afford the product **Example 161b** (401 mg, 86.4% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 465.2$.

Step 2: Example 161c


A solution of **Example 161b** (401 mg, 0.86 mmol) in HCl/EtOAc (2 mL) was stirred at room temperature for 3 h. Then the mixture was concentrated, and used at next step directly.

Step 3: Example 161

[00658] To a solution of **Example 161c** (40 mg, 0.1 mmol) and **Example 161d** (10.4 mg, 0.1 mmol) in pyridine (3 mL) was added POCl_3 (77 mg, 0.5 mmol) at 0°C . The reaction mixture was stirred at 0°C for 10 min. Then water (5 mL) was added dropwise, and the mixture was extracted with DCM (6 mL). The organic phase was dried over anhydrous Na_2SO_4 , concentrated under reduced pressure. The residue was purified by Prep-HPLC to give **Example 161** (12.1 mg, 26.9% yield) as a white solid. LCMS $[\text{M}+1]^+ = 451.2$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.01 (s, 1H), 10.95 (s, 1H), 9.11 (s, 1H), 8.53 (s, 1H), 8.00 (s, 1H), 7.63 (d, $J = 7.8$ Hz, 1H), 7.51 (d, $J = 7.8$ Hz, 1H), 7.25 (t, $J = 7.9$ Hz, 1H), 4.97 (s, 1H), 4.80 (s, 1H),

3.92 (s, 3H), 3.67 (s, 3H), 2.98 (s, 1H), 2.19 (s, 1H), 1.58 (d, $J = 23.5$ Hz, 1H), 1.21 (s, 2H), 1.10 - 1.00 (m, 4H).

Example 162:




Step 1: Example 162c

[00659] A solution of **Example 162a** (800 mg, 17.34 mmol) in 1,4-dioxane (35 mL) was treated with **Example 162b** (1.0 g, 7.34 mmol) and Na_2CO_3 (1.2 g, 11.1 mmol). The mixture was stirred at r.t. for 16 h. After reaction completed, the solvent was concentrated, the residue was suspended in DCM (5 mL), sonicated and the resulting solid was collected via filtration, dried to afford the desired product **Example 162c** (300 mg, 35.3% yield) as a white solid.

Step 2: Example 162

[00660] To a solution of **Example 162d** (58 mg, 0.5 mmol) and **Example 162c** (94 mg, 0.25 mmol) in dioxane (3 mL) were added $\text{Pd}_2(\text{dba})_3$ (46 mg, 0.05 mmol), Xantphos (30 mg, 0.05 mmol) and Cs_2CO_3 (163 mg, 0.5 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for overnight. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 162** (2.7 mg, yield: 11.8%) as a off white solid. LCMS $[\text{M}+1]^+ = 457.2$. $^1\text{H NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ 10.90 (s, 1H), 9.67 (s, 1H), 9.00 (s, 1H), 8.54 (s, 1H), 8.00 (s, 1H), 7.62 (d, $J = 7.6$ Hz, 1H), 7.49 (d, $J = 7.8$ Hz, 1H), 7.25 (d, $J = 8.4$ Hz, 1H), 5.62 (d, $J = 6.2$ Hz, 1H), 4.38 (s, 1H), 4.17 (s, 2H), 3.93 (s, 3H), 3.76 – 3.69 (m, 5H).

Example 163:



Step 1: Example 163c

[00661] To a solution of **Example 163a** (7.5 g, 43.0 mmol, 1.0 eq) in THF (100 mL) was added **Example 163b** (21.4 mL, 3.0 M in THF, 64.2 mmol, 1.5 eq) dropwise at -20°C under N₂ protection. The mixture was stirred at r.t. for 0.5 h. The reaction was quenched with saturated aqueous of NH₄Cl (80 mL) and extracted with EtOAc (150 mL*2). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 5/1) to afford the product **Example 163c** (4.0 g, 45.8% yield) as yellow oil. LCMS [M+1]⁺ = 206.1.

Step 2: Example 163d

To a solution of **Example 163c** (3.8 g, 18.0 mmol, 1.0 eq) in DCM (100 mL) was added MnO₂ (40.1 g, 450 mmol, 25.0 eq), and the reaction mixture was stirred at r.t. for 16 h. The reaction mixture was filtered and the filtrate was concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 10/1) to afford the product **Example 163d** (2.5 g, 66.5% yield) as an off-white solid. LCMS [M+1]⁺ = 204.1.

Step 3: Example 163f

[00662] To a solution of **Example 163d** (100 mg, 0.49 mmol, 1.0 eq) in dioxane (2.0 mL) were added Cs₂CO₃ (320 mg, 0.98 mmol, 2.0 eq), **Example 163e** (41.9 mg, 0.49 mmol, 1.0 eq), BINAP (61.3 mg, 0.098 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (50.9 mg, 0.049 mmol, 0.1 eq). The reaction mixture was stirred for 2 h at 80°C under N₂ protection. After cooling to r.t., the solvent was concentrated under

vacuum. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 3/1) to afford the product **Example 163f** (60 mg, 48.4% yield) as a yellow solid. LCMS $[M+1]^+ = 253.2$.

Step 4: Example 163i

[00663] To a solution of **Example 163g** (28.0 g, 243.4 mmol, 1.0 eq) in MeOH (400 mL) at 0°C (ice-water bath) was added **Example 163h** (16.0 g, 267.7 mmol, 1.1 eq) dropwise. After addition, the reaction was stirred at r.t. for 16 h. The solution was concentrated in vacuum and then swapped with THF (80 mL*4) to remove the MeOH residue. After removal of solvent by evaporation, **Example 163i** (15.0 g, 83.8% yield) was obtained as yellow oil.

Step 5: Example 163k

[00664] To a solution of **Example 163j** (40.0 g, 279.7 mmol, 1.0 eq) in dry THF (400 mL) was added LDA (181.8 mL, 2 M in THF, 363.6 mmol, 1.3 eq) at -70°C dropwise, which was stirred for additional 40 min at the same temperature after addition. I₂ (92.3 g, 363.6 mmol, 1.3 eq) in THF (100 mL) was added to the solution dropwise. After addition, the reaction was warmed to r.t. and stirred for 2 h. The mixture was quenched with saturated aqueous of NH₄Cl (200 mL), extracted with EtOAc (200 mL*3). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 80/1) to afford the desired product **Example 163k** (20.0 g, 26.6% yield) as a white solid. LCMS $[M+1]^+ = 270.1$.

Step 6: Example 163l

[00665] To a solution of **Example 163k** (15.0 g, 55.6 mmol, 1.0 eq) in NMP (50 mL) was added CuCN (9.9 g, 111.2 mmol, 2.0 eq), and the mixture was stirred for 5 h at 120°C. After cooling to room temperature, the mixture was diluted with EtOAc (200 mL) and filtered. The filtrate was washed with brine (50 mL*3), dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 50/1) to afford the desired product **Example 163l** (7.5g, 80.1% yield) as a yellow solid. LCMS $[M+1]^+ = 169.2$.

Step 7: Example 163m

[00666] To a solution of **Example 163l** (5.0 g, 29.8 mmol, 1.0 eq) and **Example 163i** (4.4 g, 59.6 mmol, 2.0 eq) in THF (150 mL) was added *t*-BuOK (7.3 g, 65.6 mmol, 2.2 eq) in portions at 0°C. After addition, the mixture was warmed to r.t. and stirred for 5 h. The mixture was concentrated in vacuum. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 1/1) to afford the product **Example 163m** (2.6 g, 42.7% yield) as a white solid. LCMS $[M+1]^+ = 225.2$.

Step 8: Example 163o


[00667] To a solution of **Example 163m** (1.1 g, 4.9 mmol, 1.0 eq) in dioxane (20 mL) were added Cs₂CO₃ (3.26 g, 9.8 mmol, 2.0 eq), **Example 163n** (1.37 g, 5.39 mmol, 1.1 eq), BINAP (623 mg, 0.98 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (517 mg, 0.49 mmol, 0.1 eq). The reaction mixture was stirred for 3 h at 90°C under N₂ protection. After the reaction was completed, the solvent was removed, and the residue was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 163o** (1.4 g, 89.3% yield) as a yellow solid. LCMS $[M+1]^+ = 326.3$.

Step 9: Example 163p

[00668] A solution of **Example 163o** (1.4 g, 4.29 mmol, 1.0 eq) in TFA (20 mL) was stirred for 1 h at 70°C. After the reaction was completed, it was concentrated in vacuum. The residue was dissolved in MeOH (20 mL) and basified with NaHCO₃ (910 mg, 8.58 mmol, 2.0 eq). The solid was filtered out, and the filtrate was concentrated in vacuum. The residue was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 163p** (750 mg, 84.9% yield) as a light gray solid. $[M+1]^+ = 206.2$.

Step 10: Example 163

[00669] To a solution of **Example 163f** (22 mg, 0.11 mmol, 1.0 eq) and **Example 163p** (30 mg, 0.12 mmol, 0.9 eq) in dioxane (2 mL) were added Cs₂CO₃ (77.3 mg, 0.22 mmol, 2.0 eq), Pd₂(dba)₃.CHCl₃ (12.3 mg, 0.011 mmol, 0.1 eq) and BINAP (15.0 mg, 0.022 mmol, 0.2 eq). The reaction mixture was stirred for 3 h at 110°C under N₂ protection. The reaction solution was filtered and the filtrate was concentrated in *vacuo*. The crude product was purified by Prep-TLC (DCM/MeOH = 30/1) to afford the desired product **Example 163** (13.8 mg, 27.6% yield) as a light yellow solid. LCMS $[M+1]^+ = 422.3$. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.36 (s, 1H), 10.91 (s, 1H), 9.71 (s, 1H), 8.96 (s, 1H), 8.67 (s, 1H), 8.14 (d, *J* = 5.4 Hz, 1H), 7.50 (d, *J* = 5.1 Hz, 1H), 4.00 (s, 3H), 3.92 (s, 3H), 3.16 (q, *J* = 7.2 Hz, 2H), 2.12-2.04 (m, 1H), 1.15 (t, *J* = 7.2 Hz, 3H), 0.89-0.82 (m, 4H).

Example 164:**Step 1: 2-methylazetidine-1-carboxamide**

[00670] To a solution of 2-methylazetidine hydrochloride (120 mg, 1.12 mmol), phenyl carbamate (184 mg, 1.34 mmol) in 1,4-dioxane (20 mL) was added NEt₃ (0.5 mL, 3.36 mmol) and stirred at r.t. for 16 hrs. After reaction completed, the solvent was concentrated, the residue was suspended in DCM (5 mL), sonicated and the resulting solid was collected via filtration, dried to afford the desired product (150 mg, 89%) as a white solid. LM-MS: $m/z = 115.1[M+H]^+$

Step 2:**4-((2-methoxy-3-(1-methyl-1H-1,2,4-triazol-3-yl)phenyl)amino)-N-(methyl-d3)-6-(2-methylazetidine-1-carboxamido)pyridazine-3-carboxamide**

To a solution of 6-chloro-4-((2-methoxy-3-(1-methyl-1H-1,2,4-triazol-3-yl)


[00671] phenyl)amino)-N-(methyl-d₃)pyridazine-3-carboxamide (244.7 mg, 0.65 mmol) and 2-methylazetidine-1-carboxamide (150 mg, 1.0 mmol) in 1,4-dioxane (10 mL) was added Pd₂(dba)₃ (60

mg, 0.065 mmol), XantPhos (75 mg, 0.13 mmol) and Cs_2CO_3 (635 mg, 1.95 mmol). The mixture was sealed, degassed by nitrogen for 3 times and stirred at 120°C for 4 hrs. When completed, the reaction was cooled to r.t. and filtered. The filtrate was purified directly by prep-HPLC to give the racemic product (125 mg, yield: 42%) as a white solid. LM-MS: $m/z = 455.3[\text{M}+\text{H}]^+$

[00672] The racemic compound was separated by SFC, and got enantiomer A (49.5 mg, R.T.: 3.008 min). ^1H NMR (400 MHz, CDCl_3) δ 11.19 (s, 1H), 8.15 (s, 1H), 8.10 (s, 1H), 7.93 (s, 1H), 7.82 (dd, 1H), 7.53 (dd, 1H), 7.31 – 7.24 (m, 2H), 4.53 (dd, 1H), 4.17 – 4.01 (m, 2H), 4.00 (s, 3H), 3.82 (s, 3H), 2.53 – 2.36 (m, 1H), 1.98 – 1.83 (m, 1H), 1.50 (d, 3H). ee% = 100% ; LM-MS: $m/z = 455.3[\text{M}+\text{H}]^+$

[00673] enantiomer B (49.9 mg, R.T.: 4.240 min). ^1H NMR (400 MHz, CDCl_3) δ 11.13 (s, 1H), 8.13 (s, 1H), 8.09 (s, 1H), 7.96 (s, 1H), 7.80 (dd, 1H), 7.53 (d, 1H), 7.32 – 7.21 (m, 2H), 4.52 (dd, 1H), 4.14 – 4.02 (m, 2H), 4.00 (s, 3H), 3.82 (s, 3H), 2.57 – 2.35 (m, 1H), 2.05 – 1.81 (m, 1H), 1.50 (d, 3H). ee% = 100% ; LM-MS: $m/z = 455.3 [\text{M}+\text{H}]^+$

Example 165:




Step 1: Example 165c

[00674] To a solution of **Example 165a** (270 mg, 3.1 mmol) in dioxane (5 mL) were added **Example 165b** (500 mg, 3.64 mmol), and TEA (795 mg, 7.87 mmol). The reaction mixture was stirred at room temperature overnight. Then the mixture was diluted with EtOAc, washed by water, brine, and dried over anhydrous Na_2SO_4 . The solution was concentrated to afford crude **Example 165c** (200 mg, 49.6% yield) and used directly for next step.

Step 2: Example 165

[00675] To a solution of **Example 165d** (100 mg, 0.26 mmol) and **Example 165c** (120 mg, 0.92 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), xantphos (30 mg, 0.052 mmol) and Cs_2CO_3 (340 mg, 1.04 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 165** (46.0 mg, 37.6% yield) as a white solid. LCMS $[\text{M}+\text{H}]^+ = 471.50$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 10.89 (s, 1H), 9.67 (s, 1H), 8.99 (s, 1H), 8.54 (s, 1H), 8.01 (s, 1H), 7.62 (d, $J = 7.9$ Hz, 1H), 7.48 (d, $J = 7.9$ Hz, 1H), 7.24 (s, 1H), 5.54 (s, 1H), 3.93 (s, 3H), 3.83 (br, 4H), 3.71 (s, 3H), 1.33 (s, 3H).

Example 166:

**Step 1: Example 166c**

[00676] To a solution of **Example 166a** (2.0 g, 8.0 mmol, 1.0 eq) in dioxane (20 mL) were added Cs_2CO_3 (5.2 g, 16 mmol, 2.0 eq), **Example 166b** (1.36 g, 16 mmol, 2.0 eq), BINAP (997 mg, 1.6 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (800 mg, 0.8 mmol, 0.1 eq). The reaction mixture was stirred for 2 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 1/1) to afford the product **Example 166c** (1.9 g, 86% yield) as yellow oil. LCMS $[\text{M}+1]^+ = 284.2$.

Step 2: Example 166e


To a solution of **Example 166c** (1.9 g, 7.0 mmol, 1.0 eq) in THF (20 mL) was added **Example 166d** (210 mL, 1.0 M in THF, 210 mmol, 30.0 eq) dropwise at 0°C under N_2 protection. The mixture was stirred for 0.5 h at r.t.. The reaction was poured into saturated aqueous of NH_4Cl (200 mL) and extracted with EtOAc (150 mL*3). The combined organic layer were washed with brine, dried over Na_2SO_4 and concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 2/1) to afford the product **Example 166e** (850 mg, 47% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 265.2$.

Step 3: Example 166

[00677] To a solution of **Example 166e** (650 mg, 2.33 mmol, 1.2 eq) in dioxane (10 mL) were added Cs_2CO_3 (1.3 g, 3.88 mmol, 2.0 eq), **Example 166f** (400 mg, 1.94 mmol, 1.0 eq), BINAP (241.7 mg, 0.388 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (200.8 mg, 0.194 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to obtain 580 mg crude product (90% purity), which was further purified by Prep-HPLC (Prep-C18, 5 μM XBridge column, 19×150 mm, Waters; gradient elution of 35% MeCN in water to 55% MeCN in

water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to afford the product **Example 166** (401.4 mg, 48% yield) as a white solid. LCMS $[\text{M}+1]^+ = 434.4$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.30 (s, 1H), 10.94 (s, 1H), 9.70 (s, 1H), 9.21 (s, 1H), 8.66 (s, 1H), 8.13 (d, $J = 5.1$ Hz, 1H), 7.49 (d, $J = 5.1$ Hz, 1H), 3.99 (s, 3H), 3.86 (s, 3H), 3.11-2.96 (m, 1H), 2.15-2.04 (m, 1H), 1.19 - 1.03 (m, 4H), 0.92 - 0.80 (m, 4H).

Example 169:



Step 1: Example 169c

[00678] To a solution of **Example 169a** (123 mg, 1.0 mmol) and **Example 169b** (206 mg, 1.5 mmol) in dioxane (3 mL) was added Na_2CO_3 (212 mg, 2.0 mmol). The reaction mixture was stirred at r.t. overnight. The mixture was filtrated, the solid was washed by DCM, and the combined filtrate was used directly at next step.

Step 2: Example 169d


[00679] To a solution of **Example 169c** was added imidazole (350 mg, 5.0 mmol) in portions, followed by addition of TBSCl (300 mg, 2.0 mmol). The reaction mixture was stirred at room temperature overnight. The mixture was concentrated, and the crude product was purified by silica gel flash column chromatography, eluted with (DCM/MeOH = 30/1) to afford the product **Example 169d** (60 mg, 24.5% yield) as a white solid.

Step 3: Example 169f

[00680] To a solution of **Example 169d** (60 mg, 0.26 mmol) in dioxane (5 mL) were added **Example 169e** (100 mg, 0.26 mmol), $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.027 mmol), Xantphos (16 mg, 0.027 mmol) and Cs_2CO_3 (130 mg, 0.40 mmol). The mixture was sealed and heated to 110°C for overnight. The mixture was filtrated and concentrated under reduced pressure to give crude **Example 169f** (190 mg, quant.).

Step 4: Example 169

[00681] To a solution of **Example 169f** (130 mg, crude, 0.22 mmol) in DCM (4 mL) was added TFA (1 mL). The mixture was stirred at r.t. for 30 min. The mixture was concentrated under reduced pressure, and the residue was purified by Prep-HPLC (Prep-C18, 5 μ M XBridge column, 19 \times 150 mm, Waters; gradient elution of 35% MeCN in water to 55% MeCN in water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to give **Example 169** (2 mg, 1.9% yield over 2 steps) as a white solid. LCMS $[\text{M}+1]^+ = 471.2$. $^1\text{H NMR}$ (300 MHz, CDCl_3) δ 10.94 (s, 1H), 8.11 - 8.03 (m, 3H), 7.77 (d, $J = 8.0$ Hz, 1H), 7.51 (d, $J = 8.0$ Hz, 1H), 4.14 (br, 2H), 4.00 (s, 3H), 3.90 (s, 2H), 3.80 (s, 5H), 2.83 (br, 1H), 1.25 (s, 1H).

Example 171:**Step 1: Example 171c**

[00682] To a solution of **Example 171a** (5.0 g, 26.3 mmol, 1.0 eq) in **Example 171b** (50 mL) was added NaH (3.15 g, 60% in mineral oil, 78.9 mmol, 3.0 eq) in portions at 0°C. The reaction mixture was stirred at r.t. for 3 h. The reaction mixture was diluted with EtOAc (100 mL), quenched with 2 N aqueous HCl (15 mL), washed with brine (50 mL), dried over Na_2SO_4 and concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 3/1) to give the desired product **Example 171c** (4.1 g, 64.0% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 244.2$.

Step 2: Example 171d

[00683] To a solution of **Example 171c** (4.1 g, 16.80 mmol, 1.0 eq) and K_2CO_3 (2.55 g, 18.48 mmol, 1.1 eq) in DMF (50 mL) was added CD_3I (2.68 g, 18.48 mmol, 1.1 eq) at 0°C. The reaction mixture was stirred at r.t. for 6 h. The mixture was diluted with EtOAc (100 mL) and washed with brine (50 mL*3). The organic layer was dried by Na_2SO_4 and concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum ether/EtOAc = 3/1) to give the desired product **Example 171d** (2.83 g, 64.5% yield) as colorless oil. LCMS $[\text{M}+1]^+ = 261.1$.

Step 3: Example 171e

[00684] To a solution of **Example 171d** (2.83 g, 10.84 mmol, 1.0 eq) in AcOH (20 mL) was added conc. HCl (40 mL). The reaction solution was heated at 100°C for 16 h. After cooled to r.t., the mixture was concentrated, diluted with H₂O (20 mL) and extracted with EtOAc (30 mL*3). The combined organic layer was washed with brine (50 mL), dried by Na₂SO₄ and concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum ether/EtOAc = 3/1) to give the desired product **Example 171e** (1.12 g, 54.7% yield) as a white solid. LCMS [M+1]⁺ = 189.2.

Step 4: Example 171f

[00685] To a solution of **Example 171e** (1.12 g, 5.94 mmol, 1.0 eq) in CH₃CN (15 mL) was added POCl₃ (2 mL). The reaction solution was heated at 85°C for 1 h. After cooled to r.t., the mixture was concentrated and diluted with EtOAc (5 mL). The solution was added to a mixed solution of EtOAc (20 mL) and saturated aqueous NaHCO₃ (20 mL). After separation, the aqueous layer was extracted with EtOAc (20 mL*3). The combined organic layer was washed with brine (20 mL), dried by Na₂SO₄ and concentrated. The crude product was purified by silica gel flash column chromatography (Petroleum ether/EtOAc = 8/1) to give the desired product **Example 171f** (1.05 g, 85.4% yield) as a white solid. LCMS [M+1]⁺ = 207.1.


Step 5: Example 171h

To a solution of **Example 171f** (100 mg, 0.48 mmol, 1.0 eq) in dioxane (4 mL) were added **Example 171g** (37 mg, 0.43 mmol, 0.9 eq), Cs₂CO₃ (315 mg, 0.97 mmol, 2.0 eq), BINAP (30 mg, 0.048 mmol, 0.1 eq) and Pd₂(dba)₃.CHCl₃ (50 mg, 0.048 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂. The mixture was filtered and the filtrate was concentrated. The crude product was purified by prep-TLC (Petroleum ether/EtOAc = 3/1) to give the desired product **Example 171h** (40.0 mg, 32.4% yield) as a light yellow solid. LCMS [M+1]⁺ = 256.3.

Step 6: Example 171

[00686] To a solution of **Example 171h** (40 mg, 0.16 mmol, 1.0 eq) in dioxane (3 mL) were added **Example 171i** (29 mg, 0.14 mol, 0.9 eq), Cs₂CO₃ (102 mg, 0.31mmol, 2.0 eq), BINAP (10 mg, 0.016 mmol, 0.1 eq) and Pd₂(dba)₃.CHCl₃ (16 mg, 0.016 mmol, 0.1 eq). The reaction solution was stirred for 6 h at 110°C under N₂. The reaction mixture was filtered and the filtrate was concentrated. The crude product was purified by prep-TLC (DCM/MeOH = 20/1) to give the desired product **Example 171** (8.2 mg, 12.3% yield) as an off-white solid. LCMS [M+1]⁺ = 424.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.05 (s, 1H), 10.90 (s, 1H), 8.89 (s, 1H), 8.57 (s, 1H), 8.03 (s, 1H), 7.65 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.53 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.26 (d, *J* = 7.8 Hz, 1H), 3.96 (s, 3H), 3.72 (s, 3H), 3.11 (s, 2H), 2.07-1.96 (m, 1H), 0.79 (d, *J* = 6.0 Hz, 4H).

Example 172:




Step 1: Example 172c

[00687] To a solution of **Example 172a** (150 mg, 0.72 mmol, 1.0 eq) in dioxane (5 mL) were added **Example 172b** (88 mg, 0.58 mmol, 0.8 eq), Cs_2CO_3 (472 mg, 1.45 mmol, 2.0 eq), BINAP (45 mg, 0.072 mmol, 0.1 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (75 mg, 0.072 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 . The mixture was filtered and the filtrate was concentrated. The crude product was purified by prep-TLC (Petroleum Ether/EtOAc = 1/1) to give the desired product **Example 172c** (62 mg, 26.5% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 323.3$.

Step 2: Example 172

[00688] To a solution of **Example 172c** (62 mg, 0.19 mmol, 1.0 eq) in dioxane (5 mL) were added **Example 172d** (33 mg, 0.19 mmol, 1.0 eq), Cs_2CO_3 (126 mg, 0.39 mmol, 2.0 eq), BINAP (24 mg, 0.039 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (20 mg, 0.019 mmol, 0.1 eq). The reaction mixture was stirred for 6 h at 110°C under N_2 . The mixture was filtered and the filtrate was concentrated. The crude product was purified by prep-TLC (DCM/MeOH = 20/1) to give the desired product **Example 172** (16.1 mg, 18.3% yield) as a white solid. LCMS $[\text{M}+1]^+ = 459.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.96 (s, 1H), 10.02 (s, 1H), 8.95 (s, 1H), 8.92 (s, 1H), 8.67 (dd, $J = 4.8, 2.1$ Hz, 1H), 8.34 (d, $J = 2.4$ Hz, 1H), 8.28 (dd, $J = 7.8, 2.1$ Hz, 1H), 7.78 (dd, $J = 8.7, 2.4$ Hz, 1H), 7.68 (d, $J = 8.7$ Hz, 1H), 7.33 (dd, $J = 7.8, 4.8$ Hz, 1H), 5.07 (s, 1H), 3.34 (s, 3H), 3.08 (s, 2H), 1.45 (s, 6H).

Example 173:



Step 1: Example 173b


[00689] To a solution of **Example 173a** (1.0 g, 3.7 mmol, 1.0 eq) in NaOH aqueous solution (11.1 mL, 1M, 11.1 mmol, 3.0 eq) was added $(\text{Boc})_2\text{O}$ (1.29 g, 5.92 mmol, 1.6 eq) in DCM (11.1 mL). The reaction mixture was stirred for 16 h at r.t.. Upon completion of the reaction, two phases were separated. The

organic layer was washed with water (10 mL), dried over Na_2SO_4 and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (1/1) to afford the product **Example 173b** (505 mg, 74% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 184.2$.

Step 2: Example 173

[00690] To a solution of **Example 173c** (150 mg, 0.39 mmol, 1.0 eq.) in dioxane (5 mL) were added Cs_2CO_3 (254.3 mg, 0.78 mmol, 2.0 eq), **Example 173b** (143.5 mg, 0.78 mmol, 2.0 eq), BINAP (48.6 mg, 0.078 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (40.4 mg, 0.039 mmol, 0.1 eq). The reaction mixture was stirred for 3 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by Prep-TLC (DCM/MeOH = 10/1) to afford the crude product 42 mg (90% purity), which was purified by Prep-HPLC (Prep-C18, $5 \mu\text{M}$ XBridge column, $19 \times 150 \text{ mm}$, Waters; gradient elution of 35% MeCN in water to 55% MeCN in water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to afford the product **Example 173** (14.2 mg, 8% yield) as a white solid. LCMS $[\text{M}+1]^+ = 431.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.48 (br, 1H), 10.97 (s, 1H), 10.22 (br, 1H), 9.04 (s, 1H), 8.56 (s, 1H), 7.66 - 7.52 (m, 2H), 7.26 (t, $J = 7.8 \text{ Hz}$, 1H), 7.01 (br, 1H), 6.74 (br, 2H), 3.95 (s, 3H), 3.72 (s, 3H), 2.97-2.81 (m, 1H), 1.18 - 0.93 (m, 4H).

Example 174:




Step 1: Example 174c

[00691] To a solution of **Example 174a** (500 mg, 2.45 mmol) in dioxane (20 mL) were added **Example 174b** (417 mg, 2.45 mmol), K_2CO_3 (676 mg, 4.9 mmol), BINAP (304 mg, 0.49 mmol) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (252 mg, 0.25 mmol). The reaction mixture was stirred at 90°C for 16 h under N_2 . The reaction mixture was concentrated and the crude was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 2/1) to afford the product **Example 174c** (170 mg, 20.5% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 338.2$.

Step 2: Example 174

[00692] To a mixture of **Example 174c** (170 mg, 0.5 mmol) in dioxane (5 mL) were added **Example 174d** (86 mg, 0.5 mmol), Cs_2CO_3 (326 mg, 1.0 mmol), BINAP (125 mg, 0.1 mmol) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (51 mg, 0.05 mmol). The reaction mixture was stirred at 110°C for 4 h under N_2 . The reaction mixture was concentrated and the residue was purified by Prep-TLC (DCM/EtOAc = 1/2) to afford the product **Example 174** (66.6 mg, 28.0% yield) as a white solid. LCMS $[\text{M}+1]^+ = 474.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 12.00 (s, 1H), 10.25 (s, 1H), 9.02 (s, 1H), 8.95 (s, 1H), 8.66 (dd, $J = 4.8, 2.1 \text{ Hz}$, 1H), 8.29 (dd, $J = 7.8, 2.1 \text{ Hz}$, 1H), 7.99 (dd, $J = 10.8, 8.1 \text{ Hz}$, 1H), 7.60 (dd, $J = 8.4, 1.8 \text{ Hz}$, 1H), 7.35 (dd, $J = 7.8, 1.5 \text{ Hz}$, 1H), 5.29 (s, 1H), 3.34 (s, 3H), 3.11 (q, $J = 7.2 \text{ Hz}$, 2H), 1.48 (s, 6H), 1.13 (t, $J = 7.2 \text{ Hz}$, 3H).

Example 175:




Step 1: Example 175c

[00693] To a solution of **Example 175a** (500 mg, 2.46 mmol, 1.0 eq) in dioxane (10 mL) were added K_2CO_3 (680 mg, 4.93 mmol, 2.0 eq), **Example 175b** (293 mg, 2.46 mmol, 1.0 eq), Xantphos (285 mg, 0.493 mmol, 0.2 eq) and $Pd_2(dba)_3 \cdot CHCl_3$ (255 mg, 0.246 mmol, 0.1 eq). The reaction mixture was stirred for 16 h at 110°C under N_2 protection. After the reaction was completed, the solvent was concentrated, and the crude product was purified by silica gel flash column chromatography, eluted with (DCM/MeOH = 30/1) to afford the product **Example 175c** (380 mg, 53.8% yield) as a yellow solid. LCMS $[M+1]^+ = 287.1$.

Step 2: Example 175

[00694] To a solution of **Example 175c** (300 mg, 1.05 mmol, 1.0 eq) in dioxane (6 mL) were added Cs_2CO_3 (683.9 mg, 2.10 mmol, 2.0 eq), **Example 175d** (193.5 mg, 0.94 mmol, 0.9 eq), BINAP (130.7 mg, 0.21 mmol, 0.2 eq) and $Pd_2(dba)_3 \cdot CHCl_3$ (108.5 mg, 0.105 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography, eluted with (DCM/MeOH = 20/1) to afford the product 160 mg (crude, 80% purity), which was further purified by Prep-HPLC (Prep-C18, 5 μM XBridge column, 19 \times 150 mm, Waters; gradient elution of 35% MeCN in water to 55% MeCN in water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to afford the product **Example 175** (58.3 mg, 12.2% yield) as a yellow solid. LCMS $[M+1]^+ = 456.3$. 1H NMR (300 MHz, $DMSO-d_6$) δ 12.49 (s, 1H), 10.53 (s, 1H), 9.78 (s, 1H), 8.96 (s, 1H), 8.68 (s, 1H), 8.36 (d, $J = 5.4$ Hz, 1H), 7.92 (t, $J = 8.1$ Hz, 1H), 7.82 (d, $J = 8.4$ Hz, 1H), 7.59 (d, $J = 7.2$ Hz, 1H), 7.50 (d, $J = 5.1$ Hz, 1H), 4.01 (s, 3H), 3.94 (s, 3H), 3.17 (q, $J = 7.2$ Hz, 2H), 1.17 (t, $J = 7.2$ Hz, 3H).

Example 176:



Step 1: Example 176c


[00695] To a solution of **Example 176a** (500 mg, 2.46 mmol, 1.0 eq) in dioxane (10 mL) were added K_2CO_3 (680 mg, 4.93 mmol, 2.0 eq), **Example 176b** (293 mg, 2.46 mmol, 1.0 eq), Xantphos (285 mg, 0.493 mmol, 0.2 eq) and $Pd_2(dba)_3 \cdot CHCl_3$ (255 mg, 0.246 mmol, 0.1 eq). The reaction mixture was

stirred for 16 h at 110°C under N₂ protection. The solvent was concentrated. The crude product was purified by silica gel flash column chromatography, eluted with (DCM/MeOH = 30/1) to afford the product **Example 176c** (310 mg, 44% yield) as a yellow solid. LCMS [M+1]⁺ = 287.1.


Step 2: Example 176

[00696] To a solution of **Example 176c** (310 mg, 1.084 mmol, 1.0 eq) in dioxane (6 mL) were added Cs₂CO₃ (683.9 mg, 2.168 mmol, 2.0 eq), **Example 176d** (167.8 mg, 0.975 mmol, 0.9 eq), BINAP (130.7 mg, 0.217 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (112.2 mg, 0.108 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography, eluted with (DCM/MeOH = 15/1) to afford the crude product 120 mg (80% purity) and further purified by Prep-HPLC to afford the product **Example 176** (61.1 mg, 13% yield) as an off-white solid. LCMS [M+1]⁺ = 423.2. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.06 (s, 1H), 10.54 (s, 1H), 9.19 (s, 1H), 8.97 (s, 1H), 8.77 (dd, *J* = 4.8, 2.1 Hz, 1H), 8.30 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.97 - 7.84 (m, 2H), 7.57 (dd, *J* = 6.6, 1.5 Hz, 1H), 7.34 (dd, *J* = 7.8, 4.8 Hz, 1H), 3.36 (s, 3H), 3.14 (q, *J* = 7.2 Hz, 2H), 1.13 (t, *J* = 7.2 Hz, 3H).


Example 177:



[00697] To a solution of **Example 177a** (371 mg, 1.0 mmol) and **Example 177b** (119 mg, 1.0 mmol) in dioxane (10 mL) were added Pd₂(dba)₃ (91.5 mg, 0.1 mmol), Xantphos (59.5 mg, 0.1 mmol) and Cs₂CO₃ (489 mg, 1.5 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 100°C for 3 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 177** (154.2 mg, 33.9% yield) as a pale yellow solid. LCMS [M+1]⁺ = 455.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.27 (s, 1H), 10.37 (s, 1H), 8.85 (s, 1H), 8.54 (s, 1H), 7.96 (s, 1H), 7.84 (t, *J* = 7.8 Hz, 1H), 7.74 (dd, *J* = 17.3, 8.3 Hz, 2H), 7.58 (d, *J* = 7.7 Hz, 1H), 7.50 (d, *J* = 7.2 Hz, 1H), 7.34 (d, *J* = 8.6 Hz, 1H), 3.93 (s, 3H), 3.73 (s, 3H), 3.09 (t, *J* = 7.0 Hz, 2H), 1.11 (t, *J* = 6.9 Hz, 3H).

Example 178:

[00698] To a solution of **Example 178a** (371 mg, 1.0 mmol) and **Example 178b** (111 mg, 1.0 mmol) in dioxane (10 mL) were added Pd₂(dba)₃ (91.5 mg, 0.1 mmol), Xantphos (59.5 mg, 0.1 mmol) and Cs₂CO₃ (489 mg, 1.5 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 3 h. When completed, the reaction was cooled to r.t, diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 178** (8.4 mg, 1.9% yield) as a pale yellow solid. LCMS [M+1]⁺ = 447.2. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.09 (s, 1H), 9.45 (s, 1H), 8.73 (s, 1H), 8.54 (s, 1H), 7.66 (d, *J* = 7.4 Hz, 1H), 7.57 (d, *J* = 7.8 Hz, 2H), 7.27 (t, *J* = 7.9 Hz, 1H), 5.90 (s, 1H), 3.93 (s, 3H), 3.72 (s, 3H), 3.57 (s, 3H), 3.04 - 2.97 (m, 2H), 2.17 (s, 3H), 1.12 - 1.08 (m, 3H).

Example 179:**Step 1: Example 179c**


[00699] To a solution of **Example 179a** (650 mg, 3.19 mmol, 1.0 eq) in dioxane (10 mL) were added NaHCO₃ (548.7 mg, 6.38 mmol, 2.0 eq), **Example 179b** (354.1 mg, 0.319 mmol, 1.0 eq), BINAP (398.7 mg, 0.638 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (331.2 mg, 0.319 mmol, 0.1 eq). The reaction mixture was stirred for 2 h at 100 °C under N₂ protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 2/1) to afford the product **Example 179c** (250 mg, 28% yield) as a yellow solid. LCMS [M+1]⁺ = 279.2.

Step 2: Example 179

[00700] To a solution of **Example 179c** (250 mg, 0.899 mmol, 1.0 eq) in dioxane (5 mL) were added Cs₂CO₃ (586 mg, 1.80 mmol, 2.0 eq), **Example 179d** (139 mg, 0.809 mmol, 0.9 eq), BINAP (112 mg, 0.180 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (90 mg, 0.087 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography, eluted with DCM/MeOH (20/1) to afford the crude product 140 mg (90% purity) and further purified by Prep-HPLC to afford the product

Example 179 (84.6 mg, 23% yield) as an off-white solid. LCMS $[M+1]^+ = 415.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.98 (s, 1H), 9.64 (s, 1H), 8.83 (s, 1H), 8.65 (dd, $J = 4.8, 1.8$ Hz, 1H), 8.53 (brs, 1H), 8.27 (dd, $J = 7.8, 2.1$ Hz, 1H), 7.31 (dd, $J = 7.8, 4.8$ Hz, 1H), 6.11 (s, 1H), 3.64 (s, 3H), 3.33 (s, 3H), 3.06 (q, $J = 7.2$ Hz, 2H), 2.24 (s, 3H), 1.11 (t, $J = 7.2$ Hz, 3H).

Example 180:




Step 1: Example 180c

[00701] To a solution of **Example 180a** (1.0 g, 4.9 mmol, 1.0 eq) in dioxane (20 mL) were added NaHCO_3 (842.8 mg, 9.8 mmol, 2.0 eq), **Example 180b** (543.9 mg, 4.9 mmol, 1.0 eq), BINAP (610.5 mg, 0.98 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (507.2 mg, 0.49 mmol, 0.1 eq). The reaction mixture was stirred for 2 h at 100°C under N_2 protection. After cooled to room temperature, the solvent was removed, and the crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 2/1) to afford the product **Example 180c** (380 mg, 28% yield) as a yellow solid. LCMS $[M+1]^+ = 279.2$.

Step 2: Example 180

[00702] To a solution of **Example 180c** (340 mg, 1.22 mmol, 1.0 eq) in dioxane (5 mL) were added Cs_2CO_3 (795.4 mg, 2.44 mmol, 2.0 eq), **Example 180d** (226.6 mg, 0.32 mmol, 1.1 eq), BINAP (152.0 mg, 0.24 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (126.3 mg, 0.12 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the crude product (220 mg) and further purified by Prep-HPLC (Prep-C18, 5 μM XBridge column, 19×150 mm, Waters; gradient elution of 35% MeCN in water to 55% MeCN in water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to afford the product **Example 180** (61.9 mg, 11% yield) as a white solid. LCMS $[M+1]^+ = 448.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 12.38 (s, 1H), 9.61 (s, 1H), 9.00 (s, 1H), 8.83 (s, 1H), 8.67 (s, 1H), 8.16 (d, $J = 5.4$ Hz, 1H), 7.47 (d, $J = 5.1$ Hz, 1H), 6.13 (s, 1H), 4.01 (s, 3H), 3.92 (s, 3H), 3.68 (s, 3H), 3.08 (q, $J = 7.2$ Hz, 2H), 2.26 (s, 3H), 1.15 (t, $J = 7.2$ Hz, 3H).

Example 181:


**Step 1: Example 181c**

[00703] To a solution of **Example 181a** (1 g, 10.0 mmol) in dioxane (15 mL) were added **Example 181b** (2 g, 14.6 mmol), and TEA (2 g, 19.8 mmol). The reaction mixture was stirred at room temperature for 16 h. Then the mixture was diluted with EtOAc, washed by water, brine, and dried over anhydrous Na_2SO_4 . The solution was concentrated to afford crude residue, which was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 1/1) to afford the product **Example 181c** (1 g, 69.8% yield) as a white solid.


Step 2: Example 181

[00704] To a solution of **Example 181d** (100 mg, 0.26 mmol) and **Example 181c** (38.3 mg, 0.26 mmol) in DMA (2.5 mL) were added $\text{Pd}_2(\text{dba})_3$ (24 mg, 0.026 mmol), Xantphos (15.6 mg, 0.026 mmol) and Cs_2CO_3 (172 mg, 0.53 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 130°C for 2 h. When completed, the reaction was cooled to r.t., diluted with MeOH (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 181** (24.3 mg, 18.9% yield) as a off white solid. LCMS $[\text{M}+1]^+ = 484.3$. ^1H NMR (400 MHz, Chloroform-*d*) δ 11.33 (s, 1H), 10.08 (s, 1H), 9.43 (s, 1H), 8.96 (s, 1H), 8.41 (s, 1H), 8.04 (d, $J = 7.8$ Hz, 1H), 7.91 (d, $J = 7.8$ Hz, 1H), 7.66 (t, $J = 7.9$ Hz, 1H), 4.41 (s, 2H), 4.35 (s, 3H), 4.18 (s, 2H), 4.13 (s, 3H), 3.40 (d, $J = 9.1$ Hz, 1H), 2.91 (s, 6H).


10.25 (s, 1H), 9.02 (s, 1H), 8.95 (s, 1H), 8.66 (dd, $J = 4.8, 2.1$ Hz, 1H), 8.29 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.99 (dd, $J = 10.8, 8.4$ Hz, 1H), 7.62 (d, $J = 7.8$ Hz, 1H), 7.35 (dd, $J = 7.8, 4.5$ Hz, 1H), 5.29 (s, 1H), 3.34 (s, 3H), 3.09 (s, 2H), 1.48 (s, 6H).

Example 185:


[00712] To a solution of **Example 185a** (50 mg, 0.15 mmol, 1.0 eq) in dioxane (2 mL) were added **Example 185b** (30 mg, 0.15 mmol, 1.0 eq), Cs_2CO_3 (96 mg, 0.29 mmol, 2.0 eq), BINAP (18 mg, 0.029 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (15 mg, 0.015 mmol, 0.1 eq). The reaction solution was stirred for 4 h at 110°C under N_2 . The mixture was filtered and the filtrate was concentrated. The crude product was purified by prep-TLC (DCM/MeOH = 20/1) to give the desired product **Example 185** (35.0 mg, 46.8% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 510.4$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.38 (s, 1H), 10.25 (s, 1H), 9.53 (s, 1H), 8.94 (s, 1H), 8.68 (s, 1H), 8.18 (d, $J = 5.1$ Hz, 1H), 8.00 (dd, $J = 10.8, 8.4$ Hz, 1H), 7.59 (d, $J = 8.1$ Hz, 1H), 7.52 (d, $J = 5.1$ Hz, 1H), 5.29 (s, 1H), 4.01 (s, 3H), 3.93 (s, 3H), 3.11 (s, 2H), 1.50 (s, 6H).

Example 188:

[00713] To a solution of **Example 188a** (100 mg, 0.27 mmol) and **Example 188b** (30 mg, 0.27 mmol) in dioxane (10 mL) were added $\text{Pd}_2(\text{dba})_3$ (24.6 mg, 0.027 mmol), Xantphos (16 mg, 0.027 mmol) and Cs_2CO_3 (131 mg, 0.40 mmol). The mixture was degassed by nitrogen for 3 times and stirred at 110°C for 4 h. When completed, the reaction was cooled to r.t, diluted with EtOAc (5 mL) and filtered. The filtrate was purified directly by Prep-HPLC to give the desired product **Example 188** (27.5 mg, 22.4% yield) as a pale yellow solid. LCMS $[\text{M}+1]^+ = 445.2$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 11.07 (s, 1H), 10.19 (s, 1H), 8.85 (s, 1H), 8.55 (s, 1H), 7.97 (d, $J = 9.2$ Hz, 1H), 7.67-7.61 (m, 3H), 7.43 (d, $J = 9.2$ Hz, 1H), 7.25 (t, $J = 8.0$ Hz, 1H), 3.94 (s, 3H), 3.73 (s, 3H), 3.08 (q, $J = 7.6$ Hz, 2H), 1.11 (t, $J = 7.1$ Hz, 3H).

Example 189:

[00714] To a solution of **Example 189b** (869 mg, 3.45 mmol, 1.1 eq) in dioxane (100 mL) were added **Example 189a** (650 mg, 3.14 mmol, 1.0 eq), Cs_2CO_3 (2.04 g, 6.27 mmol, 2.0 eq), BINAP (391 mg, 0.63 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (325 mg, 0.31 mmol, 0.1 eq). The reaction solution was stirred for 4 h at 110°C under N_2 . The solvent was concentrated, and the residue was purified by silica gel flash column chromatography, eluted with DCM/MeOH (v/v = 25/1) to give the desired product **Example 189** (1.1 g, crude, 85% purity) as a yellow solid and further purified by reverse phase preparative MPLC to give the desired product **Example 189** (512.6 mg, 38.6% yield) as a white solid. LCMS $[\text{M}+1]^+ = 424.4$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.05 (s, 1H), 10.90 (s, 1H), 8.89 (s, 1H), 8.56 (s, 1H), 8.04 (s, 1H), 7.65 (dd, $J = 7.8, 1.8$ Hz, 1H), 7.53 (dd, $J = 8.1, 1.5$ Hz, 1H), 7.26 (t, $J = 7.8$ Hz, 1H), 3.72 (s, 3H), 3.13 (q, $J = 7.2$ Hz, 2H), 2.10-1.94 (m, 1H), 1.13 (t, $J = 7.2$ Hz, 3H), 0.79 (d, $J = 6.0$ Hz, 4H).

Example 190:**Step 1: Example 190c**


[00715] To a solution of **Example 190a** (150 mg, 0.73 mmol, 1.0 eq) and **Example 190b** (79 mg, 0.73 mmol, 1.0 eq) in dioxane (3 mL) were added K_2CO_3 (201 mg, 1.46 mmol, 2.0 eq), $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (75 mg, 0.073 mmol, 0.1 eq) and BINAP (15.0 mg, 0.15 mmol, 0.2 eq). The reaction mixture was stirred for

16 h at 80°C under N₂ protection. The reaction solution was filtered and the filtrate was concentrated in vacuum. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 1/1) to afford the desired product **Example 190c** (65 mg, 32.0% yield) as a yellow solid. LCMS [M+1]⁺ = 280.2.

Step 2: Example 190

[00716] To a solution of **Example 190c** (60 mg, 0.22 mmol, 1.0 eq) in dioxane (3 mL) and **Example 190d** (37 mg, 0.22 mmol, 1.0 eq) were added Cs₂CO₃ (139.7 mg, 0.44 mmol, 2.0 eq), Pd₂(dba)₃.CHCl₃ (22.2 mg, 0.022 mmol, 0.1 eq) and BINAP (25 mg, 0.044 mmol, 0.2 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. The mixture was filtered and the filtrate was concentrated in vacuum. The crude product was purified by prep-TLC (DCM/MeOH = 30/1) to afford the desired product **Example 190** (9.2 mg, 10.3% yield) as a light yellow solid. LCMS [M+1]⁺ = 416.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.94 (s, 1H), 10.41 (s, 1H), 8.94 (s, 1H), 8.73 (s, 1H), 8.67 - 8.59 (m, 1H), 8.29 (d, *J* = 8.1 Hz, 1H), 8.09 (d, *J* = 9.0 Hz, 1H), 7.49 (d, *J* = 9.0 Hz, 1H), 7.34 (dd, *J* = 7.8, 4.8 Hz, 1H), 3.35 (s, 3H), 3.09 (s, 2H), 2.54 (s, 3H).

Example 191:




Step 1: Example 191c


[00717] To a solution of **Example 191a** (150 mg, 0.73 mmol, 1.0 eq) and **Example 191b** (89.5 mg, 0.73 mmol, 1.0 eq) in dioxane (10 mL) were added K₂CO₃ (201 mg, 1.46 mmol, 2.0 eq), Pd₂(dba)₃.CHCl₃ (75.3 mg, 0.073 mmol, 0.1 eq) and BINAP (90.6 mg, 0.15 mmol, 0.2 eq). The reaction mixture was stirred for 16 h at 80°C under N₂ protection. The mixture was filtered and the filtrate was concentrated in vacuum. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 40/1) to afford the desired product **Example 191c** (60 mg, 28.4% yield) as a yellow solid. LCMS [M+1]⁺ = 294.2.

Step 2: Example 191

[00718] To a solution of **Example 191c** (60 mg, 0.22 mmol, 1.0 eq) and **Example 191d** (35 mg, 0.22 mmol, 1.0 eq) in dioxane (3 mL) were added Cs₂CO₃ (132 mg, 0.44 mmol, 2.0 eq), Pd₂(dba)₃.CHCl₃ (21 mg, 0.022 mmol, 0.1 eq) and BINAP (25.4 mg, 0.044 mmol, 0.2 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. The mixture was filtered and the filtrate was concentrated in vacuum. The crude product was purified by prep-TLC (DCM/MeOH = 30/1) to afford the desired product **Example 191** (19.2 mg, 22.3% yield) as a light yellow solid. LCMS [M+1]⁺ = 430.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.97 (s, 1H), 10.35 (s, 1H), 9.07 (s, 1H), 8.98 (s, 1H), 8.68 (dd, *J* = 4.8, 1.8 Hz, 1H), 8.30 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.43 - 7.30 (m, 2H), 3.35 (s, 3H), 3.11 (s, 2H), 2.44 (s, 3H), 2.33 (s, 3H).

Example 193:


[00719] To a solution of **Example 193a** (750 mg, 2.93 mmol, 1.0 eq) in dioxane (25 mL) were added **Example 193b** (900 mg, 4.39 mmol, 1.5 eq), Cs_2CO_3 (1.91 g, 5.86 mmol, 2.0 eq), BINAP (365 mg, 0.59 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (303 mg, 0.29 mmol, 0.1 eq). The reaction solution was stirred for 6 h at 110°C under N_2 . The reaction solution was concentrated and the residue was purified by silica gel flash column chromatography, eluted with DCM/MeOH (v/v = 20/1) to give the desired product **Example 193** (411.8 mg, 33.1% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 425.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 12.36 (s, 1H), 10.91 (s, 1H), 9.71 (s, 1H), 8.96 (s, 1H), 8.67 (s, 1H), 8.14 (d, $J = 5.1$ Hz, 1H), 7.50 (d, $J = 5.1$ Hz, 1H), 4.00 (s, 3H), 3.92 (s, 3H), 3.14 (s, 2H), 2.13-2.01 (m, 1H), 0.91-0.79 (m, 4H).

Example 195:

[00720] To a solution of **Example 195b** (700 mg, 2.73 mmol, 1.0 eq) in dioxane (20 mL) were added Cs_2CO_3 (1.8 g, 5.46 mmol, 2.0 eq), **Example 195a** (847.6 mg, 4.09 mmol, 1.5 eq), BINAP (340.2 mg, 0.55 mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3\text{CHCl}_3$ (279.5 mg, 0.27 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the crude product 1.2 g, which was further purified by Prep-HPLC (Prep-C18, 5 μM XBridge column, 19 x 150 mm, Waters; gradient elution of 35% MeCN in water to 55% MeCN in water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to afford the product **Example 195** (401.9 mg, 35% yield) as an off-white solid. LCMS $[\text{M}+1]^+ = 427.3$. $^1\text{H NMR}$ (300 MHz, $\text{DMSO}-d_6$) δ 11.05 (s, 1H), 10.89 (s, 1H), 8.89 (s, 1H), 8.56 (s, 1H), 8.03 (s, 1H), 7.65 (dd, $J = 7.8, 1.5$ Hz, 1H),

7.53 (dd, $J = 7.8, 1.5$ Hz, 1H), 7.26 (t, $J = 7.8$ Hz, 1H), 3.72 (s, 3H), 3.11 (s, 2H), 2.08- 1.95 (m, 1H), 0.79 (d, $J = 6.3$ Hz, 4H).

Example 196:




Step 1: Example 196b

[00721] To a solution of **Example 196a** (100 mg, 0.75 mmol, 1.0 eq) in 1M NaOH aqueous solution (2.25 mL, 2.25 mmol, 3.0 eq) was added a solution of (Boc)₂O (261.6 mg, 1.2 mmol, 1.6 eq) in DCM (5 mL). The reaction mixture was stirred for 16 h at r.t.. Upon completion of the reaction, two phases were separated. The organic layer was washed with water (5 mL), dried over Na₂SO₄ and concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 20/1) to afford the product **Example 196b** (78 mg, 53% yield) as yellow oil. LCMS [M+1]⁺ = 198.2.

Step 2: Example 196

[00722] To a solution of **Example 196c** (50 mg, 0.13 mmol, 1.0 eq.) in DMA (2 mL) were added Cs₂CO₃ (84.8 mg, 0.26 mmol, 2.0 eq), **Example 196b** (51 mg, 0.26 mmol, 2.0 eq), BINAP (16.2 mg, 0.03 mmol, 0.2 eq) and Pd₂(dba)₃.CHCl₃ (13.5 mg, 0.01 mmol, 0.1 eq). The reaction mixture was stirred for 2 h at 110°C under N₂ protection. After cooled to room temperature, the reaction was diluted with EtOAc (20 mL), washed with brine, dried over Na₂SO₄ and concentrated. The crude product was purified by Prep-TLC (DCM/MeOH = 10/1) to afford the product **Example 196** (9.1 mg, 16% yield) as a yellow solid. LCMS [M+1]⁺ = 445.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.10 (br, 1H), 10.98 (s, 1H), 10.03 (brs, 1H), 9.04 (s, 1H), 8.56 (s, 1H), 7.61 (d, $J = 8.1$ Hz, 2H), 7.26 (t, $J = 7.8$ Hz, 1H), 7.08 (brs, 1H), 6.45 (brs, 1H), 3.95 (s, 3H), 3.71 (s, 3H), 2.95-2.81 (m, 1H), 2.09 (s, 3H), 1.12-0.96 (m, 4H).

Example 198:




Step 1: Example 198

[00723] To a solution of **Example 198a** (200 mg, 0.97 mmol, 1.0 eq) in dioxane (5 mL) were added K_2CO_3 (267.7 mg, 1.94 mmol, 2.0 eq), **Example 198b** (115.4 mg, 0.97 mmol, 1.0 eq) and Xantphos (112.3 mg, 0.194 mmol, 0.2 eq) and $Pd_2(dba)_3 \cdot CHCl_3$ (100.4 mg, 0.097 mmol, 0.1 eq). The reaction mixture was stirred for 6 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 198c** (120 mg, 43% yield) as a yellow solid.

LCMS $[M+1]^+ = 290.2$

Step 2: Example 198


[00724] To a solution of **Example 198c** (60 mg, 0.21 mmol, 1.0 eq) in dioxane (3 mL) were added Cs_2CO_3 (136.9 mg, 0.42 mmol, 2.0 eq), **Example 198d** (65.2 mg, 0.32 mmol, 1.5 eq), BINAP (26.2 mg, 0.042 mmol, 0.2 eq) and $Pd_2(dba)_3 \cdot CHCl_3$ (21.7 mg, 0.021 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product 60 mg crude product, which was further purified by Prep-HPLC (Prep-C18, 5 μ M XBridge column, 19 \times 150 mm, Waters; gradient elution of 35% MeCN in water to 55% MeCN in water over a 7 min period, where the aqueous phase contains 10 mM NH_4HCO_3 + 0.5% ammonia) to afford the product **Example 198** (15.4 mg, 16% yield) as an off-white solid. LCMS $[M+1]^+ = 461.3$. 1H NMR (300 MHz, $DMSO-d_6$) δ 11.29 (s, 1H), 10.40 (brs, 1H), 8.90 (s, 1H), 8.57 (s, 1H), 7.99 (s, 1H), 7.92-7.86 (m, 1H), 7.85-7.71 (m, 2H), 7.62 (dd, $J = 8.1, 1.8$ Hz, 1H), 7.54 (d, $J = 7.2$ Hz, 1H), 7.38 (t, $J = 7.8$ Hz, 1H), 3.76 (s, 3H), 3.10 (s, 2H).

Example 199:

[00725] To a solution of **Example 199a** (60 mg, 0.204 mmol, 1.2 eq) in dioxane (3 mL) were added Cs_2CO_3 (110.8 mg, 0.34 mmol, 2.0 eq), **Example 199b** (35 mg, 0.17 mmol, 1.0 eq), BINAP (21.2 mg, 0.034 mmol, 0.2 eq) and $Pd_2(dba)_3 \cdot CHCl_3$ (17.6 mg, 0.017 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N_2 protection. After cooled to room temperature, the solvent was removed. The crude product was purified by Prep-TLC (DCM/MeOH = 15/1) to afford the product **Example 199** (19.3 mg, 24% yield) a yellow solid. LCMS $[M+1]^+ = 462.3$. 1H NMR (300 MHz, $DMSO-d_6$) δ 12.48 (s, 1H), 10.52 (s, 1H), 9.78 (s, 1H), 8.96 (s, 1H), 8.67 (s, 1H), 8.36 (d, $J = 5.1$ Hz, 1H), 7.92 (t, $J = 8.1$ Hz,

1H), 7.82 (d, $J = 8.4$ Hz, 1H), 7.59 (d, $J = 6.9$ Hz, 1H), 7.50 (d, $J = 5.4$ Hz, 1H), 3.94 (s, 3H), 3.13 (s, 2H).

Example 200:



Step 1: Example 200b

[00726] To a solution of **Example 200a** (3.2 g, 18.93 mmol, 1.0 eq) in EtOH (20 mL) and H₂O (4 mL) was added NaOH (908 mg, 22.72 mmol, 1.2 eq) at r.t.. The mixture was stirred for 1 h at 50 °C. After the reaction was completed, the solvent was removed. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 20/1) to afford the product **Example 200b** (1.2 g, 33.9% yield) as a white solid. LCMS [M+1]⁺ = 187.2.

Step 2: Example 200c

[00727] A solution of **Example 200b** (1.2 g, 6.45 mmol, 1.0 eq) in DMF-DMA (3.07 g, 25.8 mmol, 4.0 eq) was stirred for 2 h at 95°C. After the reaction completed, it was concentrated under vacuum. The residue was dissolved in AcOH (5 mL) and EtOH (20 mL), and N₂H₄·H₂O (3.2 g, 80% in water, 51.6 mmol, 8.0 eq) was added dropwise at 0°C. The reaction mixture was stirred for 2 h at r.t.. After completion, the mixture was concentrated in vacuo. The crude product was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 200c** (600 mg, 44.4% yield) as a white solid. LCMS [M+1]⁺ = 211.2.

Step 3: Example 200d

[00728] To a solution of **Example 200c** (600 mg, 2.86 mmol, 1.0 eq) in DMF (10 mL) were added K₂CO₃ (789 mg, 5.72 mmol, 2.0 eq) and CD₃I (621 mg, 4.29 mmol, 1.5 eq) at 0°C. The reaction mixture was stirred for 4 h at r.t.. The reaction was diluted with EtOAc and washed with brine. The organic layer was dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography, eluted with Petroleum Ether/EtOAc (v/v = 1/3) to afford the product **Example 200d** (550 mg, 84.8% yield) as a yellow solid. LCMS [M+1]⁺ = 228.2.

Step 4: Example 200f

[00729] To a solution of **Example 200d** (550 mg, 2.42 mmol, 1.0 eq) in dioxane (10 mL) were added Cs₂CO₃ (1.57 g, 4.84 mmol, 2.0 eq), **Example 200e** (399 mg, 2.9 mmol, 1.2 eq), BINAP (302 mg, 0.48

mmol, 0.2 eq) and $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (251 mg, 0.24 mmol, 0.1 eq). The reaction mixture was stirred for 3 h at 110 °C under N_2 protection. After the reaction was completed, the solvent was removed, and the residue was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 200f** (505 mg, 63.5% yield) as a yellow solid. LCMS $[\text{M}+1]^+ = 329.3$.


Step 5: Example 200g

[00730] The solution of **Example 200f** (505 mg, 1.54 mmol, 1.0 eq) in TFA (20 mL) was stirred for 1 h at 70 °C. After the reaction was completed, it was concentrated in vacuum. The residue was dissolved in MeOH (20 mL) and basified with NaHCO_3 (910 mg, 8.58 mmol, 2.0 eq). The solid was filtered out, and the filtrate was concentrated in vacuum. The residue was purified by silica gel flash column chromatography (DCM/MeOH = 20/1) to afford the product **Example 200g** (180 mg, 56.3% yield) as a light gray solid. $[\text{M}+1]^+ = 209.4$.

Step 6: Example 200

[00731] To a solution of **Example 200g** (40 mg, 0.19 mmol, 1.0 eq) and **Example 200h** (59.1 mg, 0.23 mmol, 1.2 eq) in dioxane (2 mL) were added Cs_2CO_3 (125 mg, 0.38 mmol, 2.0 eq), $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (19.9 mg, 0.019 mmol, 0.1 eq) and BINAP (23.9 mg, 0.038 mmol, 0.2 eq). The reaction mixture was stirred for 3 h at 110°C under N_2 protection. The mixture was filtered and the filtrate was concentrated in vacuum. The crude product was purified by prep-TLC (DCM/MeOH = 30/1) to afford the desired product **Example 200** (19.6 mg, 23.9% yield) as a light yellow solid. LCMS $[\text{M}+1]^+ = 428.3$. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 12.36 (s, 1H), 10.90 (s, 1H), 9.71 (s, 1H), 8.96 (s, 1H), 8.67 (s, 1H), 8.14 (d, $J = 5.1$ Hz, 1H), 7.50 (d, $J = 5.1$ Hz, 1H), 3.92 (s, 3H), 3.14 (s, 2H), 2.06-2.08 (m, 1H), 0.83-0.88 (m, 4H).

Example 201:




Step 1: Example 201c

[00732] To a solution of **Example 201a** (1.7 g, 8.21 mmol, 1.0 eq) and **Example 201b** (1.39 g, 8.21 mmol, 1.0 eq) in dioxane (25 mL) were added NaHCO_3 (1.38 g, 16.4 mmol, 2.0 eq), $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (850 mg, 0.82 mmol, 0.1 eq) and BINAP (15.0 mg, 1.64 mmol, 0.2 eq). The reaction mixture was stirred for 16 h at 100°C under N_2 protection. The mixture was filtered and the filtrate was concentrated in vacuum. The crude product was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 1/1) to afford the desired product **Example 201c** (1.01 g, 36.2% yield) as a light brown solid. LCMS $[\text{M}+1]^+ = 341.3$.


Step 2: Example 201

[00733] To a solution of **Example 201c** (78.7 mg, 0.23 mmol, 1.2 eq) and **Example 201d** (40 mg, 0.19 mmol, 1.0 eq) in dioxane (2 mL) were added Cs_2CO_3 (125 mg, 0.38 mmol, 2.0 eq), $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$

(19.9 mg, 0.019 mmol, 0.1 eq) and BINAP (23.9 mg, 0.038 mmol, 0.2 eq). The reaction mixture was stirred for 4 h at 110°C under N₂ protection. The mixture was filtered and the filtrate was concentrated in vacuum. The crude product was purified by prep-TLC (DCM/MeOH = 30/1) to afford the desired product **Example 201** (18.9 mg, 19.3% yield) as a light yellow solid. LCMS [M+1]⁺ = 513.4. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.38 (s, 1H), 10.24 (s, 1H), 9.53 (s, 1H), 8.94 (s, 1H), 8.67 (s, 1H), 8.18 (d, *J* = 5.4 Hz, 1H), 8.00 (dd, *J* = 10.8, 8.4 Hz, 1H), 7.60 (dd, *J* = 8.4, 1.8 Hz, 1H), 7.51 (d, *J* = 5.1 Hz, 1H), 5.29 (s, 1H), 3.93 (s, 3H), 3.11 (s, 2H), 1.50 (s, 6H).

Example 202:**Example 202a****Example 202**

[00734] To a solution of **Example 202b** (750 mg, 2.20 mmol, 1.0 eq) in dioxane (25 mL) were added **Example 202a** (546 mg, 2.64 mmol, 1.2 eq), Cs₂CO₃ (1.43 g, 4.40 mmol, 2.0 eq), BINAP (274 mg, 0.44 mmol, 0.2 eq) and Pd₂(dba)₃CHCl₃ (228 mg, 0.22 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂. The solvent was concentrated; the residue was purified by silica gel flash column chromatography, eluted with DCM/MeOH (v/v = 20/1) to give the desired product **Example 202** (401.3 mg, 35.67% yield) as a light yellow solid. LCMS [M+1]⁺ = 512.4. ¹H NMR (300 MHz, DMSO-*d*₆) δ 11.15 (s, 1H), 10.11 (s, 1H), 8.87 (s, 1H), 8.57 (s, 1H), 7.96 (dd, *J* = 10.8, 8.4 Hz, 1H), 7.79 (s, 1H), 7.73 (dd, *J* = 8.4, 1.5 Hz, 2H), 7.63 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.49 (dd, *J* = 8.1, 1.8 Hz, 1H), 7.28 (t, *J* = 7.8 Hz, 1H), 5.26 (s, 1H), 3.76 (s, 3H), 3.08 (s, 2H), 1.46 (s, 6H).

Example 203:

Step 1: Example 203b

[00735] To a solution of **Example 203a** (50.0 g, 0.446 mol, 1.0 eq) in ACN (1000 mL) was added NBS (87.0 g, 0.491 mol, 1.1 eq) at 0 °C under N₂ protection. The reaction solution was stirred for 1 h at r.t.. The solvent was concentrated under vacuum. The residue was purified by silica gel flash column chromatography (Petroleum Ether/EtOAc = 5/1) to afford the product **Example 203b** (68.1 g, 80.3% yield) as a white solid. LCMS [M+1]⁺ = 191.2.

Step 2: Example 203c

[00736] To a solution of **Example 211b** (10.0 g, 52.63 mmol, 1.0 eq) in THF (200 mL) was added n-BuLi (73.68 mL, 2.5 M in hexane, 184.21 mmol, 3.5 eq) dropwise at -78 °C under N₂. The reaction was stirred for 30 min at the same temperature; A solution of Acetone-*d*₆ (16.84 g, 263.16 mmol, 5.0 eq) in THF (30 mL) was added dropwise at -78 °C. The reaction mixture was stirred for 30 min at r.t. and then quenched with saturated aqueous of NH₄Cl (200 mL) and extracted with EtOAc (300 mL*3). The combined organic layers were washed with brine (500 mL), dried over Na₂SO₄ and concentrated. The residue was purified by silica gel flash column chromatography eluted with Petroleum Ether/EtOAc (1/1) to give the desired product **Example 203c** (5.9 g, 64.0% yield) as a yellow solid. LCMS [M+1]⁺ = 177.2.

Step 3: Example 203e

[00737] To a solution of **Example 203d** (2.0 g, 9.66 mmol, 1.0 eq) in dioxane (50 mL) were added **Example 203c** (1.71 g, 9.66 mol, 1.0 eq), NaHCO₃ (1.62 g, 19.32 mmol, 2.0 eq), BINAP (1.2 g, 1.93 mmol, 0.2 eq) and Pd₂(dba)₃CHCl₃ (1.0 g, 0.97 mmol, 0.1 eq). The reaction mixture was stirred for 16 h at 100°C under N₂. The solvent was concentrated and the residue was purified by silica gel flash column chromatography eluted with Petroleum Ether/EtOAc (1/1) to give the desired product **Example 203e** (960 mg, 28.7% yield) as a yellow solid. LCMS [M+1]⁺ = 347.3.

Step 4: Example 203

[00738] To a solution of **Example 203e** (860 mg, 2.48 mmol, 1.0 eq) in dioxane (40 mL) were added **Example 203f** (512 mg, 2.97 mol, 1.2 eq), Cs₂CO₃ (1.62 g, 4.96 mmol, 2.0 eq), BINAP (309 mg, 0.50 mmol, 0.2 eq) and Pd₂(dba)₃CHCl₃ (257 mg, 0.25 mmol, 0.1 eq). The reaction mixture was stirred for 4 h at 110°C under N₂. The solvent was concentrated and the residue was purified by silica gel flash column chromatography eluted with DCM/MeOH (20/1) to give the desired product **Example 203** (425.0 mg, 35.5% yield) as an off white solid. LCMS [M+1]⁺ = 483.3. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.00 (s, 1H), 10.24 (s, 1H), 9.02 (s, 1H), 8.94 (s, 1H), 8.66 (dd, *J* = 4.8, 1.8 Hz, 1H), 8.29 (dd, *J* = 7.8, 1.8 Hz, 1H), 7.99 (dd, *J* = 10.8, 8.1 Hz, 1H), 7.60 (dd, *J* = 8.4, 2.1 Hz, 1H), 7.35 (dd, *J* = 7.8, 4.8 Hz, 1H), 5.25 (s, 1H), 3.34 (s, 3H), 3.09 (s, 2H).

Example A. TYK2 JH2 Domain Binding Assay

[00739] DiscoverX's KINOMEscan™ is a popular platform for kinase profiling, frequently used in academic and industry witnessed by publications, therefore we selected this platform as a primary cell-free screening assay to determine the relative binding potency and guide chemistry SAR.

[00740] Developed by DiscoverX, KINOMEscan™ employs proprietary active-site dependent competition binding assays to determine how compounds bind to kinases. KINOMEscan™ is based on a

competition binding assay that quantitatively measures the ability of a compound to compete with an immobilized, active site directed ligand. The assay is performed by combining three components: DNA-tagged kinase; immobilized ligand; and a test compound. The ability of the test compound to compete with the immobilized ligand is measured via quantitative PCR of the DNA tag.

[00741] Streptavidin-coated magnetic beads were treated with biotinylated small molecule ligands for 30 minutes at room temperature to generate affinity resins for kinase assays. The liganded beads were blocked with excess biotin and washed with blocking buffer (SeaBlock (Pierce), 1% BSA, 0.05% Tween 20, 1 mM DTT) to remove unbound ligand and to reduce non-specific binding. Binding reactions were assembled by combining TYK2 (JH2domain-pseudokinase), liganded affinity beads, and test compounds in 1x binding buffer (20% SeaBlock, 0.17x PBS, 0.05% Tween 20, 6 mM DTT). Test compounds were prepared as 111X stocks in 100% DMSO. All test compounds were shipped to DiscoverX in DMSO with concentration of 10 mM. Kds were determined using an 11-point 3-fold compound dilution series with three DMSO control points. Each compound was tested in duplicate. All compounds for Kd measurements are distributed by acoustic transfer (non-contact dispensing) in 100% DMSO. The compounds were then diluted directly into the assays such that the final concentration of DMSO was 0.9%. All reactions performed in polypropylene 384-well plate. Each was a final volume of 0.02 ml. The assay plates were incubated at room temperature with shaking for 1 hour and the affinity beads were washed with wash buffer (1x PBS, 0.05% Tween 20). The beads were then re-suspended in elution buffer (1x PBS, 0.05% Tween 20, 0.5 μ M non-biotinylated affinity ligand) and incubated at room temperature with shaking for 30 minutes. The kinase concentration in the eluates was measured by qPCR.

[00742] The amount of kinase measured by qPCR (Signal; y-axis) is plotted against the corresponding compound concentration in nM in log₁₀ scale (x-axis). Binding constants (Kds) were calculated with a standard dose-response curve using the Hill equation:

$$\text{Response} = \text{Background} + \frac{\text{Signal} - \text{Background}}{1 + (\text{Kd}^{\text{Hill Slope}} / \text{Dose}^{\text{Hill Slope}})}$$

[00743] The Hill Slope was set to -1.

[00744] Curves were fitted using a non-linear least square fit with the Levenberg-Marquardt algorithm.

[00745] The results are shown in table 1.

TABLE 1

Ex.	TYK2 (JH2 domain) binding Kd (nM)	Ex.	TYK2 (JH2 domain) binding Kd (nM)	Ex.	TYK2 (JH2 domain) binding Kd (nM)
1	A	42	B	82	A
2	A	43	C	83	A
3	C	44	B	84	A
4	A	45	A	85	A
5	A	46	A	86	A
6	A	47	A	87	A

Ex.	TYK2 (JH2 domain) binding Kd (nM)	Ex.	TYK2 (JH2 domain) binding Kd (nM)	Ex.	TYK2 (JH2 domain) binding Kd (nM)
7	A	48	A	88	A
8	A	49	A	89	A
9	A	50	A	90	A
10	A	51	A	91	A
11	A	52	A	92	A
12	A	53	A	93	A
13	A	54	A	94	A
14	A	55	B	95	A
15	A	56	B	96	A
16	A	57	B	97	A
17	A	58	C	98	A
18	A	59	B	99	A
19	A	60	B	100	A
20	A	61	B	101	A
21	A	62	C	102	A
22	A	63	A	103	A
23	A	64	A	104	C
24	B	65	C	116	A
25	A	66	B	108	A
26	A	67	A	118	D
27	A	68	A	123	D
28	A	69	A	126	A
29	A	70	B	127	A
30	A	71	A	145	A
32	A	72	A	147	A
33	A	73	C	159	A
34	A	74	A	163	A
35	A	75	A	166	A
36	A	76	B	195	A
37	A	77	B	200	A
38	A	78	A	198	A
39	C	79	A	199	A
40	A	80	A	202	A
41	B	81	A	201	A

A is less or equal than 1 nM;

B is more than 1 nM and less or equal than 5 nM;

C is more than 5 nM and less or equal than 10 nM;

D is more than 10 nM.


Example B: IL-12 induced pSTAT4 in human PBMC

[00746] Fresh Human PBMCs were resuspended in RPMI 1640 medium with 10% FBS. Cells were seeded in a round bottom 96-well plate at the concentration of 200,000 cells/well. A 10-point dilution series of test compound (top dose 10 μ M, 1:5 dilution) was added to the well using the liquid dispenser (Tecan D300e) and incubated for 1 hour at 37C. Then human IL-12 recombinant protein (R&D Systems) was added to the well at the final concentration of 10 ng/ml and incubated for 15 minutes at 37C. Cell lysates were prepared and analyzed by Phospho STAT4 (Tyr693) Kit (Meso Scale Discovery) following manufacturer's protocol.

[00747] For calculation of the inhibition rate, the relative pSTAT4 signal of each well = pSTAT4 signal of each well - the average pSTAT4 signal of baseline.

The inhibition% = (the average pSTAT4 signal of IL-12 treatment wells - the relative of pSTAT4 signal in each compound containing well) / the average pSTAT4 signal of IL-12 treatment wells * 100%

[00748] The curve was plotted as the inhibition% (y-axis) vs. compounds concentration (x-axis) and was fitted with log(inhibitor) vs. normalized response -- Variable slope by GraphPad Prism7.0.



Control is BMS-986165:

[00749] The results are shown in table 2.

TABLE 2

Suppression of IL12-induced p-STAT4 in human PBMC			Suppression of IL12-induced p-STAT4 in human PBMC		
Ex.	p-STAT4 IC ₅₀ (nM)	Relative IC ₅₀ to control	Ex.	p-STAT4 IC ₅₀ (nM)	Relative IC ₅₀ to control
1	+	17.4	47	+	10.6
2	+	10.1	48	+	2.6
3	++	146.6	49	+	5.1
4	+	15.6	50	+	21.8
5	+	20.8	51	+	4.1
6	+	4.4	52	+	5.6
7	++	59.8	54	+	3.1
8	+	13.7	55	++	260.4
9	+	22.9	56	++	53.8
10	+	6.4	57	+++	208.9
11	+	3.6	59	+++	188.6
12	+	4.5	60	++++	>196.1
13	+	2.9	61	+++	224.8

Suppression of IL12-induced p-STAT4 in human PBMC			Suppression of IL12-induced p-STAT4 in human PBMC		
Ex.	p-STAT4 IC ₅₀ (nM)	Relative IC ₅₀ to control	Ex.	p-STAT4 IC ₅₀ (nM)	Relative IC ₅₀ to control
14	+	2	62	+++	630.4
15	+	2.7	63	++	202.1
16	+	3.4	64	++	143.1
17	+	7.8	67	+	17.3
18	+	5.4	68	+	10.2
19	+	3	69	+	6.7
20	+	3.6	71	++	28.5
21	+	4.3	72	+	9.1
22	+	3.4	74	+	13.6
23	++	51.9	77	++	52.7
24	++	32.4	78	+	58.9
25	++	32.5	79	++	10.6
26	+	11	80	+	3.7
27	+	3	82	+	4.3
28	+	3.3	84	+	6
29	+	10.2	85	++	613.9
30	+	1.7	86	+	9.7
31	+	4.7	87	+	18.7
32	+	9.3	88	+	6
33	+	10.6	89	+	8
34	+	6.4	91	+	8.3
36	+	44.9	94	+	16.1
37	+	2	95	+	18
39	++	45.4	96	+	18.7
40	++	46.9	98	+	24.2
41	+++	468.6	99	+	9.2
42	++	147.9	100	+	19.7
43	++++	> 3039	103	+	10.8
44	++	77			

+ is less or equal than 100 nM;

++ is more than 100 nM and less or equal than 1 μ M;

+++ is more than 1 μ M and less or equal than 10 μ M;

++++ is more than 10 μ M.

Example C: INF α induced pSTAT3 or pSTAT5 in human PBMC

[00750] Fresh Human PBMCs were resuspended in RPMI 1640 medium with 10% FBS. Cells were seeded in a round bottom 96-well plate at the concentration of 200,000 cells/well. A 10-point dilution series of test compound (top dose 10 μ M, 1:5 dilution) was added to the well using the liquid dispenser (Tecan D300e) and incubated for 1 hour at 37C. Then human INF α recombinant protein (R&D Systems)

was added to the well at the final concentration of 5000 units/ml and incubated for 15 minutes at 37C. Cell lysates were prepared and analyzed by Phospho STAT3 (Tyr705) cellular kit (Cisbio) or Phospho STAT5 (Tyr693) Kit (Meso Scale Discovery) following manufacturer's protocol.

[00751] For calculation of the inhibition rate, the relative pSTAT signal of each well = pSTAT signal of each well - the average pSTAT signal of baseline.

[00752] The inhibition% = (the average pSTAT signal of INF α treatment wells - the relative of pSTAT signal in each compound containing well) / the average pSTAT signal of INF α treatment wells * 100%

[00753] The curve was plotted as the inhibition% (y-axis) vs. compounds concentration (x-axis) and was fitted with log(inhibitor) vs. normalized response -- Variable slope by GraphPad Prism7.0.

[00754] The results are shown in table 3.

TABLE 3

Ex.	p-STAT3 IC ₅₀ (nM)	p-STAT5 IC ₅₀ (nM)	Relative IC ₅₀ to control
1	B		6.4
2	B		2.7
3	C		>61.7
4	B		19.0
5	B		13.9
6		A	10.6
7	B		18.7
8	A		8.0
9	B		21.7
10		A	14.9
11		A	6.3
12	A		7.4
13	A		2.2
14		A	1.9
15		A	5.0
16		A	4.8
23	B		6.6
24	B		14.2
25	B		8.4
26	B		10.5
27		A	1.5
28		A	7.8
37		A	2.2
38		A	4.1
39	B		14.6
40	B		14.2
41	C		575.0
42	C		224.3

Ex.	p-STAT3 IC ₅₀ (nM)	p-STAT5 IC ₅₀ (nM)	Relative IC ₅₀ to control
43	C		135.5
44	C		374.8
49		A	6.5
51		A	12.8
54		A	9.3
55	B		15.5
56	C		89.5
57	C		230.5
59	C		122.2
60	D		>61.7
61	C		113.5
62	C		544.7
63	B		79.9
64	B		91.6
67	B		9.4
68	A		4.5
69	A		8.5
71	B		10.0
72	B		15.7
74	A		7.0
77	C		145.8
79	B		49.7
80	A		3.7
82	A		5.6
84	A		5.2
86	B		17.0
87	B		28.6
89		A	5.4
107		A	5.3
108		A	2.9
109		A	5.5
110		A	1.5
111		A	1.4
112		A	1.8
113		A	2.2
114		A	4.7
115		A	53.8
116		A	2.8
119		A	14.1
120		A	1.1

Ex.	p-STAT3 IC ₅₀ (nM)	p-STAT5 IC ₅₀ (nM)	Relative IC ₅₀ to control
121		A	9.1
122		A	2.7
124		A	3.1
125		A	14.4
126		A	4.8
127		A	10.7
128		C	>1000
129		A	9.6
130		A	9.3
131		A	2.1
132		A	4.3
133		A	18.8
135		C	>1000
136		A	3.3
137		A	6.1
138		A	20.8
139		A	2.3
140		A	0.8
141		A	3.2
142		A	4
143		A	3.8
144		A	24.2
145		A	2.5
147		A	5.8
148		A	3
149		A	2.6
150		A	9.7
151		A	16.3
152		A	2.8
153		A	4.8
154		A	3.0
155		A	3.8
156		A	4.4
157		A	9.5
158		A	7.2
159		A	1.6
160		A	5.5
161		A	6.1
163		A	1.7
164		A	3.9

Ex.	p-STAT3 IC ₅₀ (nM)	p-STAT5 IC ₅₀ (nM)	Relative IC ₅₀ to control
165		B	206
166		A	3.0
169		A	49.3
171		A	1.2
172		A	1.7
173		B	113.7
174		A	0.8
175		A	0.3
176		A	1.8
177		A	0.4
178		A	1.2
179		A	5.8
180		A	1.3
184		A	0.8
185		A	0.3
188		A	0.6
189		A	0.8
190		A	6.3
191		A	5.4
193		A	1.4
195		A	1.6
196		B	784
198		A	0.3
199		A	0.2
200		A	1.9
201		A	0.2
202		A	0.3
203		A	0.7

A is less or equal than 100 nM;

B is more than 100 nM and less or equal than 1 μ M;

C is more than 1 μ M and less or equal than 10 μ M;

D is more than 10 μ M.

Example D: JAK1 JH2 and JAK2 JH1 Domain Binding Assay

[00755] Similar to the method for TYK2 JH2 binding described above, JAK1 JH2 and JAK2 JH1 domain binding assay was performed using DiscoverX's KINOMEscan™, but with change of kinase domain. These assays were performed to compare the binding selectivity of test compounds to JAK1 JH2 and JAK2 JH1 domain. The results are shown in table 4.

TABLE 4

Ex.	JAK1 (JH2 domain) binding Kd (nM)	JAK2 (JH1 domain) binding Kd (nM)
control	A	C
147	A	D
38	A	C
116	A	C
166	A	C
159	A	C
163	A	B
195	A	C
200	A	C
198	A	D
199	A	D
202	A	B
201	A	B

A is less or equal than 100 nM;

B is more than 100 nM and less or equal than 1 μ M;

C is more than 1 μ M and less or equal than 10 μ M;

D is more than 10 μ M.

Example E: GM-CSF-induced pSTAT5 and IL-2-induced pSTAT5 in human PBMC in human PBMC

[00756] Similar to the method for IL-12 induced pSTAT4 in human PBMC described above, these assays were performed to check if test compounds have cross-activity to JAK1, JAK2 and JAK3 pathways in human PBMC. The procedure is as described with change of stimuli to 10 ng/ml of GM-CSF or 20 ng/ml of IL-2. The data are shown in Table 5.

TABLE 5

Ex.	GM-CSF-induced pSTAT5 IC ₅₀ (nM)	IL-2-induced pSTAT5 IC ₅₀ (nM)
control	C	B
195	C	C
200	D	C
198	C	B
199	C	B
202	B	B
201	B	B
203	C	B

A is less or equal than 100 nM;

B is more than 100 nM and less or equal than 1 μ M;

C is more than 1 μ M and less or equal than 10 μ M;

D is more than 10 μ M.

Example F: Pharmacokinetic Studies

[00757] The pharmacokinetics of test compounds were evaluated in male C57BL/6 mice, Sprague Dawley rats, Beagle dogs, and cynomolgus monkeys when administered via oral gavage and IV injection. The formulation for each test compound is summarized in the table 6. The animals were fasted overnight

before administration. Plasma samples were collected predose and at 0.5, 1, 3, 6, 9, 12, and 24 hours postdose. The samples were analyzed by LC/MS/MS and the concentration of test compound at each timepoint was determined by linear regression. Pharmacokinetic parameters were calculated from the plasma concentrations using Phoenix WinNonlin. The PK results were summarized in the tables 7-10.

TABLE 6: Drug formulations used for each test compound in different species

Ex.	Route of administration	Mouse	Rat	Dog	Monkey
control	Oral	A	A	A	A
	IV	B	B	B	B
195	Oral	A	A	A	A
	IV	C	B	B	B
200	Oral	A	A	A	A
	IV	C	B	B	B
198	Oral		A	D	D
	IV		B	C	C
199	Oral		A	D	
	IV		B	C	
202	Oral	D	A	A	
	IV	C	B	B	
201	Oral		A	D	D
	IV		B	C	C
203	Oral				D
	IV				C
Formulation A: 0.5% methylcellulose					
Formulation B: 5% DMSO/5% Solutol/90% saline					
Formulation C: 10%DMSO+10%Solutol+80% (20% SBE- β -CD)					
Formulation D: 10%DMSO+10%HS-15+40% PEG400+40% (30%SBE- β -CD)					

TABLE 7: Pharmacokinetic parameters of test compounds in C57BL/6 mice.

Ex.	Route of administration and dose	C ₀ or C _{max} (ng/mL)	AUC (h·ng·mL ⁻¹)	T _{1/2} (h)	T _{max} (h)	F%	CL(mL·kg ⁻¹ ·min ⁻¹)	Vdss(L/kg)
control	Oral (10 mg/kg)	4699	8871	2.96	0.25	86.7		
	I.V. (2 mg/kg)	3220	2045	2.19			15.6	1.56
195	Oral (10 mg/kg)	4295	16678	2.110	0.500	41.7		
	I.V. (2 mg/kg)	2950	7994	2.46			4.16	0.688
200	Oral (10 mg/kg)	4193	19295	2.010	0.250	85.1		
	I.V. (2 mg/kg)	2391	4475	4.62			7.35	1.5
202	Oral (10 mg/kg)	1899	6395	3.27	0.500	66.6		
	I.V. (2 mg/kg)	3343	1911	0.97			17.3	0.785

TABLE 8: Pharmacokinetic parameters of test compounds in Sprague Dawley rats.

Ex.	Route of administration & dose	C ₀ or C _{max} (ng/mL)	AUC (h·ng·mL ⁻¹)	T _{1/2} (h)	T _{max} (h)	F%	CL(mL·kg ⁻¹ ·min ⁻¹)	Vdss(L/kg)
control	Oral (5 mg/kg)	792±260	3197±600	2.93±1.2	1.0±0.87	32.6±5.4		
	I.V. (1 mg/kg)	1743±276	1966±166	2.42±1.7			8.41±0.7	0.933±0.12
195	Oral (5 mg/kg)	1255±226	7444±1396	2.37±0.32	1.67±0.58	46.7±8.8		
	I.V. (1 mg/kg)	2153±123	3188±603	2.14±0.12			4.96±0.91	0.824±0.11
200	Oral (5 mg/kg)	948±43	3657±684	4.22±3.6	1.00	42.7±8.0		
	I.V. (1 mg/kg)	1033±47	1714±19	1.58±0.21			9.42±0.022	1.24±0.065
198	Oral (5 mg/kg)	261±35	1187±53	2.85±2.8	1.00±0.87	16.9±0.76		
	I.V. (1 mg/kg)	1202±70	1405±29	1.22±0.12			11.8±0.22	1.10±0.065
199	Oral (5 mg/kg)	140±32	650±34	1.66±0.37	1.33±0.58	18.4±1.0		
	I.V. (1 mg/kg)	414±56	705±140	1.74±0.20			23.3±4.9	3.27±0.87
202	Oral (5 mg/kg)	50.1±11	210±60	1.29±0.24	4.00±2.0	8.35±2.4		
	I.V. (1 mg/kg)	1975±101	504±58	0.323±0.035			33.1±4.0	0.653±0.03
201	Oral (5 mg/kg)	185±138	499±502	1.14±0.2	0.42±0.14	18.7±18.9		
	I.V. (1 mg/kg)	1186±116	533±225	0.759±0.60			34.3±12	1.31±0.42

TABLE 9: Pharmacokinetic parameters of test compounds in Beagle dogs

Ex.	Route of administration & dose	C ₀ or C _{max} (ng/mL)	AUC (h·ng·mL ⁻¹)	T _{1/2} (h)	T _{max} (h)	F%	CL(mL·kg ⁻¹ ·min ⁻¹)	Vdss(L/kg)
control	Oral (2 mg/kg)	222±46	1312±579	6.21±2.9	1.67±0.58	35.5±15		
	I.V. (1 mg/kg)	559±34	1902±349	4.43±0.87			8.85±1.7	2.27±0.23
195	Oral (2 mg/kg)	285±285	1517±1334	5.73±2.02	1.33±0.58	26.2±23.0		
	I.V. (1 mg/kg)	185±8.4	2899±453	6.69±0.39			5.31±0.93	3.04±0.27
200	Oral (2 mg/kg)	247±17	2619±258	8.51±1.6	1.33±0	82.9±8.2		
	I.V. (0.5 mg/kg)	29.7±14	790±196	11.5±1.6			8.17±2.6	8.37±1.39
198	Oral (2 mg/kg)	68.8±18	253±48	1.42±0.24	1.67±0.58	54.0±10		
	I.V. (1 mg/kg)	182±47	234±94	4.29±1			68.6±28	23.1±4.59

Ex.	Route of administration & dose	C ₀ or C _{max} (ng/mL)	AUC (h·ng·mL ⁻¹)	T _{1/2} (h)	T _{max} (h)	F%	CL(mL·kg ⁻¹ ·min ⁻¹)	Vdss(L/kg)
	mg/kg)	6		25				
199	Oral (2 mg/kg)	23.4±4	67.6±25	1.36±0.05	1.33±0.6	10.7±3.9		
	I.V. (1 mg/kg)	334±96	315±188	1.78±0.59			63.2±30	5.62±1.13
202	Oral (2 mg/kg)	156±53	555±154	1.59±0.15	1.00	55.0±15		
	I.V. (1 mg/kg)	469±25	504±49	1.74±0.28			32.7±3.1	4.21±0.19
201	Oral (2 mg/kg)	103±28	442±81	5.05±0.47	2.0±0	38.5±7.1		
	I.V. (1 mg/kg)	458±33	575±158	4.54±1.26			29.3±8.01	7.41±2.64

TABLE 10: Pharmacokinetic parameters of test compounds in cynomolgus monkeys (NA: no detectable)

Ex.	Route of administration & dose	C ₀ or C _{max} (ng/mL)	AUC (h·ng·mL ⁻¹)	T _{1/2} (h)	T _{max} (h)	F%	CL(mL·kg ⁻¹ ·min ⁻¹)	Vdss(L/kg)
control	Oral (2 mg/kg)	247±13	3192±16	10.6±3.8	5.33±1.2	27.8±14		
	I.V. (1 mg/kg)	1417±2	5739±11	7.54±1.6			2.79±0.74	1.20±0.04
195	Oral (2 mg/kg)	663±49	5420±32	5.80±0.7	4.0±0	16.6±9.95		
	I.V. (1 mg/kg)	1286±1	16326±6	7.65±0.5			0.898±0.04	0.624±0.03
200	Oral (2 mg/kg)	384±22	4060±25	8.60±0.7	2.0±0	27.7±1.7		
	I.V. (1 mg/kg)	677±81	7316±20	9.35±1.5			1.87±0.21	1.45±0.18
198	Oral (2 mg/kg)	246±12	1228±43	1.28±0.1	4±0	59.4±21.1		
	I.V. (1 mg/kg)	667±82	1034±15	1.45±0.2			16.1±2.4	1.94±0.03
201	Oral (2 mg/kg)	13.9±2	67.9±21	3.08±0.6	2.33±1.5	4.67±1.5	23.2±4.3	1.47±0.1
	I.V. (1 mg/kg)	849±50	726±134	1.81±0.7				
203	Oral (2 mg/kg)	NA	NA	NA	NA	NA		
	I.V. (1 mg/kg)	787±61	595±48	0.951±0.22			27.6±1.9	0.931±0.01

Example E: Pharmaceutical Compositions*Example E1: Parenteral Composition*

[00758] To prepare a parenteral pharmaceutical composition suitable for administration by injection, 100 mg of a water-soluble salt of a compound described herein is dissolved in DMSO and then mixed

with 10 mL of 0.9% sterile saline. The mixture is incorporated into a dosage unit form suitable for administration by injection.

Example E2: Oral Composition

[00759] To prepare a pharmaceutical composition for oral delivery, 100 mg of a compound described herein is mixed with 750 mg of starch. The mixture is incorporated into an oral dosage unit form, such as a hard gelatin capsule, which is suitable for oral administration.


Example E3: Sublingual (Hard Lozenge) Composition

[00760] To prepare a pharmaceutical composition for buccal delivery, such as a hard lozenge, mix 100 mg of a compound described herein, with 420 mg of powdered sugar mixed, with 1.6 mL of light corn syrup, 2.4 mL distilled water, and 0.42 mL mint extract. The mixture is gently blended and poured into a mold to form a lozenge suitable for buccal administration.

[00761] The examples and embodiments described herein are for illustrative purposes only and in some embodiments, various modifications or changes are to be included within the purview of disclosure and scope of the appended claims.

CLAIMS


1. A compound of Formula (XIII):



Formula (XIII)

or a pharmaceutically acceptable salt or stereoisomer thereof:

wherein:



L is a bond;

Ring A is cycloalkyl, aryl, or heteroaryl; wherein the cycloalkyl, aryl, and heteroaryl are each optionally substituted with one or more independently selected R^A substituents;

Ring B is phenyl or pyridinyl;

R^3 is hydrogen, C_1 - C_6 alkyl, C_1 - C_6 deuteroalkyl, or C_1 - C_6 haloalkyl;

R^4 is hydrogen, deuterium, halogen, CN, NO_2 , C_1 - C_6 alkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 hydroxyalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $C(O)R^a$, $C(O)NR^cR^d$, $C(O)OR^b$, NR^cR^d , $NR^bC(O)R^a$, $NR^bC(O)NR^cR^d$, $NR^bC(O)OR^b$, $NHS(O)_2R^a$, OR^b , $OC(O)R^a$, $OC(O)NR^cR^d$, $OC(O)OR^b$, $P(O)R^bR^b$, SR^b , $S(O)R^a$, $S(O)_2R^a$, or $S(O)_2NR^cR^d$;

R^5 is halogen, CN, NO_2 , C_1 - C_6 alkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 hydroxyalkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $C(NCN)R^7$, $C(NCN)NR^9R^{10}$, $C(NCN)OR^8$, $C(O)R^7$, $C(O)NR^9R^{10}$, $C(O)OR^8$, NR^9R^{10} , $NR^8C(NCN)R^7$, $NR^8C(NOH)R^7$, $NR^8C(NCN)NR^9R^{10}$, $NR^8C(O)OR^8$, $NR^8S(O)R^7$, $NR^8C(O)NR^9R^{10}$, $NR^8C(NCN)OR^8$, $NR^8C(O)R^7$, $NR^8S(O)(NR^8)R^7$, $NR^8S(O)_2R^7$, OR^8 , $OC(NCN)R^7$, $OC(NCN)NR^9R^{10}$,

OC(NCN)OR⁸, OC(O)R⁷, OC(O)OR⁸, OC(O)NR⁹R¹⁰, SR⁸, S(O)R⁷, S(O)₂R⁷, S(O)₂NR⁹R¹⁰, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₁-C₆heteroalkyl, C₁-C₆alkyl(cycloalkyl), C₁-C₆alkyl(heterocycloalkyl), C₁-C₆alkyl(aryl), and C₁-C₆alkyl(heteroaryl), C₂-C₆alkenyl, C₂-C₆alkynyl, C(O)R^a, C(O)NR^cR^d, C(O)OR^b, NR^cR^d, OR^b, oxo, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)R^a, C(O)NR^cR^d, C(O)OR^b, NR^cR^d, OR^b, and oxo;

R⁷ is C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

each R⁸ is independently hydrogen, CN, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, and C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

R⁹ is hydrogen, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-

C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

R¹⁰ is hydrogen, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆hydroxyalkyl, C₁-C₆hydroxydeuteroalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

R⁸ and R⁹, taken together with the atoms to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

R⁹ and R¹⁰, taken together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

R¹¹ is C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally substituted with one or more independently selected R^{11a} substituents;

each R^{11a} is independently deuterium, halogen, CN, NO₂, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C(O)R^a, C(O)NR^cR^d, C(O)OR^b, NR^cR^d, NR^bC(O)R^a, NR^bC(O)NR^cR^d, NR^bC(O)OR^b, NHS(O)₂R^a, OR^b, OC(O)R^a, OC(O)NR^cR^d, OC(O)OR^b, SR^b, S(O)R^a, S(O)₂R^a, and S(O)₂NR^cR^d;

each R^A is independently deuterium, halogen, CN, NO₂, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C(O)R^a, C(O)NR^cR^d, C(O)OR^b, NR^cR^d, NR^bC(O)R^a, NR^bC(O)NR^cR^d, NR^bC(O)OR^b, NHS(O)₂R^a, OR^b, OC(O)R^a, OC(O)NR^cR^d, OC(O)OR^b, SR^b, S(O)R^a, S(O)₂R^a, S(O)₂NR^cR^d, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl,

heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

two R^A, taken together with the carbon to which they are attached, form an oxo;

each R^a is independently C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

each R^b is independently hydrogen, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;


each R^c is independently hydrogen, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

each R^d is independently hydrogen, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl; wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo;

R^c and R^d, taken together with the nitrogen atom to which they are attached, form a heterocycloalkyl; wherein the heterocycloalkyl is optionally substituted with one or more

substituents independently selected from the group consisting of deuterium, halogen, CN, C₁-C₆alkyl, C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo; and each R^x is independently hydrogen, deuterium, halogen, CN, NO₂, C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, C₁-C₆aminoalkyl, C₁-C₆hydroxyalkyl, C₂-C₆alkenyl, C₂-C₆alkynyl, C(O)R^a, C(O)NR^cR^d, C(O)OR^b, NR^cR^d, NR^bC(O)R^a, NR^bC(O)NR^cR^d, NR^bC(O)OR^b, NHS(O)₂R^a, OR^b, OC(O)R^a, OC(O)NR^cR^d, OC(O)OR^b, SR^b, S(O)R^a, S(O)₂R^a, and S(O)₂NR^cR^d.

2. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein





3. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein Ring A is a 5-membered heteroaryl, wherein the 5-membered heteroaryl is optionally substituted with one or more independently selected R^A substituents.
4. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein Ring B is phenyl.
5. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein Ring B is pyridinyl.
6. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R³ is hydrogen.
7. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R⁴ is OR^b.


8. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R^5 is $NR^8C(O)R^7$, $NR^8C(O)NR^9R^{10}$, or heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C_1 - C_6 alkyl, C_1 - C_6 deuteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 aminoalkyl, C_1 - C_6 hydroxyalkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 alkyl(cycloalkyl), C_1 - C_6 alkyl(heterocycloalkyl), C_1 - C_6 alkyl(aryl), C_1 - C_6 alkyl(heteroaryl), C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $C(O)R^a$, $C(O)NR^cR^d$, $C(O)OR^b$, NR^cR^d , OR^b , oxo, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, wherein each alkyl, alkenyl, alkynyl, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl is optionally and independently substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, $C(O)R^a$, $C(O)NR^cR^d$, $C(O)OR^b$, NR^cR^d , OR^b , and oxo.
9. The compound of claim 1, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein R^5 is $NR^8C(O)R^7$.
10. The compound of claim 9, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R^7 is cycloalkyl, wherein the cycloalkyl is optionally substituted with one or more substituents independently selected from the group consisting of deuterium, halogen, CN, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, $C(O)CH_3$, $C(O)OH$, $C(O)OCH_3$, NH_2 , OH , OCH_3 , and oxo.
11. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R^5 is $NR^8C(O)NR^9R^{10}$.
12. The compound of claim 11, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R^9 and R^{10} , taken together with the nitrogen atom to which they are attached, form a heterocycloalkyl, wherein the heterocycloalkyl is optionally substituted with one or more substituents independently selected from the group consisting of deuterium,


halogen, CN, C₁-C₆alkyl, and C₁-C₆haloalkyl, C(O)CH₃, C(O)OH, C(O)OCH₃, NH₂, OH, OCH₃, and oxo.


13. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein each R⁸ is independently hydrogen.
14. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R¹¹ is C₁-C₆alkyl, C₁-C₆deuteroalkyl, C₁-C₆haloalkyl, or cycloalkyl.
15. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R¹¹ is C₁-C₆deuteroalkyl.
16. The compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein each R^A is independently deuterium, halogen, C₁-C₆alkyl, C₁-C₆deuteroalkyl, or C₁-C₆haloalkyl.
17. A compound selected from the group consisting of:
























or a pharmaceutically acceptable salt or stereoisomer thereof.


18. A pharmaceutical composition comprising a pharmaceutically acceptable excipient and a therapeutically effective amount of the compound of claim 1, or a pharmaceutically acceptable salt or stereoisomer thereof.

19. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 1, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.
20. The method of claim 19, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.
21. The method of claim 20, wherein the disease or disorder is rheumatoid arthritis.
22. The method of claim 20, wherein the disease is psoriasis.
23. The method of claim 20, wherein the disease or disorder is systemic lupus erythematosus.
24. The method of claim 20, wherein the disease or disorder is ulcerative colitis.
25. The method of claim 20, wherein the disease or disorder is psoriatic arthritis.
26. The method of claim 20, wherein the disease or disorder is systemic sclerosis.
27. A compound having the structure:



or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

28. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 27, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.
29. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 27, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.
30. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

31. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 30, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

32. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 30, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

33. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

34. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 33, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

35. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 33, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

36. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

37. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 36, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

38. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 36, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

39. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

40. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 39, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

41. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 39, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

42. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

43. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 42, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

44. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 42, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

45. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

46. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 45, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

47. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 45, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

48. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

49. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 48, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

50. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 48, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

51. A compound having the structure:




or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

52. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 51, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

53. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 51, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

54. A compound having the structure:



or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof.

55. A pharmaceutical composition comprising a therapeutically effective amount of the compound of claim 54, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, and a pharmaceutically acceptable excipient.

56. A method for inhibiting tyrosine kinase 2 activity in a patient, comprising administering to a patient in need thereof a therapeutically effective amount of the compound of claim 54, or a pharmaceutically acceptable salt, stereoisomer, or solvate thereof, wherein inhibiting tyrosine kinase 2 activity is associated with treating a disease or disorder selected from Crohn's disease, rheumatoid arthritis, psoriasis, systemic lupus erythematosus, ulcerative colitis, psoriatic arthritis, and systemic sclerosis.

57. The method of claim 20, wherein the disease is Crohn's disease.