

(19) AUSTRALIAN PATENT OFFICE

(54) Title
HFA-suspension formulations containing an anticholinergic

(51)⁶ International Patent Classification(s)
A61K 9/12 (2006.01) **OBMJP** **A61K**
A61K 9/00 (2006.01) 47/12
A61K 31/46 (2006.01) 20060101ALI2006031
A61K 47/02 (2006.01) **OBMJP** **A61K**
A61K 47/10 (2006.01) 47/14
A61K 47/12 (2006.01) 20060101ALI2006031
A61K 47/14 (2006.01) **OBMJP** **A61K**
A61K 47/18 (2006.01) 47/18
A61K 47/22 (2006.01) 20060101ALI2006031
A61K 47/34 (2006.01) **OBMJP** **A61K**
A61K 47/42 (2006.01) 47/22
A61K 47/44 (2006.01) 20060101ALI2006031
A61K 47/44 (2006.01) **OBMJP** **A61K**
A61P 11/00 (2006.01) 47/34
A61P 11/06 (2006.01) 20060101ALI2006031
A61K 9/12 **OBMJP** **A61K**
20060101AFI2006031 47/42
OBMJP **A61K** 20060101ALI2006031
9/00 **OBMJP** **A61K**
20060101ALI2007072 47/44
1BMFP **A61K** 20060101ALI2006031
31/46 **OBMJP** **A61P**
20060101ALI2005100 11/00
8BMFP **A61K** 20060101ALI2006031
47/02 **OBMJP** **A61P**
20060101ALI2006031 11/06
OBMJP **A61K** 20060101ALI2006031
47/10 **OBMJP**
20060101ALI2006031 PCT/EP03/02898

(21) Application No: 2003209743

(22) Application Date: 2003 .03 .20

(87) WIPO No: W003/082252

(30) Priority Data

(31) Number	(32) Date	(33) Country
102 14 263.7	2002 .03 .28	DE

(43) Publication Date : 2003 .10 .13

(43) Publication Journal Date : 2003 .11 .20

(71) Applicant(s)
Boehringer Ingelheim Pharma GmbH & Co. KG

(72) Inventor(s)
Schmelzer, Christel

(74) Agent/Attorney
Davies Collison Cave, 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art
WO 2002/036106
WO 2002/038154

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
9. Oktober 2003 (09.10.2003)

PCT

(10) Internationale Veröffentlichungsnummer
WO 03/082252 A1

- (51) Internationale Patentklassifikation: **A61K 9/72**, 31/46
- (21) Internationales Aktenzeichen: PCT/EP03/02898
- (22) Internationales Anmeldedatum: 20. März 2003 (20.03.2003)
- (25) Einreichungssprache: Deutsch
- (26) Veröffentlichungssprache: Deutsch
- (30) Angaben zur Priorität: 102 14 263 7 28. März 2002 (28.03.2002) DE
- (71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): **BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG** [DE/DE]; Binger Strasse 173, 55216 INGELHEIM (DE).
- (72) Erfinder; und
- (75) Erfinder/Anmelder (nur für US): **SCHMELZER, Christel** [DE/DE]; Wellenstrasse 14, 55218 INGELHEIM AM RHEIN (DE).
- (74) Gemeinsamer Vertreter: **BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG**; Binger Strasse 173, 55216 INGELHEIM (DE).
- (81) Bestimmungsstaaten (national): AF, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CI, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Bestimmungsstaaten (regional): ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI-Patent (BF, BJ, CI, CG, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Veröffentlicht:
mit internationalem Recherchenbericht
- Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

WO 03/082252 A1

(54) Title: HFA-SUSPENSION FORMULATIONS CONTAINING AN ANTICHLINERGIC

(54) Bezeichnung: HFA-SUSPENSIONSFORMULIERUNGEN ENTHALTEND EIN ANTICHLINERGIKUM

(57) Abstract: The invention relates to a propellant gas formulation containing suspensions of a crystalline monohydrate of (1 α ,2 β ,4 β ,5 α ,7 β)-7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^{2,4}]nonane-bromide.

(57) Zusammenfassung: Die Erfindung betrifft Treibgasformulierungen enthaltend Suspensionen des kristallinen Monohydrats von (1 α ,2 β ,4 β ,5 α ,7 β)-7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^{2,4}]nonan-bromid.

80281pct.210


HFA suspension formulations containing an anticholinergic

- 5 The invention relates to pressurised gas preparations for metered-dose aerosols with suspension formulations of the crystalline monohydrate of (1 α ,2 β ,4 β ,5 α ,7 β)-7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^{2,4}]nonane-bromide, processes for the preparation thereof and the use thereof for preparing a pharmaceutical composition, particularly for preparing
- 10 a pharmaceutical composition with an anticholinergic activity.

Background to the invention

The compound (1 α ,2 β ,4 β ,5 α ,7 β)-7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.0^{2,4}]nonane-bromide, is known from European Patent

- 15 Application EP 418 716 A1 and has the following chemical structure:



(I)

- The compound has valuable pharmacological properties and is known by the name tiotropium bromide (BA679). Tiotropium bromide is a highly effective anticholinergic
- 20 and can therefore provide therapeutic benefit in the treatment of asthma or COPD (chronic obstructive pulmonary disease).

Tiotropium bromide is preferably administered by inhalation.

- 25 The aim of the present invention is to prepare HFA-metered-dose aerosols containing tiotropium bromide as the sole active ingredient in suspended form.

Detailed description of the invention

- It has been found that, depending on the choice of conditions which can be used
- 30 when purifying the crude product obtained after industrial manufacture, tiotropium bromide occurs in various crystalline modifications.

2003209743 19 Feb 2008

2

It has been found that these different modifications can be deliberately produced by selecting the solvents used for the crystallisation as well as by a suitable choice of the process conditions used in the crystallisation process. For the purposes of
 5 preparing the formulations according to the invention, crystalline tiotropium bromide monohydrate has proved particularly suitable.

Accordingly, the present invention relates to suspensions of crystalline tiotropium bromide monohydrate in the propellant gases HFA 227 and/or HFA 134a, optionally
 10 in admixture with one or more other propellant gases, preferably selected from the group consisting of propane, butane, pentane, dimethylether, CHClF_2 , CH_2F_2 , CF_3CH_3 , isobutane, isopentane and neopentane, and wherein tiotropium bromide is present as the only active substance.

- 15 Preferred suspensions according to the invention are those which contain as propellant gas HFA 227 on its own, a mixture of HFA 227 and HFA 134a or HFA 134a on its own. If a mixture of propellant gases HFA 227 and HFA 134a is used in the suspension formulations according to the invention, the weight ratios in which these two propellant gas components are used may be freely selected.
- 20 If in the suspension formulations according to the invention one or more other propellant gases are used in addition to the propellant gases HFA 227 and/or HFA 134a, selected from the group consisting of propane, butane, pentane, dimethylether, CHClF_2 , CH_2F_2 , CF_3CH_3 , isobutane, isopentane and neopentane, the proportion of this other propellant gas component is preferably less than 50 %, preferably less than 40%, more preferably less than 30%.

The suspensions according to the invention preferably contain between 0.001 and 0.8% tiotropium. Suspensions which contain 0.08 to 0.5%, more preferably 0.2 to 0.4% tiotropium are preferred according to the invention.

- 30 By tiotropium is meant the free ammonium cation. The propellant gas suspensions according to the invention are characterised in that they contain tiotropium in the form of the crystalline tiotropium bromide monohydrate which is exceptionally suitable for this application. Accordingly, the present invention preferably relates to suspensions which contain between 0.0012 and 1% crystalline tiotropium bromide monohydrate. Of particular interest according to the invention are suspensions which
 35 contain 0.1 to 0.62%, more preferably 0.25 to 0.5% crystalline tiotropium bromide monohydrate.

The percentages specified within the scope of the present invention are always percent by mass. If parts by mass of tiotropium are given in percent by mass, the corresponding values for the crystalline tiotropium bromide monohydrate which is preferably used within the scope of the present invention may be obtained by multiplying by a conversion factor of 1.2495.

In some cases within the scope of the present invention the term suspension formulation may be used instead of the term suspension. The two terms are to be regarded as interchangeable within the scope of the present invention.

10

The propellant-containing inhalation aerosols or suspension formulations according to the invention may also contain other ingredients such as surface-active agents (surfactants), adjuvants, antioxidants or flavourings.

15 The surface-active agents (surfactants) which may be contained in the suspensions according to the invention are preferably selected from among Polysorbate 20, Polysorbate 80, Myvacet 9-45, Myvacet 9-08, isopropylmyristate, oleic acid, propyleneglycol, polyethyleneglycol, Brij, ethyloleate, glyceryl trioleate, glyceryl monolaurate, glyceryl monooleate, glyceryl monostearate, glyceryl monoricinoleate, cetylalcohol, sterylalcohol, cetylpyridinium chloride, block polymers, natural oil, ethanol and isopropanol. Of the abovementioned suspension adjuvants Polysorbate 20, Polysorbate 80, Myvacet 9-45, Myvacet 9-08 or isopropylmyristate are preferably used. Myvacet 9-45 or isopropylmyristate are particularly preferred.

20 Where the suspensions according to the invention contain surfactants, these are preferably present in an amount of 0.0005 - 1 %, more preferably 0.005 - 0.5 %.

The adjuvants optionally contained in the suspensions according to the invention are preferably selected from among alanine, albumin, ascorbic acid, aspartame, betaine, cysteine, phosphoric acid, nitric acid, hydrochloric acid, sulphuric acid and citric acid.

30 Of these, ascorbic acid, phosphoric acid, hydrochloric acid or citric acid are preferred, while hydrochloric acid or citric acid is more preferable.

Where the suspensions according to the invention contain adjuvants, these are preferably present in an amount of 0.0001-1.0 %, preferably 0.0005-0.1 %, more preferably 0.001-0.01 %, while an amount of from 0.001-0.005 % is particularly preferred according to the invention.

35 The antioxidants optionally contained in the suspensions according to the invention are preferably selected from among ascorbic acid, citric acid, sodium edetate, editic

acid, tocopherols, butylhydroxytoluene, butylhydroxyanisol and ascorbyl palmitate, of which tocopherols, butylhydroxytoluene, butylhydroxyanisol and ascorbyl palmitate are preferred.

- 5 The flavourings which may be contained in the suspensions according to the invention are preferably selected from among peppermint, saccharine, Dentomint®, aspartame and ethereal oils (e.g. cinnamon, aniseed, menthol, camphor), of which peppermint or Dentomint® is particularly preferred.
- 10 For administration by inhalation it is necessary to prepare the active substance in finely divided form. The crystalline tiotropium bromide monohydrate which may be obtained as detailed in the experimental section is either ground (micronised or obtained in finely divided form by other technical methods known in principle in the art (such as precipitation and spray drying). Methods of micronising active
- 15 substances are known in the art. Preferably, after micronisation, the active substance has an average particle size of 0.5 to 10 µm, preferably 1 to 6 µm, more preferably 1.5 to 5 µm. Preferably, at least 50%, more preferably at least 60%, most preferably at least 70% of the particles of active substance have a particle size which is within the ranges specified above. More preferably, at least 80%, most preferably
- 20 at least 90% of the particles of active substance have a particle size within the ranges specified above.

Surprisingly, it has been found that it is also possible to prepare suspensions which contain, apart from the abovementioned propellant gases, only the active substance

25 and no other additives. Accordingly, in another aspect, the present invention relates to suspensions which contain only the active substance and no other additives.

The suspensions according to the invention may be prepared by methods known in the art. For this the ingredients of the formulation are mixed with the propellant gas

30 or gases (optionally at low temperatures) and transferred into suitable containers.

The propellant gas-containing suspensions according to the invention mentioned above may be administered using inhalers known in the art (pMDIs = pressurised metered dose inhalers). Accordingly, in another aspect, the present invention relates

35 to pharmaceutical compositions in the form of suspensions as hereinbefore described combined with one or more inhalers suitable for administering these suspensions. In addition, the present invention relates to inhalers which are characterised in that they contain the propellant gas-containing suspensions described above according to the invention. The present invention also relates to

containers (e.g. cartridges) which are fitted with a suitable valve and can be used in a suitable inhaler and which contain one of the above-mentioned propellant gas-containing suspensions according to the invention. Suitable containers (e.g. cartridges) and methods of filling these cartridges with the propellant gas-containing suspensions according to the invention are known from the prior art.

In view of the pharmaceutical activity of tiotropium the present invention further relates to the use of the suspensions according to the invention for preparing a drug for administration by inhalation or by nasal route, preferably for preparing a drug for the treatment by inhalation or by nasal route of diseases in which anticholinergics may provide a therapeutic benefit.

Most preferably, the invention further relates to the use of the suspensions according to the invention for preparing a pharmaceutical composition for the treatment by inhalation of respiratory complaints, preferably asthma or COPD.

The Examples that follow serve to illustrate the present invention more fully by way of example, without restricting it to their content.

Starting materials

20

Crystalline tiotropium bromide monohydrate:

The tiotropium obtained according to EP 418 716 A1 may be used to prepare the crystalline tiotropium bromide monohydrate. This is then reacted as described below.

25 15.0 kg of tiotropium bromide are added to 25.7 kg of water in a suitable reaction vessel. The mixture is heated to 80-90°C and stirred at constant temperature until a clear solution is formed. Activated charcoal (0.8 kg), moistened with water, is suspended in 4.4 kg of water, this mixture is added to the solution containing tiotropium bromide and rinsed with 4.3 kg of water. The mixture thus obtained is
30 stirred for at least 15 min. at 80-90°C and then filtered through a heated filter into an apparatus which has been preheated to an outer temperature of 70°C. The filter is rinsed with 8.6 kg of water. The contents of the apparatus are cooled to a temperature of 20-25°C at a rate of 3-5°C every 20 minutes. Using cold water the apparatus is cooled further to 10-15°C and crystallisation is completed by stirring for
35 at least another hour. The crystals are isolated using a suction filter drier, the crystal slurry isolated is washed with 9 L of cold water (10-15°C) and cold acetone (10-15°C). The crystals obtained are dried at 25°C for 2 hours in a nitrogen current.
Yield: 13.4 kg of tiotropium bromide monohydrate (86 % of theory).

The tiotropium bromide monohydrate obtainable using the method described above was investigated by DSC (Differential Scanning Calorimetry). The DSC diagram shows two characteristic signals. The first, relatively broad, endothermic signal between 50-120°C can be attributed to the dehydration of the tiotropium bromide monohydrate into the anhydrous form. The second, relatively sharp, endothermic peak at $230 \pm 5^\circ\text{C}$ can be put down to the melting of the substance. This data was obtained using a Mettler DSC 821 and evaluated using the Mettler STAR software package. The data was recorded at a heating rate of 10 K/min.

The crystalline tiotropium bromide monohydrate was characterised by IR spectroscopy. The data was obtained using a Nicolet FTIR spectrometer and evaluated with the Nicolet OMNIC software package, version 3.1. The measurement was carried out with 2.5 μmol of tiotropium bromide monohydrate in 300 mg of KBr. The following Table shows some of the essential bands of the IR spectrum.

Wave number (cm^{-1})	Attribution	Type of oscillation
3570, 3410	O-H	elongated oscillation
3105	Aryl C-H	elongated oscillation
1730	C=O	elongated oscillation
1260	Epoxide C-O	elongated oscillation
1035	Ester C-OC	elongated oscillation
720	Thiophene	cyclic oscillation

The monocrystal X-ray structural analysis carried out showed that the crystalline tiotropium bromide monohydrate obtainable by the above process has a simple monoclinic cell with the following dimensions:

$a = 18.0774 \text{ \AA}$, $b = 11.9711 \text{ \AA}$, $c = 9.9321 \text{ \AA}$, $\beta = 102.691^\circ$, $V = 2096.96 \text{ \AA}^3$.

These data were obtained using an AFC7R 4-circuit diffractometer (Rigaku) using monochromatic copper K_α radiation. The structural resolution and refinement of the crystal structure were obtained by direct methods (SHELXS86 Program) and FMLQ-refinement (TeXsan Program).

To prepare the suspensions according to the invention the crystalline tiotropium bromide monohydrate obtainable by the above process is micronised by methods

known *per se* in the art, to prepare the active substance in the form of the average particle size which corresponds to the specifications according to the invention.

- 5 A method of determining the average particle size of the active substance will now be described.

Determining the particle size of micronised tiotropium bromide monohydrate:

Measuring equipment and settings:

- 10 The equipment is operated according to the manufacturer's instructions.

	Measuring equipment:	HELOS Laser diffraction spectrometer, SympaTec
	Dispersing unit:	RODOS dry disperser with suction funnel, SympaTec
15	Sample quantity:	50 mg - 400 mg
	Product feed:	Vibri Vibrating channel, Messrs. Sympatec
	Frequency of vibrating channel:	40 rising to 100 %
	Duration of sample feed:	15 to 25 sec. (in the case of 200 mg)
	Focal length:	100 mm (measuring range: 0.9 - 175 μ m)
20	Measuring time:	about 15 s (in the case of 200 mg)
	Cycle time:	20 ms
	Start/stop at:	1 % on channel 28
	Dispersing gas:	compressed air
	Pressure:	3 bar
25	Vacuum:	maximum
	Evaluation method:	HRLD

Sample preparation /product feed:

About 200 mg of the test substance are weighed onto a piece of card.

- 30 Using another piece of card all the larger lumps are broken up. The powder is then sprinkled finely over the front half of the vibrating channel (starting about 1 cm from the front edge). After the start of the measurement the frequency of the vibrating channel is varied from about 40 % up to 100 % (towards the end of the measurement). The sample should be fed in as continuously as possible. However,
- 35 the quantity of product should not be too great, so as to ensure adequate dispersal. The time taken to feed in the entire 200 mg sample is about 15 to 25 sec., for example.

Examples of formulations

Suspensions containing other ingredients in addition to active substance and propellant gas:

- a) 0.02 % Tiotropium*
0.20 % Polysorbate 20
99.78 % HFA 227
- b) 0.02 % Tiotropium*
1.00 % Isopropylmyristate
98.98 % HFA 227
- c) 0.02 % Tiotropium*
0.3 % Myvacet 9-45
99.68 % HFA 227
- d) 0.04 % Tiotropium*
1.00 % Myvacet 9-08
98.96 % HFA 227
- e) 0.04 % Tiotropium*
0.04 % Polysorbate 80
99.92 % HFA 227
- f) 0.04 % Tiotropium*
0.005 % Oleic acid
99.955 % HFA 227
- g) 0.02 % Tiotropium*
0.1 % Myvacet 9-45
60.00 % HFA 227
39.88 % HFA 134a
- h) 0.02 % Tiotropium*
0.30 % Isopropylmyristate
20.00 % HFA 227
79.68 % HFA 134a
- i) 0.02 % Tiotropium*
0.01 % Oleic acid
60.00 % HFA 227
39.97 % HFA 134a

5

*used in the form of the tiotropium bromide monohydrate
(conversion factor 1.2495)

2003209743 11 Feb 2005

- 9 -

Suspensions containing only active substance and propellant gas:

- j) 0.02 % Tiotropium*
99.98 % HFA 227
- k) 0.02 % Tiotropium*
99.98 % HFA 134a
- l) 0.04 % Tiotropium*
99.96 % HFA 227
- m) 0.04 % Tiotropium*
99.96 % HFA 134a
- n) 0.02 % Tiotropium*
20.00 % HFA 227
79.98 % HFA 134a
- o) 0.02 % Tiotropium*
60.00 % HFA 227
39.98 % HFA 134a
- p) 0.04 % Tiotropium*
40.00 % HFA 227
59.96 % HFA 134a
- q) 0.04 % Tiotropium*
80.00 % HFA 227
19.96 % HFA 134a

* used in the form of the tiotropium bromide monohydrate (conversion factor 1.2495).

5

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers

10 or steps.

The reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that that prior art forms part of the common general knowledge in Australia.

2003209743 19 Feb 2008

- 10 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

- 1) Suspensions of crystalline tiotropium bromide monohydrate in the propellant gases HFA 227 and/or HFA 134a, optionally in admixture with one or more other
5 propellant gases selected from the group consisting of propane, butane, pentane, dimethylether, CHClF_2 , CH_2F_2 , CF_3CH_3 , isobutane, isopentane and neopentane and wherein tiotropium bromide is present as the only active substance.
- 2) Suspensions according to claim 1, characterised in that they contain between
10 0.001 and 0.8% tiotropium.
- 3) Suspensions according to claim 1 or 2, characterised in that they contain as other ingredients surface-active agents (surfactants), adjuvants, antioxidants and/or flavourings.
15
- 4) Suspensions according to claim 3, characterised in that they contain as surface-active agents (surfactants) one or more compounds selected from the group consisting of Polysorbate 20, Polysorbate 80, Myvacet 9-45, Myvacet 9-08, isopropylmyristate, oleic acid, propyleneglycol, polyethyleneglycol, Brij, ethyl
20 oleate, glyceryl trioleate, glyceryl monolaurate, glyceryl monooleate, glyceryl monostearate, glyceryl monoricinoleate, cetylalcohol, sterylalcohol, cetylpyridinium chloride, block polymers, natural oil, ethanol and isopropanol.
- 5) Suspensions according to claim 3, characterised in that they contain as
25 adjuvants one or more compounds selected from the group consisting of alanine, albumin, ascorbic acid, aspartame, betaine, cysteine, phosphoric acid, nitric acid, hydrochloric acid, sulphuric acid and citric acid.
- 6) Suspensions according to claim 3, characterised in that they contain as
30 antioxidants one or more compounds selected from the group consisting of ascorbic acid, citric acid, sodium edetate, editic acid, tocopherols, butylhydroxytoluene, butylhydroxyanisol and ascorbyl palmitate.
- 7) Suspensions according to claim 1 or 2, characterised in that they contain no
35 other ingredients apart from the active substance and the propellant gas or gases.
- 8) Use of a suspension according to one of claims 1 to 7 for preparing a pharmaceutical composition, preferably for preparing a pharmaceutical

2003209743 19 Feb 2008

- 11 -

composition for the treatment by inhalation or nasal route of diseases in which anticholinergics may confer a therapeutic benefit.

- 5 9) Use according to claim 8, characterised in that the diseases are respiratory complaints, preferably asthma or COPD.
- 10 10) A method of treatment of a disease in which anticholinergics may confer a therapeutic benefit, said method comprising the administration of a pharmaceutically effective amount of the suspension according to one of claims 1 to 7.
- 11) The method of claim 10 wherein said administration is by inhalation or the nasal route.
- 15 12) The method of claim 10 or claim 11 wherein said diseases are respiratory diseases.
- 20 13) The method of claim 12 wherein said respiratory diseases are asthma or COPD.
- 25 14) The suspension as claimed in any one of claims 1 to 7 substantially as described herein with reference to the Figures and /or Examples.
- 15) The use as claimed in claim 8 or claim 9 substantially as described herein with reference to the Figures and /or Examples.
- 16) The method as claimed in any one of claims 10 to 13 substantially as described herein with reference to the Figures and /or Examples.