

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 March 2012 (01.03.2012)

PCT

(10) International Publication Number
WO 2012/025123 A1

(51) International Patent Classification:
A61K 31/495 (2006.01) *A61P 25/00* (2006.01)

(21) International Application Number:
PCT/DK2011/050317

(22) International Filing Date:
22 August 2011 (22.08.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/375,885 23 August 2010 (23.08.2010) US
PA201000739 23 August 2010 (23.08.2010) DK

(71) Applicant (for all designated States except US): **H. LUNDBECK A/S** [DK/DK]; Ottiliavej 9, DK-2500 Valby (DK).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DRAGHEIM, Marianne** [DK/DK]; Troldager 2, DK-2950 Vedbæk (DK). **FLOREA, Ioana** [RO/DK]; Engløkken 16, DK-2791 Dragør (DK).

(74) Common Representative: **H. LUNDBECK A/S**; Corporate Patents and Trademarks, Ottiliavej 9, DK-2500 Valby (DK).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2012/025123 A1

(54) Title: THERAPEUTIC USES OF 1-[2-(2,4-DIMETHYL-PHENYLSULFANYL)PHENYL]PIPERAZINE

(57) Abstract: The invention provides new therapeutic uses of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]- piperazine and pharmaceutically acceptable salts thereof.

Therapeutic uses of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine

Field of the invention

The present invention relates to novel uses of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]-
5 piperazine and pharmaceutically acceptable salts thereof in the treatment of CNS diseases.

Background of the invention

1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine is a multimodal antidepressant that is thought to work through a combination of two pharmacological modes of action: reuptake inhibition and receptor activity. In vitro studies indicate that 1-[2-(2,4-dimethyl-phenylsulfanyl)-
10 phenyl]piperazine is a 5-HT₃ and 5-HT₇ receptor antagonist, 5-HT_{1B} receptor partial agonist, 5-HT_{1A} receptor agonist and inhibitor of the 5-HT transporter. *In vivo* nonclinical studies have demonstrated that 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine enhances levels of the neurotransmitters serotonin, noradrenalin, dopamine, acetylcholine and histamine in specific areas of the brain. All of these activities are considered to be of clinical relevance and potentially
15 involved in the mechanism of action of the compound.

1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and its use in the treatment of central nervous system (CNS) diseases were first disclosed in the International patent application published as WO 02/029232. Later patent applications (WO 2007/144005; WO 2008/113359; WO 2009/062517) have disclosed crystalline salts of the compound, manufacturing processes,
20 and further therapeutic uses of the compound. Poster P.2.c.040 presented at the 22nd Congress of the European College of Neuropsychopharmacology, 12-16 September 2009, Istanbul, Turkey and poster NR4-024 presented at the 162nd Annual Meeting of the American Psychiatric Association, 16-21 May 2009, San Francisco, USA disclose the results from a proof-of-concept study to evaluate the efficacy and tolerability of the compound in patients with major depressive
25 disorder (MDD). The results from this six weeks, randomised, placebo-controlled study with approximately 100 patients in each arm show that 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]-piperazine separates significantly from placebo in the treatment of depressive and anxious symptoms in patients with MDD. The posters also report that no clinically relevant changes were seen in the clinical laboratory results, vital signs, weight, or ECG parameters. A full paper

reporting the above mentioned proof-of-concept study authored by Alvares *et al* was made available on-line by *Int. J. Neuropsychopharm.* 18 July 2011.

Weight gain is a commonly observed adverse event for treatment with psychotropic drugs. The most pronounced effect seen is probably the weight gain associated with antipsychotics, such as clozapine and olanzapine, but clinically relevant weight gain is also observed for antidepressants [*Exp. Opin. Pharmacother.*, 1, 377-389, 2000].

J.Clin.Psychopharm., 8, 323-330, 1988 reports that treatment with the tricyclic antidepressants (TCA) amitriptyline, nortriptyline, imipramine and desimipramine gave rise to a dose-dependent weight gain of 0.57-1.37 kg/months. Similarly, *J. Affective. Disord.* 7, 133-138, 1984 reports that patients in treatment with the TCA's amitriptyline, nortriptyline or imipramine experienced a weight gain of 1.3-2.9 lbs/month (0.58- 1.31 kg/months). In fact, one half of the patients dropped out of the study due to excessive weight gain.

J.Clin.Psychiatry, 65, 1365-1371, 2004 reports on a 2.5 years study comparing weight gain in patients with obsessive-compulsive disorder (OCD) being treated with various antidepressants. In this study (as in many others), an increase in weight is considered potentially clinically significant (PCS) if it is 7% or above compared to baseline level. At the end of the study period the following percentages of the patients had a PCS weight increase: clomipramine (34.8%), citalopram (14.3%), fluoxetine (8.7%), fluvoxamine (10.7%), paroxetine (14.3%) and sertraline (4.5%).

Int. Clin. Psychopharm., 13, 63-73, 1998 reports on a relapse prevention study comparing the long-term (up to two years) effects of the antidepressants mirtazapine and amitriptyline. The proportion of patients showing a PCS increase in body weight was 22% with amitriptyline, 12.7% with mirtazapine and 3.6% with placebo.

J.Clin.Psychiatry, 61, 863-867, 2000 reports on a long-term study (26-32 weeks) comparing the weight effects of the antidepressants fluoxetine, sertraline and paroxetine. The proportion of patients showing a PCS increase in body weight was 25.5% with paroxetine, 4.2% with sertraline and 6.6% with fluoxetine. These observations are supported by anecdotal reports [*J.Clin.Psychiatry* 60 (suppl 21), 16-19, 1999] that 25-33% of patients on maintenance treatment (i.e. long-term) with serotonin re-uptake inhibitors (SSRI) gain a substantial amount of weight and that paroxetine may be more likely than other drugs in this class to induce weight gain.

On the other hand, a number of short-term studies with paroxetine report an absence of weight gain or even weight loss. *Acta Psychiatr. Scand.* 80 (suppl 350), 117-123, 1989 reports

that no weight gain was noted on a six week study with paroxetine. *Am. J. Psychiatry*, 160, 749-756, 2003 reports a small weight loss in a 566 patients, eight weeks study with paroxetine. *J Clin. Psychiatry*, 62, 350-536, 2001 that no substantial effects on weight was observed in a 324 patients, eight weeks trial with paroxetine.

5 *Am.J. Psychiatry*, 156, 1170-1176, 1999 reports on a long-term (one year) study on the body weight effects of fluoxetine. While fluoxetine in the acute phase induce a slight weight decrease, the long-term effect on weight is similar to placebo.

Finally, *Clin. Therapeutics*, 24, 662-672, 2002 reports on the long-term (52 weeks) body weight effects of bupropion. Bupropion gives rise to a modest, but significant decrease in body weight.

10 The above literature references show that it is difficult to predict the effects on weight associated with treatment with antidepressants. Many antidepressants do have weight gain as an adverse event, others don't. Also, it appears that it is difficult to predict the weight effects associated with long-term treatment with an antidepressant from results reported from short-term studies. This unpredictability is likely to be explained by at least two factors. Firstly, notwithstanding the severity of weight gain related to treatment with antidepressants, it is normally of a smaller magnitude than for treatment with antipsychotics. Thus, any changes in weight may take longer time to develop and become evident. Secondly, weight loss or gain may in itself be a symptom of depression. Therefore, treatment emergent weight gain (i.e. weight gain associated with the compound as such) may be confounded with weight changes resulting from the treatment of the disease; hence it may take longer time to separate the two effects. These observations seem to be supported by a recently published meta-analysis of 116 clinical studies on the effect of antidepressants on body weight [*J.lin.Psych.* 71, 1259-1272, 2010].

20 Given the important implications treatment emergent weight gain may have on patients, it is important to be able to provide treatments with antidepressants associated with no or only few weight related adverse events.

Summary of the invention

25 The present inventors have surprisingly found that long-term treatment with 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof (Compound I) is not associated with weight increase. Thus, in one embodiment, the invention relates to Compound I for use in the treatment of a CNS disease in a patient who has previously

received medication (or is still receiving it) for the treatment of said disease which medication was ceased or reduced (or has to be ceased or reduced) due to weight related adverse events.

In one embodiment, the invention provides Compound I for use in the treatment of a CNS disease in a patient, which patient is overweight.

5 In one embodiment, the invention provides Compound I for use in the treatment of a CNS disease in a patient, which patient suffers from a further disease wherein weight increase should be avoided.

10 In one embodiment, the invention provides a method for the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased or reduced (or has to be ceased or reduced) due to weight related adverse events, the method comprising the administration of a therapeutically effective amount of Compound I to a patient in need thereof.

15 In one embodiment, the invention relates to a method of treating a CNS disease in a patient who is overweight, which method comprises the administration of a therapeutically effective amount of Compound I to a patient in need thereof.

In one embodiment, the invention relates to a method for the treatment of a CNS disease in a patient who suffers from a further disease wherein weight increase should be avoided the method comprising the administration of a therapeutically effective amount of Compound I to a patient in need thereof.

20 In one embodiment, the invention provides Compound I for use in the manufacture of a medicament for the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased or reduced (or has to be ceased or reduced) due to weight related adverse events.


25 In one embodiment, the invention relates to the use of Compound I in the manufacture of a medicament for the treatment of a CNS disease in a patient who is overweight.

In one embodiment, the invention relates to the use of Compound I in the manufacture of a medicament for the treatment of a CNS disease in a patient who suffers from a further disease wherein weight increase should be avoided.

Detailed description of the invention

30 The molecular structure of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine is shown below

5



In the present context, 1-[2-(2,4-dimethylphenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof is referred to as Compound I. A particular salt, e.g. the hydrobromide salt is referred to as Compound I HBr.

5 In one embodiment, said pharmaceutically acceptable salts are acid addition salts of acids that are non-toxic. Said salts include salts made from organic acids, such as maleic, fumaric, benzoic, ascorbic, succinic, oxalic, bis-methylenesalicylic, methanesulfonic, ethanedisulfonic, acetic, propionic, tartaric, salicylic, citric, gluconic, lactic, malic, mandelic, cinnamic, citraconic, aspartic, stearic, palmitic, itaconic, glycolic, p-aminobenzoic, glutamic, benzenesulfonic, 10 theophylline acetic acids, as well as the 8-halothephyllines, for example 8-bromothephylline. Said salts may also be made from inorganic salts, such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric and nitric acids. Particular mention is made of salts made from methanesulfonic acid, maleic acid, fumaric acid, meso-tartaric acid, (+)-tartaric acid, (-)-tartaric acid, hydrochloric acid, hydrobromic acid, sulphuric acid, phosphorous acid and nitric acid. 15 Distinct mention is made of the hydrobromide salt.

Oral dosage forms, and in particular tablets, are often preferred by the patients and the medical practitioner due to the ease of administration and the consequent better compliance. For tablets, it is preferable that the active ingredients are crystalline. In one embodiment, the invention relates to the use of Compound I in a crystalline form. The crystallinity of Compound I 20 is evidenced by the XRDP shown in e.g WO 2007/144005.

As evidenced by the data provided in WO 2007/144005 (see example 4a-f) the hydrobromide salt of 1-[2-(2,4-dimethylphenylsulphanyl)-phenyl]piperazine (Compound I HBr) may exist in several forms, i.e. it is polymorphic. The polymorphic forms have different properties. The beta form Compound I HBr (as defined in WO 2007/144005) is the more stable 25 as demonstrated by the higher DSC melting point and the lower solubility. Moreover, the beta form has an attractive combination of low hygroscopicity and solubility, which makes this compound particularly suited for making tablets. Hence, in one embodiment, the invention

provides the use of the hydrobromide salt of 1-[2-(2,4-dimethylphenylsulphanyl)-phenyl]piperazine with XRDP reflections at approximately 6.89, 9.73, 13.78 and 14.62 ($^{\circ}2\theta$) (± 0.1), such as at approximately 6.89, 8.48, 9.73, 13.78, 14.62 and 24.73 ($^{\circ}2\theta$) (± 0.1).

Alternatively, if Compound I is provided in an oral drop formulation, other salts
5 characterised by higher solubility may be preferred. When a compound is administered as an oral drop formulation, a few drops of a concentrated, liquid formulation of said drug is measured out and added to a glass of water, juice or the like that the patient drinks. As an example, the antidepressant cipramil is provided as an oral drop formulation at 40 mg/ml. The DL lactate salt of Compound I has been found to have a high solubility and therefore to be particularly suited
10 for oral drop formulations. Hence, in one embodiment, the invention provides the use of the DL-lactate salt of 1-[2-(2,4-dimethylphenylsulphanyl)-phenyl]piperazine.

There are numerous reasons why weight gain should be avoided in general and in relation with the treatment of CNS diseases in particular. Obesity and overweight is associated with serious conditions, such as type II diabetes mellitus, hypertension and coronary heart diseases. A
15 person who gains 5-7.9 kg as adult is 1.9 times more likely to develop type II diabetes mellitus than a person who maintains a stable weight after the age of 18 [*J. Clin. Psych.*, 1999, 60(suppl 21), 5-9, 1999]. Hence, a primary focus in the treatment of diseases, such as type II diabetes mellitus, hypertension and coronary heart diseases is a dietary effort in order to make the patient lose weight. In more general terms, weight gain is associated with impaired physical functioning,
20 reduced quality of life and poor mental health. Treatment compliance is a further consideration. Due to the physical discomfort, the reduced self esteem and the above discussed increased health risk associated with weight increase many patients who experience treatment emergent weight gain are reluctant to take the drug as prescribed with the consequent poor treatment outcome. Hence, weight gain associated with treatment of CNS diseases is an essential factor and it is
25 important to be able to offer a treatment of CNS diseases with no or only a few weight associated adverse events.

As shown in the examples, Compound I is not associated with weight gain in long-term treatment. Therefore, Compound I is particularly well suited for long-term treatment, treatment of patients who cannot accept other medication due to weight related adverse events (i.e. as 2nd
30 line treatment), for patients who in addition to their CNS disease suffer from a disease wherein it is important to avoid weight gain, and in patients that are overweight.

In the present context, “long-term” is intended to include maintenance treatment, and it is characterised by a treatment period above 12 weeks, such as above 24 weeks, such as above 48 weeks, such as above 72 weeks. Long-term treatment includes chronic treatment, such as life-long treatment.

5 In one embodiment, the invention relates to Compound I for use in the treatment of a CNS disease in a patient, wherein said treatment is associated with little or no weight gain. In particular, said treatment is long-term.

In one embodiment, the invention relates to Compound I for use in the treatment of a CNS disease in a patient, wherein said patient loses weight. In particular, said patient has
10 previously received another medication for the treatment of said disease, wherein said other medication is associated with weight gain.

In one embodiment, the invention provides Compound I for use in the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased (or has to be ceased) due to weight
15 related adverse events. Weight gain associated adverse events include hypertension, increased blood sugar level and decreased self esteem or quality of life. In particular, said adverse event is weight gain. The previously received medication is typically a psychotropic, such as an antipsychotic or an antidepressant.

In one embodiment, the invention provides Compound I for use in the treatment of a
20 CNS disease in a patient who is overweight. In the present context, “overweight” can be defined in terms of Body Mass Index (BMI). In one embodiment, said patient has a BMI above 25, such as above 30. BMI is defined as the mass in kilogram divided by the height in meter squared.

In one embodiment, the invention relates to Compound I for use in the treatment of a
25 CNS disease in a patient who suffers from a (further) disease where weight increase should be avoided. As discussed above, such further disease includes type II diabetes, hypertension and coronary heart diseases. Additionally, it may include conditions, such as abstinence from nicotine, e.g. smoking cessation. In particular, said treatment is long-term.

According to its unique pharmacological profile, Compound I is expected to be useful in the treatment of a various CNS diseases. The International patent applications published as WO
30 03/029232, WO 2007/144005, WO 2008/113359 and WO 2009/062517 disclose examples of such CNS diseases. In particular, CNS disease as used in the present context denotes a disease selected from depression disorders, anxiety disorders, chronic pain and abuse.

Depression disorders include major depressive disorder, dysthymic disorder, depression associated with bipolar disorder and depression with anxious features (anxious depression). Anxiety disorders include generalised anxiety disorder, panic attacks, phobias, obsessive-compulsive disorder and post-traumatic stress disorder. Chronic pain includes phantom limb
5 pain, neuropathic pain, diabetic neuropathy, post-herpetic neuralgia, carpal tunnel syndrome, HIV neuropathy, migraine, tension-type headache, and fibromyalgia syndrome. Abuse includes craving for or abuse of alcohol, narcotics and drugs.

In one embodiment, the invention provides method for the long-term treatment of a CNS disease comprising the administration of a therapeutically effective amount of Compound I to a
10 patient in need thereof wherein said administration is not associated with weight gain.

In one embodiment, the invention provides a method for the treatment of a CNS disease comprising the administration of a therapeutically effective amount of Compound I to a patient in need thereof, wherein said administration is associated with weight loss. In particular, said patient has previously received another medication for the treatment of said disease, wherein said
15 other medication is associated with weight gain.

In one embodiment, the invention provides a method for the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased (or has to be ceased) due to weight related adverse events, the method comprising the administration of a therapeutically effective amount of
20 Compound I to said patient in need thereof. Weight gain associated adverse events include hypertension, increased blood sugar level and decreased self esteem or quality of life. In particular, said adverse event is weight gain. The previously received medication is typically a psychotropic, such as an antipsychotic or an antidepressant.

In one embodiment, the invention provides a method for the treatment of a CNS disease
25 in a patient who is overweight, the method comprising the administration of a therapeutically effective amount of Compound I to a patient in need thereof. In one embodiment, said patient has a BMI above 25, such as above 30.

In one embodiment, the invention provides a method for treatment of a CNS disease in a patient in a need thereof, the method comprising the steps of a) determining the BMI of said
30 patient; and b) administering a therapeutically effective amount of Compound I to said patient if the BMI is determined to be above 25, such as above 30.

In one embodiment, the invention relates to a method of treating a CNS disease in a patient in need thereof, wherein said patient is suffering from a further disease wherein weight increase should be avoided, the method comprising the administration of a therapeutically effective amount of Compound I to said patient. In one embodiment, said further disease is selected from type II diabetes mellitus, hypertension, coronary heart disease, or the patient is abstaining from nicotine. In particular, said treatment is long-term.

In one embodiment, the invention provides a method for the treatment of a CNS disease in a patient in need thereof, the method comprising the steps of a) diagnosing whether said patient is suffering from a disease wherein weight gain should be avoided; and b) administering a therapeutically effective amount of Compound I to said patient if said diagnosis shows that said patient is suffering from a disease wherein weight gain should be avoided. In particular, such disease wherein weight gain should be avoided includes type II diabetes, hypertension and coronary heart diseases. Additionally, it may include conditions, such as abstinence from nicotine, e.g. smoking cessation. In particular, said treatment is long-term.

In one embodiment, the method provides a method of promoting the sales of Compound I, said method comprising the public distribution of the information that administration of Compound I to patients is not unfavourably associated with weight gain, or that the administration of Compound I to patients is not or only to a limited extent associated with weight gain.

In one embodiment, distribution of said information is achieved by a method selected from the group consisting of verbal communication, pamphlet distribution, print media, audio tapes, magnetic media, digital media, audiovisual media, billboards, advertising, newspapers, magazines, direct mailings, radio, television, electronic mail, braille, electronic media, banner ads, fiber optics, and laser light shows.

In one embodiment, the invention provides the use of Compound I in the manufacture of a medicament for the long-term treatment of a CNS disease, wherein said treatment is associated with little or no weight gain.

In one embodiment, the invention relates to the use of Compound I for the manufacture of a medicament for the treatment of a CNS disease in a patient, wherein said patient loses weight. In particular, said patient has previously received another medication for the treatment of said disease, wherein said other medication is associated with weight gain.

In one embodiment, the invention provides the use of Compound I in the manufacture of a medicament for the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased (or has to be ceased) due to weight related adverse events. Weight gain associated adverse events include hypertension, increased blood sugar level and decreased self esteem or quality of life. In particular, said adverse event is weight gain. The previously received medication is typically a psychotropic, such as an antipsychotic or an antidepressant.

In one embodiment, the invention provides the use of Compound I in the manufacture of a medicament for the treatment of a CNS disease in a patient who is overweight. In one embodiment, said patient has a BMI above 25, such as above 30.

In one embodiment, the invention relates to the use of Compound I in the manufacture of a medicament for the treatment of a CNS disease in a patient who suffers from a further disease wherein weight increase should be avoided. As discussed above, such further disease includes type II diabetes, hypertension and coronary heart diseases. Additionally, it may include conditions, such as abstinence from nicotine, e.g. smoking cessation. In particular, said treatment is long-term.

A "therapeutically effective amount" of a compound as used herein means an amount sufficient to cure, alleviate or partially arrest the clinical manifestations of a given disease and its complications in a therapeutic intervention comprising the administration of said compound. An amount adequate to accomplish this is defined as "a therapeutically effective amount". Effective amounts for each purpose will depend on the severity of the disease or injury as well as the weight and general state of the subject. It will be understood that determining an appropriate dosage may be achieved using routine experimentation, by constructing a matrix of values and testing different points in the matrix, which is all within the ordinary skills of a trained physician.

The term "treatment" and "treating" as used herein means the management and care of a patient for the purpose of combating a condition, such as a disease or a disorder. The term is intended to include the full spectrum of treatments for a given condition from which the patient is suffering, such as administration of the active compound to alleviate the symptoms or complications, to delay the progression of the disease, disorder or condition, to alleviate or relief the symptoms and complications, and/or to cure or eliminate the disease, disorder or condition as well as to prevent the condition, wherein prevention is to be understood as the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the

administration of the active compounds to prevent the onset of the symptoms or complications. Nonetheless, prophylactic (preventive) and therapeutic (curative) treatment are two separate aspects of the invention. The patient to be treated is preferably a mammal, in particular a human being.

5 Compound I is typically administered in daily doses of 1-100 mg, such as 2- 40 mg, such as 2, 5, 10, 15, 20, 25, 30, 35 mg. A daily dose may involve once daily dosing, or dosing two or more times daily.

 Compound I is conveniently administered as a pharmaceutical composition which may be prepared by conventional methods in the art. Particular mentioning is made of tablets, which
10 may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: anhydrous calcium hydrogen phosphate, PVP, PVP-VA co-polymers, microcrystalline cellulose, sodium starch glycolate, corn starch, mannitol, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvants or
15 additives usually used for such purposes such as colourings, flavourings, preservatives etc. may be used provided that they are compatible with the active ingredients.

 Solutions for injections may be prepared by dissolving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to desired volume, sterilising the solution and filling it in suitable ampules or vials. Any suitable
20 additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

 Oral drop formulations typically comprise, in addition to the active ingredient, excipients selected from solvent, buffer, surfactant, surface tension modifier, viscosity modifier, preservative, antioxidant, colourants, tastes masker, and flavour.

25 The pharmaceutical compositions which may be used in this invention may be administered by any suitable route, for example orally in the form of tablets, capsules, powders, syrups, liquids etc., or parenterally in the form of solutions for injection. For preparing such compositions, methods well known in the art may be used, and any pharmaceutically acceptable carriers, diluents, excipients or other additives normally used in the art may be used.

30 Conveniently, Compound I is administered in unit dosage form containing said compound in an amount of about 1 to 50 mg, such as 5, 10, 15 or 20 mg. Tablets comprising Compound I may conveniently be prepared by wet granulation. Using this method, the dry solids

(active ingredients, filler, binder etc.) are blended and moistened with water or another wetting agent (e.g. an alcohol) and agglomerates or granules are built up of the moistened solids. Wet massing is continued until a desired homogenous particle size has been achieved whereupon the granulated product is dried. Compound I is typically mixed with lactose monohydrate, corn starch and copovidone in a high shear mixer together with water. Following formation of granulates, these granulates may be sieved in a sieve with a suitable sieve size, and dried. The resulting, dried granulates are then mixed with microcrystalline cellulose, croscarmellose sodium and magnesium stearate, following which the tablets are pressed. Alternatively, wet granulation of Compound I may be achieved using mannitol, corn starch and copovidone, which granulates are mixed with microcrystalline cellulose, sodium starch glycolate and magnesium stearate before tablets are pressed. Alternatively, wet granulation of Compound I may be achieved by using anhydrous calcium hydrogen phosphate, corn starch and copovidone, which granulates are mixed with microcrystalline cellulose, sodium starch glycolate (type A), talc and magnesium stearate before tablets are pressed. Copovidone is a PVP-VA copolymer. Alternatively, tablets comprising Compound I may be obtained by mixing Compound I, mannitol and microcrystalline cellulose in a fluid bed granulation dryer onto which mixture an aqueous solution of hydroxypropyl cellulose is sprayed to give a granulated powder. The obtained granulates are then mixed with microcrystalline cellulose, sodium starch glycolate and magnesium stearate. The obtained mixture can subsequently be pressed into tablets. Typically, the tablets are coated with a suitable coating material.

All references, including publications, patent applications, and patents, cited herein are hereby incorporated by reference in their entirety and to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein (to the maximum extent permitted by law), regardless of any separately provided incorporation of particular documents made elsewhere herein.

The use of the terms "a" and "an" and "the" and similar referents in the context of describing the invention are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. For example, the phrase "the compound" is to be understood as referring to various compounds used in the invention or particular described aspect, unless otherwise indicated.

Unless otherwise indicated, all exact values provided herein are representative of corresponding approximate values (e.g., all exact exemplary values provided with respect to a

particular factor or measurement can be considered to also provide a corresponding approximate measurement, modified by "about," where appropriate).

The description herein of any aspect or aspect of the invention using terms such as “comprising”, “having,” “including,” or “containing” with reference to an element or elements is intended to provide support for a similar aspect or aspect of the invention that “consists of”, “consists essentially of”, or “substantially comprises” that particular element or elements, unless otherwise stated or clearly contradicted by context (e.g., a composition described herein as comprising a particular element should be understood as also describing a composition consisting of that element, unless otherwise stated or clearly contradicted by context).

10 Examples

Example 1 Long-term effects on body weight in MDD patients

The effects on weight associated with treatment with 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine HBr was investigated in a relapse prevention study in patients with major depressive disorder. Approximately 600 patients were enrolled in an open-label study in which they received 5 or 10 mg Compound I HBr (calculated as free base). After 12 weeks, responders (N~400) were randomised to placebo or 5/10 mg Compound I HBr for a 24 weeks double-blind placebo-controlled study. The recruitment of patients took place over time, and since the end of the study was defined by the last patient having completed the 24 weeks study, some patients received up to 64 weeks double-blind placebo-controlled treatment. This study is also reported in poster NR4-14 presented at the 164th Annual Meeting of the American Psychiatric Association, May 14-18 2011, Honolulu, Hawaii, USA.

The tables below show the mean difference in weight relative to the start of the open label period and to the time of randomisation and the fraction of patients experiencing equal to or more than 7% weight gain during the study, i.e. potentially clinically significant increase.

Table 1 Difference in weight relative to start of open label period (t=0)

Treatment	Week	Number of patients	Δ weight (kg)	≥7% weight gain (%)
Placebo	12	192	0.4	0.5
Placebo	24	188	0.9	4.8
Placebo	36	136	0.8	3.7

Placebo	48	66	0.3	4.5
Placebo	60	28	0.9	10.7
Placebo	72	12	0.0	8.3
Placebo	76	3	4.3	33.3
Compound	12	204	0.0	2.0
Compound	24	199	0.3	6.5
Compound	36	157	0.6	8.9
Compound	48	90	0.7	8.9
Compound	60	52	0.4	9.6
Compound	72	19	-0.2	10.5
Compound	76	3	-2.3	0.0

Table 2 Weight difference relative to time of randomisation (t=12 weeks)

Treatment	Week	Number of patients	Δ weight (kg)	$\geq 7\%$ weight gain (%)
Placebo	24	188	0.5	2.7
Placebo	36	136	0.3	3.7
Placebo	48	66	0.1	4.5
Placebo	60	28	1.1	7.1
Placebo	72	12	0.2	16.7
Placebo	76	3	4.9	66.7
Compound	24	199	0.4	2.0
Compound	36	157	0.6	6.4
Compound	48	90	0.9	8.9
Compound	60	52	0.8	13.5
Compound	72	19	0.3	10.5
Compound	76	3	-0.8	0.0

When the data in table 1 are evaluated it should be kept in mind that patients in the placebo arm
 5 received active treatment for the first 12 weeks. Nevertheless, the data presented in tables 1 and

2 clearly show that even after long-term treatment there is no meaningful difference between placebo and 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine HBr as far as weight gain in concerned.

5 *Example 2 Long-term effects on body weight in GAD patients*

The effects on weight associated with treatment with 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine HBr was investigated in a relapse prevention study in patients with generalised anxiety disorder. 687 patients were enrolled in an open-label study in which they received 5 or 10 mg Compound I HBr (calculated as free base). After 20 weeks, responders to treatment (N=459) were randomised to placebo or 5/10 mg Compound I HBr for a 24-56 weeks double-blind placebo-controlled study. The recruitment of patients took place over time, and since the end of the study was defined by the last patient having completed at least 24 weeks, some patients received up to 56 weeks double-blind placebo-controlled treatment.

The tables below show the mean difference in weight relative to the start of the open label period and to the time of randomisation and the fraction of patients experiencing equal to or more than 7% weight gain during the study, i.e. potentially clinically significant increase.

Table 3 Difference in weight relative to start of open label period (t=0)

Treatment	Week	Number of patients	Δ weight (kg)	$\geq 7\%$ weight gain (%)
Placebo	20	227	0.1	4.0
Placebo	32	222	0.3	5.9
Placebo	44	153	0.4	8.5
Placebo	56	72	0.3	9.7
Placebo	68	20	0.3	5.0
Placebo	76	3	2.0	0.0
Compound	20	227	0.3	2.6
Compound	32	220	0.5	3.6
Compound	44	179	0.8	8.4
Compound	56	84	1.5	17.9
Compound	68	28	1.5	10.7

Compound	76	4	0.9	0.0
----------	----	---	-----	-----

Tabel 4 Weight difference relative to time of randomisation (t=20 weeks)

Treatment	Week	Number of patients	Δ weight (kg)	$\geq 7\%$ weight gain (%)
Placebo	32	223	0.2	2.7
Placebo	44	154	0.4	5.8
Placebo	56	72	0.3	5.6
Placebo	68	20	0.5	0.0
Placebo	76	3	1.6	0.0
Compound	32	221	0.3	0.9
Compound	44	180	0.5	4.4
Compound	56	84	1.1	10.7
Compound	68	28	1.7	10.7
Compound	76	4	1.6	0.0

When the data in table 3 are evaluated it should be kept in mind that patients in the placebo arm received active treatment for the first 20 weeks. The combined data presented in tables 1-4 show that even after long-term treatment there is no meaningful difference between placebo and 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine HBr as far as weight gain is concerned.

Claims

1. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof for use in the long-term treatment of a CNS disease in a patient, wherein said
5 treatment is associated with little or no weight gain.
2. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof for use in the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased (or has to be ceased) due to weight related adverse events.
- 10 3. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof use in the treatment of a CNS disease in a patient who is overweight.
4. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof according to claim 3, wherein said patient is characterised by a BMI above 25.
5. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable
15 salts thereof for use in the treatment of a CNS disease in a patient who also suffers from a disease where weight increase should be avoided.
6. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof according to claim 5, wherein said patient also suffers from type II diabetes, hypertension or coronary heart diseases, or is abstaining from nicotine.
- 20 7. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof according to any of claims 1-6, wherein said CNS disease is selected from depression, anxiety, chronic pain and abuse.
8. 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof according to any of claims 1-7 which is the hydrobromide salt.
- 25 9. A method for the long-term treatment of a CNS disease comprising the administration of a therapeutically effective amount of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and

pharmaceutically acceptable salts thereof to a patient in need thereof wherein said administration is not associated with weight gain.

10. A method for the treatment of a CNS disease in a patient in need thereof who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased (or has to be ceased) due to weight related adverse events, the method comprising the administration of a therapeutically effective amount of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to said patient.
11. A method for the treatment of a CNS disease in a patient in need thereof who is overweight, the method comprising the administration of a therapeutically effective amount of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to said patient.
12. The method according to claim 12 wherein said patient is characterised by a BMI above 25.
13. A method for treatment of a CNS disease in a patient in a need thereof, the method comprising the steps of a) determining the BMI of said patient; and b) administering a therapeutically effective amount of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to said patient if the BMI is determined to be above 25.
14. A method of treating a CNS disease in a patient in need thereof, wherein said patient is suffering from a further disease wherein weight increase should be avoided, the method comprising the administration of a therapeutically effective amount of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to said patient.
15. The method according to claim 14 wherein said further disease said patient is selected from type II diabetes, hypertension or coronary heart diseases, or said patient is abstaining from nicotine.
16. A method for the treatment of a CNS disease in a patient in need thereof, the method comprising the steps of a) diagnosing whether said patient is suffering from a disease wherein weight gain should be avoided; and b) administering a therapeutically effective amount of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to

said patient if said diagnosis shows that said patient is suffering from a disease wherein weight gain should be avoided.

17. The method according to claim 16, wherein the disease being diagnosed for in step a) is selected from type II diabetes, hypertension and coronary heart diseases, or whether said patient
5 is abstaining from nicotine.

18. The method according to any of claims 9-17, wherein said CNS disease is selected from depression, anxiety, chronic pain and abuse.

19. A method of promoting the sales of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof, said method comprising the public distribution of
10 the information that administration of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to a patient is not unfavourably associated with weight gain or that the administration of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof to a patient is not or only to limited extent associated with weight gain.

15 20. The method according to any of claims 9-19, wherein said pharmaceutically acceptable salt is the hydrobromide salt.

21. Use of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof in the manufacture of a medicament for the long-term treatment of a CNS disease wherein said treatment is not associated with weight gain.

20 22. Use of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof in the manufacture of a medicament for the treatment of a CNS disease in a patient who has previously received medication (or is still receiving it) for the treatment of said disease which medication was ceased (or has to be ceased) due to weight related adverse events.

25 23. Use of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof in the manufacture of a medicament for the treatment of a CNS disease in a patient who is overweight.

24. The use according to claim 23 wherein said patient is characterised by a BMI above 25.
25. Use of 1-[2-(2,4-dimethyl-phenylsulfanyl)phenyl]piperazine and pharmaceutically acceptable salts thereof in the manufacture of a medicament for the treatment of a CNS disease in a patient who suffers from a further disease where weight increase should be avoided.
- 5 26. The use according to claim 26 wherein said further disease is selected from type II diabetes, hypertension and coronary heart diseases, or wherein said patient is abstaining from nicotine.
27. The use according to any of claims 21-26, wherein said CNS disease is selected from depression, anxiety, chronic pain and abuse.
- 10 28. The use according to any of claims 21-27, wherein said pharmaceutically acceptable salt is the hydrobromide salt.

INTERNATIONAL SEARCH REPORT

International application No PCT/DK2011/050317

A. CLASSIFICATION OF SUBJECT MATTER INV. A61K31/495 A61P25/00 ADD.				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) A61K A61P				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, BIOSIS, EMBASE, WPI Data				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	ARTIGAS F ET AL: "P.2.c.040 A randomised, double-blind, placebo-controlled, active-referenced study of Lu AA21004 in patients with major depression", EUROPEAN NEUROPSYCHOPHARMACOLOGY, ELSEVIER SCIENCE PUBLISHERS BV, AMSTERDAM, NL, vol. 19, 1 September 2009 (2009-09-01), pages S426-S427, XP026795343, ISSN: 0924-977X [retrieved on 2009-09-01] the whole document	1-28		
X	----- WO 2007/144005 A1 (LUNDBECK & CO AS H [DK]; BANG-ANDERSEN BENNY [DK]; FALDT ANDRE [DK]; M) 21 December 2007 (2007-12-21) claims 15,17 ----- -/--	1-28		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; border: none;"><input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.</td> <td style="width: 50%; border: none;"><input checked="" type="checkbox"/> See patent family annex.</td> </tr> </table>			<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.	<input checked="" type="checkbox"/> See patent family annex.
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.	<input checked="" type="checkbox"/> See patent family annex.			
* Special categories of cited documents :				
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family			
Date of the actual completion of the international search	Date of mailing of the international search report			
26 September 2011	04/10/2011			
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Loher, Florian			

INTERNATIONAL SEARCH REPORT

International application No PCT/DK2011/050317

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2009/062517 A1 (LUNDBECK & CO AS H [DK]; TAKEDA PHARMACEUTICALS NORTH A [US]; CHON JIN) 22 May 2009 (2009-05-22) claims 19,20 <p style="text-align: center;">-----</p>	1-28

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/DK2011/050317

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007144005 A1	21-12-2007	AR 061481 A1	27-08-2008
		CN 101472906 A	01-07-2009
		EA 200970018 A1	30-06-2009
		KR 20090028712 A	19-03-2009
		US 2010297240 A1	25-11-2010
		ZA 200810017 A	28-04-2010

WO 2009062517 A1	22-05-2009	AR 069260 A1	06-01-2010
		AU 2008323390 A1	22-05-2009
		CA 2705163 A1	22-05-2009
		CN 102014908 A	13-04-2011
		CO 6270322 A2	20-04-2011
		EA 201070598 A1	29-10-2010
		EP 2219647 A1	25-08-2010
		JP 2011503125 A	27-01-2011
		KR 20100099115 A	10-09-2010
		US 2011201617 A1	18-08-2011
