



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2351671 A1 2000/06/08

(21) **2 351 671**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 1999/11/24
(87) Date publication PCT/PCT Publication Date: 2000/06/08
(85) Entrée phase nationale/National Entry: 2001/05/16
(86) N° demande PCT/PCT Application No.: US 99/28033
(87) N° publication PCT/PCT Publication No.: WO 00/32750
(30) Priorité/Priority: 1998/11/24 (60/109,687) US

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 5/06, A61L 27/38, A61K 48/00,
A61P 7/04, C12Q 1/02, C12N 5/10, C12N 5/08,
C07K 14/755
(71) Demandeurs/Applicants:
REGENTS OF THE UNIVERSITY OF MINNESOTA, US;

EMORY UNIVERSITY, US
(72) Inventeurs/Inventors:
LOLLAR, JOHN S., US;
LIN, YI, US;
HEBBEL, ROBERT P., US
(74) Agent: GOUDREAU GAGE DUBUC

(54) Titre : CELLULES ENDOTHELIALES TRANSGENIQUES CIRCULANTES
(54) Title: TRANSGENIC CIRCULATING ENDOTHELIAL CELLS

(57) **Abrégé/Abstract:**

A process is provided for expanding the population of endothelial cells obtained from peripheral blood which can be transformed with a vector comprising a DNA sequence encoding a preselected bioactive polypeptide. The resulting transgenic endothelial cells are useful to biocompatibilize implantable medical devices or can be used directly, as for gene therapy.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 5/06, 5/08, 5/10, A61K 48/00, A61L 27/38, C12Q 1/02, A61P 7/04 // C07K 14/755	A1	(11) International Publication Number: WO 00/32750 (43) International Publication Date: 8 June 2000 (08.06.00)
(21) International Application Number: PCT/US99/28033 (22) International Filing Date: 24 November 1999 (24.11.99) (30) Priority Data: 60/109,687 24 November 1998 (24.11.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/109,687 (CIP) Filed on 24 November 1998 (24.11.98) (71) Applicants (for all designated States except US): REGENTS OF THE UNIVERSITY OF MINNESOTA [US/US]; Morrill Hall, 100 Church Street S.E., Minneapolis, MN 55455 (US). EMORY UNIVERSITY [US/US]; Room 1003, Woodruff Memorial Building, 1639 Pierce Drive, Atlanta, GA 30322 (US). (71)(72) Applicants and Inventors: HEBBEL, Robert, P. [US/US]; 25 Island Road, North Oaks, MN 55127 (US). LIN, Yi [US/US]; 1410 Carling Drive, #108, St. Paul, MN 55108 (US). LOLLAR, John, S. [US/US]; 2568 Oak Crossing Drive, Decatur, GA 30033 (US).	(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg, Woessner & Kluth, P.O. Box 2938, Minneapolis, MN 55402 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	

(54) Title: TRANSGENIC CIRCULATING ENDOTHELIAL CELLS**(57) Abstract**

A process is provided for expanding the population of endothelial cells obtained from peripheral blood which can be transformed with a vector comprising a DNA sequence encoding a preselected bioactive polypeptide. The resulting transgenic endothelial cells are useful to biocompatibilize implantable medical devices or can be used directly, as for gene therapy.

TRANSGENIC CIRCULATING ENDOTHELIAL CELLS

BACKGROUND OF THE INVENTION

This invention was made with the support of NIH Grant No. HL 55174 (now
5 HL 30160). The Government has certain rights in the invention.

The endothelial cell participates in numerous functions of vascular physiology. Many factors, such as cytokines, can alter the surface of the endothelial cell and thereby modulate the role of the endothelium in coagulation, inflammation, vaso-regulation, and adhesion. See, for example, R. P. Hebbel et al., J. Lab. Clin.
10 Med., 129, 288 (1997); J. S. Pober, Am. J. Path., 133, 426 (1988); E. J. Favalaro, Immunol. Cell. Biol., 71, 571 (1993). The endothelial cell may also have a key role in the vascular pathology of sickle cell anemia, including the vaso-occlusions that cause acute painful crises. However, research in this area has been hindered by the inaccessibility of vascular endothelium in patients. For example, E. M. Levine et al.
15 (U.S. Pat. No. 5,132,223) disclosed cloning and serial cultivation of adult human endothelial cells derived from brain-dead, but heart-beating cadaver organs. K. Gupta et al., Exp. Cell. Res., 230, 244 (1997) reported the culture of microvascular endothelial cells derived from newborn human foreskin. Thus, circulating endothelial cells might provide useful material for the study of vascular pathologies,
20 for gene therapy, and for biomedical engineering applications. In previous investigations increased numbers of circulating endothelial cells have been found in sickle cell anemia and other conditions associated with vascular injury, such as that due to cytomegalovirus infection, rickettsial infection, myocardial infarction, intravascular instrumentation, and endotoxemia. See, for example, F. George et al., Blood, 80, Suppl: 12a, abstract (1992); E. Percivalle et al., J. Clin. Invest., 92, 663 (1993), F. George et al., Blood, 82, 2109 (1993); C. A. Bouvier et al., Thomb. Diath. Haemorrh. Suppl., 40, 163 (1970); F. George et al., J. Immunol. Meth., 139, 65 (1991) and R. G. Gerrity et al., Exp. Mol. Pathol., 24, 59 (1976).

WO 00/32750

PCT/US99/28033

However, in normal donors, there are only about 2-3 circulating endothelial cells per ml of blood; they have a quiescent phenotype, and about 50% of them are microvascular as evidenced by CD36 positivity. See, A. Solovey et al., New Engl. J. Med., 337, 1584 (1997), who reported using the methodology of Gupta et al., cited above, to coculture viable circulating endothelial cells identified in the blood of patients with sickle cell anemia with primary microvascular endothelial cells (MVEC). T. Asahara et al., Science, 275, 964 (1997) isolated putative endothelial cell (EC) progenitors from human peripheral blood after CD34⁺ enrichment by magnetic bead selection on the basis of cell surface antigen expression. The cells were cultured on fibronectin-coated wells in modified M-199 medium containing bovine brain extract and 20% fetal bovine serum. Q. Shi et al., Blood, 92, 362 (1998) characterized bone marrow-derived precursor endothelial cells by isolating CD34⁺ cells derived from peripheral blood using murine anti-CD34⁺ antibody binding followed by exposure to anti-mouse immunomagnetic beads. The cells were cultured in gelatin or fibronectin-coated plastic wells in M199 medium containing VEGF, FBS, bFGF and IGF.

However, due to the low concentration of CEC in blood, a need exists for a culture method and medium that will permit the rapid expansion of CEC from blood, without the attendant difficulties of isolation discussed above.

20

SUMMARY OF THE INVENTION

The present invention provides a process for expanding the population of endothelial cells (EC) present in an aliquot of peripheral mammalian blood, i.e., blood obtained from an animal or human patient. The process comprises culturing buffy coat mononuclear cells, which are readily obtained from peripheral mammalian blood. The buffy coat mononuclear cells are cultured in contact with a surface coated with collagen I, such as a coated plastic culture well, wall of a culture flask or bioreactor, in a culture medium containing an effective amount of vascular endothelial growth factor (VEGF), which medium is free of bovine brain extract. Optionally, the buffy coat mononuclear cells can thereafter be cultured in contact

30

with a surface coated with fibronectin/gelatin. The medium can also comprise heparin and/or dextran sulfate, and other growth factors conventionally employed in endothelial cell culture media. The present method accomplishes the rapid and extensive expansion of the initial population of endothelial cells and any endothelial progenitor cells present in the population of buffy coat mononuclear cells. For example, the present method typically results in an outgrowth of endothelial cells that is at least a billion-fold greater than the 20-30 endothelial cells identifiable on culture day #2, when endothelial cells are first counted. The endothelial outgrowth includes the expansion of two populations. There is a limited expansion of the mature endothelial cells of vessel wall origin that are found in fresh blood, as well as a delayed, but greater, expansion of a rarer population of marrow-derived cells, e.g., endothelial progenitor cells or angioblasts, that are also found in fresh blood.

Unexpectedly, the cultured endothelial cells have been found to be amenable to cryopreservation in conventional cryopreservation media, followed by thawing and continued culture/expansion. The present method is thus more convenient and less complex than methods based on processing and culturing adult vessel human endothelial cells, as described, for example, in E. M. Levine et al. (U.S. Pat. No. 5,132,223). It does not comprise selection of sub-populations of EC-enriched hematopoietic cells using antibodies and/or magnetic bead technology.

The present invention also comprises an isolated, purified transgenic mammalian endothelial cell comprising a recombinant DNA sequence encoding at least one biologically active preselected protein or polypeptide, such as a Factor VIII protein, and optionally, a selectable marker gene or reporter gene. Preferably, the transgenic endothelial cell is prepared by stably transforming (transfecting/transducing) a population of circulating endothelial cells outgrown from blood in accord with the present method, with a vector comprising an isolated DNA sequence encoding the protein/polypeptide of interest, operably linked to a promoter functional in said endothelial cells.

A population of said transgenic cells can be formulated into a pharmaceutical composition and administered to a mammal, such as a human patient

WO 00/32750

PCT/US99/28033

afflicted with hemophilia, preferably in combination with a pharmaceutically acceptable carrier. The carrier may be a liquid carrier (for example, a saline solution) or a solid carrier; e.g., an implant. In employing a liquid carrier, the engineered cells may be introduced, e.g., intravenously, sub-cutaneously, intramuscularly, intraperitoneally, intralesionally, and the like. The polypeptide, such as Factor VIII protein or proteins are expressed *in situ*, i.e., in the bloodstream of said mammal in an amount effective to treat (alleviate) said hemophilia. Since the transgenic endothelial cells will also migrate to the bone marrow, they are also useful for many aspects of gene therapy, apart from treating circulatory pathologies. Other uses of these transgenic EC, such as in biocompatibilization of implantable devices, diagnostics, and local drug delivery, are discussed hereinbelow.

Novel vectors useful for transforming the expanded circulating endothelial cells of the invention are also within the scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph plotting the outgrowth of endothelial cells in accord with the method of the invention. Arrows show times at which cultures were passaged. All data points are plotted as mean $\pm$ SD. Data are shown for n = 5 through culture passage 6, and n = 4 for subsequent passages.

Figure 2 is a graph plotting the outgrowth of endothelial cells that were subjected to cryopreservation.

Figure 3 depicts the DNA sequence of HSQ/eGFP (SEQ ID NO:1). (HFVIII/SQ/egfp) which comprises original HB with B-domain SQ insert based on Lind et al., Eur. J. Biochem., 232; and p21 primer sequence containing eGFP protein sequence.

Figure 4 shows the DNA sequence (SEQ ID NO:2) of HSQRENeo.

Figure 5 shows the results of G418 selection (shaded area) of outgrowth endothelial cells stably transfected with a vector encoding eGFP.

Figure 6 shows factor VIII activity in conditioned media from cells transfected with various constructs (see Example 7).

Figure 7 depicts the results of a RT-PCR analysis for human FVIII mRNA in cells transfected with various constructs.

5

DETAILED DESCRIPTION OF THE INVENTION

The endothelial cell culture medium used in the process of this invention comprises any of the media conventionally employed for the culture of this type of cell, as modified in accord with the present method, to include an effective amount
10 of endothelial cell growth factor (ECG or VEGF), and to exclude bovine brain extract. The exclusion of bovine brain extract is advantageous both from a standpoint of consistency of formulation, and from a health standpoint, in that cells are not subjected to the risk of contamination with infectious agents (viruses, prions, etc.) which may be present in such extracts.

15 Preferably, the medium contains heparin, dextran sulfate or a combination thereof. These materials are described in detail in U.S. Pat. No. 5,132,223. Of the various basal endothelial cell growth media, EGM[®]-2 (Clonetics, Inc., San Diego, CA) is particularly preferred. It is based on CCMD[®]130 culture medium plus human epidermal growth factor (hEGF), human basic fibroblast growth factor
20 (hFGF-B), human recombinant insulin-like growth factor (Long R3-IGF-1). FGF can be present at about 0.5-5 ng/ml, EGF can be present at about 0.5-10 ng/ml and IGF can be present at about 1-7.5 ng/ml of the culture medium.

The medium can also comprise hydrocortisone (0.1-2 µg/ml), heparin (1-20 µg/ml), gentamicin, amphotericin-B (0.1-0.5 mg/ml) and fetal bovine serum.
25 Vascular endothelial growth factor (VEGF) is present at about 1-100 ng/ml, preferably about 5-25 ng/ml. It is included within this concentration range in some commercially available media or supplements or is available from Collaborative Research, Inc. VEGF can also be prepared as described by Maciag et al., PNAS USA, 76, 5674 (1979).

A medium that is effective to disperse and pre-wash the buffy coat mononuclear cells prior to culturing, has been described by K. Gupta et al., Exp. Cell Res., 230, 244 (1997). This endothelial culture medium consists of MCDB 131 medium supplemented with 1 µg/ml hydrocortisone acetate, 5×10^{-4} M dibutyryl cAMP, 1.6 nM L-glutamine, 100 U/ml penicillin, 100 U/ml streptomycin, 0.25 mg/ml amphotericin B, 0.004% heparin and 20% human male serum. VEGF (0.01-100 ng/ml) can also be added to this medium.

Circulating endothelial cells can be isolated from whole blood via isolation of the buffy coat mononuclear cells by methodologies well-known to the art. For example, see A. Solovey et al., New Engl. J. Med., 337, 1584 (1997). Whole blood is anti-coagulated and diluted with a physiological salt solution containing EDTA and BSA. The diluted blood is layered onto Histopaque® 1077 (Sigma Chem. Co.) and centrifuged at ambient temperature. The buffy coat mononuclear cells are collected and washed by centrifuging with the medium described in K. Gupta et al., Exp. Cell. Research, 230, 244 (1977), described hereinabove, i.e., supplemented MCDB 131 medium.

The buffy coat mononuclear cells are resuspended in the endothelial cell culture medium (i.e., EGM®-2) and held in a suitable vessel having the walls thereof coated with collagen I. Type I collagen is commercially available from Sigma Chem. Co. (St. Louis, MO) and is obtainable from bovine Achilles tendon, calf skin, rat tail, and human placenta. See, for example, Niyiloizi et al., J. Biol. Chem., 259, 14170 (1984). On day two of culture, unattached buffy coat mononuclear cells and cell debris are removed when the medium is changed, and the small number of adhered cells exhibit endothelial cell morphology, and stain positive with mAb P1H12.

Cultured cells were evaluated by inverted phase-contrast microscopy to confirm the characteristic cobblestone morphology of the endothelial cells and to examine for the presence of cells other than endothelial cells. Cells were also analyzed for von Willebrand factor (vWF) and CD36 expression by immunofluorescence and flow cytometry. The presence of cell surface CD36 has

been used as a marker for human microvascular endothelial cells (HDMEC), as it has been shown that most MEC express CD36, but endothelial cells lining large vessels do not. Briefly, cells in suspension were fixed with paraformaldehyde, cytopun onto glass slides, washed, and incubated with either rabbit anti-human
5 vWF (1:400 dilution) or anti-CD36 mAb FA6-152 (5 µg/ml) (Immunotech, Westbrook, ME). Slides were then washed and incubated with secondary antibodies conjugated to rhodamine or fluorescein, washed, and viewed under a fluorescent microscope.

To identify endothelial cells, the antibody P1H12 was also used. This
10 murine IgG1 monoclonal antibody was obtained by immunizing mice with HUVEC, generating a hybridoma line, and separating IgG from supernatants of hybridoma cell cultures with a protein G column. For some studies, fluorescein isothiocyanate-labeled P1H12 was used, prepared with the Fluoro Tag FITC Conjugation kit (Sigma).

15 P1H12 reacts specifically with endothelial cells in blood. It stains primary HUVEC and MVEC cultures and the endothelial cells of all vessels in frozen sections of human skin, intestine, ovary, tonsil, lymph node, lung, and kidney. It does not stain any other type of cell in those tissues. It does not stain carcinoma cell lines HT-29 and COLO205, melanoma cells lines A-375 and M21, the T-cell lines
20 Jurkat and HuT78, fibroblasts, HL-60 or Chinese-hamster-ovary cells, or Epstein-Barr virus-transformed B-cell lines. It does not stain monocytes, granulocytes, red cells, platelets, T cells, or B cells from marrow or peripheral blood; nor does it react with marrow megakaryocytes or the megakaryoblast line HU3. The peripheral blood cells that do stain with P1H12 are also positive for both von Willebrand factor
25 and thrombomodulin (the combined expression of which is limited to endothelium), and they stain for flt and flk (receptors for the endothelial-specific vascular endothelial growth factor). Subgroups of P1H12-positive blood cells also stain for CD34 and two endothelial-specific activation markers (VCAM and E-selectin).

After good EC growth is obtained, preferably after about 10^3 -fold expansion,
30 or at 15-25 days, the cells are trypsinized and isolated from the supernatant by

centrifugation. The cells are then suspended in the initial culture medium in a flask coated with fibronectin/gelatin for continued culture. Clones can be derived from secondary cultures and seeded at about 10 cells/cm² of flask surface. The clones are then serially propagated in the culture medium.

- 5 It was also found that EC cultured in accord with the present method can be cryopreserved and then thawed and returned to culture without significant loss of their capacity to proliferate. To cryopreserve the cells, the cultured cells can be detached as described hereinabove and resuspended in suitable cryopreservation medium, i.e., containing a cryopreservation agent such as sugar(s), BSA,
10 dimethylsulfoxide (DMSO), glycerol, glycerol esters and the like.

Cryopreservation of hematopoietic cells, such as bone marrow and peripheral blood fractions enriched in progenitor stem cells, has become standard clinical practice for autologous bone marrow transplantation (Areman et al., Bone Marrow and Stem Cell Processing: A Manual of Current Techniques, F. A. Davis,
15 Philadelphia, PA, 1st edition (1992)). Two basic techniques are used to cryopreserve hematopoietic cells. The most commonly used technique in clinical laboratories, uses tissue culture medium combined with 95% fetal calf serum ("FCS"), and 5 v/v % dimethylsulfoxide ("DMSO"). After suspension in the cryopreservation medium, the cells are placed on ice for about 5 minutes, then at
20 -70° C overnight and finally in liquid nitrogen. This technique was developed in the early 1960s by Ashwood-Smith (see Sputtek et al., Clinical Applications of Cryobiology, Chapter 5, 127-147 (CRC Press, Boca Raton, FL, 1991)), and has been the predominant technique used in clinical practice. In early 1983, Stiff and colleagues developed a modified method for the preservation of stem cells (Stiff et
25 al., Cryobiology, 20, 17-24 (1983)). In this method, the concentration of DMSO was reduced to 5% and an additional cryoprotective agent, hydroxyethyl starch ("HES"), was added to the solution. The cells could be frozen using a controlled rate freezer or in a mechanical freezer at -80°C.

In present clinical practice, the base of most cryopreservation solutions is a tissue culture medium, potentially containing many different components. Typical tissue culture media include RPMI 1640, IMDM, AIM-5, X-VIVO 10, or α MEM.

5 The cell suspension is then frozen, i.e., at liquid nitrogen temperatures in a suitable container and stored until needed. The frozen suspension is then thawed, by immersing the container in a warm water bath and culturing can be resumed. The resulting EC growth is similar to that of cells that were not subjected to cryopreservation.

The cultured EC may also be used in gene therapy wherein a gene producing
10 a protein polypeptide, enzyme or other product is inserted into the DNA of the EC. The transgenic EC are then administered to a mammal, e.g., by infusion into a patient's bloodstream. See, for example, B. P. Luskey et al., Annals. N. Y. Acad. Sci., 612, 398 (1990), B. A. Naughton et al. (U.S. Pat. No. 4,721,096), and Anderson et al. (U.S. Pat. No. 5,399,346).

15 The gene carried by the EC can be any gene which allows the blood cells to exert a therapeutic effect that they would not ordinarily have, such as a gene encoding a clotting factor, such as Factor VIII, useful in the treatment of hemophilia. The gene can encode one or more products having therapeutic effects. Examples of suitable genes include those that encode cytokines such as TNF,
20 interleukins (interleukins 1-12), interferons (α , β , γ -interferons), T-cell receptor proteins and Fc receptors for antigen-binding domains of antibodies, such as immunoglobulins.

Additional examples of suitable genes include genes that modify EC to "target" a site in the body to which the blood cells would not ordinarily "target,"
25 thereby making possible the use of the blood cell's therapeutic properties at that site. In this fashion, blood cells such as TIL can be modified, for example, by introducing a Fab portion of a monoclonal antibody into the cells, thereby enabling the cells to recognize a chosen antigen. Likewise, blood cells having therapeutic properties can be used to target, for example, a tumor, that the blood cells would not normally
30 target. Other genes useful in cancer therapy can be used to encode chemotactic

factors which cause an inflammatory response at a specific site, thereby having a therapeutic effect. Other examples of suitable genes include genes encoding soluble CD4 which is used in the treatment of HIV infection and genes encoding α -antitrypsin, which is useful in the treatment of emphysema caused by α -antitrypsin deficiency, or genes encoding factors which promote bone growth such as bone morphogenic proteins (BMPs) or other factors in the BMP pathway.

The gene therapy of the present invention is useful in the treatment of a variety of diseases including but not limited to adenosine deaminase deficiency, sickle cell anemia, thalassemia, hemophilia, diabetes, α -antitrypsin deficiency, brain disorders such as Alzheimer's disease, and other illnesses such as growth disorders and heart diseases, for example, those caused by alterations in the way cholesterol is metabolized and defects for the immune system, as well as to repair bone fractures or treat or prevent osteoporosis.

In still another embodiment, there is provided a method of detecting the presence of human CEC present in a patient, comprising: (i) inserting into an expanded population of human CEC removed from the patient a DNA segment encoding the detectable marker under conditions such that the marker is expressed in the CEC; (ii) introducing cells resulting from step (i) into the patient; (iii) removing from the patient an aliquot of tissue (which can be, for example, normal tissue, cancerous tissue, vascular tissue, blood, lymph nodes, etc.) including cells resulting from step (ii) and their progeny; and (iv) detecting or determining the quantity of the cells resulting from step (ii) and their progeny, in said aliquot.

To effect gene therapy with a substantially pure population of human EC, the following general method may be used to insert a gene into these EC. For a review of transformation methodologies, see Friedman, Science, 244, 1275 (1989) and Lancet, Jun. 4, 1988, p. 1271. In order to introduce a normal structural gene, to correct a genetic error, a gene can first be isolated from the cells of a donor. The cells may be isolated from tissue(s), blood or other body fluids. To find a gene coding for the defective protein, DNA from the donor cell is isolated and cleaved by enzymatic digestion into segments of varying length by means known to those

skilled in the art. The segments of DNA then may be inserted individually into vectors containing the appropriate regulatory sequences for expression of a gene product. The vectors then can be screened by conventional means such as Northern blotting, if the sequence for the normal gene is known, or the expression product can
5 be screened by Western blotting.

Alternatively, if the DNA sequence of the desired gene or the sequence of the normal protein is known, the gene can be made by synthetic methods such as by using a DNA synthesizer (Applied Biosystems). In any case, the method of isolation or construction of the gene sequence can yield a "normal" gene that codes
10 for the desired gene product. Once isolated, a functional structural gene, such as a DNA sequence coding for a factor VIII protein, human factor VIII, porcine factor VIII and the like can be modified for expression *in vivo* by linkage to suitable control regions, such as promoters. Hybrid or modified genes can also be constructed, such as DNA sequences encoding factor VIII molecules, modified to
15 reduce their antigenicity and immunogenicity. DNA methods can be used to substitute elements of animal factor VIII for the corresponding elements of human factor VIII, resulting in hybrid human/animal factor VIII molecules. The preparation of transformation vectors comprising DNA sequences encoding chimeric hybrid Factor VIII molecules is disclosed in Lollar et al. (U.S. Pat. No.
20 5,744,446). DNA methods may also be used to substitute amino acid residues at selected sites, such as epitopes, to yield factor VIII constructs having improved therapeutic attributes such as reduced antigenicity or immunogenicity or greater potency in comparison to unmodified human factor VIII.

In a preferred embodiment, a hybrid human/porcine cDNA encoding factor
25 VIII, in which the porcine sequence encodes a domain or part domain, such as the A2 domain or part domain, is inserted in a mammalian expression vector, such as ReNeo, to form a hybrid factor VIII construct. Preliminary characterization of the hybrid factor VIII is accomplished by insertion of the hybrid cDNA into the ReNeo mammalian expression vector and transient expression of the hybrid protein in
30 COS-7 cells. A determination of whether active hybrid protein is expressed can

then be made. The expression vector construct is used further to stably transfect cells in culture, such as baby hamster kidney cells, using methods that are routine in the art, such as liposome-mediated transfection (Lipofectin™, Life Technologies, Inc.). Expression of recombinant hybrid factor VIII protein can be confirmed, for example, by sequencing, Northern and Western blotting, or polymerase chain reaction (PCR). Hybrid factor VIII protein in the culture media in which the transfected cells stably expressing the protein are maintained can be precipitated, pelleted, washed, and resuspended in an appropriate buffer, and the recombinant hybrid factor VIII protein purified by standard techniques, including immunoaffinity chromatography using, for example, monoclonal anti-A2-Sepharose™.

In a further embodiment, the hybrid factor VIII comprising subunit, domain, or amino acid sequence substitutions is expressed as a fusion protein from a recombinant molecule in which a sequence encoding a protein or peptide that enhances, for example, stability, secretion, detection, isolation, or the like is inserted adjacent to the factor VIII encoding sequence. Established protocols for use of homologous or heterologous species expression control sequences including, for example, promoters, operators, and regulators, in the preparation of fusion proteins are known and routinely used in the art. See Current Protocols in Molecular Biology (Ausubel, F. M. et al., eds.), Wiley Interscience, New York.

The purified hybrid factor VIII or fragment thereof can be assayed for coagulation activity by standard assays including, for example, the plasma-free factor VIII assay, the one-stage clotting assay, and the enzyme-linked immunosorbent assay using purified recombinant human factor VIII as a standard. Antigenicity and immunogenicity of factor VIII constructs may be tested *in vitro* using the standard Bethesda inhibitor assay, or *in vivo* using a knockout hemophilic mouse model.

Other vectors, including both plasmid and eukaryotic viral vectors, may be used to express a recombinant gene construct in eukaryotic cells depending on the preference and judgment of the skilled practitioner. Other vectors and expression

systems, including bacterial, yeast, and insect cell systems, can be used but are not preferred due to differences in, or lack of, glycosylation.

The same methods employed for preparing hybrid human/porcine factor VIII having subunit, domain, or amino acid sequence substitution can be used to prepare other recombinant hybrid factor VIII protein and fragments thereof and the nucleic acid sequences encoding these hybrids, such as human/non-human, non-porcine mammalian or animal/animal. As used herein the term "Factor VIII protein" includes any of these materials that are biologically active as determined by the assays listed above, i.e., that possess coagulation activity when assayed *in vitro*.

10 The murine and part of the porcine factor VIII cDNA have been cloned. Factor VIII sequences of other species for use in preparing a hybrid human/animal or animal/animal factor VIII molecule can be obtained using the known human and porcine DNA sequences as a starting point. Other techniques that can be employed include PCR amplification methods with animal tissue DNA, and use of a cDNA

15 library from the animal to clone the factor VIII sequence.

Once the DNA containing the gene is prepared, the DNA can be inserted into the population of EC isolated and expanded as above. The DNA can be inserted by 1) physical methods such as coprecipitation with calcium phosphate, electroporation, lipofection or microinjection (e.g., U.S. Pat. No. 4,873,191), and/or

20 by 2) the use of viral vectors such as adenoviruses, if the DNA is less than approximately 7-8 kB, or retroviruses for longer segments of DNA. In the latter case, the DNA of the retrovirus is cut with a restriction enzyme and the human DNA containing the desired sequence is inserted and ligated. The retrovirus containing the insertion then is transfected into the EC. The EC then can be assayed for

25 production of the desired protein. See, e.g., U.S. Pat. Nos. 4,902,783 and 5,681,746.

In general, molecular DNA cloning methods are well known in the art and are not limiting in the practice of this invention. For a further description of similar methods, see Friedmann, Science, 244, 1275 (1989) and Molecular Cloning: A Laboratory Manual (2nd ed.), Cold Spring Harbor Laboratory Press, Sambrook,

30 Fritsch and Maniatis eds. (1989).

Transgenic, i.e., transduced, endothelial cells stably incorporating and expressing heterologous DNA or RNA and therapeutic uses therefore, are described in Mulligan et al. (U.S. Pat. No. 5,674,722). In particular, retroviral vectors have been used to stably transduce endothelial cells with genetic material which includes
5 genetic material encoding a polypeptide or protein of interest not normally expressed at biologically or therapeutically significant levels in endothelial cells. The genetic material introduced in this manner can also include genetic material encoding a dominant selectable marker, such as antibiotic or herbicide resistance. Genetic material including DNA encoding a polypeptide of interest alone, such as a
10 Factor VIII protein, or DNA encoding a polypeptide of interest and a dominant selectable marker can be introduced into cultured endothelial cells. Expression of these genes by the endothelial cells into which they have been incorporated (i.e., endothelial cells transduced by the use of retroviral vectors) has also been demonstrated.

15 Endothelial cells transduced *in vitro* with the genetic material can then be transplanted using one of a variety of known methods. Such methods include, but are not limited to, the transplantation of synthetic vessels or prosthetic valves lined with transduced endothelial cells or the transplantation of a device or matrix designed to house transduced endothelial cells.

20 To administer the EC containing the desired gene, the cells may simply be introduced into the bloodstream of the patient by conventional means, such as of intravenous infusion over a period of time.

Endothelial cells which have been transduced *in vitro* are particularly useful for improving prosthetic implants (e.g., vessels made of synthetic materials such as
25 Dacron®, Gortex® and other plastics or metal-plastic laminates), including shunts, stents and grafts, which are used in vascular reconstructive surgery. For example, prosthetic arterial grafts are often used to replace diseased arteries which perfuse vital organs or limbs. However, the currently available grafts are usually made of synthetic material and are subject to many complications, the worst of which is a
30 high rate of thrombosis or occlusion. Animal studies suggest that lining the graft

with autologous endothelial cells prior to implantation may decrease, but not prevent, graft reocclusion with its attendant morbid consequences.

However, endothelial cells can be modified according to the method of the present invention in a way that improves their performance in the context of an
5 implanted graft or provides a means for local drug delivery. Examples include local delivery to the interior of the lumen of antirestenotic, antiproliferative, or thrombolytic agents to prevent intraluminal clot formation, secretion of an inhibitor of smooth muscle proliferation to prevent luminal stenosis due to smooth muscle hypertrophy, and expression and/or secretion of an endothelial cell mitogen or
10 autocrine factor to stimulate endothelial cell proliferation and improve the extent or duration of the endothelial cell lining of the graft lumen. The latter agents are termed "biocompatibilization" agents (polypeptides or proteins).

For a similar application, endothelial cells of the present invention can also be used to cover the surface of prosthetic heart valves to decrease the risk of the
15 formation of emboli by making the valve surface less thrombogenic.

Endothelial cells transduced by the method of the subject invention or a vascular implant lined with transduced endothelial cells can also be used to provide constitutive synthesis and delivery of polypeptides or proteins, which are useful in prevention or treatment of disease, such as Factor VIII proteins, to treat hemophilia.
20 In this way, the polypeptide is secreted directly into the bloodstream of the individual, from circulating cells, over an extended period of time. Currently available methods, in contrast, involve parenteral administration of the desired polypeptide.

In addition, there is no need for extensive purification of the polypeptide
25 before it is administered to an individual, as is generally necessary with an isolated polypeptide (e.g., insulin). Endothelial cells modified according to the present invention produce the polypeptide hormone as it would normally be produced.

Another advantage to the use of genetically engineered endothelial cells is that one can target the delivery of therapeutic levels of a secreted product to a
30 specific organ or limb. For example, a vascular implant lined with endothelial cells

transduced *in vitro* can be grafted into a specific organ or limb; or the endothelial cells of a particular limb, organ or vessel can be transduced *in vivo*. The secreted product of the transduced endothelial cells will be delivered in high concentrations to the perfused tissue, thereby achieving a desired effect to a targeted anatomical location. This product will then be diluted to nontherapeutic levels in the venous circulation during its return to the heart.

Another important advantage of the delivery system of this invention is that because it is a continuous delivery system, the short half lives of hormone polypeptides is not a limitation. For example, the half life of human growth hormone (HGH) is approximately 19 minutes and parathyroid hormone, approximately 2½ to 5 minutes.

The invention will be further described by reference to the following detailed examples.

15

Example 1

Culture of Endothelial Cells from Peripheral Blood

One hundred ml of fresh venous blood anti-coagulated with either heparin or citrate was diluted 1:2 with Hanks' Balanced Salt Solution (HBSS) containing 1 mM EDTA and 0.5% bovine serum albumin, carefully layered on an equal volume of Histopaque®-1077 (Sigma Chemical Co.) without disruption of the interface, and centrifuged at 400 × G for 30 minutes at room temperature. The layer containing the mononuclear cells was collected, and the cells were washed 3 times by centrifuging at 250 × G for 10 minutes, using the previously described MEC media (Gupta et al., Exp. Cell Res., 230, 244-251 (1997)) modified to contain 10% human male serum, to exclude the ECGS and comprising 1 µg/ml hydrocortisone. Washing with this culture medium is preferred over washing with buffer because the medium maintains higher levels of cell viability.

The buffy coat mononuclear cells were resuspended in EGM®-2 medium (Clonetics), and all buffy coat mononuclear cells from 100 ml of starting blood were placed into one well of a 6-well plate coated with collagen I (Becton-Dickinson).

This is defined as day #1, on which the plate is placed into an incubator at a controlled temperature (37°C) and humidified environment (5% CO₂). The EGM[®]-2 medium is changed daily. On day #2, most of the buffy coat mononuclear cells have remained unattached and they and cell debris were removed at the time of culture medium change. This left on the bottom of the culture well a small number of cells that have typical endothelial cell morphology and which stain positively with anti-endothelial monoclonal antibody P1H12. Typically, there were about 20-30 such cells identifiable on day #2, along with 100-200 other mononuclear cells.

After good cell growth was established (typically at about 20 days, when cells may or may not be confluent), they are passed into a fibronectin/gelatin-coated T25 flask, and then into fibronectin/gelatin-coated T75 flasks. To pass the cells, the wells are washed twice with calcium-free HBSS and then once with 0.5 X trypsin (i.e., 50% of the concentration for the Gibco BRL trypsin product recommended for cell lifting, because it is less harsh than the full-strength product) plus 1 mM EDTA. Cells detach after 2 minutes. Trypsin is neutralized by adding an equal volume of human serum or MEC media containing 20% human serum. Cells were collected from the solution by centrifugation at 250 × G for 5 minutes, and resuspended in EGM[®]-2 medium for continued culture.

Figure 1 is plot of the outgrowth of endothelial cells expanded using this method. The mean ± SD of five different culture experiments using five different blood donors was plotted up until culture passage 5, after which four different cultures were plotted. Error bars are not evident on the graph because of the high reproducibility of the method. As shown, the method of the invention can result in an outgrowth, and a 10¹⁸ fold expansion, of endothelial cells far greater than the 20-30 endothelial cells identifiable on culture day #2 (the day the unattached mononuclear cells and cell debris from the buffy coat are removed and any EC can be first identified).

It should be noted that this fold expansion is not a maximal expansion. It is just the degree of expansion allowed before cells are used for further experiments.

The maximal cap (if any) in expandability of these cells, therefore, has not yet been established or reached in these experiments.

The outgrowth cells have the characteristics of quiescent, microvascular EC. They have typical "cobblestone" endothelial morphology; they exhibit positive
5 expression for von Willebrand Factor, P1H12 antigen, ICAM1, $\alpha_v\beta_3$,
 β_2 -microglobulin, thrombomodulin, flk-1 (a VEGF receptor), and CD34; they are uniformly CD36-positive; they take up acetylated-LDL; they do not constitutively express tissue factor or VCAM, but do express both upon appropriate stimulation.

10

Example 2

Cryopreservation of Cultured Endothelial Cells

Vigorous EC outgrowth can be obtained even after cryopreservation. This is important because the utility of the present method is greater if cultured EC can be cryopreserved for later use.

15

To show this, EC were cryopreserved after their outgrowth had reached the capacity of two T75 flasks. To do this, the cells were detached as above and washed twice with HBSS. Cells were resuspended in 100% fetal calf serum at a concentration of one-million cells per 950 μ l serum, and then 50 μ l dimethylsulfoxide was added. The suspension was mixed quickly and placed on ice
20 for 5 minutes. Afterward, the cells were stored at -70°C overnight, and they were then transferred to liquid nitrogen.

After six weeks of storage, the cells were thawed and one million cells were plated onto T75 flask with excess (20 ml) of EGM[®]-2 medium. Medium was changed again four hours after initial plating. The cells were then grown as in
25 Example 1.

The resulting growth is similar to that of EC that were not subjected to cryopreservation, as shown in Figure 2. Again, data are shown as mean $\pm$ SD for three experiments. Cell count at origin indicates the number of cells thawed and started with. These data suggest that the endothelial outgrowth cells expanded from
30 blood resume their vigorous growth rate after cryopreservation.

Example 3

Study of Origin of CEC in Blood

The origin of the CEC identifiable in fresh blood is not known, nor is the relationship between CEC and the endothelial outgrowth from cultured blood. To answer this, 4 adults were studied who had undergone allogeneic marrow transplantation (for malignancy) 5-20 months earlier using opposite-gender donors (three female donors to three male recipients and one male donor to one female recipient). All 4 subjects are disease-free and have peripheral blood and/or marrow aspirates that are 100% donor by RFLP; two also had marrow cytogenetics showing 100% donor.

The CEC in fresh blood, plus the peripheral blood endothelial outgrowth from buffy coat mononuclear cells cultured with endothelial growth factors was studied using the culture conditions of Ex. 1. MAb P1H12 was used to identify cells as being endothelial, and fluorescence *in situ* hybridization was used to identify cells as having XX or XY genotype.

CEC in fresh blood were almost exclusively of recipient genotype (95, 100, 100, 88%), arguing for their origin from the vessel wall. However, the blood endothelial outgrowth at 26-28 days (after 100-fold expansion) in culture was mostly donor genotype (82, 86, 77, 85%), revealing predominant origination from a marrow-derived cell. Remarkably, endothelial outgrowth at only 9 days (after 5-fold expansion) in contemporaneous culture was still largely recipient genotype (78, 88%). These data indicate that the CEC detected in fresh blood are derived from vessel wall and have some, but limited, expansion potential. Conversely, blood contains transplantable marrow-derived cells that take longer to expand but have greater proliferative potential. Thus, the CEC comprise a more mature and differentiated cell population, while there also are circulating marrow-derived endothelial-generating cells that comprise a more primitive progenitor population (putative angioblasts).

Example 4**Transfection of CEC with Factor VIII Expression Vectors**

Endothelial cells were expanded from blood, as described in Example 1. For these transfection experiments, they were used after 4-5 passages. They were plated in a 6 well plate at density of 2×10^5 /well, and allowed to attach overnight. Then, cells in different wells were transfected by lipofection with one of four vectors as summarized in Table 1, below.

Table 1. Plasmid Vectors for CEC Transfection

- | | |
|----|---|
| 10 | <p>A. Two vectors from Octagen with B-domainless FVIII replaced with green fluorescent protein (GFP):</p> <p style="margin-left: 20px;">#1 HSQ/eGFP/ReNeo</p> <p style="margin-left: 20px;">#2 HSQ/eGFP/CP</p> |
| 15 | <p>B. Control vector from Octagen with GFP but no factor VIII:</p> <p style="margin-left: 20px;">#3 pEGFP-N1</p> <p>C. Vector constructed at University of Minnesota, Department of Medicine with the FVIII/GFP from Octagen inserted between XhoI and NotI sites in commercial vector pcDNA3.1(-):</p> <p style="margin-left: 20px;">#4 pcHSQ/eGFP</p> |

20 All the transfection vectors contain the neomycin resistance gene (ReNeo) as a selection marker.

Vectors 1, 2 and 4 contain an HSQ/eGFP insert which codes for a fVIII-dGFP fusion protein. This insert comprises DNA that codes in order of sequence, 1) the human fVIII human activation peptide, 2) the human fVIII A1 domain, 3) the human fVIII A2 domain, 4) the first five amino acids of the SQ B domain linker peptide, 5) the enhanced green fluorescent protein (eGFP), 6) the last nine amino acids of the SQ linker peptide, 7) the human fVIII light chain activation peptide, 8) the human fVIII A3 domain, 9) the human fVIII C1 domain, and 10) the human fVIII C2 domain.

30 The DNA sequence of HSQ/eGFP is depicted in Fig. 3 (SEQ ID NO:1). The DNA sequence contains 5094 bases, encoding the peptides listed on Table 2.

Table 2. HSQ/eGFP Peptides SEQ ID NO:1

- | | |
|----|--|
| 35 | <p>1-57 signal peptide</p> <p>58-1173 A1 domain</p> <p>1174-2274 A2 domain</p> |
|----|--|

2275-2292 SQ sequence (first part)
 2293-3012 eGFP sequence
 3013-3039 SQ sequence (second part)
 3040-3162 light chain activation peptide
 5 3163-4152 A3 domain
 4153-4611 C1 domain
 4612-5091 C2 domain
 5092-5094 stop codon

10 ReNeo is a mammalian expression vector (non-proprietary). The second
 plasmid, HSQ/eGFP/CP, contained the same fVIII construct but the vector is the
 same one used to express eGFP (see below). The eGFP protein was removed from
 p-EGFP-N1 and replaced with the HSQ/eGFP construct. The third plasmid is p-
 EGFP-N1 from Clontech, which expresses only eGFP.

15 The transfection protocol was based on the instructions supplied by Life
 Technologies Co. with the lipofectamine. Briefly, 2 micrograms of plasmid DNA
 was mixed gently with 15 microliters of lipofectamine in 200 microliters of E-
 STIM[®] basal medium (Becton Dickinson) without antibiotics. The mixture was
 allowed to stand at room temperature for 30 minutes. Expanded endothelial cells in
 20 6-well plates were washed twice with E-STIM[®] basal medium before adding the
 DNA-liposome complex. Dropwise, 1.8 ml of E-STIM[®] was added to the DNA-
 liposome complex, then the mixture was overlaid on the endothelial cells. After
 5 hours incubation, the DNA-liposome mixture was replaced with the endothelial
 culture medium of Example 1. Then, 2 days after transfection, the cells from one
 25 culture well were subcultured into one 10 cm dish in endothelial culture medium
 containing 50-100 µg/ml of Geneticin (Life Technologies Co.). This selection
 medium was changed every other day.

Transfection of endothelial cells derived from the blood of two donors has
 been accomplished to date. The number of endothelial colonies (derived from the
 30 transfection of greater than 2×10^5 cells) was checked on day 20 in selection
 medium for two donors, as shown on Table 3.

Table 3. Number of Colonies After 20 Days in Selection Medium

	<u>Vector #</u>	<u>Donor 1</u>	<u>Donor 2</u>
	1	3	9
	2	2	8
5	3	4	7
	4	1	8

On day 20, fluorescence microscopy showed all selected colonies to be positive for green fluorescent protein (GFP). On day 20, the colonies were pooled and transferred to the regular endothelial growth condition.

In the past, many basic and applied studies had to be performed on endothelial cells from other animal species because existing culture techniques permitted only restricted proliferation of human endothelial cells. The present method accomplishes at least a 10^{18} -fold expansion of circulating human EC. This method will permit peripheral blood from living donors to be used for the generation of large numbers of cultured endothelial cells. Thus, problems of human pathology involving the endothelium now can be approached directly by employing a human endothelial cell model. In addition, expanded mammalian EC should prove valuable for various clinical applications, such as *in vitro* testing of vasoactive agents, the coating of artificial graft materials and gene therapy designed to treat vascular pathologies and genetic disorders, for local drug delivery and enhanced biocompatibility.

Example 5

Use of Endothelial Cell Culture Methodology in Disease Diagnostics

Currently, there is no direct or non-invasive method to examine a patient's endothelium to assess it for either acquired or genetic defects that contribute to disease. A 25 year old male patient had been evaluated for the presence of a hypercoagulability disorder. The patient had a history of multiple thromboses, and a family history suggesting a familial pre-disposition to thrombosis. No positive diagnosis was obtained after extensive evaluation for known genetic and

predisposing causes. Therefore, the patient was evaluated for a defect in thrombomodulin. Thrombomodulin is an enzyme that is expressed on the endothelial cell surface and is an anti-thrombotic defense. A patient having a mutation in thrombomodulin that leads to poor function would have an increased risk for thrombosis.

Endothelial cells were expanded from the blood of the patient and three normal control donors using the present method. Thrombomodulin activity was assessed in a standard biochemical assay. Briefly, a known number of endothelial cells is incubated with bovine protein C (100 μ L per well of a 1 μ M solution) and human recombinant thrombin (10 μ L of 10 nM). It is incubated at 37°C for 2.5 hours. The reaction is terminated by adding EDTA and I-2581 (Chromogenix AB, Molndal, Sweden), and the chromatographic substrate S-2366 (DiaPharma Group, Inc., Franklin, OH) is added. Supernatant absorbance is read at 405 nm.

The results are given in OD units per minute per 50,000 endothelial cells. The value for 18 samples from the three normal donors was 53 ± 7 , while the value for six samples from the patient was 41 ± 5 , a significant difference ($p=0.0034$). Thus, the expanded endothelial cells from the patient had a significantly lower thrombomodulin activity than those of control patients. Therefore, a genetic mutation in thrombomodulin resulting in a thrombomodulin deficiency may be the underlying basis for this patient's disorder. Molecular biological analyses are employed to confirm the genetic basis for the thrombomodulin deficiency.

Example 6

Stable Transfection, Selection and Expansion

The plasmid pLE (Clontech), which contains sequence for enhanced green fluorescent protein (eGFP), was transfected into line PA317. Virus from one of the resulting cloned transfectants, pLE9, was employed to infect endothelial cells.

A culture of blood buffy coat mononuclear cells was set up according to the method of the invention. When cells had reached passage 4 and about a 10^6 fold expansion, they were seeded at 10^5 cells/dish, with the 10 cm dishes coated with 6

WO 00/32750

PCT/US99/28033

microgram/cm² of type 1 collagen and 250 microgram/dish of fibronectin in 1% gelatin. After two days of growth, the cells were exposed to conditioned medium from the packaging cell line in combination with 4 microgram/ml of polybrene to dishes that were 70% confluent. After three days exposure, cells were split 1:2 into
5 a 10 cm dish (coated as above), and new culture medium added. Then they were subjected to selection in media containing G418 for 19 days.

As shown in Figure 5, cultured endothelial cells prepared according to the method of the invention can survive chemical selection and be expanded thereafter, and be stably transfected to express a foreign (non-selected) gene. Moreover, the
10 outgrowth cells have preserved normal endothelial cell morphology.

Example 7

Transfection of CEC with Human Factor VIII (hFVIII) Expression Vectors

Materials and Methods

15 Transfection

Ten cm dishes and T75 flasks were coated with 6 µg/ml of type I collagen and 50 µg/ml of fibronectin. Passage 4 or 5 outgrowth endothelial cells were obtained as described in Example 1. They were seeded into the dish or flasks and allowed to grow to 40% confluence. Then the cells were transfected with hFVIII
20 vectors (Table 4) using transfection reagent Eugene6 (Roche Molecular Biochemicals) following the manufacturer's protocol. Briefly, Eugene6 was added to each vector at a ratio of 2.5:1. Ten µg of DNA was used for transfection of cells in 10 cm dishes; and 15 µg of DNA was used for cells in T75 flasks.

The Eugene6-DNA mixture was added to the cells in media containing 10%
25 human serum and incubated for 3 days. Cells reached 100% confluence after 3 days.

Table 4. Plasmid Vectors

pTracerHSQ: has HSQ (the B-domainless hFVIII without the eGFP insert)
cloned into pTracer-CMV (Invitrogen)

pcF8G: has HSQ/eGFP (eGFP replaces the B domain of hFVIII) cloned into
5 plasmid pCDNA 3.1 (Invitrogen)

pcHSQ: has HSQ cloned into plasmid pCDNA 3.1 (Invitrogen)

pTracerCMV: obtained from Invitrogen

HSQ/GFP/ReNeo: has HSQ/eGFP cloned into pRENeo (Biogen)

Transient expression

10 At the end of 3 days, the conditioned media from T75 flasks were collected
and centrifuged. The presence of hFVIII in the supernatant was detected using an
ELISA kit (American Diagnostics). Cells from T75 flask were harvested, lysed and
processed (RT-PCR, Titan One Tube RT-PCR System, Roche Molecular
Biochemicals) for detection of hFVIII mRNA. A reverse transcriptase reaction was
15 conducted at 50°C for 45 minutes, then the samples were subject to 94°C, for 2
minutes; 10 cycles of 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1
minute; 20 cycles of 94°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1
minute, plus cycle elongation of 5 seconds for each cycle elongation at 72°C for 7
minutes. The primer pair employed was hFVIIC-S (5' GCC CTT TTC TTG GAT
20 CAA GGT GG3'; SEQ ID NO: 3) and hFVIIC-AS (5' CTC CCT GAG TAG TTA
CTC CTG TG3'; SEQ ID NO: 4).

Results

Three days after infection, hFVIII was detected in the supernatant of
transfected cells (Figure 6) but not in control cells. Figure 7 shows the results of a
25 RT-PCR analysis of RNA from those cells. The expression of hFVIII mRNA is
increased in transfected cells relative to control cells.

At the end of 3 days, cells from 10 cm dishes were split into 2 10 cm dishes
coated with 6 µg/ml of type I collagen and 50 µg/ml of fibronectin. The following
day, cells were exposed to selection medium EGM-2 (Clonectis) containing 50 to
30 100 µg/ml of G418 or 50 µg/ml of Zeocin or both (Zeocin was used for selection of

WO 00/32750

PCT/US99/28033

pTracerHSQ; G418 for pcFVIII_G and HSQ/GFP/ReNeo; and both Zeocin and G418 were used for cells subjected to cotransfection). Stable expression of hFVIII is detected in these cells after chemical selection.

- 5 All patents, patent applications and publications cited hereinabove, are incorporated by reference herein. While a preferred embodiment of the present invention has been described, it should be understood that various changes, adaptations and modifications may be made therein without departing from the spirit of the invention and the scope of the appended claims.

10

WHAT IS CLAIMED IS:

1. A process for expanding the population of endothelial cells obtained from peripheral blood comprising culturing, in contact with a collagen I-coated surface, buffy coat cells obtained from peripheral mammalian blood in the presence of a cell culture medium containing an effective amount of vascular endothelial growth factor (VEGF), and which medium is free of bovine brain extract, so as to expand the population of endothelial cells in said buffy coat cells.
2. The process of claim 1 wherein the blood is human blood.
3. The process of claim 1 wherein said cell culture medium comprises heparin, dextran sulfate or mixtures thereof.
4. The process of claim 1 wherein the buffy coat cells are obtained by washing cells from a buffy coat layer obtained from human blood in cell culture medium comprising 20% human male serum.
5. The process of claim 1 wherein the cell culture medium comprises human basic fibroblast growth factor.
6. The process of claim 1 or 5 wherein the cell culture medium comprises insulin-like growth factor.
7. The process of claim 1 or 5 wherein the cell culture medium contains human epidermal growth factor.
8. The process of claim 1 wherein the cell culture medium comprises about 0.5-10 vol-% fetal bovine serum and about 95-99.5 vol-% of a cell culture medium.

9. The process of claim 1 wherein the cultured cells are trypsinized at about 10^3 -fold expansion, collected by centrifugation, resuspended in cell culture medium, and subjected to continued culture in contact with a fibronectin/gelatin-coated surface.
- 5 10. The process of claim 1 wherein the cultured cells are subjected to cryopreservation.
11. The process of claim 10 wherein the cells are frozen in a cryopreservation medium comprising fetal calf serum containing an effective amount of dimethylsulfoxide.
- 10 12. The process of claim 10 or 11 wherein the cryopreserved cells are thawed and culturing is resumed in said cell culture medium.
- 15 13. The process of claim 1 wherein the expanded population comprises microvascular endothelial cells.
14. The process of claim 13 wherein the microvascular endothelial cells are $CD34^+$, $CD36^+$ and express the P1H1 antigen.
- 20 15. A transgenic mammalian endothelial cell comprising an isolated DNA sequence encoding a recombinant Factor VIII protein.
- 25 16. The transgenic endothelial cell of claim 15 wherein the Factor VIII protein is a hybrid human/porcine protein.
17. A transgenic endothelial cell which is prepared by a process comprising stably transforming a population of circulating human endothelial cells outgrown from blood with a vector comprising an isolated DNA sequence encoding a preselected protein operably linked to a promoter functional in human endothelial cells.
- 30

18. The transgenic endothelial cell of claim 17 wherein the DNA sequence encodes a Factor VIII protein.
19. The transgenic endothelial cell of claim 17 wherein the DNA sequence comprises SEQ ID NO:1 or SEQ ID NO:2.
20. The transgenic endothelial cell of claim 18 wherein the Factor VIII protein is a hybrid human/porcine protein.
21. The transgenic endothelial cell of claim 18 wherein the Factor VIII protein is a chimeric human protein.
22. The transgenic endothelial cell of claim 17 wherein the DNA sequence further comprises a selectable marker gene or a reporter gene.
23. The transgenic endothelial cell of claim 17 which is prepared by lipofection.
24. A pharmaceutical composition comprising a population of the transgenic endothelial cells of claim 15 or 17 in combination with a pharmaceutically acceptable carrier.
25. A method of treating hemophilia comprising introducing an amount of the endothelial cells of claim 15 or 17 into the bloodstream of a mammal afflicted with hemophilia so that an effective amount of a Factor VIII protein is secreted into the bloodstream of said mammal.
26. The method of claim 25 wherein the mammal is a human.
27. An implantable medical prosthetic device comprising a surface coated with the endothelial cells of claim 17 wherein the protein is expressed in an amount effective to increase the biocompatibility of said device upon implantation into a mammal.

28. The device of claim 27 which comprises a plastic surface, a metal surface or a laminate surface.
29. The device of claim 28 which is a vascular graft.
- 5 30. The device of claim 28 which is a shunt or a stent.
31. The device of claim 28 which is a heart valve.
- 10 32. The device of claim 28 which is a controlled drug release depot.
- 15 33. A diagnostic method comprising:
detecting or determining whether an expanded population of endothelial cells obtained from a test mammal has an acquired or genetic indication or disease relative to a control expanded population of endothelial cells obtained from a mammal not at risk of the acquired or genetic indication or disease, wherein the expanded cells are obtained by culturing, in contact with a collagen I-coated surface, buffy coat cells obtained from peripheral mammalian blood in the presence of a cell culture medium containing an effective amount of vascular endothelial growth factor (VEGF), which medium is free of bovine brain extract, so as to yield an expanded population of endothelial cells.
- 20 34. The method of claim 33 wherein the indication or disease is a clotting disorder.
- 25 35. The method of claim 33 wherein the indication or disease is associated with a reduction in the activity of an enzyme.
- 30 36. The method of claim 33 wherein the indication or disease is acquired.
37. A population of mammalian endothelial cells prepared by the method of claim 1.

38. A cultured, quiescent mammalian endothelial cell.
39. A population of cultured, quiescent mammalian endothelial cells.
- 5 40. The population of cells of claim 37 or 39 at least a portion of which is stably transformed with a vector comprising an isolated DNA sequence encoding a preselected protein operably linked to a promoter functional in mammalian endothelial cells.
- 10 41. The cell of claim 38 which is stably transformed with a vector comprising an isolated DNA sequence encoding a preselected protein operably linked to a promoter functional in the cell.




Fig 1

CEC Expansion after Cryopreservation




Fig 2

3 / 14
Fig. 3

HFVIII/SQ/egfp, HF8SQgfp.gp, original HB- with B-domain SQ insert based on Lind et al, Eur J Biochem, 232, p21 primer sequence, containing eGFP protein sequence\$

ATGCAAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGCGAT
TCTGCTTT
AGTGCCACCAGAAGATACTACCTGGGTGCAGTGGAAGTGTGTCATGG
GACTATATGCAAAGTGA
TCTCGGTGAGCTGCCTGTGGACGCAAGATTTCTCCTAGAGTGCC
AAAATCTTTTCCATTCAA
CACCTCAGTCGTGTACAAAAGACTCTGTTTGTAGAATTCACGGTT
CACCTTTTCAACATCGC
TAAGCCAAGGCCACCCTGGATGGGTCTGCTAGGTCCTACCATCCA
GGCTGAGGTTTATGATAC
AGTGGTCATTACACTTAAGAACATGGCTTCCCATCCTGTCAGTCTT
CATGCTGTTGGTGTATCC
TACTGGAAAGCTTCTGAGGGAGCTGAATATGATGATCAGACCAGT
CAAAGGGAGAAAGAAGA
TGATAAAGTCTTCCCTGGTGGAAGCCATACATATGTCTGGCAGGTC
CTGAAAGAGAATGGTCC
AATGGCCTCTGACCCACTGTGCCTTACCTACTCATATCTTTCTCAT
GTGGACCTGGTAAAAGA
CTTGAATTCAGGCCTCATTGGAGCCCTACTAGTATGTAGAGAAGG
GAGTCTGGCCAAGGAAA
AGACACAGACCTTGCACAAATTTATACTACTTTTTGCTGTATTTGA
TGAAGGGAAAAGTTGGC
ACTCAGAAACAAAGAACTCCTTGATGCAGGATAGGGATGCTGCAT
CTGCTCGGGCCTGGCCT
AAAATGCACACAGTCAATGGTTATGTAAACAGGTCTCTGCCAGGT
CTGATTGGATGCCACAG
GAAATCAGTCTATTGGCATGTGATTGGAATGGGCACCACTCCTGA
AGTGCACTCAATATTCCT
CGAAGGTCACACATTTCTTGTGAGGAACCATCGCCAGGCGTCCTT
GGAAATCTCGCCAATAAC
TTTCCTTACTGCTCAAACACTCTTGATGGACCTTGGACAGTTTCTA
CTGTTTTGTGTCATATCTCT
TCCCACCAACATGATGGCATGGAAGCTTATGTCAAAGTAGACAGC
TGTCCAGAGGAACCCCA
ACTACGAATGAAAAATAATGAAGAAGCGGAAGACTATGATGATGA
TCTTACTGATTCTGAAA
TGGATGTGGTCAGGTTTGATGATGACAACTCTCCTTCCTTTATCCA
AATTCGCTCAGTTGCCA
AGAAGCATCCTAAACTTGGGTACATTACATTGCTGCTGAAGAGG
AGGACTGGGACTATGCT
CCCTTAGTCCTCGCCCCGATGACAGAAGTTATAAAAGTCAATATT
TGAACAATGGCCCTCAG
CGGATTGGTAGGAAGTACAAAAAGTCCGATTTATGGCATAACACA
GATGAAACCTTTAAGAC
TCGTGAAGCTATTTCAGCATGAATCAGGAATCTTGGGACCTTTACTT
TATGGGGAAGTTGGAGA
CACACTGTTGATTATATTTAAGAATCAAGCAAGCAGACCATATAA
CATCTACCCTCACGGAAT
CACTGATGTCCGTCCTTTGTATTCAAGGAGATTACCAAAGGTGTA
AAACATTTGAAGGATTT
TCCAATTCTGCCAGGAGAAATATTCAAATATAAATGGACAGTGAC

WO 00/32750

PCT/US99/28033

4 / 14
Fig. 1 (cont-2)

TGTAGAAGATGGGCAA
CTAAATCAGATCCTCGGTGCCTGACCCGCTATTACTCTAGTTTCGT
TAATATGGAGAGAGATC
TAGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAATC
TGATAGATCAAAGAGGAA
ACCAGATAATGTCAGACAAGAGGAATGTCATCCTGTTTTCTGTATT
TGATGAGAACCGAAGCT
GGTACCTCACAGAGAATATACAACGCTTTCTCCCAATCCAGCTG
GAGTGCAGCTTGAGGATC
CAGAGTTCCAAGCCTCCAACATCATGCACAGCATCAATGGCTATG
TTTTTGATAGTTTGCACT
TGTCAGTTTGTGTCATGAGGTGGCATACTGGTACATTCTAAGCAT
TGGAGCACAGACTGACT
TCCTTTCTGTCTTCTCTCTGGATATACCTTCAAACACAAAATGGTC
TATGAAGACACACTCAC
CCTATTCCCATTTCTCAGGAGAACTGTCTTCATGTCGATGGAAAAC
CCAGGTCTATGGATTCT
GGGGTGCCACAACCTCAGACTTTCGGAACAGAGGCATGACCGCCTT
ACTGAAGGTTTCTAGTTG
TGACAAGAACTGGTGATTATTACGAGGACAGTTATGAAGATAT
TTCAGCATACTTGCTGAG
TAAAAACAATGCCATTGAACCTAGG
AGCTTCTCTCAGAATATGGTGAGCAAGGGCGAGGAGC
TGTTACCGG GGTGGTGCCC
ATCCTGGTCG AGCTGGACGG CGACGTAAAC GGCCACAAGT
TCAGCGTGTC CGGCGAGGGC
GAGGGCGATG CCACCTACGG CAAGCTGACC CTGAAGTTCA
TCTGCACCAC CGGCAAGCTG
CCCGTGCCCT GGCCACCCCT CGTGACCACC CTGACCTACG
GCGTGCACTG CTTGAGCCGC
TACCCCGACC ACATGAAGCA GCACGACTTC TTCAAGTCCG
CCATGCCCCG AGGCTACGTC
CAGGAGCGCA CCATCTTCTT CAAGGACGAC GGCAACTACA
AGACCCGCGC CGAGGTGAAG
TTCGAGGGCG ACACCCTGGT GAACCGCATC GAGCTGAAGG
GCATCGACTT CAAGGAGGAC
GGCAACATCC TGGGGCACAA GCTGGAGTAC AACTACAACA
GCCACAACGT CTATATCATG
GCCGACAAGC AGAAGAACGG CATCAAGGTG AACTTCAAGA
TCCGCCACAA CATCGAGGAC
GGCAGCGTGC AGCTCGCCGA CCACTACCAG CAGAACACCC
CCATCGGCGA CGGCCCCGTG
CTGCTGCCCC ACAACCACTA CCTGAGCACC CAGTCCGCCC
TGAGCAAAGA CCCCACGAG
AAGCGCGATC ACATGGTCCT GCTGGAGTTC GTGACCGCCG
CCGGGATCAC TCTCGGCATG
GACGAGCTGT ACAAGTATCCACCAGTCTTGAAACGCCATCAACGG
GAAATAACTCGTACTACTCT
TCAGTCAGATCAAGAGG
AAATTGACTATGATGATACCATATCAGTTGAAATGAAGAAGGAAG
ATTTTGACATTTATGATG
AGGATGAAATCAGAGCCCCCGCAGCTTTCAAAGAAAACACGAC
ACTATTTTATGCTGCA
GTGGAGAGGCTCTGGGATTATGGGATGAGTAGCTCCCCACATGTT

Fig. 3 . cont. - 3) ⁵ / 14

CTAAGAAACAGGGCTCA
GAGTGGCAGTGTCCCTCAGTTCAAGAAAGTTGTTTTCCAGGAATTT
ACTGATGGCTCCTTTAC
TCAGCCCTTATACCGTGGAGAACTAAATGAACATTTGGGACTCCT
GGGGCCATATATAAGAG
CAGAAGTTGAAGATAATATCATGGTAACTTTCAGAAATCAGGCCT
CTCGTCCCTATTCTCT
ATTCTAGCCTTATTTCTTATGAGGAAGATCAGAGGCAAGGAGCAG
AACCTAGAAAAAACTTT
GTCAAGCCTAATGAAACCAAACTTACTTTTGGAAGTGCAACAT
CATATGGCACCCTAA
AGATGAGTTTGACTGCAAAGCCTGGGCTTATTTCTCTGATGTTGAC
CTGGAAAAAGATGTGCA
CTCAGGCCTGATTGGACCCCTTCTGGTCTGCCACACTAACACACTG
AACCTGCTCATGGGAG
ACAAGTGACAGTACAGGAATTTGCTCTGTTTTTCACCATCTTTGAT
GAGACCAAAAGCTGGTA
CTTCACTGAAAATATGGAAAGAACTGCAGGGCTCCCTGCAATAT
CCAGATGGAAGATCCCA
CTTTTAAAGAGAATTATCGCTTCCATGCAATCAATGGCTACATAAT
GGATACACTACCTGGCT
TAGTAATGGCTCAGGATCAAAGGATTCTGATGGTATCTGCTCAGCA
TGGGCAGCAATGAAAC
ATCCATTCTATTCAATTCAGTGGACATGTGTTCACTGTACGAAAAA
AAGAGGAGTATAAAATG
GCACTGTACAATCTCTATCCAGGTGTTTTTGAGACAGTGGAAATGT
TACCATCCAAAGCTGGA
ATTTGGCGGGTGGAAATGCCTTATTGGCGAGCATCTACATGCTGGG
ATGAGCACACTTTTTCTG
GTGTACAGCAATAAGTGTCACTCCCTGGGAATGGCTTCTGGA
CACATTAGAGATTTTCAG
ATTACAGCTTCAGGACAATATGGACAGTGGGCCCCAAAGCTGGCC
AGACTTCATTATTCCGG
ATCAATCAATGCCTGGAGCACCAGGAGCCCTTTTCTTGGATCAA
GGTGGATCTGTTGGCACC
AATGATTATTCACGGCATCAAGACCCAGGGTGCCCGTCAGAAGTT
CTCCAGCCTCTACATCTC
TCAGTTTATCATCATGTATAGTCTTGATGGGAAGAAGTGGCAGACT
TATCGAGGAAATTCCAC
TGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTCTCTGGGATA
AAACACAATATTTTAA
CCCTCCAATTATTGCTCGATACATCCGTTTGCACCCAACTCATTAT
AGCATTTCGCAGCACTCTT
CGCATGGAGTTGATGGGCTGTGATTTAAATAGTTGCAGCATGCCA
TTGGGAATGGAGAGTAA
AGCAATATCAGATGCACAGATTACTGCTTCATCCTACTTTACCAAT
ATGTTTGCCACCTGGTC
TCCTTCAAAGCTCGACTTCACCTCCAAGGGAGGAGTAATGCCTG
GAGACCTCAGGTGAATA
ATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGAAAG
TCACAGGAGTAACTACT
CAGGGAGTAAAATCTCTGCTTACCAGCATGTATGTGAAGGAGTTC
CTCATCTCCAGCAGTCAA
GATGGCCATCAGTGGACTCTTTTTTTTCAGAATGGCAAAGTAAAG

WO 00/32750

PCT/US99/28033

6 / 14

Fig. 3 (continued)

GTTTTTCAGGGAAATCAA
GACTCCTTCACACCTGTGGTGAACCTCTCTAGACCCACCGTTACTGA
CTCGCTACCTTCGAATT
CACCCCAGAGTTGGGTGCACCAGATTGCCCTGAGGATGGAGGTT
CTGGGCTGCGAGGCACA
GGACCTCTACTGA*

HSQRENBO


GAATTCCGGAATTCCAGCTTGCTGTGGAATGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGG
CTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAG
TCCCCAGGCTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCATAG
TCCCGCCCCCTAACTCCGCCCCATCCCGCCCCCTAACTCCGCCCCAGTTCCGCCCCATTCTCCGCCCCA
TGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAG
AAGTAGTGAGGAGGCTTTTTTGGAGGGGTCTCCTCGTATAGAACTCGGACCACTCTGAGACG
AAGGCTCGCGTCCAGGCCAGCACGAAGGAGGCTAAGTGGGAGGGGTAGCGGTCTGTGTCCTACTA
GGGGGTCCACTCGCTCCAGGGTGTGAAGACACATGTCGCCCTCTTCGGCATCAAGGAAGGTGAT
TGGTTTATAGGTGTAGGCCACGTGACCGGGTGTTCCTGAAGGGGGGTATAAAAGGGGGTGGGG
GCGCGTTCTGCTCCTCACTCTCTTCCGCATCGCTGTCTGCGAGGGCCAGCTGTTGGGCTCGCGGTT
GAGGACAACTCTTCGCGGTCTTTCCAGTACTCTTGGATCGGAAACCCGTCCGCCTCCGAACGG
TACTCCGCCACCGAGGGACCTGAGCGAGTCCGCATCGACCGGATCGGAAAACCTCTCGAGCCAC
C
ATGCAAATAGAGCTCTCCACCTGCTTCTTTCTGTGCCTTTTGCGATTCTGCTTTAGTGCCACCA
GAAGATACTACCTGGGTGCAGTGGAAGTGTATGGGACTATATGCAAAGTGATCTCGGTGAGCT
GCCTGTGGACGCAAGATTTCTCCTAGAGTGCCAAAATCTTTTCCATTCAACACCTCAGTCGTG
TACAAAAGACTCTGTTTGTAGAATTCACGGTTCACCTTTTCAACATCGCTAAGCCAAGGCCAC
CCTGGATGGGTCTGCTAGGTCTTACCATCCAGGCTGAGGTTTATGATACAGTGGTCATTACACT
TAAGAACATGGCTTCCCATCCTGTCTGCTCTTTCATGCTGTTGGTGTATCCTACTGGAAAGCTTCT
GAGGGAGCTGAATATGATGATCAGACCAGTCAAAGGGAGAAAGAAGATGATAAAGTCTTCCCTG
GTGGAAGCCATACATATGTCTGGCAGGTCTGAAAGAGAATGGTCCAATGGCCTCTGACCCACT
GTGCCTTACCTACTCATATCTTTCTCATGTGGACCTGGTAAAGACTTGAATTCAGGCCTCATT
GGAGCCCTACTAGTATGTAGAGAAGGGAGTCTGGCCAAGGAAAAGACACAGACCTTGACACAAAT
TTATACTACTTTTTTGCTGTATTTGATGAAGGGAAAAGTTGGCACTCAGAAACAAAGAACTCCTT
GATGCAGGATAGGGATGCTGCATCTGCTCGGGCCTGGCCTAAAATGCACACAGTCAATGGTTAT
GTAAACAGGTCTCTGCCAGGTCTGATTGGATGCCACAGGAAATCAGTCTATTGGCATGTGATTG
GAATGGGCACCACTCCTGAAGTGCATCAATATTCCTCGAAGGTCACACATTTCTTGTGAGGAA
CCATCGCCAGGCGTCTTGGAAATCTCGCCAATAACTTTCTTACTGCTCAAACACTCTTGATG
GACCTTGGACAGTTTCTACTGTTTTGTTCATATCTCTTCCCAACATGATGGCATGGAAGCTT
ATGTCAAAGTAGACAGCTGTCCAGAGGAACCCCACTACGAATGAAAAATAATGAAGAAGCGGA
AGACTATGATGATGATCTTACTGATTCTGAAATGGATGTGGTCAGGTTTGTATGATGACAACTCT
CCTTCCTTTATCCAAATTCGCTCAGTTGCCAAGAAGCATCCTAAACTTGGGTACATTACATTG
CTGCTGAAGAGGAGGACTGGGACTATGCTCCCTTAGTCCTCGCCCCGATGACAGAAGTTATAA
AAGTCAATATTTGAACAATGGCCCTCAGCGGATTGGTAGGAAGTACAAAAAGTCCGATTTATG
GCATACACAGATGAAACCTTTAAGACTCGTGAAGCTATTTCAGCATGAATCAGGAATCTTGGGAC
CTTTACTTTTATGGGGAAGTTGGAGACACACTGTTGATTATATTTAAGAATCAAGCAAGCAGACC
ATATAACATCTACCCTCACGGAATCACTGATGTCCGTCTTTGTATTCAAGGAGATTACCAAAA

Fig 4

GGTGTAAAACATTTGAAGGATTTTCCAATTCTGCCAGGAGAAATATTCAAATATAAATGGACAG
TGACTGTAGAAGATGGGCCAACTAAATCAGATCCTCGGTGCCTGACCCGCTATTACTCTAGTTT
CGTTAATATGGAGAGAGATCTAGCTTCAGGACTCATTGGCCCTCTCCTCATCTGCTACAAAGAA
TCTGTAGATCAAAGAGGAAACCAGATAATGTCAGACAAGAGGAATGTCATCCTGTTTTCTGTAT
TTGATGAGAACCGAAGCTGGTACCTCACAGAGAATATACAACGCTTTCTCCCAATCCAGCTGG
AGTGCAGCTTGAGGATCCAGAGTTCCAAGCCTCCAACATCATGCACAGCATCAATGGCTATGTT
TTTGATAGTTTTGCAGTTGTGAGTTTGTTCATGAGGTGGCATACTGGTACATTCTAAGCATTG
GAGCACAGACTGACTTCCTTTCTGTCTTCTCTCTGGATATACCTTCAAACACAAAATGGTCTA
TGAAGACACACTCACCTTATTCCCATTCTCAGGAGAACTGTCTTCATGTGATGGAAAACCCA
GGTCTATGGATTCTGGGGTGCCACAACCTCAGACTTTCGGAACAGAGGCATGACCGCCTTACTGA
AGGTTTCTAGTTGTGACAAGAACACTGGTGATTATTACGAGGACAGTTATGAAGATATTTTCAGC
ATACTTGCTGAGTAAAAACAATGCCATTGAACCTAGGAGCTTCTCTCAGAATCCACCAGTCTTG
AAACGCCATCAACGGGAAATAACTCGTACTACTCTTCAGTCAGATCAAGAGGAAATTGACTATG
ATGATACCATATCAGTTGAAATGAAGAAGGAAGATTTTGACATTTATGATGAGGATGAAATCA
GAGCCCCCGCAGCTTTCAAAGAAAACACGACACTATTTTATTGCTGCAGTGGAGAGGCTCTGG
GATTATGGGATGAGTAGCTCCCCACATGTTCTAAGAAACAGGGCTCAGAGTGGCAGTGTCCCTC
AGTTCAAGAAAGTTGTTTTCCAGGAATTTACTGATGGCTCCTTTACTCAGCCCTTATACCGTGG
AGAATAAATGAACATTTGGGACTCCTGGGGCCATATATAAGAGCAGAAGTTGAAGATAATATC
ATGGTAACTTTTCAGAAATCAGGCCTCTCGTCCCTATTCTTCTATTCTAGCCTTATTTCTTATG
AGGAAGATCAGAGGCAAGGAGCAGAACCTAGAAAAAACTTTGTCAAGCCTAATGAAACCAAAAC
TTACTTTTGGAAAGTGCAACATCATATGGCACCCACTAAAGATGAGTTTGACTGCAAAGCCTGG
GCTTATTTCTCTGATGTTGACCTGGAAAAAGATGTGCACTCAGGCCTGATTGGACCCCTTCTGG
TCTGCCACACTAACACACTGAACCCTGCTCATGGGAGACAAGTGACAGTACAGGAATTTGCTCT
GTTTTTCACCATCTTTGATGAGACCAAAAGCTGGTACTTCACTGAAAATATGGAAAGAACTGC
AGGGCTCCCTGCAATATCCAGATGGAAGATCCCACTTTTAAAGAGAATTATCGCTTCCATGCAA
TCAATGGCTACATAATGGATACACTACCTGGCTTAGTAATGGCTCAGGATCAAAGGATTCGATG
GTATCTGCTCAGCATGGGCAGCAATGAAAACATCCATTCTATTCAATTCAGTGGACATGTGTTT
ACTGTACGAAAAAAGAGGAGTATAAATGGCACTGTACAATCTCTATCCAGGTGTTTTTGAGA
CAGTGGAAATGTTACCATCCAAAGCTGGAATTTGGCGGGTGAATGCCTTATTGGCGAGCATCT
ACATGCTGGGATGAGCACACTTTTTCTGGTGTACAGCAATAAGTGTGCACTCCCCTGGGAATG
GCTTCTGGACACATTAGAGATTTTCAGATTACAGCTTCAGGACAATATGGACAGTGGGCCCCAA
AGCTGGCCAGACTTCATTATTCCGGATCAATCAATGCCTGGAGCACCAGGAGCCCTTTTCTTG
GATCAAGGTGGATCTGTTGGCACCAATGATTATTACGGCATCAAGACCCAGGGTGCCCGTCAG
AAGTTCTCCAGCCTCTACATCTCTCAGTTTATCATCATGTATAGTCTTGATGGGAAGAAGTGGC
AGACTTATCGAGGAAATTCCACTGGAACCTTAATGGTCTTCTTTGGCAATGTGGATTCTCTGG
GATAAAACACAATATTTTAAACCCTCCAATTATTGCTCGATACATCCGTTTGCACCCCACTCAT
TATAGCATTTCGCAGCACTCTTCGCATGGAGTTGATGGGCTGTGATTAAATAGTTGCAGCATGC
CATTGGGAATGGAGAGTAAAGCAATATCAGATGCACAGATTACTGCTTCATCCTACTTTACCAA
TATGTTTGCCACCTGGTCTCCTTCAAAGCTCGACTTCACCTCCAAGGGAGGAGTAATGCCTGG
AGACCTCAGGTGAATAATCCAAAAGAGTGGCTGCAAGTGGACTTCCAGAAGACAATGAAAGTCA
CAGGAGTAACTACTCAGGGAGTAAATCTCTGCTTACCAGCATGTATGTGAAGGAGTTCCTCAT
CTCCAGCAGTCAAGATGGCCATCAGTGGACTCTCTTTTTTTCAGAATGGCAAAGTAAAGGTTTTT
CAGGGAAATCAAGACTCCTTCACACCTGTGGTGAACCTCTTAGACCCACCGTTACTGACTCGCT
ACCTTCGAATTCACCCCCAGAGTTGGGTGCACCAGATTGCCCTGAGGATGGAGGTTCTGGGCTGC
GAGGCACAGGACCTCTACTGA
GGGCGGCGCTGCAGCACCTGCCACTGCCGTCACCTCTCCCTCCTCAGCTCCAGGGCAGTGTCC
CTCCCTGGCTTGCTTCTACCTTTGTGCTAAATCCTAGCAGACACTGCCTTGAAGCCTCCTGAA
TTAACTATCATCAGTCTGCAATTTCTTTGGTGGGGGGCCAGGAGGGTGCATCCAATTTAACTTA
ACTCTTACCTATTTTCTGCAGCTGCTCCCAGATTACTCCTTCCCTTCCAATATAACTAGGCAAAA
AGAAGTGAGGAGAAACCTGCATGAAAGCATTTCTCCCTGAAAAGTTAGGCCTCTCAGAGTCACC
ACTTCTCTGTTGTAGAAAACTATGTGATGAACTTTGAAAAAGATATTTATGATGTTAACAT
TTCAGGTAAAGCCTCATACGTTTAAAATAAACTCTCAGTTGTTTATTATCCTGATCAAGCATG
GAACAAAGCATGTTTCAGGATCAGATCAATAAATCTTGGAGTCAAAGGCAAATCATTTGGAC
AATCTGCAAATGGAGAGAATACAATACTACTACAGTAAAGTCTGTTTCTGCTTCTTACACA
TAGATATAATTATGTTATTTAGTCATTATGAGGGGCACATTCTTATCTCCAAACTAGCATTCT
TAAACTGAGAATTATAGATGGGGTTCAAGAATCCCTAAGTCCCCTGAAATTATATAAGGCATTCT
TGTATAAATGCAAATGTGCATTTTTCTGACGAGTGTCCATAGATATGGGACATATGACGTGAGC


TCAGATCTTTGTGAAGGAACCTTACTTCTGTGGTGTGACATAATTGGACAAACTACCTACAGAG
ATTTAAAGCTCTAAGGTAAATATAAAATTTTAAAGTGTATAATGTGTTAAACTACTGATTCTAA
TTGTTTGTGTATTTTAGATTCCAACCTATGGAACCTGATGAATGGGAGCAGTGGTGGAATGCCTT
TAATGAGGAAAACCTGTTTTGCTCAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGAG
TGTGAACATTCTACTCCTCCAAAAAGAAGAGAAAGGTAGAAGACCCCAAGGACTTTCCTTCAG
AATTGCTAAGTTTTTTGAGTCATGCTGTGTTTAGTAATAGAACTCTTGCTTGCTTTGCTATTTA
CACCACAAAGGAAAAAGCTGCACCTGCTATACAAGAAAATTATGGAAAAATATTCTGTAACCTTT
ATAAGTAGGCATAACAGTTATAATCATAACATACTGTTTTTTCTTACTCCACACAGGCATAGAG
TGTCTGCTATTAATAACTATGCTCAAAAATTGTGTACCTTTAGCTTTTAAATTTGTAAAGGGGT
TAATAAGGAATATTTGATGTATAGTGCCCTGACTAGAGATCATAATCAGCCATACCACATTTGT
AGAGGTTTTACTTGCTTTAAAAAACCTCCACACCTCCCCCTGAACCTGAAACATAAAATGAAT
GCAATTGTTGTTGTTAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCA
CAAATTTACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTTGTCCAAACTCATCAA
TGTATCTTATCATGTCTGGATCCTCTACGCCGACGCATCGTGGCCGGCATCACCGGCGCCACA
GGTGCGGTTGCTGGCGCCTATATCGCCGACATCACCGATGGGGAAGATCGGGCTCGCCACTTCG
GGCTCATGAGCGCTTGTTTCGGCGTGCGGTATGGTGGCAGGCCCGTGCGCGGGGGACTGTTGGGC
GCCATCTCCTTGCTATGCACCATTCCTTGCGCGCGCGGTGCTCAACGGCCTCAACCTACTACTGG
GCTGCTTCCTAATGCAGGAGTCGCATAAGGGAGAGCGTCGAAATTCTCATGTTTGACAGCTTAT
CATCGGCGCAGCACCATGGCCTGAAATAACCTCTGAAAGAGGAACTTGGTTAGGTACCTTCTGA
GGCGGAAAGAACCAGCTGTGGAATGTGTGTGTCAGTTAGGGTGTGGAAAGTCCCCAGGCTGGGGAG
CAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCAGGTGTGGAAAGTCCCCAGG
CTCCCCAGCAGGCAGAAGTATGCAAAGCATGCATCTCAATTAGTCAGCAACCATAGTCCCGCCC
CTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTTCTCGCCCCATGGCTGAC
TAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATTCCAGCCGTAGTG
AGGAGGCTTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTTCACGCTGCCGCAAGCACTCAGGGC
GCAAGGGCTGCTAAAGGAAGCGGAACACGTAGAAAGCCAGTCCGCAGAAACGGTGCTGACCCCG
GATGAATGTGAGCTACTGGGCTATCTGGACAAGGGAAAACGCAAGCGCAAAGAGAAAGCAGGTA
GCTTGCACTGGGCTTACATGGCGATAGCTAGACTGGGCGGTTTTATGGACAGCAAGCGAACCAG
AATTGCCAGCTGGGGCGCCCTCTGGTAAGGTGGGAAGCCCTGCAAAGTAAACTGGATGGCTTT
CTTGCCGCCAAGGATCTGATGGCGCAGGGGATCAAGATCTGATCAAGAGACAGGATGAGGATCG
TTTCGCATGATTGAACAAGATGGATTGCACGCAGGTTCTCCGGCCGCTTGGGTGGAGAGGCTAT
TCGGCTATGACTGGGCACAACAGACAATCGGCTGCTCTGATGCCGCCGTGTTCCGGCTGTCAGC
GCAGGGGCGCCCGGTTCTTTTGTCAAGACCGACCTGTCCGGTGCCCTGAATGAACTGCAGGAC
GAGGCAGCGCGGCTATCGTGGCTGGCCACGACGGGCGTTCTTGCGCAGCTGTGCTCGACGTTG
TCACTGAAGCGGGAAGGGACTGGCTGCTATTGGGCGAAGTGCCGGGGCAGGATCTCCTGTCATC
TCACCTTGCTCCTGCCGAGAAAGTATCCATCATGGCTGATGCAATGCGGCGGCTGCATACGCTT
GATCCGGCTACCTGCCCATTTCGACCACCAAGCGAAACATCGCATCGAGCGAGCACGTACTCGGA
TGGAAGCCGGTCTTGTGATCAGGATGATCTGGACGAAGAGCATCAGGGGCTCGCGCCAGCCGA
ACTGTTTCGCCAGGCTCAAGGCGCGCATGCCCGACGGCGAGGATCTCGTCGTGACCCATGGCGAT
GCCTGCTTGCCGAATATCATGGTGGAAAATGGCCGCTTTTCTGGATTCTCGACTGTGGCCGGC
TGGGTGTGGCGGACCGCTATCAGGACATAGCGTTGGCTACCCGTGATATTGCTGAAGAGCTTGG
CGGCGAATGGGCTGACCGCTTCTCGTGCTTTACGGTATCGCCGCTCCCGATTTCGAGCGCATC
GCCTTCTATCGCCTTCTTGACGAGTTCTTCTGAGCGGGACTCTGGGGTTCGAAATGACCGACCA
AGCGACGCCCAACCTGCCATCACGAGATTTGATTTCCACCGCCGCTTCTATGAAAGGTTGGGC
TTCGGAATCGTTTTCCGGGACGCCGGCTGGATGATCCTCCAGCGCGGGGATCTCATGCTGGAGT
TCTTCGCCCACCCCGGGCTCGATCCCTCGCGAGTTGGTTCAGCTGCTGCCTGAGGCTGGACGA
CCTCGCGGAGTTCTACCGGCAGTGCAATCCGTCCGCATCCAGGAAACCAGCAGCGGCTATCCG
CGCATCCATGCCCCCGAACTGCAGGAGTGGGGAGGCACGATGGCCGCTTTGGTCCCGGATCTTT
GTGAAGGAACCTTACTTCTGTGGTGTGACATAATTGGAGAAACTACCTACAGAGATTTAAAGCT
CTAAGGTAAATATAAAATTTTAAAGTGTATAATGTGTTAAACTACTGATTCTAATTGTTTGTGT
ATTTTAGATTCCAACCTATGGAACCTGATGAATGGGAGCAGTGGTGGAATGCCTTTAATGAGGAA
AACCTGTTTTGCTCAGAAGAAATGCCATCTAGTGATGATGAGGCTACTGCTGACTCTCAACATT
CTACTCCTCCAAAAAGAAGAGAAAGGTAGAAGACCCCAAGGACTTTCCTTCAGAATTGCTAAG
TTTTTTGAGTCATGCTGTGTTTAGTAATAGAACTCTTGCTTGCTTTGCTATTTACACCACAAAG
GAAAAAGCTGCACCTGCTATACAAGAAAATTATGGAAAAATATTCTGTAACCTTTATAAGTAGGC
ATAACAGTTATAATCATAACATACTGTTTTTTCTTACTCCACACAGGCATAGAGTGTCTGCTAT
TAATAACTATGCTCAAAAATTGTGTACCTTTAGCTTTTAAATTTGTAAAGGGTTAATAAGGAA


TATTTGATGTATAGTGCCTTGACTAGAGATCATAATCAGCCATACCACATTTGTAGAGGTTTTA
CTTGCTTTAAAAAACCTCCACACCTCCCCCTGAACCTGAAACATAAAATGAATGCAATTGTTG
TTGTTAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAATTTTAC
AAATAAAGCATTTTTTTCACTGCATTCTAGTTGTGGTTTTGTCCAAACTCATCAATGGTATCTTA
TCATGTCTGGATCTCGACCGAGCCCTTGAGAGCCTTCAACCCAGTCAGCTCCTTCCGGTGGGCG
CGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATGCAACTCGTAGGACAGG
TGCCGGCAGCGCTCTGGGTCATTTTCGGCGAGGACCGCTTTCGCTGGAGCGCGACGATGATCGG
CCTGTCGCTTGCGGTATTTCGGAATCTTGCACGCCCTCGCTCAAGCCTTCGTCACTGGTCCCGCC
ACCAAACGTTTTCGGCGAGAAGCAGGCCATTATCGCCGGCATGGCGGCCGACGCGCTGGGCTACG
TCTTGCTGGCGTTTCGCGACGCGAGGCTGGATGGCCTTCCCCATTATGATTCTTCTCGCTTCCGG
CGGCATCGGGATGCCCGCGTTGCAGGCCATGCTGTCCAGGCAGGTAGATGACGACCATCAGGGA
CAGCTTCAAGGATCGCTCGCGGCTCTTACCAGCCTAACTTCGATCACTGGACCGCTGATCGTCA
CGGCGATTTATGCCGCCTCGGCGAGCACATGGAACGGGTTGGCATGGATTGTAGGCGCCGCCCT
ATACCTTGTCTGCCTCCCCGCGTTGCGTCGCGGTGCATGGAGCCGGGCCACCTCGACCTGAATG
GAAGCCGGCGGCACCTCGCTAACGGATTACCACTCCAAGAATTGGAGCCAATCAATTCTTGCG
GAGAACTGTGAATGCGCAAACCAACCCTTGGCAGAACATATCCATCGCGTCCGCCATCTCCAGC
AGCCGCACGCGGCGCATCTCGGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGAC
GAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACC
AGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATA
CCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTCTCAATGCTCACGCTGTACCTATCTC
AGTTCGGTGTACCTCGTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACC
GCTGCGCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGTAAGACACGACTTATCGCCACT
GGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTG
AAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGC
CAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGG
TGGTTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTG
ATCTTTTCTACGGGGTCTGACGCTCAGTGGAACGAAACTCACGTTAAGGGATTTTGGTCATGA
GATTATCAAAAAGGATCTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTA
AAGTATATATGAGTAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCA
GCGATCTGTCTATTTTCGTTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATAC
GGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCC
AGATTTATCAGCAATAAACCAGCCAGCCAGAAGGGCCGAGCGCAGAAGTGGTCTTGCAACTTTA
TCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATA
GTTTGCGCAACGTTGTTGCCATTGCTGCAGGCATCGTGGTGTACGCTCGTCTGTTGGTATGGC
TTCATTTCAGCTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGTGCAAAAAA
GCGGTTAGCTCCTTCGGTCTCCGATCGTTGTGAGAAGTAAGTTGGCCGAGTGTATCACTCA
TGGTTATGGCAGCACTGCATAATTCTCTTACTGTATGCCATCCGTAAGATGCTTTTCTGTGAC
TGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCG
GCGTCAACACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTGGAAAAC
GTTCTTCGGGGCGAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCAC
TCGTGCACCCAACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACA
GGAAGGCAAAATGCCGCAAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCT
TCCTTTTTCAATATTATTGAAGCATTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGA
ATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGTGCCACCTGAC
GTCTAAGAAACCATTATTATCATGACATTAACCTATAAAAATAGGCGTATCACGAGGCCCTTTC
GTCTTCAA*




Morphology & fluorescence of PLE9
transduced cells at day 53.

Fig 5

ELISA of FVIII in conditioned media from transfected cells*Fig. 4*

ELISA of FVIII in conditioned media from transfected cells*Fig. 4*

*Fig 7*