

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07H 17/00		A2	(11) International Publication Number: WO 00/44761
			(43) International Publication Date: 3 August 2000 (03.08.00)
(21) International Application Number: PCT/IB99/02051 (22) International Filing Date: 28 December 1999 (28.12.99) (30) Priority Data: 60/117,342 27 January 1999 (27.01.99) US (71) Applicant (for all designated States except US): PFIZER PRODUCTS INC. [US/US]; Eastern Point Road, Groton, CT 06340 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KANEKO, Takushi [JP/US]; 398 Northwood Drive, Guilford, CT 06437 (US). MCMILLEN, William, Thomas [US/US]; 4-K Lakeside Drive, Ledyard, CT 06339 (US). (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: KETOLIDE ANTIBIOTICS			
(57) Abstract			
This invention relates to compounds of formula (1) and to pharmaceutically acceptable salts and solvates thereof wherein X¹, X², R², R⁸, R⁹, R¹⁰ and R¹¹ are as defined herein. The compounds of formula (1) are antibacterial and antiprotozoal agents that may be used to treat various bacterial and protozoal infections and disorders related to such infections. The invention also relates to pharmaceutical compositions containing the compounds of formula (1) and to methods of treating bacterial and protozoal infections by administering the compounds of formula (1).		(1)	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5


KETOLIDE ANTIBIOTICS**Background of the Invention**

This invention relates to novel macrolide compounds that are useful as antibacterial and antiprotozoal agents in mammals, including man, as well as in fish and birds. This invention also relates to pharmaceutical compositions containing the novel compounds and to methods of treating bacterial and protozoal infections in mammals, fish and birds by administering the novel compounds to mammals, fish and birds requiring such treatment.

Macrolide antibiotics are known to be useful in the treatment of a broad spectrum of bacterial and protozoal infections in mammals, fish and birds. Such antibiotics include various derivatives of erythromycin A such as azithromycin which is commercially available and is referred to in United States patents 4,474,768 and 4,517,359, both of which are incorporated herein by reference in their entirety. Other macrolide antibiotics are disclosed and claimed in PCT international published application WO 98/56800 (published December 17, 1998), which designates the United States; United States Patent 5,527,780, issued June 18, 1996; United States provisional application number 60/101263 (filed September 22, 1998)(attorney docket number PC 10406); United States provisional patent application number 60/111,728 (filed December 10, 1998)(attorney docket number PC 10494); PCT published application WO 98/01546 (published January 15, 1998); PCT published application WO 98/01571 (published January 15, 1998); EP published application number 949268 (published October 13, 1999); and U.S. patent 5,747,467 (issued May 5, 1998). Each of the foregoing United States patents and patent applications and published EP and PCT international patent applications are incorporated herein by reference in their entirety. Like azithromycin and other macrolide antibiotics, the novel macrolide compounds of the present invention possess activity against various bacterial and protozoal infections as described below.

Summary of the Invention

The present invention relates to compounds of the formula

**1**

5 and to pharmaceutically acceptable salts and solvates thereof, wherein:

X^1 is O, $-CR^4R^5$ - or $-NR^4$ -;

X^2 is =O or =NOR¹;

R^1 is H or C₁-C₁₀ alkyl, wherein 1 to 3 carbons of said alkyl are optionally replaced by a heteroatom selected from O, S and -N(R⁴)-, and said alkyl is optionally substituted by 1 to 3
 10 substituents independently selected from the group consisting of -C(O)O(C₁-C₁₀ alkyl), C₁-C₁₀ alkoxy, C₁-C₁₀ alkanoyl, halo, nitro, cyano, 4-10 membered heterocyclyl, C₁-C₁₀ alkyl, $-NR^4R^5$, C₆-C₁₀ aryl, $-S(O)_n(C_1-C_{10} \text{ alkyl})$ wherein n is an integer ranging from 0 to 2, and $-SO_2NR^4R^5$;

R^2 is $-(CR^4R^5)_n$ (4-10 membered heterocyclic) or $-(CR^4R^5)_n$ (C₆-C₁₀ aryl), wherein n is an integer from 0 to 6, from 1 to 3 R⁴ or R⁵ groups of the $-(CR^4R^5)_n$ - moiety of the foregoing R²
 15 groups are optionally replaced with a halo substituent, and the heterocyclic and aryl moieties of the foregoing R² groups are optionally substituted with 1 to 4 R³ groups;

each R³ is independently selected from halo, cyano, nitro, trifluoromethoxy, trifluoromethyl, azido, hydroxy, C₁-C₆ alkoxy, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, $-C(O)R^6$, $-C(O)OR^6$, $-OC(O)R^6$, $-NR^6C(O)R^7$, $-NR^6C(O)NR^1R^7$, $-NR^6C(O)OR^7$, $-C(O)NR^6R^7$, $-NR^6R^7$,
 20 $-NR^6OR^7$, $-SO_2NR^6R^7$, $-S(O)_j(C_1-C_6 \text{ alkyl})$ wherein j is an integer from 0 to 2, $-(CR^1R^2)_h$ (C₆-C₁₀ aryl), $-(CR^4R^5)_h$ (4-10 membered heterocyclic), $-(CR^4R^5)_qC(O)(CR^4R^5)_h$ (C₆-C₁₀ aryl), $-(CR^4R^5)_qC(O)(CR^4R^5)_h$ (4-10 membered heterocyclic), $-(CR^4R^5)_hO(CR^4R^5)_q$ (C₆-C₁₀ aryl), $-(CR^4R^5)_hO(CR^4R^5)_q$ (4-10 membered heterocyclic), $-(CR^4R^5)_qSO_2(CR^4R^5)_h$ (C₆-C₁₀ aryl), and $-(CR^4R^5)_qSO_2(CR^4R^5)_h$ (4-10 membered heterocyclic), wherein q and t are each independently
 25 an integer from 0 to 5, 1 or 2 ring carbon atoms of the heterocyclic moieties of the foregoing R¹⁰ groups are optionally substituted by an oxo (=O) moiety, and the alkyl, alkenyl, alkynyl, aryl and heterocyclic moieties of the foregoing R¹⁰ groups are optionally substituted by 1 to 3 substituents independently selected from halo, cyano, nitro, trifluoromethyl, trifluoromethoxy, azido, $-OR^6$, $-C(O)R^6$, $-C(O)OR^6$, $-OC(O)R^6$, $-NR^6C(O)R^7$, $-C(O)NR^6R^7$, $-NR^6R^7$, $-NR^6OR^7$, C₁-
 30 C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, $-(CR^4R^5)_h$ (C₆-C₁₀ aryl), and $-(CR^4R^5)_h$ (4-10 membered heterocyclic), wherein t is an integer from 0 to 5;

each R⁴ and R⁵ is independently selected from H and C₁-C₆ alkyl;

each R⁶ and R⁷ is independently selected from H, C₁-C₆ alkyl, $-(CR^4R^5)_h$ (C₆-C₁₀ aryl), and $-(CR^4R^5)_h$ (4-10 membered heterocyclic), wherein t is an integer from 0 to 5, 1 or 2 ring
 35 carbon atoms of the heterocyclic group are optionally substituted by an oxo (=O) moiety, and the alkyl, aryl and heterocyclic moieties of the foregoing R⁶ and R⁷ groups are optionally substituted by 1 to 3 substituents independently selected from halo, cyano, nitro, $-NR^4R^5$, trifluoromethyl, trifluoromethoxy, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, hydroxy, and C₁-C₆ alkoxy;

40 R^8 is H, $-C(O)(C_1-C_6 \text{ alkyl})$, benzyl, benzyloxycarbonyl or $(C_1-C_6 \text{ alkyl})_3\text{silyl}$;


R^9 is C₁-C₆ alkyl;

R^{10} is H or C₁-C₁₀ alkyl; and

5 R^{11} is selected from chloro, bromo, iodo, fluoro, and cyano;


with the proviso that where X^2 is =O then the R^{11} is cyano or from 1 to 3 R^4 or R^5 groups of the $-(CR^4R^5)_n$ - moiety of the R^2 groups recited above are either C_1 - C_6 alkyl or replaced with a halo substituent.

Specific embodiments of the present invention include compounds of formula 2 (which
10 is a specific embodiment within the genus of formula 1)



wherein R^{12} , R^{13} , R^{14} and R^{15} are each independently selected from H, halo, methyl and ethyl. More specific embodiments include the compounds of formula 2 wherein R^{14} and R^{15} are both H and R^{12} and R^{13} are each independently selected from H and methyl. In a preferred
15 embodiment of the compounds of formula 2, R^{12} , R^{13} , R^{14} and R^{15} are each H.

Other specific embodiments of the present invention include compounds of formula 3 (which is a specific embodiment within the genus of formula 1)



3

5

wherein R¹², R¹³, R¹⁴ and R¹⁵ are each independently selected from H, halo, methyl and ethyl with the proviso that at least one of R¹², R¹³, R¹⁴ and R¹⁵ is not H. More specific embodiments include the compounds of formula **3** wherein R¹⁴ and R¹⁵ are both H, R¹² is methyl, and R¹³ is selected from H and methyl.

10

The invention also relates to a pharmaceutical composition for the treatment of a bacterial infection or a protozoa infection, or a disorder related to a bacterial or protozoal infection, in a mammal, fish, or bird, which comprises a therapeutically effective amount of a compound of formula 1, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

15

The invention also relates to a method of treating a bacterial infection or a protozoa infection, or a disorder related to a bacterial or protozoal infection, in a mammal, fish, or bird which comprises administering to said mammal, fish or bird a therapeutically effective amount of a compound of formula **1** or a pharmaceutically acceptable salt or solvate thereof.

The term "treating", as used herein, unless otherwise indicated, means reversing, 20 alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, refers to the act of treating, as "treating" is defined immediately above.

As used herein, unless otherwise indicated, the terms or phrases "bacterial infection(s)", "protozoal infection(s)", and "disorders related to bacterial infections or protozoal infections" include the following: pneumonia, otitis media, sinusitis, bronchitis, tonsillitis, and mastoiditis related to infection by *Streptococcus pneumoniae*, *Haemophilus influenzae*, *Moraxella catarrhalis*, *Staphylococcus aureus*, *Enterococcus faecalis*, *E. faecium*, *E. casseliflavus*, *S. epidermidis*, *S. haemolyticus*, or *Peptostreptococcus* spp.; pharyngitis,

5 rheumatic fever, and glomerulonephritis related to infection by *Streptococcus pyogenes*, Groups C and G streptococci, *Corynebacterium diphtheriae*, or *Actinobacillus haemolyticum*; respiratory tract infections related to infection by *Mycoplasma pneumoniae*, *Legionella pneumophila*, *Streptococcus pneumoniae*, *Haemophilus influenzae*, or *Chlamydia pneumoniae*; blood and tissue infections, including endocarditis and osteomyelitis, caused by *S. aureus*, *S.*

10 *haemolyticus*, *E. faecalis*, *E. faecium*, *E. durans*, including strains resistant to known antibacterials such as, but not limited to, beta-lactams, vancomycin, aminoglycosides, quinolones, chloramphenicol, tetracyclines and macrolides; uncomplicated skin and soft tissue infections and abscesses, and puerperal fever related to infection by *Staphylococcus aureus*, coagulase-negative staphylococci (i.e., *S. epidermidis*, *S. hemolyticus*, etc.), *Streptococcus*

15 *pyogenes*, *Streptococcus agalactiae*, Streptococcal groups C-F (minute-colony streptococci), viridans streptococci, *Corynebacterium minutissimum*, *Clostridium* spp., or *Bartonella henselae*; uncomplicated acute urinary tract infections related to infection by *Staphylococcus aureus*, coagulase-negative staphylococcal species, or *Enterococcus* spp.; urethritis and cervicitis; sexually transmitted diseases related to infection by *Chlamydia trachomatis*, *Haemophilus*

20 *ducreyi*, *Treponema pallidum*, *Ureaplasma urealyticum*, or *Neisseria gonorrhoeae*; toxin diseases related to infection by *S. aureus* (food poisoning and toxic shock syndrome), or Groups A, B, and C streptococci; ulcers related to infection by *Helicobacter pylori*; systemic febrile syndromes related to infection by *Borrelia recurrentis*; Lyme disease related to infection by *Borrelia burgdorferi*; conjunctivitis, keratitis, and dacrocystitis related to infection by *Chlamydia*

25 *trachomatis*, *Neisseria gonorrhoeae*, *S. aureus*, *S. pneumoniae*, *S. pyogenes*, *H. influenzae*, or *Listeria* spp.; disseminated *Mycobacterium avium* complex (MAC) disease related to infection by *Mycobacterium avium*, or *Mycobacterium intracellulare*; infections caused by *Mycobacterium tuberculosis*, *M. leprae*, *M. paratuberculosis*, *M. kansasii*, or *M. chelonae*; gastroenteritis related to infection by *Campylobacter jejuni*; intestinal protozoa related to infection by *Cryptosporidium*

30 spp.; odontogenic infection related to infection by viridans streptococci; persistent cough related to infection by *Bordetella pertussis*; gas gangrene related to infection by *Clostridium perfringens* or *Bacteroides* spp.; and atherosclerosis or cardiovascular disease related to infection by *Helicobacter pylori* or *Chlamydia pneumoniae*. Bacterial infections and protozoal infections, and disorders related to such infections, which may be treated or prevented in animals include the

35 following: bovine respiratory disease related to infection by *P. haemolytica*, *P. multocida*, *Mycoplasma bovis*, or *Bordetella* spp.; cow enteric disease related to infection by *E. coli* or protozoa (i.e., coccidia, cryptosporidia, etc.); dairy cow mastitis related to infection by *S. aureus*, *Strep. uberis*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Klebsiella* spp., *Corynebacterium*, or *Enterococcus* spp.; swine respiratory disease related to infection by *A.*

40 *pleuro.*, *P. multocida*, or *Mycoplasma* spp.; swine enteric disease related to infection by *E. coli*, *Lawsonia intracellularis*, *Salmonella*, or *Serpulina hyodysenteriae*; cow footrot related to infection

- 5 by *Fusobacterium* spp.; cow metritis related to infection by *E. coli*; cow hairy warts related to infection by *Fusobacterium necrophorum* or *Bacteroides nodosus*; cow pink-eye related to infection by *Moraxella bovis*; cow premature abortion related to infection by protozoa (i.e. neosporium); urinary tract infection in dogs and cats related to infection by *E. coli*; skin and soft tissue infections in dogs and cats related to infection by *S. epidermidis*, *S. intermedius*,
10 coagulase neg. *Staphylococcus* or *P. multocida*; and dental or mouth infections in dogs and cats related to infection by *Alcaligenes* spp., *Bacteroides* spp., *Clostridium* spp., *Enterobacter* spp., *Eubacterium*, *Peptostreptococcus*, *Porphyromonas*, or *Prevotella*. Other bacterial infections and protozoal infections, and disorders related to such infections, which may be treated or prevented in accord with the method of the present invention are referred to in J. P.
15 Sanford et al., "The Sanford Guide To Antimicrobial Therapy," 26th Edition, (Antimicrobial Therapy, Inc., 1996).

The term "halo", as used herein, unless otherwise indicated, includes fluoro, chloro, bromo or iodo. Preferred halo groups are fluoro, chloro and bromo.

- The term "alkyl", as used herein, unless otherwise indicated, includes saturated
20 monovalent hydrocarbon radicals having cyclic, straight and/or branched moieties. It is to be understood that to include cyclic moieties, the alkyl group must include at least 3 carbon atoms.

The term "alkenyl", as used herein, unless otherwise indicated, includes alkyl groups as defined above having at least one carbon-carbon double bond at some point in the alkyl chain.

- The term "alkynyl", as used herein, unless otherwise indicated, includes alkyl groups as
25 defined above having at least one carbon-carbon triple bond at some point in the alkyl chain.

The term "aryl", as used herein, unless otherwise indicated, includes an organic radical derived from an aromatic hydrocarbon by removal of one hydrogen, such as phenyl or naphthyl.

- The term "4-10 membered heterocyclic", as used herein, unless otherwise indicated, includes aromatic and non-aromatic heterocyclic groups containing one or more heteroatoms each selected from O, S and N, wherein each heterocyclic group has from 4-10 atoms in its ring system. Non-aromatic heterocyclic groups include groups having only 4 atoms in their ring system, but aromatic heterocyclic groups must have at least 5 atoms in their ring system. The heterocyclic groups include benzo-fused ring systems and ring systems substituted with one or more oxo moieties. An example of a 4 membered heterocyclic group is azetidiny (derived from azetidine).
30 An example of a 5 membered heterocyclic group is thiazolyl and an example of a 10 membered heterocyclic group is quinoliny. Examples of non-aromatic heterocyclic groups are pyrrolidiny, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperaziny, azetidiny, oxetanyl, thietanyl, homopiperidiny, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny, 1,2,3,6-
35 tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl, pyrazoliny, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidiny, imidazoliny, imidazolidiny, 3-azabicyclo[3.1.0]hexanyl, 3-
40

5 azabicyclo[4.1.0]heptanyl, 3H-indolyl and quinoliziny. Examples of aromatic heterocyclic groups are pyridinyl, imidazolyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl, indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, thiadiazolyl, furazanyl, benzofurazanyl, 10 benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, and furopyridinyl. The foregoing groups, as derived from the compounds listed above, may be C-attached or N-attached where such is possible. For instance, a group derived from pyrrole may be pyrrol-1-yl (N-attached) or pyrrol-3-yl (C-attached).

The phrase "pharmaceutically acceptable salt(s)", as used herein, unless otherwise 15 indicated, includes salts of acidic or basic groups which may be present in the compounds of formula 1. The compounds of formula 1 that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds of formula 1 are those that form non-toxic acid addition salts, *i.e.*, salts containing pharmacologically acceptable anions, 20 such as the acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, ethylsuccinate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, 25 mucate, napsylate, nitrate, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodode, and valerate salts.

Those compounds of the formula 1 that are acidic in nature, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali 30 metal or alkaline earth metal salts and particularly, the sodium and potassium salts.

The present invention also includes all radiolabelled forms of the compounds of formula 1, and pharmaceutically acceptable salts thereof, wherein the radiolabel is selected from ³H, ¹¹C and ¹⁴C. Such radiolabelled compounds are useful as research or diagnostic tools.

Certain compounds of formula 1 may have asymmetric centers and therefore exist in 35 different enantiomeric forms. This invention relates to the use of all optical isomers and stereoisomers of the compounds of formula 1 and mixtures thereof. In particular, the invention includes both the E and Z isomers of the -OR¹ group (where X² is =NOR¹) connected to the nitrogen of the oxime moiety at C-9 of the macrolide ring of formula 1.

The present invention also includes isotopically-labelled compounds, and the 40 pharmaceutically acceptable salts thereof, which are identical to those recited in formula 1, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass

5 number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Compounds of the present invention, prodrugs thereof, and pharmaceutically acceptable salts of said compounds or of said prodrugs which contain the
10 aforementioned isotopes and/or other isotopes of other atoms are within the scope of this invention. Certain isotopically-labelled compounds of the present invention, for example those into which radioactive isotopes such as ^3H and ^{14}C are incorporated, are useful in drug and/or substrate tissue distribution assays. Tritiated, i.e., ^3H , and carbon-14, i.e., ^{14}C , isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with
15 heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo* half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. Isotopically labelled compounds of formula 1 of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the Schemes and/or in the Examples and Preparations
20 below, by substituting a readily available isotopically labelled reagent for a non-isotopically labelled reagent.


This invention also encompasses pharmaceutical compositions containing and methods of treating bacterial infections through administering prodrugs of compounds of the formula 1. Compounds of formula 1 having free amino, amido, hydroxy or carboxylic groups can be
25 converted into prodrugs. Prodrugs include compounds wherein an amino acid residue, or a polypeptide chain of two or more (e.g., two, three or four) amino acid residues is covalently joined through an amide or ester bond to a free amino, hydroxy or carboxylic acid group of compounds of formula 1. The amino acid residues include but are not limited to the 20 naturally occurring amino acids commonly designated by three letter symbols and also includes 4-hydroxyproline,
30 hydroxylysine, demosine, isodemosine, 3-methylhistidine, norvalin, beta-alanine, gamma-aminobutyric acid, citrulline homocysteine, homoserine, ornithine and methionine sulfone. Additional types of prodrugs are also encompassed. For instance, free carboxyl groups can be derivatized as amides or alkyl esters. Free hydroxy groups may be derivatized using groups including but not limited to hemisuccinates, phosphate esters, dimethylaminoacetates, and
35 phosphoryloxymethyloxycarbonyls, as outlined in *Advanced Drug Delivery Reviews*, 1996, 19, 115. Carbamate prodrugs of hydroxy and amino groups are also included, as are carbonate prodrugs, sulfonate esters and sulfate esters of hydroxy groups. Derivatization of hydroxy groups as (acyloxy)methyl and (acyloxy)ethyl ethers wherein the acyl group may be an alkyl ester, optionally substituted with groups including but not limited to ether, amine and carboxylic acid functionalities,
40 or where the acyl group is an amino acid ester as described above, are also encompassed. Prodrugs of this type are described in *J. Med. Chem.* 1996, 39, 10. Free amines can also be

- 5 derivatized as amides, sulfonamides or phosphonamides. All of these prodrug moieties may incorporate groups including but not limited to ether, amine and carboxylic acid functionalities.

5


Detailed Description of the Invention

The preparation of the compounds of the present invention is illustrated in the following Schemes.

Scheme 1


5

Scheme 2




5

Scheme 2 continued



5

Scheme 2 continued



5 The preparation of the compounds of the present invention is illustrated in the above schemes. Starting materials and/or final compounds of formula 1 wherein R¹⁰ is a moiety other than ethyl within the definition of R¹⁰ provided above may be prepared as described in PCT published applications WO 98/01571 (Biotica Tech. Ltd. and Pfizer Inc.) and WO 98/01546 (assigned to Biotica Tech. Ltd.). Other specific methods that relate to the synthesis of the
10 compounds of the present invention are referred to in PCT international patent application publication number WO 98/38199 (published September 3, 1998), PCT international patent application publication number WO 98/56800 (published December 17, 1998), United States provisional patent application number 60/101,263 (filed September 22, 1998), United States provisional patent application number 60/111,728 (filed December 10, 1998), European patent
15 application number EP 487,411, and European patent application number EP 799,833. In the above Schemes, all substituents are as defined for formula 1 referred to above except otherwise indicated.

 The starting materials may or may not require proper functional group protection before various modifications can take place, and deprotection after desired modifications are complete.
20 The most commonly used protecting groups for amino moieties in the macrolide compounds of this invention are benzyloxycarbonyl (Cbz) and *t*-butyloxycarbonyl (Boc) groups. Hydroxyl groups are generally protected as acetates, Cbz carbonates or with a trialkylsilyl group. The C-2' hydroxyl group is a potentially reactive hydroxyl group among the numerous hydroxyl groups present in macrolide compounds of the type claimed herein. The C-2' hydroxyl group is
25 selectively protected by treating the compound with one equivalent of acetic anhydride in dichloromethane in the absence of external base. This process selectively converts the C-2' hydroxyl group into the corresponding acetate. The hydroxyl protecting group can be removed by treating the compound with methanol at a temperature ranging from about 0°C to 40°C to about 65°C for 10 to 48 hours. Other methods of selective protection and deprotection are
30 familiar to those skilled in the art.

 With reference to Scheme 1, the compound of formula 5, wherein R¹¹ is a halo group and all other substituents are as defined above, may be prepared by treating the compound of formula 4 with a base, such as sodium hydride, potassium hydride, potassium hexamethyldisilazide (KHMDs), pyridine, sodium carbonate, or lithium diisopropylamide, preferably KHMDs, and a halogenating agent, such as N-fluorobenzenesulfoimide, SELECTFLUOR™ (marketed by Air Products and Chemicals, Inc., Allentown, Pennsylvania, United States of America) for fluorination, pyridinium tribromide or cyanogen bromide for bromination, or hexachloroethane for chlorination, in a solvent, such as in N,N-dimethylformamide (DMF), tetrahydrofuran (THF), CH₂Cl₂, or N-methylpyrrolidone, or a mixture
40 of the foregoing solvents, preferably DMF. The reaction temperature, which is highly dependent on the reagent used, can be from -78°C to 60°C. In this step, R⁸ is preferably a hydroxy

5 protecting group such as an acetyl group, a benzyl group, or a trialkylsilyl group. To provide the compound of formula 6, deprotection of the C-2' hydroxy may proceed using methanol if R^8 is an acetyl group, hydrogenation if R^8 is an benzyl group, or fluoride anion, such as tetrabutylammonium fluoride, if R^8 is a trialkylsilyl group. The compound of formula 6 corresponds to the compound of formula 1 wherein R^8 is H.

10 Scheme 2 illustrates a method of preparing the compounds of the present invention by introducing the R^{11} group at an early step in the synthesis of the final compounds. In step 1 of Scheme 2, a R^{11} halo group may be introduced according to essentially the same procedure described above for Scheme 1. In step 2 of Scheme 2, the compound of formula 9 may be prepared by treating the compound of formula 8 with a base such as 1,8-
15 diazabicyclo[5.4.0]undec-7-ene (DBU) and 1,1'-carbonyl-diimidazole (CDI) in methylene chloride. Treatment of the compound of formula 9 with hydrazine in acetonitrile at about 60°C provides the cyclic carbazate of formula 10. Treatment of the compound formula 10 with O-alkylhydroxyamine in ethanol provides the oxime of formula 11. A reductive amination with an appropriate aldehyde of the formula $R^2-C(O)H$ and deprotection, if desired, of the C-2' hydroxy
20 group provides the compound of the formula 12 which corresponds to the compound of the formula 1 wherein X^1 is $-NH-$ and X^2 is $=NOR^1$. The compound of formula 10 may also be converted to a compound of formula 14 wherein X^1 is $-NH-$, as indicated in step 4' of scheme 2 above, by treating the compound of formula 10 with an appropriate heterocycle, such as a substituted imidazole, and an α,β -unsaturated aldehyde such as acrolein in acetic acid followed
25 by reduction with sodium borohydride.

Scheme 3 illustrates preparing compounds of the formula 13 which correspond to compounds of the formula 1 wherein X^2 is $=O$. In this process, the compound of formula 9 may be prepared as described above. The compound of formula 9 may be converted to the compound of formula 13 wherein X^1 is O, CR^4R^5 , or NR^4 by treating the compound of formula 9
30 with $NH_2-X^1-R^2$ wherein X^1 is O, CR^4R^5 , or NR^4 .

The compounds of the present invention may have asymmetric carbon atoms. Such diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, for example, by chromatography or fractional crystallization. Enantiomers can be separated by converting the
35 enantiomeric mixtures into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., alcohol), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. All such isomers, including diastereomeric mixtures and pure enantiomers are considered as part of the invention.

The compounds of formula 1 that are basic in nature are capable of forming a wide variety
40 of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for administration to animals, it is often desirable in practice to initially

5 isolate the compound of formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen
10 mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The desired acid salt can also be precipitated from a solution of the free base in an organic solvent by adding to the solution an appropriate mineral or organic acid.

Those compounds of the formula 1 that are acidic in nature are capable of forming base
15 salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts may be prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula 1. Such non-toxic base salts include those
20 derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc. These salts can be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the
25 desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.

The activity of the compounds of the present invention against bacterial and protozoa
30 pathogens is demonstrated by the compound's ability to inhibit growth of defined strains of human (Assay I) or animal (Assays II and III) pathogens.

Assay I

Assay I, described below, employs conventional methodology and interpretation criteria and is designed to provide direction for chemical modifications that may lead to compounds that
35 circumvent defined mechanisms of macrolide resistance. In Assay I, a panel of bacterial strains is assembled to include a variety of target pathogenic species, including representatives of macrolide resistance mechanisms that have been characterized. Use of this panel enables the chemical structure/activity relationship to be determined with respect to potency, spectrum of activity, and structural elements or modifications that may be necessary to obviate resistance
40 mechanisms. Bacterial pathogens that comprise the screening panel are shown in the table below. In many cases, both the macrolide-susceptible parent strain and the macrolide-resistant

- 5 strain derived from it are available to provide a more accurate assessment of the compound's ability to circumvent the resistance mechanism. Strains that contain the gene with the designation of *ermA/ermB/ermC* are resistant to macrolides, lincosamides, and streptogramin B antibiotics due to modifications (methylation) of 23S rRNA molecules by an Erm methylase, thereby generally prevent the binding of all three structural classes. Two types of macrolide
- 10 efflux have been described; *msrA* encodes a component of an efflux system in staphylococci that prevents the entry of macrolides and streptogramins while *mefA/E* encodes a transmembrane protein that appears to efflux only macrolides. Inactivation of macrolide antibiotics can occur and can be mediated by either a phosphorylation of the 2'-hydroxyl (*mph*) or by cleavage of the macrocyclic lactone (esterase). The strains may be characterized using
- 15 conventional polymerase chain reaction (PCR) technology and/or by sequencing the resistance determinant. The use of PCR technology in this application is described in J. Sutcliffe et al., "Detection Of Erythromycin-Resistant Determinants By PCR", Antimicrobial Agents and Chemotherapy, 40(11), 2562-2566 (1996). The assay is performed in microtiter trays and interpreted according to Performance Standards for Antimicrobial Disk Susceptibility Tests -
- 20 Sixth Edition; Approved Standard, published by The National Committee for Clinical Laboratory Standards (NCCLS) guidelines; the minimum inhibitory concentration (MIC) is used to compare strains. Compounds are initially dissolved in dimethylsulfoxide (DMSO) as 40 mg/ml stock solutions.

Strain Designation	Macrolide Resistance Mechanism(s)
Staphylococcus aureus 1116	susceptible parent
Staphylococcus aureus 1117	ermB
Staphylococcus aureus 0052	susceptible parent
Staphylococcus aureus 1120	ermC
Staphylococcus aureus 1032	msrA, mph, esterase
Staphylococcus hemolyticus 1006	msrA, mph
Streptococcus pyogenes 0203	susceptible parent
Streptococcus pyogenes 1079	ermB
Streptococcus pyogenes 1062	susceptible parent
Streptococcus pyogenes 1061	ermB
Streptococcus pyogenes 1064	ermB
Streptococcus agalactiae 1024	susceptible parent
Streptococcus agalactiae 1023	ermB
Streptococcus pneumoniae 1016	susceptible
Streptococcus pneumoniae 1046	ermB
Streptococcus pneumoniae 1095	ermB

Streptococcus pneumoniae 1175	mefE
Streptococcus pneumoniae 0085	susceptible
Haemophilus influenzae 0131	susceptible
Moraxella catarrhalis 0040	susceptible
Moraxella catarrhalis 1055	erythromycin intermediate resistance
Escherichia coli 0266	susceptible

5

Assay II is utilized to test for activity against *Pasteurella multocida* and Assay III is utilized to test for activity against *Pasteurella haemolytica*.

Assay II

This assay is based on the liquid dilution method in microliter format. A single colony of *P. multocida* (strain 59A067) is inoculated into 5 ml of brain heart infusion (BHI) broth. The test compounds are prepared by solubilizing 1 mg of the compound in 125 μ l of dimethylsulfoxide (DMSO). Dilutions of the test compound are prepared using uninoculated BHI broth. The concentrations of the test compound used range from 200 μ g/ml to 0.098 μ g/ml by two-fold serial dilutions. The *P. multocida* inoculated BHI is diluted with uninoculated BHI broth to make a 10^4 cell suspension per 200 μ l. The BHI cell suspensions are mixed with respective serial dilutions of the test compound, and incubated at 37°C for 18 hours. The minimum inhibitory concentration (MIC) is equal to the concentration of the compound exhibiting 100% inhibition of growth of *P. multocida* as determined by comparison with an uninoculated control.

Assay III

This assay is based on the agar dilution method using a Steers Replicator. Two to five colonies isolated from an agar plate are inoculated into BHI broth and incubated overnight at 37°C with shaking (200 rpm). The next morning, 300 μ l of the fully grown *P. haemolytica* preculture is inoculated into 3 ml of fresh BHI broth and is incubated at 37°C with shaking (200 rpm). The appropriate amounts of the test compounds are dissolved in ethanol and a series of two-fold serial dilutions are prepared. Two ml of the respective serial dilution is mixed with 18 ml of molten BHI agar and solidified. When the inoculated *P. haemolytica* culture reaches 0.5 McFarland standard density, about 5 μ l of the *P. haemolytica* culture is inoculated onto BHI agar plates containing the various concentrations of the test compound using a Steers Replicator and incubated for 18 hours at 37°C. Initial concentrations of the test compound range from 100-200 μ g/ml. The MIC is equal to the concentration of the test compound exhibiting 100% inhibition of growth of *P. haemolytica* as determined by comparison with an uninoculated control.

The *in vivo* activity of the compounds of formula 1 can be determined by conventional animal protection studies well known to those skilled in the art, usually carried out in mice.

5 Mice are allotted to cages (10 per cage) upon their arrival, and allowed to acclimate for a minimum of 48 hours before being used. Animals are inoculated with 0.5 ml of a 3×10^3 CFU/ml bacterial suspension (*P. multocida* strain 59A006) intraperitoneally. Each experiment has at least 3 non-medicated control groups including one infected with 0.1X challenge dose and two infected with 1X challenge dose; a 10X challenge data group may also be used. Generally, all mice in a given study can be challenged within 30-90 minutes, especially if a repeating syringe (such as a Cornwall® syringe) is used to administer the challenge. Thirty minutes after challenging has begun, the first compound treatment is given. It may be necessary for a second person to begin compound dosing if all of the animals have not been challenged at the end of 30 minutes. The routes of administration are subcutaneous or oral doses. Subcutaneous doses are administered into the loose skin in the back of the neck whereas oral doses are given by means of a feeding needle. In both cases, a volume of 0.2 ml is used per mouse. Compounds are administered 30 minutes, 4 hours, and 24 hours after challenge. A control compound of known efficacy administered by the same route is included in each test. Animals are observed daily, and the number of survivors in each group is recorded. The *P. multocida* model monitoring continues for 96 hours (four days) post challenge.

The PD_{50} is a calculated dose at which the compound tested protects 50% of a group of mice from mortality due to the bacterial infection which would be lethal in the absence of drug treatment.

25 The compounds of formula 1, and the pharmaceutically acceptable salts and solvates thereof (hereinafter "the active compounds"), may be administered through oral, parenteral, topical, or rectal routes in the treatment or prevention of bacterial or protozoa infections. In general, these compounds are most desirably administered in dosages ranging from about 0.2 mg per kg body weight per day (mg/kg/day) to about 200 mg/kg/day in single or divided doses (i.e., from 1 to 4 doses per day), although variations will necessarily occur depending upon the species, weight and condition of the subject being treated and the particular route of administration chosen. However, a dosage level that is in the range of about 4 mg/kg/day to about 50 mg/kg/day is most desirably employed. Variations may nevertheless occur depending upon the species of mammal, fish or bird being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effects, provided that such larger doses are first divided into several small doses for administration throughout the day.

40 The active compounds may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by the routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, the active

5 compounds may be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic
10 solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the active compounds are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed
15 along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred
20 materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active compound may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

25 For parenteral administration, solutions of an active compound in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of
30 all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

Additionally, it is also possible to administer the active compounds of the present invention topically and this may be done by way of creams, jellies, gels, pastes, patches, ointments and the like, in accordance with standard pharmaceutical practice.



35 For administration to animals other than humans, such as cattle or domestic animals, the active compounds may be administered in the feed of the animals or orally as a drench composition.

The active compounds may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles.
40 Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

5 The active compounds may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide phenyl, polyhydroxyethylaspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoylresidues. Furthermore, the active compounds may be coupled to a class of biodegradable polymers useful in achieving controlled
10 release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

 The Examples provided below illustrate specific embodiments of the invention, but the
15 invention is not limited in scope to the Examples specifically exemplified.

 The Example provided below illustrates a specific embodiment of the invention, but the invention is not limited in scope to the Example specifically exemplified. In the following example, "Ac" represents an acetyl group, "Me" represents a methyl group, and "Et" represents an ethyl group.



Example 1


To a solution containing the compound of formula **30** above (wherein "Ac" represents an acetyl group)(513 mg, 0.58 mmol) in 5.8 mL of DMF were added at -78°C 1.74 mL of 0.5 M solution of KHMDS in toluene (0.87 mmol). SELECTFLUOR™ (marketed by Air Products and Chemicals, Inc., Allentown, Pennsylvania, United States of America)(236 mg , 0.87 mmol) was added to this solution after 30 minutes of stirring at -78°C. Fresh SELECTFLUOR™ (27 mg,

5 0.076 mmol) was added after 30 minutes of stirring at -78°C . After an additional 30 minutes of stirring at the same temperature, the reaction mixture was diluted with EtOAc (ethyl acetate) and washed with water and brine. Drying over sodium sulfate and removal of the solvent gave 477 mg (93%) of a compound corresponding to formula 31 above except with the C-2' hydroxy protected with an acetyl group.

10 This material was dissolved in 50 mL of MeOH and warmed to 50°C overnight. Evaporation of the solvent and chromatography on SiO_2 gave the compound of formula 31 (which corresponds to the compound of formula 2 referred to above wherein R^{12} , R^{13} , R^{14} and R^{15} are each H); NMR (CDCl_3 , δ) 8.93 (1H, d), 8.42 (1H, dd), 8.04 (1H, dd), 7.57 (1H, s), 7.35 (1H, d), 7.24 (1H, dd), 6.13 (1H, s), 4.89 (1H, dd), 4.28 (1H, d), 4.19 (2H, m), 4.07 (1H, d), 3.69 (3H, s), 3.66 (1H, s), 3.56 (1H, m), 3.48 (1H, m), 3.41 (1H, m), 3.24 (1H, m), 2.76 (1H, m), 2.60 (2H, m), 2.57 (3H, s), 2.36 (6H, s), 1.93 (2H, m), 1.74 (3H, d), 1.76-1.20 (6H, m), 1.49 (3H, s), 1.34 (3H, s), 1.27 (3H, d), 1.22 (3H, d), 1.11 (3H, d), 0.98 (3H, d), 0.83 (3H, t).

Example 2

20 Following the procedures described in Scheme 2 above, a compound corresponding to formula 1 wherein X^1 is $-\text{CH}(\text{CH}_3)(\text{CH}_2)_2-$, X^2 is $=\text{NOCH}_3$, R^8 is H, R^9 is CH_3 , R^{10} is CH_2CH_3 , R^{11} is F, and R^{12} is 4-(pyridin-3-yl)-imidazol-1-yl, was prepared. MS 874 ($\text{M} + 1$)



32

The following examples, which each had a structure according to formula 32 above, were prepared following the procedure below which describes the preparation of the compound of example 3. Examples 4 to 10 were prepared using the same procedure and using an appropriate heterocyclic compound in place of phenyl imidazole.






5




Example 3

To a solution of compound 10 (100 mg, 0.155 mmole) and phenyl imidazole (67.0 mg, 0.465 mmole) in 1.5 mL of acetic acid was added 12.7 μ L of 90% acrolein (0.171 mmol). The resulting mixture was stirred at room temperature overnight. Sodium cyanoborohydride (46.7 mg, 0.775 mmol) was then added and the solution was stirred at room temperature over night.

10 The solution was diluted with water and its pH was adjusted to 10 with 40% aq NaOH solution. The aqueous solution was extracted with methylene chloride. The combine methylene chloride layer was dried over sodium sulfate and the solvent was removed under reduced pressure. The residue was purified by TLC (89% methylene chloride-10% methanol-1% ammonium hydroxide) to give 6 mg (5% yield) of example 3.

15


Example	R	Mass Spec. (M + 1)
3		830
4		870
5		821
6		867
7		793

8	 <chem>c1ccccc1c2nn[nH]n2</chem>	832
9	 <chem>Clc1ccccc1c2nn[nH]n2</chem>	867
10	 <chem>CC(=O)NCCc1cncn1</chem>	839

5

CLAIMS

1. A compound of the formula

**1**

or a pharmaceutically acceptable salt or solvate thereof, wherein:

X^1 is O, $-CR^4R^5$ - or $-NR^4$ -;

10 X^2 is $=O$ or $=NOR^1$;

R^1 is H or C_1 - C_{10} alkyl, wherein 1 to 3 carbons of said alkyl are optionally replaced by a heteroatom selected from O, S and $-N(R^4)$ -, and said alkyl is optionally substituted by 1 to 3 substituents independently selected from the group consisting of $-C(O)O(C_1-C_{10}$ alkyl), C_1 - C_{10} alkoxy, C_1 - C_{10} alkanoyl, halo, nitro, cyano, 4-10 membered heterocyclyl, C_1 - C_{10} alkyl, $-NR^4R^5$, C_6 - C_{10} aryl, $-S(O)_n(C_1-C_{10}$ alkyl) wherein n is an integer ranging from 0 to 2, and $-SO_2NR^4R^5$;

R^2 is $-(CR^4R^5)_n$ (4-10 membered heterocyclic) or $-(CR^4R^5)_n$ (C_6 - C_{10} aryl), wherein n is an integer from 0 to 6, from 1 to 3 R^4 or R^5 groups of the $-(CR^4R^5)_n$ - moiety of the foregoing R^2 groups are optionally replaced with a halo substituent, and the heterocyclic and aryl moieties of the foregoing R^2 groups are optionally substituted with 1 to 4 R^3 groups;

20 each R^3 is independently selected from halo, cyano, nitro, trifluoromethoxy, trifluoromethyl, azido, hydroxy, C_1 - C_6 alkoxy, C_1 - C_{10} alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $-C(O)R^6$, $-C(O)OR^6$, $-OC(O)R^6$, $-NR^6C(O)R^7$, $-NR^6C(O)NR^1R^7$, $-NR^6C(O)OR^7$, $-C(O)NR^6R^7$, $-NR^6OR^7$, $-SO_2NR^6R^7$, $-S(O)_j(C_1-C_6$ alkyl) wherein j is an integer from 0 to 2, $-(CR^1R^2)_l$ (C_6 - C_{10} aryl), $-(CR^4R^5)_l$ (4-10 membered heterocyclic), $-(CR^4R^5)_qC(O)(CR^4R^5)_l$ (C_6 - C_{10} aryl), $-(CR^4R^5)_qC(O)(CR^4R^5)_l$ (4-10 membered heterocyclic), $-(CR^4R^5)_lO(CR^4R^5)_q$ (C_6 - C_{10} aryl), $-(CR^4R^5)_lO(CR^4R^5)_q$ (4-10 membered heterocyclic), $-(CR^4R^5)_qSO_2(CR^4R^5)_l$ (C_6 - C_{10} aryl), and $-(CR^4R^5)_qSO_2(CR^4R^5)_l$ (4-10 membered heterocyclic), wherein q and t are each independently an integer from 0 to 5, 1 or 2 ring carbon atoms of the heterocyclic moieties of the foregoing R^{10} groups are optionally substituted by an oxo ($=O$) moiety, and the alkyl, alkenyl, alkynyl, aryl and heterocyclic moieties of the foregoing R^{10} groups are optionally substituted by 1 to 3 substituents independently selected from halo, cyano, nitro, trifluoromethyl, trifluoromethoxy,

- 5 azido, $-OR^6$, $-C(O)R^6$, $-C(O)OR^6$, $-OC(O)R^6$, $-NR^6C(O)R^7$, $-C(O)NR^6R^7$, $-NR^6R^7$, $-NR^6OR^7$, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, $-(CR^4R^5)_t(C_6-C_{10}$ aryl), and $-(CR^4R^5)_t(4-10$ membered heterocyclic), wherein t is an integer from 0 to 5;

each R^4 and R^5 is independently selected from H and C_1 - C_6 alkyl;

- each R^6 and R^7 is independently selected from H, C_1 - C_6 alkyl, $-(CR^4R^5)_t(C_6-C_{10}$ aryl),
 10 and $-(CR^4R^5)_t(4-10$ membered heterocyclic), wherein t is an integer from 0 to 5, 1 or 2 ring carbon atoms of the heterocyclic group are optionally substituted by an oxo ($=O$) moiety, and the alkyl, aryl and heterocyclic moieties of the foregoing R^6 and R^7 groups are optionally substituted by 1 to 3 substituents independently selected from halo, cyano, nitro, $-NR^4R^5$, trifluoromethyl, trifluoromethoxy, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, hydroxy, and C_1 - C_6
 15 alkoxy;

R^8 is H, $-C(O)(C_1-C_6$ alkyl), benzyl, benzyloxycarbonyl or $(C_1-C_6$ alkyl) $_3$ silyl;


R^9 is C_1 - C_6 alkyl;

R^{10} is H or C_1 - C_{10} alkyl; and

R^{11} is selected from chloro, bromo, iodo, fluoro, and cyano;

- 20 with the proviso that where X^2 is $=O$ then the R^{11} is cyano or from 1 to 3 R^4 or R^5 groups of the $-(CR^4R^5)_n$ - moiety of the R^2 groups recited above are either C_1 - C_6 alkyl or replaced with a halo substituent.

2. A compound of the formula




2

- 25 or a pharmaceutically acceptable salt or solvate thereof, wherein R^{12} , R^{13} , R^{14} and R^{15} are each independently selected from H, halo, methyl and ethyl.

3. A compound according to claim 2 wherein R^{14} and R^{15} are both H and R^{12} and R^{13} are each independently selected from H and methyl.

- 5 4. A compound according to claim 2 wherein R^{12} , R^{13} , R^{14} and R^{15} are each H.
 5. A compound of the formula




- or a pharmaceutically acceptable salt or solvate thereof, wherein R^{12} , R^{13} , R^{14} and R^{15} are each independently selected from H, halo, methyl and ethyl with the proviso that at least one of R^{12} , R^{13} , R^{14} and R^{15} is not H.

6. A compound according to claim 5 wherein R^{14} and R^{15} are both H, R^{12} is methyl, and R^{13} is selected from H and methyl.

7. A pharmaceutical composition for the treatment of a disorder selected from a bacterial infection, a protozoal infection, and a disorder related to a bacterial infection or protozoal infection in a mammal, fish, or bird which comprises a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

8. A method of treating a disorder selected from a bacterial infection, a protozoal infection, and a disorder related to a bacterial infection or protozoal infection in a mammal, fish, or bird which comprises administering to said mammal, fish or bird a therapeutically effective amount of a compound of claim 1.

9. A method of preparing a compound according to claim 1 wherein R^{11} is a halo group which comprises treating a compound of the formula



5

wherein R^8 , R^9 , R^{10} , X^1 , X^2 and R^2 are as defined in claim 1, with a halogenating agent.