



(72) CASTELEIJN, Eric, NL
(72) VAN DIJK, Willem R., NL
(72) KLUGKIST, Jan, NL
(72) VAN WASSENAAR, Pieter Dirk, NL
(71) UNILEVER PLC, GB
(51) Int.Cl.⁶ C11D 3/386
(30) 1995/11/27 (95203261.3) EP
(54) **COMPOSITIONS DETERGENTES ENZYMATIQUES**
(54) **ENZYMATIC DETERGENT COMPOSITIONS**

(57) L'invention porte sur une composition détergente enzymatique, renfermant un ou plusieurs surfactifs et une endoglucanase qui n'est pas une cellulase de la famille 7, qui ne contient pas de domaine de fixation de cellulose, mais dont le domaine catalytique renferme au moins deux ponts disulfure. Cette endoglucanase est produite de préférence à partir de Thermomonospora fusca, ou encore à partir de mutants ou de variants de celle-ci.

(57) There is provided an enzymatic detergent composition comprising one or more surfactants and an endoglucanase which is not a Family 7 cellulase, which contains no cellulose binding domain and wherein the catalytic domain contains at least two disulphide bridges. Preferably, the endoglucanase is producible from Thermomonospora fusca or mutants or variants thereof.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C11D 3/386, C12N 9/42	A1	(11) International Publication Number: WO 97/20026 (43) International Publication Date: 5 June 1997 (05.06.97)
(21) International Application Number: PCT/EP96/05053 ² (22) International Filing Date: 15 November 1996 (15.11.96) (30) Priority Data: 95203261.3 27 November 1995 (27.11.95) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except AU BB CA GB IE KE LK LS MN MW NZ SD SG SZ TT UG): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL). (71) Applicant (for AU BB CA GB IE KE LK LS MN MW NZ SD SG SZ TT UG only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB). (72) Inventors: CASTELEIJN, Eric; Unilever Research Vlaardingen Lab., Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). VAN DIJK, Willem, R.; Unilever Research Vlaardingen Lab., Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). KLUGKIST, Jan; Unilever Research Vlaardingen Lab., Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL). VAN WASSENAAR, Pieter, Dirk; Unilever Research Vlaardingen Lab., Olivier van Noortlaan 120, NL-3133 AT Vlaardingen (NL).		(74) Agent: UNILEVER N.V.; Patent Division, P.O. Box 137, NL-3130 AC Vlaardingen (NL). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ENZYMATIC DETERGENT COMPOSITIONS (57) Abstract There is provided an enzymatic detergent composition comprising one or more surfactants and an endoglucanase which is not a Family 7 cellulase, which contains no cellulose binding domain and wherein the catalytic domain contains at least two disulphide bridges. Preferably, the endoglucanase is producible from <i>Thermomonospora fusca</i> or mutants or variants thereof.		

ENZYMATIC DETERGENT COMPOSITIONSTECHNICAL FIELD

The present invention generally relates to the field of enzymatic detergent and cleaning compositions. More in particular, the invention is concerned with enzymatic detergent compositions for fabric washing, comprising an endoglucanase.

10 BACKGROUND AND PRIOR ART

Various types of enzymes are known in the art as additives for detergent compositions. For example, detergent compositions containing proteases, lipases, amylases and cellulases and various combinations thereof have been described in the literature and several such products are currently on the market. Of these enzymes, proteases, lipases and amylases are most abundantly used. The enzymes assist in the cleaning of fabrics by degrading their natural substrates protein, fat and starch. Cellulase, on the other hand, is not added to detergent products because of its capability to break down cellulose, but rather to attain certain "care" benefits such as colour clarification, anti-pilling and reduction of the harshness of the fabric.

The harshness-reducing action of cellulase in detergent compositions was first described GB-A-1 368 599 (Unilever). DE-A-3 207 847 (Kao) discloses that the addition of cellulase to a detergent product improves its cleaning performance. EP-A-220 016 (Novo Nordisk) describes a colour clarification activity of cellulases.

Cellulases occur in nature as very complex mixtures of enzymes and in recent years several attempts have been described to isolate its various components and to produce them by means of recombinant DNA techniques. For a classification of cellulases, see Henrissat and Bairoch, Biochemical Journal **293**, 781-788 (1993). A special class of cellulases, the endoglucanases, have been described as particularly useful for detergent applications.

W-A-89/09259 (Novo Nordisk) discloses a cellulase preparation useful for reducing the harshness of cotton-

containing fabrics, comprising at least 40% of an endoglucanase component with a high endoase activity and affinity towards cellulose. W-A-91/17243 (Novo Nordisk) discloses a cellulase preparation consisting essentially of a
5 homogeneous endoglucanase which is immunoreactive with or homologous to a 43 kD endoglucanase derived from Humicola insolens DSM 1800. The pH optimum of the endoglucanase from Humicola insolens DSM 1800 is about 8. W-A-94/21801 (Genencor) discloses the production and purification of
10 endoglucanase EGIII from Trichoderma longibrachiatum. This endoglucanase is said to have a pH optimum of 5.5-6.0.

W-A-95/24471 (Novo Nordisk) discloses that certain cellulases of Family 7 (in the classification according to Henrissat), which do not comprise a carbohydrate binding
15 domain, may have enhanced activity which may result in improved soil removal from fabrics.

Thus, although various endoglucanases have been reported to have favourable properties in detergent products, there is still a need to provide alternative or improved
20 endoglucanase containing detergent compositions. In particular, the storage stability of endoglucanases, as well as their stability in the presence of proteolytic enzymes and/or bleach leave to be desired, especially in liquid detergent formulations. There is also a need for detergent
25 products having improved anti-pilling properties.

For instance, it was found that the activity of the of Endoglucanase III from Trichoderma longibrachiatum, which has a pH optimum of 5.5-6.0, is rapidly decreasing in the alkaline region. Thus, there is also a particular need for
30 endoglucanase containing detergent compositions which exhibit typical cellulase-associated benefits at alkaline pH.

It is therefor an object of the present invention to provide a detergent composition for fabric washing, containing an endoglucanase that is stable in (liquid)
35 detergents during storage, in particular in the presence of proteolytic enzymes and/or bleach. It is a further object of the present invention to provide a detergent composition

containing an endoglucanase that has satisfactory anti-pilling properties and that is effective at alkaline pH.

We have now surprisingly found that these and other objects can be achieved by using in the composition an
5 endoglucanase which contains no cellulose binding domain and wherein the catalytic domain contains at least two and preferably three, four, five or even more disulphide bridges.

In particular, such endoglucanases can be used to formulate detergent compositions which are stable and exhibit
10 anti-pilling and colour clarification properties, even at alkaline pH and in the presence of proteolytic enzyme and/or bleach. We have also found that such endoglucanases do not depend on special proteases for stability, such as described in W-A-92/18599 (Novo Nordisk) for the 43 kD endoglucanase
15 derived from Humicola insolens DSM 1800.

DEFINITION OF THE INVENTION

According to a first aspect of the invention, there is provided an enzymatic detergent composition comprising one
20 or more surfactants and an endoglucanase which is not a Family 7 cellulase, which contains no cellulose binding domain and wherein the catalytic domain contains at least two and preferably three, four, five or even more disulphide bridges. Preferably, the endoglucanase consists essentially
25 of the catalytic domain of an endoglucanase from Thermomonospora fusca, or mutants or variants thereof.

According to a second aspect, the enzymatic detergent composition additionally comprises a proteolytic enzyme and/or bleach.

30

DESCRIPTION OF THE INVENTION

The detergent composition of the present invention comprises one or more surface active ingredients or surfactants and a specific type of endoglucanase. The
35 detergent compositions containing the special endoglucanases of the invention may be in any suitable physical form, such as a powder, an aqueous or non-aqueous liquid, a paste or a

gel. However, aqueous liquid detergents and highly alkaline powders are preferred. The storage stability of the special endoglucanase of the invention in isotropic liquid detergents was found to be exceptionally good. For liquid detergents, the pH of a solution of 1 gram of the detergent composition in 1 litre of water, with a hardness of 10° German before the addition of the detergent composition, at 20°C, is in the range of 7 to 11, preferably in the pH range of 8 to 10.5, more preferably 9 to 10.2.

10

(a) The surfactant

The compositions of the invention comprise, as a first ingredient, one or more surface active ingredients or surfactants. Depending on the physical type of detergent, the surfactants are present in an amount of 0.1 - 60 % by weight of the composition. Typically, an aqueous liquid detergent composition comprises from 5% to 50%, commonly at least 10% and up to 40%, by weight of one or more surface-active compounds. Fabric washing powders usually comprise from 20% to 45% by weight of one or more detergent-active compounds.

The compositions may comprise a single type of surfactant, mostly nonionics, but usually they contain a surfactant system consisting of 30-70 % by weight (of the system) of one or more anionic surfactants and 70-30 % by weight (of the system) of one or more nonionic surfactants. The surfactant system may additionally contain amphoteric or zwitterionic detergent compounds, but this is not normally desired owing to their relatively high cost.

In general, the nonionic and anionic surfactants of the surfactant system may be chosen from the surfactants described "Surface Active Agents" Vol. 1, by Schwartz & Perry, Interscience 1949, Vol. 2 by Schwartz, Perry & Berch, Interscience 1958, in the current edition of "McCutcheon's Emulsifiers and Detergents" published by Manufacturing Confectioners Company or in "Tenside-Taschenbuch", H. Stache, 2nd Edn., Carl Hauser Verlag, 1981.

Suitable nonionic detergent compounds which may be used include, in particular, the reaction products of com-

pounds having a hydrophobic group and a reactive hydrogen atom, for example, aliphatic alcohols, acids, amides or alkyl phenols with alkylene oxides, especially ethylene oxide either alone or with propylene oxide. Specific nonionic
5 detergent compounds are C_6-C_{22} alkyl phenol-ethylene oxide condensates, generally 5 to 25 EO, i.e. 5 to 25 units of ethylene oxide per molecule, and the condensation products of aliphatic C_8-C_{18} primary or secondary linear or branched
10 alcohols with ethylene oxide, generally 5 to 40 EO.

10 Suitable anionic detergent compounds which may be used are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl radicals containing from about 8 to about 22 carbon atoms, the term alkyl being used to include the alkyl portion of higher acyl radicals.
15 Examples of suitable synthetic anionic detergent compounds are sodium and potassium alkyl sulphates, especially those obtained by sulphating higher C_8-C_{18} alcohols, produced for example from tallow or coconut oil, sodium and potassium alkyl C_9-C_{20} benzene sulphonates, particularly sodium linear
20 secondary alkyl $C_{10}-C_{15}$ benzene sulphonates; and sodium alkyl glyceryl ether sulphates, especially those ethers of the higher alcohols derived from tallow or coconut oil and synthetic alcohols derived from petroleum. The preferred anionic detergent compounds are sodium $C_{11}-C_{15}$ alkyl benzene
25 sulphonates and sodium $C_{12}-C_{18}$ alkyl sulphates.

Also applicable are surfactants such as those described in EP-A-328 177 (Unilever), which show resistance to salting-out, the alkyl polyglycoside surfactants described in EP-A-070 074, and alkyl monoglycosides.

30 Preferred surfactant systems are mixtures of anionic with nonionic detergent active materials, in particular the groups and examples of anionic and nonionic surfactants pointed out in EP-A-346 995 (Unilever). Especially preferred is surfactant system which is a mixture
35 of an alkali metal salt of a $C_{16}-C_{18}$ primary alcohol sulphate together with a $C_{12}-C_{15}$ primary alcohol containing 3-7 ethoxylate groups.

(b) The enzyme

The compositions of the invention further comprise, as a second ingredient, a specific endoglucanase enzyme which is not a Family 7 cellulase, which contains no cellulose binding domain and wherein the catalytic domain contains at least two disulphide bridges.

It has been known for some time that cellulases consist of one or more building blocks having specific functions. See for instance, Saloheimo et al. (1993) (Proceedings of the second Tricel Symposium on *Trichoderma reesei*. Foundation for Biotechnical and Industrial Fermentation Research **8**, (1993), 139-146). Ginnis et al. (Biochemistry (1993) **32**, 8157-8161) describe that the cellulose binding domain of the endoglucanase E5 from *Thermomonospora fusca* is easily cleaved off, to form an endoglucanase which contains no cellulose binding domain and starting with threonine 121. Ginnis also disclose that the catalytic domain of E5 from *Thermomonospora fusca* contains four cysteine residues which form two disulphide bridges.

Preferably, the endoglucanases of the invention is derived from a cellulase selected from the group consisting of cellulases of Family 5, Family 6, Family 9, Family 12 and Family 45. Especially preferred are endoglucanases from *Thermomonospora fusca*, or mutants or variants thereof. This soil bacterium produces six different cellulases which are referred to in the literature as E1 to E6. All six enzymes contain a cellulose binding domain ("cbd") joined to the catalytic domain ("cd") by means of a flexible linker. Three of the cellulases are endoglucanases (E1, E2 and E5), two are exocellulases (E3 and E6) and one (E4) is an exocellulase with some endoglucanase activity. These specific enzymes and their production by means of recombinant DNA techniques have been described in the literature, see e.g. by Loa et al. Journal of Bacteriology (1991) **173**, 3397-3407. Using modern recombinant DNA techniques, the skilled man will have no difficulties in preparing the derivatives of these enzymes or mutants or variants of these enzymes which do not contain a cellulose binding domain.

In the following Table we have indicated for a number of endoglucanases whether they possess a cellulose binding domain ("cbd") and the number of cysteine residues and disulphide bridges.

5

TABLE

Endoglucanase (see also Example 1)	cbd	# cys- teines	# S-S bridges
Humicola insolens, 43kD	Y	20	?
10 Trichoderma longibrachiatum EGIII	N	2	1
KAC-500	Y	0	0
T. fusca E5	Y	6	3
T. fusca E5cd	N	4	2
15 T. fusca E2	Y	6	3
T. fusca E2cd	N	4	2

The number of cysteine residues and disulphide bridges for a given endoglucanase can be determined as described in Ginnis
20 et al. (Biochemistry (1993) **32**, 8157-8161).

In the context of the present invention, "mutants or variants" of Thermomonospora fusca endoglucanases are defined as endoglucanase enzymes which closely resemble the naturally occurring Thermomonospora fusca endoglucanases,
25 but are different in one or more amino acids, e.g. by substitution, deletion or insertion of one more amino acids. They will exhibit a high degree of homology (in terms of identity of residues) of at least 70%, preferably at least 80% or 90% or even 95% with the naturally occurring
30 Thermomonospora fusca endoglucanase.

Another way of defining "homology" is, that DNA encoding the variant or mutant endoglucanase will hybridize to the same probe as the DNA coding for the naturally occurring Thermomonospora fusca endoglucanase, under certain
35 specified conditions (i.e. presoaking in 5xSSC and

prehybridizing for 1 hour at 40°C in a solution of 20% formamide, 5x Denhard't solution, 50 mM sodium phosphate, pH 6.8 and 50 µg of denaturated calf thymus DNA, followed by hybridization in the same solution supplemented with ATP for 5 18 hours at 40°C).

Preferably, the endoglucanase consists essentially of the catalytic domain of endoglucanase of a Family 5 endoglucanase, more particular of an endoglucanase from Thermomonospora fusca such as E1, E2 or E5, or mutants or 10 variants thereof.

We have found that, although the endoase activity (as measured on CMC) of the endoglucanases of the present invention may be similar to that of the corresponding endoglucanase including the cellulose binding domain, the 15 depilling activity of the endoglucanase without its cbd may be considerably reduced. As a consequence, if a detergent product contains an endoglucanase with a cbd which is gradually cleaved off during storage, the depilling activity of the product may disappear almost unnoticed because there 20 is no corresponding decrease in endoase activity. In theory one could compensate for the loss in depilling action by increasing the amount of endoglucanase in the manufacturing stage, but then the endoase activity may be initially so high that an unacceptable tensile strength loss of cotton might 25 occur when the product is used shortly after its manufacture.

The endoglucanases of the present invention can be advantageously used to formulate safe detergent products which maintain their depilling activity upon storage, because the endoglucanases of the invention are resistant to 30 degradation and retain their depilling activity when the products containing them are stored.

The enzymatic detergent compositions of the invention comprise about 0.001 to 10 milligrams of the specific active endoglucanase protein per gram of detergent 35 composition. Preferably, they comprise 0.001 to 0.2 milligrams of active endoglucanase protein per gram of detergent composition, more preferably 0.005 to 0.04 milligrams per gram. More conveniently, the active cellulase

content is measured as enzyme activity on carboxymethyl cellulose (CMC). Expressed in CMC units, the compositions contain 0.06 - 600 CMCU per gram of detergent composition, preferably 0.06 - 12.5 CMCU per gram, and more preferably 0.3
5 - 2.5 CMCU/gram. In this specification the CMCU or carboxymethyl cellulose unit is measured according to the following protocol. The substrate used is a sodium salt of carboxymethylcellulose (CMC medium viscosity, Sigma catalogue number C4888). The CMC solution is stirred overnight or
10 heated for 30 minutes at 70°C to dissolve completely in 0.2 M sodium phosphate pH 7.0. 0.8 ml of the CMC solution is incubated with 0.2 ml enzyme/wash solution for 30 minutes at 40°C. Then the reaction is stopped by addition of 3 ml PahBah reagent (see below) and the amount of reducing sugars is
15 measured (Lever, 1972) Analytical Biochemistry **47**, 273-279). For the PahBah reagent 5 gram para-hydroxy-benzoic acidhydrazide (Sigma catalogue number H9882) is dissolved in 100 ml 0.5 N HCl and diluted with 400 ml 0.5 N NaOH prior to use. A calibration curve is prepared by dissolving 0, 10, 20,
20 30 and 40 µg/ml glucose in 0.2 M sodium phosphate pH 7.0. One ml of each glucose standard solution as well as 1 ml of the sample solutions (+ CMC) is mixed with 3 ml of the PahBah reagent. All mixtures are kept at 98°C for 5 minutes and then cooled (in water with ice). After cooling to room temperature
25 the light absorbance is spectrophotometrically measured at 405 nm. A calibration curve is obtained by plotting the amount of sugar against the OD405. The amount of sugars formed in the samples is then read from the curve and recalculated in to µmoles of glucose formed per minute
30 (CMCU). The activity is expressed as CMCU per gram of detergent composition or as CMCU per gram of enzyme protein (CMCU/g). Alternatively, it can be expressed as relative figure comparing residual activity to the activity that was originally added (CMCU%).

35 Also suitable for the present invention are endoglucanases having a high degree of homology of their amino acid sequence to the endoglucanases producible from

Thermomonospora fusca, provided that they have a similar or superior enzymatic activity.

The endoglucanase of the present invention can usefully be added to the detergent composition in any
5 suitable form, i.e. the form of a granular composition, a liquid or a slurry of the enzyme, or with carrier material (e.g. as in EP-A-258 068 and the Savinase (TM) and Lipolase (TM) products of Novo Nordisk). A good way of adding the enzyme to a liquid detergent product is in the form of a
10 slurry containing 0.5 to 50 % by weight of the enzyme in a ethoxylated alcohol nonionic surfactant, such as described in EP-A-450 702 (Unilever).

(c) Other ingredients.

15 The enzymatic detergent composition of the present invention may further contain from 5 - 60%, preferably from 20 - 50% by weight of a detergency builder. This detergency builder may be any material capable of reducing the level of free calcium ions in the wash liquor and will preferably
20 provide the composition with other beneficial properties such as the generation of an alkaline pH, the suspension of soil removed from the fabric and the suspension of the fabric-softening clay material.

Examples of detergency builders include
25 precipitating builders such as the alkali metal carbonates, bicarbonates, orthophosphates, sequestering builders such as the alkali metal tripolyphosphates, alkali metal citrates or nitrilotriacetates, or ion exchange builders such as the amorphous alkali metal aluminosilicates or the zeolites.

30 It was found to be especially favourable for the enzyme activity of the detergent compositions of the present invention if they contained a builder material such that the free calcium concentration is reduced to less than 1 mM.

The enzymatic detergent compositions of present
35 invention may also comprise, in further embodiments, combinations with other cellulolytic enzymes or other endoglucanases in an amount of at most up to 50%, preferably

up to 25%, more preferably less than 5% of the total amount of cellulolytic activity in the detergent composition.

The enzymatic detergent compositions of present invention may also comprise one or more other enzymes and
5 other constituents normally used in detergent systems, including additives for detergent compositions. Such other components can be any of many known kinds, for example enzyme stabilizers, lather boosters, soil-suspending agents, soil-release polymers, hydrotropes, corrosion inhibitors, dyes,
10 perfumes, silicates, optical brighteners, suds depressants, germicides, anti-tarnishing agents, opacifiers, fabric softening agents, oxygen-liberating bleaches such as hydrogen peroxide or sodium perborate, or sodium percarbonate, diperisophthalic anhydride, bleach precursors, oxygen-
15 activating bleaches, buffers and the like.

Examples are described in GB-A-1 372 034 (Unilever), US-A-3 950 277, US-A-4 011 169, EP-A-179 533 (Procter & Gamble), EP-A-205 208 and EP-A-206 390 (Unilever), JP-A-63-078000 (1988), and Research Disclosure 29056 of June
20 1988. The formulation of detergent compositions according to the invention can be also illustrated by reference to the Examples D1 to D14 of EP-A-407 225 (Unilever).

Special advantage may be gained in such detergent compositions wherein a proteolytic enzyme or protease is also
25 present. Proteases for use together with the endoglucanase can in certain circumstances include subtilisins of, for example, BPN' type or of many of the types of subtilisin disclosed in the literature, some of which have already been proposed for detergents use, e.g. mutant proteases as
30 described in for example EP-A-130 756 or EP-A-251 446 (both Genentech), US-A-4 760 025 (Genencor), EP-A-214 435 (Henkel), W-A-87/04661 (Amgen), W-A-87/05050 (Genex), Thomas et al. (1986) in Nature 5, 316, and 5, 375-376 and in J.Mol.Biol. (1987) 193, 803-813, Russel et al. (1987) in Nature 328,
35 496-500, and others.

Furthermore, certain polymeric materials such as polyvinyl pyrrolidones typically having a MW of 5,000 to about 30,000 are useful ingredients for preventing the

transfer of labile dye stuffs between fabrics during the washing process. Especially preferred are ingredients which also provide colour care benefits. Examples hereof are polyamide-N-oxide containing polymers. Also envisaged is the addition of peroxidase enzyme in combination with hydrogen peroxide and so-called enhancing intermediates. Finally, cellulases in general are said to provide a soil-release benefit in the wash and the present endoglucanases are no exception.

10 The invention will now be further illustrated in the following Examples.

EXAMPLE 1

15 Stability of cellulases in wash solutions.

The in-wash stability of endoglucanase according to the invention was compared with several prior art cellulases under the following conditions:

20 Enzymes

Endoglucanase E5cd from Thermomonospora fusca was obtained from Prof. D.B.Wilson, Cornell University, 458 Biotechnology Building, Ithaca NY, USA. The sample was substantially pure as measured by SDS polyacrylamide gel electrophoresis. The catalytic domain E5cd started with amino acid Gly97, as published in Biochemistry **32**, 8157-8161 (1993).

25 KAC-500 is a commercial endoglucanase ex. Kao produced by *Bacillus* sp. KSM-635 (Ozaki et al., J. of Gen. Microbiology **136**, 1327-1334 (1990) and Ito et al. Agric. Biol. Chem. **53**, 1275-1281 (1989)).

EGIII endoglucanase is a cellulase ex. Genencor International Inc. produced by *Trichoderma longibrachiatum* and described in W-A-94/21801 (Genencor).

35 Celluzyme is a commercial cellulase preparation ex. Novo Nordisk A/S produced by *Humicola insolens* DSM 1800 and described in US-A-4 435 307.

Cytolase 123 is a commercial cellulase preparation ex. Genencor International produced by *Trichoderma longibrachiatum*.

5 Detergents

Detergent A = Liquid detergent without enzymes (pH 8):

	Component	% (w/w)
	NaOH	0.93
	KOH	4.12
10	Citric acid (monohydrate)	5.5
	Propylene Glycol	0.8
	Glycerol	5.00
	Borax	3.50
	Polymer Narlex DC1	1.00
15	Nonionic.7EO (Synperonic A7)	18.4
	Priolene 6907	10.0
	Lialet 123 PAS	10.0
	PVP	0.5
	Perfume	<1.0
20	Antifoam	<0.5
	Dye	<0.5
	Water	up to 100%

Detergent B = powder detergent (pH 10.1):

25	Component	% (w/w)
	Linear PAS (Na salt of Coco alcohol derived sulphate)	6.37
	Nonionic.3EO (Synperonic A3)	8.05
	Nonionic.7EO (Synperonic A7)	6.37

	Soap	2.25
	Zeolite A24	38.84
	Sodium carbonate	1.27
	Dequest 2047	1.43
5	Sodium citrate 2aq.	23.47
	Antifoam granule	3.15
	Water/salts	up to 100%

Experimental

10 The experiments are carried out in a two litre thermostatted vessel, containing 1 litre of artificially hardened water (16°FH, prepared with CaCl_2 , MgCl_2 , Ca:Mg ratio 4:1). When the water reaches a temperature of 40°C the detergent is added to a concentration of 4 g/l. The cellulase is dosed

15 after 5 minutes at a concentration of 8 mg enzyme protein per litre. Immediately after mixing a sample is taken ($t=1$). Between $t=1$ and $t=40$ minutes samples are taken and measured for residual cellulase activity.

20 Activity measurements

The substrate used is a sodium salt of carboxymethyl-cellulose (CMC medium viscosity, Sigma catalogue number C4888). The CMC solution is stirred overnight or heated for 30 minutes at 70°C to dissolve completely in 0.2 M sodium

25 phosphate pH 7.0. 0.8 ml of the CMC solution is incubated with 0.2 ml enzyme/wash solution for 30 minutes at 40°C. Then the reaction is stopped by addition of 3 ml PahBah reagent (see below) and the amount of reducing sugars is measured (Lever, 1972) Analytical Biochemistry **47**, 273-279). For the

30 PahBah reagent 5 gram para-hydroxy-benzoic acid hydrazide (Sigma catalogue number H9882) is dissolved in 100 ml 0.5 N HCl and diluted with 400 ml 0.5 N NaOH prior to use. A calibration curve is prepared by dissolving 0, 10, 20, 30 and 40 $\mu\text{g/ml}$ glucose in 0.2 M sodium phosphate pH 7.0. One ml of

35 each glucose standard solution as well as 1 ml of the sample

solutions (+ CMC) is mixed with 3 ml of the PahBah reagent. All mixtures are kept at 98°C for 5 minutes and then cooled (in water with ice). After cooling to room temperature the light absorbance is measured spectrophotometrically at 405 nm. A calibration curve is obtained by plotting the amount of sugar against the OD405. The amount of sugars formed in the samples is then read from the curve and recalculated in to μ moles of glucose formed per minute (CMCU). Usually the activity is expressed as CMCU per gram of enzyme protein (CMCU/g) or as a percentage of the activity that was originally added (CMCU%).

Results

The in wash stability of the cellulases was determined as residual activity (in CMCU%) under the conditions indicated: In detergent A:

Time (minutes)	E5cd	EGIII	KAC-500	Cytolase 123	Celluzy me
1	100	100	100	100	100
5	100	95	66	93	100
10	93	90	47	88	102
15	92	81	30	82	99
20	97	78	20	79	102
30	92	73	7	70	102
40	86	69	5	55	103

In detergent B:

Time (minutes)	E5cd	EGIII	KAC-500	Cytolase 123	Celluzy me
1	100	100	38	100	100
5	101	89	4	95	100
10	101	83	3	86	88

15	101	72	1	80	79
20	103	64	2	81	75
30	103	54	2	72	68
40	101	47	2	69	58

5

EXAMPLE 2Stability of cellulases in wash solutions in the presence of proteolytic enzyme

10 The in wash stability of several prior art cellulases was compared with that of the endoglucanase of the invention, in the presence of proteolytic enzyme, under the following conditions:

Enzymes

15 As in Example 1.

The protease tested was Savinase 6.0T a commercial enzyme ex. Novo Nordisk A/S.

Detergents

20 As in Example 1.

Experimental

Experiments are carried out in a two litre thermostatted vessel, containing 1 litre of artificially hardened water
 25 (16°FH, prepared with CaCl₂, MgCl₂, Ca:Mg ratio 4:1). When the water reaches a temperature of 40°C the detergent is added to 4 g/l and 64.2 mg/l Savinase 6.0T. The cellulase is dosed after 5 minutes at a concentration of 8 milligrams enzyme protein per litre. Immediately after mixing a sample
 30 is taken (t=1). Between t=1 and t=40 minutes samples are taken. Protease activity is immediately inhibited by addition of phenyl methyl sulphonyl fluoride (PMSF). For this a stock solution of 20 mg of PMSF (Merck catalogue number 7349) in 1 ml of ethanol is prepared. Of this stock 0.125 ml is added to
 35 1 ml of sample. Then residual cellulase activity is measured as described in Example 1.

Results

The in-the-wash stability of the cellulases is determined as residual activity (in CMCU%) under the conditions indicated:
In detergent A + Savinase:

5	Time (minutes)	E5cd	EGIII	KAC-500	Cytolase 123	Celluzy me
	1	100	100	49	100	100
	5	100	99	4	4	100
	10	97	97	5	0	101
10	15	99	94	5	0	101
	20	98	n.d.	3	n.d.	99
	25	n.d.	85	n.d.	0	n.d.
	30	99	n.d.	0	n.d.	101
	40	100	78	0	0	103

15 n.d. = not determined

In detergent B + Savinase:

20	Time (minutes)	E5cd	EGIII	Cytolase 123
	1	100	100	100
	5	100	94	7
	10	112	85	7
	15	104	75	7
	20	104	66	6
25	30	103	56	n.d.
	40	100	45	n.d.

n.d. = not determined

EXAMPLE 3Stability of cellulases in wash solutions
in the presence of bleach

The in-the-wash stability of a cellulase of the invention was
5 compared with that of several prior art cellulases, under the
following conditions:

Enzymes

As in Example 1.

Detergents

10 As in Example 1.

Experimental

Experiments are carried out in a two litre thermostatted
vessel, containing 1 litre of artificially hardened water
(16°FH, prepared with CaCl₂, MgCl₂, Ca:Mg ratio 4:1). When
15 the water reaches a temperature of 40°C the detergent B is
added at 4 g/l and a dry mix of the bleach components giving
a final concentration of 0.26 g/l TAED (83%) + 0.82 g/l
percarbonate ISO694 + 0.017 g/l Dequest 2047. The cellulase
is dosed after 5 minutes at a concentration of 8 milligrams
20 enzyme protein per litre. Immediately after mixing a sample
is taken (t=1). Between t=1 and t=60 minutes samples are
taken. 20 g/l sodium sulphite (Merck 6652) is added to each
sample to reduce the bleach system. Then residual cellulase
activity is measured as described in Example 1.

25

Results

The in wash stability of the cellulases is determined as
residual activity (in CMCU%) under the conditions indicated:

30 In detergent B + bleach:

Time (minutes)	E5cd	EGIII	Celluzyme
1	100	100	100
5	100	96	89
10	98	81	79
15	102	n.d.	n.d.

35

20	97	54	60
30	94	34	48
40	95	21	43

n.d. = not determined

5

EXAMPLE 4

Stability of cellulases in wash solutions in the presence of bleach and proteolytic enzyme

The in-wash stability of EGIII cellulases was compared with a
10 cellulase of the invention under the following conditions:

Enzymes

As in Example 2.

Detergents

As in Example 1.

15 Experimental

Experiments are carried out in a two litre thermostatted vessel, containing 1 litre of artificially hardened water (16°FH, prepared with CaCl_2 , MgCl_2 , Ca:Mg ratio 4:1). When the water reaches a temperature of 40°C the detergent B is
20 added at 4 g/l then 64.2 mg/l Savinase 6.0T and a dry mix of the bleach components giving a final concentration of 0.26 g/l TAED (83%) + 0.82 g/l percarbonate ISO694 + 0.017 g/l Dequest 2047. The cellulase is dosed after 5 minutes at a concentration of 8 milligrams enzyme protein per litre.

25 Immediately after mixing a sample is taken (t=1). Between t=1 and t=60 minutes samples are taken. 20 g/l sodium sulphite (Merck 6652) is added to each sample to reduce the bleach system and 0.4 g/l trypsin inhibitor (Sigma T-9253) to inhibit the protease. Then residual cellulase activity is
30 measured as described in Example 1.

Results

The in wash stability of the cellulases is determined as residual activity (in CMCU%) under the conditions indicated:

35

20

In detergent B + bleach + Savinase:

Time (minutes)	E5cd	EGIII
1	100	100
5	68	89
10	86	63
20	83	19
30	86	6
40	80	4

EXAMPLE 5

Stability of cellulases in wash solutions in the presence
of lipase (and protease and/or bleach)

- 15 Similar results as in shown in Examples 1,2,3 and 4 are obtained when 0.37% Lipolase 100L is present in detergent A and 0.25% Lipolase 100T is present in detergent B.

In detergent A + Lipolase:

Time (minutes)	E5cd
1	100
5	100
10	99
15	95
20	85
30	98
40	72

21

In detergent B + Lipolase:

	Time (minutes)	E5cd
	1	100
5	5	100
	10	101
	15	107
	20	112
	30	98
10	40	97

In detergent A + bleach + Savinase + Lipolase:

	Time (minutes)	E5cd
15	1	100
	5	100
	10	100
	15	n.d.
	20	93
20	30	92
	40	90
	50	88
	60	85

n.d. = not determined

25

In detergent B + bleach + Savinase + Lipolase:

Time (minutes)	E5cd
1	100
5	100
10	86
15	73
20	74
30	87
40	69

EXAMPLE 6

Depilling of cotton by cellulase

- 15 The potential to remove pills from a cotton fabric in multiple washes of several prior art cellulases was compared with that of the cellulase of the invention, under the following conditions:

Enzymes

- 20 As in Example 2, except that "E5" was a preparation obtained from Alko Oy AB (Finland) and contained a mixture of E5 and E5cd. Celluzyme was dosed at 35 mg/l cellulase protein, KAC-500 and EGIH were dosed at 35 mg/l endoglucanase protein and E5 was dosed at 35 mg/l E5 protein and 65 mg/l E5cd protein.

Detergent

Detergent C = powder detergent (pH 9.4):

Component	% (w/w)
Linear PAS (Na salt of Coco alcohol derived sulphate)	10.67
Nonionic.3EO (Synperonic A3)	4.55
Nonionic.7EO (Synperonic A7)	6.83
Soap	1.77

	Zeolite A24	36.80
	Sodium carbonate	2.12
	Dequest 2047	0.00
	Sodium citrate 2aq.	20.93
5	Citric acid	3.00
	SCMC	1.01
	Antifoam granule	4.00
	Water/salts	up to 100%

10 Fabric

Cotton interlock was supplied scoured and bleached but without optical whitener by Phoenix Calico, Ashton-under-Lyme. The fabric possessed a definite "face" as one side had been raised during manufacture by light brushing. Further processing entailed jet-dyeing using Drimarene Brilliant Blue K-2R in the presence of 50 g/l Glauber's salt and 20 g/l soda ash followed by a hot rinse, soaping at the boil in the presence of 0.2 g/l Arkopan T and 0.5 g/l soda ash, two further rinses and stentor-drying. This cloth is further referred to as blue cotton interlock.

Prepilled cotton interlock

The blue cotton interlock was prepilled by washing 15 times in a Miele Automatic W 406 TMT washing machine for 30 minutes at 40°C in demineralised water. After every 5 wash cycles fabrics were dried in a Miele Novotronic T440C tumble dryer (programma extra dry). Each machine load comprised six pieces of the interlock fabric (length 2m, width 1.2 m) together with a dummy load of mixed cotton fabrics (terry, drill, sheeting) to bring the total mass of fabric in the drum up to 2.5 kg. After 15 wash cycles about 11 pills per square centimetre were visible on the fabric surface.

Calibration standards for pill score

Blue cotton interlock was prepilled as described above but using a variable number of wash cycles. Using image analyses a series of standards were prepared with an increasing number of pills. The increase in pilling for the standards was about linear with the scale number. Standards were scaled as 0,1,2,3,4 and 5, whereby 0 is untreated unpilled fabric and 5 is severely pilled fabric. Using this scale the above described, prepilled fabric would rank as 3.5.

10

Experimental

Prepilled blue cotton interlock was cut into pieces of 7.5 * 10 cm. Each piece of cloth was washed in 90 ml of detergent C (at 5 g/l in 16°FH tap water, Ca:Mg = 4:1) with or without cellulase (at 35 mg enzyme protein/l) in a 250 ml polyethylene bottle. 20 bottles were agitated simultaneously in a Miele Automatic W 406 TMT washing machine containing 2080 gram cotton dish cloth as ballast load. Bottles were agitated for 30 minutes at 40°C. Then the cloths were taken out of the bottles and rinsed in a bowl for 5 minutes in running tap water. The pH of residual suds after the wash was measured and found to be 9.2 +/- 0.2. Then cloths were dried in a Miele Novotronic T440C tumble dryer (extra dry). Then cloths were scored and washed again in the same detergent using the same protocol. The test continued until 10 wash/dry cycles.

20

25

Scoring procedure

The dry blue cotton was assessed by three separate persons. Each cloth had to be ranked using the above described scale. For each product 4 different test pieces were washed separately according to above described protocol. Results are presented as the average score for the 4 pieces as scored by the 3 panel members.

35

Result

In a comparative experiment the pill score of blue cotton interlock was measured for several cellulases using above

described test conditions. Results are given versus the number of wash cycles.

5	Cellulase	Number of wash/dry cycles and pills scores					
		5 cycles	6 cycles	7 cycles	8 cycles	9 cycles	10 cycles
	No cellulase	3.5	3.5	3.5	3.5	3.5	3.5
	KAC-500	3.3	3.3	3.3	3.3	3.3	3.3
	Celluzyme	3.4	2.9	3.0	2.8	2.8	2.0
10	E5/E5cd	1.5	1.5	1.3	0.1	0.0	0.0
	EGIII	2.9	3.0	3.1	3.3	3.4	3.4

EXAMPLE 7

Depilling in different detergents

15 Example 6 was repeated for E5cd, KAC-500 and Celluzyme using the following detergents at 4 g/l:

	Component	% (w/w)	
		Detergent D (pH 10)	Detergent E (pH 10)
	Linear PAS (Na salt of Coco alcohol derived sulphate)	14.67	0.00
20	Na-LAS (linear alkyl benzene sulphonate)	0.00	20.14
	Nonionic.7EO (Synperonic A7)	7.99	4.52
	Soap	2.19	1.66
	Zeolite A24	29.27	35.13
25	Sodium carbonate	2.91	12.8
	Sodium bicarbonate	0.00	3.82
	Dequest 2047 (33.5% A.I.)	1.40	0.00

	Sodium citrate 2aq.	31.12	0.00
	Citric acid	0.00	2.00
	SCMC	0.87	0.60
	PVP	0.00	0.47
5	Polymer CP5	0.00	4.33
	Na-silicate	0.00	0.47
	Savinase 6.0T	0.00	0.45
	Lipolase 100T	0.00	0.27
	Antifoam granule	4.00	2.00
10	Perfume	0.00	0.36
	Water/salts	up to 100%	up to 100%

Below, the results are given versus the number of wash cycles, for Detergent D:

15	Cellulase	Number of wash/dry cycles and pills scores					
		5	6	7	8	9	10
	No cellulase	3.5	3.5	3.5	3.5	3.5	3.5
	Celluzyme, 35 mg/l	3.5	3.4	3.4	3.4	3.5	3.3
	KAC-500, 35 mg/l endoglucanase	3.3	3.0	3.1	3.1	3.1	3.0
20	10 mg "E5"/l (3.5 mg/l E5 + 6.5 mg/l E5cd)	3.5	3.3	2.8	2.6	2.6	2.6
	30 mg "E5"/l (10 mg/l E5 + 20 mg/l E5cd)	3.0	2.8	2.6	2.5	2.6	2.6
25	33 mg "E5"/l (7 mg/l E5 + 26 mg/l E5cd)	3.0	2.9	2.9	2.6	2.6	2.5
	109 mg "E5"/l (22 mg/l E5 + 87 mg/l E5cd)	3.0	2.5	2.0	1.5	0.6	0.5

Below, the results are given versus the number of wash cycles, for Detergent E:

	Cellulase	Number of wash/dry cycles and pills scores					
		5	6	7	8	9	10
5	No cellulase	3.3	3.3	3.3	3.3	3.3	3.3
	Celluzyme, 35 mg/l	3.5	3.5	3.4	3.1	3.1	3.0
	KAC-500, 35 mg/l endoglucanase	3.5	3.1	3.0	3.3	3.3	3.1
10	10 mg "E5"/l (3.5 mg/l E5 + 6.5 mg/l E5cd)	3.5	3.3	2.8	2.6	2.6	2.8
	33 mg "E5"/l (7 mg/l E5 + 26 mg/l E5cd)	3.0	2.9	2.9	2.9	2.9	2.9
15	109 mg "E5"/l (22 mg/l E5 + 87 mg/l E5cd)	3.0	2.5	1.5	1.5	0.6	0.5

EXAMPLE 8

Depilling of cotton by cellulase in the presence of protease
 Example 6 was repeated for detergent C to which a mixture of protease, lipase and amylase was added. Product dosage was 5 g/l, the pH of the wash solution was 9.0.

Enzymes:

EGIII liquid was from Genencor, as described in example 1. It was dosed at 89 mg protein per litre wash solution.
 Endoglucanase E5 derived from Thermomonospora fusca was obtained from Alko. This sample was stored for 19 months at 4°C. After storage the sample of E5 gave a single band on SDS polyacrylamide gel electrophoresis at a molecular weight of 32,000 kD. N-terminal sequencing gave an amino acid sequence of T-Q-P-G-T-G-T-P-V-E-R-Y-G-K-V. This sequence is identical to that of E5cd starting with amino acid Thr121 as published in Biochemistry **32**, 8157-8161 (1993). E5cd obtained in this way was dosed at 50 mg/l, 150 mg/l and 250 mg protein per litre wash solution.

Detergent

Composition of detergent C was the same as described in Example 6 with additional enzymes:

0.37% Savinase 12TX + 0.17% Lipolase ultra 50T + 0.05%

- 5 Termamyl 60T. These enzymes are commercial detergent enzymes sold by Novo Nordisk, Denmark.

Experimental

The experimental part was a repeat of example 6 with some modifications. 3 instead of 4 different pieces of test cloth
10 were washed for each product. Size of the test cloths was 5cm x 5cm. Each cloth was washed in 30 ml wash liquor in a 100 ml bottle. Pill scores were made at the start and from the 5th wash onwards by 3 panel members. Results are given as average score.

15

Results

Cellulase	Number of wash/dry cycles and pill scores						
	start	5	6	7	8	9	10
20 No cellulase	3.5	3.1	3.6	3.4	3.3	3.5	3.5
EGIII 89 mg/l	3.5	3.1	3.3	2.9	2.4	2	1.3
E5cd 50 mg/l	3.5	2.8	2.9	3.1	2.8	2.6	2.5
25 E5cd 150 mg/l	3.5	2.7	2.6	2.1	1.5	1.2	0.6
E5cd 250 mg/l	3.5	1.9	1.4	0.5	0.2	0.1	0.2

CLAIMS

1. An enzymatic detergent composition comprising one or more surfactants and an endoglucanase which is not a Family 7 cellulase, which contains no cellulose binding domain and
5 wherein the catalytic domain contains at least two disulphide bridges.
2. A detergent composition according to claim 1, wherein the endoglucanase is derived from a cellulase selected from
10 the group consisting of cellulases of Family 5, Family 6, Family 9, Family 12 and Family 45.
3. A detergent composition according to any one of the preceding claims, wherein the endoglucanase consists
15 essentially of a catalytic domain.
4. A detergent composition according to any one of the preceding claims, wherein the endoglucanase is producible from Thermomonospora fusca, or mutants or variants thereof.
20
5. A detergent composition according to any one of the preceding claims, wherein the endoglucanase consists essentially of the catalytic domain of endoglucanase E5 from Thermomonospora fusca, or mutants or variants thereof.
25
6. A detergent composition according to any one of the preceding claims, further comprising a proteolytic enzyme.
7. A detergent composition according to claim 6, in which
30 the proteolytic enzyme is a subtilisin protease.
8. An enzymatic detergent composition according to any one of the preceding claims, characterized in that comprises:
(a) 0.1 - 60 % by weight of one or more a surfactants; and
35 (b) 0.06 to 600 CMCU per gram of the detergent composition of active endoglucanase;
whereby the pH of a solution of 1 gram of the detergent composition in 1 litre of water, with a hardness of 10°

30

German before the addition of the detergent composition, at 20°C, is in the range of 7 to 11.

9. A detergent composition according to any one of the
5 preceding claims, further comprising a bleaching system.

10. A detergent composition according to any one of the
preceding claims, in the form of an aqueous, isotropic
liquid.

10
