

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number
WO 03/013441 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: PCT/US02/25475
- (22) International Filing Date: 10 August 2002 (10.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|-----------------------------|----|
| 60/311,523 | 10 August 2001 (10.08.2001) | US |
| 60/311,552 | 10 August 2001 (10.08.2001) | US |
| 60/311,549 | 10 August 2001 (10.08.2001) | US |
| 60/311,522 | 10 August 2001 (10.08.2001) | US |
| 60/311,524 | 10 August 2001 (10.08.2001) | US |

North, Seminole, FL 33776 (US). **DIMENNA, Phillip, A.** [US/US]; 6773 32nd Avenue North, St. Petersburg, FL 33710 (US). **GEMMA, Rocco, L.** [US/US]; 4241 Dover-Zoar Road N.E., Dover, OH 44622 (US).

(74) Agent: **MANSO, Peter, J.**; Edwards & Angell, 600 Corporate Drive, Suite 514, Ft. Lauderdale, FL 33334 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **KING PHARMACEUTICALS, INC.** [US/US]; 501 Fifth Street, Bristol, TN 37620 (US).


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRANZ, G., Andrew** [US]; 2220 So. Warson Rd., St. Louis, MO 63124 (US). **STRAUSS, Elaine, A.** [US/US]; 9557 135th St.

[Continued on next page]

(54) Title: LEVOTHYROXINE COMPOSITIONS AND METHODS



(57) Abstract: The present invention generally relates to stable pharmaceutical compositions, and methods of making and administering such compositions. In one aspect, the invention features stabilized pharmaceutical compositions that include pharmaceutically active ingredients such as levothyroxine (T4) sodium and liothyronine (T3) sodium (thyroid hormone drugs), preferably in an immediate release solid dosage form. Also provided are methods for making and using such immediate release and stabilized compositions.

**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations*
- *of inventorship (Rule 4.17(iv)) for US only*

Published:

- *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5

10

LEVOTHYROXINE COMPOSITIONS AND METHODS

15

U.S. Patent Application

This application for U.S. patent is filed as an utility application under U.S.C., Title 35, §111(a).

20 **Related U.S. Patent Applications**

This application for U.S. patent relates to U.S. provisional application, which was filed on February 15, 2001 and assigned provisional Serial No. 60/269,009 and is entitled *Stabilized Pharmaceutical and Thyroid Hormone Compositions and Method of Preparation*. This application for U.S. patent also
25 relates to U.S. provisional application, which was filed on February 15, 2001 and assigned provisional Serial No. 60/268,998 and is entitled *Manufacture of Thyroid Hormone Tablets Having Consistent Active Moiety Amounts*.

Field of the Invention

The invention generally relates to stable pharmaceutical compositions, and methods of making and administering such compositions. In one aspect, the invention features stabilized pharmaceutical compositions that include
5 pharmaceutically active ingredients such as levothyroxine (T4) sodium and liothyronine (T3) sodium (thyroid hormone drugs), preferably in an immediate release solid dosage form. Also provided are methods for making and using such immediate release and stabilized compositions.

Background

Thyroid hormone preparations of levothyroxine sodium and liothyronine sodium are pharmaceutical preparations useful to the treatment of hypothyroidism and thyroid hormone replacement therapy in mammals, for example, humans and dogs.

15 Thyroid hormone preparations are used to treat reduced or absent thyroid function of any etiology, including human or animal ailments such as myxedema, cretinism and obesity.

Hypothyroidism is a common condition. It has been reported in the United States Federal Register that Hypothyroidism has a prevalence of 0.5 percent to 1.3
20 percent in adults. In people over 60, the prevalence of primary hypothyroidism increases to 2.7 percent in men and 7.1 percent in women. Because congenital hypothyroidism may result in irreversible mental retardation, which can be avoided with early diagnosis and treatment, newborn screening for this disorder is mandatory in North America, Europe, and Japan.

25 Thyroid hormone replacement therapy can be a chronic, lifetime endeavor. The dosage is established for each patient Individually. Generally, the initial dose is small. The amount is increased gradually until clinical evaluation and laboratory tests indicate that an optimal response has been achieved. The dose

required to maintain this response is then continued. The age and general physical condition of the patient and the severity and duration of hypothyroid symptoms determine the initial dosage and the rate at which the dosage may be increased to the eventual maintenance level. It has been reported that the dosage increase
5 should be very gradual in patients with myxedema or cardiovascular disease to prevent precipitation of angina, myocardial infarction, or stroke.

It is important that thyroid hormone treatment have the correct dosage. Both under-treatment and over-treatment can have deleterious health impacts. In the case of under-treatment, a sub-optimal response and hypothyroidism could
10 result. Under-treatment has also been reported to be a potential factor in decreased cardiac contractility and increased risk of coronary artery disease. Conversely, over-treatment may result in toxic manifestations of hyperthyroidism such as cardiac pain, palpitations, or cardiac arrhythmia's. In patients with coronary heart disease, even a small increase in the dose of levothyroxine sodium may be
15 hazardous in a particular.

Hyperthyroidism is a known risk factor for osteoporosis. Several studies suggest that sub clinical hyperthyroidism in premenopausal women receiving thyroid hormone drugs for replacement or suppressive therapy is associated with bone loss. To minimize the risk of osteoporosis, it is preferable that the dose be
20 kept to the lowest effective dose.

Because of the risks associated with over-treatment or under-treatment with levothyroxine sodium, there is a need for thyroid hormone products that are consistent in potency and bioavailability. Such consistency is best accomplished by manufacturing techniques that maintain consistent amounts of the active
25 moiety during tablet manufacture.

Thyroid hormone drugs are natural or synthetic preparations containing tetraiodothyronine (T_4 , levothyroxine) or triiodothyronine (T_3 , liothyronine) or both, usually as their pharmaceutically acceptable (e.g. sodium) salts. T_4 and T_3


are produced in the human thyroid gland by the iodination and coupling of the amino acid tyrosine. T_4 contains four iodine atoms and is formed by the coupling of two molecules of diiodotyrosine (DIT). T_3 contains three atoms of iodine and is formed by the coupling of one molecule of DIT with one molecule of
5 moniodotyrosine (MIT). Both hormones are stored in the thyroid colloid as thyroglobulin. Thyroid hormone preparations belong to two categories: (1) natural hormonal preparations derived from animal thyroid, and (2) synthetic preparations. Natural preparations include desiccated thyroid and thyroglobulin.

Desiccated thyroid is derived from domesticated animals that are used for
10 food by man (either beef or hog thyroid), and thyroglobulin is derived from thyroid glands of the hog. The United States Pharmacopoeia (USP) has standardized the total iodine content of natural preparations. Thyroid USP contains not less than (NLT) 0.17 percent and not more than (NMT) 0.23 percent iodine, and thyroglobulin contains not less than (NLT) 0.7 percent of organically
15 bound iodine. Iodine content is only an indirect indicator of true hormonal biologic activity.

Synthetic forms for both T_4 and T_3 thyroid hormone are available from a number of producers. For example, liothyronine sodium (T_3) tablets are available under the trademark Cytomel[®] from King Pharmaceuticals, Inc., St. Louis,
20 Missouri. Levothyroxine sodium (T_4) is available under the tradename Levoxyl[®] from King Pharmaceuticals, Inc., under the tradename Synthroid[®] from Knoll Pharmaceutical, Mt. Olive, New Jersey, and under the tradename Unithroid[®] from Jerome Stevens Pharmaceuticals, Bohemia, New York. In addition a veterinarian preparation of levothyroxine sodium is available under the tradename Soloxine[®]
25 from King Pharmaceuticals, Inc.

Levoxyl[®] (levothyroxine sodium tablets, USP) contain synthetic crystalline L-3,3',5,5'-tetraiodothyronine sodium salt [levothyroxine (T_4) sodium]. As indicated above, the synthetic T_4 in Levoxyl[®] is identical to that produced in the

human thyroid gland. The levothyroxine (T₄) sodium in Levoxyl® has an empirical formula of C₁₅H₁₀ I₄ N NaO₄ • H₂O, molecular weight of 798.86 g/mol (anhydrous), and a structural formula as shown:



5

It is well known that the stability of thyroid hormone drugs is quite poor. They are hygroscopic and degrade in the presence of moisture or light, and under conditions of high temperature. The instability is especially notable in the presence of pharmaceutical excipients, such as carbohydrates, including lactose, sucrose, dextrose and starch, as well as certain dyes. The critical nature of the dosage requirements, and the lack of stability of the active ingredients in the popular pharmaceutical formulations, have led to a crisis which has adversely effected the most prescribed thyroid drug products. See, e.g., 62 Fed. Reg. 43535 (Aug. 14, 1997).

It is desirable, therefore, to prepare a stabilized dosage of levothyroxine and liothyronine, which will have a longer shelf life that can be used in the treatment of human or animal thyroid hormone deficiency. U.S. Patent No. 5,225,204 (the '204 patent) is directed to improving the stability of levothyroxine sodium. In one embodiment disclosed by '204, stabilized levothyroxine sodium was prepared in a dry state by mixing levothyroxine sodium with a cellulose tableting agent using geometric dilution and subsequently combining this mixture with the same or a second cellulose tableting agent, such as microcrystalline

cellulose. Other tableting aids or excipients can be used in this formulation. This '204 patent is incorporated by reference herein, in its entirety.

The microcrystalline cellulose disclosed In '204 is AVICEL 101[®], AVICEL 102[®], AVICEL 103[®], AVICEL 105[®], trademarks of FMC Company of Newark, DE., and Microcrystalline Cellulose NF, or EMCOCEL[®], a trademark
5 owned by Penwest Pharmaceuticals of Patterson, NY. These microcrystalline cellulose products are prepared by re-slurrying the cellulose and spray drying the product. This produces an α - helix spherical microcrystalline cellulose product.

U. S. Patents 5,955,105 and 6,056,975 (the continuation of '105) disclose
10 pharmaceutical preparations of levothyroxine and microcrystalline cellulose, along with other excipients. The microcrystalline cellulose products used in the '105 and '975 patents were also the α - form Avicel microcrystalline cellulose products. U. S. Patents 5,955,105 and 6,056,975 are incorporated by reference herein, in their entirety.

15 Another microcrystalline cellulose product is a β - sheet form microcrystalline cellulose having a flat needle shape, marketed under the trademark CEOLUS KG801[®] by FMC Company of Newark, Delaware. The Ceolus[®] product has different morphology, and different performance characteristics, than those of the Avicel product. The β - sheet microcrystalline
20 cellulose of the present invention is disclosed in U.S. Patent 5,574,150, which is hereby incorporated by reference. Further disclosure relating to β - sheet microcrystalline cellulose is found in *International Journal of Pharmaceutics* 182 (199) 155 which is hereby incorporated by reference.

The Ceolus[®] product (β - sheet microcrystalline cellulose) is disclosed by
25 FMC, in its product bulletin dated October 1997, as being suitable for "smaller size tablets" and "exceptional drug carrying capacity." The Ceolus[®] product was said to provide superior compressibility and drug loading capacity, that still exhibited effective flowability. The examples given in the Ceolus[®] bulletin were

of vitamin C combined with Ceolus[®] microcrystalline cellulose at levels of from 30 to 45 weight % Ceolus[®] product in the form of a tablet.

However, there have been problems using the Ceolus[®] product. For example, at higher levels of Ceolus[®] product concentration, flow problems were encountered in the process of compressing tablets, and the Ceolus[®] product was considered unsuitable for compression at higher concentrations than about 45 weight %.

There is a definite need for solid levothyroxine (T4) and/or liothyronine (T3) (thyroid hormone drugs) pharmaceutical compositions, preferably in an immediate release solid dosage form, with the T4 and T3 in the form of their sodium salts that are relatively stable. There is also a need for methods for making such immediate release and stabilized solid levothyroxine (T4) and/or liothyronine (T3) (thyroid hormone drugs) pharmaceutical compositions.

Summary of the Invention

The present invention overcomes and alleviates the above-mentioned drawbacks and disadvantages in the thyroid drug art through the discovery of novel oral levothyroxine (T4) and/or liothyronine (T3) (thyroid hormone drugs) pharmaceutical compositions and methods.

Generally speaking, the present invention relates to stabilized solid as levothyroxine (T4) sodium and/or liothyronine (T3) sodium (thyroid hormone drugs) pharmaceutical compositions and in particular, immediate release, stabilized pharmaceutical compositions that include pharmaceutically active ingredients such as levothyroxine (T4) sodium and/or liothyronine (T3) sodium (thyroid hormone drugs). Preferably but not necessarily, the novel pharmaceutical compositions are provided in a solid dosage form, such as a tablet.

The pharmaceutical compositions of the present invention are useful for, among other things, as replacement or supplemental therapy in hypothyroidism of

any etiology, except transient hypothyroidism during the recovery phase of subacute thyroiditis, suppression of pituitary TSH secretion in the treatment or prevention of various types of euthyroid goiters, including thyroid nodules, Hashimoto's thyroiditis, multinodular goiter and, as adjunctive therapy in the management of thyrotropin-dependent well-differentiated thyroid cancer in warm-blooded animals, especially humans including pediatrics.

The present invention also provides methods for making such immediate release and stabilized levothyroxine (T4) sodium and/or liothyronine (T3) sodium (thyroid hormone drugs) pharmaceutical compositions.

Also in accordance with the present invention, because of the extraordinary release characteristics of the preferred compositions, a method of administration to children and patients who have difficulty taking pills, wherein the solid composition having the appropriate dosage is simply put in an aqueous fluid, e.g., juice, where it dissolves in a matter of 1-3 minutes, and the patient can then ingest the fluid, and receive the appropriate dosage, is now made practical.

The present invention has a wide range of important uses including providing pharmaceutically active levothyroxine compositions with enhanced bioavailability, improved shelf life, and more reliable potency.

We have discovered immediate release pharmaceutical compositions that include as pharmaceutically active ingredients at least one of levothyroxine and liothyronine, preferably at least one levothyroxine salt, as the major active ingredient. Such preferred immediate release compositions desirably provide at least about 85% (w/v) dissolution of the levothyroxine salt in less than about 20 minutes as determined by standard assays disclosed herein. Surprisingly, it has been found that by combining the pharmaceutically active ingredients with specific additives in accordance with the invention, it is possible to formulate the compositions so that the ingredients are released almost immediately after ingestion or contact with an aqueous solution, e.g., in a matter of minutes.

Preferred invention compositions are stable and provide better shelf life and potency characteristics than prior pharmaceutical compositions.

The immediate release pharmaceutical compositions of the invention provide important uses and advantages. A major advantage is the stability of the active ingredients in the composition. For example, while, as indicated above, prior formulations with sugars, starches, and various types of celluloses, including micro-cellular celluloses such as the Avicel products, have experienced substantial degradation of the active ingredients, e.g. T4 sodium. To deal with this problem, pharmaceutical manufacturers have over-formulated the T4-containing pharmaceutical compositions containing such active ingredients, so that the patient can obtain at least the prescribed dosage despite the carbohydrate-induced instability of the active ingredient. However, the patient who obtains the pharmaceutical immediately after it is made, receives an over-dosage of the active compound; whereas, the patient who has received the pharmaceutical after it has sat on the pharmacy shelf for an extended period, will receive an under-dosage of the active ingredient. In either case, the patient receives the wrong dosage, with possible serious consequences.

In sharp contrast, it has been surprisingly found that the use of the β -sheet microcrystalline cellulose in the compositions of the present invention substantially increase the stability of the thyroid hormone drugs, so that the patient obtains consistent potency over an extended shelf life, compared to prior thyroid hormone drug products. In this application, the term "stabilized", as applied to levothyroxine and/or liothyronine means that the loss of potency over the shelf life of the product is less than about 0.7 % potency per month, for at least about 18 months. Preferred compositions have a loss of potency of less than about 0.5% per month for such a period, and more preferred compositions have a loss of potency of less than about 0.3% per month for such a period.

Further, the compositions of the invention provide favorable pharmacokinetic characteristics when compared to prior formulations. In particular, the immediate release pharmaceutical compositions that include levothyroxine salt have are more quickly available for absorption by the gastrointestinal (GI) tract faster and are absorbed more completely than has
5 heretofore been possible. This invention feature substantially enhances levothyroxine bioavailability, thereby improving efficacy and reliability of many standard thyroid hormone replacement strategies.

Additionally, the desirable immediate release characteristics of the present
10 invention facilitate dosing of patients who may be generally adverse to thyroid hormone replacement strategies involving solid dosing. More specifically, immediate release pharmaceutical compositions disclosed herein can be rapidly dissolved in an appropriate aqueous solution (e.g., water, saline, juice) or colloidal suspension (e.g., baby formula or milk) for convenient administration to such
15 patients. Illustrative of such patients include infants, children, and adults who may experience swallowing difficulties. The invention thus makes standard thyroid hormone replacement strategies more flexible and reliable for such patients.

Accordingly, and in one embodiment, the invention features an immediate
20 release pharmaceutical composition comprising at least one levothyroxine salt, preferably one of such a salt. At least about 80% of the levothyroxine dissolves in aqueous solution in less than about 20 minutes as determined by a standard assay, disclosed herein. Preferably, at least about 80% of the levothyroxine is dissolved in the aqueous solution by about 15 minutes from the time that the
25 composition, in pill form, is placed in the aqueous solution. More preferably, at least about 85% of the levothyroxine is released to the aqueous solution by about 10 minutes, most preferably by about 5 minutes after exposure of the composition to the aqueous solution. As shown below, compositions in accordance with the

present invention can be formulated to release 85% of the levothyroxine within 2-3 minutes after exposure to the aqueous solution.

It has been found that by combining one or more of the pharmaceutically active agents with β -form microcrystalline cellulose, it is possible to produce compositions with favorable immediate release characteristics. Without wishing to be bound to theory, it is believed that the agents do not bind well to certain grades of the β - sheet form microcrystalline cellulose. More of the agent is thus available for immediate release. In contrast, it is believed that many prior formulations have active agents that bind cellulose additives, making less available. The release characteristics of the compositions of the invention are also improved by the use of other agents, as discussed further below.

Thus in one embodiment, the present invention relates to a stabilized pharmaceutical composition comprising a pharmaceutically active ingredient, such as levothyroxine, and the β - sheet form of microcrystalline cellulose, in the form of a solid dosage. More specifically, the present invention relates to a stabilized pharmaceutical composition comprising a pharmaceutically active ingredient, such as levothyroxine sodium and/or liothyronine sodium, at least about 50 weight % of the dosage weight composed of the β - sheet form of microcrystalline cellulose, and, optionally, additional excipients, in a solid dosage form.

In another aspect, the invention provides an aqueous solution or colloidal suspension that includes at least one of the compositions of this invention, preferably between from about one to about five of same, more preferably about one of such compositions.

It has also been found that β - sheet microcrystalline cellulose grades having preferred bulk densities provide for more compact processing than use of other celluloses. That is, use of the β - sheet microcrystalline cellulose having bulk densities in accord with this invention helps to provide for higher

compression ratios (initial volume/final volume). As discussed below, other invention aspects help reduce or avoid production of damaging compression heat that has damaged prior formulations made from high compression ratios. The compositions of the present invention generally also require less compressional
5 force to form the tablets.

Accordingly, the invention also provides methods for making an immediate release pharmaceutical composition comprising at least one levothyroxine salt, preferably one of such a salt. In one embodiment, the method includes at least one and preferably all of the following steps:

10

a) mixing a levothyroxine salt with microcrystalline β -cellulose and preferably a crosscarmellose salt to make a blend; and

15

b) compressing the blend in a ratio of initial volume to final volume of between from about 2:1 to about 5:1 to make the composition, preferably about 4:1.

20

In one embodiment, the method involves preparing an oral dosage form of a pharmaceutically active ingredient comprising dry blending the pharmaceutically active ingredient and at least about 50 weight % of the β - sheet form of microcrystalline cellulose, and compressing the blend to form a solid dosage.

25

These and other objects, features, and advantages of the present invention may be better understood and appreciated from the following detailed description of the embodiments thereof, selected for purposes of illustration and shown in the accompanying figures and examples. It should therefore be understood that the

particular embodiments illustrating the present invention are exemplary only and not to be regarded as limitations of the present invention.

Brief Description of the Fig.

5 The foregoing and other objects, advantages and features of the invention, and the manner in which the same are accomplished, will become more readily apparent upon consideration of the following detailed description of the invention taken in conjunction with the accompanying Figs., which illustrate a preferred and exemplary embodiment, wherein:

10 Figures 1A – 1C illustrate various solid dosage forms such as cylindrical tablets and raised violin shaped tablets;

 Figure 2 illustrates a tableting die pair;

 Figure 3 pair; is graphical depiction of comparative dissolution data of various strengths of Levoxyl[®] tablets made in accordance with the invention.

15 Figure 4A is an HPLC chromatogram showing a levothyroxine and liothyronine standards.

 Figure 4B is an HPLC chromatograph showing results of levothyroxine sodium sample made in accordance with the present invention.

20 Figure 5A is a chromatogram showing various levothyroxine impurity standards.

 Figure 5B is a chromatograph showing results of levothyroxine sodium sample made in accordance with the present invention.

Detailed Description

25 By way of illustrating and providing a more complete appreciation of the present invention and many of the attendant advantages thereof, the following detailed description is given concerning the novel oral levothyroxine (T4) and/or

liothyronine (T3) (thyroid hormone drugs) pharmaceutical compositions and methods for use in warm-blooded animals, especially humans and children.

As discussed, the invention relates to immediate release solid pharmaceutical compositions such as stabilized pharmaceutical compositions that include pharmaceutically active ingredients such as levothyroxine (T4) sodium and liothyronine (T3) sodium (thyroid hormone drugs), preferably in a solid dosage form. Also provided are methods for making such immediate release and stabilized compositions.

Aspects of the present invention have been disclosed in U.S. Provisional Application No. 60/269,089, entitled *Stabilized Pharmaceutical and Thyroid Hormone Compositions and Method of Preparation* and filed on February 15, 2001 by Franz, G.A. et al. The disclosure of said provisional application is incorporated herein by reference.

By the phrase "immediate release" is meant a pharmaceutical composition in which one or more active agents therein demonstrates at least about 80% (w/v) dissolution, preferably between from about 90% (w/v) to about 95% (w/v), more preferably about 95% (w/v) to about 99% (w/v) or more within 15 to 20 minutes as determined by a standard dissolution test. Suitable standard dissolution tests are known in the field. See FDA, Center for Drug Research, **Guidance for Industry, In Vivo Pharmacokinetics and Bioavailability Studies and In Vitro Dissolution Testing for Levothyroxine Sodium Tablets**, available at www.fda.gov/cder/guidance/index.htm. A specifically preferred dissolution test is provided in Example 2, below.

A pharmaceutical composition of the invention is "stable" or "stabilized" if one or more of the active agents therein exhibit good stability as determined by a standard potency test. More specifically, such compositions exhibit a potency loss of less than about 15%, preferably less than about 10%, more preferably less than about 1% to about 5% as determined by the test. Potency can be evaluated

by one or a combination of strategies known in the field. See the USP. A preferred potency test compares loss or conversion of the active agent in the presence (experimental) or absence (control) of a carrier or excipient. A specifically preferred potency test is provided in Examples 1 and 3, below.

5 In preferred embodiments, the pharmaceutical compositions of the invention include, as active agent, levothyroxine (T4), preferably a salt thereof such as levothyroxine sodium USP. Such compositions typically exhibit a levothyroxine (T4) plasma Cmax of between from about 12 µg/dl to about 16 µg/dl, preferably as determined by the standard Cmax test. Preferably, the
10 In(Cmax) of the levothyroxine (T4) plasma level is between from about 1 to about 3.

The standard Cmax test can be performed by one or a combination of strategies known in the field. See e.g., the USP. A preferred Cmax test is disclosed below in Examples 8 and 9.

15 Additionally preferred compositions in accord with the invention provide a triiodothyronine (T3) plasma Cmax of between from about 0.1 ng/ml to about 10ng/ml, preferably 0.5 ng/ml to about 2ng/ml, as determined by the standard Cmax test. Typically, the In(Cmax) is between from about 0.01 to about 5. See Examples 8 and 9 for more information.

20 Further preferred compositions exhibit a levothyroxine (T4) plasma Tmax of between from about 0.5 hours to about 5 hours, preferably as determined by a standard Tmax test. The standard Tmax test can be performed by procedures generally known in the field. See e.g., the USP. A preferred Tmax test is disclosed below in Examples 8 and 9.

25 Still further preferred compositions of the invention exhibit a triiodothyronine (T3) plasma Tmax of between from about 10 hours to about 20 hours, preferably about 12 to about 16 hours as determined by the standard Tmax test.

Additionally preferred invention compositions feature a levothyroxine (T4) plasma AUC (0-t) of between from about 450 $\mu\text{g-hour/dl}$ to about 600 $\mu\text{g-hour/dl}$, preferably 500 $\mu\text{g-hour/dl}$ to about 550 $\mu\text{g-hour/dl}$ as determined by a standard AUC (0-t) test. Preferably, the $\ln[\text{AUC}(0-t)]$ is between from about 1 to about 10.

Standard methods for performing AUC (0-t) test determinations are generally known in the field. See e.g., the USP. Examples 8 and 9 below provide a specifically preferred method of determining the AUC (0-t).

Further preferred invention compositions feature a triiodothyronine (T3) AUC (0-t) of between from about 10 ng-hour/ml to about 100 ng-hour/ml , preferably 20 ng-hour/ml to about 60 ng-hour/ml , as determined by the standard AUC (0-t) test. Preferably, the $\ln[\text{AUC}(0-t)]$ is between from about 1 to about 5.

As will be appreciated, many prior pharmaceutical formulations include lactose or other sugars as a pharmaceutically acceptable carrier. It has been found however, that sugars such as lactose can react with active agents including the levothyroxine (T4) compositions of the present invention. For example, and without wishing to be bound to theory, it is believed that lactose is particularly damaging to T4 and T3 molecules via Schiff reactions. The invention address this problem by providing compositions that are essentially sugar-free. Particular invention compositions are essentially free of lactose.

Additionally, preferred pharmaceutical compositions of the invention are provided in which the active material is a non-granulated material. Prior levothyroxine compositions have been granulated in various size reduction machines to grains of less than, e.g., 5 – 20 microns average particle size in order to be effectively incorporated into the administrable pharmaceutical composition. The granulation process subjects the active material to degrading heat, which can have adverse effects on the active material, as well as reducing the activity level. Prior manufacturers purchase micronized levothyroxine manufactured under

DMF No. 4789, and then granulate it before incorporating it into the levothyroxine pharmaceutical product.

In the preferred method of the present invention, the raw material is not granulated before incorporation into the pharmaceutical composition. Rather, the
5 ingredients of the preferred pharmaceutical are mixed and the mixture is subjected to direct compression to form the pharmaceutical tablets of appropriate dosage. As a result, the activity of the active ingredient is not degraded prior to the direct compression step. Bulk levothyroxine is obtained in a fine powdered form, preferably from Biochemie GmbH, A-6250 Kundl, Austria. More importantly,
10 the use of the preferred process results in a product which is immediately dispersible in aqueous solution, to make the active ingredient available for absorption in the body. As used in this application, "non-granulated" means that the bulk USP compound is used without subjecting it to granulators or similar high energy size reduction equipment before being mixed with the other
15 pharmaceutical components and formed into the appropriate pill. Preferably, the bulk active ingredient is mixed with the appropriate amounts of other ingredients and directly compressed into pill form. Since it is not necessary to granulate the material, it is not necessary to subject it to degrading temperatures in the process of forming the pharmaceutical compositions containing the active materials. In
20 the present process we start with micronized active material, which merely needs to be blended with the B and other materials and then compressed. Others have to be granulated, and then dried, which steps interfere with the dissolution of the active material. The drying temperatures employed in manufacturing other active ingredients can cause degradation of the levothyroxine, as experienced in other
25 available thyroxine. It has been found that providing the invention compositions in a non-granulated format helps to reduce or eliminate active agent degradation, presumably by facilitating a reduction in friction, and thus degrading heat, during compression of the compositions into pills.

Practice of the invention is compatible with several β -form microcrystalline cellulose grades. Preferably, the β -form microcrystalline cellulose has a bulk density of between from about 0.10 g/cm^3 to about 0.35 g/cm^3 , more preferably between from about 0.15 g/cm^3 to about 0.25 g/cm^3 , still
5 more preferably between from about 0.17 g/cm^3 to about 0.23 g/cm^3 , most preferably between from about 0.19 g/cm^3 to about 0.21 g/cm^3 .

Further preferred grades of the β -form microcrystalline cellulose are substantially non-conductive. Preferably, the β -form microcrystalline cellulose has a conductivity of less than about $200 \text{ }\mu\text{S/cm}$, more preferably, less than about
10 $75 \text{ }\mu\text{S/cm}$, still more preferably between from about $0.5 \text{ }\mu\text{S/cm}$ to $50 \text{ }\mu\text{S/cm}$, most preferably between from about $15 \text{ }\mu\text{S/cm}$ to $30 \text{ }\mu\text{S/cm}$.

A specifically preferred β -form microcrystalline cellulose is sold by Asahi Chemical Industry Co., Ltd (Tokyo, Japan) as Ceolus (Type KG-801 and/or KG-802).

15 Additionally preferred compositions of the invention have a post-packaging potency of between from about 95% to about 120%, preferably 98% to about 110% as determined by the standard potency test.

The present invention is a pharmaceutical product that is in the form of a solid dosage, such as a sublingual lozenge, buccal tablet, oral lozenge,
20 suppository or a compressed tablet. The pharmaceutically active ingredient is dry mixed with the β -form of the microcrystalline cellulose, optionally with additional excipients, and formed into a suitable solid dosage.

Preferred tablets according to the invention have a total hardness of between from about 1 to about 30 KP, preferably about 6 to about 14 KP as
25 determined by a standard hardness test. Methods for determining tablet hardness are generally known in the field. See e.g., the USP. A preferred standard hardness test is disclosed below in Example 4.

Additionally preferred pharmaceutical compositions including those in tablet format preferably include less than about 10% total impurities, more preferably less than about 5% of same as determined by a standard impurity test.

Reference herein to the "standard impurity test" means a USP recognized
5 assay for detecting and preferably quantitating active drug degradation products. In embodiments in which levothyroxine or liothyronine break-downs are to be monitored, such products include, but are not limited to, at least one of diiodothyronine (T2), triiodothyronine (T3), levothyroxine, triiodothyroacetic
10 acid amide, triiodothyroethylamine, triiodothyroacetic acid, triiodothyroethyl alcohol, tetraiodothyroacetic acid amide, tetraiodothyroacetic acid, triiodothyroethane, and tetraiodothyroethane. Of particular interest are diiodothyronine (T2), triiodothyronine (T3), triiodothyroacetic acid, and tetraiodothyroacetic acid impurities.

A preferred impurity test for monitoring levothyroxine and liothyronine
15 breakdown products involves liquid chromatography (LC) separation and detection, more preferably HPLC. Specifically preferred impurity tests are provided below in Examples 5 and 6

Further preferred compositions in accord with the invention include one or more standard disintegrating agents, preferably crosscarmellose, more preferably
20 a salt of same. Still further preferred compositions include a pharmaceutically acceptable additive or excipient such as a magnesium salt.

The present invention can be prepared as a direct compression formula, dry granulation formula, or as a wet granulation formula, with or without preblending of the drug, although preferably with preblending.

25 The pharmaceutically active ingredient can be any type of medication which acts locally in the mouth or systemically, which is the case of the latter, can be administered orally to transmit the active medicament into the gastrointestinal tract and into the blood, fluids and tissues of the body. Alternatively, the

medicament can be of any type of medication which acts through the buccal tissues of the mouth to transmit the active ingredient directly into the blood stream thus avoiding first liver metabolism and by the gastric and intestinal fluids which often have an adverse inactivating or destructive action on many active ingredients unless they are specially protected against such fluids as by means of an enteric coating or the like. The active ingredient can also be of a type of medication which can be transmitted into the blood circulation through the rectal tissues.

Representative active medicaments include antacids, antimicrobials, coronary dilators, peripheral vasodilators, anti psychotropics, antimanics, stimulants, antihistamines, laxatives, decongestants, vitamins, gastrosedatives, antidiarrheal preparations, vasodilators, antiarrhythmics, vasoconstrictors and migraine treatments, anticoagulants and antithrombotic drugs, analgesics, antihypnotics, sedatives, anticonvulsants, neuromuscular drugs, hyper and hypoglycemic agents, thyroid and antithyroid preparations, diuretics, antispasmodics, uterine relaxants, mineral and nutritional additives, antiobesity drugs, anabolic drugs, erythropoietic drugs, antiasthmatics, expectorants, cough suppressants, mucolytics, antiuricemic drugs, and drugs or substances acting locally in the mouth.

Typical active medicaments include gastrointestinal sedatives such as metoclopramide and propantheline bromide, antacids such as aluminum trisilicate, aluminum hydroxide and cimetidine, aspirin-like drugs such as phenylbutazone, indomethacin, and naproxen, ibuprofen, flurbiprofen, diclofenac, dexamethasone, prednisone and prednisolone, coronary vasodilator drugs such as glyceryl trinitrate, isosorbide dinitrate and pentaerythritol tetranitrate, peripheral and cerebral vasodilators such as soloctidilum, vincamine, naftidrofuryl oxalate, comesylate, cyclandelate, papaverine and nicotinic acid, antimicrobials, such as erythromycin stearate, cephalixin, nalidixic acid, tetracycline hydrochloride,

ampicillin, flucloxacillin sodium, hexamine mandelate and hexamine hippurate, neuroleptic drugs such as fluazepam, diazepam, temazepam, amitriptyline, doxepin, lithium carbonate, lithium sulfate, chlorpromazine, thioridazine, trifluoperazine, fluphenazine, piperothiazine, haloperidol, maprotiline

5 hydrochloride, imipramine and desmethylinipramine, central nervous stimulants such as methylphenidate, ephedrine, epinephrine, isoproterenol, amphetamine sulfate and amphetamine hydrochloride, anitid^rugs such as diphenylhydramine, diphenylpyramine, chlorpheniramine and brompheniramine, antidiarrheal drugs such as bisacodyl and magnesium hydroxide, the laxative drug, dioctyl sodium

10 sulfosuccinate, nutritional supplements such as ascorbic acid, alpha tocopherol, thiamine and pyridoxine, antispasmodics such as dicyclomine and diphenoxylate, drugs effecting the rhythm of the heart such as verapamil, nifedepine, diltiazem, procainamide, disopyramide, bretylium tosylate, quinidine sulfate and quinidine gluconate, drugs used in the treatment of hypertension such as propranolol

15 hydrochloride, guanethidine monosulphate, methyldopa, oxprenolol hydrochloride, captopril, Actace and hydralazine, drugs used in the treatment of migraine such as ergotamine, drugs effecting coagulability of blood such as epsilon aminocaproic acid and protamine sulfate, analgesic drugs such as acetylsalicyclic acid, acetaminophen, codeine phosphate, codeine sulfate,

20 oxycodone, dihydrocodeine tartrate, oxydodeinone, morphine, heroin, nalbuphine, butorphanol tartrate, pentazocine hydrochloride, cyclazacine, pethidine, buprenorphine, scopolamine and mefenamic acid, antl^rdrugs such as phenytoin sodium and sodium valproate, neuromuscular drugs such as dantrolene sodium, substances used in the treatment of diabetes, such as tolbutamide, diabenase

25 glucagon and insulin, drugs used in the treatment of thyroid gland dysfunction such as triiodothyronine, liothyronine sodium, levothyroxilne sodium and related compounds, and propylthiouracil, diuretic drugs, such as furosemide, chlorthalidone, hydrochlorthiazide, spironolactone and triamptereⁿe, the uterine

relaxant drugritodrine, appetite suppressants such as fenfluramine hydrochloride, phentermine and diethylpropion hydrochloride, antidrugs stimulants such as aminophylline, theophylline, salbutamol, orciprenaline sulphate and terbutaline sulphate, expectorant drug such as guaiphenesin, cough suppressants such as
5 dextromethorphan and mescaline, mucolytic drugs such as carbocisteine, antiseptics such as cetylpyridinium chloride, tyrothricin and chlorhexidine, decongestant drugs such as phenylpropanolamine and pseudoephedrine, hypnotic drugs such as dichloralphenazone and nitrazepam, antidrugs H₁ blockers such as promethazine theoclate, haemopoetic drugs such as ferrous sulphate, folic acid
10 and calcium gluconate, uricosuric drugs such as sulphinpyrazine, allopurinol and probenecid and the like. It is understood that the invention is not restricted to the above medications.

The amount of pharmaceutically active ingredient in the present composition can vary widely, as desired. Preferably, the active ingredient is
15 present in a composition of the present invention in an effective dosage amount. Exemplary of a range that the active ingredient may be present in a composition in accordance with the present invention is from about 0.000001 to about 10 weight %. More preferably, the amount of active ingredient is present in the range of about 0.001 to 5 weight %.

20 Of course, it should be understood that any suitable pharmaceutically acceptable form of the active ingredient can be employed in the compositions of the present invention, i.e., the free base or a pharmaceutically acceptable salt thereof, e.g., levothyroxine sodium salt, etc.

When the pharmaceutically active moiety is levothyroxine sodium, the
25 preferred amount of the active moiety in the composition is present in the range of about 0.00005 to about 5 weight %. The more preferred range is from about 0.001 to about 1.0 weight %, and the most preferred range is from about 0.002 to about 0.6 weight % levothyroxine. The minimum amount of levothyroxine can vary, so

long as an effective amount is utilized to cause the desired pharmacological effect. Typically, the dosage forms have a content of levothyroxine in the range of about 25 to 300 micrograms per 145 milligram pill for human applications, and about 100 to 800 micrograms per 145 mg pill for veterinary applications.

5 In accordance with the present invention, a goal of levothyroxine replacement therapy is to achieve and maintain a clinical and biochemical euthyroid state, whereas a goal of suppressive therapy is to inhibit growth and/or function of abnormal thyroid tissue. A dose of levothyroxine that is adequate to achieve these goals depends of course on a variety of factors including the
10 patient's age, body weight, cardiovascular status, concomitant medical conditions, including pregnancy, concomitant medications, and the specific nature of the condition being treated. Hence, the following recommendations serve only as dosing guidelines. It should be understood by those versed in this art that dosing should be individualized and adjustments made based on periodic assessment of a
15 patient's clinical response and laboratory parameters.

 Preferably, but not necessarily, when using levothyroxine to treat, it should be taken in the morning on an empty stomach, at least one-half hour before any food is eaten. In addition, levothyroxine is preferably taken at least about 4 hours apart from drugs that are known to interfere with its absorption.

20 Due to the long half-life of levothyroxine, the peak therapeutic effect at a given dose of levothyroxine sodium may not be attained for about 4 to about 6 weeks.

 In people older than 50 years, who have been recently treated for hyperthyroidism or who have been hypothyroid for only a short time (such as a
25 few months), the average full replacement dose of levothyroxine sodium is approximately 1.7 mcg/kg/day (e.g., about 100 to about 125 mcg/day for a 70 kg adult). Older patients may require less than 1 mcg/kg/day. Levothyroxine sodium doses greater than about 200 mcg/day may or may not be required.

For most patients older than 50 years or for patients under 50 years of age with underlying cardiac disease, an initial starting dose of about 25 to about 50 mcg/day of levothyroxine sodium is recommended, with gradual increments in dose at about 6 to about 8 week intervals, as needed. The recommended starting
5 dose of levothyroxine sodium in elderly patients with cardiac disease is about 12.5 to about 25 mcg/day, with gradual dose increments at about 4 to about 6 week intervals. The levothyroxine sodium dose is generally adjusted in about 12.5 to about 25 mcg increments until the patient with primary hypothyroidism is clinically euthyroid and the serum TSH has normalized.

10 In patients with severe hypothyroidism, the recommended initial levothyroxine sodium dose is about 12.5 to about 25 mcg/day with increases of about 25 mcg/day about every 2 to about 4 weeks, accompanied by clinical and laboratory assessment, until the TSH level is normalized.

In patients with secondary (pituitary) or tertiary (hypothalamic)
15 hypothyroidism, the levothyroxine sodium dose should be titrated until the patient is clinically euthyroid and the serum free-T₄ level is restored to the upper half of the normal range.

In children, levothyroxine therapy may be instituted at full replacement doses as soon as possible. Levothyroxine compositions of the present invention
20 may be administered to infants and children who cannot swallow intact tablets by crushing the tablet and suspending the freshly crushed tablet in a small amount (5–10 mL or 1–2 teaspoons) of water. This suspension can be administered by spoon or dropper. Foods that decrease absorption of levothyroxine, such as soybean infant formula, should not be used for administering levothyroxine
25 sodium tablets.

A recommended starting dose of levothyroxine sodium in newborn infants is about 10 to about 15 mcg/kg/day. A lower starting dose (e.g., about 25 mcg/day) may be considered in infants at risk for cardiac failure, and the dose

should be increased in 4–6 weeks as needed based on clinical and laboratory response to treatment. In infants with very low (< about 5 mcg/dL) or undetectable serum T₄ concentrations, a recommended initial starting dose is about 50 mcg/day of levothyroxine sodium.

- 5 As indicated above, levothyroxine therapy is usually initiated at full replacement doses, with the recommended dose per body weight decreasing with age (see Dose Table below). However, in children with chronic or severe hypothyroidism, an initial dose of about 25 mcg/day of levothyroxine sodium is recommended with increments of 25 mcg every 2–4 weeks until the desired effect
- 10 is achieved. Hyperactivity in an older child may be minimized if the starting dose is one-fourth of the recommended full replacement dose, and the dose is then increased on a weekly basis by an amount equal to one-fourth the full-recommended replacement dose until the full recommended replacement dose is reached.

15

Dose Table: Levothyroxine Sodium Dosing Guidelines for Pediatric Hypothyroidism	
AGE	Daily Dose per Kg Body Weight^a
0–3 months	10–15 mcg/kg/day
3–6 months	8–10 mcg/kg/day
6–12 months	6–8 mcg/kg/day
1–5 years	5–6 mcg/kg/day
6–12 years	4–5 mcg/kg/day
>12 years	2–3 mcg/kg/day
Growth and puberty complete	1.7 mcg/kg/day
^a The dose should be adjusted based on clinical response and laboratory parameters.	

Levothyroxine sodium tablets, USP, in accordance with the present invention may be supplied as oval or violin-shaped, color-coded, potency marked tablets in, for example, 12 strengths as indicated in the Strength Table

5

Strength Table	Levothyroxine Tablets
Strength (mcg)	Color
25	Orange
50	White
75	Purple
88	Olive
100	Yellow
112	Rose
125	Brown
137	Dark Blue
150	Blue
175	Turquoise
200	Pink
300	Green

When the pharmaceutically active moiety is liothyronine sodium, the preferred amount of the active moiety in the composition is present in the range of about 0.000005 to 0.5 weight %. The more preferred range is from about 0.00001 to 0.1 weight %, and the most preferred range is from about 0.00004 to about 0.002 weight % liothyronine. The minimum amount of liothyronine can vary, so long as an effective amount is utilized to cause the desired pharmacological effect. Typically, the dosage forms have a content of levothyroxine in the range of about 5 to 50 micrograms per 145 milligram pill for human applications.

The β -form microcrystalline cellulose product of the present invention is prepared by forming a wet cake, drying the cake with a drum dryer, then passing

the dried product through a screen or mill for sizing which produces a β -sheet microcrystalline cellulose which has a flat needle shape, as disclosed in U.S. Patent 5,574,150. Such β -sheet microcrystalline product is available from Asahi Chemical of Japan and/or marketed by FMC Company of Newark, Del, under the trademark Ceolus®. The morphology and performance characteristics of the
5 Ceolus® product are different from those of α - form microcellulose products (for example, Avicel® and Emcocel®), and are suitable for preparing the present stabilized pharmaceutical composition.

The amount of β -form microcrystalline product used in the present
10 composition is at least 50 weight % of the final composition. Preferably, the amount of β -form microcrystalline product is in the range of about 50 to 99 weight %. Most preferably, the amount of β -form microcrystalline product is in the range of about 60 to 90 weight % of the final composition.

Other suitable excipients for the present invention include fillers such as
15 starch, alkaline inorganic salts such as trisodium phosphate, tricalcium phosphate, calcium sulfate and sodium or magnesium carbonate. The fillers can be present in the present composition in the range of about 0 to 50 weight %.

Suitable disintegrating agents include corn starch, cross-linked sodium carboxymethylcellulose (crosscarmellose) and cross-linked polyvinylpyrrolidone
20 (crospovidone). A preferred disintegrating agent is crosscarmellose. The amount of disintegrating agent used is in the range of about 0 to 50 weight %. Preferably, the disintegrating agent is in the range of about 5 to 40 weight %, more preferably about 10 to about 30 weight %. This is in substantial excess of the recommended levels of such materials. For example, the recommended loading of
25 crosscarmellose is 0.5 to about 2% by weight. However, it has been found that the higher loadings of the disintegrating agents substantially improves the ability of the product to disperse in aqueous media.

Suitable gildents for use in the present invention include colloidal silicon dioxide and talc. The amount of gildent in the present composition is from about 0 to 5 weight %, and the preferred amount is about 0 to 2 weight %.

5 Suitable lubricants include magnesium and zinc stearate, sodium stearate fumarate and sodium and magnesium lauryl sulfate. A preferred lubricant is magnesium stearate. The amount of lubricant is typically in the range of about 0 to 5 weight %, preferably in the range of about 0.1 to 3 weight %.

10 The oral pharmaceutical product is prepared by thoroughly intermixing the active moiety and the β -form of microcrystalline cellulose, along with other excipients to form the oral dosage. Food grade dyes can also be added. For example, it is common to distinguish dosages of various potency by the color characteristics of such dyes.

15 As discussed, a preferred immediate release pharmaceutical composition in tablet form includes levothyroxine sodium. In a preferred embodiment, the composition includes at least one of, preferably all of the following:

- a) between from about 0.01 mg/tablet to about 500 mg/tablet levothyroxine sodium (USP),
- 20 b) between from about 100 mg/tablet to about 110 mg/tablet of microcrystalline β -cellulose, NF (Ceolus) having a bulk density of between from about 0.10 g/cm³ to about 0.35 g/cm³,
- c) between from about 25mg/tablet to about 50mg/tablet of crosscarmellose sodium, NF
25 (Ac-di-sol); and
- d) between from about 0.5 mg/tablet to about 5mg/tablet of magnesium stearate, NF.

Preferably, the composition further comprises at least one pharmaceutically acceptable coloring agent.

More particular methods according to the invention provide compositions having less than about 5% total impurities as determined by the standard impurity
5 test. Preferably, the method further comprises forming a tablet, particularly those tablets having a raised violin configuration.

The stabilized oral dosages of thyroid hormone are prepared by forming a trituration of the active moiety (i.e. levothyroxine sodium and/or liothyronine sodium) and β -form microcrystalline cellulose. The trituration is blended with β -
10 form microcrystalline cellulose and additional excipients and compressed into oral dosages.

Design of the tableting apparatus is important, in order to maintain consistency from one oral dosage to the next. The formulation batches are a blend of solid compositions of various shapes and sizes. Blending is used to achieve a
15 measure of homogeneity. In particular the active thyroid moiety is desired to be evenly distributed throughout the batch. In a typical 410 kg batch, the amount of active moiety represents less than 1 kg of the total weight. For example, when producing 145 mg tablets with a 300 mcg dosage, approximately 0.8 kg of a 410 kg batch is the active moiety. In addition each tablet is formulated to contain
20 100% label claim potency.

It is typical for compressible medicament tablets to be formed using a 2:1 fill to compression ratio. However, for medicament tablets formed using the present invention a fill to compression ratio from 3.3:1 to 4:1 is needed to obtain
25 desired tablet density. The β -form microcrystalline cellulose has a lower bulk density, as compared to other excipients.

Higher tablet density can be accomplished by adjusting a tableting machine to increase the compression ratio. Tableting machines are commonly known to practitioners in the art and include those available from Manesty and

Stokes. It has been found that making such adjustments to the compression ratio results in poor tablet surface finish as well as inconsistent tablet weights. Instead, the design of the tableting dies should be adjusted. It has been determined that during the filling of the tableting dies, a minimum of 5-6mm die overfill. In most cases this requires replacement of the usual tableting dies with dies which are an additional 2-3 mm deep.

When using the extra-deep dies and a compression ratio of from 3.3:1 to 4.0:1, consistent weight tablets with good surface finish were produced.

Preferably, the shape of the tablet is configured to increase heat transfer away from the tablet. More preferred tablets have a surface area per tablet of between from about 0.9 in.² to about 0.15 in.², preferably about 0.115 in.², to assist such heat transfer. Additional tablet configurations are contemplated e.g., tablets that are beveled and/or include a notch. A preferred tablet shape is a raised violin configuration, as shown in Figure 1C.

The following examples are given by way of illustration only and are not to be considered limitations of this invention or many apparent variations of which are possible without departing from the spirit or scope thereof.

Example 1

20

Stability Tests

Stability testing was performed on samples of the thyroid hormone drug formulation used in manufacturing tablets with an active moiety of levothyroxine sodium. Tests were performed on direct compression formulations for dosage strength of 25 mcg. Example 1 tablets comprise the β -form microcrystalline cellulose while Control 1 tablets comprise the traditional α -form microcrystalline cellulose. The composition of Example 1 and Control 1 tablets are presented in Table 1 and stability test results in Table 2:

Table 1 - Tablet Formulation for 25mcg Dosages of Levothyroxine Sodium

Example 1 Tablet	Control 1 Tablet	Component
0.0297 mg	0.0297 mg	Levothyroxine Sodium, USP
108.55 mg		β - sheet microcrystalline cellulose
	108.55 mg	β - form microcrystalline cellulose
35.079 mg	35.079 mg	Crosscarmellose Sodium, NF
0.352 mg	0.352 mg	FD&C Yellow #6 16% (14-20%)
1.018 mg	1.018 mg	Magnesium Stearate, NF
145.0 mg	145.0 mg	Total

5

Table 2 - Stability Test - Potency at 25 °C -- % Label Claim

Elapsed Time	0	73 Days	13 months	15 months
Example 1 Tablet	106.4	105.5	104.4	102.9
Example 1 % Potency Loss	0.0	0.9 %	2.0%	3.5%
% Change per Month	0.0	0.37	0.15	0.23
Control 1 Tablet	99.2	89.5	85.0	83.2
Control 1 % Potency Loss	0.0	2.7%	14.2 %	16.0 %
% Change per Month	0.0	1.11	1.09	1.07

As seen in Table 2, the stability of pharmaceutical formulations of the present invention is improved significantly by the use of the β - sheet microcrystalline cellulose. Potency loss of the present invention after 15 months is 3.5 %, versus 16.0 % potency loss experienced in a similar formulation with the α -form microcrystalline cellulose. The average loss in potency per month in the

case of the compositions of the present invention was only about 0.2 % per month, as compared to over 1% per month for the T4 products which included β -form microcrystalline cellulose, thus demonstrating a stability which is about 3 to 4 times better than the T4 products which utilized α -form microcrystalline cellulose.

Tableting testing was performed on the formulation for Example 1 tablets. Initial results with standard die depths provided a relative standard deviation of 2.2 to 3.5% tablet weight. With the use of the herein described extra deep tablet dies, the relative standard deviation is 1.2%. Testing was performed on a Manesty tableting machine with compression ratios of from 3.3:1 to 4.0:1.

Tablet quality is also dependent upon the storage of the β -sheet microcrystalline cellulose. Best results are achieved when the cellulose is received in drums or portable containers instead of bags. The bag form suffers from compression during transportation from raw material suppliers. Test results for tableting are presented in attached Exhibit A.

Additional examples of solid dosage formulations are illustrated in Tables 3 and 4. Stability testing data of additional examples are illustrated in Table 5.

Table 3 - Tablet Formulation for Dosages of Levothyroxine Sodium (per tablet)

25 mcg Dosage	50 mcg Dosage	75 mcg Dosage	Component
0.025 mg	0.0500 mg	0.0750 mg	levothyroxine sodium
108.529 mg	108.856 mg	108.438 mg	β -form microcrystalline cellulose
35.079 mg	35.079 mg	35.079 mg	crosscarmellose sodium
0.352 mg		0.383 mg	food grade dye
1.018 mg	1.018 mg	1.018 mg	magnesium stearate
145 mg/tablet	145 mg/tablet	145 mg/tablet	Total

Table 4 - Tablet Formulation for Dosages of Levothyroxine Sodium (per tablet)

100 mcg Dosage	112 mcg Dosage	300 mcg Dosage	Component
0.100 mg	0.112 mg	0.300 mg	Levothyroxine sodium
108.406 mg	107.711 mg	108.451 mg	β -form microcrystalline cellulose
35.079 mg	35.079 mg	35.079 mg	crosscarmellose sodium
0.388 mg	1.080 mg	0.142 mg	food grade dye
1.018 mg	1.018 mg	1.1018 mg	magnesium stearate
145 mg/tablet	145 mg/tablet	145 mg/tablet	Total

5

Table 5 shows drug stability data for a number of the above formulations:

Table 5 - Stability Test - Potency at 25 ° C - % Label Claim

Levothyroxine Na	Test Interval (months)			
Test	Initi	6	12	18
25 µg Dose	26.2	25.6	25.5	25.3
% Label Claim	104.	102.	102.	101.
% of Initial Result	100.	97.5	97.3	96.6
% Change	0.0	2.6	2.8	3.6
% Change per month	0.0	0.43	0.23	0.2
50 µG Dose	51.0	49.9	48.9	48.4
% Label Claim	102.	99.7	97.7	96.7
% of Initial Result	100.	97.7	95.8	94.8
% Change	0.0	2.3	4.3	5.3
% Change per month	0.0	0.38	0.36	0.29
112 µg Dose	113.	113.	109.	105.
% Label Claim	101.	101.	97.8	94.5
% of Initial Result	100.	100.	96.6	93.4
% Change	0.0	0.3	3.4	6.7
% Change per month	0.0	0.05	0.28	0.37
200 µg Dose	202.	196.	198.	196.
% Label Claim	101.	98.4	99.3	98.3
% of Initial Result	100.	97.3	98.2	97.2
% Change	0.0	2.7	1.7	2.8
% Change per month	0.0	0.45	0.14	0.15

Thus the formulations of the present invention provide extreme stability
 5 for the levothyroxine activity over an extended shelf life for these pharmaceutical products.

Example 2

Dissolution Tests

10 The following preferred method for testing potency will sometimes be referred to herein as method number: **AM-004B**

Table 6 – Dissolution Test Procedure

Chromatographic Conditions	
Mobile Phase:	Degassed and filtered mixture of methanol and 0.1% phosphoric acid (60:40).
Column:	C ₁₈ 3.9 mm x 30 cm
Flow Rate:	2.0 ml/minute
Detector:	Deuterium set at 225 nm
Injection Volume:	800 µL
System Suitability:	Chromatograph 6 replicate injections of the standard preparation.
	1.0 RDS for the standard replicates must not be more than 4.0%.
	2.0 The tailing factor must not be more than 1.5.
Medium:	0.01 N hydrochloric acid containing 0.2% sodium lauryl sulfate; 500 ± 5 ml; 37 ± 0.5°C This solution is very foamy; excessive mixing, shaking, and pouring will make reading the meniscus on the graduated cylinder difficult.
Apparatus:	Apparatus 2 (Paddles)
Apparatus Cleaning:	The apparatus is to be cleaned immediately after use or if left idle for more than 12 hours. Clean paddles by rinsing with distilled water, methanol, and distilled water again. Blot to dry with Kimwipes. Clean vessels by rinsing with hot tap water, microdetergent, hot tap water, and distilled water. Dry using paper towels.
Paddle Speed:	50 rpm
Incubation Period:	Up to 45 minutes

Standard Solutions:	Transfer about 50 mg USP Levothyroxine RS, accurately weighed, into a 100 ml volumetric flask. Add approximately 30 ml of methanol, dissolve and dilute to volume with methanol, mix. Using this solution, standard solutions are prepared in a volumetric flask using Dissolution Media, diluting to a concentration that comes near to the theoretical concentration of the tablet in 500 ml of Dissolution Media. Use a pipette to gently add the Dissolution media to prevent foaming. *Calculate and use the <u>actual</u> concentration in % Dissolved equation
Sample Preparation:	One tablet is placed into each vessel of the dissolution apparatus. Sample each vessel after the incubation time, as stated above. Pass a portion of the sample through a 0.45 micron filter sufficient to equilibrate the filter. Filters are to be pre-qualified according to SOP (C1-730). Use a new filter for each vessel.
Procedure:	Inject 800 µl of standard and sample into the column and record the chromatograms. Measure the responses of the major peaks. Calculate the amount of Levothyroxine dissolved in each vessel by the formula below.

5

Calculations:
$$\frac{\text{Sample Area} \times 798.86 \times \text{Amt. Std. Injected}}{\text{Std. Area} \times 776.87 \times \text{Amt. Samp. Injected}} \times 100\% = \% \text{ Dissolved}$$

% Dissolved

Where 798.86 = molecular weight of Levothyroxine as Sodium Salt
776.87 = molecular weight of Levothyroxine (as Base)

10

Table 7 – Acceptance Criteria

STAGE	#TESTED	ACCEPTANCE CRITERIA Q = 70%
S-1	6	Each unit is not less than Q + 5%
S-2	6	Average of 12 units (S-1 + S-2) is equal to or greater than Q, and no unit is less than Q-15%
S-3	12	Average of 24 units (S-1 + S-2 + S-3) is equal to or greater than Q and not more than 2 units are less than Q-15%, and no unit is less than Q-25%

15

Table 8 shows comparative dissolution data for all strengths of Levoxyl[®] tablets.

Table 8 – Comparative Dissolution Data

	0 minutes	1 minute	2.5 minute	5 minutes	7.5 minutes	10 minutes
25 mcg	0.0%	84.9%	93.7%	90.9%	88.6%	84.7%
50 mcg	0.0%	82.8%	92.7%	91.8%	87.8%	84.4%
75 mcg	0.0%	78.9%	93.6%	92.2%	88.3%	84.7%
88 mcg	0.0%	79.8%	95.6%	94.1 %	90.5%	86.9%
100 mcg	0.0%	85.4%	94.8%	94.5%	90.7%	86.5%
112 mcg	0.0%	75.5%	91.1%	90.7%	87.0%	82.9%
125 mcg	0.0%	75.0%	96.5%	95.5%	91.7%	87.8%
137 mcg	0.0%	79.9%	93.9%	93.2%	89.4%	85.7%
150 mcg	0.0%	75.6%	91.9%	91.4%	88.7%	84.6%
175 mcg	0.0%	84.2%	95.7%	93.5%	90.3%	85.5%
200 mcg	0.0%	76.5%	94.9%	94.6%	91.0%	87.6%
300 mcg	0.0%	74.5%	92.1%	91.4%	87.9%	84.0%

5 Figure 4 depicts graphs showing the mean results for each of the tablet strengths of Levoxyl[®] tested. Each point is the mean of three dissolutions, testing 12 tablets per dissolution or n=36. The data is presented as percent of label claim dissolved vs. dissolution time.

10 The results demonstrate that the multi-point dissolution profiles for Levoxyl[®] tablets are similar across a wide variety of tablet strengths. Moreover, all strengths substantially exceed the requirements for immediate release oral dosage forms (i.e. at least 80% dissolved with 15 – 20 minutes). In each dosage form, these pills were over 90% dissolved within two and a half minutes.

15 The extremely rapid dispersion rates for the tablets of the present invention make possible a simplified treatment method for infants or others who have difficulty swallowing pills. In this approach, the appropriate dosage for the patient in question, in an immediate release pill made in accordance with the present invention, is simply mixed with a suitable amount, e.g. 50 – 200 ml, of aqueous fluid, such as water, soft drinks, juice, milk, etc. The immediate release

pill is easily dissolved in the fluid, optionally with stirring or shaking, and simply administered to the patient.

Example 3

5

Potency Test

The following method for testing potency of the tablets will sometimes be referred to herein as method number: **AM-003**. Alternatively, the tablet potency can be tested according to method **AM-021**. Method number: **AM-021** is the same as method number: **AM-003**, except the tablets are dissolved whole without
10 first grinding the tablets into a powder, as with method number: **AM-003**.

Method Reference:

USP 24 pp. 968-970

15

Chromatographic Conditions:

Mobile Phase: 65:35:0.05 H₂O: CAN: H₃PO₄ degassed and filtered; mobile phase composition may be altered to achieve a satisfactory resolution factor.

20

Column:

ACN, 4.6 mm x 25 to 30 cm

Flow Rate:

1.5 ml/minute

25

Detector:

Deuterium, set at 225 nm

Injection Volume:

100 ml

System Suitability:

- 5 Chromatograph 5 replicate injections of the standard preparation. Record the peak responses as directed under "Procedure".

- 1.0 RSD for the standard replicates must not be more than 2.0% for T₄.
- 10 2.0 Calculate the resolution factor R on one of the five replicates. The R-value must be greater than or equal to 5.0 to proceed. See Method QC-009.

15

Standard Preparation:

- Accurately weight 25 mg of USP Levothyroxine RS and transfer to an amber 250-ml volumetric flask. Add approximately 50ml extraction mobile phase. Let stand for 20 minutes with occasional swirling. Sonicate for 30 seconds. Gradually add more extraction solution and repeat sonication until no undissolved particles are observed. Dilute to volume with extraction solution. Mix well. The concentration of T₄ is about 100 µg/ml. Also dissolve an accurately weighed quantity of USP Liothyronine RS to yield about 100 mg/ml, done as above with USP Levothyroxine RS. Label this solution as stock T₃-A.

25

Stock Standard dilution:

1. Pipette 10.0 ml stock T₃-A into a 500 ml Type A volumetric flask.
- 30 2. Dilute to volume with Mobile Phase for a concentration of about 2 µg/ml. Mix well and label this solution as std. T₃-B.

3. Pipette 50.0 ml each from the T₄ and T₃-B stock standards and transfer into a 500-ml Type A volumetric flask

Dilute to volume with mobile phase and mix well. Label this standard as T₃/ T₄ working standard. The concentration of the working standard should be about 0.2 µg/ml T₃ and 10.0 µg/ml T₄.

Note:

Concentrations of Levothyroxine and Liothyronine require adjustments for water content.

Assay Preparation:

Weigh not less than the specified tablet quantity and calculate the average tablet weight. Crush tablets into a uniform fine powder with a mortar and pestle.

15 Tare a polypropylene weigh boat.

Accurately weigh (to 0.1 mg) a portion of the powder into the tared weigh boat using a preconditioned stainless steel scoop or spatula (either Teflon coated or uncoated). The spatula or scoop is preconditioned by dipping it into the powder. Use the *Sample Calculation* below to achieve 50 ml of a 10 µg/ml assay solution.

Record the sample weight taken. Carefully transfer the sample into an Erlenmeyer flask, reweigh the weigh boat and subtract the residual weight from the weight taken to obtain the actual sample weight. Pipette 50 ml of mobile phase into the flask. Cover the flask with parafilm, sonicate for approximately 10 seconds and vortex for approximately 235 seconds at a speed of 6 or greater. Observe sample preparation, and if clumping is noted, repeat the sonication and/or vortex steps. Centrifuge (~3,000rpm) for NLT 1 minute until a clear supernatant is achieved. Transfer a portion of the supernatant to an auto sampler vial.

For *In-Process* granulation analysis, use the theoretical tablet weight (0.1455 g) in place of (weight of tablets/number of tablets) in the formula below.

Sample Calculation:

5

$$\frac{\text{Weight of Tablets}}{\text{Number of Tablets}} \times 10\mu\text{g/ml} \times 50\text{ml} = \text{Amount to Weight Out per Assay Dose } (\mu\text{g})$$

10 Procedure:

Separately inject 100 μl of the sample onto the column. Record the responses of the analyte peak and calculate % label claim as follows.

Calculations:

15 $\text{Sample Area} \times \text{Std conc. } (\mu\text{g}) \times 50 \text{ ml} \times \text{avg. tablet weight in g} \times \frac{798.86}{776.87} = \mu\text{g/dose} \times 100 = \% \text{ Label Claim}$

Standard Area (ml) Actual Sample wt in g 776.87 Label Claim

20 Where 798.86 = molecular weight of Levothyroxine as the Sodium Salt
776.87 = molecular weight of Levothyroxine Standard Base

Results:

25 Figures 5A and 5B show HPLC chromatograms of levothyroxine and liothyronine controls (T3/T4 working standard, shown in Figure 5A) and an experimental sample made in accordance with the present invention as described above.(Figure 5B). The peaks in both chromatograms in the area of 1.325 to 3.1 correspond to materials in the solvent. The peak at about 7.2 in Fig. 5A shows the presence of T3. Fig. 5B shows the absence of T3, as well as the absence of other related products or degradation products of levothyroxine.

30

Example 4**Hardness Test**

The following preferred method for testing tablet hardness will sometimes be referred to herein as method number: **QC-005**

5

Table 9 - QC-005 Hardness Test Procedure

APPARATUS:	Van-Keel hardness tester; Please refer to equipment Profile for instrument information.
PROCEDURE:	Lay the tablet flat with the score side up onto the instrument in between the jaw area. The tablet's score line should be perpendicular to the jaw's line for the tablet to be aligned properly. Refer to alignment diagram below. For Tamil-K caplets, place the caplet onto the instrument on its side. The caplet's score line should not be laying on the flat part of the testing area as with other tablets but should not be parallel to the jaw's line for the caplet to be aligned properly. Refer to alignment diagram below. Push the test button on the control panel. The jaws will automatically move the break the tablet. The force needed to break the tablet (KP) will read out on the digital display and print out on the print tape. Specifications: 6.0 – 14.0 kiloponds
RESULTS:	Typical results range from about 9.3 to about 12.3 kiloponds.

Generally the hardness of the pills lies between about 6.0 and about 14.0 kiloponds. Preferably the pill hardness is from about 9 to about 13 kiloponds. Typical results of products made in accordance with the present invention are about 9.3, 11.3, 9.8, 10.2, 12.3, etc. Pharmaceutical tablets which incorporate granulated active ingredient are typically much higher in hardness, which may add to the difficulty of dissolving or dissoluting them. Pills which are lower in hardness generally present more problems of pill fragmentation during handling and storage.

Example 5**Impurity Tests**

The following preferred method for testing tablet impurities is sometimes referenced herein as method number: SA-004

5

Table 10 – SA 004 Impurity Test Procedure

Method Reference:	Biochemie Method No. 1417-6, Report JMI-DP-002
Equipment:	HPLC with a gradient system and a detector at a wavelength of 225 nm
Reagents:	Acetonitrile, HPLC grade Methanol, HPLC grade Water, HPLC grade Sodium Hydroxide, ACS reagent grade Sodium Hydroxide 0.1 solution: Dissolve 40g of NaOH pellets in 1000 ml HPLC grade water. Store in a plastic container. Phosphoric acid, 85% reagent grade Diiodothyronine reference material Liothyronine RS USP reference material Levothyroxine RS USP reference material Triiodothyroacetic acid reference material Tetraiodothyroacetic acid reference material <u>Solvent 1:</u> To 100.0 ml of 0.1 N Sodium Hydroxide solution add a 1:1 V/V mixture of methanol and water to make 1000 ml. <u>Solvent 2:</u> 77:23:0.1 H ₂ O: H ₃ PO ₄ ; Degassed and filtered; mobile phase composition a may be altered to achieve a satisfactory resolution factor. <u>Extraction solution:</u> Pipette 50 ml of solvent 1 into a 1000 ml volumetric flask dilute to volume with solvent 2, stopper and mix well.
Chromatography	
Column:	Nucleosil 100-10CN, 250 mm long, 4.6 mm internal diameter, at ambient temperature
System:	Gradient Elution Mobile phase A: 1000:1 H ₂ O:H ₃ PO ₄ V/V Mobile phase B: Acetonitrile Gradient program: Time min % of mobile phase A % of mobile phase B

0
77
23

13
77
23

15
65
35

24
65
35

26
77
23

Flow rate: 1.5 ml/min.

Injection Volume: 100 µl; next injection after approx. 40 min.

Detector: UV, 225 nm

System Suitability:

Chromatograph 5 replicate injections of the Reference I Standard preparation, chromatograph 2 replicate injections of the Reference II Standard. Record the peak responses as directed under "Procedure". An extraction blank is to be run after the standards.

1. The RSD must not be greater than 2.0% for each of the impurities in the standard reference solution I.
2. The resolution factor between liothyronine and levothyroxine in the standard reference solution I must not be less than 5.0.
3. The Signal to Noise ratio must not be less than 5/1 for levothyroxine and impurities in the chromatogram obtained with standard reference solution II.
4. A peak of monochlorotriiodothyronine may occur just before the levothyroxine peak. Make sure that the degree of separation between this peak and of levothyroxine is at least sufficient to permit separate evaluations.

Monochlorotriiodothyronine reference material is not available to be purchased by any vendor. Any calculation of monochlorotriiodothyronine impurity will be done by its retention time.

Standards

Preparation:

1. Stock Standard Reference Solution:
Accurately weigh 10 mg $\pm$ 0.1 mg of each Diiodothyronine, Liothyronine, Levothyroxine, Triiodothyroacetic acid and

Tetraiodothyroacetic acid reference standards into a 100ml volumetric flask. Dissolve in Solvent 1 and dilute to volume, stopper and mix well. The concentration of each component will be approximately 100mcg/mL.

2. Standard Reference solution I:

Pipette 5.0 ml of Stock Standard Reference Solution into a 100 ml volumetric flask, dilute to volume with Solvent 2, stopper and mix well. The Final concentration of each component will be approximately 5mcg/mL.

3. Standard Reference solution II (0.05%):

Pipette 2.0 ml of Standard Reference Solution I into a 100 ml volumetric flask, dilute to volume with Solvent 2, stopper and mix well. The final concentration of each component will be approximately 0.1 mcg/mL. 100

Test Preparation:

Crush not less than 20 tablets. Tare a 250 ml Erlenmeyer flask. Accurately weigh to the nearest 0.1 mg an equivalent of 500 mcg of levothyroxine sodium (+/- 10%) into a 250 ml Erlenmeyer flask. Pipette 100.0 mcg of the Extraction solution into the flask cover the flask with parafilm, sonicate, vortex and then centrifuge the solution for 1 minute each. The final concentration of the sample will be approximately 5 mcg/ml of levothyroxine. To calculate the amount to weigh for the test preparation use the following equation:

$$\frac{500 \text{ mcg} \times 0.1450 \text{ g}}{\text{tablet label claim (mcg)}} = \text{Amount to weight for the test prep}$$

* where 0.1450 g = theoretical tablet weight

Note: Analyst must keep all materials use in performing this assay until the results are calculated, checked, and recorded and it is verified that the test is acceptable. This includes the crush, the Erlenmeyer flask with Extraction solution, the centrifuge tube and the auto-sampler vial. If the analysis is running overnight, these materials should be sealed with parafilm and saved until results are obtained and the results are deemed acceptable

Procedure:

1. Separately inject 100µl of the sample preparation onto the column. Record the response of the analyte peaks and the calculate % w/w using the equations below.
2. The chromatogram may need to be reprocessed to obtain optimal integration. A copy of the sample chromatograph is to be attached to the analytical packet.
3. Peaks on the sample chromatograph with areas less than a

signal ratio of 5/1 will be considered none detected.

Calculations:

5 Diiodothyronine:

$$\frac{\text{Sample area}}{\text{Std. Area}} \times \text{Std conc. (mcg/ml)} \times \frac{100\text{ml}}{\text{Wsimpl (g)}} \times \frac{100\%}{1000000 \text{ mcg/g}} \times 1.11^* = \% \text{ w/w}$$

10 or

Sample area X Std. Conc. (mcg) X 0.01 X 1.11* = %w/w

*where 1.11 is a correction factor

15

Triiodothyroacetic Acid:

$$20 \quad \frac{\text{Sample area}}{\text{Std. Area}} \times \frac{\text{Std conc. (mcg)}}{\text{ml}} \times \frac{100\text{ml}}{\text{Wsimpl (g)}} \times \frac{100\%}{1000000 \text{ mcg/g}} = \% \text{ w/w}$$

or

$$25 \quad \frac{\text{Sample area}}{\text{Std area}} \times \frac{\text{Std. Conc. (mcg)}}{\text{ml}} \times \frac{0.01}{\text{W}_{\text{simpl}} (\text{g})} = \% \text{w/w}$$

Tetraiodothyroacetic Acid:

$$30 \quad \frac{\text{Sample area}}{\text{Std. Area}} \times \frac{\text{Std conc. (mcg)}}{\text{ml}} \times \frac{100\text{ml}}{\text{Wsimpl (g)}} \times \frac{100\%}{1000000 \text{ mcg/g}} \times 1.16^* = \% \text{ w/w}$$

or

$$35 \frac{\text{Sample area}}{\text{Std area}} \times \frac{\text{Std. Conc. (mcg)}}{\text{ml}} \times \frac{0.01}{\text{Wsimpl (g)}} \times 1.16^* = \%w/w$$

*where 1.16 is a correction factor

Limit of Detection (LOD) Values

Impurity	Limit of Detection
Diiodothyronine (T2)	0.00625%
Triiodothyroacetic Acid (Reverse T3)	0.003125%
Tetraiodothyroacetic Acid (Reverse T4)	0.003125%

Calculation of the theoretical area for 0.05% of levothyroxine sodium, based on the initial amount in mg of levathyroxine sodium in the whole sample weight.

$$\frac{(\text{Area}_{rs II})(A)(10.0)}{(0.05)(T_4 \text{ std st.})(P)(1.0283)} = \text{Theoretical area for 0.05\% of levothyroxine Na, based on the actual weight}$$

Where: Area_{rs} - is the average area of the levothyroxine in the Standard reference solution II
 A = is the initial weight of levothyroxine Na in mg represented by the sample weight.

This is calculated by using this equation: = $\frac{\text{sample weight (g)} \times \text{claim } T_4 \text{ in mcg}}{0.1450 \text{ g} \times 1000 \text{ mcg/mg}}$

10.0 = theoretical initial weight of the Levothyroxine USP reference standard

0.500 = is the theoretical initial weight of the Levothyroxine NA to be tested, in mg

T_4 std. Wt. = the initial weight of the levothyroxine USP standard in mg

P = the purity of the levothyroxine Na USP standard (%purity/100%)

1.0283 = conversion of levothyroxine into levothyroxine sodium

10

Greatest unknown impurity (individually):

$$\frac{(\text{Area}_{\text{impurity}})(T_4 \text{ std wt mg})(1.0283)(P)(100)}{(\text{Area ref std I})(A)(2000)} = \text{impurity (\%)}%$$

15

Where: $\text{Area}_{\text{impurity}}$ is the area of the greatest unknown impurity in the test solution with an area greater than the theoretical area for 0.05% of the levothyroxine Na taken into account.

20 1.0283 = conversion of levothyroxine into levothyroxine sodium

P = the purity of the levothyroxine Na USP standard (% purity/100%)

100 is the dilution of the test solution

Area ref std I is the area of the levothyroxine in the standard reference solution I

A = is the initial weight of levothyroxine Na in mg represented by the sample weight.

This is calculated by using this equation: = $\frac{\text{sample weight (g)} \times \text{claim T4 in mcg}}{0.1450\text{g} \times 1000 \text{ mcg/mg}}$

2000 is the dilution of the reference solution.

5

Total of other Unknown Impurities:

$\frac{(\text{Sum area impurities}) (\text{T4 std wt mg}) (1.0283) (P) (100)}{(\text{are ref std I}) (A) (2000)} = \text{Total Unknown impurities (\%)}$

10 Where: Sum area impurity is the sum of the areas of all the other unknown impurities in the test solution (only areas that are greater than the theoretical area for 0.05% of the levothyroxine sodium taken into account)

T4 std. wt. = the initial weight of the levothyroxine USP standard in mg

1.0283 = conversion of levothyroxine into levothyroxine sodium

15 P = the purity of the levothyroxine Na USP standard (% purity/100%)

100 is the dilution of the test solution

Area ref std I is the area of the levothyroxine in the standard reference solution I

A = is the initial weight of levothyroxine Na in mg represented by the sample weight.

20 This is calculated by using this equation: = $\frac{\text{sample weight (g)} \times \text{claim T4 in mcg}}{0.1450\text{g} \times 100 \text{ mcg/mg}}$

0.1450g X 100 mcg/mg

2000 is the dilution of the reference solution.

Results of the test are shown in Figures 6A and 6B. Figure 6A shows an
 25 example of a chromatogram of Standard Reference Solution II, with exemplary peaks at about 5.4 for diiodo-l-thyronine, 8.4 for liothyronine, 12.8 for levothyroxine, 19.3 for triiodo thyroacetic acid, and 21.9 for tetraiodo thyroacetic acid. Figure 6B shows results of an experimental sample of levothyroxine sodium, made in accordance with this invention. As can be seen, the sample had
 30 substantially only levothyroxine, with insignificant impurities.

Example 6**Liothyronine (T3) Tests**

The following preferred method for testing for Triiodothyronine is
 5 sometimes referenced herein as method number: **QC-001**

Table 11 – QC – 001 T3 Test Procedure

Method Reference	USP 24 p. 968-970
Chromatographic Conditions:	65:35:0.05 H ₂ O:CACN:H ₃ PO ₄ degassed and filtered; mobile phase composition may be altered to achieve a satisfactory resolution factor.
Mobile Phase:	
Column:	CN, 4.6 mm x 25 to 30 cm
Flow Rate:	2.0 minute/minute
Detector:	Deuterium, set at 225 nm
Injection Volume:	100 µL
System Suitability:	Chromatograph 5 replicate injections of the standard preparation. Record the peak responses as directed under "Procedure".
	1.0 RSD for the standard replicates must not be more than 2.0% for T ₄
	2.0 Calculate the resolution factor (R) on one of the five replicates. The R value must be greater than or equal to proceed. See Method QC-009.
Standard Preparation:	Accurately weigh 25 mg of USP Levothyroxine RS and transfer to a clear 250-mL volumetric flask. Pipette 87.5 ml minute of acetonitrile in the flask. Swirl and then sonicate for less than a minute. Add portions of HPLC grade water to the flask with swirling and sonicating until the material has gone into solution. Be sure that there is no particulate material present. Do not dilute to volume at this point. The solution may be cold. Place into a room temperature water bath for ten minutes to allow the sample to warm to ambient temperature. Dilute to volume with HPLC grade water. Mix well. Label this solution as stock T₄. The concentration of T₄ is about 100 µg/ml. Also dissolve an accurately weighed quantity of USP Liothyronine RS to yield about 100 µg/minute, done as above with USP Levothyroxine RS. Label this solution as stock T₃-A. Stock Standard dilution: 1. Pipette 10.0 ml stock T₃-A into a 500-mL Type A

- volumetric flask.
2. Dilute to volume with Mobile Phase for a concentration of about 2 µg/ml. Mix well and label this solution as stock std. c-B.
 3. Pipette 50.0 ml each from the T₄ and T₃ stock standards and transfer into 500-ml Type A volumetric flask.

Dilute to volume with mobile phase and mix well. Label this standard as T₃/T₄ working standard. The concentration of the working standard should be about 0.2 µg/ml T₃ and 10.0 µg/ml T₄.

Assay Preparation:

Weigh and crush not less than the specified tablet quantity and calculate the average tablet weight. Tare a polypropylene weigh boat.

Accurately weigh (to 0.1 mg) a portion of the powder into the tared weigh boat using a preconditioned stainless steel scoop or spatula (either Teflon coated or uncoated). The spatula or scoop is preconditioned by dipping it into the power. Use the *Sample Calculation* below to achieve 50 ml of a 10 µg/ml assay solution.

Record the sample weight taken. Carefully transfer the sample into an Erlenmeyer flask, reweigh the weigh boat and subtract the residual weight from the weight taken to obtain the actual sample weight. Pipette 50 ml of mobile phase into the flask. Cover the flask with parafilm, sonicate for approximately 10 seconds and vortex for approximately 35 seconds at a speed of 6 or greater. Observe sample preparation, and if clumping is noted, repeat the sonication and/or vortex steps. Centrifuge (~3,000 rpm) for NLT 1 minute until a clear supernatant is achieved. Transfer a portion of the supernatant to an autosampler vial.

For *In-Process* granulation analysis, use the theoretical tablet weight (0.1455 g) in place of (weight of tablets/number of tablets) in the formula below.

Note

Analyst must keep all materials used in performing this assay until the results are calculated, checked, and recorded, and it is verified that the test is acceptable. This includes the crush, the Erlenmeyer flask with Mobile Phase, the centrifuge tube and the autosampler vial. If the analysis is running overnight, these materials should be sealed with parafilm and saved until results are obtained and the result is deemed acceptable.

Sample Calculation: $\frac{\text{Weight of Tablets}}{\text{Number of Tablets}} \times 10 \mu\text{g/ml} \times 50 \text{ ml} = \frac{\text{Amount to Weigh Out}}{\text{per Assay}}$

Procedure: Separately inject 100 μl of the sample onto the column. Record the responses of the analyte peak.

Calculations: Calculate the content of liothyronine using the following formula:

$$5 \quad \frac{\text{Sample } T_3 \text{ Area}}{\text{Standard } T_3 \text{ Area}} \times \frac{\text{Std } T_3 \text{ conc. } (\mu\text{g})}{(\text{ml})} \times 50 \text{ ml} = \mu\text{g } T_3$$

The specification is NGT 2.0% liothyronine calculated as follows:

$$10 \quad \frac{\text{Amt } T_3 \text{ Assayed } (\mu\text{g})}{\text{Amt } T_4 \text{ Assayed } (\mu\text{g})} \times 100 = \% \text{ LIOETHYRONINE}$$

*This number is calculated using the T_4 potency results as follows:

$$15 \quad \frac{\text{Sample } T_4 \text{ Area}}{\text{Standard } T_4 \text{ Area}} \times \frac{\text{Std } T_4 \text{ conc. } (\mu\text{g})}{(\text{ml})} \times 50 \text{ ml} \times \frac{798.86}{776.87} = \mu\text{g } T_4$$

20 where 798.86 = molecular weight of Levothyroxine as the Sodium Salt
776.87 = molecular weight of Levothyroxine Standard Base

NOTE: If the single active ingredient comprises 50% or more, by weight, of the dosage unit, use Method A; otherwise use Method B.

METHOD: USP 24 <905> pp. 2000-2002.

METHOD A: Content Uniformity as Determined by Weight Variation:
Weight accurately 10 tablets, individually. From the results of the average potency of the active ingredient determined for the product (using the assay methods as stated in the individual monograph) calculate the content of active ingredient in each of the 10 tablets.

CALCULATIONS: $\text{Individual} = \frac{(\text{Avg. potency}) (\text{Individual Wt.})}{\text{Potency} \quad \text{Avg. tablet weight}}$

NOTE: If the active ingredient(s) are less than 50% by weight of the tablet content, refer to the individual test method for potency for those products.

METHOD B: Content Uniformity as Determined by Direct Assay of Active

Ingredient:

For Levothyroxine Sodium tablets the following procedure is followed.

Individually weigh 10 tablets. Place the 10 individual tablets into round bottomed test tubes or flasks of the appropriate size as outlined in the chart below. Add the appropriate volume of extraction mobile comprised of water, acetonitrile, and phosphoric acid (65:35::0.05) to each test tube or flask as indicated in the chart below. Note: All test tubes are to be capped with screw on caps and all flasks are to be covered with parafilm as soon as mobile phase is added. Allow to stand at room temperature until the tablet completely crumbles. Secure all samples in a wrist action shaker. Test tubes are to be secured horizontally. Erlenmeyer flasks are to be secured vertically. Set the wrist shaker to the setting specified in the table. Shake sample for 3 minutes. Transfer about 10 ml of the sample preparation (or the entirety of smaller samples) to a centrifuge tube. Centrifuge samples for 1 minute at about 3000 rpm. Transfer samples to autosampler vials using disposable Pasteur pipettes.

Utilize the HPLC Method for levothyroxine separation (AM-003) for obtaining dosage uniformity, sample area, and standard area results.

CALCULATIONS: Dosage Uniformity Result (% Label Claim)

$$\frac{798.86}{776.87} \times \frac{\text{Area of Sample}}{\text{Area of Std.}} \times \frac{\text{Conc. of Std.}}{\text{Conc. Of Sample (see chart below)}} \times 100 = \% \text{ Potency}$$

SPECIFICATIONS FOR METHOD A OR METHOD B

S-1 The % active ingredient for 10 tablets tested must fall in the range of 85.0% - 115.0% and the RSD of the 10 tablets must not exceed 6.0%.

NOTE: If 1 unit in S-1 fails to meet either of the specifications, but is no outside the range of 75% - 125%, test 20 more units and proceed to S-2.

S-2 When n = 30, NGT one unit outside 85.0-115.0%, none outside 75.0-125.0% and RSD NGT 7.80%.

Results:

- 5 Results for a variety of dosages, using a sample size of 120 pills, are shown in Table 12:

Table 12 – Dosage Consistency – 120 pill samples

Dosage	25 µg	100 µg	300 µg
Label Claim	103.5 %	103.1 %	102.9 %
Activity			
High	109.1 %	104.8 %	108.8%
Low	98.0 %	100.7 %	96.5 %
RSD	< 2.0 %	0.9 %	2.2 %

The results confirm an extremely low amount of variability in active material content between the 120 pills tested. Generally the variability for a 120 pill sample should be between about 90 and about 110 % of claimed activity, preferably between about 95 % and about 105%. The RSD for a 120 pill sample should not be greater than 5%, and preferably is less than 3%.

Example 7

Levothyroxine Sodium Release Specification and Analytical Methods

The specifications for levothyroxine sodium tablets are stated in: USP 24 page 969-970 and Supplement 1 page 2638. The additional requirements are in place to ensure the tablet appearance, for the individual tablet strengths, is correct and the physical characteristics ensure a quality tablet.

A. Analytical Methods:

All the test methods utilized in the testing of levothyroxine sodium meet USP system suitability requirements. All Levoxyl® batches are tested for conformance to the following specifications. The **Table 13** below lists the test parameter, specification and the test method employed.

Table 13 – USP Specifications

Test Parameter	Specification	Test Method
Tablet Potency	90.0-110.0% label claim *	AM-003
Tablet Dissolution	NULT 7580% label claim dissolved in 145 minutes	AM-004B
Liothyronine Content	NGT 2.0%	QC-001
TLC Identification	Compares to Standard	RM-054
Uniformity of Dosage Units	S-1: 85.0-115.0% RSD NGT 6.0% n=10 (if NGT 1 unit fails, but no unit is outside range of 75.0-125.0% or if RSD fails proceed to S-2) S-2: When n=30 NGT 1 unit outside 85.0-115.0%, none outside 75.0-125.0% and RSD NGT 7.8%	QC-003

5

Additional Requirements:

Test Parameter	Specification	Test Method
Tablet Hardness	6.0-14.0 KP	QC-005
Tablet Weight	142.0-149.0 mg	QC-007
Tablet Appearance	Color, imprint, score and shape conform to specific tablet parameters as specified for the individual strengths	QC-008

Example 8**Bioavailability determination of Two Levothyroxine formulations**

15 The following example was performed along lines of a 1999 FDA publication entitled *In-Vivo Pharmacokinetics and Bioavailability Studies and In-Vitro Dissolution Testing for Levothyroxine Sodium Tablets*. The example includes the following two studies.

Study 1. Single-Dose Bioavailability Study

The objective of the study was to determine the bioavailability of Levoxyl® relative to a reference (oral solution) under fasting conditions.

5 Study 2: Dosage-Form Equivalence Study

The objective of the study was to determine the dosage-form bioequivalence between three different strengths of Levoxyl® tablets (low, middle and high range).

10 Study Objective:

To determine the bioavailability of levothyroxine sodium (Levoxyl®) 0.3 mg tablets manufactured by JONES PHARMA INCORPORATED, relative to Knoll Pharmaceutical Company's levothyroxine sodium 200 µg (Synthroid®) injection given as an oral solution following a single 0.6mg dose.

15

Study Methodology:

Single-dose, randomized, open-label, two-way crossover design

Protocol Reference:

20 Guidance for Industry: In Vivo Pharmacokinetics and Bioavailability Studies and In Vitro Dissolution Testing for Levothyroxine Sodium Tablets (June 1999).

Number of Subjects:

25 A total of 30 subjects were enrolled in the study, and 27 subjects completed the study. All 30 subjects were included in the safety analysis and 27 subjects who completed the study were included in the pharmacokinetic analyses.

Diagnosis and Main Criteria for Inclusion:

All subjects enrolled in this study were judged by the investigator to be healthy volunteers who met all inclusion and exclusion criteria.

5 Test Product, Dose, Duration, Mode of Administration, and Batch Number:

The test product was levothyroxine sodium (Levoxyl®) 2 X 0.3mg tablets administered as a single oral dose. The batch number utilized in this study was TT26.

10 Reference Product, Dose, Duration, Mode of Administration, and Batch Number:

The reference product was levothyroxine sodium (Synthroid®) 2 X 500 µg injection vials (Knoll Pharmaceutical Company) reconstituted and 600 µg administered orally. The reference product used was the 500 µg injection instead
15 of 200 µg due to the unavailability of sufficient quantities of 200 µg injection to conduct the study. The batch number utilized in this study was 80130028.

Criteria for Evaluation:*Pharmacokinetics:*

20 Pharmacokinetic assessment consisted of the determination of total (bound + free) T4 and T3 concentrations in serum at specified time points following drug administration. From the serum data, the parameters AUC(0-t), Cmax, and Tmax were calculated.

25 Safety:

Safety assessment included vital signs, clinical laboratory evaluation (including TSH), physical examination, and adverse events (AEs) assessment.

Statistical Methods

Pharmacokinetics:

Descriptive statistics (arithmetic mean, standard deviation (SD), coefficient of variation (CV), standard error of the mean (SE), sample size (N),
5 minimum, and maximum) were provided for all pharmacokinetic parameters. The effects of baseline and baseline-by treatment interaction were evaluated using a parametric (normal-theory) general linear model (ANCOVA) with treatment, period, sequence, subject within sequence, $\ln(\text{baseline})$, and interaction between
10 $\ln(\text{baseline})$ and treatment as factors, applied to the $\ln$ -transformed pharmacokinetic parameters and $C_{\max}$. In the absence of significant $\ln(\text{baseline})$ and interaction between $\ln(\text{baseline})$ and treatment, these parameters were removed from the model. The two one-sided hypotheses were tested at the 5% level of significance for $\ln[\text{AUC}(0-t)]$ and $\ln(C_{\max})$ by constructing 90% confidence intervals for the ratio of Treatment A to Treatment

15

Safety:

Frequency counts of all subjects enrolled in the study, completing the study, and discontinuing early were tabulated. Descriptive statistics were calculated for continuous demographic variables, and frequency counts were
20 tabulated for categorical demographic variables for each gender and overall.

AEs were coded using the 5th Edition of the COSTART dictionary. AEs were summarized by the number and percentage of subjects experiencing each coded event. A summary of the total number of each coded event and as a percentage of total AEs was also provided.

25 Laboratory summary tables included descriptive statistics for continuous serum chemistry and hematology results at each time point. Out-of-range values were listed by subject for each laboratory parameter.

Descriptive statistics for vital sign measurements at each time point and change from baseline to each time point were calculated by treatment group. Shifts from screening to post study results for physical examinations were tabulated.

5

Pharmacokinetic Results – T4:

ANCOVA analyses indicated that the effects of In(baseline) and interaction between In(baseline) and treatment were not significant. Thus, these factors were removed from the general linear model and an ANOVA with treatment, period, sequence, and subject within sequence was applied to the In-transformed Cmax and AUC(0-t) parameters. The arithmetic means of serum T4 pharmacokinetic parameters for Treatments A and B and the statistical comparison for In-transformed parameters are summarized in the following table.

15

**Summary of the Pharmacokinetic Parameters
of Serum T4 for Treatments A and B**

Pharmacokinetic Parameters	Treatment A*		Treatment B**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
Cmax (µg/dL)	14.48	1.93	15.09	2.10	-----	-----
Tmax (hr)	2.17	0.810	1.62	0.502	-----	-----
AUC(0-t) (µg*hr/dl)	524.3	59.07	529.3	62.83	-----	-----
In (Cmax)	2.663	0.1434	2.705	0.1339	91.1-98.1	94.5
In [AUC(0-t)]	6.256	0.1167	6.265	0.1169	95.6-100.5	98.0

* Treatment A = 2 X 0.3mg Levoxyl® Tablets: test

20 **Treatment B = 0.6mg Synthroid Reconstitute Oral Solution: reference

Pharmacokinetics Results – T3:

ANCOVA analyses indicated that the effects of In(baseline) and interaction between In(baseline) and treatment were not significant and were removed from the ANOVA model, except for In(baseline) on In(Cmax) which

was significant and was kept in the model. An ANOVA with treatment, period, sequence, and subject within sequence, and In(baseline), when significant, was applied to the In-transformed Cmax and AUC(0-t) parameters. The arithmetic means of serum T3 pharmacokinetic parameters for Treatments A and B and the statistical comparison for In-transformed parameters are summarized in the following table.

**Summary of the Pharmacokinetic Parameters
of Serum T3 for Treatments A and B**

Pharmacokinetic Parameters	Treatment A*		Treatment B**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
Cmax (ug/ml)	1.165	0.156	1.140	0.119	-----	-----
Tmax (hr)	14.6	15.2	16.3	17.0	-----	-----
AUC(0-t) (ng*hr/ml)	51.25	6.163	50.07	5.311	-----	-----
In (Cmax)	0.1444	0.1289	0.1255	0.1034	96.8-103.4	100.0
In [AUC(0-t)]	3.930	0.1209	3.908	0.1059	97.7-103.8	100.7

* Treatment A = 2 X 0.3mg Levoxyl® Tablets: test

**Treatment B = 0.6mg Synthroid Reconstitute Oral Solution: reference

Comparison of total T4 and T3 pharmacokinetics following administration of Levoxyl® (Treatment A, test formulation) and Synthroid (Treatment B, reference formulation) indicated that the test formulation met the requirements for bioequivalence with the reference formulation.

The 90% confidence intervals for the comparisons of In(Cmax) and In[AUC(0-t)] for T4 and T3 were within the 80% to 125% range required for bioequivalence.

In regard to subject safety, both treatments appeared to be equally safe and well tolerated.

Example 9**Bioavailability Study To Assess Single Dose
Bioequivalence of Three Strengths of Levothyroxine**

5 The following example was performed to determine the dosage-form
bioequivalence between three different strengths of levothyroxine sodium
(Levoxyl[®]) tablets following a single 600 mcg dose.

Study Methodology:

10 Single-dose, randomized, open-label, three-way crossover design.

Protocol Reference:

Guidance for Industry: In Vivo Pharmacokinetics and Bioavailability
Studies and In Vitro Dissolution Testing for Levothyroxine Sodium Tablets (June
15 1999). This protocol was submitted in IND 59,177.

Number of Subjects:

A total of 28 subjects were enrolled in the study, and 24 subjects
completed the study. All 28 subjects were included in the safety analysis and 24
20 subjects who completed the study were included in the pharmacokinetic analyses.

Diagnosis and Main Criteria for Inclusion:

All subjects enrolled in this study were judged by the investigator to be
healthy volunteers who met all inclusion and exclusion criteria.
25

Test Product, Dose, Duration, Mode of Administration, and Batch Number:

Subjects randomized to Treatment A received a single oral dose of 12 X
50 mcg levothyroxine sodium (Levoxyl[®]) tablets, Lot No. TT24. Subjects

randomized to Treatment B received 6 X 100 mcg levothyroxine sodium (Levoxyl[®]) tablets, Lot No.TT25. Subjects randomized to Treatment C received 2 X 300 mcg levothyroxine sodium (Levoxyl[®]) tablets, Lot No. TT26. Test products were manufactured by JMI-Daniels, a subsidiary of Jones Pharma
5 Incorporated.

Pharmacokinetics:

Pharmacokinetic assessment consisted of the determination of total (bound + free) T4 and T3 concentrations in serum at specified time points following drug
10 administration. From the serum data, the parameters AUC(0-t), Cmax, and Tmax were calculated.

Safety:

Safety assessment included monitoring of sitting vital signs, clinical
15 laboratory measurements, thyroid-stimulating hormone (TSH), physical examination, electrocardiogram (ECG), and adverse events (AEs).

Statistical Methods.

Pharmacokinetics:

20 Descriptive statistics (arithmetic mean, standard deviation (SD), coefficient of variation (CV), standard error of the mean (SEM), sample size (N), minimum, and maximum) were provided for all pharmacokinetic parameters. A parametric (normal-theory) general linear model with treatment, period, sequence, and subject within sequence as factors was applied to the In-transformed Cmax
25 and AUC(0-t). The two one-sided hypotheses were tested at the 5% level of significance for In[AUC(0-t)] and In(Cmax) by constructing 90% confidence

intervals for the ratios of Treatment A to Treatment B, Treatment A to Treatment C, and Treatment B to Treatment C.

Safety:

- 5 Frequency counts of all subjects enrolled in the study, completing the study, and discontinuing early were tabulated. Descriptive statistics were calculated for continuous demographic variables, and frequency counts were tabulated for categorical demographic variables for each gender and overall.

- 10 AEs were coded using the 5th Edition of the COSTART dictionary. AEs were summarized by the number and percentage of subjects experiencing each coded event. A summary of the total number of each coded event and as a percentage of total AEs was also provided. Laboratory summary tables included descriptive statistics for continuous serum chemistry and hematology results at each time point. Out-of-range values were listed by subject for each laboratory
15 parameter. Descriptive statistics for vital sign measurements at each time point and change from baseline to each time point were calculated by treatment group. Shifts from screening to post study results for physical examinations were tabulated.

20 Pharmacokinetic Results – T4:

The arithmetic means of serum T4 pharmacokinetic parameters for Treatments A and B and the statistical comparison for the In-transformed parameters are summarized in the following table.

25

**Summary of the Pharmacokinetic Parameters
of Serum T4 for Treatments A and B**

Pharmacokinetic Parameters	Treatment A*		Treatment B**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
C _{max} (µg/dl)	13.70	1.82	14.13	1.48	-----	-----
T _{max} (hr)	2.37	1.04	1.98	0.827	-----	-----
AUC(0-t) (µg*hr/dl)	509.0	58.36	528.3	72.41	-----	-----
In (C _{max})	2.609	0.1378	2.643	0.1095	93.6-100.1	96.8
In [AUC(0-t)]	6.226	0.1200	6.261	0.1379	93.4-100.0	96.7

5 * Treatment A = 12 X 50 mcg Levoxyl® Tablets

 **Treatment B = 6 X 100 mcg Levoxyl® Tablets

 The arithmetic means of serum T4 pharmacokinetic parameters for
Treatments A and C and the statistical comparison for the In-transformed
10 parameters are summarized in the following table.

**Summary of the Pharmacokinetic Parameters
of Serum T4 for Treatments A and C**

Pharmacokinetic Parameters	Treatment A*		Treatment C**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
C _{max} (µg/dl)	13.70	1.82	14.15	1.50	-----	-----
T _{max} (hr)	2.37	1.04	2.40	1.09	-----	-----
AUC(0-t) (µg*hr/dL1)	509.0	58.36	528.7	57.13	-----	-----
In (C _{max})	2.609	0.1378	2.644	0.1085	93.6-100.1	96.8
In [AUC(0-t)]	6.226	0.1200	6.265	0.1089	93.1-99.7	96.4

15

 * Treatment A = 12 X 50 mcg Levoxyl® Tablets

 **Treatment C = 2 X 300 mcg Levoxyl® Tablets

The arithmetic means of serum T4 pharmacokinetic parameters for Treatments B and C and the statistical comparison for the In-transformed parameters are summarized in the following table.

5 Pharmacokinetic Results – T4 (Continued):

**Summary of the Pharmacokinetic Parameters
of Serum T4 for Treatments B and C**

Pharmacokinetic Parameters	Treatment B*		Treatment C**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
C _{max} (µg/dl)	14.13	1.48	14.15	1.50	-----	-----
T _{max} (hr)	1.98	0.827	2.40	1.09	-----	-----
AUC(0-t) (µg*hr/dl)	528.3	72.41	528.7	57.13	-----	-----
ln (C _{max})	2.643	0.1095	2.644	0.1085	96.7-103.4	100.0
ln [AUC(0-t)]	6.261	0.1379	6.265	0.1089	96.4-103.1	99.7

10

* Treatment B = 6 X 100 mcg Levoxyl® Tablets

**Treatment C = 2 X 300 mcg Levoxyl® Tablets

15 Pharmacokinetic Results – T3:

The arithmetic means of serum T3 pharmacokinetic parameters for Treatments A and B and the statistical comparison for the In-transformed parameters are summarized in the following table.

20

25

**Summary of the Pharmacokinetic Parameters
of Serum T3 for Treatments A and B**

Pharmacokinetic Parameters	Treatment A*		Treatment B**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
Cmax (ng/ml)	1.173	0.138	1.142	0.133	-----	-----
Tmax (hr)	12.9	19.0	12.1	16.1	-----	-----
AUC(0-t) (ng*hr/ml)	49.43	6.872	50.35	8.994	-----	-----
In (Cmax)	0.1523	0.1226	0.1264	0.1194	98.1-107.3	102.6
In [AUC(0-t)]	3.890	0.1538	3.905	0.1731	93.1-104.3	98.5

5 * Treatment A = 12 X 50 mcg Levoxyl® Tablets

 **Treatment B = 6 X 100 mcg Levoxyl® Tablets

 The arithmetic means of serum T3 pharmacokinetic parameters for
Treatments A and C and the statistical comparison for the In-transformed
10 parameters are summarized in the following table.

Pharmacokinetic Results – T3 (Continued):

**Summary of the Pharmacokinetic Parameters of
Serum T3 for Treatments A and C**

15

Pharmacokinetic Parameters	Treatment A*		Treatment C**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
Cmax (ng/ml)	1.173	0.138	1.167	0.169	-----	-----
Tmax (hr)	12.9	19.0	11.5	16.4	-----	-----
AUC(0-t) (ng*hr/ml)	49.43	6.872	49.36	7.680	-----	-----
In (Cmax)	0.1523	0.1226	0.1437	0.1491	96.3-105.4	100.7
In [AUC(0-t)]	3.890	0.1538	3.886	0.1705	94.7-106.2	100.3

* Treatment A = 12 X 50 mcg Levoxyl® Tablets

**Treatment C = 2 X 300 mcg Levoxyl® Tablets

The arithmetic means of serum T3 pharmacokinetic parameters for Treatments B and C and the statistical comparison for the In-transformed parameters are summarized in the following table.

5 **Summary of the Pharmacokinetic Parameters of Serum T3 for Treatments B and C**

Pharmacokinetic Parameters	Treatment B*		Treatment C**		90% CI	% Mean Ratio
	Arithmetic Mean	SD	Arithmetic Mean	SD		
Cmax (ng/ml)	1.142	0.133	1.167	0.169	-----	-----
Tmax (hr)	12.1	16.1	11.5	16.4	-----	-----
AUC(0-t) (ng*hr/ml)	50.35	8.994	49.36	7.680	-----	-----
In (Cmax)	0.1264	0.1194	0.1437	0.1491	93.9-102.7	98.2
In [AUC(0-t)]	3.905	0.1731	3.886	0.1705	96.2-107.8	101.8

* Treatment B = 6 X 100 mcg Levoxyl[®] Tablets

10 **Treatment C = 2 X 300 mcg Levoxyl[®] Tablets

Safety Results:

There was a total of 59 treatment-emergent AEs reported by 15 (54%) of the 28 subjects dosed with study treatment. Incidence of AEs was similar across treatments. Headache was the most frequently reported event. The majority of the AEs were mild in intensity. There was one subject who experienced a serious adverse event of chest pain, considered by the Investigator to be unrelated to treatment. No trends were noted in vital signs, clinical laboratory results, or ECGs to suggest treatment-related differences.

20 Comparison of total T4 and T3 pharmacokinetics following administration of 12 X 50 mcg Levoxyl[®] tablets (Treatment A) and 6 X 100 mcg Levoxyl[®] tablets (Treatment B) indicated that the two formulations met the requirements for bioequivalence. The 90% confidence intervals for the comparisons of In(Cmax)

and $\ln[AUC(0-t)]$ for T4 and T3 were within the 80% to 125% range required for bioequivalence.

Comparison of total T4 and T3 pharmacokinetics following administration of 12 X 50 mcg Levoxyl® tablets (Treatment A) and 2 X 300 mcg Levoxyl® tablets (Treatment C) indicated that the two formulations met the requirements for bioequivalence. The 90% confidence intervals for the comparisons of $\ln(C_{max})$ and $\ln[AUC(0-t)]$ for T4 and T3 were within the 80% to 125% range required for bioequivalence.

Comparison of total T4 and T3 pharmacokinetics following administration of 6 X 100 mcg Levoxyl® tablets (Treatment B) and 2 X 300 mcg Levoxyl® tablets (Treatment C) indicated that the two formulations met the requirements for bioequivalence. The 90% confidence intervals for the comparisons of $\ln(C_{max})$ and $\ln[AUC(0-t)]$ for T4 and T3 were within the 80% to 125% range required for bioequivalence.

The test formulations appear to be safe and generally well tolerated when given to healthy adult volunteers.

Example 10

Levothyroxine sodium (Levoxyl®) Tablet Compositions

The following preferred levothyroxine sodium compositions in tablet form were made along lines disclosed herein.

Levoxyl® 25 mcg Tablets

Color: orange; Markings: (front) dp/25 (back) LEVOXYL®

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.025 mg
β - Form Microcrystalline Cellulose, NF (Ceolus)	108.529 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
FD&C Yellow # 6	0.352 mg
Magnesium Stearate, NF	1.018 mg

Levoxyl® 50 mcg Tablets**Color: white; Markings: (front) dp/50 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.050 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.856 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Magnesium Stearate, NF	1.018 mg

5

Levoxyl® 75 mcg Tablets**Color: purple; Markings: (front) dp/75 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.075 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.438 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1609 (Blend of D&C Red # 30 and FD&C Blue # 1)	0.383 mg
Magnesium Stearate, NF	1.018 mg

10

Levoxyl® 88 mcg Tablets**Color: olive; Markings: (front) dp/88 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.088 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.311 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1607 (Blend of FD&C Yellow # 6, D&C Red # 30 and FD&C Blue # 1)	0.507 mg
Magnesium Stearate, NF	1.018 mg

15

20

Levoxyl® 100 mcg Tablets**Color: yellow; Markings: (front) dp/100 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.100 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.406 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-282 (Blend of FD&C Yellow # 6 and D&C Yellow # 10)	0.388 mg
Magnesium Stearate, NF	1.018 mg

5

Levoxyl® 112 mcg Tablets**Color: rose; Markings: (front) dp/112 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.112 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	107.711 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1610 (Blend of FD&C Yellow # 6, D&C Red # 30 and FD&C Red # 40)	1.080 mg
Magnesium Stearate, NF	1.018 mg

10

Levoxyl® 125 mcg Tablets**Color: brown; Markings: (front) dp/125 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.125 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.701 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1617 (Blend of D&C Yellow # 10 and FD&C Red # 40)	0.080 mg
Magnesium Stearate, NF	1.018 mg

15

Levoxyl® 137 mcg Tablets**Color: dark blue; Markings: (front) dp/137 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.137 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.288 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
FD&C Blue # 1	0.478 mg
Magnesium Stearate, NF	1.018 mg

5

Levoxyl® 150 mcg Tablets**Color: blue; Markings: (front) dp/150 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.150 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.645 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1612 (Blend of D&C Red # 30 and FD&C Blue # 1)	0.108 mg
Magnesium Stearate, NF	1.018 mg

10

Levoxyl® 175 mcg Tablets**Color: turquoise; Markings: (front) dp/175 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.175 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.397 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-334 (Blend of D&C Yellow # 10, and FD&C Blue # 1)	0.334 mg
Magnesium Stearate, NF	1.018 mg

15

Levoxyl® 200 mcg Tablets**Color: pink; Markings: (front) dp/200 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.200 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.515 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1613 (Blend of D&C Yellow # 10 and D&C Red # 30)	0.188 mg
Magnesium Stearate, NF	1.018 mg

5

Levoxyl® 300 mcg Tablets**Color: green; Markings: (front) dp/300 (back) LEVOXYL®**

Component	Quantity in mg/Tablet
Levothyroxine Sodium, USP	0.300 mg
β- Form Microcrystalline Cellulose, NF (Ceolus)	108.451 mg
Croscarmellose Sodium, NF (Ac-di-sol)	35.079 mg
Lake Blend # LB-1614 (Blend of FD&C Yellow # 6, D&C Yellow # 10 and FD&C Blue # 1)	0.142 mg
Magnesium Stearate, NF	1.018 mg

10 While the present invention has been described in the context of preferred
embodiments and examples, it will be readily apparent to those skilled in the art
that other modifications and variations can be made therein without departing
from the spirit or scope of the present invention. For example, the active moiety
levothyroxine sodium can be changed to liothyronine sodium and similar products
15 and still be considered as part of the claimed invention. Accordingly, it is not
Intended that the present invention be limited to the specifics of the foregoing
description of the preferred embodiments and examples, but rather as being
limited only by the scope of the invention as defined In the claims appended
hereto.

20 Having described our invention, we claim:

1. A pharmaceutical composition comprising a levothyroxine salt and a pharmaceutically acceptable carrier, wherein the pharmaceutical composition exhibits a levothyroxine (T4) plasma C_{max} of between from about 10 µg/dl to about 20 µg/dl as determined by a standard C_{max} test.
5
2. The composition of claim 1, wherein the composition exhibits a levothyroxine (T4) plasma C_{max} of between from about 12 µg/dl to about 16 µg/dl as determined by the standard C_{max} test.
- 10 3. The composition of claim 1, wherein the ln(C_{max}) of the levothyroxine (T4) plasma level is between from about 1 to about 3.
4. The composition of claim 2, wherein the ln(C_{max}) of the levothyroxine (T4) plasma level is between from about 1 to about 3.
15
5. The composition of claim 1, wherein the composition exhibits a triiodothyronine (T3) plasma C_{max} of between from about 0.1 ng/mL to about 10ng/mL as determined by the standard C_{max} test.
20
6. The composition of claim 1, wherein the composition exhibits a triiodothyronine (T3) plasma C_{max} of between from about 0.5 ng/mL to about 2ng/mL as determined by the standard C_{max} test.
- 25 7. The composition of claim 1, wherein the ln(C_{max}) is between from about 0.01 to about 5.

8. The composition of claim 1, wherein the composition exhibits a levothyroxine (T4) plasma Tmax of between from about 0.5 hours to about 5 hours as determined by a standard Tmax test.
- 5 9. The composition of claim 1, wherein the composition exhibits a levothyroxine (T4) plasma Tmax of between from about 1 hour to about 3 hours as determined by the standard Tmax test.
- 10 10. The composition of claim 1, wherein the composition exhibits a triiodothyronine (T3) plasma Tmax of between from about 10 hours to about 20 hours as determined by the standard Tmax test.
- 15 11. The composition of claim 1, wherein the composition exhibits a triiodothyronine (T3) plasma Tmax of between from about 12 hours to about 16 hours as determined by the standard Tmax test.
- 20 12. The composition of any of claims 1, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
13. The composition of any of claims 2, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 25 14. The composition of any of claims 3, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.

15. The composition of any of claims 4, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 5 16. The composition of any of claims 5, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 10 17. The composition of any of claims 6, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 15 18. The composition of any of claims 7, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 20 19. The composition of any of claims 8, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 20 20. The composition of any of claims 9, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.
- 25 21. The composition of any of claims 10, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 µg-hour/dl to about 600 µg-hour/dl.

22. The composition of claim 12, wherein the composition features a levothyroxine (T4) AUC (0-t) of between from about 500 $\mu\text{g-hour/dL}$ to about 550 $\mu\text{g-hour/dL}$.
- 5 23. The composition of claim 12, wherein the $\ln[\text{AUC}(0-t)]$ is between from about 1 to about 10.
24. The composition of claim 12, wherein the composition features a triiodothyronine (T3) AUC (0-t) of between from about 10 ng-hour/mL to about
10 100 ng-hour/mL .
25. The composition of claim 12, wherein the composition features a triiodothyronine (T3) AUC (0-t) of between from about 20 ng-hour/mL to about
15 60 ng-hour/mL .
26. The composition of claim 12, wherein the $\ln[\text{AUC}(0-t)]$ is between from about 1 to about 5.
27. The composition of claim 1, wherein the composition is essentially sugar
20 free.
28. The composition of claim 1, wherein the composition is essentially non-granular.
- 25 29. The composition of claim 1, wherein the composition further comprises microcrystalline β -cellulose.

30. The composition of claim 29, wherein the microcrystalline β -cellulose has a bulk density of between from about 0.10 g/cm³ to about 0.35 g/cm³.
31. The composition of claim 29, wherein the microcrystalline β -cellulose has a bulk density of between from about 0.19 g/cm³ to about 0.21 g/cm³.
32. The composition of claim 29, wherein the microcrystalline β -cellulose has a conductivity of less than about 200 μ S/cm.
33. The composition of claim 29, wherein the microcrystalline β -cellulose has a conductivity of between from about 0.5 μ S/cm to 50 μ S/cm.
34. The composition of claims 1, wherein the composition has a post-packaging potency of between from about 90% to about 110% as determined by a standard potency test.
35. The composition of claims 12, wherein the composition has a post-packaging potency of between from about 90% to about 110% as determined by a standard potency test.
36. The composition of claims 29, wherein the composition has a post-packaging potency of between from about 90% to about 110% as determined by a standard potency test.
37. The composition of claim 30, wherein the composition has a post-packaging potency of between from about 90% to about 110% as determined by the standard potency test.

38. The composition of claim 1, wherein the composition is formulated as a tablet.
39. The composition of claim 38, wherein the tablet has a total hardness of
5 between from about 6 to about 14 KP, as determined by a standard hardness test.
40. The composition of claim 38, wherein the tablet is configured to increase heat transfer away from the tablet.
- 10 41. The composition of claim 40, wherein the tablet has a surface area of between from about 0.9 in.² to about 0.15 in.².
42. The composition of claim 38, wherein the tablet is beveled.
- 15 43. The composition of claim 42, wherein the tablet further comprises a notch.
44. The composition of claim 43, wherein the tablet has a raised violin configuration.
- 20 45. The composition of claim 1, wherein the composition contains less than about 5% total of diiodothyronine (T2), triiodothyronine (T3), triiodothyroacetic acid amide, triiodothyroethylamine, triiodothyroacetic acid, triiodothyroethyl alcohol, tetraiodothyroacetic acid amide, tetraiodothyroacetic acid, triiodothyroethane, and tetraiodothyroethane.
- 25 46. The composition of claim 1, wherein the composition further comprises a pharmaceutically acceptable croscarmellose salt.

47. The composition of claim 1, wherein the composition further comprises a pharmaceutically acceptable magnesium salt.

5

10

48. A pharmaceutical composition in tablet form comprising levothyroxine sodium, the composition comprising:

a) between from about 1 $\mu\text{g}/\text{tablet}$ to about 1000 $\mu\text{g}/\text{tablet}$ levothyroxine sodium (USP),

15

b) between from about 100 mg/tablet to about 110 mg/tablet of microcrystalline β -cellulose, NF (Ceolus) having a bulk density of between from about 0.10 g/cm^3 to about 0.35 g/cm^3 ,

c) between from about 25mg/tablet to about 50mg/tablet of crosscarmellose sodium, NF (Ac-di-sol); and

20

d) between from about 0.5 mg/tablet to about 5mg/tablet of magnesium stearate, NF.

25

5

10

49. A method of preparing a solid dosage form of a pharmaceutically active ingredient comprising forming a blend comprising the pharmaceutically active ingredient and β -sheet form of microcrystalline cellulose, and forming therefrom a solid dosage by compressing the blend in a tableting machine.

5

50. The method of claim 49, wherein the blend is compressed in a ratio of initial volume to final volume from 3.3:1 to 4.0:1.

51. The method of claim 50, wherein the pharmaceutically active ingredient
10 comprises levothyroxine sodium.

52. The method of claim 51, wherein the tableting machine further comprises extra deep tablet dies that maintain a free clearance of at least 3.0 mm during
filling.

15

20

25

30

5

53. An immediate release solid pharmaceutical composition comprising a levothyroxine salt and a pharmaceutically acceptable carrier.

10

54. The composition of claim 53, wherein at least about 85% of the levothyroxine dissolves in aqueous solution in less than about 20 minutes as determined by a standard dissolution test.

15

55. The composition of claim 53, wherein at least about 80% of the levothyroxine dissolves in aqueous solution by about 15 minutes as determined by the standard dissolution test.

20

56. The composition of claims 53, wherein at least about 80% of the levothyroxine dissolves in aqueous solution by about 5 minutes as determined by the standard dissolution test.

25

57. The composition of claims 53, wherein the composition further comprises microcrystalline β -cellulose.

58. The composition of claim 57, wherein the microcrystalline β -cellulose has a bulk density of between from about 0.10 g/cm^3 to about 0.35 g/cm^3 .

30

59. The composition of claim 57, wherein the microcrystalline β -cellulose has a conductivity of less than about $200 \text{ }\mu\text{S/cm}$.

60. The composition of claim 53, wherein the composition is formulated as a tablet.

5 61. The composition of claim 60, wherein tablet has a total hardness of between from about 6 to about 14 KP as determined by a standard hardness test.

62. The composition of claims 53, wherein the composition further comprises a pharmaceutically acceptable croscarmellose salt.

10

63. An immediate release pharmaceutical composition in tablet form comprising levothyroxine sodium, the composition comprising:

15 a) between from about 1 $\mu\text{g}/\text{tablet}$ to about 1000 $\mu\text{g}/\text{tablet}$ levothyroxine sodium (USP),

b) between from about 100 mg/tablet to about 110 mg/tablet of microcrystalline β -cellulose, NF (Ceolus) having a bulk density of between from about 0.10 g/cm^3 to about 0.35 g/cm^3 ,

20 c) between from about 25mg/tablet to about 50mg/tablet of croscarmellose sodium, NF (Ac-di-sol); and

d) between from about 0.5 mg/tablet to about 5mg/tablet of magnesium stearate, NF.

25

5

10

15

20

64. A method of producing a dispersible pharmaceutical composition comprising
- (a) mixing an active pharmaceutically active ingredient with microcrystalline -cellulose and a crosscarmellose salt to make a blend, and
- 5 (b) compressing the blend in a ratio of initial volume to final volume of between about 2:1 to about 5:1, to produce the pharmaceutical composition in solid form.
65. The method of claim 64, wherein the ratio of initial volume to final
- 10 volume is about 4:1.
66. The method of claim 64, wherein the composition features less than about 5% total impurities as determined by the standard impurity test.
- 15 67. The method of claim 64, wherein the method further comprises forming a tablet.
68. The method of claim 67, wherein the tablet has a raised violin configuration.
- 20 69. The method of claim 64, wherein the pharmaceutically active ingredient comprises levothyroxine sodium.
70. The method of claim 67, wherein the pharmaceutically active ingredient
- 25 comprises levothyroxine sodium.
71. The method of claim 68, wherein the pharmaceutically active ingredient comprises levothyroxine sodium.

72. The method of claims 64, wherein the pharmaceutically active ingredient comprises liothyronine sodium.

5

73. The method of claims 67, wherein the pharmaceutically active ingredient comprises liothyronine sodium.

10

74. The method of claims 68, wherein the pharmaceutically active ingredient comprises liothyronine sodium.

75. A stabilized solid pharmaceutical composition comprising a levothyroxine salt and a pharmaceutically acceptable carrier, wherein the pharmaceutical composition loses less than about 0.7% of its activity per month for up to 18 months.

5

76. The composition of claim 75, wherein the pharmaceutical composition loses less than about 0.5% of its activity per month for up to 18 months.

77. The composition of claim 75, wherein the pharmaceutical composition
10 loses less than about 0.3% of its activity per month for up to 18 months.

15

20

25

78. A pharmaceutical composition in solid form comprising a pharmaceutically active ingredient and a β - sheet form of microcrystalline cellulose.
- 5 79. The pharmaceutical composition of claim 78, wherein at least about 50 weight % of the composition weight is β - sheet form of microcrystalline cellulose.
- 10 80. The pharmaceutical composition of claims 78, wherein the active ingredient is levothyroxine sodium.
81. The pharmaceutical composition of claims 79, wherein the active ingredient is levothyroxine sodium.

82. An immediate release pharmaceutical composition comprising a levothyroxine salt, wherein the composition features a levothyroxine (T4) plasma AUC (0-t) of between from about 450 $\mu\text{g}\cdot\text{hour}/\text{dl}$ to about 600 $\mu\text{g}\cdot\text{hour}/\text{dl}$.

5 83. The composition of claim 82, wherein the composition features a levothyroxine (T4) AUC (0-t) of between from about 500 $\mu\text{g}\cdot\text{hour}/\text{dL}$ to about 550 $\mu\text{g}\cdot\text{hour}/\text{dL}$.

10 84. The composition of claims 82, wherein the $\ln[\text{AUC}(0-t)]$ is between from about 1 to about 10.

15 85. The composition of claims 83, wherein the $\ln[\text{AUC}(0-t)]$ is between from about 1 to about 10.

20

25

30

35

5

86. A non-granulated sugar-free, starch-free pharmaceutical composition comprising levothyroxine and a pharmaceutically acceptable carrier, in tablet form.
- 5 87. The composition of claim 86, wherein the composition further comprises microcrystalline β -cellulose.
88. The composition of claim 86, wherein the microcrystalline β -cellulose has a bulk density of between from about 0.10 g/cm^3 to about 0.35 g/cm^3 .
- 10 89. The composition of claims 86, wherein the microcrystalline β -cellulose has a bulk density of between from about 0.15 g/cm^3 to about 0.25 g/cm^3 .
90. The composition of claim 86, wherein the microcrystalline β -cellulose has
15 a conductivity of less than about $200 \text{ }\mu\text{S/cm}$.
91. The composition of claim 86, wherein the microcrystalline β -cellulose has a conductivity of between from about $0.5 \text{ }\mu\text{S/cm}$ to $50 \text{ }\mu\text{S/cm}$.
- 20 92. The composition of claim 86, wherein tablet has a total hardness of between from about 6 to about 14 KP as determined by a standard hardness test.

93. A method of stabilizing a pharmaceutically active compound which degrades in the presence of carbohydrates, comprising admixing the compound with a β -form microcellulose.
- 5 94. The method of claim 93, further comprising mixing the compound and the β -form microcellulose with a crosscarmellose salt to form a blend.
95. The method of claim 94, further comprising compressing the blend in a ratio of initial to final volume of between about 2:1 to about 5:1 to form the
- 10 stabilized composition.
96. The method of claim 95, wherein the ratio of initial volume to final volume is about 4:1.
- 15 97. The method of claim 95, further comprising forming the composition into a tablet.
98. The method of claim 97, wherein the tablet has a raised violin configuration.

20

1/4




FIG. 1A



FIG. 1B




FIG. 1C





FIG. 2

2/4






FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B