



- (51) **International Patent Classification:**  
C12N 15/113 (2010.01) A61K 31/712 (2006.01)
- (21) **International Application Number:**  
PCT/US2017/025461
- (22) **International Filing Date:**  
31 March 2017 (31.03.2017)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
62/317,140 1 April 2016 (01.04.2016) US
- (71) **Applicant:** AVIDITY BIOSCIENCES LLC [US/US];  
10975 North Torrey Pines Road, Suite 150, La Jolla, California 92037 (US).
- (72) **Inventors:** HUANG, Hanhua; 4826 Riding Ridge Road,  
San Diego, California 92130 (US). BURKE, Rob; 321  
Trailview Road, Encinitas, California 92024 (US).
- (74) **Agent:** YANG, Xiaofan; WILSON SONSINI  
GOODRICH & ROSATI, 650 Page Mill Road, Palo Alto,  
California 94304 (US).
- (81) **Designated States** (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN,  
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA,  
MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG,  
NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,  
RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY,  
TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,  
ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,  
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,  
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,  
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,  
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,  
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, KM, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

**Published:**

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))



WO 2017/173320 A2

(54) **Title:** MYC NUCLEIC ACIDS AND USES THEREOF

(57) **Abstract:** Disclosed herein are molecules and pharmaceutical compositions that mediate RNA interference against MYC. Also described herein include methods for treating a disease or disorder that comprises a molecule or a pharmaceutical composition that mediate RNA interference against MYC.

**MYC NUCLEIC ACIDS AND USES THEREOF****CROSS-REFERENCE**

[0001] This application claims the benefit of U.S. Provisional Application No. 62/317,140, filed April 1, 2016, which application is incorporated herein by reference.

**SEQUENCE LISTING**

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on March 28, 2017, is named 45532-713\_\_601\_\_SL.txt and is 4,146 bytes in size.

**BACKGROUND OF THE DISCLOSURE**

[0003] Gene suppression by RNA-induced gene silencing provides several levels of control: transcription inactivation, small interfering RNA (siRNA)-induced mRNA degradation, and siRNA-induced transcriptional attenuation. In some instances, RNA interference (RNAi) provides long lasting effect over multiple cell divisions. As such, RNAi represents a viable method useful for drug target validation, gene function analysis, pathway analysis, and disease therapeutics.

**SUMMARY OF THE DISCLOSURE**

[0004] Disclosed herein, in certain embodiments, are molecules and pharmaceutical compositions for modulating MYC function and/or expression in a cell.

[0005] Disclosed herein, in certain embodiments, is a polynucleic acid molecule that mediates RNA interference against MYC, wherein the polynucleic acid molecule comprises at least one 2' modified nucleotide, at least one modified internucleotide linkage, or at least one inverted abasic moiety.

[0006] In some embodiments, the at least one 2' modified nucleotide comprises 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified nucleotide. In some embodiments, the at least one 2' modified nucleotide comprises locked nucleic acid (LNA) or ethylene nucleic acid (ENA). In some embodiments, the at least one inverted basic moiety is at at least one terminus. In some embodiments, the at least one modified internucleotide linkage comprises a phosphorothioate linkage or a phosphorodithioate linkage.

**[0007]** In some embodiments, the polynucleic acid molecule is at least from about 10 to about 30 nucleotides in length. In some embodiments, the polynucleic acid molecule is at least one of: from about 15 to about 30, from about 18 to about 25, from about 18 to about 24, from about 19 to about 23, or from about 20 to about 22 nucleotides in length. In some embodiments, the polynucleic acid molecule is at least about 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides in length.

**[0008]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 5% to about 100% modification, from about 10% to about 100% modification, from about 20% to about 100% modification, from about 30% to about 100% modification, from about 40% to about 100% modification, from about 50% to about 100% modification, from about 60% to about 100% modification, from about 70% to about 100% modification, from about 80% to about 100% modification, and from about 90% to about 100% modification.

**[0009]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 90% modification, from about 20% to about 90% modification, from about 30% to about 90% modification, from about 40% to about 90% modification, from about 50% to about 90% modification, from about 60% to about 90% modification, from about 70% to about 90% modification, and from about 80% to about 100% modification.

**[0010]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 80% modification, from about 20% to about 80% modification, from about 30% to about 80% modification, from about 40% to about 80% modification, from about 50% to about 80% modification, from about 60% to about 80% modification, and from about 70% to about 80% modification.

**[0011]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 70% modification, from about 20% to about 70% modification, from about 30% to about 70% modification, from about 40% to about 70% modification, from about 50% to about 70% modification, and from about 60% to about 70% modification.

**[0012]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 60% modification, from about 20% to about 60% modification, from about 30% to about 60% modification, from about 40% to about 60% modification, and from about 50% to about 60% modification.

**[0013]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 50% modification, from about 20% to about 50% modification, from about 30% to about 50% modification, and from about 40% to about 50% modification.

**[0014]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 40% modification, from about 20% to about 40% modification, and from about 30% to about 40% modification.

**[0015]** In some embodiments, the polynucleic acid molecule comprises at least one of: from about 10% to about 30% modification, and from about 20% to about 30% modification.

**[0016]** In some embodiments, the polynucleic acid molecule comprises from about 10% to about 20% modification.

**[0017]** In some embodiments, the polynucleic acid molecule comprises from about 15% to about 90%, from about 20% to about 80%, from about 30% to about 70%, or from about 40% to about 60% modifications.

**[0018]** In some embodiments, the polynucleic acid molecule comprises at least about 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% modification.

**[0019]** In some embodiments, the polynucleic acid molecule comprises at least about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22 or more modifications.

**[0020]** In some embodiments, the polynucleic acid molecule comprises at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22 or more modified nucleotides.

**[0021]** In some embodiments, the polynucleic acid molecule comprises a single strand.

**[0022]** In some embodiments, the polynucleic acid molecule comprises two or more strands.

**[0023]** In some embodiments, the polynucleic acid molecule comprises a first polynucleotide and a second polynucleotide hybridized to the first polynucleotide to form a double-stranded polynucleic acid molecule.

**[0024]** In some embodiments, the second polynucleotide comprises at least one modification.

**[0025]** In some embodiments, the first polynucleotide and the second polynucleotide are RNA molecules. In some embodiments, the first polynucleotide and the second polynucleotide are siRNA molecules.

**[0026]** In some embodiments, the first polynucleotide comprises a sequence having at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the first polynucleotide consists of a sequence selected from SEQ ID NOs: 1-12. In some embodiments, the second polynucleotide comprises a sequence having at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100%

sequence identity to SEQ ID NOs: 1-12. In some embodiments, the second polynucleotide consists of a sequence selected from SEQ ID NOs: 1-12.

**[0027]** In some embodiments, MYC is *c-MYC* gene, *L-MYC* gene, or *N-MYC* gene. In some embodiments, MYC is *c-MYC* gene.

**[0028]** Disclosed herein, in certain embodiments, is a pharmaceutical composition comprising: a) a molecule disclosed above; and b) a pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition is formulated as a nanoparticle formulation. In some embodiments, the pharmaceutical composition is formulated for parenteral, oral, intranasal, buccal, rectal, or transdermal administration.

**[0029]** Disclosed herein, in certain embodiments, is a method of treating a disease or disorder in a patient in need thereof, comprising administering to the patient a composition comprising a molecule disclosed above. In some embodiments, the disease or disorder is a cancer. In some embodiments, the cancer is a solid tumor. In some embodiments, the cancer is a hematologic malignancy. In some embodiments, the cancer comprises a MYC-associated cancer. In some embodiments, the cancer comprises bladder cancer, breast cancer, colorectal cancer, endometrial cancer, esophageal cancer, glioblastoma multiforme, head and neck cancer, kidney cancer, lung cancer, ovarian cancer, pancreatic cancer, prostate cancer, or thyroid cancer. In some embodiments, the cancer comprises acute myeloid leukemia, CLL, DLBCL, or multiple myeloma.

**[0030]** Disclosed herein, in certain embodiments, is a method of inhibiting the expression of *MYC* gene in a primary cell of a patient, comprising administering a molecule disclosed above to the primary cell. In some embodiments, the method is an *in vivo* method. In some embodiments, the patient is a human.

**[0031]** Disclosed herein, in certain embodiments, is a kit comprising a molecule disclosed above.

#### **DETAILED DESCRIPTION OF THE DISCLOSURE**

**[0032]** The MYC family of proteins encompasses a group of basic-helix-loop-helix-leucine zipper transcription factors that are involved in cell cycle progression, apoptosis and cellular transformation. In mammals, there are three members of the MYC family--c-Myc, L-Myc, and N-Myc, in which each member functions in a similar manner but display differences in potency. In addition, each *MYC* gene is located at a different chromosome, *N-MYC* (2p24.1), *C-MYC* (8q24), and *L-MYC* (1p34.3). In some instances, mutations in the MYC proteins (e.g., c-Myc, N-Myc, and L-Myc) lead to constitutive expression of the MYC mutants, thereby leading to upregulation of numerous genes involved in cell proliferation and to cancer. For example, the c-Myc mutants are observed in a broad range of solid tumors and hematologic malignancies. In contrast, N-Myc

mutants are generally observed in solid tumors of neural origin (e.g., neuroblastoma and glioma), and L-Myc mutants are primarily observed in small cell lung carcinoma. In some cases, amplifications of wild type Myc genes are also observed in cancer.

**[0033]** In some embodiments, disclosed herein are polynucleic acid molecules and pharmaceutical compositions that modulate the expression of *MYC*. In some cases, the *MYC* gene is *c-MYC*, *L-MYC*, or *N-MYC*. In some instances, the polynucleic acid molecules and pharmaceutical compositions modulate the expression of *c-MYC*. In some instances, the polynucleic acid molecules and pharmaceutical compositions modulate the expression of *L-MYC*. In some instances, the polynucleic acid molecules and pharmaceutical compositions modulate the expression of *N-MYC*. In some instances, the polynucleic acid molecules and pharmaceutical compositions modulate the expression of wild type *MYC* gene (e.g., wild type *c-MYC*, *N-MYC*, or *L-MYC*). In other instances, the polynucleic acid molecules and pharmaceutical compositions modulate the expression of mutant *MYC* (e.g., mutant *c-MYC*, *N-MYC*, or *L-MYC*).

**[0034]** In some embodiments, the polynucleic acid molecules and pharmaceutical compositions are used for the treatment of a disease or disorder (e.g., cancer or a *MYC*-associated disease or disorder). In additional embodiments, the polynucleic acid molecules and pharmaceutical compositions are used for inhibiting the expression of *MYC* (e.g., *c-MYC*, *N-MYC*, or *L-MYC*) gene in a primary cell of a patient in need thereof.

**[0035]** In additional cases, also included herein are kits that comprise one or more of polynucleic acid molecules and pharmaceutical compositions described herein.

### **Polynucleic Acid Molecule**

**[0036]** In some embodiments, a polynucleic acid molecule described herein modulates the expression of the *MYC* gene. In some embodiments, the *MYC* gene is wild type *MYC* (e.g., *c-MYC*) or *MYC* (e.g., *c-MYC*) comprising one or more mutations. In some instances, *MYC* is wild type *MYC* DNA or RNA. In some instances, *MYC* DNA or RNA comprises one or more mutations. In some instances, the polynucleic acid molecule hybridizes to a target region of wild type *MYC* DNA or RNA. In some instances, the polynucleic acid molecule is a polynucleic acid molecule that hybridizes to a target region of *MYC* DNA or RNA comprising a mutation (e.g., a substitution, a deletion, or an addition).

**[0037]** In some embodiments, *MYC* DNA or RNA comprises one or more mutations. In some embodiments, *MYC* DNA or RNA comprises one or more mutation within one or more exons. In some instances, *MYC* DNA or RNA comprises one or more mutations within exon 2 or exon 3. In some instances, *MYC* DNA or RNA comprises one or more mutations at positions corresponding

to amino acid residues 2, 7, 17, 20, 32, 44, 58, 59, 76, 115, 138, 141, 145, 146, 169, 175, 188, 200, 202, 203, 248, 251, 298, 321, 340, 369, 373, 374, 389, 395, 404, 419, 431, 439, or a combination thereof. In some embodiments, MYC DNA or RNA comprises one or more mutations at positions corresponding to amino acid residues selected from P2L, F7L, D17N, Q20E, Y32N, A44V, A44T, T58I, P59L, A76V, F115L, F138S, A141S, V145I, S146L, S169C, S175N, C188F, N200S, S202N, S203T, T248S, D251E, S298Y, Q321E, V340D, V369D, T373K, H374R, F389L, Q395H, K404N, L419M, E431K, R439Q, or a combination thereof of the MYC polypeptide.

**[0038]** In some embodiments, a polynucleic acid molecule hybridizes to a target region of MYC DNA or RNA comprising one or more mutations. In some embodiments, the polynucleic acid molecule hybridizes to a target region of MYC DNA or RNA comprising one or more mutations within an exon. In some embodiments, the polynucleic acid molecule hybridizes to a target region of MYC DNA or RNA comprising one or more mutations within exon 2 or exon 3. In some embodiments, the polynucleic acid molecule hybridizes to a target region of MYC DNA or RNA comprising one or more mutations at positions corresponding to amino acid residues 2, 7, 17, 20, 32, 44, 58, 59, 76, 115, 138, 141, 145, 146, 169, 175, 188, 200, 202, 203, 248, 251, 298, 321, 340, 369, 373, 374, 389, 395, 404, 419, 431, 439, or a combination thereof of the MYC polypeptide. In some embodiments, the polynucleic acid molecule hybridizes to a target region of MYC DNA or RNA comprising one or more mutations at positions corresponding to amino acid residues selected from P2L, F7L, D17N, Q20E, Y32N, A44V, A44T, T58I, P59L, A76V, F115L, F138S, A141S, V145I, S146L, S169C, S175N, C188F, N200S, S202N, S203T, T248S, D251E, S298Y, Q321E, V340D, V369D, T373K, H374R, F389L, Q395H, K404N, L419M, E431K, R439Q, or a combination thereof of the MYC polypeptide.

**[0039]** In some embodiments, a polynucleic acid molecule comprises a sequence having at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to a sequence listed in Table 1. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 50% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 60% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 70% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 75% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 80% sequence identity to SEQ ID NOs: 1-12. In some

embodiments, the polynucleic acid molecule comprises a sequence having at least 85% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 90% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 95% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 96% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 97% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 98% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule comprises a sequence having at least 99% sequence identity to SEQ ID NOs: 1-12. In some embodiments, the polynucleic acid molecule consists of SEQ ID NOs: 1-12.

**[0040]** In some embodiments, a polynucleic acid molecule comprises a first polynucleotide and a second polynucleotide. In some instances, the first polynucleotide comprises a sequence having at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs: 1-12. In some cases, the second polynucleotide comprises a sequence having at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs: 1-12. In some cases, the polynucleic acid molecule comprises a first polynucleotide having at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs: 1-12 and a second polynucleotide having at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to SEQ ID NOs: 1-12.

**[0041]** In some embodiments, a polynucleic acid molecule described herein comprises RNA or DNA. In some cases, the polynucleic acid molecule comprises RNA. In some instances, RNA comprises short interfering RNA (siRNA), short hairpin RNA (shRNA), microRNA (miRNA), double-stranded RNA (dsRNA), transfer RNA (tRNA), ribosomal RNA (rRNA), or heterogeneous nuclear RNA (hnRNA). In some instances, RNA comprises shRNA. In some instances, RNA comprises miRNA. In some instances, RNA comprises dsRNA. In some instances, RNA comprises tRNA. In some instances, RNA comprises rRNA. In some instances, RNA comprises hnRNA. In some instances, the RNA comprises siRNA. In some instances, the polynucleic acid molecule comprises siRNA.

**[0042]** In some embodiments, a polynucleic acid molecule is from about 10 to about 50 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 30, from about 15 to about 30, from about 18 to about 25, from about 18 to about 24, from about 19 to about 23, or from about 20 to about 22 nucleotides in length.

**[0043]** In some embodiments, a polynucleic acid molecule is about 50 nucleotides in length. In some instances, the polynucleic acid molecule is about 45 nucleotides in length. In some instances, the polynucleic acid molecule is about 40 nucleotides in length. In some instances, the polynucleic acid molecule is about 35 nucleotides in length. In some instances, the polynucleic acid molecule is about 30 nucleotides in length. In some instances, the polynucleic acid molecule is about 25 nucleotides in length. In some instances, the polynucleic acid molecule is about 20 nucleotides in length. In some instances, the polynucleic acid molecule is about 19 nucleotides in length. In some instances, the polynucleic acid molecule is about 18 nucleotides in length. In some instances, the polynucleic acid molecule is about 17 nucleotides in length. In some instances, the polynucleic acid molecule is about 16 nucleotides in length. In some instances, the polynucleic acid molecule is about 15 nucleotides in length. In some instances, the polynucleic acid molecule is about 14 nucleotides in length. In some instances, the polynucleic acid molecule is about 13 nucleotides in length. In some instances, the polynucleic acid molecule is about 12 nucleotides in length. In some instances, the polynucleic acid molecule is about 11 nucleotides in length. In some instances, the polynucleic acid molecule is about 10 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 50 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 45 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 40 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 35 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 30 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 25 nucleotides in length. In some instances, the polynucleic acid molecule is from about 10 to about 20 nucleotides in length. In some instances, the polynucleic acid molecule is from about 15 to about 25 nucleotides in length. In some instances, the polynucleic acid molecule is from about 15 to about 30 nucleotides in length. In some instances, the polynucleic acid molecule is from about 12 to about 30 nucleotides in length.

**[0044]** In some embodiments, a polynucleic acid molecule comprises a first polynucleotide. In some instances, the polynucleic acid molecule comprises a second polynucleotide. In some instances, the polynucleic acid molecule comprises a first polynucleotide and a second polynucleotide. In some instances, the first polynucleotide is a sense strand or passenger strand. In some instances, the second polynucleotide is an antisense strand or guide strand.

**[0045]** In some embodiments, a polynucleic acid molecule is a first polynucleotide. In some embodiments, the first polynucleotide is from about 10 to about 50 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 30, from about 15 to about 30, from

about 18 to about 25, from about 18 to about 24, from about 19 to about 23, or from about 20 to about 22 nucleotides in length.

**[0046]** In some instances, a first polynucleotide is about 50 nucleotides in length. In some instances, the first polynucleotide is about 45 nucleotides in length. In some instances, the first polynucleotide is about 40 nucleotides in length. In some instances, the first polynucleotide is about 35 nucleotides in length. In some instances, the first polynucleotide is about 30 nucleotides in length. In some instances, the first polynucleotide is about 25 nucleotides in length. In some instances, the first polynucleotide is about 20 nucleotides in length. In some instances, the first polynucleotide is about 19 nucleotides in length. In some instances, the first polynucleotide is about 18 nucleotides in length. In some instances, the first polynucleotide is about 17 nucleotides in length. In some instances, the first polynucleotide is about 16 nucleotides in length. In some instances, the first polynucleotide is about 15 nucleotides in length. In some instances, the first polynucleotide is about 14 nucleotides in length. In some instances, the first polynucleotide is about 13 nucleotides in length. In some instances, the first polynucleotide is about 12 nucleotides in length. In some instances, the first polynucleotide is about 11 nucleotides in length. In some instances, the first polynucleotide is about 10 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 50 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 45 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 40 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 35 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 30 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 25 nucleotides in length. In some instances, the first polynucleotide is from about 10 to about 20 nucleotides in length. In some instances, the first polynucleotide is from about 15 to about 25 nucleotides in length. In some instances, the first polynucleotide is from about 15 to about 30 nucleotides in length. In some instances, the first polynucleotide is from about 12 to about 30 nucleotides in length.

**[0047]** In some embodiments, a polynucleic acid molecule is a second polynucleotide. In some embodiments, the second polynucleotide is from about 10 to about 50 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 30, from about 15 to about 30, from about 18 to about 25, from about 18 to about 24, from about 19 to about 23, or from about 20 to about 22 nucleotides in length.

**[0048]** In some instances, a second polynucleotide is about 50 nucleotides in length. In some instances, the second polynucleotide is about 45 nucleotides in length. In some instances, the second polynucleotide is about 40 nucleotides in length. In some instances, the second

polynucleotide is about 35 nucleotides in length. In some instances, the second polynucleotide is about 30 nucleotides in length. In some instances, the second polynucleotide is about 25 nucleotides in length. In some instances, the second polynucleotide is about 20 nucleotides in length. In some instances, the second polynucleotide is about 19 nucleotides in length. In some instances, the second polynucleotide is about 18 nucleotides in length. In some instances, the second polynucleotide is about 17 nucleotides in length. In some instances, the second polynucleotide is about 16 nucleotides in length. In some instances, the second polynucleotide is about 15 nucleotides in length. In some instances, the second polynucleotide is about 14 nucleotides in length. In some instances, the second polynucleotide is about 13 nucleotides in length. In some instances, the second polynucleotide is about 12 nucleotides in length. In some instances, the second polynucleotide is about 11 nucleotides in length. In some instances, the second polynucleotide is about 10 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 50 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 45 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 40 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 35 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 30 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 25 nucleotides in length. In some instances, the second polynucleotide is from about 10 to about 20 nucleotides in length. In some instances, the second polynucleotide is from about 15 to about 25 nucleotides in length. In some instances, the second polynucleotide is from about 15 to about 30 nucleotides in length. In some instances, the second polynucleotide is from about 12 to about 30 nucleotides in length.

**[0049]** In some embodiments, a polynucleic acid molecule comprises a first polynucleotide and a second polynucleotide. In some instances, the polynucleic acid molecule further comprises a blunt terminus, an overhang, or a combination thereof. In some instances, the blunt terminus is a 5' blunt terminus, a 3' blunt terminus, or both. In some cases, the overhang is a 5' overhang, 3' overhang, or both. In some cases, the overhang comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 non-base pairing nucleotides. In some cases, the overhang comprises 1, 2, 3, 4, 5, or 6 non-base pairing nucleotides. In some cases, the overhang comprises 1, 2, 3, or 4 non-base pairing nucleotides. In some cases, the overhang comprises 1 non-base pairing nucleotide. In some cases, the overhang comprises 2 non-base pairing nucleotides. In some cases, the overhang comprises 3 non-base pairing nucleotides. In some cases, the overhang comprises 4 non-base pairing nucleotides.

**[0050]** In some embodiments, the sequence of a polynucleic acid molecule is at least 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99%, or 99.5% complementary to a

target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule is at least 50% complementary to a target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule is at least 60% complementary to a target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule is at least 70% complementary to a target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule is at least 80% complementary to a target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule is at least 90% complementary to a target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule is at least 95% complementary to a target sequence described herein. In some instances, the sequence of the polynucleic acid molecule is 100% complementary to a target sequence described herein.

**[0051]** In some embodiments, the sequence of a polynucleic acid molecule has 5 or less mismatches to a target sequence described herein. In some embodiments, the sequence of the polynucleic acid molecule has 4 or less mismatches to a target sequence described herein. In some instances, the sequence of the polynucleic acid molecule has 3 or less mismatches to a target sequence described herein. In some cases, the sequence of the polynucleic acid molecule has 2 or less mismatches to a target sequence described herein. In some cases, the sequence of the polynucleic acid molecule has 1 or less mismatches to a target sequence described herein.

**[0052]** In some embodiments, the specificity of a polynucleic acid molecule that hybridizes to a target sequence described herein is a 95%, 98%, 99%, 99.5% or 100% sequence complementarity of the polynucleic acid molecule to a target sequence. In some instances, the hybridization is a high stringent hybridization condition.


**[0053]** In some embodiments, a polynucleic acid molecule has reduced off-target effect. In some instances, “off-target” or “off-target effects” refer to any instance in which a polynucleic acid polymer directed against a given target causes an unintended effect by interacting either directly or indirectly with another mRNA sequence, a DNA sequence or a cellular protein or other moiety. In some instances, an “off-target effect” occurs when there is a simultaneous degradation of other transcripts due to partial homology or complementarity between that other transcript and the sense and/or antisense strand of the polynucleic acid molecule.

**[0054]** In some embodiments, a polynucleic acid molecule comprises natural, synthetic, or artificial nucleotide analogues or bases. In some cases, the polynucleic acid molecule comprises combinations of DNA, RNA and/or nucleotide analogues. In some instances, the synthetic or


artificial nucleotide analogues or bases comprise modifications at one or more of ribose moiety, phosphate moiety, nucleoside moiety, or a combination thereof.

**[0055]** In some embodiments, nucleotide analogues or artificial nucleotide base comprise a nucleic acid with a modification at a 2' hydroxyl group of the ribose moiety. In some instances, the modification includes an H, OR, R, halo, SH, SR, NH<sub>2</sub>, NHR, NR<sub>2</sub>, or CN, wherein R is an alkyl moiety. Exemplary alkyl moiety includes, but is not limited to, halogens, sulfurs, thiols, thioethers, thioesters, amines (primary, secondary, or tertiary), amides, ethers, esters, alcohols and oxygen. In some instances, the alkyl moiety further comprises a modification. In some instances, the modification comprises an azo group, a keto group, an aldehyde group, a carboxyl group, a nitro group, a nitroso group, a nitrile group, a heterocycle (e.g., imidazole, hydrazino or hydroxylamino) group, an isocyanate or cyanate group, or a sulfur containing group (e.g., sulfoxide, sulfone, sulfide, or disulfide). In some instances, the alkyl moiety further comprises a hetero substitution. In some instances, the carbon of the heterocyclic group is substituted by a nitrogen, oxygen, or sulfur. In some instances, the heterocyclic substitution includes but is not limited to, morpholino, imidazole, and pyrrolidino.

**[0056]** In some instances, the modification at the 2' hydroxyl group is a 2'-O-methyl modification or a 2'-O-methoxyethyl (2'-O-MOE) modification. In some cases, the 2'-O-methyl modification adds a methyl group to the 2' hydroxyl group of the ribose moiety whereas the 2'-O-methoxyethyl modification adds a methoxyethyl group to the 2' hydroxyl group of the ribose moiety. Exemplary chemical structures of a 2'-O-methyl modification of an adenosine molecule and 2'-O-methoxyethyl modification of a uridine are illustrated below.




2'-O-methyl-adenosine




2'-O-methoxyethyl uridine

**[0057]** In some instances, the modification at the 2' hydroxyl group is a 2'-O-aminopropyl modification in which an extended amine group comprising a propyl linker binds the amine group to the 2' oxygen. In some instances, this modification neutralizes the phosphate derived overall negative charge of the oligonucleotide molecule by introducing one positive charge from the amine group per sugar and thereby improves cellular uptake properties due to its zwitterionic properties. An exemplary chemical structure of a 2'-O-aminopropyl nucleoside phosphoramidite is illustrated below.




2'-O-aminopropyl nucleoside phosphoramidite

**[0058]** In some instances, the modification at the 2' hydroxyl group is a locked or bridged ribose modification (e.g., locked nucleic acid or LNA) in which the oxygen molecule bound at the 2' carbon is linked to the 4' carbon by a methylene group, thus forming a 2'-C,4'-C-oxy-methylene-linked bicyclic ribonucleotide monomer. Exemplary representations of the chemical structure of LNA are illustrated below. The representation shown to the left highlights the chemical connectivities of an LNA monomer. The representation shown to the right highlights the locked 3'-endo (<sup>3</sup>E) conformation of the furanose ring of an LNA monomer.



LNA (Locked Nucleic Acids)

**[0059]** In some instances, the modification at the 2' hydroxyl group comprises ethylene nucleic acids (ENA) such as for example 2'-4'-ethylene-bridged nucleic acid, which locks the sugar conformation into a C<sub>3'</sub>-endo sugar puckering conformation. ENA are part of the bridged nucleic acids class of modified nucleic acids that also comprises LNA. Exemplary chemical structures of the ENA and bridged nucleic acids are illustrated below.




**[0060]** In some embodiments, additional modifications at the 2' hydroxyl group include 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O- dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA).

**[0061]** In some embodiments, nucleotide analogues comprise modified bases such as, but not limited to, 5-propynyluridine, 5-propynylcytidine, 6- methyladenine, 6-methylguanine, N, N, - dimethyladenine, 2-propyladenine, 2-propylguanine, 2-aminoadenine, 1-methylinosine, 3-methyluridine, 5-methylcytidine, 5-methyluridine and other nucleotides having a modification at the 5 position, 5- (2- amino) propyl uridine, 5-halocytidine, 5-halouridine, 4-acetylcytidine, 1-methyladenosine, 2-methyladenosine, 3-methylcytidine, 6-methyluridine, 2- methylguanosine, 7-methylguanosine, 2, 2-dimethylguanosine, 5- methylaminoethyluridine, 5-methoxyuridine, deazanucleotides (such as 7-deaza- adenosine, 6-azouridine, 6-azocytidine, or 6-azothymidine), 5-methyl-2-thiouridine, other thio bases (such as 2-thiouridine and 4-thiouridine and 2-thiocytidine), dihydrouridine, pseudouridine, queuosine, archaeosine, naphthyl and substituted naphthyl groups, any O-and N-alkylated purines and pyrimidines (such as N6-methyladenosine, 5-methylcarbonylmethyluridine, uridine 5-oxyacetic acid, pyridine-4-one, or pyridine-2-one), phenyl and modified phenyl groups such as aminophenol or 2,4, 6-trimethoxy benzene, modified cytosines that act as G-clamp nucleotides, 8-substituted adenines and guanines, 5-substituted uracils and thymines, azapyrimidines, carboxyhydroxyalkyl nucleotides, carboxyalkylaminoalkyl nucleotides, and alkylcarbonylalkylated nucleotides. Modified nucleotides also include those nucleotides that are modified with respect to the sugar moiety, as well as nucleotides having sugars or analogs thereof that are not ribosyl. For example, the sugar moieties, in some cases are, or are based on, mannoses, arabinoses, glucopyranoses, galactopyranoses, 4'-thioribose, and other sugars, heterocycles, or carbocycles. The term nucleotide also includes what are known in the art as universal bases. By way of example, universal bases include but are not limited to 3-nitropyrrole, 5-nitroindole, or nebularine.


**[0062]** In some embodiments, nucleotide analogues further comprise morpholinos, peptide nucleic acids (PNAs), methylphosphonate nucleotides, thiolphosphonate nucleotides, 2'-fluoro N3-

P5'-phosphoramidites, 1', 5'- anhydrohexitol nucleic acids (HNAs), or a combination thereof. Morpholino or phosphordiamidate morpholino oligo (PMO) comprises synthetic molecules whose structure mimics natural nucleic acid structure but deviates from the normal sugar and phosphate structures. In some instances, the five member ribose ring is substituted with a six member morpholino ring containing four carbons, one nitrogen and one oxygen. In some cases, the ribose monomers are linked by a phosphordiamidate group instead of a phosphate group. In such cases, the backbone alterations remove all positive and negative charges making morpholinos neutral molecules capable of crossing cellular membranes without the aid of cellular delivery agents such as those used by charged oligonucleotides.



Morpholino

**[0063]** In some embodiments, peptide nucleic acid (PNA) does not contain sugar ring or phosphate linkage and the bases are attached and appropriately spaced by oligoglycine-like molecules, therefore, eliminating a backbone charge.




PNA


**[0064]** In some embodiments, one or more modifications optionally occur at the internucleotide linkage. In some instances, modified internucleotide linkage includes, but is not limited to, phosphorothioates; phosphorodithioates; methylphosphonates; 5'-alkylenephosphonates; 5'-methylphosphonate; 3'-alkylene phosphonates; borontrifluoridates; borano phosphate esters and selenophosphates of 3'-5'linkage or 2'-5'linkage; phosphotriesters; thionoalkylphosphotriesters; hydrogen phosphonate linkages; alkyl phosphonates; alkylphosphonothioates; arylphosphonothioates; phosphoroselenoates; phosphorodiselenoates; phosphinates;

phosphoramidates; 3'-alkylphosphoramidates; aminoalkylphosphoramidates; thionophosphoramidates; phosphoropiperazidates; phosphoroanilothioates; phosphoroanilidates; ketones; sulfones; sulfonamides; carbonates; carbamates; methylenehydrazos; methylenedimethylhydrazos; formacetals; thioformacetals; oximes; methyleneiminos; methylenemethyliminos; thioamidates; linkages with riboacetyl groups; aminoethyl glycine; silyl or siloxane linkages; alkyl or cycloalkyl linkages with or without heteroatoms of, for example, 1 to 10 carbons that are saturated or unsaturated and/or substituted and/or contain heteroatoms; linkages with morpholino structures, amides, or polyamides wherein the bases are attached to the aza nitrogens of the backbone directly or indirectly; and combinations thereof.


**[0065]** In some instances, the modification is a methyl or thiol modification such as methylphosphonate or thiolphosphonate modification. Exemplary thiolphosphonate nucleotide (left) and methylphosphonate nucleotide (right) are illustrated below.



**[0066]** In some instances, a modified nucleotide includes, but is not limited to, 2'-fluoro N3-P5'-phosphoramidites illustrated as:



**[0067]** In some instances, a modified nucleotide includes, but is not limited to, hexitol nucleic acid (or 1', 5'-anhydrohexitol nucleic acids (HNA)) illustrated as:



HNA

**[0068]** In some embodiments, one or more modifications further optionally include modifications of the ribose moiety, phosphate backbone and the nucleoside, or modifications of the nucleotide analogues at the 3' or the 5' terminus. For example, the 3' terminus optionally includes a 3' cationic group, or by inverting the nucleoside at the 3'-terminus with a 3'-3' linkage. In another alternative, the 3'-terminus is optionally conjugated with an aminoalkyl group, e.g., a 3' C5-aminoalkyl dT. In an additional alternative, the 3'-terminus is optionally conjugated with an abasic site, e.g., with an apurinic or apyrimidinic site. In some instances, the 5'-terminus is conjugated with an aminoalkyl group, e.g., a 5'-O-alkylamino substituent. In some cases, the 5'-terminus is conjugated with an abasic site, e.g., with an apurinic or apyrimidinic site.

**[0069]** In some embodiments, a polynucleic acid molecule comprises one or more artificial nucleotide analogues described herein. In some instances, the polynucleic acid molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more artificial nucleotide analogues described herein. In some embodiments, the artificial nucleotide analogues include 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiophosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or a combination thereof. In some instances, the polynucleic acid molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of the artificial nucleotide analogues selected from 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiophosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or a combination thereof. In some instances, the polynucleic acid molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of 2'-O-methyl modified nucleotides. In some instances, the polynucleic acid molecule comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of 2'-O-methoxyethyl (2'-O-MOE) modified nucleotides. In some instances, the polynucleic acid molecule

comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 25, or more of thiolphosphonate nucleotides.

**[0070]** In some instances, a polynucleic acid molecule comprises at least one of: from about 5% to about 100% modification, from about 10% to about 100% modification, from about 20% to about 100% modification, from about 30% to about 100% modification, from about 40% to about 100% modification, from about 50% to about 100% modification, from about 60% to about 100% modification, from about 70% to about 100% modification, from about 80% to about 100% modification, and from about 90% to about 100% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0071]** In some cases, a polynucleic acid molecule comprises at least one of: from about 10% to about 90% modification, from about 20% to about 90% modification, from about 30% to about 90% modification, from about 40% to about 90% modification, from about 50% to about 90% modification, from about 60% to about 90% modification, from about 70% to about 90% modification, and from about 80% to about 100% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0072]** In some cases, a polynucleic acid molecule comprises at least one of: from about 10% to about 80% modification, from about 20% to about 80% modification, from about 30% to about 80% modification, from about 40% to about 80% modification, from about 50% to about 80% modification, from about 60% to about 80% modification, and from about 70% to about 80% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0073]** In some instances, a polynucleic acid molecule comprises at least one of: from about 10% to about 70% modification, from about 20% to about 70% modification, from about 30% to about 70% modification, from about 40% to about 70% modification, from about 50% to about 70% modification, and from about 60% to about 70% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0074]** In some instances, a polynucleic acid molecule comprises at least one of: from about 10% to about 60% modification, from about 20% to about 60% modification, from about 30% to about 60% modification, from about 40% to about 60% modification, and from about 50% to about 60% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0075]** In some cases, a polynucleic acid molecule comprises at least one of: from about 10% to about 50% modification, from about 20% to about 50% modification, from about 30% to about

50% modification, and from about 40% to about 50% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0076]** In some cases, a polynucleic acid molecule comprises at least one of: from about 10% to about 40% modification, from about 20% to about 40% modification, and from about 30% to about 40% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0077]** In some cases, a polynucleic acid molecule comprises at least one of: from about 10% to about 30% modification, and from about 20% to about 30% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0078]** In some cases, a polynucleic acid molecule comprises from about 10% to about 20% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0079]** In some cases, a polynucleic acid molecule comprises from about 15% to about 90%, from about 20% to about 80%, from about 30% to about 70%, or from about 40% to about 60% modifications. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0080]** In additional cases, a polynucleic acid molecule comprises at least about 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% modification. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0081]** In some embodiments, a polynucleic acid molecule comprises at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, or more modifications. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0082]** In some instances, a polynucleic acid molecule comprises at least about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22 or more modified nucleotides. In some instances, the polynucleic acid molecule is a polynucleic acid molecule of SEQ ID NOs: 1-12.

**[0083]** In some instances, about 5 to about 100% of a polynucleic acid molecule comprise an artificial nucleotide analogue described herein. In some instances, about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% of a polynucleic acid molecule comprise an artificial nucleotide analogue described herein. In some instances, about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%,



12 comprise an artificial nucleotide analogue described herein. In some instances, about 99% of a polynucleic acid molecule of SEQ ID NOs: 1-12 comprise an artificial nucleotide analogue described herein. In some instances, about 100% of a polynucleic acid molecule of SEQ ID NOs: 1-12 comprise an artificial nucleotide analogue described herein. In some embodiments, the artificial nucleotide analogue comprises 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiolphosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or a combination thereof.

**[0084]** In some embodiments, a polynucleic acid molecule comprises from about 1 to about 25 modifications in which the modification comprises an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises from about 1 to about 25 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 1 modification in which the modification comprises an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 2 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 3 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 4 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 5 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 6 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 7 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 8 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 9 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 10 modifications in which the modifications comprise an artificial nucleotide

analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 11 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 12 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 13 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 14 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 15 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 16 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 17 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 18 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 19 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 20 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 21 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 22 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 23 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 24 modifications in which the modifications comprise an artificial nucleotide analogue described herein. In some embodiments, a polynucleic acid molecule of SEQ ID NOs: 1-12 comprises about 25 modifications in which the modifications comprise an artificial nucleotide analogue described herein.

**[0085]** In some instances, a polynucleic acid molecule that comprises an artificial nucleotide analogue comprises SEQ ID NOs: 1-12.

**[0086]** In some embodiments, a polynucleic acid molecule is assembled from two separate polynucleotides wherein one polynucleotide comprises the sense strand and the second polynucleotide comprises the antisense strand of the polynucleic acid molecule. In other embodiments, the sense strand is connected to the antisense strand via a linker molecule, which in some instances is a polynucleotide linker or a non-nucleotide linker.

**[0087]** In some embodiments, a polynucleic acid molecule comprises a sense strand and antisense strand, wherein pyrimidine nucleotides in the sense strand comprise 2'-O-methylpyrimidine nucleotides and purine nucleotides in the sense strand comprise 2'-deoxy purine nucleotides. In some embodiments, a polynucleic acid molecule comprises a sense strand and antisense strand, wherein pyrimidine nucleotides present in the sense strand comprise 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein purine nucleotides present in the sense strand comprise 2'-deoxy purine nucleotides.

**[0088]** In some embodiments, a polynucleic acid molecule comprises a sense strand and antisense strand, wherein the pyrimidine nucleotides when present in said antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and the purine nucleotides when present in said antisense strand are 2'-O-methyl purine nucleotides.

**[0089]** In some embodiments, a polynucleic acid molecule comprises a sense strand and antisense strand, wherein the pyrimidine nucleotides when present in said antisense strand are 2'-deoxy-2'-fluoro pyrimidine nucleotides and wherein the purine nucleotides when present in said antisense strand comprise 2'-deoxy-purine nucleotides.

**[0090]** In some embodiments, a polynucleic acid molecule comprises a sense strand and antisense strand, wherein the sense strand includes a terminal cap moiety at the 5'-end, the 3'-end, or both of the 5' and 3' ends of the sense strand. In other embodiments, the terminal cap moiety is an inverted deoxy abasic moiety.

**[0091]** In some embodiments, a polynucleic acid molecule comprises a sense strand and an antisense strand, wherein the antisense strand comprises a phosphate backbone modification at the 3' end of the antisense strand. In some instances, the phosphate backbone modification is a phosphorothioate.

**[0092]** In some embodiments, a polynucleic acid molecule comprises a sense strand and an antisense strand, wherein the antisense strand comprises a glyceryl modification at the 3' end of the antisense strand.

**[0093]** In some embodiments, a polynucleic acid molecule comprises a sense strand and an antisense strand, in which the sense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide

linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or about one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and in which the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

**[0094]** In some embodiments, a polynucleic acid molecule comprises a sense strand and an antisense strand, in which the sense strand comprises about 1 to about 25, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and in which the antisense strand comprises about 1 to about 25 or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 25 or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

**[0095]** In some embodiments, a polynucleic acid molecule comprises a sense strand and an antisense strand, in which the antisense strand comprises one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages, and/or about one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 10 or more, specifically about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without one or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3' and 5'-ends, being present in the same or different strand.

**[0096]** In some embodiments, a polynucleic acid molecule comprises a sense strand and an antisense strand, in which the antisense strand comprises about 1 to about 25 or more, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand comprises about 1 to about 25 or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'-O-methyl, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. In other embodiments, one or more, for example about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more pyrimidine nucleotides of the sense and/or antisense strand are chemically-modified with 2'-deoxy, 2'-O-methyl and/or 2'-deoxy-2'-fluoro nucleotides, with or without about 1 to about 5, for example about 1, 2, 3, 4, 5 or more phosphorothioate internucleotide linkages and/or a terminal cap

molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

**[0097]** In some embodiments, a polynucleic acid molecule described herein is a chemically-modified short interfering nucleic acid molecule having about 1 to about 25, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more phosphorothioate internucleotide linkages in each strand of the polynucleic acid molecule.

**[0098]** In another embodiment, a polynucleic acid molecule described herein comprises 2'-5' internucleotide linkages. In some instances, the 2'-5' internucleotide linkage(s) is at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of one or both sequence strands. In additional instances, the 2'-5' internucleotide linkage(s) is present at various other positions within one or both sequence strands, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a pyrimidine nucleotide in one or both strands of the polynucleic acid molecule comprise a 2'-5' internucleotide linkage, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a purine nucleotide in one or both strands of the polynucleic acid molecule comprise a 2'-5' internucleotide linkage.

**[0099]** In some embodiments, a polynucleic acid molecule is a single stranded polynucleic acid molecule that mediates RNAi activity in a cell or reconstituted in vitro system, wherein the polynucleic acid molecule comprises a single stranded polynucleotide having complementarity to a target nucleic acid sequence, and wherein one or more pyrimidine nucleotides present in the polynucleic acid are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein one or more purine nucleotides present in the polynucleic acid are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and a terminal cap modification, that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the antisense sequence, the polynucleic acid molecule optionally further comprising about 1 to about 4 (e.g., about 1, 2, 3, or 4) terminal 2'-deoxynucleotides at the 3'-end of the polynucleic acid molecule, wherein the terminal nucleotides further comprise one or more (e.g., 1, 2, 3, or 4) phosphorothioate internucleotide linkages, and wherein the polynucleic acid molecule optionally further comprises a terminal phosphate group, such as a 5'-terminal phosphate group.

**[0100]** In some cases, one or more artificial nucleotide analogues described herein are resistant toward nucleases such as for example ribonuclease such as RNase H, deoxyribonuclease such as DNase, or exonuclease such as 5'-3' exonuclease and 3'-5' exonuclease when compared to natural

polynucleic acid molecules. In some instances, artificial nucleotide analogues comprising 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiophosphonate nucleotides, 2'-fluoro N3-P5'-phosphoramidites, or combinations thereof are resistant toward nucleases such as for example ribonuclease such as RNase H, deoxyribonuclease such as DNase, or exonuclease such as 5'-3' exonuclease and 3'-5' exonuclease. In some instances, 2'-O-methyl modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-O-methoxyethyl (2'-O-MOE) modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-O-aminopropyl modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-deoxy modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, T-deoxy-2'-fluoro modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-O-aminopropyl (2'-O-AP) modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-O-dimethylaminoethyl (2'-O-DMAOE) modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-O-dimethylaminopropyl (2'-O-DMAP) modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE) modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, 2'-O-N-methylacetamido (2'-O-NMA) modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, LNA modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, ENA modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, HNA modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, morpholinos are nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, PNA modified polynucleic acid molecule is resistant to nucleases (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-

5' exonuclease resistant). In some instances, methylphosphonate nucleotides modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, thiolphosphonate nucleotides modified polynucleic acid molecule is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, polynucleic acid molecule comprising 2'-fluoro N3-P5'-phosphoramidites is nuclease resistant (e.g., RNase H, DNase, 5'-3' exonuclease or 3'-5' exonuclease resistant). In some instances, the 5' conjugates described herein inhibit 5'-3' exonucleolytic cleavage. In some instances, the 3' conjugates described herein inhibit 3'-5' exonucleolytic cleavage.

**[0101]** In some embodiments, one or more artificial nucleotide analogues described herein have increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. The one or more of the artificial nucleotide analogues comprising 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified, LNA, ENA, PNA, HNA, morpholino, methylphosphonate nucleotides, thiolphosphonate nucleotides, or 2'-fluoro N3-P5'-phosphoramidites have increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-methyl modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-methoxyethyl (2'-O-MOE)-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-aminopropyl modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-deoxy modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, T-deoxy-2'-fluoro modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-aminopropyl (2'-O-AP)-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-dimethylaminoethyl (2'-O-DMAOE)-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-dimethylaminopropyl (2'-O-DMAP)-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, T-O-dimethylaminoethoxyethyl (2'-O-

DMAEOE)-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, 2'-O-N-methylacetamido (2'-O-NMA)-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, LNA-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, ENA-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, PNA-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, HNA-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, morpholino-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, methylphosphonate nucleotide-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, thiophosphonate nucleotide-modified polynucleic acid molecule has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some instances, polynucleic acid molecule comprising 2'-fluoro N3-P5'-phosphoramidites has increased binding affinity toward their mRNA target relative to an equivalent natural polynucleic acid molecule. In some cases, the increased affinity is illustrated with a lower K<sub>d</sub>, a higher melt temperature (T<sub>m</sub>), or a combination thereof.

**[0102]** In some embodiments, a polynucleic acid molecule described herein is a chirally pure (or stereo pure) polynucleic acid molecule, or a polynucleic acid molecule comprising a single enantiomer. In some instances, the polynucleic acid molecule comprises L-nucleotide. In some instances, the polynucleic acid molecule comprises D-nucleotides. In some instance, a polynucleic acid molecule composition comprises less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1%, or less of its mirror enantiomer. In some cases, a polynucleic acid molecule composition comprises less than 30%, 25%, 20%, 15%, 10%, 5%, 4%, 3%, 2%, 1%, or less of a racemic mixture. In some instances, the polynucleic acid molecule is a polynucleic acid molecule described in: U.S. Patent Publication Nos: 2014/194610 and 2015/211006; and PCT Publication No.: WO2015107425.

**[0103]** In some embodiments, a polynucleic acid molecule described herein is further modified to include an aptamer-conjugating moiety. In some instances, the aptamer conjugating moiety is a DNA aptamer-conjugating moiety. In some instances, the aptamer conjugating moiety is Alphamer (Centauri Therapeutics), which comprises an aptamer portion that recognizes a specific cell-surface

target and a portion that presents a specific epitopes for attaching to circulating antibodies. In some instance, a polynucleic acid molecule described herein is further modified to include an aptamer conjugating moiety as described in: U.S. Patent Nos: 8,604,184, 8,591,910, and 7,850,975.

**[0104]** In additional embodiments, a polynucleic acid molecule described herein is modified to increase its stability. In some embodiments, the polynucleic acid molecule is RNA (e.g., siRNA), and the polynucleic acid molecule is modified to increase its stability. In some instances, the polynucleic acid molecule is modified by one or more of the modifications described above to increase its stability. In some cases, the polynucleic acid molecule is modified at the 2' hydroxyl position, such as by 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modification or by a locked or bridged ribose conformation (e.g., LNA or ENA). In some cases, the polynucleic acid molecule is modified by 2'-O-methyl and/or 2'-O-methoxyethyl ribose. In some cases, the polynucleic acid molecule also includes morpholinos, PNAs, HNA, methylphosphonate nucleotides, thiolphosphonate nucleotides, and/or 2'-fluoro N3-P5'-phosphoramidites to increase its stability. In some instances, the polynucleic acid molecule is a chirally pure (or stereo pure) polynucleic acid molecule. In some instances, the chirally pure (or stereo pure) polynucleic acid molecule is modified to increase its stability. Suitable modifications to the RNA to increase stability for delivery will be apparent to the skilled person.

**[0105]** In some embodiments, a polynucleic acid molecule described herein has RNAi activity that modulates expression of RNA encoded by *MYC*. In some instances, a polynucleic acid molecule described herein is a double-stranded siRNA molecule that down-regulates expression of *MYC*, wherein one of the strands of the double-stranded siRNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence of *MYC* or RNA encoded by *MYC* or a portion thereof, and wherein the second strand of the double-stranded siRNA molecule comprises a nucleotide sequence substantially similar to the nucleotide sequence of *MYC* or RNA encoded by *MYC* or a portion thereof. In some cases, a polynucleic acid molecule described herein is a double-stranded siRNA molecule that down-regulates expression of *MYC*, wherein each strand of the siRNA molecule comprises about 15 to 25, 18 to 24, or 19 to about 23 nucleotides, and wherein each strand comprises at least about 14, 17, or 19 nucleotides that are complementary to the nucleotides of the other strand. In some cases, a polynucleic acid molecule described herein is a double-stranded siRNA molecule that down-regulates expression of *MYC*, wherein each strand of the siRNA molecule comprises about 19 to about 23 nucleotides, and wherein each strand comprises at least about 19 nucleotides that are complementary to the nucleotides of the other

strand. In some instances, the RNAi activity occurs within a cell. In other instances, the RNAi activity occurs in a reconstituted in vitro system.

**[0106]** In some embodiments, a polynucleic acid molecule described herein has RNAi activity that modulates expression of RNA encoded by *MYC*. In some instances, a polynucleic acid molecule described herein is a single-stranded siRNA molecule that down-regulates expression of *MYC*, wherein the single-stranded siRNA molecule comprises a nucleotide sequence that is complementary to a nucleotide sequence of *MYC* or RNA encoded by *MYC* or a portion thereof. In some cases, a polynucleic acid molecule described herein is a single-stranded siRNA molecule that down-regulates expression of *MYC*, wherein the siRNA molecule comprises about 15 to 25, 18 to 24, or 19 to about 23 nucleotides. In some cases, a polynucleic acid molecule described herein is a single-stranded siRNA molecule that down-regulates expression of *MYC*, wherein the siRNA molecule comprises about 19 to about 23 nucleotides. In some instances, the RNAi activity occurs within a cell. In other instances, the RNAi activity occurs in a reconstituted in vitro system.

**[0107]** In some instances, a polynucleic acid molecule is a double-stranded polynucleotide molecule comprising self-complementary sense and antisense regions, wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region has a nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. In some instances, the polynucleic acid molecule is assembled from two separate polynucleotides, where one strand is the sense strand and the other is the antisense strand, wherein the antisense and sense strands are self-complementary (e.g., each strand comprises a nucleotide sequence that is complementary to a nucleotide sequence in the other strand; such as where the antisense strand and sense strand form a duplex or double stranded structure, for example wherein the double stranded region is about 19, 20, 21, 22, 23, or more base pairs); the antisense strand comprises nucleotide sequence that is complementary to a nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense strand comprises a nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof. Alternatively, the polynucleic acid molecule is assembled from a single oligonucleotide, where the self-complementary sense and antisense regions of the polynucleic acid molecule are linked by means of a nucleic acid based or non-nucleic acid-based linker(s).

**[0108]** In some cases, a polynucleic acid molecule is a polynucleotide with a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region comprises a nucleotide sequence that is complementary to a nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region has a nucleotide sequence corresponding to the target nucleic acid

sequence or a portion thereof. In other cases, the polynucleic acid molecule is a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region comprises nucleotide sequence that is complementary to a nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region has a nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide is processed either in vivo or in vitro to generate an active polynucleic acid molecule capable of mediating RNAi. In additional cases, the polynucleic acid molecule also comprises a single stranded polynucleotide having a nucleotide sequence complementary to a nucleotide sequence in a target nucleic acid molecule or a portion thereof (for example, where such polynucleic acid molecule does not require the presence within the polynucleic acid molecule of a nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof), wherein the single stranded polynucleotide further comprises a terminal phosphate group, such as a 5'-phosphate (see for example Martinez et al., 2002, *Cell*, 110, 563-574 and Schwarz et al., 2002, *Molecular Cell*, 10, 537-568), or 5',3'-diphosphate.

**[0109]** In some instances, an asymmetric duplex is a linear polynucleic acid molecule comprising an antisense region, a loop portion that comprises nucleotides or non-nucleotides, and a sense region that comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complimentary nucleotides to base pair with the antisense region and form a duplex with loop. For example, an asymmetric hairpin polynucleic acid molecule comprises an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g. about 19 to about 22 nucleotides) and a loop region comprising about 4 to about 8 nucleotides, and a sense region having about 3 to about 18 nucleotides that are complementary to the antisense region. In some cases, the asymmetric hairpin polynucleic acid molecule also comprises a 5'-terminal phosphate group that is chemically modified. In additional cases, the loop portion of the asymmetric hairpin polynucleic acid molecule comprises nucleotides, non-nucleotides, linker molecules, or conjugate molecules.

**[0110]** In some embodiments, an asymmetric duplex is a polynucleic acid molecule having two separate strands comprising a sense region and an antisense region, wherein the sense region comprises fewer nucleotides than the antisense region to the extent that the sense region has enough complimentary nucleotides to base pair with the antisense region and form a duplex. For example, an asymmetric duplex polynucleic acid molecule comprises an antisense region having length sufficient to mediate RNAi in a cell or in vitro system (e.g. about 19 to about 22 nucleotides) and a sense region having about 3 to about 18 nucleotides that are complementary to the antisense region.

**[0111]** In some cases, a universal base refers to nucleotide base analogs that form base pairs with each of the natural DNA/RNA bases with little discrimination between them. Non-limiting examples of universal bases include C-phenyl, C-naphthyl and other aromatic derivatives, inosine, azole carboxamides, and nitroazole derivatives such as 3-nitropyrrole, 4-nitroindole, 5-nitroindole, and 6-nitroindole as known in the art (see for example Loakes, 2001, *Nucleic Acids Research*, 29, 2437-2447).

#### *Polynucleic Acid Molecule Synthesis*

**[0112]** In some embodiments, a polynucleic acid molecule described herein is constructed using chemical synthesis and/or enzymatic ligation reactions using procedures known in the art. For example, a polynucleic acid molecule is chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the polynucleic acid molecule and target nucleic acids. Exemplary methods include those described in: U.S. Patent Nos. 5,142,047; 5,185,444; 5,889,136; 6,008,400; and 6,111,086; PCT Publication No. WO2009099942; or European Publication No. 1579015. Additional exemplary methods include those described in: Griffey et al., "2'-O-aminopropyl ribonucleotides: a zwitterionic modification that enhances the exonuclease resistance and biological activity of antisense oligonucleotides," *J. Med. Chem.* **39**(26):5100-5109 (1997)); Obika, et al. "Synthesis of 2'-O,4'-C-methylneuridine and -cytidine. Novel bicyclic nucleosides having a fixed C3, -endo sugar pucker." *Tetrahedron Letters* **38** (50): 8735 (1997); Koizumi, M. "ENA oligonucleotides as therapeutics". *Current opinion in molecular therapeutics* **8** (2): 144-149 (2006); and Abramova et al., "Novel oligonucleotide analogues based on morpholino nucleoside subunits-antisense technologies: new chemical possibilities," *Indian Journal of Chemistry* 48B:1721-1726 (2009). Alternatively, the polynucleic acid molecule is produced biologically using an expression vector into which a polynucleic acid molecule has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted polynucleic acid molecule will be of an antisense orientation to a target polynucleic acid molecule of interest).

**[0113]** In some embodiments, a polynucleic acid molecule is synthesized via a tandem synthesis methodology, wherein both strands are synthesized as a single contiguous oligonucleotide fragment or strand separated by a cleavable linker which is subsequently cleaved to provide separate fragments or strands that hybridize and permit purification of the duplex.

**[0114]** In some instances, a polynucleic acid molecule is also assembled from two distinct nucleic acid strands or fragments wherein one fragment includes the sense region and the second fragment includes the antisense region of the molecule.

**[0115]** Additional modification methods for incorporating, for example, sugar, base and phosphate modifications include: Eckstein et al., International Publication PCT No. WO 92/07065; Perrault et al. *Nature*, 1990, 344, 565-568; Pieken et al. *Science*, 1991, 253, 314-317; Usman and Cedergren, *Trends in Biochem. Sci.*, 1992, 17, 334-339; Usman et al. International Publication PCT No. WO 93/15187; Sproat, U.S. Pat. No. 5,334,711 and Beigelman et al., 1995, *J. Biol. Chem.*, 270, 25702; Beigelman et al., International PCT publication No. WO 97/26270; Beigelman et al., U.S. Pat. No. 5,716,824; Usman et al., U.S. Pat. No. 5,627,053; Woolf et al., International PCT Publication No. WO 98/13526; Thompson et al., U.S. Ser. No. 60/082,404 which was filed on Apr. 20, 1998; Karpeisky et al., 1998, *Tetrahedron Lett.*, 39, 1131; Earnshaw and Gait, 1998, *Biopolymers (Nucleic Acid Sciences)*, 48, 39-55; Verma and Eckstein, 1998, *Annu. Rev. Biochem.*, 67, 99-134; and Burlina et al., 1997, *Bioorg. Med. Chem.*, 5, 1999-2010. Such publications describe general methods and strategies to determine the location of incorporation of sugar, base, and/or phosphate modifications and the like into nucleic acid molecules without modulating catalysis.

**[0116]** In some instances, while chemical modification of the polynucleic acid molecule internucleotide linkages with phosphorothioate, phosphorodithioate, and/or 5'-methylphosphonate linkages improves stability, excessive modifications sometimes cause toxicity or decreased activity. Therefore, when designing nucleic acid molecules, the amount of these internucleotide linkages in some cases is minimized. In such cases, the reduction in the concentration of these linkages lowers toxicity, and increases efficacy and specificity of these molecules.

## **Diseases**

**[0117]** In some embodiments, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of a disease or disorder. In some instances, the disease or disorder is a cancer. In some embodiments, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of cancer. In some instances, the cancer is a solid tumor. In some instances, the cancer is a hematologic malignancy. In some instances, the cancer is a relapsed or refractory cancer, or a metastatic cancer. In some instances, the solid tumor is a relapsed or refractory solid tumor, or a metastatic solid tumor. In some cases, the hematologic malignancy is a relapsed or refractory hematologic malignancy, or a metastatic hematologic malignancy.

**[0118]** In some embodiments, the cancer is a solid tumor. Exemplary solid tumor includes, but is not limited to, anal cancer, appendix cancer, bile duct cancer (i.e., cholangiocarcinoma), bladder cancer, brain tumor, breast cancer, cervical cancer, colon cancer, cancer of Unknown Primary (CUP), esophageal cancer, eye cancer, fallopian tube cancer, gastroenterological cancer, kidney

cancer, liver cancer, lung cancer, medulloblastoma, melanoma, oral cancer, ovarian cancer, pancreatic cancer, parathyroid disease, penile cancer, pituitary tumor, prostate cancer, rectal cancer, skin cancer, stomach cancer, testicular cancer, throat cancer, thyroid cancer, uterine cancer, vaginal cancer, or vulvar cancer.

**[0119]** In some instances, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of a solid tumor. In some instances, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of anal cancer, appendix cancer, bile duct cancer (i.e., cholangiocarcinoma), bladder cancer, brain tumor, breast cancer, cervical cancer, colon cancer, cancer of Unknown Primary (CUP), esophageal cancer, eye cancer, fallopian tube cancer, gastroenterological cancer, kidney cancer, liver cancer, lung cancer, medulloblastoma, melanoma, oral cancer, ovarian cancer, pancreatic cancer, parathyroid disease, penile cancer, pituitary tumor, prostate cancer, rectal cancer, skin cancer, stomach cancer, testicular cancer, throat cancer, thyroid cancer, uterine cancer, vaginal cancer, or vulvar cancer. In some instances, the solid tumor is a relapsed or refractory solid tumor, or a metastatic solid tumor.

**[0120]** In some instances, the cancer is a hematologic malignancy. In some instances, the hematologic malignancy is a leukemia, a lymphoma, a myeloma, a non-Hodgkin's lymphoma, or a Hodgkin's lymphoma. In some instances, the hematologic malignancy comprises chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), high risk CLL, a non-CLL/SLL lymphoma, prolymphocytic leukemia (PLL), follicular lymphoma (FL), diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), Waldenström's macroglobulinemia, multiple myeloma, extranodal marginal zone B cell lymphoma, nodal marginal zone B cell lymphoma, Burkitt's lymphoma, non-Burkitt high grade B cell lymphoma, primary mediastinal B-cell lymphoma (PMBL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, B cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, splenic marginal zone lymphoma, plasma cell myeloma, plasmacytoma, mediastinal (thymic) large B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, or lymphomatoid granulomatosis.

**[0121]** In some instances, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of a hematologic malignancy. In some instances, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of a leukemia, a lymphoma, a myeloma, a non-Hodgkin's lymphoma, or a Hodgkin's lymphoma. In some instances, the hematologic malignancy comprises chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), high risk CLL, a non-CLL/SLL lymphoma, prolymphocytic leukemia (PLL), follicular lymphoma (FL), diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), Waldenström's macroglobulinemia, multiple myeloma, extranodal marginal

zone B cell lymphoma, nodal marginal zone B cell lymphoma, Burkitt's lymphoma, non-Burkitt high grade B cell lymphoma, primary mediastinal B-cell lymphoma (PMBL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, B cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, splenic marginal zone lymphoma, plasma cell myeloma, plasmacytoma, mediastinal (thymic) large B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, or lymphomatoid granulomatosis. In some cases, the hematologic malignancy is a relapsed or refractory hematologic malignancy, or a metastatic hematologic malignancy.

**[0122]** In some instances, the cancer is a MYC-associated cancer. In some instances, a polynucleic acid molecule or a pharmaceutical composition described herein is used for the treatment of a MYC-associated cancer. In some instances, the cancer is a solid tumor. In some instances, the cancer is a hematologic malignancy. In some instances, the solid tumor is a relapsed or refractory solid tumor, or a metastatic solid tumor. In some cases, the hematologic malignancy is a relapsed or refractory hematologic malignancy, or a metastatic hematologic malignancy. In some instances, the cancer comprises bladder cancer, breast cancer, colorectal cancer, endometrial cancer, esophageal cancer, glioblastoma multiforme, head and neck cancer, kidney cancer, lung cancer, ovarian cancer, pancreatic cancer, prostate cancer, thyroid cancer, acute myeloid leukemia, CLL, DLBCL, or multiple myeloma.

### **Pharmaceutical Formulation**

**[0123]** In some embodiments, the pharmaceutical formulations described herein are administered to a subject by multiple administration routes including, but not limited to, parenteral (e.g., intravenous, subcutaneous, intramuscular), oral, intranasal, buccal, rectal, or transdermal administration routes. In some instances, the pharmaceutical composition describe herein is formulated for parenteral (e.g., intravenous, subcutaneous, intramuscular) administration. In other instances, the pharmaceutical composition describe herein is formulated for oral administration. In still other instances, the pharmaceutical composition describe herein is formulated for intranasal administration.

**[0124]** In some embodiments, the pharmaceutical formulations include, but are not limited to, aqueous liquid dispersions, self-emulsifying dispersions, solid solutions, liposomal dispersions, aerosols, solid dosage forms, powders, immediate release formulations, controlled release formulations, fast melt formulations, tablets, capsules, pills, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations (e.g., nanoparticle formulations), and mixed immediate and controlled release formulations.

**[0125]** In some instances, the pharmaceutical formulation includes multiparticulate formulations. In some instances, the pharmaceutical formulation includes nanoparticle formulations. In some instances, nanoparticles comprise cMAP, cyclodextrin, or lipids. In some cases, nanoparticles comprise solid lipid nanoparticles, polymeric nanoparticles, self-emulsifying nanoparticles, liposomes, microemulsions, or micellar solutions. Additional exemplary nanoparticles include, but are not limited to, paramagnetic nanoparticles, superparamagnetic nanoparticles, metal nanoparticles, fullerene-like materials, inorganic nanotubes, dendrimers (such as with covalently attached metal chelates), nanofibers, nanohorns, nano-onions, nanorods, nanoropes, and quantum dots. In some instances, a nanoparticle is a metal nanoparticle, e.g., a nanoparticle of scandium, titanium, vanadium, chromium, manganese, iron, cobalt, nickel, copper, zinc, yttrium, zirconium, niobium, molybdenum, ruthenium, rhodium, palladium, silver, cadmium, hafnium, tantalum, tungsten, rhenium, osmium, iridium, platinum, gold, gadolinium, aluminum, gallium, indium, tin, thallium, lead, bismuth, magnesium, calcium, strontium, barium, lithium, sodium, potassium, boron, silicon, phosphorus, germanium, arsenic, antimony, and combinations, alloys, or oxides thereof.

**[0126]** In some instances, a nanoparticle includes a core or a core and a shell, as in a core-shell nanoparticle.

**[0127]** In some instances, a nanoparticle is further coated with molecules for attachment of functional elements (e.g., with one or more of a polynucleic acid molecule or binding moiety described herein). In some instances, a coating comprises chondroitin sulfate, dextran sulfate, carboxymethyl dextran, alginic acid, pectin, carrageenan, fucoidan, agarose, porphyrin, karaya gum, gellan gum, xanthan gum, hyaluronic acids, glucosamine, galactosamine, chitin (or chitosan), polyglutamic acid, polyaspartic acid, lysozyme, cytochrome C, ribonuclease, trypsinogen, chymotrypsinogen,  $\alpha$ -chymotrypsin, polylysine, polyarginine, histone, protamine, ovalbumin, dextrin, or cyclodextrin. In some instances, a nanoparticle comprises a graphene-coated nanoparticle.

**[0128]** In some cases, a nanoparticle has at least one dimension of less than about 500nm, 400nm, 300nm, 200nm, or 100nm.

**[0129]** In some instances, the nanoparticle formulation comprises paramagnetic nanoparticles, superparamagnetic nanoparticles, metal nanoparticles, fullerene-like materials, inorganic nanotubes, dendrimers (such as with covalently attached metal chelates), nanofibers, nanohorns, nano-onions, nanorods, nanoropes or quantum dots. In some instances, a polynucleic acid molecule or a binding moiety described herein is conjugated either directly or indirectly to the nanoparticle. In some instances, at least 1, 5, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, or more polynucleic acid

molecules or binding moieties described herein are conjugated either directly or indirectly to a nanoparticle.

**[0130]** In some embodiments, the pharmaceutical formulation comprise a delivery vector, e.g., a recombinant vector, for the delivery of the polynucleic acid molecule into cells. In some instances, the recombinant vector is DNA plasmid. In other instances, the recombinant vector is a viral vector. Exemplary viral vectors include vectors derived from adeno-associated virus, retrovirus, adenovirus, or alphavirus. In some instances, the recombinant vectors capable of expressing the polynucleic acid molecules provide stable expression in target cells. In additional instances, viral vectors are used that provide for transient expression of polynucleic acid molecules.

**[0131]** In some embodiments, the pharmaceutical formulations include a carrier or carrier materials selected on the basis of compatibility with the composition disclosed herein, and the release profile properties of the desired dosage form. Exemplary carrier materials include, e.g., binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like. Pharmaceutically compatible carrier materials include, but are not limited to, acacia, gelatin, colloidal silicon dioxide, calcium glycerophosphate, calcium lactate, maltodextrin, glycerine, magnesium silicate, polyvinylpyrrolidone (PVP), cholesterol, cholesterol esters, sodium caseinate, soy lecithin, taurocholic acid, phosphatidylcholine, sodium chloride, tricalcium phosphate, dipotassium phosphate, cellulose and cellulose conjugates, sugars sodium stearyl lactylate, carrageenan, monoglyceride, diglyceride, pregelatinized starch, and the like. See, e.g., *Remington: The Science and Practice of Pharmacy*, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Decker, New York, N.Y., 1980; and *Pharmaceutical Dosage Forms and Drug Delivery Systems*, Seventh Ed. (Lippincott Williams & Wilkins 1999).

**[0132]** In some instances, the pharmaceutical formulations further include pH adjusting agents or buffering agents, which include acids such as acetic, boric, citric, lactic, phosphoric, and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, sodium bicarbonate, and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in an acceptable range.

**[0133]** In some instances, the pharmaceutical formulation includes one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having sodium, potassium, or ammonium cations and chloride, citrate, ascorbate, borate,

phosphate, bicarbonate, sulfate, thiosulfate, or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite, and ammonium sulfate.

**[0134]** In some instances, the pharmaceutical formulations further include diluent which are used to stabilize compounds because they provide a more stable environment. Salts dissolved in buffered solutions (which also provide pH control or maintenance) are utilized as diluents in the art, including, but not limited to a phosphate-buffered saline solution. In certain instances, diluents increase bulk of the composition to facilitate compression or create sufficient bulk for homogenous blend for capsule filling. Such compounds include e.g., lactose, starch, mannitol, sorbitol, dextrose, microcrystalline cellulose such as Avicel<sup>®</sup>; dibasic calcium phosphate, dicalcium phosphate dihydrate; tricalcium phosphate, calcium phosphate; anhydrous lactose, spray-dried lactose; pregelatinized starch, compressible sugar, such as Di-Pac<sup>®</sup> (Amstar); mannitol, hydroxypropylmethylcellulose, hydroxypropylmethylcellulose acetate stearate, sucrose-based diluents, confectioner's sugar; monobasic calcium sulfate monohydrate, calcium sulfate dihydrate; calcium lactate trihydrate, dextrates; hydrolyzed cereal solids, amylose; powdered cellulose, calcium carbonate; glycine, kaolin; mannitol, sodium chloride; inositol, bentonite, and the like.

**[0135]** In some cases, the pharmaceutical formulations include disintegration agents or disintegrants to facilitate the breakup or disintegration of a substance. The term "disintegrate" includes both the dissolution and dispersion of the dosage form when contacted with gastrointestinal fluid. Examples of disintegration agents include a starch, e.g., a natural starch such as corn starch or potato starch, a pregelatinized starch such as National 1551 or Amijel<sup>®</sup>, or sodium starch glycolate such as Promogel<sup>®</sup> or Explotab<sup>®</sup>; a cellulose such as a wood product, methylcellulose, e.g., Avicel<sup>®</sup>, Avicel<sup>®</sup> PH101, Avicel<sup>®</sup> PH102, Avicel<sup>®</sup> PH105, Elcema<sup>®</sup> P100, Emcocel<sup>®</sup>, Vivacel<sup>®</sup>, Ming Tia<sup>®</sup>, and Solka-Floc<sup>®</sup>, methylcellulose, croscarmellose, or a cross-linked cellulose, such as cross-linked sodium carboxymethylcellulose (Ac-Di-Sol<sup>®</sup>), cross-linked carboxymethylcellulose, or cross-linked croscarmellose; a cross-linked starch such as sodium starch glycolate, a cross-linked polymer such as crospovidone; a cross-linked polyvinylpyrrolidone, alginate such as alginic acid or a salt of alginic acid such as sodium alginate, a clay such as Veegum<sup>®</sup> HV (magnesium aluminum silicate); a gum such as agar, guar, locust bean, Karaya, pectin, or tragacanth; sodium starch glycolate; bentonite; a natural sponge; a surfactant; a resin such as a cation-exchange resin; citrus pulp; sodium lauryl sulfate; sodium lauryl sulfate in combination starch; and the like.

**[0136]** In some instances, the pharmaceutical formulations include filling agents such as lactose, calcium carbonate, calcium phosphate, dibasic calcium phosphate, calcium sulfate, microcrystalline

cellulose, cellulose powder, dextrose, dextrans, dextrans, starches, pregelatinized starch, sucrose, xylitol, lactitol, mannitol, sorbitol, sodium chloride, polyethylene glycol, and the like.

**[0137]** Lubricants and glidants are also optionally included in the pharmaceutical formulations described herein for preventing, reducing, or inhibiting adhesion or friction of materials. Exemplary lubricants include, e.g., stearic acid, calcium hydroxide, talc, sodium stearyl fumarate, a hydrocarbon such as mineral oil, or hydrogenated vegetable oil such as hydrogenated soybean oil (Sterotex<sup>®</sup>), higher fatty acids and their alkali-metal and alkaline earth metal salts, such as aluminum, calcium, magnesium, zinc, stearic acid, sodium stearates, glycerol, talc, waxes, Stearowet<sup>®</sup>, boric acid, sodium benzoate, sodium acetate, sodium chloride, leucine, a polyethylene glycol (e.g., PEG-4000) or a methoxypolyethylene glycol such as Carbowax<sup>™</sup>, sodium oleate, sodium benzoate, glyceryl behenate, polyethylene glycol, magnesium or sodium lauryl sulfate, colloidal silica such as Syloid<sup>™</sup>, Cab-O-Sil<sup>®</sup>, a starch such as corn starch, silicone oil, a surfactant, and the like.

**[0138]** Plasticizers include compounds used to soften the microencapsulation material or film coatings to make them less brittle. Suitable plasticizers include, e.g., polyethylene glycols such as PEG 300, PEG 400, PEG 600, PEG 1450, PEG 3350, and PEG 800, stearic acid, propylene glycol, oleic acid, triethyl cellulose and triacetin. Plasticizers also function as dispersing agents or wetting agents.

**[0139]** Solubilizers include compounds such as triacetin, triethylcitrate, ethyl oleate, ethyl caprylate, sodium lauryl sulfate, sodium doccusate, vitamin E TPGS, dimethylacetamide, N-methylpyrrolidone, N-hydroxyethylpyrrolidone, polyvinylpyrrolidone, hydroxypropylmethyl cellulose, hydroxypropyl cyclodextrins, ethanol, n-butanol, isopropyl alcohol, cholesterol, bile salts, polyethylene glycol 200-600, glycofurool, transcutool, propylene glycol, dimethyl isosorbide and the like.

**[0140]** Stabilizers include compounds such as any antioxidation agents, buffers, acids, preservatives and the like.

**[0141]** Suspending agents include compounds such as polyvinylpyrrolidone (e.g., polyvinylpyrrolidone K12, polyvinylpyrrolidone K17, polyvinylpyrrolidone K25, or polyvinylpyrrolidone K30), vinyl pyrrolidone/vinyl acetate copolymer (S630), polyethylene glycol (e.g., the polyethylene glycol has a molecular weight of about 300 to about 6000, or about 3350 to about 4000, or about 7000 to about 5400), sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, hydroxymethylcellulose acetate stearate, polysorbate-80, hydroxyethylcellulose, sodium alginate, gums (such as, e.g., gum tragacanth and gum acacia, guar gum, xanthans, including xanthan gum), sugars, cellulosics (such as, e.g., sodium

carboxymethylcellulose, methylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose), polysorbate-80, sodium alginate, polyethoxylated sorbitan monolaurate, polyethoxylated sorbitan monolaurate, povidone, and the like.

**[0142]** Surfactants include compounds such as sodium lauryl sulfate, sodium docusate, Tween 60 or 80, triacetin, vitamin E TPGS, sorbitan monooleate, polyoxyethylene sorbitan monooleate, polysorbates, polaxomers, bile salts, glyceryl monostearate, copolymers of ethylene oxide and propylene oxide, e.g., Pluronic<sup>®</sup> (BASF), and the like. Additional surfactants include polyoxyethylene fatty acid glycerides and vegetable oils, e.g., polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, e.g., octoxynol 10, octoxynol 40. Sometimes, surfactants are included to enhance physical stability or for other purposes.

**[0143]** Viscosity enhancing agents include, e.g., methyl cellulose, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, hydroxypropylmethyl cellulose acetate stearate, hydroxypropylmethyl cellulose phthalate, carbomer, polyvinyl alcohol, alginates, acacia, chitosans, and combinations thereof.

**[0144]** Wetting agents include compounds such as oleic acid, glyceryl monostearate, sorbitan monooleate, sorbitan monolaurate, triethanolamine oleate, polyoxyethylene sorbitan monooleate, polyoxyethylene sorbitan monolaurate, sodium docusate, sodium oleate, sodium lauryl sulfate, sodium docusate, triacetin, Tween 80, vitamin E TPGS, ammonium salts, and the like.

### **Therapeutic Regimens**

**[0145]** In some embodiments, the pharmaceutical compositions described herein are administered for therapeutic applications. In some embodiments, the pharmaceutical composition is administered once per day, twice per day, three times per day or more. The pharmaceutical composition is administered daily, every day, every alternate day, five days a week, once a week, every other week, two weeks per month, three weeks per month, once a month, twice a month, three times per month, or more. The pharmaceutical composition is administered for at least 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, 18 months, 2 years, 3 years, or more.

**[0146]** In some embodiments, one or more pharmaceutical compositions are administered simultaneously, sequentially, or at an interval period of time. In some embodiments, one or more pharmaceutical compositions are administered simultaneously. In some cases, one or more pharmaceutical compositions are administered sequentially. In additional cases, one or more pharmaceutical compositions are administered at an interval period of time (e.g., the first

administration of a first pharmaceutical composition is on day one followed by an interval of at least 1, 2, 3, 4, 5, or more days prior to the administration of at least a second pharmaceutical composition).

**[0147]** In some embodiments, two or more different pharmaceutical compositions are coadministered. In some instances, the two or more different pharmaceutical compositions are coadministered simultaneously. In some cases, the two or more different pharmaceutical compositions are coadministered sequentially without a gap of time between administrations. In other cases, the two or more different pharmaceutical compositions are coadministered sequentially with a gap of about 0.5 hour, 1 hour, 2 hour, 3 hour, 12 hours, 1 day, 2 days, or more between administrations.

**[0148]** In the case wherein the patient's status does improve, upon the doctor's discretion the administration of the composition is given continuously; alternatively, the dose of the composition being administered is temporarily reduced or temporarily suspended for a certain length of time (i.e., a "drug holiday"). In some instances, the length of the drug holiday varies between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35 days, 50 days, 70 days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, or 365 days. The dose reduction during a drug holiday is from 10%-100%, including, by way of example only, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%.

**[0149]** Once improvement of the patient's conditions has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, are optionally reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained.

**[0150]** In some embodiments, the amount of a given agent that corresponds to such an amount varies depending upon factors such as the particular compound, the severity of the disease, the identity (e.g., weight) of the subject or host in need of treatment, but nevertheless is routinely determined in a manner known in the art according to the particular circumstances surrounding the case, including, e.g., the specific agent being administered, the route of administration, and the subject or host being treated. In some instances, the desired dose is conveniently presented in a single dose or as divided doses administered simultaneously (or over a short period of time) or at appropriate intervals, for example as two, three, four or more sub-doses per day.

**[0151]** The foregoing ranges are merely suggestive, as the number of variables in regard to an individual treatment regime is large, and considerable excursions from these recommended values are not uncommon. Such dosages are altered depending on a number of variables, not limited to the

activity of the compound used, the disease or condition to be treated, the mode of administration, the requirements of the individual subject, the severity of the disease or condition being treated, and the judgment of the practitioner.

**[0152]** In some embodiments, toxicity and therapeutic efficacy of such therapeutic regimens are determined by standard pharmaceutical procedures in cell cultures or experimental animals, including, but not limited to, the determination of the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between the toxic and therapeutic effects is the therapeutic index and it is expressed as the ratio between LD50 and ED50. Compounds exhibiting high therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with minimal toxicity. The dosage varies within this range depending upon the dosage form employed and the route of administration utilized.

#### **Kits/Article of Manufacture**

**[0153]** Disclosed herein, in certain embodiments, are kits and articles of manufacture for use with one or more of the compositions and methods described herein. Such kits include a carrier, package, or container that is compartmentalized to receive one or more containers such as vials, tubes, and the like, each of the container(s) comprising one of the separate elements to be used in a method described herein. Suitable containers include, for example, bottles, vials, syringes, and test tubes. In one embodiment, the containers are formed from a variety of materials such as glass or plastic.

**[0154]** The articles of manufacture provided herein contain packaging materials. Examples of pharmaceutical packaging materials include, but are not limited to, blister packs, bottles, tubes, bags, containers, bottles, and any packaging material suitable for a selected formulation and intended mode of administration and treatment.

**[0155]** For example, the container(s) include MYC nucleic acid molecule described herein. Such kits optionally include an identifying description or label or instructions relating to its use in the methods described herein.

**[0156]** A kit typically includes labels listing contents and/or instructions for use and package inserts with instructions for use. A set of instructions will also typically be included.

**[0157]** In one embodiment, a label is on or associated with the container. In one embodiment, a label is on a container when letters, numbers, or other characters forming the label are attached, molded or etched into the container itself; a label is associated with a container when it is present

within a receptacle or carrier that also holds the container, e.g., as a package insert. In one embodiment, a label is used to indicate that the contents are to be used for a specific therapeutic application. The label also indicates directions for use of the contents, such as in the methods described herein.

**[0158]** In certain embodiments, the pharmaceutical compositions are presented in a pack or dispenser device which contains one or more unit dosage forms containing a compound provided herein. The pack, for example, contains metal or plastic foil, such as a blister pack. In one embodiment, the pack or dispenser device is accompanied by instructions for administration. In one embodiment, the pack or dispenser is also accompanied with a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the drug for human or veterinary administration. Such notice, for example, is the labeling approved by the U.S. Food and Drug Administration for prescription drugs, or the approved product insert. In one embodiment, compositions containing a compound provided herein formulated in a compatible pharmaceutical carrier are also prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

### **Certain Terminology**

**[0159]** Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which the claimed subject matter belongs. It is to be understood that the general description and the detailed description are exemplary and explanatory only and are not restrictive of any subject matter claimed. In this application, the use of the singular includes the plural unless specifically stated otherwise. It must be noted that, as used in the specification, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. In this application, the use of “or” means “and/or” unless stated otherwise. Furthermore, use of the term “including” as well as other forms, such as “include”, “includes,” and “included,” is not limiting.

**[0160]** As used herein, ranges and amounts can be expressed as “about” a particular value or range. About also includes the exact amount. Hence “about 5  $\mu$ L” means “about 5  $\mu$ L” and also “5  $\mu$ L.” Generally, the term “about” includes an amount that is expected to be within experimental error, e.g.,  $\pm 5\%$ ,  $\pm 10\%$ , or  $\pm 15\%$ .

**[0161]** The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

[0162] As used herein, the terms “individual(s),” “subject(s),” and “patient(s)” mean any mammal. In some embodiments, the mammal is a human. In some embodiments, the mammal is a non-human. None of the terms require or are limited to situations characterized by the supervision (e.g. constant or intermittent) of a health care worker (e.g. a doctor, a registered nurse, a nurse practitioner, a physician’s assistant, an orderly, or a hospice worker).

### **EXAMPLES**

[0163] These examples are provided for illustrative purposes only and not to limit the scope of the claims provided herein.

#### **Example 1. Evaluation of *in vitro* potency of anti-cMYC siRNAs**

[0164] The six anti-cMYC siRNAs listed in table 1 were transfected into two human cancer cell lines, H358 (NSCLC) and HCT116 (colon cancer). Each siRNA was formulated with a Lipofectamine RNAiMAX (Life Technologies), according to the manufacturer’s “forward transfection” instructions, at a single final concentration of 0.5 nM. Cells were plated 24 h prior to transfection in duplicate within 24-well tissue culture plates. At 96 h post-transfection, RNA was harvested from cells in all wells using Strattec InviTrap® RNA Cell HTS96 kit. The concentration of each isolated RNA was determined via A260 measurement using a NanoDrop spectrophotometer. RNA samples were reverse transcribed to cDNA using the High Capacity RNA to cDNA Kit (Life Technologies) according to the manufacturer’s instructions. cDNA samples were then evaluated by qPCR using cMYC-specific probes with results normalized to endogenous PPIB and quantified using the standard  $2^{-\Delta\Delta C_t}$  method. cMYC mRNA levels were normalized to expression in vehicle controls and are reported in Table 1.

[0165] Of the 6 siRNA candidates tested, three of them achieved  $\geq 80\%$  down-regulation of cMYC in HCT116 cells, while their inhibitory activity in H358 was about 40-60%. Based upon the results from this initial single-concentration experiment, we selected a total of three candidate siRNAs (Avidity ID # 1160, 1162, and 1164) for multi-concentration testing, to determine  $IC_{50}$  values (i.e., the concentration required to reduce cMYC expression by 50%), in HCT116. Cells were transfected as above at concentrations starting from 10000 pM using serial dilutions. At 96 h post-transfection, RNA was harvested and normalized cMYC mRNA levels were quantified as described above. Results are presented in Table 1. All the three siRNAs were found to be potent at down-regulating cMYC mRNA in both the human and mouse cell lines tested, with  $IC_{50}$  values less than 1 nM.

#### **Table 1.**

Avidity ID #	Target gene	Sense Strand Sequence (5'-3') Passenger Strand (PS)2	SEQ ID NO:	Antisense Strand Sequence (5'-3') Guide Strand (GS)3	SEQ ID NO:	H358, % mRNA at 0.5nM	HCT116, % mRNA at 0.5nM	HCT116 IC50 (pM)
R-1160	cMyc	AGGAACUAUGA CCUCGACUdTsd T	1	AGUCGAGGUCAU AGUUCCUdTsdT	2	56.7	17.2	118.3
R-1161	cMyc	ACGACGAGACC UUCAUCAAdTsd T	3	UUGAUGAAGGUC UCGUCGUdTsdT	4	76.8	51.4	
R-1162	cMyc	AAGAUGAGGAA GAAAU CGAdTsd T	5	UCGAUUUCUCC UCAUCUdTsdT	6	39.3	19.2	43.4
R-1163	cMyc	AGGAAGAAAUC GAUGUUGUdTsd T	7	ACAACAUCGAUU UCUUCCUdTsdT	8	41.4	18.2	
R-1164	cMyc	AGCUUUUUUGC CCUGCGUGdTsd T	9	CACGCAGGGCAA AAAAGCUdTsdT	10	41.5	17.5	55.2
R-1165	cMyc	AGGUAGUUAUC CUUAAAAAdTsd T	11	UUUUUAAGGAU AACUACCUdTsdT	12	102.5	100.5	

siRNA Sequence with Chemical Modification Info

lower case (n) = dT = deoxy-T residue;

s = phosphorothioate backbone modification

**[0166]** While preferred embodiments of the present disclosure have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the disclosure. It should be understood that various alternatives to the embodiments of the disclosure described herein may be employed in practicing the disclosure. It is intended that the following claims define the scope of the disclosure and that methods and structures within the scope of these claims and their equivalents be covered thereby.

## CLAIMS

### WHAT IS CLAIMED IS:

1. A polynucleic acid molecule that mediates RNA interference against MYC, wherein the polynucleic acid molecule comprises at least 80% sequence identity to a sequence selected from SEQ ID NOs: 1-12, wherein the polynucleic acid molecule hybridizes to a MYC target sequence with less than 4 mismatched bases, wherein the polynucleic acid molecule comprises at least one 2' modified nucleotide, at least one modified internucleotide linkage, or at least one inverted abasic moiety, and wherein the polynucleic acid molecule is from about 10 to about 50 nucleotides in length.
2. The polynucleic acid molecule of claim 1, wherein the polynucleic acid molecule comprises at least 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to a sequence selected from SEQ ID NOs: 1-12.
3. The polynucleic acid molecule of claim 1 or 2, wherein the polynucleic acid molecule hybridizes to a MYC target sequence with less than 3 mismatched bases, less than 2 mismatched bases, or less than 1 mismatched bases.
4. The polynucleic acid molecule of claim 1 or 2, wherein the at least one 2' modified nucleotide comprises 2'-O-methyl, 2'-O-methoxyethyl (2'-O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), or 2'-O-N-methylacetamido (2'-O-NMA) modified nucleotide.
5. The polynucleic acid molecule of claim 1 or 2, wherein the at least one 2' modified nucleotide comprises locked nucleic acid (LNA) or ethylene nucleic acid (ENA).
6. The polynucleic acid molecule of claim 1 or 2, wherein the at least one inverted basic moiety is at at least one terminus.
7. The polynucleic acid molecule of claim 1 or 2, wherein the at least one modified internucleotide linkage comprises a phosphorothioate linkage or a phosphorodithioate linkage.
8. The polynucleic acid molecule of any one of the claims 1-7, wherein the polynucleic acid molecule is from about 10 to about 30 nucleotides in length.
9. The polynucleic acid molecule of any one of the claims 1-8, wherein the polynucleic acid molecule is at least 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides in length.

10. The polynucleic acid molecule of any one of the claims 1-9, wherein the polynucleic acid molecule comprises at least one of: from about 5% to about 100% modification, from about 10% to about 100% modification, from about 20% to about 100% modification, from about 30% to about 100% modification, from about 40% to about 100% modification, from about 50% to about 100% modification, from about 60% to about 100% modification, from about 70% to about 100% modification, from about 80% to about 100% modification, and from about 90% to about 100% modification.
11. The polynucleic acid molecule of any one of the claims 1-9, wherein the polynucleic acid molecule comprises at least one of: from about 10% to about 80% modification, from about 20% to about 80% modification, from about 30% to about 80% modification, from about 40% to about 80% modification, from about 50% to about 80% modification, from about 60% to about 80% modification, and from about 70% to about 80% modification.
12. The polynucleic acid molecule of any one of the claims 1-9, wherein the polynucleic acid molecule comprises at least one of: from about 10% to about 60% modification, from about 20% to about 60% modification, from about 30% to about 60% modification, from about 40% to about 60% modification, and from about 50% to about 60% modification.
13. The polynucleic acid molecule of any one of the claims 1-9, wherein the polynucleic acid molecule comprises at least 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 99% modification.
14. The polynucleic acid molecule of any one of the claims 1-9, wherein the polynucleic acid molecule comprises about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, about 12, about 13, about 14, about 15, about 16, about 17, about 18, about 19, about 20, about 21, about 22, or more modified nucleotides.
15. The polynucleic acid molecule of claim 1, wherein the polynucleic acid molecule further hybridizes to a second polynucleotide to form a double-stranded polynucleic acid molecule.
16. The polynucleic acid molecule of claim 15, wherein the second polynucleotide comprises at least one modification.
17. The polynucleic acid molecule of claim 15, wherein the second polynucleotide comprises a sequence having at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to a sequence selected from SEQ ID NOs: 1-12.
18. The polynucleic acid molecule of any one of the claims 1-17, wherein the polynucleic acid molecule is RNA.
19. The polynucleic acid molecule of any one of the claims 1-18, wherein MYC is *c-MYC* gene, *L-MYC* gene, or *N-MYC* gene.

20. The polynucleic acid molecule of any one of the claims 1-18, wherein MYC is *c-MYC* gene.
21. A pharmaceutical composition comprising:
  - a) a molecule of claims 1-20; and
  - b) a pharmaceutically acceptable excipient.
22. The pharmaceutical composition of claim 21, wherein the pharmaceutical composition is formulated as a nanoparticle formulation.
23. The pharmaceutical composition of claim 21, wherein the pharmaceutical composition is formulated for parenteral, oral, intranasal, buccal, rectal, or transdermal administration.
24. A method of treating a disease or disorder in a patient in need thereof, comprising administering to the patient a composition comprising a molecule of claims 1-20.
25. The method of claim 24, wherein the disease or disorder is a cancer.
26. The method of claim 25, wherein the cancer comprises a MYC-associated cancer.
27. The method of claim 25, wherein the cancer is a solid tumor.
28. The method of claim 25, wherein the cancer is a hematologic malignancy.
29. The method of claim 25, wherein the cancer comprises bladder cancer, breast cancer, colorectal cancer, endometrial cancer, esophageal cancer, glioblastoma multiforme, head and neck cancer, kidney cancer, lung cancer, ovarian cancer, pancreatic cancer, prostate cancer, or thyroid cancer.
30. The method of claim 25, wherein the cancer comprises acute myeloid leukemia, CLL, DLBCL, or multiple myeloma.
31. A method of inhibiting the expression of *MYC* gene in a primary cell of a patient, comprising administering a molecule of claims 1-20 to the primary cell.
32. The method of claim 31, wherein the method is an *in vivo* method.
33. The method of claim 31, wherein the patient is a human.