




(86) Date de dépôt PCT/PCT Filing Date: 2003/03/31  
(87) Date publication PCT/PCT Publication Date: 2003/11/13  
(45) Date de délivrance/Issue Date: 2011/07/05  
(85) Entrée phase nationale/National Entry: 2004/10/25  
(86) N° demande PCT/PCT Application No.: EP 2003/003331  
(87) N° publication PCT/PCT Publication No.: 2003/093235  
(30) Priorités/Priorities: 2002/04/27 (DE102 18 974.9);  
2002/08/12 (DE102 36 868.6)

(51) Cl.Int./Int.Cl. *C07D 211/72* (2006.01),  
*A61K 31/40* (2006.01), *A61K 31/4025* (2006.01),  
*A61K 31/4172* (2006.01), *A61K 31/4178* (2006.01),  
*A61K 31/4545* (2006.01), *A61P 25/06* (2006.01),  
*A61P 29/00* (2006.01), *A61P 35/00* (2006.01),  
*A61P 7/02* (2006.01), *A61P 9/00* (2006.01),  
*C07D 207/14* (2006.01), ...

(72) Inventeurs/Inventors:  
DORSCH, DIETER, DE;  
MEDERSKI, WERNER, DE;  
GLEITZ, JOHANNES, DE;  
CEZANNE, BERTRAM, DE;  
TSAKLAKIDIS, CHRISTOS, DE; ...

(54) Titre : AMIDES D'ACIDE CARBOXYLIQUE SERVANT D'INHIBITEURS DU FACTEUR DE COAGULATION XA  
(54) Title: CARBOXAMIDES



(57) Abrégé/Abstract:

Novel compounds of the formula I (see formula I) in which D, W, X, Y, T and R<sup>1</sup> are as defined in Patent Claim 1, are inhibitors of coagulation factor Xa and can be employed for the prophylaxis and/or therapy of thromboembolic diseases and for the treatment of tumours.

(51) Cl.Int./Int.Cl. (suite/continued) *C07D 207/22* (2006.01), *C07D 211/56* (2006.01), *C07D 213/73* (2006.01),  
*C07D 233/46* (2006.01), *C07D 233/48* (2006.01), *C07D 265/30* (2006.01), *C07D 401/12* (2006.01),  
*C07D 403/12* (2006.01), *C07D 409/12* (2006.01), *C07D 453/06* (2006.01)

(72) Inventeurs(suite)/Inventors(continued): BARNES, CHRISTOPHER, DE

(73) Propriétaires(suite)/Owners(continued):MERCK PATENT GESELLSCHAFT MIT BESCHRAENKTER HAFTUNG, DE


(74) Agent: FETHERSTONHAUGH & CO.

- 70 -

**Abstract**

Novel compounds of the formula I

5



10

in which

D, W, X, Y, T and R<sup>1</sup> are as defined in Patent Claim 1,

15

are inhibitors of coagulation factor Xa and can be employed for the prophylaxis and/or therapy of thromboembolic diseases and for the treatment of tumours.

20

25

30


35

- 1 -

# Carboxamides

The invention relates to compounds of the formula I

5



10 in which

D is phenyl, pyridyl or thienyl, each of which is unsubstituted or monosubstituted or polysubstituted by Hal, A,  $\text{OR}^2$ ,  $\text{N}(\text{R}^2)_2$ ,  $\text{NO}_2$ , CN,  $\text{COOR}^2$  or  $\text{CON}(\text{R}^2)_2$ ,

15

X is  $\text{NR}^3$  or O,

$\text{R}^1$  is H, Ar, Het, cycloalkyl or

A, which may be substituted by  $\text{OR}^2$ ,  $\text{S}(\text{O})_m\text{R}^2$ ,  $\text{N}(\text{R}^2)_2$ , Ar, Het, cycloalkyl, CN,  $\text{COOR}^2$  or  $\text{CON}(\text{R}^2)_2$ ,

20

$\text{R}^2$  is H, A,  $-\text{C}(\text{R}^3)_2]_n\text{-Ar}$ ,  $-\text{C}(\text{R}^3)_2]_n\text{-Het}$ ,  $-\text{C}(\text{R}^3)_2]_n\text{-cycloalkyl}$ ,  $-\text{C}(\text{R}^3)_2]_n\text{-N}(\text{R}^3)_2$  or  $-\text{C}(\text{R}^3)_2]_n\text{-OR}^3$ ,

$\text{R}^3$  is H or A,

W is  $-\text{C}(\text{R}^3)_2]_n-$ ,

Y is alkylene, cycloalkylene, Het-diyl or Ar-diyl,

25

T is a monocyclic or bicyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having from 1 to 4 N, O and/or S atoms, which is monosubstituted or disubstituted by  $=\text{S}$ ,  $=\text{NR}^2$ ,  $=\text{NOR}^2$ ,  $=\text{NCOR}^2$ ,  $=\text{NCOOR}^2$ ,  $=\text{NOCOR}^2$ ,  $=\text{N-CN}$  or  $=\text{N-NO}_2$  and may furthermore be monosubstituted, disubstituted or trisubstituted by Hal, A,  $-\text{C}(\text{R}^3)_2]_n\text{-Ar}$ ,  $-\text{C}(\text{R}^3)_2]_n\text{-Het}$ ,  $-\text{C}(\text{R}^3)_2]_n\text{-cycloalkyl}$ ,  $\text{OR}^3$ ,  $\text{N}(\text{R}^3)_2$ ,  $\text{NO}_2$ , CN,  $\text{COOR}^2$ ,  $\text{CON}(\text{R}^2)_2$ ,  $\text{NR}^2\text{COA}$ ,  $\text{NR}^2\text{CON}(\text{R}^2)_2$ ,  $\text{NR}^2\text{SO}_2\text{A}$ ,  $\text{COR}^2$ ,  $\text{SO}_2\text{NR}^2$  and/or  $\text{S}(\text{O})_m\text{A}$ ,

30

A is unbranched or branched alkyl having 1-10 carbon atoms, in which one or two  $\text{CH}_2$  groups may be replaced by O or S atoms

35

- 2 -

and/or by -CH=CH- groups and/or in addition 1-7 H atoms may be replaced by F,

- 5 Ar is phenyl, naphthyl or biphenyl, each of which is unsubstituted or monosubstituted, disubstituted or trisubstituted by Hal, A, OR<sup>3</sup>, N(R<sup>3</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, COOR<sup>3</sup>, CON(R<sup>3</sup>)<sub>2</sub>, NR<sup>3</sup>COA, NR<sup>3</sup>CON(R<sup>3</sup>)<sub>2</sub>, NR<sup>3</sup>SO<sub>2</sub>A, COR<sup>3</sup>, SO<sub>2</sub>N(R<sup>3</sup>)<sub>2</sub>, S(O)<sub>m</sub>A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-COOR<sup>2'</sup> or -O-[C(R<sup>3</sup>)<sub>2</sub>]<sub>o</sub>-COOR<sup>2'</sup>,
- 10 R<sup>2'</sup> is H, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar', -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Het', -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub> or -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>3</sup>,
- R<sup>2''</sup> is H, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar', + -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub> or -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>3</sup>,
- 15 Ar' is phenyl or benzyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal or A,
- Het is a monocyclic or bicyclic saturated, unsaturated or aromatic heterocyclic ring having from 1 to 4 N, O and/or S atoms, which may be unsubstituted or monosubstituted, disubstituted or trisubstituted by carbonyl oxygen, =S, =N(R<sup>3</sup>)<sub>2</sub>, Hal, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Het<sup>1</sup>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>2'</sup>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>2'</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-COOR<sup>2'</sup>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-CON(R<sup>2'</sup>)<sub>2</sub>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-NR<sup>2'</sup>COA, NR<sup>2'</sup>CON(R<sup>2'</sup>)<sub>2</sub>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-NR<sup>2'</sup>SO<sub>2</sub>A, COR<sup>2'</sup>, SO<sub>2</sub>NR<sup>2'</sup> and/or S(O)<sub>m</sub>A,
- 20
- 25 Het<sup>1</sup> is a monocyclic or bicyclic saturated, unsaturated or aromatic heterocyclic ring having 1 or 2 N, O and/or S atoms, which may be unsubstituted or monosubstituted or disubstituted by carbonyl oxygen, =S, =N(R<sup>3</sup>)<sub>2</sub>, Hal, A, OR<sup>2''</sup>, N(R<sup>2''</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, COOR<sup>2''</sup>, CON(R<sup>2''</sup>)<sub>2</sub>, NR<sup>2''</sup>COA, NR<sup>2''</sup>CON(R<sup>2''</sup>)<sub>2</sub>, NR<sup>2''</sup>SO<sub>2</sub>A, COR<sup>2''</sup>, SO<sub>2</sub>NR<sup>2''</sup> and/or S(O)<sub>m</sub>A,
- 30
- Hal is F, Cl, Br or I,
- n is 0, 1 or 2,
- m is 0, 1 or 2,
- 35 o is 1, 2 or 3,

- 3 -

and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios.

5 The invention had the object of finding novel compounds having valuable properties, in particular those which can be used for the preparation of medicaments.

10 It has been found that the compounds of the formula I and their salts have very valuable pharmacological properties and are well tolerated. In particular, they exhibit factor Xa-inhibiting properties and can therefore be employed for combating and preventing thromboembolic diseases, such as thrombosis, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty and claudicatio inter-

15 mittens.

The compounds of the formula I according to the invention may furthermore be inhibitors of the coagulation factors factor VIIa, factor IXa and

20 thrombin in the blood coagulation cascade.

Aromatic amidine derivatives having an antithrombotic action are disclosed, for example, in EP 0 540 051 B1, WO 98/28269, WO 00/71508,

25 WO 00/71511, WO 00/71493, WO 00/71507, WO 00/71509, WO 00/71512, WO 00/71515 oder WO 00/71516. cyclic guanidines for the treatment of thromboembolic diseases are described, for example, in WO 97/08165. Aromatic heterocyclic compounds having factor Xa-inhibitory activity are disclosed, for example, in WO 96/10022. Substituted

30 N-[(aminoiminomethyl)phenylalkyl]azaheterocyclamides as factor Xa inhibitors are described in WO 96/40679.

35 The antithrombotic and anticoagulant effect of the compounds according to the invention is attributed to the inhibitory action against activated coagulation protease, known by the name factor Xa, or to the inhibition of

- 4 -

other activated serine proteases, such as factor VIIa, factor IXa or thrombin.

5 Factor Xa is one of the proteases involved in the complex process of blood coagulation. Factor Xa catalyses the conversion of prothrombin into thrombin. Thrombin cleaves fibrinogen into fibrin monomers, which, after crosslinking, make an elementary contribution to thrombus formation. Activation of thrombin may result in the occurrence of thromboembolic diseases. However, inhibition of thrombin may inhibit the fibrin formation involved in thrombus formation. The inhibition of thrombin can be measured, for example, by the method of G.F. Cousins et al. in *Circulation* 10 **1996**, 94, 1705-1712.

15 Inhibition of factor Xa can thus prevent the formation of thrombin.

The compounds of the formula I according to the invention and their salts engage in the blood coagulation process by inhibiting factor Xa and thus 20 inhibit the formation of thrombuses.

The inhibition of factor Xa by the compounds according to the invention and the measurement of the anticoagulant and antithrombotic activity can be determined by conventional in-vitro or in-vivo methods. A suitable 25 method is described, for example, by J. Hauptmann et al. in *Thrombosis and Haemostasis* **1990**, 63, 220-223.

30 The inhibition of factor Xa can be measured, for example, by the method of T. Hara et al. in *Thromb. Haemostas.* **1994**, 71, 314-319.

Coagulation factor VIIa initiates the extrinsic part of the coagulation cascade after binding to tissue factor and contributes to the activation of factor 35

- 5 -

X to give factor Xa. Inhibition of factor VIIa thus prevents the formation of factor Xa and thus subsequent thrombin formation.

5

The inhibition of factor VIIa by the compounds according to the invention and the measurement of the anticoagulant and antithrombotic activity can be determined by conventional in-vitro or in-vivo methods. A conventional method for the measurement of the inhibition of factor VIIa is described, for example, by H. F. Ronning et al. in *Thrombosis Research* **1996**, *84*,  
10 73-81.

15

Coagulation factor IXa is generated in the intrinsic coagulation cascade and is likewise involved in the activation of factor X to give factor Xa. Inhibition of factor IXa can therefore prevent the formation of factor Xa in a different way.

20

The inhibition of factor IXa by the compounds according to the invention and the measurement of the anticoagulant and antithrombotic activity can be determined by conventional in-vitro or in-vivo methods. A suitable method is described, for example, by J. Chang et al. in *Journal of Biological Chemistry* **1998**, *273*, 12089-12094.

25

The compounds according to the invention may furthermore be used for the treatment of tumours, tumour diseases and/or tumour metastases. A correlation between tissue factor TF / factor VIIa and the development of various types of cancer has been indicated by T. Taniguchi and N.R. Lemoine in *Biomed. Health Res.* (2000), *41* (Molecular Pathogenesis of Pancreatic Cancer), 57-59.

30

The publications listed below describe an antitumoural action of TF-VII and factor Xa inhibitors for various types of tumour:

35

K.M. Donnelly et al. in *Thromb. Haemost.* 1998; *79*: 1041-1047;  
E.G. Fischer et al. in *J. Clin. Invest.* *104*: 1213-1221 (1999);

B.M. Mueller et al. in J. Clin. Invest. 101: 1372-1378 (1998);

M.E. Bromberg et al. in Thromb. Haemost. 1999; 82: 88-92

5

The compounds of the formula I can be employed as medicament active ingredients in human and veterinary medicine, in particular for the treatment and prevention of thromboembolic diseases, such as thrombosis, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, venous

10

thrombosis, pulmonary embolism, arterial thrombosis, myocardial ischaemia, unstable angina and strokes based on thrombosis.

15

The compounds according to the invention are also employed for the treatment or prophylaxis of atherosclerotic diseases, such as coronary arterial disease, cerebral arterial disease or peripheral arterial disease.

The compounds are also employed in combination with other thrombolytic agents in the case of myocardial infarction, furthermore for prophylaxis for reocclusion after thrombolysis, percutaneous transluminal angioplasty (PTCA) and coronary bypass operations.

20

The compounds according to the invention are furthermore used for the prevention of rethrombosis in microsurgery, furthermore as anticoagulants in connection with artificial organs or in haemodialysis.

25

The compounds are furthermore used in the cleaning of catheters and medical aids *in vivo* in patients, or as anticoagulants for the preservation of blood, plasma and other blood products *in vitro*. The compounds according to the invention are furthermore used for diseases in which blood coagulation makes a crucial contribution to the course of the disease or represents a source of secondary pathology, such as, for example, in cancer, including metastasis, inflammatory disorders, including arthritis, and diabetes.

30

The compounds according to the invention are furthermore used in the treatment of migraine finden (F.Morales-Asin et al., Headache, 40, 2000, 45-47).

35

26474-891

- 7 -


In the treatment of the diseases described, the compounds according to the invention are also employed in combination with other thrombolytically active compounds, such as, for example, with "tissue plasminogen activator" t-PA, modified t-PA, streptokinase or urokinase. The compounds according to the invention are given either at the same time as or before or after the other substances mentioned.

Particular preference is given to simultaneous administration with aspirin in order to prevent recurrence of the clot formation.

The compounds according to the invention are also used in combination with blood platelet glycoprotein receptor (IIb/IIIa) antagonists, which inhibit blood platelet aggregation.

The invention relates to the compounds of the formula I and salts thereof and to a process for the preparation of compounds of the formula I according to Claims 1-16 and pharmaceutically usable derivatives, solvates and stereoisomers thereof, characterised in that

a) a compound of the formula II



in which

$\text{R}^1$ , X, W, Y and T are as defined in Claim 1,

is reacted with a compound of the formula III



in which

D is as defined in Claim 1,

- 8 -


or

b) a compound of the formula IV



in which W, Y and T are as defined in Claim 1,

is reacted with a compound of the formula V



in which

L is Cl, Br, I or a free or reactively functionally modified OH group, and R<sup>1</sup>, X and D are as defined in Claim 1,

and/or

c) in that a radical T in a compound of the formula I is converted into another radical T

by, for example,

i) converting a sulfanyl compound into an imino compound,

and/or

a base or acid of the formula I is converted into one of its salts.

- 9 -

The invention also relates to the optically active forms (stereoisomers), the enantiomers, the racemates, the diastereomers and the hydrates and solvates of these compounds. The term solvates of the compounds is taken to mean adductions of inert solvent molecules onto the compounds which form owing to their mutual attractive force. Solvates are, for example, monohydrates or dihydrates or alcoholates.

The term pharmaceutically usable derivatives is taken to mean, for example, the salts of the compounds according to the invention and also so-called prodrug compounds.

The term prodrug derivatives is taken to mean, for example, compounds of the formula I which have been modified with, for example, alkyl or acyl groups, sugars or oligopeptides and which are rapidly cleaved in the organism to give the effective compounds according to the invention. These also include biodegradable polymer derivatives of the compounds according to the invention, as described, for example, in Int. J. Pharm. 115, 61-67 (1995).

The invention also relates to mixtures of the compounds of the formula I according to the invention, for example mixtures of two diastereomers, for example in the ratio 1:1, 1:2, 1:3, 1:4, 1:5, 1:10, 1:100 or 1:1000.

These are particularly preferably mixtures of stereoisomeric compounds.

For all radicals which occur more than once, for example A, their meanings are independent of one another.

Above and below, the radicals or parameters D, W, X, Y, T and R<sup>1</sup> are as defined under the formula I, unless expressly stated otherwise.

A is alkyl, is unbranched (linear) or branched, and has 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbon atoms. A is preferably methyl, furthermore ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl or tert-butyl, furthermore also pentyl, 1-, 2-

- 10 -

or 3-methylbutyl, 1,1-, 1,2- or 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-, 2-, 3- or 4-methylpentyl, 1,1-, 1,2-, 1,3-, 2,2-, 2,3- or 3,3-dimethylbutyl, 1- or 2-ethylbutyl, 1-ethyl-1-methylpropyl, 1-ethyl-2-methylpropyl, 1,1,2- or 1,2,2-trimethylpropyl, furthermore preferably, for example, trifluoromethyl.

5 A is very particularly preferably alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, preferably methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, trifluoromethyl, pentafluoroethyl or 1,1,1-trifluoroethyl.

10 Cycloalkyl is preferably cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl.

Alkylene is preferably methylene, ethylene, propylene, butylene, pentylene or hexylene, furthermore branched alkylene.

15  $\text{COR}^2$  is for example, CHO or -COA.

-COA (acyl) is preferably acetyl, propionyl, furthermore also butyryl, pentanoyl, hexanoyl or, for example, benzoyl.

20 Hal is preferably F, Cl or Br, but alternatively I.

Ar is, for example, phenyl, o-, m- or p-tolyl, o-, m- or p-ethylphenyl, o-, m- or p-propylphenyl, o-, m- or p-isopropylphenyl, o-, m- or p-tert-butylphenyl, o-, m- or p-hydroxyphenyl, o-, m- or p-nitrophenyl, o-, m- or p-aminophenyl,

25 o-, m- or p-(N-methylamino)phenyl, o-, m- or p-(N-methylaminocarbonyl)-phenyl, o-, m- or p-acetamidophenyl, o-, m- or p-methoxyphenyl, o-, m- or p-ethoxyphenyl, o-, m- or p-ethoxycarbonylphenyl, o-, m- or p-(N,N-dimethylamino)-phenyl, o-, m- or p-(N,N-dimethylaminocarbonyl)phenyl, o-, m- or p-(N-ethylamino)-phenyl, o-, m- or p-(N,N-diethylamino)-phenyl, o-, m- or p-fluorophenyl, o-, m- or p-bromophenyl, o-, m- or p-chlorophenyl, o-, m- or p-(methylsulfonamido)phenyl, o-, m- or p-(methylsulfonyl)-phenyl,

30 further preferably 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-difluorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dichlorophenyl, 2,3-, 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dibromophenyl, 2,4- or 2,5-dinitrophenyl, 2,5- or 3,4-dimethoxyphenyl, 3-nitro-4-chlorophenyl, 3-amino-4-chloro-, 2-amino-3-chloro-, 2-amino-4-

35

- 11 -

chloro-, 2-amino-5-chloro- or 2-amino-6-chlorophenyl, 2-nitro-4-N,N-dimethylamino- or 3-nitro-4-N,N-dimethylaminophenyl, 2,3-diaminophenyl, 2,3,4-, 2,3,5-, 2,3,6-, 2,4,6- or 3,4,5-trichlorophenyl, 2,4,6-trimethoxyphenyl, 2-hydroxy-3,5-dichlorophenyl, p-iodophenyl, 3,6-dichloro-4-amino-phenyl, 4-fluoro-3-chlorophenyl, 2-fluoro-4-bromophenyl, 2,5-difluoro-4-bromophenyl, 3-bromo-6-methoxyphenyl, 3-chloro-6-methoxyphenyl, 3-chloro-4-acetamidophenyl, 3-fluoro-4-methoxyphenyl, 3-amino-6-methylphenyl, 3-chloro-4-acetamidophenyl or 2,5-dimethyl-4-chlorophenyl.

10

Ar is preferably, for example, phenyl which is unsubstituted or monosubstituted, disubstituted or trisubstituted by Hal, A,  $OR^2$ ,  $SO_2A$ ,  $COOR^2$  or CN phenyl.

15

Ar is particularly preferably, for example, phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OA,  $SO_2A$ ,  $SO_2NH_2$ ,  $COOR^2$  or CN, such as, for example, phenyl, 2-methylsulfonylphenyl, 2-amino-sulfonylphenyl, 2-, 3- or 4-chlorophenyl, 4-methylphenyl, 4-bromophenyl, 3-fluoro-4-methoxyphenyl, 4-trifluoromethoxyphenyl, 4-ethoxyphenyl, 2-methoxyphenyl, 3-cyanophenyl or 4-ethoxycarbonylphenyl.

20

Ar is very particularly preferably unsubstituted phenyl, 4-chlorophenyl or 2-methylsulfonylphenyl.

25

Y is preferably Het-diyl or Ar-diyl, particularly preferably 1,4-phenylene which is unsubstituted or monosubstituted by A, Cl or F, furthermore also pyridinediyl, preferably pyridine-2,5-diyl or piperidinediyl.

Y is in particular 1,3- or 1,4-phenylene which is unsubstituted or monosubstituted by methyl, ethyl, propyl, Cl or F.

30

35

Het is, for example, 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or 5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl,

- 12 -

1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or  
-5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, pyrazinyl, 1-, 2-, 3-, 4-,  
5-, 6- or 7-indolyl, 4- or 5-isoindolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-,  
5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or  
7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or  
7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-,  
7- or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl, 3-, 4-, 5-, 6-, 7- or  
8-quinoliny, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, 5- or 6-quinoxaliny, 2-, 3-,  
5-, 6-, 7- or 8-2H-benzo-1,4-oxazinyl, furthermore preferably 1,3-benzo-  
dioxol-5-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazol-4- or -5-yl or 2,1,3-  
benzoxadiazol-5-yl.

The heterocyclic radicals may also be partially or fully hydrogenated.

Het can thus, for example, also be 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-  
dihydro-2-, -3-, -4- or 5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl,  
tetrahydro-2- or -3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-  
dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -  
2- or -4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-,  
-3- or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-  
1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or 4-piperidinyl, 2-, 3- or 4-  
morpholiny, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4-  
or -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-  
pyrimidinyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-,  
-6-, -7- or -8-quinolyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-iso-  
quinolyl, 2-, 3-, 5-, 6-, 7- or 8- 3,4-dihydro-2H-benzo-1,4-oxazinyl,  
furthermore preferably 2,3-methylenedioxyphenyl, 3,4-methylenedioxy-  
phenyl, 2,3-ethylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 3,4-(difluoro-  
methylenedioxy)phenyl, 2,3-dihydrobenzofuran-5- or 6-yl, 2,3-(2-oxo-  
methylenedioxy)phenyl or alternatively 3,4-dihydro-2H-1,5-benzodioxepin-  
6- or -7-yl, furthermore preferably 2,3-dihydrobenzofuranyl or 2,3-dihydro-  
2-oxofuranyl.

- 13 -

5 T is preferably a monocyclic or bicyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having from 1 to 3 N, O and/or S atoms which is monosubstituted or disubstituted by =S, =NR<sup>2</sup>, =NOR<sup>2</sup>, =NCOR<sup>2</sup>, =NCOOR<sup>2</sup>, =NOCOR<sup>2</sup>, =N-CN or =N-NO<sub>2</sub> and may furthermore be mono-substituted or disubstituted by Hal, A, CONH<sub>2</sub> or COOA.

10 T is, in particular, piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl or 3-methylimidazolidin-1-yl, each of which is monosubstituted or disubstituted by =NR<sup>2</sup>, =S, =NOR<sup>2</sup> or =N-CN.

15 T is furthermore particularly preferably, for example, piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted or disubstituted by =NR<sup>2</sup>, =S, =NOR<sup>2</sup> or =N-CN,  
20 where the thiadiazole ring may additionally be substituted by A, CONH<sub>2</sub> or COOA,

very particularly preferably 2-iminopiperidin-1-yl and the corresponding hydroxylamino, thioxo and =N-(CH<sub>2</sub>)<sub>1-3</sub>NA'<sub>2</sub> derivatives, where A' is alkyl  
25 having 1, 2, 3, 4, 5 or 6 carbon atoms.

In a further embodiment, T is 2-iminopiperidin-1-yl, 2-iminopyrrolidin-1-yl, 2-hydroxyiminopiperidin-1-yl, 2-hydroxyiminopyrrolidin-1-yl, 2-methoxy-  
30 iminopiperidin-1-yl, 2-methoxyiminopyrrolidin-1-yl, 2-cyaniminoimidazolidin-1-yl or 2-cyanimino-3-methylimidazolidin-1-yl.

D is in particular, for example, phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, hydroxyl, methoxy, ethoxy, hydroxy-  
35

- 14 -

carbonyl, methoxycarbonyl or ethoxycarbonyl, or is pyridyl or thienyl, each of which is unsubstituted or monosubstituted by Hal.

D is very particularly preferably 4-chlorophenyl or 3-chloro-2-pyridyl.

5  $R^1$  is preferably, for example, H, phenyl or thienyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal and/or A, or alkyl having 1-6 carbon atoms, which may be substituted by thiophene, imidazole, indole,  $S(O)_mR^2$ ,  $OR^2$ , cycloalkyl or phenyl,  
10 where the imidazole ring can have a substituent A,  $R^1$  in a further preferred embodiment is alternatively Het', where Het' is preferably a 5- or 6-membered heterocyclic ring having one or two N, O or S atoms, for example 2-thienyl or 2-furyl.

15 Specifically,  $R^1$  is preferably H, phenyl, 2-fluorophenyl, 2-chlorophenyl, alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, thienyl, methoxymethyl, (1-methyl-1*H*-imidazolyl-4-yl)methyl or methylsulfonylethyl.

20  $R^2$  is preferably, for example, H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.

$n$  is preferably 0 or 1.

$m$  is preferably 2.

25 Y is preferably, for example, 1,4-phenylene which is unsubstituted or monosubstituted by alkyl.

W is preferably absent.

30 The compounds of the formula I can have one or more centres of chirality and can therefore occur in various stereoisomeric forms. The formula I covers all these forms.

35 Accordingly, the invention relates, in particular, to compounds of the formula I in which at least one of the said radicals has one of the preferred

meanings indicated above. Some preferred groups of compounds can be expressed by the following sub-formulae Ia to Iq, which conform to the formula I and in which the radicals not designated in greater detail are as defined for the formula I, but in which

5

in Ia        D    is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OR<sup>2</sup> or COOR<sup>2</sup> or is pyridyl or thienyl, each of which is unsubstituted or monosubstituted by Hal;

10

in Ib        R<sup>2</sup>    is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms;

15

in Ic        R<sup>1</sup>    is H, phenyl or thienyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal and/or A, or is alkyl having 1-6 carbon atoms, which may be substituted by thiophene, imidazole, indole, S(O)<sub>m</sub>R<sup>2</sup>, OR<sup>2</sup>, cycloalkyl or phenyl, where the imidazole ring may have a substituent A;

20

in Id        X    is NH or O;

25

in Ie        W    is (CH<sub>2</sub>)<sub>n</sub>;

in If        Y    is Ar-diyl or Het-diyl;

30

in Ig        T    is a monocyclic or bicyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having from 1 to 3 N, O and/or S atoms which is monosubstituted or disubstituted by =S, =NR<sup>2</sup>, =NOR<sup>2</sup>, =NCOR<sup>2</sup>, =NCOOR<sup>2</sup>, =NOCOR<sup>2</sup>, =N-CN or =N-NO<sub>2</sub> and may furthermore be monosubstituted or disubstituted by Hal, A, CONH<sub>2</sub> or COOA;

35

- 16 -

- 5            in lh        T        is a monocyclic or bicyclic saturated or unsaturated hetero-  
cyclic ring having 1 or 2 N and/or O atoms which is  
monosubstituted or disubstituted by  $=NR^2$ ,  $=S$ ,  $=NOR^2$ ,  
 $=N-CN$  or  $=N-NO_2$ ;
- 10           in li        T        is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-  
4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl,  
azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-  
yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is  
monosubstituted or disubstituted by  $=NR^2$ ,  $=S$ ,  $=NOR^2$  or  
 $=N-CN$ ,  
15           where the thiadiazole ring may additionally be substituted  
by A,  $CONH_2$  or  $COOA$ ,  
A'        is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, in which 1-7  
H atoms may be replaced by F;
- 20           in lj        T        is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-  
4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl,  
azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-  
yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is  
25           monosubstituted or disubstituted by  $=NR^{2b}$ ,  $=S$ ,  $=NOR^{2b}$  or  
 $=N-CN$ ,  
where the thiadiazole ring may additionally be substituted  
by A,  $CONH_2$  or  $COOA$ ,  
30            $R^{2b}$     is H, OH or  $OA'$ ,  
A'        is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, in which 1-7  
H atoms may be replaced by F;
- 35           in lk        T        is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-  
4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl,

- 17 -

- azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl or 3-A''-imidazolidin-1-yl, each of which is monosubstituted or disubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN, R<sup>2b</sup> is H, OH or OA'',
- 5 A'' is methyl, ethyl, propyl, isopropyl or butyl;
- in II Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OA, SO<sub>2</sub>A, COOR<sup>2</sup>, SO<sub>2</sub>NH<sub>2</sub> or
- 10 CN;
- in Im D is phenyl which is unsubstituted or monosubstituted by Hal or is pyridyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,
- 15 R<sup>1</sup> is H, phenyl or thienyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal and/or A, or is alkyl having 1-6 carbon atoms, which may be substituted by thiophene, imidazole, indole, S(O)<sub>m</sub>R<sup>2</sup>, OR<sup>2</sup>,
- 20 cycloalkyl or phenyl, where the imidazole ring may have a substituent A,
- X is NH or O,
- W is absent,
- 25 Y is Ar-diyl, pyridinediyl or piperidinediyl,
- Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OA, SO<sub>2</sub>A, COOR<sup>2</sup>, SO<sub>2</sub>NH<sub>2</sub> or CN,
- 30 T is piperidin-1-yl, pyrrolidin-1-yl, 1H-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2H-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN,
- 35 where the thiadiazole ring may additionally be substituted by A, CONH<sub>2</sub> or COOA,

- 18 -

$R^{2b}$  is H, OH or  $OA''$ ,

$A''$  is methyl, ethyl, propyl, isopropyl or butyl;

5            in In        D    is phenyl which is unsubstituted or monosubstituted by Hal  
or is pyridyl or thienyl, each of which is unsubstituted or  
monosubstituted by Hal,

10                     $R^1$     is H, phenyl or thienyl, each of which is unsubstituted or  
monosubstituted or disubstituted by Hal and/or A,  
or is alkyl having 1-6 carbon atoms, which may be substi-  
tuted by thiophene, imidazole, indole,  $S(O)_mR^2$ ,  $OR^2$ ,  
cycloalkyl or phenyl,

where the imidazole ring may have a substituent A,

15                    X    is NH or O,

W    is absent,

Y    is Ar-diyl,

20                    Ar    is phenyl which is unsubstituted or monosubstituted or  
disubstituted by Hal or A,

25                    T    is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-  
4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl,  
azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-  
yl, 3- $A'$ -imidazolidin-1-yl or thiadiazolyl, each of which is  
monosubstituted by  $=NR^{2b}$ ,  $=S$ ,  $=NOR^{2b}$  or  $=N-CN$ ,

where the thiadiazole ring may additionally be substituted  
by A,  $CONH_2$  or  $COOA$ ,

30                     $R^{2b}$  is H, OH or  $OA''$ ,

$A''$  is methyl, ethyl, propyl, isopropyl or butyl;

                      in Io        D    is phenyl or thienyl, each of which is unsubstituted or  
monosubstituted by Hal,

35                     $R^1$     is H, phenyl which is unsubstituted or monosubstituted by  
Hal, or alkyl having 1-6 carbon atoms,

X    is NH or O,

- 19 -

- W is absent,  
Y is Ar-diyl,  
Ar is phenyl which is unsubstituted or monosubstituted or  
disubstituted by Hal or A,  
5 T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-  
4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl,  
azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-  
yl or 3-A''-imidazolidin-1-yl, each of which is monosubsti-  
10 tuted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN,  
R<sup>2b</sup> is H, OH or OA'',  
A'' is methyl, ethyl, propyl, isopropyl or butyl;
- 15 in Ip D is phenyl or thienyl, each of which is unsubstituted or  
monosubstituted by Hal,  
R<sup>1</sup> is H, phenyl which is unsubstituted or monosubstituted by  
Hal, alkyl having 1-6 carbon atoms or Het',  
X is NH or O,  
20 W is absent,  
Y is Ar-diyl,  
Ar is phenyl which is unsubstituted or monosubstituted or  
disubstituted by Hal or A,  
25 T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-  
4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl,  
azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-  
yl or 3-A''-imidazolidin-1-yl, each of which is monosubsti-  
30 tuted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN,  
R<sup>2b</sup> is H, OH or OA'',  
A'' is methyl, ethyl, propyl, isopropyl or butyl,  
Het' is 2-thienyl or 2-furyl;
- 35

26474-891

- 20 -

in Iq D is phenyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,

R<sup>1</sup> is H, phenyl which is unsubstituted or monosubstituted or disubstituted by Hal and/or A, alkyl having 1-6 carbon atoms or Het',

or alkyl having 1-6 carbon atoms which may be substituted by thiophene, imidazole, indole, S(O)<sub>m</sub>A", OA", cycloalkyl or phenyl,

where the imidazole ring may have a substituent A",

X is NH or O,

W is absent,

Y is Ar-diyl,

Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal or A,

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN, where the thiadiazole ring may additionally be substituted by A", CONH<sub>2</sub> or COOA",

R<sup>2b</sup> is H, OH or OA",

A" is methyl, ethyl, propyl, isopropyl or butyl,

Het' is 2-thienyl or 2-furyl,

and pharmaceutically usable derivatives, solvates and stereoisomers thereof, including mixtures thereof in all ratios.

26474-891

20a

In an embodiment of the present invention, there is provided the compound as described herein, which is

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)phenyl]-2-phenylacetamide,

5 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-imino-piperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

10 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-(2-chlorophenyl)acetamide,

15 (S)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]propionamide,

20 (R)-2-[N-(5-chloropyridin-2-yl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

25 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-imino-2*H*-pyridin-1-yl)phenyl]-2-phenylacetamide,

26474-891

20b

(R)-2-[N-(5-chlorothiophen-2-yl)carbamoyloxy]-N-[4-(2-imino-piperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(3-iminomorpholin-4-yl)phenyl]-2-phenylacetamide,

5 2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-(2-chlorophenyl)acetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

10 (S)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-pyrrolidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-(2-chlorophenyl)acetamide,

15 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-azabicyclo-[2.2.2]octan-3-imino-2-yl)phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

20 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide, hydrochloride,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

25 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]propionamide,

26474-891

20c

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide,

10 (R)-2-[3-(5-chlorothiophen-2-yl)ureido]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-methoxyiminopyrrolidin-1-yl)phenyl]-2-phenylacetamide,

15 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-thienylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-phenylacetamide,

20 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-(thiophen-2-yl)acetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]valeramide,

25 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]-2-phenylacetamide,

26474-891

20d

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]-2-thiophen-2-ylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-(2-thienyl)acetamide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-(2-thienyl)acetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

10 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]valeramide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide, citrate,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, tartrate,

15 2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-pyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

20 2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, lactate,

(R)-2-[3-(4-chloro-2-methylphenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-thienyl)acetamide,

25 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

26474-891

20e

(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-phenylacetamide,

(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]-2-(2-thienyl)acetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-imino-5-methyl-1,3,4-thiadiazol-3-yl)phenyl]-2-phenylacetamide,

10 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-(2-thienyl)acetamide,

15 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]valeramide,

20 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]valeramide,

2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]acetamide,

25 2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]acetamide,

26474-891

20f

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]acetamide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)phenyl]propionamide,

5           (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]propionamide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]propionamide,

10           (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)phenyl]-3-methylbutyramide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]-3-methylbutyramide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-3-methylbutyramide,

15           (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)phenyl]-4-methylvaleramide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]-4-methylvaleramide,

20           (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-4-methylvaleramide,

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-3-methoxypropionamide,

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]-3-methoxypropionamide,

25           (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-3-(1-methyl-1*H*-imidazol-4-yl)propionamide,

26474-891

20g

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(5-ethyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide,

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(5-aminocarbonyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide,

5                    2-[3-(4-chlorophenyl)ureido]-*N*-[4-(5-ethoxycarbonyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide, or

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]-4-methylsulfonylbutyramide,

or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any  
10    mixture thereof in all ratios.

The compounds of the formula I and also the starting materials for the preparation are, in addition, prepared by methods known per se, as described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic

- 21 -

Chemistry], Georg-Thieme-Verlag, Stuttgart), to be precise under reaction conditions which are known and suitable for the said reactions. Use can also be made here of variants which are known per se, but are not mentioned here in greater detail.

5

If desired, the starting materials can also be formed in situ so that they are not isolated from the reaction mixture, but instead are immediately converted further into the compounds of the formula I.


10

The starting compounds of the formulae II, III, IV and V are generally known. If they are novel, they can, however, be prepared by methods known per se.

15

**All** compounds of the following formula VI (where R = H or methyl; n = 3, 4 or 5) can be synthesised in accordance with the following scheme:


20



25

For example, synthesis of 1-(4-amino-2-methylphenyl)piperidine-2-thione:


30




35

- 22 -

Alternative synthesis:



Synthesis of the phenylpiperidinethione unit without a methyl group:



25 Compounds of the formula I can preferably be obtained by reacting compounds of the formula II with compounds of the formula III.

30 The reaction is generally carried out in an inert solvent, in the presence of an acid-binding agent, preferably an alkali or alkaline earth metal hydroxide, carbonate or bicarbonate or another salt of a weak acid of the alkali or alkaline earth metals, preferably of potassium, sodium, calcium or caesium. The addition of an organic base, such as triethylamine, dimethylaniline, pyridine or quinoline, or of an excess of the phenol component of

35 the formula II or of the alkylation derivative of the formula III, may also be

favourable. Depending on the conditions used, the reaction time is between a few minutes and 14 days, and the reaction temperature is between about 0° and 150°, normally between 20° and 130°.

5 Examples of suitable inert solvents are hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; alcohols, such as methanol, ethanol, iso-  
10 propanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide or dimethylformamide (DMF);  
15 nitriles, such as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); carbon disulfide; carboxylic acids, such as formic acid or acetic acid; nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate, or mixtures of the said solvents.

20

Compounds of the formula I can furthermore be obtained by reacting compounds of the formula IV with compounds of the formula V.

25 The reaction is generally carried out in an inert solvent and under conditions as indicated above.

In the compounds of the formula V, L is preferably Cl, Br, I or a free or reactively modified OH group, such as, for example, an activated ester, an imidazolidine or alkylsulfonyloxy having 1-6 carbon atoms (preferably  
30 methylsulfonyloxy or trifluoromethylsulfonyloxy) or arylsulfonyloxy having 6-10 carbon atoms (preferably phenyl- or p-tolylsulfonyloxy).

Radicals of this type for activation of the carboxyl group in typical acylation reactions are described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie  
35 [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart).

- 24 -

Activated esters are advantageously formed in situ, for example through addition of HOBt or N-hydroxysuccinimide.

5 The reaction is generally carried out in an inert solvent, in the presence of an acid-binding agent, preferably an organic base, such as DIPEA, triethylamine, dimethylaniline, pyridine or quinoline, or an excess of the carboxyl component of the formula V.

10 It may also be favourable to add an alkali or alkaline earth metal hydroxide, carbonate or bicarbonate or another salt of a weak acid of the alkali or alkaline earth metals, preferably of potassium, sodium, calcium or caesium.

15 Depending on the conditions used, the reaction time is between a few minutes and 14 days, and the reaction temperature is between about -30° and 140°, normally between -10° and 90°, in particular between about 0° and about 70°.

Suitable inert solvents are those mentioned above.

20 Compounds of the formula I can furthermore be obtained by liberating compounds of the formula I from one of their functional derivatives by treatment with a solvolysing or hydrogenolysing agent.

25 Preferred starting materials for the solvolysis or hydrogenolysis are those which conform to the formula I, but contain corresponding protected amino and/or hydroxyl groups instead of one or more free amino and/or hydroxyl groups, preferably those which carry an amino-protecting group instead of an H atom bonded to an N atom, in particular those which carry an R'-N group, in which R' is an amino-protecting group, instead of an HN group, and/or those which carry an hydroxyl-protecting group instead of the H atom of an hydroxyl group, for example those which conform to the formula I, but carry a -COOR" group, in which R" is an hydroxyl-protecting group, instead of a -COOH group.

30

35

- 25 -

Preferred starting materials are also the oxadiazole derivatives, which can be converted into the amidino compounds.

5 It is also possible for a plurality of – identical or different – protected amino and/or hydroxyl groups to be present in the molecule of the starting material. If the protecting groups present are different from one another, they can in many cases be cleaved off selectively.

10 The term "amino-protecting group" is known in general terms and relates to groups which are suitable for protecting (blocking) an amino group against chemical reactions, but which are easy to remove after the desired chemical reaction has been carried out elsewhere in the molecule. Typical  
15 of such groups are, in particular, unsubstituted or substituted acyl, aryl, aralkoxymethyl or aralkyl groups. Since the amino-protecting groups are removed after the desired reaction (or reaction sequence), their type and size is furthermore not crucial; however, preference is given to those having 1-20, in particular 1-8, carbon atoms. The term "acyl group" is to be  
20 understood in the broadest sense in connection with the present process. It includes acyl groups derived from aliphatic, araliphatic, aromatic or heterocyclic carboxylic acids or sulfonic acids, and, in particular, alkoxy-carbonyl, aryloxy-carbonyl and especially aralkoxycarbonyl groups. Examples  
25 of such acyl groups are alkanoyl, such as acetyl, propionyl and butyryl; aralkanoyl, such as phenylacetyl; aroyl, such as benzoyl and tolyl; aryloxyalkanoyl, such as POA; alkoxy-carbonyl, such as methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, BOC (tert-butoxycarbonyl) and 2-iodoethoxycarbonyl; aralkoxycarbonyl, such as CBZ ("carbo-  
30 benzoxy"), 4-methoxybenzyloxycarbonyl and FMOC; and arylsulfonyl, such as Mtr. Preferred amino-protecting groups are BOC and Mtr, furthermore CBZ, Fmoc, benzyl and acetyl.

35 The term "hydroxyl-protecting group" is likewise known in general terms and relates to groups which are suitable for protecting a hydroxyl group

- 26 -

against chemical reactions, but are easily removable after the desired chemical reaction has been carried out elsewhere in the molecule. Typical of such groups are the above-mentioned unsubstituted or substituted aryl, aralkyl or acyl groups, furthermore also alkyl groups. The nature and size  
5 of the hydroxyl-protecting groups are not crucial since they are removed again after the desired chemical reaction or reaction sequence; preference is given to groups having 1-20, in particular 1-10, carbon atoms. Examples of hydroxyl-protecting groups are, inter alia, benzyl, 4-methoxybenzyl,  
10 p-nitrobenzoyl, p-toluenesulfonyl, tert-butyl and acetyl, where benzyl and tert-butyl are particularly preferred.

The compounds of the formula I are liberated from their functional derivatives – depending on the protecting group used – for example using strong  
15 acids, advantageously using TFA or perchloric acid, but also using other strong inorganic acids, such as hydrochloric acid or sulfuric acid, strong organic carboxylic acids, such as trichloroacetic acid, or sulfonic acids, such as benzene- or p-toluenesulfonic acid. The presence of an additional  
20 inert solvent is possible, but is not always necessary. Suitable inert solvents are preferably organic, for example carboxylic acids, such as acetic acid, ethers, such as tetrahydrofuran or dioxane, amides, such as DMF, halogenated hydrocarbons, such as dichloromethane, furthermore also  
25 alcohols, such as methanol, ethanol or isopropanol, and water. Mixtures of the above-mentioned solvents are furthermore suitable. TFA is preferably used in excess without addition of a further solvent, and perchloric acid is preferably used in the form of a mixture of acetic acid and 70% perchloric  
30 acid in the ratio 9:1. The reaction temperatures for the cleavage are advantageously between about 0 and about 50°, preferably between 15 and 30° (room temperature).

35 The BOC, OBut and Mtr groups can, for example, preferably be cleaved off using TFA in dichloromethane or using approximately 3 to 5N HCl in dioxane at 15-30°, and the FMOC group can be cleaved off using an

- 27 -

approximately 5 to 50% solution of dimethylamine, diethylamine or piperidine in DMF at 15-30°.

5 Protecting groups which can be removed hydrogenolytically (for example CBZ, benzyl or the liberation of the amidino group from its oxadiazole derivative) can be cleaved off, for example, by treatment with hydrogen in the presence of a catalyst (for example a noble-metal catalyst, such as palladium, advantageously on a support, such as carbon). Suitable sol-  
10 vents here are those indicated above, in particular, for example, alcohols, such as methanol or ethanol, or amides, such as DMF. The hydrogenolysis is generally carried out at temperatures between about 0 and 100° and pressures between about 1 and 200 bar, preferably at 20-30° and 1-10  
15 bar. Hydrogenolysis of the CBZ group succeeds well, for example, on 5 to 10% Pd/C in methanol or using ammonium formate (instead of hydrogen) on Pd/C in methanol/DMF at 20-30°.

20 Examples of suitable inert solvents are hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, trifluoromethylbenzene, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol;  
25 ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide,  
30 N-methylpyrrolidone (NMP) or dimethylformamide (DMF); nitriles, such as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); carbon disulfide; carboxylic acids, such as formic acid or acetic acid; nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate, or mixtures of the said solvents.

35

- 28 -

Esters can be saponified, for example, using acetic acid or using NaOH or KOH in water, water/THF or water/dioxane, at temperatures between 0 and 100°.

5 Free amino groups can furthermore be acylated in a conventional manner using an acid chloride or anhydride or alkylated using an unsubstituted or substituted alkyl halide or reacted with  $\text{CH}_3\text{-C(=NH)-OEt}$ , advantageously in an inert solvent, such as dichloromethane or THF and/or in the presence  
10 of a base, such as triethylamine or pyridine, at temperatures between -60 and +30°.

A base of the formula I can be converted into the associated acid-addition  
15 salt using an acid, for example by reaction of equivalent amounts of the base and the acid in an inert solvent, such as ethanol, followed by evaporation. Suitable acids for this reaction are, in particular, those which give physiologically acceptable salts. Thus, it is possible to use inorganic acids, for example sulfuric acid, nitric acid, hydrohalic acids, such as hydrochloric  
20 acid or hydrobromic acid, phosphoric acids, such as orthophosphoric acid, or sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic monobasic or polybasic carboxylic, sulfonic or sulfuric acids, for example formic acid, acetic acid, propionic  
25 acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, ethanedisulfonic acid, 2-hydroxyethanesulfonic  
30 acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenemono- and -disulfonic acids, and laurylsulfuric acid. Salts with physiologically unacceptable acids, for example picrates, can be used for the isolation and/or purification of the compounds of the formula I.

35 On the other hand, compounds of the formula I can be converted into the corresponding metal salts, in particular alkali metal or alkaline earth metal

- 29 -

salts, or into the corresponding ammonium salts using bases (for example sodium hydroxide, potassium hydroxide, sodium carbonate or potassium carbonate). It is also possible to use physiologically acceptable organic bases, such as, for example, ethanolamine.

5

Compounds of the formula I according to the invention may be chiral owing to their molecular structure and may accordingly occur in various enantiomeric forms. They can therefore exist in racemic or in optically active form.

10

Since the pharmaceutical activity of the racemates or stereoisomers of the compounds according to the invention may differ, it may be desirable to use the enantiomers. In these cases, the end product or even the intermediates can be separated into enantiomeric compounds by chemical or physical measures known to the person skilled in the art or even employed as such in the synthesis.

15

In the case of racemic amines, diastereomers are formed from the mixture by reaction with an optically active resolving agent. Examples of suitable resolving agents are optically active acids, such as the R and S forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid, suitable N-protected amino acids (for example N-benzoylproline) or N-benzenesulfonylproline), or the various optically active camphorsulfonic acids. Also advantage is chromatographic enantiomer resolution with the aid of an optically active resolving agent (for example dinitrobenzoylphenylglycine, cellulose triacetate or other derivatives of carbohydrates or chirally derivatised methacrylate polymers immobilised on silica gel). Examples of suitable eluents for this purpose are aqueous or alcoholic solvent mixtures, such as, for example, hexane/isopropanol/acetonitrile, for example in the ratio 82:15:3.

25

30

35

The invention furthermore relates to the use of compounds of the formula I and/or their physiologically acceptable salts for the preparation of a medi-

- 30 -

cament (pharmaceutical preparation), in particular by non-chemical methods. They can be converted here into a suitable dosage form together with at least one solid, liquid and/or semiliquid excipient or assistant and, if desired, in combination with one or more further active ingredients.

5

The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or its pharmaceutically usable derivatives, solvates and stereoisomers, including mixtures thereof in all ratios, and optionally excipients and/or assistants.

10

These preparations can be used as medicaments in human or veterinary medicine. Suitable excipients are organic or inorganic substances which are suitable for enteral (for example oral), parenteral or topical administration and do not react with the novel compounds, for example water, vegetable oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glycerol triacetate, gelatin, carbohydrates, such as lactose or starch, magnesium stearate, talc or vaseline. Suitable for oral administration are, in particular, tablets, pills, coated tablets, capsules, powders, granules, syrups, juices or drops, suitable for rectal administration are suppositories, suitable for parenteral administration are solutions, preferably oil-based or aqueous solutions, furthermore suspensions, emulsions or implants, and suitable for topical application are ointments, creams or powders or also as nasal sprays. The novel compounds may also be lyophilised and the resultant lyophilisates used, for example, to prepare injection preparations. The preparations indicated may be sterilised and/or comprise assistants, such as lubricants, preservatives, stabilisers and/or wetting agents, emulsifying agents, salts for modifying the osmotic pressure, buffer substances, colorants and flavours and/or a plurality of further active ingredients, for example one or more vitamins.

15

20

25

30

35

The compounds of the formula I and their physiologically acceptable salts can be used for combating thromboembolic diseases, such as thrombosis,

myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases.

5 In general, the substances according to the invention are preferably administered in doses between about 1 and 500 mg, in particular between 5 and 100 mg, per dosage unit. The daily dose is preferably between about 0.02 and 10 mg/kg of body weight. However, the specific dose for  
10 each patient depends on a wide variety of factors, for example on the efficacy of the specific compound employed, on the age, body weight, general state of health, sex, on the diet, on the time and method of administration, on the excretion rate, medicament combination and severity of the particular disease to which the therapy applies. Oral administration is preferred.  
15

The invention furthermore relates to medicaments comprising at least one compound of the formula I and/or its pharmaceutically usable derivatives,  
20 solvates and stereoisomers, including mixtures thereof in all ratios, and at least one further medicament active ingredient.

The invention also relates to a set (kit) consisting of separate packs of  
25 (a) an effective amount of a compound of the formula I and/or its pharmaceutically usable derivatives, solvates and stereoisomers, including mixtures thereof in all ratios, and  
(b) 30 an effective amount of a further medicament active ingredient.

The set comprises suitable containers, such as boxes, individual bottles, bags or ampoules. The set may, for example, comprise separate ampoules each containing an effective amount of a compound of the  
35 formula I and/or its pharmaceutically usable derivatives, solvates and stereoisomers, including mixtures thereof in all ratios,

and an effective amount of a further medicament active ingredient in dissolved or lyophilised form.

5 The invention furthermore relates to the use of compounds of the formula I and/or their pharmaceutically usable derivatives, solvates and stereoisomers, including mixtures thereof in all ratios,  
for the preparation of a medicament for the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris,  
10 ris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases,  
in combination with at least one further medicament active ingredient.

15 Above and below, all temperatures are given in °C. In the following examples, 'conventional work-up' means that water is added if necessary, the pH is adjusted, if necessary, to between 2 and 10, depending on the constitution of the end product, the mixture is extracted with ethyl acetate or dichloromethane, the phases are separated, the organic phase is dried  
20 over sodium sulfate and evaporated, and the product is purified by chromatography on silica gel and/or by crystallisation. Rf values on silica gel; eluent: ethyl acetate/methanol 9:1.

Mass spectrometry (MS): EI (electron impact ionisation)  $M^+$   
25 FAB (fast atom bombardment)  $(M+H)^+$   
ESI (electrospray ionisation)  $(M+H)^+$  (unless stated otherwise)

30

35


**Example 1***Preparation of amine units:*

a)

5

10

15



20

25

30

35

15 g (78.8 mmol) of 1-(4-aminophenyl)piperidin-2-one are heated to the boil together with 16.0 g (39.5 mmol) of 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane 2,4-disulfide (Lawesson's reagent) in 100 ml of anhydrous toluene. After 45 minutes, the solvent is evaporated, and the residue is taken up in dichloromethane and 2 N HCl. The aqueous phase is extracted three times with dichloromethane and adjusted to a pH of 12 using conc. NaOH. Extraction with dichloromethane, drying over sodium sulfate and evaporation of the solvent give 1-(4-aminophenyl)piperidine-2-thione as a colourless solid, ESI 207.


1.25 ml (20.0 mmol) of iodomethane are added to a solution of 3.74 g (18.1 mmol) of 1-(4-aminophenyl)piperidine-2-thione in 30 ml of acetone, and the mixture is stirred at room temperature for 48 hours. The reaction mixture is evaporated, giving 1-(4-aminophenyl)-6-methylsulfanyl-2,3,4,5-

- 34 -

tetrahydropyridinium iodide as a brownish solid; ESI 221.

3.5 ml (25 mmol) of triethylamine are added to a solution of 2.68 g (12.1 mmol) of 1-(4-aminophenyl)-6-methylsulfanyl-2,3,4,5-tetrahydropyridinium iodide and 1.01 g (12.1 mmol) of O-methylhydroxylammonium chloride in 30 ml of ethanol, and the mixture is stirred at room temperature for 20 hours. The reaction mixture is evaporated and taken up in water, and the resultant precipitate is filtered off, giving 1-(4-aminophenyl)-piperidin-2-one O-methyl oxime as a colourless solid; ESI 220.

b)



A mixture of 40.0 g (290 mmol) of 4-nitroaniline and 40.0 ml (421 mmol) of 4-chlorobutyronitrile is heated at the boil (about 190°C) for 30 minutes using a hot-air fan. The mixture is allowed to cool and is digested with ethyl acetate, giving 1-(4-nitrophenyl)pyrrolidin-2-ylideneamine, hydrochloride, as a brown solid, ESI 206.


2.5 g of water-moist palladium on activated carbon (5% of Pd) are added to a solution of 25.0 g (103 mmol) of 1-(4-nitrophenyl)pyrrolidin-2-ylideneamine, hydrochloride, in 500 ml of methanol, and the mixture is hydrogenated at room temperature for 17 hours. The catalyst is filtered off,

- 35 -

and the filtrate is evaporated, giving 1-(4-aminophenyl)pyrrolidin-2-ylideneamine, hydrochloride, as a brownish solid; ESI 176.

2.68 g (12.7 mmol) of 1-(4-aminophenyl)pyrrolidin-2-ylideneamine, hydrochloride, are dissolved in 70 ml of methanol, 9.49 g (113 mmol) of sodium hydrogencarbonate and 4.06 g (48.6 mmol) of methoxylamine, hydrochloride, are added, and the mixture is heated at 70°C for 48 hours. The reaction mixture is evaporated, and the residue is taken up in acetone. The insoluble components are filtered off, and the filtrate is evaporated. The residue is chromatographed on a silica-gel column with ethyl acetate as eluent, giving 1-(4-aminophenyl)pyrrolidin-2-one O-methyl oxime as a brownish solid; ESI 206.

c)



23.63 g (150 mmol) of 4-chloronitrobenzene are dissolved in 92.1 ml (1.38 mol) of ethylenediamine and heated to the boil under a nitrogen atmosphere. After 24 hours, the mixture is evaporated to dryness, and the crude product is suspended in 100 ml of 1 N sodium hydroxide

- 36 -

solution and filtered off. Washing with water and drying under reduced pressure gives 25.9 g of *N*<sup>1</sup>-(4-nitrophenyl)ethane-1,2-diamine in the form of yellow crystals (ESI 182).

5 5.9 g (32.56 mmol) of *N*<sup>1</sup>-(4-nitrophenyl)ethane-1,2-diamine are suspended in 200 ml of acetonitrile, and 3.72 g of dimethyl N-cyanimido-carbonate and 6.75 g of K<sub>2</sub>CO<sub>3</sub> are added. After the mixture has been stirred at room temperature for 2 hours, conventional work-up gives  
10 5.16 g of 1-(4-nitrophenyl)imidazolidin-2-ylidene cyanamide (ESI: 232).

3.47 g (15 mmol) of 1-(4-nitrophenyl)imidazolidin-2-ylidene cyanamide are hydrogenated in 35 ml of methanol using hydrogen on 5% palladium on carbon. Conventional work-up gives 2.88 g of 1-(4-aminophenyl)imidazolidin-2-ylidene cyanamide in the form of a pale-brown solid (ESI: 202).  
15

20


25

30

35

- 37 -

*Preparation of 2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide:*



1.1 400 mg (0.63 mmol) of dibutyltin dilaurate are added to a solution  
 25 of 10.0 g (65.7 mmol) of (R)-mandelic acid and 10.0 g (65.1 mmol) of  
 4-chlorophenyl isocyanate in 100 ml of dichloromethane, and the mixture  
 is stirred at room temperature for 24 hours. The reaction mixture is intro-  
 30 duced into water and extracted with ethyl acetate. The organic phase is  
 evaporated, giving (R)-(4-chlorophenylcarbamoyloxy)phenylacetic acid as  
 a colourless solid; ESI 306.

1.2 101 mg (1.00 mmol) of 4-methylmorpholine are added to a solution  
 35 of 183 mg (0.600 mmol) of (R)-(4-chlorophenylcarbamoyloxy)phenylacetic  
 acid, 129 mg (0.588 mmol) of 1-(4-aminophenyl)piperidin-2-one O-methyl  
 oxime, 192 mg (1.00 mmol) of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbo-

- 38 -

diimide hydrochloride (DAPECI) and 91.9 mg (0.600 mmol) of hydroxy-benzotriazole hydrate (HOBt) in 1 ml of DMF, and the mixture is stirred at room temperature for 24 hours. The reaction mixture is introduced into saturated sodium hydrogencarbonate solution, and the resultant precipitate is filtered off. The precipitate is chromatographed on a silica-gel column with petroleum ether/ethyl acetate as eluent, giving (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)phenyl]-2-phenylacetamide ("AA") as a colourless solid; ESI 507.

1.3 0.3 g of water-moist Raney nickel is added to a solution of 50 mg (0.099 mmol) of [4-(2-methoxyiminopiperidin-1-yl)phenylcarbamoyl]phenylmethyl (R)-(4-chlorophenyl)carbamate in 10 ml of methanol, and the mixture is hydrogenated. The reaction mixture is filtered, and the filtrate is evaporated. The residue is chromatographed on a silica-gel column with dichloromethane/methanol as eluent, giving (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide as a colourless solid; ESI 477.

1.4 30 mg (0.063 mmol) of [4-(2-iminopiperidin-1-yl)phenylcarbamoyl]phenylmethyl (R)-(4-chlorophenyl)carbamate are dissolved in 0.063 ml of 1 N HCl and lyophilised, giving (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamid, hydrochloride ("AB") as a colourless solid; ESI 477.

The following compounds are obtained analogously:

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide, hydrochloride, ESI 491;

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenylacetamide, hydrochloride, ESI 477;

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-(2-chlorophenyl)acetamide,

5

(S)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]propionamide,

10

(R)-2-[N-(5-chloropyridin-2-yl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-phenylacetamide,

15

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-imino-2*H*-pyridin-1-yl)phenyl]-2-phenylacetamide,

20

(R)-2-[N-(5-chlorothiophen-2-yl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(3-iminomorpholin-4-yl)-phenyl]-2-phenylacetamide,

25

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-(2-chlorophenyl)acetamide, ESI 555;

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-phenylacetamide, ESI 521;

30

(S)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-phenylacetamide, ESI 521;

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-pyrrolidin-1-yl)phenyl]-2-phenylacetamide, ESI 493;

35

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-(2-chlorophenyl)acetamide, ESI 555;

- 40 -

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-azabicyclo[2.2.2]-octan-3-imino-2-yl)phenyl]-2-phenylacetamide.

5

10

15

20


25

30

35

**Example 2**

*Preparation of (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide:*



2.1 A solution of 10.0 g (66.2 mmol) of (R)-alpha-aminophenylacetic acid in 600 ml of 0.25 M aqueous sodium hydrogencarbonate solution is heated to 80°C, and 20.2 g (132 mmol) of 4-chlorophenyl isocyanate are added. The reaction mixture is stirred at this temperature for 1 hour and cooled in an ice bath, and the resultant precipitate is filtered off. The filtrate

is acidified using 2 N HCl, and the resultant precipitate is filtered off, giving (R)-[3-(4-chlorophenyl)ureido]phenylacetic acid as a colourless solid; ESI 305.

5        2.2        152 mg (1.50 mmol) of 4-methylmorpholine are added to a solution  
of 366 mg (1.20 mmol) of (R)-[3-(4-chlorophenyl)ureido]phenylacetic acid,  
248 mg (1.20 mmol) of 1-(4-amino-2-methylphenyl)pyrrolidine-2-thione,  
249 mg (1.30 mmol) of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide  
10        hydrochloride (DAPECI) and 184 mg (1.20 mmol) of hydroxybenzotriazole  
hydrate (HOBt) in 1 ml of DMF, and the mixture is stirred at room tem-  
perature for 24 hours. The reaction mixture is introduced into saturated  
sodium hydrogencarbonate solution, and the resultant precipitate is filtered  
15        off, giving (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[3-methyl-4-(2-thioxo-  
pyrrolidin-1-yl)phenyl]-2-phenylacetamide as a colourless solid; ESI 493.

2.3        78.1 µl (1.25 mmol) of iodomethane are added to a solution of  
380 mg (0.771 mmol) of (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[3-methyl-4-(2-  
20        thioxopyrrolidin-1-yl)phenyl]-2-phenylacetamide in 10 ml of acetone, and  
the mixture is stirred at room temperature for 24 hours. The reaction mix-  
ture is evaporated, and the residue is taken up in tert-butyl methyl ether.  
The resultant precipitate is filtered off, giving (R)-1-(4-{2-[3-(4-chloro-  
25        phenyl)ureido]-2-phenylacetyl-amino}-2-methylphenyl)-5-methylsulfanyl-3,4-  
dihydro-2*H*-pyrrolium iodide as a colourless solid; ESI 507.

2.4        308 mg (4.00 mmol) of ammonium acetate are added to a solution  
of 469 mg (0.739 mmol) of (R)-1-(4-{2-[3-(4-chlorophenyl)ureido]-2-phenyl-  
30        acetyl-amino}-2-methylphenyl)-5-methylsulfanyl-3,4-dihydro-2*H*-pyrrolium  
iodide in 30 ml of ethanol, and the mixture is heated at the boil for 24  
hours. The reaction mixture is evaporated, and the residue is taken up in  
water. The resultant precipitate is filtered off, giving (R)-2-[3-(4-chloro-  
35        phenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenyl-

- 43 -

acetamide as a colourless solid; ESI 476.

2.5 170 mg (0.357 mmol) of (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-  
5 iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenylacetamide are dissolved in  
2 ml of 0.1 N HCl in 2-propanol, and tert-butyl methyl ether is added. The  
resultant precipitate is filtered off, giving (R)-2-[3-(4-chlorophenyl)ureido]-  
N-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenylacetamide, hydro-  
chloride ("BA") as a colourless solid; ESI 476.

10 The following compounds are obtained analogously:

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-3-  
15 methylphenyl]-2-phenylacetamide, hydrochloride, ESI 490;

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-  
phenyl]-2-phenylacetamide,

(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-  
20 phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-  
phenyl]propionamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-  
phenyl]-2-(2-fluorophenyl)acetamide,

25 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-  
phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-  
phenyl]-2-(2-fluorophenyl)acetamide,

30 (R)-2-[3-(5-chlorothiophen-2-yl)ureido]-N-[4-(2-iminopiperidin-1-yl)-  
phenyl]-2-phenylacetamide,

(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-methoxyiminopiperidin-1-  
yl)-3-methylphenyl]-2-phenylacetamide, ESI 520;


(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-methoxyiminopyrrolidin-1-  
35 yl)phenyl]-2-phenylacetamide, ESI 492.

**Example 3**


*Preparation of (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide:*

5

10



15



20

3.1 A solution of 366 mg (1.20 mmol) of (R)-[3-(4-chlorophenyl)-ureido]phenylacetic acid, 254 mg (1.20 mmol) of 1-(4-aminophenyl)pyrrolidin-2-ylideneamine hydrochloride, 230 mg (1.20 mmol) of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (DAPECI) and 184 mg (1.20 mmol) of hydroxybenzotriazole hydrate (HOBt) in 1 ml of DMF is stirred at room temperature for 24 hours. The reaction mixture is introduced into saturated sodium hydrogencarbonate solution, and the resultant precipitate is filtered off, giving (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide as a colourless solid; ESI 462.

25

30

35

3.2 260 mg (0.563 mmol) of (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide are dissolved in 5.6 ml of 0.1 N HCl in 2-propanol, and tert-butyl methyl ether is added. The resultant precipitate is filtered off, giving (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-

- 45 -

iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, hydrochloride, as a colourless solid; ESI 462.

The following compound is obtained analogously:

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-(thiophen-2-yl)acetamide, hydrochloride, ESI 468.

#### **Example 4**

Reaction of the corresponding carboxylic acids with the amine described in Example 1 c) analogously to Example 3 gives the following compounds:

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]-2-phenylacetamide, ESI 489;

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]-2-thiophen-2-ylacetamide ("CA"), ESI 495;

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]valeramide.

The following compounds are obtained analogously:

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]valeramide;

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]-2-phenylacetamide;

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]-2-thiophen-2-ylacetamide.

**Example 5**

The following compounds are obtained analogously to Example 2:

- 5 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-(2-thienyl)acetamide, ESI 482;  
(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-(2-thienyl)acetamide, trifluoroacetate, ESI 483;  
10 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-phenylacetamide, trifluoroacetate, ESI 477;  
(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]valeramide, trifluoroacetate, ESI 443.

15

**Example 6**

The following compounds are obtained analogously to Example 1:

- 20 2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide, citrate, ESI 481;  
(R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, tartrate, ESI 463;  
25 2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-methoxyimino-piperidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide, ESI 539;  
2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-methoxyimino-pyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide, ESI 511;  
30 2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide, citrate, ESI 495;  
(R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, lactate, ESI 463.

35

**Example 7**

The following compounds are obtained analogously to Example 3:

- 5 (R)-2-[3-(4-chloro-2-methylphenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-thienyl)acetamide, ESI 482;  
(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide, tartrate, ESI 476;
- 10 (S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-phenylacetamide, D-(-)-tartrate, ESI 462;  
(S)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide, D-(-)-tartrate, ESI 476;
- 15 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)-phenyl]-2-(2-thienyl)acetamide, trifluoroacetate, ESI 469;  
(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-imino-5-methyl-1,3,4-thiadiazol-3-yl)phenyl]-2-phenylacetamide, trifluoroacetate, ESI 493;
- 20 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)-phenyl]-2-phenylacetamide, trifluoroacetate, ESI 463;  
(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]-2-phenylacetamide, ESI 502;
- 25 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]-2-(2-thienyl)acetamide, ESI 508;  
(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]valeramide, ESI 454;
- 30 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]valeramide, ESI 468;  
(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)-phenyl]valeramide, trifluoroacetate, ESI 429;
- 35 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]valeramide, trifluoroacetate, ESI 428;

- 48 -

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-  
acetamide, trifluoroacetate, ESI 387;

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-  
imidazolidin-1-yl)phenyl]acetamide, trifluoroacetate, ESI 426;

5

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-  
phenyl]acetamide, trifluoroacetate, ESI 426;

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-  
phenyl]propionamide, trifluoroacetate, ESI 401;

10

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-  
imidazolidin-1-yl)phenyl]propionamide, trifluoroacetate, ESI 440;

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-  
phenyl]propionamide, trifluoroacetate, ESI 400;

15

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-  
phenyl]-3-methylbutyramide, trifluoroacetate, ESI 429;

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-  
imidazolidin-1-yl)phenyl]-3-methylbutyramide, trifluoroacetate, ESI 468;

20

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-  
phenyl]-3-methylbutyramide, trifluoroacetate, ESI 428;

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-  
phenyl]-4-methylvaleramide, trifluoroacetate, ESI 443;

25

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-  
imidazolidin-1-yl)phenyl]-4-methylvaleramide, trifluoroacetate, ESI 482;

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-  
phenyl]-4-methylvaleramide, trifluoroacetate, ESI 442;

30

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-3-  
methoxypropionamide, trifluoroacetate, ESI 430;

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyaniminoimidazolidin-1-yl)-  
phenyl]-3-methoxypropionamide, ESI 456;

35

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyaniminoimidazolidin-1-  
yl)phenyl]-3-(1-methyl-1*H*-imidazol-4-yl)propionamide, trifluoroacetate,  
ESI 506;

- 49 -

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(5-ethyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide, trifluoroacetate, ESI 475;

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(5-aminocarbonyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide, trifluoroacetate, ESI 490;

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(5-ethoxycarbonyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide, trifluoroacetate, ESI 519;

2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)phenyl]-4-methylsulfonylbutyramide.

#### Pharmacological data

##### Affinity to receptors

Table 1

| Compound No. | FXa-IC <sub>50</sub> [M] | TF/FVIIa-IC <sub>50</sub> [M] |
|--------------|--------------------------|-------------------------------|
| "AB"         | 5.8 x 10 <sup>-8</sup>   | 9.9 x 10 <sup>-8</sup>        |
| "BA"         | 6.8 x 10 <sup>-7</sup>   | 4.9 x 10 <sup>-7</sup>        |
| "BB"         | 2.7 x 10 <sup>-8</sup>   | 2.0 x 10 <sup>-6</sup>        |
| "AA"         | 2.2 x 10 <sup>-7</sup>   | 2.9 x 10 <sup>-7</sup>        |
| "CA"         | 5.4 x 10 <sup>-8</sup>   |                               |

The following examples relate to pharmaceutical preparations:

**Example A: Injection vials**

5

A solution of 100 g of an active ingredient of the formula I and 5 g of disodium hydrogenphosphate in 3 l of bidistilled water is adjusted to pH 6.5 using 2N hydrochloric acid, sterile filtered, transferred into injection vials, lyophilised under sterile conditions and sealed under sterile conditions. Each injection vial contains 5 mg of active ingredient.

10

**Example B: Suppositories**

15

A mixture of 20 g of an active ingredient of the formula I with 100 g of soya lecithin and 1400 g of cocoa butter is melted, poured into moulds and allowed to cool. Each suppository contains 20 mg of active ingredient.

20

**Example C: Solution**

25

A solution is prepared from 1 g of an active ingredient of the formula I, 9.38 g of  $\text{NaH}_2\text{PO}_4 \cdot 2 \text{H}_2\text{O}$ , 28.48 g of  $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$  and 0.1 g of benzalkonium chloride in 940 ml of bidistilled water. The pH is adjusted to 6.8, and the solution is made up to 1 l and sterilised by irradiation. This solution can be used in the form of eye drops.

30

**Example D: Ointment**

500 mg of an active ingredient of the formula I are mixed with 99.5 g of Vaseline under aseptic conditions.

35

**Example E: Tablets**

- 51 -

5 A mixture of 1 kg of active ingredient of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 kg of talc and 0.1 kg of magnesium stearate is pressed to give tablets in a conventional manner in such a way that each tablet contains 10 mg of active ingredient.

#### **Example F: Coated tablets**

10 Tablets are pressed analogously to Example E and subsequently coated in a conventional manner with a coating of sucrose, potato starch, talc, tragacanth and dye.

#### **Example G: Capsules**

15 2 kg of active ingredient of the formula I are introduced into hard gelatine capsules in a conventional manner in such a way that each capsule contains 20 mg of the active ingredient.

20

#### **Example H: Ampoules**

25 A solution of 1 kg of active ingredient of the formula I in 60 l of bidistilled water is sterile filtered, transferred into ampoules, lyophilised under sterile conditions and sealed under sterile conditions. Each ampoule contains 10 mg of active ingredient.

30


35

26474-891

52

CLAIMS:

1. A compound of formula I



in which

- 5 D is phenyl, pyridyl or thienyl, each of which is unsubstituted or monosubstituted or polysubstituted by Hal, A, OR<sup>2</sup>, N(R<sup>2</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, COOR<sup>2</sup> or CON(R<sup>2</sup>)<sub>2</sub>,

X is NR<sup>3</sup> or O,

R<sup>1</sup> is H, Ar, Het, cycloalkyl or A, which may be substituted by OR<sup>2</sup>, S(O)<sub>m</sub>R<sup>2</sup>, N(R<sup>2</sup>)<sub>2</sub>, Ar, Het, cycloalkyl, CN, COOR<sup>2</sup> or CON(R<sup>2</sup>)<sub>2</sub>,

- 10 R<sup>2</sup> is H, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Het, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub> or -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>3</sup>,

R<sup>3</sup> is H or A,

W is -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-,

Y is alkylene, cycloalkylene, Het-diyl or Ar-diyl,

- 15 T is a monocyclic or bicyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having from 1 to 4 N, O and/or S atoms which is monosubstituted or disubstituted by =S, =NR<sup>2</sup>, =NOR<sup>2</sup>, =NCOR<sup>2</sup>, =NCOOR<sup>2</sup>, =NOCOR<sup>2</sup>, =N-CN or =N-NO<sub>2</sub> and may furthermore be monosubstituted, disubstituted or trisubstituted by Hal, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Het, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, OR<sup>3</sup>, N(R<sup>3</sup>)<sub>2</sub>, NO<sub>2</sub>, CN,  
 20 COOR<sup>2</sup>, CON(R<sup>2</sup>)<sub>2</sub>, NR<sup>2</sup>COA, NR<sup>2</sup>CON(R<sup>2</sup>)<sub>2</sub>, NR<sup>2</sup>SO<sub>2</sub>A, COR<sup>2</sup>, SO<sub>2</sub>NR<sup>2</sup> and/or S(O)<sub>m</sub>A,

26474-891

53

A is unbranched or branched alkyl having 1-10 carbon atoms, in which one or two CH<sub>2</sub> groups may be replaced by O or S atoms and/or by -CH=CH- groups and/or in addition 1-7 H atoms may be replaced by F,

Ar is phenyl, naphthyl or biphenyl, each of which is unsubstituted or

- 5 monosubstituted, disubstituted or trisubstituted by Hal, A, OR<sup>3</sup>, N(R<sup>3</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, COOR<sup>3</sup>, CON(R<sup>3</sup>)<sub>2</sub>, NR<sup>3</sup>COA, NR<sup>3</sup>CON(R<sup>3</sup>)<sub>2</sub>, NR<sup>3</sup>SO<sub>2</sub>A, COR<sup>3</sup>, SO<sub>2</sub>N(R<sup>3</sup>)<sub>2</sub>, S(O)<sub>m</sub>A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-COOR<sup>2'</sup>, or -O-[C(R<sup>3</sup>)<sub>2</sub>]<sub>o</sub>-COOR<sup>2'</sup>,

R<sup>2'</sup> is H, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar', -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Het', -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub> or -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>3</sup>,

- 10 R<sup>2''</sup> is H, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar' or -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>3</sup>)<sub>2</sub> or -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>3</sup>,

Ar' is phenyl or benzyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal or A,

Het is a monocyclic or bicyclic saturated, unsaturated or aromatic heterocyclic ring

- 15 having from 1 to 4 N, O and/or S atoms, which may be unsubstituted or monosubstituted, disubstituted or trisubstituted by carbonyl oxygen, =S, =N(R<sup>3</sup>)<sub>2</sub>, Hal, A, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Ar, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-Het<sup>1</sup>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-cycloalkyl, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-OR<sup>2'</sup>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-N(R<sup>2'</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-COOR<sup>2'</sup>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-CON(R<sup>2'</sup>)<sub>2</sub>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-NR<sup>2'</sup>COA, NR<sup>2'</sup>CON(R<sup>2'</sup>)<sub>2</sub>, -[C(R<sup>3</sup>)<sub>2</sub>]<sub>n</sub>-NR<sup>2'</sup>SO<sub>2</sub>A, COR<sup>2'</sup>, SO<sub>2</sub>NR<sup>2'</sup> and/or
- 20 S(O)<sub>m</sub>A,

Het<sup>1</sup> is a monocyclic or bicyclic saturated, unsaturated or aromatic heterocyclic ring having 1 or 2 N, O and/or S atoms, which may be unsubstituted or

- monosubstituted or disubstituted by carbonyl oxygen, =S, =N(R<sup>3</sup>)<sub>2</sub>, Hal, A, OR<sup>2''</sup>, N(R<sup>2''</sup>)<sub>2</sub>, NO<sub>2</sub>, CN, COOR<sup>2''</sup>, CON(R<sup>2''</sup>)<sub>2</sub>, NR<sup>2''</sup>COA, NR<sup>2''</sup>CON(R<sup>2''</sup>)<sub>2</sub>, NR<sup>2''</sup>SO<sub>2</sub>A, COR<sup>2''</sup>, SO<sub>2</sub>NR<sup>2''</sup> and/or S(O)<sub>m</sub>A,
- 25

Hal is F, Cl, Br or I,

n is 0, 1 or 2,

m is 0, 1 or 2, and

26474-891

54

o is 1, 2 or 3,

or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios.

2. The compound according to Claim 1, or a pharmaceutically usable  
5 derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which D is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OR<sup>2</sup> or COOR<sup>2</sup> or is pyridyl or thienyl, each of which is unsubstituted or monosubstituted by Hal.
3. The compound according to Claim 1 or 2, or a pharmaceutically  
10 usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which R<sup>2</sup> is H or alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms.
4. The compound according to any one of Claims 1 to 3, or a  
pharmaceutically usable derivative, solvate or stereoisomer thereof, or any  
mixture thereof in all ratios, in which R<sup>1</sup> is H, phenyl or thienyl, each of which is  
15 unsubstituted or monosubstituted or disubstituted by Hal and/or A, or is alkyl having 1-6 carbon atoms, which may be substituted by thiophene, imidazole, indole, S(O)<sub>m</sub>R<sup>2</sup>, OR<sup>2</sup>, cycloalkyl or phenyl, where the imidazole ring may have a substituent A.
5. The compound according to any one of Claims 1 to 4, or a  
20 pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which X is NH or O.
6. The compound according to any one of Claims 1 to 5, or a  
pharmaceutically usable derivative, solvate or stereoisomer thereof, or any  
mixture thereof in all ratios, in which W is (CH<sub>2</sub>)<sub>n</sub>.
- 25 7. The compound according to any one of Claims 1 to 6, or a  
pharmaceutically usable derivative, solvate or stereoisomer thereof, or any  
mixture thereof in all ratios, in which Y is Ar-diyl or Het-diyl.
8. The compound according to any one of Claims 1 to 7, or a  
pharmaceutically usable derivative, solvate or stereoisomer thereof, or any

26474-891

55

mixture thereof in all ratios, in which T is a monocyclic or bicyclic saturated, unsaturated or aromatic carbocyclic or heterocyclic ring having from 1 to 3 N, O and/or S atoms which is monosubstituted or disubstituted by =S, =NR<sup>2</sup>, =NOR<sup>2</sup>, =NCOR<sup>2</sup>, =NCOOR<sup>2</sup>, =NOCOR<sup>2</sup>, =N-CN or =N-NO<sub>2</sub> and may furthermore be  
 5 monosubstituted or disubstituted by Hal, A, CONH<sub>2</sub> or COOA.

9. The compounds according to any one of Claims 1 to 8, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which T is a monocyclic or bicyclic saturated or unsaturated heterocyclic ring having 1 or 2 N and/or O atoms which is  
 10 monosubstituted or disubstituted by =NR<sup>2</sup>, =S, =NOR<sup>2</sup>, =N-CN or =N-NO<sub>2</sub>.

10. The compounds according to any of Claims 1 to 9, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which:

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H* pyridin-1-yl, morpholin-4-yl, piperazin-1-yl,  
 15 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted or disubstituted by =NR<sup>2</sup>, =S, =NOR<sup>2</sup> or =N-CN, where the thiadiazole ring may additionally be substituted by A, CONH<sub>2</sub> or COOA, and

A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, in which 1-7 H atoms may be  
 20 replaced by F.

11. The compound according to any one of Claims 1 to 10, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which:

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl,  
 25 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted or disubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN, where the thiadiazole ring may additionally be substituted by A, CONH<sub>2</sub> or COOA,

R<sup>2b</sup> is H, OH or OA', and

26474-891

56

A' is alkyl having 1, 2, 3, 4, 5 or 6 carbon atoms, in which 1-7 H atoms may be replaced by F.

12. The compound according to any one of Claims 1 to 11, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which:

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl or 3-A"-imidazolidin-1-yl, each of which is monosubstituted or disubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN,

- 10 R<sup>2b</sup> is H, OH or OA", and

A" is methyl, ethyl, propyl, isopropyl or butyl.

13. The compound according to any one of Claims 1 to 12, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OA, SO<sub>2</sub>A, SO<sub>2</sub>NH<sub>2</sub> or CN.

14. The compound according to any one of Claims 1 to 11, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which:

20 D is phenyl which is unsubstituted or monosubstituted by Hal or is pyridyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,

R<sup>1</sup> is H, phenyl or thienyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal and/or A,

or is alkyl having 1-6 carbon atoms, which may be substituted by thiophene, imidazole, indole, S(O)<sub>m</sub>R<sup>2</sup>, OR<sup>2</sup>, cycloalkyl or phenyl, where the imidazole ring may have a substituent A,

X is NH or O,

W is absent,

26474-891

57

Y is Ar-diyl, pyridinediyl or piperidinediyl,

Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal, A, OA, SO<sub>2</sub>A, SO<sub>2</sub>NH<sub>2</sub> or CN,

5 T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN, where the thiadiazole ring may additionally be substituted by A, CONH<sub>2</sub> or COOA,

R<sup>2b</sup> is H, OH or OA", and

10 A" is methyl, ethyl, propyl, isopropyl or butyl.

15. The compound according to any one of Claims 1 to 11, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which

15 D is phenyl which is unsubstituted or monosubstituted by Hal or is pyridyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,

R<sup>1</sup> is H, phenyl or thienyl, each of which is unsubstituted or monosubstituted or disubstituted by Hal and/or A,

20 or is alkyl having 1-6 carbon atoms, which may be substituted by thiophene, imidazole, indole, S(O)<sub>m</sub>R<sup>2</sup>, OR<sup>2</sup>, cycloalkyl or phenyl, where the imidazole ring may have a substituent A,

X is NH or O,

W is absent,

Y is Ar-diyl,

25 Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal or A,

26474-891

58

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl, 3-A'-imidazolidin-1-yl or thiadiazolyl, each of which is monosubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN, where the thiadiazole ring may  
 5 additionally be substituted by A, CONH<sub>2</sub> or COOA,

R<sup>2b</sup> is H, OH or OA", and

A" is methyl, ethyl, propyl, isopropyl or butyl.

16. The compound according to any one of Claims 1 to 15, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any  
 10 mixture thereof in all ratios, in which:

D is phenyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,

R<sup>1</sup> is H, phenyl which is unsubstituted or monosubstituted by Hal, or alkyl having 1-6 carbon atoms,

X is NH or O,

15 W is absent,

Y is Ar-diyl,

Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal or A,

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl or 3-A"-imidazolidin-1-yl, each of which is monosubstituted by  
 20 =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN,

R<sup>2b</sup> is H, OH or OA", and

A" is methyl, ethyl, propyl, isopropyl or butyl.

26474-891

59

17. The compound according to Claim 1, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in which:

D is phenyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,

5 R<sup>1</sup> is H, phenyl which is unsubstituted or monosubstituted by Hal, alkyl having 1-6 carbon atoms, 2-thienyl or 2-furyl,

X is NH or O,

W is absent,

Y is Ar-diyl,

10 Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal or A,

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl, 1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl, imidazolidin-1-yl or 3-A"-imidazolidin-1-yl, each of which is monosubstituted by

15 =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN,

R<sup>2b</sup> is H, OH or OA", and

A" is methyl, ethyl, propyl, isopropyl or butyl.

18. The compound according to Claim 1, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, in

20 which:

D is phenyl or thienyl, each of which is unsubstituted or monosubstituted by Hal,

R<sup>1</sup> is H, phenyl which is unsubstituted or monosubstituted or disubstituted by Hal and/or A, alkyl having 1-6 carbon atoms, 2-thienyl or 2-furyl,

or R<sup>1</sup> is alkyl having 1-6 carbon atoms which may be substituted by thiophene,

25 imidazole, indole, S(O)<sub>m</sub>A", OA", cycloalkyl or phenyl, where the imidazole ring may have a substituent A",

26474-891

60

X is NH or O,

W is absent,

Y is Ar-diyl,

Ar is phenyl which is unsubstituted or monosubstituted or disubstituted by Hal or  
5 A,

T is piperidin-1-yl, pyrrolidin-1-yl, 1*H*-pyridin-1-yl, morpholin-4-yl, piperazin-1-yl,  
1,3-oxazolidin-3-yl, 2*H*-pyridazin-2-yl, azepan-1-yl, 2-azabicyclo[2.2.2]octan-2-yl,  
imidazolidin-1-yl, 3-*A'*-imidazolidin-1-yl or thiadiazolyl, each of which is  
monosubstituted by =NR<sup>2b</sup>, =S, =NOR<sup>2b</sup> or =N-CN, where the thiadiazole ring may  
10 additionally be substituted by A'', CONH<sub>2</sub> or COOA'',

R<sup>2b</sup> is H, OH or OA'', and

A'' is methyl, ethyl, propyl, isopropyl or butyl.

19. The compound according to Claim 1, which is:

(R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-  
15 methoxyiminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-imino-piperidin-1-  
yl)phenyl]-2-phenylacetamide,

(R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopiperidin-1-yl)-  
3-methylphenyl]-2-phenylacetamide,

20 (R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)-  
3-methylphenyl]-2-phenylacetamide,

(R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopiperidin-1-  
yl)phenyl]-2-(2-fluorophenyl)acetamide,

25 2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopiperidin-1-yl)-  
phenyl]-2-(2-chlorophenyl)acetamide,

26474-891

61

(S)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)-phenyl]propionamide,

5 (R)-2-[N-(5-chloropyridin-2-yl)carbamoyloxy]-N-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide,

10 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-imino-2*H*-pyridin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[N-(5-chlorothiophen-2-yl)carbamoyloxy]-N-[4-(2-imino-piperidin-1-yl)phenyl]-2-phenylacetamide,

15 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(3-iminomorpholin-4-yl)phenyl]-2-phenylacetamide,

2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-(2-chlorophenyl)acetamide,

20 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

(S)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

(R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-pyrrolidin-1-yl)phenyl]-2-phenylacetamide,

25 (R)-2-[N-(4-chlorophenyl)carbamoyloxy]-N-[4-(2-methoxyimino-piperidin-1-yl)-3-methylphenyl]-2-(2-chlorophenyl)acetamide,

26474-891

62

(R)-2-[*N*-(4-chlorophenyl)carbamoxyloxy]-*N*-[4-(2-azabicyclo-[2.2.2]octan-3-imino-2-yl)phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide, hydrochloride,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

10 (S)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]propionamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-phenylacetamide,

15 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]valeramide,

20 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-(2-fluorophenyl)acetamide,

(R)-2-[3-(5-chlorothiophen-2-yl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)phenyl]-2-phenylacetamide,

(S)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-methoxyiminopiperidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

25 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-methoxyiminopyrrolidin-1-yl)phenyl]-2-phenylacetamide,

26474-891

63

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-thienylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-phenylacetamide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-(thiophen-2-yl)acetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]valeramide,

10 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]-2-thiophen-2-ylacetamide,

15 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-3-methylphenyl]-2-(2-thienyl)acetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-(2-thienyl)acetamide,

20 (R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]-2-phenylacetamide,

(R)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)-3-methylphenyl]valeramide,

2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide, citrate,

25 (R)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, tartrate,

26474-891

64

2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-methoxyiminopiperidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-methoxyimino-pyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

5 2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-fluorophenyl)acetamide,

(*R*)-2-[*N*-(4-chlorophenyl)carbamoyloxy]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-phenylacetamide, lactate,

10 (*R*)-2-[3-(4-chloro-2-methylphenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)phenyl]-2-(2-thienyl)acetamide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

(*S*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopyrrolidin-1-yl)-phenyl]-2-phenylacetamide,

15 (*S*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminopiperidin-1-yl)-phenyl]-2-phenylacetamide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)phenyl]-2-(2-thienyl)acetamide,

20 (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-imino-5-methyl-1,3,4-thiadiazol-3-yl)phenyl]-2-phenylacetamide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-iminoimidazolidin-1-yl)phenyl]-2-phenylacetamide,

(*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-phenylacetamide,

25 (*R*)-2-[3-(4-chlorophenyl)ureido]-*N*-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-2-(2-thienyl)acetamide,

26474-891

65

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]valeramide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]valeramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]valeramide,

10 2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]acetamide,

2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]acetamide,

2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]acetamide,

15 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]propionamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]propionamide,

20 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]propionamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]-3-methylbutyramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methyl-imidazolidin-1-yl)phenyl]-3-methylbutyramide,

25 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-3-methylbutyramide,

26474-891

66

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminoimidazolidin-1-yl)phenyl]-4-methylvaleramide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-3-methylimidazolidin-1-yl)phenyl]-4-methylvaleramide,

5 (R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-4-methylvaleramide,

2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopyrrolidin-1-yl)-phenyl]-3-methoxypropionamide,

10 2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyaniminoimidazolidin-1-yl)phenyl]-3-methoxypropionamide,

(R)-2-[3-(4-chlorophenyl)ureido]-N-[4-(2-cyanimino-imidazolidin-1-yl)phenyl]-3-(1-methyl-1*H*-imidazol-4-yl)propionamide,

2-[3-(4-chlorophenyl)ureido]-N-[4-(5-ethyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide,

15 2-[3-(4-chlorophenyl)ureido]-N-[4-(5-aminocarbonyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide,

2-[3-(4-chlorophenyl)ureido]-N-[4-(5-ethoxycarbonyl-2-imino-1,3,4-thiadiazol-3-yl)phenyl]-3-methoxypropionamide, or

20 2-[3-(4-chlorophenyl)ureido]-N-[4-(2-iminopiperidin-1-yl)-phenyl]-4-methylsulfonylbutyramide,


or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios.

20. Process for the preparation of a compound of the formula I according to any one of Claims 1 to 19 or a pharmaceutically usable derivative,  
25 solvate or stereoisomer thereof, characterised in that

a) a compound of the formula II

26474-891

67



II

in which

$\text{R}^1$ , X, W, Y and T are as defined in Claim 1,

is reacted with a compound of the formula III

5



III

in which

D is as defined in Claim 1,

or

b) a compound of the formula IV


10



IV

in which W, Y and T are as defined in Claim 1,

is reacted with a compound of the formula V



V

in which

15 L is Cl, Br, I or a free or reactively functionally modified OH group,

and

$\text{R}^1$ , X and D are as defined in Claim 1,

and/or

26474-891

68

c) in that a radical T in a compound of the formula I is converted into another radical T,

and/or

a base or acid of the formula I is converted into one of its salts.

- 5 21. The process according to Claim 20, wherein in c) a sulfanyl compound is converted into an imino compound.
22. The compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, as an inhibitor of coagulation factor Xa.
- 10 23. The compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, as an inhibitor of coagulation factor VIIa.
24. A pharmaceutical preparation comprising at least one compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative,
   
15 solvate or stereoisomer thereof, or any mixture thereof in all ratios, and an excipient and/or adjuvant.
25. The pharmaceutical preparation according to Claim 24 for use in the inhibition of coagulation factor Xa.
26. The pharmaceutical preparation according to Claim 24 for use in the
   
20 inhibition of coagulation factor VIIa.
27. The pharmaceutical preparation according to Claim 24 for use in the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases.
- 25 28. A pharmaceutical preparation comprising at least one compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, and at least one further medicament active ingredient.

26474-891

69

29. The pharmaceutical preparation according to Claim 28 for use in the inhibition of coagulation factor Xa.

30. The pharmaceutical preparation according to Claim 28 for use in the inhibition of coagulation factor VIIa.

5 31. The pharmaceutical preparation according to Claim 28 for use in the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases.

10 32. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the preparation of a medicament for inhibition of coagulation factor Xa.

15 33. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the preparation of a medicament for inhibition of coagulation factor VIIa.

20 34. Use of a compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the preparation of a medicament for the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases.

25 35. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for inhibition of coagulation factor Xa.

36. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for inhibition of coagulation factor VIIa.

26474-891

70

37. Use of a compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after  
5 angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases.

38. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the preparation of a medicament for inhibition of  
10 coagulation factor Xa in combination with at least one further medicament active ingredient.

39. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the preparation of a medicament for inhibition of  
15 coagulation factor VIIa in combination with at least one further medicament active ingredient.

40. Use of a compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the preparation of a medicament for the treatment  
20 of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases, in combination with at least one further medicament active ingredient.

41. Use of a compound according to any one of claims 1 to 19, or a  
25 pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for inhibition of coagulation factor Xa in combination with at least one further medicament active ingredient.

42. Use of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any  
30 mixture thereof in all ratios, for inhibition of coagulation factor VIIa in combination with at least one further medicament active ingredient.

26474-891

71

43. Use of a compound according to any one of Claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios, for the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases, in combination with at least one further medicament active ingredient.
44. A kit consisting of separate packs of
- (a) an effective amount of a compound according to any one of claims 1 to 19, or a pharmaceutically usable derivative, solvate or stereoisomer thereof, or any mixture thereof in all ratios,
- and
- (b) an effective amount of a further medicament active ingredient.
45. The kit according to claim 44 for use in inhibiting coagulation factor Xa.
46. The kit according to claim 44 for use in inhibiting coagulation factor VIIa.
47. The kit according to claim 44 for use in the treatment of thromboses, myocardial infarction, arteriosclerosis, inflammation, apoplexia, angina pectoris, restenosis after angioplasty, claudicatio intermittens, migraine, tumours, tumour diseases and/or tumour metastases.

