

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
31 March 2005 (31.03.2005)

PCT

(10) International Publication Number
WO 2005/029085 A2

- (51) International Patent Classification⁷: **G01N 33/68**
- (21) International Application Number:
PCT/EP2004/010112
- (22) International Filing Date:
10 September 2004 (10.09.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
03021125.4 22 September 2003 (22.09.2003) EP
- (71) Applicant (for all designated States except US): **BAYER HEALTHCARE AG** [DE/DE]; 51368 Leverkusen (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **GOLZ, Stefan** [DE/DE]; Bückmannsmühle 46, 45326 Essen (DE). **BRÜGGEMEIER, Ulf** [DE/DE]; Leysiefen 20, 42799 Leichlingen (DE). **GEERTS, Andreas** [DE/DE]; Schuckertstr. 29, 42113 Wuppertal (DE).
- (74) Common Representative: **BAYER HEALTHCARE AG**; Law and Patents, Patents and Licensing, 51368 Leverkusen (DE).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2005/029085 A2

(54) Title: DIAGNOSTICS AND THERAPEUTICS FOR DISEASES ASSOCIATED WITH ORGANIC CATION TRANSPORTER SLC22A11 (SLC22A11)

(57) Abstract: The invention provides a human SLC22A11 which is associated with the cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. The invention also provides assays for the identification of compounds useful in the treatment or prevention of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. The invention also features compounds which bind to and/or activate or inhibit the activity of SLC22A11 as well as pharmaceutical compositions comprising such compounds.

**Diagnostics and Therapeutics for Diseases Associated with Organic Cation
Transporter SLC22A11 (SLC22A11)**

5 **Technical field of the invention**

The present invention is in the field of molecular biology, more particularly, the present invention relates to nucleic acid sequences and amino acid sequences of a human SLC22A11 and its regulation for the treatment of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory
10 diseases, neurological diseases and urological diseases in mammals.

Background of the invention

15 Organic anion transporting (OAT) polypeptides play an important role in the uptake of organic anions, including bile acids, bilirubin conjugates and sulfobromophthalein, in the liver [Hsiang *et al.*, (1999); Konig *et al.*, (2000); Kouzuki *et al.*, (2000)]. Thus, levels of organic anion transporting polypeptides can be modulated to affect the rate of drug clearance via hepatocellular uptake. Further, OATs have been found
20 to transport eicosanoids, taurocholate, conjugated steroids, and thyroid hormones.

SLC22A11

The nucleotide sequence of SLC22A11 is accessible in the databases by the
25 accession number BC034384 and is given in SEQ ID NO:1. The amino acid sequence of SLC22A11 depicted in SEQ ID NO:2.

As part of a pharmacogenetics project that sought to identify genes that determine drug response, Leabman *et al.* [Leabman *et al.*, 2002] screened for variation in the set
30 of 24 genes encoding membrane transporters. They identified variants by screening an ethnically diverse collection of genomic DNA samples, derived from a total of

494 chromosomes. By combining population-genetic and phylogenetic analyses, they were able to identify amino acid residues and protein domains that may be important for human fitness. A large sample set made it possible for them also to obtain information about rare variants. The 24 membrane transporters with potential roles in drug response were grouped based on transporter family: e.g., OCT1, OCT2 and OCT3 belonged to the SLC6 family; CNT1 and CNT2 belonged to the SLC28 family.

Organic anion transporting (OAT) polypeptides play an important role in the uptake of organic anions, including bile acids, bilirubin conjugates and sulfobromophthalein, in the liver [Hsiang et al. (1999), König et al. (2000), Kouzuki et al. (2000)]. Thus, levels of organic anion transporting polypeptides can be modulated to affect the rate of drug clearance via hepatocellular uptake. Further, OATs have been found to transport eicosanoids, taurocholate, conjugated steroids, and thyroid hormones. Poly-specific organic cation transporters in the liver, kidney, and intestine are critical for elimination of many endogenous amines as well as a wide array of drugs and environmental toxins [Grundemann et al., (1994)]

SLC22A11 is published in [Enomoto et al., (2002); Cha et al., (2000)] and patents WO0149728 and WO0192304

Summary of the invention

The invention relates to novel disease associations of SLC22A11 polypeptides and polynucleotides. The invention also relates to novel methods of screening for therapeutic agents for the treatment of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal. The invention also relates to pharmaceutical compositions for the treatment of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a SLC22A11 polypeptide, a SLC22A11 polynucleotide, or regulators of SLC22A11 or modulators

of SLC22A11 activity. The invention further comprises methods of diagnosing cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal.

5

Brief Description of the Drawings

Fig. 1 shows the nucleotide sequence of a SLC22A11 polynucleotide (SEQ ID NO:1).

Fig. 2 shows the amino acid sequence of a SLC22A11 polypeptide (SEQ ID NO:2).

10 Fig. 3 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:3).

Fig. 4 shows the nucleotide sequence of a primer useful for the invention (SEQ ID NO:4).

15 Fig. 5 shows a nucleotide sequence useful as a probe to detect proteins of the invention (SEQ ID NO:5).

Detailed description of the invention

Definition of terms

20

An "oligonucleotide" is a stretch of nucleotide residues which has a sufficient number of bases to be used as an oligomer, amplimer or probe in a polymerase chain reaction (PCR). Oligonucleotides are prepared from genomic or cDNA sequence and are used to amplify, reveal, or confirm the presence of a similar DNA or RNA in a particular cell or tissue. Oligonucleotides or oligomers comprise portions of a DNA sequence having at least about 10 nucleotides and as many as about 35 nucleotides, preferably about 25 nucleotides.

25

"Probes" may be derived from naturally occurring or recombinant single- or double-stranded nucleic acids or may be chemically synthesized. They are useful in detecting the presence of identical or similar sequences. Such probes may be labeled

30

with reporter molecules using nick translation, Klenow fill-in reaction, PCR or other methods well known in the art. Nucleic acid probes may be used in southern, northern or in situ hybridizations to determine whether DNA or RNA encoding a certain protein is present in a cell type, tissue, or organ.

5

A "fragment of a polynucleotide" is a nucleic acid that comprises all or any part of a given nucleotide molecule, the fragment having fewer nucleotides than about 6 kb, preferably fewer than about 1 kb.

10

"Reporter molecules" are radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents which associate with a particular nucleotide or amino acid sequence, thereby establishing the presence of a certain sequence, or allowing for the quantification of a certain sequence.

15

"Chimeric" molecules may be constructed by introducing all or part of the nucleotide sequence of this invention into a vector containing additional nucleic acid sequence which might be expected to change any one or several of the following SLC22A11 characteristics: cellular location, distribution, ligand-binding affinities, interchain affinities, degradation/turnover rate, signaling, etc.

20

"Active", with respect to a SLC22A11 polypeptide, refers to those forms, fragments, or domains of a SLC22A11 polypeptide which retain the biological and/or antigenic activity of a SLC22A11 polypeptide.

25

"Naturally occurring SLC22A11 polypeptide" refers to a polypeptide produced by cells which have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including but not limited to acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

30

“Derivative“ refers to polypeptides which have been chemically modified by techniques such as ubiquitination, labeling (see above), pegylation (derivatization with polyethylene glycol), and chemical insertion or substitution of amino acids such as ornithine which do not normally occur in human proteins.

5

“Conservative amino acid substitutions“ result from replacing one amino acid with another having similar structural and/or chemical properties, such as the replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

10

“Insertions“ or “deletions“ are typically in the range of about 1 to 5 amino acids. The variation allowed may be experimentally determined by producing the peptide synthetically while systematically making insertions, deletions, or substitutions of nucleotides in the sequence using recombinant DNA techniques.

15

A “signal sequence“ or “leader sequence“ can be used, when desired, to direct the polypeptide through a membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous sources by recombinant DNA techniques.

20

An “oligopeptide“ is a short stretch of amino acid residues and may be expressed from an oligonucleotide. Oligopeptides comprise a stretch of amino acid residues of at least 3, 5, 10 amino acids and at most 10, 15, 25 amino acids, typically of at least 9 to 13 amino acids, and of sufficient length to display biological and/or antigenic activity.

25

“Inhibitor“ is any substance which retards or prevents a chemical or physiological reaction or response. Common inhibitors include but are not limited to antisense molecules, antibodies, and antagonists.

30

“Standard expression“ is a quantitative or qualitative measurement for comparison. It is based on a statistically appropriate number of normal samples and is created to use as a basis of comparison when performing diagnostic assays, running clinical trials, or following patient treatment profiles.

5

“Animal“ as used herein may be defined to include human, domestic (e.g., cats, dogs, etc.), agricultural (e.g., cows, horses, sheep, etc.) or test species (e.g., mouse, rat, rabbit, etc.).

10

A “SLC22A11 polynucleotide“, within the meaning of the invention, shall be understood as being a nucleic acid molecule selected from a group consisting of

(i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2,

15

(ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1,

(iii) nucleic acid molecules having the sequence of SEQ ID NO: 1,

20

(iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and

(v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code;

25

wherein the polypeptide encoded by said nucleic acid molecule has SLC22A11 activity.

30

A “SLC22A11 polypeptide“, within the meaning of the invention, shall be understood as being a polypeptide selected from a group consisting of

- (i) polypeptides having the sequence of SEQ ID NO: 2,
 - (ii) polypeptides comprising the sequence of SEQ ID NO: 2,
 - 5 (iii) polypeptides encoded by SLC22A11 polynucleotides; and
 - (iv) polypeptides which show at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii);
- 10 wherein said polypeptide has SLC22A11 activity.

The nucleotide sequences encoding a SLC22A11 (or their complement) have numerous applications in techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use in the construction of oligomers for PCR, use for chromosome and gene mapping, use in the
15 recombinant production of SLC22A11, and use in generation of antisense DNA or RNA, their chemical analogs and the like. Uses of nucleotides encoding a SLC22A11 disclosed herein are exemplary of known techniques and are not intended to limit their use in any technique known to a person of ordinary skill in the art.

20 Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, e.g., the triplet genetic code, specific base pair interactions, etc.

25 It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of SLC22A11 - encoding nucleotide sequences may be produced. Some of these will only bear minimal homology to the nucleotide sequence of the known and naturally occurring SLC22A11. The invention has specifically contemplated each and every possible variation of nucleotide sequence
30 that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as

applied to the nucleotide sequence of naturally occurring SLC22A11, and all such variations are to be considered as being specifically disclosed.

Although the nucleotide sequences which encode a SLC22A11, its derivatives or its
5 variants are preferably capable of hybridizing to the nucleotide sequence of the
naturally occurring SLC22A11 polynucleotide under stringent conditions, it may be
advantageous to produce nucleotide sequences encoding SLC22A11 polypeptides or
its derivatives possessing a substantially different codon usage. Codons can be
selected to increase the rate at which expression of the peptide occurs in a particular
10 prokaryotic or eukaryotic expression host in accordance with the frequency with
which particular codons are utilized by the host. Other reasons for substantially
altering the nucleotide sequence encoding a SLC22A11 polypeptide and/or its
derivatives without altering the encoded amino acid sequence include the production
of RNA transcripts having more desirable properties, such as a greater half-life, than
15 transcripts produced from the naturally occurring sequence.

Nucleotide sequences encoding a SLC22A11 polypeptide may be joined to a variety
of other nucleotide sequences by means of well established recombinant DNA
techniques. Useful nucleotide sequences for joining to SLC22A11 polynucleotides
20 include an assortment of cloning vectors such as plasmids, cosmids, lambda phage
derivatives, phagemids, and the like. Vectors of interest include expression vectors,
replication vectors, probe generation vectors, sequencing vectors, etc. In general,
vectors of interest may contain an origin of replication functional in at least one
organism, convenient restriction endonuclease sensitive sites, and selectable markers
25 for one or more host cell systems.

Another aspect of the subject invention is to provide for SLC22A11-specific hybridi-
zation probes capable of hybridizing with naturally occurring nucleotide sequences
encoding SLC22A11. Such probes may also be used for the detection of similar
30 organic ion transporter encoding sequences and should preferably show at least 40%
nucleotide identity to SLC22A11 polynucleotides. The hybridization probes of the

subject invention may be derived from the nucleotide sequence presented as SEQ ID NO: 1 or from genomic sequences including promoter, enhancers or introns of the native gene. Hybridization probes may be labelled by a variety of reporter molecules using techniques well known in the art.

5

It will be recognized that many deletional or mutational analogs of SLC22A11 polynucleotides will be effective hybridization probes for SLC22A11 polynucleotides. Accordingly, the invention relates to nucleic acid sequences that hybridize with such SLC22A11 encoding nucleic acid sequences under stringent conditions.

10

“Stringent conditions“ refers to conditions that allow for the hybridization of substantially related nucleic acid sequences. For instance, such conditions will generally allow hybridization of sequence with at least about 85% sequence identity, preferably with at least about 90% sequence identity, more preferably with at least about 95% sequence identity. Hybridization conditions and probes can be adjusted in well-characterized ways to achieve selective hybridization of human-derived probes. Stringent conditions, within the meaning of the invention are 65°C in a buffer containing 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7 % (w/v) SDS.

15

Nucleic acid molecules that will hybridize to SLC22A11 polynucleotides under stringent conditions can be identified functionally. Without limitation, examples of the uses for hybridization probes include: histochemical uses such as identifying tissues that express SLC22A11; measuring mRNA levels, for instance to identify a sample's tissue type or to identify cells that express abnormal levels of SLC22A11; and detecting polymorphisms of SLC22A11.

25

PCR provides additional uses for oligonucleotides based upon the nucleotide sequence which encodes SLC22A11. Such probes used in PCR may be of recombinant origin, chemically synthesized, or a mixture of both. Oligomers may comprise discrete nucleotide sequences employed under optimized conditions for identification of SLC22A11 in specific tissues or diagnostic use. The same two oligomers, a

30

nested set of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for identification of closely related DNAs or RNAs.

5 Rules for designing polymerase chain reaction (PCR) primers are now established, as reviewed by PCR Protocols. Degenerate primers, i.e., preparations of primers that are heterogeneous at given sequence locations, can be designed to amplify nucleic acid sequences that are highly homologous to, but not identical with SLC22A11. Strategies are now available that allow for only one of the primers to be required to specifically hybridize with a known sequence. For example, appropriate nucleic acid
10 primers can be ligated to the nucleic acid sought to be amplified to provide the hybridization partner for one of the primers. In this way, only one of the primers need be based on the sequence of the nucleic acid sought to be amplified.

15 PCR methods for amplifying nucleic acid will utilize at least two primers. One of these primers will be capable of hybridizing to a first strand of the nucleic acid to be amplified and of priming enzyme-driven nucleic acid synthesis in a first direction. The other will be capable of hybridizing the reciprocal sequence of the first strand (if the sequence to be amplified is single stranded, this sequence will initially be hypothetical, but will be synthesized in the first amplification cycle) and of priming
20 nucleic acid synthesis from that strand in the direction opposite the first direction and towards the site of hybridization for the first primer. Conditions for conducting such amplifications, particularly under preferred stringent hybridization conditions, are well known.

25 Other means of producing specific hybridization probes for SLC22A11 include the cloning of nucleic acid sequences encoding SLC22A11 or SLC22A11 derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA
30 polymerase and the appropriate reporter molecules.

It is possible to produce a DNA sequence, or portions thereof, entirely by synthetic chemistry. After synthesis, the nucleic acid sequence can be inserted into any of the many available DNA vectors and their respective host cells using techniques which are well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into the nucleotide sequence. Alternately, a portion of sequence in which a mutation is desired can be synthesized and recombined with longer portion of an existing genomic or recombinant sequence.

SLC22A11 polynucleotides may be used to produce a purified oligo-or polypeptide using well known methods of recombinant DNA technology. The oligopeptide may be expressed in a variety of host cells, either prokaryotic or eukaryotic. Host cells may be from the same species from which the nucleotide sequence was derived or from a different species. Advantages of producing an oligonucleotide by recombinant DNA technology include obtaining adequate amounts of the protein for purification and the availability of simplified purification procedures.

Quantitative determinations of nucleic acids

An important step in the molecular genetic analysis of human disease is often the enumeration of the copy number of a nucleic acid or the relative expression of a gene in particular tissues.

Several different approaches are currently available to make quantitative determinations of nucleic acids. Chromosome-based techniques, such as comparative genomic hybridization (CGH) and fluorescent in situ hybridization (FISH) facilitate efforts to cytogenetically localize genomic regions that are altered in tumor cells. Regions of genomic alteration can be narrowed further using loss of heterozygosity analysis (LOH), in which disease DNA is analyzed and compared with normal DNA for the loss of a heterozygous polymorphic marker. The first experiments used restriction fragment length polymorphisms (RFLPs) [Johnson, (1989)], or hyper-variable minisatellite DNA [Barnes, 2000]. In recent years LOH has been performed

primarily using PCR amplification of microsatellite markers and electrophoresis of the radio labelled [Jeffreys, (1985)] or fluorescently labelled PCR products [Weber, (1990)] and compared between paired normal and disease DNAs.

5 A number of other methods have also been developed to quantify nucleic acids [Gergen, (1992)]. More recently, PCR and RT-PCR methods have been developed which are capable of measuring the amount of a nucleic acid in a sample. One approach, for example, measures PCR product quantity in the log phase of the reaction before the formation of reaction products plateaus [Thomas, (1980)].

10

A gene sequence contained in all samples at relatively constant quantity is typically utilized for sample amplification efficiency normalization. This approach, however, suffers from several drawbacks. The method requires that each sample has equal input amounts of the nucleic acid and that the amplification efficiency between samples is identical until the time of analysis. Furthermore, it is difficult using the conventional methods of PCR quantitation such as gel electrophoresis or plate
15 capture hybridization to determine that all samples are in fact analyzed during the log phase of the reaction as required by the method.

20 Another method called quantitative competitive (QC)-PCR, as the name implies, relies on the inclusion of an internal control competitor in each reaction [Piatak, (1993), BioTechniques]. The efficiency of each reaction is normalized to the internal competitor. A known amount of internal competitor is typically added to each sample. The unknown target PCR product is compared with the known competitor
25 PCR product to obtain relative quantitation. A difficulty with this general approach lies in developing an internal control that amplifies with the same efficiency than the target molecule.

5' Fluorogenic Nuclease Assays

Fluorogenic nuclease assays are a real time quantitation method that uses a probe to monitor formation of amplification product. The basis for this method of monitoring the formation of amplification product is to measure continuously PCR product accumulation using a dual-labelled fluorogenic oligonucleotide probe, an approach frequently referred to in the literature simply as the "TaqMan method" [Piatak,(1993), Science; Heid, (1996); Gibson, (1996); Holland. (1991)].

The probe used in such assays is typically a short (about 20-25 bases) oligonucleotide that is labeled with two different fluorescent dyes. The 5' terminus of the probe is attached to a reporter dye and the 3' terminus is attached to a quenching dye, although the dyes could be attached at other locations on the probe as well. The probe is designed to have at least substantial sequence complementarity with the probe binding site. Upstream and downstream PCR primers which bind to flanking regions of the locus are added to the reaction mixture. When the probe is intact, energy transfer between the two fluorophors occurs and the quencher quenches emission from the reporter. During the extension phase of PCR, the probe is cleaved by the 5' nuclease activity of a nucleic acid polymerase such as Taq polymerase, thereby releasing the reporter from the oligonucleotide-quencher and resulting in an increase of reporter emission intensity which can be measured by an appropriate detector.

One detector which is specifically adapted for measuring fluorescence emissions such as those created during a fluorogenic assay is the ABI 7700 or 4700 HT manufactured by Applied Biosystems, Inc. in Foster City, Calif. The ABI 7700 uses fiber optics connected with each well in a 96-or 384 well PCR tube arrangement. The instrument includes a laser for exciting the labels and is capable of measuring the fluorescence spectra intensity from each tube with continuous monitoring during PCR amplification. Each tube is re-examined every 8.5 seconds.

Computer software provided with the instrument is capable of recording the fluorescence intensity of reporter and quencher over the course of the amplification. The recorded values will then be used to calculate the increase in normalized reporter emission intensity on a continuous basis. The increase in emission intensity is plotted versus time, i.e., the number of amplification cycles, to produce a continuous measure of amplification. To quantify the locus in each amplification reaction, the amplification plot is examined at a point during the log phase of product accumulation. This is accomplished by assigning a fluorescence threshold intensity above background and determining the point at which each amplification plot crosses the threshold (defined as the threshold cycle number or Ct). Differences in threshold cycle number are used to quantify the relative amount of PCR target contained within each tube. Assuming that each reaction functions at 100% PCR efficiency, a difference of one Ct represents a two-fold difference in the amount of starting template. The fluorescence value can be used in conjunction with a standard curve to determine the amount of amplification product present.

Non-Probe-Based Detection Methods

A variety of options are available for measuring the amplification products as they are formed. One method utilizes labels, such as dyes, which only bind to double stranded DNA. In this type of approach, amplification product (which is double stranded) binds dye molecules in solution to form a complex. With the appropriate dyes, it is possible to distinguish between dye molecules free in solution and dye molecules bound to amplification product. For example, certain dyes fluoresce only when bound to amplification product. Examples of dyes which can be used in methods of this general type include, but are not limited to, Syber Green.TM. and Pico Green from Molecular Probes, Inc. of Eugene, Oreg., ethidium bromide, propidium iodide, chromomycin, acridine orange, Hoechst 33258, Toto-1, Yoyo-1, DAPI (4',6-diamidino-2-phenylindole hydrochloride).

Another real time detection technique measures alteration in energy fluorescence energy transfer between fluorophors conjugated with PCR primers [Livak, (1995)].

Probe-Based Detection Methods

5

These detection methods involve some alteration to the structure or conformation of a probe hybridized to the locus between the amplification primer pair. In some instances, the alteration is caused by the template-dependent extension catalyzed by a nucleic acid polymerase during the amplification process. The alteration generates a detectable signal which is an indirect measure of the amount of amplification product formed.

10

For example, some methods involve the degradation or digestion of the probe during the extension reaction. These methods are a consequence of the 5'-3' nuclease activity associated with some nucleic acid polymerases. Polymerases having this activity cleave mononucleotides or small oligonucleotides from an oligonucleotide probe annealed to its complementary sequence located within the locus.

15

The 3' end of the upstream primer provides the initial binding site for the nucleic acid polymerase. As the polymerase catalyzes extension of the upstream primer and encounters the bound probe, the nucleic acid polymerase displaces a portion of the 5' end of the probe and through its nuclease activity cleaves mononucleotides or oligonucleotides from the probe.

20

The upstream primer and the probe can be designed such that they anneal to the complementary strand in close proximity to one another. In fact, the 3' end of the upstream primer and the 5' end of the probe may abut one another. In this situation, extension of the upstream primer is not necessary in order for the nucleic acid polymerase to begin cleaving the probe. In the case in which intervening nucleotides separate the upstream primer and the probe, extension of the primer is necessary before the nucleic acid polymerase encounters the 5' end of the probe. Once contact

25

30

occurs and polymerization continues, the 5'-3' exonuclease activity of the nucleic acid polymerase begins cleaving mononucleotides or oligonucleotides from the 5' end of the probe. Digestion of the probe continues until the remaining portion of the probe dissociates from the complementary strand.

5

In solution, the two end sections can hybridize with each other to form a hairpin loop. In this conformation, the reporter and quencher dye are in sufficiently close proximity that fluorescence from the reporter dye is effectively quenched by the quencher dye. Hybridized probe, in contrast, results in a linearized conformation in which the extent of quenching is decreased. Thus, by monitoring emission changes for the two dyes, it is possible to indirectly monitor the formation of amplification product.

10

Probes

The labeled probe is selected so that its sequence is substantially complementary to a segment of the test locus or a reference locus. As indicated above, the nucleic acid site to which the probe binds should be located between the primer binding sites for the upstream and downstream amplification primers.

15

Primers

20

The primers used in the amplification are selected so as to be capable of hybridizing to sequences at flanking regions of the locus being amplified. The primers are chosen to have at least substantial complementarity with the different strands of the nucleic acid being amplified. When a probe is utilized to detect the formation of amplification products, the primers are selected in such that they flank the probe, i.e. are located upstream and downstream of the probe.

25

The primer must have sufficient length so that it is capable of priming the synthesis of extension products in the presence of an agent for polymerization. The length and composition of the primer depends on many parameters, including, for example, the

30

temperature at which the annealing reaction is conducted, proximity of the probe binding site to that of the primer, relative concentrations of the primer and probe and the particular nucleic acid composition of the probe. Typically the primer includes 15-30 nucleotides. However, the length of the primer may be more or less depending on the complexity of the primer binding site and the factors listed above.

Labels for Probes and Primers

The labels used for labeling the probes or primers of the current invention and which can provide the signal corresponding to the quantity of amplification product can take a variety of forms. As indicated above with regard to the 5' fluorogenic nuclease method, a fluorescent signal is one signal which can be measured. However, measurements may also be made, for example, by monitoring radioactivity, colorimetry, absorption, magnetic parameters, or enzymatic activity. Thus, labels which can be employed include, but are not limited to, fluorophores, chromophores, radioactive isotopes, electron dense reagents, enzymes, and ligands having specific binding partners (e.g., biotin-avidin).

Monitoring changes in fluorescence is a particularly useful way to monitor the accumulation of amplification products. A number of labels useful for attachment to probes or primers are commercially available including fluorescein and various fluorescein derivatives such as FAM, HEX, TET and JOE (all which are available from Applied Biosystems, Foster City, Calif.); lucifer yellow, and coumarin derivatives.

Labels may be attached to the probe or primer using a variety of techniques and can be attached at the 5' end, and/or the 3' end and/or at an internal nucleotide. The label can also be attached to spacer arms of various sizes which are attached to the probe or primer. These spacer arms are useful for obtaining a desired distance between multiple labels attached to the probe or primer.

In some instances, a single label may be utilized; whereas, in other instances, such as with the 5' fluorogenic nuclease assays for example, two or more labels are attached to the probe. In cases wherein the probe includes multiple labels, it is generally advisable to maintain spacing between the labels which is sufficient to permit separation of the labels during digestion of the probe through the 5'-3' nuclease activity of the nucleic acid polymerase.

Patients Exhibiting Symptoms of Disease

10 A number of diseases are associated with changes in the copy number of a certain gene. For patients having symptoms of a disease, the real-time PCR method can be used to determine if the patient has copy number alterations which are known to be linked with diseases that are associated with the symptoms the patient has.

15 SLC22A11 expression

SLC22A11 fusion proteins

20 Fusion proteins are useful for generating antibodies against SLC22A11 polypeptides and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of SLC22A11 polypeptides. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

25 A SLC22A11 fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment can comprise at least 54, 75, 100, 125, 139, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO: 2 or of a biologically active variant, such as those described above. The first
30 polypeptide segment also can comprise full-length SLC22A11.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include, but are not limited to β -galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, or herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located adjacent to the SLC22A11.

Preparation of Polynucleotides

A naturally occurring SLC22A11 polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated SLC22A11 polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprise SLC22A11 nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

SLC22A11 cDNA molecules can be made with standard molecular biology techniques, using SLC22A11 mRNA as a template. SLC22A11 cDNA molecules can thereafter be replicated using molecular biology techniques known in the art. An amplification technique, such as PCR, can be used to obtain additional copies of

polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

5 Alternatively, synthetic chemistry techniques can be used to synthesize SLC22A11 polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode SLC22A11 having, for example, an amino acid sequence shown in SEQ ID NO: 2 or a biologically active variant thereof.

Extending Polynucleotides

10

Various PCR-based methods can be used to extend nucleic acid sequences encoding human SLC22A11, for example to detect upstream sequences of SLC22A11 gene such as promoters and regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

20

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region. Primers can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

25

30 Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast

artificial chromosome DNA. In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

5 When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for
10 extension of sequence into 5' non-transcribed regulatory regions.

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic
15 separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate equipment and software (*e.g.*, GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and
20 electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

Obtaining Polypeptides

25

SLC22A11 can be obtained, for example, by purification from human cells, by expression of SLC22A11 polynucleotides, or by direct chemical synthesis.

Protein Purification

30

SLC22A11 can be purified from any human cell which expresses the receptor, including those which have been transfected with expression constructs which express SLC22A11. A purified SLC22A11 is separated from other compounds which normally associate with SLC22A11 in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis.

10 *Expression of SLC22A11 Polynucleotides*

To express SLC22A11, SLC22A11 polynucleotides can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding SLC22A11 and appropriate transcriptional and translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination.

20 A variety of expression vector/host systems can be utilized to contain and express sequences encoding SLC22A11. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (*e.g.*, baculovirus), plant cell systems transformed with virus expression vectors (*e.g.*, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (*e.g.*, Ti or pBR322 plasmids), or animal cell systems.

The control elements or regulatory sequences are those non-translated regions of the vector - enhancers, promoters, 5' and 3' untranslated regions - which interact with host cellular proteins to carry out transcription and translation. Such elements can

vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the
5 BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (*e.g.*, heat shock, RUBISCO, and storage protein genes) or from plant viruses (*e.g.*, viral promoters or leader sequences) can be cloned into the vector. In mammalian
10 cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding SLC22A11, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

15 *Bacterial and Yeast Expression Systems*

In bacterial systems, a number of expression vectors can be selected. For example, when a large quantity of SLC22A11 is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be
20 used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding SLC22A11 can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors or pGEX vectors (Promega, Madison, Wis.)
25 also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the
30 cloned polypeptide of interest can be released from the GST moiety at will.

Plant and Insect Expression Systems

If plant expression vectors are used, the expression of sequences encoding SLC22A11 can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection.

10

An insect system also can be used to express SLC22A11. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding SLC22A11 can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of SLC22A11 will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which SLC22A11 can be expressed.

15

20

Mammalian Expression Systems

A number of viral-based expression systems can be used to express SLC22A11 in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding SLC22A11 can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing SLC22A11 in infected host cells [Engelhard, 1994]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

25

30

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (*e.g.*, liposomes, polycationic amino polymers, or vesicles). Specific initiation signals also can be used to achieve more efficient translation of sequences encoding SLC22A11. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding SLC22A11, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic.

15

Host Cells

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed SLC22A11 in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (*e.g.*, CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

30 Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express SLC22A11 can be trans-

formed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1-2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced SLC22A11 sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Logan, (1984)] and adenine phosphoribosyltransferase [Wigler, (1977)] genes which can be employed in *tk* or *aprt* cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate [Lowy, (1980)], *npt* confers resistance to the aminoglycosides, neomycin and G-418 [Wigler, (1980)], and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively [Colbere-Garapin, 1981]. Additional selectable genes have been described. For example, *trpB* allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system

Detecting Polypeptide Expression

Although the presence of marker gene expression suggests that a SLC22A11 polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding SLC22A11 is inserted within a marker gene sequence, transformed cells containing sequences which encode SLC22A11 can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding SLC22A11 under the control of a

single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of SLC22A11 polynucleotide.

Alternatively, host cells which contain a SLC22A11 polynucleotide and which
5 express SLC22A11 can be identified by a variety of procedures known to those of
skill in the art. These procedures include, but are not limited to, DNA-DNA or
DNA-RNA hybridizations and protein bioassay or immunoassay techniques which
include membrane, solution, or chip-based technologies for the detection and/or
10 quantification of nucleic acid or protein. For example, the presence of a poly-
nucleotide sequence encoding SLC22A11 can be detected by DNA-DNA or DNA-
RNA hybridization or amplification using probes or fragments or fragments of
polynucleotides encoding SLC22A11. Nucleic acid amplification-based assays
involve the use of oligonucleotides selected from sequences encoding SLC22A11 to
15 detect transformants which contain a SLC22A11 polynucleotide.

A variety of protocols for detecting and measuring the expression of SLC22A11,
using either polyclonal or monoclonal antibodies specific for the polypeptide, are
known in the art. Examples include enzyme-linked immunosorbent assay (ELISA),
radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-
20 site, monoclonal-based immunoassay using monoclonal antibodies reactive to two
non-interfering epitopes on SLC22A11 can be used, or a competitive binding assay
can be employed.

A wide variety of labels and conjugation techniques are known by those skilled in the
25 art and can be used in various nucleic acid and amino acid assays. Means for
producing labeled hybridization or PCR probes for detecting sequences related to
polynucleotides encoding SLC22A11 include oligolabeling, nick translation, end-
labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences
encoding SLC22A11 can be cloned into a vector for the production of an mRNA
30 probe. Such vectors are known in the art, are commercially available, and can be
used to synthesize RNA probes *in vitro* by addition of labeled nucleotides and an

appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

Host cells transformed with SLC22A11 polynucleotides can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing SLC22A11 polynucleotides can be designed to contain signal sequences which direct secretion of soluble SLC22A11 through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound SLC22A11.

As discussed above, other constructions can be used to join a sequence encoding SLC22A11 to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and SLC22A11 also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing SLC22A11 and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography)

Maddox, (1983)], while the enterokinase cleavage site provides a means for purifying SLC22A11 from the fusion protein [Porath, (1992)].

Chemical Synthesis

5

Sequences encoding SLC22A11 can be synthesized, in whole or in part, using chemical methods well known in the art. Alternatively, SLC22A11 itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques. Protein synthesis can either be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of SLC22A11 can be separately synthesized and combined using chemical methods to produce a full-length molecule.

15

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography. The composition of a synthetic SLC22A11 can be confirmed by amino acid analysis or sequencing. Additionally, any portion of the amino acid sequence of SLC22A11 can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

20

Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce SLC22A11 polynucleotides possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences referred to herein can be engineered using methods generally known in the art to alter SLC22A11 polynucleotides for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of SLC22A11.

“Antibody” as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')₂, and Fv, which are capable of binding an epitope of SLC22A11. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acid. An antibody which specifically binds to an epitope of SLC22A11 can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmuno-

assays, immunohistochemical assays, immunoprecipitations, or other immuno-
chemical assays known in the art. Various immunoassays can be used to identify
antibodies having the desired specificity. Numerous protocols for competitive
binding or immunoradiometric assays are well known in the art. Such immunoassays
5 typically involve the measurement of complex formation between an immunogen and
an antibody which specifically binds to the SLC22A11 immunogen.

Typically, an antibody which specifically binds to SLC22A11 provides a detection
signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other
10 proteins when used in an immunochemical assay. Preferably, antibodies which
specifically bind to SLC22A11 do not detect other proteins in immunochemical
assays and can immunoprecipitate SLC22A11 from solution.

SLC22A11 can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea
15 pig, monkey, or human, to produce polyclonal antibodies. If desired, SLC22A11 can
be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and
keyhole limpet hemocyanin. Depending on the host species, various adjuvants can
be used to increase the immunological response. Such adjuvants include, but are not
limited to, Freund's adjuvant, mineral gels (*e.g.*, aluminum hydroxide), and surface
20 active substances (*e.g.*, lysolecithin, pluronic polyols, polyanions, peptides, oil
emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used
in humans, BCG (*bacilli Calmette-Guerin*) and *Corynebacterium parvum* are
especially useful.

25 Monoclonal antibodies which specifically bind to SLC22A11 can be prepared using
any technique which provides for the production of antibody molecules by
continuous cell lines in culture. These techniques include, but are not limited to, the
hybridoma technique, the human B-cell hybridoma technique, and the EBV-
hybridoma technique [Roberge, (1995)].

In addition, techniques developed for the production of “chimeric antibodies”, the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. Monoclonal and other antibodies also can be “humanized” to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such anti-
5 antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed
10 mutagenesis of individual residues or by grating of entire complementarity determining regions. Antibodies which specifically bind to SLC22A11 can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

15 Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to SLC22A11. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combi-
natorial immunoglobulin libraries. Single-chain antibodies also can be constructed
20 using a DNA amplification method, such as PCR, using hybridoma cDNA as a template. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught. A nucleotide sequence encoding a single-chain antibody can be constructed using
25 standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology.

30 Antibodies which specifically bind to SLC22A11 also can be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents. Other types of antibodies can

be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which SLC22A11 is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of SLC22A11 gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters.

Modifications of SLC22A11 gene expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the SLC22A11 gene. Oligonucleotides derived from the transcription initiation site, *e.g.*, between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Nicholls, (1993)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a SLC22A11 polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a SLC22A11 polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent SLC22A11 nucleotides, can provide sufficient targeting specificity for SLC22A11 mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular SLC22A11 polynucleotide sequence. Antisense oligonucleotides can be modified without affecting their ability to hybridize to a SLC22A11 polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense

oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art.

Ribozymes

5

Ribozymes are RNA molecules with catalytic activity [Uhlmann, (1987)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by
10 endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences. The coding sequence of a SLC22A11 polynucleotide can be used to generate ribozymes which will specifically bind to mRNA transcribed from a SLC22A11 polynucleotide. Methods of designing and
15 constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically
20 hybridizes with the target RNA.

Specific ribozyme cleavage sites within a SLC22A11 RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences
25 of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate SLC22A11 RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. The nucleotide sequences
30 shown in SEQ ID NO: 1 and its complement provide sources of suitable hybridization region sequences. Longer complementary sequences can be used to

increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

5

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease SLC22A11 expression.

10

Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells (U.S. 5,641,673). Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

15

Screening / Screening Assays

20

Regulators

Regulators as used herein, refer to compounds that affect the activity of a SLC22A11 in vivo and/or in vivo. Regulators can be agonists and antagonists of a SLC22A11 polypeptide and can be compounds that exert their effect on the SLC22A11 activity via the expression, via post-translational modifications or by other means. Agonists of SLC22A11 are molecules which, when bound to SLC22A11, increase or prolong the activity of SLC22A11. Agonists of SLC22A11 include proteins, nucleic acids, carbohydrates, small molecules, or any other molecule which activate SLC22A11.

25

30

Antagonists of SLC22A11 are molecules which, when bound to SLC22A11, decrease the amount or the duration of the activity of SLC22A11. Antagonists include

proteins, nucleic acids, carbohydrates, antibodies, small molecules, or any other molecule which decrease the activity of SLC22A11.

5 The term "modulate", as it appears herein, refers to a change in the activity of SLC22A11 polypeptide. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of SLC22A11.

10 As used herein, the terms "specific binding" or "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein recognized by the binding molecule (i.e., the antigenic determinant or epitope). For example, if an antibody is specific for epitope "A" the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing
15 free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The invention provides methods (also referred to herein as "screening assays") for identifying compounds which can be used for the treatment of cardiovascular
20 diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. The methods entail the identification of candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other molecules) which bind to SLC22A11 and/or have a stimulatory or inhibitory effect on the biological activity of SLC22A11
25 or its expression and then determining which of these compounds have an effect on symptoms or diseases regarding the cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in an *in vivo* assay.

30 Candidate or test compounds or agents which bind to SLC22A11 and/or have a stimulatory or inhibitory effect on the activity or the expression of SLC22A11 are

identified either in assays that employ cells which express SLC22A11 on the cell surface (cell-based assays) or in assays with isolated SLC22A11 (cell-free assays). The various assays can employ a variety of variants of SLC22A11 (e.g., full-length SLC22A11, a biologically active fragment of SLC22A11, or a fusion protein which includes all or a portion of SLC22A11). Moreover, SLC22A11 can be derived from any suitable mammalian species (e.g., human SLC22A11, rat SLC22A11 or murine SLC22A11). The assay can be a binding assay entailing direct or indirect measurement of the binding of a test compound or a known SLC22A11 ligand to SLC22A11. The assay can also be an activity assay entailing direct or indirect measurement of the activity of SLC22A11. The assay can also be an expression assay entailing direct or indirect measurement of the expression of SLC22A11 mRNA or SLC22A11 protein. The various screening assays are combined with an *in vivo* assay entailing measuring the effect of the test compound on the symptoms of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a membrane-bound (cell surface expressed) form of SLC22A11. Such assays can employ full-length SLC22A11, a biologically active fragment of SLC22A11, or a fusion protein which includes all or a portion of SLC22A11. As described in greater detail below, the test compound can be obtained by any suitable means, e.g., from conventional compound libraries. Determining the ability of the test compound to bind to a membrane-bound form of SLC22A11 can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the SLC22A11 -expressing cell can be measured by detecting the labeled compound in a complex. For example, the test compound can be labelled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, the test compound can be enzymatically labelled with, for example, horseradish peroxidase,

alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

5 In a competitive binding format, the assay comprises contacting SLC22A11 expressing cell with a known compound which binds to SLC22A11 to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the SLC22A11 expressing cell, wherein determining the ability of the test compound to interact with the SLC22A11 expressing cell comprises determining the ability of the test compound to preferentially bind the SLC22A11 expressing cell as compared to the known compound.

15 In another embodiment, the assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of SLC22A11 (e.g., full-length SLC22A11, a biologically active fragment of SLC22A11, or a fusion protein which includes all or a portion of SLC22A11) expressed on the cell surface with a test compound and determining the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the membrane-bound form of SLC22A11. Determining the ability of the test compound to modulate the activity of the membrane-bound form of SLC22A11 can be accomplished by any method suitable for measuring the activity of SLC22A11, e.g., any method suitable for measuring the activity of an organic ion transporter (described in greater detail below). The activity of a seven-transmembrane receptor can be measured in a number of ways, not all of which are suitable for any given receptor. Among the measures of activity are: alteration in intracellular Ca^{2+} concentration, activation of phospholipase C, alteration in intracellular inositol triphosphate (IP_3) concentration, alteration in intracellular diacylglycerol (DAG) concentration, and alteration in intracellular adenosine cyclic 3', 5'-monophosphate (cAMP) concentration.

30 Determining the ability of the test compound to modulate the activity of SLC22A11 can be accomplished, for example, by determining the ability of SLC22A11 to bind

to or interact with a target molecule. The target molecule can be a molecule with which SLC22A11 binds or interacts with in nature, for example, a molecule on the surface of a cell which expresses SLC22A11, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. The target molecule can be a component of a signal transduction pathway which facilitates transduction of an extracellular signal (e.g., a signal generated by binding of a SLC22A11 ligand, through the cell membrane and into the cell. The target SLC22A11 molecule can be, for example, a second intracellular protein which has catalytic activity or a protein which facilitates the association of downstream signaling molecules with SLC22A11.

Determining the ability of SLC22A11 to bind to or interact with a target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of a polypeptide of the invention to bind to or interact with a target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (e.g., intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (e.g., a regulatory element that is responsive to a polypeptide of the invention operably linked to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular response.

The present invention also includes cell-free assays. Such assays involve contacting a form of SLC22A11 (e.g., full-length SLC22A11, a biologically active fragment of SLC22A11, or a fusion protein comprising all or a portion of SLC22A11) with a test compound and determining the ability of the test compound to bind to SLC22A11. Binding of the test compound to SLC22A11 can be determined either directly or indirectly as described above. In one embodiment, the assay includes contacting SLC22A11 with a known compound which binds SLC22A11 to form an assay mixture, contacting the assay mixture with a test compound, and determining the

ability of the test compound to interact with SLC22A11, wherein determining the ability of the test compound to interact with SLC22A11 comprises determining the ability of the test compound to preferentially bind to SLC22A11 as compared to the known compound.

5

The cell-free assays of the present invention are amenable to use of either a membrane-bound form of SLC22A11 or a soluble fragment thereof. In the case of cell-free assays comprising the membrane-bound form of the polypeptide, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the polypeptide is maintained in solution. Examples of such solubilizing agents include but are not limited to non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton X-100, Triton X-114, Thesit, Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-
10 [(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO),
15 or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

In various embodiments of the above assay methods of the present invention, it may be desirable to immobilize SLC22A11 (or a SLC22A11 target molecule) to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to SLC22A11, or interaction of SLC22A11 with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test
20 tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, glutathione-S-transferase (GST) fusion proteins or glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical; St. Louis, Mo.) or glutathione derivatized microtitre plates,
25 which are then combined with the test compound or the test compound and either the non-adsorbed target protein or SLC22A11, and the mixture incubated under
30

conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components and complex formation is measured either directly or indirectly, for example, as described above. Alternatively, the complexes can be
5 dissociated from the matrix, and the level of binding or activity of SLC22A11 can be determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either SLC22A11 or its target
10 molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated polypeptide of the invention or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals; Rockford, Ill.), and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies reactive with
15 SLC22A11 or target molecules but which do not interfere with binding of the polypeptide of the invention to its target molecule can be derivatized to the wells of the plate, and unbound target or polypeptide of the invention trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of
20 complexes using antibodies reactive with SLC22A11 or target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with SLC22A11 or target molecule.

The screening assay can also involve monitoring the expression of SLC22A11. For
25 example, regulators of expression of SLC22A11 can be identified in a method in which a cell is contacted with a candidate compound and the expression of SLC22A11 protein or mRNA in the cell is determined. The level of expression of SLC22A11 protein or mRNA the presence of the candidate compound is compared to the level of expression of SLC22A11 protein or mRNA in the absence of the
30 candidate compound. The candidate compound can then be identified as a regulator of expression of SLC22A11 based on this comparison. For example, when expres-

sion of SLC22A11 protein or mRNA protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of SLC22A11 protein or mRNA expression. Alternatively, when expression of SLC22A11 protein or mRNA is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of SLC22A11 protein or mRNA expression. The level of SLC22A11 protein or mRNA expression in the cells can be determined by methods described below.

10 *Binding Assays*

For binding assays, the test compound is preferably a small molecule which binds to and occupies the active site of SLC22A11 polypeptide, thereby making the ligand binding site inaccessible to substrate such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules. Potential ligands which bind to a polypeptide of the invention include, but are not limited to, the natural ligands of known SLC22A11 ion transporter and analogues or derivatives thereof.

20 In binding assays, either the test compound or the SLC22A11 polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to SLC22A11 polypeptide can then be accomplished, for example, by direct counting of radioemmission, by
25 scintillation counting, or by determining conversion of an appropriate substrate to a detectable product. Alternatively, binding of a test compound to a SLC22A11 polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a SLC22A11 polypeptide. A microphysiometer (*e.g.*, Cytosensor™) is an analytical
30 instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate

can be used as an indicator of the interaction between a test compound and SLC22A11 [Haseloff, (1988)].

5 Determining the ability of a test compound to bind to SLC22A11 also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [McConnell, (1992); Sjolander, (1991)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

10

In yet another aspect of the invention, a SLC22A11-like polypeptide can be used as a “bait protein“ in a two-hybrid assay or three-hybrid assay [Szabo, (1995); U.S. 5,283,317), to identify other proteins which bind to or interact with SLC22A11 and modulate its activity.

15

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding SLC22A11 can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct a DNA sequence that encodes an unidentified protein (“prey“ or “sample“) can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the “bait“ and the “prey“ proteins are able to interact *in vivo* to form a protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with SLC22A11.

20
25
30

It may be desirable to immobilize either the SLC22A11 (or polynucleotide) or the test compound to facilitate separation of the bound form from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the SLC22A11-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach SLC22A11-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to SLC22A11 (or a polynucleotide encoding for SLC22A11) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, SLC22A11 is a fusion protein comprising a domain that allows binding of SLC22A11 to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed SLC22A11; the mixture is then incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either SLC22A11

(or a polynucleotide encoding SLC22A11) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated SLC22A11 (or a polynucleotide encoding biotinylated SLC22A11) or test compounds can be prepared from biotin-NHS (N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated plates (Pierce Chemical). Alternatively, antibodies which specifically bind to SLC22A11, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of SLC22A11, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to SLC22A11 polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of SLC22A11 polypeptide, and SDS gel electrophoresis under non-reducing conditions.

Screening for test compounds which bind to a SLC22A11 polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a SLC22A11 polypeptide or polynucleotide can be used in a cell-based assay system. A SLC22A11 polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to SLC22A11 or a polynucleotide encoding SLC22A11 is determined as described above.

25

Functional Assays

Test compounds can be tested for the ability to increase or decrease SLC22A11 activity of a SLC22A11 transporter polypeptide. The SLC22A11 activity can be measured, for example, using methods described in the specific examples, below. SLC22A11 activity can be measured after contacting either a purified SLC22A11, a

30

cell membrane preparation, or an intact cell with a test compound. A test compound which decreases SLC22A11 activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing SLC22A11 activity. A test compound which increases SLC22A11 activity by at least
5 about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing SLC22A11 activity.

One such screening procedure involves the use of melanophores which are transfected to express SLC22A11. Such a screening technique is described in PCT
10 WO 92/01810 published Feb. 6, 1992. Thus, for example, such an assay may be employed for screening for a compound which inhibits activation of the transporter polypeptide of the present invention by contacting the melanophore cells which encode the transporter with both the transporter ligand and a compound to be screened. Inhibition of the signal generated by the ligand indicates that a compound
15 is a potential antagonist for the transporter, *i.e.*, inhibits activation of the transporter. The screen may be employed for identifying a compound which activates the transporter by contacting such cells with compounds to be screened and determining whether each compound generates a signal, *i.e.*, activates the transporter.

20 Other screening techniques include the use of cells which express SLC22A11 (for example, transfected CHO cells) in a system which measures extracellular pH changes caused by transporter activation [Iwabuchi et al. (1993)]. For example, compounds may be contacted with a cell which expresses the transporter polypeptide of the present invention and a second messenger response, *e.g.*, signal transduction or
25 pH changes, can be measured to determine whether the potential compound activates or inhibits the transporter. Another such screening technique involves introducing RNA encoding SLC22A11 into *Xenopus* oocytes to transiently express the transporter. The transporter oocytes can then be contacted with the transporter ligand and a compound to be screened, followed by detection of inhibition or activation of a
30 calcium signal in the case of screening for compounds which are thought to inhibit activation of the transporter.

Gene Expression

In another embodiment, test compounds which increase or decrease SLC22A11 gene
5 expression are identified. As used herein, the term "correlates with expression of a
polynucleotide" indicates that the detection of the presence of nucleic acids, the same
or related to a nucleic acid sequence encoding SLC22A11, by northern analysis or
relatime PCR is indicative of the presence of nucleic acids encoding SLC22A11 in a
10 sample, and thereby correlates with expression of the transcript from the poly-
nucleotide encoding SLC22A11. The term "microarray", as used herein, refers to an
array of distinct polynucleotides or oligonucleotides arrayed on a substrate, such as
paper, nylon or any other type of membrane, filter, chip, glass slide, or any other
suitable solid support. A SLC22A11 polynucleotide is contacted with a test
15 compound, and the expression of an RNA or polypeptide product of SLC22A11
polynucleotide is determined. The level of expression of appropriate mRNA or
polypeptide in the presence of the test compound is compared to the level of
expression of mRNA or polypeptide in the absence of the test compound. The test
compound can then be identified as a regulator of expression based on this
20 comparison. For example, when expression of mRNA or polypeptide is greater in
the presence of the test compound than in its absence, the test compound is identified
as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively,
when expression of the mRNA or polypeptide is less in the presence of the test
compound than in its absence, the test compound is identified as an inhibitor of the
mRNA or polypeptide expression.

25

The level of SLC22A11 mRNA or polypeptide expression in the cells can be
determined by methods well known in the art for detecting mRNA or polypeptide.
Either qualitative or quantitative methods can be used. The presence of polypeptide
products of SLC22A11 polynucleotide can be determined, for example, using a
30 variety of techniques known in the art, including immunochemical methods such as
radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively,

polypeptide synthesis can be determined *in vivo*, in a cell culture, or in an *in vitro* translation system by detecting incorporation of labelled amino acids into SLC22A11.

- 5 Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses SLC22A11 polynucleotide can be used in a cell-based assay system. The SLC22A11 polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line can be used.

10

Test Compounds

Suitable test compounds for use in the screening assays of the invention can be obtained from any suitable source, e.g., conventional compound libraries. The test
15 compounds can also be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological
20 library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds [Lam, (1997)]. Examples of methods for the synthesis of molecular libraries can be found in the art. Libraries of compounds may be presented in solution or on beads, bacteria, spores, plasmids or phage.

25

Modeling of Regulators

Computer modeling and searching technologies permit identification of compounds, or the improvement of already identified compounds, that can modulate SLC22A11
30 expression or activity. Having identified such a compound or composition, the active sites or regions are identified. Such active sites might typically be ligand binding

sites, such as the interaction domain of the ligand with SLC22A11. The active site can be identified using methods known in the art including, for example, from the amino acid sequences of peptides, from the nucleotide sequences of nucleic acids, or from study of complexes of the relevant compound or composition with its natural
5 ligand. In the latter case, chemical or X-ray crystallographic methods can be used to find the active site by finding where on the factor the complexed ligand is found.

Next, the three dimensional geometric structure of the active site is determined. This can be done by known methods, including X-ray crystallography, which can
10 determine a complete molecular structure. On the other hand, solid or liquid phase NMR can be used to determine certain intramolecular distances. Any other experimental method of structure determination can be used to obtain partial or complete geometric structures. The geometric structures may be measured with a complexed ligand, natural or artificial, which may increase the accuracy of the active
15 site structure determined.

If an incomplete or insufficiently accurate structure is determined, the methods of computer based numerical modeling can be used to complete the structure or improve its accuracy. Any recognized modeling method may be used, including
20 parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models. For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups, are necessary, and can be selected from force fields
25 known in physical chemistry. The incomplete or less accurate experimental structures can serve as constraints on the complete and more accurate structures computed by these modeling methods.

Finally, having determined the structure of the active site, either experimentally, by
30 modeling, or by a combination, candidate modulating compounds can be identified by searching databases containing compounds along with information on their

molecular structure. Such a search seeks compounds having structures that match the determined active site structure and that interact with the groups defining the active site. Such a search can be manual, but is preferably computer assisted. These compounds found from this search are potential SLC22A11 modulating compounds.

5

Alternatively, these methods can be used to identify improved modulating compounds from an already known modulating compound or ligand. The composition of the known compound can be modified and the structural effects of modification can be determined using the experimental and computer modeling methods described above applied to the new composition. The altered structure is then compared to the active site structure of the compound to determine if an improved fit or interaction results. In this manner systematic variations in composition, such as by varying side groups, can be quickly evaluated to obtain modified modulating compounds or ligands of improved specificity or activity.

15

Therapeutic Indications and Methods

It was found by the present applicant that SLC22A11 is expressed in various human tissues.

20

Neurology

CNS disorders include disorders of the central nervous system as well as disorders of the peripheral nervous system.

25

CNS disorders include, but are not limited to brain injuries, cerebrovascular diseases and their consequences, Parkinson's disease, corticobasal degeneration, motor neuron disease, dementia, including ALS, multiple sclerosis, traumatic brain injury, stroke, post-stroke, post-traumatic brain injury, and small-vessel cerebrovascular disease.

30

Dementias, such as Alzheimer's disease, vascular dementia, dementia with Lewy bodies, frontotemporal dementia and Parkinsonism linked to chromosome 17,

frontotemporal dementias, including Pick's disease, progressive nuclear palsy, corticobasal degeneration, Huntington's disease, thalamic degeneration, Creutzfeld-Jakob dementia, HIV dementia, schizophrenia with dementia, and Korsakoff's psychosis, within the meaning of the definition are also considered to be CNS disorders.

Similarly, cognitive-related disorders, such as mild cognitive impairment, age-associated memory impairment, age-related cognitive decline, vascular cognitive impairment, attention deficit disorders, attention deficit hyperactivity disorders, and memory disturbances in children with learning disabilities are also considered to be CNS disorders.

Pain, within the meaning of this definition, is also considered to be a CNS disorder. Pain can be associated with CNS disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, phantom feeling, reflex sympathetic dystrophy (RSD), trigeminal neuralgiaradioculopathy, post-surgical pain, HIV/AIDS related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic neuropathy secondary to connective tissue disease), paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with peripheral nerve damage, central pain (i.e. due to cerebral ischemia) and various chronic pain i.e., lumbago, back pain (low back pain), inflammatory and/or rheumatic pain. Headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania are also CNS disorders.

Visceral pain such as pancreatitis, intestinal cystitis, dysmenorrhea, irritable Bowel syndrome, Crohn's disease, biliary colic, ureteral colic, myocardial infarction and pain syndromes of the pelvic cavity, e.g., vulvodynia, orchialgia, urethral syndrome and protatodynia are also CNS disorders.

5

Also considered to be a disorder of the nervous system are acute pain, for example postoperative pain, and pain after trauma.

The human SLC22A11 is highly expressed in the following brain tissues: Alzheimer
10 brain, Alzheimer brain cerebellum, Alzheimer cerebral cortex, postcentral gyrus, hippocampus, posteroventral thalamus, dorsomedial thalamus, spinal cord (ventral horn), spinal cord (dorsal horn), neuroblastoma IMR32 cells, glial tumor H4 cells, HEK CNS + APP, neural progenitor cells, astrocytes. The expression in brain tissues and in particular the differential expression between diseased tissue Alzheimer brain
15 and healthy tissue brain, between diseased tissue Alzheimer brain cerebellum and healthy tissue, between diseased tissue Alzheimer cerebral cortex and healthy tissue cerebral cortex, between diseased tissue HEK CNS + APP and healthy tissue demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose nervous system diseases. Additionally the activity of the human SLC22A11 can be
20 modulated to treat nervous system diseases.

Cardiovascular Disorders

Heart failure is defined as a pathophysiological state in which an abnormality of
25 cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failures such as high-output and low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

30 Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously

narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included as well as the acute treatment of MI and the prevention of complications.

5 Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina and asymptomatic ischemia.

10 Arrhythmias include all forms of atrial and ventricular tachyarrhythmias, atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, ventricular fibrillation, as well as bradycardic forms of arrhythmias.

15 Hypertensive vascular diseases include primary as well as all kinds of secondary arterial hypertension, renal, endocrine, neurogenic, others. The genes may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications arising from cardiovascular diseases.

20 Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon and venous disorders.

25 Atherosclerosis is a cardiovascular disease in which the vessel wall is remodeled, compromising the lumen of the vessel. The atherosclerotic remodeling process involves accumulation of cells, both smooth muscle cells and monocyte/macrophage inflammatory cells, in the intima of the vessel wall. These cells take up lipid, likely from the circulation, to form a mature atherosclerotic lesion. Although the formation
30 of these lesions is a chronic process, occurring over decades of an adult human life, the majority of the morbidity associated with atherosclerosis occurs when a lesion

ruptures, releasing thrombogenic debris that rapidly occludes the artery. When such an acute event occurs in the coronary artery, myocardial infarction can ensue, and in the worst case, can result in death.

5 The formation of the atherosclerotic lesion can be considered to occur in five overlapping stages such as migration, lipid accumulation, recruitment of inflammatory cells, proliferation of vascular smooth muscle cells, and extracellular matrix deposition. Each of these processes can be shown to occur in man and in animal models of atherosclerosis, but the relative contribution of each to the
10 pathology and clinical significance of the lesion is unclear.

Thus, a need exists for therapeutic methods and agents to treat cardiovascular pathologies, such as atherosclerosis and other conditions related to coronary artery disease.

15 Cardiovascular diseases include but are not limited to disorders of the heart and the vascular system like congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, peripheral vascular diseases, and atherosclerosis.

20 Too high or too low levels of fats in the bloodstream, especially cholesterol, can cause long-term problems. The risk to develop atherosclerosis and coronary artery or carotid artery disease (and thus the risk of having a heart attack or stroke) increases with the total cholesterol level increasing. Nevertheless, extremely low cholesterol
25 levels may not be healthy. Examples of disorders of lipid metabolism are hyperlipidemia (abnormally high levels of fats (cholesterol, triglycerides, or both) in the blood, may be caused by family history of hyperlipidemia), obesity, a high-fat diet, lack of exercise, moderate to high alcohol consumption, cigarette smoking, poorly controlled diabetes, and an underactive thyroid gland), hereditary hyperlipidemias
30 (type I hyperlipoproteinemia (familial hyperchylomicronemia), type II hyperlipoproteinemia (familial hypercholesterolemia), type III hyperlipoproteinemia, type IV

hyperlipoproteinemia, or type V hyperlipoproteinemia), hypolipoproteinemia, lipidoses (caused by abnormalities in the enzymes that metabolize fats), Gaucher's disease, Niemann-Pick disease, Fabry's disease, Wolman's disease, cerebrotendinous xanthomatosis, sitosterolemia, Refsum's disease, or Tay-Sachs disease.

5

Kidney disorders may lead to hypertension or hypotension. Examples for kidney problems possibly leading to hypertension are renal artery stenosis, pyelonephritis, glomerulonephritis, kidney tumors, polycystic kidney disease, injury to the kidney, or radiation therapy affecting the kidney. Excessive urination may lead to hypotension.

10

The human SLC22A11 is highly expressed in the following cardiovascular related tissues: heart, heart atrium (left), fetal aorta, artery, coronary artery endothel cells, aortic endothel cells, liver liver cirrhosis, thrombocytes, HEK CNS + APP, fetal kidney, kidney. Expression in the above mentioned tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of cardiovascular diseases. Additionally the activity of the human SLC22A11 can be modulated to treat cardiovascular diseases.

15

The human SLC22A11 is highly expressed in liver tissues: liver liver cirrhosis. Expression in liver tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of dyslipidemia disorders as an cardiovascular disorder. Additionally the activity of the human SLC22A11 can be modulated to treat - but not limited to - dyslipidemia disorders.

20

The human SLC22A11 is highly expressed in kidney tissues : HEK CNS + APP, fetal kidney, kidney. Expression in kidney tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of blood pressure disorders as an cardiovascular disorder. Additionally the activity of the human SLC22A11 can be modulated to treat - but not limited to - blood pressure disorders as hypertension or hypotension.

25
30

Hematological Disorders

Hematological disorders comprise diseases of the blood and all its constituents as well as diseases of organs and tissues involved in the generation or degradation of all the constituents of the blood. They include but are not limited to 1) Anemias, 2) Myeloproliferative Disorders, 3) Hemorrhagic Disorders, 4) Leukopenia, 5) Eosinophilic Disorders, 6) Leukemias, 7) Lymphomas, 8) Plasma Cell Dyscrasias, 9) Disorders of the Spleen in the course of hematological disorders. Disorders according to 1) include, but are not limited to anemias due to defective or deficient hem synthesis, deficient erythropoiesis. Disorders according to 2) include, but are not limited to polycythemia vera, tumor-associated erythrocytosis, myelofibrosis, thrombocythemia. Disorders according to 3) include, but are not limited to vasculitis, thrombocytopenia, heparin-induced thrombocytopenia, thrombotic thrombocytopenic purpura, hemolytic-uremic syndrome, hereditary and acquired disorders of platelet function, hereditary coagulation disorders. Disorders according to 4) include, but are not limited to neutropenia, lymphocytopenia. Disorders according to 5) include, but are not limited to hypereosinophilia, idiopathic hypereosinophilic syndrome. Disorders according to 6) include, but are not limited to acute myeloic leukemia, acute lymphoblastic leukemia, chronic myelocytic leukemia, chronic lymphocytic leukemia, myelodysplastic syndrome. Disorders according to 7) include, but are not limited to Hodgkin's disease, non-Hodgkin's lymphoma, Burkitt's lymphoma, mycosis fungoides cutaneous T-cell lymphoma. Disorders according to 8) include, but are not limited to multiple myeloma, macroglobulinemia, heavy chain diseases. In extension of the preceding idiopathic thrombocytopenic purpura, iron deficiency anemia, megaloblastic anemia (vitamin B12 deficiency), aplastic anemia, thalassemia, malignant lymphoma bone marrow invasion, malignant lymphoma skin invasion, hemolytic uremic syndrome, giant platelet disease are considered to be hematological diseases too.

The human SLC22A11 is highly expressed in the following tissues of the hematological system: Jurkat (T-cells), erythrocytes, thrombocytes, bone marrow

CD71+ cells, bone marrow CD34+ cells, bone marrow CD15+ cells, cord blood CD71+ cells, cord blood CD34+ cells, neutrophils cord blood, neutrophils peripheral blood. The expression in the above mentioned tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of hematological diseases. Additionally the activity of the human SLC22A11 can be modulated to treat hematological disorders.

Gastrointestinal and Liver Diseases

Gastrointestinal diseases comprise primary or secondary, acute or chronic diseases of the organs of the gastrointestinal tract which may be acquired or inherited, benign or malignant or metaplastic, and which may affect the organs of the gastrointestinal tract or the body as a whole. They comprise but are not limited to 1) disorders of the esophagus like achalasia, vigorous achalasia, dysphagia, cricopharyngeal incoordination, pre-esophageal dysphagia, diffuse esophageal spasm, globus sensation, Barrett's metaplasia, gastroesophageal reflux, 2) disorders of the stomach and duodenum like functional dyspepsia, inflammation of the gastric mucosa, gastritis, stress gastritis, chronic erosive gastritis, atrophy of gastric glands, metaplasia of gastric tissues, gastric ulcers, duodenal ulcers, neoplasms of the stomach, 3) disorders of the pancreas like acute or chronic pancreatitis, insufficiency of the exocrine or endocrine tissues of the pancreas like steatorrhea, diabetes, neoplasms of the exocrine or endocrine pancreas like 3.1) multiple endocrine neoplasia syndrome, ductal adenocarcinoma, cystadenocarcinoma, islet cell tumors, insulinoma, gastrinoma, carcinoid tumors, glucagonoma, Zollinger-Ellison syndrome, Vipoma syndrome, malabsorption syndrome, 4) disorders of the bowel like chronic inflammatory diseases of the bowel, Crohn's disease, ileus, diarrhea and constipation, colonic inertia, megacolon, malabsorption syndrome, ulcerative colitis, 4.1) functional bowel disorders like irritable bowel syndrome, 4.2) neoplasms of the bowel like familial polyposis, adenocarcinoma, primary malignant lymphoma, carcinoid tumors, Kaposi's sarcoma, polyps, cancer of the colon and rectum.

Liver diseases comprise primary or secondary, acute or chronic diseases or injury of the liver which may be acquired or inherited, benign or malignant, and which may affect the liver or the body as a whole. They comprise but are not limited to disorders of the bilirubin metabolism, jaundice, syndromes of Gilbert's, Crigler-Najjar, Dubin-Johnson and Rotor; intrahepatic cholestasis, hepatomegaly, portal hypertension, ascites, Budd-Chiari syndrome, portal-systemic encephalopathy, fatty liver, steatosis, Reye's syndrome, liver diseases due to alcohol, alcoholic hepatitis or cirrhosis, fibrosis and cirrhosis, fibrosis and cirrhosis of the liver due to inborn errors of metabolism or exogenous substances, storage diseases, syndromes of Gaucher's, Zellweger's, Wilson's - disease, acute or chronic hepatitis, viral hepatitis and its variants, inflammatory conditions of the liver due to viruses, bacteria, fungi, protozoa, helminths; drug induced disorders of the liver, chronic liver diseases like primary sclerosing cholangitis, alpha₁-antitrypsin-deficiency, primary biliary cirrhosis, postoperative liver disorders like postoperative intrahepatic cholestasis, hepatic granulomas, vascular liver disorders associated with systemic disease, benign or malignant neoplasms of the liver, disturbance of liver metabolism in the new-born or prematurely born.

The human SLC22A11 is highly expressed in the following tissues of the gastroenterological system: ileum, ileum chronic inflammation, rectum, liver liver cirrhosis. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue ileum chronic inflammation and healthy tissue ileum, between diseased tissue liver liver cirrhosis and healthy tissue liver demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of gastroenterological disorders. Additionally the activity of the human SLC22A11 can be modulated to treat gastroenterological disorders.

Inflammatory Diseases

Inflammatory diseases comprise diseases triggered by cellular or non-cellular mediators of the immune system or tissues causing the inflammation of body tissues

and subsequently producing an acute or chronic inflammatory condition. Examples for such inflammatory diseases are hypersensitivity reactions of type I – IV, for example but not limited to hypersensitivity diseases of the lung including asthma, atopic diseases, allergic rhinitis or conjunctivitis, angioedema of the lids, hereditary
5 angioedema, antireceptor hypersensitivity reactions and autoimmune diseases, Hashimoto's thyroiditis, systemic lupus erythematosus, Goodpasture's syndrome, pemphigus, myasthenia gravis, Grave's and Raynaud's disease, type B insulin-resistant diabetes, rheumatoid arthritis, psoriasis, Crohn's disease, scleroderma, mixed connective tissue disease, polymyositis, sarcoidosis, glomerulonephritis, acute
10 or chronic host versus graft reactions.

The human SLC22A11 is highly expressed in the following tissues of the immune system and tissues responsive to components of the immune system as well as in the following tissues responsive to mediators of inflammation: ileum chronic
15 inflammation, liver liver cirrhosis, bone marrow CD15+ cells, neutrophils cord blood, neutrophils peripheral blood. The expression in the above mentioned tissues and in particular the differential expression between diseased tissue ileum chronic inflammation and healthy tissue ileum, between diseased tissue liver liver cirrhosis and healthy tissue liver demonstrates that the human SLC22A11 or mRNA can be
20 utilized to diagnose of inflammatory diseases. Additionally the activity of the human SLC22A11 can be modulated to treat inflammatory diseases.

Disorders Related to Pulmology

25 Asthma is thought to arise as a result of interactions between multiple genetic and environmental factors and is characterized by three major features: 1) intermittent and reversible airway obstruction caused by bronchoconstriction, increased mucus production, and thickening of the walls of the airways that leads to a narrowing of the airways, 2) airway hyperresponsiveness, and 3) airway inflammation. Certain cells
30 are critical to the inflammatory reaction of asthma and they include T cells and antigen presenting cells, B cells that produce IgE, and mast cells, basophils,

eosinophils, and other cells that bind IgE. These effector cells accumulate at the site of allergic reaction in the airways and release toxic products that contribute to the acute pathology and eventually to tissue destruction related to the disorder. Other resident cells, such as smooth muscle cells, lung epithelial cells, mucus-producing cells, and nerve cells may also be abnormal in individuals with asthma and may contribute to its pathology. While the airway obstruction of asthma, presenting clinically as an intermittent wheeze and shortness of breath, is generally the most pressing symptom of the disease requiring immediate treatment, the inflammation and tissue destruction associated with the disease can lead to irreversible changes that eventually make asthma a chronic and disabling disorder requiring long-term management.

Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway obstruction due to chronic bronchitis [Botstein, 1980]. Emphysema is characterised by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of COPD is cigarette smoking, although the disease does also occur in non-smokers.

The human SLC22A11 is highly expressed in the following tissues of the respiratory system: bone marrow CD15+ cells, neutrophils cord blood, neutrophils peripheral blood, secondary bronchia, bronchial epithelial cells, bronchial smooth muscle cells. The expression in the above mentioned tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of respiratory diseases. Additionally the activity of the human SLC22A11 can be modulated to treat those diseases.

Disorders Related to Urology

Genitourinary disorders comprise benign and malign disorders of the organs constituting the genitourinary system of female and male, renal diseases like acute or chronic renal failure, immunologically mediated renal diseases like renal transplant rejection, lupus nephritis, immune complex renal diseases, glomerulopathies, nephritis, toxic nephropathy, obstructive uropathies like benign prostatic hyperplasia (BPH), neurogenic bladder syndrome, urinary incontinence like urge-, stress-, or overflow incontinence, pelvic pain, and erectile dysfunction.

10

The human SLC22A11 is highly expressed in the following urological tissues: spinal cord (ventral horn), spinal cord (dorsal horn), ureter, penis, corpus cavernosum, fetal kidney, kidney. The expression in the above mentioned tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of urological disorders. Additionally the activity of the human SLC22A11 can be modulated to treat urological disorders.

15

The human SLC22A11 is highly expressed in spinal cord tissues: spinal cord (ventral horn), spinal cord (dorsal horn). Expression in spinal cord tissues demonstrates that the human SLC22A11 or mRNA can be utilized to diagnose of incontinence as an urological disorder. The spinal cord tissues are involved in the neuronal regulation of the urological system. Additionally the activity of the human SLC22A11 can be modulated to treat - but not limited to - incontinence.

20

Applications

25

The present invention provides for both prophylactic and therapeutic methods for cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

30

The regulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of SLC22A11. An agent that modulates activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of the polypeptide, a peptide, a peptidomimetic, or
5 any small molecule. In one embodiment, the agent stimulates one or more of the biological activities of SLC22A11. Examples of such stimulatory agents include the active SLC22A11 and nucleic acid molecules encoding a portion of SLC22A11. In another embodiment, the agent inhibits one or more of the biological activities of SLC22A11. Examples of such inhibitory agents include antisense nucleic acid
10 molecules and antibodies. These regulatory methods can be performed in vitro (e.g., by culturing the cell with the agent) or, alternatively, in vivo (e.g. by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by unwanted expression or activity of SLC22A11 or a protein in the SLC22A11 signaling
15 pathway. In one embodiment, the method involves administering an agent like any agent identified or being identifiable by a screening assay as described herein, or combination of such agents that modulate say upregulate or downregulate the expression or activity of SLC22A11 or of any protein in the SLC22A11 signaling pathway. In another embodiment, the method involves administering a regulator of
20 SLC22A11 as therapy to compensate for reduced or undesirably low expression or activity of SLC22A11 or a protein in the SLC22A11 signaling pathway.

Stimulation of activity or expression of SLC22A11 is desirable in situations in which activity or expression is abnormally low and in which increased activity is likely to
25 have a beneficial effect. Conversely, inhibition of activity or expression of SLC22A11 is desirable in situations in which activity or expression of SLC22A11 is abnormally high and in which decreasing its activity is likely to have a beneficial effect.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are hereby incorporated by reference.

5 **Pharmaceutical Compositions**

This invention further pertains to novel agents identified by the above-described screening assays and uses thereof for treatments as described herein.

10 The nucleic acid molecules, polypeptides, and antibodies (also referred to herein as “active compounds”) of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language “pharmaceutically acceptable carrier” is intended to
15 include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in
20 the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

The invention includes pharmaceutical compositions comprising a regulator of SLC22A11 expression or activity (and/or a regulator of the activity or expression of a
25 protein in the SLC22A11 signaling pathway) as well as methods for preparing such compositions by combining one or more such regulators and a pharmaceutically acceptable carrier. Also within the invention are pharmaceutical compositions comprising a regulator identified using the screening assays of the invention packaged with instructions for use. For regulators that are antagonists of SLC22A11
30 activity or which reduce SLC22A11 expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, gastro-

enterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. For regulators that are agonists of SLC22A11 activity or increase SLC22A11 expression, the instructions would specify use of the pharmaceutical composition for treatment of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory
5 diseases, neurological diseases and urological diseases.

An antagonist of SLC22A11 may be produced using methods which are generally known in the art. In particular, purified SLC22A11 may be used to produce
10 antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind SLC22A11. Antibodies to SLC22A11 may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies like
15 those which inhibit dimer formation are especially preferred for therapeutic use.

In another embodiment of the invention, the polynucleotides encoding SLC22A11, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding SLC22A11 may be used in
20 situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding SLC22A11. Thus, complementary molecules or fragments may be used to modulate SLC22A11 activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense
25 oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding SLC22A11.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide
30 sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors which will express

nucleic acid sequence complementary to the polynucleotides of the gene encoding SLC22A11. These techniques are described, for example, in [Scott and Smith (1990) Science 249:386-390].

- 5 Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

10 An additional embodiment of the invention relates to the administration of a pharmaceutical composition containing SLC22A11 in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of SLC22A11, antibodies to SLC22A11, and mimetics, agonists, antagonists, or inhibitors of SLC22A11. The compositions may be administered alone or in combination with at least one other agent, such as
15 stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

20 A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the
25 following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for
30 the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral

preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

5 Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EMTM (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, a pharmaceutically acceptable polyol like glycerol, propylene glycol, liquid polyethylene glycol, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin. Sterile injectable solutions can be prepared by incorporating the active compound (e.g., a polypeptide or antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and

10
15
20
25
30

freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

5 Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed.

10

Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient
15 such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

20 For administration by inhalation, the compounds are delivered in the form of an aerosol spray from a pressurized container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For
25 transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the
30 active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

5

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration. For pharmaceutical compositions which include an antagonist of SLC22A11 activity, a compound which reduces

30

expression of SLC22A11, or a compound which reduces expression or activity of a protein in the SLC22A11 signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases. For pharmaceutical compositions which include an agonist of SLC22A11 activity, a compound which increases expression of SLC22A11, or a compound which increases expression or activity of a protein in the SLC22A11 signaling pathway or any combination thereof, the instructions for administration will specify use of the composition for cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

Diagnostics

In another embodiment, antibodies which specifically bind SLC22A11 may be used for the diagnosis of disorders characterized by the expression of SLC22A11, or in assays to monitor patients being treated with SLC22A11 or agonists, antagonists, and inhibitors of SLC22A11. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for SLC22A11 include methods which utilize the antibody and a label to detect SLC22A11 in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent joining with a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring SLC22A11, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of SLC22A11 expression. Normal or standard values for SLC22A11 expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to SLC22A11 under

conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, preferably by photometric means. Quantities of SLC22A11 expressed in subject samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values
5 establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding SLC22A11 may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and
10 PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of SLC22A11 may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of SLC22A11, and to monitor regulation of SLC22A11 levels during therapeutic intervention.

15 Polynucleotide sequences encoding SLC22A11 may be used for the diagnosis of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases associated with expression of SLC22A11. The polynucleotide sequences encoding
20 SLC22A11 may be used in Southern, Northern, or dot-blot analysis, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patient biopsies to detect altered SLC22A11 expression. Such qualitative or quantitative methods are well known in the art.

25 In a particular aspect, the nucleotide sequences encoding SLC22A11 may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding SLC22A11 may be labelled by standard methods and added to a fluid or tissue sample from a patient under
30 conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared

with a standard value. If the amount of signal in the patient sample is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding SLC22A11 in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases associated with expression of SLC22A11, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding SLC22A11, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with SLC22A11, or fragments thereof, and washed. Bound SLC22A11 is then detected by methods well known in the art. Purified SLC22A11 can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-

neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

5 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding SLC22A11 specifically compete with a testcompound for binding SLC22A11. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with SLC22A11.

10 **Determination of a Therapeutically Effective Dose**

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases SLC22A11 activity relative to
15 SLC22A11 activity which occurs in the absence of the therapeutically effective dose. For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses
20 and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g., ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental
25 animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating
30 concentrations that include the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the

patient, and the route of administration. The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

5

Normal dosage amounts can vary from 0.1 micrograms to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc. If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either *ex vivo* or *in vivo* using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, "gene gun", and DEAE- or calcium phosphate-mediated transfection.

10

15

20

25

30

If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above. Preferably, a reagent reduces expression of SLC22A11 gene or the activity of SLC22A11 by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of SLC22A11 gene or the activity of

SLC22A11 can be assessed using methods well known in the art, such as hybridization of nucleotide probes to SLC22A11-specific mRNA, quantitative RT-PCR, immunologic detection of SLC22A11, or measurement of SLC22A11 activity.

5 In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above.
10 Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects. Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses,
15 rabbits, monkeys, and most preferably, humans.

Nucleic acid molecules of the invention are those nucleic acid molecules which are contained in a group of nucleic acid molecules consisting of (i) nucleic acid molecules encoding a polypeptide comprising the amino acid sequence of SEQ ID
20 NO: 2, (ii) nucleic acid molecules comprising the sequence of SEQ ID NO: 1, (iii) nucleic acid molecules having the sequence of SEQ ID NO: 1, (iv) nucleic acid molecules the complementary strand of which hybridizes under stringent conditions to a nucleic acid molecule of (i), (ii), or (iii); and (v) nucleic acid molecules the sequence of which differs from the sequence of a nucleic acid molecule of (iii) due to
25 the degeneracy of the genetic code, wherein the polypeptide encoded by said nucleic acid molecule has SLC22A11 activity.

Polypeptides of the invention are those polypeptides which are contained in a group of polypeptides consisting of (i) polypeptides having the sequence of SEQ ID NO: 2,
30 (ii) polypeptides comprising the sequence of SEQ ID NO: 2, (iii) polypeptides encoded by nucleic acid molecules of the invention and (iv) polypeptides which show

- 76 -

at least 99%, 98%, 95%, 90%, or 80% homology with a polypeptide of (i), (ii), or (iii), wherein said purified polypeptide has SLC22A11 activity.

5 An object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a SLC22A11 polypeptide, (ii) detect binding of said test compound to said SLC22A11
10 polypeptide. E.g., compounds that bind to the SLC22A11 polypeptide are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of
15 cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a SLC22A11 polypeptide at a certain concentration of a test compound or in the absence of said test compound, (ii) determining the activity of said polypeptide at a different
20 concentration of said test compound. E.g., compounds that lead to a difference in the activity of the SLC22A11 polypeptide in (i) and (ii) are identified potential therapeutic agents for such a disease.

Another object of the invention is a method of screening for therapeutic agents useful
25 in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the activity of a SLC22A11 polypeptide at a certain concentration of a test compound, (ii) determining the activity of
30 a SLC22A11 polypeptide at the presence of a compound known to be a regulator of a SLC22A11 polypeptide. E.g., compounds that show similar effects on the activity of

the SLC22A11 polypeptide in (i) as compared to compounds used in (ii) are identified potential therapeutic agents for such a disease.

5 Other objects of the invention are methods of the above, wherein the step of contacting is in or at the surface of a cell.

Other objects of the invention are methods of the above, wherein the cell is in vitro.

10 Other objects of the invention are methods of the above, wherein the step of contacting is in a cell-free system.

Other objects of the invention are methods of the above, wherein the polypeptide is coupled to a detectable label.

15 Other objects of the invention are methods of the above, wherein the compound is coupled to a detectable label.

20 Other objects of the invention are methods of the above, wherein the test compound displaces a ligand which is first bound to the polypeptide.

Other objects of the invention are methods of the above, wherein the polypeptide is attached to a solid support.

25 Other objects of the invention are methods of the above, wherein the compound is attached to a solid support.

30 Another object of the invention is a method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) contacting a test compound with a SLC22A11

polynucleotide, (ii) detect binding of said test compound to said SLC22A11 polynucleotide. Compounds that, e.g., bind to the SLC22A11 polynucleotide are potential therapeutic agents for the treatment of such diseases.

5 Another object of the invention is the method of the above, wherein the nucleic acid molecule is RNA.

Another object of the invention is a method of the above, wherein the contacting step is in or at the surface of a cell.

10

Another object of the invention is a method of the above, wherein the contacting step is in a cell-free system.

15 Another object of the invention is a method of the above, wherein the polynucleotide is coupled to a detectable label.

Another object of the invention is a method of the above, wherein the test compound is coupled to a detectable label.

20 Another object of the invention is a method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) determining the amount of a SLC22A11 polynucleotide in a sample taken from said mammal, (ii)
25 determining the amount of SLC22A11 polynucleotide in healthy and/or diseased mammal. A disease is diagnosed, e.g., if there is a substantial similarity in the amount of SLC22A11 polynucleotide in said test mammal as compared to a diseased mammal.

30 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases,

gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which binds to a SLC22A11 polypeptide.

5 Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a SLC22A11 polypeptide.

10

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a SLC22A11 polypeptide, wherein
15 said therapeutic agent is (i) a small molecule, (ii) an RNA molecule, (iii) an antisense oligonucleotide, (iv) a polypeptide, (v) an antibody, or (vi) a ribozyme.

20

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a SLC22A11 polynucleotide.

25

Another object of the invention is a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a SLC22A11 polypeptide.

30

Another object of the invention is the use of regulators of a SLC22A11 for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal.

Another object of the invention is a method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of (i) identifying a regulator of SLC22A11, (ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal; and (iii) combining of said regulator with an acceptable pharmaceutical carrier.

Another object of the invention is the use of a regulator of SLC22A11 for the regulation of SLC22A11 activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.

The expression of human SLC22A11 in inflammation and urological related tissues (as described above) suggests a particular – but not limited to – utilization SLC22A11 for diagnosis and modulation of inflammation and urological diseases. Furthermore the above described expression suggest a – but not limited to – utilization SLC22A11 to cardiovascular diseases, gastroenterological diseases, hematological diseases, respiratory diseases and neurological diseases.

The examples below are provided to illustrate the subject invention. These examples are provided by way of illustration and are not included for the purpose of limiting the invention.

Examples

Example 1: Search for homologous sequences in public sequence data bases

5 The degree of homology can readily be calculated by known methods. Preferred methods to determine homology are designed to give the largest match between the sequences tested. Methods to determine homology are codified in publicly available computer programs such as BestFit, BLASTP, BLASTN, and FASTA. The BLAST programs are publicly available from NCBI and other sources in the internet.

10

For SLC22A11 the following hits to known sequences were identified by using the BLAST algorithm [Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ; Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402] and the following set of parameters: matrix = BLOSUM62 and low complexity filter. The following
15 databases were searched: NCBI (non-redundant database) and DERWENT patent database (Geneseq).

The following hits were found:

20

>dbj|BD092019.1| Placenta type organic anion transporter and its gene
Length = 2210, Score = 4213 bits (2191), Expect = 0.0, Identities = 2191/2191
(100%)

25

>dbj|E54995.1| Placental organic anion transporter and gene thereof
Length = 2210, Score = 4213 bits (2191), Expect = 0.0, Identities = 2191/2191
(100%)

30

>dbj|AB026116.1| Homo sapiens hOAT4 mRNA for organic anion transporter 4,
complete cds
Length = 2210, Score = 4213 bits (2191), Expect = 0.0, Identities = 2191/2191
(100%)

- >gb|AC128661.3| Mus musculus chromosome 7 clone RP24-567N23, complete sequence
Length = 421625747, Score = 4211 bits (2190), Expect = 0.0, Identities = 2190/2190
5 (100%)
- >gb|BC034384.1| Homo sapiens, solute carrier family 22 (organic anion/cationtransporter), member 11, clone MGC:34282 IMAGE:5190509,mRNA, complete cds
10 Length = 2545, Score = 4211 bits (2190), Expect = 0.0, Identities = 2190/2190 (100%)
- >NA2001A:AAD12607 Aad12607 Human protein having hydrophobic domainencoding cDNA clone HP03882. 9/2001
15 Length = 2194, Score = 4209 bits (2189), Expect = 0.0, Identities = 2189/2189 (100%)
- >emb|AX191620.1| Sequence 142 from Patent WO0149728
Length = 2194, Score = 4209 bits (2189), Expect = 0.0, Identities = 2189/2189
20 (100%)
- >NA2002:AAD27262 Aad27262 Human transporter and ion channel-9(TRICH-9) cDNA. 4/2002
Length = 2182, Score = 4169 bits (2168), Expect = 0.0, Identities = 2168/2168
25 (100%)
- >emb|AX709965.1| Sequence 36 from Patent WO0192304
Length = 2182, Score = 4169 bits (2168), Expect = 0.0, Identities = 2168/2168
30 (100%)

>NA2001A:AAK94561 Aak94561 Human full-length cDNA, SEQ ID NO:
3467.11/2001

Length = 1871, Score = 3592 bits (1868), Expect = 0.0, Identities = 1870/1871
(99%)

5

>dbj|BD127590.1| Primer for synthesizing full-length cDNA and use thereof

Length = 1871, Score = 3592 bits (1868), Expect = 0.0, Identities = 1870/1871
(99%)

10 >dbj|AK075224.1| Homo sapiens cDNA FLJ90743 fis, clone PLACE1011236,
weakly similar toMus musculus mRNA for RST

Length = 1871, Score = 3592 bits (1868), Expect = 0.0, Identities = 1870/1871
(99%)

15 >emb|AX191610.1| Sequence 132 from Patent WO0149728

Length = 1653, Score = 3178 bits (1653), Expect = 0.0, Identities = 1653/1653
(100%)

20 >NA2002:ABL55086 Abl55086 Human cDNA sequence #1 from clone
HLWBT09.10/2002

Length = 2100, Score = 2534 bits (1318), Expect = 0.0, Identities = 1320/1325
(99%)

Example 2: Expression profiling

25

Total cellular RNA was isolated from cells by one of two standard methods: 1)
guanidine isothiocyanate/Cesium chloride density gradient centrifugation [Kellogg,
(1990)]; or with the Tri-Reagent protocol according to the manufacturer's specifica-
tions (Molecular Research Center, Inc., Cincinnati, Ohio). Total RNA prepared by
30 the Tri-reagent protocol was treated with DNase I to remove genomic DNA
contamination.

For relative quantitation of the mRNA distribution of SLC22A11, total RNA from each cell or tissue source was first reverse transcribed. 85 µg of total RNA was reverse transcribed using 1 µmole random hexamer primers, 0.5 mM each of dATP, dCTP, dGTP and dTTP (Qiagen, Hilden, Germany), 3000 U RnaseQut (Invitrogen, Groningen, Netherlands) in a final volume of 680 µl. The first strand synthesis buffer and Omniscript reverse transcriptase (2 u/µl) were from (Qiagen, Hilden, Germany). The reaction was incubated at 37°C for 90 minutes and cooled on ice. The volume was adjusted to 6800 µl with water, yielding a final concentration of 12.5 ng/µl of starting RNA.

For relative quantitation of the distribution of SLC22A11 mRNA in cells and tissues the Applied Biosystems 7900 HT Sequence Detection system or Biorad iCycler was used according to the manufacturer's specifications and protocols. PCR reactions were set up to quantitate SLC22A11 and the housekeeping genes HPRT (hypoxanthine phosphoribosyltransferase), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), β-actin, and others. Forward and reverse primers and probes for SLC22A11 were designed using the Perkin Elmer ABI Primer Express™ software and were synthesized by TibMolBiol (Berlin, Germany). The SLC22A11 forward primer sequence was: Primer1 (SEQ ID NO: 3). The SLC22A11 reverse primer sequence was Primer2 (SEQ ID NO: 4). Probe1 (SEQ ID NO: 5), labelled with FAM (carboxyfluorescein succinimidyl ester) as the reporter dye and TAMRA (carboxytetramethylrhodamine) as the quencher, is used as a probe for SLC22A11. The following reagents were prepared in a total of 25 µl : 1x TaqMan buffer A, 5.5 mM MgCl₂, 200 nM of dATP, dCTP, dGTP, and dUTP, 0.025 U/µl AmpliTaq Gold™, 0.01 U/ µl AmpErase and Probe1 (SEQ ID NO: 4), SLC22A11 forward and reverse primers each at 200 nM, 200 nM SLC22A11 FAM/TAMRA-labelled probe, and 5 µl of template cDNA. Thermal cycling parameters were 2 min at 50°C, followed by 10 min at 95°C, followed by 40 cycles of melting at 95°C for 15 sec and annealing/extending at 60°C for 1 min.

Calculation of corrected CT values

5 The CT (threshold cycle) value is calculated as described in the “Quantitative determination of nucleic acids” section. The CF-value (factor for threshold cycle correction) is calculated as follows :

1. PCR reactions were set up to quantitate the housekeeping genes (HKG) for each cDNA sample.
- 10 2. CT_{HKG} -values (threshold cycle for housekeeping gene) were calculated as described in the “Quantitative determination of nucleic acids” section.
3. CT_{HKG} -mean values (CT mean value of all HKG tested on one cDNAs) of all HKG for each cDNA are calculated (n = number of HKG):
- 15

$$CT_{HKG-n}\text{-mean value} = (CT_{HKG1}\text{-value} + CT_{HKG2}\text{-value} + \dots + CT_{HKG-n}\text{-value}) / n$$

4. CT_{panel} mean value (CT mean value of all HKG in all tested cDNAs) =
 - 20
- $$(CT_{HKG1}\text{-mean value} + CT_{HKG2}\text{-mean value} + \dots + CT_{HKG-y}\text{-mean value}) / y$$
- (y = number of cDNAs)

5. CF_{cDNA-n} (correction factor for cDNA n) = $CT_{\text{panel}}\text{-mean value} - CT_{HKG-n}\text{-mean value}$
- 25
6. CT_{cDNA-n} (CT value of the tested gene for the cDNA n) + CF_{cDNA-n} (correction factor for cDNA n) = $CT_{\text{cor-cDNA-n}}$ (corrected CT value for a gene on cDNA n)

Calculation of relative expression

Definition : highest $CT_{\text{cor-cDNA-n}} \neq 40$ is defined as $CT_{\text{cor-cDNA}} [\text{high}]$

Relative Expression = $2^{(CT_{\text{cor-cDNA}}[\text{high}] - CT_{\text{cor-cDNA-n}})}$

5

Tissues

The expression of SLC22A11 was investigated in the tissues listed in table 1.

10 *Expression profile*

The results of the the mRNA-quantification (expression profiling) is shown in Table 1.

15 **Table 1:** *Relative expression of SLC22A11 in various human tissues.*

	fetal heart	1
	heart	79
	pericardium	15
20	heart atrium (right)	2
	heart atrium (left)	202
	heart ventricle (left)	85
	heart ventricle (right)	1
	heart apex	1
25	Purkinje fibers	1
	interventricular septum	1
	fetal aorta	78
	aorta	1
	aorta sclerotic	37
30	artery	108
	coronary artery	1
	coronary artery sclerotic	2
	pulmonary artery	2
	carotid artery	1
35	mesenteric artery	1
	vein	44
	pulmonic valve	73
	coronary artery endothel cells	3373
	coronary artery smooth muscle primary cells	76
40	aortic smooth muscle cells	1
	pulmonary artery smooth muscle cells	8

- 88 -

	aortic endothel cells	584
	HUVEC cells	7
	pulmonary artery endothel cells	13
	iliac artery endothel cells	15
5	skin	15
	adrenal gland	251
	thyroid	38
10	thyroid tumor	10
	pancreas	80
	pancreas liver cirrhosis	171
	esophagus	103
15	esophagus tumor	1
	stomach	1
	stomach tumor	28
	colon	50
	colon tumor	10
20	small intestine	58
	ileum	2916
	ileum tumor	47
	ileum chronic inflammation	5713
	rectum	1361
25	salivary gland	1
	fetal liver	23
	liver	105
	liver liver cirrhosis	2048
	liver lupus disease	1
30	liver tumor	174
	HEP G2 cells	329
	leukocytes (peripheral blood)	25
	Jurkat (T-cells)	333
35	bone marrow	1
	erythrocytes	405
	lymphnode	1
	thymus	4
	thrombocytes	317
40	bone marrow stromal cells	16
	bone marrow CD71+ cells	391
	bone marrow CD33+ cells	9
	bone marrow CD34+ cells	440
	bone marrow CD15+ cells	422
45	cord blood CD71+ cells	526
	cord blood CD34+ cells	898
	neutrophils cord blood	2180
	neutrophils peripheral blood	3492
	spleen	83
50	spleen liver cirrhosis	111

	skeletal muscle	55
	cartilage	92
	bone connective tissue	45
	adipose	18
5	fetal brain	18
	brain	28
	Alzheimer brain	244
	cerebellum	70
10	Alzheimer brain cerebellum	380
	cerebellum (right)	23
	cerebellum (left)	139
	cerebral cortex	9
	Alzheimer cerebral cortex	1046
15	frontal lobe	23
	Alzheimer brain frontal lobe	29
	occipital lobe	65
	parietal lobe	23
	temporal lobe	53
20	postcentral gyrus	198
	vermis cerebelli	21
	pons	1
	substantia nigra	23
	caudatum	44
25	cerebral meninges	1
	cerebral peduncles	1
	corpus callosum	11
	nucleus accumbens	1
	putamen	17
30	hippocampus	282
	thalamus	1
	posteroventral thalamus	1783
	dorsalmedial thalamus	4771
	hypothalamus	9
35	dorsal root ganglia	5
	spinal cord	56
	spinal cord (ventral horn)	4608
	spinal cord (dorsal horn)	11911
	neuroblastoma SK-N-MC cells	54
40	neuroblastoma SH-SY5Y cells	97
	neuroblastoma IMR32 cells	838
	glial tumor H4 cells	396
	HEK CNS + APP	209
	neural progenitor cells	1652
45	astrocytes	744
	retina	168
	fetal lung	72
	fetal lung fibroblast IMR-90 cells	1
50	fetal lung fibroblast MRC-5 cells	1
	lung	1

- 90 -

	lung right upper lobe	53
	lung right mid lobe	1
	lung right lower lobe	100
	lung lupus disease	1
5	lung tumor	29
	lung COPD	17
	trachea	56
	primary bronchia	1
	secondary bronchia	1911
10	bronchial epithelial cells	2034
	bronchial smooth muscle cells	568
	small airway epithelial cells	1
	cervix	87
15	testis	393
	HeLa cells (cervix tumor)	1
	placenta	6654
	uterus	6
	uterus tumor	10
20	ovary	1
	ovary tumor	8
	breast	39
	breast tumor	393
	MDA MB 231 cells (breast tumor)	7
25	mammary gland	12
	prostate	21
	prostate BPH	1
	prostate tumor	622
30	bladder	1
	ureter	182
	penis	969
	corpus cavernosum	1468
	fetal kidney	1833
35	kidney	14664
	kidney tumor	91
	renal epithelial cells	24
	HEK 293 cells	16

40 **Example 3: Antisense Analysis**

Knowledge of the correct, complete cDNA sequence coding for SLC22A11 enables its use as a tool for antisense technology in the investigation of gene function. Oligonucleotides, cDNA or genomic fragments comprising the antisense strand of a polynucleotide coding for SLC22A11 are used either in vitro or in vivo to inhibit translation of the mRNA. Such technology is now well known in the art, and

antisense molecules can be designed at various locations along the nucleotide sequences. By treatment of cells or whole test animals with such antisense sequences, the gene of interest is effectively turned off. Frequently, the function of the gene is ascertained by observing behavior at the intracellular, cellular, tissue or organismal level (e.g., lethality, loss of differentiated function, changes in morphology, etc.).

In addition to using sequences constructed to interrupt transcription of a particular open reading frame, modifications of gene expression is obtained by designing antisense sequences to intron regions, promoter/enhancer elements, or even to trans-acting regulatory genes.

Example 4: Expression of SLC22A11

Expression of SLC22A11 is accomplished by subcloning the cDNAs into appropriate expression vectors and transfecting the vectors into expression hosts such as, e.g., *E. coli*. In a particular case, the vector is engineered such that it contains a promoter for β -galactosidase, upstream of the cloning site, followed by sequence containing the amino-terminal Methionine and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is an engineered bacteriophage promoter useful for artificial priming and transcription and for providing a number of unique endonuclease restriction sites for cloning.

Induction of the isolated, transfected bacterial strain with Isopropyl- β -D-thio-galactopyranoside (IPTG) using standard methods produces a fusion protein corresponding to the first seven residues of β -galactosidase, about 15 residues of "linker", and the peptide encoded within the cDNA. Since cDNA clone inserts are generated by an essentially random process, there is probability of 33% that the included cDNA will lie in the correct reading frame for proper translation. If the cDNA is not in the proper reading frame, it is obtained by deletion or insertion of the appropriate number of bases using well known methods including in vitro

mutagenesis, digestion with exonuclease III or mung bean nuclease, or the inclusion of an oligonucleotide linker of appropriate length.

5 The SLC22A11 cDNA is shuttled into other vectors known to be useful for expression of proteins in specific hosts. Oligonucleotide primers containing cloning sites as well as a segment of DNA (about 25 bases) sufficient to hybridize to stretches at both ends of the target cDNA is synthesized chemically by standard methods. These primers are then used to amplify the desired gene segment by PCR. The resulting gene segment is digested with appropriate restriction enzymes under
10 standard conditions and isolated by gel electrophoresis. Alternately, similar gene segments are produced by digestion of the cDNA with appropriate restriction enzymes. Using appropriate primers, segments of coding sequence from more than one gene are ligated together and cloned in appropriate vectors. It is possible to optimize expression by construction of such chimeric sequences.

15 Suitable expression hosts for such chimeric molecules include, but are not limited to, mammalian cells such as Chinese Hamster Ovary (CHO) and human 293 cells., insect cells such as Sf9 cells, yeast cells such as *Saccharomyces cerevisiae* and bacterial cells such as *E. coli*. For each of these cell systems, a useful expression
20 vector also includes an origin of replication to allow propagation in bacteria, and a selectable marker such as the β -lactamase antibiotic resistance gene to allow plasmid selection in bacteria. In addition, the vector may include a second selectable marker such as the neomycin phosphotransferase gene to allow selection in transfected eukaryotic host cells. Vectors for use in eukaryotic expression hosts require RNA
25 processing elements such as 3' polyadenylation sequences if such are not part of the cDNA of interest.

30 Additionally, the vector contains promoters or enhancers which increase gene expression. Such promoters are host specific and include MMTV, SV40, and metallothionine promoters for CHO cells; trp, lac, tac and T7 promoters for bacterial hosts; and alpha factor, alcohol oxidase and PGH promoters for yeast. Transcription

enhancers, such as the rous sarcoma virus enhancer, are used in mammalian host cells. Once homogeneous cultures of recombinant cells are obtained through standard culture methods, large quantities of recombinantly produced SLC22A11 are recovered from the conditioned medium and analyzed using chromatographic methods known in the art. For example, SLC22A11 can be cloned into the expression vector pcDNA3, as exemplified herein. This product can be used to transform, for example, HEK293 or COS by methodology standard in the art. Specifically, for example, using Lipofectamine (Gibco BRL catalog no. 18324-020) mediated gene transfer.

10

Example 5: Isolation of Recombinant SLC22A11

SLC22A11 is expressed as a chimeric protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals [Appa Rao, 1997] and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Washington). The inclusion of a cleavable linker sequence such as Factor Xa or enterokinase (Invitrogen, Groningen, The Netherlands) between the purification domain and the SLC22A11 sequence is useful to facilitate expression of SLC22A11.

15
20

Example 6: Production of SLC22A11 Specific Antibodies

Two approaches are utilized to raise antibodies to SLC22A11, and each approach is useful for generating either polyclonal or monoclonal antibodies. In one approach, denatured protein from reverse phase HPLC separation is obtained in quantities up to 75 mg. This denatured protein is used to immunize mice or rabbits using standard protocols; about 100 μ g are adequate for immunization of a mouse, while up to 1 mg might be used to immunize a rabbit. For identifying mouse hybridomas, the denatured protein is radioiodinated and used to screen potential murine B-cell

25
30

hybridomas for those which produce antibody. This procedure requires only small quantities of protein, such that 20 mg is sufficient for labeling and screening of several thousand clones.

5 In the second approach, the amino acid sequence of an appropriate SLC22A11 domain, as deduced from translation of the cDNA, is analyzed to determine regions of high antigenicity. Oligopeptides comprising appropriate hydrophilic regions are synthesized and used in suitable immunization protocols to raise antibodies. The optimal amino acid sequences for immunization are usually at the C-terminus, the N-
10 terminus and those intervening, hydrophilic regions of the polypeptide which are likely to be exposed to the external environment when the protein is in its natural conformation.

Typically, selected peptides, about 15 residues in length, are synthesized using an
15 Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to keyhole limpet hemocyanin (KLH; Sigma, St. Louis, MO) by reaction with M-maleimidobenzoyl-N-hydroxysuccinimide ester, MBS. If necessary, a cysteine is introduced at the N-terminus of the peptide to permit coupling to KLH. Rabbits are immunized with the peptide-KLH complex in complete Freund's
20 adjuvant. The resulting antisera are tested for antipeptide activity by binding the peptide to plastic, blocking with 1% bovine serum albumin, reacting with antisera, washing and reacting with labeled (radioactive or fluorescent), affinity purified, specific goat anti-rabbit IgG.

25 Hybridomas are prepared and screened using standard techniques. Hybridomas of interest are detected by screening with labeled SLC22A11 to identify those fusions producing the monoclonal antibody with the desired specificity. In a typical protocol, wells of plates (FAST; Becton-Dickinson, Palo Alto, CA) are coated during incubation with affinity purified, specific rabbit anti-mouse (or suitable antispecies 1
30 g) antibodies at 10 mg/ml. The coated wells are blocked with 1% bovine serum albumin, (BSA), washed and incubated with supernatants from hybridomas. After

washing the wells are incubated with labeled SLC22A11 at 1 mg/ml. Supernatants with specific antibodies bind more labeled SLC22A11 than is detectable in the background. Then clones producing specific antibodies are expanded and subjected to two cycles of cloning at limiting dilution. Cloned hybridomas are injected into
5 pristane-treated mice to produce ascites, and monoclonal antibody is purified from mouse ascitic fluid by affinity chromatography on Protein A. Monoclonal antibodies with affinities of at least

10⁸ M⁻¹, preferably 10⁹ to 10¹⁰ M⁻¹ or stronger, are typically made by standard
10 procedures.

Example 7: Diagnostic Test Using SLC22A11 Specific Antibodies

Particular SLC22A11 antibodies are useful for investigating signal transduction and
15 the diagnosis of infectious or hereditary conditions which are characterized by differences in the amount or distribution of SLC22A11 or downstream products of an active signaling cascade.

Diagnostic tests for SLC22A11 include methods utilizing antibody and a label to
20 detect SLC22A11 in human body fluids, membranes, cells, tissues or extracts of such. The polypeptides and antibodies of the present invention are used with or without modification. Frequently, the polypeptides and antibodies are labeled by joining them, either covalently or noncovalently, with a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known
25 and have been reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent agents, chemiluminescent agents, chromogenic agents, magnetic particles and the like.

30 A variety of protocols for measuring soluble or membrane-bound SLC22A11, using either polyclonal or monoclonal antibodies specific for the protein, are known in the

art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmuno-
assay (RIA) and fluorescent activated cell sorting (FACS). A two-site monoclonal-
based immunoassay utilizing monoclonal antibodies reactive to two non-interfering
epitopes on SLC22A11 is preferred, but a competitive binding assay may be
5 employed.

Example 8: Purification of Native SLC22A11 Using Specific Antibodies

Native or recombinant SLC22A11 is purified by immunoaffinity chromatography
10 using antibodies specific for SLC22A11. In general, an immunoaffinity column is
constructed by covalently coupling the anti-TRH antibody to an activated
chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation
15 with ammonium sulfate or by purification on immobilized Protein A (Pharmacia
LKB Biotechnology, Piscataway N.J.). Likewise, monoclonal antibodies are
prepared from mouse ascites fluid by ammonium sulfate precipitation or
chromatography on immobilized Protein A. Partially purified immunoglobulin is
covalently attached to a chromatographic resin such as CnBr-activated Sepharose
20 (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is
blocked, and the derivative resin is washed according to the manufacturer's
instructions.

Such immunoaffinity columns are utilized in the purification of SLC22A11 by pre-
25 paring a fraction from cells containing SLC22A11 in a soluble form. This
preparation is derived by solubilization of whole cells or of a subcellular fraction
obtained via differential centrifugation (with or without addition of detergent) or by
other methods well known in the art. Alternatively, soluble SLC22A11 containing a
signal sequence is secreted in useful quantity into the medium in which the cells are
30 grown.

A soluble SLC22A11-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of SLC22A11 (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/protein
5 binding (e.g., a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and SLC22A11 is collected.

Example 9: Drug Screening

10 Test compounds can be screened for the ability to bind to OAT-like polypeptides or polynucleotides or to affect OAT-like activity or OAT-like gene expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well microtiter
15 plates. The wells of the microtiter plates typically require assay volumes that range from 50 to 500 microliter. In addition to the plates, many instruments, materials, pipettors, robotics, plate washers, and plate readers are commercially available to fit the 96-well format.

20 Alternatively, "free format assays," or assays that have no physical barrier between samples, can be used. For example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by [Jayawickreme *et al.*, (1994)]. The cells are placed under agarose in petri dishes, then beads that carry combinatorial compounds are placed on the surface of the
25 agarose. The combinatorial compounds are partially released the compounds from the beads. Active compounds can be visualized as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

30 Another example of a free format assay is described by Chelsky, "Strategies for Screening Combinatorial Libraries: Novel and Traditional Approaches," reported at

the First Annual Conference of The Society for Biomolecular Screening in Philadelphia, Pa. (Nov. 7-10, 1995). Chelsky placed a simple homogenous enzyme assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel. Thereafter, beads carrying
5 combinatorial compounds via a photolinker were placed inside the gel and the compounds were partially released by UV-light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

Yet another example is described by [Salmon *et al.*, (1996)]. In this example,
10 combinatorial libraries were screened for compounds that had cytotoxic effects on cancer cells growing in agar.

Another high throughput screening method is described in Beutel *et al.*, U.S. Patent 5,976,813. In this method, test samples are placed in a porous matrix. One or more
15 assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support. When samples are introduced to the porous matrix they diffuse sufficiently slowly, such that the assays can be performed without the test samples running together.

20 **Example 10: Binding Assay**

For binding assays, the test compound is preferably a small molecule which binds to a OAT-like polypeptide, thereby reducing the normal biological activity of the OAT-like polypeptide. Examples of such small molecules include, but are not limited to,
25 small peptides or peptide-like molecules. In binding assays, either the test compound or the OAT-like polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to the OAT-like polypeptide can then be accomplished, for example, by direct counting
30 of radioemmission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product.

Alternatively, binding of a test compound to a OAT-like polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a OAT-like polypeptide. A microphysiometer (*e.g.*, Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and a OAT-like polypeptide [McConnell *et al.*, (1992)].

10

Determining the ability of a test compound to bind to a OAT-like polypeptide also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [Sjolander & Urbaniczky, (1991), and Szabo *et al.*, (1995)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (*e.g.*, BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

15

In yet another aspect of the invention, a OAT-like polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent 5,283,317; [Iwabuchi *et al.*, (1993)]; and W094/10300), to identify other proteins which bind to or interact with the OAT-like polypeptide and modulate its activity.

20

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding a OAT-like polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact *in vivo* to

25
30

form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with the OAT-like polypeptide.

It may be desirable to immobilize either the OAT-like polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the OAT-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach the OAT-like polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a OAT-like polypeptide (or polynucleotide) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, the OAT-like polypeptide is a fusion protein comprising a domain that allows the OAT-like polypeptide to be bound to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed OAT-like polypeptide; the mixture is then

incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the
5 complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a OAT-like polypeptide (or polynucleotide) or a test compound can be immobilized utilizing
10 conjugation of biotin and streptavidin. Biotinylated OAT-like polypeptides (or polynucleotides) or test compounds can be prepared from biotin-NHS(N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a
15 OAT-like polypeptide, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of the OAT-like polypeptide, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

20 Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to the OAT-like polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of the OAT-like polypeptide, and SDS gel electrophoresis under non-reducing conditions.

25 Screening for test compounds which bind to a OAT-like polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a OAT-like polypeptide or polynucleotide can be used in a cell-based assay system. A OAT-like polynucleotide can be naturally occurring in the cell or can be introduced using
30 techniques such as those described above. Binding of the test compound to a OAT-like polypeptide or polynucleotide is determined as described above.

Purified OAT-like polypeptides comprising a glutathione-S-transferase protein and absorbed onto glutathione-derivatized wells of 96-well microtiter plates are contacted with test compounds from a small molecule library at pH 7.0 in a physiological
5 buffer solution. OAT-like polypeptides comprise an amino acid sequence shown in any one or more of SEQ ID NO:6 to 10. The test compounds comprise a fluorescent tag. The samples are incubated for 5 minutes to one hour. Control samples are incubated in the absence of a test compound.

10 The buffer solution containing the test compounds is washed from the wells. Binding of a test compound to a OAT-like polypeptide is detected by fluorescence measurements of the contents of the wells. A test compound which increases the fluorescence in a well by at least 15% relative to fluorescence of a well in which a test compound is not incubated is identified as a compound which binds to an ADO-
15 ribosylation factor-related polypeptide.

Example 11: Functional activity

Functional assays can be carried out as described in the specific examples, after
20 contacting either a purified OAT-like polypeptide or an intact cell with a test compound. A test compound which decreases a functional activity of a human OAT-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for decreasing OAT-like protein activity. A test compound which increases a functional activity by at least about 10,
25 preferably about 50, more preferably about 75, 90, or 100% is identified as a potential agent for increasing OAT-like protein activity.

Example 12: Effect of test compound on anoinic transport

Transport of ^3H -labeled substrate in the presence or absence of a test compound can be measured as described in [Hsiang *et al.* (1999, *supra*)]. Briefly, 293c18 cells are
5 transfected with human OATP2-like expression constructs using LipofectAMINE Plus (Life Technologies, Inc.) according to the manufacturer's instructions. The medium is removed, and the cells are washed once in serum-free DMEM. ^3H -labeled
10 substrate, either alone or in the presence of a test compound, is added in the same medium and incubated at room temperature for 5-10 minutes. The cells are quickly washed once with ice-cold DMEM containing 5% bovine serum albumin, then washed three times with ice-cold DMEM. Cells are lysed in 0.1 N NaOH. Radiolabel incorporation is determined by liquid scintillation counting.

Example 13: Rational Drug Design

15 The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact, agonists, antagonists, or inhibitors. Any of these examples are used to fashion drugs which are more active or stable forms of the polypeptide or which enhance or interfere with the
20 function of a polypeptide in vivo.

In one approach, the three-dimensional structure of a protein of interest, or of a protein-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape
25 and charges of the polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of a polypeptide is gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design efficient inhibitors. Useful examples of rational drug design include molecules
30 which have improved activity or stability or which act as inhibitors, agonists, or antagonists of native peptides.

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design is based. It is possible to
5 bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids is expected to be an analog of the original receptor. The anti-id is then used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides then act as the
10 pharmacore.

By virtue of the present invention, sufficient amount of polypeptide are made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the SLC22A11 amino acid sequence provided herein provides
15 guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

Example 14: Production of Non-human Transgenic Animals

20 Animal model systems which elucidate the physiological and behavioral roles of the SLC22A11 are produced by creating nonhuman transgenic animals in which the activity of the SLC22A11 is either increased or decreased, or the amino acid sequence of the expressed SLC22A11 is altered, by a variety of techniques. Examples of these techniques include, but are not limited to: 1) Insertion of normal
25 or mutant versions of DNA encoding a SLC22A11, by microinjection, electroporation, retroviral transfection or other means well known to those skilled in the art, into appropriately fertilized embryos in order to produce a transgenic animal or 2) homologous recombination of mutant or normal, human or animal versions of these genes with the native gene locus in transgenic animals to alter the regulation of
30 expression or the structure of these SLC22A11 sequences. The technique of homologous recombination is well known in the art. It replaces the native gene with

the inserted gene and hence is useful for producing an animal that cannot express native SLC22A11s but does express, for example, an inserted mutant SLC22A11, which has replaced the native SLC22A11 in the animal's genome by recombination, resulting in underexpression of the transporter. Microinjection adds genes to the genome, but does not remove them, and the technique is useful for producing an animal which expresses its own and added SLC22A11, resulting in overexpression of the SLC22A11.

One means available for producing a transgenic animal, with a mouse as an example, is as follows: Female mice are mated, and the resulting fertilized eggs are dissected out of their oviducts. The eggs are stored in an appropriate medium such as cesiumchloride M2 medium. DNA or cDNA encoding SLC22A11 is purified from a vector by methods well known to the one skilled in the art. Inducible promoters may be fused with the coding region of the DNA to provide an experimental means to regulate expression of the transgene. Alternatively or in addition, tissue specific regulatory elements may be fused with the coding region to permit tissue-specific expression of the transgene. The DNA, in an appropriately buffered solution, is put into a microinjection needle (which may be made from capillary tubing using a piper puller) and the egg to be injected is put in a depression slide. The needle is inserted into the pronucleus of the egg, and the DNA solution is injected. The injected egg is then transferred into the oviduct of a pseudopregnant mouse which is a mouse stimulated by the appropriate hormones in order to maintain false pregnancy, where it proceeds to the uterus, implants, and develops to term. As noted above, microinjection is not the only method for inserting DNA into the egg but is used here only for exemplary purposes.

References

- U.S. 4,522,811
U.S. 5,283,317
5 U.S. 5,565,332
U.S. 5,976,813
WO 84/03564
WO 92/01810
WO 93/03151
10 WO 94/10300
WO 94/13804
WO 01/04297
WO 01/49728
WO 01/92304
15 Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ;
Nucleic Acids Res 1997 Sep 1; 25(17): 3389-402
Appa Rao et al., 1997, Protein Expr Purif Nov, 11(2): 201-8
Barnes, 2000, Chest, 117:10S14S
Botstein et al., 1980 , Am J Hum Genet. 32: 314-31
20 Cha et al., (2000), J. Biol. Chem. 275: 4507-4512
Colbere-Garapin *et al.*, 1981, *J. Mol. Biol.* 150, 1-14
Engelhard *et al.*, 1994, *Proc. Nat. Acad. Sci.* 91, 3224-3227
Enomoto et al., (2002), Nature 417: 447-452
Gergen and Weiss , 1992, Am Rev Respir Dis 146:823-824
25 Gibson et al., 1996, Genome Research 6: 995-1001
Grundemann et al., (1994), Nature 372: 549-552, 1994.
Haseloff *et al.*, 1988 , *Nature* 334, 585-591
Heid et al., 1996, Genome Research 6: 986-994
Hsiang *et al.*, 274, *J. Biol. Chem.* 37161-68, 1999
30 Holland et al., 1991, PNAS 88: 7276-7280
Iwabuchi *et al.*, 1993, *Oncogene* 8, 1693-1696

- Jayawickreme *et al.*, 19, *Proc. Natl. Acad. Sci. U.S.A.* 1614-18 (1994).
- Jeffreys *et al.*, 1985, *Nature* 316: 76-9
- Johnson *et al.*, 1989, *Endoc. Rev.* 10, 317-331
- Kellogg *et al.*, 1990, *Anal. Biochem.* 189:202-208
- 5 Konig *et al.*, 278, *Am. J. Physiol. Gastrointest. Liver Physiol.* G156-64, 2000
- Kouzuki *et al.*, 292, *J. Pharmacol. Exp. Ther.* 505-11, 2000
- Lam, 1997, *Anticancer Drug Res.* 12(3):145-67
- Leabman *et al.*, (2003), *Proc. Nat. Acad. Sci.* 100: 5896-5901
- Livak *et al.*, 1995, *PCR Methods and Applications* 357-362
- 10 Logan, Shenk, 1984, *Proc. Natl. Acad. Sci.* 81, 3655-3659
- Lowy *et al.*, 1980, *Cell* 22, 817-23
- Maddox *et al.*, 1983, *J. Exp. Med.* 158, 1211-1216
- McConnell *et al.*, 1992, *Science* 257, 1906-1912
- Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165, 81-91
- 15 Piatak *et al.*, 1993, *BioTechniques* 14:70-81
- Piatak *et al.*, 1993, *Science* 259:1749-1754
- Porath *et al.*, 1992, *Prot. Exp. Purif.* 3, 263-281
- Roberge *et al.*, 1995, *Science* 269, 202-204
- Salmon *et al.*, *Molecular Diversity* 2, 57-63 (1996)
- 20 Sjolander, Urbaniczky, 1991, *Anal. Chem.* 63, 2338-2345
- Szabo *et al.*, 1995, *Curr. Opin. Struct. Biol.* 5, 699-705
- Thomas, 1980, *Proc. Nat. Acad. Sci.*, 77:5201-5205
- Uhlmann *et al.*, 1987, *Tetrahedron. Lett.* 215, 3539-3542
- Weber *et al.*, 1990, *Genomics* 7: 524-30
- 25 Wigler *et al.*, 1977, *Cell* 11, 223-32
- Wigler *et al.*, 1980, *Proc. Natl. Acad. Sci.* 77, 3567-70

Claims

1. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular
5 diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- i) contacting a test compound with a SLC22A11 polypeptide,
10
- ii) detect binding of said test compound to said SLC22A11 polypeptide.
2. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular
15 diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- i) determining the activity of a SLC22A11 polypeptide at a certain
20 concentration of a test compound or in the absence of said test compound,
- ii) determining the activity of said polypeptide at a different concentration of said test compound.
25
3. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular
30 diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of

- 109 -

- i) determining the activity of a SLC22A11 polypeptide at a certain concentration of a test compound,
 - ii) determining the activity of a SLC22A11 polypeptide at the presence of a compound known to be a regulator of a SLC22A11 polypeptide.
4. The method of any of claims 1 to 3, wherein the step of contacting is in or at the surface of a cell.
 5. The method of any of claims 1 to 3, wherein the cell is in vitro.
 6. The method of any of claims 1 to 3, wherein the step of contacting is in a cell-free system.
 7. The method of any of claims 1 to 3, wherein the polypeptide is coupled to a detectable label.
 8. The method of any of claims 1 to 3, wherein the compound is coupled to a detectable label.
 9. The method of any of claims 1 to 3, wherein the test compound displaces a ligand which is first bound to the polypeptide.
 10. The method of any of claims 1 to 3, wherein the polypeptide is attached to a solid support.
 11. The method of any of claims 1 to 3, wherein the compound is attached to a solid support.
 12. A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular

diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of

- 5 i) contacting a test compound with a SLC22A11 polynucleotide,
- ii) detect binding of said test compound to said SLC22A11 polynucleotide.
- 10 13. The method of claim 12 wherein the nucleic acid molecule is RNA.
14. The method of claim 12 wherein the contacting step is in or at the surface of a cell.
- 15 15. The method of claim 12 wherein the contacting step is in a cell-free system.
16. The method of claim 12 wherein polynucleotide is coupled to a detectable label.
- 20 17. The method of claim 12 wherein the test compound is coupled to a detectable label.
18. A method of diagnosing a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- 25 i) determining the amount of a SLC22A11 polynucleotide in a sample taken from said mammal,

- ii) determining the amount of SLC22A11 polynucleotide in healthy and/or diseased mammals.
- 5 19. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which binds to a SLC22A11 polypeptide.
- 10 20. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a SLC22A11 polypeptide.
- 15 21. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a therapeutic agent which regulates the activity of a SLC22A11 polypeptide, wherein said therapeutic agent is
- 20
- i) a small molecule,
 - ii) an RNA molecule,
 - 25 iii) an antisense oligonucleotide,
 - iv) a polypeptide,
 - v) an antibody, or
 - vi) a ribozyme.
- 30 22. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological

diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a SLC22A11 polynucleotide.

- 5 23. A pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising a SLC22A11 polypeptide.
- 10
24. Use of regulators of a SLC22A11 for the preparation of a pharmaceutical composition for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal.
- 15
25. Method for the preparation of a pharmaceutical composition useful for the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- 20
- i) identifying a regulator of SLC22A11,
- 25
- ii) determining whether said regulator ameliorates the symptoms of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal; and
- 30
- iii) combining of said regulator with an acceptable pharmaceutical carrier.

26. Use of a regulator of SLC22A11 for the regulation of SLC22A11 activity in a mammal having a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases.
- 5

Fig. 1**SEQ ID NO: 1**

GTTCCAAACAGCAGTTAGGTCAGCAGTCCGCTCAGCCGAGGCAGCTCT
GTTTCATGGCGTTCTCGAAGCTCTTGGAGCAAGCCGGAGGCGTGGGCCT
CTTCCAGACCCTGCAGGTGCTCACCTTCATCCTCCCCTGCCTCATGAT
ACCTTCCCAGATGCTCCTGGAGAACTTCTCAGCCGCCATCCCAGGCCA
CCGATGCTGGACACACATGCTGGACAATGGCTCTGCGGTTTCCACAAA
CATGACCCCCAAGGCCCTTCTGACCATCTCCATCCCGCCAGGCCCCAA
CCAGGGGCCCCACCAGTGCCGCCGCTTCCGCCAGCCACAGTGGCAGCT
CTTGGACCCCAATGCCACGGCCACCAGCTGGAGCGAAGCTGACACGGA
GCCGTGTGTGGACGGCTGGGTCTATGACCGCAGCGTCTTCACCTCCAC
CATCGTGGCCAAGTGGGACCTGGTGTGCAGCTCCCAGGGCTTGAAGCC
CCTAAGCCAGTCCATCTTCATGTCCGGGATCCTGGTGGGCTCCTTTAT
CTGGGGCCTCCTCTCCTACCGGTTTGGGAGGAAGCCGATGCTGAGCTG
GTGCTGCCTGCAGTTGGCCGTGGCGGGCACCAGCACCATCTTCGCCCC
AACATTCGTCATCTACTGCGGCCTGCGGTTTCGTGGCCGCTTTTGGGAT
GGCCGCATCTTTCTGAGTTCACTGACACTGATGGTGGAGTGGACCAC
GACCAGCAGGAGGGCGGTCACCATGACGGTGGTGGGATGTGCCTTCAG
CGCAGGCCAGGCGGCGCTGGGCGGCCTGGCCTTTGCCCTGCGGGACTG
GAGGACTCTCCAGCTGGCAGCATCAGTGCCCTTCTTTGCCATCTCCCT
GATATCCTGGTGGCTGCCAGAATCCGCCCGGTGGCTGATTATTAAGGG
CAAACCAGACCAAGCACTTCAGGAGCTCAGAAAGGTGGCCAGGATAAA
TGGCCACAAGGAGGCCAAGAACCTGACCATAGAGGTGCTGATGTCCAG
CGTGAAGGAGGAGGTGGCCTCTGCAAAGGAGCCGCGGTCCGTGCTGGA
CCTGTTCTGCGTGCCCGTGCTCCGCTGGAGGAGCTGCGCCATGCTGGT
GGTGAATTTCTCTCTATTGATCTCCTACTATGGGCTGGTCTTCGACCT
GCAGAGCCTGGGCCGTGACATCTTCCTCCTCCAGGCCCTCTTCGGGGC
CGTGGACTTCCTGGGCCGGGCCACCACTGCCCTCTTGCTCAGTTTCCT
TGGCCGCCGCACCATCCAGGCGGGTTCACAGGCCATGGCCGGCCTCGC

CATTCTAGCCAACATGCTGGTGCCGCAAGATTTGCAGACCCTGCGTGT
GGTCTTTGCTGTGCTGGGAAAGGGATGTTTTGGGATAAGCCTAACCTG
CCTCACCATCTACAAGGCTGAACTCTTTCCAACGCCAGTGCGGATGAC
AGCAGATGGCATTCTGCATACAGTGGGCCGGCTGGGGGCTATGATGGG
TCCCCTGATCCTGATGAGCCGCCAAGCCCTGCCCTGCTGCCTCCTCT
CCTCTATGGCGTTATCTCCATTGCTTCCAGCCTGGTTGTGCTGTTCTT
CCTCCCGGAGACCCAGGGACTTCCGCTCCCTGACACTATCCAGGACCT
GGAGAGCCAGAAATCAACAGCAGCCCAGGGCAACCGGCAAGAGGCCGT
CACTGTGGAAAGTACCTCGCTCTAGAAATTGTGCCTGCATGGAGCCCC
TTTAGTCAAAGACTCCTGGAAAGGAGTTGCCTCTTCTCCAATCAGAGC
GTGGAGGCGAGTTGGGCGACTTCAAGGGCCTGGCATGGCAGAGGCCAG
GCAGCCGTGGCCGAGTGGACAGCGTGGCCGTCTGCTGTGGCTGAAGGC
AGCTTCCACAGCTCACTCCTCTTCTCCCTGCCCTGATCAGATTCCCCA
CCTTACCCGGGCCCTACAGGAGCCTGTGCAGATGGCCATGCCCAACCA
ATAACGAGACGGTTCCCCTCCCTTTCCCTGCCAGGCTCATGTCTTTAC
ACCTTCACTCAGCCACGCCAACCAGAGACTGGGTCCAATCTCACCCC
ACCACATACAGAGCCCTCATCTGTGAAATGAGAATGATCACGTGACCC
ACCCCCCAGGGCAGGTATCAGGGTGAACCTGATCTTAGCACCGGCCAAA
TAAATGGAACCTGCTGAGAGAGCTGCCAGATAAAAAAAAAAAAAAAAAA
AA

Fig. 2**SEQ ID NO: 2**

MAFSKLLLEQAGGVGLFQTLQVLTFILPCLMIPSQMLLENFSAAIPGHR
CwthMLDNGSAVSTNMTPKALLTISIIPPGPNQGP HQCRRFRQPQWQLL
DPNATATSWSEADTEPCVDGWVYDRSVFTSTIVAKWDLVCSSQGLKPL
SQSIFMSGILVGSFIWGLLSYRFGRKPMLSWCCLQLAVAGTSTIFAPT
FVIYCGLRFVAAF GMAGIFLSSLTLMVEWTTTSRRAVTMTVVGCAFSA
GQAALGGLAFALRDWRTLQLAASVPPFAISLISWWLPESARWLI IKGK
PDQALQELRKVARINGHKEAKNLTIEVLMSSVKEEVASAKEPRSVL DL
FCVPVLRWRSCAMLVVNFSL LISYYGLVFDLQSLGRDIFLLQALFGAV
DFLGRATTALLSFLGRRTIQAGSQAMAGLAILANMLVPQDLQTLRVV
FAVLGKGCFGISLTCLTIYKAELFPTPVRMTADGILHTVGR LGAMMGP
LILMSRQALPLL PPLLYGVISIASSLVVLFFLPETQGLPLPDTIQDLE
SQKSTAAQGNRQEAVTVESTSL

Fig. 3**SEQ ID NO: 3**

5' -GGCATCTTTCTGAGTTC ACTGACA- 3'

Fig. 4**SEQ ID NO: 4**

5' -CCTGCGCTGAAGGCACAT- 3'

Fig. 5**SEQ ID NO: 5**

5' -TGGACCACGACCAGCAGGAGGG- 3'

SEQUENCE LISTING

<110> Bayer AG, BHC

<120> Diagnostics and Therapeutics for Diseases Associated with Organic Cation Transporter SLC22A11 (SLC22A11)

<130> Le A 36 892

<160> 5

<170> PatentIn version 3.1

<210> 1

<211> 2210

<212> DNA

<213> Homo sapiens

<400> 1

```

gttccaaaca gcagttaggt cagcagtccg ctcagccgag gcagctctgt tcatggcggt      60
ctcgaagctc ttggagcaag cgggagcggt gggcctcttc cagaccctgc aggtgctcac      120
cttcacctc cctgcctca tgatacctc ccagatgctc ctggagaact tctcagccgc      180
catcccaggc caccgatget ggacacacat gctggacaat ggctctgagg ttccacaaa      240
catgaccccc aaggcccttc tgaccatctc catcccacca ggccccaacc aggggccccca      300
ccagtgccgc cgcttcgcc agccacagtg gcagctcttg gaccccaatg ccacggccac      360
cagctggagc gaagctgaca cggagccgtg tgtggacggc tgggtctatg accgcagcgt      420
cttcacctcc accatcgtgg ccaagtggga cctggtgtgc agctcccagg gcttgaagcc      480
cctaaaccag tccatcttca tgtccgggat cctggtgggc tcctttatct ggggcctcct      540
ctcctaccgg tttgggagga agccgatgct gagctgggtgc tgctgcagt tggccgtggc      600
gggcaccagc accatcttcc cccaacatt cgtcatctac tgcggcctgc ggttcgtggc      660
cgcttttggg atggccggca tctttctgag ttcactgaca ctgatggtgg agtggaccac      720
gaccagcagg agggcggta ccatgacggt ggtgggatgt gccttcagcg caggccaggc      780
ggcgtggggc ggccctggcct ttgccctgcg ggactggagg actctccagc tggcagcatc      840
agtgcccttc tttgccatct ccctgatata ctggtggctg ccagaatccg cccggtggct      900
gattattaag ggcaaaccag accaagcact tcaggagctc agaaagggtg ccaggataaa      960
tggccacaag gaggccaaga acctgacat agaggtgctg atgtccagcg tgaaggagga     1020
ggtggcctct gcaaaggagc cgcggtcggg gctggacctg ttctgcgtgc cgtgctccg     1080
ctggaggagc tgcgccatgc tgggtggtgaa tttctctcta ttgatctcct actatgggct     1140
ggtcttcgac ctgcagagcc tgggcccgtga catcttctc ctccaggccc tcttcggggc     1200
cgtggacttc ctgggccggg ccaccactgc cctcttgctc agtttccttg gccgccgcac     1260
catccaggcg ggttcccagg ccatggccgg cctcgcatt ctagccaaca tgctggtgcc     1320
gcaagatttg cagaccctgc gtgtggtcct tgctgtgctg ggaaagggat gttttgggat     1380
aagcctaacc tgcctcacca tctacaaggc tgaactcttt ccaacgccag tgcggatgac     1440

```

```

agcagatggc attctgcata cagtgggccc gctgggggct atgatgggtc ccctgatcct 1500
gatgagccgc caagccctgc cctgtctgcc tctctcctc tatggcggtta tctccattgc 1560
ttccagcctg gttgtgctgt tcttctccc ggagaccag ggacttccgc tccctgacac 1620
tatccaggac ctggagagcc agaaatcaac agcagcccag ggcaaccggc aagaggccgt 1680
cactgtggaa agtacctcgc tctagaaatt gtgcttgcac ggagcccctt tagtcaaaga 1740
ctcctggaaa ggagtgcct ctctccaat cagagcgtgg aggcgagttg ggcgacttca 1800
agggcctggc atggcagagg ccaggcagcc gtggccgagt ggacagcgtg gccgtctgct 1860
gtggctgaag gcagcttcca cagctcactc ctcttctccc tgccctgatc agattcccca 1920
ccttaccggg gccctacagg agcctgtgca gatggccatg cccaaccaat aacgagacgg 1980
ttcccctccc tttccctgcc aggctcatgt ctttacacct tcactcagcc acgccaacca 2040
gagactgggt tccaatctca ccccaccaca tacagagccc tcatctgtga aatgagaatg 2100
atcacgtgac ccacccccca gggcaggtat caggggtgaac tgatcttagc accggccaaa 2160
taaatggaac ctgctgagag agctgccaga taaaaaaaa aaaaaaaaaa 2210

```

<210> 2

<211> 550

<212> PRT

<213> Homo sapiens

<400> 2

```

Met Ala Phe Ser Lys Leu Leu Glu Gln Ala Gly Gly Val Gly Leu Phe
1           5           10           15
Gln Thr Leu Gln Val Leu Thr Phe Ile Leu Pro Cys Leu Met Ile Pro
           20           25           30
Ser Gln Met Leu Leu Glu Asn Phe Ser Ala Ala Ile Pro Gly His Arg
           35           40           45
Cys Trp Thr His Met Leu Asp Asn Gly Ser Ala Val Ser Thr Asn Met
           50           55           60
Thr Pro Lys Ala Leu Leu Thr Ile Ser Ile Pro Pro Gly Pro Asn Gln
           65           70           75           80
Gly Pro His Gln Cys Arg Arg Phe Arg Gln Pro Gln Trp Gln Leu Leu
           85           90           95
Asp Pro Asn Ala Thr Ala Thr Ser Trp Ser Glu Ala Asp Thr Glu Pro
           100          105          110
Cys Val Asp Gly Trp Val Tyr Asp Arg Ser Val Phe Thr Ser Thr Ile
           115          120          125
Val Ala Lys Trp Asp Leu Val Cys Ser Ser Gln Gly Leu Lys Pro Leu
           130          135          140
Ser Gln Ser Ile Phe Met Ser Gly Ile Leu Val Gly Ser Phe Ile Trp
           145          150          155          160
Gly Leu Leu Ser Tyr Arg Phe Gly Arg Lys Pro Met Leu Ser Trp Cys
           165          170          175

```

Cys Leu Gln Leu Ala Val Ala Gly Thr Ser Thr Ile Phe Ala Pro Thr
 180 185 190
 Phe Val Ile Tyr Cys Gly Leu Arg Phe Val Ala Ala Phe Gly Met Ala
 195 200 205
 Gly Ile Phe Leu Ser Ser Leu Thr Leu Met Val Glu Trp Thr Thr Thr
 210 215 220
 Ser Arg Arg Ala Val Thr Met Thr Val Val Gly Cys Ala Phe Ser Ala
 225 230 235 240
 Gly Gln Ala Ala Leu Gly Gly Leu Ala Phe Ala Leu Arg Asp Trp Arg
 245 250 255
 Thr Leu Gln Leu Ala Ala Ser Val Pro Phe Phe Ala Ile Ser Leu Ile
 260 265 270
 Ser Trp Trp Leu Pro Glu Ser Ala Arg Trp Leu Ile Ile Lys Gly Lys
 275 280 285
 Pro Asp Gln Ala Leu Gln Glu Leu Arg Lys Val Ala Arg Ile Asn Gly
 290 295 300
 His Lys Glu Ala Lys Asn Leu Thr Ile Glu Val Leu Met Ser Ser Val
 305 310 315 320
 Lys Glu Glu Val Ala Ser Ala Lys Glu Pro Arg Ser Val Leu Asp Leu
 325 330 335
 Phe Cys Val Pro Val Leu Arg Trp Arg Ser Cys Ala Met Leu Val Val
 340 345 350
 Asn Phe Ser Leu Leu Ile Ser Tyr Tyr Gly Leu Val Phe Asp Leu Gln
 355 360 365
 Ser Leu Gly Arg Asp Ile Phe Leu Leu Gln Ala Leu Phe Gly Ala Val
 370 375 380
 Asp Phe Leu Gly Arg Ala Thr Thr Ala Leu Leu Leu Ser Phe Leu Gly
 385 390 395 400
 Arg Arg Thr Ile Gln Ala Gly Ser Gln Ala Met Ala Gly Leu Ala Ile
 405 410 415
 Leu Ala Asn Met Leu Val Pro Gln Asp Leu Gln Thr Leu Arg Val Val
 420 425 430
 Phe Ala Val Leu Gly Lys Gly Cys Phe Gly Ile Ser Leu Thr Cys Leu
 435 440 445
 Thr Ile Tyr Lys Ala Glu Leu Phe Pro Thr Pro Val Arg Met Thr Ala
 450 455 460
 Asp Gly Ile Leu His Thr Val Gly Arg Leu Gly Ala Met Met Gly Pro
 465 470 475 480
 Leu Ile Leu Met Ser Arg Gln Ala Leu Pro Leu Leu Pro Pro Leu Leu
 485 490 495
 Tyr Gly Val Ile Ser Ile Ala Ser Ser Leu Val Val Leu Phe Phe Leu
 500 505 510

