



(86) Date de dépôt PCT/PCT Filing Date: 2001/10/17
(87) Date publication PCT/PCT Publication Date: 2002/04/25
(45) Date de délivrance/Issue Date: 2013/09/24
(85) Entrée phase nationale/National Entry: 2003/04/14
(86) N° demande PCT/PCT Application No.: EP 2001/012018
(87) N° publication PCT/PCT Publication No.: 2002/033119
(30) Priorité/Priority: 2000/10/17 (IT MI2000A002240)

(51) Cl.Int./Int.Cl. *C12N 15/12* (2006.01),
A61K 31/7088 (2006.01), *A61K 38/17* (2006.01),
A61K 39/395 (2006.01), *A61K 48/00* (2006.01),
A61P 3/02 (2006.01), *C07H 21/00* (2006.01),
C07K 14/47 (2006.01), *C07K 14/705* (2006.01),
C07K 16/28 (2006.01), *C07K 7/06* (2006.01),
C12N 5/10 (2006.01), *C12P 19/34* (2006.01),
C12Q 1/68 (2006.01), *G01N 33/53* (2006.01),
G01N 33/68 (2006.01)

(72) Inventeur/Inventor:
PIETRANGELO, ANTONELLO, IT

(73) Propriétaire/Owner:
PIETRANGELO, ANTONELLO, IT


(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : MUTATIONS AFFECTANT LE GENE DE LA FERROPORTINE 1 ASSOCIE A L'HEMATOCHROMATOSE
HEREDITAIRE

(54) Title: MUTATIONS IN THE FERROPORTIN 1 GENE ASSOCIATED WITH HEREDITARY HAEMOCHROMATOSIS

(57) Abrégé/Abstract:

The present invention relates to mutations in the gene coding for ferroportin 1 associated with hereditary haemochromatosis and methods for the diagnosis of hereditary haemochromatosis based on the identification of such mutations.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 April 2002 (25.04.2002)

PCT

(10) International Publication Number
WO 02/033119 A2

(51) International Patent Classification⁷: **C12Q 1/68**,
A61K 39/395, 38/17, C12N 15/11, A61K 31/70, C07K
16/28, G01N 33/53, C12N 15/85, 5/10, C07K 7/00, C12N
15/52, C07K 14/435

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU.

(21) International Application Number: PCT/EP01/12018

(22) International Filing Date: 17 October 2001 (17.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
MI2000A002240 17 October 2000 (17.10.2000) IT

(71) Applicant and

(72) Inventor: PIETRANGELO, Antonello [IT/IT]; Via San
Martino Mugnano, 3/1, I-41100 Modena (IT).

(74) Agent: GERVASI, Gemma; Notarbartolo & Gervasi,
Corso di Porta Vittoria, 9, I-20122 Milan (IT).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— *without international search report and to be republished
upon receipt of that report*

(48) Date of publication of this corrected version:

19 September 2002

(15) Information about Correction:

see PCT Gazette No. 38/2002 of 19 September 2002, Sec-
tion II

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

(54) Title: MUTATIONS IN THE FERROPORTIN 1 GENE ASSOCIATED WITH HEREDITARY HAEMOCHROMATOSIS

(57) Abstract: The present invention relates to mutations in the gene coding for ferroportin 1 associated with hereditary haemochro-
matosis and methods for the diagnosis of hereditary haemochromatosis based on the identification of such mutations.

WO 02/033119 A2

Mutations in the ferroportin 1 gene associated with hereditary haemochromatosis

FIELD OF THE INVENTION

The present invention concerns mutations in the gene coding for ferroportin 1 associated with hereditary haemochromatosis and the identification of such mutations as a diagnostic method for hereditary haemochromatosis.

STATE OF THE ART

Haemochromatosis is a hereditary pathology characterised by an excessive accumulation of iron in the organism, which over time leads to lesions of different organs and tissues, particularly liver, myocardium, pancreas, kidney, spleen, gonads and skin. Idiopathic haemochromatosis is the most widespread hereditary disease in the Western population (incidence 1: 300) and is characterised by a recessive transmission. Recently this type of haemochromatosis has been associated with mutations of the HFE gene, located on the short arm of chromosome 6. In a study carried out on patients suffering from this pathology in fact, it was observed that 83% of the analyzed subjects had a single point mutation at this gene (C282Y) (Feder et al, Nat Genet 1996, 13: 399-408).

However, more recent studies have shown that in the Mediterranean population only 64% of patients suffering from hereditary haemochromatosis are homozygotes for the C282Y mutation. This suggests that, in the southern European population in particular, other genes besides HFE may be responsible for idiopathic haemochromatosis (Piperno et al, Gastroenterology 1998, 114: 996-1002 and Borot et al, Immunogenetics 1997, 45: 320-324).

The identification of the genetic modifications responsible for hereditary haemochromatosis is of great diagnostic and therapeutic importance. Up to now, the diagnosis of haemochromatosis occurs too late and is based on the clinical symptomatology that develops as a result of often irreversible tissue injury. Besides, diagnosis of this pathology is made difficult by the fact that its symptoms are often similar to those of other pathologies characterised by altered iron homeostasis.

The development of genetic screening methods for the early diagnosis, at presymptomatic stage, of hereditary haemochromatosis would allow a timely

phlebotomy intervention before damage to organs and tissues occurs.

Furthermore, the identification of genetic alterations associated with hereditary haemochromatosis and the understanding of the role that they play in the development of the pathology are of extreme importance for the setting up of new and improved therapeutic strategies.

SUMMARY OF THE INVENTION

The inventor has previously identified and characterised a family suffering from a form of non HFE-dependent haemochromatosis with a autosomal dominant inheritance pattern, (Pietrangelo et al, New Eng J Med 1999, 341: 725-732).

The inventor has now surprisingly found that the locus of this pathology is on the long arm of chromosome 2 (2q32) and that subjects suffering from this type of haemochromatosis have a mutation at a codon located in the hexone 3 of the gene coding for ferroportin 1, that is situated in the same chromosomal region, which is not observed in subjects who are not suffering from the pathology. This mutation leads to the substitution of an amino acid in the ferroportin 1 molecule.

Therefore the present invention refers to a nucleic acid coding for a mutated ferroportin 1 characterised by a mutation of the codon coding for the amino acid corresponding to position 77 of SEQ. ID. No. 2, a mutated ferroportin 1 protein coded by said nucleic acid and methods for the *in vitro* diagnosis of hereditary haemochromatosis based on the identification of said nucleic acid or said protein.

DESCRIPTION OF THE FIGURES

Figure 1: Outline of the strategy for the identification of the mutation by means of enzymatic digestion with MbolI, described in Example 3.

In detail:

Figure 1a shows the genomic DNA sequence in the diagnostic method, in which the framed GCC sequence is the codon coding for the amino acid alanine in position 77 of the *wild type* ferroportin that is mutated to the GAC codon in individuals suffering from hereditary haemochromatosis. The double bar (//) denotes the separation between hexone 3 and a segment of the intron 3.

Figure 1b shows the primers used in the PCR reaction of the diagnostic method described in Example 3b and the target genomic DNA sequence in which N stands for the C or A nucleotide. The forward primer has a mismatched nucleotide, which

is underlined (A instead of G).

Figure 1c shows the amplified sequence from control individuals, in the absence of the mutation.

Figure 1d shows instead the amplified DNA sequence from individuals suffering from the pathology in which, in one of the alleles, the GCC codon is mutated to the GAC codon and leads to the appearance of a consensus site for the Mbo II enzyme.

Figure 2: Results of the diagnostic analysis on healthy family members or on family members suffering from haemochromatosis.

In detail:

Figure 2a shows the relationship between the analyzed individuals (pedigree). The subjects suffering from haemochromatosis are represented in black, while the healthy ones are in white. The circles represent the female subjects and the squares males.

Figure 2b shows the restriction profiles, following digestion with Mbo II, of the DNA amplified from each individual.

As shown in Figure 2c, in the case of healthy subjects, having only the *wild type* sequence, following digestion with Mbo II, the 131 base pair amplified DNA is not digested. Since all the subjects suffering from the pathology are heterozygotes for the mutation, the amplified DNA from these patients is digested in a band of 131 base pairs (normal allele) and two bands of 94 and 37 base pairs (the latter is not visible in Figure 2b).

DETAILED DESCRIPTION OF THE INVENTION

As will be shown in detail in the examples which follow, the authors of the present invention have identified that the mutation of a particular codon located in the hexone 3 of the ferroportin 1 gene is associated with a form of hereditary haemochromatosis that is not dependent on the HFE gene.

The identified mutation leads to the expression of a mutated ferroportin 1 in which the amino acid alanine, in a position corresponding to position 77 of SEQ ID No: 2 (sequence listing enclosed), is replaced.

The inventor has surprisingly found that the above mutation is a sufficient indication of the presence of hereditary haemochromatosis not linked to the HFE

gene, and that therefore its identification is useful for the early diagnosis of this pathology. Besides, the authors of the present invention have found that hereditary haemochromatosis is correlated with the functional impairment of ferroportin 1.

Therefore, according to a first aspect the present invention relates to a nucleic acid coding for a mutated ferroportin 1 characterised in that it comprises a mutation of the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2.

The term "nucleic acid coding for a ferroportin 1" means a genomic DNA, cDNA, DNA, for example obtained through PCR, or mRNA coding for the amino acid sequence of SEQ ID NO: 2 or for an amino acid sequence having at least 90% and preferably at least 95% homology with said amino acid sequence. When the nucleic acid is DNA this may be a single or double helix.

In addition, the invention comprises a nucleic acid with a sequence which is complementary to that of the aforesaid nucleic acid. For instance, this sequence can be an antisense sequence used to stop, the expression of the gene or of the mRNA in the cells.

The mutation, according to the invention, leads to the substitution, in the molecule of *wild-type* ferroportin 1 (GenBank accession number: AF231121) of the amino acid corresponding to position 77.

The term "*wild-type* ferroportin 1" refers to a ferroportin 1 which carries out its normal and physiological role, in particular that is without mutations which alter its functionality. Besides, the numerical position of the amino acid has the sole purpose of identifying it and may vary due to the presence of variations in the amino acid sequence of the protein, for instance with changes in the species taken into consideration or because of the presence of mutations or deletions in the regions upstream of said amino acid.

The inventor has found that the substitution of the alanine in position 77 with a molecule of aspartic acid results in a structural modification in ferroportin 1 which may be shown, for example, by prediction programs of protein secondary structure, namely "PHDsec" (Rost et al, J Mol Biol 232: 584-599, 1993) and "JRED" (Cuff et al, Proteins: Structure, Function Genetics and 34: 508-519). In particular, said substitution determines the passage of the protein region that goes

from amino acid 58 to amino acid 81 (LLLTAVYGLVVAGS VLVLGXIIGD, SEQ ID NO: 8) from the alpha-helix to beta-layer configuration.

The importance of the amino acid in position 77 in determining the secondary structure of ferroportin 1 is also confirmed by its high conservation among
5 different animal species.

The substitution of alanine, a small and uncharged amino acid, with a charged and larger molecule such as aspartic acid, results in steric and electrostatic interactions that destabilize the alpha helix hydrogen bonds. Similarly, substitutions of alanine with charged amino acids such as, for instance, arginine,
10 lysine or glutamic acid or with amino acids of greater sterical hinderance, for instance histidine, can lead to a similar distortion of the molecule.

Furthermore, the ferroportin 1 region corresponding to SEQ ID NO 8 contains a post-transductional modification site of the protein, the myristilation site GAIIGD, that is altered in the mutated protein according to the invention. As is known,
15 myristilation is important in helping the interactions of polypeptides with membrane phospholipids.

Therefore, according to a second aspect, the invention relates to a mutated ferroportin 1 protein characterised by the substitution of the amino acid in a position corresponding to position 77 of SEQ ID NO: 2 and coded by the nucleic
20 acid as above.

Said substitution consists in the substitution of the amino acid alanine in a position corresponding to position 77, with an amino acid having different steric and/or electrostatic properties from those of alanine. Preferably, said amino acid is selected from the group including arginine, lysine, glutamic acid or aspartic acid,
25 among which aspartic acid is preferred.

Therefore, according to a particularly preferred application, the mutation present in the nucleic acid of the invention consists in the substitution of the codon coding for the amino acid in a position corresponding to position 77 of SEQ ID NO: 2, preferably GCC, with a codon chosen from the group including GAC and GAU,
30 among which GAC is preferred.

In a further aspect of the invention relates to peptides with an amino acid sequence of at least 6 amino acids comprising the amino acid corresponding to

position 77 of SEQ ID NO: 2 and the amino acids immediately downstream and/or upstream of this. The length and sequence of such peptides are selected on the basis of criteria known to a person skilled in the art according to the desired application, for instance, to stimulate the production in host animals of antibodies
5 specific for the mutation or to obtain peptides and/or antibodies that specifically interact with the mutated epitope.

A preferred embodiment of such peptides is the peptide corresponding to SEQ ID NO: 8, in which Xaa is preferably aspartic acid.

The present invention also relates to nucleotide fragments of the aforesaid nucleic
10 acid comprising the mutated codon, and oligonucleotides with a sequence of at least 9 nucleotides and preferably at least 15 nucleotides of the aforesaid nucleic acid including said codon.

Said fragments and oligonucleotides can be of RNA or DNA and, in the latter case, single or double helix. Preferably, the oligonucleotides of the invention are single
15 helix.

Furthermore the invention includes nucleotide fragments and oligonucleotides with sequences complementary to those of the aforesaid fragments or oligonucleotides. By "nucleotide fragment", according to the present invention, is meant a nucleic acid with a sequence corresponding to a partial sequence of the nucleic acid of the
20 invention having a length exceeding 100 base pairs.

By "oligonucleotide" according to the present invention, is meant a fragment of the nucleic acid of the invention of a maximum length of 100 base pairs.

The nucleotide fragments and oligonucleotides of the invention may be obtained, for instance, by digestion of the nucleic acid of the invention, through amplification
25 by PCR or synthesized using techniques known in the art.

The oligonucleotides and the DNA fragments of the invention are used for different purposes such as, for instance, the production of chimeric proteins or antibodies, the determination of the mutation of the invention for diagnostic purposes or the inactivation of the mutated gene for therapeutic purposes. A person skilled in the
30 art is able to choose fragments and oligonucleotides of having sequence and length suitable for the desired uses. For instance, should said fragments or oligonucleotides be used for the determination of the mutation of invention with

hybridization techniques, they must be of a length and sequence so as to be capable of hybridizing in a specific way, under stringent conditions, at a nucleic acid sequence comprising the mutated codon.

According to a preferred embodiment, the fragments and oligonucleotides of the invention are labelled, for instance, with radioisotopes, enzymes, biotin-avidin or other molecules which allow them to be visualized through specific assays.

Moreover, the invention relates to the peptides coded by such fragments and oligonucleotides.

The nucleic acid, of the invention, a fragment thereof comprising the mutation or a nucleic acid comprising such fragment can be advantageously used for the production of a recombinant mutated ferroportin 1, a fragment of it or a chimeric protein including this fragment, in order to, for instance, study the functional characteristics of the mutated protein, for example through competition studies, or produce antibodies. To this purpose, said nucleic acid or fragment is inserted into an expression vector which, in turn, is introduced into a procaryotic or eucaryotic cell using techniques well known in the art such as transfection, transformation, infection or intranuclear injection.

Vectors suitable for this purpose include plasmids, vectors of viral origin and yeast or mammalian artificial chromosomes.

Accordingly, in a further aspect, the invention relates to a recombinant vector comprising a nucleic acid or a DNA fragment according to the invention as well as to eucaryotic or procaryotic cells comprising said vector.

The nucleic acid according to the invention can be used for the preparation of eucaryotic cells, tissues or non-human animals comprising a transgene coding for the mutated ferroportin 1 of the invention. The transgene can be permanently inserted into the genome of the cell, tissue or animal or be present in extrachromosomal form.

Said cells, tissues or non-human animals are useful as models to study the function of the gene and of the protein comprising the mutation of the invention, as well as their role in the onset of hereditary haemochromatosis. This study is of particular importance for the development of new therapeutic approaches for the treatment of hereditary haemochromatosis.

In a further aspect, the invention refers to a method for the *in vitro* diagnosis of hereditary haemochromatosis in a mammal, preferably Homo Sapiens, comprising the following steps:

a) isolation of genomic DNA or RNA from a biological sample obtained from said mammal;

b) testing for the presence, in said genomic DNA or RNA, of the mutation according to the invention,

Wherein the presence of said mutation is an indication that said mammal is suffering from hereditary haemochromatosis.

Preferably said biological sample is a sample of plasma, saliva, urine, faeces, amniotic fluid or tissue.

Prior to testing, the RNA is preferably transformed into complementary DNA (cDNA) through a reverse transcription reaction.

The genomic DNA or the cDNA are analyzed directly or following *in vitro* amplification polymerase chain reaction (PCR) (Saiki et al, Science 239: 487-491, 1988) or other techniques, for instance, ligase chain reaction (LCR) (Wu et al, genomics 4: 560-569, 1989) strand displacement amplification (SDA) (Walker et al, PNAS USA 89: 392-396) or self-sustained sequence replication (3SR) (Fahy et al, PCR methods Appl. 1: 25-33, 1992).

Preferably, the genomic DNA or the cDNA is amplified through PCR using a pair of oligonucleotides (primers) suitable for the amplification of a segment of said DNA comprising the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2.

For instance, pairs of primers that can be used to amplify the cDNA are those with the nucleotide sequence of SEQ ID No: 3 and SEQ ID No: 4 whereas primers suitable for the amplification of the genomic DNA are those with the nucleotide sequence of SEQ ID No: 5 and SEQ ID No: 6.

Numerous techniques, well known in the art, can be used to determine the presence of the mutation according to the invention in the genomic DNA or the cDNA. Suitable techniques are, for instance, techniques based on the use of restriction enzymes (Kan et al, Lancet: 910-912, 1978), hybridization techniques with allele-specific oligonucleotide probes (Wallace et al, Nucl Acids Res 6: 3543-

3557, 1978) among which, for instance, hybridization with oligonucleotides immobilized on filters (Saiki et al, PNAS USA 86: 6230-6234, 1989) or micro-chips (Chee et al, Science 274: 610-614, 1996) and oligonucleotide arrays (Maskos et al, Nucl Acids Res 21: 2269-2270, 1993), allele-specific PCR (Newton et al Nucl
5 Acid Res 17: 2503-2516, 1989), mismatch repair detection (MRD) (Faham and Cox Genome Res: 474-482, 1995), Single-strand conformational polymorphism analysis (Ravnik-Glavac et al, Hum. Mol. Gen. 3: 801, 1994), gel electrophoresis in denaturant gradient (Guldborg et al., Nucl. Acids Res. 22: 880, 1994), Hot Cleavage (Cotton et al Proc.Natl. Acad Ski USA 85: 4397, 1988), DNase (Youil et
10 al, PNAS USA 92: 87-91, 1995) and RNase protection assay (Winter et al. Proc. Natl. Acad. Ski. USA, 82: 7575, 1985; Meyers et al, Science 230: 1242, 1985), allele specific primer extension (Syvanen et al, genomics 8: 684-692, 1990 and Syvanen et al, Hum Mutat 13: 1-10, 1999), genetic bit analysis (GBA) (Nikiforov et al Nucl Acid Res 22: 4167-4175, 1994), primer-ligation assay (OLA) (Landerger et
15 al, Science 241: 1077, 1988), allele specific ligation chain reaction (LCR) (Barrany PNAS USA 88: 189-193, 1991), gap-LCR (Abravaya et al Nucl Acids Res 23: 675-682, 1995) and sequencing techniques. Particularly preferred techniques for the determination of the mutation of the invention are techniques based on the use of restriction enzymes, allele specific PCR, hybridization and sequencing techniques.

20 Therefore, according to a first preferred application, the testing for the presence of the mutation according to the invention in the analyzed DNA occurs using techniques based on the use of restriction enzymes and comprises the following steps:

a) amplification of the genomic DNA or cDNA with a pair of oligonucleotides
25 suitable for the selective amplification of a segment of said DNA including the codon coding for the amino acid corresponding to position 77 of SEQ ID NO: 2 and the concurrent introduction into the amplified DNA of such a mutation which, in the presence of the mutation of the invention, creates the consensus sequence for a restriction site otherwise not present;

30 b) incubation of the amplified DNA with an enzyme which is able to recognize said restriction site; and

c) analysis of the size of the digestion products;

wherein the occurring of digestion is a sign of the presence of the mutation of the invention in the genomic or complementary DNA.

The analysis of the size of the digestion products may be carried out, for instance, through gel electrophoresis, using a marker of molecular weights, followed by
5 visualization of the DNA bands using, for example, ethidium bromide.

In order to test, for example, for the presence of the substitution of the GCC codon, coding for alanine in position 77, with the GAC codon, coding for an aspartic acid molecule, oligonucleotides can be used with the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 7. As will be shown in the examples that follow,

10 these oligonucleotides give rise, in the presence of the aforesaid substitution, to a fragment of amplified DNA with the sequence shown in Figure 1d which contains the consensus site for the Mbo II enzyme, GAAGACATCATCGGT, not present in the fragment of DNA amplified from the *wild-type* DNA (Fig.1c). The subsequent incubation of the product of the amplification reaction with Mbo II results in the
15 fragment being digested only if the mutation was present in the original sequence.

According to a further preferred application, the determination of the mutation according to the invention is performed through hybridization techniques in which fragments of the nucleic acid of the invention or oligonucleotides specific for the mutation according to the invention are used.

20 Said fragments or oligonucleotides are capable of hybridizing, in a specific way, at a sequence of the nucleic acid of the invention comprising the mutated codon also when said sequence is present together with many other sequences.

A person skilled in the art is able to select each time the hybridization conditions and the length and sequence of the fragments or the oligonucleotides most
25 suitable for the particular hybridization technique used and the kind of DNA that is being analyzed (genomic or complementary DNA, amplified or cloned in appropriate vectors).

According to a further preferred application, the diagnostic method envisages the use of allele-specific PCR, in which the genomic or complementary DNA
30 undergoes a PCR reaction in which oligonucleotides are used which are able to selectively amplify a segment of said DNA including the mutated codon and not the corresponding segment including the *wild-type* codon.

In addition the present invention also relates to nucleic acid fragments and oligonucleotides according to the invention to be used in the aforesaid methods. In particular, it refers to oligonucleotides with the nucleotide sequence corresponding to SEQ ID NO: 3, 4, 5, 6 and 7.

5 Within the scope of the present invention are also included diagnostic kits for the identification, in an individual, of the mutation according to the invention. According to a particularly preferred application, said diagnostic kits comprise oligonucleotides with a nucleotide sequence corresponding to SEQ ID NO: 6 and 7 and the Mbo II enzyme.

10 The present invention also relates to a method for the *in vitro* diagnosis of hereditary haemochromatosis in mammals including testing for the presence, in a biological sample from said mammal, of a mutated ferroportin 1 protein according to the invention, wherein the identification of said protein is an indication that the individual is suffering from hereditary haemochromatosis.

15 Preferably said testing is performed through immunological assays in which monoclonal or polyclonal antibodies are used which are able to discriminate between a molecule of mutated ferroportin, according to the invention, and a molecule of *wild-type* ferroportin.

Therefore the present invention also refers to monoclonal or polyclonal antibodies
20 which are able to specifically recognize a molecule of mutated ferroportin 1 according to the invention, or on epitope thereof comprising the mutation. Such antibodies are obtained using methods well known in the art such as those described by Harlow and Lane in Antibodies, A Laboratory Manual, Cold Spring Harbour Laboratory 1988.

25 The antibodies of the invention are particularly useful, as well as diagnostic reagents, in studying the characteristics of the protein or for therapeutic purposes. For instance, said antibodies can be used to determine the exact tissular or cellular localization of the mutated protein, to study its biochemical characteristics or to purify it by immunoaffinity.

30 Moreover, since the presence in an individual of a gene bearing the mutation of the invention and of the ferroportin 1 coded by it is correlated with the onset of hereditary haemochromatosis it is very important to have means to stop

expression of the gene or to inactivate the protein.

Therefore, the invention also relates to oligonucleotides, for instance antisense oligonucleotides, suitable for inhibiting the expression of the gene coding for mutated ferroportin 1 of the invention, and antibodies and polypeptides able to specifically alter the functionality of the mutated ferroportin 1 of the invention.

In addition, the invention relates to pharmaceutical compositions comprising said oligonucleotides, antibodies or peptides mixed with pharmaceutically acceptable excipients.

EXAMPLES

EXAMPLE 1

Identification of the chromosome and the chromosomal locus associated with non HFE-dependent hereditary haemochromatosis

A DNA sample was extracted from the peripheral blood of the proband and family members suffering from the pathology using Quiagen Blood Extraction Kit (Quiagen).

The DNA extracted was then used for a genome wide-search that was performed by using the ABI PRISM Linkage mapping set (Perkin Elmer, United States). Fluorescent oligonucleotides were used for the PCR reactions, under the conditions suggested by the manufacturer. An aliquot of each PCR reaction was then sequenced in an ABI PRISM 377 DNA sequencer and the results obtained were analyzed using GENESCAN software. The assignment of the allele was carried out using GenotyperTM software. Statistical analysis was performed based on a dominant autosomal disease with complete penetrance. The gene-disease frequency was fixed at 0.012 and all the marker alleles were considered to be equally frequent. Table 1 contains the lod score, i.e. the maximum probability of association of a specific marker with the disease.

Table 1:

Marker Z at $\theta_{\max}$

	.00	.01	.05	.1	.2	.3	.4	Zmax ($\theta_{\max}$)		
5										
	D2S2257	.20	.20	.19	.16	.11	.06	.02	.20	(.000)
	D2S364	$-\infty$	3.46	3.82	3.67	3.00	2.08	.99	3.82	(.040)
10										
	D2S350	$-\infty$	2.82	3.16	2.99	2.31	1.44	.53	3.16	(.040)
	D2S152	5.88	5.78	5.38	4.85	3.74	2.51	1.16	5.88	(.000)
15	D2S118	5.99	5.89	5.49	4.96	3.84	2.60	1.23	5.99	(.000)
	D2S280	$-\infty$	2.35	2.77	2.70	2.22	1.53	.70	2.78	(.060)
	D2S315	3.46	3.41	3.17	2.87	2.22	1.48	.66	3.46	(.000)
20										
	D2S117	$-\infty$	2.69	3.09	3.00	2.47	1.71	.76	3.09	(.050)

The higher the score, expressed logarithmically, the lower the probability that the association of the disease with the specific markers used is casual. For example, a lod score of 1.0 denotes 1 chance in 10 that the result is due to chance; a lod score of 2, one chance in 100 and so on. The fact that the markers D2S118 (5.99) and D2S152 (5.88) gave very high scores shows that the gene associated with the disease is situated in the region delimited by these markers. Ferroportin 1 is situated in this chromosomal region. No other chromosomal region gave similar results.

EXAMPLE 2

Identification of the mutation

Blood samples were taken from the proband, from 15 family members suffering from the pathology and from 25 family members not suffering from the pathology. Total RNA was isolated from macrophages obtained from each sample by extraction in guanidine-isothiocyanate and the complementary DNA was then prepared according to a standard protocol (400 ng of total RNA, 1 µg of oligodT, 1mM of dNT, 20 U of reverse transcribed AMV in 20 µl of reaction buffer; Promega) The complete ferroportin 1 sequence was then amplified from the cDNA through a PCR reaction using the following oligonucleotide pairs:

Forward primer 5'-GCTCAGGGCGTCCGCTAGGCT-3'

10 (SEQ.ID.NO 3)

reverse primer: 5'-GGCTTACACCCTCATGTTCT-3'

(SEQ ID NO: 4)

In detail, 10 ml of the product of the reverse transcription reaction were amplified in 50 µl (final volume) of 1X reaction buffer containing 200 µM dNTP, 1.5 mM MgCl₂, 0.25 µg of each of the above described oligonucleotide, 2.6 units of enzyme. A 30 cycle program was used for the amplification reaction, each of which was characterised by the following heat profile:

94°C for 1 minute,

58°C for 40 seconds,

20 75°C for 5 minutes.

From the amplified ferroportin 1 cDNA, four partially overlying DNA fragments were then obtained through a new PCR, using the following oligonucleotide pairs:

1° fragment:

Forward primer: 5'-GCTCAGGGCGTCCGCTAGGCT-3'

25 Reverse primer: 5'-CAGACACCGCAAAGTGCCACA-3'

2° fragment:

Forward primer: 5'-ACCTCGCTGGTGGTACAG-3'

Reverse primer: 5'-CCGCAAGCAAAGAGCTTG-3'

3° fragment:

30 Forward primer: 5'-GGTGCTATCTCCAGTTCCTT-3'

Reverse primer: 5'-AGACGTACTCCACGCACA-3'

4° fragment:

Forward primer: 5'-TCAGTCTCCTTTGTGGCA-3'

Reverse primer: 5'-GGCTTACACCCTCATGTTCT-3'

The four fragments obtained from the amplification were then electrophoretically
5 separated on agarose gel, purified using the Jet Sorb kit (Genenco) and
sequenced directly using the Rhodamine Sequence kit (Perkin Elmer, United
States). Sequencing revealed the presence, in subjects suffering from the
pathology, of the substitution of a C with an A in position 230 of SEQ.ID No.1
(nucleotide in position 534 of the sequence having GenBank accession number:
10 AF231121), that was not found in any of the control subjects. This substitution is
located in hexone 3 of the ferroportin 1 gene and results in the substitution of
alanine with aspartic acid at position 77 of ferroportin 1.

EXAMPLE 3**Diagnostic method**

15 a) genomic DNA of the proband, of 15 family members suffering from the
pathology and 125 control individuals, including 100 healthy volunteers and 25
family members not suffering from the pathology, was extracted from leukocytes
obtained from blood samples of the subjects to be analyzed using a blood DNA
extraction kit (Quiagen).

20 The DNA obtained was then amplified by PCR using a pair of oligonucleotides
complementary to the intronic regions flanking hexone 3 and having the following
sequence:

Forward primer: 5'-CCTTTTGATAAGGAAGCAACTTCC-3'

(SEQ ID NO: 5)

25 Reverse primer: 5'-CAGAGGTAGCTCAGGCATTGGTCC-3'

(SEQ ID NO: 6)

In detail, 200 ng of genomic DNA were amplified in 50 µl of 1X reaction buffer
containing 200 µM dNTPs, 1.5 mM MgCl₂, 20 pmoles of each oligonucleotide and
2.6 U of enzyme.

30 A 30 cycle program was used for the amplification reaction, each of which was
characterised by the following heat profile:

94°C for 1 minute,

60°C for 1 minute,

72°C for 45 seconds.

The DNA obtained was then purified using the PCR Wizard kit (Promega) and sequenced in an automatic ABI Prism 377 sequencer (Perkin Elmer, United States), with the same pair of oligonucleotides used for the PCR reaction.

The substitution of a C with an A in hexone 3 was found in the subjects suffering from the pathology, but not in the control subjects.

b) blood samples were obtained from the proband, from 15 family members suffering from the pathology and from the 125 control subjects described in example 2 and the genomic DNA was extracted using the blood DNA extraction kit (Biorad).

Since the mutation observed does not give rise to the appearance or disappearance of any restriction site, the portion of hexone 3 containing the mutation was amplified by PCR using a pair of oligonucleotides with the following sequences:

Forward primer: 5'-GTGGCAGGGTCTGTTCTGGTCCTGGAAG-3'
(SEQ.ID NO.7)

Reverse primer: 5'-CAGAGGTAGCTCAGGCATTGGTCC-3'
(SEQ.ID NO.6)

As shown in Figure 1b, the forward primer has a mismatched nucleotide (A instead of G, underlined). Amplification with the aforesaid oligonucleotides gives rise, in the presence of the mutation, to an amplified DNA fragment, having the sequence shown in Figure 1d, that contains the consensus site for the Mbo II enzyme, GAAGACATCATCGGT. On the contrary, in the absence of the mutation, a DNA fragment with the sequence shown in Figure 1d is obtained and which does not contain such a restriction site (Fig.1c).

Therefore, the subsequent incubation of the product of the amplification reaction with the Mbo II enzyme results in the digestion of the fragment only if the original sequence contained the mutation.

In detail, 10 µl of the extracted genomic DNA were amplified in a final 50 µl of 1x reaction buffer containing 200 µM dNTPs, 1.5 mM MgCl₂, 20 pmoles of each oligonucleotide and 2.6 units of enzyme. A 30 cycle program was used for the

contains the lod score, i.e. the maximum probability of association of a specific marker with the disease.

Table 1:

Marker	Z at $\theta_{\max}$							
	.00	.01	.05	.1	.2	.3	.4	Zmax ($\theta_{\max}$)
D2S2257	.20	.20	.19	.16	.11	.06	.02	.20 (.000)
D2S364	$-\infty$	3.46	3.82	3.67	3.00	2.08	.99	3.82 (.040)
D2S350	$-\infty$	2.82	3.16	2.99	2.31	1.44	.53	3.16 (.040)
D2S152	5.88	5.78	5.38	4.85	3.74	2.51	1.16	5.88 (.000)
D2S118	5.99	5.89	5.49	4.96	3.84	2.60	1.23	5.99 (.000)
D2S280	$-\infty$	2.35	2.77	2.70	2.22	1.53	.70	2.78 (.060)
D2S315	3.46	3.41	3.17	2.87	2.22	1.48	.66	3.46 (.000)
D2S117	$-\infty$	2.69	3.09	3.00	2.47	1.71	.76	3.09 (.050)

The higher the score, expressed logarithmically, the lower the probability that the association of the disease with the specific markers used is casual. For example, a lod score of 1.0 denotes 1 chance in 10 that the result is due to chance; a lod score of

2, one chance in 100 and so on. The fact that the markers D2S118 (5.99) and D2S152 (5.88) gave very high scores shows that the gene associated with the disease is situated in the region delimited by these markers. Ferroportin 1 is situated in this chromosomal region. No other chromosomal region gave similar results.

5 EXAMPLE 2

Identification of the mutation

Blood samples were taken from the proband, from 15 family members suffering from the pathology and from 25 family members not suffering from the pathology. Total RNA was isolated from macrophages obtained from each sample by extraction in
10 guanidine-isothiocyanate and the complementary DNA was then prepared according to a standard protocol (400 ng of total RNA, 1 µg of oligodT, 1mM of dNT, 20 U of reverse trascribed AMV in 20 µl of reaction buffer; Promega)

The complete ferroportin 1 sequence was then amplified from the cDNA through a PCR reaction using the following oligonucleotide pairs:

15 Forward primer 5'-GCTCAGGGCGTCCGCTAGGCT-3'

(SEQ.ID.NO 3)

reverse primer: 5'-GGCTTACACCCTCATGTTCT-3'

(SEQ ID NO: 4)

In detail, 10 ml of the product of the reverse transcription reaction were amplified in
20 50 µl (final volume) of 1X reaction buffer containing 200 µM dNTP, 1.5 mM MgCl₂, 0.25 µg of each of the above described oligonucleotide, 2.6 units of enzyme. A 30

cycle program was used for the amplification reaction, each of which was characterised by the following heat profile:

94°C for 1 minute,

58°C for 40 seconds,

5 75°C for 5 minutes.

From the amplified ferroportin 1 cDNA, four partially overlying DNA fragments were then obtained through a new PCR, using the following oligonucleotide pairs:

1° fragment:

Forward primer: 5'-GCTCAGGGCGTCCGCTAGGCT-3'

10 Reverse primer: 5'-CAGACACCGCAAAGTGCCACA-3'

2° fragment:

Forward primer: 5'-ACCTCGCTGGTGGTACAG-3'

Reverse primer: 5'-CCGCAAGCAAAGAGCTTG-3'

3° fragment:

15 Forward primer: 5'-GGTGCTATCTCCAGTTCCTT-3'

Reverse primer: 5'-AGACGTACTCCACGCACA-3'

4° fragment:

Forward primer: 5'-TCAGTCTCCTTTGTGGCA-3'

Reverse primer: 5'-GGCTTACACCCTCATGTTCT-3'

20 The four fragments obtained from the amplification were then electrophoretically separated on agarose gel, purified using the Jet Sorb kit (Genenco) and sequenced directly using the Rhodamine Sequence kit (Perkin Elmer, United States). Sequencing revealed the presence, in subjects suffering from the pathology, of the

substitution of a C with an A in position 230 of SEQ.ID No.1 (nucleotide in position 534 of the sequence having GenBank accession number: AF231121), that was not found in any of the control subjects. This substitution is located in hexone 3 of the ferroportin 1 gene and results in the substitution of alanine with aspartic acid at position 77 of ferroportin 1.

EXAMPLE 3

Diagnostic method

a) genomic DNA of the proband, of 15 family members suffering from the pathology and 125 control individuals, including 100 healthy volunteers and 25 family members not suffering from the pathology, was extracted from leukocytes obtained from blood samples of the subjects to be analyzed using a blood DNA extraction kit (Quiagen).

The DNA obtained was then amplified by PCR using a pair of oligonucleotides complementary to the intronic regions flanking hexone 3 and having the following sequence:

Forward primer: 5'-CCTTTTGATAAGGAAGCAACTTCC-3'

(SEQ ID NO: 5)

Reverse primer: 5'-CAGAGGTAGCTCAGGCATTGGTCC-3'

(SEQ ID NO: 6)

In detail, 200 ng of genomic DNA were amplified in 50 µl of 1X reaction buffer containing 200 µM dNTPs, 1.5 mM MgCl₂, 20 pmoles of each oligonucleotide and 2.6 U of enzyme.

A 30 cycle program was used for the amplification reaction, each of which was characterised by the following heat profile:

94°C for 1 minute,

60°C for 1 minute,

5 72°C for 45 seconds.

The DNA obtained was then purified using the PCR Wizard kit (Promega) and sequenced in an automatic ABI Prism 377 sequencer (Perkin Elmer, United States), with the same pair of oligonucleotides used for the PCR reaction.

10 The substitution of a C with an A in hexone 3 was found in the subjects suffering from the pathology, but not in the control subjects.

b) blood samples were obtained from the proband, from 15 family members suffering from the pathology and from the 125 control subjects described in example 2 and the genomic DNA was extracted using the blood DNA extraction kit (Biorad).

15 Since the mutation observed does not give rise to the appearance or disappearance of any restriction site, the portion of hexone 3 containing the mutation was amplified by PCR using a pair of oligonucleotides with the following sequences:

Forward primer: 5'-GTGGCAGGGTCTGTTCTGGTCCTGGAAG-3'

(SEQ.ID NO.7)

Reverse primer: 5'-CAGAGGTAGCTCAGGCATTGGTCC-3'

20 (SEQ.ID NO.6)

As shown in Figure 1b, the forward primer has a mismatched nucleotide (A instead of G, underlined). Amplification with the aforesaid oligonucleotides gives rise, in the presence of the mutation, to an amplified DNA fragment, having the sequence shown

in Figure 1d, that contains the consensus site for the Mbo II enzyme, GAAGACATCATCGGT. On the contrary, in the absence of the mutation, a DNA fragment with the sequence shown in Figure 1d is obtained and which does not contain such a restriction site (Fig.1c).

- 5 Therefore, the subsequent incubation of the product of the amplification reaction with the Mbo II enzyme results in the digestion of the fragment only if the original sequence contained the mutation.

In detail, 10 µl of the extracted genomic DNA were amplified in a final 50 µl of 1x reaction buffer containing 200 µM dNTPs, 1.5 mM MgCl₂, 20 pmoles of each
10 oligonucleotide and 2.6 units of enzyme. A 30 cycle program was used for the amplification reaction, each cycle being characterised by the following heat profile:

94°C for 1 minute,

58°C for 1 minute,

72°C for 45 seconds.

- 15 10 µl of the product obtained from the amplification were then digested with the Mbo II enzyme (Geneco) in 1X reaction buffer for 3 hours at 37°C.

The fragments obtained from the digestion were then separated on a 12% polyacrylamide gel. The samples obtained from patients suffering from the pathology resulted in 3 bands of 131, 94 and 37 base pairs. On the contrary, all the control
20 samples gave had just one band of 131 base pairs. Figure 2 shows the results obtained from the proband, from 4 family members suffering from the pathology and from 3 healthy family members. As can be seen from the figure, digestion of the

fragment of 131 base pairs into two fragments of 94 and 37 base pairs was only observed in subjects suffering from the pathology.

Also comprised in the present invention is any therapeutic operation involving the substitution of the mutated gene with the *wild type* gene. Therefore, the mutated

5 ferroportin 1 molecule is a therapeutic target for all genic therapy operations aimed at the substitution of the mutated gene.

WE CLAIM:

1. A nucleic acid coding for a ferroportin 1 protein comprising SEQ ID NO:2, wherein the amino acid at position 77 of SEQ ID NO:2 is selected from the group consisting of arginine, lysine, glutamic acid and aspartic acid, for diagnostic use of hereditary haemochromatosis.
2. The nucleic acid as claimed in claim 1 characterised in that it is cDNA.
3. The nucleic acid as claimed in claim 1 characterised in that it is mRNA.
4. The nucleic acid as claimed in claim 1 characterised in that it is genomic DNA.
5. The nucleic acid as claimed in claim 1 characterised in that said amino acid at position 77 of SEQ ID NO:2 is aspartic acid.
6. The nucleic acid as claimed in claim 5, wherein said aspartic acid is encoded by a GAC or a GAU codon.
7. The nucleic acid as claimed in claim 6, wherein said aspartic acid is encoded by the GAC codon.
8. A nucleic acid with a sequence which is complementary to that of the nucleic acid as claimed in claim 1.
9. A ferroportin 1 protein comprising SEQ ID NO:2 wherein the amino acid at position 77 of SEQ ID NO:2 is selected from the group consisting of arginine, lysine, glutamic acid and aspartic acid.
10. A fragment of the nucleic acid as claimed in claim 1, wherein said fragment comprises a sequence of at least 15 nucleotides comprising the codon coding for the amino acid corresponding to position 77 of SEQ ID NO:2.

11. A fragment with a sequence which is complementary to that of the fragment as claimed in claim 10.
12. The fragment as claimed in claim 10 or 11 characterised in that it is labelled.
13. An oligonucleotide with a sequence of at least 9 nucleotides of the nucleic acid as claimed in claim 1 and comprising the codon coding for the amino acid corresponding to position 77 of SEQ ID NO:2.
14. The oligonucleotide as claimed in claim 13 characterised in that said sequence is of at least 15 nucleotides.
15. The oligonucleotide as claimed in claim 13 characterised in that it is labelled.
16. An oligonucleotide with a sequence complementary to that of the oligonucleotide as claimed in claim 13.
17. A peptide coded by a fragment as claimed in claim 10 or by an oligonucleotide as claimed in claim 13.
18. A recombinant vector comprising the nucleotide sequence of the nucleic acid as claimed in claim 1 or of the fragment as claimed in claim 10.
19. A cell transfected or transformed with the recombinant vector as claimed in claim 18.
20. A method for the *in vitro* diagnosis of hereditary haemochromatosis in a mammal comprising:

testing for the presence of a codon coding for aspartic acid in position 77 of SEQ ID NO:2 in a genomic DNA or RNA isolated from a biological sample obtained from said mammal,

wherein the presence of said codon coding for aspartic acid in position 77 of SEQ ID NO:2 is an indication that said mammal is suffering from hereditary haemochromatosis.

21. The method as claimed in claim 20 characterised in that said mammal is a Homo sapiens.

22. The method as claimed in claim 20 characterised in that said biological sample is a sample of blood, plasma, saliva, urine, faeces, amniotic fluid or tissue.

23. The method as claimed in claim 20 characterised in that the isolated genomic DNA is amplified before said testing.

24. The method as claimed in claim 20 further comprising amplifying by PCR a segment of the genomic DNA comprising the codon coding for aspartic acid in position 77 of SEQ ID NO:2 using a pair of oligonucleotides.

25. The method as claimed in claim 24 characterised in that said oligonucleotides have the nucleotide sequence of SEQ. ID NO: 5 and SEQ. ID NO: 6.

26. The method as claimed in claim 20 characterised in that the RNA is transformed into cDNA before said testing.

27. The method as claimed in claim 26 characterised in that the cDNA is amplified before said testing.

28. The method as claimed in claim 20 further comprising transforming the RNA into cDNA and amplifying by PCR a segment of the cDNA comprising the codon coding for aspartic acid in position 77 of SEQ ID NO:2 using a pair of oligonucleotides.

29. The method as claimed in claim 28 characterised in that said oligonucleotides have the nucleotide sequence of SEQ. ID NO: 3 and SEQ. ID NO: 4.

30. The method as claimed in claim 20 characterised in that said testing is carried out using a technique selected from the group including techniques based on the use of restriction enzymes, hybridization techniques with allele-specific oligonucleotide probes, allele-specific PCR, mismatch repair detection, single-strand conformational polymorphism analysis, gel electrophoresis in denaturant gradient, Hot Cleavage, DNase and RNase protection assay, allele specific primer extension, genetic bit analysis oligonucleotide-ligation assay, allele specific ligation chain reaction and sequencing techniques.

31. The method as claimed in claim 30 characterised in that said testing is carried out using techniques based on the use of restriction enzymes, allele specific PCR, hybridization or sequencing techniques.

32. The method as claimed in claim 31 characterised in that said testing comprises the following steps:

- a) amplifying a segment of said genomic DNA or of a cDNA transformed from said RNA using oligonucleotides to produce an amplified DNA, wherein the amplified DNA includes a codon coding for aspartic acid in a position corresponding to position 77 of SEQ ID NO: 2 and a consensus sequence for a restriction site absent in a nucleotide sequence coding a protein having a sequence of SEQ ID NO:2 where the codon coding for an amino acid in position 77 of SEQ ID NO:2 encodes alanine;
- b) incubating the amplified DNA with an enzyme which is able to recognize said restriction site to produce digestion products; and
- c) analyzing the size of the digestion products;

wherein the presence of digestion products indicates the presence of the codon coding for aspartic acid in position 77 of SEQ ID NO:2 in the genomic DNA or cDNA.

33. The method as claimed in claim 32 characterised in that said oligonucleotides have the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 7 and that said enzyme is Mbo II.

34. An oligonucleotide with the nucleotide sequence of SEQ ID NO: 3, 4, 5, 6 or 7, for use as a primer to amplify a genomic DNA, cDNA or RNA sequence coding for a ferroportin 1 protein comprising SEQ ID NO:2 or segment thereof containing a codon encoding for one of an arginine, lysine, glutamic acid and aspartic acid corresponding to position 77 of SEQ ID NO:2.

35. A diagnostic kit for the determination of a codon encoding aspartic acid in position 77 of SEQ ID NO:2 in an individual comprising oligonucleotides with the nucleotide sequence of SEQ ID NO: 6 and SEQ ID NO: 7 and a Mbo II enzyme.

36. A method for the *in vitro* diagnosis of hereditary haemochromatosis in a mammal comprising testing for the presence of a ferroportin 1 protein in a biological sample obtained from said mammal, wherein said ferroportin 1 protein has an amino acid sequence of SEQ ID NO:2 in which the amino acid in position 77 of SEQ ID NO:2 is aspartic acid, in which the presence of said protein in said biological sample is an indication that said mammal is suffering from hereditary haemochromatosis.

FIGURE 1

GAC
A77D

a) Exon 3 GTGGCAGGGTCTGTTCTGGTCTGCTGGGA **GC**ATCATCGGTGACTGGGTGGACAAGAATGCTAGACTAAAG // ATTCAATGGGACCAATGCCCTGAGCTACCTCTG
CACCGTCCCAGACAAGACCAGGACCCTCTGTGTAGTAGCCACTGACCCACCTGTGTTCTTACGATCTGATTTC // TAAGTACCCCTGGTTACGGACTCGGATGGAGAC

Forward primer

5' GTGGCAGGGTCTGTTCTGGTCTGCTGGGAAG 3'

b) PCR

GTGGCAGGGTCTGTTCTGGTCTGCTGGAGNCATCATCGGTGACTGGGTGGACAAGAATGCTAGACTAAAG // ATTCAATGGGACCAATGCCCTGAGCTACCTCTG
CACCGTCCCAGACAAGACCAGGACCCTCTGTGTAGTAGCCACTGACCCACCTGTGTTCTTACGATCTGATTTC // TAAGTACCCCTGGTTACGGACTCGGATGGAGAC

1/2

GGTTACGGACTCGATGGAGAC 3' Reverse primer 5'

c) A77D +/-

GTGGCAGGGTCTGTTCTGGTCTGGAAGCCATCATCGGTGACTGGGTGGACAAGAATGCTAGACTAAAG // ATTCAATGGGACCAATGCCCTGAGCTACCTCTG
CACCGTCCCAGACAAGACCAGGACCCTCTGTGTAGTAGCCACTGACCCACCTGTGTTCTTACGATCTGATTTC // TAAGTACCCCTGGTTACGGACTCGGATGGAGAC

Consensus site for MboII

d) A77D +/-

GTGGCAGGGTCTGTTCTGGTCTGGAAGACATCATCGGTGACTGGGTGGACAAGAATGCTAGACTAAAG // ATTCAATGGGACCAATGCCCTGAGCTACCTCTG
CACCGTCCCAGACAAGACCAGGACCCTCTGTGTAGTAGCCACTGACCCACCTGTGTTCTTACGATCTGATTTC // TAAGTACCCCTGGTTACGGACTCGGATGGAGAC

Figure 2

