(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
21 June 2018 (21.06.2018)

WIPO | PCT

(10) International Publication Number
WO 2018/107244 A1

(51) International Patent Classification:
A61K 31/704 (2006.01) *A61K 39/395* (2006.01)
A61K 31/702 (2006.01) *A61P 35/00* (2006.01)
A61K 31/573 (2006.01)

(21) International Application Number:
PCT/AU2017/051402

(22) International Filing Date:
15 December 2017 (15.12.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2016905199 15 December 2016 (15.12.2016) AU

(71) Applicant: **PROGEN PG500 SERIES PTY LTD** [AU/AU]; c/- Cullens Pty Ltd, Level 32, 239 George Street, Brisbane, Queensland 4000 (AU).


(72) Inventor: **DREDGE, Keith**; c/- Cullens Pty Ltd, Level 32, 239 George Street, Brisbane, Queensland 4000 (AU).

(74) Agent: **CULLENS PTY LTD**; Level 32, 239 George Street, Brisbane, Queensland 4000 (AU).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: COMPOSITION AND USES THEREOF



Formula (I)

Figure 1



(57) Abstract: The present invention relates, *inter alia*, to methods and uses for the prevention or treatment of cancer, and to compositions, combinations and kits for the treatment or prevention of cancer. In one embodiment, the invention provides a method of treating or preventing cancer, comprising administering to a patient in need thereof an immune checkpoint modulator, such as the PD- 1 inhibitor Nivolumab, together with a compound of Formula (I), wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M, such the compound PG-545 (pixatimod).

WO 2018/107244 A1

[Continued on next page]

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

COMPOSITION AND USES THEREOF

TECHNICAL FIELD

[0001] The present invention relates, *inter alia*, to methods and uses for the prevention or treatment of cancer, and to compositions, combinations and kits for the treatment or prevention of cancer.

BACKGROUND ART

[0002] It will be clearly understood that, if a prior art publication is referred to herein, this reference does not constitute an admission that the publication forms part of the common general knowledge in the art in Australia or in any other country.

[0003] The immune system can be effective in combating malignant tumour cells that arise in the human body. If malignant tumour cells can evade the immune system they can proliferate and spread, developing into life threatening cancer. Numerous cancers have developed the capability to evade the immune system, many of them very serious diseases that affect the health and reduce the life expectancy of a large number of people worldwide.

[0004] Success has been achieved in the treatment of cancer using therapeutics that specifically target the processes employed by cancer cells to evade the immune system. One key process involves checkpoint targets (such as checkpoint receptors and checkpoint ligands). In healthy tissue, checkpoint targets are important in preventing over stimulation of the immune system that could otherwise lead to autoimmune diseases. However, many types of cancer cells interact with one or more of these checkpoint targets thus allowing them to lessen or inactivate an immune response. Checkpoint modulators are therapeutics which modulate checkpoint targets on the surface of immune cells, preventing cancer cells from selectively engaging these checkpoint targets to inactivate the immune cells.

[0005] Pharmaceuticals that target the checkpoint receptors PD-1 and CTLA-4, for example, have recently been approved to treat a variety of cancers. Nivolumab and pembrolizumab are inhibitors of PD-1 and ipilimumab inhibits CTLA-4. The cancers for which these agents have been approved include melanoma (ipilimumab, nivolumab, pembrolizumab; Franklin 2016), non-small cell lung cancer (NSCLC) (nivolumab, pembrolizumab; Chen 2016), renal cell cancer (RCC) (nivolumab; Mennitto 2016) and Hodgkin lymphoma (nivolumab; US FDA 2016). However, even in these cancers there are numerous patients who derive limited, or no benefit from treatment.

[0006] PG545 is a cholestanol-sulfotetrasaccharide conjugated small molecule compound, especially for the treatment of cancer. PG545 has been described as a heparanase inhibitor (Dredge *et al*, 2010, Hammond *et al*, 2013) but also possesses immunomodulatory properties. Use of this compound has been demonstrated to lead to a depletion of immunosuppressive pro-cancer immune cells such as tumour-associated macrophages (TAMs) and myeloid-derived suppressor cells (MDSCs) within the tumour microenvironment (TME) as evidenced by preclinical models of pancreatic cancer (Ostapoff *et al*, 2013) and a chemically-induced model of skin cancer (Boyango *et al*, 2014).


[0007] However, PG545 also drives a strong immunostimulatory activity on the dendritic cell (DC), another myeloid-derived cell, via the MyD88-dependant toll-like receptor 9 (TLR9) pathway and the subsequent activation of natural killer (NK) cells (Brennan *et al*, 2016). The authors also found that the immunostimulatory effect of PG545 was reproduced in heparanase knockout mice but that another heparanase inhibitor, muparfostat (PI-88), did not have this activity profile in normal mice (unpublished data). Taken together these results demonstrate that PG545's effect on DCs is heparanase-independent. Moreover, PG545's effect on DCs led to potent activation of NK cells but not T cells (Brennan *et al*, 2016). As illustrated in Brennan *et al*, 2016, administration of PG545 alone does not activate T cells in naïve mice (see Figure 3C), nor in mice bearing A20 lymphoma tumours (see Figure 2C and E of the Supplementary Material). Thus, the immunomodulatory activity of PG545 has not shown any interaction or impact on T cells or adaptive immunity.

SUMMARY OF INVENTION

[0008] The present invention is directed, *inter alia*, to methods, compositions or kits which may improve the immune response to cancer cells, or which may provide the consumer with a useful or commercial choice.

[0009] With the foregoing in view, the present invention in some forms resides broadly in methods, uses, combinations, kits and compositions for treating or preventing cancer.

[0010] In a first aspect, the present invention provides a method of treating or preventing cancer, comprising administering to a patient in need thereof an immune checkpoint modulator together with a compound of Formula (I):



Formula (I)


wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0011] Advantageously, it has been found that a compound of Formula (I) and an immune checkpoint modulator, especially a T cell immune checkpoint inhibitor, have a synergistic effect to accumulate T cells in tumours. This effect is especially surprising, given that a compound of Formula (I) has been shown to have no interaction or impact on T cells (Brennan *et al*, 2016). Furthermore, in some studies the immune checkpoint modulator when administered alone has been shown to provide a limited effect on the accumulation of T cells in tumours (see Spranger 2016 for example). This synergistic effect is believed to enable a patient's immune system to provide an enhanced immune response specifically against cancer cells.

[0012] In one embodiment, at least 9 of the X groups in the compound of Formula (I) is SO_3M. In another embodiment, at least 10 of the X groups in the compound of Formula (I) is SO_3M; especially at least 11 of the X groups in the compound of Formula (I) is SO_3M; more especially at least 12 of the X groups in the compound of Formula (I) is SO_3M; most especially 13 or all of the X groups in the compound of Formula (I) is SO_3M. In one embodiment, at least 75% of the X groups is SO_3M; especially at least 80% of the X groups is SO_3M; more especially at least 90% of the X groups is SO_3M; most especially 100% of the X groups is SO_3M.


[0013] In the compound of Formula (I), M may be any pharmaceutically acceptable cation. Exemplary cations include sodium, calcium, potassium, magnesium or ammonium; especially sodium or potassium; more especially sodium.

[0014] In one embodiment, the compound of Formula (I) may be a compound of Formula (II):




Formula (II)

[0015] In another embodiment, the compound of Formula (I) may be a compound of Formula (III):




Formula (III)

[0016] In another embodiment, the compound of Formula (I) may be a compound of Formula (IV):




Formula (IV)

[0017] In another embodiment, the compound of Formula (I) may be a compound of Formula (V):




Formula (V)

[0018] In another embodiment, the compound of Formula (I) may be a compound of Formula (VI):



Formula (VI)

[0019] In another embodiment, the compound of Formula (I) may be a compound of Formula (VII):



Formula (VII)

[0020] The compounds of Formulae (II) to (VII) possess asymmetric centres. These compounds may be in substantially pure isomeric form at one or more of said centres, for example greater than about 90% enantiomeric excess (ee) or diasteromeric excess (de), such as

greater than 95% ee or de, or greater than 97% ee or de, or greater than 99 % ee or de. Such isomers may be prepared by asymmetric synthesis, for example using chiral starting materials or intermediates, or by chiral resolution.

[0021] If the compound of Formula (I) is a compound of Formula (VII), and X is SO_3Na, then the compound is PG545. In one embodiment, the compound of Formula (I) is PG545. PG545 is 3β,5α-cholestanyl 2,3,4,6-tetra-*O*-sodium sulfonato-α-D-glucopyranosyl-(1→4)- 2,3,6-tri-*O*-sodium sulfonato-α-D-glucopyranosyl-(1→4)-2,3,6-tri-*O*-sodium sulfonato-α-D-glucopyranosyl-(1→4)-2,3,6-tri-*O*-sodium sulfonato-β-D-glucopyranoside, and is described in WO2009/049370.

[0022] In one embodiment, the immune checkpoint modulator is an immune checkpoint inhibitor or an immune checkpoint activator; especially an immune checkpoint inhibitor. In another embodiment, the immune checkpoint modulator targets one or more of a T cell, an Antigen-Presenting Cell (APC) / tumour cell, a Natural Killer (NK) cell, a Natural Killer T cell (NKT), a Gamma Delta T cell, an invariant T cell, or an invariant Natural Killer T cell (NKT). The immune checkpoint modulator especially targets a T cell; and is especially a T cell immune checkpoint inhibitor. The immune checkpoint modulator may target a CD8 T cell, especially an effector memory CD8 T cell or a central memory CD8 T cell. The immune checkpoint modulator may target a CD4 T cell.

[0023] The immune checkpoint modulator may target one or more of: CTLA-4, PD-1, PD-L1, LAG3, B7-H3, B7-H4, TIM3, Galectin9, BTLA, HVEM, A2aR, VISTA, KIR, TIGIT, IDO, CD47, CD28, ICOS, CD137 (4-1BB), CD137L, CD40, CD40L, OX40, OX40L, GITR and CD27. The immune checkpoint modulator may target one or more of: CTLA-4, PD-1, LAG3, TIM3, BTLA, A2aR, CD28, ICOS, CD137 (4-1BB), CD40L, OX40, GITR and CD27. The immune checkpoint modulator may target one or more of PD-1, CTLA-4 or PD-L1. The immune checkpoint modulator may target PD-1. The immune checkpoint modulator may be a PD-1 inhibitor.

[0024] The immune checkpoint modulator may be one or more of: Ipilimumab, Tremelimumab, Nivolumab, Pembrolizumab, Pidilizumab, Atezolizumab, Avelumab, Durvalumab, Enoblituzumab, Lirilumab, Epacadostat, Indoximod, Urelumab, Utomilumab, Lucatumumab, Dacetuzumab, Oxelumab, and Varlilumab. The immune checkpoint modulator may especially be Nivolumab, Pembrolizumab or Pidilizumab.

[0025] Exemplary immune checkpoint targets, cell types and agents are provided in Table 1.

Table 1: Immune Checkpoint Targets and Agents

Target	Cell Type	Inhibitory / Activating	Agent	Synonym
CTLA-4	T cell	**Inhibitory**	Ipilimumab	-
	T cell	**Inhibitory**	Tremelimumab	CP-675,206
PD-1	T cell	**Inhibitory**	Nivolumab	MDX-1106 (BMS-936558)
	T cell	**Inhibitory**	Pembrolizumab	MK-3475 (lambrolizumab)
	T cell	**Inhibitory**	Pidilizumab	CT-011
	T cell	**Inhibitory**		AMP-224
PD-L1	APC/tumour cell	**Inhibitory**	Atezolizumab	MDPL3280 (RG7446)
	APC/tumour cell	**Inhibitory**	Avelumab	MSB0010718C
	APC/tumour cell	**Inhibitory**	Durvalumab	MEDI-4736
	APC/tumour cell	**Inhibitory**	N/A	MDX-1105 (BMS-936559)
LAG3	T cell	**Inhibitory**	N/A	IMP321
	T cell	**Inhibitory**	N/A	IMP701
	T cell	**Inhibitory**	N/A	LAG525
B7-H3	APC/tumour cell	**Inhibitory**	Enoblituzumab	MGA271
B7-H4	APC/tumour cell	**Inhibitory**	N/A	Developed by AstraZeneca
TIM3	T cell	**Inhibitory**	N/A	MGB453
	T cell	**Inhibitory**	N/A	CA-327
Galectin9	APC/tumour cell	**Inhibitory**	N/A	Developed by GlycoMimetics
BTLA	T cell	**Inhibitory**	N/A	
HVEM	APC/tumour cell	**Inhibitory**	N/A	
A2aR	T cell	**Inhibitory**	N/A	CPI-444
VISTA	APC/tumour cell	**Inhibitory**	N/A	JNJ 61610588
	APC/tumour cell	**Inhibitory**	N/A	CA-170
KIR	NK cell	**Inhibitory**	Lirilumab	BMS-986015 (IPH2102)
TIGIT	NK cell	**Inhibitory**	N/A	MTIG7192A, RG6058
IDO	APC/tumour cell	**Inhibitory**	Epacadostat	INCB024360
	APC/tumour cell	**Inhibitory**	Indoximod	NLG-8189
CD47	APC/tumour cell	**Inhibitory**	N/A	Hu5F9-G4
CD28	T cell	**Activating**	N/A	TGN1412
ICOS	T cell	**Activating**	N/A	JTX-2011
CD137 (4-1BB)	T cell	**Activating**	Urelumab	BMS-663513
	T cell	**Activating**	Utomilumab	PF-05082566 (PF-2566)
CD137L	APC/tumour cell	**Activating**	N/A	
CD40	APC/tumour cell	**Activating**	N/A	SEA-CD40
	APC/tumour cell	**Activating**	Lucatumumab	HCD120
	APC/tumour cell	**Activating**	Dacetuzumab	SGN-40, huS2C6
CD40L	T cell	**Activating**	N/A	
OX40	T cell	**Activating**	N/A	Anti-OX40 mAb (9B12)
OX40L	APC/tumour cell	**Activating**	Oxelumab	
GITR	T cell	**Activating**	N/A	TRX518, INCAGN1876
	T cell	**Activating**	N/A	MK-4166
CD27	T cell	**Activating**	Varlilumab	CDX-1127

APC – Antigen-presenting cell
NK – Natural Killer cell
N/A – Not available

[0026] The patient may be a mammal, especially a human. However, the patient may be a primate or a veterinary patient, such as a dog, cat or horse.

[0027] The cancer may be a malignant or a benign cancer, especially a malignant cancer. The cancer may be a solid tumour or a haematological tumour. The haematological tumour may be a haematological malignancy such as, for example, lymphoma, myeloma or leukemia. The solid tumour may be selected from the group consisting of: melanoma, colon cancer, breast cancer, prostate cancer, liver cancer, lung cancer (including non-small cell, small cell or mesothelioma), head and neck cancer, skin cancer, renal cancer, pancreatic cancer, oesophageal cancer, cholangiocarcinoma, gastric cancer, urothelial or bladder cancer, sarcoma, uterine cancer, endometrial cancer, ovarian cancer, cervical cancer and brain cancer (including glioma and astrocytoma). The cancer may be breast cancer.

[0028] The cancer may be a tumour enriched with T cells. In another embodiment, the cancer may be a tumour including substantially no T cells, or a tumour resistant to accumulation of T cells. A tumour including substantially no T cells, or a tumour resistant to accumulation or infiltration of T cells is sometimes called a "cold" or "non-inflamed" tumour, and such terms would be known to a skilled person (see Spranger 2016, Woo 2015, for example). It is believed that the compound of Formula (I) and the immune checkpoint modulator may inflame or convert such a "cold" tumour into a "hot" or "inflamed" tumour which is enriched with T cells, thereby improving the immune response to the tumour.

[0029] As a result of administration of the immune checkpoint modulator and the compound of Formula (I), cancer may be more effectively treated or prevented, and/or a medically beneficial effect on a tumour may be provided. Administration of the immune checkpoint modulator and the compound of Formula (I) may initiate, enable, increase, enhance or prolong the activity of immune cells against the cancer and/or may provide an anti-tumour adaptive immune response.

[0030] Administration of the immune checkpoint modulator and the compound of Formula (I) may increase the number of immune cells accumulating in the tumour. Administration of the immune checkpoint modulator and the compound of Formula (I) may lead to accumulation of lymphocytes in a tumour; especially T cells; more especially Effector Memory $CD8^+$ T cells, Central Memory $CD8^+$ T cells and $CD4^+$ T cells.

[0031] As used herein, the terms "treatment" (or "treating") and "prevention" (or "preventing") are to be considered in their broadest contexts. For example, the term "treatment"

does not necessarily imply that a patient is treated until full recovery. The term "treatment" includes amelioration of the symptoms of a disease or condition, or reducing the severity of a disease or condition. Similarly, "prevention" does not necessarily imply that a subject will never contract a disease or condition. "Prevention" may be considered as reducing the likelihood of onset of a disease or condition, or preventing or otherwise reducing the risk of developing a disease or condition.

[0032] The method may include administering an effective amount of the immune checkpoint modulator and/or an effective amount of the compound of Formula (I). An "effective amount" of a compound of Formula (I) means an amount necessary to at least partly attain the desired response, or to delay the onset or progression of the cancer. The amount may vary depending on factors such as: the health and physical condition of the individual to whom the compound is administered, the taxonomic group of the individual to whom the compound is administered, the extent of treatment / prevention desired, the formulation of the composition, and the assessment of the medical situation. It is expected that the "effective amount" will fall within a broad range that can be determined through routine trials. An effective amount in relation to a human patient, for example, may lie in the range of about 0.1 ng per kg of body weight to 1 g per kg of body weight per dosage, or in the range of about 100 ng to 100 mg per kg of body weight per dosage. Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several doses may be administered daily, bi-weekly or weekly, or at other suitable time intervals, or the dose may be proportionally reduced as indicated by the circumstances. Decisions on dosage and the like would be within the skill of the medical practitioner or veterinarian responsible for the care of the patient.

[0033] The compound of Formula (I) and the immune checkpoint modulator may be administered sequentially, simultaneously or substantially simultaneously. The compound of Formula (I) may be administered prior to, or after, the immune checkpoint modulator. Either or both of the compound of Formula (I) and the immune checkpoint modulator may be administered as part of a treatment regime or cyclical dosage. The immune checkpoint modulator and the compound of Formula (I) may accumulate in the patient over time, and consequently the immune checkpoint modulator and the compound of Formula (I) may be administered hours or days apart yet still act synergistically.

[0034] While it is possible that the compound of Formula (I) and the immune checkpoint modulator may be administered as neat chemicals, they may be administered as part of a pharmaceutical composition which includes a carrier or excipient. The compound of Formula (I)

and the immune checkpoint modulator may be provided together in a single pharmaceutical composition. Alternatively, the compound of Formula (I) and the immune checkpoint modulator may be provided in separate pharmaceutical compositions. If the compound of Formula (I) and the immune checkpoint modulator are to be administered at different times, or at varying dosages, it may be advantageous to provide the compound of Formula (I) and the immune checkpoint modulator in separate pharmaceutical compositions.

[0035] Any said pharmaceutical composition may comprise the active agent (i.e. a compound of Formula (I) and/or the immune checkpoint modulator) and a pharmaceutically acceptable carrier or excipient. Any pharmaceutically acceptable carrier(s) or excipient(s) must be acceptable in the sense of being compatible with the other components in the composition and not being deleterious to the patient.

[0036] The type of pharmaceutical composition may depend upon the Absorption, Distribution, Metabolism and Excretion (ADME) profile of the active agent. For example, it may be most appropriate for compounds of Formula (I) and/or immune checkpoint inhibitors to be administered parenterally, especially intravenously, and consequently the pharmaceutical composition may be formulated for parenteral or intravenous administration. Nevertheless, if possible the pharmaceutical composition may include those suitable for oral or rectal administration, or for administration by non-intravenous routes.

[0037] Parenteral administration may include administration by one or more of the following routes: intravenously, intrathecally, cutaneously, subcutaneously, nasally, intramuscularly, intraocularly, transepithelially, vaginally, intraperitoneally and topically. Topical administration includes buccal, sub-lingual, dermal, ocular, rectal, nasal, as well as administration by inhalation or by aerosol means. For intravenous, cutaneous or subcutaneous injection, or injection at a site where treatment is desired, the active agent may be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of skill in the art would be able to prepare suitable solutions.

[0038] The nature of the pharmaceutical composition and the carrier or excipient will depend on the route of administration and the nature of the condition and the patient being treated. It is believed that the choice of a particular carrier, excipient or delivery system, and route of administration could be readily determined by a person skilled in the art. In some circumstances it may be necessary to protect the active agent by means known in the art, for example, by micro encapsulation. The route of administration should also be chosen such that the active agent reaches its site of action. The pharmaceutical composition may include any

suitable effective amount of the active agent commensurate with the intended dosage range to be employed.

[0039] The pharmaceutical composition may be in the form of a solid (including tablets, filled capsules, powders, cachets, capsules, troches, suppositories, wafers, dispersible granules and pessaries), or a liquid (including solutions, suspensions, syrups, emulsions, colloids, elixirs, creams, gels and foams). In one embodiment, the pharmaceutical composition may be in the form of a sterile injectable solution for parenteral use.

[0040] The pharmaceutically acceptable carrier or excipient may be either a solid or a liquid. The carrier or excipient may act as a diluent, buffer, stabiliser, isotonicising agent, flavouring agent, anti-oxidant, solubilizer, lubricant, suspending agent, binder, preservative, tablet disintegrating agent or an encapsulating material. Suitable carriers and excipients would be known to a skilled person. With regard to buffers, aqueous compositions may include buffers for maintaining the composition at close to physiological pH or at least within a range of about pH 6.0 to 9.0.

[0041] If the pharmaceutical composition is a powder, the active agent and a carrier or excipient may both be finely divided powders which are mixed together.

[0042] If the pharmaceutical composition is a tablet, the active component may be mixed with a suitable amount of a carrier or excipient which has the necessary binding capacity before compaction into a tablet of the desired shape and size.

[0043] Powders or tablets may include any suitable amount of the active agent, and exemplary amounts of the active agent in the powder or tablet may range from about five or ten percent to about seventy percent. Exemplary carriers or excipients for powders and tablets may include, for example, magnesium carbonate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, a low melting wax, cocoa butter and the like.

[0044] Liquid form preparations may include, for example, water, saline, water-dextrose, water-propylene glycol, petroleum, or oil (including animal, vegetable mineral or synthetic oil) solutions. For example, parenteral injection liquid preparations may be formulated as solutions in aqueous polyethylene glycol solution. Such liquid form preparations may contain at least 0.1 wt% of the active compound.

[0045] Liquid pharmaceutical compositions may be formulated in unit dose form. For example, the compositions may be presented in ampoules, pre-filled syringes, small volume

infusion or in multi-dose containers. Such compositions may include a preservative. The compositions may also include formulatory agents such as suspending, stabilising and/or dispersing agents. The composition may also be in powder form for constitution with a suitable vehicle (such as sterile water) before use. Liquid carriers and excipients may include colorants, flavours, stabilizers, buffers, artificial and natural sweeteners, dispersants, thickeners, solubilizing agents, suspending agents and the like.

[0046] Aqueous solutions for oral use may be prepared by dissolving the active agent in water and adding colourants, thickeners, flavours, and stabilizing agents, as necessary. Aqueous suspensions for oral use may be prepared by dispersing the active agent in water with viscous material, such as natural or synthetic gums, resins, methyl cellulose or other suspending agents.

[0047] For topical administration to the epidermis the compounds may be formulated as an ointment, cream or lotion, or as a transdermal patch.

[0048] The compositions may also be administered by inhalation in the form of an aerosol spray from a pressurised dispenser or container, which contains a propellant such as carbon dioxide gas, dichlorodifluoromethane, nitrogen, propane or other suitable gas or gas combination. The pharmaceutical composition may be in a form suitable for administration by inhalation or insufflation.


[0049] The pharmaceutical composition may be adapted to provide sustained release of the active agent.

[0050] The pharmaceutical composition may be in unit dosage form. In such form, the pharmaceutical composition may be prepared as unit doses containing appropriate quantities of the active agent. The unit dosage form may be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

[0051] The pharmaceutical composition may also include at least one further active ingredient, in addition to the compound of Formula (I) and/or the immune checkpoint modulator. Such ingredients may have efficacy as anti-cancer, anti-angiogenic, anti-metastatic, anti-inflammatory, anti-coagulant, antimicrobial or anti-thrombotic agents, and agents effective against elevated blood triglyceride levels and cardiovascular disease. However, the pharmaceutical composition may include any suitable further active ingredient.

[0052] In a second aspect, the present invention provides a use of a compound of Formula (I) in the manufacture of a medicament for the treatment or prevention of cancer,

wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M;

and wherein the medicament is formulated for administration with an immune checkpoint modulator.

[0053] In a third aspect, the present invention provides a use of an immune checkpoint modulator in the manufacture of a medicament for the treatment or prevention of cancer,

wherein the medicament is formulated for administration with a compound of Formula (I):




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0054] In a fourth aspect, the present invention provides a use of an immune checkpoint modulator and a compound of Formula (I) in the manufacture of a medicament for the treatment or prevention of cancer,

wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0055] In a fifth aspect, the present invention provides an immune checkpoint modulator and a compound of Formula (I) for use in the treatment or prevention of cancer;

wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0056] In a sixth aspect, the present invention provides a pharmaceutical composition

comprising a compound of Formula (I) and an immune checkpoint modulator; wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0057] Features of the second to sixth aspects of the present invention may be as described for the first aspect of the present invention. The medicament of the second to fourth aspects of the present invention may be a pharmaceutical composition, as described above.

[0058] In a seventh aspect, the present invention provides a kit comprising a compound of Formula (I) and an immune checkpoint modulator; wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0059] Features of the seventh aspect of the present invention may be as described for the first to sixth aspects of the present invention. The kit may include a single pharmaceutical

composition or at least two pharmaceutical compositions. The kit may include an immune checkpoint modulator and the compound of Formula (I) in the same pharmaceutical composition or in separate pharmaceutical compositions.

[0060] In an eighth aspect, the present invention provides a pharmaceutical combination comprising a compound of Formula (I) and an immune checkpoint modulator; wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0061] Features of the eighth aspect may be as described for the first to seventh aspects of the present invention. The pharmaceutical combination may be a pharmaceutical composition, or the pharmaceutical combination may include at least two pharmaceutical compositions. The pharmaceutical combination may include an immune checkpoint modulator and the compound of Formula (I) in the same pharmaceutical composition or in separate pharmaceutical compositions.

[0062] In a ninth aspect, the present invention provides a method of enhancing an immune response in a patient, comprising administering to a patient in need thereof an immune checkpoint modulator together with a compound of Formula (I):



Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

[0063] Features of the ninth aspect of the present invention may be as described for the first to eighth aspects of the present invention.

[0064] Any of the features described herein can be combined in any combination with any one or more of the other features described herein within the scope of the invention.

BRIEF DESCRIPTION OF DRAWINGS

[0065] Examples of the invention will now be described by way of example with reference to the accompanying Figures, in which:

[0066] Figure 1 shows the effect of treatments on 4T1.2 tumour bearing mouse body weight. PG545 was administered intraperitoneally on days 1, 8 and 15 (vertical lines) and antibodies on days 1, 4, 8, 11 and 15;

[0067] Figure 2 shows the anti-tumour activity of PD-1 antibody and PG545 alone and in combination against 4T1.2 tumours *in vivo*;

[0068] Figure 3 shows the mean tumour volume and mean weight change of the mice over the course of the study of Example 2;

[0069] Figure 4 shows a flow cytometric analysis of the bulk tumour cultures from each treatment group;

[0070] Figure 5 shows therapy-induced changes in 4T1.2 tumour and spleen associated CD8 T cell frequency;

[0071] Figures 6A and 6B shows the gating strategy for analysing the CD8 T responses to therapy in the 4T1.2 mammary tumours;

[0072] Figure 7 shows histograms demonstrating the granzyme B status for both the tumour-associated effector and central memory CD8 T cell populations in each treatment group;

[0073] Figure 8 shows therapy-induced changes in 4T1.2 tumour and spleen associated CD4 T cell frequency;

[0074] Figure 9 shows flow cytometric analysis demonstrating the tumour-associated changes in the CD4 T cell compartment within each treatment group (concatenated results shown for each treatment group (4 tumours/group));

[0075] Figure 10 shows therapy-induced changes in 4T1.2 tumour and spleen associated NK cell frequency;

[0076] Figure 11A and 11B shows the gating strategy for analysing the NK cell responses to therapy in the 4T1.2 mammary tumours;

[0077] Figure 12 shows therapy-induced changes in 4T1.2 tumour myeloid cell frequency;

[0078] Figure 13A shows the gating strategy for analysing the myeloid cell responses to therapy (Group 1 - Vehicle + Control antibody) in the 4T1.2 mammary tumours;

[0079] Figure 13B shows the gating strategy for analysing the myeloid cell responses to therapy (Group 2 - PG545 + Control antibody) in the 4T1.2 mammary tumours;

[0080] Figure 13C shows the gating strategy for analysing the myeloid cell responses to therapy (Group 3 - Vehicle + Anti-PD-1) in the 4T1.2 mammary tumours;

[0081] Figure 13D shows the gating strategy for analysing the myeloid cell responses to therapy (Group 4 - PG545 + Anti-PD-1) in the 4T1.2 mammary tumours; and

[0082] Figure 14 shows therapy-induced changes in PD-L1 expression on 4T1.2 tumour associated myeloid cells.


[0083] Preferred features, embodiments and variations of the invention may be discerned from the following Examples which provides sufficient information for those skilled in the art to perform the invention. The following Examples are not to be regarded as limiting the scope of the preceding Summary of the Invention in any way.

EXAMPLES

[0084] Examples of the present invention will now be described with reference to Figures 1 to 14.

Compounds used

[0085] PG545 (or 3β,5α-Cholestanyl 2,3,4,6-tetra-*O*-sodium sulfonato-α-D-glucopyranosyl-(1→4)- 2,3,6-tri-*O*-sodium sulfonato-α-D-glucopyranosyl-(1→4)-2,3,6-tri-*O*-sodium sulfonato-α-D-glucopyranosyl-(1→4)-2,3,6-tri-*O*-sodium sulfonato-β-D-glucopyranoside), has the structure below. PG545 has been allocated a proposed International Nonproprietary Name (INN) by the World Health Organization of pixatimod (for the acid). This compound is described in WO2009/049370. PG545 may be synthesised as provided in Ferro *et al* 2012 or in WO2009/049370.



[0086] Anti-PD-1 antibody (RMP1-14) was purchased from Bio-X-Cell (NH, USA).

[0087] Isotype Control antibody (2A3) was purchased from Bio-X-Cell (NH, USA).

Example 1 – Study in the 4T1.2 Tumour Model

[0088] Female Balb/c mice (Age 8 weeks; Walter and Eliza Hall Institute of Medical Research (WEHI), Victoria Australia) were inoculated in the intramammary fatpad with 1 x 10^5 4T1.2 cells in phosphate buffered saline. 4T1.2 is a mouse mammary tumour model.

[0089] Mice were weighed and tumours measured 2-3 times weekly using electronic callipers. Tumour volume (mm^3) was calculated as length (mm)/2 x width $(mm)^2$. One week following implantation mice with similar sized tumours (mean tumour volume 56 mm^3) were randomised into 4 groups of 6 animals (Day 1). Treatment groups were saline + control antibody, PG545 + control antibody, saline + anti-PD-1 and PG545 + anti-PD-1.

[0090] PG545 was given at 15 mg/kg weekly (0.1 ml/10g body weight) for 3 weeks and anti-PD-1 or control antibody (200 μg; 100 μL of a 2mg/mL solution) was given intraperitoneally on days 1, 4, 8, 11 and 15. Mice received a small dish containing a food supplement (Ensure mixed with food dust) daily. A dosage of 15 mg/kg PG545 was selected in view of the body weight issues observed at 20 mg/kg PG545 in these mice (i.e. issues with maximum tolerated dose at 20 mg/kg PG545. The maximum tolerated drug dose (MTD) is defined as that causing 10 % body weight loss and from which the animals recovered weight to baseline levels within 7-10 days).

[0091] The experiment was ended on day 18 or earlier if an ethical endpoint was met. Blood was collected via cardiac bleed and serum was prepared and frozen at -80°C.

[0092] The mean body weight changes following the treatments are shown in Figure 1. In Figure 1, the data represent the mean percent weight change from day 1 for each group; bars are standard error of the mean (SEM). At day 15 one mouse in the saline + control antibody group was found dead. At day 18: (i) in the PG545 + anti-PD-1 group one mouse was found dead, and the remaining mice showed signs of distress (ruffled fur); and (ii) in the PG545 + control antibody group four of the six mice were bloated and showed signs of distress (ruffled fur). It was decided to harvest the experiment on day 18.

[0093] The percentage tumour growth inhibition was determined according to the following formula: 100 x (1-ΔT/ ΔC) where ΔC and ΔT were calculated by subtracting the mean tumour volume in each group on day 1 of treatment from the mean tumour volume on the day of analysis. Statistical analysis of the *in vivo* data was performed using Graph Pad Prism Version 6.0 (Graph Pad La Jolla, CA). An ANOVA analysis was performed followed by Dunnett's post hoc test to compare the tumour growth in the treated groups to the vehicle control.

[0094] Figure 2 summarises the effect of the treatment on 4T1.2 tumour growth. Tumour growth in the PG545 + control antibody, saline (or vehicle) + anti-PD-1 and PG545 + anti-PD-1 groups was inhibited by 68%, 44% and 84%, respectively on day 18. Tumour growth in the PG545 + control antibody and PG545 + anti-PD-1 groups was significantly inhibited compared to the saline (or vehicle) + control antibody group ($P < 0.01$ for both). No significant differences were seen between any other groups. In Figure 2, tumour volumes are expressed as mean tumour volume (± SEM); ** = $P < 0.01$ compared to vehicle + control antibody.

Example 2 – Study to Evaluate Tumour Infiltrating Leukocytes

[0095] Sixteen mice from the cohort implanted in Example 1 above were randomised 1

week after inoculation into 4 groups of 4 mice as for Example 1 (mean tumour volume = 77 mm^3). PG545 was given intraperitoneally at 15 mg/kg (0.1 ml/10g body weight) on days 1 and 8 with anti-PD-1 or control antibody (100 µl) given intraperitoneally on days 1, 4 and 8.

[0096] On day 11, mice were euthanized and tumours harvested. The tumours were mechanically disaggregated and then digested in collagenase IV and DNase 1 before being filtered through a cell strainer to yield a single cell suspension. Spleens were also removed and used as background staining controls. Where appropriate, the cells were fixed and permeabilised [Ebioscience or BD Pharmingen Fixation/Permeabilisation Kits; also see the procedure provided in the technical data sheet for BD Cytofix/Cytoperm™ available from www.bdbiosciences.com]. The cells were then incubated with the antibody cocktails summarised in Table 2. Flow cytometry analysis was performed on the LSR II analyser (BD Biosciences).

Table 2: Immune Staining Cocktails

Stain 1	**Stain 2**	**Stain 3**	**Stain 4**
CD45.2-eFluor450	CD45.2-APC	CD45.2-PEcy7	CD45.2-eFluor450
TCRb-APC-cy7	TCRb-PE	CD49b-APC	Ly6C-PEcy7
CD8b-PEcy7	CD4-eFluor450	CD27-PE	CD11b-APCcy7
CD44-FITC	FoxP3-FITC	CD11b-APCcy7	Ly6G-BV711
CD62L-BV510	CD8-PEcy7	CD69-FITC	CD11c-FITC
CD69-PE	NO DAPI	CD335-eFluor450	MHCII-APC
Granzyme B		DAPI	PD-L1-PE
NO DAPI			DAPI

[0097] Staining set summary:

- Stain 1 (fixed group due to Granzyme B intracellular stain). Stopping gate for collection was set on CD45.2 for all groups to remove any bias associated with tumour size.
- Stain 2. Did not produce a clear data set and was excluded from further analysis.
- Stains 3 and 4. Stopping gate for collection was set on DAPI negative cells (viable cell collection date) for all groups to remove any bias associated with tumour size.

[0098] In this study statistical analysis was performed using the Mann Whitney U Test.

[0099] The mean tumour volume and body weight change of the mice in this study are provided in Figure 3.

[00100] A significant increase in lymphocyte frequency was observed within the PG545 +

anti-PD-1 treated tumours relative to the single agent and control treated tumours (Figure 4). This increase in 4T1.2 tumour-associated lymphocyte frequency correlated with a reduction in CD45.2 negative cells in the PG545 + anti-PD-1 treated tumours; highlighting the impact of the combination therapy on tumour and/or stromal cell viability. In Figure 4, data shows the concatenated (merged) results from 4 mice in each group. Group 1 = Vehicle (saline) + control antibody (CIg); Group 2 = PG545 + Control antibody; Group 3 = Vehicle (saline) + anti-PD-1; Group 4 = PG545 + anti-PD-1. Row 1 shows the concatenated fluorescence-activated cell sorting (FACs) blots for CD45.2 versus FSC. Row 2 shows the concatenated FACs blots for SSC versus FSC (morphology plot) on the CD45+ leukocyte gate.

[00101] The increase in tumour-associated lymphocytes in response to PG545 + anti-PD-1 treatment correlated with a significant increase in both CD8 (TCRb+CD8+) and CD4 (TCRb+CD8-) T cells (Figure 5; FACS data collected for Stain 2 was deemed uninterruptible, so the treatment-associated effects on CD4+ T effector cells and T regulatory cells was not accurately quantitated). The gating strategy used to analyse Stain 1 is shown in Figure 6 (in which concatenated results are shown for each treatment group (4 tumours/group)). Notably these therapy-induced changes in T cell frequency were not observed in the spleen (Figure 5); highlighting the tumour-specific nature of these effects. In Figure 5, each symbol represents an individual tumour. The horizontal line shown in each treatment group represents the average frequency of tumour-associated CD8 T cells for the group of 4 mice. Spleens from each treatment group were combined for analysis. Each horizontal bar represents the average splenic associated CD8 T cell frequency for each treatment group.

[00102] Within the tumour-associated CD8+ T cell compartment, a significant increase in the frequency of Central Memory $CD8^+$ T cells (CD62L+CD44+ cells) was observed (Figure 5). Within the PG545 + anti-PD-1 treated tumours the frequency of Effector Memory CD8+ T cells was elevated relative to the control and anti-PD-1 treated groups (Figure 5). However, the frequency of CD69+Granzyme B+ Effector CD8+ T cells appeared comparable across the four treatment groups (Figures 6 and 7). In Figure 7, concatenated results are shown for each treatment group (four tumours/group).

[00103] The frequency of tumour-associated $CD4^+CD44^+CD62L^+$ and $CD4^+CD44^+CD62L^-$ T cells was elevated in the PG545 + anti-PD-1 group compared to the single agent and control groups (Figure 8, Figure 9). These changes in the CD4 T cell compartment of the PG545 + anti-PD-1 treated tumours were not evident within the spleens of these mice (Figure 8), again highlighting the tumour-specific nature of these effects. In Figure 8, each symbol represents an

individual tumour. The horizontal line shown in each treatment group represents the average frequency of tumour-associated CD4 T cells for the group of four mice. Spleens from each treatment group were combined for analysis. Each horizontal bar represents the average splenic associated CD4 T cell frequency for each treatment group.

[00104] NK cell frequency is low in the 4T1.2 tumours. Co-treatment with PG545 + anti-PD-1 significantly increased the frequency of tumour-associated $CD69^+$ NK cells compared to the single agent and control treatments (Figure 10, Figures 11A-11B). Notably this effect of the PG545 + anti-PD-1 treatment was not observed within the spleens of these mice. Therapy-associated changes in NK cell expression of CD335 expression were not detected (Figure 10, Figures 11A-11B). In Figure 10, each symbol represents an individual tumour. The horizontal line shown in each treatment group represents the average frequency of tumour-associated NK cells for the group of four mice. Spleens from each treatment group were combined for analysis. Each horizontal bar represents the average splenic associated NK cell frequency for each treatment group. In Figures 11A-11B, concatenated results are shown for each treatment group (four tumours/group).

[00105] Drug-associated changes within the 4T1.2 tumour-associated myeloid populations were only observed in the neutrophil cell subset ($Ly6C^{int}CD11b^+CD11^-Ly6G^+$ cells) (Figure 12; Figures 13A-13D (in which concatenated results are shown for each treatment group, four tumours/group)). Anti-PD-1 treatment was the primary mediator of the observed reduction in tumour-associated neutrophils (Figure 12). In Figure 12, each symbol represents an individual tumour and the horizontal line shown in each treatment group represents the average frequency of tumour-associated myeloid cells for the group of four mice.

[00106] Drug-associated changes in PD-L1 expression within the 4T1.2 tumour-associated myeloid compartment were not detected (Figure 14; Figures 13A-13D). In Figure 14, each symbol represents the mean fluorescence intensity (MFI) for PD-L1 expression within an individual tumour. The horizontal line shown in each treatment group represents the average MFI for PD-L1 expression.

Conclusions

[00107] The %TGI for the PG545 and anti-PD-1 group (84%) was almost two-fold higher compared with the vehicle + anti-PD1 antibody group (44%) and higher than the PG545 + control antibody group (68%).

[00108] Immune analysis revealed significant synergy of PG545 and anti-PD-1 in the

resected 4T1.2 mammary tumours. Significant increases in tumour-associated lymphocytes were observed, with the accumulation of Effector Memory CD8+ T cells, Central Memory CD8+ T cells, and CD4+ T cells in the 4T1.2 tumours.

[00109] Synergistic effects of PG545 and anti-PD-1 were also evident on the 4T1.2 tumour-associated NK cells. Significant increases in tumour associated NK cell frequency and CD69 expression within the NK cell compartment were observed in the PG545 + anti-PD-1 treated 4T1.2 tumours.

[00110] The PG545 and anti-PD1 antibody combination enhanced the TGI almost two-fold compared with anti-PD1 alone. Moreover, synergistic increases in tumour-associated CD8+ T, CD4+ T and NK cells were identified in the 4T1.2 tumours treated with both PG545 and anti-PD1 therapy.

[00111] In the present specification and claims (if any), the word 'comprising' and its derivatives including 'comprises' and 'comprise' include each of the stated integers but does not exclude the inclusion of one or more further integers.

[00112] Reference throughout this specification to 'one embodiment' or 'an embodiment' means that a particular feature, structure, or characteristic described in connection with the embodiment is included in at least one embodiment of the present invention. Thus, the appearance of the phrases 'in one embodiment' or 'in an embodiment' in various places throughout this specification are not necessarily all referring to the same embodiment. Furthermore, the particular features, structures, or characteristics may be combined in any suitable manner in one or more combinations.

[00113] In compliance with the statute, the invention has been described in language more or less specific to structural or methodical features. It is to be understood that the invention is not limited to specific features shown or described since the means herein described comprises preferred forms of putting the invention into effect. The invention is, therefore, claimed in any of its forms or modifications within the proper scope of the appended claims (if any) appropriately interpreted by those skilled in the art.

CITATION LIST

[00114] Boyango I, Barash U, Naroditsky I, Li JP, Hammond E, Ilan N, Vlodavsky I. (2014) Heparanase cooperates with Ras to drive breast and skin tumorigenesis. Cancer Res. 74(16): 4504-14.

[00115] Brennan TV, Lin L, Brandstadter JD, Rendell VR, Dredge K, Huang X, Yang Y. (2016) Heparan sulfate mimetic PG545-mediated antilymphoma effects require TLR9-dependent NK cell activation. J Clin Invest. 126(1): 207-19.

[00116] Chen YM. (2016) Immune checkpoint inhibitors for nonsmall cell lung cancer treatment. J Chin Med Assoc. doi: 10.1016/j.jcma.2016.08.005. [Epub ahead of print].

[00117] Dredge K, Hammond E, Davis K, Li CP, Liu L, Johnstone K, Handley P, Wimmer N, Gonda TJ, Gautam A, Ferro V, Bytheway I. (2010) The PG500 series: novel heparan sulfate mimetics as potent angiogenesis and heparanase inhibitors for cancer therapy. Invest New Drugs. 28(3): 276-83. doi: 10.1007/s10637-009-9245-5.

[00118] Ferro V, Liu L, Johnstone KD, Wimmer N, Karoli K, Handley P, Rowley J, Dredge K, Li CP, Hammond E, Davis K, Sarimaa L, Harenberg J, and Bytheway I (2012) Discovery of PG545: A Highly Potent and Simultaneous Inhibitor of Angiogenesis, Tumor Growth, and Metastasis. J. Med. Chem. 55(8): 3804-3813. doi: 10.1021/jm201708h.

[00119] Hammond E, Handley P, Dredge K, Bytheway I. (2013) Mechanisms of heparanase inhibition by the heparan sulfate mimetic PG545 and three structural analogues. FEBS Open Bio. 3: 346-51. doi: 10.1016/j.fob.2013.07.007.

[00120] Mennitto A, Grassi P, Ratta R, Verzoni E, Prisciandaro M, Procopio G. (2016) Nivolumab in the treatment of advanced renal cell carcinoma: clinical trial evidence and experience. Ther Adv Urol. 8(5): 319-326.

[00121] Ostapoff KT, Awasthi N, Cenik BK, Hinz S, Dredge K, Schwarz RE, Brekken RA. (2013) PG545, an angiogenesis and heparanase inhibitor, reduces primary tumor growth and metastasis in experimental pancreatic cancer. Mol Cancer Ther. 12(7): 1190-201.


[00122] Spranger S, Sivan A, Corrales L, Gajewski TF (2016) Tumor and Host Factors Controlling Antitumor Immunity and Efficacy of Cancer Immunotherapy. Adv Immunol. 130: 75-93.

[00123] U.S. Food and Drug Administration (May 17, 2016). "Nivolumab (Opdivo) for Hodgkin Lymphoma". U.S. Food and Drug Administration. U.S. Food and Drug Administration.

[00124] Woo SR, Corrales L, Gajewski TF (2015) The STING pathway and the T cell-inflamed tumor microenvironment. Trends Immunol. 36(4): 250-256.

CLAIMS

1. A method of treating or preventing cancer, comprising administering to a patient in need thereof an immune checkpoint modulator together with a compound of Formula (I):




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

2. The method of claim 1, wherein at least 90% of the X groups is SO_3M.

3. The method of claim 1 or claim 2, wherein the compound of Formula (I) is a compound of Formula (VI):



Formula (VI).

4. The method of any one of claims 1 to 3, wherein the immune checkpoint modulator is an immune checkpoint inhibitor.

5. The method of any one of claims 1 to 4, wherein the immune checkpoint modulator targets one or more of a T cell, an Antigen-Presenting Cell (APC) / tumour cell, a Natural Killer (NK) cell, a Natural Killer T cell (NKT), a Gamma Delta T cell, an

invariant T cell, or an invariant Natural Killer T cell (NKT).

6. The method of any one of claims 1 to 5, wherein the immune checkpoint modulator is a T cell immune checkpoint inhibitor.


7. The method of any one of claims 1 to 4, wherein the immune checkpoint modulator is a PD-1 inhibitor.

8. The method of any one of claims 1 to 3, wherein the immune checkpoint modulator is Nivolumab.

9. The method of any one of claims 1 to 8, wherein the cancer is a solid tumour.

10. Use of a compound of Formula (I) in the manufacture of a medicament for the treatment or prevention of cancer,

wherein the compound of Formula (I) is:




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M;

and wherein the medicament is formulated for administration with an immune checkpoint modulator.

11. Use of an immune checkpoint modulator in the manufacture of a medicament for the treatment or prevention of cancer,

wherein the medicament is formulated for administration with a compound of Formula (I):




Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

12. The use of claim 10 or claim 11, wherein at least 90% of the X groups is SO_3M.

13. The use of any one of claims 10 to 12, wherein the compound of Formula (I) is a compound of Formula (VI):



Formula (VI).

14. The use of any one of claims 10 to 13, wherein the immune checkpoint modulator is an immune checkpoint inhibitor.

15. The use of any one of claims 10 to 14, wherein the immune checkpoint modulator targets one or more of a T cell, an Antigen-Presenting Cell (APC) / tumour cell, a Natural Killer (NK) cell, a Natural Killer T cell (NKT), a Gamma Delta T cell, an invariant T cell, or an invariant Natural Killer T cell (NKT).

16. The use of any one of claims 10 to 15, wherein the immune checkpoint modulator is a T cell immune checkpoint inhibitor.


17. The use of any one of claims 10 to 14, wherein the immune checkpoint modulator is a

PD-1 inhibitor.

18. The use of any one of claims 10 to 13, wherein the immune checkpoint modulator is Nivolumab.

19. An immune checkpoint modulator and a compound of Formula (I) for use in the treatment or prevention of cancer;


wherein the compound of Formula (I) is:



Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.


20. A pharmaceutical composition or combination comprising a compound of Formula (I) and an immune checkpoint modulator; wherein the compound of Formula (I) is:



Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.

21. A kit comprising a compound of Formula (I) and an immune checkpoint modulator; wherein the compound of Formula (I) is:



Formula (I)

wherein X is SO_3M or H, wherein M is any pharmaceutically acceptable cation; and wherein at least 70% of the X groups is SO_3M.




Figure 1




Figure 2



Vehicle + control Ab
PG545 + control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
% mean weight change
20
10
0
-10
-20
0
2
4
6
8
10
12
Day
Tumour Volume (mm3)
400
300
200
100
0
0
2
4
6
8
10
12
14
Day

Figure 3



Group 1:
Vehicle + Clg
Group 2:
PG545 + Clg
Group 3:
Vehicle + Anti-PD-1
Group 4:
PG545 + Anti-PD-1
CD45.2
FSC
SSC
FSC

Figure 4



4T1.2 Tumour Analysis
*P<0.03
CD8 T cell Analysis
% of CD45.2+ Cells
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
CD44+CD62L- CD8+ T cells
Effector CD8 T cell memory
CD44+CD62L+ CD8+ T cells
Central CD8 T cell memory
Spleen Analysis
CD8 T cell Analysis

Figure 5



Group 1
Group 2
Group 3
Group 4
CD8
CD44
CD62L
CD69
Group 1: Vehicle + Control Ab
Group 2: PG545 + Control Ab
Group 3: Vehicle + anti-PD-1
Group 4: PG545 + anti-PD-1

Figure 6A




Group 1
Group 2
Group 3
Group 4
CD69
CD62L
Granzyme B
CD62L
CD69
Granzyme B

Figure 6B



Group 1: Vehicle + Control Ab
Group 2: PG545 + Control Ab
Group 3: Vehicle + anti-PD-1
Group 4: PG545 + anti-PD-1

Figure 7



*P<0.03
4T1.2 Tumour Analysis
Spleen Analysis
CD4 T cells (Total)
CD4 T cells (Total)
% of CD45.2+ Cells
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
4T1.2 Tumour Analysis
CD4 T cells
CD44+ CD62L-
CD4 T cells
CD44+CD62L+

Figure 8



Group 1
Group 2
Group 3
Group 4
CD44
CD62L
CD69
Group 1: Vehicle + Control Ab
Group 2: PG545 + Control Ab
Group 3: Vehicle + anti-PD-1
Group 4: PG545 + anti-PD-1

Figure 9



*P<0.03
4T1.2 Tumour Analysis
Spleen Analysis
NK Cell Analysis
NK Cell Analysis
% of CD45.2+ Cells
% of Viable Cells
NK cells: CD69 Analysis
NK cell CD69
MFI: CD69
NK cells: CD335 Analysis
NK Cell CD335
MFI: CD335
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1

Figure 10



Group 1: Vehicle + Control Ab
Group 2: PG545 + Control Ab
Group 3: Vehicle + anti-PD-1
Group 4: PG545 + anti-PD-1
Group 1
Group 2
Group 3
Group 4
CD49b
FSC

Figure 11A



Group 1: Vehicle + Control Ab
Group 2: PG545 + Control Ab
Group 3: Vehicle + anti-PD-1
Group 4: PG545 + anti-PD-1
Group 1
Group 2
Group 3
Group 4
CD27
CD11b
CD69
CD335

Figure 11B



4T1.2 Tumour Analysis
*P<0.03
Inflammatory Monocytes
% of Viable Cells
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
Neutrophil Analysis
Macrophages
Ly6Cint CD11b+ CD11c+ Ly6G+
Ly6Cint CD11b+ CD11c+ Ly6G-

Figure 12




Not defined
Granulocytes
Inflammatory Monocytes
Neutrophils
Macrophages
Ly6G
CD11b
CD11c
MHCII
Ly6G+
Ly6G-
PD-L1

Figure 13A



Labels are as shown in Figure 13A

Figure 13B




Figure 13C

Labels are as shown in Figure 13A




Figure 13D

Labels are as shown in Figure 13A



4T1.2 Tumour Analysis
Inflammatory monocytes PDL1
MFI:PD-L1
6000
4000
2000
0
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
Neutrophils PD-L1
MFI:PD-L1
4000
3000
2000
1000
0
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
Macrophages PDL1
MFI:PD-L1
6000
4000
2000
0
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
CD11c+Ly6G- PD-L1
MFI:PD-L1
15000
10000
5000
0
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1
CD11c+Ly6G+ PD-L1
MFI:PD-L1
3000
2000
1000
0
Vehicle + Control Ab
PG545 + Control Ab
Vehicle + anti-PD-1
PG545 + anti-PD-1

Figure 14