

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2002248184 C1**

(54) Title
Molecules with extended half-lives, compositions and uses thereof

(51) International Patent Classification(s)
C07K 14/705 (2006.01) **C07K 19/00** (2006.01)
A61K 39/395 (2006.01) **C12N 1/15** (2006.01)
A61K 49/00 (2006.01) **C12N 1/19** (2006.01)
A61P 31/16 (2006.01) **C12N 1/21** (2006.01)
A61P 37/00 (2006.01) **C12N 5/10** (2006.01)
C07K 16/00 (2006.01) **C12N 15/09** (2006.01)
C07K 16/08 (2006.01) **G01N 33/534** (2006.01)
C07K 16/18 (2006.01) **A61K 38/00** (2006.01)
C07K 16/46 (2006.01) **A61K 39/00** (2006.01)

(21) Application No: **2002248184** (22) Date of Filing: **2001.12.12**

(87) WIPO No: **WO02/060919**

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/289,760	2001.05.09	US
60/254,884	2000.12.12	US

(43) Publication Date: **2002.08.12**
(43) Publication Journal Date: **2003.02.20**
(44) Accepted Journal Date: **2008.01.10**
(44) Amended Journal Date: **2018.01.04**

(71) Applicant(s)
MedImmune, LLC; Board of Regents, The University of Texas System

(72) Inventor(s)
Ward, Elizabeth Sally; Johnson, Leslie S.; Dall'acqua, William

(74) Agent / Attorney
Phillips Ormonde Fitzpatrick, L 16 333 Collins St, Melbourne, VIC, 3000, AU

(56) Related Art
WO1998023289
WO1999043713
WO1997034631

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
8 August 2002 (08.08.2002)

PCT

(10) International Publication Number
WO 02/060919 A2(51) International Patent Classification⁷: C07K

(21) International Application Number: PCT/US01/48432

(22) International Filing Date:
12 December 2001 (12.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/254,884 12 December 2000 (12.12.2000) US
60/289,760 9 May 2001 (09.05.2001) US(71) Applicant (for all designated States except US): MED-
IMMUNE, INC. [US/US]; 35 West Watkins Mill Road,
Gaithersburg, MD 20878 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DALL'ACQUA,
William [FR/US]; 9926 Shelburne Terrace #208, Gaithers-
burg, MD 20878 (US). JOHNSON, Leslie, S. [US/US];
20147 Laurel Hill Way, Germantown, MD 20874 (US).
WARD, Elizabeth, Sally [GB/US]; Apartment 2017, 4606
Cedar Springs, Dallas, TX 75230 (US).(74) Agents: POISSANT, Brian, M. et al.; Pennie & Edmonds
LLP, 1155 Avenue of the Americas, New York, NY 10036
(US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).**Published:**— without international search report and to be republished
upon receipt of that report(48) Date of publication of this corrected version:
3 January 2003(15) Information about Correction:
see PCT Gazette No. 01/2003 of 3 January 2003, Section IIFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: MOLECULES WITH EXTENDED HALF-LIVES, COMPOSITIONS AND USES THEREOF

(57) Abstract: The present invention provides molecules, including IgGs, non-IgG immunoglobulins, proteins and non-protein agents, that have increased *in vivo* half-lives due to the presence of an IgG constant domain, or a portion thereof that binds the FcRn, having one or more amino acid modifications that increase the affinity of the constant domain or fragment for FcRn. Such proteins and molecules with increased half-lives have the advantage that smaller amounts and or less frequent dosing is required in the therapeutic, prophylactic or diagnostic use of such molecules.

WO 02/060919 A2

5

MOLECULES WITH EXTENDED HALF-LIVES, COMPOSITIONS AND USES THEREOF

This application claims the benefit of United States provisional application Serial Nos. 60/254,884, filed December 12, 2000, and 60/289,760, filed May 9, 2001, both
10 of which are incorporated by reference herein in their entireties. This invention was made, in part, with United States Government support under award number A139167 from the National Institute of Health. The United States Government may have certain rights in the invention.

15

1. INTRODUCTION

The present invention relates to molecules whose *in vivo* half-lives are increased by modification of an IgG constant domain, or FcRn (Fc Receptor-neonate) binding domain thereof. Specifically, these molecules have amino acid modifications that increase the affinity of the constant domain or fragment thereof for the FcRn. Increasing the
20 half-life of therapeutic and diagnostic IgGs and other bioactive molecules using methods of the invention has many benefits including reducing the amount and/or frequency of dosing of these molecules, for example, in vaccines, passive immunotherapy and other therapeutic and prophylactic methods. The invention further relates to fusion proteins containing all or a portion (a FcRn binding portion) of an IgG constant domain having one or more of these
25 amino acid modifications and a non-IgG protein or non-protein molecule conjugated to such a modified IgG constant domain, where the presence of the modified IgG constant domain increases the *in vivo* half-life of the non-IgG protein or molecule.

2. BACKGROUND OF THE INVENTION

30 The use of immunoglobulins as therapeutic agents has increased dramatically in recent years and have expanded to different areas of medical treatments. Such uses include treatment of agammaglobulinemia and hypogammaglobulinemia, as immunosuppressive agents for treating autoimmune diseases and graft-vs.-host (GVH) diseases, the treatment of lymphoid malignancies, and passive immunotherapies for the
35 treatment of various systemic and infectious diseases. Also, immunoglobulins are useful as *in vivo* diagnostic tools, for example, in diagnostic imaging procedures.

One critical issue in these therapies is the persistence of immunoglobulins in the circulation. The rate of immunoglobulin clearance directly affects the amount and frequency of dosage of the immunoglobulin. Increased dosage and frequency of dosage
5 may cause adverse effects in the patient and also increase medical costs.

IgG is the most prevalent immunoglobulin class in humans and other mammals and is utilized in various types of immunotherapies and diagnostic procedures. The mechanism of IgG catabolism in the circulation has been elucidated through studies related to the transfer of passive immunity from mother to fetus/neonate through the
10 placenta or yolk sac or through colostrum (maternofetal transfer of IgG via transcytosis) in rodents (Brambell, *Lancet*, ii:1087-1093, 1966; Rodewald, *J. Cell Biol.*, 71:666-670, 1976; Morris *et al.*, In: *Antigen Absorption by the Gut*, pp. 3-22, 1978, University Park Press, Baltimore; Jones *et al.*, *J. Clin. Invest.*, 51:2916-2927, 1972).

The involvement of certain receptors in the maternofetal transmission of
15 maternal IgGs was first suggested by Brambell's group in their study on the intestinal absorption of maternal antibodies from ingested milk in newborn rats (Halliday, *Proc. R. Soc. B.*, 143:408-413, 1955; Halliday, *Proc. R. Soc. B.*, 144:427-430, 1955; Halliday, *Proc. R. Soc. B.*, 148:92-103, 1957; Morris, *Proc. R. Soc. B.*, 148:84-91, 1957; Brambell *et al.*, *Proc. R. Soc. B.*, 149:1-11, 1958; Morris, *Proc. R. Soc. B.*, 160:276-292, 1964). Brambell
20 *et al.* suggested, based on the observation that heterologous IgGs interfered with the transmission of a specific antibody, that IgG molecules from various species might have sufficiently similar structures or sequences that bind to common receptors (Brambell *et al.*, *Proc. R. Soc. B.*, 149:1-11, 1958).

A high-affinity Fc receptor, FcRn, has been implicated in this transfer
25 mechanism. The FcRn receptor has been isolated from duodenal epithelial brush borders of suckling rats (Rodewald *et al.*, *J. Cell Biol.*, 99:154s-164s, 1984; Simister *et al.*, *Eur. J. Immunol.*, 15:733-738, 1985) and the corresponding gene has been cloned (Simister *et al.*, *Nature*, 337:184, 1989 and *Cold Spring Harbor Symp. Quant. Biol.*, LIV, 571-580, 1989). The later clonings of FcRn-encoding genes from mice (Ahouse *et al.*, *J. Immunol.*,
30 151:6076-6088, 1993) and humans (Story *et al.*, *J. Exp. Med.*, 180:2377-2381, 1994) demonstrate high homology of these sequences to the rat FcRn, suggesting a similar mechanism of maternofetal transmission of IgGs involving FcRn in these species.

Meanwhile, a mechanism for IgG catabolism was also proposed by Brambell's group (Brambell *et al.*, *Nature*, 203:1352-1355, 1964; Brambell, *Lancet*,
35 ii:1087-1093, 1966). They proposed that a proportion of IgG molecules in the circulation are bound by certain cellular receptors (*i.e.*, FcRn), which are saturable, whereby the IgGs

are protected from degradation and eventually recycled into the circulation; on the other hand, IgGs which are not bound by the receptors are degraded. The proposed mechanism was consistent with the IgG catabolism observed in hypergammaglobulinemic or hypogammaglobulinemic patients. Furthermore, based on his studies as well as others (see, e.g., Spiegelberg *et al.*, *J. Exp. Med.*, 121:323-338, 1965; Edelman *et al.*, *Proc. Natl. Acad. Sci. USA*, 63:78-85, 1969), Brambell also suggested that the mechanisms involved in maternofetal transfer of IgG and catabolism of IgG may be either the same or, at least, very closely related (Brambell, *Lancet*, ii:1087-1093, 1966). Indeed, it was later reported that a mutation in the Fc-hinge fragment caused concomitant changes in catabolism, maternofetal transfer, neonatal transcytosis, and, particularly, binding to FcRn (Ghetie *et al.*, *Immunology Today*, 18(12):592-598, 1997).

These observations suggested that portions of the IgG constant domain control IgG metabolism, including the rate of IgG degradation in the serum through interactions with FcRn. Indeed, increased binding affinity for FcRn increased the serum half-life of the molecule (Kim *et al.*, *Eur. J. Immunol.*, 24:2429-2434, 1994; Popov *et al.*, *Mol. Immunol.*, 33:493-502, 1996; Ghetie *et al.*, *Eur. J. Immunol.*, 26:690-696, 1996; Junghans *et al.*, *Proc. Natl. Acad. Sci. USA*, 93:5512-5516, 1996; Israel *et al.*, *Immunol.*, 89:573-578, 1996).

Various site-specific mutagenesis experiments in the Fc region of mouse IgGs have led to identification of certain critical amino acid residues involved in the interaction between IgG and FcRn (Kim *et al.*, *Eur. J. Immunol.*, 24:2429-2434, 1994; Medesan *et al.*, *Eur. J. Immunol.*, 26:2533, 1996; Medesan *et al.*, *J. Immunol.*, 158:2211-2217, 1997). These studies and sequence comparison studies found that isoleucine at position 253, histidine at position 310, and histidine at position 435 (according to Kabat numbering, Kabat *et al.*, In: *Sequences of Proteins of Immunological Interest*, US Department of Health and Human Services, 1991, which is hereby incorporated by reference in its entirety), are highly conserved in human and rodent IgGs, suggesting their importance in IgG-FcRn binding.

Additionally, various publications describe methods for obtaining physiologically active molecules whose half-lives are modified either by introducing an FcRn-binding polypeptide into the molecules (WO 97/43316; U.S. Patent No. 5,869,046; U.S. Patent No. 5,747,035; WO 96/32478; WO 91/14438) or by fusing the molecules with antibodies whose FcRn-binding affinities are preserved but affinities for other Fc receptors have been greatly reduced (WO 99/43713) or fusing with FcRn binding domains of

antibodies (WO 00/09560; U.S. Patent No. 4,703,039). However, none of these publications disclose specific mutants in the IgG constant domain that affect half-life.

Prior studies have demonstrated that certain constant domain mutations actually reduce binding to FcRn and, thereby, reduce the IgG *in vivo* half-life. PCT publication WO 93/22332 (by Ward *et al.*) discloses various recombinant mouse IgGs whose *in vivo* half-lives are reduced by mutations between about residue 253 and about residue 434. Particularly, substitutions of isoleucine at position 253; histidine at position 310; glutamine at position 311; His at position 433; and asparagine at position 434 were found to reduce IgG half-life.

Modulation of IgG molecules by amino acid substitution, addition, or deletion to increase or reduce affinity for FcRn is also disclosed in WO 98/23289; however, the publication does not list any specific mutants that exhibit either longer or shorter *in vivo* half-lives.

In fact, only one mutant of mouse IgG1 that actually exhibited increased half-life, the triple mutation Thr252 to Ala, Thr254 to Ser, and Thr256 to Phe, has been identified (WO 97/34631).

In view of the pharmaceutical importance of increasing the *in vivo* half-lives of immunoglobulins and other bioactive molecules, there is a need to develop modified IgGs and FcRn-binding fragments thereof, (particularly modified human IgGs) that confer increased *in vivo* half-life on immunoglobulins and other bioactive molecules.

3. SUMMARY OF THE INVENTION

The present invention is based upon the inventors' identification of several mutations in the constant domain of a human IgG molecule that increase the affinity of the IgG molecule for the FcRn. In particular, the present inventors have screened libraries of human IgG1 constant domains with random amino acid mutations introduced into particular regions of the constant domain for increased affinity for FcRn. Such random mutations were made in the regions of residues 251-256, 285-290, and 308-314, all of which are in CH2 domain, and 385-389 and 428-436, which are in CH3 domain, of human IgG1 hinge-Fc regions (residues as depicted in Figure 2 (SEQ ID NO:83 or analogous residues in hinge-Fc regions of other IgG molecules as determined by sequence alignment). As used herein, all residues of the IgG constant domain are numbered according to Kabat *et al.* (Sequences of Proteins of Immunological Interest, U.S. Department of Health and Human Services, 1991, which is incorporated by reference herein in its entirety) and as presented in Figure 2 (SEQ ID NO:83), and include corresponding residues in other IgG constant domains as

determined by sequence alignment. The *in vivo* half-life, or persistence in serum or other tissues of a subject, of antibodies, and other therapeutic agents and other bioactive molecules is an important clinical parameter which determines the amount and frequency of antibody (or any other pharmaceutical molecule) administration. Accordingly, such molecules, including antibodies, with increased half-life are of significant pharmaceutical importance.

Thus, the present invention relates to a modified molecule (preferably a protein, but may be a non-protein agent) that has an increased *in vivo* half-life by virtue of the presence of a modified IgG constant domain, or FcRn-binding portion thereof (preferably the Fc or hinge-Fc domain) (preferably from a human IgG) wherein the IgG constant domain, or fragment thereof, is modified (*e.g.*, by amino acid substitution, deletion or insertion) to increase the affinity for the FcRn. In a particular embodiment, the present invention relates to modified IgGs, whose *in vivo* half-lives are extended by the modification of amino acid residues identified to be involved in the interaction of the hinge-Fc domain with the FcRn receptor. Preferably, the constant domain or fragment thereof has higher affinity for FcRn at pH 6.0 than at pH 7.4. Such modifications may also alter (*i.e.*, increase or decrease) the bioavailability (*e.g.*, transport to mucosal surfaces, or other target tissues) of the molecules. The invention also relates to other types of immunoglobulins or fragments thereof (*i.e.*, non-IgG immunoglobulins), non-immunoglobulin proteins and non-protein agents that are fused or conjugated to, or engineered to contain, an IgG constant domain, or FcRn-binding fragment thereof, having one or more such amino acid modifications.

In preferred embodiments, the present invention provides molecules, particularly, immunoglobulins whose *in vivo* half-lives are extended by the presence of an IgG constant domain, or FcRn binding fragment thereof (preferably, Fc or hinge-Fc domain), that has modifications of one or more of amino acid residues 251-256, 285-290, 308-314, 385-389, and 428-436 that increase the affinity of the constant domains or fragments thereof for FcRn. In certain embodiments, these modifications preferably exclude residues 252, 254, and 256, in particular when the IgG constant domain or fragment thereof, is murine. In particular embodiments, the modification is at one or more surface-exposed residues, and the modification is a substitution with a residue of similar charge, polarity or hydrophobicity to the residue being substituted. In preferred embodiments, the modified IgG constant domain, or fragment thereof, binds with higher affinity to FcRn at pH 6.0 than at pH 7.4. In a preferred embodiment, the constant domain, or fragment thereof, is modified by substitution of one or more of amino acid residues 251-256, 285-

290, 308-314, 385-389, and 428-436 that increase the affinity of the constant domain or FcRn-binding fragments thereof for FcRn. In certain embodiments, substitutions of residue 252 with leucine, residue 254 with serine, and/or residue 256 with phenylalanine are
5 excluded, particularly when the constant domain or fragment thereof is derived from a mouse IgG.

In specific embodiments, the invention provides immunoglobulins or other bioactive molecules that contain an IgG1 constant domain, or FcRn-binding fragment thereof (preferably Fc or hinge-Fc domain) (preferably human), having amino acid
10 modifications at one or more of position 308, 309, 311, 312, and 314, more specifically, having substitutions at one or more of positions 308, 309, 311, 312 and 314 with threonine, proline, serine, aspartic acid and leucine respectively. In another embodiment, residues at one or more of positions 308, 309, and 311 are substituted with isoleucine, proline, and glutamic acid, respectively. In yet another embodiment, residues at one or more of positions
15 308, 309, 311, 312, and 314, are substituted with threonine, proline, serine, aspartic acid, and leucine, respectively. The invention further relates to combinations of these amino acid substitutions.

Furthermore, the invention provides immunoglobulins or other bioactive molecules that contain an IgG1 constant domain, or FcRn-binding fragment thereof
20 (preferably, Fc or hinge-Fc domain) (preferably human), having amino acid modifications at one or more of positions 251, 252, 254, 255, and 256, more specifically, having substitutions at one or more of these positions. In specific embodiments, residue 251 is substituted with leucine or arginine, residue 252 is substituted with tyrosine, phenylalanine, serine, tryptophan or threonine, residue 254 is substituted with threonine or serine, residue
25 255 is substituted with leucine, glycine, isoleucine or arginine, and/or residue 256 is substituted with serine, arginine, glutamine, glutamic acid, aspartic acid, alanine, asparagine or threonine. In a more specific embodiment, residue 251 is substituted with leucine, residue 252 is substituted with tyrosine, residue 254 is substituted with threonine or serine, and/or residue 255 is substituted with arginine. In yet another specific embodiment, residue
30 252 is substituted with phenylalanine and/or residue 256 is substituted with aspartic acid. In a preferred embodiment, residue 251 is substituted with leucine, residue 252 is substituted with tyrosine, residue 254 is substituted with threonine or serine, and/or residue 255 is substituted with arginine. The invention further relates to any combination of these substitutions.

35 Furthermore, the invention provides immunoglobulins or other bioactive molecules that contain an IgG1 constant domain, or FcRn-binding fragment thereof

(preferably, Fc or hinge-Fc domain) (preferably human), having amino acid modifications at one or more of positions 251, 252, 254, 255, and 256, more specifically, having substitutions at one or more of these positions. In specific embodiments, residue 251 is substituted with leucine or arginine, residue 252 is substituted with tyrosine, phenylalanine, serine, tryptophan or threonine, residue 254 is substituted with threonine or serine, residue 255 is substituted with leucine, glycine, isoleucine or arginine, and/or residue 256 is substituted with serine, arginine, glutamine, glutamic acid, aspartic acid, alanine, asparagine or threonine. In a more specific embodiment, residue 251 is substituted with leucine, residue 252 is substituted with tyrosine, residue 254 is substituted with threonine or serine, and/or residue 255 is substituted with arginine. In yet another specific embodiment, residue 252 is substituted with phenylalanine and/or residue 256 is substituted with aspartic acid. In a preferred embodiment, residue 251 is substituted with leucine, residue 252 is substituted with tyrosine, residue 254 is substituted with threonine or serine, and/or residue 255 is substituted with arginine. The invention further relates to any combination of these substitutions.

Furthermore, the invention provides immunoglobulins or other bioactive molecules that contain an IgG1 constant domain, or FcRn-binding fragment thereof (preferably, Fc or hinge-Fc domain) (preferably human), having amino acid modifications at one or more of positions 428, 433, 434, and 436, more specifically, having substitutions at one or more of these positions. In specific embodiments, residue 428 is substituted with methionine, threonine, leucine, phenylalanine, or serine, residue 433 is substituted with lysine, arginine, serine, isoleucine, proline, glutamine, or histidine, residue 434 is substituted with phenylalanine, tyrosine, or histidine, and/or residue 436 is substituted with histidine, asparagine, arginine, threonine, lysine, methionine, or threonine. In a more specific embodiment, residues at one or more positions 433, 434, and 436 are substituted with lysine, phenylalanine, and histidine, respectively. In a preferred embodiment, residue 428 is substituted with methionine and/or residue 434 is substituted with tyrosine.

Furthermore, the invention provides immunoglobulins or other bioactive molecules that contain an IgG1 constant domain, or FcRn-binding fragment thereof (preferably, Fc or hinge-Fc domain) (preferably human), having amino acid modifications at one or more positions 385, 386, 387 and 389, more specifically, having substitutions at one or

7a

more of these positions. In specific embodiments, residue 385 is substituted with arginine, aspartic acid, serine, threonine, histidine, lysine, or alanine, residue 386 is substituted with threonine, proline, aspartic acid, serine, lysine, arginine, isoleucine, or methionine, residue 387 is substituted with arginine, histidine, serine, threonine, alanine, or proline and/or residue 389 is substituted with proline or serine. In more specific embodiments, residues at one or more positions 385, 386, 387, and 389 are substituted with arginine, threonine, arginine, and proline, respectively. In yet another specific embodiment, residues at one or more positions 385, 386, and 389 are substituted with aspartic acid, proline, and serine, respectively.

Molecules of the invention include any combination of the above-described substitutions at one or more of residues 251, 252, 254, 255, 256, 308, 309, 311, 312, 385, 386, 387, 389, 428, 433, 434 and/or 436. In a preferred embodiment, the molecule of the invention contains a Fc region, or FcRn-binding domain thereof, having one or more of the following substitutions: leucine at residue 251, tyrosine at residue 252, threonine or serine at residue 254, arginine at residue 255, threonine at residue 308, proline at residue 309, serine at residue 311, aspartic acid at residue 312, leucine at residue 314, arginine at residue 385, threonine at residue 386, arginine at residue 387, proline at residue 389, methionine at residue 428, and/or tyrosine at residue 434.

Thus, in one embodiment, the present invention provides a modified IgG comprising an IgG constant domain, or an FcRn-binding fragment thereof, wherein the constant domain or fragment comprises a substitution at amino acid residue 252 relative to a wild-type IgG constant domain, numbered according to the EU numbering index of Kabat, wherein the modified IgG has an increased half-life compared to the half-life of an IgG having the wild-type IgG constant domain, and wherein the substitution at amino acid residue 252 is a substitution with tyrosine, phenylalanine, tryptophan, or threonine.

Included within the invention are pharmaceutical compositions and methods of prophylaxis and therapy using modified immunoglobulins, proteins and other bioactive molecules of the invention having extended half-lives. Also included are methods of

30

diagnosis using modified immunoglobulins, proteins and other bioactive molecules of the invention having extended half-lives. In a specific embodiment, the invention provides an anti-respiratory syncytial virus (RSV) antibody useful to treat or prevent RSV infection, such as SYNAGIS® (see U.S. Patent No. 5,824,307 and Johnson *et al.*, *J. Infectious Disease* 176:1215-1224, 1997, both of which are incorporated by reference in their entireties), and other anti-RSV antibodies, including variants of SYNAGIS® (see United States patent Application Serial No., 09/724,396, filed November 28, 2000, United States patent Application Serial No. 09/724,531, filed November 28, 2000, United States patent Application Serial No. ___, filed November 28, 2001 (attorney docket no. 10271-047), and United States patent Application Serial No. ___, filed November 28, 2001 (attorney docket no. 10271-048), all entitled "Methods of Administering/Dosing Anti-RSV Antibodies for Prophylaxis and Treatment," all by Young *et al.*, all of which are incorporated by reference herein in their entireties, particularly the sequences of heavy and light chain variable domains and CDRs of anti-RSV antibodies disclosed therein), which has one or more amino acid modifications in the constant domain that increase the affinity of the antibody for FcRn and that has an increased *in vivo* half-life (see also, Section 5.1 *infra*).

3.1 DEFINITIONS

The term "IgG Fc region" as used herein refers the portion of an IgG molecule that correlates to a crystallizable fragment obtained by papain digestion of an IgG molecule. The Fc region consists of the C-terminal half of the two heavy chains of an IgG molecule that are linked by disulfide bonds. It has no antigen binding activity but contains the carbohydrate moiety and the binding sites for complement and Fc receptors, including the FcRn receptor (see below). The Fc fragment contains the entire second constant domain CH2 (residues 231-340 of human IgG1, according to the Kabat numbering system) (*e.g.*, SEQ ID NO:80) and the third constant domain CH3 (residues 341-447) (*e.g.*, SEQ ID NO:81).

The term "IgG hinge-Fc region" or "hinge-Fc fragment" as used herein refers to a region of an IgG molecule consisting of the Fc region (residues 231-447) and a hinge region (residues 216-230; *e.g.*, SEQ ID NO:82) extending from the N-terminus of the Fc region. An example of the amino acid sequence of the human IgG1 hinge-Fc region is SEQ ID NO:83.

The term "constant domain" refers to the portion of an immunoglobulin molecule having a more conserved amino acid sequence relative to the other portion of the immunoglobulin, the variable domain, which contains the antigen binding site. The

constant domain contains the CH1, CH2 and CH3 domains of the heavy chain and the CHL domain of the light chain.

The term "FcRn receptor" or "FcRn" as used herein refers to an Fc receptor ("n" indicates neonatal) which is known to be involved in transfer of maternal IgGs to a fetus through the human or primate placenta, or yolk sac (rabbits) and to a neonate from the colostrum through the small intestine. It is also known that FcRn is involved in the maintenance of constant serum IgG levels by binding the IgG molecules and recycling them into the serum. The binding of FcRn to IgG molecules is strictly pH-dependent with optimum binding at pH 6.0. FcRn comprises a heterodimer of two polypeptides, whose molecular weights are approximately 50 kD and 15 kD, respectively. The extracellular domains of the 50 kD polypeptide are related to major histocompatibility complex (MHC) class I α -chains and the 15 kD polypeptide was shown to be the non-polymorphic β_2 -microglobulin (β_2 -m). In addition to placenta and neonatal intestine, FcRn is also expressed in various tissues across species as well as various types of endothelial cell lines. It is also expressed in human adult vascular endothelium, muscle vasculature and hepatic sinusoids and it is suggested that the endothelial cells may be most responsible for the maintenance of serum IgG levels in humans and mice. The amino acid sequences of human FcRn and murine FcRn are indicated by SEQ ID NO:84 and SEQ ID NO:85, respectively. Homologs of these sequences having FcRn activity are also included.

The term "*in vivo* half-life" as used herein refers to a biological half-life of a particular type of IgG molecule or its fragments containing FcRn-binding sites in the circulation of a given animal and is represented by a time required for half the quantity administered in the animal to be cleared from the circulation and/or other tissues in the animal. When a clearance curve of a given IgG is constructed as a function of time, the curve is usually biphasic with a rapid α -phase which represents an equilibration of the injected IgG molecules between the intra- and extra-vascular space and which is, in part, determined by the size of molecules, and a longer β -phase which represents the catabolism of the IgG molecules in the intravascular space. The term "*in vivo* half-life" practically corresponds to the half life of the IgG molecules in the β -phase.

An "isolated" or "purified" antibody or fusion protein is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or substantially free of chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of an antibody or a fusion protein in which the antibody or the fusion protein is separated from cellular components of the cells from which it is isolated or recombinantly

produced. Thus, an antibody or a fusion protein that is substantially free of cellular material includes preparations of antibody or fusion protein having less than about 30%, 20%, 10%, or 5% (by dry weight) of contaminating protein. When the antibody or the fusion protein is recombiantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, 10%, or 5% of the volume of the protein preparation. When the antibody or the fusion protein is produced by chemical synthesis, it is preferably substantially free of chemical precursors or other chemicals, *i.e.*, it is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. Accordingly such preparations of the antibody or the fusion protein have less than about 30%, 20%, 10%, 5% (by dry weight) of chemical precursors or compounds other than the antibody or antibody fragment of interest. In a preferred embodiment of the present invention, antibodies are isolated or purified. In another preferred embodiment of the invention, fusion proteins are isolated or purified.

An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid molecule. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. An "isolated" nucleic acid molecule does not include cDNA molecules within a cDNA library. In a preferred embodiment of the invention, nucleic acid molecules encoding antibodies are isolated or purified. In another preferred embodiment of the invention, nucleic acid molecules encoding fusion proteins are isolated or purified.

The term "host cell" as used herein refers to the particular subject cell transfected with a nucleic acid molecule or infected with phagemid or bacteriophage and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.

The names of amino acids referred to herein are abbreviated either with three-letter or one-letter symbols.

To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino acid or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are

then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity = number of identical overlapping positions/total number of positions x 100%). In one embodiment, the two sequences are the same length.

The determination of percent identity between two sequences can also be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. U.S.A.* 87:2264-2268, modified as in Karlin and Altschul, 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:5873-5877. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul *et al.*, 1990, *J. Mol. Biol.* 215:403. BLAST nucleotide searches can be performed with the NBLAST nucleotide program parameters set, *e.g.*, for score=100, wordlength=12 to obtain nucleotide sequences homologous to a nucleic acid molecules of the present invention. BLAST protein searches can be performed with the XBLAST program parameters set, *e.g.*, to score=50, wordlength=3 to obtain amino acid sequences homologous to a protein molecule of the present invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, 1997, *Nucleic Acids Res.* 25:3389-3402. Alternatively, PSI-BLAST can be used to perform an iterated search which detects distant relationships between molecules (*Id.*). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs (*e.g.*, of XBLAST and NBLAST) can be used (see, *e.g.*, <http://www.ncbi.nlm.nih.gov>). Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, 1988, *CABIOS* 4:11-17. Such an algorithm is incorporated in the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

The percent identity between two sequences can be determined using techniques similar to those described above, with or without allowing gaps. In calculating percent identity, typically only exact matches are counted.

4. DESCRIPTION OF THE FIGURES

FIG. 1 shows the structure of the IgG hinge-Fc region indicating the locations of the residues identified to be involved in the interaction with the FcRn receptor (Ghetie *et al.*, *Immunology Today*, 18(12):592-598, 1997).

FIG. 2 shows the amino acid sequence of the human IgG1 hinge-Fc region (SEQ ID NO:83) containing a hinge region (SEQ ID NO:82), CH2 domain (SEQ ID NO:80), and CH3 domain (SEQ ID NO:81).

FIGS. 3 (A and B) show the amino acid sequences of (A) human FcRn (SEQ ID NO:84) and (B) mouse FcRn (SEQ ID NO:85), respectively.

FIG. 4 shows the amino acid sequence of the human IgG1 hinge-Fc region (SEQ ID NO:83), in which wild-type residues which are mutated by amino acid substitutions are indicated in underlined bold-face.

FIG. 5 shows a schematic diagram of panning process for the phage-displayed modified hinge-Fc library.

FIG. 6 shows a summary of the occurrence of selected mutant residues at the variant positions in the libraries screened.

FIGS. 7 (A-D). (A) shows the binding of murine FcRn to immobilized IgG1 having M252Y/S254T/T256E substitutions. Murine FcRn was injected at 10 different concentrations ranging from 1 nM to 556 nM over a surface on which 4000 resonance units (RU) of IgG1 had been coupled. After equilibrium was reached, residual bound protein was eluted with a pulse of PBS, pH 7.4. (B) shows the binding of human FcRn to immobilized IgG1/M252Y/S254T/T256E. Murine FcRn was injected at 8 different concentrations ranging from 71 nM to 2.86 μ M over a surface on which 1000 RU of IgG1 had been coupled. After equilibrium was reached, residual bound protein was eluted with a pulse of PBS, pH 7.4. (C) and (D) show scatchard analyses of the data in (A) and (B), respectively, after correction for nonspecific binding. R_{eq} is the corrected equilibrium response at a given concentration C. The plots are linear with correlation coefficients of 0.97 and 0.998, respectively. The apparent K_d are 24 nM and 225 nM, respectively.

FIGS. 8 (A-H). (A)-(D) show the results from BIAcore analysis of the binding of murine FcRn at pH 6.0 and pH 7.4 to (A) wild type human IgG1, (B) M252Y/S254T/T256E, (C) H433K/N434F/Y436H, and (D) G385D/G386P/N389S, respectively, after correction for nonspecific binding. Murine FcRn was injected at a concentration of 1.1 μ M over a surface on which 1000 RU of wild type IgG1, 1000 RU of M252Y/S254T/T256E, 955 RU of H433K/N434F/Y436H, and 939 RU of G385D/Q386P/N389S had been coupled. (E)-(H) show the results from BIAcore analysis

of the binding of human FcRn at pH 6.0 and pH 7.4 to (E) wild type human IgG1, (F) M252Y/S254T/T256E, (G) H433K/N434F/Y436H, and (H) G385D/Q386P/N389S, respectively, after correction for nonspecific binding. Human FcRn was injected at a
 5 concentration of 1.4 μ m over a surface on which 1000 RU of wild type IgG1, 1000 RU of M252Y/S254T/T256E, 955 RU of H433K/N434F/Y436H, and 939 RU of G385D/Q386P/N389S had been coupled.

FIG. 9 shows the space-filling model of the surface of the Fc fragment of a human IgG1 based upon the human IgG1 structure of Deisenhofer, 1981, *Biochemistry*
 10 20:2361-2370. Residues are color-coded according to the gain of free energy of stabilization of the Fc-FcRn complex: red, substitutions at these positions were found to increase affinity by a factor of at least 2.5 times in the Fc/human FcRn interaction and of at least 5 times in the Fc/mouse FcRn interaction; blue, substitutions at those positions were found to increase affinity by a factor of less than 2 times in both the Fc-human FcRn and
 15 Fc-mouse FcRn interaction. The figure was drawn using Swiss pdb viewer (Guex and Peitsch, 1997, *Electrophoresis* 18:2714-2723).

FIG. 10 shows the changes in serum concentration ([Mab] ng/ml) over time (in days) of antibody having a wild type constant domain (SYNAGIS[®]) (open squares), or constant domains with the following mutations: M252Y/S254T/T256E (open circles),
 20 G385D/Q386P/N389S (solid squares), and H433K/N434F/Y436H (solid circles). Antibody concentration was determined using anti-human IgG ELISA.

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to molecules, particularly proteins, more
 25 particularly immunoglobulins, that have an increased *in vivo* half-life and comprise an IgG constant domain, or fragment thereof that binds to an FcRn (preferably a Fc or hinge-Fc domain), that contains one or more amino acid modifications relative to a wild type IgG constant domain which modifications increase the affinity of the IgG constant domain, or fragment thereof, for the FcRn. In a preferred embodiment, the invention particularly
 30 relates to the modification of human or humanized IgGs and other bioactive molecules containing FcRn-binding portions of human IgGs, which have particular use in human therapy, prophylaxis and diagnosis.

5.1 MOLECULES WITH INCREASED *IN VIVO* HALF-LIVES

The present invention is based upon identification of amino acid modifications in particular portions of the IgG constant domain that interact with the FcRn, which modifications increase the affinity of the IgG, or fragment thereof, for the FcRn. Accordingly, the invention relates to molecules, preferably proteins, more preferably immunoglobulins, that comprise an IgG constant domain, or FcRn binding fragment thereof (preferably a Fc or hinge-Fc domain fragment), having one or more amino acid modifications (*i.e.*, substitutions, insertions or deletions) in one or more regions that interact with the FcRn, which modifications increase the affinity of the IgG or fragment thereof, for the FcRn, and also increase the *in vivo* half-life of the molecule. In preferred embodiments, the one or more amino acid modifications are made in one or more of residues 251-256, 285-290, 308-314, 385-389, and 428-436 of the IgG hinge-Fc region (for example, as in the human IgG1 hinge-Fc region depicted in Figure 4, SEQ ID NO:83), or analogous residues thereof, as determined by amino acid sequence alignment, in other IgG hinge-Fc regions. In a preferred embodiment, the amino acid modifications are made in a human IgG constant domain, or FcRn-binding domain thereof. In a certain embodiment, the modifications are not made at residues 252, 254, or 256 (*i.e.*, all are made at one or more of residues 251, 253, 255, 285-290, 308-314, 385-389, or 428-436) of the IgG constant domain. In a more preferred embodiment, the amino acid modifications are not the substitution with leucine at residue 252, with serine at 254, and/or with phenylalanine at position 256. In particular, in preferred embodiments, such modifications are not made when the IgG constant domain, hinge-Fc domain, hinge-Fc domain or other FcRn-binding fragment thereof is derived from a mouse.

The amino acid modifications may be any modification, preferably at one or more of residues 251-256, 285-290, 308-314, 385-389, and 428-436, that increases the *in vivo* half-life of the IgG constant domain, or FcRn-binding fragment thereof (*e.g.*, Fc or hinge-Fc domain), and any molecule attached thereto, and increases the affinity of the IgG, or fragment thereof, for FcRn. Preferably, the one or more modifications also result in a higher binding affinity of the constant domain, or FcRn-binding fragment thereof, for FcRn at pH 6.0 than at pH 7.4. In other embodiments, the modifications alter (*i.e.*, increase or decrease) bioavailability of the molecule, in particular, alters (*i.e.*, increases or decreases) transport (or concentration or half-life) of the molecule to mucosal surfaces (*e.g.*, of the lungs) or other portions of a target tissue. In a preferred embodiment, the amino acid modifications alter (preferably, increase) transport or concentration or half-life of the

molecule to the lungs. In other embodiments, the amino acid modifications alter (preferably, increase) transport (or concentration or half-life) of the molecule to the heart, pancreas, liver, kidney, bladder, stomach, large or small intestine, respiratory tract, lymph nodes, nervous tissue (central and/or peripheral nervous tissue), muscle, epidermis, bone, cartilage, joints, blood vessels, bone marrow, prostate, ovary, uterine, tumor or cancer tissue, etc. In a preferred embodiment, the amino acid modifications do not abolish, or, more preferably, do not alter, other immune effector or receptor binding functions of the constant domain, for example, but not limited to complement fixation, ADCC and binding to FcγRI, FcγRII, and FcγRIII, as can be determined by methods well-known and routine in the art. In another preferred embodiment, the modified FcRn binding fragment of the constant domain does not contain sequences that mediate immune effector functions or other receptor binding. Such fragments may be particularly useful for conjugation to a non-IgG or non-immunoglobulin molecule to increase the *in vivo* half-life thereof. In yet another embodiment, the effector functions are selectively altered (*e.g.*, to reduce or increase effector functions).

In preferred embodiments, the amino acid modifications are substitutions at one or more of residues 308, 309, 311, 312 and 314, particularly a substitution with threonine at position 308, proline at position 309, serine at position 311, aspartic acid at position 312, and/or leucine at position 314. Alternatively, the modification is the substitution with an isoleucine at position 308, proline at position 309, and/or a glutamic acid at position 311. In yet another embodiment, residues at one or more of positions 308, 309, 311, 312, and 314, are substituted with threonine, proline, leucine, alanine, and alanine, respectively. Accordingly, in certain embodiments the residue at position 308 is substituted with threonine or isoleucine, the residue at position 309 is substituted with proline, the residue at position 311 is substituted with serine, glutamic acid or leucine, the residue at position 312 is substituted with alanine, and/or the residue at position 314 is substituted with leucine or alanine. In a preferred embodiment, the substitution is a threonine at position 308, a proline at position 309, a serine at position 311, an aspartic acid at position 312, and/or a leucine at position 314.

In preferred embodiments, the amino acid modifications are substitutions at one or more of residues 251, 252, 254, 255, and 256. In specific embodiments, residue 251 is substituted with leucine or arginine, residue 252 is substituted with tyrosine, phenylalanine, serine, tryptophan or threonine, residue 254 is substituted with threonine or serine, residue 255 is substituted with arginine, leucine, glycine, or isoleucine, and/or residue 256 is substituted with serine, arginine, glutamine, glutamic acid, aspartic acid,

alanine, asparagine or threonine. In a more specific embodiment, residue 251 is substituted with leucine, residue 252 is substituted with tyrosine, residue 254 is substituted with threonine or serine, residue 255 is substituted with arginine, and/or residue 256 is substituted with glutamic acid.

In preferred embodiments, the amino acid modifications are substitutions at one or more of residues 428, 433, 434, and 436. In specific embodiments, residue 428 is substituted with threonine, methionine, leucine, phenylalanine, or serine, residue 433 is substituted with lysine, arginine, serine, isoleucine, proline, glutamine or histidine, residue 434 is substituted with phenylalanine, tyrosine, or histidine, and/or residue 436 is substituted with histidine, asparagine, arginine, threonine, lysine, or methionine. In a more specific embodiment, residues at position 428 and/or 434 are substituted with methionine, and/or histidine respectively.

In preferred embodiments, the amino acid modifications are substitutions at one or more of residues 385, 386, 387, and 389, more specifically, having substitutions at one or more of these positions. In specific embodiments, residue 385 is substituted with arginine, aspartic acid, serine, threonine, histidine, lysine, alanine or glycine, residue 386 is substituted with threonine, proline, aspartic acid, serine, lysine, arginine, isoleucine, or methionine, residue 387 is substituted with arginine, proline, histidine, serine, threonine, or alanine, and/or residue 389 is substituted with proline, serine or asparagine. In more specific embodiments, residues at one or more positions 385, 386, 387, and 389 are substituted with arginine, threonine, arginine, and proline, respectively. In yet another specific embodiment, residues at one or more positions 385, 386, and 389 are substituted with aspartic acid, proline, and serine, respectively.

In particular embodiments, amino acid modifications are made at one or a combination of residues 251, 252, 254, 255, 256, 308, 309, 311, 312, 314, 385, 386, 387, 389, 428, 433, 434, and/or 436, particularly where the modifications are one or more of the amino acid substitutions described immediately above for these residues.

In a preferred embodiment, the molecule of the invention contains a Fc region, or FcRn-binding domain thereof, having one or more of the following substitutions: leucine at residue 251, tyrosine at residue 252, threonine or serine at residue 254, arginine at residue 255, threonine at residue 308, proline at residue 309, serine at residue 311, aspartic acid at residue 312, leucine at residue 314, arginine at residue 385, threonine at residue 386, arginine at residue 387, proline at residue 389, methionine at residue 428, and/or tyrosine at residue 434.

In a preferred embodiment, the FcRn binding domain has a substitution at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16 or all 18 of residues 251, 252, 254, 255, 256, 308, 309, 311, 312, 314, 385, 386, 387, 389, 428, 433, 434, and/or 436.

5 Amino acid modifications can be made by any method known in the art and many such methods are well known and routine for the skilled artisan. For example, but not by way of limitation, amino acid substitutions, deletions and insertions may be accomplished using any well-known PCR-based technique. Amino acid substitutions may be made by site-directed mutagenesis (see, for example, Zoller and Smith, *Nucl. Acids Res.* 10:6487-6500, 1982; Kunkel, *Proc. Natl. Acad. Sci USA* 82:488, 1985, which are hereby
10 incorporated by reference in their entireties). Mutants that result in increased affinity for FcRn and increased *in vivo* half-life may readily be screened using well-known and routine assays, such as those described in Section 5.11, *infra*. In a preferred method, amino acid substitutions are introduced at one or more residues in the IgG constant domain or FcRn-
15 binding fragment thereof and the mutated constant domains or fragments are expressed on the surface of bacteriophage which are then screened for increased FcRn binding affinity (see, in particular, Section 5.2 and 5.11, *infra*).

Preferably, the amino acid residues to be modified are surface exposed residues. Additionally, in making amino acid substitutions, preferably the amino acid
20 residue to be substituted is a conservative amino acid substitution, for example, a polar residue is substituted with a polar residue, a hydrophilic residue with a hydrophilic residue, hydrophobic residue with a hydrophobic residue, a positively charged residue with a positively charged residue, or a negatively charged residue with a negatively charged residue. Moreover, preferably, the amino acid residue to be modified is not highly or
25 completely conserved across species and/or is critical to maintain the constant domain tertiary structure or to FcRn binding. For example, but not by way of limitation, modification of the histidine at residue 310 is not preferred.

Specific mutants of the Fc domain that have increased affinity for FcRn were isolated after the third-round panning (as described in Section 6) from a library of mutant
30 human IgG1 molecules having mutations at residues 308-314 (histidine at position 310 and tryptophan at position 313 are fixed), those isolated after the fifth-round panning of the library for residues 251-256 (isoleucine at position 253 is fixed), those isolated after fourth-round panning of the library for residues 428-436 (histidine at position 429, glutamic acid at position 430, alanine at position 431, leucine at position 432, and histidine at position 435
35 are fixed), and those isolated after sixth-round panning of the library for residues 385-389 (glutamic acid at position 388 is fixed) are listed in Table I. The wild type human IgG1 has

a sequence Val-Leu-His-Gln-Asp-Trp-Leu (SEQ ID NO:86) at positions 308-314, Leu-Met-Ile-Ser-Arg-Thr (SEQ ID NO:87) at positions 251-256, Met-His-Glu-Ala-Leu-His-Asn-His-Tyr (SEQ ID NO:88) at positions 428-436, and Gly-Gln-Pro-Glu-Asn (SEQ ID NO:89) at positions 385-389.

Table I
MUTANTS ISOLATED BY PANNING

LIBRARY	MUTANTS*
251-256	<u>Leu Tyr Ile Thr Arg Glu</u> (SEQ ID NO:90)
	<i>Leu Tyr Ile Ser Arg Thr</i> (SEQ ID NO:91)
	<i>Leu Tyr Ile Ser Arg Ser</i> (SEQ ID NO:92)
	<i>Leu Tyr Ile Ser Arg Arg</i> (SEQ ID NO:93)
	<i>Leu Tyr Ile Ser Arg Gln</i> (SEQ ID NO:94)
	<i>Leu Trp Ile Ser Arg Thr</i> (SEQ ID NO:95)
	Leu Tyr Ile Ser Leu Gln (SEQ ID NO:96)
	Leu Phe Ile Ser Arg Asp (SEQ ID NO:97)
	Leu Phe Ile Ser Arg Thr (SEQ ID NO:98)
	Leu Phe Ile Ser Arg Arg (SEQ ID NO:99)
	Leu Phe Ile Thr Gly Ala (SEQ ID NO:100)
	Leu Ser Ile Ser Arg Glu (SEQ ID NO:101)
	Arg Thr Ile Ser Ile Ser (SEQ ID NO:102)
308-314	Thr Pro His Ser Asp Trp Leu (SEQ ID NO:103)
	Ile Pro His Glu Asp Trp Leu (SEQ ID NO:104)
385-389	Arg Thr Arg Glu Pro (SEQ ID NO:105)
	<i>Asp Pro Pro Glu Ser</i> (SEQ ID NO:106)
	Ser Asp Pro Glu Pro (SEQ ID NO:107)
	Thr Ser His Glu Asn (SEQ ID NO:108)
	Ser Lys Ser Glu Asn (SEQ ID NO:109)
	His Arg Ser Glu Asn (SEQ ID NO:110)
	Lys Ile Arg Glu Asn (SEQ ID NO:111)

LIBRARY	MUTANTS*
	Gly Ile Thr Glu Ser (SEQ ID NO:112)
	Ser Met Ala Glu Pro (SEQ ID NO:113)
5 428-436	<i>Met His Glu Ala Leu Arg Tyr His His</i> (SEQ ID NO:114)
	Met His Glu Ala Leu His Phe His His (SEQ ID NO:115)
	Met His Glu Ala Leu Lys Phe His His (SEQ ID NO:116)
	Met His Glu Ala Leu Ser Tyr His Arg (SEQ ID NO:117)
10	Thr His Glu Ala Leu His Tyr His Thr (SEQ ID NO:118)
	Met His Glu Ala Leu His Tyr His Tyr (SEQ ID NO:119)

* Substituting residues are indicated in bold face

The underlined sequences in Table I correspond to sequences that occurred 10 to 20 times in the final round of panning and the sequences in italics correspond to sequences that occurred 2 to 5 times in the final round of panning. Those sequences that are neither underlined nor italicized occurred once in the final round of panning.

In one preferred embodiment, the invention provides modified immunoglobulin molecules (*e.g.*, various antibodies) that have increased *in vivo* half-life and affinity for FcRn relative to unmodified molecules (and, in preferred embodiments, altered bioavailability such as increased or decreased transport to mucosal surfaces or other target tissues). Such immunoglobulin molecules include IgG molecules that naturally contain an FcRn binding domain and other non-IgG immunoglobulins (*e.g.*, IgE, IgM, IgD, IgA and IgY) or fragments of immunoglobulins that have been engineered to contain an FcRn-binding fragment (*i.e.*, fusion proteins comprising non-IgG immunoglobulin or a portion thereof and an FcRn binding domain). In both cases the FcRn-binding domain has one or more amino acid modifications that increase the affinity of the constant domain fragment for FcRn.

The modified immunoglobulins include any immunoglobulin molecule that binds (preferably, immunospecifically, *i.e.*, competes off non-specific binding), as determined by immunoassays well known in the art for assaying specific antigen-antibody binding) an antigen and contains an FcRn-binding fragment. Such antibodies include, but are not limited to, polyclonal, monoclonal, bi-specific, multi-specific, human, humanized, chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, disulfide-linked Fvs, and fragments containing either a VL or VH domain or even a complementary

determining region (CDR) that specifically binds an antigen, in certain cases, engineered to contain or fused to an FcRn binding domain.

The IgG molecules of the invention, and FcRn-binding fragments thereof, are preferably IgG1 subclass of IgGs, but may also be any other IgG subclasses of given animals. For example, in humans, the IgG class includes IgG1, IgG2, IgG3, and IgG4; and mouse IgG includes IgG1, IgG2a, IgG2b, IgG2c and IgG3. It is known that certain IgG subclasses, for example, mouse IgG2b and IgG2c, have higher clearance rates than, for example, IgG1 (Medesan *et al.*, *Eur. J. Immunol.*, 28:2092-2100, 1998). Thus, when using IgG subclasses other than IgG1, it may be advantageous to substitute one or more of the residues, particularly in the CH2 and CH3 domains, that differ from the IgG1 sequence with those of IgG1, thereby increasing the *in vivo* half-life of the other types of IgG.

The immunoglobulins (and other proteins used herein) may be from any animal origin including birds and mammals. Preferably, the antibodies are human, rodent (e.g., mouse and rat), donkey, sheep, rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described *infra* and, for example, in U.S. Patent No. 5,939,598 by Kucherlapati *et al.*

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide or may be specific for heterologous epitopes, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, *et al.*, *J. Immunol.*, 147:60-69, 1991; U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny *et al.*, *J. Immunol.*, 148:1547-1553, 1992.

The antibodies of the invention include derivatives that are otherwise modified, *i.e.*, by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from binding antigen and/or generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, *e.g.*, by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to, specific chemical cleavage, acetylation, formylation, metabolic synthesis of

tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow *et al.*, *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, *et al.*, in: *Monoclonal Antibodies and T-Cell Hybridomas*, pp. 563-681 (Elsevier, N.Y., 1981) (both of which are incorporated herein by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art. In a non-limiting example, mice can be immunized with an antigen of interest or a cell expressing such an antigen. Once an immune response is detected, *e.g.*, antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells. Hybridomas are selected and cloned by limiting dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding the antigen. Ascites fluid, which generally contains high levels of antibodies, can be generated by inoculating mice intraperitoneally with positive hybridoma clones.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the complete light chain, and the variable region, the CH1 region and the hinge region of the heavy chain.

For example, antibodies can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains, such as Fab and Fv or disulfide-bond stabilized Fv, expressed from a repertoire or combinatorial antibody library (*e.g.*, human or murine). Phage expressing an

antigen binding domain that binds the antigen of interest can be selected or identified with antigen, *e.g.*, using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage, including fd and M13. The antigen binding domains are expressed as a recombinantly fused protein to either the phage gene III or gene VIII protein. Alternatively, the modified FcRn binding portion of immunoglobulins of the present invention can be also expressed in a phage display system. Examples of phage display methods that can be used to make the immunoglobulins, or fragments thereof, of the present invention include those disclosed in Brinkman *et al.*, *J. Immunol. Methods*, 182:41-50, 1995; Ames *et al.*, *J. Immunol. Methods*, 184:177-186, 1995; Kettleborough *et al.*, *Eur. J. Immunol.*, 24:952-958, 1994; Persic *et al.*, *Gene*, 187:9-18, 1997; Burton *et al.*, *Advances in Immunology*, 57:191-280, 1994; PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/1 1236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired fragments, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, *e.g.*, as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax *et al.*, *BioTechniques*, 12(6):864-869, 1992; and Sawai *et al.*, *AJRI*, 34:26-34, 1995; and Better *et al.*, *Science*, 240:1041-1043, 1988 (each of which is incorporated by reference in its entirety). Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patent Nos. 4,946,778 and 5,258,498; Huston *et al.*, *Methods in Enzymology*, 203:46-88, 1991; Shu *et al.*, *PNAS*, 90:7995-7999, 1993; and Skerra *et al.*, *Science*, 240:1038-1040, 1988.

For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a constant region derived from a human immunoglobulin. Methods for producing chimeric antibodies are known in the art. See *e.g.*, Morrison,

Science, 229:1202, 1985; Oi *et al.*, *BioTechniques*, 4:214 1986; Gillies *et al.*, *J. Immunol. Methods*, 125:191-202, 1989; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entireties. Humanized antibodies are antibody
5 molecules from non-human species that bind the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are
10 identified by methods well known in the art, *e.g.*, by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. See, *e.g.*, Queen *et al.*, U.S. Patent No. 5,585,089; Riechmann *et al.*, *Nature*, 332:323, 1988, which are incorporated herein by reference in their entireties. Antibodies can be humanized
15 using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101 and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology*, 28(4/5):489-498, 1991; Studnicka *et al.*, *Protein Engineering*, 7(6):805-814, 1994; Roguska *et al.*, *Proc Natl. Acad. Sci. USA*, 91:969-973, 1994), and chain shuffling
20 (U.S. Patent No. 5,565,332), all of which are hereby incorporated by reference in their entireties.

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries
25 derived from human immunoglobulin sequences. See U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645; WO 98/50433; WO 98/24893; WO 98/16654; WO 96/34096; WO 96/33735; and WO 91/10741, each of which is incorporated herein by reference in its entirety.

Human antibodies can also be produced using transgenic mice which are
30 incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.*, 13:65-93, 1995. For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, *e.g.*, PCT publications WO
35 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806;

5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entireties. In addition, companies such as Abgenix, Inc. (Freemont, CA), Medarex (NJ) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed
 5 against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, *e.g.*, a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers *et al.*, *Bio/technology*,
 10 12:899-903, 1988).

In particular embodiments, the modified antibodies have *in vivo* therapeutic and/or prophylactic uses. Examples of therapeutic and prophylactic antibodies which may be so modified include, but are not limited to, SYNAGIS® (MedImmune, MD) which is a humanized anti-respiratory syncytial virus (RSV) monoclonal antibody for the treatment of
 15 patients with RSV infection; HERCEPTIN® (Trastuzumab) (Genentech, CA) which is a humanized anti-HER2 monoclonal antibody for the treatment of patients with metastatic breast cancer; REMICADE® (infliximab) (Centocor, PA) which is a chimeric anti-TNF α monoclonal antibody for the treatment of patients with Crone's disease; REOPRO® (abciximab) (Centocor) which is an anti-glycoprotein IIb/IIIa receptor on the platelets for
 20 the prevention of clot formation; ZENAPAX® (daclizumab) (Roche Pharmaceuticals, Switzerland) which is an immunosuppressive, humanized anti-CD25 monoclonal antibody for the prevention of acute renal allograft rejection. Other examples are a humanized anti-CD18 F(ab')₂ (Genentech); CDP860 which is a humanized anti-CD18 F(ab')₂ (Celltech, UK); PRO542 which is an anti-HIV gp120 antibody fused with CD4 (Progenics/Genzyme
 25 Transgenics); Ostavir which is a human anti Hepatitis B virus antibody (Protein Design Lab/Novartis); PROTOVIR™ which is a humanized anti-CMV IgG1 antibody (Protein Design Lab/Novartis); MAK-195 (SEGARD) which is a murine anti-TNF- α F(ab')₂ (Knoll Pharma/BASF); IC14 which is an anti-CD14 antibody (ICOS Pharm); a humanized anti-VEGF IgG1 antibody (Genentech); OVAREX™ which is a murine anti-CA 125 antibody
 30 (Altarex); PANOREX™ which is a murine anti-17-IA cell surface antigen IgG2a antibody (Glaxo Wellcome/Centocor); BEC2 which is a murine anti-idiotypic (GD3 epitope) IgG antibody (ImClone System); IMC-C225 which is a chimeric anti-EGFR IgG antibody (ImClone System); VITAXIN™ which is a humanized anti- α V β 3 integrin antibody (Applied Molecular Evolution/MedImmune); Campath 1H/LDP-03 which is a humanized
 35 anti CD52 IgG1 antibody (Leukosite); Smart M195 which is a humanized anti-CD33 IgG antibody (Protein Design Lab/Kanebo); RITUXAN™ which is a chimeric anti-CD20 IgG1

antibody (IDEC Pharm/Genentech, Roche/Zettyaku); LYMPHOCIDE™ which is a humanized anti-CD22 IgG antibody (Immunomedics); Smart ID10 which is a humanized anti-HLA antibody (Protein Design Lab); ONCOLYM™ (Lym-1) is a radiolabelled murine anti-HLA DIAGNOSTIC REAGENT antibody (Techniclone); ABX-IL8 is a human anti-IL8 antibody (Abgenix); anti-CD11a is a humanized IgG1 antibody (Genetech/Xoma); ICM3 is a humanized anti-ICAM3 antibody (ICOS Pharm); IDEC-114 is a primatized anti-CD80 antibody (IDEC Pharm/Mitsubishi); ZEVALIN™ is a radiolabelled murine anti-CD20 antibody (IDEC/Schering AG); IDEC-131 is a humanized anti-CD40L antibody (IDEC/Eisai); IDEC-151 is a primatized anti-CD4 antibody (IDEC); IDEC-152 is a primatized anti-CD23 antibody (IDEC/Seikagaku); SMART anti-CD3 is a humanized anti-CD3 IgG (Protein Design Lab); 5G1.1 is a humanized anti-complement factor 5 (C5) antibody (Alexion Pharm); D2E7 is a humanized anti-TNF- α antibody (CAT/BASF); CDP870 is a humanized anti-TNF- α Fab fragment (Celltech); IDEC-151 is a primatized anti-CD4 IgG1 antibody (IDEC Pharm/SmithKline Beecham); MDX-CD4 is a human anti-CD4 IgG antibody (Medarex/Eisai/Genmab); CDP571 is a humanized anti-TNF- α IgG4 antibody (Celltech); LDP-02 is a humanized anti- α 4 β 7 antibody (LeukoSite/Genentech); OrthoClone OKT4A is a humanized anti-CD4 IgG antibody (Ortho Biotech); ANTOVA™ is a humanized anti-CD40L IgG antibody (Biogen); ANTEGREN™ is a humanized anti-VLA-4 IgG antibody (Elan); MDX-33 is a human anti-CD64 (Fc γ R) antibody (Medarex/Centeon); SCH55700 is a humanized anti-IL-5 IgG4 antibody (Celltech/Schering); SB-240563 and SB-240683 are humanized anti-IL-5 and IL-4 antibodies, respectively, (SmithKline Beecham); rhuMab-E25 is a humanized anti-IgE IgG1 antibody (Genentech/Norvartis/Tanox Biosystems); IDEC-152 is a primatized anti-CD23 antibody (IDEC Pharm); ABX-CBL is a murine anti CD-147 IgM antibody (Abgenix); BTI-322 is a rat anti-CD2 IgG antibody (Medimmune/Bio Transplant); Orthoclone/OKT3 is a murine anti-CD3 IgG2a antibody (ortho Biotech); SIMULECT™ is a chimeric anti-CD25 IgG1 antibody (Novartis Pharm); LDP-01 is a humanized anti- β ₂-integrin IgG antibody (LeukoSite); Anti-LFA-1 is a murine anti CD18 F(ab')₂ (Pasteur-Merieux/Immunotech); CAT-152 is a human anti-TGF- β ₂ antibody (Cambridge Ab Tech); and Corsevin M is a chimeric anti-Factor VII antibody (Centocor).

In specific embodiments, the invention provides modified antibodies having one or more of the mutations described herein and that immunospecifically bind RSV, *e.g.*, SYNAGIS®. The present invention also provides modified antibodies having one or more of the mutations described herein and that comprise a variable heavy (VH) and/or variable light (VL) domain having the amino acid sequence of any VH and/or VL domain listed in

Table III. The present invention further encompasses anti-RSV antibodies comprising one or more VH complementarity determining regions (CDRs) and/or one or more VL CDRs having the amino acid sequence of one or more VH CDRs and/or VL CDRS listed in Table
 5 III or one or more of the CDRs listed in Table II wherein one or more of the bolded and underlined residues has an amino acid substitution, preferably that increases the affinity of the antibody for RSV. In specific embodiments, the antibody to be modified is AFFF, p12f2, p12f4, p11d4, Ale109, A12a6, A13c4, A17d4, A4B4, A8C7, 1X-493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L215B10, A13A11, A1H5, A4B4(1),
 10 A4B4L1FR-S28R, A4B4-F52S.

Table II.
CDR Sequences of SYNAGIS®

	CDR	Sequence	SEQ ID NO:
	VH1	T <u>S</u> GMSVG	1
15	VH2	DIWWDD <u>DKKD</u> YNPSLK <u>S</u>	2
	VH3	<u>S</u> MITN <u>W</u> YFDV	3
	VL1	<u>KCQLS</u> VG ^Y MH	4
	VL2	DT <u>SKLAS</u>	5
20	VL3	FQGS <u>G</u> YP <u>F</u> T	6

25

30

35

Table III
ANTI-RSV ANTIBODIES

Antibody Name	VH Domain	VH CDR1	VH CDR2	VH CDR3	VL Domain	VL CDR1	VL CDR2	VL CDR3
SYNAGIS	SEQ ID NO:7	TSGMSVG (SEQ ID NO:1)	DIWWDKDKDYNPSLKS (SEQ ID NO:2)	SMITNWFYDV (SEQ ID NO:3)	SEQ ID NO:8	KCQLSVGYMH (SEQ ID NO:4)	DTSKLAS (SEQ ID NO:5)	FQSGYPFT (SEQ ID NO:6)
AFF	SEQ ID NO:9	TAGMSVG (SEQ ID NO:10)	DIWWDKDKDYNPSLKS (SEQ ID NO:2)	SMITNWFYDV (SEQ ID NO:12)	SEQ ID NO:13	SASSSVGYMH (SEQ ID NO:14)	DTEKLAS (SEQ ID NO:15)	FQSGYPFT (SEQ ID NO:16)
p12f2	SEQ ID NO:17	TPGMSVG (SEQ ID NO:18)	DIWWDKDKKHYNPSLKD (SEQ ID NO:19)	DMITNWFYDV (SEQ ID NO:20)	SEQ ID NO:21	SLSSRVGYMH (SEQ ID NO:22)	DTFYLS (SEQ ID NO:23)	FQSGYPFT (SEQ ID NO:6)
p12f4	SEQ ID NO:24	TPGMSVG (SEQ ID NO:18)	DIWWDGKKKHYNPSLKD (SEQ ID NO:25)	DMITNWFYDV (SEQ ID NO:20)	SEQ ID NO:26	SLSSRVGYMH (SEQ ID NO:22)	DTRGLPS (SEQ ID NO:27)	FQSGYPFT (SEQ ID NO:6)
p11d4	SEQ ID NO:28	TPGMSVG (SEQ ID NO:18)	DIWWDGKKKHYNPSLKD (SEQ ID NO:25)	DMITNWFYDV (SEQ ID NO:29)	SEQ ID NO:30	SPSSRVGYMH (SEQ ID NO:31)	DTMRLAS (SEQ ID NO:32)	FQSGYPFT (SEQ ID NO:6)
Ale109	SEQ ID NO:33	TAGMSVG (SEQ ID NO:10)	DIWWDGKKKHYNPSLKD (SEQ ID NO:25)	DMITNWFYDV (SEQ ID NO:29)	SEQ ID NO:34	SLSSRVGYMH (SEQ ID NO:22)	DTEKLSS (SEQ ID NO:35)	FQSGYPFT (SEQ ID NO:6)
A12a6	SEQ ID NO:36	TAGMSVG (SEQ ID NO:10)	DIWWDGKKDYNPSLKD (SEQ ID NO:37)	DMITNWFYDV (SEQ ID NO:20)	SEQ ID NO:38	SASSRVGYMH (SEQ ID NO:39)	DTEKLSS (SEQ ID NO:35)	FQSGYPFT (SEQ ID NO:6)
A13c4	SEQ ID NO:40	TAGMSVG (SEQ ID NO:10)	DIWWDGKKKSYNPSLKD (SEQ ID NO:41)	DMITNWFYDV (SEQ ID NO:20)	SEQ ID NO:42	SLSSRVGYMH (SEQ ID NO:22)	DTMYQSS (SEQ ID NO:43)	FQSGYPFT (SEQ ID NO:6)
A17d4	SEQ ID NO:44	TAGMSVG (SEQ ID NO:10)	DIWWDKDKSYNPSLKD (SEQ ID NO:45)	DMITNWFYDV (SEQ ID NO:20)	SEQ ID NO:46	LPSSRVGYMH (SEQ ID NO:47)	DTMYQSS (SEQ ID NO:43)	FQSGYPFT (SEQ ID NO:6)

Antibody Name	VH Domain	VH CDR1	VH CDR2	VH CDR3	VL Domain	VL CDR1	VL CDR2	VL CDR3
A4B4	SEQ ID NO:48	TAGMSVG (SEQ ID NO:10)	DIWDDKKHYNPSLKD (SEQ ID NO:19)	DMITNFIYFDV (SEQ ID NO:20)	SEQ ID NO:49	SASSRVGYMH (SEQ ID NO:39)	DTEFLDS (SEQ ID NO:50)	FQSGYPFT (SEQ ID NO:6)
A8C7	SEQ ID NO:51	TAGMSVG (SEQ ID NO:10)	DIWDDKKSYNPSLKD (SEQ ID NO:45)	DMITNFIYFDV (SEQ ID NO:29)	SEQ ID NO:52	SPSSRVGYMH (SEQ ID NO:31)	DTRYQSS (SEQ ID NO:53)	FQSGYPFT (SEQ ID NO:6)
1X-493L1FR	SEQ ID NO:7	TSGMSVG (SEQ ID NO:1)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	SMITNFIYFDV (SEQ ID NO:3)	SEQ ID NO:54	SASSSVGYMH (SEQ ID NO:14)	DTSKLAS (SEQ ID NO:5)	FQSGYPFT (SEQ ID NO:6)
H3-3F4	SEQ ID NO:55	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:29)	SEQ ID NO:56	SASSSVGYMH (SEQ ID NO:14)	DTEKLAS (SEQ ID NO:15)	FQSGYPFT (SEQ ID NO:6)
M3H9	SEQ ID NO:55	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:29)	SEQ ID NO:57	SASSSVGYMH (SEQ ID NO:14)	DTYKQTS (SEQ ID NO:58)	FQSGYPFT (SEQ ID NO:6)
Y10H6	SEQ ID NO:55	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:29)	SEQ ID NO:59	SASSSVGYMH (SEQ ID NO:14)	DTRYLSS (SEQ ID NO:125)	FQSGYPFT (SEQ ID NO:6)
DG	SEQ ID NO:78	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:79)	SEQ ID NO:56	SASSSVGYMH (SEQ ID NO:14)	DTEKLAS (SEQ ID NO:15)	FQSGYPFT (SEQ ID NO:6)
AFFF(1)	SEQ ID NO:9	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	SMITNFIYFDV (SEQ ID NO:12)	SEQ ID NO:60	SASSSVGYMH (SEQ ID NO:14)	DTEKLAS (SEQ ID NO:15)	FQSGYPFT (SEQ ID NO:61)
6H8	SEQ ID NO:78	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:79)	SEQ ID NO:62	SASSSVGYMH (SEQ ID NO:14)	DTEKLTS (SEQ ID NO:63)	FQSGYPFT (SEQ ID NO:6)
L1-7E5	SEQ ID NO:78	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:79)	SEQ ID NO:64	SASSRVGYMH (SEQ ID NO:39)	DTEKLAS (SEQ ID NO:15)	FQSGYPFT (SEQ ID NO:6)
L215B10	SEQ ID NO:78	TAGMSVG (SEQ ID NO:10)	DIWDDKKDYNPSLKS (SEQ ID NO:2)	DMITNFIYFDV (SEQ ID NO:79)	SEQ ID NO:65	SASSSVGYMH (SEQ ID NO:14)	DTEKLAS (SEQ ID NO:66)	FQSGYPFT (SEQ ID NO:6)
A13A11	SEQ ID NO:67	TAGMSVG (SEQ ID NO:10)	DIWDDKKHYNPSLKD (SEQ ID NO:19)	DMITNFIYFDV (SEQ ID NO:29)	SEQ ID NO:68	SPSSRVGYMH (SEQ ID NO:31)	DTYRHSS (SEQ ID NO:69)	FQSGYPFT (SEQ ID NO:6)

Antibody Name	VH Domain	VH CDR1	VH CDR2	VH CDR3	VL Domain	VL CDR1	VL CDR2	VL CDR3
A1H5	SEQ ID NO:70	<u>T</u> <u>A</u> <u>G</u> <u>M</u> <u>S</u> <u>V</u> <u>G</u> (SEQ ID NO:10)	<u>D</u> <u>I</u> <u>W</u> <u>W</u> <u>D</u> <u>D</u> <u>K</u> <u>K</u> <u>H</u> <u>Y</u> <u>N</u> <u>P</u> <u>S</u> <u>L</u> <u>K</u> <u>D</u> (SEQ ID NO:25)	<u>D</u> <u>M</u> <u>I</u> <u>E</u> <u>N</u> <u>W</u> <u>Y</u> <u>F</u> <u>D</u> <u>V</u> (SEQ ID NO:29)	SEQ ID NO:71	<u>S</u> <u>L</u> <u>S</u> <u>S</u> <u>S</u> <u>V</u> <u>G</u> <u>Y</u> <u>M</u> <u>H</u> (SEQ ID NO:72)	<u>D</u> <u>T</u> <u>F</u> <u>F</u> <u>H</u> <u>R</u> <u>S</u> (SEQ ID NO:73)	<u>F</u> <u>Q</u> <u>G</u> <u>S</u> <u>G</u> <u>Y</u> <u>P</u> <u>F</u> <u>T</u> (SEQ ID NO:6)
A4B4(1)	SEQ ID NO:48	<u>T</u> <u>A</u> <u>G</u> <u>M</u> <u>S</u> <u>V</u> <u>G</u> (SEQ ID NO:10)	<u>D</u> <u>I</u> <u>W</u> <u>W</u> <u>D</u> <u>D</u> <u>K</u> <u>K</u> <u>H</u> <u>Y</u> <u>N</u> <u>P</u> <u>S</u> <u>L</u> <u>K</u> <u>D</u> (SEQ ID NO:19)	<u>D</u> <u>M</u> <u>I</u> <u>E</u> <u>N</u> <u>W</u> <u>Y</u> <u>F</u> <u>D</u> <u>V</u> (SEQ ID NO:20)	SEQ ID NO:74	<u>S</u> <u>A</u> <u>S</u> <u>S</u> <u>R</u> <u>V</u> <u>G</u> <u>Y</u> <u>M</u> <u>H</u> (SEQ ID NO:39)	<u>D</u> <u>T</u> <u>L</u> <u>L</u> <u>L</u> <u>D</u> <u>S</u> (SEQ ID NO:75)	<u>F</u> <u>Q</u> <u>G</u> <u>S</u> <u>G</u> <u>Y</u> <u>P</u> <u>F</u> <u>T</u> (SEQ ID NO:6)
A4B4L1F R-S28R	SEQ ID NO:48	<u>T</u> <u>A</u> <u>G</u> <u>M</u> <u>S</u> <u>V</u> <u>G</u> (SEQ ID NO:10)	<u>D</u> <u>I</u> <u>W</u> <u>W</u> <u>D</u> <u>D</u> <u>K</u> <u>K</u> <u>H</u> <u>Y</u> <u>N</u> <u>P</u> <u>S</u> <u>L</u> <u>K</u> <u>D</u> (SEQ ID NO:19)	<u>D</u> <u>M</u> <u>I</u> <u>E</u> <u>N</u> <u>W</u> <u>Y</u> <u>F</u> <u>D</u> <u>V</u> (SEQ ID NO:20)	SEQ ID NO:11	<u>S</u> <u>A</u> <u>S</u> <u>S</u> <u>R</u> <u>V</u> <u>G</u> <u>Y</u> <u>M</u> <u>H</u> (SEQ ID NO:39)	<u>D</u> <u>T</u> <u>S</u> <u>K</u> <u>L</u> <u>A</u> <u>S</u> (SEQ ID NO:5)	<u>F</u> <u>Q</u> <u>G</u> <u>S</u> <u>G</u> <u>Y</u> <u>P</u> <u>F</u> <u>T</u> (SEQ ID NO:6)
A4B4-F52S	SEQ ID NO:48	<u>T</u> <u>A</u> <u>G</u> <u>M</u> <u>S</u> <u>V</u> <u>G</u> (SEQ ID NO:10)	<u>D</u> <u>I</u> <u>W</u> <u>W</u> <u>D</u> <u>D</u> <u>K</u> <u>K</u> <u>H</u> <u>Y</u> <u>N</u> <u>P</u> <u>S</u> <u>L</u> <u>K</u> <u>D</u> (SEQ ID NO:19)	<u>D</u> <u>M</u> <u>I</u> <u>E</u> <u>N</u> <u>W</u> <u>Y</u> <u>F</u> <u>D</u> <u>V</u> (SEQ ID NO:20)	SEQ ID NO:76	<u>S</u> <u>A</u> <u>S</u> <u>S</u> <u>R</u> <u>V</u> <u>G</u> <u>Y</u> <u>M</u> <u>H</u> (SEQ ID NO:39)	<u>D</u> <u>T</u> <u>S</u> <u>E</u> <u>L</u> <u>D</u> <u>S</u> (SEQ ID NO:77)	<u>F</u> <u>Q</u> <u>G</u> <u>S</u> <u>G</u> <u>Y</u> <u>P</u> <u>F</u> <u>T</u> (SEQ ID NO:6)

In other embodiments, the antibody is a modified anti- $\alpha_v\beta_3$ antibody, preferably a Vitaxin antibody (see, PCT publications WO 98/33919 and WO 00/78815, both by Huse *et al.*, and both of which are incorporated by reference herein in their entireties).

5 Modified IgGs of the present invention having longer half-lives than wild type may also include IgGs whose bioactive sites, such as antigen-binding sites, Fc-receptor binding sites, or complement-binding sites, are modified by genetic engineering to increase or reduce such activities compared to the wild type.

10 Modification of these and other therapeutic antibodies to increase the *in vivo* half-life permits administration of lower effective dosages and/or less frequent dosing of the therapeutic antibody. Such modification to increase *in vivo* half-life can also be useful to improve diagnostic immunoglobulins as well, for example, permitting administration of lower doses to achieve sufficient diagnostic sensitivity.

The present invention also provides fusion proteins comprising a bioactive
15 molecule and an hinge-Fc region or a fragment thereof (preferably human) having one or more modifications (*i.e.*, substitutions, deletions, or insertions) in amino acid residues identified to be involved in the interaction between the hinge-Fc region and the FcRn receptor. In particular, the present invention provides fusion proteins comprising a
20 bioactive molecule recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a CH2 domain having one or more modifications in amino acid residues 251-256, 285-290, and/or amino acid residues 308-314, and/or to a CH3 domain having one or more modifications in amino acid residues 385-389 and/or 428-436, in particular, one or more of the amino acid substitutions discussed above. The fusion
25 of a bioactive molecule to a constant domain or a fragment thereof with one or more of such modifications increases the *in vivo* half-life of the bioactive molecule.

In a preferred embodiment, fusion proteins of the invention comprise a bioactive molecule recombinantly fused or chemically conjugated to a CH2 domain having one or more amino acid residue substitutions in amino acid residues 251-256, 285-290, and/or amino acid residues 308-314, and/or to a CH3 domain having one or more
30 modifications in amino acid residues 385-389 and/or 428-436. In certain embodiments, a fusion protein comprises a CH2 domain of IgG molecule in which amino acid residues 253, 310, and 313 are not modified. In another embodiment, a fusion protein comprises a CH3 domain of IgG molecule in which amino acid residues 388, 429, 430, 431, 432, and 435 are not modified.

35 A bioactive molecule can be any polypeptide or synthetic drug known to one of skill in the art. Preferably, a bioactive molecule is a polypeptide consisting of at least 5,

preferably at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acid residues. Examples of bioactive polypeptides include, but are not limited to, various types of antibodies, cytokines (*e.g.*, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-10, IL-12, IL-15, IFN- γ , IFN- α , and IFN- β), cell adhesion molecules (5 *e.g.*, CTLA4, CD2, and CD28), ligands (*e.g.*, TNF- α , TNF- β , and an anti-angiogenic factor such as endostatin), receptors, antibodies and growth factors (*e.g.*, PDGF, EGF, NGF, and KGF).

A bioactive molecule can also be a therapeutic moiety such as a cytotoxin (10 *e.g.*, a cytostatic or cytocidal agent), a therapeutic agent or a radioactive element (*e.g.*, alpha-emitters, gamma-emitters, etc.). Examples of cytostatic or cytocidal agents include, but are not limited to, paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-15 dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, 20 dibromomannitol, streptozotocin, mitomycin C, and cisdichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

25 The present invention also provides polynucleotides comprising a nucleotide sequence encoding a modified IgG of the invention and fragments thereof which contain the modified FcRn binding sites with increased affinity and vectors comprising said polynucleotides. Furthermore, the invention includes polynucleotides that hybridize under stringent or lower stringent hybridization conditions to polynucleotides encoding modified 30 IgGs of the present invention.

The nucleotide sequence of modified IgGs and the polynucleotides encoding the same may be obtained by any methods known in the art, including general DNA sequencing method, such as dideoxy chain termination method (Sanger sequencing), and oligonucleotide priming in combination with PCR, respectively.

35

5.2. IDENTIFICATION OF MUTATIONS WITHIN THE HINGE-FC REGION OF IMMUNOGLOBULIN MOLECULES

5 One or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and 428-436 of the constant domain may be introduced utilizing any technique known to those of skill in the art. The constant domain or fragment thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and 428-436 may be screened by, for example, a binding assay to identify the constant
10 domain or fragment thereof with increased affinity for the FcRn receptor (*e.g.*, as described in section 5.11, *infra*). Those modifications in the hinge-Fc domain or the fragments thereof which increase the affinity of the constant domain or fragment thereof for the FcRn receptor can be introduced into antibodies to increase the *in vivo* half-lives of said antibodies. Further, those modifications in the constant domain or the fragment thereof
15 which increase the affinity of the constant domain or fragment thereof for the FcRn can be fused to bioactive molecules to increase the *in vivo* half-lives of said bioactive molecules (and, preferably alter (increase or decrease) the bioavailability of the molecule, for example, to increase or decrease transport to mucosal surfaces (or other target tissue) (*e.g.*, the lungs).

20 5.2.1. MUTAGENESIS

Mutagenesis may be performed in accordance with any of the techniques known in the art including, but not limited to, synthesizing an oligonucleotide having one or more modifications within the sequence of the constant domain of an antibody or a fragment thereof (*e.g.*, the CH2 or CH3 domain) to be modified. Site-specific mutagenesis
25 allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to about 75 nucleotides or more in length is preferred, with
30 about 10 to about 25 or more residues on both sides of the junction of the sequence being altered. A number of such primers introducing a variety of different mutations at one or more positions may be used to generated a library of mutants.

The technique of site-specific mutagenesis is well known in the art, as exemplified by various publications (see, *e.g.*, Kunkel *et al.*, *Methods Enzymol.*,
35 154:367-82, 1987, which is hereby incorporated by reference in its entirety). In general, site-directed mutagenesis is performed by first obtaining a single-stranded vector or melting

apart of two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as T7 DNA polymerase, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform or transfect appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

Alternatively, the use of PCRTM with commercially available thermostable enzymes such as *Taq* DNA polymerase may be used to incorporate a mutagenic oligonucleotide primer into an amplified DNA fragment that can then be cloned into an appropriate cloning or expression vector. See, e.g., Tomic *et al.*, *Nucleic Acids Res.*, 18(6):1656, 1987, and Upender *et al.*, *Biotechniques*, 18(1):29-30, 32, 1995, for PCRTM - mediated mutagenesis procedures, which are hereby incorporated in their entireties. PCRTM employing a thermostable ligase in addition to a thermostable polymerase may also be used to incorporate a phosphorylated mutagenic oligonucleotide into an amplified DNA fragment that may then be cloned into an appropriate cloning or expression vector (see e.g., Michael, *Biotechniques*, 16(3):410-2, 1994, which is hereby incorporated by reference in its entirety).

Other methods known to those of skill in art of producing sequence variants of the Fc domain of an antibody or a fragment thereof can be used. For example, recombinant vectors encoding the amino acid sequence of the constant domain of an antibody or a fragment thereof may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants.

5.2.2. PANNING

Vectors, in particular, phage, expressing constant domains or fragments thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and/or 428-436 can be screened to identify constant domains or fragments thereof

having increased affinity for FcRn to select out the highest affinity binders from a population of phage. Immunoassays which can be used to analyze binding of the constant domain or fragment thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and/or 428-436 to the FcRn include, but are not limited to, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, and fluorescent immunoassays. Such assays are routine and well known in the art (see, e.g., Ausubel *et al.*, eds, 1994, *Current Protocols in Molecular Biology*, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly herein below (but are not intended by way of limitation). BIAcore kinetic analysis can also be used to determine the binding on and off rates of a constant domain or a fragment thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and/or 428-436 to the FcRn. BIAcore kinetic analysis comprises analyzing the binding and dissociation of a constant domain or a fragment thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and/or 428-436 from chips with immobilized FcRn on their surface (see section 5.1 and the Example section *infra*).

5.2.3. SEQUENCING

Any of a variety of sequencing reactions known in the art can be used to directly sequence the nucleotide sequence encoding constant domains or fragments thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and/or 428-436. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert (*Proc. Natl. Acad. Sci. USA*, 74:560, 1977) or Sanger (*Proc. Natl. Acad. Sci. USA*, 74:5463, 1977). It is also contemplated that any of a variety of automated sequencing procedures can be utilized (*Bio/Techniques*, 19:448, 1995), including sequencing by mass spectrometry (see, e.g., PCT Publication No. WO 94/16101, Cohen *et al.*, *Adv. Chromatogr.*, 36:127-162, 1996, and Griffin *et al.*, *Appl. Biochem. Biotechnol.*, 38:147-159, 1993).

5.3. RECOMBINANT METHODS OF PRODUCING ANTIBODIES

The antibodies of the invention or fragments thereof can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

The nucleotide sequence encoding an antibody may be obtained from any information available to those of skill in the art (*i.e.*, from Genbank, the literature, or by routine cloning). If a clone containing a nucleic acid encoding a particular antibody or an epitope-binding fragment thereof is not available, but the sequence of the antibody molecule or epitope-binding fragment thereof is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (*e.g.*, an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A⁺ RNA, isolated from any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, *e.g.*, a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, *e.g.*, recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook *et al.*, 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY; and Ausubel *et al.*, eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence by, for example, introducing amino acid substitutions, deletions, and/or insertions into the epitope-binding domain regions of the antibodies and preferably, into the hinge-Fc regions of the antibodies which are involved in the interaction with the FcRn. In a preferred embodiment, antibodies having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and 428-436 are generated.

Recombinant expression of an antibody requires construction of an expression vector containing a nucleotide sequence that encodes the antibody. Once a nucleotide sequence encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably, but not necessarily, containing the heavy or light chain variable region) has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and

appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide
5 sequence encoding the constant region of the antibody molecule with one or more modifications in the amino acid residues involved in the interaction with the FcRn (see, *e.g.*, PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464). The nucleotide sequence encoding the heavy-chain variable region, light-chain variable region, both the heavy-chain and light-chain variable regions, an epitope-binding
10 fragment of the heavy- and/or light-chain variable region, or one or more complementarity determining regions (CDRs) of an antibody may be cloned into such a vector for expression.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody having an increased affinity for the FcRn and an increased *in vivo* half-life. Thus, the
15 invention includes host cells containing a polynucleotide encoding an antibody, a constant domain or a FcRn binding fragment thereof having one or more modifications in amino acid residues 251-256, 285-290, 308-314, 385-389, and/or 428-436, preferably, operably linked to a heterologous promoter.

A variety of host-expression vector systems may be utilized to express the
20 antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention *in situ*. These include, but are not limited to, microorganisms such as bacteria (*e.g.*, *E. coli* and *B. subtilis*) transformed
25 with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (*e.g.*, *Saccharomyces* and *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (*e.g.*, baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression
30 vectors (*e.g.*, cauliflower mosaic virus, CaMV; and tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (*e.g.*, Ti plasmid) containing antibody coding sequences; and mammalian cell systems (*e.g.*, COS, CHO, BHK, 293, 3T3 and NSO cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian
35 viruses (*e.g.*, the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for

the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for
5 antibodies (Foecking *et al.*, *Gene*, 45:101, 1986, and Cockett *et al.*, *Bio/Technology*, 8:2, 1990).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For
10 example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther *et al.*, *EMBO*, 12:1791, 1983), in which the antibody coding sequence may be ligated individually
15 into the vector in frame with the lacZ coding region so that a fusion protein is produced; and pIN vectors (Inouye & Inouye, *Nucleic Acids Res.*, 13:3101-3109, 1985 and Van Heeke & Schuster, *J. Biol. Chem.*, 24:5503-5509, 1989).

In an insect system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera*
20 *frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

In mammalian host cells, a number of viral-based expression systems may be utilized to express an antibody molecule of the invention. In cases where an adenovirus is
25 used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the
30 antibody molecule in infected hosts (*e.g.*, see Logan & Shenk, *Proc. Natl. Acad. Sci. USA*, 81:355-359, 1984). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These
35 exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of

appropriate transcription enhancer elements, transcription terminators, etc. (see, *e.g.*, Bitter *et al.*, *Methods in Enzymol.*, 153:516-544, 1987).

In addition, a host cell strain may be chosen which modulates the expression
5 of the antibody sequences, or modifies and processes the antibody in the specific fashion
desired. Such modifications (*e.g.*, glycosylation) and processing (*e.g.*, cleavage) of protein
products may be important for the function of the antibody. Different host cells have
characteristic and specific mechanisms for the post-translational processing and modification
of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure
10 the correct modification and processing of the antibody expressed. To this end, eukaryotic
host cells which possess the cellular machinery for proper processing of the primary
transcript, glycosylation, and phosphorylation of the gene product may be used. Such
mammalian host cells include but are not limited to CHO, VERY, BHK, HeLa, COS,
MDCK, 293, 3T3, W138, and in particular, myeloma cells such as NS0 cells, and related cell
15 lines, see, for example, Morrison *et al.*, U.S. Patent No. 5,807,715, which is hereby
incorporated by reference in its entirety.

For long-term, high-yield production of recombinant antibodies, stable
expression is preferred. For example, cell lines which stably express the antibody molecule
may be engineered. Rather than using expression vectors which contain viral origins of
20 replication, host cells can be transformed with DNA controlled by appropriate expression
control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators,
polyadenylation sites, etc.), and a selectable marker. Following the introduction of the
foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media,
and then are switched to a selective media. The selectable marker in the recombinant
25 plasmid confers resistance to the selection and allows cells to stably integrate the plasmid
into their chromosomes and grow to form foci which in turn can be cloned and expanded into
cell lines. This method may advantageously be used to engineer cell lines which express the
antibody molecule. Such engineered cell lines may be particularly useful in screening and
evaluation of compositions that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to, the
30 herpes simplex virus thymidine kinase (Wigler *et al.*, *Cell*, 11:223, 1977), hypoxanthine-
guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA*,
48:202, 1992), and adenine phosphoribosyltransferase (Lowy *et al.*, *Cell*, 22:8-17, 1980)
genes can be employed in tk⁻, hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite
35 resistance can be used as the basis of selection for the following genes: *dhfr*, which confers
resistance to methotrexate (Wigler *et al.*, *Natl. Acad. Sci. USA*, 77:357, 1980 and O'Hare *et*

al., *Proc. Natl. Acad. Sci. USA*, 78:1527, 1981); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA*, 78:2072, 1981); *neo*, which confers resistance to the aminoglycoside G-418 (Wu and Wu, *Biotherapy*, 3:87-95, 1991; Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.*, 32:573-596, 1993; Mulligan, *Science*, 260:926-932, 1993; and Morgan and Anderson, *Ann. Rev. Biochem.*, 62: 191-217, 1993; and May, *TIB TECH*, 11(5):155-2 15, 1993); and *hygro*, which confers resistance to hygromycin (Santerre *et al.*, *Gene*, 30:147, 1984). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel *et al.* (eds.), 1993, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY; Krieger, 1990, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY; in Chapters 12 and 13, Dracopoli *et al.* (eds), 1994, *Current Protocols in Human Genetics*, John Wiley & Sons, NY; and Colberre-Garapin *et al.*, *J. Mol. Biol.*, 150:1, 1981, which are incorporated by reference herein in their entireties.

The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, 1987, *The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning*, Vol.3. Academic Press, New York). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse *et al.*, *Mol., Cell. Biol.*, 3:257, 1983).

The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides or different selectable markers to ensure maintenance of both plasmids. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, *Nature*, 322:52, 1986; and Kohler, *Proc. Natl. Acad. Sci. USA*, 77:2 197, 1980). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by recombinant expression, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (*e.g.*, ion exchange,

affinity, particularly by affinity for the specific antigen after Protein A purification, and sizing column chromatography), centrifugation, differential solubility, or by any other standard techniques for the purification of proteins. Further, the antibodies of the present invention or fragments thereof may be fused to heterologous polypeptide sequences described herein or otherwise known in the art to facilitate purification.

5.3.1. ANTIBODY CONJUGATES

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to heterologous polypeptides (*i.e.*, an unrelated polypeptide; or portion thereof, preferably at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids of the polypeptide) to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. Antibodies fused or conjugated to heterologous polypeptides may also be used in *in vitro* immunoassays and purification methods using methods known in the art. See *e.g.*, PCT Publication No. WO 93/21232; EP 439,095; Naramura *et al.*, *Immunol. Lett.*, 39:91-99, 1994; U.S. Patent 5,474,981; Gillies *et al.*, *PNAS*, 89:1428-1432, 1992; and Fell *et al.*, *J. Immunol.*, 146:2446-2452, 1991, which are incorporated herein by reference in their entirety.

Antibodies can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA*, 86:821-824, 1989, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, *Cell*, 37:767 1984) and the "flag" tag (Knappik *et al.*, *Biotechniques*, 17(4):754-761, 1994).

The present invention also encompasses antibodies conjugated to a diagnostic or therapeutic agent or any other molecule for which *in vivo* half-life is desired to be increased. The antibodies can be used diagnostically to, for example, monitor the development or progression of a disease, disorder or infection as part of a clinical testing procedure to, *e.g.*, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent

materials, bioluminescent materials, radioactive materials, positron emitting metals, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99m}Tc .

An antibody may be conjugated to a therapeutic moiety such as a cytotoxin (*e.g.*, a cytostatic or cytotoxic agent), a therapeutic agent or a radioactive element (*e.g.*, alpha-emitters, gamma-emitters, etc.). Cytotoxins or cytotoxic agents include any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (*e.g.*, vincristine and vinblastine).

Further, an antibody may be conjugated to a therapeutic agent or drug moiety that modifies a given biological response. Therapeutic agents or drug moieties are not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon (IFN- α), β -interferon (IFN- β),

nerve growth factor (NGF), platelet derived growth factor (PDGF), tissue plasminogen activator (TPA), an apoptotic agent (e.g., TNF- α , TNF- β , AIM I as disclosed in PCT Publication No. WO 97/33899), AIM II (see, PCT Publication No. WO 97/34911), Fas
5 Ligand (Takahashi *et al.*, *J. Immunol.*, 6:1567-1574, 1994), and VEGI (PCT Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent (e.g., angiostatin or endostatin); or a biological response modifier such as, for example, a lymphokine (e.g., interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte
macrophage colony stimulating factor ("GM-CSF"), and granulocyte colony stimulating
10 factor ("G-CSF")), or a growth factor (e.g., growth hormone ("GH")).

Techniques for conjugating such therapeutic moieties to antibodies are well known; see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), 1985, pp. 243-56, Alan R. Liss, Inc.); Hellstrom *et al.*, "Antibodies For Drug Delivery", in
15 *Controlled Drug Delivery (2nd Ed.)*, Robinson *et al.* (eds.), 1987, pp. 623-53, Marcel Dekker, Inc.); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), 1985, pp. 475-506; "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies
20 For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), 1985, pp. 303-16, Academic Press; and Thorpe *et al.*, *Immunol. Recombinant expression vector.*, 62:119-58, 1982.

An antibody or fragment thereof, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

25 Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include,
30 but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

5.4 METHODS OF PRODUCING FUSION PROTEINS

Fusion proteins can be produced by standard recombinant DNA techniques or
35 by protein synthetic techniques, e.g., by use of a peptide synthesizer. For example, a nucleic acid molecule encoding a fusion protein can be synthesized by conventional techniques

including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and
5 reamplified to generate a chimeric gene sequence (*see, e.g., Current Protocols in Molecular Biology*, Ausubel *et al.*, eds., John Wiley & Sons, 1992). Moreover, a nucleic acid encoding a bioactive molecule can be cloned into an expression vector containing the Fc domain or a fragment thereof such that the bioactive molecule is linked in-frame to the constant domain or fragment thereof.

10 Methods for fusing or conjugating polypeptides to the constant regions of antibodies are known in the art. See, *e.g.*, U.S. Patent Nos. 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, 5,723,125, 5,783,181, 5,908,626, 5,844,095, and 5,112,946; EP 307,434; EP 367,166; EP 394,827; PCT publications WO 91/06570, WO 96/04388, WO 96/22024, WO 97/34631, and WO 99/04813; Ashkenazi *et al.*, *Proc. Natl. Acad. Sci. USA*,
15 88: 10535-10539, 1991; Traunecker *et al.*, *Nature*, 331:84-86, 1988; Zheng *et al.*, *J. Immunol.*, 154:5590-5600, 1995; and Vil *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:11337-11341, 1992, which are incorporated herein by reference in their entireties.

The nucleotide sequence encoding a bioactive molecule may be obtained from any information available to those of skill in the art (*e.g.*, from Genbank, the literature, or by
20 routine cloning), and the nucleotide sequence encoding a constant domain or a fragment thereof with increased affinity for the FcRn may be determined by sequence analysis of mutants produced using techniques described herein, or may be obtained from Genbank or the literature. The nucleotide sequence coding for a fusion protein can be inserted into an appropriate expression vector, *i.e.*, a vector which contains the necessary elements for the
25 transcription and translation of the inserted protein-coding sequence. A variety of host-vector systems may be utilized in the present invention to express the protein-coding sequence. These include but are not limited to mammalian cell systems infected with virus (*e.g.*, vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (*e.g.*, baculovirus); microorganisms such as yeast containing yeast vectors; or bacteria transformed
30 with bacteriophage, DNA, plasmid DNA, or cosmid DNA. The expression elements of vectors vary in their strengths and specificities. Depending on the host-vector system utilized, any one of a number of suitable transcription and translation elements may be used.

The expression of a fusion protein may be controlled by any promoter or enhancer element known in the art. Promoters which may be used to control the expression
35 of the gene encoding fusion protein include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature*, 290:304-310, 1981), the promoter contained in the 3'

long terminal repeat of Rous sarcoma virus (Yamamoto, *et al.*, *Cell*, 22:787-797, 1980), the herpes thymidine kinase promoter (Wagner *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 78:1441-1445, 1981), the regulatory sequences of the metallothionein gene (Brinster *et al.*, *Nature*, 296:39-42, 1982), the tetracycline (Tet) promoter (Gossen *et al.*, *Proc. Nat. Acad. Sci. USA*, 89:5547-5551, 1995); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Kamaroff, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 75:3727-3731, 1978), or the *tac* promoter (DeBoer, *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 80:21-25, 1983; see also "Useful proteins from recombinant bacteria" in *Scientific American*, 242:74-94, 1980); plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella *et al.*, *Nature*, 303:209-213, 1983) or the cauliflower mosaic virus 35S RNA promoter (Gardner, *et al.*, *Nucl. Acids Res.*, 9:2871, 1981), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella *et al.*, *Nature*, 310:115-120, 1984); promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift *et al.*, *Cell* 38:639-646, 1984; Ornitz *et al.*, 50:399-409, *Cold Spring Harbor Symp. Quant. Biol.*, 1986; MacDonald, *Hepatology* 7:425-515, 1987); insulin gene control region which is active in pancreatic beta cells (Hanahan, *Nature* 315:115-122, 1985), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl *et al.*, *Cell*, 38:647-658, 1984; Adames *et al.*, *Nature* 318:533-538, 1985; Alexander *et al.*, *Mol. Cell. Biol.*, 7:1436-1444, 1987), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder *et al.*, *Cell*, 45:485-495, 1986), albumin gene control region which is active in liver (Pinkert *et al.*, *Genes and Devel.*, 1:268-276, 1987), α -fetoprotein gene control region which is active in liver (Krumlauf *et al.*, *Mol. Cell. Biol.*, 5:1639-1648, 1985; Hammer *et al.*, *Science*, 235:53-58, 1987; α 1-antitrypsin gene control region which is active in the liver (Kelsey *et al.*, *Genes and Devel.*, 1:161-171, 1987), beta-globin gene control region which is active in myeloid cells (Mogram *et al.*, *Nature*, 315:338-340, 1985; Kollias *et al.*, *Cell*, 46:89-94, 1986; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead *et al.*, *Cell*, 48:703-712, 1987); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, *Nature*, 314:283-286, 1985); neuronal-specific enolase (NSE) which is active in neuronal cells (Morelli *et al.*, *Gen. Virol.*, 80:571-83, 1999); brain-derived neurotrophic factor (BDNF) gene control region which is active in neuronal cells (Tabuchi *et al.*, *Biochem. Biophysic. Res. Comprising.*, 253:818-823,

1998); glial fibrillary acidic protein (GFAP) promoter which is active in astrocytes (Gomes *et al.*, Braz. J. Med. Biol. Res., 32(5):619-631, 1999; Morelli *et al.*, *Gen. Virol.*, 80:571-83, 1999) and gonadotropic releasing hormone gene control region which is active in the
5 hypothalamus (Mason *et al.*, *Science*, 234:1372-1378, 1986).

In a specific embodiment, the expression of a fusion protein is regulated by a constitutive promoter. In another embodiment, the expression of a fusion protein is regulated by an inducible promoter. In accordance with these embodiments, the promoter may be a tissue-specific promoter.

10 In a specific embodiment, a vector is used that comprises a promoter operably linked to a fusion protein-encoding nucleic acid, one or more origins of replication, and, optionally, one or more selectable markers (*e.g.*, an antibiotic resistance gene).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the fusion protein
15 coding sequence may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts (*e.g.*,
20 see Logan & Shenk, *Proc. Natl. Acad. Sci. USA*, 81:355-359, 1984). Specific initiation signals may also be required for efficient translation of inserted fusion protein coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational
25 control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bitter *et al.*, *Methods in Enzymol.*, 153:516-544, 1987).

Expression vectors containing inserts of a gene encoding a fusion protein can
30 be identified by three general approaches: (a) nucleic acid hybridization, (b) presence or absence of "marker" gene functions, and (c) expression of inserted sequences. In the first approach, the presence of a gene encoding a fusion protein in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted gene encoding the fusion protein. In the second approach, the
35 recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (*e.g.*, thymidine kinase activity, resistance to

antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of a nucleotide sequence encoding a fusion protein in the vector. For example, if the nucleotide sequence encoding the fusion protein is inserted within the marker
 5 gene sequence of the vector, recombinants containing the gene encoding the fusion protein insert can be identified by the absence of the marker gene function. In the third approach, recombinant expression vectors can be identified by assaying the gene product (*i.e.*, fusion protein) expressed by the recombinant. Such assays can be based, for example, on the physical or functional properties of the fusion protein in *in vitro* assay systems, *e.g.*, binding
 10 with anti-bioactive molecule antibody.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus, expression of the genetically engineered fusion protein may be controlled.
 15 Furthermore, different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (*e.g.*, glycosylation, phosphorylation of proteins). Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system will produce an unglycosylated product and expression in
 20 yeast will produce a glycosylated product. Eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular, neuronal cell lines such as, for example, SK-N-AS, SK-N-FI, SK-N-DZ human neuroblastomas (Sugimoto
 25 *et al.*, *J. Natl. Cancer Inst.*, 73: 51-57, 1984), SK-N-SH human neuroblastoma (*Biochim. Biophys. Acta*, 704: 450-460, 1982), Daoy human cerebellar medulloblastoma (He *et al.*, *Cancer Res.*, 52: 1144-1148, 1992) DBTRG-05MG glioblastoma cells (Kruse *et al.*, 1992, *In Vitro Cell. Dev. Biol.*, 28A:609-614, 1992), IMR-32 human neuroblastoma (*Cancer Res.*, 30: 2110-2118, 1970), 1321N1 human astrocytoma (*Proc. Natl Acad. Sci. USA*, 74: 4816, 1997),
 30 MOG-G-CCM human astrocytoma (*Br. J. Cancer*, 49: 269, 1984), U87MG human glioblastoma-astrocytoma (*Acta Pathol. Microbiol. Scand.*, 74: 465-486, 1968), A172 human glioblastoma (Olopade *et al.*, *Cancer Res.*, 52: 2523-2529, 1992), C6 rat glioma cells (Benda *et al.*, *Science*, 161: 370-371, 1968), Neuro-2a mouse neuroblastoma (*Proc. Natl. Acad. Sci. USA*, 65: 129-136, 1970), NB41A3 mouse neuroblastoma (*Proc. Natl. Acad. Sci. USA*, 48: 1184-1190, 1962), SCP sheep choroid plexus (Bolin *et al.*, *J. Virol. Methods*, 48: 211-221,
 35 1994), G355-5, PG-4 Cat normal astrocyte (Haapala *et al.*, *J. Virol.*, 53: 827-833, 1985), Mpf

ferret brain (Trowbridge *et al.*, *In Vitro*, 18: 952-960, 1982), and normal cell lines such as, for example, CTX TNA2 rat normal cortex brain (Radany *et al.*, *Proc. Natl. Acad. Sci. USA*, 89: 6467-6471, 1992) such as, for example, CRL7030 and Hs578Bst. Furthermore, different
5 vector/host expression systems may effect processing reactions to different degrees.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the fusion protein may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression
10 control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched medium, and then are switched to a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid
15 into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the differentially expressed or pathway gene protein. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the differentially expressed or pathway gene protein.

A number of selection systems may be used, including but not limited to the
20 herpes simplex virus thymidine kinase (Wigler, *et al.*, *Cell*, 11:223, 1997), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA*, 48:2026, 1962), and adenine phosphoribosyltransferase (Lowy, *et al.*, 1980, *Cell*, 22:817, 1980) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also,
25 antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler, *et al.*, *Natl. Acad. Sci. USA*, 77:3567, 1980; O'Hare, *et al.*, *Proc. Natl. Acad. Sci. USA*, 78:1527, 1981); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA*, 78:2072, 1981); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin, *et al.*, *J. Mol. Biol.*, 150:1, 1981);
30 and hyg^r, which confers resistance to hygromycin (Santerre, *et al.*, *Gene*, 30:147, 1984) genes.

Once a fusion protein of the invention has been produced by recombinant expression, it may be purified by any method known in the art for purification of a protein, for example, by chromatography (*e.g.*, ion exchange, affinity, particularly by affinity for the
35 specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins.

5.5. PROPHYLACTIC AND THERAPEUTIC USES OF ANTIBODIES

The present invention encompasses antibody-based therapies which involve
5 administering antibodies to an animal, preferably a mammal, and most preferably a human, for preventing, treating, or ameliorating symptoms associated with a disease, disorder, or infection. Prophylactic and therapeutic compounds of the invention include, but are not limited to, antibodies and nucleic acids encoding antibodies. Antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

10 Antibodies of the present invention that function as antagonists of a disease, disorder, or infection can be administered to an animal, preferably a mammal and most preferably a human, to treat, prevent or ameliorate one or more symptoms associated with the disease, disorder, or infection. For example, antibodies which disrupt or prevent the interaction between a viral antigen and its host cell receptor may be administered to an
15 animal, preferably a mammal and most preferably a human, to treat, prevent or ameliorate one or more symptoms associated with a viral infection.

In a specific embodiment, an antibody or fragment thereof prevents a viral or bacterial antigen from binding to its host cell receptor by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at
20 least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to antigen binding to its host cell receptor in the absence of said antibodies. In another embodiment, a combination of antibodies prevent a viral or bacterial antigen from binding to its host cell receptor by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%,
25 at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to antigen binding to its host cell receptor in the absence of said antibodies. In a preferred embodiment, the antibody is used to treat or prevent RSV infection

Antibodies which do not prevent a viral or bacterial antigen from binding its host cell receptor but inhibit or downregulate viral or bacterial replication can also be
30 administered to an animal to treat, prevent or ameliorate one or more symptoms associated with a viral or bacterial infection. The ability of an antibody to inhibit or downregulate viral or bacterial replication may be determined by techniques described herein or otherwise known in the art. For example, the inhibition or downregulation of viral replication can be determined by detecting the viral titer in the animal.

35 In a specific embodiment, an antibody inhibits or downregulates viral or bacterial replication by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at

least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to viral or bacterial replication in absence of said antibody. In another embodiment, a combination of antibodies inhibit or downregulate viral or bacterial replication by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to viral or bacterial replication in absence of said antibodies.

Antibodies can also be used to prevent, inhibit or reduce the growth or metastasis of cancerous cells. In a specific embodiment, an antibody inhibits or reduces the growth or metastasis of cancerous cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the growth or metastasis in absence of said antibody. In another embodiment, a combination of antibodies inhibits or reduces the growth or metastasis of cancer by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the growth or metastasis in absence of said antibodies. Examples of cancers include, but are not limited to, leukemia (*e.g.*, acute leukemia such as acute lymphocytic leukemia and acute myelocytic leukemia), neoplasms, tumors (*e.g.*, fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma), heavy chain disease, metastases, or any disease or disorder characterized by uncontrolled cell growth.

Antibodies can also be used to reduce the inflammation experienced by animals, particularly mammals, with inflammatory disorders. In a specific embodiment, an antibody reduces the inflammation in an animal by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%,
5 at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the inflammation in an animal in the not administered said antibody. In another embodiment, a combination of antibodies reduce the inflammation in an animal by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%,
10 at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the inflammation in an animal in not administered said antibodies. Examples of inflammatory disorders include, but are not limited to, rheumatoid arthritis, spondyloarthropathies, inflammatory bowel disease and asthma.

15 In certain embodiments, the antibody used for treatment of inflammation (or cancer) is a modified anti- $\alpha_v\beta_3$ antibody, preferably a Vitaxin antibody (see, PCT publications WO 98/33919 and WO 00/78815, both by Huse *et al.*, and both of which are incorporated by reference herein in their entireties).

Antibodies can also be used to prevent the rejection of transplants.
20 Antibodies can also be used to prevent clot formation. Further, antibodies that function as agonists of the immune response can also be administered to an animal, preferably a mammal, and most preferably a human, to treat, prevent or ameliorate one or more symptoms associated with the disease, disorder, or infection.

One or more antibodies that immunospecifically bind to one or more antigens
25 may be used locally or systemically in the body as a therapeutic. The antibodies of this invention may also be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, *e.g.*, IL-2, IL-3 and IL-7), which, for example, serve to increase the number or activity of effector cells which interact with the antibodies. The antibodies of this invention may also be
30 advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, *e.g.*, IL-2, IL-3 and IL-7), which, for example, serve to increase the immune response. The antibodies of this invention may also be advantageously utilized in combination with one or more drugs used to treat a disease, disorder, or infection such as, for example anti-cancer agents, anti-inflammatory
35 agents or anti-viral agents. Examples of anti-cancer agents include, but are not limited to, isplatin, ifosfamide, paclitaxel, taxanes, topoisomerase I inhibitors (*e.g.*, CPT-11, topotecan,

9-AC, and GG-211), gemcitabine, vinorelbine, oxaliplatin, 5-fluorouracil (5-FU), leucovorin, vinorelbine, temodal, and taxol. Examples of anti-viral agents include, but are not limited to, cytokines (*e.g.*, IFN- α , IFN- β , IFN- γ), inhibitors of reverse transcriptase (*e.g.*, AZT, 3TC, D4T, ddC, ddI, d4T, 3TC, adefovir, efavirenz, delavirdine, nevirapine, abacavir, and other dideoxynucleosides or dideoxyfluoronucleosides), inhibitors of viral mRNA capping, such as ribavirin, inhibitors of proteases such HIV protease inhibitors (*e.g.*, amprenavir, indinavir, nelfinavir, ritonavir, and saquinavir,), amphotericin B, castanospermine as an inhibitor of glycoprotein processing, inhibitors of neuraminidase such as influenza virus neuraminidase inhibitors (*e.g.*, zanamivir and oseltamivir), topoisomerase I inhibitors (*e.g.*, camptothecins and analogs thereof), amantadine, and rimantadine. Examples of anti-inflammatory agents include, but are not limited to, nonsteroidal anti-inflammatory drugs such as COX-2 inhibitors (*e.g.*, meloxicam, celecoxib, rofecoxib, flosulide, and SC-58635, and MK-966), ibuprofen and indomethacin, and steroids (*e.g.*, deflazacort, dexamethasone and methylprednisolone).

In a specific embodiment, antibodies administered to an animal are of a species origin or species reactivity that is the same species as that of the animal. Thus, in a preferred embodiment, human or humanized antibodies, or nucleic acids encoding human or human, are administered to a human patient for therapy or prophylaxis.

In preferred embodiments, immunoglobulins having extended *in vivo* half-lives are used in passive immunotherapy (for either therapy or prophylaxis). Because of the extended half-life, passive immunotherapy or prophylaxis can be accomplished using lower doses and/or less frequent administration of the therapeutic resulting in fewer side effects, better patient compliance, less costly therapy/prophylaxis, etc. In a preferred embodiment, the therapeutic/prophylactic is an antibody that binds RSV, for example, SYNAGIS® or other anti-RSV antibody. Such anti-RSV antibodies, and methods of administration are disclosed in U.S. patent application Serial Nos. 09/724,396 and 09/724,531, both entitled "Methods of Administering/Dosing Anti-RSV Antibodies For Prophylaxis and Treatment," both by Young *et al.*, both filed November 28, 2000, and continuation-in-part applications of these applications, Serial Nos. _____ and _____, respectively (attorney docket Nos. 10271-048 and 10271-047, respectively), also entitled "Methods of Administering/Dosing Anti-RSV Antibodies for Prophylaxis and Treatment," by Young *et al.*, all which are incorporated by reference herein in their entireties. Also included are the anti-RSV antibodies described in Section 5.1, *supra*.

In a specific embodiment, fusion proteins administered to an animal are of a species origin or species reactivity that is the same species as that of the animal. Thus, in a

preferred embodiment, human fusion proteins or nucleic acids encoding human fusion proteins, are administered to a human subject for therapy or prophylaxis.

5

5.6. PROPHYLACTIC AND THERAPEUTIC USES OF FUSION PROTEINS AND CONJUGATED MOLECULES

The present invention encompasses fusion protein-based and conjugated molecule-based therapies which involve administering fusion proteins or conjugated molecules to an animal, preferably a mammal and most preferably a human, for preventing, treating, or ameliorating symptoms associated with a disease, disorder, or infection. Prophylactic and therapeutic compounds of the invention include, but are not limited to, fusion proteins and nucleic acids encoding fusion proteins and conjugated molecules. Fusion proteins and conjugated molecules may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

Fusion proteins and conjugated molecules of the present invention that function as antagonists of a disease, disorder, or infection can be administered to an animal, preferably a mammal, and most preferably a human, to treat, prevent or ameliorate one or more symptoms associated with the disease, disorder, or infection. Further, fusion proteins and conjugated molecules of the present invention that function as agonists of the immune response may be administered to an animal, preferably a mammal, and most preferably a human, to treat, prevent or ameliorate one or more symptoms associated with the disease, disorder, or infection.

One or more fusion proteins and conjugated molecules may be used locally or systemically in the body as a therapeutic. The fusion proteins and conjugated molecules of this invention may also be advantageously utilized in combination with monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, *e.g.*, IL-2, IL-3 and IL-7), which, for example, serve to increase the number or activity of effector cells which interact with the antibodies. The fusion proteins and conjugated molecules of this invention may also be advantageously utilized in combination with monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, *e.g.*, IL-2, IL-3 and IL-7), which, for example, serve to increase the immune response. The fusion proteins and conjugated molecules of this invention may also be advantageously utilized in combination with one or more drugs used to treat a disease, disorder, or infection such as, for example anti-cancer agents, anti-inflammatory agents or anti-viral agents. Examples of anti-cancer agents include, but are not limited to, isplatin, ifosfamide, paclitaxel, taxanes,

topoisomerase I inhibitors (*e.g.*, CPT-11, topotecan, 9-AC, and GG-211), gemcitabine, vinorelbine, oxaliplatin, 5-fluorouracil (5-FU), leucovorin, vinorelbine, temodal, and taxol. Examples of anti-viral agents include, but are not limited to, cytokines (*e.g.*, IFN- α , IFN- β , IFN- γ), inhibitors of reverse transcriptase (*e.g.*, AZT, 3TC, D4T, ddC, ddI, d4T, 3TC, adefovir, efavirenz, delavirdine, nevirapine, abacavir, and other dideoxynucleosides or dideoxyfluoronucleosides), inhibitors of viral mRNA capping, such as ribavirin, inhibitors of proteases such as HIV protease inhibitors (*e.g.*, amprenavir, indinavir, nelfinavir, ritonavir, and saquinavir), amphotericin B, castanospermine as an inhibitor of glycoprotein processing, inhibitors of neuraminidase such as influenza virus neuraminidase inhibitors (*e.g.*, zanamivir and oseltamivir), topoisomerase I inhibitors (*e.g.*, camptothecins and analogs thereof), amantadine, and rimantadine. Examples of anti-inflammatory agents include, but are not limited to, nonsteroidal anti-inflammatory drugs such as COX-2 inhibitors (*e.g.*, meloxicam, celecoxib, rofecoxib, flosulide, and SC-58635, and MK-966), ibuprofen and indomethacin, and steroids (*e.g.*, deflazacort, dexamethasone and methylprednisolone).

5.7. ADMINISTRATION OF ANTIBODIES OR FUSION PROTEINS

The invention provides methods of treatment, prophylaxis, and amelioration of one or more symptoms associated with a disease, disorder or infection by administering to a subject of an effective amount of an antibody of the invention, or pharmaceutical composition comprising an antibody of the invention. The invention also provides methods of treatment, prophylaxis, and amelioration of one or more symptoms associated with a disease, disorder or infection by administering to a subject an effective amount of a fusion protein or conjugated molecule of the invention, or a pharmaceutical composition comprising a fusion protein or conjugated molecules of the invention. In a preferred aspect, an antibody or fusion protein or conjugated molecule, is substantially purified (*i.e.*, substantially free from substances that limit its effect or produce undesired side-effects). In a specific embodiment, the subject is an animal, preferably a mammal such as non-primate (*e.g.*, cows, pigs, horses, cats, dogs, rats etc.) and a primate (*e.g.*, monkey such as a cynomolgous monkey and a human). In a preferred embodiment, the subject is a human.

Various delivery systems are known and can be used to administer an antibody or fusion protein or conjugated molecule of the invention, *e.g.*, encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the

antibody or fusion protein, receptor-mediated endocytosis (see, *e.g.*, Wu and Wu, *J. Biol. Chem.*, 262:4429-4432, 1987), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of administering an antibody, a fusion protein or conjugated molecule, or pharmaceutical composition include, but are not limited to, parenteral administration (*e.g.*, intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidural, and mucosal (*e.g.*, intranasal and oral routes). In a specific embodiment, antibodies, fusion proteins, conjugated molecules, or pharmaceutical compositions are administered intramuscularly, intravenously, or subcutaneously. The compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (*e.g.*, oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, pulmonary administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent. See, *e.g.*, U.S. Patent Nos. 6,019,968; 5,985,320; 5,985,309; 5,934,272; 5,874,064; 5,855,913; 5,290,540; and 4,880,078; and PCT Publication Nos. WO 92/19244; WO 97/32572; WO 97/44013; WO 98/31346; and WO 99/66903, each of which is incorporated herein by reference in its entirety. In a preferred embodiment, an antibody, a fusion protein, conjugated molecules, or a pharmaceutical composition is administered using Alkermes AIR™ pulmonary drug delivery technology (Alkermes, Inc., Cambridge, MA).

The invention also provides that an antibody, a fusion protein, or conjugated molecule is packaged in a hermetically sealed container such as an ampoule or sachette indicating the quantity of antibody, fusion protein, or conjugated molecule. In one embodiment, the antibody, fusion protein, or conjugated molecule is supplied as a dry sterilized lyophilized powder or water free concentrate in a hermetically sealed container and can be reconstituted, *e.g.*, with water or saline to the appropriate concentration for administration to a subject. Preferably, the antibody, fusion protein, or conjugated molecule is supplied as a dry sterile lyophilized powder in a hermetically sealed container at a unit dosage of at least 5 mg, more preferably at least 10 mg, at least 15 mg, at least 25 mg, at least 35 mg, at least 45 mg, at least 50 mg, or at least 75 mg. The lyophilized antibody, fusion protein, or conjugated molecule should be stored at between 2 and 8°C in its original container and the antibody, fusion protein, or conjugated molecules should be administered within 12 hours, preferably within 6 hours, within 5 hours, within 3 hours, or within 1 hour after being reconstituted. In an alternative embodiment, an antibody, fusion protein, or conjugated molecule is supplied in liquid form in a hermetically sealed container indicating the quantity and concentration of the antibody, fusion protein, or conjugated molecule.

Preferably, the liquid form of the antibody, fusion protein, or conjugated molecule is supplied in a hermetically sealed container at least 1 mg/ml, more preferably at least 2.5 mg/ml, at least 5 mg/ml, at least 8 mg/ml, at least 10 mg/ml, at least 15 mg/kg, or at least 25 mg/ml.

5 In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion, by injection, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when
10 administering an antibody or a fusion protein, care must be taken to use materials to which the antibody or the fusion protein does not absorb.

In another embodiment, the composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science*, 249:1527-1533, 1990; Treat *et al.*, in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 3 17-327; see generally *ibid.*).
15

In yet another embodiment, the composition can be delivered in a controlled release or sustained release system. Any technique known to one of skill in the art can be used to produce sustained release formulations comprising one or more antibodies, or one or more fusion proteins. See, *e.g.*, U.S. Patent No. 4,526,938; PCT publication WO 91/05548; PCT publication WO 96/20698; Ning *et al.*, "Intratumoral Radioimmunotherapy of a Human Colon Cancer Xenograft Using a Sustained-Release Gel," *Radiotherapy & Oncology*, 39:179-189, 1996; Song *et al.*, "Antibody Mediated Lung Targeting of Long-Circulating Emulsions," *PDA Journal of Pharmaceutical Science & Technology*, 50:372-397, 1995; Cleek *et al.*, "Biodegradable Polymeric Carriers for a bFGF Antibody for Cardiovascular
25 Application," *Pro. Intl. Symp. Control. Rel. Bioact. Mater.*, 24:853-854, 1997; and Lam *et al.*, "Microencapsulation of Recombinant Humanized Monoclonal Antibody for Local Delivery," *Proc. Int'l. Symp. Control Rel. Bioact. Mater.*, 24:759-760, 1997, each of which is incorporated herein by reference in its entirety. In one embodiment, a pump may be used in a controlled release system (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.*, 14:20, 1987; Buchwald *et al.*, *Surgery*, 88:507, 1980; and Saudek *et al.*, *N. Engl. J. Med.*, 321:574, 1989). In another embodiment, polymeric materials can be used to achieve controlled release of antibodies or fusion proteins (see *e.g.*, *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New
35 York (1984); Ranger and Peppas, *J. Macromol. Sci. Rev. Macromol. Chem.*, 23:61, 1983; see also Levy *et al.*, *Science*, 228:190, 1985; During *et al.*, *Ann. Neurol.*, 25:351, 1989;

Howard *et al.*, *J. Neurosurg.*, 71:105, 1989); U.S. Patent No. 5,679,377; U.S. Patent No. 5,916,597; U.S. Patent No. 5,912,015; U.S. Patent No. 5,989,463; U.S. Patent No. 5,128,326; PCT Publication No. WO 99/15154; and PCT Publication No. WO 99/20253). In yet
5 another embodiment, a controlled release system can be placed in proximity of the therapeutic target (*e.g.*, the lungs), thus requiring only a fraction of the systemic dose (see, *e.g.*, Goodson, in *Medical Applications of Controlled Release*, supra, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer,
10 *Science*, 249:1527-1533, 1990).

In a specific embodiment where the composition of the invention is a nucleic acid encoding an antibody or fusion protein, the nucleic acid can be administered *in vivo* to promote expression of its encoded antibody or fusion protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes
15 intracellular, *e.g.*, by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see *e.g.*, Joliot *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:1864-1868, 1991), etc. Alternatively, a nucleic acid can
20 be introduced intracellularly and incorporated within host cell DNA for expression by homologous recombination.

The present invention also provides pharmaceutical compositions. Such compositions comprise a prophylactically or therapeutically effective amount of an antibody, fusion protein or conjugated molecule, and a pharmaceutically acceptable carrier. In a
25 specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant (*e.g.*, Freund's complete and incomplete, mineral gels such as aluminum hydroxide, surface active substances such as
30 lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful adjuvants for humans such as BCG (Bacille Calmette-Guerin) and *Corynebacterium parvum*), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as
35 peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and

aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a prophylactically or therapeutically effective amount of the antibody or fragment thereof, or fusion protein or conjugated molecule, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection.

Generally, the ingredients of compositions of the invention are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the composition of the invention which will be effective in the treatment, prevention or amelioration of one or more symptoms associated with a disease, disorder, or infection can be determined by standard clinical techniques. The precise dose to be employed in the formulation will depend on the route of administration, the age of the subject, and the seriousness of the disease, disorder, or infection, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model (e.g., the cotton rat or Cynomolgous monkey) test systems.

For fusion proteins, the therapeutically or prophylactically effective dosage administered to a subject ranges from about 0.001 to 50 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. For antibodies, the therapeutically or prophylactically effective dosage administered to a subject is typically 0.1 mg/kg to 200 mg/kg of the subject's body weight. Preferably, the dosage administered to a subject is between 0.1 mg/kg and 20 mg/kg of the subject's body weight and more preferably the dosage administered to a subject is between 1 mg/kg to 10 mg/kg of the subject's body weight. The dosage will, however, depend upon the extent to which the *in vivo* half-life of the molecule has been increased. Generally, human antibodies and human fusion proteins have longer half-lives within the human body than antibodies of fusion proteins from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies or human fusion proteins and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies, fusion proteins, or conjugated molecules may be reduced also by enhancing uptake and tissue penetration (e.g., into the lung) of the antibodies or fusion proteins by modifications such as, for example, lipidation.

Treatment of a subject with a therapeutically or prophylactically effective amount of an antibody, fusion protein, or conjugated molecule can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with an antibody, fusion protein, or conjugated molecule in the range of between about 0.1 to 30 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. In other embodiments, the pharmaceutical composition of the invention is administered once a day, twice a day, or three times a day. In other embodiments, the pharmaceutical composition is administered once a week, twice a week, once every two weeks, once a month, once every six weeks, once every two months, twice a

year or once per year. It will also be appreciated that the effective dosage of the antibody, fusion protein, or conjugated molecule used for treatment may increase or decrease over the course of a particular treatment.

5

5.7.1. GENE THERAPY

In a specific embodiment, nucleic acids comprising sequences encoding antibodies or fusion proteins, are administered to treat, prevent or ameliorate one or more symptoms associated with a disease, disorder, or infection, by way of gene therapy. Gene
10 therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded antibody or fusion protein that mediates a therapeutic or prophylactic effect.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

15 For general reviews of the methods of gene therapy, see Goldspiel *et al.*, *Clinical Pharmacy*, 12:488-505, 1993; Wu and Wu, *Biotherapy*, 3:87-95, 1991; Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.*, 32:573-596, 1993; Mulligan, *Science*, 260:926-932, 1993; and Morgan and Anderson, *Ann. Rev. biochem.* 62:191-217, 1993; *TIBTECH* 11(5):155-215, 1993. Methods commonly known in the art of recombinant DNA technology which can be
20 used are described in Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression*, A Laboratory Manual, Stockton Press, NY (1990).

In a preferred aspect, a composition of the invention comprises nucleic acids encoding an antibody, said nucleic acids being part of an expression vector that expresses the
25 antibody in a suitable host. In particular, such nucleic acids have promoters, preferably heterologous promoters, operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a
30 desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, *Proc. Natl. Acad. Sci. USA*, 86:8932-8935, 1989; and Zijlstra *et al.*, *Nature*, 342:435-438, 1989).

In another preferred aspect, a composition of the invention comprises nucleic acids encoding a fusion protein, said nucleic acids being a part of an expression vector that
35 expression the fusion protein in a suitable host. In particular, such nucleic acids have promoters, preferably heterologous promoters, operably linked to the coding region of a

fusion protein, said promoter being inducible or constitutive, and optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the coding sequence of the fusion protein and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for
5 intrachromosomal expression of the fusion protein encoding nucleic acids.

Delivery of the nucleic acids into a subject may be either direct, in which case the subject is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids *in vitro*, then transplanted
10 into the subject. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, *e.g.*, by constructing them as part of an
15 appropriate nucleic acid expression vector and administering it so that they become intracellular, *e.g.*, by infection using defective or attenuated retroviral or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or
20 microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, *e.g.*, Wu and Wu, *J. Biol. Chem.*, 262:4429-4432, 1987) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to
25 disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, *e.g.*, PCT Publications WO 92/06180; WO 92/22635; WO 92/20316; WO 93/14188; WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by
30 homologous recombination (Koller and Smithies, *Proc. Natl. Acad. Sci. USA*, 86:8932-8935, 1989; and Zijlstra *et al.*, *Nature*, 342:435-438, 1989).

In a specific embodiment, viral vectors that contain nucleic acid sequences encoding an antibody or a fusion protein are used. For example, a retroviral vector can be used (see Miller *et al.*, *Meth. Enzymol.*, 217:581-599, 1993). These retroviral vectors contain
35 the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody or a fusion protein to

be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the nucleotide sequence into a subject. More detail about retroviral vectors can be found in Boesen *et al.*, *Biotherapy*, 6:291-302, 1994, which describes the use of a retroviral vector to deliver the *mdr 1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes *et al.*, *J. Clin. Invest.*, 93:644-651, 1994; Klein *et al.*, *Blood* 83:1467-1473, 1994; Salmons and Gunzberg, *Human Gene Therapy*, 4:129-141, 1993; and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.*, 3:110-114, 1993.

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development*, 3:499-503, 1993, present a review of adenovirus-based gene therapy. Bout *et al.*, *Human Gene Therapy*, 5:3-10, 1994, demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld *et al.*, *Science*, 252:431-434, 1991; Rosenfeld *et al.*, *Cell*, 68:143-155, 1992; Mastrangeli *et al.*, *J. Clin. Invest.*, 91:225-234, 1993; PCT Publication WO 94/12649; and Wang *et al.*, *Gene Therapy*, 2:775-783, 1995. In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (see, *e.g.*, Walsh *et al.*, *Proc. Soc. Exp. Biol. Med.*, 204:289-300, 1993, and U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a subject.

In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcellmediated gene transfer,

spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, *e.g.*, Loeffler and Behr, *Meth. Enzymol.*, 217:599-618, 1993; Cohen *et al.*, *Meth. Enzymol.*, 217:618-644, 1993; and *Clin. Pharma. Ther.*, 29:69-92, 1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a subject by various methods known in the art. Recombinant blood cells (*e.g.*, hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, *e.g.*, as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the subject.

In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody or a fusion protein are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention (see *e.g.*, PCT Publication WO 94/08598; Stemple and Anderson, *Cell*, 71:973-985, 1992; Rheinwald, *Meth. Cell Bio.*, 21A:229, 1980; and Pittelkow and Scott, *Mayo Clinic Proc.*, 61:771, 1986).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

5.8. CHARACTERIZATION AND DEMONSTRATION OF THERAPEUTIC OR PROPHYLACTIC UTILITY

Antibodies, fusion proteins, and conjugated molecules of the present invention may be characterized in a variety of ways. In particular, antibodies of the invention may be assayed for the ability to immunospecifically bind to an antigen. Such an assay may be performed in solution (*e.g.*, Houghten, *Bio/Techniques*, 13:412-421, 1992), on beads (Lam, *Nature*, 354:82-84, 1991, on chips (Fodor, *Nature*, 364:555-556, 1993), on bacteria (U.S. Patent No. 5,223,409), on spores (U.S. Patent Nos. 5,571,698; 5,403,484; and 5,223,409), on plasmids (Cull *et al.*, *Proc. Natl. Acad. Sci. USA*, 89:1865-1869, 1992) or on phage (Scott and Smith, *Science*, 249:386-390, 1990; Devlin, *Science*, 249:404-406, 1990; Cwirla *et al.*, *Proc. Natl. Acad. Sci. USA*, 87:6378-6382, 1990; and Felici, *J. Mol. Biol.*, 222:301-310, 1991) (each of these references is incorporated herein in its entirety by reference). Antibodies that have been identified to immunospecifically bind to an antigen or a fragment thereof can then be assayed for their specificity affinity for the antigen.

The antibodies of the invention or fragments thereof may be assayed for immunospecific binding to an antigen and cross-reactivity with other antigens by any method known in the art. Immunoassays which can be used to analyze immunospecific binding and cross-reactivity include, but are not limited to, competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, *e.g.*, Ausubel *et al.*, eds, 1994, *Current Protocols in Molecular Biology*, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasyolol) supplemented with protein phosphatase and/or protease inhibitors (*e.g.*, EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (*e.g.*, 1 to 4 hours) at 40°C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 40°C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to

immunoprecipitate a particular antigen can be assessed by, *e.g.*, western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (*e.g.*, pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (*e.g.*, 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (*e.g.*, PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (*e.g.*, PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, *e.g.*, an anti-human antibody) conjugated to an enzymatic substrate (*e.g.*, horseradish peroxidase or alkaline phosphatase) or radioactive molecule (*e.g.*, ^{32}P or ^{125}I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (*e.g.*, horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, *e.g.*, Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (*e.g.*, ^3H or ^{125}I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of the present invention or a fragment thereof for the antigen and the binding off-rates can be determined from the saturation data by scatchard analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with an antibody of the present invention or a fragment thereof conjugated to a labeled compound (*e.g.*, ^3H or ^{125}I) in the presence of increasing amounts of an unlabeled second antibody.

In a preferred embodiment, BIAcore kinetic analysis is used to determine the binding on and off rates of antibodies to an antigen. BIAcore kinetic analysis comprises analyzing the binding and dissociation of an antigen from chips with immobilized antibodies on their surface (see the Example section *infra*).

The antibodies of the invention as well as fusion proteins and conjugated molecules can also be assayed for their ability to inhibit the binding of an antigen to its host cell receptor using techniques known to those of skill in the art. For example, cells expressing the receptor for a viral antigen can be contacted with virus in the presence or absence of an antibody and the ability of the antibody to inhibit viral antigen's binding can be measured by, for example, flow cytometry or a scintillation counter. The antigen or the antibody can be labeled with a detectable compound such as a radioactive label (*e.g.*, ^{32}P , ^{35}S , and ^{125}I) or a fluorescent label (*e.g.*, fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine) to enable detection of an interaction between the antigen and its host cell receptor. Alternatively, the ability of antibodies to inhibit an antigen from binding to its receptor can be determined in cell-free assays. For example, virus or a viral antigen (*e.g.*, RSV F glycoprotein) can be contacted in a cell-free assay with an antibody and the ability of the antibody to inhibit the virus or the viral antigen from binding to its host cell receptor can be determined. Preferably, the antibody is immobilized on a solid support and the antigen is labeled with a detectable compound. Alternatively, the antigen is immobilized on a solid support and the antibody is labeled with a detectable compound. The antigen may be partially or completely purified (*e.g.*, partially or completely free of other polypeptides) or part of a cell lysate. Further, the antigen may be a fusion protein comprising the viral antigen and a domain such as glutathione-S-transferase. Alternatively, an antigen can be biotinylated using techniques

well known to those of skill in the art (*e.g.*, biotinylation kit, Pierce Chemicals; Rockford, IL).

5 The antibodies, fusion proteins, and conjugated molecules of the invention can also be assayed for their ability to inhibit or downregulate viral or bacterial replication using techniques known to those of skill in the art. For example, viral replication can be assayed by a plaque assay such as described, *e.g.*, by Johnson *et al.*, *Journal of Infectious Diseases*, 176:1215-1224, 1997. The antibodies, fusion proteins, and conjugated molecules of the invention of the invention can also be assayed for their ability to inhibit or
10 downregulate the expression of viral or bacterial polypeptides. Techniques known to those of skill in the art, including, but not limited to, Western blot analysis, Northern blot analysis, and RT-PCR, can be used to measure the expression of viral or bacterial polypeptides. Further, the antibodies, fusion proteins, and conjugated molecules of the invention of the invention can be assayed for their ability to prevent the formation of syncytia.

15 The antibodies, fusion proteins, conjugated molecules, and compositions of the invention are preferably tested *in vitro*, and then *in vivo* for the desired therapeutic or prophylactic activity, prior to use in humans. For example, *in vitro* assays which can be used to determine whether administration of a specific antibody, a specific fusion protein, a specific conjugated molecule, or a composition of the present invention is indicated, include
20 *in vitro* cell culture assays in which a subject tissue sample is grown in culture, and exposed to or otherwise administered an antibody, a fusion protein, conjugated molecule, or composition of the present invention, and the effect of such an antibody, a fusion protein, conjugated molecule, or a composition of the present invention upon the tissue sample is observed. In various specific embodiments, *in vitro* assays can be carried out with
25 representative cells of cell types involved in a disease or disorder, to determine if an antibody, a fusion protein, conjugated molecule, or composition of the present invention has a desired effect upon such cell types. Preferably, the antibodies, the fusion proteins, the conjugated molecules, or compositions of the invention are also tested in *in vitro* assays and animal model systems prior to administration to humans.

30 Antibodies, fusion proteins, conjugated molecules, or compositions of the present invention for use in therapy can be tested for their toxicity in suitable animal model systems, including but not limited to rats, mice, cows, monkeys, and rabbits. For *in vivo* testing for the toxicity of an antibody, a fusion protein, a conjugated molecule, or a composition, any animal model system known in the art may be used.

35 Efficacy in treating or preventing viral infection may be demonstrated by detecting the ability of an antibody, a fusion protein, a conjugated molecule, or a composition

of the invention to inhibit the replication of the virus, to inhibit transmission or prevent the virus from establishing itself in its host, or to prevent, ameliorate or alleviate one or more symptoms associated with viral infection. The treatment is considered therapeutic if there is, for example, a reduction in viral load, amelioration of one or more symptoms or a decrease in mortality and/or morbidity following administration of an antibody, a fusion protein, a conjugated molecule, or a composition of the invention. Antibodies, fusion proteins, conjugated molecules, or compositions of the invention can also be tested for their ability to inhibit viral replication or reduce viral load in *in vitro* and *in vivo* assays.

Efficacy in treating or preventing bacterial infection may be demonstrated by detecting the ability of an antibody, a fusion protein or a composition of the invention to inhibit the bacterial replication, or to prevent, ameliorate or alleviate one or more symptoms associated with bacterial infection. The treatment is considered therapeutic if there is, for example, a reduction in bacterial numbers, amelioration of one or more symptoms or a decrease in mortality and/or morbidity following administration of an antibody, a fusion protein or a composition of the invention.

Efficacy in treating cancer may be demonstrated by detecting the ability of an antibody, a fusion protein, a conjugated molecule, or a composition of the invention to inhibit or reduce the growth or metastasis of cancerous cells or to ameliorate or alleviate one or more symptoms associated with cancer. The treatment is considered therapeutic if there is, for example, a reduction in the growth or metastasis of cancerous cells, amelioration of one or more symptoms associated with cancer, or a decrease in mortality and/or morbidity following administration of an antibody, a fusion protein, a conjugated molecule, or a composition of the invention. Antibodies, fusion proteins or compositions of the invention can be tested for their ability to reduce tumor formation in *in vitro*, *ex vivo*, and *in vivo* assays.

Efficacy in treating inflammatory disorders may be demonstrated by detecting the ability of an antibody, a fusion protein, a conjugated molecule, or a composition of the invention to reduce or inhibit the inflammation in an animal or to ameliorate or alleviate one or more symptoms associated with an inflammatory disorder. The treatment is considered therapeutic if there is, for example, a reduction in inflammation or amelioration of one or more symptoms following administration of an antibody, a fusion proteins, a conjugated molecule, or a composition of the invention.

Antibodies, fusion proteins, conjugated molecules, or compositions of the invention can be tested *in vitro* and *in vivo* for the ability to induce the expression of cytokines (*e.g.*, IFN- α , IFN- β , IFN- γ , IL-2, IL-3, IL-4, IL-5, IL-6, IL10, IL-12, and IL-15) and

activation markers (*e.g.*, CD28, ICOS, and SLAM). Techniques known to those of skill in the art can be used to measure the level of expression of cytokines and activation markers. For example, the level of expression of cytokines can be measured by analyzing the level of
5 RNA of cytokines by, for example, RT-PCR and Northern blot analysis, and by analyzing the level of cytokines by, for example, immunoprecipitation followed by Western blot analysis or ELISA.

Antibodies, fusion proteins, conjugated molecules, or compositions of the invention can be tested *in vitro* and *in vivo* for their ability to modulate the biological activity
10 of immune cells, preferably human immune cells (*e.g.*, T-cells, B-cells, and Natural Killer cells). The ability of an antibody, a fusion protein, a conjugated molecule, or a composition of the invention to modulate the biological activity of immune cells can be assessed by detecting the expression of antigens, detecting the proliferation of immune cells, detecting the activation of signaling molecules, detecting the effector function of immune cells, or
15 detecting the differentiation of immune cells. Techniques known to those of skill in the art can be used for measuring these activities. For example, cellular proliferation can be assayed by ³H-thymidine incorporation assays and trypan blue cell counts. Antigen expression can be assayed, for example, by immunoassays including, but are not limited to, competitive and non-competitive assay systems using techniques such as Western blots,
20 immunohistochemistry, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays and FACS analysis. The activation of signaling molecules can be assayed, for example, by
25 kinase assays and electrophoretic shift assays (EMSAs).

Antibodies, fusion proteins, conjugated molecules, or compositions of the invention can also be tested for their ability to increase the survival period of animals, preferably mammals and most preferably humans, suffering from a disease, disorder, or infection by at least 25%, preferably at least 50%, at least 60%, at least 75%, at least 85%, at
30 least 95%, or at least 99%. Further, antibodies, fusion proteins, conjugated molecules, or compositions of the invention can be tested for their ability reduce the hospitalization period of animals, preferably mammals and most preferably humans, suffering from a disease, disorder, or infection by at least 60%, preferably at least 75%, at least 85%, at least 95%, or at least 99%. Techniques known to those of skill in the art can be used to analyze the
35 function of the antibodies or compositions of the invention *in vivo*.

5.9. **DIAGNOSTIC USES OF ANTIBODIES AND FUSION PROTEINS**

Labeled antibodies, fusion proteins, and conjugated molecules of the invention can be used for diagnostic purposes to detect, diagnose, or monitor diseases, disorders or infections. The invention provides for the detection or diagnosis of a disease, disorder or infection, comprising: (a) assaying the expression of an antigen in cells or a tissue sample of a subject using one or more antibodies that immunospecifically bind to the antigen; and (b) comparing the level of the antigen with a control level, *e.g.*, levels in normal tissue samples, whereby an increase in the assayed level of antigen compared to the control level of the antigen is indicative of the disease, disorder or infection. The invention also provides for the detection or diagnosis of a disease, disorder or infection, comprising (a) assaying the expression of an antigen in cells or a tissue sample of a subject using one or fusion proteins or conjugated molecules of the invention that bind to the antigen; and (b) comparing the level of the antigen with a control level, *e.g.*, levels in normal tissue samples, whereby an increase of antigen compared to the control level of the antigen is indicative of the disease, disorder or infection. Accordingly, the fusion protein or conjugated molecule comprises a bioactive molecule such as a ligand, cytokine or growth factor and the hinge-Fc region or fragments thereof, wherein the fusion protein or conjugated molecule is capable of binding to an antigen being detected.

Antibodies of the invention can be used to assay antigen levels in a biological sample using classical immunohistological methods as described herein or as known to those of skill in the art (*e.g.*, see Jalkanen *et al.*, *J. Cell. Biol.*, 101:976-985, 1985; Jalkanen *et al.*, *J. Cell. Biol.*, 105:3087-3096, 1987). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, alkaline phosphatase, glucose oxidase; radioisotopes, such as iodine (^{125}I , ^{131}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{121}In), and technetium ($^{99\text{m}}\text{Tc}$); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine.

Fusion proteins can be used to assay antigen levels in a biological sample using, for example, SDS-PAGE and immunoassays known to those of skill in the art.

One aspect of the invention is the detection and diagnosis of a disease, disorder, or infection in a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled antibody that immunospecifically binds to an antigen; b) waiting for a

time interval following the administration for permitting the labeled antibody to preferentially concentrate at sites in the subject where the antigen is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled antibody in the subject, such that detection of labeled antibody above the background level indicates that the subject has the disease, disorder, or infection. In accordance with this embodiment, the antibody is labeled with an imaging moiety which is detectable using an imaging system known to one of skill in the art. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

In another embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled fusion protein or conjugated molecule that binds to an antigen or some other molecule; b) waiting for a time interval following the administration for permitting the labeled fusion protein or conjugated molecule to preferentially concentrate at sites in the subject where the antigen or other molecule is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled fusion protein or conjugated molecule in the subject, such that detection of labeled fusion protein above the background level indicates that the subject has the disease, disorder, or infection. In accordance with this embodiment, the fusion protein or conjugated molecule comprises a bioactive molecule such as a ligand, cytokine or growth factor and a hinge-Fc region or a fragment thereof, wherein said fusion protein or conjugated molecule is labeled with an imaging moiety and is capable of binding to the antigen being detected.

It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc . The labeled antibody will then preferentially accumulate at the location of cells which contain the specific protein. *In vivo* tumor imaging is described in S.W. Burchiel *et al.*, "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments," Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours.

In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

In one embodiment, monitoring of a disease, disorder or infection is carried
5 out by repeating the method for diagnosing the disease, disorder or infection, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

Presence of the labeled molecule can be detected in the subject using methods known in the art for *in vivo* scanning. These methods depend upon the type of label used.
10 Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

In a specific embodiment, the molecule is labeled with a radioisotope and is
15 detected in the patient using a radiation responsive surgical instrument (Thurston *et al.*, U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the
20 molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI).

5.10. KITS

The invention also provides a pharmaceutical pack or kit comprising one or
25 more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

The present invention provides kits that can be used in the above methods. In
30 one embodiment, a kit comprises an antibody, fusion protein, or conjugated molecule, of the invention, preferably in a purified form, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated antigen as a control. Preferably, the kits of the present invention further comprise a control antibody,
35 fusion protein, or conjugated molecule which does not react with the antigen included in the kit. In another specific embodiment, the kits of the present invention contain a means for

detecting the binding of an antibody, fusion protein, or conjugated molecule, to an antigen (e.g., the antibody, fusion protein, or conjugated molecule, may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized antigen. The antigen provided in the kit may also be attached to a solid support. In a more specific embodiment the detecting means of the above-described kit includes a solid support to which antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the antigen can be detected by binding of the said reporter-labeled antibody.

5.11 *IN VITRO* AND *IN VIVO* ASSAYS FOR EXTENDED HALF-LIFE OF MODIFIED IGG HINGE-FC FRAGMENTS

The binding ability of modified IgGs and molecules comprising an IgG constant domain of FcRn fragment thereof to FcRn can be characterized by various *in vitro* assays. PCT publication WO 97/34631 by Ward discloses various methods in detail and is incorporated herein in its entirety by reference.

For example, in order to compare the ability of the modified IgG or fragments thereof to bind to FcRn with that of the wild type IgG, the modified IgG or fragments thereof and the wild type IgG can be radio-labeled and reacted with FcRn-expressing cells *in vitro*. The radioactivity of the cell-bound fractions can be then counted and compared. The cells expressing FcRn to be used for this assay are preferably endothelial cell lines including mouse pulmonary capillary endothelial cells (B10, D2.PCE) derived from lungs of B10.DBA/2 mice and SV40 transformed endothelial cells (SVEC) (Kim *et al.*, *J. Immunol.*, 40:457-465, 1994) derived from C3H/HeJ mice. However, other types of cells, such as intestinal brush borders isolated from 10- to 14-day old suckling mice, which express sufficient number of FcRn can be also used. Alternatively, mammalian cells which express recombinant FcRn of a species of choice can be also utilized. After counting the radioactivity of the bound fraction of modified IgG or that of wild type, the bound molecules can be then extracted with the detergent, and the percent release per unit number of cells can be calculated and compared.

Affinity of modified IgGs for FcRn can be measured by surface plasmon resonance (SPR) measurement using, for example, a BIAcore 2000 (BIAcore Inc.) as

described previously (Popov *et al.*, *Mol. Immunol.*, 33:493-502, 1996; Karlsson *et al.*, *J. Immunol. Methods*, 145:229-240, 1991, both of which are incorporated by reference in their entireties). In this method, FcRn molecules are coupled to a BIAcore sensor chip (*e.g.*, CM5 chip by Pharmacia) and the binding of modified IgG to the immobilized FcRn is measured at
5 a certain flow rate to obtain sensorgrams using BIA evaluation 2.1 software, based on which on- and off-rates of the modified IgG, constant domains, or fragments thereof, to FcRn can be calculated.

Relative affinities of modified IgGs or fragments thereof, and the wild type
10 IgG for FcRn can be also measured by a simple competition binding assay. Unlabeled modified IgG or wild type IgG is added in different amounts to the wells of a 96-well plate in which FcRn is immobilize. A constant amount of radio-labeled wild type IgG is then added to each well. Percent radioactivity of the bound fraction is plotted against the amount of unlabeled modified IgG or wild type IgG and the relative affinity of the modified hinge-Fc
15 can be calculated from the slope of the curve.

Furthermore, affinities of modified IgGs or fragments thereof, and the wild type IgG for FcRn can be also measured by a saturation study and the Scatchard analysis.

Transfer of modified IgG or fragments thereof across the cell by FcRn can be measured by *in vitro* transfer assay using radiolabeled IgG or fragments thereof and FcRn-
20 expressing cells and comparing the radioactivity of the one side of the cell monolayer with that of the other side. Alternatively, such transfer can be measured *in vivo* by feeding 10- to 14-day old suckling mice with radiolabeled, modified IgG and periodically counting the radioactivity in blood samples which indicates the transfer of the IgG through the intestine to the circulation (or any other target tissue, *e.g.*, the lungs). To test the dose-dependent
25 inhibition of the IgG transfer through the gut, a mixture of radiolabeled and unlabeled IgG at certain ratio is given to the mice and the radioactivity of the plasma can be periodically measured (Kim *et al.*, *Eur. J. Immunol.*, 24:2429-2434, 1994).

The half-life of modified IgG or fragments thereof can be measure by pharmacokinetic studies according to the method described by Kim *et al.* (*Eur. J. of Immuno.*
30 24:542, 1994), which is incorporated by reference herein in its entirety. According to this method, radiolabeled modified IgG or fragments thereof is injected intravenously into mice and its plasma concentration is periodically measured as a function of time, for example, at 3 minutes to 72 hours after the injection. The clearance curve thus obtained should be biphasic, that is, α -phase and β -phase. For the determination of the *in vivo* half-life of the
35 modified IgGs or fragments thereof, the clearance rate in β -phase is calculated and compared with that of the wild type IgG.

6. EXAMPLES

The following examples illustrate the production, isolation, and
5 characterization of modified hinge-Fc fragments that have longer *in vivo* half-lives.

6.1 LIBRARY CONSTRUCTION

6.1.1 REAGENTS

All chemicals were of analytical grade. Restriction enzymes and DNA-
10 modifying enzymes were purchased from New England Biolabs, Inc. (Beverly, MA).
Oligonucleotides were synthesized by MWG Biotech, Inc. (High Point, NC). pCANTAB5E
phagemid vector, anti-E-tag-horseradish peroxidase conjugate, TG1 *E. Coli* strain, IgG
Sephadex 6 Fast Flow and HiTrap protein A columns were purchased from APBiotec, Inc.
(Piscataway, NJ). VCSM13 helper phage and the Quick change mutagenesis kit were
15 obtained from Stratagene (La Jolla, CA). CJ236 *E. coli* strain was purchased from Bio-Rad
(Richmond, CA). BCA Protein Assay Reagent Kit was obtained from Pierce (Rockford, IL).
Lipofectamine 2000 was purchased from Invitrogen, Inc. (Carlsbad, CA).

6.1.2 EXPRESSION AND PURIFICATION OF MURINE AND HUMAN FcRn

20 The amino acid sequences of human and mouse FcRn are SEQ ID NOs. 84
and 85, respectively (see also Firan *et al.*, *Intern. Immunol.*, 13:993-1002, 2001 and Popov *et al.*,
Mol. Immunol., 33:521-530, 1996, both of which are incorporated herein by reference in
their entireties). Human FcRn was also obtained following isolation from human placenta
25 cDNA (Clontech, Palo Alto, CA) of the genes for human β 2-microglobulin (Kabat *et al.*,
1991, Sequences of Proteins of Immunological Interest, U.S. Public Health Service, National
Institutes of Health, Washington, DC) and codons -23 to 267 of the human α chain (Story *et al.*,
J. Exp. Med., 180:2377-2381, 1994) using standard PCR protocols. Light and heavy
chains along with their native signal sequence (Kabat *et al.*, 1991, *supra*; Story *et al.*, *supra*)
30 were cloned in pFastBac DUAL and pFastBac1 bacmids, respectively, and viral stocks
produced in *Spodoptera frugiperda* cells (Sf9) according to the manufacturer's instructions
(Invitrogen, Carlsbad, CA). High-Five cells were infected at a multiplicity of infection of 3
with the baculoviruses encoding α and β 2 chains using commercially available protocols
(Invitrogen). Recombinant human FcRn was purified as follows: supernatant of infected
35 insect cells was dialyzed into 50 mM MES (2-N-[Morpholino]ethanesulfonic acid) pH 6.0 and

applied to a 10 ml human IgG Sepharose 6 Fast Flow column (APBiotect, Piscataway, NJ). Resin was washed with 200 ml 50 mM MES pH 6.0 and FcRn eluted with 0.1 M Tris-Cl pH 8.0. Purified FcRn was dialyzed against 50 mM MES pH 6.0, flash frozen and stored at -70°C. The purity of proteins was checked by SDS-PAGE and HPLC.

6.1.3 PREPARATION OF TAA-CONTAINING ssDNA URACIL TEMPLATE

Construction of the libraries was based on a site directed mutagenesis strategy derived from the Kunkel method (Kunkel *et al.*, *Methods Enzymol.* 154:367-382, 1987). A human hinge-Fc gene spanning amino acid residues 226-478 (Kabat numbering, Kabat *et al.*, 1991, *supra*) derived from MEDI-493 human IgG1 (Johnson *et al.*, *J. Infect. Disease*, 176:1215-1224, 1997), was cloned into the pCANTAB5E phagemid vector as an SfiI/NotI fragment. Four libraries were generated by introducing random mutations at positions 251, 252, 254, 255, 256 (library 1), 308, 309, 311, 312, 314 (library 2), 385, 386, 387, 389 (library 3) and 428, 433, 434, 436 (library 4). Briefly, four distinct hinge-Fc templates were generated using PCR by overlap extension (Ho *et al.*, *Gene*, 15:51-59, 1989), each containing one TAA stop codon at position 252 (library 1), 310 (library 2), 384 (library 3) or 429 (library 4), so that only mutagenized phagemids will give rise to Fc-displaying phage.

Each TAA-containing single-stranded DNA (TAAssDNA) was then prepared as follows: a single CJ236 *E. coli* colony harboring one of the four relevant TAA-containing phagemids was grown in 10 ml 2 x YT medium supplemented with 10 µg/ml chloramphenicol and 100 µg/ml ampicillin. At OD₆₀₀ = 1, VCSM13 helper phage was added to a final concentration of 10¹⁰ pfu/ml. After 2 hours, the culture was transferred to 500 ml of 2 x YT medium supplemented with 0.25 µg/ml uridine, 10 µg/ml chloramphenicol, 30 µg/ml kanamycin, 100 µg/ml ampicillin and grown overnight at 37°C. Phage were precipitated with PEG6000 using standard protocols (Sambrook *et al.*, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, New York, Vols. 1-3) and purified using the Qiaprep Spin M13 Kit (Qiagen, Valencia, CA) according to the manufacturer's instructions. 10 to 30 µg of each uracil-containing TAAssDNA template was then combined with 0.6 µg of the following phosphorylated oligonucleotides (randomized regions underlined) in 50 mM Tris-HCl, 10 mM MgCl₂, pH 7.5 in a final volume of 250 µl:

Library 1:

5'-CATGTGACCTCAGGSNNNSNNNGATNNNSNNGGTGTCTTGGGTTTT

GGGGGG-3' (SEQ ID NO:120)

Library 2:

5'-GCACTTGTA~~CTCCTTGCCATT~~SNNCCASNN~~SNN~~GTGSNN~~SNN~~GGTGA

5 GGACGC-3' (SEQ ID NO:121)

Library 3:

5'-GGTCTTGTA~~GTTSNN~~CTCSNN~~SNN~~SNNATTGCTCTCCC-3' (SEQ ID NO:122)

Library 4:

5'-GGCTCTTCTGCGTSNNGTGSNN~~SNN~~CAGAGCCTCATGSNN~~CAC~~GGAGC

10 ATGAG-3' (SEQ ID NO:123)

where N = A, C, T or G and S = G or C.

6.1.4 SYNTHESIS OF HETERODUPLEX DNA

Appropriate, degenerate oligonucleotides were phosphorylated in the presence of T4 polynucleotide kinase using the standard protocol. Ten to 30 µg of ssDNA U template and 0.6 µg of phosphorylated oligonucleotide were combined in 50 mM Tris-HCl containing 10 mM MgCl₂, pH 7.5, to a final volume of 250 µl and incubated at 90°C for 2 minutes, 50°C for 3 minutes, and 20°C for 5 minutes. Synthesis of the heteroduplex DNA was carried out by adding 30 units of both T4 DNA ligase and T7 DNA polymerase in the presence of 0.4 mM ATP, 1 mM dNTPs and 6 mM DTT and the mixture was incubated for 4 hours at 20°C. The heteroduplex DNA thus produced was then purified and desalted using Qiagen Qiaquick DNA purification Kit (Qiagen, CA).

6.1.5 ELECTROPORATION

300 µl electrocompetent TG1 *E. coli* cells were electroporated with 1 to 5 µg of the heteroduplex DNA in a 2.5 kV field using 200 Ω and 25 µF capacitance until a library size of 1 x 10⁸ (library 1 and 2) or 1 x 10⁷ (library 3 and 4) was reached. The cells were resuspended in 2 ml SOC medium and the procedure was repeated 6 to 10 times. The diversity was assessed by titration of recombinant *E. coli*. The pulsed cells were incubated in 50 ml SOC medium for 30 minutes at 37°C under agitation, centrifuged, and resuspended in 500 ml 2xYT containing 100 µg/ml ampicillin and 10¹⁰ pfu/ml of VCSM13 helper phage. The culture was incubated overnight at 37°C and the cells were pelleted by centrifugation. The phage in the supernatant which express mutated hinge-Fc portion on its GIII-coat protein were precipitated with PEG6000 as previously described (Sambrook *et al.*, 1989, *supra*) and resuspended in 5 ml of 20 mM MES, pH 6.0.

6.2 PANNING OF THE LIBRARY

Phage were panned using an ELISA-based approach. A 96-well ELISA plate was coated with 100 μ l/well of 0.01 mg/ml murine FcRn in sodium carbonate buffer, pH 9.0, at 4°C overnight and then blocked with 4% skimmed milk at 37°C for 2 hours. In each well of the coated plate, 100-150 μ l of the phage suspension (about 10^{13} phage in total) in 20 mM MES, pH 6.0, containing 5% milk and 0.05% Tween 20, were placed and incubated at 37°C for two to three hours with agitation.

After the incubation, the wells were washed with 20 mM MES, pH 6.0, containing 0.2% Tween 20 and 0.3 M NaCl about thirty times at room temperature. The bound phage were eluted with 100 μ l/well of PBS, pH 7.4, at 37°C for 30 minutes.

The eluted phage were then added to the culture of exponentially growing *E. coli* cells and propagation was carried out overnight at 37°C in 250 ml 2xYT supplemented with 100 μ g/ml ampicillin and 10^{10} pfu/ml of VCSM13 helper phage. Propagated phage were collected by centrifugation followed by precipitation with PEG and the panning process was repeated up to a total of six times.

For the phage library containing mutations in residues 308-314 (H310 and W313 fixed), the phage expressing hinge-Fc region with higher affinities for FcRn were enriched by each panning process as shown in Table IV. The panning results of the library for the mutations in the residues 251-256 (I253 fixed) and that of the library for the mutations in the residues 428-436 (H429, E430, A431, L432, and H435 fixed), are shown in Tables V and VI, respectively. Furthermore, the panning results of the library for the mutations in the residues 385-389 (E388 fixed) is shown in Table VII.

Table IV

**PANNING OF LIBRARY (RESIDUES 308-314; H310 AND W313 FIXED)
pCANTAB5E-KUNKEL-muFcRn (MURINE FcRn)**

PANNING	OUTPUT		ENRICHMENT RATIO
	+ FcRn	- FcRn	
1st Round	1.1×10^5	0.5×10^5	2
2nd Round	1×10^4	0.2×10^4	5
3rd Round	9×10^4	0.3×10^4	30
4th Round	3×10^5	2×10^4	15

Table V
PANNING OF LIBRARY (RESIDUES 251-256; I253 FIXED)
pCANTAB5E-KUNKEL-muFcRn

PANNING	OUTPUT		ENRICHMENT RATIO
	+ FcRn	- FcRn	
1st Round	2.5×10^5	1×10^5	2.5
2nd Round	6×10^4	2×10^4	3.0
3rd Round	8×10^5	4×10^4	20
4th Round	1.2×10^6	5×10^4	24
5th Round	3.0×10^6	6×10^4	50

Table VI

**PANNING OF LIBRARY (RESIDUES 428-436;
H429, E430, A431, L432, AND H435 FIXED)
pCANTAB5E-KUNKEL- μ FcRn**

5

10

PANNING	OUTPUT		ENRICHMENT RATIO
	+ FcRn	- FcRn	
1st Round	2.3×10^5	0.9×10^5	2.5
2nd Round	3×10^4	1×10^4	3
3rd Round	2×10^5	2×10^4	10
4th Round	8×10^5	5×10^4	16

15

20

25

30

35

Table VII
PANNING OF LIBRARY (RESIDUES 385-389; E388 FIXED)
pCANTAB5E-KUNKEL-muFcRn

PANNING	OUTPUT		ENRICHMENT RATIO
	+ FcRn	- FcRn	
1st Round	4.2×10^5	3.8×10^5	1.1
2nd Round	5×10^4	0.3×10^4	17
3rd Round	3.5×10^5	1×10^4	35
4th Round	5.5×10^5	4×10^4	14
5th Round	7.5×10^5	5×10^4	15
6th Round	2×10^6	1×10^5	20

6.3 IDENTIFICATION OF ISOLATED CLONES FROM PANNING

After each panning process, phage were isolated and the nucleic acids encoding the expressed peptides which bound to FcRn were sequenced by a standard sequencing method such as by dideoxynucleotide sequencing (Sanger *et al.*, *Proc. Natl. Acad. Sci USA*, 74:5463-5467, 1977) using a ABI3000 genomic analyzer (Applied Biosystems, Foster City, CA).

As a result of panning, two mutants were isolated from the phage library containing mutations in residues 308-314 (H310 and W313 fixed), thirteen mutants from the library for residues 251-256 (I253 fixed), six mutants from the library for residues 428-436 (H429, E430, A431, L432, and H435 fixed), and nine mutants from the library for residues 385-389 (E388 fixed). The mutants isolated from the libraries are listed in Table VIII.

Table VIII
MUTANTS ISOLATED BY PANNING

LIBRARY	MUTANTS*
251-256	<u>Leu Tyr Ile Thr Arg Glu</u> (SEQ ID NO:90)
	<u>Leu Tyr Ile Ser Arg Thr</u> (SEQ ID NO:91)
	<u>Leu Tyr Ile Ser Arg Ser</u> (SEQ ID NO:92)
	<u>Leu Tyr Ile Ser Arg Arg</u> (SEQ ID NO:93)

LIBRARY	MUTANTS*
5 10	<i>Leu Tyr Ile Ser Arg Gln</i> (SEQ ID NO:94)
	<i>Leu Trp Ile Ser Arg Thr</i> (SEQ ID NO:95)
	<i>Leu Tyr Ile Ser Leu Gln</i> (SEQ ID NO:96)
	<i>Leu Phe Ile Ser Arg Asp</i> (SEQ ID NO:97)
	<i>Leu Phe Ile Ser Arg Thr</i> (SEQ ID NO:98)
	<i>Leu Phe Ile Ser Arg Arg</i> (SEQ ID NO:99)
	<i>Leu Phe Ile Thr Gly Ala</i> (SEQ ID NO:100)
	<i>Leu Ser Ile Ser Arg Glu</i> (SEQ ID NO:101)
	<i>Arg Thr Ile Ser Ile Ser</i> (SEQ ID NO:102)
15 308-314	<i>Thr Pro His Ser Asp Trp Leu</i> (SEQ ID NO:103)
	<i>Ile Pro His Glu Asp Trp Leu</i> (SEQ ID NO:104)
20 25	<i>Arg Thr Arg Glu Pro</i> (SEQ ID NO:105)
	<i>Asp Pro Pro Glu Ser</i> (SEQ ID NO:106)
	<i>Ser Asp Pro Glu Pro</i> (SEQ ID NO:107)
	<i>Thr Ser His Glu Asn</i> (SEQ ID NO:108)
	<i>Ser Lys Ser Glu Asn</i> (SEQ ID NO:109)
	<i>His Arg Ser Glu Asn</i> (SEQ ID NO:110)
	<i>Lys Ile Arg Glu Asn</i> (SEQ ID NO:111)
	<i>Gly Ile Thr Glu Ser</i> (SEQ ID NO:112)
	<i>Ser Met Ala Glu Pro</i> (SEQ ID NO:113)
428-436	<i>Met His Glu Ala Leu Arg Tyr His His</i> (SEQ ID NO:114)
30	<i>Met His Glu Ala Leu His Phe His His</i> (SEQ ID NO:115)
	<i>Met His Glu Ala Leu Lys Phe His His</i> (SEQ ID NO:116)
	<i>Met His Glu Ala Leu Ser Tyr His Arg</i> (SEQ ID NO:117)
	<i>Thr His Glu Ala Leu His Tyr His Thr</i> (SEQ ID NO:118)
	<i>Met His Glu Ala Leu His Tyr His Tyr</i> (SEQ ID NO:119)

* Substituting residues are indicated in bold face

The underlined sequences in Table VIII correspond to sequences that occurred 10 to 20 times in the final round of panning and the sequences in italics correspond to sequences that occurred 2 to 5 times in the final round of panning. Those sequences that are neither underlined nor italicized occurred once in the final round of panning.

6.4 EXPRESSION AND PURIFICATION OF SOLUBLE MUTANT HINGE-FC REGION

The genes encoding mutated hinge-Fc fragments are excised with appropriate restriction enzymes and recloned into an expression vector, for example, V β pelBhis (Ward, *J. Mol. Biol.*, 224:885-890, 1992). Vectors containing any other type of tag sequence, such as *c-myc* tag, decapeptide tag (Huse *et al.*, *Science*, 246:1275-1281, 1989), FlagTM (Immunex) tags, can be used. Recombinant clones, such as *E. coli*, are grown and induced to express soluble hinge-Fc fragments, which can be isolated from the culture media or cell lysate after osmotic shock, based on the tag used, or by any other purification methods well known to those skilled in the art and characterized by the methods as listed below.

6.5 CONSTRUCTION, PRODUCTION AND PURIFICATION OF IgG1 VARIANTS

Representative Fc mutations such as I253A, M252Y/S254T/T256E, M252W, M252Y, M252Y/T256Q, M252F/T256D, V308T/L309P/Q311S, G385D/Q386P/N389S, G385R/Q386T/P387R/N389P, H433K/N434F/Y436H, and N434F/Y436 were incorporated into the human IgG1 MEDI-493 (SYNAGIS[®]) (Johnson *et al.*, 1997, *supra*). The heavy chain was subjected to site-directed mutagenesis using a Quick Change Mutagenesis kit (Stratagene, La Jolla, CA) according to the manufacturer's instructions and sequences were verified by didoxynucleotide sequencing using a ABI3000 (Applied Biosystems, Foster City, CA) sequencer. The different constructions were expressed transiently in human embryonic kidney 293 cells using a CMV immediate-early promoter and dicistronic operon in which IgG1/V_H is cosecreted with IgG1/V_L (Johnson *et al.*, 1997, *supra*). Transfection was carried out using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) and standard protocols. IgGs were purified from the conditioned media directly on 1 ml HiTrap protein A columns according to the manufacturer's instructions (APBiotech).

6.6 CHARACTERIZATION OF MUTATED HINGE-FC REGION

6.6.1 IN VITRO CHARACTERIZATION HPLC AND SDS-PAGE

5 Following the purification, general characteristics such as molecular weight and bonding characteristics of the modified hinge-Fc fragments may be studied by various methods well known to those skilled in the art, including SDS-PAGE and HPLC.

FcRn binding assay

10 Binding activity of modified hinge-Fc fragments can be measured by incubating radio-labeled wild-type hinge-Fc or modified hinge-Fc with the cells expressing either mouse or human FcRn. Typically, endothelial cell lines such as SV40 transformed endothelial cells (SVEC) (Kim *et al.*, *J. Immunol.*, 40:457-465, 1994) are used. After incubation with the hinge-Fc fragments at 37°C for 16-18 hours, the cells are washed with
15 medium and then detached by incubation with 5 mM Na₂EDTA in 50 mM phosphate buffer, pH 7.5, for 5 minutes. The radioactivity per 10⁷ cells is measured.

 Then, the cells are resuspended in 2 ml of 2.5 mg/ml CHAPS, 0.1 M Tris-HCl pH 8.0 containing 0.3 mg/ml PMSF, 25 mg/ml pepstatin and 0.1 mg/ml aprotinin and incubated for 30 minutes at room temperature. The cell suspension is then centrifuged and
20 the supernatant separated. The radioactivity of the supernatant is measured and used to calculate the amount of the hinge-Fc fragments extracted per 10⁷ cells.

 The K_d for the interaction of wild type human IgG1 with murine and human FcRn (269 and 2527 nM, respectively) agree well with the values determined by others (265 and 2350 nM, respectively, Firan *et al.*, 2001, *supra*). The I253A mutation virtually
25 abolishes binding to human and murine FcRn, as reported by others (Kim *et al.*, *Eur. J. Immunol.*, 29:2819-2825, 1991; Shields *et al.*, *J. Biol. Chem.*, 276:6591-6604, 2001). This is not the result of misfolding of the antibody as this mutant retains the same specific activity than the wild type molecule (SYNAGIS®) in a microneutralization assay (Johnson *et al.*, 1997, *supra*; data not shown).

30 Human IgG1 mutants with increased binding affinity towards both murine and human FcRn were generated (Table VIII). Improvements in complex stability were overall less marked for the human IgG1-human FcRn pair than for the human IgG1-murine FcRn compared to wild type IgG1 were 30-($\Delta\Delta G = 2.0$ kcal/mol for N434F/Y436H) and 11-($\Delta\Delta G = 1.4$ kcal/mol for M252Y/S254Y/S254T/T256E) fold, respectively. However, ranking of
35 the most critical positions remain unchanged when comparing human and murine FcRn: the largest increases in IgG1-murine FcRn complex stability ($\Delta\Delta G > 1.3$ kcal/mol) occurred on

mutations at positions 252, 254, 256 (M252Y/S254T/T256E and M252W) and 433, 434, 436 (H433K/N434F/Y436H and N434F/Y436H). Likewise, the same mutations were found to have the most profound impact on the IgG1-human FcRn interaction and also resulted in the largest increases in complex stability ($\Delta\Delta G > 1.0$ kcal/mol). Substitutions at positions 308, 309, 311, 385, 386, 387 and 389 had little or no effect on the stability of the complexes involving human or murine FcRn ($\Delta\Delta G < 0.5$ kcal/mol). Residues at the center of the Fc-FcRn combining site contribute significantly more to improvement in complex stability than residues at the periphery (FIG. 9).

Efficient binding of human Fc to murine FcRn apparently requires the presence of several wild type Fc residues. For example, leucine is very conserved at 251, arginine at 255, aspartic acid at 310, leucine at 314 and methionine at 428 (FIG. 6). Another specificity trend is observed when one considers positions 308, 309, and 311 where threonine, proline, and serine, respectively, are very strongly favored over the corresponding wild type residues (FIG. 6). However, generation of this strong consensus sequences does not correlate with the magnitude of increase in affinity as V308T/L309P/Q311S binds less than 2-fold better than the wild type IgG1 to both human and murine FcRn (Table IX).

Increases in affinity can be strongly dependent upon residue substitution at one 'hot spot' position. For example, the single mutation M252Y causes an increase in binding to murine FcRn by 9-fold, whereas additional mutations bring little (M252Y/S254T/T256E) or no (M252Y/T256Q) added benefit. The same trend is observed for the human receptor, although to a lesser extent. Indeed, M252Y/S254T/T256E shows a marked improvement of 2.5-fold in affinity compared to M252Y. This probably reflects the differences between the binding site of human and murine FcRn (West and Bjorkman, *Biochemistry*, 39:9698-9708, 2000).

Phage-derived IgG1 mutants exhibiting a significant increase in affinity towards murine FcRn ($\Delta\Delta G > 1.3$ kcal/mol) also showed significant binding activity to the receptor at pH 7.2 when compared to wild type IgG1 (FIGs. 8A-H). IgG1 mutants with moderate increase in affinity ($\Delta\Delta G < 0.3$ kcal/mol) bound very poorly at pH 7.2 (data not shown). In contrast, IgG1 mutants with large ($\Delta\Delta G > 1.0$ kcal/mol) increase in affinity towards human FcRn exhibited only minimal binding at pH 7.4 when compared to wild type IgG1 (FIGs. 8A-H).

Table IX
DISSOCIATION CONSTANTS AND RELATIVE FREE ENERGY
CHANGES FOR THE BINDING OF IgG1/FC MUTANTS TO
MURINE AND HUMAN FcRn*

MUTANT	Dissociation Constant Fc/Murine FcRn (nM)	$\Delta\Delta G$ (kcal/mol)	Dissociation Constant Fc/Human FcRn (mM)	$\Delta\Delta G$ (kcal/mol)
wild type	269 ± 1		2527 ± 117	
I253A	NB	NA	NB	NA
M252Y/S254T/T256E	27 ± 6	1.4	225 ± 10	1.4
M252W	30 ± 1	1.3	408 ± 24	1.1
M252Y	41 ± 7	1.1	532 ± 37	0.9
M252Y/T256Q	39 ± 8	1.1	560 ± 102	0.9
M252F/T256D	52 ± 9	1.0	933 ± 170	0.6
V308T/L309P/Q311S	153 ± 23	0.3	1964 ± 84	0.1
G385D/Q386P/N389S	187 ± 10	0.2	2164 ± 331	0.1
G385R/Q386T/P387R/N389P	147 ± 24	0.4	1620 ± 61	0.3
H433K/N434F/Y436H	14 ± 2	1.8	399 ± 47	1.1
N434F/Y436H	9 ± 1	2.0	493 ± 7	1.0

*Affinity measurements were carried out by BIAcore as described above. Residue numbering is according to EU (Kabat *et al.*, 1991, *supra*). Differences in free energy changes are calculated as the differences between the ΔG s of wild type and mutant reactions ($\Delta\Delta G = \Delta G_{\text{wild type}} - \Delta G_{\text{mutant}}$). NB, no binding. NA, not-applicable.

FcRn-mediated transfer assay

This assay follows the protocol disclosed in PCT publication WO 97/34631. Radiolabeled modified hinge-Fc fragments at various concentration (1 $\mu\text{g/ml}$ -1 mg/ml) are added to the one side of the transwell and the transfer of the fragments mediated by FcRn-expressing monolayer of the cells can be quantitated by measuring the radioactivity on the other side of the transwell.

6.6.2 IN VIVO PHARMACOKINETIC STUDY

In order to determine the half-life of the modified IgG hinge-Fc, modified hinge-Fc fragments are radiolabelled with ^{125}I (approximate specific activity of 10^7 cpm/ μg) and dissolved in saline (pH 7.2). The solution is injected intravenously into BALB/c mice (Harlan, Indianapolis, IN), which have been given NaI-containing water previously to block the thyroid, in a volume not more than 150 μl and with a radioactivity of 10×10^6 - 50×10^6 cpm. The mice are bled from the retro-orbital sinus at various time points, for example, at 3 minutes to 72 hours after the injection, into heparinized capillary tubes and the plasma collected from each sample is counted for radioactivity.

To generate the data provided in FIG. 10, 10 animals were used for each molecule assayed with 2.5 μg of antibody injected per animal. Antibody serum levels were determined using an anti-human IgG ELISA (FIG. 10). There seems to be an inverse correlation between affinity to mouse FcRn and persistence in serum. This might be due to the significant amount of binding of the mutants observed at pH 7.2, which leads to the sequestration (i.e., lack of release in the serum) of the molecules. Preliminary data (not shown) suggests increased transport of the mutants to the lung. Additionally, since the mutants exhibit lower levels of binding to human FcRn than murine FcRn (see FIGS. 8A-H), antibody serum levels are expected to be higher in primates and humans.

6.6.3 SURFACE PLASMON RESONANCE ANALYSES

The interaction of soluble murine and human FcRn with immobilized human IgG1 variants was monitored by surface plasmon resonance detection using a BIAcore 3000 instrument (Pharmacia Biosensor, Uppsala, Sweden). No aggregated material which could interfere with affinity measurements (van der Merwe *et al.*, *EMBO J.*, 12:4945-4954, 1993; van der Merwe *et al.*, *Biochemistry*, 33:10149-10160, 1994) was detected by gel filtration. Protein concentrations were calculated by the bicinchoninic acid (BCA) method for both human and murine FcRn or using the 1% extinction coefficient at 280 nm of 1.5 for IgG1 wild type and variants. The latter were coupled to the dextran matrix of a CM5 sensor chip (Pharmacia Biosensor) using an Amine Coupling Kit as described (Johnson *et al.*, *supra*). The protein concentrations ranged from 3-5 $\mu\text{g}/\text{ml}$ in 10 mM sodium acetate, pH 5.0. The activation period was set for 7 minutes at a flow rate of 10 $\mu\text{l}/\text{min}$ and the immobilization period was set to between 10 and 20 minutes at a flow rate of 10 $\mu\text{l}/\text{min}$. Excess reactive esters were quenched by injection of 70 μl of 1.0 methanolamine hydrochloride, pH 8.5. This typically resulted in the immobilization of between 500 and 4000 resonance units (RU). Human and murine FcRn were buffer exchanged against 50 mM PBS buffer pH 6.0

containing 0.05% Tween 20. Dilutions were made in the same buffer. All binding experiments were performed at 25°C with concentrations ranging from 120 to 1 µg/ml at a flow rate of 5 to 10 µl/min; data were collected for 25 to 50 minutes and three 1-minute pulses of PBS buffer pH 7.2 were used to regenerate the surfaces. FcRn was also flowed over an uncoated cell and the sensorgrams from these blank runs subtracted from those obtained with IgG1-coupled chips. Runs were analyzed using the software BIAevaluation 3.1 (Pharmacia). Association constants (K_A s) were determined from Scatchard analysis by measuring the concentration of free reactants and complex at equilibrium after correction for nonspecific binding. In equilibrium binding BIAcore experiments (Karlsson *et al.*, 1991, *supra*; van der Merwe *et al.*, 1993, *supra*; van der Merwe *et al.*, 1994, *supra*; Raghavan *et al.*, *Immunity*, 1:303-315, 1994; Malchiodi *et al.*, *J. Exp. Med.*, 182:1833-1845, 1995), the concentration of the complex can be assessed directly as the steady-state response. The concentration of free analyte (human or murine FcRn) is equal to the bulk analyte concentration since analyte is constantly replenished during sample injection. The concentration of free ligand on the surface of the sensor chip can be derived from the concentration of the complex and from the total binding capacity of the surface as $K_A = R_{eq}/C(R_{max} - R_{eq})$ where C is the free analyte concentration, R_{eq} is the steady-state response, and R_{max} is the total surface binding capacity. Rearranging, the equation reads: $R_{eq}/C = K_A R_{max} - K_A R_{eq}$.

A plot of R_{eq}/C versus R_{eq} at different analyte concentrations thus gives a straight line from which K_A can be calculated (see Table IX). Errors were estimated as the standard deviation for two or three independent determinations and were <20%.

Representative mutations identified after panning libraries 1 through 4 (FIG. 6, Table VIII) were introduced into the Fc portion of a human IgG1. Injection of different concentrations of human or murine FcRn over the immobilized IgG1 variants gave concentration-dependent binding. Typical resonance profiles for equilibrium binding of the mutant M252Y/S254T/T256E to murine and human FcRn are shown in FIGs. 7A and B. To estimate apparent K_A s, concentrations of FcRn ranging from 120 to 1 µg/ml were used. In all cases, equilibrium (or near-equilibrium) binding levels were reached within 50 minutes. To estimate the increase in RU resulting from the non specific effect of protein on the bulk refractive index, binding of FcRn to an uncoated cell was measured and the sensorgrams from these blank runs subtracted from those obtained with IgG1-coupled chips. The scatchard plots for the binding of the mutant M252Y/S254T/T256E to murine and human FcRn are shown in FIGs. 7C and D. The plots were all linear, and apparent K_A s were

calculated from the relevant slopes. Measurements were carried out in duplicate or triplicate and confirmed that the immobilized IgGs retained their original binding activity.

Since there are two non-equivalent binding sites on mouse IgG1 for murine FcRn with affinities of < 130 nM and $6 \mu\text{M}$ (Sanchez *et al.*, *Biochemistry*, 38:9471-9476, 1999; Schuck *et al.*, *Mol. Immunol.*, 36:1117-1125, 1999; Ghetie and Ward, *Ann. Rev. Immunol.*, 18:739-766, 2000), the receptor was used in solution to avoid avidity effects that arise when IgG1 binds to immobilized FcRn. Consistent with this, systematically higher affinities are observed when FcRn, rather than IgG, immobilized on the biosensor chip (Popov *et al.*, 1996, *supra*; Vaughn and Bjorkman, *Biochemistry*, 36:9374-9380, 1997; Martin and Bjorkman, *Biochemistry*, 38:12639-12647; West and Bjorkman, *Biochemistry*, 39:9698-9708, 2000). Under our experimental BIAcore conditions, mainly interactions corresponding to the higher-affinity association (*i.e.* single liganded-receptor) are measured, according for the linearity of the scatchard plots (FIGs. 7C and D).

BIAcore analysis was also used to compare the affinity of wild type IgG1 and IgG1 mutants. Phage-derived IgG1 mutants exhibiting a significant increase in affinity towards murine FcRn at pH 6.0 ($\Delta\Delta G \geq 1.0$ kcal/mol) also showed significant binding to the mouse receptor at pH 7.2 with $\text{SPR signal}_{\text{pH}7.4} / \text{SPR signal}_{\text{pH}6.0} > 0.6$ at saturation. IgG1 mutants with moderate increase in affinity towards murine FcRn at pH 6.0 ($\Delta\Delta G < 0.4$ kcal/mol) bound very poorly to the mouse receptor at pH 7.2. In contrast, IgG1 mutants exhibiting large affinity increase towards human FcRn at pH 6.0 ($\Delta\Delta G \geq 1.0$ kcal/mol) only showed minimal binding to the human receptor at pH 7.4 with $\text{SPR signal}_{\text{pH}7.4} / \text{SPR signal}_{\text{pH}6.0} < 0.15$ at saturation.

Those skilled in the art will recognize, or be able to ascertain using no more routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

All publications, patents and patent applications mentioned in this specification are herein incorporated by reference into the specification to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A modified IgG comprising an IgG constant domain, or an FcRn-binding
fragment thereof, wherein the constant domain or fragment comprises a
5 substitution at amino acid residue 252 relative to a wild-type IgG constant
domain, numbered according to the EU numbering index of Kabat, wherein the
modified IgG has an increased half-life compared to the half-life of an IgG
having the wild-type IgG constant domain, and wherein the substitution at
amino acid residue 252 is a substitution with tyrosine, phenylalanine,
10 tryptophan, or threonine.
2. The modified IgG according to claim 1, further comprising one or more amino
acid substitutions at one or more of amino acid residues 251, 253-256,
285-290, 308-314, 385-389, and 428-436 relative to a wild-type IgG constant
domain.
- 15 3. The modified IgG according to claim 1, further comprising one or more amino
acid substitutions at one or more of amino acid residues 254, 256, 309, 311,
314, 428, 433, 434, or 436 relative to a wild-type IgG constant domain.
4. The modified IgG according to claim 1, further comprising an amino acid
substitution at amino acid residue 254 and an amino acid substitution at amino
20 acid residue 256.
5. The modified IgG according to claim 3 or 4, further comprising one or more
amino acid substitutions at one or more of amino acid residues 385, 386, 387,
or 389.
6. The modified IgG according to claim 4, wherein an amino acid substitution at
25 amino acid residue 254 does not include a substitution with serine.
7. The modified IgG according to claim 4 or 6, wherein an amino acid
substitution at amino acid residue 256 does not include a substitution with
phenylalanine.

8. The modified IgG according to claim 2 or 3, wherein the amino acid substitution at amino acid residue 254 is a substitution with threonine, the amino acid substitution at amino acid residue 256 is a substitution with serine, arginine, glutamine, glutamic acid, aspartic acid, or threonine, the amino acid substitution at amino acid residue 309 is a substitution with proline, the amino acid substitution at amino acid residue 311 is a substitution with serine, the amino acid substitution at amino acid residue 433 is a substitution with arginine, serine, isoleucine, proline, or glutamine, and the amino acid substitution at amino acid residue 434 is a substitution with histidine, phenylalanine, or tyrosine.
9. The modified IgG according to claim 2 or 3, wherein the amino acid substitution at amino acid residue 252 is a substitution with tyrosine, the amino acid substitution at amino acid residue 254 is a substitution with threonine, the amino acid substitution at amino acid residue 256 is glutamic acid, the amino acid substitution at amino acid residue 433 is a substitution with lysine, the amino acid substitution at amino acid residue 434 is a substitution with phenylalanine, and the amino acid substitution at amino acid residue 436 is a substitution with histidine.
10. The modified IgG according to claim 2 or 3, the amino acid substitution at amino acid residue 254 is a substitution with threonine, the amino acid substitution at amino acid residue 256 is a substitution with serine, arginine, glutamine, glutamic acid, aspartic acid, alanine, asparagine, or threonine, the amino acid substitution at amino acid residue 309 is a substitution with proline, the amino acid substitution at amino acid residue 311 is a substitution with serine, the amino acid substitution at amino acid residue 433 is a substitution with arginine, serine, isoleucine, proline, or glutamine, and the amino acid substitution at amino acid residue 434 is a substitution with histidine, phenylalanine, or tyrosine.
11. The modified IgG according to claim 4, wherein the amino acid substitution at amino acid residue 254 is a substitution with threonine and the amino acid

substitution at amino acid residue 256 is a substitution with serine, arginine, glutamine, glutamic acid, aspartic acid, or threonine.

12. The modified IgG according to claim 4, wherein the amino acid substitution at amino acid residue 252 is a substitution with tyrosine, the amino acid substitution at amino acid residue 254 is a substitution with threonine and the amino acid substitution at amino acid residue 256 is a substitution with glutamic acid.
13. The modified IgG according to claim 4, wherein the amino acid substitution at amino acid residue 254 is a substitution with threonine, and the amino acid substitution at amino acid residue 256 is a substitution with glutamic acid.
14. The modified IgG according to claim 5, wherein the amino acid substitution at amino acid residue 385 is a substitution with arginine, aspartic acid, serine, threonine, histidine, lysine, alanine or glycine, the amino acid substitution at amino acid residue 386 is a substitution with threonine, proline, aspartic acid, serine, lysine, arginine, isoleucine, or methionine, the amino acid substitution at amino acid residue 387 is a substitution with arginine, proline, histidine, serine, threonine, or alanine, and the amino acid substitution at amino acid residue 389 is a substitution with proline, serine or asparagine.
15. The modified IgG according to claim 2 or 3, wherein the amino acid substitution at amino acid residue 428 is a substitution with methionine, threonine, leucine, phenylalanine or serine.
16. The modified IgG according to claim 2 or 3, wherein the amino acid substitution at amino acid residue 314 is a substitution with leucine.
17. The modified IgG according to claim 2, wherein an amino acid residue substitution at amino acid residue 251 is a substitution with leucine or arginine an amino acid substitution at amino acid residue 254 is a substitution with threonine, an amino acid substitution at amino acid residue 255 is a substitution with leucine, glycine, isoleucine or arginine, an amino acid substitution at amino acid residue 256 is a substitution with serine, arginine, glutamine, glutamic acid, aspartic acid, or threonine, an amino acid substitution at amino

- acid residue 308 is a substitution with a threonine, an amino acid substitution at amino acid residue 309 is a substitution with proline, an amino acid substitution at amino acid residue 311 is a substitution with serine, an amino acid substitution at amino acid residue 312 is a substitution with aspartic acid, an amino acid substitution at amino acid residue 314 is a substitution with leucine, an amino acid substitution at amino acid residue 385 is a substitution with arginine, aspartic acid, serine, threonine, histidine, lysine or alanine, an amino acid substitution at amino acid residue 386 is a substitution with threonine, proline, aspartic acid, serine, lysine, arginine, isoleucine or methionine, an amino acid substitution at amino acid residue 387 is a substitution with arginine, histidine, serine, threonine, alanine or proline, an amino acid substitution at amino acid residue 389 is a substitution with proline or serine, an amino acid substitution at amino acid residue 433 is a substitution with arginine, serine, isoleucine, proline, glutamine or histidine, and an amino acid substitution at amino acid residue 434 is a substitution with phenylalanine, tyrosine or histidine, an amino acid substitution at amino acid residue 436 is a substitution with histidine, asparagine, arginine, threonine, lysine or methionine, and an amino acid substitution at amino acid residue 428 is a substitution with methionine, threonine, leucine, phenylalanine or serine.
18. The modified IgG according to any one of the preceding claims, wherein the modified IgG constant domain or FcRn-binding fragment has a higher affinity for FcRn than a wild-type IgG constant domain or FcRn-binding fragment thereof.
19. The modified IgG according to claim 18, wherein the modified IgG constant domain or FcRn-binding fragment has a higher affinity for FcRn than a wild-type IgG constant domain or FcRn-binding fragment thereof at pH 6.0 than at pH 7.4.
20. The modified IgG according to any one of the preceding claims, wherein the IgG constant domain, or FcRn-binding fragment thereof, is a human IgG constant domain, or an FcRn binding fragment thereof.

21. The modified IgG according to claim 20, wherein the modified IgG is a human or humanized IgG.
22. The modified IgG according to any one of claims 1-19, wherein the IgG constant domain, or FcRn-binding fragment thereof, is a non-human IgG constant domain or an FcRn-binding fragment thereof.
23. The modified IgG according to claim 22, wherein the modified IgG is a non-human IgG.
24. The modified IgG according to claim 21, wherein the modified IgG is a humanized anti-CD25 monoclonal antibody, a chimeric anti-CD25 monoclonal antibody, a humanized anti- $\alpha_v\beta_3$ integrin antibody, or a humanized anti-CD3 antibody.
25. The modified IgG according to claim 22, wherein the non-human IgG constant domain or FcRn-binding fragment is IgG₁, IgG_{2a}, IgG_{2b}, IgG_{2c} or IgG₃ or an FcRn-binding fragment thereof.
26. The modified IgG according to claim 22, wherein the non-human IgG constant domain or FcRn-binding fragment is a rodent, donkey, sheep, rabbit, goat, guinea pig, camel, horse or chicken IgG constant domain or an FcRn-binding fragment thereof.
27. The modified IgG according to claim 23, wherein the non-human Ig is a rodent, donkey, sheep, rabbit, goat, guinea pig, camel, horse or chicken Ig.
28. The modified IgG according to claim 20, wherein the human IgG constant domain or FcRn-binding fragment is IgG₁, IgG₂, IgG₃ or IgG₄ or an FcRn-binding fragment thereof.
29. The modified IgG according to claim 28, wherein the human IgG constant domain or FcRn-binding fragment is IgG₁ or an FcRn-binding fragment thereof.
30. The modified IgG of claim 21, wherein the human IgG is IgG₁, IgG₂, IgG₃ or IgG₄.

31. The modified IgG according to any one of the preceding claims which immunospecifically binds to an RSV antigen.
32. A modified IgG according to any one of claims 1-21 or 28-30 which comprises a heavy chain variable domain and a light chain variable domain of one of the following antibodies: palivizumab (SYNAGIS[®]), AFFF, p12f2, p12f4, p11d4, Ale109, A12a6, A13c4, A17d4, A4B4, A8C7, 1X 493L1FR, H3-3F4, M3H9, Y10H6, DG, AFFF(1), 6H8, L1-7E5, L215B10, A13A11, A1H5, A4B4(1), A4B4L1FR-S28R or A4B4-F52S, wherein the modified IgG immunospecifically binds to the RSV F antigen.
33. The modified IgG according to claim 32, which comprises the heavy chain variable domain and light chain variable domain of palivizumab (SEQ ID NOS.: 7 and 8).
34. The modified IgG according to claim 31, which comprises the variable heavy (VH) complementarily determining region (CDR)1, VH CDR2, VH CDR3, variable light (VL) CDR1, VL CDR2 and VL CDR3 of palivizumab (SEQ ID NOS.: 1-6, respectively).
35. The modified IgG according to claim 31, which comprises the VH CDR1, VH CDR2, or VH CDR3, VL CDR1, VL CDR2 and VL CDR3 of A4B4L1FR-S28R (SEQ ID NOS.: 10, 19, 20, 39, 5, and 6, respectively).
36. The modified Ig according to any one of claims 1-30, wherein the modified IgG immunospecifically binds to HER2, tumor necrosis factor-alpha (TNF- α), transforming growth factor- β (TGF- β), interleukin-4 (IL-4), interleukin-5 (IL-5), interleukin-8 (IL-8), CD2, CD3, CD4, CD11a, CD14, CD18, CD20, CD23, CD25, CD33, CD52, CD64, CD80, CD147, CD40 ligand (CD40L), vascular endothelial growth factor (VEGF), intracellular adhesion molecule-3 (ICAM-3), epithelial growth factor receptor (EGFR), $\alpha_v\beta_3$ integrin, $\alpha_4\beta_7$ integrin, human leukocyte antigen (HLA), complement factor 5 (C5), immunoglobulin E (IgE), glycoprotein II_b/III_a receptor, CA125, 17-IA cell surface antigen, Factor VII, GD3 epitope, human immunodeficiency glycoprotein 120 (HIV gp120), hepatitis B virus (HBV) or cytomeglovirus (CMV).

37. A pharmaceutical composition comprising the modified IgG according to any one of the preceding claims, and a pharmaceutically acceptable carrier.
38. A kit comprising the modified IgG of any one of the preceding claims, in a container, and instructions for use.
- 5 39. A nucleic acid comprising a nucleotide sequence encoding the modified IgG of any one of claims 32-35.
40. A nucleic acid comprising a nucleotide sequence encoding the modified IgG constant domain, or FcRn-binding fragment thereof, as defined in any one of claims 1-31.
- 10 41. A host cell comprising the nucleic acid of claim 39 or 40.
42. A method of preventing or treating a disease or disorder, comprising administering to a subject a modified IgG according to any one of the preceding claims.
- 15 43. A method of preventing or treating respiratory syncytial virus (RSV) infection, comprising administering to a subject the modified IgG according to any one of claims 31-35.
44. The method according to claim 42 or 43, wherein the subject is a human.
- 20 45. A method of diagnosing a disease or disorder, comprising: (a) contacting the modified IgG according to any one of claims 1-30 or 36 with a sample from a subject; and (b) comparing the level of the antigen to which the modified IgG immunospecifically binds to a control level, wherein an increase in level of the antigen in the sample compared to the control level is indicative of the disease or disorder.
- 25 46. A method of diagnosing a RSV infection, comprising: (a) contacting the modified IgG according to any one of claims 31-35 with a sample from a subject; and (b) comparing the level of the RSV antigen to which the modified IgG immunospecifically binds to a control level, wherein an increase in level of

the RSV antigen in the sample compared to the control level is indicative of a RSV infection.

47. A method of diagnosing a disease or disorder, comprising: (a) administering to a subject an effective amount of labeled form of the modified IgG of any one of claims 1 to 30 or 36; (b) waiting for a time interval following the administration for permitting the labeled form of the modified IgG to preferentially concentrate at sites in the subject where the antigen to which the modified IgG immunospecifically binds is expressed; (c) determining background level; and (d) detecting the labeled form of the modified IgG in the subject, wherein detection of the labeled form of the modified IgG above the background level indicates that the subject has the disease or disorder.
48. A method of diagnosing a RSV infection, comprising: (a) administering to a subject an effective amount of labeled form of the modified IgG of any one of claims 31 to 35; (b) waiting for a time interval following the administration for permitting the labeled form of the modified IgG to preferentially concentrate at sites in the subject where the RSV antigen to which the modified IgG immunospecifically binds is expressed; (c) determining background level; and (d) detecting the labeled form of the modified IgG in the subject, wherein detection of the labeled form of the modified IgG above the background level indicates that the subject has a RSV infection.
49. The method according to claim 47 or 48, wherein the subject is in a human.
50. A fusion protein comprising a non-IgG polypeptide covalently linked to a modified IgG constant domain, or a fragment thereof that binds to FcRn, said modified IgG constant domain or fragment comprising a substitution at amino acid residue 252 relative to a wild-type IgG constant domain, numbered according to the EU Index of Kabat, wherein the modified IgG has an increased half-life compared to the half-life of an IgG having the wild-type IgG constant domain, and wherein the substitution at amino acid residue 252 is a substitution with tyrosine, phenylalanine, tryptophan, or threonine.

51. The fusion protein according to claim 50, wherein the modified IgG constant domain or fragment further comprises one or more amino acid substitutions at one or more of amino acid residues 251, 253-256, 258-290, 308-314, 385-389, and 428-436 relative to a wild-type IgG constant domain.
- 5 52. The fusion protein according to claim 50, wherein the modified IgG constant domain or fragment further comprises an amino acid substitution at amino acid residue 254 and an amino acid substitution at amino acid residue 256.
- 10 53. The fusion protein according to claim 52, wherein the amino acid substitution at amino acid residue 254 is a substitution with threonine and the amino acid substitution at amino acid residue 256 is a substitution with serine, arginine, glutamine, glutamic acid, aspartic acid or threonine.
- 15 54. The fusion protein according to claim 52, wherein the amino acid substitution at amino acid residues 252 is a substitution with tyrosine, the amino acid substitution at amino acid residue 254 is a substitution with threonine and the amino acid substitution at amino acid residue 256 is a substitution with glutamic acid.
55. The fusion protein of claim 51, wherein the amino acid substitution at amino acid residue 428 is a substitution with methionine, threonine, leucine, phenylalanine or serine.
- 20 56. The fusion protein according to claim 50-55, wherein the modified IgG constant domain or FcRn-binding fragment has a higher affinity for FcRn than a wild-type IgG constant domain or FcRn-binding fragment thereof.
- 25 57. The fusion protein according to claim 56, wherein the modified IgG constant domain or FcRn-binding fragment has a higher affinity for FcRn than a wild-type IgG constant domain or FcRn-binding fragment thereof at pH 6.0 than at pH 7.4.
58. A pharmaceutical composition comprising the fusion protein according to any one of claims 50-57, and a pharmaceutically acceptable carrier.

59. A kit comprising the fusion protein of any one of claims 50-57, in a container, and instructions for use.
60. A method of preventing or treating a disease or disorder in a subject, comprising administering to a subject the fusion protein according to any one of claims 50-57.
61. A method of diagnosing a disease or disorder, comprising: (a) contacting the fusion protein according to any one of claims 50-57 with a sample from a subject; and (b) comparing the level of the antigen to which the fusion protein immunospecifically binds with a control level, wherein an increase in the level of antigen in the sample compared to the control level is indicative of the disease or disorder.
62. A method of diagnosing a disease or disorder, comprising: (a) administering to a subject an effective amount of labeled form of the fusion protein of any one of claims 50 to 57; (b) waiting for a time interval following the administration for permitting the labeled form of the fusion protein to preferentially concentrate at sites in the subject where the antigen to which the fusion protein immunospecifically binds is expressed; (c) determining background level; and (d) detecting the labeled form of the fusion protein in the subject, wherein detection of the labeled form of the fusion protein above the background level indicates that the subject has the disease or disorder.
63. The method according to claim 60, 61 or 62, wherein the subject is a human.
64. A molecule comprising a non-protein agent conjugated to a modified IgG constant domain, or a fragment thereof that binds to FcRn, said modified IgG constant domain or fragment comprising a substitution at amino acid residue 252 relative to a wild-type IgG constant domain, numbered according to the EU numbering index of Kabat, wherein the modified IgG has an increased half-life compared to the half-life of an IgG having the wild-type IgG constant domain, and wherein the substitution at amino acid residue 252 is a substitution with tyrosine, phenylalanine, tryptophan, or threonine.

65. The molecule according to claim 64, wherein the modified IgG constant domain or fragment further comprises one or more amino acid substitutions at one or more of amino acid residues 251, 253-256, 258-290, 308-314, 385-389 and 428-436 relative to a wide-type IgG constant domain.
- 5 66. The molecule according to claim 64, wherein the modified IgG constant domain or fragment further comprises an amino acid substitution at amino acid residue 254 and an amino acid substitution at amino acid residue 256.
67. The molecule according to claim 66, wherein the amino acid substitution at amino acid residue 254 is a substitution with threonine and the amino acid substitution at amino acid residue 256 is a substitution with serine, arginine, glutamine, glutamic acid, aspartic acid or threonine.
- 10
68. The molecule according to claim 66, wherein the amino acid substitution at amino acid residues 252 is a substitution with tyrosine, the amino acid substitution at amino acid residue 254 is a substitution with threonine and the amino acid substitution at amino acid residue 256 is a substitution with glutamic acid.
- 15
69. The molecule of claim 66, wherein the amino acid substitution at amino acid residue 428 is a substitution with methionine, threonine, leucine, phenylalanine or serine.
- 20 70. The molecule according to any one of claims 64 or 69, wherein the modified IgG constant domain or FcRn-binding fragment has a higher affinity for FcRn than the wild-type IgG constant domain of FcRn-binding fragment thereof.
71. The molecule of claim 70, wherein the modified IgG constant domain or FcRn-binding fragment has a higher affinity for FcRn than a wild-type IgG constant domain or FcRn-binding fragment thereof at pH 6.0 than at pH 7.4.
- 25
72. A pharmaceutical composition comprising the molecule according to any one of claims 64-71, and a pharmaceutically acceptable carrier.

73. A kit comprising the molecule according to any one of claims 64-71, in a container, and instructions for use.
74. A method of preventing or treating a disease or disorder, comprising administering to a subject the molecule according to any one of claims 64-71.
- 5 75. A method of diagnosing a disease or disorder, comprising: (a) containing the molecule of any one of claims 64-71 with a sample from a subject; and (b) comparing the level of the antigen to which the molecule immunospecifically binds with a control level, wherein an increase in the level of antigen in the sample compared to the control level is indicative of the disease or disorder.
- 10 76. A method of diagnosing a disease or disorder, comprising: a) administering to a subject an effective amount of labeled form of the molecule of any one of claims 64 to 71; b) waiting for a time interval following the administration for permitting the labeled form of the molecule to preferentially concentrate at sites in the subject where the antigen to which the molecule
- 15 immunospecifically binds is expressed; c) determining background level; and d) detecting the labeled form of the molecule in the subject, wherein detection of the labeled form of the molecule above the background level indicates that the subject has the disease or disorder.
77. The method according to claim 74, 75 or 76, wherein the subject is a human.
- 20 78. The modified IgG according to claim 1, substantially as hereinbefore described and with reference to any of the Examples and/or Figures.
79. Use of a modified IgG according to any one of claims 1 to 36 in the manufacture of a medicament for preventing or treating a disease or condition.
80. Use of a fusion protein according to any one of claims 50 to 57 in the
- 25 manufacture of a medicament for preventing or treating a disease or condition.
81. Use of a molecule according to any one of claims 64 to 71 in the manufacture of a medicament for preventing or treating a disease or condition.

82. The use according to any one of claims 79 to 81, wherein the disease or condition is respiratory syncytial virus (RSV) infection.

1/11




FIG. 1

2/11

```

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1           5           10           15
|-----Hinge-----|-----
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
          20           25           30
-----
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
          35           40           45
-----
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
          50           55           60
-----
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
65           70           75           80
-----
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
          85           90           95
-----
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
          100          105          110
-----
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
          115          120          125
-----
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
          130          135          140
-----
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
145          150          155          160
-----
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
          165          170          175
-----
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
          180          185          190
-----
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
          195          200          205
-----
Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
          210          215          220
-----
Ser Leu Ser Leu Ser Pro Gly Lys
225          230
-----

```

FIG.2

3/11

Met	Gly	Val	Pro	Arg	Pro	Gln	Pro	Trp	Ala	Leu	Gly	Leu	Leu	Leu	Phe
1				5					10					15	
Leu	Leu	Pro	Gly	Ser	Leu	Gly	Ala	Glu	Ser	His	Leu	Ser	Leu	Leu	Tyr
			20					25					30		
His	Leu	Thr	Ala	Val	Ser	Ser	Pro	Ala	Pro	Gly	Thr	Pro	Ala	Phe	Trp
		35					40					45			
Val	Ser	Gly	Trp	Leu	Gly	Pro	Gln	Gln	Tyr	Leu	Ser	Tyr	Asn	Ser	Leu
	50					55					60				
Arg	Gly	Glu	Ala	Glu	Pro	Cys	Gly	Ala	Trp	Val	Trp	Glu	Asn	Gln	Val
65					70					75					80
Ser	Trp	Tyr	Trp	Glu	Lys	Glu	Thr	Thr	Asp	Leu	Arg	Ile	Lys	Glu	Lys
				85					90					95	
Leu	Phe	Leu	Glu	Ala	Phe	Lys	Ala	Leu	Gly	Gly	Lys	Gly	Pro	Tyr	Thr
			100					105					110		
Leu	Gln	Gly	Leu	Leu	Gly	Cys	Glu	Leu	Gly	Pro	Asp	Asn	Thr	Ser	Val
	115					120						125			
Pro	Thr	Ala	Lys	Phe	Ala	Leu	Asn	Gly	Glu	Glu	Phe	Met	Asn	Phe	Asp
	130					135					140				
Leu	Lys	Gln	Gly	Thr	Trp	Gly	Gly	Asp	Trp	Pro	Glu	Ala	Leu	Ala	Ile
145					150					155					160
Ser	Gln	Arg	Trp	Gln	Gln	Gln	Asp	Lys	Ala	Ala	Asn	Lys	Glu	Leu	Thr
				165					170					175	
Phe	Leu	Leu	Phe	Ser	Cys	Pro	His	Arg	Leu	Arg	Glu	His	Leu	Glu	Arg
			180					185					190		
Gly	Arg	Gly	Asn	Leu	Glu	Trp	Lys	Glu	Pro	Pro	Ser	Met	Atg	Leu	Lys
		195					200					205			
Ala	Arg	Pro	Ser	Ser	Pro	Gly	Phe	Ser	Val	Leu	Thr	Cys	Ser	Ala	Phe
	210					215					220				
Ser	Phe	Tyr	Pro	Pro	Glu	Leu	Gln	Leu	Arg	Phe	Leu	Arg	Asn	Gly	Leu
225					230					235					240
Ala	Ala	Gly	Thr	Gly	Gln	Gly	Asp	Phe	Gly	Pro	Asn	Ser	Asp	Gly	Ser
				245					250					255	
Phe	His	Ala	Ser	Ser	Ser	Leu	Thr	Val	Lys	Ser	Gly	Asp	Glu	His	His
			260					265					270		
Tyr	Cys	Cys	Ile	Val	Gln	His	Ala	Gly	Leu	Ala	Gln	Pro	Leu	Arg	Val
	275					280						285			
Glu	Leu	Glu	Ser	Pro	Ala	Lys	Ser	Ser	Val	Leu	Val	Val	Gly	Ile	Val
	290					295					300				
Ile	Gly	Val	Leu	Leu	Leu	Thr	Ala	Ala	Ala	Val	Gly	Gly	Ala	Leu	Leu
305					310					315					320
Trp	Arg	Arg	Met	Arg	Ser	Gly	Leu	Pro	Ala	Pro	Trp	Ile	Ser	Leu	Arg
				325					330					335	
Gly	Asp	Asp	Thr	Gly	Val	Leu	Leu	Pro	Thr	Pro	Gly	Glu	Ala	Gln	Asp
			340					345					350		
Ala	Asp	Leu	Lys	Asp	Val	Asn	Val	Ile	Pro	Ala	Thr	Ala			
		355				360						365			

FIG.3A

4/11

Met	Gly	Met	Pro	Leu	Pro	Trp	Ala	Leu	Ser	Leu	Leu	Leu	Val	Leu	Leu
1				5					10					15	
Pro	Gln	Thr	Trp	Gly	Ser	Glu	Thr	Arg	Pro	Pro	Leu	Met	Tyr	His	Leu
			20					25					30		
Thr	Ala	Val	Ser	Asn	Pro	Ser	Thr	Gly	Leu	Pro	Ser	Phe	Trp	Ala	Thr
		35					40					45			
Gly	Trp	Leu	Gly	Pro	Gln	Gln	Tyr	Leu	Thr	Tyr	Asn	Ser	Leu	Arg	Gln
	50					55					60				
Glu	Ala	Asp	Pro	Cys	Gly	Ala	Trp	Val	Trp	Glu	Asn	Gln	Val	Ser	Trp
65					70					75					80
Tyr	Trp	Glu	Lys	Glu	Thr	Thr	Asp	Leu	Lys	Ser	Lys	Glu	Gln	Leu	Phe
			85						90					95	
Leu	Glu	Ala	Leu	Lys	Thr	Leu	Glu	Lys	Ile	Leu	Asn	Gly	Thr	Tyr	Thr
			100					105					110		
Leu	Gln	Gly	Leu	Leu	Gly	Cys	Glu	Leu	Ala	Ser	Asp	Asn	Ser	Ser	Val
	115					120						125			
Pro	Thr	Ala	Val	Phe	Ala	Leu	Asn	Gly	Glu	Glu	Phe	Met	Lys	Phe	Asn
	130					135					140				
Pro	Arg	Ile	Gly	Asn	Trp	Thr	Gly	Glu	Trp	Pro	Glu	Thr	Glu	Ile	Val
145					150					155					160
Ala	Asn	Leu	Trp	Met	Lys	Gln	Pro	Asp	Ala	Ala	Arg	Lys	Glu	Ser	Glu
			165						170					175	
Phe	Leu	Leu	Asn	Ser	Cys	Pro	Glu	Arg	Leu	Leu	Gly	His	Leu	Glu	Arg
			180					185					190		
Gly	Arg	Arg	Asn	Leu	Glu	Trp	Lys	Glu	Pro	Pro	Ser	Met	Arg	Leu	Lys
	195						200					205			
Ala	Arg	Pro	Gly	Asn	Ser	Gly	Ser	Ser	Val	Leu	Thr	Cys	Ala	Ala	Phe
	210					215					220				
Ser	Phe	Tyr	Pro	Pro	Glu	Leu	Lys	Phe	Arg	Phe	Leu	Arg	Asn	Gly	Leu
225					230					235					240
Ala	Ser	Gly	Ser	Gly	Asn	Cys	Ser	Thr	Gly	Pro	Asn	Gly	Asp	Gly	Ser
			245						250				255		
Phe	His	Ala	Trp	Ser	Leu	Leu	Glu	Val	Lys	Arg	Gly	Asp	Glu	His	His
		260						265					270		
Tyr	Gln	Cys	Gln	Val	Glu	His	Glu	Gly	Leu	Ala	Gln	Pro	Leu	Thr	Val
	275						280					285			
Asp	Leu	Asp	Ser	Ser	Ala	Arg	Ser	Ser	Val	Pro	Val	Val	Gly	Ile	Val
	290					295				300					
Leu	Gly	Leu	Leu	Leu	Val	Val	Val	Ala	Ile	Ala	Gly	Gly	Val	Leu	Leu
305					310					315					320
Trp	Gly	Arg	Met	Arg	Ser	Gly	Leu	Pro	Ala	Pro	Trp	Leu	Ser	Leu	Ser
			325						330				335		
Gly	Asp	Asp	Ser	Gly	Asp	Leu	Leu	Pro	Gly	Gly	Asn	Leu	Pro	Pro	Glu
			340					345				350			
Ala	Glu	Pro	Gln	Gly	Ala	Asn	Ala	Phe	Pro	Ala	Thr	Ser			
		355				360						365			

FIG.3B

5/11


```

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1          5          10          15
-----Hinge-----|-----
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
          20          25          30
-----
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
          40          45          50
-----
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val
          50          55          60
-----CH2-----
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
65          70          75          80
-----
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
          85          90          95
-----
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala
          100          105          110
-----
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
          115          120          125
-----|-----
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr
          130          135          140
-----
Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
145          150          155          160
-----CH3-----
Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
          165          170          175
-----
Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
          180          185          190
-----
Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
          195          200          205
-----
Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
          210          215          220
-----
Ser Leu Ser Leu Ser Pro Gly Lys
225          230
-----|-----

```

FIG.4

6/11



7/11




FIG.6

8/11




FIG. 7B




FIG. 7D




FIG. 7A




FIG. 7C




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D




FIG. 8E




FIG. 8F




FIG. 8G




FIG. 8H

10/11




FIG.9

11/11




FIG.10

2002248184 15 Sep 2017

SEQUENCE LISTING

<110> MEDIMMUNE, LLC
BOARD OF REGENTS, THE UNIVERSITY OF TEXAS SYSTEM

<120> MOLECULES WITH EXTENDED HALF-LIVES, COMPOSITIONS AND USES
THEREOF

<130> 10271-027-228

<140> PCT/US01/48432

<141> 2001-12-12

<150> 60/254,884

<151> 2000-12-12

<150> 60/289,760

<151> 2001-05-09

<160> 124

<170> PatentIn version 3.1

<210> 1

<211> 7

<212> PRT

<213> Homo sapiens

<400> 1

Thr Ser Gly Met Ser Val Gly
1 5

<210> 2

<211> 16

<212> PRT

<213> Homo sapiens

2002248184 15 Sep 2017

<400> 2

Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser Leu Lys Ser

1

5

10

15

5

<210> 3

<211> 10

<212> PRT

<213> Homo sapiens

10

<400> 3

Ser Met Ile Thr Asn Trp Tyr Phe Asp Val

1

5

10

<210> 4

15

<211> 10

<212> PRT

<213> Homo sapiens

<400> 4

20

Lys Cys Gln Leu Ser Val Gly Tyr Met His

1

5

10

<210> 5

<211> 7

25

<212> PRT

<213> Homo sapiens

<400> 5

Asp Thr Ser Lys Leu Ala Ser

30

1

5

<210> 6

<211> 9

<212> PRT

35

<213> Homo sapiens

<400> 6

2002248184 15 Sep 2017

3 of 59

Phe Gln Gly Ser Gly Tyr Pro Phe Thr
1 5

<210> 7

<211> 120

<212> PRT

<213> Homo sapiens

<400> 7

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Ser Met Ile Thr Asn Trp Tyr Phe Asp Val Trp Gly Ala
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 8

<211> 106

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> VL Domain

<400> 8

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Thr	Leu	Ser	Ala	Ser	Val	Gly
1				5					10					15	

Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Cys	Gln	Leu	Ser	Val	Gly	Tyr	Met
			20				25						30		

His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr
		35					40					45			

Asp	Thr	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
	50					55					60				

Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Asp
65					70					75					80

Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Phe	Gln	Gly	Ser	Gly	Tyr	Pro	Phe	Thr
				85					90					95	

Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys
		100					105		

<210> 9

<211> 120

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> VH Domain

<400> 9

15 Sep 2017
2002248184

5 of 59

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
5 20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Ser Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Ala
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 10
<211> 7
<212> PRT
<213> Homo sapiens

<400> 10
Thr Ala Gly Met Ser Val Gly
1 5

<210> 11
<211> 106
<212> PRT
<213> Homo sapiens

2002248184 15 Sep 2017

<400> 11

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 12

<211> 10

<212> PRT

<213> Homo sapiens

<400> 12

Ser Met Ile Thr Asn Phe Tyr Phe Asp Val
1 5 10

<210> 13

<211> 106

<212> PRT

<213> Homo sapiens

15 Sep 2017
2002248184

7 of 59

<400> 13

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

5 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

10 Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

15 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Phe Ser Gly Tyr Pro Phe Thr
85 90 95

20 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 14

<211> 10

25 <212> PRT

<213> Homo sapiens

<400> 14

30 Ser Ala Ser Ser Ser Val Gly Tyr Met His
1 5 10

<210> 15

<211> 7

<212> PRT

35 <213> Homo sapiens

<400> 15

2002248184 15 Sep 2017

Asp Thr Phe Lys Leu Ala Ser
1 5

<210> 16

<211> 9

<212> PRT

<213> Homo sapiens

<400> 16

Phe Gln Phe Ser Gly Tyr Pro Phe Thr
1 5

<210> 17

<211> 120

<212> PRT

<213> Homo sapiens

<400> 17

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Ala

2002248184 15 Sep 2017

9 of 59

100

105

110

Gly Thr Thr Val Thr Val Ser Ser

115

120

5

<210> 18

<211> 7

<212> PRT

<213> Homo sapiens

10

<400> 18

Thr Pro Gly Met Ser Val Gly

1

5

15

<210> 19

<211> 16

<212> PRT

<213> Homo sapiens

20

<400> 19

Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser Leu Lys Asp

1

5

10

15

<210> 20

25

<211> 10

<212> PRT

<213> Homo sapiens

<400> 20

30

Asp Met Ile Phe Asn Phe Tyr Phe Asp Val

1

5

10

<210> 21

<211> 106

35

<212> PRT

<213> Homo sapiens

2002248184 15 Sep 2017

10 of 59

<220>

<221> misc_feature

<223> VL Domain

5 <400> 21

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

10 Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

15 Asp Thr Phe Tyr Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

20 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

25 Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 22

<211> 10

<212> PRT

30 <213> Homo sapiens

<400> 22

Ser Leu Ser Ser Arg Val Gly Tyr Met His
1 5 10

35 <210> 23

<211> 7

2002248184 15 Sep 2017

<212> PRT
<213> Homo sapiens

<400> 23

5 Asp Thr Phe Tyr Leu Ser Ser
1 5

<210> 24

<211> 120

10 <212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

15 <223> VH Domain

<400> 24

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

20 Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
20 25 30

25 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
50 55 60

30 Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

35 Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
100 105 110

2002248184 15 Sep 2017

Gly Thr Thr Val Thr Val Ser Ser
115 120

5 <210> 25
<211> 16
<212> PRT
<213> Homo sapiens

10 <400> 25
Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser Leu Lys Asp
1 5 10 15

15 <210> 26
<211> 106
<212> PRT
<213> Homo sapiens

20 <400> 26
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
20 25 30

25 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

30 Asp Thr Arg Gly Leu Pro Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

35 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

2002248184 15 Sep 2017

13 of 59

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 27
<211> 7
<212> PRT
<213> Homo sapiens

<400> 27
Asp Thr Arg Gly Leu Pro Ser
1 5

<210> 28
<211> 120
<212> PRT
<213> Homo sapiens

<400> 28
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln

2002248184 15 Sep 2017

14 of 59

100

105

110

Gly Thr Thr Val Thr Val Ser Ser

115

120

5

<210> 29

<211> 10

<212> PRT

<213> Homo sapiens

10

<400> 29

Asp Met Ile Phe Asn Trp Tyr Phe Asp Val

1

5

10

15

<210> 30

<211> 106

<212> PRT

<213> Homo sapiens

20

<400> 30

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly

1

5

10

15

Asp Arg Val Thr Ile Thr Cys Ser Pro Ser Ser Arg Val Gly Tyr Met

25

20

25

30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr

35

40

45

30

Asp Thr Met Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser

50

55

60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp

65

70

75

80

35

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr

85

90

95

2002248184 15 Sep 2017

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

5 <210> 31
<211> 10
<212> PRT
<213> Homo sapiens

10 <400> 31
Ser Pro Ser Ser Arg Val Gly Tyr Met His
1 5 10

15 <210> 32
<211> 7
<212> PRT
<213> Homo sapiens

20 <400> 32
Asp Thr Met Arg Leu Ala Ser
1 5

25 <210> 33
<211> 120
<212> PRT
<213> Homo sapiens

30 <400> 33
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Pro
20 25 30

35 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

15 Sep 2017
2002248184

16 of 59

Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys His Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 34

<211> 106

<212> PRT

<213> Homo sapiens

<400> 34

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Phe Lys Leu Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr

2002248184 15 Sep 2017

17 of 59

85

90

95

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

5

<210> 35

<211> 7

<212> PRT

<213> Homo sapiens

10

<400> 35

Asp Thr Phe Lys Leu Ser Ser
1 5

15

<210> 36

<211> 120

<212> PRT

<213> Homo sapiens

20

<400> 36

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
25 20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

30

Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys Asp Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

35

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

15 Sep 2017
2002248184

	Cys	Ala	Arg	Asp	Met	Ile	Phe	Asn	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Gln
				100				105						110		
5	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser								
				115				120								
	<210>		37													
	<211>		16													
10	<212>		PRT													
	<213>		Homo sapiens													
	<400>		37													
15	Asp	Ile	Trp	Trp	Asp	Gly	Lys	Lys	Asp	Tyr	Asn	Pro	Ser	Leu	Lys	Asp
	1				5					10					15	
	<210>		38													
	<211>		106													
	<212>		PRT													
20	<213>		Homo sapiens													
	<400>		38													
25	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Thr	Leu	Ser	Ala	Ser	Val	Gly
	1			5						10					15	
	Asp	Arg	Val	Thr	Ile	Thr	Cys	Ser	Ala	Ser	Ser	Arg	Val	Gly	Tyr	Met
				20					25					30		
30	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr
			35					40					45			
	Asp	Thr	Phe	Lys	Leu	Ser	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
			50				55					60				
35	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Asp
	65					70					75				80	

2002248184 15 Sep 2017

19 of 59

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

<210> 39

<211> 10

<212> PRT

<213> Homo sapiens

<400> 39

Ser Ala Ser Ser Arg Val Gly Tyr Met His
1 5 10

<210> 40

<211> 120

<212> PRT

<213> Homo sapiens

<400> 40

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Gly Lys Lys Ser Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr

2002248184 15 Sep 2017

20 of 59

85

90

95

Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 41

<211> 16

<212> PRT

<213> Homo sapiens

<400> 41

Asp Ile Trp Trp Asp Gly Lys Lys Ser Tyr Asn Pro Ser Leu Lys Asp
1 5 10 15

<210> 42

<211> 106

<212> PRT

<213> Homo sapiens

<400> 42

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Arg Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Met Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

2002248184 15 Sep 2017

	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Phe	Gln	Gly	Ser	Gly	Tyr	Pro	Phe	Thr	
					85					90					95		
5	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys							
				100					105								
	<210>		43														
	<211>		7														
10	<212>		PRT														
	<213>		Homo sapiens														
	<400>		43														
	Asp	Thr	Met	Tyr	Gln	Ser	Ser										
15	1				5												
	<210>		44														
	<211>		120														
	<212>		PRT														
20	<213>		Homo sapiens														
	<400>		44														
	Gln	Val	Thr	Leu	Arg	Glu	Ser	Gly	Pro	Ala	Leu	Val	Lys	Pro	Thr	Gln	
	1				5					10					15		
25																	
	Thr	Leu	Thr	Leu	Thr	Cys	Thr	Phe	Ser	Gly	Phe	Ser	Leu	Ser	Thr	Ala	
				20					25					30			
	Gly	Met	Ser	Val	Gly	Trp	Ile	Arg	Gln	Pro	Pro	Gly	Lys	Ala	Leu	Glu	
30			35					40					45				
	Trp	Leu	Ala	Asp	Ile	Trp	Trp	Asp	Gly	Lys	Lys	Ser	Tyr	Asn	Pro	Ser	
		50					55					60					
35	Leu	Lys	Asp	Arg	Leu	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Lys	Asn	Gln	Val	
	65					70					75				80		

2002248184 15 Sep 2017

22 of 59

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Asp Met Ile Phe Asn Phe Tyr Phe Asp Val Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 45

<211> 16

<212> PRT

<213> Homo sapiens

<400> 45

Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser Leu Lys Asp
1 5 10 15

<210> 46

<211> 106

<212> PRT

<213> Homo sapiens

<400> 46

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Leu Pro Ser Ser Arg Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Met Tyr Gln Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp

2002248184 15 Sep 2017

23 of 59

65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Phe Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 47

<211> 10

<212> PRT

<213> Homo sapiens

<400> 47

Leu Pro Ser Ser Arg Val Gly Tyr Met His
1 5 10

<210> 48

<211> 120

<212> PRT

<213> Homo sapiens

<400> 48

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

15 Sep 2017
2002248184

	Val	Leu	Lys	Val	Thr	Asn	Met	Asp	Pro	Ala	Asp	Thr	Ala	Thr	Tyr	Tyr
					85					90					95	
5	Cys	Ala	Arg	Asp	Met	Ile	Phe	Asn	Phe	Tyr	Phe	Asp	Val	Trp	Gly	Gln
				100					105					110		
	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser								
				115				120								
10	<210>	49														
	<211>	106														
	<212>	PRT														
	<213>	Homo sapiens														
15	<400>	49														
	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Thr	Leu	Ser	Ala	Ser	Val	Gly
	1				5					10					15	
20	Asp	Arg	Val	Thr	Ile	Thr	Cys	Ser	Ala	Ser	Ser	Arg	Val	Gly	Tyr	Met
				20					25					30		
	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr
				35				40					45			
25	Asp	Thr	Phe	Phe	Leu	Asp	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
		50					55					60				
	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Asp
30	65					70					75				80	
	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Phe	Gln	Gly	Ser	Gly	Tyr	Pro	Phe	Thr
					85					90					95	
35	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys						
				100						105						

2002248184 15 Sep 2017

25 of 59

<210> 50
<211> 7
<212> PRT
<213> Homo sapiens

<400> 50
Asp Thr Phe Phe Leu Asp Ser
1 5

<210> 51
<211> 120
<212> PRT
<213> Homo sapiens

<400> 51
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
20 20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Ser Tyr Asn Pro Ser
50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Gln
100 105 110

Gly Thr Thr Val Thr Val Ser Ser

2002248184 15 Sep 2017

26 of 59

115

120

<210> 52

<211> 106

<212> PRT

<213> Homo sapiens

<400> 52

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met

20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr

35 40 45

Asp Thr Arg Tyr Gln Ser Ser Gly Val Pro Ser Arg Phe Ser Gly Ser

50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp

65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr

85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys

100 105

<210> 53

<211> 7

<212> PRT

<213> Homo sapiens

<400> 53

Asp Thr Arg Tyr Gln Ser Ser

1 5

15 Sep 2017
2002248184

<210> 54
<211> 106
<212> PRT
5 <213> Homo sapiens

<220>
<221> misc_feature
<223> VL Domain
10
<400> 54
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15
15 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30
His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45
20 Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60
Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
25 65 70 75 80
Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95
30 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 55
<211> 120
35 <212> PRT
<213> Homo sapiens

2002248184 15 Sep 2017

28 of 59

<400> 55

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

5 Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

10 Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
50 55 60

15 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

20 Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Ala
100 105 110

Gly Thr Thr Val Thr Val Ser Ser
115 120

25

<210> 56

<211> 106

<212> PRT

<213> Homo sapiens

30

<220>

<221> misc_feature

<223> VL Domain

35

<400> 56

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

15 Sep 2017
2002248184

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

5 His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Phe Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

10

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

15

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

20

<210> 57
<211> 106
<212> PRT
<213> Homo sapiens

25

<220>
<221> misc_feature
<223> VL Domain

<400> 57

30

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

35

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

15 Sep 2017
2002248184

Asp Thr Tyr Lys Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

5 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

10 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

15 <210> 58
<211> 7
<212> PRT
<213> Homo sapiens

<400> 58
20 Asp Thr Tyr Lys Gln Thr Ser
1 5

<210> 59
<211> 106
25 <212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
30 <223> VL Domain

<400> 59
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

35 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

2002248184 15 Sep 2017

	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr
			35					40					45			
5	Asp	Thr	Arg	Tyr	Leu	Ser	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
		50					55					60				
	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Asp
10	65					70					75					80
	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Phe	Gln	Gly	Ser	Gly	Tyr	Pro	Phe	Thr
					85					90					95	
	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys						
15				100					105							
	<210>	60														
	<211>	106														
	<212>	PRT														
20	<213>	Homo sapiens														
	<400>	60														
	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Thr	Leu	Ser	Ala	Ser	Val	Gly
25	1				5					10					15	
	Asp	Arg	Val	Thr	Ile	Thr	Cys	Ser	Ala	Ser	Ser	Ser	Val	Gly	Tyr	Met
				20					25					30		
	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr
30			35					40					45			
	Asp	Thr	Phe	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser
		50					55					60				
35	Gly	Ser	Gly	Thr	Glu	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Asp
	65					70					75					80

2002248184 15 Sep 2017

32 of 59

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Phe Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
5 100 105

<210> 61

<211> 9

<212> PRT

<213> Homo sapiens

<400> 61

Phe Gln Gly Ser Phe Tyr Pro Phe Thr
1 5

<210> 62

<211> 106

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> VL Domain

<400> 62

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
30 20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Phe Lys Leu Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

2002248184 15 Sep 2017

33 of 59

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
5 85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 63

<211> 7

<212> PRT

<213> Homo sapiens

<400> 63

Asp Thr Phe Lys Leu Thr Ser
1 5

<210> 64

<211> 106

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> VL Domain

<400> 64

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
30 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 35 40 45

15 Sep 2017
2002248184

34 of 59

Asp Thr Phe Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 65

<211> 106

<212> PRT

<213> Homo sapiens

<400> 65

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Phe Arg Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys

2002248184 15 Sep 2017

35 of 59

100

105

<210> 66

<211> 7

<212> PRT

<213> Homo sapiens

<400> 66

Asp Thr Phe Arg Leu Ala Ser

1 5

<210> 67

<211> 120

<212> PRT

<213> Homo sapiens

<400> 67

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln

1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala

20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu

35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys His Tyr Asn Pro Ser

50 55 60

Leu Lys Asp Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val

65 70 75 80

Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr

85 90 95

Cys Ala Arg Asp Met Ile Phe Asn Trp Tyr Phe Asp Val Trp Gly Ala

100 105 110

2002248184 15 Sep 2017

Gly Thr Thr Val Thr Val Ser Ser
115 120

5 <210> 68
<211> 106
<212> PRT
<213> Homo sapiens

10 <400> 68
Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

15 Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

20 Asp Thr Tyr Arg His Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

25 Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

30 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 69
<211> 7
<212> PRT
35 <213> Homo sapiens

<400> 69

2002248184 15 Sep 2017

Asp Thr Tyr Arg His Ser Ser
1 5

<210> 70

<211> 106

<212> PRT

<213> Homo sapiens

<400> 70

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Tyr Lys Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 71

<211> 106

<212> PRT

<213> Homo sapiens

<400> 71

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly

2002248184 15 Sep 2017

38 of 59

1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Leu Ser Ser Ser Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
35 40 45

Asp Thr Phe Phe His Arg Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 72
<211> 10
<212> PRT
<213> Homo sapiens

<400> 72
Ser Leu Ser Ser Ser Val Gly Tyr Met His
1 5 10

<210> 73
<211> 7
<212> PRT
<213> Homo sapiens

<400> 73
Asp Thr Phe Phe His Arg Ser
1 5

15 Sep 2017
2002248184

<210> 74

<211> 106

<212> PRT

5 <213> Homo sapiens

<400> 74

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
15 35 40 45

Asp Thr Leu Leu Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

20 Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
85 90 95

25 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 75

30 <211> 7

<212> PRT

<213> Homo sapiens

<400> 75

35 Asp Thr Leu Leu Leu Asp Ser
1 5

<210> 76
 <211> 106
 <212> PRT
 <213> Homo sapiens

<400> 76
 Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly
 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Arg Val Gly Tyr Met
 20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr
 35 40 45

Asp Thr Ser Phe Leu Asp Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
 50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp
 65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr
 85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

<210> 77
 <211> 7
 <212> PRT
 <213> Homo sapiens

<400> 77
 Asp Thr Ser Phe Leu Asp Ser
 1 5

<210> 78

2002248184 15 Sep 2017

<211> 120
 <212> PRT
 <213> Homo sapiens

5 <400> 78
 Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
 1 5 10 15

10 Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ala
 20 25 30

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
 35 40 45

15 Trp Leu Ala Asp Ile Trp Trp Asp Asp Lys Lys Asp Tyr Asn Pro Ser
 50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
 65 70 75 80

20 Val Leu Lys Val Thr Asn Met Asp Pro Ala Asp Thr Ala Thr Tyr Tyr
 85 90 95

25 Cys Ala Arg Asp Met Ile Thr Asn Phe Tyr Phe Asp Val Trp Gly Ala
 100 105 110

Gly Thr Thr Val Thr Val Ser Ser
 115 120

30 <210> 79
 <211> 10
 <212> PRT
 <213> Homo sapiens

35 <400> 79
 Asp Met Ile Thr Asn Phe Tyr Phe Asp Val
 1 5 10

15 Sep 2017
2002248184

<210> 80

<211> 109

<212> PRT

5 <213> Homo sapiens

<400> 80

Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
1 5 10 15

Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
20 25 30

Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
15 35 40 45

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
50 55 60

20 Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
65 70 75 80

Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
85 90 95

25 Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala
100 105

<210> 81

30 <211> 108

<212> PRT

<213> Homo sapiens

<400> 81

35 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg
1 5 10 15

15 Sep 2017
2002248184

43 of 59

Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
20 25 30

Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
5 35 40 45

Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
50 55 60

Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln
10 65 70 75 80

Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
85 90 95

Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
15 100 105

<210> 82
20 <211> 15
<212> PRT
<213> Homo sapiens

<400> 82
25 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
1 5 10 15

<210> 83
<211> 232
30 <212> PRT
<213> Homo sapiens

<400> 83
35 Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
1 5 10 15

Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro

2002248184 15 Sep 2017

44 of 59

				20					25					30			
	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	
			35					40					45				
5	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	
		50					55					60					
	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	
10	65					70					75					80	
	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	
					85					90					95		
	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	
15				100					105					110			
	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	
			115					120					125				
20	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	
		130					135					140					
	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	
25	145					150					155					160	
	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	
					165					170					175		
	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	
30				180					185					190			
	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	
			195					200					205				
35	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	
		210						215					220				

2002248184 15 Sep 2017

Ser Leu Ser Leu Ser Pro Gly Lys
225 230

5 <210> 84
<211> 365
<212> PRT
<213> Homo sapiens

10 <400> 84
Met Gly Val Pro Arg Pro Gln Pro Trp Ala Leu Gly Leu Leu Leu Phe
1 5 10 15

15 Leu Leu Pro Gly Ser Leu Gly Ala Glu Ser His Leu Ser Leu Leu Tyr
20 25 30

His Leu Thr Ala Val Ser Ser Pro Ala Pro Gly Thr Pro Ala Phe Trp
35 40 45

20 Val Ser Gly Trp Leu Gly Pro Gln Gln Tyr Leu Ser Tyr Asn Ser Leu
50 55 60

Arg Gly Glu Ala Glu Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val
65 70 75 80

25 Ser Trp Tyr Trp Glu Lys Glu Thr Thr Asp Leu Arg Ile Lys Glu Lys
85 90 95

30 Leu Phe Leu Glu Ala Phe Lys Ala Leu Gly Gly Lys Gly Pro Tyr Thr
100 105 110

Leu Gln Gly Leu Leu Gly Cys Glu Leu Gly Pro Asp Asn Thr Ser Val
115 120 125

35 Pro Thr Ala Lys Phe Ala Leu Asn Gly Glu Glu Phe Met Asn Phe Asp
130 135 140

15 Sep 2017
2002248184

46 of 59

Leu Lys Gln Gly Thr Trp Gly Gly Asp Trp Pro Glu Ala Leu Ala Ile
145 150 155 160

Ser Gln Arg Trp Gln Gln Gln Asp Lys Ala Ala Asn Lys Glu Leu Thr
165 170 175

Phe Leu Leu Phe Ser Cys Pro His Arg Leu Arg Glu His Leu Glu Arg
180 185 190

Gly Arg Gly Asn Leu Glu Trp Lys Glu Pro Pro Ser Met Arg Leu Lys
195 200 205

Ala Arg Pro Ser Ser Pro Gly Phe Ser Val Leu Thr Cys Ser Ala Phe
210 215 220

Ser Phe Tyr Pro Pro Glu Leu Gln Leu Arg Phe Leu Arg Asn Gly Leu
225 230 235 240

Ala Ala Gly Thr Gly Gln Gly Asp Phe Gly Pro Asn Ser Asp Gly Ser
245 250 255

Phe His Ala Ser Ser Ser Leu Thr Val Lys Ser Gly Asp Glu His His
260 265 270

Tyr Cys Cys Ile Val Gln His Ala Gly Leu Ala Gln Pro Leu Arg Val
275 280 285

Glu Leu Glu Ser Pro Ala Lys Ser Ser Val Leu Val Val Gly Ile Val
290 295 300

Ile Gly Val Leu Leu Leu Thr Ala Ala Ala Val Gly Gly Ala Leu Leu
305 310 315 320

Trp Arg Arg Met Arg Ser Gly Leu Pro Ala Pro Trp Ile Ser Leu Arg
325 330 335

Gly Asp Asp Thr Gly Val Leu Leu Pro Thr Pro Gly Glu Ala Gln Asp

2002248184 15 Sep 2017

47 of 59

340

345

350

Ala Asp Leu Lys Asp Val Asn Val Ile Pro Ala Thr Ala

355

360

365

<210> 85

<211> 365

<212> PRT

<213> Mus sp.

<400> 85

Met Gly Met Pro Leu Pro Trp Ala Leu Ser Leu Leu Leu Val Leu Leu

1

5

10

15

Pro Gln Thr Trp Gly Ser Glu Thr Arg Pro Pro Leu Met Tyr His Leu

20

25

30

Thr Ala Val Ser Asn Pro Ser Thr Gly Leu Pro Ser Phe Trp Ala Thr

35

40

45

Gly Trp Leu Gly Pro Gln Gln Tyr Leu Thr Tyr Asn Ser Leu Arg Gln

50

55

60

Glu Ala Asp Pro Cys Gly Ala Trp Val Trp Glu Asn Gln Val Ser Trp

65

70

75

80

Tyr Trp Glu Lys Glu Thr Thr Asp Leu Lys Ser Lys Glu Gln Leu Phe

85

90

95

Leu Glu Ala Leu Lys Thr Leu Glu Lys Ile Leu Asn Gly Thr Tyr Thr

100

105

110

Leu Gln Gly Leu Leu Gly Cys Glu Leu Ala Ser Asp Asn Ser Ser Val

115

120

125

Pro Thr Ala Val Phe Ala Leu Asn Gly Glu Glu Phe Met Lys Phe Asn

130

135

140

15 Sep 2017
2002248184

	Pro	Arg	Ile	Gly	Asn	Trp	Thr	Gly	Glu	Trp	Pro	Glu	Thr	Glu	Ile	Val	
	145					150					155					160	
5	Ala	Asn	Leu	Trp	Met	Lys	Gln	Pro	Asp	Ala	Ala	Arg	Lys	Glu	Ser	Glu	
					165					170					175		
	Phe	Leu	Leu	Asn	Ser	Cys	Pro	Glu	Arg	Leu	Leu	Gly	His	Leu	Glu	Arg	
				180					185					190			
10	Gly	Arg	Arg	Asn	Leu	Glu	Trp	Lys	Glu	Pro	Pro	Ser	Met	Arg	Leu	Lys	
			195					200					205				
	Ala	Arg	Pro	Gly	Asn	Ser	Gly	Ser	Ser	Val	Leu	Thr	Cys	Ala	Ala	Phe	
15		210					215					220					
	Ser	Phe	Tyr	Pro	Pro	Glu	Leu	Lys	Phe	Arg	Phe	Leu	Arg	Asn	Gly	Leu	
	225					230					235					240	
20	Ala	Ser	Gly	Ser	Gly	Asn	Cys	Ser	Thr	Gly	Pro	Asn	Gly	Asp	Gly	Ser	
					245					250					255		
	Phe	His	Ala	Trp	Ser	Leu	Leu	Glu	Val	Lys	Arg	Gly	Asp	Glu	His	His	
				260					265					270			
25	Tyr	Gln	Cys	Gln	Val	Glu	His	Glu	Gly	Leu	Ala	Gln	Pro	Leu	Thr	Val	
			275					280					285				
	Asp	Leu	Asp	Ser	Ser	Ala	Arg	Ser	Ser	Val	Pro	Val	Val	Gly	Ile	Val	
30		290					295					300					
	Leu	Gly	Leu	Leu	Leu	Val	Val	Val	Ala	Ile	Ala	Gly	Gly	Val	Leu	Leu	
	305					310					315					320	
35	Trp	Gly	Arg	Met	Arg	Ser	Gly	Leu	Pro	Ala	Pro	Trp	Leu	Ser	Leu	Ser	
				325						330					335		

2002248184 15 Sep 2017

49 of 59

Gly Asp Asp Ser Gly Asp Leu Leu Pro Gly Gly Asn Leu Pro Pro Glu
340 345 350

Ala Glu Pro Gln Gly Ala Asn Ala Phe Pro Ala Thr Ser
5 355 360 365

<210> 86

<211> 7

<212> PRT

<213> Homo sapiens

<400> 86

Val Leu His Gln Asp Trp Leu
1 5

<210> 87

<211> 6

<212> PRT

<213> Homo sapiens

<400> 87

Leu Met Ile Ser Arg Thr
1 5

<210> 88

<211> 9

<212> PRT

<213> Homo sapiens

<400> 88

Met His Glu Ala Leu His Asn His Tyr
1 5

<210> 89

<211> 5

<212> PRT

<213> Homo sapiens

2002248184 15 Sep 2017

5

<400> 89
Gly Gln Pro Glu Asn
1 5

10

<210> 90
<211> 6
<212> PRT
<213> Homo sapiens

<400> 90
Leu Tyr Ile Thr Arg Glu
1 5

15

<210> 91
<211> 6
<212> PRT
<213> Homo sapiens

20

<400> 91
Leu Tyr Ile Ser Arg Thr
1 5

25

<210> 92
<211> 6
<212> PRT
<213> Homo sapiens

30

<400> 92
Leu Tyr Ile Ser Arg Ser
1 5

35

<210> 93
<211> 6
<212> PRT
<213> Homo sapiens

2002248184 15 Sep 2017

<400> 93
Leu Tyr Ile Ser Arg Arg
1 5

5 <210> 94
<211> 6
<212> PRT
<213> Homo sapiens

10 <400> 94
Leu Tyr Ile Ser Arg Gln
1 5

15 <210> 95
<211> 6
<212> PRT
<213> Homo sapiens

20 <400> 95
Leu Trp Ile Ser Arg Thr
1 5

25 <210> 96
<211> 6
<212> PRT
<213> Homo sapiens

30 <400> 96
Leu Tyr Ile Ser Leu Gln
1 5

35 <210> 97
<211> 6
<212> PRT
<213> Homo sapiens

<400> 97

2002248184 15 Sep 2017

	Leu Phe Ile Ser Arg Asp
	1 5
	<210> 98
5	<211> 6
	<212> PRT
	<213> Homo sapiens
	<400> 98
10	Leu Phe Ile Ser Arg Thr
	1 5
	<210> 99
	<211> 6
15	<212> PRT
	<213> Homo sapiens
	<400> 99
	Leu Phe Ile Ser Arg Arg
20	1 5
	<210> 100
	<211> 6
	<212> PRT
25	<213> Homo sapiens
	<400> 100
	Leu Phe Ile Thr Gly Ala
	1 5
30	
	<210> 101
	<211> 6
	<212> PRT
	<213> Homo sapiens
35	
	<400> 101
	Leu Ser Ile Ser Arg Glu

2002248184 15 Sep 2017

1 5
<210> 102
<211> 6
5 <212> PRT
<213> Homo sapiens

<400> 102
Arg Thr Ile Ser Ile Ser
10 1 5

<210> 103
<211> 7
<212> PRT
15 <213> Homo sapiens

<400> 103
Thr Pro His Ser Asp Trp Leu
1 5

20
<210> 104
<211> 7
<212> PRT
<213> Homo sapiens

25
<400> 104
Ile Pro His Glu Asp Trp Leu
1 5

30 <210> 105
<211> 5
<212> PRT
<213> Homo sapiens

35 <400> 105
Arg Thr Arg Glu Pro
1 5

15 Sep 2017
2002248184

5	<210>	106
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
10	<400>	106
	Asp Pro Pro Glu Ser	
	1	5
15	<210>	107
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
	<400>	107
	Ser Asp Pro Glu Pro	
	1	5
20	<210>	108
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
25	<400>	108
	Thr Ser His Glu Asn	
	1	5
30	<210>	109
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
35	<400>	109
	Ser Lys Ser Glu Asn	
	1	5

2002248184 15 Sep 2017

	<210>	110
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
5		
	<400>	110
	His Arg Ser Glu Asn	
	1	5
10		
	<210>	111
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
15		
	<400>	111
	Lys Ile Arg Glu Asn	
	1	5
20		
	<210>	112
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
25		
	<400>	112
	Gly Ile Thr Glu Ser	
	1	5
30		
	<210>	113
	<211>	5
	<212>	PRT
	<213>	Homo sapiens
35		
	<400>	113
	Ser Met Ala Glu Pro	
	1	5
	<210>	114

2002248184 15 Sep 2017

<211> 9
<212> PRT
<213> Homo sapiens

5 <400> 114
Met His Glu Ala Leu Arg Tyr His His
1 5

10 <210> 115
<211> 9
<212> PRT
<213> Homo sapiens

15 <400> 115
Met His Glu Ala Leu His Phe His His
1 5

20 <210> 116
<211> 9
<212> PRT
<213> Homo sapiens

25 <400> 116
Met His Glu Ala Leu Lys Phe His His
1 5

30 <210> 117
<211> 9
<212> PRT
<213> Homo sapiens

35 <400> 117
Met His Glu Ala Leu Ser Tyr His Arg
1 5

<210> 118
<211> 9

2002248184 15 Sep 2017

<212> PRT

<213> Homo sapiens

<400> 118

5 Thr His Glu Ala Leu His Tyr His Thr
1 5

<210> 119

<211> 9

10 <212> PRT

<213> Homo sapiens

<400> 119

15 Met His Glu Ala Leu His Tyr His Tyr
1 5

<210> 120

<211> 54

20 <212> DNA

<213> Artificial Sequence

<220>

25 <223> oligonucleotide sequence used to construct library 1

<220>

<221> misc_feature

<222> 16, 17, 19, 20, 22, 23, 28, 29, 31, 32

30 <223> n = A,T,C or G

<400> 120

catgtgacct caggsnnnsnn snngatsnns nnggtgtcct tgggttttgg gggg 54

<210> 121

35 <211> 53

<212> DNA

<213> Artificial Sequence

2002248184 15 Sep 2017

<220>

<223> oligonucleotide sequence used to construct library 2

5 <220>

<221> misc_feature

<222> 23, 24, 29, 30, 32, 33, 38, 39, 41, 42

<223> n = A,T,C or G

10 <400> 121

gcacttgtag tccttgccat tsnnccasnn snngtgsnns nnggtgagga cgc 53

<210> 122

<211> 38

15 <212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide sequence used to construct library 3

20

<220>

<221> misc_feature

<222> 14, 15, 20, 21, 23, 24, 26, 27

<223> n = A,T,C or G

25

<400> 122

ggtcttgtag ttsnnctcsn nsnnnsnnatt gctctccc 38

<210> 123

30 <211> 53

<212> DNA

<213> Artificial Sequence

<220>

35 <223> oligonucleotide sequence used to construct library 4

<220>

15 Sep 2017

2002248184

<221> misc_feature

<222> 15, 16, 21, 22, 24, 25, 39, 40

<223> n = A,T,C or G

5 <400> 123

ggctcttctg cgtsnngtgs nnsnncagag cctcatgsnn cacggagcat gag 53

<210> 124

<211> 106

.0 <212> PRT

<213> Homo sapiens

<400> 124

Asp Ile Gln Met Thr Gln Ser Pro Ser Thr Leu Ser Ala Ser Val Gly

.5 1 5 10 15

Asp Arg Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Gly Tyr Met

20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile Tyr

35 40 45

!0 Asp Thr Tyr Lys Gln Thr Ser Gly Val Pro Ser Arg Phe Ser Gly Ser

50 55 60

Gly Ser Gly Thr Glu Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro Asp

65 70 75 80

Asp Phe Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Phe Thr

25 85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys

100 105

<210> 125

30 <211> 7

<212> PRT

<213> Homo sapiens

<400> 125

35 Asp Thr Arg Tyr Leu Ser Ser

1 5