



(86) Date de dépôt PCT/PCT Filing Date: 2012/02/23
(87) Date publication PCT/PCT Publication Date: 2012/08/30
(45) Date de délivrance/Issue Date: 2016/01/26
(85) Entrée phase nationale/National Entry: 2013/08/23
(86) N° demande PCT/PCT Application No.: IL 2012/050060
(87) N° publication PCT/PCT Publication No.: 2012/114342
(30) Priorité/Priority: 2011/02/23 (US61/445,559)

(51) Cl.Int./Int.Cl. *A61K 31/5517* (2006.01),
A61K 31/465 (2006.01), *A61P 25/00* (2006.01),
C07D 487/04 (2006.01)
(72) Inventeur/Inventor:
AMSELEM, SHIMON, IL
(73) Propriétaire/Owner:
COERULEUS LTD., IL
(74) Agent: SHAPIRO COHEN LLP

(54) Titre : COMPLEXES DE FLUMAZENIL, COMPOSITIONS LES COMPRENANT ET LEURS UTILISATIONS
(54) Title: FLUMAZENIL COMPLEXES, COMPOSITIONS COMPRISING SAME AND USES THEREOF

(57) **Abrégé/Abstract:**

The present invention provides soluble complexes of flumazenil, methods for the preparation thereof, pharmaceutical compositions comprising same and use of said compositions for alleviating or counteracting the various types of hypersomnia, drowsiness, residual effects associated with the administration of sleep/hypnotic drugs, alcohol intoxication or hepatic encephalopathy. The present invention is based in part on the unexpected discovery that flumazenil complexes are more soluble than flumazenil alone.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number

WO 2012/114342 A1

(43) International Publication Date
30 August 2012 (30.08.2012)

(51) International Patent Classification:
C07D 245/06 (2006.01)

(21) International Application Number:
PCT/IL2012/050060

(22) International Filing Date:
23 February 2012 (23.02.2012)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/445,559 23 February 2011 (23.02.2011) US

(71) Applicant (for all designated States except US): **COER-
ULEUS LTD.** [IL/IL]; Industrial Zone, P.O. Box 12,
12900 Katzrin (IL).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **AMSELEM, Shimon**
[IL/IL]; 10 Miriam Mizrahi Street, 76551 Rehovot (IL).

(74) Agents: **WEBB & CO.** et al.; P.O. Box 2189, 76121 Re-
hovot (IL).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: FLUMAZENIL COMPLEXES, COMPOSITIONS COMPRISING SAME AND USES THEREOF

(57) Abstract: The present invention provides soluble complexes of flumazenil, methods for the preparation thereof, pharmaceutical compositions comprising same and use of said compositions for alleviating or counteracting the various types of hypersomnia, drowsiness, residual effects associated with the administration of sleep/hypnotic drugs, alcohol intoxication or hepatic encephalopathy. The present invention is based in part on the unexpected discovery that flumazenil complexes are more soluble than flumazenil alone.

**WO 2012/114342 A1**

FLUMAZENIL COMPLEXES, COMPOSITIONS COMPRISING SAME AND USES THEREOF

FIELD OF THE INVENTION

5 The present invention provides soluble complexes of flumazenil, methods for the preparation thereof, pharmaceutical compositions comprising same and use of said compositions for alleviating or counteracting the various types of hypersomnia, drowsiness, residual effects associated with the administration of sleep/hypnotic drugs, sedation, alcohol intoxication and hepatic encephalopathy.

BACKGROUND OF THE INVENTION

10 Flumazenil (chemically named ethyl 8-fluoro- 5,6-dihydro- 5-methyl- 6-oxo- 4H-imidazo [1,5-a] [1,4] benzodiazepine- 3-carboxylate), also known as, Ro 15-1788, Anexate™, Lanexat™, Mazicon™ and Romazicon™, was initially disclosed in US
15 4,316,839. Flumazenil is an imidazobenzodiazepine having high affinity for the GABA_A/benzodiazepine-receptor complex, the specific binding site of benzodiazepines. As flumazenil is a competitive inhibitor of benzodiazepines, it is used for reversing benzodiazepine-induced sedation and anesthesia following therapeutic or diagnostic
20 procedures (e.g. WO 2009/114740). Flumazenil is also known to reverse the effect of non-benzodiazepine drugs, such as the imidazopyridine hypnotic zolpidem (e.g. Patat et al., Clin Pharmacol Ther., 1994, 56(4):430-6). Flumazenil in combination with hydroxyzine and gabapentin are used for treating stimulant addiction (methamphetamine) (Bond AJ, 1998, CNS Drugs, 9(1): 41-57). Flumazenil is also effective in treating hepatic encephalopathy (Als-Nielsen B et al. Cochrane Database of Systematic Reviews 2004, Issue 2. Art. No.:
25 CD002798. DOI: 10.1002/14651858.CD002798.pub2). The aforementioned therapeutic effects were achieved by intravenous administration of flumazenil in a liquid formulation.

A liquid formulation of flumazenil (Romazicon®) is currently approved for reversing the sedative effects of benzodiazepines. Due to the limited solubility of flumazenil, the Romazicon® liquid formulation contains only 0.01% flumazenil. Poor water solubility
30 frequently correlates with low drug absorption and bioavailability, and limits the amount of drug that can be administered in a pharmaceutical composition.

The use of hydrotopes, such as nicotinamide, for complexation with insoluble

compounds in order to improve their solubility is known, for example, US 6,087,353.

There remains an unmet need to formulate flumazenil derivatives that are highly soluble and thereby suitable for the preparation of highly concentrated flumazenil formulations.

SUMMARY OF THE INVENTION

The present invention is directed to pharmaceutically acceptable complexes of flumazenil, comprising flumazenil and a complexation agent. The flumazenil complexes of the invention are highly soluble while maintaining the therapeutic activity known for flumazenil, namely, ameliorating or inhibiting any type of hypersomnia and drowsiness, including endogenous hypersomnia associated with GABA hyper-activation or exogenic hypersomnia e.g, residual hypersomnia associated with the administration of sleep/hypnotic drugs and ameliorating or inhibiting alcohol intoxication, or treating hepatic encephalopathy or any other disease or disorder associated with GABA related sedation or hypnotics. Having a significantly higher solubility of flumazenil, the flumazenil complexes of the invention offer a pronounced advantage over flumazenil compositions currently known in the art. Moreover, the flumazenil complexes of invention are advantageous for the preparation of formulations suitable for self-administration, such as, sublingual administration, thereby avoiding the involvement of professional care takers and the need of admitting to a health center.

According to one aspect of the present invention, there is provided a pharmaceutically acceptable complex of ethyl 8-fluoro-5,6-dihydro-5-methyl-6-oxo- 4H- imidazo [1,5-a] [1,4] benzodiazepine- 3-carboxylate or a salt thereof and a complexation agent or a derivative thereof, wherein the complexation agent is selected from the group consisting of: nicotinamide, sodium nicotinate, sodium benzoate, sodium hydroxybenzoates, sodium salicylate, sodium gentisate, gentisic acid ethanolamide, sodium toluates, sodium aminobenzoate, sodium anthranilate, sodium butylmonoglycolsulfate and resorcinol.

2a

The present invention is based in part on the unexpected discovery that flumazenil complexes are more soluble than flumazenil alone. Accordingly, these compounds are
5 suitable for the preparation of highly concentrated pharmaceutical compositions of flumazenil. This enables administration of higher doses of flumazenil (per volume) than currently available. Typically, the concentration of flumazenil in the form of the flumazenil complexes of the invention is within the range of about 0.4 to 2%. This concentration is much
10 higher than the concentration of flumazenil in commercially available formulations. In fact, the concentration of flumazenil in the pharmaceutical compositions of the invention is higher by more than two orders of magnitude from the concentration of flumazenil in the commercially available formulations. Highly concentrated flumazenil compositions are particularly advantageous in view of the maximal recommended daily dose approved for flumazenil solution (for example, Romazicon®), which is 3 mg. As the concentration of
15 flumazenil in the commercial solution is only 0.01%, a dose of 3 mg requires administration

of a large volume of the flumazenil solution, as high as 30 ml. The present invention overcomes this deficiency by providing concentrated formulations that can be administered in lower volumes thus improving patient compliance. Administering highly concentrated formulations of flumazenil also enables applying more routes of delivery, specifically, 5 sublingual, submucosal, intramuscular, transdermal or any parenteral route of administration.

Sublingual administration of the formulations of the invention is of special advantage. Apart from being easy to administer, the sublingual route bypasses the first liver metabolism, absorbance of flumazenil is immediate, thereby fast response to therapy can be achieved.

Although being highly concentrated, the formulations of the invention are stable and no 10 precipitation occurs during storage. The stability of the formulations of the invention renders them suitable for long periods of storage.

According to a first aspect, the present invention provides a complex of ethyl 8-fluoro- 5,6-dihydro- 5-methyl- 6-oxo- 4H- imidazo [1,5-a] [1,4] benzodiazepine- 3-carboxylate (flumazenil) or a salt thereof and a complexation agent or derivatives thereof, with the 15 proviso that the complexation agent is other than cyclodextrin or meglumine.

According to another aspect, the present invention provides a complex of ethyl 8-fluoro- 5,6-dihydro- 5-methyl- 6-oxo- 4H- imidazo [1,5-a] [1,4] benzodiazepine- 3-carboxylate (flumazenil) or a salt thereof and an aromatic stacking complexation agent.

According to another embodiment, the stacking complexation agent is selected from the 20 group consisting of: benzoic acid, benzoate, hydroxybenzoic acid, hydroxybenzoate, salicylic acid, salicylate, nicotinamide, nicotinic acid, nicotinate, gentisic acid, gentisate, ethanolamide, toluic acid (ortho, meta or para), toluate (ortho, meta or para), aminobenzoic acid, aminobenzoate, anthranilic acid, anthranilate, butylmonoglycolsulfate and resorcinol.

According to yet another embodiment, the stacking complexation agent is selected from 25 the group consisting of: benzoic acid, sodium benzoate, potassium benzoate, hydroxybenzoic acid, sodium hydroxybenzoate, potassium hydroxybenzoate, salicylic acid, sodium salicylate, potassium salicylate, nicotinamide, nicotinic acid, sodium nicotinate, potassium nicotinate, gentisic acid, sodium gentisate, potassium gentisate, ethanolamide, sodium toluate, toluic acid, potassium toluate, aminobenzoic acid, sodium aminobenzoate, potassium 30 aminobenzoate, anthranilic acid, sodium anthranilate, potassium anthranilate, sodium butylmonoglycolsulfate, potassium butylmonoglycolsulfate and resorcinol.

According to yet another embodiment, the stacking complexation agent is selected from the group consisting of: sodium benzoate, sodium hydroxybenzoate, sodium salicylate,

nicotinamide, sodium nicotinate, sodium gentisate, gentisic acid ethanolamide, sodium toluate, sodium aminobenzoates, sodium anthranilate, sodium butylmonoglycolsulfate and resorcinol.

It is noted that the complexation agent of the flumazenil complex of the current invention may be any complexation agent known in the art other than cyclodextrin. However, the present invention contemplates compositions comprising the flumazenil complexes of the invention, and cyclodextrin as an additional excipient.

According to one embodiment, the complexation agent is nicotinamide or a derivative thereof. According to yet another embodiment, the nicotinamide derivative is selected from the group consisting of: 2-amino-nicotinamide derivatives, 5-phenyl-nicotinamide derivative and 6-substituted nicotinamide derivative.

According to yet another embodiment, the complexation agent:flumazenil ratio is in the range of 1:1 to 2:1.

According to another aspect, the present invention provides a flumazenil nicotinamide stacking complex.

According to yet another aspect, the present invention provides a pharmaceutical composition comprising the flumazenil complex and a pharmaceutical acceptable carrier.

According to yet another embodiment, the pharmaceutical composition is in a form selected from the group consisting of: a pill, tablet, lozenge, coated tablet, granule, capsule, hard or soft gelatin capsule, aqueous solution, alcoholic solution, oily solution, syrup, emulsion suspension pastille suppository, solution for injection or infusion, ointment, tincture, cream, lotion, powder, spray, transdermal therapeutic systems, nasal spray, buccal spray, aerosol mixture, microcapsule, implant, rod and plaster. Each possibility represents a separate embodiment of the present invention.

According to yet another embodiment, the pharmaceutical composition is in a form selected from the group consisting of: immediate release, delayed release, pulsatile release, continuous release and repetitive release. Each possibility represents a separate embodiment of the present invention.

According to yet another embodiment, the composition is in a solid dosage form. According to yet another embodiment, the composition is suitable for oral administration.

According to yet another embodiment, flumazenil concentration in the pharmaceutical composition is within the range of about 0.4 to 2 % w/w. According to yet another embodiment, the concentration of flumazenil is within the range of about 0.5 to 1.8 % w/w.

According to certain embodiments, the concentration of flumazenil is within the range of about 0.6 to 1.5 % w/w.

According to yet another embodiment, the pharmaceutical composition comprises flumazenil complex as an active ingredient, and a solubilizing agent selected from an alcohol, a glycol and a combination thereof. Each possibility represents a separate embodiment of the present invention.

According to yet another embodiment, the solubilizing agent comprises a combination of an alcohol and a glycol wherein the alcohol:glycol ratio is at least 1.5:1. According to yet another embodiment, the alcohol:glycol ratio is in the range of 1.5:1 to 5:1.

According to yet another embodiment, the solubilizing agent comprises a combination of an alcohol and a glycol wherein the concentration of the solubilizing agent is at least 40%. According to yet another embodiment, the solubilizing agent comprises a combination of an alcohol and a glycol wherein the concentration of the solubilizing agent is in the range of 40% to 60%. According to yet another embodiment, the solubilizing agent comprises ethanol and propylene glycol.

According to yet another embodiment, the pharmaceutical composition further comprises a buffering agent.

According to yet another embodiment, the pharmaceutical composition further comprises at least one agent selected from the group consisting of: a penetration enhancer, a surfactant and cyclodextrin, as an inclusion complexing agent. Each possibility represents a separate embodiment of the present invention.

According to yet another embodiment, the cyclodextrin is hydroxypropyl β -cyclodextrin (HPCD). The cyclodextrin is preferably formulated in a buffer having a pH from about 3 to about 6. In one particular embodiment, the cyclodextrin (e.g., HPCD) is formulated in a citric acid buffer having a pH of about 4.

According to yet another embodiment, the preservative is selected from the group consisting of benzyl alcohol, propylparaben, methylparaben and combinations thereof. In one embodiment, the preservative is benzyl alcohol. In another embodiment, the preservative is a combination of propylparaben and methylparaben. Each possibility represents a separate embodiment of the present invention.

According to yet another embodiment, the penetration enhancer is menthol.

According to yet another embodiment, the buffering agent is selected from the group consisting of: citric buffer, sodium chloride and combination thereof. Each possibility

represents a separate embodiment of the present invention.

According to yet another embodiment, the surfactant is a cationic surfactant. According to yet another embodiment, the surfactant is benzalkonium chloride.

According to yet another aspect, the present invention provides a method for treating a disease or disorder associated with binding of an agent to the GABA receptor, comprising administering to a patient in need thereof a pharmaceutical composition comprising as an active ingredient a flumazenil complex, and at least one pharmaceutical acceptable carrier.

According to one embodiment, treating a disease or disorder associated with binding of an agent to the GABA receptor comprises reversing the effect caused by the binding of the agent to the GABA receptor.

According to another embodiment, the agent is a benzodiazepine a non-benzodiazepine the binding of which to the GABA receptor causes sedation.

According to yet another aspect, the present invention provides a method for treating a disease or disorder comprising administering to a patient in need thereof a pharmaceutical composition comprising as an active ingredient a flumazenil complex, and at least one pharmaceutical acceptable carrier, wherein the disease or disorder is selected from the group consisting of: excessive sleepiness, alcohol intoxication and hepatic encephalopathy.

According to yet another aspect, the present invention provides a method for treating excessive sleepiness comprising administering to a patient in need thereof a pharmaceutical composition comprising as an active ingredient a flumazenil complex, and at least one pharmaceutical acceptable carrier.

According to one embodiment, the excessive sleepiness is selected from the group consisting of: excessive sleepiness associated with hypersomnia, excessive sleepiness associated with drowsiness, symptoms associated with overdose of a sleep drug, excessive sleepiness associated with sedation and anesthesia induced by a sleep drug, excessive sleepiness associated with sleep drug addiction, excessive sleepiness associated with stimulant addiction, Alzheimer's disease, anxiety, schizophrenia, specifically, schizophrenia associated with excessive sleepiness, the rebound effect of a sleep drug, balance impairment induced by a sleep drug or any combination thereof.

According to another embodiment, the excessive sleepiness is caused by alcohol intoxication.

According to yet another embodiment, the stimulant is methamphetamine. According to yet another embodiment, treating excessive sleepiness associated with stimulant addiction

further comprises administering the flumazenil complex, or the pharmaceutical composition comprising same, in combination with hydroxyzine and gabapentin.

According to yet another embodiment, the sleep-drug is selected from the group consisting of a benzodiazepine, a benzodiazepine modulator, a benzodiazepine analog, a non-
5 benzodiazepine, a 5-HT_{2A} receptor antagonist, a melatonin receptor agonist, an orexin receptor antagonist, a selective serotonin reuptake inhibitor (SSRI), an antihistamine and an herbal product.

According to yet another embodiment, the sleep drug is a sedative/hypnotic drug.

According to yet another embodiment, the sleep drug is a benzodiazepine
10 sedative/hypnotic drug. According to yet another embodiment, the benzodiazepine sleep drug is selected from the group consisting of alprazolam, bromazepam, clonazepam, clotiazepam, cloxazolam, diazepam, estazolam, etizolam, fludiazepam, flunitrazepam, flurazepam, halazepam, haloxazolam, lorazepam, medazepam, midazolam, nimetazepam, nitrazepam, olanzapine, oxazepam, quazepam, temazepam and triazolam.

15 According to yet another embodiment, treating excessive sleepiness comprises reversing benzodiazepine-induced sedation and anesthesia following therapeutic or diagnostic procedures.

According to yet another embodiment, the sleep-drug is a non-benzodiazepine sedative/hypnotic drug. According to yet another embodiment, the non-benzodiazepine sleep
20 drug is selected from the group consisting of adiplon (NG-2-73), agomelatine, almoxerant (ACT-078573), brotizolam, diphenhydramine, divaplon, doxepin, eplivanserin (SR 46349), doxylamine succinate, eszopiclone, indiplon, ocinaplon, pagoclone, pazinaclone, pruvanserin (EMD 281014), suproclone, suriclone, L-tryptophan, 5-hydroxy-L-tryptophan, melatonin, melatonin receptor agonists, such as VEC-162 and PD-6735, muramyl dipeptide, ramelteon,
25 sleep-promoting substance, uridine, volinanserin (M-100907), zaleplon, zolpidem, imidazopyridine hypnotic zolpidem, APD125, ACP-103, PD 200-390, HY10275, GW649863 and EVT-201.

According to yet another aspect, the present invention provides a method for treating alcohol intoxication comprising administering to a patient in need thereof a pharmaceutical
30 composition comprising the flumazenil complex of the invention, and at least one pharmaceutical acceptable carrier.

According to one another embodiment, the excessive sleepiness caused by alcohol intoxication is treated by any one or more of reversing the effects of alcohol intoxication,

reducing the effects of alcohol intoxication, alleviating the effects of alcohol intoxication and/or improving performance after alcohol consumption.

According to yet another aspect, the present invention provides a method for treating hepatic encephalopathy, comprising administering to a subject in need thereof the flumazenil
5 complex of the invention or a pharmaceutical composition comprising same.

According to another embodiment, the route of administration is selected from the group consisting of: oral, buccal, sublingual, transdermal, transmucosal, intranasal, intravenous (*i.v.*), intraperitoneal (*i.p.*), intramuscular (*i.m.*), subcutaneous (*s.c.*) or intra-
thecal (*i.t.*). Each possibility represents a separate embodiment of the present invention.

10 According to yet another embodiment, the methods of the invention further comprise co-administering, concomitantly or in sequence, a pharmaceutical composition comprising a flumazenil complex together with a wakefulness promoting agent. According to yet another embodiment, the wakefulness promoting agent is selected from the group consisting of modafinil, armodafinil, adrafinil, methylphenidate, nefazodone, sodium oxybate,
15 phentermine, pemoline, adrenaline, methylxantines, theobromine, caffeine and a combination thereof.

According to yet another aspect, the present invention provides a flumazenil complex for the treatment of excessive sleepiness or hepatic encephalopathy, the flumazenil complex comprises flumazenil and a complexation agent, with the proviso that the complexation agent
20 is other than cyclodextrin or meglumine.

According to yet another aspect, the present invention provides a flumazenil complex for the treatment of excessive sleepiness or hepatic encephalopathy, the flumazenil complex comprises flumazenil and an aromatic complexation agent, such as, nicotinamide derivatives.

25 According to yet another aspect, the present invention provides a pharmaceutical composition comprising a flumazenil complex as the active ingredient, for the treatment of excessive sleepiness or hepatic encephalopathy.

Other objects, features and advantages of the present invention will become clear from the following description.

30 BRIEF DESCRIPTION OF THE INVENTION

Figure 1 is a schematic presentation of a flumazenil hydrochloric acid salt.

Figure 2 exhibits a chromatogram of 0.6 mg/ml nicotinamide solution in diluents.

Figure 3 shows a chromatogram of 40 µg/ml flumazenil solution in diluents.

Figure 4 shows a chromatogram of a flumazenil nicotinamide complex in diluents.

Figure 5 presents the body weight distribution of the rats (1F-treatment and 2F-placebo) in the study of reversal diazepam-induced sedative-hypnotic effects.

Figure 6 exhibits the sleeping time of the rats (1F-treatment and 2F-placebo) following
5 treatment.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to pharmaceutically acceptable flumazenil salts comprising flumazenil and a counter-ion. In addition, the present invention provides
10 pharmaceutically acceptable flumazenil complexes comprising flumazenil with a stacking complexation agent, such as an aromatic complexation agent, wherein the complexation agent is other than meglumine and cyclodextrin.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof.
15 Examples of pharmaceutically acceptable salts include, but are not limited to, inorganic or organic acid salts of basic residues such as amines and the like. The pharmaceutically acceptable salts include conventional non-toxic salts of the parent compound formed, for example, from inorganic or organic acids, such as mineral acids (e.g., HCl) and nicotinic acids. Conventional non-toxic salts include those derived from inorganic acids, such as,
20 hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids, such as, acetic, propionic, succinic, glycolic, stearic, lactic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

25 According to particular embodiments, the salts of the present invention are acid addition salts of flumazenil. According to some embodiments, the salts of the present invention are acid addition salts of flumazenil other than the acid addition salt flumazenil HCl.

Acid addition salts may, for example, be formed by mixing a solution of the compound
30 of the present invention with a solution of a pharmaceutically acceptable acid, such as, hydrochloric acid, sulfuric acid, acetic acid, trifluoroacetic acid and benzoic acid.

The terms "acid salt" and "acid addition salts" are synonymous and refer to a class of salts formed by the partial neutralization of protic, diprotic or polyprotic acids. In diprotic or

polyprotic acids, because the parent acid is only partially neutralized, one or more replaceable protons remain. Acid salt compounds can act either as an acid or a base: addition of a suitably strong acid will restore protons, and addition of a suitably strong base will remove protons. The pH of a solution of an acid salt will depend on the relevant equilibrium constants and the amounts of any additional base or acid.

According to some embodiments, the flumazenil complex of the invention comprises flumazenil and a complexation agent, wherein the complexation agent is preferably a hydrotope, other than cyclodextrins or meglumine, preferably, an aromatic stacking complexation agent.

The term “hydrotopes” as used herein, refers to compounds which are capable of opening up the water structure associated with hydrophobic (lipophilic) and other molecules. These compounds may be used to enhance the aqueous solubility of poorly water-soluble substances. Non limiting examples of hydrotopes includes sodium benzoate, sodium hydroxybenzoate, sodium salicylate, nicotinamide, sodium nicotinate, urea, sodium gentisate, gentisic acid ethanolamide, sodium toluate, sodium aminobenzoates, sodium anthranilate, sodium butylmonoglycolsulfate, resorcinol and the like. Each possibility represents a separate embodiment of the present invention.

As used herein, the term “stacking complexation agent” refers to complexation agent that forms a layered structure consisting of planar layers of the complexation agent stacked above and below the compounds with which the complex is formed. This mechanism is particularly suitable for complexing insoluble molecules having a relatively planar configuration, such as flumazenil. Preferably, the stacking complexation agent also has a relatively planar configuration. Preferred stacking complexation agents include nicotinamide and nicotinamide derivatives.

The term “nicotinamide derivatives” as used herein is meant to include any derivative of nicotinamide capable of forming a complex with flumazenil with improved solubility compared to the solubility of uncomplexed flumazenil. Acceptable nicotinamide derivatives include, but are not limited to, the nicotinamide derivatives disclosed in any one or more of: US Patent Nos. 7,056,934 and 7,153,870; WO 2008/071665; Kuramochia et al., Bioorganic & Medicinal Chemistry, Vol. 13(12), 2005, 4022-4036; and Fonti et al., Nucl Med Biol., 1999, 26(6):681-6, among others. Further non-limiting examples of nicotinamide derivatives include 2-amino-nicotinamide derivatives, 5-phenyl-nicotinamide derivatives, and 6-substituted nicotinamide derivatives.

Complex formation, which is non-covalent in nature, may be achieved by mixing appropriate ratios of the solvent-derived composition and the hydrotope or mixtures thereof in a suitable liquid vehicle, which may be aqueous, organic or a combination of both. Additional excipients, such as, surfactants, polyols, disaccharides may be added to facilitate complexation or for assisting in dispersability. The resultant complex may be isolated as a dry powder by any process known in the art (co-precipitation and drying, evaporation of the liquid vehicle, spray drying, lyophilization etc.). Particle size may be reduced by any standard technique known in the art, if desired. The resultant hydrotope complex may be used without further modification or may be compounded into a variety of other formulations or vehicles as required.

In a particular embodiment, the flumazenil stacking complex of the invention is a flumazenil nicotinamide complex. Nicotinamide, also known as niacinamide and nicotinic acid amide, is the amide of nicotinic acid (vitamin B3/niacin) and is also known as 3-pyridinecarboxamide, nicotinic acid amide and Vitamin PP.

Nicotinamide has over the past years been given at high doses for a variety of therapeutic applications. For regulatory purposes nicotinamide is classed as a food additive rather than a drug, a category that does require the formal safety evaluation normally expected of new drugs and therapies. Overall, and as detailed hereinafter, nicotinamide is considered safe. The therapeutic index of nicotinamide is wide but at very high doses (i.e. in megadoses) reversible hepatotoxicity has been reported in animals and humans. Minor abnormalities of liver enzymes can infrequently occur at the doses used for diabetes prevention. There is no evidence of teratogenicity from animal studies and there is no evidence of oncogenicity in human. Growth inhibition was shown in rats but growth in children is unaffected. In addition, minor degrees of insulin resistance attributed to nicotinamide have been reported. High-dose nicotinamide, i.e. in excess of 3 gm/day, should be considered as a drug with toxic potential at adult doses and unsupervised use should be discouraged.

The ratio of flumazenil:nicotinamide in the stacking complex of the invention is within the range of 1:1 to 1:2. In certain embodiments the flumazenil:nicotinamide ratio in the complex of the invention is about 1:1.5. This ratio increases the solubility of flumazenil such that formulations comprising about 1% flumazenil with excellent solubility are obtained. The amount of nicotinamide in such formulations is about 1.5%. Accordingly, for example, a pharmaceutical composition comprising 3 mg of flumazenil, which is about the maximal

daily dose allowed for flumazenil, corresponds to 4.5 mg of nicotinamide. An amount of 4.5 mg nicotinamide is lower by a factor of about 670 from 3 gr. (where 3 gr. is considered high dose and potentially toxic, for nicotinamide).

5 The marginal toxicity of nicotinamide renders it advantageous over other complexation agents. For example, cyclodextrins, another family of complexation agents, which increase the solubility of poorly soluble compounds by means of inclusion, are found toxic at relatively low concentrations, although there are significant differences between the cytotoxicity of the various cyclodextrin derivatives.

10 For example, the quantity of solubilized paclitaxel (as Taxol®), a poorly soluble drug, increased with cyclodextrin concentration. However, some of the cyclodextrins tested in this study were found toxic in mice with a maximum tolerated dose of 2 g/kg body weight, which is the quantity of cyclodextrin required to administer paclitaxel at 10 mg/kg. Other cyclodextrins allowed paclitaxel administration at higher doses yet had a maximum tolerated dose of 25 mg drug/kg.

15 Another advantage attributed to the flumazenil-nicotinamide complex of the invention is long shelf life. As exemplified hereinbelow, the flumazenil complex of the invention has an exceptional stability, during days, weeks and month of storage at room temperature and even at higher temperatures.

20 Without being bound to any theory or mechanism it is worth noting that cyclodextrins and nicotinamide increase solubility by different complexation mechanisms. It has been suggested that nicotinamide complex with hydrophobic ('guest') species is made out of a stack of nicotinamide and the guest species, whereby one or more 'sheets' of nicotinamide are stacked above and below a layer of the hydrophobic species. Clearly, stack formation depends on the 3D configuration of the members of the complex. While nicotinamide is
25 substantially planar, for the formation of a stacked complex, it is generally required that the hydrophobic guest species would also be planar. Complex formation by cyclodextrins is exerted by way of inclusion. Cyclodextrins form a cavity, having an inner hydrophobic core and an outer hydrophilic core, where the cavity traps the hydrophobic species. The resulting complex exhibits modified physical and chemical properties, mostly in terms of water
30 solubility. Cyclodextrins are relatively large molecules (molecular weight ranging from almost 1000 to over 1500), with a hydrated outer surface, and under normal conditions, cyclodextrin molecules will only permeate biological membranes with considerable difficulty.

The present invention further provides a pharmaceutical composition comprising the flumazenil salt or complex as the pharmaceutically active ingredient and a pharmaceutical acceptable carrier.

5 The pharmaceutical compositions according to the present invention preferably comprise one or more pharmaceutical acceptable carriers and the active constituent(s). The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. The carriers in the pharmaceutical composition may comprise a binder, such as microcrystalline cellulose, polyvinylpyrrolidone (polyvidone or povidone), gum tragacanth, gelatine, starch, lactose or lactose monohydrate; a disintegrating agent, such as alginic acid,
10 maize starch and the like; a lubricant or surfactant, such as magnesium stearate, or sodium lauryl sulphate; a glidant, such as colloidal silicon dioxide; a sweetening agent, such as sucrose or saccharin; and/or a flavoring agent, such as peppermint, methyl salicylate, or orange flavoring. More examples of the carriers include, but not limited to, physiological saline, polyethylene glycol, ethanol, vegetable oils, isopropyl myristate, etc. For topical
15 administration, the compounds of the present invention can be formulated in the form of ointment or cream. Each possibility represents a separate embodiment of the present invention.

The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipients thereof.

20 The pharmaceutical composition comprising the flumazenil salt or complex of the invention as the active ingredient, may further comprise cyclodextrins, at least one solubilizing agent, at least one penetration enhancer, and at least one preservative. The pharmaceutical composition may further comprise additional excipients, such as, flavoring agents, among others.

25 The solubilizing agent is preferably a polar solvent such as mono- or poly-alcohols of linear or branched configuration (e.g., C1 to C8 alcohols). Non-limiting examples include methanol, ethanol, propanol, iso-propanol, n-butanol, sec-butanol, isobutanol, t-butanol, n-pentanol, 2-pentanol, 3-pentanol, neopentanol, 1-hexanol, 2-hexanol, 3-hexanol, 1-heptanol, 2-heptanol, 1-octanol, as well as any geometrical isomers, enantiomers and diastereomers of
30 any of the foregoing. Other suitable polar solvents include glycols such as ethylene glycol, propylene glycol and their polymers having a molecular weight between 400 and 1000. Each possibility represents a separate embodiment of the present invention. In one particular embodiment, the solubilizing agent is a combination of ethanol and propylene glycol.

Preferred solubilizing agents include alcohols, glycols and combinations thereof. Each possibility represents a separate embodiment of the present invention.

According to some embodiments, the solubilizing agent in the composition of the invention comprises a combination of an alcohol and a glycol wherein the alcohol:glycol ratio is at least 1.5:1 or in the range of 1.5:1 to 5:1. The solubilizing agent, preferably in the form of combination of an alcohol and a glycol, may be at least 40%, or in the range of 40% to 60%, of the total weight of the pharmaceutical composition of the invention.

The expression "cyclodextrin" as used herein means α -, β - or γ - cyclodextrin or a derivative thereof. Suitable cyclodextrin derivatives for use in the formulations of the present invention include, but are not limited to the cyclodextrin listed above, e.g. hydroxypropyl derivatives of α -, β - and γ -cyclodextrin, sulfoalkylether cyclodextrins such as sulfobutylether β -cyclodextrin, alkylated cyclodextrins such as the randomly methylated β -cyclodextrin, and various branched cyclodextrins such as glucosyl- and maltosyl β -cyclodextrin. Other cyclodextrins are described in US patent publication US 2004/0186075.

In one currently preferred embodiment, the cyclodextrin is hydroxypropyl β -cyclodextrin (HPCD). The cyclodextrin is preferably formulated in a buffer having a pH from about 3 to about 6. In one particular embodiment, the cyclodextrin (e.g., HPCD) is formulated in a citric acid buffer having a pH of about 4.

The cyclodextrin component of the formulations of the present invention may be present in an amount from about 10% to about 95% w/w, for example from about 30% to about 80%, from about 30% to about 75%, or about 60% based on the formulations of the invention. It is generally recognized that cyclodextrins act as true carriers by keeping the hydrophobic drug molecules in solution and delivering them to the surface of the biological membrane, e.g. skin, mucosa or the eye cornea, where they partition into the membrane. The relatively lipophilic membrane has low affinity for the hydrophilic cyclodextrin molecules and therefore they remain in the aqueous membrane exterior, e.g. the aqueous vehicle system, saliva or the tear fluid. Conventional penetration enhancers, such as alcohols and fatty acids, disrupt the lipid layers of the biological barrier. Cyclodextrins, on the other hand, may act as penetration enhancers by increasing drug availability at the surface of the biological barrier.

Therapeutic formulations suitable for oral administration, e.g. tablets and pills, may be obtained by compression or moulding, optionally with one or more accessory ingredients.

Compressed tablets may be prepared by mixing the constituent(s), and compressing this mixture in a suitable apparatus into tablets having a suitable size. Prior to the mixing, the phanquinone may be mixed with a binder, a lubricant, an inert diluent and/or a disintegrating agent and the further optionally present constituents may be mixed with a diluent, a lubricant and/or a surfactant. A tablet may be coated or uncoated. An uncoated tablet may be scored. A coated tablet may be coated with sugar, shellac, film or other enteric coating agents. Each possibility represents a separate embodiment of the present invention.

Therapeutic formulations suitable for parenteral administration include sterile solutions or suspensions of the active constituents. An aqueous or oily carrier may be used. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soy bean oil, mineral oil, sesame oil and the like. Formulations for parenteral administration also include a lyophilized powder comprising flumazenil and, optionally, further active constituents, that is to be reconstituted by dissolving in a pharmaceutically acceptable carrier that dissolves the active constituents, e.g. an aqueous solution of carboxymethylcellulose and lauryl sulphate. Each possibility represents a separate embodiment of the present invention.

When the pharmaceutical composition is a capsule, it may contain a liquid carrier, such as fatty oil, e.g. cacao butter.

Additional suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Each possibility represents a separate embodiment of the present invention.

The formulations of the present invention may further comprise penetration enhancers, such as menthol. Other penetration enhancers that may be used in the formulations of the present invention include, but are not limited to, anionic surfactants (e.g. sodium lauryl sulphate, sodium dodecyl sulphate), cationic surfactants (e.g. palmitoyl DL camitine chloride, cetylpyridinium chloride), nonionic surfactants (e.g. polysorbate 80, polyoxyethylene 9-lauryl ether, glyceryl monolaurate, polyoxyalkylenes, polyoxyethylene 20 cetyl ether), lipids (e.g. oleic acid), bile salts (e.g. sodium glycocholate, sodium taurocholate), chitosan or a chitosan derivative, linalool, carvacrol, thymol, citral or t-anethole, and related compounds. Each

possibility represents a separate embodiment of the present invention.

The formulations of the invention optionally further comprise at least one preservative. Any suitable preservative may be present in the formulation in the present invention. The preservative may be any pharmaceutically acceptable preservative, for example methyl 4-
5 hydroxybenzoate (methyl paraben), ethyl 4-hydroxybenzoate (ethyl paraben), propyl 4-hydroxybenzoate (propylparaben), benzyl alcohol, sorbic acid, sodium benzoate, benzoic acid, and any combination thereof. Each possibility represents a separate embodiment of the present invention.

The formulation of the invention optionally further comprises a flavoring agent in an
10 amount between 0.05 and 10 percent by weight of the total composition. In one embodiment, the flavoring agent is present in an amount between 0.1 and 2.5 percent by weight of the total composition. The flavoring agent is preferably selected from the group consisting of synthetic or natural oil of peppermint, oil of spearmint, citrus oil, fruit flavors, sweeteners (sugars, aspartame, saccharin, Estevia, etc.), and mixtures thereof. Menthol can also act as a flavoring
15 agent.

In yet another embodiment, the therapeutic compound can be delivered in a controlled release system. The term "controlled release" is used herein to refer to a pharmaceutical dosage form in which release of the active ingredient is timed or modified to a rate sufficient to maintain the desired therapeutic level over an extended period of time. The release may be
20 a "sustained release" or a "delayed release" such that release of the active ingredient from the pharmaceutical dosage form is other than promptly after administration of the dosage form, but rather is withheld or delayed following administration.

In one embodiment, a pump may be used. In another embodiment, polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in
25 proximity of the therapeutic target, i.e., the central nervous system, thus requiring only a fraction of the systemic dose. Other controlled release systems are discussed, for example, in U.S. Patent No. 5,120,548 which is directed a controlled-release drug delivery device comprised of swellable polymers. U.S. Patent No. 5,073,543 also describes controlled-release formulations containing a trophic factor entrapped by a ganglioside-liposome vehicle. U.S.
30 Patent No. 5,639,476 discloses a stable solid controlled-release formulation having a coating derived from an aqueous dispersion of a hydrophobic acrylic polymer. Biodegradable microparticles are also known for use in controlled-release formulations. U.S. Patent No. 5,733,566 describes the use of polymeric microparticles that release antiparasitic

compositions.

The controlled-release of the active ingredient may be stimulated by various inducers, for example, pH, temperature, enzymes, water, or other physiological conditions or compounds.

5 The terms "composition," "formulation" and "dosage form" are used herein interchangeably to encompass formulated preparations comprising one or more pharmacologically active drugs, and one or more pharmaceutically acceptable excipients, diluents or carriers. Compositions, formulations and dosage forms can be designed for administration by all possible administration routes to achieve the desired therapeutic
10 response. The terms used may refer to the physical format of the product which is dispensed and administered to the patient, for example, a capsule or a patch. Alternately or in addition, the terms used may refer to any of: the mode of administration, the mode of delivery or the mode of release of the drug, for example a transdermal delayed release formulation.

15 The compounds of the present invention may be formulated into injections by dissolving, suspending or emulsifying water-soluble solvent such as saline and 5% dextrose, or in water-insoluble solvents such as vegetable oils, synthetic fatty acid glyceride, higher fatty acid esters and propylene glycol. The formulations of the invention may include any of conventional additives such as dissolving agents, isotonic agents, suspending agents, emulsifiers, stabilizers and preservatives.

20 According to yet another aspect, the present invention provides a method for treating a disease or disorder associated with binding of an agent to the GABA receptor, comprising administering to a patient in need thereof a pharmaceutical composition comprising an active ingredient selected from flumazenil salt and flumazenil complex, and at least one pharmaceutical acceptable carrier. Thus, the pharmaceutical composition of the invention is
25 intended for use as an antidote for GABA related sedatives/hypnotics.

Preferably, the disease or disorder is excessive sleepiness, the agent is a benzodiazepine or a non-benzodiazepine, and the effect is sedation.

As used herein, the term "treating" encompasses substantially ameliorating, relieving, alleviating and preventing symptoms of a disease, disorder or condition in a subject.

30 As used herein, the term "administering" refers to delivery of a pharmaceutical compound to a subject by any means that does not affect the ability of the compound to perform its intended function.

The terms "sleep drug" is used herein in reference to pharmaceutical agents used for

inducing and/or maintaining sleep, in particular, prescription sleep drugs that are classified as hypnotics/sedatives. The sleep-drug may be a sedative/hypnotic drug, a benzodiazepine, a benzodiazepine modulator, a benzodiazepine analog, a non-benzodiazepine, a 5-HT_{2A} receptor antagonist, a melatonin receptor agonist, an orexin receptor antagonist, a selective serotonin reuptake inhibitor (SSRI), an antihistamine and an herbal product.

Suitable benzodiazepine sleep drugs include alprazolam, bromazepam, clonazepam, clotiazepam, cloxazolam, diazepam, estazolam, etizolam, fludiazepam, flunitrazepam, flurazepam, halazepam, haloxazolam, lorazepam, medazepam, midazolam, nimetazepam, nitrazepam, olanzapine, oxazepam, quazepam, temazepam and triazolam.

Suitable non-benzodiazepine sleep drug include adiplon, agomelatine, almoxerant, brotizolam, diphenhydramine, divaplon, doxepin, eplivanserin, doxylamine succinate, eszopiclone, indiplon, ocinaplon, pagoclone, pazinaclone, pruvanserin (EMD 281014), suproclone, suriclone, L-tryptophan, 5-hydroxy-L-tryptophan, melatonin, melatonin receptor agonists, muramyl dipeptide, ramelteon, uridine, volinanserin, zaleplon, zolpidem, imidazopyridine hypnotic zolpidem, APD125, ACP-103, PD 200-390, HY10275, GW649863 and EVT-201

As used herein, the term "effective amount" refers to an amount of a pharmaceutical compound sufficient to achieve its desired effect.

According to one embodiment, the excessive sleepiness is selected from the group consisting of: excessive sleepiness associated with hypersomnia, excessive sleepiness associated with drowsiness, symptoms associated with overdose of a sleep drug, excessive sleepiness associated with alcohol intoxication, excessive sleepiness associated with sedation and anesthesia induced by a sleep drug, excessive sleepiness associated with sleep drug addiction, excessive sleepiness associated with stimulant addiction, the rebound effect of a sleep drug, balance impairment induced by a sleep drug or any combination thereof.

According to certain embodiments, treating excessive sleepiness comprises reversing benzodiazepine-induced sedation and anesthesia following therapeutic or diagnostic procedures.

As used herein, the term "hypersomnia" refers to chronic or recurrent bouts of excessive sleepiness, characterized by one or more of near-daily diurnal sleep episodes, excessive naps, abnormally prolonged sleep intervals, a perception of non-restorative sleep, and difficulty in making the transition from sleep to wakefulness. Hypersomnia may be one or more of: shift work sleep disorder; narcolepsy; obstructive sleep apnea/hypopnea

syndrome; REM behavior disorder; frontal nocturnal dystonia; restless legs syndrome; nocturnal movement disorder; Kleine-Levin syndrome; Parkinson's disease; Alzheimer's disease, schizophrenia, anxiety, schizophrenia (in particular, schizophrenia associated with excessive sleepiness), excessive sleepiness; hypersomnia; idiopathic hypersomnia; recurrent
5 hypersomnia; endozepine related recurrent stupor; and amphetamine resistant hypersomnia.

According to yet another embodiment, the excessive sleepiness is caused by alcohol intoxication.

According to yet another aspect, the method of the invention is directed to treating alcohol intoxication.

10 The term "alcohol intoxication" means overdose of alcohol (e.g., ethanol) leading to behavioral impairment. A person is said to suffer from alcohol intoxication when the quantity of alcohol the person consumes exceeds the individual's tolerance for alcohol and produces behavioral or physical abnormalities. In other words, the person's mental and physical abilities are impaired.

15 In this respect is it noted that the term alcohol refers to the generic term for ethanol, which is a particular type of alcohol produced by the fermentation of many foodstuffs—most commonly barley, hops, and grapes. Other types of alcohol commonly available such as methanol (common in glass cleaners), isopropyl alcohol (rubbing alcohol), and ethylene glycol (automobile antifreeze solution) are highly poisonous when swallowed, even in small
20 quantities. Ethanol produces intoxication because of its depressive effects on various areas of the brain causing these impairments in a progressive order as the person gets more and more drunk. Symptoms of alcohol intoxication and/or impaired performance after alcohol consumption include disinhibition of normal social functioning (e.g., excessive talking), loss of memory, confusion, disorientation, uncoordinated movement, progressive lethargy, coma,
25 or ultimately death.

According to one another embodiment, treating excessive sleepiness caused by alcohol intoxication is selected from the group consisting of: reversing the effects of alcohol intoxication, reducing the effects of alcohol intoxication, alleviating the effects of alcohol intoxication and improving performance after alcohol consumption.

30 According to yet another aspect, the method of the invention is directed treating hepatic encephalopathy with the flumazenil formulation.

The method of the invention provides an improved treatment of hepatic encephalopathy. First, the pharmaceutical compositions of the present invention are suitable

for oral, sublingual or transdermal administration. Second, the pharmaceutical composition of the present invention includes much higher concentrations of flumazenil, of about 0.4 to 2%. This concentration is higher by two orders of magnitude from the concentration of flumazenil in the commercially available flumazenil formulations that are known to date. Thereby the pharmaceutical compositions and methods of the present invention provide advantageous patient compliance. In addition, the present invention provides attractive routes for the delivery of flumazenil (e.g. sub-mucosal and transdermal) thereby averting the need for admission to health centers.

Without being bound by any theory or mechanism, administering the salts and complexes of the present invention by submucosal or sublingual or transdermal routes, bypasses the first pass effect (through the liver). This mechanism probably confers a material advantage in the treatment of the aforementioned diseases and disorders (residual effect of hypnotics, sedation, hepatic encephalopathy, etc.). Hepatic encephalopathy refers to a complex neuropsychiatric syndrome, which may complicate acute or chronic hepatic failure.

It is characterized by changes in mental state including a wide range of neuropsychiatric symptoms ranging from minor not readily discernible signs of altered brain function, overt psychiatric and/or neurological symptoms to deep coma. Accordingly, the methods to estimate treatment effects and treatment outcomes are highly variable. The majority of hepatic encephalopathy occurs in patients with cirrhosis, often associated with spontaneous or iatrogenic portal-systemic shunting. Hepatic encephalopathy is generally considered a reversible metabolic encephalopathy. Traditionally, hepatic encephalopathy has been considered to be secondary to the accumulation of toxic products, which have not been metabolized by the liver. Various hypotheses have been suggested, e.g., alterations in the permeability of the blood-brain barrier, abnormal neurotransmitter balance, altered cerebral metabolism, and increased amounts of endogenous benzodiazepine-like compounds - the gamma-amino butyric acid (GABA)/benzodiazepine hypothesis. GABA is the principal inhibitory neurotransmitter in mammals that acts by binding to a receptor on a 'supramolecular complex' called the GABA/benzodiazepine complex, which also has binding sites for benzodiazepines and barbiturates. By binding to the GABA/benzodiazepine complex, benzodiazepines cause sedation through neural inhibition. It has been suggested that liver failure leads to the accumulation of substances that bind to the GABA/benzodiazepine complex resulting in neural inhibition which may progress to coma. Accordingly, a benzodiazepine-receptor antagonist, flumazenil, has been assessed in the

treatment of hepatic encephalopathy in the hope of reversing neuropsychiatric symptoms related to the accumulation of endogenous benzodiazepine.

As detailed above, Als-Nielsen et al. (*ibid*) disclose the use of flumazenil, administered via intravenous infusion of 1 mg flumazenil in 20 ml saline solution (0.005%) over 3-5 minutes, for treating hepatic encephalopathy. Als-Nielsen reports that flumazenil had a significant beneficial effect on short-term improvement of hepatic encephalopathy in patients with cirrhosis and a highly favorable prognosis but had no significant effect on recovery or survival.

The methods of the invention may be used in conjunction with insomnia treatment modalities, and serve to eliminate or diminish residual soporific effects associated with administration of sleep drugs. The invention is effective for counteracting excessive sleepiness induced by a wide variety of sleep drugs. Such sleep drugs include benzodiazepine and non-benzodiazepine drugs which are classified as hypnotics/sedatives, as well as other prescription and non-prescription sleep drugs, including those classified as 5-HT_{2A} receptor antagonists, melatonin receptor agonists, orexin receptor antagonists, selective serotonin reuptake inhibitors (SSRIs), and other sleep inducing agents such as antihistamines, melatonin and certain herbal products. It is to be specifically understood that a particular sleep drug may be classified under more than one of the aforementioned categories.

According to another embodiment, the route of administration is selected from the group consisting of: oral, buccal, sublingual, transdermal, transmucosal, intranasal, intravenous (*i.v.*), intraperitoneal (*i.p.*), intramuscular (*i.m.*), subcutaneous (*s.c.*) or intrathecal (*i.t.*).

For buccal administration, buccal tablets or sublingual tablets may be used. These tablets are typically small, flat and soft, designed to be placed in the side of the cheek (i.e. buccal cavity) or designed to be placed under the tongue, to be directly absorbed through the buccal mucosa for a systemic effect. Other dosage forms suitable for buccal administration are, for example, oral films administered on the gingiva or tongue.

Sublingual spray is also a buccal formulation for delivery to the sublingual mucosa in the form of a spray for a systemic effect, typically provided in spray actuators, designed to access the mucosal surfaces under the tongue or the lips.

For transdermal delivery of the composition of the invention, the composition may be provided in the form of a patch. The major approaches for transdermal delivery include use of chemical penetration enhancers; physical enhancers, such as ultrasound, iontophoresis,

electroporation, magnetophoresis, and microneedles; vesicles; particulate systems, such as those incorporating liposomes, niosomes, transfersomes, microemulsions, or solid lipid nanoparticles, as described for example in Rizwan et al., Recent Pat Drug Deliv Formul., 2009, 3(2):105-24.

5 Suitable penetration enhancers for transdermal delivery include, for example, saturated and unsaturated fatty acids and their esters, alcohols, monoglycerides, diethanolamines, N,N-dimethylamines such as linolenic acid, linolenyl alcohol, oleic acid, oleyl alcohol, stearic acid, stearyl alcohol, palmitic acid, palmityl alcohol, myristic acid, myristyl alcohol, 1-dodecanol, 2-dodecanol, lauric acid, decanol, capric acid, octanol, caprylic acid, 1-
10 dodecylazacycloheptan-2-one, ethyl caprylate, isopropyl myristate, hexamethylene lauramide, hexamethylene palmitate, capryl alcohol, decyl methyl sulfoxide, dimethyl sulfoxide, salicylic acid and its derivatives, N,N-diethyl-m-toluamide, 1-substituted azacycloalkan-2-ones, propylene glycol, polyethylene and glycol monolaurate. Any compound compatible with flumazenil, and that has transdermal permeation enhancing
15 activity may be selected.

 Creams for transdermal delivery of flumazenil typically include gelling agents, for example, hydroxy methyl cellulose, hydroxypropyl cellulose, tragacanth, sodium alginate, gelatin, methylcellulose, sodium carboxymethylcellulose and polyvinyl alcohols.

 The efficacy of the methods described herein may be assessed, for example by direct
20 observation of behavioral and physiological properties, by self-reporting, and/or by various well-known electrophysiological methods and performance skill methods. Such methods include, for example, examining electroencephalograph (EEG) activity amplitude and frequency patterns, examining electromyogram activity, and examining the amount of time during a measurement time period, in which a mammal is awake or exhibits a behavioral or
25 physiological property characteristic of wakefulness.

 Objective and subjective tests for wakefulness, alertness and performance include, for example, the Epworth Sleepiness Scale (Johns MW (1991) Sleep 14 (6): 540-5) and the Stanford Sleepiness Scale (Hoddes et al (1972) Psychophysiology 9:150).

 Additional methods used to monitor or assess alertness/drowsiness levels in a subject
30 prior to and following use of the methods disclosed herein may employ various devices for measurement of eye position or closure, assumed to correlate with alertness/drowsiness, as disclosed for example in U.S. Patent Nos. 5,689,241; 5,682,144 and 5,570,698.

 The flumazenil compositions according to the present invention may also be used in

conjunction with the administration of other pharmaceutically active compounds. In addition, the pharmaceutical composition according to the present invention may contain other pharmaceutically active compounds.

According to some embodiments, the methods of the invention further comprise co-administering, concomitantly or in sequence, the pharmaceutical composition comprising the flumazenil salt or flumazenil complex of the invention together with a wakefulness promoting agent. According to some embodiments, the wakefulness promoting agent is selected from the group consisting of modafinil, armodafinil, adrafinil, methylphenidate, nefazodone, sodium oxybate, phentermine, pemoline, adrenaline, methylxantines, theobromine, caffeine and a combination thereof.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form described by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

The following examples are presented in order to more fully illustrate certain embodiments of the invention. They should in no way, however, be construed as limiting the broad scope of the invention. One skilled in the art can readily devise many variations and modifications of the principles disclosed herein without departing from the scope of the invention.

EXAMPLES

25

Example 1. Formulations of flumazenil salts and complexes at 1% flumazenil

Formulations with 1% flumazenil, in the form of a flumazenil salt or flumazenil complex, are presented in the following Tables. All formulations were evaluated for their appearance and only clear solutions, namely, clear solutions where a precipitate ('ppt') or slight precipitate ('sl-ppt') or a few particles ('part') are not formed, were tested for pH.

Compositions comprising nicotinamide as a complexation agent and meglumine resulted in clear solutions at first. Twelve hours later, in formulations containing the

meoglumine a precipitate was formed. Formulation No. 2 containing 4.5% nicotinamide showed less precipitate (designated 'ppt' in the Tables) after 12 hours compared to formulation No. 1 containing 1.5% nicotinamide (Table 1). This result indicates that high nicotinamide concentrations may be preferred, although not necessary in order to achieve the concept of the present invention. The appropriate pH values of a formulation intended for sublingual administration are between 4.0 to 9.0. The pH values of the nicotinamide-containing formulations meet this requirement ($\text{pH} = 5.0 \pm 0.2$).

Table 1. Formulations of flumazenil-nicotinamide and flumazenil meglumine (% w/w)

Material name	1	2	3	4	5	6
Flumazenil	1	1	1	1	1	1
Ethanol	25	25	25			
Propylene glycol	25	25	25			
Citric buffer 10mM pH 4.0	47.4	44.4	48.7			
Menthol in EtOH (1/1 w/w)	0.1	0.1	0.1			
Nicotinamide/WFI (1/2 w/w)	1.5	4.5				
Benzalkonium chloride (BKC) 50% aq. solution			0.2			
Water (WFI)				98.67	98.67	98.356
HCl				0.33		
H ₂ SO ₄					0.33	
Meglumine						0.644
Appearance t=0	clear	clear	ppt	ppt	ppt	ppt
Appearance T=12h	sl-ppt	sl-ppt				
pH	5.06	5.15				

Table 2. Formulations with cyclodextrin (HPCD; % w/w)

Material name	1'	2'	3'	4'	5'	6'
Flumazenil	1	1	1	1	1	1
Ethanol	20	20	20			
Propylene glycol	20	20	20			
30% HPCD in citric buffer	57.4	54.4	58.7			
30% HPCD in water				98.67	98.67	98.36
Menthol in EtOH (1/1 w/w)	0.1	0.1	0.1			
Nicotinamide/WFI (1/2 w/w)	1.5	4.5				
BKC 50% aq. solution			0.2			
HCl				0.33		
H ₂ SO ₄					0.33	
Meglumine						0.644
Appearance t=0	ppt	ppt	ppt	ppt	ppt	ppt

Table 3. Formulations with flumazenil meglumine complex (% w/w)

Material name	7	8	9	8'	9'
Flumazenil	1	1	1	1	1
Ethanol	25	25	25	20	20
Propylene glycol	25	25	25	20	20
WFI	47.4	44.4			
30% HPCD in water				57.4	54.4
Citrate buffer			44.4		
Menthol in EtOH (1/1 w/w)	0.1	0.1	0.1	0.1	0.1
Meglumine in WFI (1/2 w/w)	1.5	4.5	4.5	1.5	4.5
Appearance t=0	clear	clear	clear	ppt	ppt
Appearance t=12 h	sl- ppt	sl- ppt	ppt		
pH	10.91	11.32	10.50		

Table 4. Formulations with flumazenil nicotinamide complex and flumazenil meglumine complex (% w/w)

Material name	10	11	12	13	14	15	16
Flumazenil	1	1	1	1	1	1	1
Ethanol	25	25	25	30	35	35	40
Propylene glycol	25	25	25	20	20	15	10
Citric buffer 10mM pH 4.0	41.5	46.9	43.9	44.5	39.5	44.5	44.5
Nicotinamide/WFI (1/2 w/w)	7.5	1.5	4.5	4.5	4.5	4.5	4.5
Meglumine in WFI (1/2 w/w)		0.6	0.6				
Appearance t=0	clear	clear	clear	clear	clear	clear	clear
pH, T-0	5.10	9.02	9.12	5.09	5.25	5.16	5.17
Appearance t=24-72h	sl-ppt	ppt	ppt	sl-ppt	clear	clear	clear
Appearance t=24-72h	10.91	11.32	10.50				

5

Example 2. Formulations of flumazenil (1.2 and 1.5 % w/w) nicotinamide complex

Formulations containing 1.2 to 1.5 % w/w flumazenil in the form of a flumazenil complex with nicotinamide were prepared and are presented in the following Table.

Table 5. Flumazenil formulations with flumazenil nicotinamide complex (% w/w)

Material name	14A	15A	16A	14B	16B	17	18	19	20
Flumazenil	1.5	1.5	1.5	1.2	1.2	1.5	1.5	1.5	1.5
Ethanol	35	35	40	35	40	40	35	40	30
Propylene glycol	20	15	10	20	10	20	20	10	30
Citric buffer 10mM pH 4.0	39	44	44	39.3	44.3	34	36	41	34
Nicotinamide/WFI (1/2 w/w)	4.5	4.5	4.5	4.5	4.5	4.5	7.5	7.5	4.5
Appearance T-0	clear	clear	clear	clear	clear	clear	clear	clear	clear
pH, T-0	5.27	5.18	5.24	5.25	5.26	5.44	5.28	5.26	5.30
Appearance T-24-72h	sl-ppt	ppt	sl-ppt	clear	clear	clear	ppt	ppt	ppt
Appearance T-1week				part	clear	part			

Example 3. Formulations of flumazenil nicotinamide complexes and menthol

Formulations containing 1.2 % w/w flumazenil based on formulation 16 B (Table 6, 5 above), were prepared with menthol and are presented in the following Table.

Table 6. Flumazenil formulations with 1.2 flumazenil (% w/w) and menthol

Material name	0.2%menthol		0.1%menthol	
	%	g	%	g
Flumazenil	1.2	0.06	1.2	0.06
Ethanol	40	2	40	2
Propylene glycol	10	0.5	10	0.5
Citric buffer 10mM pH 4.0	43.9	2.195	44.1	2.205
Nicotinamide/WFI (1/2 w/w)	4.5	0.225	4.5	0.225
Menthol/EtOH (1/1 w/w)	0.4	0.02	0.2	0.01
Appearance T-0	clear		clear	
pH, T-0	5.17		5.25	
Appearance T-24h	part		clear	

The results are similar to the results obtained in the absence of menthol (see Tables 4-5, 10 above) indicating that menthol is not necessary in order to achieve the concept of the present invention, particularly, the improved solubility of the flumazenil salts or complexes.

Example 4. HPLC analysis of flumazenil in the presence of nicotinamide

The conditions for determining by HPLC the presence of flumazenil in solutions containing flumazenil complex (1% (10 mg/g) or 1.2% (12 mg/g) flumazenil and 1.5% nicotinamide), were determined.

The details of the tested compounds were as follows:

1. Flumazenil [$C_{15}H_{14}FN_3O_3$], 303.29 g/mol (Chemagis™, purity: 99.3%,)
2. Niacinamide USP (Nicotinamide) [$C_6H_6N_2O$], 122.12 g/mol (Spectrum™, purity: 99.5%)

The HPLC (High-performance liquid chromatography) agents were HPLC grade water, methanol, acetonitrile, ammonium acetate, ammonium acetate and acetic acid glacial. Analysis was performed utilizing the column Inertsil™, G.L. Sciences, ODS-3V, 5mm, 250x4.6 mm, C.N. 5020-01802 and the pre-column: Phenomenex™ C18, 5 mm.

For Mobile Phase A, 13 mM Ammonium acetate buffer in water (pH adjusted at 4.40 with acetic acid) was utilized and acetonitrile for Mobile Phase B. Analysis was performed at column temperature of $30^\circ \pm 5^\circ\text{C}$, autosampler temperature of $10^\circ \pm 5^\circ\text{C}$, injection volume of 20 ml, PDA programmed 210 to 330 nm, sampling rate = 2, resolution = 1.2 and monitoring wavelength: 247 nm. The resulting gradient Table is given below.

Table 7. HPLC calibration

Time (min)	Flow (ml/min)	%A	%B	Curve
0	1	80	20	6
0.5	1	80	20	6
8.0	1	50	50	6
8.01	1.2	50	50	6
13.0	1.2	50	50	6
13.01	1.2	80	20	6
19.0	1	80	20	6

Samples of the following liquid formulations were prepared: Formulation 16 (Table 4; 1% flumazenil), Formulation 16B (Table 5; 1.2% flumazenil) and Formulation 16B with menthol (Table 6; 1.2% flumazenil). Solutions of flumazenil and of nicotinamide were made by mixing the dry compounds with the diluent (acetonitrile/water/acetic acid, 20/80/0.1 v/v/v).

Nicotinamide solution was injected and nicotinamide was eluted at retention time

(RT) of 3.2 min. Good resolution was obtained between nicotinamide and flumazenil, since the retention time of flumazenil is about 9.4 min. Representative chromatograms are presented in Figures 2-4.

Assay of flumazenil solutions was determined using single point calibration. Flumazenil standard solution 40ug/ml was injected five times. Flumazenil sample solutions were prepared in triplicate and injected once. The assay (%) of flumazenil in a sample was calculated according to the following formula:

$$\text{Assay (\%)} = (\text{Wstd} \times \text{P} \times \text{Dsm} \times \text{Asmp} / (\text{Dstd} \times \text{Astd} \times \text{Dose} \times \text{Wsm})) \times 100$$

Where:

- Wstd is the weight of flumazenil standard in mg;
- P is the purity of flumazenil standard (99.3%);
- Dstd is the dilution factor of the calibration solution (500);
- Dsm is the dilution of the sample solution in ml (100 ml);
- Wsm is the weight of sample solution in the sample in mg;
- Dose is the theoretical amount (%) of flumazenil in the solution;
- Astd is an average peak area of flumazenil for the five first injections of flumazenil calibration solution;
- Asmp is an average peak area of Flumazenil in the sample.

The assay results are summarized in the Table below.

Table 8. Flumazenil formulation analysis

Formulation No.	Sample weight (mg)	Peak area (%)	Assay (%)	Mean assay (%)
16	412.18	2956920	103.64	103.8
	411.98	2964443	103.96	
16B	338.43	2744305	97.63	100.9
	337.90	2863082	102.01	
	339.55	2907803	103.1	
16B with menthol	327.52	2803639	103.06	101.6
	328.91	2757927	100.95	
	332.91	2783892	100.68	

The results indicate that under the tested conditions there is no interference of nicotinamide signal with the signal of flumazenil. Accordingly, the aforementioned HPLC parameters are appropriate for evaluating the content of flumazenil in formulations comprising the flumazenil nicotinamide complex of the invention.

Example 5. HCl acid addition salt of flumazenil

Flumazenil chloride salt consisting of flumazenil and hydrochloride acid was prepared (Fig. 1). The increased solubility of the flumazenil chloride salt enabled formulating a composition comprising flumazenil at a concentration of 0.66%. The concentration of flumazenil in that composition is higher by about one order of magnitude than the concentration of flumazenil in the commercial product Romazicon® (Romazicon® contains 0.01% flumazenil).

Example 6. Stability analysis

The short term stability (STS) of flumazenil-nicotinamide formulations was tested on two flumazenil solutions (Table 9).

Table 9. Formulations participated in the STS study (% w/w)

Material	Batch No. 1	Batch No. 2
Flumazenil	1	1.2
Ethanol (absolute)	40.1	40.1
Propylene glycol	10	10
Citric acidic anhydrous	0.05	0.05
Niacinamide (nicotinamide)	1.5	1.5
Menthol	0.1	0.1
Water for injection	47.2	47.0

The formulations were packed in 5-ml glass vials, capped with Teflon stoppers and crimped with aluminium seals. Vials were stored in a stability chamber at 25°C and 40°C. Stability was evaluated at the following stability time points: 2 weeks, 1 months and 3 months. At each stability time point samples were tested for: pH, appearance and flumazenil assay.

The results are summarized in Table 10 below. The assay values for flumazenil were in the range of 90-110% and did not change over 3-months storage at 25°C and at 40°C. The pH of the formulations remained in the range of 5.13-5.39 throughout the experiment (3 months) under all temperatures. Furthermore, the solutions remained clear during storage in the stability chambers over the period and storage conditions that were tested. Thus, the results clearly indicate that the flumazenil-niacinamide formulations are stable and suitable for long shelf lives.

Table 10. STS study

Time point	Batch No. API content Storage Temperature	1		2	
		1%		1.2%	
		25°C	40°C	25°C	40°C
T-0	pH	5.17		5.20	
	Assay (%)	105.1		107.9	
T-2 weeks	Appearance	clear	clear	clear	clear
	pH	5.24	5.26	5.30	5.34
	Assay (%)	100.6	102.2	100.3	102.2
T-1 month	Appearance	clear	clear	clear	clear
	pH	5.24	5.38	5.31	5.39
	Assay (%)	99.5	100.2	99.6	98.4
T-3 months	Appearance	clear	clear	clear	clear
	pH	5.13	5.13	5.28	5.25
	Assay (%)	103.0	102.7	102.9	103.4
	Appearance	clear	clear	clear	clear

Example 7. Toxicity studies in dogs

Toxicity of the flumazenil formulation of the invention is conducted in dogs. Four
5 formulations are tested: vehicle (control), low dose, medium dose and higdose, according to
the following study design:

Table 11. Toxicity study design

	Main Study		Recovery	
	Male	Female	Male	Female
Dose				
Vehicle	3	3	2	2
Low dose	3	3	2	2
Medium dose	3	3	2	2
High dose	3	3	2	2

Flumazenil is administered once a day, sublingually. The dose is sparyed under the
10 tongue and is held there for at lest 5 minutes prior to swallowing.

All animals fast overnight (approximately 12 hours) prior to dosing and at least 2 hours
postdose. Animals are observed twice daily and undergo detailed clinical observation weekly.
The weekly observation include evaluating body weights and food consumption. Prior to
initiation of the experiment, animals undergo physical examinations by staff veterinarian.

15 All animals pretest and at termination and recovery undergo ophthalmology evaluation.
In addition, animal undergo electrocardiograms pretest, predose and postdose on day 1,
predose and postdose prior to the terminal necropsy, and prior to the recovery necropsy.

Clinical pathology (hematology, coagulation, clinical chemistry and urinalysis) is

obtained for animals pretest, and all survivors prior to the terminal and recovery necropsies. The tests are listed in Table 12 below.

Blood toxicokinetics is performed on blood samples collected on days 1 and 28 at six time points from each animal. Following necropsy, at the end of the experiment, weights of the following organs is measured: adrenals, brain, heart, kidneys, liver, lungs, ovaries with oviducts, pituitary, prostate, salivary glands, spleen, thyroid with parathyroid, thymus, testes and uterus. For all main study animals, a full set of standard tissues for slide preparation/microscopic pathology is collected (approximately 70). The target organs in recovery animals include gross lesions. A complete list of the target organs is given in Table 13 below. For toxicokinetic modeling standard parameters are evaluated, for example, AUC, $t_{1/2}$, Tmax and Cmax. Data is analyzed by standard statistical analysis.

Table 12: Standard clinical pathology tests

Standard hematology parameters evaluated	
Absolute and percent reticulocytes	Mean corpuscular hemoglobin
Erythrocyte count	Mean corpuscular hemoglobin concentration
Hematocrit	Mean corpuscular volume
Hemoglobin	Platelet count
Leukocyte count (total and differential)	Reticulocytes
Standard coagulation parameters evaluated	
Prothrombin time	Activated partial thromboplastin time
Standard clinical chemistry parameters evaluated	
Albumin/Globulin Ratio (calculated)	Glucose
Alanine aminotransferase	Phosphorus
Albumin	Potassium
Alkaline phosphatase	Sodium
Aspartate aminotransferase	Sorbitol dehydrogenase
Calcium	Total bilirubin (with direct if total exceeds 1 mg/dL)
Chloride	Total cholesterol
Creatinine	Total protein
Gamma glutamyl transferase	Triglycerides
Globulin (calculated)	Urea nitrogen
Standard urinalysis parameters evaluated	
Bilirubin	pH
Color and appearance	Protein
Glucose	Specific gravity
Ketones	Total volume
Microscopy of spun deposit	Urobilinogen
Occult blood	

Table 13: Standard microscopic tissue list

Standard microscopic pathology for dogs	
Adrenal gland	Pancreases
Aorta	Pituitary
Bone with bone marrow, femur	Parathyroid gland
Bone with bone marrow, sternum	Prostate
Bone with bone marrow, rib	Salivary gland, mandibular
Bone marrow smear	Salivary gland, parotid
Brain	Salivary gland, sublingual
Epididymis	Skeletal muscle, biceps femoris
Esophagus	Skin
Eye (with optic nerve)	Small intestine, duodenum
Gallbladder	Small intestine, ileum with Peyer's patches
Heart	Small intestine, jejunum
Joint, tibiofemoral	Spinal cord, cervical
Kidney	Spinal cord, lumbar
Large intestine, cecum	Spinal cord, thoracic
Large intestine, colon	Spleen
Large intestine, rectum	Stomach, cardia
Larynx	Stomach, fundus
Liver	Stomach, pylorus
Lung	Testis
Lymph node (mandibular)	Thymus
Lymph node (mesenteric)	Thyroid gland
Lymph node (tracheobronchial)	Tongue
Lymph node (regional, if applicable)	Trachea
Mammary gland (process females only)	Urinary bladder
Nerve (sciatic)	Uterus with cervix
Nictitans gland	Vagina
Ovary	Gross lesions
Oviducts	Tissue masses

Example 8. Reversal of diazepam-induced sedative-hypnotic effects in rats

- 5 The effect of flumazenil nicotinamide formulation A (Table 15) on reversal of diazepam induced sedative- hypnotic state was studied in SD rat model. The study end point was set by determining the sleeping time of the rats (females, 8-9 weeks of age, n=8) in each group. The experiment was conducted in two groups, of four rats each:

Table 14. Treatment groups

Group No.	N=	Treatment
1F	4	Flumazenil Formulation A
2F	4	Placebo (vehicle)

Following acclimation (5 days) inductive of sedative-hypnotic effects was induced by an intravenous injection of 20 mg/kg of benzodiazepine diazepam (Assival®) using a 24G

sized needle over approximately 15 seconds. In case an animal was not sufficiently sedated an additional injection of 5 mg/kg was administrated. The day of induction is defined as "Day 1".

The test item (Formulation A or placebo) was administered via the sublingual route twice at an interval of approximately 2 minutes once the animal had no righting reflex about 5 minutes post the second injection of the diazepam.

The components of the placebo solution are listed in Table 15.

Table 15. Placebo and flumazenil solutions for sublingual administration in rats.

Component	Formulation A	Placebo
	% w/w	
Flumazenil	1.1	-
Ethanol	40	10
Propylene glycol	10	20
30% HPCD in citric buffer		59.3
10mM pH 4.0		
Menthol in EtOH (1/1 w/w)	-	0.1
Propylparaben/Methylparaben in EtOH 0.02/0.18/10 w/w/w	-	10.2
Citric acid anyhdrous	0.05	-
Sodium citrate dehydrate	0.05	-
Nicotinamide	1.5	-
L-menthol	0.1	-
Water	47	-

10 Sleeping Time Determination was carried out as follows: upon diazepam injection the animal was placed on its back in a bedded standard cage. Sleeping time was determined when the test animal was no longer capable to maintain its Righting Reflex. This reflex is defined as the animal's failure of righting 2 times within 30 seconds. Clock time was recorded after the first administration of the test item and again when the animal woke up. The sleeping
15 time was determined as the elapsed time between first administration of the tested item and the time that the animal regained its Righting Reflex. The entire study was recorded (VCR).

Observations for signs of morbidity and mortality were performed twice a day. One animal died immediately after diazepam dosing. Individual body weight determination was made on Day 1 prior to induction (see Table 16 and Fig. 5). At study termination all animals
20 were euthanized by respiratory exposure to excess CO₂.

Table 16. Body weights

Group	Animal No.	Body weight (g)	Group	Animal No.	Body weight (g)
1F	1	183	2F	5	184
	2	187		6	175
				7	181
	4	178		8	182
Average $\pm$ sd		182 $\pm$ 3.3	Average $\pm$ sd		180.5 $\pm$ 3.3

As shown in Table 17, sleeping time following drug treatment was clearly shortened. The large variation is due to the small number of animals in the two groups.

5

Table 17. Sleeping time

Group	Animal No.	Time (min)	Group	Animal No.	Time (min)
1F	1	3.18	2F	5	40.46
	2	5.24		6	10.1
	-	-		7	3.5
	4	3.45		8	died
Average $\pm$ sd		3.96 $\pm$ 0.91	Average $\pm$ sd		18.02 $\pm$ 16.02

Example 9. Clinical study protocols**1. PK protocol**

10 The primary objective is to determine the single dose absolute bioavailability of sublingual (SL) Formulation A using the marketed IV flumazenil formulation as the comparator. The secondary objectives are to characterize the concentration time course of two dose levels of SL Formulation A to support dose selection for Phase 2 and 3 studies and to evaluate the safety and tolerability of flumazenil formulations.

15 Study design includes masking and enrolment, open label, randomized, three-way crossover study, with 15 healthy subjects $\geq$ 18 years of age, with no medication, based on the following study arms:

1. SL Formulation A 100 μ l (1.1 mg) N=5
2. SL Formulation A 200 μ l (2.2 mg) N=5
- 20 3. IV Flumazenil (Romazicon®) (0.2 mg) N=5

Study conditions: standardized as much as possible in terms of diet and fluid intake. This type of study is interventional. The secondary outcome measure is to examine the safety and tolerability of Formulation A (1.1mg and 2.2mg), where the safety endpoints include: adverse event monitoring, vital signs, physical examinations, clinical laboratory tests

and ECGs.

The time-points for blood and serum tests of flumazenil concentration are: 0, 10min, 30min, 60min, 90min, 2h, 4h, 6h and 24h. PK parameters for analysis are C_{max}, T_{max}, C_{min}, T_{min}, AUC_{0-∞}, AUC_{0-t} and T_{1/2}.

5

2. Phase 2B study: Dose range study of SL flumazenil

The study is a Phase 2B Dose Range, Efficacy and Safety of sublingual Formulation A for the indication: reversal of the residual effects of GABA related (benzodiazepine (BNZ) and nonBNZ) hypnotic drugs in insomniac patients. The residual sedative effect relates to benzodiazepine (BNZ) and nonBNZ hypnotic drugs. The study evaluates different doses of Formulation A in comparison to a placebo. Formulation A is provided as a sublingual (SL) spray formulation with flumazenil 11mg/ml.

Short-term safety and tolerability data is monitored along with psychomotor /cognitive and behavioral effects. The rationale of this study is to support study the pivotal efficacy phase 3 study, for the aforementioned indication.

Doses are determined with reference to the bioavailability study results. The study includes two groups of insomniac patients (males and females) aged ≥18y, Group A (n=60) and Group B (n=60). Each group is divided into Dose 1 (n=20), Dose 2 (n=20) and Placebo (n=20). The subjects all groups (n=20) are tested for the safety and efficacy of SL spray administration of Formulation A in the reversal of the sedative residual effect of BNZ (Group A) or Non BNZ (Group B).

Response to treatment is evaluated for each treatment arm. Good Response (GR) is R*25%, namely, an improvement of 25% in efficacy parameters of Formulation A treated patients compared to placebo/baseline. Very Good Response to treatment (VGR) is R*35%.

The secondary objectives are to evaluate the duration of action of a single dose of SL CRLS003 and to evaluate the safety of a single daily dose of Formulation A for 7 days. Study Design is randomized, double blind, dose range and placebo controlled.

The following examinations are carried out:

- a) Physical examination, clinical laboratory, vital signs.
- b) Sleep induction by BNZ/nonBNZ (Study A/B, respectively), per patient routine.
- c) All patients undergo full polysomnography.
- 5 d) At 6 AM patients are awaked.
- e) Baseline - On awakening digit symbol substitution test (DSST) is performed.
- f) Patients are randomly treated with Formulation A 1.1 or 2.2 mg or with placebo.
- g) The following tests are conducted at 10 min, 60 min, and 120 min following SL
10 Formulation A or placebo administration: DSST, immediate Word Recall Test
(iWRT), Visual Analog Scale (VAS) and Profile of Mood States (POMS).
- h) Patients will be self-treated throughout the week at home using daily hypnotics and
Formulation A/placebo (blinded) on awakening. Patients will answer questions and
write sleep and performance logs.
- i) On the 7th night, patients will report to the clinic for full polysomnography and will
15 repeat the same evaluation as in the previous visit.
- j) One week after termination, subjects will visit for safety monitoring.

Subjects meet all of the following inclusion criteria:

- 20 i. Each subject understands and voluntarily signs an informed consent form prior to
any study-mandated procedure.
- ii. Male or female aged ≥ 18 y at screening. Women of childbearing potential must
have a negative pregnancy test at the screening visit and use a reliable method of
contraception during the entire study duration (e.g., contraceptive pill; intra-
uterine device; contraceptive injection (prolonged-release gestagen); subdermal
25 implantation; vaginal ring or transdermal patch).
- iii. Body mass index ≥ 18.5 and $< 32 \text{ kg/m}^2$.
- iv. Chronic insomniac who are regularly treated by BNZ (Group A) or nonBNZ
(Group B) for at least 2 months and report residual morning symptoms.
- v. Each subject is in good health as determined by a medical history, physical
30 examination and ECG.
- vi. Negative regarding any use of illicit drug, alcohol (ethanol), stimulants.

Exclusion criteria are:

1. Using medications, except hypnotics and contraceptives, 1 month prior to screening.
2. A change in the hypnotics within the last 2 months.
3. Any sleep associated complains other than insomnia.
- 5 4. History of Epilepsy and or anti-epileptic drugs.
5. Excessive caffeine consumption (≥ 500 mg per day).
6. Pregnancy or breast feeding.
7. Night shift workers within 1 month prior to the screening visit.
8. Clinically relevant ECG abnormalities.
- 10 9. History of alcohol or drug abuse within 3 years prior to the screening visit.
10. Cognitive Behavioral Therapy (CBT) started within 1 month prior to screening.
11. Known hypersensitivity to drugs of the same class as the study treatment, or any excipients of the drug formulation.
12. Treatment with another investigational drug within 1 month prior to the screening.
- 15 13. History of severe head injury.

The study includes the following visits:

Visit 1- Screening visit: A signed and dated informed consent form is collected from each participating subject, prior to any screening procedures. Screening visit takes place up to 14 days prior to treatment visit. Subjects who are screened are assigned a 3-digit Screening number. During the screening visit, the following procedures are performed: Review
20 Inclusion / Exclusion criteria; Medical history inquiry and demographic data; Record prior medication; Neurological assessment; Conduct physical examination including vital signs, height and weight measurements; Pregnancy test; 12-lead electrocardiography.

Visits 2 and 3 - Treatment at sleep laboratory visits: prior to Formulation A
25 administration subjects undergo full physical examination, including sublingual and oral examination; vital signs measurements; adverse events and concomitant medication inquiry; Blood baseline tests. After determining that a subject is eligible, subjects receives a Randomization number and progresses to the sleep induction stage. At this stage the subject receives a sleep / hypnotic drug as follows:

30 Study A: BNZ (same drug and dose that patients use regularly)

Study B: nonBNZ (same drug and dose that patients use regularly)

Subjects sleep under full polysomnography until 6 AM. On awakening DSST is performed and the study Drug is administered: (At time 0) 0.1/0.2 ml Formulation A or

placebo (randomly and blindly). Thereafter, at 10 min., 60 min., and 120 min. post Formulation A administration, subjects are tested for psychomotor performance and behavioral tasks and vital signs. Safety measurement are conducted after performance tasks completion, as follows, full physical examination, including the sublingual and the oral cavity testing; neurological assessment; vital signs measurements; AE inquiry and blood tests.

The participants are released by a physician and are not allowed to drive until 10 hours after hypnotic administration. On the 7th day after study visit 2, visit 3 is conducted applying same procedures as in visit plan 2.

Across the study at home, patients continue using hypnotics and, on awakening taking Formulation A. Patients also keep a daily sleep-wake-performance log. The following prohibitions apply to the visits: caffeine consumption after 2:00 pm on the day before any study visit; consumption of alcohol or grapefruit (including as juice) from the day prior to each of the study visits, and for 3 consecutive days; and napping on the day before any study visit. During study any diet or non-pharmacological activity are allowed if started at least 1 month prior to the screening visit and stable until 24 hours after the last administration of study treatment.

Follow-up visit- Subjects pay a safety visit at day 7 after treatment termination and are examined, including blood tests, and report adverse effects (AEs).

Unscheduled visit - Subjects are monitored for AEs throughout the study. For any reported AEs defined as “Moderate or “Severe”, the subject is instructed to visit the clinic for safety measures as required. Causal AEs is monitored until resolved.

Study Duration: subjects participate in this study for 1 week of therapy and 1 week of follow-up. Interim analysis is carried (after 30 patients). Standard statistical methods are used to analyze the results (e.g. mean and standard deviation based on student-t test and ANOVA), where $p < 0.05$ defines significance.

3. Phase 3 pivotal efficacy study

This study is a double blind evaluation of SL Formulation A for reversal of next day residual sedative effect from hypnotic drugs used to treat insomnia. It is directed to evaluate the safety and effectiveness of SL Formulation A in insomniac patients in reversing the next day residual effect of hypnotic drugs. The study is designed to collect short-term safety and tolerability data along with psychomotor/cognitive and behavioral function. It is an interventional type of study. The indication, drugs and treatment are similar to the those

tested in the Phase 2B study.

Sample includes 300 subjects (150 per treatment group) randomized to flumazenil or placebo 1:1 ratio, stratified by BNZ or nonBNZ. Treatment response is evaluated for each treatment arm. Good Response (GR) is $R \geq 25\%$. Secondary objective include safety of SL Formulation A compared to placebo administered daily for 6 weeks. The Study Design is randomized, double blind, placebo and controlled.

All subjects are evaluated for the safety and efficacy of SL Formulation A to reverse the sedative effect of BNZ or nonBNZ applying the same routine used in Phase 2B study with the following exceptions:

- 10 a) Patients are self-treated for 3 weeks using daily hypnotics and Formulation A/placebo (blinded) on awakening.
- b) By the end of the 3rd week $\pm$ 3 nights and at the last night (6 weeks after trial start), patients visit the clinic for full polysomnography and repeat the procedures and testing described above.
- 15 c) Two weeks after the last treatment, the patient take safety monitoring and AE inquiry.

Subjects eligible for the study meet all of the aforementioned inclusion and exclusion criteria. The study includes the same type of visits listed above, with the addition of a fourth sleep laboratory visit.

- 20 4. Phase 3 double blind placebo controlled trial for the evaluation of Formulation A in the treatment of Episodic Hepatic Encephalopathy (HE) grade ≥ 2 .

The purpose of this study is to evaluate mental score/quality of life improvement in patients with episodic HE following the administration of sublingual flumazenil (Formulation A). This is an interventional type of study. The indication is: relief of cognitive impairment symptoms in patients with episodic HE. The investigational drug is Formulation A sublingual spray comprising 1.12mg /ml Flumazenil. The time to recovery from episodic HE grade ≥ 2 is defined as a decrease of Conn score to Grade < 2 , or a decrease in Asterixis.

The secondary objectives of this study are as follows:

- a) Duration and number of HE-related hospitalization.
- b) Time to any increase from baseline in Conn score.
- c) Time to any increase from baseline in Asterixis grade
- 5 d) Number connection Test (NCT or Reitan Test)
- e) Continuous reaction times to sound (CRT)
- f) Block design test (BDT)
- g) Symbol digit modalities test (SDMT)
- h) Mean change from baseline in fatigue domain score on the Chronic Liver
- 10 Disease Questionnaire (CLDQ)

The study design is randomized, placebo controlled, double blinded administered sublingual flumazenil twice a day for 3 weeks. During study, patients with episodic HE grade ≥ 2 are screened for a respond to flumazenil therapy. Baseline assessment includes in addition to the routine testing described below: Clinical assessment (physical and blood tests), EEG,

15 plasma assays for benzodiazepines and NCT.

Physical and blood tests include :

- a. Ammonia measurements, weight, renal functions electrolytes each visit.
- b. EEG
- c. A questionnaire regarding diarrhea and abdominal discomfort
- 20 d. Cognitive impairment
- e. Performance memory test each visit (e.g. NCT and Reitan Test)
- f. Quality of life evaluation (HE related hospitalization). Conn score and asterixis increase are documented.

The study population includes male and female, cirrhotic patients aged ≥ 18 y, with

25 episodic HE grade ≥ 2 .

Inclusion criteria that subjects eligible for enrolment into the study meet are:

- a) Episodic HE grade ≥ 2 .
- b) Responsiveness to flumazenil therapy.
- c) Each subject understands and voluntarily signs an informed consent form prior to any
- 30 study-mandated procedure.
- d) Male or female aged ≥ 18 at screening.
- e) Women of childbearing potential must have a negative pregnancy test at the screening visit and use a reliable method of contraception during the entire study duration (e.g.

Contraceptive pill; Intra-uterine device; Contraceptive injection (prolonged-release gestagen); Subdermal implantation; Vaginal ring or Transdermal patch).

Exclusion criteria include chronic BNZ therapy and epilepsy.

- 5 Safety assessment is based on changes from screening/baseline for clinical AEs reported by the subject, or observed by the investigator assessed by physical exam, oral and sublingual assessment, neurological assessment and vital signs.

Standard statistical methods are used to analyze the results (e.g. mean and standard deviation based on student-t test and ANOVA), where $p < 0.05$ defines significance.

10

- 15 The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without undue experimentation and without departing from the generic concept, and, therefore, such adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology employed herein is for the purpose of description and not of limitation. The means, materials, and steps for carrying out various disclosed functions may take a variety of alternative forms without departing from the invention.

CLAIMS

1. A pharmaceutically acceptable complex of ethyl 8-fluoro-5,6-dihydro-5-methyl-6-oxo- 4H- imidazo [1,5-a] [1,4] benzodiazepine- 3-carboxylate or a salt thereof and a complexation agent or a derivative thereof, wherein the complexation agent is selected from the group consisting of: nicotinamide, sodium nicotinate, sodium benzoate, sodium hydroxybenzoates, sodium salicylate, sodium gentisate, gentisic acid ethanolamide, sodium toluates, sodium aminobenzoate, sodium anthranilate, sodium butylmonoglycolsulfate and resorcinol.
2. The pharmaceutically acceptable complex of claim 1, wherein the complexation agent is nicotinamide or a derivative thereof.
3. The pharmaceutically acceptable complex of claim 1, wherein the complexation agent:flumazenil ratio is in the range of 1:1 to 2:1.
4. The pharmaceutically acceptable complex of claim 1, which is a flumazenil nicotinamide complex.
5. A pharmaceutical composition comprising the flumazenil complex of any one of claims 1 to 4 and a pharmaceutical acceptable carrier.
6. The pharmaceutical composition according to claim 5, wherein the composition is in a form selected from the group consisting of: a pill, tablet, lozenge, coated tablet, granule, capsule, hard or soft gelatin capsule, aqueous solution, alcoholic solution, oily solution, syrup, emulsion suspension pastille suppository, solution for injection or infusion, ointment, tincture, cream, lotion, powder, spray, transdermal therapeutic systems, nasal spray, aerosol mixture, microcapsule, implant, rod and plaster.
7. The pharmaceutical composition according to claim 5, wherein the composition is in a form selected from the group consisting of: immediate release, delayed release, pulsatile release, continuous release and repetitive release.
8. The pharmaceutical composition according to claim 5, wherein the concentration of flumazenil is within the range of about 0.4 to 2 % w/w of the total weight of the pharmaceutical composition.
9. The pharmaceutical composition according to claim 8, wherein the concentration of

flumazenil is within the range of about 0.5 to 1.8 % w/w of the total weight of the pharmaceutical composition.

10. The pharmaceutical composition according to claim 5, further comprising a solubilizing agent selected from an alcohol, a glycol and a combination thereof.

11. The pharmaceutical composition according to claim 10, wherein the solubilizing agent comprises a combination of an alcohol and a glycol and wherein the alcohol:glycol ratio is at least 1.5:1.

12. The pharmaceutical composition according to claim 11, wherein the alcohol:glycol ratio is in the range of 1.5:1 to 5:1.

13. The pharmaceutical composition according to claim 10, wherein the solubilizing agent comprises a combination of an alcohol and a glycol and wherein the concentration of the solubilizing agent is at least 40% of the total weight of the pharmaceutical composition.

14. The pharmaceutical composition according to claim 13, wherein the concentration of the solubilizing agent is in the range of 40% to 60% of the total weight of the pharmaceutical composition.

15. The pharmaceutical composition according to claim 10, further comprising a buffering agent.

16. The pharmaceutical composition according to claim 10, further comprising at least one agent selected from the group consisting of: a penetration enhancer, a surfactant and cyclodextrin.

17. The pharmaceutical composition according to claim 10, further comprising a preservative selected from the group consisting of: benzyl alcohol, propylparaben, methylparaben and combinations thereof.

18. The pharmaceutical composition according to claim 17, wherein the penetration enhancer is menthol.

19. The pharmaceutical composition according to claim 15, wherein the buffering agent is selected from the group consisting of: citric buffer, sodium chloride and combination thereof.

20. The pharmaceutical composition according to claim 16, wherein the surfactant is benzalkonium chloride.

1/3




Figure 1




Figure 2

2/3




Figure 3




Figure 4

3/3




Figure 5




Figure 6