



US010875825B2

(12) **United States Patent**  
**Kim et al.**

(10) **Patent No.:** **US 10,875,825 B2**  
(45) **Date of Patent:** **Dec. 29, 2020**

(54) **AMINE-BASED COMPOUND AND ORGANIC LIGHT-EMITTING DEVICE INCLUDING THE SAME**

(71) Applicant: **SAMSUNG DISPLAY CO. LTD.**,  
Yongin-si (KR)

(72) Inventors: **Jongwoo Kim**, Yongin-si (KR);  
**Jangyeol Baek**, Yongin-si (KR);  
**Eunjae Jeong**, Yongin-si (KR);  
**Sanghyun Han**, Yongin-si (KR);  
**Youngkook Kim**, Yongin-si (KR);  
**Seokhwan Hwang**, Yongin-si (KR)

(73) Assignee: **Samsung Display Co., Ltd.**, Yongin-si (KR)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 226 days.

(21) Appl. No.: **15/870,782**

(22) Filed: **Jan. 12, 2018**

(65) **Prior Publication Data**  
US 2019/0055187 A1 Feb. 21, 2019

(30) **Foreign Application Priority Data**  
Aug. 18, 2017 (KR) ..... 10-2017-0105081

(51) **Int. Cl.**  
**H01L 51/50** (2006.01)  
**C07C 211/61** (2006.01)  
(Continued)

(52) **U.S. Cl.**  
CPC ..... **C07C 211/61** (2013.01); **C07D 307/91** (2013.01); **C09K 11/06** (2013.01);  
(Continued)

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,465,115 B2 10/2002 Shi et al.  
6,596,415 B2 7/2003 Shi et al.  
(Continued)

FOREIGN PATENT DOCUMENTS

EP 3 524 661 A1 8/2019  
JP 2008-050337 A 3/2008  
(Continued)

OTHER PUBLICATIONS

EPO Extended Search Report dated Jul. 6, 2018, for corresponding European Patent Application No. 18173452.6 (8 pages).


(Continued)

*Primary Examiner* — Gregory D Clark  
(74) *Attorney, Agent, or Firm* — Lewis Roca Rothgerber Christie LLP

(57) **ABSTRACT**

An amine-based compound represented by Formula 1, in which one of R<sub>1</sub> to R<sub>10</sub> is a group represented by Formula 2, and an organic light-emitting device including the same are provided:

Formula 1




(Continued)

**10**

<b>190</b>
<b>150</b>
<b>110</b>

-continued



Formula 2

2005/0208331	A1	9/2005	Maeda
2007/0018569	A1	1/2007	Kawamura et al.
2008/0160347	A1	7/2008	Wang et al.
2013/0334518	A1	12/2013	Park et al.
2014/0131681	A1	5/2014	Ito et al.
2015/0287921	A1	10/2015	Kato et al.
2015/0303379	A1	10/2015	Lee et al.
2015/0329772	A1	11/2015	Heil et al.
2016/0155943	A1	6/2016	Sasaki et al.
2017/0317289	A1*	11/2017	Lee ..... C09K 11/06
2017/0317290	A1	11/2017	Lee et al.

When the amine-based compound is included in an organic light-emitting device as a hole transport material, the device may exhibit excellent I-V-L characteristics, including a low driving voltage, improved efficiency, and improved lifespan.

**18 Claims, 4 Drawing Sheets**

- (51) **Int. Cl.**  
*C07D 307/91* (2006.01)  
*H01L 51/00* (2006.01)  
*C09K 11/06* (2006.01)
- (52) **U.S. Cl.**  
 CPC ..... *H01L 51/006* (2013.01); *H01L 51/0058* (2013.01); *H01L 51/0073* (2013.01); *H01L 51/506* (2013.01); *H01L 51/5012* (2013.01); *C07C 2603/18* (2017.05); *C07C 2603/40* (2017.05); *C09K 2211/1011* (2013.01); *C09K 2211/1088* (2013.01); *H01L 51/5056* (2013.01)

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,849,348	B2	2/2005	Zheng et al.
8,980,442	B2	3/2015	Yabunouchi et al.
8,993,805	B2	3/2015	Nishiyama et al.
9,099,654	B2	8/2015	Wang et al.
9,893,293	B2	2/2018	Sasaki et al.

FOREIGN PATENT DOCUMENTS

JP	2015-119166	A	6/2015
KR	10-2005-0089993	A	9/2005
KR	10-2008-0031808	A	4/2008
KR	10-2008-0108115	A	12/2008
KR	10-1053466	B1	7/2011
KR	10-2011-0117716	A	10/2011
KR	10-2013-0123484	A	11/2013
KR	10-2014-0078096	A	6/2014
KR	10-2014-0119731	A	10/2014
KR	10-2014-0128878	A	11/2014
KR	10-2014-0128879	A	11/2014
KR	10-2015-0007476	A	1/2015
KR	10-2015-0066202	A	6/2015
KR	10-1530886	B1	6/2015
KR	10-2015-0074015		7/2015
KR	10-2016-0053561	A	5/2016
KR	10-2016-0067021	A	6/2016
KR	10-2018-0037717	A	4/2018
WO	WO 2014/065391	A1	5/2014
WO	WO 2014/088285	A1	6/2014
WO	WO 2014/106522	A1	7/2014
WO	WO 2015-084114	A1	6/2015
WO	WO 2016/064111	A1	4/2016
WO	WO-2016-129861	A1	8/2016

OTHER PUBLICATIONS

U.S. Office Action dated Apr. 29, 2020, issued in U.S. Appl. No. 15/869,988 (9 pages).

\* cited by examiner

**FIG. 1**

**10**

<b>190</b>
<b>150</b>
<b>110</b>

**FIG. 2**

**20**

<b>190</b>
<b>150</b>
<b>110</b>
<b>210</b>

**FIG. 3**

**30**

<b>220</b>
<b>190</b>
<b>150</b>
<b>110</b>

**FIG. 4**

**40**

<b>220</b>
<b>190</b>
<b>150</b>
<b>110</b>
<b>210</b>

1

AMINE-BASED COMPOUND AND ORGANIC  
LIGHT-EMITTING DEVICE INCLUDING  
THE SAME

CROSS-REFERENCE TO RELATED  
APPLICATION

This application claims priority to and the benefit of Korean Patent Application No. 10-2017-0105081, filed on Aug. 18, 2017, in the Korean Intellectual Property Office, the entire content of which is incorporated herein by reference.

BACKGROUND

1. Field

Aspects of one or more embodiments of the present disclosure relate to an amine-based compound and an organic light-emitting device including the same.

2. Description of the Related Art

Organic light-emitting devices are self-emission devices that have wide viewing angles, high contrast ratios, short response times, and excellent characteristics in terms of brightness, driving voltage, and response speed, compared to devices in the art.


An organic light-emitting device may include a first electrode on a substrate, and a hole transport region, an emission layer, an electron transport region, and a second electrode sequentially positioned on the first electrode. Holes provided from the first electrode may move toward the emission layer through the hole transport region, and electrons provided from the second electrode may move toward the emission layer through the electron transport region. The carriers (e.g., holes and electrons) may recombine in the emission layer to produce excitons. Then, the excitons may transition from an excited state to a ground state, thereby releasing energy in the form of light.

SUMMARY

One or more aspects of embodiments of the present disclosure provide an amine-based compound and an organic light-emitting device including the same.

Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments.

One or more aspects of embodiments of the present disclosure provides an amine-based compound represented by Formula 1:




Formula 1

2

-continued

Formula 2



In Formulae 1 and 2,

Ar<sub>1</sub>, Ar<sub>2</sub>, L<sub>1</sub>, and L<sub>2</sub> may each independently be a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

a<sub>1</sub> and a<sub>2</sub> may each independently be an integer from 0 to 5, wherein, when a<sub>1</sub> is two or more, two or more L<sub>1</sub>(s) may be identical to or different from each other, and when a<sub>2</sub> is two or more, two or more L<sub>2</sub>(s) may be identical to or different from each other,

when a<sub>1</sub> is zero, \*-(L<sub>1</sub>)<sub>a1</sub>-\* may be a single bond, and when a<sub>2</sub> is zero, \*-(L<sub>2</sub>)<sub>a2</sub>-\* may be a single bond,

R<sub>1</sub> to R<sub>10</sub> may each independently be selected from a group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

provided that one of R<sub>1</sub> to R<sub>10</sub> is a group represented by Formula 2,

R<sub>11</sub> and R<sub>12</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

b<sub>1</sub> and b<sub>2</sub> may each independently be an integer from 1 to 10, wherein, when b<sub>1</sub> is two or more, two or more R<sub>11</sub>(s)

may be identical to or different from each other, and when b2 is two or more, two or more R<sub>12</sub>(s) may be identical to or different from each other,

at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), -B(Q<sub>11</sub>)(Q<sub>12</sub>), -C(=O)(Q<sub>11</sub>), -S(=O)<sub>2</sub>(Q<sub>11</sub>), and -P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), -B(Q<sub>21</sub>)(Q<sub>22</sub>), -C(=O)(Q<sub>21</sub>), -S(=O)<sub>2</sub>(Q<sub>21</sub>), and -P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and -Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), -B(Q<sub>31</sub>)(Q<sub>32</sub>), -C(=O)(Q<sub>31</sub>), -S(=O)<sub>2</sub>(Q<sub>31</sub>), and -P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>).

Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono

group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

\* and \*' each indicate—a binding site to a neighboring atom.

Another aspect provides an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, wherein the organic layer includes an emission layer and at least one of the amine-based compound.

#### BRIEF DESCRIPTION OF THE DRAWINGS

These and/or other aspects will become apparent and more readily appreciated from the following description of the example embodiments, taken in conjunction with the accompanying drawings, in which:

FIG. 1 is a schematic view of an organic light-emitting device according to embodiments of the present disclosure;

FIG. 2 is a schematic view of an organic light-emitting device according to embodiments of the present disclosure;

FIG. 3 is a schematic view of an organic light-emitting device according to embodiments of the present disclosure; and

FIG. 4 is a schematic view of an organic light-emitting device according to embodiments of the present disclosure.

#### DETAILED DESCRIPTION

The present disclosure will now be described more fully with reference to example embodiments. Advantages, features, and how to achieve such advantages and features of the present invention will become apparent by reference to the embodiments described in detail herein, together with the accompanying drawings. This invention may, however, be embodied in many different forms and should not be construed as being limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the concept of the disclosure to those skilled in the art.

Hereinafter, embodiments are described in more detail by referring to the attached drawings, in which like reference numerals denote like elements and duplicative descriptions thereof may not be provided.

As used herein, the singular forms “a”, “an”, and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.


It will be further understood that the terms “comprises” and/or “comprising” may be used herein to specify the presence of stated features or components, but do not preclude the presence or addition of one or more other features or components.

It will be understood that when a layer, region, or component is referred to as being “on” or “onto” another layer, region, or component, it may be directly or indirectly formed on the other layer, region, or component. That is, for example, intervening layers, regions, or components may be present. In contrast, when an element is referred to as being “directly on” another element, no intervening elements are present.


5

The sizes and thicknesses of components in the drawings may be exaggerated for convenience of explanation. In other words, since sizes and thicknesses of components in the drawings are arbitrarily illustrated for convenience of explanation, the following embodiments of the present disclosure are not limited thereto.

An amine-based compound according to an embodiment of the present disclosure is represented by Formula 1 below:



Formula 1



Formula 2

$Ar_1$ ,  $Ar_2$ ,  $L_1$ , and  $L_2$  in Formulae 1 and 2 may each independently be a substituted or unsubstituted  $C_5$ - $C_{60}$  carbocyclic group or a substituted or unsubstituted  $C_2$ - $C_{60}$  heterocyclic group.

In one or more embodiments,  $Ar_1$ ,  $Ar_2$ ,  $L_1$ , and  $L_2$  may each independently be a substituted or unsubstituted  $C_5$ - $C_{30}$  carbocyclic group or a substituted or unsubstituted  $C_2$ - $C_{30}$  heterocyclic group.

In one or more embodiments,  $Ar_1$ ,  $Ar_2$ ,  $L_1$ , and  $L_2$  may each independently be selected from:

a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, a heptalene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-fluorene group, a benzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluo-  
 50 ranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, a hexacene group, a pyrrole group, an imidazole group, a pyrazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an iso-indole group, an indole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a qui-  
 55 nazoline group, a cinnoline group, a carbazole group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzoxazole group, a benzimidazole group, a furan group, a benzofuran group, a thio-  
 60 phene group, a benzothiophene group, a thiazole group, an isothiazole group, a benzothiazole group, an isoxazole group, an oxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a benzoxazole group, a dibenzofuran group, a dibenzothiophene group, a benzo-  
 carbazole group, and a dibenzocarbazole group; and

a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, a heptalene group, an indacene group, an acenaphthalene group, a fluorene group,

6


a spiro-fluorene group, a benzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluo-  
 5 ranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, a hexacene group, a pyrrole group, an imidazole group, a pyrazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an iso-indole group, an indole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a qui-  
 10 nazoline group, a cinnoline group, a carbazole group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzoxazole group, a benzimidazole group, a furan group, a benzofuran group, a thio-  
 15 phene group, a benzothiophene group, a thiazole group, an isothiazole group, a benzothiazole group, an isoxazole group, an oxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a benzoxazole group, a dibenzofuran group, a dibenzothiophene group, a benzo-  
 20 carbazole group, and a dibenzocarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a  $C_6$ - $C_{20}$  aryl group, a  $C_2$ - $C_{20}$  heteroaryl group, and —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ),

wherein  $Q_{31}$  to  $Q_{33}$  may each independently be selected  
 30 from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{60}$  alkyl group, a  $C_2$ - $C_{60}$  alkenyl group, a  $C_2$ - $C_{60}$  alkynyl group, a  $C_1$ - $C_{60}$  alkoxy group, a  $C_3$ - $C_{10}$  cycloalkyl group, a  $C_1$ - $C_{10}$  hetero-  
 cycloalkyl group, a  $C_3$ - $C_{10}$  cycloalkenyl group, a  $C_1$ - $C_{10}$  heterocycloalkenyl group, a  $C_6$ - $C_{60}$  aryl group, a  $C_1$ - $C_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl  
 40 group,


but embodiments of the present disclosure are not limited thereto.

In one or more embodiments,  $L_1$  and  $L_2$  may each independently be selected from a group represented by any of Formulae 3-1 to 3-46, but embodiments of the present disclosure are not limited thereto:


Formula 3-1



Formula 3-2











Formula 3-3






**9**  
-continued




**10**  
-continued

Formula 3-24




Formula 3-32

Formula 3-25




Formula 3-33

Formula 3-26




Formula 3-34

Formula 3-27




Formula 3-35

Formula 3-28




Formula 3-36

Formula 3-29




Formula 3-37

Formula 3-30



Formula 3-36

Formula 3-31



Formula 3-37

45

50






55

60

65

11


-continued




12

-continued


Formula 3-38



Formula 3-39



Formula 3-40



In Formulae 3-1 to 3-46,

Y<sub>1</sub> may be O, S, C(Z<sub>3</sub>)(Z<sub>4</sub>), N(Z<sub>5</sub>), or Si(Z<sub>6</sub>)(Z<sub>7</sub>),

Z<sub>1</sub> to Z<sub>7</sub> may each independently be selected from hydro-  
 40 gen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a  
 cyano group, a nitro group, an amidino group, a hydrazino  
 group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub>  
 alkoxy group, a cyclopentyl group, a cyclohexyl group, a  
 45 cycloheptyl group, a cyclopentenyl group, a cyclohexenyl  
 group, a phenyl group, a biphenyl group, a naphthyl group,  
 a fluorenyl group, a spiro-bifluorenyl group, a phenalenyl  
 group, a phenanthrenyl group, an anthracenyl group, a  
 fluoranthenyl group, a triphenylenyl group, a pyridinyl  
 50 group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl  
 group, an isoquinolinyl group, a benzoquinolinyl group, a  
 naphthyridinyl group, a quinoxalinyl group, a quinazolinyl  
 group, a carbazolyl group, a phenanthridinyl group, an  
 acridinyl group, a phenanthrolinyl group, a phenazinyl  
 group, a triazinyl group, a dibenzofuran group, a diben-  
 55 zothiofenyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>).

Formula 3-42

d<sub>2</sub> may be an integer from 0 to 2, wherein, when d<sub>2</sub> is two  
 or more, two or more Z<sub>1</sub>(s) may be identical to or different  
 from each other,

d<sub>3</sub> may be an integer from 0 to 3, wherein, when d<sub>3</sub> is two  
 60 or more, two or more Z<sub>1</sub>(s) may be identical to or different  
 from each other and two or more Z<sub>2</sub>(s) may be identical  
 to or different from each other,

d<sub>4</sub> may be an integer from 0 to 4, wherein, when d<sub>4</sub> is two  
 or more, two or more Z<sub>1</sub>(s) may be identical to or different  
 65 from each other,

d<sub>5</sub> may be an integer from 0 to 5, wherein, when d<sub>5</sub> is two  
 or more, two or more Z<sub>1</sub>(s) may be identical to or different

Formula 3-43

Formula 3-44

Formula 3-45

Formula 3-46

## 13

from each other and two or more and  $Z_2(s)$  may be identical to or different from each other,

$d_6$  may be an integer from 0 to 6, wherein, when  $d_6$  is two or more, two or more  $Z_1(s)$  may be identical to or different from each other and two or more and  $Z_2(s)$  may be identical to or different from each other,

$d_8$  may be an integer from 0 to 8, wherein, when  $d_8$  is two or more, two or more  $Z_1(s)$  may be identical to or different from each other, and

\* and \*' each indicate a binding site to a neighboring atom.

$a_1$  and  $a_2$  in Formula 2 may each independently be an integer selected from 0 to 5.  $a_1$  indicates the number of  $L_1(s)$ , wherein, when  $a_1$  is two or more, two or more  $L_1(s)$  may be identical to or different from each other.  $a_2$  indicates the number of  $L_2(s)$ , wherein, when  $a_2$  is two or more, two or more  $L_2(s)$  may be identical to or different from each other.

When  $a_1$  is zero,  $*(L_1)_{a_1}-*'$  may be a single bond, and when  $a_2$  is zero,  $*(L_2)_{a_2}-*'$  may be a single bond.


In one or more embodiments,  $a_1$  and  $a_2$  may each independently be 0 or 1, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments,  $Ar_1$  and  $Ar_2$  may each independently be selected from:


a benzene group; and

a benzene group substituted with at least one selected from deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1-C_{20}$  alkyl group, a  $C_1-C_{20}$  alkoxy group, a  $C_6-C_{20}$  aryl group, a  $C_2-C_{20}$  heteroaryl group, and  $-Si(Q_{31})(Q_{32})(Q_{33})$ , but embodiments of the present disclosure are not limited thereto.


In one or more embodiments,  $Ar_1$  and  $Ar_2$  may each independently be selected from a group represented by any of Formulae 5-1 to 5-80, but embodiments of the present disclosure are not limited thereto:




Formula 5-1



Formula 5-2




Formula 5-3




Formula 5-4

## 14


-continued




Formula 5-5




Formula 5-6




Formula 5-7




Formula 5-8



Formula 5-9











Formula 5-10



Formula 5-11

15

-continued




16

-continued


Formula 5-12

5




Formula 5-13

10




Formula 5-14

15




Formula 5-15

20




Formula 5-16

25




Formula 5-17

30




Formula 5-18

35




Formula 5-19


40



45




50




Formula 5-20

55



60



65

Formula 5-20

Formula 5-21

Formula 5-22

Formula 5-23

Formula 5-24

Formula 5-25

Formula 5-26

Formula 5-27











Formula 5-28

Formula 5-29

Formula 5-30

17

-continued




18


-continued

Formula 5-31


5



Formula 5-32 10




15




Formula 5-33

20




Formula 5-34 25




30

Formula 5-35




35

Formula 5-36




40

Formula 5-37 45




Formula 5-38 50




55

Formula 5-39




60

Formula 5-40



65



Formula 5-41

Formula 5-42

Formula 5-43

Formula 5-44

Formula 5-45

Formula 5-46

Formula 5-47

Formula 5-48











Formula 5-49

Formula 5-50

Formula 5-51

Formula 5-52


**19**  
-continued



**20**  
-continued


Formula 5-53

5




Formula 5-54

10




Formula 5-55

15



Formula 5-56

20




Formula 5-57

25


Formula 5-58

35




Formula 5-59

40




Formula 5-60

45




Formula 5-61

55



Formula 5-62

60



65

Formula 5-63

Formula 5-64

Formula 5-65

Formula 5-66

Formula 5-67

Formula 5-68










Formula 5-69

Formula 5-70

Formula 5-71

21

-continued



22

In Formulae 5-1 to 5-80,

Formula 5-72 Y<sub>31</sub> may be O, S, C(Z<sub>33</sub>)(Z<sub>34</sub>), N(Z<sub>35</sub>), or Si(Z<sub>36</sub>)(Z<sub>37</sub>),

Z<sub>31</sub> to Z<sub>37</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranly group, a dibenzothiophenyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>),

Formula 5-73 e2 may be an integer from 0 to 2, wherein, when e2 is two or more, two or more Z<sub>32</sub>(s) may be identical to or different from each other,

Formula 5-74 e3 may be an integer from 0 to 3, wherein, when e3 is two or more, two or more Z<sub>31</sub>(s) may be identical to or different from each other and two or more Z<sub>32</sub>(s) may be identical to or different from each other,

Formula 5-75 e4 may be an integer from 0 to 4, wherein, when e4 is two or more, two or more Z<sub>31</sub>(s) may be identical to or different from each other and two or more Z<sub>32</sub>(s) may be identical to or different from each other,

Formula 5-76 e5 may be an integer from 0 to 5, wherein, when e5 is two or more, two or more Z<sub>31</sub>(s) may be identical to or different from each other and two or more Z<sub>32</sub>(s) may be identical to or different from each other,

Formula 5-77 e6 may be an integer from 0 to 6, wherein, when e6 is two or more, two or more Z<sub>31</sub>(s) may be identical to or different from each other and two or more Z<sub>32</sub>(s) may be identical to or different from each other,

Formula 5-78 e7 may be an integer from 0 to 7, wherein, when e7 is two or more, two or more Z<sub>31</sub>(s) may be identical to or different from each other,

Formula 5-79 e9 may be an integer from 0 to 9, wherein, when e9 is two or more, two or more Z<sub>31</sub>(s) may be identical to or different from each other, and

\* indicates a binding site to a neighboring atom.

R<sub>1</sub> to R<sub>10</sub> in Formula 1 may each independently be selected from a group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

Formula 5-72

Formula 5-73

Formula 5-74

Formula 5-75

Formula 5-76

Formula 5-77

Formula 5-78

Formula 5-79

Formula 5-80

provided that one selected from  $R_1$  to  $R_{10}$  is a group represented by Formula 2, and

$R_{11}$  and  $R_{12}$  may each independently be selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2$ - $C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-Si(Q_1)(Q_2)(Q_3)$ ,  $-B(Q_1)(Q_2)$ ,  $-C(=O)(Q_1)$ ,  $-P(=O)(Q_1)(Q_2)$ ,  $-P(=S)(Q_1)(Q_2)$ ,  $-S(=O)(Q_1)(Q_2)$ , and  $-S(=O)_2(Q_1)(Q_2)$ ,

wherein  $Q_1$  to  $Q_3$  may each independently be the same as described herein in connection with  $Q_{31}$  to  $Q_{33}$ .


In one or more embodiments,  $R_1$  to  $R_{10}$  may each independently be selected from a group represented by Formula 2, hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and  $-Si(Q_1)(Q_2)(Q_3)$ , and

$R_{11}$  and  $R_{12}$  may each independently be selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and  $-Si(Q_1)(Q_2)(Q_3)$ , but embodiments of the present disclosure are not limited thereto.


In one or more embodiments,  $R_1$  to  $R_{10}$  may each independently be selected from a group represented by Formula 2, hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a group represented by any of Formulae 5-1 to 5-80, and  $-Si(Q_1)(Q_2)(Q_3)$ , and

$R_{11}$  and  $R_{12}$  may each independently be selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a group represented by any of Formulae 5-1 to 5-80, and  $-Si(Q_1)(Q_2)(Q_3)$ , but embodiments of the present disclosure are not limited thereto.


In one or more embodiments,  $R_{11}$  and  $R_{12}$  may each independently be selected from a group represented by any of Formulae 6-1 to 6-34, but embodiments of the present disclosure are not limited thereto:




Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5




Formula 6-6




Formula 6-7



Formula 6-8










Formula 6-9



Formula 6-10


**25**  
-continued



**26**  
-continued


Formula 6-11

5




Formula 6-12 10

15




Formula 6-13

20




Formula 6-14

25




Formula 6-15

30




Formula 6-16

35




Formula 6-17


40



45




50



Formula 6-18

55



Formula 6-18

Formula 6-19

Formula 6-20

Formula 6-21

Formula 6-22

Formula 6-23


Formula 6-24

Formula 6-25

Formula 6-26

27

-continued



In Formulae 6-1 to 6-34,  
 Ph indicates a phenyl group, and  
 \* indicates a binding site to a neighboring atom.  
 b1 and b2 in Formula 2 may each independently be an integer selected from 1 to 10. b1 indicates the number of  $R_{11}(s)$ , wherein, when b1 is two or more, two or more  $R_{11}(s)$

28

Formula 6-27

may be identical to or different from each other. b2 indicates the number of  $R_{12}(s)$ , wherein, when b2 is two or more, two or more  $R_{12}(s)$  may be identical to or different from each other.

5

In one or more embodiments, the amine-based compound represented by Formula 1 may be represented by one selected from Formulae 1-1 to 1-6, but embodiments of the present disclosure are not limited thereto:


Formula 6-28

10

Formula 1-1

Formula 6-29

15



Formula 6-30


20

Formula 1-2

25

Formula 6-31

30



Formula 6-32


35

Formula 1-3

40

Formula 6-33

45




Formula 6-34

50

Formula 1-4

55




60


65

29

-continued



Formula 1-5



Formula 1-6

Ar<sub>1</sub>, Ar<sub>2</sub>, L<sub>1</sub>, L<sub>2</sub>, a<sub>1</sub>, a<sub>2</sub>, R<sub>1</sub> to R<sub>12</sub>, b<sub>1</sub>, and b<sub>2</sub> in Formulae 1-1 to 1-6 may each be the same as described in connection with Formula 1.

In one or more embodiments, R<sub>1</sub> to R<sub>10</sub> in Formulae 1-1 to 1-6 may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an iso-propyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, and a biphenyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments,

i) when the amine-based compound represented by Formula 1 is represented by Formula 1-1, R<sub>2</sub> to R<sub>10</sub> may each be hydrogen;

ii) when the amine-based compound represented by Formula 1 is represented by Formula 1-2, R<sub>8</sub> and R<sub>9</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a methyl group, an ethyl group, a propyl group, an iso-propyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, and a biphenyl group, and R<sub>1</sub>, R<sub>3</sub> to R<sub>7</sub>, and R<sub>10</sub> may each be hydrogen;

iii) when the amine-based compound represented by Formula 1 is represented by Formula 1-3, R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> to R<sub>10</sub> may each be hydrogen;

iv) when the amine-based compound represented by Formula 1 is represented by Formula 1-4, R<sub>1</sub> to R<sub>3</sub> and R<sub>5</sub> to R<sub>10</sub> may each be hydrogen;

v) when the amine-based compound represented by Formula 1 is represented by Formula 1-5, R<sub>1</sub> to R<sub>7</sub> and R<sub>9</sub> to R<sub>10</sub> may each be hydrogen; or

v) when the amine-based compound represented by Formula 1 is represented by Formula 1-6, R<sub>2</sub> may be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a methyl group, an ethyl group, a propyl group, an iso-propyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a

30

ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, and a biphenyl group, and R<sub>1</sub>, R<sub>3</sub> to R<sub>8</sub>, and R<sub>10</sub> may each be hydrogen, but embodiments of the present disclosure are not limited thereto.

5 In one or more embodiments, Ar<sub>1</sub> and Ar<sub>2</sub> in Formulae 1-1 to 1-6 may each independently be selected from:

a benzene group; and

10 a benzene group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>2</sub>-C<sub>20</sub> heteroaryl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), but embodiments of the present disclosure are not limited thereto.

15 In one or more embodiments, R<sub>11</sub> and R<sub>12</sub> in Formulae 1-1 to 1-6 may each independently be selected from a group represented by any of Formulae 5-1 to 5-80, but embodiments of the present disclosure are not limited thereto.

20 In one or more embodiments, R<sub>11</sub> and R<sub>12</sub> in Formulae 1-1 to 1-6 may each independently be selected from a group represented by any of Formulae 6-1 to 6-34, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, in Formulae 1-1 to 1-6,

25 R<sub>11</sub> and R<sub>12</sub> may each independently be selected from a group represented by any of Formulae 6-1 to 6-34, and Ar<sub>1</sub> and Ar<sub>2</sub> may each independently be selected from:

a benzene group; and  
 a benzene group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>2</sub>-C<sub>20</sub> heteroaryl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be selected from:

30 deuterium (—D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

35 a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

31

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group,


—Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

—Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),

wherein Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and





\* indicates a binding site to a neighboring atom.

In one or more embodiments, the amine-based compound represented by Formula 1 may be selected from any of Compounds 1 to 144, but embodiments of the present disclosure are not limited thereto:




32

-continued



**33**  
-continued




6

5

10

15

20



7


25

30

35

40

45



8


50

55


60

65


**34**  
-continued




9



10







11



12

**35**

-continued




**36**

-continued

13

17

5




10

15

14

18

20




25

30

15

19

35



40

45

50


16

20





55

60

65



**37**  
-continued



**38**  
-continued


21

25

5

10

15




22

26

20

25

30




23

27

40

45

50




24

28

55





60

65







39

-continued



40

-continued



29

5

10

15

30

20

25

30

31

35

40

45

50

32

55

60

65

33





34

35

36

41

-continued




42

-continued

37

41

5



10

15


38

42

20

25

30



39


43

35

40

45

50




40

44

55





60

65



43

-continued



44

-continued


45

49

5

10

15



46

20

50

25


30

35

47

40

45



50


51

48

55

60


65



45

-continued

52




5

10

15

20

53 25




30

35

40

45

54



50

55


60

65


46

-continued


55



56







57



47

-continued



48

-continued

58

62

5

10

15

59

20

25

63

30

60

35

40

45

61




50

55

64


60

65








**49**

-continued







**50**

-continued



**51**

-continued



**52**

-continued

71

75

5

10

15

72

20

25

30

35

73

40

45





50

74

55

60

65




76

77


78

53


-continued




79



80




81




82

54


-continued




83



84



85



86

5

10

15

20

25

30

35

40

45


50

55

60


65

**55**  
-continued



87

**56**  
-continued



90

5

10

15

20

25

88


30

35

40

45

91





89

55

60


65

92



**57**

-continued



93

5


10

15

20

**58**

-continued



96

94

25

30

35

40

45

95


50

55



60

65

97




98



**59**


-continued



99

**60**

-continued



102


5

10

15

20

103




100 25

30

35

40

45



101



50

104

55





60

65



**61**

-continued




**62**

-continued

105

109

5




10

15

110

106

20



25


30

111

107

35

40



45

50


112

108

55

60


65



**63**

-continued

113




5

10

15

114




20

25

30

115




35

40

45

50

116



55


60

65


**64**

-continued


117




118



119




120




**65**

-continued


121




122



123




124



**66**

-continued

125



5

10

15

20

25

30

35

40

45


50

55


60

65


126



127




128



**67**  
-continued

129




5

10

15

130




20

25

30

131




40

45

50

132




55

60


65

**68**  
-continued


133




134



135




136




69

-continued


137




138



139



140



60


65

70

-continued

141

5



10

15

20

25

30

35


40

45

50


55

60




142

143



144



When the amine-based compound represented by Formula 1 satisfies a structure in which benzene is further condensed at a specific position of fluorene and a substituent having a cyclic structure is included at a specific position of fluorene (e.g., a fluorene group is substituted at the C9 position with a group having a cyclic structure and a phenyl ring of the fluorene group is further condensed with a

benzene group), hole hopping may be facilitated by orbital delocalization. The amine-based compound may thus have high charge mobility, and therefore, an organic light-emitting device including the amine-based compound may be driven with low power.

Also, when the amine-based compound includes one amino group within a molecule (e.g., is a monoamine compound), the amine-based compound may have a relatively low highest occupied molecular orbital (HOMO) energy level, as compared to a compound including two or more amino groups (e.g., a diamine, triamine, etc.). As such, the amine-based compound (e.g., monoamine compound) may have an energy level optimized for hole transport. Therefore, an electronic device, (for example, an organic light-emitting device) including the amine-based compound may have a low driving voltage, high efficiency, and a long lifespan.

A synthetic method for preparing the amine-based compound represented by Formula 1 should be apparent to those of ordinary skill in the art by referring to the following examples.

At least one of the amine-based compound represented by Formula 1 may be used between a pair of electrodes (e.g., in a layer positioned between a cathode and an anode) of an organic light-emitting device. For example, the amine-based compound may be included at least one selected from a hole transport region, an electron transport region, and an emission layer. In one or more embodiments, the amine-based compound of Formula 1 may be used as a material for a capping layer located outside a pair of electrodes (e.g., in a layer that is not positioned between a cathode and an anode) of an organic light-emitting device.

Accordingly, embodiments of the present disclosure provide an organic light-emitting device including: a first electrode; a second electrode facing the first electrode; and an organic layer between the first electrode and the second electrode, the organic layer including an emission layer, wherein the organic layer includes at least one amine-based compound (e.g., an amine-based compound represented by Formula 1).

The expression “(an organic layer) includes at least one amine-based compound”, as used herein, may refer to a case in which (an organic layer) includes identical amine-based compounds represented by Formula 1, and may also refer to a case in which (an organic layer) includes two or more different amine-based compounds represented by Formula 1.

In one or more embodiments, for example, the organic layer may include only Compound 1 as the amine-based compound. In this regard, Compound 1 may exist (e.g., be included) in an emission layer of the organic light-emitting device. In one or more embodiments, the organic layer may simultaneously include Compound 1 and Compound 2 (e.g., a mixture of Compound 1 and Compound 2) as the amine-based compound. In this regard, Compound 1 and Compound 2 may exist in an identical (e.g., the same) layer (for example, Compound 1 and Compound 2 may both exist in an emission layer), or may exist in different layers (for example, Compound 1 may exist in an emission layer and Compound 2 may exist in an electron transport layer).

In one or more embodiments, the first electrode may be an anode, the second electrode may be a cathode, and the organic layer may further include a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode. The hole transport region may include a hole injection layer, a hole transport layer, an emission auxiliary layer, an electron blocking layer, or any combina-

tion thereof, and the electron transport region may include a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, an electron injection layer, or any combination thereof.

In one or more embodiments, the hole transport region may include the amine-based compound (e.g., the amine-based compound represented by Formula 1).

In one or more embodiments, the hole transport region may include a hole transport layer, which includes the amine-based compound (e.g., the amine-based compound represented by Formula 1).

In one or more embodiments, the emission layer may include the amine-based compound represented by Formula 1.

In one or more embodiments, the hole transport region may include a p-dopant, and the p-dopant may have a lowest unoccupied molecular orbital (LUMO) energy level of about  $-3.5$  eV or less.

In one or more embodiments, the emission layer may include at least one selected from a styryl-based compound, an anthracene-based compound, a pyrene-based compound, and a spiro-bifluorene-based compound.

In one or more embodiments, the emission layer may be a first color light emission layer,

the organic light-emitting device may further include: i) at least one second color light emission layer or ii) at least one second color light emission layer and at least one third color light emission layer, between the first electrode and the second electrode,

a maximum emission wavelength of the first color light, a maximum emission wavelength of the second color light, and a maximum emission wavelength of the third color light may be identical to or different from each other, and,

the first color light and the second color light may be emitted in the form of mixed light, or the first color light, the second color light, and the third color light may be emitted in the form of mixed light.

The term “organic layer”, as used herein, may refer to a single layer and/or a plurality of layers positioned between the first electrode and the second electrode of an organic light-emitting device. The materials included in the “organic layer” are not limited to being organic materials.

FIG. 1 is a schematic view of an organic light-emitting device **10** according to an embodiment of the present disclosure. The organic light-emitting device **10** includes a first electrode **110**, an organic layer **150**, and a second electrode **190**.

Hereinafter, the structure of the organic light-emitting device **10** according to an embodiment of the present disclosure and a method of manufacturing the organic light-emitting device **10** will be described in connection with FIG. 1.

In FIG. 1, a substrate may be additionally positioned under the first electrode **110** or above the second electrode **190**. The substrate may be a glass substrate or a plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

The first electrode **110** may be formed by depositing or sputtering a material for forming the first electrode **110** on the substrate. When the first electrode **110** is an anode, the material for forming a first electrode **110** may be selected from materials with a high work function to facilitate hole injection.

The first electrode **110** may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. When the first electrode **110** is a transmissive electrode, a

73

material for forming a first electrode **110** may be selected from indium tin oxide (ITO), indium zinc oxide (IZO), tin oxide (SnO<sub>2</sub>), zinc oxide (ZnO), and any combination thereof, but embodiments of the present disclosure are not limited thereto. When the first electrode **110** is a semi-transmissive electrode or a reflective electrode, the material for forming the first electrode **110** may include magnesium (Mg), silver (Ag), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), or any combination thereof. However, the material for forming the first electrode **110** is not limited thereto.

The first electrode **110** may have a single-layered structure, or a multi-layered structure including two or more layers. For example, the first electrode **110** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **110** is not limited thereto.

The organic layer **150** is positioned on the first electrode **110**. The organic layer **150** may include an emission layer.

The organic layer **150** may further include a hole transport region between the first electrode **110** and the emission layer, and an electron transport region between the emission layer and the second electrode **190**.

The hole transport region may have: i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.


The hole transport region may include at least one layer selected from a hole injection layer (HIL), a hole transport layer (HTL), an emission auxiliary layer, and an electron blocking layer (EBL).

For example, the hole transport region may have a single-layered structure including a single layer including a plurality of different materials, or a multi-layered structure having a hole injection layer/hole transport layer structure, a hole injection layer/hole transport layer/emission auxiliary layer structure, a hole injection layer/emission auxiliary layer structure, a hole transport layer/emission auxiliary layer structure, or a hole injection layer/hole transport layer/electron blocking layer structure, wherein the constituting layers of each structure are sequentially stacked on the first electrode **110** in their stated orders. However, the structure of the hole transport region is not limited thereto.

The hole transport region may include the amine-based compound represented by Formula 1.


The hole transport region may include, in addition to the amine-based compound represented by Formula 1, at least one selected from m-MTDATA, TDATA, 2-TNATA, NPB (NPD), β-NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4''-tris(N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201, and a compound represented by Formula 202:

74




75


-continued




NPB




$\beta$ -NPB




TPD



Spiro-TPD




Spiro-NPB




methylated NPB

76

-continued




TAPC




HMTPD

Formula 201



Formula 202



In Formulae 201 and 202,

$L_{201}$  to  $L_{204}$  may each independently be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$L_{205}$  may be selected from  $*-O-*$ ,  $*-S-*$ ,  $*-N(Q_{201})-*$ , a substituted or unsubstituted  $C_1$ - $C_{20}$  alkylene group, a substituted or unsubstituted  $C_2$ - $C_{20}$  alkenylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

xa1 to xa4 may each independently be an integer selected from 0 to 3,

xa5 may be an integer selected from 1 to 10, and

R<sub>201</sub> to R<sub>204</sub> and Q<sub>201</sub> may each independently be selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group.

In one or more embodiments, in Formula 202, R<sub>201</sub> and R<sub>202</sub> may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group, and R<sub>203</sub> and R<sub>204</sub> may optionally be linked via a single bond, a dimethyl-methylene group, or a diphenyl-methylene group.

In one or more embodiments, in Formulae 201 and 202,

L<sub>201</sub> to L<sub>205</sub> may each independently be selected from:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylenylene group, and a pyridinylenylene group; and

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylene group, an indacenylene group, an acenaphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenalenylenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, a pentaphenylenylene group, a hexacenylenylene group, a pentacenylenylene group, a rubicenylenylene group, a coronenylenylene group, an ovalenylenylene group, a thiophenylenylene group, a furanylenylene group, a carbazolylene group, an indolylenylene group, an isoindolylenylene group, a benzofuranylenylene group, a benzothiophenylenylene group, a dibenzofuranylenylene group, a dibenzothiophenylenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, and a pyridinylenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphe-

nyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), and —N(Q<sub>31</sub>)(Q<sub>32</sub>),

wherein Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, xa1 to xa4 may each independently be 0, 1, or 2.

In one or more embodiments, xa5 may be 1, 2, 3, or 4.

In one or more embodiments, R<sub>201</sub> to R<sub>204</sub> and Q<sub>201</sub> may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenylyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenylyl group, a pentacenylyl group, a rubicenylyl group, a coronenylyl group, an ovalenylyl group, a thiophenylyl group, a furanylyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl

79

group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), and —N(Q<sub>31</sub>)(Q<sub>32</sub>),

wherein Q<sub>31</sub> to Q<sub>33</sub> are the same as described above.

In one or more embodiments, at least one of R<sub>201</sub> to R<sub>203</sub> in Formula 201 may be selected from:

a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and

a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,

but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, in Formula 202, i) R<sub>201</sub> and R<sub>202</sub> may be linked via a single bond, and/or ii) R<sub>203</sub> and R<sub>204</sub> may be linked via a single bond.

In one or more embodiments, at least one of R<sub>201</sub> to R<sub>204</sub> in Formula 202 may be selected from:

a carbazolyl group; and


a carbazolyl group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group,

but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A:


80

Formula 201A




In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A(1), but embodiments of the present disclosure are not limited thereto:

Formula 201A(1)



In one or more embodiments, the compound represented by Formula 201 may be represented by Formula 201A-1, but embodiments of the present disclosure are not limited thereto:

Formula 201A-1




In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A:


81

82

Formula 202A



In one or more embodiments, the compound represented by Formula 202 may be represented by Formula 202A-1:



In Formulae 201A, 201A(1), 201A-1, 202A, and 202A-1, L<sub>201</sub> to L<sub>203</sub>, xa1 to xa3, xa5, and R<sub>202</sub> to R<sub>204</sub> may each be the same as described above,

R<sub>211</sub> and R<sub>212</sub> may be understood by referring to the description provided herein in connection with R<sub>203</sub>, and



R<sub>213</sub> to R<sub>217</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group,

a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a terphenyl group, a phenyl group substituted with a C<sub>1</sub>-C<sub>10</sub> alkyl group, a phenyl group substituted with —F, a pentalenyl group, an indenyl group, a naphthyl group, an azulenyl group, a heptalenyl group, an indacenyl group, an acenaphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a naphthacenyl group, a picenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a rubicenyl group, a coronenyl group, an ovalenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

The hole transport region may include at least one compound selected from any of Compounds HT1 to HT39, but embodiments of the present disclosure are not limited thereto:

HT1

HT2





83

84



-continued  
HT3

HT4



HT5

HT6





85

86



-continued  
HT7

HT8




HT9

HT10




87

-continued  
HT11





88

HT12





HT13

HT14



HT15


HT16






91

-continued  
HT21




92


HT22



HT23




93




-continued  
HT24

94




HT25


HT26




HT27




HT28



HT29




95




-continued  
HT30

96




HT31


HT32




HT33



HT34



HT35





97

98



-continued  
HT36

HT37



HT38

HT39



The hole transport region may have a thickness of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, the hole injection layer may have a thickness of about 100 Å to about 9,000 Å, and for example, about 100 Å to about 1,000 Å, and the hole transport layer may have a thickness of about 50 Å to about 2,000 Å, and for example, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer, and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.

The emission auxiliary layer may increase the light-emission efficiency of the device by compensating for an optical resonance distance according to the wavelength of light emitted by an emission layer (e.g., by adjusting the optical resonance distance to match the wavelength of light emitted from the emission layer), and the electron blocking layer may block or reduce the flow of electrons from an electron transport region. The emission auxiliary layer and the electron blocking layer may each include the materials as described above.

The hole transport region may further include, in addition to these materials, a charge-generation material for the

improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

The charge-generation material may be, for example, a p-dopant.

In one or more embodiments, the p-dopant may have a lowest unoccupied molecular orbital (LUMO) energy level of about -3.5 eV or less.

The p-dopant may include at least one selected from a quinone derivative, a metal oxide, and a cyano group-containing compound, but embodiments of the present disclosure are not limited thereto.

In one of more embodiments, for example, the p-dopant may include at least one selected from:


a quinone derivative, such as tetracyanoquinodimethane (TCNQ) and 2,3,5,6-tetrafluoro-7,7,8,8-tetracyanoquinodimethane (F4-TCNQ);

a metal oxide, such as tungsten oxide or molybdenum oxide;


1,4,5,8,9,12-hexaazatriphenylene-hexacarbonitrile (HAT-CN); and

a compound represented by Formula 221,


but embodiments of the present disclosure are not limited thereto:



HAT-CN



F4-TCNQ



Formula 221

In Formula 221,

R<sub>221</sub> to R<sub>223</sub> may each independently be selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, wherein at least one selected from R<sub>221</sub> to R<sub>223</sub> may include at least one substituent selected from a cyano group, —F, —Cl, —Br, —I, a C<sub>1</sub>-C<sub>20</sub> alkyl group substituted with —F, a C<sub>1</sub>-C<sub>20</sub> alkyl group substituted with —Cl, a C<sub>1</sub>-C<sub>20</sub> alkyl group substituted with —Br, and a C<sub>1</sub>-C<sub>20</sub> alkyl group substituted with —I.

When the organic light-emitting device 10 is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, or a blue emission layer, according to a sub-pixel. In one or more embodiments, the emission layer may have a stacked structure of two or more layers selected from a red emission layer, a green emission layer, and a blue emission layer, in which the two or more layers may contact each other or may be separated from each other. In one or more embodiments, the emission layer may include two or more materials selected from a red light-emitting material, a green light-emitting material, and a blue light-emitting material, in which the two or more materials are mixed with each other in a single layer to emit white light.

The emission layer may include a host and a dopant. The dopant may include at least one selected from a phosphorescent dopant and a fluorescent dopant.

The dopant may be included in the emission layer in an amount of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.

The emission layer may have a thickness of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent emission characteristics may be obtained without a substantial increase in driving voltage.

In one or more embodiments, the host may include a compound represented by Formula 301:



In Formula 301,

Ar<sub>301</sub> may be a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group,

xb11 may be 1, 2, or 3,

L<sub>301</sub> may each independently be selected from a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylene group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

xb1 may be an integer from 0 to 5,

R<sub>301</sub> may be selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazone group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>301</sub>)(Q<sub>302</sub>)(Q<sub>303</sub>), —N(Q<sub>301</sub>)(Q<sub>302</sub>), —B(Q<sub>301</sub>)(Q<sub>302</sub>), —C(=O)(Q<sub>301</sub>), —S(=O)<sub>2</sub>(Q<sub>301</sub>), and —P(=O)(Q<sub>301</sub>)(Q<sub>302</sub>), and

xb21 may be an integer from 1 to 5,

wherein Q<sub>301</sub> to Q<sub>303</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, Ar<sub>301</sub> in Formula 301 may be selected from:

a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothioophene group; and


## 101

a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, and a dibenzothio-  
 5 phenylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>) and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),


wherein Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group. However, embodiments of the present disclosure are not limited thereto.

When xb11 in Formula 301 is two or more, two or more Ar<sub>301(s)</sub> may be linked via a single bond.

In one or more embodiments, the compound represented by Formula 301 may be represented by Formula 301-1 or 301-2:



Formula 301-1



Formula 301-2

In Formulae 301-1 and 301-2,

A<sub>301</sub> to A<sub>304</sub> may each independently be selected from a benzene group, a naphthalene group, a phenanthrene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a pyridine group, a pyrimidine group, an indene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, an indole group, a carbazole group, a benzocarbazole group, a dibenzocarbazole group, a furan group, a benzofuran group, a dibenzofuran group, a naphthofuran group, a benzonaphthofuran group, a dinaphthofuran group, a thiophene group, a benzothiophene group, a dibenzothiophene group, a naphthothiophene group, a benzonaphthothiophene group, and a dinaphthothiophene group,

X<sub>301</sub> may be O, S, or N-[(L<sub>304</sub>)<sub>xb4</sub>-R<sub>304</sub>],

R<sub>311</sub> to R<sub>314</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a

## 102

hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —N(Q<sub>31</sub>)(Q<sub>32</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),

xb22 and xb23 may each independently be 0, 1, or 2,

L<sub>301</sub>, xb1, R<sub>301</sub> and Q<sub>31</sub> to Q<sub>33</sub> may each be the same as described above,

L<sub>302</sub> to L<sub>304</sub> may each independently be the same as described in connection with L<sub>301</sub>,

xb2 to xb4 may each independently be the same as described in connection with xb1, and

R<sub>302</sub> to R<sub>304</sub> may each independently be the same as described in connection with R<sub>301</sub>.

For example, in Formulae 301, 301-1, and 301-2, L<sub>301</sub> to L<sub>304</sub> may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofura-

nylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group, an imidazolylylene group, a pyrazolylylene group, a thiazolylylene group, an isothiazolylylene group, an oxazolylylene group, an isoxazolylylene group, a thiadiazolylylene group, an oxadiazolylylene group, a pyrazinylylene group, a pyrimidinylylene group, a pyridazinylylene group, a triazinylylene group, a quinolinylylene group, an isoquinolinylylene group, a benzoquinolinylylene group, a phthalazinylylene group, a naphthyridinylylene group, a quinoxalinylylene group, a quinazolinylylene group, a cinnolinylylene group, a phenanthridinylylene group, an acridinylylene group, a phenanthrolinylylene group, a phenazinylylene group, a benzimidazolylylene group, an isobenzothiazolylylene group, a benzoxazolylylene group, an isobenzoxazolylylene group, a triazolylylene group, a tetrazolylylene group, an imidazopyridinylylene group, an imidazopyrimidinylylene group, and an azacarbazolylylene group; and







105

linyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, an azacarbazolyl group,  $-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{C}(=\text{O})(\text{Q}_{31})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$  and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ ,

wherein  $\text{Q}_{31}$  to  $\text{Q}_{33}$  may be the same as described above.







In one or more embodiments, the host may include an alkaline earth metal complex. For example, the host may be selected from a Be complex (for example, Compound H55) and a Mg complex. In some embodiments, the host may include a Zn complex.

In one or more embodiments, the host may include at least one selected from 9,10-di(2-naphthyl)anthracene (ADN), 2-methyl-9,10-bis(naphthalen-2-yl)anthracene (MADN), 9,10-di-(2-naphthyl)-2-t-butyl-anthracene (TBADN), 4,4'-bis(N-carbazolyl)-1,1'-biphenyl (CBP), 1,3-di-9-carbazolylbenzene (mCP), 1,3,5-tri(carbazol-9-yl)benzene (TCP), and any of Compounds H1 to H55, but embodiments of the present disclosure are not limited thereto:



106


-continued




5  
H5  
15  
H6  
20  
25  
H7  
30  
35  
H8  
40  
45  
H9  
50  
55  
60  
65  
H10

**107**


-continued



5




10



15


20



25

30

35



40

45

50


55

60


65

**108**


-continued




H16




H17



H18




H19



H20

109

-continued




H21

5

10

15




H22

20

25

30




H23

35

40

45



H24


55

60


65

110

-continued






H25



H26

**111**  
-continued



**112**  
-continued

H27

H30

5

10

15

20

25

H28

30

35

40

45






H29

50

55

60

65



H31


H32

H33

H34

113

-continued



H35


5

10

15

H36

20




25

30

35

H37

40




45

50

55

H38

60




65


114

-continued


H39







H40



H41



**115**  
-continued



**116**  
-continued

H42

H46

5

10

15

H43

20

H47

25

30

H48

35

H44

40

H49

45

H45

50







55

60

65


H50

H51




117


-continued




H52



H53




H54



H55

The phosphorescent dopant may include an organometallic complex represented by Formula 401 below:




Formula 401

118

-continued

Formula 402



5

10

15

In Formulae 401 and 402,

M may be selected from iridium (Ir), platinum (Pt), palladium (Pd), osmium (Os), titanium (Ti), zirconium (Zr), hafnium (Hf), europium (Eu), terbium (Tb), rhodium (Rh), and thulium (Tm),

$L_{401}$  may be selected from ligands represented by Formula 402, and  $xc1$  may be 1, 2, or 3, wherein, when  $xc1$  is two or more, two or more  $L_{401}(s)$  may be identical to or different from each other,

$L_{402}$  may be an organic ligand, and  $xc2$  may be an integer from 0 to 4, wherein, when  $xc2$  is two or more, two or more  $L_{402}(s)$  may be identical to or different from each other,

$X_{401}$  to  $X_{404}$  may each independently be nitrogen (N) or carbon (C),

$X_{401}$  and  $X_{403}$  may be linked via a single bond or a double bond, and  $X_{402}$  and  $X_{404}$  may be linked via a single bond or a double bond,

$A_{401}$  and  $A_{402}$  may each independently be selected from a  $C_5$ - $C_{60}$  carbocyclic group or a  $C_1$ - $C_{60}$  heterocyclic group,

$X_{405}$  may be a single bond,  $*-O-*$ ,  $*-S-*$ ,  $*-C(=O)-*$ ,  $*-N(Q_{411})-*$ ,  $*-C(Q_{411})(Q_{412})-*$ ,  $*-C(Q_{411})=C(Q_{412})-*$ ,  $*(Q_{411})=*$ , or  $*=C-*$ , wherein  $Q_{411}$  and  $Q_{412}$  may be hydrogen, deuterium, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group,

$X_{406}$  may be a single bond, O, or S,

$R_{401}$  and  $R_{402}$  may each independently be selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{20}$  alkyl group, a substituted or unsubstituted  $C_1$ - $C_{20}$  alkoxy group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-Si(Q_{401})(Q_{402})(Q_{403})$ ,  $-N(Q_{401})(Q_{402})$ ,  $-B(Q_{401})(Q_{402})$ ,  $-C(=O)(Q_{401})$ ,  $-S(=O)_2(Q_{401})$ , and  $-P(=O)(Q_{401})(Q_{402})$ , wherein  $Q_{401}$  to  $Q_{403}$  may each independently be selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a  $C_6$ - $C_{20}$  aryl group, and a  $C_1$ - $C_{20}$  heteroaryl group,

$xc11$  and  $xc12$  may each independently be an integer selected from 0 to 10, and

\* and \* in Formula 402 each indicate a binding site to M of Formula 401.

In one or more embodiments,  $A_{401}$  and  $A_{402}$  in Formula 402 may each independently be selected from a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, an indene group, a pyrrole group, a thiophene group, a furan group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a quinoxaline group, a quinazoline group, a carbazole group, a benzimidazole group, a benzofuran group, a benzothio-  
 5 phenene group, an isobenzothiophene group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a dibenzofuran group, and a dibenzothiophene group.

In one or more embodiments, in Formula 402, i)  $X_{401}$  may be nitrogen, and  $X_{402}$  may be carbon, or ii)  $X_{401}$  and  $X_{402}$  may each be nitrogen at the same time.

In one or more embodiments, in Formula 402,  $R_{401}$  and  $R_{402}$  may each independently be selected from:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group;

a  $C_1$ - $C_{20}$  alkyl group, and a  $C_1$ - $C_{20}$  alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a phenyl group, a naphthyl group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, and a norbornenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group;

a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a quinoxalinyl group, a quinazoliny group, a carbazolyl group, a dibenzofuranyl group, and a dibenzothiophenyl group; and  
 —Si( $Q_{401}$ )( $Q_{402}$ )( $Q_{403}$ ), —N( $Q_{401}$ )( $Q_{402}$ ), —B( $Q_{401}$ )( $Q_{402}$ ), —C(=O)( $Q_{401}$ ), —S(=O)<sub>2</sub>( $Q_{401}$ ), and —P(=O)( $Q_{401}$ )( $Q_{402}$ ),

wherein  $Q_{401}$  to  $Q_{403}$  may each independently be selected from a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl


group, a biphenyl group, and a naphthyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, when xc1 in Formula 401 is two or more, two  $A_{401}$ (s) in two or more  $L_{401}$ (s) may optionally be linked via  $X_{407}$ , which is a linking group, or two  $A_{402}$ (s) in two or more  $L_{401}$ (s) may optionally be linked via  $X_{408}$ , which is a linking group (see Compounds PD1 to PD4 and PD7).  $X_{407}$  and  $X_{408}$  may each independently be a single bond, \*—O—\*, \*—S—\*, \*—C(=O)—\*, \*—N( $Q_{413}$ )\*, \*—C( $Q_{413}$ )( $Q_{414}$ )\*, or \*—C( $Q_{413}$ )=C( $Q_{414}$ )\* (wherein  $Q_{413}$  and  $Q_{414}$  may each independently be hydro-  
 5 gen, deuterium, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group), but are not limited thereto.


$L_{402}$  in Formula 401 may be a monovalent, divalent, or trivalent organic ligand. For example,  $L_{402}$  may be selected from halogen, diketone (for example, acetylacetonate), carboxylic acid (for example, picolinate), —C(=O), isonitrile, —CN, and phosphorus (for example, phosphine and phosphite), but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, the phosphorescent dopant may be selected from, for example, any of Compounds PD1 to PD25, but embodiments of the present disclosure are not limited thereto:


PD1









PD2



PD3




**121**  
-continued



**122**  
-continued

PD4

5




PD10

10

PD5

20




PD11

25

PD6

30




PD12

35

PD7

40




PD13

45

PD8


50



55

PD9






60



PD14


65

**123**  
-continued




**124**  
-continued



PD15




PD16




PD17



PD18



PD19



65

PD20


PD21

PD22


PD23

125

-continued




PD24



PD25

The fluorescent dopant may include an arylamine compound or a styrylamine compound.

The fluorescent dopant may include a compound represented by Formula 501:



Formula 501

In Formula 501,

$\text{Ar}_{501}$  may be a substituted or unsubstituted  $\text{C}_5\text{-C}_{60}$  carbocyclic group or a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  heterocyclic group,

$\text{L}_{501}$  to  $\text{L}_{503}$  may each independently be selected from a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkylene group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkenylene group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  arylene group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group,

$\text{xd1}$  to  $\text{xd3}$  may each independently be an integer of 0 to 3,

$\text{R}_{501}$  and  $\text{R}_{502}$  may each independently be selected from a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkyl group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $\text{C}_3\text{-C}_{10}$  cycloalkenyl group, a substituted or unsubstituted  $\text{C}_1\text{-C}_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  aryl group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  aryloxy group, a substituted or unsubstituted  $\text{C}_6\text{-C}_{60}$  arylthio group, a substituted or

126

unsubstituted  $\text{C}_1\text{-C}_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and

$\text{xd4}$  may be an integer of 1 to 6.

In one or more embodiments,  $\text{Ar}_{501}$  in Formula 501 may be selected from:

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indeno-phenanthrene group; and

a naphthalene group, a heptalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, and an indeno-phenanthrene group, each substituted with at least one selected from deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1\text{-C}_{20}$  alkyl group, a  $\text{C}_1\text{-C}_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments,  $\text{L}_{501}$  to  $\text{L}_{503}$  in Formula 501 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, a pyridinylylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylylene group, a fluoranthenylylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylylene group, a hexacenylylene group, a pentacenylylene group, a thiophenylylene group, a furanylylene group, a carbazolylylene group, an indolylylene group, an isoindolylylene group, a benzofuranylylene group, a benzothiophenylylene group, a dibenzofuranylylene group, a dibenzothiophenylylene group, a benzocarbazolylylene group, a dibenzocarbazolylylene group, a dibenzosilolylylene group, and a pyridinylylene group, each substituted with at least one selected from deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1\text{-C}_{20}$  alkyl group, a  $\text{C}_1\text{-C}_{20}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl

127

group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group.

In one or more embodiments, in Formula 501, R<sub>501</sub> and R<sub>502</sub> may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group; and

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, and a pyridinyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a pyridinyl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>),


wherein Q<sub>31</sub> to Q<sub>33</sub> may each be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, xd4 in Formula 501 may be 2, but embodiments of the present disclosure are not limited thereto.


In one or more embodiments, for example, the fluorescent dopant may be selected from any of Compounds FD1 to FD22:

128


FD4



FD5




FD6




129

-continued







130

-continued



**131**  
-continued



**132**  
-continued

FD15

FD19

5

10

15

FD16

20

25

30

FD17

35

FD20

40

45

FD21


FD18

50

55

60


FD22




65 In one or more embodiments, the fluorescent dopant may be selected from the following compounds, but embodiments of the present disclosure are not limited thereto:

133


134




DPVBi




DPAVBi




TBPe




DCM



DCJTb



Coumarin 6



C545T

The electron transport region may have: i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron transport region may include at least one selected from a buffer layer, a hole blocking layer, an electron control layer, an electron transport layer, and an electron injection layer, but embodiments of the present disclosure are not limited thereto.

For example, the electron transport region may have an electron transport layer/electron injection layer structure, a hole blocking layer/electron transport layer/electron injection layer structure, an electron control layer/electron transport layer/electron injection layer structure, or a buffer layer/electron transport layer/electron injection layer structure, wherein the constituting layers of each structure are sequentially stacked from an emission layer. However, embodiments of the structure of the electron transport region are not limited thereto.

The electron transport region (for example, a buffer layer, a hole blocking layer, an electron control layer, or an electron transport layer in the electron transport region) may include a metal-free compound containing at least one  $\pi$  electron-depleted nitrogen-containing ring.

As used herein, the term " $\pi$  electron-depleted nitrogen-containing ring" may refer to a  $C_1$ - $C_{60}$  heterocyclic group having at least one  $*-N=*$  moiety as a ring-forming moiety.

For example, the " $\pi$  electron-depleted nitrogen-containing ring" may be: i) a 5-membered to 7-membered heteromonocyclic group having at least one  $*-N=*$  moiety, ii) a heteropolycyclic group in which two or more 5-membered to 7-membered heteromonocyclic groups each having at least one  $*-N=*$  moiety are condensed with each other, or iii) a heteropolycyclic group in which at least one of 5-membered to 7-membered heteromonocyclic groups, each having at least one  $*-N=*$  moiety, is condensed with at least one  $C_5$ - $C_{60}$  carbocyclic group.

Non-limiting examples of the  $\pi$  electron-depleted nitrogen-containing ring may include an imidazole, a pyrazole, a thiazole, an isothiazole, an oxazole, an isoxazole, a pyridine, a pyrazine, a pyrimidine, a pyridazine, an indazole, a purine, a quinoline, an isoquinoline, a benzoquinoline, a phthalazine, a naphthyridine, a quinoxaline, a quinazoline, a cinnoline, a phenanthridine, an acridine, a phenanthroline, a phenazine, a benzimidazole, an isobenzothiazole, a benzoxazole, an isobenzoxazole, a triazole, a tetrazole, an oxadiazole, a triazine, a thiadiazole, an imidazopyridine, an imidazopyrimidine, and an azacarbazole.

In one or more embodiments, for example, the electron transport region may include a compound represented by Formula 601:



In Formula 601,

$Ar_{601}$  may be a substituted or unsubstituted  $C_5$ - $C_{60}$  carbocyclic group or a substituted or unsubstituted  $C_1$ - $C_{60}$  heterocyclic group,

$xe11$  may be 1, 2, or 3,

$L_{601}$  may each independently be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkylene group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenylene group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenylene group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylene

group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroarylene group, a substituted or unsubstituted divalent non-aromatic condensed polycyclic group, and a substituted or unsubstituted divalent non-aromatic condensed heteropolycyclic group;

$xe1$  may be an integer from 0 to 5,

$R_{601}$  may be selected from a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3$ - $C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_1$ - $C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6$ - $C_{60}$  arylthio group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-\text{Si}(\text{Q}_{601})(\text{Q}_{602})(\text{Q}_{603})$ ,  $-\text{C}(=\text{O})(\text{Q}_{601})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{601})$ , and  $-\text{P}(=\text{O})(\text{Q}_{601})(\text{Q}_{602})$ ,

wherein  $\text{Q}_{601}$  to  $\text{Q}_{603}$  may each independently be a  $C_1$ - $C_{10}$  alkyl group, a  $C_1$ - $C_{10}$  alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, or a naphthyl group, and  $xe21$  may be an integer from 1 to 5.

In one or more embodiments, at least one of  $Ar_{601}(s)$  in the number of  $xe11$  and/or at least one of  $R_{601}(s)$  in the number of  $xe21$  may include the  $\pi$  electron-depleted nitrogen-containing ring.

In one or more embodiments, ring  $Ar_{601}$  in Formula 601 may be selected from:

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzimidazole group, an isobenzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group; and

a benzene group, a naphthalene group, a fluorene group, a spiro-bifluorene group, a benzofluorene group, a dibenzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, an indenoanthracene group, a dibenzofuran group, a dibenzothiophene group, a carbazole group, an imidazole group, a pyrazole group, a thiazole group, an isothiazole group, an oxazole group, an isoxazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a phenanthridine group, an acridine group, a phenanthroline group, a

phenazine group, a benzimidazole group, an iso-benzothiazole group, a benzoxazole group, an isobenzoxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a thiadiazole group, an imidazopyridine group, an imidazopyrimidine group, and an azacarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),


wherein Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, a C<sub>1</sub>-C<sub>10</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

When xe11 in Formula 601 is two or more, two or more Ar<sub>601</sub>(s) may be linked via a single bond.

In one or more embodiments, Ar<sub>601</sub> in Formula 601 may be an anthracene group.

In one or more embodiments, a compound represented by Formula 601 may be represented by Formula 601-1:

Formula 601-1



In Formula 601-1,

X<sub>614</sub> may be N or C(R<sub>614</sub>), X<sub>615</sub> may be N or C(R<sub>615</sub>), X<sub>616</sub> may be N or C(R<sub>616</sub>), and at least one selected from X<sub>614</sub> to X<sub>616</sub> may be N,

L<sub>611</sub> to L<sub>613</sub> may each independently be the same as described in connection with L<sub>601</sub>,

xe611 to xe613 may each independently be the same as described in connection with xe1,

R<sub>611</sub> to R<sub>613</sub> may each independently be the same as described in connection with R<sub>601</sub>,

R<sub>614</sub> to R<sub>616</sub> may each independently be selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, and a naphthyl group.

In one or more embodiments, L<sub>601</sub> and L<sub>611</sub> to L<sub>613</sub> in Formulae 601 and 601-1 may each independently be selected from:

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a dibenzothiophenylene group, a benzocarbazolylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxa-

zolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group; and

a phenylene group, a naphthylene group, a fluorenylene group, a spiro-bifluorenylene group, a benzofluorenylene group, a dibenzofluorenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylene group, a perylenylene group, a pentaphenylene group, a hexacenylene group, a pentacenylene group, a thiophenylene group, a furanylene group, a carbazolylene group, an indolylene group, an isoindolylene group, a benzofuranylene group, a benzothiophenylene group, a dibenzofuranylene group, a benzothiophenylene group, a benzocarbazolylene group, a dibenzocarbazolylene group, a dibenzosilolylene group, a pyridinylene group, an imidazolylene group, a pyrazolylene group, a thiazolylene group, an isothiazolylene group, an oxazolylene group, an isoxazolylene group, a thiadiazolylene group, an oxadiazolylene group, a pyrazinylene group, a pyrimidinylene group, a pyridazinylene group, a triazinylene group, a quinolinylene group, an isoquinolinylene group, a benzoquinolinylene group, a phthalazinylene group, a naphthyridinylene group, a quinoxalinylene group, a quinazolinylene group, a cinnolinylene group, a phenanthridinylene group, an acridinylene group, a phenanthrolinylene group, a phenazinylene group, a benzimidazolylene group, an isobenzothiazolylene group, a benzoxazolylene group, an isobenzoxazolylene group, a triazolylene group, a tetrazolylene group, an imidazopyridinylene group, an imidazopyrimidinylene group, and an azacarbazolylene group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexacenyl group, a pentacenyl group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetra-

139

zoyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, xe1 and xe611 to xe613 in Formulae 601 and 601-1 may each independently be 0, 1, or 2.

In one or more embodiments, R<sub>601</sub> and R<sub>611</sub> to R<sub>613</sub> in Formula 601 and 601-1 may each independently be selected from:

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group;

a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imida-

140


zopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a phenyl group, a biphenyl group, a terphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a perylenyl group, a pentaphenyl group, a hexaceny group, a pentaceny group, a thiophenyl group, a furanyl group, a carbazolyl group, an indolyl group, an isoindolyl group, a benzofuranyl group, a benzothiophenyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, a dibenzosilolyl group, a pyridinyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a thiadiazolyl group, an oxadiazolyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a phthalazinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a benzimidazolyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, and an azacarbazolyl group; and

—S(=O)<sub>2</sub>(Q<sub>601</sub>) and —P(=O)(Q<sub>601</sub>)(Q<sub>602</sub>),

wherein Q<sub>601</sub> and Q<sub>602</sub> may be the same as described above.




The electron transport region may include at least one compound selected from any of Compounds ET1 to ET36, but embodiments of the present disclosure are not limited thereto:

ET1



141

-continued



142

-continued

ET2

5

10

15

20

25

ET3

30

35

40

45

ET4


50

55


60

65


ET5



ET6





ET7



143

-continued



144

-continued

ET8


5

10

15

20

25



ET10

ET9

40


45

50

55

60




65






ET11

ET12




**145**  
-continued



**146**  
-continued



**147**  
-continued



**148**  
-continued

ET19


ET22

5

10

15

20



ET20


ET23

25

30

35

40



45

ET24


ET21

50


55

60

65



**149**  
-continued



ET25


5

10

15

20

**150**  
-continued




ET28

20

ET29

ET26

25




30

35

40

45





ET27

50

55


60

65



ET30

**151**  
-continued



ET31


5

10

15

20

**152**  
-continued



ET34


ET32

25


30

35

40



ET35




ET33

45


50

55

60





ET36




65

In one or more embodiments, the electron transport region may include at least one selected from 2,9-dimethyl-4,7-diphenyl-1,10-phenanthroline (BCP), 4,7-diphenyl-1,10-phenanthroline (Bphen), Alq<sub>3</sub>, BAlq, 3-(biphenyl-4-yl)-5-(4-tert-butylphenyl)-4-phenyl-4H-1,2,4-triazole (TAZ), and NTAZ:


153

Alq<sub>3</sub>

BALq



TAZ



NTAZ

The buffer layer, the hole blocking layer, and/or the electron controlling layer may each independently have a thickness of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thicknesses of the buffer layer, the hole blocking layer, and the electron control layer are within these ranges, the electron blocking layer may have excellent electron blocking characteristics or electron control characteristics without a substantial increase in driving voltage.

The electron transport layer may have a thickness of about 100 Å to about 1,000 Å, for example, about 150 Å to about 500 Å. When the thickness of the electron transport layer is within these ranges, the electron transport layer may have satisfactory electron transport characteristics without a substantial increase in driving voltage.

The electron transport region (for example, the electron transport layer in the electron transport region) may further include, in addition to the materials described above, a metal-containing material.

The metal-containing material may include at least one selected from an alkali metal complex and an alkaline


154

earth-metal complex. The alkali metal complex may include a metal ion selected from a lithium (Li) ion, a sodium (Na) ion, a potassium (K) ion, a rubidium (Rb) ion, and a cesium (Cs) ion, and the alkaline earth-metal complex may include a metal ion selected from a beryllium (Be) ion, a magnesium (Mg) ion, a calcium (Ca) ion, a strontium (Sr) ion, and a barium (Ba) ion. The alkali metal complex may include a ligand coordinated with the metal ion of the alkali metal complex or the alkaline earth-metal complex, and the ligand may be selected from a hydroxy quinoline, a hydroxy isoquinoline, a hydroxy benzoquinoline, a hydroxy acridine, a hydroxy phenanthridine, a hydroxy phenyloxazole, a hydroxy phenylthiazole, a hydroxy phenyloxadiazole, a hydroxy phenylthiadiazole, a hydroxy phenylpyridine, a hydroxy phenylbenzimidazole, a hydroxy phenylbenzothiazole, a bipyridine, a phenanthroline, and a cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

For example, the metal-containing material may include a Li complex. The Li complex may include, for example, Compound ET-D1 (lithium quinolate, LiQ) or ET-D2:

25


ET-D1



30

ET-D2

35



40

The electron transport region may include an electron injection layer that facilitates injection of electrons from the second electrode **190**. The electron injection layer may directly contact the second electrode **190**.

The electron injection layer may have: i) a single-layered structure including a single layer including a single material, ii) a single-layered structure including a single layer including a plurality of different materials, or iii) a multi-layered structure having a plurality of layers including a plurality of different materials.

The electron injection layer may include an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combination thereof.

The alkali metal may be selected from Li, Na, K, Rb, and Cs. In one or more embodiments, the alkali metal may be Li, Na, or Cs. In one or more embodiments, the alkali metal may be Li or Cs, but embodiments of the present disclosure are not limited thereto.

The alkaline earth metal may be selected from Mg, Ca, Sr, and Ba.

The rare earth metal may be selected from scandium (Sc), yttrium (Y), cerium (Ce), terbium (Tb), ytterbium (Yb), and gadolinium (Gd).

The alkali metal compound, the alkaline earth-metal compound, and the rare earth metal compound may be selected from oxides and halides (for example, fluorides, chlorides, bromides, or iodides) of the alkali metal, the alkaline earth-metal, and the rare earth metal.

The alkali metal compound may be selected from alkali metal oxides (such as  $\text{Li}_2\text{O}$ ,  $\text{Cs}_2\text{O}$ , or  $\text{K}_2\text{O}$ ) and alkali metal halides (such as  $\text{LiF}$ ,  $\text{NaF}$ ,  $\text{CsF}$ ,  $\text{KF}$ ,  $\text{LiI}$ ,  $\text{NaI}$ ,  $\text{CsI}$ , or  $\text{KI}$ ). In one or more embodiments, the alkali metal compound may be selected from  $\text{LiF}$ ,  $\text{Li}_2\text{O}$ ,  $\text{NaF}$ ,  $\text{LiI}$ ,  $\text{NaI}$ ,  $\text{CsI}$ , and  $\text{KI}$ , but embodiments of the present disclosure are not limited thereto.

The alkaline earth-metal compound may be selected from alkaline earth-metal oxides (such as  $\text{BaO}$ ,  $\text{SrO}$ ,  $\text{CaO}$ ,  $\text{Ba}_x\text{Sr}_{1-x}\text{O}$  ( $0 < x < 1$ ), or  $\text{Ba}_x\text{Ca}_{1-x}\text{O}$  ( $0 < x < 1$ )). In one or more embodiments, the alkaline earth-metal compound may be selected from  $\text{BaO}$ ,  $\text{SrO}$ , and  $\text{CaO}$ , but embodiments of the present disclosure are not limited thereto.

The rare earth metal compound may be selected from  $\text{YbF}_3$ ,  $\text{ScF}_3$ ,  $\text{ScO}_3$ ,  $\text{Y}_2\text{O}_3$ ,  $\text{Ce}_2\text{O}_3$ ,  $\text{GdF}_3$ , and  $\text{TbF}_3$ . In one or more embodiments, the rare earth metal compound may be selected from  $\text{YbF}_3$ ,  $\text{ScF}_3$ ,  $\text{TbF}_3$ ,  $\text{YbI}_3$ ,  $\text{ScI}_3$ , and  $\text{TbI}_3$ , but embodiments of the present disclosure are not limited thereto.

The alkali metal complex, the alkaline earth-metal complex, and the rare earth metal complex may respectively include an alkali metal ion, alkaline earth-metal ion, and rare earth metal ion as described above. The ligand coordinated with the metal ion in the alkali metal complex, the alkaline earth-metal complex, or the rare earth metal complex may be selected from hydroxy quinoline, hydroxy isoquinoline, hydroxy benzoquinoline, hydroxy acridine, hydroxy phenanthridine, hydroxy phenyloxazole, hydroxy phenylthiazole, hydroxy phenyloxadiazole, hydroxy phenylthiadiazole, hydroxy phenylpyridine, hydroxy phenylbenzimidazole, hydroxy phenylbenzothiazole, bipyridine, phenanthroline, and cyclopentadiene, but embodiments of the present disclosure are not limited thereto.

The electron injection layer may include (e.g., may consist of) an alkali metal, an alkaline earth metal, a rare earth metal, an alkali metal compound, an alkaline earth-metal compound, a rare earth metal compound, an alkali metal complex, an alkaline earth-metal complex, a rare earth metal complex, or any combination thereof, as described above. In one or more embodiments, the electron injection layer may further include an organic material. When the electron injection layer further includes an organic material, the alkali metal, alkaline earth metal, rare earth metal, alkali metal compound, alkaline earth-metal compound, rare earth metal compound, alkali metal complex, alkaline earth-metal complex, rare earth metal complex, or combination thereof may be homogeneously or non-homogeneously dispersed in a matrix including the organic material.

The electron injection layer may have a thickness of about 1 Å to about 100 Å, for example, about 3 Å to about 90 Å. When the thickness of the electron injection layer is within the range described above, the electron injection layer may have satisfactory electron injection characteristics without a substantial increase in driving voltage.

The second electrode **190** may be positioned on the organic layer **150** having the above-described structure. The second electrode **190** may be a cathode that is an electron injection electrode, and in this regard, the material for forming the second electrode **190** may be a material having

a low work function, and may include a metal, an alloy, an electrically conductive compound, or a combination thereof.

In one or more embodiments, the second electrode **190** may include at least one selected from lithium (Li), silver (Ag), magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), magnesium-silver (Mg—Ag), ITO, and IZO, but embodiments of the present disclosure are not limited thereto. The second electrode **190** may be a transmissive electrode, a semi-transmissive electrode, or a reflective electrode.

The second electrode **190** may have a single-layered structure, or a multi-layered structure including two or more layers.

An organic light-emitting device **20** as illustrated in FIG. **2** includes a first capping layer **210**, a first electrode **110**, an organic layer **150**, and a second electrode **190** sequentially stacked in this stated order. An organic light-emitting device **30** as illustrated in FIG. **3** includes a first electrode **110**, an organic layer **150**, a second electrode **190**, and a second capping layer **220** sequentially stacked in this stated order. An organic light-emitting device **40** as illustrated in FIG. **4** includes a first capping layer **210**, a first electrode **110**, an organic layer **150**, a second electrode **190**, and a second capping layer **220** sequentially stacked in this stated order.

In FIGS. **2** to **4**, the first electrode **110**, the organic layer **150**, and the second electrode **190** may be understood by referring to the description presented in connection with FIG. **1**.

In the organic layer **150** of each of the organic light-emitting devices **20** and **40**, light generated in the emission layer may pass through the first electrode **110** (which is a semi-transmissive or transmissive electrode) and the first capping layer **210** toward the outside. In the organic layer **150** of each of the organic light-emitting devices **30** and **40**, light generated in the emission layer may pass through the second electrode **190** (which is a semi-transmissive or transmissive electrode) and the second capping layer **220** toward the outside.

The first capping layer **210** and the second capping layer **220** may increase the external luminescent efficiency of the device according to the principle of constructive interference.


The first capping layer **210** and the second capping layer **220** may each independently be an organic capping layer including an organic material, an inorganic capping layer including an inorganic material, or a composite capping layer including an organic material and an inorganic material.

At least one selected from the first capping layer **210** and the second capping layer **220** may include at least one material selected from carbocyclic compounds, heterocyclic compounds, amine-based compounds, porphyrin derivatives, phthalocyanine derivatives, a naphthalocyanine derivatives, alkali metal complexes, and alkaline earth-based complexes. The carbocyclic compound, the heterocyclic compound, and the amine-based compound may be optionally substituted with a substituent containing at least one element selected from O, N, S, selenium (Se), silicon (Si), fluorine (F), chlorine (Cl), bromine (Br), and iodine (I). In one or more embodiments, at least one selected from the first capping layer **210** and the second capping layer **220** may include an amine-based compound.


In one or more embodiments, at least one selected from the first capping layer **210** and the second capping layer **220** may include the compound represented by Formula 201 or the compound represented by Formula 202.

157


In one or more embodiments, at least one selected from the first capping layer **210** and the second capping layer **220** may include a compound selected from any of Compounds HT28 to HT33 and Compounds CP1 to CP5, but embodiments of the present disclosure are not limited thereto:




CP1



CP2



CP3




CP4

158

-continued

CP5



The organic light-emitting device according to embodiments of the present disclosure has been described in connection with FIGS. **1** to **4**. However, embodiments of the present disclosure are not limited thereto.

The layers constituting the hole transport region, the emission layer, and the layers constituting the electron transport region may each be formed in a set or predetermined region (e.g., position) using one or more suitable methods selected from vacuum deposition, spin coating, casting, Langmuir-Blodgett (LB) deposition, ink-jet printing, laser-printing, and laser-induced thermal imaging.

When the layers constituting the hole transport region, the emission layer, and the layers constituting the electron transport region are formed by vacuum deposition, for example, the vacuum deposition may be performed at a deposition temperature of about 100 to about 500° C., at a vacuum degree of about 10<sup>-8</sup> Torr to about 10<sup>-3</sup> Torr, and at a deposition rate of about 0 Å/sec to about 100 Å/sec depending on the material to be included in each layer, and the intended structure of each layer.

When the layers constituting the hole transport region, the emission layer, and the layers constituting the electron transport region are formed by spin coating, the spin coating may be performed at a coating speed of about 2,000 rpm to about 5,000 rpm and at a heat treatment temperature of about 80° C. to about 200° C., depending on the material to be included in each layer and the intended structure of each layer.

The term “C<sub>1</sub>-C<sub>60</sub> alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and non-limiting examples thereof may include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an iso-amyl group, and a hexyl group. The term “C<sub>1</sub>-C<sub>60</sub> alkylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>1</sub>-C<sub>60</sub> alkyl group.

The term “C<sub>2</sub>-C<sub>60</sub> alkenyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon double bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and non-limiting examples thereof may include an ethenyl group, a propenyl group, and a butenyl group. The term “C<sub>2</sub>-C<sub>60</sub> alkenylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>2</sub>-C<sub>60</sub> alkenyl group.

The term “C<sub>2</sub>-C<sub>60</sub> alkynyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon triple bond in the middle or at the terminus of the C<sub>2</sub>-C<sub>60</sub> alkyl group, and non-limiting examples thereof may include an ethynyl group and a propynyl group. The

term “C<sub>2</sub>-C<sub>60</sub> alkynylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>2</sub>-C<sub>60</sub> alkynyl group.

The term “C<sub>1</sub>-C<sub>60</sub> alkoxy group” as used herein refers to a monovalent group represented by —O-A<sub>101</sub> (wherein A<sub>101</sub> is a C<sub>1</sub>-C<sub>60</sub> alkyl group), and non-limiting examples thereof may include a methoxy group, an ethoxy group, and an isopropoxy group.

The term “C<sub>3</sub>-C<sub>10</sub> cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and non-limiting examples thereof may include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkyl group.

The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group” as used herein refers to a monovalent saturated monocyclic group having at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom and 1 to 10 carbon atoms, and non-limiting examples thereof may include a 1,2,3,4-oxatriazolidinyl group, a tetrahydrofuran group, a tetrahydrothiophenyl group. The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group.

The term “C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms, at least one carbon-carbon double bond in the ring thereof, and no aromaticity, and non-limiting examples thereof may include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group.

The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, 1 to 10 carbon atoms, and at least one double bond in its ring. Non-limiting examples of the C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group may include a 4,5-dihydro-1,2,3,4-oxatriazolyl group, a 2,3-dihydrofuran group, and a 2,3-dihydrothiophenyl group. The term “C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group” as used herein refers to a divalent group having substantially the same structure as the C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group.

The term “C<sub>6</sub>-C<sub>60</sub> aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and the term “C<sub>6</sub>-C<sub>60</sub> arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Non-limiting examples of the C<sub>6</sub>-C<sub>60</sub> aryl group may include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C<sub>6</sub>-C<sub>60</sub> aryl group and the C<sub>6</sub>-C<sub>60</sub> arylene group each include two or more rings, the rings may be fused to each other.

The term “C<sub>1</sub>-C<sub>60</sub> heteroaryl group” as used herein refers to a monovalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. The term “C<sub>1</sub>-C<sub>60</sub> heteroarylene group” as used herein refers to a divalent group having a heterocyclic aromatic system that has at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, in addition to 1 to 60 carbon atoms. Non-limiting examples of the C<sub>1</sub>-C<sub>60</sub> heteroaryl group include a pyridinyl group, a pyrim-

idinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C<sub>1</sub>-C<sub>60</sub> heteroaryl group and the C<sub>1</sub>-C<sub>60</sub> heteroarylene group each include two or more rings, the rings may be fused to each other.

The term “C<sub>6</sub>-C<sub>60</sub> aryloxy group” as used herein refers to —O-A<sub>102</sub> (wherein A<sub>102</sub> is a C<sub>6</sub>-C<sub>60</sub> aryl group), and the term “C<sub>6</sub>-C<sub>60</sub> arylthio group” as used herein refers to —S-A<sub>103</sub> (wherein A<sub>103</sub> is a C<sub>6</sub>-C<sub>60</sub> aryl group).

The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) having two or more rings condensed with each other, only carbon atoms as ring-forming atoms, and no aromaticity in its entire molecular structure. A non-limiting example of the monovalent non-aromatic condensed polycyclic group is a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having substantially the same structure as the monovalent non-aromatic condensed polycyclic group.

The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 1 to 60 carbon atoms) having two or more rings condensed to each other, at least one heteroatom selected from N, O, Si, P, and S as a ring-forming atom, and no aromaticity in its entire molecular structure. A non-limiting example of the monovalent non-aromatic condensed heteropolycyclic group is a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

The term “C<sub>5</sub>-C<sub>60</sub> carbocyclic group” as used herein refers to a monocyclic or polycyclic group having 5 to 60 carbon atoms, in which the ring-forming atoms are only carbon atoms. The C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be an aromatic carbocyclic group or a non-aromatic carbocyclic group. The C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be a ring, such as benzene, a monovalent group, such as a phenyl group, or a divalent group, such as a phenylene group. In one or more embodiments, depending on the number of substituents connected to the C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the C<sub>5</sub>-C<sub>60</sub> carbocyclic group may be a trivalent group or a quadrivalent group.

The term “C<sub>1</sub>-C<sub>60</sub> heterocyclic group” as used herein refers to a group having substantially the same structure as the C<sub>1</sub>-C<sub>60</sub> carbocyclic group, except that at least one heteroatom selected from N, O, Si, P, and S may be used as a ring-forming atom in addition to carbon (the number of carbon atoms may be in a range of 1 to 60).

At least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkylene group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkylene group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenylene group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenylene group, the substituted C<sub>6</sub>-C<sub>60</sub> arylene group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroarylene group, the substituted divalent non-aromatic condensed polycyclic group, the substituted divalent non-aromatic condensed heteropolycyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub>

heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group, when present, may be selected from:

deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), -N(Q<sub>11</sub>)(Q<sub>12</sub>), -B(Q<sub>11</sub>)(Q<sub>12</sub>), -C(=O)(Q<sub>11</sub>), -S(=O)<sub>2</sub>(Q<sub>11</sub>), and -P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, and a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic

group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group,

-Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), -N(Q<sub>21</sub>)(Q<sub>22</sub>), -B(Q<sub>21</sub>)(Q<sub>22</sub>), -C(=O)(Q<sub>21</sub>), -S(=O)<sub>2</sub>(Q<sub>21</sub>), and -P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

-Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), -N(Q<sub>31</sub>)(Q<sub>32</sub>), -B(Q<sub>31</sub>)(Q<sub>32</sub>), -C(=O)(Q<sub>31</sub>), -S(=O)<sub>2</sub>(Q<sub>31</sub>), and -P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),

wherein Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> may each independently be selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group.

The term "Ph", as used herein, may refer to a phenyl group; the term "Me", as used herein, may refer to a methyl group; the term "Et", as used herein, may refer to an ethyl group; the terms "ter-Bu" or "But", as used herein, may refer

to a tert-butyl group; and the term "OMe" as used herein may refer to a methoxy group.

The term "biphenyl group" as used herein refers to "a phenyl group substituted with a phenyl group." In other words, the "biphenyl group" is a substituted phenyl group having a phenyl group as a substituent.


The term "terphenyl group" as used herein refers to "a phenyl group substituted with a biphenyl group." In other words, the "terphenyl group" is a phenyl group having, as a substituent, a phenyl group substituted with a phenyl group.

\* and \* used herein, unless defined otherwise, each refer to a binding site to a neighboring atom in a corresponding formula.


Hereinafter, a compound according to embodiments of the present disclosure and an organic light-emitting device according to embodiments of the present disclosure will be described in more detail with reference to Synthesis Examples and Examples. The wording "B was used instead of A" used in describing Synthesis Examples indicates that an identical molar equivalent of B was used in place of A.

### Synthesis Example

#### Synthesis Example 1: Synthesis of Compound 2



163



## (1) Synthesis of Intermediate 2-3

10 g of methyl 5-bromo-2-iodobenzoate (2-1) and 5 g of naphthalene-1-boronic acid (2-2) were dissolved in 100 mL of tetrahydrofuran (THF) and 25 mL of water, and 1 g of  $\text{Pd}(\text{PPh}_3)_4$  and 12 g of  $\text{K}_2\text{CO}_3$  were added dropwise thereto. The reaction solution was stirred under reflux at a temperature of 65° C. for 6 hours. After the reaction solution was cooled to room temperature, the THF was removed under reduced pressure, and the mixture was extracted three times using ethyl acetate. The organic solution was dried over anhydrous magnesium sulfate and filtered under reduced pressure. The residue obtained therefrom was purified by column chromatography to obtain 8.3 g (yield: 83%) of Intermediate 2-3.

$\text{C}_{18}\text{H}_{13}\text{BrO}_2$  M+1: 341.01.

## (2) Synthesis of Intermediate 2-4

8.3 g of Intermediate 2-3 was dissolved in 120 mL of THF, and 60 mL of  $\text{PhMgBr}$  (1M) was slowly added dropwise thereto at a temperature of 0° C. The mixture was slowly allowed to warm to room temperature and stirred for 16 hours. The reaction was then cooled to temperature of 0° C. and terminated by addition of a saturated ammonium chloride aqueous solution. The reaction mixture was extracted three times using ethyl acetate. The organic solution was dried over anhydrous magnesium sulfate and filtered under reduced pressure. Then, the residue obtained therefrom was purified by column chromatography to obtain 7.7 g (yield: 68%) of Intermediate 2-4.

$\text{C}_{29}\text{H}_{21}\text{BrO}$  M+1: 465.08.

## (3) Synthesis of Intermediate 2-5

7.7 g of Intermediate 2-4 was dissolved in 80 mL of dichloromethane, and 0.5 mL of methanesulfonic acid was added thereto and stirred for 3 hours. The reaction was

164

terminated by addition of 1 mL of triethylamine, and filtered under reduced pressure. Then, the residue obtained therefrom was purified by column chromatography to obtain 7.0 g (yield: 95%) of Intermediate 2-5.

$\text{C}_{29}\text{H}_{19}\text{Br}$  M+1: 447.07.


## (4) Synthesis of Compound 2

3 g of Intermediate 2-5, 3 g of N-phenyl-[1,1'-biphenyl]-4-amine, 0.34 g of  $\text{Pd}_2(\text{dba})_3$ , 0.1 mL of  $\text{PtBu}_3$ , and 3.6 g of  $\text{KOtBu}$  were dissolved in 60 mL of toluene and stirred at a temperature of 85° C. for 2 hours. The reaction solution was cooled to room temperature, and the reaction was terminated by water. The reaction mixture was extracted three times using ethyl acetate. The organic solution was dried over anhydrous magnesium sulfate and distilled under reduced pressure. Then, the residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 3.7 g (yield: 90%) of Compound 2.

$\text{C}_{47}\text{H}_{33}\text{N}$  M+1: 612.26.

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$ =7.72-7.50 (m, 6H), 7.48-7.38 (m, 5H), 7.25-6.94 (m, 16H), 6.77 (td, 1H), 6.55 (dt, 1H), 6.09 (m, 1H), 6.05 (m, 2H), 6.00 (dd, 1H).

## Synthesis Example 2: Synthesis of Compound 10





3.2 g (yield: 78%) of Compound 10 was obtained in substantially the same manner as in the Synthesis of Compound 2, except that 3 g of N-phenyl-[1,1'-biphenyl]-2-amine was used instead of N-phenyl-[1,1'-biphenyl]-4-amine.

$\text{C}_{47}\text{H}_{33}\text{N}$  M+1: 612.26.

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$ =7.72-7.50 (m, 6H), 7.48-7.43 (m, 4H), 7.32 (dt, 1H), 7.25-6.94 (m, 16H), 6.77 (td, 1H), 6.55 (dt, 1H), 6.09 (m, 1H), 6.05 (m, 2H), 6.00 (dd, 1H).

**165**

Synthesis Example 3: Synthesis of Compound 11





3.4 g (yield: 81%) of Compound 11 was obtained in substantially the same manner as in the Synthesis of Compound 2, except that 3 g of N-phenyldibenzo[b,d]furan-1-amine was used instead of N-phenyl-[1,1'-biphenyl]-4-amine.

$C_{47}H_{31}NO$  M+1: 626.24.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.79 (m, 1H), 7.72-7.60 (m, 5H), 7.49-7.44 (m, 2H), 7.33-7.25 (m, 4H), 7.22-7.16 (m, 4H), 7.15-6.99 (m, 9H), 6.67 (dd, 1H), 6.61 (dt, 1H), 6.63 (dd, 1H), 6.26 (m, 3H).

Synthesis Example 4: Synthesis of Compound 16



**166**  
-continued

3.0 g (yield: 69%) of Compound 16 was obtained in substantially the same manner as in the Synthesis of Compound 2, except that 3 g of 9,9-dimethyl-N-phenyl-9H-fluorene-2-amine (16-1) was used instead of N-phenyl-[1,1'-biphenyl]-4-amine, and 3 g of Intermediate 16-2 was used instead of Intermediate 2-5.

$C_{50}H_{37}N$  M+1: 651.29.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.81 (dd, 1H), 7.75 (dd, 1H), 7.62 (s, 1H), 7.55-7.52 (m, 2H), 7.47-7.41 (m, 2H), 7.36-7.25 (m, 2H), 7.23-6.93 (m, 16H), 6.82 (td, 1H), 6.65-6.60 (m, 2H), 6.36 (d, 1H), 6.11 (m, 2H), 1.61 (s, 6H).

Synthesis Example 5: Synthesis of Compound 21



2.7 g (yield: 60%) of Compound 21 was obtained in substantially the same manner as in the Synthesis of Com-


## 167

pond 2, except that 3 g of N-([1,1'-biphenyl]-4-yl)naphthalen-2-amine was used instead of N-phenyl-[1,1'-biphenyl]-4-amine, and 3 g of Intermediate 16-2 was used instead of Intermediate 2-5.

$C_{51}H_{35}N$  M+1: 662.28.


$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.81-7.75 (m, 2H), 7.70 (m, 1H), 7.66-7.60 (m, 3H), 7.59-7.34 (m, 13H), 7.28-7.05 (m, 12H), 6.99 (dt, 1H), 6.84-6.74 (m, 4H).

## Synthesis Example 6: Synthesis of Compound 34



## 168

-continued



## (1) Synthesis of Intermediate 34-2

8 g (yield: 85%) of Intermediate 34-2 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-3, except that 5 g of (4-fluoronaphthalen-1-yl)boronic acid (34-1) was used instead of naphthalene-1-boronic acid.

$C_{18}H_{12}BrFO_2$  M+1: 359.00.

## (2) Synthesis of Intermediate 34-3

8 g (yield: 74%) of Intermediate 34-3 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-4, except that 8 g of Intermediate 34-2 was used instead of Intermediate 2-3.

$C_{29}H_{20}BrFO$  M+1: 483.07.

## (3) Synthesis of Intermediate 34-4

6.5 g (yield: 84%) of Intermediate 34-4 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-5, except that 8 g of Intermediate 34-3 was used instead of Intermediate 2-4.

$C_{29}H_{18}BrF$  M+1: 465.06.


## (4) Synthesis of Compound 34

3.4 g (yield: 84%) of Compound 34 was obtained in substantially the same manner as in the Synthesis of Compound 10, except that 3 g of Intermediate 34-4 was used instead of Intermediate 2-5.

$C_{47}H_{32}FN$  M+1: 630.25.





$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.76-7.70 (m, 2H), 7.62-7.53 (m, 3H), 7.50-7.44 (m, 3H), 7.37-7.30 (m, 5H), 7.25-6.95 (m, 13H), 6.77 (dd, 1H), 6.65 (dt, 1H), 6.18 (d, 1H), 6.06 (m, 3H).

## Synthesis Example 7: Synthesis of Compound 38



169

-continued



## (1) Synthesis of Intermediate 38-3

12 g (yield: 80%) of Intermediate 38-3 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-3, except that 10 g of (4-bromonaphthalen-1-yl) boronic acid (38-2) was used instead of naphthalene-1-boronic acid.

$C_{18}H_{12}BrFO_2$  M+1: 359.00.

## (2) Synthesis of Intermediate 38-4

14 g (yield: 87%) of Intermediate 38-4 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-4, except that 12 g of Intermediate 38-3 was used instead of Intermediate 2-3.

$C_{29}H_{20}BrFO$  M+1: 483.07.

## (3) Synthesis of Intermediate 38-5

8 g (yield: 59%) of Intermediate 38-5 was obtained in substantially the same manner as in the Synthesis of Inter

170

mediate 2-5, except that 14 g of Intermediate 38-4 was used instead of Intermediate 2-4.

$C_{29}H_{18}BrF$  M+1: 465.06.





## (4) Synthesis of Compound 38

3.1 g (yield: 60%) of Compound 38 was obtained in substantially the same manner as in the Synthesis of Compound 2, except that 3 g of Intermediate 38-5 was used instead of Intermediate 2-5.

$C_{47}H_{32}FN$  M+1: 630.25.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.77 (m, 1H), 7.65-7.60 (m, 2H), 7.55-7.28 (m, 11H), 7.22-6.94 (m, 12H), 6.78 (m, 2H), 6.65 (dt, 1H), 6.43 (dd, 1H), 6.12 (m, 2H).

## Synthesis Example 8: Synthesis of Compound 54



65

## 171

## (1) Synthesis of Intermediate 54-1

5 g of 1-bromo-4-methylbenzene was dissolved in THF and cooled to a temperature of  $-78^{\circ}\text{C}$ . 12.3 mL of n-BuLi (2.5M in hexane) was slowly added dropwise thereto while maintaining the temperature. After 1 hour, 5 g of Intermediate 2-3 dissolved in THF was slowly added dropwise to the reaction vessel. When the reaction was complete, the temperature was slowly raised to room temperature, and the reaction was terminated by addition of a saturated ammonium chloride solution. The reaction mixture was extracted three times using ethyl acetate. The organic solution was dried over anhydrous magnesium sulfate and distilled under reduced pressure. The residue obtained therefrom was separated and purified by silica gel column chromatography to obtain 4.8 g (yield: 66%) of Intermediate 54-1.

$\text{C}_{31}\text{H}_{25}\text{BrO}$  M+1: 493.11.

## (2) Synthesis of Intermediate 54-2

3.6 g (yield: 75%) of Intermediate 54-2 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-5, except that 4.8 g of Intermediate 54-1 was used.

$\text{C}_{31}\text{H}_{23}\text{Br}$  M+1: 475.10


## (3) Synthesis of Compound 54

3.8 g (yield: 79%) of Compound 54 was obtained in substantially the same manner as in the Synthesis of Compound 2, except that 3 g of N-phenyldibenzo[b,d]furan-2-amine was used instead of N-phenyl-[1,1'-biphenyl]-4-amine, and 3.5 g of Intermediate 54-2 was used instead of Intermediate 2-5.

$\text{C}_{49}\text{H}_{35}\text{NO}$  M+1: 654.27.



$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$ =7.84 (m, 1H), 7.74-7.65 (m, 4H), 7.63 (m, 1H), 7.55-7.46 (m, 4H), 7.44 (m, 1H), 7.32 (dt, 1H), 7.13-6.98 (m, 7H), 6.95 (dd, 1H), 6.86 (m, 4H), 6.67 (dt, 1H), 6.42 (d, 1H), 6.30 (m, 3H), 2.34 (s, 3H), 2.32 (s, 3H).

## Synthesis Example 9: Synthesis of Compound 64



## 172

-continued



## (1) Synthesis of Intermediate 64-1

6.6 g (yield: 90%) of Intermediate 64-1 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-4, except that 5.5 g of 1-bromo-4-fluorobenzene was used.

$\text{C}_{29}\text{H}_{19}\text{BrF}_2\text{O}$  M+1: 501.06.

## (2) Synthesis of Intermediate 64-2

5.3 g (83%) of Intermediate 64-2 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-5, except that 6.6 g of Intermediate 64-1 was used instead of Intermediate 2-4.

$\text{C}_{29}\text{H}_{17}\text{BrF}_2$  M+1: 484.05.

## (3) Synthesis of Compound 64



3.4 g (yield: 80%) of Compound 64 was obtained in substantially the same manner as in the Synthesis of Compound 2, except that 3 g of 9,9-dimethyl-N-phenyl-9H-fluoren-2-amine was used instead of N-phenyl-[1,1'-diphenyl]-4-amine, and 3 g of Intermediate 64-2 was used instead of Intermediate 2-5.

$\text{C}_{50}\text{H}_{35}\text{F}_2\text{N}$  M+1: 688.27.

$^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$ =7.78 (td, 1H), 7.72-7.64 (m, 3H), 7.63 (m, 1H), 7.58 (d, 1H), 7.50 (d, 1H), 7.36-7.27 (m, 2H), 7.14-7.00 (m, 9H), 6.72 (dd, 1H), 6.66 (td, 1H), 6.51-6.46 (m, 4H), 6.42 (dd, 1H), 6.33 (dd, 2H), 6.22 (m, 2H), 1.62 (s, 6H).

## 173

## Synthesis Example 10: Synthesis of Compound 81




3.7 g (yield: 74%) of Compound 81 was obtained in substantially the same manner as in the Synthesis of Compound 21, except that 3 g of Intermediate 81-1 was used instead of Intermediate 16-2.

$C_{51}H_{35}N$  M+1: 662.28.


$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.78-7.62 (m, 7H), 7.57-7.32 (m, 11H), 7.31 (dt, 1H), 7.18-7.01 (m, 8H), 6.96 (t, 1H), 6.78-6.75 (m, 4H), 6.41 (m, 2H), 5.71 (dd, 1H).

## Synthesis Example 11: Synthesis of Compound 95



## 174

-continued





2.5 g (yield: 86%) of Compound 95 was obtained in substantially the same manner as in the Synthesis of Compound 11, except that 2.5 g of Intermediate 95-1 was used instead of Intermediate 2-5.

$C_{47}H_{31}NO$  M+1: 626.24.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.79-7.66 (m, 5H), 7.49-7.44 (m, 2H), 7.36-7.27 (m, 4H), 7.21-7.19 (m, 4H), 7.14-7.01 (m, 10H), 6.69-6.60 (m, 2H), 6.39 (dd, 1H), 6.34 (m, 2H), 6.05 (d, 1H).

## Synthesis Example 12: Synthesis of Compound 98



2.9 g (yield: 85%) of Compound 98 was obtained in substantially the same manner as in the Synthesis of Com-





## 175

pond 2, except that 2.5 g of N-phenyl-[1,1'-biphenyl]-4-amine was used, and 2.5 g of Intermediate 98-1 was used instead of 3 g of Intermediate 2-5.

$C_{47}H_{33}N$  M+1: 612.26.


$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.92 (d, 1H), 7.67-7.58 (m, 4H), 7.54-7.30 (m, 7H), 7.26-7.01 (m, 13H), 6.94 (dd, 1H), 6.83 (m, 2H), 6.65 (dt, 1H), 6.57 (dd, 1H), 6.22 (m, 2H), 5.58 (t, 1H).

## Synthesis Example 13: Synthesis of Compound 110



## 176

-continued



## (1) Synthesis of Intermediate 110-2

8.8 g (yield: 68%) of Intermediate 110-2 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-3, except that 8.4 g of methyl 2-bromobenzoate (110-1) was used instead of methyl 5-bromo-2-iodobenzoate, and 10 g of (5-bromonaphthalen-1-yl)boronic acid was used instead of naphthalene-1-boronic acid.

$C_{18}H_{13}BrO_2$ : M+1 341.01

## (2) Synthesis of Intermediate 110-3

8.5 g (yield: 71%) of Intermediate 110-3 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-4, except that 8.8 g of Intermediate 110-2 was used instead of Intermediate 2-3.

$C_{29}H_{21}BrO$ : M+1 465.08.

## (3) Synthesis of Intermediate 110-4

6.6 g (yield: 81%) of Intermediate 110-4 was obtained in substantially the same manner as in the Synthesis of Intermediate 2-5, except that 8.5 g of Intermediate 110-3 was used instead of Intermediate 2-4.

$C_{29}H_{19}Br$ : M+1 447.07.


## (4) Synthesis of Compound 110



3.1 g (yield: 71%) of Compound 110 was obtained in substantially the same manner as in the Synthesis of Compound 16, except that 2.8 g of Intermediate 110-4 was used instead of Intermediate 16-2.

$C_{50}H_{37}N$ : M+1 652.29.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.81 (td, 1H), 7.76 (td, 1H), 7.63 (d, 1H), 7.55 (td, 1H), 7.51 (dd, 1H), 7.45 (dt, 1H), 7.36-7.25 (m, 2H), 7.22-7.01 (m, 15H), 6.86 (td, 1H), 6.81 (d, 1H), 6.65-6.59 (m, 2H), 6.37 (dd, 1H), 6.31 (m, 1H), 6.12-6.08 (m, 2H), 1.61 (s, 6H).

## Synthesis Example 14: Synthesis of Compound 124







2.7 g (yield: 75%) of Compound 124 was obtained in substantially the same manner as in the Synthesis of Compound 16, except that 2.6 g of Intermediate 124-1 was used instead of Intermediate 16-2.

$C_{56}H_{41}N$ : M+1 728.32.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =8.15 (d, 1H), 8.07-8.03 (m, 2H), 7.78 (d, 1H), 7.70 (d, 1H), 7.60-7.55 (m, 2H), 7.51-7.43 (m, 2H), 7.41-7.28 (m, 4H), 7.25-7.18 (m, 4H), 7.13-7.02 (m, 10H), 6.88 (d, 1H), 6.72-6.70 (m, 2H), 6.42 (m, 2H), 6.29 (d, 1H), 6.22 (m, 2H), 1.62 (s, 6H).

#### Synthesis Example 15: Synthesis of Compound 136



2.3 g (yield: 69%) of Compound 136 was obtained in substantially the same manner as in the Synthesis of Compound 16, except that 2.3 g of Intermediate 136-1 was used instead of Intermediate 16-2.

$C_{50}H_{36}FN$ : M+1 670.28.

$^1H$  NMR (500 MHz,  $CDCl_3$ )  $\delta$ =7.78 (td, 1H), 7.74 (dd, 1H), 7.58-7.52 (m, 2H), 7.48-7.40 (m, 2H), 7.36-7.28 (m, 1H), 7.25-7.20 (m, 4H), 7.18-7.05 (m, 11H), 6.96 (d, 1H), 6.72 (dd, 1H), 6.66 (t, 1H), 6.42-6.39 (m, 2H), 6.29 (d, 1H), 6.22 (m, 2H), 1.62 (s, 6H).

#### EXAMPLES

##### Comparative Example 1

A Corning 15  $\Omega/cm^2$  (1,200 Å) ITO glass substrate was cut to a size of 50 mm×50 mm×0.7 mm, sonicated with isopropyl alcohol and pure water each for 5 minutes, and then cleaned by exposure to ultraviolet rays and ozone for 30 minutes. Then, the ITO glass substrate was provided to a vacuum deposition apparatus for use as an anode.


2-TNATA was vacuum-deposited on the ITO glass substrate to form a hole injection layer having a thickness of 600 Å, and 4,4'-bis[N-(1-naphthyl)-N-phenylamino]biphenyl (NPB) (a hole transport material) was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 300 Å.

9,10-di(naphthalen-2-yl)anthracene (DNA) (a blue fluorescent host) and 4,4'-bis[2-(4-(N,N-diphenylamino)phenyl)vinyl]biphenyl (DPAVBi) (a blue fluorescent dopant) was co-deposited on the hole transport layer at a weight ratio of 98:2 to form an emission layer having a thickness of 300 Å.


Then,  $Alq_3$  was deposited on the emission layer to form an electron transport layer having a thickness of 300 Å. LiF (an alkali metal halide) was deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å. Al was vacuum-deposited on the electron injection layer to form a cathode electrode having a thickness of 3,000 Å, thereby forming a LiF/Al electrode. In this manner, an organic light-emitting device was manufactured.

179


180




2-TNATA



NPB



DPAVBI




DNA

181

Comparative Example 2

An organic light-emitting device was manufactured in substantially the same manner as Comparative Example 1, except that Compound A was used instead of NPB in forming a hole transport layer.


Compound A



Comparative Example 3

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound B was used instead of NPB in forming a hole transport layer.

Compound B




Comparative Example 4

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound C was used instead of NPB in forming a hole transport layer.

182


Compound C



Comparative Example 5

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound D was used instead of NPB in forming a hole transport layer.


Compound D



Comparative Example 6

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound E was used instead of NPB in forming a hole transport layer.


Compound E



## 183


## Comparative Example 7

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound F was used instead of NPB in forming a hole transport layer.  
Compound F



## Comparative Example 8

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound G was used instead of NPB in forming a hole transport layer.  
Compound G



## Example 1

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 2 was used instead of NPB in forming a hole transport layer.

## Example 2

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 10 was used instead of NPB in forming a hole transport layer.

## Example 3

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example

## 184

1, except that Compound 11 was used instead of NPB in forming a hole transport layer.

## Example 4

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 16 was used instead of NPB in forming a hole transport layer.

## Example 5

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 21 was used instead of NPB in forming a hole transport layer.

## Example 6

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 34 was used instead of NPB in forming a hole transport layer.

## Example 7

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 38 was used instead of NPB in forming a hole transport layer.

## Example 8

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 54 was used instead of NPB in forming a hole transport layer.

## Example 9

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 64 was used instead of NPB in forming a hole transport layer.

## Example 10

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 81 was used instead of NPB in forming a hole transport layer.

## Example 11

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 95 was used instead of NPB in forming a hole transport layer.

## Example 12

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 98 was used instead of NPB in forming a hole transport layer.

## Example 13

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example

1, except that Compound 110 was used instead of NPB in forming a hole transport layer.

Example 14

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 124 was used instead of NPB in forming a hole transport layer.

Example 15

An organic light-emitting device was manufactured in substantially the same manner as in Comparative Example 1, except that Compound 136 was used instead of NPB in forming a hole transport layer.

The device performance (driving voltage, luminance, efficiency) at a current density of 50 mA/cm<sup>2</sup> and the half lifespan at a current density of 100 mA/cm<sup>2</sup> of the organic light-emitting devices manufactured according to Comparative Examples 1 to 8 and Examples 1 to 15 were measured using the following methods, and results thereof are shown in Table 1.

Luminance: A current-voltage meter (Keithley SMU 236) supplied power, and luminance was measured by using a luminance meter PR650.

Efficiency: A current-voltage meter (Keithley SMU 236) supplied power, and efficiency was measured by using a luminance meter PR650.

The half lifespan indicates an amount of time (hr) that lapsed when luminance was 50% of initial luminance (100%) (at 100 mA/cm<sup>2</sup>).

Referring to Table 1, when the compound according to one or more embodiments of the present disclosure was used as a hole transport material for an organic light-emitting device, the organic light-emitting devices of Examples 1-15 exhibited excellent I-V-L characteristics including a low driving voltage and improved efficiency, as compared with the organic light-emitting device of Comparative Example 1, which used NPB. In particular, the lifespan improvement effect was excellent and the lifespan was remarkably improved in Examples 1-15, which used the compound according to one or more embodiments of the present disclosure as a hole transport material. The driving voltage was improved and the efficiency and half lifespan were also excellent compared with Comparative Examples 2 to 8, which use Compounds A to G.

It should be understood that the embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as being available for other similar features or aspects in other embodiments.

As used herein, expressions such as “at least one of”, “one of”, “selected from”, “at least one selected from”, and “one selected from”, when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list. Further, the use of “may” when describing embodiments of the present disclosure refers to “one or more embodiments of the present disclosure”.

As used herein, the terms “use”, “using”, and “used” may be considered synonymous with the terms “utilize”, “utilizing”, and “utilized”, respectively. Further, the use of “may”

TABLE 1

	Hole transport material	Driving voltage (V)	Current density (mA/cm <sup>2</sup> )	Luminance (cd/m <sup>2</sup> )	Efficiency (cd/A)	Emission color	Half lifespan (hr @100 mA/cm <sup>2</sup> )
Comparative Example 1	NPB	7.01	50	2645	5.29	Blue	258
Comparative Example 2	Compound A	6.31	50	2775	5.55	Blue	283
Comparative Example 3	Compound B	6.21	50	2810	5.62	Blue	299
Comparative Example 4	Compound C	6.11	50	2755	5.51	Blue	305
Comparative Example 5	<Compound D>	6.42	50	2775	5.55	Blue	220
Comparative Example 6	<Compound E>	6.89	50	2330	4.66	Blue	210
Comparative Example 7	Compound F	6.60	50	2615	5.23	Blue	243
Comparative Example 8	Compound G	6.76	50	2310	4.62	Blue	200
Example 1	Compound 2	4.89	50	3410	6.82	Blue	412
Example 2	Compound 10	4.87	50	3435	6.87	Blue	406
Example 3	Compound 11	4.64	50	3385	6.77	Blue	419
Example 4	Compound 16	4.56	50	3460	6.92	Blue	423
Example 5	Compound 21	4.67	50	3475	6.95	Blue	410
Example 6	Compound 34	4.72	50	3440	6.88	Blue	408
Example 7	Compound 38	4.79	50	3280	6.56	Blue	420
Example 8	Compound 54	4.81	50	3360	6.72	Blue	392
Example 9	Compound 64	4.92	50	3440	6.88	Blue	381
Example 10	Compound 81	4.99	50	3460	6.92	Blue	399
Example 11	Compound 95	4.87	50	3265	6.53	Blue	378
Example 12	Compound 98	4.83	50	3265	6.53	Blue	388
Example 13	Compound 110	4.89	50	3235	6.47	Blue	391
Example 14	Compound 124	4.82	50	3280	6.56	Blue	368
Example 15	Compound 136	4.87	50	3115	6.23	Blue	368

when describing embodiments of the present disclosure refers to “one or more embodiments of the present disclosure”.


As used herein, the terms “substantially”, “about”, and similar terms are used as terms of approximation and not as terms of degree, and are intended to account for the inherent deviations in measured or calculated values that would be recognized by those of ordinary skill in the art.

Also, any numerical range recited herein is intended to include all subranges of the same numerical precision subsumed within the recited range. For example, a range of “1.0 to 10.0” is intended to include all subranges between (and including) the recited minimum value of 1.0 and the recited maximum value of 10.0, that is, having a minimum value equal to or greater than 1.0 and a maximum value equal to or less than 10.0, such as, for example, 2.4 to 7.6. Any maximum numerical limitation recited herein is intended to include all lower numerical limitations subsumed therein and any minimum numerical limitation recited in this specification is intended to include all higher numerical limitations subsumed therein. Accordingly, Applicant reserves the right to amend this specification, including the claims, to expressly recite any sub-range subsumed within the ranges expressly recited herein.


While one or more embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims and equivalents thereof.

What is claimed is:

1. An amine-based compound represented by Formula 1:



Formula 1



Formula 2

wherein, in Formulae 1 and 2,

Ar<sub>1</sub>, Ar<sub>2</sub>, L<sub>1</sub>, and L<sub>2</sub> are each independently a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

a<sub>1</sub> and a<sub>2</sub> are each independently an integer from 0 to 5, wherein, when a<sub>1</sub> is two or more, two or more L<sub>1</sub> (s) are identical to or different from each other, and when a<sub>2</sub> is two or more, two or more L<sub>2</sub> (s) are identical to or different from each other,

when a<sub>1</sub> is zero, \*(L<sub>1</sub>)<sub>a1</sub>-\* is a single bond, and when a<sub>2</sub> is zero, \*(L<sub>2</sub>)<sub>a2</sub>-\* is a single bond,

R<sub>1</sub> to R<sub>10</sub> are each independently selected from a group represented by Formula 2, hydrogen, deuterium, —F,

—Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

provided that one selected from R<sub>5</sub> to R<sub>8</sub> is a group represented by Formula 2,

R<sub>11</sub> and R<sub>12</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

b<sub>1</sub> and b<sub>2</sub> are each independently an integer from 1 to 10, wherein, when b<sub>1</sub> is two or more, two or more R<sub>11</sub> (s) are identical to or different from each other, and when b<sub>2</sub> is two or more, two or more R<sub>12</sub> (s) are identical to or different from each other,

at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:


deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a

- hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;
- a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);
- a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;
- a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and
- Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>),
- Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and
- \* and \*' each indicate a binding site to a neighboring atom.
2. The amine-based compound of claim 1, wherein Ar<sub>1</sub>, Ar<sub>2</sub>, L<sub>1</sub>, and L<sub>2</sub> are each independently selected from:
- a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, a heptalene group, an indacene group, an acenaphthalene group, a


- fluorene group, a spiro-fluorene group, a benzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, a hexacene group, a pyrrole group, an imidazole group, a pyrazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an iso-indole group, an indole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a carbazole group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzoxazole group, a benzimidazole group, a furan group, a benzofuran group, a thiophene group, a benzothiophene group, a thiazole group, an isothiazole group, a benzothiazole group, an isoxazole group, an oxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a benzoxazole group, a dibenzofuran group, a dibenzothiophene group, a benzocarbazole group, and a dibenzocarbazole group; and
- a benzene group, a pentalene group, an indene group, a naphthalene group, an azulene group, a heptalene group, an indacene group, an acenaphthalene group, a fluorene group, a spiro-fluorene group, a benzofluorene group, a phenalene group, a phenanthrene group, an anthracene group, a fluoranthene group, a triphenylene group, a pyrene group, a chrysene group, a naphthacene group, a picene group, a perylene group, a pentaphene group, a hexacene group, a pyrrole group, an imidazole group, a pyrazole group, a pyridine group, a pyrazine group, a pyrimidine group, a pyridazine group, an iso-indole group, an indole group, an indazole group, a purine group, a quinoline group, an isoquinoline group, a benzoquinoline group, a phthalazine group, a naphthyridine group, a quinoxaline group, a quinazoline group, a cinnoline group, a carbazole group, a phenanthridine group, an acridine group, a phenanthroline group, a phenazine group, a benzoxazole group, a benzimidazole group, a furan group, a benzofuran group, a thiophene group, a benzothiophene group, a thiazole group, an isothiazole group, a benzothiazole group, an isoxazole group, an oxazole group, a triazole group, a tetrazole group, an oxadiazole group, a triazine group, a benzoxazole group, a dibenzofuran group, a dibenzothiophene group, a benzocarbazole group, and a dibenzocarbazole group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>2</sub>-C<sub>20</sub> heteroaryl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>).
3. The amine-based compound of claim 1, wherein Ar<sub>1</sub> and Ar<sub>2</sub> are each independently selected from:
- a benzene group; and
- a benzene group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>2</sub>-C<sub>20</sub> heteroaryl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>).

191


4. The amine-based compound of claim 1, wherein  $L_1$  and  $L_2$  are each independently selected from groups represented by any of Formulae 3-1 to 3-46:




Formula 3-1




Formula 3-2




Formula 3-3




Formula 3-4




Formula 3-5



Formula 3-6





Formula 3-7













Formula 3-8

192

-continued




**193**  
-continued



**194**  
-continued


Formula 3-21




Formula 3-22




Formula 3-23




Formula 3-24




Formula 3-25




Formula 3-26




Formula 3-27




Formula 3-28




Formula 3-29




Formula 3-30




Formula 3-31




Formula 3-32




Formula 3-33




Formula 3-34




Formula 3-35




Formula 3-36




Formula 3-37




Formula 3-38




Formula 3-39




Formula 3-40




Formula 3-41




Formula 3-42




Formula 3-43




Formula 3-44




Formula 3-45




Formula 3-46




Formula 3-47




Formula 3-48




Formula 3-49




Formula 3-50




Formula 3-51




Formula 3-52




Formula 3-53




Formula 3-54




Formula 3-55




Formula 3-56




Formula 3-57




Formula 3-58




Formula 3-59




Formula 3-60




Formula 3-61




Formula 3-62




Formula 3-63




Formula 3-64




Formula 3-65




Formula 3-66




Formula 3-67




Formula 3-68




Formula 3-69




Formula 3-70




Formula 3-71




Formula 3-72




Formula 3-73




Formula 3-74




Formula 3-75




Formula 3-76




Formula 3-77




Formula 3-78




Formula 3-79




Formula 3-80




Formula 3-81




Formula 3-82




Formula 3-83




Formula 3-84




Formula 3-85




Formula 3-86




Formula 3-87




Formula 3-88




Formula 3-89



Formula 3-90








Formula 3-91



195


-continued




196

-continued


Formula 3-38




Formula 3-31



Formula 3-39



Formula 3-40



Formula 3-41

45

Formula 3-42

65

Formula 3-43

Formula 3-44

Formula 3-45

Formula 3-46

wherein, in Formulae 3-1 to 3-46,

$Y_1$  is O, S,  $C(Z_3)(Z_4)$ ,  $N(Z_5)$ , or  $Si(Z_6)(Z_7)$ ,

$Z_1$  to  $Z_7$  are each independently selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoanthrenyl group, a triphenylenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, and  $-Si(Q_{31})(Q_{32})(Q_{33})$ ,

$d_2$  is an integer from 0 to 2, wherein, when  $d_2$  is two or more, two or more  $Z_1$ (s) are identical to or different from each other,

$d_3$  is an integer from 0 to 3, wherein, when  $d_3$  is two or more, two or more  $Z_1$ (s) are identical to or different from each other and two or more  $Z_2$ (s) are identical to or different from each other,

$d_4$  is an integer from 0 to 4, wherein, when  $d_4$  is two or more, two or more  $Z_1$ (s) are identical to or different from each other,

197

d5 is an integer from 0 to 5, wherein, when d5 is two or more, two or more  $Z_1(s)$  are identical to or different from each other and two or more  $Z_2(s)$  are identical to or different from each other,

d6 is an integer from 0 to 6, wherein, when d6 is two or more, two or more  $Z_1(s)$  are identical to or different from each other and two or more  $Z_2(s)$  are identical to or different from each other,

d8 is an integer from 0 to 8, wherein, when d8 is two or more, two or more  $Z_1(s)$  are identical to or different from each other, and

\* and \*' each indicate a binding site to a neighboring atom.

5. The amine-based compound of claim 1, wherein:


$R_1$  to  $R_{10}$  are each independently selected from a group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and —Si( $Q_1$ )( $Q_2$ )( $Q_3$ ), and

$R_{11}$  and  $R_{12}$  are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, a substituted or unsubstituted  $C_6$ - $C_{60}$  aryl group, a substituted or unsubstituted  $C_1$ - $C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, and —Si( $Q_1$ )( $Q_2$ )( $Q_3$ ).


6. The amine-based compound of claim 1, wherein:

$R_1$  to  $R_{10}$  are each independently selected from a group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, groups represented by Formulae 5-1 to 5-80, and —Si( $Q_1$ )( $Q_2$ )( $Q_3$ ), and


$R_{11}$  and  $R_{12}$  are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1$ - $C_{60}$  alkyl group, groups represented by any of Formulae 5-1 to 5-80, and —Si( $Q_1$ )( $Q_2$ )( $Q_3$ ):



Formula 5-1



Formula 5-2




Formula 5-3


198

-continued


Formula 5-4




Formula 5-5




Formula 5-6




Formula 5-7




Formula 5-8




Formula 5-9




Formula 5-10














**199**  
-continued



**200**  
-continued




**201**  
-continued



**202**  
-continued

Formula 5-29


5



Formula 5-40

Formula 5-30


10



Formula 5-41

Formula 5-31


15



Formula 5-42

Formula 5-32


20



Formula 5-43

Formula 5-33


25



Formula 5-44

Formula 5-34


30



Formula 5-45

Formula 5-35


35



Formula 5-46

Formula 5-36


40



Formula 5-47

Formula 5-37


45



Formula 5-48

Formula 5-38


50



Formula 5-49

Formula 5-39


55



Formula 5-50

Formula 5-40


60



Formula 5-51

Formula 5-41


65



Formula 5-52

Formula 5-42


70



Formula 5-53


203

-continued




Formula 5-54

5




Formula 5-55

10




Formula 5-56

15




Formula 5-57

20




Formula 5-58

25




Formula 5-59

30




Formula 5-60

35




Formula 5-61

40




Formula 5-62

45



Formula 5-63

50




Formula 5-64


55

204


-continued




Formula 5-65




Formula 5-66




Formula 5-67




Formula 5-68




Formula 5-69




Formula 5-70




Formula 5-71




Formula 5-72



Formula 5-73








Formula 5-74



Formula 5-75

205

-continued



wherein, in Formulae 5-1 to 5-80,

$Y_{31}$  is O, S, C( $Z_{33}$ )( $Z_{34}$ ), N( $Z_{35}$ ), or Si( $Z_{36}$ )( $Z_{37}$ ),

$Z_{31}$  to  $Z_{37}$  are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazone group, a  $C_1$ - $C_{20}$  alkyl group, a  $C_1$ - $C_{20}$  alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclopentenyl group, a cyclohexenyl group, a phenyl group, a biphenyl group, a naphthyl group, a fluorenyl group, a spiro-bifluorenyl group, a spiro-fluorene-benzofluorenyl group, a benzofluorenyl group, a dibenzofluorenyl group, a phenalenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a naphthyridinyl group, a quinoxalinyl group, a quinazolinyl group, a carbazolyl group, a phenanthridinyl group, an acridinyl group, a phenanthrolinyl group, a phenazinyl group, a triazinyl group, a dibenzofuranlyl group, a dibenzothiofenyl group, and —Si( $Q_{31}$ )( $Q_{32}$ )( $Q_{33}$ ).

e2 is an integer from 0 to 2, wherein, when e2 is two or more, two or more  $Z_{32}$ (s) are identical to or different from each other,

e3 is an integer from 0 to 3, wherein, when e3 is two or more, two or more  $Z_{31}$ (s) are identical to or different

206

from each other and two or more  $Z_{32}$ (s) are identical to or different from each other,

e4 is an integer from 0 to 4, wherein, when e4 is two or more, two or more  $Z_{31}$ (s) are identical to or different from each other and two or more  $Z_{32}$ (s) are identical to or different from each other,

e5 is an integer from 0 to 5, wherein, when e5 is two or more, two or more  $Z_{31}$ (s) are identical to or different from each other and two or more  $Z_{32}$ (s) are identical to or different from each other,

e6 is an integer from 0 to 6, wherein, when e6 is two or more, two or more  $Z_{31}$ (s) are identical to or different from each other and two or more  $Z_{32}$ (s) are identical to or different from each other,

e7 is an integer from 0 to 7, wherein, when e7 is two or more, two or more  $Z_{31}$ (s) are identical to or different from each other,

e9 is an integer from 0 to 9, wherein, when e9 is two or more, two or more  $Z_{31}$ (s) are identical to or different from each other, and

\* indicates a binding site to a neighboring atom.

7. The amine-based compound of claim 1, wherein  $R_{11}$  and  $R_{12}$  are each independently selected from groups represented by any of Formulae 6-1 to 6-34:

Formula 5-76

5

Formula 5-77

10

Formula 5-78

20

Formula 5-79

25

Formula 5-80

30

35

40

45


50

55


60

65


Formula 6-1




Formula 6-2




Formula 6-3




Formula 6-4




Formula 6-5











Formula 6-6



Formula 6-7




**207**  
-continued



**208**  
-continued

Formula 6-8

5




Formula 6-9

10

Formula 6-10


15

20



Formula 6-11


25



Formula 6-12


30

35



Formula 6-13


40



Formula 6-14

45

50






Formula 6-15

55

60

65



Formula 6-16

Formula 6-17

Formula 6-18

Formula 6-19

Formula 6-20

Formula 6-21










Formula 6-22

Formula 6-23

Formula 6-24

209


-continued



210

-continued

Formula 6-25



Formula 6-26

10

Formula 6-27

15

wherein, in Formulae 6-1 to 6-34,

Ph indicates a phenyl group, and

\* indicates a binding site to a neighboring atom.


**8.** The amine-based compound of claim 1, wherein the amine-based compound is represented by Formula 1-5:

Formula 6-28

20

Formula 6-29

25



Formula 6-30

35

Formula 6-31

40

wherein, in Formula 1-5,

Ar<sub>1</sub>, Ar<sub>2</sub>, L<sub>1</sub>, L<sub>2</sub>, a<sub>1</sub>, a<sub>2</sub>, R<sub>1</sub> to R<sub>12</sub>, b<sub>1</sub>, and b<sub>2</sub> are the same as described in connection with Formulae 1 and 2.

**9.** The amine-based compound of claim 8, wherein R<sub>1</sub> to R<sub>10</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a methyl group, an ethyl group, a propyl group, an iso-propyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a ter-butyl group, a pentyl group, an iso-amyl group, a hexyl group, a phenyl group, and a biphenyl group.

Formula 6-32

50

**10.** The amine-based compound of claim 8, wherein Ar<sub>1</sub> and Ar<sub>2</sub> are each independently selected from:

a benzene group; and

a benzene group substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>20</sub> alkyl group, a C<sub>1</sub>-C<sub>20</sub> alkoxy group, a C<sub>6</sub>-C<sub>20</sub> aryl group, a C<sub>2</sub>-C<sub>20</sub> heteroaryl group, and —Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>).

Formula 6-33

60


**11.** The amine-based compound of claim 1, wherein the amine-based compound is selected from any of Compounds 109-120:

65

Formula 6-34

Formula 1-5

211



109

212

-continued


113

5

10

110

15




20

25

111

30




35

40

45

112


50




55

60


65




114



115




116



213

-continued




117

5

10

15

118




20

25


119

30



35

40



120

45


50

55


12. An organic light-emitting device comprising:  
 a first electrode;  
 a second electrode facing the first electrode; and  
 an organic layer between the first electrode and the second electrode,  
 wherein the organic layer comprises an emission layer  
 and a hole transport region between the first electrode  
 and the emission layer,  
 the hole transport region comprises a hole transport layer,  
 and  
 the hole transport layer comprises one amine-based compound  
 represented by Formula 1:

214

Formula 1



Formula 2



wherein, in Formulae 1 and 2,

$Ar_1$ ,  $Ar_2$ ,  $L_1$ , and  $L_2$  are each independently a substituted or unsubstituted  $C_5-C_{60}$  carbocyclic group or a substituted or unsubstituted  $C_2-C_{60}$  heterocyclic group,

$a_1$  and  $a_2$  are each independently an integer from 0 to 5, wherein, when  $a_1$  is two or more, two or more  $L_1$ (s) are identical to or different from each other, and when  $a_2$  is two or more, two or more  $L_2$ (s) are identical to or different from each other,

when  $a_1$  is zero,  $*(L_1)_{a_1}-*$  is a single bond, and when  $a_2$  is zero,  $*(L_2)_{a_2}-*$  is a single bond,

$R_1$  to  $R_{10}$  are each independently selected from a group represented by Formula 2, hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1-C_{60}$  alkyl group, a substituted or unsubstituted  $C_2-C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2-C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1-C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkenyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $C_6-C_{60}$  aryl group, a substituted or unsubstituted  $C_6-C_{60}$  aryloxy group, a substituted or unsubstituted  $C_6-C_{60}$  arylthio group, a substituted or unsubstituted  $C_1-C_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-Si(Q_1)(Q_2)(Q_3)$ ,  $-B(Q_1)(Q_2)$ ,  $-C(=O)(Q_1)$ ,  $-P(=O)(Q_1)(Q_2)$ ,  $-P(=S)(Q_1)(Q_2)$ ,  $-S(=O)(Q_1)(Q_2)$ , and  $-S(=O)_2(Q_1)(Q_2)$ ,

provided that one selected from  $R_1$  to  $R_{10}$  is a group represented by Formula 2,

$R_{11}$  and  $R_{12}$  are each independently selected from hydrogen, deuterium,  $-F$ ,  $-Cl$ ,  $-Br$ ,  $-I$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $C_1-C_{60}$  alkyl group, a substituted or unsubstituted  $C_2-C_{60}$  alkenyl group, a substituted or unsubstituted  $C_2-C_{60}$  alkynyl group, a substituted or unsubstituted  $C_1-C_{60}$  alkoxy group, a substituted or unsubstituted  $C_3-C_{10}$  cycloalkyl group, a substituted or unsubstituted  $C_2-C_{10}$  heterocycloalkyl group, a substi-

tuted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

b1 and b2 are each independently an integer from 1 to 10, wherein, when b1 is two or more, two or more R<sub>11</sub>(s) are identical to or different from each other, and when b2 is two or more, two or more R<sub>12</sub>(s) are identical to or different from each other,

at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an

amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), —B(Q<sub>21</sub>)(Q<sub>22</sub>), —C(=O)(Q<sub>21</sub>), —S(=O)<sub>2</sub>(Q<sub>21</sub>), and —P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

—Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), —B(Q<sub>31</sub>)(Q<sub>32</sub>), —C(=O)(Q<sub>31</sub>), —S(=O)<sub>2</sub>(Q<sub>31</sub>), and —P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>);

Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

\* and \*' each indicate a binding site to a neighboring atom.

13. The organic light-emitting device of claim 12, wherein:

the first electrode is an anode,  
the second electrode is a cathode,  
the organic layer further comprises an electron transport region between the emission layer and the second electrode,  
the hole transport region further comprises a hole injection layer, an emission auxiliary layer, an electron blocking layer, or any combination thereof, and  
the electron transport region comprises a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.


14. An organic light-emitting device comprising:

a first electrode;  
a second electrode facing the first electrode; and  
an organic layer between the first electrode and the second electrode,

wherein the organic layer comprises an emission layer, and


the emission layer comprises at least one amine-based compound represented by Formula 1:

Formula 1



217

-continued



Formula 2

wherein, in Formulae 1 and 2,

Ar<sub>1</sub>, Ar<sub>2</sub>, L<sub>1</sub>, and L<sub>2</sub> are each independently a substituted or unsubstituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group or a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group,

a1 and a2 are each independently an integer from 0 to 5, wherein, when a1 is two or more, two or more L<sub>1</sub>(s) are identical to or different from each other, and when a2 is two or more, two or more L<sub>2</sub>(s) are identical to or different from each other,

when a1 is zero, \*-(L<sub>1</sub>)<sub>a1</sub>-\* is a single bond, and when a2 is zero, \*-(L<sub>2</sub>)<sub>a2</sub>-\* is a single bond,

R<sub>1</sub> to R<sub>10</sub> are each independently selected from a group represented by Formula 2, hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>), —C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

provided that one selected from R<sub>1</sub>, R<sub>2</sub>, and R<sub>4</sub> to R<sub>10</sub> is a group represented by Formula 2,

R<sub>11</sub> and R<sub>12</sub> are each independently selected from hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkyl group, a substituted or unsubstituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a substituted or unsubstituted C<sub>2</sub>-C<sub>10</sub> heterocycloalkenyl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryl group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, a substituted or unsubstituted C<sub>6</sub>-C<sub>60</sub> arylthio group, a substituted or unsubstituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>1</sub>)(Q<sub>2</sub>)(Q<sub>3</sub>), —B(Q<sub>1</sub>)(Q<sub>2</sub>),

218

—C(=O)(Q<sub>1</sub>), —P(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), —P(=S)(Q<sub>1</sub>)(Q<sub>2</sub>), —S(=O)(Q<sub>1</sub>)(Q<sub>2</sub>), and —S(=O)<sub>2</sub>(Q<sub>1</sub>)(Q<sub>2</sub>),

b1 and b2 are each independently an integer from 1 to 10, wherein, when b1 is two or more, two or more R<sub>11</sub>(s) are identical to or different from each other, and when b2 is two or more, two or more R<sub>12</sub>(s) are identical to or different from each other,

at least one substituent of the substituted C<sub>5</sub>-C<sub>60</sub> carbocyclic group, the substituted C<sub>2</sub>-C<sub>60</sub> heterocyclic group, the substituted C<sub>1</sub>-C<sub>60</sub> alkyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkenyl group, the substituted C<sub>2</sub>-C<sub>60</sub> alkynyl group, the substituted C<sub>1</sub>-C<sub>60</sub> alkoxy group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, the substituted C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, the substituted C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryl group, the substituted C<sub>6</sub>-C<sub>60</sub> aryloxy group, the substituted C<sub>6</sub>-C<sub>60</sub> arylthio group, the substituted C<sub>1</sub>-C<sub>60</sub> heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium (-D), —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), —B(Q<sub>11</sub>)(Q<sub>12</sub>), —C(=O)(Q<sub>11</sub>), —S(=O)<sub>2</sub>(Q<sub>11</sub>), and —P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed het-

eropolycyclic group,  $-\text{Si}(\text{Q}_{21})(\text{Q}_{22})(\text{Q}_{23})$ ,  $-\text{B}(\text{Q}_{21})(\text{Q}_{22})$ ,  $-\text{C}(=\text{O})(\text{Q}_{21})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{21})$ , and  $-\text{P}(=\text{O})(\text{Q}_{21})(\text{Q}_{22})$ ; and

$-\text{Si}(\text{Q}_{31})(\text{Q}_{32})(\text{Q}_{33})$ ,  $-\text{B}(\text{Q}_{31})(\text{Q}_{32})$ ,  $-\text{C}(=\text{O})(\text{Q}_{31})$ ,  $-\text{S}(=\text{O})_2(\text{Q}_{31})$ , and  $-\text{P}(=\text{O})(\text{Q}_{31})(\text{Q}_{32})$ ,

$\text{Q}_1$  to  $\text{Q}_3$ ,  $\text{Q}_{11}$  to  $\text{Q}_{13}$ ,  $\text{Q}_{21}$  to  $\text{Q}_{23}$ , and  $\text{Q}_{31}$  to  $\text{Q}_{33}$  are each independently selected from hydrogen, deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, a  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkyl group, a  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkenyl group, a  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

\* and \*' each indicate a binding site to a neighboring atom.

15. An organic light-emitting device comprising:

an anode,

a cathode facing the anode; and

an organic layer between the anode and the cathode,

wherein the organic layer comprises:

an emission layer,

a hole transport region between the anode and the emission layer and comprising a hole injection layer,

a hole transport layer, an emission auxiliary layer, an

electron blocking layer, or any combination thereof,

an electron transport region between the emission layer


and the cathode and comprising a hole blocking layer,

an electron transport layer, an electron injection layer,


or any combination thereof, and

at least one of an amine-based compound represented

by Formula 1:



Formula 1



Formula 2

wherein, in Formulae 1 and 2,

$\text{Ar}_1$ ,  $\text{Ar}_2$ ,  $\text{L}_1$ , and  $\text{L}_2$  are each independently a substituted or unsubstituted  $\text{C}_5$ - $\text{C}_{60}$  carbocyclic group or a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{60}$  heterocyclic group,

$a_1$  and  $a_2$  are each independently an integer from 0 to 5,

wherein, when  $a_1$  is two or more, two or more  $\text{L}_1$ (s) are

identical to or different from each other, and when  $a_2$  is two or more, two or more  $\text{L}_2$ (s) are identical to or

different from each other, when  $a_1$  is zero,  $*(\text{L}_1)_{a_1}$  is a single bond, and when  $a_2$  is zero,  $*(\text{L}_2)_{a_2}$  is a single bond,

$\text{R}_1$  to  $\text{R}_{10}$  are each independently selected from a group represented by Formula 2, hydrogen, deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  aryloxy group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  arylthio group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-\text{Si}(\text{Q}_1)(\text{Q}_2)(\text{Q}_3)$ ,  $-\text{B}(\text{Q}_1)(\text{Q}_2)$ ,  $-\text{C}(=\text{O})(\text{Q}_1)$ ,  $-\text{P}(=\text{O})(\text{Q}_1)(\text{Q}_2)$ ,  $-\text{P}(=\text{S})(\text{Q}_1)(\text{Q}_2)$ ,  $-\text{S}(=\text{O})(\text{Q}_1)(\text{Q}_2)$ , and  $-\text{S}(=\text{O})_2(\text{Q}_1)(\text{Q}_2)$ ,

provided that one selected from  $\text{R}_1$  to  $\text{R}_{10}$  is a group represented by Formula 2,

$\text{R}_{11}$  and  $\text{R}_{12}$  are each independently selected from hydrogen, deuterium,  $-\text{F}$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{I}$ , a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkyl group, a substituted or unsubstituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, a substituted or unsubstituted  $\text{C}_2$ - $\text{C}_{10}$  heterocycloalkenyl group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  aryl group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  aryloxy group, a substituted or unsubstituted  $\text{C}_6$ - $\text{C}_{60}$  arylthio group, a substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group,  $-\text{Si}(\text{Q}_1)(\text{Q}_2)(\text{Q}_3)$ ,  $-\text{B}(\text{Q}_1)(\text{Q}_2)$ ,  $-\text{C}(=\text{O})(\text{Q}_1)$ ,  $-\text{P}(=\text{O})(\text{Q}_1)(\text{Q}_2)$ ,  $-\text{P}(=\text{S})(\text{Q}_1)(\text{Q}_2)$ ,  $-\text{S}(=\text{O})(\text{Q}_1)(\text{Q}_2)$ , and  $-\text{S}(=\text{O})_2(\text{Q}_1)(\text{Q}_2)$ ,

$b_1$  and  $b_2$  are each independently an integer from 1 to 10, wherein, when  $b_1$  is two or more, two or more  $\text{R}_{11}$ (s) are identical to or different from each other, and when  $b_2$  is two or more, two or more  $\text{R}_{12}$ (s) are identical to or different from each other,

at least one substituent of the substituted  $\text{C}_5$ - $\text{C}_{60}$  carbocyclic group, the substituted  $\text{C}_2$ - $\text{C}_{60}$  heterocyclic group, the substituted  $\text{C}_1$ - $\text{C}_{60}$  alkyl group, the substituted  $\text{C}_2$ - $\text{C}_{60}$  alkenyl group, the substituted  $\text{C}_2$ - $\text{C}_{60}$  alkynyl group, the substituted  $\text{C}_1$ - $\text{C}_{60}$  alkoxy group, the substituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkyl group, the substituted  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkyl group, the substituted  $\text{C}_3$ - $\text{C}_{10}$  cycloalkenyl group, the substituted  $\text{C}_1$ - $\text{C}_{10}$  heterocycloalkenyl group, the substituted  $\text{C}_6$ - $\text{C}_{60}$  aryl group, the substituted  $\text{C}_6$ - $\text{C}_{60}$  aryloxy group, the substituted  $\text{C}_6$ - $\text{C}_{60}$  arylthio group, the substituted  $\text{C}_1$ - $\text{C}_{60}$  heteroaryl group, the substituted monovalent non-aro-

matic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is selected from:

deuterium (-D), -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group;

a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, and a C<sub>1</sub>-C<sub>60</sub> alkoxy group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q<sub>11</sub>)(Q<sub>12</sub>)(Q<sub>13</sub>), -B(Q<sub>11</sub>)(Q<sub>12</sub>), -C(=O)(Q<sub>11</sub>), -S(=O)<sub>2</sub>(Q<sub>11</sub>), and -P(=O)(Q<sub>11</sub>)(Q<sub>12</sub>);

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group;

a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, and a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one selected from deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, -Si(Q<sub>21</sub>)(Q<sub>22</sub>)(Q<sub>23</sub>), -B(Q<sub>21</sub>)(Q<sub>22</sub>), -C(=O)(Q<sub>21</sub>), -S(=O)<sub>2</sub>(Q<sub>21</sub>), and -P(=O)(Q<sub>21</sub>)(Q<sub>22</sub>); and

-Si(Q<sub>31</sub>)(Q<sub>32</sub>)(Q<sub>33</sub>), -B(Q<sub>31</sub>)(Q<sub>32</sub>), -C(=O)(Q<sub>31</sub>), -S(=O)<sub>2</sub>(Q<sub>31</sub>), and -P(=O)(Q<sub>31</sub>)(Q<sub>32</sub>).

Q<sub>1</sub> to Q<sub>3</sub>, Q<sub>11</sub> to Q<sub>13</sub>, Q<sub>21</sub> to Q<sub>23</sub>, and Q<sub>31</sub> to Q<sub>33</sub> are each independently selected from hydrogen, deuterium, -F, -Cl, -Br, -I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazino group, a hydrazono group, a C<sub>1</sub>-C<sub>60</sub> alkyl group, a C<sub>2</sub>-C<sub>60</sub> alkenyl group, a C<sub>2</sub>-C<sub>60</sub> alkynyl group, a C<sub>1</sub>-C<sub>60</sub> alkoxy group, a C<sub>3</sub>-C<sub>10</sub> cycloalkyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkyl group, a C<sub>3</sub>-C<sub>10</sub> cycloalkenyl group, a C<sub>1</sub>-C<sub>10</sub> heterocycloalkenyl group, a C<sub>6</sub>-C<sub>60</sub> aryl group, a C<sub>6</sub>-C<sub>60</sub> aryloxy group, a C<sub>6</sub>-C<sub>60</sub> arylthio group, a C<sub>1</sub>-C<sub>60</sub> heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, a biphenyl group, and a terphenyl group, and

\* and \*' each indicate a binding site to a neighboring atom;

the hole transport region comprises a p-dopant, and the p-dopant has a lowest unoccupied molecular orbital (LUMO) level of about -3.5 eV or less.

16. The organic light-emitting device of claim 12, wherein the emission layer comprises at least one selected from a styryl-based compound, an anthracene-based compound, a pyrene-based compound, and a spiro-bifluorene-based compound.

17. The organic light-emitting device of claim 12, wherein:



the emission layer is a first emission layer for emitting first color light,

the organic light-emitting device further comprises, between the first electrode and the second electrode, i) at least one second emission layer for emitting second color light or ii) at least one second emission layer for emitting second color light and at least one third emission layer for emitting third color light,

a maximum emission wavelength of the first color light, a maximum emission wavelength of the second color light, and a maximum emission wavelength of the third color light are identical to or different from one another, and





the first color light and the second color light are emitted in the form of mixed light, or the first color light, the second color light, and the third color light are emitted in the form of mixed light.

18. The organic light-emitting device of claim 12, wherein the one amine-based compound is one selected from Compounds 1 to 144:



223

-continued



224

-continued

3

7

5

10

15

20

8

25

30

35

9

5

40

45

50





6

10





55

60

65



**225**  
-continued



**226**  
-continued

11

15

5

10

15

12

20

16

25

30

35

13

17

40

45

50





14

18

55


60

65



227

-continued




19

5

10

15




20

20

25

30




21

21

40

45

50



22


55

60


65

228


-continued




23



24




25



26

229

-continued



27

230

-continued

5

31

10

15

20

25

28

30

35

40







45

50

55





60

65



**231**

-continued



**232**

-continued

35

5

10

15

36

20

25

30

35

37

40

45

50





38

55

60

65

39




40

41

42

233

-continued



43

5

10

15

20

44

25

30

35  
45

40

45

50

46


55

60

65

234

-continued



47


48

49

50

235

-continued



51

236

-continued

5

10

15

20

52

25

30

35

53

40

45

50


54

55


60

65




55







56



57



**237**  
-continued



**238**  
-continued

58

5

10

15

59

20

25

30

35  
60

40

45

50



61

55

60


65

62






63

64



**239**

-continued



**240**

-continued


65

68

5

10

15




20

69

25

30

35




40

70

45

50




55





71

60

65



**241**  
-continued



**242**  
-continued

72

76

5

10

15

73

20

25

30

74

35

40

45





75

50

55





60

65



243

-continued




244

-continued

80

84

5



10


15

81

20

25

30




35

82

40

45




50

83

55


60

65



245

-continued




88

5

10

15

20



89


25

30

35

40

90 45



50


55

60


65

246


-continued



91




92



93

247

-continued



94

5


10

15

20

248

-continued



97

25


95

30

35

40

45





98

96

55

60




65



99

249

-continued



250

-continued

100

103

5

10

15

20

25

101

104

30

35

40

45

102




50

55

60





65

105



**251**

-continued




**252**

-continued

106

110

5




10

15

111

107

20



25


30

35

112

108

40




45

50

113

109

55




60

65

**253**

-continued

114




5

10

115

15




20

25

30

35

116




40

45

50

117



55


60

65


**254**

-continued


118




119



120




121




**255**  
-continued

122




**256**  
-continued


126




123




124




125



126




127







128

129







129

**257**  
-continued




**258**  
-continued



259

-continued

138



5


10

15

260

-continued

141



139

20





25

30

35


40

140



50

55



142

143

144

\* \* \* \* \*