



(43) International Publication Date  
27 September 2012 (27.09.2012)

- (51) International Patent Classification:  
G01N 33/53 (2006.01) G01N 33/574 (2006.01)  
G01N 33/68 (2006.01)
- (21) International Application Number:  
PCT/US2012/030129
- (22) International Filing Date:  
22 March 2012 (22.03.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
61/466,173 22 March 2011 (22.03.2011) US  
61/523,548 15 August 2011 (15.08.2011) US  
61/598,582 14 February 2012 (14.02.2012) US
- (71) Applicant (for all designated States except US): **THE JOHNS HOPKINS UNIVERSITY** [US/US]; 3400 North Charles Street, Baltimore, Maryland 21218 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ZHANG, Hui** [CN/US]; 10137 Lakeside Ct., Ellicott City, Maryland 21042 (US). **TIAN, Yuan** [CN/US]; 108 W. 39th Street, Apt. 20, Baltimore, Maryland 21210 (US). **CHAN, Daniel W.** [US/US]; 12925 Wexford Park, Clarksville, Maryland 21029 (US). **CHEN, Jing** [CN/US]; 105 W. 39th Street, #613, Baltimore, Maryland 21210 (US).

- (74) Agent: **GALVEZ, Guido J.**; 100 N. Charles Street, 5th Floor, Baltimore, Maryland 21201 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:  
— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: BIOMARKERS FOR AGGRESSIVE PROSTATE CANCER




FIG. 1

(57) Abstract: The present invention relates to the field of biomarkers and, more specifically, to biomarkers useful in diagnosing aggressive prostate cancer. In specific embodiments, a method for diagnosing aggressive prostate cancer in a patient comprises (a) measuring the levels of one or more biomarkers in a sample collected from the patient; and (b) comparing the levels of the one or more biomarkers with predefined levels of the same biomarkers that correlate to a patient having aggressive prostate cancer and predefined levels of the same biomarkers that correlate to a patient not having aggressive prostate cancer, wherein a correlation to one of the predefined levels provides the diagnosis. In a specific embodiment, the one or more biomarkers may comprise cathepsin-L (CTSL), periostin, microfibrillar-associated protein 4 (MFAP4), collagen XII, neprilysin, clusterin, neutrophil gelatinase associated lipocalin (NGAL), epithelial cell activating molecule (EpCAM), prostate specific antigen (PSA), membrane metallo-endopeptidase (MME) and asporin (ASPN).



WO 2012/129408 A2

## BIOMARKERS FOR AGGRESSIVE PROSTATE CANCER

### CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 61/598,582,  
5 filed February 14, 2012, U.S. Provisional Application No. 61/523,548, filed August 15, 2011,  
and U.S. Provisional Application No. 61/466,173, filed March 22, 2011, each of which is  
incorporated herein by reference in its entirety.

### STATEMENT OF GOVERNMENTAL INTEREST

This invention was made with U.S. government support under grant no.  
10 U01CA152813. The U.S. government has certain rights in the invention.

### FIELD OF THE INVENTION

The present invention relates to the field of biomarkers. More specifically,  
the present invention relates to biomarkers useful in diagnosing aggressive prostate cancer.

### BACKGROUND OF THE INVENTION

15 Prostate cancer is the most common malignancy in men and the second leading cause  
of death from cancer in the United States. Metastases are the major cause of death from  
cancer. Therefore, aggressive (AG) prostate cancer leads to a higher metastasis rate and  
requires early detection and treatment. Since the discovery of prostate-specific antigen  
(PSA), assays that detect this serum biomarker (together with digital rectal exams) have been  
20 used for the screening of prostate cancer. PSA testing has resulted in early detection and  
intervention. However, the major limitation of PSA is the low specificity and high  
prevalence of detecting benign prostatic hyperplasia, especially in older men. Early detection  
based on PSA testing also fails to distinguish aggressive prostate cancer from nonaggressive  
prostate cancer. Indeed, with the illustration of the limitations of the current PSA-based  
25 screening method, a recently published study randomly assigned 76 693 men at 10 U.S. study  
centers to receive either annual PSA screening (38 343 subjects) or usual care as the control  
(38 350 subjects); this study reported no statistical differences in prostate cancer specific  
mortality between the groups after 7–10 years of follow-up.

Besides preoperative PSA, clinical risk assessment tools for prostate cancer metastasis  
30 before surgery largely rely on the prostate biopsy Gleason score. However, the risk  
assessment based on this clinical criterion is too imprecise to be useful due to biopsy  
sampling error and interobserver grading differences. It is also unable to be used as a  
screening test for early detection of aggressive prostate cancer. Currently, aggressive prostate

cancer is under-detected and under treated while nonaggressive prostate cancer is overdetected and overtreated. Consequences of the difficulty of distinguishing the aggressive and nonaggressive prostate cancer are that prostate cancer patients suffer from unnecessary surgeries, and health care faces massive unnecessary expenditures. Therefore, reliable  
5 biomarkers to distinguish aggressive and nonaggressive prostate cancer are badly needed to prevent patients with nonaggressive prostate cancer from overtreatment and to allow patients with aggressive cancer to receive appropriate treatment earlier in the course of their disease.

#### SUMMARY OF THE INVENTION

The present invention is based, at least in part, on the identification of proteins whose  
10 expression is significantly altered in aggressive prostate tumors. The present invention is the first proteomic study to elucidate the differentially expressed proteins associated with aggressive prostate cancer and thus, are potentially useful for the diagnosis of aggressive prostate cancer. Furthermore, the present invention demonstrates that OCT-embedded frozen tissues could be used in proteomic analysis, and may facilitate the use of OCT-embedded  
15 frozen tissues in biomarker discovery.

Accordingly, in one aspect, the present invention provides methods and compositions useful in diagnosing aggressive prostate cancer. In specific embodiments, a method for aggressive prostate cancer in a patient comprises the steps of (a) measuring the levels of one or more biomarkers in a sample collected from the patient; and (b) comparing the levels of  
20 the one or more biomarkers with predefined levels of the same biomarkers that correlate to a patient having aggressive prostate cancer and predefined levels of the same biomarkers that correlate to a patient not having aggressive prostate cancer, wherein a correlation to one of the predefined levels provides the diagnosis. In a specific embodiment, the one or more biomarkers is selected from the group consisting of cathepsin-L (CTSL), clusterin, neutrophil  
25 gelatinase associated lipocalin (NGAL), epithelial cell activating molecule (EpCAM), prostate specific antigen (PSA) and membrane metallo-endopeptidase (MME). In another embodiment, the one or more biomarkers comprises CTSL, clusterin, NGAL, EpCAM, PSA and MME. In such embodiments, the one or more biomarkers further comprise one or more biomarkers selected from the group consisting of periostin, microfibrillar-associated protein 4  
30 (MFAP4), collagen XII, neprilysin, and asporin (ASPN).

In certain embodiments, the one or more biomarkers comprises CTSL, periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL, EpCAM, PSA MME, ASPN, or combinations thereof. In other embodiments, the one or more biomarkers is selected from the group consisting of CTSL, periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL,

EpCAM, PSA, MME and ASPN. In further embodiments, the one or more biomarkers comprises CTSL, periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL, EpCAM, PSA, MME and ASPN.

In one embodiment, the one or more biomarkers is CTSL. In another embodiment, the one or more biomarkers is periostin. Alternatively, the one or more biomarkers can be MFAP4. In yet another embodiment, the one or more biomarkers is collagen XII. In one embodiment, the one or more biomarkers is neprilysin. In another embodiment, the one or more biomarkers is clusterin. In a specific embodiment, the one or more biomarkers is NGAL. In another specific embodiment, the one or more biomarkers is EpCAM. The one or more biomarkers can be PSA. Alternatively, the one or more biomarkers can be MME. In yet a further embodiment, the one or more biomarkers is ASPN.

In the methods of the present invention, the one or more biomarkers further comprises one or more biomarkers from FIG. 9. In other embodiments, the one or more biomarkers further comprises one or more biomarkers from FIG. 6. In alternative embodiments, the one or more biomarkers further comprises one or more biomarkers from FIG. 7. In other embodiments, the one or more biomarkers further comprises one or more biomarkers from FIGS. 6, 7 and 9. Indeed, the present invention provides methods that utilize any combination of the foregoing biomarkers.

A patient sample can be any type of traditional sample taken from a patient. In certain embodiments, the sample is a blood, plasma, or serum sample. In a specific embodiment, the sample is a blood sample. In another specific embodiment, the sample is a plasma sample. In yet another embodiment, the sample is a serum sample. Alternatively, the sample can be a tissue sample. For example, the sample could be an embedded tissue sample.

In the methods described herein, the measuring step can be performed using an immunoassay. In other embodiments, the measuring step is performed using mass spectrometry. In a specific embodiment, the mass spectrometry technique is multiple reaction monitoring mass spectrometry (MRM-MS).

In certain embodiments, the correlation to a patient not having aggressive prostate cancer refers to a patient having non-aggressive prostate cancer. In other embodiments, the correlation to a patient not having aggressive prostate cancer refers to a patient not having cancer.

The present invention further provides a method for diagnosing aggressive prostate cancer in a patient comprising the steps of (a) collecting a sample from the patient; (b)

measuring the levels of a panel of biomarkers in the sample collected from the patient, wherein the panel of biomarkers comprises one or more of CTSL, clusterin, NGAL, EpCAM, PSA and MME; and (c) comparing the levels of the panel of biomarkers with predefined levels of the same panel of biomarkers that correlate to a patient having aggressive prostate cancer and predefined levels of the same panel of biomarkers that correlate to a patient not  
5 having aggressive prostate cancer, wherein a correlation to one of the predefined levels provides the diagnosis. In a specific embodiment, the panel of biomarkers further comprises one or more of MFAP4, collagen XII, neprilysin, and ASPN.

The present invention also provides methods for determining the aggressive prostate cancer status in a patient. In a specific embodiment, the method comprises the steps of (a)  
10 collecting a sample from the patient; (b) measuring the levels of a panel of biomarkers in the sample collected from the patient, wherein the panel of biomarkers comprises one or more of CTSL, clusterin, NGAL, EpCAM, PSA and MME; and (c) comparing the levels of the panel of biomarkers with predefined levels of the same panel of biomarkers that correlate to one or  
15 more aggressive prostate cancer statuses selected from the group consisting of having aggressive prostate cancer, not having aggressive prostate cancer, progressing aggressive prostate cancer, and regressing aggressive prostate cancer, wherein a correlation to one of the predefined levels determines the aggressive prostate cancer status of the patient. In certain embodiments, the panel of biomarkers further comprises one or more of MFAP4, collagen  
20 XII, neprilysin, and ASPN.

In yet another aspect, the present invention provides kits useful in diagnosing cancer. In certain embodiments, a diagnostic kit for diagnosing aggressive prostate cancer in a patient comprises (a) a substrate for collecting a biological sample from the patient; and  
(b) means for measuring the levels of one or more of CTSL, periostin, MFAP4, collagen XII,  
25 neprilysin, clusterin, NGAL, EpCAM, PSA MME and ASPN. In other embodiments, the kit further comprises means for measuring the levels of one or more of the biomarkers in FIGS 6, 7 and 9.

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 present the workflow of identifying altered glycoprotein changes related to  
30 aggressive prostate carcinoma.

FIG. 2 shows the statistical analysis of protein expression in aggressive (AG) prostate tumor and nonaggressive (NAG) prostate tumor:(FIG. 2A) microfibrillar-associated protein 4 (peptide, vDLEDFEnNTAYAk); (FIG. 2B) microfibrillar-associated protein 4 (peptide, fnGSVSFFR); (FIG. 2C) periostin; and (FIG. 2D) cathepsin L.

FIG. 3 presents the analysis of candidate proteins using Western blots. Periostin showed upregulated expression in all four AG-prostate tumors, while cathepsin L was up-regulated in three out of four AG-prostate tumors, compared to NAG-prostate tumors. IgG was detected evenly in each sample and used as a quantitation control.

5 FIG. 4 shows the analysis of identified proteins using immunohistochemistry. FIG. 4A: immunohistochemical analysis of prostate primary tumor and its adjacent normal tissue with antibody specific to cathepsin L. Cathepsin L was expressed in epithelial cells. Arrows indicate the epithelial cell staining of antibody against cathepsin L. FIG. 4B: immunohistochemical analysis of prostate primary tumor and its adjacent normal tissue with antibody  
10 specific to periostin. Staining exhibited a low background in the normal prostate samples but revealed an overexpression in peritumoral stroma of Gleason 3 tumors and strong overexpression in the peritumoral stroma of Gleason 4 tumors. Arrows point to the peritumoral stroma staining of antibody against periostin.

FIG. 5 is a table showing examples of glycoproteins overexpressed in aggressive  
15 (AG) prostate tumors.

FIG. 6 is a table listing glycopeptides identified from the mass spectrometry analysis as potential biomarkers.

FIG. 7 is a table listing 102 unique N-linked glycopeptides identified (with 95% confidence) from mass spectrometry analysis, representing 79 unique glycoproteins.

20 In FIG. 8, iTRAQ report tags show the increased glycoprotein expression in aggressive (AG) prostate tumor compared to non-aggressive (NAG) prostate tumor. AG-prostate cancer tissues were labeled with iTRAQ Tag 113, 114, 115 and 121 and NAG-prostate cancer tissues were labeled with iTRAQ Tag 116, 117, 118 and 119. FIG. 8A: microfibrillar-associated protein 4 (peptide: vDLEDFEnNTAYAk); FIG. 8B: microfibrillar-associated protein 4 (peptide:  
25 fnGSVSFFR); FIG. 8C: periostin; and FIG. 8D: cathepsin L.

FIG. 9 is a list of 42 proteins identified in the discovery study, with an average ratio between aggressive and non-aggressive tissue and the p value of student's t-test.

FIG. 10 shows the types of tissue specimens used in the validation study.

FIG. 11 shows Cathepsin-L (CTSL) levels in non-aggressive tumor (NAG) vs.  
30 aggressive tumor (AG).

FIG. 12 shows Epithelial Cell Activating Molecule (EpCAM) levels in normal prostate, NAG, AG, and Met.

FIG. 13 shows expression of CTSL and EpCAM in multiple mets by subject.

FIG. 14 shows human Neutrophil Gelatinase Associated Lipocalin (NGAL) levels in normal prostate, NAG, AG, and Met.

FIG. 15 shows Clusterin levels in normal prostate, NAG, AG, and Met.

FIG. 16 shows Neprilysin levels in normal prostate, NAG, AG, and Met.

5 FIG. 17 shows total prostate specific antigen (PSA) levels in NAG, AG, normal prostate and Met.

FIG. 18 shows CTSL levels in NAG vs. AG (no adjustment, adjusted for tumor nuclei, and adjusted for tumor area).

10 FIG. 19 shows EpCAM levels in NAG vs. AG (no adjustment, adjusted for tumor nuclei, and adjusted for tumor area).

FIG. 20 shows biomarker levels in metastatic tumors by site for CTSL (FIG. 20A), Clusterin (FIG. 20B), EpCAM (FIG. 20C), NGAL (FIG. 20D), MME (FIG. 20E), and PSA (FIG. 20F). Sites include subdural, lymph node (LN), liver, bone, prostate, and other.

15 FIG. 21 shows biomarker levels in castrations sensitive vs. castration resistant lymph node metastatic prostate tumors.

FIG. 22 is a Western blot 5 AG and 4 NAG and showing significant Asporin (ASPEN) expression in some AG subjects.

20 FIG. 23. shows immunostaining results for ASPEN. Immunostaining subject PB663 (AG17, intermediate ASPEN expression in Western blot (FIG. 23)) showed strong ASPEN staining for high grade Pca with no/weak staining for Gleason 3 Pca and normal prostate gland.

#### DETAILED DESCRIPTION OF THE INVENTION

It is understood that the present invention is not limited to the particular methods and components, etc., described herein, as these may vary. It is also to be understood that the terminology used herein is used for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention. It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include the plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to a "protein" is a reference to one or more proteins, and includes equivalents thereof known to those skilled in the art and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Specific methods, devices, and materials are described, although any methods and

materials similar or equivalent to those described herein can be used in the practice or testing of the present invention.

All publications cited herein are hereby incorporated by reference including all journal articles, books, manuals, published patent applications, and issued patents. In addition, the meaning of certain terms and phrases employed in the specification, examples, and appended claims are provided. The definitions are not meant to be limiting in nature and serve to provide a clearer understanding of certain aspects of the present invention.

#### I. Definitions

As used herein, the term “antibody” is used in reference to any immunoglobulin molecule that reacts with a specific antigen. It is intended that the term encompass any immunoglobulin (e.g., IgG, IgM, IgA, IgE, IgD, etc.) obtained from any source (e.g., humans, rodents, non-human primates, caprines, bovines, equines, ovines, etc.). Specific types/examples of antibodies include polyclonal, monoclonal, humanized, chimeric, human, or otherwise-human-suitable antibodies. “Antibodies” also includes any fragment or derivative of any of the herein described antibodies.

As used herein, the term “comparing” refers to making an assessment of how the proportion, level or cellular localization of one or more biomarkers in a sample from a patient relates to the proportion, level or cellular localization of the corresponding one or more biomarkers in a standard or control sample. For example, “comparing” may refer to assessing whether the proportion, level, or cellular localization of one or more biomarkers in a sample from a patient is the same as, more or less than, or different from the proportion, level, or cellular localization of the corresponding one or more biomarkers in standard or control sample. More specifically, the term may refer to assessing whether the proportion, level, or cellular localization of one or more biomarkers in a sample from a patient is the same as, more or less than, different from or otherwise corresponds (or not) to the proportion, level, or cellular localization of predefined biomarker levels/ratios that correspond to, for example, a patient having aggressive prostate cancer, not having aggressive prostate cancer (e.g., non-aggressive prostate cancer or no cancer), is responding to treatment for aggressive prostate cancer, is not responding to treatment for aggressive prostate cancer, is/is not likely to respond to a particular aggressive prostate cancer treatment, or having /not having another disease or condition. In a specific embodiment, the term “comparing” refers to assessing whether the level of one or more biomarkers of the present invention in a sample from a patient is the same as, more or less than, different from other otherwise correspond (or not) to levels/ratios of the same biomarkers in a control sample (e.g., predefined levels/ratios that

correlate to uninfected individuals, non-aggressive prostate cancer, standard aggressive prostate cancer levels/ratios, etc.).

In another embodiment, the term “comparing” refers to making an assessment of how the proportion, level or cellular localization of one or more biomarkers in a sample from a patient relates to the proportion, level or cellular localization of another biomarker in the same sample. For example, a ratio of one biomarker to another from the same patient sample can be compared. In another embodiment, a level of one biomarker in a sample (e.g., a post-translationally modified biomarker protein) can be compared to the level of the same biomarker (e.g., unmodified biomarker protein) in the sample. In a specific embodiment, the proportion of a glycosylated biomarker protein can be compared to the unmodified protein, both of which are measured in the same patient sample. Ratios of modified:unmodified biomarker proteins can be compared to other protein ratios in the same sample or to predefined reference or control ratios.

As used herein, the terms “indicates” or “correlates” (or “indicating” or “correlating,” or “indication” or “correlation,” depending on the context) in reference to a parameter, e.g., a modulated proportion, level, or cellular localization in a sample from a patient, may mean that the patient has aggressive prostate cancer. In specific embodiments, the parameter may comprise the level of one or more biomarkers of the present invention. A particular set or pattern of the amounts of one or more biomarkers may indicate that a patient has aggressive prostate cancer (i.e., correlates to a patient having aggressive prostate cancer). In other embodiments, a correlation could be that the ratio of a post-translationally modified protein (e.g., glycosylation) to the unmodified protein indicates (or a change in the ratio over time or as compared to a reference/control ratio) that the patient has aggressive prostate cancer. In specific embodiments, a correlation could be the ratio of a glycosylated peptide to the non-glycosylated form, or any other combination in which a change in one peptide causes or is accompanied by a change in another.

In other embodiments, a particular set or pattern of the amounts of one or more biomarkers may be correlated to a patient being unaffected (i.e., indicates a patient does not have aggressive prostate cancer, a patient has non-aggressive prostate cancer, or a patient does not have cancer). In certain embodiments, “indicating,” or “correlating,” as used according to the present invention, may be by any linear or non-linear method of quantifying the relationship between levels/ratios of biomarkers to a standard, control or comparative value for the assessment of the diagnosis, prediction of aggressive prostate cancer or aggressive prostate cancer progression, assessment of efficacy of clinical treatment,

identification of a patient that may respond to a particular treatment regime or pharmaceutical agent, monitoring of the progress of treatment, and in the context of a screening assay, for the identification of an anti-aggressive prostate cancer therapeutic.

The terms “patient,” “individual,” or “subject” are used interchangeably herein, and  
5 refer to a mammal, particularly, a human. The patient may have mild, intermediate or severe disease. The patient may be treatment naïve, responding to any form of treatment, or refractory. The patient may be an individual in need of treatment or in need of diagnosis based on particular symptoms or family history. In some cases, the terms may refer to treatment in experimental animals, in veterinary application, and in the development of  
10 animal models for disease, including, but not limited to, rodents including mice, rats, and hamsters; and primates.

The terms “measuring” and “determining” are used interchangeably throughout, and refer to methods which include obtaining a patient sample and/or detecting the level of a biomarker(s) in a sample. In one embodiment, the terms refer to obtaining a patient sample  
15 and detecting the level of one or more biomarkers in the sample. In another embodiment, the terms “measuring” and “determining” mean detecting the level of one or more biomarkers in a patient sample. Measuring can be accomplished by methods known in the art and those further described herein. The term “measuring” is also used interchangeably throughout with the term “detecting.”

The terms “sample,” “patient sample,” “biological sample,” and the like, encompass a variety of sample types obtained from a patient, individual, or subject and can be used in a diagnostic or monitoring assay. The patient sample may be obtained from a healthy subject, a diseased patient or a patient having associated symptoms of aggressive prostate cancer. Moreover, a sample obtained from a patient can be divided and only a portion may be used  
25 for diagnosis. Further, the sample, or a portion thereof, can be stored under conditions to maintain sample for later analysis. The definition specifically encompasses blood and other liquid samples of biological origin (including, but not limited to, peripheral blood, serum, plasma, cerebrospinal fluid, urine, saliva, stool and synovial fluid), solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom and the progeny thereof. In  
30 a specific embodiment, a sample comprises a blood sample. In another embodiment, a sample comprises a plasma sample. In yet another embodiment, a serum sample is used.

The definition of “sample” also includes samples that have been manipulated in any way after their procurement, such as by centrifugation, filtration, precipitation, dialysis, chromatography, treatment with reagents, washed, or enriched for certain cell populations.

The terms further encompass a clinical sample, and also include cells in culture, cell supernatants, tissue samples, organs, and the like. Samples may also comprise fresh-frozen and/or formalin-fixed, paraffin-embedded tissue blocks, such as blocks prepared from clinical or pathological biopsies, prepared for pathological analysis or study by  
5 immunohistochemistry. In certain embodiments, a sample comprises an optimal cutting temperature (OCT)-embedded frozen tissue sample.

The terms “specifically binds to,” “specific for,” and related grammatical variants refer to that binding which occurs between such paired species as enzyme/substrate, receptor/agonist, antibody/antigen, and lectin/carbohydrate which may be mediated by  
10 covalent or non-covalent interactions or a combination of covalent and non-covalent interactions. When the interaction of the two species produces a non-covalently bound complex, the binding which occurs is typically electrostatic, hydrogen-bonding, or the result of lipophilic interactions. Accordingly, “specific binding” occurs between a paired species where there is interaction between the two which produces a bound complex having the  
15 characteristics of an antibody/antigen or enzyme/substrate interaction. In particular, the specific binding is characterized by the binding of one member of a pair to a particular species and to no other species within the family of compounds to which the corresponding member of the binding member belongs. Thus, for example, an antibody typically binds to a single epitope and to no other epitope within the family of proteins. In some embodiments,  
20 specific binding between an antigen and an antibody will have a binding affinity of at least  $10^{-6}$  M. In other embodiments, the antigen and antibody will bind with affinities of at least  $10^{-7}$  M,  $10^{-8}$  M to  $10^{-9}$  M,  $10^{-10}$  M,  $10^{-11}$  M, or  $10^{-12}$  M.

Various methodologies of the instant invention include a step that involves comparing a value, level, feature, characteristic, property, etc. to a “suitable control,” referred to  
25 interchangeably herein as an “appropriate control” or a “control sample.” A “suitable control,” “appropriate control” or a “control sample” is any control or standard familiar to one of ordinary skill in the art useful for comparison purposes. In one embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc., determined in a cell, organ, or patient, e.g., a control or normal cell, organ, or patient,  
30 exhibiting, for example, normal traits. For example, the biomarkers of the present invention may be assayed for levels/ratios in a sample from an unaffected individual (UI) or a normal control individual (NC) (both terms are used interchangeably herein). In another embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, ratio, etc. determined prior to performing a therapy (e.g., aggressive

prostate cancer treatment) on a patient. In yet another embodiment, a transcription rate, mRNA level, translation rate, protein level/ratio, biological activity, cellular characteristic or property, genotype, phenotype, etc., can be determined prior to, during, or after administering a therapy into a cell, organ, or patient. In a further embodiment, a “suitable control” or  
5 “appropriate control” is a predefined value, level, feature, characteristic, property, ratio, etc. A “suitable control” can be a profile or pattern of levels/ratios of one or more biomarkers of the present invention that correlates to aggressive prostate cancer, to which a patient sample can be compared. The patient sample can also be compared to a negative control, i.e., a profile that correlates to not having aggressive prostate cancer.

## 10 II. Detection of Aggressive Prostate Cancer Biomarkers

### A. Detection by Mass Spectrometry

In one aspect, the biomarkers of the present invention may be detected by mass spectrometry, a method that employs a mass spectrometer to detect gas phase ions. Examples of mass spectrometers are time-of-flight, magnetic sector, quadrupole filter, ion trap, ion  
15 cyclotron resonance, Orbitrap, hybrids or combinations of the foregoing, and the like.

In particular embodiments, the biomarkers of the present invention are detected using selected reaction monitoring (SRM) mass spectrometry techniques. Selected reaction monitoring (SRM) is a non-scanning mass spectrometry technique, performed on triple quadrupole-like instruments and in which collision-induced dissociation is used as a means to  
20 increase selectivity. In SRM experiments two mass analyzers are used as static mass filters, to monitor a particular fragment ion of a selected precursor ion. The specific pair of mass-over-charge ( $m/z$ ) values associated to the precursor and fragment ions selected is referred to as a "transition" and can be written as parent  $m/z \rightarrow$  fragment  $m/z$  (e.g.  $673.5 \rightarrow 534.3$ ). Unlike common MS based proteomics, no mass spectra are recorded in a SRM analysis. Instead, the  
25 detector acts as counting device for the ions matching the selected transition thereby returning an intensity distribution over time. Multiple SRM transitions can be measured within the same experiment on the chromatographic time scale by rapidly toggling between the different precursor/fragment pairs (sometimes called multiple reaction monitoring, MRM). Typically, the triple quadrupole instrument cycles through a series of transitions and  
30 records the signal of each transition as a function of the elution time. The method allows for additional selectivity by monitoring the chromatographic coelution of multiple transitions for a given analyte. The terms SRM/MRM are occasionally used also to describe experiments conducted in mass spectrometers other than triple quadrupoles (e.g. in trapping instruments) where upon fragmentation of a specific precursor ion a narrow mass range is scanned in MS2

mode, centered on a fragment ion specific to the precursor of interest or in general in experiments where fragmentation in the collision cell is used as a means to increase selectivity. In this application, the terms SRM and MRM or also SRM/MRM can be used interchangeably, since they both refer to the same mass spectrometer operating principle. As  
5 a matter of clarity, the term MRM is used throughout the text, but the term includes both SRM and MRM, as well as any analogous technique, such as e.g. highly-selective reaction monitoring, hSRM, LC-SRM or any other SRM/MRM-like or SRM/MRM-mimicking approaches performed on any type of mass spectrometer and/or, in which the peptides are fragmented using any other fragmentation method such as e.g. CAD (collision-activated  
10 dissociation (also known as CID or collision-induced dissociation), HCD (higher energy CID), ECD (electron capture dissociation), PD (photodissociation) or ETD (electron transfer dissociation)).

In another specific embodiment, the mass spectrometric method comprises matrix assisted laser desorption/ionization time-of-flight (MALDI-TOF MS or MALDI-TOF). In  
15 another embodiment, method comprises MALDI-TOF tandem mass spectrometry (MALDI-TOF MS/MS). In yet another embodiment, mass spectrometry can be combined with another appropriate method(s) as may be contemplated by one of ordinary skill in the art. For example, MALDI-TOF can be utilized with trypsin digestion and tandem mass spectrometry as described herein.

In an alternative embodiment, the mass spectrometric technique comprises surface  
20 enhanced laser desorption and ionization or "SELDI," as described, for example, in U.S. Patents No. 6,225,047 and No. 5,719,060. Briefly, SELDI refers to a method of desorption/ionization gas phase ion spectrometry (e.g. mass spectrometry) in which an analyte (here, one or more of the biomarkers) is captured on the surface of a SELDI mass  
25 spectrometry probe. There are several versions of SELDI that may be utilized including, but not limited to, Affinity Capture Mass Spectrometry (also called Surface-Enhanced Affinity Capture (SEAC)), and Surface-Enhanced Neat Desorption (SEND) which involves the use of probes comprising energy absorbing molecules that are chemically bound to the probe surface (SEND probe). Another SELDI method is called Surface-Enhanced Photolabile  
30 Attachment and Release (SEPAR), which involves the use of probes having moieties attached to the surface that can covalently bind an analyte, and then release the analyte through breaking a photolabile bond in the moiety after exposure to light, e.g., to laser light (see, U.S. Patent No. 5,719,060). SEPAR and other forms of SELDI are readily adapted to detecting a biomarker or biomarker panel, pursuant to the present invention.

In another mass spectrometry method, the biomarkers can be first captured on a chromatographic resin having chromatographic properties that bind the biomarkers. For example, one could capture the biomarkers on a cation exchange resin, such as CM Ceramic HyperD F resin, wash the resin, elute the biomarkers and detect by MALDI. Alternatively, this method could be preceded by fractionating the sample on an anion exchange resin before application to the cation exchange resin. In another alternative, one could fractionate on an anion exchange resin and detect by MALDI directly. In yet another method, one could capture the biomarkers on an immuno-chromatographic resin that comprises antibodies that bind the biomarkers, wash the resin to remove unbound material, elute the biomarkers from the resin and detect the eluted biomarkers by MALDI or by SELDI.

#### B. Detection by Immunoassay

In other embodiments, the biomarkers of the present invention can be detected and/or measured by immunoassay. Immunoassay requires biospecific capture reagents, such as antibodies, to capture the biomarkers. Many antibodies are available commercially. Antibodies also can be produced by methods well known in the art, e.g., by immunizing animals with the biomarkers. Biomarkers can be isolated from samples based on their binding characteristics. Alternatively, if the amino acid sequence of a polypeptide biomarker is known, the polypeptide can be synthesized and used to generate antibodies by methods well-known in the art.

The present invention contemplates traditional immunoassays including, for example, sandwich immunoassays including ELISA or fluorescence-based immunoassays, immunoblots, Western Blots (WB), as well as other enzyme immunoassays. Nephelometry is an assay performed in liquid phase, in which antibodies are in solution. Binding of the antigen to the antibody results in changes in absorbance, which is measured. In a SELDI-based immunoassay, a biospecific capture reagent for the biomarker is attached to the surface of an MS probe, such as a pre-activated protein chip array. The biomarker is then specifically captured on the biochip through this reagent, and the captured biomarker is detected by mass spectrometry.

Although antibodies are useful because of their extensive characterization, any other suitable agent (e.g., a peptide, an aptamer, lectin, or a small organic molecule) that specifically binds a biomarker of the present invention is optionally used in place of the antibody in the above described immunoassays. For example, an aptamer that specifically binds all neurogranin and/or one or more of its breakdown products might be used. Aptamers are nucleic acid-based molecules that bind specific ligands. Methods for making aptamers

with a particular binding specificity are known as detailed in U.S. Patents No. 5,475,096; No. 5,670,637; No. 5,696,249; No. 5,270,163; No. 5,707,796; No. 5,595,877; No. 5,660,985; No. 5,567,588; No. 5,683,867; No. 5,637,459; and No. 6,011,020.

C. Detection by Electrochemicaluminescent Assay

5 In several embodiments, the biomarker biomarkers of the present invention may be detected by means of an electrochemicaluminescent assay developed by Meso Scale Discovery (Gaithersburg, MD). Electrochemiluminescence detection uses labels that emit light when electrochemically stimulated. Background signals are minimal because the stimulation mechanism (electricity) is decoupled from the signal (light). Labels are stable,  
10 non-radioactive and offer a choice of convenient coupling chemistries. They emit light at ~620 nm, eliminating problems with color quenching. *See* U.S. Patents No. 7,497,997; No. 7,491,540; No. 7,288,410; No. 7,036,946; No. 7,052,861; No. 6,977,722; No. 6,919,173; No. 6,673,533; No. 6,413,783; No. 6,362,011; No. 6,319,670; No. 6,207,369; No. 6,140,045; No. 6,090,545; and No. 5,866,434. *See also* U.S. Patent Applications Publication No.  
15 2009/0170121; No. 2009/006339; No. 2009/0065357; No. 2006/0172340; No. 2006/0019319; No. 2005/0142033; No. 2005/0052646; No. 2004/0022677; No. 2003/0124572; No. 2003/0113713; No. 2003/0003460; No. 2002/0137234; No. 2002/0086335; and No. 2001/0021534.

D. Other Methods for Detecting Biomarkers

20 The biomarkers of the present invention can be detected by other suitable methods. Detection paradigms that can be employed to this end include optical methods, electrochemical methods (voltametry and amperometry techniques), atomic force microscopy, and radio frequency methods, e.g., multipolar resonance spectroscopy. Illustrative of optical methods, in addition to microscopy, both confocal and non-confocal,  
25 are detection of fluorescence, luminescence, chemiluminescence, absorbance, reflectance, transmittance, and birefringence or refractive index (e.g., surface plasmon resonance, ellipsometry, a resonant mirror method, a grating coupler waveguide method or interferometry).

Furthermore, a sample may also be analyzed by means of a biochip. Biochips  
30 generally comprise solid substrates and have a generally planar surface, to which a capture reagent (also called an adsorbent or affinity reagent) is attached. Frequently, the surface of a biochip comprises a plurality of addressable locations, each of which has the capture reagent bound there. Protein biochips are biochips adapted for the capture of polypeptides. Many protein biochips are described in the art. These include, for example, protein biochips

produced by CIPHERGEN Biosystems, Inc. (Fremont, CA.), Invitrogen Corp. (Carlsbad, CA), Affymetrix, Inc. (Fremont, CA), Zyomyx (Hayward, CA), R&D Systems, Inc. (Minneapolis, MN), Biacore (Uppsala, Sweden) and Procognia (Berkshire, UK). Examples of such protein biochips are described in the following patents or published patent applications: U.S. Patent  
5 No. 6,537,749; U.S. Patent No. 6,329,209; U.S. Patent No. 6,225,047; U.S. Patent No. 5,242,828; PCT International Publication No. WO 00/56934; and PCT International Publication No. WO 03/048768.

### III. Determination of a Patient's Aggressive Prostate Cancer Status

A. The present invention relates to the use of biomarkers to diagnose aggressive prostate  
10 cancer. More specifically, the biomarkers of the present invention can be used in diagnostic tests to determine, qualify, and/or assess aggressive prostate cancer or status, for example, to diagnose aggressive prostate cancer, in an individual, subject or patient. In particular embodiments, aggressive prostate cancer status can include determining a patient's aggressive prostate cancer status, for example, to diagnose aggressive prostate cancer, in an  
15 individual, subject or patient. More specifically, the biomarkers to be detected in diagnosing aggressive prostate cancer include cathepsin-L (CTSL), periostin, microfibrillar-associated protein 4 (MFAP4), collagen XII, nephrilysin, clusterin, neutrophil gelatinase associated lipocalin (NGAL), epithelial cell activating molecule (EpCAM), prostate specific antigen (PSA), membrane metallo-endopeptidase (MME) and asporin (ASPN). FIGS. 6, 7 and 9 also  
20 list biomarkers useful in the methods of the present invention. Other biomarkers known in the relevant art may be used in combination with the biomarkers described herein.

#### B. Biomarker Panels

The biomarkers of the present invention can be used in diagnostic tests to assess, determine, and/or qualify (used interchangeably herein) aggressive prostate cancer status in a  
25 patient. The phrase "aggressive prostate cancer status" includes any distinguishable manifestation of the condition, including not having aggressive prostate cancer. For example, aggressive prostate cancer status includes, without limitation, the presence or absence of aggressive prostate cancer in a patient, the risk of developing aggressive prostate cancer, the stage or severity of aggressive prostate cancer, the progress of aggressive prostate cancer  
30 (e.g., progress of aggressive prostate cancer over time) and the effectiveness or response to treatment of aggressive prostate cancer (e.g., clinical follow up and surveillance of aggressive prostate cancer after treatment). Based on this status, further procedures may be indicated, including additional diagnostic tests or therapeutic procedures or regimens.

The power of a diagnostic test to correctly predict status is commonly measured as the sensitivity of the assay, the specificity of the assay or the area under a receiver operated characteristic (“ROC”) curve. Sensitivity is the percentage of true positives that are predicted by a test to be positive, while specificity is the percentage of true negatives that are predicted by a test to be negative. An ROC curve provides the sensitivity of a test as a function of 1-specificity. The greater the area under the ROC curve, the more powerful the predictive value of the test. Other useful measures of the utility of a test are positive predictive value and negative predictive value. Positive predictive value is the percentage of people who test positive that are actually positive. Negative predictive value is the percentage of people who test negative that are actually negative.

In particular embodiments, the biomarker panels of the present invention may show a statistical difference in different aggressive prostate cancer statuses of at least  $p < 0.05$ ,  $p < 10^{-2}$ ,  $p < 10^{-3}$ ,  $p < 10^{-4}$  or  $p < 10^{-5}$ . Diagnostic tests that use these biomarkers may show an ROC of at least 0.6, at least about 0.7, at least about 0.8, or at least about 0.9.

The biomarkers can be differentially present in UI (NC or non-aggressive prostate cancer) and aggressive prostate cancer, and, therefore, are useful in aiding in the determination of aggressive prostate cancer status. In certain embodiments, the biomarkers are measured in a patient sample using the methods described herein and compared, for example, to predefined biomarker levels/ratios and correlated to aggressive prostate cancer status. In particular embodiments, the measurement(s) may then be compared with a relevant diagnostic amount(s), cut-off(s), or multivariate model scores that distinguish a positive aggressive prostate cancer status from a negative aggressive prostate cancer status. The diagnostic amount(s) represents a measured amount of a biomarker(s) above which or below which a patient is classified as having a particular aggressive prostate cancer status. For example, if the biomarker(s) is/are up-regulated compared to normal (e.g., no cancer or non-aggressive prostate cancer) during aggressive prostate cancer, then a measured amount(s) above the diagnostic cutoff(s) provides a diagnosis of aggressive prostate cancer. Alternatively, if the biomarker(s) is/are down-regulated during aggressive prostate cancer, then a measured amount(s) at or below the diagnostic cutoff(s) provides a diagnosis of non-aggressive prostate cancer. The opposite may hold true as well (i.e., expression of the biomarker is lower/downregulated in progressive prostate cancer vs. no cancer or non-aggressive prostate cancer) As is well understood in the art, by adjusting the particular diagnostic cut-off(s) used in an assay, one can increase sensitivity or specificity of the diagnostic assay depending on the preference of the diagnostician. In particular

embodiments, the particular diagnostic cut-off can be determined, for example, by measuring the amount of biomarkers in a statistically significant number of samples from patients with the different aggressive prostate cancer statuses, and drawing the cut-off to suit the desired levels of specificity and sensitivity.

5           In other embodiments, ratios of post-translationally modified biomarkers (e.g., glycosylated, citrullination, oxidation, methylation, phosphorylation, cysteinylolation, s-nitrosation, s-glutathylation, or a combination thereof) to the corresponding unmodified biomarkers are useful in aiding in the determination of aggressive prostate cancer status. In certain embodiments, the biomarker ratios are indicative of diagnosis. In other embodiments, a biomarker ratio can be compared to another biomarker ratio in the same sample or to a set  
10 of biomarker ratios from a control or reference sample.

Indeed, as the skilled artisan will appreciate there are many ways to use the measurements of two or more biomarkers in order to improve the diagnostic question under investigation. In a quite simple, but nonetheless often effective approach, a positive result is  
15 assumed if a sample is positive for at least one of the markers investigated.

Furthermore, in certain embodiments, the values measured for markers of a biomarker panel are mathematically combined and the combined value is correlated to the underlying diagnostic question. Biomarker values may be combined by any appropriate state of the art mathematical method. Well-known mathematical methods for correlating a marker  
20 combination to a disease status employ methods like discriminant analysis (DA) (e.g., linear-, quadratic-, regularized-DA), Discriminant Functional Analysis (DFA), Kernel Methods (e.g., SVM), Multidimensional Scaling (MDS), Nonparametric Methods (e.g., k-Nearest-Neighbor Classifiers), PLS (Partial Least Squares), Tree-Based Methods (e.g., Logic Regression, CART, Random Forest Methods, Boosting/Bagging Methods), Generalized Linear Models  
25 (e.g., Logistic Regression), Principal Components based Methods (e.g., SIMCA), Generalized Additive Models, Fuzzy Logic based Methods, Neural Networks and Genetic Algorithms based Methods. The skilled artisan will have no problem in selecting an appropriate method to evaluate a biomarker combination of the present invention. In one embodiment, the method used in a correlating a biomarker combination of the present  
30 invention, e.g. to diagnose aggressive prostate cancer, is selected from DA (e.g., Linear-, Quadratic-, Regularized Discriminant Analysis), DFA, Kernel Methods (e.g., SVM), MDS, Nonparametric Methods (e.g., k-Nearest-Neighbor Classifiers), PLS (Partial Least Squares), Tree-Based Methods (e.g., Logic Regression, CART, Random Forest Methods, Boosting Methods), or Generalized Linear Models (e.g., Logistic Regression), and Principal

Components Analysis. Details relating to these statistical methods are found in the following references: Ruczinski et al., 12 J. OF COMPUTATIONAL AND GRAPHICAL STATISTICS 475-511 (2003); Friedman, J. H., 84 J. OF THE AMERICAN STATISTICAL ASSOCIATION 165-75 (1989); Hastie, Trevor, Tibshirani, Robert, Friedman, Jerome, The Elements of Statistical Learning, Springer Series in Statistics (2001); Breiman, L., Friedman, J. H., Olshen, R. A., Stone, C. J. Classification and regression trees, California: Wadsworth (1984); Breiman, L., 45 MACHINE LEARNING 5-32 (2001); Pepe, M. S., The Statistical Evaluation of Medical Tests for Classification and Prediction, Oxford Statistical Science Series, 28 (2003); and Duda, R. O., Hart, P. E., Stork, D. G., Pattern Classification, Wiley Interscience, 2nd Edition (2001).

10 C. Determining Risk of Developing Aggressive Prostate Cancer

In a specific embodiment, the present invention provides methods for determining the risk of developing aggressive prostate cancer in a patient. Biomarker percentages, ratios, amounts or patterns are characteristic of various risk states, e.g., high, medium or low. The risk of developing aggressive prostate cancer is determined by measuring the relevant biomarkers and then either submitting them to a classification algorithm or comparing them with a reference amount, i.e., a predefined level or pattern of biomarkers that is associated with the particular risk level.

D. Determining Aggressive Prostate Cancer Severity

In another embodiment, the present invention provides methods for determining the severity of aggressive prostate cancer in a patient. Each grade or stage of aggressive prostate cancer likely has a characteristic level of a biomarker or relative levels/ratios of a set of biomarkers (a pattern or ratio). The severity of aggressive prostate cancer is determined by measuring the relevant biomarkers and then either submitting them to a classification algorithm or comparing them with a reference amount, i.e., a predefined level or pattern of biomarkers that is associated with the particular stage.

E. Determining Aggressive Prostate Cancer Prognosis

In one embodiment, the present invention provides methods for determining the course of aggressive prostate cancer in a patient. Aggressive prostate cancer course refers to changes in aggressive prostate cancer status over time, including aggressive prostate cancer progression (worsening) and aggressive prostate cancer regression (improvement). Over time, the amount or relative amount (e.g., the pattern or ratio) of the biomarkers changes. For example, biomarker "X" may be increased with aggressive prostate cancer, while biomarker "Y" may be decreased with aggressive prostate cancer. Therefore, the trend of these biomarkers, either increased or decreased over time toward aggressive prostate cancer or non-

aggressive prostate cancer indicates the course of the condition. Accordingly, this method involves measuring the level of one or more biomarkers in a patient at least two different time points, e.g., a first time and a second time, and comparing the change, if any. The course of aggressive prostate cancer is determined based on these comparisons.

5 F. Patient Management

In certain embodiments of the methods of qualifying aggressive prostate cancer status, the methods further comprise managing patient treatment based on the status. Such management includes the actions of the physician or clinician subsequent to determining aggressive prostate cancer status. For example, if a physician makes a diagnosis of aggressive prostate cancer, then a certain regime of monitoring would follow. An assessment of the course of aggressive prostate cancer using the methods of the present invention may then require a certain aggressive prostate cancer therapy regimen. Alternatively, a diagnosis of non-aggressive prostate cancer might be followed with further testing to determine a specific disease that the patient might be suffering from. Also, further tests may be called for if the diagnostic test gives an inconclusive result on aggressive prostate cancer status.

10 G. Determining Therapeutic Efficacy of Pharmaceutical Drug

In another embodiment, the present invention provides methods for determining the therapeutic efficacy of a pharmaceutical drug. These methods are useful in performing clinical trials of the drug, as well as monitoring the progress of a patient on the drug. Therapy or clinical trials involve administering the drug in a particular regimen. The regimen may involve a single dose of the drug or multiple doses of the drug over time. The doctor or clinical researcher monitors the effect of the drug on the patient or subject over the course of administration. If the drug has a pharmacological impact on the condition, the amounts or relative amounts (e.g., the pattern, profile or ratio) of one or more of the biomarkers of the present invention may change toward a non-aggressive prostate cancer profile. Therefore, one can follow the course of one or more biomarkers in the patient during the course of treatment. Accordingly, this method involves measuring one or more biomarkers in a patient receiving drug therapy, and correlating the biomarker levels/ratios with the aggressive prostate cancer status of the patient (e.g., by comparison to predefined levels/ratios of the biomarkers that correspond to different aggressive prostate cancer statuses). One embodiment of this method involves determining the levels/ratios of one or more biomarkers for at least two different time points during a course of drug therapy, e.g., a first time and a second time, and comparing the change in levels/ratios of the biomarkers, if any. For example, the levels/ratios of one or more biomarkers can be measured before and after drug

administration or at two different time points during drug administration. The effect of therapy is determined based on these comparisons. If a treatment is effective, then the level/ratio of one or more biomarkers will trend toward normal, while if treatment is ineffective, the level/ratio of one or more biomarkers will trend toward aggressive prostate cancer indications.

#### H. Generation of Classification Algorithms for Qualifying Aggressive Prostate Cancer Status

In some embodiments, data that are generated using samples such as “known samples” can then be used to “train” a classification model. A “known sample” is a sample that has been pre-classified. The data that are used to form the classification model can be referred to as a “training data set.” The training data set that is used to form the classification model may comprise raw data or pre-processed data. Once trained, the classification model can recognize patterns in data generated using unknown samples. The classification model can then be used to classify the unknown samples into classes. This can be useful, for example, in predicting whether or not a particular biological sample is associated with a certain biological condition (e.g., diseased versus non-diseased).

Classification models can be formed using any suitable statistical classification or learning method that attempts to segregate bodies of data into classes based on objective parameters present in the data. Classification methods may be either supervised or unsupervised. Examples of supervised and unsupervised classification processes are described in Jain, “Statistical Pattern Recognition: A Review”, IEEE Transactions on Pattern Analysis and Machine Intelligence, Vol. 22, No. 1, January 2000, the teachings of which are incorporated by reference.

In supervised classification, training data containing examples of known categories are presented to a learning mechanism, which learns one or more sets of relationships that define each of the known classes. New data may then be applied to the learning mechanism, which then classifies the new data using the learned relationships. Examples of supervised classification processes include linear regression processes (e.g., multiple linear regression (MLR), partial least squares (PLS) regression and principal components regression (PCR)), binary decision trees (e.g., recursive partitioning processes such as CART), artificial neural networks such as back propagation networks, discriminant analyses (e.g., Bayesian classifier or Fischer analysis), logistic classifiers, and support vector classifiers (support vector machines).

Another supervised classification method is a recursive partitioning process. Recursive partitioning processes use recursive partitioning trees to classify data derived from unknown samples. Further details about recursive partitioning processes are provided in U.S. Patent Application No. 2002 0138208 A1 to Paulse et al., “Method for analyzing mass  
5 spectra.”

In other embodiments, the classification models that are created can be formed using unsupervised learning methods. Unsupervised classification attempts to learn classifications based on similarities in the training data set, without pre-classifying the spectra from which the training data set was derived. Unsupervised learning methods include cluster analyses. A  
10 cluster analysis attempts to divide the data into “clusters” or groups that ideally should have members that are very similar to each other, and very dissimilar to members of other clusters. Similarity is then measured using some distance metric, which measures the distance between data items, and clusters together data items that are closer to each other. Clustering techniques include the MacQueen’s K-means algorithm and the Kohonen’s Self-Organizing  
15 Map algorithm.

Learning algorithms asserted for use in classifying biological information are described, for example, in PCT International Publication No. WO 01/31580 (Barnhill et al., “Methods and devices for identifying patterns in biological systems and methods of use thereof”), U.S. Patent Application Publication No. 2002/0193950 (Gavin et al. “Method or  
20 analyzing mass spectra”), U.S. Patent Application Publication No. 2003/0004402 (Hitt et al., “Process for discriminating between biological states based on hidden patterns from biological data”), and U.S. Patent Application Publication No. 2003/0055615 (Zhang and Zhang, “Systems and methods for processing biological expression data”).

The classification models can be formed on and used on any suitable digital computer.  
25 Suitable digital computers include micro, mini, or large computers using any standard or specialized operating system, such as a Unix, Windows® or Linux™ based operating system. In embodiments utilizing a mass spectrometer, the digital computer that is used may be physically separate from the mass spectrometer that is used to create the spectra of interest, or it may be coupled to the mass spectrometer.

30 The training data set and the classification models according to embodiments of the invention can be embodied by computer code that is executed or used by a digital computer. The computer code can be stored on any suitable computer readable media including optical or magnetic disks, sticks, tapes, etc., and can be written in any suitable computer programming language including R, C, C++, visual basic, etc.

The learning algorithms described above are useful both for developing classification algorithms for the biomarkers already discovered, and for finding new biomarker biomarkers. The classification algorithms, in turn, form the base for diagnostic tests by providing diagnostic values (e.g., cut-off points) for biomarkers used singly or in combination.

5 IV. Kits for the Detection of Aggressive Prostate Cancer Biomarkers

In another aspect, the present invention provides kits for qualifying aggressive prostate cancer status, which kits are used to detect the biomarkers described herein. In a specific embodiment, the kit is provided as an ELISA kit comprising antibodies to the biomarkers of the present invention including, but not limited to, CTSL, periostin, MFAP4,  
10 collagen XII, neprilysin, clusterin, NGAL, EpCAM, PSA MME, ASPN, as well as the proteins/peptides listed in FIGS. 6, 7 and 9, and combinations of all of the foregoing.

The ELISA kit may comprise a solid support, such as a chip, microtiter plate (e.g., a 96-well plate), bead, or resin having biomarker capture reagents attached thereon. The kit may further comprise a means for detecting the biomarkers, such as antibodies, and a  
15 secondary antibody-signal complex such as horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG antibody and tetramethyl benzidine (TMB) as a substrate for HRP.

The kit for qualifying aggressive prostate cancer status may be provided as an immuno-chromatography strip comprising a membrane on which the antibodies are immobilized, and a means for detecting, e.g., gold particle bound antibodies, where the  
20 membrane, includes NC membrane and PVDF membrane. The kit may comprise a plastic plate on which a sample application pad, gold particle bound antibodies temporally immobilized on a glass fiber filter, a nitrocellulose membrane on which antibody bands and a secondary antibody band are immobilized and an absorbent pad are positioned in a serial manner, so as to keep continuous capillary flow of blood serum.

In certain embodiments, a patient can be diagnosed by adding blood or blood serum  
25 from the patient to the kit and detecting the relevant biomarkers conjugated with antibodies, specifically, by a method which comprises the steps of: (i) collecting blood or blood serum from the patient; (ii) separating blood serum from the patient's blood; (iii) adding the blood serum from patient to a diagnostic kit; and, (iv) detecting the biomarkers conjugated with  
30 antibodies. In this method, the antibodies are brought into contact with the patient's blood. If the biomarkers are present in the sample, the antibodies will bind to the sample, or a portion thereof. In other kit and diagnostic embodiments, blood or blood serum need not be collected from the patient (i.e., it is already collected). Moreover, in other embodiments, the sample may comprise a tissue sample or a clinical sample.

The kit can also comprise a washing solution or instructions for making a washing solution, in which the combination of the capture reagents and the washing solution allows capture of the biomarkers on the solid support for subsequent detection by, e.g., antibodies or mass spectrometry. In a further embodiment, a kit can comprise instructions for suitable operational parameters in the form of a label or separate insert. For example, the instructions may inform a consumer about how to collect the sample, how to wash the probe or the particular biomarkers to be detected, etc. In yet another embodiment, the kit can comprise one or more containers with biomarker samples, to be used as standard(s) for calibration.

Without further elaboration, it is believed that one skilled in the art, using the preceding description, can utilize the present invention to the fullest extent. The following examples are illustrative only, and not limiting of the remainder of the disclosure in any way whatsoever.

#### EXAMPLES

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices, and/or methods described and claimed herein are made and evaluated, and are intended to be purely illustrative and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for herein. Unless indicated otherwise, parts are parts by weight, temperature is in degrees Celsius or is at ambient temperature, and pressure is at or near atmospheric. There are numerous variations and combinations of reaction conditions, e.g., component concentrations, desired solvents, solvent mixtures, temperatures, pressures and other reaction ranges and conditions that can be used to optimize the product purity and yield obtained from the described process. Only reasonable and routine experimentation will be required to optimize such process conditions.

#### Materials and Methods

Materials. Hydrazide resin and sodium periodate were from Bio-Rad (Hercules, CA); sequencing grade trypsin was purchased from Promega (Madison, WI); PNGase F was from New England Biolabs (Ipswich, MA); C18 columns were from Waters (Milford, MA); iTRAQ reagent was from Applied Biosystems (Foster City, CA); rabbit antihuman periostin antibody and mouse antihuman cathepsin L antibody for Western blot analysis were from Abcam (Cambridge, U.K.); goat anticathepsin L antibody for immunohistochemistry was from R&D systems (Minneapolis, MN); BCA assay kit, HRP-labeled secondary antibodies,

and the Novex ECL Chemiluminescent Substrate Reagent Kit were from Pierce (Rockford, IL); LSAB+ System-AP kit from Dako (Carpinteria, CA); and all other chemicals were purchased from Sigma-Aldrich (St. Louis, MO).

Prostate Cancer Tissues. Samples and clinical information were obtained as part of a  
5 Johns Hopkins Medicine IRB approved study. Tissues from four nonaggressive and four aggressive primary prostate tumors were analyzed. The nonaggressive and aggressive primary cancer tissues were collected from radical prostatectomy specimens at Johns Hopkins Hospital and Johns Hopkins Bayview Medical Center. Under the NCI-funded Johns Hopkins prostate cancer SPORE project, annual follow-up data is obtained from consenting  
10 subjects, allowing clinical follow up data to be used in conjunction with pathologic data. The cancer tissues were selected from cases with known Gleason score and clinical outcome following surgery. The nonaggressive cancer tissues were microdissected from primary tissues with radical prostatectomy Gleason scores of  $3 + 3 = 6$  and without evidence of cancer recurrence (by PSA or other clinical data) after 11–15 years of follow-up. The aggressive  
15 cancer tissues were similarly microdissected from primary tissues of tumors with Gleason scores of  $5 + 4 = 9$  (3 cases) or  $5 + 3 = 8$  (one case) and either seminal vesicle or pelvic lymph node involvement (or both). Two of the men with aggressive prostate cancer died of prostate cancer 2 and 6 years after surgery, 1 died of myocardial infarction 5 years after surgery with known prostate cancer recurrence, and 1 is thought to have been surgically  
20 cured of aggressive prostate cancer as he shows no evidence of recurrence 15 years after surgery.

Glycopeptide Isolation. OCT-embedded prostate frozen tissues were sectioned and stained with hematoxylin and eosin. The microscopic view of a histologic specimen was used for cryostat microdissection to remove the nontumor tissue. The proteins from each  
25 prostate cancer tissue were collected from one 10  $\mu$ m OCT tumor section using lysis buffer (50 mM Tris, pH8.0, 150 mM NaCl, 0.1% SDS, 0.5% Na Deoxycholate, 1% Triton X 100). BCA assay was performed to determine the protein concentration. For each tissue specimen, 100  $\mu$ g of proteins were used in the following experiment. Urea (8 M) and 5 mM TCEP as the final concentration were added to the sample and incubated at 60 °C for 2 h. Iodoaceta-  
30 mide (10 mM) as the final concentration was added to the sample and incubated for 30 min. at room temperature in the dark. The solution was diluted 8-fold with 100 mM  $\text{KH}_2\text{PO}_4$  (pH 8.0). An amount of 2  $\mu$ g of trypsin (Promega) was added to the solution and incubated at 37°C overnight with shaking. Silver staining was used to determine whether the trypsin digestion was complete, which was indicated by the disappearance of the upper protein bands

and appearance of lower peptide bands (<10 kDa). After trypsin digestion, samples were centrifuged at 13 000 rpm for 5 min to remove any particulate matter. The peptides were cleaned by C18 column and followed by N-linked glycopeptide isolation using the method of solid phase extraction of glycopeptides (SPEG). The enriched N-linked glycopeptides were concentrated by C18 columns and dried down and resuspended in 10  $\mu$ L of 0.4% acetic acid.

iTRAQ Labeling. Glycopeptides (10  $\mu$ L) from each sample were labeled with iTRAQ 8plex (AB SCIEX) according to the manufacturer's instruction. Peptides of four AG-prostate tumors were labeled by iTRAQ with 113, 114, 115, and 121, respectively, while peptides of four NAG-prostate tumors were labeled by iTRAQ with 116, 117, 118, and 119. Labeled peptides were then mixed and cleaned by a SCX column.

LC-MS/MS Analysis. iTRAQ labeled glycopeptides were separated on a C18 column (75  $\mu$ m  $\times$  10 cm, 5  $\mu$ m, 120  $\text{\AA}$ , Magic C18, MicromBioresources, Auburn, CA) at 750 nL/min in 15 min loading time. The peptides were then eluted at 300 nL/min by using a gradient of 100 min at the voltage of 2.0 kV. Eluting peptides were sprayed into an LTQ Orbitrap Velos mass spectrometer (Thermo Scientific) with a 1  $\mu$ m emitter tip (New Objective, Woburn, MA) using a 5–40% solution B (90% acetonitrile in 0.1% formic acid) gradient.

Identification of Protein and Glycosylation Sites. MS/MS spectra were searched with MASCOT using Proteome Discoverer (version 1.0) (Thermo Fisher) against the human subdatabase of NCBI Reference Sequence (RefSeq) (version 40, released on April 16, 2010) containing 29 704 sequences. The precursor mass tolerance was set as 15 ppm while the fragment mass tolerance was set as 0.05 Da. The enzyme was set to trypsin, allowing one missed cleavage, and the flexible modifications were set as deamidation (NQ) and oxidation (M). Carbamidomethylation (C) was set as a fixed modification. The criterion of peptide probability score is  $\geq 0.95$  (decoy target FDR, 0.05) so that low probability protein identifications can be filtered out. The N-linked glycosylation site is the Asp contained within the consensus N-linked glycosylation motif.

Statistical Analysis. To determine whether there was a significant difference in the glycoproteins of the AG and NAG-prostate cancer groups, *P*-values of the iTRAQ tag intensities representing the different groups were calculated using two-tailed *t*-tests.

Western Blot Analysis. A total of 25  $\mu$ g of proteins was resolved on SDS-PAGE and transferred electrophoretically onto a 0.2  $\mu$ m nitrocellulose membrane. The membrane was blocked by 5% nonfat milk/0.1% TBS-Tween 20 at room temperature for 2 h. The membrane was then probed with primary antibody (rabbit antihuman periostin at 1:2000,

mouse antihuman cathepsin L at 1:200) at 4°C overnight, followed by a three-time wash of 0.1% TBS-Tween 20. HRP conjugated secondary antibody was added at 1:2000 and incubated at room temperature for 1 h. Three washes of 0.1% TBS-Tween 20 were performed. The signal was visualized by a superSignal West Femto Maximum Sensitivity Substrate (Pierce).

Immunohistochemistry Staining. Rabbit antihuman periostin and goat anti-antibody were used in a dilution of 1:500 and 1:50 to stain sections from formalin-fixed and paraffin-embedded tissue specimens and were detected using the Dako LSAB+ System-AP kit according to the manufacturer's instructions.

## Results

Example 1: Quantitative Glycoproteomic Profiling of Aggressive (AG) Prostate Tumors and Nonaggressive (NAG) Prostate Tumors. The scheme of the present study is shown in FIG. 1. Briefly, to identify glycoproteins associated with AG-prostate cancer, glycoproteins from OCT-embedded AG-prostate tumor and NAG-prostate tumor tissues were quantitatively analyzed. The candidate proteins were further verified using Western blot and immunohistochemistry.

Glycopeptides were isolated from four AG-prostate tumors and four NAG-prostate tumors using solid-phase extraction of glycopeptides (SPEG). To determine the relative abundance of glycoproteins in AG and NAG-prostate cancer tissues, the glycopeptides isolated from each specimen were labeled with iTRAQ prior to LC-MS/MS analysis. From mass spectrometry analysis of glycopeptides, 102 unique N-linked glycopeptides were identified with 95% confidence, representing 79 unique glycoproteins (FIG. 7).

Example 2: Glycoprotein Changes Associated with Aggressive Prostate Tumors. To identify the glycoproteins associated with aggressive prostate tumors, the relative abundance of each glycopeptide in tissues of four cases of AG prostate cancer and four cases of NAG-prostate cancer were determined by iTRAQ labeling and tandem mass spectrometry (FIG. 7). With the use of the peak intensity of the iTRAQ reporter tags which represented the relative abundance of the peptides, *t* tests were performed to identify the glycopeptides and the glycoproteins associated with AG-prostate tumors. According to iTRAQ results, glycopeptides from three glycoproteins, microfibrillar-associated protein 4, periostin, and cathepsin L, showed a significant difference between the AG-prostate cancer group (iTRAQ Tag 113, 114, 115, and 121) and the NAG-prostate cancer group (iTRAQ Tag 116, 117, 118, and 119) (with *p*-values of less than 0.05) (FIGS. 2, 5, 7 and 8). Two peptides, vDLEDFEnNTAYAk and fnGSVSFFR (lower case v, f, and k represent the iTRAQ labeled

N-termini and Lys, lower case n in the nXT/S motif represents the formerly glycosylated Asp and deaminated after SPEG isolation), were identified from microfibrillar-associated protein 4. The quantitation of the two peptides was consistent: the average ratio of the AG-prostate tumor group vs the reference channel (iTRAQ tag 113, labeled as one of the AG-prostate tumors) was  $1.29 \pm 0.22$ , while the average ratio of the NAG-prostate tumor group was  $0.40 \pm 0.21$ .

Example 3: Verification of Glycoproteomic Results Using Western Blot. To verify the proteomic results, the four cases of AG-prostate tumors and four cases of NAG-prostate tumors were analyzed by Western blot assay. Periostin showed overexpression in all four AG-prostate tumors, while cathepsin L was elevated in three out of four AG-prostate tumors, compared to the NAG-prostate tumors (FIG. 3). A specific band for microfibrillar-associated protein 4 could not be identified by Western blot to verify its expression. IgG was detected evenly in each sample and was used as a quantitation control. These observations supported the glycoproteomic results for periostin and cathepsin L.

Example 4: Analysis of Overexpressed Proteins Using Immunohistochemistry. Cathepsin L and periostin were further analyzed in primary prostate tumors and normal prostate tissues using immunohistochemistry. FIG. 4A shows the cathepsin L staining in primary prostate tumor and its adjacent normal tissue. Staining was detected in both normal and tumor epithelial cells. However, the percentage of cells stained with cathepsin L was increased in the prostate tumor (FIG. 4A) with apparent increased staining in aggressive cancer, consistent with iTRAQ results (data not shown).

FIG. 4B shows the immunohistochemistry staining of periostin in primary prostate cancer tissues. The staining exhibited a low background in the normal prostate samples but revealed an overexpression in the peritumoral stroma of Gleason 3 tumors and strong overexpression in the peritumoral stroma of Gleason 4 tumors. This indicates that periostin expression may correlate with aggressive prostate cancer.

### Discussion

Several glycoproteins were identified in this study and showed significant altered expression in aggressive prostate tumors compared to nonaggressive prostate tumors. To the present inventors' knowledge, this is the first glycoproteomic study to elucidate the differentially expressed proteins associated with aggressive prostate cancer to identify proteins that are potentially useful for diagnosis of aggressive prostate cancer. Furthermore, though OCT-embedded tissues have been used in proteomic analyses, this study was the first to demonstrate the use of OCT-embedded frozen tissues in MS-based glycoproteomic

analysis. These findings may facilitate the usage of OCT-embedded frozen tissues in biomarker discovery.

Tumor invasion and metastasis is a complicated and multifaceted process which may include intravasation, survival in the circulatory system, arrest and extravasation into a new tissue, initiation and maintenance of tumor cell growth, and reactivation of angiogenesis. In this multistep process, interactions between cancer cells and stromal cells as well as between cancer cells and the extracellular matrix (ECM) are required. Therefore, component changes in the ECM within the tumor microenvironment could play a fundamental impact on the metastatic process.

Cathepsin L acts as endopeptidase, which can degrade many intracellular and extracellular proteins and thereby modify their function. Cathepsin L has been reported to be upregulated in a variety of malignancies including breast, lung, colon, gastric, head, and neck carcinomas, melanomas, and gliomas. Furthermore, overexpression of cathepsin L correlates positively with malignancy. Several studies have revealed that cathepsin L plays an important role in the processes of invasion and migration. The data show that cell-cell adhesion is diminished and degradation of ECM is increased when extracellular activity of cathepsin L increases, which suggests that cathepsin L can increase metastatic tumor development. Studies have also found that inhibition of cathepsin L mRNA decreased tumor growth of murine myeloma and that anticathepsin L ScFv (single chain variable fragment) inhibited the tumorigenic and metastatic phenotype of human melanoma.

Periostin is a unique and important ECM protein involved in cell development and adhesion. Periostin has been shown to interact with many other ECM proteins, including fibronectin, collagen V, tenascin-C and periostin itself. The epithelial-mesenchymal transition (EMT), which gives epithelial cancer cells invasive and metastatic potential, is one of the critical steps of tumor metastasis, and it has been shown that periostin can facilitate cell migration and differentiation during EMT. Periostin has also been found to be overexpressed in various types of human cancer, such as breast cancer, colon cancer, lung cancer, pancreatic cancer, prostate cancer, and ovarian cancer. Most recently, periostin was found to be overexpressed in prostate cancer tissue compared to benign prostate hyperplasia. However, our study is the first to discover the overexpression of periostin in aggressive prostate cancer using quantitative glycoproteomics and mass spectrometry. Consistent with the present glycoproteomics findings, periostin was found to promote tumor metastasis in colon cancer, melanoma, head and neck squamous cell carcinoma, gastric cancer, and lymph node metastases. For example, the average expression level of periostin is not increased in primary

tumors of melanoma, whereas periostin overexpression is found in around 60% of metastatic melanoma tumors in the liver or lymph nodes.

Microfibrillar-associated protein-4 (MFAP4) is an ECM protein that is upregulated in aggressive prostate cancer. MFAP4 binds to collagen and contains a C-terminal fibrinogen-like domain and an N-terminal integrin-binding motif. The fibrinogen-like domain is  
5 responsible for the carbohydrate binding activity. The N-terminal part of MFAP4 includes one cysteine-residue and a ligand motif Arg-Gly-Asp (RGD) for cell surface integrins. The function of MFAP4 is not yet clear. Human recombinant MFAP4 was reported to bind the collagen domain of surfactant protein A (SP-A), which suggests that MFAP4 may be  
10 involved in inflammatory processes. In a study of MFAP4 in catfish, Niu et al. reported that MFAP4 may play a novel role in teleost immune responses.

In this study, the couplegulation of cathepsin L and several ECM proteins, such as periostin, and microfibrillar-associated protein-4, in aggressive prostate tumor may indicate that there are some interactions among ECM proteins to facilitate the process of prostate  
15 tumor metastasis. Further studies will facilitate the understanding of the mechanisms involved in the function, regulation, and biological activities of cathepsin L and ECM proteins in tumor metastasis.

There has been a concern about using OCT-embedded frozen tissue in mass spectrometry analysis. OCT contains several chemicals to lower freezing temperature, such  
20 as PEG, which may suppress ionization and cause contamination of mass spectrometry analysis. Somiari and co-workers presented the first high-throughput proteomic analysis of human breast infiltrating ductal carcinoma using OCT-embedded biopsies, using two-dimensional gel electrophoresis (2-D DIGE) technology. Asomugha et al. used the same technologies for identification of crystalline modifications in the human lens cortex and  
25 nucleus. 2-D DIGE could separate the proteins from the OCT chemicals so as to avoid contaminating the mass spectrometers. However, to the best of our knowledge, directly analyzing OCT samples with mass spectrometry has not been reported. Our strategy of using solid phase extraction of N-glycopeptide overcomes this difficulty by chemical immobilization of glycopeptides and subsequent removing OCT in the washing step so that  
30 the glycopeptides isolated from OCT-embedded tissues could be isotopically labeled and directly injected to mass spectrometry instruments for protein identification and quantification. In our study, glycopeptides were readily identified from the OCT-embedded frozen tissues and no contamination was detected in the spectrum of mass spectrometry analysis. These results demonstrate (1) the protein glycosylation is preserved in the OCT-

embedded frozen tissues; (2) the OCT in the sample can be removed during the procedure of glycopeptide capture to eliminate the interference of OCT to the mass spectrometry analysis. Therefore, OCT-embedded frozen tissues can be used for glycoproteomic analysis and can provide invaluable specimen sources for identification of glycoprotein biomarkers.

5

We claim:

1. A method for aggressive prostate cancer in a patient comprising the steps of:
  - a. measuring the levels of one or more biomarkers in a sample collected from the  
5 patient; and
  - b. comparing the levels of the one or more biomarkers with predefined levels of  
the same biomarkers that correlate to a patient having aggressive prostate cancer and  
predefined levels of the same biomarkers that correlate to a patient not having aggressive  
prostate cancer, wherein a correlation to one of the predefined levels provides the diagnosis.  
10
2. The method of claim 1, wherein the one or more biomarkers is selected from the  
group consisting of cathepsin-L (CTSL), clusterin, neutrophil gelatinase associated lipocalin  
(NGAL), epithelial cell activating molecule (EpCAM), prostate specific antigen (PSA) and  
membrane metallo-endopeptidase (MME).  
15
3. The method of claim 1, wherein the one or more biomarkers comprises CTSL,  
clusterin, NGAL, EpCAM, PSA and MME.
4. The method of claims 2 or 3, wherein the one or more biomarkers further comprise  
20 one or more biomarkers selected from the group consisting of periostin, microfibrillar-  
associated protein 4 (MFAP4), collagen XII, neprilysin, and asporin (ASPN).
5. The method of claim 1, wherein the one or more biomarkers comprises CTSL,  
periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL, EpCAM, PSA MME, ASPN,  
25 or combinations thereof.
6. The method of claim 1, wherein the one or more biomarkers is selected from the  
group consisting of CTSL, periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL,  
EpCAM, PSA, MME and ASPN.  
30
7. The method of claim 1, wherein the one or more biomarkers comprises CTSL,  
periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL, EpCAM, PSA, MME and  
ASPN.

8. The method of any one of claims 2-6, wherein the one or more biomarkers further comprises one or more biomarkers from FIG. 9.
9. The method of any one of claims 2-6, wherein the one or more biomarkers further  
5 comprises one or more biomarkers from FIG. 6.
10. The method of any one of claims 2-6, wherein the one or more biomarkers further comprises one or more biomarkers from FIG. 7.
- 10 11. The method of any one of claims 2-6, wherein the one or more biomarkers further comprises one or more biomarkers from FIGS. 6, 7 and 9.
12. The method of claim 1, wherein the sample is a blood, plasma, or serum sample.
- 15 13. The method of claim 12, wherein the sample is a blood sample.
14. The method of claim 12, wherein the sample is a plasma sample.
15. The method of claim 12, wherein the sample is a serum sample.
- 20 16. The method of claim 1, wherein the sample is a tissue sample.
17. The method of claim 16, wherein the sample is an embedded tissue sample.
- 25 18. The method of any of claims 1-17, wherein the measuring step is performed using an immunoassay.
19. The method of any of claims 1-17, wherein the measuring step is performed using mass spectrometry.
- 30 20. The method of claim 18, wherein the mass spectrometry is multiple reaction monitoring mass spectrometry (MRM-MS).
21. The method of claim 1, wherein the correlation to a patient not having aggressive  
35 prostate cancer refers to a patient having non-aggressive prostate cancer.

22. The method of claim 1, wherein the correlation to a patient not having aggressive prostate cancer refers to a patient not having cancer.

23. A method for diagnosing aggressive prostate cancer in a patient comprising the steps  
5 of:

a. collecting a sample from the patient;

b. measuring the levels of a panel of biomarkers in the sample collected from the patient, wherein the panel of biomarkers comprises one or more of CTSL, clusterin, NGAL, EpCAM, PSA and MME; and

10 c. comparing the levels of the panel of biomarkers with predefined levels of the same panel of biomarkers that correlate to a patient having aggressive prostate cancer and predefined levels of the same panel of biomarkers that correlate to a patient not having aggressive prostate cancer, wherein a correlation to one of the predefined levels provides the diagnosis.

15

24. The method of claim 23, wherein the panel of biomarkers further comprises one or more of MFAP4, collagen XII, neprilysin, and ASPN.

25. A method for determining the aggressive prostate cancer status in a patient  
20 comprising the steps of:

a. collecting a sample from the patient;

b. measuring the levels of a panel of biomarkers in the sample collected from the patient, wherein the panel of biomarkers comprises one or more of CTSL, clusterin, NGAL, EpCAM, PSA and MME; and

25 c. comparing the levels of the panel of biomarkers with predefined levels of the same panel of biomarkers that correlate to one or more aggressive prostate cancer statuses selected from the group consisting of having aggressive prostate cancer, not having aggressive prostate cancer, progressing aggressive prostate cancer, and regressing aggressive prostate cancer, wherein a correlation to one of the predefined levels determines the  
30 aggressive prostate cancer status of the patient.

26. The method of claim 25, wherein the panel of biomarkers further comprises one or more of MFAP4, collagen XII, neprilysin, and ASPN.

27. A diagnostic kit for diagnosing aggressive prostate cancer in a patient comprising:
- a. a substrate for collecting a biological sample from the patient; and
  - b. means for measuring the levels of one or more of CTSL, periostin, MFAP4, collagen XII, neprilysin, clusterin, NGAL, EpCAM, PSA MME and ASPN.

5

28. The kit of claim 27, further comprising means for measuring the levels of one or more of the biomarkers in FIGS 6, 7 and 9.

10




FIG. 1




FIG. 2




FIG. 3




FIG. 4

Table 1. Glycoproteins Overexpressed in Aggressive (AG) Prostate Tumors

proteins	peptides <sup>a</sup>	AG-tumor <sup>b</sup>				NAG-tumor <sup>b</sup>				ratio of AG/NAG <sup>c</sup>
		113	114	115	121	116	117	118	119	
microfibrillar-associated protein 4	vDLEDPEaNTAYak	1.00	0.95	1.50	1.30	0.31	0.20	0.41	0.53	3.28
microfibrillar-associated protein 4	fnGSVVSFFR	1.00	1.56	1.29	1.70	0.14	0.34	0.38	0.87	3.22
periostin	eVnDTLLYNELk	1.00	0.57	1.24	0.54	0.26	0.17	0.34	0.52	2.62
cathepsin L	ySVAnDTGfVDIPk	1.00	0.52	0.96	1.33	0.38	0.44	0.78	0.62	1.85

<sup>a</sup> Lower case v, e, y, f, and k represent the iTRAQ labeled N-termini and Lys, lower case n in the nXT/S motif represents the formerly glycosylated Asp and delaminated after SPEP isolation. <sup>b</sup> The number showed the ratio of different channel vs 113. <sup>c</sup> The ratio of AG/NAG uses the average of each group.

FIG. 5

Protein Name	Swiss-Prot	Peptide Sequence
Seven transmembrane helix receptor	Q8NGK5	M.YLVAVVGN#VTILAVVK.I
Kallikrein 5 precursor	Q9Y337	R.DSCQGDSSGGPVVCN#GSLQ.G
Glandular kallikrein 1 precursor	P06870	E.PEN#FSFPDDLQCVDLK.I
Procollagen C-proteinase enhancer protein precursor	Q15113	G.QTPN#YTRPVFLCGGDVK.G
Glutamate decarboxylase 1	Q9BU91	-.MASSTPSSSATSSNAGADPN#TT.N
Thyroxine-binding globulin precursor	P05543	K.TTTVQVPMMHQMEQYYHLVDMELN#CTVLQMDYSK.N
Alpha-1-acid glycoprotein 2 precursor	P19652	P.ITN#ATLDR.I
scavenger receptor class B, member 2	Q14108	F.N#VTNPPEILR.G
Alpha-2-macroglobulin precursor	P01023	R.GNEANYYSN#ATTDEHGLVQFSIN#TTNVMGTSLTVR.V
Membrane copper amine oxidase	Q16853	K.EEEPSSSSVFNQNDPWAPTVDVDFSDFINN#ETIAGK.D
Splice isoform 1 of P24821 Tenascin precursor	P24821	K.GFEESEPVSGSFTTALDGPSGLVTAN#ITDSEALAR.W
Kallikrein 13 precursor	Q9UKR3	K.VLNTN#GTSGFLPGGYTCFPH.S
Presenilin-like protein 2	Q8TCT8	A.QEAILHASGN#GTTK.D
Beta-hexosaminidase beta chain precursor	P07686	R.GLETFSQLVYQDSYGTFTIN#ESTIIDSPR.F
Alpha-1-antitrypsin precursor	P01009	R.QLAHQSN#STNIFFSPVSIATAFAMLSLGTK.A
Splice isoform Beta-1A of P05556 Integrin beta-1 precursor	P05556	K.SCGECIQAGPNCGWCTN#STFLQEGMPTSAR.C
Prolargin precursor	P51888	R.IHYLYLQNNFITELPVESFQN#ATGLR.W
Alpha-galactosidase A precursor	P06280	F.QKPN#YTEIR.Q
Acid phosphatase, prostate	Q96QM0	K.VYDPLYCESVHN#FTLPSWATEDTMTK.L
CD44 isoform RC	O95370	D.LN#ITCR.F
Splice isoform 1 of Q9NZ08 Adipocyte-derived leucine aminopeptidase precursor	Q9NZ08	K.CFDAMEVDALN#SSHPVSTPVENPAQIR.E
Integrin alpha-1	P56199	K.EN#MTFGSTLVNPNNGGFLACGLYAYR.C
Splice isoform TGN51 of O43493 Trans-Golgi network integral membrane protein 2 precursor	O43493	K.QEEAGVRPSAGN#VSTHPSLSQRPGGSTK.S
Prostate stem cell antigen precursor	O43653	K.AQVSNEDECLQVEN#CTQLGEQCWTAR.I

FIG. 6

Protein Name	Swiss-Prot	Peptide Sequence
Splice isoform A of P07585 Decorin precursor	P07585	R.IADTN#ITSIPQGLPPSLTELHLDGNK.I
Splice isoform 1 of P24821 Tenascin precursor	P24821	R.QSGVN#ATLPEENQPVVFNHVYNIK.L
Acid phosphatase, prostate	Q96QM0	R.N#ETQHEPYPLMLPGCSPSCPLER.F
Splice isoform PSMA-1 of Q04609 Glutamate carboxypeptidase II	Q04609	K.FLYN#FTQIPHLAGTEQNFQLAK.Q
Carboxypeptidase H precursor	P16870	R.DLQGNPIAN#ATISVEGIDHDVTSK.D
Kallikrein 10 precursor	O43240	K.LLPLLMAQLWAAEAALLPQN#DTR.L
Polymeric-immunoglobulin receptor precursor (Poly-Ig receptor) (PIGR) [Contains: Secretory component]	P01833	K.QIGLYPVLVIDSSGYVNPV#YTGR.I
Clusterin precursor	P10909	R.QLEEFNL#QSSPFYFWMNGDR.I
membrane alanine aminopeptidase precursor	P15144	K.GPSTPLPEDPNWN#VTEFHHTPK.M
Nidogen-2 precursor	Q14112	K.TNIQGQVPYVPAN#FTAHSIPYK.E
RPE-spondin	Q96J64	R.CSGDGLDSDGN#QTLHWQAIGNPR.C
Integrin alpha-1	P56199	K.QTQVQVQYGEN#VTHEFNLNK.Y
Polymeric-immunoglobulin receptor precursor (Poly-Ig receptor) (PIGR) [Contains: Secretory component]	P01833	R.LSLLLEPGN#GTFTVILNQLTSR.D
Serotransferrin precursor	P02787	R.QQQHLFGSN#VTDCSGNFCLFR.S
BA209J19.1.1	Q9H1L6	R.VN#LTTNTIAVTQTLNAAAYNNR.F
Splice isoform 1 of Q9UBX7 Kallikrein 11 precursor	Q9UBX7	K.CENAYPGN#ITDTMVCASVQEGGK.D
Splice isoform 1 of P12111 Collagen alpha 3(VI) chain precursor	P12111	R.QLINALQIN#NTAVGHALVLPAGR.D
Splice isoform A of P07585 Decorin precursor	P07585	K.LGLSFNSISAVDN#GSLANTPHLR.E
CD166 antigen precursor	Q13740	K.IIISPEEN#VTLTCTAENQLER.T
Laminin alpha-5 chain precursor	O15230	R.AQQLLAN#STALEEAMLQEQQR.L
Alpha-2-HS-glycoprotein precursor	P02765	K.AALAAFNAQNN#GSNFQLEEISR.A
Membrane copper amine oxidase	Q16853	R.QPQPN#VSELVVGPLPHPSYMR.D
Putative transmembrane protein NMB precursor	Q14956	R.VSVNTAN#VTLGPQLMEVTVYR.R
Acid ceramidase precursor	Q13510	K.ILAPAYFILGGN#QSGEGCVITR.D
Chymase precursor	P23946	R.SITVTLGAHN#ITEEEDTWQK.L

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Complement C3 precursor [Contains: C3a anaphylatoxin]	P01024	K.TVLTPATNHMGN#VTFTIPANR.E
Apolipoprotein D precursor	P05090	R.ADGTVNQIEGEATPVN#LTEPAK.L
Fibrillin 1 precursor	P35555	R.NYYADN#QTCDGELLFN#MTK.K
Hypothetical protein FLJ20421	Q9NX62	K.QVALQTFGN#QTTIIPAGGAGYK.V
Splice isoform 1 of P02751 Fibronectin precursor	P02751	K.LDAPTNLQFVN#ETDSTVLVR.W
Cell surface glycoprotein MUC18 precursor	P43121	K.CGLSQSQGN#LSHVDWFSVHK.E
Hypothetical protein FLJ90478	Q8NC56	K.CIPVMEAEQYIAN#VTSSSSAK.F
BA209J19.1.1	Q9H1L6	R.SLGSGGSVSQLFSN#FTGSVDDR.G
Unknown		R.TLN#QSSDELQLSMGNAMFVK.E
glycoprotein 2 (zymogen granule membrane)	P55259	R.QDLN#SSDVHSLQPQLDCGPR.E
CD166 antigen precursor	Q13740	K.LGDCISEDSYPDGN#ITWYR.N
Dipeptidyl-peptidase II precursor	Q9UHL4	R.ALAGLVYN#ASGSEHCYDIYR.L
similar to embryonic blastocoelar extracellular matrix protein precursor	Q7Z341	K.GSSSSEPMVPPQSHHN#DSSEV.-
Plasma protease C1 inhibitor precursor	P05155	R.ASSNPN#ATSSSSQDPESLQDR.G
Splice isoform 1 of P12111 Collagen alpha 3(VI) chain precursor	P12111	R.VAVVQHAPSESVDN#ASMPPVK.V
EW12	Q96EW3	R.IGPGEPELLCN#VSGALPPAGR.H
Basement membrane-specific heparan sulfate proteoglycan core protein precursor	P98160	R.SLTQGSLIVGDLAPVN#GTSQGK.F
Splice isoform A of Q9UL36 Zinc finger protein 236	Q9UL36	R.CDQCPQTFNVEFN#LTLHK.C
Ceruloplasmin precursor	P00450	K.EN#LTAPGSDSAVFFEQGTTR.I
Integrin alpha-5 precursor	P08648	K.EPLSDPVGTCYLSTDN#FTR.I
Plasma protease C1 inhibitor precursor	P05155	R.VLSN#NSDANLELINTWVAK.N
Dipeptidyl peptidase IV	P27487	R.IQN#YSVMDICDYDESSGR.W
biotinidase precursor	P43251	K.NPVGLIGAEN#ATGETDPSHSK.F
Splice isoform OA3-323 of Q08722 Leukocyte surface antigen CD47 precursor	Q08722	K.SDAVSHTGN#YTCEVTELTR.E

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 3	P08754	R.EYQLN#DSASYLNDLDR.I
Costimulatory molecule	Q9BXR1	K.QLVHSFAEQDQGSAYAN#R.T
Membrane copper amine oxidase	Q16853	R.IQMLSFAGEPLPQN#SSMAR.G
Vacuolar ATP synthase subunit S1 precursor	Q15904	K.QPVSPVIHPPVSYN#DTAPR.I
HLA class II histocompatibility antigen, DR-1 beta chain precursor	P01912	R.FLEYSTSECHFFN#GTER.V
Splice isoform 1 of P20151 Glandular kallikrein 2 precursor	P20151	R.VPVSHSFPHPLYN#MSLLK.H
Alpha-2-glycoprotein 1, zinc	P25311	K.DIVEYYN#DSN#GSHVLQGR.F
Hypothetical protein	Q9H670	R.TYLVN#SSDSGSSQTESPSSK.Y
Dihydropyridine-sensitive L-type, calcium channel alpha-2/delta subunits precursor	P54289	K.QSCITEQTQYFFDN#DSK.S
Tripeptidyl-peptidase I precursor	O14773	K.FLSSSPHLPSSYFN#ASGR.A
Tetraspanin 1	O60635	K.DYGSQEDFTQVWN#TTMK.G
Myosin heavy chain, smooth muscle isoform	P35749	K.NMDPLNDN#VTSLN#ASSDK.F
Lactotransferrin precursor (Lactoferrin) [Contains: Lactoferroxin A; Lactoferroxin B; Lactoferroxin C]	P02788	R.TAGWNIPMGLLFN#QTGSCK.F
Tetraspanin 1	O60635	K.CCGFTN#YTDFFEDSPYFK.E
150 kDa oxygen-regulated protein precursor	Q9Y4L1	K.EN#GTDTVQEEEESPAEGSK.D
Laminin beta-1 chain precursor	P07942	R.VN#ASTTEPN#STVEQSALMR.D
Splice isoform 1 of O00584 Ribonuclease 6 precursor	O00584	K.QDQQLQN#CTEPGEQSPK.Q
scavenger receptor class B, member 2	Q14108	R.TMVPVMYLN#ESVHIDK.E
MAC-2 binding protein precursor	Q08380	K.EPGSN#VTMSVDAECVPMVR.D
Podocalyxin-like protein 1 precursor	O00592	K.QLVLN#LTGNTLCAGGASDEK.L
Fibrillin 1 precursor	P35555	K.AWGTPCEMCPAVN#TSEYK.I
Plasma kallikrein precursor	P03952	K.IYPGVDFGGEELN#VTFVK.G

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
monogenic, audiogenic seizure susceptibility 1 homolog	Q8TF58	R.DSGTGLMMSVN#FSTQELR.S
Probable serine carboxypeptidase CPVL precursor	Q9H3G5	R.QAIHVG#QTFNDGTIVEK.Y
Splice isoform LAMP-2A of P13473 Lysosome-associated membrane glycoprotein 2 precursor	P13473	K.VASVININPN#TTHSTGSCR.S
scavenger receptor class B, member 2	Q14108	K.CNMIN#GTDGDSFHPLTK.D
Histidine-rich glycoprotein precursor	P04196	R.VIDFN#CTTSSVSSALANTK.D
Splice isoform 1 of Q9UBX7 Kallikrein 11 precursor	Q9UBX7	R.TATESFPHPGFN#NSLPNK.D
Splice isoform PSMA-1 of Q04609 Glutamate carboxypeptidase II	Q04609	R.GVAYINADSSIEGN#YTLR.V
Laminin gamma-1 chain precursor	P11047	R.TLAGEN#QTAFEIEELNR.K
Polymeric-immunoglobulin receptor precursor (Poly-Ig receptor) (PIGR) [Contains: Secretory component]	P01833	K.WN#NTGCCQALPSQDEGPSK.A
Splice isoform 1 of P24821 Tenascin precursor	P24821	R.LN#YSLPTGQWVGVQLPR.N
Prostaglandin-H2 D-isomerase precursor	P41222	K.SVVAPATDGGLN#LTSTFLR.K
Dihydropyridine-sensitive L-type, calcium channel alpha-2/delta subunits precursor	P54289	R.SLDNDNYVFTAPYFN#K.S
Hypothetical protein	Q86V91	K.QDILN#NSLTTLSQDITK.V
Splice isoform 1 of Q92859 Neogenin precursor	Q92859	R.TLSDVPSAAPQN#LSLEVR.N
Fibrinogen beta chain precursor [Contains: Fibrinopeptide B]	P02675	R.GTAGNALMDGASQLMGEN#R.T
Ceruloplasmin precursor	P00450	K.EHEGAIYPDN#TTDFQR.A
Laminin beta-2 chain precursor	P55268	R.N#TSAASTAQLVEATEELR.R
Lysosomal alpha-glucosidase precursor	P10253	K.LEN#LSSSEMGYTATLTR.T
Alpha-1-acid glycoprotein 2 precursor	P19652	R.QNQCFYN#SSYLNVR.E
Tumor endothelial marker 7-related precursor	Q96PD9	R.VN#LSFDFPFYGHFLR.E

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Alpha-1-acid glycoprotein 1 precursor	P02763	R.QDQCIYN#TTYLNVQR.E
scavenger receptor class B, member 2	Q14108	K.ANIQFGDN#GTTISAVSNK.A
Cartilage oligomeric matrix protein precursor	P49747	R.CN#DTIPEYETHQLR.Q
Splice isoform 1 of Q16563 Pantophysin	Q16563	K.GQTEIQVNCPPAVTEN#K.T
O-acetyltransferase	Q96PB1	K.IDAYNEAAVSILN#SSTR.N
Intercellular adhesion molecule-2 precursor	P13598	K.AAPAPQEATATFN#STADR.E
Splice isoform PSMA-1 of Q04609 Glutamate carboxypeptidase II	Q04609	K.VPYNVGPFTGN#FSTQK.V
BA209J19.1.1	Q9H1L6	K.LN#DTTLQVLNTWYTK.Q
Antithrombin-III precursor	P01008	K.LGACN#DTLQQLMEVFK.F
Hypothetical protein	Q8WUM6	R.NPCTSEQN#CTSPFSYK.N
Prostate-specific membrane antigen-like protein	Q9HBA9	K.VSYNVGPFTGN#FSTQK.V
Laminin beta-2 chain precursor	P55268	R.AN#TSALAVPSPVNSASAR.H
Tumor-associated calcium signal transducer 1	Q96C47	K.QCN#GTSTCWCVNTAGVR.R
Carboxypeptidase D precursor	O75976	K.DLDTDFTNN#ASQPETK.A
Haptoglobin precursor	P00738	K.VVLHPN#YSQVDIGLIK.L
Lysosome-associated membrane glycoprotein 1 precursor	P11279	K.N#MTFDLPSDATVVLN#R.S
Cadherin-13 precursor	P55290	K.IN#NTHALVSLQLNLNK.A
Endoplasmic precursor	P14625	R.EEEAIQLDGLN#ASQIR.E
EMILIN 1 precursor	Q9Y6C2	R.LEQLGGLLAN#VSGELGGR.L
Hypothetical protein	Q96C67	K.AMN#ASAN#ITSDGVEVLGK.M
Complement C4 precursor [Contains: C4a anaphylatoxin]	P01028	R.FSDGLESN#SSTQFEVK.K
Splice isoform 2C2A' of P12110 Collagen alpha 2(VI) chain precursor	P12110	R.GTFTDCALAN#MTEQIR.Q
Kallikrein 5 precursor	Q9Y337	K.DVRPIN#VSSHCPASAGTK.C
FLJ00343 protein	Q96C61	R.VAN#PSGN#LTETYVQDR.G
Splice isoform B of P23142 Fibulin-1 precursor	P23142	R.CATPHGDN#ASLEATFVK.R
Alpha-1-antitrypsin precursor	P01009	K.YLGN#ATAIFFLPDEGK.L

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Metalloproteinase inhibitor 1 precursor	P01033	K.FVGTPEVN#QTTLYQR.Y
Cleft lip and palate associated transmembrane protein 1	Q9BSS5	K.DYYPIN#ESLASPLR.V
Laminin alpha-5 chain precursor	O15230	R.LN#TTGVSAGCTADLLVGR.A
Integrin alpha-1	P56199	K.LDLPVN#TSIPN#VTEVK.E
Protein FAM3D precursor	Q96BQ1	R.GLNIALVN#GTTGAVLGQK.A
Carboxypeptidase B-like protein similar to collagen type XIV	Q9P2Y6	K.QVHFFVN#ASDVDNVK.A
Integral membrane protein 2B (Transmembrane protein BRI) [Contains: ABri/ADan amyloid peptide]	Q05707	R.SFMVNV#WTHAPGNVEK.Y
Kallikrein 6 precursor	Q9Y287	K.CYVIPLN#TSIVMPPR.N
laminin alpha 2 subunit precursor	Q92876	R.DCSAN#TTSCHILGWGK.T
CD59 glycoprotein precursor	P24043	R.YMQN#LTVEQPIEVK.K
Hypothetical protein FLJ90516	P13987	K.TAVN#CSSDFDACLITK.A
Apolipoprotein B-100 precursor (Apo B-100) [Contains: Apolipoprotein B-48 (Apo B-48)]	Q8NC34	R.TALFPDLAQGN#ASLR.L
Clusterin precursor	P04114	R.FN#SSYLQGTNQTGR.Y
Laminin gamma-1 chain precursor	P10909	R.LAN#LTQGEDQYYLR.V
Hemopexin precursor	P11047	K.TAN#DTSTEAYNLLLR.T
Splice isoform 2A of Q02487 Desmocollin 2A/2B precursor	P02790	K.ALPQPQN#VTSLLGCTH.-
Protein-lysine 6-oxidase precursor	Q02487	K.NGIYN#ITVLASDQGGR.T
hypothetical protein DKFZp313G1735	P28300	R.DPGAAPVGAAN#ASAQQPR.T
Lutheran blood group glycoprotein precursor	Q8N3Q8	K.QSIGQN#YSNVIANLR.W
Dopamine beta-monoxygenase precursor	P50895	R.TQN#FTLLVQGSPELK.T
MAC-2 binding protein precursor	P09172	R.SLEAIN#GSGLQMGLQR.V
EMILIN 1 precursor	Q08380	R.TVIRPFYLTN#SSGVD.-
Alpha-2-HS-glycoprotein precursor	Q9Y6C2	R.ETN#TTSQMQAALLEK.L
FLJ00343 protein	P02765	K.VCQDCPLLAPLN#DTR.V
	Q96C61	K.VTAQGPGLPSGNIAN#K.T

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Splice isoform Short of Q99715 Collagen alpha 1(XII) chain precursor	Q99715	R.NLQVYN#ATSNSLTVK.W
Biglycan precursor	P21810	K.LLQVVYLHSNN#ITK.V
apolipoprotein F	Q13790	R.QGGVN#ATQVLIQHLR.G
Splice isoform 1 of Q92932 Receptor-type protein-tyrosine phosphatase N2 precursor	Q92932	K.VSANVQN#VTTEDVEK.A
Blood plasma glutamate carboxypeptidase precursor	Q9UNM8	K.IVVYNQPYIN#YSR.T
Angiotensin-converting enzyme, somatic isoform precursor	P12821	R.VTN#DTESDINYLK.M
similar to hypothetical protein 6720478C22		R.TLLN#ASCDNMLMGIK.S
similar to Peroxidasin CG12002- PA	Q92626	R.QGEHLSN#STSAFSTR.S
Fibrillin 1 precursor	P35555	R.VLPVN#VTDYCQLVR.Y
Integrin alpha-V precursor	P06756	K.AN#TTQPGIVEGGQVLK.C
Integrin alpha-5 precursor	P08648	R.VTGLN#CTTNHPINPK.G
Desmoglein 2 precursor	Q14126	K.IN#ATDADEPNTLNSK.I
Cation-dependent mannose-6- phosphate receptor precursor	P20645	R.EAGN#HTSGAGLVQIN#K.S
solute carrier family 39 (zinc transporter), member 6	Q8IXR3	R.NTNENPQECFN#ASK.L
Transcobalamin I precursor	P20061	K.MN#DTIFGFTMEER.S
Beta-1,3-N- acetylgalactosaminyltransferase	Q8TDY1	R.QDFLDTYNN#LTLK.T
Glutaminy-peptide cyclotransferase precursor	Q16769	K.NYHQPAILN#SSALR.Q
Laminin gamma-1 chain precursor	P11047	R.CDQCEENYFYN#R.S
Splice isoform Short of Q99715 Collagen alpha 1(XII) chain precursor	Q99715	K.EAGN#ITTDGYEILGK.L
Integrin-like protein	Q8WWJ8	K.LTN#NSNQFQTEVGK.Q
Hypothetical protein KIAA0315	O15031	K.SCVAVTSAQPQN#MSR.R
Biglycan precursor	P21810	R.MIEN#GSLFLPTLR.E
Low density lipoprotein receptor related protein-deleted in tumor	Q8WY28	R.VGMDGTN#QSVVIETK.I
Lysosomal alpha-glucosidase precursor	P10253	R.GVFITN#ETGQPLIGK.V

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Splice isoform Sap-mu-0 of P07602 Proactivator polypeptide precursor [Contains: Saposin A (Protein A); Saposin B (Sphingolipid activator protein 1) (SAP-1) (Cerebroside sulfate activator) (CSAct) (Dispersin) (Sulfatide/GM1 activator); Saposin C (Co-beta-glucosidase) (A1 activator) (Glucosylceramidase activator) (Sphingolipid activator protein 2) (SAP-2); Saposin D (Protein C) (Component C)]	P07602	R.TN#STFVQALVEHVK.E
DJ68D18.2.5 (CD44 antigen CTL2 gene)	Q9H5A6	K.AFN#STLPTMAQMEK.A
Fibroleukin precursor	Q8IWA5	K.TCNPETFPSSN#ESR.Q
Laminin gamma-1 chain precursor	Q14314	R.LHVGNYN#GTAGDALR.F
EMILIN 3 precursor	P11047	K.IPAIN#QTITEANEK.T
Splice isoform Long of P36980 Complement factor H-related protein 2 precursor	Q9H8L6	R.QVEELQVN#HTALR.E
EMILIN 1 precursor	P36980	R.LQNNENN#ISCV.R.G
Apolipoprotein B-100 precursor (Apo B-100) [Contains: Apolipoprotein B-48 (Apo B-48)]	Q9Y6C2	R.LGALN#SSLQLEDR.L
Cathepsin L precursor	P04114	K.YDFN#SSMLYSTAK.G
Stromal cell-derived receptor-1 alpha	P07711	K.YSVAN#DTGFVDIPK.Q
150 kDa oxygen-regulated protein precursor	Q9Y639	K.ENGMPMDIVN#TSGR.F
Asporin precursor	Q9Y4L1	R.AEPPLN#ASASDQGEK.V
Splice isoform 1 of P08603 Complement factor H precursor	Q9BXN1	K.ITDIEN#GSLANIPR.V
M130 antigen cytoplasmic variant 2 precursor	P08603	R.ISEEN#ETTCYMGK.W
Splice isoform Lamin A of P02545 Lamin A/C	Q07900	K.EDAAVN#CTDISVQK.T
Splice isoform V0 of P13611 Versican core protein precursor	P02545	R.TALIN#STGEEVAMR.K
glycoprotein 2 (zymogen granule membrane)	P13611	R.FEN#QTGFPPPSR.F
	P55259	R.DPN#CSSILQTEER.N

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Melanoma-associated chondroitin sulfate proteoglycan	Q92675	R.LDPTVLDAGELAN#R.T
Splice isoform Beta-1A of P05556 Integrin beta-1 precursor	P05556	K.CHEGN#GTFECGACR.C
Integrin alpha-5 precursor	P08648	K.NLN#NSQSDVVSFR.L
Inter-alpha-trypsin inhibitor heavy chain H2 precursor	P19823	K.GAFISN#FSMTVDGK.T
Biliary glycoprotein	Q13857	K.LSQGN#TTLSINPVK.R
solute carrier family 39 (zinc transporter), member 6	Q8IXR3	R.YGEN#NLSVEGFR.K
Acid ceramidase precursor	Q13510	R.TVLEN#STSYEEAK.N
Apolipoprotein D precursor	P05090	R.CIQAN#YSLMENGK.I
Splice isoform 1 of P24821 Tenascin precursor	P24821	K.VEAAQN#LTLPGSLR.A
Beta-2-glycoprotein I precursor	P02749	R.VYKPSAGN#NSLYR.D
Sel-1 homolog precursor	Q9UBV2	R.EASIVGEN#ETYPR.A
Integrin alpha-1	P56199	R.SEN#ASLVLSSSNQK.R
Di-N-acetylchitobiase precursor	Q01459	K.QIN#SSISGNLWDK.D
Haptoglobin precursor	P00738	K.NLFLN#HSEN#ATAK.D
Splice isoform 1 of Q8NBK3 Sulfatase modifying factor 1 precursor	Q8NBK3	K.FVN#STGYLTEAEK.F
Complement C2 precursor	P06681	K.QSVAHFVALN#GSK.L
Carboxypeptidase H precursor	P16870	K.GN#ETIVNLIHSTR.I
Hypothetical protein	Q8N382	R.LAVTN#TTMTGTVLK.M
Splice isoform 1 of Q12860 Contactin precursor	Q12860	K.AN#STGTLVITDPTR.I
Kallikrein 5 precursor	Q9Y337	K.VLQCLN#ISVLSQK.R
Skeletal muscle LIM-protein 1	Q13642	K.CLHPLAN#ETFVAK.D
Hypothetical protein FLJ11273	Q8N4L0	R.LNN#ITIIGPLDMK.Q
Plasma kallikrein precursor	P03952	R.IYSGILN#LSDITK.D
Splice isoform A of O60245 Protocadherin 7 precursor	O60245	K.IDN#LTGELSTSER.R
Desmoglein 2 precursor	Q14126	K.DTGELN#VTSILDR.E
Splice isoform HMW of P01042 Kininogen precursor (Alpha-2-thiol proteinase inhibitor) [Contains: Bradykinin]	P01042	K.LNAENN#ATFYFK.I
Laminin alpha-5 chain precursor	O15230	R.LN#ASIADLQSQLR.S

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Leucine-rich alpha-2-glycoprotein precursor	P02750	K.LPPGLLAN#FTLLR.T
Angiotensin-converting enzyme, somatic isoform precursor	P12821	R.QQYNALLSN#MSR.I
Adipocyte plasma membrane-associated protein	Q9HDC9	R.AGPN#GTLFVADAYK.G
laminin alpha 2 subunit precursor	P24043	K.QEGILYVDGASN#R.T
Serotransferrin precursor	P02787	K.CGLVPVLAENYN#K.S
Splice isoform Short of Q13459 Myosin IXb	Q13459	K.LGFSSPYEGVLN#K.S
Transferrin receptor protein 1	P02786	K.QNNGAFN#ETLFR.N
Similar to RIKEN cDNA 1300012G16	Q8NHP8	R.SDLNPAN#GSYPFK.A
Human full-length cDNA 5-PRIME end of clone CS0DM009YC13 of fetal liver of Homo sapiens	P29622	K.FLN#DTMAVYEAK.L
Splice isoform A of P12314 High affinity immunoglobulin gamma Fc receptor I precursor	P12314	R.ITSASVN#DSGEYR.C
Polymeric-immunoglobulin receptor precursor (Poly-Ig receptor) (PIGR) [Contains: Secretory component]	P01833	K.VPGN#VTAVLGETLK.V
Splice isoform OSF-2OS of Q15063 Osteoblast specific factor 2 precursor	Q8IZF9	K.EVN#DTLLVNELK.S
Laminin alpha-5 chain precursor	O15230	R.DN#ATLQATLHAAR.D
Splice isoform 1 of Q12884 Seprase	Q12884	K.SVN#ASNYGLSPDR.Q
HLA class II histocompatibility antigen, DRB1-4 beta chain precursor	P13760	K.HECHFFN#GTER.V
Splice isoform 1 of Q02083 N-acylsphingosine amidohydrolase-like precursor	Q02083	R.FN#VSLDSVPELR.W
Bactericidal/permeability-increasing protein-like 1 precursor	Q8N4F0	R.SDDNLLN#TSALGR.L
Nidogen-2 precursor	Q14112	R.IHQN#ITYQVCR.H
Splice isoform 2C2A' of P12110 Collagen alpha 2(VI) chain precursor	P12110	R.N#MTLFSDLVAEK.F

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Splice isoform 1 of P08603 Complement factor H precursor	P08603	K.MDGASN#VTCINSR.W
Integrin alpha-1	P56199	R.YN#HTGQVIIYR.M
CD63 antigen	P08962	K.CCGAAN#YTDWEK.I
Urokinase receptor-associated protein UPARAP	Q9UBG0	R.VTPACN#TSLPAQR.W
Splice isoform HMW of P01042 Kininogen precursor (Alpha-2- thiol proteinase inhibitor) [Contains: Bradykinin]	P01042	R.ITYSIVQTN#CSK.E
Myosin light chain 1, slow- twitch muscle A isoform	P14649	R.ALGQN#PTNAEVLK.V
maba1	Q8N1G3	K.MSVCTDN#VTDLR.I
MAC-2 binding protein precursor	Q08380	R.ALGFEN#ATQALGR.A
Hemopexin precursor	P02790	R.SWPAVGN#CSSALR.W
Low-density lipoprotein receptor-related protein 1 precursor	Q07954	R.FN#STEYQVVTR.V
Procollagen-lysine,2- oxoglutarate 5-dioxygenase 3 precursor	O60568	K.EQYIHEN#YSR.A
Lysosomal acid phosphatase precursor	P11117	R.QTPEYQN#ESSR.N
Complement component C8 alpha chain precursor	P07357	R.GGSSGWSGGLAQN#R.S
O-acetyltransferase	Q96PB1	K.MN#ITSIAPLLEK.L
Unknown		K.SECHFFN#GTER.V
Transmembrane 4 superfamily, member 3	P19075	R.IVN#ETLYENTK.L
Splice isoform 1 of O75882 Attractin precursor	O75882	K.IDSTGN#VTNELR.V
Splice isoform H17 of P05164 Myeloperoxidase precursor	P05164	R.SCPACPGSN#ITIR.N
Lymphocyte antigen	Q30161	R.QECYAFN#GTQR.F
membrane metallo- endopeptidase	P08473	K.EIAN#ATAKPEDR.N
Membrane copper amine oxidase	Q16853	R.YLYLASN#HSNK.W
similar to agrin precursor	Q96IC1	K.NELMLN#SSLMR.I
Dipeptidyl-peptidase I precursor	P53634	R.DVN#CSVMPQEK.K

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Low-density lipoprotein receptor-related protein 1 precursor	Q07954	R.LN#GTDPIVAADSK.R
Hypothetical protein NT2RP3000266	Q8NBJ4	K.AVLVNN#ITTGER.L
FLJ00343 protein	Q96C61	K.LPQLPITN#FSR.D
membrane alanine aminopeptidase precursor	P15144	K.AEFN#ITLIHPK.D
Beta-sarcoglycan	Q16585	R.ITSN#ATSDLNIK.V
MAC-2 binding protein precursor	Q08380	K.AAIPSAIDTN#SSK.S
Splice isoform Long of P08236 Beta-glucuronidase precursor	P08236	K.VVAN#GTGTQGQLK.V
Sad1/unc-84-like protein 2	Q9UH99	K.ALSPN#STISSAPK.D
Splice isoform 1 of P18206 Vinculin	P18206	K.AVAGN#ISDPGLQK.S
Splice isoform 1 of Q9UBX7 Kallikrein 11 precursor	Q9UBX7	R.CAN#ITIEHQK.C
5'-nucleotidase precursor	P21589	K.LDN#YSTQELGK.T
Lymphatic endothelium-specific hyaluronan receptor LYVE-1	Q9Y5Y7	K.ANQQLN#FTEAK.E
Fibromodulin precursor	Q06828	R.LYLDHNN#LTR.M
Splice isoform OA3-323 of Q08722 Leukocyte surface antigen CD47 precursor	Q08722	R.DIYTFD GALN#K.S
Similar to I factor	Q8WW88	K.FLNN#GTCTAEGK.F
Collagen alpha 1(VI) chain precursor	P12109	R.N#FTAADW GQSR.D
Low-density lipoprotein receptor-related protein 1 precursor	Q07954	R.IETILLN#GTDR.K
similar to RIKEN cDNA 2010200I23		R.VN#ITYNYPVR.A
Transmembrane protease, serine 2 precursor	O15393	R.CIACGVNLN#SSR.Q
immunoglobulin J chain	P01591	R.EN#ISDPTSPLR.T
Myosin regulatory light chain	O14950	K.LN#GTD PEDVIR.N
Integrin alpha-1	P56199	K.VYVYALN#QTR.F
Carboxypeptidase D precursor	O75976	R.FANEYPN#ITR.L
Laminin gamma-1 chain precursor	P11047	R.VN#NTLSSQISR.L
Splice isoform Short of Q99715 Collagen alpha 1(XII) chain precursor	Q99715	K.MLEAYN#LTEK.N

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Dipeptidyl peptidase IV	P27487	K.LDFIILN#ETK.F
Hypothetical protein KIAA0960	O43376	R.CPN#SSALQEV.R.S
Afamin precursor	P43652	R.DIENFN#STQK.F
membrane metallo- endopeptidase	P08473	R.SCIN#ESAIDSR.G
150 kDa oxygen-regulated protein precursor	Q9Y4L1	R.VFGSQN#LTTVK.L
Complement C2 precursor	P06681	K.TMFPN#LTDVR.E
Beta-2-glycoprotein I precursor	P02749	K.LGN#WSAMPSC.A
ASH1	Q9NR48	R.EGATAN#VSEGEK.K
Factor VII active site mutant immunoconjugate	Q96PQ8	R.EEQYN#STYR.V
membrane alanine aminopeptidase precursor	P15144	K.LN#YTLSQGHR.V
Beta-galactosidase precursor	P16278	R.NNVITLN#ITGK.A
Neutrophil gelatinase- associated lipocalin precursor	P80188	K.SYN#VTSVLF.R.K
Lysosomal acid phosphatase precursor	P11117	R.YEQLQN#ETR.Q
Transmembrane 9 superfamily protein member 3 precursor	Q9HD45	R.IVDVN#LTSEGK.V
Ig gamma-4 chain C region	P01861	R.EEQFN#STYR.V
Human full-length cDNA clone CS0DI019YF20 of placenta of Homo sapiens	Q86TT2	R.EEQYN#STFR.V
Lysosomal alpha-glucosidase precursor	P10253	R.N#NTIVNELVR.V
FLJ00385 protein	Q8NF17	R.EEQFN#STFR.V
Neuroserpin precursor	Q99574	K.DAN#LTGLSDNK.E
CD63 antigen	P08962	K.NN#HTASILDR.M
Integrin alpha-1	P56199	R.N#TTFNVESTK.K
Follistatin-related protein 1 precursor	Q12841	K.GSN#YSEILDK.Y
biotinidase precursor	P43251	R.FN#DTEVLQR.L
Splice isoform LAMP-2A of P13473 Lysosome-associated membrane glycoprotein 2 precursor	P13473	R.VQPFN#VTQGK.Y
Beta-hexosaminidase alpha chain precursor	P06865	K.SAEGTFFIN#K.T
Cathepsin D precursor	P07339	K.GSLSYLN#VTR.K

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Complement C4 precursor [Contains: C4a anaphylatoxin]	P01028	R.GLN#VTLSTGR.N
Transmembrane 4 superfamily, member 6	O43657	K.QYN#STGDYR.S
laminin alpha 2 subunit precursor	P24043	K.LSAIPN#DTAAK.L
similar to Prostaglandin F2 receptor negative regulator precursor (Prostaglandin F2- alpha receptor regulatory protein) (Prostaglandin F2-alpha receptor associated protein) (CD9 partner 1) (CD9P-1)	Q9P2B2	K.LEN#WTDASR.V
vimentin	P08670	R.QDVDN#ASLAR.L
AD-017 protein	Q9P0I5	R.QN#ITNQLEK.W
Low-density lipoprotein receptor-related protein 1 precursor	Q07954	K.DN#ATDSVPLR.T
TSLC1-like 2	Q8NFZ8	R.QTLFFN#GTR.A
Splice isoform LAMP-2A of P13473 Lysosome-associated membrane glycoprotein 2 precursor	P13473	K.WQMN#FTVR.Y
Alpha-2-glycoprotein 1, zinc	P25311	R.FGCEIENN#R.S
Mucin 2 precursor	Q02817	K.YN#NTVEIVK.V
membrane alanine aminopeptidase precursor	P15144	R.N#ATLVNEADK.L
Splice isoform Beta-1A of P05556 Integrin beta-1 precursor	P05556	K.NGVN#GTGENGR.K
Procollagen-lysine,2- oxoglutarate 5-dioxygenase 2 precursor	O00469	R.YN#CSIESPR.K
Fibrillin 2 precursor	P35556	R.N#CTDIDECR.I
Hypothetical protein FLJ21551	Q9H717	K.QVIQN#VTHK.D
similar to RIKEN cDNA 2010200I23		R.QGN#YSAGLPR.G
Tissue factor precursor	P13726	K.VN#VTVEDER.T
Microfibril-associated glycoprotein 4 precursor	P55083	R.FN#GSVSFFR.G
CTL2 gene	Q8IWA5	K.N#ITDLVEGAK.K
Hypothetical protein	Q8N348	R.GFLN#FTLAR.A

FIG. 6 cont'd.

Protein Name	Swiss-Prot	Peptide Sequence
Splice isoform Vascular of P49961 Ectonucleoside triphosphate diphosphohydrolase 1	P49961	K.VVN#VSDLYK.T
Laminin gamma-1 chain precursor	P11047	K.LLNN#LTSIK.I
Fibroleukin precursor	Q14314	R.LDGSTN#FTR.T
Ceroid-lipofuscinosis neuronal protein 5	O75503	K.NIETN#YTR.I
Hypothetical protein DKFZp686C02218	Q7Z374	K.TPLTAN#ITK.S
Splice isoform 1 of P24821 Tenascin precursor	P24821	R.N#TTSYVLR.G
Hypothetical protein	Q9Y4J9	K.LN#VSDLYK.L
Clusterin precursor	P10909	K.EDALN#ETR.E
Splice isoform 1 of P12111 Collagen alpha 3(VI) chain precursor	P12111	K.QN#LTVTDR.V
Mimecan precursor	P20774	K.AN#DTSYIR.D
similar to hypothetical protein	Q9UF56	R.N#LSSLDLR.H
Vitronectin precursor (Serum spreading factor) (S-protein) (V75) [Contains: Vitronectin V65 subunit; Vitronectin V10 subunit; Somatomedin B]	P04004	K.N#GSLFAFR.G
Splice isoform PSMA-1 of Q04609 Glutamate carboxypeptidase II	Q04609	K.N#FTEIASK.F
Plasma protease C1 inhibitor precursor	P05155	R.DTFVN#ASR.T
Golgi apparatus protein 1 precursor	Q92896	K.LN#LTTDPK.F
Acid phosphatase, prostate	Q96QM0	K.FLN#ESYK.H
ADAM 10 precursor	O14672	R.IN#TTADEK.D
Prostate-specific membrane antigen-like protein	Q9HBA9	K.IYN#ISMK.H
Afamin precursor	P43652	K.FN#ETTEK.S
47 kDa heat shock protein precursor	P29043	R.SLSN#STAR.N
BA209J19.1.1	Q9H1L6	K.LLN#LTVR.I
Splice isoform 2A of Q02487 Desmocollin 2A/2B precursor	Q02487	R.AN#YTILK.G

FIG. 6 cont'd.

<b>Protein Name</b>	<b>Swiss-Prot</b>	<b>Peptide Sequence</b>
Basement membrane-specific heparan sulfate proteoglycan core protein precursor	P98160	R.ALVN#FTR.S
Alpha-1-acid glycoprotein 1 precursor	P02763	R.EN#GTISR.Y
Splice isoform LAMP-2A of P13473 Lysosome-associated membrane glycoprotein 2 precursor	P13473	R.LN#SSTIK.Y
EMILIN 1 precursor	Q9Y6C2	R.LN#LTAAR.L
alpha-1-antichymotrypsin, precursor	P01011	K.LINDYVKN#GTR.G
Prostate specific antigen precursor	P07288	R.N#KSVILLGR.H

FIG. 6 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
cathepsin L [Homo sapiens]	gi3929737	ySVAnDT GFVDIPk	N-Term(iTRAQ8plex), N5(Deamidated), K14(iTRAQ8plex)	3	712.38525	2135.14121	-3.23	94.18	0.824	0.956	0.377	0.443	0.784	0.618	1.332
Chain A, Crystal Structure Of The C2 Fragment Of Streptococcal Protein G In Complex With The Fc Domain Of Human Igg	gi1065199	eqQYnSTY R	N-Term(iTRAQ8plex), Q2(Deamidated), N5(Deamidated)	3	498.90378	1494.69678	-3.09	47.82	1.360	1.209	0.758	0.814	1.413	1.579	1.141
PREDICTED: CD63 molecule isoform 1 [Macaca mulatta]	gi10909713_1	mHTASIL DR	N-Term(iTRAQ8plex), N1(Deamidated), N2(Deamidated)	2	723.87830	1446.74932	0.19	61.29	0.981	0.931	1.461	2.661	3.034	0.650	1.282
lysosome-associated membrane protein 2	gi1094569	lnSSTIk	N-Term(iTRAQ8plex), N2(Deamidated), K7(iTRAQ8plex)	2	686.41791	1371.82854	-1.33	62.51	0.806	0.337	2.319	0.800	1.360	0.360	0.477

FIG. 7

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	$\Delta$ M [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
lysosome-associated membrane protein 2	gi1094569	vQPFnVT QGk	N-Term(iTRAQ8plex), N5(Deamidated), K10(iTRAQ8plex)	2	863.99768	1726.98808	-3.90	84.75	1.323	0.624	3.095	1.661	2.668	0.653	1.194
Biotinidase precursor (Biotinase)	gi1168695	fnDTEVL QR	N-Term(iTRAQ8plex), N2(Deamidated)	2	713.87793	1426.74858	0.42	85.39	1.120	0.800	0.748	0.860	0.965	1.266	0.807
acid alpha-glucosidase preproprotein [Homo sapiens]	gi11939389_1	gVFITnET GQPLIGk	N-Term(iTRAQ8plex), N6(Deamidated), K15(iTRAQ8plex)	3	728.42102	2183.24851	-2.17	94.09	0.895	0.625	0.698	0.762	0.846	0.727	0.847
acid alpha-glucosidase preproprotein [Homo sapiens]	gi11939389_1	lEnLSSSE mGYTATL TR	N-Term(iTRAQ8plex), N3(Deamidated), M9(Oxidation)	3	732.03253	2194.08304	-0.95	95.21	0.841	0.477	0.457	0.552	1.492	2.490	1.029
acid alpha-glucosidase preproprotein [Homo sapiens]	gi11939389_1	nNTIVNE LVR	N-Term(iTRAQ8plex), N1(Deamidated)	2	738.91858	1476.82988	-1.70	90.69	0.474	0.239	0.255	0.116	0.311	0.311	0.314

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
Chain A, Structure Of Human Neutral Endopeptidase Complexed With Phosphoramidon	gi12084341	scInESAIDSR	N-Term(iTRAQ8plex), C2(Carbamidomethyl), N4(Deamidated)	2	778.87817	1556.74907	-2.41	63.31	0.580	0.169	0.690	3.057	2.068	0.480	0.273
Laminin subunit gamma-1 precursor (Laminin B2 chain)	gi126369	iPAImQITTEANEK	N-Term(iTRAQ8plex), N5(Deamidated), K14(iTRAQ8plex)	3	717.73785	2151.19901	-5.93	94.14	0.671	0.520	0.363	0.569	0.599	0.588	0.658
Laminin subunit gamma-1 precursor (Laminin B2 chain)	gi126369	ILNnLTSIk	N-Term(iTRAQ8plex), N4(Deamidated), K9(iTRAQ8plex)	3	542.33923	1625.00315	-3.85	97.17	0.569	0.424	0.191	0.664	0.461	0.512	0.461
Laminin subunit gamma-1 precursor (Laminin B2 chain)	gi126369	tAdYTSIEAYNLLLR	N-Term(iTRAQ8plex), N3(Deamidated)	2	994.01373	1987.02019	-4.23	95.33	1.021	1.092	0.440	0.600	0.458	3.968	0.463
Laminin subunit gamma-1 precursor (Laminin B2 chain)	gi126369	vnNTLSSQISR	N-Term(iTRAQ8plex), N2(Deamidated)	2	762.41846	1523.82964	-2.28	77.07	1.016	0.933	0.687	0.522	0.844	2.666	0.681

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
Acid phosphatase 2, lysosomal [Homo sapiens]	gi13111975	yEQLQmE TR	N-Term(iTRAQ8plex), N6(Deamidated)	2	743.37640	1485.74553	-2.13	69.85	0.846	0.624	1.416	0.562	1.642	1.289	1.996
Tetraspanin 8 [Homo sapiens]	gi13528897	iVnETLYE NTK	N-Term(iTRAQ8plex), N3(Deamidated), K11(iTRAQ8plex)	3	645.02661	1933.06528	-4.43	89.95	0.630	0.531	0.518	0.583	0.650	0.474	0.373
HPX protein [Homo sapiens]	gi13529281	sWPAVGn cSSALR	N-Term(iTRAQ8plex), N7(Deamidated), C8(Carbamidomethyl)	2	855.43036	1709.85344	-2.84	86.88	1.132	0.827	1.042	1.114	2.231	5.946	1.299
type VI collagen alpha 2 chain precursor [Homo sapiens]	gi13603394	nmTLFSD LVAEk	N-Term(iTRAQ8plex), N1(Deamidated), M2(Oxidation), K12(iTRAQ8plex)	3	665.02771	1993.06858	-4.35	96.99	0.651	0.423		0.739	0.554	1.983	0.361

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
cysteine-rich fibroblast growth factor receptor	gi1373019	InLTTDPK	N-Term(iTRAQ8plex), N2(Deamidated), K8(iTRAQ8plex)	2	755.94385	1510.88042	-8.79	82.68	1.173	0.971	1.224	1.015	1.088	1.168	1.273
EMILIN1 protein [Homo sapiens]	gi14043093	eInTTSQmQAALLLEk	N-Term(iTRAQ8plex), N3(Deamidated), M8(Oxidation), K15(iTRAQ8plex)	3	764.07062	2290.19730	-3.67	82.36	1.276	1.444	0.765		0.776	0.334	1.128
HSP90B1 protein [Homo sapiens]	gi14327942	eEEAIQLDGLnASQIR	N-Term(iTRAQ8plex), N11(Deamidated)	3	697.69934	2091.08347	-1.78	93.40	0.817	0.551	0.418	0.760	0.611	0.645	0.730
alpha-N-acetylglucosaminidase	gi1479983	vFPQVnV Tk	N-Term(iTRAQ8plex), N6(Deamidated), K9(iTRAQ8plex)	3	547.66205	1640.97159	-7.07	89.01	0.412	0.352	0.275	0.424	0.401	0.358	0.421
ceruloplasmin [Homo sapiens]	gi1620909	eHEGAIYPDnTTDFQR	N-Term(iTRAQ8plex), N10(Deamidated)	3	733.34686	2198.02604	-1.98	81.13	1.105	0.655	0.518	1.160	1.378	0.664	1.314

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
ceruloplasmin [Homo sapiens]	gi1620909	eNLTA PGS DSAVFFE QGTR	N-Term(iTRAQ8plex), N2(Deamidated)	3	811.40076	2432.18772	-0.27	94.45	0.692	0.469	0.421	0.868	0.430	0.414	0.631
ASAHI protein [Homo sapiens]	gi16877108	tVLEnSIS YEEAk	N-Term(iTRAQ8plex), N5(Deamidated), K13(iTRAQ8plex)	2	#####	2080.08403	-3.17	84.46	0.863	0.964	1.476	0.633	2.183	2.681	1.612
Cl-inhibitor	gi179617	dTFVnAS R	N-Term(iTRAQ8plex), N5(Deamidated)	2	607.81750	1214.62773	-3.42	53.93	0.679	0.994	0.687	1.434	0.646	0.363	0.642
complement H factor	gi180498	iSEEnETT cYmGk	N-Term(iTRAQ8plex), N5(Deamidated), C9(Carbamidomethyl), M11(Oxidation), K13(iTRAQ8plex)	3	729.68347	2187.03586	-2.16	78.48	1.005	0.285	0.808	0.709	0.966	0.532	0.503
fibronectin receptor alpha-subunit precursor [Homo sapiens]	gi182710	nLnNSQS DVV/SFR	N-Term(iTRAQ8plex), N3(Deamidated)	2	892.95593	1784.90459	-1.95	83.95	3.362	1.367	2.164	2.433	3.213	15.073	1.254

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
collagen VI- alpha-1 chain [Homo sapiens]	gi1915900	nFTAADW GQSR	N- Term(iTRAQ8pl ex), N1(Deamidated)	2	779.38062	1557.75395	-3.85	86.43	1.170	1.370	1.058	0.246	1.125	0.494	0.511
collagen VI- alpha-1 chain [Homo sapiens]	gi1915900	mFTAAD WGQSR	N- Term(iTRAQ8pl ex), N2(Deamidated)	3	571.95801	1713.85947	-0.94	81.15	0.783	0.466	0.478	0.489	0.686	0.498	0.576
integrin beta 1 isoform 1B precursor [Homo sapiens]	gi19743815	nPcTSEQn cTSPFSYk	N- Term(iTRAQ8pl ex), C3(Carbamidom ethyl), N8(Deamidated) , C9(Carbamidom ethyl), K16(iTRAQ8ple x)	3	843.72949	2529.17392	-4.23	84.21	1.153	1.170	2.067	1.980	1.189	7.544	1.202
Chain A, Fc Fragment Of Rituximab Bound To A Minimized Version Of The B- Domain From Protein A Called Z34c	gi20664303	eEQYnST YR	N- Term(iTRAQ8pl ex), N5(Deamidated)	3	498.90378	1494.69678	-3.09	47.82	1.360	1.209	0.758	0.814	1.413	1.579	1.141

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	$\Delta$ M [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
attractin isoform 2 [Homo sapiens]	gi21750863	iDSTGnVT NELR	N-Term(iTRAQ8plex), N6(Deamidated)	2	812.42651	1623.84575	-2.10	84.49	1.183	1.039	1.055	0.864	1.400	1.832	1.171
unnamed protein product [Homo sapiens]	gi21751668	mIEhGSLs FLPTLR	N-Term(iTRAQ8plex), M1(Oxidation), N4(Deamidated)	2	950.01038	1899.01348	-3.42	97.49	1.222	0.775	0.243	0.279	0.595	2.318	1.740
unnamed protein product [Homo sapiens]	gi21753029	anTSALA VSPVSN SASAR	N-Term(iTRAQ8plex), N2(Deamidated)	3	702.37561	2105.11228	-0.85	84.93	1.059	0.582	0.677	0.654	0.878	0.597	0.942
unnamed protein product [Homo sapiens]	gi21755029	rAnTSALA VSPVSN SASAR	N-Term(iTRAQ8plex), N3(Deamidated)	3	754.40979	2261.21482	-0.17	80.01	1.024	0.807	0.350	0.493	0.907	0.888	0.778
unnamed protein product [Homo sapiens]	gi21754713	vnETEmDI Ak	N-Term(iTRAQ8plex), N2(Deamidated), M6(Oxidation), K10(iTRAQ8plex)	3	592.31494	1774.93027	-2.85	76.22	1.131	0.533	0.642	0.863	0.739	0.797	0.721

FIG. 8

carboxypeptidase D [Homo sapiens]	gi21903712	fANIEYPnI TR	N- Term(iTRAQ8pl ex), N7(Deamidated)	2	765.39386	1529.78044	-6.36	87.02	1.023	0.760	0.960	0.942	1.421	1.059	1.403
aminopeptidase N	gi228379	lnYTL SQG HR	N- Term(iTRAQ8pl ex), N2(Deamidated)	3	498.60516	1493.80094	-0.31	77.54	1.197	0.573	0.768	1.217	1.030	0.660	0.647
microfibrillar-associated protein 4 [Homo sapiens]	gi23111005	fnGSV SFF R	N- Term(iTRAQ8pl ex), N2(Deamidated)	3	455.90692	1365.70621	-3.12	96.00	1.560	1.290	0.140	0.340	0.381	0.870	1.698
microfibrillar-associated protein 4 [Homo sapiens]	gi23111005	vDLEDFE nNTAYAk	N- Term(iTRAQ8pl ex), N8(Deamidated) , K14(iTRAQ8ple x)	2	#####	2238.13164	-3.14	94.09	0.948	1.502	0.314	0.198	0.410	0.525	1.297
immunoglobulin heavy chain, constant region [Homo sapiens]	gi2414494	eEQFnSTY R	N- Term(iTRAQ8pl ex), N5(Deamidated)	2	739.85406	1478.70085	-3.82	61.70	2.046	1.742	1.079	1.186	1.929	0.666	8.762

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
alpha2-HS glycoprotein [Homo sapiens]	gi2521981	vcQDePLL APLnDTR	N-Term(iTRAQ8plex), C2(Carbamidomethyl), C5(Carbamidomethyl), N12(Deamidated)	2	#####	2077.02764	-4.04	96.14	1.751	1.157	1.248	0.861	3.085	10.023	1.656
Chain A, Human Beta-Glucuronidase At 2.6 A Resolution	gi2554776	vVAnGTG TQGQLk	N-Term(iTRAQ8plex), N4(Deamidated), K13(iTRAQ8plex)	3	628.03027	1882.07627	-4.87	67.90	0.785	0.601	0.997	0.311	0.729	0.447	1.271
Chain U, Complex Of Active Site Inhibited Human Blood Coagulation Factor Viiia With Human Recombinant Soluble Tissue Factor	gi2554907	vnVTVED ER	N-Term(iTRAQ8plex), N2(Deamidated)	2	683.36029	1365.71330	-2.24	64.96	0.663	0.534	0.481	9.714	0.663	0.509	0.647
immunoglobulin gamma 2 heavy chain constant region [Homo sapiens]	gi25987831	eEQFnSTF R	N-Term(iTRAQ8plex), N5(Deamidated)	2	731.85754	1462.70781	-2.59	81.09	2.254	1.475	0.368	0.727	0.823	0.416	3.565

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	$\Delta$ M [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
KIAA0533 protein [Homo sapiens]	gi3043590	dnATLQA TLHAAR	N-Term(iTRAQ8plex), N2(Deamidated)	3	562.97168	1686.90049	-4.27	82.38	0.657	0.416	0.293	0.536	0.473	0.709	0.556
KIAA0533 protein [Homo sapiens]	gi3043590	InTTGVSA GcTADLL VGR	N-Term(iTRAQ8plex), N2(Deamidated), C10(Carbamido methyl)	3	704.04431	2110.11838	3.19	95.02	0.677	0.432	0.343	0.488	0.525	0.536	0.648
lysosomal membrane glycoprotein-2	gi307110	vQPFnVT QGk	N-Term(iTRAQ8plex), N5(Deamidated), K10(iTRAQ8plex)	2	863.99762	1726.98796	-3.97	84.09	1.002	0.916	2.291	0.616	1.637	1.825	1.470
lysosomal membrane glycoprotein-2	gi307110	wQmnFTV R	N-Term(iTRAQ8plex), M3(Oxidation), N4(Deamidated)	2	701.85553	1402.70378	-3.79	88.66	0.697	0.434	0.622	0.474	1.051	1.425	1.367

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
integrin, alpha 1 precursor [Homo sapiens]	gi31657142	nITFNVE STk	N-Term(iTRAQ8plex), N1(Deamidated), K10(iTRAQ8plex)	3	583.98602	1749.94352	-2.52	76.28	1.046	1.569	0.506	0.820	0.492	0.678	0.492
integrin, alpha 1 precursor [Homo sapiens]	gi31657142	sEnASLVL SSSNQk	N-Term(iTRAQ8plex), N3(Deamidated), K14(iTRAQ8plex)	3	691.71124	2073.11917	-4.46	83.53	1.406	0.929	0.699	1.137	1.287	2.055	0.832
integrin, alpha 1 precursor [Homo sapiens]	gi31657142	vYVYALn QTR	N-Term(iTRAQ8plex), N7(Deamidated)	2	766.42200	1531.83672	-3.58	91.43	1.125	1.033	0.817	0.889	0.985	1.860	1.256
hypothetical protein [Homo sapiens]	gi31873561	tmFPnLTD VR	N-Term(iTRAQ8plex), M2(Oxidation), N5(Deamidated)	2	757.89307	1514.77886	-2.52	89.88	1.021	0.562	0.482	0.582	0.787	1.264	1.159
sulfated glycoprotein-2	gi338057	eDALnETR	N-Term(iTRAQ8plex), N5(Deamidated)	2	626.81970	1252.63213	-0.13	52.81	1.080	1.046	0.794	0.927	0.770	0.210	2.112

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
sulfated glycoprotein-2	gi338057	IAnLTQGE DQYYLR	N-Term(iTRAQ8plex), N3(Deamidated)	2	995.01166	1989.01604	-3.55	96.65	1.288	1.308	0.433	1.265	0.639	0.349	1.944
transferrin	gi339469	gLVPLA ENYnk	N-Term(iTRAQ8plex), N11(Deamidated), K12(iTRAQ8plex)	2	963.55560	1926.10393	-6.09	93.21	1.543	0.901	1.791	0.961	1.590	2.292	1.311
inter-alpha-trypsin inhibitor heavy chain ITIH1 [Homo sapiens]	gi33989	aNLSSQAL R	N-Term(iTRAQ8plex), N2(Deamidated)	2	632.85986	1264.71245	-3.03	58.64	0.701	1.299	0.511	0.227	0.494	0.167	0.930
HP protein [Homo sapiens]	gi34785974	nLFLnHSE nATAk	N-Term(iTRAQ8plex), N5(Deamidated), N9(Deamidated), K13(iTRAQ8plex)	2	#####	2069.10307	-4.49	81.01	4.767	0.999	0.337	6.168	1.962	2.450	10.207

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
HP protein [Homo sapiens]	gi34785974	vVLLHPnY SQVDIGLI k	N-Term(iTRAQ8plex), N6(Deamidated), K16(iTRAQ8plex)	3	802.13641	2404.39469	-4.72	95.49	0.936	0.290		0.685	0.381	0.264	1.232
Chain A, Crystal Structure Of A Human Igm Rheumatoid Factor Fab In Complex With Its Autoantigen Igg Fc	gi3659940	eqQFnSTY R	N-Term(iTRAQ8plex), Q2(Deamidated), N5(Deamidated)	2	739.85406	1478.70085	-3.81	61.70	2.046	1.742	1.079	1.186	1.929	0.666	8.762
Chain A, Structural Origins Of L(+)-Tartarate Inhibition Of Human Prostatic Acid Phosphatase	gi3660342	flnESYk	N-Term(iTRAQ8plex), N3(Deamidated), K7(iTRAQ8plex)	2	755.41895	1509.83061	-6.82	87.63	5.341	4.310	9.322	5.055	25.338	14.592	7.422

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
Chain A, Structural Origins Of L(+)-Tartrate Inhibition Of Human Prostatic Acid Phosphatase	gi3660342	kFLnESYk	N-Term(iTRAQ8plex), K1(iTRAQ8plex), N4(Deamidated), K8(iTRAQ8plex)	4	486.28677	1942.12526	-8.23	82.97	0.550	0.418	0.576	0.453	1.088	0.909	0.677
beta-2-glycoprotein I [Homo sapiens]	gi3688372	vYkFSA Gn NSLYR	N-Term(iTRAQ8plex), K3(iTRAQ8plex), N8(Deamidated)	4	520.29242	2078.14785	-0.59	80.54	0.724	0.711	0.543	0.622	0.916	0.501	0.863
CD47 antigen isoform 2 precursor [Homo sapiens]	gi38683836	dIYTFDG ALnk	N-Term(iTRAQ8plex), N10(Deamidated), K11(iTRAQ8plex)	2	933.50287	1865.99846	-6.47	93.48	1.467	1.214	1.553	0.987	2.002	2.630	1.994
Immunoglobulin J chain	gi400044	enISDPTS PLR	N-Term(iTRAQ8plex), N2(Deamidated)	2	767.40289	1533.79851	-5.04	80.69	1.002	2.457	0.451	2.781	1.629	0.401	0.753

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
desmoglein 2 [Homo sapiens]	gi416178	inATDAD EPNTLNS k	N-Term(iTRAQ8plex), N2(Deamidated), K15(iTRAQ8plex)	3	738.05304	2212.14457	-4.89	81.10	1.311	1.657	2.464	1.224	1.582	1.389	0.811
desmoglein 2 [Homo sapiens]	gi416178	yVQnGTY TVk	N-Term(iTRAQ8plex), N4(Deamidated), K10(iTRAQ8plex)	3	594.66321	1781.97507	-8.02	79.76	0.939	0.521	0.450	0.449	1.508	0.399	0.441
amine oxidase, copper containing 3 precursor [Homo sapiens]	gi4502119	iQmLSFA GEPLPQn SSmAR	N-Term(iTRAQ8plex), M3(Oxidation), N14(Deamidated), M17(Oxidation)	2	#####	2414.19438	-2.25	93.59	1.870	0.424	2.283	1.264	2.892	8.984	1.530
carboxypeptidase E preproprotein [Homo sapiens]	gi4503009	gnETIVNL IHSTR	N-Term(iTRAQ8plex), N2(Deamidated)	3	586.99048	1758.95688	-4.73	91.80	0.850	0.498	0.421	0.535	0.919	0.924	0.932
cathepsin D preproprotein [Homo sapiens]	gi4503143	gSLSYLnV TR	N-Term(iTRAQ8plex), N7(Deamidated)	2	707.89374	1414.78020	-2.94	87.17	0.679	0.849	0.843	0.467	0.979	0.792	1.296

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
fibromodulin [Homo sapiens]	gi453157	IYLDHNn LTR	N-Term(iTRAQ8plex), N7(Deamidated)	3	521.94971	1563.83457	-5.56	83.79	0.475	0.542	0.281	0.249	0.538	0.658	0.480
fibronin 1 [Homo sapiens]	gi46559356	vLPVnVT DYcQLVR	N-Term(iTRAQ8plex), N5(Deamidated), C10(Carbamidomethyl)	2	991.03967	1981.07207	-0.49	97.34	1.480	1.603	0.497	0.370	0.955	1.986	1.707
Chain B, Human Zinc-Alpha-2-Glycoprotein	gi4699583	dIVEYYn DSnGSHV LQGR	N-Term(iTRAQ8plex), N7(Deamidated), N10(Deamidated)	3	791.37982	2372.12491	-2.49	91.75	1.518	0.696	1.309	1.131	2.278	2.654	2.008
Chain B, Human Zinc-Alpha-2-Glycoprotein	gi4699583	fGcEIENn R	N-Term(iTRAQ8plex), C3(Carbamidomethyl), N8(Deamidated)	2	722.34436	1443.68144	-1.78	75.82	2.501	1.115	7.889	1.718	8.566	15.306	6.280

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
HYOU1 protein [Homo sapiens]	gi47938913	vFGSQnLT TVK	N-Term(iTRAQ8plex), N6(Deamidated), K11(iTRAQ8plex)	2	902.02911	1803.05095	2.05	88.05	0.701	0.863	1.989	3.174	1.510	0.956	1.214
GOLPH2 [Homo sapiens]	gi48146519	aVLVnnIT TGER	N-Term(iTRAQ8plex), N5(Deamidated), N6(Deamidated)	2	796.93860	1592.86992	-6.17	83.92	0.804	0.971	2.525	1.236	1.900	1.321	1.950
galectin 3 binding protein [Homo sapiens]	gi5031863	aAIPsALD TnSSk	N-Term(iTRAQ8plex), N10(Deamidated), K13(iTRAQ8plex)	3	628.68817	1884.04996	-1.86	83.88	1.055	1.094	1.557	3.905	1.464	0.636	1.177
galectin 3 binding protein [Homo sapiens]	gi5031863	aLGFEnAT QALGR	N-Term(iTRAQ8plex), N6(Deamidated)	2	826.94916	1652.89104	0.03	94.18	0.604	0.642	2.405	4.645	1.847	0.520	0.575
myelin protein zero-like 2 precursor [Homo sapiens]	gi5032247	vLEAVnG TDAR	N-Term(iTRAQ8plex), N6(Deamidated)	2	725.39380	1449.78032	-3.31	86.48	0.717	0.747	0.958	0.743	1.058	0.995	0.950

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
hypothetical protein [Homo sapiens]	gi51476184	anYTTLk	N-Term(iTRAQ8plex), N2(Deamidated), K7(iTRAQ8plex)	2	716.43402	1431.86077	-4.17	83.78	1.165	0.642	1.361	1.002	1.488	0.675	0.877
hypothetical protein [Homo sapiens]	gi51476184	nGIYnITV L/ASDQGG R	N-Term(iTRAQ8plex), N1(Deamidated), N5(Deamidated)	2	992.51953	1984.03179	1.44	96.81	1.543	2.182	1.271	0.834	3.191	2.185	2.101
hypothetical protein [Homo sapiens]	gi52545569	anDTSYIR	N-Term(iTRAQ8plex), N2(Deamidated)	2	622.82288	1244.63848	-3.18	52.21	1.196	2.917	1.601	0.629	1.176	0.531	1.234
hypoxia up-regulated 1 precursor [Homo sapiens]	gi5453832	aEPPLnAS ASDQGEk	N-Term(iTRAQ8plex), N6(Deamidated), K15(iTRAQ8plex)	3	708.37085	2123.09800	-4.57	63.22	1.237	1.541	2.315	0.793	1.963	1.257	2.284
SM-11044 binding protein (SMBP)(EP70-P-iso) [Homo sapiens]	gi55957667	iVDVnLTS EGk	N-Term(iTRAQ8plex), N5(Deamidated), K11(iTRAQ8plex)	2	892.51147	1784.01567	-5.89	89.54	0.949	1.232	1.951	1.265	2.083	2.083	1.995

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
heparan sulfate proteoglycan 2 [Homo sapiens]	gi55960426	aLVnFTR	N-Term(iTRAQ8plex), N4(Deamidated)	2	563.32977	1125.65227	-4.18	86.34	1.318	1.454	1.051	0.728	1.157	3.401	1.216
OMM protein (Ig gamma3) heavy chain	gi567112	eEQYnSTFR	N-Term(iTRAQ8plex), N5(Deamidated)	2	739.85516	1478.70305	-2.33	59.55	6.379	0.779	0.363	0.590	2.364	1.206	4.700
TIMP metalloproteinase inhibitor 1 [Homo sapiens]	gi57210055	IVGTPEVnQTLLYQR	N-Term(iTRAQ8plex), N8(Deamidated)	3	686.69812	2058.07981	-0.56	91.43	0.856	0.475	0.780	0.812	0.760	0.481	0.559
ceroid-lipofuscinosis, neuronal 5 [Homo sapiens]	gi5729772	nIEtnYTR	N-Term(iTRAQ8plex), N5(Deamidated)	2	658.34198	1315.67668	-2.18	53.89	1.571	0.725	2.216	1.781	2.470	0.797	0.792
fibrinogen-like 2 precursor [Homo sapiens]	gi5730075	IDGSTnFT R	N-Term(iTRAQ8plex), N6(Deamidated)	2	658.34167	1315.67607	-2.66	77.23	1.402	1.097	0.503	0.320	0.721	0.362	0.713

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
serine (or cysteine) proteinase inhibitor, clade A, member 3 precursor variant [Homo sapiens]	gi62087700	tLnQSSDE LQLSmGN AmFvk	N-Term(iTRAQ8plex), N3(Deamidated), M13(Oxidation), M17(Oxidation), K20(iTRAQ8plex)	4	714.36304	2854.43032	-4.22	91.25	0.798	0.268	0.278	0.706	0.571	0.949	0.645
periostin [Homo sapiens]	gi62824474	eVnDTLL VNELk	N-Term(iTRAQ8plex), N3(Deamidated), K12(iTRAQ8plex)	2	998.56995	1996.13262	-4.85	95.34	0.570	1.241	0.263	0.172	0.341	0.519	0.539
unnamed protein product [Homo sapiens]	gi7020027	iTDIEhGS LANIPR	N-Term(iTRAQ8plex), N6(Deamidated)	2	909.49719	1817.98711	-2.19	96.18	0.619	0.542	0.423	0.253	0.409	0.836	0.334
Chain A, Crystal Structure Of Human Neutrophil Gelatinase Associated Lipocalin Monomer	gi7245433	sYnVTSV LFR	N-Term(iTRAQ8plex), N3(Deamidated)	2	745.90845	1490.80962	-4.05	96.53	1.128	0.655	0.397	0.399	1.623	6.372	0.627

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
Chain A, The Human Complement Component C3	gi78101267	tVLTTPAT NHmGnVT FTTPANR	N-Term(iTRAQ8plex), M10(Oxidation), N12(Deamidated)	3	859.45270	2576.34354	-0.37	89.00	2.595	0.418		0.662	1.743	3.932	3.628
Chain A, The Crystal Structure Of Cellular Repressor Of Ela-Stimulated Genes (Creg)	gi83753668	vnETExDI Ak	N-Term(iTRAQ8plex), N2(Deamidated), X6(M), X6(Oxidation), K10(iTRAQ8plex)	3	592.31494	1774.93027	-2.85	76.22	1.131	0.533	0.642	0.863	0.739	0.797	0.721
alpha-albumin	gi857675	dIENFnST Qk	N-Term(iTRAQ8plex), N6(Deamidated), K10(iTRAQ8plex)	3	602.31836	1804.94052	-7.32	81.21	0.888	0.757	0.526	0.707	0.740	0.490	0.571
Ig alpha-2 chain C region (allotype A2m(2)) - human (fragment)	gi87783	qPLTAnITk	N-Term(iTRAQ8plex), N6(Deamidated), K9(iTRAQ8plex)	2	784.47571	1567.94414	-4.72	83.56	1.139	2.259	0.582	2.435	1.693	0.569	0.652

FIG. 7 cont'd.

Protein Name	Protein Accessions	Peptides	Modifications	Charge	m/z [Da]	MH+ [Da]	ΔM [ppm]	RT [min]	114/113	115/113	116/113	117/113	118/113	119/113	121/113
CD47 glycoprotein=ovarian tumour marker OA3 homolog [human, erythrocytes, Peptide Partial, 15 aa, segment 1 of 2]	gi894281	dIYTFDG ALxk	N-Term(iTRAQ8plex), X10(Deamidated), X10(N), K11(iTRAQ8plex)	2	933.50287	1865.99846	-6.47	93.48	1.467	1.214	1.553	0.987	2.002	2.630	1.994
collagen, type XII, alpha 1 short isoform precursor [Homo sapiens]	gi93141049	eAGnITTD GYEILGk	N-Term(iTRAQ8plex), N4(Deamidated), K15(iTRAQ8plex)	3	730.72620	2190.16404	-5.03	94.69	1.443	1.235	0.900	0.746	1.406	2.492	2.974
collagen, type XII, alpha 1 short isoform precursor [Homo sapiens]	gi93141049	mLEAYnL TEk	N-Term(iTRAQ8plex), M1(Oxidation), N6(Deamidated), K10(iTRAQ8plex)	3	612.99811	1836.97977	-4.13	87.67	1.174	0.988	1.280	0.476	1.650	0.849	1.170
collagen, type XII, alpha 1 short isoform precursor [Homo sapiens]	gi93141049	nLQVYnA TSNSLTV k	N-Term(iTRAQ8plex), N6(Deamidated), K15(iTRAQ8plex)	3	754.42090	2261.24814	-5.14	88.80	4.820	2.600	0.260	0.298	0.291	0.462	0.770

FIG. 7 cont'd.




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D

<b>Description</b>	<b>AG/NAG</b>	<b>t-TEST</b>
Cathepsin L	1.85	0.01
Human Prostatic Acid Phosphatase	0.28	0.02
Microfibrillar-associated protein 4	5.16	0.03
Fibrinogen-like protein 2	1.64	0.03
Periodontal ligament-specific periostin	2.43	0.04
Heparan sulfate proteoglycan 2	2.52	0.04
Galectin 3 binding protein	3.15	0.04
Biglycan	2.74	0.04
HYOU1 protein	0.5	0.05
Acid phosphatase 2, lysosomal	1.76	0.07
Ceroid-lipofuscinosis, neuronal 5	0.75	0.08
Collagen VI-alpha-1 chain	1.67	0.1
Chain A, Human Beta-Glucuronidase At 2.6 A Resolution	1.85	0.11
C1-inhibitor	1.52	0.11
Human Neutral Endopeptidase Complexed With Phosphoramidon	1.39	0.12
SM-11044 binding protein (SMBP)(EP70-P-iso)	1.4	0.12
Desmocollin	1.31	0.13
Cathepsin D	1.33	0.13
Fibronectin receptor alpha-subunit	0.63	0.13
Fibrillin 1	1.52	0.13
Mimecan	1.8	0.13
fibromodulin [Homo sapiens]	1.44	0.13
collagen, type XII, alpha 1 short isoform	3.79	0.13
Integrin beta 1 isoform 1B	0.43	0.13
Type VI collagen alpha 2 chain	2.3	0.13
ASAHI protein	1.81	0.14
Laminin alpha-5	1.36	0.14
Golgi phosphoprotein 2	0.82	0.15
Lysosomal alpha-glucosidase	1.33	0.16
Tetraspanin 8	1.44	0.16
clusterin isoform 2 preproprotein	1.52	0.16
Amine oxidase, copper containing 3	0.5	0.18
Versican	1.22	0.18
Aminopeptidase N	0.7	0.19
Human Neutrophil Gelatinase Associated Lipocalin Monomer	0.39	0.19
Human Zinc-Alpha-2-Glycoprotein	1.7	0.2
Lysosome-associated membrane protein 2	1.71	0.2
Attractin	0.85	0.21
Asporin	1.3	0.24
HP protein	1.54	0.27
Desmoglein 2	0.88	0.29
CD63	1.13	0.38

FIG. 9

# Tissue Specimen for Validation




FIG. 10




FIG. 11A

AG/NAG	
LC-MS/MS	ELISA
1.85	0.69

FIG. 11B




FIG. 12

# Expression of Multiple Mets by Subject




FIG. 13




FIG. 14A

AG/NAG	
LC-MS/MS	ELISA
0.39	0.68

FIG. 14B




FIG. 15A

AG/NAG	
LC-MS/MS	ELISA
1.52	1.18

FIG. 15B




FIG. 16A




FIG. 16B




FIG. 17




FIG. 18A

ID	CV	
	NAG	AG
No Adjustment	0.4	0.31
Adjust for Tumor Nuclei	0.34	0.42
Adjust for Tumor Area	0.32	0.5

FIG. 18B




FIG. 19A

	CV	
	NAG	AG
No Adjustment	0.55	0.50
Adjust for Tumor Nuclei	0.42	0.47
Adjust for Tumor Area	0.47	0.50

FIG. 19B




FIG. 20A




FIG. 20B




FIG. 20C




FIG. 20D




FIG. 20E




FIG. 20F

**Castration Sensitive vs. Castration Resistant Lymph Node Metastatic Prostate Tumor**




FIG. 21




FIG. 22

No/Weak staining on Norm/NAG Pca and Strong staining on AG Pca

Sandwich ELISA Under Optimization-ASPN Continued  
Subject PB633




FIG. 23