[19]	INTELLECTUAL PROPERTY PHILIPPINES			
[12]	INVENTION PUBLICATION			
[11]	Publication Number:	12012501006	Document Code:	B1
[22]	Publication Date:	8/9/2014		
[21]	Application Number:	12012501006	Document Code:	A
[22]	Date Filed:	22/5/2012		
[54]	Title:	COMPOSITIONS AND METHODS RELATING TO ANTI-IGF-1 RECEPTOR ANTIBODIES		
[71]	Applicant(s):	AMGEN INC		
[72]	Inventor(s):	CALZONE FRANK J DESHPANDE RAJENDRA V TSAI MEI-MEI		
[30]	Priority Data:	22/12/2004 US20040638961P		
[51]	International Class 8:	C12N 15/00 20060101ALI20180625BHPH; C12P 19/34 20060101ALI20180625BHPH; C12N 15/13 20060101ALI20180625BHPH; C07K 16/28 20060101ALI20180625BHPH; C07H 21/04 20060101AFI20180625BHPH;		
[57]	Abstract:	The present invention provides compositions and methods relating to or derived from anti-IGF-1R antibodies. In particular embodiments, the invention provides fully human, humanized, or chimeric anti-IGF-1R antibodies that bind human IGF-1R, IGF-1R-binding fragments and derivatives of such antibodies, and IGF-1R-binding polypeptides comprising such fragments. Other embodiments provide nucleic acids encoding such antibodies, antibody fragments and derivatives and polypeptides, cells comprising such polynucleotides, methods of making such antibodies, antibody fragments and derivatives and polypeptides, and methods of using such antibodies, antibody fragments and derivatives and polypeptides, including methods of treating or diagnosing subjects having IGF-1R-related disorders or conditions.		

embodiment, the isolated antigen binding protein comprises an amino acid sequence selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR1 sequence of L1-L52 as shown in Figure 4; and b. a heavy chain CDR2 sequence of H1-H52 as shown in Figure 8. In another embodiment, the isolated antigen binding protein comprises a CDR1 sequence of L1-L52 as shown in Figure 4. In another embodiment, the isolated antigen binding protein comprises a sequence selected from the group consisting of: a. a light chain CDR1 sequence selected from the group consisting of: i. RSSQSLLHSNGYNYLD; ii. RASQ(G/S)(I/V)(G/S)X(Y/F)L(A/N); and iii. RSSQS(L/I)XXXXX; b. a light chain CDR2 sequence selected from the group consisting of: i. LGSNRAS; ii. AASTLQS; and iii. EDNXRPS; c. a heavy chain CDR1 sequence selected from the group consisting of: i. SSNWWS; ii. XYYWS; and iii. SYAM(S/H); and d. a heavy chain CDR2 sequence selected from the group consisting of: i. (E/I)(I/V)(Y/N)(H/Y)SGST(N/Y)YNPSLKS; and ii. XIS(G/S)SG(G/S)STYYADSVKG; wherein amino acid residue symbols enclosed in parentheses identify alternative residues for the same position in a sequence, each X is independently any amino acid residue, and each Z is independently a glycine residue, an alanine residue, a valine residue, a leucine residue, an isoleucine residue, a proline residue, a phenylalanine residue, a methionine residue, a tryptophan residue, or a cysteine residue. In another embodiment, the isolated antigen binding protein comprises a heavy chain CDR3 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises a heavy chain CDR3 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises a heavy chain CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises two amino acid sequences selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a

total of six amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises three amino acid sequences selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of six amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises four amino acid sequences selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a

total of six amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises five amino acid sequences selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of six amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises: a. a light chain CDR1 sequence that differs by no more than a total of six amino acid additions, substitutions, and/or deletions from a

CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain comprising: i. a light chain CDR1 sequence shown in Figure 4; ii. a light chain CDR2 sequence shown in Figure 5; and iii. a light chain CDR3 sequence shown in Figure 6; b. a heavy chain variable domain comprising: i. a heavy chain CDR1 sequence shown in Figure 7; ii. a heavy chain CDR2 sequence shown in Figure 8; and iii. a heavy chain CDR3 sequence shown in Figure 9; or c. the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises either: a. light chain CDR1, CDR2, and CDR3 sequences that each is identical to the CDR1, CDR2, and CDR3 sequences, respectively, of the same light chain variable domain sequence selected from the group consisting of L1-L52; b. heavy chain CDR1, CDR2, and CDR3 sequences that each is identical to the CDR1, CDR2, and CDR3 sequences, respectively, of the same heavy chain variable domain sequence selected from the group consisting of H1-H52; or c. the light chain CDR1, CDR2, and CDR3 sequences of (a) and the heavy chain CDR1, CDR2, and CDR3 sequences of (b).

In another aspect, the present invention provides an isolated antigen binding protein comprising either: a. a light chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least

80% identical to a light chain variable domain sequence of L1-L52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 15 contiguous amino acid residues of a light chain variable domain sequence of L1-L52 as shown in Figure 2; iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 80% identical to a polynucleotide sequence encoding a light chain variable domain sequence of L1-L52 as shown in Figure 1; and iv. a sequence of amino acids encoded by a polynucleotide sequence that hybridizes under moderately stringent conditions to the complement of a polynucleotide consisting of a light chain variable domain sequence of L1-L52 as shown in Figure 1; b. a heavy chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 80% identical to a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 15 contiguous amino acid residues of a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 80% identical to a polynucleotide sequence encoding a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; and iv. a sequence of amino acids encoded by a polynucleotide sequence that hybridizes under moderately stringent conditions to the complement of a polynucleotide consisting of a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; or c. the light chain variable domain of (a) and the heavy chain variable domain of (b); wherein said antigen binding protein binds to human IGF-1R. In one embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 85% identical to a light chain variable domain sequence of L1-L52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 25 contiguous amino acid residues of a light chain variable domain sequence of L1-L52 as shown in Figure 2; iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 85% identical to a polynucleotide sequence encoding a light chain variable domain sequence of L1-L52 as shown in Figure 1; and iv. a sequence of amino acids encoded by a polynucleotide sequence that hybridizes under highly stringent conditions to the complement of a

polynucleotide consisting of a light chain variable domain sequence of L1-L52 as shown in Figure 1; b. a heavy chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 85% identical to a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 25 contiguous amino acid residues of a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 85% identical to a polynucleotide sequence encoding a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; and iv. a sequence of amino acids encoded by a polynucleotide sequence that hybridizes under highly stringent conditions to the complement of a polynucleotide consisting of a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; or c) the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 90% identical to a light chain variable domain sequence of L1-L52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 35 contiguous amino acid residues of a light chain variable domain sequence of L1-L52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 90% identical to a polynucleotide sequence encoding a light chain variable domain sequence of L1-L52 as shown in Figure 1; and b. a heavy chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 90% identical to a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 35 contiguous amino acid residues of a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 90% identical to a polynucleotide sequence encoding a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; or c) the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain sequence

selected from the group consisting of: i. a sequence of amino acids at least 95% identical to a light chain variable domain sequence of L1-L52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 50 contiguous amino acid residues of a light chain variable domain sequence of L1-L52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 95% identical to a polynucleotide sequence encoding a light chain variable domain sequence of L1-L52 as shown in Figure 1; and b. a heavy chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 95% identical to a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 50 contiguous amino acid residues of a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 95% identical to a polynucleotide sequence encoding a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; or c) the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 97% identical to a light chain variable domain sequence of L1-L52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 75 contiguous amino acid residues of a light chain variable domain sequence of L1-L52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 97% identical to a polynucleotide sequence encoding a light chain variable domain sequence of L1-L52 as shown in Figure 1; and b. a heavy chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 97% identical to a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 75 contiguous amino acid residues of a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 97% identical to a polynucleotide sequence encoding a heavy chain variable domain sequence of H1-H52 as

shown in Figure 1; or c) the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 99% identical to a light chain variable domain sequence of L1-L52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 90 contiguous amino acid residues of a light chain variable domain sequence of L1-L52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 99% identical to a polynucleotide sequence encoding a light chain variable domain sequence of L1-L52 as shown in Figure 1; and b. a heavy chain variable domain sequence selected from the group consisting of: i. a sequence of amino acids at least 99% identical to a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; ii. a sequence of amino acids comprising at least 90 contiguous amino acid residues of a heavy chain variable domain sequence of H1-H52 as shown in Figure 2; and iii. a sequence of amino acids encoded by a polynucleotide sequence that is at least 99% identical to a polynucleotide sequence encoding a heavy chain variable domain sequence of H1-H52 as shown in Figure 1; or c. the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises either: a. a light chain variable domain sequence selected from the group consisting of L1-L52 as shown in Figure 2; b. a heavy chain variable domain sequence selected from the group consisting of H1-H52 as shown in Figure 3; or c. the light chain variable domain of (a) and the heavy chain variable domain of (b). In another embodiment, the isolated antigen binding protein comprises a combination of a light chain variable domain and a heavy chain variable domain selected from the group of combinations consisting of: L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20, H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47,

L48H48, L49H49, L50H50, L51H51, and L52H52. In another embodiment, the isolated antigen binding protein further comprises: a. the kappa light chain constant sequence of Figure 13, b. the IgG1 heavy chain constant sequence of Figure 13, or c. the kappa light chain constant sequence of Figure 13 and the IgG1 heavy chain constant sequence of Figure 13. In another embodiment, the isolated antigen binding protein, when bound to IGF-1R: a. inhibits IGF-1R; b. activates IGF-1R; c. cross-competes with a reference antibody for binding to IGF-1R; d. binds to the same epitope of IGF-1R as said reference antibody; e. binds to IGF-1R with substantially the same Kd as said reference antibody; or f. binds to IGF-1R with substantially the same off rate as said reference antibody; wherein said reference antibody comprises a combination of light chain and heavy chain variable domain sequences selected from the group of combinations consisting of L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20, H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47, L48H48, L49H49, L50H50, L51H51, and L52H52. In another embodiment, the isolated antigen binding protein, when bound to a human IGF-1R, inhibits binding of IGF-1 and/or IGF-2 to said human IGF-1R. In another embodiment, the isolated antigen binding protein inhibits the growth of a cancer cell by greater than about 80% in the presence of a growth stimulant selected from the group consisting of serum, IGF-1, and IGF-2. In another embodiment, said cancer cell is an MCF-7 human breast cancer cell. In another embodiment, the isolated antigen binding protein binds to human IGF-1R with a selectivity that is at least fifty times greater than its selectivity for human insulin receptor. In another embodiment, the isolated antigen binding protein inhibits tumor growth *in vivo*. In another embodiment, the isolated antigen binding protein inhibits IGF-1R mediated tyrosine phosphorylation. In another embodiment, the isolated antigen binding protein specifically binds to the IGF-1R of a non-human primate, a cynomologous monkey, a chimpanzee, a non-primate mammal, a rodent, a mouse, a rat, a hamster, a

guinea pig, a cat, or a dog. In another embodiment, the isolated antigen binding protein comprises: a. a human antibody; b. a humanized antibody; c. a chimeric antibody; d. a monoclonal antibody; e. a polyclonal antibody; f. a recombinant antibody; g. an antigen-binding antibody fragment; h. a single chain antibody; i. a diabody; j. a triabody; k. a tetrabody; l. a Fab fragment; m. a $F(ab')_2$ fragment; n. a domain antibody; o. an IgD antibody; p. an IgE antibody; q. an IgM antibody; r. an IgG1 antibody; s. an IgG2 antibody; t. an IgG3 antibody; u. an IgG4 antibody; or v. an IgG4 antibody having at least one mutation in a hinge region that alleviates a tendency to form intra-H chain disulfide bond.

In another aspect, the present invention provides an isolated polynucleotide comprising a sequence that encodes the light chain, the heavy chain, or both of said antigen binding protein. In one embodiment, said polynucleotide comprises a light chain variable domain nucleic acid sequence of Figure 1 and/or a heavy chain variable domain nucleic acid sequence of Figure 1. In another embodiment, a plasmid comprises said isolated polynucleotide. In another embodiment, said plasmid is an expression vector. In another embodiment, an isolated cell comprises said polynucleotide. In another embodiment, a chromosome of said cell comprises said polynucleotide. In another embodiment, said cell is a hybridoma. In another embodiment, an expression vector comprises said polynucleotide. In another embodiment, said cell is a CHO cell. In another embodiment, the present invention provides a method of making an antigen binding protein that binds human IGF-1R, comprising incubating said isolated cell under conditions that allow it to express said antigen binding protein.

In another aspect, the present invention provides a pharmaceutical composition comprising the antigen binding protein. In one embodiment, the present invention provides a method of treating a condition in a subject comprising administering to said subject said pharmaceutical composition, wherein said condition is treatable by reducing the activity of IGF-1R in said subject. In another embodiment, said subject is a human being. In another embodiment, said condition is multiple myeloma, a liquid tumor, liver cancer, a thymus disorder, a T-cell mediated autoimmune disease, an

endocronological disorder, ischemia, or a neurodegenerative disorder. In another embodidment, said liquid tumor is selected from the group consisting of acute lymphocytic leukemia (ALL) and chronic myelogenous leukemia (CML); wherein said liver cancer is selected from the group consisting of hepatoma, hepatocellular carcinoma, cholangiocarcinoma, angiosarcomas, hemangiosarcomas, hepatoblastoma; wherein said thymus disorder is selected from the group consisting of thymoma and thyroiditis, wherein said T-cell mediated autoimmune disease is selected from the group consisting of Multiple Sclerosis, Rheumatoid Arthritis, Systemic Lupus Erythematosus (SLE), Grave's Disease, Hashimoto's Thyroiditis, Myasthenia Gravis, Auto-Immune Thyroiditis, Bechet's Disease, wherein said endocrinological disorder is selected from the group consisting of Type II Diabetes, hyperthyroidism, hypothyroidism, thyroiditis, hyperadrenocorticism, and hypoadrenocorticism; wherein said ischemia is post cardiac infarct ischemia, or wherein said neurodegenerative disorder is Alzheimer's Disease. In another embodiment, said condition is selected from the group consisting of acromegaly, bladder cancer, Wilm's tumor, ovarian cancer, pancreatic cancer, benign prostatic hyperplasia, breast cancer, prostate cancer, bone cancer, lung cancer, colorectal cancer, cervical cancer, synovial sarcoma, diarrhea associated with metastatic carcinoid, vasoactive intestinal peptide secreting tumors, gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels, inappropriate microvascular proliferation, glioblastoma, medulloblastoma, head and neck squamous cell cancer, oral cancer, oral leukoplakia, prostate intraepithelial neoplasia, anal cancer, esophageal cancer, gastric cancer, bone cancer, metastatic cancer, polycythemia rubra vera, a benign condition related to oxidative stress, retinopathy of prematurity, Acute Respiratory Distress Syndrome, an overdose of acetaminophen, bronchopulmonary dysplasia, cystic fibrosis, lung fibrosis, and diabetic retinopathy. In another embodiment, the method further comprising administering to said subject a second treatment. In another embodiment, said second treatment is administered to said subject before and/or simultaneously with and/or after said pharmaceutical composition is administered to said subject. In another embodiment, said second treatment comprises radiation treatment, surgery, or

a second pharmaceutical composition. In another embodiment, said second pharmaceutical composition comprises an agent selected from the group consisting of a corticosteroid, an anti-emetic, ondansetron hydrochloride, granisetron hydrochloride, metroclopramide, domperidone, haloperidol, cyclizine, lorazepam, prochlorperazine, dexamethasone, levomepromazine, tropisetron, a cancer vaccine, a GM-CSF inhibiting agent, a GM-CSF DNA vaccine, a cell-based vaccine, a dendritic cell vaccine, a recombinant viral vaccine, a heat shock protein (HSP) vaccine, an allogeneic tumor vaccine, an autologous tumor vaccine, an analgesic, ibuprofen, naproxen, choline magnesium trisalicylate, an oxycodone hydrochloride, an anti-angiogenic agent, an anti-vascular agent, bevacizumab, an anti-VEGF antibody, an anti-VEGF receptor antibody, a soluble VEGF receptor fragment, an anti-TWEAK antibody, an anti-TWEAK receptor antibody, a soluble TWEAK receptor fragment, AMG 706, AMG 386, an anti-proliferative agent, a farnesyl protein transferase inhibitor, an $\alpha v\beta 3$ inhibitor, an $\alpha v\beta 5$ inhibitor, a p53 inhibitor, a Kit receptor inhibitor, a ret receptor inhibitor, a PDGFR inhibitor, a growth hormone secretion inhibitor, an angiopoietin inhibitor, a tumor infiltrating macrophage-inhibiting agent, a c-fms inhibiting agent, an anti-c-fms antibody, an CSF-1 inhibiting agent, an anti-CSF-1 antibody, a soluble c-fms fragment, pegvisomant, gemcitabine, panitumumab, irinothecan, and SN-38. In another embodiment, said method comprises administering to said subject a third treatment. In another embodiment, said condition is a cancer, said second treatment comprises administering panitumumab, and said third treatment comprises administering gemcitabine. In another embodiment, said condition is selected from the group consisting of acromegaly, bladder cancer, Wilm's tumor, ovarian cancer, pancreatic cancer, benign prostatic hyperplasia, breast cancer, prostate cancer, bone cancer, lung cancer, colorectal cancer, cervical cancer, synovial sarcoma, diarrhea associated with metastatic carcinoid, vasoactive intestinal peptide secreting tumors, gigantism, psoriasis, atherosclerosis, smooth muscle restenosis of blood vessels, inappropriate microvascular proliferation, glioblastoma, medulloblastoma, head and neck squamous cell cancer, oral cancer, oral leukoplakia, prostate intraepithelial neoplasia, anal cancer, esophageal cancer, gastric cancer, bone cancer,

metastatic cancer, polycythemia rubra vera, a benign condition related to oxidative stress, retinopathy of prematurity, Acute Respiratory Distress Syndrome, an overdose of acetaminophen, bronchopulmonary dysplasia, cystic fibrosis, lung fibrosis, and diabetic retinopathy.

In another aspect, the present invention provides a method of increasing the longevity of a subject comprising administering to said subject said pharmaceutical composition.

In another aspect, the present invention provides a method of decreasing IGF-1R activity in a subject in need thereof comprising administering to said subject said pharmaceutical composition.

In another aspect, the present invention provides a method of decreasing IGF-1R signaling in a subject in need thereof comprising administering to said subject said pharmaceutical composition.

In another aspect, the present invention provides a method of inhibiting the binding of IGF-1 and/or IGF-2 to IGF-1R in a subject in need thereof comprising administering to said subject said pharmaceutical composition.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides compositions, kits, and methods relating to molecules that bind to the Insulin-Like Growth Factor Receptor ("IGF-1R"), including molecules that agonize or antagonize IGF-1R, such as anti-IGF-1R antibodies, antibody fragments, and antibody derivatives, *e.g.*, antagonistic anti-IGF-1R antibodies, antibody fragments, or antibody derivatives. Also provided are nucleic acids, and derivatives and fragments thereof, comprising a sequence of nucleotides that encodes all or a portion of a polypeptide that binds to IGF-1R, *e.g.*, a nucleic acid encoding all or part of an anti-IGF-1R antibody, antibody fragment, or antibody derivative, plasmids and vectors comprising such nucleic acids, and cells or cell lines comprising such nucleic acids and/or vectors and plasmids. The provided methods include, for example, methods of making, identifying, or isolating molecules that bind to IGF-1R, such as anti-IGF-1R antibodies, methods of determining whether a molecule binds to IGF-1R, methods of determining whether a molecule agonizes or antagonizes IGF-1R, methods of making compositions,

such as pharmaceutical compositions, comprising a molecule that binds to IGF-1R, and methods for administering a molecule that binds IGF-1R to a subject, for example, methods for treating a condition mediated by IGF-1R, and for agonizing or antagonizing a biological activity of IGF-1R, IGF-1, and/or IGF-2 *in vivo* or *in vitro*.

Polynucleotide and polypeptide sequences are indicated using standard one- or three-letter abbreviations. Unless otherwise indicated, polypeptide sequences have their amino termini at the left and their carboxy termini at the right and single-stranded nucleic acid sequences, and the top strand of double-stranded nucleic acid sequences, have their 5' termini at the left and their 3' termini at the right. A particular polypeptide or polynucleotide sequence also can be described by explaining how it differs from a reference sequence.

Polynucleotide and polypeptide sequences of particular light and heavy chain variable domains are shown in Figures 1, 2 and 3,, where they are labeled, for example, L1 ("light chain variable domain 1"), H1 ("heavy chain variable domain 1"), *etc.* Antibodies comprising a light chain and heavy chain from Figures 2 and 3 are indicated by combining the name of the light chain and the name of the heavy chain variable domains. For example, "L4H7," indicates an antibody comprising the light chain variable domain of L4 and the heavy chain variable domain of H7.

Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, *e.g.*, Sambrook *et al.* Molecular Cloning: A Laboratory Manual, 2d ed.,

Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989) and Ausubel *et al.*, Current Protocols in Molecular Biology, Greene Publishing Associates (1992), and Harlow and Lane Antibodies: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990), which are incorporated herein by reference. Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The terminology used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques can be used for chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

The following terms, unless otherwise indicated, shall be understood to have the following meanings:

The term "isolated molecule" (where the molecule is, for example, a polypeptide, a polynucleotide, or an antibody) is a molecule that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) is substantially free of other molecules from the same species (3) is expressed by a cell from a different species, or (4) does not occur in nature. Thus, a molecule that is chemically synthesized, or synthesized in a cellular system different from the cell from which it naturally originates, will be "isolated" from its naturally associated components. A molecule also may be rendered substantially free of naturally associated components by isolation, using purification techniques well known in the art. Molecule purity or homogeneity may be assayed by a number of means well known in the art. For example, the purity of a polypeptide sample may be assayed using polyacrylamide gel electrophoresis and staining of the gel to visualize the polypeptide using techniques well known in the art. For certain purposes, higher resolution may be provided by using HPLC or other means well known in the art for purification.

The terms "IGF-1R inhibitor" and "IGF-1R antagonist" are used interchangeably. Each is a molecule that detectably inhibits at least one

function of IGF-1R. Conversely, an "IGF-1R agonist" is a molecule that detectably increases at least one function of IGF-1R. The inhibition caused by an IGF-1R inhibitor need not be complete so long as it is detectable using an assay. Any assay of a function of IGF-1R can be used, examples of which are provided herein. Examples of functions of IGF-1R that can be inhibited by an IGF-1R inhibitor, or increased by an IGF-1R agonist, include binding to IGF-1, IGF-12, and/or another IGF-1R-activating molecule, kinase activity, downstream signaling, and so on. Examples of types of IGF-1R inhibitors and IGF-1R agonists include, but are not limited to, IGF-1R binding polypeptides such as antigen binding proteins (e.g., IGF-1R inhibiting antiben binding proteins), antibodies, antibody fragments, and antibody derivatives.

The terms "peptide," "polypeptide" and "protein" each refers to a molecule comprising two or more amino acid residues joined to each other by peptide bonds. These terms encompass, *e.g.*, native and artificial proteins, protein fragments and polypeptide analogs (such as muteins, variants, and fusion proteins) of a protein sequence as well as post-translationally, or otherwise covalently or non-covalently, modified proteins. A peptide, polypeptide, or protein may be monomeric or polymeric.

The term "polypeptide fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion as compared to a corresponding full-length protein. Fragments can be, for example, at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 50, 70, 80, 90, 100, 150 or 200 amino acids in length. Fragments can also be, for example, at most 1,000, 750, 500, 250, 200, 175, 150, 125, 100, 90, 80, 70, 60, 50, 40, 30, 20, 15, 14, 13, 12, 11, or 10 amino acids in length. A fragment can further comprise, at either or both of its ends, one or more additional amino acids, for example, a sequence of amino acids from a different naturally-occurring protein (*e.g.*, an Fc or leucine zipper domain) or an artificial amino acid sequence (*e.g.*, an artificial linker sequence).

Polypeptides of the invention include polypeptides that have been modified in any way and for any reason, for example, to: (1) reduce susceptibility to proteolysis, (2) reduce susceptibility to oxidation, (3) alter binding affinity for forming protein complexes, (4) alter binding affinities, and

(4) confer or modify other physicochemical or functional properties. Analogs include muteins of a polypeptide. For example, single or multiple amino acid substitutions (*e.g.*, conservative amino acid substitutions) may be made in the naturally occurring sequence (*e.g.*, in the portion of the polypeptide outside the domain(s) forming intermolecular contacts. A "conservative amino acid substitution" is one that does not substantially change the structural characteristics of the parent sequence (*e.g.*, a replacement amino acid should not tend to break a helix that occurs in the parent sequence, or disrupt other types of secondary structure that characterize the parent sequence or are necessary for its functionality). Examples of art-recognized polypeptide secondary and tertiary structures are described in Proteins, Structures and Molecular Principles (Creighton, Ed., W. H. Freeman and Company, New York (1984)); Introduction to Protein Structure (C. Branden and J. Tooze, eds., Garland Publishing, New York, N.Y. (1991)); and Thornton *et at.* Nature 354:105 (1991), which are each incorporated herein by reference.

The present invention also provides non-peptide analogs of IGF-1R binding polypeptides. Non-peptide analogs are commonly used in the pharmaceutical industry as drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed "peptide mimetics" or "peptidomimetics". Fauchere, J. Adv. Drug Res. 15:29 (1986); Veber and Freidinger TINS p.392 (1985); and Evans *et al.* J. Med. Chem. 30:1229 (1987), which are incorporated herein by reference. Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (*i.e.*, a polypeptide that has a desired biochemical property or pharmacological activity), such as a human antibody, but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: --CH_2NH--, --CH_2S--, --CH_2--CH_2--, --CH=CH-(*cis* and *trans*), --$COCH_2$--, --$CH(OH)CH_2$--, and --CH_2SO--, by methods well known in the art. Systematic substitution of one or more amino acids of a consensus sequence with a D-amino acid of the same type (*e.g.*, D-lysine in place of L-lysine) may also be used to generate more stable peptides. In addition, constrained peptides

comprising a consensus sequence or a substantially identical consensus sequence variation may be generated by methods known in the art (Rizo and Gierasch Ann. Rev. Biochem. 61:387 (1992), incorporated herein by reference), for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

A "variant" of a polypeptide (*e.g.*, an antibody) comprises an amino acid sequence wherein one or more amino acid residues are inserted into, deleted from and/or substituted into the amino acid sequence relative to another polypeptide sequence. Variants of the invention include fusion proteins.

A "derivative" of a polypeptide is a polypeptide (*e.g.*, an antibody) that has been chemically modified, *e.g.*, via conjugation to another chemical moiety such as, for example, polyethylene glycol, albumin (*e.g.*, human serum albumin), phosphorylation, and glycosylation. Unless otherwise indicated, the term "antibody" includes, in addition to antibodies comprising two full-length heavy chains and two full-length light chains, derivatives, variants, fragments, and muteins thereof, examples of which are described below.

An "antigen binding protein" is a protein comprising a portion that binds to an antigen and, optionally, a scaffold or framework portion that allows the antigen binding portion to adopt a conformation that promotes binding of the antigen binding protein to the antigen. Examples of antigen binding proteins include antibodies, antibody fragments (e.g., an antigen binding portion of an antibody), antibody derivatives, and antibody analogs. The antigen binding protein can comprise, for example, an alternative protein scaffold or artificial scaffold with grafted CDRs or CDR derivatives. Such scaffolds include, but are not limited to, antibody-derived scaffolds comprising mutations introduced to, for example, stabilize the three-dimensional structure of the antigen binding protein as well as wholly synthetic scaffolds comprising, for example, a biocompatible polymer. See, for example, Korndorfer et al., 2003, Proteins: Structure, Function, and Bioinformatics, Volume 53, Issue 1:121-129; Roque et al., 2004, Biotechnol. Prog. 20:639-654. In addition, peptide antibody mimetics ("PAMs") can be

used, as well as scaffolds based on antibody mimetics utilizing fibronection components as a scaffold.

An antigen binding protein can have, for example, the structure of a naturally occurring immunoglobulin. An "immunoglobulin" is a tetrameric molecule. In a naturally occurring immunoglobulin, each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Human light chains are classified as kappa and lambda light chains. Heavy chains are classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 more amino acids. See generally, Fundamental Immunology Ch. 7 (Paul, W., ed., 2nd ed. Raven Press, N.Y. (1989)) (incorporated by reference in its entirety for all purposes). The variable regions of each light/heavy chain pair form the antibody binding site such that an intact immunoglobulin has two binding sites.

Naturally occurring immunoglobulin chains exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. From N-terminus to C-terminus, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat *et al.* in *Sequences of Proteins of Immunological Interest*, 5th Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication no. 91-3242, 1991.

An "antibody" refers to an intact immunoglobulin or to an antigen binding portion thereof that competes with the intact antibody for specific binding, unless otherwise specified. Antigen binding portions may be produced by recombinant DNA techniques or by enzymatic or chemical cleavage of intact antibodies. Antigen binding portions include, *inter alia*,

Fab, Fab', $F(ab')_2$, Fv, domain antibodies (dAbs), and complementarity determining region (CDR) fragments, single-chain antibodies (scFv), chimeric antibodies, diabodies, triabodies, tetrabodies, and polypeptides that contain at least a portion of an immunoglobulin that is sufficient to confer specific antigen binding to the polypeptide.

A Fab fragment is a monovalent fragment having the V_L, V_H, C_L and C_H1 domains; a $F(ab')_2$ fragment is a bivalent fragment having two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment has the V_H and C_H1 domains; an Fv fragment has the V_L and V_H domains of a single arm of an antibody; and a dAb fragment has a V_H domain, a V_L domain, or an antigen-binding fragment of a V_H or V_L domain (US Pat. No. 6,846,634, 6,696,245, US App. Pub. No. 05/0202512, 04/0202995, 04/0038291, 04/0009507, 03/0039958, Ward *et al.*, Nature 341:544-546, 1989).

A single-chain antibody (scFv) is an antibody in which a V_L and a V_H region are joined via a linker (*e.g.*, a synthetic sequence of amino acid residues) to form a continuous protein chain wherein the linker is long enough to allow the protein chain to fold back on itself and form a monovalent antigen binding site (see, *e.g.*, Bird *et al.*, 1988, Science 242:423-26 and Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-83). Diabodies are bivalent antibodies comprising two polypeptide chains, wherein each polypeptide chain comprises V_H and V_L domains joined by a linker that is too short to allow for pairing between two domains on the same chain, thus allowing each domain to pair with a complementary domain on another polypeptide chain (see, *e.g.*, Holliger *et al.*, 1993, Proc. Natl. Acad. Sci. USA 90:6444-48, and Poljak *et al.*, 1994, Structure 2:1121-23). If the two polypeptide chains of a diabody are identical, then a diabody resulting from their pairing will have two identical antigen binding sites. Polypeptide chains having different sequences can be used to make a diabody with two different antigen binding sites. Similarly, tribodies and tetrabodies are antibodies comprising three and four polypeptide chains, respectively, and forming three and four antigen binding sites, respectively, which can be the same or different.

Complementarity determining regions (CDRs) and framework regions (FR) of a given antibody may be identified using the system described by

Kabat *et al.* in Sequences of Proteins of Immunological Interest, 5th Ed., US Dept. of Health and Human Services, PHS, NIH, NIH Publication no. 91-3242, 1991. One or more CDRs may be incorporated into a molecule either covalently or noncovalently to make it an antigen binding protein. An antigen binding protein may incorporate the CDR(s) as part of a larger polypeptide chain, may covalently link the CDR(s) to another polypeptide chain, or may incorporate the CDR(s) noncovalently. The CDRs permit the antigen binding protein to specifically bind to a particular antigen of interest.

An antigen binding protein may have one or more binding sites. If there is more than one binding site, the binding sites may be identical to one another or may be different. For example, a naturally occurring human immunoglobulin typically has two identical binding sites, while a "bispecific" or "bifunctional" antibody has two different binding sites.

The term "human antibody" includes all antibodies that have one or more variable and constant regions derived from human immunoglobulin sequences. In one embodiment, all of the variable and constant domains are derived from human immunoglobulin sequences (a fully human antibody). These antibodies may be prepared in a variety of ways, examples of which are described below, including through the immunization with an antigen of interest of a mouse that is genetically modified to express antibodies derived from human heavy and/or light chain-encoding genes.

A humanized antibody has a sequence that differs from the sequence of an antibody derived from a non-human species by one or more amino acid substitutions, deletions, and/or additions, such that the humanized antibody is less likely to induce an immune response, and/or induces a less severe immune response, as compared to the non-human species antibody, when it is administered to a human subject. In one embodiment, certain amino acids in the framework and constant domains of the heavy and/or light chains of the non-human species antibody are mutated to produce the humanized antibody. In another embodiment, the constant domain(s) from a human antibody are fused to the variable domain(s) of a non-human species. In another embodiment, one or more amino acid residues in one or more CDR sequences of a non-human antibody are changed to reduce the likely immunogenicity of

the non-human antibody when it is administered to a human subject, wherein the changed amino acid residues either are not critical for immunospecific binding of the antibody to its antigen, or the changes to the amino acid sequence that are made are conservative changes, such that the binding of the humanized antibody to the antigen is not significantly worse than the binding of the non-human antibody to the antigen. Examples of how to make humanized antibodies may be found in U.S. Pat. Nos. 6,054,297, 5,886,152 and 5,877,293.

The term "chimeric antibody" refers to an antibody that contains one or more regions from one antibody and one or more regions from one or more other antibodies. In one embodiment, one or more of the CDRs are derived from a human anti-IGF-1R antibody. In another embodiment, all of the CDRs are derived from a human anti-IGF-1R antibody. In another embodiment, the CDRs from more than one human anti-IGF-1R antibodies are mixed and matched in a chimeric antibody. For instance, a chimeric antibody may comprise a CDR1 from the light chain of a first human anti-IGF-1R antibody, a CDR2 and a CDR3 from the light chain of a second human anti-IGF-1R antibody, and the CDRs from the heavy chain from a third anti-IGF-1R antibody. Further, the framework regions may be derived from one of the same anti-IGF-1R antibodies, from one or more different antibodies, such as a human antibody, or from a humanized antibody. In one example of a chimeric antibody, a portion of the heavy and/or light chain is identical with, homologous to, or derived from an antibody from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is/are identical with, homologous to, or derived from an antibody (-ies) from another species or belonging to another antibody class or subclass. Also included are fragments of such antibodies that exhibit the desired biological activity (*i.e.*, the ability to specifically bind IGF-1R). See, *e.g.*, U.S. Patent No. 4,816,567 and Morrison, 1985, Science 229:1202-07.

A "neutralizing antibody" or "an inhibitory antibody" is an antibody that inhibits the binding of IGF-1R to IGF-I and/or IGF-2 when an excess of the anti-IGF-1R antibody reduces the amount of IGF-I and/or IGF-2 bound to IGF-1R by at least about 20% using the assay described in Example 9. In

various embodiments, the antibody reduces the amount of IGF-I and/or IGF-2 bound to IGF-1R by at least 30%, 40%, 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 99%, and 99.9%.

An "activating antibody" is an antibody that activates IGF-1R by at least about 20% when added to a cell, tissue or organism expressing IGF-1R, where "100% activation" is the level of activation achieved under physiological conditions by the same molar amount of IGF-1 and/or IGF-2. In various embodiments, the antibody activates IGF-1R activity by at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 125%, 150%, 175%, 200%, 250%, 300%, 350%, 400%, 450%, 500%, 750%, or 1000%.

Fragments or analogs of antibodies can be readily prepared by those of ordinary skill in the art following the teachings of this specification and using techniques well-known in the art. Preferred amino- and carboxy-termini of fragments or analogs occur near boundaries of functional domains. Structural and functional domains can be identified by comparison of the nucleotide and/or amino acid sequence data to public or proprietary sequence databases. Computerized comparison methods can be used to identify sequence motifs or predicted protein conformation domains that occur in other proteins of known structure and/or function. Methods to identify protein sequences that fold into a known three-dimensional structure are known. See, *e.g.*, Bowie *et al.*, 1991, Science 253:164.

A "CDR grafted antibody" is an antibody comprising one or more CDRs derived from an antibody of a particular species or isotype and the framework of another antibody of the same or different species or isotype.

A "multi-specific antibody" is an antibody that recognizes more than one epitope on one or more antigens. A subclass of this type of antibody is a "bi-specific antibody" which recognizes two distinct epitopes on the same or different antigens.

An antigen binding protein "specifically binds" to an antigen (*e.g.*, human IGF-1R) if it binds to the antigen with a dissociation constant of 1 nanomolar or less.

An "antigen binding domain," "antigen binding region," or "antigen binding site" is a portion of an antigen binding protein that contains amino

acid residues (or other moieties) that interact with an antigen and contribute to the antigen binding protein's specificity and affinity for the antigen. For an antibody that specifically binds to its antigen, this will include at least part of at least one of its CDR domains.

An "epitope" is the portion of a molecule that is bound by an antigen binding protein (e.g., by an antibody). An epitope can comprise non-contiguous portions of the molecule (*e.g.*, in a polypeptide, amino acid residues that are not contiguous in the polypeptide's primary sequence but that, in the context of the polypeptide's tertiary and quaternary structure, are near enough to each other to be bound by an antigen binding protein).

The "percent identity" of two polynucleotide or two polypeptide sequences is determined by comparing the sequences using the GAP computer program (a part of the GCG Wisconsin Package, version 10.3 (Accelrys, San Diego, CA)) using its default parameters.

The terms "polynucleotide," "oligonucleotide" and "nucleic acid" are used interchangeably throughout and include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs (*e.g.*, peptide nucleic acids and non-naturally occurring nucleotide analogs), and hybrids thereof. The nucleic acid molecule can be single-stranded or double-stranded. In one embodiment, the nucleic acid molecules of the invention comprise a contiguous open reading frame encoding an antibody, or a fragment, derivative, mutein, or variant thereof, of the invention.

Two single-stranded polynucleotides are "the complement" of each other if their sequences can be aligned in an anti-parallel orientiation such that every nucleotide in one polynucleotide is opposite its complementary nucleotide in the other polynucleotide, without the introduction of gaps, and without unpaired nucleotides at the 5' or the 3' end of either sequence. A polynucleotide is "complementary" to another polynucleotide if the two polynucleotides can hybridize to one another under moderately stringent conditions. Thus, a polynucleotide can be complementary to another polynucleotide without being its complement.

A "vector" is a nucleic acid that can be used to introduce another nucleic acid linked to it into a cell. One type of vector is a "plasmid," which refers to a linear or circular double stranded DNA molecule into which additional nucleic acid segments can be ligated. Another type of vector is a viral vector (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), wherein additional DNA segments can be introduced into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors comprising a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. An "expression vector" is a type of vector that can direct the expression of a chosen polynucleotide.

A nucleotide sequence is "operably linked" to a regulatory sequence if the regulatory sequence affects the expression (*e.g.*, the level, timing, or location of expression) of the nucleotide sequence. A "regulatory sequence" is a nucleic acid that affects the expression (*e.g.*, the level, timing, or location of expression) of a nucleic acid to which it is operably linked. The regulatory sequence can, for example, exert its effects directly on the regulated nucleic acid, or through the action of one or more other molecules (*e.g.*, polypeptides that bind to the regulatory sequence and/or the nucleic acid). Examples of regulatory sequences include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Further examples of regulatory sequences are described in, for example, Goeddel, 1990, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA and Baron *et al.*, 1995, Nucleic Acids Res. 23:3605–06.

A "host cell" is a cell that can be used to express a nucleic acid, *e.g.*, a nucleic acid of the invention. A host cell can be a prokaryote, for example, *E. coli*, or it can be a eukaryote, for example, a single-celled eukaryote (*e.g.*, a yeast or other fungus), a plant cell (*e.g.*, a tobacco or tomato plant cell), an animal cell (*e.g.*, a human cell, a monkey cell, a hamster cell, a rat cell, a mouse cell, or an insect cell) or a hybridoma. Examples of host cells include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (see Gluzman *et*

al., 1981, Cell 23:175), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells or their derivatives such as Veggie CHO and related cell lines which grow in serum-free media (see Rasmussen *et al.*, 1998, Cytotechnology 28:31) or CHO strain DX-B11, which is deficient in DHFR (see Urlaub *et al.*, 1980, Proc. Natl. Acad. Sci. USA 77:4216-20), HeLa cells, BHK (ATCC CRL 10) cell lines, the CV1/EBNA cell line derived from the African green monkey kidney cell line CV1 (ATCC CCL 70) (see McMahan *et al.*, 1991, EMBO J. 10:2821), human embryonic kidney cells such as 293, 293 EBNA or MSR 293, human epidermal A431 cells, human Colo205 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HL-60, U937, HaK or Jurkat cells. Typically, a host cell is a cultured cell that can be transformed or transfected with a polypeptide-encoding nucleic acid, which can then be expressed in the host cell. The phrase "recombinant host cell" can be used to denote a host cell that has been transformed or transfected with a nucleic acid to be expressed. A host cell also can be a cell that comprises the nucleic acid but does not express it at a desired level unless a regulatory sequence is introduced into the host cell such that it becomes operably linked with the nucleic acid. It is understood that the term host cell refers not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to, *e.g.*, mutation or environmental influence, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

IGF-1R

IGF-1R is a transmembrane receptor tyrosine kinase (Blume-Jensen *et al.*, 2001, Nature 411:355-65). The human IGF-1R is synthesized as a 1367 amino acid precursor polypeptide that includes a 30 amino acid signal peptide removed during translocation into the endoplasmic reticulum (Swiss-Prot: P08069). The IGF-1R proreceptor is glycosylated and cleaved by a protease at positions 708-711 (counting from the first amino acid following the signal peptide sequence) during maturation in the ER-golgi resulting in the formation

of an α-chain (1-707) and a β-chain (712-1337) that remain linked by disulfide bonds (Bhaumick *et al.*, 1981, Proc Natl Acad Sci USA 78:4279-83, Chernausek *et al.*, 1981, Biochemistry 20:7345-50, Jacobs *et al.*, 1983, Proc Natl Acad Sci USA 80:1228-31, LeBon *et al.*, 1986, J Biol Chem 261:7685-89, Elleman, *et al.*, 2000, Biochem J 347:771-79). The predominant form of the IGF-1R (and INSR) that exists on the cell-surface is a proteolytically processed and glycosylated $(\alpha\beta)_2$ dimer joined covalently by one or more disulfide bonds.

The extracellular portion of the IGF-1R consists of the α-chain and 191 amino acids of the β-chain (712-905). The receptor contains a single transmembrane spanning sequence (906-929) and a 408-residue cytoplasmic domain that includes a functional tyrosine kinase (Rubin *et al.*, 1983, Nature 305:438-440). Comparative sequence analysis has revealed that the IGF-1R is composed of 11 distinct structural motifs (reviewed by Adams *et al.*, 2000, Cell Mol Life Sci 57:1050-93, Marino-Buslje *et al.*, 1998, FEBS Ltrs 441:331-36, Ward *et al.*, 2001, BMC Bioinformatics 2:4). The N-terminal half of the extracellular domain contains two homologous domains referred to as L1 (1-151) and L2 (299-461) (Ward *et al.*, 2001, *supra*) separated by a cysteine-rich (CR) region (152-298) consisting of several structural modules with disulfide linkages that align with repeating units present in the TNF receptor and laminin (Ward *et al.*, 1995, Proteins 22:141-53). The crystal structure of the L1—CR-L2 domain has been solved (Garrett *et al.*, 1998, Nature 394:395-99). The L2 domain is followed by three fibronectin type III domains (Marino-Buslje *et al.*, 1998, *supra*, Mulhern *et al.*, 1998, Trends Biochem Sci 23:465-66, Ward *et al.*, 1999, Growth Factors 16:315-22). The first FnIII domain (FnIII-1, 461-579) is 118 amino acids in length. The second FnIII domain (FnIII-2, 580-798) is disrupted by a major insert sequence (ID) of about 120 amino acids in length. The ID domain includes a furin protease cleavage site that separates the α and β chains of the mature receptor. The third FnIII domain (FnIII-3) is located entirely in the β-chain (799-901) terminating several residues before the transmembrane sequence. The catalytic domain of the IGF-1R tyrosine kinase is located between amino acids positions 973-1229, and its structure has been solved (Favelyukis *et al.*, 2001, Nature

Structural Biol 8:1058-63, Pautsch *et al.*, 2001, Structure 9:955-65). The kinase is flanked by two regulatory regions, the juxtamembrane region (930-972) and a 108 amino acid C-terminal tail (1220-1337) (Surmacz *et al.*, 1995, Experimental Cell Res 218:370-80, Hongo *et al.*, 1996, Oncogene 12:1231-38). The two regulatory regions contain tyrosine residues that serve as docking sites for signal transducing proteins when phosphorylated by the activated IGF-1R tyrosine kinase (reviewed by Baserga (ed.), 1998 *The IGF-1 Receptor in Normal and Abnormal Growth*, Hormones and Growth Factors in Development and Neoplasia, Wiley-Liss, Inc., Adams *et al.*, 2000, Cell Mol Life Sci 57:1050-93).

The IGF-1R amino acid sequence is about 70% identical to the insulin receptor (INSR; Swiss-Prot: P06213). The highest homology between the receptors is located in the tyrosine kinase domain (84%); the lowest identity is in the CR region and the C-terminus. The IGF-1R is also highly related (~ 55% identical) to the insulin related receptor (IRR; Swiss-Prot: P14616).

Human IGF-1R can be activated by the insulin-like growth factors, IGF-1 and IGF-2 and insulin (INS) (Hill *et al.*, 1985, Pediatric Research 19:879-86). IGF-1 and IGF-2 are encoded nonallelic genes (Brissenden *et al.*, 1984, Nature 310: 781-8, Bell *et al.*, 1985, Proceedings of the National Academy of Sciences of the United States of America 82: 6450-54), and both genes express alternative proteins related by differential RNA splicing and protein processing. The most common and well-studied mature forms of IGF-1 and IGF-2 are respectively 70 and 67 amino acids in length (Jansen *et al.*, 1983, Nature 306:609-11, Dull *et al.*, 1984, Nature 310: 777-81). These proteins (and their isoforms) are identical at 11/21 positions to the insulin A-peptide, and identical at 12/30 positions with the insulin B-peptide.

IGF-1R is expressed in all cells types in the normal adult animal except for liver hepatocytes and mature B-cells. Human blood plasma contains high concentrations of IGF-1 and IGF-2, and IGF-1 can be detected in most tissues. The receptor is an integral component of the physiological mechanism controlling organ size and homeostasis. Without being bound to a particular theory, the "Somatomedin Hypothesis" states that Growth Hormone (GH) mediated somatic growth that occurs during childhood and adolescence is

dependent on the endocrine form of IGF-1 that is mainly produced and secreted by the liver (Daughaday, 2000, Pediatric Nephrology 14: 537-40). The synthesis of hepatic IGF-1 is stimulated by GH release in the pituitary in response to hypothalamic GHRH (GH releasing hormone). The serum concentration of IGF-1 increases over 100 fold between ages 5-15 in humans. The bioavailability of IGF-1 is regulated by IGF binding protein 3 (IGFBP3) with approximately 99% of the growth factor compartmentalized in the bound state. Primary IGF-1 deficiency arising form partial gene deletions, and secondary IGF-1 deficiency resulting from defects in GH production or signaling are not lethal (Woods, 1999, *IGF Deficiency* in Contemporary Endocrinology: The IGF System, R. a.. R. Rosenfeld, C. Jr. Totowa, ed.s, Humana Press, NJ: 651-74). The affected individuals exhibit growth retardation at birth, grow slowly and can face certain CNS abnormalities.

IGF-1R signaling promotes cell growth and survival through the IRS adapter protein-dependent activation of the PI3Kinase/Akt pathway. IGF-1R transmits a signal to its major substrates, IRS-1 through IRS-4 and the Shc proteins (Blakesley *et al.*, 1999, *IGF-1 receptor function: transducing the IGF-1 signal into intracellular events* in The IGF System, R. G. a. R. Rosenfeld, Jr. C.T. Totowa, ed.s, Humana Press, NJ: 143-63). This results in activation of the Ras/Raf/MAP kinase and PI3 Kinase/Akt signaling pathways. However, induction of Akt-mediated cell survival via IRS is the dominant pathway response upon IGF stimulation of most cells. See Figure 10.

Antigen binding proteins

In one aspect, the present invention provides antigen binding proteins (*e.g.*, antibodies, antibody fragments, antibody derivatives, antibody muteins, and antibody variants), that bind to IGF-1R, *e.g.*, human IGF-1R.

Antigen binding proteins in accordance with the present invention include antigen binding proteins that inhibit a biological activity of IGF-1R. Examples of such biological activities include binding a signaling molecule (*e.g.*, IGF-1 and/or IGF-2), and transducing a signal in response to binding a signaling molecule.

Different antigen binding proteins may bind to different domains or epitopes of IGF-1R or act by different mechanisms of action. Examples include but are not limited to antigen binding proteins that interfere with binding of IGF-1 and/or IGF-2 to IGF-1R or that inhibit signal transduction. The site of action may be, for example, intracellular (*e.g.*, by interfering with an intracellular signaling cascade) or extracellular. An antigen binding protein need not completely inhibit an IGF-1 and/or IGF-2 induced activity to find use in the present invention; rather, antigen binding proteins that reduce a particular activity of IGF-1 and/or IGF-2 are contemplated for use as well. (Discussions herein of particular mechanisms of action for IGF-1R-binding antigen binding proteins in treating particular diseases are illustrative only, and the methods presented herein are not bound thereby.)

It has been observed that IGF-1 and IGF-2 each exhibits biphasic binding to IGF-1R. High affinity binding has been reported to have a K_D in the range of 0.2 nM; high affinity binding, about ten fold higher. Thus, in one embodiment, the present invention provides an IGF-1R inhibitor that inhibits both the high and low affinity binding of IGF-1 and/or IGF-2 to IGF-R. It has been suggested that the high affinity binding, rather than the low affinity binding, of IGF-1 and/or IGF-2 to IGF-1R is required for the conformation change that activates the tyrosine kinase activity of IGF-1R. Thus, in another embodiment, the IGF-1R inhibitor preferentially inhibits the high affinity binding of IGF-1 and/or IGF-2 to IGF-1R as compared to the low affinity binding.

In another aspect, the present invention provides antigen binding proteins that comprise a light chain variable region selected from the group consisting of L1 through L52 and/or a heavy chain variable region selected from the group consisting of H1 through H52, and fragments, derivatives, muteins, and variants thereof (see Figures 2 and 3). Such an antigen binding protein can be denoted using the nomenclature "LxHy", wherein "x" corresponds to the number of the light chain variable region and "y" corresponds to the number of the heavy chain variable region as they are labeled in Figures 2 and 3. For example, L2H1 refers to an antigen binding protein with a light chain variable region comprising the amino acid sequence

COMPOSITIONS AND METHODS RELATING TO ANTI-IGF-1 RECEPTOR ANTIBODIES

This application is a divisional application of Application No. 1-2007-501296 filed on June 20, 2007 and now Philippine Patent No. 1-2007-501296 issued on January 7, 2013.

FIELD OF THE INVENTION

This application provides compositions and methods relating to anti-IGF-1 receptor antibodies.

BACKGROUND OF THE INVENTION

Insulin-like growth factors 1 and 2 (IGF-1 and IGF-2, respectively) promote the differentiation and proliferation of a wide variety of mammalian cell types.

IGF-1 and IGF-2 both circulate widely throughout the body in plasma. They exert their effects on cells by binding to and activating the IGF-1 receptor (IGF-1R). IGF-1R is a member of the family of tyrosine kinase growth factor receptors. Its amino acid sequence is about 70% identical to that of the insulin receptor.

Abnormal IGF-1, IGF-2, and/or IGF-1R activities are associated with a number of medical conditions, including various types of cancer, growth defects (*e.g.*, acromegaly, gigantism, and small stature), psoriasis, atherosclerosis, post angioplasty smooth muscle restonsis of blood vessels, diabetes, microvasular proliferation, neuropathy, loss of muscle mass, and osteoporosis.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides nucleotide sequences encoding light chain variable domains L1 through L52 and heavy chain variable domains H1 through H52.

Figure 2 provides amino acid sequences of light chain variable domains L1 through L52. CDR and FR regions are indicated.

06192015000093

2015 MAY 25

of L2 and a heavy chain variable region comprising the amino acid sequence of H1, as shown in Figures 2 and 3. Figures 2 and 3 also indicate the location of the CDR and framework regions of each of these variable domain sequences. The CDR regions of each light and heavy chain also are grouped by type and by sequence similarity in Figures 4 through 9. Antigen binding proteins of the invention include, for example, antigen binding proteins having a combination of light chain and heavy chain variable domains selected from the group of combinations consisting of L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47, L48H48, L49H49, L50H50, L51H51, and L52H52.

In one embodiment, the present invention provides an antigen binding protein comprising a light chain variable domain comprising a sequence of amino acids that differs from the sequence of a light chain variable domain selected from the group consisting of L1 through L52 only at 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 residues, wherein each such sequence difference is independently either a deletion, insertion, or substitution of one amino acid residue. In another embodiment, the light-chain variable domain comprises a sequence of amino acids that is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% identical to the sequence of a light chain variable domain selected from the group consisting of L1 through L52. In another embodiment, the light chain variable domain comprises a sequence of amino acids that is encoded by a nucleotide sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% identical to a nucleotide sequence that encodes a light chain variable domain selected from the group consisting of L1 through L52. In another embodiment, the light chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to the complement of a polynucleotide that encodes a light chain variable domain selected from the group consisting of L1 through L52. In another embodiment, the light chain variable domain

comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to the complement of a polynucleotide that encodes a light chain variable domain selected from the group consisting of L1 through L52. In another embodiment, the light chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to a complement of a light chain polynucleotide selected from Figure 1.

In another embodiment, the present invention provides an antigen binding protein comprising a heavy chain variable domain comprising a sequence of amino acids that differs from the sequence of a heavy chain variable domain selected from the group consisting of H1 through H52 only at 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 residue(s), wherein each such sequence difference is independently either a deletion, insertion, or substitution of one amino acid residue. In another embodiment, the heavy chain variable domain comprises a sequence of amino acids that is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% identical to the sequence of a heavy chain variable domain selected from the group consisting of H1 through H52. In another embodiment, the heavy chain variable domain comprises a sequence of amino acids that is encoded by a nucleotide sequence that is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% identical to a nucleotide sequence that encodes a heavy chain variable domain selected from the group consisting of H1 through H52. In another embodiment, the heavy chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to the complement of a polynucleotide that encodes a heavy chain variable domain selected from the group consisting of H1 through H52. In another embodiment, the heavy chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes under moderately stringent conditions to the complement of a polynucleotide that encodes a heavy chain variable domain selected from the group consisting of H1 through H52. In another embodiment, the heavy chain variable domain comprises a sequence of amino acids that is encoded by a polynucleotide that hybridizes

under moderately stringent conditions to a complement of a heavy chain polynucleotide selected from Figure 1.

Particular embodiments of antigen binding proteins of the present invention comprise one or more amino acid sequences that are identical to the amino acid sequences of one or more of the CDRs and/or FRs illustrated in Figures 2 through 9. In one embodiment, the antigen binding protein comprises a light chain CDR1 sequence illustrated in Figure 4. In another embodiment, the antigen binding protein comprises a light chain CDR2 sequence illustrated in Figure 5. In another embodiment, the antigen binding protein comprises a light chain CDR3 sequence illustrated in Figure 6. In another embodiment, the antigen binding protein comprises a heavy chain CDR1 sequence illustrated in Figure 7. In another embodiment, the antigen binding protein comprises a heavy chain CDR2 sequence illustrated in Figure 8. In another embodiment, the antigen binding protein comprises a heavy chain CDR3 sequence illustrated in Figure 9. In another embodiment, the antigen binding protein comprises a light chain FR1 sequence illustrated in Figure 2. In another embodiment, the antigen binding protein comprises a light chain FR2 sequence illustrated in Figure 2. In another embodiment, the antigen binding protein comprises a light chain FR3 sequence illustrated in Figure 2. In another embodiment, the antigen binding protein comprises a light chain FR4 sequence illustrated in Figure 2. In another embodiment, the antigen binding protein comprises a heavy chain FR1 sequence illustrated in Figure 3. In another embodiment, the antigen binding protein comprises a heavy chain FR2 sequence illustrated in Figure 3. In another embodiment, the antigen binding protein comprises a heavy chain FR3 sequence illustrated in Figure 3. In another embodiment, the antigen binding protein comprises a heavy chain FR4 sequence illustrated in Figure 3.

In one embodiment, the present invention provides an antigen binding protein that comprises one or more CDR sequences that differ from a CDR sequence shown in Figures 2 through 9 by no more than 5, 4, 3, 2, or 1 amino acid residues.

In one embodiment, the present invention provides an antigen binding protein that comprises at least one CDR from L1-L52 and/or H1-H52, as

shown in Figures 2 through 9, and at least one CDR sequence from an anti-IGF-1R antibody described in US Pat. App. Pub. Nos. 03/0235582, 04/0228859, 04/0265307, 04/0886503, 05/0008642, 05/0084906, 05/0186203, 05/0244408, PCT Pub. Nos. WO 03/059951, WO 03/100008, WO 04/071529A2, WO 04/083248, WO 04/087756, WO 05/016967, WO 05/016970, or WO 05/058967 (each of which is incorporated herein by reference in its entirety for all purposes) wherein the antigen binding protein binds to IGF-1 receptor. In another embodiment, the antigen binding protein comprises 2, 3, 4, or 5 CDR sequences from L1-L52 and/or H1-H52, as shown in Figures 2 through 9. In another embodiment, the antigen binding protein comprises 2, 3, 4, or 5 CDR sequences from an anti-IGF-1R antibody described in US Pat. App. Pub. Nos. 03/0235582, 04/0228859, 04/0265307, 04/0886503, 05/0008642, 05/0084906, 05/0186203, 05/0244408, PCT Pub. Nos. WO 03/059951, WO 03/100008, WO 04/071529A2, WO 04/083248, WO 04/087756, WO 05/016967, WO 05/016970, or WO 05/058967. In another embodiment, at least one of the antigen binding protein's CDR3 sequences is a CDR3 sequence from L1-L52 and/or H1-H52, as shown in Figures 2, 3, 6, and 9. In another embodiment, the antigen binding protein's light chain CDR3 sequence is a light chain CDR3 sequence from L1-L52 as shown in Figures 2 and 6 and the antigen binding protein's heavy chain CDR3 sequence is a heavy chain sequence from H1-H52 as shown in Figures 3 and 9. In another embodiment, the antigen binding protein comprises 1, 2, 3, 4, or 5 CDR sequences that each independently differs by 6, 5, 4, 3, 2, 1, or 0 single amino acid additions, substitutions, and/or deletions from a CDR sequence of L1-L52 and/or H1-H52, and the antigen binding protein further comprises 1, 2, 3, 4, or 5 CDR sequences that each independently differs by 6, 5, 4, 3, 2, 1, or 0 single amino acid additions, substitutions, and/or deletions from a CDR sequence of US Pat. App. Pub. Nos. 03/0235582, 04/0228859, 04/0265307, 04/0886503, 05/0008642, 05/0084906, 05/0186203, 05/0244408, PCT Pub. Nos. WO 03/059951, WO 03/100008, WO 04/071529A2, WO 04/083248, WO 04/087756, WO 05/016967, WO 05/016970, or WO 05/058967. In another embodiment, the CDR sequence(s) from US Pat. App. Pub. Nos. 03/0235582, 04/0228859, 04/0265307, 04/0886503, 05/0008642, 05/0084906,

05/0186203, 05/0244408, PCT Pub. Nos. WO 03/059951, WO 03/100008, WO 04/071529A2, WO 04/083248, WO 04/087756, WO 05/016967, WO 05/016970, or WO 05/058967. In another embodiment, the CDR sequence(s) are from (an) antibody(-ies) that bind(s) to the L2 portion of the extracellular domain of IGF-1 receptor. In another embodiment, the antigen binding protein does not comprise a light chain CDR3 sequence and/or a heavy chain CDR3 sequence from an anti-IGF-1R antibody from US Pat. App. Pub. Nos. 03/0235582, 04/0228859, 04/0265307, 04/0886503, 05/0008642, 05/0084906, 05/0186203, 05/0244408, PCT Pub. Nos. WO 03/059951, WO 03/100008, WO 04/071529A2, WO 04/083248, WO 04/087756, WO 05/016967, WO 05/016970, or WO 05/058967.

In one embodiment, the present invention provides an antigen binding protein that comprises a light chain CDR1 comprising the sequence $RSSQSLLHX_1X_2GYNX_3LX_4$ (SEQ ID NO:236), wherein X_1 is a serine or a threonine residue, X_2 is an asparagine, serine, or histidine residue, X_3 is a tyrosine or a phenylalanine residue, and X_4 is an aspartate or an asparagine residue. In another embodiment, the light chain CDR1 comprises the sequence $TRSSGX_1IX_2X_3NYVQ$ (SEQ ID NO:237), wherein X_1 is a serine or an aspartate residue, X_2 is an alanine or an aspartate residue, and X_3 is a serine or an asparagine residue. In another embodiment, the light chain CDR1 comprises the sequence $RASQX_1X_2X_3X_4X_5LX_6$ (SEQ ID NO:238), wherein X_1 is a glycine or a serine residue, X_2 is an isoleucine, valine, or proline residue, and X_3 is a serine, glycine, or tyrosine residue, X_4 is any amino acid residue, X_5 is a phenylalanine, tyrosine, asparagine, or tryptophan residue, and X_6 is an alanine or an asparagine residue. In another embodiment, X_2 is an isoleucine or valine residue, X_3 is a glycine or serine residue, X_4 is an arginine, serine, asparagine, serine, tyrosine, or isoleucine residue, and X_5 is a phenylalanine or a tyrosine residue.

In one embodiment, the present invention provides an antigen binding protein that comprises a light chain CDR2 comprising the sequence $LX_1X_2X_3RX_4S$ (SEQ ID NO:239), wherein X_1 is a glycine or a valine residue, X_2 is a serine or a phenylalanine residue, X_3 is an asparagine, tyrosine, or threonine residue, and X_4 is an alanine or an aspartate residue. In another

embodiment, the CDR2 comprises the sequence $AX_1SX_2LX_3S$ (SEQ ID NO:240), wherein X_1 is an alanine or a threonine residue, X_2 is a threonine or a glycine residue, and X_3 is a glutamine or a glutamate residue. In another embodiment, the CDR2 comprises the sequence $X_1X_2NX_3RPS$ (SEQ ID NO:241), wherein X_1 is a glutamate, glutamine, or glycine residue, X_2 is an aspartate or lysine residue, and X_3 is any amino acid residue.

In one embodiment, the present invention provides an antigen binding protein that comprises a light chain CDR3 comprising the sequence $MX_1X_2X_3X_4X_5PX_6X_7$ (SEQ ID NO:242), wherein X_1 is a glutamine or glutamate residue, X_2 is an alanine, glycine, serine, or threonine residue, X_3 is a leucine or threonine residue, X_4 is a glutamine, glutamate, or histidine residue, X_5 is a threonine, tryptophan, methionine, or valine residue, X_6 is a nonpolar side chain residue, and X_7 is a threonine, serine, or alanine residue. In another embodiment, the CDR3 comprises the sequence $QQX_1X_2X_3X_4PX_5T$ (SEQ ID NO:243), wherein X_1 is an arginine, serine, leucine, or alanine residue, X_2 is an asparagine, serine, or histidine residue, X_3 is a serine or an asparagine residue, X_4 is a nonpolar side chain residue, and X_5 is a leucine, isoleucine, tyrosine, or tryptophan residue. In another embodiment, the CDR3 comprises the sequence $QSYX_1SX_2NX_3X_4V$ (SEQ ID NO:244), wherein X_1 is an aspartate or a glutamine residue, X_2 is a serine or an aspartate residue, X_3 is a glutamine, valine, or tryptophan residue, and X_4 is an arginine residue or no residue.

In one embodiment, the present invention provides an antigen binding protein that comprises a heavy chain CDR1 comprising the sequence $X_1X_2X_3WWS$ (SEQ ID NO:245), wherein X_1 is a serine residue or no residue, X_2 is a serine or asparagine residue, and X_3 is an asparagine residue and an isoleucine residue. In another embodiment, the heavy chain CDR1 comprises the sequence X_1X_2YWS (SEQ ID NO:246), wherein X_1 is a glycine, asparagine, or aspartate residue, and X_2 is a tyrosine or phenylalanine residue. In another embodiment, the heavy chain CDR1 comprises the sequence $SYX_1X_2X_3$ (SEQ ID NO:247), wherein X_1 is an alanine or glycine residue, X_2 is a methionine or isoleucine residue, and X_3 is a serine or histidine residue.

In one embodiment, the present invention provides an antigen binding protein that comprises a heavy chain CDR2 comprising the sequence $X_1X_2X_3X_4X_5GX_6TX_7YNPSLX_8S$ (SEQ ID NO:248), wherein X_1 is a glutamate, tyrosine, or serine residue, X_2 is a isoleucine or valine residue, X_3 is a tyrosine, asparagine, or serine residue, X_4 is a histidine, tyrosine, aspartate, or proline residue, X_5 is a serine or arginine residue, X_6 is a serine or asparagine residue, X_7 is an asparagine or tyrosine residue, and X_8 is a lysine or glutamate residue. In another embodiment, the heavy chain CDR2 comprises the sequence $X_1ISX_2X_3X_4X_5X_6X_7YYADSVKG$ (SEQ ID NO:249), wherein X_1 is a threonine, alanine, valine, or tyrosine residue, X_2 is a glycine, serine, or tyrosine residue, X_3 is a serine, asparagine, or aspartate residue, X_4 is a glycine or serine residue, X_5 is a glycine, serine, or aspartate residue, X_6 is a serine, threonine, or asparagine residue, and X_7 is a threonine, lysine, or isoleucine residue.

In one embodiment, the present invention provides an antigen binding protein that comprises a heavy chain CDR3 comprising the sequence $X_1X_2X_3X_4X_5X_6X_7X_8X_9FDI$ (SEQ ID NO:250), wherein X_1 is a glutamate residue or no residue, X_2 is tyrosine, glycine, or serine residue or no residue, X_3 is a serine, asparagine, tryptophan, or glutamate residue, or no residue, X_4 is a serine, aspartate, tryptophan, alanine, arginine, threonine, glutamine, leucine, or glutamate residue, or no residue, X_5 is a serine, glycine, asparagine, threonine, tryptophan, alanine, valine, or isoleucine residue, X_6 is an arginine, glutamine, tyrosine, valine, alanine, glycine, serine, phenylalanine, or tryptophan residue, X_7 is a leucine, asparagine, aspartate, threonine, tryptophan, tyrosine, valine, alanine, or histidine residue, X_8 is an aspartate, serine, asparagine, or glutamine residue, and X_9 is an alanine or a proline residue. In another embodiment, the heavy chain CDR3 comprises the sequence $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}X_{11}MDV$ (SEQ ID NO:251), wherein X_1 is an alanine residue, or no residue, X_2 is a glutamate, tyrosine, or glycine residue, or no residue, X_3 is a serine or arginine residue, or no residue, X_4 is an aspartate, glycine, serine, or valine residue, or no residue, X_5 is a serine, glycine, or aspartate residue, or no residue, X_6 is a glycine, phenylalanine, aspartate, serine, tryptophan, or tyrosine residue, or no residue, X_7 is a

tyrosine, tryptophan, serine, or aspartate residue, or no residue, X_8 is an aspartate, arginine, serine, glycine, tyrosine, or tryptophan residue, X_9 is a tyrosine, isoleucine, leucine, phenylalanine, or lysine residue, X_{10} is a tyrosine, phenylalanine, aspartate, or glycine residue, and X_{11} is a glycine, tyrosine, or asparagine residue. In another embodiment, the heavy chain CDR3 comprises the sequence $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}Y$ (SEQ ID NO:252), wherein X_1 is an aspartate or valine residue, or no residue, X_2 is a glycine, tyrosine, arginine, or aspartate residue, or no residue, X_3 is an asparagine, leucine, glycine, isoleucine, serine, valine, phenylalanine, or tyrosine residue, or no residue, X_4 is a leucine, serine, tryptophan, alanine, tyrosine, isoleucine, glycine, or aspartate residue, or no residue, X_5 is a glycine, alanine, tyrosine, serine, aspartate, or leucine residue, X_6 is a valine, alanine, glycine, threonine, proline, histidine, or glutamine residue, X_7 is a glutamate, glycine, serine, aspartate, glycine, valine, tryptophan, histidine, or arginine residue, X_8 is a glutamine, alanine, glycine, tyrosine, proline, leucine, aspartate, or serine residue, X_9 is a nonpolar side chain residue, and X_{10} is an aspartate or alanine residue. In another embodiment, the heavy chain CDR3 comprises the sequence $X_1X_2X_3X_4X_5X_6X_7X_8X_9X_{10}YFDX_{11}$ (SEQ ID NO:253), wherein X_1 is a glycine residue, or no residue, X_2 is a proline residue, or no residue, X_3 is an arginine or aspartate residue, or no residue, X_4 is a histidine or proline residue, X_5 is an arginine or glycine residue, X_6 is an arginine, serine, or phenylalanine residue, X_7 is an aspartate or serine residue, X_8 is a glycine, tryptophan, or tyrosine residue, X_9 is a tyrosine or alanine residue, X_{10} is an asparagine or tryptophan residue, and X_{11} is an asparagine or leucine residue. In another embodiment, the heavy chain CDR3 comprises the sequence $X_1X_2X_3X_4DSSX_5X_6X_7X_8X_9X_{10}X_{11}X_{12}$ (SEQ ID NO:254), wherein X_1 is a phenylalanine residue, or no residue, X_2 is an asparagine or glycine residue, or no residue, X_3 is a tyrosine or a leucine residue, or no residue, X_4 is a tyrosine or glycine residue, or no residue, X_5 is a glycine, serine, or valine residue, X_6 is a tyrosine, phenylalanine, tryptophan, or glutamine residue, or no residue, X_7 is a tyrosine, glycine, or isoleucine residue, or no residue, X_8 is a tyrosine, leucine, or glycine residue, or no residue, X_9 is a methionine, glycine, or phenylalanine residue, or no residue, X_{10} is an aspartate or methionine residue,

or no residue, X_{11} is a valine, aspartate, or tyrosine residue, or no residue, and X_{12} is a valine residue, or no residue.

In one embodiment, the present invention provides an isolated antigen binding protein, comprising either: a. a light chain CDR3 comprising a sequence selected from the group consisting of: i. a light chain CDR3 sequence selected from the group consisting of the light chain CDR3 sequences of L1-L52 as shown in Figure 6; ii. MQALQTPZT; iii. QQ(R/S)(N/S)(S/N)ZPLT; and iv. QSYDSSNXJV; b. a heavy chain CDR3 comprising a sequence selected from the group consisting of: i. a heavy chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, or deletions from a CDR3 sequence selected from the group consisting of the heavy chain CDR3 sequences of H1-H52 as shown in Figure 9; ii. SRLDAFDI; iii. SXYDYYGMDV; iv. HRXDXAWYFDL; and v. DSSG; or c. the light chain CDR3 sequence of (a) and the heavy chain CDR3 sequence of (b); wherein amino acid residue symbols enclosed in parentheses identify alternative residues for the same position in a sequence, each X is independently any amino acid residue, each Z is independently a glycine residue, an alanine residue, a valine residue, a leucine residue, an isoleucine residue, a proline residue, a phenylalanine residue, a methionine residue, a tryptophan residue, or a cysteine residue, each J is independently a glutamine residue, an arginine residue, a valine residue, or a tryptophan residue, and the antigen binding protein binds to human IGF-1R.

The nucleotide sequences of Figure 1, or the amino acid sequences of Figures 2 through 9, can be altered, for example, by random mutagenesis or by site-directed mutagenesis (*e.g.*, oligonucleotide-directed site-specific mutagenesis) to create an altered polynucleotide comprising one or more particular nucleotide substitutions, deletions, or insertions as compared to the non-mutated polynucleotide. Examples of techniques for making such alterations are described in Walder *et al.*, 1986,Gene 42:133; Bauer *et al.*1985, Gene 37:73; Craik, BioTechniques, January 1985, 12-19; Smith *et al.*, 1981, *Genetic Engineering: Principles and Methods*, Plenum Press; and U.S. Patent Nos. 4,518,584 and 4,737,462. These and other methods can be used to make, for example, derivatives of anti-IGF-1R antibodies that have a desired

property, for example, increased affinity, avidity, or specificity for IGF-1R, increased activity or stability *in vivo* or *in vitro*, or reduced *in vivo* side-effects as compared to the underivatized antibody.

Other derivatives of anti- IGF-1R antibodies within the scope of this invention include covalent or aggregative conjugates of anti-IGF-1R antibodies, or fragments thereof, with other proteins or polypeptides, such as by expression of recombinant fusion proteins comprising heterologous polypeptides fused to the N-terminus or C-terminus of an anti- IGF-1R antibody polypeptide. For example, the conjugated peptide may be a heterologous signal (or leader) polypeptide, *e.g.*, the yeast alpha-factor leader, or a peptide such as an epitope tag. Antigen binding protein-containing fusion proteins can comprise peptides added to facilitate purification or identification of antigen binding protein (*e.g.*, poly-His). An antigen binding protein also can be linked to the FLAG peptide Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys (DYKDDDDK) (SEQ ID NO:255) as described in Hopp *et al.*, *Bio/Technology* 6:1204, 1988, and U.S. Patent 5,011,912. The FLAG peptide is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody (mAb), enabling rapid assay and facile purification of expressed recombinant protein. Reagents useful for preparing fusion proteins in which the FLAG peptide is fused to a given polypeptide are commercially available (Sigma, St. Louis, MO).

Oligomers that contain one or more antigen binding proteins may be employed as IGF-1R antagonists. Oligomers may be in the form of covalently-linked or non-covalently-linked dimers, trimers, or higher oligomers. Oligomers comprising two or more antigen binding protein are contemplated for use, with one example being a homodimer. Other oligomers include heterodimers, homotrimers, heterotrimers, homotetramers, heterotetramers, *etc.*

One embodiment is directed to oligomers comprising multiple antigen binding proteins joined *via* covalent or non-covalent interactions between peptide moieties fused to theantigen binding proteins. Such peptides may be peptide linkers (spacers), or peptides that have the property of promoting oligomerization. Leucine zippers and certain polypeptides derived from

Figure 5 provides amino acid sequences of the light chain CDR2 regions of light chain variable domains L1 through L52. Consensus sequences for groups of related CDR sequences are also provided.

Figure 6 provides amino acid sequences of the light chain CDR3 regions of light chain variable domains L1 through L52. Consensus sequences for groups of related CDR sequences are also provided.

Figure 7 provides amino acid sequences of the heavy chain CDR1 regions of heavy chain variable domains H1 through H52. Consensus sequences for groups of related CDR sequences are also provided.

Figure 8 provides amino acid sequences of the heavy chain CDR2 regions of heavy chain variable domains H1 through H52. Consensus sequences for groups of related CDR sequences are also provided.

Figure 9 provides amino acid sequences of the heavy chain CDR3 regions of heavy chain variable domains H1 through H52. Consensus sequences for groups of related CDR sequences are also provided.

Figure 10 provides the amino acid sequence of a human IGF-1R extracellular domain fused to a human IgG1 Fc region (underlined) with an intervening caspace-3 cleavage site (bold).

Figure 11 provides the amino acid sequence of a human insulin receptor extracellular domain fused to a human IgG1 Fc region (underlined).

Figure 12 provides the protein sequence of a human IGF-1R extracellular domain (including signal peptide) fused at the C-terminus with chicken avidin. The initiating met in the IGF-1R ECD is designated position 1 in this figure.

Figure 13 provides the polypeptide sequence of a human kappa light chain antibody constant region and a human IgG1 heavy chain antibody constant region.

Figure 14 provides a graph illustrating that four phage-displayed antibodies bind significantly better to an IGF-1R-Fc molecule than they bind to an insulin-receptor-Fc or a murine Fc.

Figure 15 provides graphs illustrating the ability of certain antibodies to compete for binding to IGF-1R with IGF-1 and IGF-2.

antibodies are among the peptides that can promote oligomerization of antigen binding proteins attached thereto, as described in more detail below.

In particular embodiments, the oligomers comprise from two to four antigen binding proteins. The antigen binding proteins of the oligomer may be in any form, such as any of the forms described above, *e.g.*, variants or fragments. Preferably, the oligomers comprise antigen binding proteins that have IGF-1R binding activity.

In one embodiment, an oligomer is prepared using polypeptides derived from immunoglobulins. Preparation of fusion proteins comprising certain heterologous polypeptides fused to various portions of antibody-derived polypeptides (including the Fc domain) has been described, *e.g.*, by Ashkenazi *et al.*, 1991, PNAS USA 88:10535; Byrn *et al.*, 1990, Nature 344:677; and Hollenbaugh *et al.*, 1992 "Construction of Immunoglobulin Fusion Proteins", in *Current Protocols in Immunology*, Suppl. 4, pages 10.19.1 - 10.19.11.

One embodiment of the present invention is directed to a dimer comprising two fusion proteins created by fusing an IGF-1R binding fragment of an anti- IGF-1R antibody to the Fc region of an antibody. The dimer can be made by, for example, inserting a gene fusion encoding the fusion protein into an appropriate expression vector, expressing the gene fusion in host cells transformed with the recombinant expression vector, and allowing the expressed fusion protein to assemble much like antibody molecules, whereupon interchain disulfide bonds form between the Fc moieties to yield the dimer.

The term "Fc polypeptide" as used herein includes native and mutein forms of polypeptides derived from the Fc region of an antibody. Truncated forms of such polypeptides containing the hinge region that promotes dimerization also are included. Fusion proteins comprising Fc moieties (and oligomers formed therefrom) offer the advantage of facile purification by affinity chromatography over Protein A or Protein G columns.

One suitable Fc polypeptide, described in PCT application WO 93/10151 (hereby incorporated by reference), is a single chain polypeptide extending from the N-terminal hinge region to the native C-terminus of the Fc

region of a human IgG1 antibody. Another useful Fc polypeptide is the Fc mutein described in U.S. Patent 5,457,035 and in Baum *et al.*, 1994, EMBO J. 13:3992-4001. The amino acid sequence of this mutein is identical to that of the native Fc sequence presented in WO 93/10151, except that amino acid 19 has been changed from Leu to Ala, amino acid 20 has been changed from Leu to Glu, and amino acid 22 has been changed from Gly to Ala. The mutein exhibits reduced affinity for Fc receptors.

In other embodiments, the variable portion of the heavy and/or light chains of an anti- IGF-1R antibody may be substituted for the variable portion of an antibody heavy and/or light chain.

Alternatively, the oligomer is a fusion protein comprising multiple antigen binding proteins, with or without peptide linkers (spacer peptides). Among the suitable peptide linkers are those described in U.S. Patents 4,751,180 and 4,935,233.

Another method for preparing oligomeric antigen binding proteins involves use of a leucine zipper. Leucine zipper domains are peptides that promote oligomerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz *et al.*, 1988, Science 240:1759), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble oligomeric proteins are described in PCT application WO 94/10308, and the leucine zipper derived from lung surfactant protein D (SPD) described in Hoppe *et al.*, 1994, FEBS Letters 344:191, hereby incorporated by reference. The use of a modified leucine zipper that allows for stable trimerization of a heterologous protein fused thereto is described in Fanslow *et al.*, 1994, Semin. Immunol. 6:267-78. In one approach, recombinant fusion proteins comprising an anti-IGF-1R antibody fragment or derivative fused to a leucine zipper peptide are expressed in suitable host cells, and the soluble oligomeric anti- IGF-1R antibody fragments or derivatives that form are recovered from the culture supernatant.

In one aspect, the present invention provides antigen binding proteins that interfere with the binding of IGF-1 and/or IGF-2 to an IGF-1R. Such antigen binding proteins can be made against IGF-1R, or a fragment, variant or derivative thereof, and screened in conventional assays for the ability to interfere with binding of IGF-1 and/or IGF-2 to IGF-1R. Examples of suitable assays are assays that test the antigen binding proteins for the ability to inhibit binding of IGF-1 and/or IGF-2 to cells expressing IGF-1R, or that test antigen binding proteins for the ability to reduce a biological or cellular response that results from the binding of IGF-1 and/or IGF-2 to cell surface IGF-1R receptors.

In another aspect, the present invention provides an antigen binding protein that blocks the binding of IGF-1 and/or IGF-2 to IGF-1R but does not significantly block the binding of insulin to insulin receptor (INS-R). In one embodiment, the antigen binding protein does not bind to INS-R. In another embodiment, the antigen binding protein binds to the INS-R with such a low affinity that it does not effectively block the binding of insulin to INS-R. In another embodiment, the antigen binding protein binds to INS-R, but antigen binding protein-bound INS-R can still bind to insulin. In another embodiment, the antigen binding protein's selectivity for IGF-1R is at least 50 times greater than its selectivity for insulin receptor. In another embodiment, the selectivity of the antigen binding protein is more than 100 times greater than its selectivity for insulin receptor.

In another aspect, the present invention provides an antigen binding protein that demonstrates species selectivity. In one embodiment, the antigen binding protein binds to one or more mammalian IGF-1R, for example, to human IGF-1R and one or more of mouse, rat, guinea pig, hamster, gerbil, cat, rabbit, dog, goat, sheep, cow, horse, camel, and non-human primate IGF-1R. In another embodiment, the antigen binding protein binds to one or more primate IGF-1R, for example, to human IGF-1R and one or more of cynomologous, marmoset, rhesus, and chimpanzee IGF-1R. In another embodiment, the antigen binding protein binds specifically to human, cynomologous, marmoset, rhesus, or chimpanzee IGF-1R. In another embodiment, the antigen binding protein does not bind to one or more of

mouse, rat, guinea pig, hamster, gerbil, cat, rabbit, dog, goat, sheep, cow, horse, camel, and non-human primate IGF-1R. In another embodiment, the antigen binding protein does not bind to a New World monkey species such as a marmoset. In another embodiment, the antigen binding protein does not exhibit specific binding to any naturally occurring protein other than IGF-1R. In another embodiment, the antigen binding protein does not exhibit specific binding to any naturally occurring protein other than mammalian IGF-1R. In another embodiment, the antigen binding protein does not exhibit specific binding to any naturally occurring protein other than primate IGF-1R. In another embodiment, the antigen binding protein does not exhibit specific binding to any naturally occurring protein other than human IGF-1R. In another embodiment, the antigen binding protein specifically binds to mouse, rat, cynomolgus monkey, and human IGF-1R. In another embodiment, the antigen binding protein specifically binds to mouse, rat, cynomolgus monkey, and human IGF-1R with a similar binding affinity. In another embodiment, the antigen binding protein blocks binding of human IGF-1 and IGF-2 with mouse, rat, cynomolgus monkey, and human IGF-1R. In another embodiment, the antigen binding protein blocks binding of human IGF-1 and IGF-2 with mouse, rat, cynomolgus monkey, and human IGF-1R with similar K_i. In another embodiment, the antigen binding protein blocks binding of human IGF-1 and IGF-2 with mouse, rat, cynomolgus monkey, and human IGF-1R with a K_i of between about 0.57 and about 0.61 nM.

One may determine the selectivity of an antigen binding protein for an IGF-1R using methods well known in the art and following the teachings of the specification. For example, one may determine the selectivity using Western blot, FACS, ELISA or RIA.

In another aspect, the present invention provides an IGF-1R binding antigen binding protein (for example, an anti-IGF-1R antibody), that has one or more of the following characteristics: binds to both human and murine IGF-1R, inhibits the binding of both IGF-1 and IGF-2 to human IGF-1R, inhibits the binding of both IGF-1 and IGF-2 to murine IGF-1R, preferentially inhibits the high affinity binding of IGF-1 and/or of IGF-2 to IGF-1R, binds to the L2 domain of IGF-1R, causes relatively little down-regulation of cell-

surface expressed IGF-1R after 17 hours of exposure (as compared to MAB391 (R&D systems, Minneapolis, MN); e.g., amount of IGF-1R is reduced by less than 20%), causes a level of down-regulation of cell-surface expressed IGF-1R on Colo-205 or MiaPaCa-2 xenograft tumor cells in mice as MAB391 after four weeks of once weekly doses of 200 micrograms.

Antigen-binding fragments of antigen binding proteins of the invention may be produced by conventional techniques. Examples of such fragments include, but are not limited to, Fab and $F(ab')_2$ fragments. Antibody fragments and derivatives produced by genetic engineering techniques also are contemplated.

Additional embodiments include chimeric antibodies, *e.g.*, humanized versions of non-human (*e.g.*, murine) monoclonal antibodies. Such humanized antibodies may be prepared by known techniques, and offer the advantage of reduced immunogenicity when the antibodies are administered to humans. In one embodiment, a humanized monoclonal antibody comprises the variable domain of a murine antibody (or all or part of the antigen binding site thereof) and a constant domain derived from a human antibody. Alternatively, a humanized antibody fragment may comprise the antigen binding site of a murine monoclonal antibody and a variable domain fragment (lacking the antigen-binding site) derived from a human antibody. Procedures for the production of chimeric and further engineered monoclonal antibodies include those described in Riechmann *et al.*, 1988, Nature 332:323, Liu *et al.*, 1987, Proc. Nat. Acad. Sci. USA 84:3439, Larrick *et al.*, 1989, Bio/Technology 7:934, and Winter *et al.*, 1993, TIPS 14:139. In one embodiment, the chimeric antibody is a CDR grafted antibody. Techniques for humanizing antibodies are discussed in, *e.g.*, U.S. Pat. App. No. 10/194,975 (published February 27, 2003), U.S. Pat. No.s 5,869,619, 5,225,539, 5,821,337, 5,859,205, Padlan *et al.*, 1995, FASEB J. 9:133-39, and Tamura *et al.*, 2000, J. Immunol. 164:1432-41.

Procedures have been developed for generating human or partially human antibodies in non-human animals. For example, mice in which one or more endogenous immunoglobulin genes have been inactivated by various means have been prepared. Human immunoglobulin genes have been

introduced into the mice to replace the inactivated mouse genes. Antibodies produced in the animal incorporate human immunoglobulin polypeptide chains encoded by the human genetic material introduced into the animal. In one embodiment, a non-human animal, such as a transgenic mouse, is immunized with an IGF-1R polypeptide, such that antibodies directed against the IGF-1R polypeptide are generated in the animal. One example of a suitable immunogen is a soluble human IGF-1R, such as a polypeptide comprising the extracellular domain of the protein of Figure 10, or other immunogenic fragment of the protein of Figure 10. Examples of techniques for production and use of transgenic animals for the production of human or partially human antibodies are described in U.S. Patents 5,814,318, 5,569,825, and 5,545,806, Davis *et al.*, 2003, *Production of human antibodies from transgenic mice* in Lo, ed. Antibody Engineering: Methods and Protocols, Humana Press, NJ:191-200, Kellermann *et al.*, 2002, Curr Opin Biotechnol. 13:593-97, Russel *et al.*, 2000, Infect Immun. 68:1820-26, Gallo *et al.*, 2000, Eur J Immun. 30:534-40, Davis *et al.*, 1999, Cancer Metastasis Rev. 18:421-25, Green, 1999, J Immunol Methods. 231:11-23, Jakobovits, 1998, Advanced Drug Delivery Reviews 31:33-42, Green *et al.*, 1998, J Exp Med. 188:483-95, Jakobovits A, 1998, Exp. Opin. Invest. Drugs. 7:607-14, Tsuda *et al.*, 1997, Genomics. 42:413-21, Mendez *et al.*, 1997, Nat Genet. 15:146-56, Jakobovits, 1994, Curr Biol. 4:761-63, Arbones *et al.*, 1994, Immunity. 1:247-60, Green *et al.*, 1994, Nat Genet. 7:13-21, Jakobovits *et al.*, 1993, Nature. 362:255-58, Jakobovits *et al.*, 1993, Proc Natl Acad Sci U S A. 90:2551-55. Chen, J., M. Trounstine, F. W. Alt, F. Young, C. Kurahara, J. Loring, D. Huszar. "Immunoglobulin gene rearrangement in B cell deficient mice generated by targeted deletion of the JH locus." International Immunology 5 (1993): 647-656, Choi *et al.*, 1993, Nature Genetics 4: 117-23, Fishwild *et al.*, 1996, Nature Biotechnology 14: 845-51, Harding *et al.*, 1995, Annals of the New York Academy of Sciences, Lonberg *et al.*, 1994, Nature 368: 856-59, Lonberg, 1994, *Transgenic Approaches to Human Monoclonal Antibodies* in Handbook of Experimental Pharmacology 113: 49-101, Lonberg *et al.*, 1995, Internal Review of Immunology 13: 65-93, Neuberger, 1996, Nature Biotechnology 14: 826, Taylor *et al.*, 1992, Nucleic Acids Research 20: 6287-

95, Taylor *et al.*, 1994, International Immunology 6: 579-91, Tomizuka *et al.*, 1997, Nature Genetics 16: 133-43, Tomizuka *et al.*, 2000, Proceedings of the National Academy of Sciences USA 97: 722-27, Tuaillon *et al.*, 1993, Proceedings of the National Academy of Sciences USA 90: 3720-24, and Tuaillon *et al.*, 1994, Journal of Immunology 152: 2912-20.

In another aspect, the present invention provides monoclonal antibodies that bind to IGF-1R. Monoclonal antibodies may be produced using any technique known in the art, *e.g.*, by immortalizing spleen cells harvested from the transgenic animal after completion of the immunization schedule. The spleen cells can be immortalized using any technique known in the art, *e.g.*, by fusing them with myeloma cells to produce hybridomas. Myeloma cells for use in hybridoma-producing fusion procedures preferably are non-antibody-producing, have high fusion efficiency, and enzyme deficiencies that render them incapable of growing in certain selective media which support the growth of only the desired fused cells (hybridomas). Examples of suitable cell lines for use in mouse fusions include Sp-20, P3-X63/Ag8, P3-X63-Ag8.653, NS1/1.Ag 4 1, Sp210-Ag14, FO, NSO/U, MPC-11, MPC11-X45-GTG 1.7 and S194/5XX0 Bul; examples of cell lines used in rat fusions include R210.RCY3, Y3-Ag 1.2.3, IR983F and 4B210. Other cell lines useful for cell fusions are U-266, GM1500-GRG2, LICR-LON-HMy2 and UC729-6.

In one embodiment, a hybridoma cell line is produced by immunizing an animal (*e.g.*, a transgenic animal having human immunoglobulin sequences) with an IGF-1R immunogen; harvesting spleen cells from the immunized animal; fusing the harvested spleen cells to a myeloma cell line, thereby generating hybridoma cells; establishing hybridoma cell lines from the hybridoma cells, and identifying a hybridoma cell line that produces an antibody that binds an IGF-1R polypeptide. Such hybridoma cell lines, and anti-IGF-1R monoclonal antibodies produced by them, are encompassed by the present invention.

Monoclonal antibodies secreted by a hybridoma cell line can be purified using any technique known in the art. Hybridomas or mAbs may be further screened to identify mAbs with particular properties, such as the ability

to block an IGF-1 and/or IGF-2 induced activity. Examples of such screens are provided in the examples below.

Molecular evolution of the complementarity determining regions (CDRs) in the center of the antibody binding site also has been used to isolate antibodies with increased affinity, for example, antibodies having increased affinity for c-erbB-2, as described by Schier *et al.*, 1996, J. Mol. Biol. 263:551. Accordingly, such techniques are useful in preparing antibodies to IGF-1R.

Antigen binding proteins directed against an IGF-1R can be used, for example, in assays to detect the presence of IGF-1R polypeptides, either *in vitro* or *in vivo*. The antigen binding proteins also may be employed in purifying IGF-1R proteins by immunoaffinity chromatography. Those antigen binding proteins that additionally can block binding of IGF-1 and/or IGF-2 to IGF-1R may be used to inhibit a biological activity that results from such binding. Blocking antigen binding proteins can be used in the methods of the present invention. Such antigen binding proteins that function as IGF-1 and/or IGF-2 antagonists may be employed in treating any IGF-1 and/or IGF-2-induced condition, including but not limited to cancer. In one embodiment, a human anti- IGF-1R monoclonal antibody generated by procedures involving immunization of transgenic mice is employed in treating such conditions.

Antigen binding proteins may be employed in an *in vitro* procedure, or administered *in vivo* to inhibit an IGF-1 and/or IGF-2-induced biological activity. Disorders caused or exacerbated (directly or indirectly) by the interaction of IGF-1 and/or IGF-2 with cell surface IGF-1R, examples of which are provided above, thus may be treated. In one embodiment, the present invention provides a therapeutic method comprising *in vivo* administration of an IGF-1 and/or IGF-2 blocking antigen binding protein to a mammal in need thereof in an amount effective for reducing an IGF-1 and/or IGF-2-induced biological activity.

Antigen binding proteins of the invention include partially human and fully human monoclonal antibodies that inhibit a biological activity of IGF-1 and also inhibit a biological activity of IGF-2. One embodiment is directed to a human monoclonal antibody that at least partially blocks binding of IGF-1

and of IGF-2 to a cell that expresses human IGF-1R. In one embodiment, the antibodies are generated by immunizing a transgenic mouse with an IGF-1R immunogen. In another embodiment, the immunogen is a human IGF-1R polypeptide (e.g., a soluble fragment comprising all or part of the IGF-1R extracellular domain). Hybridoma cell lines derived from such immunized mice, wherein the hybridoma secretes a monoclonal antibody that binds IGF-1R, also are provided herein.

Although human, partially human, or humanized antibodies will be suitable for many applications, particularly those involving administration of the antibody to a human subject, other types of antigen binding proteins will be suitable for certain applications. The non-human antibodies of the invention can be, for example, derived from any antibody-producing animal, such as mouse, rat, rabbit, goat, donkey, or non-human primate (such as monkey (*e.g.*, cynomologous or rhesus monkey) or ape (*e.g.*, chimpanzee)). Non-human antibodies of the invention can be used, for example, in *in vitro* and cell-culture based applications, or any other application where an immune response to the antibody of the invention does not occur, is insignificant, can be prevented, is not a concern, or is desired. In one embodiment, a non-human antibody of the invention is administered to a non-human subject. In another embodiment, the non-human antibody does not elicit an immune response in the non-human subject. In another embodiment, the non-human antibody is from the same species as the non-human subject, *e.g.*, a mouse antibody of the invention is administered to a mouse. An antibody from a particular species can be made by, for example, immunizing an animal of that species with the desired immunogen (*e.g.*, a soluble IGF-1R polypeptide) or using an artificial system for generating antibodies of that species (*e.g.*, a bacterial or phage display-based system for generating antibodies of a particular species), or by converting an antibody from one species into an antibody from another species by replacing, *e.g.*, the constant region of the antibody with a constant region from the other species, or by replacing one or more amino acid residues of the antibody so that it more closely resembles the sequence of an antibody from the other species. In one embodiment, the antibody is a chimeric antibody

comprising amino acid sequences derived from antibodies from two or more different species.

Antigen binding proteins may be prepared by any of a number of conventional techniques. For example, they may be purified from cells that naturally express them (e.g., an antibody can be purified from a hybridoma that produces it), or produced in recombinant expression systems, using any technique known in the art. See, for example, *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Kennet *et al.* (eds.), Plenum Press, New York (1980); and *Antibodies: A Laboratory Manual*, Harlow and Land (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, (1988).

Any expression system known in the art can be used to make the recombinant polypeptides of the invention. In general, host cells are transformed with a recombinant expression vector that comprises DNA encoding a desired polypeptide. Among the host cells that may be employed are prokaryotes, yeast or higher eukaryotic cells. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or bacilli. Higher eukaryotic cells include insect cells and established cell lines of mammalian origin. Examples of suitable mammalian host cell lines include the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman *et al.*, 1981, Cell 23:175), L cells, 293 cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, BHK (ATCC CRL 10) cell lines, and the CVI/EBNA cell line derived from the African green monkey kidney cell line CVI (ATCC CCL 70) as described by McMahan *et al.*, 1991, EMBO J. 10: 2821. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described by Pouwels *et al.* (*Cloning Vectors: A Laboratory Manual*, Elsevier, New York, 1985).

The transformed cells can be cultured under conditions that promote expression of the polypeptide, and the polypeptide recovered by conventional protein purification procedures. One such purification procedure includes the use of affinity chromatography, *e.g.*, over a matrix having all or a portion (*e.g.*, the extracellular domain) of IGF-1R bound thereto. Polypeptides contemplated for use herein include substantially homogeneous recombinant

Figure 16 provides graphs illustrating the ability of certain antibodies to inhibit the growth of 32D hu IGF-1R+IRS-1 cells.

Figure 17 provides graphs illustrating the ability of certain antibodies to inhibit the growth of Balb/C 3T3 hu IGF-1R cells.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides an isolated antigen binding protein comprising either: a. a light chain CDR3 comprising a sequence selected from the group consisting of: i. a light chain CDR3 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR3 sequence selected from the group consisting of the light chain CDR3 sequences of L1-L52 as shown in Figure 6; ii. M X_1 X_2 X_3 X_4 X_5 P X_6 X_7; iii. Q Q X_8 X_9 X_{10} X_{11} P X_{12} T; and iv. Q S Y X_{13} X_{14} X_{15} N X_{16} X_{17} X_{18}; b. a heavy chain CDR3 comprising a sequence selected from the group consisting of: i. a heavy chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence selected from the group consisting of the heavy chain CDR3 sequences of H1-H52 as shown in Figure 9; ii. X_{19} X_{20} X_{21} X_{22} X_{23} X_{24} X_{25} X_{26} X_{27} F D I; iii. X_{28} X_{29} X_{30} X_{31} X_{32} X_{33} X_{34} X_{35} X_{36} X_{37} X_{38} M D V; iv. D S S X_{39}; or c. the light chain CDR3 sequence of (a) and the heavy chain CDR3 sequence of (b); wherein X_1 is a glutamine residue or a glutamate residue, X_2 is an alanine residue, a glycine residue, a threonine residue, or a serine residue, X_3 is a leucine residue, a phenylalanine residue, or a threonine residue, X_4 is glutamine residue, a glutamate residue, or a histidine residue, X_5 is a threonine residue, a methionine residue, a tryptophan residue, or a valine residue, X_6 is a glycine residue, an alanine residue, a valine residue, a leucine residue, an isoleucine residue, a proline residue, a phenylalanine residue, a methionine residue, a tryptophan residue, or a cysteine residue, X_7 is threonine residue, an alanine residue, or a serine residue, X_8 is an arginine residue, a serine residue, a leucine residue, or an alanine residue, X_9 is an asparagine residue, a serine residue, or a histidine residue, X_{10} is an asparagine residue or a serine residue, X_{11} is a tryptophan residue, a valine residue, a tyrosine residue, a proline residue, or a phenylalanine residue, X_{12} is a leucine

mammalian anti- IGF-1R antibody polypeptides substantially free of contaminating endogenous materials.

Antigen binding proteins may be prepared, and screened for desired properties, by any of a number of known techniques. Certain of the techniques involve isolating a nucleic acid encoding a polypeptide chain (or portion thereof) of an antigen binding protein of interest (e.g., an anti-IGF-1R antibody), and manipulating the nucleic acid through recombinant DNA technology. The nucleic acid may be fused to another nucleic acid of interest, or altered (*e.g.*, by mutagenesis or other conventional techniques) to add, delete, or substitute one or more amino acid residues, for example.

In one aspect, the present invention provides antigen-binding fragments of an anti-IGF-1R antibody of the invention. Such fragments can consist entirely of antibody-derived sequences or can comprise additional sequences. Examples of antigen-binding fragments include Fab, F(ab')2, single chain antibodies, diabodies, triabodies, tetrabodies, and domain antibodies. Other examples are provided in Lunde *et al.*, 2002, Biochem. Soc. Trans. 30:500-06.

Single chain antibodies may be formed by linking heavy and light chain variable domain (Fv region) fragments via an amino acid bridge (short peptide linker), resulting in a single polypeptide chain. Such single-chain Fvs (scFvs) have been prepared by fusing DNA encoding a peptide linker between DNAs encoding the two variable domain polypeptides (V_L and V_H). The resulting polypeptides can fold back on themselves to form antigen-binding monomers, or they can form multimers (*e.g.*, dimers, trimers, or tetramers), depending on the length of a flexible linker between the two variable domains (Kortt *et al.*, 1997, Prot. Eng. 10:423; Kortt *et al.*, 2001, Biomol. Eng. 18:95-108). By combining different V_L and V_H-comprising polypeptides, one can form multimeric scFvs that bind to different epitopes (Kriangkum *et al.*, 2001, Biomol. Eng. 18:31-40). Techniques developed for the production of single chain antibodies include those described in U.S. Patent No. 4,946,778; Bird, 1988, Science 242:423; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879; Ward *et al.*, 1989, Nature 334:544, de Graaf *et al.*, 2002, Methods Mol Biol. 178:379-87. Single chain antibodies derived from antibodies

provided herein include, but are not limited to, scFvs comprising the variable domain combinations L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47, L48H48, L49H49, L50H50, L51H51, and L52H52) are encompassed by the present invention.

Antigen binding proteins (e.g., antibodies, antibody fragments, and antibody derivatives) of the invention can comprise any constant region known in the art. The light chain constant region can be, for example, a kappa- or lambda-type light chain constant region, *e.g.*, a human kappa- or lambda-type light chain constant region. The heavy chain constant region can be, for example, an alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant regions, *e.g.*, a human alpha-, delta-, epsilon-, gamma-, or mu-type heavy chain constant region. In one embodiment, the light or heavy chain constant region is a fragment, derivative, variant, or mutein of a naturally occurring constant region.

Techniques are known for deriving an antibody of a different subclass or isotype from an antibody of interest, *i.e.*, subclass switching. Thus, IgG antibodies may be derived from an IgM antibody, for example, and *vice versa.* Such techniques allow the preparation of new antibodies that possess the antigen-binding properties of a given antibody (the parent antibody), but also exhibit biological properties associated with an antibody isotype or subclass different from that of the parent antibody. Recombinant DNA techniques may be employed. Cloned DNA encoding particular antibody polypeptides may be employed in such procedures, *e.g.*, DNA encoding the constant domain of an antibody of the desired isotype. See also Lantto *et al.*, 2002, Methods Mol. Biol.178:303-16.

In one embodiment, an antigen binding protein of the invention comprises the IgG1 heavy chain domain of Figure 13 or a fragment of the IgG1 heavy chain domain of Figure 13. In another embodiment, an antigen

binding protein of the invention comprises the kappa light chain constant chain region of Figure 13 or a fragment of the kappa light chain constant region of Figure 13. In another embodiment, an antigen binding protein of the invention comprises both the IgG1 heavy chain domain, or a fragment thereof, of Figure 13 and the kappa light chain domain, or a fragment thereof, of Figure 13.

Accordingly, the antigen binding proteins of the present invention include those comprising, for example, the variable domain combinations L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47, L48H48, L49H49, L50H50, L51H51, and L52H52, having a desired isotype (for example, IgA, IgG1, IgG2, IgG3, IgG4, IgM, IgE, and IgD) as well as Fab or $F(ab')_2$ fragments thereof. Moreover, if an IgG4 is desired, it may also be desired to introduce a point mutation (CPSCP -> CPPCP) in the hinge region as described in Bloom *et al.*, 1997, Protein Science 6:407, incorporated by reference herein) to alleviate a tendency to form intra-H chain disulfide bonds that can lead to heterogeneity in the IgG4 antibodies.

Moreover, techniques for deriving antigen binding proteins having different properties (*i.e.*, varying affinities for the antigen to which they bind) are also known. One such technique, referred to as chain shuffling, involves displaying immunoglobulin variable domain gene repertoires on the surface of filamentous bacteriophage, often referred to as phage display. Chain shuffling has been used to prepare high affinity antibodies to the hapten 2-phenyloxazol-5-one, as described by Marks *et al.*, 1992, BioTechnology, 10:779.

In particular embodiments, antigen binding proteins of the present invention have a binding affinity (K_a) for IGF-1R of at least 10^6, measured as described in the Examples. In other embodiments, the antigen binding proteins exhibit a K_a of at least 10^7, at least 10^8, at least 10^9, or at least 10^{10}.

In another embodiment, the present invention provides an antigen binding protein that has a low dissociation rate from IGF-1R. In one embodiment, the antigen binding protein has a K_{off} of $1x10^{-4}$ s^{-1} or lower. In another embodiment, the K_{off} is $5x10^{-5}$ s^{-1} or lower. In another embodiment, the K_{off} is substantially the same as an antibody having a combination of light chain and heavy chain variable domain sequences selected from the group of combinations consisting of L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47, L48H48, L49H49, L50H50, L51H51, and L52H52. In another embodiment, the antigen binding protein binds to IGF-1R with substantially the same K_{off} as an antibody that comprises one or more CDRs from an antibody having a combination of light chain and heavy chain variable domain sequences selected from the group of combinations consisting of L1H1, L2H2, L3H3, L4H4, L5H5, L6H6, L7H7, L8H8, L9H9, L10H10, L11H11, L12H12, L13H13, L14H14, L15H15, L16H16, L17H17, L18H18, L19H19, L20H20, L21H21, L22H22, L23H23, L24H24, L25H25, L26H26, L27H27, L28H28, L29H29, L30H30, L31H31, L32H32, L33H33, L34H34, L35H35, L36H36, L37H37, L38H38, L39H39, L40H40, L41H41, L42H42, L43H43, L44H44, L45H45, L46H46, L47H47, L48H48, L49H49, L50H50, L51H51, and L52H52. In another embodiment, the antigen binding protein binds to IGF-1R with substantially the same K_{off} as an antibody that comprises one of the amino acid sequences illustrated in Figures 2 through 9. In another embodiment, the antigen binding protein binds to IGF-1R with substantially the same K_{off} as an antibody that comprises one or more CDRs from an antibody that comprises one of the amino acid sequences illustrated in Figures 2 through 9.

In another aspect, the present invention provides an antigen binding protein that binds to the L2 domain of human IGF-1R. Antigen binding proteins that bind to the L2 domain can be made using any technique known in

the art. For example, such antigen binding proteins can be isolated using the full-length IGF-1R polypeptide (e.g., in a membrane-bound preparation), a soluble extracellular domain fragment of IGF-1R (an example of which is provided in Example 1), or a smaller fragment of the IGF-1R extracellular domain comprising or consisting of the L2 domain (examples of which are provided in Example 10). Antigen binding proteins so isolated can be screened to determine their binding specificity using any method known in the art (an example of which is provided in Example 10).

In another aspect, the present invention provides an antigen binding protein that binds to human IGF-1R expressed on the surface of a cell and, when so bound, inhibits IGF-1R signaling activity in the cell without causing a significant reduction in the amount of IGF-1R on the surface of the cell. Any method for determining or estimating the amount of IGF-1R on the surface and/or in the interior of the cell can be used. In one embodiment, the present invention provides an antigen binding protein that binds to the L2 domain of a human IGF-1R expressed on the surface of a cell and, when so bound, inhibits IGF-1R signaling activity in the cell without significantly increasing the rate of internalization of the IGF-1R from the surface of the cell. In other embodiments, binding of the antigen binding protein to the IGF-1R-expressing cell causes less than about 75%, 50%, 40%, 30%, 20%, 15%, 10%, 5%, 1%, or 0.1% of the cell-surface IGF-1R to be internalized. In another aspect, binding of the antigen binding protein to the IGF-1R-expressing cell causes a gradual reduction in the amount of IGF-1R on the cell surface such that within a few hours of contacting the cell with the antigen binding protein, little or no decrease in cell surface IGF-1R is detected, but, after several days or weeks of exposure of the cell to the antigen binding protein, a marked decrease in cell surface IGF-1R is detected.

In another aspect, the present invention provides an antigen binding protein having a half-life of at least one day *in vitro* or *in vivo* (*e.g.*, when administered to a human subject). In one embodiment, the antigen binding protein has a half-life of at least three days. In another embodiment, the antigen binding protein has a half-life of four days or longer. In another embodiment, the antigen binding protein has a half-life of eight days or longer.

In another embodiment, the antigen binding protein is derivatized or modified such that it has a longer half-life as compared to the underivatized or unmodified antigen binding protein. In another embodiment, the antigen binding protein contains one or more point mutations to increase serum half life, such as described in WO 00/09560, published Feb.24, 2000, incorporated by reference.

The present invention further provides multi-specific antigen binding proteins, for example, bispecific antigen binding protein, *e.g.*, antigen binding protein that bind to two different epitopes of IGF-1R, or to an epitope of IGF-1R and an epitope of another molecule, via two different antigen binding sites or regions. Moreover, bispecific antigen binding protein as disclosed herein can comprise an IGF-1R binding site from one of the herein-described antibodies and a second IGF-1R binding region from another of the herein-described antibodies, including those described herein by reference to other publications. Alternatively, a bispecific antigen binding protein may comprise an antigen binding site from one of the herein described antibodies and a second antigen binding site from another IGF-1R antibody that is known in the art, or from an antibody that is prepared by known methods or the methods described herein.

Numerous methods of preparing bispecific antibodies are known in the art, and discussed in US Patent Application 09/839,632, filed April 20, 2001 (incorporated by reference herein). Such methods include the use of hybrid-hybridomas as described by Milstein *et al.*, 1983, Nature 305:537, and others (U.S. Patent 4,474,893, U.S. Patent 6,106,833), and chemical coupling of antibody fragments (Brennan *et al.*,1985, Science 229:81; Glennie *et al.*,1987, J. Immunol. 139:2367; U.S. Patent 6,010,902). Moreover, bispecific antibodies can be produced via recombinant means, for example by using leucine zipper moieties *(i.e.*, from the Fos and Jun proteins, which preferentially form heterodimers; Kostelny *et al.*, 1992, J. Immnol. 148:1547) or other lock and key interactive domain structures as described in U.S. Patent 5,582,996. Additional useful techniques include those described in Kortt *et al.*, 1997, *supra*; U.S. Patent 5,959,083; and U.S. Patent 5,807,706.

In another aspect, the antigen binding protein of the present invention comprises a derivative of an antibody. The derivatized antibody can comprise any molecule or substance that imparts a desired property to the antibody, such as increased half-life in a particular use. The derivatized antibody can comprise, for example, a detectable (or labeling) moiety (*e.g.*, a radioactive, colorimetric, antigenic or enzymatic molecule, a detecable bead (such as a magnetic or electrodense (*e.g.*, gold) bead), or a molecule that binds to another molecule (*e.g.*, biotin or streptavidin)), a therapeutic or diagnostic moiety (*e.g.*, a radioactive, cytotoxic, or pharmaceutically active moiety), or a molecule that increases the suitability of the antibody for a particular use (*e.g.*, administration to a subject, such as a human subject, or other *in vivo* or *in vitro* uses). Examples of molecules that can be used to derivatize an antibody include albumin (*e.g.*, human serum albumin) and polyethylene glycol (PEG). Albumin-linked and PEGylated derivatives of antibodies can be prepared using techniques well known in the art. In one embodiment, the antibody is conjugated or otherwise linked to transthyretin (TTR) or a TTR variant. The TTR or TTR variant can be chemically modified with, for example, a chemical selected from the group consisting of dextran, poly(n-vinyl pyurrolidone), polyethylene glycols, propropylene glycol homopolymers, polypropylene oxide/ethylene oxide co-polymers, polyoxyethylated polyols and polyvinyl alcohols. US Pat. App. No. 20030195154.

In another aspect, the present invention provides methods of screening for a molecule that binds to IGF-1R using the antigen binding proteins of the present invention. Any suitable screening technique can be used. In one embodiment, an IGF-1R molecule, or a fragment thereof to which an antigen binding protein of the present invention binds, is contacted with the antigen binding protein of the invention and with another molecule, wherein the other molecule binds to IGF-1R if it reduces the binding of the antigen binding protein to IGF-1R. Binding of the antigen binding protein can be detected using any suitable method, *e.g.*, an ELISA. Detection of binding of the antigen binding protein to IGF-1R can be simplified by detectably labeling the antigen binding protein, as discussed above. In another embodiment, the IGF-

1R-binding molecule is further analyzed to determine whether it inhibits IGF-1R, IGF-1, and/or IGF-2-mediated signaling.

Nucleic acids

In one aspect, the present invention provides isolated nucleic acid molecules. The nucleic acids comprise, for example, polynucleotides that encode all or part of an antigen binding protein , for example, one or both chains of an antibody of the invention, or a fragment, derivative, mutein, or variant thereof, polynucleotides sufficient for use as hybridization probes, PCR primers or sequencing primers for identifying, analyzing, mutating or amplifying a polynucleotide encoding a polypeptide, anti-sense nucleic acids for inhibiting expression of a polynucleotide, and complementary sequences of the foregoing. The nucleic acids can be any length. They can be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 1,500, 3,000, 5,000 or more nucleotides in length, and/or can comprise one or more additional sequences, for example, regulatory sequences, and/or be part of a larger nucleic acid, for example, a vector. The nucleic acids can be single-stranded or double-stranded and can comprise RNA and/or DNA nucleotides, and artificial variants thereof (*e.g.*, peptide nucleic acids).

Nucleic acids encoding antibody polypeptides (*e.g.*, heavy or light chain, variable domain only, or full length) may be isolated from B-cells of mice that have been immunized with IGF-1R. The nucleic acid may be isolated by conventional procedures such as polymerase chain reaction (PCR).

Figure 1 provides nucleic acid sequences encoding the variable regions of the heavy and light chain variable regions shown in Figures 2 and 3. The skilled artisan will appreciate that, due to the degeneracy of the genetic code, each of the polypeptide sequences in Figures 2 through 9 also is encoded by a large number of other nucleic acid sequences. The present invention provides each degenerate nucleotide sequence encoding each antigen binding protein of the invention.

The invention further provides nucleic acids that hybridize to other nucleic acids (*e.g.*, nucleic acids comprising a nucleotide sequence of Figure

1) under particular hybridization conditions. Methods for hybridizing nucleic acids are well-known in the art. See, *e.g.*, Current Protocols in Molecular Biology, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. As defined herein, a moderately stringent hybridization condition uses a prewashing solution containing 5X sodium chloride/sodium citrate (SSC), 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization buffer of about 50% formamide, 6X SSC, and a hybridization temperature of 55° C (or other similar hybridization solutions, such as one containing about 50% formamide, with a hybridization temperature of 42° C), and washing conditions of 60° C, in 0.5X SSC, 0.1% SDS. A stringent hybridization condition hybridizes in 6X SSC at 45° C, followed by one or more washes in 0.1X SSC, 0.2% SDS at 68° C. Furthermore, one of skill in the art can manipulate the hybridization and/or washing conditions to increase or decrease the stringency of hybridization such that nucleic acids comprising nucleotide sequences that are at least 65, 70, 75, 80, 85, 90, 95, 98 or 99% identical to each other typically remain hybridized to each other. The basic parameters affecting the choice of hybridization conditions and guidance for devising suitable conditions are set forth by, for example, Sambrook, Fritsch, and Maniatis (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., chapters 9 and 11; and Current Protocols in Molecular Biology, 1995, Ausubel *et al.*, eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4), and can be readily determined by those having ordinary skill in the art based on, for example, the length and/or base composition of the DNA.

Changes can be introduced by mutation into a nucleic acid, thereby leading to changes in the amino acid sequence of a polypeptide (*e.g.*, an antigen binding protein) that it encodes. Mutations can be introduced using any technique known in the art. In one embodiment, one or more particular amino acid residues are changed using, for example, a site-directed mutagenesis protocol. In another embodiment, one or more randomly selected residues is changed using, for example, a random mutagenesis protocol. However it is made, a mutant polypeptide can be expressed and screened for a desired property (*e.g.*, binding to IGF-1R or blocking the binding of IGF-1 and/or IGF-2 to IGF-1R).

Mutations can be introduced into a nucleic acid without significantly altering the biological activity of a polypeptide that it encodes. For example, one can make nucleotide substitutions leading to amino acid substitutions at non-essential amino acid residues. In one embodiment, a nucleotide sequence provided in Figure 1, or a desired fragment, variant, or derivative thereof, is mutated such that it encodes an amino acid sequence comprising one or more deletions or substitutions of amino acid residues that are shown in Figures 2 through 9 to be residues where two or more sequences differ. In another embodiment, the mutagenesis inserts an amino acid adjacent to one or more amino acid residues shown in Figures 2 through 9 to be residues where two or more sequences differ. Alternatively, one or more mutations can be introduced into a nucleic acid that selectively change the biological activity (*e.g.*, binding of IGF-1R, inhibiting IGF-1 and/or IGF-2, *etc.*) of a polypeptide that it encodes. For example, the mutation can quantitatively or qualitatively change the biological activity. Examples of quantitative changes include increasing, reducing or eliminating the activity. Examples of qualitative changes include changing the antigen specificity of an antigen binding protein.

In another aspect, the present invention provides nucleic acid molecules that are suitable for use as primers or hybridization probes for the detection of nucleic acid sequences of the invention. A nucleic acid molecule of the invention can comprise only a portion of a nucleic acid sequence encoding a full-length polypeptide of the invention, for example, a fragment that can be used as a probe or primer or a fragment encoding an active portion (*e.g.*, an IGF-1R binding portion) of a polypeptide of the invention.

Probes based on the sequence of a nucleic acid of the invention can be used to detect the nucleic acid or similar nucleic acids, for example, transcripts encoding a polypeptide of the invention. The probe can comprise a label group, *e.g.*, a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used to identify a cell that expresses the polypeptide.

In another aspect, the present invention provides vectors comprising a nucleic acid encoding a polypeptide of the invention or a portion thereof. Examples of vectors include, but are not limited to, plasmids, viral vectors,

residue, a tyrosine residue, or an isoleucine residue, X_{13} is an aspartate residue or a glutamine residue, X_{14} is a serine residue or a proline residue, X_{15} is a serine residue, a tyrosine residue, an aspartate residue, or an alanine residue, X_{16} is a glutamine residue, an arginine residue, a valine residue, or a tryptophan residue, X_{17} is an arginine residue, a valine residue, an isoleucine residue, or no residue, X_{18} is a valine residue or no residue, X_{19} is a glutamate residue or no residue, X_{20} is a tyrosine residue, a glycine residue, a serine residue, or no residue, X_{21} is a serine residue, an asparagine residue, a tryptophan residue, a glutamate residue, as aspartate residue, or no residue, X_{22} is a serine residue, an aspartate residue, a tryptophan residue, an alanine residue, an arginine residue, a threonine residue, a glutamine residue, a leucine residue, a glutamate residue, or no residue, X_{23} is a serine residue, a glycine residue, an asparagine residue, a threonine residue, a tryptophan residue, a valine residue, an alanine residue, or an isoleucine residue, X_{24} is an arginine residue, a glutamine residue, a tyrosine residue, a valine residue, an alanine residue, a glycine residue, a serine residue, a phenylalanine residue, or a tryptophan residue, X_{25} is an asparagine residue, a leucine residue, an aspartate residue, a threonine residue, a tryptophan residue, a tyrosine residue, a valine residue, an alanine residue, or a histidine residue, X_{26} is an aspartate residue, a serine residue, an asparagine residue, or a glutamine residue, X_{27} is an alanine residue or a proline residue, X_{28} is an alanine residue or no residue, X_{29} is a glutamate residue, a tyrosine residue, a glycine residue, or no residue, X_{30} is an arginine residue, a serine residue, or no residue, X_{31} is a glycine residue, an aspartate residue, a valine residue, a serine residue, or no residue, X_{32} is a serine residue, an aspartate residue, a glycine residue, or no residue, X_{33} is a phenylalanine residue, an aspartate residue, a tyrosine residue, a glycine residue, a serine residue, a histidine residue, a tryptophan residue, or no residue, X_{34} is a tryptophan residue, an aspartate residue, a tyrosine residue, a serine residue, or no residue, X_{35} is an aspartate residue, a glutamate residue, an arginine residue, a serine residue, a glycine residue, a tyrosine residue, or a tryptophan residue, X_{36} is a tyrosine residue, a lysine residue, an isoleucine residue, a leucine residue or a phenylalanine residue, X_{37} is a tyrosine residue, a serine residue, a phenylalanine residue, an aspartate residue, or a glycine

non-episomal mammalian vectors and expression vectors, for example, recombinant expression vectors.

The recombinant expression vectors of the invention can comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell. The recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operably linked to the nucleic acid sequence to be expressed. Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cells (*e.g.*, SV40 early gene enhancer, Rous sarcoma virus promoter and cytomegalovirus promoter), those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences, see Voss *et al.*, 1986, Trends Biochem. Sci. 11:287, Maniatis *et al.*, 1987, Science 236:1237, incorporated by reference herein in their entireties), and those that direct inducible expression of a nucleotide sequence in response to particular treatment or condition (*e.g.*, the metallothionin promoter in mammalian cells and the tet-responsive and/or streptomycin responsive promoter in both prokaryotic and eukaryotic systems (see *id.*). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, *etc.* The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein.

In another aspect, the present invention provides host cells into which a recombinant expression vector of the invention has been introduced. A host cell can be any prokaryotic cell (for example, *E. coli*) or eukaryotic cell (for example, yeast, insect, or mammalian cells (*e.g.*, CHO cells)). Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, for resistance to

antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die), among other methods.

Indications

In one aspect, the present invention provides methods of treating a subject. The method can, for example, have a generally salubrious effect on the subject, *e.g.*, it can increase the subject's expected longevity. Alternatively, the method can, for example, treat, prevent, cure, relieve, or ameliorate ("treat") a disease, disorder, condition, or illness ("a condition"). Among the conditions to be treated in accordance with the present invention are conditions characterized by inappropriate expression or activity of IGF-1, IGF-2, and/or IGF-1R. In some such conditions, the expression or activity level is too high, and the treatment comprises administering an IGF-1R antagonist as described herein. In other such conditions, the expression or activity level is too low, and the treatment comprises administering an IGF-1R agonist as described herein.

One example of a type of condition that can be treated using the methods and compositions of the present invention is a condition that involves cell growth, for example, a cancerous condition. Thus, in one embodiment, the present invention provides compositions and methods for treating a cancerous condition. The cancerous condition can be any cancerous condition that can be treated using the compositions comprised herein, for example, IGF-1R antagonizing antigen binding proteins such as anti-IGF-1R antibodies, antibody fragments, or antibody derivatives. Examples of cancerous conditions include, for example, Acute Lymphoblastic Leukemia, Adrenocortical Carcinoma, AIDS-Related Cancers, AIDS-Related Lymphoma, Anal Cancer, Childhood Cerebellar Astrocytoma, Childhood Cerebral Astrocytoma, Basal Cell Carcinoma, Extrahepatic Bile Duct Cancer, Bladder Cancer, Osteosarcoma/Malignant Fibrous Histiocytoma Bone Cancer,

Brain Tumors (e.g., Brain Stem Glioma, Cerebellar Astrocytoma, Cerebral Astrocytoma/Malignant Glioma, Ependymoma, Medulloblastoma, Supratentorial Primitive Neuroectodermal Tumors, Visual Pathway and Hypothalamic Glioma), Breast Cancer, Bronchial Adenomas/Carcinoids, Burkitt's Lymphoma, Carcinoid Tumor, Gastrointestinal Carcinoid Tumor, Carcinoma of Unknown Primary, Primary Central Nervous System, Cerebellar Astrocytoma, Cerebral Astrocytoma/Malignant Glioma, Cervical Cancer, Childhood Cancers, Chronic Lymphocytic Leukemia, Chronic Myelogenous Leukemia, Chronic Myeloproliferative Disorders, Colon Cancer, Colorectal Cancer, Cutaneous T-Cell Lymphoma, Endometrial Cancer, Ependymoma, Esophageal Cancer, Ewing's Family of Tumors, Extracranial Germ Cell Tumor, Extragonadal Germ Cell Tumor, Extrahepatic Bile Duct Cancer, Intraocular Melanoma Eye Cancer, Retinoblastoma Eye Cancer, Gallbladder Cancer, Gastric (Stomach) Cancer, Gastrointestinal Carcinoid Tumor, Germ Cell Tumors (e.g., Extracranial, Extragonadal, and Ovarian), Gestational Trophoblastic Tumor, Glioma (e.g., Adult, Childhood Brain Stem, Childhood Cerebral Astrocytoma, Childhood Visual Pathway and Hypothalamic), Hairy Cell Leukemia, Head and Neck Cancer, Hepatocellular (Liver) Cancer, Hodgkin's Lymphoma, Hypopharyngeal Cancer, Hypothalamic and Visual Pathway Glioma, Intraocular Melanoma, Islet Cell Carcinoma (Endocrine Pancreas), Kaposi's Sarcoma, Kidney (Renal Cell) Cancer, Laryngeal Cancer, Leukemia (e.g., Acute Lymphoblastic, Acute Myeloid, Chronic Lymphocytic, Chronic Myelogenous, and Hairy Cell), Lip and Oral Cavity Cancer, Liver Cancer, Non-Small Cell Lung Cancer, Small Cell Lung Cancer, Lymphoma (e.g., AIDS-Related, Burkitt's, Cutaneous T-Cell, Hodgkin's, Non-Hodgkin's, and Primary Central Nervous System), Waldenström's Macroglobulinemia, Malignant Fibrous Histiocytoma of Bone/Osteosarcoma, Medulloblastoma, Melanoma, Intraocular (Eye) Melanoma, Merkel Cell Carcinoma, Mesothelioma, Metastatic Squamous Neck Cancer with Occult Primary, Multiple Endocrine Neoplasia Syndrome, Multiple Myeloma/Plasma Cell Neoplasm, Mycosis Fungoides, Myelodysplastic Syndromes, Myelodysplastic/Myeloproliferative Diseases, Myelogenous Leukemia, Chronic Myeloid Leukemia, Multiple Myeloma, Chronic Myeloproliferative

Disorders, Nasal Cavity and Paranasal Sinus Cancer, Nasopharyngeal Cancer, Neuroblastoma, Oral Cancer, Oropharyngeal Cancer, Osteosarcoma/Malignant Fibrous Histiocytoma of Bone, Ovarian Cancer, Ovarian Epithelial Cancer, Ovarian Germ Cell Tumor, Ovarian Low Malignant Potential Tumor, Pancreatic Cancer, Islet Cell Pancreatic Cancer, Paranasal Sinus and Nasal Cavity Cancer, Parathyroid Cancer, Penile Cancer, Pheochromocytoma, Pineoblastoma, Pituitary Tumor, Plasma Cell Neoplasm/Multiple Myeloma, Pleuropulmonary Blastoma, Primary Central Nervous System Lymphoma, Prostate Cancer, Rectal Cancer, Renal Cell (Kidney) Cancer, Renal Pelvis and Ureter Transitional Cell Cancer, Retinoblastoma, Rhabdomyosarcoma, Salivary Gland Cancer, Soft Tissue Sarcoma, Uterine Sarcoma, Sezary Syndrome, non-Melanoma Skin Cancer, Merkel Cell Skin Carcinoma, Small Intestine Cancer, Soft Tissue Sarcoma, Squamous Cell Carcinoma, Cutaneous T-Cell Lymphoma, Testicular Cancer, Thymoma, Thymic Carcinoma, Thyroid Cancer, Gestational Trophoblastic Tumor, Carcinoma of Unknown Primary Site, Cancer of Unknown Primary Site, Urethral Cancer, Endometrial Uterine Cancer, Uterine Sarcoma, Vaginal Cancer, Visual Pathway and Hypothalamic Glioma, Vulvar Cancer, Waldenström's Macroglobulinemia, and Wilms' Tumor.

Four different groups have studied a total of 425 breast cancers, mostly ductal in origin, and 48 normal tissues or benign specimens by radioimmunoassay ("RIA") or immunohistochemistry ("IHC") (Papa *et al.*, 1993, Cancer Research 53: 3736-40, Happerfield *et al.*, 1997, Journal of Pathology 183: 412-17; Ellis *et al.*, 1998, Breast Cancer Research & Treatment 52: 175-84, Lee *et al.*, 1998, Breast Cancer Research & Treatment 47: 295-302, Schnarr *et al.*, 2000, International Journal of Cancer 89: 506-13). These studies suggest that elevated IGF-1R expression, on the order of 5-10 fold, is associated with favorable prognosis and biomarkers (ER+ PR+), suggesting that estrogen and IGF cooperate in the maintenance or progression of well differentiated tumor. Similarly, estrogen has been shown to be essential for the growth and survival of the ER+ MCF-7 breast cancer cell line, and in this context IGF-1R is up-regulated by estrogen treatment (reviewed in Ellis *et al.*, 1998, Breast Cancer Research & Treatment 52: 175-84). Thus, in

one embodiment, the present invention provides a method of treating breast cancer in a subject in need of such treatment, comprising administering to the subject an effective amount of an IGF-1R antagonist as described herein. In another embodiment, the method further comprises administering a hormone inhibitor, *e.g.*, an estrogen inhibitor.

A retrospective IGF-1R IHC analysis has been reported for a collection of 12 colonic adenomas, 36 primary colorectal adenocarcinomas and 27 corresponding metastases, and 34 adjacent normal tissues (Hakam *et al.*, 1999, Human Pathology. 30: 1128-33). The frequency of moderate to strong IHC staining appeared to dramatically increase with higher stage and tumor grade (0% normal vs. 93 % metastases). The results are consistent with RNA analysis by RNAse protection assay ("RPA") (Freier *et al.*, 1999, Gut 44: 704-08). Thus, in one embodiment, the present invention provides a method of treating colon cancer in a subject in need of such treatment, comprising administering to the subject an effective amount of an IGF-1R antagonist as described herein.

High plasma IGF-1 and reduced IGFbp3 in men 40-80 years old is associated with increased prostate cancer risk (Chan *et al.*, 1998, Science 279: 563-6). High IGF-1 is associated with a risk of other cancers including breast (Hankinson *et al.*, 1998, Lancet 351: 1393-96), colon (Ma *et al.*, 1999, Journal of the National Cancer Institute 91: 620-25) and lung (Yu *et al.*, 1999, Journal of the National Cancer Institute 91: 151-56). In transgenic mouse models, tumor incidence is increased by IGF-1 overexpression in diverse locations (Bol *et al.*, 1997, Oncogene 14: 1725-34; DiGiovanni *et al.*, 2000, Cancer Research 60: 1561-70; DiGiovanni *et al.*, 2000, Proceedings of the National Academy of Sciences of the United States of America 97: 3455-60, Hadsell *et al.*, 2000, Oncogene 19: 889-98). These mouse studies point to a role for both serum and stromal produced IGF-1. Thus, in one embodiment, the present invention provides a method of treating a subject in need of such treatment, comprising administering to the subject an effective amount of an antagonist of IGF-1R as described herein, wherein the antagonist inhibits the activation of IGF-1R by IGF-1. In another embodiment, the subject has cancer. In

another embodiment, the subject has a tumor. In another embodiment, the cancer is prostate, breast, colon or lung cancer.

It has been observed that bone is the major source of IGF-1 in the body. Thus, in one aspect, the present invention provides compositions and methods for inhibiting IGF-1R in a bone of a subject. In one embodiment, an IGF-1R inhibitor of the present invention is administered to a subject that has, or is at risk for developing, a tumor in a bone. The tumor can be, for example, a primary tumor or a metastatic tumor. The treatment optionally further comprises administering to the subject one or more additional therapeutic and/or palliative treatments, for example, an anti-tumor treatment (*e.g.*, chemotherapy, radiation therapy, or anti-hormone therapy) or a treatment that inhibits bone turnover (*e.g.*, denosumab (Amgen Inc., Thousand Oaks, CA)).

IGF-2 is overexpressed in a variety of tumors and stromal tissues. IGF-2 levels appear especially high (as much as 40 fold) in primary liver cancers (Cariani *et al.*, 1988, Cancer Research 48: 6844-49) and adenocarcinoma of the colon (Freier *et al.*, 1999, Gut 44: 704-08). Many of the overgrowth disorders are associated with an increased incidence of childhood tumors. Five to ten percent of individuals with either the prenatal growth disorder Beckwith-Weidmann Syndrome (BWS) or hemihyperplasia develop tumors such as nephroblastoma, adrenal carcinoma, and neuroblastoma (reviewed by Morison *et al.*, 1998, Molecular Medicine Today 4: 110-05). The tumor-predisposing factor in these children appears to be the mosaic loss of maternal IGF-2 gene imprinting, or duplication of the paternal chromosomal arm (11p) that carries IGF-2. Both alterations would increase the level of IGF-2 expression. IGF-2 overexpression as a result of mosaic uniparental disomy or loss of IGF-2 imprinting has also been detected in Wilms tumors. Growth disorders are not observed in these children even though the IGF-2 gene alterations also occur in some normal tissues, perhaps reflecting the tissue distribution of the affected cells. Imprinting of the maternal IGF-2 gene also occurs in mice, and the effects of IGF-2 overexpression are consistent with the human situation (Cariani *et al.*, 1991, Journal of Hepatology 13: 220-26, Schirmacher *et al.*, 1992, Cancer Research 52: 2549-56; Harris *et al.*, 1998, Oncogene 16: 203-09). The incidence of

tumors and organomegaly increases in mice that transgenically express excess IGF-2 (Christofori *et al.*, 1994, Nature 369: 414-18, Ward *et al.*, 1994, Proceedings of the National Academy of Sciences of the United States of America 91: 10365-9, Wolf *et al.*, 1994, Endocrinology 135: 1877-86, Bates *et al.*, 1995, British Journal of Cancer 72: 1189-93, Hassan *et al.*, 2000, Cancer Research 60: 1070-76). Local IGF-2 overexpression increases the spontaneous appearance of prostate, mammary, intestinal, liver and epidermal tumors. Plasma specific expression using liver promoters elevate hepatocellular carcinomas and lymphoma. Thus, in one embodiment, the present invention provides a method of treating a subject in need of such treatment, comprising administering to the subject an effective amount of an antagonist of IGF-1R as described herein, wherein the antagonist inhibits the activation of IGF-1R by IGF-2. In another embodiment, the subject has cancer. In another embodiment, the subject has a tumor. In another embodiment, the subject has liver cancer, adenocarcinoma of the colon, Beckwith-Weidmann Syndrome, hemihyperplasia, nephroblastoma, adrenal carcinoma, neuroblastoma, mosaic loss of maternal IGF-2 gene imprinting, duplication of the paternal chromosomal arm (11p), increased IGF-2 expression, a tumor (*e.g.*, a prostate, mammary, intestinal, liver, epidermal, or Wilms tumor), organomegaly, hepatocellular carcinoma, or lymphoma.

In another aspect, the invention provides methods of preventing or inhibiting a cancer from spreading to another part of the body, or of treating a cancer that has spread to another part of the body. In one embodiment, the cancer has spread to a regional lymph node. In another embodiment, the cancer is metastatic. The primary tumor can be any kind of tumor, for example, an adenocarcinoma tumor (*e.g.*, a prostate adenocarcinoma tumor, a breast carcinoma tumor, or a renal cell carcinoma tumor), a non-small cell or small cell lung cancer tumor, a thyroid cancer tumor, *etc.* The site of the metastatic tumor can be anywhere in the body. It can be, for example, in bone, the lymph system, lung, brain, eye, skin, pancrease, or liver. In one particular embodiment, a subject having a tumor disease is treated with an effective amount of an IGF-1R inhibiting composition of the present invention such that the primary tumor is prevented from metastasizing. In another

particular embodiment, a subject having a primary tumor is treated with an effective amount of an IGF-1R inhibiting composition of the present invention such that the primary tumor is inhibited from metastasizing. In another particular embodiment, a subject having a metastatic tumor is treated with an effective amount of an IGF-1R inhibiting composition of the present invention such that growth or spreading of the secondary tumor is inhibited. In another particular embodiment, a subject having a metastatic tumor is treated with an effective amount of an IGF-1R inhibiting composition of the present invention such that the secondary tumor is reduced in size. In a more particular embodiment, the primary tumor is an adenocarcinoma tumor, a non-small cell lung tumor, a small cell lung tumor, or a thyroid cancer. In another more particular embodiment, the metastatic tumor is in a bone. In another more particular embodiment, a metastatic tumor is prevented or inhibited from forming in a bone. In another more particularly defined embodiment, the method comprises treating the subject with an IGF-1R inhibiting composition of the present invention and one or more other treatments (e.g., a treatment that kills or inhibits the growth of cancer cells, such as radiation, hormonal therapy, or chemotherapy, or a treatment that inhibits the turnover of bone, such as denosumab), non-limiting examples of which are provided herein. The one or more other treatments can include, for example the standard of care for the subject's particular condition and/or palliative care.

Without being bound to any particular theory, tumor cells appear to depend on the PI3 Kinase/Akt signaling pathway to resist the apoptosis-inducing activity of chemotherapeutics, radiation, and anti-hormone therapy. Thus, in one embodiment, the present invention provides methods of treating a subject in need of such treatment comprising administering to the subject an IGF-1R antagonist of the present invention and a chemotherapeutic, radiation, and/or an anti-hormone therapy. This concept has been validated experimentally in cell culture models and rodent tumor models by antisense and dominant negative mutations (reviewed by Baserga *et al.*, 1997, Biochimica et Biophysica Acta 1332: F105-26, Baserga, 2000, Oncogene 19: 5574-81). In one embodiment, the chemotherapeutic agents is selected from the group consisting of mitotic inhibitors, alkylating agents, anti-metabolites,

intercalating antibiotics, growth factor inhibitors, cell cycle inhibitors, enzymes, topoisomerase inhibitors, anti-survival agents, biological response modifiers, anti-hormones, e.g. anti-androgens, and anti-angiogenesis agents.

One example of a chemotherapeutic agent that can be administered in combination with an IGF-1 receptor inhibitor of the invention is CPT-11. CPT-11 (Irinotecan hydorchloride trihydrate) is a semi synthetic, water soluble derivative of camptothecin, a plant alkaloid. CPT-11 and an associated metabolite called SN38 inhibit topoisomerase 1 (TOPO1). This enzyme introduces reversible single-strand breaks in DNA that allow unwinding and permit DNA replication to proceed. Inhibition of TOPO1 prevents religation of single-strand breaks after DNA replication resulting in greatly increased chromosomal fragmentation. This DNA damage promotes cell death by apoptosis through the action of p53 and other systems that monitor genome integrity. The cytotoxic effect of CPT-11 is generally limited to cells that are replicating DNA (S-Phase). Quiescent cells are largely unaffected.

In another embodiment, the present invention provides treating a subject in need thereof with an effective amount of an IGF-1R antagonist of the present invention and with an effective amount of an apoptosis-inducing agent.

In another embodiment, an anti-angiogenesis agent, such as an MMP-2 (matrix-metalloproteinase 2) inhibitor, an MMP-9 (matrix-metalloproteinase 9) inhibitor, and/or a COX-II (cyclooxygenase II) inhibitor, is used in conjunction with a compound of the invention. Examples of useful COX-II inhibitors include CELEBREX™ (alecoxib), BEXTRA™ (valdecoxib), and VIOXX™ (rofecoxib). Examples of useful matrix metalloproteinase inhibitors are described in WO 96/33172 (published Oct. 24, 1996), WO 96/27583 (published Mar. 7, 1996), European Patent Application No. 97304971.1 (filed Jul. 8, 1997), European Patent Application No. 99308617.2 (filed Oct. 29, 1999), WO 98/07697 (published Feb. 26, 1998), WO 98/03516 (published Jan. 29, 1998), WO 98/34918 (published Aug. 13, 1998), WO 98/34915 (published Aug. 13, 1998), WO 98/33768 (published Aug. 6, 1998), WO 98/30566 (published Jul. 16, 1998), European Patent Publication 606,046 (published Jul. 13, 1994), European Patent Publication 931,788 (published Jul.

28, 1999), WO 90/05719 (published May 31, 1990), WO 99/52910 (published Oct. 21, 1999), WO 99/52889 (published Oct. 21, 1999), WO 99/29667 (published Jun. 17, 1999), PCT International Application No. PCT/IB98/01113 (filed Jul. 21, 1998), European Patent Application No. 99302232.1 (filed Mar. 25, 1999), Great Britain patent application number 9912961.1 (filed Jun. 3, 1999), U.S. Provisional Application No. 60/148,464 (filed Aug. 12, 1999), U.S. Pat. No. 5,863,949 (issued Jan. 26, 1999), U.S. Pat. No. 5,861,510 (issued Jan. 19, 1999), and European Patent Publication 780,386 (published Jun. 25, 1997), all of which are incorporated herein in their entireties by reference. In one embodiment, the MMP inhibitor is one that does not demonstrate arthralgia. In another embodiment, the MMP inhibitor selectively inhibits MMP-2 and/or MMP-9 relative to other matrix-metalloproteinases (*i.e.*, MMP-1, MMP-3, MMP-4, MMP-5, MMP-6, MMP-7, MMP-8, MMP-10, MMP-11, MMP-12, and MMP-13). Some specific examples of MMP inhibitors useful in the present invention are AG-3340, RO 32-3555, RS 13-0830, and the compounds recited in the following list: 3-[[4-(4-fluoro-phenoxy)-benzene- sulfonyl]-(1-hydroxycarbamoyl-cyclopentyl)-amino]-propionic acid; 3-exo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-bicyclo[3.2.1]o- ctane-3-carboxylic acid hydroxyamide; (2R, 3R) 1-[4-(2-chloro-4-fluoro-ben- zyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-piperidine-2-carboxylic acid hydroxyamide; 4-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-py- ran-4-carboxylic acid hydroxyamide; 3-[[4-(4-fluoro-phenoxy)-benzenesulfon- yl]-(1-hydroxycarbamoyl-cyclobutyl)-amino]-propionic acid; 4-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-tetrahydro-pyran-4-carboxyl-ic acid hydroxyamide; (R) 3-[4-(4-chloro-phenoxy)-benzenesulfonylamino]-te- trahydro-pyran-3-carboxylic acid hydroxyamide; (2R, 3R) 1-[4-(4-fluoro-2-methyl-benzyloxy)-benzenesulfonyl]-3-hydroxy-3-methyl-pi- peridine-2-carboxylic acid hydroxyamide; 3-[[4-(4-fluoro-phenoxy)-benzenes- ulfonyl]-(1-hydroxycarbamoyl-1-methyl-ethyl)-amino]-propionic acid; 3-[[4-(4-fluoro-phenoxy)-benzenesulfonyl]-(4-hydroxycarbamoyl-tetrahydro-- pyran-4-yl)-amino]-propionic acid; 3-exo-3-[4-(4-chloro-phenoxy)-benzenesu-lfonylamino]-8-oxa-icyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; 3-

residue, X_{38} is a glycine residue, an asparagine residue, or a tyrosine residue, X_{39} is a valine residue, a glycine residue, or a serine residue, and said antigen binding protein binds specifically to human IGF-1R. In one embodiment, the isolated antigen binding protein comprises an amino acid sequence selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of six amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises an amino acid sequence selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of five amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a

endo-3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-8-oxa-icyclo[3.2.1]octane-3-carboxylic acid hydroxyamide; and (R) 3-[4-(4-fluoro-phenoxy)-benzenesulfonylamino]-tetrahydro-furan-3-carboxylic acid hydroxyamide; and pharmaceutically acceptable salts, solvates, derivatives, and other preparations of the compounds.

Sporadic mutations that inactivate the PETN gene product occur relatively frequently in most human cancers (Yamada *et al.*, 2001, J Cell Sci 114:2375-82, Hill *et al.*, 2002, Pharmacol Therapeut 93:243-51). Loss of PTEN causes the Akt phosphorylated state to persist through loss of the ability to down-regulate stimulatory signals originating from IGF-1R and other sources. The status of the p53 tumor suppressor also influences the activity of the IGF-1R signaling system. In the ground state, the basal or constitutive transcription of IGF-1R is repressed by p53 via an indirect mechanism. Activation of Akt promotes the phosphorylation of mdm2, which then binds the p53 tumor suppressor and promotes its degradation (Mayo *et al.*, 2002, TIBS 27:462-67), resulting in increased IGF-1R expression. A similar outcome is observed when p53 is inactivated by mutation. When transiently expressed in Saos-2 (a human osteosarcoma cell line) and RD (a rhabdomyosarcoma cell line), wild-type p53 is able to suppress the activity of a cotransfected IGF-1R promoter construct, whereas tumor-derived, mutant versions of p53 have no effect. It has been proposed that the increased level of IGF-1R promotes the resistance to apoptosis associated with p53 loss in malignant cells (Werner *et al.*, 2000, Cell Mol Life Sci 57:932-42). Thus, in one embodiment, the present invention provides a method of treating a cancerous condition in a subject in need of such treatment comprising administering to the subject an effective amount of an IGF-1R antagonist as described herein, wherein the cancerous condition is characterized by cells that have a reduced expression or activity of p53.

The WT1 (Wilms kidney tumor suppressor 1 protein) also has been shown to bind and repress the IGF-1R promoter. Thus, in one embodiment, the present invention provides a method of treating a cancerous condition in a subject in need of such treatment comprising administering to the subject an effective amount of an IGF-1R antagonist as described herein wherein the

cancerous condition is characterized by a reduced expression or activity of WT1.

The proliferation of normal fibroblasts has been shown to require, under defined culture conditions, the combined action of IGF and a stromal growth factor (*e.g.* PDGF, EGF) to ramp-up Ras/Raf/Map Kinase and promote cell cycle entry (the G0 to G1 transition). Fibroblasts derived from IGF-1R (-/-) mice do not respond to growth factor alone, or most oncogenes (*e.g.* oncogenic Ras) that activate the Ras/Raf/Map Kinase pathway. Thus, in one embodiment, the present invention provides a method of treating a subject in need of such treatment comprising administering to the subject an IGF-1R antagonist as described herein and an agent that targets a growth factor and/or a growth factor receptor, such as a growth factor receptor tyrosine kinase, *e.g.*, the EGFR, HER-2, bcr-abl, VEGFR, Kit, raf, mTOR, CDK1/2, VEGFR2, PKCβ, Mek, and/or KDR. Examples of molecules that target such growth factors and/or receptors include panitumumab (Abgenix, Fremont, CA/Amgen, Thousand Oaks, CA), HERCEPTIN™ (Genentech, South San Francisco, CA), GLEEVEC™ (Novartis, East Hanover, NJ), IRESSA™ (AstraZeneca, Wilmington, DE), ERBITUX™, (ImClone, New York, NY), AVASTIN™, (Genentech), PTK787 (Novartis), SU11248 (Pfizer, New York, NY), TARCEVA™ (OSI Pharmaceuticals, Melville, NY), 43-9006 (Bayer, West Haven, CT), CCI-779 (Wyeth, Madison, NJ), RAD001 (Novartis), BMS-387032 (Bristol-Myers Squibb, New York, NY), IMC-1C11 (ImClone), LY333531 (Eli Lilly, Indianapolis, IN), PD 184352 (Pfizer), 2C4 (Genentech), and GW2016 (GlaxoSmithKline, Research Triangle Park, NC).

The role of IGF-1R in hematological malignancies has been reviewed by (Novak *et al.*, 2003, *Insulin-Like Growth Factors and Hematological Malignancies* in Insulin-Like Growth Factors, LeRoith *et al.*, ed.s, Landes Bioscience). A functional role for the IGF-1R in hematopoietic malignancies is demonstrated by, for example, the ability of IGF-1R monoclonal antibodies to block transformed cell growth in culture. IGF-I has been found to enhance growth of freshly isolated human acute myelogenous leukemia and acute lymphoblastic leukemia blasts. With respect to T cell malignancies, IGF-I has been shown to influence the growth of murine lymphoma cells bearing a pre-T

cell phenotype and, immature and mature primary human T lineage acute lymphoblastic leukemia cells were found to express high numbers of IGF-1R. Thus, in one embodiment, the present invention provides methods of treating a hematological malignancy in a subject in need thereof comprising administering to the subject an antagonist of IGF-1R as described herein. In another embodiment, the malignancy is an acute myelogenous leukemia, an acute lymphoblastic leukemia, or a T cell malignancy.

In another aspect, the present invention provides methods of identifying subjects who are more likely to benefit from treatment using the compositions and/or methods of treatment of the present invention. Such methods can enable a caregiver to better tailor a therapeutic regimen to a particular subject's needs and reduce the likelihood of an ineffective or counterproductive course of treatment. In one embodiment, the present invention provides a method of determining whether a subject is a candidate for treatment using a composition or method as described herein comprising determining whether a target cell type in the subject expresses IGF-1R, wherein if the target cell type expresses IGF-1R, then the subject is a candidate for treatment. In another embodiment, the method comprises determining the approximate average number of IGF-1R molecules per target cell, wherein 10^2, 10^3, 10^4, 10^5, or 10^6 IGF-1R per cell indicates that the subject is a candidate for treatment. The approximate average number of IGF-1R molecules per target cell can be determined using any technique known in the art, for example, by staining a sample comprising cells of the target cell type with an IGF-1R binding molecule, and detecting the amount of IGF-1R binding molecule bound to the sample, where the amount of IGF-1R binding molecule detected is proportional to the average number of IGF-1R molecules in the sample. In another embodiment, the method comprises comparing the approximate average number of IGF-1R molecules per target cell to a reference standard, wherein if the approximate average number of IGF-1R molecules per target cell is greater than the reference standard, then the subject is more likely to benefit from treatment using the compositions and/or methods of treatment of the present invention. In another embodiment, the target cell type is a cancerous cell type. In another embodiment, the target cell

type is a colon cancer cell type, a breast cancer cell type, an NSCLC cell type, or a leukemic cell type.

In another embodiment, a subject who is a candidate for treatment is identified by detecting IGF-1 and/or IGF-2 in the target cell type, or in the stratum of the target cell type. In another embodiment, the target cell type is a cancerous cell type. In another embodiment, the target cell type is a colon cancer cell type, a breast cancer cell type, an NSCLC cell type, or a leukemic cell type.

In another embodiment, a subject who is a candidate for treatment is identified by detecting activity of IGF-1R-mediated signaling in the target cell type, wherein IGF-1R-mediated signaling in the target cell type indicates that the subject is a candidate for treatment. Examples of molecules that can be monitored for IGF-1R-dependent changes are shown in Figure 10, such as molecules in the PI3/Akt pathway, *e.g.*, IGF-1R, IRS adapters, Akt, *etc.* Such molecules can be monitored for, for example, a change in phosphorylation status, *e.g.*, an increase in phosphorylation. Phosphospecific antibodies that recognize the activated forms of these protein markers are highly developed, and these reagents have proven to be reliable for immunoblot detection in experimental systems.

The compositions and/or methods of the present invention also can be used, for example, in cosmetic treatments, in veterinary treatments, to increase longevity, to treat reproductive defects, and to treat a variety of growth related disorders.

Therapeutic methods and administration of antigen binding proteins

Certain methods provided herein comprise administering an IGF-1R binding antigen binding protein to a subject, thereby reducing an IGF-1-induced biological response that plays a role in a particular condition. In particular embodiments, methods of the invention involve contacting endogenous IGF-1R with an IGF-1R binding antigen binding protein, *e.g.*, via administration to a subject or in an *ex vivo* procedure.

The term "treatment" encompasses alleviation or prevention of at least one symptom or other aspect of a disorder, or reduction of disease severity,

and the like. An antigen binding protein need not effect a complete cure, or eradicate every symptom or manifestation of a disease, to constitute a viable therapeutic agent. As is recognized in the pertinent field, drugs employed as therapeutic agents may reduce the severity of a given disease state, but need not abolish every manifestation of the disease to be regarded as useful therapeutic agents. Similarly, a prophylactically administered treatment need not be completely effective in preventing the onset of a condition in order to constitute a viable prophylactic agent. Simply reducing the impact of a disease (for example, by reducing the number or severity of its symptoms, or by increasing the effectiveness of another treatment, or by producing another beneficial effect), or reducing the likelihood that the disease will occur or worsen in a subject, is sufficient. One embodiment of the invention is directed to a method comprising administering to a patient an IGF-1R antagonist in an amount and for a time sufficient to induce a sustained improvement over baseline of an indicator that reflects the severity of the particular disorder.

As is understood in the pertinent field, pharmaceutical compositions comprising the molecules of the invention are administered to a subject in a manner appropriate to the indication. Pharmaceutical compositions may be administered by any suitable technique, including but not limited to parenterally, topically, or by inhalation. If injected, the pharmaceutical composition can be administered, for example, via intra-articular, intravenous, intramuscular, intralesional, intraperitoneal or subcutaneous routes, by bolus injection, or continuous infusion. Localized administration, *e.g.* at a site of disease or injury is contemplated, as are transdermal delivery and sustained release from implants. Delivery by inhalation includes, for example, nasal or oral inhalation, use of a nebulizer, inhalation of the antagonist in aerosol form, and the like. Other alternatives include eyedrops; oral preparations including pills, syrups, lozenges or chewing gum; and topical preparations such as lotions, gels, sprays, and ointments.

Use of antigen binding proteins in *ex vivo* procedures also is contemplated. For example, a patient's blood or other bodily fluid may be contacted with an antigen binding protein that binds IGF-1R *ex vivo*. The

antigen binding protein may be bound to a suitable insoluble matrix or solid support material.

Advantageously, antigen binding proteins are administered in the form of a composition comprising one or more additional components such as a physiologically acceptable carrier, excipient or diluent. Optionally, the composition additionally comprises one or more physiologically active agents, for example, a second IGF-1 receptor-inhibiting substance, an anti-angiogenic substance, a chemotherapeutic substance, an analgesic substance, *etc.*, non-exclusive examples of which are provided herein. In various particular embodiments, the composition comprises one, two, three, four, five, or six physiologically active agents in addition to an IGF-1R binding antigen binding protein

In one embodiment, the pharmaceutical composition comprise an antigen binding protein of the invention together with one or more substances selected from the group consisting of a buffer, an antioxidant such as ascorbic acid, a low molecular weight polypeptide (such as those having fewer than 10 amino acids), a protein, an amino acid, a carbohydrate such as glucose, sucrose or dextrins, a chelating agent such as EDTA, glutathione, a stabilizer, and an excipient. Neutral buffered saline or saline mixed with conspecific serum albumin are examples of appropriate diluents. In accordance with appropriate industry standards, preservatives such as benzyl alcohol may also be added. The composition may be formulated as a lyophilizate using appropriate excipient solutions (*e.g.*, sucrose) as diluents. Suitable components are nontoxic to recipients at the dosages and concentrations employed. Further examples of components that may be employed in pharmaceutical formulations are presented in Remington's Pharmaceutical Sciences, 16th Ed. (1980) and 20th Ed. (2000), Mack Publishing Company, Easton, PA.

Kits for use by medical practitioners include an IGF-1 receptor-inhibiting substance of the invention and a label or other instructions for use in treating any of the conditions discussed herein. In one embodiment, the kit includes a sterile preparation of one or more IGF-1R binding antigen binding

proteins, which may be in the form of a composition as disclosed above, and may be in one or more vials.

Dosages and the frequency of administration may vary according to such factors as the route of administration, the particular antigen binding proteins employed, the nature and severity of the disease to be treated, whether the condition is acute or chronic, and the size and general condition of the subject. Appropriate dosages can be determined by procedures known in the pertinent art, *e.g.* in clinical trials that may involve dose escalation studies.

An IGF-1 receptor inhibiting substance of the invention may be administered, for example, once or more than once, *e.g.*, at regular intervals over a period of time. In particular embodiments, an antigen binding protein is administered over a period of at least a month or more, *e.g.*, for one, two, or three months or even indefinitely. For treating chronic conditions, long-term treatment is generally most effective. However, for treating acute conditions, administration for shorter periods, *e.g.* from one to six weeks, may be sufficient. In general, the antigen binding protein is administered until the patient manifests a medically relevant degree of improvement over baseline for the chosen indicator or indicators.

Particular embodiments of the present invention involve administering an antigen binding protein at a dosage of from about 1 ng of antigen binding protein per kg of subject's weight per day ("1ng/kg/day") to about 10 mg/kg/day, more preferably from about 500 ng/kg/day to about 5 mg/kg/day, and most preferably from about 5 μg/kg/day to about 2 mg/kg/day, to a subject. In additional embodiments, an antigen binding protein is administered to adults one time per week, two times per week, or three or more times per week, to treat an IGF-1 and/or IGF-2 mediated disease, condition or disorder, *e.g.*, a medical disorder disclosed herein. If injected, the effective amount of antigen binding protein per adult dose may range from 1-20 mg/m^2, and preferably is about 5-12 mg/m^2. Alternatively, a flat dose may be administered; the amount may range from 5-100 mg/dose. One range for a flat dose is about 20-30 mg per dose. In one embodiment of the invention, a flat dose of 25 mg/dose is repeatedly administered by injection. If a route of administration other than injection is used, the dose is appropriately adjusted

in accordance with standard medical practices. One example of a therapeutic regimen involves injecting a dose of about 20-30 mg of antigen binding protein to one to three times per week over a period of at least three weeks, though treatment for longer periods may be necessary to induce the desired degree of improvement. For pediatric subjects (age 4-17), one exemplary suitable regimen involves the subcutaneous injection of 0.4 mg/kg, up to a maximum dose of 25 mg of antigen binding protein administered two or three times per week.

Particular embodiments of the methods provided herein involve subcutaneous injection of from 0.5 mg to 10 mg, preferably from 3 to 5 mg, of an antigen binding protein, once or twice per week. Another embodiment is directed to pulmonary administration (*e.g.*, by nebulizer) of 3 or more mg of antigen binding protein once a week.

Examples of therapeutic regimens provided herein comprise subcutaneous injection of an antigen binding protein once a week, at a dose of 1.5 to 3 mg, to treat a condition in which IGF-1R signaling plays a role. Examples of such conditions are provided herein and include, for example, cancer, acromegaly and other overgrowth disorders, diabetes, obesity, macular degeneration, and aging. Weekly administration of antigen binding protein is continued until a desired result is achieved, *e.g.*, the subject's symptoms subside. Treatment may resume as needed, or, alternatively, maintenance doses may be administered.

Other examples of therapeutic regimens provided herein comprise subcutaneous or intravenous administration of a dose of 1, 3, 5, 6, 7, 8, 9, 10, 11, 12, 15, or 20 milligrams of an IGF-1R inhibitor of the present invention per kilogram body mass of the subject (mg/kg). The dose can be administered once to the subject, or more than once at a certain interval, for example, once a day, three times a week, twice a week, once a week, three times a month, twice a month, once a month, once every two months, once every three months, once every six months, or once a year. The duration of the treatment, and any changes to the dose and/or frequency of treatment, can be altered or varied during the course of treatment in order to meet the particular needs of the subject.

In another embodiment, an antigen binding protein is administered to the subject in an amount and for a time sufficient to induce an improvement, preferably a sustained improvement, in at least one indicator that reflects the severity of the disorder that is being treated. Various indicators that reflect the extent of the subject's illness, disease or condition may be assessed for determining whether the amount and time of the treatment is sufficient. Such indicators include, for example, clinically recognized indicators of disease severity, symptoms, or manifestations of the disorder in question. In one embodiment, an improvement is considered to be sustained if the subject exhibits the improvement on at least two occasions separated by two to four weeks. The degree of improvement generally is determined by a physician, who may make this determination based on signs, symptoms, biopsies, or other test results, and who may also employ questionnaires that are administered to the subject, such as quality-of-life questionnaires developed for a given disease.

Elevated levels of IGF-1 and/or IGF-2 are associated with a number of disorders, including, for example, cancer (*e.g.*, lung, prostate, breast and colon cancers), and acromegaly and other overgrowth disorders (*e.g.*, constitutionally tall children). Subjects with a given disorder may be screened, to identify those individuals who have elevated IGF-1 and/or IGF-2 levels, thereby identifying the subjects who may benefit most from treatment with an IGF-1R binding antigen binding protein. Thus, treatment methods provided herein optionally comprise a first step of measuring a subject's IGF-1 and/or IGF-2 levels. An antigen binding protein may be administered to a subject in whom IGF-1 and/or IGF-2 levels are elevated above normal. In one embodiment, the present invention provides a method of treating an overgrowth disorder (*e.g.*, acromegaly) comprising administering to a subject in need thereof an antigen binding protein of the present invention and pegvisomant.

A subject's levels of IGF-1 and/or IGF-2 may be monitored before, during and/or after treatment with an antigen binding protein, to detect changes, if any, in their levels. For some disorders, the incidence of elevated IGF-1 and/or IGF-2 levels may vary according to such factors as the stage of

the disease or the particular form of the disease. Known techniques may be employed for measuring IGF-1 and/or IGF-2 levels, *e.g.*, in a subject's serum. IGF-1 and/or IGF-2 levels in blood samples may be measured using any suitable technique, for example, ELISA.

Particular embodiments of methods and compositions of the invention involve the use of an antigen binding protein and one or more additional IGF-1R antagonists, for example, two or more antigen binding proteins of the invention, or an antigen binding protein of the invention and one or more other IGF-1R antagonists. In further embodiments, antigen binding protein are administered alone or in combination with other agents useful for treating the condition with which the patient is afflicted. Examples of such agents include both proteinaceous and non-proteinaceous drugs. When multiple therapeutics are co-administered, dosages may be adjusted accordingly, as is recognized in the pertinent art. "Co-administration" and combination therapy are not limited to simultaneous administration, but also include treatment regimens in which an antigen binding protein is administered at least once during a course of treatment that involves administering at least one other therapeutic agent to the patient.

Examples of other agents that may be co-administered with an antigen binding protein are other antigen binding proteins or therapeutic polypeptides that are chosen according to the particular condition to be treated. Alternatively, non-proteinaceous drugs that are useful in treating one of the particular conditions discussed above may be co-administered with an IGF-1R antagonist.

Combination therapy

In another aspect, the present invention provides a method of treating a subject with an IGF-1R inhibiting antigen binding protein and one or more other treatments. In one embodiment, such a combination therapy achieves synergy or an additive effect by, for example, attacking multiple sites or molecular targets in a tumor. Types of combination therapies that can be used in connection with the present invention include inhibiting or activating (as appropriate) multiple nodes in a single disease-related pathway, multiple

heavy chain CDR3 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises an amino acid sequence selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of four amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR2 sequence of L1-L52 as shown in Figure 5; c. a light chain CDR3 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR3 sequence of L1-L52 as shown in Figure 6; d. a heavy chain CDR1 sequence of H1-H52 as shown in Figure 7; e. a heavy chain CDR2 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and f. a heavy chain CDR3 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises an amino acid sequence selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of three amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a light chain CDR3 sequence of L1-L52 as shown in Figure 6; c. a heavy chain CDR2 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR2 sequence of H1-H52 as shown in Figure 8; and d. a heavy chain CDR3 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR3 sequence of H1-H52 as shown in Figure 9. In another embodiment, the isolated antigen binding protein comprises an amino acid sequence selected from the group consisting of: a. a light chain CDR1 sequence that differs by no more than a total of two amino acid additions, substitutions, and/or deletions from a CDR1 sequence of L1-L52 as shown in Figure 4; b. a heavy chain CDR2 sequence that differs by no more than a total of one amino acid addition, substitution, or deletion from a CDR2 sequence of H1-H52 as shown in Figure 8; and c. a heavy chain CDR3 sequence of H1-H52 as shown in Figure 9. In another

pathways in a target cell, and multiple cell types within a target tissue (e.g., within a tumor). For example, an IGF-1R inhibitor of the present invention can be combined with a treatment that inhibits IGF-1, promotes apoptosis, inhibits angiogenesis, or inhibits macrophage. In another embodiment, a targeted agent, that, when used by itself, fails to elicit a therapeutically desired effect, could be used to, for example, sensitize cancer cells or augment treatment effect of other agents. In another embodiment, an IGF-1R inhibitor according to the invention is used in combination with a cytotoxic drug or other targeted agent that induces apoptosis. In another embodiment, an IGF-1R inhibitor is used in combination with one or more agents that inhibit different targets that are involved in cell survival (*e.g.*, PKB, mTOR), different receptor tyrosine kinases (*e.g.*, ErbB1, ErbB2, c-Met, c-kit), or different cell types (*e.g.*, KDR inhibitors, c-fms). In another embodiment, an IGF-1R inhibitor of the invention is added to the existing standard of care for a particular condition. Examples of therapeutic agents include, but are not limited to, gemcitabine, taxol, taxotere, and CPT-11.

In another embodiment, a combination therapy method comprises administering to the subject two, three, four, five, six, or more of the IGF-1R agonists or antagonists described herein. In another embodiment, the method comprises administering to the subject two or more treatments that together inhibit or activate (directly or indirectly) IGF-1R-mediated signal transduction. Examples of such methods include using combinations of two or more IGF-1R inhibiting antigen binding progeins, of an IGF-1R inhibiting antigen binding protein and one or more other IGF-1, IGF-2, and/or IGF-1R agonists or antagonists (*e.g.*, IGF-1 and/or IGF-2 binding polypeptides, IGF-1R binding polypeptides, IGF-1 and/or IGF-2 derivatives, anti-IGF-1 and/or IGF-2 antibodies, anti-sense nucleic acids against IGF-1, IGF-2, and/or IGF-1R, or other molecules that bind to IGF-1, IGF-2, and/or IGF-1R polypeptides or nucleic acids), or of an IGF-1R inhibiting antigen binding protein and one or more other treatments (*e.g.*, surgery, ultrasound, radiotherapy, chemotherapy, or treatment with another anti-cancer agent), as described, for example, in US Pat. No. 5,473,054 (issued Dec. 5, 1995), 6,051,593 (issued April 18, 2000), 6,084,085 (issued July 4, 2000), 6,506,763 (issued Jan. 14,

2003), US Pat. App. Pub. No.s 03/0092631 (published May 15, 2003), 03/0165502 (published Sept. 4, 2003), 03/0235582 (published Dec. 25, 2003), 04/0886503 (published May 6, 2004), 05/0272637 (published Dec. 8, 2005), PCT Pub. Ser. No.s WO 99/60023 (published Nov. 25, 1999), WO 02/053596 (published July 11, 2002), WO 02/072780 (published Sept. 19, 2002), WO 03/027246 (published March 3, 2003), WO 03/020698 (published March 13, 2003), WO 03/059951 (published July 24, 2003), WO 03/100008 (published Dec. 4, 2003), WO 03/106621 (published Dec. 24, 2003), WO 04/071529 (published August 26, 2004), WO 04/083248 (published Sept. 30, 2004), WO 04/087756 (published Oct. 14, 2004), WO 05/112969 (published Dec. 1, 2005), Kull *et al.*, 1983, J Biol Chem 258:6561-66, Flier *et al.*, 1986, Proc Natl Acad Sci USA 83:664-668, Conover *et al.*, 1987, J Cell Physiol 133:560-66, Rohlik *et al.*, 1987, Biochem Biophys Res Comm 149:276-81, Arteaga *et al.*, 1989, J Clinical Investigation 84:1418-23, Arteaga *et al.*, 1989, Cancer Res 49:6237-41, Gansler *et al.*, 1989, American J Pathol 135:961-66, Gustafson *et al.*, 1990, J Biol Chem 265:18663-67, Steele-Perkins *et al.*, 1990, Biochem Biophys Res Comm 171:1244-51, Cullen *et al.*, 1992, Mol Endocrinol 6:91-100, Soos *et al.*, 1992, J Biol Chem 267:12955-63, Xiong *et al.*, 1992, Proc Natl Acad Sci USA 89:5356-60, Brunner *et al.*, 1993, Euro J Cancer 29A:562-69, Furlanetto *et al.*, 1993, Cancer Res 53:2522-26, Li *et al.*, 1993, Biochem Biophys Res Comm 196:92-98, Kalebic *et al.*, 1994, Cancer Res 54:5531-34, Lahm *et al.*, 1994, Intl J Cancer 58:452-59, Zia *et al.*, 1996, J Cell Biochem Supp 24:269-75, Jansson *et al.*, 1997, J Biol Chem 272:8189-97, Scotlandi *et al.*, 1998, Cancer Res 58:4127-31, Logie *et al.*, 1999, Li *et al.*, 2000, Cancer Immunol Immunotherapy 49:243-52, J Mol Endocrinol 23:23-32, De Meyts *et al.*, 2002, Nature Reviews 1:769-83, Hailey *et al.*, 2002, Mol Cancer Therapeutics 1:1349-53, Maloney *et al.*, 2003, Cancer Research 63:5073-83, Burtrum *et al.*, 2003, Cancer Research 63:8912-21, and Karavitaki *et al.*, 2004, Hormones 3:27-36, (each incorporated herein by reference in its entirety) may be employed in methods and compositions of the present invention. Furthermore, one or more anti-IGF-1R antibodies or antibody derivatives can be used in combination with one or more molecules or other treatments, wherein the other molecule(s) and/or treatment(s) do not

directly bind to or affect IGF-1R, IGF-1, or IGF-2, but which combination is effective for treating or preventing a condition, such as cancer or an overgrowth disorder (*e.g.*, acromegaly). In one embodiment, one or more of the molecule(s) and/or treatment(s) treats or prevents a condition that is caused by one or more of the other molecule(s) or treatment(s) in the course of therapy, *e.g.*, nausea, fatigue, alopecia, cachexia, insomnia, *etc.* In every case where a combination of molecules and/or other treatments is used, the individual molecule(s) and/or treatment(s) can be administered in any order, over any length of time, which is effective, *e.g.*, simultaneously, consecutively, or alternately. In one embodiment, the method of treatment comprises completing a first course of treatment with one molecule or other treatment before beginning a second course of treatment. The length of time between the end of the first course of treatment and beginning of the second course of treatment can be any length of time that allows the total course of therapy to be effective, *e.g.*, seconds, minutes, hours, days, weeks, months, or even years.

In another embodiment, the method comprises administering one or more of the IGF-1R antagonists described herein and one or more other treatments (*e.g.*, a therapeutic or palliative treatment), for example, anti-cancer treatments (such as surgery, ultrasound, radiotherapy, chemotherapy, or treatment with another anti-cancer agent). Where a method comprises administering more than one treatment to a subject, it is to be understood that the order, timing, number, concentration, and volume of the administrations is limited only by the medical requirements and limitations of the treatment, *i.e.*, two treatments can be administered to the subject, *e.g.*, simultaneously, consecutively, alternately, or according to any other regimen. Examples of agents that can be administered in combination with the IGF-1R antagonists described herein include, but are not limited to, neutrophil-boosting agents, irinothecan, SN-38, gemcitabine, herstatin, or an IGF-1R-binding herstatin derivative (as described, for example, in US Pat. App. No. 05/0272637), AVASTIN® (Genentech, South San Francisco, CA), HERCEPTIN® (Genentech), RITUXAN® (Genentech), ARIMIDEX® (AstraZeneca, Wilmington, DE), IRESSA® (AstraZeneca), BEXXAR® (Corixa, Seattle,

WA), ZEVALIN® (Biogen Idec, Cambridge, MA), ERBITUX® (Imclone Systems Inc., New York, NY), GEMZAR® (Eli Lilly and Co., Indianapolis, IN), CAMPTOSAR® (Pfizer, New York, NY), GLEEVEC® (Novartis), SU-11248 (Pfizer), BMS-354825 (Bristol-Myers Squibb), panitumumab (Abgenix, Fremont, CA/Amgen Inc., Thousand Oaks, CA), and denosumab (Amgen Inc., Thousand Oaks, CA).

The following examples, both actual and prophetic, are provided for the purpose of illustrating specific embodiments or features of the instant invention and do not limit its scope.

EXAMPLE 1: Preparation of Antibodies

This example demonstrates a method of preparing antibodies recognizing the IGF-1 receptor.

IGF-1 receptor polypeptides may be employed as immunogens in generating monoclonal antibodies by conventional techniques. It is recognized that polypeptides in various forms may be employed as immunogens, *e.g.*, full length proteins, fragments thereof, fusion proteins thereof such as Fc fusions, cells expressing the recombinant protein on the cell surface, *etc.*

To summarize an example of such a procedure, an IGF-1R immunogen emulsified in complete Freund's adjuvant is injected subcutaneously into Lewis rats, in amounts ranging from 10-100 µl. Three weeks later, the immunized animals are boosted with additional immunogen emulsified in incomplete Freund's adjuvant and boosted every three weeks thereafter. Serum samples are periodically taken by retro-orbital bleeding or tail-tip excision for testing by dot-blot assay, ELISA (enzyme-linked immunosorbent assay), or inhibition of binding of ^{125}I-IGF-1 or ^{125}I-IGF-2 to extracts of IGF-1R-expressing cells. Following detection of an appropriate antibody titer, positive animals are given a final intravenous injection of antigen in saline. Three to four days later, the animals are sacrificed, splenocytes harvested, and fused to the murine myeloma cell line AG8653. The resulting hybridoma cell lines are plated in multiple microtiter plates in a HAT selective medium (hypoxanthine, aminopterin, and thymidine) to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

Hybridoma clones thus generated are screened for reactivity with IGF-1R. Initial screening of hybridoma supernatants utilizes an antibody capture and binding of partially purified ^{125}I-IGF-1 receptor. Hybridomas that are positive in this screening method are tested by a modified antibody capture to detect hybridoma cells lines that are producing blocking antibody. Hybridomas that secrete a monoclonal antibody capable of inhibiting ^{125}I-IGF-1 binding to cells expressing IGF-1R are thus detected. Such hydridomas then are injected into the peritoneal cavities of nude mice to produce ascites containing high concentrations (>1 mg/ml) of anti-IGF-1R monoclonal antibody. The resulting monoclonal antibodies may be purified by ammonium sulfate precipitation followed by gel exclusion chromatography, and/or affinity chromatography based on binding of antibody to Protein G.

Similar methods can be used to generate human antibodies in transgenic mice. See, *e.g.*, Chen *et al.*, 1993, lnternat. Immunol. 5: 647-56; Chen *et al.*, 1993, EMBO J. 12: 821-30; Choi *et al.*, 1993, Nature Genetics 4: 117-23; Fishwild *et al.*, 1996, Nature Biotech. 14: 845-51; Harding et al., 1995, Annals New York Acad. Sci.; Lonberg *et al.*, 1994, Nature 368: 856-59; Lonberg, 1994, Handbook Exper.l Pharmacol. 113: 49-101; Lonberg *et al.*, 1995, Internal Rev. Immunol. 13: 65-93; Morrison, 1994, Nature 368: 812-13; Neuberger, 1996, Nature Biotech. 14: 826; Taylor *et al.*, 1992, Nuc. Acids Res. 20: 6287-95; Taylor *et al.*, 1994, lnternat. Immunol. 6: 579-91; Tomizuka *et al.*, 1997, Nature Genetics 16: 133-43; Tomizuka *et al.*, 2000, Proc. Nat. Acad. Sci. USA 97: 722-27; Tuaillon et al., 1993, Proc. Nat. Acad. Sci. USA 90: 3720-24; Tuaillon *et al.*, 1994, J. Immunol. 152: 2912-20; Russel *et al.*, 2000, Infection and Immunity April 2000: 1820-26; Gallo *et al.*, 2000, Eur. J. Immunol. 30: 534-40; Davis *et al.*, 1999, Cancer Metastasis Rev. 18:421-25; Green, 1999, J. Immunol. Methods 231:11-23; Jakobovits, 1998, Advanced Drug Delivery Rev. 31:33-42; Green *et al.*, 1998, J. Exp. Med. 188: 483-95; Jakobovits, 1998, Exp. Opin. Invest. Drugs 7: 607-14; Tsuda *et al.*, 1997, Genomics 42: 413-21; Mendez *et al.*, 1997, Nature Genetics 15: 146-56; Jakobovits, 1996, Weir's Handbook of Experimental Immunology, The Integrated Immune System Vol. IV, 194.1-194.7; Mendez *et al.*,1995, Genomics 26: 294-307; Jakobovits, 1994, Current Biol. 4: 761-63; Arbones,

1994, Immunity 1: 247-60; Green *et al.*, 1994, Nature Genetics 7: 13-21; Jakobovits *et al.*, 1993, Nature 362: 255-58; Jakobovits *et al.*, 1993,. Proc. Nat. Acad. Sci. USA 90: 2551-55.

EXAMPLE 2: Isolation of Human IGF-1R(ECD)-C3-muIgG1

This example provides a method of making a soluble fragment of IGF-1R useful for raising antibodies.

Cloning of pDSRα:huIGF-1R(ECD)-C3-muIgG1Fc

Primers 2830-36:

5' AGCAAGCTTCCACCATGAAGTCTGGCTCCGGAGGAGG 3' SEQ ID NO:256)

and 2830-38:

5' ATTTGTCGACTTCGTCCAGATGGATGAAGTTTTCAT 3', SEQ ID NO:257)

were used to amplify the human IGF-1R extracellular domain (1-906) cDNA sequence. The primers included a Kozak translation initiation sequence (underlined above) preceding the start codon, restriction sites for subsequent subcloning, and a caspace-3 site, which is inserted next to the extracellular domain C-terminus. PCR was performed on a PerkinElmer 2400 (PerkinElmer, Torrance, CA) under the following conditions: 1 cycle at 95° C for 2 min, 23 cycles at 95° C for 30 sec, 58.5° C for 30 sec, and 72° C for 3 min, and 1 cycle at 72° C for 10 min. Final reaction conditions were 1X *pfu* TURBO® buffer (Stratagene, La Jolla, CA), 200 μM dNTPs, 2 μM each primer, 5 U *pfu* TURBO® (Stratagene) and 1 ng template DNA. The PCR product was purified using a Clontech Nucleospin Column (Clontech, Palo Alto, CA) according to the manufacturers instructions, digested with *Hind* III and *Sal* I (Roche, Indianapolis, IN) and gel purified. The human IGF-1R insert was ligated into *Hind* III/*Sal* I digested pDSRα-muIgG1. Integrity of the insert was confirmed by DNA sequencing. The sequence of the protein encoded by the resulting open reading frame (IGF-1R-C3-muFc) is shown in Figure 10. The final expression vector, pDSRα:huIGF1R(ECD)-C3-muIgG1Fc, is described in Table 1.

Table 1

pDSRα:huIGF1R(ECD)-C3-muIgG1Fc

Plasmid Base

Pair Number:

11-3496	HuIGF1R (Caspase 3 site)-muIgG1Fc atgaagtctggctccggaggagggtccccgacctcgctgtgggggctcctgtttctctccgccgcgctctc gctctggccgacgagtggagaaatctgcgggccaggcatcgacatccgcaacgactatcagcagctgaa gcgcctggagaactgcacggtgatcgagggctacctccacatcctgctcatctccaaggccgaggactac cgcagctaccgcttccccaagctcacggtcattaccgagtacttgctgctgttccgagtggctggcctcgag agcctcggagacctcttccccaacctcacggtcatccgcggctggaaactcttctacaactacgccctggtc atcttcgagatgaccaatctcaaggatattgggctttacaacctgaggaacattactcgggggccatcagg attgagaaaaatgctgacctctgttacctctccactgtggactggtccctgatcctggatgcggtgtccaataa ctacattgtggggaataagcccccaaaggaatgtggggacctgtgtccagggaccatggaggagaagcc gatgtgtgagaagaccaccatcaacaatgagtacaactaccgctgctggaccacaaaccgctgccagaaa atgtgcccaagcacgtgtgggaagcgggcgtgcaccgagaacaatgagtgctgccaccccgagtgcctg ggcagctgcagcgcgcctgacaacgacacggcctgtgtagcttgccgccactactactatgccggtgtctg tgtgcctgcctgcccgcccaacacctacaggtttgagggctggcgctgtgtggaccgtgacttctgcgcca acatcctcagcgccgagagcagcgactccgaggggtttgtgatccacgacggcgagtgcatgcaggagt gcccctcgggcttcatccgcaacggcagccagagcatgtactgcatcccttgtgaaggtccttgcccgaag gtctgtgaggaagaaaagaaaacaaagaccattgattctgttacttctgctcagatgctccaaggatgcacc atcttcaagggcaatttgctcattaacatccgacgggggaataacattgcttcagagctggagaacttcatgg ggctcatcgaggtggtgacgggctacgtgaagatccgccattctcatgccttggtctccttgtccttcctaaa aaaccttcgcctcatcctaggagaggagcagctagaagggaattactccttctacgtcctcgacaaccaga acttgcagcaactgtgggactgggaccaccgcaacctgaccatcaaagcagggaaaatgtactttgctttc aatcccaaattatgtgtttccgaaatttaccgcatggaggaagtgacggggactaaagggcgccaaagcaa aggggacataaacaccaggaacaacggggagagagcctcctgtgaaagtgacgtcctgcatttcacctcc accaccacgtcgaagaatcgcatcatcataacctggcaccggtaccggccccctgactacagggatctcat cagcttcaccgtttactacaaggaagcacccttaagaatgtcacagagtatgatgggcaggatgcctgcgg ctccaacagctggaacatggtggacgtggacctcccgcccaacaaggacgtggagcccggcatcttacta catgggctgaagccctggactcagtacgccgtttacgtcaaggctgtgaccctcaccatggtggagaacga ccatatccgtggggccaagagtgagatcttgtacattcgcaccaatgcttcagttccttccattcccttggacg ttctttcagcatcgaactcctcttctcagttaatcgtgaagtggaaccctcctctctgcccaacggcaacctg agttactacattgtgcgctggcagcggcagcctcaggacggctacctttaccggcacaattactgctccaaa gacaaaatccccatcaggaagtatgccgacggcaccatcgacattgaggaggtcacagagaaccccaag actgaggtgtgtggtggggagaaagggccttgctgcgcctgccccaaaactgaagccgagaagcaggc cgagaaggaggaggctgaataccgcaaagtctttgagaatttcctgcacaactccatcttcgtgcccagac ctgaaaggaagcggagagatgtcatgcaagtggccaacaccaccatgtccagccgaagcaggaacacc acggccgcagacacctacaacatcactgacccggaagagctggagacagagtacccttttctttgagagca

	gagtggataacaaggagagaactgtcatttctaaccttcggcctttcacattgtaccgcatcgatatccacag ctgcaaccacgaggctgagaagctgggctgcagcgcctccaacttcgtctttgcaaggactatgcccgca gaaggagcagatgacattcctgggccagtgacctgggagccaaggcctgaaaactccatctttttaaagtg gccggaacctgagaatcccaatggattgattctaatgtatgaaataaaatacggatcacaagttgaggatca gcgagaatgtgtgtccagacaggaatacaggaagtatggaggggccaagctaaaccggctaaacccgg ggaactacacagcccggattcaggccacatctctctctgggaatgggtcgtggacagatcctgtgttcttcta tgtccaggccaaaacaggatatgaaaacttcatccatctggacgaagtcgacggttgtaagccttgcatatgt acagtcccagaagtatcatctgtcttcatcttccccccaaagcccaaggatgtgctcaccattactctgactcc taaggtcacgtgtgttgtggtagacatcagcaaggatgatcccgaggtccagttcagctggtttgtagatgat gtggaggtgcacacagctcagacgcaacccccgggaggagcagttcaacagcactttccgctcagtcagt gaacttcccatcatgcaccaggactggctcaatggcaaggagttcaaatgcagggtaaacagtgcagcttt ccctgcccccatcgagaaaaccatctccaaaaccaaaggcagaccgaaggctccacaggtgtacaccatt ccacctcccaaggagcagatggccaaggataaagtcagtctgacctgcatgataacagacttcttccctga agacattactgtggagtggcagtggaatgggcagccagcggagaactacaagaacactcagcccatcat ggacacagatggctcttacttcgtctacagcaagctcaatgtgcagaagagcaactgggaggcaggaaat actttcacctgctctgtgttacatgagggcctgcacaaccaccatactgagaagagcctctcccactctcctg gtaaa (SEQ ID NO:258)
3507 to 4391	A transcription termination/polyadenylation signal from the α-subunit of the bovine pituitary glycoprotein hormone (α-FSH) (Goodwin *et al.*, 1983, *Nucleic Acids Res.* 11:6873-82; Genbank Accession Number X00004)
4600 to 5163	A mouse dihydrofolate reductase (DHFR) minigene containing the endogenous mouse DHFR promoter, the cDNA coding sequences, and the DHFR transcription termination/polyadenylation signals (Gasser *et al.*, 1982, *Proc. Natl. Acad. Sci. U. S. A.* 79:6522-6; Nunberg *et al.*, 1980, *Cell* 19:355-64; Setzer *et al.*, 1982, *J. Biol. Chem.* 257:5143-7; McGrogan *et al.*, 1985, *J. Biol. Chem.* 260:2307-14)
6389 to 7246	pBR322 sequences containing the ampicillin resistance marker gene and the origin for replication of the plasmid in *E. coli* (Genbank Accession Number J01749)
7459 to 7802	An SV40 early promoter, enhancer and origin of replication (*Takebe et al.*, 1988, *Mol. Cell Biol.* 8:466-72, Genbank Accession Number J02400)
7809 to 8065	A translational enhancer element from the HTLV-1 LTR domain (Seiki *et al.*, 1983, *Proc. Natl. Acad. Sci. U. S. A.* 80:3618-22, Genbank Accession Number J02029)
8109 to 8205	An intron from the SV40 16S, 19S splice donor/acceptor signals (Okayama and Berg, 1983. *Mol. Cell Biol.* 3:280-9, Genbank Accession Number J02400)

Expression of hu IGF-1R(ECD)-C3-muIgG1Fc

Fifteen micrograms of linearized expression vector pDSRα:huIGF1R(ECD)-C3-muIgG1Fc was transfected into AM-1/D CHOd- cells using LT1 lipofection reagent (PanVera Corp., Madison, WI), and cells cultured under conditions to allow expression and secretion of protein into the cell media. Twenty-four colonies were selected after 10-14 days on DHFR selection medium (Dulbecco's Modified Eagles Medium (Invitrogen) supplemented with 10% dialyzed fetal bovine serum, 1x penicillin-streptomycin (Invitrogen)) and expression levels evaluated by western blot. To perform this assay, 0.5 ml of serum free medium was added to a single well confluent cells cultured in a 24 well plate (Falcon). The conditioned medium was recovered after 48hr. Samples for western blotting were run in 10% Tris-glycine gel (Novex), and blotted on 0.45 μm Nitrocellulose membrane (Invitrogen), using the Mini Trans-Blot cell (Biorad). The blotted membranes were incubated with rabbit anti-mouse IgG Fc antibody, conjugated with Horseradish Peroxidase (Pierce). The clone expressing the highest level of IGF-1R(ECD)-C3-muIgG1Fc was expanded in DHFR selection medium and 2 x 10^7 cells were inoculated into 50 roller bottles each (Corning) in 250 ml of high-glucose DMEM (Invitrogen), 10% dialyzed FBS (Invitrogen), 1x glutamine (Invitrogen), 1x Non essential amino acids (Invitrogen), 1x sodium pyruvate (Invitrogen). Medium was gassed with 10% CO_2/balance air for 5 seconds before capping the roller bottle. Roller bottles were kept at 37° C on roller racks spinning at 0.75 rpm.

When cells reached approximately 85-90% confluency (after approximately 5-6 days in culture), growth medium was discarded, cells washed with 100 ml PBS and 200 ml production medium was added (50 % DMEM (Invitrogen)/ 50 % F12 (Invitrogen), 1x glutamine (Invitrogen), 1x non-essential amino acids (Invitrogen), 1x sodium pyruvate (Invitrogen), 1.5% DMSO (Sigma)). The conditioned medium was harvested and replaced at one week intervals. The resulting 30 liters of conditioned medium were filtered through a 0.45 μm cellulose acetate filter (Corning, Acton, MA).

Purification of hu IGF-1R(ECD)-C3-muIgG1Fc

The resulting filtrate from the conditioned medium was concentrated 20-fold using a spiral-wound cartridge (molecular weight cut-off = 10 kDa), then diluted 1:1 with 3 M KCl, 1 M glycine, pH 9.0 to bring the final salt concentration to 1.5 M KCl, 0.5 M glycine, pH 9.0. This sample was applied to a rProtein A-Sepharose column (Amersham Pharmacia Biotech, Uppsala, Sweden) which had been equilibrated in 1.5 M KCl, 0.5 M glycine, pH 9.0. The column was washed with 40 column volumes of the same buffer, then eluted with 20 column volumes of 0.1 M glycine-HCl, pH 2.8. Five-mL fractions were collected and immediately neutralized with 1 mL of 1 M Tris-HCl, pH 7.5. Fractions containing huIGF1R(ECD)-C3-muIgGFc were identified by SDS-PAGE, pooled, and dialyzed against phosphate-buffered saline. The yield was 2.4 mg/L of conditioned medium. The major protein species detected were the mature α and β chains and murine Fc, each of which appeared to be properly glycosylated based on their elevated and heterogeneous molecular weights. Unprocessed IGF-1R(ECD), as well as glycosylated but not proteolytically cleaved IGF-1R(CED), was also present in the preparation. The shift in bands to higher molecular weights under non-reducing conditions indicates that disulfide linkages joined the α □and β chains. Amino-terminal sequencing of the final product indicated that 60% of the protein was correctly processed between the α- and β-chains of IGF-1R(ECD), while 40% remained unprocessed.

EXAMPLE 3: Isolation of Human INSR(ECD)-muIgG1

This example presents a method of cloning and expressing a soluble fragment of the human insulin receptor.

Cloning of pDSRα:huINSR(ECD)-muIgG1Fc

Primers 2830-40:

5' AGCAAGCTTCCACCATGGGCACCGGGGGCCGG 3' SEQ ID NO:259

(*Hin*d III site underlined) and 2830-41:

5' ATTTGTCGACTTTTGCAATATTTGACGGGACGTCTAA 3' SEQ ID NO:260

(*Sal* I site underlined) were used to amplify the human INSR extracellular domain (1-929) from and INSR parental plamid encoding the B form of the

What is claimed is:

What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence, or its complement, wherein said nucleotide sequence encodes the light chain, the heavy chain, or both the light chain and the heavy chain of an anti-IGF-1R antibody or an antigen binding fragment of said anti-IGF-1R antibody, wherein:

a. the light chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises the amino acid sequence of SEQ ID NO:32 and the heavy chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises the amino acid sequence of SEQ ID NO:136; or

b. the light chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises:

i. the CDR 1 sequence of residues 24 through 39 of SEQ ID NO:32; and

ii. the CDR 2 sequence of residues 55 through 61 of SEQ ID NO:32; and

iii. the CDR 3 sequence of residues 94 through 102 of SEQ ID NO:32;

and the heavy chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises:

i. the CDR 1 sequence of residues 31 through 36 of SEQ ID NO:136; and

ii. the CDR 2 sequence of residues 51 through 66 of SEQ ID NO:136; and

iii. the CDR 3 sequence of residues 99 through 108 of SEQ ID NO:136; or

c. said nucleotide sequence encodes both the light chain and the heavy chain of said anti-IGF-IR antibody, or of said antigen binding fragment of said anti-IGF1R antibody, and wherein the light chain variable region of said anti-IGF-1R antibody or antigen binding fragment is at least 90% identical to SEQ ID NO:32 and the heavy

c. the light chain of said anti-IGF-1R antibody or antigen binding fragment comprises the kappa light chain constant sequence of SEQ ID NO:234 and the heavy chain of said anti-IGF-1R antibody or antigen binding fragment further comprises the IgG1 heavy chain constant sequence of SEQ ID NO:235.

7. The isolated polynucleotide of claim 1, wherein:

a. the nucleotide sequence that encodes the light chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises a sequence of nucleotides that is at least 90% identical to the nucleotide sequence of SEQ ID NO:31; or

b. the nucleotide sequence that encodes the heavy chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises a sequence of nucleotides that is at least 90% identical to the nucleotide sequence of SEQ ID NO:135; or

c. said nucleotide sequence that encodes the light chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises a sequence of nucleotides that is at least 90% identical to the nucleotide sequence of SEQ ID NO:31 and said nucleotide sequence that encodes the heavy chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises a sequence of nucleotides that is at least 90% identical to the nucleotide sequence of SEQ ID NO:135.

8. The isolated polynucleotide of claim 7, wherein said nucleotide sequence that encodes the light chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises a sequence of nucleotides that is at least 95% identical to the nucleotide sequence of SEQ ID NO:31 and said nucleotide sequence that encodes the heavy chain variable region of said anti-IGF-1R antibody or antigen binding fragment comprises a sequence of nucleotides that is at least 95% identical to the nucleotide sequence of SEQ ID NO:135.

FIG. 1

L1 (SEQ ID NO:1)

```
       GAT GTTGTGATGA CTCAGTCTCC ACTCTCCCTG CCCGTCACCC CTGGAGAGCC GGCCTCCATC
TCCTGCAGGT CTAGTCAGAG CCTCCTGCAT AGTAGTGGAT ACAACTATTT GGATTGGTAC CTGCAGAAGC
CAGGGCAGTC TCCACAGCTC CTGATCTATT TGGGTTCTAA TCGGGCCTCC GGGGTCCCTG ACAGGTTCAG
TGGCAGTGGA TCAGGCACAG ATTTTACACT GAAAATCAGC AGAGTGGAGG CTGAGGATGT TGGGGTTTAT
TACTGCATGC AAGCTCTACA AACTCCGATC ACCTTCGGCC AAGGGACACG ACTGGAGATT AAA
```

L2 (SEQ ID NO:3)

```
       GAT GTTGTGATGA CTCAGTCTCC ACTCTCCCTG CCCGTCACCC CTGGAGAGCC GGCCTCCATC
TCCTGCAGGT CTAGTCAGAG CCTCCTGCAT AGTAATGGAT ACAACTATTT GGATTGGTAC CTGCAGAAGC
CAGGGCAGTC TCCACAGCTC CTGATCTATT TGGGTTCTAA TCGGGCCTCC GGGGTCCCTG ACAGGTTCAG
TGGCAGTGGA TCAGGCACAG ATTTTACACT GAAAATCAGC AGAGTGGAGG CTGAGGATGT TGGGGTTTAT
TACTGCATGC AAGCTCTACA AACTCCGATC ACCTTCGGCC AAGGGACACG ACTGGAGATT AAA
```

L3 (SEQ ID NO:5)

```
       GAT GTTGTGATGA CTCAGTCTCC ACTCTCCCTG CCCGTCACCC CTGGAGAGCC GGCCTCCATC
TCCTGCAGGT CTAGTCAGAG CCTCCTGCAT AGTAATGGAT ACAACTATTT GGATTGGTAC CTGCAGAAGC
CAGGGCAGTC TCCACAGCTC CTGATCTATT TGGGTTCTAA TCGGGCCTCC GGGGTCCCTG ACAGGTTCAG
TGGCAGTGGA TCAGGCACAG ATTTTACACT GAAAATCAGC AGAGTGGAGG CTGAGGATGT TGGGGTTTAT
TACTGCATGC AAGCTCTACA AACTCCACTC ACTTTCGGCG GCGGGACCAA GGTGGAGATC AAA
```

L4 (SEQ ID NO:7)

```
        GA AATTGTGATG ACGCAGTCTC CACTCTCCCT GCCCGTCACC CCTGGAGAGC CGGCCTCCAT
CTCCTGCAGG TCTAGTCAGA GCCTCCTGCA TAGTAATGGA TACAACTATT TGGATTGGTA CCTGCAGAAG
CCAGGGCAGT CTCCACAGCT CCTGATCTAT TTGGGTTCTA ATCGGGCCTC CGGGGTCCCT GACAGGTTCA
GTGGCAGTGG ATCAGGCACA GATTTTACAC TGAAAATCAG CAGAGTGGAG GCTGAGGATG TTGGGGTTTA
TTACTGCATG CAAGCTCTAC AAACTCCTCA CACTTTCGGC GGAGGGACCA AGGTGGAGAT CAAA
```

L5 (SEQ ID NO:9)

```
      GAAA TTGTGCTGAC TCAGTCTCCA CTCTCCCTGC CCGTCACCCC TGGAGAGCCG GCCTCCATCT
CCTGCAGGTC TAGTCAGAGC CTCCTGCATA GTAATGGATA CAACTATTTG GATTGGTACC TGCAGAAGCC
AGGGCAGTCT CCACAGCTCC TGATCTATTT GGGTTCTAAT CGGGCCTCCG GGGTCCCTGA CAGGTTCAGT
GGCAGTGGAT CAGGCACAGA TTTTACACTG AAAATCAGCA GAGTGGAGGC TGAGGATGTT GGGGTTTATT
ACTGCATGCA AGCTCTACAA ACCCCTCTCA CTTTCGGCCC TGGGACCAAA GTGGATATCA AA
```

L6 (SEQ ID NO:11)

```
       GAT GTTGTGATGA CTCAGTCTCC ACTCTCCCTG GCCGTCACCC CTGGAGAGCC GGCCTCCATC
TCCTGCAGGT CTAGTCAGAG CCTCCTGCAT AGTAATGGAT ACAACTATTT GGATTGGTAC CTGCAGAAGC
CAGGGCAGTC TCCACAGCTC CTGATCTATT TGGGTTCTAA TCGGGCCTCC GGGGTCCCTG ACAGGTTCAG
TGGCAGTGGA TCAGGCACAG ATTTTACACT GAAAATCAGC AGAGTGGAGG CTGAGGATGT TGGGGTTTAT
TACTGCATGC AAGCTCTACA AACTCCGCTC ACTTTCGGCG GAGGGACCAA GGTGGAGATC AAA
```

L7 (SEQ ID NO:13)

```
       GAT GTTGTGATGA CTCAGTCTCC ACTCTCCCTG CCCGTCACCC CTGGAGAGCC GGCCTCCATC
TCCTGCAGGT CTAGTCAGAG CCTCCTGCAT AGTAATGGAT ACAACTATTT GGATTGGTAC CTGCAGAAGC
CAGGGCAGTC TCCACAGCTC CTGATCTATT TGGGTTCTAA TCGGGCCTCC GGGGTCCCTG ACAGGTTCAG
TGGCAGTGGA TCAGGCACAG ATTTTACACT GAAAATCAGC AGAGTGGAGG CTGAGGATGT TGGGGTTTAT
TACTGCATGC AAGCTCTACA AACTCCTCTC ACTTTCGGCG GAGGGACCAA GGTGGAGATC AAA
```

L8 (SEQ ID NO:15)

```
 GATGTTGTG ATGACTCAGT CTCCAGTCTC CCTGCCCGTC ACCCCTGGAG AGCCGGCCTC CATCTCCTGC
AGGTCTAGTC AGAGCCTCCT GCATAGTAAT GGATACAACT ATTTGGATTG GTACCTGCAG AAGCCAGGGC
AGTCTCCACA GCTCCTGATC TATTTGGGTT CTAATCGGGC CTCCGGGGTC CCTGACAGGT TCAGTGGCAG
TGGATCAGGC ACAGATTTTA CACTGAAAAT CAGCAGAGTG GAGGCTGAAG ATGTTGGGGT TTATTACTGT
ATGCAAGCTC TACAAACCCC CCTCACTTTC GGCGGAGGGA CCAAGGTGGA GATCAAA
```

L9 (SEQ ID NO:17)

```
      GATG TTGTGATGAC TCAGTCTCCA CTCTCCCTGC CCGTCACCCC TGGAGAGCCG GCCTCCATCT
CCTGCAGGTC TAGTCAGAGC CTCCTGCATA GTAATGGATA CAACTATTTG GATTGGTACC TGCAGAAGCC
AGGGCAGTCT CCACAGCTCC TGATCTATTT GGGTTCTAAT CGGGCCTCCG GGGTCCCTGA CAGGTTCAGT
GGCAGTGGAT CAGGCACAGA TTTTACACTG AAAATCAGCA GAGTGGAGGC TGAGGATGTT GGGGTTTATT
ACTGCATGCA AGCTCTACAA ACTCCGTTCA CCTTCGGCCA AGGGACACGA CTGGAGATTA AA
```

AMGEN INC.
Applicant
SAPALO VELEZ BUNDANG & BULILAN

By:

Attorneys

FIG. 1E

L37 (SEQ ID NO:73)
GATGTTGTGA TGACTCAGTC TCCACTCTCC CTGCCCGTCA CCCCTGGGGA GCCGGCCTCC ATCTCCTGCA
GGTCTAGTCA GAGCCTCCTG CATAGTAATG GATACAACTA TTTGGATTGG TACCTGCAGA AGCCAGGGCA
GTCTCCACAG CTCCTGATCT ATTTGGGTTC TAACCGGGAC TCTGGGGTCC CAGACAGATT CAGCGGCAGT
GGGTCAGGCA CTGATTTCAC ACTGAAAATC AGCAGGGTGG AGGCTGAGGA TGTTGGGGTT TATTACTGCA
TGCAAGGTAC ACACTGGCCG TACACTTTTG GCCAGGGGAC CAGGCTGGAG ATCAAA

L38 (SEQ ID NO:75)
GATGTTGT GATGACTCAG TCTCCACTCT CCCTGCCCGT CACCCCTGGA GAGTCGGCCT CCATCTCCTG
CAGGTCTAGT CAGAGCCTCC TGCATAGTAA TGGATACAAC TTTTTGGATT GGTACCTGCA GAAGCCAGGG
CAGTCTCCAC AGCTCCTGAT CTATTTGGGT TCTAATCGGG CCTCCGGGGT CCCTGACAGG TTCAGTGGCA
GTGGATCAGG CACAGATTTT ACACTGAAAA TCAGCAGAGT GGAGGCTGAG GATGTTGGGG TTTATTACTG
CATGCAAGCT CTACAAACTC CTCTCACTTT CGGCGGAGGG ACCAAGGTGG AGATCAAA

L39 (SEQ ID NO:77)
GA TGTTGTGATG ACTCAGTCTC CACTCTCCCT GCCCGTCACC CCTGGAGAGC CGGCCTCCAT
CTCCTGCAGG TCTAGTCAGA GCCTCCTGCA TAGTAATGGA TACAACTATT TGGATTGGTA CCTGCAGAAG
CCAGGGCAGT CTCCACAGCT CCTGATCTAT TTGGGTTCTA ATCGGGCCTC CGGGGTCCCT GACAGGTTCA
GTGGCAGTGG ATCAGGCACA GATTTTACAC TGAAAATCAG CAGAGTGGAG GCTGAGGATG TTGGGGTTTA
TTACTGCATG CAAGCTCTAC AAACCCCCCT CACTTTCGGC GGAGGGACCA AGGTGGAGAT CAAA

L40 (SEQ ID NO:79)
GAAACGAC ACTCACGCAG TCTCCAGCCA CCCTGTCTTT GTCTCCAGGG CAAAGAGCCA CCCTCTCCTG
CAGGGCCAGT CAGAGTGTCT ACAACTACTT AGCCTGGTAC CAACAGAAGC CTGGCCAGGC TCCCAGGCTC
CTCATCTATG ATGCATCCAG AAGGGCAACT GGCATCCCAG CCAGGTTCAG TGGCAGTGGG TCTGGGACAG
ACTTCACTCT CACCATCAGC AGCCTAGAGC CTGAAGATTT TGCAGTTTAT TACTGTCAGC AGCGTAACAA
CTGGCCGCTC ACTTTCGGTG GAGGGACCAA GGTGGAGATC AAA

L41 (SEQ ID NO:81)
GACAT CCAGTTGACC CAGTCTCCAT CCTCCCTGTC TGCTTCTGTT GGAGACAGCG TCACCATCTC
TTGCCGGGCA AGTCAGAGTC CTGGCATCTT TTTAAATTGG TATCAGCAGA TACCAGGGAA AGCCCCTAAA
CTCCTGATCT ACGCTACATC CACTCTGGAA AGTGGGGTCC CCCCCAGGTT CACCGGCAGT GGATCTGGGA
CAGATTTCAC TCTCACCATC AGCAGTCTGC AACCTGAGGA CTTTGCAACT TACTACTGTC AACAGAGTAA
CAGTGTTCCG CTCACTTTCG GCGGCGGGAC CAAGGTGGAG ATCAAA

L42 (SEQ ID NO:83)
GATGT TGTGATGACT CAGTCTCCAC TCTCCCTGCC CGTCACCCCT GGAGAGCCGG CCTCCATCTC
CTGCAGGTCT AGTCAGAGCC TCCTGCATAG TAATGGATAC AACTATTTGG ATTGGTACCT GCAGAAGCCA
GGGCAGTCTC CACAGCTCCT GATCTATTTG GGTTCTAATC GGGCCTCCGG GGTCCCTGAC AGGTTCAGTG
GCAGTGGATC AGGCACAGAT TTTACACTAA AAATCAGCAG AGTGGAGGCT GAGGATGTTG GGGTTTATTA
CTGCATGCAA GCTCTACAAA CTCCTCTAAC CTTCGGCCAA GGGACACGAC TGGAGATTAA A

L43 (SEQ ID NO:85)
GAAATT GTGATGACGC AGTCTCCAGC CACCCTGTCT GTGTCTCCAG GGGAAAGAGC CACCTTCTCC
TGTAGGGCCA GTCAGAGTGT TGGCAGCAAC TTAGCCTGGT ACCAGCAGAA ACCTGGCCAG GCTCCCAGGC
TCCTCATCTA TGATGCATCC AACAGGGCCA CTGGCATCCC AGCCAGGTTC AGTGGCAGTG GGTCTGGGAC
AGACTTCACT CTCACCATCA GCAGACTGGA GCCTGAAGAT TTTGCAGTGT ATTACTGTCA GCAGCGTAGC
AACTGGCCCC TCACTTTCGG CGGAGGGACC AAGGTGGAGA TCAAA

L44 (SEQ ID NO:87)
GATGT TGTGATGACT CAGTCTCCAC TCTCCCTGCC CGTCACCCCT GGAGAGCCGG CCTCCATCTC
CTGCAGGTCT AGTCAGAGCC TCCTGCATAG TAATGGATAC AACTATTTGG ATTGGTACCT GCAGAAGCCA
GGGCAGTCTC CACAGCTCCT GATCTATTTG GGTTCTAATC GGGCCTCCGG GGTCCCTGAC AGGTTCAGTG
GCAGTGGATC AGGCACAGAT TTTACACTGA AAATCAGCAG AGTGGAGGCT GAGGATGTTG GGGTTTATTA
CTGCATGCAA GCTCTACAAA CTCCGCTCAC TTTCGGCGGA GGGACCAAGG TGGAGATCAA A


L45 (SEQ ID NO:89)
GAT GTTGTGATGA CTCAGTCTCC ACTCTCCCTG CCCGTCACCC CTGGAGAGCC GGCCTCCATC
TCCTGCAGGT CTAGTCAGAG CCTCCTGCAT AGTAATGGAT ACAACTATTT GGATTGGTAC CTGCAGAAGC
CAGGGCAGTC TCCACAGCTC CTGATCTACT TGGGTTCTAC TCGGGCCTCC GGCGTCCCTG ACAGGTTCAG
TGGCAGTGGA TCAGGCACAG ATTTTACACT GAAAATCAGC AGAGTGGAGG CTGAGGATGT TGGGGTTTAT
TACTGCATGC AAGCTCTACA AACTCCTTAC ACTTTCGGCG GAGGGACCAA GGTGGAGATC AAA

AMGEN INC.
Applicant
SAPALO VELEZ BUNDANG & BULILAN

By:

Attorneys

FIG. 2B



AMGEN INC.
Applicant
SAPALO VELEZ BUNDANG & BULILAN
Attorneys

By:

FIG. 11

```
   1 MGTGGRRGAA AAPLLVAVAA LLLGAAGHLY PGEVCPGMDI RNNLTRLHEL ENCSVIEGHL
  61 QILLMFKTRP EDFRDLSFPK LIMITDYLLL FRVYGLESLK DLFPNLTVIR GSRLFFNYAL
 121 VIFEMVHLKE LGLYNLMNIT RGSVRIEKNN ELCYLATIDW SRILDSVEDN HIVLNKDDNE
 181 ECGDICPGTA KGKTNCPATV INGQFVERCW THSHCQKVCP TICKSHGCTA EGLCCHSECL
 241 GNCSQPDDPT KCVACRNFYL DGRCVETCPP PYYHFQDWRC VNFSFCQDLH HKCKNSRRQG
 301 CHQYVIHNNK CIPECPSGYT MNSSNLLCTP CLGPCPKVCH LLEGEKTIDS VTSAQELRGC
 361 TVINGSLIIN IRGGNNLAAE LEANLGLIEE ISGYLKIRRS YALVSLSFFR KLRLIRGETL
 421 EIGNYSFYAL DNQNLRQLWD WSKHNLTTTQ GKLFFHYNPK LCLSEIHKME EVSGTKGRQE
 481 RNDIALKTNG DKASCENELL KFSYIRTSFD KILLRWEPYW PPDFRDLLGF MLFYKEAPYQ
 541 NVTEFDGQDA CGSNSWTVVD IDPPLRSNDP KSQNHPGWLM RGLKPWTQYA IFVKTLVTFS
 601 DERRTYGAKS DIIYVQTDAT NPSVPLDPIS VSNSSSQIIL KWKPPSDPNG NITHYLVFWE
 661 RQAEDSELFE LDYCLKGLKL PSRTWSPPFE SEDSQKHNQS EYEDSAGECC SCPKTDSQIL
 721 KELEESSFRK TFEDYLHNVV FVPRKTSSGT GAEDPRPSRK RRSLGDVGNV TVAVPTVAAF
 781 PNTSSTSVPT SPEEHRPFEK VVNKESLVIS GLRHFTGYRI ELQACNQDTP EERCSVAAYV
 841 SARTMPEAKA DDIVGPVTHE IFENNVVHLM WQEPKEPNGL IVLYEVSYRR YGDEELHLCV
 901 SRKHFALERG CRLRGLSPGN YSVRIRATSL AGNGSWTEPT YFYVTDYLDV PSNIAKVDGC
 961 KPCICTVPEV SSVFIFPPKP KDVLTITLTP KVTCVVVDIS KDDPEVQFSW FVDDVEVHTA
1021 QTQPREEQFN STFRSVSELP IMHQDWLNGK EFKCRVNSAA FPAPIEKTIS KTKGRPKAPQ
1081 VYTIPPPKEQ MAKDKVSLTC MITDFFPEDI TVEWQWNGQP AENYKNTQPI MDTDGSYFVY
1141 SKLNVQKSNW EAGNTFTCSV LHEGLHNHHT EKSLSHSPGK
```

AMGEN INC.
Applicant
SAPALO VELEZ BUNDANG & BULILAN

By:

Attorneys