

(19) **DANMARK**



Patent- og  
Varemærkestyrelsen

(10) **DK/EP 3673084 T3**

(12) **Oversættelse af  
europæisk patentskrift**

- 
- (51) Int.Cl.: **C 12 Q 1/6844 (2018.01)** **C 12 Q 1/686 (2018.01)**
- (45) Oversættelsen bekendtgjort den: **2021-11-08**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2021-08-11**
- (86) Europæisk ansøgning nr.: **19707096.4**
- (86) Europæisk indleveringsdag: **2019-02-19**
- (87) Den europæiske ansøgnings publiceringsdag: **2020-07-01**
- (86) International ansøgning nr.: **GB2019050443**
- (87) Internationalt publikationsnr.: **WO2019162657**
- (30) Prioritet: **2018-02-20 GB 201802744**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Longas Technologies Pty Ltd, Level 3 , 56 Pitt Street, Sydney NSW 2000, Australien**
- (72) Opfinder: **MONAHAN, Leigh G, c/o Longas Technologies Pty Ltd, Level 3, 56 Pitt Street, Sydney, New South Wales 2000, Australien**  
**TO, Joyce, c/o Longas Technologies Pty Ltd, Level 3, 56 Pitt Street, Sydney, New South Wales 2000, Australien**  
**BURKE, Catherine M., c/o Longas Technologies Pty Ltd, Level 3, 56 Pitt Street, Sydney, New South Wales 2000, Australien**  
**IMELFORT, Michael, c/o Longas Technologies Pty Ltd, Level 3, 56 Pitt Street, Sydney, New South Wales 2000, Australien**  
**DARLING, Aaron E, c/o Longas Technologies Pty Ltd, Level 3, 56 Pitt Street, Sydney, New South Wales 2000, Australien**
- (74) Fuldmægtig i Danmark: **Marks & Clerk (Luxembourg) LLP, 44 rue de la Vallée, B.P. 1775, L-1017 Luxembourg, Luxembourg**
- (54) Benævnelse: **FREMANGSMÅDE TIL INDFØRING AF MUTATIONER**
- (56) Fremdragne publikationer:  
**JUNG-SUK CHOI ET AL: "The use of modified and non-natural nucleotides provide unique insights into pro-mutagenic replication catalyzed by polymerase eta", NUCLEIC ACIDS RESEARCH, vol. 44, no. 3, 18 February 2016 (2016-02-18), pages 1022-1035, XP055462517, ISSN: 0305-1048, DOI: 10.1093/nar/gkv1509**  
**MASAYASU KUWAHARA ET AL: "Study on Suitability of KOD DNA Polymerase for Enzymatic Production of Artificial Nucleic Acids Using Base/Sugar Modified Nucleoside Triphosphates", MOLECULES, vol. 15, no. 11, 12 November 2010 (2010-11-12), pages 8229-8240, XP055078441, DOI: 10.3390/molecules15118229**  
**PETRIE KATHERINE L ET AL: "Deep sequencing analysis of mutations resulting from the incorporation of dNTP analogs", NUCLEIC ACIDS RESEARCH, vol. 38, no. 22, December 2010 (2010-12), pages 8095-8104, XP002791491,**  
**SHEN BINZHANG ED - DUCRY ET AL: "PCR approaches to DNA mutagenesis and recombination: An**

Fortsættes ...

overview", DNA REPAIR PROTOCOLS. METHODS IN MOLECULAR BIOLOGY, HUMANA PRESS, US, vol. 192, 1 January 2002 (2002-01-01), pages 167-174, XP008093989, ISBN: 978-1-62703-541-5

ASHRAF M. ELSHAWADFY ET AL: "DNA polymerase hybrids derived from the family-B enzymes of *Pyrococcus furiosus* and *Thermococcus kodakarensis*: improving performance in the polymerase chain reaction", FRONTIERS IN MICROBIOLOGY, vol. 5, 27 May 2014 (2014-05-27), XP055200889, DOI: 10.3389/fmicb.2014.00224

YAMASHITA MAMI ET AL: "Characterization of Recombinant *Thermococcus kodakaraensis*(KOD) DNA Polymerases Produced Using Silkworm-Baculovirus Expression Vector System", MOLECULAR BIOTECHNOLOGY, HUMANA PRESS, BOSTON, vol. 59, no. 6, 8 May 2017 (2017-05-08), pages 221-233, XP036243650, ISSN: 1073-6085, DOI: 10.1007/S12033-017-0008-9 [retrieved on 2017-05-08]

# DESCRIPTION

## Field of the invention

**[0001]** This invention relates to a method for introducing mutations into one or more nucleic acid molecule(s), a use of a low bias DNA polymerase in a method for introducing mutations into one or more nucleic acid molecule(s).

## Background of the invention

**[0002]** DNA polymerases can be used to introduce mutations into nucleic acid sequences. This can be useful in multiple applications. For example mutagenesis techniques can be useful in applications including sequencing assisted by mutagenesis (SAM) techniques and for introducing mutations into protein sequences to find mutations that affect the activity of the protein.

**[0003]** Mutations may be introduced using DNA polymerases that have low fidelity. Low fidelity DNA polymerases make mistakes during replication that result in the introduction of mutations. However, many low fidelity DNA polymerases only introduce mutations at a rate of less than 2% per mutation reaction (round of replication), and for some applications higher mutagenesis rates are useful. In addition, low fidelity DNA polymerases may introduce mutations in a biased manner. Such DNA polymerases can be referred to as high bias DNA polymerases.

**[0004]** Mutations may be introduced by replicating sequences, using DNA polymerases, in the presence of nucleotide analogs such as dPTP. DNA polymerases may incorporate the nucleotide analogs in place of a natural nucleotide. Then, in a subsequent cycle of replication, the nucleotide analog can pair with a natural nucleotide that was not present in the original sequence, thereby introducing a mutation. Introducing mutations by replicating sequences in the presence of nucleotide analogs can be used to achieve higher mutations rates.

**[0005]** Commonly used DNA polymerases (such as Taq polymerase) can be used to incorporate nucleotide analogs in place of a natural nucleotide. However, these polymerases are high bias polymerases. High bias DNA polymerases may display two possible biases: mutation bias and template amplification bias.

**[0006]** Some high bias polymerases have high mutation bias, as they do not mutate all four natural nucleotides (adenine, cytosine, guanine and thymine) uniformly at random. For example, high bias DNA polymerases may mutate some nucleotides with a greater frequency than others. Adenine/thymine pairs are connected by two hydrogen bonds, whereas guanine/cytosine pairs are connected by three hydrogen bonds. Thus, it is possible that high bias DNA polymerases are more likely to introduce mutations into adenine/thymine pairs than

guanine/cytosine pairs.

**[0007]** High bias polymerases, having high mutation bias, may fail to incorporate nucleotide analogs randomly. For example, high bias polymerases may favour replacing certain bases with nucleotide analogs. dPTP can interconvert between two different tautomeric forms, an imino form and an amino form. The imino tautomer can form Watson-Crick base pairs with adenine, whilst the amino form can form Watson-Crick base pairs with guanine (Kong Thoo Lin P, Brown D M (1989). "Synthesis and duplex stability of oligonucleotides containing cytosine-thymine analogues ". *Nucleic Acids Research*. 17: 10373-10383; Stone M J et al. (1991). "Molecular basis for methoxyamine-initiated mutagenesis: 1H nuclear magnetic resonance studies of base-modified oligodeoxynucleotides. " *Journal of Molecular Biology*. 222: 711-723; Nedderman A N R et al. (1993). "Molecular basis for methoxyamine initiated mutagenesis: 1H nuclear magnetic resonance studies of oligonucleotide duplexes containing base-modified cytosine residues". *Journal of Molecular Biology*. 230: 1068-1076; Moore M H et al. (1995). "Direct observation of two base-pairing modes of a cytosine-thymine analogue with guanine in a DNA-form duplex. Significance for base analogue mutagenesis". *Journal of Molecular Biology*. 251: 665-673). This effectively means that replication in the presence of dPTPs can be used to introduce substitutions in place of adenine, cytosine, guanine or thymine in a nucleotide sequence. However, in aqueous solution, the ratio of the imino to amino forms of dPTP has been shown to be around 10:1 (Harris V H et al. (2003). "The effect of tautomeric constant on the specificity of nucleotide incorporation during DNA replication: support for the rare tautomer hypothesis of substitution mutagenesis ". *Journal of Molecular Biology*. 326: 1389-1401). Accordingly, when a polymerase such as Taq polymerase is used to introduce mutations using dPTP, it introduces substitutions of adenine and thymine much more frequently than substitutions of guanine and cytosine (Zaccolo M et al. (1996). "An approach to random mutagenesis of DNA using mixtures of triphosphate derivatives of nucleoside analogues". *Journal of Molecular Biology*. 255: 589-603; Harris V H et al. (2003). "The effect of tautomeric constant on the specificity of nucleotide incorporation during DNA replication: support for the rare tautomer hypothesis of substitution mutagenesis". *Journal of Molecular Biology*. 326: 1389-1401).

**[0008]** Secondly, high bias polymerases may demonstrate template amplification bias, *i.e.* they may replicate some template nucleic acid molecules with a higher success rate per PCR cycle than others. Over many cycles of PCR this bias can create extreme differences in copy number amongst templates. Regions of a template nucleic acid molecule may form secondary structures or may contain a higher proportion of some nucleotides (for example guanine or cytosine nucleotides) than others. A high bias polymerase may be more effective to amplify, for example, guanine and cytosine rich template nucleic acid molecules compared to adenine and thymine rich template nucleic acid molecules, or may be more effective to amplify template nucleic acid molecules that do not form secondary structures.

**[0009]** Many of the applications of mutagenesis are more effective if mutagenesis can be performed with low bias (both mutation bias and template amplification).

**[0010]** The accurate assembly of genome sequences has proven difficult as many second generation sequencing platforms are only capable of sequencing short nucleic acid fragments, and require the target nucleic acid sequences to be amplified during the sequencing process in order to provide sufficient nucleic acid molecules for the sequencing step. If the user desires to sequence a larger nucleic acid sequence, this can be achieved by sequencing regions of the target nucleic acid molecules. The user must then computationally assemble the sequence of the full nucleic acid sequence from the sequences of the regions.

**[0011]** Assembling a nucleic acid sequence using sequences of regions can be difficult. In particular, where long regions of the sequences are very similar to one another it may be difficult to determine whether sequences of two regions are both sequences of replicates of the same original template nucleic acid molecule or correspond to sequences from two different original template nucleic acid molecules. Similarly, it may be difficult to determine whether sequences of two regions correspond to sequences of replicates of the same portion of a template nucleic acid molecule, or actually correspond to two different repeats within the template nucleic acid molecule. These difficulties can be circumvented by introducing mutations into the target nucleic acid molecules prior to amplification. The user may then identify that fragments having the same mutation patterns are likely to have originated from the same portion of the same original template nucleic acid molecule. This type of sequencing method is sometimes referred to as sequencing aided by mutagenesis (SAM).

**[0012]** Shen et al, DNA REPAIR PROTOCOLS. METHODS IN MOLECULAR BIOLOGY, HUMANA PRESS, US, (20020101), vol. 192, ISBN 978-1-62703-541-5, pages 167 - 174 describes PCR approaches to DNA mutagenesis and recombination. Choi et al, NUCLEIC ACIDS RESEARCH, (20160218), vol. 44, no. 3, doi:10.1093/nar/gkv1509, ISSN 0305-1048, pages 1022 - 1035 describes that the use of modified and non-natural nucleotides provide unique insights into pro-mutagenic replication catalyzed by polymerase eta. Masayasu et al, MOLECULES, (20101112), vol. 15, no. 11, doi:10.3390/molecules15118229, pages 8229 - 8240 describes a study on suitability of KOD DNA Polymerase for enzymatic production of artificial nucleic acids using base/sugar modified nucleoside triphosphates. Petrie et al, NUCLEIC ACIDS RESEARCH, (201012), vol. 38, no. 22, pages 8095 - 8104 describes deep sequencing analysis of mutations resulting from the incorporation of dNTP analogs. Elshawadfy et al FRONTIERS IN MICROBIOLOGY, (20140527), vol. 5 is directed to DNA polymerase hybrids derived from the family-B enzymes of *Pyrococcus furiosus* and *Thermococcus kodakarensis*: improving performance in the polymerase chain reaction. Yamashita et al, MOLECULAR BIOTECHNOLOGY, HUMANA PRESS, BOSTON, vol. 59, no. 6, doi:10.1007/S12033-017-0008-9, ISSN 1073-6085, (20170508), pages 221 - 233 is directed to the characterization of recombinant *Thermococcus kodakaraensis* (KOD) DNA polymerases produced using silkworm-baculovirus expression vector system.

### **Summary of the Invention**

**[0013]** The sequencing methods described above are more effective when the mutations that

are introduced into the target nucleic acid molecules are uniformly random. If the mutations are uniformly random, then the likelihood, for example, that any given portion of a template nucleic acid molecule would have a unique mutation pattern is higher. Thus, there is a need for the identification of DNA polymerases that are able to introduce mutations uniformly at random (have low mutation bias).

**[0014]** In addition, sequencing methods using DNA polymerases having high template amplification bias may be limited. DNA polymerases having high template amplification bias will replicate and/or mutate some target nucleic acid molecules better than others, and so a sequencing method that uses such a high bias DNA polymerase may not be able to sequence some target nucleic acid molecules well.

**[0015]** The present inventors have identified polymerases that are low bias polymerases (have both low template amplification bias and low mutation bias), and so are particularly useful in a method for introducing mutations into at least one target nucleic acid molecule.

**[0016]** The user may wish to use the methods of the invention on more than one sample at once. In such cases, it would be advantageous for the user to be able to identify which target nucleic acid molecule came from which original sample. Such identification could be achieved by labelling the target nucleic acid molecules with sample tags. However, the sample tags may, themselves, be mutated during the method and so the present inventors have determined how to design sample tags that can be distinguished from one another even if they are mutated.

**[0017]** The user may also wish to ensure that the methods of the invention are used to mutate and amplify long target nucleic acid molecules in preference compared to short nucleic acid molecules. The present inventors have found that this can be achieved by introducing special primer binding sites into each end of the target nucleic acid molecules.

**[0018]** Thus, in a first aspect of the invention, there is provided a method for introducing substitution mutations into at least one target DNA molecule comprising:

1. a. providing at least one sample comprising at least one target DNA molecule; and
2. b. amplifying the at least one target DNA molecule using a low bias high fidelity DNA polymerase having low template amplification bias;

wherein the step of amplifying the at least one target DNA molecule is carried out in the presence of a nucleotide analog and comprises at least 2 rounds of replicating the at least one target DNA molecule, wherein in a first round of replication the DNA polymerase incorporates the nucleotide analog in place of a nucleotide and in a second round of replication, the nucleotide analog pairs with a natural nucleotide to introduce a substitution mutation in the complementary strand.

**[0019]** In a second aspect of the invention, there is provided a use of a low bias high fidelity DNA polymerase having low template amplification bias in a method for introducing mutations into at least one target DNA molecule, wherein the method comprises:

1. a. providing at least one sample comprising at least one target DNA molecule; and
2. b. amplifying the at least one target DNA molecule using the DNA polymerase;

wherein the step of amplifying the at least one target DNA molecule is carried out in the presence of a nucleotide analog and comprises at least 2 rounds of replicating the at least one target DNA molecule, wherein in a first round of replication the DNA polymerase incorporates the nucleotide analog in place of a nucleotide and in a second round of replication, the nucleotide analog pairs with a natural nucleotide to introduce a substitution mutation in the complementary strand.

**[0020]** In a third aspect of the invention, there is provided a method for determining a sequence of at least one target DNA molecule comprising steps of:

1. a. performing the methods of the invention to provide at least one mutated target DNA molecule;
2. b. optionally fragmenting and/or amplifying the at least one mutated target DNA molecule to provide at least one fragmented and/or amplified mutated target DNA molecule;
3. c. sequencing regions of the at least one mutated target DNA molecule of the at least one fragmented and/or amplified mutated target DNA molecule to provide mutated sequence reads; and
4. d. assembling a sequence for at least a portion of the at least one target DNA molecule using the mutated sequence reads.

**[0021]** In a fourth aspect of the invention, there is provided a method for engineering a protein comprising steps of:

1. a. performing the methods of the invention to provide at least one mutated target DNA molecule;
2. b. inserting the at least one mutated target DNA molecule into a vector; and
3. c. expressing a protein encoded by the at least one mutated target DNA molecule; optionally wherein:
  1. i) the method comprises steps of:
    1. a. amplifying the at least one target DNA molecule comprising the nucleotide analog in the absence of the nucleotide analog to provide at least one mutated target DNA molecule;
    2. b. inserting the at least one mutated target DNA molecule into a vector; and
    3. c. expressing a protein encoded by the at least one mutated target DNA molecule; and/or
  2. ii) the method further comprises a step of testing the activity or assessing the structure of the protein encoded by the at least one mutated target DNA molecule; and/or
  3. iii) the vector is a plasmid, a virus, a cosmid, or an artificial chromosome; and/or

4. iv) the step of expressing a protein encoded by the at least one mutated target DNA molecule is achieved by transforming bacterial cells, transfecting eukaryotic cells or transducing eukaryotic cells with the vector.

### **Brief description of the Figures**

#### **[0022]**

Figure 1 shows the level of mutation achieved with three different polymerases in the presence or absence of dPTP. Panel A shows data obtained using Taq (Jena Biosciences), panel B shows data obtained using LongAmp (New England Biolabs) and panel C shows data using Primestar GXL (Takara). The dark grey bars show the results obtained in the absence of dPTP and the pale grey bars show the results obtained in the presence of 0.5 mM dPTP.

Figure 2 describes the mutation rates obtained by dPTP mutagenesis using a *Thermococcus* polymerase (Primestar GXL; Takara) on templates with diverse G+C content. The median observed rate of mutations was ~7% for low GC templates from *S. aureus* (33% GC), while the median for other templates was about 8%.

Figure 3 is a sequence listing.

Figure 4 depicts self annealing of nucleic acid molecules when a first primer binding site and a second primer binding site that anneal to one another are used.

Figure 6 depicts the sizes of target nucleic acid molecules amplified using adapters that anneal to one another (right line) or using standard adapters (left line).

Figure 7 provides a pictorial representation of mutation using the nucleotide analog dPTP (referred to as "P" in Figure 7).

### **Detailed Description of the Invention**

#### ***General definitions***

**[0023]** Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by a person skilled in the art to which this invention belongs.

**[0024]** In general, the term "*comprising*" is intended to mean including, but not limited to. For

example, the phrase "*a method for introducing mutations into at least one target nucleic acid molecule comprising*" certain steps should be interpreted to mean that the method includes the recited steps, but that additional steps may be performed.

**[0025]** In some embodiments of the invention, the word "*comprising*" is replaced with the phrase "*consisting of*". The term "*consisting of*" is intended to be limiting. For example, the phrase "*a method for introducing mutations into at least one target nucleic acid molecule consisting of*" certain steps should be understood to mean that the method includes the recited steps, and that no additional steps are performed.

**[0026]** For the purpose of this invention, in order to determine the percent identity of two sequences (such as two polynucleotide sequences), the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in a first sequence for optimal alignment with a second sequence). The nucleotide or amino acid residues at each of the positions are then compared. When a position in the first sequence is occupied by the same residue as the corresponding position in the second sequence, then the residues are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity = number of identical positions/total number of positions x 100). Typically the sequence comparison is carried out over the length of the reference sequence. For example, to assess whether a test sequence is at least 95% identical to SEQ ID NO. 2 (the reference sequence), the skilled person would carry out an alignment over the length of SEQ ID NO. 2, and identify how many positions in the test sequence were identical to those of SEQ ID NO. 2. If at least 80% of the positions are identical, the test sequence is at least 80% identical to SEQ ID NO. 2. If the sequence is shorter than SEQ ID NO. 2, the gaps should be considered to be non-identical positions.

**[0027]** The skilled person is aware of different computer programs that are available to determine the homology or identity between two sequences. For instance, a comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In an embodiment, the percent identity between two amino acid or nucleic acid sequences is determined using the Needleman and Wunsch (1970) algorithm which has been incorporated into the GAP program in the Accelrys GCG software package (available at <http://www.accelrys.com/products/gcg/>), using either a Blosum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

***A method for introducing mutations into at least one target nucleic acid molecule***

**[0028]** In one aspect, the invention provides a method for introducing mutations into at least one target nucleic acid molecule. In a further aspect, the invention provides a use of a low bias DNA polymerase in a method for introducing mutations into at least one target nucleic acid molecule.

[0029] The mutations may be substitution mutations, insertion mutations or deletion mutations. For the purposes of the present invention, the term "*substitution mutation*" should be interpreted to mean that a nucleotide is replaced with a different nucleotide. For example, the conversion of the sequence ATCC to the sequence AGCC is a substitution mutation. For the purposes of the present invention, the term "*insertion mutation*" should be interpreted to mean that at least one nucleotide is added to a sequence. For example, conversion of the sequence ATCC to the sequence ATTCC is an example of an insertion mutation (with an additional T nucleotide being inserted). For the purposes of the present invention, the term "*deletion mutation*" should be interpreted to mean that at least one nucleotide is removed from a sequence. For example, conversion of the sequence ATTCC to ATCC is an example of a deletion mutation (with a T nucleotide being removed). Preferably the mutations are substitution mutations.

[0030] For the purposes of the present invention, a "*nucleic acid molecule*" refers to a polymeric form of nucleotides of any length. The nucleotides may be deoxyribonucleotides, ribonucleotides or analogs thereof. In the methods and uses of the invention, the target nucleic acid molecule is made up of deoxyribonucleotides, *i.e.* the target nucleic acid molecule is a DNA molecule. Thus, for the purpose of the present invention, references to "*target nucleic acid molecule*" can be considered to refer to a "*target DNA molecule*".

[0031] The at least one "*target nucleic acid molecule*" can be any nucleic acid molecule into which the user of the method would like to introduce mutations. The target nucleic acid molecule may form part of a larger nucleic acid molecule such as a chromosome. The target nucleic acid molecule may comprise a gene, multiple genes or a fragment of a gene. The target nucleic acid molecule may be greater than 1 kbp, greater than 1.5 kbp, greater than 2 kbp, greater than 4 kbp, greater than 5 kbp, greater than 7 kbp, greater than 8 kbp, between 1 kbp and 50 kbp, or between 1 kbp and 20 kbp in size.

[0032] The term "*at least one target nucleic acid molecule*" is considered to be interchangeable with the term "*at least one target nucleic acid molecules*".

[0033] The "*at least one target nucleic acid molecule*" can be single stranded, or may be part of a double stranded complex. The at least one target nucleic acid molecule is made up of deoxyribonucleotides, and so it may form part of a double stranded DNA complex. In which case, one strand (for example the coding strand) will be considered to be the at least one target nucleic acid molecule, and the other strand is a nucleic acid molecule that is complementary to the at least one target nucleic acid molecule.

[0034] The method for introducing mutations into at least one target nucleic acid molecule may comprise:

1. a. providing at least one sample comprising at least one target nucleic acid molecule;  
and
2. b. amplifying the at least one target nucleic acid molecule using a low bias DNA

polymerase.

***Providing at least one sample comprising at least one target nucleic acid molecule***

[0035] The method for introducing mutations into at least one target nucleic acid molecule may comprise a step of providing at least one sample comprising at least one target nucleic acid molecule.

[0036] The at least one sample may comprise any sample that comprises at least one target nucleic acid molecule. The at least one sample may be obtained from any source. For example, the at least one sample may comprise a sample of nucleic acids derived from a human, for example a sample extracted from a skin swab of a human patient.

[0037] Alternatively, the at least one sample may be derived from other sources such as a sample from a water supply. Such a sample could contain billions of template nucleic acid molecules. It would be possible to mutate each of these billions of target nucleic acid molecules simultaneously using the methods of the invention, and so there is no upper limit on the number of target nucleic acid molecules which could be used in the methods of the invention.

[0038] In an embodiment, step a. comprises providing more than one sample. For example, step a. may comprise providing 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 15, 20, 25, 50, 75, or 100 samples. Optionally, step a. comprises providing less than 2000, less than 1000, less than 750, or less than 500 samples. In a further embodiment, step a. comprises providing between 2 and 100, between 2 and 75, between 2 and 50, between 2 and 25, between 5 and 15, or between 7 and 15 samples.

***Amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase***

[0039] The methods of the invention comprise amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase.

[0040] Amplifying the at least one target nucleic acid molecule refers to replicating the at least one target nucleic acid molecule to provide at least one nucleic acid molecule that is complementary to the at least one target nucleic acid molecule and/or replicates of the at least one target nucleic acid molecule. Amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase, increases the number of replicates of the at least one target nucleic acid molecule, and introduces mutations into the at least one target nucleic acid molecule. Since mutations are introduced, the replicates are not necessarily identical to the original at least one target nucleic acid molecule. The original at least one target nucleic acid

molecule and the replicates of the at least one target nucleic acid molecule may be referred to collectively as "*at least one mutated target nucleic acid molecule*".

**[0041]** For example, amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase may comprise incubating the sample comprising the at least one target nucleic acid molecule with the low bias DNA polymerase and suitable primers under conditions suitable for the low bias DNA polymerase to catalyse the generation of replicates of the at least one target nucleic acid molecule.

**[0042]** Suitable primers comprise short nucleic acid molecules complementary to regions flanking the at least one target nucleic acid molecules or to regions flanking nucleic acid molecules that are complementary to the at least one target nucleic acid molecule. For example, if the target nucleic acid molecule is part of a chromosome, the primers may be complementary to regions of the chromosome immediately 3' to the 3' end of the target nucleic acid molecule and nucleic acid molecules complementary to regions immediately 5' to the 5' end of the target nucleic acid molecule, or the primers will be complementary to regions of the chromosome immediately 3' to the 3' end of a nucleic acid molecule complementary to the target nucleic acid molecule and nucleic acid molecules complementary to regions immediately 5' to the 5' end of a nucleic acid molecule complementary to the target nucleic acid molecule. Alternatively, the user may introduce primer binding sites (short nucleic acid sequences) into regions flanking the at least one target nucleic acid molecules. This is described in more detail in the section entitled "*barcodes, samples and adapters*".

**[0043]** Suitable conditions include a temperature at which the low bias DNA polymerase can catalyse the generation of replicates of the at least one target nucleic acid molecule. For example, a temperature of between 40°C and 90°C, between 50°C and 80°C, between 60°C and 70°C, or around 68°C may be used.

**[0044]** The step of amplifying the at least one target nucleic acid molecule comprises multiple rounds of replication. For example, the step of amplifying the at least one target nucleic acid molecule comprises:

1. i) a round of replicating the at least one target nucleic acid molecule to provide at least one nucleic acid molecule that is complementary to the at least one target nucleic acid molecule; and
2. ii) a round of replicating the at least one target nucleic acid molecule to provide replicates of the at least one target nucleic acid molecule.

**[0045]** Optionally, the step of amplifying the at least one target nucleic acid molecule comprises at least 4, at least 6, at least 8, or at least 10 rounds of replicating the at least one target nucleic acid molecule. Some of these rounds of replicating the at least one target nucleic acid molecule take place in the presence of nucleotide analogs. Optionally, the step of amplifying the at least one target nucleic acid molecule comprises at least 1, at least 2, at least

3, at least 4, at least 5, or at least 6 rounds of replication at a temperature between 60°C and 80°C.

[0046] Optionally, the step of amplifying the at least one target nucleic acid molecule is carried out using the polymerase chain reaction (PCR). PCR is a process that involves multiple rounds of the following steps for replicating a nucleic acid molecule:

1. a) melting;
2. b) annealing;
3. c) extension; and
4. d) elongation.

[0047] The nucleic acid molecule (such as the at least one target nucleic acid molecule) is mixed with suitable primers and a polymerase, such as a low bias DNA polymerase of the invention. In the melting step, the nucleic acid molecule is heated to a temperature above 90°C such that a double-stranded nucleic acid molecule will denature (separate into two strands). In the annealing step, the nucleic acid molecule is cooled to a temperature below 75°C, for example between 55°C and 70°C, around 55°C, or around 68°C to allow the primers to anneal to the nucleic acid molecule. In the extension step, the nucleic acid molecule is heated to a temperature greater than 60°C to allow the DNA polymerase to catalyse primer extension, the addition of nucleotides complementary to the template strand. In the elongation step, the nucleic acid molecule is heated to a temperature at which the DNA polymerase has high activity, such as a temperature between 60°C and 70°C, to catalyse addition of further complementary nucleic acids in order to complete the new nucleic acid strand.

[0048] Optionally, the method of the invention comprises multiple rounds of PCR using the low bias DNA polymerase.

#### ***The low bias DNA polymerase***

[0049] The methods of the invention comprise a step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase. As discussed below, the low bias DNA polymerase used in the methods and uses of the invention is a low bias high fidelity DNA polymerase, and so references to the "*low bias DNA polymerase*" can be considered to be references to a "*low bias high fidelity DNA polymerase*".

[0050] According to the present invention, a "*low bias DNA polymerase*" is a DNA polymerase that (a) exhibits low mutation bias, and/or (b) exhibits low template amplification bias. The low bias DNA polymerase used in the methods and uses of the invention has low template amplification bias.

#### ***Low mutation bias***

**[0051]** A low bias DNA polymerase that exhibits low mutation bias is a DNA polymerase that is able to mutate adenine and thymine, adenine and guanine, adenine and cytosine, thymine and guanine, thymine and cytosine, or guanine and cytosine at similar rates. In an embodiment, the low bias DNA polymerase is able to mutate adenine, thymine, guanine, and cytosine at similar rates.

**[0052]** Optionally, the low bias DNA polymerase is able to mutate adenine and thymine, adenine and guanine, adenine and cytosine, thymine and guanine, thymine and cytosine, or guanine and cytosine at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively. Preferably, the low bias DNA polymerase is able to mutate guanine and adenine at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively. Preferably, the low bias DNA polymerase is able to mutate thymine and cytosine at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively.

**[0053]** In such embodiments, in a step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase, the DNA polymerase mutates adenine and thymine, adenine and guanine, adenine and cytosine, thymine and guanine, thymine and cytosine, or guanine and cytosine nucleotides in the at least one target nucleic acid molecule at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively. Preferably, the low bias DNA polymerase mutates guanine and adenine nucleotides in the at least one target nucleic acid molecule at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively. Preferably, the low bias DNA polymerase mutates thymine and cytosine nucleotides in the at least one target nucleic acid molecule at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively.

**[0054]** Optionally, the low bias DNA polymerase is able to mutate adenine, thymine, guanine, and cytosine at a rate ratio of 0.5-1.5:0.5-1.5:0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4:0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2:0.8-1.2:0.8-1.2, or around 1:1:1:1 respectively. Preferably, the low bias DNA polymerase is able to mutate adenine, thymine, guanine and cytosine at a rate ratio of 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3.

**[0055]** In such embodiments, in a step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase, the DNA polymerase may mutate adenine, thymine, guanine, and cytosine nucleotides in the at least one target nucleic acid molecule at a rate ratio of 0.5-1.5:0.5-1.5:0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4:0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2:0.8-1.2:0.8-1.2, or around 1:1:1:1 respectively. Preferably, the low bias DNA polymerase mutates adenine, thymine, guanine, and cytosine nucleotides in the at least one target nucleic acid molecule at a rate ratio of 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3.

**[0056]** The adenine, thymine, cytosine, and/or guanine may be substituted with another nucleotide. For example, if the low bias DNA polymerase is able to mutate adenine, amplifying the at least one target nucleic acid molecule in the presence of the low bias DNA polymerase may substitute at least one adenine nucleotide in the nucleic acid molecule with thymine, guanine, or cytosine. Similarly, if the low bias DNA polymerase is able to mutate thymine, amplifying the at least one target nucleic acid molecule in the presence of the low bias DNA polymerase may substitute at least one thymine nucleotide with adenine, guanine, or cytosine. If the low bias DNA polymerase is able to mutate guanine, amplifying the at least one target nucleotide in the presence of the low bias DNA polymerase may substitute at least one guanine nucleotide with thymine, adenine, or cytosine. If the low bias DNA polymerase is able to mutate cytosine, amplifying the at least one target nucleotide in the presence of the low bias DNA polymerase may substitute at least one cytosine nucleotide with thymine, guanine, or adenine.

**[0057]** The low bias DNA polymerase may not be able to substitute a nucleotide directly, but it may still be able to mutate that nucleotide by replacing the corresponding nucleotide on the complementary strand. For example, if the target nucleic acid molecule comprises thymine, there will be an adenine nucleotide present in the corresponding position of the at least one nucleic acid molecule that is complementary to the at least one target nucleic acid molecule. The low bias DNA polymerase may be able to replace the adenine nucleotide of the at least one nucleic acid molecule that is complementary to the at least one target nucleic acid molecule with a guanine and so, when the at least one nucleic acid molecule that is complementary to the at least one target nucleic acid molecule is replicated, this will result in a cytosine being present in the corresponding replicated at least one target nucleic acid molecule where there was originally a thymine (a thymine to cytosine substitution).

**[0058]** In an embodiment, the low bias DNA polymerase mutates between 1% and 15%, between 2% and 10%, or around 8% of the nucleotides in the at least one target nucleic acid. In such embodiments, the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase is carried out in such a way that between 1% and 15%, between 2% and 10%, or around 8% of the nucleotides in the at least one target nucleic acid are mutated. For example, if the user wishes to mutate around 8% of the nucleotides in the target nucleic acid molecule, and the low bias DNA polymerase mutates around 1% of the nucleotides per round of replication, the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase may comprise 8 rounds of replication.

**[0059]** In an embodiment, the low bias DNA polymerase is able to mutate between 0% and 3%, between 0% and 2%, between 0.1% and 5%, between 0.2% and 3%, or around 1.5% of the nucleotides in the at least one target nucleic acid molecule per round of replication. In an embodiment, the low bias DNA polymerase mutates between 0% and 3%, between 0% and 2%, between 0.1% and 5%, between 0.2% and 3%, or around 1.5% of the nucleotides in the at least one target nucleic acid molecule per round of replication. The actual amount of mutation that takes place each round may vary, but may average to between 0% and 3%, between 0% and 2%, between 0.1% and 5%, between 0.2% and 3%, or around 1.5%.

***Whether a DNA polymerase is able to mutate a nucleotide and, if so, at what rate***

**[0060]** Whether the low bias DNA polymerase is able to mutate a certain percentage of the nucleotides in the at least one target nucleic acid molecule per round of replication can be determined by amplifying a nucleic acid molecule of known sequence in the presence of the low bias DNA polymerase for a set number of rounds of replication. The resulting amplified nucleic acid molecule can then be sequenced, and the percentage of nucleotides that are mutated per round of replication calculated. For example, the nucleic acid molecule of known sequence can be amplified using 10 rounds of PCR in the presence of the low bias DNA polymerase. The resulting nucleic acid molecule can then be sequenced. If the resulting nucleic acid molecule comprises 10% nucleotides that are different in corresponding nucleotides in the original known sequence, then the user would understand that the low bias DNA polymerase is able to mutate 1% of the nucleotides in the at least one target nucleic acid molecule on average per round of replication. Similarly, to see whether the low bias DNA polymerase mutates a certain percentage of the nucleotides in the at least one target nucleic acid molecule in a given method, the user could perform the method on a nucleic acid molecule of known sequence and use sequencing to determine the percentage of nucleotides that are mutated once the method is completed.

**[0061]** The low bias DNA polymerase is able to mutate a nucleotide such as adenine, if, when used to amplify a nucleic acid molecule, it provides a nucleic acid molecule in which some instances of that nucleotide are substituted or deleted. Preferably, the term "*mutate*" refers to introduction of substitution mutations, and in some embodiments the term "*mutate*" can be replaced with "*introduces substitutions of*".

**[0062]** The low bias DNA polymerase mutates a nucleotide such as adenine in at least one target nucleic acid molecule in the method of the invention if, when the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase is carried out, this step results in a mutated at least one target nucleic acid molecule in which some instances of that nucleotide are mutated. For example, if the low bias DNA polymerase mutates adenine in the at least one target nucleic acid molecule, when the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase is carried out, this step results in a mutated at least one target nucleic acid molecule in which at least one adenine has been substituted or deleted.

**[0063]** To determine whether a DNA polymerase is able to introduce certain mutations, the skilled person merely needs to test the DNA polymerase using a nucleic acid molecule of known sequence. A suitable nucleic acid molecule of known sequence is a fragment from a bacterial genome of known sequence, such as *E.coli* MG1655. The skilled person could amplify the nucleic acid molecule of known sequence using PCR in the presence of the low bias DNA polymerase. The skilled person could then sequence the amplified nucleic acid molecule and determine whether its sequence is the same as the original known sequence. If

not, the skilled person could determine the nature of the mutations. For example, if the skilled person wished to determine whether a DNA polymerase is able to mutate adenine using a nucleotide analog, the skilled person could amplify the nucleic acid molecule of known sequence using PCR in the presence of the nucleotide analog, and sequence the resulting amplified nucleic acid molecule. If the amplified DNA has mutations in positions corresponding to adenine nucleotides in the known sequence, then the skilled person would know that the DNA polymerase could mutate adenine using a nucleotide analog.

**[0064]** Rate ratios can be calculated in a similar manner. For example, if the skilled person wishes to determine the rate ratio at which guanine and cytosine nucleotides are mutated, the skilled person could amplify a nucleic acid molecule having a known sequence using PCR in the presence of the low bias DNA polymerase. The skilled person could then sequence the resulting amplified nucleic acid molecule and identify how many of the guanine nucleotides have been substituted or deleted and how many of the cytosine nucleotides have been substituted or deleted. The rate ratio is the ratio of the number of guanine nucleotides that have been substituted or deleted to the number of cytosine nucleotides that have been substituted or deleted. For example, if 16 guanine nucleotides have been replaced or deleted and 8 cytosine nucleotides have been replaced or deleted, the guanine and cytosine nucleotides have been mutated at a rate ratio of 16:8 or 2:1 respectively.

#### ***Using nucleotide analogs***

**[0065]** The low bias DNA polymerase may not be able to replace nucleotides with other nucleotides directly (at least not with high frequency), but the low bias DNA polymerase may still be able to mutate a nucleic acid molecule using a nucleotide analog. The low bias DNA polymerase may be able to replace nucleotides with other natural nucleotides (i.e. cytosine, guanine, adenine or thymine) or with nucleotide analogs.

**[0066]** The low bias DNA polymerase is a high fidelity DNA polymerase. High fidelity DNA polymerases tend to introduce very few mutations in general, as they are highly accurate. However, the present inventors have found that some high fidelity DNA polymerases may still be able to mutate a target nucleic acid molecule, as they may be able to introduce nucleotide analogs into a target nucleic acid molecule.

**[0067]** In an embodiment, in the absence of nucleotide analogs, the high fidelity DNA polymerase introduces less than 0.01%, less than 0.0015%, less than 0.001%, between 0% and 0.0015%, or between 0% and 0.001% mutations per round of replication.

**[0068]** The low bias DNA polymerase is able to incorporate nucleotide analogs into the at least one target nucleic acid molecule. The low bias DNA polymerase incorporates nucleotide analogs into the at least one target nucleic acid molecule. The low bias DNA polymerase can mutate adenine, thymine, guanine, and/or cytosine using a nucleotide analog. The low bias DNA polymerase mutates adenine, thymine, guanine, and/or cytosine in the at least one target

nucleic acid molecule using a nucleotide analog. The DNA polymerase replaces guanine, cytosine, adenine and/or thymine with a nucleotide analog. The DNA polymerase can replace guanine, cytosine, adenine and/or thymine with a nucleotide analog.

**[0069]** Incorporating nucleotide analogs into the at least one target nucleic acid molecule can be used to mutate nucleotides, as they may be incorporated in place of existing nucleotides and they may pair with nucleotides in the opposite strand. For example dPTP can be incorporated into a nucleic acid molecule in place of a pyrimidine nucleotide (may replace thymine or cytosine); please see Figure 7. Once in a nucleic acid strand, it may pair with adenine when in an imino tautomeric form. Thus, when a complementary strand is formed, that complementary strand may have an adenine present at a position complementary to the dPTP. Similarly, once in a nucleic acid strand, it may pair with guanine when in an amino tautomeric form. Thus, when a complementary strand is formed, that complementary strand may have a guanine present at a position complementary to the dPTP.

**[0070]** For example, if a dPTP is introduced into the at least one target nucleic acid molecule of the invention, when an at least one nucleic acid molecule complementary to the at least one target nucleic acid molecule is formed, the at least one nucleic acid molecule complementary to the at least one target nucleic acid molecule will comprise an adenine or a guanine at a position complementary to the dPTP in the at least one target nucleic acid molecule (depending on whether the dPTP is in its amino or imino form). When the at least one nucleic acid molecule complementary to the at least one target nucleic acid molecule is replicated, the resulting replicate of the at least one target nucleic acid molecule will comprise a thymine or a cytosine in a position corresponding to the dPTP in the at least one target nucleic acid molecule. Thus, a mutation to thymine or cytosine can be introduced into the mutated at least one target nucleic acid molecule.

**[0071]** Alternatively, if a dPTP is introduced in at least one nucleic acid molecule complementary to the at least one target nucleic acid molecule, when a replicate of the at least one target nucleic acid molecule is formed, the replicate of the at least one target nucleic acid molecule will comprise an adenine or a guanine at a position complementary to the dPTP in the at least one nucleic acid molecule complementary to the at least one target nucleic acid molecule (depending on the tautomeric form of the dPTP). Thus, a mutation to adenine or guanine can be introduced into the mutated at least one target nucleic acid molecule.

**[0072]** In an embodiment, the low bias DNA polymerase can replace cytosine or thymine with a nucleotide analog. In a further embodiment, the low bias DNA polymerase introduces guanine or adenine nucleotides using a nucleotide analog at a rate ratio of 0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2, or around 1:1 respectively. The guanine or adenine nucleotides may be introduced by the low bias DNA polymerase pairing them opposite a nucleotide analog such as dPTP. In a further embodiment, the low bias DNA polymerase introduces guanine or adenine nucleotides using a nucleotide analog at a rate ratio of 0.7-1.3:0.7-1.3 respectively.

**[0073]** The skilled person can determine, using conventional methods, whether the low bias DNA polymerase is able to incorporate nucleotide analogs into the at least one target nucleic acid molecule or mutate adenine, thymine, guanine, and/or cytosine in the at least one target nucleic acid molecule using a nucleotide analog using conventional methods.

**[0074]** For example, in order to determine whether the low bias DNA polymerase is able to incorporate nucleotide analogs into the at least one target nucleic acid molecule, the skilled person could amplify a nucleic acid molecule using a low bias DNA polymerase for two rounds of replication. The first round of replication should take place in the presence of the nucleotide analog, and the second round of replication should take place in the absence of the nucleotide analog. The resulting amplified nucleic acid molecules could be sequenced to see whether mutations have been introduced, and if so, how many mutations. The user should repeat the experiment without the nucleotide analog, and compare the number of mutations introduced with and without the nucleotide analog. If the number of mutations that have been introduced with the nucleotide analog is significantly higher than the number of mutations that have been introduced without the nucleotide analog, the user can conclude that the low bias DNA polymerase is able to incorporate nucleotide analogs. Similarly, the skilled person can determine whether a DNA polymerase incorporates nucleotide analogs or mutates adenine, thymine, guanine, and/or cytosine using a nucleotide analog. The skilled person merely need perform the method in the presence of nucleotide analogs, and see whether the method leads to mutations at positions originally occupied by adenine, thymine, guanine, and/or cytosine.

**[0075]** The method comprises a step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase, where the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase is carried out in the presence of the nucleotide analog, and the step of amplifying the at least one target nucleic acid molecule provides at least one target nucleic acid molecule comprising the nucleotide analog.

**[0076]** Suitable nucleotide analogs include dPTP (2'-deoxy-P-nucleoside-5'-triphosphate), 8-Oxo-dGTP (7,8-dihydro-8-oxoguanine), 5Br-dUTP (5-bromo-2'-deoxy-uridine-5'-triphosphate), 2OH-dATP (2-hydroxy-2'-deoxyadenosine-5'-triphosphate), dKTP (9-(2-Deoxy-(3-D-ribofuranosyl)-N6-methoxy-2,6,-diaminopurine-5'-triphosphate) and dITP (2'-deoxyinosine 5'-triphosphate). The nucleotide analog may be dPTP. The nucleotide analogs may be used to introduce the substitution mutations described in Table 1.

**Table 1**

Nucleotide	Substitution
8-oxo-dGTP	A:T to C:G and T:A to G:C
dPTP	A:T to G:C and G:C to A:T
5Br-dUTP	A:T to G:C and T:A to C:G
2OH-dATP	A:T to C:G, G:C to T:A and A:T to G:C
dITP	A:T to G:C and G:C to A:T
dKTP	A:T to G:C and G:C to A:T

**[0077]** The different nucleotide analogs can be used, alone or in combination, to introduce different mutations into the at least one target nucleic acid molecule. Accordingly, the low bias DNA polymerase may introduce guanine to adenine substitution mutations, cytosine to thymine substitution mutations, adenine to guanine substitution mutations, and thymine to cytosine substitution mutations using a nucleotide analog. The low bias DNA polymerase may be able to introduce guanine to adenine substitution mutations, cytosine to thymine substitution mutations, adenine to guanine substitution mutations, and thymine to cytosine substitution mutations, optionally using a nucleotide analog.

**[0078]** The low bias DNA polymerase may be able to introduce guanine to adenine substitution mutations, cytosine to thymine substitution mutations, adenine to guanine substitution mutations, and thymine to cytosine substitution mutations at a rate ratio of 0.5-1.5:0.5-1.5:0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4:0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2:0.8-1.2:0.8-1.2, or around 1:1:1:1 respectively. Preferably, the low bias DNA polymerase is able to introduce guanine to adenine substitution mutations, cytosine to thymine substitution mutations, adenine to guanine substitution mutations, and thymine to cytosine substitution mutations at a rate ratio of 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3 respectively. Suitable methods for determining whether the low bias DNA polymerase is able to introduce substitution mutations and at what rate ratio are described under the heading "*whether a DNA polymerase is able to mutate a nucleotide and, if so, at what rate*".

**[0079]** In some methods the low bias DNA polymerase introduces guanine to adenine substitution mutations, cytosine to thymine substitution mutations, adenine to guanine substitution mutations, and thymine to cytosine substitution mutations at a rate ratio of 0.5-1.5:0.5-1.5:0.5-1.5:0.5-1.5, 0.6-1.4:0.6-1.4:0.6-1.4:0.6-1.4, 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3, 0.8-1.2:0.8-1.2:0.8-1.2:0.8-1.2, or around 1:1:1:1 respectively. Preferably, the low bias DNA polymerase introduces guanine to adenine substitution mutations, cytosine to thymine substitution mutations, adenine to guanine substitution mutations, and thymine to cytosine substitution mutations at a rate ratio of 0.7-1.3:0.7-1.3:0.7-1.3:0.7-1.3 respectively. Suitable methods for determining whether substitution mutations are introduced and at what rate ratio are described under the heading "*whether a DNA polymerase is able to mutate a nucleotide and, if so, at what rate*".

**[0080]** Generally, when a low bias DNA polymerase uses a nucleotide analog to introduce a mutation, this requires more than one round of replication. In the first round of replication the low bias DNA polymerase introduces the nucleotide analog in place of a nucleotide, and in a second round of replication, that nucleotide analog pairs with a natural nucleotide to introduce a substitution mutation in the complementary strand. The second round of replication may be carried out in the presence of the nucleotide analog. However, the method may further comprise a step of amplifying the at least one target nucleic acid molecule comprising nucleotide analogs in the absence of nucleotide analogs. The step of amplifying the at least one target nucleic acid molecule comprising nucleotide analogs in the absence of nucleotide analogs may be carried out using the low bias DNA polymerase.

[0081] Optionally, the method provides a mutated at least one target nucleic acid molecule and the method comprises a further step of amplifying the mutated at least one target nucleic acid molecule using the low bias DNA polymerase.

***Low template amplification bias***

[0082] The low bias DNA polymerase has low template amplification bias. A low bias DNA polymerase has low template amplification bias if it is able to amplify different target nucleic acid molecules with similar degrees of success per cycle. High bias DNA polymerases may struggle to amplify template nucleic acid molecules that comprise a high G:C content or contain a large degree of secondary structure. In an embodiment, the low bias DNA polymerase of the invention has low template amplification bias for template nucleic acid molecules that are less than 25 000, less than 10 000, between 1 and 15 000, or between 1 and 10 000 nucleotides in length.

[0083] In an embodiment, to determine whether a DNA polymerase has low template amplification bias, the skilled person could amplify a range of different sequences using the DNA polymerase, and see whether the different sequences are amplified at different levels by sequencing the resultant amplified DNA. For example, the skilled person could select a range of short (possibly 50 nucleotide) nucleic acid molecules having different characteristics, including a nucleic acid molecule having high GC content, a nucleic acid molecule having low GC content, a nucleic acid molecule having a large degree of secondary structure and a nucleic acid molecule have a low degree of second structure. The user could then amplify those sequences using the DNA polymerase and quantify the level at which each of the nucleic acid molecules is amplified to. In an embodiment, if the levels are within 25%, 20%, 10%, or 5% of one another, then the DNA polymerase has low template amplification bias.

[0084] Alternatively, in an embodiment, a DNA polymerase has low template amplification bias if it is able to amplify 7-10 kbp fragments with a Kolmogorov-Smirnov D of less than 0.1, less than 0.09, or less than 0.08. The Kolmogorov-Smirnov D with which a particular low bias DNA polymerase is able to amplify 7-10 kbp fragments may be determined using an assay provided in Example 4.

[0085] The low bias DNA polymerase is a high fidelity DNA polymerase. A high fidelity DNA polymerase is a DNA polymerase which is not highly error-prone, and so does not generally introduce a large number of mutations when used to amplify a target nucleic acid molecule in the absence of nucleotide analogs. High fidelity DNA polymerases are not generally used in methods for introducing mutations, as it is generally considered that error-prone DNA polymerases are more effective. However, the present application demonstrates that certain high fidelity polymerases are able to introduce mutations using a nucleotide analog, and that those mutations may be introduced with lower bias compared to error-prone DNA polymerases such as Taq polymerase.

**[0086]** High fidelity DNA polymerases have an additional advantage. High fidelity DNA polymerases can be used to introduce mutations when used with nucleotide analogs, but in the absence of nucleotide analogs they can replicate a target nucleic acid molecule highly accurately. This means that the user can mutate the at least one target nucleic acid molecule to high effect and amplify the mutated at least one target nucleic acid molecule with high accuracy using the same DNA polymerase. If a low fidelity DNA polymerase is used to mutate the target nucleic acid molecule, it may need to be removed from the reaction mixture before the target nucleic acid molecule is amplified.

**[0087]** High fidelity DNA polymerases may have a proof-reading activity. A proof-reading activity may help the DNA polymerase to amplify a target nucleic acid sequence with high accuracy. For example, a low bias DNA polymerase may comprise a proof-reading domain. A proof reading domain may confirm whether a nucleotide that has been added by the polymerase is correct (checks that it correctly pairs with the corresponding nucleic acid of the complementary strand) and, if not, excises it from the nucleic acid molecule. The inventors have surprisingly found that in some DNA polymerases, the proof-reading domain will accept pairings of natural nucleotides with nucleotide analogs. The structure and sequence of suitable proof-reading domains are known to the skilled person. DNA polymerases that comprise a proof-reading domain include members of DNA polymerase families I, II and III, such as Pfu polymerase (derived from *Pyrococcus furiosus*), T4 polymerase (derived from bacteriophage T4) and the Thermococcal polymerases that are described in more detail below.

**[0088]** In an embodiment, in the absence of nucleotide analogs, the high fidelity DNA polymerase introduces less than 0.01%, less than 0.0015%, less than 0.001%, between 0% and 0.0015%, or between 0% and 0.001% mutations per round of replication.

**[0089]** In addition, the low bias DNA polymerase may comprise a processivity enhancing domain. A processivity enhancing domain allows a DNA polymerase to amplify a target nucleic acid molecule more quickly. This is advantageous as it allows the methods of the invention to be performed more quickly.

### ***Thermococcal polymerases***

**[0090]** In an embodiment, the low bias DNA polymerase is a fragment or variant of a polypeptide comprising SEQ ID NO. 2, SEQ ID NO. 4, SEQ ID NO. 6 or SEQ ID NO.7. The polypeptides of SEQ ID NO. 2, 4, 6 and 7 are thermococcal polymerases. The polymerases of SEQ ID NO. 2, SEQ ID NO. 4, SEQ ID NO. 6 or SEQ ID NO. 7 are low bias DNA polymerases having high fidelity, and they can mutate target nucleic acid molecules by incorporating a nucleotide analog such as dPTP. The polymerases of SEQ ID NO. 2, SEQ ID NO. 4, SEQ ID NO. 6 or SEQ ID NO. 7 are particularly advantageous as they have low mutation bias and low template amplification bias. They are also highly processive and are high fidelity polymerases

comprising a proof-reading domain, meaning that, in the absence of nucleotide analogs, they can amplify mutated target nucleic acid molecules quickly and accurately.

**[0091]** The low bias DNA polymerase may comprise a fragment of at least 400, at least 500, at least 600, at least 700, or at least 750 contiguous amino acids of:

1. a. a sequence of SEQ ID NO. 2;
2. b. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 2;
3. c. a sequence of SEQ ID NO. 4;
4. d. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 4;
5. e. a sequence of SEQ ID NO. 6;
6. f. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 6;
7. g. a sequence of SEQ ID NO. 7; or
8. h. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 7.

**[0092]** Preferably, the low bias DNA polymerase comprises a fragment of at least 700 contiguous amino acids of:

1. a. a sequence of SEQ ID NO. 2;
2. b. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 2;
3. c. a sequence of SEQ ID NO. 4;
4. d. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 4;
5. e. a sequence of SEQ ID NO. 6;
6. f. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 6;
7. g. a sequence of SEQ ID NO. 7; or
8. h. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 7.

**[0093]** The low bias DNA polymerase may comprise:

1. a. a sequence of SEQ ID NO. 2;
2. b. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 2;
3. c. a sequence of SEQ ID NO. 4;
4. d. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 4;
5. e. a sequence of SEQ ID NO. 6;
6. f. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 6;
7. g. a sequence of SEQ ID NO. 7; or
8. h. a sequence at least 95%, at least 98%, or at least 99% identical to SEQ ID NO. 7.

**[0094]** Preferably, the low bias DNA polymerase comprises:

1. a. a sequence of SEQ ID NO. 2;
2. b. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 2;
3. c. a sequence of SEQ ID NO. 4;
4. d. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 4;
5. e. a sequence of SEQ ID NO. 6;
6. f. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 6;
7. g. a sequence of SEQ ID NO. 7; or
8. h. a sequence at least 98%, or at least 99% identical to SEQ ID NO. 7.

**[0095]** The low bias DNA polymerase may be a thermococcal polymerase, or derivative thereof. The DNA polymerases of SEQ ID NO 2, 4, 6 and 7 are thermococcal polymerases. Thermococcal polymerases are advantageous, as they are generally high fidelity polymerases that can be used to introduce mutations using a nucleotide analog with low mutation and template amplification bias.

**[0096]** A thermococcal polymerase is a polymerase having the polypeptide sequence of a polymerase isolated from a strain of the *Thermococcus* genus. A derivative of a thermococcal polymerase may be a fragment of at least 400, at least 500, at least 600, at least 700, or at least 750 contiguous amino acids of a thermococcal polymerase, or at least 95%, at least 98%, at least 99%, or 100% identical to a fragment of at least 400, at least 500, at least 600, at least 700, or at least 750 contiguous amino acids of a thermococcal polymerase. The derivative of a thermococcal polymerase may be at least 95%, at least 98%, at least 99%, or 100% identical to a thermococcal polymerase. The derivative of a thermococcal polymerase may be at least 98% identical to a thermococcal polymerase.

**[0097]** A thermococcal polymerase from any strain may be effective in the context of the present invention. In an embodiment, the thermococcal polymerase is derived from a thermococcal strain selected from the group consisting of *T. kodakarensis*, *T. celer*, *T. siculi*, and *T. sp* KS-1. Thermococcal polymerases from these strains are described in SEQ ID NO. 2, SEQ ID NO. 4, SEQ ID NO. 6 and SEQ ID NO. 7.

**[0098]** Optionally, the low bias DNA polymerase is a polymerase that has high catalytic activity at temperatures between 50°C and 90°C, between 60°C and 80°C, or around 68°C.

#### ***Barcodes, sample tags and adapters***

**[0099]** The method may further comprise introducing barcodes into the target nucleic acid molecules. For the purposes of the present invention, a barcode is a degenerate or randomly generated sequence of nucleotides. The term "barcode" is synonymous with the terms "unique molecular identifiers" (UMIs) or "unique molecular tags" (UMTs). The method may comprise introducing 1, 2 or more barcodes into the target nucleic acid molecules. In a preferred

embodiment, the method comprises introducing a variety of barcodes into the target nucleic acid molecules, such that, after the barcodes are introduced, most of the original target nucleic acid molecules comprise unique barcodes compared to other original target nucleic acid molecules.

**[0100]** Introducing barcodes into the target nucleic acid molecules may be useful if the method for introducing mutations of the invention is used as part of a method for determining a sequence. The use of barcodes may help the user to identify which of the original at least one target nucleic acid molecules each sequence of at least one of target nucleic acid molecule (or amplified or fragmented at least one target nucleic acid molecule) was derived from. If the barcodes used in each original target nucleic acid molecule are different, the user can sequence the barcodes or the target nucleic acid molecules, and sequences of target nucleic acid molecules comprising the same barcodes are likely to be sequences of target nucleic acid molecules that originated from the same original target nucleic acid molecule.

**[0101]** The method for introducing mutations into at least one target nucleic acid molecule may comprise introducing sample tags into the target nucleic acid molecules. A sample tag is a short series of nucleic acids of known (specified) sequence. For example, the method of the invention may be performed on multiple target nucleic acid molecules taken from different samples. Those samples may be pooled, but prior to pooling, a sample tag may be introduced into the target nucleic acid molecules in a sample (the target nucleic acid molecules are labelled with a sample tag). Target nucleic acid molecules from different samples may be labelled with different sample tags. Optionally, target nucleic acid molecules from the same sample are labelled with the same sample tag or a sample tag from the same sub-group of sample tags. For example, if the user decides to use two samples, the target nucleic acid molecules in the first sample may be labelled with a first sample tag having a specified sequence and the target nucleic acid molecules in the second sample may be tagged with a second sample tag having a second specified sequence. Similarly, if the user decides to use two samples, the target nucleic acid molecules in the first sample may be labelled with a sample tag from a first sub-group of sample tags and the target nucleic acid molecules in the second sample may be labelled with a sample tag from a second sub-group of sample tags. The user would understand that any target nucleic acid molecules comprising the first sample tag or a sample tag from the first sub-group of sample tags originated from the first sample, and any target nucleic acid molecules comprising the second sample tag or a sample tag from the second sub-group of sample tags originated from the second sample. It is possible to determine which tag has been used to label a target nucleic acid sequence by sequencing the target nucleic acid sequence. Suitable sequencing methods are described in more detail below.

**[0102]** In an embodiment, the sample tags are introduced (the target nucleic acid molecules are labelled with a sample tag) prior to the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase. This is advantageous as it means that samples may be pooled at an early stage in the method, reducing handling time, the number of reagents required and the possibility of introducing sample handling mistakes. However, if the

sample tags are introduced prior to the step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase, it is possible that the sample tags will be mutated by the low bias DNA polymerase. The present inventors have designed groups of sample tags that are designed such that they may be distinguished from one another even if they have been mutated.

**[0103]** In an embodiment, a group of sample tags is used and target nucleic acid molecules from different samples are labelled with different sample tags from the group. Target nucleic acid molecules from the same sample may be labelled with the same sample tag from the group or with a sample tag from the same sub-group of sample tags from the group. For example, if the group of sample tags comprises sample tags named A, B, C and D, all target nucleic acid molecules in a first sample may be labelled using A or A/B, and all the target nucleic acid molecules in a second sample may be labelled using C or C/D. Each sample tag in the group of sample tags may differ from substantially all other sample tags in the group by at least 1 low probability mutation difference. Each sample tag in the group of sample tags may differ from all other sample tags in the group by at least 1 low probability mutation difference.

**[0104]** In an aspect, the disclosure provides a group of sample tags, wherein each sample tag in the group differs from substantially all other sample tags in the group by at least 1 low probability mutation difference. Each sample tag may differ from all other sample tags in the group by at least 1 low probability mutation difference.

**[0105]** By the term *"differs from substantially all other sample tags in the group by at least 1 low probability mutation difference"* we mean that each tag has been designed such that if the sample tags are mutated by at least 1 low probability mutation, the tags will still be different to one another almost (substantially all or all other tags). In an embodiment, the term *"substantially all other sample tags"* refers to at least 90%, at least 95%, or at least 98% of the other sample tags. A low probability mutation is a mutation that occurs infrequently in the method for introducing mutations of the invention. For example, a low probability mutation may be a transversion mutation, or an indel mutation. Transversion mutations and indel mutations occur infrequently when the method for introducing mutations of the invention is performed using dPTP as a nucleotide analog. A transversion mutation is a replacement of a purine nucleotide with a pyrimidine nucleotide (adenine to cytosine, adenine to thymine, guanine to cytosine or guanine to thymine), or a pyrimidine nucleotide with a purine nucleotide (cytosine to adenine, cytosine to guanine, thymine to adenine, or thymine to guanine). An indel mutation is a deletion mutation or an insertion mutation. Suitable tags may be designed computationally using statistical methods. For example, the skilled person would be able to determine what type of mutation is a low probability mutation in a method for introducing mutations of the invention. The skilled person can perform the method for introducing mutations of the invention, and determine the types of mutations that have been introduced by sequencing the nucleic acid molecule product. The mutations that occur most frequently are high probability mutations, and the mutations that occur least frequently are low probability mutations.

**[0106]** The user could generate suitable sample tags using the method for designing a group

of sample tags.

**[0107]** Optionally, each sample tag differs from substantially all other sample tags in the group by at least 2, at least 3, at least 4, at least 5, between 3 and 50, between 3 and 25, or between 3 and 10 low probability mutation differences. Optionally, each sample tag differs from all other sample tags in the group by at least 2, at least 3, at least 4, at least 5, between 3 and 50, between 3 and 25, or between 3 and 10 low probability mutation differences.

**[0108]** Each sample tag may differ from substantially all other sample tags in the group by at least 2 high probability mutation differences. A high probability mutation difference, is a mutation that occurs frequently in a method for introducing mutations of the invention. For example, a high probability mutation difference may be a transition mutation. A transition mutation is a replacement of a purine nucleotide with another purine nucleotide (adenine to guanine or guanine to adenine), or a pyrimidine nucleotide with another pyrimidine nucleotide (cytosine to thymine or thymine to cytosine).

**[0109]** Each sample tag may differ from all other sample tags in the group by at least 2 high probability mutation differences, *i.e.* each sample tag has been designed such that if the sample tags are mutated by at least 2 high probability mutations, the tags will still be different to one another.

**[0110]** Optionally, each sample tag differs from substantially all other sample tags in the group by at least 3, between 2 and 50, between 3 and 25, or between 3 and 10 high probability mutation differences. Optionally, each sample tag differs from all other sample tags in the group by at least 3, between 2 and 50, between 5 and 25, or between 5 and 10 high probability mutation differences.

**[0111]** In an embodiment, each sample tag is at least 8 nucleotides, at least 10 nucleotides, at least 12 nucleotides, between 8 and 50 nucleotides, between 10 and 50 nucleotides, or between 10 and 50 nucleotides in length.

**[0112]** Suitable sample tags are those of SEQ ID NOs: 8-136.

**[0113]** The method may further comprise introducing adapters into each of the target nucleic acid molecules. The adapters may comprise a primer binding site. For the purposes of the invention, primer binding sites are known sequences of nucleotides that are sufficiently long for primers to specifically hybridise to. Optionally, the primer binding sites are at least 8, at least 10, at least 12, between 8 and 50, or between 10 and 25 nucleotides in length.

**[0114]** The method may comprise introducing a first adapter at the 3' end of the at least one target nucleic acid molecule and a second adapter at the 5' end of the at least one target nucleic acid molecule, wherein the first adapter and the second adapter can anneal to one another.

**[0115]** Also described herein is a method for preferentially amplifying nucleic acid molecules that are larger than 1 kbp in length comprising:

1. a. providing at least one sample comprising target nucleic acid molecules;
2. b. introducing a first adapter at the 3' end of the target nucleic acid molecules and a second adapter at the 5' end of the target nucleic acid molecules; and
3. c. amplifying the target nucleic acid molecules using primers that are complementary to a portion of the first adapter,

wherein the first adapter and the second adapter can anneal to one another.

**[0116]** The second adapter may comprise a portion that is complementary to a first primer binding site and the first adapter may comprise the first primer binding site.

**[0117]** The present inventors have found that by introducing a first adapter and a second adapter that can anneal to one another into the at least one target nucleic acid molecule, they can ensure that the methods of the invention preferentially amplify and/or mutate long target nucleic acid molecules. If the first adapter can anneal to the second adapter, then they may do so in the methods of the invention resulting in a self-annealed at least one target nucleic acid molecule (as indicated in Figure 5). Self-annealed target nucleic acid molecules are not replicated and so will not be amplified and/or mutated by the methods of the invention. The likelihood that the first adapter and the second adapter anneal to one another during the methods of the invention will be higher for shorter target nucleic acid molecules than for longer target nucleic acid molecules. For these reasons, the addition of a first adapter and a second adapter to the at least one target nucleic acid molecule of the invention can be used to preferentially amplify larger at least one target nucleic acid molecules.

**[0118]** The method for preferentially amplifying nucleic acid molecules may be a method for preferentially amplifying target nucleic acid molecules that are longer than 1.5 kbp. The method may further comprise a step of sequencing the target nucleic acid molecules. Examples of possible sequencing methods include Maxam Gilbert Sequencing, Sanger Sequencing, nanopore sequencing or sequencing comprising bridge PCR. In a typical embodiment, the sequencing steps involve bridge PCR. Optionally, the bridge PCR step is carried out using an extension time of greater than 5, greater than 10, greater than 15 or greater than 20 seconds. An example of the use of bridge PCR is in Illumina Genome Analyzer Sequencers.

**[0119]** It is possible for a user to determine whether a first adapter and a second adapter can anneal to one another. In an embodiment, the user may identify whether a first adapter and a second adapter can anneal to one another by providing a nucleic acid molecule comprising the first adapter, and seeing whether a primer comprising the second adapter is capable of initiating replication of the nucleic acid molecule under PCR conditions.

**[0120]** Alternatively, in an embodiment, the first adapter and the second adapter can be considered to be able to anneal to one another if they hybridise under the following conditions:

equimolar concentrations of the two primers are combined (e.g. 50  $\mu\text{M}$ ), then incubated at a high temperature such as 95°C for 5 minutes to ensure that the primers are single-stranded. The solution is then slowly cooled to room temperature (25°C) over a period of approximately 45 minutes.

**[0121]** The methods may comprise amplifying the target nucleic acid molecules using primers that are identical to one another, or substantially identical to one another. The primers may be complementary to a portion of the first adapter. Two primers are "*substantially identical*" to one another if they have an identical sequence, or a sequence that differs by 1, 2 or 3 nucleotides. In a preferred embodiment, the methods of the invention comprise amplifying the target nucleic acid molecules using primers that are identical in sequence or differ by a single nucleotide difference.

**[0122]** In an embodiment, the first adapter and the second adapter comprise sequences that are complementary to one another, or substantially complementary to one another. The first adapter may be substantially complementary to the second adapter if the first adapter is complementary to a nucleic acid molecule that is at least 80%, at least 90%, at least 95%, or at least 99% identical to the second adapter.

**[0123]** The user may use primers that comprise primer binding sites, and these primers may be used to preferentially amplify replicates of the at least one target nucleic acid molecule that were generated in the last round of replication. For example, a first set of primers comprising a third primer binding site may be used in a round of replication. In a further round of replication a second set of primers may be used that bind to the third primer binding site. The second set of primers will only replicate replicates of the at least one target nucleic acid molecule that were generated in a previous round of replication, using the first set of primers.

**[0124]** Third and further sets of primers may be used. Preferentially replicating replicates of a previous round of replication is advantageous as it can ensure that each amplified target nucleic acid molecule comprises a high level of mutation (since only at least one target nucleic acid molecules that have been exposed to at least one round of amplification by the low bias DNA polymerase will be replicated).

**[0125]** Accordingly, the methods of the invention may comprise:

1. (a) introducing a first adapter comprising a first primer binding site at the 3' end of the at least one target nucleic acid molecule or target nucleic acid molecules and a second adapter comprising a portion that is complementary to the first primer binding site at the 5' end of the at least one target nucleic acid molecule or target nucleic acid molecules, wherein the first adapter and the second adapter can anneal to one another;
2. (b) amplifying the target nucleic acid molecules using a first set of primers that are complementary to the first primer binding site and comprise a second primer binding site, optionally using a low bias DNA polymerase; and
3. (c) amplifying the target nucleic acid molecules using a second set of primers that are

complementary to the second primer binding site, optionally using a low bias DNA polymerase.

**[0126]** The second set of primers may comprise a third primer binding site, and further amplification steps may be carried out using a third or further sets of primers that are complementary to the third or further primer binding sites.

**[0127]** The barcodes, sample tags and/or adapters may be introduced using any suitable method including PCR, tagmentation and physical shearing or restriction digestion of target nucleic acids combined with subsequent adapter ligation (optionally sticky-end ligation). For example, PCR can be carried out on the at least one target template nucleic acid molecule using a first set of primers capable of hybridising to the at least one target nucleic acid molecule. The barcodes, sample tags and adapters may be introduced into each of the at least one target nucleic acid molecules by PCR using primers comprising a portion (a 5' end portion) comprising a barcode, a sample tag and/or an adapter, and a portion (a 3' end portion) having a sequence that is capable of hybridising to (optionally complementary to) the at least one target nucleic acid molecule. Such primers will hybridise to a target nucleic acid molecule, PCR primer extension will then provide a nucleic acid molecule which comprises a barcode, sample tag and/or an adapter. A further cycle of PCR with these primers can be used to add a barcode, sample tag and/or an adapter to the other end of the at least one target nucleic acid molecule. The primers may be degenerate, *i.e.* the 3' end portion of the primers may be similar but not identical to one another.

**[0128]** The barcodes, sample tags and/or adapters may be introduced using tagmentation. The barcodes, sample tags and/or adapters can be introduced using direct tagmentation, or by introducing a defined sequence by tagmentation followed by two cycles of PCR using primers that comprise a portion capable of hybridising to the defined sequence, and a portion comprising a barcode, a sample tag and/or an adapter. The barcodes, sample tags and/or adapters can be introduced by restriction digestion of the original at least one target nucleic acid molecule followed by ligation of nucleic acids comprising the barcode, sample tag and/or an adapter. The restriction digestion of the original at least one nucleic acid molecule should be performed such that the digestion results in a nucleic acid molecule comprising the region to be sequenced (the at least one target template nucleic acid molecule). The barcodes, sample tags and/or adapters may be introduced by shearing the at least one target nucleic acid molecule, followed by end repair, A-tailing and then ligation of nucleic acids comprising the barcode, sample tag and/or an adapter.

***A method for determining a sequence of at least one target nucleic acid molecule***

**[0129]** One aspect of the invention relates to a method for determining a sequence of at least one target nucleic acid molecule comprising the method for introducing mutations of the

invention.

**[0130]** As described above, the method for introducing mutations of the invention can be useful as part of a method for determining a sequence of at least one target nucleic acid molecule, as the mutations can enable the skilled person to assemble sequences.

**[0131]** As described in the background section, sequencing methods can be improved by incorporating steps that introduce mutations into at least one target nucleic acid molecule that is to be sequenced. A user will often amplify and/or fragment the at least one target nucleic acid molecule prior to sequencing it. The user will then assemble a consensus sequence for at least one of the target nucleic acid molecules from the sequences of regions of the amplified or fragmented at least one target nucleic acid molecule. Introducing mutations into the at least one target nucleic acid molecules prior to amplification or fragmentation can help the user to identify which of the original at least one template nucleic acid molecules each sequence of regions of amplified or fragmented at least one target nucleic acid molecule was derived from, and so improve the accuracy of the consensus sequences.

**[0132]** The more random the mutations that are introduced, the easier it is to identify which of the original at least one target nucleic acid molecule each sequence of amplified or fragmented at least one target nucleic acid molecule was derived from. The method of introducing mutations of the invention, which utilises a low bias DNA polymerase, can be used to introduce mutations in a substantially random way, and so is ideal for inclusion in a method for determining a sequence of at least one target nucleic acid molecule.

**[0133]** The method for determining a sequence of at least one target nucleic acid molecule may comprise steps of:

1. a. performing the method for introducing mutations into at least one target nucleic acid molecule of the invention to provide at least one mutated target DNA molecule;
2. b. optionally fragmenting and/or amplifying the at least one mutated target DNA molecule to provide at least one fragmented and/or amplified mutated target DNA molecule;
3. c. sequencing regions of the least one mutated target DNA molecule of the at least one fragmented and/or amplified mutated target DNA molecule to provide mutated sequence reads; and
4. d. assembling a sequence for at least a portion of the at least one target DNA molecule using the mutated sequence reads.

**[0134]** In general, sequencing steps can be carried out using any method of sequencing. Examples of possible sequencing methods include Maxam Gilbert Sequencing, Sanger Sequencing, nanopore sequencing, or sequencing comprising bridge PCR. In a typical embodiment, the sequencing steps involve bridge PCR. Optionally, the bridge PCR step is carried out using an extension time of greater than 5, greater than 10, greater than 15 or

greater than 20 seconds. An example of the use of bridge PCR is in Illumina Genome Analyzer Sequencers.

**[0135]** The method may comprise sequencing regions of at least one mutated target nucleic acid molecule to provide mutated sequence reads. The regions may correspond to a fragment that may comprise a substantial portion of the at least one mutated target nucleic acid molecule. It may be that the entire at least one mutated target nucleic acid molecule cannot be sequenced for some reason, but the user may still find the sequence of a portion of the at least one mutated target nucleic acid molecule to be useful. The regions of the at least one mutated target nucleic acid molecule may comprise the entire length of the at least one mutated target nucleic acid molecule.

**[0136]** The method may comprise assembling a sequence for at least a portion of the at least one target nucleic acid molecule from the mutated sequence reads. The sequence may be assembled by aligning the mutated sequence reads and grouping together reads that share the same mutation pattern. A sequence will be assembled from mutated sequence reads in the same group. The assembly may be carried out using software such as Clustal W2, IDBA-UD or SOAPdenovo.

**[0137]** A step of amplifying the at least one mutated target nucleic acid molecule could be performed by any suitable amplification technique such as PCR. Suitably, the PCR is carried out using the low bias DNA polymerase under conditions such as those described under the heading "*amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase*".

**[0138]** A step of fragmenting the at least one mutated target nucleic acid molecule could be carried out using any appropriate method. For example, fragmentation can be carried out using restriction digestion or using PCR with primers complementary to at least one internal region of the at least one mutated target nucleic acid molecule. Preferably, fragmentation is carried out using a technique that produces arbitrary fragments. The term "*arbitrary fragment*" refers to a randomly generated fragment, for example a fragment generated by tagmentation. Fragments generated using restriction enzymes are not "*arbitrary*" as restriction digestion occurs at specific DNA sequences defined by the restriction enzyme that is used. Even more preferably, fragmentation is carried out by tagmentation. If fragmentation is carried out by tagmentation, the tagmentation reaction optionally introduces an adapter region into the at least one mutated target nucleic acid molecule. This adapter region is a short DNA sequence which may encode, for example, adapters to allow the at least one mutated target nucleic acid molecule to be sequenced using Illumina technology.

**[0139]** The fragmentation step may comprise a further step of enriching the at least one mutated fragmented target nucleic acid molecule. The step of enriching the at least one mutated fragmented target nucleic acid molecule may be carried out by PCR. Suitably, the PCR is carried out using the low bias DNA polymerase under conditions such as those described under the heading "*amplifying the at least one target nucleic acid molecule using a*

*low bias DNA polymerase* ".

**A method for engineering a protein**

**[0140]** The method for introducing mutations of the invention may be useful as part of a method for engineering a protein. For example, protein engineering may involve searching for mutations that increase or decrease the activity of a protein, or change its structure. As part of protein engineering, a user may wish to randomly mutate the protein and see how the mutations effect the activity or structure of the protein. The present method is a method that results in highly random mutagenesis, and so can advantageously be used as part of a method for engineering a protein.

**[0141]** Accordingly, in one aspect of the invention there is provided a method for engineering a protein comprising the method for introducing mutations of the invention.

**[0142]** The method comprises steps of:

1. a. performing a method for introducing mutations of the invention to provide at least one mutated target DNA molecule;
2. b. inserting the at least one mutated target DNA molecule into a vector; and
3. c. expressing a protein encoded by the at least one mutated target nucleic acid molecule.

**[0143]** The method may comprise steps of:

1. a. performing a method for introducing mutations of the invention to provide at least one mutated target nucleic acid molecule;
2. b. amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase in the presence of a nucleotide analog to provide target nucleic acid molecules comprising a nucleotide analog;
3. c. amplifying the target nucleic acid molecules comprising a nucleotide analog in the absence of nucleotide analogs to provide at least one mutated target nucleic acid molecule;
4. d. inserting the at least one mutated target nucleic acid molecule into a vector; and
5. e. expressing a protein encoded by the at least one mutated target nucleic acid molecule.

**[0144]** Any suitable vector can be used. Optionally the vector is a plasmid, a virus, a cosmid or an artificial chromosome. Typically, the vector further comprises a control sequence operably linked to the inserted sequence, thus allowing for expression of a polypeptide. Preferably, the

vector further comprises appropriate initiators, promoters, enhances and other elements which may be necessary and which are positioned in the correct orientation, in order to allow for expression of a polypeptide.

**[0145]** Optionally, the step of expressing the at least one mutated target nucleic acid molecule is achieved by transforming bacterial cells, transfecting eukaryotic cells or transducing eukaryotic cells with the vector. Optionally, the bacterial cells are *Escherichia coli* (*E.coli*) cells.

**[0146]** For example, the step of expressing the at least one mutated target nucleic acid molecule may comprise inserting the at least one mutated target nucleic acid molecule into a plasmid vector and transforming *E.coli* with the plasmid. The plasmid may comprise control elements suitable for expressing in *E.coli* such as a lac or T7 promoter (Dubendorff JW, Studier FW (1991). "Controlling basal expression in an inducible T7 expression system by blocking the target T7 promoter with lac repressor". Journal of Molecular Biology. 219 (1): 45-59.)). Suitable expression techniques are described in Sambrook, J. et al., (1989) Molecular Cloning: A Laboratory Manual Second Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.

**[0147]** Alternatively, the step of expressing the at least one mutated target nucleic acid molecule may comprise expressing fragments produced directly from the step of amplifying the target nucleic acid molecules using an *in vitro* method.

**[0148]** The method may further comprise a step of testing the activity or assessing the structure of the protein encoded by the at least one mutated target nucleic acid molecule.

**[0149]** The step of testing the activity or assessing the structure of the protein encoded by the at least one mutated target nucleic acid molecule may be carried out using any number of well-known techniques. For example, the skilled person would be aware of suitable techniques for assessing the structure of a protein, including nuclear magnetic resonance (NMR) techniques, microscopy techniques such as cryo-electron microscopy, small angle x-ray scattering techniques, or X-ray crystallography.

**[0150]** Similarly, the skilled person would be aware of techniques that could be used for assessing the activity of a protein. The method used will depend on the protein that is encoded by the at least one mutated target nucleic acid molecule. For example, if the protein that is encoded by the at least one mutated target nucleic acid molecule is a blood clotting factor, the skilled person would test the protein for clotting activity, for example using a chromogenic clotting assay. Alternatively, if the protein that is encoded by the at least one mutated target nucleic acid molecule is an enzyme, the skilled person could test the activity of the enzyme by measuring the rate at which it catalyses its reaction, for example by measuring reduction in concentration of a starting product or increase in concentration of an end product of the reaction catalysed by the enzyme.

***A method for designing a group of sample tags***

**[0151]** Also described herein is a method for designing a group of sample tags suitable for use in a method for introducing mutations into at least one target nucleic acid molecule comprising:

1. a. analysing the method for introducing mutations into at least one target nucleic acid molecule and determining the average number of low probability mutations that take place during the method for introducing mutations into at least one target nucleic acid molecule; and
2. b. determining sequences for a group of sample tags wherein each sample tag differs from substantially all sample tags in the group by more low probability mutation differences than the average number of low probability mutations that take place during the method for introducing mutations into at least one target nucleic acid molecule.

**[0152]** For example, the user may generate a first putative sample tag by using a computer programme to generate a random sequence. The first putative sample tag is added to the group of sample tags. The user may then generate a second putative sample tag in the same manner, and compare the sequence of the second putative sample tag to the first putative sample tag to see whether the second sample tag differs from the first sample tag such that even if the relevant number of low probability mutations were introduced into the second putative sample tag it would still differ from the first putative sample tag. If yes, then the second putative sample tag is added to the group of sample tags. If no, then the second putative sample tag is discarded. This may be repeated for third and further putative sample tags.

**[0153]** As discussed above, it is advantageous for sample tags to be added to at least one target nucleic acid molecule in a method for introducing mutations into at least one target nucleic acid molecule. However, if the sample tags are added prior to the mutations being introduced, this may mean that the sample tags are mutated and cannot then be used to distinguish target nucleic acid molecules that originated from the same or different samples. This can be avoided by designing the sample tags such that even if they are mutated they are sufficiently different from one another for the user to be able to distinguish between them.

**[0154]** The method may further comprise:

1. a.
  1. (i) analysing the method for introducing mutations into at least one target nucleic acid molecule and determining the average number of high probability mutations that take place during the method for introducing mutations into at least one target nucleic acid molecule; and
  2. (ii) determining sequences for a group of sample tags wherein each sample tag differs from substantially all sample tags in the group by more high probability mutation differences than the average number of high probability mutations that take place during the method for introducing mutations into at least one target

nucleic acid molecule.

**[0155]** A low probability mutation may be a transversion mutation or an indel mutation. A high probability mutation may be a transition mutation.

**[0156]** The method may be a computer implemented method.

**[0157]** Also described herein is a computer-readable medium configured to perform the method for designing a group of sample tags suitable for use in a method for introducing mutations into at least one target nucleic acid molecule.

**[0158]** Also described herein is a group of sample tags obtainable by the method for designing sample tags of the disclosure. Optionally, the group of sample tags are obtained by the method for designing sample tags of the disclosure.

#### ***Using dNTPs at unequal concentrations***

**[0159]** The step of amplifying the at least one target nucleic acid using a low bias DNA polymerase in the methods of the invention may be carried out using dNTPs at unequal concentrations.

**[0160]** Described herein is a method for introducing mutations into at least one target nucleic acid molecule comprising:

1. a. providing at least one sample comprising at least one target nucleic acid molecule;  
and
2. b. introducing mutations into the at least one target nucleic acid molecule by amplifying the at least one target nucleic acid molecule using a DNA polymerase to provide a mutated at least one target nucleic acid molecule,

wherein step b. is carried out using dNTPs at unequal concentrations.

**[0161]** In order to be able to amplify the at least one target nucleic acid using a DNA polymerase (such as a low bias DNA polymerase), the target nucleic acid may be exposed to the DNA polymerase and dNTPs under conditions suitable for DNA replication to take place, for example in a PCR machine. If a step of amplifying the at least one target nucleic acid is carried out using dNTPs at unequal concentrations, the target nucleic acid is exposed to a DNA polymerase (such as a low bias DNA polymerase) and dNTPs, wherein the concentrations of the dNTPs are different relative to one another.

**[0162]** The term dNTPs is intended to refer to deoxynucleotides. Specifically, in the context of the present application, the term "dNTPs" is intended to refer to a solution comprising dTTP

(deoxythymidine triphosphate) or dUTP (deoxyuridine), dGTP (deoxyguanine triphosphate), dCTP (deoxycytidine triphosphate), and dATP (deoxyadenosine triphosphate). Optionally, "dNTPs" refers to a solution comprising dTTP (deoxythymidine triphosphate), dGTP (deoxyguanine triphosphate), dCTP (deoxycytidine triphosphate), and dATP (deoxyadenosine triphosphate).

**[0163]** By the phrase "dNTPs at unequal concentrations" is meant that the four dNTPs are present in solution at different concentrations relative to one another. For example, one dNTP may be present at a higher concentration compared to (than) the other three dNTPs, two dNTPs may be present at a higher concentration compared to (than) the other two dNTPs, or three dNTPs may be present at a higher concentration compared to (than) the other one dNTP.

**[0164]** dGTP may be present at a higher concentration compared to (than) dCTP, dTTP and dATP, dGTP may be present at a higher concentration compared to (than) dTTP and dATP, dGTP may be present at a higher concentration compared to (than) dATP, dGTP may be present at a higher concentration compared to (than) dTTP, dCTP may be present at a higher concentration compared to (than) dGTP, dTTP and dATP, dCTP may be present at a higher concentration compared to (than) dTTP and dATP, dCTP may be present at a higher concentration compared to (than) dATP, dCTP may be present at a higher concentration compared to (than) dTTP, dTTP may be present at a higher concentration compared to (than) dGTP, dCTP and dATP, dTTP may be present at a higher concentration compared to (than) dGTP and dCTP, dTTP may be present at a higher concentration compared to (than) dCTP, dTTP may be present at a higher concentration compared to (than) dGTP, dATP may be present at a higher concentration compared to (than) dGTP, dTTP and dCTP, dATP may be present at a higher concentration compared to (than) dGTP and dCTP, dATP may be present at a higher concentration compared to (than) dGTP, dATP may be present at a higher concentration compared to dGTP, dCTP and dATP may be present at a higher concentration compared to (than) dGTP and dCTP, or dGTP and dCTP may be present at a higher concentration compared to (than) dATP and dTTP.

**[0165]** The user may prepare solutions of dNTPs at unequal concentrations in any convenient manner. dATP, dTTP, dGTP and dCTP solutions are readily commercially available, and the user merely needs to mix these in an appropriate ratio.

**[0166]** Optionally, the method:

1. (i) comprises a further step of amplifying the at least one target nucleic acid molecule comprising nucleotide analogs in the absence of nucleotide analogs and the further step of amplifying the at least one target nucleic acid molecule comprising nucleotide analogs in the absence of nucleotide analogs is carried out using dNTPs at unequal concentrations; or
2. (ii) provides a mutated at least one target nucleic acid molecule, and comprises a further step of amplifying the mutated at least one target nucleic acid molecule using the low

bias DNA polymerase and the further step of amplifying the mutated at least one target nucleic acid molecule using the low bias DNA polymerase is carried out using dNTPs at unequal concentrations.

**[0167]** Optionally, introducing mutations into the at least one target nucleic acid molecule by amplifying the at least one target nucleic acid molecule using a DNA polymerase to provide a mutated at least one target nucleic acid molecule is carried out in the presence of a nucleotide analog. Optionally, the method for introducing mutations into at least one target nucleic acid molecule comprises a step of amplifying the mutated at least one target nucleic acid molecule in the absence of the nucleotide analog, and optionally this step is carried out using dNTPs at unequal concentrations.

**[0168]** When a nucleotide analog is used to introduce mutations into at least one target nucleic acid molecule, this will generally involve two amplification steps. In the first amplification step, the nucleotide analog is incorporated into the target nucleic acid molecule (a mutation step). In the second amplification step, the nucleotide analog pairs with a natural nucleotide, thereby introducing a mutation into one strand of the target nucleic acid molecule (a recovery step). When the target nucleic acid molecule is further amplified, this mutation will be transmitted to both strands of the target nucleic acid molecule. Optionally, both the first (mutation) amplification step and the second (recovery) amplification step may be carried out using dNTPs at unequal concentrations. Optionally the dNTPs at unequal concentrations are different in the first (mutation) amplification step and the second (recovery) amplification step. For example, the dNTPs at unequal concentrations may comprise dTTP at a lower concentration than other dNTPs in the first (mutation) amplification step and the dNTPs at unequal concentrations may comprise dATP at a lower concentration than other dNTPs in the second (recovery) amplification step. The step of amplifying the at least one target nucleic acid molecule using a low bias DNA polymerase or steps that provide a mutated at least one target nucleic acid molecule may correspond to one or more "*mutation steps*". A further step of amplifying the at least one target nucleic acid molecule comprising nucleotide analogs in the absence of nucleotide analogs or a further step of amplifying the mutated at least one target nucleic acid molecule may correspond to one or more "*recovery steps*".

**[0169]** Optionally, the nucleotide analog is dPTP.

**[0170]** In an embodiment, dNTPs at unequal concentrations are used to alter the profile of mutations that are introduced. The dNTPs at unequal concentrations are used in methods comprising introducing mutations into at least one target nucleic acid molecule. Thus, the methods result in target nucleic acid molecules comprising mutations (such as the mutated target nucleic acid molecules described herein). The number of mutations, type of mutations, and position of each mutations that are introduced into a given target nucleic acid molecule by the methods may be referred to as the "*profile of mutations*" that is introduced. The term "*type of mutation*" is intended to refer to the nature of the mutation, *i.e.* is it a substitution mutation,

an addition mutation or a deletion mutation, and if it is a substitution mutation what was the starting nucleotide and what was the starting nucleotide mutated to (e.g. an A to G mutation has an A starting nucleotide which is mutated to G)?

**[0171]** The user may determine the "*profile of mutations*" that is introduced by a given method by replicating a test target nucleic acid molecule, then subjecting some of the replicates to the methods comprising introducing mutations of the invention, but reserving some of the replicates (without mutating them). The user may then sequence the replicates that have been subjected to the methods comprising introducing mutations of the invention, and the reserved replicates. Finally, the user can align the sequences of the replicates that have been subjected to the methods comprising introducing mutations of the invention, and the reserved replicates to determine the number of mutations, type of mutations and position of each mutation that have been introduced. Alternatively, the user may use a test target nucleic acid molecule of known sequence. The user will then merely need to subject the test target nucleic acid molecule to the methods comprising introducing mutations of the invention, and then sequence the resultant mutated target nucleic acid molecule to see what profile of mutations has been introduced.

**[0172]** The user may wish to alter the mutation profile in a number of ways. For example, as discussed above, it is advantageous to be able to reduce mutation bias. Accordingly, in an embodiment, dNTPs at unequal concentrations are used to reduce bias in the profile of mutations that are introduced. In a further embodiment, the method is a method for introducing mutations in a low bias mutation profile.

**[0173]** The present application demonstrates that using dNTPs at unequal concentrations can be used to reduce bias in the profile of mutations that are introduced. For example, if a DNA polymerase (such as a low bias DNA polymerase described above) is used to mutate a target nucleic acid molecule, and introduces a higher number of G to A mutations compared to other mutations, the user can reduce the concentration of dATPs relative to other dNTPs, and this may decrease the frequency at which A nucleotides are incorporated in place of dGTPs and so decrease the number of G to A mutations.

**[0174]** Similarly, if a nucleotide analog is used when introducing mutations into a target nucleic acid molecule, altering the relative concentrations of the dNTPs can be used to alter the mutation profile. For example, dPTP can be used to introduce G to A, C to T, A to G and T to C mutations. As described in more detail above, dPTP can replace a T nucleotide or a C nucleotide, and depending on whether the dPTP is in its amino or imino form, it can subsequently pair with an A nucleotide or a G nucleotide. This leads to two scenarios. In the first scenario, the dPTP replaces T in (for example) the sense strand (mutation step), it can then pair with A (no mutation) or G (A to G mutation) in the anti sense strand. If dPTP replaces T and pairs with G in the antisense strand, the mutant G will pair with a C to introduce a T to C mutation in a replicate of the sense strand (recovery step). Conversely, dPTP may replace T in the antisense strand, which may lead to an A to G mutation in the sense strand and a T to C mutation in a replicate of the antisense strand. In the second scenario, the dPTP replaces C in

the (for example) sense strand, it can then pair with A (G to A mutation) or G (no mutation) in the antisense strand (mutation step). If dPTP replaces C and pairs with A in the antisense strand, the mutant A will pair with a T to introduce a C to T mutation in a replicate of the sense strand (recovery step). Conversely, dPTP may replace C in the antisense strand, which may lead to a G to A mutation in the sense strand and a C to T mutation in a replicate of the antisense strand.

**[0175]** The present application demonstrates that if the rate of G to A and C to T mutations is higher than the rate of A to G and T to C mutations, then reducing the concentration of dTTPs compared to the other dNTPs (and preferably compared to the concentration of dCTP) will encourage dPTP to be incorporated in place of dTTP, increasing the instances of the first scenario set out above relative to the second scenario, meaning that the A to G and T to C mutations introduced in the first scenario will be increased. Similarly, the present application demonstrates that if the level of dATPs is reduced during the recovery step, then the level of G to A and C to T mutations increases. This is because in scenario 2 above, if dATP is present at a lower concentration compared to the other dNTPs (and preferably compared to the concentration of dGTP), this will mean that dPTP that has incorporated in place of a C nucleotide will pair more frequently with G and fewer G to A or C to T mutations will be introduced. The two scenarios are set out in Figure 7.

**[0176]** Even the low bias DNA polymerases disclosed herein introduce mutations into a target nucleic acid molecule with a small bias. The present application demonstrates that using unequal concentrations of dNTPs with a low bias DNA polymerase can virtually eliminate any mutation bias.

**[0177]** Based on the information provided in the present application, it is within the abilities of the skilled person to determine how altering the concentrations of various dNTPs will affect the mutation profile depending on whether a nucleotide analog is used, and if so which one. Accordingly, in some embodiments, the methods which use dNTPs at unequal concentrations comprise a step of identifying a dNTP whose level should be increased or decreased in order to reduce bias in the profile of mutations that are introduced.

**[0178]** Optionally, the dNTPs at unequal concentrations comprise dTTP at a lower concentration than other dNTPs. As described above, this can increase the rate of T to C and A to G mutations that are introduced when dPTP is used as a nucleotide analog. Optionally, the dNTPs at unequal concentrations comprise dTTP at a concentration less than 75%, less than 70%, less than 60%, less than 55%, between 25% and 75%, between 25% and 70, between 25% and 60%, or around 50% of the concentration of dATP, dCTP or dGTP. Optionally, the dNTPs at unequal concentrations comprise dTTP at a concentration less than 60% of the concentration of dCTP. Optionally, the dNTPs at unequal concentrations comprise dTTP at a concentration between 25% and 60% of the concentration of dCTP.

**[0179]** Optionally, the dNTPs at unequal concentrations comprises dATP at a lower concentration compared to other dNTPs. As described above, this can decrease the rate of G

to A or C to T mutations that are introduced when dPTP is used as a nucleotide analog. Optionally, the dNTPs at unequal concentrations comprises dATP at a concentration less than 75%, less than 70%, less than 60%, less than 55%, between 25% and 75%, between 25% and 70, between 25% and 60%, or around 50% of the concentration of dTTP, dCTP or dGTP. Optionally, the dNTPs at unequal concentrations comprises dATP at a concentration less than 75%, less than 70%, less than 60%, less than 55%, between 25% and 75%, between 25% and 70, between 25% and 60%, or around 50% of the concentration of dGTP. Optionally, the dNTPs at unequal concentrations comprises dATP at a concentration less than 60% of the concentration of dGTP. Optionally, the dNTPs at unequal concentrations dNTPs comprises dATP at a concentration between 25% and 60% of the concentration of dGTP.

**[0180]** As set out in the two scenarios above, when using dPTP as a nucleotide analog, reducing dTTPs increases T to C and A to G mutations by encouraging the replacement of T nucleotides in the target nucleic acid molecule with dPTP. Thus, dNTPs at unequal concentrations which comprise dTTP at a lower concentration than other dNTPs are preferably used in a mutagenesis step (for example a step of PCR in the presence of dPTPs). Similarly, when using dPTP as a nucleotide analog, reducing dATPs reduces the number of dPTPs that have replaced C nucleotides and pair with dATP and so increases G to A and C to T mutations. Since dPTP pairing with dATP tends to occur during a recovery step, reducing dATPs during the recovery step increases the number of G to A and C to T mutations. Optionally, therefore, the step of amplifying the at least one target nucleic acid molecule comprising nucleotide analogs in the absence of nucleotide analogs or amplifying the mutated at least one target nucleic acid molecule in the absence of the nucleotide analog is carried out using dNTPs at unequal concentrations, and the dNTPs at unequal concentrations comprises dATP at a lower concentration compared to other dNTPs.

## **Examples**

### **Example 1 - Mutating nucleic acid molecules using PrimeStar GXL of other polymerases**

**[0181]** DNA molecules were fragmented to the appropriate size (e.g. 10 kb) and a defined sequence priming site (adapter) was attached on each end using tagmentation.

**[0182]** The first step is a tagmentation reaction to fragment the DNA. 50 ng high molecular weight genomic DNA in 4µl or less volume of one or more bacterial strains was subjected to tagmentation under the following conditions. 50 ng DNA is combined with 4 µl Nextera Transposase (diluted to 1:50), and 8 µl 2X tagmentation buffer (20mM Tris [pH7.6], 20mM MgCl, 20% (v/v) dimethylformamide) in a total volume of 16 µl. The reaction was incubated at 55°C for 5 minutes, 4µl of NT buffer (or 0.2% SDS) was added to the reaction and the reaction was incubated at room temperature for 5 minutes.

[0183] The fragmentation reaction was cleaned using SPRIselect beads (Beckman Coulter) following the manufacturer's instructions for a left side size selection using 0.6 volume of beads, and the DNA was eluted in molecular grade water.

[0184] This was followed by PCR with a combination of standard dNTPs and dPTP for a limited 6 cycles. Using Primestar GXL, 12.5 ng of fragmented and purified DNA was added to a total reaction volume of 25µl, containing 1 x GXL buffer, 200 µM each of dATP, dTTP, dGTP and dCTP, as well as 0.5 mM dPTP, and 0.4 µM custom primers (Table 2).

Table 2:

i7 custom index primer	CAAGCAGAAGACGGCA	NNN	XXX	GTCTCGTGG
	TACGAGAT	NNN	X	GCTCGG
i5 custom index primer	AATGATACGGCGACCA	XXX	NNN	TCGTCGGCA
	CCGAGATCTACAC	XXX	N	GCGTC

Table 2. Custom primers used for mutagenesis PCR on 10kbp templates. XXXXXX is a defined, sample-specific 6-8nt barcode sequence. NNNNNN is a 6nt region of random nucleotides.

[0185] The reaction was subject to the following thermal cycling in the presence of Primestar GXL. Initial gap extension at 68°C for 3 minutes, followed by 6 cycles of 98°C for 10 seconds, 55°C for 15 seconds and 68°C for 10 minutes.

[0186] The next stage is a PCR without dPTP, to remove dPTP from the templates and replace them with a transition mutation ("recovery PCR"). PCR reactions were cleaned with SPRIselect beads to remove excess dPTP and primers, then subjected to a further 10 rounds (minimum 1 round, maximum 20) of amplification using primers that anneal to the fragment ends introduced during the dPTP incorporation cycles (Table 3).

Table 3

i7 flow cell primer	CAAGCAGAAGAC GGCATACGA
i5 flow cell primer	AATGATACGGCG ACCACCGA

[0187] This was followed by a gel extraction step to size select amplified and mutated fragments in a desired size range, for example from 7-10 kb. The gel extraction can be done manually or via an automated system such as a BluePippin. This was followed by an additional round of PCR for 16-20 cycles ("*enrichment PCR*").

[0188] After amplifying a defined number of long mutated templates, random fragmentation of the templates was carried out to generate a group of overlapping shorter fragments for sequencing. Fragmentation was performed by tagmentation.

[0189] Long DNA fragments from the previous step were subject to a standard tagmentation reaction (e.g. Nextera XT or Nextera Flex), except that the reaction was split into three pools for the PCR amplification. This enables selective amplification of fragments derived from each end of the original template (including the sample barcode) as well as internal fragments from the long template that have been newly tagmented at both ends. This effectively creates three pools for sequencing on an Illumina instrument (e.g. MiSeq or HiSeq).

[0190] The method was repeated using a standard Taq (Jena Biosciences) and a blend of Taq and a proofreading polymerase (DeepVent) called LongAmp (New England Biolabs).

[0191] The data obtained from this experiment is depicted in Figure 1. No dPTP was used as a control. Reads were mapped against the *E. coli* genome, and a median mutation rate of ~ 8% was achieved.

### **Example 2 - Comparison of mutation frequencies of different DNA polymerases**

[0192] Mutagenesis was performed with a range of different DNA polymerases (Table 4). Genomic DNA from *E. coli* strain MG1655 was tagmented to produce long fragments and bead cleaned as described in the method of Example 1. This was followed by "mutagenesis PCR" for 6 cycles in the presence of 0.5 mM dPTP, SPRIselect bead purification and an additional 14-16 cycles of "recovery PCR" in the absence of dPTP. The resulting long mutated templates were then subjected to a standard tagmentation reaction (see Example 1) and "internal" fragments were amplified and sequenced on an Illumina MiSeq instrument.

[0193] The mutation rates are described in Table 4, which normalized frequencies of base substitution via dPTP mutagenesis reactions as measured using Illumina sequencing of DNA from the known reference genome. For Taq polymerase, only ~12% of mutations occur at template G+C sites, even when used in buffer optimised for Thermococcus polymerases. Thermococcus-like polymerases result in 58-69% of mutations at template G+C sites, while polymerase derived from Pyrococcus gives 88% of mutations at template G+C sites.

[0194] Enzymes were obtained from Jena Biosciences (Taq), Takara (Primestar variants),

Merck Millipore (KOD DNA Polymerase) and New England Biolabs (Phusion).

[0195] Taq was tested with the supplied buffer, and also with Primestar GXL Buffer (Takara) for this experiment. All other reactions were carried out with the standard supplied buffer for each polymerase.

**Table 4**

Polymerase <sup>1</sup>	Origin	Mutation frequency (% of total observed mutations)				
		A → G	T → C	G → A	C → T	Other (transversion)
Taq (standard buffer)	<i>Thermus aquaticus</i>	43.1	41.7	6.3	6.1	2.7
Taq ( <i>Thermococcus</i> buffer <sup>2</sup> )	<i>Thermus aquaticus</i>	48.9	47.5	2.9	0.7	0.0
Primestar GXL	<i>Thermococcus</i>	21.5	20.1	29.5	28.9	0.0
Primestar HS	<i>Thermococcus</i>	16.3	15.2	30.1	38.4	0.0
Primestar Max	<i>Thermococcus</i>	16.5	14.6	33.2	35.7	0.0
KOD DNA polymerase	<i>Thermococcus</i>	20.5	16.1	31.8	31.5	0.0
Phusion	<i>Pyrococcus</i>	5.4	6.4	44.1	44.1	0.0

### **Example 3 - determining dPTP mutagenesis rates**

[0196] We performed dPTP mutagenesis on a range of genomic DNA samples with different levels of G+C content (33-66%) using a *Thermococcus* polymerase (Primestar GXL; Takara) under a single set of reaction conditions. Mutagenesis and sequencing was performed as described in the method of example 3, except that 10 cycles of "recovery PCR" were performed. As predicted, mutation rates were roughly similar between samples (median rate 7-8%) despite the diversity of G+C content (figure 2).

### **Example 4 - measuring template amplification bias**

[0197] Template amplification bias was measured for two polymerases: Kapa HiFi, which is a proofreading polymerase commonly used in Illumina sequencing protocols, and PrimeStar GXL, which is a KOD family polymerase known for its ability to amplify long fragments. In the first experiment Kapa HiFi was used to amplify a limited number of *E. coli* genomic DNA templates with sizes around 2kbp. The ends of these amplified fragments were then sequenced. A similar experiment was done with PrimeStar GXL on fragments around 7-10kbp from *E. coli*. The positions of each end sequence read were determined by mapping to the *E.*

*coli* reference genome. The distances between neighboring fragment ends was measured. These distances were compared to a set of distances randomly sampled from the uniform distribution. The comparison was carried out via the nonparametric Kolmogorov-Smirnov test, D. When two samples come from the same distribution, the value of D approaches zero. For the low bias PrimeStar polymerase, we observed D=0.07 when measured on 50,000 fragment ends, compared to a uniform random sample of 50,000 genomic positions. For the Kapa HiFi polymerase we observed D=0.14 on 50,000 fragment ends.

**Example 5 - Using two identical primer binding sites and a single primer sequence for preferential amplification of longer templates**

[0198] As described above, tagmentation can be used to fragment DNA molecules and simultaneously introduce primer binding sites (adapters) onto the ends of the fragments. The Nextera tagmentation system (Illumina) utilises transposase enzymes loaded with one of two unique adapters (referred to here as X and Y). This generates a random mixture of products, some with identical end sequences (X-X, Y-Y) and some with unique ends (X-Y). Standard Nextera protocols use two distinct primer sequences to selectively amplify "X-Y" products containing different adapters on each end (as required for sequencing with Illumina technology). However, it is also possible to use a single primer sequence to amplify "X-X" or "Y-Y" fragments with identical end adapters.

[0199] To generate long mutated templates containing identical end adapters, 50 ng of high molecular weight genomic DNA (*E. coli* strain MG1655) was first subjected to tagmentation and then cleaned with SPRIselect beads as described in Example 1. This was followed by 5 cycles of "mutagenesis PCR" with a combination of standard dNTPs and dPTP, which was performed as detailed in Example 1 except that a single primer sequence was used (Table 5).

[0200] The PCR reaction was cleaned with SPRIselect beads to remove excess dPTP and primers, then subjected to a further 10 cycles of "recovery PCR" in the absence of dPTP to replace dPTP in the templates with transition mutations. Recovery PCR was performed with a single primer that anneals to the fragment ends introduced during the dPTP incorporation cycles, thereby enabling selective amplification of mutated templates generated in the previous PCR step.

Table 5:

Primer name	Step	Sequence								
single_mut	mutagenesis	<table border="1"> <tr> <td>TCGGTCTGCGCCTC</td> <td>NNN</td> <td>XXXXXXXX</td> <td>GTCTCGTGG</td> </tr> <tr> <td>TAGC</td> <td></td> <td>XXXXXX</td> <td>GCTCGGAG</td> </tr> </table>	TCGGTCTGCGCCTC	NNN	XXXXXXXX	GTCTCGTGG	TAGC		XXXXXX	GCTCGGAG
TCGGTCTGCGCCTC	NNN	XXXXXXXX	GTCTCGTGG							
TAGC		XXXXXX	GCTCGGAG							
single_rec	recovery	<table border="1"> <tr> <td>CAAGCAGAAGACG</td> <td>TCGGTCTGCGCCTCTAGC</td> </tr> <tr> <td>GCATACGAGAT</td> <td></td> </tr> </table>	CAAGCAGAAGACG	TCGGTCTGCGCCTCTAGC	GCATACGAGAT					
CAAGCAGAAGACG	TCGGTCTGCGCCTCTAGC									
GCATACGAGAT										

Primer name	Step	Sequence
-------------	------	----------

**Table 5.** Primers used to generate mutated templates with the same basic adapter structure on both ends. Primer "single mut" was used for mutagenesis PCR on DNA fragments generated by Nextera tagmentation. This primer contains a 5' portion that introduces an additional primer binding site at the fragment ends. Primer "single rec" is capable of annealing to this site, and was used during recovery PCR to selectively amplify mutated templates generated with the single\_mut primer. XXXXXXXXXXXXXXX is a defined, sample-specific 13nt barcode sequence. NNN is a 3nt region of random nucleotides.

**[0201]** As a control, mutated templates with different adapters on each end were generated using an identical protocol to that described above, except that two distinct primer sequences were used during both mutagenesis PCR (shown in Table 2) and recovery PCR (Table 3). Final PCR products were cleaned with SPRIselect beads and analysed on a High Sensitivity DNA Chip using the 2100 Bioanalyzer System (Agilent). As shown in Figure xxx, the templates generated with identical end adapters were significantly longer on average than the control sample containing dual adapters. Control templates could be detected down to a minimum size of ~800 bp, while no templates below 2000 bp were observed for the single adapter sample.

**[0202]** Mutated templates with identical end adapters (blue) and control templates with dual adapters were run on an Agilent 2100 Bioanalyzer (High Sensitivity DNA Kit) to compare size profiles. The use of identical end adapters inhibits the amplification of templates < 2kbp. The data is presented in Figure 6.

### **Example 8 - Further reducing the mutation bias of *Thermococcus* polymerases by altering natural dNTP levels during PCR**

**[0203]** Although *Thermococcus* polymerases generate a much more balanced mutation profile compared to other DNA polymerases, they do exhibit a small amount of bias towards mutations at G and C sites (see Table 4). To eliminate this residual bias, we tested the effect of altering the concentrations of natural dNTPs during the mutagenesis and recovery PCR steps to influence the relative incorporation rates of the different nucleotides.

**[0204]** First, long mutated templates were prepared from bacterial genomic DNA (*E. coli* strain MG1655) using the approach outlined in Example 5, except that the concentration of individual nucleotides in the PCR reactions were varied. This was achieved by adding individual solutions of the four natural nucleotides (purchased from New England Biolabs) separately to the PCR mixture, either at a standard final concentration of 200  $\mu$ M or at a lower concentration of 160  $\mu$ M (80% relative to standard) or 100  $\mu$ M (50%). Only one nucleotide was varied per reaction. As a control, all natural nucleotides were added to the same final concentration of 200  $\mu$ M, using an equimolar dNTP mixture provided with the Primestar GXL polymerase (Takara). Five mutagenesis PCR cycles and twelve recovery cycles were performed using primers shown in

Table 5. The resulting long mutated templates were then subjected to a standard tagmentation reaction (see Example 1) and "internal" fragments were amplified and sequenced on an Illumina MiSeq instrument. Mutation frequencies were determined by comparison against the known reference sequence.

[0205] As shown in Table 6 changes in the concentration of individual dNTPs during mutagenesis and/or recovery PCR altered the observed profile of mutations. Importantly, limiting the amount of dTTP by 50% during mutagenesis was found to produce virtually identical mutation frequencies for each nucleotide (Table 3). This confirms that the residual mutation bias of *Thermococcus* polymerases can be eliminated through changes in dNTP levels.

**Table 6.**

Treatment	Mutation frequency (% of total observed mutations)			
	A → G	T → C	G → A	C → T
Equimolar dNTP control	17.4	16.8	32.1	33.7
80% dTTP (mutagenesis)	13.9	13.8	36.1	36.2
50% dTTP (mutagenesis)	23.7	24.9	25.3	26.2
80% dATP (recovery)	13.4	12.5	36.7	37.3
50% dATP (recovery)	18.9	19.1	31.4	30.6
80% dTTP (mutagenesis) and 80% dATP (recovery)	17.8	15.0	34.0	33.2
50% dTTP (mutagenesis) and 50% dATP (recovery)	34.4	34.7	15.4	15.5

## SEQUENCE LISTING

[0206]

<110> LONGAS TECHNOLOGIES PTY LTD

<120> ENZYME

<130> N411620WO

<140> TBC

<141> 2019-02-19

<150> GB 1802744.1

<151> 2018-02-20

<160> 136

<170> PatentIn version 3.5

<210> 1

<211> 2325

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA polymerase from Thermococcus sp. KS-1

<400> 1

```

atgatcctcg acactgacta cataactgag aatggaaaac ccgtcataag gattttcaag      60
aaggagaacg gcgagtttaa gattgagtac gataggactt ttgaacccta catttacgcc      120
ctcctgaagg acgattctgc cattgaggag gtcaagaaga taaccgccga gaggcacgga      180
acggttgtaa cggtaagcg  ggctgaaaag gttcagaaga agttcctcgg gagaccagtt      240
gaggtctgga aactctactt tactcaccct caggacgtcc cagcgataag ggacaagata      300
cgagagcatc cagcagttat tgacatctac gagtacgaca tacccttcgc caagcgctac      360
ctcatagaca agggattagt gccaatggaa ggcgacgagg agctgaaaat gcttgccttt      420
gatatcgaga cgctctacca tgagggcgag gagttcggcg aggggccaat ccttatgata      480
agctacgccg acgaggaagg ggcagggtg ataacgtgga agaacgcgga tctgccctac      540
gttgacgtcg tctcgacgga gagggagatg ataaagcgct tcctaaaggt ggtcaaagag      600
aaagatcctg acgtcctaataacctacaac ggcgacaact tcgacttcgc ctacctaaaa      660
aaacgctgtg aaaagcttg  aataaaacttc acgctcgaa  gggacggaag cgagccgaag      720
attcagagga tgggcgacag gtttgccgtc gaagtgaagg gacggataca cttcgatctc      780
tatcctgtga taagacggac gataaacctg cccacataca cgcttgaggc cgtttatgaa      840
gccgtcttcg gtcagccgaa ggagaaggtc tacgctgagg agatagctac agcttgggag      900
agcgggtaag gccttgagag agtagccaga tactcgatgg aagatgcgaa ggtcacatac      960
gagcttggga aggagttttt ccctatggag gccagcttt  ctcgcttaat cggccagtcc     1020
ctctgggacg tctcccgtc  cagcactggc aacctcgttg agtggttcct cctcaggaag     1080
gcctacgaga ggaatgagct ggccccgaac aagcccgatg aaaaggagct ggccagaaga     1140
cgacagagct atgaaggagg ctatgtaaaa gagcccgaga gagggttg  ggagaacata     1200
gtgtacctag attttagatc tctgtacccc tcaatcatca tcaccacaa  cgtctcgccg     1260
gatactctca acaggaagg  atgcaaggaa tatgacgttg cccccagg  cggtcacccg     1320
ttctgcaagg acttcccagg atttatcccg agcctgcttg gagacctcct agaggagagg     1380
cagaagataa agaagaagat gaaggccacg attgaccoga tcgagaggaa gctcctcgat     1440
tacaggcaga gggccatcaa gatcctggcc aacagctact acggttacta cggctatgca     1500
agggcgcgct ggtactgcaa ggagtgtgca gagagcgtaa cggcctgggg aagggagtac     1560
ataacgatga ccatcagaga gatagaggaa aagtacggct ttaaggtaat ctacagcgac     1620
accgacggat tttttgccac aatacctgga gccgatgctg aaaccgtcaa aaagaaggcg     1680

```

atggagttcc tcaagtatat caacgcaaaa ctcccgggcg cgcttgagct cgagtacgag 1740  
 ggcttctaca aacgcggctt ctctgtcacg aagaagaagt acgcggtgat agacgaggaa 1800  
 ggcaagataa caacgcgcgg acttgagatt gtgaggcgcg actggagcga gatagcgaaa 1860  
 gagacgcagg cgagggttct tgaagctttg ctaaaggacg gtgacgtcga gaaggccgtg 1920  
 aggatagtca aagaagttac cgaaaagctg agcaagtacg aggttccgcc ggagaagctg 1980  
 gtgatccacg agcagataac gagggattta aaggactaca aggcaaccgg tccccagtt 2040  
 gccgttgcca agaggttggc gcgcgagagga gtcaaaatac gccctggaac ggtgataagc 2100  
 tacatcgtgc tcaagggctc tgggaggata ggcgacaggg cgataccggt cgacgagttc 2160  
 gacccgacga agcacaagta cgacgccgag tactacattg agaaccaggt tctcccagcc 2220  
 gttgagagaa ttctgagagc ctctcggttac cgcaaggaag acctgcgcta ccagaagacg 2280  
 agacaggttg gtctgggagc ctggctgaag ccgaagggaa cttga 2325

<210> 2

<211> 774

<212> PRT

<213> Artificial Sequence

<220>

<223> DNA polymerase from Thermococcus sp. KS-1

<400> 2

Met Ile Leu Asp Thr Asp Tyr Ile Thr Glu Asn Gly Lys Pro Val Ile  
 1 5 10 15  
 Arg Ile Phe Lys Lys Glu Asn Gly Glu Phe Lys Ile Glu Tyr Asp Arg  
 20 25 30  
 Thr Phe Glu Pro Tyr Ile Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
 35 40 45  
 Glu Glu Val Lys Lys Ile Thr Ala Glu Arg His Gly Thr Val Val Thr  
 50 55 60  
 Val Lys Arg Ala Glu Lys Val Gln Lys Lys Phe Leu Gly Arg Pro Val  
 65 70 75 80  
 Glu Val Trp Lys Leu Tyr Phe Thr His Pro Gln Asp Val Pro Ala Ile  
 85 90 95  
 Arg Asp Lys Ile Arg Glu His Pro Ala Val Ile Asp Ile Tyr Glu Tyr  
 100 105 110  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Val Pro  
 115 120 125  
 Met Glu Gly Asp Glu Glu Leu Lys Met Leu Ala Phe Asp Ile Glu Thr  
 130 135 140  
 Leu Tyr His Glu Gly Glu Glu Phe Ala Glu Gly Pro Ile Leu Met Ile  
 145 150 155 160

Ser Tyr Ala Asp Glu Glu Gly Ala Arg Val Ile Thr Trp Lys Asn Ala  
 165 170 175

Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Arg Glu Met Ile Lys  
 180 185 190

Arg Phe Leu Lys Val Val Lys Glu Lys Asp Pro Asp Val Leu Ile Thr  
 195 200 205

Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Lys Arg Cys Glu  
 210 215 220

Lys Leu Gly Ile Asn Phe Thr Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 225 230 235 240

Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 245 250 255

His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Ile Asn Leu Pro Thr  
 260 265 270

Tyr Thr Leu Glu Ala Val Tyr Glu Ala Val Phe Gly Gln Pro Lys Glu  
 275 280 285

Lys Val Tyr Ala Glu Glu Ile Ala Thr Ala Trp Glu Ser Gly Glu Gly  
 290 295 300

Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Tyr  
 305 310 315 320

Glu Leu Gly Lys Glu Phe Phe Pro Met Glu Ala Gln Leu Ser Arg Leu  
 325 330 335

Ile Gly Gln Ser Leu Trp Asp Val Ser Arg Ser Ser Thr Gly Asn Leu  
 340 345 350

Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 355 360 365

Pro Asn Lys Pro Asp Glu Lys Glu Leu Ala Arg Arg Arg Gln Ser Tyr  
 370 375 380

Glu Gly Gly Tyr Val Lys Glu Pro Glu Arg Gly Leu Trp Glu Asn Ile  
 385 390 395 400

Val Tyr Leu Asp Phe Arg Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 405 410 415

Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Lys Glu Tyr Asp  
 420 425 430

Val Ala Pro Gln Val Gly His Arg Phe Cys Lys Asp Phe Pro Gly Phe  
 435 440 445

Ile Pro Ser Leu Leu Gly Asp Leu Leu Glu Glu Arg Gln Lys Ile Lys  
 450 455 460

Lys Lys Met Lys Ala Thr Ile Asp Pro Ile Glu Arg Lys Leu Leu Asp  
 465 470 475 480

Tyr Arg Gln Arg Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr  
 485 490 495

Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Lys Glu Cys Ala Glu Ser  
 500 505 510

Val Thr Ala Trp Gly Arg Glu Tyr Ile Thr Met Thr Ile Arg Glu Ile  
 515 520 525

Glu Glu Lys Tyr Gly Phe Lys Val Ile Tyr Ser Asp Thr Asp Gly Phe  
 530 535 540

Phe Ala Thr Ile Pro Gly Ala Asp Ala Glu Thr Val Lys Lys Lys Ala  
 545 550 555 560

Met Glu Phe Leu Lys Tyr Ile Asn Ala Lys Leu Pro Gly Ala Leu Glu  
 565 570 575

Leu Glu Tyr Glu Gly Phe Tyr Lys Arg Gly Phe Phe Val Thr Lys Lys  
 580 585 590

Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 595 600 605

Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln Ala  
 610 615 620

Arg Val Leu Glu Ala Leu Leu Lys Asp Gly Asp Val Glu Lys Ala Val  
 625 630 635 640

Arg Ile Val Lys Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 645 650 655

Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Asp Leu Lys Asp  
 660 665 670

Tyr Lys Ala Thr Gly Pro His Val Ala Val Ala Lys Arg Leu Ala Ala  
 675 680 685

Arg Gly Val Lys Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 690 695 700

Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 705 710 715 720

Asp Pro Thr Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn Gln  
 725 730 735

Val Leu Pro Ala Val Glu Arg Ile Leu Arg Ala Phe Gly Tyr Arg Lys  
 740 745 750



```

agggcgaggt ggtactgcag ggagtgcgcg gagagcgta ccgcctgggg cagggagtac      1560
atcgataggg tcatcagga gctcgaggag aagttcggct tcaagtgct ctacgcggac      1620
acggacggac tgcacgccac gatccccggg gcggaccccg ggaccgtcaa ggagaggcg      1680
agggggttcc tgagatacat caaccccaag ctccccggcc tcctggagct cgagtacgag      1740

gggttctacc tgagggggtt cttcgtgacg aagaagaagt acgcggtcat agacgaggag      1800
ggcaagataa ccacgcgcg cctcgagata gtcaggcggg actggagcga ggtggccaag      1860
gagacgcagg cgagggtcct ggaggcgata ctgaggcacg gtgacgtcga ggaggccgtt      1920
agaatcgtca gggaggtaac cgaaaagctg agcaagtacg aggttccgcc ggagaaactg      1980
gtgatccacg agcagataac gagggatttg agggactaca aagccacggg accgcacgtg      2040
gcggtggcga agcgcctggc cgggaggggg gtaaggatac gccccgggac ggtgataagc      2100
tacatcgtcc tcaagggctc cggaaggata ggggacaggg cgattccctt cgacgagttc      2160
gacccgacta agcacaggta cgacgccgac tactacatcg agaaccaggt tctgccagcc      2220
gtcgagagga tcctgaaggc cttcggctac cgcaaggagg acctgaaata ccagaagacg      2280
aggcaggtgg gcctgggtgc gtggctcaac gcggggaagg ggtga                        2325

```

<210> 4

<211> 774

<212> PRT

<213> Artificial Sequence

<220>

<223> DNA polymerase from Thermococcus celer

<400> 4

```

Met Ile Leu Asp Ala Asp Tyr Ile Thr Glu Asp Gly Lys Pro Val Val
1           5           10           15

```

```

Arg Ile Phe Arg Lys Glu Lys Gly Glu Phe Arg Ile Asp Tyr Asp Arg
          20           25           30

```

```

Asp Phe Glu Pro Tyr Ile Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile
          35           40           45

```

```

Glu Glu Val Lys Arg Ile Thr Val Glu Arg His Gly Lys Ala Val Arg
          50           55           60

```

```

Val Lys Arg Val Glu Lys Val Glu Lys Lys Phe Leu Asn Arg Pro Ile
65           70           75           80

```

```

Glu Val Trp Lys Leu Tyr Phe Asn His Pro Gln Asp Val Pro Ala Ile
          85           90           95

```

```

Arg Asp Glu Ile Arg Lys His Pro Ala Val Val Asp Ile Tyr Glu Tyr
          100          105          110

```

```

Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Val Pro
          115          120          125

```

Met Glu Gly Glu Glu Glu Leu Lys Leu Met Ala Phe Asp Ile Glu Thr  
 130 135 140

Leu Tyr His Glu Gly Asp Glu Phe Gly Glu Gly Pro Ile Leu Met Ile  
 145 150 155 160

Ser Tyr Ala Asp Gly Asp Gly Ala Arg Val Ile Thr Trp Lys Lys Ile  
 165 170 175

Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Lys Glu Met Ile Lys  
 180 185 190

Arg Phe Leu Gln Val Val Lys Glu Lys Asp Pro Asp Val Leu Val Thr  
 195 200 205

Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Arg Arg Ser Glu  
 210 215 220

Glu Leu Gly Leu Lys Phe Ile Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 225 230 235 240

Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 245 250 255

His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Val Asn Leu Pro Thr  
 260 265 270

Tyr Thr Leu Glu Ala Val Tyr Glu Ala Ile Phe Gly Arg Pro Lys Glu  
 275 280 285

Lys Val Tyr Ala Gly Glu Ile Val Glu Ala Trp Glu Thr Gly Glu Gly  
 290 295 300

Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Phe  
 305 310 315 320

Glu Leu Gly Arg Glu Phe Phe Pro Met Glu Ala Gln Leu Ser Arg Leu  
 325 330 335

Ile Gly Gln Gly Leu Trp Asp Val Ser Arg Ser Ser Thr Gly Asn Leu  
 340 345 350

Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 355 360 365

Pro Asn Lys Pro Ser Gly Arg Glu Val Glu Ile Arg Arg Arg Gly Tyr  
 370 375 380

Ala Gly Gly Tyr Val Lys Glu Pro Glu Arg Gly Leu Trp Glu Asn Ile  
 385 390 395 400

Val Tyr Leu Asp Phe Arg Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 405 410 415

Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Glu Asn Tyr Asp  
 420 425 430

Val Ala Pro Gln Val Gly His Lys Phe Cys Lys Asp Phe Pro Gly Phe  
 435 440 445

Ile Pro Ser Leu Leu Gly Gly Leu Leu Glu Glu Arg Gln Lys Ile Lys  
 450 455 460

Arg Arg Met Lys Ala Ser Val Asp Pro Val Glu Arg Lys Leu Leu Asp  
 465 470 475 480

Tyr Arg Gln Arg Ala Ile Lys Ile Leu Ala Asn Ser Phe Tyr Gly Tyr  
 485 490 495

Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Arg Glu Cys Ala Glu Ser  
 500 505 510

Val Thr Ala Trp Gly Arg Glu Tyr Ile Asp Arg Val Ile Arg Glu Leu  
 515 520 525

Glu Glu Lys Phe Gly Phe Lys Val Leu Tyr Ala Asp Thr Asp Gly Leu  
 530 535 540

His Ala Thr Ile Pro Gly Ala Asp Ala Gly Thr Val Lys Glu Arg Ala  
 545 550 555 560

Arg Gly Phe Leu Arg Tyr Ile Asn Pro Lys Leu Pro Gly Leu Leu Glu  
 565 570 575

Leu Glu Tyr Glu Gly Phe Tyr Leu Arg Gly Phe Phe Val Thr Lys Lys  
 580 585 590

Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 595 600 605

Glu Ile Val Arg Arg Asp Trp Ser Glu Val Ala Lys Glu Thr Gln Ala  
 610 615 620

Arg Val Leu Glu Ala Ile Leu Arg His Gly Asp Val Glu Glu Ala Val  
 625 630 635 640

Arg Ile Val Arg Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 645 650 655

Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Asp Leu Arg Asp  
 660 665 670

Tyr Lys Ala Thr Gly Pro His Val Ala Val Ala Lys Arg Leu Ala Gly  
 675 680 685

Arg Gly Val Arg Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 690 695 700

Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 705 710 715 720

Asp Pro Thr Lys His Arg Tyr Asp Ala Asp Tyr Tyr Ile Glu Asp Gln

Val Leu Pro Ala Val Glu Arg Ile Leu Lys Ala Phe Gly Tyr Arg Lys  
 725 730 735

Val Leu Pro Ala Val Glu Arg Ile Leu Lys Ala Phe Gly Tyr Arg Lys  
 740 745 750

Glu Asp Leu Lys Tyr Gln Lys Thr Arg Gln Val Gly Leu Gly Ala Trp  
 755 760 765

Leu Asn Ala Gly Lys Gly  
 770

<210> 5

<211> 2328

<212> DNA

<213> Artificial Sequence

<220>

<223> DNA polymerase from Thermococcus siculi

<400> 5

```

atgatcctcg acacggacta catcacggaa gatgggaaac ccgtcataag gatattcaag      60
aaagagaacg gcgagttcaa gatcgagtac gacaggactt ttgaacccta catctacgcc      120
ctcctgaagg acgactccgc gattgaggat gttaaaaaga taaccgccga gaggcacgga      180
acggtggtga aggtcaagcg cgccgaaaag gtgcagaaga agttcctagg caggccggtt      240
gaagtctgga agctctactt caccaccccc caagatgtcc cggcgataag ggacaagatt      300
aggaagcatc cagctgtaat tgacatctac gagtacgaca taccattcgc caagcgetac      360
ctcatcgaca agggcctgat tccgatggag ggtgaagaag agcttaagat gctcgccttc      420
gacattgaga cgctctacca tgagggtgag gagttcgccg aggggcctat tctgatgata      480

agctacgccg acgagagcga ggcacgcgtc atcacctgga agaaaatcga cctcccctac      540
gttgacgtcg tctcaacgga gaaggagatg ataaagcgct tcctccgctg tgtgaaggag      600
aaagatcccg atgtcctcat aacctacaac ggcgacaact tcgacttcgc ctacctgaag      660
aagcgtgtg aaaagcttgg aataaacttc ctcttgaa gggacgggag cgagccgaag      720
atccagagaa tgggtgaccg cttcgccggt gaggtgaagg ggaggataca cttcgacctc      780
tatcctgtaa taaggcgcac gataaacctg ccgacctaca tgcttgaggc agtctacgag      840
gccatctttg ggaagccaaa ggagaagggt tacgccgagg agatagccac cgcttgggaa      900
accggagagg gccttgagag ggtggctcgc tactctatgg aggacgcgaa ggtcacgttt      960
gagcttgaa aggagttctt cccgatggag gcccaacttt cgaggttggt cggccagagc     1020
ttctgggatg tcgcgcgctc aagcacgggc aatctggtcg agtggttcct cctcaggaag     1080
gcctacgaga ggaacgagct ggctccaaac aagccctctg gaagggaata tgacgagagg     1140
cgcggtgatg acgccggcgg ctacgtcaag gaaccgaaa agggcctgtg ggagaacata     1200
gtctacctcg actataaatc tctctacccc tcaatcatca tcaccacaaa cgtctcgcctc     1260
gataccctca accgcgaggg ctgtaaggag tatgacgtag ctccacaggt cggccaccgc     1320
ttctgcaagg actttccagg cttcatcccc agcctgctcg gggatctcct ggaggagagg     1380
    
```

cagaagataa agaggaagat gaaggcaaca attgaccoga tcgagagaaa gtccttgat 1440  
 tacaggcaac gggccatcaa gatccttcta aatagttttt acggctacta cggctacgca 1500  
 agggctcgct ggtactgcaa ggagtgtgcc gagagcgta cggcatgggg aagggaatat 1560  
 atcaccatga caatcagga aatagaagag aagtatggct ttaaagtact ttatgcggac 1620  
 actgacggct tcttcgcgac gattcccggg gaagatgccg agaccatcaa aaagagggcg 1680  
 atggagttcc tcaagtacat aaacgcaaaa ctccccggtg cgctcgaact tgagtacgag 1740  
 gacttctaca ggcgcggtt ctctgcacc aagaagaaat acgcggttat cgacgaggag 1800  
 ggcaagataa caacgcgcgg gctggagatc gtcaggcgcg actggagcga gatagccaag 1860  
 gagacgcagg cgcgggttct ggaggccctt ctgaaggacg gtgacgtcga agaggccgtg 1920  
 agcatagtca aagaagtgac cgagaagctg agcaagtacg aggttccgcc ggagaagctc 1980  
 gttatccacg agcagataac gcgcgagctg aaggactaca aggcaacggg accacacgtg 2040  
 gcgatagcga agaggttagc cgcgagaggc gtcaaaatcc gcccggggac agtcatcagc 2100  
 tacatcgtgc tcaagggctc cgggaggata ggcgacaggg cgattccctt cgacgagttc 2160  
 gaccccacga agcacaagta cgatgcagag tactacatcg agaaccaggt tctacctgcc 2220  
 gtcgagagga ttctgaaggc ctctcgctat cgcggtgagg agctcagata ccagaagacg 2280  
 aggcaggttg gacttggggc gtggctgaag ccgaagggga aggggtga 2328

<210> 6

<211> 775

<212> PRT

<213> Artificial Sequence

<220>

<223> DNA polymerase from *Thermococcus siculi*

<400> 6

Met Ile Leu Asp Thr Asp Tyr Ile Thr Glu Asp Gly Lys Pro Val Ile  
1 5 10 15

Arg Ile Phe Lys Lys Glu Asn Gly Glu Phe Lys Ile Glu Tyr Asp Arg  
20 25 30

Thr Phe Glu Pro Tyr Ile Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
35 40 45

Glu Asp Val Lys Lys Ile Thr Ala Glu Arg His Gly Thr Val Val Lys  
50 55 60

Val Lys Arg Ala Glu Lys Val Gln Lys Lys Phe Leu Gly Arg Pro Val  
65 70 75 80

Glu Val Trp Lys Leu Tyr Phe Thr His Pro Gln Asp Val Pro Ala Ile  
85 90 95

Arg Asp Lys Ile Arg Lys His Pro Ala Val Ile Asp Ile Tyr Glu Tyr  
100 105 110

Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Ile Pro  
 115 120 125

Met Glu Gly Glu Glu Glu Leu Lys Met Leu Ala Phe Asp Ile Glu Thr  
 130 135 140

Leu Tyr His Glu Gly Glu Glu Phe Ala Glu Gly Pro Ile Leu Met Ile  
 145 150 155 160

Ser Tyr Ala Asp Glu Ser Glu Ala Arg Val Ile Thr Trp Lys Lys Ile  
 165 170 175

Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Lys Glu Met Ile Lys  
 180 185 190

Arg Phe Leu Arg Val Val Lys Glu Lys Asp Pro Asp Val Leu Ile Thr  
 195 200 205

Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Lys Arg Cys Glu  
 210 215 220

Lys Leu Gly Ile Asn Phe Leu Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 225 230 235 240

Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 245 250 255

His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Ile Asn Leu Pro Thr  
 260 265 270

Tyr Met Leu Glu Ala Val Tyr Glu Ala Ile Phe Gly Lys Pro Lys Glu  
 275 280 285

Lys Val Tyr Ala Glu Glu Ile Ala Thr Ala Trp Glu Thr Gly Glu Gly  
 290 295 300

Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Phe  
 305 310 315 320

Glu Leu Gly Lys Glu Phe Phe Pro Met Glu Ala Gln Leu Ser Arg Leu  
 325 330 335

Val Gly Gln Ser Phe Trp Asp Val Ala Arg Ser Ser Thr Gly Asn Leu  
 340 345 350

Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 355 360 365

Pro Asn Lys Pro Ser Gly Arg Glu Tyr Asp Glu Arg Arg Gly Gly Tyr  
 370 375 380

Ala Gly Gly Tyr Val Lys Glu Pro Glu Lys Gly Leu Trp Glu Asn Ile  
 385 390 395 400

Val Tyr Leu Asp Tyr Lys Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 405 410 415

Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Lys Glu Tyr Asp  
 420 425 430

Val Ala Pro Gln Val Gly His Arg Phe Cys Lys Asp Phe Pro Gly Phe  
 435 440 445

Ile Pro Ser Leu Leu Gly Asp Leu Leu Glu Glu Arg Gln Lys Ile Lys  
 450 455 460

Arg Lys Met Lys Ala Thr Ile Asp Pro Ile Glu Arg Lys Leu Leu Asp  
 465 470 475 480

Tyr Arg Gln Arg Ala Ile Lys Ile Leu Leu Asn Ser Phe Tyr Gly Tyr  
 485 490 495

Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Lys Glu Cys Ala Glu Ser  
 500 505 510

Val Thr Ala Trp Gly Arg Glu Tyr Ile Thr Met Thr Ile Arg Glu Ile  
 515 520 525

Glu Glu Lys Tyr Gly Phe Lys Val Leu Tyr Ala Asp Thr Asp Gly Phe  
 530 535 540

Phe Ala Thr Ile Pro Gly Glu Asp Ala Glu Thr Ile Lys Lys Arg Ala  
 545 550 555 560

Met Glu Phe Leu Lys Tyr Ile Asn Ala Lys Leu Pro Gly Ala Leu Glu  
 565 570 575

Leu Glu Tyr Glu Asp Phe Tyr Arg Arg Gly Phe Phe Val Thr Lys Lys  
 580 585 590

Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 595 600 605

Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln Ala  
 610 615 620

Arg Val Leu Glu Ala Leu Leu Lys Asp Gly Asp Val Glu Glu Ala Val  
 625 630 635 640

Ser Ile Val Lys Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 645 650 655

Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Glu Leu Lys Asp  
 660 665 670

Tyr Lys Ala Thr Gly Pro His Val Ala Ile Ala Lys Arg Leu Ala Ala  
 675 680 685

Arg Gly Val Lys Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 690 695 700

Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 705 710 715 720

Asp Pro Thr Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn Gln  
 725 730 735

Val Leu Pro Ala Val Glu Arg Ile Leu Lys Ala Phe Gly Tyr Arg Gly  
 740 745 750

Glu Glu Leu Arg Tyr Gln Lys Thr Arg Gln Val Gly Leu Gly Ala Trp  
 755 760 765

Leu Lys Pro Lys Gly Lys Gly  
 770 775

<210> 7

<211> 774

<212> PRT

<213> Artificial Sequence

<220>

<223> DNA polymerase from Thermococcus kodakarensis

<400> 7

Met Ile Leu Asp Thr Asp Tyr Ile Thr Glu Asp Gly Lys Pro Val Ile  
 1 5 10 15

Arg Ile Phe Lys Lys Glu Asn Gly Glu Phe Lys Ile Glu Tyr Asp Arg  
 20 25 30

Thr Phe Glu Pro Tyr Phe Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
 35 40 45

Glu Glu Val Lys Lys Ile Thr Ala Glu Arg His Gly Thr Val Val Thr  
 50 55 60

Val Lys Arg Val Glu Lys Val Gln Lys Lys Phe Leu Gly Arg Pro Val  
 65 70 75 80

Glu Val Trp Lys Leu Tyr Phe Thr His Pro Gln Asp Val Pro Ala Ile  
 85 90 95

Arg Asp Lys Ile Arg Glu His Pro Ala Val Ile Asp Ile Tyr Glu Tyr  
 100 105 110

Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Val Pro  
 115 120 125

Met Glu Gly Asp Glu Glu Leu Lys Met Leu Ala Phe Asp Ile Glu Thr  
 130 135 140

Leu Tyr Glu Glu Gly Glu Glu Phe Ala Glu Gly Pro Ile Leu Met Ile  
 145 150 155 160

Ser Tyr Ala Asp Glu Glu Gly Ala Arg Val Ile Thr Trp Lys Asn Val  
 165 170 175

Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Arg Glu Met Ile Lys





<210> 8

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 8

tagaattgaa gaa 13

<210> 9

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 9

tggccatagc tac 13

<210> 10

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 10

gtcatctgcg acc 13

<210> 11

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 11

ttcgcgcttg gac 13

<210> 12

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 12

cgcgaaccgt tag 13

<210> 13

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 13

ttgcagcctc taa 13

<210> 14

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 14

tctactagta cga 13

<210> 15

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 15

gtaggttcta ctg 13

<210> 16

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 16  
gccaatatca agt 13

<210> 17  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 17  
ctatcttgct ggt 13

<210> 18  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 18  
gttctcatag gta 13

<210> 19  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 19  
gtctatgaac caa 13

<210> 20  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 20  
cggagcgctt att 13

<210> 21

<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 21  
tatgcatga gga 13

<210> 22  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 22  
atacgactcg gag 13

<210> 23  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 23  
gatggaactc agc 13

<210> 24  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 24  
ggacctgcat gaa 13

<210> 25  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 25

tagactggaa ctt 13

<210> 26

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 26

gaattacctc gtt 13

<210> 27

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 27

aggatcaggc tac 13

<210> 28

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 28

acgcgtagaa gag 13

<210> 29

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 29

cttcgagact tac 13

<210> 30

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 30

gacggctaac tcc 13

<210> 31

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 31

ttagcattct ctt 13

<210> 32

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 32

gcaaggcata gta 13

<210> 33

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 33

acctagatat gga 13

<210> 34

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 34

acgccaaggc gta 13

<210> 35

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 35

tatgacggat ccg 13

<210> 36

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 36

cctccattag aga 13

<210> 37

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 37

attgaatact ctg 13

<210> 38

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 38

gagatgagaa gaa 13

<210> 39

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 39

tctgagtagc cgg 13

<210> 40

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 40

aataggtagt acg 13

<210> 41

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 41

gtcgaagaag tcc 13

<210> 42

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 42

tactgcatct cgt 13

<210> 43  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 43  
gacgtattag agc 13

<210> 44  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 44  
cctgcattat tcg 13

<210> 45  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 45  
acgaatgatg ctc 13

<210> 46  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 46  
tactagcaga gat 13

<210> 47  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 47

ctcctcatct tcc 13

<210> 48

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 48

tcctctgcbc tgc 13

<210> 49

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 49

ccttctcagt ccg 13

<210> 50

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 50

cagcttcata gcg 13

<210> 51

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 51

ttgactctcg cgc 13

<210> 52

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 52

tatcctgagc gat 13

<210> 53

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 53

aacgcctagc cga 13

<210> 54

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 54

ccgaagacgt cat 13

<210> 55

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 55

gagttctcca gat 13

<210> 56

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 56

tgcatccgctt 13

<210> 57

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 57

cctgaactca agt 13

<210> 58

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> sample tag sequence

<400> 58

ggtcgtatgc gta 13

<210> 59

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 59

aggcctctct acc 13

<210> 60

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 60  
gtactccatc caa 13

<210> 61

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 61  
cagcggacgc gct 13

<210> 62

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 62  
atctctctta gca 13

<210> 63

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 63  
aagcaataat aat 13

<210> 64

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 64  
aaggcgactc cga 13

<210> 65  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 65  
acgtctctag gag 13

<210> 66  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 66  
ccatcagacc tct 13

<210> 67  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 67  
acttaatcgt act 13

<210> 68  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 68  
tggaattctc caa 13

<210> 69  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 69

ccatacgatc agg 13

<210> 70

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 70

ttatggagca ata 13

<210> 71

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 71

gctcggcggt cga 13

<210> 72

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 72

ttggccagtc gct 13

<210> 73

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 73

cagatacgta gag 13

<210> 74

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 74

aatgctatta tcc 13

<210> 75

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 75

gcagcatgcc gat 13

<210> 76

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 76

ggagagttac ctc 13

<210> 77

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 77

gagagtccat gat 13

<210> 78

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 78

caatctattc tga 13

<210> 79

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 79

gctcttagta tcc 13

<210> 80

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 80

ccatagttat ggt 13

<210> 81

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 81

tgcgagatcg aag 13

<210> 82

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 82  
agagaagtcg agt 13

<210> 83  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 83  
ggtaactcca tat 13

<210> 84  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 84  
tgctattcca ggc 13

<210> 85  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 85  
aaccgaggg ctc 13

<210> 86  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 86  
ttctagagat acc 13

<210> 87  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 87  
ttcgctcaag tat 13

<210> 88  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 88  
cagagaaggc gca 13

<210> 89  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 89  
tagaattggc ctc 13

<210> 90  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 90  
ggccattctc cag 13

<210> 91  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 91

tccaacgcgc gtt 13

<210> 92

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 92

gccgcagatt acg 13

<210> 93

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 93

gcagttcgaa cgc 13

<210> 94

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 94

ttctctctgc agg 13

<210> 95

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 95

taagctacca gcg 13

<210> 96

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 96

ctgcatgagg ttg 13

<210> 97

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 97

ttgcctagcg agg 13

<210> 98

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 98

caactgaatt agg 13

<210> 99

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 99

aagcggctct ctt 13

<210> 100

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 100

aatggaagga ccg 13

<210> 101

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 101

gagttagtaa gtt 13

<210> 102

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 102

ttcctaattc caa 13

<210> 103

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 103

gttctggttc gct 13

<210> 104

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 104  
gttcatctct tcc 13

<210> 105

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 105  
attccgagga aga 13

<210> 106

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 106  
cttagccgag aga 13

<210> 107

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 107  
gtctgctacg ctt 13

<210> 108

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 108  
atggcgccgc gca 13

<210> 109

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 109

taattggta tct 13

<210> 110

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 110

tcggtataa gtc 13

<210> 111

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 111

tgctgagaa cgt 13

<210> 112

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 112

agatgcggtt aac 13

<210> 113

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 113

atggaatagg cga 13

<210> 114

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 114

agagatgcga tcg 13

<210> 115

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 115

ctccaactaa cgt 13

<210> 116

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 116

gccttgctac tgg 13

<210> 117

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 117

cttcgtctct acg 13

<210> 118

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 118

acgctcatag cct 13

<210> 119

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 119

gtcgaagata agg 13

<210> 120

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 120

gccggagtcc tcg 13

<210> 121

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 121

tatacggcga cct 13

<210> 122

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 122

aggtagatat tcg 13

<210> 123

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 123

ttaaggctact gct 13

<210> 124

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 124

cggatctggt ata 13

<210> 125

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 125

gaggtctcgg agg 13

<210> 126

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 126

ggcatcgatg gac 13

<210> 127

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 127

gatctccgat ata 13

<210> 128

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 128

gattcggaat act 13

<210> 129

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 129

ctgcatccg gcc 13

<210> 130

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 130

gatccggtg caa 13

<210> 131  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 131  
cgtcaggctt gac 13

<210> 132  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 132  
tcggcaaggc gag 13

<210> 133  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 133  
gaacggcgaa cgc 13

<210> 134  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Sample tag sequence

<400> 134  
cctcaagcgg act 13

<210> 135  
<211> 13  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 135

gaagccagat ggt 13

<210> 136

<211> 13

<212> DNA

<213> Artificial Sequence

<220>

<223> Sample tag sequence

<400> 136

tgctcatacc aat 13

## REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

### Patent documents cited in the description

- [GB1802744A \[0206\]](#)

### Non-patent literature cited in the description

- **KONG THOO LIN PBROWN D M** Synthesis and duplex stability of oligonucleotides containing cytosine-thymine analogues *Nucleic Acids Research*, 1989, vol. 17, 10373-

10383 [0007]

- **STONE M J et al.** Molecular basis for methoxyamine-initiated mutagenesis: H nuclear magnetic resonance studies of base-modified oligodeoxynucleotides *Journal of Molecular Biology*, 1991, vol. 222, 711-723 [0007]
- **NEDDERMAN A N R et al.** Molecular basis for methoxyamine initiated mutagenesis: H nuclear magnetic resonance studies of oligonucleotide duplexes containing base-modified cytosine residues *Journal of Molecular Biology*, 1993, vol. 230, 1068-1076 [0007]
- **MOORE M H et al.** Direct observation of two base-pairing modes of a cytosine-thymine analogue with guanine in a DNA-form duplex. Significance for base analogue mutagenesis *Journal of Molecular Biology*, 1995, vol. 251, 665-673 [0007]
- **HARRIS V H et al.** The effect of tautomeric constant on the specificity of nucleotide incorporation during DNA replication: support for the rare tautomer hypothesis of substitution mutagenesis *Journal of Molecular Biology*, 2003, vol. 326, 1389-1401 [0007] [0007]
- **ZACCOLO M et al.** An approach to random mutagenesis of DNA using mixtures of triphosphate derivatives of nucleoside analogues *Journal of Molecular Biology*, 1996, vol. 255, 589-603 [0007]
- **SHEN et al.** DNA REPAIR PROTOCOLS. METHODS IN MOLECULAR BIOLOGY HUMANA PRESS 2002 0101 vol. 192, 167-174 [0012]
- **CHOI et al.** NUCLEIC ACIDS RESEARCH, 2016, vol. 44, 30305-10481022-1035 [0012]
- **MASAYASU et al.** MOLECULES, 2010, vol. 15, 118229-8240 [0012]
- **PETRIE et al.** NUCLEIC ACIDS RESEARCH, 2010, vol. 38, 228095-8104 [0012]
- **ELSHAWADFY et al.** FRONTIERS IN MICROBIOLOGY, 2014, vol. 5, [0012]
- **YAMASHITA et al.** MOLECULAR BIOTECHNOLOGY HUMANA PRESS 2017 0508 vol. 59, 221-233 [0012]
- **DUBENDORFF JW STUDIER FW** Controlling basal expression in an inducible T7 expression system by blocking the target T7 promoter with lac repressor" *Journal of Molecular Biology*, 1991, vol. 219, 145-59 [0146]
- **SAMBROOK, J. et al.** *Molecular Cloning: A Laboratory Manual* Cold Spring Harbor Laboratory Press 1989 0000 [0146]

## FREM GANGSMÅDE TIL IND FØRING AF MUTATIONER

## PATENTKRAV

1. Fremgangsmåde til indføring af substitutionsmutationer i mindst ét mål-DNA-molekyle, hvilken fremgangsmåde omfatter:
  - 5 a. tilvejebringelse af mindst én prøve, der omfatter mindst ét mål-DNA-molekyle; og
  - b. amplifikation af det mindst ene mål-DNA-molekyle ved hjælp af en DNA-polymerase med lav bias og høj nøjagtighed med lav template-amplifikations-bias;

10 hvor trinnet med amplifikation af det mindst ene mål-DNA-molekyle udføres i tilstedeværelse af en nukleotidanalogue og omfatter mindst 2 runder med replikation af det mindst ene mål-DNA-molekyle, hvor i en første replikationsrunde DNA-polymerasen inkorporerer nukleotidanalogen i stedet for et nukleotid og i en anden replikationsrunde nukleotidanalogen parres med et naturligt nukleotid for at indføre en substitutionsmutation i den komplementære streng.
2. Anvendelse af en DNA-polymerase med lav bias og høj nøjagtighed med lav template-amplifikations-bias i en fremgangsmåde til indføring af mutationer i mindst ét mål-DNA-molekyle, hvor  
15 fremgangsmåden omfatter:
  - a. tilvejebringelse af mindst én prøve, der omfatter mindst ét mål-DNA-molekyle; og
  - b. amplifikation af det mindst ene mål-DNA-molekyle ved hjælp af DNA-polymerasen;

20 hvor trinnet med amplifikation af det mindst ene mål-DNA-molekyle udføres i tilstedeværelse af en nukleotidanalogue og omfatter mindst 2 runder med replikation af det mindst ene mål-DNA-molekyle, hvor i en første replikationsrunde DNA-polymerasen inkorporerer nukleotidanalogen i stedet for et nukleotid og i en anden replikationsrunde nukleotidanalogen parres med et naturligt nukleotid for at indføre en substitutionsmutation i den komplementære streng.
3. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor DNA-polymerasen muterer adenin-, thymine-, guanin- og cytosinnukleotider i det mindst ene mål-DNA-molekyle  
25 ved et rateforhold på henholdsvis 0,5-1,5:0,5-1,5:0,5-1,5:0,5-1,5, 0,6-1,4:0,6-1,4:0,6-1,4:0,6-1,4, 0,7-1,3:0,7-1,3:0,7-1,3:0,7-1,3, 0,8-1,2:0,8-1,2:0,8-1,2:0,8-1,2, eller omtrent 1:1:1:1, eventuelt hvor DNA-polymerasen muterer adenin-, thymine-, guanin- og cytosinnukleotider i det mindst ene mål-DNA-molekyle ved et rateforhold på henholdsvis 0,7-1,3:0,7-1,3:0,7-1,3:0,7-1,3.
4. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor DNA-polymerasen muterer mellem 1 % og 15 %, mellem 2% og 10 %, eller omtrent 8 % af nukleotiderne i det  
30 mindst ene mål-DNA-molekyle, og/eller hvor DNA-polymerasen muterer mellem 0 % og 3%, eller mellem 0 % og 2% af nukleotiderne i det mindst ene mål-DNA-molekyle pr. replikationsrunde.
5. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor DNA-polymerasen muterer adenin, thymine, guanin og/eller cytosin i det mindst ene mål-DNA-molekyle ved hjælp  
35 af en nukleotidanalogue, og/eller hvor DNA-polymerasen erstatter guanin, cytosin, adenin og/eller thymine med en nukleotidanalogue, og/eller hvor DNA-polymerasen indfører guanin- eller adeninnukleotider ved hjælp af en nukleotidanalogue ved et rateforhold på henholdsvis 0,5-1,5:0,5-1,5, 0,6-1,4:0,6-1,4, 0,7-1,3:0,7-1,3, 0,8-1,2:0,8-1,2, eller omtrent 1:1, og/eller hvor DNA-polymerasen indfører guanin- eller adeninnukleotider ved hjælp af en nukleotidanalogue ved et rateforhold på henholdsvis 0,7-1,3:0,7-1,3.

6. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af kravene 1-5, hvor nukleotidanalogen er dPTP, eventuelt hvor DNA-polymerasen indfører guanin-til-adeninsubstitutionsmutationer, cytosin-til-thyminsubstitutionsmutationer, adenin-til-guaninsubstitutionsmutationer og thymin-til-cytosinsubstitutionsmutationer, eventuelt hvor DNA-polymerasen indfører guanin-til-adeninsubstitutionsmutationer, cytosin-til-thyminsubstitutionsmutationer, adenin-til-guaninsubstitutionsmutationer og thymin-til-cytosinsubstitutionsmutationer ved et rateforhold på 0,5-1,5:0,5-1,5:0,5-1,5:0,5-1,5, 0,6-1,4:0,6-1,4:0,6-1,4:0,6-1,4, 0,7-1,3:0,7-1,3:0,7-1,3:0,7-1,3, 0,8-1,2:0,8-1,2:0,8-1,2:0,8-1,2, eller omtrent 1:1:1:1, eventuelt hvor DNA-polymerasen indfører guanin-til-adeninsubstitutionsmutationer, cytosin-til-thyminsubstitutionsmutationer, adenin-til-guaninsubstitutionsmutationer og thymin-til-cytosinsubstitutionsmutationer ved et rateforhold på henholdsvis 0,7-1,3:0,7-1,3:0,7-1,3:0,7-1,3.
7. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af kravene 1-6, hvor, i fravær af nukleotidanaloger, DNA-polymerasen med høj præcision indfører mindre end 0,01 %, mindre end 0,0015 %, mindre end 0,001 %, mellem 0 % og 0,0015 %, eller mellem 0 % og 0,001 % mutationer pr. replikationsrunde.
8. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af kravene 1 til 7, hvor fremgangsmåden omfatter et yderligere trin med amplifikation af det mindst ene mål-DNA-molekyle, der omfatter nukleotidanaloger, i fravær af nukleotidanaloger, eventuelt hvor trinnet med amplifikation af det mindst ene mål-DNA-molekyle, der omfatter nukleotidanaloger, i fravær af nukleotidanaloger udføres ved hjælp af DNA-polymerasen.
9. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor fremgangsmåden tilvejebringer et muteret mindst ét mål-DNA-molekyle og fremgangsmåden endvidere omfatter et yderligere trin med amplifikation af det muterede mindst ene mål-DNA-molekyle ved hjælp af DNA-polymerasen.
10. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor DNA-polymerasen omfatter et korrekturlæsningsdomæne og/eller et progressivitetsforbedrende domæne, og/eller hvor DNA-polymerasen omfatter et fragment af mindst 400, mindst 500, mindst 600, mindst 700 eller mindst 750 tilstødende aminosyrer fra:
- a. en sekvens ifølge SEQ ID NO: 2;
  - b. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 2;
  - c. en sekvens ifølge SEQ ID NO: 4;
  - d. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 4;
  - e. en sekvens ifølge SEQ ID NO: 6;
  - f. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 6;
  - g. en sekvens ifølge SEQ ID NO: 7; eller

- 3 -

- h. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 7;
- eventuelt hvor DNA-polymerasen omfatter:
- a. en sekvens ifølge SEQ ID NO: 2;
- 5 b. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 2;
- c. en sekvens ifølge SEQ ID NO: 4;
- d. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 4;
- 10 e. en sekvens ifølge SEQ ID NO: 6;
- f. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 6;
- g. en sekvens ifølge SEQ ID NO: 7; eller
- h. en sekvens, der er mindst 95 %, mindst 98 %, eller mindst 99 % identisk med SEQ ID NO: 7;
- 15 7;
- eventuelt hvor:
- i) DNA-polymerasen omfatter en sekvens, der er mindst 98 % identisk med SEQ ID NO: 2;
- ii) DNA-polymerasen omfatter en sekvens, der er mindst 98 % identisk med SEQ ID NO: 4;
- iii) DNA-polymerasen omfatter en sekvens, der er mindst 98 % identisk med SEQ ID NO: 6;
- 20 eller
- iv) DNA-polymerasen omfatter en sekvens, der er mindst 98 % identisk med SEQ ID NO: 7; og/eller
- hvor DNA-polymerasen med lav bias er en termokokpolymerase, eller et derivat deraf, eventuelt hvor termokokpolymerasen er afledt af en termokokstamme, der er valgt fra gruppen bestående af *T. kodakarensis*, *T. siculi*, *T. celer* og *T. sp KS-1*.
- 25 *kodakarensis*, *T. siculi*, *T. celer* og *T. sp KS-1*.
11. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, der endvidere omfatter:
- A) indføring af stregkoder og/eller prøvetags i det mindst ene mål-DNA-molekyle, eventuelt hvor:
- 30 a) en gruppe afprøvetags anvendes og mål-DNA-molekyler fra forskellige prøver mærkes med forskellige prøvetags fra gruppen, eventuelt hvor hvert prøvetag adskiller sig fra i alt væsentligt samtlige andre prøvetags i gruppen med mindst 1 lavt sandsynlig mutationsforskel eller mindst 3 højt sandsynlige mutationsforskelle, eventuelt hvor hvert prøvetag adskiller sig fra i alt væsentligt samtlige andre prøvetags i gruppen med mindst 3 lavt sandsynlige mutationsforskelle og/eller hvert prøvetag adskiller sig fra i alt
- 35 væsentligt samtlige andre prøvetags i gruppen med mellem 3 og 25, eller mellem 3 og 10 lavt sandsynlige mutationsforskelle og/eller den lave sandsynlighedsmutation er en transversionsmutation eller en indel-mutation og/eller hvor den højt sandsynlige mutation er en overgangsmutation;
- eller

- b) hvert prøvetag adskiller sig fra i alt væsentligt samtlige andre prøvetags i gruppen med mindst 5 højt sandsynlige mutationsforskelle, eventuelt hvor hvert prøvetag adskiller sig fra i alt væsentligt samtlige andre prøvetags i gruppen med mellem 5 og 25, eller mellem 5 og 10 højt sandsynlige mutationsforskelle, eventuelt hvor den højt sandsynlige mutation er en overgangsmutation;
- 5 og/eller
- B) indføring af adaptere i hvert af det mindst ene mål-DNA-molekyle, eventuelt hvor indføring af adaptere i hvert af det mindst ene mål-DNA-molekyle omfatter indføring af en første adapter ved 3'-enden af det mindst ene mål-DNA-molekyle og en anden adapter ved 5'-enden af det mindst ene mål-DNA-molekyle, hvor den første adapter og den anden adapter kan anneale til hinanden, eventuelt hvor det mindst
- 10 ene mål-DNA-molekyle amplificeres ved anvendelse af primere, der er identiske med hinanden og komplementære til en del af den første adapter, og/eller hvor den første adapter er komplementær til et DNA-molekyle, der er mindst 80 %, mindst 90 %, mindst 95 %, mindst 99 % eller 100 % identisk med den anden adapter, og/eller hvor primerne omfatter et andet primerbindingssted, og fremgangsmåden omfatter
- 15 amplifikation af det mindst ene mål-DNA-molekyle ved hjælp af primerne, fjernelse af primerne og yderligere amplifikation af det mindst ene mål-DNA-molekyle ved hjælp af et andet sæt af primere, der annealer til det andet primerbindingssted;
- og/eller
- C) indføring af stregkoder, prøvetags og adaptere i hvert af mål-DNA-molekyle, og/eller hvor stregkoderne, prøvetags og/eller adaptere indføres ved tagmentering eller ved klipning eller ligering.
- 20 12. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor det mindst ene mål-DNA-molekyle er større end 1 kbp, større end 1,5 kbp, større end 2 kbp, større end 4 kbp, større end 5 kbp, større end 7 kbp eller større end 8 kbp.
13. Fremgangsmåde til bestemmelse af en sekvens af mindst ét mål-DNA-molekyle, hvilken fremgangsmåde omfatter følgende trin:
- 25 a. udførelse af fremgangsmåden ifølge et hvilket som helst af kravene 1 eller 3-12 til at tilvejebringe mindst ét muteret mål-DNA-molekyle;
- b. eventuelt fragmentering og/eller amplifikation af det mindst ene muterede mål-DNA-molekyle for at tilvejebringe mindst ét fragmenteret og/eller amplificeret muteret mål-DNA-molekyle;
- c. sekvensering af områder af det mindst ene muterede mål-DNA-molekyle eller det mindst
- 30 ene fragmenterede og/eller amplificerede muterede mål-DNA-molekyle for at tilvejebringe muterede sekvenslæsninger; og
- d. sammensætning af en sekvens for mindst en del af det mindst ene mål-DNA-molekyle ved hjælp af de muterede sekvenslæsninger.
14. Fremgangsmåde til protein-engineering, hvilken fremgangsmåde omfatter følgende trin:
- 35 a. udførelse af fremgangsmåden ifølge et hvilket som helst af kravene 1 eller 3-12 for at tilvejebringe mindst ét muteret mål-DNA-molekyle;
- b. indsætning af det mindst ene muterede mål-DNA-molekyle i en vektor og
- c. ekspresion af et protein, der er kodet for af det mindst ene muterede mål-DNA-molekyle; eventuelt hvor:

- 5 -

- i) fremgangsmåden omfatter følgende trin:
- a. amplifikation af det mindst ene mål-DNA-molekyle, der omfatter nukleotidanalogen, i fravær af nukleotidanalogen for at tilvejebringe mindst ét muteret mål-DNA-molekyle;
  - b. indsætning af det mindst ene muterede mål-DNA-molekyle i en vektor; og
  - 5 c. ekspresion af et protein, der er kodet for af det mindst ene muterede mål-DNA-molekyle; og/eller
- ii) fremgangsmåden endvidere omfatter et trin med teste af aktiviteten eller vurdering af proteinets struktur, der er kodet for af det mindst ene muterede mål-DNA-molekyle; og/eller
- iii) vektoren er et plasmid, et virus, et cosmid eller et kunstigt kromosom; og/eller
- 10 iv) trinnet med ekspresion af et protein, der er kodet for af det mindst ene muterede mål-DNA-molekyle, opnås ved transformation af bakterieceller, transfektion af eukaryotiske celler eller transduktion af eukaryotiske celler med vektoren.
- 15 15. Fremgangsmåde eller anvendelse ifølge et hvilket som helst af de foregående krav, hvor trinnet med amplifikation af det mindst ene mål-DNA-molekyle ved anvendelse af en DNA-polymerase med lav bias og høj nøjagtighed udføres ved hjælp af dNTP'er ved uens koncentrationer; og/eller
- 15 I) hvor:
- A)
- i) fremgangsmåden omfatter et yderligere trin med amplifikation af det mindst ene mål-DNA-molekyle, der omfatter nukleotidanaloger, i fravær af nukleotidanaloger og det yderligere trin med amplifikation af det mindst ene mål-DNA-molekyle, der omfatter nukleotidanaloger, i fravær af nukleotidanaloger udføres ved hjælp af dNTP'er ved uens koncentrationer; eller
  - 20 ii) fremgangsmåden tilvejebringer et muteret mindst ét mål-DNA-molekyle, fremgangsmåden omfatter et yderligere trin med amplifikation af det muterede mindst ene mål-DNA-molekyle ved hjælp af DNA-polymerasen eventuelt i fravær af nukleotidanaloger og det yderligere trin med amplifikation af det muterede mindst ene mål-DNA-molekyle ved hjælp af DNA-polymerasen udføres ved hjælp af dNTP'er ved uens koncentrationer; og/eller
- 25 B) dNTP'er ved uens koncentrationer anvendes til at ændre profilen af mutationer, der indføres, eventuelt hvor dNTP'er ved uens koncentrationer anvendes til at reducere bias i den mutationsprofil, der indføres; og/eller
- 30 C) dNTP'erne ved uens koncentrationer omfatter dATP, dCTP, dTTP og dGTP og ét eller to af dATP, dCTP, dTTP eller dGTP er ved en lavere koncentration sammenlignet med andre dNTP'er; og/eller
- D) anvendelse af dNTP'er ved uens koncentrationer omfatter et trin med identifikation af et dNTP, hvis niveau skal øges eller reduceres for at reducere bias i den mutationsprofil, der indføres; og/eller
- E) dNTP'erne ved uens koncentrationer omfatter dTTP ved en lavere koncentration end andre
- 35 dNTP'er, eventuelt hvor dNTP'erne ved uens koncentrationer omfatter dTTP ved en koncentration, der er lavere end 75 %, lavere end 70 %, lavere end 60 %, lavere end 55 %, mellem 25 % og 75 %, mellem 25 % og 70, mellem 25 % og 60 %, eller omtrent 50 % af koncentrationen af dATP, dCTP eller dGTP, eventuelt hvor dNTP'erne ved uens koncentrationer omfatter dTTP ved en koncentration, der er lavere end 75 %, lavere end 70 %, lavere end 60 %, lavere end 55 %, mellem 25 % og 75 %, mellem 25 % og 70 %, mellem 25 % og

60 %, eller omtrent 50 % af koncentrationen af dCTP, eventuelt hvor dNTP'erne ved uens koncentrationer omfatter dTTP ved en koncentration, der er lavere end 60 % af koncentrationen af dCTP, eventuelt dNTP'erne ved uens koncentrationer omfatter dTTP ved en koncentration mellem 25 % og 60 % af koncentrationen af dCTP;

5 og/eller

F) dNTP'erne ved uens koncentrationer omfatter dATP ved en lavere koncentration sammenlignet med andre dNTP'er, eventuelt hvor dNTP'erne ved uens koncentrationer omfatter dATP ved en koncentration, der er lavere end 75 %, lavere end 70 %, lavere end 60 %, lavere end 55 %, mellem 25 % og 75 %, mellem 25 % og 70 %, mellem 25 % og 60 % eller omtrent 50 % af koncentrationen af dTTP, dCTP eller dGTP, eventuelt hvor dNTP'erne ved uens koncentrationer af dNTP'er omfatter dATP ved en koncentration, der er lavere end 75 %, lavere end 70 %, lavere end 60 %, lavere end 55 %, mellem 25 % og 75 %, mellem 25 % og 70 %, mellem 25 % og 60 % eller omtrent 50 % af koncentrationen af dGTP, eventuelt hvor dNTP'erne ved uens koncentrationer omfatter dATP ved en koncentration, der er lavere end 60 % af koncentrationen af dGTP, eventuelt hvor dNTP'erne ved uens koncentrationer omfatter dATP ved en koncentration mellem 25 % og 60 % af koncentrationen af dGTP;

15


og/eller

II) fremgangsmåden er en fremgangsmåde til indføring af mutationer i en mutationsprofil med lav bias.

# DRAWINGS

Figure 1

Panel A:



Panel B:




Figure 1 (Cont.)

Panel C:




Figure 2




FIGURE 3

SEQ ID NO: 1 – nucleotide sequence of DNA polymerase from *Thermococcus* sp. KS-1

```

atgatcctcg acactgacta cataactgag aatggaaaac ccgtcataag gattttcaag
aaggagaacg gcgagttaa gattgagtac gataggactt ttgaacccta catttacgcc
ctcctgaagg acgattctgc cattgaggag gtcaagaaga taaccgccga gaggaacgga
acggttgtaa cggttaagcg ggctgaaaag gttcagaaga agttcctcgg gagaccagtt
gagggtctgga aactctactt tactcaccct caggacgtcc cagcgataag ggacaagata
cgagagcatc cagcagttat tgacatctac gagtacgaca tacccttcgc caagcgctac
ctcatagaca agggattagt gccaatggaa ggcgacgagg agctgaaaat gcttgccctt
gatatcgaga cgctctacca tgagggcgag gagttcgccg aggggccaat ccttatgata
agctacgccg acgaggaagg ggccagggtg ataacgtgga agaaccggga tctgccctac
gttgacgtcg tctcgacgga gagggagatg ataaagcgt tcttaagggt ggtcaaagag
aaagatcctg acgtccta atacctacaac ggcgacaact tcgacttcgc ctacctaaaa
aaacgctgtg aaaagcttgg aataaacttc acgctcggaa gggacggaag cgagccgaag
attcagagga tgggcgacag gtttgccctc gaagtgaagg gacggataca cttcgatctc
tatcctgtga taagacggac gataaacctg cccacataca cgcttgaggc cgtttatgaa
gccgtcttcg gtcagccgaa ggagaaggtc tacgctgagg agatagctac agcttgggag
agcggtgaa gcttgagag agtagccaga tactcgatgg aagatgcgaa ggtcacatac
gagcttggga aggagtttt cctatggag gccagcttt ctcgctta at cggccagtcc
ctctgggacg tctccgctc cagcactggc aacctcgttg agtggttcct cctcaggaag
gcctacgaga ggaatgagct ggccccgaac aagccgatg aaaaggagct ggccagaaga
cgacagagct atgaaggagg ctatgtaaaa gagcccgaga gagggttggt ggagaaacata
gtgtacctag attttagatc tctgtacccc tcaatcatca tcaccacaaa cgtctcgccg
gatactctca acaggaagg atgcaaggaa tatgacggtt cccccagggt cggtaaccgc
ttctgcaagg acttccagg atttatccc agcctgcttg gagacctcct agaggagagg
cagaagataa agaagaagat gaagggccacg attgacccga tcgagaggaa gctcctcgat
tacaggcaga gggccatcaa gatcctggcc aacagctact acggttacta cggctatgca
agggcgcgct ggtactgcaa ggagtgtgca gagagcgtaa cggcctgggg aagggagtac
ataacgatga ccatcagaga gatagaggaa aagtacggct ttaaggta at ctacagcgac
accgacggat tttttgccac aatacctgga gccgatgctg aaaccgtcaa aaagaaggcg
atggagtctc tcaagtatat caacgccaaa ctcccggggc cgcttgagct cgagtaacgag
ggcttctaca aacgcccgtt ctctgctcac aagaagaagt acgcggtgat agacgaggaa
ggcaagataa caacgcgccc acttgagatt gtgaggcgcg actggagcga gatagcga
gagacgcagg cgagggttct tgaagctttg ctaaaggacg gtgacgtcga gaaggccgtg
aggatagtca aagaagttac cgaaaagctg agcaagtac aggttccgcc ggagaagctg
gtgatccacg agcagataac gagggattta aaggactaca aggcaaccgg tccccacggt
gccgttgcca agagggttggc cgcgagagga gtcaaaatac gccttggaac ggtgataagc
tacatcgtgc tcaagggctc tgggaggata ggcgacaggg cgataaccgtt cgacgagttc
gaccocgacg agcacaagta cgacgcgag tactacattg agaaccagggt tctcccagcc
gttgagagaa ttctgagagc cttcggttac cgcaaggaag acctgcgcta ccagaagacg
agacaggttg gtctgggagc ctggctgaag ccgaaggga cttga

```

SEQ ID NO: 2 – polypeptide sequence of DNA polymerase from *Thermococcus* sp. KS-1

Met Ile Leu Asp Thr Asp Tyr Ile Thr Glu Asn Gly Lys Pro Val Ile  
 Arg Ile Phe Lys Lys Glu Asn Gly Glu Phe Lys Ile Glu Tyr Asp Arg  
 Thr Phe Glu Pro Tyr Ile Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
 Glu Glu Val Lys Lys Ile Thr Ala Glu Arg His Gly Thr Val Val Thr  
 Val Lys Arg Ala Glu Lys Val Gln Lys Lys Phe Leu Gly Arg Pro Val  
 Glu Val Trp Lys Leu Tyr Phe Thr His Pro Gln Asp Val Pro Ala Ile  
 Arg Asp Lys Ile Arg Glu His Pro Ala Val Ile Asp Ile Tyr Glu Tyr  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Val Pro  
 Met Glu Gly Asp Glu Glu Leu Lys Met Leu Ala Phe Asp Ile Glu Thr  
 Leu Tyr His Glu Gly Glu Glu Phe Ala Glu Gly Pro Ile Leu Met Ile  
 Ser Tyr Ala Asp Glu Glu Gly Ala Arg Val Ile Thr Trp Lys Asn Ala  
 Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Arg Glu Met Ile Lys  
 Arg Phe Leu Lys Val Val Lys Glu Lys Asp Pro Asp Val Leu Ile Thr  
 Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Lys Arg Cys Glu  
 Lys Leu Gly Ile Asn Phe Thr Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Ile Asn Leu Pro Thr  
 Tyr Thr Leu Glu Ala Val Tyr Glu Ala Val Phe Gly Gln Pro Lys Glu  
 Lys Val Tyr Ala Glu Glu Ile Ala Thr Ala Trp Glu Ser Gly Glu Gly  
 Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Tyr  
 Glu Leu Gly Lys Glu Phe Phe Pro Met Glu Ala Gln Leu Ser Arg Leu  
 Ile Gly Gln Ser Leu Trp Asp Val Ser Arg Ser Ser Thr Gly Asn Leu  
 Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 Pro Asn Lys Pro Asp Glu Lys Glu Leu Ala Arg Arg Arg Gln Ser Tyr  
 Glu Gly Gly Tyr Val Lys Glu Pro Glu Arg Gly Leu Trp Glu Asn Ile  
 Val Tyr Leu Asp Phe Arg Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Lys Glu Tyr Asp  
 Val Ala Pro Gln Val Gly His Arg Phe Cys Lys Asp Phe Pro Gly Phe  
 Ile Pro Ser Leu Leu Gly Asp Leu Leu Glu Glu Arg Gln Lys Ile Lys  
 Lys Lys Met Lys Ala Thr Ile Asp Pro Ile Glu Arg Lys Leu Leu Asp  
 Tyr Arg Gln Arg Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr  
 Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Lys Glu Cys Ala Glu Ser  
 Val Thr Ala Trp Gly Arg Glu Tyr Ile Thr Met Thr Ile Arg Glu Ile  
 Glu Glu Lys Tyr Gly Phe Lys Val Ile Tyr Ser Asp Thr Asp Gly Phe  
 Phe Ala Thr Ile Pro Gly Ala Asp Ala Glu Thr Val Lys Lys Lys Ala  
 Met Glu Phe Leu Lys Tyr Ile Asn Ala Lys Leu Pro Gly Ala Leu Glu  
 Leu Glu Tyr Glu Gly Phe Tyr Lys Arg Gly Phe Phe Val Thr Lys Lys  
 Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln Ala  
 Arg Val Leu Glu Ala Leu Leu Lys Asp Gly Asp Val Glu Lys Ala Val  
 Arg Ile Val Lys Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Asp Leu Lys Asp  
 Tyr Lys Ala Thr Gly Pro His Val Ala Val Ala Lys Arg Leu Ala Ala  
 Arg Gly Val Lys Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 Asp Pro Thr Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn Gln  
 Val Leu Pro Ala Val Glu Arg Ile Leu Arg Ala Phe Gly Tyr Arg Lys  
 Glu Asp Leu Arg Tyr Gln Lys Thr Arg Gln Val Gly Leu Gly Ala Trp  
 Leu Lys Pro Lys Gly Thr

SEQ ID NO: 3 – nucleic acid sequence of DNA polymerase from *Thermococcus celer*

atgatacctcg acgctgacta catcacccgaa gatgggaagc ccgtcgtgag gatattcagg  
 aaggagaagg gccagttcag aatcgactac gacagggact tccagcccta catctacgcc  
 ctccctgaagg acgattcggc catcgaggag gtgaagagga taaccgttga gccccaaggg  
 aaggccgtca gggttaagcg ggtggagaag gtcgaaaaga agtcctcaa caggccgata  
 gaggtctgga agctctactt caatcacccg caggacggtc cggcgataag ggaccgata  
 aggaagcatc cggccgtcgt tgatatctac gagtacgaca tccccttcgc caagcgctac  
 ctcatcgata aggggctcgt cccgatggag ggggaggagg agctcaaact gatggccttc  
 gacatcgaga ccctctacca cgagggagac gaggtcgggg agggccgat cctgatgata  
 agctacgccg acggggacgg gccgaggggc ataacctgga agaagatcga cctcccctac  
 gtcgacgtcg tctcgaccga gaaggagatg ataaagcgtc tcctccaggt ggtgaaggag  
 aaggaccgcg acgtgctcgt aacttacaac ggcgacaact tcgacttcgc ctacctgaag  
 agacgtccg aggagcttgg attgaagttc atcctcggga gggacgggag cgagcccaag  
 atccagcgca tggcgacccg cttcgccgtc gaggtgaagg ggaggataca cttcgacctc  
 taccgggtga taaggcgcac cgtgaacctg ccgacctaca cgctcgaggc ggtctacgag  
 gccatcttcg ggaggccaaa ggagaaggctc tacgcccggg agatagtgga ggctgggaa  
 accggcgagg gtcttgagag ggttgcccgc tactccatgg aggacgcaaa ggttaccttc  
 gagctcggga gggagttctt cccgatggag gccagctct cgaggctcat cggccagggt  
 ctctgggacg tctccgctc gagcacccgc aacctggtcg agtggttcct cctgaggaag  
 gcctacgaga ggaacgaact ggccccgaac aagccgagcg gccgggaagt ggagatcagg  
 aggcgtggct acgcccgttg ttacgttaag gagccggaga ggggtttatg ggagaaatc  
 gtgtacctcg actttcgctc tctttacccc tccatcatca taaccacaaa cgtctcgccc  
 gataccetaa acagggaggg ctgtgagaac tacgacgtcg cccccaggt gggcataag  
 ttctgcaaag attttcggg cttcatcccg agcctgctcg gaggcctgct tgaggagagg  
 cagaagataa agcggaggat gaaggcctct gtggatcccg ttgagcggaa gctcctcgat  
 tacagcgaga gggccatcaa gatactggcc aacagcttct acggatacta cggctacgcg  
 agggcgaggt ggtactgcag ggagtgcgcg gagagcgtta ccgctgggg cagggagtac  
 atcgataggg tcatcagggg gctcgaggag aagttcggct tcaagggtgct ctaccgggac  
 acggacggac tgcacgccac gatccccggg gccgacgccg ggaccgtcaa ggagagggcg  
 aggggggttc tgagatacat caacccccag ctccccggcc tcctggagct cgagtacgag  
 gggttctacc tgagggggtt cttcgtgacg aagaagaagt acgcggtcat agacgaggag  
 ggcaagataa ccacgcgagg cctcgagata gtcaggcggg actggagcga ggtggccaag  
 gagacgcagg cgagggctct ggagggcagata ctgaggcacg gtgacgtcga ggagccggtt  
 agaatcgtca gggaggtaac cgaaaagctg agcaagtacg aggttcgccc ggagaaactg  
 gtgatccacg agcagataac gagggatttg agggactaca aagccacggg accgcaagtg  
 gcggtggcga agcgcctggc cgggaggggg gtaaggatac gcccggggac ggtgataagc  
 tacatcgtec tcaagggtc cggaaggata ggggacaggg cgattccctt cgacgagttc  
 gaccgcacta agcacaggta cgacgccgac tactacatcg agaaccaggt tctgccagcc  
 gtcgagagga tctgaaggc cttcggctac cgcaaggagg acctgaaata ccagaagacg  
 aggcaggtgg gcctgggtgc gtggctcaac gccgggaagg ggtga

SEQ ID NO: 4 – polypeptide sequence of DNA polymerase from *Thermococcus celer*

Met Ile Leu Asp Ala Asp Tyr Ile Thr Glu Asp Gly Lys Pro Val Val  
 Arg Ile Phe Arg Lys Glu Lys Gly Glu Phe Arg Ile Asp Tyr Asp Arg  
 Asp Phe Glu Pro Tyr Ile Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
 Glu Glu Val Lys Arg Ile Thr Val Glu Arg His Gly Lys Ala Val Arg  
 Val Lys Arg Val Glu Lys Val Glu Lys Lys Phe Leu Asn Arg Pro Ile  
 Glu Val Trp Lys Leu Tyr Phe Asn His Pro Gln Asp Val Pro Ala Ile  
 Arg Asp Glu Ile Arg Lys His Pro Ala Val Val Asp Ile Tyr Glu Tyr  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Val Pro  
 Met Glu Gly Glu Glu Glu Leu Lys Leu Met Ala Phe Asp Ile Glu Thr  
 Leu Tyr His Glu Gly Asp Glu Phe Gly Glu Gly Pro Ile Leu Met Ile  
 Ser Tyr Ala Asp Gly Asp Gly Ala Arg Val Ile Thr Trp Lys Lys Ile  
 Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Lys Glu Met Ile Lys  
 Arg Phe Leu Gln Val Val Lys Glu Lys Asp Pro Asp Val Leu Val Thr  
 Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Arg Arg Ser Glu  
 Glu Leu Gly Leu Lys Phe Ile Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Val Asn Leu Pro Thr  
 Tyr Thr Leu Glu Ala Val Tyr Glu Ala Ile Phe Gly Arg Pro Lys Glu  
 Lys Val Tyr Ala Gly Glu Ile Val Glu Ala Trp Glu Thr Gly Glu Gly  
 Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Phe  
 Glu Leu Gly Arg Glu Phe Phe Pro Met Glu Ala Gln Leu Ser Arg Leu  
 Ile Gly Gln Gly Leu Trp Asp Val Ser Arg Ser Ser Thr Gly Asn Leu  
 Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 Pro Asn Lys Pro Ser Gly Arg Glu Val Glu Ile Arg Arg Arg Gly Tyr  
 Ala Gly Gly Tyr Val Lys Glu Pro Glu Arg Gly Leu Trp Glu Asn Ile  
 Val Tyr Leu Asp Phe Arg Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Glu Asn Tyr Asp  
 Val Ala Pro Gln Val Gly His Lys Phe Cys Lys Asp Phe Pro Gly Phe  
 Ile Pro Ser Leu Leu Gly Gly Leu Leu Glu Glu Arg Gln Lys Ile Lys  
 Arg Arg Met Lys Ala Ser Val Asp Pro Val Glu Arg Lys Leu Leu Asp  
 Tyr Arg Gln Arg Ala Ile Lys Ile Leu Ala Asn Ser Phe Tyr Gly Tyr  
 Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Arg Glu Cys Ala Glu Ser  
 Val Thr Ala Trp Gly Arg Glu Tyr Ile Asp Arg Val Ile Arg Glu Leu  
 Glu Glu Lys Phe Gly Phe Lys Val Leu Tyr Ala Asp Thr Asp Gly Leu  
 His Ala Thr Ile Pro Gly Ala Asp Ala Gly Thr Val Lys Glu Arg Ala  
 Arg Gly Phe Leu Arg Tyr Ile Asn Pro Lys Leu Pro Gly Leu Leu Glu  
 Leu Glu Tyr Glu Gly Phe Tyr Leu Arg Gly Phe Phe Val Thr Lys Lys  
 Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 Glu Ile Val Arg Arg Asp Trp Ser Glu Val Ala Lys Glu Thr Gln Ala  
 Arg Val Leu Glu Ala Ile Leu Arg His Gly Asp Val Glu Glu Ala Val  
 Arg Ile Val Arg Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Asp Leu Arg Asp  
 Tyr Lys Ala Thr Gly Pro His Val Ala Val Ala Lys Arg Leu Ala Gly  
 Arg Gly Val Arg Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 Asp Pro Thr Lys His Arg Tyr Asp Ala Asp Tyr Tyr Ile Glu Asn Gln  
 Val Leu Pro Ala Val Glu Arg Ile Leu Lys Ala Phe Gly Tyr Arg Lys  
 Glu Asp Leu Lys Tyr Gln Lys Thr Arg Gln Val Gly Leu Gly Ala Trp  
 Leu Asn Ala Gly Lys Gly

SEQ ID NO: 5 – nucleic acid sequence of DNA polymerase from *Thermococcus siculi*

atgatacctcg acacggacta catcacggaa gatgggaaac ccgtcataag gatattcaag  
aaagagaacg gcgagttcaa gatcgagtac gacaggactt ttgaacccta catctacgcc  
ctcctgaagg acgactccgc gattgaggat gttaaaaaga taaccgccga gaggcaogga  
acgggtgggga aggtcaagcg cgcgaaaagc gtgcagaaga agttcctagg caggccgggt  
gaagtctgga agctctactt caccaccccc caagatgtcc cggcgataag ggacaagatt  
aggaagcatc cagctgtaat tgacatctac gagtacgaca taccattcgc caagcgctac  
ctcatcgaca agggcctgat tccgatggag ggtgaagaag agcttaagat gctcgccttc  
gacattgaga cgctctacca tgaggggtgag gaggttcgcc aggggcctat tctgatgata  
agctacgccg acgagagcga ggcacgcgtc atcacctgga agaaaatcga cctccccac  
gttgacgtcg tctcaacgga gaaggagatg ataaagcgtt tcctccgcgt tgtgaaggag  
aaagatcccc atgtectcat aacctacaac ggcgacaact tcgacttcgc ctacctgaag  
aagcgtgtg aaaagcttgg aataaacttc ctcttggaa gggacgggag cgagccgaag  
atccagagaa tgggtgaccg cttcgcgctt gaggtgaagg ggaggataca cttcgacctc  
tacctgtaa taaggcgcac gataaacctg ccgacctaca tgcttgaggc agtctacgag  
gcatctttt ggaagccaaa ggagaagggt tactctatgg aggacgcgaa ggtcacggtt  
accggagagg gccttgagag ggtggctcgc tactctatgg aggacgcgaa ggtcacggtt  
gagcttggaa aggagttctt cccgatggag gcccaacttt cgaggttggt cggccagagc  
ttctgggatg tcgcgcgctc aagcacgggc aatctggctg agtggttcct cctcaggaag  
gcctacgaga ggaacgagct ggctccaaac aagccctctg gaagggaaata tgacgagagg  
cgcggtggat acgcggcgg ctacgtcaag gaaccggaaa agggcctgtg ggagaacata  
gtctacctcg actataaatc tctctacccc tcaatcatca tcaccacaaa cgtctcgcct  
gataccctca acccgagagg ctgtaaggag tatgacgtag ctccacaggt cggccaccgc  
ttctgcaagg actttccagg cttcatcccc agcctgctcg gggatctcct ggaggagagg  
cagaagataa agaggaagat gaaggcaaca attgaccgga tcgagagaaa gctccttgat  
tacaggcaac gggccatcaa gatcctteta aatagttttt acggctacta cggctacgca  
agggctcgtt ggtactgcaa ggagtgtgcc gagagcggtt cggcatgggg aagggaatat  
atcaccatga caatcagggg aatagaagag aagtatggct ttaaagtact ttatcgggac  
actgacggct tcttcgcgac gattccccgg gaagatgccg agaccatcaa aaagagggcg  
atggagttcc tcaagtacat aaacgccaaa ctccccggtg cgtcgaact tgagtacgag  
gacttctaca ggcggggctt cttcgtcacc aagaagaaat acgcggttat cgacgaggag  
ggcaagataa caacgcgcg gctggagatc gtcaggcgcg actggagcga gatagccaag  
gagacgcagg cgcgggttct ggaggccctt ctgaaggacg gtgacgtcga agagccggtg  
agcatagtca aagaagtgac cgagaagctg agcaagtacg aggttccgcc ggagaagctc  
gttatccacg agcagataac gcgcgagctg aaggactaca aggcaacggg accacacgtg  
gcatagcga agaggttagc cgcgagaggc gtcaaaaatcc gcccggggac agtcatcagc  
tacatcgtgc tcaagggtc cgggaggata ggcgacaggg cgattccctt cgacgagttc  
gacccacga agcacaagta cgatgcagag tactacatcg agaaccaggt tctacctgcc  
gtcgagagga ttctgaaggc cttcggctat cgcggtgagg agctcagata ccagaagacg  
aggcaggttg gacttggggc gtggctgaag ccgaagggga aggggtga

SEQ ID NO: 6 – polypeptide sequence of DNA polymerase from *Thermococcus siculi*

Met Ile Leu Asp Thr Asp Tyr Ile Thr Glu Asp Gly Lys Pro Val Ile  
 Arg Ile Phe Lys Lys Glu Asn Gly Glu Phe Lys Ile Glu Tyr Asp Arg  
 Thr Phe Glu Pro Tyr Ile Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
 Glu Asp Val Lys Lys Ile Thr Ala Glu Arg His Gly Thr Val Val Lys  
 Val Lys Arg Ala Glu Lys Val Gln Lys Lys Phe Leu Gly Arg Pro Val  
 Glu Val Trp Lys Leu Tyr Phe Thr His Pro Gln Asp Val Pro Ala Ile  
 Arg Asp Lys Ile Arg Lys His Pro Ala Val Ile Asp Ile Tyr Glu Tyr  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Ile Pro  
 Met Glu Gly Glu Glu Glu Leu Lys Met Leu Ala Phe Asp Ile Glu Thr  
 Leu Tyr His Glu Gly Glu Glu Phe Ala Glu Gly Pro Ile Leu Met Ile  
 Ser Tyr Ala Asp Glu Ser Glu Ala Arg Val Ile Thr Trp Lys Lys Ile  
 Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Lys Glu Met Ile Lys  
 Arg Phe Leu Arg Val Val Lys Glu Lys Asp Pro Asp Val Leu Ile Thr  
 Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Lys Arg Cys Glu  
 Lys Leu Gly Ile Asn Phe Leu Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Ile Asn Leu Pro Thr  
 Tyr Met Leu Glu Ala Val Tyr Glu Ala Ile Phe Gly Lys Pro Lys Glu  
 Lys Val Tyr Ala Glu Glu Ile Ala Thr Ala Trp Glu Thr Gly Glu Gly  
 Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Phe  
 Glu Leu Gly Lys Glu Phe Phe Pro Met Glu Ala Gln Leu Ser Arg Leu  
 Val Gly Gln Ser Phe Trp Asp Val Ala Arg Ser Ser Thr Gly Asn Leu  
 Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 Pro Asn Lys Pro Ser Gly Arg Glu Tyr Asp Glu Arg Arg Gly Gly Tyr  
 Ala Gly Gly Tyr Val Lys Glu Pro Glu Lys Gly Leu Trp Glu Asn Ile  
 Val Tyr Leu Asp Tyr Lys Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Lys Glu Tyr Asp  
 Val Ala Pro Gln Val Gly His Arg Phe Cys Lys Asp Phe Pro Gly Phe  
 Ile Pro Ser Leu Leu Gly Asp Leu Leu Glu Glu Arg Gln Lys Ile Lys  
 Arg Lys Met Lys Ala Thr Ile Asp Pro Ile Glu Arg Lys Leu Leu Asp  
 Tyr Arg Gln Arg Ala Ile Lys Ile Leu Leu Asn Ser Phe Tyr Gly Tyr  
 Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Lys Glu Cys Ala Glu Ser  
 Val Thr Ala Trp Gly Arg Glu Tyr Ile Thr Met Thr Ile Arg Glu Ile  
 Glu Glu Lys Tyr Gly Phe Lys Val Leu Tyr Ala Asp Thr Asp Gly Phe  
 Phe Ala Thr Ile Pro Gly Glu Asp Ala Glu Thr Ile Lys Lys Arg Ala  
 Met Glu Phe Leu Lys Tyr Ile Asn Ala Lys Leu Pro Gly Ala Leu Glu  
 Leu Glu Tyr Glu Asp Phe Tyr Arg Arg Gly Phe Phe Val Thr Lys Lys  
 Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln Ala  
 Arg Val Leu Glu Ala Leu Leu Lys Asp Gly Asp Val Glu Glu Ala Val  
 Ser Ile Val Lys Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Glu Leu Lys Asp  
 Tyr Lys Ala Thr Gly Pro His Val Ala Ile Ala Lys Arg Leu Ala Ala  
 Arg Gly Val Lys Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 Asp Pro Thr Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn Gln  
 Val Leu Pro Ala Val Glu Arg Ile Leu Lys Ala Phe Gly Tyr Arg Gly  
 Glu Glu Leu Arg Tyr Gln Lys Thr Arg Gln Val Gly Leu Gly Ala Trp  
 Leu Lys Pro Lys Gly Lys Gly

SEQ ID NO: 7 - polypeptide sequence of DNA polymerase from *Thermococcus kodakarensis*

Met Ile Leu Asp Thr Asp Tyr Ile Thr Glu Asp Gly Lys Pro Val Ile  
 Arg Ile Phe Lys Lys Glu Asn Gly Glu Phe Lys Ile Glu Tyr Asp Arg  
 Thr Phe Glu Pro Tyr Phe Tyr Ala Leu Leu Lys Asp Asp Ser Ala Ile  
 Glu Glu Val Lys Lys Ile Thr Ala Glu Arg His Gly Thr Val Val Thr  
 Val Lys Arg Val Glu Lys Val Gln Lys Lys Phe Leu Gly Arg Pro Val  
 Glu Val Trp Lys Leu Tyr Phe Thr His Pro Gln Asp Val Pro Ala Ile  
 Arg Asp Lys Ile Arg Glu His Pro Ala Val Ile Asp Ile Tyr Glu Tyr  
 Asp Ile Pro Phe Ala Lys Arg Tyr Leu Ile Asp Lys Gly Leu Val Pro  
 Met Glu Gly Asp Glu Glu Leu Lys Met Leu Ala Phe Asp Ile Glu Thr  
 Leu Tyr Glu Glu Gly Glu Glu Phe Ala Glu Gly Pro Ile Leu Met Ile  
 Ser Tyr Ala Asp Glu Glu Gly Ala Arg Val Ile Thr Trp Lys Asn Val  
 Asp Leu Pro Tyr Val Asp Val Val Ser Thr Glu Arg Glu Met Ile Lys  
 Arg Phe Leu Arg Val Val Lys Glu Lys Asp Pro Asp Val Leu Ile Thr  
 Tyr Asn Gly Asp Asn Phe Asp Phe Ala Tyr Leu Lys Lys Arg Cys Glu  
 Lys Leu Gly Ile Asn Phe Ala Leu Gly Arg Asp Gly Ser Glu Pro Lys  
 Ile Gln Arg Met Gly Asp Arg Phe Ala Val Glu Val Lys Gly Arg Ile  
 His Phe Asp Leu Tyr Pro Val Ile Arg Arg Thr Ile Asn Leu Pro Thr  
 Tyr Thr Leu Glu Ala Val Tyr Glu Ala Val Phe Gly Gln Pro Lys Glu  
 Lys Val Tyr Ala Glu Glu Ile Thr Thr Ala Trp Glu Thr Gly Glu Asn  
 Leu Glu Arg Val Ala Arg Tyr Ser Met Glu Asp Ala Lys Val Thr Tyr  
 Glu Leu Gly Lys Glu Phe Leu Pro Met Glu Ala Gln Leu Ser Arg Leu  
 Ile Gly Gln Ser Leu Trp Asp Val Ser Arg Ser Ser Thr Gly Asn Leu  
 Val Glu Trp Phe Leu Leu Arg Lys Ala Tyr Glu Arg Asn Glu Leu Ala  
 Pro Asn Lys Pro Asp Glu Lys Glu Leu Ala Arg Arg Arg Gln Ser Tyr  
 Glu Gly Gly Tyr Val Lys Glu Pro Glu Arg Gly Leu Trp Glu Asn Ile  
 Val Tyr Leu Asp Phe Arg Ser Leu Tyr Pro Ser Ile Ile Ile Thr His  
 Asn Val Ser Pro Asp Thr Leu Asn Arg Glu Gly Cys Lys Glu Tyr Asp  
 Val Ala Pro Gln Val Gly His Arg Phe Cys Lys Asp Phe Pro Gly Phe  
 Ile Pro Ser Leu Leu Gly Asp Leu Leu Glu Arg Gln Lys Ile Lys  
 Lys Lys Met Lys Ala Thr Ile Asp Pro Ile Glu Arg Lys Leu Leu Asp  
 Tyr Arg Gln Arg Ala Ile Lys Ile Leu Ala Asn Ser Tyr Tyr Gly Tyr  
 Tyr Gly Tyr Ala Arg Ala Arg Trp Tyr Cys Lys Glu Cys Ala Glu Ser  
 Val Thr Ala Trp Gly Arg Glu Tyr Ile Thr Met Thr Ile Lys Glu Ile  
 Glu Glu Lys Tyr Gly Phe Lys Val Ile Tyr Ser Asp Thr Asp Gly Phe  
 Phe Ala Thr Ile Pro Gly Ala Asp Ala Glu Thr Val Lys Lys Lys Ala  
 Met Glu Phe Leu Lys Tyr Ile Asn Ala Lys Leu Pro Gly Ala Leu Glu  
 Leu Glu Tyr Glu Gly Phe Tyr Glu Arg Gly Phe Phe Val Thr Lys Lys  
 Lys Tyr Ala Val Ile Asp Glu Glu Gly Lys Ile Thr Thr Arg Gly Leu  
 Glu Ile Val Arg Arg Asp Trp Ser Glu Ile Ala Lys Glu Thr Gln Ala  
 Arg Val Leu Glu Ala Leu Leu Lys Asp Gly Asp Val Glu Lys Ala Val  
 Arg Ile Val Lys Glu Val Thr Glu Lys Leu Ser Lys Tyr Glu Val Pro  
 Pro Glu Lys Leu Val Ile His Glu Gln Ile Thr Arg Asp Leu Lys Asp  
 Tyr Lys Ala Thr Gly Pro His Val Ala Val Ala Lys Arg Leu Ala Ala  
 Arg Gly Val Lys Ile Arg Pro Gly Thr Val Ile Ser Tyr Ile Val Leu  
 Lys Gly Ser Gly Arg Ile Gly Asp Arg Ala Ile Pro Phe Asp Glu Phe  
 Asp Pro Thr Lys His Lys Tyr Asp Ala Glu Tyr Tyr Ile Glu Asn Gln  
 Val Leu Pro Ala Val Glu Arg Ile Leu Arg Ala Phe Gly Tyr Arg Lys  
 Glu Asp Leu Arg Tyr Gln Lys Thr Arg Gln Val Gly Leu Ser Ala Trp  
 Leu Lys Pro Lys Gly Thr

## SAMPLE TAG SEQUENCES

SEQ ID Number	Sample tag sequence
8	TAGAATTGAAGAA
9	TGGCCATAGCTAC
10	GTCATCTGCGACC
11	TTCGCGCTTGGAC
12	CGCGAACCGTTAG
13	TTGCAGCCTCTAA
14	TCTACTAGTACGA
15	GTAGGTTCTACTG
16	GCCAATATCAAGT
17	CTATCTTGCTGGT
18	GTTCTCATAGGTA
19	GTCTATGAACCAA
20	CGGAGCGCTTATT
21	TATGCCATGAGGA
22	ATACGACTCGGAG
23	GATGGAACTCAGC
24	GGACCTGCATGAA
25	TAGACTGGAACTT
26	GAATTACCTCGTT
27	AGGATCAGGCTAC
28	ACGCGTAGAAGAG
29	CTTCGAGACTTAC
30	GACGGCTAACTCC
31	TTAGCATTCTCTT

32	GCAAGGCATAGTA
33	ACCTAGATAATGGA
34	ACGCCAAGGCGTA
35	TATGACGGATCCG
36	CCTCCATTAGAGA
37	ATTGAATACTCTG
38	GAGATGAGAAGAA
39	TCTGAGTAGCCGG
40	AATAGGTAGTACG
41	GTCGAAGAAGTCC
42	TACTGCATCTCGT
43	GACGTATTAGAGC
44	CCTGCATTATTCCG
45	ACGAAATGATGCTC
46	TACTAGCAGAGAT
47	CTCCTCATCTTCC
48	TCCTCTGCGCTGC
49	CCTTCTCAGTCCG
50	CAGCTTCATAGCG
51	TTGACTCTCGCGC
52	TATCCATGAGCGAT
53	AACGCCTAGCCGA
54	CCGAAGACGTCAT
55	GAGTTCTCCAGAT
56	TGCATCCGCGCTT
57	CCTGAACTCAAGT
58	GGTCGTATGCGTA

59	AGGCCTCTCTACC
60	GTACTCCA'CCAA
61	CAGCGGACGCGCT
62	ATCTCTCTTAGCA
63	AAGCAATAATAAT
64	AAGGCGACTCCGA
65	ACGTCTCTAGGAG
66	CCATCAGACCTCT
67	ACTTAATCGTACT
68	TGGAA'CT'CCAA
69	CCATACGATCAGG
70	TTATGGAGCAATA
71	GCTCGGCGTTCGA
72	T'GGCCAGTCGCT
73	CAGATACGTAGAG
74	AATGCTATTATCC
75	GCAGCATGCCGAT
76	GGAGAGTTACCTC
77	GAGAGTCCATGAT
78	CAATCTATTCTGA
79	GC'CT'AGTATCC
80	CCATAGTTATGGT
81	TGCGAGATCGAAG
82	AGAGAAGTCGAGT
83	GGTAACTCCATAT
84	TGCTATTCCAGGC
85	AACCGCGAGGC'CT

86	TTCTAGAGATACC
87	TTCGCTCAAGTAT
88	CAGAGAAGGCGCA
89	TAGAATTGGCCTC
90	GGCCATTCTCCAG
91	TCCAACGCGCGTT
92	GCCGCAGATTACG
93	GCAGTTCGAACGC
94	TTC'TC'TC'GCAGG
95	TAAGCTACCAGCG
96	CTGCATGAGGTTG
97	TTGCCTAGCGAGG
98	CAACTGAATTAGG
99	AAGCGGTCC'TC'T'
100	AAIGGAAGGACCG
101	GAGTTAGTAAGTT
102	TTCCTAATTCCAA
103	GTTCTGGTTCGCT
104	GTTCACTCTTCC
105	ATTCCGAGGAAGA
106	CTTAGCCGAGAGA
107	GTCTGCTACGCTT
108	ATGGCGCCGCGCA
109	TAATTGGTTATCT
110	TCGGTTATAAGTC
111	TGCCTGAGAACGT
112	AGATGCGGTTAAC

113	ATGGAATAGGCGA
114	AGAGATGCGATCG
115	CTCCAAC TAACGT
116	GCC TIGC TAC TGG
117	CTTCGTCTCTACG
118	ACGCTCATAGCCT
119	GTCGAAGATAAGG
120	GCCGGAGTCC TCG
121	TATACGGCGACCT
122	AGGTAGATATTCG
123	TTAAGG TAC TGT
124	CGGATCTGGTATA
125	GAGGTCTCGGAGG
126	GGCATCGATGGAC
127	GATCTCCGATATA
128	GATTCGGAATACT
129	CTGCGATCCGGCC
130	GATCCGGTTGCAA
131	CGTCAGGCTTGAC
132	TCGGCAAGGCGAG
133	GAACGGCGAACGC
134	CCTCAAGCGGACT
135	GAAGCCAGATGGT
136	TGCTCATACCAAT

FIGURE 5





Figure 6






FIGURE 7

— C —  
 — G —