

(12) **DEMANDE DE BREVET CANADIEN**  
**CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/09/06

(87) Date publication PCT/PCT Publication Date: 2020/03/19

(85) Entrée phase nationale/National Entry: 2021/03/05

(86) N° demande PCT/PCT Application No.: US 2019/049859

(87) N° publication PCT/PCT Publication No.: 2020/055672

(30) Priorité/Priority: 2018/09/10 (US62/729,337)


(51) Cl.Int./Int.Cl. *C07D 487/14* (2006.01),  
*A61K 31/551* (2006.01), *A61P 35/00* (2006.01),  
*C07D 487/22* (2006.01), *C07D 498/14* (2006.01),  
*C07D 498/22* (2006.01)

(71) Demandeur/Applicant:  
ARRAY BIOPHARMA INC., US

(72) **Inventeurs/Inventors:**  
BLAKE, JAMES F., US;  
DAI, DONGHUA, US;  
HAAS, JULIA, US;  
JIANG, YUTONG, US;  
KAHN, DEAN, US;  
KOLAKOWSKI, GABRIELLE R., US;

(54) Titre : COMPOSES HETEROCYCLIQUES CONDENSES COMME INHIBITEURS DE KINASES RET

(54) Title: FUSED HETEROCYCLIC COMPOUNDS AS RET KINASE INHIBITORS



(I)

\_\_\_\_\_ (AA)

(57) **Abrégé/Abstract:**

Provided herein are compounds of the Formula (I): (I) and tautomers, stereoisomers and pharmaceutically acceptable salts and solvates thereof, wherein R<sup>x</sup>, R<sup>y</sup>, W, X, Y, Z, Ring A and (AA) have the meanings given in the specification, which are inhibitors of

(72) **Inventeurs(suite)/Inventors(continued)**: MCFADDIN, ELIZABETH A., US; MCKENNEY, MEGAN L., US; METCALF, ANDREW T., US; MORENO, DAVID A., US; PRIGARO, BRETT, US; RAMANN, GINELLE A., US; REN, LI, US; WALLS, SHANE M., US; ZHANG, HAILONG, US

(74) **Agent**: GOWLING WLG (CANADA) LLP

(57) **Abrégé(suite)/Abstract(continued)**:

RET kinase and are useful in the treatment and prevention of diseases which can be treated with a RET kinase inhibitor, including RET-associated diseases and disorders.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

19 March 2020 (19.03.2020)



(10) International Publication Number

**WO 2020/055672 A1**

## (51) International Patent Classification:

C07D 487/14 (2006.01) C07D 498/22 (2006.01)

C07D 487/22 (2006.01) A61P 35/00 (2006.01)

C07D 498/14 (2006.01) A61K 31/551 (2006.01)

## (21) International Application Number:

PCT/US2019/049859

## (22) International Filing Date:

06 September 2019 (06.09.2019)

## (25) Filing Language:

English

## (26) Publication Language:

English

## (30) Priority Data:

62/729,337 10 September 2018 (10.09.2018) US

(71) Applicant: **ARRAY BIOPHARMA INC.** [US/US]; 3200 Walnut St., Boulder, Colorado 80301 (US).


(72) Inventors: **BLAKE, James F.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **DAI, Donghua**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **HAAS, Julia**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **JIANG, Yutong**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **KAHN, Dean**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **KOLAKOWSKI, Gabrielle R.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **MCFADDIN, Elizabeth A.**; c/o Eli Lilly and Company,

P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **MCKENNEY, Megan L.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **MET-CALF, Andrew T.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **MORENO, David A.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **PRIGARO, Brett**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **RAMANN, Ginnelle A.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **REN, Li**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **WALLS, Shane M.**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US). **ZHANG, Hailong**; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US).

(74) Agent: **ASAM, Michael R.** et al.; c/o Eli Lilly and Company, P.O. Box 6288, Indianapolis, Indiana 46206-6288 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

## (54) Title: FUSED HETEROCYCLIC COMPOUNDS AS RET KINASE INHIBITORS



(I)

----- (AA)

(57) Abstract: Provided herein are compounds of the Formula (I): (I) and tautomers, stereoisomers and pharmaceutically acceptable salts and solvates thereof, wherein R<sup>x</sup>, R<sup>y</sup>, W, X, Y, Z, Ring A and (AA) have the meanings given in the specification, which are inhibitors of RET kinase and are useful in the treatment and prevention of diseases which can be treated with a RET kinase inhibitor, including RET-associated diseases and disorders.

**WO 2020/055672 A1** 

SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,  
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Declarations under Rule 4.17:**

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

**Published:**

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*



## DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME        1    DE    2  
CONTENANT LES PAGES    1    À    318

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

## JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME        1    OF    2  
CONTAINING PAGES    1    TO    318

NOTE: For additional volumes, please contact the Canadian Patent Office

NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:

## FUSED HETEROCYCLIC COMPOUNDS AS RET KINASE INHIBITORS

## BACKGROUND

**[0001]** The present disclosure relates to novel compounds which exhibit Rearranged during Transfection (RET) kinase inhibition, pharmaceutical compositions comprising the compounds, processes for making the compounds, and the use of the compounds in therapy. More particularly, it relates to fused heterocyclic compounds useful in the treatment and prevention of diseases which can be treated with a RET kinase inhibitor, including RET-associated diseases and disorders.

**[0002]** RET is a single-pass transmembrane receptor belonging to the tyrosine kinase superfamily that is required for normal development, maturation and maintenance of several tissues and cell types (Mulligan, L. M., *Nature Reviews Cancer*, 2014, 14, 173-186). The extracellular portion of the RET kinase contains four calcium-dependent cadherin-like repeats involved in ligand binding and a juxtamembrane cysteine-rich region necessary for the correct folding of the RET extracellular domain, while the cytoplasmic portion of the receptor includes two tyrosine kinase subdomains.

**[0003]** RET signaling is mediated by the binding of a group of soluble proteins of the glial cell line-derived neurotrophic factor (GDNF) family ligands (GFLs), which also includes neurturin (NTRN), artemin (ARTN) and persephin (PSPN) (Arighi et al., *Cytokine Growth Factor Rev.*, 2005, 16, 441-67). Unlike other receptor tyrosine kinases, RET does not directly bind to GFLs and requires an additional co-receptor: that is, one of four GDNF family receptor- $\alpha$  (GFR $\alpha$ ) family members, which are tethered to the cell surface by a glycosylphosphatidylinositol linkage. GFLs and GFR $\alpha$  family members form binary complexes that in turn bind to RET and recruit it into cholesterol-rich membrane subdomains, which are known as lipid rafts, where RET signaling occurs.


**[0004]** Upon binding of the ligand-co-receptor complex, RET dimerization and autophosphorylation on intracellular tyrosine residues recruits adaptor and signaling proteins to stimulate multiple downstream pathways. Adaptor protein binding to these docking sites leads to activation of Ras-MAPK and PI3K-Akt/mTOR signaling pathways or to recruitment of the CBL family of ubiquitin ligases that functions in RET downregulation of the RET-mediated functions.

**[0005]** Aberrant RET expression and/or activity have been demonstrated in different cancers and in gastrointestinal disorders such as irritable bowel syndrome (IBS).

## SUMMARY OF THE INVENTION

**[0006]** It has now been found that certain fused heterocyclic compounds are inhibitors of RET kinase, and are useful for treating diseases such as proliferative diseases such as cancers.

**[0007]** Accordingly, provided herein is a compound of the Formula I:



**[0008]** and tautomers, stereoisomers, and pharmaceutically acceptable salts and solvates thereof, wherein  $R^x$ ,  $R^y$ , W, X, Y, Z, Ring A and ----- are as defined herein.

**[0009]** Also provided herein is a pharmaceutical composition comprising a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, in admixture with a pharmaceutically acceptable diluent or carrier.

**[0010]** Also provided herein is a method of inhibiting cell proliferation, in vitro or in vivo, the method comprising contacting a cell with an effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof as defined herein.

**[0011]** Also provided herein is a method of treating a RET-associated disease or disorder in a patient in need of such treatment, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof as defined herein.

**[0012]** Also provided herein is a method of treating cancer and/or inhibiting metastasis associated with a particular cancer in a patient in need of such treatment, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein.

**[0013]** Also provided herein is a method of treating irritable bowel syndrome (IBS) and/or pain associated with IBS in a patient in need of such treatment, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein.

**[0014]** Also provided is a method of providing supportive care to a cancer patient, including preventing or minimizing gastrointestinal disorders, such as diarrhea, associated with

treatment, including chemotherapeutic treatment, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein.

**[0015]** Also provided herein is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof as defined herein for use in therapy.

**[0016]** Also provided herein is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein for use in the treatment of cancer and/or inhibiting metastasis associated with a particular cancer.

**[0017]** Also provided herein is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein for use in the treatment of irritable bowel syndrome (IBS) or pain associated with IBS.

**[0018]** Also provided is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein for use providing supportive care to a cancer patient, including preventing or minimizing gastrointestinal disorders, such as diarrhea, associated with treatment, including chemotherapeutic treatment.

**[0019]** Also provided herein is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof for use in the inhibition of RET kinase activity.

**[0020]** Also provided herein is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof as defined herein, for use in the treatment of a RET-associated disease or disorder.

**[0021]** Also provided herein is the use of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, as defined herein in the manufacture of a medicament for the treatment of cancer and/or inhibiting metastasis associated with a particular cancer.

**[0022]** Also provided herein is the use of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, as defined herein in the manufacture of a medicament for the treatment of irritable bowel syndrome (IBS) or pain associated with IBS.

**[0023]** Also provided herein is the use of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, as defined herein in the manufacture of a medicament for providing supportive care to a cancer patient, including preventing or minimizing gastrointestinal disorders, such as diarrhea, associated with treatment, including chemotherapeutic treatment.

**[0024]** Also provided herein is a use of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, as defined herein in the manufacture of a medicament for the inhibition of RET kinase activity.

**[0025]** Also provided herein is the use of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, as defined herein, in the manufacture of a medicament for the treatment of a RET-associated disease or disorder.

**[0026]** Also provided herein is a method for treating cancer in a patient in need thereof, the method comprising (a) determining if the cancer is associated with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same (e.g., a RET-associated cancer); and (b) if the cancer is determined to be associated with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same (e.g., a RET-associated cancer), administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof.

**[0027]** Also provided herein is a pharmaceutical combination for treating cancer (e.g., a RET-associated cancer, such as a RET-associated cancer having one or more RET inhibitor resistance mutations) in a patient in need thereof, which comprises (a) a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, (b) an additional therapeutic agent, and (c) optionally at least one pharmaceutically acceptable carrier, wherein the compound of Formula I or the pharmaceutically acceptable salt or solvate thereof and the additional therapeutic are formulated as separate compositions or dosages for simultaneous, separate or sequential use for the treatment of cancer, wherein the amounts of the compound of Formula I or a pharmaceutically acceptable salt or solvate thereof and of the additional therapeutic agent are together effective in treating the cancer. Also provided herein is a pharmaceutical composition comprising such a combination. Also provided herein is the use of such a combination for the preparation of a medicament for the treatment of cancer. Also provided herein is a commercial package or product comprising such a combination as a combined preparation for simultaneous, separate or sequential use; and to a method of treatment of cancer a patient in need thereof.

**[0028]** Also provided herein is a method for reversing or preventing acquired resistance to an anticancer drug, comprising administering a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, to a patient at risk for developing or having acquired resistance to an anticancer drug. In some embodiments, the patient is administered a dose of the anticancer drug (e.g., at substantially the same time as a dose of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof is administered to the patient).

**[0029]** Also provided herein is a method of delaying and/or preventing development of cancer resistant to an anticancer drug in an individual, comprising administering to the individual an effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, before, during, or after administration of an effective amount of the anticancer drug.

**[0030]** Also provided herein is a method of treating an individual with cancer who has an increased likelihood of developing resistance to an anticancer drug, comprising administering to the individual (a) an effective amount of a compound of Formula I before, during, or after administration of (b) an effective amount of the anticancer drug.

**[0031]** Also provided are methods of treating an individual with a RET-associated cancer that has one or more RET inhibitor resistance mutations that increase resistance of the cancer to a first RET inhibitor (e.g., one or more amino acid substitutions in the kinase domain (e.g., amino acid positions 700 to 1012 in a wildtype RET protein), a gatekeeper amino acid (e.g., amino acid position 804 in a wildtype RET protein), the P-loop (e.g., amino acid positions 730-737 in a wildtype RET protein), the X-DFG residue (e.g., amino acid position 891 in a wildtype RET protein), ATP cleft solvent front amino acids (e.g., amino acid positions 806-811 in a wildtype RET protein), the activation loop (e.g., amino acid positions 891-916 in a wildtype RET protein), the C-helix and loop preceeding the C-helix (e.g., amino acid positions 768-788 in a wildtype RET protein), and/or the ATP binding site (e.g., amino acid positions 730-733, 738, 756, 758, 804, 805, 807, 811, 881, and 892 in a wildtype RET protein) (e.g., a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D, and/or one or more RET inhibitor resistance mutations listed in Tables 3 and 4), that include administering a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, before, during, or after administration of another anticancer drug (e.g., a second RET kinase inhibitor). See also J. Kooistra, G. K. Kanev, O. P. J. Van Linden, R. Leurs, I. J. P. De Esch, and C. De Graaf, "KLIFS: A structural kinase-ligand interaction database," *Nucleic Acids Res.*, vol. 44, no. D1, pp. D365-D371, 2016; and O. P. J. Van Linden, A. J. Kooistra, R. Leurs, I. J. P. De Esch, and C. De Graaf, "KLIFS: A knowledge-based structural database to navigate kinase-ligand interaction space," *J. Med. Chem.*, vol. 57, no. 2, pp. 249-277, 2014, both of which are incorporated by reference in their entirety herein. In some embodiments, a wildtype RET protein is the exemplary wildtype RET protein described herein.

**[0032]** Also provided are methods of treating an individual with a RET-associated cancer that include administering a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, before, during, or after administration of another anticancer drug (e.g., a first RET kinase inhibitor or another kinase inhibitor).

**[0033]** Also provided herein is a method for treating irritable bowel syndrome (IBS) in a patient in need thereof, the method comprising (a) determining if the IBS is associated with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same; and (b) if the IBS is determined to be associated with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, administering to the patient a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof.

**[0034]** Also provided herein is a pharmaceutical combination for treating irritable bowel syndrome (IBS) in a patient in need thereof, which comprises administering (a) a compound of General Formula I or a pharmaceutically acceptable salt or solvate thereof, (b) an additional therapeutic agent, and (c) optionally at least one pharmaceutically acceptable carrier, for simultaneous, separate or sequential

use for the treatment of IBS, wherein the amounts of the compound of Formula I or a pharmaceutically acceptable salt or solvate thereof and of the additional therapeutic agent are together effective in treating the IBS. Also provided herein is a pharmaceutical composition comprising such a combination. Also provided herein is the use of such a combination for the preparation of a medicament for the treatment of the IBS. Also provided herein is a commercial package or product comprising such a combination as a combined preparation for simultaneous, separate or sequential use; and to a method of treatment of the IBS a patient in need thereof.

**[0035]** Also provided herein is a process for preparing a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof.


**[0036]** Also provided herein is a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof obtained by a process of preparing the compound as defined herein.

**[0037]** Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

**[0038]** Other features and advantages of the invention will be apparent from the following detailed description and figures, and from the claims.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0039]** Provided herein is a compound of Formula I:



**[0040]** and tautomers, stereoisomers, and pharmaceutically acceptable salts and solvates thereof, wherein:

**[0041]** W is CR<sup>3</sup>R<sup>4</sup>,

**[0042]** ----- represents a single or double bond, wherein when ----- is a double bond, then X is CR<sup>5</sup> and Y is CR<sup>6</sup>, and when ----- is a single bond, then X is O, CR<sup>7</sup>R<sup>8</sup> or NR<sup>9</sup> and Y is C(=O), SO<sub>2</sub>, CR<sup>10</sup>R<sup>11</sup> or O, and

**[0043]** Z is C or N;

**[0044]**

**[0045]** R<sup>3</sup> is hydrogen;

**[0046]** R<sup>4</sup> is hydrogen;

**[0047]** R<sup>5</sup> is hydrogen;

**[0048]** R<sup>6</sup> is hydrogen;

**[0049]** R<sup>7</sup> is hydrogen or C1-C6 alkyl;

**[0050]** R<sup>8</sup> is hydrogen;

**[0051]** R<sup>9</sup> is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)-;

**[0052]** R<sup>10</sup> is hydrogen, hydroxy, C1-C6 alkoxy, or R<sup>a</sup>R<sup>b</sup>N- wherein R<sup>a</sup> and R<sup>b</sup> are independently H or C1-C6 alkyl;

**[0053]** R<sup>11</sup> is hydrogen, C1-C6 alkyl, or fluoroC1-C6 alkyl;

**[0054]** hetCyc<sup>1</sup> is a 5-6 membered saturated heterocyclic ring having 1-2 ring heteroatoms independently selected from N and SO<sub>2</sub>, wherein said heterocyclic ring is optionally substituted with C1-C6 alkyl or (Ar<sup>2</sup>)C1-C6 alkyl-;

**[0055]** Ar<sup>1</sup> is phenyl optionally substituted with one or more substituents independently selected from halogen and C1-C6 alkoxy;

**[0056]** hetAr<sup>1</sup> is a 5-6 membered heteroaryl ring having 1-2 ring nitrogen atoms, wherein said heteroaryl ring is optionally substituted with C1-C6 alkyl;

**[0057]** Ar<sup>2</sup> is phenyl optionally substituted with one or more substituents independently selected from halogen and C1-C6 alkoxy;

**[0058]** Ring A, including the atoms at the points of attachment to the ring containing Z, Y, X, and W, is (i) a monocyclic aryl ring when Z is C, wherein said ring is optionally substituted with halogen,



C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH-, (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O when Z is C, or when Z is N, a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said rings are optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or (iii) when Z is N, a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH-;

**[0059]** R<sup>c</sup> is C3-C6 cycloalkyl or (C3-C6 cycloalkyl)CH<sub>2</sub>-;

**[0060]** R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>;

**[0061]** Cyc<sup>1</sup> is C3-C6 cycloalkyl optionally substituted with C1-C6 alkyl;

**[0062]** hetAr<sup>2</sup> is (i) pyrazolyl optionally substituted with 1-2 substituents independently selected from C1-C6 alkyl, fluoroC1-C6 alkyl, (di-C1-C6 alkyl)NC(=O)CH<sub>2</sub>-, (hetCyc<sup>2</sup>)CH<sub>2</sub>CH<sub>2</sub>-, phenyl, and (C1-C6 alkoxy)C1-C6 alkyl, or (ii) pyridyl;

**[0063]** hetCyc<sup>2</sup> is a 6-membered saturated heterocyclic ring having 2 ring heteroatoms independently selected from N and O;

**[0064]** R<sup>x</sup> is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>;

**[0065]** Cyc<sup>2</sup> is a 3-6 membered saturated carbocyclic ring optionally substituted with 1-2 substituents independently selected from halogen, C1-C6 alkyl, hydroxy, hydroxyC1-C6 alkyl, and (C1-C6 alkoxy)C(=O)-;

**[0066]** Cyc<sup>3</sup> is cyclopentenyl;

**[0067]** hetCyc<sup>3</sup> is a 5-membered saturated heterocyclic ring having a ring heteroatom selected from N and O and optionally substituted with (C1-C6 alkyl)C(=O)-;

**[0068]** Ar<sup>3</sup> is phenyl optionally substituted with (C1-C6 alkoxy)C(=O)-;

**[0069]** hetAr<sup>3</sup> is a 5-6 membered heteroaryl ring having 1-2 ring nitrogen atoms and optionally substituted with C1-C6 alkyl or C1-C6 alkoxy; and

**[0070]** R<sup>y</sup> is hydrogen or halogen.

**[0071]** For complex chemical names employed herein, a substituent group is typically named before the group to which it attaches. For example, methoxyethyl comprises an ethyl backbone with a methoxy substituent.

**[0072]** The term "halogen" means -F (sometimes referred to herein as "fluoro" or "fluoros"), -Cl, -Br and -I.

**[0073]** The term "C1-C6 alkyl" as used herein refers to saturated linear or branched-chain monovalent hydrocarbon radicals of one to six carbon atoms. Examples include, but are not limited to,

methyl, ethyl, 1-propyl, isopropyl, 1-butyl, isobutyl, sec-butyl, tert-butyl, 2-methyl-2-propyl, pentyl, neopentyl, and hexyl.

**[0074]** The term "fluoroC1-C6 alkyl" as used herein refers to a C1-C6 alkyl radical as defined herein, wherein one to three hydrogen atoms is replaced with one to three fluoro atoms, respectively. Examples include, but are not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, 2-fluoroethyl, 2,2-difluoroethyl, and 2,2,2-trifluoroethyl.

**[0075]** The term "C1-C6 alkoxy" as used herein refers to saturated linear or branched-chain monovalent alkoxy radicals of one to six carbon atoms, wherein the radical is on the oxygen atom. Examples include methoxy, ethoxy, propoxy, isopropoxy, butoxy and tert-butoxy.

**[0076]** The term "(C1-C6 alkoxy)C1-C6 alkyl" as used herein refers to saturated linear or branched-chain monovalent radicals of one to six carbon atoms, wherein one of the carbon atoms is substituted with a C1-C6 alkoxy group as defined herein. Examples include methoxymethyl ( $\text{CH}_3\text{OCH}_2-$ ) and methoxyethyl ( $\text{CH}_3\text{OCH}_2\text{CH}_2-$ ).

**[0077]** The term "hydroxyC1-C6 alkyl", as used herein refers to saturated linear or branched-chain monovalent alkyl radicals of one to six or two to six carbon atoms, respectively, wherein one of the carbon atoms is substituted with a hydroxy group.

**[0078]** The term "C3-C6 cycloalkyl" as used herein refers to cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.


**[0079]** The term "(hetCyc<sup>1</sup>)C1-C6 alkyl-" as used herein refers to saturated linear or branched-chain monovalent alkyl radicals of one to six carbon atoms, wherein one of the carbon atoms is substituted with a hetCyc<sup>1</sup> group, wherein hetCyc<sup>1</sup> is as defined herein.

**[0080]** The term "(Ar<sup>1</sup>)C1-C6 alkyl-" as used herein refers to saturated linear or branched-chain monovalent alkyl radicals of one to six carbon atoms, wherein one of the carbon atoms is substituted with an Ar<sup>1</sup> group, wherein Ar<sup>1</sup> is as defined herein.

**[0081]** The term "(hetAr<sup>1</sup>)C1-C6 alkyl-" as used herein refers to saturated linear or branched-chain monovalent alkyl radicals of one to six carbon atoms, wherein one of the carbon atoms is substituted with a hetAr<sup>1</sup> group, wherein hetAr<sup>1</sup> is as defined herein.


**[0082]** The term "compound," as used herein is meant to include all stereoisomers, geometric isomers, tautomers, and isotopes of the structures depicted. Compounds herein identified by name or structure as one particular tautomeric form are intended to include other tautomeric forms unless otherwise specified.

**[0083]** The term "tautomer" as used herein refers to compounds whose structures differ markedly in arrangement of atoms, but which exist in easy and rapid equilibrium, and it is to be understood that compounds provided herein may be depicted as different tautomers, and when compounds have tautomeric forms, all tautomeric forms are intended to be within the scope of the invention, and the naming of the compounds does not exclude any tautomer. An example of a tautomeric forms includes the following example:




**[0084]** It will be appreciated that certain compounds provided herein may contain one or more centers of asymmetry and may therefore be prepared and isolated in a mixture of isomers such as a racemic mixture, or in an enantiomerically pure form.

**[0085]** Where a dashed line (----) appears in a structure, the dashed line represents a bond that is optionally present, indicating, together with the single bond to which it is adjacent, either a single or double bond. Accordingly, when  is a single bond, the general structure of Formula I may be represented as:



**[0086]** and when  is a double bond, the general structure of Formula I may be represented as:





**[0087]** It is to be understood that the ring in compounds of Formula I comprising atoms W, X, Y and Z does not contain two adjacent oxygen atoms or two adjacent S atoms.

[0088] In one embodiment of Formula I, R<sup>x</sup> is hydrogen.

[0089] In one embodiment of Formula I, R<sup>x</sup> is C1-C6 alkyl. In one embodiment, R<sup>x</sup> is isopropyl.


[0090] In one embodiment of Formula I, R<sup>x</sup> is fluoroC1-C6 alkyl. In one embodiment, R<sup>x</sup> is 1,3-difluoropropan-2-yl or 1,1,1-trifluoropropan-2-yl.

[0091] In one embodiment of Formula I, R<sup>x</sup> is Cyc<sup>2</sup>. In one embodiment, R<sup>x</sup> is cyclopropyl, cyclobutyl, cyclopentyl optionally substituted with 1-2 substituents independently selected from halogen, C1-C6 alkyl, hydroxy, hydroxyC1-C6 alkyl, and (C1-C6 alkoxy)C(=O)-. Non-limiting examples when R<sup>x</sup> is Cyc<sup>2</sup> include the structures:



[0092] In one embodiment of Formula I, R<sup>x</sup> is Cyc<sup>3</sup>. In one embodiment, R<sup>x</sup> is cyclopentenyl.

[0093] In one embodiment of Formula I, R<sup>x</sup> is hetCyc<sup>3</sup>. Non-limiting examples include the structures:




[0094] In one embodiment of Formula I, R<sup>x</sup> is Ar<sup>3</sup>. Non-limiting examples include phenyl and (4-methoxycarbonyl)phenyl.

[0095] In one embodiment of Formula I, R<sup>x</sup> is hetAr<sup>3</sup>. Non-limiting examples include pyridin-2-yl, pyridin-3-yl, 6-methoxypyridin-3-yl and 1-methylpyrazol-4-yl.

[0096] In one embodiment of Formula I, R<sup>y</sup> is hydrogen.


**[0097]** In one embodiment of Formula I,  $R^y$  is halogen. In one embodiment of Formula I,  $R^y$  is chloro.

**[0098]** In one embodiment of Formula I, Z is carbon and Ring A, including the atoms at the points of attachment to the ring containing Z, Y, X, and W, is a monocyclic aryl ring optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  and  $R^d$  are as defined for Formula I. Accordingly, when Ring A is a monocyclic aryl ring, and Z is carbon, Ring A may be represented by formula (i):




(i)

**[0099]** wherein the bond labeled "a" indicates the point of attachment to Y. Non-limiting examples when Z is C, Ring A is a monocyclic aryl ring and  $R^y$  is hydrogen or halogen include the structures:




**[00100]** In one embodiment, Ring A is a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O when Z is C, or when Z is N, a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said rings are optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

**[00101]** In one embodiment, Z is N and Ring A is a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said heteroaryl ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment when Z is N, Ring A can be represented by the formula (ii):






**[00102]** wherein the bond labeled "a" indicates the point of attachment to Y, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. Non-limiting examples include the structures:




**[00103]** wherein the bond labeled "a" indicates the point of attachment to Y.

**[00104]** In one embodiment, Z is C and Ring A is a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, and R<sup>y</sup> is hydrogen or halogen. In one embodiment, Z is C, Ring A is a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, and R<sup>y</sup> is hydrogen. In one embodiment when Z is C, Ring A can be represented by formulas (iii-1), (iii-2) and (iii-3):




**[00105]** wherein the bond labeled "a" indicates the point of attachment to Y, wherein each of said formulas (iii-1), (iii-2) and (iii-3) is optionally substituted C1-C6 alkyl or C3-C6 cycloalkyl. Non-limiting examples include the structures:





**[00106]** wherein the bond labeled "a" indicates the point of attachment to Y.

**[00107]** In one embodiment, Z is N, R<sup>y</sup> is hydrogen or halogen, and Ring A is a bicyclic heteroaryl ring having one ring nitrogen heteroatom that results when Z is N, wherein said bicyclic heteroaryl ring is optionally substituted with R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH-. In one embodiment, Z is N and Ring A is a 5,6-bicyclic heteroaryl ring having one ring nitrogen heteroatom that results when Z is N, wherein said bicyclic heteroaryl ring is optionally substituted with R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH-. In one embodiment when Z is N and Ring A is a bicyclic heteroaryl ring, Ring A can be represented by formula (iv):




**[00108]** wherein the bond labeled "a" indicates the point of attachment to Y and formula (iv) is optionally substituted with R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH-. Non-limiting examples when Z is N, Ring A is a bicyclic heteroaryl ring, and R<sup>y</sup> is hydrogen or halogen include the structures:










[00109] wherein the bond labeled "a" indicates the point of attachment to Y.


[00110] In one embodiment of Formula I, W is  $\text{CR}^3\text{R}^4$ , X is  $\text{CR}^7\text{R}^8$ , Y is  $\text{C}(=\text{O})$ , Z is N, and  $\text{-----}$  is a single bond, wherein Formula I can be represented by Formula I-A:



**I-A**

**[00111]** wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^7$  and  $R^8$  are as defined for Formula I and Ring A is (ii) a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or (iii) a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^d\text{NHC}(=\text{O})-$ , (C1-C6 alkoxy) $\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $\text{C}(=\text{O})\text{NH}-$ .

**[00112]** In one embodiment, Ring A is a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^d\text{NHC}(=\text{O})-$ , (C1-C6 alkoxy) $\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $\text{C}(=\text{O})\text{NH}-$ . In one embodiment, Ring A is a 5,6-bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^d\text{NHC}(=\text{O})-$ , (C1-C6 alkoxy) $\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $\text{C}(=\text{O})\text{NH}-$ . In one embodiment, Ring A is a 5,6-bicyclic heteroaryl ring optionally substituted with  $R^d\text{NHC}(=\text{O})-$ . In one embodiment, Ring A is a 5,6-bicyclic heteroaryl ring optionally substituted with  $R^d\text{NHC}(=\text{O})-$  wherein  $R^d$  is C3-C6 cycloalkyl. In one embodiment of Formula **I-A**, Ring A is selected from the structures:




**[00113]** wherein the bond labeled "a" indicates the point of attachment to Y.

**[00114]** In one embodiment of Formula **I-A**,  $R^7$  and  $R^8$  are hydrogen.

**[00115]** In one embodiment of Formula **I-A**,  $R^x$  is C1-C6 alkyl.


**[00116]** In one embodiment of Formula **I-A**,  $R^y$  is hydrogen.

**[00117]** In one embodiment of Formula I, W is  $\text{CR}^3\text{R}^4$ , X is  $\text{NR}^9$ , Y is  $\text{C}(=\text{O})$ , Z is N, and  $\text{-----}$  is a single bond, wherein Formula I can be represented by Formula **I-B**:

**I-B**


**[00118]** wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ , and  $R^9$  are as defined for Formula I and Ring A is (ii) a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or (iii) a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^d\text{NHC}(=\text{O})-$ ,  $(\text{C1-C6 alkoxy})\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or  $(\text{C1-C6 alkyl})\text{C}(=\text{O})\text{NH}-$ .


**[00119]** In one embodiment of Formula I-B, Ring A is a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is a pyrazolyl ring optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-B, Ring A is selected from the structures:




**[00120]** wherein the bond labeled "a" indicates the point of attachment to Y.

**[00121]** In one embodiment of Formula I-B, Ring A is a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^d\text{NHC}(=\text{O})-$ ,  $(\text{C1-C6 alkoxy})\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or  $(\text{C1-C6 alkyl})\text{C}(=\text{O})\text{NH}-$ . In one embodiment of Formula I-B, Ring A is a 5,6-bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^d\text{NHC}(=\text{O})-$ ,  $(\text{C1-C6 alkoxy})\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or  $(\text{C1-C6 alkyl})\text{C}(=\text{O})\text{NH}-$ . In one embodiment of Formula I-B, Ring A is selected from the structures:







[00122] wherein the bond labeled "a" indicates the point of attachment to Y.


[00123] In one embodiment of Formula I-B,  $R^9$  is hydrogen.

[00124] In one embodiment of Formula I-B,  $R^x$  is C1-C6 alkyl.

[00125] In one embodiment of Formula I-B,  $R^y$  is hydrogen.


[00126] In one embodiment of Formula I-B,  $R^y$  is halogen. In one embodiment of Formula I-B,  $R^y$  is chloro.

[00127] In one embodiment, W is  $CR^3R^4$ , X is  $NR^9$ , Y is  $C(=O)$ , Z is C, and ----- is a single bond, wherein Formula I can be represented by Formula I-C:




[00128] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ , and  $R^9$  is as defined for Formula I, and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

[00129] In one embodiment of Formula I-C, Ring A is a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I. In one embodiment of Formula I-C, Ring A is selected from the structures:



[00130] wherein the bond labeled "a" indicates the point of attachment to Y.

[00131] In one embodiment of Formula I-C, Ring A is a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-C, Ring A is an isoxazolyl ring or a pyrazolyl ring optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-C, Ring A is selected from the structures:



[00132] wherein the bond labeled "a" indicates the point of attachment to Y. In one embodiment of Formula I-C, Ring A is 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, with the exception that Ring A does not include the structures:



[00133] wherein the bond labeled "a" indicates the point of attachment to Y.

**[00134]** In one embodiment of Formula I-C, R<sup>x</sup> is hydrogen, C1-C6 alkyl, or Cyc<sup>2</sup>.


**[00135]** In one embodiment of Formula I-C, R<sup>x</sup> is C1-C6 alkyl.

**[00136]** In one embodiment of Formula I-C, R<sup>9</sup> is hydrogen, C1-C6 alkyl, (C1-C6 alkoxy)C1-C6 alkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, or (Ar<sup>1</sup>)C1-C6 alkyl-.

**[00137]** In one embodiment of Formula I-C, R<sup>y</sup> is hydrogen.

**[00138]** In one embodiment of Formula I-C, R<sup>y</sup> is halogen. In one embodiment of Formula I-C, R<sup>y</sup> is chloro.


**[00139]** In one embodiment, W is CR<sup>3</sup>R<sup>4</sup>, X is O, Y is C(=O), Z is C, and ----- is a single bond, wherein Formula I can be represented by Formula I-D



**[00140]** wherein R<sup>x</sup>, R<sup>y</sup>, R<sup>3</sup>, and R<sup>4</sup> are as defined for Formula I and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH- wherein R<sup>c</sup> is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

**[00141]** In one embodiment of Formula I-D, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-D, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-D, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-D, Ring A is:






[00142] wherein the bond labeled "a" indicates the point of attachment to Y.

[00143] In one embodiment of Formula I-D,  $R^x$  is C1-C6 alkyl.


[00144] In one embodiment of Formula I-D,  $R^y$  is hydrogen.

[00145] In one embodiment, W is  $CR^3R^4$ , X is  $NR^9$ , Y is  $SO_2$ , Z is C, and  $\text{-----}$  is a single bond, wherein Formula I can be represented by Formula I-E:




[00146] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ , and  $R^9$  are as defined for Formula I, and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.


[00147] In one embodiment of Formula I-E, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-E, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-E, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-E, Ring A is:




- [00148] wherein the bond labeled "a" indicates the point of attachment to Y.
- [00149] In one embodiment of Formula I-E,  $R^x$  is C1-C6 alkyl.
- [00150] In one embodiment of Formula I-E,  $R^9$  is hydrogen or C1-C6 alkyl.
- [00151] In one embodiment of Formula I-E,  $R^9$  is hydrogen.
- [00152] In one embodiment of Formula I-E,  $R^9$  is C1-C6 alkyl.
- [00153] In one embodiment of Formula I-E,  $R^y$  is hydrogen.
- [00154] In one embodiment, W is  $CR^3R^4$ , X is  $CR^7R^8$ , Y is  $SO_2$ , Z is C, and ----- is a single bond, wherein the compound has the structure of Formula I-F:



- [00155] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^7$ , and  $R^8$  are as defined for Formula I and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.
- [00156] In one embodiment of Formula I-F, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-F, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-F, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-F, Ring A is:




- [00157] wherein the bond labeled "a" indicates the point of attachment to Y.
- [00158] In one embodiment of Formula I-F,  $R^x$  is C1-C6 alkyl.
- [00159] In one embodiment of Formula I-F,  $R^7$  and  $R^8$  are hydrogen.
- [00160] In one embodiment of Formula I-F,  $R^y$  is hydrogen.
- [00161] In one embodiment, W is  $CR^3R^4$ , X is  $CR^7R^8$ , Y is  $C(=O)$ , Z is C, and ----- is a single bond, wherein the compound has the structure of Formula I-G:



I-G

- [00162] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^7$ , and  $R^8$  are as defined for Formula I and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.
- [00163] In one embodiment of Formula I-G, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-G, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-G, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-G, Ring A is:




- [00164] wherein the bond labeled "a" indicates the point of attachment to Y.

[00165] In one embodiment of Formula I-G,  $R^x$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or hetCyc<sup>3</sup>.

[00166] In one embodiment of Formula I-G,  $R^7$  and  $R^8$  are hydrogen.


[00167] In one embodiment of Formula I-G,  $R^y$  is hydrogen.

[00168] In one embodiment, W is  $CR^3R^4$ , X is  $CR^7R^8$ , Y is  $CR^{10}R^{11}$ , Z is C, and ----- is a single bond, wherein the compound may be represented by the structure of Formula I-H:



[00169] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^7$ ,  $R^8$ ,  $R^{10}$ , and  $R^{11}$  are as defined for Formula I and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

[00170] In one embodiment of Formula I-H, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-H, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-H, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-H, Ring A is:




[00171] wherein the bond labeled "a" indicates the point of attachment to Y.

[00172] In one embodiment of Formula I-H,  $R^x$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>.

[00173] In one embodiment of Formula I-H,  $R^7$  and  $R^8$  are hydrogen.


[00174] In one embodiment of Formula I-H,  $R^y$  is hydrogen.

[00175] In one embodiment, W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is CR<sup>10</sup>R<sup>11</sup>, Z is C, and ----- is a single bond, wherein the compound may be represented by the structure of Formula I-I:



[00176] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^9$ ,  $R^{10}$ , and  $R^{11}$  are as defined for Formula I and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH- wherein R<sup>c</sup> is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

[00177] In one embodiment of Formula I-I, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-I, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-I, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-I, Ring A is:



[00178] wherein the bond labeled "a" indicates the point of attachment to Y.


[00179] In one embodiment of Formula I-I,  $R^x$  is C1-C6 alkyl.

**[00180]** In one embodiment of Formula I-I,  $R^9$  is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, C3-C6 cycloalkyl,  $Ar^1$ ,  $hetAr^1$ , (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-,  $H_2NC(=O)-$ ,  $Ar^1CH_2C(=O)-$ , (C3-C6 cycloalkyl)C(=O)- or (C1-C6 alkyl)SO<sub>2</sub>-, wherein  $Ar^1$  and  $hetAr^1$  are as defined for Formula I.

**[00181]** In one embodiment of Formula I-I,  $R^{10}$  and  $R^{11}$  are hydrogen.


**[00182]** In one embodiment of Formula I-I,  $R^y$  is hydrogen.

**[00183]** In one embodiment, W is  $CR^3R^4$ , X is O, Y is  $CR^{10}R^{11}$ , Z is C, and ----- is a single bond, wherein the compound may be represented by the structure of Formula I-J:



**[00184]** wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^{10}$ , and  $R^{11}$  are as defined for Formula I, and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

**[00185]** In one embodiment of Formula I-J, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-J, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-J, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-J, Ring A is:




**[00186]** wherein the bond labeled "a" indicates the point of attachment to Y.

[00187] In one embodiment of Formula I-J,  $R^x$  is C1-C6 alkyl.

[00188] In one embodiment of Formula I-J,  $R^{10}$  and  $R^{11}$  are hydrogen.


[00189] In one embodiment of Formula I-J,  $R^y$  is hydrogen.

[00190] In one embodiment,  $\text{-----}$  is a double bond, W is  $CR^3R^4$ , X is  $CR^5$ , Y is  $CR^6$ ,  $R^5$  and  $R^6$  are hydrogen, and Z is C, wherein the compound may be represented by the structure of Formula I-K:



[00191] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$  are as defined for Formula I and Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)-$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$  wherein  $R^c$  is as defined for Formula I or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl.

[00192] In one embodiment of Formula I-K, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment of Formula I-K, Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-K, Ring A is an isoxazolyl ring, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl. In one embodiment, Ring A is an isoxazolyl ring optionally substituted with C3-C6 cycloalkyl. In one embodiment of Formula I-K, Ring A is:




[00193] wherein the bond labeled "a" indicates the point of attachment to Y.

[00194] In one embodiment of Formula I-K,  $R^x$  is C1-C6 alkyl.

[00195] In one embodiment of Formula I-K,  $R^y$  is hydrogen.

[00196] In one embodiment, W is  $CR^3R^4$ , X is  $CR^7R^8$ , Y is  $CR^{10}R^{11}$ , Z is N, and ----- is a single bond, wherein the compound may be represented by the structure of Formula I-L:



[00197] wherein  $R^x$ ,  $R^y$ ,  $R^3$ ,  $R^4$ ,  $R^7$ ,  $R^8$ ,  $R^{10}$ , and  $R^{11}$  are as defined for Formula I and Ring A is (ii) a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or (iii) a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^dNHC(=O)-$ , (C1-C6 alkoxy) $C(=O)-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $C(=O)NH-$ .

[00198] In one embodiment of Formula I-L, Ring A is a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^dNHC(=O)-$ , (C1-C6 alkoxy) $C(=O)-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $C(=O)NH-$ .

[00199] In one embodiment of Formula I-L, Ring A is a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with  $R^dNHC(=O)-$ , (C1-C6 alkoxy) $C(=O)-$ , halogen, or (C1-C6 alkyl) $C(=O)NH-$ .

[00200] In one embodiment of Formula I-L,  $R^7$  and  $R^8$  are hydrogen.

[00201] In one embodiment of Formula I-L,  $R^{10}$  and  $R^{11}$  are hydrogen.

[00202] In one embodiment of Formula I-L,  $R^x$  is C1-C6 alkyl.

[00203] In one embodiment of Formula I-L,  $R^y$  is hydrogen.

[00204] In one embodiment of Formula I-L,  $R^y$  is halogen.

[00205] The compounds of Formula I include pharmaceutically acceptable salts thereof. In addition, the compounds of Formula I also include other salts of such compounds which are not necessarily pharmaceutically acceptable salts, and which may be useful as intermediates for preparing and/or purifying compounds of Formula I and/or for separating enantiomers of compounds of Formula I. Non-limiting examples of pharmaceutically acceptable salts of compounds of Formula I include trifluoroacetic acid and hydrochloride salts.




**[00206]** It will further be appreciated that the compounds of Formula I or their salts may be isolated in the form of solvates, and accordingly that any such solvate is included within the scope of the present invention. For example, compounds of Formula I and salts thereof can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like.

**[00207]** In one embodiment, the compounds of Formula I include the compounds of Examples 1-151 and stereoisomers and pharmaceutically acceptable salts and solvates thereof. In one embodiment, the compounds of Examples 1-151 are in the free base form. In one embodiment, the compounds of Examples 1-151 are in the salt form. In one embodiment, the compounds of Examples 1-151 are trifluoroacetate salts. In one embodiment, the compounds of Examples 1-151 are in the form of hydrochloride salts.

**[00208]** The term "pharmaceutically acceptable" indicates that the compound, or salt or composition thereof is compatible chemically and/or toxicologically with the other ingredients comprising a formulation and/or the patient being treated therewith.


**[00209]** Compounds provided herein may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. That is, an atom, in particular when mentioned in relation to a compound according to Formula I, comprises all isotopes and isotopic mixtures of that atom, either naturally occurring or synthetically produced, either with natural abundance or in an isotopically enriched form. For example, when hydrogen is mentioned, it is understood to refer to  $^1\text{H}$ ,  $^2\text{H}$ ,  $^3\text{H}$  or mixtures thereof; when carbon is mentioned, it is understood to refer to  $^{11}\text{C}$ ,  $^{12}\text{C}$ ,  $^{13}\text{C}$ ,  $^{14}\text{C}$  or mixtures thereof; when nitrogen is mentioned, it is understood to refer to  $^{13}\text{N}$ ,  $^{14}\text{N}$ ,  $^{15}\text{N}$  or mixtures thereof; when oxygen is mentioned, it is understood to refer to  $^{14}\text{O}$ ,  $^{15}\text{O}$ ,  $^{16}\text{O}$ ,  $^{17}\text{O}$ ,  $^{18}\text{O}$  or mixtures thereof; and when fluoro is mentioned, it is understood to refer to  $^{18}\text{F}$ ,  $^{19}\text{F}$  or mixtures thereof. The compounds provided herein therefore also comprise compounds with one or more isotopes of one or more atoms, and mixtures thereof, including radioactive compounds, wherein one or more non-radioactive atoms has been replaced by one of its radioactive enriched isotopes. Radiolabeled compounds are useful as therapeutic agents, e.g., cancer therapeutic agents, research reagents, e.g., assay reagents, and diagnostic agents, e.g., in vivo imaging agents. All isotopic variations of the compounds provided herein, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

**[00210]** For illustrative purposes, Schemes 1-25 show general methods for preparing the compounds provided herein as well as key intermediates. For a more detailed description of the individual reaction steps, see the Examples section below. Those skilled in the art will appreciate that other synthetic routes may be used to synthesize the inventive compounds. Although specific starting materials and reagents are depicted in the Schemes and discussed below, other starting materials and reagents can be easily substituted to provide a variety of derivatives and/or reaction conditions. In addition, many of the compounds prepared by the methods described below can be further modified in light of this disclosure using conventional chemistry well known to those skilled in the art.




Scheme 1

[00211] Scheme 1 shows a process for preparing compounds of Formula I-C (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is C(=O), Z is C, and ----- is a single bond, R<sup>9</sup> is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl-, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




**[00212]** wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>2</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00213]** Compound 1, which is commercially available, may be reacted with a reagent having the formula R<sup>x</sup>-X wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and X is a leaving group such as a halogen, to provide compound 2. Compound 2 may be reacted with hydroxylamine hydrochloride to provide oxime compound 3. The oxime moiety of compound 3 may undergo cycloaddition to form an isoxazole ring upon treatment with a compound having the formula HC≡C-R<sup>2</sup> wherein R<sup>2</sup> is C1-C6 alkyl or C3-C6 cycloalkyl to provide compound 4. Compound 4 may be treated with LDA and ethyl formate to provide the carbaldehyde-substituted compound 5. The carbaldehyde group of compound 5 may be reduced upon treatment of compound 5 with a reducing agent such as sodium borohydride to provide compound 6. The isoxazole ring of compound 6 may be iodinated upon treatment with N-iodosuccinimide in the presence of a catalytic amount of TFA to provide compound 7. The alcohol group of compound 7 may be protected with a suitable hydroxy protecting group P<sup>1</sup> such as a t-butyldimethyl silyl group to provide compound 8. The iodo group of compound 8 may be replaced with a carboxylate group upon treatment, for example, with a reagent of formula (R''O)<sub>2</sub>CO where R'' is C1-C6 alkyl in the presence of an organometallic reagent such as n-butyl lithium to provide compound 9. The hydroxy protecting group P<sup>1</sup> may be removed under standard conditions, for example in the presence of TBAF (tetrabutylammonium fluoride), to provide compound 10. The alcohol group of compound 10 may be displaced by an azide group upon treatment with diphenylphosphoryl azide to provide compound 11. The azide group of compound 11 may be reduced upon treatment with triphenylphosphine to provide compound 12. Reaction of compound 12 with ammonium hydroxide results in the displacement of the chloro group with an amino group as well as an intramolecular cyclization to provide compound 13, which is a compound of Formula I-C wherein R<sup>9</sup> is hydrogen. Compound 13 may be reacted with a reagent having the formula R<sup>9</sup>-X wherein R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group (e.g., t-butyldimethyl silyl), (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl- and X is a leaving group such as a halogen, to provide compound 14 which is a compound of Formula I-C, following removal of protecting groups where applicable, wherein R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl- or (hetAr<sup>1</sup>)C1-C6 alkyl-.




Scheme 2

[00214] Scheme 2 shows a process for preparing compounds of Formula I-C (i.e., a compound of Formula I wherein  $W$  is  $CR^3R^4$ ,  $X$  is  $NR^9$ ,  $Y$  is  $C(=O)$ ,  $Z$  is  $C$ , and  $-----$  is a single bond), wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl-, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, ( $Ar^1$ )C1-C6 alkyl-, or (het $Ar^1$ )C1-C6 alkyl-,  $R^x$  is as defined for Formula I, and Ring A is




[00215] wherein the bond labeled "a" indicates the point of attachment to  $Y$  and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl.

[00216] Compound 9 (wherein  $R^x$ ,  $P^1$  and  $R^Z$  are as defined in Scheme 1), which may be prepared according to Scheme 1, may be reacted with an amine reagent having the formula  $P^2NH_2$  wherein  $P^2$  is a suitable amino protecting group such as DMB (2,4-dimethoxybenzyl) to provide compound 15. The hydroxy protecting group  $P^1$  of compound 15 may be removed under suitable reaction conditions such as treatment with TBAF to provide compound 16. The alcohol group of compound 16 may be converted to a mesylate leaving group upon treatment with methanesulfonyl chloride to provide compound 17. Compound 17 may be undergo mesylate displacement and cyclization upon treatment with a reagent having the formula  $R^9NH_2$  wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl-, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, ( $Ar^1$ )C1-C6 alkyl-, or (het $Ar^1$ )C1-C6 alkyl- to provide compound 18. The amino protecting group  $P^2$  may be removed under standard conditions, for example in the presence of TFA, to provide compound 19, which is a compound of Formula I-C wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl-, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, ( $Ar^1$ )C1-C6 alkyl-, or (het $Ar^1$ )C1-C6 alkyl-.


**Scheme 3**

**[00217]** Scheme 3 shows a process for preparing a compound of Formula **I-D** (i.e., a compound of Formula **I** wherein W is CR<sup>3</sup>R<sup>4</sup>, X is O, Y is C(=O), Z is C, and ----- is a single bond) wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




**[00218]** wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00219]** Compound 15, (wherein R<sup>x</sup>, R<sup>z</sup>, R'', P<sup>1</sup> and P<sup>2</sup> are as defined in Scheme 2) which may be prepared according to Scheme 2, may undergo an intramolecular cyclization upon treatment with TBAF to provide compound 20. The amino protecting group P<sup>2</sup> may be removed under standard conditions, for example in the presence of TFA, to provide compound 21, which is a compound of Formula **I-D**.



Scheme 4


**[00220]** Scheme 4 shows a process for preparing a compound of Formula I-E (i.e., a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is SO<sub>2</sub>, Z is C, and — is a single bond) wherein R<sup>9</sup> is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is



**[00221]** wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.


**[00222]** Compound 8 (wherein R<sup>x</sup>, R<sup>z</sup> and P<sup>1</sup> are as defined in Scheme 1), which may be prepared according to Scheme 1, may be reacted with Burgess Reagent (methyl N-(triethylammoniumsulfonyl)carbamate) to provide compound 22. The alcohol group of compound 22 may be converted to a mesylate leaving group upon treatment with methanesulfonyl chloride to provide compound 22a. Compound 22a may undergo mesylate displacement and intramolecular cyclization upon treatment with ammonium hydroxide to provide compound 23, which is a compound of Formula I-E wherein R<sup>9</sup> is hydrogen. Compound 23 may be reacted with a reagent having the formula R<sup>9</sup>-X wherein R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group (e.g., tert-butyldimethylsilyl), (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl- and X is a leaving group such as a halogen, to provide compound 24 which is a compound of Formula I-E, following

removal of protecting groups where applicable, wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl- or (hetAr<sup>1</sup>)C1-C6 alkyl-.



**Scheme 5**


[00223] Scheme 5 shows a process for preparing a compound of Formula I-F (i.e., a compound of Formula I wherein  $W$  is  $\text{CR}^3\text{R}^4$ ,  $X$  is  $\text{CR}^7\text{R}^8$ ,  $Y$  is  $\text{SO}_2$ ,  $Z$  is  $\text{C}$ , and  $\text{-----}$  is a single bond), wherein  $R^7$  and  $R^8$  are hydrogen,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is



[00224] wherein the bond labeled "a" indicates the point of attachment to  $Y$  and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl.


[00225] Compound 8 (wherein  $R^x$ ,  $R^z$  and  $P^1$  are as defined in Scheme 1), which may be prepared according to Scheme 1, may be reacted with dimethyl disulfide in the presence of an organometallic reagent (e.g., *n*-butyl lithium). Oxidation of the intermediate sulfide to a sulfone upon treatment with mCPBA (meta-chloroperoxybenzoic acid) can provide compound 25. The hydroxy protecting group  $P^1$  of

compound 25 may be removed under standard conditions to provide the intermediate alcohol (not shown), which may be converted to the bromide upon treatment with  $\text{PBr}_3$  to provide compound 26. Compound 26 may undergo an intramolecular cyclization upon treatment with a strong non-nucleophilic base such as LHMDS (lithium hexamethyldisilazide) to provide compound 27. The chloro group of compound 27 may be displaced by an amino group upon treatment of compound 27 with ammonium hydroxide to provide compound 28, which is a compound of Formula I-F.



Scheme 6

[00226] Scheme 6 shows a process for preparing a compound of Formula I-G (i.e., a compound of Formula I wherein W is  $\text{CR}^3\text{R}^4$ , X is  $\text{CR}^7\text{R}^8$ , Y is  $\text{C}(=\text{O})$ , Z is C, and — is a single bond), wherein  $\text{R}^7$  and  $\text{R}^8$  are hydrogen,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




[00227] wherein the bond labeled "a" indicates the point of attachment to Y and  $\text{R}^z$  is C1-C6 alkyl or C3-C6 cycloalkyl.


[00228] Compound 5 (wherein  $\text{R}^x$  and  $\text{R}^z$  are as defined for Scheme 1), which may be prepared according to Scheme 1, may be converted to compound 29 upon treatment with benzyl



(triphenylphosphoranylidene)acetate. The double bond of compound 29 may be reduced under standard hydrogenation conditions (e.g., palladium catalyzed hydrogenation conditions) to provide compound 30 with concomitant cleavage of the benzyl group. The chloro group of compound 30 may be displaced by an amino group upon treatment of compound 30 with ammonium hydroxide to provide compound 31. Compound 31 may be cyclized upon treatment with polyphosphoric acid to provide compound 32, which is a compound of Formula I-G.



[00229] Scheme 7 shows processes for preparing compounds of Formula I-H (i.e., a compound of Formula I wherein W is  $CR^3R^4$ , X is  $CR^7R^8$ , Y is  $CR^{10}R^{11}$ , Z is C, and ----- is a single bond), wherein  $R^7$  and  $R^8$  are hydrogen,  $R^x$  is as defined for Formula I,  $R^{10}$  and  $R^{11}$  are as defined below for Scheme 7, and Ring A is




[00230] wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl.

[00231] Compound 32 (wherein  $R^x$  and  $R^Z$  are as defined in Scheme 6), which may be prepared according to Scheme 6, may be reacted with a Grignard reagent having the formula  $R^{11}MgX$  wherein  $R^{11}$  is C1-C6 alkyl and X is Br, I or Cl to provide compound 33, which is a compound of Formula I-H wherein  $R^{10}$  is hydroxy and  $R^{11}$  is C1-C6 alkyl.

[00232] Alternatively, compound 32 may be reacted with  $TMSCF_3$  (trifluoromethyltrimethylsilane) to provide compound 34, which is a compound of Formula I-H wherein  $R^{10}$  is hydroxy and  $R^{11}$  is  $CF_3$ .


[00233] Alternatively, compound 32 may be reacted with a reagent having the formula  $R^aR^bNH$  wherein  $R^a$  is hydrogen and  $R^b$  is H or C1-C6 alkyl in the presence of a reductant (e.g., sodium triacetoxyborohydride) to provide a compound of Formula I-H wherein  $R^{10}$  is  $R^aR^bN$ - where  $R^a$  is hydrogen and  $R^b$  is H or C1-C6 alkyl, and  $R^{11}$  is hydrogen.

[00234] Alternatively, compound 32 may be treated under standard hydrogenation conditions for reducing a ketone to an alcohol (e.g., using sodium borohydride) to provide compound 32a, which is a compound of Formula I-H wherein  $R^{10}$  is OH and  $R^{11}$  is hydrogen. Compound 32a may be reacted with a reagent having the formula (C1-C6 alkyl)-X wherein X is a halogen to provide compound 32b, which is a compound of Formula I-H wherein  $R^{10}$  is C1-C6 alkoxy and  $R^{11}$  is hydrogen.




Scheme 8

**[00235]** Scheme 8 shows a process for preparing compounds of Formula I-K (i.e., compounds of Formula I wherein W is  $CR^3R^4$ , X is  $CR^5$ , Y is  $CR^6$ , Z is C, and ----- is a double bond), wherein  $R^5$  and  $R^6$  are hydrogen,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




**[00236]** wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00237]** Compound 32a (wherein  $R^x$  and  $R^Z$  are as defined in Scheme 6), which may be prepared according to Scheme 7, may be protected with an amino protecting group (e.g., dimethylformamide dimethyl acetal) to provide compound 37 where P is an amino protecting group. Treatment of compound 37 with iodomethane in the presence of a base (e.g., diisopropylethylamine), followed by removal of the amino protecting group under standard conditions provides compound 38, which is a compound of Formula I-K.



[00238] Scheme 9 shows a process for preparing compounds of Formula I-I (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is CR<sup>10</sup>R<sup>11</sup>, Z is C, and — is a single bond), wherein R<sup>10</sup> and R<sup>11</sup> are hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, R<sup>9</sup> is as defined below for Scheme 9, and Ring A is




[00239] wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.

[00240] Compound 3 (wherein R<sup>x</sup> is as defined for Scheme 1), which may be prepared according to Scheme 1, may undergo cycloaddition to form an isoxazole ring upon treatment with a compound

having the formula  $R^2C\equiv C-COOEt$  wherein  $R^2$  is C1-C6 alkyl or C3-C6 cycloalkyl to provide compound 39. Compound 39 may be treated with a suitable ester-reducing agent such as DIBAL-H to provide compound 40. The alcohol group of compound 40 may be protected with a suitable hydroxy protecting group  $P^3$ , for example, a tert-butyldimethylsilyl protecting group upon treatment with TBSCl, to provide compound 41. Compound 41 may be treated with LDA and ethyl formate to provide the carbaldehyde-substituted compound 42. Compound 42 may be reacted with an amine of the formula  $(P^4)_2N$  where  $P^4$  is a suitable amino protecting group (e.g., a benzyl protecting group such as 4-methoxybenzyl) to provide compound 43. The hydroxy protecting group  $P^3$  of compounds 43 may be removed under standard conditions to provide compound 44. The alcohol group of compound 44 may be oxidized to a carbaldehyde under suitable oxidation conditions (e.g., by treatment with Dess-Martin periodinane), to provide compound 45. Compound 45 may be treated with a reagent having the formula  $R^9NH_2$  where  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup> or (hetAr<sup>1</sup>)C1-C6 alkyl-, in the presence of a reducing agent (e.g., sodium triacetoxyborohydride) to provide compound 46. Removal of the amino protecting groups  $P^4$  of compound 46 provides compound 47, which is a compound of Formula I-I wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup> or (hetAr<sup>1</sup>)C1-C6 alkyl-.


**[00241]** Alternatively, compound 45 may be cyclized upon treatment with ammonium acetate in the presence of a reducing agent (e.g., sodium triacetoxyborohydride) to provide compound 48. Compound 48 may be treated with (C1-C6 alkyl)SO<sub>2</sub>Cl, which after removal of the amino protecting groups  $P^4$  provides compound 49a, which is a compound of Formula I-I wherein  $R^9$  is (C1-C6 alkyl)SO<sub>2</sub>-.

**[00242]** Alternatively, compound 48 may be treated with a reagent having the formula  $R^{9'}C(=O)Cl$ , wherein  $R^{9'}$  is C1-C6 alkyl or fluoroC1-C6 alkyl, to provide compound 49b, which is a compound of Formula I-I wherein  $R^9$  is C1-C6 alkyl or fluoroC1-C6 alkyl.




**Scheme 10**

**[00243]** Scheme 10 shows a process for preparing compounds of Formula I-J (i.e., compounds of Formula I wherein W is  $\text{CR}^3\text{R}^4$ , X is O, Y is  $\text{CR}^{10}\text{R}^{11}$ , Z is C, and  $\text{-----}$  is a single bond), wherein  $\text{R}^{10}$  and  $\text{R}^{11}$  are hydrogen,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $\text{Cyc}^2$ , or  $\text{Cyc}^3$ , and Ring A is



**[00244]** wherein the bond labeled "a" indicates the point of attachment to Y and  $\text{R}^z$  is C1-C6 alkyl or C3-C6 cycloalkyl.


**[00245]** Compound 43 (wherein  $\text{R}^z$ ,  $\text{P}^3$ ,  $\text{P}^4$  and  $\text{R}^x$  are as defined for Scheme 9), which may be prepared according to Scheme 9, may undergo an intramolecular cyclization upon treatment with trimethylsilyl trifluoromethanesulfonate ( $\text{CF}_3\text{SO}_3\text{TMS}$ ), followed by treatment with triethylsilane, to provide compound 50, which is a compound of Formula I-J.



Scheme 11

[00246] Scheme 11 shows a process for preparing compounds of Formula I-C (i.e. compounds of Formula I wherein W is  $CR^3R^4$ , X is  $NR^9$ , Y is  $C(=O)$ , Z is C, and  $-----$  is a single bond), wherein Ring A is a 6-membered aryl ring optionally substituted with one or more substituents independently selected from halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^cNHC(=O)$ ,  $R^cC(=O)NH-$ , or  $R^cNHC(=O)NH-$ , wherein  $R^c$  is C3-C6 cycloalkyl or (C3-C6 cycloalkyl) $CH_2-$ ,  $R^9$  is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-, and  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>. Compound 51, which is commercially available, may be reacted with an alcohol having the formula  $R^xOH$ , wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, under standard Mitsunobu reaction conditions, to provide compound 52. The ester group of compound 52 may be reduced under standard conditions (e.g., by treating with a reducing agent such as DIBAL-H) to provide compound 53. Compound 53 may be iodinated upon treatment with N-iodosuccinimide to provide compound 54. Compound 54 may be treated with a suitable alcohol protecting reagent to provide compound 55 where  $P^5$  is a hydroxy protecting group (e.g., t-butyldimethylsilyl). Compound 55 may be treated with a suitably protected


amine to provide compound 56 where P<sup>6</sup> is an amino protecting group (e.g., a benzyl group, e.g., 2,4-dimethoxybenzyl). Compound 56 may be treated with a boronic ester reagent (1) wherein n is 0, 1, 2, 3 or 4, each R<sup>2</sup> is independently selected from halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH-, wherein R<sup>c</sup> is C3-C6 cycloalkyl or (C3-C6 cycloalkyl)CH<sub>2</sub>-, R' is C1-C6 alkyl, and each R' is independently H or (1-6C)alkyl, or each R' together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl), using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions, to provide compound 57. The hydroxy protecting group of compound 57 may be removed under standard conditions to provide compound 58. The alcohol group of compound 58 may be displaced by an azide group upon treatment with diphenylphosphoryl azide to provide compound 59. The azide group of compound 59 may be reduced upon treatment with triphenylphosphine to provide compound 60. Compound 60 may undergo an intramolecular cyclization upon heating at elevated temperatures to provide compound 61a, which is a of Formula I-C wherein R<sup>9</sup> is hydrogen. Compound 61a may be reacted with a reagent having the formula R<sup>9</sup>-X wherein R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group (e.g., tert-butylidimethylsilyl), (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl- and X is a leaving group such as a halogen, to provide compound 61b which is a compound of Formula I-C, following removal of protecting groups where applicable, wherein R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl- or (hetAr<sup>1</sup>)C1-C6 alkyl-.



Scheme 12




[00247] Scheme 12 shows a process for preparing compounds of Formula I-A (i.e., compounds of Formula I wherein W is  $\text{CR}^3\text{R}^4$ , X is  $\text{CR}^7\text{R}^8$ , Y is  $\text{C}(=\text{O})$ , Z is N, and ----- is a single bond) wherein  $\text{R}^7$  and  $\text{R}^8$  are hydrogen,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




[00248] wherein the bond labeled "a" indicates the point of attachment to Y, n is 0, 1, 2, 3 or 4, and each  $\text{R}^2$  is independently selected from  $\text{R}^d\text{NHC}(=\text{O})-$ , (C1-C6 alkoxy) $\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $\text{C}(=\text{O})\text{NH}-$  wherein  $\text{R}^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup> $\text{CH}_2-$ , phenyl or hetAr<sup>2</sup>.

[00249] Compound 56 (wherein  $\text{R}^x$ ,  $\text{P}^5$  and  $\text{P}^6$  are as defined in Scheme 10), which may be prepared according to Scheme 10, may be reacted with a compound of formula (2) wherein n is 0, 1, 2, 3 or 4, each  $\text{R}^2$  is independently selected from  $\text{R}^d\text{NHC}(=\text{O})-$ , (C1-C6 alkoxy) $\text{C}(=\text{O})-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $\text{C}(=\text{O})\text{NH}-$  wherein  $\text{R}^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup> $\text{CH}_2-$ , phenyl or hetAr<sup>2</sup>, and each  $\text{R}^1$  is independently H or (1-6C)alkyl, or each  $\text{R}^1$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl) using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions, to provide compound 62. The hydroxy protecting group  $\text{P}^5$  may be removed under standard conditions to provide compound 63. The hydroxy group of compound 63 may be oxidized to a carbaldehyde under suitable oxidation conditions (e.g., by treatment with Dess-Martin periodinane), to provide compound 64. Compound 64 may undergo a Wittig olefination upon treatment with ethyl 2-(triphenyl- $\lambda^5$ -phosphanylidene)acetate to provide compound 65. The double bond of compound 65 may be reduced under standard conditions (e.g., copper-catalyzed hydrogenation reaction conditions) to provide compound 66. Compound 66 may be converted to compound 67 under standard ester hydrolysis conditions. Compound 67 may undergo an intramolecular cyclization upon treatment with di-tert-butyl dicarbonate ( $\text{Boc}_2\text{O}$ ) to provide compound 68. Removal of the amino protecting group  $\text{P}^6$  provides compound 69, which is a compound of Formula I-A.




**[00250]** Scheme 13 shows a process for preparing compounds of Formula I-C (i.e., a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is C(=O), Z is C, and ----- is a single bond), wherein R<sup>9</sup> is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)-, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is



**[00251]** wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>2</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00252]** Compound 4 (wherein R<sup>x</sup>, R<sup>2</sup>, and R<sup>9</sup> are as defined in Scheme 1), which may be prepared according to Scheme 1, may be methylated upon treatment with methyl iodide in the presence of a strong base such as n-butyl lithium to provide compound 70. Compound 70 may be iodinated upon treatment with N-iodosuccinimide in the presence of a catalytic amount of TFA to provide compound 71. Compound 71 may be converted to compound 72 upon treatment with ethyl chloroformate and isopropylmagnesium(II) lithium chloride. Compound 72 may be treated with N-bromosuccinimide in the presence of AIBN to provide a brominated intermediate (not shown); subsequent treatment with ammonium hydroxide results in an intramolecular cyclization to provide 13, which is a compound Formula I-C wherein R<sup>9</sup> is hydrogen. Compound 13 may be reacted with a reagent having the formula

$R^9$ -X wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group (e.g., tert-butyldimethylsilyl), (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)- and X is a leaving group such as a halogen, to provide compound 14 which is a compound of Formula I-C wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)-, after removal of protecting groups where applicable.




Scheme 14

[00253] Scheme 14 shows processes for preparing compounds of Formula I-C (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is C(=O), Z is C, and ----- is a single bond), wherein R<sup>9</sup> is hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A are as defined for Formula I.

[00254] Compound 56 (wherein R<sup>x</sup>, P<sup>5</sup> and P<sup>6</sup> are as defined in Scheme 11), which may be prepared according to Scheme 11, may be reacted with 4,4,5,5-tetramethyl-1,3,2-dioxaborolane in the

presence of triphenylphosphine to provide compound 73. Compound 73 may be coupled to a reagent having formula (3) wherein Ring A is as defined for Formula I, X is a leaving group such as halogen, and R is C1-C6 alkyl, using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions, to provide compound 74. Alternatively, compound 74 may be prepared by coupling compound 56 with a boronic acid or boronic ester reagent (4) wherein Ring A is as defined for Formula I, R is C1-C6 alkyl, and each R' is independently H or (1-6C)alkyl, or each R' together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl), using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions. The hydroxy protecting group of compound 74 may be removed under standard conditions to provide compound 75. The alcohol group of compound 75 may be displaced by an azide group upon treatment with diphenylphosphoryl azide to provide compound 76. The azide group of compound 76 may be reduced upon treatment with triphenylphosphine to provide compound 76a. Reaction of with ammonium hydroxide results an intramolecular cyclization of compound 76a, which after removal of the amino protecting group provides compound 77, which is a compound of Formula I-C wherein R<sup>9</sup> is hydrogen.




Scheme 15

**[00255]** Scheme 15 shows a process for preparing compounds of Formula I-B (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is C(=O), Z is N, and ----- is a single bond) wherein R<sup>9</sup> is hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is a monocyclic heteroaryl ring or a bicyclic heteroaryl ring as defined for Formula I.

**[00256]** Compound 56 (wherein R<sup>x</sup>, P<sup>5</sup> and P<sup>6</sup> are as defined for Scheme 11), which may be prepared according to Scheme 11, may be treated with a boronic ester reagent (5) wherein Ring A is a

monocyclic heteroaryl ring or a bicyclic heteroaryl ring as defined for Formula I, and each R<sup>1</sup> is independently H or (1-6C)alkyl, or each R<sup>1</sup> together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl), using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions, to provide compound 78. The hydroxy protecting group P<sup>5</sup> of compound 78 may be removed under standard conditions to provide compound 79. The alcohol group of compound 79 may be displaced by an azide group upon treatment with diphenylphosphoryl azide to provide compound 80. The azide group of compound 80 may be reduced upon treatment with triphenylphosphine to provide compound 81. Ring closure may be effected upon treatment of compound 89 with di-tert-butyl dicarbonate in the presence of DMAP, after which removal of the amino protecting group P<sup>6</sup> provides compound 82, which is a compound of Formula I-B wherein R<sup>9</sup> is hydrogen.




Scheme 16

[00257] Scheme 16 shows a process for preparing compounds of Formula I-A (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is CR<sup>7</sup>R<sup>8</sup>, Y is C(=O), Z is N, and — is a single bond) wherein R<sup>7</sup> and R<sup>8</sup> are hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is a monocyclic heteroaryl ring or a bicyclic heteroaryl ring as defined for Formula I.


[00258] Compound 79 (wherein Ring A, R<sup>x</sup> and P<sup>6</sup> are as defined in Scheme 15), which may be prepared according to Scheme 15, may be oxidized to a carbonyl compound under suitable oxidation conditions (e.g., by treatment with Dess-Martin periodinane), to provide compound 83. Compound 83 may undergo a Wittig olefination upon treatment with ethyl 2-(triphenyl-λ<sup>5</sup>-phosphanylidene)acetate to

provide compound 84. The double bond of compound 84 may be reduced under standard conditions (e.g., using copper-catalyzed hydrogenation reaction conditions) to provide compound 85. Compound 85 may be converted to compound 86 under standard ester hydrolysis conditions. Compound 86 may undergo an intramolecular cyclization upon treatment with di-tert-butyl dicarbonate to provide compound 87, which is a compound of Formula I-A, after removal of the amino protecting group.




Scheme 17

**[00259]** Scheme 17 shows a process for preparing compounds of Formula **I-B** (i.e., compounds of Formula **I** wherein  $W$  is  $CR^3R^4$ ,  $X$  is  $NR^9$ ,  $Y$  is  $C(=O)$ ,  $Z$  is  $N$ , and  $-----$  is a single bond) wherein  $R^9$  is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




**[00260]** wherein the bond labeled "a" indicates the point of attachment to  $Y$ ,  $n$  is 0, 1, 2, 3 or 4, and each  $R^Z$  is independently selected from  $R^dNHC(=O)-$ , (C1-C6 alkoxy) $C(=O)-$ , C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl) $C(=O)NH-$ .

**[00261]** Compound 63 (wherein  $R^x$ ,  $P^6$ ,  $R^Z$  and  $n$  are as defined for Scheme 12), which may be prepared according to Scheme 12, may be treated with diphenylphosphoryl azide to provide compound 88. The azide group of compound 88 may be reduced upon treatment with triphenylphosphine to provide compound 89. Ring closure may be effected upon treatment of compound 89 with di-tert-butyl dicarbonate in the presence of DMAP to afford 90a, after which removal of the amino protecting group under standard conditions provides compound 90b, which is a compound of Formula **I-B** wherein  $R^9$  is hydrogen. Compound 90a may be treated with a reagent of formula  $R^x-X$ , wherein  $X$  is a leaving atom (e.g., halo) or leaving group (e.g., mesylate); and  $X$  is C1-C6 alkyl, to provide compound 90c, which is a compound of Formula **I-B** wherein  $R^9$  is C1-C6 alkyl, after removal of the amino protecting group.



Scheme 18

[00262] Scheme 18 shows processes for preparing compounds of Formula I-B (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>, Y is C(=O), Z is N, and ----- is a single bond), wherein R<sup>9</sup> is hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




[00263] wherein the bond labeled "a" indicates the point of attachment to Y, each R<sup>2</sup> is independently selected from C1-C6 alkyl and C3-C6 cycloalkyl, n is 0, 1 or 2, and ring A is further optionally substituted with one occurrence of halo.

[00264] Compound 56 (wherein R<sup>x</sup>, P<sup>5</sup>, and P<sup>6</sup> are as defined for Scheme 12), which may be prepared according to Scheme 12, may be reacted with 4,4,5,5-tetramethyl-1,3,2-dioxaborolane in the presence of triphenylphosphine to provide compound 73. Compound 73 may be coupled with a reagent having formula (6) wherein each R<sup>2</sup> is independently selected from halogen, C3-C6 cycloalkyl and C1-C6 alkoxy, and n is 0, 1 or 2, using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions, to provide compound 91. Alternatively, compound 91 may be prepared by coupling compound 56 with a boronic ester reagent (7) wherein each R<sup>2</sup> is independently selected from halogen, C1-C6 alkyl, C3-C6 cycloalkyl and C1-C6 alkoxy, n is 0, 1 or 2, and each R<sup>1</sup> is independently H or (1-6C)alkyl, or each R<sup>1</sup> together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl), using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction




conditions. Compound 91 may be converted to compound 92 using appropriate conditions to remove both the amino protecting group  $P^6$  and the hydroxy protecting group  $P^5$ . The alcohol group of compound 92 may be displaced by an azide group upon treatment with diphenylphosphoryl azide to provide compound 93. The azide group of compound 93 may be reduced upon treatment with triphenylphosphine to provide compound 94. Ring closure may be effected upon treatment of compound 94 with bis(4-nitrophenyl) carbonate to provide compound 96, which is a compound of Formula I-B wherein  $R^9$  is hydrogen.



Scheme 19


[00265] Scheme 19 shows a process for preparing compounds of Formula I-C (i.e., compounds of Formula I wherein  $W$  is  $CR^3R^4$ ,  $X$  is  $NR^9$ ,  $Y$  is  $C(=O)$ ,  $Z$  is  $C$ , and  $-----$  is a single bond), wherein  $R^9$  is hydrogen,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




[00266] wherein the bond labeled "a" indicates the point of attachment to  $Y$ , and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl.

[00267] Compound 73 (wherein  $R^x$ ,  $P^5$  and  $P^6$  are as defined for Scheme 18), which may be prepared according to Scheme 18), may be coupled with a reagent having the formula (8) wherein  $R^Z$  is on one of the ring nitrogen atoms and is selected from C1-C6 alkyl and C3-C6 cycloalkyl and  $R''$  is C1-C6 alkyl, using appropriate palladium-catalyzed cross-coupling reaction conditions, e.g., Suzuki coupling reaction conditions, to provide compound 96. The hydroxy protecting group of compound 96 may be removed under standard conditions to provide compound 97. The alcohol group of compound 97 may be displaced by an azide group upon treatment with diphenylphosphoryl azide to provide compound 98. The azide group of compound 98 may be reduced upon treatment with triphenylphosphine to provide compound 98a. Compound 98a can undergo an intramolecular cyclization upon treatment with a base

at elevated temperatures, after which removal of the amino protecting group under standard conditions provides compound 99, which is a compound of Formula I-C wherein  $R^9$  is hydrogen.



**[00268]** Scheme 20 shows a process for preparing compounds of Formula I-C (i.e., compounds of Formula I wherein  $W$  is  $CR^3R^4$ ,  $X$  is  $NR^9$ ,  $Y$  is  $C(=O)$ ,  $Z$  is  $C$ , and  $-----$  is a single bond), wherein  $R^9$  is hydrogen,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




**[00269]** wherein the bond labeled "a" indicates the point of attachment to  $Y$  and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00270]** The ring nitrogen atom of Compound 1, which is commercially available, may be protected by reacting compound 1 with a suitable reagent to provide compound 100 wherein PG is an

amino protecting group (e.g., SEM). Compound 100 may be reacted in a similar manner described for the conversion of compound 2 to compound 4 in Scheme 1 to provide compound 101 wherein R<sup>2</sup> is C1-C6 alkyl or C3-C6 cycloalkyl. Compound 101 may be methylated upon treatment with methyl iodide in the presence of a strong base such as n-butyl lithium to provide compound 102. Compound 102 may be iodinated upon treatment with N-iodosuccinimide in the presence of a catalytic amount of TFA to provide compound 103. Compound 103 may be converted to compound 104 upon treatment with ethyl chloroformate and isopropylmagnesium(II) lithium chloride. Compound 104 may be treated with N-bromosuccinimide in the presence of AIBN to provide the corresponding methyl bromide intermediate (not shown), followed by treatment with ammonium hydroxide which results in an intramolecular cyclization. Removal of the amino protecting group PG provides compound 13a, which is a compound Formula I-C wherein R<sup>9</sup> is hydrogen.


**[00271]** Alternatively, the amino protecting group PG of compound 104 may be removed under standard conditions to provide compound 105. Compound 105 may be reacted with a reagent having the formula R<sup>x</sup>-OH wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, using Mitsunobu reaction conditions to provide compound 106. Compound 106 may be cyclized upon treatment with N-bromosuccinimide in the presence of AIBN to provide compound 13.

**[00272]**



Scheme 21

[00273] Scheme 21 shows a process for preparing a compound of Formula I-G (i.e., a compound of Formula I wherein W is  $\text{CR}^3\text{R}^4$ , X is  $\text{CR}^7\text{R}^8$ , Y is  $\text{C}(=\text{O})$ , Z is C, and ----- is a single bond), wherein  $\text{R}^7$  and  $\text{R}^8$  are hydrogen,  $\text{R}^x$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, and Ring A is



[00274] wherein the bond labeled "a" indicates the point of attachment to Y, and  $\text{R}^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl.


[00275] Compound 1, which is commercially available, may be reacted with a reagent having the formula PG-X wherein PG is a suitable amino protecting group (e.g., SEM), and X is a leaving group (e.g.,

halo, e.g., Cl) to provide compound 2a. Compound 2a may be reacted with hydroxylamine hydrochloride to provide oxime compound 3a. The oxime moiety of compound 3a may undergo cycloaddition to form an isoxazole ring upon treatment with a compound having the formula  $\text{HC}\equiv\text{C}-\text{R}^2$  wherein  $\text{R}^2$  is C1-C6 alkyl or C3-C6 cycloalkyl to provide compound 4a. Compound 4a may be treated with n-butyl lithium and methyl iodide to provide the methyl-substituted compound 70a. The isoxazole ring of compound 70a may be iodinated upon treatment with N-iodosuccinimide in the presence of a catalytic amount of TFA to provide compound 71a. The iodo group in compound 71a may undergo metal halogen exchange with an organometallic reagent (e.g.,  $\text{iPrMgCl}_2\text{Li}$ ) followed by trapping with acetic anhydride to provide an acetyl substituted intermediate (not shown), whereupon treatment of this intermediate with N-bromosuccinimide and AIBN under radical bromination conditions can afford compound 107. Treatment of compound 107 with a strong non-nucleophilic base such as LHMDs (lithium hexamethyldisilazide) allows an intramolecular cyclization, thereby affording compound 108. Removal of the protecting group on 108 provides compound 109. Compound 109 may be treated with a reagent of formula  $\text{R}^x-\text{X}$  wherein X is a leaving group such as halogen, a reagent of formula  $\text{R}^x-\text{OH}$ , or a reagent with formula  $\text{R}^x\text{B}(\text{OR}')_2$  wherein each  $\text{R}'$  is independently H or (1-6C)alkyl, or each  $\text{R}'$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl) to provide compound 110, wherein in each of  $\text{R}^x-\text{X}$  and  $\text{R}^x\text{B}(\text{OR}')_2$ ,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>; and in  $\text{R}^x-\text{OH}$ ,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, or hetCyc<sup>3</sup>. As non-limiting examples of the foregoing, compound 109 may be reacted with  $\text{R}^x-\text{X}$  under transition-metal (e.g., palladium or copper) catalyzed cross-coupling conditions or under nucleophilic substitution conditions (e.g., base and/or heat); compound 109 may be reacted with  $\text{R}^x\text{B}(\text{OR}')_2$  under transition-metal (e.g., copper) catalyzed cross-coupling conditions; and compound 109 may be reacted with  $\text{R}^x-\text{OH}$  under Mitsunobu conditions. The chloro group in 110 may be displaced with ammonium hydroxide to provide compound 32, which is a compound of Formula I-G, wherein  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, and  $\text{R}^2$  is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00276]** Alternatively, compound 109 may be reacted with ammonium hydroxide to provide compound 111, which is a compound of Formula I-G wherein  $\text{R}^x$  is hydrogen, and  $\text{R}^2$  is C1-C6 alkyl or C3-C6 cycloalkyl. Compound 111 may be treated with a reagent of formula  $\text{R}^x-\text{X}$  wherein X is a leaving group such as halogen, a reagent of formula  $\text{R}^x-\text{OH}$ , or a reagent with formula  $\text{R}^x\text{B}(\text{OR}')_2$  wherein each  $\text{R}'$  is independently H or (1-6C)alkyl, or each  $\text{R}'$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl) to provide compound 32, which is a compound of Formula I-G, wherein  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, and  $\text{R}^2$  is C1-C6 alkyl or C3-C6 cycloalkyl. In each of  $\text{R}^x-\text{X}$  and  $\text{R}^x\text{B}(\text{OR}')_2$ ,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>; and in  $\text{R}^x-\text{OH}$ ,  $\text{R}^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, or hetCyc<sup>3</sup>. As non-limiting examples of the foregoing, compound 111 may be reacted with  $\text{R}^x-\text{X}$  under transition-metal (e.g., palladium or copper) catalyzed cross-coupling conditions or nucleophilic substitution conditions (e.g., base and/or heat); compound 111 may be reacted with  $\text{R}^x\text{B}(\text{OR}')_2$  under transition-metal (e.g., copper) catalyzed cross-coupling conditions; and compound 111 may be reacted with  $\text{R}^x-\text{OH}$  under Mitsunobu conditions.


**[00277]** As another alternative, compound 109 may be reacted with a reagent of formula  $\text{P}^6-\text{NH}_2$  wherein  $\text{P}^6$  is a suitable amino protecting group (e.g., a benzyl group, e.g., 2,4-dimethoxybenzyl, DMB) to

provide compound 112. Compound 112 may be treated with a reagent of formula  $R^x-X$  wherein  $X$  is a leaving group such as halogen, a reagent of formula  $R^x-OH$ , or a reagent with formula  $R^x-B(OR')_2$  wherein each  $R'$  is independently H or (1-6C)alkyl, or each  $R'$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl) to provide compound 113, wherein in each of  $R^x-X$ , and  $R^x-B(OR')_2$ ,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>; and in  $R^x-OH$ ,  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, or hetCyc<sup>3</sup>. As non-limiting examples of the foregoing, compound 112 may be reacted with  $R^x-X$  under transition-metal (e.g., palladium or copper) catalyzed cross-coupling conditions or nucleophilic substitution conditions (e.g., base and/or heat); compound 112 may be reacted with  $R^x-B(OR')_2$  under transition-metal (e.g., copper) catalyzed cross-coupling conditions; and compound 112 may be reacted with  $R^x-OH$  under Mitsunobu conditions. Removal of the amino protecting group on compound 113 affords compound 32, which is a compound of Formula I-G, wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl.



Scheme 22

[00278] Scheme 22 shows a process for preparing a compound of Formula I-H (i.e., a compound of Formula I wherein  $W$  is  $CR^3R^4$ ,  $X$  is  $CR^7R^8$ ,  $Y$  is  $CR^{10}R^{11}$ ,  $Z$  is C, and  $-----$  is a single bond), wherein  $R^7$  and  $R^8$  are hydrogen,  $R^x$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>,  $R^{10}$  and  $R^{11}$  are as defined below for Scheme 22, and Ring A is




**[00279]** wherein the bond labeled "a" indicates the point of attachment to Y, and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.

**[00280]** Ketone-containing compound 110 (wherein R<sup>x</sup> and R<sup>z</sup> are as defined for Scheme 21), which may be prepared according to Scheme 21, may be reduced under standard conditions to provide compound 110a, which may be reacted with ammonium hydroxide to afford compound 36, which is a compound of Formula **I-H**, wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, R<sup>10</sup> is hydroxyl, and R<sup>11</sup> is hydrogen.


**[00281]** Alternatively, compound 32 (wherein R<sup>x</sup> and R<sup>z</sup> are as defined for Scheme 21), which may be prepared according to Scheme 21, may be subjected under ketone reduction conditions (e.g., with sodium borohydride) to provide compound 36 which is a compound of Formula **I-H**, wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, R<sup>10</sup> is hydroxyl, and R<sup>11</sup> is hydrogen.

**[00282]** Alternatively, compound 111 (wherein R<sup>z</sup> is as defined for Scheme 21), which may be prepared according to Scheme 21, may be subjected under ketone reduction conditions (e.g., with sodium borohydride) to provide compound 36 which is a compound of Formula **I-H**, wherein R<sup>x</sup> is hydrogen, R<sup>10</sup> is hydroxyl, and R<sup>11</sup> is hydrogen.



Scheme 23

[00283] Scheme 23 shows a process for preparing a compound of Formula I-H (i.e., a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is CR<sup>7</sup>R<sup>8</sup>, Y is CR<sup>10</sup>R<sup>11</sup>, Z is C, and ----- is a single bond), wherein R<sup>7</sup> and R<sup>8</sup> are hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, R<sup>10</sup> and R<sup>11</sup> are as defined below for Scheme 23, and Ring A is




[00284] wherein the bond labeled "a" indicates the point of attachment to Y, and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl.

[00285] Compound 114 may be treated with a reagent of formula R<sup>x</sup>NH<sub>2</sub> wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup> to provide compound 115. Iodination of compound 115 (e.g., with N-iodosuccinimide, NIS) can afford compound 116, whereupon treatment of 116 with 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane e.g., under palladium catalyzed cross-coupling (e.g., Suzuki coupling) conditions can provide compound 117. Oxidative cleavage of the olefin moiety in 117 allows the formation of aldehyde compound 2 which can react with hydroxylamine hydrochloride to




provide oxime compound 3. Compound 3 may undergo cycloaddition with a reagent of formula  $R^z \equiv$  wherein  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl to provide compound 4. Exposure of compound 4 to the presence of a strong base (e.g., *n*-butyl lithium) and iodomethane can lead to compound 70 which can be iodinated at the isoxazole (e.g., with NIS) to afford compound 71. Compound 71 can undergo metal-halogen exchange with an organometallic reagent (e.g., *i*PrMgCl<sub>2</sub>Li, Ac<sub>2</sub>O) wherein trapping with acetic anhydride can provide compound 100a. Compound 100a may be treated with *N*-bromosuccinimide (NBS) and AIBN to afford bromination product 100b. Treatment of 100b with a strong non-nucleophilic base e.g., LHMDS (lithium hexamethyldisilazide) can provide compound 118. The ketone moiety in compound 118 may be reduced under standard conditions (e.g., with sodium borohydride) to provide compound 119, whereupon treatment with ammonium hydroxide can displace the chloro group to afford compound 36 which is a compound of Formula I-H wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>;  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl;  $R^{10}$  is OH; and  $R^{11}$  is hydrogen. Compound 36 may be subjected to chiral resolution to provide enantiomerically enriched compounds 36a and 36b both of which are compounds of Formula I-H wherein  $R^x$ ,  $R^z$ ,  $R^{10}$ , and  $R^{11}$  are as defined for compound 36. Alternatively, compound 118 can be subjected to chiral reduction conditions to afford compound 119a or 119b selectively. The chloro group in compound 119a or 119b can be displaced with ammonium hydroxide to provide compound 36a and 36b, wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>;  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl;  $R^{10}$  is OH; and  $R^{11}$  is hydrogen.



Scheme 24

[00286] Scheme 24 shows a process for preparing a compound of Formula I-A (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is CR<sup>7</sup>R<sup>8</sup>, Y is C(=O), Z is N, and ----- is a single bond) wherein R<sup>7</sup> and R<sup>8</sup> are hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is as defined for Formula I, wherein ring A is substituted with R<sup>d</sup>NHC(=O)-; and R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>.

**[00287]** Compound 53 (wherein  $R^x$  is as defined in Scheme 11) which may be prepared according to Scheme 11 may be subjected under alcohol oxidation conditions (e.g., in the presence of Dess-Martin periodinane) to afford aldehyde 120. Compound 120 can be subjected under Wittig olefination conditions (e.g., with benzyl 2-(triphenyl- $\lambda$ 5-phosphanylidene)acetate) to provide compound 121, whereupon reduction of the olefin in 121 can afford compound 122. The chloro group in 122 can be displaced with a nucleophile of formula  $P^6-NH_2$  wherein  $P^6$  is an amino protecting group (e.g., benzyl, e.g., 2,4-dimethoxybenzyl, DMB) to provide compound 123. Iodination of compound 123 (e.g., with N-iodosuccinimide) can lead to compound 124. Compound 124 can be coupled with boronic acid or boronic ester 125 under appropriate palladium-catalyzed cross coupling conditions (e.g., Suzuki coupling) to provide compound 126. In compound 125, each  $R^1$  is independently H or (1-6C)alkyl, or each  $R^1$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl); and  $R''$  is C1-C6 alkyl. Hydrolysis of the ester groups in 126 provides dicarboxylic acid compound 127 whereupon treatment of 127 under standard amide coupling conditions allows intramolecular cyclization to provide compound 128. Compound 128 may be coupled with a reagent of formula  $R^d-NH_2$  wherein  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup> under standard amide coupling conditions. Subsequent removal of the amino protecting group (e.g., with TFA) provides compound 129 which is a compound of Formula I-A, wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; ring A is substituted with  $R^dNHC(=O)$ , and  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>.



Scheme 25

[00288] Scheme 25 shows a process for preparing a compound of Formula I-L (i.e., compounds of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>, X is CR<sup>7</sup>R<sup>8</sup>, Y is CR<sup>10</sup>R<sup>11</sup>, Z is N, and — is a single bond) wherein R<sup>7</sup> and R<sup>8</sup> are hydrogen, R<sup>10</sup> and R<sup>11</sup> are hydrogen, R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is as defined for Formula I, wherein ring A is substituted with R<sup>d</sup>NHC(=O)- or (C1-C6 alkyl)C(=O)NH-; and R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>.

[00289] Compound 120 (wherein R<sup>x</sup> is as defined for Scheme 24) which may be prepared according to Scheme 24 may be subjected to Wittig olefination reaction (e.g., with ethyl 2-(triphenyl-λ5-phosphanylidene)acetate) to provide compound 130. Reduction of the olefin moiety in 130 provides compound 131, whereupon iodination (e.g., with N-iodosuccinimide) affords compound 132. Reduction of the ester moiety in 132 (e.g., with DIBAL-H) provides alcohol-containing compound 133. Protection of the alcohol in 133 with a reagent of formula P<sup>5</sup>-X wherein P<sup>5</sup> is a suitable hydroxy protecting group (e.g., a silyl protecting group, e.g., tert-butyldimethyl silyl) and X is a leaving group (e.g., halogen) affords

compound 134. Treatment of 134 with  $P^6-NH_2$  wherein  $P^6$  is an amino protecting group (e.g., benzyl, e.g., 2,4-dimethoxybenzyl, DMB) can provide compound 135. Subjecting 135 to appropriate palladium-catalyzed cross-coupling (e.g., Suzuki coupling) conditions with a boronic acid or boronic ester of formula 125 can provide compound 136. In compound 125, each  $R^1$  is independently H or (1-6C)alkyl, or each  $R^1$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl); and  $R''$  is C1-C6 alkyl. Removal of the hydroxy protecting group in 136 (e.g., with TBAF) can provide compound 137. Exposure of 137 to mesyl chloride allows methanesulfonylation of the primary alcohol group. Therefore, the NH group of ring A can displace this ensuing mesylate intramolecularly upon treatment with a base (e.g., cesium carbonate) to provide compound 138. Hydrolysis of the ester group in 138 affords 139, whereupon coupling with  $R^dNH_2$  under standard amide coupling conditions followed by removal of the amino protecting group provides compound 140 which is a compound of Formula I-L, wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; ring A is substituted with  $R^dNHC(=O)$ , and  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>. Alternatively, compound 140 can be obtained from compound 139 by sequential removal of the amino protecting group and coupling with  $R^dNH_2$  under standard amide coupling conditions.

**[00290]** Further, compound 139 may be treated with diphenylphosphoryl azide (DPPA) and trimethylamine in a Curtius-type rearrangement. Treatment of the intermediate (not shown) to (C1-C6 alkylCO)<sub>2</sub>O followed by removal of the amino protecting group can provide compound 141 which is a compound of Formula I-L, wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and ring A is substituted with (C1-C6 alkyl)C(=O)NH-.

**[00291]** The term "amino protecting group" as used herein refers to a derivative of the groups commonly employed to block or protect an amino group while reactions are carried out on other functional groups on the compound. Examples of suitable protecting groups for use in any of the processes described herein include carbamates, amides, alkyl and aryl groups, benzyl groups and substituted benzyl groups, imines, as well as many N-heteroatom derivatives which can be removed to regenerate the desired amine group. Non-limiting examples of amino protecting groups are 2,4-dimethoxybenzyl (DMB), acetyl, trifluoroacetyl, t-butyloxycarbonyl ("Boc"), benzyloxycarbonyl ("CBz") [2-(trimethylsilyl)ethoxy]methyl (SEM), dimethylformamide dimethyl acetal, and 9-fluorenylmethylenecarbonyl ("Fmoc"). Further examples of these groups, and other protecting groups, are found in T. W. Greene, et al. Greene's Protective Groups in Organic Synthesis. New York: Wiley Interscience, 2006.


**[00292]** Nitrogen atoms in compounds described in any of the above methods may be protected with one or more of any convenient nitrogen protecting group, for example, as described in Greene & Wuts, eds., "Protecting Groups in Organic Synthesis", 2<sup>nd</sup> ed. New York; John Wiley & Sons, Inc., 1991. A nitrogen protecting group can be any "amino protecting group" as described above. Examples of nitrogen protecting groups include acyl and alkoxycarbonyl groups, such as t-butyloxycarbonyl (BOC), phenoxycarbonyl, and [2-(trimethylsilyl)ethoxy]methyl (SEM).

**[00293]** Hydroxy groups may be protected with any convenient hydroxy protecting group, for example, as described in T. W. Greene, et al., Greene's Protective Groups in Organic Synthesis. New

York: Wiley Interscience, 2006. Examples include benzyl, trityl, silyl ethers (e.g., tert-butyldimethylsilyl), and the like.


**[00294]** Accordingly, further provided herein is a process for preparing of a compound of Formula I or a pharmaceutically acceptable salt thereof as defined herein which comprises:

**[00295]** (a) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is  $C(=O)$ ; Z is N; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, subjecting a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, Ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Ring A in Formula I, and  $P^6$  is an amino protecting group, to an intramolecular cyclization; and removing the amino protecting group  $P^6$  on the resulting intermediate;


**[00296]** (b) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is  $C(=O)$ ; Z is N; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, wherein ring A is substituted with  $NHR^dC(=O)-$ , and  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>, coupling a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, and  $P^6$  is an amino protecting group, with a compound of formula  $R^dNH_2$  wherein  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>;

and removing the amino protecting group  $P^6$ ;


**[00297]** (c) for a compound of formula I, wherein W is  $CR^3R^4$ ; X is  $NR^9$ ; Y is  $C(=O)$ ; Z is N, and ----- is a single bond;  $R^9$  is hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is a monocyclic heteroaryl ring or a bicyclic heteroaryl ring as defined for Formula I, reacting a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; Ring A is a monocyclic heteroaryl ring or a bicyclic heteroaryl ring as defined for Formula I; and  $P^6$  is an amino protecting group, with Boc<sub>2</sub>O; and


removing the amino protecting group  $P^6$ ;

**[00298]** (d) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>; X is NR<sup>9</sup>; Y is C(=O); Z is N; ----- is a single bond; R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl-, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-; R<sup>x</sup> is C1-




C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is

wherein the bond labeled "a" indicates the point of attachment to Y, n is 0, 1, 2, 3 or 4, and each R<sup>z</sup> is independently selected from R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH- wherein R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>, reacting a compound of the following formula:




wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; n is 0, 1, 2, 3 or 4, and each R<sup>z</sup> is independently selected from R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH- wherein R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>, with a compound of formula R<sup>9</sup>-X, wherein X is a leaving group; and R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a hydroxy protecting group, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-; and removing the hydroxy protecting group if present;

**[00299]** (e) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>; Y is C(=O); Z is N; ----- is a single bond); R<sup>9</sup> is hydrogen; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y, each  $R^Z$  is independently selected from C1-C6 alkyl and C3-C6 cycloalkyl,  $n$  is 0, 1 or 2, and Ring A is further optionally substituted with one occurrence of halo, reacting a compound of the following formula




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, each  $R^Z$  is independently selected from halogen, C1-C6 alkyl and C3-C6 cycloalkyl,  $n$  is 0, 1 or 2, and the pyrazolyl ring is further optionally substituted with one occurrence of halo, with a reagent of formula  $(p\text{-NO}_2\text{C}_6\text{H}_5\text{O})_2\text{CO}$ ;

**[00300]** (f) for a compound of Formula I, wherein W is  $\text{CR}^3\text{R}^4$ ; X is  $\text{NR}^9$ ; Y is  $\text{C}(=\text{O})$ , Z is C; ----- is a single bond;  $R^9$  is hydrogen;  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl with ammonium hydroxide;

**[00301]** (g) for a compound of Formula I, wherein W is  $\text{CR}^3\text{R}^4$ ; X is  $\text{NR}^9$ ; Y is  $\text{C}(=\text{O})$ , Z is C; ----- is a single bond;  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl-, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-;  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with a compound of formula  $R^9$ -X wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl- the hydroxy group is protected with a suitable hydroxy protecting group, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-; and removing the hydroxy protecting group if present;

**[00302]** (h) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>, X is NR<sup>9</sup>; Y is C(=O); Z is C; and ----- is a single bond;  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-;  $R^X$  is as defined for Formula I; and Ring A is



wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, removing the amino protecting group P<sup>2</sup> on a compound of the following formula:




wherein  $R^X$  is as defined for Formula I,  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl,  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-, and P<sup>2</sup> is an amino protecting group;

**[00303]** (i) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>; X is NR<sup>9</sup>; Y is C(=O); Z is C; ----- is a single bond; Ring A is a 6-membered aryl ring optionally substituted with one or more substituents independently selected from halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>C</sup>NHC(=O)-, R<sup>C</sup>C(=O)NH-, or




$R^c\text{NHC(=O)NH-}$  wherein  $R^c$  is C3-C6 cycloalkyl or (C3-C6 cycloalkyl) $\text{CH}_2-$ ;  $R^9$  is hydrogen; and  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $\text{Cyc}^2$ , or  $\text{Cyc}^3$ , subjecting a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $\text{Cyc}^2$ , or  $\text{Cyc}^3$ ,  $n$  is 0, 1, 2, 3, or 4, each  $R^z$  is independently halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^c\text{NHC(=O)-}$ ,  $R^c\text{C(=O)NH-}$ , or  $R^c\text{NHC(=O)NH-}$  wherein  $R^c$  is C3-C6 cycloalkyl or (C3-C6 cycloalkyl) $\text{CH}_2-$ , and  $P^6$  is an amino protecting group, to an intramolecular cyclization;


and removing the amino protecting group  $P^6$ ;

**[00304]** (j) for a compound of Formula I, wherein  $W$  is  $\text{CR}^3\text{R}^4$ ;  $X$  is  $\text{NR}^9$ ;  $Y$  is  $\text{C(=O)}$ ;  $Z$  is  $\text{C}$ ; ----- is a single bond; Ring A is a 6-membered aryl ring optionally substituted with one or more substituents independently selected from halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^c\text{NHC(=O)-}$ ,  $R^c\text{C(=O)NH-}$ , or  $R^c\text{NHC(=O)NH-}$  wherein  $R^c$  is C3-C6 cycloalkyl or (C3-C6 cycloalkyl) $\text{CH}_2-$ ;  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-; and  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $\text{Cyc}^2$ , or  $\text{Cyc}^3$ , reacting a compound of the following formula:




wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $\text{Cyc}^2$ , or  $\text{Cyc}^3$ ,  $n$  is 0, 1, 2, 3, or 4, and each  $R^z$  is independently selected from halogen, C3-C6 cycloalkyl, C1-C6 alkoxy,  $R^c\text{NHC(=O)-}$ ,  $R^c\text{C(=O)NH-}$ , or  $R^c\text{NHC(=O)NH-}$  wherein  $R^c$  is C3-C6 cycloalkyl or (C3-C6 cycloalkyl) $\text{CH}_2-$ , with a compound of formula  $R^9\text{-X}$ , wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-, and  $X$  is a leaving group; and removing the hydroxy protecting group if present;

**[00305]** (k) for a compound of Formula I wherein  $W$  is  $\text{CR}^3\text{R}^4$ ;  $X$  is  $\text{NR}^9$ ;  $Y$  is  $\text{C(=O)}$ ;  $Z$  is  $\text{C}$ ; ----- is a single bond;  $R^9$  is hydrogen;  $R^x$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl,  $\text{Cyc}^2$ , or  $\text{Cyc}^3$ ; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y; and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:



wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, with N-bromosuccinimide and AIBN; and reacting the resulting intermediate with ammonium hydroxide; or


reacting a compound of the following formula:




wherein PG is an amino protecting group, and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, with N-bromosuccinimide and AIBN; reacting the resulting intermediate with ammonium hydroxide; and

removing the amino protecting group PG on the resulting intermediate;

**[00306]** (I) for a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>; X is NR<sup>9</sup>; Y is C(=O); Z is C; ----- is a single bond; R<sup>9</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl-, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)-; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with a compound of formula  $R^9-X$  wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group, (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)-, and X is a leaving group; and removing the hydroxy protecting group if present;


**[00307]** (m) for a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>; X is NR<sup>9</sup>; Y is C(=O); Z is C; and ----- is a single bond); R<sup>9</sup> is hydrogen; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is as defined for Formula I, subjecting a compound of the following formula:




wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, P<sup>6</sup> is an amino protecting group, and R'' is C1-C6 alkyl, to an intramolecular cyclization; and

removing the amino protecting group P<sup>6</sup>;

**[00308]** (n) for a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>; X is O; Y is C(=O); Z is C; and ----- is a single bond; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, removing the amino protecting group P<sup>2</sup> on a compound of the following formula:




wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, and P<sup>2</sup> is an amino protecting group;

**[00309]** (o) for a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>; X is NR<sup>9</sup>; Y is SO<sub>2</sub>; Z is C, and ----- is a single bond; R<sup>9</sup> is hydrogen; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with ammonium hydroxide;

**[00310]** (p) for a compound of Formula I wherein W is  $CR^3R^4$ ; X is  $NR^9$ ; Y is  $SO_2$ ; Z is C, and ----- is a single bond;  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-;  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with a compound of formula  $R^9-X$  wherein  $R^9$  is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl- wherein the hydroxy group is protected with a suitable hydroxy protecting group, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, (hetCyc<sup>1</sup>)C1-C6 alkyl-, (Ar<sup>1</sup>)C1-C6 alkyl-, or (hetAr<sup>1</sup>)C1-C6 alkyl-; and removing the hydroxy protecting group if present;

**[00311]** (q) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is  $SO_2$ ; Z is C; and ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl with ammonium hydroxide;

**[00312]** (r) for a compound of Formula I wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is C(=O); Z is C; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with polyphosphoric acid;

**[00313]** (s) for a compound of Formula I wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is C(=O); Z is C; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^X$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^x$  is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with ammonium hydroxide;

**[00314]** (t) for a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>; X is CR<sup>7</sup>R<sup>8</sup>; Y is C(=O); Z is C; ----- is a single bond; R<sup>7</sup> and R<sup>8</sup> are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, removing the amino protecting group P<sup>6</sup> on a compound of the following formula:




wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>,  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, and P<sup>6</sup> is an amino protecting group;

**[00315]** (u) for a compound of Formula I wherein W is CR<sup>3</sup>R<sup>4</sup>; X is CR<sup>7</sup>R<sup>8</sup>; Y is C(=O); Z is C; ----- is a single bond; R<sup>7</sup> and R<sup>8</sup> are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with a compound of formula  $R^X-X$  wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $Cyc^2$ ,  $Cyc^3$ ,  $hetCyc^3$ ,  $Ar^3$  or  $hetAr^3$ , a compound of formula  $R^X-OH$  wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $Cyc^2$ ,  $Cyc^3$ , or  $hetCyc^3$ , or a compound of formula  $R^XB(OR')_2$  wherein  $R^X$  is C1-C6 alkyl, fluoroC1-C6 alkyl,  $Cyc^2$ ,  $Cyc^3$ ,  $hetCyc^3$ ,  $Ar^3$  or  $hetAr^3$ , and each  $R'$  is independently H or (1-6C)alkyl, or each  $R'$  together with the atoms to which they are connected form a 5-6 membered ring optionally substituted with 1-4 substituents selected from (C1-C3 alkyl);

**[00316]** (v) for a compound of formula I, wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is  $CR^{10}R^{11}$ ; Z is C; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^X$  is as defined for Formula I,  $R^{10}$  and  $R^{11}$  are as defined for Formula I, and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, functionalizing a compound of the following formula:




wherein  $R^X$  is as defined for Formula I, and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl; and optionally subjecting the resulting product to chiral separation;

**[00317]** (w) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is  $CR^{10}R^{11}$ ; Z is C; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^X$  is as defined for Formula I,  $R^{10}$  is hydroxyl,  $R^{11}$  is hydrogen, and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^Z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^x$  is as defined for Formula I, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, with ammonium hydroxide; and

optionally subjecting the resulting product to chiral separation;

**[00318]** (x) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $CR^7R^8$ ; Y is  $CR^{10}R^{11}$ ; Z is C; ----- is a single bond;  $R^7$  and  $R^8$  are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>,  $R^{10}$  is hydroxyl, and  $R^{11}$  is hydrogen, and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, subjecting a compound of the following formula:



wherein  $R^x$  is as defined for Formula I, and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, to chiral separation; and


reacting the resulting intermediate with ammonium hydroxide;

**[00319]** (y) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $NR^9$ ; Y is  $CR^{10}R^{11}$ ; Z is C; ----- is a single bond;  $R^{10}$  and  $R^{11}$  are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>;  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup> or (hetAr<sup>1</sup>)C1-C6 alkyl-, and Ring A is



wherein the bond labeled "a" indicates the point of attachment to Y and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:







wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>,  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, and each  $P^4$  is independently an amino protecting group, with a compound of formula  $R^9NH_2$ , wherein  $R^9$  is C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-,  $Ar^1$ , ( $Ar^1$ )C1-C6 alkyl-, hetAr<sup>1</sup> or (hetAr<sup>1</sup>)C1-C6 alkyl-, in the presence of a reducing agent; and

removing the amino protecting groups  $P^4$  on the resulting intermediate;

**[00320]** (z) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is  $NR^9$ ; Y is  $CR^{10}R^{11}$ ; Z is C; ----- is a single bond;  $R^{10}$  and  $R^{11}$  are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>;  $R^9$  is (C1-C6 alkyl)SO<sub>2</sub> or  $R^9C(=O)$  wherein  $R^9$  is C1-C6 alkyl or fluoroC1-C6 alkyl, and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>,  $R^z$  is C1-C6 alkyl or C3-C6 cycloalkyl, and each  $P^4$  is independently an amino protecting group, with a compound of formula (C1-C6 alkyl)SO<sub>2</sub>Cl or  $R^9C(=O)Cl$  wherein  $R^9$  is C1-C6 alkyl or fluoroC1-C6 alkyl; and

removing the amino protecting groups  $P^4$  on the resulting intermediate;

**[00321]** (aa) for a compound of Formula I, wherein W is  $CR^3R^4$ ; X is O; Y is  $CR^{10}R^{11}$ ; Z is C; ----- is a single bond;  $R^{10}$  and  $R^{11}$  are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:




wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, each P<sup>4</sup> is independently an amino protecting group, and P<sup>3</sup> is a hydroxy protecting group, with CF<sub>3</sub>SO<sub>3</sub>TMS and Et<sub>3</sub>SiH;

**[00322]** (bb) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>; X is CR<sup>5</sup>; Y is CR<sup>6</sup>; Z is C; ----- is a double bond; R<sup>5</sup> and R<sup>6</sup> are hydrogen; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; and Ring A is




wherein the bond labeled "a" indicates the point of attachment to Y and R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, reacting a compound of the following formula:



wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, R<sup>z</sup> is C1-C6 alkyl or C3-C6 cycloalkyl, and P is an amino protecting group, with iodomethane in the presence of a base; and

removing the amino protecting group P on the resulting intermediate;


**[00323]** (cc) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>; X is CR<sup>7</sup>R<sup>8</sup>; Y is CR<sup>10</sup>R<sup>11</sup>; Z is N; ----- is a single bond; R<sup>7</sup> and R<sup>8</sup> are hydrogen; R<sup>10</sup> and R<sup>11</sup> are hydrogen; R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, wherein ring A is substituted with NHR<sup>d</sup>C(=O)-, and R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>, removing the amino protecting group P<sup>6</sup> on a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, and  $P^6$  is an amino protecting group;

and coupling the resulting intermediate with a compound of formula  $R^dNH_2$  wherein  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>;


**[00324]** (dd) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>; X is CR<sup>7</sup>R<sup>8</sup>; Y is CR<sup>10</sup>R<sup>11</sup>; Z is N; ----- is a single bond; R<sup>7</sup> and R<sup>8</sup> are hydrogen; R<sup>10</sup> and R<sup>11</sup> are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, wherein ring A is substituted with NHR<sup>d</sup>C(=O)-, and  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>, coupling a compound of the following formula:



wherein  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, and  $P^6$  is an amino protecting group, with a compound of formula  $R^dNH_2$  wherein  $R^d$  is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>;

and removing the amino protecting group  $P^6$ ; or

**[00325]** (ee) for a compound of Formula I, wherein W is CR<sup>3</sup>R<sup>4</sup>; X is CR<sup>7</sup>R<sup>8</sup>; Y is CR<sup>10</sup>R<sup>11</sup>; Z is N; ----- is a single bond; R<sup>7</sup> and R<sup>8</sup> are hydrogen; R<sup>10</sup> and R<sup>11</sup> are hydrogen;  $R^x$  is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>; ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, wherein ring A is substituted with (C1-C6 alkyl)C(=O)NH-, reacting a compound of the following formula:



wherein R<sup>x</sup> is C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, or Cyc<sup>3</sup>, ring A is a monocyclic heteroaryl ring or bicyclic heteroaryl ring as defined for Formula I, and P<sup>6</sup> is an amino protecting group, with diphenylphosphoryl azide (DPPA);

reacting the resulting intermediate with a compound of formula (C1-C6 alkylCO)<sub>2</sub>O; and

removing the amino protecting group P<sup>6</sup> on the resulting intermediate,

**[00326]** optionally removing any remaining protecting groups, and optionally forming a pharmaceutically acceptable salt thereof.

**[00327]** The ability of test compounds to act as RET inhibitors may be demonstrated by the assay described in Examples A, B, C, and D. IC<sub>50</sub> values are shown in **Table 5**.

**[00328]** In some embodiments, the compounds provided herein exhibit potent and selective RET inhibition. For example, the compounds provided herein exhibit nanomolar potency against wild type RET and a RET kinase encoded by a RET gene including an activating mutation or a RET kinase inhibitor resistance mutation, including, for example, the KIF5B-RET fusion, G810R and G810S ATP cleft front mutations, M918T activating mutation, and V804M, V804L, and V804E gatekeeper mutations, with minimal activity against related kinases.

**[00329]** In some embodiments, the compounds provided herein exhibit nanomolar potency against an altered RET fusion protein encoded by a RET gene encoding the RET fusion protein (e.g. any of the RET fusion proteins described herein including, without limitation, CCDC6-RET or KIF5B-RET) which RET gene includes a RET kinase inhibitor resistance mutation (e.g., any of the RET mutations described herein including, without limitation, V804M, V804L, or V804E) such that the altered RET protein is a RET fusion protein that exhibits RET kinase resistance due to the presence of a RET kinase inhibitor resistance amino acid substitution or deletion. Non-limiting examples include CCDC6-RET-V804M and KIF5B-RET-V804M. In some embodiments, the compounds provided herein exhibit nanomolar potency against an altered RET protein encoded by a RET gene that includes a RET mutation (e.g. any of the RET mutations described herein including, without limitation, C634W or M918T) and that includes a RET kinase inhibitor resistance mutation (e.g., any of the RET kinase inhibitor resistance mutations described herein including, without limitation, V804M, V804L, or V804E) such that the altered RET protein includes a RET substitution caused by the RET mutation (e.g., a RET primary mutation) and the altered RET protein exhibits RET kinase resistance due to the presence of a RET kinase inhibitor resistance amino acid substitution or deletion.

**[00330]** In some embodiments, the compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt thereof, selectively target a RET kinase. For example, a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, can selectively target a RET kinase over another kinase or non-kinase target.

**[00331]** In some embodiments, a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, exhibits at least a 30-fold selectivity for a RET kinase over another kinase. For example, a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, exhibits at least a 40-fold selectivity; at least a 50-fold selectivity; at least a 60-fold selectivity; at least a 70-fold selectivity; at least a 80-fold selectivity;

at least a 90-fold selectivity; at least 100-fold selectivity; at least 200-fold selectivity; at least 300-fold selectivity; at least 400-fold selectivity; at least 500-fold selectivity; at least 600-fold selectivity; at least 700-fold selectivity; at least 800-fold selectivity; at least 900-fold selectivity; or at least 1000-fold selectivity for a RET kinase over another kinase. In some embodiments, selectivity for a RET kinase over another kinase is measured in a cellular assay (e.g., a cellular assay as provided herein).

**[00332]** In some embodiments, the compounds provided herein can exhibit selectivity for a RET kinase over a KDR kinase (e.g., VEGFR2). In some embodiments, the selectivity for a RET kinase over a KDR kinase is observed without loss of potency for a RET kinase encoded by a RET gene including an activating mutation or a RET kinase inhibitor resistance mutation (e.g., a gatekeeper mutant). In some embodiments, the selectivity over a KDR kinase is at least 10-fold (e.g., at least a 40-fold selectivity; at least a 50-fold selectivity; at least a 60-fold selectivity; at least a 70-fold selectivity; at least a 80-fold selectivity; at least a 90-fold selectivity; at least 100-fold selectivity; at least 150-fold selectivity; at least 200-fold selectivity; at least 250-fold selectivity; at least 300-fold selectivity; at least 350-fold selectivity; or at least 400-fold selectivity) as compared to the inhibition of KIF5B-RET (e.g., the compounds are more potent against KIF5B-RET than KDR). In some embodiments, the selectivity for a RET kinase over a KDR kinase is about 30-fold. In some embodiments, the selectivity for a RET kinase over a KDR kinase is at least 100-fold. In some embodiments, the selectivity for a RET kinase over a KDR kinase is at least 150-fold. In some embodiments, the selectivity for a RET kinase over a KDR kinase is at least 400-fold. Without being bound by any theory, potent KDR kinase inhibition is believed to be a common feature among multikinase inhibitors (MKIs) that target RET and may be the source of the dose-limiting toxicities observed with such compounds.

**[00333]** In some embodiments, inhibition of V804M is similar to that observed for wild-type RET. For example, inhibition of V804M is within about 2-fold (e.g., about 5-fold, about 7-fold, about 10-fold) of inhibition of wild-type RET (e.g., the compounds were similarly potent against wild-type RET and V804M). In some embodiments, selectivity for a wildtype or V804M RET kinase over another kinase is measured in an enzyme assay (e.g., an enzyme assay as provided herein). In some embodiments, the compounds provided herein exhibit selective cytotoxicity to RET-mutant cells.

**[00334]** In some embodiments, inhibition of G810S and/or G810R is similar to that observed for wild-type RET. For example, inhibition of G810S and/or G810R is within about 2-fold (e.g., about 5-fold, about 7-fold, about 10-fold) of inhibition of wild-type RET (e.g., the compounds were similarly potent against wild-type RET and G810S and/or G810R). In some embodiments, selectivity for a wildtype or G810S and/or G810R RET kinase over another kinase is measured in an enzyme assay (e.g., an enzyme assay as provided herein). In some embodiments, the compounds provided herein exhibit selective cytotoxicity to RET-mutant cells.

**[00335]** In some embodiments, the compounds provided herein exhibit brain and/or central nervous system (CNS) penetrance. Such compounds are capable of crossing the blood brain barrier and inhibiting a RET kinase in the brain and/or other CNS structures. In some embodiments, the compounds provided herein are capable of crossing the blood brain barrier in a therapeutically effective amount. For example, treatment of a patient with cancer (e.g., a RET-associated cancer such as a RET-associated brain or CNS cancer) can include administration (e.g., oral administration) of the compound to the patient. In some such embodiments, the compounds provided herein are useful for treating a primary

brain tumor or metastatic brain tumor. For example, a RET-associated primary brain tumor or metastatic brain tumor.

**[00336]** In some embodiments, the compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, exhibit one or more of high GI absorption, low clearance, and low potential for drug-drug interactions.

**[00337]** Compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof are useful for treating diseases and disorders which can be treated with a RET kinase inhibitor, such as RET-associated diseases and disorders, e.g., proliferative disorders such as cancers, including hematological cancers and solid tumors (e.g., advanced solid tumors and/or RET-fusion positive solid tumors), and gastrointestinal disorders such as IBS.

**[00338]** As used herein, terms "treat" or "treatment" refer to therapeutic or palliative measures. Beneficial or desired clinical results include, but are not limited to, alleviation, in whole or in part, of symptoms associated with a disease or disorder or condition, diminishment of the extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state (e.g., one or more symptoms of the disease), and remission (whether partial or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival as compared to expected survival if not receiving treatment.

**[00339]** As used herein, the terms "subject," "individual," or "patient," are used interchangeably, refers to any animal, including mammals such as mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, primates, and humans. In some embodiments, the patient is a human. In some embodiments, the subject has experienced and/or exhibited at least one symptom of the disease or disorder to be treated and/or prevented. In some embodiments, the subject has been identified or diagnosed as having a cancer with a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same (a RET-associated cancer) (e.g., as determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit). In some embodiments, the subject has a tumor that is positive for a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same (e.g., as determined using a regulatory agency-approved assay or kit). The subject can be a subject with a tumor(s) that is positive for a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same (e.g., identified as positive using a regulatory agency-approved, e.g., FDA-approved, assay or kit). The subject can be a subject whose tumors have a dysregulation of a RET gene, a RET protein, or expression or activity, or a level of the same (e.g., where the tumor is identified as such using a regulatory agency-approved, e.g., FDA-approved, kit or assay). In some embodiments, the subject is suspected of having a RET-associated cancer. In some embodiments, the subject has a clinical record indicating that the subject has a tumor that has a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same (and optionally the clinical record indicates that the subject should be treated with any of the compositions provided herein). In some embodiments, the patient is a pediatric patient.

**[00340]** The term "pediatric patient" as used herein refers to a patient under the age of 21 years at the time of diagnosis or treatment. The term "pediatric" can be further be divided into various subpopulations including: neonates (from birth through the first month of life); infants (1 month up to

two years of age); children (two years of age up to 12 years of age); and adolescents (12 years of age through 21 years of age (up to, but not including, the twenty-second birthday)). Berhman RE, Kliegman R, Arvin AM, Nelson WE. *Nelson Textbook of Pediatrics*, 15th Ed. Philadelphia: W.B. Saunders Company, 1996; Rudolph AM, et al. *Rudolph's Pediatrics*, 21st Ed. New York: McGraw-Hill, 2002; and Avery MD, First LR. *Pediatric Medicine*, 2nd Ed. Baltimore: Williams & Wilkins; 1994. In some embodiments, a pediatric patient is from birth through the first 28 days of life, from 29 days of age to less than two years of age, from two years of age to less than 12 years of age, or 12 years of age through 21 years of age (up to, but not including, the twenty-second birthday). In some embodiments, a pediatric patient is from birth through the first 28 days of life, from 29 days of age to less than 1 year of age, from one month of age to less than four months of age, from three months of age to less than seven months of age, from six months of age to less than 1 year of age, from 1 year of age to less than 2 years of age, from 2 years of age to less than 3 years of age, from 2 years of age to less than seven years of age, from 3 years of age to less than 5 years of age, from 5 years of age to less than 10 years of age, from 6 years of age to less than 13 years of age, from 10 years of age to less than 15 years of age, or from 15 years of age to less than 22 years of age.

**[00341]** In certain embodiments, compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof are useful for preventing diseases and disorders as defined herein (for example, autoimmune diseases, inflammatory diseases, and cancer). The term "preventing" as used herein means the prevention of the onset, recurrence or spread, in whole or in part, of the disease or condition as described herein, or a symptom thereof.

**[00342]** The term "RET-associated disease or disorder" as used herein refers to diseases or disorders associated with or having a dysregulation of a RET gene, a RET kinase (also called herein RET kinase protein), or the expression or activity or level of any (e.g., one or more) of the same (e.g., any of the types of dysregulation of a RET gene, a RET kinase, a RET kinase domain, or the expression or activity or level of any of the same described herein). Non-limiting examples of a RET-associated disease or disorder include, for example, cancer and gastrointestinal disorders such as irritable bowel syndrome (IBS).

**[00343]** The term "RET-associated cancer" as used herein refers to cancers associated with or having a dysregulation of a RET gene, a RET kinase (also called herein RET kinase protein), or expression or activity, or level of any of the same. Non-limiting examples of a RET-associated cancer are described herein.

**[00344]** The phrase "dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same" refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a RET kinase domain and a fusion partner, a mutation in a RET gene that results in the expression of a RET protein that includes a deletion of at least one amino acid as compared to a wildtype RET protein, a mutation in a RET gene that results in the expression of a RET protein with one or more point mutations as compared to a wildtype RET protein, a mutation in a RET gene that results in the expression of a RET protein with at least one inserted amino acid as compared to a wildtype RET protein, a gene duplication that results in an increased level of RET protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of RET protein in a cell), an alternative spliced version of a RET mRNA that results in a

RET protein having a deletion of at least one amino acid in the RET protein as compared to the wild-type RET protein), or increased expression (e.g., increased levels) of a wildtype RET kinase in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same, can be a mutation in a RET gene that encodes a RET protein that is constitutively active or has increased activity as compared to a protein encoded by a RET gene that does not include the mutation. For example, a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of RET that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not RET). In some examples, dysregulation of a RET gene, a RET protein, or expression or activity or level of any of the same can be a result of a gene translocation of one RET gene with another non-RET gene. Non-limiting examples of fusion proteins are described in Table 1. Non-limiting examples of RET kinase protein point mutations/insertions/deletions are described in Tables 2 and 2a. Additional examples of RET kinase protein mutations (e.g., point mutations) are RET inhibitor resistance mutations. Non-limiting examples of RET inhibitor resistance mutations are described in Tables 3 and 4.

**[00345]** In some embodiments, dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same can be caused by an activating mutation in a RET gene (see, e.g., chromosome translocations that result in the expression of any of the fusion proteins listed in Table 1). In some embodiments, dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same can be caused by a genetic mutation that results in the expression of a RET kinase that has increased resistance to inhibition by a RET kinase inhibitor and/or a multi-kinase inhibitor (MKI), e.g., as compared to a wildtype RET kinase (see, e.g., the amino acid substitutions in Tables 3 and 4). In some embodiments, dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same can be caused by a mutation in a nucleic acid encoding an altered RET protein (e.g., a RET fusion protein or a RET protein having a mutation (e.g., a primary mutation)) that results in the expression of an altered RET protein that has increased resistance to inhibition by a RET kinase inhibitor and/or a multi-kinase inhibitor (MKI), e.g., as compared to a wildtype RET kinase (see, e.g., the amino acid substitutions in Tables 3 and 4). The exemplary RET kinase point mutations, insertions, and deletions shown in Tables 2 and 2a can be caused by an activating mutation and/or can result in the expression of a RET kinase that has increased resistance to inhibition by a RET kinase inhibitor and/or a multi-kinase inhibitor (MKI).

**[00346]** The term “activating mutation” describes a mutation in a RET kinase gene that results in the expression of a RET kinase that has an increased kinase activity, e.g., as compared to a wildtype RET kinase, e.g., when assayed under identical conditions. For example, an activating mutation can result in the expression of a fusion protein that includes a RET kinase domain and a fusion partner. In another example, an activating mutation can be a mutation in a RET kinase gene that results in the expression of a RET kinase that has one or more (e.g., two, three, four, five, six, seven, eight, nine, or ten) amino acid substitutions (e.g., any combination of any of the amino acid substitutions described herein) that has increased kinase activity, e.g., as compared to a wildtype RET kinase, e.g., when assayed under identical conditions. In another example, an activating mutation can be a mutation in a RET kinase gene that results in the expression of a RET kinase that has one or more (e.g., two, three, four, five, six, seven,



eight, nine, or ten) amino acids deleted, e.g., as compared to a wildtype RET kinase, e.g., when assayed under identical conditions. In another example, an activating mutation can be a mutation in a RET kinase gene that results in the expression of a RET kinase that has at least one (e.g., at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 12, at least 14, at least 16, at least 18, or at least 20) amino acid inserted as compared to a wildtype RET kinase, e.g., the exemplary wildtype RET kinase described herein, e.g., when assayed under identical conditions. Additional examples of activating mutations are known in the art.

**[00347]** The term "wildtype" or "wild-type" describes a nucleic acid (e.g., a RET gene or a RET mRNA) or protein (e.g., a RET protein) that is found in a subject that does not have a RET-associated disease, e.g., a RET-associated cancer (and optionally also does not have an increased risk of developing a RET-associated disease and/or is not suspected of having a RET-associated disease), or is found in a cell or tissue from a subject that does not have a RET-associated disease, e.g., a RET-associated cancer (and optionally also does not have an increased risk of developing a RET-associated disease and/or is not suspected of having a RET-associated disease).

**[00348]** The term "regulatory agency" refers to a country's agency for the approval of the medical use of pharmaceutical agents with the country. For example, a non-limiting example of a regulatory agency is the U.S. Food and Drug Administration (FDA).

**[00349]** Provided herein is a method of treating cancer (e.g., a RET-associated cancer) in a patient in need of such treatment, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof. For example, provided herein are methods for treating a RET-associated cancer in a patient in need of such treatment, the method comprising a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the patient; and b) administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same includes one or more fusion proteins. Non-limiting examples of RET gene fusion proteins are described in Table 1. In some embodiments, the fusion protein is KIF5B-RET. In some embodiments, the dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same includes one or more RET kinase protein point mutations/insertions. Non-limiting examples of RET kinase protein point mutations/insertions/deletions are described in Tables 2 and 2a. In some embodiments, the RET kinase protein point mutations/insertions/deletions are selected from the group consisting of M918T, M918V, C634W, V804L, V804M, G810S, and G810R. In some embodiments, the RET kinase protein point mutations/insertions/deletions occur in a RET fusion protein (e.g., any of the RET gene fusion proteins described in Table 1). In some embodiments, a compound of Formula I is selected from: i) Examples 1-20, ii) Examples 21-40, iii) Examples 41-60, iv) Examples 61-80, v) Examples 81-100, vi) Examples 101-120, vii) Examples 121-140, or viii) Examples 141-151.

**[00350]** In some embodiments of any of the methods or uses described herein, the cancer (e.g., RET-associated cancer) is a hematological cancer. In some embodiments of any of the methods or uses described herein, the cancer (e.g., RET-associated cancer) is a solid tumor (e.g., an advanced solid tumor

and/or a RET-fusion positive solid tumor). In some embodiments of any of the methods or uses described herein, the cancer (e.g., RET-associated cancer) is a lung cancer (e.g., small cell lung carcinoma or non-small cell lung carcinoma), thyroid cancer (e.g., papillary thyroid cancer, medullary thyroid cancer (e.g., sporadic medullary thyroid cancer or hereditary medullary thyroid cancer), differentiated thyroid cancer, recurrent thyroid cancer, or refractory differentiated thyroid cancer), thyroid adenoma, endocrine gland neoplasms, lung adenocarcinoma, bronchioles lung cell carcinoma, multiple endocrine neoplasia type 2A or 2B (MEN2A or MEN2B, respectively), pheochromocytoma, parathyroid hyperplasia, breast cancer, mammary cancer, mammary carcinoma, mammary neoplasm, colorectal cancer (e.g., metastatic colorectal cancer), papillary renal cell carcinoma, ganglioneuromatosis of the gastroenteric mucosa, inflammatory myofibroblastic tumor, or cervical cancer. In some embodiments of any of the methods or uses described herein, the cancer (e.g., RET-associated cancer) is selected from the group of: acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), cancer in adolescents, adrenocortical carcinoma, anal cancer, appendix cancer, astrocytoma, atypical teratoid/rhabdoid tumor, basal cell carcinoma, bile duct cancer, bladder cancer, bone cancer, brain stem glioma, brain tumor, breast cancer, bronchial tumor, Burkitt lymphoma, carcinoid tumor, unknown primary carcinoma, cardiac tumors, cervical cancer, childhood cancers, chordoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative neoplasms, neoplasms by site, neoplasms, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, cutaneous angiosarcoma, bile duct cancer, ductal carcinoma in situ, embryonal tumors, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, Ewing sarcoma, extracranial germ cell tumor, extragonadal germ cell tumor, extrahepatic bile duct cancer, eye cancer, fallopian tube cancer, fibrous histiocytoma of bone, gallbladder cancer, gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumors (GIST), germ cell tumor, gestational trophoblastic disease, glioma, hairy cell tumor, hairy cell leukemia, head and neck cancer, thoracic neoplasms, head and neck neoplasms, CNS tumor, primary CNS tumor, heart cancer, hepatocellular cancer, histiocytosis, Hodgkin's lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumors, pancreatic neuroendocrine tumors, Kaposi sarcoma, kidney cancer, Langerhans cell histiocytosis, laryngeal cancer, leukemia, lip and oral cavity cancer, liver cancer, lung cancer, lymphoma, macroglobulinemia, malignant fibrous histiocytoma of bone, osteocarcinoma, melanoma, Merkel cell carcinoma, mesothelioma, metastatic squamous neck cancer, midline tract carcinoma, mouth cancer, multiple endocrine neoplasia syndromes, multiple myeloma, mycosis fungoides, myelodysplastic syndromes, myelodysplastic/myeloproliferative neoplasms, neoplasms by site, neoplasms, myelogenous leukemia, myeloid leukemia, multiple myeloma, myeloproliferative neoplasms, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, neuroblastoma, non-Hodgkin's lymphoma, non-small cell lung cancer, lung neoplasm, pulmonary cancer, pulmonary neoplasms, respiratory tract neoplasms, bronchogenic carcinoma, bronchial neoplasms, oral cancer, oral cavity cancer, lip cancer, oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, papillomatosis, paraganglioma, paranasal sinus and nasal cavity cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pheochromocytoma, pituitary cancer, plasma cell neoplasm, pleuropulmonary blastoma, pregnancy-associated breast cancer, primary central nervous system lymphoma, primary peritoneal cancer, prostate cancer, rectal cancer, colon cancer, colonic neoplasms, renal cell cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, sarcoma, Sezary syndrome, skin cancer, Spitz tumors, small cell lung cancer, small intestine cancer, soft tissue sarcoma, squamous cell carcinoma, squamous neck cancer, stomach cancer, T-cell lymphoma,

testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, transitional cell cancer of the renal pelvis and ureter, unknown primary carcinoma, urethral cancer, uterine cancer, uterine sarcoma, vaginal cancer, vulvar cancer, and Wilms' tumor.

**[00351]** In some embodiments, a hematological cancer (e.g., hematological cancers that are RET-associated cancers) is selected from the group consisting of leukemias, lymphomas (non-Hodgkin's lymphoma), Hodgkin's disease (also called Hodgkin's lymphoma), and myeloma, for instance, acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), acute promyelocytic leukemia (APL), chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), chronic myelomonocytic leukemia (CMML), chronic neutrophilic leukemia (CNL), acute undifferentiated leukemia (AUL), anaplastic large-cell lymphoma (ALCL), prolymphocytic leukemia (PML), juvenile myelomonocytic leukemia (JMML), adult T-cell ALL, AML with trilineage myelodysplasia (AML/TMDS), mixed lineage leukemia (MLL), myelodysplastic syndromes (MDSs), myeloproliferative disorders (MPD), and multiple myeloma (MM). Additional examples of hematological cancers include myeloproliferative disorders (MPD) such as polycythemia vera (PV), essential thrombocytopenia (ET) and idiopathic primary myelofibrosis (IMF/IPF/PMF). In one embodiment, the hematological cancer (e.g., the hematological cancer that is a RET-associated cancer) is AML or CMML.

**[00352]** In some embodiments, the cancer (e.g., the RET-associated cancer) is a solid tumor. Examples of solid tumors (e.g., solid tumors that are RET-associated cancers) include, for example, thyroid cancer (e.g., papillary thyroid carcinoma, medullary thyroid carcinoma), lung cancer (e.g., lung adenocarcinoma, small-cell lung carcinoma), pancreatic cancer, pancreatic ductal carcinoma, breast cancer, colon cancer, colorectal cancer, prostate cancer, renal cell carcinoma, head and neck tumors, neuroblastoma, and melanoma. See, for example, Nature Reviews Cancer, 2014, 14, 173-186.

**[00353]** In some embodiments, the cancer is selected from the group consisting of lung cancer, papillary thyroid cancer, medullary thyroid cancer, differentiated thyroid cancer, recurrent thyroid cancer, refractory differentiated thyroid cancer, multiple endocrine neoplasia type 2A or 2B (MEN2A or MEN2B, respectively), pheochromocytoma, parathyroid hyperplasia, breast cancer, colorectal cancer, papillary renal cell carcinoma, ganglioneuromatosis of the gastroenteric mucosa, and cervical cancer.

**[00354]** In some embodiments, the patient is a human.

**[00355]** Compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof are also useful for treating a RET-associated cancer.

**[00356]** Accordingly, also provided herein is a method for treating a patient diagnosed with or identified as having a RET-associated cancer, e.g., any of the exemplary RET-associated cancers disclosed herein, comprising administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof as defined herein.

**[00357]** Dysregulation of a RET kinase, a RET gene, or the expression or activity or level of any (e.g., one or more) of the same can contribute to tumorigenesis. For example, a dysregulation of a RET

kinase, a RET gene, or expression or activity or level of any of the same can be a translocation, overexpression, activation, amplification, or mutation of a RET kinase, a RET gene, or a RET kinase domain. Translocation can include a gene translocation resulting in the expression of a fusion protein that includes a RET kinase domain and a fusion partner. For example, a fusion protein can have increased kinase activity as compared to a wildtype RET protein. In some embodiments, a mutation in a RET gene can involve mutations in the RET ligand-binding site, extracellular domains, kinase domain, and in regions involved in protein:protein interactions and downstream signaling. In some embodiments, a mutation (e.g., an activating mutation) in a RET gene can result in the expression of a RET kinase having one or more (e.g., two, three, four, five, six, seven, eight, nine, or ten) amino acid substitutions (e.g., one or more amino acid substitutions in the kinase domain (e.g., amino acid positions 723 to 1012 in a wildtype RET protein), a gatekeeper amino acid (e.g., amino acid position 804 in a wildtype RET protein), the P-loop (e.g., amino acid positions 730-737 in a wildtype RET protein), the DFG motif (e.g., amino acid positions 892-894 in a wildtype RET protein), ATP cleft solvent front amino acids (e.g., amino acid positions 758, 811, and 892 in a wildtype RET protein), the activation loop (e.g., amino acid positions 891-916 in a wildtype RET protein), the C-helix and loop preceeding the C-helix (e.g., amino acid positions 768-788 in a wildtype RET protein), and/or the ATP binding site (e.g., amino acid positions 730-733, 738, 756, 758, 804, 805, 807, 811, 881, and 892 in a wildtype RET protein). In some embodiments, a mutation can be a gene amplification of a RET gene. In some embodiments, a mutation (e.g., an activating mutation) in a RET gene can result in the expression of a RET kinase or RET receptor that lacks at least one amino acid (e.g., at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 12, at least 14, at least 16, at least 18, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, or at least 50 amino acids) as compared to a wildtype RET protein. In some embodiments, dysregulation of a RET kinase can be increased expression (e.g., increased levels) of a wildtype RET kinase in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). In some embodiments, a mutation (e.g., an activating mutation) in a RET gene can result in the expression of a RET kinase or RET receptor that has at least one amino acid (e.g., at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 12, at least 14, at least 16, at least 18, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, or at least 50 amino acids) inserted as compared to a wildtype RET protein. In some embodiments, dysregulation of a RET kinase can be increased expression (e.g., increased levels) of a wildtype RET kinase in a mammalian cell (e.g., as compared to a control non-cancerous cell), e.g., due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling. Other dysregulations can include RET mRNA splice variants. In some embodiments, the wildtype RET protein is the exemplary wildtype RET protein described herein.

**[00358]** In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes overexpression of wild-type RET kinase (e.g., leading to autocrine activation). In some embodiments, the dysregulation of a RET gene, a RET kinase protein, or expression or activity or level of any of the same, includes overexpression, activation, amplification, or mutation in a chromosomal segment comprising the RET gene or a portion thereof, including, for example, the kinase domain portion, or a portion capable of exhibiting kinase activity.

**[00359]** In some embodiments, the dysregulation of a RET gene, a RET kinase protein, or expression or activity or level of any of the same, includes one or more chromosome translocations or

inversions resulting in a RET gene fusion. In some embodiments, the dysregulation of a RET gene, a RET kinase protein, or expression or activity or level of any of the same, is a result of genetic translocations in which the expressed protein is a fusion protein containing residues from a non-RET partner protein, and includes a minimum of a functional RET kinase domain.

**[00360]** Non-limiting examples of RET fusion proteins are shown in **Table 1**.

**Table 1. Exemplary RET Fusion Partners and Cancers**

| <b>Fusion Partner</b>                      | <b>Non-limiting Exemplary RET-Associated Cancer(s)</b>                                                                                                                                                                                                                                                                                                                                                  |
|--------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| BCR                                        | Chronic Myelomonocytic Leukemia (CMML)                                                                                                                                                                                                                                                                                                                                                                  |
| CLIP1                                      | Adenocarcinoma                                                                                                                                                                                                                                                                                                                                                                                          |
| KIF5B                                      | NSCLC, Ovarian Cancer, Spitzoid Neoplasms; Lung Adenocarcinoma <sup>3, 4, 14, 28</sup> ; Adenosquamous Carcinomas <sup>15</sup>                                                                                                                                                                                                                                                                         |
| CCDC6 (also called PTC1, D10S170, or H4)   | NSCLC, Colon Cancer, Papillary Thyroid Cancer; Adenocarcinomas; Lung Adenocarcinoma; Metastatic Colorectal Cancer <sup>5</sup> ; Adenosquamous Carcinomas <sup>15</sup> , Breast Cancer <sup>30</sup>                                                                                                                                                                                                   |
| PTC1ex9 (a novel CCDC6 rearrangement)      | Metastatic papillary thyroid cancer <sup>2</sup>                                                                                                                                                                                                                                                                                                                                                        |
| NCOA4 (also called PTC3, ELE1, and RFG)    | Papillary Thyroid Cancer <sup>21</sup> , NSCLC, Colon Cancer, Salivary Gland Cancer, Metastatic Colorectal Cancer <sup>5</sup> ; Lung Adenocarcinoma <sup>15</sup> ; Adenosquamous Carcinomas <sup>15</sup> Diffuse Sclerosing Variant of Papillary Thyroid Cancer <sup>16</sup> , Breast Cancer <sup>30</sup> , Acinic Cell Carcinoma <sup>32</sup> , Mammary Analog Secretory Carcinoma <sup>33</sup> |
| TRIM33 (also called PTC7, RFG7, and TIF1G) | NSCLC, Papillary Thyroid Cancer, Lung Adenocarcinoma <sup>46</sup> , Various <sup>22</sup>                                                                                                                                                                                                                                                                                                              |
| ERC1 (also called ELKS and RAB61P2)        | Papillary Thyroid Cancer, Breast Cancer                                                                                                                                                                                                                                                                                                                                                                 |

| <b>Fusion Partner</b>                | <b>Non-limiting Exemplary RET-Associated Cancer(s)</b>            |
|--------------------------------------|-------------------------------------------------------------------|
| FGFR10P                              | CMML, Primary Myelofibrosis with secondary Acute Myeloid Leukemia |
| MBD1(also known as PCM1)             | Papillary Thyroid Cancer                                          |
| PRKAR1A (also called PTC2)           | Papillary Thyroid Cancer                                          |
| TRIM24 (also called PTC6)            | Papillary Thyroid Cancer                                          |
| KTN1 (also called PTC8 )             | Papillary Thyroid Cancer                                          |
| GOLGA5 (also called PTC5)            | Papillary Thyroid Cancer, Spitzoid Neoplasms                      |
| HOOK3                                | Papillary Thyroid Cancer                                          |
| KIAA1468 (also called PTC9 and RFG9) | Papillary Thyroid Cancer, Lung Adenocarcinoma <sup>8,12</sup>     |
| TRIM27 (also called RFP)             | Papillary Thyroid Cancer                                          |
| AKAP13                               | Papillary Thyroid Cancer                                          |
| FKBP15                               | Papillary Thyroid Cancer, Acute Myeloid Leukemia <sup>46</sup>    |
| SPECC1L                              | Papillary Thyroid Cancer; Thyroid Gland Carcinoma                 |
| TBL1XR1                              | Papillary Thyroid Cancer; Thyroid Gland Carcinoma                 |
| CEP55                                | Diffuse Gastric Cancer <sup>7</sup>                               |
| CUX1                                 | Lung Adenocarcinoma                                               |
|                                      |                                                                   |
| ACBD5                                | Papillary Thyroid Carcinoma                                       |
| MYH13                                | Medullary Thyroid Carcinoma <sup>1</sup>                          |

| <b>Fusion Partner</b>                  | <b>Non-limiting Exemplary RET-Associated Cancer(s)</b>                                                 |
|----------------------------------------|--------------------------------------------------------------------------------------------------------|
| Uncharacterized                        | Inflammatory Myofibroblastic Tumor <sup>6</sup>                                                        |
| PIBF1                                  | Bronchiolus Lung Cell Carcinoma <sup>9</sup>                                                           |
| KIAA1217 (also called SKT)             | Papillary Thyroid Cancer <sup>10, 13</sup><br>Lung Adenocarcinoma <sup>14</sup><br>NSCLC <sup>14</sup> |
| MPRIP                                  | NSCLC <sup>11</sup>                                                                                    |
| HRH4-RET                               | Thyroid Cancer and/or Paillary Thyroid Carcinoma <sup>17</sup>                                         |
| Ria-RET                                | Thyroid Cancer and/or Papillary Thyroid Carcinoma <sup>17</sup>                                        |
| RFG8                                   | Papillary Thyroid Carcinoma <sup>18</sup>                                                              |
| FOXP4                                  | Lung Adenocarcinoma <sup>19</sup>                                                                      |
| MYH10                                  | Infantile Myofibromatosis <sup>20</sup>                                                                |
| HTIF1                                  | Various <sup>22</sup>                                                                                  |
| H4L                                    | Various <sup>22</sup>                                                                                  |
| PTC4 (a novel NCO4/ELE1 rearrangement) | Papillary Thyroid Cancer <sup>23</sup>                                                                 |
| FRMD4A                                 | NSCLC <sup>24</sup>                                                                                    |
| SQSTM1                                 | Papillary Thyroid Carcinoma <sup>25</sup>                                                              |
| AFAP1L2                                | Papillary Thyroid Carcinoma <sup>25</sup>                                                              |
| AFAP1                                  | NSCLC <sup>31</sup>                                                                                    |
| PPFIBP2                                | Papillary Thyroid Carcinoma <sup>25</sup>                                                              |
| EML4                                   | NSCLC                                                                                                  |
| PARD3                                  | NSCLC <sup>27</sup>                                                                                    |
| RASGEF1A                               | Breast Cancer <sup>30</sup>                                                                            |
| TEL                                    | <i>In vitro</i> <sup>34</sup>                                                                          |



| <b>Fusion Partner</b> | <b>Non-limiting Exemplary RET-Associated Cancer(s)</b>     |
|-----------------------|------------------------------------------------------------|
| RUFY1                 | Colorectal Cancer <sup>35</sup>                            |
| OLFM4                 | Small-Bowel Cancer <sup>36</sup>                           |
| UEVLD                 | Papillary Thyroid Carcinoma <sup>29</sup>                  |
| DLG5                  | Non-Anaplastic Thyroid (NAT) Cancer <sup>37</sup>          |
| RRBP1                 | Colon Cancer <sup>38</sup>                                 |
| ANK3                  | Papillary Thyroid Carcinoma <sup>39</sup>                  |
| PICALM                | NSCLC <sup>40</sup>                                        |
| MYO5C                 | NSCLC <sup>41</sup>                                        |
| EPHA5                 | NSCLC <sup>40</sup>                                        |
| RUFY2                 | Lung Cancer <sup>42</sup>                                  |
| KIF13A                | Lung Adenocarcinoma <sup>43</sup> ,<br>NSCLC <sup>45</sup> |
| TNIP1                 | Colorectal Cancer <sup>44</sup>                            |
| SNRNP70               | Colorectal Cancer <sup>44</sup>                            |
| MRLN                  | Thyroid Carcinoma <sup>46</sup>                            |
| LMNA                  | Spitzoid Melanoma <sup>47</sup>                            |
| RUFY3                 | Papillary Thyroid Carcinoma                                |
| TFG                   |                                                            |
| MYO5A                 | Pigmented spindle cell nevus (PSCN) of Reed <sup>48</sup>  |
| ADD3                  | Lung adenocarcinoma <sup>49</sup>                          |
| JMJD1C                | NSCLC <sup>50</sup>                                        |
| RPMS                  |                                                            |
| DOCK1                 |                                                            |
| TAF3                  |                                                            |

<sup>1</sup> Grubbs et al., *J. Clin. Endocrinol. Metab.* 100:788-793, 2015.

- <sup>2</sup> Halkova et al., *Human Pathology* 46:1962-1969, 2015.
- <sup>3</sup> U.S. Patent No. 9,297,011
- <sup>4</sup> U.S. Patent No. 9,216,172
- <sup>5</sup> Le Rolle et al., *Oncotarget*. 6(30):28929-37, 2015.
- <sup>6</sup> Antonescu et al., *Am J Surg Pathol*. 39(7):957-67, 2015.
- <sup>7</sup> U.S. Patent Application Publication No. 2015/0177246.
- <sup>8</sup> U.S. Patent Application Publication No. 2015/0057335.
- <sup>9</sup> Japanese Patent Application Publication No. 2015/109806A.
- <sup>10</sup> Chinese Patent Application Publication No. 105255927A.
- <sup>11</sup> Fang, et al. *Journal of Thoracic Oncology* 11.2 (2016): S21-S22.
- <sup>12</sup> European Patent Application Publication No. EP3037547A1.
- <sup>13</sup> Lee et al., *Oncotarget*. DOI: 10.18632/oncotarget.9137, e-published ahead of printing, 2016.
- <sup>14</sup> Saito et al., *Cancer Science* 107:713-720, 2016.
- <sup>15</sup> Pirker et al., *Transl. Lung Cancer Res*. 4(6):797-800, 2015.
- <sup>16</sup> Joung et al., *Histopathology* 69(1):45-53, 2016.
- <sup>17</sup> PCT Patent Application Publication No. WO 2016/141169.
- <sup>18</sup> Klugbauer et al., *Cancer Res.*, 60(24):7028-32, 2000.
- <sup>19</sup> Bastien et al., *Journal of Molecular Diagnostics*, 18(6):1027, Abstract Number: S120, 2016 Annual Meeting of the Association for Molecular Pathology, Charlotte, NC, 2016.
- <sup>20</sup> Rosenzweig et al., *Pediatr Blood Cancer*, doi:10.1002/pbc.26377, 2016.
- <sup>21</sup> Su et al., *PLoS One*, 11(11): e0165596, 2016.
- <sup>22</sup> U.S. Patent No. 9,487,491.
- <sup>23</sup> Fugazzola et al., *Oncogene*, 13(5):1093-7, 1996.
- <sup>24</sup> Velcheti et al., *J Thorac Oncol.*, 12(2):e15-e16. doi: 10.1016/j.jtho.2016.11.274, 2017.
- <sup>25</sup> Kato et al, *Clin Cancer Res*. 2017 Apr 15;23(8):1988-1997. doi: 10.1158/1078-0432.CCR-16-1679. Epub 2016 Sep 28.
- <sup>26</sup> Drilon, Alexander, et al. "A phase 1/1b study of RXDX-105, an oral RET and BRAF inhibitor, in patients with advanced solid tumors." Aug 8 (2016): 7.
- <sup>27</sup> Sabari et al., *Oncoscience*, Advance Publications, www.impactjournals.com/oncoscience/files/papers/1/345/345.pdf, 2017.

- <sup>28</sup> U.S. Patent Application Publication No. 2017/0014413.
- <sup>29</sup> Lu et al., *Oncotarget*, 8(28):45784-45792, doi: 10.18632/oncotarget.17412, 2017.
- <sup>30</sup> Hirshfield et al., *Cancer Research*, (February 2017) Vol. 77, No. 4, Supp. 1. Abstract Number: P3-07-02. Meeting Info: 39th Annual CTRC-AACR San Antonio Breast Cancer Symposium. San Antonio, TX, United States. 06 Dec 2016-10 Dec 2016.
- <sup>31</sup> Morgensztern et al., *Journal of Thoracic Oncology*, (January 2017) Vol. 12, No. 1, Supp. 1, pp. S717-S718, Abstract Number: P1.07-035, Meeting Info: 17th World Conference of the International Association for the Study of Lung Cancer, IASLC 2016. Vienna, Austria. 04 Dec 2016.
- <sup>32</sup> Dogan et al., *Laboratory Investigation*, (February 2017) Vol. 97, Supp. 1, pp. 323A. Abstract Number: 1298, Meeting Info: 106th Annual Meeting of the United States and Canadian Academy of Pathology, USCAP 2017. San Antonio, TX, United States.
- <sup>33</sup> Dogan et al., *MODERN PATHOLOGY*, Vol. 30, Supp. [2], pp. 323A-323A. MA 1298, 2017.
- <sup>34</sup> PCT Patent Application Publication No. WO 2017/146116.
- <sup>35</sup> PCT Patent Application Publication No. WO 2017/122815.
- <sup>36</sup> Reeser et al., *J. Mol. Diagn.*, 19(5):682-696, doi: 10.1016/j.jmoldx.2017.05.006, 2017.
- <sup>37</sup> Ibrahimasic et al., *Clin. Cancer Res.*, doi: 10.1158/1078-0432.CCR-17-1183, 2017.
- <sup>38</sup> Kloosterman et al., *Cancer Res.*, 77(14):3814-3822. doi: 10.1158/0008-5472.CAN-16-3563, 2017.
- <sup>39</sup> Chai et al., *Oncology Reports*, 35(2): 962-970. doi: 10.3892/or.2015.4466, 2015.
- <sup>40</sup> Gautschi et al. *Journal of Clinical Oncology*, 35(13) 1403-1410. doi: 10.1200/JCO.2016.70.9352, 2017.
- <sup>41</sup> Lee et al. *Annals of Oncology*, 28(2), 292-297. doi: 10.1093/annonc/mdw559, 2016.
- <sup>42</sup> Zheng et al. *Nature Medicine*, 20(12), 1479-1484. doi: 10.1038/nm.3729, 2014.
- <sup>43</sup> Zhang et al. *Lung Cancer*, 118, 27-29. doi: 10.1016/j.lungcan.2017.08.019, 2018.
- <sup>44</sup> Morano et al. *Molecular Cancer Therapeutics*, (January 2018) Vol. 17, No. 1, Supp. Supplement 1. Abstract Number: B049. Meeting Info: AACR-NCI-EORTC International Conference: Molecular Targets and Cancer Therapeutics 2017.
- <sup>45</sup> Wang et al. *Journal of Thoracic Oncology*, (November 2017) Vol. 12, No. 11, Supp. Supplement 2, pp. S2105. Abstract Number: P2.02-018. Meeting Info: 18th World Conference on Lung Cancer of the International Association for the Study of Lung Cancer, IASLC 2017. Yokohama, Japan. 15 Oct 2017-18 Oct 2017.
- <sup>46</sup> Gao et al. *Cell Reports*, 23(1), 227-238. doi: 10.1016/j.celrep.2018.03.050, 2018.
- <sup>47</sup> U.S. Patent Application Publication No. 2016/0010068.
- <sup>48</sup> VandenBoom, et al. *Am. J. Surg. Pathol.* 42(8): 1042-1051, 2018. doi: 10.1097/PAS.0000000000001074

<sup>49</sup> Cao, et al. *Onco. Targets. Ther.* 2018(11):2637-2646, 2018. doi: 10.2147/OTT.S155995

<sup>50</sup> Luo, et al. *Int. J. Cancer*, 2018. epub ahead of print. doi: 10.1002/ijc.31542

**[00361]** In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes one or more deletions (e.g., deletion of an amino acid at position 4), insertions, or point mutation(s) in a RET kinase. In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes a deletion of one or more residues from the RET kinase, resulting in constitutive activity of the RET kinase domain.

**[00362]** In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes at least one point mutation in a RET gene that results in the production of a RET kinase that has one or more amino acid substitutions, insertions, or deletions as compared to the wild-type RET kinase (see, for example, the point mutations listed in **Table 2**).

**Table 2. RET Kinase Protein Amino Acid Substitutions/Insertions/Deletions<sup>A</sup>**

|                                                   |
|---------------------------------------------------|
| Amino acid position 2                             |
| Amino acid position 3                             |
| Amino acid position 4                             |
| Amino acid position 5                             |
| Amino acid position 6                             |
| Amino acid position 7                             |
| Amino acid position 8                             |
| Amino acid position 11                            |
| Amino acid position 12                            |
| Amino acid position 13                            |
| Amino acid position 20                            |
| Amino acid position 32 (e.g., S32L)               |
| Amino acid position 34 (e.g., D34S)               |
| Amino acid position 40 (e.g., L40P)               |
| Amino acid position 45 (e.g., A45A) <sup>39</sup> |
| Amino acid position 56 (e.g., L56M) <sup>30</sup> |
| Amino acid position 64 (e.g., P64L)               |
| Amino acid position 67 (e.g., R67H)               |

|                                                                                                |
|------------------------------------------------------------------------------------------------|
| Amino acid position 77 (e.g., R77C) <sup>65</sup>                                              |
| Amino acid position 114 (e.g., R114H)                                                          |
| Amino acid position 136 (e.g., glutamic acid to stop codon)                                    |
| Amino acid position 145 (e.g., V145G)                                                          |
| Amino acid position 177 (e.g., R177L) <sup>67</sup>                                            |
| Amino acid position 180 (e.g., arginine to stop codon)                                         |
| Amino acid position 200                                                                        |
| Amino acid position 270 (e.g., P270L) <sup>65</sup>                                            |
| Amino acid position 278 (e.g., T278N) <sup>57</sup>                                            |
| Amino acid position 292 (e.g., V292M)                                                          |
| Amino acid position 294                                                                        |
| Amino acid position 321 (e.g., G321R)                                                          |
| Amino acid position 330 (e.g., R330Q)                                                          |
| Amino acid position 338 (e.g., T338I)                                                          |
| Amino acid position 360 (e.g., R360W)                                                          |
| Amino acid position 373 (e.g., alanine to frameshift)                                          |
| Δ Amino acid positions 378 – 385 with insertion of one amino acid (e.g., D378 – G385>E)        |
| Amino acid position 393 (e.g., F393L)                                                          |
| Amino acid position 423 (e.g., G423R) <sup>27</sup>                                            |
| Amino acid position 428 (e.g., E428K) <sup>57</sup>                                            |
| Amino acid position 432 (e.g., A432A <sup>39</sup> )                                           |
| Amino acid position 446 (e.g., G446R) <sup>28</sup>                                            |
| Δ Amino acid positions 505-506 (6-Base Pair In-Frame Germline Deletion in Exon 7) <sup>3</sup> |
| Amino acid position 510 (e.g., A510V)                                                          |
| Amino acid position 511 (e.g., E511K)                                                          |
| Amino acid position 513 (e.g., G513D) <sup>7*</sup>                                            |
| Amino acid position 515 (e.g., C515S, C515W <sup>4</sup> )                                     |

|                                                                                                       |
|-------------------------------------------------------------------------------------------------------|
| Amino acid position 525 (e.g., R525W) <sup>7*</sup>                                                   |
| Amino acid position 531 (e.g., C531R, or 9 base pair duplication <sup>2</sup> )                       |
| Amino acid position 532 (e.g., duplication) <sup>2</sup>                                              |
| Amino acid position 533 (e.g., G533C, G533S)                                                          |
| Amino acid position 534 (e.g., L534L) <sup>6</sup>                                                    |
| Amino acid position 550 (e.g., G550E)                                                                 |
| Amino acid position 591 (e.g., V591I)                                                                 |
| Amino acid position 593 (e.g., G593E)                                                                 |
| Amino acid position 595 (e.g., E595D and E595A) <sup>18</sup>                                         |
| Amino acid position 600 (e.g., R600Q)                                                                 |
| Amino acid position 602 (e.g., I602V) <sup>6</sup>                                                    |
| Amino acid position 603 (e.g., K603Q, K603E <sup>2</sup> )                                            |
| Amino acid position 606 (e.g., Y606C)                                                                 |
| Amino acid position 609 (e.g., C609Y, C609S, C609G, C609R, C609F, C609W, C609C <sup>32</sup> )        |
| Amino acid position 611 (e.g., C611R, C611S, C611G, C611Y, C611F, C611W)                              |
| Amino acid position 616 (e.g., E616Q) <sup>23</sup>                                                   |
| $\Delta$ Amino acid position 616 <sup>64</sup>                                                        |
| Amino acid position 618 (e.g., C618S, C618Y, C618R, C618G, C618F, C618W, stop <sup>56</sup> )         |
| Amino acid position 619 (e.g., F619F)                                                                 |
| Amino acid position 620 (e.g., C620S, C620W, C620R, C620G, C620L, C620Y, C620F, C620A <sup>47</sup> ) |
| Amino acid position 622 (e.g., P622L) <sup>68</sup>                                                   |
| Amino acid position 623 (e.g., E623K)                                                                 |
| Amino acid position 624 (e.g., D624N)                                                                 |
| Amino acid position 630 (e.g., C630A, C630R, C630S, C630Y, C630F, C630W)                              |
| $\Delta$ Amino acid position 630 <sup>56</sup>                                                        |

|                                                                                                                                                                                                                                                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Amino acid position 631 (e.g., D631N, D631Y, D631A, D631G, D631V, D631E, )                                                                                                                                                                         |
| Δ Amino acid position 631 <sup>69</sup>                                                                                                                                                                                                            |
| Amino acid positions 631-633>V (i.e., residues 631-633 are replaced with a single valine residue)                                                                                                                                                  |
| Amino acid positions 631-633>A (i.e., residues 631-633 are replaced with a single alanine residue)                                                                                                                                                 |
| Amino acid positions 631-633>E (i.e., residues 631-633 are replaced with a single glutamic acid residue)                                                                                                                                           |
| Δ Amino acid positions 631-633 (e.g., D631 – L633)                                                                                                                                                                                                 |
| Δ Amino acid positions 631-634 (e.g., D631-C634)                                                                                                                                                                                                   |
| Amino acid position 632 (e.g., E632K, E632G <sup>5, 11</sup> , E632V <sup>62</sup> , 632 to frameshift <sup>47</sup> )                                                                                                                             |
| Δ Amino acid positions 632-633 (e.g., E632 – L633 in either the somatic cells, or a 6-Base Pair In-Frame Germline Deletion in Exon 11 <sup>9</sup> )                                                                                               |
| Amino acid positions 632-639>HR (i.e., residues 632-639 are replaced with two residues, histidine and arginine)                                                                                                                                    |
| Amino acid position 633 (e.g., L633R <sup>62</sup> , 9 base pair duplication <sup>2</sup> )                                                                                                                                                        |
| Amino acid position 634 (e.g., C634W, C634Y, C634S, C634R, C634F, C634G, C634L, C634A, or C634T, a 9 base pair deletion <sup>62</sup> , a 9 base pair duplication <sup>56</sup> , or a 12 base pair duplication <sup>2</sup> ) (e.g., causing MTC) |
| Δ Amino acid position 634 <sup>56</sup>                                                                                                                                                                                                            |
| Amino acid position 632/633/634 (E632V/L633R/634 9 base pair deletion) <sup>62</sup>                                                                                                                                                               |
| Amino acid position 635 (e.g., R635G or an insertion ELCR <sup>2</sup> )                                                                                                                                                                           |
| Amino acid position 636 (e.g., T636P <sup>2</sup> , T636M <sup>4</sup> )                                                                                                                                                                           |
| Amino acid position 638 (e.g., isoleucine to frameshift <sup>47</sup> )                                                                                                                                                                            |
| Amino acid position 640 (e.g., A640G)                                                                                                                                                                                                              |
| Amino acid position 634/640 (e.g., C634R/A640G) <sup>56</sup>                                                                                                                                                                                      |
| Amino acid position 641 (e.g., A641S, A641T <sup>8</sup> )                                                                                                                                                                                         |

|                                                                                            |
|--------------------------------------------------------------------------------------------|
| Amino acid position 634/641 (e.g., C634S/A641S) <sup>56</sup>                              |
| Amino acid position 639/641 (e.g., A639G/A641R) <sup>56</sup>                              |
| Amino acid position 644 (e.g., T644M) <sup>59</sup>                                        |
| Amino acid position 648 (e.g., V648I)                                                      |
| Amino acid position 649 (e.g., S649L) <sup>28</sup>                                        |
| Amino acid position 661 (e.g., H661H) <sup>6</sup>                                         |
| Amino acid position 664 (e.g., A664D)                                                      |
| Amino acid position 665 (e.g., H665Q)                                                      |
| Amino acid position 666 (e.g., K666E, K666M, K666N, K666R)                                 |
| Amino acid position 675 (T675T, silent nucleotide change) <sup>18</sup>                    |
| Amino acid position 679 (e.g., P679P) <sup>6</sup>                                         |
| Amino acid position 680 (e.g., A680T, alanine to frameshift) <sup>6</sup>                  |
| Amino acid position 686 (e.g., S686N)                                                      |
| Amino acid position 689 (e.g., S689T) <sup>18</sup>                                        |
| Amino acid position 691 (e.g., G691S)                                                      |
| Amino acid position 694 (e.g., R694Q)                                                      |
| Amino acid position 700 (e.g., M700L)                                                      |
| Amino acid position 706 (e.g., V706M, V706A)                                               |
| Amino acid position 713 splice variant (e.g., E713K (e.g., a splice variant)) <sup>6</sup> |
| Amino acid position 714 (e.g., D714Y) <sup>57</sup>                                        |
| Amino acid position 727 (e.g., G727E) <sup>6</sup>                                         |
| Amino acid position 732 (e.g., E732K) <sup>20</sup>                                        |
| Amino acid position 734 (e.g., E734K) <sup>48</sup>                                        |
| Amino acid position 736 (e.g., G736R) <sup>6</sup>                                         |
| Amino acid position 738 (e.g., V738V) <sup>6</sup>                                         |
| Amino acid position 742 (e.g., T742M) <sup>51</sup>                                        |
| Amino acid position 748 (e.g., G748C)                                                      |



|                                                                                                                                                                                         |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Amino acid position 749 (e.g., R749T <sup>36</sup> )                                                                                                                                    |
| Amino acid position 750 (e.g., A750P, A750G <sup>6</sup> )                                                                                                                              |
| Amino acid position 752 (e.g., Y752Y) <sup>6</sup>                                                                                                                                      |
| Amino acid position 751 (e.g., G751G) <sup>6</sup>                                                                                                                                      |
| Amino acid position 762 (e.g., E762Q <sup>36</sup> )                                                                                                                                    |
| Amino acid position 765 (e.g., S765P, S765F)                                                                                                                                            |
| Amino acid position 766 (e.g., P766S, P766M <sup>6</sup> )                                                                                                                              |
| Amino acid position 768 (e.g., E768Q, E768D, E768N <sup>46</sup> )                                                                                                                      |
| Amino acid position 769 (e.g., L769L <sup>6</sup> )                                                                                                                                     |
| Amino acid position 770 (e.g., R770Q)                                                                                                                                                   |
| Amino acid position 771 (e.g., D771N)                                                                                                                                                   |
| Amino acid position 777 (e.g., N777S)                                                                                                                                                   |
| Amino acid position 778 (e.g., V778I)                                                                                                                                                   |
| Amino acid position 781 (e.g., Q781R)                                                                                                                                                   |
| Amino acid position 788 (e.g., I788I <sup>32</sup> )                                                                                                                                    |
| Amino acid position 790 (e.g., L790F)                                                                                                                                                   |
| Amino acid position 768/790 (e.g., E768D/L790T) <sup>40</sup>                                                                                                                           |
| Amino acid position 791 (e.g., Y791F, Y791N <sup>24</sup> )                                                                                                                             |
| Amino acid position 634/791 (e.g., C634Y/Y791F) <sup>55</sup>                                                                                                                           |
| Amino acid position 790/791 (e.g., L790F/Y791F) <sup>55</sup>                                                                                                                           |
| Amino acid position 802                                                                                                                                                                 |
| Amino acid position 804 (e.g., V804L <sup>15, 16</sup> , V804M <sup>15, 16</sup> , V804E <sup>12</sup> ) (e.g., causing MTC)                                                            |
| Amino acid position 778/804 <sup>50</sup> (e.g., V778I/V804M <sup>54</sup> )                                                                                                            |
| Amino acid position 781/804 (e.g., Q781R/V804M) <sup>41</sup>                                                                                                                           |
| Amino acid position 805 (e.g., E805K)                                                                                                                                                   |
| Amino acid position 804/805 (e.g., V804M/E805K) <sup>17</sup>                                                                                                                           |
| Amino acid position 806 (e.g., Y806F, Y806S <sup>12</sup> , Y806G, Y806C <sup>2, 12, 14</sup> , Y806E <sup>14</sup> , Y806H <sup>12</sup> , Y806N <sup>12</sup> , Y806Y <sup>32</sup> ) |

|                                                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------|
| Amino acid position 804/806 (e.g., V804M/Y806C) <sup>38</sup>                                                             |
| Amino acid position 810 (e.g., G810R <sup>12</sup> , G810S <sup>12</sup> , G810A <sup>13</sup> , G810C, G810V, and G810D) |
| Amino acid position 818 (e.g., E818K)                                                                                     |
| Amino acid position 819 (e.g., S819I)                                                                                     |
| Amino acid position 820 (e.g., R820L) <sup>57</sup>                                                                       |
| Amino acid position 823 (e.g., G823E)                                                                                     |
| Amino acid position 826 (e.g., Y826M, Y826S) <sup>10</sup>                                                                |
| Amino acid position 828 (e.g., G828R) <sup>57</sup>                                                                       |
| Amino acid position 833 (e.g., R833C)                                                                                     |
| Amino acid position 836 (e.g., S836S) <sup>19</sup>                                                                       |
| Amino acid position 841 (e.g., P841L, P841P)                                                                              |
| Amino acid position 843 (e.g., E843D)                                                                                     |
| Amino acid position 844 (e.g., R844W, R844Q, R844L)                                                                       |
| Amino acid position 845 (e.g., A845A) <sup>63</sup>                                                                       |
| Amino acid position 848 (e.g., M848T)                                                                                     |
| Amino acid position 852 (e.g., I852M)                                                                                     |
| Amino acid position 853 (e.g., S853T) <sup>57</sup>                                                                       |
| Amino acid position 865 (e.g., L865V) <sup>12</sup>                                                                       |
| Amino acid position 866 (e.g., A866W) <sup>33</sup>                                                                       |
| Amino acid position 867 (e.g., E867K) <sup>37</sup>                                                                       |
| Amino acid position 870 (e.g., L870F) <sup>12</sup>                                                                       |
| Amino acid position 873 (e.g., R873W, R873Q <sup>42</sup> )                                                               |
| Amino acid position 876 (e.g., A876V)                                                                                     |
| Amino acid position 881 (e.g., L881V)                                                                                     |
| Amino acid position 882                                                                                                   |
| Amino acid position 883 (e.g., A883F, A883S, A883T, A883Y <sup>53</sup> , A883V)                                          |
| Amino acid position 884 (e.g., E884K, E884V <sup>35</sup> )                                                               |

|                                                                                                     |
|-----------------------------------------------------------------------------------------------------|
| Amino acid position 886 (e.g., R886W)                                                               |
| Amino acid position 891 (e.g., S891A, S891S <sup>32</sup> , S891L <sup>35</sup> )                   |
| Amino acid position 893 (e.g., F893L) <sup>42</sup>                                                 |
| Amino acid position 894 (e.g., G894S) <sup>43</sup>                                                 |
| Amino acid position 897 (e.g., R897Q, R897P)                                                        |
| Amino acid position 898 (e.g., D898V, D898Y <sup>66</sup> )                                         |
| Δ Amino acid position 898                                                                           |
| Δ Amino acid positions 898-902 <sup>58</sup>                                                        |
| Δ Amino acid positions 899-902 <sup>47</sup>                                                        |
| Δ Amino acid positions 898-901 <sup>47</sup>                                                        |
| Δ Amino acid positions 632-633/Δ Amino acid positions 898-901 <sup>47</sup>                         |
| Amino acid position 900 (e.g., Y900F) <sup>22</sup>                                                 |
| Amino acid position 901 (e.g., E901K)                                                               |
| Amino acid position 904 (e.g., S904F, S904S, S904C <sup>2</sup> , S904T <sup>57</sup> )             |
| Amino acid position 691/904 (e.g., G691S/S904S) <sup>49</sup>                                       |
| Amino acid position 804/904 (e.g., V804M/S904C) <sup>38</sup>                                       |
| Amino acid position 905 (e.g., Y905F) <sup>22</sup>                                                 |
| Amino acid position 907 (e.g., K907E, K907M)                                                        |
| Amino acid position 908 (e.g., R908K)                                                               |
| Amino acid position 911 (e.g., G911D, G911G (e.g., a splice variant) <sup>6</sup> )                 |
| Amino acid position 912 (e.g., R912P, R912Q)                                                        |
| Amino acid position 918 (e.g., M918T <sup>2</sup> , M918V, M918L <sup>6</sup> ) (e.g., causing MTC) |
| Amino acid position 591/918 (e.g., V591I/M918T) <sup>61</sup>                                       |
| Amino acid position 620/918 (e.g., C620F/M918T) <sup>47</sup>                                       |
| Amino acid position 891/918 (e.g., S891A/M918T) <sup>47</sup>                                       |
| Δ Amino acid position 898-901/M918T <sup>47</sup>                                                   |

|                                                                         |
|-------------------------------------------------------------------------|
| Amino acid position 919 (e.g., A919V, A919P <sup>52</sup> )             |
| Amino acid position 768/919 <sup>54</sup>                               |
| Amino acid position 921 (e.g., E921K, E921D)                            |
| Amino acid position 911/918/921 (e.g., G911E/M918T/E921K) <sup>61</sup> |
| Amino acid position 922 (e.g., S922P, S922Y)                            |
| Amino acid position 924 (e.g., F924S) <sup>6</sup>                      |
| Amino acid position 930 (e.g., T930M)                                   |
| Amino acid position 961 (e.g., F961L)                                   |
| Amino acid position 972 (e.g., R972G)                                   |
| Amino acid position 973 (e.g., P973T) <sup>57</sup>                     |
| Amino acid position 977 (e.g., S977R) <sup>37</sup>                     |
| Amino acid position 981 (e.g., Y981F) <sup>22</sup>                     |
| Amino acid position 982 (e.g., R982C) <sup>70</sup>                     |
| Amino acid position 634/691/982 (e.g., C634R/G691S/R982C) <sup>45</sup> |
| Amino acid position 1009 (e.g., M1009V)                                 |
| Amino acid position 1015 (e.g., Y1015F) <sup>22</sup>                   |
| Amino acid position 1017 (e.g., D1017N)                                 |
| Amino acid position 1041 (e.g., V1041G)                                 |
| Amino acid position 1047 (e.g., P1047S) <sup>65</sup>                   |
| Amino acid position 1051 (e.g., A1051T) <sup>57</sup>                   |
| Δ Amino acid position 1059 <sup>57</sup>                                |
| Amino acid position 1064 (e.g., M1064T)                                 |
| Amino acid position 1096 (e.g., Y1096F) <sup>21</sup>                   |
| Amino acid position 1105 (e.g., A1105V) <sup>57</sup>                   |
| Amino acid position 1109 (e.g., M1109T) <sup>34</sup>                   |
| RET+3 <sup>1</sup>                                                      |
| (In-Frame Deletion in Exons 6 and 11) <sup>25</sup>                     |

|                                                                      |
|----------------------------------------------------------------------|
| (3bp In-Frame Deletion in Exon 15) <sup>26</sup>                     |
| Nucleotide position 2136+2 (e.g., 2136+2T>G) <sup>29</sup>           |
| (del632-636 ins6) <sup>31</sup>                                      |
| Amino acid positions 791 and 852 (e.g., Y791F + I852M) <sup>31</sup> |
| Amino acid positions 634 and 852 (e.g., C634R + I852M) <sup>31</sup> |
| c.1893_1895del <sup>44</sup>                                         |

<sup>A</sup> The RET kinase mutations shown may be activating mutations and/or confer increased resistance of the RET kinase to a RET kinase inhibitor and/or a multi-kinase inhibitor (MKI), e.g., as compared to a wildtype RET kinase.

<sup>1</sup> U.S. Patent Application Publication No. 2014/0272951.

<sup>2</sup> Krampitz et al., *Cancer* 120:1920-1931, 2014.

<sup>3</sup> Latteyer, et al., *J. Clin. Endocrinol. Metab.* 101(3):1016-22, 2016.

<sup>4</sup> Silva, et al. *Endocrine* 49.2:366-372, 2015.

<sup>5</sup> Scollo, et al., *Endocr. J.* 63(1):87-91, 2016.

<sup>6</sup> Jovanovic, et al., *Prilozi* 36(1):93-107, 2015.

<sup>7</sup> Qi, et al., *Oncotarget*. 6(32):33993-4003, 2015. \*R525W and G513D appear to act in combination with S891A to enhance oncogenic activity.

<sup>8</sup> Kim, et al. *ACTA ENDOCRINOLOGICA-BUCHAREST* 11.2, 189-194, 2015.

<sup>9</sup> Cecchirini, et al. *Oncogene*, 14, 2609-2612, 1997.

<sup>10</sup> Karrasch, et al. *Eur. Thyroid J.*, 5(1):73-7, 2016.

<sup>11</sup> Scollo et al., *Endocr. J.* 63:87-91, 2016.

<sup>12</sup> PCT Patent Application Publication No. WO 2016/127074.

<sup>13</sup> Huang et al., *Mol. Cancer Ther.*, 2016 Aug 5. pii: molcanther.0258.2016. [Epub ahead of print].

<sup>14</sup> Carlomagno, et al., *Endocr. Rel. Cancer* 16(1):233-41, 2009.

<sup>15</sup> Yoon et al., *J. Med. Chem.* 59(1):358-73, 2016.

<sup>16</sup> U.S. Patent No. 8,629,135.

<sup>17</sup> Cranston, et al., *Cancer Res.* 66(20):10179-87, 2006.

<sup>18</sup> Kheiroddin et al., *Clin. Lab.* 62(5):871-6, 2016.

<sup>19</sup> Ceolin et al., *PLoS One*. 11(2): e0147840, doi: 10.1371/journal.pone.0147840, 2016.

- <sup>20</sup> Mamedova et al., Summer Undergraduate Research Programs (SURP) Student Abstracts, University of Oklahoma Health Sciences Center, 2016.
- <sup>21</sup> Liu et al., *J. Biol. Chem.*, 271(10): 5309-12, 1995.
- <sup>22</sup> Kato et al., *Cancer Res.*, 62: 2414-22, 2002.
- <sup>23</sup> Grey et al., *Endocrine Pathology*, doi:10.1007/s12022-016-9451-6, 2016.
- <sup>24</sup> De Almeida et al., *Endocrine Reviews*, 2016, Vol. 37, No. 2, Supp. Supplement 1. Abstract Number: SUN-068; 98<sup>th</sup> Annual Meeting and Expo of the Endocrine Society, ENDO 2016. Boston, MA, US. 01 Apr 2016-04 Apr 2016.
- <sup>25</sup> Vanden et al., *Annals of Oncology*, 2016, Vol. 27, Supp. Supplement 6. Abstract Number: 427PD; 41<sup>st</sup> European Society for Medical Oncology Congress, ESMP 2016. Copenhagen, Denmark. 07 Oct 2016-11 Oct 2016.
- <sup>26</sup> Romei et al., *European Thyroid Journal* (August 2016) Vol. 5, Supp. Supplement 1, pp. 75; 39<sup>th</sup> Annual Meeting of the European Thyroid Association, ETA 2016. Copenhagen, Denmark. 03 Sep 2016-06 Sep 2016.
- <sup>27</sup> Lee et al., *Oncotarget*, 8(4): 6579–6588, doi: 10.18632/oncotarget.14172, 2017.
- <sup>28</sup> Zhang et al., *Laboratory Investigation*, (February 2017) Vol. 97, Supp. 1, pp. 209A. Abstract Number: 840, Meeting Info: 106th Annual Meeting of the United States and Canadian Academy of Pathology, USCAP 2017. San Antonio, TX, United States.
- <sup>29</sup> Borecka et al., *European Journal of Cancer*, (July 2016) Vol. 61, No. 1, pp. S26, Abstract Number: 162, Meeting Info: 24th Biennial Congress of the European Association for Cancer Research, EACR 2016. Manchester, United Kingdom.
- <sup>30</sup> Corsello et al., *Endocrine Reviews*, (JUN 2014) Vol. 35, No. 3, Suppl. S, pp. SUN-0322, Meeting Info.: 96th Annual Meeting and Expo of the Endocrine-Society, Chicago, IL, USA, June 21-24, 2014.
- <sup>31</sup> Gazizova et al., *Endocrine Reviews*, (JUN 2014) Vol. 35, No. 3, Suppl. S, pp. SAT-0304, Meeting Info.: 96th Annual Meeting and Expo of the Endocrine-Society, Chicago, IL, USA, June 21-24, 2014.
- <sup>32</sup> Sromek et al., *Endocr Pathol.*, doi: 10.1007/s12022-017-9487-2, 2017.
- <sup>33</sup> U.S. Patent Application Publication No. 2017/0267661.
- <sup>34</sup> Davila et. al., *Rare Tumors*, 2017; 9(2): 6834. doi:10.4081/rt.2017.6834.
- <sup>35</sup> U.S. Patent Application Publication No. 2018/0009818.
- <sup>36</sup> PCT Patent Application Publication No. WO 2017/197051
- <sup>37</sup> European Patent Application Publication No. 3271848
- <sup>38</sup> Roskoski and Sadeghi-Nejad, *Pharmacol. Res.*, 128, 1-17. doi: 10.1016/j.phrs.2017.12.021, 2018.
- <sup>39</sup> Kaczmarek-Ryś, et al. *Endocrine-related cancer* 25(4): 421-436. doi: 10.1530/ERC-17-0452, 2018.

- <sup>40</sup> Raue, et al. *J. Clin Endocrinol Metab*, 103(1): 235-243. doi: 10.1210/jc.2017-01884, 2018.
- <sup>41</sup> Nakao, et al. *Head and Neck*, 35: E363-E368. doi: 10.1002/hed.23241, 2013.
- <sup>42</sup> Attié, et al. *Human Molecular Genetics* 4(8): 1381-1386. doi: 10.1093/hmg/4.8.1381, 1995.
- <sup>43</sup> Fitze, et al. *Lancet*, 393(9313): 1200-1205. doi: 10.1016/S0140-6736(02)08218-1, 2002.
- <sup>44</sup> Weng, et al. *Zhonghua Nei Ke Za Zhi*, 57(2):134-137. doi: 10.3760/cma.j.issn.0578-1426.2018.02.010, 2018.
- <sup>45</sup> Chen, et al. *Medical Journal of Chinese People's Liberation Army* 38.4 (2013): 308-312.
- <sup>46</sup> Gudernova, et al. *eLife*, 6:e21536. doi: 10.7554/eLife.21536, 2017.
- <sup>47</sup> Romei, et al. *Oncotarget*, 9(11): 9875-9884. doi: 10.18632/oncotarget.23986, 2018.
- <sup>48</sup> Plaza-Menacho. *Endocr Relat Cancer*, 25(2):T79-T90. doi: 10.1530/ERC-17-0354, 2017.
- <sup>49</sup> Guerin, et al. *Endocr Relat Cancer*, 25(2):T15-T28. doi: 10.1530/ERC-17-0266, 2017.
- <sup>50</sup> Roy et al. *Oncologist*, 18(10): 1093-1100. doi: 10.1634/theoncologist.2013-0053, 2013
- <sup>51</sup> U.S. Patent Application Publication No. 2017/0349953
- <sup>52</sup> Santoro, et al. *Endocrinology*, 145(12), 5448-5451, 2004. doi: 10.1210/en.2004-0922
- <sup>53</sup> U.S. Patent No. 9,006,256
- <sup>54</sup> Yeganeh, et al. *Asian Pac J Cancer Prev*, 16(6), 2107-17. doi: 10.7314/APJCP.2015.16.6.2107
- <sup>55</sup> Mulligan, L. M, *Nature Reviews Cancer*, 14(3), 173, 2014, doi: 10.1038/nrc3680
- <sup>56</sup> Arighi, et al, *Cytokine & Growth Factor Reviews*, 16(4-5), 441-467, 2005. doi: 10.1016/j.cytogfr.2005.05.010
- <sup>57</sup> Dabir, et al, *Journal of Thoracic Oncology*, 9(9), 1316-1323, 2014. doi: 10.1097/JTO.0000000000000234
- <sup>58</sup> Uchino, et al, *Cancer Science*, 90(11), 1231-1237, 1999. doi: 10.1111/j.1349-7006.1999.tb00701.x
- <sup>59</sup> Krampitz. *Cancer*, 120(13), 1920-1931, 2014: 10.1002/cncr.28661
- <sup>60</sup> Jhiang et al, *Thyroid* 6(2), 1996. doi: 10.1089/thy.1996.6.115
- <sup>61</sup> Dvořáková, et al, *Thyroid*, 16(3), 311-316, 2006. doi: 10.1089/thy.2006.16.311
- <sup>62</sup> Severskaya et al, *Genomics Transcriptomics Proteomics*, 40(3) 425-435.
- <sup>63</sup> Elisei, et al, *Journal of Genetic Syndromes & Gene Therapy*, 5(1), 1, 2014. doi: 10.4172/2157-7412.1000214
- <sup>64</sup> Ahmed et al, *The Journal of Molecular Diagnostics*, 7(2), 283-288, 2005. doi: 10.1016/S1525-1578(10)60556-9

- <sup>65</sup> Oliveira, et al. *J. Exp. Clin. Cancer Res.* 37(84), 2018. doi: 10.1186/s13046-018-0746-y
- <sup>66</sup> Yi, et al. *Case Rep. Endocrinol.* 2018:8657314, 2018. doi: 10.1155/2018/8657914
- <sup>67</sup> Huang, et al. *Cell.* 173(2): 355-370, 2018. doi: 10.1016/j.cell.2018.03.039
- <sup>68</sup> Bosic, et al. *Pathology.* 50(3):327-332, 2018. doi: 10.1016/j.pathol.2017.10.011
- <sup>69</sup> Yao, et al. *Zhonghua Yi Xue Za Zhi.* 87(28):1962-1965, 2007. PMID: 17923033
- <sup>70</sup> Quintela-Fandino, et al. *Mol. Oncol.* 8(8):1719-1728, 2014. doi: 10.1016/j.molonc.2014.07.005

**[00363]** In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes at least one point mutation in a RET gene that results in the production of a RET kinase that has one or more amino acid substitutions, insertions, or deletions as compared to the wild-type RET kinase (see, for example, the point mutations listed in **Table 2a**).

**[00364]** **Table 2a RET Kinase Protein Amino Acid Substitutions/Insertions/Deletions<sup>A</sup>**

|                                       |
|---------------------------------------|
| Amino acid position 20                |
| Amino acid position 32 (e.g., S32L)   |
| Amino acid position 34 (e.g., D34S)   |
| Amino acid position 40 (e.g., L40P)   |
| Amino acid position 64 (e.g., P64L)   |
| Amino acid position 67 (e.g., R67H)   |
| Amino acid position 114 (e.g., R114H) |
| Amino acid position 145 (e.g., V145G) |
| Amino acid position 200               |
| Amino acid position 292 (e.g., V292M) |
| Amino acid position 294               |
| Amino acid position 321 (e.g., G321R) |
| Amino acid position 330 (e.g., R330Q) |
| Amino acid position 338 (e.g., T338I) |
| Amino acid position 360 (e.g., R360W) |
| Amino acid position 393 (e.g., F393L) |



|                                                                                  |
|----------------------------------------------------------------------------------|
| Amino acid position 432                                                          |
| Δ Amino acid residues 505-506 (6-Base Pair In-Frame Germline Deletion in Exon 7) |
| Amino acid position 510 (e.g., A510V)                                            |
| Amino acid position 511 (e.g., E511K)                                            |
| Amino acid position 513 (e.g., G513D)                                            |
| Amino acid position 515 (e.g., C515S, C515W <sup>4</sup> )                       |
| Amino acid position 525 (e.g., R525W)                                            |
| Amino acid position 531 (e.g., C531R, or 9 base pair duplication)                |
| Amino acid position 532 (e.g., duplication)                                      |
| Amino acid position 533 (e.g., G533C, G533S)                                     |
| Amino acid position 550 (e.g., G550E)                                            |
| Amino acid position 591 (e.g., V591I)                                            |
| Amino acid position 593 (e.g., G593E)                                            |
| Amino acid position 595 (e.g., E595D and E595A)                                  |
| Amino acid position 600 (e.g., R600Q)                                            |
| Amino acid position 602 (e.g., I602V)                                            |
| Amino acid position 603 (e.g., K603Q, K603E)                                     |
| Amino acid position 606 (e.g., Y606C)                                            |
| Amino acid position 609 (e.g., C609Y, C609S, C609G, C609R, C609F, C609W)         |
| Amino acid position 611 (e.g., C611R, C611S, C611G, C611Y, C611F, C611W)         |
| Amino acid position 616 (e.g., E616Q)                                            |
| Amino acid position 618 (e.g., C618S, C618Y, C618R, C618G, C618F, C618W)         |
| Amino acid position 620 (e.g., C620S, C620W, C620R, C620G, C620L, C620Y, C620F)  |
| Amino acid position 623 (e.g., E623K)                                            |
| Amino acid position 624 (e.g., D624N)                                            |

|                                                                                                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Amino acid position 630 (e.g., C630A, C630R, C630S, C630Y, C630F, C630W)                                                                                                  |
| Amino acid position 631 (e.g., D631N, D631Y, D631A, D631G, D631V, D631E, )                                                                                                |
| Amino acid position 632 (e.g., E632K, E632G)                                                                                                                              |
| $\Delta$ Amino acid residues 632-633 (6-Base Pair In-Frame Germline Deletion in Exon 11)                                                                                  |
| Amino acid position 633 (e.g., 9 base pair duplication)                                                                                                                   |
| Amino acid position 634 (e.g., C634W, C634Y, C634S, C634R, C634F, C634G, C634L, C634A, or C634T, or an insertion ELCR, or a 12 base pair duplication) (e.g., causing MTC) |
| Amino acid position 635 (e.g., R635G)                                                                                                                                     |
| Amino acid position 636 (e.g., T636P, T636M)                                                                                                                              |
| Amino acid position 640 (e.g., A640G)                                                                                                                                     |
| Amino acid position 641 (e.g., A641S, A641T)                                                                                                                              |
| Amino acid position 648 (e.g., V648I)                                                                                                                                     |
| Amino acid position 649 (e.g., S649L)                                                                                                                                     |
| Amino acid position 664 (e.g., A664D)                                                                                                                                     |
| Amino acid position 665 (e.g., H665Q)                                                                                                                                     |
| Amino acid position 666 (e.g., K666E, K666M, K666N, K666R)                                                                                                                |
| Amino acid position 686 (e.g., S686N)                                                                                                                                     |
| Amino acid position 689 (e.g., S689T)                                                                                                                                     |
| Amino acid position 691 (e.g., G691S)                                                                                                                                     |
| Amino acid position 694 (e.g., R694Q)                                                                                                                                     |
| Amino acid position 700 (e.g., M700L)                                                                                                                                     |
| Amino acid position 706 (e.g., V706M, V706A)                                                                                                                              |
| Amino acid position 713 splice variant (e.g., E713K)                                                                                                                      |
| Amino acid position 732 (e.g., E732K)                                                                                                                                     |
| Amino acid position 736 (e.g., G736R)                                                                                                                                     |

|                                                                                 |
|---------------------------------------------------------------------------------|
| Amino acid position 748 (e.g., G748C)                                           |
| Amino acid position 750 (e.g., A750P)                                           |
| Amino acid position 765 (e.g., S765P)                                           |
| Amino acid position 766 (e.g., P766S, P766M)                                    |
| Amino acid position 768 (e.g., E768Q, E768D)                                    |
| Amino acid position 769 (e.g., L769L)                                           |
| Amino acid position 770 (e.g., R770Q)                                           |
| Amino acid position 771 (e.g., D771N)                                           |
| Amino acid position 777 (e.g., N777S)                                           |
| Amino acid position 778 (e.g., V778I)                                           |
| Amino acid position 781 (e.g., Q781R)                                           |
| Amino acid position 790 (e.g., L790F)                                           |
| Amino acid position 791 (e.g., Y791F, Y791N)                                    |
| Amino acid position 802                                                         |
| Amino acid position 804 (e.g., V804L, V804M, V804E) (e.g., causing MTC)         |
| Amino acid position 805 (e.g., E805K)                                           |
| Amino acid position 804/805 (e.g., V804M/E805K)                                 |
| Amino acid position 806 (e.g., Y806F, Y806S, Y806G, Y806C, Y806E, Y806H, Y806N) |
| Amino acid position 810 (e.g., G810R, G810S, G810A, G810C, G810V, and G810D)    |
| Amino acid position 818 (e.g., E818K)                                           |
| Amino acid position 819 (e.g., S819I)                                           |
| Amino acid position 823 (e.g., G823E)                                           |
| Amino acid position 826 (e.g., Y826M, Y826S)                                    |
| Amino acid position 833 (e.g., R833C)                                           |
| Amino acid position 836 (e.g., S836S)                                           |
| Amino acid position 841 (e.g., P841L, P841P)                                    |
| Amino acid position 843 (e.g., E843D)                                           |

|                                                                            |
|----------------------------------------------------------------------------|
| Amino acid position 844 (e.g., R844W, R844Q, R844L)                        |
| Amino acid position 848 (e.g., M848T)                                      |
| Amino acid position 852 (e.g., I852M)                                      |
| Amino acid position 865 (e.g., L865V)                                      |
| Amino acid position 870 (e.g., L870F)                                      |
| Amino acid position 873 (e.g., R873W)                                      |
| Amino acid position 876 (e.g., A876V)                                      |
| Amino acid position 881 (e.g., L881V)                                      |
| Amino acid position 882                                                    |
| Amino acid position 883 (e.g., A883F, A883S, A883T)                        |
| Amino acid position 884 (e.g., E884K)                                      |
| Amino acid position 886 (e.g., R886W)                                      |
| Amino acid position 891 (e.g., S891A)                                      |
| Amino acid position 897 (e.g., R897Q)                                      |
| Amino acid position 898 (e.g., D898V)                                      |
| Amino acid position 900 (e.g., Y900F)                                      |
| Amino acid position 901 (e.g., E901K)                                      |
| Amino acid position 904 (e.g., S904F, S904S, S904C)                        |
| Amino acid position 907 (e.g., K907E, K907M)                               |
| Amino acid position 908 (e.g., R908K)                                      |
| Amino acid position 911 (e.g., G911D)                                      |
| Amino acid position 912 (e.g., R912P, R912Q)                               |
| Amino acid position 918 (e.g., M918T, M918V, M918L)<br>(e.g., causing MTC) |
| Amino acid position 919 (e.g., A919V)                                      |
| Amino acid position 921 (e.g., E921K)                                      |
| Amino acid position 922 (e.g., S922P, S922Y)                               |
| Amino acid position 930 (e.g., T930M)                                      |
| Amino acid position 961 (e.g., F961L)                                      |

|                                         |
|-----------------------------------------|
| Amino acid position 972 (e.g., R972G)   |
| Amino acid position 982 (e.g., R982C)   |
| Amino acid position 1009 (e.g., M1009V) |
| Amino acid position 1015 (e.g., Y1015F) |
| Amino acid position 1017 (e.g., D1017N) |
| Amino acid position 1041 (e.g., V1041G) |
| Amino acid position 1064 (e.g., M1064T) |
| Amino acid position 1096 (e.g., Y1096F) |
| RET+3                                   |
| (In-Frame Deletion in Exons 6 and 11)   |
| (3bp In-Frame Deletion in Exon 15)      |

<sup>A</sup>The RET kinase mutations shown above may be activating mutations and/or may confer increased resistance of the RET kinase to a RET inhibitor and/or a multi-kinase inhibitor (MKI), e.g., as compared to a wildtype RET kinase.

**[00365]** In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes a splice variation in a RET mRNA which results in an expressed protein that is an alternatively spliced variant of RET having at least one residue deleted (as compared to the wild-type RET kinase) resulting in a constitutive activity of a RET kinase domain.

**[00366]** A “RET kinase inhibitor” as defined herein includes any compound exhibiting RET inhibition activity. In some embodiments, a RET kinase inhibitor is selective for a RET kinase. Exemplary RET kinase inhibitors can exhibit inhibition activity (IC<sub>50</sub>) against a RET kinase of less than about 1000 nM, less than about 500 nM, less than about 200 nM, less than about 100 nM, less than about 50 nM, less than about 25 nM, less than about 10 nM, or less than about 1 nM as measured in an assay as described herein. In some embodiments, a RET kinase inhibitor can exhibit inhibition activity (IC<sub>50</sub>) against a RET kinase of less than about 25 nM, less than about 10 nM, less than about 5 nM, or less than about 1 nM as measured in an assay as provided herein.

**[00367]** As used herein, a “first RET kinase inhibitor” or “first RET inhibitor” is a RET kinase inhibitor as defined herein, but which does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as defined herein. As used herein, a “second RET kinase inhibitor” or “second RET inhibitor” is a RET kinase inhibitor as defined herein, but which does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as defined herein. When both a first and a second

RET inhibitor are present in a method provided herein, the first and second RET kinase inhibitor are different.

**[00368]** In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes at least one point mutation in a RET gene that results in the production of a RET kinase that has one or more amino acid substitutions or insertions or deletions in a RET gene that results in the production of a RET kinase that has one or more amino acids inserted or removed, as compared to the wild-type RET kinase. In some cases, the resulting RET kinase is more resistant to inhibition of its phosphotransferase activity by one or more first RET kinase inhibitor(s), as compared to a wildtype RET kinase or a RET kinase not including the same mutation. Such mutations, optionally, do not decrease the sensitivity of the cancer cell or tumor having the RET kinase to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof (e.g., as compared to a cancer cell or a tumor that does not include the particular RET inhibitor resistance mutation). In such embodiments, a RET inhibitor resistance mutation can result in a RET kinase that has one or more of an increased  $V_{max}$ , a decreased  $K_m$  for ATP, and an increased  $K_D$  for a first RET kinase inhibitor, when in the presence of a first RET kinase inhibitor, as compared to a wildtype RET kinase or a RET kinase not having the same mutation in the presence of the same first RET kinase inhibitor.

**[00369]** In other embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, includes at least one point mutation in a RET gene that results in the production of a RET kinase that has one or more amino acid substitutions as compared to the wild-type RET kinase, and which has increased resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as compared to a wildtype RET kinase or a RET kinase not including the same mutation. In such embodiments, a RET inhibitor resistance mutation can result in a RET kinase that has one or more of an increased  $V_{max}$ , a decreased  $K_m$ , and a decreased  $K_D$  in the presence of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as compared to a wildtype RET kinase or a RET kinase not having the same mutation in the presence of the same compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00370]** Examples of RET inhibitor resistance mutations can, e.g., include point mutations, insertions, or deletions in and near the ATP binding site in the tertiary structure of RET kinase (e.g., amino acid positions 730-733, 738, 756, 758, 804, 805, 807, 810, 811, 881, and 892 of a wildtype RET kinase, e.g., the exemplary wildtype RET kinase described herein), including but not limited to a gatekeeper residue (e.g., amino acid position 804 in a wildtype RET kinase), P-loop residues (e.g., amino acid positions 730-737 in a wildtype RET kinase), residues in or near the DFG motif (e.g., amino acid positions 888-898 in a wildtype RET kinase), and ATP cleft solvent front amino acid residues (e.g., amino acid positions 758, 811, and 892 of a wildtype RET kinase). Additional examples of these types of mutations include changes in residues that may affect enzyme activity and/or drug binding including but are not limited to residues in the activation loop (e.g., amino acid positions 891-916 of a wildtype RET kinase), residues near or interacting with the activation loop, residues contributing to active or inactive enzyme conformations, changes including mutations, deletions, and insertions in the loop proceeding the C-helix and in the C-helix (e.g., amino acid positions 768-788 in a wildtype RET protein). In some embodiments, the wildtype RET protein is the exemplary wildtype RET kinase described herein. Specific residues or

residue regions that may be changed (and are RET inhibitor resistance mutations) include but are not limited to those listed in Table 3, with numbering based on the human wildtype RET protein sequence (e.g., SEQ ID NO: 1). As can be appreciated by those skilled in the art, an amino acid position in a reference protein sequence that corresponds to a specific amino acid position in SEQ ID NO: 1 can be determined by aligning the reference protein sequence with SEQ ID NO: 1 (e.g., using a software program, such as ClustalW2). Additional examples of RET inhibitor resistance mutation positions are shown in Table 4. Changes to these residues may include single or multiple amino acid changes, insertions within or flanking the sequences, and deletions within or flanking the sequences. See also J. Kooistra, G. K. Kanev, O. P. J. Van Linden, R. Leurs, I. J. P. De Esch, and C. De Graaf, "KLIFS: A structural kinase-ligand interaction database," *Nucleic Acids Res.*, vol. 44, no. D1, pp. D365–D371, 2016, which is incorporated by reference in its entirety herein.

**[00371]** Exemplary Sequence of Mature Human RET Protein (SEQ ID NO: 1)

```

MAKATSGAAG LRLLLLLLLP LLGKVALGLY FSRDAYWEKL YVDQAAGTPL LYVHALRDAP EEVPSFRLGQ
HLYGTYRTRL HENNWICIQE DTGLLYLNRS LDHSSWEKLS VRNRGFPLLT VYLKVFLSPT SLREGEQWP
GCARVYFSFF NTSFPACSSL KPRELCFPET RPSFRIENR PPGTFHQFRL LPVQFLCPNI SVAYRLLIGE
GLPFRCAPDS LEVSTRWALD REQREKYELV AVCTVHAGAR EEVVMVFPV TVYDEDDSDAP TFPAGVDTAS
AVVEFKRKED TVVATLRVFD ADVVPASGEL VRRYTSTLLP GDTWAQQTFR VEHWPNETSV QANGSFVRAT
VHDYRLVLNR NLSISENRTM QLAVLVNDS FQGPAGVLL LHFNVSVLPV SLHLPSTYSL SVSRRARRFA
QIGKVCVENC QAFSGINVQY KLHSSGANCS TLGVVTS AED TSGILFVNDT KALRRPKCAE LHYMVVATDQ
QTSRQAQAL LVTVEGSYVA EEAGCPLSCA VSKRRLECEE CGGLGSPTGR CEWRQGDGKG ITRNFSTCSP
STKTCPDGHC DVVETQDINI CPQDCLRGS VGGHEPGEPR GIKAGYGCN CFPEEEKCFE EPEDIQDPLC
DELCRTVIAA AVLFSSFIVSV LLSAFCIHCY HKFAHKPPIS SAEMTFRRPA QAFPVSYS S GARRPSLD SM
ENQVSVD AFK ILEDPKWEFP RKNLVLGKTL GEGEFGKVVK ATAFHLKGRA GYTTVAVKML KENASPS ELR
DLLSEFNVLK QVNHPHVIKL YGACSQDGPL LLIVEYAKYG SLRGFLRESR KVGPGYLGSG GSRNSSSLDH
PDERALTMGD LISFAWQISQ GMQYLAEMKL VHRDLAARNI LVAEGRKMKI SDFGLSRDVY EEDSYVKRSQ
GRIPVKWMAI ESLFDHIYTT QSDVWSFGVL LWEIVTLGGN PYPGIPPERL FNLLKTGHRM ERPDNCSEEM
YRLMLQCWKQ EPDKRPVFAD ISKDLEKMMV KRRDYLDLAA STPSDSL IYD DGLSEEETPL VDCNNAPLPR
ALPSTWIENK LYGMSDPNWP GESPVPLTRA DGTNTGFPRY PNDSVYANWM LSPSAAKLMD TFDS

```

**[00372]** In some embodiments, a RET inhibitor resistance mutation can include a dysregulation of a MET gene, a MET kinase, or the expression or activity or level of any of the same.

**[00373]** The phrase "dysregulation of a MET gene, a MET kinase, or the expression or activity or level of any of the same" refers to a genetic mutation (e.g., a MET gene translocation that results in the expression of a fusion protein, a deletion in a MET gene that results in the expression of a RET protein that includes a deletion of at least one amino acid as compared to the wild-type RET protein, or a mutation in a MET gene that results in the expression of a RET protein with one or more point mutations, or an alternative spliced version of a MET mRNA that results in a MET protein that results in the deletion of at least one amino acid in the MET protein as compared to the wild-type MET protein), or a MET gene amplification that results in overexpression of a MET protein or an autocrine activity resulting from the overexpression of a MET gene a cell, that results in a pathogenic increase in the activity of a kinase domain of a MET protein (e.g., a constitutively active kinase domain of a MET protein) in a cell. As another example, a dysregulation of a MET gene, a MET protein, or expression or activity, or level of any of the same, can be a mutation in a MET gene that encodes a MET protein that is constitutively active or has increased activity as compared to a protein encoded by a MET gene that

does not include the mutation. For example, a dysregulation of a MET gene, a MET protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of MET that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not MET). In some examples, dysregulation of a MET gene, a MET protein, or expression or activity, can be a result of a gene translocation of one MET gene with another non-MET gene.

**[00374]** The term "wildtype MET" or "wild-type MET" describes a nucleic acid (e.g., a MET gene or a MET mRNA) or protein (e.g., a MET protein) that is found in a subject that does not have a MET-associated cancer (and optionally also does not have an increased risk of developing a MET-associated cancer and/or is not suspected of having a MET-associated cancer), or is found in a cell or tissue from a subject that does not have a MET-associated cancer (and optionally also does not have an increased risk of developing a MET-associated cancer and/or is not suspected of having a MET-associated cancer). The term "MET-associated cancer" as used herein refers to cancers associated with or having a dysregulation of a MET gene, a MET kinase, or expression or activity, or level of any of the same.

**[00375]** In some embodiments, compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof are useful in treating patients that develop cancers with RET inhibitor resistance mutations (e.g., that result in an increased resistance to a first RET inhibitor, e.g., a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D, and/or one or more RET inhibitor resistance mutations listed in Tables 3 and 4) by either dosing in combination or as a subsequent or additional (e.g., follow-up) therapy to existing drug treatments (e.g., other RET kinase inhibitors; e.g., first and/or second RET kinase inhibitors). Exemplary first and second RET kinase inhibitors are described herein. In some embodiments, a first or second RET kinase inhibitor can be selected from the group consisting of cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864.

**[00376]** In some embodiments, compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof are useful for treating a cancer that has been identified as having one or more RET inhibitor resistance mutations (that result in an increased resistance to a first or second RET inhibitor, e.g., a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or e.g., a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D). In some embodiments, the one or more RET inhibitor resistance mutations occur in a nucleic acid sequence encoding a RET fusion protein (e.g. any of the RET gene fusion proteins described in Table 1) resulting in a RET fusion protein that exhibits RET kinase inhibitor resistance. In some embodiments, the one or more RET inhibitor resistance mutations occurs in a nucleic acid sequence encoding a mutant RET protein (e.g. a mutant RET protein having any of the mutations described in Table 2) resulting in a mutant RET protein that exhibits RET kinase resistance. Non-limiting examples of RET inhibitor resistance mutations are listed in Tables 3 and 4.



**Table 3. RET Inhibitor Resistance Mutations**

| <b>Exemplary RET Resistance Mutations</b>                                                                                                  |
|--------------------------------------------------------------------------------------------------------------------------------------------|
| Amino acid position 634 (e.g., C634W) <sup>10</sup>                                                                                        |
| Amino acid position 732 (e.g., E732K) <sup>7</sup>                                                                                         |
| Amino acid position 788 (e.g., I788N) <sup>8</sup>                                                                                         |
| Amino acid position 790 (e.g., L790F) <sup>9</sup>                                                                                         |
| Amino acid position 804 (e.g., V804M <sup>1, 2</sup> , V804L <sup>1, 2</sup> , V804E <sup>6</sup> )                                        |
| Amino acid position 778/804 <sup>13</sup>                                                                                                  |
| Amino acid position 804/805 (e.g., V804M/E805K) <sup>3</sup>                                                                               |
| Amino acid position 806 (e.g., Y806C <sup>4, 6</sup> , Y806E <sup>4</sup> , Y806S <sup>6</sup> , Y806H <sup>6</sup> , Y806N <sup>6</sup> ) |
| Amino acid position 804/806 (e.g., V804M/Y806C) <sup>11</sup>                                                                              |
| Amino acid position 810 (e.g., G810A <sup>5</sup> , G810R <sup>6</sup> , G810S <sup>6</sup> , G810C, G810V, and G810D)                     |
| Amino acid position 865 (e.g., L865V <sup>6</sup> )                                                                                        |
| Amino acid position 870 (e.g., L870F <sup>6</sup> )                                                                                        |
| Amino acid position 891 (e.g., S891A) <sup>10</sup>                                                                                        |
| Amino acid position 904 (e.g., S904F) <sup>12</sup>                                                                                        |
| Amino acid position 804/904 (e.g., V804M/S904C) <sup>11</sup>                                                                              |
| Amino acid position 918 (e.g., M918T) <sup>10</sup>                                                                                        |

<sup>1</sup> Yoon et al., *J. Med. Chem.* 59(1):358-73, 2016.<sup>2</sup> U.S. Patent No. 8,629,135.<sup>3</sup> Cranston, et al., *Cancer Res.* 66(20):10179-87, 2006.<sup>4</sup> Carlomagno, et al., *Endocr. Rel. Cancer* 16(1):233-41, 2009.<sup>5</sup> Huang et al., *Mol. Cancer Ther.*, 2016 Aug 5. pii: molcanther.0258.2016. [Epub ahead of print].<sup>6</sup> PCT Patent Application Publication No. WO 2016/127074.<sup>7</sup> Mamedova et al., Summer Undergraduate Research Programs (SURP) Student Abstracts, University of Oklahoma Health Sciences Center, 2016.<sup>8</sup> Plenker et al., *Sci. Transl. Med.*, 9(394), doi: 10.1126/scitranslmed.aah6144, 2017.

<sup>9</sup> Kraft et al, *Cancer Research*, 2017, Vol. 77, No. 13, Supp. Supplement 1. Abstract Number: 4882; American Association for Cancer Research Annual Meeting 2017. Washington, DC, United States. 01 Apr 2017-05 Apr 2017.

<sup>10</sup> U.S. Patent Application Publication No. 2018/0022732.

<sup>11</sup> Roskoski and Sadeghi-Nejad, *Pharmacol. Res.*, 128, 1-17. doi: 10.1016/j.phrs.2017.12.021, 2018.

<sup>12</sup> Nakaoku, et al. *Nat Commun*, 9(1), 625. doi: 10.1038/s41467-018-02994-7, 2018.

<sup>13</sup> Roy et al. *Oncologist*, 18(10): 1093-1100. doi: 10.1634/theoncologist.2013-0053, 2013.

**Table 4. Additional Exemplary Amino Acid Positions of RET Inhibitor Resistance Mutations**

| RET Amino Acid and Position | Exemplary Mutation | Mechanistic Resistance Rationale                     |
|-----------------------------|--------------------|------------------------------------------------------|
| L730                        | P                  | Steric hindrance and/or active conformational effect |
| G731                        | V                  | Steric hindrance and/or active conformational effect |
| E732                        | K                  | Steric hindrance and/or active conformational effect |
| G733                        | V                  | Steric hindrance and/or active conformational effect |
| E734                        | K                  | Steric hindrance and/or active conformational effect |
| L760                        | M                  | Active conformational effect                         |
| K761                        | E                  | Active conformational effect                         |
| E762                        | K                  | Active conformational effect                         |
| N763                        | D                  | Active conformational effect                         |
| A764                        | V                  | Active conformational effect                         |
| S765                        | N                  | Active conformational effect                         |
| P766                        | A                  | Active conformational effect                         |
| S767                        | C                  | Active conformational effect                         |
| E768                        | K                  | Active conformational effect                         |
| L779                        | M                  | Steric hindrance and/or active conformational effect |
| I788                        | M                  | Steric hindrance and/or active conformational effect |
| M868                        | R                  | Steric hindrance and/or active conformational effect |

|      |   |                                                      |
|------|---|------------------------------------------------------|
| K869 | E | Steric hindrance and/or active conformational effect |
| L870 | Q | Steric hindrance and/or active conformational effect |
| V871 | M | Steric hindrance and/or active conformational effect |
| H872 | R | Steric hindrance and/or active conformational effect |
| R873 | P | Steric hindrance and/or active conformational effect |
| D874 | Y | Steric hindrance and/or active conformational effect |
| L881 | R | Steric hindrance and/or active conformational effect |
| L895 | M | Active conformational effect                         |
| S896 | N | Active conformational effect                         |
| R897 | C | Active conformational effect                         |
| D898 | Y | Active conformational effect                         |
| V899 | G | Active conformational effect                         |
| Y900 | D | Active conformational effect                         |
| E901 | K | Active conformational effect                         |
| E902 | K | Active conformational effect                         |
| D903 | Y | Active conformational effect                         |
| S904 | C | Active conformational effect                         |
| Y905 | D | Active conformational effect                         |
| V906 | M | Active conformational effect                         |
| K907 | E | Active conformational effect                         |
| R908 | P | Active conformational effect                         |
| S909 | C | Active conformational effect                         |
| Q910 | R | Active conformational effect                         |
| G911 | C | Active conformational effect                         |
| R912 | P | Active conformational effect                         |

**[00377]** The oncogenic role of RET was first described in papillary thyroid carcinoma (PTC) (Grieco et al., *Cell*, 1990, 60, 557-63), which arises from follicular thyroid cells and is the most common

thyroid malignancy. Approximately 20-30% of PTC harbor somatic chromosomal rearrangements (translocations or inversions) linking the promoter and the 5' portions of constitutively expressed, unrelated genes to the RET tyrosine kinase domain (Greco et al., *Q. J. Nucl. Med. Mol. Imaging*, 2009, 53, 440-54), therefore driving its ectopic expression in thyroid cells. Fusion proteins generated by such rearrangements are termed "RET/PTC" proteins. For example, RET/PTC 1 is a fusion between CCDD6 and RET that is commonly found in papillary thyroid carcinomas. Similarly, both RET/PTC3 and RET/PTC4 are fusions of ELE1 and RET that are commonly found in papillary thyroid carcinomas, although the fusion events resulting RET/PTC3 and RET/PTC4 lead to different proteins with different molecular weights (see e.g., Fugazzola et al., *Oncogene*, 13(5):1093-7, 1996). Some RET fusions associated with PTC are not referred to as "RET/PTC", but instead are referred to as the the fusion protein itself. For example, fusion between RET and both ELKS and PCM1 are found in PTCs, but the fusion proteins are referred to as ELKS-RET and PCM1-RET (see e.g., Romei and Elisei, *Front. Endocrinol. (Lausanne)*, 3:54, doi: 10.3389/fendo.2012.00054, 2012). The role of RET-PTC rearrangements in the pathogenesis of PTC has been confirmed in transgenic mice (Santoro et al., *Oncogene*, 1996, 12, 1821-6). To date, a variety of fusion partners have been identified, from PTC and other cancer types, all providing a protein/protein interaction domain that induces ligand-independent RET dimerization and constitutive kinase activity (see, e.g., Table 1). Recently, a 10.6 Mb pericentric inversion in chromosome 10, where RET gene maps, has been identified in about 2% of lung adenocarcinoma patients, generating different variants of the chimeric gene KIF5B-RET (Ju et al., *Genome Res.*, 2012, 22, 436-45; Kohno et al., 2012, *Nature Med.*, 18, 375-7; Takeuchi et al., *Nature Med.*, 2012, 18, 378-81; Lipson et al., 2012, *Nature Med.*, 18, 382-4). The fusion transcripts are highly expressed and all the resulting chimeric proteins contain the N-terminal portion of the coiled-coil region of KIF5B, which mediates homodimerization, and the entire RET kinase domain. None of RET positive patients harbor other known oncogenic alterations (such as EGFR or K-Ras mutation, ALK translocation), supporting the possibility that KIF5B-RET fusion could be a driver mutation of lung adenocarcinoma. The oncogenic potential of KIF5B-RET has been confirmed by transfecting the fusion gene into cultured cell lines: similarly to what has been observed with RET-PTC fusion proteins, KIF5B-RET is constitutively phosphorylated and induces NIH-3T3 transformation and IL-3 independent growth of BA-F3 cells. However, other RET fusion proteins have been identified in lung adenocarcinoma patients, such as the CCDC6-RET fusion protein, which has been found to play a key role in the proliferation of the human lung adenocarcinoma cell line LC-2/ad (*Journal of Thoracic Oncology*, 2012, 7(12):1872-1876). RET inhibitors have been shown to be useful in treating lung cancers involving RET rearrangements (Drilon, A.E. et al. *J Clin Oncol* 33, 2015 (suppl; abstr 8007)). RET fusion proteins have also been identified in patients having colorectal cancer (Song Eun-Kee, et al. *International Journal of Cancer*, 2015, 136: 1967-1975).

**[00378]** Besides rearrangements of the RET sequence, gain of function point mutations of RET proto-oncogene are also driving oncogenic events, as shown in medullary thyroid carcinoma (MTC), which arises from parafollicular calcitonin-producing cells (de Groot, et al., *Endocrine Rev.*, 2006, 27, 535-60; Wells and Santoro, *Clin. Cancer Res.*, 2009, 15, 7119-7122). Around 25% of MTC are associated with multiple endocrine neoplasia type 2 (MEN2), a group of inherited cancer syndromes affecting neuroendocrine organs caused by germline activating point mutations of RET. In MEN2 subtypes (MEN2A, MEN2B and Familial MTC/FMTC) RET gene mutations have a strong phenotype-genotype correlation defining different MTC aggressiveness and clinical manifestations of the disease. In MEN2A syndrome mutations involve one of the six cysteine residues (mainly C634) located in the cysteine-rich

extracellular region, leading to ligand-independent homodimerization and constitutive RET activation. Patients develop MTC at a young age (onset at 5-25 years) and may also develop pheochromocytoma (50%) and hyperparathyroidism. MEN2B is mainly caused by M918T mutation, which is located in the kinase domain. This mutation constitutively activates RET in its monomeric state and alters substrate recognition by the kinase. MEN2B syndrome is characterized by an early onset (< 1 year) and very aggressive form of MTC, pheochromocytoma (50% of patients) and ganglioneuromas. In FMTC the only disease manifestation is MTC, usually occurring at an adult age. Many different mutations have been detected, spanning the entire RET gene. The remaining 75% of MTC cases are sporadic and about 50% of them harbor RET somatic mutations: the most frequent mutation is M918T that, as in MEN2B, is associated with the most aggressive phenotype. Somatic point mutations of RET have also been described in other tumors such as colorectal cancer (Wood et al., *Science*, 2007, 318, 1108-13) and small cell lung carcinoma (*Jpn. J. Cancer Res.*, 1995, 86, 1127-30). In some embodiments, the MTC is RET-fusion positive MTC.

**[00379]** RET signaling components have been found to be expressed in primary breast tumors and to functionally interact with estrogen receptor-cc pathway in breast tumor cell lines (Boulay et al., *Cancer Res.* 2008, 68, 3743-51; Plaza-Menacho et al., *Oncogene*, 2010, 29, 4648-57), while RET expression and activation by GDNF family ligands could play an important role in perineural invasion by different types of cancer cells (Ito et al., *Surgery*, 2005, 138, 788-94; Gil et al., *J. Natl. Cancer Inst.*, 2010, 102, 107-18; Iwahashi et al., *Cancer*, 2002, 94, 167-74).

**[00380]** RET is also expressed in 30-70% of invasive breast cancers, with expression being relatively more frequent in estrogen receptor-positive tumors (Plaza-Menacho, I., et al., *Oncogene*, 2010, 29, 4648-4657; Essegir, S., et al., *Cancer Res.*, 2007, 67, 11732-11741; Morandi, A., et al., *Cancer Res.*, 2013, 73, 3783-3795; Gattelli, A., *EMBO Mol. Med.*, 2013, 5, 1335-1350).

**[00381]** The identification of RET rearrangements has been reported in a subset of (patient-derived xenograft) PDX established from colorectal cancer. Although the frequency of such events in colorectal cancer patients remains to be defined, these data suggest a role of RET as a target in this indication (Gozgit et al., AACR Annual Meeting 2014). Studies have shown that the RET promoter is frequently methylated in colorectal cancers, and heterozygous missense mutations, which are predicted to reduce RET expression, are identified in 5-10% of cases, which suggests that RET might have some features of a tumor suppressor in sporadic colon cancers (Luo, Y., et al., *Oncogene*, 2013, 32, 2037-2047; Sjoblom, T., et al., *Science*, 2006, 268-274; Cancer Genome Atlas Network, *Nature*, 2012, 487, 330-337).

**[00382]** An increasing number of tumor types are now being shown to express substantial levels of wild-type RET kinase that could have implications for tumor progression and spread. RET is expressed in 50-65% of pancreatic ductal carcinomas, and expression is more frequent in metastatic and higher grade tumors (Ito, Y., et al., *Surgery*, 2005, 138, 788-794; Zeng, Q., et al., *J. Int. Med. Res.* 2008, 36, 656-664).

**[00383]** In neoplasms of hematopoietic lineages, RET is expressed in acute myeloid leukemia (AML) with monocytic differentiation, as well as in CMML (Gattei, V. et al., *Blood*, 1997, 89, 2925-2937; Gattei, V., et al., *Ann. Hematol.*, 1998, 77, 207-210; Camos, M., *Cancer Res.* 2006, 66, 6947-6954). Recent studies have identified rare chromosomal rearrangements that involve RET in patients with

chronic myelomonocytic leukemia (CMML). CMML is frequently associated with rearrangements of several tyrosine kinases, which result in the expression of chimeric cytosolic oncoproteins that lead to activation of RAS pathways (Kohlmann, A., et al., *J. Clin. Oncol.* 2010, 28, 2858-2865). In the case of RET, gene fusions that link RET with BCR (BCR-RET) or with fibroblast growth factor receptor 1 oncogene partner (FGFR1OP-RET) were transforming in early hematopoietic progenitor cells and could shift maturation of these cells towards monocytic paths, probably through the initiation of RET-mediated RAS signaling (Ballerini, P., et al., *Leukemia*, 2012, 26, 2384-2389).

**[00384]** RET expression has also been shown to occur in several other tumor types, including prostate cancer, small-cell lung carcinoma, melanoma, renal cell carcinoma, and head and neck tumors (Narita, N., et al., *Oncogene*, 2009, 28, 3058-3068; Mulligan, L. M., et al., *Genes Chromosomes Cancer*, 1998, 21, 326-332; Flavin, R., et al., *Urol. Oncol.*, 2012, 30, 900-905; Dawson, D. M., *J Natl Cancer Inst*, 1998, 90, 519-523).

**[00385]** In neuroblastoma, RET expression and activation by GFLs has roles in tumor cell differentiation, potentially collaborating with other neurotrophic factor receptors to down regulate N-Myc, the expression of which is a marker of poor prognosis (Hofstra, R. M., W., et al., *Hum. Genet.* 1996, 97, 362-364; Petersen, S. and Bogenmann, E., *Oncogene*, 2004, 23, 213-225; Brodeur, G. M., *Nature Ref. Cancer*, 2003, 3, 203-216).

**[00386]** Multitargeted inhibitors which cross react with RET are known (Borrello, M.G., et al., *Expert Opin. Ther. Targets*, 2013, 17(4), 403-419; International Patent Application Nos. WO 2014/141187, WO 2014/184069, and WO 2015/079251). Such multitargeted inhibitors (or multikinase inhibitors or MKIs) can also be associated with development of RET inhibitor resistance mutations. See, for example, Q. Huang et al., "Preclinical Modeling of KIF5B-RET Fusion Lung Adenocarcinoma.," *Mol. Cancer Ther.*, no. 18, pp. 2521-2529, 2016; Yasuyuki Kaneta et al., Abstract B173: Preclinical characterization and antitumor efficacy of DS-5010, a highly potent and selective RET inhibitor, *Mol Cancer Ther* January 1 2018 (17) (1 Supplement) B173; DOI:10.1158/1535-7163.TARG-17-B173, both of which are incorporated by reference in their entirety herein.

**[00387]** Accordingly, provided herein are methods for treating a patient diagnosed with (or identified as having) a cancer that include administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Also provided herein are methods for treating a patient identified or diagnosed as having a RET-associated cancer that include administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof. In some embodiments, the patient that has been identified or diagnosed as having a RET-associated cancer through the use of a regulatory agency-approved, e.g., FDA-approved test or assay for identifying dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, in a patient or a biopsy sample from the patient or by performing any of the non-limiting examples of assays described herein. In some embodiments, the test or assay is provided as a kit. In some embodiments, the cancer is a RET-associated cancer. For example, the RET-associated cancer can be a cancer that includes one or more RET inhibitor resistance mutations.

**[00388]** Also provided are methods for treating cancer in a patient in need thereof, the method comprising: (a) detecting a RET-associated cancer in the patient; and (b) administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof. Some embodiments of these methods further include administering to the subject another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or an immunotherapy). In some embodiments, the subject was previously treated with a first RET inhibitor or previously treated with another anticancer treatment, e.g., at least partial resection of the tumor or radiation therapy. In some embodiments, the patient is determined to have a RET-associated cancer through the use of a regulatory agency-approved, e.g., FDA-approved test or assay for identifying dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, in a patient or a biopsy sample from the patient or by performing any of the non-limiting examples of assays described herein. In some embodiments, the test or assay is provided as a kit. In some embodiments, the cancer is a RET-associated cancer. For example, the RET-associated cancer can be a cancer that includes one or more RET inhibitor resistance mutations.

**[00389]** Also provided are methods of treating a patient that include performing an assay on a sample obtained from the patient to determine whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, and administering (e.g., specifically or selectively administering) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof to the patient determined to have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. Some embodiments of these methods further include administering to the subject another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy). In some embodiments of these methods, the subject was previously treated with a first RET inhibitor or previously treated with another anticancer treatment, e.g., at least partial resection of a tumor or radiation therapy. In some embodiments, the patient is a patient suspected of having a RET-associated cancer, a patient presenting with one or more symptoms of a RET-associated cancer, or a patient having an elevated risk of developing a RET-associated cancer. In some embodiments, the assay utilizes next generation sequencing, pyrosequencing, immunohistochemistry, or break apart FISH analysis. In some embodiments, the assay is a regulatory agency-approved assay, e.g., FDA-approved kit. In some embodiments, the assay is a liquid biopsy. Additional, non-limiting assays that may be used in these methods are described herein. Additional assays are also known in the art. In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations.

**[00390]** Also provided is a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof for use in treating a RET-associated cancer in a patient identified or diagnosed as having a RET-associated cancer through a step of performing an assay (e.g., an in vitro assay) on a sample obtained from the patient to determine whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, where the presence of a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, identifies that the patient has a RET-associated cancer.

Also provided is the use of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof for the manufacture of a medicament for treating a RET-associated cancer in a patient identified or diagnosed as having a RET-associated cancer through a step of performing an assay on a sample obtained from the patient to determine whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same where the presence of dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, identifies that the patient has a RET-associated cancer. Some embodiments of any of the methods or uses described herein further include recording in the patient's clinical record (e.g., a computer readable medium) that the patient is determined to have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, through the performance of the assay, should be administered a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof. In some embodiments, the assay utilizes next generation sequencing, pyrosequencing, immunohistochemistry, or break apart FISH analysis. In some embodiments, the assay is a regulatory agency-approved assay, e.g., FDA-approved kit. In some embodiments, the assay is a liquid biopsy. In some embodiments, the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations.

**[00391]** Also provided is a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, for use in the treatment of a cancer in a patient in need thereof or a patient identified or diagnosed as having a RET-associated cancer. Also provided is the use of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof for the manufacture of a medicament for treating a cancer in a patient identified or diagnosed as having a RET-associated cancer. In some embodiments, the cancer is a RET-associated cancer, for example, a RET-associated cancer having one or more RET inhibitor resistance mutations. In some embodiments, a patient is identified or diagnosed as having a RET-associated cancer through the use of a regulatory agency-approved, e.g., FDA-approved, kit for identifying dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, in a patient or a biopsy sample from the sample. As provided herein, a RET-associated cancer includes those described herein and known in the art.

**[00392]** In some embodiments of any of the methods or uses described herein, the patient has been identified or diagnosed as having a cancer with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. In some embodiments of any of the methods or uses described herein, the patient has a tumor that is positive for a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. In some embodiments of any of the methods or uses described herein, the patient can be a patient with a tumor(s) that is positive for a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. In some embodiments of any of the methods or uses described herein, the patient can be a patient whose tumors have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. In some embodiments of any of the methods or uses described herein, the patient is suspected of having a RET-associated cancer (e.g., a cancer having one or more RET inhibitor resistance mutations). In some embodiments, provided herein are methods for treating a RET-associated cancer in a patient in need of such treatment, the method comprising a) detecting a dysregulation of a RET gene, a RET kinase, or the



expression or activity or level of any of the same in a sample from the patient; and b) administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same includes one or more fusion proteins. Non-limiting examples of RET gene fusion proteins are described in Table 1. In some embodiments, the fusion protein is KIF5B-RET. In some embodiments, the dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same includes one or more RET kinase protein point mutations/insertions/deletions. Non-limiting examples of RET kinase protein point mutations/insertions/deletions are described in Tables 2 and 2a. In some embodiments, the RET kinase protein point mutations/insertions/deletions are selected from the group consisting of M918T, M918V, C634W, V804L, V804M, G810S, and G810R. In some embodiments, the dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations. Non-limiting examples of RET inhibitor resistance mutations are described in Tables 3 and 4. In some embodiments, the RET inhibitor resistance mutation is V804M. In some embodiments, the RET inhibitor resistance mutation is G810S. In some embodiments, the RET inhibitor resistance mutation is G810R. In some embodiments, the cancer with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit. In some embodiments, the tumor that is positive for a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is a tumor positive for one or more RET inhibitor resistance mutations. In some embodiments, the tumor with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit.

**[00393]** In some embodiments of any of the methods or uses described herein, the patient has a clinical record indicating that the patient has a tumor that has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same (e.g., a tumor having one or more RET inhibitor resistance mutations). In some embodiments, the clinical record indicates that the patient should be treated with one or more of the compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or compositions provided herein. In some embodiments, the cancer with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is a cancer having one or more RET inhibitor resistance mutations. In some embodiments, the cancer with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit. In some embodiments, the tumor that is positive for a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is a tumor positive for one or more RET inhibitor resistance mutations. In some embodiments, the tumor with a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same is determined using a regulatory agency-approved, e.g., FDA-approved, assay or kit.

**[00394]** Also provided are methods of treating a patient that include administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to a patient having a clinical record that indicates that the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. Also provided is the use of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a

pharmaceutically acceptable salt or solvate thereof for the manufacture of a medicament for treating a RET-associated cancer in a patient having a clinical record that indicates that the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. Some embodiments of these methods and uses can further include: a step of performing an assay (e.g., an in vitro assay) on a sample obtained from the patient to determine whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, and recording the information in a patient's clinical file (e.g., a computer readable medium) that the patient has been identified to have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same. In some embodiments, the assay is an in vitro assay. For example, an assay that utilizes next generation sequencing, immunohistochemistry, or break apart FISH analysis. In some embodiments, the assay is a regulatory agency-approved, e.g., FDA-approved, kit. In some embodiments, the assay is a liquid biopsy. In some embodiments, the dysregulation of a RET gene, RET kinase, or expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations.

**[00395]** Also provided herein is a method of treating a subject. In some embodiments, the method includes performing an assay on a sample obtained from the subject to determine whether the subject has a dysregulation of a RET gene, a RET protein, or expression or level of any of the same. In some such embodiments, the method also includes administering to a subject determined to have a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the method includes determining that a subject has a dysregulation of a RET gene, a RET protein, or expression or level of any of the same via an assay performed on a sample obtained from the subject. In such embodiments, the method also includes administering to a subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the dysregulation in a RET gene, a RET kinase protein, or expression or activity of the same is a gene or chromosome translocation that results in the expression of a RET fusion protein (e.g., any of the RET fusion proteins described herein). In some embodiments, the RET fusion can be selected from a KIF5B-RET fusion and a CCDC6-RET fusion. In some embodiments, the dysregulation in a RET gene, a RET kinase protein, or expression or activity or level of any of the same is one or more point mutation in the RET gene (e.g., any of the one or more of the RET point mutations described herein). The one or more point mutations in a RET gene can result, e.g., in the translation of a RET protein having one or more of the following amino acid substitutions: M918T, M918V, C634W, V804L, V804M, G810S, and G810R. In some embodiments, the dysregulation in a RET gene, a RET kinase protein, or expression or activity or level of any of the same is one or more RET inhibitor resistance mutations (e.g., any combination of the one or more RET inhibitor resistance mutations described herein). Some embodiments of these methods further include administering to the subject another anticancer agent (e.g., a second RET inhibitor a second compound of Formula I

(e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy).

**[00396]** In some embodiments, the compounds provided herein exhibit brain and/or central nervous system (CNS) penetrance. Such compounds are capable of crossing the blood brain barrier and inhibiting a RET kinase in the brain and/or other CNS structures. In some embodiments, the compounds provided herein are capable of crossing the blood brain barrier in a therapeutically effective amount. For example, treatment of a patient with cancer (e.g., a RET-associated cancer such as a RET-associated brain or CNS cancer) can include administration (e.g., oral administration) of the compound to the patient. In some such embodiments, the compounds provided herein are useful for treating a primary brain tumor or metastatic brain tumor. For example, the compounds can be used in the treatment of one or more of gliomas such as glioblastoma (also known as glioblastoma multiforme), astrocytomas, oligodendrogliomas, ependymomas, and mixed gliomas, meningiomas, medulloblastomas, gangliogliomas, schwannomas (neurilemmomas), and craniopharyngiomas (see, for example, the tumors listed in Louis, D.N. et al. *Acta Neuropathol* 131(6), 803-820 (June 2016)). In some embodiments, the brain tumor is a primary brain tumor. In some embodiments, the patient has previously been treated with another anticancer agent, e.g., another RET inhibitor (e.g., a compound that is not a compound of General Formula I) or a multi-kinase inhibitor. In some embodiments, the brain tumor is a metastatic brain tumor. In some embodiments, the patient has previously been treated with another anticancer agent, e.g., another RET inhibitor (e.g., a compound that is not a compound of General Formula I) or a multi-kinase inhibitor.

**[00397]** Also provided are methods (e.g., in vitro methods) of selecting a treatment for a patient identified or diagnosed as having a RET-associated cancer. Some embodiments can further include administering the selected treatment to the patient identified or diagnosed as having a RET-associated cancer. For example, the selected treatment can include administration of a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Some embodiments can further include a step of performing an assay on a sample obtained from the patient to determine whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, and identifying and diagnosing a patient determined to have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, as having a RET-associated cancer. In some embodiments, the cancer is a RET-associated cancer having one or more RET inhibitor resistance mutations. In some embodiments, the patient has been identified or diagnosed as having a RET-associated cancer through the use of a regulatory agency-approved, e.g., FDA-approved, kit for identifying dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, in a patient or a biopsy sample from the patient. In some embodiments, the RET-associated cancers is a cancer described herein or known in the art. In some embodiments, the assay is an in vitro assay. For example, an assay that utilizes the next generation sequencing, immunohistochemistry, or break apart FISH analysis. In some embodiments, the assay is a regulatory agency-approved, e.g., FDA-approved, kit. In some embodiments, the assay is a liquid biopsy.

**[00398]** Also provided herein are methods of selecting a treatment for a patient, wherein the methods include a step of performing an assay on a sample obtained from the patient to determine

whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same (e.g., one or more RET inhibitor resistance mutations), and identifying or diagnosing a patient determined to have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, as having a RET-associated cancer. Some embodiments further include administering the selected treatment to the patient identified or diagnosed as having a RET-associated cancer. For example, the selected treatment can include administration of a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the patient identified or diagnosed as having a RET-associated cancer. In some embodiments, the assay is an in vitro assay. For example, an assay that utilizes the next generation sequencing, immunohistochemistry, or break apart FISH analysis. In some embodiments, the assay is a regulatory agency-approved, e.g., FDA-approved, kit. In some embodiments, the assay is a liquid biopsy.

**[00399]** Also provided are methods of selecting a patient for treatment, wherein the methods include selecting, identifying, or diagnosing a patient having a RET-associated cancer, and selecting the patient for treatment including administration of a therapeutically-effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, identifying or diagnosing a patient as having a RET-associated cancer can include a step of performing an assay on a sample obtained from the patient to determine whether the patient has a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, and identifying or diagnosing a patient determined to have a dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, as having a RET-associated cancer. In some embodiments, the method of selecting a patient for treatment can be used as a part of a clinical study that includes administration of various treatments of a RET-associated cancer. In some embodiments, a RET-associated cancer is a cancer having one or more RET inhibitor resistance mutations. In some embodiments, the assay is an in vitro assay. For example, an assay that utilizes the next generation sequencing, immunohistochemistry, or break apart FISH analysis. In some embodiments, the assay is a regulatory agency-approved, e.g., FDA-approved, kit. In some embodiments, the assay is a liquid biopsy. In some embodiments, the dysregulation of the RET gene, the RET kinase, or expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations.

**[00400]** In some embodiments of any of the methods or uses described herein, an assay used to determine whether the patient has a dysregulation of a RET gene, or a RET kinase, or expression or activity or level of any of the same, using a sample from a patient can include, for example, next generation sequencing, immunohistochemistry, fluorescence microscopy, break apart FISH analysis, Southern blotting, Western blotting, FACS analysis, Northern blotting, and PCR-based amplification (e.g., RT-PCR and quantitative real-time RT-PCR). As is well-known in the art, the assays are typically performed, e.g., with at least one labelled nucleic acid probe or at least one labelled antibody or antigen-binding fragment thereof. Assays can utilize other detection methods known in the art for detecting dysregulation of a RET gene, a RET kinase, or expression or activity or levels of any of the same (see, e.g., the references cited herein). In some embodiments, the dysregulation of the RET gene, the RET kinase, or expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations. In some embodiments, the sample is a biological sample or a biopsy sample (e.g., a paraffin-embedded biopsy sample) from the patient. In some embodiments, the patient is a patient

suspected of having a RET-associated cancer, a patient having one or more symptoms of a RET-associated cancer, and/or a patient that has an increased risk of developing a RET-associated cancer).

**[00401]** In some embodiments, dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same can be identified using a liquid biopsy (variously referred to as a fluid biopsy or fluid phase biopsy). See, e.g., Karachialiou et al., “Real-time liquid biopsies become a reality in cancer treatment”, *Ann. Transl. Med.*, 3(3):36, 2016. Liquid biopsy methods can be used to detect total tumor burden and/or the dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same. Liquid biopsies can be performed on biological samples obtained relatively easily from a subject (e.g., via a simple blood draw) and are generally less invasive than traditional methods used to detect tumor burden and/or dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same. In some embodiments, liquid biopsies can be used to detect the presence of dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same at an earlier stage than traditional methods. In some embodiments, the biological sample to be used in a liquid biopsy can include, blood, plasma, urine, cerebrospinal fluid, saliva, sputum, broncho-alveolar lavage, bile, lymphatic fluid, cyst fluid, stool, ascites, and combinations thereof. In some embodiments, a liquid biopsy can be used to detect circulating tumor cells (CTCs). In some embodiments, a liquid biopsy can be used to detect cell-free DNA. In some embodiments, cell-free DNA detected using a liquid biopsy is circulating tumor DNA (ctDNA) that is derived from tumor cells. Analysis of ctDNA (e.g., using sensitive detection techniques such as, without limitation, next-generation sequencing (NGS), traditional PCR, digital PCR, or microarray analysis) can be used to identify dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same.

**[00402]** In some embodiments, ctDNA derived from a single gene can be detected using a liquid biopsy. In some embodiments, ctDNA derived from a plurality of genes (e.g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more, or any number of genes in between these numbers) can be detected using a liquid biopsy. In some embodiments, ctDNA derived from a plurality of genes can be detected using any of a variety of commercially-available testing panels (e.g., commercially-available testing panels designed to detect dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same). Liquid biopsies can be used to detect dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same including, without limitation, point mutations or single nucleotide variants (SNVs), copy number variants (CNVs), genetic fusions (e.g., translocations or rearrangements), insertions, deletions, or any combination thereof. In some embodiments, a liquid biopsy can be used to detect a germline mutation. In some embodiments, a liquid biopsy can be used to detect a somatic mutation. In some embodiments, a liquid biopsy can be used to detect a primary genetic mutation (e.g., a primary mutation or a primary fusion that is associated with initial development of a disease, e.g., cancer). In some embodiments, a liquid biopsy can be used to detect a genetic mutation that develops after development of the primary genetic mutation (e.g., a resistance mutation that arises in response to a treatment administered to a subject). In some embodiments, a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same identified using a liquid biopsy is also present in a cancer cell that is present in the subject (e.g., in a tumor). In some embodiments, any of the types of dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same described herein can be detected using a liquid biopsy. In some embodiments, a genetic mutation

identified via a liquid biopsy can be used to identify the subject as a candidate for a particular treatment. For example, detection of dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in the subject can indicate that the subject will be responsive to a treatment that includes administration of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00403]** Liquid biopsies can be performed at multiple times during a course of diagnosis, a course of monitoring, and/or a course of treatment to determine one or more clinically relevant parameters including, without limitation, progression of the disease, efficacy of a treatment, or development of resistance mutations after administering a treatment to the subject. For example, a first liquid biopsy can be performed at a first time point and a second liquid biopsy can be performed at a second time point during a course of diagnosis, a course of monitoring, and/or a course of treatment. In some embodiments, the first time point can be a time point prior to diagnosing a subject with a disease (e.g., when the subject is healthy), and the second time point can be a time point after subject has developed the disease (e.g., the second time point can be used to diagnose the subject with the disease). In some embodiments, the first time point can be a time point prior to diagnosing a subject with a disease (e.g., when the subject is healthy), after which the subject is monitored, and the second time point can be a time point after monitoring the subject. In some embodiments, the first time point can be a time point after diagnosing a subject with a disease, after which a treatment is administered to the subject, and the second time point can be a time point after the treatment is administered; in such cases, the second time point can be used to assess the efficacy of the treatment (e.g., if the genetic mutation(s) detected at the first time point are reduced in abundance or are undetectable) or to determine the presence of a resistance mutation that has arisen as a result of the treatment. In some embodiments, a treatment to be administered to a subject can include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00404]** In some embodiments, the efficacy of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, can be determined by assessing the allele frequency of a dysregulation of a RET gene in cfDNA obtained from a patient at different time points, e.g., cfDNA obtained from the patient at a first time point and cfDNA obtained from the patient at a second time point, where at least one dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L) is administered to the patient between the first and second time points. Some embodiments of these methods can further include administering to the patient the at least one dose of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, between the first and second time points. For example, a reduction (e.g., a 1% to about a 99% reduction, a 1% to about a 95% reduction, a 1% to about a 90% reduction, a 1% to about a 85% reduction, a 1% to about a 80% reduction, a 1% to about a 75% reduction, a 1% reduction to about a 70% reduction, a 1% reduction to about a 65% reduction, a 1% reduction to about a 60% reduction, a 1% reduction to about a 55% reduction, a 1% reduction to about a 50% reduction, a 1% reduction to about a 45% reduction, a 1% reduction to about a 40% reduction, a 1% reduction to about a 35% reduction, a 1% reduction to about a 30% reduction, a 1% reduction to about a 25% reduction, a 1% reduction to about a 20% reduction, a 1% reduction to about a 15% reduction, a 1% reduction to about a 10% reduction, a 1% to about a 5% reduction, about a 5% to about a 99% reduction, about a 10% to about a 99% reduction, about a 15% to about a 99% reduction, about a 20% to about a 99% reduction, about a 25% to about a 99% reduction, about a 30% to about a

99% reduction, about a 35% to about a 99% reduction, about a 40% to about a 99% reduction, about a 45% to about a 99% reduction, about a 50% to about a 99% reduction, about a 55% to about a 99% reduction, about a 60% to about a 99% reduction, about a 65% to about a 99% reduction, about a 70% to about a 99% reduction, about a 75% to about a 95% reduction, about a 80% to about a 99% reduction, about a 90% reduction to about a 99% reduction, about a 95% to about a 99% reduction, about a 5% to about a 10% reduction, about a 5% to about a 25% reduction, about a 10% to about a 30% reduction, about a 20% to about a 40% reduction, about a 25% to about a 50% reduction, about a 35% to about a 55% reduction, about a 40% to about a 60% reduction, about a 50% reduction to about a 75% reduction, about a 60% reduction to about 80% reduction, or about a 65% to about a 85% reduction) in the allele frequency (AF) of the dysregulation of a RET gene in the cfDNA obtained from the patient at the second time point as compared to the allele frequency (AF) of the dysregulation of a RET gene in the cfDNA obtained from the patient at the first time point indicates that the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, was effective in the subject. In some embodiments, the AF is reduced such that the level is below the detection limit of the instrument. Alternatively, an increase in the allele frequency (AF) of the dysregulation of a RET gene in the cfDNA obtained from the patient at the second time point as compared to the allele frequency (AF) of the dysregulation of a RET gene in the cfDNA obtained from the patient at the first time point indicates that the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, was not effective in the subject (e.g., the subject has developed a resistance mutation to the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof). Some embodiments of these methods can further include, administering additional doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, to a patient in which a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, was determined to be effective. Some embodiments of these methods can further include, administering a different treatment (e.g., a treatment that does not include the administration of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy) to a patient in which a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, was determined not to be effective.

**[00405]** In some examples of these methods, the time difference between the first and second time points can be about 1 day to about 1 year, about 1 day to about 11 months, about 1 day to about 10 months, about 1 day to about 9 months, about 1 day to about 8 months, about 1 day to about 7 months, about 1 day to about 6 months, about 1 day to about 5 months, about 1 day to about 4 months, about 1 day to about 3 months, about 1 day to about 10 weeks, about 1 day to about 2 months, about 1 day to about 6 weeks, about 1 day to about 1 month, about 1 day to about 25 days, about 1 day to about 20 days, about 1 day to about 15 days, about 1 day to about 10 days, about 1 day to about 5 days, about 2 days to about 1 year, about 5 days to about 1 year, about 10 days to about 1 year, about 15 days to about 1 year, about 20 days to about 1 year, about 25 days to about 1 year, about 1 month to about 1 year, about 6 weeks to about 1 year, about 2 months to about 1 year, about 3 months to about 1 year, about 4 months to about 1 year, about 5 months to about 1 year, about 6 months to about 1 year, about 7 months to about 1 year, about 8 months to about 1 year, about 9 months to about 1 year, about 10 months to about 1 year, about 11 months to about 1 year, about 1 day to about 7 days, about 1 day to about 14 days, about 5 days to about 10 days, about 5 days to about 20 days, about 10 days to about 20 days, about 15 days to about 1 month, about 15 days to about 2 months, about 1 week to about 1

month, about 2 weeks to about 1 month, about 1 month to about 3 months, about 3 months to about 6 months, about 4 months to about 6 months, about 5 months to about 8 months, or about 7 months to about 9 months. In some embodiments of these methods, the patient can be previously identified as having a cancer having a dysregulated RET gene (e.g., any of the examples of a dysregulated RET gene described herein). In some embodiments of these methods, a patient can have been previously diagnosed as having any of the types of cancer described herein. In some embodiments of these methods, the patient can have one or more metastases (e.g., one or more brain metastases).

**[00406]** In some of the above embodiments, the cfDNA comprises ctDNA such as RET-associated ctDNA. For example, the cfDNA is ctDNA such as RET-associated ctDNA. In some embodiments, at least some portion of cfDNA is determined to be RET-associated ctDNA, for example, a sequenced and/or quantified amount of the total cfDNA is determined to have a RET fusion and/or a RET resistance mutation.

**[00407]** In the field of medical oncology it is normal practice to use a combination of different forms of treatment to treat each patient with cancer. In medical oncology the other component(s) of such conjoint treatment or therapy in addition to compositions provided herein may be, for example, surgery, radiotherapy, and chemotherapeutic agents, such as other kinase inhibitors, signal transduction inhibitors and/or monoclonal antibodies. For example, a surgery may be open surgery or minimally invasive surgery. Compounds of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof therefore may also be useful as adjuvants to cancer treatment, that is, they can be used in combination with one or more additional therapies or therapeutic agents, for example, a chemotherapeutic agent that works by the same or by a different mechanism of action. In some embodiments, a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, can be used prior to administration of an additional therapeutic agent or additional therapy. For example, a patient in need thereof can be administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof for a period of time and then under go at least partial resection of the tumor. In some embodiments, the treatment with one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof reduces the size of the tumor (e.g., the tumor burden) prior to the at least partial resection of the tumor. In some embodiments, a patient in need thereof can be administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof for a period of time and under one or more rounds of radiation therapy. In some embodiments, the treatment with one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof reduces the size of the tumor (e.g., the tumor burden) prior to the one or more rounds of radiation therapy.

**[00408]** In some embodiments, a patient has a cancer (e.g., a locally advanced or metastatic tumor) that is refractory or intolerant to standard therapy (e.g., administration of a chemotherapeutic agent, such as a first RET inhibitor or a multikinase inhibitor, immunotherapy, or radiation (e.g., radioactive iodine)). In some embodiments, a patient has a cancer (e.g., a locally advanced or metastatic tumor) that is refractory or intolerant to prior therapy (e.g., administration of a chemotherapeutic agent, such as a first RET inhibitor or a multikinase inhibitor, immunotherapy, or radiation (e.g., radioactive iodine)). In some embodiments, a patient has a cancer (e.g., a locally advanced or metastatic tumor) that has no



standard therapy. In some embodiments, a patient is RET-kinase inhibitor naïve. For example, the patient is naïve to treatment with a selective RET-kinase inhibitor. In some embodiments, a patient is not RET-kinase inhibitor naïve.

**[00409]** In some embodiments, a patient has undergone prior therapy. In some embodiments, a patient having NSCLC (e.g., a RET-fusion positive NSCLC) has received treatment with a platinum-based chemotherapy, PD-1/PDL1 immunotherapy, or both prior to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, a patient having a thyroid cancer (e.g., a RET-fusion positive thyroid cancer) has received treatment with one or more of sorafenib, lenvatinib, and radioactive iodine prior to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, a patient having a colorectal cancer (e.g., a RET-fusion positive colorectal cancer) has received treatment with a fluoropyrimidine-based chemotherapy, with or without ant-VEGF-directed therapy or anti-EGFR-directed therapy, prior to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, a patient having a pancreatic cancer (e.g., a RET-fusion positive pancreatic cancer) has received treatment with one or more of a fluoropyrimidine-based chemotherapy, a gemcitabine-based chemotherapy, and a S-1 chemotherapy prior to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, a patient having a breast cancer (e.g., a RET-fusion positive breast cancer) has received treatment with one or more of anthracycline, taxane, HER2-directed therapy, and hormonal therapy prior to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, a patient having a MTC (e.g., a RET-fusion positive MTC cancer) has received treatment with one or more of caboxantinib and vandetanib prior to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00410]** In some embodiments of any the methods described herein, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is administered in combination with a therapeutically effective amount of at least one additional therapeutic agent selected from one or more additional therapies or therapeutic (e.g., chemotherapeutic) agents.

**[00411]** Non-limiting examples of additional therapeutic agents include: other RET-targeted therapeutic agents (i.e. a first or second RET kinase inhibitor), other kinase inhibitors (e.g., receptor tyrosine kinase-targeted therapeutic agents (e.g., Trk inhibitors or EGFR inhibitors)), signal transduction pathway inhibitors, checkpoint inhibitors, modulators of the apoptosis pathway (e.g. obataclax); cytotoxic chemotherapeutics, angiogenesis-targeted therapies, immune-targeted agents, including immunotherapy, and radiotherapy.

**[00412]** In some embodiments, the other RET-targeted therapeutic is a multikinase inhibitor exhibiting RET inhibition activity. In some embodiments, the other RET-targeted therapeutic inhibitor is selective for a RET kinase. Exemplary RET kinase inhibitors can exhibit inhibition activity ( $IC_{50}$ ) against a RET kinase of less than about 1000 nM, less than about 500 nM, less than about 200 nM, less than about 100 nM, less than about 50 nM, less than about 25 nM, less than about 10 nM, or less than about 1 nM as measured in an assay as described herein. In some embodiments, a RET kinase inhibitors can exhibit


inhibition activity (IC<sub>50</sub>) against a RET kinase of less than about 25 nM, less than about 10 nM, less than about 5 nM, or less than about 1 nM as measured in an assay as provided herein.

**[00413]** Non-limiting examples of RET-targeted therapeutic agents (e.g., a first RET inhibitor or a second RET inhibitor) include alectinib (9-Ethyl-6,6-dimethyl-8-[4-(morpholin-4-yl)piperidin-1-yl]-11-oxo-6,11-dihydro-5H-benzo[b]carbazole-3-carbonitrile); amuvatinib (MP470, HPK56) (N-(1,3-benzodioxol-5-ylmethyl)-4-([1]benzofuro[3,2-d]pyrimidin-4-yl)piperazine-1-carbothioamide); apatinib (YN968D1) (N-[4-(1-cyanocyclopentyl) phenyl-2-(4-picolyl)amino-3-Nicotinamide methanesulphonate]; cabozantinib (Cometriq XL-184) (N-(4-((6,7-Dimethoxyquinolin-4-yl)oxy)phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide); dovitinib (TKI258; GFKI-258; CHIR-258) ((3Z)-4-amino-5-fluoro-3-[5-(4-methylpiperazin-1-yl)-1,3-dihydrobenzimidazol-2-ylidene]quinolin-2-one); famitinib (5-[2-(diethylamino)ethyl]-2-[(Z)-(5-fluoro-2-oxo-1H-indol-3-ylidene)methyl]-3-methyl-6,7-dihydro-1H-pyrrolo[3,2-c]pyridin-4-one); fedratinib (SAR302503, TG101348) (N-(2-Methyl-2-propanyl)-3-([5-methyl-2-([4-[2-(1-pyrrolidinyl)ethoxy]phenyl)amino]-4-pyrimidinyl]amino)benzenesulfonamide); foretinib (XL880, EXEL-2880, GSK1363089, GSK089) (N1'-[3-fluoro-4-[[6-methoxy-7-(3-morpholinopropoxy)-4-quinolyl]oxy]phenyl]-N1-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide); fostamantinib (R788) (2H-Pyrido[3,2-b]-1,4-oxazin-3(4H)-one, 6-[[5-fluoro-2-[(3,4,5-trimethoxyphenyl)amino]-4-pyrimidinyl]amino]-2,2-dimethyl-4-[(phosphonoxy)methyl]-, sodium salt (1:2)); ilorasertib (ABT-348) (1-(4-(4-amino-7-(1-(2-hydroxyethyl)-1H-pyrazol-4-yl)thieno[3,2-c]pyridin-3-yl)phenyl)-3-(3-fluorophenyl)urea); lenvatinib (E7080, Lenvima) (4-[3-chloro-4-(cyclopropylaminocarbonyl)aminophenoxy]-7-methoxy-6-quinolinecarboxamide); motesanib (AMG 706) (N-(3,3-Dimethyl-2,3-dihydro-1H-indol-6-yl)-2-[(pyridin-4-ylmethyl)amino]pyridine-3-carboxamide); nintedanib (3-Z-[1-(4-(N-(4-methyl-piperazin-1-yl)-methylcarbonyl)-N-methyl-amino)-anilino]-1-phenyl-methylene]-6-methoxycarbonyl-2-indolinone); ponatinib (AP24534) (3-(2-Imidazo[1,2-b]pyridazin-3-ylethynyl)-4-methyl-N-[4-[(4-methylpiperazin-1-yl)methyl]-3-(trifluoromethyl)phenyl]benzamide); PP242 (a TORKinib) (2-[4-Amino-1-(1-methylethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl]-1H-indol-5-ol); quizartinib (1-(5-(tert-Butyl)isoxazol-3-yl)-3-(4-(7-(2-morpholinoethoxy)benzo[d]imidazo[2,1-b]thiazol-2-yl)phenyl)urea); regorfenib (BAY 73-4506, stivarga) (4-[4-([4-Chloro-3-(trifluoromethyl)phenyl]carbonyl)amino]-3-fluorophenoxy]-N-methylpyridine-2-carboxamide hydrate); RDX-105 (CEP-32496, agerafenib) (1-(3-((6,7-dimethoxyquinazolin-4-yl)oxy)phenyl)-3-(5-(1,1,1-trifluoro-2-methylpropan-2-yl)isoxazol-3-yl)urea); semaxanib (SU5416) ((3Z)-3-[(3,5-dimethyl-1H-pyrrol-2-yl)methylidene]-1,3-dihydro-2H-indol-2-one); sitravatinib (MGCD516, MG516) (N-(3-Fluoro-4-[[2-(5-[(2-methoxyethyl)amino]methyl)-2-pyridinyl]thieno[3,2-b]pyridin-7-yl]oxy)phenyl)-N'-(4-fluorophenyl)-1,1-cyclopropanedicarboxamide); sorafenib (BAY 43-9006) (4-[4-[[[4-chloro-3-(trifluoromethyl)phenyl]amino]carbonyl]amino]phenoxy]-N-methyl-2-pyridinecarboxamide); vandetanib (N-(4-bromo-2-fluorophenyl)-6-methoxy-7-[(1-methylpiperidin-4-yl)methoxy]quinazolin-4-amine); vatalanib (PTK787, PTK/ZK, ZK222584) (N-(4-chlorophenyl)-4-(pyridin-4-ylmethyl)phthalazin-1-amine); AD-57 (N-[4-[4-amino-1-(1-methylethyl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl]phenyl]-N'-[3-(trifluoromethyl)phenyl]-urea); AD-80 (1-[4-(4-amino-1-propan-2-ylpyrazolo[3,4-d]pyrimidin-3-yl)phenyl]-3-[2-fluoro-5-(trifluoromethyl)phenyl]urea); AD-81 (1-(4-(4-amino-1-isopropyl-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)-3-(4-chloro-3-(trifluoromethyl)phenyl)urea); ALW-II-41-27 (N-(5-((4-(4-ethylpiperazin-1-yl)methyl)-3-(trifluoromethyl)phenyl)carbonyl)-2-methylphenyl)-5-(thiophen-2-yl)nicotinamide); BPR1K871 (1-(3-chlorophenyl)-3-(5-(2-((7-(3-(dimethylamino)propoxy)quinazolin-4-yl)amino)ethyl)thiazol-2-yl)urea); CLM3 (1-phenethyl-N-(1-phenylethyl)-1H-pyrazolo[3,4-d]pyrimidin-4-

amine); EBI-907 (N-(2-chloro-3-(1-cyclopropyl-8-methoxy-3H-pyrazolo[3,4-c]isoquinolin-7-yl)-4-fluorophenyl)-3-fluoropropane-1-sulfonamide); NVP-AST-487 (N-[4-[(4-ethyl-1-piperazinyl)methyl]-3-(trifluoromethyl)phenyl]-N'-[4-[[6-(methylamino)-4-pyrimidinyl]oxy]phenyl]-urea); NVP-BBT594 (BBT594) (5-((6-acetamidopyrimidin-4-yl)oxy)-N-(4-((4-methylpiperazin-1-yl)methyl)-3-(trifluoromethyl)phenyl)indoline-1-carboxamide); PD173955 (6-(2,6-dichlorophenyl)-8-methyl-2-(3-methylsulfanylanilino)pyrido[2,3-d]pyrimidin-7-one); PP2 (4-amino-5-(4-chlorophenyl)-7-(dimethylethyl)pyrazolo[3,4-d]pyrimidine); PZ-1 (N-(5-(tert-butyl)isoxazol-3-yl)-2-(4-(5-(1-methyl-1H-pyrazol-4-yl)-1Hbenzo[d]imidazol-1-yl)phenyl)acetamide); RPI-1 (1,3-dihydro-5,6-dimethoxy-3-[(4-hydroxyphenyl)methylene]-H-indol-2-one); (3E)-3-[(4-hydroxyphenyl)methylidene]-5,6-dimethoxy-1H-indol-2-one); SGI-7079 (3-[2-[[3-fluoro-4-(4-methyl-1-piperazinyl)phenyl]amino]-5-methyl-7H-pyrrolo[2,3-d]pyrimidin-4-yl]-benzeneacetoneitrile); SPP86 (1-Isopropyl-3-(phenylethynyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine); SU4984 (4-[4-[(E)-(2-oxo-1H-indol-3-ylidene)methyl]phenyl]piperazine-1-carbaldehyde); sunitinib (SU11248) (N-(2-Diethylaminoethyl)-5-[(Z)-(5-fluoro-2-oxo-1H-indol-3-ylidene)methyl]-2,4-dimethyl-1H-pyrrole-3-carboxamide); TG101209 (N-tert-butyl-3-(5-methyl-2-(4-(4-methylpiperazin-1-yl)phenylamino)pyrimidin-4-ylamino)benzenesulfonamide); Withaferin A ((4 $\beta$ ,5 $\beta$ ,6 $\beta$ ,22R)-4,27-Dihydroxy-5,6:22,26-diepoxyergosta-2,24-diene-1,26-dione); XL-999 ((Z)-5-((1-ethylpiperidin-4-yl)amino)-3-((3-fluorophenyl)(5-methyl-1H-imidazol-2-yl)methylene)indolin-2-one); BPR1J373 (a 5-phenylthiazol-2-ylamine-pyrimidine derivative); CG-806 (CG'806); DCC-2157; GTX-186; HG-6-63-01 ((E)-3-(2-(4-chloro-1H-pyrrolo[2,3-b]pyridin-5-yl)vinyl)-N-(4-((4-ethylpiperazin-1-yl)methyl)-3-(trifluoromethyl)phenyl)-4-methylbenzamide); SW-01 (Cyclobenzaprine hydrochloride); XMD15-44 (N-(4-((4-ethylpiperazin-1-yl)methyl)-3-(trifluoromethyl)phenyl)-4-methyl-3-(pyridin-3-ylethynyl)benzamide (generated from structure)); Y078-DM1 (an antibody drug conjugate composed of a RET antibody (Y078) linked to a derivative of the cytotoxic agent maytansine); Y078-DM4 (an antibody drug conjugate composed of a RET antibody (Y078) linked to a derivative of the cytotoxic agent maytansine); ITRI-305 (DON5TB, DIB003599); BLU-667 (((1S,4R)-N-(S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide); BLU6864; DS-5010; GSK3179106; GSK3352589; and NMS-E668.


**[00414]** Further examples of RET-targeted therapeutics (e.g., a first RET kinase inhibitor or a second RET kinase inhibitor) include 5-amino-3-(5-cyclopropylisoxazol-3-yl)-1-isopropyl-1H-pyrazole-4-carboxamide; 3-(5-cyclopropylisoxazol-3-yl)-1-isopropyl-1H-pyrazolo[3,4-d]pyrimidin-4-amine; 3-((6,7-Dimethoxyquinazolin-4-yl)amino)-4-fluoro-2-methylphenol; N-(5-(tert-butyl)isoxazol-3-yl)-2-(4-(imidazo[1,2-a]pyridin-6-yl)phenyl)acetamide; N-(5-(tert-butyl)isoxazol-3-yl)-2-(3-(imidazo[1,2-b]pyridazin-6-yloxy)phenyl)acetamide; N-(2-fluoro-5-trifluoromethylphenyl)-N'-{4'-[(2''-benzamido)pyridin-4''-ylamino]phenyl}urea; 2-amino-6-[[2-(4-chlorophenyl)-2-oxoethyl]sulfanyl]-4-(3-thienyl)pyridine-3,5-dicarbonitrile; and 3-arylureidobenzylidene-indolin-2-ones.

**[00415]** Additional examples of other RET kinase inhibitors include those described in U.S. Patent Nos. 9,150,517 and 9,149,464, and International Publication No. WO 2014075035, all of which are hereby incorporated by reference. For example, in some embodiments the other RET inhibitor is a compound of formula I:



wherein  $R_1$  is  $C_6$ - $C_{24}$ alkyl or polyethylene glycol; or a pharmaceutically acceptable salt form thereof. In some embodiments, the other RET inhibitor is 4-{5-[bis-(chloroethyl)-amino]-1-methyl-1H-benzimidazol-2-yl}butyric acid dodecyl ester.

**[00416]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO 2016127074, which is hereby incorporated by reference. For example, in some embodiments, the other RET inhibitor is a compound of Formula (I) or a pharmaceutically acceptable salt thereof, wherein:



wherein Rings A and B are each independently selected from aryl, heteroaryl, cycloalkyl and heterocyclyl;

each  $L^1$  and  $L^2$  is independently selected from a bond, -(C1-C6 alkylene)-, -(C2-C6alkenylene)-, -(C2-C6 alkynylene)-, -(C1-C6 haloalkylene)-, -(C1-C6 heteroalkylene)-, -C(O)-, -O-, -S-, -S(O), -S(O)<sub>2</sub>-, -N(R<sup>1</sup>)-, -O-(C1-C6 alkylene)-, -(C1-C6 alkylene)-O-, -N(R<sup>1</sup>)-C(O)-, -C(O)N(R<sup>1</sup>)-, -(C1-C6 alkylene)-N(R<sup>1</sup>)-, -N(R<sup>1</sup>)-(C1-C6 alkylene)-, -N(R<sup>1</sup>)-C(O)-(C1-C6 alkylene)-, -(C1-C6 alkylene)-N(R<sup>1</sup>)-C(O)-, -C(O)-N(R<sup>1</sup>)-(C1-C6 alkylene)-, -(C1-C6 alkylene)-C(O)-N(R<sup>1</sup>)-, -N(R<sup>1</sup>)-S(O)<sub>2</sub>-, -S(O)<sub>2</sub>-N(R<sup>1</sup>)-, -N(R<sup>1</sup>)-S(O)<sub>2</sub>-(C1-C6 alkylene)-, and -S(O)<sub>2</sub>-N(R<sup>1</sup>)-(C1-C6 alkylene)-; wherein each alkylene, alkenylene, alkynylene, haloalkylene, and heteroalkylene is independently substituted with 0-5 occurrences of  $R^1$ ;

each  $R^A$  and  $R^B$  is independently selected from C1-C6 alkyl, C1-C6 alkoxy, halo, C1-C6 haloalkyl, C1-C6 hydroxyalkyl, C1-C6 heteroalkyl, and -N(R<sup>1</sup>)(R<sup>1</sup>); wherein each alkyl, alkoxy, haloalkyl, hydroxyalkyl, and hydroxyalkyl is independently substituted with 0-5 occurrences of  $R_a$ ;

each  $R^C$  and  $R^D$  is independently selected from C1-C6 alkyl, C2-C6 alkenyl, C2-C6 alkynyl, C1-C6 alkoxy, halo, C1-C6 heteroalkyl, C1-C6 haloalkyl, C1-C6 haloalkoxy, C1-C6 hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, heterocyclylalkyl, nitro, cyano, -C(O)R<sup>1</sup>, -OC(O)R<sup>1</sup>, -C(O)OR<sup>1</sup>, -(C1-C6 alkylene)-C(O)R<sup>1</sup>, -SR<sup>1</sup>, -S(O)<sub>2</sub>R<sup>1</sup>, -S(O)<sub>2</sub>-N(R<sup>1</sup>)(R<sup>1</sup>), -(C1-C6 alkylene)-S(O)<sub>2</sub>R<sup>1</sup>, -(C1-C6 alkylene)-S(O)<sub>2</sub>-N(R<sup>1</sup>)(R<sup>1</sup>), -N(R<sup>1</sup>)(R<sup>1</sup>)-C(O)-N(R<sup>1</sup>)(R<sup>1</sup>)-N(R<sup>1</sup>)-C(O)R<sup>1</sup>, -N(R<sup>1</sup>)-C(O)OR<sup>1</sup>, -(C1-C6 alkylene)-N(R<sup>1</sup>)-C(O)R<sup>1</sup>, -N(R<sup>1</sup>)S(O)<sub>2</sub>R<sup>1</sup>, and -P(O)(R<sup>1</sup>)(R<sup>1</sup>); wherein each of alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, haloalkyl, haloalkoxy, hydroxyalkyl, cycloalkyl, aryl, heteroaryl, aryloxy, aralkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of  $R^a$ ; or 2  $R^C$  or 2  $R^D$  together with the carbon atom(s)

to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of  $R^a$ ;

each  $R^1$  is independently selected from hydrogen, hydroxyl, halo, thiol, C1-C6 alkyl, C1-C6 thioalkyl, C1-C6 alkoxy, C1-C6 haloalkyl, C1-C6 hydroxyalkyl, C1-C6 heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl, wherein each of alkyl, thioalkyl, alkoxy, haloalkyl, hydroxyalkyl, heteroalkyl, cycloalkyl, cycloalkylalkyl, heteroarylalkyl, heterocyclyl, and heterocyclylalkyl is independently substituted with 0-5 occurrences of  $R^b$ , or 2  $R^1$  together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring independently substituted with 0-5 occurrences of  $R^b$ ;


each  $R^a$  and  $R^b$  is independently C1-C6 alkyl, halo, hydroxyl, C1-C6 haloalkyl, C1-C6 heteroalkyl, C1-C6 hydroxyalkyl, C1-C6 alkoxy, cycloalkyl, heterocyclyl, or cyano, wherein each of alkyl, haloalkyl, heteroalkyl, hydroxyalkyl, alkoxy, cycloalkyl and heterocyclyl is independently substituted with 0-5 occurrences of  $R^1$ ;


each  $R^1$  is C1-C6 alkyl, C1-C6 heteroalkyl, halo, hydroxyl, C1-C6 haloalkyl, C1-C6 hydroxyalkyl, cycloalkyl or cyano; or 2  $R^1$ , together with the atom(s) to which they are attached form a cycloalkyl or heterocyclyl ring;








$m$  is 0, 1, 2, or 3;








$n$  is 0, 1, or 2; and

$p$  and  $q$  are each independently 0, 1, 2, 3, or 4. For example, a RET inhibitor can be selected from the group consisting of:
























, or a pharmaceutically acceptable salt thereof.

**[00417]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO 2016075224, which is hereby incorporated by reference. For example, in some embodiments, the other RET inhibitor is a compound of Formula (II) or a pharmaceutically acceptable salt thereof, wherein:



R1 and R2 are independently hydrogen or an optionally substituted group selected from straight or branched (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>3</sub>-C<sub>6</sub>) cycloalkyl and COR', wherein R' is an optionally substituted group selected from straight or branched (C<sub>1</sub>-C<sub>6</sub>) alkyl and (C<sub>3</sub>-C<sub>6</sub>) cycloalkyl;

R3 is hydrogen or an optionally substituted group selected from straight or branched (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>2</sub>-C<sub>6</sub>) alkenyl, (C<sub>2</sub>-C<sub>6</sub>) alkynyl, (C<sub>3</sub>-C<sub>6</sub>) cycloalkyl, aryl, heteroaryl and a 3- to 7-membered heterocyclyl ring;

R4 is hydrogen or an optionally substituted group selected from straight or branched (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>2</sub>-C<sub>6</sub>) alkenyl, aryl, heteroaryl or heterocyclyl;

A is a 5- or 6-membered heteroaryl ring or a phenyl ring;

B is a 5- or 6-membered ring selected from heteroaryl, (C<sub>5</sub>-C<sub>6</sub>) cycloalkyl and heterocyclyl ring or a phenyl ring; wherein ring A and ring B are fused together to form a bicyclic system comprising a 6-


membered aromatic or 5- to 6-membered heteroaromatic ring fused with a 6-membered aromatic or 5- to 6-membered heteroaromatic, (C<sub>5</sub>-C<sub>6</sub>) cycloalkyl or heterocyclyl ring;

Y is carbon or nitrogen;

X is hydrogen, halogen, hydroxyl, cyano or an optionally substituted group selected from straight or branched (C<sub>1</sub>-C<sub>6</sub>) alkyl and (C<sub>1</sub>-C<sub>6</sub>) alkoxy; and

R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or an optionally substituted group selected from straight or branched (C<sub>1</sub>-C<sub>6</sub>) alkyl, (C<sub>3</sub>-C<sub>6</sub>) cycloalkyl, heterocyclyl, aryl and heteroaryl.

**[00418]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO 2015079251, which is hereby incorporated by reference. For example, in some embodiments, the other RET inhibitor is a compound of Formula (III) or a pharmaceutically acceptable salt or solvate thereof, wherein:



X is NH, NR<sub>x</sub>, O or S, wherein R<sub>x</sub> is (1-3C)alkyl;


R<sub>1</sub> is selected from halo (e.g., fluoro, chloro, or bromo), trifluoromethyl, (1-4C)alkyl (e.g., methyl), (1-4C)alkoxy or (3-6C)cycloalkyl, wherein an alkyl, alkoxy or cycloalkyl group is optionally substituted with one or more fluoro;

R<sub>2</sub> is selected from hydrogen, halo (e.g., fluoro, chloro or bromo), hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, (1-6C)alkyl (e.g., methyl), (3-8C)cycloalkyl, or (1-4C)alkoxy (e.g., OMe), wherein an alkyl, cycloalkyl or alkoxy group is optionally substituted with one or more fluoro;

R<sub>3</sub> is selected from hydrogen, halo (e.g. fluoro, chloro or bromo), hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, (1-6C)alkyl (e.g., methyl), (3-8C)cycloalkyl, or (1-4C)alkoxy (e.g., OMe), wherein an alkyl, cycloalkyl or alkoxy group is optionally substituted with one or more fluoro;

R<sub>4</sub> is selected from hydrogen, halo (e.g., fluoro, chloro or bromo), hydroxyl, cyano, trifluoromethyl, trifluoromethoxy, (1-6C)alkyl (e.g., methyl), (3-8C)cycloalkyl, or (1-4C)alkoxy (e.g., OMe), wherein an alkyl, cycloalkyl or alkoxy group is optionally substituted with one or more fluoro;

R<sub>5</sub> is selected from hydrogen or a group defined by the formula:




wherein

$L_5$  is absent or a linear or branched (1-4C)alkylene;

$X_5$  is absent or -C(O)O-, -O-, -C(O)-, -OC(O)-, -CH(QR<sub>5L</sub>)-, -N(R<sup>i</sup>)-, -N(R<sub>5L</sub>)-C(O)-, -N(R<sub>5L</sub>)-C(O)O-, -C(O)-N(R<sub>5L</sub>)-, -S-, -SO-, -SO<sub>2</sub>-, -S(O)<sub>2</sub>N(R<sub>5L</sub>)-, or -N(R<sub>5L</sub>)SO<sub>2</sub>- wherein R<sub>5L</sub> is selected from hydrogen or methyl; and

$Q_5$  is (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, (3-8C)cycloalkyl, (3-8C)cycloalkyl-(1-4C)alkyl, aryl, aryl-(1-4C)alkyl, heteroaryl, heteroaryl-(1-4C)alkyl, heterocyclyl or heterocyclyl-(1-4C)alkyl;

$R_6$  is selected from hydrogen, or a group defined by the formula:



wherein

$L_6$  is absent or a linear or branched (1-4C)alkylene;


$X_6$  is absent or selected from -O-, -C(O)-, -C(O)O-, -OC(O)-, -CH(OR<sub>6L</sub>)-, -N(R<sub>6L</sub>)-, -N(R<sub>6L</sub>)-C(O)-, -N(R<sub>6L</sub>)-C(O)O-, -C(O)-N(R<sub>6L</sub>)-, -S-, -SO-, -SO<sub>2</sub>-, -S(O)<sub>2</sub>N(R<sub>6L</sub>)-, or -N(R<sub>6L</sub>)SO<sub>2</sub>- wherein R<sub>6L</sub> is selected from hydrogen or (1-3C)alkyl;

$Q_6$  is hydrogen, (1-8C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (3-8C)cycloalkyl, (3-8C)cycloalkyl-(1-6C)alkyl, aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl, heterocyclyl-(1-6C)alkyl,

or  $Q_6$  and  $R_{L6}$  are linked such that, together with the nitrogen atom to which they are attached, they form a heterocyclic ring;

wherein  $R_6$  is optionally substituted (e.g. substituted on  $L_6$  and/or  $Q_6$ ) with one or more (1-6C)alkyl, (1-6C)alkanoyl, OR<sub>6X</sub>, SR<sub>6X</sub>, S(O)R<sub>6X</sub>, S(O)<sub>2</sub>R<sub>6X</sub>, C(O)OR<sub>6X</sub> or C(O)NR<sub>6X</sub>R'<sub>6X</sub>, wherein R<sub>6X</sub> and R'<sub>6X</sub> are independently hydrogen, (1-8C)alkyl, or R<sub>6X</sub> and R'<sub>6X</sub> are linked such that, together with the nitrogen atom to which they are attached, they form a heterocyclic ring; and

$R_7$  is selected from hydrogen, (1-6C)alkoxy, or a group defined by the formula:



wherein

$L_7$  is absent or a linear or branched (1-4C)alkylene;

$X_7$  is absent or selected from -O-, -C(O)-, -C(O)O-, -OC(O)-, -CH(OR<sub>6L</sub>)-, -N(R<sub>7L</sub>)-, -N(R<sub>7L</sub>)-C(O)-, -N(R<sub>7L</sub>)-C(O)O-, -C(O)-N(R<sub>7L</sub>)-, -S-, -SO-, -SO<sub>2</sub>-, -S(O)<sub>2</sub>N(R<sub>7L</sub>)-, or -N(R<sub>7L</sub>)SO<sub>2</sub>- wherein R<sub>7L</sub> is selected from hydrogen or (1-3C)alkyl;


$Q_7$  is hydrogen, (1-8C)alkyl, (2-8C)alkenyl, (2-8C)alkynyl, (3-8C)cycloalkyl, (3-8C)cycloalkyl-(1-6C)alkyl, aryl, aryl-(1-6C)alkyl, heteroaryl, heteroaryl-(1-6C)alkyl, heterocyclyl, heterocyclyl-(1-6C)alkyl,

or  $Q_7$  and R<sub>7L</sub> are linked such that, together with the nitrogen atom to which they are attached, they form a heterocyclic ring;

wherein R<sub>7</sub> is optionally substituted (e.g., substituted on L<sub>7</sub> and/or Q<sub>7</sub>) with one or more halo, hydroxyl, nitro, cyano, (1-8C)alkyl, (1-8C)alkanoyl, OR<sub>7X</sub>, SR<sub>7X</sub>, S(O)R<sub>7X</sub>, S(O)<sub>2</sub>R<sub>7X</sub>, C(O)OR<sub>7X</sub> or C(O)NR<sub>7X</sub>R'<sub>7X</sub>, wherein R<sub>7X</sub> and R'<sub>7X</sub> are independently hydrogen, (1-8C)alkyl, or R<sub>7X</sub> and R'<sub>7X</sub> are linked such that, together with the nitrogen atom to which they are attached, they form a heterocyclic ring; or


R<sub>7</sub> is optionally substituted with one or more groups selected from oxo, (1-4C)haloalkyl, (1-4C)hydroxyalkyl, C(O)R<sub>7Y</sub> or NR<sub>7Y</sub>R'<sub>7Y</sub>, wherein R<sub>7Y</sub> and R'<sub>7Y</sub> are independently hydrogen or (1-8C)alkyl.

**[00419]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO2017178845, which is hereby incorporated by reference. For example, in some embodiments, the other RET inhibitor is a compound of Formula (IV) or a pharmaceutically acceptable salt thereof, wherein:



(IV)

HET is selected from one of the following:



wherein  denotes the point of attachment;

$R_1$  is selected from hydrogen, (1-4C)haloalkyl, (1-4C)haloalkoxy or a group of the formula:



wherein:

L is absent or (1-5C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;

Y is absent or O, S, SO, SO<sub>2</sub>, N(R<sub>a</sub>), C(O), C(O)O, OC(O), C(O)N(R<sub>a</sub>), N(R<sub>a</sub>)C(O), N(R<sub>a</sub>)C(O)N(R<sub>b</sub>), N(R<sub>a</sub>)C(O)O, OC(O)N(R<sub>a</sub>), S(O)<sub>2</sub>N(R<sub>a</sub>), or N(R<sub>a</sub>)SO<sub>2</sub>, wherein R<sub>a</sub> and R<sub>b</sub> are each independently selected from hydrogen or (1-4C)alkyl; and

Q is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, aryl, (3-10C)cycloalkyl, (3-10C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Q is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, (1-4C)aminoalkyl, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl,

mercapto, ureido,  $\text{NR}_c\text{R}_d$ ,  $\text{OR}_c$ ,  $\text{C(O)R}_c$ ,  $\text{C(O)OR}_c$ ,  $\text{OC(O)R}_c$ ,  $\text{C(O)N(R}_d\text{)R}_c$ ,  $\text{N(R}_d\text{)C(O)R}_c$ ,  $\text{S(O)}_p\text{R}_c$  (where  $p$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_d\text{)R}_c$ ,  $\text{N(R}_d\text{)SO}_2\text{R}_c$ ,  $\text{Si(R}_e\text{)(R}_d\text{)R}_c$  or  $(\text{CH}_2)_q\text{NR}_c\text{R}_d$  (where  $q$  is 1, 2 or 3); wherein  $\text{R}_c$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from hydrogen, (1-6C)alkyl or (3-6C)cycloalkyl; or  $\text{R}_c$  and  $\text{R}_d$  are linked such that, together with the nitrogen atom to which they are attached, they form a 4-7 membered heterocyclic ring which is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano or hydroxy; or

Q is optionally substituted by a group of the formula:



wherein:

$\text{L}_1$  is absent or (1-3C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;

$\text{L}_{\text{Q1}}$  is absent or selected from O, S, SO,  $\text{SO}_2$ ,  $\text{N(R}_f\text{)}$ ,  $\text{C(O)}$ ,  $\text{C(O)O}$ ,  $\text{OC(O)}$ ,  $\text{C(O)N(R}_f\text{)}$ ,  $\text{N(R}_f\text{)C(O)}$ ,  $\text{N(R}_f\text{)C(O)N(R}_g\text{)}$ ,  $\text{N(R}_f\text{)C(O)O}$ ,  $\text{OC(O)N(R}_f\text{)}$ ,  $\text{S(O)}_2\text{N(R}_f\text{)}$ , or  $\text{N(R}_f\text{)SO}_2$ , wherein  $\text{R}_f$  and  $\text{R}_g$  are each independently selected from hydrogen or (1-2C)alkyl; and

$\text{W}_1$  is hydrogen, (1-6C)alkyl, aryl, aryl(1-2C)alkyl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein  $\text{W}_1$  is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl, mercapto, ureido, aryl, heteroaryl, heterocyclyl, (3-6C)cycloalkyl,  $\text{NR}_h\text{R}_i$ ,  $\text{OR}_h$ ,  $\text{C(O)R}_h$ ,  $\text{C(O)OR}_h$ ,  $\text{OC(O)R}_h$ ,  $\text{C(O)N(R}_i\text{)R}_h$ ,  $\text{N(R}_i\text{)C(O)R}_h$ ,  $\text{S(O)}_r\text{R}_h$  (where  $r$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_i\text{)R}_h$ ,  $\text{N(R}_i\text{)SO}_2\text{R}_h$  or  $(\text{CH}_2)_s\text{NR}_i\text{R}_h$  (where  $s$  is 1, 2 or 3); wherein  $\text{R}_h$  and  $\text{R}_i$  are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl;

$\text{R}_{1a}$  and  $\text{R}_{1b}$  are each selected from H, (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl or mercapto;


W is selected from O, S or  $\text{NR}_{\text{W1}}$ , wherein  $\text{R}_{\text{W1}}$  is selected from H or (1-2C)alkyl;

$\text{X}_1$ ,  $\text{X}_2$ ,  $\text{X}_3$  and  $\text{X}_4$  are independently selected from CH,  $\text{CR}_2$  or N;

$\text{R}_2$  is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, nitro, aryl, heteroaryl, heterocyclyl, cycloalkyl, (2-4C)alkynyl,  $\text{NR}_j\text{R}_k$ ,  $\text{OR}_j$ ,  $\text{C(O)R}_j$ ,  $\text{C(O)OR}_j$ ,  $\text{OC(O)R}_j$ ,  $\text{C(O)N(R}_k\text{)R}_j$ ,  $\text{N(R}_k\text{)C(O)R}_j$ ,  $\text{N(R}_k\text{)C(O)N(R}_j\text{)}$ ,  $\text{S(O)}_{r1}\text{R}_k$  (where  $r_1$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_j\text{)R}_k$ ,  $\text{N(R}_j\text{)SO}_2\text{R}_k$  or  $(\text{CH}_2)_v\text{NR}_j\text{R}_k$  (where  $v$  is 1, 2 or 3); wherein  $\text{R}_j$  and  $\text{R}_k$  are each independently selected from hydrogen or (1-4C)alkyl; and wherein said (1-4C)alkyl, aryl, heteroaryl, heterocyclyl or cycloalkyl is optionally substituted by one or more substituents selected from halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, nitro, phenyl, (2-4C)alkynyl,  $\text{NR}_{j1}\text{R}_{k1}$ ,  $\text{OR}_{j1}$ ,  $\text{C(O)R}_{j1}$ ,  $\text{C(O)OR}_{j1}$ ,  $\text{OC(O)R}_{j1}$ ,  $\text{C(O)N(R}_{k1}\text{)R}_{j1}$ ,  $\text{N(R}_{k1}\text{)C(O)R}_{j1}$ ,  $\text{S(O)}_{r2}\text{R}_h$  (where  $r_2$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_{j1}\text{)R}_{k1}$ ,  $\text{N(R}_{j1}\text{)SO}_2\text{R}_{k1}$  or  $(\text{CH}_2)_{v1}\text{NR}_{j1}\text{R}_{k1}$  (where  $v_1$  is 1, 2 or 3); and wherein  $\text{R}_{j1}$  and  $\text{R}_{k1}$  are each independently selected from hydrogen or (1-4C)alkyl; and

$R_3$  is selected from halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, nitro, (2-4C)alkynyl,  $NR_iR_m$ ,  $OR_i$ ,  $C(O)R_i$ ,  $C(O)OR_i$ ,  $OC(O)R_i$ ,  $C(O)N(R_m)R_i$ ,  $N(R_m)C(O)R_i$ , or  $(CH_2)_yNR_iR_m$  (where  $y$  is 1, 2 or 3); wherein said (1-4C)alkyl is optionally substituted by one or more substituents selected from amino, hydroxy, (1-2C)alkoxy or halo; and wherein  $R_i$  and  $R_m$  are each independently selected from hydrogen or (1-4C)alkyl.


**[00420]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO2017178844, which is hereby incorporated by reference. For example, in some embodiments, the other RET inhibitor is a compound of Formula (V) or a pharmaceutically acceptable salt thereof, wherein:



(V)

HET is selected from one of the following:





wherein  denotes the point of attachment;

R<sub>1</sub> is selected from hydrogen, (1-4C)haloalkyl, (1-4C)haloalkoxy or a group of the formula:



wherein:


L is absent or (1-5C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;

Y is absent or O, S, SO, SO<sub>2</sub>, N(R<sub>a</sub>), C(O), C(O)O, OC(O), C(O)N(R<sub>a</sub>), N(R<sub>a</sub>)C(O), N(R<sub>a</sub>)C(O)N(R<sub>b</sub>), N(R<sub>a</sub>)C(O)O, OC(O)N(R<sub>a</sub>), S(O)<sub>2</sub>N(R<sub>a</sub>), or N(R<sub>a</sub>)SO<sub>2</sub>, wherein R<sub>a</sub> and R<sub>b</sub> are each independently selected from hydrogen or (1-4C)alkyl; and

Q is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, aryl, (3-10C)cycloalkyl, (3-10C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Q is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, (1-4C)aminoalkyl, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl,

mercapto, ureido,  $\text{NR}_c\text{R}_d$ ,  $\text{OR}_c$ ,  $\text{C(O)R}_c$ ,  $\text{C(O)OR}_c$ ,  $\text{OC(O)R}_c$ ,  $\text{C(O)N(R}_d\text{)R}_c$ ,  $\text{N(R}_d\text{)C(O)R}_c$ ,  $\text{S(O)}_y\text{R}_c$  (where  $y$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_d\text{)R}_c$ ,  $\text{N(R}_d\text{)SO}_2\text{R}_c$ ,  $\text{Si(R}_d\text{)(R}_c\text{)R}_e$  or  $(\text{CH}_2)_z\text{NR}_c\text{R}_d$  (where  $z$  is 1, 2 or 3); wherein  $\text{R}_c$ ,  $\text{R}_d$  and  $\text{R}_e$  are each independently selected from hydrogen, (1-6C)alkyl or (3-6C)cycloalkyl; or  $\text{R}_c$  and  $\text{R}_d$  can be linked such that, together with the nitrogen atom to which they are attached, they form a 4-7 membered heterocyclic ring which is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano or hydroxyl; or

Q is optionally substituted by a group of the formula:



wherein:

$\text{L}_1$  is absent or (1-3C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;

$\text{L}_{\text{Q1}}$  is absent or selected from O, S, SO,  $\text{SO}_2$ ,  $\text{N(R}_f\text{)}$ ,  $\text{C(O)}$ ,  $\text{C(O)O}$ ,  $\text{OC(O)}$ ,  $\text{C(O)N(R}_f\text{)}$ ,  $\text{N(R}_f\text{)C(O)}$ ,  $\text{N(R}_g\text{)C(O)N(R}_f\text{)}$ ,  $\text{N(R}_f\text{)C(O)O}$ ,  $\text{OC(O)N(R}_f\text{)}$ ,  $\text{S(O)}_2\text{N(R}_f\text{)}$ , or  $\text{N(R}_f\text{)SO}_2$ , wherein  $\text{R}_f$  and  $\text{R}_g$  are each independently selected from hydrogen or (1-2C)alkyl; and

$\text{Z}_1$  is hydrogen, (1-6C)alkyl, aryl, aryl(1-2C)alkyl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein  $\text{Z}_1$  is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl, mercapto, ureido, aryl, heteroaryl, heterocycyl, (3-6C)cycloalkyl,  $\text{NR}_h\text{R}_i$ ,  $\text{OR}_h$ ,  $\text{C(O)R}_h$ ,  $\text{C(O)OR}_h$ ,  $\text{OC(O)R}_h$ ,  $\text{C(O)N(R}_i\text{)R}_h$ ,  $\text{N(R}_i\text{)C(O)R}_h$ ,  $\text{S(O)}_{y^a}\text{R}_h$  (where  $y^a$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_i\text{)R}_h$ ,  $\text{N(R}_i\text{)SO}_2\text{R}_h$  or  $(\text{CH}_2)_{z^a}\text{NR}_i\text{R}_h$  (where  $z^a$  is 1, 2 or 3); wherein  $\text{R}_h$  and  $\text{R}_i$  are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl;

$\text{R}_{1a}$  and  $\text{R}_{1b}$  are each selected from hydrogen, (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl or mercapto;

W is selected from O, S or  $\text{NR}_j$ , wherein  $\text{R}_j$  is selected from H or (1-2C)alkyl;

$\text{X}_1$  and  $\text{X}_2$  are each independently selected from N or  $\text{CR}_k$ ;

wherein

$\text{R}_k$  is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, amino, (1-4C)alkylamino, (1-4C)dialkylamino, cyano, (2C)alkynyl,  $\text{C(O)R}_{k1}$ ,  $\text{C(O)OR}_{k1}$ ,  $\text{OC(O)R}_{k1}$ ,  $\text{C(O)N(R}_{k2}\text{)R}_{k1}$ ,  $\text{N(R}_{k2}\text{)C(O)R}_{k1}$ ,  $\text{S(O)}_{y^b}\text{R}_{k1}$  (where  $y^b$  is 0, 1 or 2),  $\text{SO}_2\text{N(R}_{k2}\text{)R}_{k1}$ ,  $\text{N(R}_{k2}\text{)SO}_2\text{R}_{k1}$  or  $(\text{CH}_2)_{z^b}\text{NR}_{k1}\text{R}_{k2}$  (where  $z^b$  is 1, 2 or 3); wherein said (1-4C)alkyl is optionally substituted by one or more substituents selected from amino, hydroxy, (1-2C)alkoxy or halo; and

$\text{R}_{k1}$  and  $\text{R}_{k2}$  are each independently selected from hydrogen or (1-4C)alkyl;

$\text{X}_3$  is selected from N or  $\text{CR}_m$ ;

wherein


$R_m$  is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, amino, (1-4C)alkylamino, (1-4C)dialkylamino, cyano, (2C)alkynyl,  $C(O)R_{m1}$ ,  $C(O)OR_{m1}$ ,  $OC(O)R_{m1}$ ,  $C(O)N(R_{m2})R_{m1}$ ,  $N(R_{m2})C(O)R_{m1}$ ,  $S(O)_{y^c}R_{m1}$  (where  $y^c$  is 0, 1 or 2),  $SO_2N(R_{m2})R_{m1}$ ,  $N(R_{m2})SO_2R_{m1}$  or  $(CH_2)_{z^c}NR_{m1}R_{m2}$  (where  $z^c$  is 1, 2 or 3); wherein said (1-4C)alkyl is optionally substituted by one or more substituents selected from amino, hydroxy, (1-2C)alkoxy or halo; and

$R_{m1}$  and  $R_{m2}$  are each independently selected from hydrogen or (1-4C)alkyl;

$R_o$  is selected from halo, (1-4C)alkyl, (1-4C)alkoxy, amino, (1-4C)alkylamino, (1-4C)dialkylamino, cyano, (2C)alkynyl,  $C(O)R_{o1}$ ,  $C(O)OR_{o1}$ ,  $OC(O)R_{o1}$ ,  $C(O)N(R_{o2})R_{o1}$ ,  $N(R_{o2})C(O)R_{o1}$ ,  $S(O)_{y^d}R_{o1}$  (where  $y^d$  is 0, 1 or 2),  $SO_2N(R_{o2})R_{o1}$ ,  $N(R_{o2})SO_2R_{o1}$  or  $(CH_2)_{z^d}NR_{o1}R_{o2}$  (where  $z^d$  is 1, 2 or 3); wherein said (1-4C)alkyl is optionally substituted by one or more substituents selected from amino, hydroxy, (1-2C)alkoxy or halo; and

$R_{o1}$  and  $R_{o2}$  are each independently selected from hydrogen or (1-4C)alkyl;

$R_2$  is selected from hydrogen, (1-4C)alkyl or a group of the formula:



wherein:

$L_2$  is absent or (1-3C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;

$Y_2$  is absent or  $C(O)$ ,  $C(O)O$ ,  $C(O)N(R_p)$ , wherein  $R_p$  is selected from hydrogen or (1-4C)alkyl; and

$Q_2$  is hydrogen, (1-6C)alkyl, aryl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein  $Q_2$  is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl,  $NR_qR_r$ ,  $OR_q$ , wherein  $R_q$  and  $R_r$  are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl;

$R_3$  is selected from a group of the formula:




wherein:

$Y_3$  is  $C(O)$ ,  $C(O)N(R_y)$ ,  $C(O)N(R_y)O$ ,  $N(R_y)(O)C$ ,  $C(O)O$ ,  $OC(O)$ ,  $N(R_y)C(O)N(R_{y1})$ ,  $SO_2N(R_y)$ ,  $N(R_y)SO_2$ , oxazolyl, triazolyl, oxadiazolyl, thiazolyl, imidazolyl, thiadiazolyl, pyridinyl, pyrazolyl, pyrrolol or tetrazolyl, wherein  $R_y$  and  $R_{y1}$  are independently selected from hydrogen or (1-2C)alkyl; and

$Q_3$  is hydrogen, (1-6C)alkyl, aryl, aryl(1-2C)alkyl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein  $Q_3$  is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl,  $NR_zR_{aa}$ ,  $OR_z$ , wherein  $R_z$

and  $R_{aa}$  are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl; or  $Q_3$  is optionally substituted by a group of the formula:



wherein:

$L_4$  is absent or (1-3C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;


$L_{Q4}$  is absent or selected from O, S, SO, SO<sub>2</sub>, N( $R_{ab}$ ), C(O), C(O)O, OC(O), C(O)N( $R_{ab}$ ), N( $R_{ab}$ )C(O), N( $R_{ac}$ )C(O)N( $R_{ab}$ ), N( $R_{ab}$ )C(O)O, OC(O)N( $R_{ab}$ ), S(O)<sub>2</sub>N( $R_{ab}$ ), or N( $R_{ab}$ )SO<sub>2</sub>, wherein  $R_{ab}$  and  $R_{ac}$  are each independently selected from hydrogen or (1-2C)alkyl; and

$Z_4$  is hydrogen, (1-6C)alkyl, aryl, aryl(1-2C)alkyl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein  $Z_4$  is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl, mercapto, ureido, aryl, heteroaryl, heterocyclyl, (3-6C)cycloalkyl, NR<sub>ad</sub>R<sub>ae</sub>, OR<sub>ad</sub>, C(O)R<sub>ad</sub>, C(O)OR<sub>ad</sub>, OC(O)R<sub>ad</sub>, C(O)N(R<sub>ae</sub>)R<sub>ad</sub>, N(R<sub>ae</sub>)C(O)R<sub>ad</sub>, S(O)<sub>y<sup>e</sup></sub>R<sub>ad</sub> (where  $y^e$  is 0, 1 or 2), SO<sub>2</sub>N(R<sub>ae</sub>)R<sub>ad</sub>, N(R<sub>ae</sub>)SO<sub>2</sub>R<sub>ad</sub> or (CH<sub>2</sub>)<sub>z<sup>e</sup></sub>NR<sub>ad</sub>R<sub>ae</sub> (where  $z^e$  is 1, 2 or 3); wherein R<sub>ad</sub> and R<sub>ae</sub> are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl; or

$Q_3$  and  $R_y$  are linked such that, together with the nitrogen atom to which they are attached, they form a 4-7 membered heterocyclic ring which is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano or hydroxyl;


with the proviso that only one or two of  $X_1$ ,  $X_2$  or  $X_3$  can be N.

**[00421]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO 2017145050, which is hereby incorporated by reference. For example, in some embodiments, the other RET has the Formula (VI) or is a pharmaceutically acceptable salt thereof.




(VI)

**[00422]** Additional examples of other RET kinase inhibitors include those described in International Publication No. WO 2016038552 is hereby incorporated by reference. For example, in some embodiments, the other RET has the Formula (VII), or the Formula (VIII), or is a pharmaceutically acceptable salt thereof.



(VII)




(VIII)

**[00423]** Yet other therapeutic agents include RET inhibitors such as those described, for example, in U.S. Patent Nos. 9,738,660; 9,801,880; 9,682,083; 9,789,100; 9,550,772; 9,493,455; 9,758,508; 9,604,980; 9,321,772; 9,522,910; 9,669,028; 9,186,318; 8,933,230; 9,505,784; 8,754,209; 8,895,744; 8,629,135; 8,815,906; 8,354,526; 8,741,849; 8,461,161; 8,524,709; 8,129,374; 8,686,005; 9,006,256; 8,399,442; 7,795,273; 7,863,288; 7,465,726; 8,552,002; 8,067,434; 8,198,298; 8,106,069; 6,861,509; 8,299,057; 9,150,517; 9,149,464; 8,299,057; and 7,863,288; U.S. Publication Nos. 2018/0009818; 2018/0009817; 2017/0283404; 2017/0267661; 2017/0298074; 2017/0114032; 2016/0009709; 2015/0272958; 2015/0238477; 2015/0099721; 2014/0371219; 2014/0137274; 2013/0079343; 2012/0283261; 2012/0225057; 2012/0065233; 2013/0053370; 2012/0302567; 2011/0189167; 2016/0046636; 2013/0012703; 2011/0281841; 2011/0269739; 2012/0271048; 2012/0277424; 2011/0053934; 2011/0046370; 2010/0280012; 2012/0070410; 2010/0081675; 2010/0075916; 2011/0212053; 2009/0227556; 2009/0209496; 2009/0099167; 2010/0209488; 2009/0012045; 2013/0303518; 2008/0234267; 2008/0199426; 2010/0069395; 2009/0312321; 2010/0173954; 2011/0195072; 2010/0004239; 2007/0149523; 2017/0281632; 2017/0226100; 2017/0121312; 2017/0096425; 2017/0044106; 2015/0065468; 2009/0069360; 2008/0275054; 2007/0117800; 2008/0234284; 2008/0234276; 2009/0048249; 2010/0048540; 2008/0319005; 2009/0215761; 2008/0287427; 2006/0183900; 2005/0222171; 2005/0209195; 2008/0262021; 2008/0312192; 2009/0143399; 2009/0130229; 2007/0265274; 2004/0185547; and 2016/0176865; and International Publication Nos. WO 2017/079140; WO 2017/145050; WO 2017/097697; WO 2017/049462; WO 2017/043550; WO 2017/027883; WO 2017/013160; WO 2017/009644; WO 2016/168992; WO 2016/137060; WO 2016/127074; WO 2016/075224; WO 2016/038552; WO 2015/079251; WO 2014/086284; WO 2013/042137; WO 2013/036232; WO 2013/016720; WO 2012/053606; WO 2012/047017; WO 2007/109045; WO 2009/042646; WO 2009/023978; WO 2009/017838; WO 2017/178845; WO 2017/178844; WO 2017/146116; WO 2017/026718; WO 2016/096709; WO 2007/057397; WO 2007/057399; WO 2007/054357; WO 2006/130613; WO 2006/089298; WO 2005/070431; WO 2003/020698; WO 2001/062273; WO 2001/016169; WO 1997/044356; WO

2007/087245; WO 2005/044835; WO 2014/075035; and WO 2016/038519; and *J. Med.Chem.* 2012, 55 (10), 4872-4876, all of which are hereby incorporated by reference in their entireties.

**[00424]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is a compound of the Formula II:



or a pharmaceutically acceptable salt or solvate thereof, wherein:

X<sup>1</sup> is CH, CCH<sub>3</sub>, CF, CCl or N;

X<sup>2</sup> is CH, CF or N;

X<sup>3</sup> is CH, CF or N;

X<sup>4</sup> is CH, CF or N;

wherein zero, one or two of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> is N;

A is H, Cl, CN, Br, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub> or cyclopropyl;

B is hetAr<sup>1</sup>;

hetAr<sup>1</sup> is a 5-membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, S and O, wherein said heteroaryl ring is optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C6 alkyl, hydroxyC1-C6 alkyl, fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, cyanoC1-C6 alkyl, (C1-C6 alkoxy)C1-C6 alkyl, (C1-C4 alkoxy)CH<sub>2</sub>C(=O)-, (C1-C4 alkoxy)C(=O)C1-C3 alkyl, C3-C6 cycloalkyl, (R<sup>a</sup>R<sup>b</sup>N)C1-C6 alkyl, (R<sup>a</sup>R<sup>b</sup>N)C(=O)C1-C6 alkyl, (C1-C6 alkylSO<sub>2</sub>)C1-C6 alkyl, hetCyc<sup>a</sup>, and 4-methoxybenzyl;

R<sup>a</sup> and R<sup>b</sup> are independently H or C1-C6 alkyl;

hetCyc<sup>a</sup> is a 4-6 membered heterocyclic ring having a ring heteroatom selected from N and O, wherein said heterocyclic ring is optionally substituted with halogen, C1-C6 alkyl, fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, (C1-C6 alkoxy)C1-C6 alkyl, di(C1-C3 alkyl)NCH<sub>2</sub>C(=O), (C1-C6 alkoxy)C(=O) or (C1-C6 alkoxy)CH<sub>2</sub>C(=O);

D is hetCyc<sup>1</sup>, hetCyc<sup>2</sup>, hetCyc<sup>3</sup> or hetCyc<sup>9</sup>;

hetCyc<sup>1</sup> is a 4-6 membered heterocyclic ring having 1-2 ring atoms selected from N and O, wherein said heterocyclic ring is optionally substituted with one or more substituents independently selected

from the group consisting of C1-C3 alkyl, fluoroC1-C3 alkyl, difluoroC1-C3 alkyl, trifluoroC1-C3 alkyl and OH, or said heterocyclic ring is substituted with a C3-C6 cycloalkylidene ring, or said heterocyclic ring is substituted with an oxo group;

hetCyc<sup>2</sup> is a 7-8 membered bridged heterocyclic ring having 1-3 ring heteroatoms independently selected from N and O, wherein said heterocyclic ring is optionally substituted with C1-C3 alkyl;

hetCyc<sup>3</sup> is a 7-11 membered heterospirocyclic ring having 1-2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C3 alkyl;

hetCyc<sup>9</sup> is a fused 9-10 membered heterocyclic ring having 1-3 ring nitrogen atoms and optionally substituted with oxo;

E is

- (a) hydrogen,
- (b) OH,
- (c) R<sup>a</sup>R<sup>b</sup>N-, wherein R<sup>a</sup> is H or C1-C6 alkyl and R<sup>b</sup> is H, C1-C6 alkyl or phenyl;
- (d) C1-C6 alkyl optionally substituted with one to three fluoros,
- (e) hydroxyC1-C6 alkyl- optionally substituted with one to three fluoros,
- (f) C1-C6 alkoxy optionally substituted with one to three fluoros,
- (g) hydroxy(C1-C6 alkoxy) optionally substituted with one to three fluoros,
- (h) (C1-C6 alkoxy)hydroxy C1-C6 alkyl- optionally substituted with one to three fluoros,
- (i) (C1-C6 alkyl)C(=O)- optionally substituted with one to three fluoros,
- (j) (hydroxy C1-C6 alkyl)C(=O)- optionally substituted with one to three fluoros,
- (k) (C1-C6 alkoxy)C(=O)-,
- (l) (C1-C6 alkoxy)(C1-C6 alkyl)C(=O)-,
- (m) HC(=O)-,
- (n) Cyc<sup>1</sup>,
- (o) Cyc<sup>1</sup>C(=O)-,
- (p) Cyc<sup>1</sup>(C1-C6 alkyl)C(=O)- wherein said alkyl portion is optionally substituted with one or more groups independently selected from the group consisting of OH, fluoro, C1-C3 alkoxy and R<sup>c</sup>R<sup>d</sup>N-, where R<sup>c</sup> and R<sup>d</sup> are independently H or C1-C6 alkyl,
- (q) hetCyc<sup>4</sup>,
- (r) hetCyc<sup>4</sup>C(=O)-,
- (s) hetCyc<sup>4</sup>(C1-C3 alkyl)C(=O)-,

- (t)  $(\text{hetCyc}^4)\text{C}(=\text{O})\text{C1-C2 alkyl-}$ ,
- (u)  $\text{hetCyc}^4\text{C}(=\text{O})\text{NH-}$ ,
- (v)  $\text{Ar}^2$ ,
- (w)  $\text{Ar}^2\text{C}(=\text{O})-$ ,
- (x)  $\text{Ar}^2\text{C1-C6 alkyl-}$ ,
- (y)  $(\text{Ar}^2)\text{hydroxy C2-C6 alkyl-}$ ,
- (z)  $\text{Ar}^2(\text{C1-C3 alkyl})\text{C}(=\text{O})-$  wherein said alkyl portion is optionally substituted with one or two groups independently selected from the group consisting of OH, C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl, C1-C6 alkoxy and  $\text{R}^e\text{R}^f\text{N-}$ , where  $\text{R}^e$  and  $\text{R}^f$  are independently H or C1-C6 alkyl, or  $\text{R}^e$  and  $\text{R}^f$  together with the nitrogen to which they are attached form a 5-6 membered azacyclic ring optionally having an additional ring heteroatom selected from N and O,
- (aa)  $\text{hetAr}^2\text{C}(=\text{O})-$ ,
- (bb)  $(\text{hetAr}^2)\text{hydroxyC2-C6 alkyl-}$ ,
- (cc)  $\text{hetAr}^2(\text{C1-C3 alkyl})\text{C}(=\text{O})-$ , wherein said alkyl portion is optionally substituted with one or two groups independently selected from the group consisting of OH, C1-C6 alkyl, hydroxyC1-C6 alkyl, C1-C6 alkoxy and  $\text{R}^e\text{R}^f\text{N-}$ , wherein  $\text{R}^e$  and  $\text{R}^f$  are independently H or C1-C6 alkyl or  $\text{R}^e$  and  $\text{R}^f$  together with the nitrogen to which they are attached form a 5-6 membered azacyclic ring optionally having an additional ring heteroatom selected from N and O,
- (dd)  $\text{R}^1\text{R}^2\text{NC}(=\text{O})-$ ,
- (ee)  $\text{R}^1\text{R}^2\text{N}(\text{C1-C3 alkyl})\text{C}(=\text{O})-$ , wherein said alkyl portion is optionally substituted with phenyl,
- (ff)  $\text{R}^1\text{R}^2\text{NC}(=\text{O})\text{C1-C2 alkyl-}$ ,
- (gg)  $\text{R}^1\text{R}^2\text{NC}(=\text{O})\text{NH-}$ ,
- (hh)  $\text{CH}_3\text{SO}_2(\text{C1-C6 alkyl})\text{C}(=\text{O})-$ ,
- (ii)  $(\text{C1-C6 alkyl})\text{SO}_2-$ ,
- (jj)  $(\text{C3-C6 cycloalkyl})\text{CH}_2\text{SO}_2-$ ,
- (kk)  $\text{hetCyc}^5\text{-SO}_2-$ ,
- (ll)  $\text{R}^4\text{R}^5\text{NSO}_2-$ ,
- (mm)  $\text{R}^6\text{C}(=\text{O})\text{NH-}$ ,
- (nn)  $\text{hetCyc}^6$ ,
- (oo)  $\text{hetAr}^2\text{C1-C6 alkyl-}$ ,



- (pp) (hetCyc<sup>4</sup>)C1-C6 alkyl-,
- (qq) (C1-C6 alkoxy)C1-C6 alkyl- optionally substituted with 1-3 fluoros,
- (rr) (C3-C6 cycloalkoxy)C1-C6 alkyl-,
- (ss) (C3-C6 cycloalkyl)C1-C6 alkyl-, wherein said cycloalkyl is optionally substituted with 1-2 fluoros,
- (tt) (R<sup>g</sup>R<sup>h</sup>N)C1-C6 alkyl-, wherein R<sup>g</sup> and R<sup>h</sup> are independently H or C1-C6 alkyl,
- (uu) Ar<sup>2</sup>-O-,
- (vv) (C1-C6 alkylSO<sub>2</sub>)C1-C6 alkyl-,
- (ww) (C1-C6 alkoxy)C(=O)NHC1-C6 alkyl-,
- (xx) (C3-C6 cycloalkoxy)C(=O)-,
- (yy) (C3-C6 cycloalkyl)SO<sub>2</sub>-, wherein said cycloalkyl is optionally substituted with C1-C6 alkyl,
- (zz) Ar<sup>4</sup>CH<sub>2</sub>OC(=O)-,
- (aaa) (N-(C1-C3 alkyl)pyridinonyl)C1-C3 alkyl-, and
- (bbb) (Ar<sup>4</sup>SO<sub>2</sub>)C1-C6 alkyl-;

Cyc<sup>1</sup> is a C3-C6 cycloalkyl, wherein (a) said cycloalkyl is optionally substituted with one or more substituents independently selected from the group consisting of OH, halogen, C1-C6 alkoxy, CN, hydroxyC1-C6 alkyl, (C1-C6 alkoxy)C1-C6 alkyl, and C1-C6 alkyl optionally substituted with 1-3 fluoros, or (b) said cycloalkyl is substituted with phenyl, wherein said phenyl is optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C3 alkyl, C1-C3 alkoxy and CF, or (c) said cycloalkyl is substituted with a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N and O, wherein said heteroaryl ring is optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C3 alkyl, C1-C3 alkoxy and CF<sub>3</sub>;

Ar<sup>2</sup> is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C6 alkyl, C1-C6 alkoxy (optionally substituted with 1-3 fluoros), fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, CN, a 5-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O, and R<sup>i</sup>R<sup>j</sup>N- wherein R<sup>i</sup> and R<sup>j</sup> are independently H or C1-C6 alkyl;

hetAr<sup>2</sup> is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S and optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C6 alkyl, C1-C6 alkoxy (optionally substituted with 1-3 fluoros), fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, hydroxyC1-C6 alkyl, (C3-C6)cycloalkyl, (C1-C6 alkoxy)C1-C6 alkyl, CN, OH, and R'R''N-, wherein R' and R'' are independently H or C1-C3 alkyl;

hetCyc<sup>4</sup> is (a) a 4-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N, O and S wherein said S is optionally oxidized to SO<sub>2</sub>, (b) a 7-8 membered bridged heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O, (c) a 6-12 membered fused bicyclic heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O and optionally independently substituted with 1-2 C1-C6 alkyl substituents, or (d) a 7-10 membered spirocyclic heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O, wherein each of said heterocyclic rings is optionally substituted with one or more substituents independently selected from the group consisting of halogen, OH, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy, (C1-C6 alkoxy)C1-C6 alkyl, (C3-C6)cycloalkyl, (C1-C6 alkyl)C(=O)-, a 5-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O, and phenyl wherein said phenyl is optionally substituted with one or more substituents selected from halogen, C1-C6 alkyl and C1-C6 alkoxy;

hetCyc<sup>5</sup> is a 5-6 membered heterocyclic ring having a ring heteroatom selected from O and N;

hetCyc<sup>6</sup> is a 5 membered heterocyclic ring having one or two ring heteroatoms independently selected from N and O, wherein said ring is substituted with oxo and wherein said ring is further optionally substituted with one or more substituents independently selected from the group consisting of OH and C1-C6 alkyl;

R<sup>1</sup> is H, C1-C6 alkyl or (C1-C6 alkoxy)C1-C6 alkyl;

R<sup>2</sup> is H, C1-C6 alkyl (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl (optionally substituted with 1-3 fluoros), Cyc<sup>3</sup>, hydroxyC1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C(=O), hetCyc<sup>7</sup>, Ar<sup>3</sup>, Ar<sup>3</sup>C1-C3 alkyl-, hydroxyC1-C6 alkoxy or (3-6C cycloalkyl)CH<sub>2</sub>O-;

Cyc<sup>3</sup> is a 3-6 membered carbocyclic ring optionally substituted with 1-2 groups independently selected from the group consisting of C1-C6 alkoxy, OH and halogen;

hetCyc<sup>7</sup> is a 5-6 membered heterocyclic ring having a ring heteroatom selected from O and N wherein said ring is optionally substituted with C1-C6 alkyl;

Ar<sup>3</sup> is phenyl optionally substituted with one or more substituents independently selected from halogen, C1-C3 alkyl, C1-C3 alkoxy, fluoroC1-C3 alkyl, difluoroC1-C3 alkyl and trifluoroC1-C3 alkyl;


R<sup>4</sup> and R<sup>5</sup> are independently H or C1-C6 alkyl;

R<sup>6</sup> is C1-C6 alkyl, hydroxyC1-C6 alkyl, C1-C6 alkoxy, (C1-C6 alkoxy)C1-C6 alkyl, phenyl or hetCyc<sup>8</sup>;

hetCyc<sup>8</sup> is a 5-6 membered heterocyclic ring having a ring heteroatom selected from O and N, wherein said heterocyclic ring is optionally substituted with C1-C6 alkyl; and

Ar<sup>4</sup> is phenyl optionally substituted with one or more halogens.

**[00425]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is a compound of the Formula III:



III

or a pharmaceutically acceptable salt or solvate thereof, wherein:

$X^1$  is CH or N;

$X^2$  is CH or N;

$X^3$  is CH or N;

$X^4$  is CH or N;

wherein one or two of  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  is N;

A is CN;

B is  $\text{hetAr}^1$ ;

$\text{hetAr}^1$  is a 5-membered heteroaryl ring having 1-3 ring nitrogen atoms, wherein said heteroaryl ring is optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C6 alkyl, hydroxyC1-C6 alkyl, fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, cyanoC1-C6 alkyl, (C1-C6 alkoxy)C1-C6 alkyl, (C1-C4 alkoxy)CH<sub>2</sub>C(=O)-, (C1-C4 alkoxy)C(=O)C1-C3 alkyl, C3-C6 cycloalkyl, ( $R^a R^b$ N)C1-C6 alkyl, ( $R^a R^b$ N)C(=O)C1-C6 alkyl, (C1-C6 alkylSO<sub>2</sub>)C1-C6 alkyl, and 4-methoxybenzyl;

$R^a$  and  $R^b$  are independently H or C1-C6 alkyl;

D is  $\text{hetCyc}^1$ ;

$\text{hetCyc}^1$  is a 4-6 membered heterocyclic ring having 1-2 ring nitrogen atoms, wherein said heterocyclic ring is optionally substituted with one or more substituents independently selected from the group consisting of C1-C3 alkyl, fluoroC1-C3 alkyl, difluoroC1-C3 alkyl, trifluoroC1-C3 alkyl and OH, or said heterocyclic ring is substituted with a C3-C6 cycloalkylidene ring, or said heterocyclic ring is substituted with an oxo group;

E is

(w)  $\text{Ar}^2\text{C}(=\text{O})-$ ,

(x)  $\text{Ar}^2\text{C1-C6 alkyl-}$ ,

(z)  $\text{Ar}^2(\text{C1-C3 alkyl})\text{C}(=\text{O})-$  wherein said alkyl portion is optionally substituted with one or two groups independently selected from the group consisting of OH, C1-C6 alkyl (optionally

substituted with 1-3 fluoros), hydroxyC1-C6 alkyl, C1-C6 alkoxy and  $R^eR^fN-$ , where  $R^e$  and  $R^f$  are independently H or C1-C6 alkyl, or  $R^e$  and  $R^f$  together with the nitrogen to which they are attached form a 5-6 membered azacyclic ring optionally having an additional ring heteroatom selected from N and O,

(cc)  $hetAr^2(C1-C3 \text{ alkyl})C(=O)-$ , wherein said alkyl portion is optionally substituted with one or two groups independently selected from the group consisting of OH, C1-C6 alkyl, hydroxyC1-C6 alkyl, C1-C6 alkoxy and  $R^eR^fN-$ , wherein  $R^e$  and  $R^f$  are independently H or C1-C6 alkyl or  $R^e$  and  $R^f$  together with the nitrogen to which they are attached form a 5-6 membered azacyclic ring optionally having an additional ring heteroatom selected from N and O,

(dd)  $R^1R^2NC(=O)-$ ,

(oo)  $hetAr^2C1-C6 \text{ alkyl}-$ ,

$Ar^2$  is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C6 alkyl, C1-C6 alkoxy (optionally substituted with 1-3 fluoros), fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, CN, a 5-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O, and  $R^iR^jN-$  wherein  $R^i$  and  $R^j$  are independently H or C1-C6 alkyl;

$hetAr^2$  is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S and optionally substituted with one or more substituents independently selected from the group consisting of halogen, C1-C6 alkyl, C1-C6 alkoxy (optionally substituted with 1-3 fluoros), fluoroC1-C6 alkyl, difluoroC1-C6 alkyl, trifluoroC1-C6 alkyl, hydroxyC1-C6 alkyl, (C3-C6)cycloalkyl, (C1-C6 alkoxy)C1-C6 alkyl, CN, OH, and  $R'R''N-$ , wherein  $R'$  and  $R''$  are independently H or C1-C3 alkyl;


$R^1$  is H, C1-C6 alkyl or (C1-C6 alkoxy)C1-C6 alkyl; and

$R^2$  is H, C1-C6 alkyl (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C(=O), hydroxyC1-C6 alkoxy or (3-6C cycloalkyl)CH<sub>2</sub>O.

**[00426]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-

yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof.

**[00427]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is a compound of the Formula **IV**:



**IV**

or a pharmaceutically acceptable salt or solvate thereof, wherein:

$X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  are independently CH, CF, CCH<sub>3</sub> or N, wherein zero, one or two of  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  is N;

A is H, CN, Cl, CH<sub>3</sub>-, CH<sub>3</sub>CH<sub>2</sub>-, cyclopropyl, -CH<sub>2</sub>CN or -CH(CN)CH<sub>3</sub>;

B is

- (a) hydrogen,
- (b) C1-C6 alkyl optionally substituted with 1-3 fluoros,
- (c) hydroxyC2-C6 alkyl-, wherein the alkyl portion is optionally substituted with 1-3 fluoros or a C3-C6 cycloalkylidene ring,
- (d) dihydroxyC3-C6 alkyl-, wherein the alkyl portion is optionally substituted with a C3-C6 cycloalkylidene ring,
- (e) (C1-C6 alkoxy)C1-C6 alkyl- optionally substituted with 1-3 fluoros,
- (f) (R<sup>1</sup>R<sup>2</sup>N)C1-C6 alkyl- wherein said alkyl portion is optionally substituted with OH and wherein R<sup>1</sup> and R<sup>2</sup> are independently H or C1-C6 alkyl (optionally substituted with 1-3 fluoros);
- (g) hetAr<sup>1</sup>C1-C3 alkyl-, wherein hetAr<sup>1</sup> is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S and is optionally substituted with one or more independently selected C1-C6 alkyl substituents;
- (h) (C3-C6 cycloalkyl)C1-C3 alkyl-, wherein said cycloalkyl is optionally substituted with OH,
- (i) (hetCyc<sup>a</sup>)C1-C3 alkyl-,

- (j) hetCyc<sup>a</sup>-,
- (k) C3-C6 cycloalkyl-, wherein said cycloalkyl is optionally substituted with OH,
- (l) (C1-C4 alkyl)C(=O)O-C1-C6 alkyl-, wherein each of the C1-C4 alkyl and C1-C6 alkyl portions is optionally and independently substituted with 1-3 fluoros, or
- (m) (R<sup>1</sup>R<sup>2</sup>N)C(=O)C1-C6 alkyl-, wherein R<sup>1</sup> and R<sup>2</sup> are independently H or C1-C6 alkyl (optionally substituted with 1-3 fluoros);

hetCyc<sup>a</sup>- is a 4-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O and optionally substituted with one or more substituents independently selected from OH, C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl-, C1-C6 alkoxy, (C1-C6 alkyl)C(=O)-, (C1-C6 alkoxy)C1-C6 alkyl-, and fluoro, or wherein hetCyc<sup>a</sup> is substituted with oxo;

Ring D is (i) a saturated 4-7 membered heterocyclic ring having two ring nitrogen atoms, (ii) a saturated 7-8 membered bridged heterocyclic ring having two ring nitrogen atoms and optionally having a third ring heteroatom which is oxygen, (iii) a saturated 7-11 membered heterospirocyclic ring having two ring nitrogen atoms, or (iv) a saturated 9-10 membered bicyclic fused heterocyclic ring having two ring nitrogen atoms, wherein each of said rings is optionally substituted with (a) one to four groups independently selected from halogen, OH, C1-C3 alkyl which is optionally substituted with 1-3 fluoros, or C1-C3 alkoxy which is optionally substituted with 1-3 fluoros, (b) a C3-C6 cycloalkylidene ring, or (c) an oxo group;

E is

- (a) hydrogen,
- (b) C1-C6 alkyl optionally substituted with 1-3 fluoros,
- (c) (C1-C6 alkoxy)C1-C6 alkyl- optionally substituted with 1-3 fluoros,
- (d) (C1-C6 alkyl)C(=O)-, wherein said alkyl portion is optionally substituted with 1-3 fluoros or with a R<sup>g</sup>R<sup>h</sup>N- substituent wherein R<sup>g</sup> and R<sup>h</sup> are independently H or C1-C6 alkyl,
- (e) (hydroxyC2-C6 alkyl)C(=O)- optionally substituted with 1-3 fluoros,
- (f) (C1-C6 alkoxy)C(=O)-,
- (g) (C3-C6 cycloalkyl)C(=O)-, wherein said cycloalkyl is optionally substituted with one or more substituents independently selected from C1-C6 alkyl, C1-C6 alkoxy, OH, and (C1-C6 alkoxy)C1-C6 alkyl-, or said cycloalkyl is substituted with a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N and O,
- (h) Ar<sup>1</sup>C1-C6 alkyl-,
- (i) Ar<sup>1</sup>(C1-C6 alkyl)C(=O)-, wherein said alkyl portion is optionally substituted with OH, hydroxyC1-C6 alkyl-, C1-C6 alkoxy, R<sup>m</sup>R<sup>n</sup>N- or R<sup>m</sup>R<sup>n</sup>N-CH<sub>2</sub>-, wherein each R<sup>m</sup> and R<sup>n</sup> is independently H or C1-C6 alkyl,

- (j) hetAr<sup>2</sup>C1-C6 alkyl-, wherein said alkyl portion is optionally substituted with 1-3 fluoros,
- (k) hetAr<sup>2</sup>(C1-C6 alkyl)C(=O)- wherein said alkyl portion is optionally substituted with OH, hydroxyC1-C6 alkyl- or C1-C6 alkoxy,
- (l) hetAr<sup>2</sup>C(=O)-,
- (m) hetCyc<sup>1</sup>C(=O)-,
- (n) hetCyc<sup>1</sup>C1-C6 alkyl-,
- (o) R<sup>3</sup>R<sup>4</sup>NC(=O)-,
- (p) Ar<sup>1</sup>N(R<sup>3</sup>)C(=O)-,
- (q) hetAr<sup>2</sup>N(R<sup>3</sup>)C(=O)-,
- (r) (C1-C6 alkyl)SO<sub>2</sub>-, wherein the alkyl portion is optionally substituted with 1-3 fluoros,
- (s) Ar<sup>1</sup>SO<sub>2</sub>-,
- (t) hetAr<sup>2</sup>SO<sub>2</sub>-,
- (u) N-(C1-C6 alkyl)pyridinonyl,
- (v) Ar<sup>1</sup>C(=O)-;
- (w) Ar<sup>1</sup>O-C(=O)-,
- (x) (C3-C6 cycloalkyl)(C1-C6 alkyl)C(=O)-,
- (y) (C3-C6 cycloalkyl)(C1-C6 alkyl)SO<sub>2</sub>-, wherein the alkyl portion is optionally substituted with 1-3 fluoros,
- (z) Ar<sup>1</sup>(C1-C6 alkyl)SO<sub>2</sub>-,
- (aa) hetCyc<sup>1</sup>-O-C(=O)-,
- (bb) hetCyc<sup>1</sup>CH<sub>2</sub>C(=O)-,
- (cc) hetAr<sup>2</sup>, or
- (dd) C3-C6 cycloalkyl;

Ar<sup>1</sup> is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), R<sup>e</sup>R<sup>f</sup>N- wherein R<sup>e</sup> and R<sup>f</sup> are independently H, C1-C6 alkyl, (R<sup>p</sup>R<sup>q</sup>N)C1-C6 alkoxy- wherein R<sup>p</sup> and R<sup>q</sup> are independently H or C1-C6 alkyl, and (hetAr<sup>a</sup>)C1-C6 alkyl- wherein hetAr<sup>a</sup> is a 5-6 membered heteroaryl ring having 1-2 ring nitrogen atoms, or Ar<sup>1</sup> is a phenyl ring fused to a 5-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O;


hetAr<sup>2</sup> is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S or a 9-10 membered bicyclic heteroaryl ring having 1-3 ring nitrogen atoms, wherein hetAr<sup>2</sup> is optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros), R<sup>e</sup>R<sup>f</sup>N- wherein R<sup>e</sup> and R<sup>f</sup> are independently H or C1-C6 alkyl, OH, (C1-C6 alkoxy)C1-C6 alkoxy- and C3-C6 cycloalkyl;

hetCyc<sup>1</sup> is a 4-6 membered saturated heterocyclic ring having 1-2 ring heteroatoms independently selected from N, O and S wherein said heterocyclic ring is optionally substituted with one or more substituents independently selected from C1-C6 alkoxy and halogen;

R<sup>3</sup> is H or C1-C6 alkyl; and

R<sup>4</sup> is C1-C6 alkyl.

**[00428]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is a compound of the Formula **V**:



**V**

or a pharmaceutically acceptable salt and solvate thereof, wherein:

X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> are independently CH or N, wherein zero, one or two of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> is N;

A is CN;

B is

(b) C1-C6 alkyl optionally substituted with 1-3 fluoros,

(c) hydroxyC2-C6 alkyl-, wherein the alkyl portion is optionally substituted with 1-3 fluoros or a C3-C6 cycloalkylidene ring,

(e) (C1-C6 alkoxy)C1-C6 alkyl- optionally substituted with 1-3 fluoros,

(f) (R<sup>1</sup>R<sup>2</sup>N)C1-C6 alkyl-, wherein said alkyl portion is optionally substituted with OH and wherein R<sup>1</sup> and R<sup>2</sup> are independently H or C1-C6 alkyl (optionally substituted with 1-3 fluoros);



(g)  $\text{hetAr}^1\text{C1-C3 alkyl-}$ , wherein  $\text{hetAr}^1$  is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S and is optionally substituted with one or more independently selected C1-C6 alkyl substituents; or

(i)  $(\text{hetCyc}^a)\text{C1-C3 alkyl-}$ ,

$\text{hetCyc}^a$  is a 4-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O and optionally substituted with one or more substituents independently selected from OH, C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl-, C1-C6 alkoxy, (C1-C6 alkyl)C(=O)-, (C1-C6 alkoxy)C1-C6 alkyl- and fluoro, or wherein  $\text{hetCyc}^a$  is substituted with oxo;

Ring D is (i) a saturated 4-7 membered heterocyclic ring having two ring nitrogen atoms, or (ii) a saturated 7-9 membered bridged heterocyclic ring having two ring nitrogen atoms and optionally having a third ring heteroatom which is oxygen, wherein each of said rings is optionally substituted with (a) one to four groups independently selected from halogen, OH, C1-C3 alkyl which is optionally substituted with 1-3 fluoros, or C1-C3 alkoxy which is optionally substituted with 1-3 fluoros, (b) a C3-C6 cycloalkylidene ring, or (c) an oxo group;

E is

(h)  $\text{Ar}^1\text{C1-C6 alkyl-}$ ,

(j)  $\text{hetAr}^2\text{C1-C6 alkyl-}$ , wherein the alkyl portion is optionally substituted with 1-3 fluoros, or

(l)  $\text{hetAr}^2\text{C(=O)-}$ ,


$\text{Ar}^1$  is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros),  $\text{R}^e\text{R}^f\text{N-}$  wherein  $\text{R}^e$  and  $\text{R}^f$  are independently H or C1-C6 alkyl,  $(\text{R}^p\text{R}^q\text{N})\text{C1-C6 alkoxy-}$  wherein  $\text{R}^p$  and  $\text{R}^q$  are independently H or C1-C6 alkyl, and  $(\text{hetAr}^3)\text{C1-C6 alkyl-}$  wherein  $\text{hetAr}^3$  is a 5-6 membered heteroaryl ring having 1-2 ring nitrogen atoms, or  $\text{Ar}^1$  is a phenyl ring fused to a 5-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O; and

$\text{hetAr}^2$  is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S or a 9-10 membered bicyclic heteroaryl ring having 1-3 ring nitrogen atoms, wherein  $\text{hetAr}^2$  is optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros),  $\text{R}^e\text{R}^f\text{N-}$  wherein  $\text{R}^e$  and  $\text{R}^f$  are independently H or C1-C6 alkyl, OH, (C1-C6 alkoxy)C1-C6 alkoxy- and C3-C6 cycloalkyl.

**[00429]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-

yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof.

**[00430]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is a compound of Formula VI:



VI

or a pharmaceutically acceptable salt or solvate thereof, wherein:

$X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  are independently CH,  $CCH_3$ , CF or N, wherein zero, one or two of  $X^1$ ,  $X^2$ ,  $X^3$  and  $X^4$  is N;

A is H, CN, Cl, methyl, ethyl or cyclopropyl;

B is:

- (a) hydrogen,
- (b) C1-C6 alkyl optionally substituted with 1-3 fluoros,
- (c) hydroxyC2-C6 alkyl- wherein the alkyl portion is optionally substituted with a C3-C6 cycloalkylidene ring,
- (d) dihydroxyC3-C6 alkyl- wherein the alkyl portion is optionally substituted with a C3-C6 cycloalkylidene ring,
- (e) (C1-C6 alkoxy)C1-C6 alkyl- optionally substituted with 1-3 fluoros,
- (f)  $(R^1R^2N)C1-C6$  alkyl- where  $R^1$  and  $R^2$  are independently selected from H, C1-C6 alkyl (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)- and (C1-C6 alkoxy)C(=O)-;

- (g)  $\text{hetAr}^1\text{C1-C3 alkyl-}$ , where  $\text{hetAr}^1$  is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S and is optionally substituted with one or more independently selected C1-C6 alkyl substituents;
- (h)  $(\text{C3-C6 cycloalkyl})\text{C1-C3 alkyl-}$ , wherein said cycloalkyl is optionally substituted with OH,
- (i)  $(\text{hetCyc}^a)\text{C1-C3 alkyl-}$ ,
- (j)  $\text{hetCyc}^a$ ,
- (k)  $(\text{R}^1\text{R}^2\text{N})\text{C(=O)C1-C6 alkyl-}$ , where  $\text{R}^1$  and  $\text{R}^2$  are independently selected from H and C1-C6 alkyl;
- (l)  $(\text{R}^1\text{R}^2\text{N})\text{C(=O)-}$ , where  $\text{R}^1$  and  $\text{R}^2$  are independently selected from H and C1-C6 alkyl, or
- (m)  $\text{hetCyc}^a\text{C(=O)C1-C6 alkyl-}$ ;

$\text{hetCyc}^a$  is a 4-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O and optionally substituted with one or more substituents independently selected from OH, C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl, halogen,  $(\text{C1-C6 alkyl})\text{C(=O)-}$ , C1-C6 alkoxy, oxo and  $(\text{C1-C6 alkoxy})\text{C(=O)-}$ ;

Ring D is (i) a saturated monocyclic 4-7 membered heterocyclic ring having one ring heteroatom which is nitrogen, (ii) a saturated 7-8 membered bridged heterocyclic ring having one ring heteroatom which is nitrogen, or (iii) a saturated 7-11 membered heterospirocyclic ring system having one ring heteroatom which is nitrogen;

each  $\text{R}^a$  is independently C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl or  $(\text{C1-C6 alkoxy})\text{C1-C6 alkyl-}$ ;

$\text{R}^b$  is (a) hydroxy, (b) cyclopropyl, (c)  $\text{hetCyc}^b\text{CH}_2\text{-}$ , (d)  $\text{R}^i\text{R}^j\text{NC(=O)CH}_2\text{OCH}_2\text{-}$  where  $\text{R}^i$  and  $\text{R}^j$  are independently H or C1-C6 alkyl, (e)  $\text{R}^c\text{R}^d\text{N-}$ , (f)  $\text{R}^c\text{R}^d\text{NCH}_2\text{-}$ , (g) C1-C6 alkoxy-, (h)  $(\text{C1-C4 alkyl})\text{-C(=O)NH-}$  wherein said alkyl portion is optionally substituted with  $\text{hetCyc}^b$ ,  $\text{hetAr}^a$ , C1-C6 alkoxy- or  $\text{R}^i\text{R}^j\text{N-}$ , or said alkyl portion is optionally substituted with two substituents independently selected from  $\text{R}^i\text{R}^j\text{N-}$  and OH, where each  $\text{R}^i$  and  $\text{R}^j$  is independently hydrogen or C1-C6 alkyl, (i)  $(\text{R}^i\text{R}^j\text{N})\text{C1-C6 alkoxy}(\text{CH}_2)_n\text{-}$  where n is 0 or 1 and  $\text{R}^i$  and  $\text{R}^j$  are independently hydrogen or C1-C6 alkyl, (j)  $\text{hetCyc}^b(\text{C1-C3 alkyl})\text{OCH}_2\text{-}$ , (k)  $\text{hetCyc}^b\text{C(=O)NH-}$  or (l)  $\text{hetAr}^a\text{C(=O)NH-}$ ;

$\text{hetCyc}^b$  is a 4-6 membered heterocyclic ring, a 7-8 membered bridged heterocyclic ring, or a 7-10 membered heterospirocyclic ring, each ring having 1-2 ring heteroatoms independently selected from N and O, wherein  $\text{hetCyc}^b$  is optionally substituted with one or more substituents independently selected from OH, fluoro, C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl- (optionally substituted with 1-3 fluoros),  $(\text{C1-C6 alkoxy})\text{C1-C6 alkyl-}$ ,  $(\text{C1-C6 alkoxy})\text{C(=O)-}$ , C1-C6 alkoxy, and  $\text{R}^i\text{R}^j\text{N-}$  where  $\text{R}^i$  and  $\text{R}^j$  are independently hydrogen or C1-C6 alkyl;

$\text{hetAr}^a$  is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S wherein  $\text{hetAr}^a$  is optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), and C1-C6 alkoxy (optionally substituted with 1-3 fluoros),

R<sup>c</sup> is hydrogen or C1-C6 alkyl;

R<sup>d</sup> is hydrogen, C1-C6 alkyl (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C(=O)-, hydroxyC1-C6 alkyl (optionally substituted with 1-3 fluoros), (hydroxyC1-C6 alkyl)C(=O)-, (C1-C6 alkyl)C(=O)-, (R<sup>k</sup>R<sup>l</sup>N)C1-C6 alkyl- where R<sup>k</sup> and R<sup>l</sup> are independently H or C1-C6 alkyl, R<sup>m</sup>R<sup>n</sup>NC(=O)C1-C6 alkyl- where R<sup>m</sup> and R<sup>n</sup> are independently H or C1-C6 alkyl, PhCH<sub>2</sub>- wherein the phenyl is optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros), C3-C6 cycloalkyl, hydroxyC1-C6 alkyl, (C1-C6 alkyl)SO<sub>2</sub>-, R<sup>e</sup>R<sup>f</sup>N- and (R<sup>e</sup>R<sup>f</sup>N)C1-C6 alkyl- where each R<sup>e</sup> and R<sup>f</sup> is independently H or C1-C6 alkyl, (C1-C6 alkoxy)C1-C6 alkyl-, or hetCyc<sup>c</sup> where hetCyc<sup>c</sup> is a 4-6 membered heterocyclic ring having a ring heteroatom selected from N and O and optionally substituted with C1-C6 alkyl;




n is 0, 1, 2, 3, 4, 5 or 6;

m is 0 or 1;

E is:

- (a) hydrogen,
- (b) hydroxy,
- (c) C1-C6 alkyl optionally substituted with 1-3 fluoros,
- (d) Ar<sup>1</sup>C1-C6 alkyl- wherein said alkyl portion is optionally substituted with 1-3 fluoros,
- (e) hetAr<sup>2</sup>C1-C6 alkyl-,
- (f) (C1-C6 alkoxy)C1-C6 alkoxy-,
- (g) Ar<sup>1</sup>O-,
- (h) hetAr<sup>2</sup>-O-,
- (i) Ar<sup>1</sup>NR<sup>g</sup>- where R<sup>g</sup> is H or C1-C6 alkyl,
- (j) hetAr<sup>2</sup>NR<sup>g</sup>- where R<sup>g</sup> is H or C1-C6 alkyl,
- (k) R<sup>3</sup>C(=O)NR<sup>g</sup>- where R<sup>g</sup> is H or C1-C6 alkyl;
- (l) Ar<sup>1</sup>C(=O)NR<sup>g</sup>- where R<sup>g</sup> is H or C1-C6 alkyl,
- (m) hetAr<sup>2</sup>C(=O)NR<sup>g</sup>(CH<sub>2</sub>)<sub>p</sub>- where p is 0 or 1 and R<sup>g</sup> is H or C1-C6 alkyl,
- (n) R<sup>4</sup>R<sup>5</sup>NC(=O)-,
- (o) Ar<sup>1</sup>NR<sup>g</sup>C(=O)-, where R<sup>g</sup> is H or C1-C6 alkyl,
- (p) hetAr<sup>2</sup>NR<sup>g</sup>C(=O)-, where R<sup>g</sup> is H or C1-C6 alkyl,

- (q)  $\text{Ar}^1(\text{C1-C6 alkyl})\text{C}(=\text{O})-$  wherein said alkyl portion is optionally substituted with OH, hydroxy(C1-C6 alkyl), C1-C6 alkoxy or  $\text{NH}_2$ ,
- (r)  $\text{hetCyc}^5\text{C}(=\text{O})-$ ,
- (s)  $\text{R}^4\text{R}^5\text{NC}(=\text{O})\text{NR}^g-$  where  $\text{R}^g$  is H or C1-C6 alkyl, or
- (t)  $(\text{C1-C6 alkyl})\text{SO}_2-$ ;
- (u)  $\text{Ar}^1(\text{C1-C6 alkyl})\text{C}(=\text{O})\text{NR}^g-$  where  $\text{R}^g$  is H or C1-C6 alkyl,
- (v)  $\text{hetAr}^4\text{C}(=\text{O})\text{NR}^g-$  where  $\text{R}^g$  is H or C1-C6 alkyl,
- (w)  $\text{hetAr}^2\text{-S}(=\text{O})-$ ,
- (x)  $(\text{C3-C6 cycloalkyl})\text{CH}_2\text{SO}_2-$ ,
- (y)  $\text{Ar}^1(\text{C1-C6 alkyl})\text{SO}_2-$ ,
- (z)  $\text{hetAr}^2\text{SO}_2-$ ,
- (aa)  $\text{Ar}^1$ ,
- (bb)  $\text{hetAr}^2$ ,
- (cc)  $\text{hetCyc}^5$ ,
- (dd) C1-C6 alkoxy,
- (ee)  $\text{Ar}^1(\text{C1-C6 alkyl})\text{-O-}$ ,
- (ff)  $\text{hetAr}^2(\text{C1-C6 alkyl})\text{-O-}$ ,
- (gg)  $\text{hetAr}^2\text{-O-C1-C6 alkyl-}$ ,
- (hh)  $\text{Ar}^1(\text{C1-C6 alkyl})\text{NR}^g-$  where  $\text{R}^g$  is H or C1-C6 alkyl,
- (ii)  $\text{hetAr}^2\text{-S-}$ ,
- (jj)  $\text{Ar}^2\text{SO}_2\text{NR}^g(\text{CH}_2)_p-$  where p is 0 or 1 and  $\text{R}^g$  is H or C1-C6 alkyl,
- (kk)  $(\text{C1-C6 alkoxy})\text{C}(=\text{O})-$ ,
- (ll)  $(\text{C1-C6 alkyl})\text{NR}^g\text{C}(=\text{O})\text{O-}$  where  $\text{R}^g$  is H or C1-C6 alkyl,
- (mm)  $(\text{C1-C6 alkyl})\text{NR}^g\text{SO}_2-$  where  $\text{R}^g$  is H or C1-C6 alkyl,
- (nn)  $\text{hetCyc}^5\text{C}(=\text{O})\text{NR}^g-$  where  $\text{R}^g$  is H or C1-C6 alkyl,
- (oo)  $\text{Q-NR}^h(\text{C1-C3 alkyl})\text{C}(=\text{O})\text{NR}^g-$  where  $\text{R}^g$  and  $\text{R}^h$  are independently H or C1-C6 alkyl and Q is H, C1-C6 alkyl or  $(\text{C1-C6 alkyl})\text{OC}(=\text{O})-$ ,



(ss)  $R^s R^h N$ - where  $R^s$  and  $R^h$  are independently H or C1-C6 alkyl,

(tt) (C3-C6 cycloalkyl)C(=O)NR<sup>g</sup>- where the cycloalkyl is optionally and independently substituted with one or more halogens,

(uu) (C1-C6 alkyl)C(=O)NR<sup>g</sup>CH<sub>2</sub>- where  $R^g$  is H or C1-C6 alkyl, or

(vv) C1-C6 alkyl)SO<sub>2</sub>NR<sup>g</sup>- where  $R^g$  is H or C1-C6 alkyl;

Ar<sup>1</sup> is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros), C3-C6 cycloalkyl, hydroxyC1-C6 alkyl, (C1-C6 alkyl)SO<sub>2</sub>-, R<sup>e</sup>R<sup>f</sup>N- and (R<sup>e</sup>R<sup>f</sup>N)C1-C6 alkyl- where each R<sup>e</sup> and R<sup>f</sup> is independently H or C1-C6 alkyl;

hetAr<sup>2</sup> is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S, or a 9-10 membered bicyclic heteroaryl having 1-2 ring nitrogen atoms, wherein hetAr<sup>2</sup> is optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros) and hydroxyC1-C6 alkoxy-;

hetCyc<sup>5</sup> is a 4-6 membered saturated heterocyclic ring having 1-2 ring heteroatoms independently selected from N, O and S wherein said heterocyclic ring is optionally substituted with one or more substituents independently selected from C1-C6 alkoxy and oxo;

R<sup>3</sup> is C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl-, C1-C6 alkoxy, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)CH<sub>2</sub>-, (C3-C6 cycloalkyl)O-, (C3-C6 cycloalkyl)CH<sub>2</sub>O-, hetCyc<sup>7</sup>O-, Ph-O-, or

(C1-C6 alkoxy)C1-C6 alkyl-; wherein each of said C3-C6 cycloalkyl moieties is optionally substituted with C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy, OH or R'R"N- where R' and R" are independently hydrogen or C1-C6 alkyl;

R<sup>4</sup> is H or C1-C6 alkyl;

R<sup>5</sup> is Ar<sup>2</sup>, hetAr<sup>3</sup>, Ar<sup>2</sup>CH<sub>2</sub>-, hetCyc<sup>6</sup>-CH<sub>2</sub>-, hydroxyC1-C6 alkyl-, (C3-C6 cycloalkyl)CH<sub>2</sub>-, or C1-C6 alkyl optionally substituted with 1-3 fluoros;

Ar<sup>2</sup> is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros), C3-C6 cycloalkyl, and R<sup>g</sup>R<sup>h</sup>N- where R<sup>g</sup> and R<sup>h</sup> are independently H or C1-C6 alkyl, or Ar<sup>2</sup> is phenyl fused to a 6 membered heterocyclic ring having a ring nitrogen atom and optionally substituted with C1-C6 alkyl;


hetAr<sup>3</sup> is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S and optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), and (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros);

hetAr<sup>4</sup> is pyridin-4(1H)-onyl or pyridin-2(1H)-onyl optionally substituted with one or more substituents independently selected from C1-C6 alkyl and halogen;

hetCyc<sup>6</sup> is a 5-7 membered heterocyclic ring having 1-3 ring heteroatoms independently selected from N, O and S; and

hetCyc<sup>7</sup> is a 5-7 membered heterocyclic ring having 1-3 ring heteroatoms independently selected from N, O and S.

**[00431]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is a compound of the Formula **VII**:



**VII**

or a pharmaceutically acceptable salt or solvate thereof, wherein:

X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> are independently CH or N, wherein zero, one or two of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup> and X<sup>4</sup> is N;

A is CN;

B is:

(b) C1-C6 alkyl optionally substituted with 1-3 fluoros,

(c) hydroxyC2-C6 alkyl- wherein the alkyl portion is optionally substituted with a C3-C6 cycloalkylidene ring, or

(i) (hetCyc<sup>a</sup>)C1-C3 alkyl-;

hetCyc<sup>a</sup> is a 4-6 membered heterocyclic ring having 1-2 ring heteroatoms independently selected from N and O and optionally substituted with one or more substituents independently selected from OH, C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl, halogen, (C1-C6 alkyl)C(=O)-, C1-C6 alkoxy, oxo, and (C1-C6 alkoxy)C(=O)-;

Ring D is a saturated monocyclic 4-7 membered heterocyclic ring having one ring heteroatom which is nitrogen;

each R<sup>a</sup> is independently C1-C6 alkyl (optionally substituted with 1-3 fluoros);

R<sup>b</sup> is (a) hydroxy;

n is 0 or 1;

m is 0 or 1;

E is:

(e) hetAr<sup>2</sup>C1-C6 alkyl-,

(h) hetAr<sup>2</sup>-O-,

(k) R<sup>3</sup>C(=O)NR<sup>g</sup>- where R<sup>g</sup> is H or C1-C6 alkyl,

(l) Ar<sup>1</sup>C(=O)NR<sup>g</sup>- where R<sup>g</sup> is H or C1-C6 alkyl, or

(m) hetAr<sup>2</sup>C(=O)NR<sup>g</sup>(CH<sub>2</sub>)<sub>p</sub>- where p is 0 or 1 and R<sup>g</sup> is H or C1-C6 alkyl;

Ar<sup>1</sup> is phenyl optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros), C3-C6 cycloalkyl, hydroxyC1-C6 alkyl, (C1-C6 alkyl)SO<sub>2</sub>-, R<sup>e</sup>R<sup>f</sup>N- and (R<sup>e</sup>R<sup>f</sup>N)C1-C6 alkyl- where each R<sup>e</sup> and R<sup>f</sup> is independently H or C1-C6 alkyl;

hetAr<sup>2</sup> is a 5-6 membered heteroaryl ring having 1-3 ring heteroatoms independently selected from N, O and S, or a 9-10 membered bicyclic heteroaryl having 1-2 ring nitrogen atoms, wherein hetAr<sup>2</sup> is optionally substituted with one or more substituents independently selected from the group consisting of halogen, CN, C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy (optionally substituted with 1-3 fluoros), (C1-C6 alkoxy)C1-C6 alkyl- (optionally substituted with 1-3 fluoros) and hydroxyC1-C6 alkoxy-; and



R<sup>3</sup> is C1-C6 alkyl (optionally substituted with 1-3 fluoros), hydroxyC1-C6 alkyl-, C1-C6 alkoxy, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)CH<sub>2</sub>-, (C3-C6 cycloalkyl)O-, (C3-C6 cycloalkyl)CH<sub>2</sub>O-, hetCyc<sup>7</sup>O-, Ph-O-, or (C1-C6 alkoxy)C1-C6 alkyl-; wherein each of said C3-C6 cycloalkyl moieties is optionally substituted with C1-C6 alkyl (optionally substituted with 1-3 fluoros), C1-C6 alkoxy, OH, or R'R''N- where R' and R'' are independently hydrogen or C1-C6 alkyl.

**[00432]** In some embodiments, a RET inhibitor (e.g., a first RET inhibitor or a second RET inhibitor) is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof.

**[00433]** Non-limiting examples of receptor tyrosine kinase (e.g., Trk) targeted therapeutic agents, include afatinib, cabozantinib, cetuximab, crizotinib, dabrafenib, entrectinib, erlotinib, gefitinib, imatinib, lapatinib, lestaurtinib, nilotinib, pazopanib, panitumumab, pertuzumab, sunitinib, trastuzumab, I-((3S,4R)-4-(3-fluorophenyl)-I-(2-methoxyethyl)pyrrolidin-3-yl)-3-(4-methyl-3-(2-methylpyrimidin-5-yl)-I-phenyl)-IH-pyrazol-5-yl)urea, AG 879, AR-772, AR-786, AR-256, AR-618, AZ-23, AZ623, DS-6051, Gö 6976, GNF-5837, GTx-186, GW 441756, LOXO-101, MGCD516, PLX7486, RXDX101, VM-902A, TPX-0005, and TSR-011. Additional Trk targeted therapeutic agents include those described in U.S. Patent No. 8,450,322; 8,513,263; 8,933,084; 8,791,123; 8,946,226; 8,450,322; 8,299,057; and 8,912,194; U.S. Publication No. 2016/0137654; 2015/0166564; 2015/0051222; 2015/0283132; and 2015/0306086; International Publication No. WO 2010/033941; WO 2010/048314; WO 2016/077841; WO 2011/146336; WO 2011/006074; WO 2010/033941; WO 2012/158413; WO 2014078454; WO 2014078417; WO 2014078408; WO 2014078378; WO 2014078372; WO 2014078331; WO 2014078328; WO 2014078325; WO 2014078323; WO 2014078322; WO 2015175788; WO 2009/013126; WO 2013/174876; WO 2015/124697; WO 2010/058006; WO 2015/017533; WO 2015/112806; WO 2013/183578; and WO 2013/074518, all of which are hereby incorporated by reference in their entireties.

**[00434]** Further examples of Trk inhibitors can be found in U.S. Patent No. 8,637,516, International Publication No. WO 2012/034091, U.S. Patent No. 9,102,671, International Publication No. WO 2012/116217, U.S. Publication No. 2010/0297115, International Publication No. WO 2009/053442, U.S. Patent No. 8,642,035, International Publication No. WO 2009092049, U.S. Patent No. 8,691,221, International Publication No. WO2006131952, all of which are incorporated by reference in their

entireties herein. Exemplary Trk inhibitors include GNF-4256, described in *Cancer Chemother. Pharmacol.* 75(1):131-141, 2015; and GNF-5837 (N-[3-[[2,3-dihydro-2-oxo-3-(1H-pyrrol-2-ylmethylene)-1H-indol-6-yl]amino]-4-methylphenyl]-N'-[2-fluoro-5-(trifluoromethyl)phenyl]-urea), described in *ACS Med. Chem. Lett.* 3(2):140-145, 2012, each of which is incorporated by reference in its entirety herein.

**[00435]** Additional examples of Trk inhibitors include those disclosed in U.S. Publication No. 2010/0152219, U.S. Patent No. 8,114,989, and International Publication No. WO 2006/123113, all of which are incorporated by reference in their entireties herein. Exemplary Trk inhibitors include AZ623, described in *Cancer* 117(6):1321-1391, 2011; AZD6918, described in *Cancer Biol. Ther.* 16(3):477-483, 2015; AZ64, described in *Cancer Chemother. Pharmacol.* 70:477-486, 2012; AZ-23 ((S)-5-Chloro-N2-(1-(5-fluoropyridin-2-yl)ethyl)-N4-(5-isopropoxy-1H-pyrazol-3-yl)pyrimidine-2,4-diamine), described in *Mol. Cancer Ther.* 8:1818-1827, 2009; and AZD7451; each of which is incorporated by reference in its entirety.

**[00436]** A Trk inhibitor can include those described in U.S. Patent Nos. 7,615,383; 7,384,632; 6,153,189; 6,027,927; 6,025,166; 5,910,574; 5,877,016; and 5,844,092, each of which is incorporated by reference in its entirety.

**[00437]** Further examples of Trk inhibitors include CEP-751, described in *Int. J. Cancer* 72:672-679, 1997; CT327, described in *Acta Derm. Venereol.* 95:542-548, 2015; compounds described in International Publication No. WO 2012/034095; compounds described in U.S. Patent No. 8,673,347 and International Publication No. WO 2007/022999; compounds described in U.S. Patent No. 8,338,417; compounds described in International Publication No. WO 2016/027754; compounds described in U.S. Patent No. 9,242,977; compounds described in U.S. Publication No. 2016/0000783; sunitinib (N-(2-diethylaminoethyl)-5-[(Z)-(5-fluoro-2-oxo-1H-indol-3-ylidene)methyl]-2,4-dimethyl-1H-pyrrole-3-carboxamide), as described in *PLoS One* 9:e95628, 2014; compounds described in International Publication No. WO 2011/133637; compounds described in U.S. Patent No. 8,637,256; compounds described in *Expert. Opin. Ther. Pat.* 24(7):731-744, 2014; compounds described in *Expert Opin. Ther. Pat.* 19(3):305-319, 2009; (R)-2-phenylpyrrolidine substituted imidazopyridazines, e.g., GNF-8625, (R)-1-(6-(6-(2-(3-fluorophenyl)pyrrolidin-1-yl)imidazo[1,2-b]pyridazin-3-yl)-[2,4'-bipyridin]-2'-yl)piperidin-4-ol as described in *ACS Med. Chem. Lett.* 6(5):562-567, 2015; GTx-186 and others, as described in *PLoS One* 8(12):e83380, 2013; K252a ((9S-(9 $\alpha$ ,10 $\beta$ ,12 $\alpha$ ))-2,3,9,10,11,12-hexahydro-10-hydroxy-10-(methoxycarbonyl)-9-methyl-9,12-epoxy-1H-diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one), as described in *Mol. Cell Biochem.* 339(1-2):201-213, 2010; 4-aminopyrazolylpyrimidines, e.g., AZ-23 (((S)-5-chloro-N2-(1-(5-fluoropyridin-2-yl)ethyl)-N4-(5-isopropoxy-1H-pyrazol-3-yl)pyrimidine-2,4-diamine)), as described in *J. Med. Chem.* 51(15):4672-4684, 2008; PHA-739358 (danusertib), as described in *Mol. Cancer Ther.* 6:3158, 2007; Gö 6976 (5,6,7,13-tetrahydro-13-methyl-5-oxo-12H-indolo[2,3-a]pyrrolo[3,4-c]carbazole-12-propanenitrile), as described in *J. Neurochem.* 72:919-924, 1999; GW441756 ((3Z)-3-[(1-methylindol-3-yl)methylidene]-1H-pyrrolo[3,2-b]pyridin-2-one), as described in *IJAE* 115:117, 2010; milciclib (PHA-848125AC), described in *J. Carcinog.* 12:22, 2013; AG-879 ((2E)-3-[3,5-Bis(1,1-dimethylethyl)-4-hydroxyphenyl]-2-cyano-2-propenethioamide); altiratinib (N-(4-((2-(cyclopropanecarboxamido)pyridin-4-yl)oxy)-2,5-difluorophenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide); cabozantinib (N-(4-((6,7-Dimethoxyquinolin-4-yl)oxy)phenyl)-N'-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide); lestaurtinib ((5S,6S,8R)-6-Hydroxy-6-(hydroxymethyl)-5-methyl-7,8,14,15-tetrahydro-5H-16-oxa-4b,8a,14-triaza-5,8-

methanodibenzo[b,h]cycloocta[jkl]cyclopenta[e]-as-indacen-13(6H)-one); dovatinib (4-amino-5-fluoro-3-[6-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]quinolin-2(1H)-one mono 2-hydroxypropanoate hydrate); sitravatinib (N-(3-fluoro-4-((2-(5-(((2-methoxyethyl)amino)methyl)pyridin-2-yl)thieno[3,2-b]pyridin-7-yl)oxy)phenyl)-N-(4-fluorophenyl)cyclopropane-1,1-dicarboxamide); ONO-5390556; regorafenib (4-[4-({[4-Chloro-3-(trifluoromethyl)phenyl]carbonyl}amino)-3-fluorophenoxy]-N-methylpyridine-2-carboxamide hydrate); and VSR-902A; all of the references above are incorporated by reference in their entireties herein.

**[00438]** The ability of a Trk inhibitor to act as a TrkA, TrkB, and/or Trk C inhibitor may be tested using the assays described in Examples A and B in U.S. Patent No. 8,513,263, which is incorporated herein by reference.

**[00439]** In some embodiments, the receptor tyrosine kinase inhibitor is an epidermal growth factor receptor tyrosine kinase inhibitor (EGFR). For example, EGFR inhibitors can include osimertinib (merelectinib, Tagrisso), erlotinib (Tarceva), gefitinib (Iressa), cetuximab (Erbix), necitumumab (Portrazza), neratinib (Nerlynx), lapatinib (Tykerb), panitumumab (Vectibix), and vandetanib (Caprelsa). In some embodiments, the EGFR inhibitor is osimertinib.

**[00440]** In some embodiments, signal transduction pathway inhibitors include Ras-Raf-MEK-ERK pathway inhibitors (e.g., binimetinib, selumetinib, encorafenib, sorafenib, trametinib, and vemurafenib), PI3K-Akt-mTOR-S6K pathway inhibitors (e.g. everolimus, rapamycin, perifosine, temsirolimus), and other kinase inhibitors, such as baricitinib, brigatinib, capmatinib, danusertib, ibrutinib, milciclib, quercetin, regorafenib, ruxolitinib, semaxanib, AP32788, BLU285, BLU554, INCB39110, INCB40093, INCB50465, INCB52793, INCB54828, MGCD265, NMS-088, NMS-1286937, PF 477736 ((R)-amino-N-[5,6-dihydro-2-(1-methyl-1H-pyrazol-4-yl)-6-oxo-1H-pyrrolo[4,3,2-ef][2,3]benzodiazepin-8-yl]-cyclohexanecarboxamide), PLX3397, PLX7486, PLX8394, PLX9486, PRN1008, PRN1371, RXDX103, RXDX106, RXDX108, and TG101209 (N-tert-butyl-3-(5-methyl-2-(4-(4-methylpiperazin-1-yl)phenylamino)pyrimidin-4-ylamino)benzenesulfonamide).

**[00441]** Non-limiting examples of checkpoint inhibitors include ipilimumab, tremelimumab, nivolumab, pidilizumab, MPDL3208A, MEDI4736, MSB0010718C, BMS-936559, BMS-956559, BMS-935559 (MDX-1105), AMP-224, and pembrolizumab.

**[00442]** In some embodiments, cytotoxic chemotherapeutics are selected from arsenic trioxide, bleomycin, cabazitaxel, capecitabine, carboplatin, cisplatin, cyclophosphamide, cytarabine, dacarbazine, daunorubicin, docetaxel, doxorubicin, etoposide, fluorouracil, gemcitabine, irinotecan, lomustine, methotrexate, mitomycin C, oxaliplatin, paclitaxel, pemetrexed, temozolomide, and vincristine.

**[00443]** Non-limiting examples of angiogenesis-targeted therapies include aflibercept and bevacizumab.

**[00444]** The term “immunotherapy” refers to an agent that modulates the immune system. In some embodiments, an immunotherapy can increase the expression and/or activity of a regulator of the immune system. In some embodiments, an immunotherapy can decrease the expression and/or activity of a regulator of the immune system. In some embodiments, an immunotherapy can recruit and/or enhance the activity of an immune cell.

**[00445]** In some embodiments, the immunotherapy is a cellular immunotherapy (e.g., adoptive T-cell therapy, dendritic cell therapy, natural killer cell therapy). In some embodiments, the cellular immunotherapy is sipuleucel-T (APC8015; Provenge™; Plosker (2011) *Drugs* 71(1): 101-108). In some embodiments, the cellular immunotherapy includes cells that express a chimeric antigen receptor (CAR). In some embodiments, the cellular immunotherapy is a CAR-T cell therapy. In some embodiments, the CAR-T cell therapy is tisagenlecleucel (Kymriah™).

**[00446]** In some embodiments, the immunotherapy is an antibody therapy (e.g., a monoclonal antibody, a conjugated antibody). In some embodiments, the antibody therapy is bevacizumab (Mvasti™, Avastin®), trastuzumab (Herceptin®), avelumab (Bavencio®), rituximab (MabThera™, Rituxan®), edrecolomab (Panorex), daratumumab (Darzalex®), olaratumab (Lartruvo™), ofatumumab (Arzerra®), alemtuzumab (Campath®), cetuximab (Erbix™), oregovomab, pembrolizumab (Keytruda®), dinutuximab (Unituxin®), obinutuzumab (Gazyva®), tremelimumab (CP-675,206), ramucirumab (Cyramza®), ublituximab (TG-1101), panitumumab (Vectibix®), elotuzumab (Empliciti™), avelumab (Bavencio®), necitumumab (Portrazza™), cirmutuzumab (UC-961), ibritumomab (Zevalin®), isatuximab (SAR650984), nimotuzumab, fresolimumab (GC1008), lirilumab (INN), mogamulizumab (Poteligeo®), ficlatuzumab (AV-299), denosumab (Xgeva®), ganitumab, urelumab, pidilizumab or amatuximab.

**[00447]** In some embodiments, the immunotherapy is an antibody-drug conjugate. In some embodiments, the antibody-drug conjugate is gemtuzumab ozogamicin (Mylotarg™), inotuzumab ozogamicin (Besponsa®), brentuximab vedotin (Adcetris®), ado-trastuzumab emtansine (TDM-1; Kadcyla®), mirvetuximab soravtansine (IMGN853) or anetumab ravtansine.

**[00448]** In some embodiments, the immunotherapy includes blinatumomab (AMG103; Blincyto®) or midostaurin (Rydapt).

**[00449]** In some embodiments, the immunotherapy includes a toxin. In some embodiments, the immunotherapy is denileukin diftitox (Ontak®).

**[00450]** In some embodiments, the immunotherapy is a cytokine therapy. In some embodiments, the cytokine therapy is an interleukin 2 (IL-2) therapy, an interferon alpha (IFNα) therapy, a granulocyte colony stimulating factor (G-CSF) therapy, an interleukin 12 (IL-12) therapy, an interleukin 15 (IL-15) therapy, an interleukin 7 (IL-7) therapy or an erythropoietin-alpha (EPO) therapy. In some embodiments, the IL-2 therapy is aldesleukin (Proleukin®). In some embodiments, the IFNα therapy is IntronA® (Roferon-A®). In some embodiments, the G-CSF therapy is filgrastim (Neupogen®).

**[00451]** In some embodiments, the immunotherapy is an immune checkpoint inhibitor. In some embodiments, the immunotherapy includes one or more immune checkpoint inhibitors. In some embodiments, the immune checkpoint inhibitor is a CTLA-4 inhibitor, a PD-1 inhibitor or a PD-L1 inhibitor. In some embodiments, the CTLA-4 inhibitor is ipilimumab (Yervoy®) or tremelimumab (CP-675,206). In some embodiments, the PD-1 inhibitor is pembrolizumab (Keytruda®) or nivolumab (Opdivo®). In some embodiments, the PD-L1 inhibitor is atezolizumab (Tecentriq®), avelumab (Bavencio®) or durvalumab (Imfinzi™).

**[00452]** In some embodiments, the immunotherapy is mRNA-based immunotherapy. In some embodiments, the mRNA-based immunotherapy is CV9104 (see, e.g., Rausch et al. (2014) Human Vaccin Immunother 10(11): 3146-52; and Kubler et al. (2015) J. Immunother Cancer 3:26).

**[00453]** In some embodiments, the immunotherapy is bacillus Calmette-Guerin (BCG) therapy.

**[00454]** In some embodiments, the immunotherapy is an oncolytic virus therapy. In some embodiments, the oncolytic virus therapy is talimogene alherparepvec (T-VEC; Imlygic®).

**[00455]** In some embodiments, the immunotherapy is a cancer vaccine. In some embodiments, the cancer vaccine is a human papillomavirus (HPV) vaccine. In some embodiments, the HPV vaccine is Gardasil®, Gardasil9® or Cervarix®. In some embodiments, the cancer vaccine is a hepatitis B virus (HBV) vaccine. In some embodiments, the HBV vaccine is Engerix-B®, Recombivax HB® or Gl-13020 (Tarmogen®). In some embodiments, the cancer vaccine is Twinrix® or Pediarix®. In some embodiments, the cancer vaccine is BiovaxID®, Oncophage®, GVAX, ADXS11-001, ALVAC-CEA, PROSTVAC®, Rindopepimut®, CimaVax-EGF, lapuleucel-T (APC8024; Neuvence™), GRNVAC1, GRNVAC2, GRN-1201, hepcortespenlisimut-L (Hepko-V5), DCVAX®, SCIB1, BMT CTN 1401, PrCa VBIR, PANVAC, ProstAtak®, DPX-Survivac, or viagenpumatucl-L (HS-110).

**[00456]** In some embodiments, the immunotherapy is a peptide vaccine. In some embodiments, the peptide vaccine is neli pepimut-S (E75) (NeuVax™), IMA901, or SurVaxM (SVN53-67). In some embodiments, the cancer vaccine is an immunogenic personal neoantigen vaccine (see, e.g., Ott et al. (2017) Nature 547: 217-221; Sahin et al. (2017) Nature 547: 222-226). In some embodiments, the cancer vaccine is RGS4K, or NEO-PV-01. In some embodiments, the cancer vaccine is a DNA-based vaccine. In some embodiments, the DNA-based vaccine is a mammaglobin-A DNA vaccine (see, e.g., Kim et al. (2016) OncoImmunology 5(2): e1069940).

**[00457]** In some embodiments, immune-targeted agents are selected from aldesleukin, interferon alfa-2b, ipilimumab, lambrolizumab, nivolumab, prednisone, and sipuleucel-T.

**[00458]** Non-limiting examples of radiotherapy include radioiodide therapy, external-beam radiation, and radium 223 therapy.

**[00459]** Additional kinase inhibitors include those described in, for example, U.S. Patent No. 7,514,446; 7,863,289; 8,026,247; 8,501,756; 8,552,002; 8,815,901; 8,912,204; 9,260,437; 9,273,051; U.S. Publication No. US 2015/0018336; International Publication No. WO 2007/002325; WO 2007/002433; WO 2008/080001; WO 2008/079906; WO 2008/079903; WO 2008/079909; WO 2008/080015; WO 2009/007748; WO 2009/012283; WO 2009/143018; WO 2009/143024; WO 2009/014637; WO 2009/152083; WO 2010/111527; WO 2012/109075; WO 2014/194127; WO 2015/112806; WO 2007/110344; WO 2009/071480; WO 2009/118411; WO 2010/031816; WO 2010/145998; WO 2011/092120; WO 2012/101032; WO 2012/139930; WO 2012/143248; WO 2012/152763; WO 2013/014039; WO 2013/102059; WO 2013/050448; WO 2013/050446; WO 2014/019908; WO 2014/072220; WO 2014/184069; and WO 2016/075224 all of which are hereby incorporated by reference in their entireties.

**[00460]** Further examples of kinase inhibitors include those described in, for example, WO 2016/081450; WO 2016/022569; WO 2016/011141; WO 2016/011144; WO 2016/011147; WO 2015/191667; WO 2012/101029; WO 2012/113774; WO 2015/191666; WO 2015/161277; WO 2015/161274; WO 2015/108992; WO 2015/061572; WO 2015/058129; WO 2015/057873; WO 2015/017528; WO/2015/017533; WO 2014/160521; and WO 2014/011900, each of which is hereby incorporated by reference in its entirety.

**[00461]** Further examples of kinase inhibitors include luminespib (AUY-922, NVP-AUY922) (5-(2,4-dihydroxy-5-isopropylphenyl)-N-ethyl-4-(4-(morpholinomethyl)phenyl)isoxazole-3-carboxamide) and doramapimod (BIRB-796) (1-[5-tert-butyl-2-(4-methylphenyl)pyrazol-3-yl]-3-[4-(2-morpholin-4-ylethoxy)naphthalen-1-yl]urea).

**[00462]** Accordingly, also provided herein is a method of treating cancer, comprising administering to a patient in need thereof a pharmaceutical combination for treating cancer which comprises (a) a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, (b) an additional therapeutic agent, and (c) optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use for the treatment of cancer, wherein the amounts of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof and the additional therapeutic agent are together effective in treating the cancer.

**[00463]** In some embodiments, the additional therapeutic agent(s) includes any one of the above listed therapies or therapeutic agents which are standards of care in cancers wherein the cancer has a dysregulation of a RET gene, a RET protein, or expression or activity, or level of any of the same.

**[00464]** These additional therapeutic agents may be administered with one or more doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, or pharmaceutical composition thereof, as part of the same or separate dosage forms, via the same or different routes of administration, and/or on the same or different administration schedules according to standard pharmaceutical practice known to one skilled in the art.

**[00465]** Also provided herein is (i) a pharmaceutical combination for treating a cancer in a patient in need thereof, which comprises (a) a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, (b) at least one additional therapeutic agent (e.g., any of the exemplary additional therapeutic agents described herein or known in the art), and (c) optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use for the treatment of cancer, wherein the amounts of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof and of the additional therapeutic agent are together effective in treating the cancer; (ii) a pharmaceutical composition comprising such a combination; (iii) the use of such a combination for the preparation of a medicament for the treatment of cancer; and (iv) a commercial package or product comprising such a combination as a combined preparation for simultaneous, separate or sequential use; and to a method of treatment of cancer in a patient in need thereof. In one embodiment the patient is a human. In some embodiments, the

cancer is a RET-associated cancer. For example, a RET-associated cancer having one or more RET inhibitor resistance mutations.

**[00466]** The term "pharmaceutical combination", as used herein, refers to a pharmaceutical therapy resulting from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term "fixed combination" means that a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and at least one additional therapeutic agent (e.g., a chemotherapeutic agent), are both administered to a patient simultaneously in the form of a single composition or dosage. The term "non-fixed combination" means that a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and at least one additional therapeutic agent (e.g., chemotherapeutic agent) are formulated as separate compositions or dosages such that they may be administered to a patient in need thereof simultaneously, concurrently or sequentially with variable intervening time limits, wherein such administration provides effective levels of the two or more compounds in the body of the patient. These also apply to cocktail therapies, e.g. the administration of three or more active ingredients

**[00467]** Accordingly, also provided herein is a method of treating a cancer, comprising administering to a patient in need thereof a pharmaceutical combination for treating cancer which comprises (a) a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, (b) an additional therapeutic agent, and (c) optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use for the treatment of cancer, wherein the amounts of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof and the additional therapeutic agent are together effective in treating the cancer. In one embodiment, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, and the additional therapeutic agent are administered simultaneously as separate dosages. In one embodiment, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, and the additional therapeutic agent are administered as separate dosages sequentially in any order, in jointly therapeutically effective amounts, e.g. in daily or intermittently dosages. In one embodiment, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, and the additional therapeutic agent are administered simultaneously as a combined dosage. In some embodiments, the cancer is a RET-associated cancer. For example, a RET-associated cancer having one or more RET inhibitor resistance mutations. In some embodiments, the additional therapeutic agent is crizotinib. In some embodiments, the additional therapeutic agent is osimertinib. In some embodiments, the patient has been administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, prior to administration of the pharmaceutical composition. In some embodiments, the cancer is a lung cancer (e.g., a RET-associated lung cancer).

**[00468]** Also provided herein is a method of treating a disease or disorder mediated by RET in a patient in need of such treatment, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition

thereof. In some embodiments, the disease or disorder mediated by RET is a dysregulation of RET gene, a RET kinase, or expression or activity or level of any of the same. For example the dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same includes one or more RET inhibitor resistance mutations. A disease or disorder mediated by RET can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of RET, including overexpression and/or abnormal activity levels. In one embodiment, the disease is cancer (e.g., a RET-associated cancer). In one embodiment, the cancer is any of the cancers or RET-associated cancers described herein. In some embodiments, the additional therapeutic agent is crizotinib. In some embodiments, the additional therapeutic agent is osimertinib. In some embodiments, the patient has been administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, prior to administration of the pharmaceutical composition. In some embodiments, the cancer is a lung cancer (e.g., a RET-associated lung cancer).

**[00469]** Although the genetic basis of tumorigenesis may vary between different cancer types, the cellular and molecular mechanisms required for metastasis appear to be similar for all solid tumor types. During a metastatic cascade, the cancer cells lose growth inhibitory responses, undergo alterations in adhesiveness and produce enzymes that can degrade extracellular matrix components. This leads to detachment of tumor cells from the original tumor, infiltration into the circulation through newly formed vasculature, migration and extravasation of the tumor cells at favorable distant sites where they may form colonies. A number of genes have been identified as being promoters or suppressors of metastasis. For example, overexpression of glial cell-derived neurotrophic factor (GDNF) and its RET receptor tyrosine kinase have been correlated with cancer proliferation and metastasis. See, e.g., Zeng, Q. et al. *J. Int. Med. Res.* (2008) 36(4): 656-64.

**[00470]** Accordingly, also provided herein are methods for inhibiting, preventing, aiding in the prevention, or decreasing the symptoms of metastasis of a cancer in a patient in need thereof, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or a pharmaceutical composition thereof. Such methods can be used in the treatment of one or more of the cancers described herein. See, e.g., US Publication No. 2013/0029925; International Publication No. WO 2014/083567; and US Patent No. 8,568,998. See also, e.g., Hezam K et al., *Rev Neurosci* 2018 Jan 26;29:93-98; Gao L, et al., *Pancreas* 2015 Jan;44:134-143; Ding K et al., *J Biol Chem* 2014 Jun 6; 289:16057-71; and Amit M et al., *Oncogene* 2017 Jun 8; 36:3232-3239. In some embodiments, the cancer is a RET-associated cancer. In some embodiments, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is used in combination with an additional therapy or another therapeutic agent, including a chemotherapeutic agent, such as a kinase inhibitor. For example, a first or second RET kinase inhibitor. In some embodiments, the additional therapeutic agent is crizotinib. In some embodiments, the additional therapeutic agent is osimertinib. In some embodiments,



the patient has been administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, prior to administration of the pharmaceutical composition. In some embodiments, the cancer is a lung cancer (e.g., a RET-associated lung cancer).

**[00471]** The term “metastasis” is an art known term and means the formation of an additional tumor (e.g., a solid tumor) at a site distant from a primary tumor in a subject or patient, where the additional tumor includes the same or similar cancer cells as the primary tumor.

**[00472]** Also provided are methods of decreasing the risk of developing a metastasis or an additional metastasis in a patient having a RET-associated cancer that include: selecting, identifying, or diagnosing a patient as having a RET-associated cancer, and administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the patient selected, identified, or diagnosed as having a RET-associated cancer. Also provided are methods of decreasing the risk of developing a metastasis or an additional metastasis in a patient having a RET-associated cancer that includes administering a therapeutically effective amount of a Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to a patient having a RET-associated cancer. The decrease in the risk of developing a metastasis or an additional metastasis in a patient having a RET-associated cancer can be compared to the risk of developing a metastasis or an additional metastasis in the patient prior to treatment, or as compared to a patient or a population of patients having a similar or the same RET-associated cancer that has received no treatment or a different treatment. In some embodiments, the RET-associated cancer is a RET-associated cancer having one or more RET inhibitor resistance mutations. In some embodiments, the additional therapeutic agent is crizotinib. In some embodiments, the additional therapeutic agent is osimertinib. In some embodiments, the patient has been administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, prior to administration of the pharmaceutical composition. In some embodiments, the cancer is a lung cancer (e.g., a RET-associated lung cancer).

**[00473]** The phrase “risk of developing a metastasis” means the risk that a subject or patient having a primary tumor will develop an additional tumor (e.g., a solid tumor) at a site distant from a primary tumor in a subject or patient over a set period of time, where the additional tumor includes the same or similar cancer cells as the primary tumor. Methods for reducing the risk of developing a metastasis in a subject or patient having a cancer are described herein.

**[00474]** The phrase “risk of developing additional metastases” means the risk that a subject or patient having a primary tumor and one or more additional tumors at sites distant from the primary tumor (where the one or more additional tumors include the same or similar cancer cells as the primary tumor) will develop one or more further tumors distant from the primary tumor, where the further tumors include the same or similar cancer cells as the primary tumor. Methods for reducing the risk of developing additional metastasis are described herein.

**[00475]** In some embodiments, the presence of one or more RET inhibitor resistance mutations in a tumor causes the tumor to be more resistant to treatment with a first RET inhibitor. Methods useful

when a RET inhibitor resistance mutation causes the tumor to be more resistant to treatment with a first RET inhibitor are described below. For example, provided herein are methods of treating a subject having a cancer that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and administering to the identified subject a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is administered in combination with the first RET inhibitor. Also provided are methods of treating a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations that include administering to the subject a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is administered in combination with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D.

**[00476]** For example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00477]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance

mutation; and (d) administering a compound selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00478]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a compound selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00479]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00480]** As another example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a

sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method

comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L) selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00481]** As another example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second therapeutic agent, wherein the second therapeutic agent is selected from the group consisting of crizotinib and osimertinib, as a monotherapy or in conjunction with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions of Table 2 in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a second therapeutic agent, wherein the second therapeutic agent is selected from the group consisting of crizotinib and osimertinib, as a monotherapy or in conjunction with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments of the above, the RET-associated cancer is a lung cancer.

**[00482]** In some embodiments, the presence of one or more RET inhibitor resistance mutations in a tumor causes the tumor to be more resistant to treatment with a first RET inhibitor. Methods useful when a RET inhibitor resistance mutation causes the tumor to be more resistant to treatment with a first RET inhibitor are described below. For example, provided herein are methods of treating a subject having a cancer that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and administering to the identified subject a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is administered in combination with the first RET inhibitor. Also provided are methods of treating a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations that include administering to the subject a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is administered in combination with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D.

**[00483]** For example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET

inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00484]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L) selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00485]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-

yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L) selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00486]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00487]** As another example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any



one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-

3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: ((S)-4-(6-(4-(2-hydroxy-3-phenylpropanoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(2-(pyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2,6-difluorobenzoyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-

carbonitrile 2,2,2-trifluoroacetate; 4-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N,N-diethylpiperazine-1-carboxamide; 1-(5-(3-cyano-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-N-(2-methoxy-3-methylbutyl)piperidine-4-carboxamide; 4-(6-(4-(2-(5-fluoropyridin-2-yl)acetyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile bis(2,2,2-trifluoroacetate); 4-(6-(4-(2,6-difluorobenzyl)piperazin-1-yl)pyridine-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(2-methoxybenzyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(1-methyl-1H-pyrazol-4-yl)-4-(6-(4-(pyridine-2-ylmethyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(4-(6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)-6-(1-methyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00488]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00489]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00490]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-

morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00491]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a compound of Formula I selected from Examples 1-151 or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00492]** As another example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b)

administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151 or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-

(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a second RET inhibitor, wherein the second

RET inhibitor is selected from the group consisting of: 4-(6-(4-benzylpiperazin-1-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxyethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (R)-6-(2-hydroxypropoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-methoxyethoxy)-4-(6-(4-((6-methoxypyridin-3-yl)methyl)piperazin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(6-(6-methoxynicotinoyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-(dimethylamino)ethoxy)-4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-(2-morpholinoethoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; 4-(6-(6-((6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyridin-3-yl)-6-((1-methyl-1H-imidazol-4-yl)methoxy)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 6-ethoxy-4-(5-(6-((5-fluoro-6-methoxypyridin-3-yl)methyl)-3,6-diazabicyclo[3.1.1]heptan-3-yl)pyrazin-2-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00493]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at



least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00494]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00495]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-

fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00496]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first RET inhibitor, wherein the first RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the first RET inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00497]** As another example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any

one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a

pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; (S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a second RET inhibitor, wherein the second RET inhibitor is selected from the group consisting of: N-(1-(5-(3-cyano-6-(2-hydroxy-2-methylpropoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)benzamide; 6-ethoxy-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)azetidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-((6-methoxypyridazin-3-yl)oxy)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile;

(S)-6-(2-hydroxy-2-methylpropoxy)-4-(6-(3-(pyridin-2-yloxy)pyrrolidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)-5-fluoro-2-methylbenzamide; 3-chloro-N-(1-(5-(3-cyano-6-((3-fluoro-1-methylazetidin-3-yl)methoxy)pyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-4-methylpiperidin-4-yl)picolinamide; N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyridin-2-yl)-3-hydroxypiperidin-4-yl)-3-methylbutanamide; 6-(2-hydroxy-2-methylpropoxy)-4-(6-(4-hydroxy-4-(pyridin-2-ylmethyl)piperidin-1-yl)pyridin-3-yl)pyrazolo[1,5-a]pyridine-3-carbonitrile; and 3-chloro-N-((3S,4S)-1-(5-(3-cyano-6-ethoxypyrazolo[1,5-a]pyridin-4-yl)pyrazin-2-yl)-3-hydroxypiperidin-4-yl)picolinamide; or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00498]** In some embodiments provided herein, circulating tumor DNA can be used to monitor the responsiveness of a patient to a particular therapy (e.g., a first RET inhibitor, a second RET inhibitor, or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof). For example, prior to starting treatment with a therapy as described herein (e.g., a first RET inhibitor, a second RET inhibitor, or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof), a biological sample can be obtained from the subject and the level of circulating tumor DNA determined in the biological sample. This sample can be considered a base-line sample. The subject can then be administered one or more doses of a therapy as described herein (e.g., a first RET inhibitor, a second RET inhibitor, or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof) and the levels of circulating tumor DNA can be monitored (e.g., after the first dose, second dose, third dose, etc. or after one week, two weeks, three weeks, four weeks, etc.). If the level of circulating tumor DNA is lower than the baseline sample (e.g., a 1% to about a 99% reduction, a 1% to about a 95% reduction, a 1% to about a 90% reduction, a 1% to about a 85% reduction, a 1% to about a 80% reduction, a 1% to about a 75% reduction, a 1% reduction to about a 70% reduction, a 1% reduction to about a 65% reduction, a 1% reduction to about a 60% reduction, a 1% reduction to about a 55% reduction, a 1% reduction to about a 50% reduction, a 1% reduction to about a 45% reduction, a 1% reduction to about a 40% reduction, a 1% reduction to about a 35% reduction, a 1% reduction to about a 30% reduction, a 1% reduction to about a 25% reduction, a 1% reduction to about a 20% reduction, a 1% reduction to about a 15% reduction, a 1% reduction to about a 10% reduction, a 1% to about a 5% reduction, about a 5% to about a 99% reduction, about a 10% to about a 99% reduction, about a 15% to about a 99% reduction, about a 20% to about a 99% reduction, about a 25% to about a 99% reduction, about a 30% to about a 99% reduction, about a 35% to about a 99% reduction, about a 40% to about a 99% reduction, about a 45% to about a 99% reduction, about a 50% to about a 99% reduction, about a 55% to about a 99% reduction, about a 60% to about a 99% reduction, about a 65% to about a 99% reduction, about a 70% to about a 99% reduction, about a 75% to about a 95% reduction, about a 80% to about a 99% reduction, about a 90% reduction to about a 99% reduction, about a 95% to about a 99% reduction, about a 5% to about a 10% reduction, about a 5% to about a 25% reduction, about a 10% to about a 30% reduction, about a 20% to about a 40% reduction, about a 25% to about a 50% reduction, about a 35% to about a 55% reduction, about a 40% to about a 60% reduction, about a 50% reduction to about a 75%

reduction, about a 60% reduction to about 80% reduction, or about a 65% to about a 85% reduction etc.), this is indicative of responsiveness to the therapy. In some embodiments, the level of circulating tumor DNA is reduced such that it is below the detection limit of the instrument. In some embodiments, the level of circulating tumor DNA in a biological sample obtained from the patient (n) is compared to the sample taken just previous (n-1). If the level of circulating tumor DNA in the n sample is lower than the n-1 sample (e.g., a 1% to about a 99% reduction, a 1% to about a 95% reduction, a 1% to about a 90% reduction, a 1% to about a 85% reduction, a 1% to about a 80% reduction, a 1% to about a 75% reduction, a 1% reduction to about a 70% reduction, a 1% reduction to about a 65% reduction, a 1% reduction to about a 60% reduction, a 1% reduction to about a 55% reduction, a 1% reduction to about a 50% reduction, a 1% reduction to about a 45% reduction, a 1% reduction to about a 40% reduction, a 1% reduction to about a 35% reduction, a 1% reduction to about a 30% reduction, a 1% reduction to about a 25% reduction, a 1% reduction to about a 20% reduction, a 1% reduction to about a 15% reduction, a 1% reduction to about a 10% reduction, a 1% to about a 5% reduction, about a 5% to about a 99% reduction, about a 10% to about a 99% reduction, about a 15% to about a 99% reduction, about a 20% to about a 99% reduction, about a 25% to about a 99% reduction, about a 30% to about a 99% reduction, about a 35% to about a 99% reduction, about a 40% to about a 99% reduction, about a 45% to about a 99% reduction, about a 50% to about a 99% reduction, about a 55% to about a 99% reduction, about a 60% to about a 99% reduction, about a 65% to about a 99% reduction, about a 70% to about a 99% reduction, about a 75% to about a 95% reduction, about a 80% to about a 99% reduction, about a 90% reduction to about a 99% reduction, about a 95% to about a 99% reduction, about a 5% to about a 10% reduction, about a 5% to about a 25% reduction, about a 10% to about a 30% reduction, about a 20% to about a 40% reduction, about a 25% to about a 50% reduction, about a 35% to about a 55% reduction, about a 40% to about a 60% reduction, about a 50% reduction to about a 75% reduction, about a 60% reduction to about 80% reduction, or about a 65% to about a 85% reduction, etc.), this is indicative of responsiveness to the therapy. In some embodiments, the level of circulating tumor DNA is reduced such that it is below the detection limit of the instrument. In the case of responsiveness to therapy, the subject can to be administered one or more doses of the therapy and the circulating tumor DNA can be continued to be monitored.

**[00499]** If the level of circulating tumor DNA in the sample is higher than the baseline (e.g., a 1% to about a 99% increase, a 1% to about a 95% increase, a 1% to about a 90% increase, a 1% to about a 85% increase, a 1% to about a 80% increase, a 1% to about a 75% increase, a 1% increase to about a 70% increase, a 1% increase to about a 65% increase, a 1% increase to about a 60% increase, a 1% increase to about a 55% increase, a 1% increase to about a 50% increase, a 1% increase to about a 45% increase, a 1% increase to about a 40% increase, a 1% increase to about a 35% increase, a 1% increase to about a 30% increase, a 1% increase to about a 25% increase, a 1% increase to about a 20% increase, a 1% increase to about a 15% increase, a 1% increase to about a 10% increase, a 1% to about a 5% increase, about a 5% to about a 99% increase, about a 10% to about a 99% increase, about a 15% to about a 99% increase, about a 20% to about a 99% increase, about a 25% to about a 99% increase, about a 30% to about a 99% increase, about a 35% to about a 99% increase, about a 40% to about a 99% increase, about a 45% to about a 99% increase, about a 50% to about a 99% increase, about a 55% to about a 99% increase, about a 60% to about a 99% increase, about a 65% to about a 99% increase, about a 70% to about a 99% increase, about a 75% to about a 95% increase, about a 80% to about a 99% increase, about a 90% increase to about a 99% increase, about a 95% to about a 99% increase, about a 5% to

about a 10% increase, about a 5% to about a 25% increase, about a 10% to about a 30% increase, about a 20% to about a 40% increase, about a 25% to about a 50% increase, about a 35% to about a 55% increase, about a 40% to about a 60% increase, about a 50% increase to about a 75% increase, about a 60% increase to about 80% increase, or about a 65% to about a 85% increase, etc.), this can be indicative of resistance to the therapy. If the level of circulating tumor DNA in the n sample is higher than the n-1 sample (e.g., a 1% to about a 99% increase, a 1% to about a 95% increase, a 1% to about a 90% increase, a 1% to about a 85% increase, a 1% to about a 80% increase, a 1% to about a 75% increase, a 1% increase to about a 70% increase, a 1% increase to about a 65% increase, a 1% increase to about a 60% increase, a 1% increase to about a 55% increase, a 1% increase to about a 50% increase, a 1% increase to about a 45% increase, a 1% increase to about a 40% increase, a 1% increase to about a 35% increase, a 1% increase to about a 30% increase, a 1% increase to about a 25% increase, a 1% increase to about a 20% increase, a 1% increase to about a 15% increase, a 1% increase to about a 10% increase, a 1% to about a 5% increase, about a 5% to about a 99% increase, about a 10% to about a 99% increase, about a 15% to about a 99% increase, about a 20% to about a 99% increase, about a 25% to about a 99% increase, about a 30% to about a 99% increase, about a 35% to about a 99% increase, about a 40% to about a 99% increase, about a 45% to about a 99% increase, about a 50% to about a 99% increase, about a 55% to about a 99% increase, about a 60% to about a 99% increase, about a 65% to about a 99% increase, about a 70% to about a 99% increase, about a 75% to about a 95% increase, about a 80% to about a 99% increase, about a 90% increase to about a 99% increase, about a 95% to about a 99% increase, about a 5% to about a 10% increase, about a 5% to about a 25% increase, about a 10% to about a 30% increase, about a 20% to about a 40% increase, about a 25% to about a 50% increase, about a 35% to about a 55% increase, about a 40% to about a 60% increase, about a 50% increase to about a 75% increase, about a 60% increase to about 80% increase, or about a 65% to about a 85% increase etc.), this can be indicative of resistance to the therapy. When resistance to therapy is suspected, the subject can undergo one or more of imaging, biopsy, surgery, or other diagnostic tests. In some embodiments, when resistance to the therapy is suspected, the subject can be administered (either as a monotherapy or in combination with the previous therapy) a compound capable of treating a RET inhibitor resistance (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as provided herein). See, for example, Cancer Discov; 7(12); 1368–70 (2017); and Cancer Discov; 7(12); 1394–403 (2017).

**[00500]** In some embodiments provided herein, a protein biomarker can be used to monitor the responsiveness of a patient to a particular therapy (e.g., a first RET inhibitor, a second RET inhibitor, or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof). For example, prior to starting treatment with a therapy as described herein (e.g., a first RET inhibitor, a second RET inhibitor, or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof), a biological sample can be obtained from the subject and the level of a protein biomarker can be determined in the biological sample. This sample can be considered a base-line sample. The subject can then be administered one or more doses of a therapy as described herein (e.g., a first RET inhibitor, a second RET inhibitor, or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof) and the levels of the protein biomarker can be monitored (e.g., after the first dose, second dose, third dose, etc. or after one week, two weeks, three weeks, four weeks, etc.). If the level of the protein biomarker is lower than the baseline sample (e.g., a 1% to about a 99% reduction, a 1% to about a 95% reduction, a 1% to about

a 90% reduction, a 1% to about a 85% reduction, a 1% to about a 80% reduction, a 1% to about a 75% reduction, a 1% reduction to about a 70% reduction, a 1% reduction to about a 65% reduction, a 1% reduction to about a 60% reduction, a 1% reduction to about a 55% reduction, a 1% reduction to about a 50% reduction, a 1% reduction to about a 45% reduction, a 1% reduction to about a 40% reduction, a 1% reduction to about a 35% reduction, a 1% reduction to about a 30% reduction, a 1% reduction to about a 25% reduction, a 1% reduction to about a 20% reduction, a 1% reduction to about a 15% reduction, a 1% reduction to about a 10% reduction, a 1% to about a 5% reduction, about a 5% to about a 99% reduction, about a 10% to about a 99% reduction, about a 15% to about a 99% reduction, about a 20% to about a 99% reduction, about a 25% to about a 99% reduction, about a 30% to about a 99% reduction, about a 35% to about a 99% reduction, about a 40% to about a 99% reduction, about a 45% to about a 99% reduction, about a 50% to about a 99% reduction, about a 55% to about a 99% reduction, about a 60% to about a 99% reduction, about a 65% to about a 99% reduction, about a 70% to about a 99% reduction, about a 75% to about a 95% reduction, about a 80% to about a 99% reduction, about a 90% reduction to about a 99% reduction, about a 95% to about a 99% reduction, about a 5% to about a 10% reduction, about a 5% to about a 25% reduction, about a 10% to about a 30% reduction, about a 20% to about a 40% reduction, about a 25% to about a 50% reduction, about a 35% to about a 55% reduction, about a 40% to about a 60% reduction, about a 50% reduction to about a 75% reduction, about a 60% reduction to about 80% reduction, or about a 65% to about a 85% reduction etc.), this is indicative of responsiveness to the therapy. In some embodiments, the level of the protein biomarker is reduced such that it is below the detection limit of the instrument. In some embodiments, the level of the protein biomarker in a biological sample obtained from the patient (n) is compared to the sample taken just previous (n-1). If the level of the protein biomarker in the n sample is lower than the n-1 sample (e.g., a 1% to about a 99% reduction, a 1% to about a 95% reduction, a 1% to about a 90% reduction, a 1% to about a 85% reduction, a 1% to about a 80% reduction, a 1% to about a 75% reduction, a 1% reduction to about a 70% reduction, a 1% reduction to about a 65% reduction, a 1% reduction to about a 60% reduction, a 1% reduction to about a 55% reduction, a 1% reduction to about a 50% reduction, a 1% reduction to about a 45% reduction, a 1% reduction to about a 40% reduction, a 1% reduction to about a 35% reduction, a 1% reduction to about a 30% reduction, a 1% reduction to about a 25% reduction, a 1% reduction to about a 20% reduction, a 1% reduction to about a 15% reduction, a 1% reduction to about a 10% reduction, a 1% to about a 5% reduction, about a 5% to about a 99% reduction, about a 10% to about a 99% reduction, about a 15% to about a 99% reduction, about a 20% to about a 99% reduction, about a 25% to about a 99% reduction, about a 30% to about a 99% reduction, about a 35% to about a 99% reduction, about a 40% to about a 99% reduction, about a 45% to about a 99% reduction, about a 50% to about a 99% reduction, about a 55% to about a 99% reduction, about a 60% to about a 99% reduction, about a 65% to about a 99% reduction, about a 70% to about a 99% reduction, about a 75% to about a 95% reduction, about a 80% to about a 99% reduction, about a 90% reduction to about a 99% reduction, about a 95% to about a 99% reduction, about a 5% to about a 10% reduction, about a 5% to about a 25% reduction, about a 10% to about a 30% reduction, about a 20% to about a 40% reduction, about a 25% to about a 50% reduction, about a 35% to about a 55% reduction, about a 40% to about a 60% reduction, about a 50% reduction to about a 75% reduction, about a 60% reduction to about 80% reduction, or about a 65% to about a 85% reduction, etc.), this is indicative of responsiveness to the therapy. In some embodiments, the level of the protein biomarker is reduced such that it is below the detection limit of the instrument. In the case of



responsiveness to therapy, the subject can to be administered one or more doses of the therapy and the protein biomarker can be continued to be monitored.

**[00501]** If the level of the protein biomarker in the sample is higher than the baseline (e.g., a 1% to about a 99% increase, a 1% to about a 95% increase, a 1% to about a 90% increase, a 1% to about a 85% increase, a 1% to about a 80% increase, a 1% to about a 75% increase, a 1% increase to about a 70% increase, a 1% increase to about a 65% increase, a 1% increase to about a 60% increase, a 1% increase to about a 55% increase, a 1% increase to about a 50% increase, a 1% increase to about a 45% increase, a 1% increase to about a 40% increase, a 1% increase to about a 35% increase, a 1% increase to about a 30% increase, a 1% increase to about a 25% increase, a 1% increase to about a 20% increase, a 1% increase to about a 15% increase, a 1% increase to about a 10% increase, a 1% to about a 5% increase, about a 5% to about a 99% increase, about a 10% to about a 99% increase, about a 15% to about a 99% increase, about a 20% to about a 99% increase, about a 25% to about a 99% increase, about a 30% to about a 99% increase, about a 35% to about a 99% increase, about a 40% to about a 99% increase, about a 45% to about a 99% increase, about a 50% to about a 99% increase, about a 55% to about a 99% increase, about a 60% to about a 99% increase, about a 65% to about a 99% increase, about a 70% to about a 99% increase, about a 75% to about a 95% increase, about a 80% to about a 99% increase, about a 90% increase to about a 99% increase, about a 95% to about a 99% increase, about a 5% to about a 10% increase, about a 5% to about a 25% increase, about a 10% to about a 30% increase, about a 20% to about a 40% increase, about a 25% to about a 50% increase, about a 35% to about a 55% increase, about a 40% to about a 60% increase, about a 50% increase to about a 75% increase, about a 60% increase to about 80% increase, or about a 65% to about a 85% increase, etc.), this can be indicative of resistance to the therapy. If the level of the protein biomarker in the n sample is higher than the n-1 sample (e.g., a 1% to about a 99% increase, a 1% to about a 95% increase, a 1% to about a 90% increase, a 1% to about a 85% increase, a 1% to about a 80% increase, a 1% to about a 75% increase, a 1% increase to about a 70% increase, a 1% increase to about a 65% increase, a 1% increase to about a 60% increase, a 1% increase to about a 55% increase, a 1% increase to about a 50% increase, a 1% increase to about a 45% increase, a 1% increase to about a 40% increase, a 1% increase to about a 35% increase, a 1% increase to about a 30% increase, a 1% increase to about a 25% increase, a 1% increase to about a 20% increase, a 1% increase to about a 15% increase, a 1% increase to about a 10% increase, a 1% to about a 5% increase, about a 5% to about a 99% increase, about a 10% to about a 99% increase, about a 15% to about a 99% increase, about a 20% to about a 99% increase, about a 25% to about a 99% increase, about a 30% to about a 99% increase, about a 35% to about a 99% increase, about a 40% to about a 99% increase, about a 45% to about a 99% increase, about a 50% to about a 99% increase, about a 55% to about a 99% increase, about a 60% to about a 99% increase, about a 65% to about a 99% increase, about a 70% to about a 99% increase, about a 75% to about a 95% increase, about a 80% to about a 99% increase, about a 90% increase to about a 99% increase, about a 95% to about a 99% increase, about a 5% to about a 10% increase, about a 5% to about a 25% increase, about a 10% to about a 30% increase, about a 20% to about a 40% increase, about a 25% to about a 50% increase, about a 35% to about a 55% increase, about a 40% to about a 60% increase, about a 50% increase to about a 75% increase, about a 60% increase to about 80% increase, or about a 65% to about a 85% increase etc.), this can be indicative of resistance to the therapy. When resistance to therapy is suspected, the subject can undergo one or more of imaging, biopsy, surgery, or other diagnostic tests. In some embodiments, when resistance to the therapy is suspected, the subject can be administered

(either as a monotherapy or in combination with the previous therapy) a compound capable of treating a RET inhibitor resistance (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as provided herein).

**[00502]** In some embodiments, one or more protein biomarkers are monitored. The particular protein biomarkers to be monitored can depend on the type of cancer and can be readily identified by one having ordinary skill in the art. Non-limiting examples of protein biomarkers include: CA 125, carcinoembryonic antigen (CEA), calcitonin, thyroglobulin, adrenocorticotrophic hormone (ACTH), cortisol, CA 19-9, prolactin, hepatocyte growth factor, osteopontin, myeloperoxidase, tissue inhibitor of metalloproteinases 1, angiopoietin-1 (Ang-1), cytokeratin 19 (CK-19), tissue inhibitor of metalloproteinase-1 (TIMP-1), chitinase 3 like-1 (YKL-40), galectin-3 (GAL-3), CYFRA 21-1 (cytokeratins), EPCAM (epithelial cell adhesion molecule), ProGRP (pro-gastrin-releasing peptide), and CEACAM (carcinoembryonic antigen). See, for example, Cohen JD, Li L, Wang Y, et al. Detection and localization of surgically resectable cancers with a multi-analyte blood test. *Science*; Published online 18 January 2018. pii: eaar3247. DOI: 10.1126/science.aar3247; Fawaz M Makki et al. Serum biomarkers of papillary thyroid cancer. *J Otolaryngol Head Neck Surg*. 2013; 42(1): 16; and Tatiana N. Zamay et al. Current and Prospective Protein Biomarkers of Lung Cancer. *Cancers* (Basel). 2017 Nov; 9(11): 155. In some embodiments, the biomarkers include one or more of CEA, calcitonin, thyroglobulin, ACTH, and cortisol. In some embodiments, the cancer is medullary thyroid cancer and the protein biomarkers include CEA and calcitonin. In some embodiments, the cancer is non-medullary thyroid cancer and the protein biomarker include thyroglobulin. In some embodiments, the biomarkers are ACTH and cortisol (e.g., when a patient as Cushing's disease related to their cancer).

**[00503]** Also provided herein are methods of treating a RET-associated cancer in a subject that include (a) administering one or more (e.g., two or more, three or more, four or more, five or more, or ten or more) doses of a first RET kinase inhibitor to a subject identified or diagnosed as having a RET-associated cancer (e.g., any of the types of RET-associated cancers described herein)(e.g., identified or diagnosed as having a RET-associated cancer using any of the exemplary methods described herein or known in the art); (b) after step (a), determining a level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from the subject; (c) administering a therapeutically effective amount of a second RET inhibitor or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to a subject identified as having about the same or an elevated level of circulating tumor DNA as compared to a reference level of circulating tumor DNA (e.g., any of the reference levels of circulating tumor DNA described herein). In some examples of these methods, the reference level of circulating tumor DNA is a level of circulating tumor DNA in a biological sample obtained from the subject prior to step (a). Some embodiments of these methods further include determining the level of circulating tumor DNA in the biological sample obtained from the subject prior to step (a). In some examples of these methods, the reference level of circulating tumor DNA is a threshold level of circulating tumor DNA (e.g., an average level of circulating tumor DNA in a population of subjects having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment, or a level of circulating tumor DNA in a subject having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a

placebo, or not yet receiving therapeutic treatment). In some examples of these methods, the first RET inhibitor is selected from the group of: cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864.

**[00504]** Also provided herein are methods of treating a RET-associated cancer in a subject that include administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, to a subject (i) identified or diagnosed as having a RET-associated cancer (e.g., any of the types of RET-associated cancers described herein) (e.g., identified or diagnosed as having a RET-associated cancer using any of the exemplary methods described herein or known in the art), (ii) previously administered one or more (e.g., two or more, three or more, four or more, five or more, or ten or more) doses of a second RET kinase inhibitor, and (ii) after the prior administration of the one or more doses of the second RET kinase inhibitor, identified as having about the same or an elevated level of circulating tumor DNA as compared to a reference level of circulating tumor DNA (e.g., any of the reference levels of circulating tumor DNA described herein or known in the art). In some embodiments of these methods, the reference level of circulating tumor DNA is a level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, plasma, or serum) obtained from the subject prior to the administration of the one or more doses of the second RET kinase inhibitor. Some embodiments of these methods further include determining the level of circulating tumor DNA in the biological sample obtained from the subject prior to administration of the one or more doses of the second RET kinase inhibitor. In some examples of these methods, the reference level of circulating tumor DNA is a threshold level of circulating tumor DNA (e.g., an average level of circulating tumor DNA in a population of subjects having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment, or a level of circulating tumor DNA in a subject having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment). In some embodiments of these methods, the second RET kinase inhibitor is selected from the group consisting of: cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864.

**[00505]** Also provided herein are methods of treating a RET-associated cancer in a subject that include: (a) administering one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy to a subject identified or diagnosed as having a RET-associated cancer (e.g., any of the types of RET-associated cancer described herein) (e.g., a subject identified or diagnosed as having a RET-associated cancer using any of the methods described herein or known in the art); (b) after step (a), determining a level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from the subject; (c) administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and an additional therapeutic agent or treatment (e.g., any of the additional therapeutic agents or treatments of a RET-associated cancer described herein or known in the art) to a subject identified as having about the same or an elevated level of circulating tumor DNA as compared to a reference level of circulating tumor DNA (e.g., any of the exemplary reference levels of circulating tumor DNA described herein or known in the art). In some embodiments of these methods, the additional therapeutic agent

is a second RET kinase inhibitor (e.g., a RET kinase inhibitor selected from the group of: cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864. In some examples of any of these methods, the additional therapeutic agent or treatment comprises one or more of: radiation therapy, a chemotherapeutic agent (e.g., any of the exemplary chemotherapeutic agents described herein or known in the art), a checkpoint inhibitor (e.g., any of the exemplary checkpoint inhibitors described herein or known in the art), surgery (e.g., at least partial resection of the tumor) and one or more other kinase inhibitors (e.g., any of the exemplary kinase inhibitors described herein or known in the art). In some examples of these methods, the reference level of circulating tumor DNA is a level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from the subject prior to step (a). In some examples of these methods, the reference level of circulating tumor DNA is a threshold level of circulating tumor DNA (e.g., an average level of circulating tumor DNA in a population of subjects having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment, or a level of circulating tumor DNA in a subject having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment).

**[00506]** Also provided herein are methods of treating a RET-associated cancer in a subject that include: administering a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and an additional therapeutic agent or treatment to a subject (i) identified or diagnosed as having a RET-associated cancer (e.g., any of the types of RET-associated cancer described herein) (e.g., a subject identified or diagnosed as having a RET-associated cancer using any of the methods described herein or known in the art), (ii) previously administered one or more doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy, and (ii) after administration of the one or more (e.g., two or more, three or more, four or more, five or more, or ten or more) doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy, identified as having about the same or an elevated level of circulating tumor DNA as compared to a reference level of circulating tumor DNA (e.g., any of the exemplary reference levels of circulating tumor DNA described herein). In some embodiments of these methods, the reference level of circulating tumor DNA is a level of circulating tumor DNA in a biological sample obtained from the subject prior to administration of the one or more (e.g., two or more, three or more, four or more, five or more, or ten or more) doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy. Some embodiments of these methods further include determining the level of circulating tumor DNA in the biological sample obtained from the subject prior to administration of the one or more doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy. In some examples of these methods, the reference level of circulating tumor DNA is a threshold level of circulating tumor DNA (e.g., an average level of circulating tumor DNA in a population of subjects having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment, or a level of circulating tumor DNA in a subject having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment). In some embodiments of this method, the additional therapeutic agent is a second RET

kinase inhibitor (e.g., a second RET kinase inhibitor selected from the group of cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864. In some embodiments of these methods, the additional therapeutic agent or treatment includes one or more of radiation therapy, a chemotherapeutic agent (e.g., any of the exemplary chemotherapeutic agents described herein or known in the art), a checkpoint inhibitor (e.g., any of the exemplary checkpoint inhibitors described herein or known in the art), surgery (e.g., at least partial resection of the tumor), and one or more other kinase inhibitors (e.g., any of the kinase inhibitors described herein or known in the art).

**[00507]** Also provided herein are methods of selecting a treatment for a subject that include: selecting a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, for a subject (i) identified or diagnosed as having a RET-associated cancer (e.g., any of the RET-associated cancers described herein) (e.g., a subject identified or diagnosed as having a RET-associated cancer using any of the methods described herein or known in the art), (ii) previously administered one or more (e.g., two or more, three or more, four or more, five or more, or ten or more) doses of a second RET kinase inhibitor (e.g., any of the RET kinase inhibitors described herein or known in the art), and (ii) after administration of the one or more doses of the second RET kinase inhibitor, identified as having about the same or an elevated level of circulating tumor DNA as compared to a reference level of circulating tumor DNA. In some embodiments of any of these methods, the reference level of circulating tumor DNA is a level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from the subject prior to administration of the one or more doses of the second RET kinase inhibitor. Some embodiments of these methods further include determining the level of circulating tumor DNA in the biological sample obtained from the subject prior to administration of the one or more doses of the second RET kinase inhibitor. In some examples of these methods, the reference level of circulating tumor DNA is a threshold level of circulating tumor DNA (e.g., an average level of circulating tumor DNA in a population of subjects having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment, or a level of circulating tumor DNA in a subject having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment). In some embodiments of any these methods, the second RET kinase inhibitor is selected from the group of cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864.

**[00508]** Also provided herein are methods of selecting a treatment for a subject that include selecting a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and an additional therapeutic agent or treatment for a subject (i) identified or diagnosed as having a RET-associated cancer (e.g., any of the RET-associated cancers described herein or known in the art) (e.g., a subject diagnosed or identified as having a RET-associated cancer using any of the methods described herein or known in the art), (ii) previously administered one or more doses (e.g., two or more, three or more, four or more, five or more, or ten or more) of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy, and (ii) after administration of the one or more doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, identified as having about the same

or an elevated level of circulating tumor DNA as compared to a reference level of circulating tumor DNA. In some embodiments of these methods, the reference level of circulating tumor DNA is a level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from the subject prior to administration of the one or more doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy. Some embodiments further include determining the level of circulating tumor DNA in the biological sample obtained from the subject prior to administration of the one or more doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy. In some examples of these methods, the reference level of circulating tumor DNA is a threshold level of circulating tumor DNA (e.g., an average level of circulating tumor DNA in a population of subjects having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment, or a level of circulating tumor DNA in a subject having a similar RET-associated cancer and having a similar stage of the RET-associated cancer, but receiving a non-effective treatment or a placebo, or not yet receiving therapeutic treatment). In some embodiments of any of these methods, the additional therapeutic agent is a second RET kinase inhibitor (e.g., a second RET kinase inhibitor selected from the group of: cabozantinib, vandetanib, alectinib, apatinib, sitravatinib, sorafenib, lenvatinib, ponatinib, dovitinib, sunitinib, foretinib, BLU667, and BLU6864. In some embodiments of any of the methods described herein, the additional therapeutic agent or treatment includes one or more of radiation therapy, a chemotherapeutic agent (e.g., any of the examples of a chemotherapeutic agent described herein or known in the art), a checkpoint inhibitor (e.g., any of the checkpoint inhibitors described herein or known in the art), surgery (e.g., at least partial resection of the tumor), and one or more other kinase inhibitors (e.g., any of the other kinase inhibitors described herein or known in the art).

**[00509]** Also provided herein are methods of determining the efficacy of a treatment in a subject that include: (a) determining a first level of circulating tumor DNA in a biological sample (e.g., a biological sample including blood, serum, or plasma) obtained from a subject identified or diagnosed as having a RET-associated cancer at a first time point; (b) administering a treatment including one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject, after the first time point and before a second time point; (c) determining a second level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from the subject at the second time point; and (d) identifying that the treatment is effective in a subject determined to have a decreased second level of circulating tumor DNA as compared to the first level of circulating tumor DNA; or identifying the treatment is not effective in a subject determined to have about the same or an elevated second level of circulating tumor DNA as compared to the first level of circulating tumor DNA. In some embodiments of these methods, the first time point and the second time point are about 1 week to about 1 year apart (e.g., about 1 week to about 10 months, about 1 week to about 8 months, about 1 week to about 6 months, about 1 week to about 4 months, about 1 week to about 3 months, about 1 week to about 2 months, about 1 week to about 1 month, or about 1 week to about 2 weeks).

**[00510]** Also provided herein are methods of determining whether a subject has developed resistance to a treatment that include: (a) determining a first level of circulating tumor DNA in a biological sample (e.g., a biological sample comprising blood, serum, or plasma) obtained from a subject

identified or diagnosed as having a RET-associated cancer at a first time point; (b) administering a treatment including one or more (e.g., two or more, three or more, four or more, five or more, or ten or more) doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject, after the first time point and before a second time point; (c) determining a second level of circulating tumor DNA in a biological sample obtained from the subject at the second time point; and (d) determining that a subject having a decreased second level of circulating tumor DNA as compared to the first level of circulating tumor DNA has not developed resistance to the treatment; or determining that a subject having about the same or an elevated second level of circulating tumor DNA as compared to the first level of circulating tumor DNA has developed resistance to the treatment. In some embodiments of these methods, the first time point and the second time point are about 1 week to about 1 year apart (e.g., about 1 week to about 10 months, about 1 week to about 8 months, about 1 week to about 6 months, about 1 week to about 4 months, about 1 week to about 3 months, about 1 week to about 2 months, about 1 week to about 1 month, or about 1 week to about 2 weeks).

**[00511]** Exemplary methods for detecting circulating tumor DNA are described in Moati et al., *Clin. Res. Hepatol. Gastroenterol.* April 4, 2018; Oussalah et al., *EBioMedicine* March 28, 2018; Moon et al., *Adv. Drug Deliv. Rev.* April 4, 2018; Solassaol et al., *Clin. Chem. Lab. Med.* April 7, 2018; Arriola et al., *Clin. Transl. Oncol.* April 5, 2018; Song et al., *J. Circ. Biomark.* March 25, 2018; Aslibekyan et al., *JAMA Cardiol.* April 4, 2018; Isbell et al., *J. Thorac. Cardiovasc. Surg.* March 13, 2018; Boeckx et al., *Clin. Colorectal Cancer* February 22, 2018; Anunobi et al., *J. Surg. Res.* March 28, 2018; Tan et al., *Medicine* 97(13):e0197, 2018; Reithdorf et al., *Transl. Androl. Urol.* 6(6):1090-1110, 2017; Volckmar et al., *Genes Chromosomes Cancer* 57(3):123-139, 2018; and Lu et al., *Chronic Dis. Transl. Med.* 2(4):223-230, 2016. Additional methods for detecting circulating tumor DNA are known in the art.

**[00512]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a multikinase inhibitor, wherein the multikinase inhibitor is selected from vandetanib or cabozantinib; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the multikinase inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00513]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a first multikinase inhibitor, wherein the multikinase inhibitor is selected from the group consisting of: vandetanib or cabozantinib; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at

least one RET inhibitor resistance mutation; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the multikinase inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00514]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a multikinase inhibitor, wherein the multikinase inhibitor is selected from the group consisting of: vandetanib or cabozantinib; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the multikinase inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00515]** In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a multikinase inhibitor, wherein the multikinase inhibitor is selected from the group consisting of vandetanib or cabozantinib; or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the multikinase inhibitor of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00516]** As another example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a multikinase inhibitor (e.g., vandetanib or cabozantinib, as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the



subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering a multikinase inhibitor (e.g., vandetanib or cabozantinib), as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering a multikinase inhibitor (e.g., vandetanib or cabozantinib), as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering a multikinase inhibitor (e.g., vandetanib or cabozantinib) as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00517]** Also, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering additional doses of the

compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation) if the subject has a cancer cell that has at least one RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation) if the subject has a cancer cell that has at least one RET inhibitor resistance mutation. In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 151 or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation of Tables 3 or 4; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation) if the subject has a cancer cell that has at least one RET inhibitor resistance mutation. In some embodiments, a second RET inhibitor selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668 is administered in step (d). In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has the RET inhibitor resistance mutation V804M, G810S, or G810R; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of

Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation) if the subject has a cancer cell that has at least one RET inhibitor resistance mutation. In some embodiments, a second RET inhibitor selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668 is administered in step (d).

**[00518]** Also, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) detecting at least one RET inhibitor resistance mutation in a cancer cell in a sample obtained from the subject; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation). In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) detecting at least one RET inhibitor resistance mutation in a cancer cell in a sample obtained from the subject; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation). In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions/deletions of Tables 2 and 2a in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151 or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) detecting at least one RET inhibitor resistance mutation of Tables 3 or 4 in a cancer cell in a sample obtained from the subject; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation). In some embodiments, a second RET inhibitor selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-

methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668 is administered in step (d). In some embodiments, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting the fusion protein KIF5B-RET in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I selected from Examples 1-151, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprise (after (b)) (c) detecting the RET inhibitor resistance mutation V804M, G810S, or G810R in a cancer cell in a sample obtained from the subject; and (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject as a monotherapy or in conjunction with another anticancer agent (e.g., a second RET inhibitor, a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy) or anticancer therapy (e.g., surgery or radiation). In some embodiments, a second RET inhibitor selected from the group consisting of alectinib, cabozantinib, lenvatinib, nintedanib, ponatinib, regorfenib, sorafenib, sunitinib, vandetanib, RXDX-105 (agerafenib), BLU-667 ((1S,4R)-N-((S)-1-(6-(4-fluoro-1H-pyrazol-1-yl)pyridin-3-yl)ethyl)-1-methoxy-4-(4-methyl-6-((5-methyl-1H-pyrazol-3-yl)amino)pyrimidin-2-yl)cyclohexane-1-carboxamide), BLU6864, DS-5010, GSK3179106, GSK3352589, and NMS-E668 is administered in step (d).

**[00519]** Further provided herein is a method for treating lung cancer in a patient in need thereof, the method comprising administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, crizotinib, osimertinib, or any combination thereof.

**[00520]** In some embodiments, the lung cancer is a RET-associated cancer. For example, the method can include: (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the methods further comprises (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation (e.g., a MET dysregulation such as a MET gene amplification); and (d) administering a second therapeutic agent, wherein the second therapeutic agent is crizotinib, as a monotherapy or in conjunction with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some such embodiments, the method comprises (a) detecting one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions of Table 2 in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In further embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation (e.g., a MET dysregulation such as a MET gene amplification); and (d) administering a second therapeutic agent, wherein the second therapeutic agent is crizotinib, as a monotherapy or in conjunction with a

compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (e) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (b) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation.

**[00521]** In some embodiments, the lung cancer is an EGFR-associated cancer. For example, the method can include: (a) detecting a dysregulation of an EGFR gene, an EGFR kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of an EGFR inhibitor (e.g., osimertinib). In some embodiments, the methods further comprises (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has at least one dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same (e.g., a RET gene fusion); and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with the EGFR inhibitor (e.g., osimertinib) to the subject if the subject has a cancer cell that has at least one dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same (e.g., a RET gene fusion); or (e) administering additional doses of the EGFR inhibitor (e.g., osimertinib) of step (b) to the subject if the subject has a cancer cell that does not have a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same (e.g., a RET gene fusion). In some such embodiments, the method comprises (a) detecting a dysregulation of an EGFR gene, an EGFR kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of osimertinib. In further embodiments, the methods further comprise (after (b)) (c) determining whether a cancer cell in a sample obtained from the subject has one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions of Table 2; and (d) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, as a monotherapy or in conjunction with osimertinib to the subject if the subject has a cancer cell that has one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions of Table 2; or (e) administering additional doses of the osimertinib of step (b) to the subject if the subject has a cancer cell that does not have one or more fusion proteins of Table 1 and/or one or more RET kinase protein point mutations/insertions of Table 2.

**[00522]** The term “EGFR-associated cancer” as used herein refers to cancers associated with or having a dysregulation of a EGFR gene, a EGFR kinase, or expression or activity, or level of any of the same.

**[00523]** The phrase “dysregulation of a EGFR gene, a EGFR kinase, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a EGFR gene translocation that results in the expression of a fusion protein, a deletion in a EGFR gene that results in the expression of a EGFR protein that includes a deletion of at least one amino acid as compared to the wild-type EGFR protein, or a mutation in a EGFR gene that results in the expression of a EGFR protein with one or more point mutations, or an alternative spliced version of a EGFR mRNA that results in a EGFR protein that results in the deletion of at least one amino acid in the EGFR protein as compared to the wild-type EGFR protein), or a EGFR gene amplification that results in overexpression of a EGFR protein or an autocrine activity resulting from the overexpression of a EGFR gene a cell, that results in a pathogenic increase in the

activity of a kinase domain of a EGFR protein (*e.g.*, a constitutively active kinase domain of a EGFR protein) in a cell. As another example, a dysregulation of a EGFR gene, a EGFR protein, or expression or activity, or level of any of the same, can be a mutation in a EGFR gene that encodes a EGFR protein that is constitutively active or has increased activity as compared to a protein encoded by a EGFR gene that does not include the mutation. For example, a dysregulation of a EGFR gene, a EGFR protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of EGFR that includes a functional kinase domain, and a second portion of a partner protein (*i.e.*, that is not EGFR). In some examples, dysregulation of a EGFR gene, a EGFR protein, or expression or activity, can be a result of a gene translocation of one EGFR gene with another non-EGFR gene.

**[00524]** The term “wildtype EGFR” or “wild-type EGFR” describes a nucleic acid (*e.g.*, a EGFR gene or a EGFR mRNA) or protein (*e.g.*, a EGFR protein) that is found in a subject that does not have a EGFR-associated cancer (and optionally also does not have an increased risk of developing a EGFR-associated cancer and/or is not suspected of having a EGFR-associated cancer), or is found in a cell or tissue from a subject that does not have a EGFR-associated cancer (and optionally also does not have an increased risk of developing a EGFR-associated cancer and/or is not suspected of having a EGFR-associated cancer).

**[00525]** Also provided are methods of selecting a treatment for a subject having a cancer that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and selecting a treatment that includes administration of a compound of Formula I (*e.g.*, any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a first RET inhibitor. In some embodiments, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof is administered in combination with the first RET inhibitor. Also provided are methods of selecting a treatment for a subject having a cancer that include: selecting a treatment that includes administration of a compound of Formula I (*e.g.*, any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof for a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations. Also provided are methods of selecting a subject having a cancer for a treatment that does not include a first RET inhibitor as a monotherapy that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and selecting the identified subject for a treatment that includes a compound of Formula I (*e.g.*, any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Also provided are methods of selecting a subject having a cancer for a treatment that does not include a first RET inhibitor as a monotherapy that include: selecting a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations for a treatment that includes administration of a compound of Formula I (*e.g.*, any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. In some embodiments, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, *e.g.*, V804M, V804L, or V804E, or a substitution amino acid position 810, *e.g.*, G810S, G810R, G810C, G810A, G810V, and G810D.

**[00526]** Also provided are methods of determining the likelihood that a subject having a cancer (*e.g.*, a RET-associated cancer) will have a positive response to treatment with a first RET inhibitor as a

monotherapy that include: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that a subject having a cancer cell that has one or more RET inhibitor resistance mutations has a decreased likelihood of having a positive response (i.e. an increased likelihood of having a negative response) to treatment with a first RET inhibitor as a monotherapy. Also provided are methods of determining the likelihood that a subject having a cancer (e.g., a RET-associated cancer) will have a positive response to treatment with a first RET inhibitor as a monotherapy that include: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that a subject not having a cancer cell that has one or more RET inhibitor resistance mutations has an increased likelihood of having a positive response to treatment with a first RET inhibitor as a monotherapy as compared to a subject having a cancer cell that has one or more RET inhibitor resistance mutations. Also provided are methods of predicting the efficacy of treatment with a first RET inhibitor as a monotherapy in a subject having cancer that include: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that treatment with a first RET inhibitor as a monotherapy is less likely to be effective in a subject having a cancer cell in a sample obtained from the subject that has one or more RET inhibitor resistance mutations. Also provided are methods of predicting the efficacy of treatment with a first RET inhibitor as a monotherapy in a subject having cancer that include: determining that treatment with a first RET inhibitor as a monotherapy is less likely to be effective in a subject having a cancer cell in a sample obtained from the subject that has one or more RET inhibitor resistance mutations. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D.

**[00527]** Also provided are methods of treating a subject having a cancer that include: (a) administering one or more doses of a first RET inhibitor to the subject for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (c) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (d) administering additional doses of the first RET inhibitor of step (a) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where the subject is administered additional doses of the first RET inhibitor of step (a), the subject can also be administered another anticancer agent (e.g., a second RET inhibitor or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy). In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments of step (c), another RET inhibitor can be the first RET inhibitor administered in step (a). In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some

embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D.

**[00528]** Also provided are methods of treating a subject having a cancer that include: (a) administering one or more doses of a first RET inhibitor to the subject for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (c) administering a second RET inhibitor as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (d) administering additional doses of the first RET inhibitor step (a) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where the subject is administered additional doses of the first RET inhibitor of step (a), the subject can also be administered another anticancer agent. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof). In some embodiments, the additional anticancer agent is an immunotherapy.

**[00529]** Also provided are methods of treating a subject having a cancer (e.g., a RET-associated cancer) that include: (a) determining whether a cancer cell in a sample obtained from a subject having a cancer and previously administered one or more doses of a first RET inhibitor, has one or more RET inhibitor resistance mutations; and (b) administering a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (c) administering additional doses of the first RET inhibitor previously administered to the subject if the subject has cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where the subject is administered additional doses of the first RET inhibitor previously administered to the subject, the subject can also be administered another anticancer agent (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or immunotherapy). In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an



immunotherapy. In some embodiments of step (b), another anticancer agent can be the first RET inhibitor administered in step (a).

**[00530]** Also provided are methods of treating a subject having a cancer that include: (a) determining whether a cancer cell in a sample obtained from a subject having a cancer and previously administered one or more doses of a first RET inhibitor has one or more RET inhibitor resistance mutations; and (b) administering a second RET inhibitor as a monotherapy or in conjunction with another anticancer agent to the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (c) administering additional doses of the first RET inhibitor previously administered to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where the subject is administered additional doses of the first RET inhibitor previously administered to the subject, the subject can also be administered another anticancer agent. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments of (b), another anticancer agent can be the first RET inhibitor administered in step (a).

**[00531]** In some embodiments, a RET-associated cancer as described herein can occur in a subject along with a dysregulation of another gene, another protein, or the expression or activity or level of any of the same.

**[00532]** For example, a RET-associated cancer that exhibits a RET fusion can occur in a subject along with one or more of: a dysregulation of a MET gene, a MET protein, or the expression or activity or level of any of the same; a dysregulation of a PIK3CA gene, a PIK3CA protein, or the expression or activity or level of any of the same; a dysregulation of a KRAS gene, a KRAS protein, or the expression or activity or level of any of the same; a dysregulation of a EGFR gene, a EGFR protein, or the expression or activity or level of any of the same (e.g., an amplification of a EGFR gene); a dysregulation of a FGFR2 gene, a FGFR2 protein, or the expression or activity or level of any of the same (eg., a fusion of an FGFR2 gene or an FGFR2 protein); a dysregulation of a CDK4 gene, a CDK4 protein, or the expression or activity or level of any of the same (e.g., an amplification of a CDK4 gene); a dysregulation of a mTOR gene, a mTOR protein, or the expression or activity or level of any of the same; a dysregulation of a CDKN2A gene, a CDKN2A protein, or the expression or activity or level of any of the same (e.g., a deletion in a CDKN2A gene or a CDKN2A protein); a dysregulation of a CDKN2B gene, a CDKN2B protein, or the expression or activity or level of any of the same (e.g., a deletion in a CDKN2B gene or a CDKN2B protein); a dysregulation of a NF1 gene, a NF1 protein, or the expression or activity or level of any of the same; a dysregulation of a MYC gene, a MYC protein, or the expression or activity or level of any of the same (e.g., an amplification in a MYC gene); a dysregulation of a MDM2 gene, a MDM2 protein, or the expression or activity or level of any of the same (e.g., an amplification in a MDM2 gene); a

dysregulation of a GNAS gene, a GNAS protein, or the expression or activity or level of any of the same; a dysregulation of a BRCA2 gene, a BRCA2 protein, or the expression or activity or level of any of the same.

**[00533]** In some embodiments, a RET-associated cancer that exhibits a mutation of a RET gene and/or a RET protein can occur in a subject along with one or more of: a dysregulation of a PIK3CA gene, a PIK3CA protein, or the expression or activity or level of any of the same; a dysregulation of a KRAS gene, a KRAS protein, or the expression or activity or level of any of the same; a dysregulation of a EGFR gene, a EGFR protein, or the expression or activity or level of any of the same; a dysregulation of a FGFR1 gene, a FGFR1 protein, or the expression or activity or level of any of the same (e.g., an amplification of a FGFR1 gene); a dysregulation of a FGFR2 gene, a FGFR2 protein, or the expression or activity or level of any of the same (e.g., an amplification of a FGFR2 gene); a dysregulation of a FGFR3 gene, a FGFR3 protein, or the expression or activity or level of any of the same (e.g., a fusion of a FGFR3 gene or a FGFR3 protein); a dysregulation of a ERBB2 gene, a ERBB2 protein, or the expression or activity or level of any of the same (e.g., an amplification of ERBB2 gene); and a dysregulation of a KIT gene, a KIT protein, or the expression or activity or level of any of the same.

**[00534]** In some embodiments, a RET-associated cancer that exhibits an amplification of a RET gene can occur in a patient along with one or more additional kinase amplifications. For example, an amplification in a FGFR1 gene; an amplification in a FGFR2 gene; an amplification in a FGFR3 gene; an amplification of a FGFR4 gene; an amplification of a CDK4 gene; and an amplification in a CDK6 gene.

**[00535]** In some embodiments, wherein a RET-associated cancer as described herein can occur in a subject along with a dysregulation in another kinase, the methods described herein can further comprise administration of an additional therapeutic agent that targets and/or treats the dysregulation in the other kinase. For example, provided herein are methods for treating a RET-associated cancer in a subject in need of such treatment, the method comprising (a) detecting a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of any of the same in a sample from the subject; and (b) administering to the subject a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the method further comprises (c) detecting a dysregulation in another kinase in a sample from the subject; and (d) administering to the subject a therapeutic agent that targets and/or treats the dysregulation in the other kinase. In some embodiments, the administration of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof is done concurrently, sequentially, or serially. In some embodiments, the detecting steps (a) and (c) can be done simultaneously or sequentially in any order.

**[00536]** Additional therapeutic agents that target and/or treat the dysregulation of the other kinase can include any known inhibitor of the other kinase. Examples of such agents are as follows:

**[00537]** Exemplary PARP inhibitors include: 3-aminobenzamide (INO-1001), 5-aminoisoquinoline, ABT472, ABT767, AG140361, AG14032, ANG2864, ANG3186, AZD2281, AZD2461, BGP-15, BSI101, BSI401, CEP6800, CEP8983, CK102, CEP9722 (prodrug of CEP8983), CPH101 with CPH102, DR2313, E7016 (GPI-21016), E7449, GP16150, IMP4297, IMP04149, INO1002, INO1003, JPI283, JPI289, KU0687, KU58948, niraparib (MK-4827), NT125, olaparib (AZD2281), ONO-1924H, ONO2231, pamiparib (BGB-

290), PJ-34, rucaparib (AG014699), SC10914, SOMCL9112, talazoparib (BMN-673), and veliparib (ABT-888).

**[00538]** Exemplary CDK 4/6 inhibitors include: palbociclib (PD0332991), abemaciclib (LY2835219), ribociclib (LEE011), trilaciclib (G1T28), voruciclib, and G1T38.

**[00539]** Exemplary ERBB2 (HER2/neu) inhibitors include: afatinib, afatinib, dacomitinib (PF-00299804), DS8201-a, erlontinib, gefitinib, KU004, lapatinib, lapatinib ditosylate, MM-111, mubritinib (TAK-165), neratinib, pyrotinib (HTI-1001), tucatinib (ONT-380, ARRY-380), 7C3, cetuximab, HER2-BsAb, hersintuzumab, margetuximab, MI130004, NeuVax, paitumumab, pertuzumab, SYD985, trastuzumab, and trastuzumab emtansine.

**[00540]** Exemplary inhibitors of amplified ERBB2 (HER2/neu) include dacomitinib (PF-00299804), lapatinib, neratinib, pertuzumab, trastuzumab, and trastuzumab emtansine.

**[00541]** Exemplary EGFR inhibitors include: AC0010, afatinib, AP26113, ASP8273, avatinib, avitinib, AZD3759, BMS-690514, brigatinib, canertinib, Cap-701, CHMFL-EGFR-202, CUDC-101, dacomitinib, EAI045, EGF816, erlontinib, erlotinib, gefitinib, GNS-1481, GNS-1486, Gö6976, HS-10296, icotinib, KU004, lapatinib, nazartinib, neratinib, olmutinib (HM61713, BI 1482694), osimertinib, osimertinib (AZD9291), pelitinib, PF-06747775, PKC412, pyrotinib (HTI-1001), rocilentinib, vandetanib, varlitinib, XL647, 7C3, cetuximab, depatuxizumab mafodotin (ABT-414), matuzumab, nimotuzumab, panitumumab, and zalutumumab.

**[00542]** Exemplary wild-type EGFR inhibitors include: afatinib, BMS-690514, canertinib, CUDC-101, dacomitinib, erlotinib, gefitinib, lapatinib, neratinib, pelitinib, vandetanib, varlitinib, XL647, cetuximab, matuzumab, nimotuzumab, panitumumab, and zalutumumab.

**[00543]** Exemplary inhibitors of mutated EGFR include: AC0010, afatinib, AP26113, ASP8273, avatinib, avitinib, AZD3759, BMS-690514, brigatinib, canertinib, Cap-701, CHMFL-EGFR-202, CUDC-101, dacomitinib, EAI045, EGF816, GNS-1481, GNS-1486, Gö6976, HS-10296, icotinib, nazartinib, neratinib, olmutinib (HM61713, BI 1482694), osimertinib (AZD9291), PF-06747775, PKC412, rocilentinib, vandetanib, varlitinib, and cetuximab.

**[00544]** An exemplary inhibitor of amplified EGFR is depatuxizumab mafodotin (ABT-414).

**[00545]** Exemplary inhibitors of FGFR include: ASP5878, AZD4547, BGJ398, BLU9931, brivatinib, cediranib, DEBIO 1347, derazantinib (ARQ-087), dovitinib (CHIR258), E7090, ENMD-2076, erdafitinib (JNJ-42756293), FGF 401, FIIN-1, FRIN-1, INCB054828, L16H50, lenvatinib, lucitanib, LY2874455, nintedanib, NP603, orantinib (SU6668), pazopanib, PBI05204, PD173074, ponatinib, PRN1371, regorafenib, rogaratinib (BAY-1163877), S49076, SOMCL-085, SU5402, sunitinib, TAS-120, FP-1039, GAL-F2, GAL-FR21, GAL-FR22, GAL-FR23, GP369, hLD1.vb, LD1, MFGR1877S, MM-161, PRO-001, and R3Mab.

**[00546]** Exemplary inhibitors of FGFR fusions include: BGJ398, DEBIO 1347, derazantinib (ARQ-087), E7090, erdafitinib (JNJ-42756293), lucitanib, and TAS-120.

**[00547]** Exemplary inhibitors of FGFR1, FGFR2, and FGFR3 include: AZD4547, BGJ398, DEBIO 1347, E7090, INCB054828, S49076, SOMCL-085, and TAS-120.

**[00548]** Exemplary inhibitors of FGF4 include: BLU-554, BLU9931, NVP-FGF401, and hLD1.vb.

**[00549]** Exemplary inhibitors of amplified FGFR1 include: AZD4547, BGJ398, DEBIO 1347, derazantinib (ARQ-087), erdafitinib (JNJ-42756293), INCB054828, and lucitanib.

**[00550]** Exemplary inhibitors of amplified FGFR2 include: AZD4547, DEBIO 1347, derazantinib (ARQ-087), lucitanib, regorafenib, and TAS-120.

**[00551]** An exemplary inhibitor of amplified FGFR3 is AZD4547.

**[00552]** Exemplary MEK inhibitors include: AZD8330 (ARRY-424704), AZD6244 (ARRY-142866), BI-847325, binimetinib, BIX02188, BIX02189, CH4987655, CH5126766, CI-1040, cobimetinib (GDC-0973), EBI-1051, G-573, G8935, GDC-0623, Myricetin, nobiletin, PD0325901, PD184161, PD318088, PD98059, PD334581, pimasertib (AS-703026), refametinib (RDEA119, BAY 869766), selumetinib (AZD6244), SL-327, TAK-733, trametinib, and U0126.

**[00553]** Exemplary KRAS inhibitors include: 0375-0604, a covalent quinazoline-based switch II pocket (SIIP) compound, ARS-1620, AZD4785, and LP1.

**[00554]** Exemplary PI3K inhibitors include: 3-methyladenine, A66, alpelisib (BYL719), AMG319, apitolisib (GDC-0980, RG7422), AS-252424, AS-604850, AS-605240, AZD6842, AZD8186, AZD8835, BGT226 (NVP-BGT226), buparlisib (BKM120), CAY10505, CH5132799, copanlisib (BAY 80-6946), CUDC-907, CZC24832, dactolisib (BEZ235, NVP-BEZ235), DS7423, duvelisib (IPI-145, INK1197), GDC-0032, GDC-0084, GDC-0326, gedatolisib (PF-05212384, PKI-5587), GNE-317, GS-9820, GSK1059615, GSK2292767, GSK2636771, HS-173, IC-87114, Idelalisib (CAL-101, GS-1101), IPI-145, IPI-3063, IPI-549, LY294002, LY3023414, nemiralisib (GSK2269557), omipalisib (GSK2126458, GSK458), PF-04691502, PF-4989216, PI-103, PI-3065, pictilisib (GDC-0941), PIK-293, PIK-294, PIK-75, PIK-90, PIK-93, PIK-III, pilaralisib (XL147), PKI-587, PP-110, PQR309, PQR309, PW-12, PX-866, quercetin, S14161, SAR245409 (XL765), SAR260301, SAR405, serabelisib (INK-1117, MLN-1117, TAK-1117), SF-1126, SF-2523, SN32976, taselisib (GDC-0032), TB101110, TG100-115, TG100-713, TGR-1202, TGX-221, umbralisib (TGR-1202), voxalisib (XL765, SAR245409), VPS34-IN1, VS-5584 (SB2343), WJD008, wortmannin, and ZSTK474.

**[00555]** Exemplary KIT inhibitors include: AMG 706, amuvatinib (MP-470), APck110, axitinib (AG-013736), AZD2932, dasatinib (BMS-354825), dovitinib (TKI-258, CHIR-258), EXEL-0862, imatinib, KI-328, masitinib (AB1010), midostaurin, MLN518, motesanib, N3-(6-aminopyridin-3-yl)-N1-(2-cyclopentylethyl)-4-methylisophthalamide, nilotinib, OSI-930, pazopanib (GW786034), pexidartinib (PLX3397), PKC412, PLX647, PP1, quizartinib (AC220), regorafenib (BAY 73-4506), semaxinib (SU 5416), sitravatinib (MGCD516), sorafenib, STI571, SU11248, SU9529, sunitinib, telatinib, tivozanib (AV-951), tyrphostin AG 1296, VX-322, and WBZ\_4.

**[00556]** Exemplary MDM2 inhibitors include: (-)-parthenolide, ALRN6924, AM-8553, AMG232, CGM-097, DS-3032b, GEM240, HDM201, HLI98, idasanutlin (RG-7338), JapA, MI-219, MI-219, MI-319, MI-77301 (SAR405838), MK4828, MK-8242, MX69, NSC 207895 (XI-006), Nutlin-3, Nutlin-3a, Nutlin-3b, NVP-CFC218, NVP-CGM097, PXn727/822, RG7112, RO2468, RO5353, RO5503781, serdemetan (JNJ-26854165), SP-141, and YH239-EE.

**[00557]** Exemplary inhibitors of amplified MDM2 include: AM-8553, AMG232, DS-3032b, MI-77301 (SAR405838), NSC 207895 (XI-006), Nutlin-3a, NVP-CFC218, NVP-CGM097, and RG7112.

**[00558]** Exemplary inhibitors of MET include: (–)-Oleocanthol, ABBV-399, AMG-208, AMG-337, AMG-458, BAY-853474, BMS-754807, BMS-777607, BMS-794833, cabozantinib (XL184, BMS-907351), capmatinib (INCB28060), crizotinib (PF-02341066), DE605, foretinib (GSK1363089, XL880), glesatinib (MGCD265), golvatinib (E7050), INCB028060, JNJ-38877605, KRC-408, merestinib (LY2801653), MK-2461, MK8033, NPS-1034, NVP-BVU972, PF-04217903, PHA-665752, S49076, savolitinib (AZD6094, HMPL-504), SGX-523, SU11274, TAS-115, tepotinib (EMD 1214063, MSC2156119J), volitinib, CE-355621, and Onartuzumab.

**[00559]** Exemplary inhibitors of mTOR include: anthracimycin, apitolisib (GDC-0980, RG7422), AZD-8055, BGT226 (NVP-BGT226), CC-223, CZ415, dactolisib (BEZ235, NVP-BEZ235), DS7423, everolimus (RAD001), GDC-0084, GDC-0349, gedatolisib (PF-05212384, PKI-5587), GSK1059615, INK128, KU-0063794, LY3023414, MLN0128, omipalisib (GSK2126458, GSK458), OSI-027, OSU-53, Palomid 529 (P529), PF-04691502, PI-103, PKI-587, PP242, PQR309, ridafarolimus (AP-23573), sapanisertib (INK 128, MLN0128), SAR245409 (XL765), SF-1126, SF2523, sirolimus (rapamycin), SN32976, TAK228, temsirolimus (CCI-779, NSC 683864), Torin 1, Torin 2, torkinib (PP242), umirolimus, vistusertib (AZD2014), voxtalisib (XL765, SAR245409), VS-5584, VS-5584 (SB2343), WAY-600, WYE-125132 (WYE-132), WYE-354, WYE-687, XL388, and zotarolimus (ABT-578).

**[00560]** Exemplary inhibitors of MYC include: 10058-F4, 10074-G5, and KSI-3716.

**[00561]** The phrase “dysregulation of a gene, a protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a kinase domain and a fusion partner, a mutation in a gene that results in the expression of a protein that includes a deletion of at least one amino acid as compared to a wildtype protein, a mutation in a gene that results in the expression of a protein with one or more point mutations as compared to a wildtype protein, a mutation in a gene that results in the expression of a protein with at least one inserted amino acid as compared to a wildtype protein, a gene duplication that results in an increased level of protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of protein in a cell), an alternative spliced version of a mRNA that results in a protein having a deletion of at least one amino acid in the protein as compared to the wild-type protein), or increased expression (e.g., increased levels) of a wildtype protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a gene, a protein, or expression or activity, or level of any of the same, can be a mutation in a gene that encodes a protein that is constitutively active or has increased activity as compared to a protein encoded by a gene that does not include the mutation. For example, a dysregulation of a gene, a protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not the primary protein). In some examples, dysregulation of a gene, a protein, or expression or activity or level of any of the same can be a result of a gene translocation of one gene with a different gene.

**[00562]** Treatment of a patient having a cancer with a multi-kinase inhibitor (MKI) or target-specific kinase inhibitor (e.g., a BRAF inhibitor, a EGFR inhibitor, a MEK inhibitor, an ALK inhibitor, a ROS1 inhibitor, a MET inhibitor, an aromatase inhibitor, a RAF inhibitor, or a RAS inhibitor) can result in dysregulation of a RET gene, a RET kinase, or the expression or activity or level of the same in the cancer, and/or resistance to a RET inhibitor. See, e.g., Bhinge et al., *Oncotarget* 8:27155-27165, 2017; Chang et al., *Yonsei Med. J.* 58:9-18, 2017; and Lopez-Delisle et al., doi: 10.1038/s41388-017-0039-5, *Oncogene* 2018.

**[00563]** Treatment of a patient having a cancer with a RET inhibitor in combination with a multi-kinase inhibitor or a target-specific kinase inhibitor (e.g., a BRAF inhibitor, a EGFR inhibitor, a MEK inhibitor, an ALK inhibitor, a ROS1 inhibitor, a MET inhibitor, an aromatase inhibitor, a RAF inhibitor, or a RAS inhibitor) can have increased therapeutic efficacy as compared to treatment of the same patient or a similar patient with the RET inhibitor as a monotherapy, or the multi-kinase inhibitor or the target-specific kinase inhibitor as a monotherapy. See, e.g., Tang et al., doi: 10.1038/modpathol.2017.109, *Mod. Pathol.* 2017; Andreucci et al., *Oncotarget* 7:80543-80553, 2017; Nelson-Taylor et al., *Mol. Cancer Ther.* 16:1623-1633, 2017; and Kato et al., *Clin. Cancer Res.* 23:1988-1997, 2017.

**[00564]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) and previously administered a multi-kinase inhibitor (MKI) or a target-specific kinase inhibitor (e.g., a BRAF inhibitor, a EGFR inhibitor, a MEK inhibitor, an ALK inhibitor, a ROS1 inhibitor, a MET inhibitor, an aromatase inhibitor, a RAF inhibitor, or a RAS inhibitor) (e.g., as a monotherapy) that include: administering to the patient (i) a therapeutically effective dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy, or (ii) a therapeutically effective dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and a therapeutically effective dose of the previously administered MKI or the previously administered target-specific kinase inhibitor.

**[00565]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) previously administered a MKI or a target specific kinase inhibitor (e.g., a BRAF inhibitor, a EGFR inhibitor, a MEK inhibitor, an ALK inhibitor, a ROS1 inhibitor, a MET inhibitor, an aromatase inhibitor, a RAF inhibitor, or a RAS inhibitor) (e.g., as a monotherapy) that include: identifying a patient having a cancer cell that has a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy, or (ii) a therapeutically effective dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and a therapeutically effective dose of the previously administered MKI or the previously administered target-specific kinase inhibitor.

**[00566]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: administering to a patient a therapeutically effective amount of a MKI or a target-specific kinase inhibitor (e.g., a BRAF inhibitor, a EGFR inhibitor, a MEK inhibitor, an ALK inhibitor, a ROS1 inhibitor, a MET inhibitor, an aromatase inhibitor, a RAF inhibitor, or a RAS inhibitor) (e.g., as a monotherapy) for a first period of time; after the period of time, identifying a patient having a

cancer cell that has a dysregulation of a RET gene, a RET kinase, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy, or (ii) a therapeutically effective dose of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and a therapeutically effective dose of the previously administered MKI or the previously administered target-specific kinase inhibitor.

**[00567]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of a BRAF gene, a BRAF kinase, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a BRAF inhibitor (e.g., any of the BRAF inhibitors described herein or known in the art).

**[00568]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of a BRAF gene, a BRAF kinase, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a BRAF inhibitor (e.g., any of the BRAF inhibitors described herein or known in the art).

**[00569]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of an EGFR gene, an EGFR protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of an EGFR inhibitor (e.g., any of the EGFR inhibitors described herein or known in the art).

**[00570]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of an EGFR gene, an EGFR protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of an EGFR inhibitor (e.g., any of the EGFR inhibitors described herein or known in the art).

**[00571]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of a MEK gene, a MEK protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a MEK inhibitor (e.g., any of the MEK inhibitors described herein or known in the art).

**[00572]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of a MEK gene, a MEK protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a MEK inhibitor (e.g., any of the MEK inhibitors described herein or known in the art).

**[00573]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of an ALK gene, an ALK protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of an ALK inhibitor (e.g., any of the ALK inhibitors described herein or known in the art).

**[00574]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of an ALK gene, an ALK protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount an ALK inhibitor (e.g., any of the ALK inhibitors described herein or known in the art).

**[00575]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of a ROS gene, a ROS protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a ROS inhibitor (e.g., any of the ROS inhibitors described herein or known in the art).

**[00576]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of a ROS gene, a ROS protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount a ROS inhibitor (e.g., any of the ROS inhibitors described herein or known in the art).

**[00577]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of a MET gene, a MET protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a MET inhibitor (e.g., any of the MET inhibitors described herein or known in the art).

**[00578]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of a MET gene, a MET protein, or the expression or activity or level of the same; and administering to



the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount a MET inhibitor (e.g., any of the MET inhibitors described herein or known in the art).

**[00579]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of an aromatase gene, an aromatase protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of an aromatase inhibitor (e.g., any of the aromatase inhibitors described herein or known in the art).

**[00580]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of an aromatase gene, an aromatase protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount an aromatase inhibitor (e.g., any of the aromatase inhibitors described herein or known in the art).

**[00581]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of a RAF gene, a RAF protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a RAF inhibitor (e.g., any of the RAF inhibitors described herein or known in the art).

**[00582]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of a RAF gene, a RAF protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount a RAF inhibitor (e.g., any of the RAF inhibitors described herein or known in the art).

**[00583]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that has dysregulation of a RAS gene, a RAS protein, or the expression or activity or level of the same that include administering to the patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount of a RAS inhibitor (e.g., any of the RAS inhibitors described herein or known in the art).

**[00584]** Provided herein are methods of treating a patient having a cancer (e.g., any of the cancers described herein) that include: identifying a patient having a cancer cell that has dysregulation of a RAS gene, a RAS protein, or the expression or activity or level of the same; and administering to the identified patient (i) a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and (ii) a therapeutically effective amount a RAS inhibitor (e.g., any of the RAS inhibitors described herein or known in the art).

**[00585]** The phrase “dysregulation of a BRAF gene, a BRAF protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a BRAF kinase domain and a fusion partner, a mutation in a BRAF gene that results in the expression of a BRAF protein that includes a deletion of at least one amino acid as compared to a wildtype BRAF protein, a mutation in a BRAF gene that results in the expression of a BRAF protein with one or more point mutations as compared to a wildtype BRAF protein, a mutation in a BRAF gene that results in the expression of a BRAF protein with at least one inserted amino acid as compared to a wildtype BRAF protein, a gene duplication that results in an increased level of BRAF protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of BRAF protein in a cell), an alternative spliced version of a BRAF mRNA that results in a BRAF protein having a deletion of at least one amino acid in the BRAF protein as compared to the wild-type BRAF protein), or increased expression (e.g., increased levels) of a wildtype BRAF protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a BRAF gene, a BRAF protein, or expression or activity, or level of any of the same, can be a mutation in a BRAF gene that encodes a BRAF protein that is constitutively active or has increased activity as compared to a protein encoded by a BRAF gene that does not include the mutation. For example, a dysregulation of a BRAF gene, a BRAF protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a BRAF protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not BRAF). In some examples, dysregulation of a BRAF gene, a BRAF protein, or expression or activity or level of any of the same can be a result of a gene translocation of one BRAF gene with another non-BRAF gene.

**[00586]** Non-limiting examples of a BRAF inhibitor include dabrafenib, vemurafenib (also called RG7204 or PLX4032), sorafenib tosylate, PLX-4720, GDC-0879, BMS-908662 (Bristol-Meyers Squibb), LGX818 (Novartis), PLX3603 (Hofmann-LaRoche), RAF265 (Novartis), RO5185426 (Hofmann-LaRoche), and GSK2118436 (GlaxoSmithKline). Additional examples of a BRAF inhibitor are known in the art.

**[00587]** The phrase “dysregulation of an EGFR gene, an EGFR protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including an EGFR kinase domain and a fusion partner, a mutation in an EGFR gene that results in the expression of an EGFR protein that includes a deletion of at least one amino acid as compared to a wildtype EGFR protein, a mutation in an EGFR gene that results in the expression of an EGFR protein with one or more point mutations as compared to a wildtype EGFR protein, a mutation in an EGFR gene that results in the expression of an EGFR protein with at least one inserted amino acid as compared to a wildtype EGFR protein, a gene duplication that results in an increased level of EGFR protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of EGFR protein in a cell), an alternative spliced version of a EGFR mRNA that results in an EGFR protein having a deletion of at least one amino acid in the EGFR protein as compared to the wild-type EGFR protein), or increased expression (e.g., increased levels) of a wildtype EGFR protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of an EGFR gene, an EGFR protein, or expression or activity, or level of any of the same,

can be a mutation in an EGFR gene that encodes an EGFR protein that is constitutively active or has increased activity as compared to a protein encoded by an EGFR gene that does not include the mutation. For example, a dysregulation of an EGFR gene, an EGFR protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a EGFR protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not EGFR). In some examples, dysregulation of an EGFR gene, an EGFR protein, or expression or activity or level of any of the same can be a result of a gene translocation of one EGFR gene with another non-EGFR gene.

**[00588]** Non-limiting examples of an EGFR inhibitor include gefitinib, erlotinib, brigatinib, lapatinib, neratinib, icotinib, afatinib, dacomitinib, poziotinib, vandetanib, afatinib, AZD9291, CO-1686, HM61713, AP26113, CI-1033, PKI-166, GW-2016, EKB-569, PDI-168393, AG-1478, CGP-59326A. Additional examples of an EGFR inhibitor are known in the art.

**[00589]** The phrase “dysregulation of a MEK gene, a MEK protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a MEK kinase domain and a fusion partner, a mutation in a MEK gene that results in the expression of a MEK protein that includes a deletion of at least one amino acid as compared to a wildtype MEK protein, a mutation in a MEK gene that results in the expression of a MEK protein with one or more point mutations as compared to a wildtype MEK protein, a mutation in a MEK gene that results in the expression of a MEK protein with at least one inserted amino acid as compared to a wildtype MEK protein, a gene duplication that results in an increased level of MEK protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of MEK protein in a cell), an alternative spliced version of a MEK mRNA that results in a MEK protein having a deletion of at least one amino acid in the MEK protein as compared to the wild-type MEK protein), or increased expression (e.g., increased levels) of a wildtype MEK protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a MEK gene, a MEK protein, or expression or activity, or level of any of the same, can be a mutation in a MEK gene that encodes a MEK protein that is constitutively active or has increased activity as compared to a protein encoded by a MEK gene that does not include the mutation. For example, a dysregulation of a MEK gene, a MEK protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a MEK protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not MEK). In some examples, dysregulation of a MEK gene, a MEK protein, or expression or activity or level of any of the same can be a result of a gene translocation of one MEK gene with another non-MEK gene.

**[00590]** Non-limiting examples of a MEK inhibitor include mekinist, trametinib (GSK1120212), cobimetinib (XL518), binimetinib (MEK162), selumetinib, PD-325901, CI-1040, PD035901, TAK-733, PD098059, U0126, AS703026/MSK1935369, XL-518/GDC-0973, BAY869766/RDEA119, and GSK1120212. Additional examples of a MEK inhibitor are known in the art.

**[00591]** The phrase “dysregulation of an ALK gene, an ALK protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results

in the expression of a fusion protein including an ALK kinase domain and a fusion partner, a mutation in an ALK gene that results in the expression an ALK protein that includes a deletion of at least one amino acid as compared to a wildtype ALK protein, a mutation in an ALK gene that results in the expression of an ALK protein with one or more point mutations as compared to a wildtype ALK protein, a mutation in an ALK gene that results in the expression of an ALK protein with at least one inserted amino acid as compared to a wildtype ALK protein, a gene duplication that results in an increased level of ALK protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of ALK protein in a cell), an alternative spliced version of an ALK mRNA that results in an ALK protein having a deletion of at least one amino acid in the ALK protein as compared to the wild-type ALK protein), or increased expression (e.g., increased levels) of a wildtype ALK protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of an ALK gene, an ALK protein, or expression or activity, or level of any of the same, can be a mutation in an ALK gene that encodes an ALK protein that is constitutively active or has increased activity as compared to a protein encoded by an ALK gene that does not include the mutation. For example, a dysregulation of an ALK gene, an ALK protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of an ALK protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not ALK). In some examples, dysregulation of an ALK gene, an ALK protein, or expression or activity or level of any of the same can be a result of a gene translocation of one ALK gene with another non-ALK gene.

**[00592]** Non-limiting examples of an ALK inhibitor include crizotinib (Xalkori), ceritinib (Zykadia), alectinib (Alecensa), dalantercept, ACE-041 (Brigatinib) (AP26113), entrectinib (NMS-E628), PF-06463922 (Pfizer), TSR-011 (Tesar), CEP-37440 (Teva), CEP-37440 (Teva), X-396 (Xcovery), and ASP-3026 (Astellas). Additional examples of an ALK inhibitor are known in the art.

**[00593]** The phrase “dysregulation of a ROS1 gene, a ROS1 protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a ROS1 kinase domain and a fusion partner, a mutation in a ROS1 gene that results in the expression a ROS1 protein that includes a deletion of at least one amino acid as compared to a wildtype ROS1 protein, a mutation in a ROS1 gene that results in the expression of a ROS1 protein with one or more point mutations as compared to a wildtype ROS1 protein, a mutation in a ROS1 gene that results in the expression of a ROS1 protein with at least one inserted amino acid as compared to a wildtype ROS1 protein, a gene duplication that results in an increased level of ROS1 protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of ROS1 protein in a cell), an alternative spliced version of a ROS1 mRNA that results in a ROS1 protein having a deletion of at least one amino acid in the ROS1 protein as compared to the wild-type ROS1 protein), or increased expression (e.g., increased levels) of a wildtype ROS1 protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a ROS1 gene, a ROS1 protein, or expression or activity, or level of any of the same, can be a mutation in a ROS1 gene that encodes a ROS1 protein that is constitutively active or has increased activity as compared to a protein encoded by a ROS1 gene that does not include the mutation. For example, a

dysregulation of a ROS1 gene, a ROS1 protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a ROS1 protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not ROS1). In some examples, dysregulation of a ROS1 gene, a ROS1 protein, or expression or activity or level of any of the same can be a result of a gene translocation of one ROS1 gene with another non-ROS1 gene.

**[00594]** Non-limiting examples of a ROS1 inhibitor include crizotinib, entrectinib (RXDX-101), lorlatinib (PF-06463922), certinib, TPX-0005, DS-605, and cabozantinib. Additional examples of a ROS1 inhibitor are known in the art.

**[00595]** The phrase “dysregulation of a MET gene, a MET protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a MET kinase domain and a fusion partner, a mutation in a MET gene that results in the expression a MET protein that includes a deletion of at least one amino acid as compared to a wildtype MET protein, a mutation in a MET gene that results in the expression of a MET protein with one or more point mutations as compared to a wildtype MET protein, a mutation in a MET gene that results in the expression of a MET protein with at least one inserted amino acid as compared to a wildtype MET protein, a gene duplication that results in an increased level of MET protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of MET protein in a cell), an alternative spliced version of a MET mRNA that results in a MET protein having a deletion of at least one amino acid in the MET protein as compared to the wild-type MET protein), or increased expression (e.g., increased levels) of a wildtype MET protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a MET gene, a MET protein, or expression or activity, or level of any of the same, can be a mutation in a MET gene that encodes a MET protein that is constitutively active or has increased activity as compared to a protein encoded by a MET gene that does not include the mutation. For example, a dysregulation of a MET gene, a MET protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a MET protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not MET). In some examples, dysregulation of a MET gene, a MET protein, or expression or activity or level of any of the same can be a result of a gene translocation of one MET gene with another non-MET gene.

**[00596]** Non-limiting examples of a MET inhibitor include crizotinib, cabozantinib, JNJ-38877605, PF-04217903 (Pfizer), MK-2461, GSK 1363089, AMG 458 (Amgen), tivantinib, INCB28060 (Incyte), PF-02341066 (Pfizer), E7050 (Eisai), BMS-777607 (Bristol-Meyers Squibb), JNJ-38877605 (Johnson & Johnson), ARQ197 (ArQule), GSK/1363089/XL880 (GSK/Exelixis), and XL174 (BMS/Exelixis). Additional examples of a MET inhibitor are known in the art.

**[00597]** The phrase “dysregulation of an aromatase gene, an aromatase protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a mutation in an aromatase gene that results in the expression an aromatase protein that includes a deletion of at least one amino acid as compared to a wildtype aromatase protein, a mutation in an aromatase gene that

results in the expression of an aromatase protein with one or more point mutations as compared to a wildtype aromatase protein, a mutation in an aromatase gene that results in the expression of an aromatase protein with at least one inserted amino acid as compared to a wildtype aromatase protein, a gene duplication that results in an increased level of aromatase protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of aromatase protein in a cell), an alternative spliced version of an aromatase mRNA that results in an aromatase protein having a deletion of at least one amino acid in the aromatase protein as compared to the wild-type aromatase protein), or increased expression (e.g., increased levels) of a wildtype aromatase in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of an aromatase gene, an aromatase protein, or expression or activity, or level of any of the same, can be a mutation in an aromatase gene that encodes an aromatase protein that is constitutively active or has increased activity as compared to a protein encoded by an aromatase gene that does not include the mutation.

**[00598]** Non-limiting examples of an aromatase inhibitor include Arimidex (anastrozole), Aromasin (exemestane), Femara (letrozole), Teslac (testolactone), and formestane. Additional examples of an aromatase inhibitor are known in the art.

**[00599]** The phrase “dysregulation of a RAF gene, a RAF protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a RAF kinase domain and a fusion partner, a mutation in a RAF gene that results in the expression a RAF protein that includes a deletion of at least one amino acid as compared to a wildtype RAF protein, a mutation in a RAF gene that results in the expression of a RAF protein with one or more point mutations as compared to a wildtype RAF protein, a mutation in a RAF gene that results in the expression of a RAF protein with at least one inserted amino acid as compared to a wildtype RAF protein, a gene duplication that results in an increased level of RAF protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of RAF protein in a cell), an alternative spliced version of a RAF mRNA that results in a RAF protein having a deletion of at least one amino acid in the RAF protein as compared to the wild-type RAF protein), or increased expression (e.g., increased levels) of a wildtype RAF protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a RAF gene, a RAF protein, or expression or activity, or level of any of the same, can be a mutation in a RAF gene that encodes a RAF protein that is constitutively active or has increased activity as compared to a protein encoded by a RAF gene that does not include the mutation. For example, a dysregulation of a RAF gene, a RAF protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a RAF protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not RAF). In some examples, dysregulation of a RAF gene, a RAF protein, or expression or activity or level of any of the same can be a result of a gene translocation of one RAF gene with another non-RAF gene.

**[00600]** Non-limiting examples of a RAF inhibitor include sorafenib, vemurafenib, dabrafenib, BMS-908662/XL281, GSK2118436, RAF265, RO5126766, and RO4987655. Additional examples of a RAF inhibitor are known in the art.

**[00601]** The phrase “dysregulation of a RAS gene, a RAS protein, or the expression or activity or level of any of the same” refers to a genetic mutation (e.g., a chromosomal translocation that results in the expression of a fusion protein including a RAS kinase domain and a fusion partner, a mutation in a RAS gene that results in the expression a RAS protein that includes a deletion of at least one amino acid as compared to a wildtype RAS protein, a mutation in a RAS gene that results in the expression of a RAS protein with one or more point mutations as compared to a wildtype RAS protein, a mutation in a RAS gene that results in the expression of a RAS protein with at least one inserted amino acid as compared to a wildtype RAS protein, a gene duplication that results in an increased level of RAS protein in a cell, or a mutation in a regulatory sequence (e.g., a promoter and/or enhancer) that results in an increased level of RAS protein in a cell), an alternative spliced version of a RAS mRNA that results in a RAS protein having a deletion of at least one amino acid in the RAS protein as compared to the wild-type RAS protein), or increased expression (e.g., increased levels) of a wildtype RAS protein in a mammalian cell due to aberrant cell signaling and/or dysregulated autocrine/paracrine signaling (e.g., as compared to a control non-cancerous cell). As another example, a dysregulation of a RAS gene, a RAS protein, or expression or activity, or level of any of the same, can be a mutation in a RAS gene that encodes a RAS protein that is constitutively active or has increased activity as compared to a protein encoded by a RAS gene that does not include the mutation. For example, a dysregulation of a RAS gene, a RAS protein, or expression or activity, or level of any of the same, can be the result of a gene or chromosome translocation which results in the expression of a fusion protein that contains a first portion of a RAS protein that includes a functional kinase domain, and a second portion of a partner protein (i.e., that is not RAS). In some examples, dysregulation of a RAS gene, a RAS protein, or expression or activity or level of any of the same can be a result of a gene translocation of one RAS gene with another non-RAS gene.

**[00602]** Non-limiting examples of a RAS inhibitor include Kobe0065 and Kobe2602. Additional examples of a RAS inhibitor are known in the art.

**[00603]** Non-limiting examples of multi-kinase inhibitors (MKIs) include dasatinib and sunitinib.

**[00604]** In some embodiments, provided herein are methods of treating a subject having a cancer that include: (a) administering one or more doses of a first RET inhibitor or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has at least one dysregulation of a gene, a protein, or the expression or activity or level of any of the same, wherein the gene or protein is selected from the group consisting of EGFR, MET, ALK, ROS1, KRAS, BRAF, RAS, PIK3CA, and HER2; and (c) 1) administering a second RET inhibitor as a monotherapy or in conjunction with another anticancer agent, 2) administering additional doses of the first RET inhibitor or a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof in combination with an inhibitor targeting the gene or protein (e.g., an inhibitor of EGFR, MET, ALK, ROS1, KRAS, BRAF, RAS, PIK3CA, and HER2), or 3) stopping administration of the RET inhibitor of step a) and administering an inhibitor targeting the gene or protein (e.g., an inhibitor of EGFR, MET, ALK, ROS1, KRAS, BRAF, RAS, PIK3CA, and HER2) to the subject if the subject has a cancer cell that has at least one dysregulation of a gene, a protein, or the expression or activity or level of the same, wherein the gene or protein is selected from the group

consisting of EGFR, MET, ALK, ROS1, KRAS, BRAF, RAS, PIK3CA, and HER2; or (d) administering additional doses of the first RET inhibitor step (a) to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same, wherein the gene or protein is selected from the group consisting of EGFR, MET, ALK, ROS1, KRAS, BRAF, RAS, PIK3CA, and HER2 confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor or the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the tumor is a NSCLC tumor and the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same are selected from targetable mutations in EGFR or MET, targetable rearrangements involving ALK or ROS1, or activating mutations in KRAS. In some embodiments, the tumor is a thyroid (non-MTC) tumor and the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same are selected from targetable mutations in BRAF or activating mutations in RAS. In some embodiments, the tumor is a MTC tumor and the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same are selected from targetable mutations in ALK or activating mutations in RAS. In some embodiments, the tumor is a pancreatic tumor and the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same is an activating mutations in KRAS. In some embodiments, the tumor is a colorectal tumor and the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same are selected from targetable mutations in BRAF or PIK3CA or an activating mutation in RAS. In some embodiments, the tumor is a breast tumor and the one or more dysregulations of a gene, a protein, or the expression or activity or level of any of the same are selected from targetable mutations in PIK3CA or alteration in HER2.

**[00605]** Also provided are methods of selecting a treatment for a subject having a cancer that include (a) administering one or more doses of a first RET inhibitor to the subject for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (c) selecting a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent for the subject if the subject has a cancer cell that has one or more RET inhibitor resistance mutations; or (d) selecting additional doses of the first RET inhibitor of step (a) for the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, when additional doses of the first RET inhibitor of step (a) are selected for the subject, the method can further include selecting doses of another anticancer agent for the subject. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments of step (c), another RET inhibitor can be the first RET inhibitor administered in step (a).



**[00606]** Also provided are methods of selecting a treatment for a subject having a cancer that include (a) administering one or more doses of a first RET inhibitor to the subject for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has at least one RET inhibitor resistance mutation; and (c) selecting a second RET inhibitor as a monotherapy or in conjunction with another anticancer agent if the subject has a cancer cell that has one or more RET inhibitor resistance mutations; or (d) selecting additional doses of the first RET inhibitor of step (a) for the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, when additional doses of the first RET inhibitor of step (a) are selected for the subject, the method can further include selecting doses of another anticancer agent for the subject. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments, another RET can be the first RET inhibitor administered in step (a).

**[00607]** Also provided are methods of selecting a treatment for a subject having a cancer that include (a) determining whether a cancer cell in a sample obtained from a subject having a cancer and previously administered one or more doses of a first RET inhibitor has one or more RET inhibitor resistance mutations; (b) selecting a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent for the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (c) selecting additional doses of the first RET inhibitor previously administered to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, when additional doses of the first RET inhibitor previously administered to the subject are selected for the subject, the method can further include selecting doses of another anticancer agent (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof or immunotherapy) for the subject. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments of step (c), another RET inhibitor can be the first RET inhibitor administered in step (a).

**[00608]** Also provided are methods of selecting a treatment for a subject having a cancer that include (a) determining whether a cancer cell in a sample obtained from a subject having a cancer and previously administered one or more doses of a first RET inhibitor has one or more RET inhibitor resistance mutations; (b) selecting a second RET inhibitor as a monotherapy or in conjunction with another anticancer agent for the subject if the subject has a cancer cell that has at least one RET inhibitor resistance mutation; or (c) selecting additional doses of the first RET inhibitor previously administered to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, when additional doses of the first RET inhibitor previously administered to the subject are selected for the subject, the method can further include selecting doses of another anticancer agent (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or an immunotherapy) for the subject. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments, another RET can be the first RET inhibitor administered in step (a).

**[00609]** Also provided are methods of determining a subject's risk for developing a cancer that has some resistance to a first RET inhibitor that include: determining whether a cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and identifying a subject having a cell that has one or more RET inhibitor resistance mutations, as having an increased likelihood of developing a cancer that has some resistance to the first RET inhibitor. Also provided are methods of determining a subject's risk for developing a cancer that has some resistance to a first RET inhibitor that include: identifying a subject having a cell that has one or more RET inhibitor resistance mutations, as having an increased likelihood of developing a cancer that has some resistance to the first RET inhibitor. Also provided are methods of determining the presence of a cancer that has some resistance to a first RET inhibitor that include: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that the subject having a cancer cell that has one or more RET inhibitor resistance mutations has a cancer that has some resistance to the first RET inhibitor. Also provided are methods of determining the presence of a cancer that has some resistance to a first RET inhibitor in a subject that include: determining that a subject having a cancer cell that has one or more RET inhibitor resistance mutations, has a cancer that has some resistance to the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with the first RET inhibitor. In some embodiments, the one or more RET inhibitor resistance mutations include one or more RET inhibitor resistance mutations listed in Tables 3 and 4. For example, the one or more RET inhibitor resistance mutations can include a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D.

**[00610]** In some embodiments of any of the methods described herein, a RET inhibitor resistance mutation that confers increased resistance to a cancer cell or tumor to treatment with a first RET inhibitor can be any of the RET inhibitor resistance mutations listed in Table 3 or 4 (e.g., a substitution at amino acid position 804, e.g., V804M, V804L, or V804E, or a substitution at amino acid position 810, e.g., G810S, G810R, G810C, G810A, G810V, and G810D).

**[00611]** In some embodiments, the presence of one or more RET inhibitor resistance mutations in a tumor causes the tumor to be more resistant to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Methods useful when a RET inhibitor resistance mutation causes the tumor to be more resistant to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof are described below. For example, provided herein are methods of treating a subject having a cancer that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and administering to the identified subject a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy (e.g., a second RET kinase inhibitor). Also provided are methods of treating a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations that include administering to the subject a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy (e.g., a second RET kinase inhibitor). In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00612]** Also provided are methods of selecting a treatment for a subject having a cancer that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and selecting a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy for the identified subject (e.g., a second RET kinase inhibitor). Also provided are methods of selecting a treatment for a subject having a cancer that include: selecting a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy (e.g., a second RET kinase inhibitor) for a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations. Also provided are methods of selecting a subject having a cancer for a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy (e.g., a second RET kinase inhibitor) that include: identifying a subject having a cancer cell that has one or more RET inhibitor resistance mutations; and selecting the identified subject for a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy (e.g., a second RET kinase inhibitor). Also provided are methods of selecting a subject having a cancer for a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy (e.g., a second RET kinase inhibitor) that include: selecting a subject identified as having a cancer cell that has one or more RET inhibitor resistance mutations for a treatment that does not include a compound of Formula I (e.g., any one of Formulas I-A

to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00613]** Also provided are methods of determining the likelihood that a subject having a cancer will have a positive response to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy that include: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that the subject having the cancer cell that has one or more RET inhibitor resistance mutations has a decreased likelihood of having a positive response to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy. Also provided are methods of determining the likelihood that a subject having cancer will have a positive response to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy that include: determining that a subject having a cancer cell that has one or more RET inhibitor resistance mutations has a decreased likelihood of having a positive response to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy. Also provided are methods of predicting the efficacy of treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy in a subject having cancer that include: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy is less likely to be effective in a subject having a cancer cell in a sample obtained from the subject that has one or more RET inhibitor resistance mutations. Also provided are methods of predicting the efficacy of treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy in a subject having cancer that include: determining that treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy is less likely to be effective in a subject having a cancer cell in a sample obtained from the subject that has one or more RET inhibitor resistance mutations. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00614]** Also provided are methods of treating a subject having a cancer that include: (a) administering one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and (c) administering a second RET inhibitor or a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to a subject having a cancer cell that has one or more RET inhibitor resistance mutations; or (d) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (a) to a subject having a cancer cell that does

not have a RET inhibitor resistance mutation. In some embodiments, where the subject is administered additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (a), the subject can also be administered another anticancer agent or a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments, another RET can be the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof administered in step (a).

**[00615]** Also provided are methods of treating a subject having a cancer that include: (a) determining whether a cancer cell in a sample obtained from a subject having a cancer and previously administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, has one or more RET inhibitor resistance mutations; (b) administering a second RET inhibitor or a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent to a subject having a cancer cell that has one or more RET inhibitor resistance mutations; or (c) administering additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof previously administered to a subject having a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where the subject is administered additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (a), the subject can also be administered another anticancer agent. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments, another RET can be the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof administered in step (a).

**[00616]** Also provided are methods of selecting a treatment for a subject having a cancer that include: (a) administering one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to the subject for a period of time; (b) after (a), determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and (c) selecting a second RET inhibitor or a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent for the subject if the subject has a cancer cell that has a RET inhibitor resistance mutation; or (d) selecting additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (a) for the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where additional doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof of step (a) are selected for the subject, the

method can also include further selecting another anticancer agent. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments, another RET can be the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof administered in step (a).

**[00617]** Also provided are methods of selecting a treatment for a subject having a cancer that include: (a) determining whether a cancer cell in a sample obtained from a subject having a cancer and previously administered one or more doses of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, has one or more RET inhibitor resistance mutations; (b) selecting a second RET inhibitor or a second compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as a monotherapy or in conjunction with another anticancer agent for the subject if the subject has a cancer cell that has a RET inhibitor resistance mutation; or (c) selecting additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof previously administered to the subject if the subject has a cancer cell that does not have a RET inhibitor resistance mutation. In some embodiments, where additional doses of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof of step (a) are selected for the subject, the method can also include further selecting another anticancer agent. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the additional anticancer agent is any anticancer agent known in the art. For example, the additional anticancer agent is another RET inhibitor (e.g., a second RET inhibitor). In some embodiments, the additional anticancer agent is an immunotherapy. In some embodiments, another RET can be the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof administered in step (a).

**[00618]** Also provided are methods of determining a subject's risk for developing a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof that include: determining whether a cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and identifying the subject if the subject has a cell that has one or more RET inhibitor resistance mutations as having an increased likelihood of developing a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Also provided are methods of determining a subject's risk for developing a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof that include: identifying a subject having a cell that has one or more RET inhibitor resistance mutations as having an increased likelihood of developing a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Also provided are methods of determining the presence of a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or

solvate thereof that includes: determining whether a cancer cell in a sample obtained from the subject has one or more RET inhibitor resistance mutations; and determining that the subject having the cancer cell that has one or more RET inhibitor resistance mutations has a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. Also provided are methods of determining the presence of a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof in a subject that include: determining that a subject having a cancer cell that has one or more RET inhibitor resistance mutations has a cancer that has some resistance to a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the one or more RET inhibitor resistance mutations confer increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00619]** In some embodiments of any of the methods described herein, a RET inhibitor resistance mutation that confers increased resistance to a cancer cell or tumor to treatment with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, can be any of the RET inhibitor resistance mutations listed in Table 3 or 4.

**[00620]** Methods of determining the level of resistance of a cancer cell or a tumor to a RET inhibitor (e.g., any of the RET inhibitors described herein or known in the art) can be determined using methods known in the art. For example, the level of resistance of a cancer cell to a RET inhibitor can be assessed by determining the  $IC_{50}$  of a RET inhibitor (e.g., any of the RET inhibitors described herein or known in the art) on the viability of a cancer cell. In other examples, the level of resistance of a cancer cell to a RET inhibitor can be assessed by determining the growth rate of the cancer cell in the presence of a RET inhibitor (e.g., any of the RET inhibitors described herein). In other examples, the level of resistance of a tumor to a RET inhibitor can be assessed by determining the mass or size of one or more tumors in a subject over time during treatment with a RET inhibitor (e.g., any of the RET inhibitors described herein). In other examples, the level of resistance of a cancer cell or a tumor to a RET inhibitor can be indirectly assessed by determining the activity of a RET kinase including one or more of the RET inhibitor resistance mutations (i.e., the same RET kinase expressed in a cancer cell or a tumor in a subject). The level of resistance of a cancer cell or tumor having one or more RET inhibitor resistance mutations to a RET inhibitor is relative to the level of resistance in a cancer cell or tumor that does not have a RET inhibitor resistance mutation (e.g., a cancer cell or tumor that does not have the same RET inhibitor resistance mutations, a cancer cell or a tumor that does not have any RET inhibitor resistance mutations, or a cancer cell or a tumor that expresses a wildtype RET protein). For example, the determined level of resistance of a cancer cell or a tumor having one or more RET inhibitor resistance mutations can be greater than about 1%, greater than about 2%, greater than about 3%, greater than about 4%, greater than about 5%, greater than about 6%, greater than about 7%, greater than about 8%, greater than about 9%, greater than about 10%, greater than about 11%, greater than about 12%, greater than about 13%, greater than about 14%, greater than about 15%, greater than about 20%, greater than about 25%, greater than about 30%, greater than about 35%, greater than about 40%, greater than about 45%, greater than about 50%, greater than about 60%, greater than about 70%, greater than about 80%, greater than about 90%, greater than about 100%, greater than about 110%, greater than about 120%, greater than about 130%, greater than about 140%, greater than about 150%,

greater than about 160%, greater than about 170%, greater than about 180%, greater than about 190%, greater than about 200%, greater than about 210%, greater than about 220%, greater than about 230%, greater than about 240%, greater than about 250%, greater than about 260%, greater than about 270%, greater than about 280%, greater than about 290%, or greater than about 300% of the level of resistance in a cancer cell or tumor that does not have a RET inhibitor resistance mutation (e.g., a cancer cell or tumor that does not have the same RET inhibitor resistance mutations, a cancer cell or a tumor that does not have any RET inhibitor resistance mutations, or a cancer cell or a tumor that expresses a wildtype RET protein).

**[00621]** RET is thought to play an important role in the development and survival of afferent nociceptors in the skin and gut. RET kinase knock-out mice lack enteric neurons and have other nervous system anomalies suggesting that a functional RET kinase protein product is necessary during development (Taraviras, S. et al., *Development*, 1999, 126:2785-2797). Moreover population studies of patients with Hirschsprung's disease characterized by colonic obstruction due to lack of normal colonic enervation have a higher proportion of both familial and sporadic loss of function RET mutations (Butler Tjaden N., et al., *Transl. Res.*, 2013, 162: 1-15). Irritable bowel syndrome (IBS) is a common illness affecting 10-20% of individuals in developed countries and is characterized by abnormal bowel habits, bloating and visceral hypersensitivity (Camilleri, M., *N. Engl. J. Med.*, 2012, 367: 1626-1635). While the etiology of IBS is unknown it is thought to result from either a disorder between the brain and gastrointestinal tract, a disturbance in the gut microbiome or increased inflammation. The resulting gastrointestinal changes affect normal bowel transit resulting in either diarrhea or constipation. Furthermore in many IBS patients the sensitization of the peripheral nervous system results in visceral hypersensitivity or allodynia (Keszthelyi, D., *Eur. J. Pain*, 2012, 16: 1444-1454). See, e.g., U.S. Publication No. 2015/0099762.

**[00622]** Accordingly, provided herein are methods for treating a patient diagnosed with (or identified as having) an irritable bowel syndrome (IBS) including diarrhea-predominant, constipation- predominant or alternating stool pattern, functional bloating, functional constipation, functional diarrhea, unspecified functional bowel disorder, functional abdominal pain syndrome, chronic idiopathic constipation, functional esophageal disorders, functional gastroduodenal disorders, functional anorectal pain, and inflammatory bowel disease that include administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00623]** Also provided herein are methods for treating a patient identified or diagnosed as having a RET-associated irritable bowel syndrome (IBS) (e.g., a patient that has been identified or diagnosed as having a RET-associated irritable bowel syndrome (IBS) through the use of a regulatory agency-approved, e.g., FDA-approved, kit for identifying dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, in a patient or a biopsy sample from the patient) that include administering to the patient a therapeutically



effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00624]** Also provided herein are methods for treating pain associated with IBS that include administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof is administered in combination with another therapeutic agent useful for treating one or more symptoms of IBS.

**[00625]** Also provided are methods for treating an irritable bowel syndrome (IBS) in a patient in need thereof, the method comprising: (a) determining if the irritable bowel syndrome (IBS) in the patient is a RET-associated IBS (e.g., using a regulatory-agency approved, e.g., FDA-approved, kit for identifying dysregulation of a RET gene, a RET kinase, or expression or activity or level of any of the same, in a patient or a biopsy sample from the patient, or by performing any of the non-limiting examples of assays described herein); and (b) if the IBS is determined to be a RET-associated IBS, administering to the patient a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof.

**[00626]** In some embodiments, the compounds of the present invention are useful for treating irritable bowel syndrome (IBS) in combination with one or more additional therapeutic agents or therapies effective in treating the irritable bowel syndrome that work by the same or a different mechanism of action. The at least one additional therapeutic agent may be administered with a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof as part of the same or separate dosage forms, via the same or different routes of administration, and on the same or different administration schedules according to standard pharmaceutical practice known to one skilled in the art.

**[00627]** Non-limiting examples of additional therapeutics for the treatment of irritable bowel syndrome (IBS) include probiotics, fiber supplements (e.g., psyllium, methylcellulose), anti-diarrheal medications (e.g., loperamide), bile acid binders (e.g., cholestyramine, colestipol, colesevelam), anticholinergic and antispasmodic medications (e.g., hyoscyamine, dicyclomine), antidepressant medications (e.g., tricyclic antidepressant such as imipramine or nortriptyline or a selective serotonin reuptake inhibitor (SSRI) such as fluoxetine or paroxetine), antibiotics (e.g., rifaximin), alosetron, and lubiprostone.

**[00628]** Accordingly, also provided herein are methods of treating irritable bowel syndrome (IBS), comprising administering to a patient in need thereof a pharmaceutical combination for treating IBS which comprises (a) a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, (b) an additional therapeutic agent, and (c) optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use for the treatment of IBS, wherein the amounts of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof and the additional therapeutic agent are together effective in treating the IBS. In one

embodiment, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, and the additional therapeutic agent are administered simultaneously as separate dosages. In one embodiment, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, and the additional therapeutic agent are administered as separate dosages sequentially in any order, in jointly therapeutically effective amounts, e.g. in daily or intermittently dosages. In one embodiment, compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, and the additional therapeutic agent are administered simultaneously as a combined dosage.

**[00629]** Also provided herein is (i) a pharmaceutical combination for treating irritable bowel syndrome in a patient in need thereof, which comprises (a) a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, (b) at least one additional therapeutic agent (e.g., any of the exemplary additional therapeutic agents described herein for treating irritable bowel syndrome or known in the art), and (c) optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use for the treatment of irritable bowel syndrome, wherein the amounts of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof and of the additional therapeutic agent are together effective in treating the irritable bowel syndrome; (ii) a pharmaceutical composition comprising such a combination; (iii) the use of such a combination for the preparation of a medicament for the treatment of irritable bowel syndrome; and (iv) a commercial package or product comprising such a combination as a combined preparation for simultaneous, separate or sequential use; and to a method of treatment of irritable bowel syndrome in a patient in need thereof. In one embodiment the patient is a human.

**[00630]** The term "pharmaceutical combination", as used herein, refers to a pharmaceutical therapy resulting from the mixing or combining of more than one active ingredient and includes both fixed and non-fixed combinations of the active ingredients. The term "fixed combination" means that a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and at least one additional therapeutic agent (e.g., an agent effective in treating irritable bowel syndrome), are both administered to a patient simultaneously in the form of a single composition or dosage. The term "non-fixed combination" means that a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof and at least one additional therapeutic agent (e.g., an agent effective in treating irritable bowel syndrome) are formulated as separate compositions or dosages, such that they may be administered to a patient in need thereof simultaneously, concurrently or sequentially with variable intervening time limits, wherein such administration provides effective levels of the two or more compounds in the body of the patient. In one embodiment, the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof and the additional therapeutic agent are formulated as separate unit dosage forms, wherein the separate dosage forms are suitable for either sequential or simultaneous administration. These also apply to cocktail therapies, e.g. the administration of three or more active ingredients.

**[00631]** In some embodiments, a compound provided herein can be used as an agent for supportive care for a patient undergoing cancer treatment. For example, a compound of

Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, can be useful to reduce one or more symptoms associated with treatment with one or more cancer therapies such as diarrhea or constipations complications and/or abdominal pain. See, for example, U.S. Publication No. 2015/0099762 and Hoffman, J.M. et al. *Gastroenterology* (2012) 142:844-854. Accordingly, a compound, or a pharmaceutically acceptable salt thereof, or composition provided herein can be administered to a patient to address one or more complications associated with cancer treatment (e.g., gastrointestinal complications such as diarrhea, constipation, or abdominal pain).

**[00632]** In some embodiments, a therapeutically effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, can be administered to a patient undergoing cancer treatment (e.g., a patient experiencing an adverse event associated with cancer treatment such as an immune-related adverse event or a gastrointestinal complication including diarrhea, constipation, and abdominal pain). For example, a compound provided herein, or a pharmaceutically acceptable salt thereof, can be used in the treatment of colitis or IBS associated with administration of a checkpoint inhibitor; see, e.g., Postow, M.A. et al. *Journal of Clinical Oncology* (2015) 33: 1974-1982. In some such embodiments, a compound provided herein, or a pharmaceutically acceptable salt thereof, can be formulated to exhibit low bioavailability and/or be targeted for delivery in the gastrointestinal tract. See, for example, US Patent No. 6,531,152.

**[00633]** Also provided is a method for inhibiting RET kinase activity in a cell, comprising contacting the cell with a compound of Formula I. In one embodiment, the contacting is in vitro. In one embodiment, the contacting is in vivo. In one embodiment, the contacting is in vivo, wherein the method comprises administering an effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to a subject having a cell having RET kinase activity. In some embodiments, the cell is a cancer cell. In one embodiment, the cancer cell is any cancer as described herein. In some embodiments, the cancer cell is a RET-associated cancer cell. In some embodiments, the cell is a gastrointestinal cell.

**[00634]** Also provided is a method for inhibiting RET kinase activity in a mammalian cell, comprising contacting the cell with a compound of Formula I. In one embodiment, the contacting is in vitro. In one embodiment, the contacting is in vivo. In one embodiment, the contacting is in vivo, wherein the method comprises administering an effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof to a mammal having a cell having RET kinase activity. In some embodiments, the mammalian cell is a mammalian cancer cell. In one embodiment, the mammalian cancer cell is any cancer as described herein. In some embodiments, the mammalian cancer cell is a RET-associated cancer cell. In some embodiments, the mammalian cell is a gastrointestinal cell.

**[00635]** As used herein, the term "contacting" refers to the bringing together of indicated moieties in an in vitro system or an in vivo system. For example, "contacting" a RET kinase with a compound provided herein includes the administration of a compound provided herein to an individual

or patient, such as a human, having a RET kinase, as well as, for example, introducing a compound provided herein into a sample containing a cellular or purified preparation containing the RET kinase.

**[00636]** Also provided herein is a method of inhibiting cell proliferation, in vitro or in vivo, the method comprising contacting a cell with an effective amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition thereof as defined herein.

**[00637]** The phrase "effective amount" means an amount of compound that, when administered to a patient in need of such treatment, is sufficient to (i) treat a RET kinase-associated disease or disorder, (ii) attenuate, ameliorate, or eliminate one or more symptoms of the particular disease, condition, or disorder, or (iii) delay the onset of one or more symptoms of the particular disease, condition, or disorder described herein. The amount of a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof that will correspond to such an amount will vary depending upon factors such as the particular compound, disease condition and its severity, the identity (e.g., weight) of the patient in need of treatment, but can nevertheless be routinely determined by one skilled in the art.

**[00638]** When employed as pharmaceuticals, the compound of Formula I (e.g., any one of Formulas I-A to I-L), including pharmaceutically acceptable salts or solvates thereof, can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration can be topical (including transdermal, epidermal, ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral or parenteral. Oral administration can include a dosage form formulated for once-daily or twice-daily (BID) administration. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus dose, or can be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration can include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable.

**[00639]** Also provided herein are pharmaceutical compositions which contain, as the active ingredient, a compound of Formula I (e.g., any one of Formulas I-A to I-L) or pharmaceutically acceptable salt or solvate thereof, in combination with one or more pharmaceutically acceptable carriers (excipients). For example, a pharmaceutical composition prepared using a compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof. In some embodiments, the composition is suitable for topical administration. In making the compositions provided herein, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups,

aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders. In some embodiments, the composition is formulated for oral administration. In some embodiments, the composition is a solid oral formulation. In some embodiments, the composition is formulated as a tablet or capsule.

**[00640]** Further provided herein are pharmaceutical compositions containing a compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof with a pharmaceutically acceptable carrier. Pharmaceutical compositions containing a compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof as the active ingredient can be prepared by intimately mixing the compound of Formula I or a pharmaceutically acceptable salt or solvate thereof with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier can take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral). In some embodiments, the composition is a solid oral composition.

**[00641]** Suitable pharmaceutically acceptable carriers are well known in the art. Descriptions of some of these pharmaceutically acceptable carriers can be found in *The Handbook of Pharmaceutical Excipients*, published by the American Pharmaceutical Association and the Pharmaceutical Society of Great Britain.

**[00642]** Methods of formulating pharmaceutical compositions have been described in numerous publications such as *Pharmaceutical Dosage Forms: Tablets, Second Edition, Revised and Expanded*, Volumes 1-3, edited by Lieberman et al; *Pharmaceutical Dosage Forms: Parenteral Medications*, Volumes 1-2, edited by Avis et al; and *Pharmaceutical Dosage Forms: Disperse Systems*, Volumes 1-2, edited by Lieberman et al; published by Marcel Dekker, Inc.

**[00643]** In preparing the compositions in oral dosage form, any of the usual pharmaceutical media can be employed. Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like. Solid oral preparations can also be coated with substances such as sugars or be enteric-coated so as to modulate major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other ingredients can be added to increase solubility or preservation. Injectable suspensions or solutions can also be prepared utilizing aqueous carriers along with appropriate additives. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described herein.

**[00644]** The compositions comprising a compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof can be formulated in a unit dosage form, each dosage containing from about 5 to about 1,000 mg (1 g), more usually about 100 mg to about 500 mg, of the active ingredient. The term “unit dosage form” refers to physically discrete units suitable as unitary dosages for human subjects and other patients, each unit containing a predetermined quantity of active material (*i.e.*, a compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof) calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

**[00645]** In some embodiments, the compositions provided herein contain from about 5 mg to about 50 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 5 mg to about 10 mg, about 10 mg to about 15 mg, about 15 mg to about 20 mg, about 20 mg to about 25 mg, about 25 mg to about 30 mg, about 30 mg to about 35 mg, about 35 mg to about 40 mg, about 40 mg to about 45 mg, or about 45 mg to about 50 mg of the active ingredient.

**[00646]** In some embodiments, the compositions provided herein contain from about 50 mg to about 500 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 50 mg to about 100 mg, about 100 mg to about 150 mg, about 150 mg to about 200 mg, about 200 mg to about 250 mg, about 250 mg to about 300 mg, about 350 mg to about 400 mg, or about 450 mg to about 500 mg of the active ingredient. In some embodiments, the compositions provided herein contain about 10 mg, about 20 mg, about 80 mg, or about 160 mg of the active ingredient.

**[00647]** In some embodiments, the compositions provided herein contain from about 500 mg to about 1,000 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 500 mg to about 550 mg, about 550 mg to about 600 mg, about 600 mg to about 650 mg, about 650 mg to about 700 mg, about 700 mg to about 750 mg, about 750 mg to about 800 mg, about 800 mg to about 850 mg, about 850 mg to about 900 mg, about 900 mg to about 950 mg, or about 950 mg to about 1,000 mg of the active ingredient.

**[00648]** The daily dosage of the compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof can be varied over a wide range from 1.0 to 10,000 mg per adult human per day, or higher, or any range therein. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 160, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.1 mg/kg to about 1000 mg/kg of body weight per day, or any range therein. Preferably, the range is from about 0.5 to about 500 mg/kg of body weight per day, or any range therein. More preferably, from about 1.0 to about 250 mg/kg of body weight per day, or any range therein. More preferably, from about 0.1 to about 100 mg/kg of body weight per day, or any range therein. In an example, the range can be from about 0.1 to about 50.0 mg/kg of body weight per day, or any amount or range therein. In another example, the range can be from about 0.1 to about 15.0 mg/kg of body weight per day, or any range therein. In yet another example, the range can be from about 0.5 to about 7.5 mg/kg of body weight per day, or any amount or range therein. Pharmaceutical

compositions containing a compound of Formula I (e.g., any one of Formulas I-A to I-L) or a pharmaceutically acceptable salt or solvate thereof can be administered on a regimen of 1 to 4 times per day or in a single daily dose.

**[00649]** The active compound may be effective over a wide dosage range and is generally administered in a pharmaceutically effective amount. Optimal dosages to be administered can be readily determined by those skilled in the art. It will be understood, therefore, that the amount of the compound actually administered will usually be determined by a physician, and will vary according to the relevant circumstances, including the mode of administration, the actual compound administered, the strength of the preparation, the condition to be treated, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient response, age, weight, diet, time of administration and severity of the patient's symptoms, will result in the need to adjust dosages.

**[00650]** In some embodiments, the compounds provided herein can be administered in an amount ranging from about 1 mg/kg to about 100 mg/kg. In some embodiments, the compound provided herein can be administered in an amount of about 1 mg/kg to about 20 mg/kg, about 5 mg/kg to about 50 mg/kg, about 10 mg/kg to about 40 mg/kg, about 15 mg/kg to about 45 mg/kg, about 20 mg/kg to about 60 mg/kg, or about 40 mg/kg to about 70 mg/kg. For example, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 25 mg/kg, about 30 mg/kg, about 35 mg/kg, about 40 mg/kg, about 45 mg/kg, about 50 mg/kg, about 55 mg/kg, about 60 mg/kg, about 65 mg/kg, about 70 mg/kg, about 75 mg/kg, about 80 mg/kg, about 85 mg/kg, about 90 mg/kg, about 95 mg/kg, or about 100 mg/kg. In some embodiments, such administration can be once-daily or twice-daily (BID) administration.

**[00651]** In some embodiments, the compounds provided herein can be administered in an amount of about about 10 mg twice a day (BID), 20 mg BID, about 40 mg BID, about 60 mg BID, about 80 mg BID, about 120 mg BID, about 160 mg BID, and about 240 mg BID. In some embodiments, each dose is administered at least six hours after the previous dose. In some embodiments, each dose is administered at least twelve hours after the previous dose.

**[00652]** In some embodiments, a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof exhibits pH dependent solubility at lower pH values. Accordingly, patients also receiving proton pump inhibitors (PPIs) and/or antacids may need to adjust the dosage of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof (e.g., increase the dose of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof). In some embodiments, the isoform of cytochrome P450 (CYP) that metabolizes a compound of Formula I (e.g., any one of Formulas I-A to I-L), or a pharmaceutically acceptable salt or solvate thereof, is CYP3A4. Accordingly, patients also receiving agents that inhibit or induce CYP3A4 may need to adjust the dosage of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof (e.g., increase the dose of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, in the case of a CYP3A4 inducer or decrease the dose of the compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, in the case of a CYP3A4 inhibitor).

**[00653]**

**[00654]** One skilled in the art will recognize that both *in vivo* and *in vitro* trials using suitable, known and generally accepted cell and/or animal models are predictive of the ability of a test compound to treat or prevent a given disorder.

**[00655]** One skilled in the art will further recognize that human clinical trials including first-in-human, dose ranging and efficacy trials, in healthy patients and/or those suffering from a given disorder, can be completed according to methods well known in the clinical and medical arts.

**[00656]** Provided herein are pharmaceutical kits useful, for example, in the treatment of RET-associated diseases or disorders, such as cancer or irritable bowel syndrome (IBS), which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound provided herein. Such kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit.

#### EXAMPLES

**[00657]** The following examples illustrate the invention.

**[00658]** Biological Examples

**[00659]** **Example A**

**[00660]** **RET Enzyme Assay**

**[00661]** The potency of compounds inhibiting several different RET kinase forms (Wild Type, V804M, G810R, & G810S) were determined using CisBio's HTRF® KinEASE™-TK assay technology. The kinases were incubated with 250 nM TK-substrate biotin (CisBio, part of cat # 62TKOPEC) at 1 mM ATP along with test compounds in a buffer consisting of 25 mM HEPES, pH 7.4, 10 mM MgCl<sub>2</sub>, 0.01% Triton X-100, and 2% DMSO in a volume of 8 µL. Compounds were typically prepared as a three-fold serial dilution in DMSO and added to the assay to give the appropriate final concentration. After a 30-min incubation at 22 °C, the reaction was quenched by adding 8 µL of quench solution containing 31.25 nM Sa-XL665 and 1x TK-Ab-Cryptate in HTRF detection buffer (all from CisBio, part of cat # 62TKOPEC). After a 1 hour incubation at 22 °C, the extent of reaction was determined using a PerkinElmer EnVision multimode plate reader via HTRF dual wavelength detection, and the percent of control (POC) was calculated using a ratiometric emission factor. One hundred POC was determined using DMSO only samples (no compound present), and 0 POC was determined using pre-quenched control reactions. A 4-parameter logistic curve was fit to the POC values as a function of the concentration of compound, and the IC<sub>50</sub> value was the point where the best-fit curve crossed 50 POC. Enzyme lots and concentrations



used as shown in the Table below, and the averaged IC<sub>50</sub> values for the compounds tested in these assay are provided in Table 5.

[00662]

| Enzyme form | Vendor               | Lot Number | Enzyme Concentration (nM) |
|-------------|----------------------|------------|---------------------------|
| Wild Type   | Eurofins             | 3654890-B  | 0.25                      |
| V804M       | Millipore            | D8KN029U-C | 0.2                       |
| G810R       | Array BioPharma Inc. | 160713     | 2.5                       |
| G810S       | Array BioPharma Inc. | 170322A    | 0.25                      |

[00663]      **Example B**

[00664]      **RET cell assay**

[00665]      The cellular potency of compounds inhibiting RET kinase were determined in HEK-293 cells expressing a Kif5b-RET fusion protein. Briefly, HEK-293 cells expressing a Kif5b-RET fusion protein were plated at 50K cells /well in 96 well poly-D-Lysine coated plates the day prior to the assay. The cells were incubated for 1 hour with test compound in DMEM (Dulbecco's Modified Eagle Medium) at a final DMSO concentration of 0.5%. Compounds were typically prepared in a three-fold serial dilution in DMSO and added to the assay to give the appropriate final concentration. After 1 hour the media was removed, the cells were fixed with 3.8% formaldehyde for 20 min, washed with PBS, and permeabilized for 10 min with 100% methanol. The plates were then washed with PBS-0.05% Tween20, and blocked with LI-COR Blocking solution (LI-COR Catalog No. 927-40000) for 1 hour. Plates were washed with PBS-0.05% Tween20, then incubated with anti-phospho-RET(Tyr1062) (Santa Cruz Catalog No. sc-20252-R) antibody and anti-GAPDH (Millipore Catalog No. MAB374) antibody for 2 hours. The plates were washed with PBS-0.05% Tween20, and incubated with anti-rabbit 680 (Molecular Probes Catalog No. A21109) and anti-mouse 800 (LI-COR Catalog No. 926-32210) secondary antibodies for 1 hour. All antibodies were diluted in LI-COR Block containing 0.05% Tween. The plates were washed with PBS-0.05% Tween20, 100  $\mu$ L PBS is added to each well, and the plates were read on a LI-COR Aeries fluorescent plate reader. The phospho-RET signal was normalized to the GAPDH signal. 100 POC (percent of control) was determined using no test compounds and 0 POC was determined using 1  $\mu$ M of a control inhibitor. The POC values were fit to a 4 parameter logistic curve. The IC<sub>50</sub> value is the point where the curve crosses 50 POC. IC<sub>50</sub> values for the compounds tested in these assay are provided in Table 5.

[00666]      **Example C**

[00667]      **RET G810R and G810S mutant assay**

**[00668]** The potency of a compound inhibiting G810R or G810S mutant RET kinase was determined using CisBio's HTRF® KinEASE™-TK assay technology. The assays contained G810R or G810S mutant RET produced at Array Biopharma, Inc. (1 nM enzyme). The kinase was incubated with 250 nM TK-substrate biotin (CisBio, part of Catalog No. 62TK0PEC) and 1 mM ATP along with test compound in a buffer consisting of 25 mM HEPES, pH 7.4, 10 mM MgCl<sub>2</sub>, 0.01% Triton X-100, and 2% DMSO in a volume of 8 µL. Compounds were typically prepared as a three-fold serial dilution in DMSO and added to the assay to give the appropriate final concentration. After a 60-min incubation at 22 °C, the reaction was quenched by adding 8 µL of quench solution containing 31.25 nM Sa-XL665 and 1x TK-Ab-Cryptate in HTRF detection buffer (all from CisBio, part of Catalog No. 62TK0PEC). After a 1-h incubation at 22 °C, the extent of reaction was determined using a PerkinElmer EnVision multimode plate reader via HTRF dual wavelength detection, and the percent of control (POC) was calculated using a ratiometric emission factor. One hundred POC was determined using no test compounds, and 0 POC was determined using pre-quenched control reactions. A 4-parameter logistic curve was fit to the POC values as a function of the concentration of compound, and the IC<sub>50</sub> value is the point where the best-fit curve crossed 50 POC. IC<sub>50</sub> values for the compounds tested in these assay are provided in Table 5.

**[00669] Example D**

**[00670] RET G810R and G810S mutant cell assay**

**[00671]** The cellular potency of compounds inhibiting RET kinase were determined in HEK-293 cells expressing a G810R or G810S mutant RET Kif5b-RET fusion protein. Briefly, HEK-293 cells expressing a G810R or G810S mutant RET Kif5b-RET fusion protein were plated at 50K cells /well in 96 well poly-D-Lysine coated plates the day prior to the assay. The cells were incubated for 1 hour with test compound in DMEM (Dulbecco's Modified Eagle Medium) at a final DMSO concentration of 0.5%. Compounds were typically prepared in a three-fold serial dilution in DMSO and added to the assay to give the appropriate final concentration. After 1 hour the media was removed, the cells were fixed with 3.8% formaldehyde for 20 min, washed with PBS, and permeabilized for 10 min with 100% methanol. The plates were then washed with PBS-0.05% Tween20, and blocked with LI-COR Blocking solution (LI-COR Catalog No. 927-40000) for 1 hour. Plates were washed with PBS-0.05% Tween20, then incubated with anti-phospho-RET(Tyr1062) (Santa Cruz Catalog No. sc-20252-R) antibody and anti-GAPDH (Millipore Catalog No. MAB374) antibody for 2 hours. The plates were washed with PBS-0.05% Tween20, and incubated with anti-rabbit 680 (Molecular Probes Catalog No. A21109) and anti-mouse 800 (LI-COR Catalog No. 926-32210) secondary antibodies for 1 hour. All antibodies were diluted in LI-COR Block containing 0.05% Tween. The plates were washed with PBS-0.05% Tween20, 100 µL PBS was added to each well, and the plates were read on a LI-COR Aeries fluorescent plate reader. The phospho-RET signal was normalized to the GAPDH signal. 100 POC (percent of control) was determined using no test compounds and 0 POC was determined using 1 µM of a control inhibitor. The POC values are fit to a 4 parameter logistic curve. The IC<sub>50</sub> value is the point where the curve crosses 50 POC. IC<sub>50</sub> values for the compounds tested in these assay are provided in Table 5.

**Table 5.** IC<sub>50</sub>'s of compounds tested in the assay of Example A.

| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 1   | 52                                          | 202                                                     | 145                                                     | 117                                                     | 17                                                     | 179                                                     | N/A                                                     |
| 2   | 114                                         | 324                                                     | 243                                                     | 248                                                     | 99                                                     | 406                                                     | N/A                                                     |
| 3   | 16                                          | 105                                                     | 151                                                     | 84                                                      | 40                                                     | 195                                                     | 81                                                      |
| 4   | 16                                          | 106                                                     | 157                                                     | 135                                                     | 60                                                     | 290                                                     | N/A                                                     |
| 5   | 17                                          | 99                                                      | 157                                                     | 138                                                     | 75                                                     | 459                                                     | N/A                                                     |
| 6   | 58                                          | 297                                                     | 143                                                     | 178                                                     | 64                                                     | 287                                                     | N/A                                                     |
| 7   | 91                                          | 340                                                     | 171                                                     | 356                                                     | 135                                                    | 660                                                     | N/A                                                     |
| 8   | 41                                          | 227                                                     | 95                                                      | 123                                                     | 91                                                     | 302                                                     | N/A                                                     |
| 9   | 103                                         | 533                                                     | 224                                                     | 298                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 10  | 20                                          | 106                                                     | 114                                                     | 46                                                      | 28                                                     | 254                                                     | N/A                                                     |
| 11  | 136                                         | 516                                                     | 345                                                     | 507                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 12  | 47                                          | 399                                                     | 176                                                     | 112                                                     | 143                                                    | 705                                                     | N/A                                                     |
| 13  | 36                                          | 275                                                     | 137                                                     | 72                                                      | 119                                                    | 596                                                     | N/A                                                     |
| 14  | 112                                         | 865                                                     | 252                                                     | 293                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 15  | 58                                          | 607                                                     | 361                                                     | 307                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 16  | 205                                         | 1794                                                    | 354                                                     | 648                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 17  | 49                                          | 241                                                     | 533                                                     | 433                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 18  | 153                                         | 692                                                     | 253                                                     | 397                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 19  | 45                                          | 202                                                     | 103                                                     | 215                                                     | 53                                                     | 262                                                     | 184                                                     |
| 20  | 53                                          | 254                                                     | 135                                                     | 123                                                     | 28                                                     | 160                                                     | 111                                                     |
| 21  | 9                                           | 44                                                      | 31                                                      | N/A                                                     | 28                                                     | 87                                                      | 83                                                      |
| 22  | 31                                          | 323                                                     | 102                                                     | 75                                                      | 65                                                     | 496                                                     | N/A                                                     |

| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 23  | 18                                          | 189                                                     | 63                                                      | 54                                                      | 20                                                     | 133                                                     | 107                                                     |
| 24  | 25                                          | 92                                                      | 70                                                      | 87                                                      | 40                                                     | 181                                                     | N/A                                                     |
| 25  | 14                                          | 123                                                     | 37                                                      | N/A                                                     | 7                                                      | 142                                                     | N/A                                                     |
| 26  | 20                                          | 145                                                     | 62                                                      | 45                                                      | 22                                                     | 196                                                     | N/A                                                     |
| 27  | 142                                         | 595                                                     | 308                                                     | 581                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 28  | 307                                         | 2833                                                    | 641                                                     | 1025                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 29  | 8                                           | 26                                                      | 27                                                      | 25                                                      | 12                                                     | 71                                                      | 37                                                      |
| 30  | 93                                          | 259                                                     | 180                                                     | 300                                                     | 155                                                    | 2161                                                    | N/A                                                     |
| 31  | 571                                         | 2207                                                    | 1315                                                    | 1322                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 32  | 188                                         | 10000                                                   | 1232                                                    | 10000                                                   | N/A                                                    | N/A                                                     | N/A                                                     |
| 33  | 1185                                        | 10000                                                   | 10000                                                   | 10000                                                   | N/A                                                    | N/A                                                     | N/A                                                     |
| 34  | 227                                         | 805                                                     | 288                                                     | 397                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 35  | 47                                          | 114                                                     | 124                                                     | 110                                                     | 35                                                     | 831                                                     | N/A                                                     |
| 36  | 58                                          | 235                                                     | 130                                                     | 217                                                     | 53                                                     | 144                                                     | 255                                                     |
| 37  | 318                                         | 10000                                                   | 1284                                                    | 1222                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 38  | 342                                         | 3122                                                    | 649                                                     | 921                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 39  | 893                                         | 10000                                                   | 3066                                                    | 4137                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 40  | 621                                         | 2205                                                    | 724                                                     | 973                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 41  | 198                                         | 659                                                     | 415                                                     | 564                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 42  | 72                                          | 594                                                     | 166                                                     | 176                                                     | 45                                                     | 385                                                     | N/A                                                     |
| 43  | 104                                         | 187                                                     | 90                                                      | 223                                                     | 105                                                    | 298                                                     | 213                                                     |
| 44  | 104                                         | 248                                                     | 196                                                     | 245                                                     | 142                                                    | 1888                                                    | N/A                                                     |
| 45  | 194                                         | 10000                                                   | 275                                                     | 557                                                     | 344                                                    | 814                                                     | N/A                                                     |
| 46  | 330                                         | 10000                                                   | 400                                                     | 973                                                     | N/A                                                    | N/A                                                     | N/A                                                     |

| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 47  | 112                                         | 9371                                                    | 269                                                     | 303                                                     | 134                                                    | 1949                                                    | N/A                                                     |
| 48  | 2443                                        | 10000                                                   | 7876                                                    | 10000                                                   | N/A                                                    | N/A                                                     | N/A                                                     |
| 49  | 8422                                        | 10000                                                   | 10000                                                   | 10000                                                   | N/A                                                    | N/A                                                     | N/A                                                     |
| 50  | 2183                                        | 10000                                                   | 2570                                                    | 5082                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 51  | 7364                                        | 10000                                                   | 10000                                                   | 10000                                                   | N/A                                                    | N/A                                                     | N/A                                                     |
| 52  | 1055                                        | 10000                                                   | 877                                                     | 3199                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 53  | 16                                          | 34                                                      | 102                                                     | 94                                                      | 64                                                     | 485                                                     | N/A                                                     |
| 54  | 13                                          | 41                                                      | 70                                                      | 54                                                      | 22                                                     | 103                                                     | 92                                                      |
| 55  | 8                                           | 21                                                      | 31                                                      | 29                                                      | 10                                                     | 46                                                      | 40                                                      |
| 56  | 9                                           | 51                                                      | 79                                                      | 42                                                      | 7                                                      | 50                                                      | 37                                                      |
| 57  | 29                                          | 178                                                     | 123                                                     | 176                                                     | 38                                                     | 131                                                     | 121                                                     |
| 58  | 23                                          | 66                                                      | 64                                                      | 159                                                     | 49                                                     | 186                                                     | 181                                                     |
| 59  | 275                                         | 749                                                     | 308                                                     | 859                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 60  | 21                                          | 98                                                      | 45                                                      | 40                                                      | 29                                                     | 157                                                     | 101                                                     |
| 61  | 31                                          | 136                                                     | 89                                                      | 58                                                      | 33                                                     | 281                                                     | 101                                                     |
| 62  | 34                                          | 138                                                     | 96                                                      | 74                                                      | 52                                                     | 349                                                     | 192                                                     |
| 63  | 61                                          | 211                                                     | 118                                                     | 108                                                     | 72                                                     | 439                                                     | 352                                                     |
| 64  | 39                                          | 182                                                     | 169                                                     | 76                                                      | 47                                                     | 119                                                     | 90                                                      |
| 65  | 43                                          | 163                                                     | 64                                                      | 133                                                     | 45                                                     | 621                                                     | 290                                                     |
| 66  | 1792                                        | 6834                                                    | 2185                                                    | 2302                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 67  | 23                                          | 92                                                      | 53                                                      | 69                                                      | 60                                                     | 467                                                     | N/A                                                     |
| 68  | 386                                         | 10000                                                   | 485                                                     | 1165                                                    | 1775                                                   | 4747                                                    | N/A                                                     |
| 69  | 34                                          | 10000                                                   | 47                                                      | 65                                                      | 103                                                    | 394                                                     | N/A                                                     |
| 70  | 92                                          | 10000                                                   | 183                                                     | 312                                                     | 166                                                    | 607                                                     | 388                                                     |

| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 71  | 1161                                        | 10000                                                   | 3735                                                    | 4753                                                    | 2529                                                   | 6525                                                    | 3623                                                    |
| 72  | 512                                         | 10000                                                   | 3091                                                    | 2629                                                    | 3957                                                   | 16180                                                   | 6351                                                    |
| 73  | 531                                         | 10000                                                   | 2738                                                    | 2478                                                    | 13910                                                  | 16667                                                   | 16667                                                   |
| 74  | 353                                         | 6004                                                    | 2083                                                    | 289                                                     | N/A                                                    | 2944                                                    | 918                                                     |
| 75  | 82                                          | 2031                                                    | 421                                                     | 51                                                      | N/A                                                    | 170                                                     | 191                                                     |
| 76  | 117                                         | 10000                                                   | 118                                                     | 292                                                     | 143                                                    | 845                                                     | 570                                                     |
| 77  | 27                                          | 10000                                                   | 135                                                     | 73                                                      | 13                                                     | 89                                                      | 32                                                      |
| 78  | 638                                         | 10000                                                   | 507                                                     | 720                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 79  | 356                                         | 10000                                                   | 270                                                     | 591                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 80  | 195                                         | 10000                                                   | 320                                                     | 404                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 81  | 19                                          | 5488                                                    | 94                                                      | 52                                                      | 11                                                     | 76                                                      | 39                                                      |
| 82  | 358                                         | 10000                                                   | 420                                                     | 821                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 83  | 63                                          | 10000                                                   | 58                                                      | 97                                                      | 293                                                    | 1062                                                    | 812                                                     |
| 84  | 182                                         | 10000                                                   | 244                                                     | 634                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 85  | 389                                         | 10000                                                   | 501                                                     | 960                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 86  | 37                                          | 8320                                                    | 150                                                     | 98                                                      | 73                                                     | N/A                                                     | 169                                                     |
| 87  | 47                                          | 5783                                                    | 202                                                     | 204                                                     | 32                                                     | 89                                                      | 71                                                      |
| 88  | 62                                          | 6257                                                    | 288                                                     | 297                                                     | 50                                                     | 53                                                      | 44                                                      |
| 89  | 311                                         | 10000                                                   | 544                                                     | 1583                                                    | 233                                                    | N/A                                                     | 269                                                     |
| 90  | 353                                         | 10000                                                   | 1059                                                    | 1597                                                    | 222                                                    | N/A                                                     | 408                                                     |
| 91  | 166                                         | 10000                                                   | 624                                                     | 596                                                     | 66                                                     | N/A                                                     | 98                                                      |
| 92  | 114                                         | 10000                                                   | 285                                                     | 168                                                     | 56                                                     | N/A                                                     | 301                                                     |
| 93  | 1263                                        | 10000                                                   | 1914                                                    | 5573                                                    | 1000                                                   | N/A                                                     | 1000                                                    |
| 94  | 568                                         | 10000                                                   | 840                                                     | 2001                                                    | 388                                                    | N/A                                                     | 297                                                     |

| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 95  | 546                                         | 10000                                                   | 816                                                     | 2080                                                    | 318                                                    | N/A                                                     | 376                                                     |
| 96  | 27                                          | 10000                                                   | 187                                                     | 393                                                     | 85                                                     | 142                                                     | 116                                                     |
| 97  | 73                                          | 4457                                                    | 478                                                     | 457                                                     | 26                                                     | 47                                                      | 47                                                      |
| 98  | 2212                                        | 10000                                                   | 4069                                                    | 7952                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 99  | 881                                         | 10000                                                   | 2210                                                    | 3251                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 100 | 233                                         | 2856                                                    | 1605                                                    | 914                                                     | 10                                                     | N/A                                                     | 15                                                      |
| 101 | 135                                         | 10000                                                   | 1416                                                    | 805                                                     | 43                                                     | 75                                                      | 62                                                      |
| 102 | 128                                         | 483                                                     | 760                                                     | 183                                                     | 35                                                     | N/A                                                     | 86                                                      |
| 103 | 43                                          | 6096                                                    | 190                                                     | 196                                                     | 41                                                     | N/A                                                     | 141                                                     |
| 104 | 54                                          | 8234                                                    | 186                                                     | 259                                                     | 36                                                     | N/A                                                     | 92                                                      |
| 105 | 639                                         | 10000                                                   | 742                                                     | 2584                                                    | 1355                                                   | 15535                                                   | N/A                                                     |
| 106 | 404                                         | 10000                                                   | 2425                                                    | 3266                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 107 | 10000                                       | 10000                                                   | 10000                                                   | 10000                                                   | N/A                                                    | N/A                                                     | N/A                                                     |
| 108 | 10000                                       | 10000                                                   | 8385                                                    | 10000                                                   | 12859                                                  | 16667                                                   | N/A                                                     |
| 109 | 1776                                        | 10000                                                   | 1810                                                    | 2268                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 110 | 411                                         | 1742                                                    | 214                                                     | 284                                                     | N/A                                                    | 537                                                     | 201                                                     |
| 111 | 3443                                        | 6237                                                    | 3296                                                    | 2708                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 112 | 1776                                        | 3735                                                    | 2017                                                    | 958                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 113 | 102                                         | 169                                                     | 106                                                     | 63                                                      | 513                                                    | N/A                                                     | 274                                                     |
| 114 | 357                                         | 1051                                                    | 435                                                     | 282                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 115 | 346                                         | 1090                                                    | 179                                                     | 153                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 116 | 1296                                        | 1059                                                    | 238                                                     | 491                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 117 | 373                                         | 863                                                     | 300                                                     | 304                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 118 | 131                                         | 99                                                      | 80                                                      | 75                                                      | 125                                                    | 206                                                     | 108                                                     |

| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 119 | 24                                          | 79                                                      | 45                                                      | 67                                                      | 66                                                     | 593                                                     | 131                                                     |
| 120 | 10                                          | 20                                                      | 74                                                      | 24                                                      | 15                                                     | 35                                                      | 33                                                      |
| 121 | 41                                          | 120                                                     | 87                                                      | 84                                                      | 74                                                     | N/A                                                     | 118                                                     |
| 122 | 14                                          | 38                                                      | 28                                                      | 26                                                      | 31                                                     | N/A                                                     | 77                                                      |
| 123 | 39                                          | 87                                                      | 103                                                     | 86                                                      | 21                                                     | N/A                                                     | 42                                                      |
| 124 | 24                                          | 80                                                      | 89                                                      | 49                                                      | 19                                                     | 44                                                      | 33                                                      |
| 125 | 11                                          | 19                                                      | 41                                                      | 19                                                      | 7                                                      | 16                                                      | 16                                                      |
| 126 | 6                                           | 30                                                      | 15                                                      | 9                                                       | 5                                                      | 29                                                      | 26                                                      |
| 127 | 6                                           | 21                                                      | 28                                                      | 23                                                      | 9                                                      | 57                                                      | 52                                                      |
| 128 | 88                                          | 349                                                     | 98                                                      | 160                                                     | 157                                                    | N/A                                                     | 435                                                     |
| 129 | 8                                           | 21                                                      | 23                                                      | 19                                                      | 18                                                     | 46                                                      | 65                                                      |
| 130 | 12                                          | 64                                                      | 40                                                      | 32                                                      | 34                                                     | N/A                                                     | 131                                                     |
| 131 | 14                                          | 41                                                      | 97                                                      | 18                                                      | 14                                                     | 30                                                      | 60                                                      |
| 132 | 1709                                        | 1936                                                    | 4462                                                    | 2318                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 133 | 2642                                        | 6204                                                    | 6142                                                    | 3402                                                    | N/A                                                    | N/A                                                     | N/A                                                     |
| 134 | 112                                         | 355                                                     | 560                                                     | 488                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 135 | 45                                          | 152                                                     | 150                                                     | 229                                                     | 54                                                     | N/A                                                     | 103                                                     |
| 136 | 201                                         | 551                                                     | 274                                                     | 817                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 137 | 38                                          | 60                                                      | 52                                                      | 155                                                     | 67                                                     | N/A                                                     | 280                                                     |
| 138 | 143                                         | 376                                                     | 272                                                     | 646                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 139 | 75                                          | 10000                                                   | 1032                                                    | 474                                                     | 16                                                     | 49                                                      | 36                                                      |
| 140 | 554                                         | 8965                                                    | 876                                                     | 3072                                                    | 884                                                    | N/A                                                     | 1000                                                    |
| 141 | 139                                         | 415                                                     | 211                                                     | 405                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 142 | 124                                         | 505                                                     | 157                                                     | 425                                                     | N/A                                                    | N/A                                                     | N/A                                                     |




| Ex# | RET Enz<br>FRET_WT<br>IC <sub>50</sub> (nM) | RET<br>V804M<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810R<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | RET<br>G810S<br>Enz<br>FRET<br>IC <sub>50</sub><br>(nM) | KIF5B-RET<br>pTYR1062<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810R<br>Cell IC <sub>50</sub><br>(nM) | KIF5B-<br>RET<br>G810S<br>Cell IC <sub>50</sub><br>(nM) |
|-----|---------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|--------------------------------------------------------|---------------------------------------------------------|---------------------------------------------------------|
| 143 | 53                                          | 438                                                     | 109                                                     | 261                                                     | 151                                                    | N/A                                                     | 447                                                     |
| 144 | 24                                          | 57                                                      | 61                                                      | 101                                                     | 40                                                     | N/A                                                     | 177                                                     |
| 145 | 20                                          | 31                                                      | 85                                                      | 101                                                     | 73                                                     | 120                                                     | 93                                                      |
| 146 | 26                                          | 87                                                      | 43                                                      | 116                                                     | 63                                                     | 98                                                      | 68                                                      |
| 147 | 14                                          | 32                                                      | 74                                                      | 94                                                      | 21                                                     | 56                                                      | 90                                                      |
| 148 | 183                                         | 216                                                     | 154                                                     | 758                                                     | N/A                                                    | N/A                                                     | N/A                                                     |
| 149 | 7                                           | 22                                                      | 70                                                      | 41                                                      | 2                                                      | N/A                                                     | 20                                                      |
| 150 | 12                                          | 15                                                      | 84                                                      | 46                                                      | 4                                                      | 40                                                      | 18                                                      |
| 151 | 9                                           | 27                                                      | 86                                                      | 55                                                      | 14                                                     | 113                                                     | 57                                                      |

## [00672] Synthetic examples

### Preparation of Synthetic Intermediates

#### Intermediate P1



#### 4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde

Step 1: Preparation of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde. To a mixture of 4-chloro-7H-pyrrolo[2,3-d]pyrimidine-5-carboxaldehyde (25.0 g, 138 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (53.8 g, 165 mmol) in DMF (275 mL) was added 2-bromopropane (15.5 mL, 165 mmol). The reaction was stirred overnight at 60 °C. It was diluted with EtOAc (100 mL), washed with water (2x100mL) and brine (1x), then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated *in vacuo*. The crude residue was purified by silica


chromatography (0-40% acetone in DCM) to afford the title compound as a pale yellowish solid (21.7 g, 70% yield). MS (apci)  $m/z$  = 224.1 (M+H).

**Step 2: Preparation of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime.** A solution of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (4.75 g, 21.2 mmol) in EtOH (21 mL) was treated with hydroxylamine hydrochloride (3.69 g, 53.1 mmol) and NaOAc (4.36 g, 53.1 mmol), then stirred for 1 h at RT. The mixture was then diluted with DCM, and the solids were filtered off. The filtrate was concentrated *in vacuo*, and the residue was purified by silica chromatography (0-20% MeOH in DCM) to afford the title compound (4.98 g, 98% yield). LCMS  $m/z$  = 239.1 (M+H).

**Step 3: Preparation of 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-Cyclopropyl-isoxazole.** A mixture of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime (12.4 g, 52.0 mmol), 1-chloropyrrolidine-2,5-dione (7.63 g, 57.1 mmol),  $\text{KHCO}_3$  (6.76 g, 67.5 mmol) and ethynylcyclopropane (13.2 mL, 156 mmol) in DMF (260 mL) was stirred for 2 d at RT. The mixture was poured into water (800 mL) and filtered. The solids collected were rinsed with hexanes (200 mL), and dried *in vacuo* to afford the title compound (15.7 g, quantitative yield). MS (apci)  $m/z$  = 303.1 (M+H).

**Step 4: Preparation of 4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]-pyrimidine-6-carbaldehyde.** To a solution of 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (5.27 g, 17.4 mmol) and diisopropylamine (0.98 mL, 6.96 mmol) in THF (58 mL) was slowly added *n*-BuLi (2.5 M in THF; 9.75 mL, 24.37 mmol) at -78 °C. After stirring 30 min, the mixture was cannulated into a solution of ethyl formate (4.2 mL, 52.23 mmol) in THF (5 mL) at -78 °C. The reaction was stirred for 1 h, after which it was quenched with saturated  $\text{NH}_4\text{Cl}_{(\text{aq})}$ . The mixture was partitioned between EtOAc and water. After phase separation, the aqueous phase was extracted with EtOAc (2x). The combined organic extracts were washed with brine, then dried over anhydrous  $\text{Na}_2\text{SO}_4(\text{s})$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (10-50% EtOAc in hexanes) to give the title compound (3.7 g, 64% yield). MS (apci)  $m/z$  = 331.1 (M+H).

## Intermediate P2



## 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole


**Step 1: Preparation of (4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol.** A solution of 4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P1**) (3.54 g, 10.7 mmol) in methanol (36 mL) was cooled to 0 °C, and treated with  $\text{NaBH}_4$  (0.40 g, 10.7 mmol). After 1 h stirring, the reaction was diluted with EtOAc,

and washed with water (2x) and brine (2x). The organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound as white solid (3.6 g, quantitative yield). MS (apci)  $m/z = 333.1$  (M+H).

**Step 2: Preparation of (4-chloro-5-(5-cyclopropyl-4-iodoisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol.** A suspension of (4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (3.56 g, 10.7 mmol) in ACN (70 mL) was treated sequentially with NIS (3.61 g, 16.0 mmol) and TFA (7.13 mL, 10.7 mmol). After 90 min stirring at RT, the reaction was diluted with EtOAc, washed successively with water (1x), saturated  $\text{NaHCO}_{3(aq)}$  (1x), 10%  $\text{Na}_2\text{S}_2\text{O}_3$ , and brine. The organic extracts then were dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound as white solid (4.9 g, quantitative yield). MS (apci)  $m/z = 459.0$  (M+H).


**Step 3: Preparation of 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole.** A solution of (4-chloro-5-(5-cyclopropyl-4-iodoisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (5.53 g, 12.1 mmol) in DMF (80 mL) was treated with TBDMSCl (2.73 g, 18.1 mmol) and 1H-imidazole (1.64 g, 24.1 mmol). After stirring overnight, the reaction was diluted with EtOAc, washed with water and brine, then dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered and concentrated *in vacuo*. The crude was purified by silica chromatography (0-25% EtOAc in hexanes) to give the title compound as a clear yellowish oil (6.2 g, 90% yield). MS (apci)  $m/z = 573.1$  (M+H).

### Intermediate P3




### methyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate

A cold (-78 °C) solution of 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole (**Intermediate P2**; 594 mg, 1.04 mmol) in THF (6.912 mL) was treated dropwise with n-BuLi (2.5 M in THF; 580.6  $\mu\text{L}$ , 1.451 mmol). The resulting mixture was stirred for 30 min at -78 °C. Subsequently, the mixture was treated slowly with dimethyl carbonate (349.4  $\mu\text{L}$ , 4.147 mmol), and then stirred for 1 h at -10 °C. The reaction mixture was quenched with saturated  $\text{NH}_4\text{Cl}_{(aq)}$ , and allowed to warm to RT. The quenched mixture was partitioned between EtOAc and water. After phase-separation, the organic extracts were washed with brine, then dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo*. The residue was purified by silica chromatography (10-50% hexanes: EtOAc) to afford the title compound (279 mg, 53% yield). MS (apci)  $m/z = 505.2$  (M+H).

**Intermediate P4**

ethyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate

The title compound (1.1 g, 51% yield) was prepared, worked up and purified using a similar procedure to that described for methyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P3**), replacing dimethyl carbonate with diethyl carbonate. MS (apci)  $m/z$  = 519.2 (M+H).

**Intermediate P5**

ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate


Step 1: Preparation of ethyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate. A solution of ethyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P4**; 700 mg, 1.35 mmol) and (2,4-dimethoxyphenyl)methylamine (607.7  $\mu$ L, 4.045 mmol) in DMSO (8.99 mL) was stirred overnight at 60 °C. After cooling to RT, the reaction mixture was partitioned between EtOAc and water. Following phase separation, the organic extracts were washed sequentially with additional water (3x) and brine. The combined organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (10-90% hexanes: EtOAc as the gradient eluent) to afford the title compound (826 mg, 94% yield). MS (apci)  $m/z$  = 650.3 (M+H).

Step 2: Preparation of ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate. TBAF (1.0 M THF) (1.65 mL, 1.65 mmol) was added to a cold (0 °C) solution of ethyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-

carboxylate (826 mg, 1.27 mmol) and acetic acid (17.4 M; 219  $\mu$ L, 3.81 mmol) in THF (8.47 mL). After being allowed to slowly warm to RT, the reaction mixture was stirred overnight at RT. The mixture was then partitioned between DCM and water. Following phase separation, the organic extracts were washed with brine, then dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound (446 mg, 66% yield). MS (apci)  $m/z$  = 536.2 (M+H).

Step 3: Preparation of ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate. A cold (0 °C) solution of ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate (446 mg, 0.833 mmol) and methanesulfonyl chloride (67.7  $\mu$ L, 0.874 mmol) in DCM (8.33 mL) was treated with DIEA (580  $\mu$ L, 3.33 mmol), and stirred overnight at RT. The reaction mixture was then partitioned between DCM and water. Following phase separation, the organic extracts were washed with brine, then dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound, which was used directly in the next step without further purifications, assuming quantitative yield.


#### Intermediate P6



methyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate

The title compound was prepared in a similar fashion as described for **Intermediate P5**, replacing ethyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P4**) in step 1 with methyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P3**) instead.

#### Intermediate P7




Ethyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate

Step 1: Preparation of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde. A solution of 4-chloro-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (3.6 g, 19.8 mmol) in DMF (40 mL) was treated with  $\text{Cs}_2\text{CO}_3$  (7.1 g, 21.8 mmol) and 2-bromopropane (2.2 mL, 23.8 mmol). The reaction mixture was then stirred overnight at 50 °C. After cooling to RT, the reaction was diluted with water (100 mL) and filtered. The filter cake was rinsed with water (30 mL) and then air-dried to afford the title compound as off-white solid (3.1 g, 70% yield). MS (apci)  $m/z$  = 224.1 (M+H).

Step 2: Preparation of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime. A mixture of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (1.15 g, 5.14 mmol),  $\text{NH}_2\text{OH}\cdot\text{HCl}$  (0.39 g, 5.66 mmol) and NaOAc (0.46 g, 5.66 mmol) in EtOH (26 mL) was stirred for 30 min at 50 °C. After cooling to RT, the reaction was diluted with DCM (50 mL), and filtered through a celite pad. The filtrate was concentrated *in vacuo* to afford the title compound as yellowish solid (1.25 g, quantitative yield). MS (apci)  $m/z$  = 239.1 (M+H).

Step 3: Preparation of ethyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate. In a pressure vessel, a solution of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime (96.6 mg, 0.40 mmol) in 1,2-dimethoxyethane (8.1 mL) was treated with NCS (59.4 mg, 0.44 mmol). The vessel was then sealed, and the reaction was stirred overnight at RT. Ethyl 3-cyclopropylpropionate (**Intermediate R1**) (559  $\mu\text{L}$ , 4.05 mmol) and  $\text{KHCO}_3$  (122 mg, 1.21 mmol) were then introduced, and the reaction was again sealed and heated at 65 °C for 2 h. The reaction mixture then was filtered. The filtrate was concentrated and purified by silica chromatography (0-70% EtOAc in hexanes as the gradient eluent) to afford the title compound (82.4 mg, 54% yield). MS (apci)  $m/z$  = 375.1 (M+H).

**Intermediate P8**




5-(4-(((tert-butyldimethylsilyl)oxy)methyl)-5-cyclopropylisoxazol-3-yl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde

**Step 1: Preparation of (3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)methanol.** A cold (-78 °C) solution of ethyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P7**; 128 mg, 0.342 mmol) in DCM (1.366  $\mu$ L) was treated dropwise with DIBAL-H (25 wt.% in toluene; 459  $\mu$ L, 0.68 mmol). After stirring the resulting mixture for 30 min at -78 °C, additional DIBAL-H (230  $\mu$ L, 0.34 mmol) was introduced. The mixture was stirred for an additional 30 min at -78 °C before quenching with water. The aqueous phase of the quench solidified at -78 °C, and the organic phase was separated by decanting. The organic extracts were washed successively with water and brine, then passed through a phase separator column and concentrated *in vacuo* to afford the title compound (100.6 mg, 89% yield). MS (apci)  $m/z$  = 333.1 (M+H).

**Step 2: Preparation of 4-(((tert-butyldimethylsilyl)oxy)methyl)-3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole.** A solution of (3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)methanol (501.4 mg, 1.51 mmol) in DMF (6.027 mL) was treated with tert-butyldichlorodimethylsilane (454.2 mg, 3.013 mmol) and 1H-imidazole (307.7 mg, 4.520 mmol) then stirred for 45 min at RT. The resulting mixture was quenched with water and extracted with EtOAc. The combined organic extracts were washed successively with water and brine then dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-50% EtOAc in Hexanes as the gradient eluent) to afford the title compound (597.3 mg, 89% yield). MS (apci)  $m/z$  = 447.2 (M+H).

**Step 3: Preparation of 5-(4-(((tert-butyldimethylsilyl)oxy)methyl)-5-cyclopropylisoxazol-3-yl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde.** A cold (-78 °C) solution of 4-(((tert-butyldimethylsilyl)oxy)methyl)-3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (338 mg, 0.756 mmol) in THF (7.561 mL) was treated with LDA (2 M THF) (1.13 mL, 2.27 mmol). After stirring the mixture for 30 min at -78 °C, ethyl formate (123.1  $\mu$ L, 1.512 mmol) was added. The resulting mixture was stirred overnight at -20 °C. After being allowed to warm to RT, the reaction mixture was quenched with saturated  $\text{NH}_4\text{Cl}_{(aq)}$ . The quenched mixture was diluted with water and extracted with EtOAc. The organic extracts were washed with brine, dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-50% EtOAc in hexanes as the gradient eluent) to afford the title compound (179.3 mg, 50% yield). MS (apci)  $m/z$  = 475.2 (M+H).


**Intermediate P9**



4-(bis(4-methoxybenzyl)amino)-5-(4-(((tert-butyldimethylsilyl)oxy)methyl)-5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde

A solution of 5-(4-(((tert-butyldimethylsilyl)oxy)methyl)-5-cyclopropylisoxazol-3-yl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P8**; 106 mg, 0.22 mmol) in DMSO (444  $\mu$ L) was treated with bis(4-methoxybenzyl)amine (336.2 mg, 1.110 mmol) and DIEA (387.9  $\mu$ L, 2.221 mmol). The resulting mixture was stirred for 1 h at 95  $^{\circ}$ C. After subsequently cooling to RT, the reaction was quenched with water and extracted with EtOAc (x3). The combined organic extracts were washed with brine, dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-50% EtOAc in hexanes as the gradient eluent) to afford the title compound (140 mg, 91% yield). MS (apci)  $m/z$  = 696.3 (M+H).

**Intermediate P10**




3-(4-(bis(4-methoxybenzyl)amino)-6-formyl-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carbaldehyde

Step 1: Preparation of 4-(bis(4-methoxybenzyl)amino)-5-(5-cyclopropyl-4-(hydroxymethyl)isoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde. A solution of 4-(bis(4-methoxybenzyl)amino)-5-(4-(((tert-butyldimethylsilyl)oxy)methyl)-5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P9**; 170.2 mg, 0.2446 mmol) in THF (2.446 mL) was treated with TBAF (1.0 M in THF) (269  $\mu$ L, 0.27 mmol). After stirring for 20 min at RT, the reaction mixture was quenched with saturated  $\text{NH}_4\text{Cl}_{(aq)}$  and extracted with EtOAc. The combined organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound (141.4 mg, 99% yield). MS (apci)  $m/z$  = 582.3 (M+H).

Step 2: Preparation of 3-(4-(bis(4-methoxybenzyl)amino)-6-formyl-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carbaldehyde. A solution of 4-(bis(4-methoxybenzyl)amino)-5-(5-cyclopropyl-4-(hydroxymethyl)isoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (141.4 mg, 0.2431 mmol) in DCM (2.431 mL) was treated with DMP (154.7 mg, 0.3646 mmol) and stirred for 30 min at RT. The reaction mixture then was washed with saturated  $\text{NaHCO}_{3(aq)}$ . The organic extracts were passed through a phase separator column, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-70% EtOAc in hexanes as the gradient eluent) to afford the title compound (106.5 mg, 76% yield). MS (apci)  $m/z$  = 580.2 (M+H).

**Intermediate P11**






(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol

Step 1: Preparation of ethyl 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylate. A solution of ethyl 4-chloro-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylate (1.56 g, 6.91 mmol) in THF (70 mL) was treated with iPrOH (794  $\mu$ L, 10.4 mmol) and  $\text{PPh}_3$  (2.72 g, 10.4 mmol), and the resulting mixture was then cooled to 0  $^\circ\text{C}$ . The cold (0  $^\circ\text{C}$ ) mixture was treated dropwise with DIAD (2.04 mL, 10.4 mmol), after which it was stirred for 16 h at RT. The resulting mixture was concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-60% hexanes-EtOAc as the gradient eluent) to afford the title compound (1.76 g, 95% yield). MS (apci)  $m/z$  = 268.1 (M+H).

Step 2: Preparation of (4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol. A cold (-78  $^\circ\text{C}$ ) solution of ethyl 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylate (1.76 g, 6.57 mmol) in DCM (66 mL) was treated with DIBAL-H (25 wt.% in toluene; 13.3 mL, 19.7 mmol), and stirred for 30 min at -78  $^\circ\text{C}$ . The cold (-78  $^\circ\text{C}$ ) reaction mixture was quenched with the addition of an aqueous solution of sodium potassium tartrate (1.0 M, ~50 mL). The quenched mixture then was stirred for 60 h at RT, before diluting with additional water. The biphasic mixture was extracted with DCM (2x). The organic extracts were combined, dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-60% hexanes-Acetone as the gradient eluent) to afford the title product (1.43 g, 96% yield). MS (apci)  $m/z$  = 226.1 (M+H).


**Intermediate P12**



4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde

A solution of (4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (**Intermediate P11**) (3.0 g, 13 mmol) in DCM (66 mL) was treated with 1,1,1-tris(acetyloxy)-1,1-dihydro-1,2-benziodoxol-3-(1H)-one (Dess-Martin periodinane) (8.5 g, 20 mmol) at RT, and the resulting mixture was stirred for 1 h. The reaction was then washed with sat.  $\text{NaHCO}_3$  (3x) and water (1x), dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to afford the title compound as white solid, which was directly used in the next step without further purifications, assuming quantitative yield. MS (apci)  $m/z$  = 224.1 (M+H).

**Intermediate P13**




6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine

Step 1: Preparation of (4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol. A solution of (4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (**Intermediate P11**) (1.43 g, 6.34 mmol) in DMF (25 mL) was treated with NIS (1.57 g, 6.97 mmol), and the mixture was then stirred for 16 h at RT. The resulting mixture was diluted with water and extracted with EtOAc (2x). The combined organic extracts were washed with water (3x) and brine (1x), and then dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo* to afford the title compound (2.12 g, 95% yield). MS (apci) *m/z* = 352.0 (M+H).

Step 2: Preparation of 6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine. A solution of (4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (2.12 g, 6.03 mmol) in DMF (30 mL) was treated with TBDMSCl (1.36 g, 9.05 mmol) and imidazole (821 mg, 12.1 mmol), and then stirred for 1 h at RT. The resulting mixture was quenched with water, and extracted with EtOAc (2x). The combined organic extracts were washed with water (3x) and brine (1x), then dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-50% hexanes-EtOAc as the gradient eluent) to afford the title compound (2.58 g, 92% yield). MS (apci) *m/z* = 466 (M+H).


**Intermediate P14**



6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine

A solution of 6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine (**Intermediate P13**; 1.49 g, 3.20 mmol) in DMSO (13 mL) was treated with 2,4-dimethoxybenzylamine (1.201 mL, 7.996 mmol), and the resulting mixture was then stirred for 16 h at 60 °C. After cooling to RT, the reaction mixture was diluted with water, and extracted with EtOAc (2x). The combined organic extracts were washed with water (3x) and brine (1x), then dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-75% MTBE in hexanes) to afford the title compound (1.88 g, 99% yield). MS (apci) *m/z* = 597.1 (M+H).


**Intermediate P15**



6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-7-isopropyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine

To a mixture of 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (0.30 g, 0.50 mmol) (**Intermediate P14**) and Pd(PPh<sub>3</sub>)<sub>4</sub> (58 mg, 0.05 mmol) in 1,4-dioxane (2.5 mL) was added Et<sub>3</sub>N (0.1 mL, 0.75 mmol) under nitrogen. The mixture was degassed and purged with N<sub>2</sub> before 4,4,5,5-tetramethyl-1,3,2-dioxaborolane (95 µL, 0.65 mmol) was introduced. The resulting mixture was heated at 90 °C for 3 h then cooled to rt, after which it was concentrated, and the resulting residue was purified by silica chromatography (0-50% EtOAc in hexanes) to afford the title product as light yellowish solid (0.24 g, 80%). MS (apci) m/z = 597.4 (M+H).

**Intermediate P16**



Benzyl 3-(4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate

Step 1: Preparation of benzyl (E)-3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)acrylate. To a solution of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P12**) (2.1 g, 9.34 mmol) in DCM (47 mL) was added benzyl 2-(triphenyl-λ<sup>5</sup>-phosphanylidene)acetate (4.22 g, 10.3 mmol) at RT and stirred for 30 min. The reaction was partitioned between DCM and water. After phase-separation, the organic layer was washed with brine then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated


and purified by silica chromatography (10-90% EtOAc in hexanes) to afford the title compound (1.6 g, 47%). MS (apci)  $m/z$  = 356.1 (M+H).

**Step 2: Preparation of benzyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate.** To a solution of  $\text{Cu}(\text{OAc})_2$  (0.080 g, 0.44 mmol) and Xantphos (0.51 g, 0.88 mmol) in THF (20 mL) was added phenylsilane (5.4 mL, 43.9 mmol) at RT. The resulting mixture was stirred for 5 min. Then a solution of benzyl (E)-3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)acrylate (1.56 g, 4.39 mmol) in THF (10 mL) was introduced, and stirring was continued for another 1 h. The reaction was partitioned between EtOAc and water. After phase-separation, the organic layer was washed with brine then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (10-90% EtOAc in hexanes) to afford the title compound (1.24 g, 79%). MS (apci)  $m/z$  = 358.1 (M+H).

**Step 3: Preparation of benzyl 3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate.** A mixture of benzyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate (1.24 g, 3.47 mmol) and (2,4-dimethoxyphenyl)methanamine (1.6 mL, 10.4 mmol) in DMSO (23 mL) was heated at 90 °C overnight. After cooled to RT, the reaction was partitioned between EtOAc and water. After phase-separation, the organic layer was washed with brine then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (10-90% EtOAc in hexanes) to afford the title compound (0.50 g, 30%). MS (apci)  $m/z$  = 489.2 (M+H).

**Step 4: Preparation of benzyl 3-(4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate.** To a solution of benzyl 3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate (148 mg, 0.30 mmol) in DMF (3 mL) was added NIS (72 mg, 0.32 mmol) at RT and stirred for 2 h. The reaction was partitioned between EtOAc and water. After phase-separation, the organic layer was washed with brine then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (10-90% EtOAc in hexanes) to afford the title product (108 mg, 58%). MS (apci)  $m/z$  = 615.2 (M+H).

#### Intermediate P17



#### 3-(4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole

**Step 1: Preparation of 4-chloro-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde.** Under an atmosphere of  $\text{N}_2$ , n-BuLi (2.5 M, hexanes) (9.2 mL, 23.0 mmol) was dropwise added to a suspension of 5-bromo-4-chloro-7H-pyrrolo[2,3-d]pyrimidine (2.32 g, 9.98 mmol) in THF (50 mL) cooled to -78 °C. The reaction was stirred for 1 h before DMF (1.00 mL, 13.0 mmol) was introduced dropwise. After stirring at -78 °C for another 30 min, the reaction was slowly warmed to RT, then quenched with saturated  $\text{NH}_4\text{Cl}_{(\text{aq})}$  and


water. The biphasic mixture was concentrated *in vacuo* to remove the organic solvents, and the resulting suspension was filtered. The solids collected were rinsed with water and hexanes, and then dried *in vacuo* to afford the title compound (1.52 g, 84% yield).

**Step 2: Preparation of 4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde.** A solution of 4-chloro-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (400 mg, 2.2 mmol) in DMF (11 mL) was treated with 60 wt.% NaH (110 mg, 2.75 mmol) and stirred for 15 min at RT. The resulting yellow solution was treated dropwise with benzenesulfonyl chloride (340  $\mu$ L, 2.6 mmol). After stirring for 20 min at RT, the reaction mixture was quenched with ice (2 g) and water (20 mL). The quenched mixture was stirred at RT for 3 min, and then the resulting suspension was filtered. The solids collected were rinsed with water (10 mL) and hexanes (10 mL), and then air dried to afford the title compound (650 mg, 92% yield).

**Step 3: Preparation of 4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime.** A mixture of 4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (650 mg, 2.0 mmol),  $\text{NH}_2\text{OH}\cdot\text{HCl}$  (170 mg, 2.4 mmol) and NaOAc (200 mg, 2.4 mmol) in EtOH (20 mL) was stirred for 20 h at RT. The resulting mixture was diluted with DCM (20 mL), and the resulting suspension was filtered through a pad of Celite®. The filtrate was concentrated *in vacuo* to afford the title compound (680 mg, quantitative yield).

**Step 4: Preparation of 3-(4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole.** A solution of 4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime (170 mg, 0.505 mmol), NCS (80.9 mg, 0.606 mmol),  $\text{KHCO}_3$  (92 mg, 1.52 mmol) and cyclopropylacetylene (128  $\mu$ L, 1.51 mmol) in DMF (5.048 mL) was stirred for 45 min at 50 °C. After cooling to RT, the reaction mixture was diluted with EtOAc and then quenched with water. The resulting suspension was filtered, and the solids were rinsed with EtOAc and water. The biphasic filtrate was separated, and the organic extracts were concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-30% EtOAc in hexanes) to afford the title compound (101 mg, 50% yield). MS (apci)  $m/z$  = 401.0 (M+H).

#### Intermediate P18




#### 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole

To a solution of tetrabutylammonium fluoride (TBAF) in THF (1 M) (14.6 mL, 14.6 mmol) was added 3-(4-chloro-7-(phenylsulfonyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (**Intermediate P17**) (2.93 g, 7.31 mmol). The mixture was heated at 60 °C for 1 h, then cooled to RT, diluted with EtOAc (50 mL) and washed with water (2  $\times$  30 mL). After phase-separation, the organic layer was dried ( $\text{Na}_2\text{SO}_4$ ),

filtered and concentrated to give the title product as beige solid, assuming quantitative yield. MS (apci neg)  $m/z$  = 259.1 (M-H).

#### Intermediate P19



3-(4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole

#### Method A:

**Step 1: Preparation of 4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde.** To a suspension of 4-chloro-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (1.82 g, 10.0 mmol) in DMF (33 mL) was added NaH (60 wt%) (0.44 g, 11.0 mmol) at 0 °C. After 20 min stirring 2-(trimethylsilyl)ethoxymethyl chloride (SEM-Cl) (2.13 mL, 12.0 mmol) was introduced. The reaction was allowed to warm to RT and stirred for 3 h. This was followed by the addition of more NaH (0.24 g, 6 mmol) and SEM-Cl (0.84 g, 5 mmol). After another 30 min of stirring, the reaction mixture was quenched with ice and NH<sub>4</sub>Cl (sat.), and then extracted with EtOAc. The combined organic extracts were washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and purified by silica chromatography (0-25% EtOAc in hexanes) to give the title product as yellowish solid (1.33 g, 43%). MS (apci)  $m/z$  = 312.1 (M+H).

**Step 2: Preparation of 4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime.** A mixture of 4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (1.33 g, 4.26 mmol), hydroxylamine hydrochloride (0.33 g, 4.7 mmol) and NaOAc (0.38 g, 4.7 mmol) in EtOH (21 mL) was stirred at 40 °C for 1 h, after which it was concentrated. The residue thus obtained was taken up in DCM and filtered through a short pad of celite. The filtrate was concentrated to afford the title product as white solid, assuming quantitative yield. LCMS  $m/z$  = 327.1 (M+H).


**Step 3: Preparation of 3-(4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole.** A mixture of 4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime (1.37 g, 4.2 mmol), NCS (0.617 g, 4.62 mmol), KHCO<sub>3</sub> (0.631 g, 6.30 mmol) and cyclopropylacetylene (1.07 mL, 12.6 mmol) in DMF (21 mL) was stirred at RT for 5 d. The reaction was diluted with water and extracted with EtOAc (3x). The combined organic extracts were washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and purified by silica chromatography (0-30% EtOAc in hexanes) to afford the title product (0.83 g, 50%). MS (apci)  $m/z$  = 391.2 (M+H).

#### Method B:

To a solution of 3-(4-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (**Intermediate P18**) (5.87 g, 22.52 mmol) in DMF (113 mL) was added NaH (60 wt%) (1.08 g, 27.0 mmol) followed by (2-(chloromethoxy)ethyl)trimethylsilane (SEM-Cl) (4.4 mL, 24.8 mmol) at RT. After 30 min stirring, the

reaction was diluted with water and extracted with EtOAc (2x). The combined organic extracts were washed with brine, dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (0-50% EtOAc in hexanes) to afford the title product as clear brownish oil (2.7 g, 31%). MS (apci)  $m/z$  = 391.1 (M+H).

#### Intermediate P20




#### 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole

Step 1: Preparation of 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole. To a solution of 3-(4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (**Intermediate P19**) (1.96 g, 5.01 mmol) in THF (25 mL) was added n-BuLi (2.5 M THF) (3.0 mL, 7.5 mmol) at  $-78^\circ\text{C}$ . After 10 min stirring, iodomethane (0.63 mL, 10.0 mmol) was introduced, and the resulting solution was stirred at  $-78^\circ\text{C}$  for 30 min before warming up to  $-20^\circ\text{C}$ . The reaction was quenched with  $\text{NH}_4\text{Cl}$  (sat.) and extracted with EtOAc. The combined organic extracts were dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to give the title product as brownish oil, assuming quantitative yield. MS (apci)  $m/z$  = 405.1 (M+H).

Step 2: Preparation of 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole. A solution of 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (2.0 g, 5.0 mmol) in MeCN (25 mL) and treated with NIS (1.69 g, 7.52 mmol) and TFA (1.16 mL, 15.0 mmol). The resulting mixture was stirred at RT for 3 h, after which it was quenched with 10%  $\text{Na}_2\text{S}_2\text{O}_3$  (aq) (5 mL) and sat.  $\text{NaHCO}_3$  (20 mL). This was followed by extraction with EtOAc (3x). The combined organic extracts were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (0-30% EtOAc in hexanes) to afford the title product as yellowish oil (1.5 g, 56%). LCMS  $m/z$  = 531.0 (M+H).


#### Intermediate P21



#### Ethyl 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate

To a solution of 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole (**Intermediate P20**) (1.50 g, 2.83 mmol) in THF (14 mL) was dropwise added isopropylmagnesium(II) lithium chloride (1.3 M) (2.8 mL, 3.67 mmol) at 0 °C. After 10 min stirring ethyl chloroformate (0.54 mL, 5.65 mmol) was introduced, and the resulting mixture was warmed to RT and stirred for 1 h. The reaction was quenched with NH<sub>4</sub>Cl (sat.) and extracted with EtOAc. The combined organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and purified by silica chromatography (0-30% EtOAc in hexanes) to afford the title product along with a side product 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (ca. 25%) as yellowish oil. The title product was used directly in the next step without further purifications (1.2 g, 90%). MS (apci) m/z = 477.2 (M+H).


#### Intermediate P22



#### Ethyl 3-(4-chloro-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate

A solution of ethyl 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P21**) (150 mg, 0.31 mmol) in TFA (0.5 mL) was stirred at RT for 16 h. The mixture was then concentrated and the resulting residue purified by reverse phase chromatography (0-95% MeCN in water with 0.1% TFA). The combined product fractions were partitioned between sat. NaHCO<sub>3</sub> and DCM. After phase-separation the organic layer was concentrated to afford the title product as off-white solid (82 mg, 75%). MS (apci) m/z = 347.0 (M+H).

#### Intermediate P23




#### 11-amino-3-cyclopropyl-7-((2-(trimethylsilyl)ethoxy)methyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one

A mixture of ethyl 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P21**) (189 mg, 0.40 mmol), NBS (84.6 mg, 0.48 mmol) and AIBN (9.8 mg, 0.059 mmol) in 1,2-DCE (4 mL) was flushed with nitrogen, sealed in a pressure vessel and stirred at 80 °C for 2 h. Upon cooling to RT, the reaction was concentrated. The residue thus obtained was then taken up in 1,4-dioxane (4 mL) and NH<sub>4</sub>OH (28-30%) (2 mL). The mixture was re-sealed in the vessel and stirred at 90 °C for 16 h, then cooled to RT and concentrated. The crude residue



was subjected to reverse phase chromatography (0-95% MeCN in water with 0.1% TFA). The combined product fractions were neutralized with 10% Na<sub>2</sub>CO<sub>3</sub> (aq) and extracted with DCM. The organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to afford the title product as off-white solid (15 mg, 8.8%). MS (apci) m/z = 427.1 (M+H).

#### Intermediate P24




#### (4-((2,4-dimethoxybenzyl)amino)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol

Step 1: Preparation of 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine. A mixture of 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (**Intermediate P14**) (218 mg, 0.36 mmol), (PPh<sub>3</sub>)<sub>2</sub>Pd(II)Cl<sub>2</sub> (51.3 mg, 0.073 mmol), 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole (178 mg, 0.73 mmol) and Na<sub>2</sub>CO<sub>3</sub> (2 N aq) (365 µL, 0.73 mmol) in dioxane (5.2 mL) was sparged with Ar and stirred at 80 °C overnight. After cooled to RT, the reaction was partitioned between DCM and water. The aqueous was phase extracted with 4:1 DCM:IPA (3x). The combined organic extracts were concentrated and purified using silica chromatography (0-100% EtOAc in hexanes) to give the title compound (164 mg, 76% yield). MS (apci) m/z = 586.3 (M+H).

Step 2: Preparation of (4-((2,4-dimethoxybenzyl)amino)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol. To a solution of 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (164 mg, 0.28 mmol) in THF (2.8 mL) was added acetic acid (32 µL, 0.56 mmol) at 0°C, followed by tetrabutylammonium fluoride (TBAF) (1 M, THF; 308 µL, 0.31 mmol). The reaction was allowed to warm to RT and stirred for 3 h before it was concentrated and purified using silica chromatography (0-100% EtOAc in hexanes) to give the title compound (109 mg, 82% yield). MS (apci) m/z = 472.2 (M+H).

#### Intermediate P25




6-(aminomethyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine

Step 1: Preparation of 6-(azidomethyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine. To a solution of 4-((2,4-dimethoxybenzyl)amino)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (**Intermediate P24**) (99 mg, 0.21 mmol) in toluene (2.1 mL) was added 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) (38  $\mu$ L, 0.25 mmol) at 0 °C, followed by diphenylphosphoryl azide (DPPA) (54  $\mu$ L, 0.25 mmol). The reaction was allowed to warm to rt. After stirring overnight, the reaction mixture was diluted with EtOAc and washed with water and brine, then dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated. The residue was purified using silica chromatography (0-100% EtOAc in hexanes) to afford the title compound (71 mg, 68% yield). MS (apci)  $m/z$  = 497.2 (M+H).

Step 2: Preparation of 6-(aminomethyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine. To a solution of 6-(azidomethyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (70 mg, 0.14 mmol) in 1:1 THF:water (1.4 mL) was added  $\text{PPh}_3$  (74 mg, 0.28 mmol) at RT and stirred for 16 h. The reaction was diluted with 4:1 DCM:IPA and washed with sat.  $\text{NaHCO}_3$ . The combined organic phase was then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (0-100% EtOAc in hexanes) to afford the title product (51 mg, 77%). MS (apci)  $m/z$  = 471.2 (M+H).

**Intermediate P26**




Methyl 2-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate

A mixture of 6-methoxycarbonyl-1H-indole-2-boronic acid pinacol ester (852 mg, 2.83 mmol),  $(\text{PPh}_3)_2\text{Pd(II)Cl}_2$  (99.3 mg, 0.141 mmol), 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-

dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (**Intermediate P14**) (422 mg, 0.707 mmol) and Na<sub>2</sub>CO<sub>3</sub> (2 N aq) (707  $\mu$ L, 1.41 mmol) in dioxane (10 mL) was sparged with Ar and heated at 80 °C for 2 h. The reaction was partitioned between DCM and water then extracted with 4:1 DCM:IPA (3x). The combined organic extracts were concentrated and purified using silica chromatography (0-40% EtOAc in hexanes) to yield the title product (415 mg, 91%). MS (apci) m/z = 644.3 (M+H).


#### Intermediate P27



#### 2-(4-((2,4-Dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylic acid

To a solution of methyl 2-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate (**Intermediate P26**) (339 mg, 0.53 mmol) in THF (10 mL) and MeOH (1 mL) was added LiOH (2 M aq) (1.3 mL, 2.6 mmol) at RT. The reaction mixture was stirred overnight. Additional LiOH (2 M aq) (1.3 mL, 2.6 mmol) was then introduced and the reaction was stirred for another 48 h before it was acidified with HCl (1 M aq) to pH ~5. It was then diluted with water (3 mL) and extracted with DCM (3 x 5 mL). The combined organic extracts were concentrated and purified with silica chromatography (0-10% MeOH in DCM) to yield the title product (140 mg, 50%). MS (apci) m/z = 516.2 (M+H).


#### Intermediate P28



#### Methyl 2-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate

The title product was prepared in a similar fashion as described for the synthesis of 4-((2,4-dimethoxybenzyl)amino)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (**Intermediate P24**), replacing 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole with methyl 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-6-carboxylate during step 1 instead. MS (apci)  $m/z = 530.2$  (M+H).


#### Intermediate P29



Methyl 2-(6-(aminomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate

The title product was prepared in a similar fashion as described for the synthesis of 6-(aminomethyl)-N-(2,4-dimethoxybenzyl)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (**Intermediate P25**), replacing 4-((2,4-dimethoxybenzyl)amino)-5-(1H-indol-2-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (**Intermediate P24**) with methyl 2-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate (**Intermediate P28**) during step 1 instead. MS (apci)  $m/z = 529.2$  (M+H).

#### Intermediate P30



1-((2,4-dimethoxybenzyl)amino)-5-isopropyl-8-oxo-5,6,7,8-tetrahydropyrimido[5'',4'':4',5']pyrrolo[3',2':3,4]azepino[1,2-a]indole-11-carboxylic acid


Step 1: Preparation of methyl 2-(6-(3-(benzyloxy)-3-oxopropyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate. A mixture of methyl 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indole-6-carboxylate (0.318 g, 1.05 mmol), benzyl 3-(4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate (**Intermediate**

**P16** (0.108 g, 0.176 mmol),  $\text{PdCl}_2(\text{PPh}_3)_2$  (0.0123 g, 0.0176 mmol) and  $\text{Na}_2\text{CO}_3$  (2 N, aq) (0.27 mL, 0.54 mmol) in dioxane (0.88 mL) was heated to 90 °C overnight. The reaction mixture was then filtered, concentrated and purified by silica chromatography (1-15% MeOH in DCM) to give the title compound (0.116 g, 99%). MS (apci)  $m/z$  = 662.3 (M+H).

Step 2: Preparation of 2-(6-(2-carboxyethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylic acid. A mixture of methyl 2-(6-(3-(benzyloxy)-3-oxopropyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylate (0.10 g, 0.15 mmol) and LiOH (2 N, aq) (0.23 mL, 0.46 mmol) in MeOH (0.76 mL) was refluxed overnight. The mixture was concentrated and purified by silica chromatography (1-15% MeOH in DCM with 1% AcOH) to give the title compound (0.058 g, 69%). MS (apci)  $m/z$  = 558.2 (M+H).

Step 3: Preparation of 1-((2,4-dimethoxybenzyl)amino)-5-isopropyl-8-oxo-5,6,7,8-tetrahydropyrimido[5'',4':4',5']pyrrolo[3',2':3,4]azepino[1,2-a]indole-11-carboxylic acid. A mixture of 2-(6-(2-carboxyethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-1H-indole-6-carboxylic acid (0.034 g, 0.061 mmol), BOC-anhydride (0.050 mL, 0.213 mmol), DMAP (0.022 g, 0.18 mmol) and 2,6-lutidine (0.021 mL, 0.183 mmol) in MeCN (0.61 mL) was stirred at RT for 2 h. The reaction was partitioned between DCM and water. After phase-separation, the organic layer was washed with brine then dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to afford the title compound, which was used directly in the next step without further purifications, assuming quantitative yield. MS (apci)  $m/z$  = 540.2 (M+H).

#### Intermediate P31



#### 12-((2,4-Dimethoxybenzyl)amino)-8-isopropyl-5-oxo-5,6,7,8-tetrahydrobenzo[c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepine-3-carboxylic acid

Step 1: Preparation of dimethyl 4-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate. A mixture of 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-7-isopropyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-7H-pyrrolo[2,3-d]pyrimidin-4-amine (**Intermediate P15**) (159 mg, 0.27 mmol), dimethyl 4-bromoisophthalate (109 mg, 0.40 mmol),  $\text{Pd}(\text{PPh}_3)_4$  (31 mg, 0.027 mmol), and  $\text{K}_2\text{CO}_3$  (111 mg, 0.80 mmol) in 4:1 dioxane:water (2.7 mL) was sparged with argon, sealed in a pressure vessel and stirred at 80 °C for 16 h. After cooled to RT, the reaction was diluted with 4:1 DCM:IPA and washed with water. The organic layer was then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica chromatography

(5-95% acetone in hexanes) to afford the title compound (125 mg, 70%). MS (apci)  $m/z$  = 663.3 (M+H).

**Step 2: Preparation of dimethyl 4-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate.** To a solution of dimethyl 4-(6-(((tert-butyl)dimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate (125 mg, 0.18 mmol) in THF (1.9 mL) was added acetic acid (27  $\mu$ L, 0.47 mmol) followed by TBAF (1 M THF) (470  $\mu$ L, 0.47 mmol) at RT and stirred for 48 h. The reaction was concentrated and purified by silica chromatography (5-95% acetone in hexanes) to afford the title compound (92 mg, 89%). MS (apci)  $m/z$  = 549.2 (M+H).


**Step 3: Preparation of dimethyl 4-(6-(azidomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate.** To a solution of dimethyl 4-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate (92 mg, 0.17 mmol) in toluene (1.7 mL) was added DBU (50  $\mu$ L, 0.34 mmol) followed by DPPA (72  $\mu$ L, 0.34 mmol) at 0 °C. The reaction was warmed to RT and stirred for 2 h before it was diluted with EtOAc and washed with water and brine, then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica chromatography (5-95% acetone in hexanes) to give the title compound (75 mg, 78%). MS (apci)  $m/z$  = 574.2 (M+H).

**Step 4: Preparation of dimethyl 4-(6-(aminomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate.** To a solution of dimethyl 4-(6-(azidomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate (75 mg, 0.13 mmol) in 1:1 THF:water (1.3 mL) was added  $\text{PPh}_3$  (68 mg, 0.26 mmol) at RT and stirred for 16 h. The reaction was diluted with 4:1 DCM:IPA and washed with sat.  $\text{NaHCO}_3$ , then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by reverse phase chromatography (5-95% MeCN in water with 0.1% TFA). The combined product fractions were partitioned between 4:1 DCM:IPA and sat.  $\text{NaHCO}_3$ , then dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated to give the title compound (52 mg, 73%). MS (apci)  $m/z$  = 548.3 (M+H).

**Step 5: Preparation of methyl 12-((2,4-dimethoxybenzyl)amino)-8-isopropyl-5-oxo-5,6,7,8-tetrahydrobenzo[c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepine-3-carboxylate.** A solution of dimethyl 4-(6-(aminomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isophthalate (52 mg, 0.095 mmol) in dioxane (1.9 mL) was stirred at 90 °C for 24 h. After cooled to RT, the reaction was concentrated and purified by silica chromatography (5-95% acetone in DCM) to afford the title compound, which was directly used in the next step assuming quantitative yield. MS (apci)  $m/z$  = 516.2 (M+H).

**Step 6: Preparation of 12-((2,4-dimethoxybenzyl)amino)-8-isopropyl-5-oxo-5,6,7,8-tetrahydrobenzo[c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepine-3-carboxylic acid.** To a solution of methyl 12-((2,4-dimethoxybenzyl)amino)-8-isopropyl-5-oxo-5,6,7,8-tetrahydrobenzo[c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepine-3-carboxylate (49 mg, 0.095 mmol) in THF (1.9 mL) was added KOH (2 N aq) (475  $\mu$ L, 0.95 mmol) at RT and stirred for 16 h. The reaction was then diluted with water and acidified to pH~5 using 1.0 M HCl, followed by extracting with 4:1 DCM:IPA (2x). The combined organics were dried over  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated to afford the title product (33 mg, 70%). MS (apci)  $m/z$  = 502.2 (M+H).

#### Intermediate P32



Tert-butyl ((4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methyl)carbamate


**Step 1: Preparation of (4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol.** A mixture of 6-(((tert-butyl dimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (**Intermediate P14**) (1.30 g, 2.18 mmol), acetic acid (0.125 mL, 2.18 mmol) and TBAF (1 M THF) (2.40 mL, 2.40 mmol) in THF (14.5 mL) was stirred at rt for 2 h. The reaction was diluted with water and extracted with EtOAc. The combined organic extracts were washed with brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated, and purified by silica chromatography (1-10% MeOH in DCM) to afford the title compound (1.05 g, 99%). MS (apci) m/z = 483.1 (M+H).

**Step 2: Preparation of 6-(azidomethyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine.** To a solution of (4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methanol (1.05 g, 2.18 mmol) in toluene (14.5 mL) cooled to 0 °C was added diphenyl phosphorazidate (0.94 mL, 4.35 mmol), followed by 2,3,4,6,7,8,9,10-octahydropyrimido[1,2-a]azepine (0.65 mL, 4.35 mmol). The reaction was warmed up to rt and stirred overnight. It was then diluted with water and extracted with DCM. The combined organic extracts were washed with brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated, and purified by silica chromatography (1-10% MeOH in DCM) to afford the title compound (1.09 g, 99%). MS (apci) m/z = 508.1 (M+H).

**Step 3: Preparation of 6-(aminomethyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine.** A mixture of 6-(azidomethyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (1.09 g, 2.160 mmol) and PPh<sub>3</sub> (1.13 g, 4.32 mmol) in THF (8.6 mL) and water (8.6 mL) was stirred at rt overnight. It was then diluted with water and extracted with DCM. The combined organic extracts were washed with brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated, and purified by silica chromatography (1-10% MeOH in DCM) to afford the title compound (0.44 g, 42%). MS (apci) m/z = 482.1 (M+H).

**Step 4: Preparation of tert-butyl ((4-((2,4-dimethoxybenzyl)amino)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)methyl)carbamate.** To a solution of 6-(aminomethyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (56 mg, 0.12 mmol) in THF (2.3 mL) was added di-tert-butyl dicarbonate (27 µL, 0.12 mmol) at rt and stirred for 2 h. The reaction was then diluted with water (20 mL) and extracted with DCM (3 × 20 mL). The combined organic extracts were washed with brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated, and purified by silica chromatography (0-100% EtOAc in hexanes) to afford the title compound (59 mg, 88%). MS (apci) m/z = 582.2 (M+H).

**Intermediate P33**



6-(3-((tert-butyldimethylsilyl)oxy)propyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine

Step 1: Preparation of ethyl (E)-3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)acrylate. To a solution of 4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P12**) (14.9 g, 32.0 mmol) in DCM (300 mL) was added carbethoxymethylene triphenylphosphorane (13.4 g, 38.4 mmol) at rt and stirred for 16 h. The reaction was concentrated and purified by silica chromatography (5 to 75% acetone in hexanes) to afford the title compound (8 g, 85%). MS (apci)  $m/z$  = 294.1 (M+H).

Step 2: Preparation of ethyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate. To a solution of  $\text{Cu}(\text{OAc})_2$  (0.17 g, 0.95 mmol) in THF (50 mL) was added (9,9-dimethyl-9H-xanthene-4,5-diyl)bis(diphenylphosphane) (1.10 g, 1.90 mmol) followed by phenylsilane (5.85 mL, 47.5 mmol). After 15 min stirring at rt, a solution of ethyl (E)-3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)acrylate (2.79 g, 9.50 mmol) in THF (50 mL) was introduced, and stirring continued at rt for 3 h. The reaction was cooled to 0 °C and treated slowly with sat.  $\text{NaHCO}_3$ . It was diluted with water and extracted with DCM (2x). The combined organic extracts was dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica chromatography (5 to 75% EtOAc in hexanes) to afford the title compound (2.7 g, 97%). MS (apci)  $m/z$  = 296.1 (M+H).

Step 3: Preparation of ethyl 3-(4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate. To a solution of ethyl 3-(4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate (2.46 g, 8.32 mmol) in DMF (55 mL) was added N-iodosuccinimide (1.96 g, 8.73 mmol) at rt and stirred for 60 h. The reaction was diluted with water and extracted with EtOAc (2x). The combined organic extracts were washed with water and brine, then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica chromatography (5 to 75% EtOAc in hexanes) to afford the title compound (3.05 g, 87%). MS (apci)  $m/z$  = 422.0 (M+H).


Step 4: Preparation of 3-(4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propan-1-ol. To a solution of ethyl 3-(4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoate (5.06 g, 12.0 mmol) in THF (60 mL) cooled to -78 °C was added DIBAL-H (25 wt%) (24.2 mL, 36.0 mmol). After stirred at -78 °C for 30 min, the reaction was quenched with water and allowed to warm to rt and stirred overnight. After phase-separation, the aqueous phase was extracted with DCM (2x). The combined organic extracts was washed with water, then dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to afford the title compound as white solid (2.9 g, 64%). MS (apci)  $m/z$  = 380.0 (M+H).



Step 5: Preparation of 6-(3-((tert-butyldimethylsilyl)oxy)propyl)-4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine. To a solution of 3-(4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propan-1-ol (2.9 g, 7.6 mmol) in DMF (38 mL) were added imidazole (1.0 g, 15 mmol) and tert-butyldimethylsilyl chloride (1.7 g, 11 mmol). After stirred at rt for 1 h, the reaction was diluted with EtOAc, washed with water and brine, then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica chromatography (0 to 50% EtOAc in hexanes) to afford the title compound as clear oil (3.6 g, 95%). MS (apci)  $m/z = 494.1$  (M+H).

Step 6: Preparation of 6-(3-((tert-butyldimethylsilyl)oxy)propyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine. To a solution of 6-(3-((tert-butyldimethylsilyl)oxy)propyl)-4-chloro-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine (1.53 g, 3.1 mmol) in DMSO (15.5 mL) was added (2,4-dimethoxyphenyl)methanamine (1.16 mL, 7.74 mmol). The reaction was heated to 60 °C and stirred overnight. After cooled to rt, the reaction was diluted with EtOAc, washed with water and brine, then dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated, and purified by silica chromatography (0-100% MTBE in hexanes) to afford the title product as clear oil (1.8 g, 92%). MS (apci)  $m/z = 625.2$  (M+H).


#### Intermediate P34



#### 1-((2,4-dimethoxybenzyl)amino)-5-isopropyl-5,6,7,8-tetrahydropyrimido[5'',4'':4',5']pyrrolo[3',2':3,4]azepino[1,2-a]indole-11-carboxylic acid

To a solution of methyl 1-((2,4-dimethoxybenzyl)amino)-5-isopropyl-5,6,7,8-tetrahydropyrimido[5'',4'':4',5']pyrrolo[3',2':3,4]azepino[1,2-a]indole-11-carboxylate (**Example 86**, Step 3) (458 mg, 0.85 mmol) in THF (17 mL) and MeOH (17 mL) was added LiOH (2 M aq) (4.2 mL, 8.4 mmol) at ambient temperature and stirred overnight. The reaction was neutralized with sat.  $\text{NH}_4\text{Cl}$  (aq) (20 mL) and extracted with DCM (3 x 50 mL) through a Phase Separator frit. The combined organic extracts were concentrated and purified using silica chromatography (0-100% EtOAc in hexanes followed by 0-10% MeOH in EtOAc) to yield the title product (319 mg, 72%). MS (apci)  $m/z = 526.2$  (M+H).

#### Intermediate P35




11-chloro-3-cyclopropyl-7-((2-(trimethylsilyl)ethoxy)methyl)-6,7-dihydroisoxazolo[4'',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one

Step 1: Preparation of 1-(3-(6-(bromomethyl)-4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)ethan-1-one. To a solution of 3-(4-chloro-6-methyl-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole (**Intermediate P20**) (2.9 g, 5.46 mmol) in THF (27 mL) cooled to -78 °C was dropwise added isopropylmagnesium chloride lithium chloride complex (1.3 M THF) (5.46 mL, 7.10 mmol), maintaining an internal temperature below -60 °C. After addition, stirring was continued for 20 min before acetic anhydride (0.77 mL, 8.2 mmol) was introduced at -70 °C. The resulting mixture was warmed to rt and quenched with sat. NH<sub>4</sub>Cl (50 mL) and water (50 mL), after which it was extracted with EtOAc (100 mL). The combined organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), and then filtered through a short pad of celite, rinsing the pad with 50% EtOAc in hexanes (30 mL). The filtrate was concentrated, taken up in DCE (27 mL) and treated with AIBN (90 mg, 0.55 mmol) and NBS (1.2 g, 6.74 mmol). The resulting mixture was heated at 80 °C for 90 min. After cooled to rt, the reaction was washed with Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (10% aq), water and brine, then dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and purified with silica chromatography (0-30% EtOAc in hexanes) to give the title compound as white solid (1.38 g, 48%). MS (apci) m/z = 525.1 (M+H).

Step 2: Preparation of 11-chloro-3-cyclopropyl-7-((2-(trimethylsilyl)ethoxy)methyl)-6,7-dihydroisoxazolo[4'',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one. To a solution of LHMDs (1 M THF) (5.90 mL, 5.90 mmol) in THF (22 mL) cooled to -78 °C was dropwise added a solution of 1-(3-(6-(bromomethyl)-4-chloro-7-((2-(trimethylsilyl)ethoxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)ethan-1-one (1.15 g, 2.19 mmol) in THF (8 mL). The resulting mixture was slowly warmed to -40 °C over 1 h, then to -20 °C over 30 min. It was quenched with NH<sub>4</sub>Cl (sat.) and water, then extracted with EtOAc. The combined organic extracts were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, concentrated and purified with silica chromatography (0-40% EtOAc in hexanes) to give the title product as yellowish solid (0.53 g, 54%). MS (apci) m/z = 445.2 (M+H).


**Intermediate P36**



11-chloro-3-cyclopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one

A solution of 11-chloro-3-cyclopropyl-7-((2-(trimethylsilyl)ethoxy)methyl)-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one (**Intermediate P35**) (0.52 g, 1.2 mmol) in DCM (2.9 mL) and TFA (5.8 mL) was stirred at rt for 2 h, then concentrated to near dryness. It was taken up in dioxane and re-concentrated. The residue was dissolved in 1,4-dioxane (5.8 mL), followed by addition of NH<sub>4</sub>OH (28% aq) (2.9 mL). The mixture was concentrated to near dryness and treated with water (15 mL), then vacuum-filtered, rinsing the solid with water and hexanes, giving the title product as yellowish solid after air-drying (0.37 g, 101%). MS (apci) m/z = 315.1 (M+H).


**Intermediate P37**



3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one

A mixture of 11-chloro-3-cyclopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one (**Intermediate P36**) (158 mg, 0.50 mmol), 2,4-dimethoxybenzylamine (302  $\mu$ L, 2.0 mmol) and DMA (2.5 mL) was stirred at 75 °C for 1 h. After cooled to rt, the reaction was quenched with water (10 mL) and citric acid (10% aq) (2 mL) then filtered, rinsing the collected solid with water (2 mL) and hexanes (2 mL). The solid was further triturated with aid of sonication in a mixture of water (10 mL) and citric acid (10% aq) (1 mL). The resulting suspension was filtered, rinsed with water and dried to afford the title product as yellowish solid (190 mg, 85%). MS (apci) m/z = 446.2 (M+H).

**Intermediate P38**



4-Chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde

Step 1: Preparation of 4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine. To a solution of 2-(4,6-dichloropyrimidin-5-yl)acetaldehyde (5.0 g, 26.2 mmol) in 2-methoxyethan-1-ol (26 mL) in a


pressure vessel was added DIEA (22.8 mL, 131 mmol) and 1-methylcyclopropan-1-amine hydrochloride (3.1 g, 28.8 mmol). The reaction was sealed and stirred at 70 °C overnight. After cooled to rt, the reaction was diluted with water and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (5-40% EtOAc in hexanes) to afford the title compound (4.6 g, 84%). MS (apci)  $m/z$  = 208.1 (M+H).

Step 2: Preparation of 4-chloro-5-iodo-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine. To a solution of 4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine (4.6 g, 22.1 mmol) in DMF (55 mL) was added 1-iodopyrrolidine-2,5-dione (5.97 g, 26.5 mmol). The reaction was covered with aluminum foil and stirred at rt for 3 h. It was then diluted with sat. sodium thiosulfate (aq) and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (1-10% MeOH in DCM) to afford the title compound (5.79 g, 78.5%). MS (apci)  $m/z$  = 334.0 (M+H).

Step 3: Preparation of 4-chloro-7-(1-methylcyclopropyl)-5-vinyl-7H-pyrrolo[2,3-d]pyrimidine. A solution of 4-chloro-5-iodo-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine (5.76 g, 17.3 mmol), 4,4,5,5-tetramethyl-2-vinyl-1,3,2-dioxaborolane (3.08 mL, 18.1 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (1.20 g, 1.04 mmol), and potassium phosphate (2 N, aq) (34.5 mL, 69.1 mmol) in 1,4-dioxane (57.6 mL) was sparged with argon and heated under N<sub>2</sub> at 90 °C for 3 h. After cooled to rt, the reaction was diluted with sat. NaHCO<sub>3</sub> (aq) and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (1-5% MeOH in DCM) to afford the title compound (4.0 g, 99%). MS (apci)  $m/z$  = 234.1 (M+H).

Step 4: Preparation of 4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde. To a solution of 4-chloro-7-(1-methylcyclopropyl)-5-vinyl-7H-pyrrolo[2,3-d]pyrimidine (4.0 g, 17.1 mmol) in THF (86 mL) and water (34 mL) cooled to 0 °C was added 4-methylmorpholine N-oxide (2.0 g, 17.1 mmol) and OsO<sub>4</sub> (4%) (10.9 mL, 1.71 mmol). After 1 h stirring at 0 °C, sodium periodate (7.322 g, 34.23 mmol) was introduced, and stirring continued for 2 h. The reaction was then diluted with sat. sodium thiosulfate (aq) and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (1-10% MeOH in DCM) to afford the title product (2.6 g, 65%). MS (apci)  $m/z$  = 236.0 (M+H).

#### Intermediate P39



#### 3-(4-Chloro-6-methyl-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole


Step 1: Preparation of (E)-4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime. To a solution of 4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde (**Intermediate P38**) (2.63 g, 11.16 mmol) in ethanol (37.2 mL) was added hydroxylamine

(50%) (0.72 mL, 11.72 mmol) at 0 °C. The reaction mixture was allowed to warm to rt and stirred for 90 min before it was diluted with water and extracted with DCM. The combined organic extracts were concentrated to afford the title compound, which was used directly in the next step assuming quantitative yield. MS (apci)  $m/z$  = 251.1 (M+H).

**Step 2: Preparation of 3-(4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole.** To a solution of (E)-4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidine-5-carbaldehyde oxime (2.8 g, 11.2 mmol), potassium hydrogen carbonate (1.45 g, 14.5 mmol), and 1-chloropyrrolidine-2,5-dione (1.64 g, 12.3 mmol) in DMF (55.8 mL) was added ethynylcyclopropane (2.84 mL, 33.5 mmol) at rt. The reaction mixture was stirred overnight, after which it was diluted with sat. sodium thiosulfate (aq) and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (1-7% MeOH in DCM) to afford the title compound (3.14 g, 89%). MS (apci)  $m/z$  = 315.1 (M+H).

**Step 3: Preparation of 3-(4-chloro-6-methyl-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole.** To a solution of 3-(4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (3.14 g, 9.98 mmol) in THF (50 mL) cooled to -78 °C was added lithium diisopropylamide (1.96 M, THF) (8.1 mL, 16.0 mmol), followed by dropwise addition of iodomethane (2.48 mL, 39.9 mmol). The reaction was stirred at -78 °C for 20 min and then quenched with sat.  $\text{NH}_4\text{Cl}$  (aq). The reaction was diluted with sat.  $\text{NaHCO}_3$  (aq) and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (5-40% EtOAc in hexanes) to afford the title product (1.99 g, 61%). MS (apci)  $m/z$  = 329.1 (M+H).

#### Intermediate P40



#### 11-Chloro-3-cyclopropyl-7-(1-methylcyclopropyl)-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one

**Step 1: Preparation of 3-(4-chloro-6-methyl-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole.** A solution of 3-(4-chloro-6-methyl-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole (**Intermediate P39**) (1.99 g, 6.05 mmol), 1-iodopyrrolidine-2,5-dione (2.04 g, 9.08 mmol), and 2,2,2-trifluoroacetic acid (1.40 mL, 18.2 mmol) in acetonitrile (30 mL) was stirred for 90 min. The reaction was diluted with 1:1 sat.  $\text{Na}_2\text{S}_2\text{O}_3$  (aq) and sat.  $\text{NaHCO}_3$  (aq), after which it was extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (1-5% MeOH in DCM) to afford the title compound (2.7 g, 98%). MS (apci)  $m/z$  = 455.0 (M+H).


**Step 2: Preparation of 1-(3-(4-chloro-6-methyl-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)ethan-1-one.** To a solution of 3-(4-chloro-6-methyl-7-(1-

methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole (2.3 g, 5.0 mmol) in THF (25 mL) cooled to 0 °C was added isopropylmagnesium(II) lithium chloride (1.2 M THF) (5.5 mL, 6.6 mmol) and acetic anhydride (0.72 mL, 7.6 mmol). The reaction was allowed to warm to rt and stirred for 15 min. It was then quenched with sat.  $\text{NH}_4\text{Cl}$  (aq), then diluted with sat.  $\text{NaHCO}_3$  (aq) and extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (5-100% EtOAc in hexanes) to afford the title compound (1.3 g, 70%). MS (apci)  $m/z$  = 371.1 (M+H).

Step 3: Preparation of 1-(3-(6-(bromomethyl)-4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)ethan-1-one. A solution of 1-(3-(4-chloro-6-methyl-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)ethan-1-one (1.45 g, 3.91 mmol), 1-bromopyrrolidine-2,5-dione (0.84 g, 4.69 mmol), and (E)-2,2'-(diazene-1,2-diyl)-bis(2-methylpropanenitrile) (0.038 g, 0.24 mmol) in 1,2-dichloroethane (20 mL) was heated to 75 °C for 90 min. The reaction was diluted with sat.  $\text{Na}_2\text{S}_2\text{O}_3$  (aq) then extracted with DCM. The combined organic extracts were concentrated and purified by silica chromatography (5-70% EtOAc in hexanes) to afford the title compound (1.30 g, 2.89 mmol, 74 % yield). MS (apci)  $m/z$  = 451.0 (M+H).

Step 4: Preparation of 11-chloro-3-cyclopropyl-7-(1-methylcyclopropyl)-6,7-dihydroisoxazolo[4',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one. To a solution of 1-(3-(6-(bromomethyl)-4-chloro-7-(1-methylcyclopropyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)ethan-1-one (1.32 g, 2.94 mmol) in THF (294 mL) cooled to -78 °C was added LiHMDS (1 M THF) (5.88 mL, 5.88 mmol). The reaction was warmed to -40 °C and stirred for 1 h. The reaction was then quenched with sat.  $\text{NH}_4\text{Cl}$ , then diluted with sat.  $\text{NaHCO}_3$  (aq) and extracted with DCM. The combined organic extracts were concentrated and purified first by silica chromatography (5-80% EtOAc in hexanes), followed by reverse-phase chromatography (5 to 95% MeCN in water with 0.1% TFA). The combined product fractions was diluted with sat.  $\text{NaHCO}_3$  (aq) and extracted with DCM. The combined organic extracts was dried ( $\text{Na}_2\text{SO}_4$ ), filtered, and concentrated to afford the title product (0.48 g, 45%). MS (apci)  $m/z$  = 369.1 (M+H).


#### Intermediate R1



#### Ethyl 3-cyclopropylpropiolate

A solution of ethynylcyclopropane (0.78 g, 11.8 mmol) in THF (8 mL) was sparged with  $\text{Ar}_{(\text{g})}$  and cooled to -78 °C, followed by dropwise addition of n-BuLi (2.5 M in THF; 5.2 mL, 13.0 mmol). The mixture was stirred for 30 min at -78 °C and for 45 min at -10 °C. The reaction mixture then was re-cooled to -78 °C before ethyl carbonochloridate (1.24 mL, 13.0 mmol) was added dropwise. The resulting mixture was allowed to warm to RT, then was stirred overnight before partitioning between EtOAc and saturated  $\text{NH}_4\text{Cl}_{(\text{aq})}$ . After phase-separation, the organic layer was washed with water, then dried over anhydrous  $\text{Na}_2\text{SO}_{4(\text{s})}$ , filtered, and concentrated *in vacuo* to afford the title compound (1.6 g, 98% yield). MS (apci)  $m/z$  = 139.1 (M+H).


#### Intermediate R2



Methyl 2-bromo-5-(2-cyclopropylacetamido)benzoate

To a mixture of methyl 5-amino-2-bromobenzoate (1.0 g, 4.3 mmol), cyclopropylacetic acid (0.48 mL, 5.2 mmol) and HATU (4.9 g, 13.0 mmol) in DCM (22 mL) was added DIEA (3.8 mL, 21.7 mmol) at RT, and the reaction mixture was stirred overnight. The reaction was concentrated and purified by silica chromatography (0-100% acetone in DCM) to afford the title product (1.36 g, 100%). MS (apci)  $m/z$  = 312.0 (M+H).


**Intermediate R3**



Methyl 2-bromo-5-(3-cyclopropylureido)benzoate

To a solution of triphosgene (297 mg, 1.0 mmol) in DCM (15 mL) was added a solution of methyl 5-amino-2-bromobenzoate (576 mg, 2.5 mmol) and DIEA (2.2 mL, 12.5 mmol) in DCM (10 mL) at 0 °C. After 1 h stirring, cyclopropylamine (208  $\mu$ L, 3.0 mmol) was introduced and the reaction was warmed to RT then stirred for 16 h. The reaction mixture was diluted with sat.  $\text{NaHCO}_3$  and extracted with DCM (2x). The combined organics were dried ( $\text{Na}_2\text{SO}_4$ ), filtered, concentrated and purified by silica chromatography (5 to 95% acetone in hexanes) to afford the title product (702 mg, 90%).  $^1\text{H-NMR}$  ( $d^6$ -DMSO)  $\delta$  8.65 (br s, 1H), 7.93 (m, 1H), 7.56 (d, 1H), 7.46 (m, 1H), 6.52 (br s, 1H), 3.84 (s, 3H), 2.53 (m, 1H), 0.63 (m, 2H), 0.41 (m, 2H).

**Intermediate R4**




(R)-1,1,1-trifluoropropan-2-yl trifluoromethanesulfonate

To a solution of (R)-1,1,1-trifluoropropan-2-ol (90 mg, 0.79 mmol) and pyridine (67  $\mu$ L, 0.83 mmol) in DCM (1.6 mL) was slowly added triflic anhydride (133  $\mu$ L, 0.79 mmol) at 0 °C. The reaction was slowly

warmed to rt and stirred for 15 min, after which it was filtered. The filter cake was rinsed with DCM to yield the title product as a 0.4 M solution, which was directly used in the next step assuming quantitative yield.


#### Intermediate R5



#### (S)-1,1,1-trifluoropropan-2-yl trifluoromethanesulfonate

The title product was prepared in a similar fashion as described in **Intermediate R4**, replacing (R)-1,1,1-trifluoropropan-2-ol with (S)-1,1,1-trifluoropropan-2-ol instead.

#### Intermediate R6




#### 3,3-difluorocyclobutyl trifluoromethanesulfonate

The title product was prepared in a similar fashion as described in **Intermediate R4**, replacing (R)-1,1,1-trifluoropropan-2-ol with 3,3-difluorocyclobutan-1-ol instead.

### Preparation of Synthetic Examples

#### Example 1



#### 11-amino-3-cyclopropyl-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one

Step 1: Preparation of 3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one. A mixture of ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate (**Intermediate P5**; 27 mg, 0.044 mmol) and DIEA (15.3  $\mu$ L, 0.0880 mmol) in DMF (440  $\mu$ L) was treated with ethylamine (2 M in THF; 110  $\mu$ L, 0.22 mmol). The resulting mixture was stirred for 2 h at RT and then overnight at 90 °C. After cooling to RT, the reaction mixture was partitioned between DCM and water. The organic extracts were washed successively with water (2x) and brine, then dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound (22 mg, 97% yield). MS (apci)  $m/z$  = 517.2 (M+H).



Step 2: Preparation of 11-amino-3-cyclopropyl-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one. A solution of 3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one (22 mg, 0.043 mmol) in TFA (1 mL, 12.9 mmol) and DCM (1 mL) was stirred for 1 h at RT. Subsequently, the mixture was concentrated *in vacuo*, then partitioned between DCM and saturated NaHCO<sub>3(aq)</sub>. Following phase separation, the organic extracts were washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (1-10% DCM:MeOH as the gradient eluent) to afford the title compound (3.5 mg, 22 % yield). MS (apci) m/z = 367.2 (M+H).


The compounds in the following Table were prepared using a similar two-step procedure as described for the synthesis of 11-amino-3-cyclopropyl-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one (**Example 1**), replacing ethylamine with the appropriate amine in Step 1, followed by protection group(s) removal with the appropriate reagent(s). Reactions were monitored by LCMS, and the products purified by chromatography using appropriate eluents.

Table A

| Ex # | Structure | Chemical Name                                                                                                                       | MS apci (m/z) |
|------|-----------|-------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 2    |           | 11-amino-3-cyclopropyl-5-(2-hydroxypropyl)-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one | 397.2 (M+H)   |
| 3    |           | 11-amino-3-cyclopropyl-5-(2-hydroxyethyl)-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one  | 383.2 (M+H)   |
| 4    |           | 11-amino-3-cyclopropyl-7-isopropyl-5-propyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one            | 381.2 (M+H)   |

| Ex # | Structure | Chemical Name                                                                                                                                          | MS apci (m/z) |
|------|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 5    |           | 11-amino-3-cyclopropyl-5-(cyclopropylmethyl)-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                  | 393.2 (M+H)   |
| 6    |           | 11-amino-3-cyclopropyl-7-isopropyl-5-(2-methoxyethyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                     | 397.2 (M+H)   |
| 7    |           | 11-amino-5-benzyl-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                               | 429.2 (M+H)   |
| 8    |           | 11-amino-3-cyclopropyl-7-isopropyl-5-phenethyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                            | 443.2 (M+H)   |
| 9    |           | 11-amino-3-cyclopropyl-5-(2-(1,1-dioxidothiomorpholino)ethyl)-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one | 500.2 (M+H)   |

## Example 10



11-amino-5-(3-chlorophenethyl)-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one

The title compound was prepared in a similar fashion as described in the synthesis of 11-amino-3-cyclopropyl-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one (**Example 1**), replacing ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate (**Intermediate P5**) with

methyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate


(**Intermediate P6**), and ethylamine with 2-(3-chlorophenyl)ethan-1-amine in step 1 instead. MS (apci)  $m/z = 477.2$  (M+H).


The compounds in the following Table were prepared in a similar fashion as described for the synthesis of 11-amino-5-(3-chlorophenethyl)-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one (**Example 10**), replacing 2-(3-chlorophenyl)ethan-1-amine with the appropriate amine in Step 1.

**Table B**

| Ex # | Structure | Chemical Name                                                                                                                                          | MS apci (m/z) |
|------|-----------|--------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 11   |           | 11-amino-3-cyclopropyl-7-isopropyl-5-(2-(4-isopropylpiperazin-1-yl)ethyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one | 493.3 (M+H)   |

| Ex # | Structure | Chemical Name                                                                                                                                         | MS apci (m/z) |
|------|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 12   |           | 11-amino-5-(4-chlorophenethyl)-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                 | 477.2 (M+H)   |
| 13   |           | 11-amino-3-cyclopropyl-7-isopropyl-5-(3-methoxyphenethyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                | 473.2 (M+H)   |
| 14   |           | 11-amino-3-cyclopropyl-7-isopropyl-5-(4-methoxyphenethyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one                | 473.2 (M+H)   |
| 15   |           | 11-amino-3-cyclopropyl-7-isopropyl-5-(2-(pyridin-2-yl)ethyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one             | 444.2 (M+H)   |
| 16   |           | 11-amino-3-cyclopropyl-7-isopropyl-5-(2-(1-methyl-1H-pyrazol-4-yl)ethyl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one | 447.2 (M+H)   |

| Ex # | Structure                                                                         | Chemical Name                                                                                                                               | MS apci (m/z) |
|------|-----------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 17   |  | 11-amino-5-(1-benzylpyrrolidin-3-yl)-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one | 498.3 (M+H)   |

**Example 18**


11-amino-3-cyclopropyl-7-isopropyl-5-(1-methyl-1H-pyrazol-4-yl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one

Step 1: Preparation of ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((1-methyl-1H-pyrazol-4-yl)amino)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate. A mixture of ethyl To a solution of ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((methylsulfonyl)oxy)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate (**Intermediate PS**; 32 mg, 0.052 mmol) in DMF (0.6 mL) was added 1-methyl-1H-pyrazol-4-amine (25.3 mg, 0.26 mmol) followed by DIEA (19.1  $\mu$ L, 0.110 mmol) and stirred overnight at RT, then at 90°C for 18 h. After cooled to RT, the reaction was partitioned between water and DCM. The combined organic extracts were concentrated and purified by silica chromatography (0-4% MeOH in DCM) to afford the title product (15 mg, 47% yield). MS (apci) m/z = 615.3 (M+H).

Step 2: Preparation of 3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-7-isopropyl-5-(1-methyl-1H-pyrazol-4-yl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one. To ethyl 5-cyclopropyl-3-(4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6-(((1-methyl-1H-pyrazol-4-yl)amino)methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)isoxazole-4-carboxylate (15 mg, 0.024 mmol) in MeOH (0.25 mL) was added LiOH (2 M, aq) (61  $\mu$ L, 0.12 mmol) and stirred for 2 h. The reaction was partitioned between water and DCM. After phase-separation and extracting the aqueous with DCM (2x), the combined organic extracts were concentrated to give the title product (12 mg, 86 % yield). MS (apci) m/z = 569.2 (M+H).

Step 3: Preparation of 11-amino-3-cyclopropyl-7-isopropyl-5-(1-methyl-1H-pyrazol-4-yl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one. To a solution of 3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-7-isopropyl-5-(1-methyl-1H-pyrazol-4-yl)-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one (12 mg, 0.021 mmol) in DCM (0.5 mL) was added triethylsilane (0.034 mL, 0.21 mmol) and 2,2,2-trifluoroacetic acid (0.5 mL, 0.021 mmol) and stirred overnight. The reaction mixture was then concentrated and partitioned between DCM and sat. NaHCO<sub>3</sub>(aq). After phase-separation and extracting the aqueous with DCM (2x), the combined organic extracts were concentrated and purified by silica chromatography (0-5% MeOH in DCM) to afford the title product (7 mg, 79% yield) MS (apci) m/z = 419.2 (M+H).

#### Example 19



#### 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one 2,2,2-trifluoroacetate

Step 1: Preparation of ethyl 3-(4-chloro-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate. A cold (0 °C) solution of ethyl 3-(6-(((tert-butyl)dimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P4**; 120 mg, 0.231 mmol) and acetic acid (14.7 M; 13.9 µL, 0.243 mmol) in THF (2.31 mL) was treated with TBAF (1.0 M in THF; 243 µL, 0.243 mmol). The resulting solution was stirred for 4 h at RT, after which the mixture was partitioned between DCM and saturated NaHCO<sub>3</sub>(aq). Following phase separation, the organic extracts were washed with brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>(s), filtered, and concentrated *in vacuo* to afford the title compound (93 mg, 99% yield). MS (apci) m/z = 405.1 (M+H).


Step 2: Preparation of ethyl 3-(6-(azidomethyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate. A solution of ethyl 3-(4-chloro-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (30 mg, 0.074 mmol) in toluene (741 µL) was treated sequentially with diphenylphosphoryl azide (32 µL, 0.15 mmol) and DBU (22 µL, 0.15 mmol), then stirred for 20 h at RT. Subsequently, the reaction mixture was diluted with EtOAc, and washed with water. The organic extracts then were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>(s), filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-50% EtOAc in hexanes) to afford the title compound (20 mg, 63% yield). MS (apci) m/z = 430.1 (M+H).

Step 3: Preparation of ethyl 3-(6-(aminomethyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate. A solution of ethyl 3-(6-(azidomethyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (20 mg, 0.047 mmol) in THF (931 µL)

and water (93  $\mu$ L) was treated with  $\text{PPh}_3$  (24 mg, 0.093 mmol). The resulting solution was stirred for 1 h at RT before introducing additional THF (500  $\mu$ L) and water (500  $\mu$ L). After stirring for an additional 2 h at RT, the solution was concentrated *in vacuo* to afford the title compound (19 mg, quantitative yield). MS (apci)  $m/z$  = 404.2 (M+H).

Step 4: Preparation of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one 2,2,2-trifluoroacetate. In a sealed tube, a mixture of ethyl 3-(6-(aminomethyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (19 mg, 0.047 mmol),  $\text{NH}_4\text{OH}$  (28-30%) (2 mL) and 1,4-dioxane (1 mL) was stirred for 2 h at 80 °C and then for 2 h at 100 °C. After cooling to RT, the solution was concentrated *in vacuo*. The crude residue was purified by C18 reverse phase chromatography (0-95% ACN in water with 0.1% TFA) to afford the title compound as the TFA salt (10 mg, 47% yield). MS (apci)  $m/z$  = 339.2 (M+H).

#### Example 20




11-amino-3-cyclopropyl-7-isopropyl-5-methyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one

#### Method A:

The title product was prepared by as similar fashion as described for the synthesis of 11-amino-3-cyclopropyl-5-ethyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one (**Example 1**), replacing ethylamine with methyl amine in Step 1 instead. MS (apci)  $m/z$  = 353.1 (M+H).

#### Method B:

A suspension of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-4(5H)-one 2,2,2-trifluoroacetate (**Example 19**; 15 mg, 0.033 mmol) in DMF (554  $\mu$ L) was treated with 60 wt.% NaH dispersion in mineral oil (2.1 mg, 0.053 mmol), and stirred for 10 min at RT. The resulting solution was cooled to 0 °C then treated with iodomethane (3.3  $\mu$ L, 0.053 mmol), and stirred for 1 h at 0 °C. After being allowed to warm to RT, the reaction mixture was directly purified by C18 reverse phase chromatography (0-95% ACN in water with 0.1% TFA) to afford the title compound as the TFA salt. The TFA salt then was dissolved in MeOH (500  $\mu$ L), and treated with saturated  $\text{NaHCO}_3(\text{aq})$  (100  $\mu$ L) and water (2 mL). The resulting mixture was concentrated *in vacuo* to about half of the original volume (1.5 mL). The resulting suspension was allowed to settle, and the supernatant was cannulated from the solid precipitate. The solids then were dried *in vacuo* to afford the title compound (4 mg, 34% yield). MS (apci)  $m/z$  = 353.1 (M+H).

**Example 21**


11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[3,4-f]pyrimido[5',4':4,5]pyrrolo[2,3-d][1,2]thiazepine 4,4-dioxide 2,2,2-trifluoroacetate

Step 1: Preparation of methyl ((3-(4-chloro-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)sulfonyl)carbamate, tetrabutylammonium salt. A cold (-78 °C) solution of 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole (**Intermediate P2**; 387 mg, 0.675 mmol) in THF (3.38 mL) was treated with *n*-BuLi (2.5 M in THF; 324 µL, 0.811 mmol), and stirred for 10 min at -78 °C. The resultant red-orange solution then was treated with Burgess reagent (145 mg, 0.608 mmol), and the solution was allowed to warm to -30 °C over a period of 1 h. Subsequently, the mixture was quenched with saturated NH<sub>4</sub>Cl<sub>(aq)</sub> and water. After being allowed to warm to RT, the quenched mixture was extracted with EtOAc. The combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The resulting foam residue was then dissolved in THF (6.5 mL), and treated with TBAF (1.0 M in THF; 878 µL, 0.878 mmol). After stirring for 15 min at RT, the reaction mixture was diluted with EtOAc, and washed with water. The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-50% EtOAc in hexanes followed by 0-10% MeOH in DCM) to afford the title compound as the tetrabutylammonium salt (67 mg, 21% yield). MS (apci) *m/z* = 470.1 (M+H).

Step 2: Preparation of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[3,4-f]pyrimido[5',4':4,5]pyrrolo[2,3-d][1,2]thiazepine 4,4-dioxide 2,2,2-trifluoroacetate. A solution of methyl ((3-(4-chloro-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazol-4-yl)sulfonyl)carbamate, tetrabutylammonium salt (48 mg, 0.067 mmol) in DCM (2.0 mL) was treated sequentially with MsCl (23.9 µL, 0.306 mmol) and DIPEA (89.2 µL, 0.511 mmol), then stirred for ~15 h at RT. The reaction mixture then was concentrated *in vacuo*. The resulting residue was dissolved in 1,4-dioxane (1.0 mL) and NH<sub>4</sub>OH<sub>(aq)</sub> (28-30%) (511 µL). The resulting mixture was transferred to a sealed tube, and stirred for 30 min at 80 °C. After cooling to RT, the resulting solution was concentrated *in vacuo*. The crude residue was purified by C18 reverse phase chromatography (0-95% ACN in water with 0.1% TFA). The desired fractions were combined and concentrated *in vacuo* to about 5 mL, at which point solid precipitated from the solution. The solids were collected by filtration and then dried *in vacuo* to afford the title compound as the TFA salt (23.5 mg, 71% yield). MS (apci) *m/z* = 375.2 (M+H).

**Example 22**






11-amino-3-cyclopropyl-7-isopropyl-5-methyl-6,7-dihydro-5H-isoxazolo[3,4-f]pyrimido[5',4':4,5]pyrrolo[2,3-d][1,2]thiazepine 4,4-dioxide

A suspension of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[3,4-f]pyrimido[5',4':4,5]pyrrolo[2,3-d][1,2]thiazepine 4,4-dioxide 2,2,2-trifluoroacetate (**Example 21**; 11 mg, 0.0225 mmol) in DMF (322  $\mu$ L) was treated with 60 wt.% NaH dispersion in mineral oil (1.98 mg, 0.0496 mmol), and stirred for 10 min at RT. The resulting solution was cooled to 0°C, then treated with iodomethane (1.55  $\mu$ L, 0.0248 mmol) and allowed to warm to RT. After stirring for 1 h at RT, the resulting mixture was directly purified by C18 reverse phase chromatography (0-95% ACN in water with 0.1% TFA) to afford the title compound as the TFA salt. The TFA salt then was dissolved in MeOH, and the solution was passed through a basic resin, PL-HCO<sub>3</sub> MP, filter plug. The filtrate was concentrated *in vacuo* to afford the title compound (1.5 mg, 17% yield). MS (apci)  $m/z$  = 389.3 (M+H).

**Example 23**



11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[4'',3'':2',3']thiepine[4',5':4,5]pyrrolo[2,3-d]pyrimidine 4,4-dioxide 2,2,2-trifluoroacetate

Step 1: Preparation of 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-(methylsulfonyl)isoxazole. A cold (-78 °C) solution of 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-iodoisoxazole (**Intermediate P2**; 130 mg, 0.227 mmol) in THF (1.51 mL) was treated with n-BuLi (2.5 M in THF; 127  $\mu$ L, 0.318 mmol), and stirred for 15 min at -78 °C. The resultant yellow-orange solution then was treated with DMDS (40.3  $\mu$ L, 0.454 mmol), and the solution was allowed to warm to -30 °C over a period of 1 h. Subsequently, the mixture was quenched with saturated NH<sub>4</sub>Cl<sub>(aq)</sub>. After being allowed to warm to RT, the quenched mixture was extracted with EtOAc. The combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The resulting residue then was dissolved in DCM (2.27 mL). The solution was cooled to 0 °C and treated with mCPBA (70 wt.%; 117 mg, 0.476 mmol). After overnight stirring at 0 °C, additional mCPBA (70 wt.%; 27.9 mg, 0.113 mmol) was


introduced, and stirring was resumed for another 1.5 h at 0 °C. The reaction was quenched with 10% Na<sub>2</sub>S<sub>2</sub>O<sub>3(aq)</sub> (2 mL) and saturated NaHCO<sub>3(aq)</sub> (5 mL) and extracted with DCM. The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-50% EtOAc in hexanes) to afford the title compound (72 mg, 60% yield). MS (apci) m/z = 525.2 (M+H).

Step 2: Preparation of 3-(6-(bromomethyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-(methylsulfonyl)isoxazole. A solution of 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-(methylsulfonyl)isoxazole (65 mg, 0.12 mmol) in THF (1.2 mL) was treated with TBAF (1.0 M THF) (186 µL, 0.19 mmol), and stirred for 15 min at RT. The reaction mixture then was diluted with EtOAc and washed with water. The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The residue was dissolved in DCM (1.2 mL) and treated with PBr<sub>3</sub> (97.4 mg, 0.36 mmol). The reaction mixture was stirred for 1 h at RT, before quenching with saturated NaHCO<sub>3(aq)</sub>. The quenched mixture was extracted with DCM, and the organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-30% EtOAc in hexanes) to afford the title compound (50 mg, 85% yield). MS (apci) m/z = 473.0 (M+H).

Step 3: Preparation of 11-chloro-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[4'',3'':2',3']thiepino[4',5':4,5]pyrrolo[2,3-d]pyrimidine 4,4-dioxide. A cold (-78 °C) solution of 3-(6-(bromomethyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropyl-4-(methylsulfonyl)isoxazole (50 mg, 0.11 mmol) in THF (2.1 mL) was treated with LiHMDS (1 M in THF; 317 µL, 0.32 mmol). The resulting blue solution was stirred for 1 h at -78 °C before the mixture was quenched with saturated NH<sub>4</sub>Cl<sub>(aq)</sub> and water. After being allowed to warm to RT, the quenched mixture was extracted with EtOAc. The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo* to afford the title compound (43 mg, quantitative yield). MS (apci) m/z = 393.0 (M+H).

Step 4: Preparation of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[4'',3'':2',3']thiepino[4',5':4,5]pyrrolo[2,3-d]pyrimidine 4,4-dioxide 2,2,2-trifluoroacetate. In a sealed tube, a solution of 11-chloro-3-cyclopropyl-7-isopropyl-6,7-dihydro-5H-isoxazolo[4'',3'':2',3']thiepino[4',5':4,5]pyrrolo[2,3-d]pyrimidine 4,4-dioxide (43 mg, 0.11 mmol) in 1,4-dioxane (1.1 mL) and NH<sub>4</sub>OH<sub>(aq)</sub> (28-30%) (1.1 mL) was stirred for 45 min at 100 °C. After cooling to RT, the resulting solution was concentrated *in vacuo*. The crude residue was purified by C18 reverse phase chromatography (0-95% ACN in water with 0.1% TFA) to afford the title compound as the TFA salt (8 mg, 15% yield). MS (apci) m/z = 374.3 (M+H).

#### Example 24




11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-4H-isoxazolo[3'',4'':5',6']oxepino[4',3':4,5]pyrrolo[2,3-d]pyrimidin-4-one

Step 1: Preparation of methyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate. A solution of methyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-chloro-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (**Intermediate P3**; 279 mg, 0.552 mmol) and (2,4-dimethoxyphenyl)methanamine (207.5  $\mu$ L, 1.381 mmol) in DMSO (3.683 mL) was stirred overnight at 60 °C. After cooling to RT, the reaction mixture was diluted with EtOAc, then successively washed with water (4x) and brine. The organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo* to afford the title compound (351 mg, 100% yield). MS (apci)  $m/z$  = 636.3 (M+H).

Step 2: Preparation of 3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6,7-dihydro-4H-isoxazolo[3'',4'':5',6']oxepino[4',3':4,5]pyrrolo[2,3-d]pyrimidin-4-one. A solution methyl 3-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carboxylate (351 mg, 0.552 mmol) in THF (3.68 mL) was treated with TBAF (1 M in THF; 607  $\mu$ L, 0.607 mmol), and stirred for 2 h at RT. The reaction mixture then was diluted with EtOAc and extracted successively with water and brine. The organic extracts were dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (10-90% EtOAc in hexanes) to afford the title compound (140 mg, 52% yield). MS (apci)  $m/z$  = 490.2 (M+H).

Step 3: 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydro-4H-isoxazolo[3'',4'':5',6']oxepino[4',3':4,5]pyrrolo[2,3-d]pyrimidin-4-one. A mixture of 3-cyclopropyl-11-((2,4-dimethoxybenzyl)amino)-7-isopropyl-6,7-dihydro-4H-isoxazolo[3'',4'':5',6']oxepino[4',3':4,5]pyrrolo[2,3-d]pyrimidin-4-one (54 mg, 0.11 mmol) and triethylsilane (17.6  $\mu$ L, 0.110 mmol) in TFA (1 mL) and DCM (1 mL) was stirred for 1 h at RT. The resulting mixture was concentrated *in vacuo*, and the residue was partitioned between DCM and saturated  $\text{NaHCO}_{3(aq)}$ . The organics extracts then were washed with brine, dried over anhydrous  $\text{Na}_2\text{SO}_{4(s)}$ , filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (1-10% DCM: MeOH) to afford the title compound (19.5 mg, 52% yield). MS (apci)  $m/z$  = 340.1 (M+H).

#### Example 25



11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one


Step 1: Preparation of benzyl (E)-3-(4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)acrylate. A solution of 4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P1**; 100 mg, 0.302 mmol) in DCM (3.02 mL) was treated with benzyl (triphenylphosphoranylidene)acetate (149 mg, 0.363 mmol), and stirred for 1 h at RT. The resulting solution was concentrated *in vacuo*, and the residue was purified silica chromatography (0-25% EtOAc in hexanes) to afford the title compound (137 g, 98% yield). MS (apci)  $m/z$  = 463.1 (M+H).

Step 2: Preparation of 3-(4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoic acid. A solution of benzyl (E)-3-(4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)acrylate (137 mg, 0.296 mmol) in EtOAc (2.96 mL) was added 10% Pd/C (15.7 mg, 0.0148 mmol). The mixture was stirred under a balloon of  $H_{2(g)}$  for 1 h at RT before introducing EtOH (3 mL) and additional 10% Pd/C (31.4 mg, 0.0296 mmol). The resulting mixture was placed back under a balloon of  $H_{2(g)}$  and stirred for an additional 1.5 h at RT. The mixture was filtered through a short pad of celite, and the filtrate was concentrated *in vacuo* to afford title compound (100 mg), which was used directly in the next step without further purifications. MS (apci)  $m/z$  = 375.1 (M+H).

Step 3: Preparation of 3-(4-amino-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoic acid. A solution of crude 3-(4-chloro-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoic acid (100 mg, 0.267 mmol) and  $NH_4OH_{(aq)}$  (28-30%) (3 mL) in 1,4-dioxane (2 mL) was stirred for 16 h at 100 °C. After cooling to RT, the solution was concentrated *in vacuo* to afford a product mixture containing the title compound, which was used directly in the next step without further purifications. MS (apci)  $m/z$  = 356.2 (M+H).

Step 4: Preparation of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one. A solution of the crude mixture from the previous step containing 3-(4-amino-5-(5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-6-yl)propanoic acid (17 mg, 0.048 mmol) in polyphosphoric acid (PPA) (1 mL) was stirred for 1 h at 100 °C. After cooling to RT, the mixture was neutralized with saturated  $NaHCO_{3(aq)}$  and extracted with DCM (3x30 mL). The combined organic extracts were concentrated *in vacuo*, and the residue was purified by silica chromatography (0-10% MeOH/DCM) to afford the title compound (7.4 mg, 45% yield). MS (apci)  $m/z$  = 338.1 (M+H).


**Example 26**



11-amino-3-cyclopropyl-7-isopropyl-4-methyl-4,5,6,7-tetrahydroisoxazolo[4'',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4-ol


A solution of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one (**Example 25**; 7 mg, 0.02 mmol) in THF (200  $\mu$ L) was treated with anhydrous  $\text{CeCl}_3$  (26 mg, 0.11 mmol) at 0  $^\circ\text{C}$ , followed by  $\text{MeMgBr}$  (1.4M in toluene/THF 75:25) (300  $\mu$ L, 0.4 mmol). After stirred for 1.5 h at 0  $^\circ\text{C}$ , the reaction mixture was cooled to -78  $^\circ\text{C}$  before introducing  $\text{MeLi}$  (1.6 M  $\text{Et}_2\text{O}$ ) (133  $\mu$ L, 0.21 mmol). After stirred for 1 h at -78  $^\circ\text{C}$ , the reaction mixture was quenched with  $\text{MeOH}$ , allowed to warm to RT and then partitioned between DCM and water (5 mL each). Following phase separation, the aqueous phase was back extracted with additional DCM (2 x 5 mL). The DCM extracts were combined and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-10%  $\text{MeOH}/\text{DCM}$ ) to afford the title compound (4.5 mg, 60% yield). MS (apci)  $m/z$  = 354.2 ( $\text{M}+\text{H}$ ).

**Example 27**




3-cyclopropyl-7-isopropyl-N4-methyl-4,5,6,7-tetrahydroisoxazolo[4'',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidine-4,11-diamine

A solution of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3'':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one (**Example 25**; 10 mg, 0.03 mmol) in DCE (200  $\mu$ L) was treated with methylamine (2 M THF) (74  $\mu$ L, 0.15 mmol). The reaction mixture was stirred for 30 min at RT, and was then treated with  $\text{NaBH}(\text{OAc})_3$  (38 mg, 0.18 mmol). The resulting mixture was stirred for 45 min at RT, then overnight at 50  $^\circ\text{C}$ . Subsequently,  $\text{AcOH}$  (10 drops) and additional methylamine (300  $\mu$ L, 0.61 mmol) were introduced, and the reaction mixture was stirred for 30 min at 50  $^\circ\text{C}$ . The resulting mixture was treated with additional  $\text{NaBH}(\text{OAc})_3$  (38 mg, 0.18 mmol). This mixture was stirred overnight at 50  $^\circ\text{C}$ , then for 2 days at 65  $^\circ\text{C}$  and finally for 3 days at RT. The reaction then was concentrated *in vacuo*, and purified by silica chromatography (0-10%  $\text{MeOH}/\text{DCM}$ ) to afford the title compound (2.8 mg, 27% yield). MS (apci)  $m/z$  = 353.2 ( $\text{M}+\text{H}$ ).

**Example 28**


11-amino-3-cyclopropyl-7-isopropyl-4-(trifluoromethyl)-4,5,6,7-tetrahydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4-ol

A solution of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one (**Example 25**; 18 mg, 0.054 mmol) in THF (400  $\mu$ L) was treated with TMSCF<sub>3</sub> (50  $\mu$ L, 0.33 mmol) and CsF (4 mg, 0.027 mmol). The reaction mixture was stirred for 1.5 h at RT, and then HCl<sub>(aq)</sub> (5 M; 542  $\mu$ L, 2.71 mmol) was added. The resulting mixture was stirred for 15 min at RT, then neutralized with saturated NaHCO<sub>3(aq)</sub> (5 mL). The neutralized mixture was extracted with DCM (3 x 5 mL). The organic extracts were concentrated *in vacuo*, and the residue was purified first by silica chromatography (0-10% MeOH/DCM), then by C18 reverse-phase chromatography (5-90% MeCN/water with 0.1% TFA) to afford the title compound as the TFA salt. The TFA salt was dissolved in DCM (5 mL) and extracted with saturated NaHCO<sub>3(aq)</sub> (5 mL). The aqueous phase then was back extracted with DCM (2 x 5 mL). The DCM extracts were combined, dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo* to afford the title compound (7 mg, 32% yield). MS (apci) *m/z* = 408.1 (M+H).

**Example 29**

11-amino-3-cyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4-ol

A solution of 11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4(5H)-one (**Example 25**; 18 mg, 0.055 mmol) in MeOH (600  $\mu$ L) was treated with NaBH<sub>4</sub> (6.19 mg, 0.164 mmol), and the resulting mixture was stirred for 15 min at RT. The reaction mixture then was concentrated *in vacuo*, and the residue was purified by silica chromatography (0-10% MeOH/DCM) to afford the title compound (13.6 mg, 74% yield). MS (apci) *m/z* = 340.2 (M+H).


**Example 30**

3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-11-amine

Step 1: Preparation of (E)-N'-(3-cyclopropyl-4-hydroxy-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-11-yl)-N,N-dimethylformimidamide. A solution of 11-amino-3-cyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4'',3''':6',7']cyclohepta-[1',2':4,5]pyrrolo[2,3-d]pyrimidin-4-ol (**Example 29**; 22.2 mg, 0.0654 mmol) in EtOH (300  $\mu$ L) was treated with 1,1-dimethoxy-N,N-dimethylmethanamine (87.2  $\mu$ L, 0.654 mmol) and 2,2,2-trifluoroacetic acid (1.5  $\mu$ L, 0.02 mmol) then stirred for 3 h at 65°C. After cooling to RT, the reaction mixture was concentrated *in vacuo* to afford the title compound, which was used directly in the next step without further purifications, assuming quantitative yield.

Step 2: Preparation of 3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-11-amine. A solution crude (E)-N'-(3-cyclopropyl-4-hydroxy-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4'',3''':6',7']cyclohepta[1',2':4,5]pyrrolo[2,3-d]pyrimidin-11-yl)-N,N-dimethylformimidamide (25 mg, 0.063 mmol) in DMF (300  $\mu$ L) was treated with iodomethane (39.5  $\mu$ L, 0.634 mmol) and DIEA (277  $\mu$ L, 1.58 mmol), and stirred for 45 min at RT. The reaction mixture was partitioned between DCM (5 mL) and saturated NaHCO<sub>3(aq)</sub> (5 mL). Following phase separation, the aqueous extracts were back extracted with DCM (2 x 5 mL). The DCM extracts were combined and concentrated *in vacuo*. The crude residue was treated with 4 N HCl (1 mL), and stirred overnight at 65 °C. After cooling to RT, the reaction mixture was partitioned between DCM (10 mL) and saturated NaHCO<sub>3(aq)</sub> (10 mL). Following phase separation, the aqueous extracts were back extracted with DCM (2 x 10 mL). The DCM extracts were combined and concentrated *in vacuo*. The residue was purified by silica chromatography (0-10% MeOH/DCM) to afford the title compound (2 mg, 10% yield). MS (apci) m/z = 322.2 (M+H).

**Example 31**



3-cyclopropyl-7-isopropyl-5-methyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine




Step 1: Preparation of 3-cyclopropyl-7-isopropyl-N,N-bis(4-methoxybenzyl)-5-methyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine. A solution of 3-(4-(bis(4-methoxybenzyl)amino)-6-formyl-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carbaldehyde (**Intermediate P10**; 25.7 mg, 0.0443 mmol) in DCE (887  $\mu$ L) was treated with  $\text{NaBH}(\text{OAc})_3$  (37.6 mg, 0.177 mmol), stirred for 15 min at RT, and then treated dropwise with methylamine (2.0 M THF) (44.3  $\mu$ L, 0.088 mmol). After stirring for 30 min at RT, the resulting mixture was diluted with DCM and washed with saturated  $\text{NaHCO}_3(\text{aq})$ . Following phase separation, the aqueous extracts were back extracted with DCM. The DCM extracts were combined, then passed through a phase separator column and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-100% [15% MeOH + 1.5%  $\text{NH}_4\text{OH}$  in DCM] in DCM) to afford the title compound (11.8 mg, 46% yield). MS (apci)  $m/z$  = 579.4 (M+H).


Step 2: Preparation of 3-cyclopropyl-7-isopropyl-5-methyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine. A solution of 3-cyclopropyl-7-isopropyl-N,N-bis(4-methoxybenzyl)-5-methyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (11.8 mg, 0.0204 mmol) in DCM (408  $\mu$ L) was treated with TFA (78.5  $\mu$ L, 1.02 mmol), and stirred for 30 min at RT. The reaction mixture was then treated with triethylsilane (32.6  $\mu$ L, 0.204 mmol), followed by stirring for an additional 30 min at RT before TFOH was introduced (1.81  $\mu$ L, 0.0204 mmol). The resulting mixture was stirred for 10 min at RT, and then neutralized with saturated  $\text{NaHCO}_3(\text{aq})$ . The neutralized mixture was extracted with DCM. The organic extracts were combined, passed through a phase separator column and concentrated *in vacuo*. After drying the residue under high-vacuum for 1 h, the title compound was cleanly isolated (4.1 mg, 59% yield). MS (apci)  $m/z$  = 399.2 (M+H).

The compounds in the following Table were prepared in a similar fashion as described for the synthesis of 3-cyclopropyl-7-isopropyl-5-methyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (**Example 31**), replacing methylamine with the appropriate amine in Step 1 instead.

**Table C**



| Ex # | Structure                                                                           | Chemical Name                                                                                                                             | MS<br>apci<br>(m/z) |
|------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|---------------------|
| 32   |    | 3,5-dicyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine 2,2,2-trifluoroacetate   | 365.2<br>(M+H)      |
| 33   |   | 3-cyclopropyl-7-isopropyl-5-phenyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine                     | 401.3<br>(M+H)      |
| 34   |  | 3-cyclopropyl-7-isopropyl-5-(1-methyl-1H-pyrazol-3-yl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine | 405.3<br>(M+H)      |

**Example 35**

3-cyclopropyl-7-isopropyl-5-(methylsulfonyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine 2,2,2-trifluoroacetate


Step 1: Preparation of 3-cyclopropyl-7-isopropyl-N,N-bis(4-methoxybenzyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine. A solution of 3-(4-(bis(4-methoxybenzyl)amino)-6-formyl-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-

carbaldehyde (**Intermediate P10**; 80.7 mg, 0.139 mmol) in DCE (2.78 mL) was treated with NaBH(OAc)<sub>3</sub> (148 mg, 0.696 mmol), stirred for 15 min at RT, and then treated dropwise with ammonium acetate (32.2 mg, 0.418 mmol). After stirring for 90 min at RT, the mixture was treated with AcOH (10 drops), and stirred for 1 h at 60 °C. After cooling to RT, the reaction mixture was diluted with DCM and washed with saturated NaHCO<sub>3(aq)</sub>. Following phase separation, the aqueous extracts were back extracted with DCM. The DCM extracts were combined, then passed through a phase separator column and concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-100% [15% MeOH + 1.5% NH<sub>4</sub>OH in DCM] in DCM) to afford the title compound (11.6 mg, 15% yield). MS (apci) *m/z* = 565.3 (M+H).


Step 2: Preparation of 3-cyclopropyl-7-isopropyl-5-(methylsulfonyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine 2,2,2-trifluoroacetate. A solution of 3-cyclopropyl-7-isopropyl-N,N-bis(4-methoxybenzyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (11.6 mg, 0.0205 mmol) in DCM (205 µL, 0.0205 mmol) was treated with MsCl (3.53 mg, 0.0308 mmol) and DIEA (17.9 µL, 0.103 mmol), then stirred overnight at RT. The resulting mixture then was treated with TfOH (182.4 µL, 2.06 mmol) and TFA (a few drops). The resulting mixture was stirred for 120 min at RT, then extracted with DCM and saturated NaHCO<sub>3(aq)</sub>. The organic extracts were concentrated *in vacuo* and purified by C18 reverse phase chromatography (10-90% ACN in water with 0.1% TFA) to afford the title compound (1.6 mg, 15% yield). MS (apci) *m/z* = 403.1 (M+H).

The compounds in the following Table were prepared by a similar fashion as described in the synthesis of 3-cyclopropyl-7-isopropyl-5-(methylsulfonyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine 2,2,2-trifluoroacetate (**Example 35**), replacing MsCl with the appropriate electrophile reagent (e.g. an acyl chloride) in Step 2 instead.

Table D

| Ex # | Structure                                                                           | Chemical Name                                                                                                                                        | MS apci (m/z) |
|------|-------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 36   |  | 1-(11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(4H)-yl)ethan-1-one 2,2,2-trifluoroacetate | 367.1 (M+H)   |

| Ex # | Structure | Chemical Name                                                                                                                                         | MS apci (m/z) |
|------|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| 37   |           | 1-(11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(4H)-yl)propan-1-one 2,2,2-trifluoroacetate | 381.3 (M+H)   |
| 38   |           | (11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(4H)-yl)(cyclopropyl)methanone                | 393.3 (M+H)   |
| 39   |           | 1-(11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(4H)-yl)-2-phenylethan-1-one                | 443.3 (M+H)   |


**Example 40**

2-(11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(4H)-yl)ethan-1-ol 2,2,2-trifluoroacetate

A solution of 3-cyclopropyl-7-isopropyl-N,N-bis(4-methoxybenzyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (Step 1, **Example 35**) (25 mg, 0.044 mmol) in DCM (443  $\mu$ L) was treated with (2-bromoethoxy)(tert-butyl)dimethylsilane (16 mg, 0.066 mmol) and DIEA (39  $\mu$ L, 0.22 mmol). The reaction mixture was then stirred at RT overnight, after which DMF (1 mL) was introduced, and the reaction vessel was sealed. The reaction was heated to 65°C for 16 h to reach

completion. After cooled to RT, the reaction mixture was treated with trifluoromethanesulfonic acid (197  $\mu$ L, 2.2 mmol) then stirred for 20 min. It was partitioned between  $\text{Na}_2\text{CO}_3$  (aq) and DCM. The combined organic extracts were concentrated and purified by prep HPLC (5-50% ACN in  $\text{H}_2\text{O}$  with 0.1% TFA) to afford the title product (1.6 mg, 7.5 % yield). MS (apci)  $m/z$  = 369.2 (M+H).

#### Example 41




#### 3-cyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine

Step 1: Preparation of 3-cyclopropyl-5-(2,4-dimethoxybenzyl)-7-isopropyl-N,N-bis(4-methoxybenzyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine. A solution of 3-(4-(bis(4-methoxybenzyl)amino)-6-formyl-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-cyclopropylisoxazole-4-carbaldehyde (**Intermediate P10**) (169 mg, 0.29 mmol) in DCE (5.8 mL) was treated with  $\text{NaBH}(\text{OAc})_3$  (309.0 mg, 1.46 mmol) and stirred at rt for 5 min, followed by portion-wise addition of (2,4-dimethoxyphenyl)methanamine (0.22 mL, 1.45 mmol). After 5 h stirring at RT, the mixture was concentrated. The residue was taken up in DCM and washed with sat.  $\text{NaHCO}_3$ . The aqueous was extracted with DCM, and the combined organic extracts were concentrated and purified by silica chromatography (0-70% EtOAc in hexanes) to afford the title compound (130 mg, 62 % yield). MS (apci)  $m/z$  = 715.3 (M+H).

Step 2: Preparation of 3-cyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine. A solution of 3-cyclopropyl-5-(2,4-dimethoxybenzyl)-7-isopropyl-N,N-bis(4-methoxybenzyl)-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (109 mg, 0.15 mmol) in DCM (3 mL) was treated with TFA (234  $\mu$ L, 3.0 mmol) and  $\text{Et}_3\text{SiH}$  (244  $\mu$ L, 1.52 mmol). After 2 d stirring at RT, the reaction mixture was neutralized with sat.  $\text{NaHCO}_3$  then extracted into DCM. The combined organic extracts were concentrated and purified by silica chromatography (0-100% [15% MeOH + 1.5%  $\text{NH}_4\text{OH}$  in DCM] in DCM) to afford the title product (18.7 mg, 35 % yield). MS (apci)  $m/z$  = 325.2 (M+H).


#### Example 42



1-(11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(4H)-yl)-3,3,3-trifluoropropan-1-one 2,2,2-trifluoroacetate

A solution of 3-cyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (**Example 41**) (14 mg, 0.043 mmol) in DCM (0.86 mL) was treated with 3,3,3-trifluoropropanoyl chloride (6.7  $\mu$ L, 0.065 mmol) and DIEA (38  $\mu$ L, 0.22 mmol) then stirred at rt overnight. The reaction was diluted with water and extracted with DCM. The combined organic extracts were concentrated and purified by prep HPLC (10-90% ACN in H<sub>2</sub>O with 0.1% TFA) to afford the title product (1.3 mg, 7 % yield). MS (apci)  $m/z$  = 435.2 (M+H).


**Example 43**



11-amino-3-cyclopropyl-7-isopropyl-6,7-dihydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepine-5(4H)-carboxamide 2,2,2-trifluoroacetate

To a solution of 3-cyclopropyl-7-isopropyl-4,5,6,7-tetrahydroisoxazolo[4,3-c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-11-amine (**Example 41**) (1.2 mg, 0.0037 mmol) in THF (1 mL) was added trimethylsilyl isocyanate (0.015 mL, 0.11 mmol) at rt. The reaction mixture was stirred for 3 d, after which it was directly purified by reverse phase chromatography (0-95% MeCN in water with 0.1% TFA) to afford the title compound as white solid (1.3 mg, 73%). MS (apci)  $m/z$  = 368.2 (M+H).


**Example 44**



3-cyclopropyl-7-isopropyl-6,7-dihydro-4H-isoxazolo[3'',4'':5',6']oxepino[4',3':4,5]pyrrolo[2,3-d]pyrimidin-11-amine

A cold (-78 °C) solution of 4-(bis(4-methoxybenzyl)amino)-5-(4-(((tert-butyldimethylsilyl)oxy)methyl)-5-cyclopropylisoxazol-3-yl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidine-6-carbaldehyde (**Intermediate P9**; 8.00 mg, 0.0115 mmol) in DCM (1.15 mL) was treated with TMSOTf (1.11 µL, 0.00575 mmol). After stirring for 1.5 h at -78 °C, the reaction mixture was treated with triethylsilane (2.02 µL, 0.0126 mmol). The resulting mixture was stirred for an additional 1.5 h at -78 °C, and it was then allowed to warm to RT. The reaction mixture was then partitioned between water and DCM. Following phase separation, the aqueous extracts were back extracted with DCM (3x). The DCM extracts were combined, passed through a phase separator column, and then concentrated *in vacuo*. The crude residue was purified by silica chromatography (0-10% MeOH in DCM with 0.1% NH<sub>4</sub>OH) to afford the title compound (2.34 mg, 63% yield). MS (apci) *m/z* = 326.1 (M+H).

**Example 45**



12-amino-3-chloro-8-isopropyl-7,8-dihydrobenzo[c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(6H)-one

Step 1: Preparation of methyl 2-(6-(((tert-butyldimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-chlorobenzoate. In a pressure vessel, solution of 6-(((tert-butyldimethylsilyl)oxy)methyl)-N-(2,4-dimethoxybenzyl)-5-iodo-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-4-amine (**Intermediate P14**; 299 mg, 0.501 mmol) in 4:1 dioxane:water (4.2 mL) was treated with 4-chloro-2-(methoxycarbonyl)phenylboronic acid (322 mg, 1.50 mmol), **PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>** (87.9 mg, 0.125 mmol), and K<sub>2</sub>CO<sub>3(s)</sub> (208 mg, 1.50 mmol). After sparging the mixture with Ar<sub>(g)</sub>, the vessel was sealed. The resulting mixture was stirred for 16 h at 90 °C. After cooling to RT, the mixture was treated with additional 4-chloro-2-(methoxycarbonyl)phenylboronic acid (107 mg, 0.499 mmol) and **PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub>** (35 mg, 0.0499 mmol), and sparged with Ar<sub>(g)</sub>. The vessel was sealed, and the mixture was stirred for an additional 1 h at 90 °C. After cooling to RT, the reaction mixture was diluted with 4:1 DCM: iPrOH, and washed with water (1x). The organic extracts were dried over anhydrous

Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-95% DCM-EtOAc) to afford the title compound (160.6 mg, 50% yield). MS (apci) m/z = 639.3 (M+H).

Step 2: Preparation of methyl 5-chloro-2-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzoate. A cold (0 °C) solution of methyl 2-(6-(((tert-butyl)dimethylsilyl)oxy)methyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-chlorobenzoate (160.6 mg, 0.2512 mmol) in THF (2.5 mL) was treated sequentially with AcOH (15.82 µL, 0.2763 mmol) and TBAF (1.0 M THF) (76.3 µL, 0.28 mmol). The resulting mixture was stirred for 16 h at RT, before introducing additional TBAF (276 µL, 0.28 mmol). The mixture was stirred for an additional 24 h at RT, and then was concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-95% DCM-EtOAc) to afford the title compound (114.8 mg, 87% yield). MS (apci) m/z = 525.2 (M+H).

Step 3: Preparation of methyl 2-(6-(azidomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-chlorobenzoate. A cold (0 °C) solution of methyl 5-chloro-2-(4-((2,4-dimethoxybenzyl)amino)-6-(hydroxymethyl)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)benzoate (114.8 mg, 0.2187 mmol) in toluene (2.2 mL) was treated sequentially with DBU (39.24 µL, 0.2624 mmol) and DPPA (56.55 µL, 0.2624 mmol), then warmed to RT, and stirred for 2 h. The resulting mixture was diluted with EtOAc, and washed successively with water (1x) and brine (1x). The combined organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by silica chromatography (5-95% hexanes-Acetone) to afford the title compound (95.7 mg, 80% yield). MS (apci) m/z = 550.2 (M+H).

Step 4: Preparation of methyl 2-(6-(aminomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-chlorobenzoate. A solution of methyl 2-(6-(azidomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-chlorobenzoate (95.7 mg, 0.174 mmol) in 1:1 THF:water (1.7 mL) was treated with PPh<sub>3</sub> (91.3 mg, 0.348 mmol), then stirred for 16 h at RT. Subsequently, the reaction mixture was diluted with 4:1 DCM:iPrOH, and washed with saturated NaHCO<sub>3(aq)</sub> (1x). The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo*. The crude residue was purified by C18 reverse-phase chromatography (5-95% ACN/water with 0.1% TFA) to afford the title compound as the TFA salt. The TFA salt was dissolved in 4:1 DCM:iPrOH and extracted with saturated NaHCO<sub>3(aq)</sub>. The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo* to afford the title compound (78.8 g, 86% yield). MS (apci) m/z = 524.2 (M+H).

Step 5: 12-amino-3-chloro-8-isopropyl-7,8-dihydrobenzo[c]pyrimido[5',4':4,5]pyrrolo[3,2-e]azepin-5(6H)-one. A solution of methyl 2-(6-(aminomethyl)-4-((2,4-dimethoxybenzyl)amino)-7-isopropyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-5-chlorobenzoate (78.8 mg, 0.150 mmol) in dioxane (3.0 mL) was stirred for 24 h at 90 °C. Subsequently, the mixture was cooled to RT, and concentrated *in vacuo*. The residue thus obtained was re-dissolved in TFA (1.5 mL), stirred for 1 h at RT, and concentrated *in vacuo*. The crude residue was purified by C18 reverse phase chromatography (5-95% water-ACN with 0.1% TFA) to afford the title compound as the TFA salt. The TFA salt was dissolved in 4:1 DCM:iPrOH, and extracted with saturated NaHCO<sub>3(aq)</sub>. The organic extracts were dried over anhydrous Na<sub>2</sub>SO<sub>4(s)</sub>, filtered, and concentrated *in vacuo* to afford the title compound (35.9 mg, 70% yield). MS (apci) m/z = 342.1 (M+H).

## DEMANDE OU BREVET VOLUMINEUX

LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVET COMPREND PLUS D'UN TOME.

CECI EST LE TOME        1    DE    2  
CONTENANT LES PAGES    1    À    318

NOTE : Pour les tomes additionels, veuillez contacter le Bureau canadien des brevets

## JUMBO APPLICATIONS/PATENTS

THIS SECTION OF THE APPLICATION/PATENT CONTAINS MORE THAN ONE VOLUME

THIS IS VOLUME        1    OF    2  
CONTAINING PAGES    1    TO    318

NOTE: For additional volumes, please contact the Canadian Patent Office


NOM DU FICHIER / FILE NAME :

NOTE POUR LE TOME / VOLUME NOTE:



## WHAT IS CLAIMED IS:

1. A compound of Formula I



I

wherein,

W is CR<sup>3</sup>R<sup>4</sup>,

----- represents a single or a double bond, wherein when ----- is a double bond, then X is CR<sup>5</sup> and Y is CR<sup>6</sup>, and when ----- is a single bond, then X is O, CR<sup>7</sup>R<sup>8</sup> or NR<sup>9</sup> and Y is C(=O), SO<sub>2</sub>, CR<sup>10</sup>R<sup>11</sup> or O, and

Z is C or N;

R<sup>3</sup> is hydrogen;

R<sup>4</sup> is hydrogen;

R<sup>5</sup> is hydrogen;

R<sup>6</sup> is hydrogen;

R<sup>7</sup> is hydrogen or C1-C6 alkyl;

R<sup>8</sup> is hydrogen;

R<sup>9</sup> is hydrogen, C1-C6 alkyl, hydroxyC1-C6 alkyl-, (C1-C6 alkyl)SO<sub>2</sub>-, (C1-C6 alkoxy)C1-C6 alkyl, C3-C6 cycloalkyl, (C3-C6 cycloalkyl)C1-C6 alkyl-, hetCyc<sup>1</sup>, (hetCyc<sup>1</sup>)C1-C6 alkyl-, Ar<sup>1</sup>, (Ar<sup>1</sup>)C1-C6 alkyl-, hetAr<sup>1</sup>, (hetAr<sup>1</sup>)C1-C6 alkyl-, (C1-C6 alkyl)C(=O)-, (fluoroC1-C6 alkyl)C(=O)-, H<sub>2</sub>NC(=O)-, Ar<sup>1</sup>CH<sub>2</sub>C(=O)- or (C3-C6 cycloalkyl)C(=O)-;

R<sup>10</sup> is hydrogen, hydroxy, C1-C6 alkoxy, or R<sup>a</sup>R<sup>b</sup>N- where R<sup>a</sup> and R<sup>b</sup> are independently H or C1-C6 alkyl;

R<sup>11</sup> is hydrogen, C1-C6 alkyl, or fluoroC1-C6 alkyl;

hetCyc<sup>1</sup> is a 5-6 membered saturated heterocyclic ring having 1-2 ring heteroatoms independently selected from N and SO<sub>2</sub>, wherein said heterocyclic ring is optionally substituted with C1-C6 alkyl or (Ar<sup>2</sup>)C1-C6 alkyl-;

Ar<sup>1</sup> is phenyl optionally substituted with one or more substituents independently selected from halogen and C1-C6 alkoxy;

hetAr<sup>1</sup> is a 5-6 membered heteroaryl ring having 1-2 ring nitrogen atoms, wherein said heteroaryl ring is optionally substituted with C1-C6 alkyl;

Ar<sup>2</sup> is phenyl optionally substituted with one or more substituents independently selected from halogen and C1-C6 alkoxy;

Ring A, including the atoms at the points of attachment to the ring containing Z, Y, X, and W, is:

(i) a monocyclic aryl ring when Z is C, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH-,

(ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O when Z is C, or when Z is N, a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said rings are optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or

(iii) when Z is N, a bicyclic heteroaryl ring having one ring nitrogen atom which is Z, wherein said ring is optionally substituted with R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH-;

R<sup>c</sup> is C3-C6 cycloalkyl or (C3-C6 cycloalkyl)CH<sub>2</sub>-;

R<sup>d</sup> is C1-C6 alkyl, C1-C6 alkoxy, Cyc<sup>1</sup>, Cyc<sup>1</sup>CH<sub>2</sub>-, phenyl or hetAr<sup>2</sup>;

Cyc<sup>1</sup> is C3-C6 cycloalkyl optionally substituted with C1-C6 alkyl;

hetAr<sup>2</sup> is (i) pyrazolyl optionally substituted with 1-2 substituents independently selected from C1-C6 alkyl, fluoroC1-C6 alkyl, (di-C1-C6 alkyl)NC(=O)CH<sub>2</sub>-, (hetCyc<sup>2</sup>)CH<sub>2</sub>CH<sub>2</sub>-, phenyl, and (C1-C6 alkoxy)C1-C6 alkyl, or (ii) pyridyl;

hetCyc<sup>2</sup> is a 6-membered saturated heterocyclic ring having 2 ring heteroatoms independently selected from N and O;

R<sup>x</sup> is hydrogen, C1-C6 alkyl, fluoroC1-C6 alkyl, Cyc<sup>2</sup>, Cyc<sup>3</sup>, hetCyc<sup>3</sup>, Ar<sup>3</sup> or hetAr<sup>3</sup>;

Cyc<sup>2</sup> is a 3-6 membered saturated carbocyclic ring optionally substituted with 1-2 substituents independently selected from halogen, C1-C6 alkyl, hydroxy, hydroxyC1-C6 alkyl, and (C1-C6 alkoxy)C(=O)-;

Cyc<sup>3</sup> is cyclopentenyl;

hetCyc<sup>3</sup> is a 5-membered saturated heterocyclic ring having a ring heteroatom selected from N and O and optionally substituted with (C1-C6 alkyl)C(=O)-;

Ar<sup>3</sup> is phenyl optionally substituted with (C1-C6 alkoxy)C(=O)-;

hetAr<sup>3</sup> is a 5-6 membered heteroaryl ring having 1-2 ring nitrogen atoms and optionally substituted with C1-C6 alkyl or C1-C6 alkoxy; and

R<sup>y</sup> is hydrogen or halogen;


or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, wherein Ring A is a monocyclic aryl ring when Z is C, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH-, or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 1, wherein Ring A is a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O when Z is C, or when Z is N, a 5-membered heteroaryl ring having two ring nitrogen atoms, one of which is Z, wherein said rings are optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 1, wherein Ring A is a bicyclic heteroaryl ring having one ring nitrogen atom that results when Z is N, wherein said ring is optionally substituted with R<sup>d</sup>NHC(=O)-, (C1-C6 alkoxy)C(=O)-, C1-C6 alkyl, C1-C6 alkoxy, fluoroC1-C6 alkyl, CN, halogen, or (C1-C6 alkyl)C(=O)NH-, or a pharmaceutically acceptable salt thereof.

5. A compound according to claim 1, wherein W is CR<sup>3</sup>R<sup>4</sup>, X is CR<sup>7</sup>R<sup>8</sup>, Y is CR<sup>10</sup>R<sup>11</sup>, Z is C, and ----- is a single bond, wherein the compound is represented by the structure of Formula I-H:



wherein Ring A is (i) a monocyclic aryl ring, wherein said ring is optionally substituted with halogen, C3-C6 cycloalkyl, C1-C6 alkoxy, R<sup>c</sup>NHC(=O)-, R<sup>c</sup>C(=O)NH-, or R<sup>c</sup>NHC(=O)NH- or (ii) a 5-membered monocyclic heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is

optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or a pharmaceutically acceptable salt thereof.


6. A compound according to claim 5, wherein Ring A is a 5-membered heteroaryl ring having 2 ring heteroatoms independently selected from N and O, wherein said ring is optionally substituted with C1-C6 alkyl or C3-C6 cycloalkyl, or a pharmaceutically acceptable salt thereof.

7. A compound according to claim 5 or 6, wherein R<sup>7</sup> and R<sup>8</sup> are hydrogen, or a pharmaceutically acceptable salt thereof.

8. A compound according to any one of claims 1-7, wherein R<sup>x</sup> is C1-C6 alkyl, or a pharmaceutically acceptable salt thereof.


9. A compound according to claim 1, selected from Examples 1-151, or a pharmaceutically acceptable salt thereof.

10. A compound according to claim 1, of the formula:




, or a pharmaceutically acceptable salt thereof.

11. A compound according to claim 1, of the formula:



12. A pharmaceutical composition, comprising a compound according to any one of claims 1-10 in admixture with a pharmaceutically acceptable diluent or carrier.
13. A method for treating cancer in a subject in need thereof, the method comprising administering a pharmaceutical composition comprising a therapeutically effective amount of a compound according to any one of claims 1-11 or a pharmaceutically acceptable salt thereof or a pharmaceutical composition according to claim 12.
14. The method of claim 13, wherein the cancer is a RET-associated cancer.
15. The method of claim 13, wherein the cancer is selected from the group consisting of: lung cancer, papillary thyroid cancer, medullary thyroid cancer, differentiated thyroid cancer, recurrent thyroid cancer, refractory differentiated thyroid cancer, multiple endocrine neoplasia type 2A or 2B (MEN2A or MEN2B, respectively), pheochromocytoma, parathyroid hyperplasia, breast cancer, colorectal cancer, papillary renal cell carcinoma, ganglioneuromatosis of the gastroenteric mucosa, and cervical cancer.
16. The method of claim 14, wherein the cancer is medullary thyroid cancer.
17. The method of claim 14, wherein the cancer is lung cancer and the lung cancer is small cell lung carcinoma, non-small cell lung cancer, bronchioles lung cell carcinoma, RET fusion lung cancer, or lung adenocarcinoma.
18. The method of claim 17, wherein the lung cancer is RET fusion lung cancer.
19. The method of any one of claims 13-18, further comprising administering an additional therapy or therapeutic agent to the subject.
20. Use of a compound or a pharmaceutically acceptable salt thereof according to any one of claims 1-10 for the manufacture of a medicament for treating a RET-associated cancer in a subject.

21. The use of claim 20, wherein the RET-associated cancer is selected from the group consisting of: lung cancer, papillary thyroid cancer, medullary thyroid cancer, differentiated thyroid cancer, recurrent thyroid cancer, refractory differentiated thyroid cancer, multiple endocrine neoplasia type 2A or 2B (MEN2A or MEN2B, respectively), pheochromocytoma, parathyroid hyperplasia, breast cancer, colorectal cancer, papillary renal cell carcinoma, ganglioneuromatosis of the gastroenteric mucosa, and cervical cancer.
22. The use of claim 21, wherein the RET associated cancer is medullary thyroid cancer.
23. The use of claim 21, wherein the RET associated cancer is lung cancer and the lung cancer is small cell lung carcinoma, non-small cell lung cancer, bronchioles lung cell carcinoma, RET fusion lung cancer, or lung adenocarcinoma.
24. The use of claim 22, wherein the lung cancer is RET fusion lung cancer.
25. A compound, or a pharmaceutically acceptable salt thereof, according to any one of claims 1-11 for use in therapy.
26. A compound, or a pharmaceutically acceptable salt thereof, according to any one of claims 1-11 for use in the treatment of a RET-associated cancer.



(I)



(AA)