



(51) International Patent Classification:

C07K 16/46 (2006.01) C07K 16/12 (2006.01)
A61K 39/395 (2006.01) C07K 16/24 (2006.01)
A61P 35/00 (2006.01) C07K 16/28 (2006.01)
A61P 37/02 (2006.01) C07K 16/30 (2006.01)

(21) International Application Number:

PCT/CA2013/000471

(22) International Filing Date:

10 May 2013 (10.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/645,555 10 May 2012 (10.05.2012) US

(71) Applicant: **ZYMEWORKS INC.** [CA/CA]; 540 - 1385
West 8th Avenue, Vancouver, British Columbia V6H 3V9
(CA).

(72) Inventors: **SPRETER VON KREUDENSTEIN, Thomas**; 1007-1330 Harwood Street, Vancouver, British
Columbia V6E 1S8 (CA). **ESCOBAR-CABRERA, Eric**;
8562 Woodridge Place, Burnaby, British Columbia V5A
4B3 (CA). **NG, Gordon Yiu Kon**; 1535 West 64th Avenue,
Vancouver, British Columbia V6P 2N8 (CA). **DIXIT, Surjit Bhimarao**; 51, 11651 Kingfisher Drive, Richmond,

British Columbia V7E 3N5 (CA). **LARIO, Paula Irene**;
201 - 2080 East Kent Avenue South, Vancouver, British
Columbia V5P 4Y2 (CA). **POON, David Kai Yuen**; 9320
Railway Avenue, Richmond, British Columbia V7E 2B6
(CA). **D'ANGELO, Igor Edmondo Paolo**; 15 Alder
Drive, Port Moody, British Columbia, V3H 5M3 (CA).

(74) Agents: **SALISBURY, Clare** et al.; Gowling Lafleur
Henderson LLP, 2300 - 550 Burrard Street, Vancouver,
British Columbia V6C 2B5 (CA).


(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,

[Continued on next page]

(54) Title: HETEROMULTIMER CONSTRUCTS OF IMMUNOGLOBULIN HEAVY CHAINS WITH MUTATIONS IN THE FC DOMAIN

Figure 1



(57) Abstract: Provided herein are isolated heteromultimers comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to that of a native Fc homodimer; and wherein said isolated heteromultimer is devoid of immunoglobulin light chains and optionally devoid of immunoglobulin CH1 region. These novel molecules comprise complexes of heterogeneous components designed to alter the natural way antibodies behave and that find use in therapeutics.



TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, **Published:**

EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,

LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,

GW, ML, MR, NE, SN, TD, TG).

— *with international search report (Art. 21(3))*

HETEROMULTIMER CONSTRUCTS OF IMMUNOGLOBULIN HEAVY CHAINS WITH MUTATIONS IN THE F_c DOMAIN

5

Cross Reference to Related Applications

This application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Patent Application No. 61/645,555, filed May 10, 2012 which is herein incorporated by reference in its entirety.

10

Field of the Invention

The present disclosure generally provides polypeptide heterodimers, compositions thereof, and methods for making and using such polypeptide heterodimers. More specifically, provided herein are thermo-stable antibody constructs, said constructs comprising heterodimeric F_c domain, wherein said constructs are devoid of immunoglobulin light chains. In certain embodiments, the antibody constructs are multi-specific and/or multivalent. In certain embodiments, the antibody constructs are devoid of an immunoglobulin first constant (CH1) region.

15

20

Background of the Invention

Bi-specific therapeutics are antibody-based molecules that can simultaneously bind two separate and distinct targets or different epitopes of the same antigen. Bi-specific antibodies are comprised of the immunoglobulin domain based entities and try to structurally and functionally mimic components of the antibody molecule. One use of bi-specific antibodies has been to redirect cytotoxic immune effector cells for enhanced killing of tumor cells, such as by antibody dependent cellular cytotoxicity (ADCC). In this context, one arm of the bi-specific antibody binds an antigen on the tumor cell, and the other binds a determinant expressed on effector cells. By cross-linking tumor and effector cells, the bi-specific antibody not only brings the effector cells within the proximity of the tumor cells but also simultaneously triggers their activation, leading to effective tumor cell-killing. Bi-specific antibodies have also been used to enrich chemo- or radiotherapeutic agents in tumor tissues to minimize detrimental effects to normal tissue. In this setting, one arm of the bi-specific antibody binds an antigen expressed on the cell targeted for destruction, and the other arm delivers a chemotherapeutic drug, radioisotope, or toxin. Going beyond bi-specifics,

25

30

35

there is a need for protein therapeutics that achieve their efficacy by targeting multiple modalities concurrently. Such complex and novel biological effects can be obtained with protein therapeutics with multi-target binding and multi-functional aspects designed into the protein.

5

A robust scaffold that provides a framework to fuse other functional war-heads or target protein binding domains in order to design these multifunctional and multi-target binding therapeutics is required. Ideally, the scaffold should not only provide the framework but also makes available a number of other therapeutically relevant and valuable features to the designed therapeutic. A major obstacle in the general development of antibody based bi-specific and multifunctional therapeutics has been the difficulty of producing materials of sufficient quality and quantity for both preclinical and clinical studies.

10

15

Antigen-binding polypeptides that lack a light chain (i.e. comprising a single variable domains) are known in the art and include those derived from camelids or cartilaginous fish, for example. These types of antigen-binding polypeptides have been shown to have many advantages as antigen-binding fragments, for example, they are more thermostable, can penetrate tumors and cross the blood-brain-barrier, and they can bind to epitopes that other antigen-binding polypeptide fragments (such as Fabs and scFvs) cannot. Thus, monovalent or bi-specific antibodies that have this type of antigen-binding polypeptide fragment have been developed. However, existing technologies for preparing such monovalent or bi-specific antibodies are not ideal, and results in products that lack the purity and/or stability required to manufacture them in the amounts and quality necessary for therapeutic and clinical applications. There remains a need in the art for polypeptide constructs that comprise single variable domains as protein binding domains that are linked to a variant Fc region, said variant Fc comprising CH3 domains, which have been modified to select for heterodimers with an increased stability and purity, with Fc effector activities.

20

25

30

SUMMARY OF THE INVENTION

35

There is provided according to one aspect an isolated heteromultimer comprising: at least one immunoglobulin single domain antigen-binding construct attached to at

least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and wherein said isolated heteromultimer is devoid of immunoglobulin light chains.

5

Provided herein is an isolated heteromultimer comprising: at least one single domain antigen-binding construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide; wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations that promote the formation of said heterodimer Fc region with stability comparable to a native homodimeric Fc region; and wherein said isolated heteromultimer is devoid of immunoglobulin light chain and immunoglobulin first constant (CH1) region.

10

15

In certain embodiments is the isolated heteromultimer provided herein, comprising one single domain antigen-binding construct attached to one monomeric Fc polypeptide.

20

In some embodiments, is the isolated heteromultimer described herein comprising one single domain antigen-binding construct attached to one monomeric Fc polypeptide, and a second single domain antigen-binding construct attached to the other monomeric Fc polypeptide. In some embodiments is the isolated heteromultimer provided herein, wherein both single-domain antigen-binding constructs bind to the same epitope. In some embodiments is the isolated heteromultimer provided herein, wherein said one single domain antigen-binding construct binds to one epitope, and the second single domain antigen-binding construct binds to a different epitope. In some embodiments, the single domain antigen binding construct is selected from single domain antibodies (sdAb or VH), camelid nanobodies (V_HH), cartilaginous fish (V_{NAR}), SH3-derived fynomers, and fibronectin-derived binding domains. Provided in some embodiments herein is an isolated heteromultimer described herein, wherein the single domain antigen-binding construct is a camelid nanobody (V_HH). In certain embodiments, the single domain antigen-binding construct binds to one or more cytokines or chemokines selected from IL2, IFN α -2a/b, IFN-1a/b, IL-21, IL-17a, TNF, IL23, VEGF, and ANG2. In some embodiments, the single domain antigen-binding construct binds to one or more tumor associated antigens such as EpCam, EGFR, VEGFR, CEA, or GP100. In

25

30

35

select embodiments, the single domain antigen-binding construct binds to one or more immunoregulatory antigens such as CD16, CD30, CD137, CD22, CD52, CD80, CD23, CD2, CD4, CD40, KIR, CD32b, CD25, LAG3, or B7-H3. In an embodiment, the single domain antigen-binding construct binds to one or more bacterial toxins
5 such as Clostridium difficile toxin A, Clostridium difficile toxin B.

Provided herein are isolated heteromultimer described herein wherein the single domain antigen-binding construct binds to EGFR1. In an embodiment the single domain antigen-binding construct binds to the EGFR1 mutated variant EGFRvIII.

10 Provided herein is an isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant CH3 region comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and wherein said isolated heteromultimer is devoid of immunoglobulin light chains.

15 Provided is an isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant constant domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and wherein said isolated
20 heteromultimer is devoid of immunoglobulin light chains.

Also provided is an isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant constant domain comprising amino acid mutations that promote the formation of said heterodimer with
25 stability comparable to a native homodimeric Fc region; and wherein said isolated heteromultimer is devoid of immunoglobulin light chain and immunoglobulin first constant (CH1) region. Provided herein is the isolated heteromultimer described herein wherein said single domain antigen-binding construct is derived from a camelid or a cartilaginous fish. In an embodiment is the isolated heteromultimer
30 described herein wherein said camelid is a llama. Also provided is the isolated heteromultimer described herein, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability wherein said amino acid mutations promote the formation of heterodimer Fc region with increased stability as compared to a CH3

domain that does not comprise amino acid mutations, and wherein the variant CH3 domain has a melting temperature (T_m) of about 70 °C or greater. In an embodiment is the isolated heteromultimer described herein, wherein the heterodimer Fc region does not comprise an additional disulfide bond in the CH3 domain relative to a wild type Fc region. In a further embodiment is the isolated heteromultimer described herein, wherein the heterodimer Fc region comprises an additional disulfide bond in the variant CH3 domain relative to a wild type Fc region, with the proviso that the melting temperature (T_m) of about 70 °C or greater for the CH3 domain is in the absence of the additional disulfide bond. In another embodiment is the isolated heteromultimer described herein, wherein the heterodimer Fc region has a purity greater than about 90%. In a further embodiment is the isolated heteromultimer described herein, wherein the heterodimer Fc region has a purity of about 98% or greater. In another embodiment is the isolated heteromultimer described herein, wherein the T_m is about 74 °C or greater. In an embodiment is the isolated heteromultimer, wherein a first Fc polypeptide comprises amino acid modification at positions F405 and Y407 and a second Fc polypeptide comprises amino acid modification at position T394. In another embodiment is the isolated heteromultimer wherein the first Fc polypeptide comprises one or more amino acid modifications selected from L351Y, Y405A and Y407V, and the second Fc polypeptide comprises one or more amino acid modifications selected from T366L, T366I, K392L, K392M and T394W. In a further embodiment is the isolated heteromultimer described herein, wherein a first Fc polypeptide comprises amino acid modifications at positions D399 and Y407 and a second Fc polypeptide comprises amino acid modification at positions K409 and T411.

Provided herein is an isolated heteromultimer described herein, said heterodimer Fc region comprising: a first monomeric Fc polypeptide comprising a first modified CH3 domain comprising at least three amino acid modifications as compared to a wild-type CH3 domain polypeptide, and a second monomeric Fc polypeptide comprising a second modified CH3 domain comprising at least three amino acid modifications as compared to a wild-type CH3 domain polypeptide; wherein one of said first and second CH3 domain comprises an amino acid modification of K392J wherein J is selected from L, I, M or an amino acid with a side chain volume not substantially larger than the side chain volume of K; wherein said first and second modified CH3 domain polypeptides preferentially form a heterodimeric CH3 domain with a melting temperature (T_m) of at least about 74°C and a purity of at least 95%; and wherein at

least one amino acid modification is not of an amino acid which is at the interface between said first and said second CH3 domain polypeptides.

In an embodiment is the isolated heteromultimer, comprising at least one T350X modification, wherein X is a natural or non-natural amino acid selected from valine, isoleucine, leucine, methionine, and derivatives or variants thereof. In a further embodiment is the isolated heteromultimer comprising at least one T350V modification. In some embodiments each of said first and second Fc polypeptides further comprises a T350V modification. In some embodiments the Fc heterodimer domain has a T_m of about 77°C or greater.

- 10 In some embodiments is the isolated heteromultimer described herein, wherein at least one monomeric Fc polypeptide comprises the modification S400Z, wherein Z is selected from a positively charged amino acid and a negatively charged amino acid. In an embodiment is the isolated heteromultimer, wherein said first Fc polypeptide comprises an amino acid modification selected from S400E, S400D, S400K and S400R. In another embodiment, one of said first and second Fc polypeptides comprises the amino acid modification selected from S400E and S400R, and the other Fc polypeptide comprises an amino acid modification at position N390. In a further embodiment is the isolated heteromultimer, comprising the modification N390Z, wherein Z is selected from a positively charged amino acid and a negatively charged amino acid. In an embodiment is the isolated heteromultimer described herein, said second Fc polypeptide comprising the amino acid modification N390R or N390K.

- 25 In an embodiment is provided an isolated heteromultimer described herein, wherein said first Fc polypeptide is a modified CH3 domain polypeptide comprising the amino acid modification S400E and said second Fc polypeptide is a modified CH3 domain polypeptide comprising the amino acid modification N390R. In an embodiment is the isolated heteromultimer described herein, one said Fc polypeptide comprising the amino acid modification Q347R and the other Fc polypeptide comprising the amino acid modification K360E.

- 30 Provided is an isolated heteromultimer described herein, wherein the Fc polypeptide comprises at least one amino acid modification selected from T366V, T366I, T366A, T366M, T366L, K409F, T411E and T411D, and the second Fc polypeptide comprises at least one amino acid modification selected from L351Y, Y407A, Y407I, Y407V, D399R and D399K. In an embodiment is the isolated heteromultimer wherein the

heterodimer Fc region further comprises a variant CH2 domain comprising asymmetric amino acid modifications to promote selective binding of a Fcgamma receptor.

5 In an embodiment is the isolated heteromultimer described herein, wherein the variant CH2 domain selectively binds FcgammaIIIa receptor as compared to wild-type CH2 domain.

In some embodiments is the isolated heteromultimer provided herein comprising a Fc construct based on a type G immunoglobulin (IgG). In certain embodiments is the isolated heteromultimer, wherein said IgG is one of IgG1, IgG2 IgG3 and IgG4. In
10 some embodiments is the isolated heteromultimer comprising a Fc construct based on Immunoglobulin M (IgM), Immunoglobulin A (IgA), Immunoglobulin D (IgD), or Immunoglobulin E (IgE).

Provided in some embodiments is the isolated heteromultimer described herein, wherein said heteromultimer is a bispecific antibody or a multispecific antibody.

15 Provided in some embodiments is the isolated heteromultimer described herein, wherein at least one single domain antigen binding construct binds EGFR or EGFRvIII. In some embodiments is the isolated heteromultimer described herein wherein said EGFR or EGFRvIII binding construct is derived from an antibody or fragment thereof. In certain embodiments, said EGFR or EGFRvIII binding construct
20 is a heavy chain antibody construct. In some embodiments said heavy chain antibody construct is a camelid construct. In an embodiment, said camelid construct comprises the sequence shown in Figure 44.

Provided is a composition comprising the isolated heteromultimer described herein and a pharmaceutically acceptable carrier.

25 Provided is a mammalian host cell comprising nucleic acid encoding the isolated heteromultimer described herein.

In an embodiment is the isolated heteromultimer described herein, wherein the single domain antigen-binding construct competes for binding with at least one therapeutic antibody. In an embodiment is the isolated heteromultimer, wherein said at least one
30 therapeutic antibody is selected from the group consisting of abagovomab, adalimumab, alemtuzumab, aurograb, bapineuzumab, basiliximab, belimumab, bevacizumab, briakinumab, canakinumab, catumaxomab, certolizumab pegol,

cetuximab, daclizumab, denosumab, efalizumab, galiximab, gemtuzumab
ozogamicin, golimumab, ibritumomab tiuxetan, infliximab, ipilimumab, lumiliximab,
mepolizumab, motavizumab, muromonab, mycograb, natalizumab, nimotuzumab,
ocrelizumab, ofatumumab, omalizumab, palivizumab, panitumumab, pertuzumab,
5 ranibizumab, reslizumab, rituximab, teplizumab, tocilizumab/atlizumab, tositumomab,
trastuzumab, ProxiniumTM, RencarexTM, ustekinumab, and zalutumumab.

In an embodiment is a method of treating cancer in a patient having a cancer
characterized by a cancer antigen, said method comprising administering to said
patient a therapeutically effective amount of an isolated heteromultimer described
10 herein. In an embodiment is the method of treating cancer, wherein said cancer is
characterized by overexpression of EGFR or EGFRvIII.

Provided is a method of treating cancer cells expressing EGFR or EGFRvIII,
comprising contacting said cells with an amount of a heteromultimer provided herein.
In some embodiments is the said cancer cell is at least one of a breast cancer cell, a
15 lung cancer cell, an anal cancer cell and a glioblastoma.

In some embodiments is the method of treating cancer described herein, comprising
administration of said heteromultimer, in addition to another therapeutic molecule. In
some embodiments, said therapeutic molecule is conjugated to the heteromultimer.

Provided is a method of treating immune disorders in a patient having an immune
20 disorder characterized by an immune antigen, said method comprising administering
to said patient a therapeutically effective amount of an isolated heteromultimer
described herein.

In an aspect provided herein is an isolated heteromultimer comprising: at least one
immunoglobulin single domain antigen-binding construct attached to at least one
25 monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a
variant CH3 domain comprising amino acid mutations that promote the formation of
said heterodimer with stability comparable to a native homodimeric Fc; and wherein
said isolated heteromultimer is devoid of immunoglobulin first constant (CH1) region
and immunoglobulin light chains.

30 In an aspect provided herein is an isolated heteromultimer comprising: at least one
immunoglobulin single domain antigen-binding construct attached to at least one
monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a

variant CH3 domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and wherein said isolated heteromultimer is devoid of immunoglobulin first and second constant domains (CH1 & CH2) and immunoglobulin light chains.

5

In certain embodiments of the isolated heteromultimers provided herein, the variant CH3 domain has a melting temperature (T_m) of about 70 °C or greater. In certain embodiments, the variant CH3 domain has a melting temperature (T_m) of at least about 75 °C. In some embodiments, the variant CH3 domain has a melting

10

In some embodiments of the isolated heteromultimers provided herein, the heterodimer Fc region further comprises a variant CH2 domain comprising at least one asymmetric amino acid modification to promote selective binding to certain Fcγ receptors. In one embodiment the variant CH2 domain selectively binds Fcγ3a receptor as compared to wild-type CH2 domain.

15

In a particular embodiment of the invention, the heteromultimers described herein are the product of the expression in a prokaryotic or in a eukaryotic host cell, of a DNA or of a cDNA having the sequence of an immunoglobulin devoid of immunoglobulin first constant (CH1) region. In certain embodiments, the at least one immunoglobulin heavy chain variable region is from an immunoglobulin devoid of light chains, said immunoglobulin obtainable from lymphocytes or other cells of Camelids such as but not restricted to Dromedaries, Bactrian camels, llamas, alpacas, vicuñas, and guanacos. In a specific embodiment, the at least one immunoglobulin heavy chain variable region is from an immunoglobulin devoid of light chains, said immunoglobulin obtained from lymphocytes or other cells of a llama.

20

25

In a particular embodiment of the invention, the heteromultimers described herein are the product of the expression in a prokaryotic or in a eukaryotic host cell, of a DNA or of a cDNA. In certain embodiments, at least one immunoglobulin heavy chain variable region and/or Fc heterodimer is from an immunoglobulin devoid of light chains obtainable from cartilaginous fishes such as but not restricted to sharks, rays, skates, ghost sharks, ratfish, elephantfish and rabbitfish. In a specific embodiment, at least one immunoglobulin heavy chain variable region and/or Fc heterodimer is from an immunoglobulin devoid of light chains, said immunoglobulin obtained from a shark.

30

35

There is provided in another aspect an isolated heteromultimer comprising at least one immunoglobulin single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations, wherein said isolated heteromultimer is devoid of immunoglobulin first constant (CH1) region, immunoglobulin light chains and optionally devoid of immunoglobulin second constant (CH2) region, wherein the variant CH3 domain has a melting temperature (T_m) of about 70 °C or greater, and wherein said variant CH3 domain results in the formation of heterodimer Fc region with stability comparable to a CH3 domain in native IgG1 antibody.

In one embodiment, the heterodimer Fc region does not comprise an additional disulfide bond in the CH3 domain relative to a wild type Fc region. In an alternative embodiment, the heterodimer Fc region comprises at least one additional disulfide bond in the variant CH3 domain relative to a wild type Fc region, with the proviso that the melting temperature (T_m) of about 70 °C or greater is in the absence of the additional disulfide bond. In another embodiment, the heterodimer Fc region comprises at least one additional disulfide bond in the variant CH3 domain relative to a wild type Fc region, and wherein the variant CH3 domain has a melting temperature (T_m) of about 77.5 °C or greater.

Provided in one embodiment, an isolated heteromultimer comprising at least one immunoglobulin single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations, wherein said isolated heteromultimer is devoid of immunoglobulin light chains and immunoglobulin first constant (CH1) domain and optionally devoid of the immunoglobulin second constant (CH2) domain, wherein the variant CH3 domain has a melting temperature (T_m) of about 70 °C or greater and the heterodimer Fc region is formed with a purity greater than about 90%, or the heterodimer Fc region is formed with a purity of about 95% or greater or the heterodimer Fc region is formed with a purity of about 98% or greater.

In certain embodiments, the immunoglobulins described herein, which are devoid of light chains, are such that the variable domains of their heavy chains (V_H) have properties differing from those of the V_H in four-chain immunoglobulin. In some embodiments, the variable domain of a heavy-chain immunoglobulin described

herein has no interaction sites for V_L such as in the case of heavy chain immunoglobulins from cartilaginous fish. In some embodiments, the variable domain of a heavy-chain immunoglobulin described herein has no normal interaction sites with V_L or with C_{H1} domain, neither of which exists in heavy chain immunoglobulins from Camelids and some cartilaginous fish.

In certain embodiments of the heteromultimer described herein, the at least one immunoglobulin heavy chain is attached to at least one monomer of the heterodimer Fc region by means of a linker. In certain embodiments, the at least one single domain antigen-binding construct is attached to at least one monomer of the heterodimer Fc region by means of a hinge region. Linkers and/or hinge regions of variable lengths are utilized in the heteromultimers provided herein. One of skill in the art can appreciate that the length of the linker and/or hinge region will participate to the determination of the distance separating the antigen binding sites. In certain embodiments provided herein, the hinge region comprises from 0 to 50 amino acids. In certain embodiments, the sequence of hinge region is as follows:

GTNEVCKCPKCP

In an embodiment, the sequence of the hinge region is:

EPKIPQPQPKPQPQPQPQPKPQPKPEPECTCPKCP

In certain embodiments, at least one single domain antigen-binding construct and/or modified Fc region utilized in the heteromultimer constructs described herein comprises type G immunoglobulins for instance immunoglobulins which are defined as immunoglobulins of class 2 (IgG2) or immunoglobulins of class 3 (IgG3). In some embodiments, at least one single domain antigen-binding construct and/or modified Fc region utilized in the heteromultimer constructs described herein comprises immunoglobulin M, or IgM. In some embodiments, at least one single domain antigen-binding construct and/or modified Fc region utilized in the heteromultimer constructs described herein comprises immunoglobulin A, or IgA. In some embodiments, at least one single domain antigen-binding construct and/or modified Fc region utilized in the heteromultimer constructs described herein comprises immunoglobulin D, or IgD. In some embodiments, at least one single domain antigen-binding construct and/or modified Fc region utilized in the heteromultimer constructs described herein comprises Immunoglobulin E, or IgE.

In certain embodiments, the heteromultimers described herein comprise an Fc portion that is derived from IgG (e.g. an IgG1, IgG2, IgG3 or IgG4) or IgA (or IgM,

IgD). According to one specific aspect certain embodiments comprise a modified "IgE-derived Fc portion" (i.e. an Fc portion that is derived from IgE).

5 In certain embodiments, the hinge connecting the variable domain of the heavy chain and the Fc portion of the heavy chain is derived from the hinge sequence of an IgG isotype (e.g. an IgG1, IgG2, IgG3 or IgG4) or IgA (or IgM, IgD, IgE).

10 In certain embodiments described herein, the variable heavy chains may be a domain antibody (or an amino acid sequence that is suitable for use as a domain antibody), a single domain antibody (or an amino acid sequence that is suitable for use as a single domain antibody), a "dAb" (or an amino acid sequence that is suitable for use as a dAb) or a Nanobody (as defined herein, and including but not limited to a V_{HH} sequence); other single variable domains, or any suitable fragment of any one thereof. For a general description of (single) domain antibodies, reference is
15 also made to the prior art cited above, as well as to EP 0 368 684. For the term "dAb's", reference is for example made to Ward et al. (Nature 1989 Oct. 12; 341 (6242): 544-6), to Holt et al. (Trends Biotechnol., 2003, 21(11):484-490); as well as to for example WO 04/068820, WO 06/030220, WO 06/003388 and other published patent applications of Domantis Ltd. In some embodiments, the variable heavy
20 chains comprise single domain antibodies or single variable domains can be derived from certain species of shark (for example, the so-called "IgNAR domains", see for example WO 05/18629).

25 Provided herein are isolated heteromultimers comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region;
wherein the heterodimer Fc region comprises a variant constant domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and wherein said isolated heteromultimer is
30 devoid of immunoglobulin light chains.

In an embodiment is provided an isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant
35 constant domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc region; and

wherein said isolated heteromultimer is devoid of immunoglobulin light chain and immunoglobulin first constant (CH1) region.

In certain embodiments, the heteromultimers described herein comprise an Fc portion or constant domain that is derived from antibodies from any animal origin (e.g. human, murine, donkey, sheep, rabbit, goat, guinea pig, camel, horse, or chicken) which maintain sequence similarity of the constant domains (Figures 40A-40B). In a specific embodiment the constant domain is derived from cartilaginous fishes such as but not restricted to sharks, rays, skates, ghost sharks, rattfish, elephantfish and rabbitfish. In specific embodiments, the heteromultimers described herein comprise a constant domain that is derived from antibodies from camels such as but not restricted to Dromedaries, Bactrian camels, llamas, alpacas, vicuñas, and guanacos.

single domain antigen-binding constructs according to certain embodiments provided herein are obtainable by purification from serum of camelids such as but not restricted to llamas using processes of purification well known in the art.

In certain embodiments provided herein, the variable region of immunoglobulins of the heteromultimers comprises frameworks (FW) and complementarity determining regions (CDR), especially 4 frameworks and 3 complementarity regions. It is distinguished from the four-chain immunoglobulins especially by the fact that this variable region can itself contain an antigen binding site or several, without contribution of the variable region of a light chain which is absent.

Also provided in certain embodiments is an isolated heteromultimer comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising one or more amino acid mutations that result in the formation of heterodimer Fc region with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of about 70 °C or greater or the T_m is about 71°C or greater or the T_m is about 74°C or greater. In another embodiment, the heterodimer Fc region is formed in solution with a purity of about 98% or greater, and T_m about 73°C or wherein the heterodimer Fc region is formed with a purity of about 90% or greater, and T_m about 75°C.

Provided in certain embodiments is an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a

heterodimer Fc region, wherein the heterodimer Fc region comprises a first and a second CH3 domain polypeptides, wherein at least one of said first and second CH3 domain polypeptides comprises amino acid modification T350V. Provided in certain embodiments is an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modification T350V and a second CH3 domain polypeptide also comprising amino acid modification T350V. Provided in certain embodiments is an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modification at positions F405 and Y407 and a second CH3 domain polypeptide comprising amino acid modification at position T394. In certain embodiments, a first CH3 domain polypeptide comprises amino acid modifications at positions D399 and Y407 and a second CH3 domain polypeptide comprises amino acid modification at positions K409 and T411. Provided in certain embodiments is an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modifications L351Y and Y407A and a second CH3 domain polypeptide comprising amino acid modifications T366A and K409F. In one aspect, the first CH3 domain polypeptide or the second CH3 domain polypeptide comprises a further amino acid modification at position T411, D399, S400, F405, N390, or K392. The amino acid modification at position T411 is selected from T411N, T411R, T411Q, T411K, T411D, T411E or T411W. The amino acid modification at position D399 is selected from D399R, D399W, D399Y or D399K. The amino acid modification at position S400 is selected from S400E, S400D, S400R, or S400K. The amino acid modification at position F405 is selected from F405I, F405M, F405T, F405S, F405V or F405W. The amino acid modification at position N390 is selected from N390R, N390K or N390D. The amino acid modification at position K392 is selected from K392V, K392M, K392R, K392L, K392F or K392E.

In certain embodiments is provided an isolated heteromultimer comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modifications T350V and L351Y and a second CH3 domain polypeptide also comprising amino acid modifications T350V and L351Y.

In another embodiment is provided an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modification Y407A and a second CH3 domain polypeptide comprising amino acid modifications T366A and K409F. In one aspect the first CH3 domain polypeptide or the second CH3 domain polypeptide comprises further amino acid modifications K392E, T411E, D399R and S400R. In another aspect, the first CH3 domain polypeptide comprises amino acid modification D399R, S400R and Y407A and the second CH3 domain polypeptide comprises amino acid modification T366A, K409F, K392E and T411E. In a further embodiment the variant CH3 domain has a melting temperature (T_m) of about 74°C or greater and the heterodimer has a purity of about 95% or greater.

Provided in another embodiment is an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising an amino acid modification at positions L351 and amino acid modification Y407A and a second CH3 domain polypeptide comprises an amino acid modification at position T366 and amino acid modification K409F. In one aspect the amino acid modification at position L351 is selected from L351Y, L351I, L351D, L351R or L351F. In another aspect, the amino acid modification at position Y407 is selected from Y407A, Y407V or Y407S. In yet another aspect the amino acid modification at position T366 is selected from T366A, T366I, T366L, T366M, T366Y, T366S, T366C, T366V or T366W. In one embodiment the variant CH3 domain has a melting temperature (T_m) of about 75°C or greater and the heterodimer has a purity of about 90% or greater.

Provided in another embodiment is an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising an amino acid modification at position F405 and amino acid modifications L351Y and Y407V and a second CH3 domain polypeptide comprises amino acid modification T394W. In one aspect the first CH3 domain polypeptide or the second CH3 domain polypeptide comprise an amino acid modification at positions K392, T411, T366, L368 or S400. The amino acid modification at position F405 is F405A, F405I, F405M, F405T, F405S, F405V or

F405W. The amino acid modification at position K392 is K392V, K392M, K392R, K392L, K392F or K392E. The amino acid modification at position T411 is T411N, T411R, T411Q, T411K, T411D, T411E or T411W. The amino acid modification at position S400 is S400E, S400D, S400R or S400K. The amino acid modification at position T366 is T366A, T366I, T366L, T366M, T366Y, T366S, T366C, T366V or T366W. The amino acid modification at position L368 is L368D, L368R, L368T, L368M, L368V, L368F, L368S and L368A.

In another embodiment is provided an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modifications L351Y, F405A and Y407V and a second CH3 domain polypeptide comprises amino acid modification T394W. In one aspect, the second CH3 domain polypeptide comprises amino acid modification T366L or T366I.

In yet another embodiment is provided an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising at least one of amino acid modifications Y349C, F405A and Y407V and a second CH3 domain polypeptide comprises amino acid modifications T366I, K392M and T394W.

In certain embodiments are provided an isolated heteromultimer comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modifications L351Y, F405A and Y407V and a second CH3 domain polypeptide comprises amino acid modifications K392M and T394W, and one of T366L and T366I.

In another embodiment is provided an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modifications F405A and Y407V and a second CH3 domain polypeptide comprises amino acid modifications T366L and T394W.

In another embodiment is provided an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a first CH3 domain polypeptide comprising amino acid modifications F405A and Y407V and a
5 second CH3 domain polypeptide comprises amino acid modifications T366I and T394W. In certain embodiments of the heteromultimer is provided bi-specific antibody or a multispecific antibody.

In another embodiment is provided a composition comprising a heteromultimer of the
10 invention and a pharmaceutically acceptable carrier.

In another embodiment is provided a host cell comprising nucleic acid encoding the heteromultimer of the invention.

In certain embodiments is provided heteromultimer, wherein target binding by the heteromultimer is competitive to at least one other therapeutic antibody. In one aspect the therapeutic antibody is selected from the group consisting of
15 abagovomab, adalimumab, alemtuzumab, aurograb, bapineuzumab, basiliximab, belimumab, bevacizumab, briakinumab, canakinumab, catumaxomab, certolizumab, pegol, cetuximab, daclizumab, denosumab, efalizumab, galiximab, gemtuzumab
20 ozogamicin, golimumab, ibritumomab tiuxetan, infliximab, ipilimumab, lumiliximab, mepolizumab, motavizumab, muromonab, mycograb, natalizumab, nimotuzumab, ocrelizumab, ofatumumab, omalizumab, palivizumab, panitumumab, pertuzumab, ranibizumab, reslizumab, rituximab, teplizumab, tocilizumab/atlizumab, tositumomab,
25 trastuzumab, Proxinium, Rencarex, ustekinumab, and zalutumumab.

In another embodiment of the heteromultimer of the invention is provided a method of treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering to said patient a therapeutically effective amount of
30 a heteromultimer.

In another embodiment of the heteromultimer of the invention is provided a method of treating immune disorders in a patient having an immune disorder characterized by an immune antigen, said method comprising administering to said patient a
35 therapeutically effective amount of a heteromultimer.

In yet another embodiment is provided an isolated heteromultimer comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability and wherein the variant CH3 domains are selected from the variants listed in Table 1, Table 6 or Table 7.

In certain embodiments are provided heteromultimers that comprise modified anti-idiotypes antibodies belonging to the heavy chain immunoglobulin classes. Such anti-idiotypes can be produced against human or animal idiotypes. A property of these anti-idiotypes is that they can be used as idiotypic vaccines, in particular for vaccination against glycoproteins or glycolipids and where the carbohydrate determines the epitope.

Provided herein are cells or organisms in which heteromultimers described herein have been cloned, said heteromultimers comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region. Such cells or organisms can be used for the purpose of producing heteromultimers having a desired preselected specificity, or corresponding to a particular repertoire. They can also be produced for the purpose of modifying the metabolism of the cell which expresses them. In an embodiment, the heteromultimers comprising at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region is produced in plant cells, especially in transgenics plants.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 is a graphical 3-D structure of a wild type antibody showing the CH3 (top), CH2 (middle) and receptor regions. The dotted line rectangle on the left hand side is expanded to the right hand side showing two regions, Region 1 and Region 2, of the target area of CH3;

Figure 2 is a graphical 3-D representation of showing the wild type residue at position 368;

Figure 3 is a graphical 3-D representation of Region 1 showing mutated position 368;

Figure 4 is a graphical 3-D representation of additional mutations in Region 2;

Figure 5 is a table of in silico calculations for clash score, interface area difference, packing different, electrostatic energy difference and overall "affinity score" for the first three variants AZ1, AZ2 and AZ3;

Figure 6 shows a graphical 3-D image showing variants AZ2 and AZ3, which are "built onto" variant AZ1;

Figure 7 show graphical 3-D representations of AZ2 and AZ3 variants;

Figure 8 shows a table as in Figure 5 but for AZ1, AZ2 and AZ3 heterodimers, and potential homodimers. Affinity score is not shown for homodimers, as it is not relevant;

Figure 9 is a graphical representation of a 3-D representation of wild type (left) and mutated AZ4 (right);

Figure 10 is a table as Figure 5 showing in silico calculations for AZ4 heterodimer and potential homodimers;

Figure 11 is a graphical representation of CH3 variants AZ5 (left) and AZ6 (right);

Figure 12 is a table as described for Figure 5 showing in silico data for AZ4, AZ5 and AZ6;

Figure 13 is a graphical 3-D representation of an antibody on the left, with a drawing of the possibilities of binding characteristics at the receptor region using a heterodimeric approach;

Figure 14 is a schematic representation of the IgG molecule;

Figure 15 shows multiple sequence alignment of Fc γ receptors. Genbank/Uniprot Sequence ID's: Fc γ RIIA (sp P12318), Fc γ RIIB (sp P31994), Fc γ RIIC (gi 126116592), Fc γ RIIIA (sp P08637), Fc γ RIIIB (sp O75015);

5 **Figure 16** is a schematic of the crystal structure of Fc-Fc γ RIIIb Complex [PDB ID: 1T83, Radaev & Sun]. A 1:1 complex of the Fc and Fc γ receptor is observed with an asymmetric contact between the two chains of Fc and the Fc γ R;

10 **Figure 17** shows a schematic of multifunctional molecules based on the asymmetric Fc scaffold formed by heterodimeric variants described herein: Asymmetric Fc Scaffold and Asymmetric Fc- Monomeric IgG Arm;

15 **Figure 18** shows a schematic of multifunctional molecules based on the asymmetric Fc scaffold formed by heterodimeric variants described herein: Asymmetric Fc- Monospecific IgG arms and Asymmetric Fc – Bi-specific IgG Arms (Common Light Chain);

20 **Figure 19** shows an illustration of multifunctional molecules based on the the asymmetric Fc scaffold formed by heterodimeric variants described herein. Asymmetric Fc- Bi-specific IgG Arms and a functional molecule such as toxin;

Figure 20 illustrates multifunctional molecules based on the asymmetric Fc scaffold formed by heterodimeric variants described herein: Asymmetric Fc- Single scFv arm and Asymmetric Fc- bi-specific scFv Arms.

25 **Figure 21** illustrations of alternative multifunctional molecules based on the asymmetric Fc scaffold formed by the heterodimeric variants described herein: Asymmetric Fc- Trispecific scFv Arms and Asymmetric Fc-tetraspecific scFv arms.

30 **Figure 22** displays asymmetric design of mutations on one face of the Fc for better Fc γ R selectivity introduces a productive side for Fc γ R interactions and a non-productive face with wild type like interactions. Mutations on the non-productive face of the Fc can be introduced to block interactions with FcR and bias polarity of the Fc so as to interact on the productive face only.

35 **Figure 23** shows the amino acid sequence for wild-type human IgG1.

Figure 24 Shows the iterative process of the Fc heterodimer design, combining positive and negative design strategies as described in detail below.

Figures 25A-25 C show the in vitro assay used to determine heterodimer purity. The assay is based on a full length monospecific antibody scaffold with two Fc heavy chains of different molecular weight; Heavy chain A has a C-terminal HisTag (His) and heavy chain B a C-terminal, cleavable mRFP Tag (RFP). The two heavy chains A (His) and B (RFP) are expressed in different relative ratios together with a fixed amount of light chain, giving rise to 3 possible dimer species with different molecular weight: a) Homodimer Chain A(His)/ Chain A (His) (~150kDa); b) Heterodimer Chain A (His)/ Chain B (RFP) (~175kDa); c) Homodimer Chain B (RFP)/ Chain B (RFP) (~200kDa). After expression, as described in Example 2, the ratio of heterodimer vs. the two homodimers was determined by non-reducing SDS-PAGE, which allows separation of the 3 dimer species by molecular weight. SDS-PAGE gels were stained with Coomassie Brilliant Blue. **Figure 25A:** Variants tested were WT Chain A (His) only; WT chain B (RFP) only; WT chain A (His) plus chain B (RFP); Control 1 chain A (His) plus chain B (RFP), which has a reported heterodimer purity of >95%. The composition of the dimer bands was verified by Western Blot with antibodies directed against the IgG-Fc (anti-Fc), the mRFP Tag (anti-mRFP) and the HisTag (anti-His), as illustrated above. The SDS-PAGE shows a single band for the His/His homodimer, a double band for the His/RFP heterodimer and multiple bands for the RFP homodimer. The multiple bands are an artifact of the mRFP Tag and have been confirmed not to influence the physical properties of the Fc heterodimer. **Figure 25B:** The SDS-PAGE assay was validated with the published Fc heterodimer variants Controls 1-4 as controls, See, Table A. The variants were expressed with different relative ratios of chain A (His) vs chain B (RFP): Specifically, Ratio 1:3 is equivalent to a LC,HC_His,HC_mRFP ratio of 25%,10%,65%; Ratio 1:1 of 25%,20%,55% and Ratio 3:1 of 25%, 40%,35% respectively (the apparent 1:1 expression of chain A (His) to chain B (RFP) has been determined to be close to 20%/55% (His/RFP) for WT Fc). **Figure 25C** shows a non-reducing SDS-PAGE assay to determine heterodimer purity of Scaffold 1 variants. The heteromultimers were expressed with different relative ratios of chain A (His) vs chain B (RFP) and analyzed by non-reducing SDS-PAGE as described in Figure 2. Specifically, Ratio 1:3 is equivalent to a LC,HC_His,HC_mRFP ratio of 25%,10%,65%; Ratio 1:1 of 25%,20%,55% and Ratio 3:1 of 25%, 40%,35% respectively (the apparent 1:1 expression of chain A (His) to chain B (RFP) has been determined to be close to 20%/55% (His/RFP) for WT Fc).

Figures 26A-26B show Fc Heterodimer variants expressed with a specific ratio of chain A (His) vs chain B (RFP) (See, Table 2), purified by Protein A affinity chromatography and analyzed by non-reducing SDS-PAGE as described in Figures 25A-25C. **Figure 26A** illustrates classification of heterodimers based on purity as observed by visual inspection of the SDS-PAGE results. For comparison the equivalent amount of Protein A purified product was loaded on the gel. This definition of purity based on non-reducing SDS-PAGE has been confirmed by LC/MS on selected variants (see Figure 28). **Figure 26B** provides exemplary SDS-PAGE results of selected Protein A purified heterodimer variants (AZ94, AZ86, AZ70, AZ33 and AZ34).

Figures 27A-27B illustrate DSC analyses to determine the melting temperature of the heterodimeric CH3-CH3 domain formed by the Heterodimer variants described herein. Two independent methods were used to determine the melting temperatures. **Figure 27A** provides thermograms fitted to 4 independent non-2-state-transitions and optimized to yield values for the CH2 and Fab transitions close to the reported literature values for Herceptin of ~72°C (CH2) and ~82°C (Fab). **Figure 27B** shows the normalized and baseline corrected thermograms for the heterodimer variants were subtracted from the WT to yield a positive and negative difference peak for only the CH3 transition.

Figure 28 illustrates the LC/MS analysis of example variant AZ70 as described in the example 2. The expected (calculated average) masses for the glycosylated heterodimer and homodimers are indicated. The region consistent with the heterodimer mass contains major peaks corresponding to the loss of a glycine (-57 Da) and the addition of 1 or 2 hexoses (+162 Da and +324 Da, respectively). The Heterodimer purity is classified as >90% if there are no significant peaks corresponding to either of the homodimers.

Figures 29A-29D shows the CH3 interface of Fig29AWT Fc; Fig29BAZ6; Fig29CAZ33; Fig29DAZ19. The comprehensive in silico analysis, as described in the detailed description section, and the comparison of the variants to the WT indicated that one of the reasons for the lower than WT stability of the initial AZ33 heterodimer is the loss of the core interaction/packing of Y407 and T366. The initial AZ33 shows non-optimal packing at this hydrophobic core as illustrated Fig29B, suggesting that optimization of this region, particularly at position T366 would improve the stability of AZ33. This is illustrated in Fig 29C and Fig29D with T366I and T366L. The

experimental data correlates with this structural analysis and shows that T366L gives the greatest improvement in T_m. See, Example 5.

Figure 30 illustrates the utility and importance of the conformational dynamics analysis, exemplified at the initial Scaffold 1 variant AZ8. The structure after in silico mutagenesis (backbone conformation close to WT) is superimposed with a representative structure of a 50ns Molecular Dynamics simulation analysis. The figure highlights the large conformational difference in the loop region D399-S400 of AZ8 variant vs. WT, which in turn exposes the hydrophobic core to solvent and causes decreased stability of the AZ8 heterodimer.

Figures 31A-31C illustrate how the information from the comprehensive in silico analysis and the MD simulation was used in the described positive design strategy. As illustrated in Figure 30, one of the reasons for the lower than WT stability of AZ8 is the weakened interaction of the loop 399-400 to 409, which is mainly due to the loss of the F405 packing interactions (see comparison of **Fig 31A** (WT) vs **Fig 31B** (AZ8)). One of the positive design strategies was optimization of the hydrophobic packing of area, to stabilize the 399-400 loop conformation. This was achieved by the K392M mutation that is illustrated in Fig 31C. **Fig 31C** represents the heterodimer AZ33, which has a T_m of 74° vs. 68° of the initial negative design variant AZ8.

Figures 32A-32B illustrate the dynamics of the Fc molecule observed using principal component analysis of a molecular dynamics trajectory. **Fig 32A** shows a backbone trace of the Fc structure as reference. **Fig 32B** and C represent an overlay of dynamics observed along the top 2 principal modes of motion in the Fc structure. The CH2 domains of chain A and B exhibits significant opening /closing motion relative to each other while the CH3 domains are relatively rigid. Mutations at the CH3 interface impact the relative flexibility and dynamics of this open/close motion in the CH2 domains.

Figures 33A-33C illustrate the hydrophobic core packing of two Scaffold-2 variants vs. WT. **Fig 33A** WT Fc; **Fig 33B** AZ63; and **Fig 33C** AZ70. The comprehensive in-silico analysis of the initial Scaffold-2 variant suggested that loss of the core WT interactions of Y407-T366 is one of the reasons for the lower than WT stability for the initial Scaffold-2 variants. The loss of Y407-T366 is partially compensated by the mutations K409F, but as illustrated in Fig 33B, particularly the T366A mutation leaves a cavity in the hydrophobic core, which destabilizes the variant vs. WT. Targeting this

hydrophobic core by additional mutations T366V_L351Y, as shown by AZ70 in **Fig33C**, proved to be successful; AZ70 has an experimentally determined T_m of 75.5°C. See, Table 4 and Example 6.

5 **Figures 34A-34C** illustrate the interactions of the loop 399-400 of two Scaffold-2 variants vs. the WT: **Fig 34A** WT Fc; **Fig 34B** AZ63; and **Fig 34C** AZ94. The comprehensive in-silico analysis of the initial Scaffold-2 variant suggested that loss of the WT salt-bridge K409-D399 (**Fig 34A**) due to the mutation K409F and the hence unsatisfied D399 (**Fig34B**) causes a more 'open' conformation of the 399-400 loop.

10 This leads furthermore to a greater solvent exposure of the hydrophobic core and a further destabilization of the variant vs WT. One of the strategies employed to stabilize the 399-400 loop and compensate for the loss of the K409-D399 interaction was the design of additional salt bridges D399R-T411E and S400R-K392E as illustrated in **Fig34C** for variant AZ94. Experimental data showed a purity of >95%

15 and T_m of 74°C. See, Table 4 and Example 6. Further, although AZ94 has a considerably higher purity and stability compared to the initial Scaffold-2 variant (purity <90%, T_m 71°C), the hydrophobic core mutations of AZ94 are less preferred than the 'best' hydrophobic core mutations identified in variant AZ70 (Figure 33). Since the mutations at the hydrophobic core in AZ70 (T366V_L351Y) are distal from

20 the salt-bridge mutations of AZ94 at the loop 399-400, the combination of AZ70 amino acid mutations and the additional AZ94 mutations, is expected to have a higher melting temperature than AZ70 or AZ94. This combination can be tested as described in Examples 1-4.

25 **Figure 35** illustrates the association constant ($K_a(M^{-1})$) of homodimeric IgG1 Fc, the heterodimeric variants het1(Control 1):

A:Y349C_T366S_L368A_Y407V/B:S354C_T366W and het2(Control 4):

A:K409D_K392D/B:D399K_D356K binding to the six Fcγ receptors. The heterodimers tend to show slightly altered binding to the Fcγ receptors

30 compared to the wild type IgG1 Fc. See, Example 7

Figure 36A Shows the relative binding strength of a wild type IgG1 Fc and its various homodimeric and asymmetric mutant forms to the IIbF, IIbY and IIaR receptors, based on the wild type binding strength as reference. (Homo Fc + S267D) refers to the binding strength of a homodimeric Fc with the S267D mutation on both chains.

35 (Het Fc + asym S267D) refers to the binding strength of a heterodimeric Fc with the S267D mutation introduced in one of the two chains in Fc. The average of the

binding strength obtained by introducing the mutation on either of the two Fc chains is reported. Introduction of this mutation on one chain reduced the binding strength to roughly half the strength observed for the same mutation in a homodimeric manner.

The (Het Fc + asym S267D + asym E269K) refers to the binding strength of a heterodimeric Fc with both the S267D and E269K mutations introduced in an asymmetric manner on one of the two Fc chains. The E269K mutation blocks the interaction of the FcγR to one of the faces of the Fc and is able to bring down the binding strength by roughly half of what was observed for the asymmetric S267D variant (Het Fc+S267D) by itself. The Het Fc here is comprised of CH3 mutations as indicated for the variant het2 (Control 4) in Figure 35.

Figure 36B Shows the association constant ($K_a(M^{-1})$) of various Fc's and its variants with a number of FcγRIIa, FcγRIIb and FcγRIIIa allotypes. The K_a of wild type IgG1 Fc to various Fcγ receptors is represented as columns with horizontal shade. The bars with vertical shades (homodimer base2) represent the K_a of homodimeric Fc with the mutations S239D/D265S/I332E/S298A. The columns with the slanted shade represent the K_a of heterodimeric Fc with asymmetric mutations A:S239D/D265S/I332E/E269K and B:S239D/D265S/S298A in the CH2 domain. The introduction of asymmetric mutations is able to achieve increased selectivity between the IIIa and IIa/IIb receptors. The Heterodimeric Fc here is comprised of CH3 mutations as indicated for the variant het2 (Control 4) in Figure 35.

Figure 36C Shows the association constant ($K_a(M^{-1})$) for wild type IgG1 and three other constructs involving homodimeric or asymmetric mutations in the CH2 domain of the Fc region. The K_a of wild type Fc is represented in the column shaded with grids. The K_a of Fc variant with the base mutation S239D/K326E/A330L/I332E/S298A introduced in a homodimeric manner (homodimer base1) on both the chains of Fc is shown with the slanted patterned column. Introduction of related mutations in an asymmetric manner in chains A and B of a heterodimeric Fc (hetero base1) is shown with the horizontal lines. The column with vertical shaded lines represents the asymmetric variant including the E269K mutation (hetero base 1+PD). The Heterodimeric Fc here is comprised of CH3 mutations as indicated for the variant het2 (Control 4) in Figure 35.

Figure 37 - Table 6 is a list of variants CH3 domains based on the third design phase as described in Example 5 for Scaffold 1.

Figure 38 - Table 7 is a list of variant CH3 domains based on the third design phase as described in Example 6 for scaffold 2.

Figure 39A-39B illustrate Purity determination of variants without any C-terminal

- 5 Tags using LC/MS. **Fig 39A** shows the LC/MS spectra of one representative variant (AZ162: L351Y_F405A_Y407V / T366L_K392L_T394W). The variant was expressed by transient co-expression as described in the Examples using 3 different HeavyChain-A to HeavyChain-B ratios of 1:1.5 (AZ133-1), 1:1 (AZ133-2) and 1.5:1 (AZ133-3). The samples were purified and deglycosylated with Endo S for 1 hr at
- 10 37°C. Prior to MS analysis the samples were injected onto a Poros R2 column and eluted in a gradient with 20-90% ACN, 0.2% FA in 3 minutes. The peak of the LC column was analyzed with a LTQ-Orbitrap XL mass spectrometer (Cone Voltage: 50 V' Tube lens: 215 V; FT Resolution: 7,500) and integrated with the software Promass to generate molecular weight profiles. **Fig 39B** shows the LC/MS spectra of the
- 15 Control 2 sample, which represents the Knobs-into-Holes variant. The variant was expressed by transient co-expression as described in the Examples using 3 different HeavyChain-A to HeavyChain-B ratios of 1:1.5 (Control 2-1), 1:1 (Control 2-2) and 1.5:1 (Control 2-3). The samples were purified and deglycosylated with Endo S for 1 hr at 37°C. Prior to MS analysis the samples were injected onto a Poros R2 column
- 20 and eluted in a gradient with 20-90% ACN, 0.2% FA in 3 minutes. The peak of the LC column was analyzed with a LTQ-Orbitrap XL mass spectrometer (Cone Voltage: 50 V' Tube lens: 215 V; FT Resolution: 7,500) and integrated with the software Promass to generate molecular weight profiles.

- 25 **Figure 40** provides multiple sequence alignment of CH3 domain sequences. The CH3 sequences of the different species are numbered according to the human IgG1 reference and the Eu numbering scheme. The residues mutated to achieve heterodimer formation are indicated by * for Chain_A and + for Chain_B. The mutations for Scaffold 1 include: Chain_A: (350V)_351Y_405A_407V and Chain_B: (350V)_366L_392L_394W. The corresponding positions for each of the species, as
- 30 illustrated by gray and black boxes in the sequence alignment, can be mutated to form heterodimers in an IgG1 like manner with high purity. Species legend:
- IgG1_human (Homo sapiens); IgG2a_camel Camelus dromedarius (Arabian camel); IgG3_camel Camelus dromedarius (Arabian camel); IgG2a_llama Lama glama (Llama); IgG3_llama Lama glama (Llama); IgG2a_mouse Mus musculus (house mouse); IgG1_mouse Mus musculus (house mouse); IgG3_mouse Mus musculus (house mouse); IgG_rabbit Oryctolagus cuniculus (rabbit); IgG1_sheep Ovis aries
- 35

(sheep); IgG2a_rat *Rattus norvegicus* (Norway rat); IgG1_rat *Rattus norvegicus* (Norway rat); IgY_chicken *Gallus gallus* (Chicken).

Figure 41 depicts SDS-PAGE analysis of an exemplary heteromultimer

5

Figure 42 depicts UPLC analysis of an exemplary heteromultimer.

Figure 43 depicts the ability of an exemplary heteromultimer to bind to EGFR.

10 **Figure 44** depicts the amino acid sequence of an sdab that binds EGFR

DETAILED DESCRIPTION

Provided herein are isolated heteromultimers that comprise at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer FC region that comprises modified CH3 domain comprising specific amino acid modifications to promote heteromultimer formation, wherein said isolated heteromultimers are devoid of immunoglobulin light chains, and optionally devoid of immunoglobulin first constant (CH1) region and immunoglobulin second constant (CH2) region. In some embodiments, the modified CH3 domains comprise specific amino acid modifications to promote heterodimer formation (See, for example Tables 1.1-1.3). In another embodiment the modified CH3 domains comprise specific amino acid modifications to promote heterodimer formation with increased stability (See, for example Table 4, Table 6 and Table 7). Stability is measured as the melting temperature (T_m) of the CH3 domain and an increased stability refers to a T_m of about 70°C or greater. The CH3 domains form part of the Fc region of a heteromultimeric, multispecific antibody. Provided herein in one embodiment are heteromultimers comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a modified or variant CH3 domain comprising amino acid mutations to promote heterodimer formation wherein the variant CH3 domains are selected from the variants listed in Table 1. In a second embodiment, provided are heteromultimers comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of about 70°C or greater.

35

Amino acid modifications utilized to generate a modified CH3 domain include, but are not limited to, amino acid insertions, deletions, substitutions, and rearrangements.

The modifications of the CH3 domain and the modified CH3 domains are referred to herein collectively as "CH3 modifications", "modified CH3 domains", "variant CH3 domains" or "CH3 variants". In certain embodiments, the modified CH3 domains are incorporated into a molecule of choice. Accordingly, in one embodiment are provided molecules, for instance polypeptides, such as immunoglobulins (e.g., antibodies) and other binding proteins, comprising an Fc region (as used herein "Fc region" and similar terms encompass any heavy chain constant region domain comprising at least a portion of the CH3 domain) incorporating a modified CH3 domain. Molecules comprising Fc regions comprising a modified CH3 domain (e.g., a CH3 domain comprising one or more amino acid insertions, deletions, substitutions, or rearrangements) are referred to herein as "Fc variants", "heterodimers" or "heteromultimers". The present Fc variants comprise a CH3 domain that has been asymmetrically modified to generate heterodimer Fc variants or regions. The heteromultimer is comprised of two heavy chain polypeptides - Chain A and Chain B, which can be used interchangeably provided that each Fc region comprises one Chain A and one Chain B polypeptide, and provided that at least one of Chain A and Chain B comprises a heavy chain variable region. The amino acid modifications are introduced into the CH3 in an asymmetric fashion resulting in a heterodimer when two modified CH3 domains form an Fc variant (See, e.g., Table 1). As used herein, asymmetric amino acid modifications are any modification wherein an amino acid at a specific position on one polypeptide (e.g., "Chain A") is different from the amino acid on the second polypeptide (e.g., "Chain B") at the same position of the heterodimer or Fc variant. This can be a result of modification of only one of the two amino acids or modification of both amino acids to two different amino acids from Chain A and Chain B. It is understood that the variant CH3 domains comprise one or more asymmetric amino acid modifications.

Definitions

In the present description, any concentration range, percentage range, ratio range, or integer range is to be understood to include the value of any integer within the recited range and, when appropriate, fractions thereof (such as one tenth and one hundredth of an integer), unless otherwise indicated. As used herein, "about" means $\pm 10\%$ of the indicated range, value, sequence, or structure, unless otherwise indicated. It should be understood that the terms "a" and "an" as used herein refer to "one or

more" of the enumerated components unless otherwise indicated or dictated by its context. The use of the alternative (e.g., "or") should be understood to mean either one, both, or any combination thereof of the alternatives. As used herein, the terms "include" and "comprise" are used synonymously. In addition, it should be

5 understood that the individual single chain polypeptides or heterodimers derived from various combinations of the structures and substituents (e.g., variant CH3 domains) described herein are disclosed by the present application to the same extent as if each single chain polypeptide or heterodimer were set forth individually. Thus, selection of particular components to form individual single chain polypeptides or
10 heterodimers is within the scope of the present disclosure.

The "first polypeptide" is any polypeptide that is to be associated with a second polypeptide, also referred to herein as "Chain A". The first and second polypeptide meet at an "interface". The "second polypeptide" is any polypeptide that is to be
15 associated with the first polypeptide via an "interface", also referred to herein as "Chain B". At least one of said first and second polypeptides comprise at least one heavy chain variable domain. In certain embodiments, the at least one heavy chain variable domain is obtained from a heavy chain antibody. In some embodiments, the heavy chain antibody is obtained from a cartilaginous fish such as a shark or a
20 camelid such as a llama. The "interface" comprises those "contact" amino acid residues in the first polypeptide that interact with one or more "contact" amino acid residues in the interface of the second polypeptide. As used herein, the interface comprises the CH3 domain of an Fc region. In some embodiments, the Fc region is derived from an IgG antibody such as, but not restricted to a human IgG₁ antibody. In
25 certain embodiments, the at least one heavy chain variable domain is connected to the CH3 domain by means of a linker.

As used herein, "isolated" heteromultimer means a heteromultimer that has been identified and separated and/or recovered from a component of its natural cell culture
30 environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the heteromultimer, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes.

As used herein, "stability comparable to a native Fc homodimer" means that the Fc
35 heterodimer has a stability analogous to that of the native homodimer. In certain embodiments, the stability of the heterodimer is within $\pm 5^{\circ}\text{C}$ of the corresponding

native homodimeric Fc. In some embodiments, the stability of the heterodimer is within $\pm 2^{\circ}\text{C}$ of the corresponding native homodimeric Fc.

The heteromultimers described herein are generally purified to substantial
5 homogeneity. The phrases “substantially homogeneous”, “substantially homogeneous form” and “substantial homogeneity” are used to indicate that the product is substantially devoid of by-products originated from undesired polypeptide combinations (e.g. homodimers). Expressed in terms of purity, substantial
10 homogeneity means that the amount of by-products does not exceed 10%, and preferably is below 5%, more preferably below 1%, most preferably below 0.5%, wherein the percentages are by weight.

Terms understood by those in the art of antibody technology are each given the meaning acquired in the art, unless expressly defined differently herein. Antibodies
15 are known to have variable regions, a hinge region, and constant domains.

Immunoglobulin structure and function are reviewed, for example, in Harlow et al, Eds., *Antibodies: A Laboratory Manual*, Chapter 14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, 1988).

20 The “Fab fragment” of an antibody (also referred to as fragment antigen binding) contains the constant domain (CL) of the light chain and the first constant domain (CH1) of the heavy chain along with the variable domains VL and VH on the light and heavy chains respectively. The variable domains comprise the complementarity determining loops (CDR, also referred to as hypervariable region) that are involved in
25 antigen binding. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region.

“Single-chain Fv” or “scFv” antibody fragments comprise the VH and VL domains of an antibody, wherein these domains are present in a single polypeptide chain. In
30 one embodiment, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen binding. For a review of scFv see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994). HER2 antibody scFv fragments are described in
35 WO93/16185; U.S. Pat. No. 5,571,894; and U.S. Pat. No. 5,587,458.

“Humanized” forms of non-human (e.g., rodent) antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., *Nature* 321:522-525 (1986); Riechmann et al., *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

Heteromultimers

The present invention provides heteromultimers comprising at least one single domain antigen-binding (sdAg) construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide attached to an immunoglobulin heterodimer Fc region that comprises amino acid modifications that promote the formation of a heterodimeric Fc region with stability comparable to that of a native immunoglobulin homodimeric Fc region, and are devoid of IgG light chains and IgG CH1 regions. The heteromultimers can be monovalent and monospecific, bivalent and monospecific, or bivalent and bi-specific. The heteromultimers described herein, comprising the heterodimer Fc region described herein, have an intrinsic stability comparable to wild-type IgG1, and are formed with a purity of greater than about 85%.

In one embodiment the heteromultimer is a heterodimer comprising two heavy chains.

The heteromultimers described herein allow for easier construction and manufacturability of multi-functional, bi-specific antibodies compared to scFv or Fab comprising antibody formats. Since the heteromultimers described here are devoid of IgG light chains, the "light chain scrambling" problem inherent to making a bi-specific antibody comprising two light chains is avoided.

Heteromultimers as described herein also demonstrate

- a) high affinity binding of antigen by the single domain antigen binding construct;
- b) good heteromultimer quality and biophysical stability (lack of aggregation) of the constituent single domain antigen-binding construct, and
- 10 c) high antibody titre in Chinese hamster ovary (CHO) cell expression systems and manufacturability compared to other heteromultimer constructs comprising scFv and Fab formats.

These properties allow the heteromultimers described here to be used in a variety of applications including the development of bi-specific and multifunctional therapeutic antibodies and diagnostic or targeting reagents.

In one embodiment, the isolated heteromultimer comprises at least one single domain antigen-binding construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations that promote the formation of said heterodimer Fc region with stability comparable to a native homodimeric Fc region; and wherein said isolated heteromultimer is devoid of immunoglobulin light chain and immunoglobulin first constant (CH1) region.

It is contemplated that the heteromultimer described herein may have only one single domain antigen-binding construct attached to the heterodimer Fc region. Thus, in one embodiment, the isolated heteromultimer comprises one single domain antigen-binding construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide.

It is further contemplated that the isolated heteromultimer described herein may comprise two single domain antigen-binding constructs attached to the heteromultimer Fc region. In another embodiment, the isolated heteromultimer

comprises one single domain antigen-binding construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the one single domain antigen-binding construct is attached to one monomeric Fc polypeptide and a second single domain antigen-binding construct is attached to the second monomeric Fc polypeptide.

Single domain antigen binding constructs

As indicated above, the heteromultimers described herein comprise at least one single domain antigen-binding construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide.

Single domain antigen-binding constructs include binding polypeptides that can specifically or selectively bind target polypeptides. The term "specific binding" as used herein, refers to high-affinity binding of the antigen binding construct to the antigen as observed in the equilibrium dissociation constant K_d . K_d is the equilibrium dissociation constant and equal to k_{off}/k_{on} . k_{off} describes the dissociation rate of antigen binding construct complexed to the antigen and k_{on} describes the association rate of antigen binding construct to the antigen. The term "selective binding" as used herein, refers to the differing affinities with which a ligand binds to receptors or targets, such that a ligand shows higher affinity for one target over another; Selective ligands bind to a very limited types of receptors, whereas non-selective ligands bind to several types of receptors. As is known in the art, specific and selective binding to target receptor can empower a therapeutic comprising the single domain antigen binding construct to recognize and treat diseased cells that express the target.

Examples of suitable single domain antigen-binding constructs include those that are devoid of antibody light chains such as single domain antibodies (sdAb or V_H), camelid nanobodies (V_HH), shark V_{NAR} , SH3-derived fynomers, and fibronectin-derived binding domains such as adnectins and DARPins (designed ankyrin repeat proteins). These single domain antigen-binding constructs have been shown to exhibit properties such as the ability to bind to alternative or cryptic epitopes that may not be accessible by traditional Fabs due to their size and structural conformation.

In one embodiment, the single domain antigen-binding construct is an immunoglobulin heavy chain variable region or variable heavy chain selected from a

domain antibody (or an amino acid sequence that is suitable for use as a domain antibody), a single domain antibody (or an amino acid sequence that is suitable for use as a single domain antibody), a "dAb" (or an amino acid sequence that is suitable for use as a dAb) or a Nanobody (as defined herein, and including but not limited to a V_{HH} sequence); other single variable domains, or any suitable fragment of any one thereof. For a general description of (single) domain antibodies, reference is also made to the art cited above, as well as to EP 0 368 684. For the term "dAb's", reference is for example made to Ward et al. (Nature 1989 Oct. 12; 341 (6242): 544-6), to Holt et al. (Trends Biotechnol., 2003, 21(11):484-490); as well as to for example WO 04/068820, WO 06/030220, WO 06/003388 and other published patent applications of Domantis Ltd. In some embodiments, the variable heavy chains comprise single domain antibodies or single variable domains can be derived from certain species of shark (for example, the so-called "IgNAR domains", see for example WO 05/18629).

Thus, in one embodiment, the isolated heteromultimer comprises a single domain antigen-binding construct selected from single domain antibodies, camelid nanobodies, shark V_{NAR}, SH3-derived fynomers, and fibronectin-derived binding domains such as adnectins and DARPins. In one embodiment, the isolated heteromultimer comprises a single domain antigen-binding construct selected from single domain antibodies, camelid nanobodies and cartilaginous fish V_{NAR}. In one embodiment, the isolated heteromultimer comprises a single domain antigen-binding construct that is an SH3-derived fynomer or a fibronectin-derived binding domain. In one embodiment, the fibronectin-derived binding domain is a DARPin or adnectin. SH3-derived fynomers, adnectins and DARPins are known in the art (see for example Grabulovski et al. J Biol Chem. (2007) 282(5):3196-204; Lipovšek Protein Engineering, Design and Selection (2011) 24(1-2): 3-9, and Tamaskovic, Methods in Enzymology (2012), 503, 101-134).

In one embodiment, the single domain antigen-binding construct is a single domain antibody. Single domain antibodies are antibody fragments that consist of a single monomeric variable antibody domain. Suitable examples of domain antibodies are known in the art and include, for example, (see for example Figure 4 of Wesolowski et al. *Med Microbiol Immunol.* (2009), 198(3): 157-174 and Figure 1A of Barthelemy et al. *J. Biol. Chem.* (2008) 138, 3639-54).

In one embodiment, the single domain antigen-binding construct is a camelid nanobody (V_hH). Camelid nanobodies are antibody fragments derived from heavy

chain antibodies found in camelids that are devoid of light chain and heavy chain CH1 constant domain. In one embodiment, the single domain antigen-binding construct is a cartilaginous fish V_{NAR} . Such cartilaginous fish V_{NAR} include V_{NARS} antibody fragments derived from heavy chain antibodies found in sharks. These

5 antibody fragments are also devoid of light chains and heavy chain CH1 constant domain. Examples of known camelid and shark single domain antigen-binding constructs are identified in Table 1 of Wesolowski *et al*, *Med Microbiol Immunol* (2009) 198:157-174. For example, single domain antigen-binding constructs that target membrane proteins include ART2.2 from Immune llama; CD16 from Immune

10 llama; EGFR from Immune camel, immune llama; CEA Cancer immunotherapy from Immune llama; MUC 1 Tumor targeting Immune from camel and llama, and CD105 (endoglin) from Immune camel. Single domain antigen-binding constructs that target secretory proteins include TNF from Immune llama and alpaca; PSA from Immune dromedary; von Willebrand factor from Immune llama; Amyloid A peptide from

15 Immune dromedary and alpaca; Lysozyme from Immune dromedary; IgG from Immune llama; and Serum albumin from Immune llama. Single domain antigen-binding constructs that target intracellular proteins include Bax from Non-immune llama; HIF-1 from Non-immune llama, PABPN1 from Immune and non-immune llama.

20 Single domain antigen-binding constructs suitable for use in the heteromultimers described herein can be obtained from naturally occurring sources such as camelids (including camels, llamas, and alpacas, for example), and sharks. Single domain antigen-binding constructs may also be obtained by screening libraries such as phage-display libraries in order to select for single domain antigen-binding constructs

25 that bind to a target of interest. Methods of screening such libraries in order to select target-specific single domain antigen-binding constructs are known in the art (see for example Groot *et al*, in *Lab Invest* (2006) 86:345-56, and Verheesen *et al* in *Methods Mol Biol* (2012) 911:81-104).

The nucleotide and/or amino acid sequences of specific single domain antigen-binding constructs are known in the art or are accessible in published sequence databases, for example, GenBank, SwissProt, or EMBL for example, thus facilitating the preparation of the heteromultimers comprising single domain antigen-binding constructs as described herein.

30

Selection of targets

As indicated above, the single domain antigen-binding constructs are able to selectively and/or specifically bind to a target antigen. The target antigen is selected based on the intended use of the heteromultimer. In one embodiment, the target cell is a cell that is activated or amplified in a cancer, an infectious disease, an autoimmune disease, or in an inflammatory disease. In one embodiment, where the heteromultimer binds to EGFR1, the target cell is a cell that is activated or amplified in a cancer, an autoimmune disease, or in an inflammatory disease.

In another embodiment, the target cell is one that is activated or amplified when a subject is suffering from an infection with a pathogenic organism, such as bacteria or fungi.

In another embodiment, the heteromultimers are used to target a cell expressing a target antigen that is not typically accessible by traditional antigen-binding moieties such as Fabs. The nature of the single domain antigen-binding construct of the heteromultimer allows for binding to targets such as, for example, highly conserved residues such as CD4, for example, that are protected from the humoral immune system by conformational masking and steric occlusion, and to target antigen smaller epitopes.

Specific targets

In one embodiment, the heteromultimers according to the invention target one or more cytokines or chemokines such as, for example, IL2, IFN α -2a/b, IFN-1a/b, IL-21, IL-17a, TNF, IL23, VEGF, or ANG2. In another embodiment, the heteromultimers according to the invention target one or more tumor associated antigens such as EpCam, EGFR, VEGFR, CEA, or GP100. In another embodiment, the heteromultimers according to the invention target immunoregulatory antigens such as CD16, CD30, CD137, CD22, CD52, CD80, CD23, CD2, CD4, CD40, KIR, CD32b, CD25, LAG3, or B7-H3.

In one embodiment, the heteromultimers provided herein target one or more bacterial toxins such as Clostridium difficile toxin A, Clostridium difficile toxin B.

In certain embodiments, the heteromultimers provided herein are useful to target one or more target antigen selected from EGFR, IGF1R, ICAM-1, Clostridium difficile toxin A, Clostridium difficile toxin B, ICAM-1, Bax-protein, CDC50A, and CD3 isoforms inclusive of the epsilon isoform. In certain embodiments are heteromultimers described herein, comprising at least one single domain antigen-

binding construct that targets one or more of EGFR, IGF1R, ICAM-1, Clostridium difficile toxin A, Clostridium difficile toxin B, ICAM-1, Bax-protein, CDC50A, and CD3 isoforms inclusive of the epsilon isoform. In certain embodiments are heteromultimers described herein, comprising at least one single domain antigen-binding construct derived from a llama heavy chain antibody, wherein said variable heavy chain targets one or more of EGFR, IGF1R, ICAM-1, Clostridium difficile toxin A, Clostridium difficile toxin B, ICAM-1, Bax-protein, CDC50A, and CD3 isoforms inclusive of the epsilon isoform. In certain embodiments are multispecific heteromultimers comprising single domain antigen-binding constructs that target one or more of EGFR, IGF1R, ICAM-1, Clostridium difficile toxin A, Clostridium difficile toxin B, ICAM-1, Bax-protein, CDC50A, and CD3 isoforms inclusive of the epsilon isoform. In certain embodiments are bi-specific heteromultimers described herein, wherein said heteromultimers comprise single domain antigen-binding constructs derived from camelid heavy chain antibodies that target one or more of EGFR, IGF1R, ICAM-1, Clostridium difficile toxin A, Clostridium difficile toxin B, ICAM-1, Bax-protein, CDC50A, and CD3 isoforms inclusive of the epsilon isoform. In certain embodiments are bi-specific heteromultimers described herein, wherein said heteromultimers comprise single domain antigen-binding constructs derived from llama heavy chain antibodies that target one or more of EGFR, IGF1R, ICAM-1, Clostridium difficile toxin A, Clostridium difficile toxin B, ICAM-1, Bax-protein, CDC50A, and CD3 isoforms inclusive of the epsilon isoform.

In one embodiment, the single domain antigen-binding construct is one that has a neutralizing activity on the target antigen. The term "neutralizing activity," as used herein in the context of a single domain antigen-binding construct, refers the ability of the single domain antigen-binding construct to block binding of a cognate ligand to the target antigen. In another embodiment, the single domain antigen-binding construct is one that does not have neutralizing activity on the target antigen. In one embodiment, the heteromultimer comprises an EGFR single domain antigen-binding construct that is non-neutralizing. In another embodiment, the heteromultimer comprises an EGFR single domain antigen-binding construct that is neutralizing. Examples of neutralizing EGFR single domain antigen-binding constructs are found in Omidfar *et al.* (2012) 31:1015-1026.

In one embodiment where the heteromultimer according to the invention comprises two single domain antigen-binding constructs, both single domain antigen-binding constructs bind to the same antigen. In another embodiment, where the

heteromultimer according to the invention comprises two single domain antigen-binding constructs, both single domain antigen-binding constructs bind to the same epitope. In another embodiment, where the heteromultimer according to the invention comprises two single domain antigen-binding constructs, one single domain antigen-binding construct binds to one target and the second single domain antigen-binding construct binds to a different target. In still another embodiment, where the heteromultimer according to the invention comprises two single domain antigen-binding constructs, one single domain antigen-binding construct binds to one epitope and the second single domain antigen-binding construct binds to a different epitope.

10 **Immunoglobulin heterodimer Fc region**

As indicated above, the heteromultimers described herein comprise at least one single domain antigen-binding (sdAg) construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide attached to an immunoglobulin heterodimer Fc region that comprises amino acid modifications that promote the formation of a heterodimeric Fc region with stability comparable to that of a native immunoglobulin homodimeric Fc region, and are devoid of IgG light chains and IgG CH1 regions.

Immunoglobulin heterodimer Fc regions are further described as follows. As indicated elsewhere herein, in one embodiment are provided molecules, for instance polypeptides, such as immunoglobulins (e.g., antibodies) and other binding proteins, comprising an Fc region (as used herein "Fc region" and similar terms encompass any heavy chain constant region domain comprising at least a portion of the CH3 domain) incorporating a modified CH3 domain. Molecules comprising Fc regions comprising a modified CH3 domain (e.g., a CH3 domain comprising one or more amino acid insertions, deletions, substitutions, or rearrangements) are referred to herein as "Fc variants", "heterodimers," "variant Fc heterodimers" or "heteromultimers". The present Fc variants comprise a CH3 domain that has been asymmetrically modified to generate heterodimer Fc variants or regions. The heteromultimer is comprised of two heavy chain polypeptides, or two monomeric Fc polypeptides - Chain A and Chain B, which can be used interchangeably provided that each Fc region comprises one Chain A and one Chain B polypeptide, and provided that at least one of Chain A and Chain B comprises a heavy chain variable region.

The design of variant Fc heterodimers from wildtype homodimers is illustrated by the concept of positive and negative design in the context of protein engineering by balancing stability vs. specificity, wherein mutations are introduced with the goal of driving heterodimer formation over homodimer formation when the polypeptides are expressed in cell culture conditions. Negative design strategies maximize unfavorable interactions for the formation of homodimers, by either introducing bulky sidechains on one chain and small sidechains on the opposite, for example the knobs-into-holes strategy developed by Genentech (Ridgway JB, Presta LG, Carter P. 'Knobs-into-holes' engineering of antibody CH3 domains for heavy chain heterodimerization. *Protein Eng.* 1996 Jul;9(7):617-21; Atwell S, Ridgway JB, Wells JA, Carter P. Stable heterodimers from remodeling the domain interface of a homodimer using a phage display library. *J Mol Biol.* 270(1):26-35 (1997))), or by electrostatic engineering that leads to repulsion of homodimer formation, for example the electrostatic steering strategy developed by Amgen (Gunashekar K, et al. Enhancing antibody Fc heterodimer formation through electrostatic steering effects: applications to bi-specific molecules and monovalent IgG. *JBC* 285 (25): 19637-19646 (2010)). In these two examples, negative design asymmetric point mutations were introduced into the wild-type CH3 domain to drive heterodimer formation. To date, only negative design strategies have been used to develop Fc heterodimers. Published results show that heterodimers designed using only a negative design approach leads to high specificity with >95% heterodimers, but destabilizes the complex considerably (*Supra*). These negative design heterodimers possess a melting temperature, of the modified CH3 domain, of 69°C or less, absent additional disulfide bonds as compared to the wild type. See, Table A below.

25

Table A: Published Fc Heterodimer Antibodies.

	Chains	Engineering Approach	Source	Purity	T _m °C
Wild-Type	-				81-83
	-				
Control 4	Y349C_T366S_L368A_Y407V	Knobs-into-holes plus disulfide	Genentech (Merchant et al.)	95%	>77**
	S354C_T366W				
Control 3	K409D_K392D	Electrostatic steering	Amgen (Gunasekaran et al.)	<80%	NP
	D399K				
Control 2	T366S_L368A_Y407V	Knobs-into-holes	Genentech (Atwell et al.)	95%	69
	T366W				
Control 1	K409D_K392D	Electrostatic steering	Amgen (Gunasekaran et al.)	100%*	67
	D399K_E356K				
Control 5	IgG-IgA chimera	Strand Exchange	EMD Serono (Muda et al.)	>90%	68

5 * We observed a purity of >90% for Control 1 in our assay system, but not 100% as previously reported in the literature.

** We observed a T_m greater than 77°C for control 4 in our assay system; the T_m for this variant has not been published in the literature.

NP - The T_m for Control 3 has not been published and it was not tested in our assays
10 systems.

The melting temperature for wild-type IgG1 is shown as a range from 81-83 as the values in the literature vary depending on the assay system used, we report a value of 81.5°C in our assay system.

15 In contrast to negative design, a general concept used to engineer proteins is positive design. In this instance amino acid modifications are introduced into polypeptides to maximize favorable interactions within or between proteins. This

strategy assumes that when introducing multiple mutations that specifically stabilize the desired heterodimer while neglecting the effect on the homodimers, the net effect will be better specificity for the desired heterodimer interactions over the homodimers and hence a greater heterodimer specificity. It is understood in the context of protein engineering that positive design strategies optimize the stability of the desired protein interactions, but rarely achieve >90% specificity (Havranek JJ & Harbury PB.

Automated design of specificity in molecular recognition. *Nat Struct Biol.* 10(1):45-52 (2003); Bolon DN, Grant RA, Baker TA, Sauer RT. Specificity versus stability in computational protein design. *Proc Natl Acad Sci U S A.* 6;102(36):12724-9 (2005);

Huang PS, Love JJ, Mayo SL. A de novo designed protein protein interface *Protein Sci.* 16(12):2770-4 (2007)). Prior to this disclosure positive design strategies have not been used to design Fc heterodimers as more attention was devoted to specificity as compared to stability for therapeutic antibody manufacturing and development. In addition, beneficial positive design mutations can be hard to predict. Other

methodologies for improving stability, such as additional disulfide bonds, have been tried to improve stability in Fc heterodimers with limited success on improvements to the molecule. (See, Table A) This may be because all engineered Fc CH3 domain disulfide bonds are solvent exposed, which results in a short lifetime of the disulfide bond and therefore a significant impact on the long-term stability of the heterodimer - especially when the engineered CH3 domain has a T_m of less than 70°C without the additional disulfide bond (as in Control 4 which has a T_m of 69°C without the disulfide (see Control 2)). It is contemplated that other methodologies to improve stability, such as disulfide bonds, can also be used with the present Fc variants, provided the intrinsic stability (measured as melting temperature) of the CH3 domain is 70°C or greater without the disulfide bond, in particular when the intrinsic stability (measured as melting temperature) of the CH3 domain is 72°C or greater without the disulfide bond.

Therefore, we herein disclose a novel method for designing Fc heterodimers that results in both stable and highly specific heterodimer formation. This design method combines both negative and positive design strategies along with structural and computational modeling guided protein engineering techniques. This powerful method has allowed us to design novel combinations of mutations in the IgG1 CH3 domain wherein using only standard cell culture conditions heterodimers were formed with more than 90% purity compared to homodimers and the resulting heterodimers had a melting temperature of 70°C or greater. In exemplary embodiments, the Fc variant heterodimers have a melting temperature of 73°C or

greater and a purity of greater than 98%. In other exemplary embodiments, the Fc variant heterodimers have a melting temperature of 75°C or greater and a purity of greater than 90%. In certain embodiments of the heterodimer Fc variants described herein, the Fc variant heterodimers have a melting temperature of 77°C or greater
5 and a purity of greater than 98%. In some embodiments of the heterodimer Fc variants described herein, the Fc variant heterodimers have a melting temperature of 78°C or greater and a purity of greater than 98%. In certain embodiments of the heterodimer Fc variants described herein, the Fc variant heterodimers have a melting temperature of 79°C or greater and a purity of greater than 98%. In certain
10 embodiments of the heterodimer Fc variants described herein, the Fc variant heterodimers have a melting temperature of 80°C or greater and a purity of greater than 98%. In certain embodiments of the heterodimer Fc variants described herein, the Fc variant heterodimers have a melting temperature of 81°C or greater and a purity of greater than 98%.

15 In certain embodiments, an isolated heteromultimer comprising at least one heavy chain variable domain and a heterodimer Fc region is provided wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the
20 variant CH3 domain has a melting temperature (T_m) of 70°C or greater. As used herein "increased stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 70°C or greater. In certain embodiments, "increased stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 72°C or
25 greater. In certain embodiments, "increased stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 74°C or greater. In certain embodiments, "increased stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 75°C or greater. In certain embodiments, "increased
30 stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 76°C or greater. In certain embodiments, "increased stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 78°C or greater. In certain embodiments, "increased stability" or "stable heterodimer", refers
35 to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 79°C or greater. In certain embodiments, "increased stability" or "stable heterodimer", refers to a variant CH3 domain, in heterodimer formation, with a

melting temperature of about 80°C or greater. In certain embodiments, “increased stability” or “stable heterodimer”, refers to a variant CH3 domain, in heterodimer formation, with a melting temperature of about 81°C or greater. In addition, it is understood that the term “to promote heterodimer formation” refers herein to the amino acid mutations in the CH3 domain that result in greater than 90% heterodimer formation compared to homodimer formation.

In a further embodiment, this increased stability is in the absence of an additional disulfide bond. Specifically, the increased stability is in the absence of an additional disulfide bond in the CH3 domain. In one embodiment, the variant CH3 domain does not comprise an additional disulfide bond as compared to wild-type CH3 domain. In an alternative embodiment, the variant CH3 comprises at least one disulfide bond as compared to wild-type CH3 domain, provided that the variant CH3 has a melting temperature of 70°C or greater in the absence of the disulfide bond. In one embodiment, the variant CH3 domain comprises at least one disulfide bond as compared to wild-type CH3 domain, and the variant CH3 domain has a melting temperature (T_m) of about 77.5 °C or greater. In an embodiment, the variant CH3 domain comprises at least one disulfide bond as compared to wild-type CH3 domain, and the variant CH3 domain has a melting temperature (T_m) of about 78°C or greater. In another embodiment, the variant CH3 domain comprises at least one disulfide bond as compared to wild-type CH3 domain, and the variant CH3 domain has a melting temperature (T_m) of greater than about 78°C, or greater than about 78.5°C, or greater than about 79°C, or greater than about 79.5°C, or greater than about 80°C, or greater than about 80.5°C, or greater than about 81°C, or greater than about 81.5°C, or greater than about 82°C, or greater than about 82.5°C, or greater than about 83°C.

In one embodiment, the variant CH3 domain has a melting temperature of greater than about 70°C, or greater than about 70.5°C, or greater than about 71°C, or greater than about 71.5°C, or greater than about 72°C, or greater than about 72.5°C, or greater than about 73°C, or greater than about 73.5°C, or greater than about 74°C, or greater than about 74.5°C, or greater than about 75°C, or greater than about 75.5°C, or greater than about 76°C, or greater than about 76.5°C, or greater than about 77°C, or greater than about 77.5°C, or greater than about 78°C, or greater than about 78.5°C, or greater than about 79°C, or greater than about 79.5°C, or greater than about 80°C, or greater than about 80.5°C, or greater than about 81°C, or greater than about 81.5°C, or greater than about 82°C, or greater than about 82.5°C, or greater

than about 83°C. In another embodiment, the variant CH3 domain has a melting temperature of about 70°C, or about 70.5°C, or about 71°C, or about 71.5°C, or about 72°C, or about 72.5°C, or about 73°C, or about 73.5°C, or about 74°C, or about 74.5°C, or about 75°C, or about 75.5°C, or about 76°C, or about 76.5°C, or about 77°C, or about 77.5°C, or about 78°C, or about 78.5°C, or about 79°C, or about 79.5°C, or about 80°C, or about 80.5°C, or about 81°C. In yet another embodiment, the variant CH3 domain has a melting temperature of about 70°C to about 81°C, or about 70.5°C to about 81°C, or about 71°C to about 81°C, or about 71.5°C to about 81°C, or about 72°C to about 81°C, or about 72.5°C to about 81°C, or about 73°C to about 81°C, or about 73.5°C to about 81°C, or about 74°C to about 81°C, or about 74.5°C to about 81°C, or about 75°C to about 81°C, or about 75.5°C to about 81°C, or about 76°C to about 81°C, or about 76.5°C to about 81°C, or about 77°C to about 81°C, or about 77.5°C to about 81°C, or about 78°C to about 81°C, or about 78.5°C to about 82°C, or about 79°C to about 81°C. In yet another embodiment, the variant CH3 domain has a melting temperature of about 71°C to about 76°C, or about 72°C to about 76°C, or about 73°C to about 76°C, or about 74°C to about 76°C.

In addition to improved stability, the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation. It is understood that these amino acid mutations to promote heterodimer formation are as compared to homodimer formation. This heterodimer formation as compared to homodimer formation is referred jointly herein as “purity” or “specificity” or “heterodimer purity” or “heterodimer specificity”. It is understood that the heterodimer purity refers to the percentage of desired heterodimer formed as compared to homodimer species formed in solution under standard cell culture conditions prior to selective purification of the heterodimer species. For instance, a heterodimer purity of 90% indicates that 90% of the dimer species in solution is the desired heterodimer. In one embodiment, the Fc variant heterodimers have a purity of greater than about 90%, or greater than about 91%, or greater than about 92%, or greater than about 93%, or greater than about 94%, or greater than about 95%, or greater than about 96%, or greater than about 97%, or greater than about 98%, or greater than about 99%. In another embodiment, the Fc variant heterodimers have a purity of about 90%, or about 91%, or about 92%, or about 93%, or about 94%, or about 95%, or about 96%, or about 97%, or about 98%, or about 99%, or about 100%.

In a specific embodiment, is provided an isolated heteromultimer comprising at least one heavy chain variable domain and a heterodimer Fc region, wherein the

heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of 70 °C or greater and the resulting heterodimer has a purity greater than 90%. In one aspect, the resulting heterodimer has a purity greater than 98% and the variant CH3 domain has a melting temperature of greater than about 70°C, or greater than about 71°C, or greater than about 72°C, or greater than about 73°C, or greater than about 74°C, or greater than about 75°C, or greater than about 76°C, or greater than about 77°C, or greater than about 78°C, or greater than about 79°C, or greater than about 80°C or greater than about 81°C. In a further aspect, the variant CH3 domain has a melting temperature of 70°C or greater and the resulting Fc variant heterodimer has a purity greater than about 90%, or greater than about 91%, or greater than about 92%, or greater than about 93%, or greater than about 94%, or greater than about 95%, or greater than about 96%, or greater than about 97%, or greater than about 98%, or greater than about 99%.

In order to design these heteromultimers comprising Fc variants with improved stability and purity we employed an iterative process of computational design and experimental screening to select the most successful combinations of positive and negative design strategies (See, Figure 24).

Specifically, in the initial design phase different negative design Fc variant heterodimers were made and tested for expression and stability as described in Examples 1-3. The initial design phase included Fc variant heterodimers AZ1-AZ16 (See, Table 1). From this initial set of negative design Fc variant heterodimers, which were expected to have low stability (e.g., a T_m of less than 71°C), the Fc variant heterodimers with greater than 90% purity and a melting temperature of about 68°C or greater were selected for further development. This included Fc variant heterodimers AZ6, AZ8 and AZ15. In the second design phase, those selected Fc variant heterodimers were further modified to drive both stability and purity using positive design strategies following a detailed computational and structural analysis. The selected Fc variant heterodimers (AZ6, AZ8, and AZ15) were each analyzed with computational methods and comprehensive structure function analysis to identify the structural reasons these Fc variants had a lower stability than the wild-type Fc homodimer, which is 81°C for IgG1. See, Table 4 for the list of Fc variant heterodimers and the T_m values.

In certain embodiments, the variant CH3 domain is selected from AZ1, or AZ2, or AZ3, or AZ4, or AZ5, or AZ6, or AZ7, or AZ8, or AZ9, or AZ10, or AZ11, or AZ12, or AZ13, or AZ14, or AZ15, or AZ16. In selected embodiments, the variant CH3 domain is AZ6, or AZ8 or AZ15.

5

The computational tools and structure-function analysis included, but were not limited to molecular dynamic analysis (MD), sidechain/backbone re-packing, Knowledge Base Potential (KBP), cavity and (hydrophobic) packing analysis (LJ, CCSD, SASA, dSASA(carbon/all-atom)), electrostatic-GB calculations, and coupling analysis. (See, Figure 24 for an overview of the computational strategy)

10

An aspect of the protein engineering approach relied on combining structural information of the Fc IgG protein derived from X-ray crystallography with computational modeling and simulation of the wild type and variant forms of the CH3 domain. This allowed us to gain novel structural and physico-chemical insights about the potential role of individual amino acids and their cooperative action. These structural and physico-chemical insights, obtained from multiple variant CH3 domains, along with the resulting empirical data pertaining to their stability and purity helped us develop an understanding for the relationship between purity and stability of the heterodimer. In order to execute our simulations we started by building complete and realistic models and refining the quality of the wild type Fc structure of an IgG1 antibody. Protein structures derived from X-ray crystallography are lacking in detail regarding certain features of the protein in aqueous medium under physiological condition and our refinement procedures addressed these limitations. These include building missing regions of the protein structure, often flexible portions of the protein such as loops and some residue side chains, evaluating and defining the protonation states of the neutral and charged residues and placement of potential functionally relevant water molecules associated with the protein.

15

20

25

30

35

Molecular dynamics (MD) algorithms are one tool we used, by simulating the protein structure, to evaluate the intrinsic dynamic nature of the Fc homodimer and the variant CH3 domains in an aqueous environment. Molecular dynamics simulations track the dynamic trajectory of a molecule resulting from motions arising out of interactions and forces acting between all the atomic entities in the protein and its local environment, in this case the atoms constituting the Fc and its surrounding water molecules. Following molecular dynamics simulations, various aspects of the trajectories were analyzed to gain insight into the structural and dynamic

characteristics of the Fc homodimer and variant Fc heterodimer, which we used to identify specific amino acid mutations to improve both purity and stability of the molecule.

5 Therefore, the generated MD trajectories were studied using methods such as the principal component analysis to reveal the intrinsic low frequency modes of motion in the Fc structure. This provides insight into the potential conformational sub-states of the protein (See, Figure 32). While the critical protein-protein interactions between chain A and B in the Fc region occur at the interface of the CH3 domains, our
10 simulations indicated that this interface acts as a hinge in a motion that involves the “opening” and “closing” of the N-terminal ends of the CH2 domains relative to each other. The CH2 domain interacts with FcγR’s at this end as seen in figure 16. Thus, while not wishing to be bound by a theory, it appears that introduction of amino acid mutations at the CH3 interface impacts the magnitude and nature of the open/close
15 motion at the N-terminal end of the Fc and therefore how the Fc interacts with the FcγR’s. See, example 4 and Table 5.

The generated MD trajectories were also studied to determine the mutability of specific amino acid residue positions in the Fc structure based on profiling their
20 flexibility and analysis of their environment. This algorithm allowed us to identify residues that could affect protein structure and function, providing unique insight into residue characteristics and mutability for subsequent design phases of the variant CH3 domains. This analysis also enabled us to compare multiple simulations, and assess mutability based on outliers following profiling.

25 The generated MD trajectories were also studied to determine correlated residue motions in the protein and the formation of networks of residues as a result of coupling between them. Finding dynamic correlations and networks of residues within the Fc structure is a critical step in understanding the protein as a dynamic
30 entity and for developing insight into the effects of mutations at distal sites. See, e.g. Example 6

Thus, we studied in detail the impact of mutations on the local environment of the site of mutation. The formation of a well packed core at the CH3 interface between chain
35 A and B is critical for the spontaneous pairing of the two chains in a stable Fc structure. Good packing is the result of strong structural complementarity between interacting molecular partners coupled with favorable interactions between the

contacting groups. The favorable interactions result from either buried hydrophobic contacts well removed from solvent exposure and/or from the formation of complementary electrostatic contacts between hydrophilic polar groups. These hydrophobic and hydrophilic contacts have entropic and enthalpic contributions to the free energy of dimer formation at the CH3 interface. We employ a variety of algorithms to accurately model the packing at the CH3 interface between chain A and chain B and subsequently evaluate the thermodynamic properties of the interface by scoring a number of relevant physicochemical properties.

We employed a number of protein packing methods including flexible backbones to optimize and prepare model structures for the large number of variants we computationally screened. Following packing we evaluated a number of terms including contact density, clash score, hydrogen bonds, hydrophobicity and electrostatics. The use of the solvation models allowed us to more accurately address the effect of solvent environment and contrast the free energy differences following mutation of specific positions in the protein to alternate residue types. Contact density and clash score provide a measure of complementarity, a critical aspect of effective protein packing. These screening procedures are based on the application of knowledge-based potentials or coupling analysis schemes relying on pair-wise residue interaction energy and entropy computations.

This comprehensive in-silico analysis provided a detailed understanding of the differences of each Fc variant compared to wild-type with respect to interface hotspots, sites of asymmetry, cavities and poorly packed regions, structural dynamics of individual sites and sites of local unfolding. These combined results of the described computational analysis identified specific residues, sequence/structural motifs and cavities that were not optimized and in combination responsible for the lower stability (e.g., T_m of 68°C) and/or lower specificity of <90% purity. In the second design phase we used targeted positive design to specifically address these hypothesis by additional point-mutations and tested these by in-silico engineering using the above described methodology and analysis (See, Figure 24). The Fc variant heterodimers designed to improve stability and purity for each targeted design in phase two (Fc variant heterodimers AZ17-AZ101) were validated experimentally for expression and stability as described in Examples 1-4.

In certain embodiments, provided herein are isolated heteromultimers comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3

domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain is AZ17, or AZ18, or AZ19, or AZ20, or AZ21, or AZ22, or AZ23, or AZ24, or AZ25, or AZ26, or AZ27, or AZ28, or AZ29, or AZ30, or AZ31, or AZ32, or AZ33, or AZ34, or AZ35, or AZ36, or AZ37, or AZ38, or AZ39, or AZ40, or AZ41, or AZ42, or AZ43, or AZ44, or AZ45, or AZ46, or AZ47, or AZ48, or AZ49, or AZ50, or AZ51, or AZ52, or AZ53, or AZ54, or AZ55, or AZ56, or AZ57, or AZ58, or AZ59, or AZ60, or AZ61, or AZ62, or AZ63, or AZ64, or AZ65, or AZ66, or AZ67, or AZ68, or AZ69, or AZ70, or AZ71, or AZ72, or AZ73, or AZ74, or AZ75, or AZ76, or AZ77, or AZ78, or AZ79, or AZ80, or AZ81, or AZ82, or AZ83, or AZ84, or AZ85, or AZ86, or AZ87, or AZ88, or AZ89, or AZ90, or AZ91, or AZ92, or AZ93, or AZ94, or AZ95, or AZ96, or AZ97, or AZ98, or AZ99, or AZ100 or AZ101. In an exemplary embodiment, the variant CH3 domain is AZ17, or AZ18, or AZ19, or AZ20, or AZ21, or AZ22, or AZ23, or AZ24, or AZ25, or AZ26, or AZ27, or AZ28, or AZ29, or AZ30, or AZ31, or AZ32, or AZ33, or AZ34, or AZ38, or AZ42, or AZ43, or AZ44, or AZ45, or AZ46, or AZ47, or AZ48, or AZ49, or AZ50, or AZ52, or AZ53, or AZ54, or AZ58, or AZ59, or AZ60, or AZ61, or AZ62, or AZ63, or AZ64, or AZ65, or AZ66, or AZ67, or AZ68, or AZ69, or AZ70, or AZ71, or AZ72, or AZ73, or AZ74, or AZ75, or AZ76, or AZ77, or AZ78, or AZ79, or AZ81, or AZ82, or AZ83, or AZ84, or AZ85, or AZ86, or AZ87, or AZ88, or AZ89, or AZ91, or AZ92, or AZ93, or AZ94, or AZ95, or AZ98, or AZ99, or AZ100 or AZ101. In a specific embodiment, the variant CH3 domain is AZ33 or AZ34. In another embodiment, the variant CH3 domain is AZ70 or AZ90.

In an exemplary embodiment, the heteromultimer comprises a first and a second polypeptide, wherein the first polypeptide comprises amino acid modifications L351Y, F405A, and Y407V and wherein the second polypeptide comprises amino acid modifications T366I, K392M and T394W. In another embodiment, a first polypeptide comprises amino acid modifications L351Y, S400E, F405A and Y407V and the second polypeptide comprises amino acid modifications T366I, N390R, K392M and T394W.

This iterative process of computational structure-function analysis, targeted engineering and experimental validation was used to design the remaining Fc variants listed in Table 1 in subsequent design phases and resulting in heteromultimers with a purity greater than 90% and an increased stability with a CH3 domain melting temperature greater than 70°C. In certain embodiments, the Fc variants comprise amino acid mutations selected from AZ1 to AZ 136. In further

embodiments, the Fc variants comprise amino acid mutations selected from the Fc variants listed in Table 4.

From the first and second design phases two core scaffolds were identified, Scaffold 1 and Scaffold 2, wherein additional amino acid modifications were introduced into these scaffolds to fine tune the purity and stability of the Fc variant heterodimers. See Example 5 for a detailed description of the development of Scaffold 1 including AZ8, AZ17-62 and the variants listed in Table 6. See Example 6 for a detailed description of the development of Scaffold 2 including AZ15 and AZ63-101 and the variants listed in Table 7.

The core mutations of Scaffold 1 comprise L351Y_F405A_Y407V / T394W. Scaffold 1a comprises the amino acid mutations T366I_K392M_T394W/F405A_Y407V and Scaffold 1b comprises the amino acid mutations T366L_K392M_T394W/F405A_Y407V. See, Example 5.

In certain embodiments, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications L351Y, F405A and Y407V and the second polypeptide comprises amino acid modification T394W. In one aspect the heteromultimer further comprises point mutations at positions F405 and/or K392. These mutations at position K392 include, but are not limited to, K392V, K392M, K392R, K392L, K392F or K392E. These mutations at position F405 include, but are not limited to, F405I, F405M, F405S, F405S, F405V or F405W. In another aspect, the heteromultimer further comprises point mutations at positions T411 and/or S400. These mutations at position T411 include, but are not limited to, T411N, T411R, T411Q, T411K, T411D, T411E or T411W. These mutations at position S400 include, but are not limited to, S400E, S400D, S400R or S400K. In yet another embodiment, the heteromultimer comprises a first and second polypeptide wherein the first polypeptide comprises amino acid modifications L351Y, F405A and Y407V and the second polypeptide comprises amino acid modification T394W, wherein the first and/or second polypeptide comprises further amino acid modifications at positions T366 and/or L368. These mutations at position T366 include, but are not limited to, T366A, T366I, T366L, T366M, T366Y, T366S, T366C, T366V or T366W. In an exemplary embodiment, the amino acid mutation at position T366 is T366I. In another exemplary embodiment, the amino acid mutation at position T366 is T366L.

The mutations at position L368 include, but are not limited to, L368D, L368R, L368T, L368M, L368V, L368F, L368S and L368A.

In certain embodiments, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications L351Y, F405A and Y407V and the second polypeptide comprises amino acid modifications T366L and T394W. In another embodiment, the heteromultimer comprises a first and second polypeptide wherein the first polypeptide comprises amino acid modifications L351Y, F405A and Y407V and the second polypeptide comprises amino acid modifications T366I and T394W.

In certain other embodiments, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications L351Y, F405A and Y407V and the second polypeptide comprises amino acid modifications T366L, K392M and T394W. In another embodiment, the heteromultimer comprises a first and second polypeptide wherein the first polypeptide comprises amino acid modifications L351Y, F405A and Y407V and the second polypeptide comprises amino acid modifications T366I, K392M and T394W.

In yet another embodiment, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications F405A and Y407V and the second polypeptide comprises amino acid modifications T366L, K392M and T394W. In another embodiment, the heteromultimer comprises a first and second polypeptide wherein the first polypeptide comprises amino acid modifications F405A and Y407V and the second polypeptide comprises amino acid modifications T366I, K392M and T394W.

In certain embodiments, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications F405A and Y407V and the second polypeptide comprises amino acid modifications T366L and T394W. In another embodiment, the heteromultimer comprises a first and second polypeptide wherein the first polypeptide comprises amino acid modifications F405A and Y407V and the second polypeptide comprises amino acid modifications T366I and T394W.

In an exemplary embodiment, provided herein are isolated heteromultimers comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a heteromultimer comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the heteromultimer has a melting temperature (T_m) of about 74°C or greater. In another embodiment, provided herein are isolated heteromultimers comprising a heterodimer Fc region, wherein the heteromultimer comprises at least one heavy chain variable domain and a variant CH3 region comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of about 74°C or greater and the heterodimer has a purity of about 98% or greater.

In certain embodiments, the isolated heteromultimer comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) greater than 70 °C and the variant CH3 domains are selected from Table 6.

The core mutations of Scaffold 2 comprise L351Y_Y407A / T366A_K409F. Scaffold 2a comprises the amino acid mutations L351Y_Y407A / T366V_K409F and Scaffold 2b comprises the amino acid mutations Y407A / T366A_K409F. See, Example 6.

In certain embodiments, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications L351Y and Y407A and the second polypeptide comprises amino acid modifications T366A and K409F. In one aspect the heteromultimer further comprises point mutations at positions T366, L351, and Y407. These mutations at position T366 include, but are not limited to, T366I, T366L, T366M, T366Y, T366S, T366C, T366V or T366W. In a specific embodiment, the mutation at position T366 is T366V. The mutations at position L351 include, but are not limited to, L351I, L351D, L351R or L351F. The mutations at position Y407 include, but are not limited to, Y407V or Y407S. See, CH3 variants AZ63-AZ70 in Table 1 and Table 4 and Example 6.

In an exemplary embodiment, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first

polypeptide comprises amino acid modifications L351Y and Y407A and the second polypeptide comprises amino acid modification T366V and K409F.

In an exemplary embodiment, provided herein are isolated heteromultimers comprising at least one single domain antigen-binding construct, and a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of about 75.5°C or greater; and wherein the heteromultimer is devoid of immunoglobulin light chains, immunoglobulin CH1 and optionally devoid of immunoglobulin CH2 regions.

In another embodiment, provided herein are isolated heteromultimers comprising at least one heavy chain variable region and a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of about 75°C or greater and the heterodimer has a purity of about 90% or greater.

In other certain embodiments, the heteromultimer comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modifications L351Y and Y407A and the second polypeptide comprises amino acid modification T366A and K409F, wherein the variant CH3 domain comprises one or more amino acid modifications at positions T411, D399, S400, F405, N390, and/or K392. These mutations at position D399 include, but are not limited to, D399R, D399W, D399Y or D399K. The mutations at position T411 includes, but are not limited to, T411N, T411R, T411Q, T411K, T411D, T411E or T411W. The mutations at position S400 includes, but are not limited to, S400E, S400D, S400R, or S400K. The mutations at position F405 includes, but are not limited to, F405I, F405M, F405S, F405V or F405W. The mutations at position N390 include, but are not limited to, N390R, N390K or N390D. The mutations at position K392 include, but are not limited to, K392V, K392M, K392R, K392L, K392F or K392E. See, CH3 variants AZ71-101 in Table 1 and Table 4 and Example 6.

In an exemplary embodiment, the heteromultimer is a heterodimer that comprises a first and second polypeptide (also referred to herein as Chain A and Chain B) wherein the first polypeptide comprises amino acid modification Y407A and the second polypeptide comprises amino acid modification T366A and K409F. In one aspect, this heterodimer further comprises the amino acid modifications K392E,

T411E, D399R and S400R. In a further embodiment, the heteromultimer comprises a first and second polypeptide wherein the first polypeptide comprises amino acid modification D399R, S400R and Y407A and the second polypeptide comprises amino acid modification T366A, K409F, K392E and T411E.

- 5 In an exemplary embodiment, provided herein are isolated heteromultimers comprising at least one heavy chain variable region and a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) of about 74°C or
10 greater, and wherein the heteromultimer is devoid of immunoglobulin light chains and CH₁ domains. In another embodiment, provided herein are isolated heteromultimers comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting
15 temperature (T_m) of about 74°C or greater and the heterodimer has a purity of about 95% or greater, and wherein the heteromultimer is devoid of immunoglobulin light chains and CH₁ domains.

- In certain embodiments, provided herein are isolated heteromultimers comprising a
20 heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) greater than 70 °C and the variant CH3 domains are selected from Table 7, and wherein the heteromultimer is devoid of immunoglobulin light chains and CH₁
25 domains.

- Furthermore, this new method of designing heteromultimers comprising at least one single domain antigen-binding construct and Fc variant heterodimers with improved stability and purity can be applied to other classes and isotypes of Fc regions. In
30 certain embodiments, the Fc region is a human IgG Fc region. In further embodiments, the human IgG Fc region is a human IgG1, IgG2, IgG3, or IgG4 Fc region. In some embodiments the Fc regions is from an immunoglobulin selected from the group consisting of IgG, IgA, IgD, IgE and IgM. In some embodiments, the IgG is of subtype selected from the group consisting of IgG1, IgG2a, IgG2b, IgG3
35 and IgG4.

Table 1.1: CH3 domain amino acid modifications for the generation of Fc variant heterodimers.

Wild-Type IgG1	Variant	Chain	Fc Mutations							
		A	-	-	-	-	-	-	-	-
		B	-	-	-	-	-	-	-	-
CH3 Variants	AZ1	A	L368D	K370Q	-	-	-	-	-	-
		B	E357R	L368R	-	-	-	-	-	-
	AZ2	A	L351I	L368D	K370Q	-	-	-	-	-
		B	E357R	L368R	-	-	-	-	-	-
	AZ3	A	L351D	L368D	K370Q	-	-	-	-	-
		B	E357R	L368R	-	-	-	-	-	-
	AZ4	A	L368D	K370E	-	-	-	-	-	-
		B	E357R	L368R	-	-	-	-	-	-
	AZ5	A	L368D	K370E	-	-	-	-	-	-
		B	E357K	L368R	-	-	-	-	-	-
	AZ6	A	V397S	F405A	Y407V	-	-	-	-	-
		B	K392V	T394W	-	-	-	-	-	-
	AZ7	A	L351R	V397S	F405A	Y407V	-	-	-	-
		B	K392V	T394W	-	-	-	-	-	-
	AZ8	A	L351Y	V397S	F405A	Y407V	-	-	-	-
		B	K392V	T394W	-	-	-	-	-	-
	AZ9	A	V397S	F405A	Y407V	-	-	-	-	-
		B	L368R	K392V	T394W	-	-	-	-	-
	AZ10	A	V397T	F405I	-	-	-	-	-	-
		B	K392V	T394H	-	-	-	-	-	-
	AZ11	A	E357W	S364F	-	-	-	-	-	-
		B	Y349A	L351Y	K370I	-	-	-	-	-

AZ12	A	E357H	S364F	-	-	-	-	-
	B	L351Y	K370I	-	-	-	-	-
AZ13	A	E357W	S364F	-	-	-	-	-
	B	Y349A	L351Y	K370F	-	-	-	-
AZ14	A	E357H	S364F	-	-	-	-	-
	B	L351Y	K370F	-	-	-	-	-
AZ15	A	E357L	T366A	K409F	T411N	-	-	-
	B	L351Y	Y407A	-	-	-	-	-
AZ16	A	E357L	T366A	K409Y	T411N	-	-	-
	B	L351Y	L368T	Y407A	-	-	-	-
AZ17	A	L351Y	F405A	Y407V	-	-	-	-
	B	T366I	T394W	-	-	-	-	-
AZ18	A	L351Y	V397T	F405M	Y407V	-	-	-
	B	T366I	T394W	-	-	-	-	-
AZ19	A	L351Y	V397T	F405M	Y407V	-	-	-
	B	T366L	T394W	-	-	-	-	-
AZ20	A	L351Y	V397T	F405M	Y407V	-	-	-
	B	T366M	T394W	-	-	-	-	-
AZ21	A	L351Y	L368M	V397T	F405I	Y407V	-	-
	B	T366L	T394W	-	-	-	-	-
AZ22	A	L351Y	L368M	V397T	F405I	Y407V	-	-
	B	T366M	T394W	-	-	-	-	-
AZ23	A	L351Y	V397T	F405M	Y407V	-	-	-
	B	L351I	T366I	T394W	-	-	-	-
AZ24	A	L351Y	V397T	L398D	F405M	Y407V	-	-
	B	S354E	T366I	T394W	-	-	-	-
AZ25	A	L351Y	V397T	L398D	S400E	F405M	Y407V	-
	B	T366I	N390R	T394W	-	-	-	-
AZ26	A	R344H	L351Y	V397T	S400E	F405M	Y407V	-
	B	Q362R	T366I	T394W	-	-	-	-
AZ27	A	R344H	L351Y	V397T	D401E	F405M	Y407V	-
	B	Q362R	T366I	T394W	-	-	-	-
AZ28	A	Q347R	L351Y	V397T	F405M	Y407V	-	-
	B	S354E	K360E	T366I	T394W	-	-	-

AZ29	A	Q347R	L351Y	V397T	F405M	Y407V	-	-
	B	S354N	K360E	T366I	T394W	-	-	-
AZ30	A	T350V	L351Y	V397T	S400E	F405M	Y407V	-
	B	T350V	T366I	T394W	T411R	-	-	-
AZ31	A	R344H	L351Y	V397T	L398D	F405M	Y407V	-
	B	T366I	T394W	T411R	-	-	-	-
AZ32	A	Q347R	T350V	L351Y	V397T	F405M	Y407V	-
	B	T350V	K360E	T366I	T394W	T411R	-	-
AZ33	A	L351Y	F405A	Y407V	-	-	-	-
	B	T366I	K392M	T394W	-	-	-	-
AZ34	A	L351Y	S400E	F405A	Y407V	-	-	-
	B	T366I	N390R	K392M	T394W	-	-	-
AZ35	A	L351Y	K370Q	G371D	F405M	Y407V	-	-
	B	Q362R	T366I	T394W	K409R	T411Q	-	-
AZ36	A	L351Y	K370Q	G371D	F405S	Y407V	-	-
	B	Q362R	T366I	T394W	K409R	T411Q	-	-
AZ37	A	R344H	L351Y	K370Q	G371D	L398D	F405M	Y407V
	B	Q362R	T366I	T394W	K409R	T411Q	-	-
AZ38	A	R344H	L351Y	K370Q	G371D	S400E	F405M	Y407V
	B	Q362R	T366I	N390R	T394W	K409R	T411Q	-
AZ39	A	L351Y	K370Q	G371D	F405M	Y407V	-	-
	B	T366I	T394W	T411R	-	-	-	-
AZ40	A	L351Y	K370Q	G371D	F405M	Y407V	-	-
	B	T366I	T394W	K409M	T411R	-	-	-
AZ41	A	R344H	L351Y	K370Q	G371D	L398D	F405M	Y407V
	B	T366I	T394W	K409M	T411R	-	-	-
AZ42	A	R344H	L351Y	K370Q	G371D	S400E	F405M	Y407V
	B	T366I	N390R	T394W	K409M	T411R	-	-
AZ43	A	L351Y	K370T	G371D	F405I	Y407V	-	-
	B	E357Q	S364R	T394W	-	-	-	-
AZ44	A	L351Y	K370T	G371D	F405M	Y407V	-	-
	B	E357Q	S364R	T394W	K409I	-	-	-
AZ45	A	R344H	L351Y	K370T	G371D	S400E	F405M	Y407V
	B	E357Q	S364R	T366I	N390R	T394W	K409I	-

AZ46	A	R344H	L351Y	K370T	G371D	F405M	Y407V	-
	B	E357Q	S364R	T366I	T394W	K409I	T411R	-
AZ47	A	L351Y	K370A	G371S	D399R	F405S	Y407V	-
	B	E357Q	Q362R	T364Y	T366I	T394W	K409S	-
AZ48	A	L351Y	V397S	D399W	F405M	Y407V	-	-
	B	Q362R	T366I	T394W	K409M	-	-	-
AZ49	A	L351Y	V397S	D399Y	F405M	Y407V	-	-
	B	Q362R	T366I	T394W	K409I	-	-	-
AZ50	A	R344H	L351Y	V397T	L398D	D399W	F405M	Y407V
	B	Q362R	T366I	T394W	K409M	-	-	-
AZ51	A	R344H	L351Y	V397T	D399W	S400E	F405M	Y407V
	B	Q362R	T366I	T394W	K409M	-	-	-
AZ52	A	L368V	K370F	F405I	Y407V	-	-	-
	B	E357Q	S364Y	T366I	T394W	-	-	-
AZ53	A	L368V	K370Y	F405I	Y407V	-	-	-
	B	E357Q	S364Y	T394W	-	-	-	-
AZ54	A	R344H	L368V	K370Y	F405M	Y407V	-	-
	B	E357Q	Q362R	S364Y	T394W	-	-	-
AZ55	A	L368V	K370Y	S400E	F405M	Y407V	-	-
	B	E357Q	S364Y	N390R	T394W	-	-	-
AZ56	A	L368V	K370Y	L398D	F405M	Y407V	-	-
	B	E357Q	S364Y	T394W	T411R	-	-	-
AZ57	A	R344H	L351Y	K370Y	F405M	Y407V	-	-
	B	E357Q	Q362R	T364T	T366I	T394W	-	-
AZ58	A	L368V	V397T	F405M	Y407V	-	-	-
	B	T366Y	T394W	-	-	-	-	-
AZ59	A	L368V	K370Q	V397T	F405M	Y407V	-	-
	B	T366Y	T394W	-	-	-	-	-
AZ60	A	R344H	L368V	V397T	S400E	F405M	Y407V	-
	B	Q362R	T366Y	T394W	-	-	-	-
AZ61	A	L368V	V397T	S400E	F405M	Y407V	-	-
	B	T366Y	N390R	T394W	-	-	-	-
AZ62	A	L368V	V397T	L398D	F405M	Y407V	-	-
	B	T366Y	T394W	T411R	-	-	-	-

AZ63	A	T366A	K409F	-	-	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ64	A	T366A	K409F	-	-	-	-	-
	B	L351Y	Y407A	-	-	-	-	-
AZ65	A	T366A	K409F	-	-	-	-	-
	B	L351F	Y407A	-	-	-	-	-
AZ66	A	T366S	K409F	-	-	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ67	A	T366C	K409F	-	-	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ68	A	T366L	K409F	-	-	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ69	A	T366M	K409F	-	-	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ70	A	T366V	K409F	-	-	-	-	-
	B	L351Y	Y407A	-	-	-	-	-
AZ71	A	T366A	K409F	-	-	-	-	-
	B	L351I	T366S	L368F	Y407A	-	-	-
AZ72	A	T366A	K409F	-	-	-	-	-
	B	D399W	Y407A	-	-	-	-	-
AZ73	A	T366A	K409F	-	-	-	-	-
	B	D399W	S400D	Y407A	-	-	-	-
AZ74	A	T366A	K409F	-	-	-	-	-
	B	D399W	S400E	Y407A	-	-	-	-
AZ75	A	T366A	K409F	T411R	-	-	-	-
	B	D399W	S400D	Y407A	-	-	-	-
AZ76	A	T366A	K409F	T411R	-	-	-	-
	B	G371D	D399W	Y407A	-	-	-	-
AZ77	A	T366A	K409F	T411R	-	-	-	-
	B	K370Q	G371D	D399W	Y407A	-	-	-
AZ78	A	T366A	N390R	K409F	-	-	-	-
	B	D399Y	S400D	Y407A	-	-	-	-
AZ79	A	Q362R	T366A	K409F	T411K	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ80	A	Q362R	T366A	K409F	T411R	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ81	A	Q362K	T366A	K409F	T411R	-	-	-
	B	Y407A	-	-	-	-	-	-

AZ82	A	T366A	N390K	K392R	K409F	T411R	-	-
	B	S400E	Y407A	-	-	-	-	-
AZ83	A	T366A	N390K	K392R	K409F	T411K	-	-
	B	S400E	Y407A	-	-	-	-	-
AZ84	A	T366A	N390K	K409F	T411R	-	-	-
	B	S400D	Y407A	-	-	-	-	-
AZ85	A	T366A	K392L	K409F	T411D	-	-	-
	B	D399R	Y407A	-	-	-	-	-
AZ86	A	T366A	K392L	K409F	T411E	-	-	-
	B	D399R	Y407A	-	-	-	-	-
AZ87	A	T366A	K392L	K409F	T411D	-	-	-
	B	D399K	Y407A	-	-	-	-	-
AZ88	A	T366A	K392L	K409F	T411E	-	-	-
	B	D399K	Y407A	-	-	-	-	-
AZ89	A	T366A	K392M	K409F	T411E	-	-	-
	B	D399R	Y407A	-	-	-	-	-
AZ90	A	T366A	K392M	K409F	T411D	-	-	-
	B	D399R	Y407A	-	-	-	-	-
AZ91	A	T366A	K392F	K409F	T411D	-	-	-
	B	D399R	F405V	Y407A	-	-	-	-
AZ92	A	T366A	K409F	T411E	-	-	-	-
	B	D399R	S400E	Y407A	-	-	-	-
AZ93	A	T366A	K409F	T411E	-	-	-	-
	B	D399R	S400D	Y407A	-	-	-	-
AZ94	A	T366A	K392E	K409F	T411E	-	-	-
	B	D399R	S400R	Y407A	-	-	-	-
AZ95	A	T366A	K392E	K409F	T411D	-	-	-
	B	D399R	S400R	Y407A	-	-	-	-
AZ96	A	Q362E	T366A	K409F	T411W	-	-	-
	B	D399R	Y407A	-	-	-	-	-
AZ97	A	Q362D	T366A	K409F	T411W	-	-	-
	B	D399R	Y407A	-	-	-	-	-
AZ98	A	S364Y	T366A	K409F	T411R	-	-	-
	B	Y407A	-	-	-	-	-	-
AZ99	A	T366V	K409W	-	-	-	-	-
	B	L368V	Y407S	-	-	-	-	-
AZ100	A	T366V	K409W	-	-	-	-	-
	B	L351Y	L368S	Y407A	-	-	-	-
AZ101	A	T366V	K409W	-	-	-	-	-
	B	L351Y	Y407A	-	-	-	-	-
AZ102	A	E357Q	S364F	K392E	-	-	-	-
	B	K370F	V397R	S400R	-	-	-	-

AZ103	A	E357Q	S364F	K392E	V397E	-	-	-
	B	K370F	V397R	S400R	-	-	-	-
AZ104	A	E357Q	S364F	N390D	K392E	-	-	-
	B	K370F	V397R	S400K	-	-	-	-
AZ105	A	E357Q	S364F	K370E	G371W	-	-	-
	B	E357Q	K360R	S364N	K370F	-	-	-
AZ106	A	S354R	D356K	E357Q	S364F	-	-	-
	B	S354E	K370F	K439E	-	-	-	-
AZ107	A	Q347R	E357Q	S364F	-	-	-	-
	B	Q347E	K360E	K370F	-	-	-	-
AZ108	A	E357Q	S364F	K370E	-	-	-	-
	B	E357R	K370F	-	-	-	-	-
AZ109	A	E357Q	S364F	L368D	K370E	-	-	-
	B	E357R	K370F	-	-	-	-	-
AZ110	A	E357Q	S364F	K370T	G371D	-	-	-
	B	E357Q	S364R	K370F	-	-	-	-
AZ111	A	E357Q	S364Y	K392E	-	-	-	-
	B	K370F	V397R	S400K	-	-	-	-
AZ112	A	E357Q	S364Y	K392E	-	-	-	-
	B	L368A	K370F	V397R	S400K	-	-	-
AZ113	A	K409F	T411E	-	-	-	-	-
	B	L368V	D399R	S400D	-	-	-	-
AZ114	A	K409F	T411E	-	-	-	-	-
	B	L368V	D399K	S400D	-	-	-	-
AZ115	A	K409F	-	-	-	-	-	-
	B	L368V	D399Y	-	-	-	-	-
AZ116	A	E357Q	K409F	T411R	-	-	-	-
	B	L368A	K370F	-	-	-	-	-
AZ117	A	S354R	D356K	K409F	T411R	-	-	-
	B	S354E	L368V	S400E	K439E	-	-	-
AZ118	A	K360E	K370E	-	-	-	-	-
	B	Y349R	E357R	-	-	-	-	-
AZ119	A	K360E	K370E	-	-	-	-	-
	B	Y349K	E357R	-	-	-	-	-
AZ120	A	S354E	K360E	K370E	-	-	-	-
	B	Y349R	E357R	-	-	-	-	-
AZ121	A	K360E	L368D	K370E	-	-	-	-
	B	Y349R	E357R	-	-	-	-	-
AZ122	A	K360E	L368D	K370E	-	-	-	-
	B	Y349R	E357R	T411R	-	-	-	-
AZ123	A	K360E	K370T	G371D	-	-	-	-
	B	Y349R	E357Q	S364R	-	-	-	-

AZ124	A	K360E	K370T	G371D	-	-	-	-
	B	Y349R	E357Q	S364K	-	-	-	-
AZ125	A	S364E	K370T	G371D	-	-	-	-
	B	E357Q	S364R	G371R	-	-	-	-
AZ126	A	S364E	K370T	G371D	-	-	-	-
	B	E357Q	S364R	G371K	-	-	-	-
AZ127	A	G371D	T411E	-	-	-	-	-
	B	G371R	T411R	-	-	-	-	-
AZ128	A	G371D	T411E	-	-	-	-	-
	B	G371K	T411R	-	-	-	-	-
AZ129	A	Y349C	L351Y	V397T	F405M	Y407V	-	-
	B	S354C	T366I	T394W	-	-	-	-
AZ130	A	L351Y	S354C	V397T	F405M	Y407V	-	-
	B	Y349C	T366I	T394W	-	-	-	-
AZ132	A	L368A	F405W	Y407V	-	-	-	-
	B	T366W	-	-	-	-	-	-

Table 1.2: CH3 domain amino acid modifications for the generation of Fc variant heterodimers. The DSC melting temperature of the CH3 domain was estimated as shown in Figures 29A-29B and described in the Examples.

Heterodimer Purity (%)	CH3 Tm (°C)	Mutations (Chain A)	Mutations (Chain B)
>98	70.5	F405A_Y407V	T366L_T394W
>98	73.5	F405A_Y407V	T366L_K392M_T394W
>98	76.5	T350V_F405A_Y407V	T350V_T366L_K392M_T394W
>98	78.7	L351Y_F405A_Y407V	T366L_K392M_T394W
>98	79.5	T350V_L351Y_F405A_Y407V	T350V_T366L_K392M_T394W
>98	81.8	T350V_L351Y_F405A_Y407V	T350V_T366L_K392L_T394W
>98	81	T350V_L351Y_S400R_F405A_Y407V	T350V_T366L_K392M_T394W
>98	79.5	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392M_T394W
>98	77.5	T350V_L351Y_S400E_F405V_Y407V	T350V_T366L_N390R_K392M_T394W
>98	77	T350V_L351Y_S400E_F405T_Y407V	T350V_T366L_N390R_K392M_T394W
>98	78	T350V_L351Y_S400E_F405S_Y407V	T350V_T366L_N390R_K392M_T394W
>98	76.5	T350V_S400E_F405A_Y407V	T350V_T366L_N390R_K392M_T394W
>98	76.5	T350V_L351Y_S400E_F405A_Y407V	T350V_L351Y_T366L_N390R_K392M_T394W
>98	81.5	Q347R_T350V_L351Y_S400E_F405A_Y407V	T350V_K360E_T366L_N390R_K392M_T394W
>98	80.5	T350V_L351Y_S400R_F405A_Y407V	T350V_T366L_N390D_K392M_T394W
>98	79.5	T350V_L351Y_S400R_F405A_Y407V	T350V_T366L_N390E_K392M_T394W
>98	81.5	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392L_T394W
>98	76.5	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392F_T394W

>98	73.5	Y349C_F405A_Y407V	S354C_T366L_T394W
>98	78	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W
>98	82	Y349C_T350V_L351Y_S400E_F405A_Y407V	T350V_S354C_T366L_N390R_K392M_T394W
>98	82	Y349C_T350V_S400E_F405A_Y407V	T350V_S354C_T366L_N390R_K392M_T394W
>98	76	L351Y_F405A_Y407V	T366L_K392M_T394W
>98	81.5	Y349C_T350V_F405A_Y407V	T350V_S354C_T366L_K392M_T394W

Table 1.3: CH3 domain amino acid modifications for the generation of Fc variant heterodimers. The Kd in the table above were determined as described in the Examples and Figure 35

CD16a(F158) Kd [M]	CD32b(Y163) Kd [M]	Mutations(Chain-A)	Mutations(Chain-B)
4.4E-07	1.7E-06	Herceptin WT	
4.5E-07	9.0E-07	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392M_T394W
3.7E-07	7.0E-07	T350V_L351Y_S400E_F405V_Y407V	T350V_T366L_N390R_K392M_T394W
3.9E-07	6.7E-07	T350V_L351Y_S400E_F405T_Y407V	T350V_T366L_N390R_K392M_T394W
4.2E-07	8.3E-07	T350V_L351Y_S400E_F405S_Y407V	T350V_T366L_N390R_K392M_T394W
4.5E-07	1.0E-06	T350V_S400E_F405A_Y407V	T350V_T366L_N390R_K392M_T394W
3.7E-07	7.1E-07	T350V_L351Y_S400E_F405A_Y407V	T350V_L351Y_T366L_N390R_K392M_T394W
4.2E-07	9.2E-07	Q347R_T350V_L351Y_S400E_F405A_Y407V	T350V_K360E_T366L_N390R_K392M_T394W
4.3E-07	8.9E-07	T350V_L351Y_S400R_F405A_Y407V	T350V_T366L_K392M_T394W
4.3E-07	9.4E-07	T350V_L351Y_S400R_F405A_Y407V	T350V_T366L_N390D_K392M_T394W
4.2E-07	8.9E-07	T350V_L351Y_S400R_F405A_Y407V	T350V_T366L_N390E_K392M_T394W
4.4E-07	9.1E-07	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392L_T394W
3.6E-07	7.1E-07	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392F_T394W
4.6E-07	1.1E-06	F405A_Y407V	T366L_K392M_T394W
4.3E-07	1.0E-06	T350V_F405A_Y407V	T350V_T366L_K392M_T394W
4.8E-07	1.1E-06	F405A_Y407V	T366L_T394W
5.1E-07	1.2E-06	D399C_F405A_Y407V	T366L_K392C_T394W
5.8E-07	1.2E-06	Y349C_F405A_Y407V	S354C_T366L_T394W
6.3E-07	1.3E-06	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W
4.2E-07	9.5E-07	Y349C_T350V_L351Y_S400E_F405A_Y407V	T350V_S354C_T366L_N390R_K392M_T394W
4.4E-07	1.1E-06	Y349C_T350V_S400E_F405A_Y407V	T350V_S354C_T366L_N390R_K392M_T394W
4.2E-07	1.2E-06	L351Y_F405A_Y407V	T366L_K392M_T394W
4.2E-07	1.3E-06	L351Y_F405A_Y407V	T366L_K392M_T394W
4.6E-07	1.2E-06	T350V_L351Y_F405A_Y407V	T350V_T366L_K392M_T394W
4.6E-07	1.3E-06	Y349C_T350V_F405A_Y407V	T350V_S354C_T366L_K392M_T394W
4.2E-07	1.1E-06	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392M_T394W
3.6E-07	9.9E-07	T350V_L351Y_F405A_Y407V	T350V_T366L_K392L_T394W

Fc region definition

The Fc region as defined herein comprises a CH3 domain or fragment thereof, and may additionally comprise one or more addition constant region domains, or fragments thereof, including hinge, CH1, or CH2. It will be understood that the numbering of the Fc amino acid residues is that of the EU index as in Kabat et al., 1991, NIH Publication 91-3242, National Technical Information Service, Springfield, Va. The "EU index as set forth in Kabat" refers to the EU index numbering of the human IgG1 Kabat antibody. For convenience, **Table B** provides the amino acids numbered according to the EU index as set forth in Kabat of the CH2 and CH3 domain from human IgG1.

Table B

CH2 Domain						CH3 Domain					
EU No.	Amino Acid	EU No.	Amino Acid	EU No.	Amino Acid	EU No.	Amino Acid	EU No.	Amino Acid	EU No.	Amino Acid
231	A	271	P	311	Q	341	G	381	W	421	N
232	P	272	E	312	D	342	Q	382	E	422	V
233	E	273	V	313	W	343	P	383	S	423	F
234	L	274	K	314	L	344	R	384	N	424	S
235	L	275	F	315	N	345	E	385	G	425	C
236	G	276	N	316	G	346	P	386	Q	426	S
237	G	277	W	317	K	347	Q	387	P	427	V
238	P	278	Y	318	E	348	V	388	E	428	M
239	S	279	V	319	Y	349	Y	389	N	429	H
240	V	280	D	320	K	350	T	390	N	430	E
241	F	281	G	321	C	351	L	391	Y	431	A
242	L	282	V	322	K	352	P	392	K	432	L
243	F	283	E	323	V	353	P	393	T	433	H
244	P	284	V	324	S	354	S	394	T	434	N
245	P	285	H	325	N	355	R	395	P	435	H
246	K	286	N	326	K	356	D	396	P	436	Y
247	P	287	A	327	A	357	E	397	V	437	T
248	K	288	K	328	L	358	L	398	L	438	Q
249	D	289	T	329	P	359	T	399	D	439	K
250	T	290	K	330	A	360	K	400	S	440	S
251	L	291	P	331	P	361	N	401	D	441	L
252	M	292	R	332	I	362	Q	402	G	442	S
253	I	293	E	333	E	363	V	403	S	443	L
254	S	294	E	334	K	364	S	404	F	444	S
255	R	295	Q	335	T	365	L	405	F	445	P
256	T	296	Y	336	I	366	T	406	L	446	G
257	P	297	N	337	S	367	C	407	Y	447	K

258	E	298	S	338	K		368	L	408	S		
259	V	299	T	339	A		369	V	409	K		
260	T	300	Y	340	K		370	K	410	L		
261	C	301	R				371	G	411	T		
262	V	302	V				372	F	412	V		
263	V	303	V				373	Y	413	D		
264	V	304	S				374	P	414	K		
265	D	305	V				375	S	415	S		
266	V	306	L				376	D	416	R		
267	S	307	T				377	I	417	W		
268	H	308	V				378	A	418	Q		
269	E	309	L				379	V	419	Q		
270	D	310	H				380	E	420	G		

In certain embodiments, the heteromultimers comprise an Fc region that comprises a CH2 domain. In some embodiments, the CH2 domain is a variant CH2 domain. In some
5 embodiments, the variant CH2 domains comprise asymmetric amino acid substitutions in the first and/or second polypeptide chain. In some embodiments, the heteromultimer comprises asymmetric amino acid substitutions in the CH2 domain such that one chain of said heteromultimer selectively binds an Fc receptor.

FcγR selectivity

10 In one aspect, this application describes a molecular design for achieving exquisite FcγR selectivity profiles via the design of an asymmetric scaffold built on a heterodimeric Fc. This scaffold allows for asymmetric mutations in the CH2 domain to achieve a variety of novel selectivity profiles. Further, the scaffold has inherent features for the engineering of multifunctional (bi, tri, tetra or penta functional) therapeutic molecules. In certain
15 embodiments, the asymmetric scaffold is optimized for pH dependent binding properties to the neonatal Fc receptor (FcRn) to enable better recycling of the molecule and enhance its half life and related pharmacokinetic properties.

The asymmetric scaffold can be optimized for binding to the functionally relevant FcγRI
20 receptor allotypes. FcγRI is a prominent marker on macrophages that are involved in chronic inflammatory disorders such as Rheumatoid Arthritis, Atopic Dermatitis, Psoriasis and a number of pulmonary diseases.

The asymmetric scaffold can be optimized for protein A binding. Protein A binding is often employed for separation and purification of antibody molecules. Mutations can be introduced in the asymmetric scaffold to avoid aggregation of the therapeutic during storage.

5

Therefore, it is specifically contemplated that the heteromultimers comprising heavy chain variable region and Fc variants of the invention may contain inter alia one or more additional amino acid residue substitutions, mutations and/or modifications which result in an antibody with preferred characteristics including but not limited to: increased serum half
10 life, increase binding affinity, reduced immunogenicity, increased production, enhanced or reduced ADCC or CDC activity, altered glycosylation and/or disulfide bonds and modified binding specificity.

***In vivo* half life**

15 It is contemplated that the heteromultimers described herein may have other altered characteristics including increased in vivo half-lives (e.g., serum half-lives) in a mammal; in particular a human, increased stability in vivo (e.g., serum half-lives) and/or in vitro (e.g., shelf-life) and/or increased melting temperature (T_m), relative to a comparable molecule. In one embodiment, a heteromultimer of the invention has an in vivo half-life of greater
20 then 15 days, greater than 20 days, greater than 25 days, greater than 30 days, greater than 35 days, greater than 40 days, greater than 45 days, greater than 2 months, greater than 3 months, greater than 4 months, or greater than 5 months. In another embodiment, a heteromultimer of the invention has an in vitro half-live (e.g, liquid or powder formulation) of greater then 15 days, greater than 30 days, greater than 2 months, greater than 3
25 months, greater than 6 months, or greater than 12 months, or greater than 24 months, or greater than 36 months, or greater than 60 months.

It will also be appreciated by one skilled in the art that the heteromultimers of the invention may have altered immunogenicity when administered to a subject. Accordingly, it is
30 contemplated that the variant CH3 domain, which minimize the immunogenicity of the Fc variant are generally more desirable for therapeutic applications.

Altered effector function

The heteromultimers of the present invention may be combined with other modifications, including but not limited to modifications that alter effector function. The invention encompasses combining a heteromultimer of the invention with other Fc modifications to provide additive, synergistic, or novel properties in antibodies or Fc fusion proteins. Such modifications may be in the hinge, or CH2, (or CH3 provided it does not negatively alter the stability and purity properties of the present variant CH3 domains) domains or a combination thereof. It is contemplated that the heteromultimers of the invention enhance the property of the modification with which they are combined.

10 FcγR binding

As part of the characterization of the heteromultimers they were tested for their binding affinity to FcγRIIIA (CD16a) and FcγRIIB(CD32b) reported as a ratio in comparison to wild-type IgG1. (See, Example 4 and Table 5) In this instance it was possible to evaluate the impact of the CH3 domain mutations on binding to these activating and inhibitory Fc receptors. In one embodiment, provided herein are isolated heteromultimers comprising a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) greater than 70 °C, wherein the heterodimer binding to CD16a is about the same as compared to wild-type homodimer. In certain embodiments the heterodimer binding to CD16a is increased as compared to wild-type homodimer. In an alternative embodiment, the heterodimer binding to CD16a is reduced as compared to wild-type homodimer.

In certain embodiments, provided herein are isolated heteromultimers comprising at least one single domain antigen-binding construct and a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) greater than 70 °C, wherein the heterodimer binding to CD32b is about the same as compared to wild-type homodimer, and wherein the heteromultimer is devoid of immunoglobulin light chains and immunoglobulin CH1 and optionally devoid of immunoglobulin CH2 regions. In certain embodiments the heterodimer binding to CD32b is increased as compared to wild-type homodimer. In an alternative

embodiment, the heterodimer binding to CD32b is reduced as compared to wild-type homodimer.

One of skill in the art will understand that instead of reporting the K_D of binding CD16a and CD32b as a ratio Fc variant to wild-type homodimer, the K_D could be reported as a ratio of Fc variant binding to CD16a to Fc variant binding to CD32b (data not shown). This ratio would provide an indication of the variant CH3 domain mutation on ADCC, either unchanged, increased to decreased compared to wild-type, described below in more detail.

The affinities and binding properties of the heteromultimers of the invention for an FcγR are initially determined using in vitro assays (biochemical or immunological based assays) known in the art for determining Fc-FcγR interactions, i.e., specific binding of an Fc region to an FcγR including but not limited to ELISA assay, surface plasmon resonance assay, immunoprecipitation assays (See section entitled "Characterization and Functional Assays" infra) and other methods such as indirect binding assays, competitive inhibition assays, fluorescence resonance energy transfer (FRET), gel electrophoresis and chromatography (e.g., gel filtration). These and other methods may utilize a label on one or more of the components being examined and/or employ a variety of detection methods including but not limited to chromogenic, fluorescent, luminescent, or isotopic labels. A detailed description of binding affinities and kinetics can be found in Paul, W. E., ed., Fundamental Immunology, 4th Ed., Lippincott-Raven, Philadelphia (1999), which focuses on antibody-immunogen interactions.

It is contemplated that the binding properties of the molecules of the invention are also characterized by in vitro functional assays for determining one or more FcγR mediator effector cell functions (See section entitled "Characterization and Functional Assays" infra). In certain embodiments, the molecules of the invention have similar binding properties in *in vivo* models (such as those described and disclosed herein) as those in in vitro based assays. However, the present invention does not exclude molecules of the invention that do not exhibit the desired phenotype in *in vitro* based assays but do exhibit the desired phenotype *in vivo*.

The invention encompasses heteromultimers comprising Fc variants that bind FcγRIIIA (CD16a) with increased affinity, relative to a comparable molecule. In a specific embodiment, the Fc variants of the invention bind FcγRIIIA with increased affinity and bind FcγRIIB (CD32b) with a binding affinity that is either unchanged or reduced, relative to a comparable molecule. In yet another embodiment, the Fc variants of the invention have a ratio of FcγRIIIA/FcγRIIB equilibrium dissociation constants (K_D) that is decreased relative to a comparable molecule.

Also encompassed by the present invention are heteromultimers comprising Fc variants that bind FcγRIIIA (CD16a) with decreased affinity, relative to a comparable molecule. In a specific embodiment, the Fc variants of the invention bind FcγRIIIA with decreased affinity, relative to a comparable molecule and bind FcγRIIB with a binding affinity that is unchanged or increased, relative to a comparable molecule.

In one embodiment, the heteromultimers comprise Fc variants bind with increased affinity to FcγRIIIA. In a specific embodiment, said Fc variants have affinity for FcγRIIIA that is at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 30 fold, or at least 40 fold, or at least 50 fold, or at least 60 fold, or at least 70 fold, or at least 80 fold, or at least 90 fold, or at least 100 fold, or at least 200 fold greater than that of a comparable molecule. In other embodiments, the Fc variants have an affinity for FcγRIIIA that is increased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

In another embodiment, the Fc variant has an equilibrium dissociation constant (K_D) for an Fc ligand (e.g., FcγR, C1q) that is decreased between about 2 fold and 10 fold, or between about 5 fold and 50 fold, or between about 25 fold and 250 fold, or between about 100 fold and 500 fold, or between about 250 fold and 1000 fold relative to a comparable molecule.

In another embodiment, said Fc variants have an equilibrium dissociation constant (K_D) for FcγRIIIA that is reduced by at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 30 fold, or at least 40 fold, or at least

50 fold, or at least 60 fold, or at least 70 fold, or at least 80 fold, or at least 90 fold, or at least 100 fold, or at least 200 fold, or at least 400 fold, or at least 600 fold, relative to a comparable molecule. In another embodiment, the Fc variants have an equilibrium dissociation constant (K_D) for FcγRIIIA that is reduced by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

10 In one embodiment, the Fc variant binds to FcγRIIB with an affinity that is unchanged or reduced. In a specific embodiment, said Fc variants have affinity for FcγRIIB that is unchanged or reduced by at least 1 fold, or by at least 3 fold, or by at least 5 fold, or by at least 10 fold, or by at least 20 fold, or by at least 50 fold, or by at least 100 fold, relative to a comparable molecule. In other embodiments, the Fc variants have an affinity for FcγRIIB that is unchanged or reduced by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

20 In another embodiment, the Fc variants have an equilibrium dissociation constant (K_D) for FcγRIIB that is unchanged or increased by at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 30 fold, or at least 40 fold, or at least 50 fold, or at least 60 fold, or at least 70 fold, or at least 80 fold, or at least 90 fold, or at least 100 fold, or at least 200 fold relative to a comparable molecule. In another specific embodiment, the Fc variants have an equilibrium dissociation constant (K_D) for FcγRIIB that is unchanged or increased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

30 In still another embodiment, the Fc variants bind FcγRIIIA with increased affinity, relative to a comparable molecule and bind FcγRIIB with an affinity that is unchanged or reduced, relative to a comparable molecule. In a specific embodiment, the Fc variants have affinity for FcγRIIIA that is increased by at least 1 fold, or by at least 3 fold, or by at least 5 fold, or by at least 10 fold, or by at least 20 fold, or by at least 50 fold, or by at least 100 fold,

relative to a comparable molecule. In another specific embodiment, the Fc variants have affinity for FcγRIIB that is either unchanged or is reduced by at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 50 fold, or at least 100 fold, relative to a comparable molecule. In other embodiments, the Fc variants have an affinity for FcγRIIIA that is increased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule and the Fc variants have an affinity for FcγRIIB that is either unchanged or is increased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

In yet another embodiment, the Fc variants have a ratio of FcγRIIIA/FcγRIIB equilibrium dissociation constants (K_D) that is decreased relative to a comparable molecule. In a specific embodiment, the Fc variants have a ratio of FcγRIIIA/FcγRIIB equilibrium dissociation constants (K_D) that is decreased by at least 1 fold, or by at least 3 fold, or by at least 5 fold, or by at least 10 fold, or by at least 20 fold, or by at least 50 fold, or by at least 100 fold, relative to a comparable molecule. In another specific embodiment, the Fc variants have a ratio of FcγRIIIA/FcγRIIB equilibrium dissociation constants (K_D) that is decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

In another embodiment, the Fc variants bind FcγRIIIA with a decreased affinity, relative to a comparable molecule. In a specific embodiment, said Fc variants have affinity for FcγRIIIA that is reduced by at least 1 fold, or by at least 3 fold, or by at least 5 fold, or by at least 10 fold, or by at least 20 fold, or by at least 50 fold, or by at least 100 fold, relative to a comparable molecule. In other embodiments, the Fc variants have an affinity for FcγRIIIA that is decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

In still another embodiment, the Fc variants bind FcγRIIIA with decreased affinity and bind

FcγRIIB with an affinity that is either unchanged or increased, relative to a comparable molecule. In a specific embodiment, the Fc variants have affinity for FcγRIIIA that is reduced by at least 1 fold, or by at least 3 fold, or by at least 5 fold, or by at least 10 fold, or by at least 20 fold, or by at least 50 fold, or by at least 100 fold relative to a comparable molecule. In another specific embodiment, the Fc variants have affinity for FcγRIIB that is at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 50 fold, or at least 100 fold, greater than that of a comparable molecule. In other embodiments, the Fc variants have an affinity for FcγRIIIA that is decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule and the Fc variants have an affinity for FcγRIIB that is increased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule.

In still another embodiment, the Fc variants have an equilibrium dissociation constant (K_D) for FcγRIIIA that are increased by at least 1 fold, or by at least 3 fold, or by at least 5 fold or by at least 10 or by at least 20 fold, or by at least 50 fold when compared to that of a comparable molecule. In a specific embodiment, said Fc variants have equilibrium dissociation constant (K_D) for FcγRIIB that are decreased at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 50 fold or at least 100 fold, relative to a comparable molecule.

CH2 variations for fcγR selectivity

The Fc-FcγR protein-protein interaction in this complex indicates that the two chains in the heteromultimer interact with two distinct sites on the FcγR molecule. Although there is symmetry in the two heavy chains in the natural Fc molecules, the local FcγR environment around residues on one chain is different from the FcγR residues surrounding the same residue position on the opposite Fc chain. The two symmetry related positions interact with different selection of FcγR residues.

Given the asymmetry in the association of Fc to Fc γ R, concurrent mutations in chain A and B of the Fc molecule do not impact the interactions with Fc γ R in a symmetric manner.

When introducing mutations to optimize interactions on one chain of the Fc with its local

5 Fc γ R environment, in a homodimeric Fc structure, the corresponding mutation in the second chain may be favorable, unfavorable or non-contributing to the required Fc γ R binding and selectivity profile.

Using a structure and computation guided approach, asymmetric mutations are

10 engineered in the two chains of the Fc to overcome these limitations of traditional Fc engineering strategies, which introduce the same mutations on both the chains of Fc. One can achieve better binding selectivity between the receptors if the two chains of Fc are optimized independently for enhanced binding to their corresponding face of the receptor molecule.

15

For instance, mutations at a particular position on one chain of the Fc can be designed to enhance selectivity to a particular residue, a positive design effort, while the same residue position can be mutated to unfavorably interact with its local environment in an alternate Fc γ receptor type, a negative design effort, hence achieving better selectivity between the

20 two receptors. In certain embodiments, is provided a method for designing asymmetric amino acid modifications in the CH2 domain that selectively bind one Fc gamma receptor as compared to a different Fc gamma receptor (e.g., selectively bind Fc γ RIIIa instead of Fc γ RIIb). In other certain embodiments, is provided a method for the design of asymmetric amino acid modifications in the CH2 domain of a variant Fc

25 heterodimer comprising amino acid modifications in the CH3 domain to promote heterodimer formation. In another embodiment, is provided a method to design selectivity for the different Fc gamma receptors based on a variant Fc heterodimer comprising asymmetric amino acid modifications in the CH2 domain. In yet another embodiment, is provided a method for designing asymmetric amino acid modifications that bias binding of

30 the Fc gamma receptors to one face of the Fc molecule. In other certain embodiments, is provided a method for designing polarity drivers that bias the Fc gamma receptors to

interact with only one face of the variant Fc heterodimer comprising asymmetric amino acid modifications in the CH2 domain.

The asymmetric design of mutations in the CH2 domain can be tailored to recognize the FcγR on one face of the Fc molecule. This constitutes the productive face of the asymmetric Fc scaffold while the opposite face presents wild type like interaction propensity without the designed selectivity profile and can be considered a non-productive face. A negative design strategy can be employed to introduce mutations on the non-productive face to block FcγR interactions to this side of the asymmetric Fc scaffold, thereby forcing the desired interaction tendencies to the Fcγ receptors.

Table E: Potentially Interesting Selectivity Profiles of Fc for different Fcγ Receptors

Variant Selectivity	Receptor Binding		
	FcγRIIIa F/V	FcγRIIa H/R	FcγRIIb F/Y
	↑ / -	×	×
	×	↑ / -	×
	×	×	↑ / -
	↑ / -	↑ / -	×
	↑ / -	×	↑ / -
	×	↑ / -	↑ / -

(↑/-) indicates a variant which exhibits an increased or wild type like binding to the particular receptor type or one of its allotype. (×) indicates no noticeable binding to the receptor or a subset allotype.

Certain embodiments provided herein relate to fusion polypeptides comprising a binding domain fused to an Fc region, wherein the Fc region comprising a variant CH3 domain, comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) greater than 70 °C. It is specifically contemplated that molecules comprising a heterodimer comprising a

variant CH3 domain may be generated by methods well known to one skilled in the art. Briefly, such methods include but are not limited to, combining a variable region or binding domain with the desired specificity (e.g., a variable region isolated from a phage display or expression library or derived from a human or non-human antibody or a binding domain of a receptor) with a variant Fc heterodimers. Alternatively, one skilled in the art may generate a variant Fc heterodimer by modifying the CH3 domain in the Fc region of a molecule comprising an Fc region (e.g., an antibody).

“Antibody-dependent cell-mediated cytotoxicity” or “ADCC” refers to a form of cytotoxicity in which secreted antibody bound onto Fc receptors (FcRs) present on certain cytotoxic cells (e.g., Natural Killer (NK) cells, neutrophils, and macrophages) enables these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. Specific high-affinity IgG antibodies directed to the surface of target cells “arm” the cytotoxic cells and are absolutely required for such killing. Lysis of the target cell is extracellular, requires direct cell-to-cell contact, and does not involve complement.

The ability of any particular antibody to mediate lysis of the target cell by ADCC can be assayed. To assess ADCC activity an antibody of interest is added to target cells in combination with immune effector cells, which may be activated by the antigen antibody complexes resulting in cytolysis of the target cell. Cytolysis is generally detected by the release of label (e.g. radioactive substrates, fluorescent dyes or natural intracellular proteins) from the lysed cells. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Specific examples of in vitro ADCC assays are described in Wisecarver et al., 1985, 79:277; Bruggemann et al., 1987, J Exp Med 166:1351; Wilkinson et al., 2001, J Immunol Methods 258:183; Patel et al., 1995 J Immunol Methods 184:29 and herein (see section entitled “Characterization and Functional Assays” infra). Alternatively, or additionally, ADCC activity of the antibody of interest may be assessed in vivo, e.g., in an animal model such as that disclosed in Clynes et al., 1998, PNAS USA 95:652.

The present invention further provides heteromultimers comprising Fc variants with enhanced CDC function. In one embodiment, the Fc variants have increased CDC activity.

In one embodiment, the Fc variants have CDC activity that is at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 10 fold, or at least 50 fold, or at least 100 fold greater than that of a comparable molecule. In another embodiment, the Fc variants bind C1q with an affinity that is at least 2 fold, or at least 3 fold, or at least 5 fold, or at least 7 fold, or at least 10 fold, or at least 20 fold, or at least 50 fold, or at least 100 fold, greater than that of a comparable molecule. In yet another embodiment, the Fc variants have CDC activity that is increased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule. In a specific embodiment, the Fc variants of the invention bind C1q with increased affinity; have enhanced CDC activity and specifically bind to at least one antigen.

The present invention also provides heteromultimers comprising Fc variants with reduced CDC function. In one embodiment, the Fc variants have reduced CDC activity. In one embodiment, the Fc variants have CDC activity that is at least 2 fold, or at least 3 fold, or at least 5 fold or at least 10 fold or at least 50 fold or at least 100 fold less than that of a comparable molecule. In another embodiment, an Fc variant binds C1q with an affinity that is reduced by at least 1 fold, or by at least 3 fold, or by at least 5 fold, or by at least 10 fold, or by at least 20 fold, or by at least 50 fold, or by at least 100 fold, relative to a comparable molecule. In another embodiment, the Fc variants have CDC activity that is decreased by at least 10%, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or at least 150%, or at least 200%, relative to a comparable molecule. In a specific embodiment, Fc variants bind to C1q with decreased affinity have reduced CDC activity and specifically bind to at least one antigen.

In some embodiments, the Fc variants comprise one or more engineered glycoforms, i.e., a carbohydrate composition that is covalently attached to a molecule comprising an Fc region. Engineered glycoforms may be useful for a variety of purposes, including but not limited to enhancing or reducing effector function. Engineered glycoforms may be generated by any method known to one skilled in the art, for example by using engineered or variant expression strains, by co-expression with one or more enzymes, for example

β (1,4)-N-acetylglucosaminyltransferase III (GnT111), by expressing a molecule comprising an Fc region in various organisms or cell lines from various organisms, or by modifying carbohydrate(s) after the molecule comprising Fc region has been expressed. Methods for generating engineered glycoforms are known in the art, and include but are not limited to those described in Umana et al, 1999, Nat. Biotechnol 17:176-180; Davies et al., 20017 Biotechnol Bioeng 74:288-294; Shields et al, 2002, J Biol Chem 277:26733-26740; Shinkawa et al., 2003, J Biol Chem 278:3466-3473) U.S. Pat. No. 6,602,684; U.S. Ser. No. 10/277,370; U.S. Ser. No. 10/113,929; PCT WO 00/61739A1; PCT WO 01/292246A1; PCT WO 02/311140A1; PCT WO 02/30954A1; Potillegent™ technology (Biowa, Inc. Princeton, N.J.); GlycoMAb™ glycosylation engineering technology (GLYCART biotechnology AG, Zurich, Switzerland). See, e.g., WO 00061739; EA01229125; US 20030115614; Okazaki et al., 2004, JMB, 336: 1239-49.

It is contemplated that Fc variants include antibodies comprising a variable region and a heterodimer Fc region, wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability, wherein the variant CH3 domain has a melting temperature (T_m) greater than 70 °C. The Fc variants which are antibodies may be produced “de novo” by combining a variable domain, or fragment thereof, that specifically binds at least one antigen with a heterodimer Fc region comprising a variant CH3 domain. Alternatively, heterodimer Fc variants may be produced by modifying the CH3 domain of an Fc region containing antibody that binds an antigen.

Heteromultimers of the invention may be monospecific, bi-specific, trispecific or have greater multispecificity. Multispecific antibodies may specifically bind to different epitopes of desired target molecule or may specifically bind to both the target molecule as well as a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., International Publication Nos. WO 94/04690; WO 93/17715; WO 92/08802; WO 91/00360; and WO 92/05793; Tutt, et al., 1991, J. Immunol. 147:60-69; U.S. Pat.Nos.; 4,474,893; 4,714,681; 4,925,648; 5,573,920 and 5,601,819 and Kostelny et al., 1992, J. Immunol.148:1547).

Various embodiments of multifunctional targeting molecules can be designed on the basis

of this asymmetric scaffold as shown in Figure 20.

Bi-specific or multi-specific antibodies

Multispecific heteromultimers are based on antibodies that have binding specificities for at least two different antigens. While such molecules normally will only bind two antigens (i.e. bi-specific antibodies, BsAbs), antibodies with additional specificities such as trispecific antibodies are encompassed by the instant invention. Examples of BsAbs include without limitation those with one arm directed against a tumor cell antigen and the other arm directed against a cytotoxic molecule, or both arms are directed against two different tumor cell antigens, or both arms are directed against two different soluble ligands, or one arm is directed against a soluble ligand and the other arm is directed against a cell surface receptor, or both arms are directed against two different cell surface receptors. Methods for making bi-specific antibodies are known in the art.

According to a different approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences. The fusion may be with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. DNAs encoding the immunoglobulin heavy chain fusions, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of the three polypeptide chains used in the construction provide the optimum yields. See, Example 1 and Table 2. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when, the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no particular significance.

Bi-specific antibodies include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Pat. No. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known

in the art, and are disclosed in U.S. Pat. No. 4,676,980, along with a number of cross-linking techniques.

Antibodies with more than two valencies incorporating variant CH3 domains and resulting Fc heterodimers of the invention are contemplated. For example, trispecific antibodies can be prepared. See, e.g., Tutt et al. J. Immunol. 147: 60 (1991).

Serum half-life

Antibodies of the present invention also encompass those that have half-lives (e.g., serum half-lives) in a mammal, (e.g., a human), of greater than 15 days, greater than 20 days, greater than 25 days, greater than 30 days, greater than 35 days, greater than 40 days, greater than 45 days, greater than 2 months, greater than 3 months, greater than 4 months, or greater than 5 months. The increased half-lives of the antibodies of the present invention in a mammal, (e.g., a human), results in a higher serum titer of said antibodies or antibody fragments in the mammal, and thus, reduces the frequency of the administration of said antibodies or antibody fragments and/or reduces the concentration of said antibodies or antibody fragments to be administered. Antibodies having increased in vitro half-lives can be generated by techniques known to those of skill in the art. For example, antibodies with increased in vivo half-lives can be generated by modifying (e.g., substituting, deleting or adding) amino acid residues identified as involved in the interaction between the Fc domain and the FcRn receptor (see, e.g., International Publication Nos. WO 97/34631; WO 04/029207; U.S. Pat. No. 6,737,056 and U.S. Patent Publication No. 2003/0190311).

Therapeutic antibodies

In a specific embodiment the variant Fc heterodimer comprising at least an immunoglobulin heavy chain variable region and a variant CH3 domain is a multi-specific antibody, wherein the heteromultimer is devoid of immunoglobulin light chains and immunoglobulin CH1 region and optionally devoid of immunoglobulin CH2 region (referred to herein as an antibody of the invention), the antibody of the invention specifically binds an antigen of interest. In particular the antibody of the invention is a bi-specific antibody. In one embodiment, an antibody of the invention specifically binds a polypeptide antigen. In another embodiment, an antibody of the invention specifically binds a nonpolypeptide

antigen. In yet another embodiment, administration of an antibody of the invention to a mammal suffering from a disease or disorder can result in a therapeutic benefit in that mammal.

- 5 Also provided are antibodies of the invention that specifically bind cancer antigens including, but not limited to, ALK receptor (pleiotrophin receptor), pleiotrophin, KS 1/4 pan-carcinoma antigen; ovarian carcinoma antigen (CA125); prostatic acid phosphate; prostate specific antigen (PSA); melanoma-associated antigen p97; melanoma antigen gp75; high molecular weight melanoma antigen (HMW-MAA); prostate specific membrane antigen;
- 10 carcinoembryonic antigen (CEA); polymorphic epithelial mucin antigen; human milk fat globule antigen; colorectal tumor-associated antigens such as: CEA, TAG-72, CO17-1A, GICA 19-9, CTA-1 and LEA; Burkitt's lymphoma antigen-38.13; CD19; human B-lymphoma antigen-CD20; CD33; melanoma specific antigens such as ganglioside GD2, ganglioside GD3, ganglioside GM2 and ganglioside GM3; tumor-specific transplantation
- 15 type cell-surface antigen (TSTA); virally-induced tumor antigens including T-antigen, DNA tumor viruses and Envelope antigens of RNA tumor viruses; oncofetal antigen-alpha-fetoprotein such as CEA of colon, 5T4 oncofetal trophoblast glycoprotein and bladder tumor oncofetal antigen; differentiation antigen such as human lung carcinoma antigens L6 and L20; antigens of fibrosarcoma; human leukemia T cell antigen-Gp37; neoglycoprotein;
- 20 sphingolipids; breast cancer antigens such as EGFR (Epidermal growth factor receptor); NY-BR-16; NY-BR-16 and HER2 antigen (p185HER2); polymorphic epithelial mucin (PEM); malignant human lymphocyte antigen-APO-1; differentiation antigen such as I antigen found in fetal erythrocytes; primary endoderm I antigen found in adult erythrocytes; preimplantation embryos; I(Ma) found in gastric adenocarcinomas; M18, M39 found in
- 25 breast epithelium; SSEA-1 found in myeloid cells; VEP8; VEP9; Myl; Va4-D5; D₁56-22 found in colorectal cancer; TRA-1-85 (blood group H); SCP-1 found in testis and ovarian cancer; C14 found in colonic adenocarcinoma; F3 found in lung adenocarcinoma; AH6 found in gastric cancer; Y hapten; Ley found in embryonal carcinoma cells; TL5 (blood group A); EGF receptor found in A431 cells; E₁ series (blood group B) found in pancreatic
- 30 cancer; FC10.2 found in embryonal carcinoma cells; gastric adenocarcinoma antigen; CO-514 (blood group Lea) found in Adenocarcinoma; NS-10 found in adenocarcinomas; CO-43 (blood group Leb); G49 found in EGF receptor of A431 cells; MH2 (blood group AL_{Leb}/Ley) found in colonic adenocarcinoma; 19.9 found in colon cancer; gastric cancer

mucins; T₅A₇ found in myeloid cells; R₂₄ found in melanoma; 4.2, G_{D3}, D1.1, OFA-1, G_{M2}, OFA-2, G_{D2}, and M1:22:25:8 found in embryonal carcinoma cells and SSEA-3 and SSEA-4 found in 4 to 8-cell stage embryos; Cutaneous Tcell Lymphoma antigen; MART-1 antigen; Sialy Tn (STn) antigen; Colon cancer antigen NY-CO-45; Lung cancer antigen NY-LU-12
 5 valiant A; Adenocarcinoma antigen ART1; Paraneoplastic associated brain-testis-cancer antigen (onconeural antigen MA2; paraneoplastic neuronal antigen); Neuro-oncological ventral antigen 2 (NOVA2); Hepatocellular carcinoma antigen gene 520; TUMOR-ASSOCIATED ANTIGEN CO-029; Tumor-associated antigens MAGE-C1 (cancer/testis antigen CT7), MAGE-B1 (MAGE-XP antigen), MAGE-B2 (DAM6), MAGE-2, MAGE-4-a,
 10 MAGE-4-b and MAGE-X2; Cancer-Testis Antigen (NY-EOS-1) and fragments of any of the above-listed polypeptides.

In certain embodiments, the heteromultimer described herein is competitive to at least one domain of at least one therapeutic antibody. In some embodiments, the therapeutic
 15 antibody binds a cancer target antigen. In an embodiment, the therapeutic antibody may be selected from the group consisting of abagovomab, adalimumab, alemtuzumab, aurograb, bapineuzumab, basiliximab, belimumab, bevacizumab, briakinumab, canakinumab, catumaxomab, certolizumab pegol, cetuximab, daclizumab, denosumab, efalizumab, galiximab, gemtuzumab ozogamicin, golimumab, ibritumomab tiuxetan,
 20 infliximab, ipilimumab, lumiliximab, mepolizumab, motavizumab, muromonab, mycograb, natalizumab, nimotuzumab, ocrelizumab, ofatumumab, omalizumab, palivizumab, panitumumab, pertuzumab, ranibizumab, reslizumab, rituximab, teplizumab, tocilizumab/atlizumab, tositumomab, trastuzumab, ProxiniumTM, RencarexTM, ustekinumab, zalutumumab, and any other antibodies.

25 **Antibody derivatives**

Antibodies of the invention include derivatives that are modified (i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment). For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation,
 30 derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to, specific chemical cleavage,

acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

Antibodies or fragments thereof with increased in vivo half-lives can be generated by
5 attaching polymer molecules such as high molecular weight polyethyleneglycol (PEG) to the antibodies or antibody fragments. PEG can be attached to the antibodies or antibody fragments with or without a multifunctional linker either through site-specific conjugation of the PEG to the N- or C-terminus of said antibodies or antibody fragments or via epsilon-amino groups present on lysine residues. Linear or branched polymer derivatization that
10 results in minimal loss of biological activity will be used. The degree of conjugation will be closely monitored by SDS-PAGE and mass spectrometry to ensure proper conjugation of PEG molecules to the antibodies. Unreacted PEG can be separated from antibody-PEG conjugates by, e.g., size exclusion or ion-exchange chromatography.

15 Further, antibodies can be conjugated to albumin in order to make the antibody or antibody fragment more stable in vivo or have a longer half life *in vivo*. The techniques are well known in the art, see e.g., International Publication Nos. WO 93/15199, WO 93/15200, and WO 01/77137; and European Patent No. EP 413,622. The present invention encompasses the use of antibodies or fragments thereof conjugated or fused to one or more moieties,
20 including but not limited to, peptides, polypeptides, proteins, fusion proteins, nucleic acid molecules, small molecules, mimetic agents, synthetic drugs, inorganic molecules, and organic molecules.

The present invention encompasses the use of antibodies or fragments thereof
25 recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a heterologous protein or polypeptide (or fragment thereof, for example, to a polypeptide of at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids) to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker
30 sequences. For example, antibodies may be used to target heterologous polypeptides to particular cell types, either in vitro or in vivo, by fusing or conjugating the antibodies to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to heterologous polypeptides may also be used in in vitro immunoassays and purification

methods using methods known in the art. See e.g., International publication No. WO 93/21232; European Patent No. EP 439,095; Naramura et al., 1994, Immunol. Lett. 39:91-99; U.S. Pat.No. 5,474,981; Gillies et al., 1992, PNAS 89:1428-1432; and Fell et al., 1991, J. Immunol. 146:2446-2452.

5

The present invention further includes compositions comprising heterologous proteins, peptides or polypeptides fused or conjugated to antibody fragments. For example, the heterologous polypeptides may be fused or conjugated to a Fab fragment, Fd fragment, Fv fragment, F(ab)₂ fragment, a VH domain, a VL domain, a VH CDR, a VL CDR, or fragment thereof. Methods for fusing or conjugating polypeptides to antibody portions are well known in the art. See, e.g., U.S. Pat. Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851 and 5,112,946; European Patent Nos. EP 307,434 and EP 367,166; International publication Nos. WO 96/04388 and WO 91/06570; Ashkenazi et al., 1991, Proc. Natl. Acad. Sci. USA 88: 10535-10539; Zheng et al., 1995, J. Immunol. 154:5590-5600; and Vil et al., 1992, Proc. Natl. Acad. Sci. USA 89:11337-11341.

10

15

20

25

30

Additional fusion proteins, e.g. of antibodies that specifically bind an antigen (e.g., supra), may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to alter the activities of antibodies of the invention or fragments thereof (e.g., antibodies or fragments thereof with higher affinities and lower dissociation rates). See, generally, U.S. Pat. Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252 and 5,837,458, and Patten et al., 1997, Curr. Opin. Biotechnol. 8:724-33; Harayama, 1998, Trends Biotechnol. 16(2): 76-82; Hansson, et al., 1999, J. Mol. Biol. 287:265-76; and Lorenzo and Blasco, 1998, Biotechniques 24(2): 308-313. Antibodies or fragments thereof, or the encoded antibodies or fragments thereof, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. One or more portions of a polynucleotide encoding an antibody or antibody fragment, which portions specifically bind to an antigen may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

Drug conjugation

The present invention further encompasses uses of heteromultimers comprising variant Fc heterodimers or fragments thereof conjugated to a therapeutic agent or a cytotoxin.

An antibody or fragment thereof may be conjugated to a therapeutic moiety such as a
5 cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include ribonuclease, monomethylauristatin E and F, paclitaxel, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy
10 anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, epirubicin, and cyclophosphamide and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa
15 chlorambucil, melphalan, carmustine (BCNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cisdichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and
20 vinblastine). A more extensive list of therapeutic moieties can be found in PCT publications WO 03/075957.

Methods for fusing or conjugating antibodies to polypeptide moieties are known in the art. See, e.g., U.S. Pat. Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851 and
25 5,112,946; EP 307,434; EP 367,166; PCT Publications WO 96/04388 and WO 91/06570; Ashkenazi et al., 1991, PNAS USA 88:10535; Zheng et al., 1995, J Immunol 154:5590; and Vil et al., 1992, PNAS USA 89:11337. The fusion of an antibody to a moiety does not necessarily need to be direct, but may occur through linker sequences. Such linker molecules are commonly known in the art and described in Denardo et al., 1998, Clin
30 Cancer Res 4:2483; Peterson et al., 1999, Bioconjug Chem 10:553; Zimmerman et al., 1999, Nucl Med Biol 26:943; Garnett, 2002, Adv Drug Deliv Rev 53:171.

Recombinant expression

Recombinant expression of a heteromultimer, derivative, analog or fragment thereof, (e.g.,

an antibody or fusion protein of the invention), requires construction of an expression vector containing a polynucleotide that encodes the heteromultimer (e.g., antibody, or fusion protein). Once a polynucleotide encoding the heteromultimer (e.g., antibody, or fusion protein) has been obtained, the vector for the production of the heteromultimer (e.g.,
5 antibody, or fusion protein) may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing a heteromultimer (e.g., antibody, or fusion protein) encoding nucleotide sequence are described herein. Methods that are well known to those skilled in the art can be used to construct expression vectors containing the heteromultimer (e.g.,
10 antibody, or fusion protein) coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding a heteromultimer of the invention, operably linked to a promoter. Such vectors may include
15 the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., International Publication No. WO 86/05807; International Publication No. WO 89/01036; and U.S. Pat. No. 5,122,464 and the variable domain of the antibody, or a polypeptide for generating an Fc variant may be cloned into such a vector for expression of the full length antibody chain (e.g. heavy or light chain), or complete Fc variant comprising a fusion of a
20 non-antibody derived polypeptide and an Fc region incorporating at least the variant CH3 domain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an Fc variant of
25 the invention. Thus, the invention includes host cells containing a polynucleotide encoding a heteromultimer of the invention, operably linked to a heterologous promoter. In specific embodiments for the expression of heteromultimers comprising double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

30 A variety of host-expression vector systems may be utilized to express the heteromultimers of the invention (e.g., antibody or fusion protein molecules) (see, e.g., U.S. Pat. No. 5,807,715). Such host-expression systems represent vehicles by which the

coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an heteromultimer of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli* and *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing heteromultimer coding sequences; yeast (e.g., *Saccharomyces Pichia*) transformed with recombinant yeast expression vectors containing Fc variant coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing heteromultimer coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing Fc variant coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, NS0, and 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). In certain embodiments, bacterial cells such as *Escherichia coli*, or eukaryotic cells, are used for the expression of a heteromultimer, which is a recombinant antibody or fusion protein molecules. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., 1986, *Gene* 45:101; and Cockett et al., 1990, *Bio/Technology* 8:2). In a specific embodiment, the expression of nucleotide sequences encoding an Fc variant of the invention (e.g., antibody or fusion protein) is regulated by a constitutive promoter, inducible promoter or tissue specific promoter.

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the heteromultimer (e.g., antibody or fusion protein) being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an Fc variant, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited to, the *E. coli* expression vector pUR278 (Ruther et al., 1983, *EMBO* 12:1791), in which the heteromultimer coding

sequence may be ligated individually into the vector in frame with the lac Z coding region so that a lac Z-fusion protein is produced; pIN vectors (Inouye & Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke & Schuster, 1989, J. Biol. Chem. 24:5503-5509); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione 5-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The heteromultimer (e.g., antibody or fusion protein) coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the heteromultimer (e.g., antibody or fusion protein) coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the heteromultimer (e.g., antibody or fusion protein) in infected hosts (e.g., see Logan & Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:355-359). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see, e.g., Bittner et al., 1987, Methods in Enzymol. 153:516-544).

The expression of a heteromultimer (e.g., antibody or fusion protein) may be controlled by any promoter or enhancer element known in the art. Promoters which may be used to control the expression of the gene encoding a heteromultimer (e.g., antibody or fusion protein) include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, 1981, *Nature* 290:304-310), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto, et al., 1980, *Cell* 22:787-797), the herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1441-1445), the regulatory sequences of the metallothionein gene (Brinster et al., 1982, *Nature* 296:39-42), the tetracycline (Tet) promoter (Gossen et al., 1995, *Proc. Nat. Acad. Sci. USA* 89:5547-5551); prokaryotic expression vectors such as the β -lactamase promoter (Villa-Kamaroff et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731), or the tac promoter (DeBoer et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25; see also "Useful proteins from recombinant bacteria" in *Scientific American*, 1980, 242:74-94); plant expression vectors comprising the nopaline synthetase promoter region (Herrera-Estrella et al., *Nature* 303:209-213) or the cauliflower mosaic virus 35S RNA promoter (Gardner et al., 1981, *Nucl. Acids Res.* 9:2871), and the promoter of the photosynthetic enzyme ribulose biphosphate carboxylase (Herrera-Estrella et al., 1984, *Nature* 310:115-120); promoter elements from yeast or other fungi such as the Gal 4 promoter, the ADC (alcohol dehydrogenase) promoter, PGK (phosphoglycerol kinase) promoter, alkaline phosphatase promoter, and the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: elastase I gene control region which is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); insulin gene control region which is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122), immunoglobulin gene control region which is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444), mouse mammary tumor virus control region which is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495), albumin gene control region which is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276), alpha-fetoprotein gene control region which is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 235:53-58; alpha 1-antitrypsin gene control region which is active in the liver

(Kelsey et al., 1987, *Genes and Devel.* 1:161-171), beta-globin gene control region which is active in myeloid cells (Mogam et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94; myelin basic protein gene control region which is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712); myosin light chain-2 gene control region which is active in skeletal muscle (Sani, 1985, *Nature* 314:283-286); neuronal-specific enolase (NSE) which is active in neuronal cells (Morelli et al., 1999, *Gen. Virol.* 80:571-83); brain-derived neurotrophic factor (BDNF) gene control region which is active in neuronal cells (Tabuchi et al., 1998, *Biochem. Biophys. Res. Com.* 253:818-823); glial fibrillary acidic protein (GFAP) promoter which is active in astrocytes (Gomes et al., 1999, *Braz J Med Biol Res* 32(5): 619-631; Morelli et al., 1999, *Gen. Virol.* 80:571-83) and gonadotropic releasing hormone gene control region which is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378).

Expression vectors containing inserts of a gene encoding a heteromultimer of the invention (e.g., antibody or fusion protein) can be identified by three general approaches: (a) nucleic acid hybridization, (b) presence or absence of "marker" gene functions, and (c) expression of inserted sequences. In the first approach, the presence of a gene encoding a peptide, polypeptide, protein or a fusion protein in an expression vector can be detected by nucleic acid hybridization using probes comprising sequences that are homologous to an inserted gene encoding the peptide, polypeptide, protein or the fusion protein, respectively. In the second approach, the recombinant vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., thymidine kinase activity, resistance to antibiotics, transformation phenotype, occlusion body formation in baculovirus, etc.) caused by the insertion of a nucleotide sequence encoding an antibody or fusion protein in the vector. For example, if the nucleotide sequence encoding the heteromultimer (e.g., antibody or fusion protein) is inserted within the marker gene sequence of the vector, recombinants containing the gene encoding the antibody or fusion protein insert can be identified by the absence of the marker gene function. In the third approach, recombinant expression vectors can be identified by assaying the gene product (e.g., antibody or fusion protein) expressed by the recombinant. Such assays can be based, for example, on the physical or functional properties of the fusion protein in *in vitro* assay systems, e.g., binding with anti-bioactive molecule antibody.

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus, expression of the genetically engineered fusion protein may be controlled.

5 Furthermore, different host cells have characteristic and specific mechanisms for the translational and post-translational processing and modification (e.g., glycosylation, phosphorylation of proteins). Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. For example, expression in a bacterial system will produce an unglycosylated product and
10 expression in yeast will produce a glycosylated product. Eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript (e.g., glycosylation, and phosphorylation) of the gene product may be used. Such mammalian host cells include, but are not limited to, CHO, VERY, BHK, Hela, COS, MDCK, 293, 3T3, WI38, NS0, and in particular, neuronal cell lines such as, for example, SK-N-AS, SK-N-FI, SK-N-
15 DZ human neuroblastomas (Sugimoto et al., 1984, J. Natl. Cancer Inst. 73: 51-57), SK-N-SH human neuroblastoma (Biochim. Biophys. Acta, 1982, 704: 450-460), Daoy human cerebellar medulloblastoma (He et al., 1992, Cancer Res. 52: 1144-1148) DBTRG-05MG glioblastoma cells (Kruse et al., 1992, In Vitro Cell. Dev. Biol. 28A: 609-614), IMR-32 human neuroblastoma (Cancer Res., 1970, 30: 2110-2118), 1321N1 human astrocytoma
20 (Proc. Natl. Acad. Sci. USA, 1977, 74: 4816), MOG-G-CCM human astrocytoma (Br. J. Cancer, 1984, 49: 269), U87MG human glioblastoma-astrocytoma (Acta Pathol. Microbiol. Scand., 1968, 74: 465-486), A172 human glioblastoma (Olopade et al., 1992, Cancer Res. 52: 2523-2529), C6 rat glioma cells (Benda et al., 1968, Science 161: 370-371), Neuro-2a mouse neuroblastoma (Proc. Natl. Acad. Sci. USA, 1970, 65: 129-136), NB41A3 mouse
25 neuroblastoma (Proc. Natl. Acad. Sci. USA, 1962, 48: 1184-1190), SCP sheep choroid plexus (Bolin et al., 1994, J. Virol. Methods 48: 211-221), G355-5, PG-4 Cat normal astrocyte (Haapala et al., 1985, J. Virol. 53: 827-833), Mpf ferret brain (Trowbridge et al., 1982, In Vitro 18: 952-960), and normal cell lines such as, for example, CTX TNA2 rat normal cortex brain (Radany et al., 1992, Proc. Natl. Acad. Sci. USA 89: 6467-6471) such
30 as, for example, CRL7030 and Hs578Bst. Furthermore, different vector/host expression systems may effect processing reactions to different extents.

For long-term, high-yield production of recombinant proteins, stable expression is often

preferred. For example, cell lines that stably express a heteromultimer of the invention (e.g., antibody or fusion protein) may be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched medium, and then are switched to a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci that in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express a heteromultimer that specifically binds to an Antigen. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the activity of a heteromultimer (e.g., a polypeptide or a fusion protein) that specifically binds to an antigen.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes can be employed in tk-, hgprt- or aprt-cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler et al., 1980, Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hygro, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147) genes.

Purification

Once a heteromultimer (e.g., antibody, or a fusion protein) of the invention has been produced by recombinant expression, it may be purified by any method known in the art for purification of a protein, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins.

The heteromultimer is generally recovered from the culture medium as a secreted polypeptide, although it also may be recovered from host cell lysate when directly produced without a secretory signal. If the heteromultimer is membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100).

When the heteromultimer is produced in a recombinant cell other than one of human origin, it is completely free of proteins or polypeptides of human origin. However, it is necessary to purify the heteromultimer from recombinant cell proteins or polypeptides to obtain preparations that are substantially homogeneous as to the heteromultimer. As a first step, the culture medium or lysate is normally centrifuged to remove particulate cell debris.

Heteromultimers having antibody constant domains can be conveniently purified by hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography, with affinity chromatography being the preferred purification technique. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin Sepharose, chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the polypeptide to be recovered. The suitability of protein A as an affinity ligand depends on the species and isotype of the immunoglobulin Fc domain that is used. Protein A can be used to purify immunoglobulin Fc regions that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$ heavy chains (Lindmark et al., J. Immunol. Meth. 62:1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss et al., EMBO J. 5:1567-1575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. The conditions for binding an immunoadhesin to the protein A or G affinity column are dictated entirely by the characteristics of the Fc domain; that is, its species and isotype. Generally, when the proper ligand is chosen, efficient binding occurs directly from unconditioned culture fluid. Bound variant Fc heterodimers can be efficiently eluted either at acidic pH (at or above 3.0), or in a neutral pH buffer containing a mildly chaotropic salt. This affinity chromatography step can result in a variant Fc

heterodimer preparation that is >95% pure.

The expression levels of a heteromultimer (e.g., antibody or fusion protein) can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of
5 vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol. 3. (Academic Press, New York, 1987)). For example, when a marker in the vector system expressing an antibody or fusion protein is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody
10 gene, production of the antibody or fusion protein will also increase (Crouse et al., 1983, Mol. Cell. Biol. 3:257).

Characterization and Functional Assays

15 Fc variants (e.g., antibodies or fusion proteins) of the present invention may be characterized in a variety of ways. In one embodiment, purity of the variant Fc heterodimers is assessed using techniques well known in the art including, but not limited to, SDS-PAGE gels, western blots, densitometry or mass spectrometry. Protein stability
20 can be characterized using an array of techniques, not limited to, size exclusion chromatography, UV Visible and CD spectroscopy, mass spectroscopy, differential light scattering, bench top stability assay, freeze thawing coupled with other characterization techniques, differential scanning calorimetry, differential scanning fluorimetry, hydrophobic interaction chromatography, isoelectric focusing, receptor binding assays or relative
25 protein expression levels. In an exemplary embodiment, stability of the variant Fc heterodimers is assessed by melting temperature of the variant CH3 domain, as compared to wild-type CH3 domain, using techniques well known in the art such as Differential Scanning Calorimetry or differential scanning fluorimetry.

30 Fc variants of the present invention may also be assayed for the ability to specifically bind to a ligand, (e.g., FcγRIIIA, FcγRIIB, C1q). Such an assay may be performed in solution (e.g., Houghten, Bio/Techniques, 13:412-421, 1992), on beads (Lam, Nature, 354:82-84, 1991, on chips (Fodor, Nature, 364:555-556, 1993), on bacteria (U.S. Pat. No. 5,223,409)

on plasmids (Cull et al., Proc. Natl. Acad. Sci. USA, 89:1865-1869, 1992) or on phage (Scott and Smith, Science, 249:386-390, 1990; Devlin, Science, 249:404-406, 1990; Cwirla et al., Proc. Natl. Acad. Sci. USA, 87:6378-6382, 1990; and Felici, J. Mol. Biol., 222:301-310, 1991). Molecules that have been identified to specifically bind to a ligand, (e.g.,
5 FcγRIIIA, FcγRIIB, C1q or to an antigen) can then be assayed for their affinity for the ligand.

Fc variants of the invention may be assayed for specific binding to a molecule such as an antigen (e.g., cancer antigen and cross-reactivity with other antigens) or a ligand (e.g.,
10 FcγR) by any method known in the art. Immunoassays which can be used to analyze specific binding and cross-reactivity include, but are not limited to, competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions,
15 immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al., eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York).

20 The binding affinity of the Fc variants of the present invention to a molecule such as an antigen or a ligand, (e.g., FcγR) and the off-rate of the interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled ligand, such as FcγR (e.g., 3H or 125I with a molecule of interest (e.g., Fc variants of the present invention) in the presence
25 of increasing amounts of unlabeled ligand, such as FcγR, and the detection of the molecule bound to the labeled ligand. The affinity of the molecule of the present invention for the ligand and the binding off-rates can be determined from the saturation data by scatchard analysis.

30 *Affinity maturation*

As is known in the art, once a single domain antigen-binding construct has been identified, and its affinity for the target antigen measured, if necessary, the affinity of the single domain antigen-binding construct for its target antigen can be improved by affinity

maturation according to methods known in the art. One exemplary method for affinity maturation of an antigen-binding domain where the crystal structure of the target antigen to the antibody is available is described as follows. Structures of the antigen:antibody complex are used for modeling. Molecular dynamics (MD) can be employed to evaluate the intrinsic dynamic nature of the WT complex in an aqueous environment. Mean field and dead-end elimination methods along with flexible backbones can be used to optimize and prepare model structures for the mutants to be screened. Following packing a number of features will be scored including contact density, clash score, hydrophobicity and electrostatics. Generalized Born method will allow accurate modeling of the effect of solvent environment and compute the free energy differences following mutation of specific positions in the protein to alternate residue types. Contact density and clash score will provide a measure of complementarity, a critical aspect of effective protein packing. The screening procedure employs knowledge-based potentials as well as coupling analysis schemes relying on pair-wise residue interaction energy and entropy computations.

The kinetic parameters of an Fc variant may also be determined using any surface plasmon resonance (SPR) based assays known in the art (e.g., BIAcore kinetic analysis). For a review of SPR-based technology see Mullet et al., 2000, *Methods* 22: 77-91; Dong et al., 2002, *Review in Mol. Biotech.*, 82: 303-23; Fivash et al., 1998, *Current Opinion in Biotechnology* 9: 97-101; Rich et al., 2000, *Current Opinion in Biotechnology* 11: 54-61.

Additionally, any of the SPR instruments and SPR based methods for measuring protein-protein interactions described in U.S. Pat. Nos. 6,373,577; 6,289,286; 5,322,798; 5,341,215; 6,268,125 are contemplated in the methods of the invention.

Fluorescence activated cell sorting (FACS), using any of the techniques known to those skilled in the art, can be used for characterizing the binding of Fc variants to a molecule expressed on the cell surface (e.g., FcγRIIIA, FcγRIIB). Flow sorters are capable of rapidly examining a large number of individual cells that contain library inserts (e.g., 10-100 million cells per hour) (Shapiro et al., *Practical Flow, Cytometry*, 1995). Flow cytometers for sorting and examining biological cells are well known in the art. Known flow cytometers are described, for example, in U.S. Pat. Nos. 4,347,935; 5,464,581; 5,483,469; 5,602,039; 5,643,796; and 6,211,477. Other known flow cytometers are the FACS Vantage™ system manufactured by Becton Dickinson and Company, and the COPAS™ system

manufactured by Union Biometrica.

The Fc variants of the invention can be characterized by their ability to mediate FcγR-mediated effector cell function. Examples of effector cell functions that can be assayed include, but are not limited to, antibody-dependent cell mediated cytotoxicity (ADCC), phagocytosis, opsonization, opsonophagocytosis, C1q binding, and complement dependent cell mediated cytotoxicity (CDC). Any cell-based or cell free assay known to those skilled in the art for determining effector cell function activity can be used (For effector cell assays, see Perussia et al., 2000, *Methods Mol. Biol.* 121: 179-92; Baggiolini et al., 1998 *Experientia*, 44(10): 841-8; Lehmann et al., 2000 *J. Immunol. Methods*, 243(1-2): 229-42; Brown E J. 1994, *Methods Cell Biol.*, 45: 147-64; Munn et al., 1990 *J. Exp. Med.*, 172: 231-237, Abdul-Majid et al., 2002 *Scand. J. Immunol.* 55: 70-81; Ding et al., 1998, *Immunity* 8:403-411).

In particular, the Fc variants of the invention can be assayed for FcγR-mediated ADCC activity in effector cells, (e.g., natural killer cells) using any of the standard methods known to those skilled in the art (See e.g., Perussia et al., 2000, *Methods Mol. Biol.* 121: 179-92). An exemplary assay for determining ADCC activity of the molecules of the invention is based on a ⁵¹Cr release assay comprising of: labeling target cells with [⁵¹Cr]Na₂CrO₄ (this cell-membrane permeable molecule is commonly used for labeling since it binds cytoplasmic proteins and although spontaneously released from the cells with slow kinetics, it is released massively following target cell necrosis); opsonizing the target cells with the Fc variants of the invention; combining the opsonized radiolabeled target cells with effector cells in a microtitre plate at an appropriate ratio of target cells to effector cells; incubating the mixture of cells for 16-18 hours at 37° C.; collecting supernatants; and analyzing radioactivity. The cytotoxicity of the molecules of the invention can then be determined, for example using the following formula: % lysis=(experimental cpm–target leak cpm)/(detergent lysis cpm–target leak cpm)×100%. Alternatively, % lysis=(ADCC–AICC)/(maximum release–spontaneous release). Specific lysis can be calculated using the formula: specific lysis=% lysis with the molecules of the invention–% lysis in the absence of the molecules of the invention. A graph can be generated by varying either the target:effector cell ratio or antibody concentration.

Method to characterize the ability of the Fc variants to bind C1q and mediate complement dependent cytotoxicity (CDC) are well known in the art. For example, to determine C1q binding, a C1q binding ELISA may be performed. An exemplary assay may comprise the following: assay plates may be coated overnight at 4 C with polypeptide variant or starting polypeptide (control) in coating buffer. The plates may then be washed and blocked.

Following washing, an aliquot of human C1q may be added to each well and incubated for 2 hrs at room temperature. Following a further wash, 100 uL of a sheep anti-complement C1q peroxidase conjugated antibody may be added to each well and incubated for 1 hour at room temperature. The plate may again be washed with wash buffer and 100 ul of substrate buffer containing OPD (O-phenylenediamine dihydrochloride (Sigma)) may be added to each well. The oxidation reaction, observed by the appearance of a yellow color, may be allowed to proceed for 30 minutes and stopped by the addition of 100 ul of 4.5 NH₂ SO₄. The absorbance may then read at (492-405) nm.

To assess complement activation, a complement dependent cytotoxicity (CDC) assay may be performed, (e.g. as described in Gazzano-Santoro et al., 1996, J. Immunol. Methods 202:163). Briefly, various concentrations of Fc variant and human complement may be diluted with buffer. Cells which express the antigen to which the Fc variant binds may be diluted to a density of about 1×10⁶ cells/ml. Mixtures of the Fc variant, diluted human complement and cells expressing the antigen may be added to a flat bottom tissue culture 96 well plate and allowed to incubate for 2 hrs at 37 C. and 5% CO₂ to facilitate complement mediated cell lysis. 50 uL of alamar blue (Accumed International) may then be added to each well and incubated overnight at 37 C. The absorbance is measured using a 96-well fluorometer with excitation at 530 nm and emission at 590 nm. The results may be expressed in relative fluorescence units (RFU). The sample concentrations may be computed from a standard curve and the percent activity, relative to a comparable molecule (i.e., a molecule comprising an Fc region with an unmodified or wild type CH3 domain) is reported for the Fc variant of interest.

Complement assays may be performed with guinea pig, rabbit or human serum. Complement lysis of target cells may be detected by monitoring the release of intracellular enzymes such as lactate dehydrogenase (LDH), as described in Korzeniewski et al., 1983, Immunol. Methods 64(3): 313-20; and Decker et al., 1988, J. Immunol Methods 115(1): 61-

9; or the release of an intracellular label such as europium, chromium 51 or indium 111 in which target cells are labeled.

Pharmacokinetic stability

- 5 In certain embodiments, the heteromultimers provided herein exhibits pharmacokinetic (PK) or *in vivo* stability properties comparable with commercially available therapeutic antibodies. In one embodiment, the heteromultimers described herein exhibit PK properties similar to known therapeutic antibodies, with respect to serum concentration, $t_{1/2}$, beta half-life, and/or CL.
- 10 In some embodiments, active transport processes such as uptake by neonatal Fc receptor (FcRn) also impact antibody biodistribution among other binding proteins. In one embodiment, the heteromultimers bind FcRn with similar affinity compared to commercially available therapeutic antibodies.

Humanized single domain antigen-binding constructs

- 15 In some embodiments it may be necessary to humanize the single domain antigen-binding construct prior to use in the heteromultimer described herein, for example, if the single domain antigen-binding construct is obtained from a camelid or shark. "Humanized" forms of non-human antibodies are chimeric antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human
- 20 immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity, affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore,
- 25 humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human immunoglobulin and all or
- 30 substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin

constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., *Nature* 321:522-525 (1986); Riechmann et al., *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

Methods of treatment

5 The present invention encompasses administering one or more heteromultimer of the invention (e.g., antibodies) to an animal, in particular a mammal, specifically, a human, for preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection.

10 In one embodiment, the heteromultimers described herein exert their therapeutic effects through Fc effector function activity. In one embodiment, the heteromultimers described herein are conjugated to a cytotoxic drug molecule, and exert their therapeutic effects through internalization of the heteromultimer and cytotoxic drug molecule into the target cell.

15 In one embodiment, the heteromultimers are used to treat infections of pathogenic organisms, such as bacteria or fungi.

The heteromultimers of the invention are particularly useful for the treatment or prevention
20 of a disease or disorder where an altered efficacy of effector cell function (e.g., ADCC, CDC) is desired. The heteromultimers and compositions thereof are particularly useful for the treatment or prevention of primary or metastatic neoplastic disease (i.e., cancer), and infectious diseases. Molecules of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein. As detailed below, the
25 molecules of the invention can be used in methods of treating or preventing cancer (particularly in passive immunotherapy), autoimmune disease, inflammatory disorders or infectious diseases.

The heteromultimers of the invention may also be advantageously utilized in combination
30 with other therapeutic agents known in the art for the treatment or prevention of a cancer, autoimmune disease, inflammatory disorders or infectious diseases. In a specific embodiment, heteromultimers of the invention may be used in combination with

monoclonal or chimeric antibodies, lymphokines, or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), which, for example, serve to increase the number or activity of effector cells which interact with the molecules and, increase immune response. The heteromultimers of the invention may also be advantageously utilized in combination with one or more drugs used to treat a disease, disorder, or infection such as, for example anti-cancer agents, anti-inflammatory agents or anti-viral agents.

Accordingly, the present invention provides methods for preventing, treating, or ameliorating one or more symptoms associated with cancer and related conditions by administering one or more heteromultimers of the invention. Although not intending to be bound by any mechanism of actions, a heteromultimer of the invention that binds FcγRIIIA and/or FcγRIIA with a greater affinity than a comparable molecule, and further binds FcγRIIB with a lower affinity than a comparable molecule, and/or said Heteromultimer has an enhanced effector function, e.g., ADCC, CDC, phagocytosis, opsonization, etc. will result in the selective targeting and efficient destruction of cancer cells.

The invention further encompasses administering one or more heteromultimers of the invention in combination with other therapies known to those skilled in the art for the treatment or prevention of cancer, including but not limited to, current standard and experimental chemotherapies, hormonal therapies, biological therapies, immunotherapies, radiation therapies, or surgery. In some embodiments, the molecules of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or more anti-cancer agents, therapeutic antibodies or other agents known to those skilled in the art for the treatment and/or prevention of cancer. Examples of dosing regimes and therapies which can be used in combination with the heteromultimers of the invention are well known in the art and have been described in detail elsewhere (see for example, PCT publications WO 02/070007 and WO 03/075957).

Cancers and related disorders that can be treated or prevented by methods and compositions of the present invention include, but are not limited to, the following: Leukemias, lymphomas, multiple myelomas, bone and connective tissue sarcomas, brain tumors, breast cancer, adrenal cancer, thyroid cancer, pancreatic cancer, pituitary cancers, eye cancers, vaginal cancers, vulvar cancer, cervical cancers, uterine cancers,

ovarian cancers, esophageal cancers, stomach cancers, colon cancers, rectal cancers, liver cancers, gallbladder cancers, cholangiocarcinomas, lung cancers, testicular cancers, prostate cancers, penal cancers; oral cancers, salivary gland cancers pharynx cancers, skin cancers, kidney cancers, bladder cancers (for a review of such disorders, see

- 5 Fishman et al., 1985, *Medicine*, 2d Ed., J.B. Lippincott Co., Philadelphia and Murphy et al., 1997, *Informed Decisions: The Complete Book of Cancer Diagnosis, Treatment, and Recovery*, Viking Penguin, Penguin Books U.S.A., Inc., United States of America).

- 10 In one embodiment, where the heteromultimer comprises a single domain antigen-binding construct that bind to EGFR1, EGFR1 or the mutant EGFR variant III (EGFRvIII) expressing cells, the heteromultimer can be used to treat cancers that overexpress EGFR1 or cancer cells that are resistant to treatment by binding to EGFRvIII.

- 15 The invention further contemplates engineering any of the antibodies known in the art for the treatment and/or prevention of cancer and related disorders, so that the antibodies comprise an Fc region incorporating a variant CH3 domain of the invention.

- 20 In a specific embodiment, a molecule of the invention (e.g., an antibody comprising a variant Fc heterodimer inhibits or reduces the growth of primary tumor or metastasis of cancerous cells by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 45%, at least 35%, at least 30%, at least 25%, at least 20%, or at least 10% relative to the growth of primary tumor or metastasis in the absence of said molecule of the invention.

- 25 The present invention encompasses the use of one or more heteromultimers of the invention for preventing, treating, or managing one or more symptoms associated with an inflammatory disorder in a subject. Although not intending to be bound by any mechanism of actions, heteromultimers with enhanced affinity for FcγRIIB will lead to a dampening of the activating receptors and thus a dampening of the immune response and have
30 therapeutic efficacy for treating and/or preventing an autoimmune disorder. Furthermore, antibodies binding more than one target, such as bi-specific antibodies comprising a variant Fc heterodimer, associated with an inflammatory disorder may provide synergist effects over monovalent therapy.

The invention further encompasses administering the heteromultimers of the invention in combination with a therapeutically or prophylactically effective amount of one or more anti-inflammatory agents. The invention also provides methods for preventing, treating, or
5 managing one or more symptoms associated with an autoimmune disease further comprising, administering to said subject a heteromultimer of the invention in combination with a therapeutically or prophylactically effective amount of one or more immunomodulatory agents. Examples of autoimmune disorders that may be treated by administering the heteromultimers of the invention include, but are not limited to, alopecia
10 areata, ankylosing spondylitis, antiphospholipid syndrome, autoimmune Addison's disease, autoimmune diseases of the adrenal gland, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune oophoritis and orchitis, autoimmune thrombocytopenia, Behcet's disease, bullous pemphigoid, cardiomyopathy, celiac sprue-dermatitis, chronic fatigue immune dysfunction syndrome (CFIDS), chronic inflammatory demyelinating
15 polyneuropathy, Churg-Strauss syndrome, cicatricial pemphigoid, CREST syndrome, cold agglutinin disease, Crohn's disease, discoid lupus, essential mixed cryoglobulinemia, fibromyalgia-fibromyositis, glomerulonephritis, Graves' disease, Guillain-Barre, Hashimoto's thyroiditis, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), IgA neuropathy, juvenile arthritis, lichen planus, lupus erythematosus, Meniere's
20 disease, mixed connective tissue disease, multiple sclerosis, type 1 or immune-mediated diabetes mellitus, myasthenia gravis, pemphigus vulgaris, pernicious anemia, polyarteritis nodosa, polychondritis, polyglandular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, psoriatic arthritis, Raynaud's phenomenon, Reiter's syndrome, Rheumatoid arthritis,
25 sarcoidosis, scleroderma, Sjogren's syndrome, stiff-man syndrome, systemic lupus erythematosus, lupus erythematosus, takayasu arteritis, temporal arteritis/giant cell arteritis, ulcerative colitis, uveitis, vasculitides such as dermatitis herpetiformis vasculitis, vitiligo, and Wegener's granulomatosis. Examples of inflammatory disorders include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive
30 pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthropathy, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections. Some autoimmune disorders are associated with an inflammatory condition, thus, there is

overlap between what is considered an autoimmune disorder and an inflammatory disorder. Therefore, some autoimmune disorders may also be characterized as inflammatory disorders. Examples of inflammatory disorders which can be prevented, treated or managed in accordance with the methods of the invention include, but are not limited to, asthma, encephalitis, inflammatory bowel disease, chronic obstructive pulmonary disease (COPD), allergic disorders, septic shock, pulmonary fibrosis, undifferentiated spondyloarthritis, undifferentiated arthropathy, arthritis, inflammatory osteolysis, and chronic inflammation resulting from chronic viral or bacteria infections.

10 Heteromultimers of the invention can also be used to reduce the inflammation experienced by animals, particularly mammals, with inflammatory disorders. In a specific embodiment, an Fc of the invention reduces the inflammation in an animal by at least 99%, at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, at least 50%, at least 45%, at least 40%, at least 35%, at least 30%, at least 25%, at
15 least 20%, or at least 10% relative to the inflammation in an animal, which is not administered the said molecule.

The invention further contemplates engineering any of the antibodies known in the art for the treatment and/or prevention of autoimmune disease or inflammatory disease, so that
20 the antibodies comprise a variant Fc heterodimer of the invention.

The invention also encompasses methods for treating or preventing an infectious disease in a subject comprising administering a therapeutically or prophylactically effective amount of one or more heteromultimers of the invention. Infectious diseases that can be treated or
25 prevented by the heteromultimers of the invention are caused by infectious agents including but not limited to viruses, bacteria, fungi, protozoa, and viruses.

Viral diseases that can be treated or prevented using the heteromultimers of the invention in conjunction with the methods of the present invention include, but are not limited to,
30 those caused by hepatitis type A, hepatitis type B, hepatitis type C, influenza, varicella, adenovirus, herpes simplex type I (HSV-I), herpes simplex type II (HSV-II), rinderpest, rhinovirus, echovirus, rotavirus, respiratory syncytial virus, papilloma virus, papova virus, cytomegalovirus, echinovirus, arbovirus, hantavirus, coxsackie virus, mumps virus,

measles virus, rubella virus, polio virus, small pox, Epstein Barr virus, human immunodeficiency virus type I (HIV-I), human immunodeficiency virus type II (HIV-II), and agents of viral diseases such as viral meningitis, encephalitis, dengue or small pox.

5 Bacterial diseases that can be treated or prevented using the heteromultimers of the invention in conjunction with the methods of the present invention, that are caused by bacteria include, but are not limited to, mycobacteria rickettsia, mycoplasma, neisseria, S. pneumonia, *Borrelia burgdorferi* (Lyme disease), *Bacillus anthracis* (anthrax), tetanus, streptococcus, staphylococcus, mycobacterium, tetanus, pertissus, cholera, plague, 10 diphtheria, chlamydia, *S. aureus* and legionella. Protozoal diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present invention, that are caused by protozoa include, but are not limited to, leishmania, kokzidioa, trypanosoma or malaria. Parasitic diseases that can be treated or prevented using the molecules of the invention in conjunction with the methods of the present 15 invention, that are caused by parasites include, but are not limited to, chlamydia and rickettsia.

In some embodiments, the heteromultimers of the invention may be administered in combination with a therapeutically or prophylactically effective amount of one or additional 20 therapeutic agents known to those skilled in the art for the treatment and/or prevention of an infectious disease. The invention contemplates the use of the molecules of the invention in combination with other molecules known to those skilled in the art for the treatment and or prevention of an infectious disease including, but not limited to, antibiotics, antifungal agents and anti-viral agents.

25 The invention provides methods and pharmaceutical compositions comprising heteromultimers of the invention (e.g., antibodies, polypeptides). The invention also provides methods of treatment, prophylaxis, and amelioration of one or more symptoms associated with a disease, disorder or infection by administering to a subject an effective 30 amount of at least one Heteromultimer of the invention, or a pharmaceutical composition comprising at least one Heteromultimer of the invention. In a one aspect, the Heteromultimer, is substantially purified (i.e., substantially free from substances that limit its effect or produce undesired side-effects this includes homodimers and other cellular

material). In a specific embodiment, the subject is an animal, such as a mammal including non-primates (e.g., cows, pigs, horses, cats, dogs, rats etc.) and primates (e.g., monkey such as, a cynomolgous monkey and a human). In a specific embodiment, the subject is a human. In yet another specific embodiment, the antibody of the invention is from the same species as the subject.

The route of administration of the composition depends on the condition to be treated. For example, intravenous injection may be preferred for treatment of a systemic disorder such as a lymphatic cancer or a tumor that has metastasized. The dosage of the compositions to be administered can be determined by the skilled artisan without undue experimentation in conjunction with standard dose-response studies. Relevant circumstances to be considered in making those determinations include the condition or conditions to be treated, the choice of composition to be administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms. Depending on the condition, the composition can be administered orally, parenterally, intranasally, vaginally, rectally, lingually, sublingually, buccally, intrabuccally and/or transdermally to the patient.

Accordingly, compositions designed for oral, lingual, sublingual, buccal and intrabuccal administration can be made without undue experimentation by means well known in the art, for example, with an inert diluent or with an edible carrier. The composition may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the pharmaceutical compositions of the present invention may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums, and the like.

Tablets, pills, capsules, troches and the like may also contain binders, recipients, disintegrating agent, lubricants, sweetening agents, and/or flavoring agents. Some examples of binders include microcrystalline cellulose, gum tragacanth and gelatin. Examples of excipients include starch and lactose. Some examples of disintegrating agents include alginic acid, cornstarch, and the like. Examples of lubricants include magnesium stearate and potassium stearate. An example of a glidant is colloidal silicon dioxide. Some examples of sweetening agents include sucrose, saccharin, and the like. Examples of flavoring agents include peppermint, methyl salicylate, orange flavoring, and

the like. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

The pharmaceutical compositions of the present invention can be administered

5 parenterally, such as, for example, by intravenous, intramuscular, intrathecal and/or subcutaneous injection. Parenteral administration can be accomplished by incorporating the compositions of the present invention into a solution or suspension. Such solutions or suspensions may also include sterile diluents, such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol and/or other synthetic solvents.

10 Parenteral formulations may also include antibacterial agents, such as, for example, benzyl alcohol and/or methyl parabens, antioxidants, such as, for example, ascorbic acid and/or sodium bisulfite, and chelating agents, such as EDTA. Buffers, such as acetates, citrates and phosphates, and agents for the adjustment of tonicity, such as sodium chloride and dextrose, may also be added. The parenteral preparation can be enclosed in ampules,

15 disposable syringes and/or multiple dose vials made of glass or plastic. Rectal administration includes administering the composition into the rectum and/or large intestine. This can be accomplished using suppositories and/or enemas. Suppository formulations can be made by methods known in the art. Transdermal administration includes percutaneous absorption of the composition through the skin. Transdermal

20 formulations include patches, ointments, creams, gels, salves, and the like. The compositions of the present invention can be administered nasally to a patient. As used herein, nasally administering or nasal administration includes administering the compositions to the mucous membranes of the nasal passage and/or nasal cavity of the patient.

25 The pharmaceutical compositions of the invention may be used in accordance with the methods of the invention for preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection. It is contemplated that the pharmaceutical compositions of the invention are sterile and in suitable form for administration to a subject.

30 In one embodiment the compositions of the invention are pyrogen-free formulations that are substantially free of endotoxins and/or related pyrogenic substances. Endotoxins include toxins that are confined inside a microorganism and are released when the

microorganisms are broken down or die. Pyrogenic substances also include fever-inducing, thermostable substances (glycoproteins) from the outer membrane of bacteria and other microorganisms. Both of these substances can cause fever, hypotension and shock if administered to humans. Due to the potential harmful effects, it is advantageous to remove even low amounts of endotoxins from intravenously administered pharmaceutical drug solutions. The Food & Drug Administration ("FDA") has set an upper limit of 5 endotoxin units (EU) per dose per kilogram body weight in a single one hour period for intravenous drug applications (The United States Pharmacopeial Convention, Pharmacopeial Forum 26 (1):223 (2000)). When therapeutic proteins are administered in amounts of several hundred or thousand milligrams per kilogram body weight, as can be the case with monoclonal antibodies, it is advantageous to remove even trace amounts of endotoxin. In a specific embodiment, endotoxin and pyrogen levels in the composition are less than 10 EU/mg, or less than 5 EU/mg, or less than 1 EU/mg, or less than 0.1 EU/mg, or less than 0.01 EU/mg, or less than 0.001 EU/mg.

The invention provides methods for preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection, said method comprising: (a) administering to a subject in need thereof a dose of a prophylactically or therapeutically effective amount of a composition comprising one or more heteromultimers and (b) administering one or more subsequent doses of said heteromultimers, to maintain a plasma concentration of the Heteromultimer at a desirable level (e.g., about 0.1 to about 100 µg/ml), which continuously binds to an antigen. In a specific embodiment, the plasma concentration of the Heteromultimer is maintained at 10 µg/ml, 15 µg/ml, 20 µg/ml, 25 µg/ml, 30 µg/ml, 35 µg/ml, 40 µg/ml, 45 µg/ml or 50 µg/ml. In a specific embodiment, said effective amount of Heteromultimer to be administered is between at least 1 mg/kg and 8 mg/kg per dose. In another specific embodiment, said effective amount of Heteromultimer to be administered is between at least 4 mg/kg and 5 mg/kg per dose. In yet another specific embodiment, said effective amount of Heteromultimer to be administered is between 50 mg and 250 mg per dose. In still another specific embodiment, said effective amount of Fc valiant to be administered is between 100 mg and 200 mg per dose.

The present invention also encompasses protocols for preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection which a

heteromultimer is used in combination with a therapy (e.g., prophylactic or therapeutic agent) other than a heteromultimer and/or variant fusion protein. The invention is based, in part, on the recognition that the heteromultimers of the invention potentiate and synergize with, enhance the effectiveness of, improve the tolerance of, and/or reduce the side effects caused by, other cancer therapies, including current standard and experimental chemotherapies. The combination therapies of the invention have additive potency, an additive therapeutic effect or a synergistic effect. The combination therapies of the invention enable lower dosages of the therapy (e.g., prophylactic or therapeutic agents) utilized in conjunction with heteromultimers for preventing, treating, or ameliorating one or more symptoms associated with a disease, disorder, or infection and/or less frequent administration of such prophylactic or therapeutic agents to a subject with a disease disorder, or infection to improve the quality of life of said subject and/or to achieve a prophylactic or therapeutic effect. Further, the combination therapies of the invention reduce or avoid unwanted or adverse side effects associated with the administration of current single agent therapies and/or existing combination therapies, which in turn improves patient compliance with the treatment protocol. Numerous molecules which can be utilized in combination with the heteromultimers of the invention are well known in the art. See for example, PCT publications WO 02/070007; WO 03/075957 and U.S. Patent Publication 2005/064514.

Industrial Uses

In view of the biophysical stability of both the single domain antigen-binding construct and the heterodimeric Fc region of the heteromultimer, it is contemplated that the heteromultimers described herein can also be used in industrial applications in which single domain antigen-binding construct fragments (such as isolated V_HH) themselves have utility (see de Marco (2011) *Microbial Cell factories* 10:44). Thus, in one embodiment, the heteromultimer according to the invention may be used to identify and detoxify toxins, as reagents for immunodetection, purification and bioseparation, as crystallography chaperones, or as tools for studying protein aggregation and activity regulation.

Kits

The present invention provides kits comprising one or more heteromultimers with altered binding affinity to FcγRs and/or C1q and altered ADCC and/or CDC activity that specifically bind to an antigen conjugated or fused to a detectable agent, therapeutic agent or drug, in one or more containers, for use in monitoring, diagnosis, preventing, treating, or
5 ameliorating one or more symptoms associated with a disease, disorder, or infection.

EXAMPLES

The examples below are given so as to illustrate the practice of this invention. They are not intended to limit or define the entire scope of this invention.

Example 1: Generation of bivalent monospecific Antibodies with Heterodimer Fc domains.

The genes encoding the antibody heavy chains were constructed via gene synthesis using codons optimized for human/mammalian expression. The sequences were generated from a known Her2/neu binding Ab (Carter P. et al. (1992) Humanization of an anti P185 Her2 antibody for human cancer therapy. *Proc Natl Acad Sci* **89**, 4285.) and the Fc was an IgG1 isotype (SEQ ID NO:1). The final gene products were sub-cloned into the mammalian expression vector pTT5 (NRC-BRI, Canada) (Durocher, Y., Perret, S. & Kamen, A. High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human HEK293-EBNA1 cells. *Nucleic acids research* **30**, E9 (2002)). The mutations in the CH3 domain were introduced via site-directed mutagenesis of the pTT5 template vectors. See *Table 1* and *Table 6* and *Table 7* for a list of the variant CH3 domain mutations made.

In order to estimate the formation of heterodimers and determine the ratio of homodimers vs. heterodimers the two heterodimer heavy chains were designed with C-terminal extensions of different size (specifically, chain A with C-terminal HisTag and chain B with C-terminal mRFP plus StrepTagII). This difference in molecular weight allows differentiation of homodimers vs. heterodimer in non-reducing SDS-PAGE as illustrated in *FIGURE 25A*.

The HEK293 cells were transfected in exponential growth phase (1.5 to 2 million cells/mL) with aqueous 1mg/mL 25kDa polyethylenimine (PEI, Polysciences) at a PEI:DNA ratio of 2.5:1. (Raymond C. et al. A simplified polyethylenimine-mediated transfection process for large-scale and high-throughput applications. *Methods*. **55**(1):44-51 (2011)). In order to determine the optimal concentration range for forming heterodimers, the DNA was transfected in three separate ratios of the two heavy chains. For example, this was done in 2ml culture volume and transfection DNA, comprised of 5% GFP, 45% salmon sperm

DNA, 25% light chain and 25% total heavy chains, where the heavy chain A plasmid (with C-terminal His-Tag) and the heavy chain B plasmid (with C-terminal StrepTagII plus RFP) at 65%/55%/35% or 10%/20%/40%) were sampled at 3 different relative ratios (chain_A(His)/chain_B(mRFP)) of 10%/65%; 20%/55%; 40%/35% (the apparent 1:1 expression ratio of a WT_His/WT_mRFP heterodimer was determined to be close to the DNA ratio 20%/55%). At 4 to 48 hours after transfection in F17 serum-free media (Gibco), TN1 peptone is added to a final concentration of 0.5%. Expressed antibody was analyzed by SDS-PAGE to determine the best ratio of heavy to light chain for optimal heterodimer formation (See figure 25B and C).

A selected DNA ratio, for example 50% light chain plasmid, 25% heavy chain A plasmid, 25% heavy chain B of AZ33 and AZ34, with 5% GFP, and 45% salmon sperm DNA was used to transfect 150mL of cell culture as described above. Transfected cells were harvested after 5-6 days with the culture medium collected after centrifugation at 4000rpm and clarified using a 0.45µm filter. See *Table 2* below, for a list of the percentage of light and heavy chain A and B plasmids used in the scale up transfection assays for each of the antibodies with CH3 mutations generated for further analysis, including determination of purity and melting temperature.

Table 2:

Variant	LC/HCA/HC B	Variant	LC/HCA/HC B	Variant	LC/HCA/HC B	Variant	LC/HCA/HC B
Wild-Type	50%,50%	AZ47	50%,25%,25%	AZ77	40%,20%,40%	AZ98	50%,20%,30%
AZ12	50%,25%,25%	AZ48	40%,25%,35%	AZ78	50%,20%,30%	AZ100	50%,20%,30%
AZ14	50%,25%,25%	AZ49	50%,25%,25%	AZ79	25%,35%,40%	AZ101	50%,20%,30%
AZ15	50%,25%,25%	AZ63	50%,20%,30%	AZ81	25%,35%,40%	AZ106	25%,35%,40%
AZ17	50%,25%,25%	AZ64	50%,20%,30%	AZ82	50%,20%,30%	AZ114	25%,20%,55%
AZ19	50%,25%,25%	AZ65	50%,20%,30%	AZ83	50%,20%,30%	AZ115	25%,20%,55%
AZ20	50%,25%,25%	AZ66	50%,20%,30%	AZ84	50%,20%,30%	AZ122	25%,20%,55%
AZ21	50%,25%,25%	AZ67	50%,20%,30%	AZ85	50%,25%,25%	AZ123	40%,20%,40%
AZ25	50%,25%,25%	AZ68	50%,20%,30%	AZ86	40%,15%,45%	AZ124	40%,20%,40%
AZ29	50%,25%,25%	AZ69	50%,20%,30%	AZ87	50%,25%,25%	AZ129	40%,30%,30%
AZ30	50%,25%,25%	AZ70	50%,20%,30%	AZ88	50%,25%,25%	AZ130	40%,30%,30%
AZ32	50%,25%,25%	AZ71	40%,20%,40%	AZ89	40%,15%,45%		
AZ33	50%,25%,25%	AZ72	40%,20%,40%	AZ91	50%,25%,25%		
AZ34	50%,25%,25%	AZ73	40%,20%,40%	AZ92	40%,20%,40%		
AZ42	50%,25%,25%	AZ74	40%,20%,40%	AZ93	40%,20%,40%		
AZ44	50%,25%,25%	AZ75	40%,20%,40%	AZ94	50%,25%,25%		
AZ46	50%,25%,25%	AZ76	40%,20%,40%	AZ95	50%,20%,30%		

Example 2: Purification of bivalent monospecific Antibodies with Heterodimer Fc domains.

The clarified culture medium was loaded onto a MabSelect SuRe (GE Healthcare) protein-A column and washed with 10 column volumes of PBS buffer at pH 7.2. The antibody was eluted with 10 column volumes of citrate buffer at pH 3.6 with the pooled fractions containing the antibody neutralized with TRIS at pH 11. The protein was finally desalted using an Econo-Pac 10DG column (Bio-Rad). The C-terminal mRFP tag on the heavy chain B was removed by incubating the antibody with enterokinase (NEB) at a ratio of 1:10,000 overnight in PBS at 25°C. The antibody was purified from the mixture by gel filtration. For gel filtration, 3.5mg of the antibody mixture was concentrated to 1.5mL and loaded onto a Sephadex 200 HiLoad 16/600 200 pg column (GE Healthcare) via an AKTA Express FPLC at a flow-rate of 1mL/min. PBS buffer at pH 7.4 was used at a flow-rate of 1mL/min. Fractions corresponding to the purified antibody were collected, concentrated to ~1mg/mL and stored at -80°C.

Formation of heterodimers, as compared to homodimers, was assayed using non-reducing SDS-PAGE and mass spectrometry. Protein A purified antibody was run on a 4-12% gradient SDS-PAGE, non-reducing gel to determine the percentage of heterodimers formed prior to enterokinase (EK) treatment (See, Figure 26). For mass spectrometry, all Trap LC/MS (ESI-TOF) experiments were performed on an Agilent 1100 HPLC system interfaced with a Waters Q-TOF2 mass spectrometer. Five µg of gel filtration purified antibody was injected into a Protein MicroTrap (1.0 by 8.0 mm), washed with 1% acetonitrile for 8 minutes, a gradient from 1 to 20% acetonitrile/0.1% formic acid for 2 minutes, then eluted with a 20 to 60% acetonitrile/0.1% formic acid gradient for 20 minutes. Eluate (30-50µL/min) was directed to the spectrometer with spectrum acquired every second (m/z 800 to 4,000). (See, Figure 28) Variants having greater than 90% heterodimers were selected for further analysis, with the exception of AZ12 and AZ14 which each had greater than 85% heterodimer formation.

Example 3: Stability determination of bivalent monospecific antibodies with Heterodimer Fc domains using Differential Scanning Calorimetry (DSC).

All DSC experiments were carried out using a GE VP-Capillary instrument. The proteins were buffer-exchanged into PBS (pH 7.4) and diluted to 0.4 to 0.5mg/mL with 0.137mL loaded into the sample cell and measured with a scan rate of 1°C/min from 20 to 100°C. Data was analyzed using the Origin software (GE Healthcare) with the PBS buffer background subtracted. (See, Figure 27). See Table 3 for a list of variants tested and a melting temperature determined. See Table 4 for a list of the variants with a melting temperature of 70°C and above and the specific T_m for each variant.

Table 3: Melting temperature measurements of variant CH3 domains in an IgG1 antibody having 90% or more heterodimer formation compared to homodimer formation

Variant	T _m °C	Variant	T _m °C	Variant	T _m °C	Variant	T _m °C
Wild-Type	81	AZ29	70	AZ63	71.5	AZ87	71
Control 1	69	AZ30	71	AZ64	74	AZ88	72
Control 2	69	AZ31	68	AZ65	73	AZ89	72.5
AZ3	65	AZ32	71.5	AZ66	72.5	AZ91	71.5
AZ6	68	AZ33	74	AZ67	72	AZ92	71.5
AZ8	68	AZ34	73.5	AZ68	72	AZ93	71.5
AZ12	77	AZ38	69	AZ69	71	AZ94	73.5
AZ14	77	AZ42	70	AZ70	75.5	AZ95	72
AZ15	71.5	AZ43	67	AZ71	71	AZ98	70
AZ16	68.5	AZ44	71.5	AZ72	70.5	AZ99	69
AZ17	71	AZ46	70.5	AZ73	71	AZ100	71.5
AZ18	69.5	AZ47	70.5	AZ74	71	AZ101	74

AZ19	70.5	AZ48	70.5	AZ75	70	AZ106	74
AZ20	70	AZ49	71	AZ76	71.5	AZ114	71
AZ21	70	AZ50	69	AZ77	71	AZ115	70
AZ22	69	AZ52	68	AZ78	70	AZ117	69.5
AZ23	69	AZ53	68	AZ79	70	AZ122	71
AZ24	69.5	AZ54	67	AZ81	70.5	AZ123	70
AZ25	70.5	AZ58	69	AZ82	71	AZ124	70
AZ26	69	AZ59	69	AZ83	71	AZ125	69
AZ27	68	AZ60	67	AZ84	71.5	AZ126	69
AZ28	69.5	AZ61	69	AZ85	71.5	AZ129	70.5
		AZ62	68	AZ86	72.5	AZ130	71

Table 4: Melting temperature measurements of select variant CH3 domains in an IgG1 antibody

Variant	T _m °C	Variant	T _m °C	Variant	T _m °C	Variant	T _m °C
Wild-Type	81.5	AZ42	70	AZ73	71	AZ91	71.5
Control 1	69	AZ44	71.5	AZ74	71	AZ92	71.5
Control 2	69	AZ46	70.5	AZ75	70	AZ93	71.5
AZ12	>77	AZ47	70.5	AZ76	71.5	AZ94	73.5
AZ14	>77	AZ48	70.5	AZ77	71	AZ95	72
AZ15	71.5	AZ49	71	AZ78	70	AZ98	70

AZ17	71	AZ63	71.5	AZ79	70	AZ100	71.5
AZ19	70.5	AZ64	74	AZ81	70.5	AZ101	74
AZ20	70	AZ65	73	AZ82	71	AZ106	74
AZ21	70	AZ66	72.5	AZ83	71	AZ114	71
AZ25	70.5	AZ67	72	AZ84	71.5	AZ115	70
AZ29	70	AZ68	72	AZ85	71.5	AZ122	71
AZ30	71	AZ69	71	AZ86	72.5	AZ123	70
AZ32	71.5	AZ70	75.5	AZ87	71	AZ124	70
AZ33	74	AZ71	71	AZ88	72	AZ129	70.5
AZ34	73.5	AZ72	70.5	AZ89	72.5	AZ130	71

Example 4: Evaluation of FcγR Binding using Surface Plasmon Resonance

All binding experiments were carried out using a BioRad ProteOn XPR36 instrument at 25°C with 10mM HEPES, 150mM NaCl, 3.4mM EDTA, and 0.05% Tween 20 at pH

7.4. Recombinant HER-2/neu(p185, ErbB-2 (eBiosciences, Inc.)) was captured on the activated GLM sensorchip by injecting 4.0μg/mL in 10mM NaOAc (pH 4.5) at 25μL/min until approx. 3000 resonance units (RUs) were immobilized with the remaining active groups quenched. 40μg/mL of purified anti-HER-2/neu antibodies comprising the variant CH3 domains were indirectly captured on the sensorchip by binding the Her-2/neu protein when injected at 25μL/min for 240s (resulting in approx. 500RUs) following a buffer injection to establish a stable baseline. FcγR (CD16a(f allotype) and CD32b) concentrations (6000, 2000, 667, 222, and 74.0nM) were injected at 60μL/min for 120s with a 180s dissociation phase to obtain a set of binding sensograms. Resultant K_D values were determined from binding isotherms using the Equilibrium Fit model with reported values as the mean of three independent runs. Comparisons were made with the wild-type IgG1 Fc domain and binding is expressed as a ratio of the WT K_D to the variant K_D (See, Table 5).

Table 5: Ratio of K_D wild-type IgG1 to variant CH3 domain antibody binding independently to CD16a and CD32b

Variant	CD16a Ratio WT/Variant	CD32b Ratio WT/Variant	Variant	CD16a Ratio WT/Variant	CD32b Ratio WT/Variant
Control 1	1.28	1.68	AZ64	0.95	0.9
Control 2	1.1	1.13	AZ65	0.93	0.9
AZ3	1.75	1.87	AZ66	1.26	1.19
AZ6	1.38	1	AZ67	1.21	1.13
AZ8	1.75	1.64	AZ68	1.02	1.1
AZ12	N/A	N/A	AZ69	0.96	1.05

AZ14	N/A	N/A	AZ70	1.06	1.11
AZ15	0.72	0.59	AZ71	0.89	0.95
AZ16	0.95	0.64	AZ72	1.04	1.02
AZ17	2.28	2.37	AZ73	1.09	1.07
AZ18	1.53	1.7	AZ74	1.25	1.17
AZ19	1.55	1.89	AZ75	1.34	1.22
AZ20	2.56	1.93	AZ76	0.99	1
AZ21	2.41	3.28	AZ77	1	1.08
AZ22	2.02	2.37	AZ78	0.9	1
AZ23	1	2.16	AZ79	1.01	0.8
AZ24	1.79	2.26	AZ81	1.01	0.84
AZ25	2.02	2.37	AZ82	0.97	0.94
AZ26	2.38	2.59	AZ83	0.94	0.94
AZ27	2.27	2.38	AZ84	0.93	1
AZ28	1.45	2.15	AZ85	1.01	1.14
AZ29	1.62	2.13	AZ86	1.22	1.18
AZ30	1.61	2.38	AZ87	1.03	1.1
AZ31	1.63	2.29	AZ88	1.11	1.15
AZ32	1.82	2.48	AZ89	1.12	1.24
AZ33	1.91	1.89	AZ91	1.11	1.11

AZ34	1.88	1.88	AZ92	1.21	1.24
AZ38	1.78	1.44	AZ93	1.21	1.18
AZ42	1.28	1.09	AZ94	1.17	1.19
AZ43	1.63	1.73	AZ95	0.86	0.96
AZ44	2.76	3.07	AZ98	0.79	0.82
AZ46	2.16	2.66	AZ99	1.16	1.15
AZ47	1.76	2.12	AZ100	1.13	1.12
AZ48	2.02	1.59	AZ101	1.24	1.23
AZ49	2.09	2.9	AZ106	0.76	0.64
AZ50	2.33	1.86	AZ114	1.3	0.84
AZ52	1.55	1.5	AZ115	1.13	0.82
AZ53	1.87	1.27	AZ117	0.89	1
AZ54	1.36	1.64	AZ122	0.89	0.92
AZ58	2.33	1.48	AZ123	0.85	0.92
AZ59	1.18	1.57	AZ124	0.99	1.09
AZ60	1.51	1.23	AZ125	1	1
AZ61	1.41	1.75	AZ126	0.86	0.9
AZ62	1.53	1.88	AZ129	1.91	2.57
AZ63	0.9	0.95	AZ130	1.91	2.54

Example 5: Rational design of heteromultimers using Fc_CH3 engineering – Scaffold 1 (1a and 1b) and the development of AZ17-62 and AZ133-AZ2438

To improve the initial AZ8 for stability and purity, the structural and computational strategies described above were employed. (See, Figure 24) For example, the in depth structure-function analysis of AZ8 provided a detailed understanding for each of the introduced mutations of AZ8, L351Y_V397S_F405A_Y407V / K392V_T394W compared to wild-type human IgG1 and indicated that the important core heterodimer mutations were L351Y_F405A_Y407V / T394W, while V397S, K392V were not relevant for heterodimer formation. The core mutations (L351Y_F405A_Y407V / T394W) are herein referred to as "Scaffold 1" mutations. The analysis furthermore revealed that the important interface hotspots that are lost with respect to wild-type (WT) homodimer formation are the interactions of WT-F405-K409, Y407-T366 and the packing of Y407-Y407 and -F405 (See, Figure 29). This was reflected in the packing, cavity and MD analysis, which showed a large conformational difference in the loop region D399-S400-D401 (See, Figure 30) and the associated β -sheets at K370. This resulted in the loss of the interchain interactions K409-D399 (See, Figure 30) and weakening of the strong K370 hydrogen bond to E357 (K370 is no longer in direct contact with S364 and E357, but is entirely solvent exposed). In the WT IgG1 CH3 domain these regions tether the interface at the rim protects the core interactions from bulk solvent competition and increases the dynamic occurrence of favorable hydrophobic van der Waals interactions. The consequence was a lower buried surface area of AZ8 compared to WT and a higher solvent accessibility of the hydrophobic core. This indicated the most important factors for the lower stability of AZ8 compared to WT stability was a) the loss of the WT-F405-K409 interaction and packing of F405, and b) the loss of the strong packing interaction of Y407-Y407 and Y407-T366. See, Figure 29

Consequently, we identified the key residues/sequence motifs responsible for the low stability of AZ8 compared to WT. To improve the stability and heterodimer specificity of AZ8 the subsequent positive design engineering efforts were therefore specifically focused on stabilizing the loop conformation of positions 399-401 in a more 'closed' - WT like conformation (See, Figure 30) and compensating for the overall slightly decreased (looser) packing of the hydrophobic core at positions T366 and L368 (See, Figure 29).

To achieve this stabilization of the loop conformation of positions 399-401 the described computational approach was used to evaluate our different targeted design ideas. Specifically, three different independent options for Heteromultimer AZ8 were analyzed to optimize the identified key regions for improving stability. First, the cavity close to position K409 and F405A was evaluated for better hydrophobic packing to both protect the hydrophobic core and stabilize the loop conformation of 399-400 (See, Figure 30). Those included, but were not limited to additional point mutations at positions F405 and K392. Second, options for improving the electrostatic interactions of positions 399-409 were evaluated, to stabilize the loop conformation of 399-400 and protect the hydrophobic core. This included, but was not limited to additional point mutations at positions T411 and S400. Third, the cavity at the core packing positions T366, T394W and L368 was evaluated to improve the core hydrophobic packing (See, Figure 29). Those included, but were not limited to additional point mutations at positions T366 and L368. The different independent positive design ideas were tested *in-silico* and the best-ranked variants using the computational tools (AZ17-AZ62) were validated experimentally for expression and stability as described in Examples 1-4. See Table 4 for a list of heteromultimers from this design phase with a melting temperature of 70°C or greater.

Heteromultimer AZ33 is an example of the development of an Heteromultimer wherein Scaffold 1 was modified resulting in Scaffold 1a mutations to improve stability and purity. This Heteromultimer was designed based on AZ8 with the goal of improving the hydrophobic packing at positions 392-394-409 and 366 to both protect the hydrophobic core and stabilize the loop conformation of 399-400. This Heteromultimer AZ33 heterodimer has two additional point mutations different from the core mutations of AZ8, K392M and T366I. The mutations T366I_K392M_T394W/F405A_Y407V are herein referred to as "Scaffold 1a" mutations. The mutation K392M was designed to improve the packing at the cavity close to position K409 and F405A to protect the hydrophobic core and stabilize the loop conformation of 399-400 (See, Figure 31). T366I was designed to improve the core hydrophobic packing and to eliminate the formation of homodimers of the T394W chain (See, Figure 29). The experimental data for AZ33 showed significantly improved stability over the initial negative design Heteromultimer AZ8 (T_m 68°C) wherein AZ33 has a T_m of 74°C and a heterodimer content of >98%. (See, Figure 25C)

Development of heteromultimers using Scaffold 1 mutations in phase three design of Heteromultimer heterodimers

Although AZ33 provides a significant stability and specificity (or purity) improvement over the initial starting variant AZ8, our analysis indicates that further improvements to the stability of the heterodimer can be made with further amino acid modifications using the experimental data of AZ33 and the above described design methods. The different design ideas have been independently tested for expression and stability, but the independent design ideas are transferable and the most successful heterodimer will contain a combination of the different designs. Specifically, for the optimization of AZ8 packing mutations at the cavity close to K409-F405A-K392 have been evaluated independently from mutations that optimize the core packing at residues L366T-L368. These two regions 366-368 and 409-405-392 are distal from each other and are considered independent. Heteromultimer AZ33 for example has been optimized for packing at 409-405-392, but not at 366-368, because these optimization mutations were separately evaluated. The comparison of the 366-368 mutations suggests that T366L has an improved stability over T366 and also T366I, the point mutation used in the development of Heteromultimer AZ33. Consequently, the presented experimental data immediately suggest further optimization of AZ33 by introducing T366L instead of T366I, for example. Therefore, the amino acid mutations in the CH3 domain T366L_K392M_T394W/F405A_Y407V are herein referred to as "Scaffold 1b" mutations.

In a similar manner the complete experimental data has been analyzed to identify point mutations that can be used to further improve the current Fc variant heterodimer AZ33. These identified mutations were analyzed by the above described computational approach and ranked to yield the list of additional Fc variant heterodimers based on AZ33 as shown in Table 6.

Example 6: Rational design of heteromultimers using Fc_CH3 engineering – Scaffold 2 (a and b) and, the development of AZ63-101 and AZ2199-AZ2524

To improve the initial negative design phase Heteromultimer AZ15 for stability and purity,

the structural and computational strategies described above were employed (See, Figure 24). For example, the in depth structure-function analysis of Heteromultimer AZ15 provided a detailed understanding for each of the introduced mutations of AZ15, L351Y_Y407A / E357L_T366A_K409F_T411N compared to wild-type (WT) human IgG1 and indicated that the important core heterodimer mutations were L351Y_Y407A / T366A_K409F, while E357L, T411N were not directly relevant for heterodimer formation and stability. The core mutations (L351Y_Y407A / T366A_K409F) are herein referred to as "Scaffold 2" mutations. The analysis furthermore revealed that the important interface hotspots that are lost with respect to wild-type (WT) homodimer formation are the salt bridge D399-K409, the hydrogen bond Y407-T366 and the packing of Y407-Y407. Our detailed analysis, provided below, describes how we improved the stability of our original Heteromultimer AZ15 and the positions and amino acid modifications made to achieve these heteromultimers with improved stability.

Development of heteromultimers using Scaffold 2 mutations and the further development of Scaffold 2a mutations.

Our in-silico analysis indicated a non-optimal packing of the Heteromultimer AZ15 mutations K409F_T366A_Y407A and an overall decreased packing of the hydrophobic core due to the loss of the WT-Y407-Y407 interactions. The positive design efforts in the subsequent engineering phase were focused on point mutations to compensate for these packing deficits in the initial Fc variant AZ15. The targeted residues included positions T366, L351, and Y407. Different combinations of these were tested in-silico and the best-ranked Fc variants using the computational tools (AZ63-AZ70) were validated experimentally for expression and stability as described in Examples 1-4.

Fc variant AZ70 is an example of the development of a Heteromultimer wherein Scaffold 2 was modified resulting in Scaffold 2a mutations to improve stability and purity. This Heteromultimer was designed based on AZ15 with the goal of achieving better packing at

the hydrophobic core as described above. Heteromultimer AZ70 has the same Scaffold 2 core mutations (L351Y_Y407A / T366A_K409F) as described above except that T366 was mutated to T366V instead of T366A (FIGURE 33). The L351Y mutation improves the 366A_409F/407A variant melting temperature from 71.5°C to 74°C, and the additional change from 366A to 366V improves the T_m to 75.5°C. (See, AZ63, AZ64 and AZ70 in Table 4, with a T_m of 71.5°C, 74°C and 75.5°C, respectively) The core mutations (L351Y_Y407A / T366V_K409F) are herein referred to as "Scaffold 2a" mutations. The experimental data for Fc variant AZ70 showed significantly improved stability over the initial negative design Fc variant AZ15 (T_m 71°C) wherein AZ70 has a T_m of 75.5°C and a heterodimer content of >90% (FIGURE 33 and 27).

Development of heteromultimers using Scaffold 2 mutations and the further development of Scaffold 2b mutations.

The Molecular Dynamics simulation (MD) and packing analysis showed a preferred more 'open' conformation of the loop 399-400, which was likely due to the loss of the WT salt bridge K409-D399. This also results in the unsatisfied D399, which in turn preferred a compensating interaction with K392 and induced a more 'open' conformation of the loop. This more 'open' loop conformation results in an overall decreased packing and higher solvent accessibility of the core CH3 domain interface residues, which in turn significantly destabilized the heterodimer complex. Therefore, one of the targeted positive design efforts was the tethering of this loop in a more 'closed', WT-like conformation by additional point mutations that compensate for the loss of the D399-K409 salt bridge and the packing interactions of K409. The targeted residues included positions T411, D399, S400, F405, N390, K392 and combinations thereof. Different packing, hydrophobic- and electrostatic positive engineering strategies were tested in silico with respect to the above positions and the best-ranked heteromultimers determined using the computational tools (AZ71-AZ101) were validated experimentally for expression and stability as described in Examples 1-4.

Heteromultimer AZ94 is an example of the development of an Fc variant wherein Scaffold 2 is modified resulting in Scaffold 2b mutations along with additional point mutations to improve stability and purity. This Fc variant was designed based on AZ15 with the goal of tethering loop 399-400 in a more 'closed', WT-like conformation and compensating for the loss of the D399-K409 salt bridge as described above. Fc variant AZ94 has four additional point mutations to Scaffold 2 (L351Y_Y407A / T366A_K409F) and returns L351Y to wild-type L351 leaving (Y407A / T366A_K409F) as the core mutations for this Fc variant. The core mutations Y407A / T366A_K409F are herein referred to as "Scaffold 2b" mutations. The four additional point mutations of AZ94 are K392E_T411E / D399R_S400R. The mutations T411E / D399R were engineered to form an additional salt bridge and compensate for the loss of the K409 / D399 interaction (FIGURE 34). Additionally, this salt bridge was designed to prevent homodimer formation by disfavoring charge-charge interactions in both potential homodimers. The additional mutations K392E / S400R were intended to form another salt bridge and hence further tether the 399_400 loop in a more 'closed', WT-like conformation (FIGURE 34). The experimental data for AZ94 showed improved stability and purity over the initial negative design Fc variant AZ15 (T_m 71°C, >90% purity) wherein Fc variant AZ94 has a T_m of 74°C and a heterodimer content or purity of >95% .

Development of heteromultimers using Scaffold 2 mutations in phase three design of heterodimers

Both Fc variants AZ70 and AZ94 provide a significant improvement in stability and purity over the initial negative design Fc variant AZ15, but our analysis and the comparison of AZ70 and AZ94 directly indicate that further improvements to the stability of the Fc variant heterodimer can be made with further amino acid modifications. For example, Fc variants AZ70 and AZ94 were designed to target two distinct non-optimized regions in the initial variant AZ15, which was accomplished by improving the packing at the hydrophobic

core and making mutations outside of the core interface residues resulting in additional salt bridges and hydrogen bonding to stabilize the loop conformation of positions 399-401. The additional point mutations of Fc variants AZ70 and AZ94 are distal from each other and are therefore independent and transferable to other Fc variants designed around the same Scaffold 2 core mutations, including 2a and 2b mutations. Specifically, AZ70 only carries the optimized core mutations L351Y_Y407A / T366A_K409F, but no additional salt bridges, whereas AZ94 comprises four additional electrostatic mutations (K392E_T411E / D399R_S400R), but has one less mutation in the hydrophobic core interface (Y407A / T366A_K409F). These Scaffold 2b mutations are less stable than AZ70 (See, for example AZ63, which has equivalent core mutations as AZ94 and T_m of 72°C), but are compensated for by the addition of K392E_T411E / D399R_S400R mutations. The presented experimental stability and purity data indicates that combining the mutations of AZ70, which optimizes the hydrophobic core, and the electrostatic mutations of AZ94 should further improve stability and purity of the heterodimers that comprise the Fc variants. In a similar manner the complete experimental data for Scaffold 2 Fc variants (AZ63-101) has been analyzed to identify point mutations that can be used to further improve the Fc variant heterodimers AZ70 and AZ94. These identified mutations were further analyzed by the above described computational approach and ranked to yield the list of additional Fc variant heterodimers based on AZ70 and AZ94 as shown in Table 7.

Example 7: Effect of Heterodimeric CH3 on FcγR binding

As a prototypical example of heterodimeric Fc activity with FcγR, we have tested two variant antibodies with heterodimeric Fc region A:K409D_K392D/B:D399K_D356K (Control 1 (het 1 in Figure 35)) and A:Y349C_T366S_L368A_Y407V/B:S354C_T366W (Control 4 (het 2 in Figure 35)) with Her2 binding Fab arms in an SPR assay described in Example 4 for FcγR binding. As shown in Figure 35, we observe that both the heterodimeric Fc regions bind the different Fcγ receptors with the same relative strength as the wild type IgG1 Fc region, but overall, the heterodimeric Fc region bound each of the FcγR's slightly better than the wild type antibody. This indicates that mutations

at the CH3 interface of Fc can impact the binding strength of the Fc region for Fcγ receptors across the CH2 domains as observed in our molecular dynamics simulations and analysis.

Example 8: Effect of Asymmetric mutations in CH2 of a heterodimeric Fc on FcγR binding

Mutation of Serine at position 267 in the CH2 domain of the Fc region to an Aspartic acid (S267D) is known to enhance binding to FcγIIbF, IIbY & IIaR receptors when introduced in a homodimeric manner in the two chains of CH2 domain. This mutation can be introduced on only one of the CH2 domains in an heterodimeric Fc molecule to gain roughly half the improvement in binding strength relative to when this mutation is introduced in a homodimeric CH2 Fc as the data presented in Figure 36A indicates. On the other hand, the E269K mutation in a homodimeric CH2 domain of Fc prevents binding of the Fc region to FcγR. We present a scheme for enhanced manipulation of the binding strength of the Fc region for the FcγReceptors by the asymmetric introduction of these favorable and unfavorable mutations on one of the two chains in the CH2 domain of the Fc. The introduction of E269K mutation in an asymmetric manner on one CH2 chain in a heterodimeric Fc acts as a polarity driver by blocking binding of the FcγR at the face where it is present, while letting the other face of the Fc interact with the FcγR in a normal manner. The results from this experimentation are presented in Figure 36A. The opportunity to selectively alter the binding strength via both the chains of Fc in an independent manner provides increased opportunity to manipulate the binding strength and selectivity between Fc and FcγReceptors. Thus, such asymmetric design of mutations in the CH2 domain allows us to introduce positive and negative design strategies to favor or disfavor certain binding models, providing greater opportunity to introduce selectivity.

In a subsequent experiment, we have altered the selectivity profile of the base Fc mutant S239D_D265S_I332E_S298A that shows increased binding strength to the FcγIIIaF and IIIaV receptors while continuing to exhibit weaker binding to the FcγIIaR, IIbF and IIbY receptors. This is shown in the binding profile shown in Figure 36B. By introducing asymmetric mutations E269K in chain A and avoiding the I332E mutation in chain B, we are able to generate a novel FcγR binding profile that further weakens IIa and IIb receptor binding and makes the Fc more specific for the IIIa receptor binding.

In another example shown in Figure 36C, asymmetric mutations are highlighted relative to the homodimeric Fc involving the mutation S239D/K326E/A330L/I332E/S298A in the CH2 domain. Relative to the wild type IgG1 Fc, this variant show increased binding to the IIIa receptor but also binds the IIa and IIb receptors slightly stronger than the wild type Fc. Introduction of these mutations in an asymmetric manner A:S239D/K326E/A330L/I332E and B:S298A while reducing the IIIa binding, also increases the IIa/IIb receptor binding, loosing selectivity in the process. By introducing an asymmetric E269K mutation in this heterodimeric variant, i.e. A:S239D/K326E/A330L/I332E/E269K and B:S298A, we are able to reduce the IIa/IIb binding back to wild type levels. This highlights the fact that the use of asymmetric mutations in the CH2 domain of Fc is able to provide significant opportunity to design improved FcγR selectivity.

The reagents employed in the examples are commercially available or can be prepared using commercially available instrumentation, methods, or reagents known in the art. The foregoing examples illustrate various aspects of the invention and practice of the methods of the invention. The examples are not intended to provide an exhaustive description of the many different embodiments of the invention. Thus, although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, those of ordinary skill in the art will realize readily that many changes and modifications can be made thereto without departing from the spirit or scope of the appended claims.

Example 9: FcRn binding determined by SPR.

Binding to FcRn was determined by SPR in two different orientations.

1. Flowing of the heterodimer variant over immobilized FcRn: In this experiment, high density surfaces approx 5000 RUs were made using standard NHS/EDC coupling. 100nM of WT and each variant was injected in triplicate at 50 uL min for 120s with 600s dissociation in MES pH6 running buffer.
2. Flowing of FcRn over indirectly captured heterodimer variants: In this SPR experiment, a goat anti-human IgG surface was used to indirectly capture the antibodies (approximately 400RUs each), followed by an injection of a 3-fold FcRn dilution series (6000nM high conc). Running buffer was 10mM MES / 150mM NaCl / 3.4 mM EDTA / 0.05

Tween20 at pH6. There was no significant binding of FcRn to the goat polyclonal surface. All variants show similar to WT sensograms. Table 8 below shows the Kd determined by the indirect immobilization with flowing FcRn (2.).

Kd [M] – pH6.0	Kd [M] – pH7.5	Mutations (Chain-A)	Mutations (Chain-B)
3.7E-06	-	Herceptin WT	
4.E-06	-	L351Y_F405A_Y407V	T366I_K392M_T394W
5.E-06	-	L351Y_F405A_Y407V	T366L_K392M_T394W
4.3E-06	-	T350V_L351Y_F405A_Y407V	T350V_T366L_K392M_T394W
4.1E-06	-	Y349C_T350V_F405A_Y407V	T350V_S354C_T366L_K392M_T394W
5.E-06	-	T350V_L351Y_S400E_F405A_Y407V	T350V_T366L_N390R_K392M_T394W
3.9E-06	-	T350V_L351Y_F405A_Y407V	T350V_T366L_K392L_T394W

Example 10: Preparation of an exemplary heteromultimers

The following heteromultimer comprising one single domain antigen-binding construct was prepared:

1. v1323, a monovalent anti-EGFR antibody (EG2), where the EGFR binding domain is a camelid V_HH on chain A, and the Fc region is a heterodimer having the mutations T350V_L351Y_F405A_Y407V in Chain A, and T350V_T366L_K392L_T394W in Chain B

This construct was prepared and expressed as follows.

The genes encoding the antibody heavy chains and Fc regions were constructed via gene synthesis using codons optimized for human/mammalian expression. The sdAb sequence encoding the anti-EGFR sdAb was generated from a known EGFR binding antibody EG2 (Bell et al. (2010) Differential tumor-targeting abilities of three single-domain antibody formats. *Cancer Letters* 289:81).

The final gene products were sub-cloned into the mammalian expression vector pTT5 (NRC-BRI, Canada) and expressed in CHO cells (Durocher, Y., Perret, S. & Kamen,

A. High-level and high-throughput recombinant protein production by transient transfection of suspension-growing CHO cells. Nucleic acids research 30, E9 (2002)).

The CHO cells were transfected in exponential growth phase (1.5 to 2 million cells/mL) with aqueous 1mg/mL 25kDa polyethylenimine (PEI, Polysciences) at a PEI:DNA ratio of 2.5:1.(Raymond C. et al. A simplified polyethylenimine-mediated transfection process for large-scale and high-throughput applications. Methods. 55(1):44-51 (2011)). The DNA was transfected in optimal DNA ratios of the chain A (CHA) and chain B (CHB) to allow for heterodimer formation (e.g. CHA/CHB ratio = 50:50) Transfected cells were harvested after 5-6 days with the culture medium collected after centrifugation at 4000rpm and clarified using a 0.45µm filter.

Example 11: Purification and characterization of an exemplary heteromultimer

The heteromultimer described in Example 1 was expressed and purified by protein A chromatography as described below.

Protein A purification

The clarified culture medium was loaded onto a MabSelect SuRe (GE Healthcare) protein-A column and washed with 10 column volumes of PBS buffer at pH 7.2. The antibody was eluted with 10 column volumes of citrate buffer at pH 3.6 with the pooled fractions containing the antibody neutralized with TRIS at pH 11.

UPLC-SEC analysis of heterodimer purity post Protein-A purification

The purity of v1323 was determined using UPLC-SEC under standard conditions described below:

Column: Waters BEH200 SEC, 1.7 µm particles, 4.6 x 150 mm

Solvent: 25 mM NaPO₄, 150 mM NaCl, pH 7.00 @ 23.4°C, 20.00 mS/cm @ 23.4°C

Flow rate: 0.4 ml/min, ~4280 psi

Temperature: 30°C

Samples: v1323 120607-KB

3 injections (~2 µg per injection)

Figures 41 and 42 depict the results of SDS-PAGE and UPLC-SEC analysis, respectively, for the exemplary heteromultimer after Protein-A purification. Figure 41 illustrates the relative purity post Protein-A purification and shows that v1323 contained no detectable contaminant species and did not require additional purification by SEC. Figure 42 contains UPLC-SEC analysis that supports the observations in Figure 41, illustrating that e.g. 1323 is > 97% heterodimer purity post Protein-A purification.

Table 9 provides a summary of the purification procedure and yield for v1323.

Table 9: Yield and Purification process for v1323

variant	purification	Conc. mg/ml	Total mg per 50ml
1323	protA	1.3	1.95 (1.5 ml)

Example 12: Heteromultimers maintain EGFR-ED binding domain properties

The ability of v1323 comprising an EGFR binding domain to bind to the extracellular domain (ED) of EGFR was tested by Surface Plasmon Resonance (SPR) using a ProteOn XPR36 system from BIO-RAD.

Approximately 3000 RU of anti-human IgG 25ug/ml was immobilized on a GLC chip using standard amine coupling. Exemplary SDACs were captured on the anti-human IgG immobilized chip to capture level of approximately 700 RU. Recombinant human EGFR-ED was diluted in running buffer and injected at a flow rate of 50 µl /min for 2 minutes, followed by dissociation for another 4 minutes. Sensograms were fit globally to a 1: 1 Langmuir binding model. All experiments were conducted at room temperature.

The SPR curves for v1323 are shown in Figure 43. The results show that v1323 binds to EGFR-ED with high affinity in the low nanomolar range between 3-4.5 nM.

Example 13: Ability of v1323 to bind to A549, BxPc3 and U87 cells

The ability of v1323 to bind to EGFR expressed on the surface of a cell was tested using A549, BxPc3 and U87 cells by a fluorescent cell binding assay as described below.

Binding of the exemplary bi-specific SDACs to the surface of A549, BxPc3 and U87 cells was determined by flow cytometry. Cells were washed with PBS and resuspended in DMEM at 1×10^5 cells/ 100 μ l. 100 μ l cell suspension was added into each microcentrifuge tube, followed by 10 μ l/ tube of the antibody variants. The tubes were incubated for 2hr 4°C on a rotator. The microcentrifuge tubes were centrifuged for 2min 2000RPM at room temperature and the cell pellets washed with 500 μ l media. Each cell pellet was resuspended 100 μ l of fluorochrome-labelled secondary antibody diluted in media to 2 μ g/ sample. The samples were then incubated for 1hr at 4°C on a rotator. After incubation, the cells were centrifuged for 2 min at 2000 RPM and washed in media. The cells were resuspended in 500 μ l media, filtered in tube containing 5 μ l propidium iodide (PI) and analyzed on a BD LSRII flow cytometer according to the manufacturer's instructions.

The results of this experiment indicated that v1323 was able to bind to the cells tested (data not shown).

All publications, patents and patent applications mentioned in this specification are herein incorporated by reference into the specification to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated herein by reference.

CLAIMS

We Claim:

1. An isolated heteromultimer comprising: at least one single domain antigen-binding construct and an immunoglobulin heterodimer Fc region, said immunoglobulin heterodimer Fc region comprising two monomeric Fc polypeptides, wherein the single domain antigen-binding construct is attached to one monomeric Fc polypeptide;

wherein the heterodimer Fc region comprises a variant CH3 domain comprising amino acid mutations that promote the formation of said heterodimer Fc region with stability comparable to a native homodimeric Fc region; and

wherein said isolated heteromultimer is devoid of immunoglobulin light chain and immunoglobulin first constant (CH1) region.

2. The isolated heteromultimer according to claim 1, comprising one single domain antigen-binding construct attached to one monomeric Fc polypeptide.
3. The isolated heteromultimer according to claim 1 comprising one single domain antigen-binding construct attached to one monomeric Fc polypeptide, and a second single domain antigen-binding construct attached to the other monomeric Fc polypeptide.
4. The isolated heteromultimer according to claim 3, wherein both single-domain antigen-binding constructs bind to the same epitope.
5. The isolated heteromultimer according to claim 3, wherein said one single domain antigen-binding construct binds to one epitope, and the second single domain antigen-binding construct binds to a different epitope.
6. The isolated heteromultimer according to any one of claims 1 to 5, wherein the single domain antigen binding construct is selected from single domain antibodies

(sdAb or VH), camelid nanobodies (V_{H}), cartilaginous fish (V_{NAR}), SH3-derived fynomers, and fibronectin-derived binding domains.

7. The isolated heteromultimer according to any one of claim 1 to 6, wherein the single domain antigen-binding construct is a camelid nanobody (V_{H}).
8. The isolated heteromultimer according to any one of claims 1 to 7, wherein the single domain antigen-binding construct binds to one or more cytokines or chemokines selected from IL2, IFN α -2a/b, IFN-1a/b, IL-21, IL-17a, TNF, IL23, VEGF, and ANG2.
9. The isolated heteromultimer of any one of claims 1 to 7, wherein the single domain antigen-binding construct binds to one or more tumor associated antigens such as EpCam, EGFR, VEGFR, CEA, or GP100.
10. The isolated heteromultimer of any one of claims 1 to 7, wherein the single domain antigen-binding construct binds to one or more immunoregulatory antigens such as CD16, CD30, CD137, CD22, CD52, CD80, CD23, CD2, CD4, CD40, KIR, CD32b, CD25, LAG3, or B7-H3.
11. The isolated heteromultimer of any one of claims 1 to 7, wherein the single domain antigen-binding construct binds to one or more bacterial toxins such as Clostridium difficile toxin A, Clostridium difficile toxin B.
12. The isolated heteromultimer according to any one of claims 1 to 7, wherein the single domain antigen-binding construct binds to EGFR1.
13. The isolated heteromultimer of any one of claims 1 to 7, wherein the single domain antigen-binding construct binds to the EGFR1 mutated variant EGFRVIII.
14. An isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region; wherein the heterodimer Fc region comprises a variant CH3 region comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and wherein said isolated heteromultimer is devoid of immunoglobulin light chains.

15. An isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region;
wherein the heterodimer Fc region comprises a variant constant domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc; and
wherein said isolated heteromultimer is devoid of immunoglobulin light chains.
16. An isolated heteromultimer comprising: at least one single domain antigen-binding construct attached to at least one monomer of a heterodimer Fc region;
wherein the heterodimer Fc region comprises a variant constant domain comprising amino acid mutations that promote the formation of said heterodimer with stability comparable to a native homodimeric Fc region; and
wherein said isolated heteromultimer is devoid of immunoglobulin light chain and immunoglobulin first constant (CH1) region.
17. The isolated heteromultimer of any one of claims 14 to 16 wherein said single domain antigen-binding construct is derived from a camelid or a cartilaginous fish.
18. The isolated heteromultimer of claim 17 wherein said camelid is a llama.
19. The isolated heteromultimer according to any one of claims 1-18, wherein the heterodimer Fc region comprises first and a second Fc polypeptide that form a variant CH3 domain comprising amino acid mutations to promote heterodimer formation with increased stability wherein said amino acid mutations promote the formation of heterodimer Fc region with increased stability as compared to a CH3 domain that does not comprise amino acid mutations, and wherein the variant CH3 domain has a melting temperature (T_m) of about 70 °C or greater.
20. The isolated heteromultimer according to any one of claims 1-19, wherein the heterodimer Fc region does not comprise an additional disulfide bond in the CH3 domain relative to a wild type Fc region.

21. The isolated heteromultimer according to Claim 19 or 20, wherein the heterodimer Fc region comprises an additional disulfide bond in the variant CH3 domain relative to a wild type Fc region, with the proviso that the melting temperature (T_m) of about 70 °C or greater for the CH3 domain is in the absence of the additional disulfide bond.
22. The isolated heteromultimer according to any one of Claims 19-21, wherein the heterodimer Fc region has a purity greater than about 90%.
23. The isolated heteromultimer according to any one of Claims 19-22, wherein the heterodimer Fc region has a purity of about 98% or greater.
24. The isolated heteromultimer according to any one of Claims 19-23, wherein the T_m is about 74 °C or greater.
25. The isolated heteromultimer according to any of Claims 19-24, wherein a first CH3 domain polypeptide comprises amino acid modification at positions F405 and Y407 and a second CH3 domain polypeptide comprises amino acid modification at position T394.
26. The isolated heteromultimer according to any of Claims 19-25 wherein the first Fc polypeptide comprises one or more amino acid modifications selected from L351Y, Y405A and Y407V, and the second Fc polypeptide comprises one or more amino acid modifications selected from T366L, T366I, K392L, K392M and T394W.
27. The isolated heteromultimer according to any of Claims 19-26, wherein a first Fc polypeptide comprises amino acid modifications at positions D399 and Y407 and a second Fc polypeptide comprises amino acid modification at positions K409 and T411.

28. The isolated heteromultimer according to any of Claims 1-27, said heterodimer Fc region comprising:
- a first monomeric Fc polypeptide comprising a first modified CH3 domain comprising at least three amino acid modifications as compared to a wild-type CH3 domain polypeptide, and a second monomeric Fc polypeptide comprising a second modified CH3 domain comprising at least three amino acid modifications as compared to a wild-type CH3 domain polypeptide;
- wherein one of said first and second CH3 domain comprises an amino acid modification of K392J wherein J is selected from L, I, M or an amino acid with a side chain volume not substantially larger than the side chain volume of K; wherein said first and second modified CH3 domain polypeptides preferentially form a heterodimeric CH3 domain with a melting temperature (T_m) of at least about 74°C and a purity of at least 95%; and
- wherein at least one amino acid modification is not of an amino acid which is at the interface between said first and said second CH3 domain polypeptides.
29. The isolated heteromultimer according to any of claims 19-28, comprising at least one T350X modification, wherein X is a natural or non-natural amino acid selected from valine, isoleucine, leucine, methionine, and derivatives or variants thereof.
30. The isolated heteromultimer according to any of claims 19-29, comprising at least one T350V modification.
31. The isolated heteromultimer according to any of claims 19-30, wherein each of said first and second Fc polypeptides further comprises a T350V modification.
32. The isolated heteromultimer according to claim 31, wherein the modified CH3 domain has a T_m of about 77°C or greater.
33. The isolated heteromultimer according to any of claims 19-32, wherein at least one monomeric Fc polypeptide comprises the modification S400Z, wherein Z is selected from a positively charged amino acid and a negatively charged amino acid.

34. The isolated heteromultimer according to claim 33, wherein said first Fc polypeptide comprises an amino acid modification selected from S400E, S400D, S400K and S400R.
35. The isolated heteromultimer according to claim 34, wherein one of said first and second Fc polypeptides comprises the amino acid modification selected from S400E and S400R, and the other Fc polypeptide comprises an amino acid modification at position N390.
36. The isolated heteromultimer according to claim 35, comprising the modification N390Z, wherein Z is selected from a positively charged amino acid and a negatively charged amino acid.
37. The isolated heteromultimer according to claim 35, said second Fc polypeptide comprising the amino acid modification N390R or N390K.
38. The isolated heteromultimer according to any one of claims 19-37, wherein said first Fc polypeptide is a modified CH3 domain polypeptide comprising the amino acid modification S400E and said second Fc polypeptide is a modified CH3 domain polypeptide comprising the amino acid modification N390R.
39. The isolated heteromultimer according to any of claims 1-38, one said Fc polypeptide comprising the amino acid modification Q347R and the other Fc polypeptide comprising the amino acid modification K360E.
40. The isolated heteromultimer according to any of claims 1-39, wherein the Fc polypeptide comprises at least one amino acid modification selected from T366V, T366I, T366A, T366M, T366L, K409F, T411E and T411D, and the second Fc polypeptide comprises at least one amino acid modification selected from L351Y, Y407A, Y407I, Y407V, D399R and D399K.
41. The isolated heteromultimer according to any one of Claims 1-40 wherein the heterodimer Fc region further comprises a variant CH2 domain comprising

asymmetric amino acid modifications to promote selective binding of a Fcgamma receptor.

42. The isolated heteromultimer according to any one of Claims 1-41, wherein the variant CH2 domain selectively binds FcgammaIIIa receptor as compared to wild-type CH2 domain.
43. The isolated heteromultimer according to any of Claims 1-42 comprising a Fc construct based on a type G immunoglobulin (IgG).
44. The isolated heteromultimer according to Claim 43, wherein said IgG is one of IgG1, IgG2 IgG3 and IgG4.
45. The isolated heteromultimer according to any of Claims 1-44 comprising a Fc construct based on Immunoglobulin M (IgM), Immunoglobulin A (IgA), Immunoglobulin D (IgD), or Immunoglobulin E (IgE).
46. The isolated heteromultimer according to any of Claim 1-45, wherein said heteromultimer is a bispecific antibody or a multispecific antibody.
47. The isolated heteromultimer according to any of claims 1-46, wherein at least one single domain antigen binding construct binds EGFR or EGFRvIII.
48. The isolated heteromultimer of claim 47 wherein said EGFR or EGFRvIII binding construct is derived from an antibody or fragment thereof.
49. The isolated heteromultimer of claim 47 wherein said EGFR or EGFRvIII binding construct is a heavy chain antibody construct.
50. The isolated heteromultimer of claim 49, wherein said heavy chain antibody construct is a camelid construct.


51. The isolated heteromultimer of claim 50, wherein said camelid construct comprises the sequence shown in Figure 44.
52. A composition comprising the isolated heteromultimer according to any of claims 1-51 and a pharmaceutically acceptable carrier.
53. A mammalian host cell comprising nucleic acid encoding the isolated heteromultimer according to any of claims 1-51.
54. The isolated heteromultimer according to any one of claims 1-51, wherein the single domain antigen-binding construct competes for binding with at least one therapeutic antibody.
55. The isolated heteromultimer according to claim 54, wherein said at least one therapeutic antibody is selected from the group consisting of abagovomab, adalimumab, alemtuzumab, aurograb, bapineuzumab, basiliximab, belimumab, bevacizumab, briakinumab, canakinumab, catumaxomab, certolizumab pegol, cetuximab, daclizumab, denosumab, efalizumab, galiximab, gemtuzumab, ozogamicin, golimumab, ibritumomab tiuxetan, infliximab, ipilimumab, lumiliximab, mepolizumab, motavizumab, muromonab, mycograb, natalizumab, nimotuzumab, ocrelizumab, ofatumumab, omalizumab, palivizumab, panitumumab, pertuzumab, ranibizumab, reslizumab, rituximab, teplizumab, tocilizumab/atlizumab, tositumomab, trastuzumab, ProxiniumTM, RencarexTM, ustekinumab, and zalutumumab.
56. A method of treating cancer in a patient having a cancer characterized by a cancer antigen, said method comprising administering to said patient a therapeutically effective amount of an isolated heteromultimer of any of claims 1-51.
57. The method of claim 56, wherein said cancer is characterized by overexpression of EGFR or EFGRvIII.

58. A method of treating cancer cells expressing EGFR or EGFRvIII, comprising contacting said cells with an amount of a heteromultimer provided in one of claims 47-51.
59. The method of claim 58, wherein said cancer cell is at least one of a breast cancer cell, a lung cancer cell, an anal cancer cell and a glioblastoma.
60. The method of any of claims 56-59, comprising administration of said heteromultimer, in addition to another therapeutic molecule.
61. The method of claim 60, wherein said therapeutic molecule is conjugated to the heteromultimer.
62. A method of treating immune disorders in a patient having an immune disorder characterized by an immune antigen, said method comprising administering to said patient a therapeutically effective amount of an isolated heteromultimer of any of claims 1-51.

Figure 1




FIGURE 2
AZ1



- L368 faces a critical natural cavity

FIGURE 3
ZYMEWORKS VARIANT: AZ1



- L368 faces a critical natural cavity
- L368R increases stability by improving packing in Region 1

FIGURE 4
ZYMEWORKS VARIANT: AZ1




FIGURE 5

ZYMEWORKS VARIANTS: AZ1

	clash score ($\frac{\text{kcal}}{\text{mol}}$) - wt. +	Δ interface area ($\text{\AA}^2$) - wt. +	Δ packing ($\text{\AA}^2$) - wt. +	Δ electrostatics ($\frac{\text{kcal}}{\text{mol}}$) - wt. +	affinity score - wt. +
Heterodimer	Control 1	-26.2	-46.0	-0.4	12.3
	AZ1	-14.6	9.9	0.1	-12.3
Homodimer	Control (chain A)	-33.6	9.3	-0.5	94.4
	Control (chain B)	-11.5	-77.9	-0.1	69.5
	AZ1 (chain A)	-36.6	23.1	-0.2	87.3
	AZ1 (chain B)	-21.3	-50.8	-0.1	30.8


green

Favorable

red

Unfavorable

FIGURE 6
ZYMEWORKS VARIANT: AZ2, AZ3




- Variants AZ 2 and 3 build on AZ 1
- L351I (AZ2) is designed and favors heterodimer formation by improving packing


FIGURE 7

ZYMEWORKS VARIANT: AZ2, AZ3

L368R_E357R / L368D_K370Q_L351I



L368R_E357R / L368D_K370Q_L351D



● Chain A ● Chain B

- Variants AZ 2 and 3 build on AZ 1
- L351I (AZ2) is designed and favors heterodimer formation by improving packing
- L351D (AZ3) is designed and favors heterodimers formation by improving electrostatic interactions

FIGURE 8

ZYMEWORKS VARIANTS: AZ1, AZ2, AZ3

	clash score (kcal/mol) - wt. +	Δ interface area (Å ²) - wt. +	Δ packing (Å ²) - wt. +	Δ electrostatics (kcal/mol) - wt. +	affinity score - wt. +
Heterodimer	Control	-26.2	-46.0	-0.4	12.3
	AZ1	-14.6	9.9	0.1	-12.3
	AZ2	-3.1	0.7	0.0	-17.2
	AZ3	-12.1	-16.1	0.1	-6.4
Homodimer	Control (chain A)	-33.6	9.3	-0.5	94.4
	Control (chain B)	-11.5	-77.9	-0.1	69.5
	AZ1 (chain A)	-36.6	23.1	-0.2	87.3
	AZ1 (chain B)	-21.3	-50.8	-0.1	30.8
	AZ2 (chain B)	76.4	-123.5	-0.3	52.2
	AZ3 (chain B)	-16.8	-80.2	-0.4	60.6


Favorable

Unfavorable

green

red

FIGURE 9
ZYMEWORKS VARIANT: AZ4



- T394W is negatively designed at the interface core to drive heterodimer formation
- Surrounding mutations are designed to optimize packing and rescue stability

FIGURE 10

ZYMEWORKS VARIANT: AZ4

	clash score ($\frac{\text{kcal}}{\text{mol}}$)		Δ interface area ($\text{\AA}^2$)		Δ packing ($\text{\AA}^3$)		Δ electrostatics ($\frac{\text{kcal}}{\text{mol}}$)		affinity score	
	-	wt. +	-	wt. +	-	wt. +	-	wt. +	-	wt. +
Control 2	-10.8		8.4		-0.1		8.1		0.7	
AZ4	-3.5		-31.1		0.2		3.0		-0.7	

Heterodimer

Control 2 (chain A)	154.2	-4.7	N/A	15.6	N/A	N/A
Control 2 (chain B)	8.3	-71.7	-0.3	2.3	N/A	N/A
AZ4 (chain A)	553.6	1.1	N/A	32.1	N/A	N/A
AZ4 (chain B)	5.6	-115.5	-0.5	6.9	N/A	N/A

Homodimer

green

Favorable

red


Unfavorable

FIGURE 11

ZYMEWORKS VARIANT: AZ5, AZ6

L351R_V397S_F405A_Y407V / K392V_T394W

V397S_F405A_Y407V / K392V_T394W_L368R



- The possible destabilizing effect of negative design (AZ4) is rescued by adding:
 - L351R in AZ5 to improve electrostatic interactions in Region 2
 - L368R in AZ6 to improve packing and electrostatic interactions in Region 1

FIGURE 12

ZYMEWORKS VARIANT: AZ4, AZ5, AZ6

	clash score (kcal/mol) - wt. +	Δ interface area ($\text{\AA}^2$) - wt. +	Δ packing ($\text{\AA}$) - wt. +	Δ electrostatics (kcal/mol) - wt. +	affinity score - wt. +
Heterodimer	Control 2	8.4	-0.1	8.1	0.7
	AZ4	-31.1	0.2	3.0	-0.7
	AZ5	-25.1	0.2	-5.5	-0.8
	AZ6	-32.5	0.0	-11.7	-0.3
Homodimer	Control 2 (chain A)	-4.7	N/A	15.6	N/A
	Control 2 (chain B)	-71.7	-0.3	2.3	N/A
	AZ4 (chain A)	1.1	N/A	32.1	N/A
	AZ4 (chain B)	-115.5	-0.5	6.9	N/A
	AZ5 (chain A)	-44.3	N/A	35.5	N/A
	AZ6 (chain B)	-100.9	-0.3	-16.1	N/A

green

red

Favorable

Unfavorable

Figure 13




Figure 14




Figure 15

FcγRIIA/1-317	1	---	MTM	QMSQNVCPRN	ML	QP	DSQAAAP	K	K	P	57
FcγRIIB/1-310	1	M	ILSFLPVLAT	DWADCKSPQ	H	H	A	T	R	A	66
FcγRIIC/1-323	1	M	ILSFLPVLAT	DWADCKSPQ	H	H	A	T	R	A	66
FcγRIIA/1-254	1	---	---	---	WQ	WQ	MRTED	P	F	F	42
FcγRIIB/1-233	1	---	---	---	WQ	WQ	MRTED	P	F	F	42
Right click to add annotation											
Alignment has no annotations											
FcγRIIA/1-317	58	A	R	I	I	I	R	F	K	R	123
FcγRIIB/1-310	67	R	T	I	I	I	R	F	K	R	132
FcγRIIC/1-323	67	R	T	I	I	I	R	F	K	R	132
FcγRIIA/1-254	43	K	A	A	DN	T	ES	S	Q	ASSY	108
FcγRIIB/1-233	43	K	A	A	DN	T	ES	S	Q	ASSY	108
Right click to add annotation											
Alignment has no annotations											
FcγRIIA/1-317	124	H	H	K	K	K	HL	T	T	H	189
FcγRIIB/1-310	133	H	H	K	K	K	RS	N	N	H	198
FcγRIIC/1-323	133	H	H	K	K	K	RS	N	N	H	198
FcγRIIA/1-254	109	A	R	V	K	E	F	H	N	S	174
FcγRIIB/1-233	109	A	R	V	K	E	F	H	N	S	174
Right click to add annotation											
Alignment has no annotations											
FcγRIIA/1-317	190	Y	L	K	K	V	VA	I	A	A	255
FcγRIIB/1-310	199	Y	L	K	K	A	VA	T	A	A	261
FcγRIIC/1-323	199	Y	L	K	K	A	VA	T	A	A	261
FcγRIIA/1-254	175	L	SKN	ET	N	TQ	ST	SS	Q	S	228
FcγRIIB/1-233	175	L	SKN	ET	N	TQ	ST	SS	Q	S	228
Right click to add annotation											
Alignment has no annotations											
FcγRIIA/1-317	256	AAQF	P	R	Q	M	I	A	R	K	317
FcγRIIB/1-310	262	REM	T	---	---	L	E	K	A	N	310
FcγRIIC/1-323	262	AAQF	P	R	Q	M	I	A	R	K	310
FcγRIIA/1-254	229	---	---	---	---	---	---	---	---	---	323
FcγRIIB/1-233	229	---	---	---	---	---	---	---	---	---	323
Right click to add annotation											
Alignment has no annotations											

Figure 16




Figure 17




Figure 18




Figure 19






Figure 20







Figure 21




Figure 22

Figure 23: Parental wild-type heavy chain IgG1 sequence

GGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS
RDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGDSFFLYSKLTVDKS
RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK

Figure 24: Iterative strategy of heterodimer Fc design




Figure 25A: Non-reducing SDS-PAGE assay to determine heterodimer purity




Figure 25B & C: Non-reducing SDS-PAGE assay to determine heterodimer purity




Figure 26:
Heterodimer Purity Determination after Protein A purification using non-reducing
SDS-PAGE




Figure 27: Sample DSC Data Analysis to Determine T_m:




Figure 28: LC/MS Data Analysis to Determine Heterodimer Purity




Figure 29: In silico analysis of initial negative design variants and independent positive design optimizations: Scaffold 1 – hydrophobic core



Figure 30 In silico analysis of initial negative design variants and independent positive design optimizations: Scaffold 1 – stabilizing 399-400 loop conformation




Figure 31 In silico analysis of initial negative design variants and independent positive design optimizations: Scaffold 1 – stabilizing 399-400 loop conformation




Figure 32




Figure 33 In silico analysis of initial negative design variants and independent positive design optimizations: Scaffold 2 – hydrophobic packing



Figure 34 In silico analysis of initial negative design variants and independent positive design optimizations: Scaffold 2 – stabilizing 399-400 loop conformation




Figure 35




Figure 36A




Figure 36B




Figure 36C




FIGURE 37

Table 6

Variant	Fc mutations chain A	Fc mutations chain B
AZ133	L351Y_F405A_Y407V	T366L_K392M_T394W
AZ134	Q347R_T350V_L351Y_F405A_Y407V	T350V_K360E_T366L_K392M_T394W_T411R
AZ135	L351L_F405A_Y407I	T366L_K392L_T394W
AZ136	L351L_F405A_Y407I	T366L_K392M_T394W
AZ137	L351L_F405A_Y407I	T366L_K392F_T394W
AZ138	L351L_F405A_Y407V	T366L_K392L_T394W
AZ139	L351L_F405A_Y407V	T366L_K392M_T394W
AZ140	L351L_F405A_Y407V	T366L_K392F_T394W
AZ141	L351L_F405S_Y407I	T366L_K392L_T394W
AZ142	L351L_F405S_Y407I	T366L_K392M_T394W
AZ143	L351L_F405S_Y407I	T366L_K392F_T394W
AZ144	L351L_F405S_Y407V	T366L_K392L_T394W
AZ145	L351L_F405S_Y407V	T366L_K392M_T394W
AZ146	L351L_F405S_Y407V	T366L_K392F_T394W
AZ147	L351L_F405T_Y407I	T366L_K392L_T394W
AZ148	L351L_F405T_Y407I	T366L_K392M_T394W
AZ149	L351L_F405T_Y407I	T366L_K392F_T394W
AZ150	L351L_F405T_Y407V	T366L_K392L_T394W
AZ151	L351L_F405T_Y407V	T366L_K392M_T394W
AZ152	L351L_F405T_Y407V	T366L_K392F_T394W
AZ153	L351L_F405V_Y407I	T366L_K392L_T394W
AZ154	L351L_F405V_Y407I	T366L_K392M_T394W
AZ155	L351L_F405V_Y407I	T366L_K392F_T394W
AZ156	L351L_F405V_Y407V	T366L_K392L_T394W
AZ157	L351L_F405V_Y407V	T366L_K392M_T394W
AZ158	L351L_F405V_Y407V	T366L_K392F_T394W
AZ159	L351Y_F405A_Y407I	T366L_K392L_T394W
AZ160	L351Y_F405A_Y407I	T366L_K392M_T394W
AZ161	L351Y_F405A_Y407I	T366L_K392F_T394W
AZ162	L351Y_F405A_Y407V	T366L_K392L_T394W
AZ163	L351Y_F405A_Y407V	T366L_K392M_T394W
AZ164	L351Y_F405A_Y407V	T366L_K392F_T394W
AZ165	L351Y_F405S_Y407I	T366L_K392L_T394W
AZ166	L351Y_F405S_Y407I	T366L_K392M_T394W
AZ167	L351Y_F405S_Y407I	T366L_K392F_T394W
AZ168	L351Y_F405S_Y407V	T366L_K392L_T394W
AZ169	L351Y_F405S_Y407V	T366L_K392M_T394W
AZ170	L351Y_F405S_Y407V	T366L_K392F_T394W
AZ171	L351Y_F405T_Y407I	T366L_K392L_T394W
AZ172	L351Y_F405T_Y407I	T366L_K392M_T394W
AZ173	L351Y_F405T_Y407I	T366L_K392F_T394W
AZ174	L351Y_F405T_Y407V	T366L_K392L_T394W
AZ175	L351Y_F405T_Y407V	T366L_K392M_T394W
AZ176	L351Y_F405T_Y407V	T366L_K392F_T394W
AZ177	L351Y_F405V_Y407I	T366L_K392L_T394W
AZ178	L351Y_F405V_Y407I	T366L_K392M_T394W
AZ179	L351Y_F405V_Y407I	T366L_K392F_T394W
AZ180	L351Y_F405V_Y407V	T366L_K392L_T394W
AZ181	L351Y_F405V_Y407V	T366L_K392M_T394W
AZ182	L351Y_F405V_Y407V	T366L_K392F_T394W
AZ183	Q347E_L351L_F405A_Y407I	T366L_K392L_T394W
AZ184	Q347E_L351L_F405A_Y407I	T366L_K392M_T394W
AZ185	Q347E_L351L_F405A_Y407I	T366L_K392F_T394W
AZ186	Q347E_L351L_F405A_Y407V	T366L_K392L_T394W

AZ187	Q347E_L351L_F405A_Y407V	T366L_K392M_T394W
AZ188	Q347E_L351L_F405A_Y407V	T366L_K392F_T394W
AZ189	Q347E_L351L_F405S_Y407I	T366L_K392L_T394W
AZ190	Q347E_L351L_F405S_Y407I	T366L_K392M_T394W
AZ191	Q347E_L351L_F405S_Y407I	T366L_K392F_T394W
AZ192	Q347E_L351L_F405S_Y407V	T366L_K392L_T394W
AZ193	Q347E_L351L_F405S_Y407V	T366L_K392M_T394W
AZ194	Q347E_L351L_F405S_Y407V	T366L_K392F_T394W
AZ195	Q347E_L351L_F405T_Y407I	T366L_K392L_T394W
AZ196	Q347E_L351L_F405T_Y407I	T366L_K392M_T394W
AZ197	Q347E_L351L_F405T_Y407I	T366L_K392F_T394W
AZ198	Q347E_L351L_F405T_Y407V	T366L_K392L_T394W
AZ199	Q347E_L351L_F405T_Y407V	T366L_K392M_T394W
AZ200	Q347E_L351L_F405T_Y407V	T366L_K392F_T394W
AZ201	Q347E_L351L_F405V_Y407I	T366L_K392L_T394W
AZ202	Q347E_L351L_F405V_Y407I	T366L_K392M_T394W
AZ203	Q347E_L351L_F405V_Y407I	T366L_K392F_T394W
AZ204	Q347E_L351L_F405V_Y407V	T366L_K392L_T394W
AZ205	Q347E_L351L_F405V_Y407V	T366L_K392M_T394W
AZ206	Q347E_L351L_F405V_Y407V	T366L_K392F_T394W
AZ207	Q347E_L351Y_F405A_Y407I	T366L_K392L_T394W
AZ208	Q347E_L351Y_F405A_Y407I	T366L_K392M_T394W
AZ209	Q347E_L351Y_F405A_Y407I	T366L_K392F_T394W
AZ210	Q347E_L351Y_F405A_Y407V	T366L_K392L_T394W
AZ211	Q347E_L351Y_F405A_Y407V	T366L_K392M_T394W
AZ212	Q347E_L351Y_F405A_Y407V	T366L_K392F_T394W
AZ213	Q347E_L351Y_F405S_Y407I	T366L_K392L_T394W
AZ214	Q347E_L351Y_F405S_Y407I	T366L_K392M_T394W
AZ215	Q347E_L351Y_F405S_Y407I	T366L_K392F_T394W
AZ216	Q347E_L351Y_F405S_Y407V	T366L_K392L_T394W
AZ217	Q347E_L351Y_F405S_Y407V	T366L_K392M_T394W
AZ218	Q347E_L351Y_F405S_Y407V	T366L_K392F_T394W
AZ219	Q347E_L351Y_F405T_Y407I	T366L_K392L_T394W
AZ220	Q347E_L351Y_F405T_Y407I	T366L_K392M_T394W
AZ221	Q347E_L351Y_F405T_Y407I	T366L_K392F_T394W
AZ222	Q347E_L351Y_F405T_Y407V	T366L_K392L_T394W
AZ223	Q347E_L351Y_F405T_Y407V	T366L_K392M_T394W
AZ224	Q347E_L351Y_F405T_Y407V	T366L_K392F_T394W
AZ225	Q347E_L351Y_F405V_Y407I	T366L_K392L_T394W
AZ226	Q347E_L351Y_F405V_Y407I	T366L_K392M_T394W
AZ227	Q347E_L351Y_F405V_Y407I	T366L_K392F_T394W
AZ228	Q347E_L351Y_F405V_Y407V	T366L_K392L_T394W
AZ229	Q347E_L351Y_F405V_Y407V	T366L_K392M_T394W
AZ230	Q347E_L351Y_F405V_Y407V	T366L_K392F_T394W
AZ231	Q347R_L351L_F405A_Y407I	K360D_T366L_K392L_T394W
AZ232	Q347R_L351L_F405A_Y407I	K360D_T366L_K392M_T394W
AZ233	Q347R_L351L_F405A_Y407I	K360D_T366L_K392F_T394W
AZ234	Q347R_L351L_F405A_Y407I	K360E_T366L_K392L_T394W
AZ235	Q347R_L351L_F405A_Y407I	K360E_T366L_K392M_T394W
AZ236	Q347R_L351L_F405A_Y407I	K360E_T366L_K392F_T394W
AZ237	Q347R_L351L_F405A_Y407V	K360D_T366L_K392L_T394W
AZ238	Q347R_L351L_F405A_Y407V	K360D_T366L_K392M_T394W
AZ239	Q347R_L351L_F405A_Y407V	K360D_T366L_K392F_T394W
AZ240	Q347R_L351L_F405A_Y407V	K360E_T366L_K392L_T394W
AZ241	Q347R_L351L_F405A_Y407V	K360E_T366L_K392M_T394W
AZ242	Q347R_L351L_F405A_Y407V	K360E_T366L_K392F_T394W
AZ243	Q347R_L351L_F405S_Y407I	K360D_T366L_K392L_T394W
AZ244	Q347R_L351L_F405S_Y407I	K360D_T366L_K392M_T394W

AZ245	Q347R_L351L_F405S_Y407I	K360D_T366L_K392F_T394W
AZ246	Q347R_L351L_F405S_Y407I	K360E_T366L_K392L_T394W
AZ247	Q347R_L351L_F405S_Y407I	K360E_T366L_K392M_T394W
AZ248	Q347R_L351L_F405S_Y407I	K360E_T366L_K392F_T394W
AZ249	Q347R_L351L_F405S_Y407V	K360D_T366L_K392L_T394W
AZ250	Q347R_L351L_F405S_Y407V	K360D_T366L_K392M_T394W
AZ251	Q347R_L351L_F405S_Y407V	K360D_T366L_K392F_T394W
AZ252	Q347R_L351L_F405S_Y407V	K360E_T366L_K392L_T394W
AZ253	Q347R_L351L_F405S_Y407V	K360E_T366L_K392M_T394W
AZ254	Q347R_L351L_F405S_Y407V	K360E_T366L_K392F_T394W
AZ255	Q347R_L351L_F405T_Y407I	K360D_T366L_K392L_T394W
AZ256	Q347R_L351L_F405T_Y407I	K360D_T366L_K392M_T394W
AZ257	Q347R_L351L_F405T_Y407I	K360D_T366L_K392F_T394W
AZ258	Q347R_L351L_F405T_Y407I	K360E_T366L_K392L_T394W
AZ259	Q347R_L351L_F405T_Y407I	K360E_T366L_K392M_T394W
AZ260	Q347R_L351L_F405T_Y407I	K360E_T366L_K392F_T394W
AZ261	Q347R_L351L_F405T_Y407V	K360D_T366L_K392L_T394W
AZ262	Q347R_L351L_F405T_Y407V	K360D_T366L_K392M_T394W
AZ263	Q347R_L351L_F405T_Y407V	K360D_T366L_K392F_T394W
AZ264	Q347R_L351L_F405T_Y407V	K360E_T366L_K392L_T394W
AZ265	Q347R_L351L_F405T_Y407V	K360E_T366L_K392M_T394W
AZ266	Q347R_L351L_F405T_Y407V	K360E_T366L_K392F_T394W
AZ267	Q347R_L351L_F405V_Y407I	K360D_T366L_K392L_T394W
AZ268	Q347R_L351L_F405V_Y407I	K360D_T366L_K392M_T394W
AZ269	Q347R_L351L_F405V_Y407I	K360D_T366L_K392F_T394W
AZ270	Q347R_L351L_F405V_Y407I	K360E_T366L_K392L_T394W
AZ271	Q347R_L351L_F405V_Y407I	K360E_T366L_K392M_T394W
AZ272	Q347R_L351L_F405V_Y407I	K360E_T366L_K392F_T394W
AZ273	Q347R_L351L_F405V_Y407V	K360D_T366L_K392L_T394W
AZ274	Q347R_L351L_F405V_Y407V	K360D_T366L_K392M_T394W
AZ275	Q347R_L351L_F405V_Y407V	K360D_T366L_K392F_T394W
AZ276	Q347R_L351L_F405V_Y407V	K360E_T366L_K392L_T394W
AZ277	Q347R_L351L_F405V_Y407V	K360E_T366L_K392M_T394W
AZ278	Q347R_L351L_F405V_Y407V	K360E_T366L_K392F_T394W
AZ279	Q347R_L351Y_F405A_Y407I	K360D_T366L_K392L_T394W
AZ280	Q347R_L351Y_F405A_Y407I	K360D_T366L_K392M_T394W
AZ281	Q347R_L351Y_F405A_Y407I	K360D_T366L_K392F_T394W
AZ282	Q347R_L351Y_F405A_Y407I	K360E_T366L_K392L_T394W
AZ283	Q347R_L351Y_F405A_Y407I	K360E_T366L_K392M_T394W
AZ284	Q347R_L351Y_F405A_Y407I	K360E_T366L_K392F_T394W
AZ285	Q347R_L351Y_F405A_Y407V	K360D_T366L_K392L_T394W
AZ286	Q347R_L351Y_F405A_Y407V	K360D_T366L_K392M_T394W
AZ287	Q347R_L351Y_F405A_Y407V	K360D_T366L_K392F_T394W
AZ288	Q347R_L351Y_F405A_Y407V	K360E_T366L_K392L_T394W
AZ289	Q347R_L351Y_F405A_Y407V	K360E_T366L_K392M_T394W
AZ290	Q347R_L351Y_F405A_Y407V	K360E_T366L_K392F_T394W
AZ291	Q347R_L351Y_F405S_Y407I	K360D_T366L_K392L_T394W
AZ292	Q347R_L351Y_F405S_Y407I	K360D_T366L_K392M_T394W
AZ293	Q347R_L351Y_F405S_Y407I	K360D_T366L_K392F_T394W
AZ294	Q347R_L351Y_F405S_Y407I	K360E_T366L_K392L_T394W
AZ295	Q347R_L351Y_F405S_Y407I	K360E_T366L_K392M_T394W
AZ296	Q347R_L351Y_F405S_Y407I	K360E_T366L_K392F_T394W
AZ297	Q347R_L351Y_F405S_Y407V	K360D_T366L_K392L_T394W
AZ298	Q347R_L351Y_F405S_Y407V	K360D_T366L_K392M_T394W
AZ299	Q347R_L351Y_F405S_Y407V	K360D_T366L_K392F_T394W
AZ300	Q347R_L351Y_F405S_Y407V	K360E_T366L_K392L_T394W
AZ301	Q347R_L351Y_F405S_Y407V	K360E_T366L_K392M_T394W
AZ302	Q347R_L351Y_F405S_Y407V	K360E_T366L_K392F_T394W

AZ303	Q347R_L351Y_F405T_Y407I	K360D_T366L_K392L_T394W
AZ304	Q347R_L351Y_F405T_Y407I	K360D_T366L_K392M_T394W
AZ305	Q347R_L351Y_F405T_Y407I	K360D_T366L_K392F_T394W
AZ306	Q347R_L351Y_F405T_Y407I	K360E_T366L_K392L_T394W
AZ307	Q347R_L351Y_F405T_Y407I	K360E_T366L_K392M_T394W
AZ308	Q347R_L351Y_F405T_Y407I	K360E_T366L_K392F_T394W
AZ309	Q347R_L351Y_F405T_Y407V	K360D_T366L_K392L_T394W
AZ310	Q347R_L351Y_F405T_Y407V	K360D_T366L_K392M_T394W
AZ311	Q347R_L351Y_F405T_Y407V	K360D_T366L_K392F_T394W
AZ312	Q347R_L351Y_F405T_Y407V	K360E_T366L_K392L_T394W
AZ313	Q347R_L351Y_F405T_Y407V	K360E_T366L_K392M_T394W
AZ314	Q347R_L351Y_F405T_Y407V	K360E_T366L_K392F_T394W
AZ315	Q347R_L351Y_F405V_Y407I	K360D_T366L_K392L_T394W
AZ316	Q347R_L351Y_F405V_Y407I	K360D_T366L_K392M_T394W
AZ317	Q347R_L351Y_F405V_Y407I	K360D_T366L_K392F_T394W
AZ318	Q347R_L351Y_F405V_Y407I	K360E_T366L_K392L_T394W
AZ319	Q347R_L351Y_F405V_Y407I	K360E_T366L_K392M_T394W
AZ320	Q347R_L351Y_F405V_Y407I	K360E_T366L_K392F_T394W
AZ321	Q347R_L351Y_F405V_Y407V	K360D_T366L_K392L_T394W
AZ322	Q347R_L351Y_F405V_Y407V	K360D_T366L_K392M_T394W
AZ323	Q347R_L351Y_F405V_Y407V	K360D_T366L_K392F_T394W
AZ324	Q347R_L351Y_F405V_Y407V	K360E_T366L_K392L_T394W
AZ325	Q347R_L351Y_F405V_Y407V	K360E_T366L_K392M_T394W
AZ326	Q347R_L351Y_F405V_Y407V	K360E_T366L_K392F_T394W
AZ327	Q347K_L351L_F405A_Y407I	K360D_T366L_K392L_T394W
AZ328	Q347K_L351L_F405A_Y407I	K360D_T366L_K392M_T394W
AZ329	Q347K_L351L_F405A_Y407I	K360D_T366L_K392F_T394W
AZ330	Q347K_L351L_F405A_Y407I	K360E_T366L_K392L_T394W
AZ331	Q347K_L351L_F405A_Y407I	K360E_T366L_K392M_T394W
AZ332	Q347K_L351L_F405A_Y407I	K360E_T366L_K392F_T394W
AZ333	Q347K_L351L_F405A_Y407V	K360D_T366L_K392L_T394W
AZ334	Q347K_L351L_F405A_Y407V	K360D_T366L_K392M_T394W
AZ335	Q347K_L351L_F405A_Y407V	K360D_T366L_K392F_T394W
AZ336	Q347K_L351L_F405A_Y407V	K360E_T366L_K392L_T394W
AZ337	Q347K_L351L_F405A_Y407V	K360E_T366L_K392M_T394W
AZ338	Q347K_L351L_F405A_Y407V	K360E_T366L_K392F_T394W
AZ339	Q347K_L351L_F405S_Y407I	K360D_T366L_K392L_T394W
AZ340	Q347K_L351L_F405S_Y407I	K360D_T366L_K392M_T394W
AZ341	Q347K_L351L_F405S_Y407I	K360D_T366L_K392F_T394W
AZ342	Q347K_L351L_F405S_Y407I	K360E_T366L_K392L_T394W
AZ343	Q347K_L351L_F405S_Y407I	K360E_T366L_K392M_T394W
AZ344	Q347K_L351L_F405S_Y407I	K360E_T366L_K392F_T394W
AZ345	Q347K_L351L_F405S_Y407V	K360D_T366L_K392L_T394W
AZ346	Q347K_L351L_F405S_Y407V	K360D_T366L_K392M_T394W
AZ347	Q347K_L351L_F405S_Y407V	K360D_T366L_K392F_T394W
AZ348	Q347K_L351L_F405S_Y407V	K360E_T366L_K392L_T394W
AZ349	Q347K_L351L_F405S_Y407V	K360E_T366L_K392M_T394W
AZ350	Q347K_L351L_F405S_Y407V	K360E_T366L_K392F_T394W
AZ351	Q347K_L351L_F405T_Y407I	K360D_T366L_K392L_T394W
AZ352	Q347K_L351L_F405T_Y407I	K360D_T366L_K392M_T394W
AZ353	Q347K_L351L_F405T_Y407I	K360D_T366L_K392F_T394W
AZ354	Q347K_L351L_F405T_Y407I	K360E_T366L_K392L_T394W
AZ355	Q347K_L351L_F405T_Y407I	K360E_T366L_K392M_T394W
AZ356	Q347K_L351L_F405T_Y407I	K360E_T366L_K392F_T394W
AZ357	Q347K_L351L_F405T_Y407V	K360D_T366L_K392L_T394W
AZ358	Q347K_L351L_F405T_Y407V	K360D_T366L_K392M_T394W
AZ359	Q347K_L351L_F405T_Y407V	K360D_T366L_K392F_T394W
AZ360	Q347K_L351L_F405T_Y407V	K360E_T366L_K392L_T394W

AZ361	Q347K_L351L_F405T_Y407V	K360E_T366L_K392M_T394W
AZ362	Q347K_L351L_F405T_Y407V	K360E_T366L_K392F_T394W
AZ363	Q347K_L351L_F405V_Y407I	K360D_T366L_K392L_T394W
AZ364	Q347K_L351L_F405V_Y407I	K360D_T366L_K392M_T394W
AZ365	Q347K_L351L_F405V_Y407I	K360D_T366L_K392F_T394W
AZ366	Q347K_L351L_F405V_Y407I	K360E_T366L_K392L_T394W
AZ367	Q347K_L351L_F405V_Y407I	K360E_T366L_K392M_T394W
AZ368	Q347K_L351L_F405V_Y407I	K360E_T366L_K392F_T394W
AZ369	Q347K_L351L_F405V_Y407V	K360D_T366L_K392L_T394W
AZ370	Q347K_L351L_F405V_Y407V	K360D_T366L_K392M_T394W
AZ371	Q347K_L351L_F405V_Y407V	K360D_T366L_K392F_T394W
AZ372	Q347K_L351L_F405V_Y407V	K360E_T366L_K392L_T394W
AZ373	Q347K_L351L_F405V_Y407V	K360E_T366L_K392M_T394W
AZ374	Q347K_L351L_F405V_Y407V	K360E_T366L_K392F_T394W
AZ375	Q347K_L351Y_F405A_Y407I	K360D_T366L_K392L_T394W
AZ376	Q347K_L351Y_F405A_Y407I	K360D_T366L_K392M_T394W
AZ377	Q347K_L351Y_F405A_Y407I	K360D_T366L_K392F_T394W
AZ378	Q347K_L351Y_F405A_Y407I	K360E_T366L_K392L_T394W
AZ379	Q347K_L351Y_F405A_Y407I	K360E_T366L_K392M_T394W
AZ380	Q347K_L351Y_F405A_Y407I	K360E_T366L_K392F_T394W
AZ381	Q347K_L351Y_F405A_Y407V	K360D_T366L_K392L_T394W
AZ382	Q347K_L351Y_F405A_Y407V	K360D_T366L_K392M_T394W
AZ383	Q347K_L351Y_F405A_Y407V	K360D_T366L_K392F_T394W
AZ384	Q347K_L351Y_F405A_Y407V	K360E_T366L_K392L_T394W
AZ385	Q347K_L351Y_F405A_Y407V	K360E_T366L_K392M_T394W
AZ386	Q347K_L351Y_F405A_Y407V	K360E_T366L_K392F_T394W
AZ387	Q347K_L351Y_F405S_Y407I	K360D_T366L_K392L_T394W
AZ388	Q347K_L351Y_F405S_Y407I	K360D_T366L_K392M_T394W
AZ389	Q347K_L351Y_F405S_Y407I	K360D_T366L_K392F_T394W
AZ390	Q347K_L351Y_F405S_Y407I	K360E_T366L_K392L_T394W
AZ391	Q347K_L351Y_F405S_Y407I	K360E_T366L_K392M_T394W
AZ392	Q347K_L351Y_F405S_Y407I	K360E_T366L_K392F_T394W
AZ393	Q347K_L351Y_F405S_Y407V	K360D_T366L_K392L_T394W
AZ394	Q347K_L351Y_F405S_Y407V	K360D_T366L_K392M_T394W
AZ395	Q347K_L351Y_F405S_Y407V	K360D_T366L_K392F_T394W
AZ396	Q347K_L351Y_F405S_Y407V	K360E_T366L_K392L_T394W
AZ397	Q347K_L351Y_F405S_Y407V	K360E_T366L_K392M_T394W
AZ398	Q347K_L351Y_F405S_Y407V	K360E_T366L_K392F_T394W
AZ399	Q347K_L351Y_F405T_Y407I	K360D_T366L_K392L_T394W
AZ400	Q347K_L351Y_F405T_Y407I	K360D_T366L_K392M_T394W
AZ401	Q347K_L351Y_F405T_Y407I	K360D_T366L_K392F_T394W
AZ402	Q347K_L351Y_F405T_Y407I	K360E_T366L_K392L_T394W
AZ403	Q347K_L351Y_F405T_Y407I	K360E_T366L_K392M_T394W
AZ404	Q347K_L351Y_F405T_Y407I	K360E_T366L_K392F_T394W
AZ405	Q347K_L351Y_F405T_Y407V	K360D_T366L_K392L_T394W
AZ406	Q347K_L351Y_F405T_Y407V	K360D_T366L_K392M_T394W
AZ407	Q347K_L351Y_F405T_Y407V	K360D_T366L_K392F_T394W
AZ408	Q347K_L351Y_F405T_Y407V	K360E_T366L_K392L_T394W
AZ409	Q347K_L351Y_F405T_Y407V	K360E_T366L_K392M_T394W
AZ410	Q347K_L351Y_F405T_Y407V	K360E_T366L_K392F_T394W
AZ411	Q347K_L351Y_F405V_Y407I	K360D_T366L_K392L_T394W
AZ412	Q347K_L351Y_F405V_Y407I	K360D_T366L_K392M_T394W
AZ413	Q347K_L351Y_F405V_Y407I	K360D_T366L_K392F_T394W
AZ414	Q347K_L351Y_F405V_Y407I	K360E_T366L_K392L_T394W
AZ415	Q347K_L351Y_F405V_Y407I	K360E_T366L_K392M_T394W
AZ416	Q347K_L351Y_F405V_Y407I	K360E_T366L_K392F_T394W
AZ417	Q347K_L351Y_F405V_Y407V	K360D_T366L_K392L_T394W
AZ418	Q347K_L351Y_F405V_Y407V	K360D_T366L_K392M_T394W

AZ419	Q347K_L351Y_F405V_Y407V	K360D_T366L_K392F_T394W
AZ420	Q347K_L351Y_F405V_Y407V	K360E_T366L_K392L_T394W
AZ421	Q347K_L351Y_F405V_Y407V	K360E_T366L_K392M_T394W
AZ422	Q347K_L351Y_F405V_Y407V	K360E_T366L_K392F_T394W
AZ423	L351L_S400D_F405A_Y407I	T366L_N390R_K392L_T394W
AZ424	L351L_S400D_F405A_Y407I	T366L_N390R_K392M_T394W
AZ425	L351L_S400D_F405A_Y407I	T366L_N390R_K392F_T394W
AZ426	L351L_S400D_F405A_Y407I	T366L_N390K_K392L_T394W
AZ427	L351L_S400D_F405A_Y407I	T366L_N390K_K392M_T394W
AZ428	L351L_S400D_F405A_Y407I	T366L_N390K_K392F_T394W
AZ429	L351L_S400D_F405A_Y407V	T366L_N390R_K392L_T394W
AZ430	L351L_S400D_F405A_Y407V	T366L_N390R_K392M_T394W
AZ431	L351L_S400D_F405A_Y407V	T366L_N390R_K392F_T394W
AZ432	L351L_S400D_F405A_Y407V	T366L_N390K_K392L_T394W
AZ433	L351L_S400D_F405A_Y407V	T366L_N390K_K392M_T394W
AZ434	L351L_S400D_F405A_Y407V	T366L_N390K_K392F_T394W
AZ435	L351L_S400D_F405S_Y407I	T366L_N390R_K392L_T394W
AZ436	L351L_S400D_F405S_Y407I	T366L_N390R_K392M_T394W
AZ437	L351L_S400D_F405S_Y407I	T366L_N390R_K392F_T394W
AZ438	L351L_S400D_F405S_Y407I	T366L_N390K_K392L_T394W
AZ439	L351L_S400D_F405S_Y407I	T366L_N390K_K392M_T394W
AZ440	L351L_S400D_F405S_Y407I	T366L_N390K_K392F_T394W
AZ441	L351L_S400D_F405S_Y407V	T366L_N390R_K392L_T394W
AZ442	L351L_S400D_F405S_Y407V	T366L_N390R_K392M_T394W
AZ443	L351L_S400D_F405S_Y407V	T366L_N390R_K392F_T394W
AZ444	L351L_S400D_F405S_Y407V	T366L_N390K_K392L_T394W
AZ445	L351L_S400D_F405S_Y407V	T366L_N390K_K392M_T394W
AZ446	L351L_S400D_F405S_Y407V	T366L_N390K_K392F_T394W
AZ447	L351L_S400D_F405T_Y407I	T366L_N390R_K392L_T394W
AZ448	L351L_S400D_F405T_Y407I	T366L_N390R_K392M_T394W
AZ449	L351L_S400D_F405T_Y407I	T366L_N390R_K392F_T394W
AZ450	L351L_S400D_F405T_Y407I	T366L_N390K_K392L_T394W
AZ451	L351L_S400D_F405T_Y407I	T366L_N390K_K392M_T394W
AZ452	L351L_S400D_F405T_Y407I	T366L_N390K_K392F_T394W
AZ453	L351L_S400D_F405T_Y407V	T366L_N390R_K392L_T394W
AZ454	L351L_S400D_F405T_Y407V	T366L_N390R_K392M_T394W
AZ455	L351L_S400D_F405T_Y407V	T366L_N390R_K392F_T394W
AZ456	L351L_S400D_F405T_Y407V	T366L_N390K_K392L_T394W
AZ457	L351L_S400D_F405T_Y407V	T366L_N390K_K392M_T394W
AZ458	L351L_S400D_F405T_Y407V	T366L_N390K_K392F_T394W
AZ459	L351L_S400D_F405V_Y407I	T366L_N390R_K392L_T394W
AZ460	L351L_S400D_F405V_Y407I	T366L_N390R_K392M_T394W
AZ461	L351L_S400D_F405V_Y407I	T366L_N390R_K392F_T394W
AZ462	L351L_S400D_F405V_Y407I	T366L_N390K_K392L_T394W
AZ463	L351L_S400D_F405V_Y407I	T366L_N390K_K392M_T394W
AZ464	L351L_S400D_F405V_Y407I	T366L_N390K_K392F_T394W
AZ465	L351L_S400D_F405V_Y407V	T366L_N390R_K392L_T394W
AZ466	L351L_S400D_F405V_Y407V	T366L_N390R_K392M_T394W
AZ467	L351L_S400D_F405V_Y407V	T366L_N390R_K392F_T394W
AZ468	L351L_S400D_F405V_Y407V	T366L_N390K_K392L_T394W
AZ469	L351L_S400D_F405V_Y407V	T366L_N390K_K392M_T394W
AZ470	L351L_S400D_F405V_Y407V	T366L_N390K_K392F_T394W
AZ471	L351L_S400E_F405A_Y407I	T366L_N390R_K392L_T394W
AZ472	L351L_S400E_F405A_Y407I	T366L_N390R_K392M_T394W
AZ473	L351L_S400E_F405A_Y407I	T366L_N390R_K392F_T394W
AZ474	L351L_S400E_F405A_Y407I	T366L_N390K_K392L_T394W
AZ475	L351L_S400E_F405A_Y407I	T366L_N390K_K392M_T394W
AZ476	L351L_S400E_F405A_Y407I	T366L_N390K_K392F_T394W

AZ477	L351L_S400E_F405A_Y407V	T366L_N390R_K392L_T394W
AZ478	L351L_S400E_F405A_Y407V	T366L_N390R_K392M_T394W
AZ479	L351L_S400E_F405A_Y407V	T366L_N390R_K392F_T394W
AZ480	L351L_S400E_F405A_Y407V	T366L_N390K_K392L_T394W
AZ481	L351L_S400E_F405A_Y407V	T366L_N390K_K392M_T394W
AZ482	L351L_S400E_F405A_Y407V	T366L_N390K_K392F_T394W
AZ483	L351L_S400E_F405S_Y407I	T366L_N390R_K392L_T394W
AZ484	L351L_S400E_F405S_Y407I	T366L_N390R_K392M_T394W
AZ485	L351L_S400E_F405S_Y407I	T366L_N390R_K392F_T394W
AZ486	L351L_S400E_F405S_Y407I	T366L_N390K_K392L_T394W
AZ487	L351L_S400E_F405S_Y407I	T366L_N390K_K392M_T394W
AZ488	L351L_S400E_F405S_Y407I	T366L_N390K_K392F_T394W
AZ489	L351L_S400E_F405S_Y407V	T366L_N390R_K392L_T394W
AZ490	L351L_S400E_F405S_Y407V	T366L_N390R_K392M_T394W
AZ491	L351L_S400E_F405S_Y407V	T366L_N390R_K392F_T394W
AZ492	L351L_S400E_F405S_Y407V	T366L_N390K_K392L_T394W
AZ493	L351L_S400E_F405S_Y407V	T366L_N390K_K392M_T394W
AZ494	L351L_S400E_F405S_Y407V	T366L_N390K_K392F_T394W
AZ495	L351L_S400E_F405T_Y407I	T366L_N390R_K392L_T394W
AZ496	L351L_S400E_F405T_Y407I	T366L_N390R_K392M_T394W
AZ497	L351L_S400E_F405T_Y407I	T366L_N390R_K392F_T394W
AZ498	L351L_S400E_F405T_Y407I	T366L_N390K_K392L_T394W
AZ499	L351L_S400E_F405T_Y407I	T366L_N390K_K392M_T394W
AZ500	L351L_S400E_F405T_Y407I	T366L_N390K_K392F_T394W
AZ501	L351L_S400E_F405T_Y407V	T366L_N390R_K392L_T394W
AZ502	L351L_S400E_F405T_Y407V	T366L_N390R_K392M_T394W
AZ503	L351L_S400E_F405T_Y407V	T366L_N390R_K392F_T394W
AZ504	L351L_S400E_F405T_Y407V	T366L_N390K_K392L_T394W
AZ505	L351L_S400E_F405T_Y407V	T366L_N390K_K392M_T394W
AZ506	L351L_S400E_F405T_Y407V	T366L_N390K_K392F_T394W
AZ507	L351L_S400E_F405V_Y407I	T366L_N390R_K392L_T394W
AZ508	L351L_S400E_F405V_Y407I	T366L_N390R_K392M_T394W
AZ509	L351L_S400E_F405V_Y407I	T366L_N390R_K392F_T394W
AZ510	L351L_S400E_F405V_Y407I	T366L_N390K_K392L_T394W
AZ511	L351L_S400E_F405V_Y407I	T366L_N390K_K392M_T394W
AZ512	L351L_S400E_F405V_Y407I	T366L_N390K_K392F_T394W
AZ513	L351L_S400E_F405V_Y407V	T366L_N390R_K392L_T394W
AZ514	L351L_S400E_F405V_Y407V	T366L_N390R_K392M_T394W
AZ515	L351L_S400E_F405V_Y407V	T366L_N390R_K392F_T394W
AZ516	L351L_S400E_F405V_Y407V	T366L_N390K_K392L_T394W
AZ517	L351L_S400E_F405V_Y407V	T366L_N390K_K392M_T394W
AZ518	L351L_S400E_F405V_Y407V	T366L_N390K_K392F_T394W
AZ519	L351L_F405A_Y407I	T366L_K392L_T394W_K409L_T411R
AZ520	L351L_F405A_Y407I	T366L_K392L_T394W_K409L_T411K
AZ521	L351L_F405A_Y407I	T366L_K392L_T394W_K409M_T411R
AZ522	L351L_F405A_Y407I	T366L_K392L_T394W_K409M_T411K
AZ523	L351L_F405A_Y407I	T366L_K392M_T394W_K409L_T411R
AZ524	L351L_F405A_Y407I	T366L_K392M_T394W_K409L_T411K
AZ525	L351L_F405A_Y407I	T366L_K392M_T394W_K409M_T411R
AZ526	L351L_F405A_Y407I	T366L_K392M_T394W_K409M_T411K
AZ527	L351L_F405A_Y407I	T366L_K392F_T394W_K409L_T411R
AZ528	L351L_F405A_Y407I	T366L_K392F_T394W_K409L_T411K
AZ529	L351L_F405A_Y407I	T366L_K392F_T394W_K409M_T411R
AZ530	L351L_F405A_Y407I	T366L_K392F_T394W_K409M_T411K
AZ531	L351L_F405A_Y407V	T366L_K392L_T394W_K409L_T411R
AZ532	L351L_F405A_Y407V	T366L_K392L_T394W_K409L_T411K
AZ533	L351L_F405A_Y407V	T366L_K392L_T394W_K409M_T411R
AZ534	L351L_F405A_Y407V	T366L_K392L_T394W_K409M_T411K

AZ535	L351L_F405A_Y407V	T366L_K392M_T394W_K409L_T411R
AZ536	L351L_F405A_Y407V	T366L_K392M_T394W_K409L_T411K
AZ537	L351L_F405A_Y407V	T366L_K392M_T394W_K409M_T411R
AZ538	L351L_F405A_Y407V	T366L_K392M_T394W_K409M_T411K
AZ539	L351L_F405A_Y407V	T366L_K392F_T394W_K409L_T411R
AZ540	L351L_F405A_Y407V	T366L_K392F_T394W_K409L_T411K
AZ541	L351L_F405A_Y407V	T366L_K392F_T394W_K409M_T411R
AZ542	L351L_F405A_Y407V	T366L_K392F_T394W_K409M_T411K
AZ543	L351L_F405S_Y407I	T366L_K392L_T394W_K409L_T411R
AZ544	L351L_F405S_Y407I	T366L_K392L_T394W_K409L_T411K
AZ545	L351L_F405S_Y407I	T366L_K392L_T394W_K409M_T411R
AZ546	L351L_F405S_Y407I	T366L_K392L_T394W_K409M_T411K
AZ547	L351L_F405S_Y407I	T366L_K392M_T394W_K409L_T411R
AZ548	L351L_F405S_Y407I	T366L_K392M_T394W_K409L_T411K
AZ549	L351L_F405S_Y407I	T366L_K392M_T394W_K409M_T411R
AZ550	L351L_F405S_Y407I	T366L_K392M_T394W_K409M_T411K
AZ551	L351L_F405S_Y407I	T366L_K392F_T394W_K409L_T411R
AZ552	L351L_F405S_Y407I	T366L_K392F_T394W_K409L_T411K
AZ553	L351L_F405S_Y407I	T366L_K392F_T394W_K409M_T411R
AZ554	L351L_F405S_Y407I	T366L_K392F_T394W_K409M_T411K
AZ555	L351L_F405S_Y407V	T366L_K392L_T394W_K409L_T411R
AZ556	L351L_F405S_Y407V	T366L_K392L_T394W_K409L_T411K
AZ557	L351L_F405S_Y407V	T366L_K392L_T394W_K409M_T411R
AZ558	L351L_F405S_Y407V	T366L_K392L_T394W_K409M_T411K
AZ559	L351L_F405S_Y407V	T366L_K392M_T394W_K409L_T411R
AZ560	L351L_F405S_Y407V	T366L_K392M_T394W_K409L_T411K
AZ561	L351L_F405S_Y407V	T366L_K392M_T394W_K409M_T411R
AZ562	L351L_F405S_Y407V	T366L_K392M_T394W_K409M_T411K
AZ563	L351L_F405S_Y407V	T366L_K392F_T394W_K409L_T411R
AZ564	L351L_F405S_Y407V	T366L_K392F_T394W_K409L_T411K
AZ565	L351L_F405S_Y407V	T366L_K392F_T394W_K409M_T411R
AZ566	L351L_F405S_Y407V	T366L_K392F_T394W_K409M_T411K
AZ567	L351L_F405V_Y407I	T366L_K392L_T394W_K409L_T411R
AZ568	L351L_F405V_Y407I	T366L_K392L_T394W_K409L_T411K
AZ569	L351L_F405V_Y407I	T366L_K392L_T394W_K409M_T411R
AZ570	L351L_F405V_Y407I	T366L_K392L_T394W_K409M_T411K
AZ571	L351L_F405V_Y407I	T366L_K392M_T394W_K409L_T411R
AZ572	L351L_F405V_Y407I	T366L_K392M_T394W_K409L_T411K
AZ573	L351L_F405V_Y407I	T366L_K392M_T394W_K409M_T411R
AZ574	L351L_F405V_Y407I	T366L_K392M_T394W_K409M_T411K
AZ575	L351L_F405V_Y407I	T366L_K392F_T394W_K409L_T411R
AZ576	L351L_F405V_Y407I	T366L_K392F_T394W_K409L_T411K
AZ577	L351L_F405V_Y407I	T366L_K392F_T394W_K409M_T411R
AZ578	L351L_F405V_Y407I	T366L_K392F_T394W_K409M_T411K
AZ579	L351L_F405V_Y407V	T366L_K392L_T394W_K409L_T411R
AZ580	L351L_F405V_Y407V	T366L_K392L_T394W_K409L_T411K
AZ581	L351L_F405V_Y407V	T366L_K392L_T394W_K409M_T411R
AZ582	L351L_F405V_Y407V	T366L_K392L_T394W_K409M_T411K
AZ583	L351L_F405V_Y407V	T366L_K392M_T394W_K409L_T411R
AZ584	L351L_F405V_Y407V	T366L_K392M_T394W_K409L_T411K
AZ585	L351L_F405V_Y407V	T366L_K392M_T394W_K409M_T411R
AZ586	L351L_F405V_Y407V	T366L_K392M_T394W_K409M_T411K
AZ587	L351L_F405V_Y407V	T366L_K392F_T394W_K409L_T411R
AZ588	L351L_F405V_Y407V	T366L_K392F_T394W_K409L_T411K
AZ589	L351L_F405V_Y407V	T366L_K392F_T394W_K409M_T411R
AZ590	L351L_F405V_Y407V	T366L_K392F_T394W_K409M_T411K
AZ591	L351Y_S400D_F405A_Y407I	T366L_N390R_K392L_T394W
AZ592	L351Y_S400D_F405A_Y407I	T366L_N390R_K392M_T394W

AZ593	L351Y_S400D_F405A_Y407I	T366L_N390R_K392F_T394W
AZ594	L351Y_S400D_F405A_Y407I	T366L_N390K_K392L_T394W
AZ595	L351Y_S400D_F405A_Y407I	T366L_N390K_K392M_T394W
AZ596	L351Y_S400D_F405A_Y407I	T366L_N390K_K392F_T394W
AZ597	L351Y_S400D_F405A_Y407V	T366L_N390R_K392L_T394W
AZ598	L351Y_S400D_F405A_Y407V	T366L_N390R_K392M_T394W
AZ599	L351Y_S400D_F405A_Y407V	T366L_N390R_K392F_T394W
AZ600	L351Y_S400D_F405A_Y407V	T366L_N390K_K392L_T394W
AZ601	L351Y_S400D_F405A_Y407V	T366L_N390K_K392M_T394W
AZ602	L351Y_S400D_F405A_Y407V	T366L_N390K_K392F_T394W
AZ603	L351Y_S400D_F405S_Y407I	T366L_N390R_K392L_T394W
AZ604	L351Y_S400D_F405S_Y407I	T366L_N390R_K392M_T394W
AZ605	L351Y_S400D_F405S_Y407I	T366L_N390R_K392F_T394W
AZ606	L351Y_S400D_F405S_Y407I	T366L_N390K_K392L_T394W
AZ607	L351Y_S400D_F405S_Y407I	T366L_N390K_K392M_T394W
AZ608	L351Y_S400D_F405S_Y407I	T366L_N390K_K392F_T394W
AZ609	L351Y_S400D_F405S_Y407V	T366L_N390R_K392L_T394W
AZ610	L351Y_S400D_F405S_Y407V	T366L_N390R_K392M_T394W
AZ611	L351Y_S400D_F405S_Y407V	T366L_N390R_K392F_T394W
AZ612	L351Y_S400D_F405S_Y407V	T366L_N390K_K392L_T394W
AZ613	L351Y_S400D_F405S_Y407V	T366L_N390K_K392M_T394W
AZ614	L351Y_S400D_F405S_Y407V	T366L_N390K_K392F_T394W
AZ615	L351Y_S400D_F405T_Y407I	T366L_N390R_K392L_T394W
AZ616	L351Y_S400D_F405T_Y407I	T366L_N390R_K392M_T394W
AZ617	L351Y_S400D_F405T_Y407I	T366L_N390R_K392F_T394W
AZ618	L351Y_S400D_F405T_Y407I	T366L_N390K_K392L_T394W
AZ619	L351Y_S400D_F405T_Y407I	T366L_N390K_K392M_T394W
AZ620	L351Y_S400D_F405T_Y407I	T366L_N390K_K392F_T394W
AZ621	L351Y_S400D_F405T_Y407V	T366L_N390R_K392L_T394W
AZ622	L351Y_S400D_F405T_Y407V	T366L_N390R_K392M_T394W
AZ623	L351Y_S400D_F405T_Y407V	T366L_N390R_K392F_T394W
AZ624	L351Y_S400D_F405T_Y407V	T366L_N390K_K392L_T394W
AZ625	L351Y_S400D_F405T_Y407V	T366L_N390K_K392M_T394W
AZ626	L351Y_S400D_F405T_Y407V	T366L_N390K_K392F_T394W
AZ627	L351Y_S400D_F405V_Y407I	T366L_N390R_K392L_T394W
AZ628	L351Y_S400D_F405V_Y407I	T366L_N390R_K392M_T394W
AZ629	L351Y_S400D_F405V_Y407I	T366L_N390R_K392F_T394W
AZ630	L351Y_S400D_F405V_Y407I	T366L_N390K_K392L_T394W
AZ631	L351Y_S400D_F405V_Y407I	T366L_N390K_K392M_T394W
AZ632	L351Y_S400D_F405V_Y407I	T366L_N390K_K392F_T394W
AZ633	L351Y_S400D_F405V_Y407V	T366L_N390R_K392L_T394W
AZ634	L351Y_S400D_F405V_Y407V	T366L_N390R_K392M_T394W
AZ635	L351Y_S400D_F405V_Y407V	T366L_N390R_K392F_T394W
AZ636	L351Y_S400D_F405V_Y407V	T366L_N390K_K392L_T394W
AZ637	L351Y_S400D_F405V_Y407V	T366L_N390K_K392M_T394W
AZ638	L351Y_S400D_F405V_Y407V	T366L_N390K_K392F_T394W
AZ639	L351Y_S400E_F405A_Y407I	T366L_N390R_K392L_T394W
AZ640	L351Y_S400E_F405A_Y407I	T366L_N390R_K392M_T394W
AZ641	L351Y_S400E_F405A_Y407I	T366L_N390R_K392F_T394W
AZ642	L351Y_S400E_F405A_Y407I	T366L_N390K_K392L_T394W
AZ643	L351Y_S400E_F405A_Y407I	T366L_N390K_K392M_T394W
AZ644	L351Y_S400E_F405A_Y407I	T366L_N390K_K392F_T394W
AZ645	L351Y_S400E_F405A_Y407V	T366L_N390R_K392L_T394W
AZ646	L351Y_S400E_F405A_Y407V	T366L_N390R_K392M_T394W
AZ647	L351Y_S400E_F405A_Y407V	T366L_N390R_K392F_T394W
AZ648	L351Y_S400E_F405A_Y407V	T366L_N390K_K392L_T394W
AZ649	L351Y_S400E_F405A_Y407V	T366L_N390K_K392M_T394W
AZ650	L351Y_S400E_F405A_Y407V	T366L_N390K_K392F_T394W

AZ651	L351Y_S400E_F405S_Y407I	T366L_N390R_K392L_T394W
AZ652	L351Y_S400E_F405S_Y407I	T366L_N390R_K392M_T394W
AZ653	L351Y_S400E_F405S_Y407I	T366L_N390R_K392F_T394W
AZ654	L351Y_S400E_F405S_Y407I	T366L_N390K_K392L_T394W
AZ655	L351Y_S400E_F405S_Y407I	T366L_N390K_K392M_T394W
AZ656	L351Y_S400E_F405S_Y407I	T366L_N390K_K392F_T394W
AZ657	L351Y_S400E_F405S_Y407V	T366L_N390R_K392L_T394W
AZ658	L351Y_S400E_F405S_Y407V	T366L_N390R_K392M_T394W
AZ659	L351Y_S400E_F405S_Y407V	T366L_N390R_K392F_T394W
AZ660	L351Y_S400E_F405S_Y407V	T366L_N390K_K392L_T394W
AZ661	L351Y_S400E_F405S_Y407V	T366L_N390K_K392M_T394W
AZ662	L351Y_S400E_F405S_Y407V	T366L_N390K_K392F_T394W
AZ663	L351Y_S400E_F405T_Y407I	T366L_N390R_K392L_T394W
AZ664	L351Y_S400E_F405T_Y407I	T366L_N390R_K392M_T394W
AZ665	L351Y_S400E_F405T_Y407I	T366L_N390R_K392F_T394W
AZ666	L351Y_S400E_F405T_Y407I	T366L_N390K_K392L_T394W
AZ667	L351Y_S400E_F405T_Y407I	T366L_N390K_K392M_T394W
AZ668	L351Y_S400E_F405T_Y407I	T366L_N390K_K392F_T394W
AZ669	L351Y_S400E_F405T_Y407V	T366L_N390R_K392L_T394W
AZ670	L351Y_S400E_F405T_Y407V	T366L_N390R_K392M_T394W
AZ671	L351Y_S400E_F405T_Y407V	T366L_N390R_K392F_T394W
AZ672	L351Y_S400E_F405T_Y407V	T366L_N390K_K392L_T394W
AZ673	L351Y_S400E_F405T_Y407V	T366L_N390K_K392M_T394W
AZ674	L351Y_S400E_F405T_Y407V	T366L_N390K_K392F_T394W
AZ675	L351Y_S400E_F405V_Y407I	T366L_N390R_K392L_T394W
AZ676	L351Y_S400E_F405V_Y407I	T366L_N390R_K392M_T394W
AZ677	L351Y_S400E_F405V_Y407I	T366L_N390R_K392F_T394W
AZ678	L351Y_S400E_F405V_Y407I	T366L_N390K_K392L_T394W
AZ679	L351Y_S400E_F405V_Y407I	T366L_N390K_K392M_T394W
AZ680	L351Y_S400E_F405V_Y407I	T366L_N390K_K392F_T394W
AZ681	L351Y_S400E_F405V_Y407V	T366L_N390R_K392L_T394W
AZ682	L351Y_S400E_F405V_Y407V	T366L_N390R_K392M_T394W
AZ683	L351Y_S400E_F405V_Y407V	T366L_N390R_K392F_T394W
AZ684	L351Y_S400E_F405V_Y407V	T366L_N390K_K392L_T394W
AZ685	L351Y_S400E_F405V_Y407V	T366L_N390K_K392M_T394W
AZ686	L351Y_S400E_F405V_Y407V	T366L_N390K_K392F_T394W
AZ687	L351Y_F405A_Y407I	T366L_K392L_T394W_K409L_T411R
AZ688	L351Y_F405A_Y407I	T366L_K392L_T394W_K409L_T411K
AZ689	L351Y_F405A_Y407I	T366L_K392L_T394W_K409M_T411R
AZ690	L351Y_F405A_Y407I	T366L_K392L_T394W_K409M_T411K
AZ691	L351Y_F405A_Y407I	T366L_K392M_T394W_K409L_T411R
AZ692	L351Y_F405A_Y407I	T366L_K392M_T394W_K409L_T411K
AZ693	L351Y_F405A_Y407I	T366L_K392M_T394W_K409M_T411R
AZ694	L351Y_F405A_Y407I	T366L_K392M_T394W_K409M_T411K
AZ695	L351Y_F405A_Y407I	T366L_K392F_T394W_K409L_T411R
AZ696	L351Y_F405A_Y407I	T366L_K392F_T394W_K409L_T411K
AZ697	L351Y_F405A_Y407I	T366L_K392F_T394W_K409M_T411R
AZ698	L351Y_F405A_Y407I	T366L_K392F_T394W_K409M_T411K
AZ699	L351Y_F405A_Y407V	T366L_K392L_T394W_K409L_T411R
AZ700	L351Y_F405A_Y407V	T366L_K392L_T394W_K409L_T411K
AZ701	L351Y_F405A_Y407V	T366L_K392L_T394W_K409M_T411R
AZ702	L351Y_F405A_Y407V	T366L_K392L_T394W_K409M_T411K
AZ703	L351Y_F405A_Y407V	T366L_K392M_T394W_K409L_T411R
AZ704	L351Y_F405A_Y407V	T366L_K392M_T394W_K409L_T411K
AZ705	L351Y_F405A_Y407V	T366L_K392M_T394W_K409M_T411R
AZ706	L351Y_F405A_Y407V	T366L_K392M_T394W_K409M_T411K
AZ707	L351Y_F405A_Y407V	T366L_K392F_T394W_K409L_T411R
AZ708	L351Y_F405A_Y407V	T366L_K392F_T394W_K409L_T411K

AZ709	L351Y_F405A_Y407V	T366L_K392F_T394W_K409M_T411R
AZ710	L351Y_F405A_Y407V	T366L_K392F_T394W_K409M_T411K
AZ711	L351Y_F405S_Y407I	T366L_K392L_T394W_K409L_T411R
AZ712	L351Y_F405S_Y407I	T366L_K392L_T394W_K409L_T411K
AZ713	L351Y_F405S_Y407I	T366L_K392L_T394W_K409M_T411R
AZ714	L351Y_F405S_Y407I	T366L_K392L_T394W_K409M_T411K
AZ715	L351Y_F405S_Y407I	T366L_K392M_T394W_K409L_T411R
AZ716	L351Y_F405S_Y407I	T366L_K392M_T394W_K409L_T411K
AZ717	L351Y_F405S_Y407I	T366L_K392M_T394W_K409M_T411R
AZ718	L351Y_F405S_Y407I	T366L_K392M_T394W_K409M_T411K
AZ719	L351Y_F405S_Y407I	T366L_K392F_T394W_K409L_T411R
AZ720	L351Y_F405S_Y407I	T366L_K392F_T394W_K409L_T411K
AZ721	L351Y_F405S_Y407I	T366L_K392F_T394W_K409M_T411R
AZ722	L351Y_F405S_Y407I	T366L_K392F_T394W_K409M_T411K
AZ723	L351Y_F405S_Y407V	T366L_K392L_T394W_K409L_T411R
AZ724	L351Y_F405S_Y407V	T366L_K392L_T394W_K409L_T411K
AZ725	L351Y_F405S_Y407V	T366L_K392L_T394W_K409M_T411R
AZ726	L351Y_F405S_Y407V	T366L_K392L_T394W_K409M_T411K
AZ727	L351Y_F405S_Y407V	T366L_K392M_T394W_K409L_T411R
AZ728	L351Y_F405S_Y407V	T366L_K392M_T394W_K409L_T411K
AZ729	L351Y_F405S_Y407V	T366L_K392M_T394W_K409M_T411R
AZ730	L351Y_F405S_Y407V	T366L_K392M_T394W_K409M_T411K
AZ731	L351Y_F405S_Y407V	T366L_K392F_T394W_K409L_T411R
AZ732	L351Y_F405S_Y407V	T366L_K392F_T394W_K409L_T411K
AZ733	L351Y_F405S_Y407V	T366L_K392F_T394W_K409M_T411R
AZ734	L351Y_F405S_Y407V	T366L_K392F_T394W_K409M_T411K
AZ735	L351Y_F405V_Y407I	T366L_K392L_T394W_K409L_T411R
AZ736	L351Y_F405V_Y407I	T366L_K392L_T394W_K409L_T411K
AZ737	L351Y_F405V_Y407I	T366L_K392L_T394W_K409M_T411R
AZ738	L351Y_F405V_Y407I	T366L_K392L_T394W_K409M_T411K
AZ739	L351Y_F405V_Y407I	T366L_K392M_T394W_K409L_T411R
AZ740	L351Y_F405V_Y407I	T366L_K392M_T394W_K409L_T411K
AZ741	L351Y_F405V_Y407I	T366L_K392M_T394W_K409M_T411R
AZ742	L351Y_F405V_Y407I	T366L_K392M_T394W_K409M_T411K
AZ743	L351Y_F405V_Y407I	T366L_K392F_T394W_K409L_T411R
AZ744	L351Y_F405V_Y407I	T366L_K392F_T394W_K409L_T411K
AZ745	L351Y_F405V_Y407I	T366L_K392F_T394W_K409M_T411R
AZ746	L351Y_F405V_Y407I	T366L_K392F_T394W_K409M_T411K
AZ747	L351Y_F405V_Y407V	T366L_K392L_T394W_K409L_T411R
AZ748	L351Y_F405V_Y407V	T366L_K392L_T394W_K409L_T411K
AZ749	L351Y_F405V_Y407V	T366L_K392L_T394W_K409M_T411R
AZ750	L351Y_F405V_Y407V	T366L_K392L_T394W_K409M_T411K
AZ751	L351Y_F405V_Y407V	T366L_K392M_T394W_K409L_T411R
AZ752	L351Y_F405V_Y407V	T366L_K392M_T394W_K409L_T411K
AZ753	L351Y_F405V_Y407V	T366L_K392M_T394W_K409M_T411R
AZ754	L351Y_F405V_Y407V	T366L_K392M_T394W_K409M_T411K
AZ755	L351Y_F405V_Y407V	T366L_K392F_T394W_K409L_T411R
AZ756	L351Y_F405V_Y407V	T366L_K392F_T394W_K409L_T411K
AZ757	L351Y_F405V_Y407V	T366L_K392F_T394W_K409M_T411R
AZ758	L351Y_F405V_Y407V	T366L_K392F_T394W_K409M_T411K
AZ759	D399C_F405A_Y407I	T366L_K392C_T394W_K409L_T411L
AZ760	D399C_F405A_Y407I	T366L_K392C_T394W_K409L_T411T
AZ761	D399C_F405A_Y407I	T366L_K392C_T394W_K409L_T411L
AZ762	D399C_F405A_Y407I	T366L_K392C_T394W_K409L_T411T
AZ763	D399C_F405A_Y407I	T366L_K392C_T394W_K409K_T411L
AZ764	D399C_F405A_Y407I	T366L_K392C_T394W_K409K_T411T
AZ765	D399C_F405A_Y407I	T366L_K392C_T394W_K409M_T411L
AZ766	D399C_F405A_Y407I	T366L_K392C_T394W_K409M_T411T

AZ767	D399C_F405A_Y407V	T366L_K392C_T394W_K409I_T411L
AZ768	D399C_F405A_Y407V	T366L_K392C_T394W_K409I_T411T
AZ769	D399C_F405A_Y407V	T366L_K392C_T394W_K409L_T411L
AZ770	D399C_F405A_Y407V	T366L_K392C_T394W_K409L_T411T
AZ771	D399C_F405A_Y407V	T366L_K392C_T394W_K409K_T411L
AZ772	D399C_F405A_Y407V	T366L_K392C_T394W_K409K_T411T
AZ773	D399C_F405A_Y407V	T366L_K392C_T394W_K409M_T411L
AZ774	D399C_F405A_Y407V	T366L_K392C_T394W_K409M_T411T
AZ775	D399C_F405S_Y407I	T366L_K392C_T394W_K409I_T411L
AZ776	D399C_F405S_Y407I	T366L_K392C_T394W_K409I_T411T
AZ777	D399C_F405S_Y407I	T366L_K392C_T394W_K409L_T411L
AZ778	D399C_F405S_Y407I	T366L_K392C_T394W_K409L_T411T
AZ779	D399C_F405S_Y407I	T366L_K392C_T394W_K409K_T411L
AZ780	D399C_F405S_Y407I	T366L_K392C_T394W_K409K_T411T
AZ781	D399C_F405S_Y407I	T366L_K392C_T394W_K409M_T411L
AZ782	D399C_F405S_Y407I	T366L_K392C_T394W_K409M_T411T
AZ783	D399C_F405S_Y407V	T366L_K392C_T394W_K409I_T411L
AZ784	D399C_F405S_Y407V	T366L_K392C_T394W_K409I_T411T
AZ785	D399C_F405S_Y407V	T366L_K392C_T394W_K409L_T411L
AZ786	D399C_F405S_Y407V	T366L_K392C_T394W_K409L_T411T
AZ787	D399C_F405S_Y407V	T366L_K392C_T394W_K409K_T411L
AZ788	D399C_F405S_Y407V	T366L_K392C_T394W_K409K_T411T
AZ789	D399C_F405S_Y407V	T366L_K392C_T394W_K409M_T411L
AZ790	D399C_F405S_Y407V	T366L_K392C_T394W_K409M_T411T
AZ791	D399C_F405V_Y407I	T366L_K392C_T394W_K409I_T411L
AZ792	D399C_F405V_Y407I	T366L_K392C_T394W_K409I_T411T
AZ793	D399C_F405V_Y407I	T366L_K392C_T394W_K409L_T411L
AZ794	D399C_F405V_Y407I	T366L_K392C_T394W_K409L_T411T
AZ795	D399C_F405V_Y407I	T366L_K392C_T394W_K409K_T411L
AZ796	D399C_F405V_Y407I	T366L_K392C_T394W_K409K_T411T
AZ797	D399C_F405V_Y407I	T366L_K392C_T394W_K409M_T411L
AZ798	D399C_F405V_Y407I	T366L_K392C_T394W_K409M_T411T
AZ799	D399C_F405V_Y407V	T366L_K392C_T394W_K409I_T411L
AZ800	D399C_F405V_Y407V	T366L_K392C_T394W_K409I_T411T
AZ801	D399C_F405V_Y407V	T366L_K392C_T394W_K409L_T411L
AZ802	D399C_F405V_Y407V	T366L_K392C_T394W_K409L_T411T
AZ803	D399C_F405V_Y407V	T366L_K392C_T394W_K409K_T411L
AZ804	D399C_F405V_Y407V	T366L_K392C_T394W_K409K_T411T
AZ805	D399C_F405V_Y407V	T366L_K392C_T394W_K409M_T411L
AZ806	D399C_F405V_Y407V	T366L_K392C_T394W_K409M_T411T
AZ807	L351L_D399R_F405A_Y407I	T366L_K392L_T394W_K409I_T411E
AZ808	L351L_D399R_F405A_Y407I	T366L_K392L_T394W_K409M_T411E
AZ809	L351L_D399R_F405A_Y407I	T366L_K392M_T394W_K409L_T411E
AZ810	L351L_D399R_F405A_Y407I	T366L_K392M_T394W_K409M_T411E
AZ811	L351L_D399R_F405A_Y407I	T366L_K392F_T394W_K409L_T411E
AZ812	L351L_D399R_F405A_Y407I	T366L_K392F_T394W_K409M_T411E
AZ813	L351L_D399R_F405A_Y407V	T366L_K392L_T394W_K409L_T411E
AZ814	L351L_D399R_F405A_Y407V	T366L_K392L_T394W_K409M_T411E
AZ815	L351L_D399R_F405A_Y407V	T366L_K392M_T394W_K409L_T411E
AZ816	L351L_D399R_F405A_Y407V	T366L_K392M_T394W_K409M_T411E
AZ817	L351L_D399R_F405A_Y407V	T366L_K392F_T394W_K409L_T411E
AZ818	L351L_D399R_F405A_Y407V	T366L_K392F_T394W_K409M_T411E
AZ819	L351L_D399R_F405M_Y407I	T366L_K392L_T394W_K409V_T411E
AZ820	L351L_D399R_F405M_Y407I	T366L_K392M_T394W_K409V_T411E
AZ821	L351L_D399R_F405M_Y407I	T366L_K392F_T394W_K409V_T411E
AZ822	L351L_D399R_F405M_Y407V	T366L_K392L_T394W_K409V_T411E
AZ823	L351L_D399R_F405M_Y407V	T366L_K392M_T394W_K409V_T411E
AZ824	L351L_D399R_F405M_Y407V	T366L_K392F_T394W_K409V_T411E

AZ825	L351L_D399R_F405S_Y407I	T366L_K392L_T394W_K409L_T411E
AZ826	L351L_D399R_F405S_Y407I	T366L_K392L_T394W_K409M_T411E
AZ827	L351L_D399R_F405S_Y407I	T366L_K392M_T394W_K409L_T411E
AZ828	L351L_D399R_F405S_Y407I	T366L_K392M_T394W_K409M_T411E
AZ829	L351L_D399R_F405S_Y407I	T366L_K392F_T394W_K409L_T411E
AZ830	L351L_D399R_F405S_Y407I	T366L_K392F_T394W_K409M_T411E
AZ831	L351L_D399R_F405S_Y407V	T366L_K392L_T394W_K409L_T411E
AZ832	L351L_D399R_F405S_Y407V	T366L_K392L_T394W_K409M_T411E
AZ833	L351L_D399R_F405S_Y407V	T366L_K392M_T394W_K409L_T411E
AZ834	L351L_D399R_F405S_Y407V	T366L_K392M_T394W_K409M_T411E
AZ835	L351L_D399R_F405S_Y407V	T366L_K392F_T394W_K409L_T411E
AZ836	L351L_D399R_F405S_Y407V	T366L_K392F_T394W_K409M_T411E
AZ837	L351L_D399R_F405V_Y407I	T366L_K392L_T394W_K409L_T411E
AZ838	L351L_D399R_F405V_Y407I	T366L_K392L_T394W_K409M_T411E
AZ839	L351L_D399R_F405V_Y407I	T366L_K392M_T394W_K409L_T411E
AZ840	L351L_D399R_F405V_Y407I	T366L_K392M_T394W_K409M_T411E
AZ841	L351L_D399R_F405V_Y407I	T366L_K392F_T394W_K409L_T411E
AZ842	L351L_D399R_F405V_Y407I	T366L_K392F_T394W_K409M_T411E
AZ843	L351L_D399R_F405V_Y407V	T366L_K392L_T394W_K409L_T411E
AZ844	L351L_D399R_F405V_Y407V	T366L_K392L_T394W_K409M_T411E
AZ845	L351L_D399R_F405V_Y407V	T366L_K392M_T394W_K409L_T411E
AZ846	L351L_D399R_F405V_Y407V	T366L_K392M_T394W_K409M_T411E
AZ847	L351L_D399R_F405V_Y407V	T366L_K392F_T394W_K409L_T411E
AZ848	L351L_D399R_F405V_Y407V	T366L_K392F_T394W_K409M_T411E
AZ849	L351L_S400R_F405A_Y407I	T366L_N390N_K392L_T394W_T411N
AZ850	L351L_S400R_F405A_Y407I	T366L_N390N_K392L_T394W_T411S
AZ851	L351L_S400R_F405A_Y407I	T366L_N390N_K392L_T394W_T411T
AZ852	L351L_S400R_F405A_Y407I	T366L_N390N_K392M_T394W_T411N
AZ853	L351L_S400R_F405A_Y407I	T366L_N390N_K392M_T394W_T411S
AZ854	L351L_S400R_F405A_Y407I	T366L_N390N_K392M_T394W_T411T
AZ855	L351L_S400R_F405A_Y407I	T366L_N390N_K392F_T394W_T411N
AZ856	L351L_S400R_F405A_Y407I	T366L_N390N_K392F_T394W_T411S
AZ857	L351L_S400R_F405A_Y407I	T366L_N390N_K392F_T394W_T411T
AZ858	L351L_S400R_F405A_Y407I	T366L_N390D_K392L_T394W_T411N
AZ859	L351L_S400R_F405A_Y407I	T366L_N390D_K392L_T394W_T411S
AZ860	L351L_S400R_F405A_Y407I	T366L_N390D_K392L_T394W_T411T
AZ861	L351L_S400R_F405A_Y407I	T366L_N390D_K392M_T394W_T411N
AZ862	L351L_S400R_F405A_Y407I	T366L_N390D_K392M_T394W_T411S
AZ863	L351L_S400R_F405A_Y407I	T366L_N390D_K392M_T394W_T411T
AZ864	L351L_S400R_F405A_Y407I	T366L_N390D_K392F_T394W_T411N
AZ865	L351L_S400R_F405A_Y407I	T366L_N390D_K392F_T394W_T411S
AZ866	L351L_S400R_F405A_Y407I	T366L_N390D_K392F_T394W_T411T
AZ867	L351L_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411N
AZ868	L351L_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411L
AZ869	L351L_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411S
AZ870	L351L_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411T
AZ871	L351L_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411N
AZ872	L351L_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411L
AZ873	L351L_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411S
AZ874	L351L_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411T
AZ875	L351L_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411N
AZ876	L351L_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411L
AZ877	L351L_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411S
AZ878	L351L_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411T
AZ879	L351L_S400R_F405A_Y407V	T366L_N390N_K392L_T394W_T411N
AZ880	L351L_S400R_F405A_Y407V	T366L_N390N_K392L_T394W_T411S
AZ881	L351L_S400R_F405A_Y407V	T366L_N390N_K392L_T394W_T411T
AZ882	L351L_S400R_F405A_Y407V	T366L_N390N_K392M_T394W_T411N

AZ883	L351L_S400R_F405A_Y407V	T366L_N390N_K392M_T394W_T411S
AZ884	L351L_S400R_F405A_Y407V	T366L_N390N_K392M_T394W_T411T
AZ885	L351L_S400R_F405A_Y407V	T366L_N390N_K392F_T394W_T411N
AZ886	L351L_S400R_F405A_Y407V	T366L_N390N_K392F_T394W_T411S
AZ887	L351L_S400R_F405A_Y407V	T366L_N390N_K392F_T394W_T411T
AZ888	L351L_S400R_F405A_Y407V	T366L_N390D_K392L_T394W_T411N
AZ889	L351L_S400R_F405A_Y407V	T366L_N390D_K392L_T394W_T411S
AZ890	L351L_S400R_F405A_Y407V	T366L_N390D_K392L_T394W_T411T
AZ891	L351L_S400R_F405A_Y407V	T366L_N390D_K392M_T394W_T411N
AZ892	L351L_S400R_F405A_Y407V	T366L_N390D_K392M_T394W_T411S
AZ893	L351L_S400R_F405A_Y407V	T366L_N390D_K392M_T394W_T411T
AZ894	L351L_S400R_F405A_Y407V	T366L_N390D_K392F_T394W_T411N
AZ895	L351L_S400R_F405A_Y407V	T366L_N390D_K392F_T394W_T411S
AZ896	L351L_S400R_F405A_Y407V	T366L_N390D_K392F_T394W_T411T
AZ897	L351L_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411N
AZ898	L351L_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411L
AZ899	L351L_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411S
AZ900	L351L_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411T
AZ901	L351L_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411N
AZ902	L351L_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411L
AZ903	L351L_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411S
AZ904	L351L_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411T
AZ905	L351L_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411N
AZ906	L351L_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411L
AZ907	L351L_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411S
AZ908	L351L_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411T
AZ909	L351L_S400R_F405S_Y407I	T366L_N390N_K392L_T394W_T411N
AZ910	L351L_S400R_F405S_Y407I	T366L_N390N_K392L_T394W_T411S
AZ911	L351L_S400R_F405S_Y407I	T366L_N390N_K392L_T394W_T411T
AZ912	L351L_S400R_F405S_Y407I	T366L_N390N_K392M_T394W_T411N
AZ913	L351L_S400R_F405S_Y407I	T366L_N390N_K392M_T394W_T411S
AZ914	L351L_S400R_F405S_Y407I	T366L_N390N_K392M_T394W_T411T
AZ915	L351L_S400R_F405S_Y407I	T366L_N390N_K392F_T394W_T411N
AZ916	L351L_S400R_F405S_Y407I	T366L_N390N_K392F_T394W_T411S
AZ917	L351L_S400R_F405S_Y407I	T366L_N390N_K392F_T394W_T411T
AZ918	L351L_S400R_F405S_Y407I	T366L_N390D_K392L_T394W_T411N
AZ919	L351L_S400R_F405S_Y407I	T366L_N390D_K392L_T394W_T411S
AZ920	L351L_S400R_F405S_Y407I	T366L_N390D_K392L_T394W_T411T
AZ921	L351L_S400R_F405S_Y407I	T366L_N390D_K392M_T394W_T411N
AZ922	L351L_S400R_F405S_Y407I	T366L_N390D_K392M_T394W_T411S
AZ923	L351L_S400R_F405S_Y407I	T366L_N390D_K392M_T394W_T411T
AZ924	L351L_S400R_F405S_Y407I	T366L_N390D_K392F_T394W_T411N
AZ925	L351L_S400R_F405S_Y407I	T366L_N390D_K392F_T394W_T411S
AZ926	L351L_S400R_F405S_Y407I	T366L_N390D_K392F_T394W_T411T
AZ927	L351L_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411N
AZ928	L351L_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411L
AZ929	L351L_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411S
AZ930	L351L_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411T
AZ931	L351L_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411N
AZ932	L351L_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411L
AZ933	L351L_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411S
AZ934	L351L_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411T
AZ935	L351L_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411N
AZ936	L351L_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411L
AZ937	L351L_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411S
AZ938	L351L_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411T
AZ939	L351L_S400R_F405S_Y407V	T366L_N390N_K392L_T394W_T411N
AZ940	L351L_S400R_F405S_Y407V	T366L_N390N_K392L_T394W_T411S

AZ941	L351L_S400R_F405S_Y407V	T366L_N390N_K392L_T394W_T411T
AZ942	L351L_S400R_F405S_Y407V	T366L_N390N_K392M_T394W_T411N
AZ943	L351L_S400R_F405S_Y407V	T366L_N390N_K392M_T394W_T411S
AZ944	L351L_S400R_F405S_Y407V	T366L_N390N_K392M_T394W_T411T
AZ945	L351L_S400R_F405S_Y407V	T366L_N390N_K392F_T394W_T411N
AZ946	L351L_S400R_F405S_Y407V	T366L_N390N_K392F_T394W_T411S
AZ947	L351L_S400R_F405S_Y407V	T366L_N390N_K392F_T394W_T411T
AZ948	L351L_S400R_F405S_Y407V	T366L_N390D_K392L_T394W_T411N
AZ949	L351L_S400R_F405S_Y407V	T366L_N390D_K392L_T394W_T411S
AZ950	L351L_S400R_F405S_Y407V	T366L_N390D_K392L_T394W_T411T
AZ951	L351L_S400R_F405S_Y407V	T366L_N390D_K392M_T394W_T411N
AZ952	L351L_S400R_F405S_Y407V	T366L_N390D_K392M_T394W_T411S
AZ953	L351L_S400R_F405S_Y407V	T366L_N390D_K392M_T394W_T411T
AZ954	L351L_S400R_F405S_Y407V	T366L_N390D_K392F_T394W_T411N
AZ955	L351L_S400R_F405S_Y407V	T366L_N390D_K392F_T394W_T411S
AZ956	L351L_S400R_F405S_Y407V	T366L_N390D_K392F_T394W_T411T
AZ957	L351L_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411N
AZ958	L351L_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411L
AZ959	L351L_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411S
AZ960	L351L_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411T
AZ961	L351L_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411N
AZ962	L351L_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411L
AZ963	L351L_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411S
AZ964	L351L_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411T
AZ965	L351L_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411N
AZ966	L351L_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411L
AZ967	L351L_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411S
AZ968	L351L_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411T
AZ969	L351L_S400R_F405T_Y407I	T366L_N390N_K392L_T394W_T411N
AZ970	L351L_S400R_F405T_Y407I	T366L_N390N_K392L_T394W_T411S
AZ971	L351L_S400R_F405T_Y407I	T366L_N390N_K392L_T394W_T411T
AZ972	L351L_S400R_F405T_Y407I	T366L_N390N_K392M_T394W_T411N
AZ973	L351L_S400R_F405T_Y407I	T366L_N390N_K392M_T394W_T411S
AZ974	L351L_S400R_F405T_Y407I	T366L_N390N_K392M_T394W_T411T
AZ975	L351L_S400R_F405T_Y407I	T366L_N390N_K392F_T394W_T411N
AZ976	L351L_S400R_F405T_Y407I	T366L_N390N_K392F_T394W_T411S
AZ977	L351L_S400R_F405T_Y407I	T366L_N390N_K392F_T394W_T411T
AZ978	L351L_S400R_F405T_Y407I	T366L_N390D_K392L_T394W_T411N
AZ979	L351L_S400R_F405T_Y407I	T366L_N390D_K392L_T394W_T411S
AZ980	L351L_S400R_F405T_Y407I	T366L_N390D_K392L_T394W_T411T
AZ981	L351L_S400R_F405T_Y407I	T366L_N390D_K392M_T394W_T411N
AZ982	L351L_S400R_F405T_Y407I	T366L_N390D_K392M_T394W_T411S
AZ983	L351L_S400R_F405T_Y407I	T366L_N390D_K392M_T394W_T411T
AZ984	L351L_S400R_F405T_Y407I	T366L_N390D_K392F_T394W_T411N
AZ985	L351L_S400R_F405T_Y407I	T366L_N390D_K392F_T394W_T411S
AZ986	L351L_S400R_F405T_Y407I	T366L_N390D_K392F_T394W_T411T
AZ987	L351L_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411N
AZ988	L351L_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411L
AZ989	L351L_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411S
AZ990	L351L_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411T
AZ991	L351L_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411N
AZ992	L351L_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411L
AZ993	L351L_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411S
AZ994	L351L_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411T
AZ995	L351L_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411N
AZ996	L351L_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411L
AZ997	L351L_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411S
AZ998	L351L_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411T

AZ999	L351L_S400R_F405T_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1000	L351L_S400R_F405T_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1001	L351L_S400R_F405T_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1002	L351L_S400R_F405T_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1003	L351L_S400R_F405T_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1004	L351L_S400R_F405T_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1005	L351L_S400R_F405T_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1006	L351L_S400R_F405T_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1007	L351L_S400R_F405T_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1008	L351L_S400R_F405T_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1009	L351L_S400R_F405T_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1010	L351L_S400R_F405T_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1011	L351L_S400R_F405T_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1012	L351L_S400R_F405T_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1013	L351L_S400R_F405T_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1014	L351L_S400R_F405T_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1015	L351L_S400R_F405T_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1016	L351L_S400R_F405T_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1017	L351L_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1018	L351L_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1019	L351L_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1020	L351L_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1021	L351L_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1022	L351L_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1023	L351L_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1024	L351L_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1025	L351L_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1026	L351L_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1027	L351L_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1028	L351L_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1029	L351L_S400R_F405V_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1030	L351L_S400R_F405V_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1031	L351L_S400R_F405V_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1032	L351L_S400R_F405V_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1033	L351L_S400R_F405V_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1034	L351L_S400R_F405V_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1035	L351L_S400R_F405V_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1036	L351L_S400R_F405V_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1037	L351L_S400R_F405V_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1038	L351L_S400R_F405V_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1039	L351L_S400R_F405V_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1040	L351L_S400R_F405V_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1041	L351L_S400R_F405V_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1042	L351L_S400R_F405V_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1043	L351L_S400R_F405V_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1044	L351L_S400R_F405V_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1045	L351L_S400R_F405V_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1046	L351L_S400R_F405V_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1047	L351L_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1048	L351L_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1049	L351L_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1050	L351L_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1051	L351L_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1052	L351L_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1053	L351L_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1054	L351L_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1055	L351L_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1056	L351L_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411N

AZ1057	L351L_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1058	L351L_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1059	L351L_S400R_F405V_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1060	L351L_S400R_F405V_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1061	L351L_S400R_F405V_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1062	L351L_S400R_F405V_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1063	L351L_S400R_F405V_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1064	L351L_S400R_F405V_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1065	L351L_S400R_F405V_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1066	L351L_S400R_F405V_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1067	L351L_S400R_F405V_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1068	L351L_S400R_F405V_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1069	L351L_S400R_F405V_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1070	L351L_S400R_F405V_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1071	L351L_S400R_F405V_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1072	L351L_S400R_F405V_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1073	L351L_S400R_F405V_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1074	L351L_S400R_F405V_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1075	L351L_S400R_F405V_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1076	L351L_S400R_F405V_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1077	L351L_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1078	L351L_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1079	L351L_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1080	L351L_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1081	L351L_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1082	L351L_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1083	L351L_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1084	L351L_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1085	L351L_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1086	L351L_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1087	L351L_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1088	L351L_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1089	L351L_S400E_F405A_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1090	L351L_S400E_F405A_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1091	L351L_S400E_F405A_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1092	L351L_S400E_F405A_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1093	L351L_S400E_F405A_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1094	L351L_S400E_F405A_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1095	L351L_S400E_F405A_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1096	L351L_S400E_F405A_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1097	L351L_S400E_F405A_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1098	L351L_S400E_F405A_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1099	L351L_S400E_F405A_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1100	L351L_S400E_F405A_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1101	L351L_S400E_F405A_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1102	L351L_S400E_F405A_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1103	L351L_S400E_F405A_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1104	L351L_S400E_F405A_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1105	L351L_S400E_F405A_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1106	L351L_S400E_F405A_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1107	L351L_S400E_F405A_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1108	L351L_S400E_F405A_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1109	L351L_S400E_F405A_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1110	L351L_S400E_F405A_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1111	L351L_S400E_F405A_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1112	L351L_S400E_F405A_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1113	L351L_S400E_F405A_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1114	L351L_S400E_F405A_Y407I	T366L_N390K_K392F_T394W_T411L

AZ1115	L351L_S400E_F405A_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1116	L351L_S400E_F405A_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1117	L351L_S400E_F405A_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1118	L351L_S400E_F405A_Y407V	T366L_N390R_K392L_T394W_T411S
AZ1119	L351L_S400E_F405A_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1120	L351L_S400E_F405A_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1121	L351L_S400E_F405A_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1122	L351L_S400E_F405A_Y407V	T366L_N390R_K392F_T394W_T411N
AZ1123	L351L_S400E_F405A_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1124	L351L_S400E_F405A_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1125	L351L_S400E_F405A_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1126	L351L_S400E_F405A_Y407V	T366L_N390N_K392L_T394W_T411L
AZ1127	L351L_S400E_F405A_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1128	L351L_S400E_F405A_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1129	L351L_S400E_F405A_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1130	L351L_S400E_F405A_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1131	L351L_S400E_F405A_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1132	L351L_S400E_F405A_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1133	L351L_S400E_F405A_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1134	L351L_S400E_F405A_Y407V	T366L_N390K_K392L_T394W_T411N
AZ1135	L351L_S400E_F405A_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1136	L351L_S400E_F405A_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1137	L351L_S400E_F405A_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1138	L351L_S400E_F405A_Y407V	T366L_N390K_K392M_T394W_T411L
AZ1139	L351L_S400E_F405A_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1140	L351L_S400E_F405A_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1141	L351L_S400E_F405A_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1142	L351L_S400E_F405A_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1143	L351L_S400E_F405S_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1144	L351L_S400E_F405S_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1145	L351L_S400E_F405S_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1146	L351L_S400E_F405S_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1147	L351L_S400E_F405S_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1148	L351L_S400E_F405S_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1149	L351L_S400E_F405S_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1150	L351L_S400E_F405S_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1151	L351L_S400E_F405S_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1152	L351L_S400E_F405S_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1153	L351L_S400E_F405S_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1154	L351L_S400E_F405S_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1155	L351L_S400E_F405S_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1156	L351L_S400E_F405S_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1157	L351L_S400E_F405S_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1158	L351L_S400E_F405S_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1159	L351L_S400E_F405S_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1160	L351L_S400E_F405S_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1161	L351L_S400E_F405S_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1162	L351L_S400E_F405S_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1163	L351L_S400E_F405S_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1164	L351L_S400E_F405S_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1165	L351L_S400E_F405S_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1166	L351L_S400E_F405S_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1167	L351L_S400E_F405S_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1168	L351L_S400E_F405S_Y407I	T366L_N390K_K392F_T394W_T411L
AZ1169	L351L_S400E_F405S_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1170	L351L_S400E_F405S_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1171	L351L_S400E_F405S_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1172	L351L_S400E_F405S_Y407V	T366L_N390R_K392L_T394W_T411S

AZ1173	L351L_S400E_F405S_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1174	L351L_S400E_F405S_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1175	L351L_S400E_F405S_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1176	L351L_S400E_F405S_Y407V	T366L_N390R_K392F_T394W_T411N
AZ1177	L351L_S400E_F405S_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1178	L351L_S400E_F405S_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1179	L351L_S400E_F405S_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1180	L351L_S400E_F405S_Y407V	T366L_N390N_K392L_T394W_T411L
AZ1181	L351L_S400E_F405S_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1182	L351L_S400E_F405S_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1183	L351L_S400E_F405S_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1184	L351L_S400E_F405S_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1185	L351L_S400E_F405S_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1186	L351L_S400E_F405S_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1187	L351L_S400E_F405S_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1188	L351L_S400E_F405S_Y407V	T366L_N390K_K392L_T394W_T411N
AZ1189	L351L_S400E_F405S_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1190	L351L_S400E_F405S_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1191	L351L_S400E_F405S_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1192	L351L_S400E_F405S_Y407V	T366L_N390K_K392M_T394W_T411L
AZ1193	L351L_S400E_F405S_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1194	L351L_S400E_F405S_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1195	L351L_S400E_F405S_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1196	L351L_S400E_F405S_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1197	L351L_S400E_F405T_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1198	L351L_S400E_F405T_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1199	L351L_S400E_F405T_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1200	L351L_S400E_F405T_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1201	L351L_S400E_F405T_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1202	L351L_S400E_F405T_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1203	L351L_S400E_F405T_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1204	L351L_S400E_F405T_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1205	L351L_S400E_F405T_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1206	L351L_S400E_F405T_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1207	L351L_S400E_F405T_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1208	L351L_S400E_F405T_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1209	L351L_S400E_F405T_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1210	L351L_S400E_F405T_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1211	L351L_S400E_F405T_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1212	L351L_S400E_F405T_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1213	L351L_S400E_F405T_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1214	L351L_S400E_F405T_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1215	L351L_S400E_F405T_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1216	L351L_S400E_F405T_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1217	L351L_S400E_F405T_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1218	L351L_S400E_F405T_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1219	L351L_S400E_F405T_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1220	L351L_S400E_F405T_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1221	L351L_S400E_F405T_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1222	L351L_S400E_F405T_Y407I	T366L_N390K_K392F_T394W_T411L
AZ1223	L351L_S400E_F405T_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1224	L351L_S400E_F405T_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1225	L351L_S400E_F405T_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1226	L351L_S400E_F405T_Y407V	T366L_N390R_K392L_T394W_T411S
AZ1227	L351L_S400E_F405T_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1228	L351L_S400E_F405T_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1229	L351L_S400E_F405T_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1230	L351L_S400E_F405T_Y407V	T366L_N390R_K392F_T394W_T411N

AZ1231	L351L_S400E_F405T_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1232	L351L_S400E_F405T_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1233	L351L_S400E_F405T_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1234	L351L_S400E_F405T_Y407V	T366L_N390N_K392L_T394W_T411L
AZ1235	L351L_S400E_F405T_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1236	L351L_S400E_F405T_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1237	L351L_S400E_F405T_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1238	L351L_S400E_F405T_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1239	L351L_S400E_F405T_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1240	L351L_S400E_F405T_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1241	L351L_S400E_F405T_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1242	L351L_S400E_F405T_Y407V	T366L_N390K_K392L_T394W_T411N
AZ1243	L351L_S400E_F405T_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1244	L351L_S400E_F405T_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1245	L351L_S400E_F405T_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1246	L351L_S400E_F405T_Y407V	T366L_N390K_K392M_T394W_T411L
AZ1247	L351L_S400E_F405T_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1248	L351L_S400E_F405T_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1249	L351L_S400E_F405T_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1250	L351L_S400E_F405T_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1251	L351L_S400E_F405V_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1252	L351L_S400E_F405V_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1253	L351L_S400E_F405V_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1254	L351L_S400E_F405V_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1255	L351L_S400E_F405V_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1256	L351L_S400E_F405V_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1257	L351L_S400E_F405V_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1258	L351L_S400E_F405V_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1259	L351L_S400E_F405V_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1260	L351L_S400E_F405V_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1261	L351L_S400E_F405V_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1262	L351L_S400E_F405V_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1263	L351L_S400E_F405V_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1264	L351L_S400E_F405V_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1265	L351L_S400E_F405V_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1266	L351L_S400E_F405V_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1267	L351L_S400E_F405V_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1268	L351L_S400E_F405V_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1269	L351L_S400E_F405V_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1270	L351L_S400E_F405V_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1271	L351L_S400E_F405V_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1272	L351L_S400E_F405V_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1273	L351L_S400E_F405V_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1274	L351L_S400E_F405V_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1275	L351L_S400E_F405V_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1276	L351L_S400E_F405V_Y407I	T366L_N390K_K392F_T394W_T411L
AZ1277	L351L_S400E_F405V_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1278	L351L_S400E_F405V_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1279	L351L_S400E_F405V_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1280	L351L_S400E_F405V_Y407V	T366L_N390R_K392L_T394W_T411S
AZ1281	L351L_S400E_F405V_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1282	L351L_S400E_F405V_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1283	L351L_S400E_F405V_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1284	L351L_S400E_F405V_Y407V	T366L_N390R_K392F_T394W_T411N
AZ1285	L351L_S400E_F405V_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1286	L351L_S400E_F405V_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1287	L351L_S400E_F405V_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1288	L351L_S400E_F405V_Y407V	T366L_N390N_K392L_T394W_T411L

AZ1289	L351L_S400E_F405V_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1290	L351L_S400E_F405V_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1291	L351L_S400E_F405V_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1292	L351L_S400E_F405V_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1293	L351L_S400E_F405V_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1294	L351L_S400E_F405V_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1295	L351L_S400E_F405V_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1296	L351L_S400E_F405V_Y407V	T366L_N390K_K392L_T394W_T411N
AZ1297	L351L_S400E_F405V_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1298	L351L_S400E_F405V_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1299	L351L_S400E_F405V_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1300	L351L_S400E_F405V_Y407V	T366L_N390K_K392M_T394W_T411L
AZ1301	L351L_S400E_F405V_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1302	L351L_S400E_F405V_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1303	L351L_S400E_F405V_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1304	L351L_S400E_F405V_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1305	L351L_S400K_F405A_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1306	L351L_S400K_F405A_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1307	L351L_S400K_F405A_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1308	L351L_S400K_F405A_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1309	L351L_S400K_F405A_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1310	L351L_S400K_F405A_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1311	L351L_S400K_F405A_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1312	L351L_S400K_F405A_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1313	L351L_S400K_F405A_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1314	L351L_S400K_F405A_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1315	L351L_S400K_F405A_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1316	L351L_S400K_F405A_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1317	L351L_S400K_F405A_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1318	L351L_S400K_F405A_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1319	L351L_S400K_F405A_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1320	L351L_S400K_F405A_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1321	L351L_S400K_F405A_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1322	L351L_S400K_F405A_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1323	L351L_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1324	L351L_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1325	L351L_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1326	L351L_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1327	L351L_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1328	L351L_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1329	L351L_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1330	L351L_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1331	L351L_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1332	L351L_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1333	L351L_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1334	L351L_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1335	L351L_S400K_F405A_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1336	L351L_S400K_F405A_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1337	L351L_S400K_F405A_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1338	L351L_S400K_F405A_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1339	L351L_S400K_F405A_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1340	L351L_S400K_F405A_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1341	L351L_S400K_F405A_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1342	L351L_S400K_F405A_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1343	L351L_S400K_F405A_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1344	L351L_S400K_F405A_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1345	L351L_S400K_F405A_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1346	L351L_S400K_F405A_Y407V	T366L_N390D_K392L_T394W_T411T

AZ1347	L351L S400K F405A Y407V	T366L N390D K392M T394W T411N
AZ1348	L351L S400K F405A Y407V	T366L N390D K392M T394W T411S
AZ1349	L351L S400K F405A Y407V	T366L N390D K392M T394W T411T
AZ1350	L351L S400K F405A Y407V	T366L N390D K392F T394W T411N
AZ1351	L351L S400K F405A Y407V	T366L N390D K392F T394W T411S
AZ1352	L351L S400K F405A Y407V	T366L N390D K392F T394W T411T
AZ1353	L351L S400K F405A Y407V	T366L N390E K392L T394W T411N
AZ1354	L351L S400K F405A Y407V	T366L N390E K392L T394W T411L
AZ1355	L351L S400K F405A Y407V	T366L N390E K392L T394W T411S
AZ1356	L351L S400K F405A Y407V	T366L N390E K392L T394W T411T
AZ1357	L351L S400K F405A Y407V	T366L N390E K392M T394W T411N
AZ1358	L351L S400K F405A Y407V	T366L N390E K392M T394W T411L
AZ1359	L351L S400K F405A Y407V	T366L N390E K392M T394W T411S
AZ1360	L351L S400K F405A Y407V	T366L N390E K392M T394W T411T
AZ1361	L351L S400K F405A Y407V	T366L N390E K392F T394W T411N
AZ1362	L351L S400K F405A Y407V	T366L N390E K392F T394W T411L
AZ1363	L351L S400K F405A Y407V	T366L N390E K392F T394W T411S
AZ1364	L351L S400K F405A Y407V	T366L N390E K392F T394W T411T
AZ1365	L351L S400K F405S Y407I	T366L N390N K392L T394W T411N
AZ1366	L351L S400K F405S Y407I	T366L N390N K392L T394W T411S
AZ1367	L351L S400K F405S Y407I	T366L N390N K392L T394W T411T
AZ1368	L351L S400K F405S Y407I	T366L N390N K392M T394W T411N
AZ1369	L351L S400K F405S Y407I	T366L N390N K392M T394W T411S
AZ1370	L351L S400K F405S Y407I	T366L N390N K392M T394W T411T
AZ1371	L351L S400K F405S Y407I	T366L N390N K392F T394W T411N
AZ1372	L351L S400K F405S Y407I	T366L N390N K392F T394W T411S
AZ1373	L351L S400K F405S Y407I	T366L N390N K392F T394W T411T
AZ1374	L351L S400K F405S Y407I	T366L N390D K392L T394W T411N
AZ1375	L351L S400K F405S Y407I	T366L N390D K392L T394W T411S
AZ1376	L351L S400K F405S Y407I	T366L N390D K392L T394W T411T
AZ1377	L351L S400K F405S Y407I	T366L N390D K392M T394W T411N
AZ1378	L351L S400K F405S Y407I	T366L N390D K392M T394W T411S
AZ1379	L351L S400K F405S Y407I	T366L N390D K392M T394W T411T
AZ1380	L351L S400K F405S Y407I	T366L N390D K392F T394W T411N
AZ1381	L351L S400K F405S Y407I	T366L N390D K392F T394W T411S
AZ1382	L351L S400K F405S Y407I	T366L N390D K392F T394W T411T
AZ1383	L351L S400K F405S Y407I	T366L N390E K392L T394W T411N
AZ1384	L351L S400K F405S Y407I	T366L N390E K392L T394W T411L
AZ1385	L351L S400K F405S Y407I	T366L N390E K392L T394W T411S
AZ1386	L351L S400K F405S Y407I	T366L N390E K392L T394W T411T
AZ1387	L351L S400K F405S Y407I	T366L N390E K392M T394W T411N
AZ1388	L351L S400K F405S Y407I	T366L N390E K392M T394W T411L
AZ1389	L351L S400K F405S Y407I	T366L N390E K392M T394W T411S
AZ1390	L351L S400K F405S Y407I	T366L N390E K392M T394W T411T
AZ1391	L351L S400K F405S Y407I	T366L N390E K392F T394W T411N
AZ1392	L351L S400K F405S Y407I	T366L N390E K392F T394W T411L
AZ1393	L351L S400K F405S Y407I	T366L N390E K392F T394W T411S
AZ1394	L351L S400K F405S Y407I	T366L N390E K392F T394W T411T
AZ1395	L351L S400K F405S Y407V	T366L N390N K392L T394W T411N
AZ1396	L351L S400K F405S Y407V	T366L N390N K392L T394W T411S
AZ1397	L351L S400K F405S Y407V	T366L N390N K392L T394W T411T
AZ1398	L351L S400K F405S Y407V	T366L N390N K392M T394W T411N
AZ1399	L351L S400K F405S Y407V	T366L N390N K392M T394W T411S
AZ1400	L351L S400K F405S Y407V	T366L N390N K392M T394W T411T
AZ1401	L351L S400K F405S Y407V	T366L N390N K392F T394W T411N
AZ1402	L351L S400K F405S Y407V	T366L N390N K392F T394W T411S
AZ1403	L351L S400K F405S Y407V	T366L N390N K392F T394W T411T
AZ1404	L351L S400K F405S Y407V	T366L N390D K392L T394W T411N

AZ1405	L351L_S400K_F405S_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1406	L351L_S400K_F405S_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1407	L351L_S400K_F405S_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1408	L351L_S400K_F405S_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1409	L351L_S400K_F405S_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1410	L351L_S400K_F405S_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1411	L351L_S400K_F405S_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1412	L351L_S400K_F405S_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1413	L351L_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1414	L351L_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1415	L351L_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1416	L351L_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1417	L351L_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1418	L351L_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1419	L351L_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1420	L351L_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1421	L351L_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1422	L351L_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1423	L351L_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1424	L351L_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1425	L351L_S400K_F405T_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1426	L351L_S400K_F405T_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1427	L351L_S400K_F405T_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1428	L351L_S400K_F405T_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1429	L351L_S400K_F405T_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1430	L351L_S400K_F405T_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1431	L351L_S400K_F405T_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1432	L351L_S400K_F405T_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1433	L351L_S400K_F405T_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1434	L351L_S400K_F405T_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1435	L351L_S400K_F405T_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1436	L351L_S400K_F405T_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1437	L351L_S400K_F405T_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1438	L351L_S400K_F405T_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1439	L351L_S400K_F405T_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1440	L351L_S400K_F405T_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1441	L351L_S400K_F405T_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1442	L351L_S400K_F405T_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1443	L351L_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1444	L351L_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1445	L351L_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1446	L351L_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1447	L351L_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1448	L351L_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1449	L351L_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1450	L351L_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1451	L351L_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1452	L351L_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1453	L351L_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1454	L351L_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1455	L351L_S400K_F405T_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1456	L351L_S400K_F405T_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1457	L351L_S400K_F405T_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1458	L351L_S400K_F405T_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1459	L351L_S400K_F405T_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1460	L351L_S400K_F405T_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1461	L351L_S400K_F405T_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1462	L351L_S400K_F405T_Y407V	T366L_N390N_K392F_T394W_T411S

AZ1463	L351L_S400K_F405T_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1464	L351L_S400K_F405T_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1465	L351L_S400K_F405T_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1466	L351L_S400K_F405T_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1467	L351L_S400K_F405T_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1468	L351L_S400K_F405T_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1469	L351L_S400K_F405T_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1470	L351L_S400K_F405T_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1471	L351L_S400K_F405T_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1472	L351L_S400K_F405T_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1473	L351L_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1474	L351L_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1475	L351L_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1476	L351L_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1477	L351L_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1478	L351L_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1479	L351L_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1480	L351L_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1481	L351L_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1482	L351L_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1483	L351L_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1484	L351L_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1485	L351L_S400K_F405V_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1486	L351L_S400K_F405V_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1487	L351L_S400K_F405V_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1488	L351L_S400K_F405V_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1489	L351L_S400K_F405V_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1490	L351L_S400K_F405V_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1491	L351L_S400K_F405V_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1492	L351L_S400K_F405V_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1493	L351L_S400K_F405V_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1494	L351L_S400K_F405V_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1495	L351L_S400K_F405V_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1496	L351L_S400K_F405V_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1497	L351L_S400K_F405V_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1498	L351L_S400K_F405V_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1499	L351L_S400K_F405V_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1500	L351L_S400K_F405V_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1501	L351L_S400K_F405V_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1502	L351L_S400K_F405V_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1503	L351L_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1504	L351L_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1505	L351L_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1506	L351L_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1507	L351L_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1508	L351L_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1509	L351L_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1510	L351L_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1511	L351L_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1512	L351L_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1513	L351L_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1514	L351L_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1515	L351L_S400K_F405V_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1516	L351L_S400K_F405V_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1517	L351L_S400K_F405V_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1518	L351L_S400K_F405V_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1519	L351L_S400K_F405V_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1520	L351L_S400K_F405V_Y407V	T366L_N390N_K392M_T394W_T411T

AZ1521	L351L_S400K_F405V_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1522	L351L_S400K_F405V_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1523	L351L_S400K_F405V_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1524	L351L_S400K_F405V_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1525	L351L_S400K_F405V_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1526	L351L_S400K_F405V_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1527	L351L_S400K_F405V_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1528	L351L_S400K_F405V_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1529	L351L_S400K_F405V_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1530	L351L_S400K_F405V_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1531	L351L_S400K_F405V_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1532	L351L_S400K_F405V_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1533	L351L_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1534	L351L_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1535	L351L_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1536	L351L_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1537	L351L_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1538	L351L_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1539	L351L_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1540	L351L_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1541	L351L_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1542	L351L_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1543	L351L_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1544	L351L_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1545	L351Y_D399R_F405A_Y407I	T366L_K392L_T394W_K409L_T411E
AZ1546	L351Y_D399R_F405A_Y407I	T366L_K392L_T394W_K409M_T411E
AZ1547	L351Y_D399R_F405A_Y407I	T366L_K392M_T394W_K409L_T411E
AZ1548	L351Y_D399R_F405A_Y407I	T366L_K392M_T394W_K409M_T411E
AZ1549	L351Y_D399R_F405A_Y407I	T366L_K392F_T394W_K409L_T411E
AZ1550	L351Y_D399R_F405A_Y407I	T366L_K392F_T394W_K409M_T411E
AZ1551	L351Y_D399R_F405A_Y407V	T366L_K392L_T394W_K409L_T411E
AZ1552	L351Y_D399R_F405A_Y407V	T366L_K392L_T394W_K409M_T411E
AZ1553	L351Y_D399R_F405A_Y407V	T366L_K392M_T394W_K409L_T411E
AZ1554	L351Y_D399R_F405A_Y407V	T366L_K392M_T394W_K409M_T411E
AZ1555	L351Y_D399R_F405A_Y407V	T366L_K392F_T394W_K409L_T411E
AZ1556	L351Y_D399R_F405A_Y407V	T366L_K392F_T394W_K409M_T411E
AZ1557	L351Y_D399R_F405M_Y407I	T366L_K392L_T394W_K409V_T411E
AZ1558	L351Y_D399R_F405M_Y407I	T366L_K392M_T394W_K409V_T411E
AZ1559	L351Y_D399R_F405M_Y407I	T366L_K392F_T394W_K409V_T411E
AZ1560	L351Y_D399R_F405M_Y407V	T366L_K392L_T394W_K409V_T411E
AZ1561	L351Y_D399R_F405M_Y407V	T366L_K392M_T394W_K409V_T411E
AZ1562	L351Y_D399R_F405M_Y407V	T366L_K392F_T394W_K409V_T411E
AZ1563	L351Y_D399R_F405S_Y407I	T366L_K392L_T394W_K409L_T411E
AZ1564	L351Y_D399R_F405S_Y407I	T366L_K392L_T394W_K409M_T411E
AZ1565	L351Y_D399R_F405S_Y407I	T366L_K392M_T394W_K409L_T411E
AZ1566	L351Y_D399R_F405S_Y407I	T366L_K392M_T394W_K409M_T411E
AZ1567	L351Y_D399R_F405S_Y407I	T366L_K392F_T394W_K409L_T411E
AZ1568	L351Y_D399R_F405S_Y407I	T366L_K392F_T394W_K409M_T411E
AZ1569	L351Y_D399R_F405S_Y407V	T366L_K392L_T394W_K409L_T411E
AZ1570	L351Y_D399R_F405S_Y407V	T366L_K392L_T394W_K409M_T411E
AZ1571	L351Y_D399R_F405S_Y407V	T366L_K392M_T394W_K409L_T411E
AZ1572	L351Y_D399R_F405S_Y407V	T366L_K392M_T394W_K409M_T411E
AZ1573	L351Y_D399R_F405S_Y407V	T366L_K392F_T394W_K409L_T411E
AZ1574	L351Y_D399R_F405S_Y407V	T366L_K392F_T394W_K409M_T411E
AZ1575	L351Y_D399R_F405V_Y407I	T366L_K392L_T394W_K409L_T411E
AZ1576	L351Y_D399R_F405V_Y407I	T366L_K392L_T394W_K409M_T411E
AZ1577	L351Y_D399R_F405V_Y407I	T366L_K392M_T394W_K409L_T411E
AZ1578	L351Y_D399R_F405V_Y407I	T366L_K392M_T394W_K409M_T411E

AZ1579	L351Y_D399R_F405V_Y407I	T366L_K392F_T394W_K409L_T411E
AZ1580	L351Y_D399R_F405V_Y407I	T366L_K392F_T394W_K409M_T411E
AZ1581	L351Y_D399R_F405V_Y407V	T366L_K392L_T394W_K409L_T411E
AZ1582	L351Y_D399R_F405V_Y407V	T366L_K392L_T394W_K409M_T411E
AZ1583	L351Y_D399R_F405V_Y407V	T366L_K392M_T394W_K409L_T411E
AZ1584	L351Y_D399R_F405V_Y407V	T366L_K392M_T394W_K409M_T411E
AZ1585	L351Y_D399R_F405V_Y407V	T366L_K392F_T394W_K409L_T411E
AZ1586	L351Y_D399R_F405V_Y407V	T366L_K392F_T394W_K409M_T411E
AZ1587	L351Y_S400R_F405A_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1588	L351Y_S400R_F405A_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1589	L351Y_S400R_F405A_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1590	L351Y_S400R_F405A_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1591	L351Y_S400R_F405A_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1592	L351Y_S400R_F405A_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1593	L351Y_S400R_F405A_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1594	L351Y_S400R_F405A_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1595	L351Y_S400R_F405A_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1596	L351Y_S400R_F405A_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1597	L351Y_S400R_F405A_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1598	L351Y_S400R_F405A_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1599	L351Y_S400R_F405A_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1600	L351Y_S400R_F405A_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1601	L351Y_S400R_F405A_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1602	L351Y_S400R_F405A_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1603	L351Y_S400R_F405A_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1604	L351Y_S400R_F405A_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1605	L351Y_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1606	L351Y_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1607	L351Y_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1608	L351Y_S400R_F405A_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1609	L351Y_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1610	L351Y_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1611	L351Y_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1612	L351Y_S400R_F405A_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1613	L351Y_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1614	L351Y_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1615	L351Y_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1616	L351Y_S400R_F405A_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1617	L351Y_S400R_F405A_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1618	L351Y_S400R_F405A_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1619	L351Y_S400R_F405A_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1620	L351Y_S400R_F405A_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1621	L351Y_S400R_F405A_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1622	L351Y_S400R_F405A_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1623	L351Y_S400R_F405A_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1624	L351Y_S400R_F405A_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1625	L351Y_S400R_F405A_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1626	L351Y_S400R_F405A_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1627	L351Y_S400R_F405A_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1628	L351Y_S400R_F405A_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1629	L351Y_S400R_F405A_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1630	L351Y_S400R_F405A_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1631	L351Y_S400R_F405A_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1632	L351Y_S400R_F405A_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1633	L351Y_S400R_F405A_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1634	L351Y_S400R_F405A_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1635	L351Y_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1636	L351Y_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411L

AZ1637	L351Y_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1638	L351Y_S400R_F405A_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1639	L351Y_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1640	L351Y_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1641	L351Y_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1642	L351Y_S400R_F405A_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1643	L351Y_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1644	L351Y_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1645	L351Y_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1646	L351Y_S400R_F405A_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1647	L351Y_S400R_F405S_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1648	L351Y_S400R_F405S_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1649	L351Y_S400R_F405S_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1650	L351Y_S400R_F405S_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1651	L351Y_S400R_F405S_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1652	L351Y_S400R_F405S_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1653	L351Y_S400R_F405S_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1654	L351Y_S400R_F405S_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1655	L351Y_S400R_F405S_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1656	L351Y_S400R_F405S_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1657	L351Y_S400R_F405S_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1658	L351Y_S400R_F405S_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1659	L351Y_S400R_F405S_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1660	L351Y_S400R_F405S_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1661	L351Y_S400R_F405S_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1662	L351Y_S400R_F405S_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1663	L351Y_S400R_F405S_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1664	L351Y_S400R_F405S_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1665	L351Y_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1666	L351Y_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1667	L351Y_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1668	L351Y_S400R_F405S_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1669	L351Y_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1670	L351Y_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1671	L351Y_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1672	L351Y_S400R_F405S_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1673	L351Y_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1674	L351Y_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1675	L351Y_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1676	L351Y_S400R_F405S_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1677	L351Y_S400R_F405S_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1678	L351Y_S400R_F405S_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1679	L351Y_S400R_F405S_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1680	L351Y_S400R_F405S_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1681	L351Y_S400R_F405S_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1682	L351Y_S400R_F405S_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1683	L351Y_S400R_F405S_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1684	L351Y_S400R_F405S_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1685	L351Y_S400R_F405S_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1686	L351Y_S400R_F405S_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1687	L351Y_S400R_F405S_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1688	L351Y_S400R_F405S_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1689	L351Y_S400R_F405S_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1690	L351Y_S400R_F405S_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1691	L351Y_S400R_F405S_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1692	L351Y_S400R_F405S_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1693	L351Y_S400R_F405S_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1694	L351Y_S400R_F405S_Y407V	T366L_N390D_K392F_T394W_T411T

AZ1695	L351Y_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1696	L351Y_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1697	L351Y_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1698	L351Y_S400R_F405S_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1699	L351Y_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1700	L351Y_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1701	L351Y_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1702	L351Y_S400R_F405S_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1703	L351Y_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1704	L351Y_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1705	L351Y_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1706	L351Y_S400R_F405S_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1707	L351Y_S400R_F405T_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1708	L351Y_S400R_F405T_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1709	L351Y_S400R_F405T_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1710	L351Y_S400R_F405T_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1711	L351Y_S400R_F405T_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1712	L351Y_S400R_F405T_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1713	L351Y_S400R_F405T_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1714	L351Y_S400R_F405T_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1715	L351Y_S400R_F405T_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1716	L351Y_S400R_F405T_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1717	L351Y_S400R_F405T_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1718	L351Y_S400R_F405T_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1719	L351Y_S400R_F405T_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1720	L351Y_S400R_F405T_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1721	L351Y_S400R_F405T_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1722	L351Y_S400R_F405T_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1723	L351Y_S400R_F405T_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1724	L351Y_S400R_F405T_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1725	L351Y_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1726	L351Y_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1727	L351Y_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1728	L351Y_S400R_F405T_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1729	L351Y_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1730	L351Y_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1731	L351Y_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1732	L351Y_S400R_F405T_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1733	L351Y_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1734	L351Y_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1735	L351Y_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1736	L351Y_S400R_F405T_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1737	L351Y_S400R_F405T_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1738	L351Y_S400R_F405T_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1739	L351Y_S400R_F405T_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1740	L351Y_S400R_F405T_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1741	L351Y_S400R_F405T_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1742	L351Y_S400R_F405T_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1743	L351Y_S400R_F405T_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1744	L351Y_S400R_F405T_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1745	L351Y_S400R_F405T_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1746	L351Y_S400R_F405T_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1747	L351Y_S400R_F405T_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1748	L351Y_S400R_F405T_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1749	L351Y_S400R_F405T_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1750	L351Y_S400R_F405T_Y407V	T366L_N390D_K392M_T394W_T411S
AZ1751	L351Y_S400R_F405T_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1752	L351Y_S400R_F405T_Y407V	T366L_N390D_K392F_T394W_T411N

AZ1753	L351Y_S400R_F405T_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1754	L351Y_S400R_F405T_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1755	L351Y_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1756	L351Y_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1757	L351Y_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1758	L351Y_S400R_F405T_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1759	L351Y_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1760	L351Y_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1761	L351Y_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1762	L351Y_S400R_F405T_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1763	L351Y_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1764	L351Y_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1765	L351Y_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1766	L351Y_S400R_F405T_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1767	L351Y_S400R_F405V_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1768	L351Y_S400R_F405V_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1769	L351Y_S400R_F405V_Y407I	T366L_N390N_K392L_T394W_T411T
AZ1770	L351Y_S400R_F405V_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1771	L351Y_S400R_F405V_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1772	L351Y_S400R_F405V_Y407I	T366L_N390N_K392M_T394W_T411T
AZ1773	L351Y_S400R_F405V_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1774	L351Y_S400R_F405V_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1775	L351Y_S400R_F405V_Y407I	T366L_N390N_K392F_T394W_T411T
AZ1776	L351Y_S400R_F405V_Y407I	T366L_N390D_K392L_T394W_T411N
AZ1777	L351Y_S400R_F405V_Y407I	T366L_N390D_K392L_T394W_T411S
AZ1778	L351Y_S400R_F405V_Y407I	T366L_N390D_K392L_T394W_T411T
AZ1779	L351Y_S400R_F405V_Y407I	T366L_N390D_K392M_T394W_T411N
AZ1780	L351Y_S400R_F405V_Y407I	T366L_N390D_K392M_T394W_T411S
AZ1781	L351Y_S400R_F405V_Y407I	T366L_N390D_K392M_T394W_T411T
AZ1782	L351Y_S400R_F405V_Y407I	T366L_N390D_K392F_T394W_T411N
AZ1783	L351Y_S400R_F405V_Y407I	T366L_N390D_K392F_T394W_T411S
AZ1784	L351Y_S400R_F405V_Y407I	T366L_N390D_K392F_T394W_T411T
AZ1785	L351Y_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411N
AZ1786	L351Y_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411L
AZ1787	L351Y_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411S
AZ1788	L351Y_S400R_F405V_Y407I	T366L_N390E_K392L_T394W_T411T
AZ1789	L351Y_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411N
AZ1790	L351Y_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411L
AZ1791	L351Y_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411S
AZ1792	L351Y_S400R_F405V_Y407I	T366L_N390E_K392M_T394W_T411T
AZ1793	L351Y_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411N
AZ1794	L351Y_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411L
AZ1795	L351Y_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411S
AZ1796	L351Y_S400R_F405V_Y407I	T366L_N390E_K392F_T394W_T411T
AZ1797	L351Y_S400R_F405V_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1798	L351Y_S400R_F405V_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1799	L351Y_S400R_F405V_Y407V	T366L_N390N_K392L_T394W_T411T
AZ1800	L351Y_S400R_F405V_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1801	L351Y_S400R_F405V_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1802	L351Y_S400R_F405V_Y407V	T366L_N390N_K392M_T394W_T411T
AZ1803	L351Y_S400R_F405V_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1804	L351Y_S400R_F405V_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1805	L351Y_S400R_F405V_Y407V	T366L_N390N_K392F_T394W_T411T
AZ1806	L351Y_S400R_F405V_Y407V	T366L_N390D_K392L_T394W_T411N
AZ1807	L351Y_S400R_F405V_Y407V	T366L_N390D_K392L_T394W_T411S
AZ1808	L351Y_S400R_F405V_Y407V	T366L_N390D_K392L_T394W_T411T
AZ1809	L351Y_S400R_F405V_Y407V	T366L_N390D_K392M_T394W_T411N
AZ1810	L351Y_S400R_F405V_Y407V	T366L_N390D_K392M_T394W_T411S

AZ1811	L351Y_S400R_F405V_Y407V	T366L_N390D_K392M_T394W_T411T
AZ1812	L351Y_S400R_F405V_Y407V	T366L_N390D_K392F_T394W_T411N
AZ1813	L351Y_S400R_F405V_Y407V	T366L_N390D_K392F_T394W_T411S
AZ1814	L351Y_S400R_F405V_Y407V	T366L_N390D_K392F_T394W_T411T
AZ1815	L351Y_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411N
AZ1816	L351Y_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411L
AZ1817	L351Y_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411S
AZ1818	L351Y_S400R_F405V_Y407V	T366L_N390E_K392L_T394W_T411T
AZ1819	L351Y_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411N
AZ1820	L351Y_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411L
AZ1821	L351Y_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411S
AZ1822	L351Y_S400R_F405V_Y407V	T366L_N390E_K392M_T394W_T411T
AZ1823	L351Y_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411N
AZ1824	L351Y_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411L
AZ1825	L351Y_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411S
AZ1826	L351Y_S400R_F405V_Y407V	T366L_N390E_K392F_T394W_T411T
AZ1827	L351Y_S400E_F405A_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1828	L351Y_S400E_F405A_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1829	L351Y_S400E_F405A_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1830	L351Y_S400E_F405A_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1831	L351Y_S400E_F405A_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1832	L351Y_S400E_F405A_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1833	L351Y_S400E_F405A_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1834	L351Y_S400E_F405A_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1835	L351Y_S400E_F405A_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1836	L351Y_S400E_F405A_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1837	L351Y_S400E_F405A_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1838	L351Y_S400E_F405A_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1839	L351Y_S400E_F405A_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1840	L351Y_S400E_F405A_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1841	L351Y_S400E_F405A_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1842	L351Y_S400E_F405A_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1843	L351Y_S400E_F405A_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1844	L351Y_S400E_F405A_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1845	L351Y_S400E_F405A_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1846	L351Y_S400E_F405A_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1847	L351Y_S400E_F405A_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1848	L351Y_S400E_F405A_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1849	L351Y_S400E_F405A_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1850	L351Y_S400E_F405A_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1851	L351Y_S400E_F405A_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1852	L351Y_S400E_F405A_Y407I	T366L_N390K_K392F_T394W_T411L
AZ1853	L351Y_S400E_F405A_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1854	L351Y_S400E_F405A_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1855	L351Y_S400E_F405A_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1856	L351Y_S400E_F405A_Y407V	T366L_N390R_K392L_T394W_T411S
AZ1857	L351Y_S400E_F405A_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1858	L351Y_S400E_F405A_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1859	L351Y_S400E_F405A_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1860	L351Y_S400E_F405A_Y407V	T366L_N390R_K392F_T394W_T411N
AZ1861	L351Y_S400E_F405A_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1862	L351Y_S400E_F405A_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1863	L351Y_S400E_F405A_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1864	L351Y_S400E_F405A_Y407V	T366L_N390N_K392L_T394W_T411L
AZ1865	L351Y_S400E_F405A_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1866	L351Y_S400E_F405A_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1867	L351Y_S400E_F405A_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1868	L351Y_S400E_F405A_Y407V	T366L_N390N_K392M_T394W_T411S

AZ1869	L351Y_S400E_F405A_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1870	L351Y_S400E_F405A_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1871	L351Y_S400E_F405A_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1872	L351Y_S400E_F405A_Y407V	T366L_N390K_K392L_T394W_T411N
AZ1873	L351Y_S400E_F405A_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1874	L351Y_S400E_F405A_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1875	L351Y_S400E_F405A_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1876	L351Y_S400E_F405A_Y407V	T366L_N390K_K392M_T394W_T411L
AZ1877	L351Y_S400E_F405A_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1878	L351Y_S400E_F405A_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1879	L351Y_S400E_F405A_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1880	L351Y_S400E_F405A_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1881	L351Y_S400E_F405S_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1882	L351Y_S400E_F405S_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1883	L351Y_S400E_F405S_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1884	L351Y_S400E_F405S_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1885	L351Y_S400E_F405S_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1886	L351Y_S400E_F405S_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1887	L351Y_S400E_F405S_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1888	L351Y_S400E_F405S_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1889	L351Y_S400E_F405S_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1890	L351Y_S400E_F405S_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1891	L351Y_S400E_F405S_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1892	L351Y_S400E_F405S_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1893	L351Y_S400E_F405S_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1894	L351Y_S400E_F405S_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1895	L351Y_S400E_F405S_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1896	L351Y_S400E_F405S_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1897	L351Y_S400E_F405S_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1898	L351Y_S400E_F405S_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1899	L351Y_S400E_F405S_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1900	L351Y_S400E_F405S_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1901	L351Y_S400E_F405S_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1902	L351Y_S400E_F405S_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1903	L351Y_S400E_F405S_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1904	L351Y_S400E_F405S_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1905	L351Y_S400E_F405S_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1906	L351Y_S400E_F405S_Y407I	T366L_N390K_K392F_T394W_T411L
AZ1907	L351Y_S400E_F405S_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1908	L351Y_S400E_F405S_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1909	L351Y_S400E_F405S_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1910	L351Y_S400E_F405S_Y407V	T366L_N390R_K392L_T394W_T411S
AZ1911	L351Y_S400E_F405S_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1912	L351Y_S400E_F405S_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1913	L351Y_S400E_F405S_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1914	L351Y_S400E_F405S_Y407V	T366L_N390R_K392F_T394W_T411N
AZ1915	L351Y_S400E_F405S_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1916	L351Y_S400E_F405S_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1917	L351Y_S400E_F405S_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1918	L351Y_S400E_F405S_Y407V	T366L_N390N_K392L_T394W_T411L
AZ1919	L351Y_S400E_F405S_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1920	L351Y_S400E_F405S_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1921	L351Y_S400E_F405S_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1922	L351Y_S400E_F405S_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1923	L351Y_S400E_F405S_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1924	L351Y_S400E_F405S_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1925	L351Y_S400E_F405S_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1926	L351Y_S400E_F405S_Y407V	T366L_N390K_K392L_T394W_T411N

AZ1927	L351Y_S400E_F405S_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1928	L351Y_S400E_F405S_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1929	L351Y_S400E_F405S_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1930	L351Y_S400E_F405S_Y407V	T366L_N390K_K392M_T394W_T411L
AZ1931	L351Y_S400E_F405S_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1932	L351Y_S400E_F405S_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1933	L351Y_S400E_F405S_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1934	L351Y_S400E_F405S_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1935	L351Y_S400E_F405T_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1936	L351Y_S400E_F405T_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1937	L351Y_S400E_F405T_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1938	L351Y_S400E_F405T_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1939	L351Y_S400E_F405T_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1940	L351Y_S400E_F405T_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1941	L351Y_S400E_F405T_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1942	L351Y_S400E_F405T_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1943	L351Y_S400E_F405T_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1944	L351Y_S400E_F405T_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1945	L351Y_S400E_F405T_Y407I	T366L_N390N_K392L_T394W_T411L
AZ1946	L351Y_S400E_F405T_Y407I	T366L_N390N_K392L_T394W_T411S
AZ1947	L351Y_S400E_F405T_Y407I	T366L_N390N_K392M_T394W_T411N
AZ1948	L351Y_S400E_F405T_Y407I	T366L_N390N_K392M_T394W_T411L
AZ1949	L351Y_S400E_F405T_Y407I	T366L_N390N_K392M_T394W_T411S
AZ1950	L351Y_S400E_F405T_Y407I	T366L_N390N_K392F_T394W_T411N
AZ1951	L351Y_S400E_F405T_Y407I	T366L_N390N_K392F_T394W_T411L
AZ1952	L351Y_S400E_F405T_Y407I	T366L_N390N_K392F_T394W_T411S
AZ1953	L351Y_S400E_F405T_Y407I	T366L_N390K_K392L_T394W_T411N
AZ1954	L351Y_S400E_F405T_Y407I	T366L_N390K_K392L_T394W_T411L
AZ1955	L351Y_S400E_F405T_Y407I	T366L_N390K_K392L_T394W_T411S
AZ1956	L351Y_S400E_F405T_Y407I	T366L_N390K_K392M_T394W_T411N
AZ1957	L351Y_S400E_F405T_Y407I	T366L_N390K_K392M_T394W_T411L
AZ1958	L351Y_S400E_F405T_Y407I	T366L_N390K_K392M_T394W_T411S
AZ1959	L351Y_S400E_F405T_Y407I	T366L_N390K_K392F_T394W_T411N
AZ1960	L351Y_S400E_F405T_Y407I	T366L_N390K_K392F_T394W_T411L
AZ1961	L351Y_S400E_F405T_Y407I	T366L_N390K_K392F_T394W_T411S
AZ1962	L351Y_S400E_F405T_Y407V	T366L_N390R_K392L_T394W_T411N
AZ1963	L351Y_S400E_F405T_Y407V	T366L_N390R_K392L_T394W_T411L
AZ1964	L351Y_S400E_F405T_Y407V	T366L_N390R_K392L_T394W_T411S
AZ1965	L351Y_S400E_F405T_Y407V	T366L_N390R_K392M_T394W_T411N
AZ1966	L351Y_S400E_F405T_Y407V	T366L_N390R_K392M_T394W_T411L
AZ1967	L351Y_S400E_F405T_Y407V	T366L_N390R_K392M_T394W_T411S
AZ1968	L351Y_S400E_F405T_Y407V	T366L_N390R_K392F_T394W_T411N
AZ1969	L351Y_S400E_F405T_Y407V	T366L_N390R_K392F_T394W_T411L
AZ1970	L351Y_S400E_F405T_Y407V	T366L_N390R_K392F_T394W_T411S
AZ1971	L351Y_S400E_F405T_Y407V	T366L_N390N_K392L_T394W_T411N
AZ1972	L351Y_S400E_F405T_Y407V	T366L_N390N_K392L_T394W_T411L
AZ1973	L351Y_S400E_F405T_Y407V	T366L_N390N_K392L_T394W_T411S
AZ1974	L351Y_S400E_F405T_Y407V	T366L_N390N_K392M_T394W_T411N
AZ1975	L351Y_S400E_F405T_Y407V	T366L_N390N_K392M_T394W_T411L
AZ1976	L351Y_S400E_F405T_Y407V	T366L_N390N_K392M_T394W_T411S
AZ1977	L351Y_S400E_F405T_Y407V	T366L_N390N_K392F_T394W_T411N
AZ1978	L351Y_S400E_F405T_Y407V	T366L_N390N_K392F_T394W_T411L
AZ1979	L351Y_S400E_F405T_Y407V	T366L_N390N_K392F_T394W_T411S
AZ1980	L351Y_S400E_F405T_Y407V	T366L_N390K_K392L_T394W_T411N
AZ1981	L351Y_S400E_F405T_Y407V	T366L_N390K_K392L_T394W_T411L
AZ1982	L351Y_S400E_F405T_Y407V	T366L_N390K_K392L_T394W_T411S
AZ1983	L351Y_S400E_F405T_Y407V	T366L_N390K_K392M_T394W_T411N
AZ1984	L351Y_S400E_F405T_Y407V	T366L_N390K_K392M_T394W_T411L

AZ1985	L351Y_S400E_F405T_Y407V	T366L_N390K_K392M_T394W_T411S
AZ1986	L351Y_S400E_F405T_Y407V	T366L_N390K_K392F_T394W_T411N
AZ1987	L351Y_S400E_F405T_Y407V	T366L_N390K_K392F_T394W_T411L
AZ1988	L351Y_S400E_F405T_Y407V	T366L_N390K_K392F_T394W_T411S
AZ1989	L351Y_S400E_F405V_Y407I	T366L_N390R_K392L_T394W_T411N
AZ1990	L351Y_S400E_F405V_Y407I	T366L_N390R_K392L_T394W_T411L
AZ1991	L351Y_S400E_F405V_Y407I	T366L_N390R_K392L_T394W_T411S
AZ1992	L351Y_S400E_F405V_Y407I	T366L_N390R_K392M_T394W_T411N
AZ1993	L351Y_S400E_F405V_Y407I	T366L_N390R_K392M_T394W_T411L
AZ1994	L351Y_S400E_F405V_Y407I	T366L_N390R_K392M_T394W_T411S
AZ1995	L351Y_S400E_F405V_Y407I	T366L_N390R_K392F_T394W_T411N
AZ1996	L351Y_S400E_F405V_Y407I	T366L_N390R_K392F_T394W_T411L
AZ1997	L351Y_S400E_F405V_Y407I	T366L_N390R_K392F_T394W_T411S
AZ1998	L351Y_S400E_F405V_Y407I	T366L_N390N_K392L_T394W_T411N
AZ1999	L351Y_S400E_F405V_Y407I	T366L_N390N_K392L_T394W_T411L
AZ2000	L351Y_S400E_F405V_Y407I	T366L_N390N_K392L_T394W_T411S
AZ2001	L351Y_S400E_F405V_Y407I	T366L_N390N_K392M_T394W_T411N
AZ2002	L351Y_S400E_F405V_Y407I	T366L_N390N_K392M_T394W_T411L
AZ2003	L351Y_S400E_F405V_Y407I	T366L_N390N_K392M_T394W_T411S
AZ2004	L351Y_S400E_F405V_Y407I	T366L_N390N_K392F_T394W_T411N
AZ2005	L351Y_S400E_F405V_Y407I	T366L_N390N_K392F_T394W_T411L
AZ2006	L351Y_S400E_F405V_Y407I	T366L_N390N_K392F_T394W_T411S
AZ2007	L351Y_S400E_F405V_Y407I	T366L_N390K_K392L_T394W_T411N
AZ2008	L351Y_S400E_F405V_Y407I	T366L_N390K_K392L_T394W_T411L
AZ2009	L351Y_S400E_F405V_Y407I	T366L_N390K_K392L_T394W_T411S
AZ2010	L351Y_S400E_F405V_Y407I	T366L_N390K_K392M_T394W_T411N
AZ2011	L351Y_S400E_F405V_Y407I	T366L_N390K_K392M_T394W_T411L
AZ2012	L351Y_S400E_F405V_Y407I	T366L_N390K_K392M_T394W_T411S
AZ2013	L351Y_S400E_F405V_Y407I	T366L_N390K_K392F_T394W_T411N
AZ2014	L351Y_S400E_F405V_Y407I	T366L_N390K_K392F_T394W_T411L
AZ2015	L351Y_S400E_F405V_Y407I	T366L_N390K_K392F_T394W_T411S
AZ2016	L351Y_S400E_F405V_Y407V	T366L_N390R_K392L_T394W_T411N
AZ2017	L351Y_S400E_F405V_Y407V	T366L_N390R_K392L_T394W_T411L
AZ2018	L351Y_S400E_F405V_Y407V	T366L_N390R_K392L_T394W_T411S
AZ2019	L351Y_S400E_F405V_Y407V	T366L_N390R_K392M_T394W_T411N
AZ2020	L351Y_S400E_F405V_Y407V	T366L_N390R_K392M_T394W_T411L
AZ2021	L351Y_S400E_F405V_Y407V	T366L_N390R_K392M_T394W_T411S
AZ2022	L351Y_S400E_F405V_Y407V	T366L_N390R_K392F_T394W_T411N
AZ2023	L351Y_S400E_F405V_Y407V	T366L_N390R_K392F_T394W_T411L
AZ2024	L351Y_S400E_F405V_Y407V	T366L_N390R_K392F_T394W_T411S
AZ2025	L351Y_S400E_F405V_Y407V	T366L_N390N_K392L_T394W_T411N
AZ2026	L351Y_S400E_F405V_Y407V	T366L_N390N_K392L_T394W_T411L
AZ2027	L351Y_S400E_F405V_Y407V	T366L_N390N_K392L_T394W_T411S
AZ2028	L351Y_S400E_F405V_Y407V	T366L_N390N_K392M_T394W_T411N
AZ2029	L351Y_S400E_F405V_Y407V	T366L_N390N_K392M_T394W_T411L
AZ2030	L351Y_S400E_F405V_Y407V	T366L_N390N_K392M_T394W_T411S
AZ2031	L351Y_S400E_F405V_Y407V	T366L_N390N_K392F_T394W_T411N
AZ2032	L351Y_S400E_F405V_Y407V	T366L_N390N_K392F_T394W_T411L
AZ2033	L351Y_S400E_F405V_Y407V	T366L_N390N_K392F_T394W_T411S
AZ2034	L351Y_S400E_F405V_Y407V	T366L_N390K_K392L_T394W_T411N
AZ2035	L351Y_S400E_F405V_Y407V	T366L_N390K_K392L_T394W_T411L
AZ2036	L351Y_S400E_F405V_Y407V	T366L_N390K_K392L_T394W_T411S
AZ2037	L351Y_S400E_F405V_Y407V	T366L_N390K_K392M_T394W_T411N
AZ2038	L351Y_S400E_F405V_Y407V	T366L_N390K_K392M_T394W_T411L
AZ2039	L351Y_S400E_F405V_Y407V	T366L_N390K_K392M_T394W_T411S
AZ2040	L351Y_S400E_F405V_Y407V	T366L_N390K_K392F_T394W_T411N
AZ2041	L351Y_S400E_F405V_Y407V	T366L_N390K_K392F_T394W_T411L
AZ2042	L351Y_S400E_F405V_Y407V	T366L_N390K_K392F_T394W_T411S

AZ2043	L351Y_S400K_F405A_Y407I	T366L_N390N_K392L_T394W_T411N
AZ2044	L351Y_S400K_F405A_Y407I	T366L_N390N_K392L_T394W_T411S
AZ2045	L351Y_S400K_F405A_Y407I	T366L_N390N_K392L_T394W_T411T
AZ2046	L351Y_S400K_F405A_Y407I	T366L_N390N_K392M_T394W_T411N
AZ2047	L351Y_S400K_F405A_Y407I	T366L_N390N_K392M_T394W_T411S
AZ2048	L351Y_S400K_F405A_Y407I	T366L_N390N_K392M_T394W_T411T
AZ2049	L351Y_S400K_F405A_Y407I	T366L_N390N_K392F_T394W_T411N
AZ2050	L351Y_S400K_F405A_Y407I	T366L_N390N_K392F_T394W_T411S
AZ2051	L351Y_S400K_F405A_Y407I	T366L_N390N_K392F_T394W_T411T
AZ2052	L351Y_S400K_F405A_Y407I	T366L_N390D_K392L_T394W_T411N
AZ2053	L351Y_S400K_F405A_Y407I	T366L_N390D_K392L_T394W_T411S
AZ2054	L351Y_S400K_F405A_Y407I	T366L_N390D_K392L_T394W_T411T
AZ2055	L351Y_S400K_F405A_Y407I	T366L_N390D_K392M_T394W_T411N
AZ2056	L351Y_S400K_F405A_Y407I	T366L_N390D_K392M_T394W_T411S
AZ2057	L351Y_S400K_F405A_Y407I	T366L_N390D_K392M_T394W_T411T
AZ2058	L351Y_S400K_F405A_Y407I	T366L_N390D_K392F_T394W_T411N
AZ2059	L351Y_S400K_F405A_Y407I	T366L_N390D_K392F_T394W_T411S
AZ2060	L351Y_S400K_F405A_Y407I	T366L_N390D_K392F_T394W_T411T
AZ2061	L351Y_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411N
AZ2062	L351Y_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411L
AZ2063	L351Y_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411S
AZ2064	L351Y_S400K_F405A_Y407I	T366L_N390E_K392L_T394W_T411T
AZ2065	L351Y_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411N
AZ2066	L351Y_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411L
AZ2067	L351Y_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411S
AZ2068	L351Y_S400K_F405A_Y407I	T366L_N390E_K392M_T394W_T411T
AZ2069	L351Y_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411N
AZ2070	L351Y_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411L
AZ2071	L351Y_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411S
AZ2072	L351Y_S400K_F405A_Y407I	T366L_N390E_K392F_T394W_T411T
AZ2073	L351Y_S400K_F405A_Y407V	T366L_N390N_K392L_T394W_T411N
AZ2074	L351Y_S400K_F405A_Y407V	T366L_N390N_K392L_T394W_T411S
AZ2075	L351Y_S400K_F405A_Y407V	T366L_N390N_K392L_T394W_T411T
AZ2076	L351Y_S400K_F405A_Y407V	T366L_N390N_K392M_T394W_T411N
AZ2077	L351Y_S400K_F405A_Y407V	T366L_N390N_K392M_T394W_T411S
AZ2078	L351Y_S400K_F405A_Y407V	T366L_N390N_K392M_T394W_T411T
AZ2079	L351Y_S400K_F405A_Y407V	T366L_N390N_K392F_T394W_T411N
AZ2080	L351Y_S400K_F405A_Y407V	T366L_N390N_K392F_T394W_T411S
AZ2081	L351Y_S400K_F405A_Y407V	T366L_N390N_K392F_T394W_T411T
AZ2082	L351Y_S400K_F405A_Y407V	T366L_N390D_K392L_T394W_T411N
AZ2083	L351Y_S400K_F405A_Y407V	T366L_N390D_K392L_T394W_T411S
AZ2084	L351Y_S400K_F405A_Y407V	T366L_N390D_K392L_T394W_T411T
AZ2085	L351Y_S400K_F405A_Y407V	T366L_N390D_K392M_T394W_T411N
AZ2086	L351Y_S400K_F405A_Y407V	T366L_N390D_K392M_T394W_T411S
AZ2087	L351Y_S400K_F405A_Y407V	T366L_N390D_K392M_T394W_T411T
AZ2088	L351Y_S400K_F405A_Y407V	T366L_N390D_K392F_T394W_T411N
AZ2089	L351Y_S400K_F405A_Y407V	T366L_N390D_K392F_T394W_T411S
AZ2090	L351Y_S400K_F405A_Y407V	T366L_N390D_K392F_T394W_T411T
AZ2091	L351Y_S400K_F405A_Y407V	T366L_N390E_K392L_T394W_T411N
AZ2092	L351Y_S400K_F405A_Y407V	T366L_N390E_K392L_T394W_T411L
AZ2093	L351Y_S400K_F405A_Y407V	T366L_N390E_K392L_T394W_T411S
AZ2094	L351Y_S400K_F405A_Y407V	T366L_N390E_K392L_T394W_T411T
AZ2095	L351Y_S400K_F405A_Y407V	T366L_N390E_K392M_T394W_T411N
AZ2096	L351Y_S400K_F405A_Y407V	T366L_N390E_K392M_T394W_T411L
AZ2097	L351Y_S400K_F405A_Y407V	T366L_N390E_K392M_T394W_T411S
AZ2098	L351Y_S400K_F405A_Y407V	T366L_N390E_K392M_T394W_T411T
AZ2099	L351Y_S400K_F405A_Y407V	T366L_N390E_K392F_T394W_T411N
AZ2100	L351Y_S400K_F405A_Y407V	T366L_N390E_K392F_T394W_T411L

AZ2101	L351Y_S400K_F405A_Y407V	T366L_N390E_K392F_T394W_T411S
AZ2102	L351Y_S400K_F405A_Y407V	T366L_N390E_K392F_T394W_T411T
AZ2103	L351Y_S400K_F405S_Y407I	T366L_N390N_K392L_T394W_T411N
AZ2104	L351Y_S400K_F405S_Y407I	T366L_N390N_K392L_T394W_T411S
AZ2105	L351Y_S400K_F405S_Y407I	T366L_N390N_K392L_T394W_T411T
AZ2106	L351Y_S400K_F405S_Y407I	T366L_N390N_K392M_T394W_T411N
AZ2107	L351Y_S400K_F405S_Y407I	T366L_N390N_K392M_T394W_T411S
AZ2108	L351Y_S400K_F405S_Y407I	T366L_N390N_K392M_T394W_T411T
AZ2109	L351Y_S400K_F405S_Y407I	T366L_N390N_K392F_T394W_T411N
AZ2110	L351Y_S400K_F405S_Y407I	T366L_N390N_K392F_T394W_T411S
AZ2111	L351Y_S400K_F405S_Y407I	T366L_N390N_K392F_T394W_T411T
AZ2112	L351Y_S400K_F405S_Y407I	T366L_N390D_K392L_T394W_T411N
AZ2113	L351Y_S400K_F405S_Y407I	T366L_N390D_K392L_T394W_T411S
AZ2114	L351Y_S400K_F405S_Y407I	T366L_N390D_K392L_T394W_T411T
AZ2115	L351Y_S400K_F405S_Y407I	T366L_N390D_K392M_T394W_T411N
AZ2116	L351Y_S400K_F405S_Y407I	T366L_N390D_K392M_T394W_T411S
AZ2117	L351Y_S400K_F405S_Y407I	T366L_N390D_K392M_T394W_T411T
AZ2118	L351Y_S400K_F405S_Y407I	T366L_N390D_K392F_T394W_T411N
AZ2119	L351Y_S400K_F405S_Y407I	T366L_N390D_K392F_T394W_T411S
AZ2120	L351Y_S400K_F405S_Y407I	T366L_N390D_K392F_T394W_T411T
AZ2121	L351Y_S400K_F405S_Y407I	T366L_N390E_K392L_T394W_T411N
AZ2122	L351Y_S400K_F405S_Y407I	T366L_N390E_K392L_T394W_T411L
AZ2123	L351Y_S400K_F405S_Y407I	T366L_N390E_K392L_T394W_T411S
AZ2124	L351Y_S400K_F405S_Y407I	T366L_N390E_K392L_T394W_T411T
AZ2125	L351Y_S400K_F405S_Y407I	T366L_N390E_K392M_T394W_T411N
AZ2126	L351Y_S400K_F405S_Y407I	T366L_N390E_K392M_T394W_T411L
AZ2127	L351Y_S400K_F405S_Y407I	T366L_N390E_K392M_T394W_T411S
AZ2128	L351Y_S400K_F405S_Y407I	T366L_N390E_K392M_T394W_T411T
AZ2129	L351Y_S400K_F405S_Y407I	T366L_N390E_K392F_T394W_T411N
AZ2130	L351Y_S400K_F405S_Y407I	T366L_N390E_K392F_T394W_T411L
AZ2131	L351Y_S400K_F405S_Y407I	T366L_N390E_K392F_T394W_T411S
AZ2132	L351Y_S400K_F405S_Y407I	T366L_N390E_K392F_T394W_T411T
AZ2133	L351Y_S400K_F405S_Y407V	T366L_N390N_K392L_T394W_T411N
AZ2134	L351Y_S400K_F405S_Y407V	T366L_N390N_K392L_T394W_T411S
AZ2135	L351Y_S400K_F405S_Y407V	T366L_N390N_K392L_T394W_T411T
AZ2136	L351Y_S400K_F405S_Y407V	T366L_N390N_K392M_T394W_T411N
AZ2137	L351Y_S400K_F405S_Y407V	T366L_N390N_K392M_T394W_T411S
AZ2138	L351Y_S400K_F405S_Y407V	T366L_N390N_K392M_T394W_T411T
AZ2139	L351Y_S400K_F405S_Y407V	T366L_N390N_K392F_T394W_T411N
AZ2140	L351Y_S400K_F405S_Y407V	T366L_N390N_K392F_T394W_T411S
AZ2141	L351Y_S400K_F405S_Y407V	T366L_N390N_K392F_T394W_T411T
AZ2142	L351Y_S400K_F405S_Y407V	T366L_N390D_K392L_T394W_T411N
AZ2143	L351Y_S400K_F405S_Y407V	T366L_N390D_K392L_T394W_T411S
AZ2144	L351Y_S400K_F405S_Y407V	T366L_N390D_K392L_T394W_T411T
AZ2145	L351Y_S400K_F405S_Y407V	T366L_N390D_K392M_T394W_T411N
AZ2146	L351Y_S400K_F405S_Y407V	T366L_N390D_K392M_T394W_T411S
AZ2147	L351Y_S400K_F405S_Y407V	T366L_N390D_K392M_T394W_T411T
AZ2148	L351Y_S400K_F405S_Y407V	T366L_N390D_K392F_T394W_T411N
AZ2149	L351Y_S400K_F405S_Y407V	T366L_N390D_K392F_T394W_T411S
AZ2150	L351Y_S400K_F405S_Y407V	T366L_N390D_K392F_T394W_T411T
AZ2151	L351Y_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411N
AZ2152	L351Y_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411L
AZ2153	L351Y_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411S
AZ2154	L351Y_S400K_F405S_Y407V	T366L_N390E_K392L_T394W_T411T
AZ2155	L351Y_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411N
AZ2156	L351Y_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411L
AZ2157	L351Y_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411S
AZ2158	L351Y_S400K_F405S_Y407V	T366L_N390E_K392M_T394W_T411T

AZ2159	L351Y_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411N
AZ2160	L351Y_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411L
AZ2161	L351Y_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411S
AZ2162	L351Y_S400K_F405S_Y407V	T366L_N390E_K392F_T394W_T411T
AZ2163	L351Y_S400K_F405T_Y407I	T366L_N390N_K392L_T394W_T411N
AZ2164	L351Y_S400K_F405T_Y407I	T366L_N390N_K392L_T394W_T411S
AZ2165	L351Y_S400K_F405T_Y407I	T366L_N390N_K392L_T394W_T411T
AZ2166	L351Y_S400K_F405T_Y407I	T366L_N390N_K392M_T394W_T411N
AZ2167	L351Y_S400K_F405T_Y407I	T366L_N390N_K392M_T394W_T411S
AZ2168	L351Y_S400K_F405T_Y407I	T366L_N390N_K392M_T394W_T411T
AZ2169	L351Y_S400K_F405T_Y407I	T366L_N390N_K392F_T394W_T411N
AZ2170	L351Y_S400K_F405T_Y407I	T366L_N390N_K392F_T394W_T411S
AZ2171	L351Y_S400K_F405T_Y407I	T366L_N390N_K392F_T394W_T411T
AZ2172	L351Y_S400K_F405T_Y407I	T366L_N390D_K392L_T394W_T411N
AZ2173	L351Y_S400K_F405T_Y407I	T366L_N390D_K392L_T394W_T411S
AZ2174	L351Y_S400K_F405T_Y407I	T366L_N390D_K392L_T394W_T411T
AZ2175	L351Y_S400K_F405T_Y407I	T366L_N390D_K392M_T394W_T411N
AZ2176	L351Y_S400K_F405T_Y407I	T366L_N390D_K392M_T394W_T411S
AZ2177	L351Y_S400K_F405T_Y407I	T366L_N390D_K392M_T394W_T411T
AZ2178	L351Y_S400K_F405T_Y407I	T366L_N390D_K392F_T394W_T411N
AZ2179	L351Y_S400K_F405T_Y407I	T366L_N390D_K392F_T394W_T411S
AZ2180	L351Y_S400K_F405T_Y407I	T366L_N390D_K392F_T394W_T411T
AZ2181	L351Y_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411N
AZ2182	L351Y_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411L
AZ2183	L351Y_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411S
AZ2184	L351Y_S400K_F405T_Y407I	T366L_N390E_K392L_T394W_T411T
AZ2185	L351Y_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411N
AZ2186	L351Y_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411L
AZ2187	L351Y_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411S
AZ2188	L351Y_S400K_F405T_Y407I	T366L_N390E_K392M_T394W_T411T
AZ2189	L351Y_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411N
AZ2190	L351Y_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411L
AZ2191	L351Y_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411S
AZ2192	L351Y_S400K_F405T_Y407I	T366L_N390E_K392F_T394W_T411T
AZ2193	L351Y_S400K_F405T_Y407V	T366L_N390N_K392L_T394W_T411N
AZ2194	L351Y_S400K_F405T_Y407V	T366L_N390N_K392L_T394W_T411S
AZ2195	L351Y_S400K_F405T_Y407V	T366L_N390N_K392L_T394W_T411T
AZ2196	L351Y_S400K_F405T_Y407V	T366L_N390N_K392M_T394W_T411N
AZ2197	L351Y_S400K_F405T_Y407V	T366L_N390N_K392M_T394W_T411S
AZ2198	L351Y_S400K_F405T_Y407V	T366L_N390N_K392M_T394W_T411T
AZ2199	L351Y_S400K_F405T_Y407V	T366L_N390N_K392F_T394W_T411N
AZ2200	L351Y_S400K_F405T_Y407V	T366L_N390N_K392F_T394W_T411S
AZ2201	L351Y_S400K_F405T_Y407V	T366L_N390N_K392F_T394W_T411T
AZ2202	L351Y_S400K_F405T_Y407V	T366L_N390D_K392L_T394W_T411N
AZ2203	L351Y_S400K_F405T_Y407V	T366L_N390D_K392L_T394W_T411S
AZ2204	L351Y_S400K_F405T_Y407V	T366L_N390D_K392L_T394W_T411T
AZ2205	L351Y_S400K_F405T_Y407V	T366L_N390D_K392M_T394W_T411N
AZ2206	L351Y_S400K_F405T_Y407V	T366L_N390D_K392M_T394W_T411S
AZ2207	L351Y_S400K_F405T_Y407V	T366L_N390D_K392M_T394W_T411T
AZ2208	L351Y_S400K_F405T_Y407V	T366L_N390D_K392F_T394W_T411N
AZ2209	L351Y_S400K_F405T_Y407V	T366L_N390D_K392F_T394W_T411S
AZ2210	L351Y_S400K_F405T_Y407V	T366L_N390D_K392F_T394W_T411T
AZ2211	L351Y_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411N
AZ2212	L351Y_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411L
AZ2213	L351Y_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411S
AZ2214	L351Y_S400K_F405T_Y407V	T366L_N390E_K392L_T394W_T411T
AZ2215	L351Y_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411N
AZ2216	L351Y_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411L

AZ2217	L351Y_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411S
AZ2218	L351Y_S400K_F405T_Y407V	T366L_N390E_K392M_T394W_T411T
AZ2219	L351Y_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411N
AZ2220	L351Y_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411L
AZ2221	L351Y_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411S
AZ2222	L351Y_S400K_F405T_Y407V	T366L_N390E_K392F_T394W_T411T
AZ2223	L351Y_S400K_F405V_Y407I	T366L_N390N_K392L_T394W_T411N
AZ2224	L351Y_S400K_F405V_Y407I	T366L_N390N_K392L_T394W_T411S
AZ2225	L351Y_S400K_F405V_Y407I	T366L_N390N_K392L_T394W_T411T
AZ2226	L351Y_S400K_F405V_Y407I	T366L_N390N_K392M_T394W_T411N
AZ2227	L351Y_S400K_F405V_Y407I	T366L_N390N_K392M_T394W_T411S
AZ2228	L351Y_S400K_F405V_Y407I	T366L_N390N_K392M_T394W_T411T
AZ2229	L351Y_S400K_F405V_Y407I	T366L_N390N_K392F_T394W_T411N
AZ2230	L351Y_S400K_F405V_Y407I	T366L_N390N_K392F_T394W_T411S
AZ2231	L351Y_S400K_F405V_Y407I	T366L_N390N_K392F_T394W_T411T
AZ2232	L351Y_S400K_F405V_Y407I	T366L_N390D_K392L_T394W_T411N
AZ2233	L351Y_S400K_F405V_Y407I	T366L_N390D_K392L_T394W_T411S
AZ2234	L351Y_S400K_F405V_Y407I	T366L_N390D_K392L_T394W_T411T
AZ2235	L351Y_S400K_F405V_Y407I	T366L_N390D_K392M_T394W_T411N
AZ2236	L351Y_S400K_F405V_Y407I	T366L_N390D_K392M_T394W_T411S
AZ2237	L351Y_S400K_F405V_Y407I	T366L_N390D_K392M_T394W_T411T
AZ2238	L351Y_S400K_F405V_Y407I	T366L_N390D_K392F_T394W_T411N
AZ2239	L351Y_S400K_F405V_Y407I	T366L_N390D_K392F_T394W_T411S
AZ2240	L351Y_S400K_F405V_Y407I	T366L_N390D_K392F_T394W_T411T
AZ2241	L351Y_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411N
AZ2242	L351Y_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411L
AZ2243	L351Y_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411S
AZ2244	L351Y_S400K_F405V_Y407I	T366L_N390E_K392L_T394W_T411T
AZ2245	L351Y_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411N
AZ2246	L351Y_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411L
AZ2247	L351Y_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411S
AZ2248	L351Y_S400K_F405V_Y407I	T366L_N390E_K392M_T394W_T411T
AZ2249	L351Y_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411N
AZ2250	L351Y_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411L
AZ2251	L351Y_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411S
AZ2252	L351Y_S400K_F405V_Y407I	T366L_N390E_K392F_T394W_T411T
AZ2253	L351Y_S400K_F405V_Y407V	T366L_N390N_K392L_T394W_T411N
AZ2254	L351Y_S400K_F405V_Y407V	T366L_N390N_K392L_T394W_T411S
AZ2255	L351Y_S400K_F405V_Y407V	T366L_N390N_K392L_T394W_T411T
AZ2256	L351Y_S400K_F405V_Y407V	T366L_N390N_K392M_T394W_T411N
AZ2257	L351Y_S400K_F405V_Y407V	T366L_N390N_K392M_T394W_T411S
AZ2258	L351Y_S400K_F405V_Y407V	T366L_N390N_K392M_T394W_T411T
AZ2259	L351Y_S400K_F405V_Y407V	T366L_N390N_K392F_T394W_T411N
AZ2260	L351Y_S400K_F405V_Y407V	T366L_N390N_K392F_T394W_T411S
AZ2261	L351Y_S400K_F405V_Y407V	T366L_N390N_K392F_T394W_T411T
AZ2262	L351Y_S400K_F405V_Y407V	T366L_N390D_K392L_T394W_T411N
AZ2263	L351Y_S400K_F405V_Y407V	T366L_N390D_K392L_T394W_T411S
AZ2264	L351Y_S400K_F405V_Y407V	T366L_N390D_K392L_T394W_T411T
AZ2265	L351Y_S400K_F405V_Y407V	T366L_N390D_K392M_T394W_T411N
AZ2266	L351Y_S400K_F405V_Y407V	T366L_N390D_K392M_T394W_T411S
AZ2267	L351Y_S400K_F405V_Y407V	T366L_N390D_K392M_T394W_T411T
AZ2268	L351Y_S400K_F405V_Y407V	T366L_N390D_K392F_T394W_T411N
AZ2269	L351Y_S400K_F405V_Y407V	T366L_N390D_K392F_T394W_T411S
AZ2270	L351Y_S400K_F405V_Y407V	T366L_N390D_K392F_T394W_T411T
AZ2271	L351Y_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411N
AZ2272	L351Y_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411L
AZ2273	L351Y_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411S
AZ2274	L351Y_S400K_F405V_Y407V	T366L_N390E_K392L_T394W_T411T

AZ2275	L351Y_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411N
AZ2276	L351Y_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411L
AZ2277	L351Y_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411S
AZ2278	L351Y_S400K_F405V_Y407V	T366L_N390E_K392M_T394W_T411T
AZ2279	L351Y_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411N
AZ2280	L351Y_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411L
AZ2281	L351Y_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411S
AZ2282	L351Y_S400K_F405V_Y407V	T366L_N390E_K392F_T394W_T411T
AZ2283	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409I_T411L
AZ2284	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409I_T411T
AZ2285	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409L_T411L
AZ2286	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409L_T411T
AZ2287	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409K_T411L
AZ2288	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409K_T411T
AZ2289	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409M_T411L
AZ2290	Y349C_D399C_F405A_Y407I	S354C_T366L_K392C_T394W_K409M_T411T
AZ2291	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409I_T411L
AZ2292	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409I_T411T
AZ2293	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409L_T411L
AZ2294	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409L_T411T
AZ2295	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409K_T411L
AZ2296	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409K_T411T
AZ2297	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409M_T411L
AZ2298	Y349C_D399C_F405A_Y407V	S354C_T366L_K392C_T394W_K409M_T411T
AZ2299	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409I_T411L
AZ2300	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409I_T411T
AZ2301	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409L_T411L
AZ2302	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409L_T411T
AZ2303	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409K_T411L
AZ2304	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409K_T411T
AZ2305	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409M_T411L
AZ2306	Y349C_D399C_F405S_Y407I	S354C_T366L_K392C_T394W_K409M_T411T
AZ2307	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409I_T411L
AZ2308	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409I_T411T
AZ2309	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409L_T411L
AZ2310	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409L_T411T
AZ2311	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409K_T411L
AZ2312	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409K_T411T
AZ2313	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409M_T411L
AZ2314	Y349C_D399C_F405S_Y407V	S354C_T366L_K392C_T394W_K409M_T411T
AZ2315	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409I_T411L
AZ2316	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409I_T411T
AZ2317	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409L_T411L
AZ2318	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409L_T411T
AZ2319	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409K_T411L
AZ2320	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409K_T411T
AZ2321	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409M_T411L
AZ2322	Y349C_D399C_F405V_Y407I	S354C_T366L_K392C_T394W_K409M_T411T
AZ2323	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409I_T411L
AZ2324	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409I_T411T
AZ2325	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409L_T411L
AZ2326	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409L_T411T
AZ2327	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409K_T411L
AZ2328	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409K_T411T
AZ2329	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409M_T411L
AZ2330	Y349C_D399C_F405V_Y407V	S354C_T366L_K392C_T394W_K409M_T411T
AZ2331	L351L_K370T_G371D_F405A_Y407I	T366L_K392L_T394W_K409M_T411R
AZ2332	L351L_K370T_G371D_F405A_Y407I	T366L_K392M_T394W_K409M_T411R

78/96

79/96

FIGURE 38

Table 7

Variant	Fc mutations chain A	Fc mutations chain B
AZ2439	T366V_K392L_K409F_T411E	L351Y_D399R_Y407A
AZ2440	T366V_K392E_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2441	T366I_K409F	L351Y_Y407A
AZ2442	T366I_K409F	L351Y_Y407I
AZ2443	T366I_K409F	L351Y_Y407V
AZ2444	T366M_K409F	L351Y_Y407A
AZ2445	T366M_K409F	L351Y_Y407I
AZ2446	T366M_K409F	L351Y_Y407V
AZ2447	T366V_K409F	L351Y_Y407A
AZ2448	T366V_K409F	L351Y_Y407I
AZ2449	T366V_K409F	L351Y_Y407V
AZ2450	T366I_T394S_K409W	L351Y_F405L_Y407A
AZ2451	T366I_T394S_K409W	L351Y_F405L_Y407I
AZ2452	T366I_T394S_K409W	L351Y_F405L_Y407L
AZ2453	T366I_T394S_K409W	L351Y_F405L_Y407M
AZ2454	T366I_T394S_K409W	L351Y_F405L_Y407V
AZ2455	T366I_T394S_K409W	L351Y_F405M_Y407A
AZ2456	T366I_T394S_K409W	L351Y_F405M_Y407I
AZ2457	T366I_T394S_K409W	L351Y_F405M_Y407L
AZ2458	T366I_T394S_K409W	L351Y_F405M_Y407M
AZ2459	T366I_T394S_K409W	L351Y_F405M_Y407V
AZ2460	T366I_T394S_K409W	L351Y_F405F_Y407A
AZ2461	T366I_T394S_K409W	L351Y_F405F_Y407I
AZ2462	T366I_T394S_K409W	L351Y_F405F_Y407L
AZ2463	T366I_T394S_K409W	L351Y_F405F_Y407M
AZ2464	T366I_T394S_K409W	L351Y_F405F_Y407V
AZ2465	T366I_T394V_K409W	L351Y_F405L_Y407A
AZ2466	T366I_T394V_K409W	L351Y_F405L_Y407I
AZ2467	T366I_T394V_K409W	L351Y_F405L_Y407L
AZ2468	T366I_T394V_K409W	L351Y_F405L_Y407M
AZ2469	T366I_T394V_K409W	L351Y_F405L_Y407V
AZ2470	T366I_T394V_K409W	L351Y_F405M_Y407A
AZ2471	T366I_T394V_K409W	L351Y_F405M_Y407I
AZ2472	T366I_T394V_K409W	L351Y_F405M_Y407L
AZ2473	T366I_T394V_K409W	L351Y_F405M_Y407M
AZ2474	T366I_T394V_K409W	L351Y_F405M_Y407V
AZ2475	T366I_T394V_K409W	L351Y_F405F_Y407A
AZ2476	T366I_T394V_K409W	L351Y_F405F_Y407I
AZ2477	T366I_T394V_K409W	L351Y_F405F_Y407L
AZ2478	T366I_T394V_K409W	L351Y_F405F_Y407M
AZ2479	T366I_T394V_K409W	L351Y_F405F_Y407V
AZ2480	T366M_T394S_K409W	L351Y_F405L_Y407A
AZ2481	T366M_T394S_K409W	L351Y_F405L_Y407I
AZ2482	T366M_T394S_K409W	L351Y_F405L_Y407L
AZ2483	T366M_T394S_K409W	L351Y_F405L_Y407M
AZ2484	T366M_T394S_K409W	L351Y_F405L_Y407V
AZ2485	T366M_T394S_K409W	L351Y_F405M_Y407A
AZ2486	T366M_T394S_K409W	L351Y_F405M_Y407I
AZ2487	T366M_T394S_K409W	L351Y_F405M_Y407L
AZ2488	T366M_T394S_K409W	L351Y_F405M_Y407M
AZ2489	T366M_T394S_K409W	L351Y_F405M_Y407V
AZ2490	T366M_T394S_K409W	L351Y_F405F_Y407A
AZ2491	T366M_T394S_K409W	L351Y_F405F_Y407I
AZ2492	T366M_T394S_K409W	L351Y_F405F_Y407L

AZ2493	T366M_T394S_K409W	L351Y_F405F_Y407M
AZ2494	T366M_T394S_K409W	L351Y_F405F_Y407V
AZ2495	T366M_T394V_K409W	L351Y_F405L_Y407A
AZ2496	T366M_T394V_K409W	L351Y_F405L_Y407I
AZ2497	T366M_T394V_K409W	L351Y_F405L_Y407L
AZ2498	T366M_T394V_K409W	L351Y_F405L_Y407M
AZ2499	T366M_T394V_K409W	L351Y_F405L_Y407V
AZ2500	T366M_T394V_K409W	L351Y_F405M_Y407A
AZ2501	T366M_T394V_K409W	L351Y_F405M_Y407I
AZ2502	T366M_T394V_K409W	L351Y_F405M_Y407L
AZ2503	T366M_T394V_K409W	L351Y_F405M_Y407M
AZ2504	T366M_T394V_K409W	L351Y_F405M_Y407V
AZ2505	T366M_T394V_K409W	L351Y_F405F_Y407A
AZ2506	T366M_T394V_K409W	L351Y_F405F_Y407I
AZ2507	T366M_T394V_K409W	L351Y_F405F_Y407L
AZ2508	T366M_T394V_K409W	L351Y_F405F_Y407M
AZ2509	T366M_T394V_K409W	L351Y_F405F_Y407V
AZ2510	T366V_T394S_K409W	L351Y_F405L_Y407A
AZ2511	T366V_T394S_K409W	L351Y_F405L_Y407I
AZ2512	T366V_T394S_K409W	L351Y_F405L_Y407L
AZ2513	T366V_T394S_K409W	L351Y_F405L_Y407M
AZ2514	T366V_T394S_K409W	L351Y_F405L_Y407V
AZ2515	T366V_T394S_K409W	L351Y_F405M_Y407A
AZ2516	T366V_T394S_K409W	L351Y_F405M_Y407I
AZ2517	T366V_T394S_K409W	L351Y_F405M_Y407L
AZ2518	T366V_T394S_K409W	L351Y_F405M_Y407M
AZ2519	T366V_T394S_K409W	L351Y_F405M_Y407V
AZ2520	T366V_T394S_K409W	L351Y_F405F_Y407A
AZ2521	T366V_T394S_K409W	L351Y_F405F_Y407I
AZ2522	T366V_T394S_K409W	L351Y_F405F_Y407L
AZ2523	T366V_T394S_K409W	L351Y_F405F_Y407M
AZ2524	T366V_T394S_K409W	L351Y_F405F_Y407V
AZ2525	T366V_T394V_K409W	L351Y_F405L_Y407A
AZ2526	T366V_T394V_K409W	L351Y_F405L_Y407I
AZ2527	T366V_T394V_K409W	L351Y_F405L_Y407L
AZ2528	T366V_T394V_K409W	L351Y_F405L_Y407M
AZ2529	T366V_T394V_K409W	L351Y_F405L_Y407V
AZ2530	T366V_T394V_K409W	L351Y_F405M_Y407A
AZ2531	T366V_T394V_K409W	L351Y_F405M_Y407I
AZ2532	T366V_T394V_K409W	L351Y_F405M_Y407L
AZ2533	T366V_T394V_K409W	L351Y_F405M_Y407M
AZ2534	T366V_T394V_K409W	L351Y_F405M_Y407V
AZ2535	T366V_T394V_K409W	L351Y_F405F_Y407A
AZ2536	T366V_T394V_K409W	L351Y_F405F_Y407I
AZ2537	T366V_T394V_K409W	L351Y_F405F_Y407L
AZ2538	T366V_T394V_K409W	L351Y_F405F_Y407M
AZ2539	T366V_T394V_K409W	L351Y_F405F_Y407V
AZ2540	T366I_K392C_K409F_T411I	L351Y_D399C_Y407A
AZ2541	T366I_K392C_K409F_T411I	L351Y_D399C_Y407I
AZ2542	T366I_K392C_K409F_T411I	L351Y_D399C_Y407V
AZ2543	T366I_K392C_K409F_T411L	L351Y_D399C_Y407A
AZ2544	T366I_K392C_K409F_T411L	L351Y_D399C_Y407I
AZ2545	T366I_K392C_K409F_T411L	L351Y_D399C_Y407V
AZ2546	T366I_K392C_K409F_T411T	L351Y_D399C_Y407A
AZ2547	T366I_K392C_K409F_T411T	L351Y_D399C_Y407I
AZ2548	T366I_K392C_K409F_T411T	L351Y_D399C_Y407V
AZ2549	T366I_K392L_K409F_T411E	L351Y_D399R_Y407A
AZ2550	T366I_K392L_K409F_T411E	L351Y_D399R_Y407I

AZ2551	T366I_K392L_K409F_T411E	L351Y_D399R_Y407V
AZ2552	T366M_K392C_K409F_T411I	L351Y_D399C_Y407A
AZ2553	T366M_K392C_K409F_T411I	L351Y_D399C_Y407I
AZ2554	T366M_K392C_K409F_T411I	L351Y_D399C_Y407V
AZ2555	T366M_K392C_K409F_T411L	L351Y_D399C_Y407A
AZ2556	T366M_K392C_K409F_T411L	L351Y_D399C_Y407I
AZ2557	T366M_K392C_K409F_T411L	L351Y_D399C_Y407V
AZ2558	T366M_K392C_K409F_T411T	L351Y_D399C_Y407A
AZ2559	T366M_K392C_K409F_T411T	L351Y_D399C_Y407I
AZ2560	T366M_K392C_K409F_T411T	L351Y_D399C_Y407V
AZ2561	T366M_K392L_K409F_T411E	L351Y_D399R_Y407A
AZ2562	T366M_K392L_K409F_T411E	L351Y_D399R_Y407I
AZ2563	T366M_K392L_K409F_T411E	L351Y_D399R_Y407V
AZ2564	T366V_K392C_K409F_T411I	L351Y_D399C_Y407A
AZ2565	T366V_K392C_K409F_T411I	L351Y_D399C_Y407I
AZ2566	T366V_K392C_K409F_T411I	L351Y_D399C_Y407V
AZ2567	T366V_K392C_K409F_T411L	L351Y_D399C_Y407A
AZ2568	T366V_K392C_K409F_T411L	L351Y_D399C_Y407I
AZ2569	T366V_K392C_K409F_T411L	L351Y_D399C_Y407V
AZ2570	T366V_K392C_K409F_T411T	L351Y_D399C_Y407A
AZ2571	T366V_K392C_K409F_T411T	L351Y_D399C_Y407I
AZ2572	T366V_K392C_K409F_T411T	L351Y_D399C_Y407V
AZ2573	T366V_K392L_K409F_T411E	L351Y_D399R_Y407A
AZ2574	T366V_K392L_K409F_T411E	L351Y_D399R_Y407I
AZ2575	T366V_K392L_K409F_T411E	L351Y_D399R_Y407V
AZ2576	T366I_K392D_K409F_T411D	L351Y_D399R_S400S_Y407A
AZ2577	T366I_K392D_K409F_T411D	L351Y_D399R_S400S_Y407I
AZ2578	T366I_K392D_K409F_T411D	L351Y_D399R_S400S_Y407V
AZ2579	T366I_K392D_K409F_T411D	L351Y_D399K_S400S_Y407A
AZ2580	T366I_K392D_K409F_T411D	L351Y_D399K_S400S_Y407I
AZ2581	T366I_K392D_K409F_T411D	L351Y_D399K_S400S_Y407V
AZ2582	T366I_K392D_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2583	T366I_K392D_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2584	T366I_K392D_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2585	T366I_K392D_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2586	T366I_K392D_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2587	T366I_K392D_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2588	T366I_K392D_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2589	T366I_K392D_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2590	T366I_K392D_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2591	T366I_K392D_K409F_T411E	L351Y_D399K_S400S_Y407A
AZ2592	T366I_K392D_K409F_T411E	L351Y_D399K_S400S_Y407I
AZ2593	T366I_K392D_K409F_T411E	L351Y_D399K_S400S_Y407V
AZ2594	T366I_K392E_K409F_T411D	L351Y_D399R_S400S_Y407A
AZ2595	T366I_K392E_K409F_T411D	L351Y_D399R_S400S_Y407I
AZ2596	T366I_K392E_K409F_T411D	L351Y_D399R_S400S_Y407V
AZ2597	T366I_K392E_K409F_T411D	L351Y_D399K_S400S_Y407A
AZ2598	T366I_K392E_K409F_T411D	L351Y_D399K_S400S_Y407I
AZ2599	T366I_K392E_K409F_T411D	L351Y_D399K_S400S_Y407V
AZ2600	T366I_K392E_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2601	T366I_K392E_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2602	T366I_K392E_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2603	T366I_K392E_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2604	T366I_K392E_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2605	T366I_K392E_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2606	T366I_K392E_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2607	T366I_K392E_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2608	T366I_K392E_K409F_T411E	L351Y_D399R_S400S_Y407V

AZ2609	T366I_K392E_K409F_T411E	L351Y_D399K_S400S_Y407A
AZ2610	T366I_K392E_K409F_T411E	L351Y_D399K_S400S_Y407I
AZ2611	T366I_K392E_K409F_T411E	L351Y_D399K_S400S_Y407V
AZ2612	T366M_K392D_K409F_T411D	L351Y_D399R_S400S_Y407A
AZ2613	T366M_K392D_K409F_T411D	L351Y_D399R_S400S_Y407I
AZ2614	T366M_K392D_K409F_T411D	L351Y_D399R_S400S_Y407V
AZ2615	T366M_K392D_K409F_T411D	L351Y_D399K_S400S_Y407A
AZ2616	T366M_K392D_K409F_T411D	L351Y_D399K_S400S_Y407I
AZ2617	T366M_K392D_K409F_T411D	L351Y_D399K_S400S_Y407V
AZ2618	T366M_K392D_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2619	T366M_K392D_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2620	T366M_K392D_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2621	T366M_K392D_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2622	T366M_K392D_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2623	T366M_K392D_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2624	T366M_K392D_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2625	T366M_K392D_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2626	T366M_K392D_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2627	T366M_K392D_K409F_T411E	L351Y_D399K_S400S_Y407A
AZ2628	T366M_K392D_K409F_T411E	L351Y_D399K_S400S_Y407I
AZ2629	T366M_K392D_K409F_T411E	L351Y_D399K_S400S_Y407V
AZ2630	T366M_K392E_K409F_T411D	L351Y_D399R_S400S_Y407A
AZ2631	T366M_K392E_K409F_T411D	L351Y_D399R_S400S_Y407I
AZ2632	T366M_K392E_K409F_T411D	L351Y_D399R_S400S_Y407V
AZ2633	T366M_K392E_K409F_T411D	L351Y_D399K_S400S_Y407A
AZ2634	T366M_K392E_K409F_T411D	L351Y_D399K_S400S_Y407I
AZ2635	T366M_K392E_K409F_T411D	L351Y_D399K_S400S_Y407V
AZ2636	T366M_K392E_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2637	T366M_K392E_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2638	T366M_K392E_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2639	T366M_K392E_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2640	T366M_K392E_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2641	T366M_K392E_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2642	T366M_K392E_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2643	T366M_K392E_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2644	T366M_K392E_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2645	T366M_K392E_K409F_T411E	L351Y_D399K_S400S_Y407A
AZ2646	T366M_K392E_K409F_T411E	L351Y_D399K_S400S_Y407I
AZ2647	T366M_K392E_K409F_T411E	L351Y_D399K_S400S_Y407V
AZ2648	T366V_K392D_K409F_T411D	L351Y_D399R_S400S_Y407A
AZ2649	T366V_K392D_K409F_T411D	L351Y_D399R_S400S_Y407I
AZ2650	T366V_K392D_K409F_T411D	L351Y_D399R_S400S_Y407V
AZ2651	T366V_K392D_K409F_T411D	L351Y_D399K_S400S_Y407A
AZ2652	T366V_K392D_K409F_T411D	L351Y_D399K_S400S_Y407I
AZ2653	T366V_K392D_K409F_T411D	L351Y_D399K_S400S_Y407V
AZ2654	T366V_K392D_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2655	T366V_K392D_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2656	T366V_K392D_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2657	T366V_K392D_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2658	T366V_K392D_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2659	T366V_K392D_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2660	T366V_K392D_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2661	T366V_K392D_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2662	T366V_K392D_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2663	T366V_K392D_K409F_T411E	L351Y_D399K_S400S_Y407A
AZ2664	T366V_K392D_K409F_T411E	L351Y_D399K_S400S_Y407I
AZ2665	T366V_K392D_K409F_T411E	L351Y_D399K_S400S_Y407V
AZ2666	T366V_K392E_K409F_T411D	L351Y_D399R_S400S_Y407A

AZ2667	T366V_K392E_K409F_T411D	L351Y_D399R_S400S_Y407I
AZ2668	T366V_K392E_K409F_T411D	L351Y_D399R_S400S_Y407V
AZ2669	T366V_K392E_K409F_T411D	L351Y_D399K_S400S_Y407A
AZ2670	T366V_K392E_K409F_T411D	L351Y_D399K_S400S_Y407I
AZ2671	T366V_K392E_K409F_T411D	L351Y_D399K_S400S_Y407V
AZ2672	T366V_K392E_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2673	T366V_K392E_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2674	T366V_K392E_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2675	T366V_K392E_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2676	T366V_K392E_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2677	T366V_K392E_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2678	T366V_K392E_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2679	T366V_K392E_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2680	T366V_K392E_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2681	T366V_K392E_K409F_T411E	L351Y_D399K_S400S_Y407A
AZ2682	T366V_K392E_K409F_T411E	L351Y_D399K_S400S_Y407I
AZ2683	T366V_K392E_K409F_T411E	L351Y_D399K_S400S_Y407V
AZ2684	S354C_T366I_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407A
AZ2685	S354C_T366I_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407I
AZ2686	S354C_T366I_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407V
AZ2687	S354C_T366I_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407A
AZ2688	S354C_T366I_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407I
AZ2689	S354C_T366I_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407V
AZ2690	S354C_T366I_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407A
AZ2691	S354C_T366I_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407I
AZ2692	S354C_T366I_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407V
AZ2693	S354C_T366M_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407A
AZ2694	S354C_T366M_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407I
AZ2695	S354C_T366M_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407V
AZ2696	S354C_T366M_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407A
AZ2697	S354C_T366M_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407I
AZ2698	S354C_T366M_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407V
AZ2699	S354C_T366M_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407A
AZ2700	S354C_T366M_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407I
AZ2701	S354C_T366M_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407V
AZ2702	S354C_T366V_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407A
AZ2703	S354C_T366V_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407I
AZ2704	S354C_T366V_K392C_K409F_T411I	Y349C_L351Y_D399C_Y407V
AZ2705	S354C_T366V_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407A
AZ2706	S354C_T366V_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407I
AZ2707	S354C_T366V_K392C_K409F_T411L	Y349C_L351Y_D399C_Y407V
AZ2708	S354C_T366V_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407A
AZ2709	S354C_T366V_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407I
AZ2710	S354C_T366V_K392C_K409F_T411T	Y349C_L351Y_D399C_Y407V
AZ2711	Q362E_T366I_K392L_K409F_T411E	L351Y_G371G_D399R_Y407A
AZ2712	Q362E_T366I_K392L_K409F_T411E	L351Y_G371G_D399R_Y407I
AZ2713	Q362E_T366I_K392L_K409F_T411E	L351Y_G371G_D399R_Y407V
AZ2714	Q362E_T366I_K392L_K409F_T411E	L351Y_G371S_D399R_Y407A
AZ2715	Q362E_T366I_K392L_K409F_T411E	L351Y_G371S_D399R_Y407I
AZ2716	Q362E_T366I_K392L_K409F_T411E	L351Y_G371S_D399R_Y407V
AZ2717	Q362E_T366M_K392L_K409F_T411E	L351Y_G371G_D399R_Y407A
AZ2718	Q362E_T366M_K392L_K409F_T411E	L351Y_G371G_D399R_Y407I
AZ2719	Q362E_T366M_K392L_K409F_T411E	L351Y_G371G_D399R_Y407V
AZ2720	Q362E_T366M_K392L_K409F_T411E	L351Y_G371S_D399R_Y407A
AZ2721	Q362E_T366M_K392L_K409F_T411E	L351Y_G371S_D399R_Y407I
AZ2722	Q362E_T366M_K392L_K409F_T411E	L351Y_G371S_D399R_Y407V
AZ2723	Q362E_T366V_K392L_K409F_T411E	L351Y_G371G_D399R_Y407A
AZ2724	Q362E_T366V_K392L_K409F_T411E	L351Y_G371G_D399R_Y407I

AZ2725	Q362E_T366V_K392L_K409F_T411E	L351Y_G371G_D399R_Y407V
AZ2726	Q362E_T366V_K392L_K409F_T411E	L351Y_G371S_D399R_Y407A
AZ2727	Q362E_T366V_K392L_K409F_T411E	L351Y_G371S_D399R_Y407I
AZ2728	Q362E_T366V_K392L_K409F_T411E	L351Y_G371S_D399R_Y407V
AZ2729	T366I_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2730	T366I_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2731	T366I_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2732	T366I_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2733	T366I_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2734	T366I_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2735	T366I_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2736	T366I_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2737	T366I_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2738	T366I_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2739	T366I_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2740	T366I_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2741	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2742	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2743	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2744	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2745	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2746	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2747	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2748	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2749	T366I_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2750	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2751	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2752	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2753	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2754	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2755	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2756	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2757	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2758	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2759	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2760	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2761	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2762	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2763	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2764	T366I_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2765	T366I_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2766	T366I_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2767	T366I_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2768	T366I_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2769	T366I_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2770	T366I_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2771	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2772	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2773	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2774	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2775	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2776	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2777	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2778	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2779	T366I_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2780	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2781	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2782	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407V

AZ2783	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2784	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2785	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2786	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2787	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2788	T366I_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2789	T366I_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2790	T366I_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2791	T366I_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2792	T366I_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2793	T366I_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2794	T366I_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2795	T366I_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2796	T366I_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2797	T366I_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2798	T366I_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2799	T366I_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2800	T366I_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2801	T366M_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2802	T366M_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2803	T366M_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2804	T366M_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2805	T366M_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2806	T366M_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2807	T366M_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2808	T366M_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2809	T366M_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2810	T366M_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2811	T366M_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2812	T366M_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2813	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2814	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2815	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2816	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2817	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2818	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2819	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2820	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2821	T366M_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2822	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2823	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2824	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2825	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2826	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2827	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2828	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2829	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2830	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2831	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2832	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2833	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2834	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2835	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2836	T366M_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2837	T366M_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2838	T366M_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2839	T366M_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2840	T366M_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A

AZ2841	T366M_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2842	T366M_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2843	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2844	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2845	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2846	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2847	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2848	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2849	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2850	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2851	T366M_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2852	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2853	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2854	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2855	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2856	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2857	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2858	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2859	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2860	T366M_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2861	T366M_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2862	T366M_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2863	T366M_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2864	T366M_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2865	T366M_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2866	T366M_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2867	T366M_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2868	T366M_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2869	T366M_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2870	T366M_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2871	T366M_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2872	T366M_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2873	T366V_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2874	T366V_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2875	T366V_N390R_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2876	T366V_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2877	T366V_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2878	T366V_N390R_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2879	T366V_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2880	T366V_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2881	T366V_N390R_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2882	T366V_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2883	T366V_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2884	T366V_N390R_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2885	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2886	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2887	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2888	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2889	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2890	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2891	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2892	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2893	T366V_N390N_K392I_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2894	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2895	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2896	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2897	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2898	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I

AZ2899	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2900	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2901	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2902	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2903	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2904	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2905	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2906	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2907	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2908	T366V_N390N_K392L_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2909	T366V_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2910	T366V_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2911	T366V_N390N_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2912	T366V_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2913	T366V_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2914	T366V_N390N_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2915	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2916	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2917	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2918	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2919	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2920	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2921	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2922	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2923	T366V_N390D_K392I_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2924	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407A
AZ2925	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407I
AZ2926	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400R_Y407V
AZ2927	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407A
AZ2928	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407I
AZ2929	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400K_Y407V
AZ2930	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407A
AZ2931	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407I
AZ2932	T366V_N390D_K392L_K409F_T411E	L351Y_D399R_S400S_Y407V
AZ2933	T366V_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2934	T366V_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2935	T366V_N390K_K392L_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2936	T366V_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2937	T366V_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2938	T366V_N390K_K392L_K409F_T411E	L351Y_D399R_S400E_Y407V
AZ2939	T366V_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407A
AZ2940	T366V_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407I
AZ2941	T366V_N390K_K392K_K409F_T411E	L351Y_D399R_S400D_Y407V
AZ2942	T366V_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407A
AZ2943	T366V_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407I
AZ2944	T366V_N390K_K392K_K409F_T411E	L351Y_D399R_S400E_Y407V

Figure 39A




Figure 39B




FIGURE: 40-A

Scaffold 1 Sequence Alignment:




FIGURE 41




FIGURE 42




FIGURE 43




FIGURE 44

QVKLEESGGGLVQAGDSL RVSCAASGRDFSDYVMGWFRQAPGKEREFVAAISRNG
LTTRYADSVKGRFTISRDNNDKNMVYLQMNSLKPEDTAVYYCAVNSAGTYVSPRSREY
DYWGQGTQVTVSS

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2013/000471

A. CLASSIFICATION OF SUBJECT MATTER IPC: <i>C07K 16/46</i> (2006.01) , <i>A61K 39/395</i> (2006.01) , <i>A61P 35/00</i> (2006.01) , <i>A61P 37/02</i> (2006.01) , <i>C07K 16/12</i> (2006.01) , <i>C07K 16/24</i> (2006.01), <i>C07K 16/28</i> (2006.01) , <i>C07K 16/30</i> (2006.01) According to International Patent Classification (IPC) or to both national classification and IPC																	
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) <i>C07K 16/46</i> (2006.01) , <i>A61K 39/395</i> (2006.01) , <i>A61P 35/00</i> (2006.01) , <i>A61P 37/02</i> (2006.01) , <i>C07K 16/12</i> (2006.01) , <i>C07K 16/24</i> (2006.01), <i>C07K 16/28</i> (2006.01) , <i>C07K 16/30</i> (2006.01) Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used) Databases: TotalPatent, CAPlus, Biosis, Intellect Keywords: sdAb, bispecific, knob, dAb, V_HH, nanobody, HCAb, single domain																	
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>Y</td> <td>BELL, A. et al. "Differential tumor-targeting abilities of three single-domain antibody formats" CANCER LETTERS. 28 August 2009 (28-08-2009), V. 289(1), p.81-90. ISSN: 1872-7980. whole document</td> <td>1-55</td> </tr> <tr> <td>Y</td> <td>MERCHANT, A.M. et al. "An efficient route to human bispecific IgG" NATURE BIOTECHNOLOGY. July 1998 (07-1998). V. 16. p. 677-81. ISSN 1087-0156. whole document</td> <td>1-55</td> </tr> <tr> <td>P, Y</td> <td>WO2012058768 A1 (CABRERA, E.E. et al.) 10 May 2012 (10-05-2012) whole document</td> <td>1-55</td> </tr> <tr> <td>P, X</td> <td>WO2013/002362 A1 (MIMOTO, F. et al.) 03 January 2013 (03-01-2013) whole document</td> <td>1-55</td> </tr> </tbody> </table>			Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	Y	BELL, A. et al. "Differential tumor-targeting abilities of three single-domain antibody formats" CANCER LETTERS. 28 August 2009 (28-08-2009), V. 289(1), p.81-90. ISSN: 1872-7980. whole document	1-55	Y	MERCHANT, A.M. et al. "An efficient route to human bispecific IgG" NATURE BIOTECHNOLOGY. July 1998 (07-1998). V. 16. p. 677-81. ISSN 1087-0156. whole document	1-55	P, Y	WO2012058768 A1 (CABRERA, E.E. et al.) 10 May 2012 (10-05-2012) whole document	1-55	P, X	WO2013/002362 A1 (MIMOTO, F. et al.) 03 January 2013 (03-01-2013) whole document	1-55
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.															
Y	BELL, A. et al. "Differential tumor-targeting abilities of three single-domain antibody formats" CANCER LETTERS. 28 August 2009 (28-08-2009), V. 289(1), p.81-90. ISSN: 1872-7980. whole document	1-55															
Y	MERCHANT, A.M. et al. "An efficient route to human bispecific IgG" NATURE BIOTECHNOLOGY. July 1998 (07-1998). V. 16. p. 677-81. ISSN 1087-0156. whole document	1-55															
P, Y	WO2012058768 A1 (CABRERA, E.E. et al.) 10 May 2012 (10-05-2012) whole document	1-55															
P, X	WO2013/002362 A1 (MIMOTO, F. et al.) 03 January 2013 (03-01-2013) whole document	1-55															
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex. <table border="1"> <tbody> <tr> <td>* Special categories of cited documents :</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"E" earlier application or patent but published on or after the international filing date</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"&" document member of the same patent family</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </tbody> </table>			* Special categories of cited documents :	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means		"P" document published prior to the international filing date but later than the priority date claimed				
* Special categories of cited documents :	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family																
"O" document referring to an oral disclosure, use, exhibition or other means																	
"P" document published prior to the international filing date but later than the priority date claimed																	
Date of the actual completion of the international search 18 July 2013 (18-07-2013)		Date of mailing of the international search report 15 August 2013 (15-08-2013)															
Name and mailing address of the ISA/CA Canadian Intellectual Property Office Place du Portage I, C114 - 1st Floor, Box PCT 50 Victoria Street Gatineau, Quebec K1A 0C9 Facsimile No.: 001-819-953-2476		Authorized officer Rena Oulton (819) 997-4567															

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. ☒ Claim Nos. : 56-62

because they relate to subject matter not required to be searched by this Authority, namely :

Although claims 56-62 encompass a method of treatment of the human/animal body which this Authority is not obliged to search under Rule 39.1(iv) of the PCT, the search has been carried out based on the alleged effects of the compounds referred to therein.

2. ☒ Claim Nos. : 1-62

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :

See supplemental sheet.

3. ☐ Claim Nos. :

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☐ No protest accompanied the payment of additional search fees.

Continuation of Box No. II, part 2:

The claims encompass a very large number of possible embodiments while the description discloses, and provides support for, only a relatively small proportion of those embodiments (see Example 10). As such, the search has been directed only towards claimed embodiments that relate to specifically disclosed compounds, or prepared or tested compositions and a structural generalization of these. The extent of the search was limited to the specifically disclosed compound of Example 10 (anti-EGFR antibody single domain antibody, or sdAb, fused to an Fc domain which is a heterodimer having the mutations T350V_L351Y_F405A_Y407V in Chain A, and T350V_T366L_K392L_T394W in Chain B) as well as a structural generalization of these (single domain antigen-binding construct fused with a heterodimeric Fc having a variant constant domain).

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2013/000471

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
WO2012058768A1	10 May 2012 (10-05-2012)	AU2011325833A1 CA2815266A1 US2012149876A1 WO2012058768A8	30 May 2013 (30-05-2013) 10 May 2012 (10-05-2012) 14 June 2012 (14-06-2012) 28 June 2012 (28-06-2012)
WO2013002362A1	03 January 2013 (03-01-2013)	TW201305215A	01 February 2013 (01-02-2013)