



(86) Date de dépôt PCT/PCT Filing Date: 2011/06/01
(87) Date publication PCT/PCT Publication Date: 2011/12/08
(45) Date de délivrance/Issue Date: 2018/08/21
(85) Entrée phase nationale/National Entry: 2012/11/30
(86) N° demande PCT/PCT Application No.: EP 2011/059139
(87) N° publication PCT/PCT Publication No.: 2011/151412
(30) Priorité/Priority: 2010/06/01 (US12/791,681)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),
A61P 35/00 (2006.01), *C07K 16/28* (2006.01)
(72) Inventeurs/Inventors:
GOETSCH, LILIANE, FR;
WURCH, THIERRY, FR;
BES, CEDRIC, FR
(73) Propriétaire/Owner:
PIERRE FABRE MEDICAMENT, FR
(74) Agent: TORYS LLP

(54) Titre : NOUVEL ANTICORPS ANTI-cMET
(54) Title: NOVEL ANTI-cMET ANTIBODY

(57) **Abrégé/Abstract:**

The present invention relates to a novel divalent antibody capable of binding specifically to the human c-Met receptor and/or capable of specifically inhibiting the tyrosine kinase activity of said receptor, preferably both in a ligand-dependent and in a ligand-independent manner as well as the amino acid and nucleic acid sequences coding for said antibody. More preferably said antibody comprises a modified hinge region and exhibits an improved antagonistic activity. More particularly, the antibody according to the invention is capable of inhibiting the c-Met dimerization. The invention likewise comprises the use of said antibody as a medicament for the prophylactic and/or therapeutic treatment of cancers, preferably for cancer characterized by a ligand- independent activation of c-Met, or any pathology connected with the over expression of said receptor as well as in processes or kits for diagnosis of illnesses connected with the over-expression of c-Met. The invention finally comprises products and/or compositions comprising such an antibody in combination with other antibodies and/or chemical compounds directed against other growth factors involved in tumor progression or metastasis and/or compounds and/or anti-cancer agents or agents conjugated with toxins and their use for the prevention and/or the treatment of certain cancers.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
8 December 2011 (08.12.2011)



(10) International Publication Number
WO 2011/151412 A1

(51) International Patent Classification:

A61K 39/395 (2006.01) *C07K 16/28* (2006.01)
A61P 35/00 (2006.01)

(21) International Application Number:

PCT/EP2011/059139

(22) International Filing Date:

1 June 2011 (01.06.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12/791,681 1 June 2010 (01.06.2010) US

(71) Applicant (for all designated States except US):
PIERRE FABRE MEDICAMENT [FR/FR]; 45, place
Abel Gance, F-92100 Boulogne-Billancourt (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GOETSCH, Liliane**
[FR/FR]; 15 route de Cluses, F-74130 Ayze (FR).
WURCH, Thierry [FR/FR]; 7 route des Etoles, Le Clos
Bellevue, F-74140 Machilly (FR). **BES, Cédric** [FR/FR];
12 boulevard des Acanthes, F-30250 Villevieille (FR).

(74) Agent: **WARCOIN, AHNER, TEXIER, LE FORESTI-
ER, CALLON DE LAMARCK, COLLIN, TETAZ,
FAIVRE PETIT- Cabinet Regimbeau**; 20, rue de
Chazelles, F-75847 Paris Cedex 17 (FR).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,

HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of
the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))
- with (an) indication(s) in relation to deposited biological
material furnished under Rule 13bis separately from the
description (Rules 13bis.4(d)(i) and 48.2(a)(viii))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: NOVEL ANTI-CMET ANTIBODY

(57) Abstract: The present invention relates to a novel divalent antibody capable of binding specifically to the human c-Met receptor and/or capable of specifically inhibiting the tyrosine kinase activity of said receptor, preferably both in a ligand-dependent and in a ligand-independent manner as well as the amino acid and nucleic acid sequences coding for said antibody. More preferably said antibody comprises a modified hinge region and exhibits an improved antagonistic activity. More particularly, the antibody according to the invention is capable of inhibiting the c-Met dimerization. The invention likewise comprises the use of said antibody as a medicament for the prophylactic and/or therapeutic treatment of cancers, preferably for cancer characterized by a ligand-independent activation of c-Met, or any pathology connected with the over expression of said receptor as well as in processes or kits for diagnosis of illnesses connected with the over-expression of c-Met. The invention finally comprises products and/or compositions comprising such an antibody in combination with other antibodies and/or chemical compounds directed against other growth factors involved in tumor progression or metastasis and/or compounds and/or anti-cancer agents or agents conjugated with toxins and their use for the prevention and/or the treatment of certain cancers.



WO 2011/151412 A1

NOVEL ANTI-cMET ANTIBODY

The present invention relates to a novel divalent antibody capable of binding specifically to the human c-Met receptor and/or capable of specifically inhibiting the tyrosine kinase activity of said receptor, preferably both in a ligand-dependent and in a ligand-independent manner as well as the amino acid and nucleic acid sequences coding for said antibody. More preferably said antibody comprises a modified hinge region and exhibits an improved antagonistic activity. More particularly, the antibody according to the invention is capable of inhibiting the c-Met dimerization. The invention likewise comprises the use of said antibody as a medicament for the prophylactic and/or therapeutic treatment of cancers, preferably for cancer characterized by a ligand-independent activation of c-Met, or any pathology connected with the overexpression of said receptor as well as in processes or kits for diagnosis of illnesses connected with the over-expression of c-Met. The invention finally comprises products and/or compositions comprising such an antibody in combination with other antibodies and/or chemical compounds directed against other growth factors involved in tumor progression or metastasis and/or compounds and/or anti-cancer agents or agents conjugated with toxins and their use for the prevention and/or the treatment of certain cancers.

Receptor tyrosine kinase (RTK) targeted agents such as trastuzumab, cetuximab, bevacizumab, imatinib and gefitinib inhibitors have illustrated the interest of targeting this protein class for treatment of selected cancers.

c-Met, is the prototypic member of a sub-family of RTKs which also includes RON and SEA. The c-Met RTK family is structurally different from other RTK families and is the only known high-affinity receptor for hepatocyte growth factor (HGF), also called scatter factor (SF) [D.P. Bottaro et al., Science 1991, 251:802-804; L. Naldini et al., Eur. Mol. Biol. Org. J. 1991, 10:2867-2878]. c-Met and HGF are widely expressed in a variety of tissue and their expression is normally restricted to cells of epithelial and mesenchymal origin respectively [M.F. Di Renzo et al., Oncogene 1991, 6:1997-2003; E. Sonnenberg et al., J. Cell. Biol. 1993, 123:223-235]. They are both required for normal mammalian development and have been shown to be particularly important in cell migration, morphogenic differentiation, and organization of the three-dimensional

tubular structures as well as growth and angiogenesis [F. Baldt et al., Nature 1995, 376:768-771; C. Schmidt et al., Nature. 1995:373:699-702; Tsarfaty et al., Science 1994, 263:98-101]. While the controlled regulation of c-Met and HGF have been shown to be important in mammalian development, tissue maintenance and repair [Nagayama
5 T., Nagayama M., Kohara S., Kamiguchi H., Shibuya M., Katoh Y., Itoh J., Shinohara Y., Brain Res. 2004, 5;999(2):155-66; Tahara Y., Ido A., Yamamoto S., Miyata Y., Uto H., Hori T., Hayashi K., Tsubouchi H., J Pharmacol Exp Ther. 2003, 307(1):146-51], their dysregulation is implicated in the progression of cancers.

Aberrant signalling driven by inappropriate activation of c-Met is one of the
10 most frequent alteration observed in human cancers and plays a crucial role in tumorigenesis and metastasis [Birchmeier et al., Nat. Rev. Mol. Cell Biol. 2003, 4:915-925; L. Trusolino and Comoglio P. M., Nat Rev. Cancer. 2002, 2(4):289-300].

Inappropriate c-Met activation can arise by ligand-dependent and independent mechanisms, which include overexpression of c-Met, and/or paracrine or autocrine
15 activation, or through gain in function mutation [J.G. Christensen, Burrows J. and Salgia R., Cancer Letters. 2005, 226:1-26]. However an oligomerization of c-Met receptor, in presence or in absence of the ligand, is required to regulate the binding affinity and binding kinetics of the kinase toward ATP and tyrosine-containing peptide substrates [Hays JL, Watowich SJ, Biochemistry, 2004 Aug 17, 43:10570-8]. Activated
20 c-Met recruits signalling effectors to its multidocking site located in the cytoplasm domain, resulting in the activation of several key signalling pathways, including Ras-MAPK, PI3K, Src and Stat3 [Gao CF, Vande Woude GF, Cell Res. 2005, 15(1):49-51; Furge KA, Zhang YW, Vande Woude GF, Oncogene. 2000, 19(49):5582-9]. These pathways are essential for tumour cell proliferation, invasion and angiogenesis and for
25 evading apoptosis [Furge KA, Zhang YW, Vande Woude GF, Oncogene, 2000, 19(49):5582-9; Gu H., Neel BG, Trends Cell Biol. 2003 Mar, 13(3):122-30; Fan S., Ma YX, Wang JA, Yuan RQ, Meng Q., Cao Y., Laterra JJ, Goldberg ID, Rosen EM, Oncogene. 2000 Apr 27, 19(18):2212-23]. In addition, a unique facet of the c-Met signalling relative to other RTK is its reported interaction with focal adhesion
30 complexes and non kinase binding partners such as $\alpha 6\beta 4$ integrins [Trusolino L., Bertotti A., Comoglio PM, Cell. 2001, 107:643-54], CD44v6 [Van der Voort R., Taher TE, Wielenga VJ, Spaargaren M., Prevo R., Smit L., David G., Hartmann G., Gherardi

E., Pals ST, J. Biol. Chem. 1999, 274(10):6499-506], Plexin B1 or semaphorins [Giordano S., Corso S., Conrotto P., Artigiani S., Gilestro G., Barberis D., Tamagnone L., Comoglio PM, Nat Cell Biol. 2002, 4(9):720-4; Conrotto P., Valdembri D., Corso S., Serini G., Tamagnone L., Comoglio PM, Bussolino F., Giordano S., Blood. 2005, 105(11):4321-9; Conrotto P., Corso S., Gamberini S., Comoglio PM, Giordano S., Oncogene. 2004, 23:5131-7] which may further add to the complexity of regulation of cell function by this receptor. Finally recent data demonstrate that c-Met could be involved in tumor resistance to gefitinib or erlotinib suggesting that combination of compound targeting both EGFR and c-Met might be of significant interest [Engelman
10 JA et al., Science, 2007, 316:1039-43].

In the past few years, many different strategies have been developed to attenuate c-Met signalling in cancer cell lines. These strategies include i) neutralizing antibodies against c-Met or HGF/SF [Cao B., Su Y., Oskarsson M., Zhao P., Kort EJ, Fisher RJ, Wang LM, Vande Woude GF, Proc Natl Acad Sci U S A. 2001, 98(13):7443-8; Martens T., Schmidt NO, Eckerich C., Fillbrandt R., Merchant M., Schwall R., Westphal M., Lamszus K., Clin Cancer Res. 2006, 12(20):6144-52] or the use of HGF/SF antagonist NK4 to prevent ligand binding to c-Met [Kuba K., Matsumoto K., Date K., Shimura H., Tanaka M., Nakamura T., Cancer Res., 2000, 60:6737-43], ii) small ATP binding site inhibitors to c-Met that block kinase activity [Christensen JG, Schreck R., Burrows J., Kuruganti P., Chan E, Le P., Chen J., Wang X., Ruslim L., Blake R., Lipson KE, Ramphal J., Do S., Cui JJ, Cherrington JM, Mendel DB, Cancer Res. 2003, 63:7345-55], iii) engineered SH2 domain polypeptide that interferes with access to the multidocking site and RNAi or ribozyme that reduce receptor or ligand expression. Most of these approaches display a selective inhibition of c-Met resulting in tumor inhibition and showing that c-Met could be of interest for therapeutic intervention
25 in cancer.

Within the molecules generated for c-Met targeting, some are antibodies. The most extensively described is the anti-c-Met 5D5 antibody generated by Genentech [WO 96/38557] which behaves as a potent agonist when added alone in various models and as an antagonist when used as a Fab fragment. A monovalent engineered form of this antibody described as one armed 5D5 (OA5D5) and produced as a recombinant protein in *E. Coli* is also the subject of a patent application [WO 2006/015371] by
30

Genentech. However, this molecule that could not be considered as an antibody because of its particular scaffold, displays also mutations that could be immunogenic in humans. In terms of activity, this unglycosylated molecule is devoided of effector functions and finally, no clear data demonstrate that OA5D5 inhibits dimerization of c-Met.

5 Moreover, when tested in the G55 *in vivo* model, a glioblastoma cell line that expresses c-Met but not HGF mRNA and protein and that grows independently of the ligand, the one armed anti-c-Met had no significant effect on G55 tumor growth suggesting that OA5D5 acts primarily by blocking HGF binding and is not able to target tumors activated independently of HGF [Martens T. et al, Clin. Cancer Res., 2006,

10 12(20):6144-6152].

Another antibody targeting c-Met is described by Pfizer as an antibody acting “predominantly as c-Met antagonist, and in some instance as a c-Met agonist” [WO 2005/016382]. No data showing any effect of Pfizer antibodies on c-Met dimerization is described in this application.

15 One of the innovative aspects of the present invention is to generate a chimeric and/or humanized monoclonal antibody without intrinsic agonist activity and inhibiting c-Met dimerization. More particularly, an innovative aspect of the present invention is to generate a chimeric and/or humanized monoclonal antibody with antagonist activity and inhibiting c-Met dimerization.

20 In addition of targeting ligand-dependent tumors, this approach will also impair ligand-independent activations of c-Met due to its overexpression or mutations of the intra cellular domains which remained dependent to oligomerization for signalling. Another aspect of the activity of this antibody could be a steric hindrance for c-Met interaction with its partners that will result in impairment of c-Met functions. This

25 antibody is humanized and engineered preferentially, but not limited, as human IgG1 to get effector functions such as ADCC and CDC in addition to functions linked to the specific blockade of the c-Met receptor.

Surprisingly, for the first time, inventors have managed to generate a chimeric and/or humanized monoclonal antagonist antibody capable of binding to c-Met but also

30 capable of inhibiting the c-Met dimerization, said monoclonal antibody being divalent contrary to existing antagonist antibodies directed against c-Met. If it is true that, in the prior art, it is sometimes suggested that an antibody capable of inhibiting the

dimerization of c-Met with its partners could be an interesting one, it has never been disclosed, or clearly suggested, an antibody capable of doing so. Moreover, regarding antibody specificity, it was not evident at all to succeed in the generation of such an active divalent antibody.

5 As it was explained before, the inhibition of the c-Met dimerization is a capital aspect of the invention as such antibodies will present a real interest for a larger population of patients. Not only ligand-dependent activated c-Met cancer, as it was the case until the present invention, but also ligand-independent activated c-Met cancer could be treated with antibodies generated by the process of the present invention.

10 Antibodies were evaluated by BRET analysis on cells expressing both c-Met-RLuc/c-Met-YFP and selected on their ability to inhibit at least 40 %, preferably 45 %, 50 %, 55 % and most preferably 60 % of the BRET signal.

The BRET technology is known as being representative of the protein dimerization [Angers et al., PNAS, 2000, 97:3684-89].

15 The BRET technology is well known by the man skill in the art and will be detailed in the following examples. More particularly, BRET (Bioluminescence Resonance Energy Transfer) is a non-radiative energy transfer occurring between a bioluminescent donor (Renilla Luciferase (Rluc)) and a fluorescent acceptor, a mutant of GFP (Green Fluorescent Protein) or YFP (Yellow fluorescent protein). In the present
20 case EYFP (Enhanced Yellow Fluorescent Protein) was used. The efficacy of transfer depends on the orientation and the distance between the donor and the acceptor. Then, the energy transfer can occur only if the two molecules are in close proximity (1-10 nm). This property is used to generate protein-protein interaction assays. Indeed, in order to study the interaction between two partners, the first one is genetically fused to
25 the Renilla Luciferase and the second one to the yellow mutant of the GFP. Fusion proteins are generally, but not obligatory, expressed in mammalian cells. In presence of its membrane permeable substrate (coelenterazine), Rluc emits blue light. If the GFP mutant is closer than 10 nm from the Rluc, an energy transfer can occur and an additional yellow signal can be detected. The BRET signal is measured as the ratio
30 between the light emitted by the acceptor and the light emitted by the donor. So the BRET signal will increase as the two fusion proteins are brought into proximity or if a conformational change brings Rluc and GFP mutant closer.

If the BRET analysis consists in a preferred embodiment, any method known by the man skilled in the art can be used to measure c-Met dimerization. Without limitation, the following technologies can be mentioned: FRET (Fluorescence Resonance Energy Transfer), HTRF (Homogenous Time resolved Fluorescence), FLIM (Fluorescence Lifetime Imaging Microscopy) or SW-FCCS single wavelength fluorescence cross-correlation spectroscopy).

Other classical technologies could also be used, such as Co-immunoprecipitation, Alpha screen, Chemical cross-linking, Double-Hybrid, Affinity Chromatography, ELISA or Far western blot.

The terms “antibody”, “antibodies” or “immunoglobulin” are used interchangeably in the broadest sense and include monoclonal antibodies (e.g., full length or intact monoclonal antibodies), polyclonal antibodies, multivalent antibodies or multispecific antibodies (e.g., bispecific antibodies so long as they exhibit the desired biological activity).

More particularly, such molecule consists in a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (or domain) (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g. effector cells) and the first component (C1q) of the classical complement system.

The heavy chains of immunoglobulins can be divided into three functional regions: the Fd region, the hinge region, and the Fc region (fragment crystallizable). The

Fd region comprises the VH and CH1 domains and, in combination with the light chain, forms Fab - the antigen-binding fragment. The Fc fragment is responsible for the immunoglobulin effector functions, which includes, for example, complement fixation and binding to cognate Fc receptors of effector cells. The hinge region, found in IgG, 5 IgA, and IgD immunoglobulin classes, acts as a flexible spacer that allows the Fab portion to move freely in space relative to the Fc region. The hinge domains are structurally diverse, varying in both sequence and length among immunoglobulin classes and subclasses.

According to crystallographic studies, the immunoglobulin hinge region can be 10 further subdivided structurally and functionally into three regions: the upper hinge, the core, and the lower hinge (Shin et al., Immunological Reviews 130:87, 1992). The upper hinge includes amino acids from the carboxyl end of CH1 to the first residue in the hinge that restricts motion, generally the first cysteine residue that forms an interchain disulfide bond between the two heavy chains. The length of the upper hinge 15 region correlates with the segmental flexibility of the antibody. The core hinge region contains the inter-heavy chain disulfide bridges. The lower hinge region joins the amino terminal end of, and includes residues in the CH2 domain. The core hinge region of human IgG1 contains the sequence Cys-Pro-Pro-Cys that, when dimerized by disulfide bond formation, results in a cyclic octapeptide believed to act as a pivot, thus conferring 20 flexibility. Conformational changes permitted by the structure and flexibility of the immunoglobulin hinge region polypeptide sequence may affect the effector functions of the Fc portion of the antibody.

The term « Monoclonal Antibody » is used in accordance with its ordinary meaning to denote an antibody obtained from a population of substantially 25 homogeneous antibodies, i.e. the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts. In other words, a monoclonal antibody consists in a homogenous antibody resulting from the proliferation of a single clone of cells (e.g., hybridoma cells, eukaryotic host cells transfected with DNA encoding the homogenous antibody, 30 prokaryotic host cells transformed with DNA encoding the homogenous antibody, etc.), and which is generally characterized by heavy chains of a single class and subclass, and light chains of a single type. Monoclonal antibodies are highly specific, being directed

against a single antigen. Furthermore, in contrast to polyclonal antibodies preparations that typically include different antibodies directed against different determinants, or epitope, each monoclonal antibody is directed against a single determinant on the antigen.

5 In the present description, the terms polypeptides, polypeptide sequences, amino acid sequences, peptides and proteins attached to antibody compounds or to their sequence are interchangeable.

The invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization and comprising
10 a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3 or a sequence having at least 80 % identity after optimum alignment with sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7 or a sequence having at least 80% identity after optimum
15 alignment with sequences SEQ ID Nos. 5, 6 or 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 56.

In a preferred embodiment, the present invention is directed to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the
20 c-Met dimerization,

said antibody comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7, said antibody further comprising a hinge region comprising the
25 amino acid sequence SEQ ID No. 56;

for use for the prevention or the treatment of a patient in need thereof having a cancer characterized by ligand-independent activation of c-Met, preferably a cancer further characterized by overexpression of c-Met, said c-Met overexpression resulting more preferably from genic amplification of c-Met, and, also more preferably, resulting in
30 ligand-independent activation of c-Met.

In this aspect, the present invention comprises a method for the prevention or the treatment of a patient in need thereof having a cancer characterized by ligand-

independent activation of c-Met, preferably a cancer further characterized by overexpression of c-Met, said c-Met overexpression resulting more preferably from genic amplification of c-Met, and, also more preferably, resulting in ligand-independent activation of c-Met, said method comprising the step of administering a composition
5 comprising a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization, said antibody comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7, said
10 antibody further comprising a hinge region comprising the amino acid sequence SEQ ID No. 56.

In this particular aspect and in a more preferred embodiment, said cancer characterized by ligand-independent activation of c-Met, and preferably further characterized by overexpression of c-Met, resulting more preferably from genic
15 amplification of c-Met, and, also more preferably, resulting in ligand-independent activation of c-Met, is selected from the group consisting of renal cell carcinoma and gastric cancer.

More particularly, the invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, as above described characterized in that said
20 hinge region comprises the amino acid sequence SEQ ID No. 57.

In other words, the invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization and comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3 or a sequence having at
25 least 80% identity after optimum alignment with sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 5, 6 or 7, said antibody being further characterized in that it also comprises a hinge region comprising the
30 amino acid sequence SEQ ID No. 57.

More particularly, the invention relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, as above described characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 21.

In other words, the invention also relates to a monoclonal antibody, or a divalent functional fragment or derivative thereof, capable to inhibit the c-Met dimerization and comprising a heavy chain comprising CDR-H1, CDR-H2 and CDR-H3 with respectively the amino acid sequences SEQ ID Nos. 1, 2 and 3 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 1, 2 and 3; and a light chain comprising CDR-L1, CDR-L2 and CDR-L3 with respectively the amino acid sequences SEQ ID Nos. 5, 6 and 7 or a sequence having at least 80% identity after optimum alignment with sequences SEQ ID Nos. 5, 6 or 7, said antibody being further characterized in that it also comprises a hinge region comprising the amino acid sequence SEQ ID No. 21.

As it will be apparent for the man skilled in the art, the consensus sequences SEQ ID Nos. 57 and 21 are comprised in the consensus sequence SEQ ID No. 56.

Table 1

	#01	#02	#03	#04	#05	#06	#07	#08	#09	#10	#11	#12	#13	#14
SEQ ID NO 56	x1	x2	x3	C	x5	x6	x7	x8	x9	C	x11	x12	C	x14
SEQ ID NO 57	x1	x2	x3	C	x5	x6	x7	x8	x9	C	P	P	C	P
SEQ ID NO 21	x1	x2	x3	C	x5	-	C	x8	x9	C	x11	x12	C	x14

For SEQ ID No. 56:

20 X1: P, R, C, - X5: D, C, G, - X8: H, V, K, - X12: P, -
 X2: K, C, R, - X6: K, C, - X9: T, C, E, P, - X14: P, T
 X3: S, C, D, - X7: T, C, - X11: P, I

The expression “functional fragments and derivatives” will be defined in details later in the present specification.

By CDR regions or CDR(s), it is intended to indicate the hypervariable regions of the heavy and light chains of the immunoglobulins as defined by IMGT.

The IMGT unique numbering has been defined to compare the variable domains whatever the antigen receptor, the chain type, or the species [Lefranc M.-P.,

Immunology Today 18, 509 (1997); Lefranc M.-P., The Immunologist, 7, 132-136 (1999); Lefranc, M.-P., Pommié, C., Ruiz, M., Giudicelli, V., Foulquier, E., Truong, L., Thouvenin-Contet, V. and Lefranc, Dev. Comp. Immunol., 27, 55-77 (2003)]. In the IMGT unique numbering, the conserved amino acids always have the same position, for instance cysteine 23 (1st-CYS), tryptophan 41 (CONSERVED-TRP), hydrophobic amino acid 89, cysteine 104 (2nd-CYS), phenylalanine or tryptophan 118 (J-PHE or J-TRP). The IMGT unique numbering provides a standardized delimitation of the framework regions (FR1-IMGT: positions 1 to 26, FR2-IMGT: 39 to 55, FR3-IMGT: 66 to 104 and FR4-IMGT: 118 to 128) and of the complementarity determining regions: CDR1-IMGT: 27 to 38, CDR2-IMGT: 56 to 65 and CDR3-IMGT: 105 to 117. As gaps represent unoccupied positions, the CDR-IMGT lengths (shown between brackets and separated by dots, e.g. [8.8.13]) become crucial information. The IMGT unique numbering is used in 2D graphical representations, designated as IMGT Colliers de Perles [Ruiz, M. and Lefranc, M.-P., Immunogenetics, 53, 857-883 (2002); Kaas, Q. and Lefranc, M.-P., Current Bioinformatics, 2, 21-30 (2007)], and in 3D structures in IMGT/3Dstructure-DB [Kaas, Q., Ruiz, M. and Lefranc, M.-P., T cell receptor and MHC structural data. Nucl. Acids. Res., 32, D208-D210 (2004)].

Three heavy chain CDRs and 3 light chain CDRs exist. The term CDR or CDRs is used here in order to indicate, according to the case, one of these regions or several, or even the whole, of these regions which contain the majority of the amino acid residues responsible for the binding by affinity of the antibody for the antigen or the epitope which it recognizes.

By “percentage of identity” between two nucleic acid or amino acid sequences in the sense of the present invention, it is intended to indicate a percentage of nucleotides or of identical amino acid residues between the two sequences to be compared, obtained after the best alignment (optimum alignment), this percentage being purely statistical and the differences between the two sequences being distributed randomly and over their entire length. The comparisons of sequences between two nucleic acid or amino acid sequences are traditionally carried out by comparing these sequences after having aligned them in an optimum manner, said comparison being able to be carried out by segment or by “comparison window”. The optimum alignment of the sequences for the comparison can be carried out, in addition to manually, by means

of the local homology algorithm of Smith and Waterman (1981) [Ad. App. Math. 2:482], by means of the local homology algorithm of Neddleman and Wunsch (1970) [J. Mol. Biol. 48: 443], by means of the similarity search method of Pearson and Lipman (1988) [Proc. Natl. Acad. Sci. USA 85:2444], by means of computer software using
5 these algorithms (GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI, or else by BLAST N or BLAST P comparison software).

The percentage of identity between two nucleic acid or amino acid sequences is determined by comparing these two sequences aligned in an optimum manner and in
10 which the nucleic acid or amino acid sequence to be compared can comprise additions or deletions with respect to the reference sequence for an optimum alignment between these two sequences. The percentage of identity is calculated by determining the number of identical positions for which the nucleotide or the amino acid residue is identical between the two sequences, by dividing this number of identical positions by
15 the total number of positions in the comparison window and by multiplying the result obtained by 100 in order to obtain the percentage of identity between these two sequences.

For example, it is possible to use the BLAST program, "BLAST 2 sequences" (Tatusova et al., "Blast 2 sequences - a new tool for comparing protein and nucleotide
20 sequences", FEMS Microbiol Lett. 174:247-250) available on the site <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>, the parameters used being those given by default (in particular for the parameters "open gap penalty": 5, and "extension gap penalty": 2; the matrix chosen being, for example, the matrix "BLOSUM 62" proposed by the program), the percentage of identity between the two sequences to be compared
25 being calculated directly by the program.

By amino acid sequence having at least 80 %, preferably 85 %, 90 %, 95 % and 98 % identity with a reference amino acid sequence, those having, with respect to the reference sequence, certain modifications, in particular a deletion, addition or substitution of at least one amino acid, a truncation or an elongation are preferred. In the
30 case of a substitution of one or more consecutive or nonconsecutive amino acid(s), the substitutions are preferred in which the substituted amino acids are replaced by "equivalent" amino acids. The expression "equivalent amino acids" is aimed here at

indicating any amino acid capable of being substituted with one of the amino acids of the base structure without, however, essentially modifying the biological activities of the corresponding antibodies and such as will be defined later, especially in the examples. These equivalent amino acids can be determined either by relying on their structural homology with the amino acids which they replace, or on results of comparative trials of biological activity between the different antibodies capable of being carried out.

By way of example, mention is made of the possibilities of substitution capable of being carried out without resulting in a profound modification of the biological activity of the corresponding modified antibody.

As non limitative example, the following table 2 is giving substitution possibilities conceivable with a conservation of the biological activity of the modified antibody. The reverse substitutions are also, of course, possible in the same conditions.

Table 2

Original residue	Substitution(s)
Ala (A)	Val, Gly, Pro
Arg (R)	Lys, His
Asn (N)	Gln
Asp (D)	Glu
Cys (C)	Ser
Gln (Q)	Asn
Glu (G)	Asp
Gly (G)	Ala
His (H)	Arg
Ile (I)	Leu
Leu (L)	Ile, Val, Met
Lys (K)	Arg
Met (M)	Leu
Phe (F)	Tyr
Pro (P)	Ala
Ser (S)	Thr, Cys
Thr (T)	Ser
Trp (W)	Tyr
Tyr (Y)	Phe, Trp
Val (V)	Leu, Ala

It must be understood here that the invention does not relate to the antibodies in natural form, that is to say they are not in their natural environment but that they have been able to be isolated or obtained by purification from natural sources, or else obtained by genetic recombination, or by chemical synthesis, and that they can then
5 contain unnatural amino acids as will be described further on.

It must also be understood, as previously mentioned, that the invention concerns more particularly a chimeric and/or a humanized divalent antibody, or any divalent functional fragment or derivative, with an antagonistic activity. Divalent antibodies of the prior art are agonists or partial agonists. The monoclonal antibody of the invention,
10 including a modified hinge as previously described, i.e. including a hinge region comprising the amino acid sequence SEQ ID No. 56, 57 or 21, is novel and presents the particularity to have a improved antagonistic activity compared to the chimeric or humanized antibody 224G11 without such a modified hinge as it will appear from the following examples.

15 Contrary to the prior art, inventors have obtained an improved antagonistic activity without modifying the format of the antibody. Actually, in the closest prior art represented by the antibody 5D5, it has been necessary to develop a monovalent fragment of the antibody to generate an antagonistic activity. In the present application, by the use of the hinge of the invention, it is possible for the first time to obtain a full
20 divalent antibody with increased antagonistic activity, and this contrary to the general knowledge.

In a preferred embodiment, the antibody of the invention comprises a hinge region comprising an amino acid sequence selected from the group consisting of SEQ ID Nos. 22 to 28 and 58 to 72, or a sequence having at least 80% identity after optimum
25 alignment with sequences SEQ ID Nos. 22 to 28 and 58 to 72.

For more clarity, the following tables 3 and 4 regroup the amino acids and nucleotides sequences of the different preferred hinges of the invention.

Table 3

SEQ ID No.	Amino acids	SEQ ID No.	Nucleotides
22	RKCCVECPPCP	29	AGGAAGTGCTGTGTGGAGTGCCCCCCTGCCCA
23	PRDCGCKPCICT	30	CCCCGGGACTGTGGGTGCAAGCCTTGCATTTGTACC
24	PKSCGCKPCICT	31	CCCAAGAGCTGTGGGTGCAAGCCTTGCATTTGTACC
25	PKSCGCKPCICP	32	CCAAAGAGCTGCGGCTGCAAGCCTTGTATCTGTCCC
26	PRDCGCKPCPPCP	33	CCACGGGACTGTGGCTGCAAGCCCTGCCCTCCGTGTCCA
27	PRDCGCHTCPPCP	34	CCCAGAGACTGTGGGTGTACACCTGCCCTCCTTGTCTT
28	PKSCDCHCPCP	35	CCCAAAGCTGCGATTGCCACTGTCCTCCATGTCCA

Table 4

SEQ ID No.	Amino acids	SEQ ID No.	Nucleotides
58	CKSCDKTHTCPPCP	73	TGCAAGAGCTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
59	PCSCDKTHTCPPCP	74	CCCTGCAGCTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
60	PKCCDKTHTCPPCP	75	CCCAAGTGCTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
61	PKSCCKTHTCPPCP	76	CCTAAGAGCTGTTGCAAGACCCACACCTGTCCCCCCTGCCCT
62	PKSCDCTHTCPPCP	77	CCCAAGAGCTGCGACTGCACCCACACCTGTCCCCCCTGCCCT
63	PKSCDKCHTCPPCP	78	CCCAAGAGCTGCGACAAGTGCCACACCTGTCCCCCCTGCCCT
64	PKSCDKTHCCPPCP	79	CCCAAGAGCTGCGACAAGACCCACTGCTGTCCCCCCTGCCCT
65	KCDKTHTCPPCP	80	AAGTGCGACAAGACCCACACCTGTCCCCCCTGCCCT
66	PKSCDCHTCPPCP	81	CCCAAGAGCTGCGACTGCCACACCTGTCCCCCCTGCCCT
67	PKSCDCTHCPPCP	82	CCCAAGAGCTGCGACTGCACCCACTGCCCCCCCTGCCCT
68	PCSCKHTCPPCP	83	CCCTGCAGCTGCAAGCACACCTGTCCCCCCTGCCCT
69	PSCCTHTCPPCP	84	CCTAGCTGCTGCACCCACACCTGTCCCCCCTGCCCT
70	PSCDKHCCPPCP	85	CCCAGCTGCGACAAGCACTGCTGCCCCCCTGCCCT
71	PKSCTCPPCP	86	CCCAAGAGCTGCACCTGTCCCCCTTGTCTT
72	PKSCDKCVECPCP	87	CCCAAGAGCTGCGATAAGTGCGTGGAGTGCCCCCCTTGTCTT

According a first approach, the antibody will be defined by its heavy chain sequence. More particularly, the antibody of the invention, or one of its functional fragments or derivatives, is characterized in that it comprises a heavy chain comprising at least one CDR chosen from CDRs comprising the amino acid sequences SEQ ID Nos. 1 to 3.

The mentioned sequences are the following ones:

SEQ ID No. 1: GYIFTAYT
 SEQ ID No. 2: IKPNNGLA
 SEQ ID No. 3: ARSEITTEFDY

According to a preferred aspect, the antibody of the invention, or one of its functional fragments or derivatives, comprises a heavy chain comprising at least one, preferably two, and most preferably three, CDR(s) chosen from CDR-H1, CDR-H2 and CDR-H3, wherein:

- 5 - CDR-H1 comprises the amino acid sequence SEQ ID No. 1,
- CDR-H2 comprises the amino acid sequence SEQ ID No. 2,
- CDR-H3 comprises the amino acid sequence SEQ ID No. 3.

In a second approach, the antibody will be now defined by its light chain sequence. More particularly, according to a second particular aspect of the invention,
10 the antibody, or one of its functional fragments or derivatives, is characterized in that it comprises a light chain comprising at least one CDR chosen from CDRs comprising the amino acid sequence SEQ ID Nos. 5 to 7.

The mentioned sequences are the following ones:

- SEQ ID No. 5: ESVDSYANSF
- 15 SEQ ID No. 6: RAS
- SEQ ID No. 7: QQSKEDPLT

According to another preferred aspect, the antibody of the invention, or one of its functional fragments or derivatives, comprises a light chain comprising at least one,
20 preferably two, and most preferably three, CDR(s) chosen from CDR-L1, CDR-L2 and CDR-L3, wherein:

- CDR-L1 comprises the amino acid sequence SEQ ID No. 5,
- CDR-L2 comprises the amino acid sequence SEQ ID No. 6,
- CDR-L3 comprises the amino acid sequence SEQ ID No. 7.

25 The murine hybridoma capable of secreting monoclonal antibodies according to the present invention, especially hybridoma of murine origin, was deposited at the CNCM (Institut Pasteur, Paris, France) on 03/14/2007 under the number CNCM I-3731.

In the present application, IgG1 are preferred to get effector functions, and most preferably ADCC and CDC.

30 The skilled artisan will recognize that effector functions include, for example, C1q binding; complement dependent cytotoxicity (CDC); Fc receptor binding;

antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; and down regulation of cell surface receptors (e.g. B cell receptor; BCR).

The antibodies according to the present invention are preferably specific monoclonal antibodies, especially of murine, chimeric or humanized origin, which can
5 be obtained according to the standard methods well known to the person skilled in the art.

In general, for the preparation of monoclonal antibodies or their functional fragments or derivatives, especially of murine origin, it is possible to refer to techniques which are described in particular in the manual "Antibodies" (Harlow and Lane,
10 Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor NY, pp. 726, 1988) or to the technique of preparation from hybridomas described by Kohler and Milstein (Nature, 256:495-497, 1975).

The monoclonal antibodies according to the invention can be obtained, for example, from an animal cell immunized against the c-Met, or one of its fragments
15 containing the epitope specifically recognized by said monoclonal antibodies according to the invention. Said c-Met, or one of its said fragments, can especially be produced according to the usual working methods, by genetic recombination starting with a nucleic acid sequence contained in the cDNA sequence coding for the c-Met or by peptide synthesis starting from a sequence of amino acids comprised in the peptide
20 sequence of the c-Met.

The monoclonal antibodies according to the invention can, for example, be purified on an affinity column on which the c-Met or one of its fragments containing the epitope specifically recognized by said monoclonal antibodies according to the invention has previously been immobilized. More particularly, said monoclonal
25 antibodies can be purified by chromatography on protein A and/or G, followed or not followed by ion-exchange chromatography aimed at eliminating the residual protein contaminants as well as the DNA and the LPS, in itself followed or not followed by exclusion chromatography on Sepharose™ gel in order to eliminate the potential aggregates due to the presence of dimers or of other multimers. In an even more
30 preferred manner, the whole of these techniques can be used simultaneously or successively.

The antibody of the invention, or a divalent functional fragment or derivative thereof, consists preferably of a chimeric antibody.

By chimeric antibody, it is intended to indicate an antibody which contains a natural variable (light chain and heavy chain) region derived from an antibody of a given species in combination with the light chain and heavy chain constant regions of an antibody of a species heterologous to said given species (e.g. mouse, horse, rabbit, dog, cow, chicken, etc.).

The antibodies or their fragments of chimeric type according to the invention can be prepared by using the techniques of genetic recombination. For example, the chimeric antibody can be produced by cloning a recombinant DNA containing a promoter and a sequence coding for the variable region of a non-human, especially murine, monoclonal antibody according to the invention and a sequence coding for the constant region of human antibody. A chimeric antibody of the invention encoded by such a recombinant gene will be, for example, a mouse-man chimera, the specificity of this antibody being determined by the variable region derived from the murine DNA and its isotype determined by the constant region derived from the human DNA. For the methods of preparation of chimeric antibodies, it is possible, for example, to refer to the documents Verhoeven et al. (BioEssays, 8:74, 1988), Morrison et al. (Proc. Natl. Acad. Sci. USA 82:6851-6855, 1984) or US 4,816,567.

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a chimeric heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 46 or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 46.

SEQ ID No. 46: EVQLQQSGPELVKPGASVKISCKTSGYIFTAYTMHWVRQSLGE
SLDWIGGIKPNNGLANYNQKFKGKATLTVDKSSSTAYMDLRSLTSEDSAVYYC
ARSEITTEFDYWGQGTALTSS

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a chimeric light chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 47 or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 47.

SEQ ID No. 47: DIVLTQSPASLAVSLGQRATISCRASESVDSYANSFMHWYQQKP
 GQPPKLLIYRASNL ESGIPARFSGSGSRTDFTLTINPVEADDVATYYCQQSKEDP
 LTFGSGTKLEMKR

5 More particularly, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG2chim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 22.

10 In the present application, the use of square brackets is not necessary and, as an example, the reference [224G11] [IgG2chim] must be considered as identical to 224G11IgG2chim. In a same way, to indicate that the antibody is a murine one, the expression murine or the letter m can be added; to indicate that the antibody is a chimeric one, the expression chim or the letter c can be added and ; to indicate that the
 15 antibody is a humanized one, the expression hum, hz, Hz or the letter h can be added. As an example, the chimeric antibody 224G11IgG2 can be referred as c224G11IgG2, c224G11[IgG2], c[224G11]IgG2, c[224G11][IgG2], 224G11IgG2chim, 224G11[IgG2chim], [224G11]IgG2chim or [224G11][IgG2chim].

In another aspect, a preferred antibody, or a divalent functional fragment or
 20 derivative thereof, according to the invention and named [224G11] [TH7chim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

In the present application, the reference TH7 must be considered as identical to
 25 C7 Δ 6-9 or TH7C7 Δ 6-9. The symbol Δ means deletion.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [MHchim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID
 30 No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 23.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [MUP9Hchim],

comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 26.

5 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [MMCHchim], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 24.

10 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C1], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 58.

15 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C2], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 59.

20 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C3], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 60.

25 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C5], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 61.

30 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C6], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a

light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 62.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C7], comprises a
5 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 63.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C9], comprises a
10 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 64.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [Δ 1-3], comprises a
15 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 65.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C7 Δ 6], comprises a
20 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 66.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C6 Δ 9], comprises a
25 heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 67.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C2 Δ 5-7], comprises
30 a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 68.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C5Δ2-6], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a
5 hinge region comprising the amino acid sequence SEQ ID No. 69.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [C9Δ2-7], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID No. 47, and a
10 hinge region comprising the amino acid sequence SEQ ID No. 70.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [Δ5-6-7-8], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID
15 No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 71.

In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG1/IgG2], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 46, a light chain variable domain comprising the amino acid sequence SEQ ID
20 No. 47, and a hinge region comprising the amino acid sequence SEQ ID No. 72.

The antibody of the invention, or a divalent functional fragment or derivative thereof, consists preferably of a human antibody.

The term “human antibody” includes all antibodies that have one or more variable and constant region derived from human immunoglobulin sequences. In a
25 preferred embodiment, all of the variable and constant domains (or regions) are derived from human immunoglobulin sequence (fully human antibody). In other words, it includes any antibody which have variable and constant regions (if present) derived from human germline immunoglobulin sequences, i.e. which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has
30 been made using any techniques for making human antibodies known by the man skill in the art.

In one embodiment, the human monoclonal antibodies are produced by a hybridoma which includes a B cell obtained from a transgenic non-human animal, e.g., a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

5 As example for such transgenic mouse, it can be mentioned the XENOMOUSETM which is an engineered mouse strain that comprises large fragments of the human immunoglobulin loci and is deficient in mouse antibody production (Green et al., 1994, Nature Genetics, 7:13-21). The XENOMOUSETM produces an adult-like human repertoire of fully human antibodies, and generate antigen-specific
10 human monoclonal antibodies. A second generation XENOMOUSETM contains approximately 80% of the human antibody repertoire (Green & Jakobovits, 1998, J. Exp. Med., 188:483-495).

Any other technique known by the man skill in the art, such as phage display technique, can also be used for the generation of human antibody according to the
15 invention.

The antibody of the invention, or a divalent functional fragment or derivative thereof, consists preferably of a humanized antibody.

By the expression “humanized antibody”, it is intended to indicate an antibody which contains CDR regions derived from an antibody of nonhuman origin, the other
20 parts of the antibody molecule being derived from one (or from several) human antibodies. Moreover, some of the residues of the segments of the skeleton (called FR) can be modified in order to conserve the affinity of the binding (Jones et al., Nature, 321:522-525, 1986; Verhoeven et al., Science, 239:1534-1536, 1988; Riechmann et al., Nature, 332:323-327, 1988).

25 The humanized antibodies according to the invention or their fragments can be prepared by techniques known to the person skilled in the art (such as, for example, those described in the documents Singer et al., J. Immun. 150:2844-2857, 1992; Mountain et al., Biotechnol. Genet. Eng. Rev., 10: 1-142, 1992; or Bebbington et al., Bio/Technology, 10:169-175, 1992).

30 Other humanization method are known by the man skill in the art as, for example, the “CDR Grafting” method described by Protein Design Lab (PDL) in the patent applications EP 0 451261, EP 0 682 040, EP 0 9127, EP 0 566 647 or US

5,530,101, US 6,180,370, US 5,585,089 and US 5,693,761. The following patent applications can also be mentioned: US 5,639,641; US 6,054,297; US 5,886,152 and US 5,877,293.

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a humanized heavy chain variable domain of sequence comprising the amino acid sequence SEQ ID No. 4 or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 4.

SEQ ID No. 4: QVQLVQSGAEVKKPGASVKVSCKASGYIFTAYTMHWVRQAPG
QGLEWMGWIKPNNGLANYAQKFQGRVTMTRDTSISTAYMELSRRLRSDDTAVY
10 YCARSEITTEFDYWGQGTLVTVSS

More particularly, said antibody, or a functional fragment or derivative thereof, comprises a humanized light chain variable domain selected from the group of sequences comprising the amino acid sequence SEQ ID No. 8, 9 or 10 or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 8, 9 or 10.

SEQ ID No. 8: DIVLTQSPDSLAVSLGERATINCKSSESVD SYANSFMHWYQQKP
GQPPKLLIYRASTRESGVPDRFSGSGSRTDFTLTISLQAEDVAVYYCQQSKEDP
LTFGGGTKVEIKR

SEQ ID No. 9: DIVMTQSPDSLAVSLGERATINCKSSESVD SYANSFMHWYQQKP
20 GQPPKLLIYRASTRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQSKEDP
LTFGGGTKVEIKR

SEQ ID No. 10: DIVMTQSPDSLAVSLGERATINCKSSESVD SYANSFLHWYQQKP
GQPPKLLIYRASTRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCQQSKEDP
LTFGGGTKVEIKR

25

More particularly, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG2Hz1], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 8, and a hinge region comprising the amino acid sequence SEQ ID No. 22.

30 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG2Hz2],

comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 9, and a hinge region comprising the amino acid sequence SEQ ID No. 22.

5 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [IgG2Hz3], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 10, and a hinge region comprising the amino acid sequence SEQ ID No. 22.

10 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [TH7Hz1], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 8, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

15 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [TH7z2], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 9, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

20 In another aspect, a preferred antibody, or a divalent functional fragment or derivative thereof, according to the invention and named [224G11] [TH7Hz3], comprises a heavy chain variable domain comprising the amino acid sequence SEQ ID No. 4, a light chain variable domain comprising the amino acid sequence SEQ ID No. 10, and a hinge region comprising the amino acid sequence SEQ ID No. 28.

25 In another aspect, antibodies of the invention can be described by their total heavy and light chains, respectively.

As example, the antibody [224G11] [IgG2chim] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 50, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 50, and a complete light chain comprising the amino acid sequence SEQ ID
30 No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [TH7chim] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 51, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 51, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C1] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 88, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 88, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 89, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 89, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 90, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 90, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C5] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 91, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 91, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C6] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 92, or a

sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 92, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

5 As another example, the antibody [224G11] [C7] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 93, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 93, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the
10 sequence SEQ ID No. 52.

 As another example, the antibody [224G11] [C9] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 94, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 94, and a complete light chain comprising the amino acid sequence SEQ ID No.
15 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

 As another example, the antibody [224G11] [Δ 1-3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 95, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ
20 ID No. 95, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

 As another example, the antibody [224G11] [C7 Δ 6] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 96, or a
25 sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 96, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

 As another example, the antibody [224G11] [C6 Δ 9] of the invention comprises a
30 complete heavy chain comprising the amino acid sequence SEQ ID No. 97, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 97, and a complete light chain comprising the amino acid sequence SEQ ID No.

52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C2Δ5-7] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 98, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 98, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C5Δ2-6] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 99, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 99, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [C9Δ2-7] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 100, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 100, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [Δ5-6-7-8] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 101, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 101, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80% identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [IgG1/IgG2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 102, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 102, and a complete light chain comprising the amino acid sequence SEQ ID No. 52, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 52.

As another example, the antibody [224G11] [IgG2Hz1] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 36, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 36 and a complete light chain comprising the amino acid sequence SEQ ID
5 No. 38, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 38.

As another example, the antibody [224G11] [IgG2Hz2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 36, or a sequence having at least 80 % identity after optimum alignment with the sequence
10 SEQ ID No. 36 and a complete light chain comprising the amino acid sequence SEQ ID No. 39, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 39.

As another example, the antibody [224G11] [IgG2Hz3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 36, or a sequence having at least 80 % identity after optimum alignment with the sequence
15 SEQ ID No. 36 and a complete light chain comprising the amino acid sequence SEQ ID No. 40, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 40.

As another example, the antibody [224G11] [TH7Hz1] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 37, or a sequence having at least 80 % identity after optimum alignment with the sequence
20 SEQ ID No. 37 and a complete light chain comprising the amino acid sequence SEQ ID No. 38, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 38.

As another example, the antibody [224G11] [TH7Hz2] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 37, or a sequence having at least 80 % identity after optimum alignment with the sequence
25 SEQ ID No. 37 and a complete light chain comprising the amino acid sequence SEQ ID No. 39, or a sequence having at least 80 % identity after optimum alignment with the
30 sequence SEQ ID No. 39.

As another example, the antibody [224G11] [TH7Hz3] of the invention comprises a complete heavy chain comprising the amino acid sequence SEQ ID No. 37,

or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 37 and a complete light chain comprising the amino acid sequence SEQ ID No. 40, or a sequence having at least 80 % identity after optimum alignment with the sequence SEQ ID No. 40.

5 Other examples of antibodies, or derivatives thereof, according to the invention comprises complete heavy chains comprising an amino acid sequence selected in the group consisting of SEQ ID Nos. 88 to 102 (corresponding nucleotide sequences are SEQ ID Nos. 103 to 117).

By “functional fragment” of an antibody according to the invention, it is
10 intended to indicate in particular an antibody fragment, such as Fv, scFv (sc for single chain), Fab, F(ab')₂, Fab', scFv-Fc fragments or diabodies, or any fragment of which the half-life time would have been increased by chemical modification, such as the addition of poly(alkylene) glycol such as poly(ethylene) glycol (“PEGylation”) (pegylated fragments called Fv-PEG, scFv-PEG, Fab-PEG, F(ab')₂-PEG or Fab'-PEG)
15 (“PEG” for Poly(Ethylene) Glycol), or by incorporation in a liposome, said fragments having at least one of the characteristic CDRs of sequence SEQ ID Nos. 1 to 3 and 5 to 7 according to the invention, and, especially, in that it is capable of exerting in a general manner an even partial activity of the antibody from which it is descended, such as in particular the capacity to recognize and to bind to the c-Met, and, if necessary, to inhibit
20 the activity of the c-Met.

Preferably, said functional fragments will be constituted or will comprise a partial sequence of the heavy or light variable chain of the antibody from which they are derived, said partial sequence being sufficient to retain the same specificity of binding as the antibody from which it is descended and a sufficient affinity, preferably at least
25 equal to 1/100, in a more preferred manner to at least 1/10, of that of the antibody from which it is descended, with respect to the c-Met. Such a functional fragment will contain at the minimum 5 amino acids, preferably 6, 7, 8, 9, 10, 12, 15, 25, 50 and 100 consecutive amino acids of the sequence of the antibody from which it is descended.

Preferably, these functional fragments will be fragments of Fv, scFv, Fab,
30 F(ab')₂, F(ab'), scFv-Fc type or diabodies, which generally have the same specificity of binding as the antibody from which they are descended. In a more preferred embodiment of the invention, these fragments are selected among divalent fragments

such as F(ab')₂ fragments. According to the present invention, antibody fragments of the invention can be obtained starting from antibodies such as described above by methods such as digestion by enzymes, such as pepsin or papain and/or by cleavage of the disulfide bridges by chemical reduction. In another manner, the antibody fragments
5 comprised in the present invention can be obtained by techniques of genetic recombination likewise well known to the person skilled in the art or else by peptide synthesis by means of, for example, automatic peptide synthesizers such as those supplied by the company Applied Biosystems, etc.

By "divalent fragment", it must be understood any antibody fragments
10 comprising two arms and, more particularly, F(ab')₂ fragments.

By "derivatives" of an antibody according to the invention, it is meant a binding protein comprising a protein scaffold and at least one of the CDRs selected from the original antibody in order to maintain the binding capacity. Such compounds are well known by the man skilled in the art and will be described in more details in the
15 following specification.

More particularly, the antibody, or one of its functional fragments or derivatives, according to the invention is characterized in that said derivative consists in a binding protein comprising a scaffold on which at least one CDR has been grafted for the conservation of the original antibody paratopic recognizing properties.

20 One or several sequences through the 6 CDR sequences described in the invention can be presented on a protein scaffold. In this case, the protein scaffold reproduces the protein backbone with appropriate folding of the grafted CDR(s), thus allowing it (or them) to maintain their antigen paratopic recognizing properties.

The man skilled in the art knows how to select the protein scaffold on which at
25 least one CDR selected from the original antibody could be grafted. More particularly, it is known that, to be selected, such scaffold should display several features as follow (Skerra A., J. Mol. Recogn., 13, 2000, 167-187):

- phylogenetically good conservation,
- robust architecture with a well known three-dimensional molecular
30 organization (such as, for example, crystallography or NMR),
- small size,
- no or only low degree of post-translational modifications,

- easy to produce, express and purify.

Such protein scaffold can be, but without limitation, structure selected from the group consisting in fibronectin and preferentially the tenth fibronectin type III domain (FNfn10), lipocalin, anticalin (Skerra A., J. Biotechnol., 2001, 74(4):257-75), the
5 protein Z derivative from the domain B of staphylococcal protein A, thioredoxin A or any protein with repeated domain such as “ankyrin repeat” (Kohl et al., PNAS, 2003, vol. 100, No.4, 1700-1705), “armadillo repeat”, “leucin-rich repeat” or “tetratricopeptide repeat”.

It could also be mentioned scaffold derivative from toxins (such as, for example,
10 scorpion, insect, plant or mollusc toxins) or protein inhibitors of neuronal nitric oxide synthase (PIN).

As non limitative example of such hybrid constructions, it can be mentioned the insertion of the CDR-H1 (heavy chain) of an anti-CD4 antibody, i.e. the 13B8.2 antibody, into one of the exposed loop of the PIN. The binding properties of the
15 obtained binding protein remain similar to the original antibody (Bes et al., BBRC 343, 2006, 334-344). It can also be mentioned the grafting of the CDR-H3 (heavy chain) of an anti-lysozyme VHH antibody on a loop of neocarzinostatine (Nicaise et al., 2004).

As above mentioned, such protein scaffold can comprise from 1 to 6 CDR(s) from the original antibody. In a preferred embodiment, but without any limitation, the
20 man skilled in the art would select at least a CDR from the heavy chain, said heavy chain being known to be particularly implicated in the antibody specificity. The selection of the CDR(s) of interest will be evident for the man of the art with known method (BES et al., FEBS letters 508, 2001, 67-74).

As an evidence, these examples are not limitative and any other scaffold known
25 or described must be included in the present specification.

According to a novel aspect, the present invention relates to an isolated nucleic acid, characterized in that it is chosen from the following nucleic acids:

- a) a nucleic acid, DNA or RNA, coding for an antibody, or one of its functional fragments or derivatives, according to the invention;
- 30 b) a nucleic acid sequence comprising the sequences SEQ ID No. 11, SEQ ID No. 12, SEQ ID No. 13 and the sequences SEQ ID No. 15, SEQ ID No. 16 and SEQ ID No. 17;

c) a nucleic acid sequence comprising the sequences SEQ ID No. 14 and SEQ ID No. 18, 19 or 20;

d) the corresponding RNA nucleic acids of the nucleic acids as defined in b) or c);

5 e) the complementary nucleic acids of the nucleic acids as defined in a), b) and c); and

f) a nucleic acid of at least 18 nucleotides capable of hybridizing under conditions of high stringency with at least one of the CDRs of sequence SEQ ID Nos. 11 to 13 and 15 to 17.

10 According to still another aspect, the present invention relates to an isolated nucleic acid, characterized in that it is chosen from the following nucleic acids:

- a nucleic acid, DNA or RNA, coding for an antibody, or one of its functional fragments or derivatives, according to the present invention and wherein the nucleic sequence coding for the hinge region of said antibody comprises or has a sequence
15 selected from the group consisting of the sequences SEQ ID Nos. 29 to 35 and SEQ ID Nos. 73 to 87.

By nucleic acid, nucleic or nucleic acid sequence, polynucleotide, oligonucleotide, polynucleotide sequence, nucleotide sequence, terms which will be employed indifferently in the present invention, it is intended to indicate a precise
20 linkage of nucleotides, which are modified or unmodified, allowing a fragment or a region of a nucleic acid to be defined, containing or not containing unnatural nucleotides, and being able to correspond just as well to a double-stranded DNA, a single-stranded DNA as to the transcription products of said DNAs.

It must also be understood here that the present invention does not concern the
25 nucleotide sequences in their natural chromosomal environment, that is to say in the natural state. It concerns sequences which have been isolated and/or purified, that is to say that they have been selected directly or indirectly, for example by copy, their environment having been at least partially modified. It is thus likewise intended to indicate here the isolated nucleic acids obtained by genetic recombination by means, for
30 example, of host cells or obtained by chemical synthesis.

A hybridization under conditions of high stringency signifies that the temperature conditions and ionic strength conditions are chosen in such a way that they

allow the maintenance of the hybridization between two fragments of complementary DNA. By way of illustration, conditions of high stringency of the hybridization step for the purposes of defining the polynucleotide fragments described above are advantageously the following.

5 The DNA-DNA or DNA-RNA hybridization is carried out in two steps: (1) prehybridization at 42°C for 3 hours in phosphate buffer (20 mM, pH 7.5) containing 5 x SSC (1 x SSC corresponds to a 0.15 M NaCl + 0.015 M sodium citrate solution), 50 % of formamide, 7 % of sodium dodecyl sulfate (SDS), 10 x Denhardt's, 5 % of dextran sulfate and 1 % of salmon sperm DNA; (2) actual hybridization for 20 hours at
10 a temperature dependent on the size of the probe (i.e.: 42°C, for a probe size > 100 nucleotides) followed by 2 washes of 20 minutes at 20°C in 2 x SSC + 2% of SDS, 1 wash of 20 minutes at 20°C in 0.1 x SSC + 0.1 % of SDS. The last wash is carried out in 0.1 x SSC + 0.1 % of SDS for 30 minutes at 60°C for a probe size > 100 nucleotides. The hybridization conditions of high stringency described above for a polynucleotide of
15 defined size can be adapted by the person skilled in the art for oligonucleotides of greater or smaller size, according to the teaching of Sambrook et al. (1989, Molecular cloning: a laboratory manual. 2nd Ed. Cold Spring Harbor).

The invention likewise relates to a vector comprising a nucleic acid according to the present invention.

20 The invention aims especially at cloning and/or expression vectors which contain a nucleotide sequence according to the invention.

The vectors according to the invention preferably contain elements which allow the expression and/or the secretion of the translated nucleotide sequences in a determined host cell. The vector must therefore contain a promoter, signals of initiation
25 and termination of translation, as well as appropriate regions of regulation of transcription. It must be able to be maintained in a stable manner in the host cell and can optionally have particular signals which specify the secretion of the translated protein. These different elements are chosen and optimized by the person skilled in the art as a function of the host cell used. To this effect, the nucleotide sequences according to the
30 invention can be inserted into autonomous replication vectors in the chosen host, or be integrative vectors of the chosen host.

Such vectors are prepared by methods currently used by the person skilled in the art, and the resulting clones can be introduced into an appropriate host by standard methods, such as lipofection, electroporation, thermal shock, or chemical methods.

The vectors according to the invention are, for example, vectors of plasmidic or viral origin. They are useful for transforming host cells in order to clone or to express the nucleotide sequences according to the invention.

The invention likewise comprises the host cells transformed by or comprising a vector according to the invention.

The host cell can be chosen from prokaryotic or eukaryotic systems, for example bacterial cells but likewise yeast cells or animal cells, in particular mammalian cells. It is likewise possible to use insect cells or plant cells.

The invention likewise concerns animals, except man, which comprise at least one cell transformed according to the invention.

According to another aspect, a subject of the invention is a process for production of an antibody, or one of its functional fragments according to the invention, characterized in that it comprises the following stages:

- a) culture in a medium and appropriate culture conditions of a host cell according to the invention; and
- b) the recovery of said antibodies, or one of their functional fragments, thus produced starting from the culture medium or said cultured cells.

The cells transformed according to the invention can be used in processes for preparation of recombinant polypeptides according to the invention. The processes for preparation of a polypeptide according to the invention in recombinant form, characterized in that they employ a vector and/or a cell transformed by a vector according to the invention, are themselves comprised in the present invention. Preferably, a cell transformed by a vector according to the invention is cultured under conditions which allow the expression of said polypeptide and said recombinant peptide is recovered.

As has been said, the host cell can be chosen from prokaryotic or eukaryotic systems. In particular, it is possible to identify nucleotide sequences according to the invention, facilitating secretion in such a prokaryotic or eukaryotic system. A vector according to the invention carrying such a sequence can therefore advantageously be

used for the production of recombinant proteins, intended to be secreted. In effect, the purification of these recombinant proteins of interest will be facilitated by the fact that they are present in the supernatant of the cell culture rather than in the interior of the host cells.

5 It is likewise possible to prepare the polypeptides according to the invention by chemical synthesis. Such a preparation process is likewise a subject of the invention. The person skilled in the art knows the processes of chemical synthesis, for example the techniques employing solid phases [Steward et al., 1984, Solid phase peptide synthesis, Pierce Chem. Company, Rockford, 111, 2nd ed., (1984)] or techniques using partial
10 solid phases, by condensation of fragments or by a classical synthesis in solution. The polypeptides obtained by chemical synthesis and being able to contain corresponding unnatural amino acids are likewise comprised in the invention.

The antibodies, or one of their functional fragments or derivatives, capable of being obtained by a process according to the invention are likewise comprised in the
15 present invention.

The invention also concerns the antibody of the invention as a medicament.

The invention likewise concerns a pharmaceutical composition comprising by way of active principle a compound consisting of an antibody, or one of its functional fragments according to the invention, preferably mixed with an excipient and/or a
20 pharmaceutically acceptable vehicle.

Another complementary embodiment of the invention consists in a composition such as described above which comprises, moreover, as a combination product for simultaneous, separate or sequential use, an anti-tumoral antibody.

Most preferably, said second anti-tumoral antibody could be chosen through
25 anti-IGF-IR, anti-EGFR, anti-HER2/neu, anti-VEGFR, anti-VEGF, etc., antibodies or any other anti-tumoral antibodies known by the man skilled in the art. It is evident that the use, as second antibody, of functional fragments or derivatives of above mentioned antibodies is part of the invention.

As a most preferred antibody, anti-EGFR antibodies are selected such as for
30 example the antibody C225 (Erbix).

“Simultaneous use” is understood as meaning the administration of the two compounds of the composition according to the invention in a single and identical pharmaceutical form.

“Separate use” is understood as meaning the administration, at the same time, of
5 the two compounds of the composition according to the invention in distinct pharmaceutical forms.

“Sequential use” is understood as meaning the successive administration of the two compounds of the composition according to the invention, each in a distinct pharmaceutical form.

10 In a general fashion, the composition according to the invention considerably increases the efficacy of the treatment of cancer. In other words, the therapeutic effect of the anti-c-Met antibodies according to the invention is potentiated in an unexpected manner by the administration of a cytotoxic agent. Another major subsequent advantage produced by a composition according to the invention concerns the possibility of using
15 lower efficacious doses of active principle, which allows the risks of appearance of secondary effects to be avoided or to be reduced, in particular the effects of the cytotoxic agent.

In addition, this composition according to the invention would allow the expected therapeutic effect to be attained more rapidly.

20 The composition of the invention can also be characterized in that it comprises, moreover, as a combination product for simultaneous, separate or sequential use, a cytotoxic/cytostatic agent.

By “anti-cancer therapeutic agents” or “cytotoxic/cytostatic agents”, it is intended a substance which, when administered to a subject, treats or prevents the
25 development of cancer in the subject’s body. As non limitative example of such agents, it can be mentioned alkylating agents, anti-metabolites, anti-tumor antibiotics, mitotic inhibitors, chromatin function inhibitors, anti-angiogenesis agents, anti-estrogens, anti-androgens or immunomodulators.

Such agents are, for example, cited in the 2001 edition of VIDAL, on the page
30 devoted to the compounds attached to the cancerology and hematology column “Cytotoxics”, these cytotoxic compounds cited with reference to this document are cited here as preferred cytotoxic agents.

More particularly, the following agents are preferred according to the invention.

“Alkylating agent” refers to any substance which can cross-link or alkylate any molecule, preferably nucleic acid (e.g., DNA), within a cell. Examples of alkylating agents include nitrogen mustard such as mechlorethamine, chlorambucol, melphalen, 5 chlorydrate, pipobromen, prednimustin, disodic-phosphate or estramustine; oxazophorins such as cyclophosphamide, altretamine, trofosfamide, sulfofosfamide or ifosfamide; aziridines or imine-ethylenes such as thiotepa, triethylenamine or altetramine; nitrosourea such as carmustine, streptozocin, fotemustin or lomustine; alkyle-sulfonates such as busulfan, treosulfan or improsulfan; triazenes such as 10 dacarbazine; or platinum complexes such as cis-platinum, oxaliplatin and carboplatin.

“Anti-metabolites” refer to substances that block cell growth and/or metabolism by interfering with certain activities, usually DNA synthesis. Examples of anti-metabolites include methotrexate, 5-fluoruracil, floxuridine, 5-fluorodeoxyuridine, capecitabine, cytarabine, fludarabine, cytosine arabinoside, 6-mercaptopurine (6-MP), 15 6-thioguanine (6-TG), chlorodesoxyadenosine, 5-azacytidine, gemcitabine, cladribine, deoxycoformycin and pentostatin.

“Anti-tumor antibiotics” refer to compounds which may prevent or inhibit DNA, RNA and/or protein synthesis. Examples of anti-tumor antibiotics include doxorubicin, daunorubicin, idarubicin, valrubicin, mitoxantrone, dactinomycin, mithramycin, 20 plicamycin, mitomycin C, bleomycin, and procarbazine.

“Mitotic inhibitors” prevent normal progression of the cell cycle and mitosis. In general, microtubule inhibitors or taxoides such as paclitaxel and docetaxel are capable of inhibiting mitosis. Vinca alkaloid such as vinblastine, vincristine, vindesine and vinorelbine are also capable of inhibiting mitosis.

25 “Chromatin function inhibitors” or “topoisomerase inhibitors” refer to substances which inhibit the normal function of chromatin modeling proteins such as topoisomerase I or topoisomerase II. Examples of chromatin function inhibitors include, for topoisomerase I, camptothecine and its derivatives such as topotecan or irinotecan, and, for topoisomerase II, etoposide, etoposide phosphate and teniposide.

30 “Anti-angiogenesis agent” refers to any drug, compound, substance or agent which inhibits growth of blood vessels. Exemplary anti-angiogenesis agents include, but are by no means limited to, razoxin, marimastat, batimastat, prinomastat, tanomastat,

ilomastat, CGS-27023A, halofuginon, COL-3, neovastat, BMS-275291, thalidomide, CDC 501, DMXAA, L-651582, squalamine, endostatin, SU5416, SU6668, interferon-alpha, EMD121974, interleukin-12, IM862, angiostatin and vitaxin.

“Anti-estrogen” or “anti-estrogenic agent” refer to any substance which reduces, antagonizes or inhibits the action of estrogen. Examples of anti-estrogen agents are tamoxifen, toremifene, raloxifene, droloxifene, idoxifene, anastrozole, letrozole, and exemestane.

“Anti-androgens” or “anti-androgen agents” refer to any substance which reduces, antagonizes or inhibits the action of an androgen. Examples of anti-androgens are flutamide, nilutamide, bicalutamide, spironolactone, cyproterone acetate, finasteride and cimitidine.

“Immunomodulators” are substances which stimulate the immune system.

Examples of immunomodulators include interferon, interleukin such as aldesleukine, OCT-43, denileukin diflitox and interleukin-2, tumoral necrosis factors such as tasonermin or other immunomodulators such as lentinan, sizofiran, roquinimex, pidotimod, pegademase, thymopentine, poly I:C or levamisole in conjunction with 5-fluorouracil.

For more detail, the man skilled in the art could refer to the manual edited by the “Association Française des Enseignants de Chimie Thérapeutique” and entitled “Traité de chimie thérapeutique”, vol. 6, Médicaments antitumoraux et perspectives dans le traitement des cancers, edition TEC & DOC, 2003.

Can also be mentioned as chemical agents or cytotoxic agents, all kinase inhibitors such as, for example, gefitinib or erlotinib.

In a particularly preferred embodiment, said composition as a combination product according to the invention is characterized in that said cytotoxic agent is coupled chemically to said antibody for simultaneous use.

In order to facilitate the coupling between said cytotoxic agent and said antibody according to the invention, it is especially possible to introduce spacer molecules between the two compounds to be coupled, such as poly(alkylene) glycols like polyethylene glycol, or else amino acids, or, in another embodiment, to use active derivatives of said cytotoxic agents into which would have been introduced functions capable of reacting with said antibody according to the invention. These coupling

techniques are well known to the person skilled in the art and will not be expanded upon in the present description.

The invention relates, in another aspect, to a composition characterized in that one, at least, of said antibodies, or one of their functional fragments or derivatives, is
5 conjugated with a cell toxin and/or a radioelement.

Preferably, said toxin or said radioelement is capable of inhibiting at least one cell activity of cells expressing the c-Met, in a more preferred manner capable of preventing the growth or the proliferation of said cell, especially of totally inactivating said cell.

10 Preferably also, said toxin is an enterobacterial toxin, especially *Pseudomonas* exotoxin A.

The radioelements (or radioisotopes) preferably conjugated to the antibodies employed for the therapy are radioisotopes which emit gamma rays and preferably iodine¹³¹, yttrium⁹⁰, gold¹⁹⁹, palladium¹⁰⁰, copper⁶⁷, bismuth²¹⁷ and antimony²¹¹. The
15 radioisotopes which emit beta and alpha rays can likewise be used for the therapy.

By toxin or radioelement conjugated to at least one antibody, or one of its functional fragments, according to the invention, it is intended to indicate any means allowing said toxin or said radioelement to bind to said at least one antibody, especially by covalent coupling between the two compounds, with or without introduction of a
20 linking molecule.

Among the agents allowing binding in a chemical (covalent), electrostatic or noncovalent manner of all or part of the components of the conjugate, mention may particularly be made of benzoquinone, carbodiimide and more particularly EDC (1-ethyl-3-[3-dimethyl-aminopropyl]-carbodiimide hydrochloride), dimaleimide, dithiobis-
25 nitrobenzoic acid (DTNB), N-succinimidyl S-acetyl thio-acetate (SATA), the bridging agents having one or more phenylazide groups reacting with the ultraviolets (U.V.) and preferably N-[4-(azidosalicylamino)butyl]-3'-(2'-pyridyldithio)-propionamide (APDP), N-succinimid-yl 3-(2-pyridyldithio)propionate (SPDP), 6-hydrazino-nicotinamide (HYNIC).

30 Another form of coupling, especially for the radioelements, can consist in the use of a bifunctional ion chelator.

Among these chelates, it is possible to mention the chelates derived from EDTA (ethylenediaminetetraacetic acid) or from DTPA (diethylenetriaminepentaacetic acid) which have been developed for binding metals, especially radioactive metals, and immunoglobulins. Thus, DTPA and its derivatives can be substituted by different groups on the carbon chain in order to increase the stability and the rigidity of the ligand-metal complex (Krejcarek et al. (1977); Brechbiel et al. (1991); Gansow (1991); US patent 4,831,175).

For example diethylenetriaminepentaacetic acid (DTPA) and its derivatives, which have been widely used in medicine and in biology for a long time either in their free form, or in the form of a complex with a metallic ion, have the remarkable characteristic of forming stable chelates with metallic ions and of being coupled with proteins of therapeutic or diagnostic interest such as antibodies for the development of radioimmunoconjugates in cancer therapy (Meases et al., 1984; Gansow et al., 1990).

Likewise preferably, said at least one antibody forming said conjugate according to the invention is chosen from its functional fragments, especially the fragments amputated of their Fc component such as the scFv fragments.

As already mentioned, in a preferred embodiment of the invention, said cytotoxic/cytostatic agent or said toxin and/or a radioelement is coupled chemically to at least one of the elements of said composition for simultaneous use.

The present invention comprises the described composition as a medicament.

The present invention moreover comprises the use of the composition according to the invention for the preparation of a medicament.

In another aspect, the invention deals with the use of an antibody, or one of its functional fragments or derivatives, and/or of a composition as above described for the preparation of a medicament intended to inhibit the growth and/or the proliferation of tumor cells.

Another aspect of the invention consists in the use of an antibody, or one of its functional fragments or derivatives and/or of a composition, as described above or the use above mentioned, for the preparation of a medicament intended for the prevention or for the treatment of cancer.

Is also comprised in the present invention a method intended to inhibit the growth and/or the proliferation of tumor cells in a patient comprising the administration

to a patient in need thereof of an antibody, or one of its functional fragments or derivatives according to the invention, an antibody produced by an hybridoma according to the invention or a composition according to the invention.

5 The present invention further comprises a method for the prevention or the treatment of cancer in a patient in need thereof, comprising the administration to the patient of an antibody, or one of its functional fragments or derivatives according to the invention, an antibody produced by an hybridoma according to the invention or a composition according to the invention.

10 In a particular preferred aspect, said cancer is a cancer chosen from prostate cancer, osteosarcomas, lung cancer, breast cancer, endometrial cancer, glioblastoma or colon cancer.

As explained before, an advantage of the invention is to allow the treatment of HGF dependent and independent Met-activation related cancers.

15 In a particular aspect, the invention comprises the use of an antibody, or one of its functional fragments or derivatives and/or of a composition, as described above or the use above mentioned, for the preparation of a medicament intended for the prevention or for the treatment of a patient in need thereof having a cancer characterized by overexpression of c-Met (for example, due to a genic amplification of c-Met) resulting in a ligand-independent constitutive activation of said c-Met receptor .

20 Preferably, said cancer characterized by a genic amplification of c-Met resulting in a ligand-independent constitutive activation of said c-Met receptor is selected from the group consisting of renal cell carcinoma and gastric cancer.

25 The invention, in yet another aspect, encompasses a method of *in vitro* diagnosis of illnesses induced by an overexpression or an underexpression of the c-Met receptor starting from a biological sample in which the abnormal presence of c-Met receptor is suspected, said method being characterized in that it comprises a step wherein said biological sample is contacted with an antibody of the invention, it being possible for said antibody to be, if necessary, labeled.

30 Preferably, said illnesses connected with an abnormal presence of c-Met receptor in said diagnosis method will be cancers.

Said antibody, or one of its functional fragments, can be present in the form of an immunoconjugate or of a labelled antibody so as to obtain a detectable and/or quantifiable signal.

The antibodies labelled according to the invention or their functional fragments include, for example, antibodies called immunoconjugates which can be conjugated, for example, with enzymes such as peroxidase, alkaline phosphatase, beta-D-galactosidase, glucose oxydase, glucose amylase, carbonic anhydrase, acetylcholinesterase, lysozyme, malate dehydrogenase or glucose 6-phosphate dehydrogenase or by a molecule such as biotin, digoxigenin or 5-bromodeoxyuridine. Fluorescent labels can be likewise conjugated to the antibodies or to their functional fragments according to the invention and especially include fluorescein and its derivatives, fluorochrome, rhodamine and its derivatives, GFP (GFP for "Green Fluorescent Protein"), dansyl, umbelliferone etc. In such conjugates, the antibodies of the invention or their functional fragments can be prepared by methods known to the person skilled in the art. They can be coupled to the enzymes or to the fluorescent labels directly or by the intermediary of a spacer group or of a linking group such as a polyaldehyde, like glutaraldehyde, ethylenediaminetetraacetic acid (EDTA), diethylene-triaminepentaacetic acid (DPTA), or in the presence of coupling agents such as those mentioned above for the therapeutic conjugates. The conjugates containing labels of fluorescein type can be prepared by reaction with an isothiocyanate.

Other conjugates can likewise include chemoluminescent labels such as luminol and the dioxetanes, bio-luminescent labels such as luciferase and luciferin, or else radioactive labels such as iodine¹²³, iodine¹²⁵, iodine¹²⁶, iodine¹³³, bromine⁷⁷, technetium^{99m}, indium¹¹¹, indium^{113m}, gallium⁶⁷, gallium⁶⁸, ruthenium⁹⁵, ruthenium⁹⁷, ruthenium¹⁰³, ruthenium¹⁰⁵, mercury¹⁰⁷, mercury²⁰³, rhenium^{99m}, rhenium¹⁰¹, rhenium¹⁰⁵, scandium⁴⁷, tellurium^{121m}, tellurium^{122m}, tellurium^{125m}, thulium¹⁶⁵, thulium¹⁶⁷, thulium¹⁶⁸, fluorine¹⁸, yttrium¹⁹⁹, iodine¹³¹. The methods known to the person skilled in the art existing for coupling the therapeutic radioisotopes to the antibodies either directly or via a chelating agent such as EDTA, DTPA mentioned above can be used for the radioelements which can be used in diagnosis. It is likewise possible to mention labelling with Na[I¹²⁵] by the chloramine T method [Hunter W.M. and Greenwood F.C. (1962) Nature 194:495] or else with technetium^{99m} by the technique of Crockford *et al.*

(US patent 4,424,200) or attached via DTPA as described by Hnatowich (US patent 4,479,930).

Thus, the antibody, or a functional fragment or derivative thereof, according to the invention can be employed in a process for the detection and/or the quantification of
5 an overexpression or of an underexpression, preferably an overexpression, of the c-Met receptor in a biological sample, characterized in that it comprises the following steps:

- a) the contacting of the biological sample with an antibody, or a functional fragment or derivative thereof, according to the invention; and
- b) the demonstration of the c-Met/antibody complex possibly formed.

10 In a particular embodiment, the antibody, or a functional fragment or derivative thereof, according to the invention, can be employed in a process for the detection and/or the quantification of the c-Met receptor in a biological sample, for the monitoring of the efficacy of a prophylactic and/or therapeutic treatment of c-Met-dependent cancer.

15 More generally, the antibody or a functional fragment or derivative thereof, according to the invention can be advantageously employed in any situation where the expression of the c-Met- receptor must be observed in a qualitative and/or quantitative manner.

Preferably, the biological sample is formed by a biological fluid, such as serum,
20 whole blood, cells, a tissue sample or biopsies of human origin.

Any procedure or conventional test can be employed in order to carry out such a detection and/or dosage. Said test can be a competition or sandwich test, or any test known to the person skilled in the art dependent on the formation of an immune complex of antibody-antigen type. Following the applications according to the
25 invention, the antibody or a functional fragment or derivative thereof can be immobilized or labelled. This immobilization can be carried out on numerous supports known to the person skilled in the art. These supports can especially include glass, polystyrene, poly-propylene, polyethylene, dextran, nylon, or natural or modified cells. These supports can be either soluble or insoluble.

30 By way of example, a preferred method brings into play immunoenzymatic processes according to the ELISA technique, by immunofluorescence, or radio-immunoassay (RIA) technique or equivalent.

Thus, the present invention likewise comprises the kits or sets necessary for carrying out a method of diagnosis of illnesses induced by an overexpression or an underexpression of the c-Met receptor or for carrying out a process for the detection and/or the quantification of an overexpression or of an underexpression of the c-Met
5 receptor in a biological sample, preferably an overexpression of said receptor, characterized in that said kit or set comprises the following elements:

- a) an antibody, or a functional fragment or derivative thereof, according to the invention;
- b) optionally, the reagents for the formation of the medium favorable to the
10 immunological reaction;
- c) optionally, the reagents allowing the demonstration of c-Met/antibody complexes produced by the immunological reaction.

A subject of the invention is likewise the use of an antibody or a composition according to the invention for the preparation of a medicament intended for the specific
15 targeting of a biologically active compound to cells expressing or overexpressing the c-Met receptor.

It is intended here by biologically active compound to indicate any compound capable of modulating, especially of inhibiting, cell activity, in particular their growth, their proliferation, transcription or gene translation.

20 A subject of the invention is also an *in vivo* diagnostic reagent comprising an antibody according to the invention, or a functional fragment or derivative thereof, preferably labelled, especially radiolabelled, and its use in medical imaging, in particular for the detection of cancer connected with the expression or the overexpression by a cell of the c-Met receptor.

25 The invention likewise relates to a composition as a combination product or to an anti-c-Met/toxin conjugate or radioelement, according to the invention, as a medicament.

Preferably, said composition as a combination product or said conjugate according to the invention will be mixed with an excipient and/or a pharmaceutically
30 acceptable vehicle.

In the present description, pharmaceutically acceptable vehicle is intended to indicate a compound or a combination of compounds entering into a pharmaceutical

composition not provoking secondary reactions and which allows, for example, facilitation of the administration of the active compound(s), an increase in its lifespan and/or in its efficacy in the body, an increase in its solubility in solution or else an improvement in its conservation. These pharmaceutically acceptable vehicles are well
5 known and will be adapted by the person skilled in the art as a function of the nature and of the mode of administration of the active compound(s) chosen.

Preferably, these compounds will be administered by the systemic route, in particular by the intravenous route, by the intramuscular, intradermal, intraperitoneal or subcutaneous route, or by the oral route. In a more preferred manner, the composition
10 comprising the antibodies according to the invention will be administered several times, in a sequential manner.

Their modes of administration, dosages and optimum pharmaceutical forms can be determined according to the criteria generally taken into account in the establishment of a treatment adapted to a patient such as, for example, the age or the body weight of
15 the patient, the seriousness of his/her general condition, the tolerance to the treatment and the secondary effects noted.

Other characteristics and advantages of the invention appear in the continuation of the description with the examples and the figures wherein:

Figure 1: Effect of irrelevant IgG1 Mabs from mouse and human origin and PBS
20 on c-Met receptor phosphorylation on A549 cells.

Figures 2A and 2B: Effect of murine and humanized 224G11 Mabs produced as a human IgG1/kappa isotype on c-Met receptor phosphorylation on A549 cells.

Figure 2A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

25 Figure 2B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 3A and 3B: Comparison between murine 224G11 Mab and chimeric 224G11 Mabs containing various engineered hinge regions, on c-Met receptor phosphorylation on A549 cells.

30 Figure 3A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 3B: antagonist effect calculated as percentage of inhibition of the maximal

stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 4A and 4B: Comparison between murine 224G11 Mab and chimeric and humanized 224G11 Mabs produced as a human IgG2/kappa isotype, on c-Met receptor phosphorylation on A549 cells.

5 Figure 4A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 4B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 5A and 5B: Comparison between murine 224G11 Mab and chimeric and
10 humanized 224G11 Mabs produced as an engineered hinge mutant TH7IgG1/kappa, on c-Met receptor phosphorylation on A549 cells.

Figure 5A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 5B: antagonist effect calculated as percentage of inhibition of the maximal
15 stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figures 6A and 6B, Figures 7A and 7B, Figures 8A and 8B, Figures 9A and 9B, Figures 10A and 10B: BRET models with Figures A: c-Met dimerization model; and Figures B: c-Met activation model.

Figure 11: c-Met recognition by chimeric and humanized 224G11 forms.



20 Figure 12: Effect of murine and chimeric antibodies on HGF-induced proliferation of NCI-H441 cells *in vitro*. NCI-H441 cells were plated in serum-free medium. 24 hours after plating m224G11 and [224G11]chim were added either in absence or in presence of HGF. Black arrows indicate the wells plated with cells alone either in absence  or in presence  of HGF. A murine IgG1 (mIgG1) was introduced
25 as an isotype control.

Figure 13: *In vivo* comparison of murine and IgG1 chimeric 224G11 Mabs on the NCI-H441 xenograft model.

Figures 14A and 14B: Effect of the murine 224G11 Mab and of various chimeric and humanized versions of this antibody on HGF-induced proliferation of
30 NCI-H441 cells *in vitro*. NCI-H441 cells were plated in serum-free medium. Twenty four hours after plating antibody to be tested were added either in absence or in presence of HGF. In panel (Figure 14A), the murine m224G11, chimeric IgG1

[224G11]chim, humanized IgG1 [224G11] [Hz1], [224G11] [Hz2], [224G11] [Hz3] versions were shown. In panel (Figure 14B), the murine m224G11 and various chimeric IgG1 forms ([224G11] chim, [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim], [224G11] [TH7 chim]) were presented. Black arrows indicate the wells plated with cells alone either in absence  or in presence  of HGF. A murine IgG1 was introduced as a negative control for agonist activity. The m5D5 was used as a dose-dependent full agonist control.

Figure 15: Effect of the murine 224G11 Mab and of various chimeric and humanized versions of this antibody on HGF-induced proliferation of NCI-H441 cells in vitro. NCI-H441 cells were plated in serum-free medium. Twenty four hours after plating antibody to be tested were added either in absence or in presence of HGF. The murine m224G11, [224G11] chim, [224G11] [TH7 chim]) IgG1 chimeric forms and [224G11] [TH7 Hz1], [224G11] [TH7 Hz3],) were presented. Black arrows indicate the wells plated with cells alone either in absence  or in presence  of HGF. A murine IgG1 was introduced as a negative control for agonist activity. The m5D5 was used as a dose-dependent full agonist control.

Figures 16A-16C: *In vivo* comparison of murine, chimeric and humanized 224G11 Mabs on the NCI-H441 xenograft model.

Figure 17A: agonist effect calculated as percentage versus maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 17B: antagonist effect calculated as percentage of inhibition of the maximal stimulation of c-Met phosphorylation by HGF [100 ng/ml].

Figure 18: BRET models with c-Met activation model.

Figures 19A-19B: Effect of m224G11 and h224G11 on c-Met degradation on A549 cells. A) Mean of 4 independent experiments +/- s.e.m. B) Western blot image representative of the 4 independent experiments performed.

Figures 20A-20B: Effect of m224G11 and h224G11 on c-Met degradation on NCI-H441 cells. A) Mean of 4 independent experiments +/- s.e.m. B) Western blot image representative of the 4 independent experiments performed.

Figure 21: Set up of an ELISA to evaluate c-Met shedding.

Figure 22: *In vitro* evaluation of c-Met shedding on NCI-H441 cells treated for 5 days with m224G11. mIgG1 is an irrelevant antibody used as an isotype control.

Figure 23: *In vitro* evaluation of c-Met shedding on amplified Hs746T, MKN45 and EBC-1 cell lines treated for 5 days with m224G11. mIgG1 is an irrelevant antibody used as an isotype control. PMA is a shedding inducer used as a positive control.

Figure 24: *In vitro* evaluation of c-Met shedding on NCI-H441 and amplified Hs746T, MKN45 and EBC-1 cell lines treated for 5 days with m224G11. mIgG1 is an irrelevant antibody used as an isotype control. PMA is a shedding inducer used as a positive control.

Figure 25: Study of intrinsic phosphorylation of h224G11 on Hs746T cell line.

Figures 26A-26B: Study of intrinsic phosphorylation of h224G11 on NCI-H441 cell line. A) phospho-ELISA and B) Western analysis.

Figures 27A-27B: Study of intrinsic phosphorylation of h224G11 on Hs578T cell line. A) phospho-ELISA and B) Western analysis.

Figures 28A-28B: Study of intrinsic phosphorylation of h224G11 on NCI-H125 cell line. A) phospho-ELISA and B) Western analysis.

Figures 29A-29B: Study of intrinsic phosphorylation of h224G11 on T98G cell line. A) phospho-ELISA and B) Western analysis.

Figures 30A-30B: Study of intrinsic phosphorylation of h224G11 on MDA-MB-231 cell line. A) phospho-ELISA and B) Western analysis.

Figures 31A-31B: Study of intrinsic phosphorylation of h224G11 on PC3 cell line. A) phospho-ELISA and B) Western analysis.

Figure 32: Study of intrinsic phosphorylation of h224G11 on HUVEC cells.

Figure 33: In vivo comparison of the wild type murine 224G11 antibody with a chimeric hinge-engineered 224G11[C2D5-7] Mabs on the NCI-H441 xenograft model.

Figures 34A-34B: ADCC induction by h224G11 on both Hs746T and NCI-H441 cells. ⁵¹Cr-labeled Hs746T (A) or NCI-H441 (B) cells loaded (bold squares) or not (empty squares) with h224G11 were mixed with different ratio of human NK cells and incubated for 4 hr. Cells were harvested and cpm of ⁵¹Cr released by lysis was counted. The results are plotted as percentage of lysis against the effector/target cell ratio. NL for non loaded cells.

Figures 35A-35C: h224G11 staining in tumor xenograft which expressed various level of c-Met (A: Hs746T amplified cell line for c-Met, B: NCI-H441 high level of c-Met expression and C: MCF-7 low level of c-Met).

Figures 36A-36D: *In vivo* activity of h224G11 on the Hs746T xenograft model. A) FACS analysis showing c-Met expression on Hs746T cells compared to other cell lines expressing c-Met. B) QRTPCR showing c-Met amplification. C) c-Met phosphorylation in absence or in presence of HGF. D) *In vivo* model.

5 Figures 37A-37D: *In vivo* activity of h224G11 on the MKN-45 xenograft model. A) FACS analysis showing c-Met expression on MKN45 cells compared to other cell lines expressing c-Met. B) QRTPCR showing c-Met amplification. C) c-Met phosphorylation in absence or in presence of HGF. D) *In vivo* model.

Figures 38A-38D: *In vivo* activity of h224G11 on the EBC-1 xenograft model. 10 A) FACS analysis showing c-Met expression on EBC-1 cells compared to other cell lines expressing c-Met. B) QRTPCR showing c-Met amplification. C) c-Met phosphorylation in absence or in presence of HGF. D) *In vivo* model.

Figures 39A-39H: IHC examination of Hs746T tumors in mice treated with the h224G11 Mab compared to control-treated tumors. (A) and (B) panels correspond to 15 isotype control antibodies. (C) and (D) show the expression of c-Met on tumor sections from mice treated with PBS or h224G11 respectively demonstrating that a chronic treatment with h224G11 induced a dramatic down regulation of c-Met. Ki-67 staining is significantly decreased on treated tumors (F) compared to PBS control samples (E). A significant apoptosis was observed in tumor treated with h224G11 (H) while no 20 apoptosis was noticed in control tumors (G).

Example 1: Generation of antibodies against c-Met

To generate anti-c-Met antibodies 8 weeks old BALB/c mice were immunized either 3 to 5 times subcutaneously with a CHO transfected cell line that express c-Met on its plasma membrane (20×10^6 cells/dose/mouse) or 2 to 3 times with a c-Met 25 extracellular domain fusion protein (10-15 μ g/dose/mouse) (R&D Systems, Catalog # 358MT) or fragments of this recombinant protein mixed with complete Freund adjuvant for the first immunization and incomplete Freund adjuvant for the following ones. Mixed protocols in which mice received both CHO-cMet cells and recombinant proteins were also performed. Three days before cell fusion, mice were boosted i.p. or i.v. with 30 the recombinant protein or fragments. Then spleens of mice were collected and fused to SP2/0-Ag14 myeloma cells (ATCC) and subjected to HAT selection. Four fusions were performed. In general, for the preparation of monoclonal antibodies or their functional

fragments, especially of murine origin, it is possible to refer to techniques which are described in particular in the manual "Antibodies" (Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor NY, pp. 726, 1988) or to the technique of preparation of hybridomas described by Kohler and
5 Milstein (Nature, 256:495-497, 1975).

Obtained hybridomas were initially screened by ELISA on the c-Met recombinant protein and then by FACS analysis on A549 NSCLC, BxPC3 pancreatic, and U87-MG glioblastoma cell lines to be sure that the produced antibodies will be able to also recognize the native receptor on tumor cells. Positive reactors on these 2 tests
10 were amplified, cloned and a set of hybridomas was recovered, purified and screened for its ability to inhibit *in vitro* cell proliferation in the BxPC3 model.

For that purpose 50 000 BxPC3 cells were plated in 96 well plates in RPMI medium, 2 mM L. Glutamine, without SVF. 24 hours after plating, antibodies to be tested were added at a final concentration ranging from 0.0097 to 40 µg/ml 60 min
15 before addition of 100 ng/ml of hHGF. After 3 days, cells were pulsed with 0.5 µCi of [³H]thymidine for 16 hours. The magnitude of [³H]thymidine incorporated into trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results were expressed as raw data to really evaluate the intrinsic agonistic effect of each Mab.

20 Then antibodies inhibiting at least 50% cell proliferation were evaluated for their activity on c-Met dimerization and activation BRET analysis on transfected cells. c-Met receptor activity was quantified by measuring the Gab1 signalling molecule recruitment on activated c-Met. For that purpose, CHO stable cell lines expressing C-Met-Rluc or C-Met-Rluc and C-Met-K1100A-YFP for c-Met dimerization or C-Met-Rluc and a
25 mutated form of Gab1 [Maroun et al., Mol. Cell. Biol., 1999, 19:1784-1799] fused to YFP for c-Met activation were generated. Cells were distributed in white 96 well microplates in DMEM-F12/FBS 5 % culture medium one or two days before BRET experiments. Cells were first cultured at 37°C with CO₂ 5 % in order to allow cell attachment to the plate. Cells were then starved with 200 µl DMEM/well overnight.
30 Immediately prior to the experiment, DMEM was removed and cells quickly washed with PBS. Cells were incubated in PBS in the presence or absence of antibodies to be tested or reference compounds, 10 min at 37°C prior to the addition of coelenterazine

with or without HGF in a final volume of 50 μ l. After incubation for further 10 minutes at 37°C, light-emission acquisition at 485 nm and 530 nm was initiated using the Mithras luminometer (Berthold) (1s/wave length/well repeated 15 times).

BRET ratio has been defined previously [Angers et al., Proc. Natl. Acad. Sci. USA, 2000, 97:3684-3689] as: [(emission at 530 nm) - (emission at 485 nm) X Cf] / (emission at 485 nm), where Cf corresponds to (emission at 530 nm) / (emission at 485 nm) for cells expressing Rluc fusion protein alone in the same experimental conditions. Simplifying this equation shows that BRET ratio corresponds to the ratio 530/485 nm obtained when the two partners were present, corrected by the ratio 530/485 nm obtained under the same experimental conditions, when only the partner fused to R. reniformis luciferase was present in the assay. For the sake of readability, results are expressed in milliBRET units (mBU); mBU corresponds to the BRET ratio multiplied by 1000.

After this second *in vitro* test, the antibody 224G11 i) without intrinsic activity as a whole molecule in the functional test of proliferation, ii) inhibiting significantly BxPC3 proliferation and iii) inhibiting c-Met dimerization was selected. In the experiments, the 5D5 Mab, generated by Genentech, and available at the ATCC, was added as a control for the intrinsic agonistic activity.

20 **Example 2: Humanization process of mouse 224G11 Mab by CDR-grafting** 1° Humanization of the light chain variable domain (VL)

As a preliminary step, the nucleotide sequence of 224G11 VL was compared to the murine germline gene sequences included in the IMGT database (<http://imgt.cines.fr>). Murine IGKV3-5*01 and IGKJ4*01 germline genes showing a sequence identity of 99.31 % for the V region and 94.28 % for the J region, respectively, have been identified. Regarding these high homologies, the 224G11VL nucleotide sequence has been used directly to search for human homologies, instead of corresponding mouse germlines.

In a second step, the human germline gene displaying the best identity with the 224G11VL has been searched to identify the best human candidate for the CDR grafting. For optimization of the selection, alignments between the amino acid sequences have been performed. The human IGKV4-1*01 germline gene yielded a

sequence identity of 67.30 %, but showed a different length for CDR1 (10 amino acids in 224G11 VL and 12 amino acids in IGKV4-1*01). For the J region, the human IGKJ4*02 germline gene (sequence identity of 77.14 %) was selected.

In a next step, mouse 224G11 VL CDR regions were engrafted into the above
5 selected human framework sequences. Each amino acid position was analyzed for several criteria such as participation in VH/VL interface, in antigen binding or in CDR structure, localization of the residue in the 3D structure of the variable domain, CDR anchors, residues belonging to the Vernier zone. Three humanized versions, corresponding to SEQ ID No. 8, SEQ ID No. 9 and SEQ ID No. 10 were constructed,
10 and containing respectively four (4, 39, 40, 84), two (39, 40) or one (40) murine residues in their FR regions and the CDRs corresponding to mouse 224G11 VL.

2°) Humanization of the heavy chain variable domain (VH)

As a preliminary step, the nucleotidic sequence of the 224G11 VH was compared to the murine germline genes sequences included in the IMGT database
15 (<http://imgt.cines.fr>).

Murine IGHV1-18*01, IGHD2-4*01 and IGHJ2*01 germline genes with a sequence identity of 92.70 % for the V region, 75.00 % for the D region and 89.36 % for the J region, respectively, have been identified. Regarding these high homologies, it has been decided to use directly the 224G11VH nucleotide sequences to search for
20 human homologies, instead of corresponding mouse germlines.

In a second step, the human germline gene displaying the best identity with the 224G11 VH has been searched to identify the best human candidate for the CDR grafting. To this end, the nucleotidic sequence of 224G11 VH has been aligned with the human germline genes sequences belonging to the IMGT database. The human IGHV1-
25 2*02 V sequence exhibited a sequence identity of 75.00 % at the nucleotide level and 64.30 % at the amino acid level. Looking for homologies for the J region led to the identification of the human IGHJ4*04 germline gene with a sequence identity of 78.72 %.

In a next step, mouse 224G11 VH CDR regions were engrafted into the above
30 selected human framework sequences. Each amino acid position was analyzed for several criteria such as participation in VH/VL interface, in antigen binding or in CDR structure, localization of the residue in the 3D structure of the variable domain, CDR

anchors, residues belonging to the Vernier zone. One fully humanized form, corresponding to SEQ ID 4 was constructed; it contains exclusively human residues in its FR regions and the CDRs corresponding to mouse 224G11 VH.

5 **Example 3: Engineering of improved hinge mutants**

It is well known by the skilled artisan that the hinge region strongly participates in the flexibility of the variable domain of immunoglobulins (see Brekke et al., 1995; Roux et al., 1997). During the chimerization process of 224G11 Mab, the mouse constant domain IGHG1 was replaced by the equivalent IGHG1 portion of human
10 origin. Since the amino acid sequence of the hinge region were highly divergent, “murinization” of the hinge region was performed in order to keep its length and rigidity. Since the human IGHG2 hinge region corresponds to the closest homologue of the mouse IGHG1 hinge, this sequence was as well considered. A series of 7 different hinge sequences were constructed (SEQ ID Nos. 22 to 28) by incorporating portions of
15 the mouse IGHG1 and the human IGHG2 hinges into the human IGHG1 hinge portion.

Another series of hinge mutants was designed and constructed (SEQ ID Nos. 58 to 72) to evaluate the influence of either an additional cysteine and its position along the hinge domain, deletion of 1, 2, 3 or 4 amino acids along the hinge domain and a combination of these two parameters (cysteine addition and amino acid deletion).

20

Example 4: Production of humanized 224G11 Mab and engineered hinge Mab formats

All above described Mab forms containing either chimeric, humanized and/or engineered hinge regions were produced upon transient transfection and by using the
25 HEK293/EBNA system with a pCEP4 expression vector (InVitrogen, US).

The entire nucleotide sequences corresponding to the humanized versions of the variable domain of 224G11 Mab light (SEQ ID No. 18, SEQ ID No. 19 and SEQ ID No. 20) and heavy (SEQ ID No. 14) chains were synthesized by global gene synthesis (Genecust, Luxembourg). They were subcloned into a pCEP4 vector (InVitrogen, US)
30 carrying the entire coding sequence of the constant domain [CH1-Hinge-CH2-CH3] of a human IgG1 or IgG2 immunoglobulin. Modification of the hinge region was performed by exchanging a {*NheII-BclI*} restriction fragment by the equivalent portion carrying

the desired modifications, each respective {*NheI-BclI*} fragment being synthesized by global gene synthesis (Genecust, LU). All cloning steps were performed according to conventional molecular biology techniques as described in the Laboratory manual (Sambrook and Russel, 2001) or according to the supplier's instructions. Each genetic
5 construct was fully validated by nucleotide sequencing using Big Dye terminator cycle sequencing kit (Applied Biosystems, US) and analyzed using a 3100 Genetic Analyzer (Applied Biosystems, US).

Suspension-adapted HEK293 EBNA cells (Invitrogen, US) were routinely grown in 250 ml flasks in 50 ml of serum-free medium Excell 293 (SAFC Biosciences)
10 supplemented with 6 mM glutamine on an orbital shaker (110 rpm rotation speed). Transient transfection was performed with $2 \cdot 10^6$ cells/ml using linear 25 kDa polyethyleneimine (PEI) (Polysciences) prepared in water at a final concentration of 1 mg/ml mixed and plasmid DNA (final concentration of 1.25 μ g/ml for heavy to light chain plasmid ratio of 1:1). At 4 hours post-transfection, the culture was diluted with
15 one volume of fresh culture medium to achieve a final cell density of 10^6 cells/ml. Cultivation process was monitored on the basis of cell viability and Mab production. Typically, cultures were maintained for 4 to 5 days. Mabs were purified using a conventional chromatography approach on a Protein A resin (GE Healthcare, US).

All different forms of Mabs were produced at levels suitable with functional
20 evaluations. Productivity levels are typically ranging between 15 and 30 mg/l of purified Mabs.

Example 5: Evaluation of c-Met phosphorylation status by a Phospho-c-Met-specific ELISA assay

25 This functional assay allows to monitor modulation c-Met phosphorylation status either by Mabs alone or in the co-presence of HGF.

A549 cells were seeded in a 12MW plate in complete growth medium [F12K + 10 % FCS]. Cells were starved for 16 hours before stimulation with HGF [100 ng/ml], and each Mab to be tested was added at its final concentration of 30 μ g/ml 15 minutes
30 prior to ligand stimulation. Ice-cold lysis buffer was added 15 minutes after the addition of HGF to stop the phosphorylation reaction. Cells were scraped mechanically and cell lysates were collected by centrifugation at 13000 rpm for 10 min. at 4°C and correspond

to the supernatant phase. Protein content was quantified using a BCA kit (Pierce) and stored at -20°C until use. The phosphorylation status of c-Met was quantified by ELISA. A goat anti-c-Met Mab (R&D, ref AF276) was used as a capture antibody (overnight coating at 4°C) and after a saturation step with a TBS-BSA 5% buffer (1
5 hour at room temperature (RT)), 25 μg of protein lysates were added to each well of the coated 96MW plate. After a 90 minutes incubation at RT, plates were washed four time and the detection antibody was added (anti-phospho-c-Met Mab, directed against the phopshorylated Tyr residues at position 1230, 1234 and 1235). After an additional 1 hour incubation and 4 washes, an anti-rabbit antibody coupled to HRP (Biosource) was
10 added for 1 hour at RT, and the luminescence detection was performed by adding Luminol. Luminescence readings were on a Mithras LB920 multimode plate reader (Berthold).

Both basal and HGF [100 ng/ml]-induced c-Met receptor phosphorylation level were unaffected neither by PBS treatment, nor by the addition of mouse or human Mabs
15 which do not target human c-Met receptor (Figure 1). On the other hand, mouse (m) 224G11 Mab strongly inhibited HGF [100 ng/ml]-induced c-Met phosphorylation (Figure 2B) without altering by itself receptor phosphorylation (Figure 2A). Surprisingly, the chimeric form of 224G11 Mab (224G11chim/IgG1), meaning variable domain (VH+VL) from m224G11 combined with human constant domain IgG1/kappa
20 yielded strong (17 % of maximal HGF effect, Figure 2A) agonist activity associated with a reduced antagonist efficacy (54 % inhibition of HGF maximal effect compared to the m224G11 that yields 75% inhibition of HGF maximum effect, Figure 2B). Three humanized forms of 224G11 Mab, [224G11]Hz1/IgG1, [224G11]Hz2/IgG1 and [224G11]Hz3/IgG1, also constructed on a human IgG1/kappa backbone, yielded also
25 decreased antagonist efficacy and significant agonist activity (11 to 24 % of maximal HGF level) as compared to mouse 224G11 (Figures 2A and 2B). A series of engineered versions of the heavy chain hinge domain were constructed and assayed in the c-Met receptor phosphorylation assay. As shown in Figure 3A, an important reduction of the agonist effect associated with the hIgG1/kappa isotype was observed for both the IgG2-
30 based construct and for engineered IgG1/kappa constructs [MH, MUP9H and TH7]. A concomitant increase in antagonist efficacy was as well obtained. The hIgG1/kappa-based TH7 hinge mutant, with the most human sequence, was selected to complete the

humanization process. In a next step, three humanized versions of 224G11 Mab variable domain were generated by combination to either a human IgG2/kappa or an IgG1/kappa-based TH7 engineered hinge constant domain. For the hIgG2/kappa humanized constructs, the humanized version Hz3 yielded strong agonism (Figure 4A),
5 and for all three humanized versions, the antagonist efficacy was below that observed with murine 224G11 Mab and comparable to the chimeric hIgG1-based Mab (56-57 % inhibition of HGF effect, Figure 4B). On the other hand, combination of the three humanized versions Hz1, Hz2 or Hz3 to the engineered IgG1/TH7 mutant almost fully restored the properties of mouse 224G11 Mab in terms of weak agonist activity (5-6 %
10 of HGF effect) and strong antagonist efficacy (68 to 72 % inhibition of HGF effect) of c-Met receptor phosphorylation (Figures 5A and 5B). These variants were highly improved as compared to chimeric IgG1-based 224G11 Mab but also to IgG2-based humanized forms.

A second series of engineered versions of the heavy chain hinge domain was
15 constructed and assayed in the c-Met receptor phosphorylation assay. As shown in figure 17A, all those new versions (c224G11[C2], c224G11[C3], c224G11[C5], c224G11[C6], c224G11[C7], c224G11[Δ1-3], c224G11[C7Δ6], c224G11[C6Δ9], c224G11[C2Δ5-7], c224G11[C5Δ2-6], c224G11[C9Δ2-7] and c224G11[Δ5-6-7-8]) exhibited weaker agonist effect than c224G11 since their agonism activities are
20 comprised between 6 and 14% of the HGF effect compared to 23% for c224G11. As c224G11[TH7], all those new versions exhibited a concomitant increase in antagonist efficacy [figure 17B]. Those results showed that engineering of the heavy chain domain by point mutation and/or deletion could modify agonistic/antagonistic properties of an antibody.

25

Example 6: BRET analysis

In a first set of experiments, it had been control that irrelevant mouse IgG1, human IgG1 and human IgG2 had no effect of HGF induced BRET signal in both BRET models (representative experiment out of 12 independent experiments; Figure 6).
30 These Mabs are forthwith cited as controls.

The effect of a IgG1 chimeric form of mouse 224G11 Mab ([224G11]chim) on both c-Met dimerization and c-met activation BRET model was evaluated. While mouse

224G11 Mab inhibited 59.4% of the HGF induced BRET signal on c-Met dimerization model, [224G11]chim Mab inhibited only 28.9% (Figure 7A). [224G11]chim antibody was also less effective in inhibiting HGF induced c-Met activation since [224G11]chim and m224G11 antibodies inhibited respectively 34.5% and 56.4% of HGF induced BRET signal (Figure 7B). Moreover, m224G11 alone had no effect on c-Met activation while [224G11]chim had a partial agonist effect on c-Met activation corresponding to 32.9% of the HGF induced signal. This partial agonist effect of the [224G11]chim was also seen on c-Met dimerization BRET model since [224G11]chim alone induced a BRET increase corresponding to 46.6% of HGF-induced signal *versus* 21.3% for m224G11 (Figure 7A).

In Figures 8A and 8B, hinge mutated chimeric forms of 224G11 antibody showed a greater inhibitory effect on HGF induced BRET signal than [224G11]chim since they showed a 59.7%, 64.4%, 53.2% and 73.8% inhibition of the HGF induced activation BRET signal (Figure 8B) and 61.8%, 64.4% 52.5% and 64.4% inhibition of the HGF induced c-Met dimerization BRET signal (Figure 8A) for [224G11][MH chim], [224G11][MUP9H chim], [224G11][MMCH chim] and [224G11][TH7 chim] respectively. Contrary to [224G11]chim, which had a partial agonist effect on c-Met activation, hinge mutated chimerical forms of 224G11 antibody showed no significant effect on c-Met activation alone (5.1%, 7.6%, -2.0% and -6.9% respectively) as observed for m224G11.

In Figure 9B, like the [224G11] [TH7 chim], the 3 humanized versions of 224G11 IgG1 antibody with the TH7 hinge induced no significant increased of BRET signal in activation model when tested alone and showed a strong inhibition of HGF induced BRET signal: 59.9%, 41.8% and 57.9% for the Hz1, Hz2 and Hz3 forms respectively. Moreover, [224G11] [TH7 Hz1], [224G11] [TH7 Hz2] and [224G11] [TH7 Hz3] inhibited HGF induced BRET signal on dimerization model of 52.2%, 35.8% and 49.4% respectively (figure 9A).

Contrary to [224G11]chim, the chimeric form of 224G11 IgG2 antibody ([224G11] [IgG2 chim]) showed no partial agonist effect alone and inhibited 66.3% of the HGF effect on c-Met activation model (Figure 10B). On c-Met dimerization model, [224G11] [IgG2 chim] inhibited 62.4% of the HGF induced BRET signal (Figure 10A).

The agonist efficacy of the second series of engineered versions of the heavy

chain hinge domain was evaluated in c-Met activation BRET model (Figure 18). In contrast to c224G11, which had a partial agonist effect on c-Met activation, c224G11[C2], c224G11[C3], c224G11[C5], c224G11[C6], c224G11[C7], c224G11[Δ1-3], c224G11[C7Δ6], c224G11[C6Δ9], c224G11[C2Δ5-7],
5 c224G11[C5Δ2-6], c224G11[C9Δ2-7] and c224G11[Δ5-6-7-8] hinge mutated chimeric forms of 224G11 antibody showed no significant effect on c-Met activation alone.

Example 7: c-Met recognition by chimeric and humanized 224G11 forms

A direct ELISA has been set up to determine the binding ability of the various
10 chimeric and humanized forms on the recombinant c-Met. Briefly recombinant dimeric c-Met from R&D Systems was coated at 1.25 μg/ml on 96-well Immunlon II plates. After an overnight incubation at 4°C, wells were saturated with a 0.5% gelatine/PBS solution. Plates were then incubated for 1 hour at 37°C before addition of 2 fold dilutions of antibodies to be tested. Plates were incubated an additional hour before
15 addition of a goat anti-mouse IgG HRP for detecting the murine antibody and a goat anti-human Kappa light chain HRP for chimeric and humanized antibody recognition. Plates were incubated for one hour and the peroxidase substrate TMB Uptima was added for 5 mn before neutralization with H₂SO₄ 1M. Results presented in Figure 11 showed that all tested forms were comparable for c-Met recognition.

20

Example 8: Effect of murine and chimeric 224G11 on HGF-induced proliferation of NCI-H441 cells in vitro

NCI-H441 cells from ATCC were routinely cultured in RPMI 1640 medium (Invitrogen Corporation, Scotland, UK), 10% FCS (Invitrogen Corporation), 1% L-
25 Glutamine (Invitrogen corporation). For proliferation assays, cells were split 3 days before use so that they were in the confluent phase of growth before plating. NCI-H441 cells were plated in 96-well tissue culture plates at a density of 3.75x10⁴ cells/well in 200 μl of serum free medium (RPMI 1640 medium plus 1% L-Glutamine). Twenty four hours after plating, antibodies to be tested were added to NCI-H441 and incubated at
30 37°C for thirty minutes before adding HGF at a final concentration of 400 ng/ml (5 nM) for 142 additional hours. The dose range tested for each antibody is from 10 to 0.0097 μg/ml (final concentration in each well). In this experiment, a murine IgG1 Mab

was added as a murine isotype control and the tested antibodies were the following one: m224G11 and its human IgG1 chimeric form identified as [224G11]chim. Wells plated with cells alone +/- HGF were also included. Then cells were pulsed with 0.25 μ Ci of [^{3} H]Thymidine (Amersham Biosciences AB, Uppsala, Sweden) for 7 hours and 30 minutes. The magnitude of [^{3} H]Thymidine incorporated in trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results are expressed as non transformed cpm data to better evaluate the potential intrinsic agonist activity that could occur with anti-c-Met Mabs when added alone to tumour cell.

Results described in Figure 12 demonstrated that, as expected, the murine antibody m224G11 displayed no agonist effect when added alone to cancer cells whatever the tested dose. No significant inhibition of the HGF-induced proliferation was observed with the isotype control regarding to the cpm variations observed for this compound in this experiment. When added alone, the m224G11 antibody did not show any agonist effect compared to the mIgG1 isotype control Mab or cells alone. A dose dependent anti-proliferative activities reaching 78% was observed for m224G11 (% inhibition calculation: $100 - [(cpm\ cells + Mab\ to\ be\ tested - mean\ cpm\ background\ mIgG1) \times 100 / (mean\ cpm\ cells + HGF - mean\ cpm\ cells\ alone)]$). Surprisingly, the chimeric form of the 224G11 Mabs induced a significant, dose dependent agonist effect when added alone. This agonist effect had an impact on the in vitro inhibition of HGF-induced proliferation that shifted from 78% for the murine 224G11 to 50% for its chimeric form. To determine whether such "lower" in vitro intrinsic agonist activity was compatible with an unchanged in vivo effect, both m224G11 and [224G11]chim were produced for in vivo testing. As, in previous studies, the 30 μ g/mice dose had demonstrated a significant in vivo activity, that dose was selected for in vivo evaluation.

Example 9: *In vivo* comparison of murine and chimeric 224G11 Mabs on the NCI-H441 xenograft model

NCI-H441 is derived from papillary lung adenocarcinoma, expresses high levels of c-Met, and demonstrates constitutive phosphorylation of c-Met RTK.

To evaluate the *in vivo* effect of antibodies on the NCI-H441 xenograft model, six to eight weeks old athymic mice were housed in sterilized filter-topped cages, maintained in sterile conditions and manipulated according to French and European

guidelines. Mice were injected subcutaneously with 9×10^6 cells. Then, six days after cell implantation, tumors were measurable (approximately 100 mm^3), animals were divided into groups of 6 mice with comparable tumor size and treated first with a loading dose of $60 \mu\text{g}$ of antibody/mice and then twice a week with $30 \mu\text{g}/\text{dose}$ of each antibody to be tested. The mice were followed for the observation of xenograft growth rate. Tumor volume was calculated by the formula: $\pi (\text{Pi})/6 \times \text{length} \times \text{width} \times \text{height}$. Results described in Figure 13 demonstrate that the murine Mab devoided of agonist activity *in vivo* behave, as expected, as potent antagonist even at the low tested dose. In contrast to what observed with the murine Mab, the chimeric one displayed a very transient *in vivo* activity and tumor completely escaped to the treatment at D20 post cell injection. This experiment demonstrates clearly that the increase of *in vitro* agonist effect that resulted in a decrease of antagonist activity was also responsible for a significant *in vivo* loss of antagonist activity.

Example 10: Effect of the murine 224G11 Mab and of various chimeric and humanized versions of this antibody on HGF-induced proliferation of NCI-H441 cells *in vitro*

NCI-H441 cells from ATCC were routinely cultured in RPMI 1640 medium (Invitrogen Corporation, Scotland, UK), 10% FCS (Invitrogen Corporation), 1% L-Glutamine (Invitrogen Corporation). For proliferation assays, cells were split 3 days before use so that they were in the confluent phase of growth before plating. NCI-H441 cells were plated in 96-well tissue culture plates at a density of 3.75×10^4 cells/well in 200 μl of serum free medium (RPMI 1640 medium plus 1% L-Glutamine). Twenty four hours after plating, antibodies to be tested were added to NCI-H441 and incubated at 37°C for thirty minutes before adding HGF at a final concentration of 400 ng/ml (5 nM) for 142 additional hours. The dose range tested for each antibody is from 10 to 0.0097 $\mu\text{g}/\text{ml}$ (final concentration in each well). In this experiment, murine IgG1 Mab was added as a murine isotype control and as an agonist negative control. The tested antibodies were the following one: i) m224G11, ii) its human IgG1 chimeric forms respectively identified as [224G11] chim, [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim], [224G11] [TH7 chim] iii) its humanized IgG1 forms respectively described as [224G11] [Hz1], [224G11] [Hz2], [224G11] [Hz3]. Wells

plated with cells alone +/- HGF were also included. The 5D5 whole antibody from Genentech commercially available at the ATCC as an hybridoma cell line was introduced as a full agonist positive control and thereafter called m5D5. Then cells were pulsed with 0.25 μ Ci of [^{3} H]Thymidine (Amersham Biosciences AB, Uppsala, Sweden) for 7 hours and 30 minutes. The magnitude of [^{3} H]Thymidine incorporated in trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results are expressed as non transformed cpm data to better evaluate the potential intrinsic agonist activity that could occur with anti-c-Met Mabs when added alone to tumour cell.

Results described in Figure 14A demonstrated that as expected neither the isotype control nor the m224G11 displayed any agonist activity on NCI-H441 proliferation. The isotype control was without effect on HGF-induced cell proliferation whereas m224G11 showed a 66% inhibition when added at the final concentration of 10 μ g/ml. The m5D5 used as an agonist control showed, as expected, a full dose dependent agonist effect when added alone to the cells. As already observed, the [224G11] chim Mab displayed a significant dose- dependent agonist effect and, a decreased inhibitory activity of this chimeric form was observed: 19% instead of 66% for the murine form. When added alone, the 3 IgG1 humanized Mabs demonstrated dose dependent agonist effects compared to the m224G11 form. [224G11] [Hz1], [224G11] [Hz2] and [224G11] [Hz3] had comparable antagonist activities about 46, 30 and 35%. These activities are significantly lower than the one observed for m224G11. In Figure 14B, various IgG1 chimeric forms were tested. Compared to [224G11] chim form which displayed a dose-dependent agonist effect when added alone to NCI-H441 cells, the [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim], [224G11] [TH7 chim] forms were without significant intrinsic agonist effect. Their antagonist activity was higher than the one observed for the m224G11 Mab (57%) with inhibitions reaching 79, 78, 84 and 93% respectively for [224G11] [MH chim], [224G11] [MUP9H chim], [224G11] [MMCH chim] and [224G11] [TH7 chim].

Example 11: *In vitro* effect of various IgG1 humanized form of the 224G11 Mab

NCI-H441 cells from ATCC were routinely cultured in RPMI 1640 medium

(Invitrogen Corporation, Scotland, UK), 10% FCS (Invitrogen Corporation), 1% L-Glutamine (Invitrogen Corporation). For proliferation assays, cells were split 3 days before use so that they were in the confluent phase of growth before plating. NCI-H441 cells were plated in 96-well tissue culture plates at a density of 3.75×10^4 cells/well in 200 μ l of serum free medium (RPMI 1640 medium plus 1% L-Glutamine). Twenty four hours after plating, antibodies to be tested were added to NCI-H441 and incubated at 37°C for thirty minutes before adding HGF at a final concentration of 400 ng/ml (5 nM) for 142 additional hours. The dose range tested for each antibody is from 10 to 0.0097 μ g/ml (final concentration in each well). In this experiment, murine IgG1 Mab was added as a background negative control for agonist activity and the tested antibodies were the following one: i) m224G11, ii) its human IgG1 chimeric forms respectively identified as [224G11] chim, [224G11] [TH7 chim] iii) its humanized IgG1 forms respectively described as [224G11] [TH7 Hz1], [224G11] [TH7 Hz3]. Wells plated with cells alone +/- HGF were also included. The 5D5 whole antibody from Genentech commercially available at the ATCC as an hybridoma cell line was introduced as a full agonist positive control and thereafter called m5D5. Then cells were pulsed with 0.25 μ Ci of [^{3} H]Thymidine (Amersham Biosciences AB, Uppsala, Sweden) for 7 hours and 30 minutes. The magnitude of [^{3} H]Thymidine incorporated in trichloroacetic acid-insoluble DNA was quantified by liquid scintillation counting. Results are expressed as non transformed cpm data to better evaluate the potential intrinsic agonist activity that could occur with anti-c-Met Mabs when added alone to tumour cell.

Figure 15 showed that the m224G11 Mab displayed the usual inhibitory effect (74% inhibition). The chimeric IgG1 form [224G11] chim had as expected a dose dependent intrinsic agonist effect and a lower antagonist effect compared to the murine form: 33% versus 74% inhibition. The [224G11] [TH7 chim] had a very weak agonist activity in this experiment. However it displayed a high inhibitory effect (81 %) close to the one noticed for the murine Mab. The 2 humanized forms had no intrinsic agonist effect and had an antagonist activity close to the ones observed for the murine Mab or the [224G11] [TH7 chim] with respectively 67 and 76% inhibition for [224G11] [TH7 Hz1] and [224G11] [TH7 Hz3].

Example 12: *In vivo* comparison of murine, chimeric and humanized 224G11 Mabs bearing either the wild type or the TH7- engineered hinge (NCI-H441 xenograft model).

NCI-H441 is derived from papillary lung adenocarcinoma, expresses high levels
5 of c-Met, and demonstrates constitutive phosphorylation of c-Met RTK.

To evaluate the necessity of hinge engineering to save *in vivo* activity of the 224G11 murine antibody, six to eight weeks old athymic mice were housed in sterilized filter-topped cages, maintained in sterile conditions and manipulated according to French and European guidelines. Mice were injected subcutaneously with 9×10^6 NCI-
10 H441 cells. Then, six days after cell implantation, tumors were measurable (approximately 100 mm^3), animals were divided into groups of 6 mice with comparable tumor size and treated first with a loading dose of 2 mg of antibody/mice and then twice a week with a 1 mg/dose of each antibody to be tested. Ten antibodies were evaluated in this experiment including the m224G11, the chimeric form displaying the wild type
15 hinge (c224G11), the TH7-engineered chimeric form (224G11[TH7 chim]), three humanized form bearing the wild type hinge (224G11[IgG1 Hz1], 224G11[IgG1 Hz2] and 224G11[IgG1 Hz3]) and the three corresponding TH7-engineered forms (224G11[TH7 Hz1], 224G11[TH7 Hz2] and 224G11[TH7 Hz3]). Mice were followed for the observation of xenograft growth rate.

20 Tumor volume was calculated by the formula: $\pi (\text{Pi})/6 \times \text{length} \times \text{width} \times \text{height}$.

Results described in figure 16 demonstrate that the murine Mab devoid of any agonist activity *in vitro* behave, as expected, as potent *in vivo* antagonist. In contrast to what observed with the murine Mab, both chimeric and humanized forms bearing the
25 wild type hinge displayed only a very transient *in vivo* activity. In any cases the substitution of the wild type hinge by the TH7-engineered one resulted in a complete restoration of the *in vivo* activity observed with murine antibodies. This experiment demonstrates clearly that the increase of *in vitro* agonist effect that resulted in a decrease of antagonist activity was also responsible of a significant *in vivo* loss of
30 antagonist activity. It also demonstrates that the use of a TH7-engineered region instead of the wild type one is needed for keeping the *in vivo* properties of the murine Mab.

Example 13: Effect of m224G11 and its humanized form h224G11 on c-Met downregulation *in vitro*

In the following examples, for the avoidance of doubt, the expression h224G11 refers to the humanized form 224G11 [TH7 Hz3] of the antibody of the invention.

5 Two cell lines have been selected to address the activity of anti-c-Met antibodies on c-Met receptor degradation. A549 (#HTB-174) and NCI-H441 (#CCL-185) are two NSCLC cell lines from the ATCC collection. NCI-H441 cells were seeded in RPMI 1640 + 1% L-glutamine + 10% heat-inactivated FBS, at 3×10^4 cells/cm² in six-well plates for 24 h at 37°C in a 5% CO₂ atmosphere. A549 cells were seeded in F12K +
10 10% heat-inactivated FBS, at 2×10^4 cells/cm² in six-well plates for 24 h at 37°C in a 5% CO₂ atmosphere.

Then, cells were washed twice with phosphate buffer saline (PBS) before being serum-starved for 24 additional hours. Anti-c-Met antibodies (10 µg/ml), irrelevant mIgG1(10 µg/ml), or HGF (400 ng/mL) were added in serum-free DMEM medium at
15 37°C. After either 4 hours or 24 hours of incubation, the medium was gently removed and cells washed twice with cold PBS. Cells were lysed with 500 µL of ice-cold lysis buffer [50 mM Tris-HCl (pH 7.5); 150 mM NaCl; 1% Nonidet P40; 0.5% deoxycholate; and 1 complete protease inhibitor cocktail tablet plus 1% antiphosphatases]. Cell lysates were shaken for 90 min at 4°C and cleared at 15 000 rpm for 10 minutes. At this stage,
20 cell lysates could be stored at -20°C until needed for western blot analysis. Protein concentration was quantified using BCA. Whole cell lysates (5µg in 20 µl) were separated by SDS-PAGE and transferred to nitrocellulose membrane. Membranes were saturated for 1 h at RT with TBS-Tween 20 0.1%(TBST); 5% non-fat dry milk and probed with anti-c-Met antibody (dilution 1/1000) overnight at 4°C in TBST-5% non-
25 fat dry milk. Antibodies were diluted in tris-buffered saline-0.1% tween 20 (v/v) (TBST) with 1 % non-fat dry milk. Then, membranes were washed with TBST and incubated with peroxylase-conjugated secondary antibody (dilution 1:1000) for 1 h at RT. Immunoreactive proteins were visualized with ECL (Pierce # 32209). After c-Met visualization, membranes were washed once again with TBST and incubated for 1 h at
30 RT with mouse anti-GAPDH antibody (dilution 1/200 000) in TBST-5% non-fat dry milk. Then, membranes were washed in TBST and incubated with peroxylase-conjugated secondary antibodies, for 1 h at RT. Membranes were washed and GAPDH

was revealed using ECL. Band intensity was quantified by densitometry.

Results presented in figures 19A and 20A demonstrated that both m224G11 and h224G11 are able to significantly downregulate c-Met, in a dose-dependant way, in both A549 and NCI-H441 cell lines. The downregulation is already significant after a 4 hour incubation time and still increased at 24 hour. Histograms presented in figures 19A and 20A corresponds to mean values of respectively 4 and 3 independent experiments. Western blot images corresponding to one significant experiment were included in figures 19B and 20B.

Example 14: Effect of m224G11 and its humanized form h224G11 on c-Met shedding *in vitro*

Soluble shedded forms of the c-Met receptor occur naturally in the serum of mice xenografted with human tumor or in serum of human patient carrying tumors expressing c-Met. Moreover, antibodies directed against c-Met such as the DN30 Mab, are described as shedding inducers of c-Met in *in vitro* experiments. To determine whether the m224G11 has such a property, cells were seeded in six-well plates in 10% FCS medium. When they reached approximately 80% confluence, medium was removed and fresh complete culture medium +/- compounds to be tested was added. Cells were incubated 72 additional hours with either m224G11, an isotype control mIgG1 or PBS. PMA (phorbol myristate acetate) was introduced as a shedding inducer. HGF was also tested on cells to determine the impact of c-Met ligand on natural occurring shedding. Then supernatants were collected and filtered on 0.2 μ m before use in an ELISA test which soluble forms of c-Met were captured with an anti-c-Met antibody that does not recognize the same epitope as either m224G11 or the c11E1 (Figure 21). Moreover, cells from each well were washed once with PBS and lysed to determine protein concentration. For the ELISA, 224D10 was used as a capture antibody and after plate saturation, filtered supernatants from six well plates were added in the ELISA test. A monomeric c-Met form was used as a positive control. After supernatant incubation, plates were washed to remove the unbound c-Met and c11E1 was used to detect c-Met captured by the 224G11 Mab. The revelation of the test was finally performed by addition of an HRP-conjugated anti-hFc polyclonal antibody.

Results shown in figure 22 indicate that a natural shedding of c-Met occurred

when cells were cultured for 72 hours in vitro. No effect of the mIgG1 was observed. However, the addition of m224G11 seemed to inhibit c-Met shedding. These results were confirmed for 3 other cells lines (Hs746T, EBC1 and MKN45) in figure 23. In that second experiment, the PMA, added as a positive shedding inducer, increased significantly, as expected, c-Met shedding at least in 2 cell lines (Hs746T and MKN45). Finally, in a third experiment (Figure 24), HGF was introduced as a control. No additional shedding was induced by HGF compared as cells alone or cells + mIgG1. Once again, a significant inhibition of c-Met shedding was observed with m224G11.

10 **Example 15: Intrinsic effect of h224G11 Ab on various cell lines**

In previous experiments described in this patent, it has been demonstrated that in contrast to what was observed with other antibodies such as 5D5, the m224G11 and its humanized form h224G11 do not display significant intrinsic activity tumor cell lines. To extend this property to other cell lines, western blot and phospho-ELISA experiments were performed with the antibody alone, added for various times, on a set of cancer cell lines, with variable levels of c-Met expression, including Hs746T, NCI-H441, Hs578T, NCI-H125, T98G, MDA-MB-231, PC3. The same test was also performed in a normal cell: HUVEC.

Method for the phospho cMet ELISA assay was already described in example 5 of the present patent application. For the western analysis, protein lysates were made from pelleted cells by incubation in lysis buffer with proteases and phosphatase inhibitors [10nM Tris (pH 7.4), 150 mM NaCl, 1mM EDTA, 1 mM EGTA, 0.5% Nonidet P40, 100 mM sodium fluoride, 10 mM sodium pyrophosphate, 2 mM sodium orthovanadate, 2 mM PMSF, 10 mg/ml leupeptin, 10 mg/ml aprotinin] at 4°C. Protein lysates were cleared of cellular debris by centrifugation, resolved by electrophoresis on 8% SDS-PAGE gels, and electrotransferred to a nitrocellulose membrane. For c-Met experiments, lysates were immunoprecipitated for specific protein of interest before electrophoresis and transfer.

Results presented in figures 25 to 32 demonstrate once again that no intrinsic activity of the h224G11 antibody was observed in the tested cells.

Example 16: *In vivo* comparison of the murine wild type 224G11 with a chimeric hinge-engineered 224G11 form described as 224G11[C2D5-7] (NCI-H441 xenograft model)

5 NCI-H441 is derived from papillary lung adenocarcinoma, expresses high levels of c-Met, and demonstrates constitutive phosphorylation of c-Met RTK.

To evaluate the necessity of hinge engineering to save *in vivo* activity of the 224G11 murine antibody, six to eight weeks old athymic mice were housed in sterilized filter-topped cages, maintained in sterile conditions and manipulated according to
10 French and European guidelines. Mice were injected subcutaneously with 9×10^6 NCI-H441 cells. Then, six days after cell implantation, tumors were measurable (approximately 100 mm^3), animals were divided into groups of 6 mice with comparable tumor size and treated first with a loading dose of 2 mg of antibody/mice and then twice a week with a 1 mg/dose of each antibody to be tested. Mice were followed for the
15 observation of xenograft growth rate. Tumor volume was calculated by the formula: $\pi (\text{Pi})/6 \times \text{length} \times \text{width} \times \text{height}$. Results described in figure 33 demonstrate that the murine Mab devoid of any agonist activity *in vitro* behave, as expected, as a potent *in vivo* antagonist. As suggested by the results obtained *in vitro*, in phosphorylation assays, the c224G11[C2D5-7] hinge-engineered antibody, that did not display a
20 significant agonist effect, demonstrate a strong *in vivo* activity, comparable to the one of the m224G11 on the NCI-H441 xenograft model.

Example 17: Evaluation of h224G11 in an ADCC test

As h224G11 is of IgG1 isotype, ADCC could be part of its *in vivo* efficacy in
25 human. An *in vitro* [^{51}Cr] release cytotoxicity assay was performed using either Hs746T or NCI-H441 cells as target cells and NK cells purified from human peripheral blood mononuclear lymphocytes.

Briefly, one million Hs746T or NCI-H441 target cells were incubated with or without 20 μg of h224G11 Ab in presence of 100 μCi of $^{51}\text{Chromium}$ (Perkin Elmer)
30 for 1 hr. Then, 4×10^3 cells were plated with an increasing number of human natural killer (NK) cells isolated from peripheral blood mononuclear cells (PBMC) using a negative selection (Stemcell Technologies). Cells were incubated together for 4

additional hours at 37°C. Percent of cell lysis was calculated following the formula:
[(experimental ^{51}Cr release - spontaneous ^{51}Cr release)/(full ^{51}Cr release - spontaneous
 ^{51}Cr release)] X 100. Spontaneous release represents the counts obtained when the
target cells were cultured in absence of natural killer cells. Full release represents the
counts obtained when the target cells were lysed with 1% Triton X-100. h224G11
significantly enhanced lysis of both Hs746T (Figure 34A) and NCI-H441 (Figure 34B)
cells by 62.9% and 63.2%, respectively, at a ratio NK/Target cells of 100.

Example 18: Immunohistochemical Studies (IHC)

Procedures of Paraffin Embedded Tumors IHC Staining: 8 to 12 μM sections of
frozen tumor were and immediately fixed in pre cooled acetone -20°C for 3 minutes.
Slides were then cooled at room temperature for 30 minutes to 1 hour. After 2 washes in
PBS the Endogenous peroxidase activity was blocked using Peroxidase Blocking
Reagent (Dako K4007) for five minutes. Sections were washed with PBS and incubated
in avidin/biotin blocking reagent (Dako X0590) just before saturation of the non
specific sites in PBS-BSA 4% for 30 minutes at room temperature. Then, slides were
incubated with the biotinylated h224G11 (50 to 10 $\mu\text{g}/\text{ml}$) or human biotinylated
IgG1/kappa (50 to 10 $\mu\text{g}/\text{ml}$, the Binding Site) as negative control 2 hours at room
temperature.

Sections were washed with PBS and incubated with Streptavidin-peroxydase
complex universal (Dako K0679) for 30 to 45 minutes. 3-Amino-9-Ethylcarbazole was
used for development of a red reaction product (Sigma). The slides were immersed in
hematoxylin for 4 minutes to counterstain (Dako S3309).

Results are represented in Figures 35A-35C.

h224G11 differentially stains the cell membrane of various tumor types. In this
immunohistochemistry procedure, the red reaction product correlates to positive
staining of the cell membrane and lack of red reaction product correlates to negative
staining and no visualization of the cell membrane. The IgG control, human IgG1/kappa
is an isotype matched control.

**Example 19: *In vivo* activity of anti-c-Met Mabs on three cell lines amplified
for c-Met and xenografted in mice.**

Hs746T (ATCC) and MKN45 (JCRB), two stomach carcinoma cells, and, EBC-1 (JCRB), a lung squamous cell carcinoma, are 3 cell lines that over-express c-Met as shown by FACS analysis in Figures 36A, 37A and 38A. These cell lines are also described in the literature as amplified cell lines for c-Met. To confirm this latter point, c-Met gene amplification was determined by quantitative real-time PCR using $\Delta\Delta C_t$ method. Fifty ng of genomic DNA from each indicated cell line, extracted using the QIAampTM DNA midi kit (Qiagen, Les Ulis, FR) was used. The reference gene was DNA topoisomerase III alpha located on chromosome 17. Primers for c-Met gene were located on exon 14. The corresponding primers for both genes are those described by Smolen et al. (PNAS, 2006, 103:2316-2321) and yielded similar PCR efficiencies. The QPCR protocol consists in a 10 min cycle at 95°C, followed by 40 successive cycles of 15 sec at 95°C/45 sec at 95°C. Data were first normalized to the reference gene, and then each cell line was normalized to MCF-7 cells containing a single copy of c-Met gene per haploid genome. The results presented in figures 36B, 37B and 38B demonstrate that the 3 cell lines display a significant amplification of c-Met ranking from 5 to 10-fold gene amplification.

As it has been described that such cells are constitutively activated for c-Met, a Western blot assay was performed to determine the level of c-Met phosphorylation of each cell line either in absence or in presence of HGF (50 ng/ml). For that purpose cells were cultured overnight in serum-free medium and then treated or not with 50 ng/ml of HGF for 2 min. Cells were lysed with lysis buffer supplemented with protease inhibitors. Cell lysates were incubated at 4°C for 90 min, then centrifuged and supernatants were collected. Protein concentrations were determined using the BCA protein assay (Pierce) and samples were immunoblotted. Antibodies used include the anti-c-Met antibody, clone 25H2 from Cell Signaling and the anti-P-c-Met Y1230, 1234, 1235 from Biosource. Immunoblotting with the phospho-specific c-Met antibody against Y1230/1234/1235 showed strong baseline phosphorylation of the receptor in each amplified cell line (Figures 36C, 37C and 38C). No additional phosphorylation is observed when HGF was added to these cells. This last observation confirms that c-Met was activated in an HGF-independent way in Hs746T, MKN45 and EBC-1 cell lines.

In order to test our anti-c-Met antibodies *in vivo* on ligand-independent cell lines, Hs746T, MKN45 and EBC-1 cells were routinely cultured in complete DMEM,

RPMI 1640 or EMEM medium respectively. Cells were split two days before engraftment so that they were in exponential phase of growth. Seven million cells were engrafted to either SCID (for Hs746T and EBC-1) or CD1 Swiss nude mice (for MKN45). Five to seven days after implantation, tumors were measurable and animals
5 were divided into groups of 6 mice with comparable tumor size. Mice were treated i.p. with a loading dose of 2 mg of h224G11 Mab /mouse and then twice a week with 1 mg of antibody/mouse. Tumor volume was measured twice a week and calculated by the formula: $\pi/6 \times \text{length} \times \text{width} \times \text{height}$. Statistical analysis was performed at each measured time using a Mann-Whitney test.

10 This experiment demonstrates that h224G11 reduced significantly tumor growth of Hs746T, MKN45 and EBC1 (Figures 36D, 37D and 38D), indicating that the humanized antibody is able to inhibit tumor growth of gastric and lung ligand-independent cell lines. Before these results were obtained, it would not have been apparent how the presence of the modified hinge in this humanized antibody would
15 affect its ligand-independent inactivation of c-Met, because the altered hinge also affected its agonist/antagonist properties.

Example 20: *Ex vivo* analysis of h224G11 activity on the Hs746T xenograft model in SCID mice.

20 In order to determine more precisely mechanisms of action that resulted in Hs746T tumor xenograft inhibition, the effect of 3 i. p. injections of h224G11 was evaluated on tumor mitotic index (Ki67), apoptosis (activated caspase 3) and c-Met expression using immunohistochemistry (IHC) methods. Three mice were analyzed for each condition. A significant down regulation of c-Met was observed on treated tumors
25 compared to control one (Fig.39C and D respectively). Panel 39A and B corresponds to isotypes controls. In addition a significant decrease in Ki67 levels (Figures 39E and F) and an increase in activated caspase 3 staining (Figures 39G and H) was noticed on all treated tumors indicating that h224G11 was able to inhibit significantly cell proliferation and to induce cell apoptosis within the tumor.

0-1	Form PCT/RO/134 (SAFE) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared Using	PCT Online Filing Version 3.5.000.225 MT/FOP 20020701/0.20.5.20
0-2	International Application No.	
0-3	Applicant's or agent's file reference	358326D27117

1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	16
1-2	line	27
1-3	Identification of deposit	
1-3-1	Name of depositary institution	CNCM Collection nationale de cultures de micro-organismes
1-3-2	Address of depositary institution	Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cedex 15, France
1-3-3	Date of deposit	14 March 2007 (14.03.2007)
1-3-4	Accession Number	CNCM I-3731
1-5	Designated States for Which Indications are Made	All designations

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	YES
0-4-1	Authorized officer	De Jong-de Koster, Marion

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	14 June 2011 (14.06.2011)
0-5-1	Authorized officer	WAGNER, Nathalie

Claims

1. A monoclonal antibody, or a divalent functional fragment thereof containing at least its 6 CDRs and its hinge region, that inhibits c-Met dimerization, said antibody comprising a heavy chain comprising the amino acid sequence SEQ ID No. 37 and a
5 light chain comprising the amino acid sequence SEQ ID No. 40, for use for the prevention or the treatment of a cancer characterized by overexpression of c-Met resulting from genic amplification of c-Met, and wherein said cancer is renal cell carcinoma or gastric cancer.
2. A composition comprising:
 - 10 • a compound consisting of a monoclonal antibody, or a divalent functional fragment thereof containing at least its 6 CDRs and its hinge region, that inhibits c-Met dimerization, said antibody comprising a heavy chain comprising the amino acid sequence SEQ ID No. 37 and a light chain comprising the amino acid sequence SEQ ID No. 40, and
 - 15 • (i) an excipient, (ii) a pharmaceutically acceptable vehicle, or (iii) an excipient and a pharmaceutically acceptable vehicle,

for use for the prevention or the treatment of a cancer characterized by overexpression of c-Met resulting from genic amplification of c-Met, and wherein said cancer is renal cell carcinoma or gastric cancer.
- 20 3. Use for the prevention or the treatment of a cancer characterized by overexpression of c-Met resulting from genic amplification of c-Met, wherein said cancer is renal cell carcinoma or gastric cancer, of a monoclonal antibody, or a divalent functional fragment thereof containing at least its 6 CDRs and its hinge region, that inhibits c-Met dimerization, said antibody comprising a heavy chain
25 comprising the amino acid sequence SEQ ID No. 37 and a light chain comprising the amino acid sequence SEQ ID No. 40.
4. Use for the prevention or the treatment of a cancer characterized by overexpression of c-Met resulting from genic amplification of c-Met, wherein said cancer is renal cell carcinoma or gastric cancer, of a composition comprising:
 - 30 • a compound consisting of a monoclonal antibody, or a divalent functional fragment thereof containing at least its 6 CDRs and its hinge region, that

inhibits c-Met dimerization, said antibody comprising a heavy chain comprising the amino acid sequence SEQ ID No. 37 and a light chain comprising the amino acid sequence SEQ ID No. 40, and

- (i) an excipient, (ii) a pharmaceutically acceptable vehicle, or (iii) an excipient and a pharmaceutically acceptable vehicle.

5

1/39




FIGURE 1

2/39




FIGURE 2A




FIGURE 2B

3/39




FIGURE 3A




FIGURE 3B

4/39




FIGURE 4A




FIGURE 4B

5/39




FIGURE 5A




FIGURE 5B

6/39




FIGURE 6A




FIGURE 6B

7/39




FIGURE 7A




FIGURE 7B

8/39




FIGURE 8A




FIGURE 8B

9/39




FIGURE 9A




FIGURE 9B

10/39




FIGURE 10A




FIGURE 10B

11/39





FIGURE 11

12/39




FIGURE 12




FIGURE 13

13/39




FIGURE 14A

14/39




FIGURE 14B

15/39




FIGURE 15




FIGURE 16




FIGURE 17A




FIGURE 17B

19/39






FIGURE 18

20/39

A**B****FIGURE 19**






FIGURE 20

22/39

ELISA**FIGURE 21****FIGURE 22**

23/39




FIGURE 23




FIGURE 24




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28




FIGURE 29




FIGURE 30




FIGURE 31




FIGURE 32

33/39




FIGURE 33

34/39




FIGURE 34

35/39




FIGURE 35




Figure 36




Figure 37




Figure 38

39/39

Figure 39

