



(86) Date de dépôt PCT/PCT Filing Date: 2004/10/28
(87) Date publication PCT/PCT Publication Date: 2005/05/26
(45) Date de délivrance/Issue Date: 2012/05/01
(85) Entrée phase nationale/National Entry: 2006/05/08
(86) N° demande PCT/PCT Application No.: EP 2004/012175
(87) N° publication PCT/PCT Publication No.: 2005/047278
(30) Priorité/Priority: 2003/11/11 (DE10352499.1)

(51) Cl.Int./Int.Cl. *C07D 401/04* (2006.01),
A61K 31/517 (2006.01), *A61P 31/12* (2006.01)


(72) Inventeurs/Inventors:
WUNBERG, TOBIAS, DE;
BAUMEISTER, JUDITH, BE;
JESKE, MARIO, DE;
NELL, PETER, DE;
NIKOLIC, SUSANNE, DE;
SUESSMEIER, FRANK, DE;
...

(73) Propriétaire/Owner:
AICURIS GMBH & CO. KG, DE

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : DIHYDROQUINAZOLINES II SUBSTITUEES

(54) Title: SUBSTITUTED DIHYDROQUINAZOLINES II




(I)

(57) Abrégé/Abstract:

The invention relates to substituted dihydroquinazolines and to a method for the production thereof, the use thereof for treating and/or preventing diseases and for producing drugs for treating and/or preventing diseases, in particular for the use of the inventive dihydroquinazolines in the form of antiviral agents, in particular against cytomegaloviruses. (see formula I)

(72) **Inventeurs(suite)/Inventors(continued)**: ZIMMERMANN, HOLGER, DE; GROSSER, ROLF, DE; HEWLETT, GUY, DE;
KELDENICH, JOERG, DE; LANG, DIETER, DE; HENNINGER, KERSTIN, DE

The invention relates to substituted dihydroquinazolines and to a method for the production thereof, the use thereof for treating and/or preventing diseases and for producing drugs for treating and/or preventing diseases, in particular for the use of the inventive dihydroquinazolines in the form of antiviral agents, in particular against cytomegaloviruses.



(1).

Substituted dihydroquinazolines II

The invention relates to substituted dihydroquinazolines and to processes for their preparation, to their use for the treatment and/or prophylaxis of diseases and to their
5 use for preparing medicaments for the treatment and/or prophylaxis of diseases, in particular for use as antiviral agents, especially against cytomegaloviruses.

The synthesis of dihydroquinazolines is described in Saito T. et al., Tetrahedron Lett., 1996, 37, 209-212, and in Wang F. et al., Tetrahedron Lett., 1997, 38, 8651-
10 8654. Furthermore, Y.S. Lee et al., *Bioorg. Med. Chem. Lett.*, **2004**, 14, 3379-3384, have described dihydroquinazolines as calcium channel inhibitors.


Agents with antiviral activity and a different structure are available on the market; however, the therapies currently available with ganciclovir, valganciclovir, foscarnet
15 and cidofovir are associated with severe side effects, for example nephrotoxicity, neutropenia or thrombocytopenia. In addition, it is always possible for resistance to develop. Novel agents for an effective therapy are therefore desirable.

One object of the present invention is therefore to provide novel compounds having
20 the same or improved antiviral action for the treatment of viral infective diseases in humans and animals.

It has been found, surprisingly, that the substituted dihydroquinazolines described in the present invention have high antiviral activity.

25

The invention provides compounds of the formula



in which

5 R^1 , R^2 and R^3 independently of one another represent hydrogen, alkyl, alkoxy, carboxyl, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, trifluoromethyl, halogen, cyano, hydroxyl or nitro,

10 R^4 and R^5 independently of one another represent hydrogen, alkyl, alkoxy, alkylthio, cyano, halogen, nitro, trifluoromethyl or trifluoromethoxy,

R^6 represents alkyl, cyano, halogen, nitro or trifluoromethyl,

R^7 and R^8 independently of one another represent hydrogen, halogen, alkyl or alkoxy and

15 R^9 represents aryl or 1,3-benzodioxol-5-yl, where aryl and 1,3-benzodioxol-5-yl may be substituted by 1 to 3 substituents, where the substituents independently of one another are selected from the group consisting of alkoxy, alkylthio, carboxyl, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, trifluoromethyl, halogen, carbamoyl, 20 cyano, hydroxyl, amino, alkylamino, nitro and optionally hydroxyl-substituted alkyl,

and their salts, their solvates and the solvates of their salts.

Compounds according to the invention are the compounds of the formula (I) and their salts, solvates and solvates of the salts, compounds mentioned hereinbelow as exemplary embodiment(s) and their salts, solvates and solvates of the salts, if the compounds mentioned hereinbelow, embraced by formula (I), are not already salts,
5 solvates and solvates of the salts.

The compounds according to the invention may, depending on their structure, exist in stereoisomeric forms (enantiomers, diastereomers). The invention therefore relates to the enantiomers or diastereomers and respective mixtures thereof. The stereo-
10 isomerically pure constituents can be isolated in a known manner from such mixtures of enantiomers and/or diastereomers.

If the compounds according to the invention can exist in tautomeric forms, the present invention provides all tautomeric forms.

15

Salts in the context of the present invention are preferably physiologically acceptable salts of the compounds according to the invention. Also provided, however, are salts which for their part are not suitable for pharmaceutical applications but which can be used, for example, for isolating or purifying the compounds according to the
20 invention.

Physiologically acceptable salts of the compounds according to the invention include acid addition salts of mineral acids, carboxylic acids and sulphonic acids, for example salts of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric
25 acid, methanesulphonic acid, ethanesulphonic acid, toluenesulphonic acid, benzenesulphonic acid, naphthalenedisulphonic acid, acetic acid, trifluoroacetic acid, propionic acid, lactic acid, tartaric acid, malic acid, citric acid, fumaric acid, maleic acid and benzoic acid.

30 Physiologically acceptable salts of the compounds according to the invention also include salts of customary bases, such as, by way of example and preferably, alkali metal salts (for example sodium and potassium salts), alkaline earth metal salts (for example calcium and magnesium salts) and ammonium salts derived from ammonia

or organic amines having 1 to 16 carbon atoms, such as, by way of example and preferably, ethylamine, diethylamine, triethylamine, ethyldiisopropylamine, monoethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, dimethylaminoethanol, procaine, dibenzylamine, N-methylmorpholine, arginine, lysine, ethylenediamine and N-methylpiperidine.

In the context of the invention, solvates are those forms of the compounds according to the invention which form a complex in the solid or liquid state by coordination with solvent molecules. Hydrates are a special form of solvates in which the coordination takes place with water.

For the purposes of the present invention, unless specified otherwise, the substituents have the following meanings:

Alkyl per se and "alk" and "alkyl" in alkoxy, alkylthio, alkylamino, alkylcarbonyl and alkoxy carbonyl represent a straight-chain or branched alkyl radical having generally 1 to 6 ("C₁-C₆-alkyl"), preferably 1 to 4, particularly preferably 1 to 3, carbon atoms, by way of example and preferably methyl, ethyl, n-propyl, isopropyl, tert-butyl, n-pentyl and n-hexyl.

Alkoxy represents, by way of example and preferably, methoxy, ethoxy, n-propoxy, isopropoxy, tert-butoxy, n-pentoxy and n-hexoxy.

Alkylthio represents, by way of example and preferably, methylthio, ethylthio, n-propylthio, isopropylthio, tert-butylthio, n-pentylthio and n-hexylthio.

Alkylcarbonyl represents, by way of example and preferably, acetyl and propanoyl.

Alkylamino represents an alkylamino radical having one or two alkyl substituents (chosen independently of one another), by way of example and preferably methylamino, ethylamino, n-propylamino, isopropylamino, tert-butylamino, n-pentylamino, n-hexylamino, N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, N-methyl-N-n-propylamino, N-isopropyl-N-n-propylamino, N-tert-butyl-

N-methlamino, *N*-ethyl-*N*-*n*-pentylamino and *N*-*n*-hexyl-*N*-methlamino. C₁-C₃-Alkylamino is, for example, a monoalkylamino radical having 1 to 3 carbon atoms or is a dialkylamino radical having 1 to 3 carbon atoms each per alkyl substituent.

- 5 Alkoxycarbonyl represents, by way of example and preferably, methoxycarbonyl, ethoxycarbonyl, *n*-propoxycarbonyl, isopropoxycarbonyl, *tert*-butoxycarbonyl, *n*-pentoxycarbonyl and *n*-hexoxycarbonyl.

- 10 Aryl represents a mono- to tricyclic aromatic carbocyclic radical having generally 6 to 14 carbon atoms; by way of example and preferably phenyl, naphthyl and phenanthrenyl.

Halogen represents fluorine, chlorine, bromine and iodine.

- 15 A symbol * on a carbon atom means that the compound, with respect to the configuration at this carbon atom, is present in enantiomerically pure form which, for the purposes of the present invention, is to be understood as meaning an enantiomeric excess of more than 90% (> 90% ee).

- 20 In the context of the present invention, preference is given to compounds of the formula (I)

in which

- 25 R¹, R² and R³ independently of one another represent hydrogen, fluorine, chlorine, cyano, hydroxyl or aminocarbonyl,

R⁴ and R⁵ independently of one another represent hydrogen, fluorine, alkyl or alkoxy,

30

R⁶ represents chlorine, nitro, trifluoromethyl, methyl, isopropyl or *tert*-butyl,

- R^7 and R^8 independently of one another represent hydrogen or C_1 - C_3 -alkyl and
- R^9 represents phenyl or 1,3-benzodioxol-5-yl, where phenyl may be substituted by 1 to 3 substituents, where the substituents independently of one another are selected from the group consisting of C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, carboxyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, trifluoromethyl, fluorine, chlorine, bromine, cyano, hydroxyl, amino, C_1 - C_6 -alkylamino and nitro,
- and their salts, their solvates and the solvates of their salts.
- In the context of the present invention, preference is given to compounds of the formula (I)
- in which
- R^1 , R^2 and R^3 independently of one another represent hydrogen, methyl, fluorine, chlorine, cyano, hydroxyl or aminocarbonyl,
- R^4 and R^5 independently of one another represent hydrogen, fluorine, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy,
- R^6 represents chlorine, nitro, trifluoromethyl, methyl, isopropyl or tert-butyl,
- R^7 and R^8 independently of one another represent hydrogen or C_1 - C_3 -alkyl and
- R^9 represents phenyl or 1,3-benzodioxol-5-yl, where phenyl may be substituted by 1 to 3 substituents, where the substituents independently of one another are selected from the group consisting of C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, carboxyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, trifluoromethyl, fluorine, chlorine, bromine, cyano, hydroxyl, amino, C_1 - C_6 -alkylamino and nitro,

and their salts, their solvates and the solvates of their salts.

In the context of the present invention, preference is also given to compounds of the formula (I)

5

in which

R^1 and R^2 represent hydrogen,

10 R^3 represents fluorine,

R^4 and R^5 independently of one another represent hydrogen, fluorine or alkoxy,

R^6 represents trifluoromethyl,

15

R^7 and R^8 represent hydrogen, and

R^9 represents phenyl, where phenyl may be substituted by 1 or 2
substituents, where the substituents independently of one another are
20 selected from the group consisting of methyl, methoxy, fluorine and
chlorine,

and their salts, their solvates and the solvates of their salts.

25 In the context of the present invention, preference is also given to compounds of the formula (I)

in which

30 R^1 and R^2 are hydrogen,

R^3 is fluorine,

R^4 and R^5 independently of one another are hydrogen, fluorine or methoxy,

R^6 is trifluoromethyl,

5 R^7 and R^8 are hydrogen and

R^9 is phenyl, where phenyl may be substituted by 1 or 2 substituents, where the substituents independently of one another are selected from the group consisting of methyl, methoxy, ethoxy, fluorine and chlorine,

10

and their salts, their solvates and the solvates of their salts.

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^1 and R^2 represent hydrogen.

15

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^3 is attached to the carbon atom in position 6 or position 8 of the quinazoline skeleton.

20 In the context of the present invention, preference is also given to compounds of the formula (I) in which R^3 is attached to the carbon atom in position 8 of the quinazoline skeleton.

25 In the context of the present invention, preference is also given to compounds of the formula (I) in which R^3 represents fluorine, in particular fluorine which is attached to the carbon atom in position 8 of the quinazoline skeleton.

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^4 and R^5 represent hydrogen.

30

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^4 represents hydrogen and R^5 represents fluorine or alkoxy.

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^4 represents hydrogen and R^5 represents methoxy.

5 In the context of the present invention, preference is also given to compounds of the formula (I) in which R^6 represents trifluoromethyl.

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^6 represents methyl, isopropyl or tert-butyl.

10 In the context of the present invention, preference is also given to compounds of the formula (I) in which R^6 represents isopropyl or tert-butyl.

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^7 and R^8 represent hydrogen.

15

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^9 represents phenyl, where phenyl may be substituted by 1 or 2 substituents, where the substituents independently of one another are selected from the group consisting of methyl, methoxy, fluorine and chlorine.


20

In the context of the present invention, preference is also given to compounds of the formula (I) in which R^9 represents phenyl, where phenyl is substituted by fluorine in the para-position to the point of attachment to the piperidine ring.

25 In the context of the present invention, preference is also given to compounds of the formula (I) in which R^9 represents phenyl, where phenyl is substituted by chlorine, methyl or methoxy in the meta-position to the point of attachment to the piperidine ring.

30 In the context of the present invention, preference is also given to compounds of the formula (I) in which R^9 represents phenyl, where phenyl is substituted by methyl in the meta-position to the point of attachment to the piperidine ring and by fluorine in the para-position to the point of attachment to the piperidine ring.

The invention furthermore provides a process for preparing the compounds of the formula (I), where compounds of the formula



5

in which

$R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8$ and R^9 are as defined above, and

10

R^{10} represents alkyl, preferably methyl or ethyl,

are reacted with bases.


15 The reaction is generally carried out in inert solvents, preferably in a temperature range of from room temperature to the reflux of the solvents at atmospheric pressure.

Bases are, for example, alkali metal hydroxides, such as sodium hydroxide, lithium hydroxide or potassium hydroxide, or alkali metal carbonates, such as caesium carbonate, sodium carbonate or potassium carbonate, if appropriate in aqueous
20 solution; preference is given to sodium hydroxide in water.

Inert solvents are, for example, halogenated hydrocarbons, such as ethers, such as 1,2-dimethoxyethane, dioxane, tetrahydrofuran, glycol dimethyl ether or diethylene
25 glycol dimethyl ether, alcohols, such as methanol, ethanol, n-propanol, isopropanol,

n-butanol or tert-butanol, or of mixtures of solvents; preference is given to dioxane or tetrahydrofuran.

The compounds of the formula (II) are known or can be prepared by reacting
5 compounds of the formula




in which

R^1 , R^2 , R^3 , R^7 , R^8 , R^9 and R^{10} are as defined above,

10

with compounds of the formula



in which

15

R^4 , R^5 and R^6 are as defined above,

in the presence of phosphorus oxychloride.


20 The reaction is generally carried out in inert solvents, preferably in a temperature range of from 50°C to the reflux of the solvents at atmospheric pressure.

Inert solvents are, for example, hydrocarbons, such as benzene, xylene, toluene, hexane, cyclohexane or mineral oil fractions; preference is given to toluene.

- 5 Alternatively, the compounds of the formula (II) can be prepared in a two-step synthesis process. In the first step, the compounds of the formula (III) are heated with phosphorus oxychloride in an inert solvent, preferably toluene, under reflux at atmospheric pressure. The solvent is removed. In the second step, the compounds obtained in this manner are reacted with compounds of the formula (IV) in an inert
10 solvent, preferably toluene, likewise under reflux at atmospheric pressure.

The compounds of the formula (IV) are known or can be synthesized by known processes from the appropriate starting materials.


- 15 The compounds of the formula (III) are known or can be prepared by reacting compounds of the formula



in which

- 20 R^1 , R^2 , R^3 and R^{10} are as defined above,

with compounds of the formula



in which

R^7 , R^8 and R^9 are as defined above, and

5

R^{11} represents halogen, preferably chlorine, bromine or iodine, or hydroxyl.

In the case that R^{11} represents hydroxyl,

10 the reaction is generally carried out in inert solvents, in the presence of customary condensing agents, if appropriate in the presence of a base, preferably in a temperature range of from room temperature to 50°C at atmospheric pressure.

Inert solvents are, for example, halogenated hydrocarbons, such as methylene
 15 chloride, trichloromethane, carbon tetrachloride, trichloroethane, tetrachloroethane, 1,2-dichloroethane or trichloroethylene, ethers, such as diethyl ether, methyl tert-butyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, glycol dimethyl ether or diethylene glycol dimethyl ether, hydrocarbons, such as benzene, xylene, toluene, hexane, cyclohexane or mineral oil fractions, or carboxamides, such as
 20 dimethylformamide or dimethylacetamide, alkylnitriles, such as acetonitrile, or heteroaromatic compounds, such as pyridine, or ethyl acetate; preference is given to tetrahydrofuran, 1,2-dichloroethane or methylene chloride.

Customary condensing agents are, for example, carbodiimides, such as, for example,
 25 N,N'-diethyl-, N,N'-dipropyl-, N,N'-diisopropyl-, N,N'-dicyclohexylcarbodiimide, N-(3-dimethylaminoisopropyl)-N'-ethylcarbodiimide hydrochloride (EDC), N-cyclohexylcarbodiimide-N'-propyloxymethyl-polystyrene (PS-carbodiimide) or carbonyl compounds, such as carbonyldiimidazole, or 1,2-oxazolium compounds, such as 2-ethyl-5-phenyl-1,2-oxazolium 3-sulphate, or 2-tert-butyl-5-methyl-

isoxazolium perchlorate, or acylamino compounds, such as 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline, or propanephosphonic anhydride, or isobutyl chloroformate, or bis(2-oxo-3-oxazolidinyl)phosphoryl chloride or benzo-
5 triazolyloxytri(dimethylamino)phosphonium hexafluorophosphate, or O-(benzo-
triazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), 2-(2-oxo-1-(2H)-pyridyl)-1,1,3,3-tetramethyluronium tetrafluoroborate (TPTU) or O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU),
or 1-hydroxybenzotriazole (HOBt), or benzotriazol-1-yloxy-
tris(dimethylamino)phosphonium hexafluorophosphate (BOP), or mixtures of these.

10

Bases are, for example, alkali metal carbonates, such as, for example, sodium carbonate or potassium carbonate, or sodium bicarbonate or potassium bicarbonate, or organic bases, such as trialkylamines, for example triethylamine, N-methylmorpholine, N-methylpiperidine, 4-dimethylaminopyridine or
15 diisopropylethylamine.

Particular preference is given to the combination of N-(3-dimethylaminoisopropyl)-N'-ethylcarbodiimide hydrochloride (EDC), 1-hydroxybenzotriazole (HOBt) and triethylamine and dimethylformamide or carbonyldiimidazole in 1,2-dichloroethane.

20

In the case that R¹¹ represents halogen,

the reaction is generally carried out in inert solvents, if appropriate in the presence of a base, preferably in a temperature range of from 0°C to 50°C at atmospheric
25 pressure.

Inert solvents are, for example, halogenated hydrocarbons, such as methylene chloride, trichloromethane, carbon tetrachloride, trichloroethane, tetrachloroethane, 1,2-dichloroethane or trichloroethylene, ethers, such as diethyl ether, methyl tert-
30 butyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, glycol dimethyl ether or diethylene glycol dimethyl ether, hydrocarbons, such as benzene, xylene, toluene, hexane, cyclohexane or mineral oil fractions, or carboxamides, such as dimethylformamide or dimethylacetamide, alkylnitriles, such as acetonitrile, or

heteroaromatic compounds, such as pyridine, or ethyl acetate; preference is given to tetrahydrofuran, dioxane or methylene chloride.


Bases are, for example, alkali metal carbonates, such as caesium carbonate, sodium
5 carbonate or potassium carbonate, alkali metal acetates, such as sodium acetate, or other bases, such as triethylamine or diisopropylethylamine, preferably diisopropylethylamine or triethylamine.

The compounds of the formula (VI) are known or can be synthesized by known
10 processes from the appropriate starting materials.

The compounds of the formula (V) are known or can be prepared by known processes from the appropriate starting materials, for example by a Heck reaction or a Wittig-Horner reaction according to the synthesis schemes below:


15

Heck reaction:



Wittig-Horner reaction:

20




The starting materials required for this purpose are known or can be synthesized by known processes from the appropriate starting materials.

The preparation of the compounds according to the invention can be illustrated by the synthesis scheme below.

Synthesis scheme:

5



The compounds of the general formula (I) according to the invention show a surprising range of effects which could not have been predicted. They show an antiviral effect on representatives of the group of the Herpes viridae (Herpes viruses), especially on cytomegaloviruses (CMV), in particular on the human cytomegalovirus (HCMV).

Areas of indication which may be mentioned by way of example are:

- 1) Treatment and prophylaxis of HCMV infections in AIDS patients (retinitis, pneumonitis, gastrointestinal infections).

- 2) Treatment and prophylaxis of cytomegalovirus infections in bone-marrow and organ transplantation patients who develop often life-threatening HCMV pneumonitis or encephalitis, and gastrointestinal and systemic HCMV infections.
- 5
- 3) Treatment and prophylaxis of HCMV infections in neonates and infants.
- 4) Treatment of an acute HCMV infection in pregnant women.
- 10 5) Treatment of HCMV infection in immunosuppressed patients associated with cancer and cancer therapy.
- 6) Treatment of HCMV-positive cancer patients with the target to reduce HCMV-mediated tumour progression (cf. J. Cinatl et al., *FEMS Microbiology Reviews* **2004**, 28, 59-77).
- 15

The present invention furthermore provides the use of the compounds according to the invention for the treatment and/or prophylaxis of diseases, especially viral infections, in particular with the viruses mentioned above, and the infective diseases

20 caused thereby. Hereinbelow, a viral infection is to be understood as meaning both an infection with a virus and a disease caused by an infection with a virus.

The present invention furthermore provides the use of the compounds according to the invention for the treatment and/or prophylaxis of disorders, in particular the

25 disorders mentioned above.

The present invention furthermore provides the use of the compounds according to the invention for preparing a medicament for the treatment and/or prophylaxis of disorders, in particular the disorders mentioned above.

30

The compounds according to the invention are preferably used for preparing medicaments suitable for the prophylaxis and/or treatment of infections with a representative of the group of the Herpes viridae, in particular a cytomegalovirus, in

particular the human cytomegalovirus.

The present invention furthermore provides a method for the treatment and/or prophylaxis of disorders, in particular the disorders mentioned above, using an anti-virally effective amount of the compounds according to the invention.

- 5 The present invention furthermore provides medicaments comprising at least one compound according to the invention and at least one or more further active compounds, in particular for the treatment and/or prophylaxis of the disorders mentioned above. Active compounds which may be mentioned by way of example and by way of preference as being suitable for combinations are: antiviral active
10 compounds, such as gancyclovir or acyclovir.

The compounds according to the invention can act systemically and/or locally. For this purpose, they can be administered in a suitable way, such as, for example, by the oral, parenteral, pulmonary, nasal, sublingual, lingual, buccal, rectal, dermal, transdermal, conjunctival or otic route, or as implant or stent.

- 15 For these administration routes, it is possible to administer the compounds according to the invention in suitable administration forms.

- Suitable for oral administration are administration forms which work as described in the prior art and deliver the compounds according to the invention rapidly and/or in modified form, which comprise the compounds according to the invention in
20 crystalline and/or amorphous and/or dissolved form, such as, for example, tablets (uncoated and coated tablets, for example tablets provided with enteric coatings or coatings whose dissolution is delayed or which are insoluble and which control the release of the compound according to the invention), tablets which rapidly decompose in the oral cavity, or films/wafers, films/lyophilizates, capsules (for
25 example hard or soft gelatin capsules), sugar-coated tablets, granules, pellets, powders, emulsions, suspensions, aerosols or solutions.

Parenteral administration can take place with avoidance of an absorption step (for example intravenously, intraarterially, intracardially, intraspinally or intralumbally) or with inclusion of absorption (for example intramuscularly, subcutaneously,

intracutaneously, percutaneously or intraperitoneally). Administration forms suitable for parenteral administration are, inter alia, preparations for injection and infusion in the form of solutions, suspensions, emulsions, lyophilizates or sterile powders.

Examples suitable for the other administration routes are pharmaceutical forms for
5 inhalation (inter alia powder inhalers, nebulizers), nasal drops/solutions/sprays; tablets to be administered lingually, sublingually or buccally, films/wafers or capsules, suppositories, preparations for the eyes or ears, vaginal capsules, aqueous suspensions (lotions, shaking mixtures), lipophilic suspensions, ointments, creams, transdermal therapeutic systems, milk, pastes, foams, dusting powders, implants or
10 stents.

The active compounds according to the invention can be converted into the stated administration forms. This can take place in a manner known per se by mixing with inert nontoxic, pharmaceutically acceptable auxiliaries. These auxiliaries include, inter alia, carriers (for example microcrystalline cellulose, lactose, mannitol),
15 solvents (for example liquid polyethylene glycols), emulsifiers and dispersants or wetting agents (for example sodium dodecyl sulphate, polyoxysorbitan oleate), binders (for example polyvinylpyrrolidone), synthetic and natural polymers (for example albumin), stabilizers (for example antioxidants, such as, for example, ascorbic acid), colorants (for example inorganic pigments, such as, for example, iron
20 oxides) and flavour- and/or odour-masking agents.

The present invention furthermore provides medicaments comprising at least one compound according to the invention, usually together with one or more inert non-toxic, pharmaceutically acceptable auxiliaries, and their use for the purposes mentioned above.

25 In general, it has proved advantageous to administer on intravenous administration amounts of from about 0.001 to 10 mg/kg, preferably from about 0.01 to 5 mg/kg, of body weight to achieve effective results, and the dosage on oral administration is from about 0.01 to 25 mg/kg, preferably from 0.1 to 10 mg/kg, of body weight.

It may nevertheless be necessary, where appropriate, to deviate from the amounts
30 mentioned, depending on the body weight, the administration route, the individual

response to the active compound, the mode of preparation and the time or interval over which administration takes place. Thus, in some cases it may be sufficient to make do with less than the aforementioned minimal amount, whereas in other cases the upper limit mentioned must be exceeded. In the event of administration of larger
5 amounts, it may be advisable to divide these into a plurality of individual doses over the day.

The percentage data in the following tests and examples are percentages by weight unless otherwise indicated; parts are parts by weight. Solvent ratios, dilution ratios and concentration data of liquid/liquid solutions are in each case based on volume.

A. Examples**Abbreviations used:**

BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
CDCl ₃	deuteriochloroform
CD ₃ CN	deuteroacetonitrile
TLC	thin-layer chromatography
DCI	direct chemical ionization (in MS)
DCM	dichloromethane
DIEA	N,N-diisopropylethylamine (Hünig base)
DMSO	dimethyl sulphoxide
DMF	N,N-dimethylformamide
EA	ethyl acetate
EI	electron impact ionization (in MS)
ESI	electrospray ionization (in MS)
m.p.	melting point
sat.	saturated
h	hour
HPLC	high-pressure, high-performance liquid chromatography
conc.	concentrated
LC-MS	liquid chromatography-coupled mass spectroscopy
LDA	lithium diisopropylamide
MS	mass spectroscopy
NMR	nuclear magnetic resonance spectroscopy
RP-HPLC	reverse phase HPLC
RT	room temperature
R _t	retention time (in HPLC)
THF	tetrahydrofuran

30009-26

- 22 -

General LCMS and HPLC methods:

Method 1 (HPLC): Instrument: HP 1100 with DAD detection; column: KromasilTM RP-18, 60 mm × 2 mm, 3.5 µm; mobile phase A: 5 ml of HClO₄/l of water, mobile phase B: acetonitrile; gradient: 0.0 min 2%B, 0.5 min 2%B, 4.5 min 90%B, 6.5 min 90%B; flow rate: 0.75 ml/min; oven: 30°C; UV detection: 210 nm.

Method 2 (HPLC, separation of enantiomers): chiral silica gel selector KBD 6136 (10 µm, 350 × 30 mm) based on the selector poly(N-methacryloyl-L-leucine-l-menthylamide); temperature: 24°C; flow rate: 50 ml/min; UV detection: 254 nm; sample application in ethyl acetate; elution mixtures of isohexane (A)/ethyl acetate (B), for example: gradient: → 0.0 min 40%B → 9.0 min 40%B → 9.01 min 100%B → 12.0 min 100%B → 12.01 min 40%B → 15 min 40%B.

Method 3 (LCMS): MS instrument type: Micromass ZQ; HPLC instrument type: HP 1100 Series; UV DAD; column: Phenomenex Synergi 2µ Hydro-RP Mercury 20 mm × 4 mm; mobile phase A: 1 l of water + 0.5 ml of 50% strength formic acid, mobile phase B: 1 l of acetonitrile + 0.5 ml of 50% strength formic acid; gradient: 0.0 min 90%A → 2.5 min 30%A → 3.0 min 5%A → 4.5 min 5%A; flow rate: 0.0 min 1 ml/min, 2.5 min/3.0 min/4.5 min 2 ml/min; oven: 50°C; UV detection: 210 nm.

Method 4 (HPLC, preparative separation): column: CromSil C18, 250 mm × 30 mm; mobile phase A: water, mobile phase B: acetonitrile; gradient: 3 min 10%B → 31 min 90%B → 34 min 90%B → 34.01 min 10%B; run time: 38 min; flow rate: 50 ml/min; UV detection: 210 nm.

Method 5 (LCMS): MS instrument type: MicromassTM ZQ; HPLC instrument type: Waters Alliance 2795; column: Phenomenex Synergi 2µ Hydro-RP Mercury 20 mm × 4 mm; mobile phase A: 1 l of water + 0.5 ml of 50% strength formic acid, mobile phase B: 1 l of acetonitrile + 0.5 ml of 50% strength formic acid; gradient: 0.0 min 90%A → 2.5 min 30%A → 3.0 min 5%A → 4.5 min 5%A; flow rate:

30009-26

- 23 -

0.0 min 1 ml/min, 2.5 min/3.0 min/4.5 min 2 ml/min; oven: 50°C; UV detection: 210 nm.

Method 6 (HPLC): instrument: HP 1100 with DAD detection; column: Kromasil
5 RP-18, 60 mm × 2 mm, 3.5 µm; mobile phase A: 5 ml of HClO₄/l of water, mobile phase B: acetonitrile; gradient: 0.0 min 2%B, 0.5 min 2%B, 4.5 min 90%B, 9 min 90%B; flow rate: 0.75 ml/min; oven: 30°C; UV detection: 210 nm.

Method 7 (LC-MS): instrument: Micromass Platform LCZ with HPLC AgilentTM
10 Series 1100; column: Grom-SIL120 ODS-4 HE, 50 mm × 2.0 mm, 3 µm; mobile phase A: 1 l of water + 1 ml of 50% strength formic acid, mobile phase B: 1 l of acetonitrile + 1 ml of 50% strength formic acid; gradient: 0.0 min 100%A → 0.2 min 100%A → 2.9 min 30%A → 3.1 min 10%A → 4.5 min 10%A; oven: 55°C; flow rate: 0.8 ml/min; UV detection: 210 nm.

Working Examples**General procedure [A]: synthesis of substituted 2-aminocinnamic acid derivatives from 2-halo-substituted anilines by Heck coupling**


5

In a one-necked flask, 1.0 equivalent of an aryl halide is initially charged in acetonitrile with 1.6 equivalents of methyl acrylate, 2.0 equivalents of triethylamine, 0.03 equivalents of palladium(II) acetate and 0.03 equivalents of tri-*o*-tolylphosphine (about 1M solution). The mixture is stirred under reflux for 48 hours. After the
10 reaction has ended (the reaction is monitored by TLC), the solvent is removed. The residue is purified chromatographically on silica gel using cyclohexane/ethyl acetate = 8:2 v/v.

Example 1A

15

Methyl (2E)-3-[2-amino-3-fluorophenyl]propenoate




Starting with 42.00 g (221.04 mmol) of 2-bromo-6-fluoroaniline, the general
20 procedure [A] gives 29.66 g (68% of theory) of product.

HPLC (method 1): $R_t = 4.14$ min

MS (ESI-pos): $m/z = 196$ ($M+H$)⁺

Example 2A

Methyl 2-amino-3-[(1E)-3-methoxy-3-oxo-1-propenyl]benzoate



5

Starting with 2.00 g (8.69 mmol) of methyl 2-amino-3-bromobenzoate, the general procedure [A] gives 1.29 g (60% of theory) of product.


HPLC (method 1): $R_t = 4.42$ min

10

MS (ESI-pos): $m/z = 236$ ($M+H$)⁺

Example 3A

15 Methyl (2E)-3-(2-amino-3,5-difluorophenyl)-2-propenoate




Starting with 3.00 g (14.42 mmol) of 2-bromo-4,6-difluoroaniline, the general procedure [A] gives 1.41 g (45% of theory) of product.

HPLC (method 1): $R_t = 4.23$ min

MS (ESI-pos): $m/z = 214$ ($M+H$)⁺

5 **Example 4A**

Methyl 4-amino-3-[(1E)-3-methoxy-3-oxo-1-propenyl]benzoate



Starting with 25.00 g (90.23 mmol) of methyl 4-amino-3-iodobenzoate, the general
10 procedure [A] gives 24.31 g (92% of theory) of product.


HPLC (method 1): $R_t = 4.71$ min

MS (ESI-pos): $m/z = 278$ ($M+H$)⁺

15

Example 5A

Methyl (2E)-3-[2-amino-5-cyanophenyl]-2-propenoate



Starting with 1.90 g (9.64 mmol) of 3-bromo-4-aminobenzonitrile, the general procedure [A] gives 1.28 g (50% of theory) of product.

HPLC (method 1): $R_t = 2.85$ min

5

MS (DCI-pos): $m/z = 220$ ($M + NH_4$)⁺

General procedure [B]: Synthesis of substituted 2-nitrocinnamic acid derivatives from 2-halo-substituted benzaldehydes by Wittig-Horner reaction


10

In a 100 ml one-necked flask, 27.5 mmol of methyl diethylphosphonoacetate, 25.0 mmol of the benzaldehyde and 27.5 mmol of lithium hydroxide are suspended in tetrahydrofuran. After the reaction has ended (the reaction is monitored by TLC), an identical volume of water is added to the mixture. The aqueous phase is extracted
15 three times with ethyl acetate. The combined organic phases are then washed with saturated sodium chloride solution and dried over sodium sulphate, and the solvent is removed. Without further purification, the product is dried under high vacuum at RT. In cases where many impurities are present, the product may be purified by column chromatography on silica gel using cyclohexane/ethyl acetate.

20

Example 6A

Methyl (2E)-3-(3-methoxy-2-nitrophenyl)-2-propenoate




25 Starting with 2.00 g (11.04 mmol) of 3-methoxy-2-nitrobenzaldehyde, the general procedure [B] gives 2.46 g (92% of theory) of product.

HPLC (method 1): $R_t = 4.37$ min

MS (ESI-pos): $m/z = 238$ ($M+H$)⁺

5 **Example 7A**

Methyl (2E)-3-(5-fluoro-2-nitrophenyl)-2-propenoate



Starting with 20.0 g (118.3 mmol) of 5-fluoro-2-nitrobenzaldehyde, the general
10 procedure [B] gives 7.25 g (27% of theory) of product.


MS (DCI): $m/z = 243$ ($M+NH_4$)⁺

15 **General procedure [C]: preparation of a 2-nitrobenzaldehyde from a benzyl
halide**

10.0 mmol of benzyl halide, 4.1 g of molecular sieve 4 Å and 20.0 mmol of
N-methylmorpholine *N*-oxide are suspended in 45 ml of acetonitrile. Until complete
conversion (the reaction is monitored by TLC), the mixture is stirred at RT. After the
20 reaction has ended, the molecular sieve is filtered off, the solvent is evaporated and
the residue is taken up again in ethyl acetate. This solution is washed initially with
1N hydrochloric acid and then with saturated sodium chloride solution. The organic
phase is separated off and then dried over sodium sulphate, and the solvent is
evaporated again. According to an analytical examination, the crude product is
25 sufficiently pure and can directly be reacted further.

Example 8A

2-Fluoro-6-nitrobenzaldehyde



Starting with 2.00 g (8.55 mmol) of 3-fluoro-6-nitrobenzyl bromide, the general procedure [C] gives 1.09 g (75% of theory) of product.


5 HPLC (method 1): $R_t = 3.58$ min

General procedure [D]: reduction of the nitro group of the 2-nitrocinnamic acid derivatives

- 10 Under argon, 25 mmol of the nitro compound and 125 mmol of tin(II) chloride dihydrate are initially charged in 60 ml of absolute ethanol or methanol in a 250 ml two-necked flask. This suspension is stirred under reflux for 30 minutes, and a clear solution is formed. The solution is then allowed to cool to room temperature and subsequently poured into ice-water. Using either solid sodium bicarbonate or a
- 15 saturated sodium carbonate solution, the pH is adjusted to pH = 7-8. 60 ml of ethyl acetate are then added, and the precipitated tin salts are filtered off through kieselguhr (a layer of a thickness of about 1 cm). The organic phase is separated off and the aqueous phase is re-extracted once with ethyl acetate. The organic phases are combined, washed once with saturated sodium chloride solution and dried over
- 20 sodium sulphate, and the solvent is concentrated to about half of its original volume. Activated carbon corresponding to 1% of the weight of the nitro compound is then added, and the mixture is heated under reflux for 30 minutes (the colour of the solution changes). The activated carbon is filtered off and the solvent is evaporated.
- 25 The residue that remains is an oil which, on drying at RT under high vacuum, forms crystals. Without further purification, it is used directly for the next step.

Example 9A

Methyl 3-[2-amino-6-fluorophenyl]propenoate



- 5 Starting with 7.25 g (32.2 mmol) of the nitro compound from Example 7A, the general procedure [D] gives 5.0 g (58% of theory) of product.

HPLC (method 1): $R_t = 3.33$ min


10 **General procedure [E]: synthesis of the N-arylpiperidine-4-carboxylic esters by Buchwald-Hartwig chemistry**

- The reaction is carried out under argon in a flask which was thoroughly dried by heating. 24.7 mmol of the bromobenzene, 74 mmol of piperidine-4-carboxylic ester
15 and 34.5 mmol of sodium tert-butoxide are initially charged in 100 ml of absolute toluene, 0.25 mmol of tris(dibenzylideneacetone)dipalladium and 0.74 mmol of BINAP are added, and the reaction mixture is heated to 120°C and heated under reflux for 16 hours. The reaction is terminated and the reaction mixture is extracted successively once with water and twice with 1N hydrochloric acid. The acidic
20 aqueous phase is then adjusted to pH 8 using 1N aqueous sodium hydroxide solution and extracted three times with dichloromethane. The combined organic phases are dried over sodium sulphate and filtered, and the solvent is removed under reduced pressure. The crude product obtained in this manner can be reacted further directly, without further purification.

25

Example 10A

Methyl N-(4-fluoro-3-methylphenyl)piperidine-4-carboxylate










starting with 10.6 g (74.0 mmol) of methyl piperidine-4-carboxylate and 4.67 g (24.7 mmol) of 5-bromo-2-fluorotoluene, the general procedure [E] gives 2.74 g (40% of theory) of product.

5

HPLC (method 1): $R_t = 3.48$ min

Examples 11A to 17A from the table below can be prepared in accordance with the general procedure [E].

Ex. No.	Structure	Starting material A Amount of starting material A Starting material B Amount of starting material B	Yield	HPLC R _t [min] (method)
11A		methyl piperidine-4-carboxylate 10.6 g (74.0 mmol) 4-fluoro-bromobenzene 4.32 g (24.7 mmol)	5.77 g (62% of theory)	3.25 (1)
12A		methyl piperidine-4-carboxylate 10.6 g (74.0 mmol) 3-fluoro-bromobenzene 4.32 g (24.7 mmol)	3.19 g (54% of theory)	3.41 (1)
13A		methyl piperidine-4-carboxylate 10.6 g (74.0 mmol) 3,4-dioxolan-bromobenzene 4.96 g (24.7 mmol)	3.75 (44% of theory)	3.27 (1)
14A		methyl piperidine-4-carboxylate 22.4 g (156.7 mmol) 3-chlorobromobenzene 10.00 g (52.3 mmol)	6.78 g (51% of theory)	3.74(1)
15A		methyl piperidine-4-carboxylate 5.02 g (35.1 mmol) 3-methyl-bromobenzene 2.00 g (11.7 mmol)	0.59 g (9% of theory)	3.46 (1)

16A		ethyl piperidine-4-carboxylate 5.00 g (31.8 mmol) 3-methoxy-bromobenzene 1.98 g (10.6 mmol)	1.66 g (59% of theory)	3.60 (1)
17A		ethyl piperidine-4-carboxylate 3.68 g (23.41 mmol) 4-bromo-1-fluoro-2-methoxybenzene 1.60 g (7.80 mmol)	1.32 g (60% of theory)	3.60 (1)


General procedure [F]: hydrolysis of the N-arylpiperidine-4-carboxylic esters

- 5 1.0 equivalent of the N-arylpiperidine-4-carboxylic ester is dissolved in dioxane, and 2.0 equivalents of 1N aqueous sodium hydroxide solution are added. The mixture is stirred at 80°C for 16 hours, and after the reaction has ended (the reaction is monitored by analytical HPLC) the mixture is concentrated. The residue is then taken up in water and adjusted to pH = 5 using 1N hydrochloric acid. The resulting
- 10 precipitate is filtered off, washed with a little water and cyclohexane and dried at room temperature under high vacuum. If the purity of the product is not high enough, the product is purified by preparative HPLC on an RP phase.

Example 18A

15

N-(4-Fluoro-4-methylphenyl)piperidine-4-carboxylic acid








Starting with 2.57 g (10.7 mmol) of the ester from Example 10A, the general procedure [F] gives 1.48 g (58% of theory) of product.

5 HPLC (method 1): $R_t = 3.35$ min

Examples 19A to 25A from the table below can be prepared in accordance with the general procedure [F].

Ex. No.	Structure	Starting material Amount of starting material	Yield	HPLC R_t [min] (method)
19A	<chem>OC(=O)C1CCN(C1)c2ccc(F)cc2</chem>	Example 11A 5.7 g (24.0 mmol)	2.06 g (38% of theory)	2.77 (1)
20A	<chem>OC(=O)C1CCN(C1)c2ccccc2F</chem>	Example 12A 3.19 g (13.4 mmol)	2.5 g (83% of theory)	3.05 (1)


21A		Example 13A 3.75 g (14.2 mmol)	2.85 g (80% of theory)	2.98 (1)
22A		Example 14A 6.73 g (26.5 mmol)	5.55 g (76% of theory)	3.36 (1)
23A		Example 15A 0.59 g (1.09 mmol)	0.13 g (48% of theory)	1.13 (3)
24A		Example 16A 1.57 g (5.96 mmol)	1.12 g (80% of theory)	3.10 (1)
25A		Example 17A 5.72 g (20.33 mmol)	4.98 g (97% of theory)	3.20 (1)

General procedure [G]: acylation of the 2-aminocinnamic esters with N-arylpiperidine-4-carboxylic acid

6.3 mmol of the N-arylpiperidine-4-carboxylic acid are initially charged in 75 ml of dichloromethane, a drop of DMF is added and 18.7 mmol of oxalyl chloride are, with ice-cooling, added to the mixture. After the addition has ended, the reaction mixture is heated under reflux for 1 h and cooled, and the solvent is removed under reduced pressure. The residue is taken up in 60 ml of dichloromethane and, with ice-cooling, added dropwise to a solution of 5.7 mmol of 2-aminocinnamic ester and 18.7 mmol of pyridine or triethylamine in 20 ml of dichloromethane. After the addition has ended, the mixture is heated under reflux for 2 h. The solvent is then removed under reduced pressure and the residue is taken up in dichloromethane. The organic phase is extracted twice with water and once with saturated sodium bicarbonate solution and dried over sodium sulphate. The sodium sulphate is filtered off and the solvent is removed under reduced pressure. The residue is triturated with diethyl ether and a few drops of ethyl acetate, and the precipitate is filtered off and washed with diethyl ether. The product is dried under reduced pressure. If the product is not pure enough for further reactions, it is purified chromatographically.

Example 26A



Methyl (2E)-3-fluoro-2-({[1-(4-fluoro-3-methyl)piperidin-4-yl]carbonyl}amino)-phenyl acrylate








Starting with 1.48 g (6.23 mmol) of the carboxylic acid from Example 18A and 1.11 g (5.67 mmol) of aniline from Example 1A, the general procedure [G] gives 7.77 g (79% of theory) of product.

5 HPLC (method 1): $R_t = 4.05$ min

Examples 27A to 33A from the table below can be prepared in accordance with the general procedure [G].

Ex. No.	Structure	Aniline Amount of aniline Acid Amount of acid	Yield	HPLC R_t [min] (method)
27A		Example 1A 4.81 g (24.67 mmol) Example 19A 5.8 g (25.98 mmol)	4.70 g (48% of theory)	3.80 (6)
28A		Example 1A 1.99 g (10.2 mmol) Example 20A 2.5 g (11.2 mmol)	3.75 g (92% of theory)	3.97 (1)

29A		Example 1A 2.03 g (10.4 mmol) Example 21A 2.85 g (11.4 mmol)	1.79 g (39% of theory)	3.89 (1)
30A		Example 1A 1.9 g (9.9 mmol) Example 22A 3.0 g (10.9 mmol)	2.78 g (68% of theory)	4.20 (1)
31A		Example 1A 0.22 g (1.11 mmol) Example 23A 0.29 g (1.20 mmol)	0.21 g (46% of theory)	3.99 (6)
32A		Example 1A 0.87 g (4.45 mmol) Example 24A 1.1 g (4.68 mmol)	0.47 g (26% of theory)	4.00 (1)

33A		Example 1A 0.73 g (3.76 mmol) Example 25A 2.40 g (9.48 mmol)	0.25 g (15% of theory)	4.00 (1)
-----	--	---	------------------------------	----------


General procedure [H]: cyclization of the 2-aminoacylcinnamic esters with anilines

- 5 At room temperature, 1.2 mmol of the 2-aminoacylcinnamic ester and 7.24 mmol of phosphorus oxychloride are initially charged in 10 ml of toluene. With vigorous stirring, the mixture is heated under reflux (bath temperature 120-125°C) for 16 h. The solvent is then distilled off under reduced pressure and codistilled once with toluene. The mixture is once again taken up in 10 ml of toluene, and 3.6 mmol of the
- 10 aniline are added. The mixture is heated at reflux for 24 h. The solvent is then removed under reduced pressure and the residue is taken up in dichloromethane and extracted twice with 1N hydrochloric acid. The organic phase is dried over sodium sulphate and filtered, and the solvent is removed under reduced pressure. The product is purified chromatographically on silica gel or by preparative HPLC
- 15 (method 4).

Example 34A

Methyl {8-fluoro-2-[1-(4-fluoro-3-methylphenyl)piperidin-4-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazolin-4-yl}acetate

20



Starting with 0.6 g (1.45 mmol) of the 2-acylaminocinnamic ester from Example 26A, the general procedure [H] and purification by preparative HPLC (method 4) and chromatography on silica gel using cyclohexane/ethyl acetate 8:2 (v/v) gives
 5 355 mg (39% of theory) of product.


HPLC (method 1): $R_t = 4.52$ min

MS: $m/z = 588$ ($M+H$)⁺

10

Example 35A

Methyl {8-fluoro-2-[1-(4-fluoro-3-methylphenyl)piperidin-4-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazolin-4-yl} acetate




15

Starting with 355 mg (0.6 mmol) of racemate from Example 34A, chromatographic separation of the enantiomers (method 2) gives 148 mg (42% of theory) of the compound as enantiomer B.

Example 36A

Methyl {8-fluoro-2-[1-(4-fluorophenyl)piperidin-4-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazolin-4-yl} acetate



5

Starting with 300 mg (0.75 mmol) of the 2-acylamino-cinnamic ester from Example 27A, the general procedure [H] gives 298 mg (69% of theory) of product.


HPLC (method 1): $R_t = 4.30$ min

10

MS (ESI pos): $m/z = 574$ ($M+H$)⁺

Example 37A

15 Methyl {8-fluoro-3-(2-methoxy-5-(trifluoromethyl)phenyl)-2-[1-(3-methylphenyl)-piperidin-4-yl]-3,4-dihydroquinazolin-4-yl} acetate



Starting with 0.057 g (0.14 mmol) of the 2-acylaminocinnamic ester from Example 31A, the general procedure [H] and purification by preparative HPLC (method 4) gives 32 mg (35% of theory) of product.

5 HPLC (method 1): $R_t = 4.44$ min

MS: $m/z = 570$ (M+H)⁺

Working Examples


10

General procedure [I]: hydrolysis of the quinazolylic esters

1.0 equivalent of the quinazolylic ester is dissolved in dioxane, and 5.0 equivalents of 1N aqueous sodium hydroxide solution are added. The mixture is stirred at 80°C for 16 hours and, after the reaction has ended (monitoring of reaction by means of analytical HPLC), the mixture is concentrated. The residue is then taken up in water and adjusted to pH = 5 using 1N hydrochloric acid. The resulting precipitate is filtered off, washed with a little water and diethyl ether and dried under high vacuum or in a drying cabinet. If the purity of the product is not high enough, the product is purified by preparative HPLC on an RP phase (method 4) or by chromatography on silica gel.

Example 1

25 {8-Fluoro-2-[1-(4-fluoro-3-methylphenyl)piperidin-4-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazolin-4-yl}acetic acid hydrochloride



Starting with 50 mg (0.085 mmol) of the methyl ester from Example 34A, the general procedure [I] and concentration of the product from a mixture of methanol/1N hydrochloric acid gives 13 mg (24% of theory) of product.

5


HPLC (method 1): $R_t = 4.27$ min

MS (ESI pos): $m/z = 574$ ($M+H$)⁺

- 10 ¹H-NMR (400 MHz, CD₃CN): δ [ppm] = 8.05-7.82 (m, 2H), 7.65-7.38 (m, 3H), 7.30 (d, 1H), 7.29-7.12 (m, 2H), 7.00 (d, 1H), 5.40 and 5.25 (2s, 1H), 4.00-3.75 (2s, 3H), 3.75-2.70 (m, 9H), 2.70-2.00 (m, 2H). (The ¹H-NMR indicates the presence of rotational isomers.)

15 Example 2

{8-Fluoro-2-[1-(4-fluoro-3-methylphenyl)piperidin-4-yl]-3-[2-methoxy-5-(trifluoromethyl)phenyl]-3,4-dihydroquinazolin-4-yl}acetic acid hydrochloride



Starting with 148 mg (0.25 mmol) of the methyl ester from Example 35A, the general procedure [I] and concentration of the product from a mixture of methanol/1N hydrochloric acid gives 62 mg (40% of theory) of product.

5

HPLC (method 1): $R_t = 4.27$ min


MS (ESI pos): $m/z = 574$ (M-HCl+H)⁺

- 10 ¹H-NMR (400 MHz, CD₃CN): δ [ppm] = 8.00-7.55 (m, 2H), 7.55-6.75 (m, 7H), 5.25 and 5.05 (2s, 1H), 3.95 and 3.75 (2s, 3H), 3.70-3.48 (d, 2H), 3.05-2.00 (m, 7H). (The ¹H-NMR indicates the presence of rotational isomers.)

Example 3

15

{8-Fluoro-3-[2-methoxy-5-(trifluoromethyl)phenyl]-2-[1-(3-methylphenyl)piperidin-4-yl]-3,4-dihydroquinazolin-4-yl}acetic acid



Starting with 41 mg (0.07 mmol) of the methyl ester from Example 37A, the general procedure [I] and concentration of the product from a mixture of methanol/1N hydrochloric acid and subsequent purification by chromatography (method 4) gives 13 mg (31% of theory) of product.

5

HPLC (method 7): $R_t = 2.64$ min


MS (ESI pos): $m/z = 556$ ($M+H$)⁺

- 10 ¹H-NMR (400 MHz, D₂O): δ [ppm] = 7.94-7.80 (m, 3H), 7.42-6.93 (m, 7H), 5.45-5.27 (m, 2H), 3.90 (s, 3H), 3.85-3.54 (m, 2H), 3.27-3.02 (m, 2H), 2.96-2.68 (m, 3H), 2.38-1.93 (m, 6H). (The ¹H-NMR indicates the presence of rotational isomers.)

Example 4

15

{8-Fluoro-2-[1-(4-fluorophenyl)piperidin-4-yl]-3-[2-methoxy-5-(trifluoromethyl)-phenyl]-3,4-dihydroquinazolin-4-yl}acetic acid







- 20 Starting with 50.00 mg (0.09 mmol) of the methyl ester from Example 36A, the general procedure [I] gives 30 mg (62% of theory) of product.






HPLC (method 1): $R_t = 4.20$ min






MS (DCI/NH₃): $m/z = 560$ ($M+H$)⁺





¹H-NMR (400 MHz, CDCl₃): δ [ppm] = 7.85-7.50 (m, 2H), 7.35-6.60 (m, 8H), 4.95 (m, 1H), 3.95 and 3.75 (2s, 3H), 3.20-1.50 (m, 10H). (The ¹H-NMR indicates the presence of rotational isomers.)



- 5 Examples 5 to 24 from the table below can be prepared according to the general procedures [H] and [I]. If required, hydrochlorides can be obtained after the reaction by concentrating the product from a mixture of methanol/1N hydrochloric acid.

Ex. No.	Structure	Molecular weight	Isomer	HPLC R_t [min] (method)	MS m/z
5		559.5	enantiomer B	4.20 (1)	560 [M+H] ⁺
6		576	enantiomer B	4.65 (1)	576 [M+H] ⁺
7		522	enantiomer B	4.78 (1)	522 [M+H] ⁺
8		542.4	enantiomer B	4.64 (1)	542 [M+H] ⁺

Ex. No.	Structure	Molecular weight	Isomer	HPLC R _t [min] (method)	MS m/z
9		540	racemate	4.13 (1)	540 [M+H] ⁺
10		553	racemate	4.50 (1)	553 [M+H] ⁺
11		547.6	racemate	4.40 (6)	548 [M+H] ⁺
12		576	racemate	4.77 (1)	576 [M+H] ⁺
13		588.5	racemate	3.96 (6)	552 [M-HCl +H] ⁺

Ex. No.	Structure	Molecular weight	Isomer	HPLC R _t [min] (method)	MS m/z
14	 <chem>COc1ccc(C(F)(F)F)cc1Oc2nc3c(nc(=O)c3cc2F)C4CCN(C4)c5ccc(F)cc5.Cl</chem> x HCl	596	racemate	4.53 (1)	560 [M-HCl +H] ⁺
15	 <chem>COc1ccc(Cl)cc1Oc2nc3c(nc(=O)c3cc2F)C4CCN(C4)c5ccc(Cl)cc5</chem>	542.4	racemate	4.67 (1)	542 [M+H] ⁺
16	 <chem>COc1cc(C(F)(F)F)c(F)cc1Oc2nc3c(nc(=O)c3cc2F)C4CCN(C4)c5ccc(F)cc5</chem>	547.5	racemate	4.10 (6)	548 [M+H] ⁺
17	 <chem>COc1cc(Cl)ccc1Oc2nc3c(nc(=O)c3cc2F)C4CCN(C4)c5ccc(Cl)cc5</chem>	522	racemate	4.70 (1)	522 [M+H] ⁺
18	 <chem>COc1cc(C)ccc1Oc2nc3c(nc(=O)c3cc2F)C4CCN(C4)c5ccc(F)cc5.Cl</chem> x HCl	556.1	racemate	4.10 (1)	520 [M-HCl +H] ⁺

Ex. No.	Structure	Molecular weight	Isomer	HPLC R _t [min] (method)	MS m/z
19		547.5	racemate	4.10 (1)	548 [M+H] ⁺
20		562.4	racemate	4.39 (1)	526 [M-HCl +H] ⁺
21		547.5	racemate	4.20 (1)	548 [M+H] ⁺
22		559.5	racemate	4.20 (1)	560 [M+H] ⁺

Ex. No.	Structure	Molecular weight	Isomer	HPLC R _t [min] (method)	MS m/z
23		575.6	racemate	4.20 (6)	576 [M+H] ⁺
24		571.6	racemate	4.20 (1)	572 [M+H] ⁺

B. Assessment of the physiological activity

The in vitro effect of the compounds of the invention can be shown in the following assays:

Anti-HCMV (anti-human cytomegalovirus) cytopathogenicity tests

The test compounds are employed as 50 millimolar (mM) solutions in dimethyl sulphoxide (DMSO). Ganciclovir, Foscarnet and Cidofovir are used as reference compounds. After addition of in each case 2 μ l of the 50, 5, 0.5 and 0.05 mM DMSO stock solutions to 98 μ l portions of cell culture medium in row 2 A-H for duplicate determinations, 1:2 dilutions are carried out with 50 μ l portions of medium up to row 11 of the 96-well plate. The wells in rows 1 and 12 each contain 50 μ l of medium. Then 150 μ l portions of a suspension of 1×10^4 cells (human prepuce fibroblasts [NHDF]) are pipetted into each of the wells (row 1 = cell control) and, in rows 2-12, a mixture of HCMV-infected and uninfected NHDF cells (M.O.I. = 0.001 – 0.002),

i.e. 1-2 infected cells per 1000 uninfected cells. Row 12 (without substance) serves as virus control. The final test concentrations are 250-0.0005 μM . The plates are incubated at 37°C/5% CO₂ for 6 days, i.e. until all the cells are infected in the virus controls (100% cytopathogenic effect [CPE]). The wells are then fixed and stained
5 by adding a mixture of formalin and Giemsa's dye (30 minutes), washed with double-distilled water and dried in a drying cabinet at 50°C. The plates are then assessed visually using an overhead microscope (Plaque Multiplier from Technomara).

The following data can be acquired from the test plates:

10 CC₅₀ (NHDF) = substance concentration in μM at which no visible cytostatic effects on the cells are evident by comparison with the untreated cell control;

EC₅₀ (HCMV) = substance concentration in μM which inhibits the CPE (cytopathic effect) by 50% compared with the untreated virus control;

SI (selectivity index) = CC₅₀ (NHDF) / EC₅₀ (HCMV).

15 Representative in vitro data for the effects of the compounds of the invention are shown in Table A:

Table A

Example No.	NHDF CC₅₀ [μM]	HCMV EC₅₀ [μM]	SI HCMV
1	24	0.03	800
2	31	0.04	775
3	31	0.1	310
4	36	0.15	237

The suitability of the compounds of the invention for the treatment of HCMV
20 infections can be shown in the following animal model:

30009-26

- 53 -

HCMV Xenograft Gelfoam® model

Animals: 3-4-week old female immunodeficient mice (16-18 g), Fox Chase SCID or Fox Chase SCID-NOD or SCID beige, are purchased from commercial breeders (Bomholtgaard, Jackson). The animals are housed under sterile conditions (including
5 bedding and feed) in isolators.

Virus growing: Human cytomegalovirus (HCMV), Davis strain, is grown in vitro on human embryonic prepuce fibroblasts (NHDF cells). After the NHDF cells have been infected with a multiplicity of infection (M.O.I.) of 0.01, the virus-infected cells are harvested 5-7 days later and stored in the presence of minimal essential medium
10 (MEM), 10% foetal calf serum (FCS) with 10% DMSO at -40°C. After serial ten-fold dilutions of the virus-infected cells, the titre is determined on 24-well plates of confluent NHDF cells after vital staining with Neutral Red, or fixation and staining with a formalin/Giemsa mixture (as described above).

Preparation of the sponges, transplantation, treatment and evaluation: Collagen
15 sponges 1×1×1 cm in size (Gelfoam®; from Peasel & Lorey, order No. 407534; K.T. Chong et al., Abstracts of 39th Interscience Conference on Antimicrobial Agents and Chemotherapy, 1999, p. 439; P.M. Kraemer et al., Cancer Research 1983, (43): 4822-4827) are initially wetted with phosphate-buffered saline (PBS), the trapped air bubbles are removed by degassing, and then stored in MEM + 10% FCS. 1× 10⁶
20 virus-infected NHDF cells (infection with HCMV Davis M.O.I. = 0.01) are detached 3 hours after infection and added in a drop of 20 µl of MEM, 10% of FCS, to a moist sponge. 12-13 hours later, the infected sponges are optionally applied to 25 µl of PBS /0.1% BSA/1 mM DTT with 5 ng/µl basic fibroblast growth factor (bFGF) and incubated for 1 hour. For the transplantation, the immunodeficient mice are
25 anaesthetized with Avertin or a mixture of xylazine/azepromazine and ketamine, the fur on the back is removed using a shaver, the epidermis is opened 1-2 cm, unstressed and the moist sponges are transplanted under the dorsal skin. The surgical wound is closed with tissue glue. 24 hours after the transplantation, the mice are, over a period of 8 days, treated with substance perorally three times a day (7.00 h
30 and 14.00 h and 19.00 h), two times a day (8.00 h and 17.00 h) or once a day (14.00 h). The dose is 3 or 10 or 30 or 100 mg/kg of body weight, the volume

administered is 10 ml/kg of body weight. The substances are formulated in the form of a 0.5% strength Tylose suspension, optionally with 2% DMSO. 9 days after transplantation and 16 hours after the last administration of substance, the animals are painlessly sacrificed and the sponge is removed. The virus-infected cells are released from the sponge by collagenase digestion (330 U/1.5 ml) and stored in the presence of MEM, 10% foetal calf serum, 10% DMSO at -140°C. Evaluation takes place after serial ten-fold dilutions of the virus-infected cells by determining the titre on 24-well plates of confluent NHDF cells after vital staining with Neutral Red or after fixation and staining with a formalin/Giemsa mixture (as described above). The number of infectious virus particles after the substance treatment compared with the placebo-treated control group is determined.

30009-26

- 55 -

C. Exemplary embodiments of pharmaceutical compositions

The compounds of the invention can be converted into pharmaceutical preparations in the following ways:

Tablet:

- 5 **Composition:** 100 mg of the compound of Example 1, 50 mg of lactose (monohydrate), 50 mg of corn starch (native), 10 mg of polyvinylpyrrolidone (PVP 25) (from BASF, Ludwigshafen, Germany) and 2 mg of magnesium stearate.

Tablet weight 212 mg. Diameter 8 mm, radius of curvature 12 mm.

- 10 **Production:** The mixture of active ingredient, lactose and starch is granulated with a 5% strength solution (m/m) of the PVP in water. The granules are then dried and mixed with the magnesium stearate for 5 min. This mixture is compressed using a conventional tablet press (see above for format of the tablet). A guideline for the compressive force used for the compression is 15 kN.

Suspension which can be administered orally:


- 15 **Composition:** 1000 mg of the compound of Example 1, 1000 mg of ethanol (96%), 400 mg of RhodigelTM (xanthan gum from FMC, Pennsylvania, USA) and 99 g of water.

10 ml of oral suspension are equivalent to a single dose of 100 mg of the compound of the invention.

- 20 **Production:** The Rhodigel is suspended in ethanol, and the active ingredient is added to the suspension. The water is added while stirring. The mixture is stirred for about 6 h until the swelling of the Rhodigel is complete.

Patent Claims

1. Compound of the formula



5 in which

10 R^1 , R^2 and R^3 independently of one another represent hydrogen, alkyl, alkoxy, carboxyl, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, trifluoromethyl, halogen, cyano, hydroxyl or nitro,

15 R^4 and R^5 independently of one another represent hydrogen, alkyl, alkoxy, alkylthio, cyano, halogen, nitro, trifluoromethyl or trifluoromethoxy,

R^6 represents alkyl, cyano, halogen, nitro or trifluoromethyl,

R^7 and R^8 independently of one another represent hydrogen, halogen, alkyl or alkoxy and

20 R^9 represents aryl or 1,3-benzodioxol-5-yl, where aryl and 1,3-benzodioxol-5-yl may be substituted by 1 to 3 substituents, where the substituents independently of one another are selected from the group consisting of alkoxy, alkylthio, carboxyl, alkylcarbonyl, alkoxy carbonyl, aminocarbonyl, trifluoromethyl, halogen, carbamoyl, cyano, hydroxyl, amino,

25

alkylamino, nitro and optionally hydroxyl-substituted alkyl,

or one of its salts, its solvates or the solvates of its salts.

5 2. Compound according to Claim 1, characterized in that

R^1 , R^2 and R^3 independently of one another represent hydrogen, methyl, fluorine, chlorine, cyano, hydroxyl or aminocarbonyl,

10 R^4 and R^5 independently of one another represent hydrogen, fluorine, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy,

R^6 represents chlorine, nitro, trifluoromethyl, methyl, isopropyl or tert-butyl,

15

R^7 and R^8 independently of one another represent hydrogen or C_1 - C_3 -alkyl and

20 R^9 represents phenyl or 1,3-benzodioxol-5-yl, where phenyl may be substituted by 1 to 3 substituents, where the substituents independently of one another are selected from the group consisting of C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, carboxyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, trifluoromethyl, fluorine, chlorine, bromine, cyano, hydroxyl, amino, C_1 - C_6 -alkylamino and nitro.

25

3. Compound according to Claim 1 or 2, characterized in that

30 R^1 and R^2 are hydrogen,

R^3 is fluorine,

R^4 and R^5 independently of one another are hydrogen, fluorine or

30009-26


- 58 -

methoxy,

 R^6 is trifluoromethyl, R^7 and R^8 are hydrogen and

R^9 is phenyl, where phenyl may be substituted by 1 or 2 substituents, where the
 5 substituents independently of one another are selected from the group consisting of
 methyl, methoxy, ethoxy, fluorine and chlorine.

4. Process for preparing a compound of the formula (I) according to
 Claim 1, characterized in that a compound of the formula



10 in which

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 and R^9 are as defined in Claim 1, and

R^{10} represents alkyl,

is reacted with a base.


5. Process according to Claim 4 wherein R^{10} is methyl or ethyl.

15 6. Compound according to any one of Claims 1 to 3 for the treatment
 and/or prophylaxis of diseases.

30009-26

- 59 -

7. Use of a compound according to any one of Claims 1 to 3 for preparing a medicament for the treatment and/or prophylaxis of diseases.
8. Use of a compound according to any one of Claims 1 to 3 for preparing a medicament for the treatment and/or prophylaxis of viral infections.
- 5 9. Use according to Claim 8, characterized in that the viral infection is an infection with the human cytomegalovirus (HCMV) or another representative of the group of the Herpes viridae.
10. Medicament, comprising a compound as defined in any one of Claims 1 to 3 in combination with a further active compound.
- 10 11. Medicament, comprising a compound according to any one of Claims 1 to 3 in combination with an inert nontoxic, pharmaceutically acceptable auxiliary.
12. Medicament according to Claim 11 for the treatment and/or prophylaxis of viral infections.
13. Use of an antivirally effective amount of at least one compound
- 15 according to any one of Claims 1 to 3, or a medicament according to any one of Claims 10 to 12 for controlling viral infections in humans and animals.



(I)