



(51) International Patent Classification:

A61K 31/16 (2006.01) *A61P 35/02* (2006.01)
A61K 31/167 (2006.01) *A61P 35/00* (2006.01)

(21) International Application Number:

PCT/US2014/031803

(22) International Filing Date:

26 March 2014 (26.03.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/805,542 27 March 2013 (27.03.2013) US

(71) Applicant: EMORY UNIVERSITY [US/US]; Office of Technology Transfer, 1599 Clifton Road NE, 4th Floor, Atlanta, Georgia 30322 (US).

(72) Inventors: KHOURY, Hanna Jean; 3021 Cravey Trail, Atlanta, Georgia 30345 (US). ANTUN, Ana; 1826 N. Akin, Atlanta, Georgia 30345 (US).

(74) Agents: MASON, James C. et al.; Emory University, Office of Technology Transfer, 1599 Clifton Road NE, 4th Floor, Atlanta, Georgia 30322 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with declaration under Article 17(2)(a); without abstract; title not checked by the International Searching Authority

(54) Title: USES OF 6-AMINOHEXANOIC ACID TO MANAGE BLEEDING CONDITIONS

(57) Abstract:



WO 2014/160742 A2

USES OF 6-AMINOHEXANOIC ACID TO MANAGE BLEEDING CONDITIONS**CROSS REFERENCE OF RELATED APPLICATIONS**

This Application claims priority to U.S. Provisional Application Number 61/805,542
5 filed the 27th of March 2013, hereby incorporated by reference in its entirety.

BACKGROUND

Thrombocytopenia is frequently chronic, severe, and occasionally refractory to
platelet transfusions in patients with marrow failure syndromes and hematological
10 malignancies. Despite prophylactic allogeneic platelet transfusions, maintaining platelet
counts in a safe range is often difficult in this patient population. Bleeding complications
occur in approximately 20% of patients with acute myeloid leukemia and in up to 58% of
hematopoietic stem cell transplants recipients. Thus, there is a need to identify improved
therapeutic methods to control thrombocytopenia.

15 AMICAR[®] is an FDA approved product containing epsilon aminocaproic acid
(EACA) (6-aminohexanoic acid) reported to enhance hemostasis when fibrinolysis
contributes to bleeding. For the treatment of acute bleeding syndromes due to elevated
fibrinolytic activity, the label of AMICAR[®] indicates using 5 AMICAR[®] 1000 mg tablets or
10 AMICAR[®] 500 mg tablets (5 g) or 20 milliliter of AMICAR[®] oral solution (5 g) that is
20 to be administered during the first hour of treatment.

Kalmadi et al. report that EACA reduces transfusion requirements in patients with
thrombocytopenic hemorrhage. Cancer, 2006, 107, 136-140. Schwartz et al., report EACA
in the treatment of patients with acute promyelocytic leukemia and acquired alpha-2-
plasmin inhibitor deficiency. Ann Intern Med, 1986, 105(6):873-7. See also Wassenaar et
25 al., Hematol Oncol, 2008, 26(4):241-6; Garewal and Durie, Scand J Haematol, 1985, 35,
497-500; Deysine and Clifton, Annals of the New York Academy of Sciences, 1964, 115,
291-297; Mannucci, New England Journal of Medicine, 1998, 339, 245-253; and U.S.
Patent 5,415,863.

Antun et al. report epsilon aminocaproic acid prevents bleeding in severely
30 thrombocytopenic patients with hematological malignancies. Cancer. 2013, 1;119(21):3784-
7.

References cited herein are not an admission of prior art.

SUMMARY

This disclosure relates to the use of EACA (6-aminohexanoic acid) as an antifibrinolytic for the prevention of thrombocytopenic bleeds. In certain embodiment, the disclosure relates to methods of treating or preventing bleeding in patients with
5 thrombocytopenia resulting from hematological malignancies by administering 6-aminohexanoic acid, salts, or prodrugs thereof prophylactically to a subject in need thereof. In certain embodiments, subjects with stable chronic thrombocytopenia previously managed with platelet transfusion represent a selected group of patients that benefit from prophylaxis.

In certain embodiments, the disclosure contemplates methods of treating or
10 preventing excessive bleeding comprising administering e.g., once, twice, or three times daily, a pharmaceutical composition comprising 6-aminohexanoic acid, salt, or prodrug thereof to a subject in need thereof, e.g., human subject, diagnosed with cancer or a hematological disorder.

In certain embodiments, the subject is on a chemotherapy or immunotherapy.

15 In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.1 to 4.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

20 In certain embodiments, the subject receives about 2.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof daily, e.g., wherein administration is twice daily a pharmaceutical composition comprising about 1.0 gram of 6-aminohexanoic acid or molar equivalent salt thereof.

In certain embodiments, the administration is sustained as long as or until a
25 measured platelet count is greater than $20 \times 10^9/L$, $25 \times 10^9/L$, $30 \times 10^9/L$, $35 \times 10^9/L$, or $40 \times 10^9/L$.

In certain embodiments, the administration is sustained for greater than 20, 30, or 40 days.

In certain embodiments, the hematological disorder is chronic immune
30 thrombocytopenia, aplastic anemia, or a hematological malignancy such as leukemia, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), chronic myelogenous leukemia (CML), acute monocytic leukemia (AMoL), lymphoma, Hodgkin's lymphoma, or Non-Hodgkin's lymphoma.

In certain embodiments, the subject received an autologous or allogeneic hematopoietic stem cell transplantation, bone marrow transplant, or peripheral stem cell transplantation.

In certain embodiments, the administration is in combination with a platelet transfusion or not done in combination with a platelet transfusion.

In certain embodiments, the subject is less than 18, 12, or 6 years of age.

In certain embodiments, the administration is oral solution or tablet.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows a table of the characteristics of 44 thrombocytopenic patients who received epsilon aminocaproic acid for the prevention of bleeding. CML is chronic myeloid leukemia, AML is acute myeloid leukemia, ALL is acute lymphoblastic leukemia, MDS is myelodysplastic syndrome, CLL is chronic lymphocytic leukemia, ITP is immune thrombocytopenic purpura. *Traumatic brain injury in a car accident **One patient had 2 episodes of hematuria.

DETAILED DISCUSSION

Before the present disclosure is described in greater detail, it is to be understood that this disclosure is not limited to particular embodiments described, and as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present disclosure will be limited only by the appended claims.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present disclosure, the preferred methods and materials are now described.

All publications and patents cited in this specification are herein incorporated by reference as if each individual publication or patent were specifically and individually indicated to be incorporated by reference and are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited. The citation of any publication is for its disclosure prior to the filing date and should not be construed as an admission that the present disclosure is not entitled to antedate such publication by virtue of prior disclosure. Further, the dates of publication

provided could be different from the actual publication dates that may need to be independently confirmed.

As will be apparent to those of skill in the art upon reading this disclosure, each of the individual embodiments described and illustrated herein has discrete components and features which may be readily separated from or combined with the features of any of the other several embodiments without departing from the scope or spirit of the present disclosure. Any recited method can be carried out in the order of events recited or in any other order that is logically possible.

Embodiments of the present disclosure will employ, unless otherwise indicated, techniques of medicine, organic chemistry, biochemistry, molecular biology, pharmacology, and the like, which are within the skill of the art. Such techniques are explained fully in the literature.

It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. In this specification and in the claims that follow, reference will be made to a number of terms that shall be defined to have the following meanings unless a contrary intention is apparent.

Prior to describing the various embodiments, the following definitions are provided and should be used unless otherwise indicated.

As used herein, "salts" refer to derivatives of the disclosed compounds where the parent compound is modified making acid or base salts thereof. Examples of salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines, alkylamines, or dialkylamines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. In preferred embodiment the salts are conventional nontoxic pharmaceutically acceptable salts including the quaternary ammonium salts of the parent compound formed, and non-toxic inorganic or organic acids. Preferred salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2- acetoxymaleic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

"Subject" refers any animal, preferably a human patient, livestock, or domestic pet.

The term "prodrug" refers to an agent that is converted into a biologically active form in vivo. Prodrugs are often useful because, in some situations, they may be easier to

administer than the parent compound. An ester of a carboxylic acid group is contemplated prodrug. They may, for instance, be bioavailable by oral administration whereas the parent compound is not. The prodrug may also have improved solubility in pharmaceutical compositions over the parent drug. A prodrug may be converted into the parent drug by various mechanisms, including enzymatic processes and metabolic hydrolysis.

As used herein, the terms "prevent" and "preventing" include the prevention of the recurrence, spread or onset. It is not intended that the present disclosure be limited to complete prevention. In some embodiments, the onset is delayed, or the severity is reduced.

As used herein, the terms "treat" and "treating" are not limited to the case where the subject (e.g., patient) is cured and the condition or disease is eradicated. Rather, embodiments, of the present disclosure also contemplate treatment that merely reduces symptoms, and/or delays conditions or disease progression.

As used herein, the term "combination with" when used to describe administration with an additional treatment means that the agent may be administered prior to, together with, or after the additional treatment, or a combination thereof.

EACA Prevents Bleeding in Severely Thrombocytopenic Patients with Hematological Malignancies

Individuals with hematological malignancies (leukemia) and bone marrow failure often have low platelet count as a result of the effects of their underlying disease or the effects of chemotherapy on the blood forming cells in bone marrow, i.e., thrombocytopenia. A common diagnostic criteria used for thrombocytopenia is a low platelet count compared to the normal range of platelets. As the platelet count falls associated bleeding complications tend to increase. These include bleeding in the mouth and gum, bruising, nosebleeds, rash (pinpoint red spots called petechiae) and excessive, uncontrolled bleeding from cuts or wounds. Thrombocytopenia-associated bleeding remains a significant problem (up to 50% of diagnosed patients have bleeding complications). The standard of care is prophylactic platelet transfusion.

In the absence of an intervention that increases platelet production or accelerates platelet recovery, recurrent platelet transfusions remain the option for patients with severe chronic thrombocytopenia associated with hematological malignancies. This approach necessitates frequent visits to the medical center, is hampered by platelet shortages, and ultimately often leads to alloimmunization and platelet refractoriness. Additionally, this approach is ineffective in preventing significant bleedings in 20-50% of patients.

In one study disclosed herein, EACA was administered in patients with transfusion-dependent thrombocytopenia. Patients were managed in the outpatient setting despite their severe thrombocytopenia and received EACA as a substitution for prophylactic, threshold directed platelet transfusion. Patients reported rapid disappearance of petechiae and resolution of subcutaneous bleeds shortly after initiation of EACA. With a median duration of EACA administration of 47 days, but ranging up to 209 days, no major spontaneous bleeds were observed in these patients, whose median platelet counts were $8 \times 10^9/L$ and for whom no prophylactic platelet transfusions were administered. Two grade 3-4 bleedings occurred in the setting of a trauma and from a pre-existing condition. Only 16% of the patients received platelet transfusions (median of 4 units).

EACA is safe and is associated with a low risk of major spontaneous bleeding in severe and refractory thrombocytopenic patients with hematological disorders not receiving prophylactic platelet transfusion.

15 Administration

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once, twice, or three times daily doses, a pharmaceutical composition comprising 6-aminohexanoic acid, salt, or prodrug thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.1 to 4.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the subject receives about 2.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof daily, e.g., wherein administration is twice daily a pharmaceutical composition comprising about 1.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.5 to 4.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.5 to 3.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed
5 with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.5 to 2.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed
10 with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.75 to 1.5 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed
15 with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering twice daily a pharmaceutical composition comprising about 0.5 to 1.5 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a
20 hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.75 to 1.25 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed
25 with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.75 to 4.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed
30 with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.75 to 3.0 grams of 6-aminohexanoic acid

or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 0.75 to 2.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the subject receives about 1.5 to 2.5 grams of 6-aminohexanoic acid or molar equivalent salt thereof daily.

In certain embodiments, the subject receives about 1.5 to 3.5 grams of 6-aminohexanoic acid or molar equivalent salt thereof daily.

In certain embodiments, the subject receives about 1.5 to 4.5 grams of 6-aminohexanoic acid or molar equivalent salt thereof daily.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 4.0 to 5.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 5.0 to 50 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 5.0 to 30 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 5.0 to 25 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about 5.0 to 10 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about less than 1, 2, 5, 15, 20, 25, 30, or 50 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the disclosure contemplates methods of treating or preventing excessive bleeding comprising administering e.g., once or twice daily, a pharmaceutical composition comprising about more than 1, 2, 5, 15, 20, 25, 30, or 50 grams of 6-aminohexanoic acid or molar equivalent salt thereof to a subject in need thereof, e.g., human subject, diagnosed with a hematological disorder.

In certain embodiments, the administration is sustained according to doses provided herein until a measured platelet count is greater than $20 \times 10^9/L$, $25 \times 10^9/L$, $30 \times 10^9/L$, $35 \times 10^9/L$, or $40 \times 10^9/L$.

In certain embodiments, the administration is sustained for greater than 20, 30, or 40 days.

Combination therapies

In certain embodiments, the administration of 6-aminohexanoic acid is in combination with a platelet transfusion or not done in combination with a platelet transfusion.

In certain embodiments, the administration is in combination with a chemotherapy or immunotherapy.

Chemotherapy may entail administration of 6-aminohexanoic acid in combination with gefitinib, erlotinib, docetaxel, cis-platin, 5-fluorouracil, gemcitabine, tegafur, raltitrexed, methotrexate, cytosine arabinoside, hydroxyurea, adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin, vincristine, vinblastine, vindesine, vinorelbine taxol, taxotere, etoposide, teniposide, amsacrine, topotecan, camptothecin bortezomib anegrilide, tamoxifen, toremifene, raloxifene, droloxifene, idoxifene fulvestrant, bicalutamide, flutamide,

nilutamide, cyproterone, goserelin, leuprorelin, buserelin, megestrol, anastrozole, letrozole, vorazole, exemestane, finasteride, marimastat, trastuzumab, cetuximab, dasatinib, imatinib, bevacizumab, combretastatin, thalidomide, and/or lenalidomide or combinations thereof.

Immunotherapy may entail administration of 6-amionhexanoic acid in combination
5 with cyclophosphamide, abatacept, abetimus, adalimumab, afelimomab, alefacept, anakinra, anti-IL-6, anti-thymocyte globulin, ascomycin, azathioprine, basiliximab, belimumab, briakinumab, certolizumab pegol, ciclosporin, daclizumab, eculizumab, efalizumab, eritoran, etanercept, everolimus, fingolimod, gliotoxin, gusperimus, infliximab, laquinimod, leflunomide, lenalidomide, mapracorat, mepolizumab, methotrexate,
10 mizoribine, muromonab-CD3, mycophenolate mofetil, mycophenolic acid, natalizumab, neurovax, pegsunercept, pimecrolimus, pomalidomide, ridaforolimus, rilonacept, secukinumab, sirolimus, tacrolimus, teriflunomide, thalidomide, tocilizumab, umirolimus, ustekinumab, voclosporin, zotarolimus or combinations thereof.

In certain embodiments, the administration of 6-amionhexanoic acid is in
15 combination with chlorambucil or cyclophosphamide, and optionally a corticosteroid such as prednisone or prednisolone.

In certain embodiments, the administration of 6-amionhexanoic acid is in combination with imatinib.

In certain embodiments, the administration of 6-amionhexanoic acid is in
20 combination with cladribine.

In certain embodiments, the administration of 6-amionhexanoic acid is in combination with pentostatin.

In certain embodiments, the administration of 6-amionhexanoic acid is in combination with rituximab or interferon-alpha.

25 In certain embodiments, the administration of 6-amionhexanoic acid is in combination with rituximab, cyclophosphamide, hydroxydaunorubicin, vincristine and optionally prednisone or prednisolone.

In certain embodiments, the administration of 6-amionhexanoic acid is in combination with cyclophosphamide, vincristine and optionally prednisone or prednisolone.

30 In certain embodiments, the administration of 6-amionhexanoic acid is in combination with cyclophosphamide, hydroxydaunorubicin, vincristine and optionally prednisone or prednisolone and optionally bendamustine.

In certain embodiments, in any of the above combinations, 6-amionhexanoic acid may be administered at any of the doses and regularity as provided herein.

EXPERIMENTAL

Example 1: Patients with Hematological Malignancies

5 Patients with chronic transfusion dependent thrombocytopenia who received prophylactic EACA in the outpatient setting were identified through a computerized database search. After initiation of EACA, prophylactic threshold-driven platelet transfusion was discontinued and patients received only therapeutic platelet transfusion in case of bleeding, except when admitted to the inpatient ward. In the inpatient ward, 10 prophylactic platelets were routinely infused when platelets were $< 10 \times 10^9/L$. The number of platelets transfused in the hospital in the absence of bleeding was censored. Patient demographics, diagnosis, treatment, platelet counts, duration of thrombocytopenia, transfusion history, bleeding and outcomes were captured onto study-specific case report forms. Bleeding was captured regardless of patient's location (out-patient or wards) and 15 retrospectively scored based on the ITP WHO classification: grade 1=petechial bleed, grade 2=clinically mild blood loss, grade 3=gross blood loss requiring transfusion, and grade 4=retinal, cerebral, or debilitating blood loss. Statistics were descriptive. Characteristics of the patients are summarized in Figure 1

Ninety-six percent had a hematological malignancy and 2 had a non-malignant 20 hematological disorder (chronic immune thrombocytopenia, aplastic anemia). Median age was 61 (range, 17-82). Median platelet count at the time of initiation of EACA was $8 \times 10^9/L$ (range, 1-19). Prior to initiation of EACA, median duration of thrombocytopenia was 273 days (range, 20-1463) and patients were transfused 2-3 times a week, in the outpatient setting, when platelet count were $< 20 \times 10^9/L$. Thirty patients (68%) had a median of 4 25 (range, 1-12) hospitalizations of a median of 12 days duration (range, 4-23) during the course of treatment with EACA mainly for chemotherapy, but also for the management of neutropenic fever. Seventy one percent had relapsed/refractory disease; while 29% had a stable disease or were actively receiving chemotherapy at the time of EACA was administered. Ten (23%) were refractory to platelet transfusions and had platelet counts 30 continuously $< 10 \times 10^9/L$.

EACA was prescribed at the oral dose of 1 gram twice daily until sustained untransfused platelet count was $> 30 \times 10^9/L$. Median duration of administration of EACA was 47 days (range, 7-209). Patients were monitored in the outpatient clinic, with a CBC obtained at least once a week. While on EACA, 7/44(16%) patients received a median of 4

(range, 1-8) platelets transfusions to treat 10 bleeding episodes. Median score for these 10 bleeding episodes was 2 (2-4), with 1 patient experiencing a traumatic intracranial hemorrhage secondary to motor vehicle accident (grade 4), and 1 patient suffering an episode of grade 3 hematochezia from a pre-existing rectal ulcer. The other 5 patients had 8
5 grade 2 mucocutaneous bleeds: hemorrhoid, epistaxis, oral, bladder and vaginal. Eleven patients (25%) had 19 episodes of minor mucocutaneous (epistaxis, bruises, oral, and vaginal) that resolved spontaneously and did not require platelet transfusions. Median bleeding score for these 19 bleeding events was 1 (1-2). Of interest, only 4 of the 10 platelet refractory patients had bleeding episodes that were mucocutaneous (3 grade 1, 5 grade 2
10 and 1 grade 3 bleeding).

EACA was overall well tolerated and no patients discontinued EACA due to intolerance. One patient discontinued EACA due to the onset of hematuria (2%), 18 (41%) due to platelet recovery, 25 (57%) due to referral to hospice or death. No patient
15 experienced a venous thromboembolic event.

CLAIMS

1. A method of preventing excessive bleeding comprising administering daily a pharmaceutical composition comprising about 0.5 to 4.0 grams of 6-aminohexanoic acid or molar equivalent salt or prodrug thereof to a subject in need thereof.
2. The method of Claim 1, wherein the subject receives about 2.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof daily.
3. The method of Claims 1-2, wherein administration is twice daily a pharmaceutical composition comprising about 1.0 grams of 6-aminohexanoic acid or molar equivalent salt thereof.
4. The method of Claims 1-3, wherein the administration is sustained until a measured platelet count is greater than $20 \times 10^9/L$.
5. The method of Claims 1-4, wherein the administration is sustained for greater than 20 days.
6. The method of Claims 1-5, wherein the subject is diagnosed with a hematological disorder.
7. The method of Claims 1-6, wherein the hematological disorder is a hematological malignancy selected from leukemia, acute lymphoblastic leukemia (ALL), acute myelogenous leukemia (AML), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), chronic myelogenous leukemia (CML), acute monocytic leukemia (AMoL), lymphoma, Hodgkin's lymphoma, or Non-Hodgkin's lymphoma.
8. The method of Claims 1-7, wherein the subject received an autologous or allogeneic hematopoietic stem cell transplantation, bone marrow transplant or peripheral stem cell transplantation.
9. The method of Claims 1-8, wherein the administration is not done in combination with a platelet transfusion.

Age, years median (range)	61 (17-82)
Male/female	29/15
Platelet counts $\times 10^9/L$ when EACA started, median (range)	8 (1 – 19)
Patients with platelet refractoriness n (%)	10 (23%)
Duration of thrombocytopenia in days, median (range)	273 (20-1463)
Diagnosis n (%)	
AML	9 (21)
AML	14 (32)
ALL	1 (2)
MDS	15 (35)
CLL	1 (2)
ITP	1 (2)
Lymphoma	1 (2)
Aplastic anemia	1 (2)
Myelofibrosis	1 (2)
Disease Status of patients at the time EACA started	
Active treatment, n (%)	13 (29)
Relapsed/refractory, n (%)	31 (71)
Duration of EACA therapy, median(range) in days	47 (7-209)
Patients without bleeding on EACA, n (%)	26 (59)
Patients with bleeding episodes while on EACA, n (%)	18 (41)
Patients requiring platelet transfusions for bleeds, n (%)	7 (16%)
Grade and location of the 10 bleeding episodes requiring platelet transfusions:	
Grade 4 – CNS*	1
Grade 3 – rectal	1
Grade 2 – hemorrhoids	1
Grade 2 – hematuria **	2
Grade 2 – epistaxis	1
Grade 2 – oral	3
Grade 2 – vaginal	1
Grade and location of the 19 bleeding episodes not requiring platelet transfusions:	
Grade 1- epistaxis	7
Grade 1 – skin bruises	4
Grade 1 – gingival bleed	2
Grade 1 - vaginal	1
Grade 2 – gingival	2
Grade 2 – vaginal	1
Grade 2 - epistaxis	2
Reason for discontinuing EACA, n (%)	
Death	25 (57)
Increased platelets count	18 (41)
Hematuria	1 (2)

FIG. 1

PATENT COOPERATION TREATY

PCT

DECLARATION OF NON-ESTABLISHMENT OF INTERNATIONAL SEARCH REPORT
(PCT Article 17(2)(a), Rules 13ter.1(c) and (d) and 39)

Applicant's or agent's file reference 13023-PCT	IMPORTANT DECLARATION	Date of mailing (<i>day/month/year</i>) 07 July 2014 (07.07.2014)
International application No. PCT/US2014/031803	International filing date (<i>day/month/year</i>) 26 March 2014 (26.03.2014)	(Earliest) Priority date (<i>day/month/year</i>) 27 March 2013 (27.03.2013)
International Patent Classification (IPC) or both national classification and IPC A61K 31/16(2006.01)i, A61K 31/167(2006.01)i, A61P 35/02(2006.01)i, A61P 35/00(2006.01)i		
Applicant EMORY UNIVERSITY		

This International Searching Authority hereby declares, according to Article 17(2)(a), that **no international search report will be established** on the international application for the reasons indicated below.


1. ☒ The subject matter of the international application relates to:
 - a. ☐ scientific theories
 - b. ☐ mathematical theories
 - c. ☐ plant varieties
 - d. ☐ animal varieties
 - e. ☐ essentially biological processes for the production of plants and animals, other than microbiological processes and the products of such processes
 - f. ☐ schemes, rules or methods of doing business
 - g. ☐ schemes, rules or methods of performing purely mental acts
 - h. ☐ schemes, rules or methods of playing games
 - i. ☒ methods for treatment of the human body by surgery or therapy
 - j. ☐ methods for treatment of the animal body by surgery or therapy
 - k. ☐ diagnostic methods practised on the human or animal body
 - l. ☐ mere presentation of information
 - m. ☐ computer programs for which this International Searching Authority is not equipped to search prior art
2. ☐ The failure of the following parts of the international application to comply with prescribed requirements prevents a meaningful search from being carried out:

☐ the description
 ☐ the claims
 ☐ the drawings
3. ☐ A meaningful search could not be carried out without the sequence listing; the applicant did not, within the prescribed time limit:

☐ furnish a sequence listing on paper complying with the standard provided for in Annex C of the Administrative Instructions, and such listing was not available to the International Searching Authority in a form and manner acceptable to it.

☐ furnish a sequence listing in electronic form complying with the standard provided for in Annex C of the Administrative Instructions, and such listing was not available to the International Searching Authority in a form and manner acceptable to it.

☐ pay the required late furnishing fee for the furnishing of a sequence listing in response to an invitation under Rule 13ter.1(a) or (b).
4. Further comments:

Name and mailing address of ISA/KR  International Application Division Korean Intellectual Property Office 189 Cheongsa-ro, Seo-gu, Daejeon Metropolitan City, 302-701, Republic of Korea Facsimile No. +82-42-472-7140	Authorized officer MOON, Sun Heup Telephone No. +82-42-481-8242	
--	---	---