



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2665205 C 2015/06/30

(11)(21) **2 665 205**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2007/10/05
(87) Date publication PCT/PCT Publication Date: 2008/04/10
(45) Date de délivrance/Issue Date: 2015/06/30
(85) Entrée phase nationale/National Entry: 2009/04/02
(86) N° demande PCT/PCT Application No.: GB 2007/003800
(87) N° publication PCT/PCT Publication No.: 2008/041004
(30) Priorité/Priority: 2006/10/06 (GB0619853.5)

(51) Cl.Int./Int.Cl. *G01N 1/34* (2006.01),
C12N 5/00 (2006.01), *G01N 33/569* (2006.01)

(72) Inventeurs/Inventors:
DURRANT, IAN, GB;
DAY, TONI, GB;
O'KEEFFE, AISLING, GB;
BAMPTON, MAXINE, GB

(73) Propriétaire/Owner:
OXFORD IMMUNOTEC LIMITED, GB

(74) Agent: GOUDREAU GAGE DUBUC

(54) Titre : PROCEDE DE PREPARATION
(54) Title: PREPARATION METHOD

(57) **Abrégé/Abstract:**

A method of storing and purifying cells from a whole blood sample, comprising storing the sample before and/or after purification of cells for at least 6 hours, and purifying a population of lymphocytes and antigen presenting cells for use in a cell-mediated immunoassay (CMI assay), by a method incorporating a positive or negative affinity selection step.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 April 2008 (10.04.2008)

PCT

(10) International Publication Number
WO 2008/041004 A3

(51) International Patent Classification:

G01N 1/34 (2006.01) **G01N 33/569** (2006.01)
A61K 35/14 (2006.01) **C12N 5/00** (2006.01)

(21) International Application Number:

PCT/GB2007/003800

(22) International Filing Date: 5 October 2007 (05.10.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

0619853.5 6 October 2006 (06.10.2006) GB

(71) Applicant (for all designated States except US): **OXFORD IMMUNOTEC LIMITED** [GB/GB]; 91 Milton Park, Abingdon, Oxfordshire OX14 4RY (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DURRANT, Ian** [GB/GB]; 46 Eskdale Road, Stoke Mandeville, Buckinghamshire HP22 5UJ (GB). **DAY, Toni** [GB/GB]; 30 Wilson Way, Kidlington, Oxford, Oxfordshire OX5 1TN (GB). **O'KEEFFE, Aisling** [IE/GB]; 4 Hiliat Fields, Drayton, Oxfordshire OX14 4JE (GB). **BAMPTON, Maxine** [GB/GB]; 22 Sandleigh Road, Wootton, Abingdon, Oxfordshire OX13 6DP (GB).

(74) Agents: **ROQUES, Sarah, Elizabeth** et al.; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5JJ (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

18 December 2008

(54) Title: PREPARATION METHOD

(57) Abstract: A method of storing and purifying cells from a whole blood sample, comprising storing the sample before and/or after purification of cells for at least 6 hours, and purifying a population of lymphocytes and antigen presenting cells for use in a cell-mediated immunoassay (CMI assay), by a method incorporating a positive or negative affinity selection step.



WO 2008/041004 A3

PREPARATION METHOD

Field of the Invention

The present invention relates to methods for storing and purifying cells from a whole blood sample. In particular, the invention relates to methods for storing and purifying cells from whole blood samples, such that a population of lymphocytes and antigen presenting cells is effectively stabilised and purified for use in assays which measure cell mediated immune responses *in vitro*.

Background to the Invention

Cell Mediated Immune (CMI) responses are commonly used to define the immune status of an individual. Typically, in the art of clinical immunology, the term CMI response encompasses *in vivo* skin testing, lymphocyte proliferation assays, and the *in vitro* detection of cytokines produced by peripheral blood mononuclear cells (PBMC) in the presence of a specific antigen. The invention described herein addresses improved methods for manipulating, stabilising and preparing cells derived from isolated whole blood samples for use in assay techniques designed to measure one class of CMI response, namely the *in vitro* cytokine-based CMI response to a specific antigen (hereinafter referred to as a "CMI Assay").

The cells of the immune system are capable of producing immune effector molecules such as cytokines following stimulation by an antigen. CMI Assays involve incubating a cell sample with an antigen and measuring for the presence or quantity of an immune effector molecule such as a cytokine to provide an indication of the ability of the individual to generate a cell-mediated immune response to the selected antigen. Cells for use in a CMI Assay also include isolated populations of lymphocytes (particularly T Cells) and Antigen Presenting Cells (APCs). APCs are involved in processing the antigen in order that the latter may be recognised by T Cell receptors on the surface of each T Cell. Antigen-induced cytokines may be released into the assay medium and detected directly by, for example, ELISA methods, or quantified in terms of the frequency of cytokine-secreting T Cells using ELISPOT methods.

The filter immunoplaque assay, otherwise called the enzyme-linked immunospot assay (ELISPOT), was initially developed to detect and quantitate

-2-

individual antibody-secreting B cells. At the time it was developed, the technique provided a rapid and versatile alternative to conventional plaque-forming cell assays. Recent modifications have improved the sensitivity of the ELISPOT such that cells producing as few as 100 molecules of a specific protein per second can be detected.

5 These assays take advantage of the relatively high concentration of a given proteinaceous cell product (such as a cytokine) in the environment immediately surrounding the protein-secreting cell. These cell products are captured and detected using high-affinity antibodies. The ELISPOT assay is reviewed in Current Protocols in Immunology, Unit 6.19 pages 6.19. 1-8.

10 The ELISPOT assay typically involves six steps: (1) coating a purified cytokine-specific antibody to a membrane-backed microtiter plate; (2) blocking the plate to prevent non-specific absorption of any other proteins; (3) incubating the cytokine-secreting cells with appropriate reagents; (4) removal of cells and reagents; (5) adding a labelled second anti-cytokine antibody; and (6) detecting the antibody-
15 cytokine complex on the membrane.

Current methods to prepare cells from a whole blood sample for use in a CMI assay involve the isolation of peripheral blood mononuclear cells (PBMCs) using a Ficoll gradient. In accordance with these methods, in order to be effective in the CMI assay, lymphocytes and APCs must be purified from the whole blood sample as soon
20 as possible, and in particular within 8 hours of collection of the blood sample from an individual.

Summary of the Invention

The present inventors have identified methods of storing and purifying cells
25 from a whole blood sample for use in cell mediated immunoassays (CMI assay), such as an ELISPOT assay. Methods are also identified for treating whole blood samples prior to storage. These methods allow whole blood samples to be stored for example for at least 6 hours, such as up to 48 hours before use in a CMI assay, such as an ELISPOT assay.

30 In accordance with the present invention, there is provided a method of storing and purifying cells from a whole blood sample, comprising storing the sample before and/or after purification of cells for at least 6 hours, and purifying a population

of lymphocytes and APCs for use in cell-mediated immunoassay (CMI assay), by a method incorporating a positive or negative affinity selection step.

In accordance with another aspect of the present invention, there is provided a method of treating and storing a whole blood sample comprising treating said whole blood sample by dilution with growth medium; and storing the treated whole blood sample for at least 6 hours, wherein after storage a population of lymphocytes and apcs can be obtained for use in a CMI assay. After storage, PBMCs or a suitable preparation of lymphocytes and APCs can be isolated from the diluted whole blood sample for use in a CMI assay, such as an ELISPOT assay.

10 In another aspect of the present invention, there is provided a method of maintaining the peptide specific cytokine response of isolated T cells derived from whole blood in an ELISPOT assay, comprising separating a population of cells comprising T cells and APCs from said whole blood sample using a positive or negative affinity selection.

15 In a further aspect of the present invention, there is provided a method of preparing lymphocytes and APCs suitable for use in an ELISPOT assay, wherein said method comprises removing red blood cells by filtration.

In a further aspect, there is provided a method of performing an *in vitro* cell-mediated immunoassay comprising:

- 20 (i) storing a whole blood sample from an individual for at least 6 hours;
- (ii) purifying a population of T cells and antigen presenting cells from granulocytes which are present in said stored sample by a method incorporating a positive or negative affinity selection step; and
- (iii) using said population of cells in a cell-mediated immunoassay to
- 25 measure antigen-specific responses in T cells.

In a further aspect, there is provided a kit for carrying out the above-mentioned method comprising:

- (i) a cytokine specific antibody attached to a microtiter plate, and
- (ii) one or more of an anti-CD15 ligand, an anti-CD4 ligand, an anti-CD8
- 30 ligand, an anti-CD19 ligand and an anti-CD14 ligand.

Detailed Description of the Invention

The present invention provides a method of storing and purifying cells from whole blood samples for use in a CMI assay such as an ELISPOT assay. The present invention also provides a method of treating a whole blood sample by dilution with
5 cell growth medium and storing the treated whole blood sample. The methods allow the peptide-specific cytokine response of isolated lymphocytes and APCs from said blood sample to be sufficient for use in a CMI assay such as an ELISPOT assay, and preferably to be useful in a diagnostic method.

In accordance with the present invention, a cell mediated immunoassay (CMI
10 assay) refers to an *in vitro* assay to measure the cytokine-based cell mediated immune response to a specific antigen. Such an assay uses a sample obtained from an individual, wherein the sample comprises cells of the immune system. The cells of the immune system are capable of producing immune effector molecules such as cytokines following stimulation by an antigen. Such assays include incubating the
15 sample with an antigen and measuring for the presence or quantity of the immune.

-4-

effector molecules such as cytokines to provide an indication of the ability of the individual to generate a cell mediated immune response to the selected antigen. In accordance with the present invention, the CMI assay is preferably an ELISPOT assay. Cells for use in the CMI assay such as an ELISPOT assay include isolated
5 populations of lymphocytes and APCs, such as a population of T cells and APCs.

The isolated preparation of lymphocytes and APCs and in particular T cells and APCs prepared in accordance with the present methods provides a preparation that can be used in a CMI assay such as an ELISPOT assay for use in diagnosis. Such preparations generate responses sufficient to allow diagnosis. Such responses
10 may be defined as 50% higher than background. Alternatively, such responses can be defined as being at least 50% of the maximal response that can be obtained using a freshly isolated whole blood sample.

For example, in a typical assay, 2.5×10^5 viable PBMCs may be used per well in an ELISPOT assay. In an ELISPOT assay for tuberculosis, typically, a negative
15 control will have less than 5 spots. In this case, a positive or reactive sample will have 6 or more spots more than the negative control. If the negative control has 6 or more spots, a reactive sample is indicated if it contains more than two times the negative control spot count.

In one embodiment the methods of the present invention use a positive
20 affinity selection step or a negative affinity selection step to purify cells from a whole blood sample, before or after storage. A positive affinity selection step is used to bind those cells of interest to a solid support and thus separate bound cells from non-bound cells. The purified bound cells are retained, and the non-bound cells are discarded.

25 In the alternative, a negative affinity selection step is used in order to remove those cells which are not required or undesirable in the cell preparation for use in the CMI assay, such as an ELISPOT assay. Thus the cells bound to the solid support are discarded and the purified unbound cells are recovered and retained.

The method will result in purification of both lymphocytes (such as T cells)
30 and APCs. The purification may comprise positive or negative affinity selection of both lymphocytes and APCs, in which case there will typically be an enrichment in the sample of both of these cell types, i.e. both the ratio of T cells to total number of

cells and the ratio of APCs to total number of cells will increase in the sample as a result of carrying out the method of the invention.

In accordance with one aspect of the invention, the whole blood sample is stored before and/or after purifying cells from the whole blood sample by a method including a positive or negative affinity selection step. For example, the whole blood sample may be stored prior to isolation of the selected cell population. After storage for at least 6 hours, at least 8 hours, up to 10, 12, 18, 24, 36 or 48 hours, the whole blood sample is treated (i.e. subjected to the affinity selection) to isolate the selected cell population, which can then be used in the CMI assay, such as an ELISPOT assay.

10 In the alternative, the whole blood sample is treated (i.e. subjected to the affinity selection) shortly after it has been obtained, for example, within 1 hour, up to 6 hours, or up to 8 hours after it has been obtained to isolate the selected cell population. Such a cell population is then stored prior to use in the CMI assay, such as the ELISPOT assay.

15 Preferably, the whole blood sample is stored between -5°C and 40°C, typically between 2°C and 8°C or between 18°C and 25°C. In a particularly preferred embodiment, the whole blood sample is stored at room temperature, such as at about 18°C or 20°C or from 18°C to 25°C or from 20°C to 25°C.

In one embodiment the sample is not frozen at any stage of any of the methods mentioned herein. In one embodiment the sample is not frozen between the stages of it being taken from the individual and being stored and purified (by a method of the invention); and/or the sample is not frozen between the stages of it being stored and purified and being used in a CMI assay.

20

In one aspect of the present invention, the purification method comprises a negative affinity selection step to remove granulocytes and optionally red blood cells from the whole blood sample in order to obtain the preparation of lymphocytes and APCs for use in an ELISPOT assay.

25

Thus, in accordance with the one aspect of the present invention, a method is provided in which both granulocytes and red blood cells are removed from the whole blood sample in a single step. In accordance with this method, the whole blood sample is contacted with an antibody preparation comprising anti-CD66b and glycophorin A antibodies. The antibodies serve to aggregate red blood cells and granulocytes. The aggregated red blood cells and granulocytes can be removed from

30

-6-

the sample, for example by centrifugation. The sample may also be subjected to a Ficoll gradient prior to centrifugation.

In another aspect of the invention, the negative affinity selection comprises the use of a solid support having a ligand bound thereto which binds to a cell surface protein present on the surface of granulocytes, which does not bind to cell surface proteins present on the surface of lymphocytes and APCs used in the ELISPOT assay. For example, a solid support can be provided comprising an anti-CD15 ligand to bind to CD15+ cells from the sample and/or anti-CD66b to remove CD66b+ cells from the sample.

10 In accordance with this aspect of the present invention, a whole blood sample, or a blood sample which has been treated to remove red blood cells is brought into contact with the solid support comprising an anti-CD15 ligand or anti-CD66b ligand, under conditions which allow binding of granulocytes. The preparation of lymphocytes or APCs having granulocytes removed therefrom can be obtained
15 directly, for example by contacting the sample with the solid support and collecting any material which does not bind the solid support.

In the alternative, the solid support having granulocytes or other unwanted cells bound thereto can be separated from the remainder of the sample. For example, the solid support can comprise beads such as magnetic beads. Once the beads have
20 been contacted with the sample and under conditions to allow binding of granulocytes to the solid support, the beads can be separated from the sample to leave a preparation of lymphocytes and APCs having granulocytes removed therefrom. In a preferred embodiment, the solid support comprises magnetic beads which can be separated from the remainder of the sample, for example by the application of a magnetic field.

25 It will be understood that other ligands, particularly antibodies, capable of binding to a cell surface marker on the surface of granulocytes, may be used for negative affinity selection. For example, the following antibodies may also be used to remove granulocytes by this method : anti-CD16b and / or anti-CD88.

Preferably, in accordance with this aspect of the present invention red blood
30 cells are also removed from the whole blood sample. This can be done before, after or at the same time as removal of CD15+ expressing cells or granulocytes.

-7-

Red blood cells can be removed from the sample by any suitable method. For example, red blood cells in the sample may be removed by lysis of the red blood cells (Simon et al., Immunol. Commun. 1983, Vol. 12, pp. 301-314).

5 In accordance with one aspect of the invention, red blood cells are removed by filtration. In accordance with this aspect of the invention, a method is provided for preparation of lymphocytes and APCs for use in a CMI assay such as an ELISPOT assay which comprises filtration to remove red blood cells. Such methods can be used on their own or with one or more of the other methods of the invention, such as negative affinity selection with anti-CD15+ ligands to remove granulocytes.

10 The filtration methods of the present application include applying a whole blood sample to a filter. The filters are selected such that red blood cells pass through the filter while lymphocytes and APCs are retained on the filter. In accordance with this method, the filters having lymphocytes and APCs thereon can be used directly in the CMI assay such as the ELISPOT assay. Alternatively, the lymphocytes and APCs
15 can be collected, for example by washing the filter or back flushing the filter and collecting the recovered lymphocytes and APCs.

In accordance with another aspect of the present invention, a preparation of lymphocytes and APCs are purified from a whole blood sample by a method involving a positive affinity step, wherein optionally the affinity selection step selects
20 all types of T cells (fore example irrespective of whether they are CD4 or CD8 or irrespective of the epitope which they recognise).

In one aspect, the whole blood sample is contacted with a solid support having attached thereto ligands which bind to cell surface proteins present on the surface of the lymphocytes and APCs. The ligands may be antibodies.

25 In a preferred aspect of the present invention, such ligands are selected from anti-CD4, anti-CD8, anti-CD19, anti-CD45, anti-CD45RC and anti-CD14 ligands. For example, a solid support can be provided to bind to CD4+, CD8+ CD14+ and CD19+ cells in the whole blood sample or a subset such as CD4+, CD8+ and CD19+ cells. Isolation of such cells provides a preparation of lymphocytes and APCs which
30 are suitable for use in the ELISPOT assay. Such a preparation may encompass a mixture of CD4+ and CD8+ T lymphocytes, B cells and monocytes. In one embodiment, anti-CD4 and/or anti-CD8 ligands are not used in the purification step,

-8-

and preferably anti-CD4 antibodies and/or anti-CD8 antibodies are not used in the purification step.

In a preferred embodiment, the ligands on the solid support are selected so that undesirable cells in the final preparation, such as granulocytes are not retained on the solid support. For example, the ligands are selected so that they are specific for lymphocyte and APCs and do not bind to granulocytes. The solid support can be provided with anti-CD4, anti-CD8 and anti-CD19 ligands or other appropriate ligands in approximately the same ratios. Alternatively, the ratio of each individual ligand can be selected, for example to reflect the proportion of each cell present in whole blood. In the alternative, the ratio of ligands, such as CD4:CD8:CD19 ligands can be selected to bind and isolate lymphocytes and APCs in a desired ratio, such as one particularly suitable for use in an ELISPOT assay.

Methods of affinity selection using solid supports having ligands bound thereto to bind to selected proteins are well known in the art. In general, a whole blood sample is brought into contact with the solid support comprising the ligands of interest, under conditions which allow binding of cells to the ligands via the relevant cell surface protein. The solid support can be separated from the remainder of the sample in order to separate bound cells from non-bound material. Washing steps may be included, for example to rinse non-bound material from the solid support.

In a preferred aspect of the present invention, the solid support comprises magnetic beads. Such magnetic beads can easily be separated from a sample, for example by the application of a magnetic field. Where the solid support is provided in the form of magnetic beads or other types of bead, each bead may have a single ligand present thereon, such as anti-CD4 ligand. Beads which individually carry each ligand, such as anti-CD4, anti-CD8, anti-CD14 or anti-CD19 can be combined together to produce a solid support which will bind to the desired cells, such as CD4+, CD8+, CD14+ and CD19+ cells. In the alternative, each magnetic bead may be provided with more than one ligand such as both anti-CD4 and anti-CD8 ligands or anti-CD4 and anti-CD19 ligands or a combination of all ligands. In a particularly preferred embodiment, the ratio of ligands, such as anti-CD4: anti-CD8: anti-CD19 is selected to correlate to the ratio of lymphocytes and APCs desired in the final preparation for use in the ELISPOT assay. The quantity of magnetic beads can also

-9-

be selected in order to isolate a selected number of cells, to facilitate further processing of the preparation for use in the ELISPOT assay.

The preparation of selected cells, and in particular lymphocytes and APCs that are bound to the solid support can be used in an assay directly. In particular,
5 magnetic beads having bound thereto the relevant cells can be used in a CMI assay, such as an ELISPOT assay. Preferably, the beads having cells bound thereto are washed prior to use in the assay. In the alternative, once the selected cells have been isolated from the remainder of the whole blood sample, the cells bound to the solid support can be dissociated from the solid support for use in the assay.

10 In accordance with another aspect of the present invention, a whole blood sample is treated by dilution with cell growth medium and stored before being used in the CMI assay such as an ELISPOT assay. The diluted whole blood sample can be stored for at least 6 hours after collection for example for at least 8 hours after collection from an individual, preferably up to 10, 12, 18, 24, 36 or 48 hours after
15 collection.

In accordance with this aspect of the present invention, the whole blood sample is stabilized or treated prior to storage. For example, the method comprises diluting a whole blood sample with cell growth medium, such that when the whole blood sample or an isolated preparation of T cells and APCs obtained from said
20 stored whole blood sample is used in an ELISPOT assay up to 48 hours after collection of the sample a viable response is maintained.

In accordance with the present invention, the whole blood sample can be diluted with cell growth medium in any appropriate ratio. In a preferred embodiment, the ratio of whole blood sample to cell growth medium is 1:1. Alternatively, the ratio
25 of whole blood sample to cell growth medium can be 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, up to 10:1, 20:1 or 30:1. Alternatively, the ratio of whole blood sample to cell growth medium can be 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9 or 1:10, up to 1:15, 1:20 or 1:30.

In accordance with one embodiment, the diluting medium is added to the
30 whole blood sample, but is not subjected to shaking or mechanical mixing. The whole blood sample is preferably stored in a standard heparinized tube. Preferably, the whole blood sample is stored in the dark.

-10-

The cell growth medium for use in the invention can be any appropriate cell growth medium. For example, the medium can be a commercially available medium such as AIM-V (Invitrogen Paisley UK), or RPMI 1640TM (Sigma Aldrich Corp, St. Louis, MO, USA). In a preferred embodiment, the cell growth medium is serum-free medium, such as AIM-V.

In accordance with one aspect of the invention said treated whole blood sample is used or prepared for use in a CMI assay after storage. Typically, a preparation of PBMCs, or T lymphocytes and APCs are purified from said treated whole blood sample after storage for use in the CMI assay. Any suitable method can be used to obtain the isolated preparation of cells for use in a CMI assay, such as an ELISPOT assay. Typically, PBMCs can be isolated, for example, using a Ficoll gradient. More preferably, red blood cells and granulocytes are removed from the treated whole blood sample, for example to isolate a preparation of lymphocytes and APCs, by a method including an affinity selection step.

Treatment by dilution may also be used in another aspect of the invention to treat an isolated preparation of lymphocytes and APCs prior to storage of the preparation before its use in a CMI assay.

In a further aspect of the invention, whole blood may be stored, with or without dilution, for several hours, preferably for at least 6 hours, or up to 12 hours, or from 12-48 hours. Such storage may be at 2-8 ° C, or at room temperature. Following storage, a preparation of lymphocytes and APCs may be prepared by affinity selection as described above, and optionally the peptide specific response of T cells in the preparation may be measured in a CMI assay. In this manner, the *in vitro* activity of T cells introduced into the CMI assay is stabilised and maintained. Accordingly, this approach reduces or obviates a marked reduction in T Cell activity *in vitro* which is observed in preparations of lymphocytes and APCs isolated using, for example, standard Ficoll gradients from a whole blood sample, when the blood sample is stored for several hours prior to the isolation step.

The present invention also provides a kit for storing a whole blood sample comprising a standard heparinized tube and cell growth medium. The invention also provides a kit for carrying out an ELISPOT assay on cells which have been stored and purified by a method as disclosed herein, wherein the kit comprises:

-11-

- (i) a cytokine-specific antibody, optionally attached to a microtiter plate, and
- (ii) one or more of the ligands mentioned herein which can be used in affinity selection, optionally bound to a solid support, and
- 5 (iii) optionally instructions for carrying out the ELISPOT assay and/or method of storing and purifying the cells.

Further, the invention also provides a kit for purifying a population of lymphocytes and APCs according to the invention, wherein the kit comprises one or more of the ligands mentioned herein, optionally bound to a solid support. The kit
10 may also comprise instructions for carrying out the method in accordance with the invention.

In other embodiments of the invention, the whole blood sample is treated using a positive or negative affinity selection step to isolate a population of T cells and APCs for use in the CMI assay, such as an ELISPOT assay without storing the
15 samples. In a further embodiment, the positive or negative affinity selection methods may be used to isolate a preparation of cells for use in a CMI assay such as an ELISPOT assay, such cells being derived from whole blood samples which have been treated by dilution and stored.

Any of the above embodiments may be carried out on a sample from a human,
20 such as a human that is suspected of having a disease which can be diagnosed by ELISPOT.

The invention is hereinafter described in more detail by reference to the following Examples.

25 Example 1 – Treatment by Dilution with Culture Medium

Immediately after venipuncture, one 10ml Lithium Heparin Vacutainer (BD Biosciences, Oxford, UK) of whole blood from each of a total of 78 donors and was processed over the space of 6 months, using the T-SPOT.*TB* assay kit (Oxford Immunotec, Abingdon, UK). The method was carried out according to the
30 manufacturer's instructions accompanying the kit. This method comprises : a) a standard Ficoll method for preparation of Peripheral Blood Mononuclear Cells (PBMC) – see under Sample Collection and Preparation, Procedure 2 “alternative blood collection methods” and Note 2.; b) viable cells were counted using the Trypan

-12-

Blue Dye Exclusion method (Freshney, R. (1987) Culture of Animal Cells: A Manual of Basic Technique, p.117, Alan R. Liss, Inc., New York) ; c) cells were diluted in GIBCO AIM-V (Invitrogen, Paisley, UK) to give 250,000 cells per 100 μ l. ; d) 250,000 (100 μ l) of the Ficoll-extracted cells were added to each microtiter plate well pre-coated with anti-gamma interferon, together with one of the following :
5 phytohaemagglutinin (PHA) positive control; or specific test antigens from M. tuberculosis (Panel A; Panel B); or from cytomegalovirus / epstein barr virus / influenza virus (CEF; Mabtech, Sweden), which antigen is not provided in the T-SPOT.TB kit; e) the plate was incubated for 16 to 20 hours (37°C, 5% CO₂), washed
10 four times with phosphate buffered saline (PBS) before adding an alkaline phosphatase-conjugated secondary antibody and incubating for 1 hour at 2-8°C; f) the wells were washed four times, and 5-bromo-4-chloro-3'-indoylphosphate/Nitro Blue Tetrazolium (BCIP/NBT) then added and incubated for 7 minutes. After the spots had developed, the reaction was stopped using distilled water; g) after drying at 37°C
15 for 4 hours, the number of spot forming cells (SFC) for each well was recorded and analysed.

The remaining blood from each donor was diluted 1:1 in AIM-V and stored at room temperature (18-25°C) in the dark overnight for 24 hours.

On Day 2, the T-SPOT.TB assay was performed on the stored blood including
20 the standard Ficoll method for preparation of PBMC. The T-SPOT.TB assay was then conducted to determine the SFC counts, using a positive control (PHA) and a nil control (lacking antigen). The nil control was subtracted from the test antigen well SFC count. If any of the antigen wells contained six or more spots and the nil control contains zero spots, then the sample was deemed 'reactive' for that antigen under test.
25 The Tables below show the relationship between fresh and diluted and stored blood. Reactive and non-reactive results are shown using Panel A, Panel B and CEF wells for each donor blood sample both before and after overnight blood storage (following 1:1 dilution in AIM-V) at room temperature (RT). The percentage of the total number of donors is also indicated. CEF antigen was dissolved in AIM V, and was
30 present in the assay at a level of 5 μ g / ml.

Panel A Antigen		
	Fresh Blood	
Diluted / Stored Blood	Reactive	Non-Reactive
Reactive	3 (3.8%)	1 (1.3%)
Non-Reactive	2 (2.6%)	72 (92.3%)

96.2% clinical concordance

Panel B Antigen		
	Fresh Blood	
Diluted / Stored Blood	Reactive	Non-Reactive
Reactive	5 (6.7%)	0 (0%)
Non-Reactive	3 (4%)	67 (89.3%)

96% clinical concordance

CEF Antigen		
	Fresh Blood	
Diluted / Stored Blood	Reactive	Non-Reactive
Reactive	28 (77.8%)	0 (0%)
Non-Reactive	5 (13.9%)	3 (8.3%)

5

Results demonstrate 86.1% clinical concordance (assuming that 6 spots greater than nil control indicates reactivity).

Thus, the addition of medium to the blood sample enables a positive SFC response to be measured in PBMC after 24 hours storage, a response that is similar to that seen when measured on blood processed immediately after collection. Blood collected and stored for 24 hours without the addition of medium demonstrated a significant loss in the measurable CMI response (results not shown).

10

Example 2 – Positive Affinity Selection

15

On Day 1, and within 2 hours of venipuncture, two 10ml Lithium Heparin Vacutainers of whole blood from each of nine donors were placed in storage, one in the dark at RT (18-25°C) and the other in the fridge (2-8°C).

20

At each time point, either 0-2 hours (fresh) or 24 hours (stored) post venipuncture for each donor, 3 mls of whole blood were diluted 1:1 with wash buffer (PBS/0.1% BSA/ 2mM EDTA) and mixed with 80 µl of a cocktail of specific magnetic Dynabeads, namely 111.45 (CD4); 111.47 (CD8); 111.49 (CD14); and 111.43 (CD19), which were mixed in equal proportions (Invitrogen). These ‘beads’

-14-

positively select cells based on the presence of specific differentiation (CD) markers :
 CD4 and CD8 for T cell subsets; CD14 (monocytes) and CD19 (B cells and
 APC's). The samples were mixed at RT for 12 mins, and the beads were separated
 by placing sample tubes in a magnetic particle concentrator (MPC: Invitrogen,
 5 Paisley, UK) for 2 mins. The supernatant was discarded, and the beads were washed
 twice in PBS and then separated again on the MPC. The beads were then
 resuspended in 1ml AIM-V and cells were counted using the Trypan Blue Dye
 Exclusion Method. Cells were diluted in AIM-V to give 250,000 cells per 100 μ l.

In parallel, at each time point, either 0-2 hours (fresh) or 24 hours (stored)
 10 post-venipuncture for each donor, 5 ml of whole blood was processed immediately
 using the standard Ficoll method as indicated in Example 1.

All processed samples were then assayed using the T.SPOT.TB assay kit (see
 Example 1), using antigens CEF (5 μ g / ml), or PPD (1 μ g / ml), or PPD / CEF (1 μ g
 / ml and 5 μ g / ml respectively) in the assay wells, none of which are provided in the
 15 T-SPOT.TB kit; PPD (Product Code RT23) was obtained from Statens Serum Institut
 (Copenhagen, Denmark); for CEF, see Example 1. PPD and CEF were used to
 identify CD4+ and CD8+ T cells respectively.

The following Table, incorporating the average of the results for all donors,
 displays the SFC counts measured by ELISPOT for cells selected from fresh or stored
 20 blood using CD4, CD8, CD14 & CD19-conjugated magnetic beads compared to use
 of Ficoll only. The blood stored at RT for 24 hours and selected using beads has
 produced comparable results to the 'fresh bead' samples, and yield better results than
 when 'Ficoll only' samples are stored at RT, and both 'Ficoll only' and 'beads' are
 stored at 4°C .

25

Antigen	Fresh		Stored (24 Hours)			
	Ficoll	Beads	Ficoll (4°C)	Ficoll (22°C)	Beads (4°C)	Beads (22°C)
PPD	100	78	15	41	53	70
CEF	100	194	13	45	111	149
PPD/CEF	100	136	14	43	82	109
PPD		71.40	18.21	35.06	67.34	49.67
CEF		109.20	18.51	37.24	87.54	109.40
PPD/CEF		81.80	1.52	2.85	40.89	56.25

-15-

The results were normalised, assuming that 'fresh' samples, processed with Ficoll only, yield an SFC count of 100 % in the ELISPOT assay. The top three rows show the mean SFC counts, and the bottom three rows show the standard deviations.

5 Example 3 – Negative Affinity Selection

Immediately after venipuncture, two 10 ml Lithium Heparin Vacutainers of whole blood were collected from each of 4 donors over the space of 1 month.

For one sample from each donor, a negative selection step was undertaken using MACSi anti-CD15 magnetic beads (Miltenyi Biotech, Bergisch Gladbach, Germany). The beads were resuspended thoroughly before use to ensure a homogeneous dispersion. 400 µl of beads were added to 4 ml of whole blood in a 15 ml conical tube. The tubes were incubated at RT using the MACSimix (Miltenyi Biotech) tube rotator at medium speed (8 rpm) for 15 minutes. The tubes were placed in the MACSiMag separator (Miltenyi Biotech), and the beads allowed to adhere to the wall of the tube for 2 minutes. Retaining the tube in the magnet, the supernatants were removed using a pipette and retained in a fresh tube. The tubes containing the supernatants were again placed in the MACSiMag separator and any residual beads allowed to adhere to the wall of the tube for 2 minutes. Retaining the tube in the magnet, the supernatant was transferred to a 50 ml conical tube. 16ml of freshly prepared 1X red blood cell (RBC) lysis buffer (prepared by dilution of 10X lysis buffer: 1.55M NH₄Cl, 100mM KHCO₃, 10mM EDTA) were added and the tubes incubated at room temperature for 5 minutes. The samples were centrifuged at 300xg for 10 minutes. The pelleted cells were washed twice in lysis buffer (5 ml) and centrifuged each time at 300xg for 5 minutes. The cells were washed once in RPMI and diluted in 1ml of AIM-V for use in an ELISPOT assay.

In parallel with this method, and using a second sample from each donor, 5ml of whole blood was processed by the standard Ficoll method as indicated in Example 1.

Purified PBMC samples were then assayed in the T.SPOT.TB assay, as outlined in Example 1, using CEF (5 µg/ml) as antigen for Donor 1,2 and 4 samples; and Panel A and B antigens (33 µg/ml) for the Donor 3 samples.

The SFC count obtained with PBMCs (shown in the Tables below) prepared using a negative selection approach with MACSi anti-CD15 magnetic beads

-16-

demonstrated high equivalence to those PBMCs that were prepared using Ficoll gradients. This antibody-based method of PBMC isolation proved to be technically feasible and produced highly reproducible results in the T-SPOT.*TB* assay.

		Donor 1		Donor 2	
		Ficoll	MACSi	Ficoll	MACSi
CEF	-	240	239	100	111
	+	248	235	98	124
Mean		244	237	99	117.5
Std dev		5.66	2.83	1.41	9.19

		Donor 3	
		Ficoll	MACSi
-	-	0	0
	+	>100	>100
panel A	-	41	31
	+	45	42
	+	44	38
panel B	-	179	99
	+	157	106
	+	151	83
Mean			
Panel A		43.3	37
Panel B		162.3	96
St Dev			
Panel A		2.08	5.57
Panel B		14.74	11.79

		Donor 4	
		Ficoll	MACSi
-	-	0	0
	+	0	0
CEF	-	116	100
	+	116	100
	+	187	240
CEF	-	210	214
	+	180	189
	+	180	189
Mean			
-		0	0
+		116	100
CEF		192.3	214.3
St Dev			
-		0	0
+		0	0
CEF		15.70	25.50

(N.B. For Donors 3 and 4, (-) wells indicate absence of antigen; (+) wells indicate presence of PHA (5 µg / ml).

Example 4 – Negative affinity Selection

Two separate experiments were performed, each using pooled whole blood from a group of either 15 or 11 donors.

For each experiment : a) on Day 1, immediately after venipuncture, one 10ml Lithium Heparin Vacutainer of whole blood from each donor pool was processed according to the standard Ficoll method (see Example 1). Viable PBMC were counted using the Trypan Blue Dye Exclusion Method. Cells were diluted in cell culture medium (AIM-V) to give 250,000 cells per 100 µl; b) the remaining blood was stored in the dark, undiluted, either in a fridge at 2-8°C or at RT (18-25°C) for 24 hours; c) on Day 2, stored blood samples were processed according to the standard Ficoll method for PBMC preparation , and viable cells were counted and diluted as described above (see paragraph 1, this Example).

In parallel samples processed in each experiment, both on Day 1 (fresh) and Day 2 (stored) for each donor, 4.5 mls of whole blood was added to 15 ml centrifuge tubes, and 225 µl of RosetteSep (StemCell, Vancouver, BC, Canada) was then added to each tube to bind the RBC to the granulocytes using a tetramer complex containing

-17-

anti-CD66b and glycophorin A antibodies. Samples were mixed gently and left to incubate for 20 minutes at RT. Samples were then diluted 1:1 with RPMI, and the PBMCs were isolated from the granulocyte-depleted cells by the standard Ficoll method (see Example 1), thus allowing the RBC and granulocytes to be separated from lymphocytes and APCs. The latter cells were then washed and processed according to the manufacturer's instructions accompanying the kit – see Example 1.

Results are shown below. In the two separate experiments performed, one used CEF (5 µg/ml) as antigen, and the other used PPD (1 µg/ml) in the assay. Top Table: CEF-induced SFC counts (mean of 15 donors) for Ficoll and RosetteSep treatment at 4°C and RT. Bottom Table: PPD-induced SFC counts (mean of 11 donors) for Ficoll and RosetteSep treatment following storage at 4°C and RT. The results demonstrate that samples stored at 4°C and then subjected to anti-granulocyte antibody treatment followed by Ficoll purification of PBMC are equivalent to standard 'fresh' samples, and also yield higher SFC counts than when samples are stored at RT prior to anti-granulocyte antibody treatment.

CEF	Fresh Ficoll	Ficoll, 4°C	RosetteSep, 4°C	Ficoll, RT	RosetteSep, RT
Mean	80	54	78	39	60
sum	4870	3252	4759	2388	3681
% diff from fresh Ficoll		-33.2	-2.3	-51	-24.4

n = 15 donors, 61 replicates

PPD	Fresh Ficoll	Ficoll, 4°C	RosetteSep, 4°C	Ficoll, RT	RosetteSep, RT
Mean	22.6	9.4	21.4	8	17.4
sum	722	301	685	256	558
% diff from fresh Ficoll		-58.31	-5.12	-64.54	-22.71

n = 11 donors, 32 replicates

Example 5 - Filtration

The feasibility of using back flushed leukocyte filter membranes (PALL Medical) and also the stability of the cells captured on such membranes was investigated at T0 and T24 hours post venipuncture. The captured cells were stored on the membranes in a fridge (2-8°C) from one donor and RT (18-25°C) for the second donor. This membrane separation method was performed along side a standard Ficoll PBMC separation method at each time point and temperature.

The membranes that were required to be stored followed steps 1-3 (see below) and then the appropriate storage condition and duration subjected on them. The

membrane holder devices were then removed from the storage area and subjected to steps 4-7 (see below) at the relevant time point. The Ficoll control storage samples were kept in the dark at RT and were diluted 1:1 in AIMV cell culture media until required for processing via the standard Ficoll separation method. The membrane procedure is described below:

Steps 1-3 consisted of; 10 ml of WASH BUFFER 1 being filtered through the membrane, using the transfer syringe, into the waste bottle. 10ml of whole blood sample together with 50 ml of WASH BUFFER 1 was mixed first and then filtered through the LK4 membrane, using the transfer syringe, in to a waste bottle. 50 ml of cooled WASH BUFFER 1 was filtered through the LK4 membrane, using the transfer syringe, into the waste bottle.

Steps 4-7 consisted of: the filter membrane being inverted either immediately or post 24h (depending on time point). 50 ml of cooled REMOVING BUFFER 1 was filtered back through the membrane, using the back flush syringe, into a 50 ml falcon tube. The cell suspension was spun at 2000 rpm for 5 mins at RT and then re-suspended in 10mls of AIMV. The cells suspension was again spun at 2000 rpm for 5 mins at RT and then re-suspended in 1 ml of AIMV media and counted according to the standard Ficoll separation method. (WASH BUFFER 1: 500 ml AIMV media buffered with 20mM HEPES and 1% [w/v] Dextran T40) (REMOVING BUFFER 1: 500 ml AIMV media buffered with 20 mM HEPES and 0.1 % [w/v] Dextran T40).

The T-SPOT.*TB* assay was then performed as indicated in Example 1 and also in the manufactures guidelines accompanying every kit.

The following table describes a summary of results from the T-SPOT.*TB* tests carried out for this investigation. A comparison of the CEF spot counts with regard to time and temperature of the membrane experiments is shown. This was carried out in terms of the membranes performance in relation to the relevant Ficoll controls carried out on the same day and also their relation at T0.

	Date	Donor	Method	Temp	CEF Count (ave, n=4)	Percentage of Ficoll Control	Percentage of Ficoll control at T0
T0	04/05/2006	0002	Ficoll	(RT)	45.5	~	~
			LK4	2-8oC	11	24.2	~
		0008	Ficoll	(RT)	44.4	~	~
			LK4	RT	21.25	47.9	~

-19-

T24	05/04/2006	0002	Ficoll	(RT)	47.5	~	104.4
			LK4	2-8oC	10.5	22.1	23.1
		0008	Ficoll	(RT)	33.75	~	76.0
			LK4	RT	4.75	14.1	10.7

It can be seen from the above table, the membranes used with fresh whole blood, stored at 2-8°C and 18-25°C (RT) at both T0 and T24 have provided PBMCs which produce sufficient CEF signal for the use in the ELISPOT assay. The concept of using back flushed filter membranes to harvest PBMCs capable of eliciting a response in the ELISPOT assay has been proven in this case.

Example 6 – Storage Time Before Negative Affinity Selection

Two separate experiments were performed, using samples of whole blood from each of 15 donors.

For each experiment: a) on Day 1, immediately after venipuncture, one 10ml Lithium Heparin Vacutainer of whole blood from each donor was processed ('fresh' samples), according to the standard Ficoll method for PBMC preparation; b) the remaining blood from each donor was stored in the dark, undiluted, in a fridge at 2-8°C (stored samples) for various times (16 to 72 hours); c) on Day 2, the stored samples were incubated with RosetteSep (StemCell, Vancouver, BC, Canada) as described in Example 4, then processed according to the standard Ficoll method for PBMC preparation. Counting and dilution of PBMC was then carried out, and samples processed, according to Example 4, except that in the two separate experiments performed, one used CEF (5 µg/ml) as antigen, and the other used CEF / PPD (5 µg/ml and 1 µg/ml respectively) in the assay.

The results shown below represent the cumulative SFC counts recorded for all individual donor samples, after each had been subjected to varying processing conditions, using either CEF antigen (n=10) or using both CEF and PPD antigens (n=5), based on summing the mean of the result of three separate ELISPOT assays for each processed sample.

The data demonstrate that samples stored for between 16 and 24 hours at 2-8°C can subsequently be 'stabilised' with anti-granulocyte antibody prior to being processed with Ficoll, whereas longer storage times followed by the same purification conditions give rise to a gradual reduction in SFC counts in the ELISPOT assay.

-20-

Antigen	Fresh Ficoll	RosetteSep, Stored at 2-8° C				
		16 hrs	20 hrs	24 hrs	48 hrs	72 hrs
CEF	940	900	846	892	661	354
CEF / PPD	481	502	476	584	324	108

Similar results are obtained when samples are stored at RT for up to 48 hours prior to being 'stabilised' and processed with Ficoll.

CLAIMS

1. A method of performing an *in vitro* cell-mediated immunoassay comprising:
 - (i) storing a whole blood sample from an individual for at least 6 hours;
 - (ii) purifying a population of T cells and antigen presenting cells from granulocytes which are present in said stored sample by a method incorporating a positive or negative affinity selection step; and
 - (iii) using said population of cells in a cell-mediated immunoassay to measure antigen-specific responses in T cells.
2. The method according to claim 1, wherein said sample is stored for at least 10 hours.
3. The method according to claim 2, wherein said sample is stored for at least 12 hours.
4. The method according to claim 3, wherein said sample is stored between 12 and 36 hours, 12 to 48 hours or 24 to 48 hours.
5. The method according to any one of claims 1 to 4, wherein said sample is stored between 2-8°C.
6. The method according to any one of claims 1 to 4, wherein said sample is stored between 18-25°C.
7. The method according to any one of claims 1 to 6, wherein said purification method comprises a negative affinity selection step to remove granulocytes.
8. The method according to claim 7, wherein said method comprises contacting the sample with an antibody preparation comprising anti-CD66b and glycophorin A antibodies to aggregate red blood cells and granulocytes, and removing said red blood cells and granulocytes from the preparation by centrifugation.

9. The method according to claim 7, wherein said granulocytes are removed by contacting said sample with a solid support comprising an anti-CD15 ligand to remove CD15+ cells from the sample.
10. The method according to claim 9, wherein said solid support comprises magnetic beads and wherein the method comprises contacting the sample with the magnetic beads to bind CD15+ cells to the beads, and separating the beads having the CD15+ cells bound thereto from the sample.
11. The method according to any one of claims 9 to 10, wherein red blood cells in the sample are removed by lysis or by filtration.
12. The method according to any one of claims 1 to 11, wherein the purification leads to an increase in the sample of both the ratio of T cells to total cells and the ratio of antigen presenting cells to total cells; and/or wherein the sample is not frozen at any point before storage or purification or use in a CMI assay.
13. The method according to any one of claims 1 to 6, that comprises a positive affinity selection step for said cells.
14. The method according to claim 13, wherein the affinity selection step selects all types of T cells.
15. The method according to claim 13 or 14, wherein said method comprises contacting said sample with a solid support having attached thereto ligands which bind to cell surface proteins present on the surface of said T cells and antigen presenting cells to thereby isolate the preparation of T cells and antigen presenting cells.
16. The method according to claim 15, wherein said solid support comprises one or more of anti-CD4, anti-CD8, anti-CD19 and anti-CD14 ligands to isolate CD4+, CD8+, CD19+ and/or CD14+ cells from the whole blood sample.

-23-

17. The method according to claim 15 or 16, wherein said preparation of T cells and antigen presenting cells are separated from the solid support prior to use in an assay.
18. The method according to claim 15 or 16, wherein said preparation of T cells and antigen presenting cells are retained on said solid support for use in an assay.
19. The method according to any one of claims 15 to 18, wherein said solid support comprises magnetic beads.
20. The method according to claim 19, wherein each said magnetic bead has bound thereto an anti-CD4, anti-CD8 and anti-CD19 ligand.
21. The method according to claim 19, wherein each magnetic bead has bound thereto one or more of anti-CD4, anti-CD8 or anti-CD19, and said magnet beads are mixed together to provide the solid support binding to CD4+, CD8+ and CD19+ cells.
22. The method according to any one of claims 1 to 21, wherein the cell-mediated immunoassay is an ELISPOT assay.
23. A kit for carrying out the method of claim 22 comprising:
 - (i) a cytokine specific antibody attached to a microtiter plate, and
 - (ii) one or more of an anti-CD15 ligand, an anti-CD4 ligand, an anti-CD8 ligand, an anti-CD19 ligand, and an anti-CD14 ligand.
24. The kit of claim 23, further comprising instructions for carrying out said ELISPOT assay.