(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
22 February 2024 (22.02.2024)

WIPO | PCT

(10) International Publication Number
WO 2024/040249 A1

(51) International Patent Classification:
C07K 14/555 (2006.01) *C07K 14/715* (2006.01)

(21) International Application Number:
PCT/US2023/072516

(22) International Filing Date:
18 August 2023 (18.08.2023)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

63/399,048	18 August 2022 (18.08.2022)	US
63/383,797	15 November 2022 (15.11.2022)	US
63/481,312	24 January 2023 (24.01.2023)	US

(71) Applicant: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).

(72) Inventors: **WEICK, Eva-Maria**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **BLOCH, Nicolin**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **GARG, Vidur**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **ULLMAN, Erica**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **ZHANG, Tong**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **LIN, Chia-Yang**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **WU, Jiaxi**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **SMITH, Eric**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).

(74) Agent: **BRUNER, J. Kyle** et al.; BIOSPARK INTELLECTUAL PROPERTY LAW, 1 Broadway, 14th Floor, Cambridge, Massachusetts 02142 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CV, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, MG, MK, MN, MU, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, CV, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SC, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, ME, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:
— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:
— *with international search report (Art. 21(3))*
— *with sequence listing part of description (Rule 5.2(a))*

WO 2024/040249 A1

(54) Title: INTERFERON RECEPTOR AGONISTS AND USES THEREOF

(57) Abstract: The present disclosure provides interferon receptor agonists with improved safety profiles and therapeutic indices. The interferon receptor agonists are attenuated through masking and/or reduced receptor binding as compared to a wild-type interferon. IFN receptor agonists optionally further comprise a targeting moiety, *e.g.*, a targeting moiety that recognizes a tumor- or immune cell-associated antigen and directs the interferon receptor agonist to a tumor site and/or tumor-reactive immune cells. The disclosure further provides pharmaceutical compositions comprising the interferon receptor agonists, and methods of use of the interferon receptor agonists in therapy, as well as nucleic acids encoding the interferon receptor agonists, recombinant cells that express the interferon receptor agonists and methods of producing the interferon receptor agonists.

DESCRIPTION

INTERFERON RECEPTOR AGONISTS AND USES THEREOF

1. CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of U.S. provisional application no. 63/399,048, filed August 18, 2022, U.S. provisional application no. 63/383,797, filed November 15, 2022, and U.S. provisional application no. 63/481,312, filed January 24, 2023, the contents of each of which are incorporated herein in their entireties by reference thereto.

2. SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically and is hereby incorporated by reference in its entirety. Said copy, created on August 15, 2023, is named RGN-024WO_SL.xml and is 386,251 bytes in size.

3. BACKGROUND

[0003] Type I interferons (IFNs) are thought to directly suppress tumor cell proliferation. Type I IFNs have utility in treatment of several types of cancer, including hematological tumors (chronic myeloid leukemia, hairy cell leukemia, multiple myeloma, and non-Hodgkin's lymphomas) and solid tumors (melanoma, renal carcinoma, and Kaposi's sarcoma). See, *e.g.*, Zitvogel *et al.*, 2015, Nat Rev Immunol 15:405-414 and Antonelli *et al.*, 2015, Cytokine Growth Factor Rev 26:121-131.

[0004] A particular advantage of Type I IFN treatment is its ability to intervene at multiple points in the generation of anti-tumor immune responses, including stimulation of the innate and adaptive cytotoxic lymphocyte populations, negative regulation of suppressive cell types, its impact on tumor cells by inhibiting proliferation, and by modulating apoptosis, differentiation, migration and cell surface antigen expression (Parker *et al.*, 2016, Nature Reviews Cancer 16:131–144).

[0005] One of the biggest barriers to the use of Type I IFNs in the clinic is the severe side effects associated with such treatments. The most frequently encountered side effects are flu-like symptoms, hematological toxicity, elevated transaminases, nausea, fatigue, and psychiatric sequelae. These side effects hamper reaching and maintaining the doses

needed for maximal therapeutic effect, and their occurrence can outweigh clinical benefit of Type I IFN treatment entirely (Lotrich, 2009, Dialogues Clin Neurosci 11:417-425). Type I IFNs signal through IFNAR1/IFNAR2 complex that are expressed on most cells and tissues in the body. Therefore, the ability to preferentially or specifically deliver active Type I IFNs to tumor-reactive immune cells (see, *e.g.*, Diamond *et al.*, 2011, J Exp Med. 208(10):1989–2003; Cauwels *et al.*, 2018, Cancer Res. 78 (2): 463–474) or to the tumor microenvironment is imperative for continued clinical use of Type I IFN. Strategies are needed to modify Type I IFN in order to obtain new forms of the drug which preferentially exert their activity on tumor-reactive immune cells and/or at the tumor, and also to reduce side effects on normal IFNAR-expressing cells.

[0006] Thus, there is a need in the art for novel Type I IFN therapies with improved therapeutic efficacy and safety profiles.

4. SUMMARY

[0007] The present disclosure relates to Fc-coupled IFN receptor agonists with IFN sequences (IFN moieties) having improved safety profiles as compared to wild-type Type I interferon molecules by virtue of attenuation by (i) masking, *e.g.*, masking with an interferon receptor (IFNR) moiety; (ii) one or more mutations in the IFN sequence, *e.g.*, one or more amino acid substitutions and/or truncations; (iii) use of native IFN sequences with low receptor affinities (*e.g.*, native IFN sequences with lower receptor affinities than IFNα2b and/or IFNβ); or (iv) any combination of two or all three of (i), (ii) and (iii).

[0008] The IFN receptor agonists may further comprise, *e.g.*, N-terminal to one or both Fc domains, a targeting moiety (or a component thereof, *e.g.*, one chain of a Fab). The targeting moiety comprises an antigen-binding domain (ABD) that can, for example, bind to a target molecule present on the tumor surface (*e.g.*, a tumor associated antigen) or other component in the tumor microenvironment (*e.g.*, extracellular matrix (ECM) or tumor lymphocytes), dendritic cells or natural killer cells.

[0009] The IFN receptor agonists, particularly those with an IFNR masking moiety, may be activatable by virtue of inclusion of one or more protease-cleavable linkers whose cleavage (*e.g.*, by a protease in the tumor environment) release the IFN moiety from the masking moiety.

[0010] Exemplary IFN moieties that can be used in the IFN receptor agonists of the disclosure are described in Section 6.3.

[0011] Exemplary masking moieties that can be used in the IFN receptor agonists of the disclosure are described in Section 6.4.

[0012] Protease-cleavable linkers that can be used in the IFN receptor agonists of the disclosure are described in Section 6.5.

[0013] Non-cleavable linkers that can be used in the IFN receptor agonists of the disclosure are described in Section 6.6.

[0014] Targeting moieties that can be used in the IFN receptor agonists of the disclosure are described in Section 6.7 and targeting moiety formats are disclosed in Section 6.8.

[0015] Fc domains that can be incorporated into the IFN receptor agonists of the disclosure are described in Section 6.9.

[0016] Exemplary IFN receptor agonists of the disclosure are described in Section 6.2 and numbered embodiments 1 to 227 and 284 to 344.

[0017] The disclosure further provides nucleic acids encoding the IFN receptor agonists of the disclosure. The nucleic acids encoding the IFN receptor agonists can be a single nucleic acid (*e.g.*, a vector encoding all polypeptide chains of an IFN receptor agonist) or a plurality of nucleic acids (*e.g.*, two or more vectors encoding the different polypeptide chains of an IFN receptor agonist). The disclosure further provides host cells and cell lines engineered to express the nucleic acids and IFN receptor agonists of the disclosure. The disclosure further provides methods of producing an IFN receptor agonist of the disclosure. Exemplary nucleic acids, host cells, and cell lines, and methods of producing an IFN receptor agonist are described in Section 6.10 and numbered embodiments 228 to 230 and 345 to 347.

[0018] The disclosure further provides pharmaceutical compositions comprising the IFN receptor agonists of the disclosure. Exemplary pharmaceutical compositions are described in Section 6.11 and numbered embodiment 231 and 348.

[0019] Further provided herein are methods of using the IFN receptor agonists and the pharmaceutical compositions of the disclosure, *e.g.*, for treating cancer. Exemplary

methods are described in Section 6.12 and numbered embodiments 232 to 283 and 349 to 356.

5. BRIEF DESCRIPTION OF THE FIGURES

[0020] FIGS. 1 is a cartoon representing a Type I IFN-IFN receptor complex.

[0021] FIGS. 2A-2X are cartoons representing IFN receptor agonists (FIGS. 2B-2X) and their constituent components (FIG. 2A). As used in FIGS. 2A-2X, "IFN" refers generally to any IFN moiety, "IFNAR1" refers generally to any IFNAR1 moiety, and "IFNAR2" refers generally to an IFNAR2 moiety. Although the IFN receptor agonists are shown without targeting moieties, targeting moieties can be incorporated into these IFN receptor agonists, *e.g.*, as shown in FIGS. 3A-3X.

[0022] FIGS. 3A-3X are cartoons representing IFN receptor agonists (FIGS. 3B-3X) and their constituent components (FIG. 3A). As used in FIGS. 3A-3X, "IFN" refers generally to any IFN moiety, "IFNAR1" refers generally to any IFNAR1 moiety, and "IFNAR2" refers generally to an IFNAR2 moiety. Although shown as having targeting moieties in the forms of Fabs, the Fabs can be replaced by other types of targeting moieties (*e.g.*, scFvs).

[0023] FIG. 4A-4B list exemplary IFN molecules that can be incorporated into the IFN receptor agonists of the disclosure.

[0024] FIGS. 5A-5D are the size exclusion ultra-performance liquid chromatography (SE-UPLC) profiles of exemplary IFN molecules that can be incorporated into the IFN receptor agonists of the disclosure.

[0025] FIGS. 6A-6C show the *in vitro* activity of exemplary IFN molecules that can be incorporated into the IFN receptor agonists of the disclosure. The cartoon images in **FIG. 6A** represent the N- and C-terminus Fc-fusions of IFN. **FIG. 6B** is a graph showing the *in vitro* activity of exemplary IFN molecules, Fc-IFNα2b, IFNα2b-Fc, and Fc-IFNα2b x Fc, in comparison to unlinked IFNα2b. **FIG. 6C** is a graph showing the activity of Fc-IFN molecules in comparison to different unlinked IFNs.

[0026] FIGS. 7A-7D are the SE-UPLC profiles of exemplary mutant IFN molecules that may be incorporated into the IFN receptor agonist constructs of the disclosure. **FIG. 7A** illustrates the SE-UPLC profile of a mutant IFN molecule, Fc-IFNα2bR33A. **FIG. 7B** illustrates the SE-

UPLC profile of a mutant IFN molecule, Fc-IFNα2bR149A. **FIG. 7C** illustrates the SE-UPLC profile of a mutant IFN molecule, Fc-IFNα2bR120A. **FIG. 7D** illustrates the SE-UPLC profile of a mutant IFN molecule, Fc-IFNα2bS152A.

[0027] FIGS. 8A-8B depict the *in vitro* activity of exemplary mutant IFN molecules that may be incorporated into the IFN receptor agonist constructs of the disclosure. The cartoon images in **FIG. 8A** represent the overall structure of wildtype (WT) or mutant (Mut) Fc-IFN molecules. **FIG. 8B** is a graph showing the in vitro activity of Fc-IFNα2b molecules with mutations affecting either the IFNAR1 or IFNAR2 interface.

[0028] FIGS. 9A-9F are the SE-UPLC profiles of some of the exemplary IFN receptor agonists shown in FIGS. 4A and 4B.

[0029] FIG. 10 is a graph showing the *in vitro* activity in reporter KG-1a cells of some of the exemplary IFN molecules shown in FIGS. 4A and 4B.

[0030] FIGS. 11A-11B are graphs showing the *in vitro* activity of exemplary IFN molecules and receptor agonists in reporter KG-1a cells. **FIG. 11A** shows the effect of receptor-masking on IFN activity using a homodimer format, whereas **FIG. 11B** illustrates the differences in reporter activity between different exemplary heterodimeric knob-in-hole (KiH) Fc-IFN molecules.

[0031] FIGS. 12A-12B are graphs showing the *in vitro* activity as measured by pSTAT flow cytometry analysis of exemplary IFN molecules and receptor agonists in murine hIFNAR $CD8^+$ T cells. **FIG. 12A** shows the effect of receptor-masking on IFN activity using a homodimer format, whereas **FIG. 12B** illustrates the differences in activity between different exemplary heterodimeric KiH Fc-IFN molecules as in FIG. 11B.

[0032] FIGS. 13A-13B are graphs showing the *in vitro* activity as measured by pSTAT flow cytometry analysis of exemplary IFN molecules and receptor agonists in murine hIFNAR $CD11b^+$ cells. **FIG. 13A** shows the effect of receptor-masking on IFN activity using a homodimer format and **FIG. 13B** shows the differences in activity between the same set of heterodimeric KiH Fc-IFN molecules in FIG. 11B.

[0033] FIGS. 14A-14B are graphs showing the *in vitro* activity as measured by pSTAT flow cytometry analysis of exemplary IFN molecules and receptor agonists in murine hIFNAR $CD4^+$ T cells. **FIG. 14A** shows the effect of receptor-masking on IFN activity using a

homodimer format and **FIG. 14B** shows the differences in activity between heterodimeric KiH Fc-IFN molecules shown in FIG. 11B.

[0034] FIGS. 15A-15B are graphs showing the *in vitro* activity as measured by pSTAT flow cytometry analysis of exemplary IFN molecules and receptor agonists in murine hIFNAR NK cells. **FIG. 15A** shows the effect of receptor-masking on IFN activity using a homodimer format and **FIG. 15B** shows the differences in activity between the heterodimeric KiH Fc-IFN molecules shown in FIG. 11B.

[0035] FIGS. 16A-16B are graphs showing the *in vitro* activity of exemplary IFN molecules and receptor agonists in two distinct types of PBMC cells as measured by pSTAT flow cytometry analysis. **FIG. 16A** shows the activity of IFN molecules in PBMC $CD8^+$ cells and **FIG. 16B** shows the activity of the same IFN molecules in FIG. 16A using PBMC NK cells.

[0036] FIG. 17 is a graph showing the *in vitro* activity of exemplary single-masked and dual-masked monovalent IFN molecules and receptor agonists in reporter KG-1a cells.

[0037] FIG. 18 is a graph showing the *in vitro* activity of exemplary single-masked and dual-masked bivalent IFN molecules and receptor agonists in reporter KG-1a cells.

[0038] FIGS. 19A-19D are graphs showing the effects of PDL1 targeting on *in vitro* activity of exemplary IFN molecules and receptor agonists in reporter KG-1a cells. **FIG. 19A** shows the activity of monovalent IFNAR2-masked IFN molecules and controls in PDL1 OE KG-1a cells. **FIG. 19B** shows the activity of monovalent IFNAR2-masked IFN molecules and controls in PDL1 KO KG-1a cells. **FIG. 19C** shows the activity of a bivalent IFNAR2-masked IFN molecule and controls in PDL1 OE KG-1a cells. **FIG. 19D** shows the activity of a bivalent IFNAR2-masked IFN molecule and controls in PDL1 KO KG-1a cells.

[0039] FIGS. 20A-20F show the effect of linker length on *in vitro* activity of exemplary dual-masked monovalent IFN molecules and receptor agonists in reporter KG-1a cells. **FIGS. 20A-2D** are cartoons representing dual-masked monovalent IFN receptor agonists with varying lengths of linkers between the IFNα2b and masking moieties. **FIG. 20E** shows the effect of linker length on the activity of monovalent dual-masked IFN molecules and controls in PDL1 OE KG-1a cells. **FIG. 20F** shows the effect of linker length on the activity of monovalent dual-masked IFN molecules and controls in PDL1 KO KG-1a cells.

6. DETAILED DESCRIPTION

6.1. DEFINITIONS

[0040] As used herein, the following terms are intended to have the following meanings:

[0041] **ABD chain, targeting moiety chain:** Targeting moieties and antigen binding sites (ABD's) within them can exist as one (*e.g.*, in the case of an scFv or scFab) polypeptide chain or form through the association of more than one polypeptide chains (*e.g.*, in the case of a Fab or an Fv). As used herein, the terms "ABD chain" and "targeting moiety chain" refer to all or a portion of an ABD or targeting moiety that exists on a single polypeptide chain. The use of the term "ABD chain" or "targeting moiety chain" is intended for convenience and descriptive purposes only and does not connote a particular configuration or method of production. Further, the reference to an ABD or targeting moiety when describing an IFN receptor agonist encompasses an ABD chain or targeting moiety chain unless the context dictates otherwise. Thus, when describing an IFN receptor agonist in which an Fc domain is operably linked to a targeting moiety, the Fc domain may be covalently linked directly or indirectly (*e.g.*, via a linker) through a peptide bond to, *e.g.*, (1) a first ABD or targeting moiety chain of a Fab or Fv (with the other components of the Fab or Fv on a second, associated ABD or targeting moiety chain) or (2) an ABD or targeting moiety chain containing an scFv or scFab.

[0042] **About, Approximately:** The terms "about", "approximately" and the like are used throughout the specification in front of a number to show that the number is not necessarily exact (*e.g.*, to account for fractions, variations in measurement accuracy and/or precision, timing, *etc.*). It should be understood that a disclosure of "about X" or "approximately X" where X is a number is also a disclosure of "X." Thus, for example, a disclosure of an embodiment in which one sequence has "about X% sequence identity" to another sequence is also a disclosure of an embodiment in which the sequence has "X% sequence identity" to the other sequence.

[0043] **Activate, activation:** The terms "activation", "activation", and the like in conjunction with an IFN receptor agonist of the disclosure refers to the protease-mediated enzymatic cleavage of a protease-cleavable linker that results in the release of an IFN moiety from a masking moiety, *e.g.*, a receptor-based masking moiety as described herein.

[0044] **And, or:** Unless indicated otherwise, an "or" conjunction is intended to be used in its correct sense as a Boolean logical operator, encompassing both the selection of features in the alternative (A or B, where the selection of A is mutually exclusive from B) and the selection of features in conjunction (A or B, where both A and B are selected). In some places in the text, the term "and/or" is used for the same purpose, which shall not be construed to imply that "or" is used with reference to mutually exclusive alternatives.

[0045] **Antibody:** The term "antibody" as used herein refers to a polypeptide (or set of polypeptides) of the immunoglobulin family that is capable of binding an antigen non-covalently, reversibly and specifically. For example, a naturally occurring "antibody" of the IgG type is a tetramer comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain (abbreviated herein as CL). The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (*e.g.*, effector cells) and the first component (Clq) of the classical complement system. The term "antibody" includes, but is not limited to, monoclonal antibodies, human antibodies, humanized antibodies, camelized antibodies, chimeric antibodies, bispecific or multispecific antibodies and anti-idiotypic (anti-id) antibodies. The antibodies can be of any isotype/class (*e.g.*, IgG, IgE, IgM, IgD, IgA and IgY) or subclass (*e.g.*, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2). Both the light and heavy chains are divided into regions of structural and functional homology. The terms "constant" and "variable" are used functionally. In this regard, it will be appreciated that the variable domains of both the light (VL) and heavy (VH) chain portions determine antigen recognition and specificity. Conversely, the constant domains of the light chain (CL) and the

heavy chain (CH1, CH2 or CH3) confer important biological properties such as secretion, transplacental mobility, Fc receptor binding, complement binding, and the like. By convention the numbering of the constant region domains increases as they become more distal from the antigen-binding domain or amino-terminus of the antibody. The N-terminus is a variable region and at the C-terminus is a constant region; the CH3 and CL domains represent the carboxy-terminus of the heavy and light chain, respectively, of natural antibodies. For convenience, and unless the context dictates otherwise, the reference to an antibody also refers to antibody fragments as well as engineered antibodies that include non-naturally occurring antigen-binding domains and/or antigen-binding domains having non-native configurations.

[0046] **Antigen-binding domain:** The term "antigen-binding domain" or "ABD" as used herein refers to a portion of an antibody or antibody fragment (*e.g.*, a targeting moiety) that has the ability to bind to an antigen non-covalently, reversibly and specifically. Examples of an antibody fragment that can comprise an ABD include, but are not limited to, a single-chain Fv (scFv), a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; a F(ab)2 fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; a Fd fragment consisting of the VH and CH1 domains; a Fv fragment consisting of the VL and VH domains of a single arm of an antibody; a dAb fragment (Ward *et al.*, 1989, Nature 341:544-546), which consists of a VH domain; and an isolated complementarity determining region (CDR). Thus, the term "antibody fragment" encompasses both proteolytic fragments of antibodies (*e.g.*, Fab and $F(ab)_2$ fragments) and engineered proteins comprising one or more portions of an antibody (*e.g.*, an scFv). Antibody fragments can also be incorporated into single domain antibodies, maxibodies, minibodies, intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv (see, *e.g.*, Hollinger and Hudson, 2005, Nature Biotechnology 23: 1126-1136).

[0047] **Associated:** The term "associated" in the context of an IFN receptor agonist refers to a functional relationship between two or more polypeptide chains. In particular, the term "associated" means that two or more polypeptides are associated with one another, *e.g.*, non-covalently through molecular interactions or covalently through one or more disulfide bridges or chemical cross-linkages, so as to produce a functional IFN receptor agonist. Examples of associations that might be present in an IFN receptor agonist of the disclosure

include (but are not limited to) associations between Fc domains to form an Fc region (homodimeric or heterodimeric as described in Section 6.9), associations between VH and VL regions in a Fab or Fv, and associations between CH1 and CL in a Fab.

[0048] **Cancer:** The term "cancer" refers to a disease characterized by the uncontrolled (and often rapid) growth of aberrant cells. Cancer cells can spread locally or through the bloodstream and lymphatic system to other parts of the body. Examples of various cancers are described herein and include but are not limited to, breast cancer, prostate cancer, ovarian cancer, cervical cancer, skin cancer, pancreatic cancer, colorectal cancer, renal cancer, liver cancer, brain cancer, adrenal gland cancer, autonomic ganglial cancer, biliary tract cancer, bone cancer, endometrial cancer, eye cancer, fallopian tube cancer, genital tract cancers, large intestinal cancer, cancer of the meninges, esophageal cancer, peritoneal cancer, pituitary cancer, penile cancer, placental cancer, pleura cancer, salivary gland cancer, small intestinal cancer, stomach cancer, testicular cancer, thymus cancer, thyroid cancer, upper aerodigestive cancers, urinary tract cancer, vaginal cancer, vulva cancer, lymphoma, leukemia, lung cancer and the like, *e.g.*, any TAA-positive cancers of any of the foregoing types.

[0049] **Complementarity Determining Region:** The terms "complementarity determining region" or "CDR," as used herein, refer to the sequences of amino acids within antibody variable regions which confer antigen specificity and binding affinity. For example, in general, there are three CDRs in each heavy chain variable region (*e.g.*, CDR-H1, CDR-H2, and CDR- H3) and three CDRs in each light chain variable region (CDR-L1, CDR-L2, and CDR-L3). The precise amino acid sequence boundaries of a given CDR can be determined using any of a number of well-known schemes, including those described by Kabat *et al.*, 1991, "Sequences of Proteins of Immunological Interest," 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD ("Kabat" numbering scheme), Al-Lazikani *et al.*, 1997, JMB 273:927-948 ("Chothia" numbering scheme) and ImMunoGenTics (IMGT) numbering (Lefranc, 1999, The Immunologist 7:132-136; Lefranc *et al.*, 2003, Dev. Comp. Immunol. 27:55-77 ("IMGT" numbering scheme). For example, for classic formats, under Kabat, the CDR amino acid residues in the heavy chain variable domain (VH) are numbered 31-35 (CDR-H1), 50-65 (CDR- H2), and 95-102 (CDR-H3); and the CDR amino acid residues in the light chain variable domain (VL) are numbered 24-34 (CDR-L1), 50-56 (CDR-

L2), and 89-97 (CDR-L3). Under Chothia, the CDR amino acids in the VH are numbered 26-32 (CDR-H1), 52-56 (CDR-H2), and 95-102 (CDR-H3); and the amino acid residues in VL are numbered 26-32 (CDR-L1), 50-52 (CDR-L2), and 91-96 (CDR-L3). By combining the CDR definitions of both Kabat and Chothia, the CDRs consist of amino acid residues 26-35 (CDR-H1), 50-65 (CDR-H2), and 95-102 (CDR-H3) in human VH and amino acid residues 24-34 (CDR-L1), 50-56 (CDR-L2), and 89-97 (CDR-L3) in human VL. Under IMGT the CDR amino acid residues in the VH are numbered approximately 26-35 (CDR-H1), 51-57 (CDR-H2) and 93-102 (CDR-H3), and the CDR amino acid residues in the VL are numbered approximately 27-32 (CDR-L1), 50-52 (CDR-L2), and 89-97 (CDR-L3) (numbering according to "Kabat"). Under IMGT, the CDR regions of an antibody can be determined using the program IMGT/DomainGap Align.

[0050] **Constant domain:** The terms "constant domain" refers to a CH1, CH2, CH3 or CL domain of an immunoglobulin.

[0051] The term "CH1 domain" refers to the heavy chain constant region linking the variable domain to the hinge in a heavy chain constant domain. In some embodiments, the term "CH1 domain" refers to the region of an immunoglobulin molecule spanning amino acids 118 to 215 (EU numbering). The term "CH1 domain" encompasses wildtype CH1 domains as well as variants thereof (*e.g.*, non-naturally-occurring CH1 domains or modified CH1 domains). For example, the term "CH1 domain" includes wildtype IgG1, IgG2, IgG3 and IgG4 CH1 domains and variants thereof having 1, 2, 3, 4, 5, 1-3, 1-5, 3-5 and/or at most 5, 4, 3, 2, or 1 mutations, *e.g.*, substitutions, deletions and/or additions. Exemplary CH1 domains include CH1 domains with mutations that modify a biological activity of an antibody, such as ADCC, CDC or half-life.

[0052] The term "CH2 domain" refers to the heavy chain constant region linking the hinge to the CH3 domain in a heavy chain constant domain. In some embodiments, the term "CH2 domain" refers to the region of an immunoglobulin molecule spanning amino acids 238 to 340 (EU numbering). The term "CH2 domain" encompasses wildtype CH2 domains as well as variants thereof (*e.g.*, non-naturally-occurring CH2 domains or modified CH2 domains). For example, the term "CH2 domain" includes wildtype IgG1, IgG2, IgG3 and IgG4 CH2 domains and variants thereof having 1, 2, 3, 4, 5, 1-3, 1-5, 3-5 and/or at most 5, 4, 3, 2, or 1 mutations, *e.g.*, substitutions, deletions and/or additions. Exemplary CH2 domains include

CH2 domains with mutations that modify a biological activity of an antibody, such as ADCC, CDC, purification, dimerization and half-life.

[0053] The term "CH3 domain" refers to the heavy chain constant region that is C-terminal to the CH2 domain in a heavy chain constant domain. In some embodiments, the term "CH3 domain" refers to the region of an immunoglobulin molecule spanning amino acids 341 to 447 (EU numbering). The term "CH3 domain" encompasses wildtype CH3 domains as well as variants thereof (*e.g.*, non-naturally-occurring CH3 domains or modified CH3 domains). For example, the term "CH3 domain" includes wildtype IgG1, IgG2, IgG3 and IgG4 CH3 domains and variants thereof having 1, 2, 3, 4, 5, 1-3, 1-5, 3-5 and/or at most 5, 4, 3, 2, or 1 mutations, *e.g.*, substitutions, deletions and/or additions. Exemplary CH3 domains include CH3 domains with mutations that modify a biological activity of an antibody, such as ADCC, CDC, purification, dimerization and half-life.

[0054] The term "CL domain" refers to the constant region of an immunoglobulin light chain. The term "CL domain" encompasses wildtype CL domains (*e.g.*, kappa or lambda light chain constant regions) as well as variants thereof (*e.g.*, non-naturally-occurring CL domains or modified CL domains). For example, the term "CL domain" includes wildtype kappa and lambda constant domains and variants thereof having 1, 2, 3, 4, 5, 1-3, 1-5, 3-5 and/or at most 5, 4, 3, 2, or 1 mutations, *e.g.*, substitutions, deletions and/or additions.

[0055] **Effector Function:** The term "effector function" refers to an activity of an antibody molecule that is mediated by binding through a domain of the antibody other than the antigen-binding domain, usually mediated by binding of effector molecules. Effector function includes complement-mediated effector function, which is mediated by, for example, binding of the C1 component of the complement to the antibody. Activation of complement is important in the opsonization and lysis of cell pathogens. The activation of complement also stimulates the inflammatory response and may also be involved in autoimmune hypersensitivity. Effector function also includes Fc receptor (FcR)-mediated effector function, which may be triggered upon binding of the constant domain of an antibody to an Fc receptor (FcR). Binding of antibody to Fc receptors on cell surfaces triggers a number of important and diverse biological responses including engulfment and destruction of antibody-coated particles, clearance of immune complexes, lysis of antibody-coated target cells by killer cells (called antibody- dependent cell-mediated cytotoxicity, or ADCC), release

of inflammatory mediators, placental transfer and control of immunoglobulin production. An effector function of an antibody may be altered by altering, *e.g.*, enhancing or reducing, the affinity of the antibody for an effector molecule such as an Fc receptor or a complement component. Binding affinity will generally be varied by modifying the effector molecule binding site, and in this case, it is appropriate to locate the site of interest and modify at least part of the site in a suitable way. It is also envisaged that an alteration in the binding site on the antibody for the effector molecule need not alter significantly the overall binding affinity but may alter the geometry of the interaction rendering the effector mechanism ineffective as in non-productive binding. It is further envisaged that an effector function may also be altered by modifying a site not directly involved in effector molecule binding, but otherwise involved in performance of the effector function.

[0056] **Epitope:** An epitope, or antigenic determinant, is a portion of an antigen recognized by an antibody or other antigen-binding moiety as described herein. An epitope can be linear or conformational.

[0057] **Fab:** The term "Fab" refers to a pair of polypeptide chains, the first comprising a variable heavy (VH) domain of an antibody operably linked (typically N-terminal to) to a first constant domain (referred to herein as C1), and the second comprising variable light (VL) domain of an antibody N-terminal operably linked (typically N-terminal) to a second constant domain (referred to herein as C2) capable of pairing with the first constant domain. In a native antibody, the VH is N-terminal to the first constant domain (CH1) of the heavy chain and the VL is N-terminal to the constant domain of the light chain (CL). The Fabs of the disclosure can be arranged according to the native orientation or include domain substitutions or swaps that facilitate correct VH and VL pairings. For example, it is possible to replace the CH1 and CL domain pair in a Fab with a CH3-domain pair to facilitate correct modified Fab-chain pairing in heterodimeric molecules. It is also possible to reverse CH1 and CL, so that the CH1 is attached to VL and CL is attached to the VH, a configuration generally known as Crossmab. The term "Fab" encompasses single chain Fabs.

[0058] **Fc Domain and Fc Region:** The term "Fc domain" refers to a portion of the heavy chain that pairs with the corresponding portion of another heavy chain. In some embodiments an Fc domain comprises a CH2 domain followed by a CH3 domain, with or without a hinge region N-terminal to the CH2 domain. The term "Fc region" refers to the

region of formed by association of two heavy chain Fc domains. The two Fc domains within the Fc region may be the same or different from one another. In a native antibody the Fc domains are typically identical, but one or both Fc domains might be modified to allow for heterodimerization, *e.g.,* via a knob-in-hole interaction.

[0059] **Fv:** The term "Fv" refers to the minimum antibody fragment derivable from an immunoglobulin that contains a complete target recognition and binding site. This region consists of a dimer of one heavy and one light chain variable domain in a tight, noncovalent association (VH-VL dimer). It is in this configuration that the three CDRs of each variable domain interact to define a target binding site on the surface of the VH-VL dimer. Often, the six CDRs confer target binding specificity to the antibody. However, in some instances even a single variable domain (or half of an Fv comprising only three CDRs specific for a target) can have the ability to recognize and bind target. The reference to a VH-VL dimer herein is not intended to convey any particular configuration. When present on a single polypeptide chain (*e.g.*, a scFv), the VH and be N- terminal or C-terminal to the VL.

[0060] **Half Antibody:** The term "half antibody" refers to a molecule that comprises at least one Fc domain and can associate with another molecule comprising an Fc domain through, *e.g.*, a disulfide bridge or molecular interactions. A half antibody can be composed of one polypeptide chain or more than one polypeptide chains (*e.g.*, the two polypeptide chains of a Fab). An example of a half antibody is a molecule comprising a heavy and light chain of an antibody (*e.g.*, an IgG antibody). Another example of a half antibody is a molecule comprising a first polypeptide comprising a VL domain and a CL domain, and a second polypeptide comprising a VH domain, a CH1 domain, a hinge domain, a CH2 domain, and a CH3 domain, wherein said VL and VH domains form an ABD. Yet another example of a half antibody is a polypeptide comprising an scFv domain, a CH2 domain and a CH3 domain.

[0061] The IFN receptor agonists of the disclosure typically comprise two half antibodies, each comprising an IFN moiety masked by one or two receptor moieties, *e.g.*, IFNR masking moieties. The one or two masking moieties can be in the same half antibody or the other half antibody as the IFN moiety, as exemplified in the embodiments illustrated in FIGS. 2B-2X and set forth Table 2. In some embodiments, the IFN moiety and/or the masking moiety has an adjacent protease-cleavable linker, the cleavage of which releases the masking moiety from the IFN moiety, thereby activating the IFN moiety. Exemplary positions of

protease cleavable linkers in the half antibodies are shown in Table 1. One or both half antibodies in the IFN receptor agonists may further comprise a targeting moiety, such as an scFv or Fab. Exemplary IFN receptor agonists comprising targeting moieties are illustrated in FIGS. 3B-3X and set forth in Table 3.

[0062] The term "half antibody" is intended for descriptive purposes only and does not connote a particular configuration or method of production. Descriptions of a half antibody as a "first" half antibody, a "second" half antibody, a "left" half antibody, a "right" half antibody or the like are merely for convenience and descriptive purposes.

[0063] **Host cell or recombinant host cell:** The terms "host cell" or "recombinant host cell" refer to a cell that has been genetically-engineered, *e.g.*, through introduction of a heterologous nucleic acid. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein. A host cell may carry the heterologous nucleic acid transiently, *e.g.*, on an extrachromosomal heterologous expression vector, or stably, *e.g.*, through integration of the heterologous nucleic acid into the host cell genome. For purposes of expressing a IFN receptor agonist of the disclosure, a host cell is preferably a cell line of mammalian origin or mammalian-like characteristics, such as monkey kidney cells (COS, *e.g.*, COS-1, COS-7), HEK293), baby hamster kidney (BHK, *e.g.*, BHK21), Chinese hamster ovary (CHO), NSO, PerC6, BSC-1 , human hepatocellular carcinoma cells (*e.g.*, Hep G2), SP2/0, HeLa, Madin-Darby bovine kidney (MDBK), myeloma and lymphoma cells, or derivatives and/or engineered variants thereof. The engineered variants include, *e.g.*, derivatives that grow at higher density than the original cell lines and/or glycan profile modified derivatives and and/or site- specific integration site derivatives.

[0064] **Interferon:** The term "interferon" as used herein refers to a full-length interferon or to a modified interferon, for example a truncated and/or mutant interferon. In some embodiments, the modified interferon is attenuated as compared to the corresponding wildtype interferon(*e.g.*, retains less than 50%, less than 40%, less than 30%, less than 20%, or less than 10%, less than 1%, less than 0.1% or less than 0.05% activity in an *in*

vitro luciferase reporter assay as described in Section 8.2.3). In some embodiments, the modified interferon is attenuated by a range bounded by any two of the foregoing values, *e.g.*, 0.05%-50%, 0.1%-20%, 0.1%-10%, 0.05%-5%, 1%-20%, and so on and so forth. In other embodiments the modified interferon substantially retains the biological activity of the corresponding wildtype interferon (*e.g.*, retains at least 50% activity in an *in vitro* luciferase reporter assay as described in Section 8.2.3). Interferons include Type I interferons (*e.g.*, interferon-α and interferon-β) as well as Type II interferons (*e.g.*, interferon-γ).

[0065] **Linker:** The term "linker" as used herein refers to a protease-cleavable linker or a non-cleavable linker.

[0066] **Non-cleavable linker:** A non-cleavable linker as used herein refers to a peptide whose amino acid sequence lacks a substrate sequence for a protease, *e.g.*, a protease as described in Section 6.5.1, that recognizes and cleaves a specific sequence motif, *e.g.*, a substrate as described in Section 6.5.2.

[0067] **Operably linked:** The term "operably linked" refers to a functional relationship between two or more peptide or polypeptide domains or nucleic acid (*e.g.*, DNA) segments. In the context of a fusion protein or other polypeptide, the term "operably linked" means that two or more amino acid segments are linked so as to produce a functional polypeptide. For example, in the context of an IFN receptor agonist of the disclosure, separate components (*e.g.*, an Fc domain and an IFN moiety) can be operably linked directly or through peptide linker sequences. In the context of a nucleic acid encoding a fusion protein, such as a half antibody of an IFN receptor agonist of the disclosure, "operably linked" means that the two nucleic acids are joined such that the amino acid sequences encoded by the two nucleic acids remain in-frame. In the context of transcriptional regulation, the term refers to the functional relationship of a transcriptional regulatory sequence to a transcribed sequence. For example, a promoter or enhancer sequence is operably linked to a coding sequence if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system.

[0068] **Polypeptide, Peptide and Protein:** The terms "polypeptide", "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues.

[0069] **Proprotein:** A "proprotein" is a protein precursor that is inactive, and which can be activated by proteolysis by a protease. Thus, proproteins are "protease activatable".

[0070] **Protease:** The term "protease" as used herein refers to any enzyme that catalyzes hydrolysis of a peptide bond. Generally, the proteases useful in the present disclosure, *e.g.*, the proteases described in Section 6.5.1, recognize and cleaves a specific sequence motif, *e.g.*, a substrate as described in Section 6.5.2. Preferably, the proteases are expressed at higher levels in cancer tissues as compared to normal tissues.

[0071] **Protease-cleavable linker:** As used herein, the term "protease-cleavable linker" or "PCL" refers to a peptide whose amino acid sequence contains one or more (*e.g.*, two, three or more) substrate sequences for one or more proteases. Exemplary protease-cleavable linkers are described in Section 6.5 and exemplary protease-cleavable linker sequences are disclosed in Section 6.5.4.

[0072] **Recognize:** The term "recognize" as used herein refers to an antibody or antibody fragment (*e.g.*, a targeting moiety) that finds and interacts (*e.g.*, binds) with its epitope.

[0073] **Single Chain Fab or scFab:** The term "single chain Fab" or "scFab" as used herein refers an ABD comprising a VH domain, a CH1 domain, a VL domain, a CL domain and a linker. In some embodiments, the foregoing domains and linker are arranged in one of the following orders in a N-terminal to C-terminal orientation: (a) VH-CH1-linker-VL-CL, (b) VL-CL-linker-VH-CH1, (c) VH-CL-linker-VL-CH1 or (d) VL-CH1-linker-VH-CL. Linkers are suitably non-cleavable linkers of at least 30 amino acids, preferably between 32 and 50 amino acids. Single chain Fab fragments are typically stabilized via the natural disulfide bond between the CL domain and the CH1 domain. In addition, these single chain Fab molecules might be further stabilized by generation of interchain disulfide bonds via insertion of cysteine residues (*e.g.*, at position 44 in the VH domain and position 100 in the VL domain according to Kabat numbering).

[0074] **Single Chain Fv or scFv:** The term "single-chain Fv" or "scFv" as used herein refers to ABDs comprising the VH and VL domains of an antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains which enables the scFv to form the desired structure for antigen-binding. For a review of scFv see Pluckthun in The Pharmacology of Monoclonal Antibodies, vol. 113, Rosenburg and Moore eds. (1994), Springer-Verlag, New York, pp. 269-315. The VH and VL and be arranged in the N- to C-

terminal order VH-VL or VL-VH, typically separated by a linker, for example a linker as set forth in Table E.

[0075] Spacer: As used herein, the term "spacer" refers to a peptide, the amino acid sequence of which is not a substrate for a protease, incorporated into a linker containing a substrate. A spacer can be used to separate the substrate from other domains in a molecule, for example an ABD. In some aspects, residues in the spacer minimize aminopeptidase and/or exopeptidase action to prevent cleavage of N-terminal amino acids.

[0076] Specifically (or selectively) binds: The term "specifically (or selectively) binds" to an antigen or an epitope refers to a binding reaction that is determinative of the presence of a cognate antigen or an epitope in a heterogeneous population of proteins and other molecules. The binding reaction can be but need not be mediated by an antibody or antibody fragment. The term "specifically binds" does not exclude cross-species reactivity. For example, an antigen-binding domain (*e.g.*, an antigen-binding fragment of an antibody) that "specifically binds" to an antigen from one species may also "specifically bind" to that antigen in one or more other species. Thus, such cross-species reactivity does not itself alter the classification of an antigen-binding domain as a "specific" binder. In certain embodiments, an antigen-binding domain of the disclosure that specifically binds to a human antigen has cross-species reactivity with one or more non-human mammalian species, *e.g.*, a primate species (including but not limited to one or more of *Macaca fascicularis*, *Macaca mulatta*, and *Macaca nemestrina*) or a rodent species, *e.g.*, *Mus musculus*.

[0077] Subject: The term "subject" includes human and non-human animals. Non-human animals include all vertebrates, *e.g.*, mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, and reptiles. In preferred embodiments, the subject is human.

[0078] Substrate: The term "substrate" refers to peptide sequence on which a protease will act and within which the protease will cleave a peptide bond.

[0079] Target Molecule: The term "target molecule" as used herein refers to any biological molecule (*e.g.,* protein, carbohydrate, lipid or combination thereof) expressed on a cell surface or in the extracellular matrix that can be specifically bound by a targeting moiety in an IFN receptor agonist of the disclosure.

[0080] **Targeting Moiety:** The term "targeting moiety" as used herein refers to any molecule or binding portion (*e.g.*, an immunoglobulin or an antigen binding fragment) thereof that can bind to a cell surface or extracellular matrix molecule at a site to which an IFN receptor agonist of the disclosure is to be localized, for example on tumor cells or on lymphocytes in the tumor microenvironment. In some embodiments, the targeting moiety binds to a TAA. In other embodiments, the targeting moiety binds to a TCA. The targeting moiety can also have a functional activity in addition to localizing an IFN receptor agonist to a particular site. For example, a targeting moiety that binds to a checkpoint inhibitor such as PD1 can also exhibit anti-tumor activity or enhance the anti-tumor activity by IFN, for example by inhibiting PD1 signaling.

[0081] **T-Cell Antigen, TCA:** The term "T-cell antigen" or "TCA" refers to a molecule (typically a protein, carbohydrate, lipid or some combination thereof) that is expressed on the surface of a T-lymphocyte and is useful for the preferential targeting of a pharmacological agent to a particular site. In some embodiments, the site is cancer tissue and/or the T-cell antigen is a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, or a checkpoint inhibitor expressed on a T-lymphocyte.

[0082] **Tumor:** The term "tumor" is used interchangeably with the term "cancer" herein, *e.g.*, both terms encompass solid and liquid, *e.g.*, diffuse or circulating, tumors. As used herein, the term "cancer" or "tumor" includes premalignant, as well as malignant cancers and tumors.

[0083] **Tumor-Associated Antigen, TAA:** The term "tumor-associated antigen" or "TAA" refers to a molecule (typically a protein, carbohydrate, lipid or some combination thereof) that is expressed on the surface of a cancer cell, either entirely or as a fragment (*e.g.*, MHC/peptide), and which is useful for the preferential targeting of a pharmacological agent to the cancer cell. In some embodiments, a TAA is a marker expressed by both normal cells and cancer cells, *e.g.*, a lineage marker. In some embodiments, a TAA is a cell surface molecule that is overexpressed in a cancer cell in comparison to a normal cell, for instance, 1-fold over expression, 2-fold overexpression, 3-fold overexpression or more in comparison to a normal cell. In some embodiments, a TAA is a cell surface molecule that is inappropriately synthesized in the cancer cell, for instance, a molecule that contains deletions, additions or mutations in comparison to the molecule expressed on a normal cell.

In some embodiments, a TAA will be expressed exclusively on the cell surface of a cancer cell, entirely or as a fragment (*e.g.*, MHC/peptide), and not synthesized or expressed on the surface of a normal cell. Accordingly, the term "TAA" encompasses antigens that are specific to cancer cells, sometimes known in the art as tumor-specific antigens (TSAs).

[0084] **Treat, Treatment, Treating:** As used herein, the terms "treat", "treatment" and "treating" refer to the reduction or amelioration of the progression, severity and/or duration of a disorder, or the amelioration of one or more symptoms (preferably, one or more discernible symptoms) of a disorder resulting from the administration of one or more IFN receptor agonists of the disclosure. In some embodiments, the disorder is a proliferative disorder and the terms "treat", "treatment" and "treating" refer to the amelioration of at least one measurable physical parameter of a proliferative disorder, such as growth of a tumor, not necessarily discernible by the patient. In other embodiments the terms "treat", "treatment" and "treating" refer to the inhibition of the progression of a proliferative disorder, either physically by, *e.g.*, stabilization of a discernible symptom, physiologically by, *e.g.*, stabilization of a physical parameter, or both. In other embodiments the terms "treat", "treatment" and "treating" refer to the reduction or stabilization of tumor size or cancerous cell count.

[0085] **Universal Light Chain, UCL**: The term "universal light chain" or "ULC" as used herein refers to a light chain variable region (VL) that can pair with more than on heavy chain variable region (VL). In the context of a targeting moiety, the term "universal light chain" or "ULC" refers to a light chain polypeptide capable of pairing with the heavy chain region of the targeting moiety and also capable of pairing with other heavy chain regions. ULCs can also include constant domains, *e.g.*, a CL domain of an antibody. Universal light chains are also known as "common light chains.

[0086] **VH:** The term "VH" refers to the variable region of an immunoglobulin heavy chain of an antibody, including the heavy chain of an Fv, scFv, dsFv or Fab.

[0087] **VL:** The term "VL" refers to the variable region of an immunoglobulin light chain, including the light chain of an Fv, scFv, dsFv or Fab.

6.2. IFN receptor agonists

[0088] The present disclosure relates to IFN receptor agonists comprising an IFN moiety that is attenuated as compared to wild-type interferon. The IFN moiety may be attenuated by (i) masking by a Type I interferon receptor (IFNR) moiety (*e.g.*, as described in Section 6.4); (ii) one or more mutations in the IFN moiety as compared to wild-type interferon, *e.g.*, one or more amino acid substitutions and/or truncations (*e.g.*, as described in Section 6.3); (iii) use of native IFN sequences with a low receptor affinity; or (iv) any combination of two or all three of (i), (ii) and (iii).

[0089] Generally, the IFN receptor agonists are composed of two half antibodies, comprising a pair of Fc domains that associate to form an Fc region (typically comprising hinge sequences).

[0090] In the IFN receptor agonists of the disclosure, the two half antibodies together comprise at least one interferon (IFN moiety) but may include two or more IFN moieties.

[0091] The IFN moieties in the IFN receptor agonists may each be masked by one or two interferon receptor (IFNR) moieties, *e.g.*, an interferon alpha receptor 1 (IFNAR1) and/or interferon alpha receptor 2 (IFNAR2) moiety.

[0092] In some embodiments, the IFN receptor agonists further comprise targeting moieties, *e.g.*, antigen binding domains of antibodies, that target the IFN receptor agonists to a selected tissue, *e.g.*, cancer tissue.

[0093] Exemplary IFN receptor agonists are illustrated in FIGS. 2B-2X and 3B-3X.

[0094] Table 1 below describes exemplary half antibodies that can be incorporated into the IFN receptor agonists of the disclosure. As evident from Table 1, each half antibody may include one or more polypeptide chains. For convenience when describing combinations of half antibodies in the IFN receptor agonists of the disclosure, each half antibody described in Table 1 is referred to herein as an "Exemplary Monomer".

TABLE 1

Exemplary Monomer	Nomenclature	Polypeptide Chain(s)
1	Fc	Chain 1: Fc domain
1T	TM-Fc	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain

TABLE 1		
Exemplary Monomer	**Nomenclature**	**Polypeptide Chain(s)**
		Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
2	Fc-IFN	Chain 1: Fc domain – optional linker 1*– IFN moiety
2T	TM-Fc-IFN	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2* – IFN moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
3	IFN-Fc	Chain 1: IFN moiety - optional linker 1* – Fc domain
4	Fc-IFNR	Chain 1: Fc domain – optional linker 1* – IFNR moiety
4A	Fc-IFNAR1 or Fc-R1	Chain 1: Fc domain – optional linker 1* – IFNAR1 moiety
4B	Fc-IFNAR2 or Fc-R2	Chain 1: Fc domain – optional linker 1* – IFNAR2 moiety
4T	TM-Fc-IFNR	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2* – IFNR moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
4AT	TM-Fc-IFNAR1 or TM-Fc-R1	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2* – IFNAR1 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
4BT	TM-Fc-IFNAR2 or TM-Fc-R2	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 – Fc domain – optional linker 2* – IFNAR2 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
5	Fc-IFNR-IFN	Chain 1: Fc domain – optional linker 1 – IFNR moiety – optional linker 2* – IFN moiety
5A	Fc-IFNAR1-IFN or Fc-R1-IFN	Chain 1: Fc domain – optional linker 1 – IFNAR1 moiety – optional linker 2* – IFN moiety
5B	Fc-IFNAR2-IFN or Fc-R2-IFN	Chain 1: Fc domain – optional linker 1 – IFNAR2 moiety – optional linker 2* – IFN moiety

TABLE 1		
Exemplary Monomer	**Nomenclature**	**Polypeptide Chain(s)**
5T	TM-Fc-IFNR-IFN	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2 – IFNR moiety – optional linker 3* – IFN moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
5AT	TM-Fc-IFNAR1-IFN or TM-Fc-R1-IFN	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2 – IFNAR1 moiety – optional linker 3* – IFN moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL)
5BT	TM-Fc-IFNAR2-IFN or TM-Fc-R2-IFN	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2 – IFNAR2 moiety – optional linker 3* – IFN moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL)
6	Fc-IFN-IFNR	Chain 1: Fc domain – optional linker 1** – IFN moiety – optional linker 2** – IFNR moiety
6A	Fc-IFN-IFNAR1 or Fc-IFN-R1	Chain 1: Fc domain – optional linker 1** – IFN moiety – optional linker 2** – IFNAR1 moiety
6B	Fc-IFN-IFNAR2 or Fc-IFN-R2	Chain 1: Fc domain – optional linker 1** – IFN moiety – optional linker 2** – IFNAR2 moiety
6T	TM- Fc-IFN-IFNR	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2** – IFN moiety – optional linker 3** – IFNR moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
6AT	TM-Fc-IFN-IFNAR1 or TM-Fc-IFN-R1	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2** – IFN moiety – optional linker 3** – IFNAR1 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
6BT	TM-Fc-IFN-IFNAR2 or TM-Fc-IFN-R2	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2** – IFN moiety – optional linker 3** – IFNAR2 moiety

TABLE 1		
Exemplary Monomer	**Nomenclature**	**Polypeptide Chain(s)**
		Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
7	Fc-IFNR	Chain 1: Fc domain – optional linker 1* – IFNR moiety
7A	Fc-IFNAR1 or Fc-R1	Chain 1: Fc domain – optional linker 1* – IFNAR1 moiety
7B	Fc-IFNAR2 or Fc-R2	Chain 1: Fc domain – optional linker 1* – IFNAR2 moiety
7T	TM-Fc-IFNR	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2* – IFNR moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
7AT	TM-Fc-IFNAR1 or TM-Fc-R1	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2* – IFNAR1 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
7BT	TM-Fc-IFNAR2 or TM-Fc-R2	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2* – IFNAR2 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
8	Fc-IFNR-IFN-IFNR	Chain 1: Fc domain – optional linker 1 – IFNR moiety – optional linker 2** – IFN moiety – optional linker 3** – IFNR moiety
8A	Fc-IFNAR1-IFN-IFNAR2 or Fc-R1-IFN-R2	Chain 1: Fc domain – optional linker 1 – IFNAR1 moiety – optional linker 2** – IFN moiety – optional linker 3** – IFNAR2 moiety
8B	Fc-IFNAR2-IFN-IFNAR1 or Fc-R2-IFN-R1	Chain 1: Fc domain – optional linker 1 – IFNAR2 moiety – optional linker 2** – IFN moiety – optional linker 3** – IFNAR1 moiety
8T	TM-Fc-IFNR-IFN-IFNR	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2 – IFNR moiety – optional linker 3** – IFN moiety – optional linker 4** – IFNR moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL)
8AT	TM-Fc-IFNAR1-IFN-IFNAR2 or TM-Fc-R1-IFN-R2	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2 – IFNAR1 moiety

TABLE 1		
Exemplary Monomer	**Nomenclature**	**Polypeptide Chain(s)**
		– optional linker 3** – IFN moiety – optional linker 4** – IFNAR2 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL)
8BT	TM-Fc-IFNAR2-IFN-IFNAR1 or TM-Fc-R2-IFN-R1	Chain 1: Targeting moiety (or targeting moiety component) – optional linker 1 - Fc domain – optional linker 2 – IFNAR2 moiety – optional linker 3** – IFN moiety – optional linker 4** – IFNAR1 moiety Chain 2 (optional): Targeting moiety component (*e.g.*, VL or VL-CL)
9	IFNR-IFN-Fc	Chain 1: IFNR moiety – optional linker 1** – IFN moiety – optional linker 2** - Fc domain
9A	IFNAR1-IFN-Fc or IFNR1-IFN-Fc	Chain 1: IFNAR1 moiety – optional linker 1** – IFN moiety – optional linker 2** - Fc domain
9B	IFNAR2-IFN-Fc or IFNR2-IFN-Fc	Chain 1: IFNAR2 moiety – optional linker 1** – IFN moiety – optional linker 2** - Fc domain
10	IFNR-IFN-IFNR-Fc	Chain 1: IFNR moiety – optional linker 1** – IFN moiety – optional linker 2** – IFNR moiety – optional linker 3 – Fc domain
10A	IFNAR1-IFN-IFNAR2-Fc or R1-IFN-R2-Fc	Chain 1: IFNAR1 moiety – optional linker 1** – IFN moiety – optional linker 2** – IFNAR2 moiety – optional linker 3 – Fc domain
10B	IFNAR2-IFN-IFNAR1-Fc or R2-IFN-R1-Fc	Chain 1: IFNAR2 moiety – optional linker 1** – IFN moiety – optional linker 2** – IFNAR1 moiety – optional linker 3 – Fc domain

[0095] The IFN receptor agonists may further include one or two protease-cleavable linkers (PCLs) in each half antibody, with other linkers being non-cleavable. In some embodiments, all linkers are non-cleavable. Exemplary protease-cleavable linkers are described in Section 6.5 and non-cleavable linkers described in Section 6.6. In the Exemplary Monomers of Table 1, linkers identified by an asterisk are optionally protease-cleavable linkers, and linkers identified by two asterisks indicate two linkers in a particular half antibody that can be protease-cleavable linkers. Where two linkers in an Exemplary Monomer are identified as being optionally protease-cleavable, in some embodiments the N-terminal linker is protease-cleavable, in other embodiments the C-terminal linker is protease-cleavable, and in yet other

embodiments both linkers are protease-cleavable. The Fc domains in the polypeptide chains described in Table 1 preferably comprise a hinge domain as set forth in Section 6.9.3.

[0096] Table 2 below shows Exemplary Monomers pairings that can be utilized in the IFN receptor agonists of the disclosure and their constituents. Additional components not specifically recited, *e.g.*, targeting moieties, may be incorporated into the IFN receptor agonists.

Table 2

	Left Half-Antibody	**Right Half-Antibody**	**Illustrative Figure**
1.	Exemplary Monomer 2	Exemplary Monomer 2	2B
2.	Exemplary Monomer 5A	Exemplary Monomer 5A	2C
3.	Exemplary Monomer 6A	Exemplary Monomer 6A	2D
4.	Exemplary Monomer 5B	Exemplary Monomer 5B	2E
5.	Exemplary Monomer 6B	Exemplary Monomer 6B	2F
6.	Exemplary Monomer 2	Exemplary Monomer 1	2G
7.	Exemplary Monomer 2	Exemplary Monomer 7A	2H
8.	Exemplary Monomer 2	Exemplary Monomer 7B	2I
9.	Exemplary Monomer 5A	Exemplary Monomer 1	2J
10.	Exemplary Monomer 5B	Exemplary Monomer 1	2W
11.	Exemplary Monomer 6A	Exemplary Monomer 1	2X
12.	Exemplary Monomer 6B	Exemplary Monomer 1	2K
13.	Exemplary Monomer 8A	Exemplary Monomer 8A	2L
14.	Exemplary Monomer 8B	Exemplary Monomer 8B	2M
15.	Exemplary Monomer 8A	Exemplary Monomer 1	2N
16.	Exemplary Monomer 8B	Exemplary Monomer 1	2O
17.	Exemplary Monomer 5A	Exemplary Monomer 4B	2P

Table 2			
	Left Half-Antibody	**Right Half-Antibody**	**Illustrative Figure**
18.	Exemplary Monomer 5B	Exemplary Monomer 4A	2Q
19.	Exemplary Monomer 1	Exemplary Monomer 3	2R
20.	Exemplary Monomer 1	Exemplary Monomer 9A	2S
21.	Exemplary Monomer 1	Exemplary Monomer 9B	2T
22.	Exemplary Monomer 1	Exemplary Monomer 10A	2U
23.	Exemplary Monomer 1	Exemplary Monomer 10B	2V

[0097] In some embodiments, this configuration is advantageously utilized for IFN receptor agonists comprising a targeting moiety that binds to a TAA or ECM target molecule that is expressed in the tumor environment. Without intending to be bound by theory, the inventors believe that in this configuration, where one or more of the linkers are protease-cleavable linkers, the targeting moiety targets the IFN receptor agonist to the tumor environment, where proteases cleave the protease-cleavable linkers resulting in the release of an IFN protein comprising the IFN moiety and linker sequences. This locally activated IFN protein then induces an immune response against the cancer cells. Table 3 below shows additional Exemplary Monomers pairings that can be utilized in the IFN receptor agonists of the disclosure. The IFN receptor agonists identified in Table 3 comprise one or two targeting moieties.

Table 3			
	Left Half-Antibody	**Right Half-Antibody**	**Illustrative Figure**
1.	Exemplary Monomer 2T	Exemplary Monomer 2T	3B
2.	Exemplary Monomer 5AT	Exemplary Monomer 5AT	3C
3.	Exemplary Monomer 6AT	Exemplary Monomer 6AT	3D
4.	Exemplary Monomer 5BT	Exemplary Monomer 5BT	3E

Table 3			
	Left Half-Antibody	**Right Half-Antibody**	**Illustrative Figure**
5.	Exemplary Monomer 6BT	Exemplary Monomer 6BT	3F
6.	Exemplary Monomer 2T	Exemplary Monomer 1T	3G
7.	Exemplary Monomer 2T	Exemplary Monomer 7AT	3H
8.	Exemplary Monomer 2T	Exemplary Monomer 7BT	3I
9.	Exemplary Monomer 5AT	Exemplary Monomer 1T	3J
10.	Exemplary Monomer 5BT	Exemplary Monomer 1T	3W
11.	Exemplary Monomer 6AT	Exemplary Monomer 1T	3X
12.	Exemplary Monomer 6BT	Exemplary Monomer 1T	3K
13.	Exemplary Monomer 8AT	Exemplary Monomer 8AT	3L
14.	Exemplary Monomer 8BT	Exemplary Monomer 8BT	3M
15.	Exemplary Monomer 8AT	Exemplary Monomer 1T	3N
16.	Exemplary Monomer 8BT	Exemplary Monomer 1T	3O
17.	Exemplary Monomer 5AT	Exemplary Monomer 4BT	3P
18.	Exemplary Monomer 5BT	Exemplary Monomer 4AT	3Q
19.	Exemplary Monomer 1T	Exemplary Monomer 3	3R
20.	Exemplary Monomer 1T	Exemplary Monomer 9A	3S
21.	Exemplary Monomer 1T	Exemplary Monomer 9B	3T
22.	Exemplary Monomer 1T	Exemplary Monomer 10A	3U
23.	Exemplary Monomer 1T	Exemplary Monomer 10B	3V

[0098] Sequence and length of hinge and linker sequences can be varied, as can the sequence of the IFN moiety (containing either the full-length or N- and/or C-terminal truncated IFN sequences as well as amino acid substitutions). Exemplary IFN moieties are

described in Section 6.3 and include IFNα- and IFNβ-based moieties as described in Sections 6.3.1 and 6.3.2 below as well as other Type I IFN-based moieties as described in Section 6.3.3. Exemplary IFN receptor moieties are disclosed in Section 6.4. Exemplary protease cleavable linker sequences are disclosed in Section 6.5. Exemplary non-cleavable linker and hinge sequences are disclosed in Sections 6.6 and 6.9.3, respectively. Exemplary targeting moieties are disclosed in Section 6.7. Exemplary Fc domains, including Fc domains suitable for heterodimerization when the two half antibodies of an IFN receptor agonist are not identical, are described in Section 6.9.

6.3. The IFN Moiety

[0099] There are two major classes of IFNs: Type I (IFN-α subtypes, IFN-β, etc.) and Type II (IFN-γ). Additional IFNs (IFN-like cytokines; IFN-λ subtype) have also been identified.

[0100] The IFN moiety of the disclosure may comprise any wild type or modified (*e.g.*, truncated and/or mutant) IFN or IFN-like cytokine sequence but preferably is a Type I IFN moiety. Type I IFNs bind a heterodimeric plasma membrane receptor IFNAR made of IFNAR1 and IFNAR2 that is ubiquitously expressed in all nucleated cells. Ligand binding is initiated by high-affinity receptor subunit IFNAR2 (Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001). As such, Type I IFNs are able to act on virtually all cells of the body. Sixteen Type I interferon subtypes have been identified, which vary in their intrinsic variability in affinity to IFNAR2 and activity.

[0101] In some embodiments, the Type I IFN moiety is an interferon-α (IFN α) moiety. In other embodiments, Type I IFN moiety is an interferon-β (IFNβ) moiety.

[0102] In other embodiments, the Type I IFN moiety is an interferon-ω (IFNω), interferon-ε (IFNε) or interferon-κ (IFNκ) moiety.

[0103] The Type I IFN moiety may comprise a sequence that varies from a wild-type IFN sequence by one or more mutations, *e.g.*, substitutions, deletions, or insertions. Substitutions that attenuate IFN activity by reducing receptor binding may suitably be used. Amino acids with N- or C-terminal deletions (or truncations) may also be used, *e.g.*, a truncation of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids from the N-and/or C-termini of a mature Type I IFN. Without being bound by theory, the present inventors believe that the terminal truncations impose additional steric constraints on the IFN moiety

and reduce IFN activity until cleavage of a protease-cleavable linker in the IFN receptor agonists.

[0104] Further details of exemplary Type I IFN moieties are provided below.

6.3.1. Interferon-α Moieties

[0105] The IFNα gene is a member of the alpha interferon gene cluster on chromosome 9. The encoded cytokine is a member of the Type I interferon family that is produced in response to viral infection as a key part of the innate immune response with potent antiviral, antiproliferative and immunomodulatory properties. IFNα refers to a family of proteins, with at least 15 known subtypes of human IFNα. The major subtypes identified are IFNα1, IFNα2, IFNα8, IFNα10, IFNα14 and IFNα21.

[0106] The IFNα1 gene has two allelic variants: IFNα 1a and IFNα1b. The amino acid sequence of human IFNα1a is assigned UniProtKB accession number P01562, reproduced below with the signal peptide underlined:

MASPFALLMV LVVLSCKSSC SLGCDLPETH SLDNRRTLML LAQMSRISPS SCLMDRHDFG FPQEEFDGNQ FQKAPAISVL HELIQQIFNL FTTKDSSAAW DEDLLDKFCT ELYQQLNDLE ACVMQEERVG ETPLMNADSI LAVKKYFRRI TLYLTEKKYS PCAWEVVRAE IMRSLSLSTN LQERLRRKE (SEQ ID NO: 1)

[0107] The human IFNα1b gene differs the IFNα1a allelic variant by one base change in the coding region, leading to a single change in amino acid sequence (Val114 instead of Ala114 in the mature protein, corresponding to Val137 instead of Ala137 in the full-length polypeptide).

[0108] There are three allelic variants of IFNα2 alleles, IFNα2a, IFNα2b and IFNα2c. Allele IFNα2b is the predominant allele while allele IFNα2a is less predominant and IFNα2c only a minor allelic variant. The amino acid sequence of human IFNα2 is assigned UniProtKB accession number P01563. The sequence of the IFNα2b allele is reproduced below with the signal peptide underlined:

MALTFALLVA LLVLSCKSSC SVGCDLPQTH SLGSRRTLML LAQMRRISLF SCLKDRHDFG FPQEEFGNQF QKAETIPVLH EMIQQIFNLF STKDSSAAWD ETLLDKFYTE LYQQLNDLEA CVIQGVGVTE TPLMKEDSIL AVRKYFQRIT LYLKEKKYSP CAWEVVRAEI MRSFSLSTNL QESLRSKE (SEQ ID NO: 2)

[0109] IFNα2b has an arginine (R) at position 23 of the mature protein while IFNα2a has a lysine (K). Thus, in some embodiments, the IFNα2 moiety has an arginine at the position corresponding to position 23 of the mature protein. In other embodiments, the IFNα2 moiety has a lysine at the position corresponding to position 23 of the mature protein.

[0110] In various aspects, the IFNα moiety comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to the amino acid sequence of mature IFNα1a, IFNα1b, and/or IFNα2b, IFNα2a, or IFNα2c or a fragment thereof having a truncation of up to 15 amino acids at its N- and/or C-terminus (*e.g.*, a truncation of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids from the N-and/or C-termini of mature IFNα1a, IFNα1b, and/or IFNα2b, IFNα2a, or IFNα2c).

[0111] In some embodiments, the IFNα moiety has one or more amino acid substitutions, *e.g.*, substitutions that alter IFNAR binding and/or agonism. Exemplary substitutions are found in WO 2013/107791, U.S. Pat. No. 8,258,263, WO2007/000769A2, WO2008/124086, WO2010/030671, WO2018/144999A1, and WO2015/007520, WO 2013/059885, WO2020156467A1, WO2021/126929A1. In some embodiments, the IFNα moiety comprises:

a) one or more substitutions selected from L15A, A19W, R22A, R23A, L26A, F27A, L30A, L30V, K31A, D32A, R33K or R33A or R33Q, H34A, D35A, Q40A, H57Y, E58N, Q61S, F64A, N65A, T69A, L80A, D82E, Y85A, T86I, Y89A, D114R or D114A, L117A, R120A or R120E or R120K, K121E, R125A, K133A, K134A, R144A, A145G or A145M, M148A, R149A, R149K, S152A, L153A, N156A; and/or

b) one or more substitutions at amino acids 57-89 and 159-165 described in WO2007000769A2; and/or

c) one or more amino acid substitutions at 9, 17, 47, 65, 66, 117, 123, 128, 147, and 157 to alanine, glycine, or threonine described in WO2021126929A1.

[0112] The amino acid positions of the foregoing substitutions are given with reference to mature IFNα2b.

[0113] In further embodiments, the IFNα moiety comprises one or more amino acid substitutions set forth in Table 4. Table 4 sets forth IFNα substitutions identified by reference to the amino acid position within the sequence of IFNα2.

Table 4
Exemplary IFNα mutations (with reference to the sequence of mature IFNα2)

IFNα sequence mutation(s)	Source / impact of substitution(s)
L15A	WO2018014068A9
R22A	WO2018014068A9
R23A	WO2018014068A9
S25A	WO2018014068A9
L26A	Thomas *et al.*, 2011, Cell, 146(4):621-632
F27A	Thomas *et al.*, 2011, Cell, 146(4):621-632
L30A	Thomas *et al.*, 2011, Cell, 146(4):621-632
L30V	WO2018014068A9
K31A	WO2018014068A9
D32A	WO2018014068A9
R33K	WO2018014068A9
R33Q	WO2018014068A9
R33A	WO2013059885A2
H34A	WO2018014068A9
D35E	WO2016201337A1
Q40A	WO2018014068A9
H57A	Thomas *et al.*, 2011, Cell, 146(4):621-632
H57S	WO2013134138A1
E58A	Thomas *et al.*, 2011, Cell, 146(4):621-632
Q61A	WO2016201337A1
Q90A	Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001
E96A	Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001
D114R	WO2018014068A9
L117A	WO2018014068A9
R120A	WO2018014068A9
R120E	WO2018014068A9
Q124R	WO2022015711A1
R125A	WO2018014068A9
R125E	WO2018014068A9
K131A	WO2018014068A9
E132A	WO2018014068A9

Table 4 Exemplary IFNα mutations (with reference to the sequence of mature IFNα2)	
IFNα sequence mutation(s)	**Source / impact of substitution(s)**
K133A	WO2018014068A9
K134A	WO2018014068A9
L135A	WO2013134138A1
R144A	WO2022015711A1
R144D	WO2018014068A9
R144E	WO2018014068A9
R144G	WO2018014068A9
R144H	WO2018014068A9
R144I	WO2018014068A9
R144K	WO2018014068A9
R144L	WO2013059885A2
R144N	WO2018014068A9
R144Q	WO2018014068A9
R144S	WO2013059885A2
R144T	WO2013059885A2
R144V	WO2018014068A9
R144Y	WO2013059885A2
A145D	WO2013059885A2
A145E	WO2018014068A9
A145G	WO2018014068A9, WO2022015711A1
A145H	WO2013059885A2
A145I	WO2018014068A9
A145K	WO2013059885A2
A145L	WO2018014068A9
A145M	WO2018014068A9
A145N	WO2018014068A9
A145Q	WO2018014068A9
A145S	WO2018014068A9
A145T	WO2018014068A9
A145V	WO2018014068A9
A145Y	WO2013059885A2
M148A	WO2018014068A9
R149A	WO2022015711A1, Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001
R149K	Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001

Table 4 Exemplary IFNα mutations (with reference to the sequence of mature IFNα2)	
IFNα sequence mutation(s)	**Source / impact of substitution(s)**
S152A	Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001, WO2022015711A1
L153A	WO2018014068A9
N156A	WO2018014068A9
R162A	WO2016201337A1
E165D	WO2016201337A1
L30A, H57Y, E58N, Q61S	WO2018014068A9
R33A, H57Y, E58N, Q61S	WO2013059885A2
H57S, E58S, Q61S	Alter IFNα-IFNAR1 binding affinity at Site 1
H57Y, E58N, Q61S	WO2007000769A2
N65A, L80A, Y85A, Y89A	WO2018014068A9
N65A, L80A, Y85A, Y89A, D114A	WO2018014068A9
N65A, L80A, Y85A, Y89A, L174A	WO2018014068A9
N65A, L80A, Y85A, Y89A, R120A	WO2018014068A9
Y85A, Y89A, D114A	WO2018014068A9
Q90A, R120A	Piehler *et al.*, 2012, Immunological Reviews, doi.org/10.1111/imr.12001
D114A, R120A	WO2018014068A9
L117A, R120A	WO2018014068A9
L117A, R120A, K121A	WO2018014068A9
R120A, K121A	WO2018014068A9
R120E, K121E	WO2018014068A9
R144A, H57Y, E58N, Q61S	WO2013059885A2
M148A, H57Y, E58N, Q61S	WO2018014068A9
R149A, R162A	WO2013134138A1
L153A, H57Y, E58N, Q61S	WO2018014068A9
Deletion of residues L161 to E165	WO2018014068A9

[0114] In some embodiments, the IFNα moiety comprises an amino acid sequence comprising the amino acid substitution R33A or R33K, Q90A, E96A, R120A, A145M, R149A or R149K, S152A, or any combination of two or more of the foregoing, *e.g.*, Q90A + R120A or A145M + R149K.

[0115] The sequences of exemplary IFNα moieties that can be utilized in the IFN receptor agonists of the disclosure are set forth in Table 5 below:

Table 5
Exemplary IFNα Moieties

Construct	Sequence	SEQ ID NO:
IFNα1a	CDLPETHSLDNRRTLMLLAQMSRISPSSCLMDRHDFGFPQEEFDG NQFQKAPAISVLHELIQQIFNLFTTKDSSAAWDEDLLDKFCTELYQQ LNDLEACVMQEERVGETPLMNADSILAVKKYFRRITLYLTEKKYSP CAWEVVRAEIMRSLSLSTNLQERLRRKE	3
IFNα1b	CDLPETHSLDNRRTLMLLAQMSRISPSSCLMDRHDFGFPQEEFDG NQFQKAPAISVLHELIQQIFNLFTTKDSSAAWDEDLLDKFCTELYQQ LNDLEACVMQEERVGETPLMNVDSILAVKKYFRRITLYLTEKKYSP CAWEVVRAEIMRSLSLSTNLQERLRRKE	4
IFNα2a	CDLPQTHSLGSRRTLMLLAQMRKISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	5
IFNα2b	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	6
IFNα2b (R33A)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDAHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	7
IFNα2b (R33K)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDKHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	8
IFNα2b (Q90A)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYAQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	9
IFNα2b (E96A)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLAACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	10
IFNα2b (R120A)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVAKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE	11
IFNα2b (A145M)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRMEIMRSFSLSTNLQESLRSKE	12

Table 5 Exemplary IFNα Moieties		
Construct	**Sequence**	**SEQ ID NO:**
IFNα2b (R149A)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMASFSLSTNLQESLRSKE	13
IFNα2b (R149K)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMKSFSLSTNLQESLRSKE	14
IFNα2b (S152A)	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFALSTNLQESLRSKE	15
ΔN-IFN-ΔC	THSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDLEA CVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVV RAEIMRSFSLSTNLQ	16
IFN-ΔC	CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGN QFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQL NDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQ	17

6.3.2. Interferon-β Moieties

[0116] Interferon-β (IFNβ) is a cytokine that is naturally produced by the immune system in response to biological and chemical stimuli. IFNβ is a glycosylate, secreted monomer having a molecular weight of around 22kDa that is produced in large quantities by fibroblasts and as such it is also known as fibroblast interferon. IFNβ binds to the IFNAR receptor composed of the IFNAR1 and IFNAR2 dimers to induce signaling via the JAK/STAT pathway and other pathways. IFNβ can also function by binding to IFNAR1 alone and signal independently of the Jak-STAT pathways (Ivashkiv and Donlin, 2014, Nat Rev Immunol. 14(1):36-49).

[0117] IFNβ contains 5 α-helices designated A (YNLLGFLQRSSNFQCQKLL (SEQ ID NO: 18)), B (KEDAALTIYEMLQNIFAIF (SEQ ID NO: 19)), C (ETIVENLLANVYHQINHLKTVLEEKL (SEQ ID NO: 20)), D (SSLHLKRYYGRILHYLKA (SEQ ID NO: 21)), and E (HCAWTIVRVEILRNFYFINRLT (SEQ ID NO: 22)). The five α-helices are interconnected by loops of 2-28 residues designated AB, BC, CD and DE loops. It has

been reported that the A helix in the AB loop and the E helix in the DE loop are involved in the binding of IFNβ to the IFNAR receptor.

[0118] Two types of IFNβ have been described: Interferon-β1 (IFNβ1) and Interferon-β3 (IFNβ3) (Schirmer and Neumann, 2019. Cytokines. In: Nijkamp and Parnham's Principles of Immunopharmacology. Springer, Cham.).

[0119] The amino acid sequence of human IFNβ precursor is listed under GenBank: accession number AAA36040.1 and reproduced below (with the signal peptide underlined):

> MTNKCLLQIA LLLCFSTTAL SMSYNLLGFL QRSSNFQCQK LLWQLNGRLE YCLKDRMNFD IPEEIKQLQQ FQKEDAALTI YEMLQNIFAI FRQDSSSTGW NETIVENLLA NVYHQINHLK TVLEEKLEKE DFTRGKLMSS LHLKRYYGRI LHYLKAKEYS HCAWTIVRVE ILRNFYFINR LTGYLRN (SEQ ID NO: 23)

[0120] In various aspects, the IFNβ moiety comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to the amino acid sequence of mature IFNβ1 or a fragment thereof having a truncation of up to 15 amino acids at its N- and/or C-terminus (*e.g.*, a truncation of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids from the N- and/or C-termini of IFNβ1).

[0121] In various embodiments, the IFNβ moiety comprises one or more amino acid substitutions and/or deletions as compared to IFNβ1. In some embodiments, the substitution is a C17S (with reference to the mature IFNβ1) and the deletions are one of the C- terminal truncations described in US 2009/0025106 A1as IFN-ΔI , IFNA2, IFNA3, IFNA4, IFNA5, IFNA6, IFN-Δ7, IFN-Δδ, IFNA9, and IFN-ΔI O.

6.3.3. Other Type I Interferons

[0122] In certain aspects, the Type I IFN moiety is other than an IFNα or IFNβ moiety, *e.g.*, an interferon-ω (IFNω), interferon-ε (IFNε) or interferon-κ (IFNκ) moiety.

[0123] Human IFNω is identified by UniProt accession no. P05000 and the IFNω1 allele has the amino acid sequence set forth below, with the signal sequence underlined:

> MALLFPLLAALVMTSYSPVGSLGCDLPQNHGLLSRNTLVLLHQMRRISPFLCLKDRR DFRFPQEMVKGSQLQKAHVMSVLHEMLQQIFSLFHTERSSAAWNMTLLDQLHTGL

HQQLQHLETCLLQVVGEGESAGAISSPALTLRRYFQGIRVYLKEKKYSDCAWEVVR MEIMKSLFLSTNMQERLRSKDRDLGSS (SEQ ID NO: 24)

[0124] In various aspects, the IFNω moiety comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to the amino acid sequence of mature IFNω1 or a fragment thereof having a truncation of up to 15 amino acids at its N- and/or C-terminus (*e.g.*, a truncation of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids from the N- and/or C-termini of IFNω1).

[0125] Human IFNε is identified by UniProt accession no. Q86WN2 and has the amino acid sequence set forth below, with the signal sequence underlined:

MIIKHFFGTVLVLLASTTIFSLDLKLIIFQQRQVNQESLKLLNKLQTLSIQQCLPHRKNF LLPQKSLSPQQYQKGHTLAILHEMLQQIFSLFRANISLDGWEENHTEKFLIQLHQQL EYLEALMGLEAEKLSGTLGSDNLRLQVKMYFRRIHDYLENQDYSTCAWAIVQVEIS RCLFFVFSLTEKLSKQGRPLNDMKQELTTEFRSPR (SEQ ID NO: 25)

[0126] In various aspects, the IFNε moiety comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to the amino acid sequence of mature IFNε or a fragment thereof having a truncation of up to 15 amino acids at its N- and/or C-terminus (*e.g.*, a truncation of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids from the N- and/or C-termini of IFNε).

[0127] Human IFNκ is identified by UniProt accession no. Q9P0W0and has the amino acid sequence set forth below, with the signal sequence underlined:

MSTKPDMIQKCLWLEILMGIFIAGTLSLDCNLLNVHLRRVTWQNLRHLSSMSNSFPV ECLRENIAFELPQEFLQYTQPMKRDIKKAFYEMSLQAFNIFSQHTFKYWKERHLKQI QIGLDQQAEYLNQCLEEDKNENEDMKEMKENEMKPSEARVPQLSSLELRRYFHRI DNFLKEKKYSDCAWEIVRVEIRRCLYYFYKFTALFRRK (SEQ ID NO: 26)

[0128] In various aspects, the IFNκ moiety comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about

94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99% or 100% sequence identity to the amino acid sequence of mature IFNκ or a fragment thereof having a truncation of up to 15 amino acids at its N- and/or C-terminus (*e.g.*, a truncation of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids from the N- and/or C-termini of IFNκ).

6.4. IFN Masking Moieties

[0129] The present disclosure provides IFN receptor agonists with the IFN moiety masked by one or more receptor moieties, thereby attenuating IFN activity. All human type I interferons bind to a cell surface receptor (IFN alpha receptor, IFNAR) which is a heterodimer consisting of two transmembrane proteins, IFNAR1 and IFNAR2 (see, *e.g.*, Novick *et al.*, 1994, Cell 77:391), both of which may be used to mask the IFN moiety in the IFN receptor agonists of the disclosure. Thus, in some embodiments, the masking moiety is an IFNAR1 moiety. In other embodiments, the masking moiety is an IFNAR2 moiety. Exemplary IFNAR1 moieties are disclosed in Section 6.4.1 and exemplary IFNAR2 moieties are disclosed in Section 6.4.2.

6.4.1. IFNAR1 Moiety

[0130] IFNAR1 is the lower affinity IFN receptor and belongs to the type II spiral-type cytokine receptors. It includes an extracellular domain that is composed of 4 type III fibronectin domains referred to as "subdomains" (SDs), a transmembrane domain and an intracellular domain of 100 amino acids. The four subdomains of IFNAR1 fold into domain 1 (SD1+SD2) and domain 2 (SD3+SD4).

[0131] The sequence of human IFNAR1 has the UniProt identifier P17181. The sequence of human IFNAR1 is reproduced below:

MMVVLLGATTLVLVAVAPWVLSAAAGG**KNLKSPQKVEVDIIDDNFILRWNRSDESV GNVTFSFDYQKTGMDNWIKLSGCQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTS SWYEVDSFTPFRKAQ**IGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSLVI WKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKIGVYSPVHCIKTTVEN *ELPPPENIEVSVQNQNYVLKWDYTYANMTFQVQWLHAFLKRNPGNHLYKWKQIPD CENVKTTQCVFPQNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQ*afllppvfnirslsds fhiyigapkqsgntpviqdypliyeiifwentsnaerkiiekktdvtvpnlkpltvycvkarahtmdeklnkssvfsdavce ktkpgNTSKIWLIVGICIALFALPFVIYAAKVFLRCINYVFFPSLKPSSSIDEYFSEQPLKN

LLLSTSEEQIEKCFIIENISTIATVEETNQTDEDHKKYSSQTSQDSGNYSNEDESESK TSEELQQDFV (SEQ ID NO: 27)

[0132] The signal sequence (single underline) corresponds to amino acids 1-27, the SD1 domain (bold) corresponds to amino acids 28-127, the SD2 domain (double underline) corresponds to amino acids 128-227, the SD3 domain (italics) corresponds to amino acids 231-329, the SD4 domain (lowercase) corresponds to amino acids 330-432, and the extracellular domain corresponds to amino acids 28-436 of the full length human IFNAR1 protein reproduced above.

[0133] An IFNAR1 moiety is an amino acid sequence comprising at least 70% sequence identity, *e.g.*, at least 70%, 71 %, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81 %, 82%, 83%, 84%, 85%, 86%,87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity, or 100% sequence identity, to an IFN-binding portion of a mammalian, *e.g.*, human, IFNAR1. In some embodiments, the IFN-binding portion comprises the SD2 and SD3 domains. In various aspects, the IFN-binding portion comprises (i) only the SD2 and SD3 domains; (ii) the SD1, SD2 and SD3 domains; (iii) the SD2, SD3 and SD4 domains; (iv) the SD1, SD2, SD3 and SD4 domains; or (v) the entire extracellular domain of IFNAR1.

6.4.2. IFNAR2 Moiety

[0134] IFNAR2 is the high affinity IFN receptor, adopting a two-domain D1/D2 receptor structure.

[0135] The sequence of human IFNAR2 has the UniProt identifier P48551. The sequence of human IFNAR2 is reproduced below:

MLLSQNAFIFRSLNLVLMVYISLVFG**ISYDSPDYTDESCTFKISLRNFRSILSWELKN HSIVPTHYTLLYTIMSKPEDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFS GNTTLFSCSHNFWLAIDMSFEP**PEFEIVGFTNHINVMVKFPSIVEEELQFDLSLVIEE QSEGIVKKHKPEIKGNMSGNFTYIIDKLIPNTNYCVSVYLEHSDEQAVIKSPLKCTLLP PGQESESAESAKIGGIITVFLIALVLTSTIVTLKWIGYICLRNSLPKVLNFHNFLAWPFP NLPPLEAMDMVEVIYINRKKKVWDYNYDDESDSDTEAAPRTSGGGYTMHGLTVRP LGQASATSTESQLIDPESEEEPDLPEVDVELPTMPKDSPQQLELLSGPCERRKSPL QDPFPEEDYSSTEGSGGRITFNVDLNSVFLRVLDDEDSDDLEAPLMLSSHLEEMVD

PEDPDNVQSNHLLASGEGTQPTFPSPSSEGLWSEDAPSDQSDTSESDVDLGDGYI MR (SEQ ID NO: 28)

[0136] The signal sequence (single underline) corresponds to amino acids 1-26, the D1 domain (bold) corresponds to amino acids 27-136, the D2 domain (double underline) corresponds to amino acids 137-232, and the extracellular domain corresponds to amino acids 27-243 of the full length human IFNAR2 protein reproduced above.

[0137] An IFNAR2 moiety is an amino acid sequence comprising at least 70% sequence identity, *e.g.*, at least 70%, 71 %, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81 %, 82%, 83%, 84%, 85%, 86%,87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity, or 100% sequence identity, to an IFN-binding portion of a mammalian, *e.g.*, human, IFNAR2. In some embodiments, the IFN-binding portion comprises the D1 domain. In various aspects, the IFN-binding portion comprises (i) only the D1 domain; (ii) the D1 and D2 domains; or (iii) the entire extracellular domain of IFNAR2.

6.5. Protease-Cleavable Linkers

[0138] The IFN receptor agonists optionally include one or two protease-cleavable linkers (PCLs) in at least one half antibody (or both half antibodies), with other linkers being non-cleavable. In some embodiments, a linker adjacent to an IFN moiety is a protease-cleavable linker. This may lead to for example release of the receptor mask from the IFN moiety upon cleavage of the PCL with the IFN moiety retained on the C-terminus of the half antibody (*e.g.*, in the configuration designated in Table 1 as Fc-IFN-IFNR). Alternatively, for other configurations (*e.g.*, the configuration designated in Table 1 as Fc-IFNR-IFN), incorporating a PCL adjacent to the IFN moiety would result in release of IFN upon cleavage, with the IFNR moiety being retained in the half-antibody. In yet other embodiments, the linker between an Fc domain and an IFNR moiety is a PCL, configured such that cleavage of releases the IFNR moiety while retaining the IFN moiety in a half antibody (*e.g.*, configurations such as Fc-IFN x Fc-R1, where cleavage releases the IFNR moiety while retaining the IFN moiety in the other half antibody).

[0139] A protease-cleavable linker can range from 8 amino acids to 100 or more amino acids. In various embodiments, the protease-cleavable linker ranges from 8 amino acids to 15 amino acids, from 10 amino acids to 20 amino acids, 20 amino acids to 80, and in certain aspects a non-cleavable peptide linker ranges from 20 amino acids to 60 amino acids, 20

amino acids to 40 amino acids, from 30 amino acids to 50 amino acids, from 20 amino acids to 80 amino acids, or from 30 amino acids to 70 amino acids in length.

[0140] The protease-cleavable linkers comprise one or more substrate sequences for one or more proteases, for example one or more of the proteases set forth in Section 6.5.1. The one or more substrate sequences, *e.g.*, one or more of the substrate sequences set forth in Section 6.5.2, are typically (but not necessarily) flanked by one or more spacer sequences, *e.g.*, spacer sequences as described in Section 6.5.3. Each protease-cleavable linker can include one, two, three or more substrate sequences. The spacer sequences can be adjoining, overlapping, or separated by spacer sequences. Preferably, the C- and N-termini of the protease-cleavable linkers contain spacer sequences.

[0141] In various aspects of IFN receptor agonists comprising four protease-cleavable linkers, the first and third protease-cleavable linkers are cleavable by the same protease and/or the second and fourth protease-cleavable linkers are cleavable by the same protease. In some embodiments, the protease is a protease set forth in Table A.

[0142] In further aspects of IFN receptor agonists comprising four protease-cleavable linkers, the first and third protease-cleavable linkers comprise the same substrate sequence(s) and/or the second and fourth protease-cleavable linkers comprise the same substrate sequence(s). In some embodiments, the substrate sequence(s) are set forth in Table B. In further embodiments, the first and third protease-cleavable linkers also comprise the same spacer sequence(s) and/or the second and fourth protease-cleavable linkers also comprise the same spacer sequence(s). In some embodiments, the spacer sequence(s) are set forth in Table C.

[0143] In further aspects IFN receptor agonists comprising four protease-cleavable linkers, the first and third linkers comprise the same linker sequence(s) and/or the second and fourth linkers comprise the same linker sequence(s). In some embodiments, the linker sequence(s) are set forth in Table D.

[0144] In some embodiments of IFN receptor agonists comprising four protease-cleavable linkers, the first and third protease-cleavable linkers are the same as the second and fourth protease-cleavable linkers.

[0145] In other embodiments, the first and third protease-cleavable linkers are different from the second and fourth protease-cleavable linkers.

[0146] In the foregoing aspects and embodiments, the different linkers may be cleavable by the same protease, different proteases, or when a linker comprises multiple substrate sequences, the different linkers may be cleavable by multiple proteases, one or more of which are common and one or more of which are different.

[0147] Exemplary protease-cleavable linker sequences are set forth in Section 6.5.4.

6.5.1. Proteases

[0148] Exemplary protease whose substrate sequences can be incorporated into the protease-cleavable linkers are set forth in Table A below.

TABLE A
Exemplary Proteases for Substrate Cleavage

ADAMS, ADAMTS, *e.g.*	Caspases, *e.g.,*	MMP24
ADAM8	Caspase 1	MMP26
ADAM9	Caspase 2	MMP27
ADAM10	Caspase 3	
ADAM12	Caspase 4	
ADAM15	Caspase 5	
ADAM17/TACE	Caspase 6	
ADAMDEC1	Caspase 7	
ADAMTS1	Caspase 8	Cysteine proteinases, *e.g.*,
ADAMTS4	Caspase 9	Cruzipain
ADAMTS5	Caspase 10	Legumain
	Caspase 14	Otubain-2
Aspartate proteases, *e.g.,*		
BACE	Cysteine cathepsins, *e.g.,*	KLKs, *e.g.*,
Renin	Cathepsin B	KLK4
	Cathepsin C	KLK5
Aspartic cathepsins, *e.g.*,	Cathepsin K	KLK6
Cathepsin D	Cathepsin L	KLK7
Cathepsin E	Cathepsin S	KLK8
	Cathepsin V/L2	KLK10
NS3/4A	Cathepsin X/Z/P	KLK11
PACE4		KLK13
Plasmin	MMPs, *e.g.*,	KLK14
PSA	MMP1	
tPA	MMP2	Metallo proteinases, *e.g.*,
Thrombin	MMP3	Meprin

TABLE A Exemplary Proteases for Substrate Cleavage		
Tryptase uPA	MMP7 MMP8 MMP9 MMP10	Neprilysin PSMA BMP-1
Type II Transmembrane Serine Proteases (TTSPs), *e.g.*,		
DESC1 DPP-4 FAP Hepsin Matriptase-2 MT/SP1/Matriptase	MMP11 MMP12 MMP13 MMP14 MMP15 MMP16 MMP17	Serine proteases, *e.g.*, activated protein C Cathepsin A Cathepsin G Chymase coagulation factor proteases (*e.g.*, FVIIa, FIXa, FXa, FXIa, FXIIa)
TMPRSS2 TMPRSS3 TMPRSS4	MMP19 MMP20 MMP23	Human Neutrophil Elastase Lactoferrin

[0149] In particular embodiments, the protease is matrix metalloprotease (MMP)-2, MMP-9, legumain asparaginyl endopeptidase, thrombin, fibroblast activation protease (FAP), MMP-1, MMP-3, MMP-7, MMP-8, MMP-12, MMP-13, MMP-14, membrane type 1 matrix metalloprotease (MT1-MMP), plasmin, transmembrane protease, serine (TMPRSS-3/4), cathepsin A, cathepsin B, cathepsin D, cathepsin E, cathepsin F, cathepsin H, cathepsin K, cathepsin L, cathepsin L2, cathepsin O, cathepsin S, caspase 1, caspase 2, caspase 3, caspase 4, caspase 5, caspase 6, caspase 7, caspase 8, caspase 9, caspase 10, caspase 11, caspase 12, caspase 13, caspase 14, human neutrophil elastase, urokinase/urokinase-type plasminogen activator (uPA), a disintegrin and metalloprotease (ADAM)10, ADAM12, ADAM17, ADAM with thrombospondin motifs (ADAMTS), ADAMTS5, beta secretase (BACE), granzyme A, granzyme B, guanidinobenzoatase, hepsin, matriptase, matriptase 2, meprin, neprilysin, prostate-specific membrane antigen (PSMA), tumor necrosis factor-converting enzyme (TACE), kallikrein-related peptidase (KLK)3, KLK5, KLK7, KLK11, NS3/4 protease of hepatitis C virus (HCV-NS3/4), tissue plasminogen activator (tPA), calpain, calpain 2, glutamate carboxypeptidase II, plasma kallikrein, AMSH-like protease, AMSH, γ-secretase component, antiplasmin cleaving enzyme (APCE), decysin 1, apoptosis-related cysteine peptidase, or N-acetylated alpha-linked acidic dipeptidase-like 1.

6.5.2. Substrates

[0150] Exemplary substrate sequences that are cleavable by a tumor protease and can be incorporated into the protease-cleavable linkers are set forth in Table B below.

TABLE B
Substrate Sequences for Protease-Cleavable Linkers

Substrate Sequence	Designation	Cleaving Protease	SEQ ID NO:
$(DE)_8$RPLALWRS$(DR)_8$	SU1	MMP7	29
AARGPAIH	SU2		30
AAYHLVSQ	SU3	Collagenase	31
AGLGISST	SU4	Collagenase	32
AGLGVVER	SU5	Collagenase	33
ALAL	SU6	Lysosomal Enzyme	34
ALFFSSPP	SU7		35
ALFKSSFP	SU8		36
ALLLALL	SU9	TOP	37
AQFVLTEG	SU10	Collagenase	38
AQNLLGMV	SU11		39
AVGLLAPP	SU12	Serine protease	40
DAFK	SU13	Urokinase plasminogen activator (uPA)	41
DEVD	SU14	Caspase-3	42
DEVDP	SU15	Caspase-3	43
DPRSFL	SU16	Thrombin	44
DVAQFVLT	SU17	Collagenase	45
DVLK	SU18	Plasmin	46
DWLYWPGI	SU19		47
EDDDDKA	SU20	Enterokinase	48
EP(Cit)G(Hof)YL	SU21	MMP2, MMP9, MMP14	49
EPQALAMS	SU22	Collagenase	50
ESLPVVAV	SU23	Collagenase	51
ESPAYYTA	SU24	MMP	52
F(Pip)RS	SU25	Thrombin	
FK	SU26	Lysosomal Enzyme	
FPRPLGITGL	SU27		53
FRLLDWQW	SU28		54
GFLG	SU29	Lysosomal Enzyme	55
GGAANLVRGG	SU30	MMP11	56
GGGRR	SU31	Urokinase plasminogen activator (uPA)	57
GGPRGLPG	SU32	Cathepsin K	58
GGQPSGMWGW	SU33		59

TABLE B Substrate Sequences for Protease-Cleavable Linkers			
Substrate Sequence	**Designation**	**Cleaving Protease**	**SEQ ID NO:**
GGSIDGR	SU34	Factor Xa	60
GGWHTGRN	SU35		61
GIAGQ	SU36	Collagenase	62
GKAFRR	SU37	Kallikrein 2	63
GPAGLYAQ	SU38		64
GPAGMKGL	SU39		65
GPEGLRVG	SU40	Collagenase	66
GPLGIAGI	SU41	Collagenase	67
GPLGVRG	SU42		68
GPQGIAGQ	SU43	Collagenase	69
GPQGLLGA	SU44	Collagenase	70
GPRSFG	SU45		71
GPRSFGL	SU46		72
GPSHLVLT	SU47		73
GVSQNYPIVG	SU48	HIV Protease	74
GVVQASCRLA	SU49	CMV Protease	75
GWEHDG	SU50	Interleukin 1β converting enzyme	76
HSSKLQ	SU51	Prostate Specific Antigen	77
HSSKLQEDA	SU52	Prostate Specific Antigen	78
HSSKLQL	SU53	Prostate Specific Antigen	79
HTGRSGAL	SU54		80
IDGR	SU55	Factor Xa	81
IEGR	SU56	Factor Xa	82
ILPRSPAF	SU57		83
IPVSLRSG	SU58	MMP	84
ISSGL	SU59	MMP	85
ISSGLL	SU60	MMP	86
ISSGLLS	SU61	MMP	87
ISSGLLSS	SU62	MMP	88
ISSGLSS	SU63	MMP	89
KGSGDVEG	SU64	Caspase-3	90
KQEQNPGST	SU65	FAP	91
KRALGLPG	SU66	MMP7	92
LAAPLGLL	SU67		93
LAPLGLQRR	SU68		94
LAQKLKSS	SU69		95

TABLE B Substrate Sequences for Protease-Cleavable Linkers			
Substrate Sequence	**Designation**	**Cleaving Protease**	**SEQ ID NO:**
LAQRLRSS	SU70		96
LEATA	SU71	MMP9	97
LKAAPRWA	SU72		98
LLAPSHRA	SU73		99
LPGGLSPW	SU74		100
LSGRSANI	SU75	Serine protease	101
LSGRSANP	SU76	Serine protease	102
LSGRSDDH	SU77	Serine protease	103
LSGRSDIH	SU78	Serine protease	104
LSGRSDNH	SU79	Serine protease	105
LSGRSDNI	SU80	Serine protease	106
LSGRSDNP	SU81	Serine protease	107
LSGRSDQG	SU82	Serine protease	108
LSGRSDQH	SU83	Serine protease	109
LSGRSDTH	SU84	Serine protease	110
LSGRSDYH	SU85	Serine protease	111
LSGRSGNH	SU86	Serine protease	112
LVLASSSFGY	SU87	Herpes Simplex Virus Protease	113
MDAFLESS	SU88	Collagenase	114
MGLFSEAG	SU89		115
MIAPVAYR	SU90		116
MVLGRSLL	SU91		117
NLL	SU92	Cathepsin B	
NTLSGRSENHSG	SU93		118
NTLSGRSGNHGS	SU94		119
PAGLWLDP	SU95		120
PGGPAGIG	SU96		121
PIC(Et)FF	SU97	Cathepsin D	122
PLGC(me)AG	SU98	MMP	123
PLGL	SU99		124
PLGLAG	SU100	MMP	125
PLGLAX	SU101	MMP	126
PLGLWA	SU102	MMP	127
PLGLWSQ	SU103	MMP	128
PLTGRSGG	SU104		129
PMAKK	SU105		130
PPRSFL	SU106	Thrombin	131
PR(S/T)(L/I)(S/T)	SU107	MMP9	
PRFRIIGG	SU108	Plasmin	132

TABLE B Substrate Sequences for Protease-Cleavable Linkers			
Substrate Sequence	**Designation**	**Cleaving Protease**	**SEQ ID NO:**
PVGYTSSL	SU109		133
PVQPIGPQ	SU110	Collagenase	134
QALAMSAI	SU111	Collagenase	135
QGRAITFI	SU112		136
QNQALRMA	SU113		137
RGPA	SU114		138
RGPAFNPM	SU115		139
RGPATPIM	SU116		140
RKSSIIRMRDVVL	SU117	Plasmin	141
RLQLKAC	SU118	MMP	142
RLQLKL	SU119	MMP	143
RMHLRSLG	SU120		144
RPSPMWAY	SU121		145
RQARVVNG	SU122	Matriptase	146
SAGFSLPA	SU123		147
SAPAVESE	SU124	Collagenase	148
SARGPSRW	SU125		149
SGEPAYYTA	SU126		150
SGGPLGVR	SU127		151
SGRIGFLRTA	SU128	MMP14	152
SGRSA	SU129	Urokinase plasminogen activator (uPA)	153
SGRSANPRG	SU130		154
SMLRSMPL	SU131		155
SPLPLRVP	SU132		156
SPLTGRSG	SU133		157
SPRSIMLA	SU134		158
SSRGPAYL	SU135		159
SSRHRRALD	SU136	Plasmin	160
SSSFDKGKYKKGDDA	SU137	Plasmin	161
SSSFDKGKYKRGDDA	SU138	Plasmin	162
STFPFGMF	SU139		163
TARGPSFK	SU140		164
TGRGPSWV	SU141		165
TSGRSANP	SU142		166
TSTSGRSANPRG	SU143		167
VAGRSMRP	SU144		168
VAQFVLTE	SU145	Collagenase	169
VHMPLGFLGP	SU146		170
VPLSLYSG	SU147	MMP9	171

TABLE B Substrate Sequences for Protease-Cleavable Linkers			
Substrate Sequence	**Designation**	**Cleaving Protease**	**SEQ ID NO:**
VVPEGRRS	SU148		172
WATPRPMR	SU149		173
YGAGLGVV	SU150	Collagenase	174
HPVGLLAR	SU151		175

6.5.3. Spacers

[0151] Exemplary spacer sequences that can be incorporated into the protease-cleavable linkers are set forth in Table C below. In addition to the spacer sequences set forth in Table C, any of the non-cleavable linker sequences described in Section 6.6, *e.g.*, the non-cleavable linker sequences set forth in Table E, or portions thereof can be used as spacer sequences. In some embodiments, spacer sequences are absent entirely from the protease-cleavable linkers.

TABLE C Spacer Sequences for Protease-Cleavable Linkers		
Spacer Sequence	**Designation**	**SEQ ID NO:**
$(GGGGS)_n$	SP1	176
$(GGGS)_n$	SP2	177
$(GGS)_n$	SP3	178
$(GS)_n$	SP4	179
$(GSGGS)_n$	SP5	180
GGGGSGGGGS	SP6	181
GGGGSGGGGSGGGGS	SP7	182
GGGGSGGGGSGGGGSGGGGS	SP8	183
GGGKSGGGKSGGGKS	SP9	184
GGGKSGGKGSGKGGS	SP10	185
GGGS	SP11	186
GGGSG	SP12	187
GGKGSGGKGSGGKGS	SP13	188
GGSGGGGSGGGGS	SP14	189
GGSGGS	SP15	190
GGSGGSGGSGS	SP16	191
GSGGG	SP17	192
GSGSG	SP18	193
GSS	SP19	
GSSG	SP20	194
GSSGGSGGSG	SP21	195

TABLE C Spacer Sequences for Protease-Cleavable Linkers		
Spacer Sequence	**Designation**	**SEQ ID NO:**
GSSGGSGGSGG	SP22	196
GSSGGSGGSGGS	SP23	197
GSSGGSGGSGGSG	SP24	198
GSSGGSGGSGGSGGGS	SP25	199
GSSGGSGGSGS	SP26	200
GSSGT	SP27	201
GSSSG	SP28	202
QGQSGQ	SP29	203
QGQSGQG	SP30	204
QGQSGS	SP31	205
QSGQ	SP32	206
QSGQG	SP33	207
QSGS	SP34	208
SGQ	SP35	
SGQG	SP36	209
SGS	SP37	
$(G)_n$	SP38	210

[0152] In some embodiments, as used in Table C above, n is an integer from 1 to 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10.

6.5.4. Exemplary Protease-Cleavable Linkers

[0153] Exemplary protease-cleavable linkers comprising one or more substrate sequences as well as spacer sequences are set forth in Table D below.

TABLE D Protease-Cleavable Linker Sequences			
Linker Sequence	**Designation**	**Cleaving Protease(s)**	**SEQ ID NO:**
GGGISSGLLSGRSDNHGGGISSGLLSGRSDNHGGS	PCL1		211
GGGISSGLLSGRSDNHGGGISSGLLSGRSDNHGGS GGGISSGLLSGRSDNHGGGISSGLLSGRSDNHGGS	PCL2		212
GGSGGSIPVSLRSGGGISSGLLSGRSDNHGGSGGS	PCL3		213
GGSGGSVPLSLYSGGGISSGLLSGRSDNHGGSGGS	PCL4		214

TABLE D Protease-Cleavable Linker Sequences			
Linker Sequence	**Designation**	**Cleaving Protease(s)**	**SEQ ID NO:**
GGSHPVGLLARGGGHPVGLLAR GGGHPVGLLARGS	PCL5		215
GGSHPVGLLARGGGHPVGLLAR GGSGRSAGGSGRSA	PCL6		216
AVGLLAPPGGLSGRSANI	PCL7	ADAM17_2, FAPa_1, CTSL1_1	217
AVGLLAPPGGLSGRSANP	PCL8	FAPa_1, ADAM17_2, CTSL1_1	218
AVGLLAPPGGLSGRSDDH	PCL9	MMP14_1, MMP14_1, MMP14_1	219
AVGLLAPPGGLSGRSDIH	PCL10	MMP14_1, MMP14_1, MMP14_1	220
AVGLLAPPGGLSGRSDNH	PCL11	MMP14_1, MMP14_1	221
AVGLLAPPGGLSGRSDNI	PCL12	MMP14_1, CTSL1_1, ADAM17_2	222
AVGLLAPPGGLSGRSDNP	PCL13	CTSL1_1, ADAM17_2, FAPa_1	223
AVGLLAPPGGLSGRSDQH	PCL14		224
AVGLLAPPGGLSGRSDTH	PCL15	FAPa_1, CTSL1_1, ADAM17_2	225
AVGLLAPPGGLSGRSDYH	PCL16		226
AVGLLAPPGGTSTSGRSANPRG	PCL17		227
AVGLLAPPSGRSANPRG	PCL18		228
AVGLLAPPTSGRSANPRG	PCL19		229
GGALFKSSFPGPAGLYAQPLAQK LKSSGGK	PCL20	CTSL1_1, MMP14_1, ADAM17_2	230
GGGGSGGGGSGGGGSFVGGTG GGGSGGGGSGGS	PCL21		231
GGGGSGGGGSGGGGSISSGLLS GRSDNHGGSGGS	PCL22		232
GGGGSGGGGSGGGGSVPLSLYS GGGSGGSGGSGS	PCL23		233
GGGGSGGGGSGPLGLWSQGGG GSGGGGSGGGGSGG	PCL24		234
GGGGSGGGGSKKAAPGGGGSG GGGSGGGGSGGS	PCL25		235
GGGGSGGGGSKKAAPVNGGGG GSGGGGSGGGGS	PCL26		236

TABLE D Protease-Cleavable Linker Sequences			
Linker Sequence	**Designation**	**Cleaving Protease(s)**	**SEQ ID NO:**
GGGGSGGGGSPMAKKGGGGSG GGGSGGGGSGGS	PCL27		237
GGGGSGGGGSPMAKKVNGGGG GSGGGGSGGGGS	PCL28		238
GGGGSGGGGSQARAKGGGGSG GGGSGGGGSGGS	PCL29		239
GGGGSGGGGSQARAKVNGGGG GSGGGGSGGGGS	PCL30		240
GGGGSGGGGSRQARVVNGGGG GSGGGGSGGGGS	PCL31		241
GGGGSGGGGSRQARVVNGGGG GSVPLSLYSGGGGGSGGGGS	PCL32		242
GGGGSGGGGSRQARVVNSVPLS LYSGGGGGSGGGGS	PCL33		243
GGGGSGGGGSVHMPLGFLGPG GGGSGGGGSGGS	PCL34		244
GGGGSVHMPLGFLGPGRSRGSF PGGGGS	PCL35		245
GGGGSVHMPLGFLGPPMAKKGG GGSGGGGSGGS	PCL36		246
GGGGSVHMPLGFLGPRQARVVN GGGGSGGGGS	PCL37		247
GGGGSVHMPLGFLGPRQARVVN GGGGSGGGGSGG	PCL38		248
GGPLAQKLKSSALFKSSFPGPAG LYAQGGR	PCL39	ADAM17_2, CTSL1_1, MMP14_1	249
GLSGRSDNHGGAVGLLAPP	PCL40		250
GLSGRSDNHGGVHMPLGFLGP	PCL41		251
ISSGLLSGRSANI	PCL42	MMP, Serine protease	252
ISSGLLSGRSANP	PCL43	MMP, Serine protease	253
ISSGLLSGRSANPRG	PCL44	MMP, Serine protease	254
ISSGLLSGRSDDH	PCL45	MMP, Serine protease	255
ISSGLLSGRSDIH	PCL46	MMP, Serine protease	256
ISSGLLSGRSDNH	PCL47	MMP, Serine protease	257
ISSGLLSGRSDNI	PCL48	CTSL1_1, MMP14_1	258
ISSGLLSGRSDNP	PCL49	MMP, Serine protease	259
ISSGLLSGRSDQH	PCL50	MMP, Serine protease	260
ISSGLLSGRSDTH	PCL51	MMP, Serine protease	261
ISSGLLSGRSDYH	PCL52	MMP, Serine protease	262

TABLE D Protease-Cleavable Linker Sequences			
Linker Sequence	**Designation**	**Cleaving Protease(s)**	**SEQ ID NO:**
ISSGLLSGRSGNH	PCL53	MMP, Serine protease	263
ISSGLLSSGGSGGSLSGRSDNH	PCL54		264
ISSGLLSSGGSGGSLSGRSGNH	PCL55		265
KGGPGGPAGIGPLAQRLRSSALF KSSFPGR	PCL56	FAPa_1, ADAM17_1, CTSL1_1	266
KSGPGGPAGIGALFFSSPPLAQKL KSSGGR	PCL57	FAPa_1, CTSL1_2, ADAM17_2	267
LSGRSDNHGGAVGLLAPP	PCL58		268
LSGRSDNHGGSGGSISSGLLSS	PCL59		269
LSGRSDNHGGSGGSQNQALRMA	PCL60		270
LSGRSDNHGGVHMPLGFLGP	PCL61		271
LSGRSGNHGGSGGSISSGLLSS	PCL62		272
LSGRSGNHGGSGGSQNQALRMA	PCL63		273
QNQALRMAGGSGGSLSGRSDNH	PCL64		274
QNQALRMAGGSGGSLSGRSGNH	PCL65		275
RGGALFKSSFPLAQKLKSSGPAG LYAQGGK	PCL66	CTSL1_1, ADAM17_2, MMP14_1	276
RGGGPAGLYAQPLAQKLKSSALF KSSFPGG	PCL67	MMP14_1, ADAM17_2, CTSL1_1	277
SGGFPRSGGSFNPRTFGSKRKR RGSRGGGG	PCL68	thrombin, factor Xa, hepsin	278
SGPLAQKLKSSGPAGLYAQALFK SSFPGSK	PCL69	ADAM17_2, MMP14_1, CTSL1_1	279
TSTSGRSANPRGGGAVGLLAPP	PCL70		280
TSTSGRSANPRGGGVHMPLGFL GP	PCL71		281
VHMPLGFLGPGGLSGRSDNH	PCL72		282
VHMPLGFLGPGGTSTSGRSANP RG	PCL73		283
SGRSAGGGSGRSAGGGSGRSA	PCL74	uPA	284
HPVGLLARGGGHPVGLLARGGG SGRSAGGGSGRSA	PCL75	MPA (MMP-2 and uPA)	285
GPLGVRGK	PCL76	MMP-2	286
HPVGLLAR	PCL77	MMP-2	175
GPQGIAGQ	PCL78	MMP-2, MMP-9, and to some degree MT1-MMP	69
VPMSMRGG	PCL79	MMP-9 and MMP-2	287
IPVSLRSG	PCL80	MMP-2, and to some degree MMP-9 or MMP-7	84

TABLE D Protease-Cleavable Linker Sequences			
Linker Sequence	**Designation**	**Cleaving Protease(s)**	**SEQ ID NO:**
RPFSMIMG	PCL81	MMP-9 and MMP-7, to some degree MMP-3	288
VPLSLTMG	PCL82	MMP-7, to some degree MMP-9, MMP-2, MPT-1-MMP	289
VPLSLYSG	PCL83	MMP-2, MMP-9, MMP-7	171
IPESLRAG	PCL84	MMP-2, MMP-7, MMP-9, to some degree MPT-1-MMP	290
VPLSLYSGGGISSGLLSGRSDNH	PCL85		291
GGGISSGLLSGRSDNHGGGS	PCL86		292
GGGHPVGLLARGGGS	PCL87		293
GGGSGGGSGGGGISSGLLSGRSDNHGGGSGGGSGGS	PCL88		294
GGGGISSGLLSGRSDNHGGGISSGLLSGRSDNHGGS	PCL89		295
GGGSGGSIPVSLRSGGGISSGLLSGRSDNHGGSGGS	PCL90		296
GGGSGGSVPLSLYSGGGISSGLLSGRSDNHGGSGGS	PCL91		297
GGGSHPVGLLARGGGHPVGLLARGGGHPVGLLARGS	PCL92		298
GGGSHPVGLLARGGGHPVGLLARGGSGRSAGGSGRS	PCL93		299
GISSGLLSGRSDNHG	PCL94		300
GGGSISSGLLSGRSDNHGGGS	PCL95		301

[0154] In certain aspects, the protease-cleavable linker comprises an amino acid sequence having up to 5, up to 4, up to 3, up to 2 or up to 1 amino acid substitution(s) as compared to the sequence set forth in Table D. Thus, in some embodiments, the protease-cleavable linker comprises or consists of any amino acid sequence in Table D with 1-5 amino acid substitutions as compared to the sequence set forth in Table D.

6.6. Non-Cleavable Linkers

[0155] In certain aspects, the present disclosure provides IFN receptor agonists in which two or more components of an IFN receptor agonist are connected to one another by a

peptide linker. By way of example and not limitation, linkers can be used to connect an Fc domain and a targeting moiety, different domains within a targeting moiety (*e.g.,* VH and VL domains in an scFv), an Fc domain and an IFN or IFNR moiety, or an IFN moiety and an IFNR moiety.

[0156] Preferably, all linkers in the IFN receptor agonist other than the specified protease-cleavable linkers (when present) are non-cleavable linkers (NCLs).

[0157] A non-cleavable linker can range from 2 amino acids to 60 or more amino acids, and in certain aspects a non-cleavable peptide linker ranges from 3 amino acids to 50 amino acids, from 4 to 30 amino acids, from 5 to 25 amino acids, from 10 to 25 amino acids, 10 amino acids to 60 amino acids, from 12 amino acids to 20 amino acids, from 20 amino acids to 50 amino acids, or from 25 amino acids to 35 amino acids in length.

[0158] In particular aspects, a non-cleavable linker is at least 5 amino acids, at least 6 amino acids or at least 7 amino acids in length and optionally is up to 30 amino acids, up to 40 amino acids, up to 50 amino acids or up to 60 amino acids in length.

[0159] In some embodiments of the foregoing, the non-cleavable linker ranges from 5 amino acids to 50 amino acids in length, *e.g.*, ranges from 5 to 50, from 5 to 45, from 5 to 40, from 5 to 35, from 5 to 30, from 5 to 25, or from 5 to 20 amino acids in length. In other embodiments of the foregoing, the non-cleavable linker ranges from 6 amino acids to 50 amino acids in length, *e.g.*, ranges from 6 to 50, from 6 to 45, from 6 to 40, from 6 to 35, from 6 to 30, from 6 to 25, or from 6 to 20 amino acids in length. In yet other embodiments of the foregoing, the non-cleavable linker ranges from 7 amino acids to 50 amino acids in length, *e.g.*, ranges from 7 to 50, from 7 to 45, from 7 to 40, from 7 to 35, from 7 to 30, from 7 to 25, or from 7 to 20 amino acids in length.

[0160] Charged (*e.g.,* charged hydrophilic linkers) and/or flexible non-cleavable linkers are particularly preferred.

[0161] Examples of flexible non-cleavable linkers that can be used in the IFN receptor agonists of the disclosure include those disclosed by Chen *et al.*, 2013, Adv Drug Deliv Rev. 65(10): 1357-1369 and Klein *et al.*, 2014, Protein Engineering, Design & Selection 27(10): 325-330. Particularly useful flexible non-cleavable linkers are or comprise repeats of glycines and serines, *e.g.,* a monomer or multimer of G_nS (SEQ ID NO: 302) or SG_n (SEQ ID

NO: 303), where n is an integer from 1 to 10, *e.g.,* 1 2, 3, 4, 5, 6, 7, 8, 9 or 10. In one embodiment, the non-cleavable linker is or comprises a monomer or multimer of repeat of G_4S (SEQ ID NO: 304) *e.g.,* $(GGGGS)_n$ (SEQ ID NO: 304).

[0162] Polyglycine non-cleavable linkers can suitably be used in the IFN receptor agonists of the disclosure. In some embodiments, a peptide non-cleavable linker comprises two consecutive glycines (2Gly), three consecutive glycines (3Gly), four consecutive glycines (4Gly (SEQ ID NO: 305)), five consecutive glycines (5Gly (SEQ ID NO: 306)), six consecutive glycines (6Gly (SEQ ID NO: 307)), seven consecutive glycines (7Gly (SEQ ID NO: 308)), eight consecutive glycines (8Gly (SEQ ID NO: 309)) or nine consecutive glycines (9Gly (SEQ ID NO: 310)).

[0163] Exemplary non-cleavable linker sequences are set forth in Table E below.

TABLE E
Non-Cleavable Linker Sequences

Linker Sequence	**Designation**	**SEQ ID NO:**
GGGGSLALGPGGGGGSLALGPGGGGGSLALGPGGS	NCL1	311
GGGGSGGGGSGGGGSGGGGSGGGGS	NCL2	312
$(GGGGS)_n$, optionally where n=1-10, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10	NCL3	176
$(GGGS)_n$, optionally where n=1-10, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10	NCL4	177
$(GGS)_n$, optionally where n=1-10, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10	NCL5	178
$(GS)_n$, optionally where n=1-10, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10	NCL6	179
$(GSGGS)_n$, optionally where n=1-10, *e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10	NCL7	180
ADAAP	NCL8	318
ADAAPTVSIFP	NCL9	319
ADAAPTVSIFPP	NCL10	320
AKTTAP	NCL11	321
AKTTAPSVYPLAP	NCL12	322
AKTTPKLEEGEFSEARV	NCL13	323
AKTTPKLGG	NCL14	324
AKTTPP	NCL15	325
AKTTPPSVTPLAP	NCL16	326
ASTKGP	NCL17	327
ASTKGPSVFPLAPASTKGPSVFPLAP	NCL18	328
EGKSSGSGSESKST	NCL19	329
GEGESGEGESGEGES	NCL20	330
GEGESGEGESGEGESGEGES	NCL21	331
GEGGSGEGGSGEGGS	NCL22	332
GENKVEYAPALMALS	NCL23	333

TABLE E Non-Cleavable Linker Sequences		
Linker Sequence	**Designation**	**SEQ ID NO:**
GGEGSGGEGSGGEGS	NCL24	334
GGGESGGEGSGEGGS	NCL25	335
GGGESGGGESGGGES	NCL26	336
GGGGSGGGGS	NCL27	181
GGGGSGGGGSGGGGS	NCL28	182
GGGGSGGGGSGGGGSGGGGS	NCL29	183
GGGKSGGGKSGGGKS	NCL30	184
GGGKSGGKGSGKGGS	NCL31	185
GGGS	NCL32	186
GGGSG	NCL33	187
GGKGSGGKGSGGKGS	NCL34	188
GGSG	NCL35	337
GGSGG	NCL36	338
GGSGGGGSG	NCL37	339
GGSGGGGSGGGGS	NCL38	189
GHEAAAVMQVQYPAS	NCL39	340
GKGGSGKGGSGKGGS	NCL40	341
GKGKSGKGKSGKGKS	NCL41	342
GKGKSGKGKSGKGKSGKGKS	NCL42	343
GKPGSGKPGSGKPGS	NCL43	344
GKPGSGKPGSGKPGSGKPSGS	NCL44	345
GPAKELTPLKEAKVS	NCL45	346
GSAGSAAGSGEF	NCL46	347
GSGGG	NCL47	192
GSGSG	NCL48	193
GSS	NCL49	
GSSG	NCL50	194
GSSGGSGGSG	NCL51	195
GSSGGSGGSGG	NCL52	196
GSSGGSGGSGGS	NCL53	197
GSSGGSGGSGGSG	NCL54	198
GSSGGSGGSGGSGGGS	NCL55	199
GSSGGSGGSGS	NCL56	200
GSSGT	NCL57	201
GSSSG	NCL58	202
GSTSGSGKPGSGEGSTKG	NCL59	348
GTAAAGAGAAGGAAAGAAG	NCL60	349
GTSGSSGSGSGGSGSGGGG	NCL61	350
IRPRAIGGSKPRVA	NCL62	351

TABLE E Non-Cleavable Linker Sequences		
Linker Sequence	**Designation**	**SEQ ID NO:**
KESGSVSSEQLAQFRSLD	NCL63	352
KTTPKLEEGEFSEAR	NCL64	353
PRGASKSGSASQTGSAPGS	NCL65	354
QPKAAP	NCL66	355
QPKAAPSVTLFPP	NCL67	356
RADAAAA$(G4S)_4$	NCL68	357
RADAAAAGGPGS	NCL69	358
RADAAP	NCL70	359
RADAAPTVS	NCL71	360
SAKTTP	NCL72	361
SAKTTPKLEEGEFSEARV	NCL73	362
SAKTTPKLGG	NCL74	363
STAGDTHLGGEDFD	NCL75	364
TVAAP	NCL76	365
TVAAPSVFIFPP	NCL77	366
TVAAPSVFIFPPTVAAPSVFIFPP	NCL78	367
AGSGNSSGSGGSGGSGNSSGSGGSPVPSTPPTPSPSTPPTPSPSAS	NCL79	368
GGGGSAS	NCL80	369
GGGGSGGGGSAS	NCL81	370
GGGGSGGGGSGGGGSAS	NCL82	371
GGGGSGGGGSGGGGSGGGGSGGGGSAS	NCL83	372
GGGGSGGGGSGGGGSGGGGSGGGGSGGGGSAS	NCL84	373
AGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSAS	NCL85	374
AGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSAS	NCL86	375
AGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSAS	NCL87	376

[0164] In certain aspects, the IFN receptor agonist of the disclosure may comprise a polypeptide chain comprising, in an N- to C-terminal orientation, a targeting moiety (or targeting moiety chain), a hinge domain, and an Fc domain. Thus, the hinge domain can be said to constitute a type of linker. Exemplary hinge domains are set forth in Section 6.9.3.

6.7. Targeting Moiety

[0165] The incorporation of targeting moieties in the IFN receptor agonists of the disclosure permits the delivery of high concentrations of IFN into the tumor microenvironment with a

concomitant reduction of systemic exposure, resulting in fewer side effects than obtained with untargeted IFN molecules.

[0166] It is anticipated that any type of target molecule present or capable of driving the IFN receptor agonist at a particular locale or tissue may be targeted by the IFN receptor agonists of the disclosure. In some embodiments, the IFN receptor agonists are intended to treat cancer, *e.g.*, by inducing a local immune response against tumor tissue. Accordingly, the targeting molecule can be any local tumor and associated target molecule. The target molecules recognized by the targeting moieties of the IFN receptor agonists of the disclosure are generally found, for example, on the surfaces of activated T cells, on the surfaces of tumor cells, on the surfaces of dendritic or other antigen-presenting cells, on the surfaces of natural killer (NK) cells, on the surfaces of virus-infected cells, on the surfaces of other diseased cells, free in blood serum, in the extracellular matrix (ECM), or immune cells present in the target site, *e.g.*, tumor reactive lymphocytes, dendritic cells or other antigen presenting cells, or natural killer cells.

[0167] In various embodiments, the target molecule is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen. The skilled artisan would recognize that the foregoing categories of target molecules are not mutually exclusive and thus a given target molecule may fall into more than one of the foregoing categories of target molecules. For example, some molecules may be considered both TAAs and ECM proteins, and other molecules may be considered both TCAs and checkpoint inhibitors.

[0168] Exemplary types of cancers that may be targeted include acute lymphoblastic leukemia, acute myelogenous leukemia, biliary cancer, B-cell leukemia, B-cell lymphoma, biliary cancer, bone cancer, brain cancer, breast cancer, triple-negative breast cancer, cervical cancer, Burkitt lymphoma, chronic lymphocytic leukemia, chronic myelogenous leukemia, colorectal cancer, endometrial cancer, esophageal cancer, gall bladder cancer, gastric cancer, gastrointestinal tract cancer, glioma, hairy cell leukemia, head and neck cancer, Hodgkin's lymphoma, liver cancer, lung cancer, medullary thyroid cancer, melanoma, multiple myeloma, ovarian cancer, non-Hodgkin's lymphoma, pancreatic cancer,

prostate cancer, pulmonary tract cancer, renal cancer, sarcoma, skin cancer, testicular cancer, urothelial cancer, and other urinary bladder cancers. However, the skilled artisan will realize that TAAs and other target molecules associated with the tumor microenvironment are known for virtually any type of cancer.

[0169] Non-limiting examples of ECM antigens include syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (*e.g.*, nectin-4), tenascin, collagen (*e.g.*, collagen type X) and matrixin.

[0170] Other target molecules are cell surface molecules of tumor or viral lymphocytes, for example T-cell co-stimulatory proteins such as CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, and B7-H3.

[0171] In particular embodiments, the target molecules are checkpoint inhibitors, for example CTLA-4, PD1, PDL1, PDL2, B7-H3, B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK1, CHK2. In particular embodiments, the target molecule is PD1. In other embodiments, the target molecule is LAG3. In yet other embodiments the target molecule is PDL1.

[0172] In certain embodiments, the target molecules are on the surfaces of dendritic cells or other antigen-presenting cells, such as XCR1, Clec9a, CD1c, CD11c, CD14, PDL1, macrophage mannose receptor (CD206), and DEC-205.

[0173] In further embodiments, the target molecules are on the surfaces of natural killer (NK) cells such as CD335, CD38, CD2, NKG2D, NKp44, NKp30, CD16, LFA-1, CD27, KIR, NKH1A, and NKp46.

[0174] The antibodies and antigen-binding portions generally bind to specific antigenic determinants and are able to direct the IFN receptor agonist to a target site, for example to a specific type of tumor cell or tumor stroma that bears the antigenic determinant. In particular embodiments, the targeting moiety recognizes a tumor-associated antigen (TAA). Preferably, the TAA is a human TAA. The antigen may or may not be present on normal cells. In certain embodiments, the TAA is preferentially expressed or upregulated on tumor cells as compared to normal cells. In other embodiments, the TAA is a lineage marker. Exemplary TAAs include Fibroblast Activation Protein (FAP), the A1 domain of Tenascin-C

(TNC A1), the A2 domain of Tenascin-C (TNC A2), the Extra Domain B of Fibronectin (EDB), the Melanoma-associated Chondroitin Sulfate Proteoglycan (MCSP), MART-1/Melan-A, gp100, Dipeptidyl peptidase IV (DPPIV), adenosine deaminase-binding protein (ADAbp), cyclophilin b, colorectal associated antigen (CRC)-C017-1A/GA733, Carcinoembryonic Antigen (CEA) and its immunogenic epitopes CAP-1 and CAP-2, etv6, aml1, Prostate Specific Antigen (PSA) and its immunogenic epitopes PSA-1, PSA-2, and PSA-3, prostate-specific membrane antigen (PSMA), T-cell receptor/CD3-zeta chain, MAGE-family of tumor antigens (*e.g.*, MAGE-A1, MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A5, MAGE-A6, MAGE-A7, MAGE-A8, MAGE-A9, MAGE-A10, MAGE-A11, MAGE-A12, MAGE-Xp2 (MAGE-B2), MAGE-Xp3 (MAGE-B3), MAGE-Xp4 (MAGE-B4), MAGE-C1, MAGE-C2, MAGE-C3, MAGE-C4, MAGE-C5), GAGE-family of tumor antigens (*e.g.*, GAGE-1, GAGE-2, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE-7, GAGE-8, GAGE-9), BAGE, RAGE, LAGE-1, NAG, GnT-V, MUM-1, CDK4, tyrosinase, p53, MUC family, HER2/neu, p21ras, RCAS1, α-fetoprotein, E-cadherin, α-catenin, β-catenin and γ-catenin, p120ctn, gp100 Pmel117, PRAME, NY-ESO-1, cdc27, adenomatous polyposis coli protein (APC), fodrin, Connexin 37, Ig-idiotype, p15, gp75, GM2 and GD2 gangliosides, viral products such as human papilloma virus proteins, Smad family of tumor antigens, Imp-1, P1A, EBV-encoded nuclear antigen (EBNA)-1, brain glycogen phosphorylase, SSX-1, SSX-2 (HOM-MEL-40), SSX-1, SSX-4, SSX-5, SCP-1 and CT-7, c-erbB-2, Her2, EGFR, IGF-1R, CD2 (T-cell surface antigen), CD3 (heteromultimer associated with the TCR), CD22 (B-cell receptor), CD23 (low affinity IgE receptor), CD30 (cytokine receptor), CD33 (myeloid cell surface antigen), CD40 (tumor necrosis factor receptor), IL-6R-(IL6 receptor), CD20, MCSP, PDGFβR (β-platelet-derived growth factor receptor), ErbB2 epithelial cell adhesion molecule (EpCAM), EGFR variant III (EGFRvIII), CD19, disialoganglioside GD2, ductal-epithelial mucine, gp36, TAG-72, glioma-associated antigen, β-human chorionic gonadotropin, alphafetoprotein (AFP), lectin-reactive AFP, thyroglobulin, MN-CA IX, human telomerase reverse transcriptase, RU1, RU2 (AS), intestinal carboxyl esterase, mut hsp70-2, M-CSF, prostase, prostase specific antigen (PSA), PAP, LAGA-1a, p53, prostein, PSMA, surviving and telomerase, prostate-carcinoma tumor antigen-1 (PCTA-1), ELF2M, neutrophil elastase, ephrin B2, insulin growth factor (IGF1)-I, IGF-II, IGFI receptor, 5T4, ROR1, Nkp30, NKG2D, tumor stromal antigens, the extra domain A (EDA) and extra domain B (EDB) of fibronectin and the A1 domain of tenascin-C(TnC A1).

[0175] Suitable targeting moiety formats are described in Section 6.8. The targeting moiety is preferably an antigen binding moiety, for example an antibody or an antigen-binding portion of an antibody, *e.g.*, an scFv, as described in Section 6.8.2 or a Fab, as described in Section 6.8.1.

[0176] In some embodiments, the targeting moieties target the exemplary target molecules set forth in Table F below, together with references to exemplary antibodies or antibody sequences upon which the targeting moiety can be based.

TABLE F
Exemplary Target Molecules

Target	**Antibody Name and/or Binding Sequences**
1-92-LFA-3	Amevive™ (alefacept)
5T4	GEN1044
Activin Receptor Type II	Bimagrumab VH: SEQ ID Nos:107, 109 of US Patent No. 8,388,968 B2 VL: SEQ ID Nos:93, 95 of US Patent No. 8,388,968 B2
B7-H3	Obrindatamab (MGD009)
B7-H3 (CD276)	Enoblituzumab (MGA271)
B7-H3 (CD276)	MGC018
B7-H3 (CD276)	MGA012
B7-H3 (CD276)	8H9
B7-H3 (CD276)	VH: the VH sequence of the heavy chain of SEQ ID NO:21, 26 or 31 of US 2021/0171641 A1. VL: the VL sequence of the light chain of SEQ ID NO:20, 22 or 30 of US 2021/0171641 A1.
B7-H3 (CD276)	VH: the VH sequence of the heavy chain of SEQ ID NO:21, 29 or 37 of US 2019/0002563 A1. VL: the VL sequence of the light chain of SEQ ID NO:17, 25 or 33 of US 2019/0002563 A1.
B7-H3 (CD276)	VH: the VH sequence of the heavy chain of SEQ ID NO:146, 147 or 148 of US Patent No. 10,640,563. VL: the VL sequence of the light chain of SEQ ID NO:143, 144 or 145 of US Patent No. 10,640,563.
BAFF/B Lymphocyte Stimulator	Benlysta™ (velimumab)
BAFF/B Lymphocyte Stimulator	VH: amino acids 1-123 of SEQ ID NO:327 of US Patent No. 7,138,501 VL: amino acids 139-249 of SEQ ID NO:327 of US Patent No. 7,138,501.
BAFF/B Lymphocyte Stimulator	VH: amino acids 1-126 of SEQ ID NO:1321 of US Patent No. 7,605,236;

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	VL: amino acids 143-251 of SEQ ID NO:1049 of US Patent No. 7,605,236.
BAFF/B Lymphocyte Stimulator	Belimumab
BCMA	VH: the VH sequence of the heavy chain of SEQ ID NO. 126 of US 2021/0206865 A1 VL: the VL sequence of the light chain of SEQ ID NO. 129 or SEQ ID NO. 132 of US 2021/0206865 A1
CA125	Igobumab
CA125	OvaRex™ (oregobumab)
Cadherin	The antibodies described in US Pub. No. US 2006/0039915.
N-cadherin	An antibody that binds to the amino acid sequence of SEQ ID NO: 10, 17 or 18 of US Pub. No. US 2010/0278821.
CD11a	Raptiva™ (efalizumab) Sequence in Werther *et al.*, 1996, The Journal of Immunology 157(11): 4986-4995.
CD19	Blincyto™ (blinatumomab)
CD19	SGN-CD19A
CD20	Bexxar™ (tositumomab) VH: the VH sequence of the heavy chain of SEQ ID NO:124 of US Patent Pub. US 2017/0002060 A1 VL: the VL sequence of the light chain of SEQ ID NO:125 of US Patent Pub. US 2017/0002060 A1
CD20	Zevalin™ (ibritumomab tiuxetan) VH: SEQ ID NO:9 of US Patent No. 5,736,137 VL: SEQ ID NO:6 of US Patent No. 5,736,137
CD20	Rituxan™ (rituximab) VH: SEQ ID NO:9 of US Patent No. 5,736,137 VL: SEQ ID NO:6 of US Patent No. 5,736,137
CD20	Ocrevus™ (ocrelizumab)
CD20	Okaratuzumab
CD20	Arzerra™ (ofatumumab) VH: SEQ ID NO:2 of US Patent No. 8,529,902 VL: SEQ ID NO:4 of US Patent No. 8,529,902
CD20	Gazyva™ (obinutuzumab)
CD20	VH: SEQ ID NO: 4 of US 2021/0206870 A1 VL of SEQ ID NO:6 of US 2021/0206870 A1
CD20	Epcoritamab

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
CD22	Belimumab
CD22	Epratuzumab
CD22	Besponsa ™ (inotuzumab ozogamicin)
CD22	Lumoxiti ™ (moxetumumab pasudox)
CD22	pinatuzumab vedotin
CD25	Zenapax™ (daclizumab) VH: SEQ ID NO:9 of US Patent No. 7,060,269 VL: SEQ ID NO:10 of US Patent No. 7,060,269
CD30	Adcetris™ (brentuximab vedotin) VH: SEQ ID NO:2 of US Patent No. 7,090,843 VL: SEQ ID NO:10 of US Patent No. 7,090,843
CD33	Myelotarg™ (gemtuzumab) Sequence in Man Sung, *et al.*, 1993, Molecular immunology 30:1361-1367
CD33	Lintuzumab
CD38	Darzalex™ (daratumumab)
CD38	IB4, HB7 CS/2, clone 90 and NIM-R5 as disclosed in PCT Pub. WO2015/009726A2 and references cited therein.
CD40	Lukatumumab
CD40	Dacetuzumab
CD40L	Hu5c8 (ruplizumab)
CD44v6	vibatuzumab mertansine
CD52	Campath™ (alemtuzumab) VH: SEQ ID NO:1 of US Patent Pub. US 2017/0002060 A1 VL: SEQ ID NO:2 of US Patent Pub. US 2017/0002060 A1
CD70	Blenrep ™ (borsetuzumab mafodotin)
CD123	Flotetuzumab
CD206	Anti-CD206 antibodies having a VH a of SEQ ID NO. 2 and a VL of SEQ ID NO:4 of WO2003/040169A2
CD221	Tepezza ™ (teprotumumab)
CEA	Hybri-Ceaker® (altumomab pentetate)
CEA	Scintimun™ (besilesomab)
CEA	CEA-CIDE™ (labetuzumab))
CEA	CEA-Scan ™ (arcitumomab)
CEA	hMN-15 CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID Nos:4-6 of US Patent No. 8,771,690 B2

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID Nos:1-3 of US Patent No. 8,771,690 B2
CEA	CEA binding portion of RO6958688/RG7802 from clinical trial NCT02324257
CEA	Cibisatamab
CEA	CEA binding portion of MEDI-565/MT110/AMG211 from clinical trials NCT01284231 and NCT02291614 VH: SEQ ID NO:49 or 51 of PCT Publication No. WO 2013/012414 A1 VL: SEQ ID NO:48 of PCT Publication No. WO 2013/012414 A1.
CEA	Rabetuzumab
CEA	Atezolizumab
CEA	Cibisatamab
CEA	MEDI-565 (AMG211, MT111)
CEA	RO6958688
CEA	VH: SEQ ID No. 9 described in WO2022/048883A1 VL: SEQ ID No. 10 described in WO2022/048883A1
CLDN18.2	AMG910
Clec9a	Anti-Clec9a antibodies having VH and VL amino acids of SEQ ID Nos. 43 and 48 of PCT Pub. No. WO2009/026660A1
Clec9a	Anti-Clec9a antibodies having a VH a of SEQ ID NO. 38 and a VL of SEQ ID NO:37 or SEQ ID NO:43 of PCT Pub. No. WO2022/073062A1 Anti-Clec9a antibodies having a VH a of SEQ ID NO. 8 and a VL of SEQ ID NO:7 of PCT Pub. No. WO2022/073062A1
Collagen alpha-4 chain	TRC093 (MT293)
Collagen	The collagen binding antibody fragment described in Liang *et al.*, 2016, Sci. Rep. 5, 18205; doi: 10.1038/srep18205 (2016).
Collagen Type I	Cetuximab (Erbitux)
Collagen type X	The amino acid sequences of SEQ ID NO: 1 or 2 of PCT Pub No. WO 2019/020797.
Collagen type X	The amino acid sequences of SEQ ID NO: 1 of PCT Pub No. WO 2014/180992.
Collagen type X	Antibody X34 as described in I. Girkontaite et al., "Immunolocalization of type X collagen in normal fetal and

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	adult osteoarthritic cartilage with monoclonal antibodies," Matrix Biol 15, 231-238 (1996).
Collagen type X	Antibodies X53 or 1H8 or ARC0659 or JF0961 collagen X polyclonal antibody sold under catalog number PA5-115039 or PA5-116871 or PA5-97603 or PA5-49198 from ThermoFisher Scientific.
Collagen type X	Antibody sold under catalog number RDI-COLL10abr from RDI.
Complement C5	Soliris™ (eculizumab) VH: amino acids 1-122 of SEQ ID NO: 10 of US Patent No. 6,355,245 VL: amino acids 3-110 of SEQ ID NO: 9 of US Patent No. 6,355,245
CTLA-4	Yervoy™ (ipilimumab) VH: SEQ ID NO: 17 of WO 2001/014424 A2 VL: SEQ ID NO: 7 of WO 2001/014424 A2
CTLA-4	(tremelimumab)
CTLA-4	Orencia™ (abatacept)
DEC-205	Anti-DEC-205 antibodies having a VH/VL pair of SEQ ID NOS. 4/10, 16/22, 28/34, 40/46, 52/58, and 76/82 of PCT Pub. WO2009/061996A2
DLL3	AMG757
EGFR	Erbitux™ (cetuximab) VH: SEQ ID NO: 11 of US Patent No. 6,217,866 VL: SEQ ID NO: 13 of US Patent No. 6,217,866
EGFR	Vectibix™ (panitumumab) VH: SEQ ID NO: 37 of US Patent No. 6,235,883 VL: SEQ ID NO: 38 of US Patent No. 6,235,883
EGFR	Zalutumumab VH: SEQ ID NO: 64 of WO 2018/140831 A2 VL: SEQ ID NO: 69 of WO 2018/140831 A2
EGFR	Mapatumumab
EGFR	Matuzumab
EGFR	Nimotuzumab VH: SEQ ID NO: 51 of WO 2018/140831 A2 VL: SEQ ID NO: 56 of WO 2018/140831 A2
EGFR	ICR62
EGFR	mAb 528
EGFR	CH806
EGFRv3	AMG596
EGFRv3	AMG404

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
EGFR/CD64	MDX-447
EpCAM	Panorex ™ (edrecolomab) VH: SEQ ID NO:129 of WO 2018/140831 A2 VL: SEQ ID NO:134 of WO 2018/140831 A2
EpCAM	Adecatumumab VH: SEQ ID NO:142 of WO 2018/140831 A2 VL: SEQ ID NO:147 of WO 2018/140831 A2
EpCAM	tucotuzumab celmoleukin
EpCAM	citatuzumab bogatox
EpCAM	EP1629013 B1 VH: SEQ ID Nos: 80, 84, 88, 92 or 96 VL: SEQ ID Nos: 82, 86, 90, 94 or 98
EpCAM	G8.8 HC: SEQ ID NO: 4 of US Patent Pub. No. US 2020/0317806 A1 HL: SEQ ID NO: 3 of US Patent Pub. No. US 2020/0317806 A1
EpCAM	VH: SEQ ID Nos: 17-22 of WO 2021/211510 A2. VL: SEQ ID NO: 15-16 of WO 2021/211510 A2.
EpCAM	Removab ™(catumaxomab)
EpCAM	Vicineum ™ (oportuzumab monatox)
EpCAM	M701
F protein of RSV	Synagic™ (palivizumab)
GD2	3F8
Glycoprotein receptor IIb/IIIa	ReoPro™ (abiciximab)
gpA33	MGD007
GPC3	ERY974
GUCY2C	PF-07062119
Heparanase	An antibody selected from HP130, HP 239, HP 108.264, HP 115.140, HP 152.197, HP 110.662, HP 144.141, HP 108.371, HP 135.108, HP 151.316, HP 117.372, HP 37/33, HP3/17, HP 201 or HP 102 or an amino acid sequence of SEQ ID NO: 1-11 described in US Patent Pub. US 2004/0170631.
Her2	Herceptin™ (trastuzumab)
Her2	Aldesleukin (proleukine)
Her2	Sargramustim (Leucine)
Her2	M802
Her2	Runimotamab (BTRC4017A, R07227780)
Her2	ISB1302
Her2-neu	Perjeta™ (pertuzumab)

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	VH: SEQ ID NO:16 of WO 2013/096812 A1. VL: SEQ ID NO:15 of WO 2013/096812 A1.
Her2-neu	Rexomun™ (ertumaxomab)
IgE	Xolair™ (omalizumab)
IGFIR	(figitumumab)
IL1β	Ilaris™ (canakinumab) VH: SEQ ID NO:1 of US Patent No. 7,446,175. VL: SEQ ID NO: 2 of US Patent No. 7,446,175
IL12/IFN3	Stelara™ (ustekinumab)
IL1Ra	Antril™, Kineret™ (ankinra)
IFNR	Simulect™ (basiliximab) VH: SEQ ID NO: 3 of US Patent No. 6,383,487 VL: SEQ ID NO: 6 of US Patent No. 6,383,487
IL6	Clazakizumab
IL6 receptor	Actemra™ (tocilizumab) VH: SEQ ID NO: 31 of US Patent No. 7,479,543 VL: SEQ ID NO: 29 of US Patent No. 7,479,543
IL12/IFN3 p40 subunit	Stelara™ (ustekinumab) VH: SEQ ID NO: 7 of US Patent No. 6,902,734 VL: SEQ ID NO: 8 of US Patent No. 6,902,734
Integrin α4	Tysabri™ (natalizumab) VH: SEQ ID Nos: 11-13 of US Patent No. 5,840,299 VL: SEQ ID Nos: 7-8 of US Patent No. 5,840,299
Integrin α4 β7	Entyvio ™ (vedolizumab) HC: SEQ ID NO: 2 of US Patent Pub. US 2012/0282249. LC: SEQ ID NO: 4 of US Patent Pub. US 2012/0282249.
Integrin α5 β1	VH: SEQ ID NO: 2 of European Patent No. 1 755 659. VL: SEQ ID NO: 4 of European Patent No. 1 755 659.
Integrin β1	VH: SEQ ID NO:2, 6, 8, 10, 12, 14, 29-43 or 91-100 of US Patent Pub. US 2022/0089744. VL: SEQ ID NO:4, 16, 18, 20, 22, 44-57 or 107-116 of US Patent Pub. US 2022/0089744.
KIR	Anti-KIR antibodies having a VH of SEQ ID NO: 5 and a VL of SEQ ID NO: 3 of PCT Pub. WO2014/066532A1
KIR	Anti-KIR antibodies having a VH of SEQ ID NO: 1 and a VL of SEQ ID NO: 2 of PCT Pub. WO2012/160448A2 Anti-KIR antibodies having a VH of SEQ ID NO: 3 and a VL of SEQ ID NO: 4 of PCT Pub. WO2012/160448A2
LAG3	Relatlimab (BMS-98016)
LAG3	Sym022

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
LAG3	HLX26
LAG3	TSR-033
LAG3	ABL501
LAG3	INCAGN02385
LAG3	Fianlimab (REGN3767)
LAG3	RO7247669
LAG3	EMB-02
LAG3	FS118
LAG3	GSK2831781
LAG3	IBI323
LAG3	IBI110
LAG3	LAG525
LAG3	XmAb®22841
LAG3	LBL-007
LAG3	VH: SEQ ID NO:1, 8, 10 or 12 of US Patent No. 9,902,772. VL: SEQ ID NO:2, 3, 4, 5, 6, 7, 9, 11, 13 or 14 of US Patent No. 9,902,772.
LAG3	VH: SEQ ID NO: 182 of US Patent Pub. US 2021/0095026. VL: SEQ ID NO: 88 of US Patent Pub. US 2021/0095026.
LAG3	Antibodies having VH/VL amino acid sequences of SEQ ID Nos 23/24, ¾ and 11/12 of US Pub. US2022/0056126A1.
Laminin	Lam-89 from Sigma Aldrich
Mesothelin	Amatuximab
Mesothelin	HPN536
MUC1	civatuzumab tetraxetane
MUC1	Pankomab ™ (gatipotuzumab)
MUC1	Femtumumab
MUC1	Cantuzumab ravtansine
MUC16 (CA125)	Anti-MUC16 antibodies having VH and VL sequences having the amino acid sequences of any one of the following SEQ ID NO: pairs from US 2018/0118848A1: 18/26; 82/858; 98/170
MUC17	AMG199
Nectin-4	Enfortumab (ASP7465, ASG-22CE, ASG-22ME) VH: SEQ ID NO:3 of PCT Pub. WO 2021/151984. VL: SEQ ID NO: 4 of PCT Pub. WO 2021/151984.
Nectin-4	SBT290
Nectin-4	VH: SEQ ID NO:1 of US Patent No. 11,274,160. VL: SEQ ID NO:2 of US Patent No. 11,274,160.
NGF	(tanezumab)

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
NKH1A	The monoclonal antibody deposited with ATCC and assigned accession no. HB8564, as described in US Patent No. 4,772,552A
NKP46	Anti-NKP46 antibodies having CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID Nos:4, 6 and 8 and CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID Nos:12, 14 and 16 of PCT Pub. WO2018/047154A1
Osteopontin	HC: SEQ ID NO: 22 of PCT Pub. WO 2021/030209. LC: SEQ ID NO: 24 of PCT Pub. WO 2021/030209.
PD1	MDX-1106 / BMS-936558 (nivolumab), a human IgG4 mAb with the structure described in WHO Drug Information, Vol. 27, No. 1, pages 68-69 (2013) and whose heavy and light chain sequences are disclosed in FIG. 7 of US Pub. No. US20190270812A1 HC: SEQ ID NO: 23 of US Pub. No. US20190270812A1 LC: SEQ ID NO: 24 of US Pub. No. US20190270812A1
PD1	MK-3475 (pembrolizumab), a humanized IgG4 mAb with the structure described in WHO Drug Information, Vol. 27, No. 2, pages 161-162 (2013) and whose heavy and light chain sequences are disclosed in FIG. 6 of US Pub. No. US20190270812A1 HC: SEQ ID NO: 21 of US Pub. No. US20190270812A1 LC: SEQ ID NO: 22 of US Pub. No. US20190270812A1
PD1	REGN2810 (disclosed as H4H7798N in US Pub No. 20150203579) HC: SEQ ID NO: 330 of US Pub. No. 20150203579 LC: SEQ ID NO: 331 of US Pub. No. 20150203579
PD1	Anti-PD1 antibodies having CDR H1-H3 and CDR L1-L3 sequences corresponding to the following SEQ ID Nos. of US Patent No. 11,034,765 B2: a) SEQ ID Nos: 18, 19, 20, 21, 22, and 23, respectively; b) SEQ ID Nos: 24, 25, 26, 27, 28, and 29, respectively; c) SEQ ID Nos: 30, 31, 32, 33, 34, and 35, respectively; d) SEQ ID Nos: 36, 37, 38, 39, 40, and 41, respectively; e) SEQ ID Nos: 42, 43, 44, 45, 46, and 47, respectively; f) SEQ ID Nos: 48, 49, 50, 51, 52, and 53, respectively; g) SEQ ID Nos: 54, 55, 56, 57, 58, and 59, respectively; and h) SEQ ID Nos: 60, 61, 62, 63, 64, and 65, respectively.
PD1	Anti-PD1 antibodies disclosed in Tables 1-3 of PCT Pub. WO2015112800A1, including but not limited to anti-PD1 antibodies having VH/VL pairs having SEQ ID Nos: 2/10, 18/26, 34/42, 50/58, 66/74, 82/90, 98/106, 1 14/122,

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	130/138, 146/154, 162/170, 178/186, 194/202, 210/202, 218/202, 226/202, 234/202, 242/202, 250/202, 258/202, 266/202, 274/202, 282/202, 290/202, 298/186, 306/186 and 314/186 of PCT Pub. WO2015112800A1.
PD1	Anti-PD1 antibodies disclosed in US Patent No. 10,294,299 B2 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 164/178 SEQ ID Nos. 165/179 SEQ ID Nos. 166/180 SEQ ID Nos. 167/181 SEQ ID Nos. 168/182 SEQ ID Nos. 169/183 SEQ ID Nos. 170/184 SEQ ID Nos. 171/185 SEQ ID Nos. 172/186 SEQ ID Nos. 173/187 SEQ ID Nos. 174/188 SEQ ID Nos. 175/189 SEQ ID Nos. 176/190 SEQ ID Nos. 177/190
PD1	MEDI-0680 (AMP-514)
PD1	PDR001 (spartalizumab), a humanized IgG4 mAb whose heavy and light chain sequences are disclosed as BAP049-Clone-E in US Patent No: 9683048 B2. HC: SEQ ID NO: 91 of US Patent No: 9,683,048 LC: SEQ ID NO: 72 of US Patent No: 9,683,048
PD1	BGB-108
PD1	h409A11, described in WO2008/156712 HC: SEQ ID NO: 31 of PCT Pub. WO2008/156712 LC: SEQ ID NO: 36 of PCT Pub. WO2008/156712
PD1	h409A16, described in WO2008/156712 HC: SEQ ID NO: 31 of PCT Pub. WO2008/156712 LC: SEQ ID NO: 37 of PCT Pub. WO2008/156712
PD1	h409A17, described in WO2008/156712 HC: SEQ ID NO: 31 of PCT Pub. WO2008/156712 LC: SEQ ID NO: 38 of PCT Pub. WO2008/156712
PD1	Anti-PD1 antibodies described in US Patent No. 7,488,802 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 2/4 SEQ ID Nos. 6/8 SEQ ID Nos. 10/12

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	SEQ ID Nos. 14/16 SEQ ID Nos. 47/49
PD1	Anti-PD1 antibodies described in US Patent No. 7,521,051 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 2/4 SEQ ID Nos. 6/8 SEQ ID Nos. 10/12 SEQ ID Nos. 14/16 SEQ ID Nos. 47/49
PD1	Anti-PD1 antibodies described in US Patent No. 8,008,449 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 1/8 SEQ ID Nos. 2/9 SEQ ID Nos. 3/10 SEQ ID Nos. 4/11 SEQ ID Nos. 5/12 SEQ ID Nos. 6/13 SEQ ID Nos. 7/14
PD1	Anti-PD1 antibodies described in US Patent No. 8,354,509 as having the following SEQ ID NO. pairs for heavy and light chain full-length domains: SEQ ID Nos. 31/36 SEQ ID Nos. 31/37 SEQ ID Nos. 31/38
PD1	Anti-PD1 antibodies described in US Patent No. 8,168,757 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 4/5 SEQ ID Nos. 12/13 SEQ ID Nos. 18/19 SEQ ID Nos.40/41 SEQ ID Nos.47/48 SEQ ID Nos.26/27 SEQ ID Nos.34/35 SEQ ID Nos.55/56 SEQ ID Nos.67/68
PD1	Anti-PD1 antibodies described in PCT Pub. No. WO2004/004771
PD1	Anti-PD1 antibodies described in PCT Pub. No. WO2004/056875 as having the following SEQ ID NO. pairs for heavy and light chain variable domains:

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	SEQ ID Nos. 2/4 SEQ ID Nos. 6/8 SEQ ID Nos. 10/12 SEQ ID Nos. 14/16 SEQ ID Nos. 47/49
PD1	Anti-PD1 antibodies described in PCT Pub. No. WO2004/072286
PD1	VH: SEQ ID NO: 25, 26, 27, 28, or 29 of US Pub. No. US2011/0271358 VL: SEQ ID NO: 30, 31, 32, or 33 of US Pub. No. US2011/0271358
PD1	SHR-1210 (Camrelizumab) described in PCT Publication No: WO 2015/085847 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 9 LC: SEQ ID NO: 10
PDL1	Durvalumab (MEDI4736) HC: SEQ ID NO: 26 of PCT application No. WO2020225552 LC: SEQ ID NO: 27 of PCT application No. WO2020225552
PDL1	Atezolizumab (Tecentriq, MPDL3280A, RG7446) HC: SEQ ID NO: 20 of US Patent No. 8,217,149 LC: SEQ ID NO: 21 of US Patent No. 8,217,149
PDL1	MDX 1105 (BMS-936559)
PDL1	Anti-PDL1 antibodies described in U.S. Patent No. 7,943,743 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos: 1/11 SEQ ID Nos: 2/12 SEQ ID Nos: 3/13 SEQ ID Nos: 4/14 SEQ ID Nos: 5/15 SEQ ID Nos: 6/16 SEQ ID Nos: 7/17 SEQ ID Nos: 8/18 SEQ ID Nos: 9/19 SEQ ID Nos: 10/20
PDL1	Avelumab, described in US Patent No: 9,624,298 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 24 LC: SEQ ID NO: 25

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
PDL1	ZKAB001 (Socazolimab)
PDL1	TQB2450 (APL-502 or CBT-502)
PDL1	HLX20 CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID Nos: 52, 56, and 77 of PCT Pub. No. 2018/080812 CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID Nos: 65, 42, and 71 of PCT Pub. No. 2018/080812
PDL1	KN035 (Envafolimab) is a nanobody described as Hu56V2 in US Patent No. 11,225,522 as having the VHH SEQ ID NO: 34
PDL1	LY3434172
PDL1	LY3300054 (Iodapolimab) described in PCT Pub No: WO 2017/034916 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 10 L: SEQ ID NO: 11
PDL1	LDP (Iesabelimab, ADG104) described in CN Patent No: 114225023 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 10 LC: SEQ ID NO: 9
PDL1	EMB-09
PDL1	ABL501
PDL1	INBRX-105
PDL1	STI-3031 (IMC-001) described in US Patent No: 10,118,963 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 1 LC: SEQ ID NO: 2
PDL1	BGB-A333 (garivulimab) described in US Patent No: 11,512,132 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 22 LC: SEQ ID NO: 23
PDL1	HLX301
PDL1	Y101D
PDL1	ES101
PDL1	IBI322
PDL1	VH: SEQ ID NO:46, 48, 50 or 52 of US Patent No. 11,168,144.

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	VL: SEQ ID NO:58, 137 or 12 of US Patent No. 11, 168,144.
PDL1	VH: SEQ ID NO:23, 124, 126, 127, 128, 130, 140 or 145 of US Patent No. 11,208,486. VL: SEQ ID NO:24 or 125 of US Patent No. 11,208,486.
Phosphatidylserine	(bavituximab)
PSCA	GEM3PSCA
PSMA	huJ591
PSMA	Anti-PSMA antibodies having VH and VL sequences having the amino acid sequences of any one of the following SEQ ID NO: pairs from WO 2017/023761A1: 2/1642; 10/1642; 18/1642; 26/1642; 34/1642; 42/1642; 50/1642; 58/1642; 66/1642; 74/1642; 82/1642; 90/1642; 98/1642; 106/1642; 1 14/1642; 122/130; and 138/146.
PSMA	An antibody such as: PSMA 3.7, PSMA 3.8, PSMA 3.9, PSMA 3.11, PSMA 5.4, PSMA 7.1, PSMA 7.3, PSMA 10.3, PSMA 1.8.3, PSMA A3.1.3, PSMA A3.3.1, Abgenix 4.248.2, Abgenix 4.360.3, Abgenix 4.7.1, Abgenix 4.4.1, Abgenix 4.177.3, Abgenix 4.16.1, Abgenix 4.22.3, Abgenix 4.28.3, Abgenix 4.40.2, Abgenix 4.48.3, Abgenix 4.49.1, Abgenix 4.209.3, Abgemx 4.219.3, Abgenix 4.288.1, Abgenix 4.333.1, Abgemx 4.54.1, Abgenix 4.153.1, Abgenix 4.232.3, Abgenix 4.292.3, Abgenix 4.304.1, Abgenix 4.78.1 and Abgenix 4.152.1 described in WO2003034903A2 A hybridoma cell line such as: PSMA 3.7 (PTA-3257), PSMA 3.8, PSMA 3.9 (PTA- 3258), PSMA 3.11 (PTA-3269), PSMA 5.4 (PTA-3268), PSMA 7.1 (PTA-3292), PSMA 7.3 (PTA-3293), PSMA 10.3 (PTA-3247) , PSMA 1.8.3 (PTA-3906), PSMA A3.1.3 (PTA- 3904), PSMA A3.3.1 (PTA-3905), Abgenix 4.248.2 (PTA-4427), Abgenix 4.360.3 (PTA- 4428), Abgenix 4.7.1 (PTA-4429), Abgenix 4.4.1 (PTA-4556), Abgenix 4.177.3 (PTA-4557), Abgenix 4.16.1 (PTA-4357), Abgenix 4.22.3 (PTA-4358), Abgenix 4.28.3 (PTA-4359), Abgenix 4.40.2 (PTA-4360), Abgenix 4.48.3 (PTA-4361), Abgenix 4.49.1 (PTA-4362), Abgenix 4.209.3 (PTA-4365), Abgenix 4.219.3 (PTA-4366), Abgenix 4.288.1 (PTA-4367), Abgenix 4.333.1 (PTA-4368), Abgenix 4.54.1 (PTA-4363), Abgenix 4.153.1 (PTA-4388), Abgenix 4.232.3 (PTA-4389), Abgenix 4.292.3 (PTA-4390), Abgenix 4.304.1 (PTA-4391),

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	Abgenix 4.78.1 (PTA-4652), and Abgemx 4.152.1(PTA-4653) described in WO 2003/034903A2. VH of SEQ ID Nos: 2-7 described in WO 2003/034903A2 VL of SEQ ID Nos: 8-13 described in WO 2003/034903A2
PMSA	VH: SEQ ID Nos: 225, 239, 253, 267, 281, 295, 309, 323, 337, 351, 365, 379, 393, 407, 421, 435, 449, 463, 477, 491, 505, 519, 533, 547, 561, 575, 589, 603 or 617 described in WO 2011/121110A1. VL SEQ ID Nos: 230, 244, 258, 272, 286, 300, 314, 328, 342, 356, 370, 384, 398, 412, 426, 440, 454, 468, 482, 496, 510, 524, 538, 552, 566, 580, 594, 608 or 622 described in WO 2011/121110A1. VH and VL SEQ ID Nos: 235, 249, 263, 277, 291, 305, 319, 333, 347, 361, 375, 389, 403, 417, 431, 445, 459, 473, 487, 501, 515, 529, 543, 557, 571, 585, 599, 613 or 627 described in WO 2011/121110A1.
PMSA	An anti-PMSA antibody having a VL amino acid sequence of any one of SEQ ID Nos: 229-312 of US 2022/0119525 A1 and a VH of SEQ ID NO: 217 of US 2022/0119525 A1.
PMSA	ES414
PMSA	BAY2010112 (pasotuxizumab)
PMSA	CCW702
PMSA	JNJ-63898081
PMSA	CC-1
PMSA	Acapatamab
PSMA	HPN424
RAAG12	RAV12
RANKL	Prolia™ (denosumab) VH: SEQ ID NO:51 of US Patent Pub. 2017/0002060 VL: SEQ ID NO:52 of US Patent Pub. 2017/0002060
SLAMF7	Empliciti™ (elotuzumab)
SSTR2	XmAb®18087
STEAP1	VHCDR1 SEQ ID Nos: 14, 33, 182, 184 or 185 described in US20210179731A1. VHCDR2 SEQ ID Nos: 15, 21, 34, 182, 184 or 185 described in US20210179731A1. VHCDR3 SEQ ID Nos: 16 and 35 described in US20210179731A1. VH SEQ ID Nos: 182 or 184 described in US20210179731A1.

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	VLCDR1 SEQ ID Nos: 11 or 30 described in US20210179731A1. VLCDR2 SEQ ID Nos: 12 or 31 described in US20210179731A1. VLCDR3 SEQ ID Nos: 13 or 32 described in US20210179731A1. VL SEQ ID Nos: 183 or 186 described in US20210179731A1.
STEAP1	AMG509
STEAP2	Anti-STEAP 2 antibodies having CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2 and CDR-L3 sequences selected from SEQ ID NOS: (1) 4-6-8-12-14-16; (2) 20-22-24-28-30-32; (3) 36-38-40-44-46-48; (4) 52-54-56-60-62-64; (5) 68-70-72-60-62-64; (6) 76-78-80-60-62-64; (7) 84-86-88-60-62-64; (8) 92-94-96-60-62-64; (9) 100-102-104-60-62-64; (10) 108-110-112-116-118-120; (11) 124-126-128-132-134-136; (12) 140-142-144-148-150-152; (13) 156-158-160-164-166-168; (14) 172-174-176-180-182-184; (15) 188-190-192-196-198-200; (16) 204-206-208-212-214-216; (17) 220-222-224-228-230-232; (18) 236-238-240-244-246-248; (19) 252-254-256-260-262-264; (20) 268-270-272-276-278-280; (21) 284-286-288-292-294-296; (22) 300-302-304-308-310-312; (23) 316-318-320-324-326-328; (24) 332-334-336-340-342-344; (25) 348-350-352-356-358-360; (26) 364-366-368-372-374-376; and (27) 380-382-384-388-390-392 of US Patent No. 10,772,972 B2. Anti-STEAP 2 antibodies having (a) a VH comprising the amino acid of any one of SEQ ID Nos: 2, 18, 34, 50, 66, 74, 82, 90, 98, 106, 122, 138, 154, 170, 186, 202, 218, 234, 250, 266, 282, 298, 314, 330, 346, 362, and 378 of US Patent No. 10,772,972 B2; and (b) a VL comprising the amino acid sequence of any one of SEQ ID Nos: 10; 26; 42; 58; 114; 130; 146; 162; 178; 194; 210; 226, 242; 258; 274; 290; 306; 322; 338; 354; 370; and 386 of US Patent No. 10,772,972 B2. Anti-STEAP 2 antibodies having a VH/VL pair comprising the amino acid sequences of any of the following pairs of SEQ ID Nos of US Patent No. 10,772,972 B2: 2/10; 18/26;

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	34/42; 50/58; 66/58; 74/58; 82/58; 90/58; 98/58; 106/114; 122/130; 138/146; 154/162; 170/178; 186/194; 202/210; 218/226; 234/242; 250/258; 266/274; 282/290; 298/306; 314/322; 330/338; 346/354; 362/370; and 378/386.
Syndecan-1 (CD 138)	The B-B4 antibody described in Wijdenes et al. (1996) Br. J. Haematol., 94: 318-323
Syndecan-4	The amino acid sequence of amino acids 93 and 121 of SEQ ID NO: 1 or the amino acid sequence of amino acids 92 and 122 of SEQ ID NO:2 described in European Patent Pub. EP 2 603 236.
TGFβ	GC1008
TNFR	Enbrel™ (etanercept)
TNFα	Remicade™ (infliximab) VH: SEQ ID NO:2 of Int. Patent Publication WO201/3087911 A1 VH: SEQ ID NO:3 of Int. Patent Publication WO2013/ A1087911
TNFα	Humira™ (adalimumab) VH: SEQ ID NO:4 of US Patent No. 6,258,562 VL: SEQ ID NO:3 of US Patent No. 6,258,562
TNFα	Cimzia ™ (certolizumab pegol) VH: SEQ ID NO:14 of US Patent No. 7,012,135 VL: SEQ ID NO:9 of US Patent No. 7,012,135
TNFα	Simponi ™ (golimumab) VH: SEQ ID NO:7 of US Patent No. 7,250,165 VL: SEQ ID NO:8 of US Patent No. 7,250,165
VEGF	Avastin™ (bevacizumab) VH: SEQ ID NO:9 of US Patent No. 7,060,269 VL: SEQ ID NO:10 of US Patent No. 7,060,269
VEGF	Lucentis™ (ranibizumab) VH: SEQ ID NO:4 of US Patent No. 9,914,770 VL: SEQ ID NO:2 of US Patent No. 9,914,770
XCR1	Anti-XCR1 antibodies disclosed in US Patent No. 9,371, 389 B2, including: - The antibodies designated 2H6, 5G7, 11H2, HK1L2 and HK5L5 - Antibodies having CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID Nos:53-55 and CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID Nos:56-58.

TABLE F Exemplary Target Molecules	
Target	**Antibody Name and/or Binding Sequences**
	- Antibodies having CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID Nos:41-43 and CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID Nos:44-46.

[0177] In some aspects, the targeting moiety competes with an antibody set forth in Table F for binding to the target molecule. In further aspects, the targeting moiety comprises CDRs having CDR sequences of an antibody set forth in Table F. In some embodiments, the targeting moiety comprises all 6 CDR sequences of the antibody set forth in Table F. In other embodiments, the targeting moiety comprises at least the heavy chain CDR sequences (CDR-H1, CDR-H2, CDR-H3) of an antibody set forth in Table F and the light chain CDR sequences of a universal light chain. In further aspects, a targeting moiety comprises a VH comprising the amino acid sequence of the VH of an antibody set forth in Table F. In some embodiments, the targeting moiety further comprises a VL comprising the amino acid sequence of the VL of the antibody set forth in Table F. In other embodiments, the targeting moiety further comprises a universal light chain VL sequence.

[0178] In some embodiments, the target molecule is PDL1. Table F-1 below provides exemplary anti-PDL1 antibodies and/or antibody sequences upon which the targeting moiety can be based, e.g., which can be incorporated into a targeting moiety for use in the interferon receptor agonists of the disclosure.

TABLE F-1 Exemplary PDL1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
PDL1	Durvalumab (MEDI4736) HC: SEQ ID NO: 26 of PCT application No. WO2020225552 LC: SEQ ID NO: 27 of PCT application No. WO2020225552
PDL1	Atezolizumab (Tecentriq, MPDL3280A, RG7446) HC: SEQ ID NO: 20 of US Patent No. 8,217,149 LC: SEQ ID NO: 21 of US Patent No. 8,217,149
PDL1	MDX 1105 (BMS-936559)
PDL1	Anti-PDL1 antibodies described in U.S. Patent No. 7,943,743 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos: 1/11

TABLE F-1 Exemplary PDL1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
	SEQ ID Nos: 2/12 SEQ ID Nos: 3/13 SEQ ID Nos: 4/14 SEQ ID Nos: 5/15 SEQ ID Nos: 6/16 SEQ ID Nos: 7/17 SEQ ID Nos: 8/18 SEQ ID Nos: 9/19 SEQ ID Nos: 10/20
PDL1	Avelumab, described in US Patent No: 9,624,298 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 24 LC: SEQ ID NO: 25
PDL1	ZKAB001 (Socazolimab)
PDL1	TQB2450 (APL-502 or CBT-502)
PDL1	HLX20 CDR-H1, CDR-H2 and CDR-H3 sequences of SEQ ID Nos: 52, 56, and 77 of PCT Pub. No. 2018/080812 CDR-L1, CDR-L2 and CDR-L3 sequences of SEQ ID Nos: 65, 42, and 71 of PCT Pub. No. 2018/080812
PDL1	KN035 (Envafolimab) is a nanobody described as Hu56V2 in US Patent No. 11,225,522 as having the VHH SEQ ID NO: 34
PDL1	LY3434172
PDL1	LY3300054 (Iodapolimab) described in PCT Pub No: WO 2017/034916 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 10 L: SEQ ID NO: 11
PDL1	LDP (Iesabelimab, ADG104) described in CN Patent No: 114225023 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 10 LC: SEQ ID NO: 9
PDL1	EMB-09
PDL1	ABL501
PDL1	INBRX-105
PDL1	STI-3031 (IMC-001) described in US Patent No: 10,118,963 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 1 LC: SEQ ID NO: 2
PDL1	BGB-A333 (garivulimab) described in US Patent No: 11,512,132 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 22

TABLE F-1 Exemplary PDL1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
	LC: SEQ ID NO: 23
PDL1	HLX301
PDL1	Y101D
PDL1	ES101
PDL1	IBI322
PDL1	VH: SEQ ID NO:46, 48, 50 or 52 of US Patent No. 11,168,144. VL: SEQ ID NO:58, 137 or 12 of US Patent No. 11, 168,144.
PDL1	VH: SEQ ID NO:23, 124, 126, 127, 128, 130, 140 or 145 of US Patent No. 11,208,486. VL: SEQ ID NO:24 or 125 of US Patent No. 11,208,486.

[0179] In some aspects, the targeting moiety competes with an anti-PDL1 antibody set forth in Table F-1 for binding to PDL1. In further aspects, the targeting moiety comprises CDRs having CDR sequences of an anti-PDL1 antibody set forth in Table F-1. In some embodiments, the targeting moiety comprises all 6 CDR sequences of the anti-PDL1 antibody set forth in Table F-1. In other embodiments, the targeting moiety comprises at least the heavy chain CDR sequences (CDR-H1, CDR-H2, CDR-H3) of an anti-PDL1 antibody set forth in Table F-1 and the light chain CDR sequences of a universal light chain. In further aspects, a targeting moiety comprises a VH comprising the amino acid sequence of the VH of an anti-PDL1 antibody set forth in Table F-1. In some embodiments, the targeting moiety further comprises a VL comprising the amino acid sequence of the VL of the anti-PDL1 antibody set forth in Table F-1. In other embodiments, the targeting moiety further comprises a universal light chain VL sequence.

[0180] In some embodiments, the target molecule is PD1. Table F-2 below provides exemplary anti-PD1 antibodies and/or antibody sequences upon which the targeting moiety can be based, e.g., which can be incorporated into a targeting moiety for use in the interferon receptor agonists of the disclosure.

TABLE F-2 Exemplary PD1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
PD1	MDX-1106 / BMS-936558 (nivolumab), a human IgG4 mAb with the structure described in WHO Drug Information, Vol. 27, No. 1, pages 68-69 (2013) and

TABLE F-2 Exemplary PD1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
	whose heavy and light chain sequences are disclosed in FIG. 7 of US Pub. No. US20190270812A1 HC: SEQ ID NO: 23 of US Pub. No. US20190270812A1 LC: SEQ ID NO: 24 of US Pub. No. US20190270812A1
PD1	MK-3475 (pembrolizumab), a humanized IgG4 mAb with the structure described in WHO Drug Information, Vol. 27, No. 2, pages 161-162 (2013) and whose heavy and light chain sequences are disclosed in FIG. 6 of US Pub. No. US20190270812A1 HC: SEQ ID NO: 21 of US Pub. No. US20190270812A1 LC: SEQ ID NO: 22 of US Pub. No. US20190270812A1
PD1	REGN2810 (disclosed as H4H7798N in US Pub No. 20150203579) HC: SEQ ID NO: 330 of US Pub. No. 20150203579 LC: SEQ ID NO: 331 of US Pub. No. 20150203579
PD1	Anti-PD1 antibodies having CDR H1-H3 and CDR L1-L3 sequences corresponding to the following SEQ ID Nos. of US Patent No. 11,034,765 B2: a) SEQ ID Nos: 18, 19, 20, 21, 22, and 23, respectively; b) SEQ ID Nos: 24, 25, 26, 27, 28, and 29, respectively; c) SEQ ID Nos: 30, 31, 32, 33, 34, and 35, respectively; d) SEQ ID Nos: 36, 37, 38, 39, 40, and 41, respectively; e) SEQ ID Nos: 42, 43, 44, 45, 46, and 47, respectively; f) SEQ ID Nos: 48, 49, 50, 51, 52, and 53, respectively; g) SEQ ID Nos: 54, 55, 56, 57, 58, and 59, respectively; and h) SEQ ID Nos: 60, 61, 62, 63, 64, and 65, respectively.
PD1	Anti-PD1 antibodies disclosed in Tables 1-3 of PCT Pub. WO2015112800A1, including but not limited to anti-PD1 antibodies having VH/VL pairs having SEQ ID Nos: 2/10, 18/26, 34/42, 50/58, 66/74, 82/90, 98/106, 1 14/122, 130/138, 146/154, 162/170, 178/186, 194/202, 210/202, 218/202, 226/202, 234/202, 242/202, 250/202, 258/202, 266/202, 274/202, 282/202, 290/202, 298/186, 306/186 and 314/186 of PCT Pub. WO2015112800A1.
PD1	Anti-PD1 antibodies disclosed in US Patent No. 10,294,299 B2 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 164/178 SEQ ID Nos. 165/179 SEQ ID Nos. 166/180 SEQ ID Nos. 167/181 SEQ ID Nos. 168/182 SEQ ID Nos. 169/183 SEQ ID Nos. 170/184 SEQ ID Nos. 171/185 SEQ ID Nos. 172/186 SEQ ID Nos. 173/187 SEQ ID Nos. 174/188 SEQ ID Nos. 175/189

TABLE F-2 Exemplary PD1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
	SEQ ID Nos. 176/190 SEQ ID Nos. 177/190
PD1	MEDI-0680 (AMP-514)
PD1	PDR001 (spartalizumab), a humanized IgG4 mAb whose heavy and light chain sequences are disclosed as BAP049-Clone-E in US Patent No: 9683048 B2. HC: SEQ ID NO: 91 of US Patent No: 9,683,048 LC: SEQ ID NO: 72 of US Patent No: 9,683,048
PD1	BGB-108
PD1	h409A11, described in WO2008/156712 HC: SEQ ID NO: 31 of PCT Pub. WO2008/156712 LC: SEQ ID NO: 36 of PCT Pub. WO2008/156712
PD1	h409A16, described in WO2008/156712 HC: SEQ ID NO: 31 of PCT Pub. WO2008/156712 LC: SEQ ID NO: 37 of PCT Pub. WO2008/156712
PD1	h409A17, described in WO2008/156712 HC: SEQ ID NO: 31 of PCT Pub. WO2008/156712 LC: SEQ ID NO: 38 of PCT Pub. WO2008/156712
PD1	Anti-PD1 antibodies described in US Patent No. 7,488,802 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 2/4 SEQ ID Nos. 6/8 SEQ ID Nos. 10/12 SEQ ID Nos. 14/16 SEQ ID Nos. 47/49
PD1	Anti-PD1 antibodies described in US Patent No. 7,521,051 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 2/4 SEQ ID Nos. 6/8 SEQ ID Nos. 10/12 SEQ ID Nos. 14/16 SEQ ID Nos. 47/49
PD1	Anti-PD1 antibodies described in US Patent No. 8,008,449 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 1/8 SEQ ID Nos. 2/9 SEQ ID Nos. 3/10 SEQ ID Nos. 4/11 SEQ ID Nos. 5/12 SEQ ID Nos. 6/13 SEQ ID Nos. 7/14

TABLE F-2 Exemplary PD1 antibodies and/or binding sequences	
Target	**Antibody Name and/or Binding Sequences**
PD1	Anti-PD1 antibodies described in US Patent No. 8,354,509 as having the following SEQ ID NO. pairs for heavy and light chain full-length domains: SEQ ID Nos. 31/36 SEQ ID Nos. 31/37 SEQ ID Nos. 31/38
PD1	Anti-PD1 antibodies described in US Patent No. 8,168,757 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 4/5 SEQ ID Nos. 12/13 SEQ ID Nos. 18/19 SEQ ID Nos.40/41 SEQ ID Nos.47/48 SEQ ID Nos.26/27 SEQ ID Nos.34/35 SEQ ID Nos.55/56 SEQ ID Nos.67/68
PD1	Anti-PD1 antibodies described in PCT Pub. No. WO2004/004771
PD1	Anti-PD1 antibodies described in PCT Pub. No. WO2004/056875 as having the following SEQ ID NO. pairs for heavy and light chain variable domains: SEQ ID Nos. 2/4 SEQ ID Nos. 6/8 SEQ ID Nos. 10/12 SEQ ID Nos. 14/16 SEQ ID Nos. 47/49
PD1	Anti-PD1 antibodies described in PCT Pub. No. WO2004/072286
PD1	VH: SEQ ID NO: 25, 26, 27, 28, or 29 of US Pub. No. US2011/0271358 VL: SEQ ID NO: 30, 31, 32, or 33 of US Pub. No. US2011/0271358
PD1	SHR-1210 (Camrelizumab) described in PCT Publication No: WO 2015/085847 as having the following heavy and light chain variable domains: HC: SEQ ID NO: 9 LC: SEQ ID NO: 10

[0181] In some aspects, the targeting moiety competes with an anti-PD1 antibody set forth in Table F-2 for binding to PD1. In further aspects, the targeting moiety comprises CDRs having CDR sequences of an anti-PD1 antibody set forth in Table F-2. In some embodiments, the targeting moiety comprises all 6 CDR sequences of the anti-PD1 antibody set forth in Table F-2. In other embodiments, the targeting moiety comprises at least the heavy chain CDR sequences (CDR-H1, CDR-H2, CDR-H3) of an anti-PD1 antibody set forth in Table F-2 and the light chain CDR sequences of a universal light chain. In further

aspects, a targeting moiety comprises a VH comprising the amino acid sequence of the VH of an anti-PD1 antibody set forth in Table F-2. In some embodiments, the targeting moiety further comprises a VL comprising the amino acid sequence of the VL of the anti-PD1 antibody set forth in Table F-2. In other embodiments, the targeting moiety further comprises a universal light chain VL sequence.

[0182] Where the target molecule is a checkpoint inhibitor, in some embodiments, the checkpoint inhibitor targeting moiety is non-blocking or poorly-blocking of ligand-receptor binding. Examples of non-blocking or poorly-blocking anti-PD1 antibodies includes antibodies having VH/VL amino acid sequences of SEQ ID NOs: 2/10 of PCT Pub. No. WO2015/112800A1; SEQ ID NOs: 16/17 of US Patent No. 11,034,765 B2; SEQ ID NOs. 164/178, 165/179, 166/180, 167/181, 168/182, 169/183, 170/184, 171/185, 172/186, 173/187, 174/188, 175/189, 176/190 and 177/190 of US Patent No. 10,294,299 B2. Examples of non-blocking or poorly-blocking anti-LAG3 antibodies includes antibodies having VH/VL amino acid sequences of SEQ ID NOs 23/24, 3/4 and 11/12 of US Pub. US2022/0056126A1.

[0183] Additional target molecules that can be targeted by the IFN receptor agonists are disclosed in Table I below and in, *e.g.*, Hafeez *et al.*, 2020, Molecules 25:4764, doi:10.3390/molecules25204764, particularly in Table 1. Table 1 of Hafeez *et al.* is incorporated by reference in its entirety here.

6.8. Targeting Moiety Formats

[0184] In certain aspects, the targeting moiety of an IFN receptor agonist of the disclosure can be any type of antibody or fragment thereof that retains specific binding to an antigenic determinant. In one embodiment the targeting moiety is an immunoglobulin molecule or fragment thereof, particularly an IgG class immunoglobulin molecule, more particularly an IgG_1 or IgG_4 immunoglobulin molecule. Antibody fragments include, but are not limited to, VH (or V_H) fragments, VL (or V_L) fragments, Fab fragments, $F(ab')_2$ fragments, scFv fragments, Fv fragments, minibodies, diabodies, triabodies, and tetrabodies.

6.8.1. Fab

[0185] Fab domains were traditionally produced by proteolytic cleavage of immunoglobulin molecules using enzymes such as papain. The Fab domains can comprise constant domain

and variable region sequences from any suitable species, and thus can be murine, chimeric, human or humanized.

[0186] Fab domains typically comprise a CH1 domain attached to a VH domain which pairs with a CL domain attached to a VL domain. In a wild-type immunoglobulin, the VH domain is paired with the VL domain to constitute the Fv region, and the CH1 domain is paired with the CL domain to further stabilize the binding site. A disulfide bond between the two constant domains can further stabilize the Fab domain.

[0187] For the IFN receptor agonists of the disclosure, particularly when the light chains of the targeting moieties are not common or universal light chains, it is advantageous to use Fab heterodimerization strategies to permit the correct association of Fab domains belonging to the same targeting moiety and minimize aberrant pairing of Fab domains belonging to different targeting moieties. For example, the Fab heterodimerization strategies shown in Table G below can be used:

TABLE G
Fab Heterodimerization Strategies

STRATEGY	VH	CH1	VL	CL	REFERENCE
CrossMabCH1-CL	WT	CL domain	WT	CH1 domain	Schaefer *et al.*, 2011, Cancer Cell 2011; 20:472-86; PMID:22014573.
orthogonal Fab VHVRD1CH1CRD2 - VLVRD1Cλ CRD2	39K, 62E	H172A, F174G	1R, 38D, (36F)	L135Y, S176W	Lewis *et al.*, 2014, Nat Biotechnol 32:191-8
orthogonal Fab VHVRD2CH1wt - VLVRD2Cλ wt	39Y	WT	38R	WT	Lewis *et al.*, 2014, Nat Biotechnol 32:191-8
TCR CαCβ	39K	TCR Cα	38D	TCR Cβ	Wu *et al.*, 2015, MAbs 7:364-76

TABLE G Fab Heterodimerization Strategies					
STRATEGY	**VH**	**CH1**	**VL**	**CL**	**REFERENCE**
CR3	WT	T192E	WT	N137K, S114A	Golay *at al.*, 2016, J Immunol 196:3199-211.
MUT4	WT	L143Q, S188V	WT	V133T, S176V	Golay *at al.*, 2016, J Immunol 196:3199-211.
DuetMab	WT	F126C	WT	S121C	Mazor *et al.*, 2015, MAbs 7:377-89; Mazor *et al.*, 2015, MAbs 7:461-669.
Domain exchanged	WT	CH3 + knob or hole mutation	WT	CH3 + hole or knob mutation	Wozniak-Knopp *et al.*, 2018, PLoSONE13(4):e0195442

[0188] Accordingly, in certain embodiments, correct association between the two polypeptides of a Fab is promoted by exchanging the VL and VH domains of the Fab for each other or exchanging the CH1 and CL domains for each other, *e.g.,* as described in WO 2009/080251.

[0189] Correct Fab pairing can also be promoted by introducing one or more amino acid modifications in the CH1 domain and one or more amino acid modifications in the CL domain of the Fab and/or one or more amino acid modifications in the VH domain and one or more amino acid modifications in the VL domain. The amino acids that are modified are typically part of the VH:VL and CH1:CL interface such that the Fab components preferentially pair with each other rather than with components of other Fabs.

[0190] In one embodiment, the one or more amino acid modifications are limited to the conserved framework residues of the variable (VH, VL) and constant (CH1, CL) domains as indicated by the Kabat numbering of residues. Almagro, 2008, Frontiers in Bioscience 13:1619-1633 provides a definition of the framework residues on the basis of Kabat, Chothia, and IMGT numbering schemes.

[0191] In one embodiment, the modifications introduced in the VH and CH1 and/or VL and CL domains are complementary to each other. Complementarity at the heavy and light chain interface can be achieved on the basis of steric and hydrophobic contacts, electrostatic/charge interactions or a combination of the variety of interactions. The complementarity between protein surfaces is broadly described in the literature in terms of lock and key fit, knob into hole, protrusion and cavity, donor and acceptor *etc.*, all implying the nature of structural and chemical match between the two interacting surfaces.

[0192] In one embodiment, the one or more introduced modifications introduce a new hydrogen bond across the interface of the Fab components. In one embodiment, the one or more introduced modifications introduce a new salt bridge across the interface of the Fab components. Exemplary substitutions are described in WO 2014/150973 and WO 2014/082179, the contents of which are hereby incorporated by reference.

[0193] In some embodiments, the Fab domain comprises a 192E substitution in the CH1 domain and 114A and 137K substitutions in the CL domain, which introduces a salt-bridge between the CH1 and CL domains (see, *e.g.,* Golay *et al.*, 2016, J Immunol 196:3199-211).

[0194] In some embodiments, the Fab domain comprises a 143Q and 188V substitutions in the CH1 domain and 113T and 176V substitutions in the CL domain, which serves to swap hydrophobic and polar regions of contact between the CH1 and CL domain (see, *e.g.,* Golay *et al.*, 2016, J Immunol 196:3199-211).

[0195] In some embodiments, the Fab domain can comprise modifications in some or all of the VH, CH1, VL, CL domains to introduce orthogonal Fab interfaces which promote correct assembly of Fab domains (Lewis *et al.*, 2014 Nature Biotechnology 32:191-198). In an embodiment, 39K, 62E modifications are introduced in the VH domain, H172A, F174G modifications are introduced in the CH1 domain, 1 R, 38D, (36F) modifications are introduced in the VL domain, and L135Y, S176W modifications are introduced in the CL domain. In another embodiment, a 39Y modification is introduced in the VH domain and a 38R modification is introduced in the VL domain.

[0196] Fab domains can also be modified to replace the native CH1:CL disulfide bond with an engineered disulfide bond, thereby increasing the efficiency of Fab component pairing. For example, an engineered disulfide bond can be introduced by introducing a 126C in the CH1 domain and a 121 C in the CL domain (see, *e.g.,* Mazor *et al.*, 2015, MAbs 7:377-89).

[0197] Fab domains can also be modified by replacing the CH1 domain and CL domain with alternative domains that promote correct assembly. For example, Wu *et al.*, 2015, MAbs 7:364-76, describes substituting the CH1 domain with the constant domain of the T cell receptor and substituting the CL domain with the b domain of the T cell receptor, and pairing these domain replacements with an additional charge-charge interaction between the VL and VH domains by introducing a 38D modification in the VL domain and a 39K modification in the VH domain.

[0198] In lieu of, or in addition to, the use of Fab heterodimerization strategies to promote correct VH-VL pairings, the VL of common light chain (also referred to as a universal light chain) can be used for each unique ABD in the IFN receptor agonists of the disclosure. In various embodiments, employing a common light chain as described herein reduces the number of inappropriate species in the IFN receptor agonists as compared to employing original cognate VLs. In various embodiments, the VL domains of ABDs are identified from monospecific antibodies comprising a common light chain. In various embodiments, the VH regions of the ABDs in the IFN receptor agonists comprise human heavy chain variable gene segments that are rearranged *in vivo* within mouse B cells that have been previously engineered to express a limited human light chain repertoire, or a single human light chain, cognate with human heavy chains and, in response to exposure with an antigen of interest, generate an antibody repertoire containing a plurality of human VHs that are cognate with one or one of two possible human VLs, wherein the antibody repertoire specific for the antigen of interest. Common light chains are those derived from a rearranged human Vκ1-39Jκ5 sequence or a rearranged human Vκ3-20Jκ1 sequence, and include somatically mutated (*e.g.,* affinity matured) versions. See, for example, U.S. Patent No. 10,412,940.

6.8.2. scFv

[0199] Single chain Fv or "scFv" antibody fragments comprise the VH and VL domains of an antibody in a single polypeptide chain, are capable of being expressed as a single chain polypeptide, and retain the specificity of the intact antibodies from which they are derived. Generally, the scFv polypeptide further comprises a polypeptide linker between the VH and VL domain that enables the scFv to form the desired structure for target binding. Examples of linkers suitable for connecting the VH and VL chains of an scFv are the non-cleavable linkers identified in Section 6.6.

[0200] Unless specified, as used herein an scFv may have the VL and VH variable regions in either order, *e.g.*, with respect to the N-terminal and C-terminal ends of the polypeptide, the scFv may comprise VL-linker-VH or may comprise VH-linker-VL.

[0201] The scFv can comprise VH and VL sequences from any suitable species, such as murine, human or humanized VH and VL sequences.

[0202] To create an scFv-encoding nucleic acid, the VH and VL-encoding DNA fragments are operably linked to another fragment encoding a linker, *e.g.*, encoding any of the linkers described in Section 6.6 (typically a repeat of a sequence containing the amino acids glycine and serine, such as the amino acid sequence (Gly4~Ser)3 (SEQ ID NO: 182), such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see, *e.g.*, Bird *et al.*, 1988, Science 242:423-426; Huston *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; McCafferty *et al.*, 1990, Nature 348:552-554).

6.9. Fc Regions

[0203] The IFN receptor agonists of the disclosure typically include a pair of Fc domains that associate to form an Fc region. In native antibodies, Fc regions comprise hinge regions at their N-termini to form a constant domain. Throughout this disclosure, the reference to an Fc domain encompasses an Fc domain with a hinge domain at its N-terminus unless specified otherwise.

[0204] The Fc domains can be derived from any suitable species operably linked to an ABD or component thereof. In one embodiment the Fc domain is derived from a human Fc domain. In preferred embodiments, the targeting moiety or component thereof is fused to an IgG Fc molecule. A targeting moiety or component thereof may be fused to the N-terminus or the C-terminus of the IgG Fc domain or both.

[0205] The Fc domains can be derived from any suitable class of antibody, including IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3 and IgG4), and IgM. In one embodiment, the Fc domain is derived from IgG1, IgG2, IgG3 or IgG4. In one embodiment the Fc domain is derived from IgG1. In one embodiment the Fc domain is derived from IgG4. Exemplary sequences of Fc domains from IgG1, IgG2, IgG3, and IgG4 are provided in Table Y, below.

TABLE Y Fc Sequences		
Fc	**Sequence**	**SEQ ID NO**
hIgG1 Fc (amino acids 99-330 of UniprotKB P01857-1)	EPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTC VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS LSLSPGK	410
hIgG2 Fc (amino acids 99-326 of UniprotKB P01859-1)	ERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVV DVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVSVLTV VHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPP SREEMTKNQVSLTCLVKGFYPSDISVEWESNGQPENNYKTTPPML DSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLS LSPGK	411
hIgG3 Fc (amino acids 99-377 of UniprotKB P01860-1)	ELKTPLGDTTHTCPRCPEPKSCDTPPPCPRCPEPKSCDTPPPCPR CPEPKSCDTPPPCPRCPAPELLGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVQFKWYVDGVEVHNAKTKPREEQYNSTFRVV SVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKTKGQPREPQVY TLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESSGQPENNYNTT PPMLDSDGSFFLYSKLTVDKSRWQQGNIFSCSVMHEALHNRFTQ KSLSLSPGK	412
hIgG4 Fc (amino acids 99-327 of UniprotKB P01861-1)	ESKYGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVV DVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTV LHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVL DSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLS LSLGK	413

[0206] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at eat least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO: 410. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO: 410 (*e.g.,* between 90% and 99% sequence identity to SEQ ID NO: 410), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (*e.g.*, as described in Section 6.9.1)

and/or one or more substitutions that promote Fc heterodimerization (*e.g.*, as described in Section 6.9.2).

[0207] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO: 411. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO: 411 (*e.g.,* between 90% and 99% sequence identity to SEQ ID NO: 411), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (*e.g.*, as described in Section 6.9.1) and/or one or more substitutions that promote Fc heterodimerization (*e.g.*, as described in Section 6.9.2).

[0208] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at eat least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO: 412. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO: 412 (*e.g.,* between 90% and 99% sequence identity to SEQ ID NO: 412), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (*e.g.*, as described in Section 6.9.1) and/or one or more substitutions that promote Fc heterodimerization (*e.g.*, as described in Section 6.9.2).

[0209] In some aspects, an Fc domain comprises an amino acid sequence having at least about 90%, at least about 91%, at least about 92%, about at least 93%, at least about 94%, at eat least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% sequence identity to SEQ ID NO: 413. In cases where an Fc domain comprises at least 90% sequence identity and less than 100% sequence identity to SEQ ID NO: 413 (*e.g.,* between 90% and 99% sequence identity to SEQ ID NO: 413), an Fc domain may also comprise one or more amino acid substitutions described herein, for example one or more substitutions that reduce effector function (*e.g.*, as described in Section 6.9.1)

and/or one or more substitutions that promote Fc heterodimerization (*e.g.*, as described in Section 6.9.2).

[0210] The two Fc domains within the Fc region can be the same or different from one another. In a native antibody the Fc domains are typically identical, but for the purpose of producing multispecific binding molecules, *e.g.,* the IFN receptor agonists of the disclosure and MBMs produced by their activation, the Fc domains might advantageously be different to allow for heterodimerization, as described in Section 6.9.2 below.

[0211] In native antibodies, the heavy chain Fc domain of IgA, IgD and IgG is composed of two heavy chain constant domains (CH2 and CH3) and that of IgE and IgM is composed of three heavy chain constant domains (CH2, CH3 and CH4). These dimerize to create an Fc region.

[0212] In IFN receptor agonists of the present disclosure, the Fc region, and / or the Fc domains within it, can comprise heavy chain constant domains from one or more different classes of antibody, for example one, two or three different classes.

[0213] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG1.

[0214] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG2.

[0215] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG3.

[0216] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG4.

[0217] In one embodiment the Fc region comprises a CH4 domain from IgM. The IgM CH4 domain is typically located at the C-terminus of the CH3 domain.

[0218] In one embodiment the Fc region comprises CH2 and CH3 domains derived from IgG and a CH4 domain derived from IgM.

[0219] It will be appreciated that the heavy chain constant domains for use in producing an Fc region for the IFN receptor agonists of the present disclosure may include variants of the naturally occurring constant domains described above. Such variants may comprise one or

more amino acid variations compared to wild type constant domains. In one example the Fc region of the present disclosure comprises at least one constant domain that varies in sequence from the wildtype constant domain. It will be appreciated that the variant constant domains may be longer or shorter than the wild-type constant domain. Preferably the variant constant domains are at least 60% identical or similar to a wild-type constant domain. In another example the variant constant domains are at least 70% identical or similar. In another example the variant constant domains are at least 80% identical or similar. In another example the variant constant domains are at least 90% identical or similar. In another example the variant constant domains are at least 95% identical or similar.

[0220] IgM and IgA occur naturally in humans as covalent multimers of the common H2L2 antibody unit. IgM occurs as a pentamer when it has incorporated a J-chain, or as a hexamer when it lacks a J-chain. IgA occurs as monomer and dimer forms. The heavy chains of IgM and IgA possess an 18 amino acid extension to the C-terminal constant domain, known as a tailpiece. The tailpiece includes a cysteine residue that forms a disulfide bond between heavy chains in the polymer, and is believed to have an important role in polymerization. The tailpiece also contains a glycosylation site. In certain embodiments, the IFN receptor agonists of the present disclosure do not comprise a tailpiece.

[0221] The Fc domains that are incorporated into the IFN receptor agonists of the present disclosure may comprise one or more modifications that alter the functional properties of the proteins, for example, binding to Fc-receptors such as FcRn or leukocyte receptors, binding to complement, modified disulfide bond architecture, or altered glycosylation patterns. Exemplary Fc modifications that alter effector function are described in Section 6.9.1.

[0222] The Fc domains can also be altered to include modifications that improve manufacturability of asymmetric IFN receptor agonists, for example by allowing heterodimerization, which is the preferential pairing of non-identical Fc domains over identical Fc domains. Heterodimerization permits the production of IFN receptor agonists in which different polypeptide components are connected to one another by an Fc region containing Fc domains that differ in sequence. Examples of heterodimerization strategies are exemplified in Section 6.9.2.

[0223] It will be appreciated that any of the modifications mentioned above can be combined in any suitable manner to achieve the desired functional properties and/or combined with other modifications to alter the properties of the IFN receptor agonists.

6.9.1. Fc Domains with Altered Effector Function

[0224] In some embodiments, the Fc domain comprises one or more amino acid substitutions that reduces binding to an Fc receptor and/or effector function.

[0225] In a particular embodiment the Fc receptor is an Fcγ receptor. In one embodiment the Fc receptor is a human Fc receptor. In one embodiment the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. In one embodiment the effector function is one or more selected from the group of complement dependent cytotoxicity (CDC), antibody-dependent cell-mediated cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP), and cytokine secretion. In a particular embodiment, the effector function is ADCC.

[0226] In one embodiment, the Fc domain (*e.g.*, an Fc domain of an IFN receptor agonist half antibody) or the Fc region (*e.g.*, one or both Fc domains of an IFN receptor agonist that can associate to form an Fc region) comprises an amino acid substitution at a position selected from the group of E233, L234, L235, N297, P331 and P329 (numberings according to Kabat EU index). In a more specific embodiment, the Fc domain or the Fc region comprises an amino acid substitution at a position selected from the group of L234, L235 and P329 (numberings according to Kabat EU index). In some embodiments, the Fc domain or the Fc region comprises the amino acid substitutions L234A and L235A (numberings according to Kabat EU index). In one such embodiment, the Fc domain or region is an Igd Fc domain or region, particularly a human Igd Fc domain or region. In one embodiment, the Fc domain or the Fc region comprises an amino acid substitution at position P329. In a more specific embodiment, the amino acid substitution is P329A or P329G, particularly P329G (numberings according to Kabat EU index). In one embodiment, the Fc domain or the Fc region comprises an amino acid substitution at position P329 and a further amino acid substitution at a position selected from E233, L234, L235, N297 and P331 (numberings according to Kabat EU index). In a more specific embodiment, the further amino acid substitution is E233P, L234A, L235A, L235E, N297A, N297D or P331S. In particular

embodiments, the Fc domain or the Fc region comprises amino acid substitutions at positions P329, L234 and L235 (numberings according to Kabat EU index). In more particular embodiments, the Fc domain comprises the amino acid mutations L234A, L235A and P329G ("P329G LALA", "PGLALA" or "LALAPG").

[0227] Typically, the same one or more amino acid substitution is present in each of the two Fc domains of an Fc region. Thus, in a particular embodiment, each Fc domain of the Fc region comprises the amino acid substitutions L234A, L235A and P329G (Kabat EU index numbering), *i.e.* in each of the first and the second Fc domains in the Fc region the leucine residue at position 234 is replaced with an alanine residue (L234A), the leucine residue at position 235 is replaced with an alanine residue (L235A) and the proline residue at position 329 is replaced by a glycine residue (P329G) (numbering according to Kabat EU index).

[0228] In one embodiment, the Fc domain is an IgG1 Fc domain, particularly a human IgG1 Fc domain. In some embodiments, the IgG1 Fc domain is a variant IgG1 comprising D265A, N297A mutations (EU numbering) to reduce effector function.

[0229] In another embodiment, the Fc domain is an IgG4 Fc domain with reduced binding to Fc receptors. Exemplary IgG4 Fc domains with reduced binding to Fc receptors may comprise an amino acid sequence selected from Table H below: In some embodiments, the Fc domain includes only the bolded portion of the sequences shown below:

TABLE H

Fc Domain	**Sequence**	**SEQ ID NO:**
SEQ ID NO:1 of WO2014/121087	Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val	313

TABLE H		
Fc Domain	**Sequence**	**SEQ ID NO:**
	Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys	
SEQ ID NO:2 of WO2014/121087	Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys	314
SEQ ID NO:30 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val **Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn** **Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His**	315

TABLE H		
Fc Domain	**Sequence**	**SEQ ID NO:**
	Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys	
SEQ ID NO:31 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val **Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn** **Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys**	316
SEQ ID NO:37 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val **Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys**	317

TABLE H		
Fc Domain	**Sequence**	**SEQ ID NO:**
	Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys	
SEQ ID NO:38 of WO2014/121087	Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val **Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn Arg Phe Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys**	409

[0230] In a particular embodiment, the IgG4 with reduced effector function comprises the bolded portion of the amino acid sequence of SEQ ID NO:31 of WO2014/121087, sometimes referred to herein as IgG4s or hIgG4s, having the amino acid sequence: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYV DGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAK

GQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDS
DGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK (SEQ ID NO: 414).

[0231] For heterodimeric Fc regions, it is possible to incorporate a combination of the variant IgG4 Fc sequences set forth above, for example an Fc region comprising an Fc domain comprising the amino acid sequence of SEQ ID NO:30 of WO2014/121087 (or the bolded portion thereof) and an Fc domain comprising the amino acid sequence of SEQ ID NO:37 of WO2014/121087 (or the bolded portion thereof) or an Fc region comprising an Fc domain comprising the amino acid sequence of SEQ ID NO:31 of WO2014/121087 (or the bolded portion thereof) and an Fc domain comprising the amino acid sequence of SEQ ID NO:38 of WO2014/121087 (or the bolded portion thereof).

6.9.2. Fc Heterodimerization Variants

[0232] Certain IFN receptor agonists entail dimerization between two Fc domains that, unlike a native immunoglobulin, are operably linked to non-identical N-terminal or C-terminal regions. Inadequate heterodimerization of two Fc domains to form an Fc region has can be an obstacle for increasing the yield of desired heterodimeric molecules and represents challenges for purification. A variety of approaches available in the art can be used in for enhancing dimerization of Fc domains that might be present in the IFN receptor agonists of the disclosure, for example as disclosed in EP 1870459A1; U.S. Patent No. 5,582,996; U.S. Patent No. 5,731,168; U.S. Patent No. 5,910,573; U.S. Patent No. 5,932,448; U.S. Patent No. 6,833,441; U.S. Patent No. 7,183,076; U.S. Patent Application Publication No. 2006204493A1; and PCT Publication No. WO 2009/089004A1.

[0233] In some embodiments, the present disclosure provides IFN receptor agonists comprising Fc heterodimers, *i.e.*, Fc regions comprising heterologous, non-identical Fc domains. Typically, each Fc domain in the Fc heterodimer comprises a CH3 domain of an antibody. The CH3 domains are derived from the constant region of an antibody of any isotype, class or subclass, and preferably of IgG (IgG1, IgG2, IgG3 and IgG4) class, as described in the preceding section.

[0234] Heterodimerization of the two different heavy chains at CH3 domains give rise to the desired IFN receptor agonist, while homodimerization of identical heavy chains will reduce yield of the desired IFN receptor agonist. Thus, in a preferred embodiment, the polypeptides

that associate to form an IFN receptor agonist of the disclosure will contain CH3 domains with modifications that favor heterodimeric association relative to unmodified Fc domains.

[0235] In a specific embodiment said modification promoting the formation of Fc heterodimers is a so-called "knob-into-hole" or "knob-in-hole" modification, comprising a "knob" modification in one of the Fc domains and a "hole" modification in the other Fc domain. The knob-into-hole technology is described *e.g.*, in U.S. Patent No. 5,731,168; US 7,695,936; Ridgway *et al.*, 1996, Prot Eng 9:617-621, and Carter, 2001, Immunol Meth 248:7-15. Generally, the method involves introducing a protuberance ("knob") at the interface of a first polypeptide and a corresponding cavity ("hole") in the interface of a second polypeptide, such that the protuberance can be positioned in the cavity so as to promote heterodimer formation and hinder homodimer formation. Protuberances are constructed by replacing small amino acid side chains from the interface of the first polypeptide with larger side chains (*e.g.*, tyrosine or tryptophan). Compensatory cavities of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino acid side chains with smaller ones (*e.g.*, alanine or threonine).

[0236] Accordingly, in some embodiments, an amino acid residue in the CH3 domain of the first subunit of the Fc domain is replaced with an amino acid residue having a larger side chain volume, thereby generating a protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and an amino acid residue in the CH3 domain of the second subunit of the Fc domain is replaced with an amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable. Preferably said amino acid residue having a larger side chain volume is selected from the group consisting of arginine (R), phenylalanine (F), tyrosine (Y), and tryptophan (W). Preferably said amino acid residue having a smaller side chain volume is selected from the group consisting of alanine (A), serine (S), threonine (T), and valine (V). The protuberance and cavity can be made by altering the nucleic acid encoding the polypeptides, *e.g.*, by site-specific mutagenesis, or by peptide synthesis. An exemplary substitution is Y470T.

[0237] In a specific such embodiment, in the first Fc domain the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in the Fc domain the tyrosine residue at position 407 is replaced with a valine residue (Y407V) and optionally the threonine residue at position 366 is replaced with a serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A) (numbering according to Kabat EU index). In a further embodiment, in the first Fc domain additionally the serine residue at position 354 is replaced with a cysteine residue (S354C) or the glutamic acid residue at position 356 is replaced with a cysteine residue (E356C) (particularly the serine residue at position 354 is replaced with a cysteine residue), and in the second Fc domain additionally the tyrosine residue at position 349 is replaced by a cysteine residue (Y349C) (numbering according to Kabat EU index). In a particular embodiment, the first Fc domain comprises the amino acid substitutions S354C and T366W, and the second Fc domain comprises the amino acid substitutions Y349C, T366S, L368A and Y407V (numbering according to Kabat EU index).

[0238] In some embodiments, electrostatic steering (*e.g.*, as described in Gunasekaran *et al.*, 2010, J Biol Chem 285(25): 19637-46) can be used to promote the association of the first and the second Fc domains of the Fc region.

[0239] As an alternative, or in addition, to the use of Fc domains that are modified to promote heterodimerization, an Fc domain can be modified to allow a purification strategy that enables selections of Fc heterodimers. In one such embodiment, one polypeptide comprises a modified Fc domain that abrogates its binding to Protein A, thus enabling a purification method that yields a heterodimeric protein. See, for example, U.S. Patent No. 8,586,713. As such, the IFN receptor agonists comprise a first CH3 domain and a second Ig CH3 domain, wherein the first and second Ig CH3 domains differ from one another by at least one amino acid, and wherein at least one amino acid difference reduces binding of the IFN receptor agonist to Protein A as compared to a corresponding IFN receptor agonist lacking the amino acid difference. In one embodiment, the first CH3 domain binds Protein A and the second CH3 domain contains a mutation/modification that reduces or abolishes Protein A binding such as an H95R modification (by IMGT exon numbering; H435R by EU numbering). The second CH3 may further comprise a Y96F modification (by IMGT; Y436F by EU). This class of modifications is referred to herein as “star” mutations.

[0240] In some embodiments, the Fc can contain one or more mutations (*e.g.*, knob and hole mutations) to facilitate heterodimerization as well as star mutations to facilitate purification.

6.9.3. Hinge Domains

[0241] The IFN receptor agonists of the disclosure can comprise an Fc domain comprising a hinge domain at its N-terminus. The hinge region can be a native or a modified hinge region. Hinge regions are typically found at the N-termini of Fc regions. The term "hinge domain", unless the context dictates otherwise, refers to a naturally or non-naturally occurring hinge sequence that in the context of a single or monomeric polypeptide chain is a monomeric hinge domain and in the context of a dimeric polypeptide (*e.g.*, a homodimeric or heterodimeric IFN receptor agonist formed by the association of two Fc domains) can comprise two associated hinge sequences on separate polypeptide chains. Sometimes, the two associated hinge sequences are referred to as a "hinge region". In certain embodiments of IFN receptor agonists, additional iterations of hinge regions may be incorporated into the polypeptide sequence.

[0242] A native hinge region is the hinge region that would normally be found between Fab and Fc domains in a naturally occurring antibody. A modified hinge region is any hinge that differs in length and/or composition from the native hinge region. Such hinges can include hinge regions from other species, such as human, mouse, rat, rabbit, shark, pig, hamster, camel, llama or goat hinge regions. Other modified hinge regions may comprise a complete hinge region derived from an antibody of a different class or subclass from that of the heavy chain Fc domain or Fc region. Alternatively, the modified hinge region may comprise part of a natural hinge or a repeating unit in which each unit in the repeat is derived from a natural hinge region. In a further alternative, the natural hinge region may be altered by converting one or more cysteine or other residues into neutral residues, such as serine or alanine, or by converting suitably placed residues into cysteine residues. By such means the number of cysteine residues in the hinge region may be increased or decreased. Other modified hinge regions may be entirely synthetic and may be designed to possess desired properties such as length, cysteine composition and flexibility.

[0243] A number of modified hinge regions have already been described for example, in U.S. Patent No. 5,677,425, WO 99/15549, WO 2005/003170, WO 2005/003169, WO

2005/003170, WO 98/25971 and WO 2005/003171 and these are incorporated herein by reference.

[0244] In one embodiment, an IFN receptor agonist of the disclosure comprises an Fc region in which one or both Fc domains possesses an intact hinge domain at its N-terminus.

[0245] In various embodiments, positions 233-236 within a hinge region may be G, G, G and unoccupied; G, G, unoccupied, and unoccupied; G, unoccupied, unoccupied, and unoccupied; or all unoccupied, with positions numbered by EU numbering.

[0246] In some embodiments, the IFN receptor agonists of the disclosure comprise a modified hinge region that reduces binding affinity for an Fcγ receptor relative to a wild-type hinge region of the same isotype (*e.g.*, human IgG1 or human IgG4).

[0247] In embodiment, the IFN receptor agonists of the disclosure comprise an Fc region in which each Fc domain possesses an intact hinge domain at its N-terminus, where each Fc domain and hinge domain is derived from IgG4, and each hinge domain comprises the modified sequence CPPC (SEQ ID NO: 377). The core hinge region of human IgG4 contains the sequence CPSC (SEQ ID NO: 378) compared to IgG1 that contains the sequence CPPC (SEQ ID NO: 377). The serine residue present in the IgG4 sequence leads to increased flexibility in this region, and therefore a proportion of molecules form disulfide bonds within the same protein chain (an intrachain disulfide) rather than bridging to the other heavy chain in the IgG molecule to form the interchain disulfide. (Angel *et al.*, 1993, Mol Immunol 30(1):105-108). Changing the serine residue to a proline to give the same core sequence as IgG1 allows complete formation of inter-chain disulfides in the IgG4 hinge region, thus reducing heterogeneity in the purified product. This altered isotype is termed IgG4P.

6.9.3.1. Chimeric Hinge Sequences

[0248] The hinge domain can be a chimeric hinge domain.

[0249] For example, a chimeric hinge may comprise an “upper hinge” sequence, derived from a human IgG1, a human IgG2 or a human IgG4 hinge region, combined with a “lower hinge” sequence, derived from a human IgG1, a human IgG2 or a human IgG4 hinge region.

[0250] In particular embodiments, a chimeric hinge region comprises the amino acid sequence EPKSCDKTHTCPPCPAPPVA (SEQ ID NO: 379) (previously disclosed as SEQ

ID NO:8 of WO2014/121087, which is incorporated by reference in its entirety herein) or ESKYGPPCPPCPAPPVA (SEQ ID NO: 380) (previously disclosed as SEQ ID NO:9 of WO2014/121087). Such chimeric hinge sequences can be suitably linked to an IgG4 CH2 region (for example by incorporation into an IgG4 Fc domain, for example a human or murine Fc domain, which can be further modified in the CH2 and/or CH3 domain to reduce effector function, for example as described in Section 6.9.1).

6.9.3.2. Hinge Sequences with Reduced Effector Function

[0251] In further embodiments, the hinge region can be modified to reduce effector function, for example as described in WO2016161010A2, which is incorporated by reference in its entirety herein. In various embodiments, the positions 233-236 of the modified hinge region are G, G, G and unoccupied; G, G, unoccupied, and unoccupied; G, unoccupied, unoccupied, and unoccupied; or all unoccupied, with positions numbered by EU numbering (as shown in FIG. 1 of WO2016161010A2). These segments can be represented as GGG-, GG--, G--- or ---- with "-" representing an unoccupied position.

[0252] Position 236 is unoccupied in canonical human IgG2 but is occupied by in other canonical human IgG isotypes. Positions 233-235 are occupied by residues other than G in all four human isotypes (as shown in FIG. 1 of WO2016161010A2).

[0253] The hinge modification within positions 233-236 can be combined with position 228 being occupied by P. Position 228 is naturally occupied by P in human IgG1 and IgG2 but is occupied by S in human IgG4 and R in human IgG3. An S228P mutation in an IgG4 antibody is advantageous in stabilizing an IgG4 antibody and reducing exchange of heavy chain light chain pairs between exogenous and endogenous antibodies. Preferably positions 226-229 are occupied by C, P, P and C respectively.

[0254] Exemplary hinge regions have residues 226-236, sometimes referred to as middle (or core) and lower hinge, occupied by the modified hinge sequences designated GGG-(233-236), GG--(233-236), G---(233-236) and no G(233-236). Optionally, the hinge domain amino acid sequence comprises CPPCPAPGGG-GPSVF (SEQ ID NO: 381) (previously disclosed as SEQ ID NO:1 of WO2016161010A2), CPPCPAPGG--GPSVF (SEQ ID NO: 382) (previously disclosed as SEQ ID NO:2 of WO2016161010A2), CPPCPAPG---GPSVF (SEQ ID NO: 383) (previously disclosed as SEQ ID NO:3 of WO2016161010A2), or

CPPCPAP----GPSVF (SEQ ID NO: 384) (previously disclosed as SEQ ID NO:4 of WO2016161010A2).

[0255] The modified hinge regions described above can be incorporated into a heavy chain constant region, which typically include CH2 and CH3 domains, and which may have an additional hinge segment (*e.g.*, an upper hinge) flanking the designated region. Such additional constant region segments present are typically of the same isotype, preferably a human isotype, although can be hybrids of different isotypes. The isotype of such additional human constant regions segments is preferably human IgG4 but can also be human IgG1, IgG2, or IgG3 or hybrids thereof in which domains are of different isotypes. Exemplary sequences of human IgG1, IgG2 and IgG4 are shown in FIGS. 2-4 of WO2016161010A2.

[0256] In specific embodiments, the modified hinge sequences can be linked to an IgG4 CH2 region (for example by incorporation into an IgG4 Fc domain, for example a human or murine Fc domain, which can be further modified in the CH2 and/or CH3 domain to reduce effector function, for example as described in Section 6.9.1).

6.10. Nucleic Acids and Host Cells

[0257] In another aspect, the disclosure provides nucleic acids encoding the IFN receptor agonists of the disclosure. In some embodiments, the IFN receptor agonists are encoded by a single nucleic acid. In other embodiments, the IFN receptor agonists can be encoded by a plurality (*e.g.*, two, three, four or more) nucleic acids.

[0258] A single nucleic acid can encode an IFN receptor agonist that comprises a single polypeptide chain, an IFN receptor agonist that comprises two or more polypeptide chains, or a portion of an IFN receptor agonist that comprises more than two polypeptide chains (for example, a single nucleic acid can encode two polypeptide chains of an IFN receptor agonist comprising three, four or more polypeptide chains, or three polypeptide chains of an IFN receptor agonist comprising four or more polypeptide chains). For separate control of expression, the open reading frames encoding two or more polypeptide chains can be under the control of separate transcriptional regulatory elements (*e.g.*, promoters and/or enhancers). The open reading frames encoding two or more polypeptides can also be controlled by the same transcriptional regulatory elements and separated by internal ribosome entry site (IRES) sequences allowing for translation into separate polypeptides.

[0259] In some embodiments, an IFN receptor agonist comprising two or more polypeptide chains is encoded by two or more nucleic acids. The number of nucleic acids encoding an IFN receptor agonist can be equal to or less than the number of polypeptide chains in the IFN receptor agonist (for example, when more than one polypeptide chains are encoded by a single nucleic acid).

[0260] The nucleic acids of the disclosure can be DNA or RNA (*e.g.*, mRNA).

[0261] In another aspect, the disclosure provides host cells and vectors containing the nucleic acids of the disclosure. The nucleic acids may be present in a single vector or separate vectors present in the same host cell or separate host cell, as described in more detail herein below.

6.10.1. Vectors

[0262] The disclosure provides vectors comprising nucleotide sequences encoding an IFN receptor agonist or a component thereof described herein, for example one or two of the polypeptide chains of a half antibody of an IFN receptor agonist. The vectors include, but are not limited to, a virus, plasmid, cosmid, lambda phage or a yeast artificial chromosome (YAC).

[0263] Numerous vector systems can be employed. For example, one class of vectors utilizes DNA elements which are derived from animal viruses such as, for example, bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (Rous Sarcoma Virus, MMTV or MOMLV) or SV40 virus. Another class of vectors utilizes RNA elements derived from RNA viruses such as Semliki Forest virus, Eastern Equine Encephalitis virus and Flaviviruses.

[0264] Additionally, cells which have stably integrated the DNA into their chromosomes can be selected by introducing one or more markers which allow for the selection of transfected host cells. The marker may provide, for example, prototropy to an auxotrophic host, biocide resistance (*e.g.*, antibiotics), or resistance to heavy metals such as copper, or the like. The selectable marker gene can be either directly linked to the DNA sequences to be expressed or introduced into the same cell by co-transformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include splice signals, as well as transcriptional promoters, enhancers, and termination signals.

[0265] Once the expression vector or DNA sequence containing the constructs has been prepared for expression, the expression vectors can be transfected or introduced into an appropriate host cell. Various techniques may be employed to achieve this, such as, for example, protoplast fusion, calcium phosphate precipitation, electroporation, retroviral transduction, viral transfection, gene gun, lipid-based transfection or other conventional techniques. Methods and conditions for culturing the resulting transfected cells and for recovering the expressed polypeptides are known to those skilled in the art and may be varied or optimized depending upon the specific expression vector and mammalian host cell employed, based upon the present description.

6.10.2. Cells

[0266] The disclosure also provides host cells comprising a nucleic acid of the disclosure.

[0267] In one embodiment, the host cells are genetically engineered to comprise one or more nucleic acids described herein.

[0268] In one embodiment, the host cells are genetically engineered by using an expression cassette. The phrase "expression cassette," refers to nucleotide sequences, which are capable of affecting expression of a gene in hosts compatible with such sequences. Such cassettes may include a promoter, an open reading frame with or without introns, and a termination signal. Additional factors necessary or helpful in effecting expression may also be used, such as, for example, an inducible promoter.

[0269] The disclosure also provides host cells comprising the vectors described herein.

[0270] The cell can be, but is not limited to, a eukaryotic cell, a bacterial cell, an insect cell, or a human cell. Suitable eukaryotic cells include, but are not limited to, Vero cells, HeLa cells, COS cells, CHO cells, HEK293 cells, BHK cells and MDCKII cells. Suitable insect cells include, but are not limited to, Sf9 cells.

6.11. Pharmaceutical Compositions

[0271] The IFN receptor agonists of the disclosure may be in the form of compositions comprising the IFN receptor agonist and one or more carriers, excipients and/or diluents. The compositions may be formulated for specific uses, such as for veterinary uses or pharmaceutical uses in humans. The form of the composition (*e.g.*, dry powder, liquid formulation, *etc.*) and the excipients, diluents and/or carriers used will depend upon the

intended uses of the IFN receptor agonist and, for therapeutic uses, the mode of administration.

[0272] For therapeutic uses, the compositions may be supplied as part of a sterile, pharmaceutical composition that includes a pharmaceutically acceptable carrier. This composition can be in any suitable form (depending upon the desired method of administering it to a patient). The pharmaceutical composition can be administered to a patient by a variety of routes such as orally, transdermally, subcutaneously, intranasally, intravenously, intramuscularly, intratumorally, intrathecally, topically or locally. The most suitable route for administration in any given case will depend on the particular IFN receptor agonist, the subject, and the nature and severity of the disease and the physical condition of the subject. Typically, the pharmaceutical composition will be administered intravenously or subcutaneously.

[0273] Pharmaceutical compositions can be conveniently presented in unit dosage forms containing a predetermined amount of an IFN receptor agonist of the disclosure per dose. The quantity of IFN receptor agonist included in a unit dose will depend on the disease being treated, as well as other factors as are well known in the art. Such unit dosages may be in the form of a lyophilized dry powder containing an amount of IFN receptor agonist suitable for a single administration, or in the form of a liquid. Dry powder unit dosage forms may be packaged in a kit with a syringe, a suitable quantity of diluent and/or other components useful for administration. Unit dosages in liquid form may be conveniently supplied in the form of a syringe pre-filled with a quantity of IFN receptor agonist suitable for a single administration.

[0274] The pharmaceutical compositions may also be supplied in bulk from containing quantities of IFN receptor agonist suitable for multiple administrations.

[0275] Pharmaceutical compositions may be prepared for storage as lyophilized formulations or aqueous solutions by mixing an IFN receptor agonist having the desired degree of purity with optional pharmaceutically-acceptable carriers, excipients or stabilizers typically employed in the art (all of which are referred to herein as “carriers”), *i.e.*, buffering agents, stabilizing agents, preservatives, isotonifiers, non-ionic detergents, antioxidants, and other miscellaneous additives. See, Remington’s Pharmaceutical Sciences, 16th edition

(Osol, ed. 1980). Such additives should be nontoxic to the recipients at the dosages and concentrations employed.

[0276] Buffering agents help to maintain the pH in the range which approximates physiological conditions. They may be present at a wide variety of concentrations but will typically be present in concentrations ranging from about 2 mM to about 50 mM. Suitable buffering agents for use with the present disclosure include both organic and inorganic acids and salts thereof such as citrate buffers (*e.g.*, monosodium citrate-disodium citrate mixture, citric acid-trisodium citrate mixture, citric acid-monosodium citrate mixture, *etc.*), succinate buffers (*e.g.*, succinic acid-monosodium succinate mixture, succinic acid-sodium hydroxide mixture, succinic acid-disodium succinate mixture, *etc.*), tartrate buffers (*e.g.*, tartaric acid-sodium tartrate mixture, tartaric acid-potassium tartrate mixture, tartaric acid-sodium hydroxide mixture, *etc.*), fumarate buffers (*e.g.*, fumaric acid-monosodium fumarate mixture, fumaric acid-disodium fumarate mixture, monosodium fumarate-disodium fumarate mixture, *etc.*), gluconate buffers (*e.g.*, gluconic acid-sodium glyconate mixture, gluconic acid-sodium hydroxide mixture, gluconic acid-potassium glyconate mixture, *etc.*), oxalate buffer (*e.g.*, oxalic acid-sodium oxalate mixture, oxalic acid-sodium hydroxide mixture, oxalic acid-potassium oxalate mixture, *etc.*), lactate buffers (*e.g.*, lactic acid-sodium lactate mixture, lactic acid-sodium hydroxide mixture, lactic acid-potassium lactate mixture, *etc.*) and acetate buffers (*e.g.*, acetic acid-sodium acetate mixture, acetic acid-sodium hydroxide mixture, *etc.*). Additionally, phosphate buffers, histidine buffers and trimethylamine salts such as Tris can be used.

[0277] Preservatives may be added to retard microbial growth and can be added in amounts ranging from about 0.2%-1 % (w/v). Suitable preservatives for use with the present disclosure include phenol, benzyl alcohol, meta-cresol, methyl paraben, propyl paraben, octadecyldimethylbenzyl ammonium chloride, benzalconium halides (*e.g.*, chloride, bromide, and iodide), hexamethonium chloride, and alkyl parabens such as methyl or propyl paraben, catechol, resorcinol, cyclohexanol, and 3-pentanol. Isotonicifiers sometimes known as “stabilizers” can be added to ensure isotonicity of liquid compositions of the present disclosure and include polyhydric sugar alcohols, for example trihydric or higher sugar alcohols, such as glycerin, erythritol, arabitol, xylitol, sorbitol and mannitol. Stabilizers refer to a broad category of excipients which can range in function from a bulking agent to an

additive which solubilizes the therapeutic agent or helps to prevent denaturation or adherence to the container wall. Typical stabilizers can be polyhydric sugar alcohols (enumerated above); amino acids such as arginine, lysine, glycine, glutamine, asparagine, histidine, alanine, ornithine, L-leucine, 2-phenylalanine, glutamic acid, threonine, *etc.*, organic sugars or sugar alcohols, such as lactose, trehalose, stachyose, mannitol, sorbitol, xylitol, ribitol, myoinisitol, galactitol, glycerol and the like, including cyclitols such as inositol; polyethylene glycol; amino acid polymers; sulfur containing reducing agents, such as urea, glutathione, thioctic acid, sodium thioglycolate, thioglycerol, a-monothioglycerol and sodium thio sulfate; low molecular weight polypeptides (*e.g.*, peptides of 10 residues or fewer); proteins such as human serum albumin, bovine serum albumin, gelatin or immunoglobulins; hydrophylic polymers, such as polyvinylpyrrolidone monosaccharides, such as xylose, mannose, fructose, glucose; disaccharides such as lactose, maltose, sucrose and trehalose; and trisaccacharides such as raffinose; and polysaccharides such as dextran. Stabilizers may be present in amounts ranging from 0.5 to 10 wt % per wt of IFN receptor agonist.

[0278] Non-ionic surfactants or detergents (also known as "wetting agents") may be added to help solubilize the glycoprotein as well as to protect the glycoprotein against agitation-induced aggregation, which also permits the formulation to be exposed to shear surface stressed without causing denaturation of the protein. Suitable non-ionic surfactants include polysorbates (20, 80, *etc.*), polyoxamers (184, 188, *etc.*), and pluronic polyols. Non-ionic surfactants may be present in a range of about 0.05 mg/mL to about 1.0 mg/mL, for example about 0.07 mg/mL to about 0.2 mg/mL.

[0279] Additional miscellaneous excipients include bulking agents (*e.g.*, starch), chelating agents (*e.g.*, EDTA), antioxidants (*e.g.*, ascorbic acid, methionine, vitamin E), and cosolvents.

[0280] The IFN receptor agonists of the disclosure can be formulated as pharmaceutical compositions comprising the IFN receptor agonists, for example containing one or more pharmaceutically acceptable excipients or carriers. To prepare pharmaceutical or sterile compositions comprising the IFN receptor agonists of the present disclosure, a IFN receptor agonist preparation can be combined with one or more pharmaceutically acceptable excipient or carrier.

[0281] For example, formulations of IFN receptor agonists can be prepared by mixing IFN receptor agonists with physiologically acceptable carriers, excipients, or stabilizers in the form of, *e.g.*, lyophilized powders, slurries, aqueous solutions, lotions, or suspensions (see, *e.g.*, Hardman *et al.,* 2001 , Goodman and Gilman's The Pharmacological Basis of Therapeutics, McGraw-Hill, New York, N.Y.; Gennaro, 2000, Remington: The Science and Practice of Pharmacy, Lippincott, Williams, and Wilkins, New York, N.Y.; Avis, *et al.* (eds.),1993, Pharmaceutical Dosage Forms: General Medications, Marcel Dekker, NY; Lieberman, *et al.* (eds.), 1990, Pharmaceutical Dosage Forms: Tablets, Marcel Dekker, NY; Lieberman, *et al.* (eds.), 1990, Pharmaceutical Dosage Forms: Disperse Systems, Marcel Dekker, NY; Weiner and Kotkoskie, 2000, Excipient Toxicity and Safety, Marcel Dekker, Inc., New York, N.Y.).

[0282] An effective amount for a particular subject may vary depending on factors such as the condition being treated, the overall health of the subject, the method route and dose of administration and the severity of side effects (see, *e.g.*, Maynard, *et al.* (1996) A Handbook of SOPs for Good Clinical Practice, Interpharm Press, Boca Raton, Fla.; Dent (2001) Good Laboratory and Good Clinical Practice, Urch Publ., London, UK).

[0283] A composition of the present disclosure may also be administered via one or more routes of administration using one or more of a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Selected routes of administration for IFN receptor agonists include intravenous, intramuscular, intradermal, intraperitoneal, subcutaneous, spinal or other general routes of administration, for example by injection or infusion. General administration may represent modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion. Alternatively, a composition of the disclosure can be administered via a non-general route, such as a topical, epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically. In one embodiment, the IFN receptor agonists

are administered by infusion. In another embodiment, the IFN receptor agonist of the disclosure is administered subcutaneously.

6.11.1. Pharmaceutical Compositions for Delivery of IFN receptor Agonist Encoding Nucleic Acids

[0284] An IFN receptor agonist of the disclosure can be delivered by any method useful for gene therapy, for example as mRNA or through viral vectors encoding the IFN receptor agonist under the control of a suitable promoter.

[0285] Exemplary viral vectors include recombinant adenovirus and adeno-associated virus vectors (rAAV). rAAV vectors are based on the defective and nonpathogenic parvovirus adeno-associated type 2 virus. Most such vectors are derived from a plasmid that retains only the AAV inverted terminal repeats flanking the transgene expression cassette. Efficient gene transfer and stable transgene delivery due to integration into the genomes of the transduced cell are key features for this vector system. AAV serotypes useful for delivering IL27 transgenes AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV8, AAV 8.2, AAV9, and AAV rh10 and pseudotyped AAV such as AAV2/8, AAV2/5 and AAV2/6.

[0286] AAV may be manufactured at a clinical scale by a number of different processes. Examples of systems that can be used include (1) plasmid DNA transfection in mammalian cells, (2) Ad infection of stable mammalian cell lines, (3) infection of mammalian cells with recombinant herpes simplex viruses (rHSVs), and (4) infection of insect cells (Sf9 cells) with recombinant baculoviruses (reviewed by Penaud-Budloo *et al.*, 2018, Mol Ther Methods Clin Dev. 8: 166-180).

[0287] Replication-deficient recombinant adenoviral vectors (Ad) can be produced at high titer and readily infect a number of different cell types. Most adenovirus vectors are engineered such that a transgene replaces the Ad Ela, Elb, and/or E3 genes; subsequently the replication defective vector is propagated in human 293 cells that supply deleted gene function in trans. Ad vectors can transduce multiple types of tissues in vivo, including non-dividing, differentiated cells such as those found in liver, kidney and muscle. Conventional Ad vectors have a large carrying capacity.

[0288] Packaging cells are used to form virus particles that are capable of infecting a host cell. Such cells include 293 cells, which package adenovirus, and w2 cells or PA317 cells, which package retrovirus. Viral vectors used in gene therapy are usually generated by a

producer cell line that packages a nucleic acid vector into a viral particle. The vectors typically contain the minimal viral sequences required for packaging and subsequent integration into a host (if applicable), other viral sequences being replaced by an expression cassette encoding the protein to be expressed. The missing viral functions are supplied in trans by the packaging cell line. For example, AAV vectors used in gene therapy typically only possess inverted terminal repeat (ITR) sequences from the AAV genome which are required for packaging and integration into the host genome. Viral DNA is packaged in a cell line, which contains a helper plasmid encoding the other AAV genes, namely rep and cap, but lacking ITR sequences. The cell line is also infected with adenovirus as a helper. The helper virus promotes replication of the AAV vector and expression of AAV genes from the helper plasmid. The helper plasmid is not packaged in significant amounts due to a lack of ITR sequences. Contamination with adenovirus can be reduced by, e.g., heat treatment to which adenovirus is more sensitive than AAV.

[0289] The nucleic acid molecule (e.g., mRNA) or virus can be formulated as the sole pharmaceutically active ingredient in a pharmaceutical composition or can be combined with other active agents for the particular disorder treated. Optionally, other medicinal agents, pharmaceutical agents, carriers, adjuvants, diluents can be included in the compositions provided herein. For example, any one or more of a wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives, antioxidants, chelating agents and inert gases also can be present in the compositions. Exemplary other agents and excipients that can be included in the compositions include, for example, water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite; oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, α-tocopherol; and metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid and phosphoric acid.

6.12. Therapeutic Indications and Methods of Use

[0290] The present disclosure provides methods for using and applications for the IFN receptor agonists of the disclosure.

[0291] The IFN receptor agonists of the disclosure can be used to stimulate the immune response in a variety of applications.

[0292] In certain aspects, the disclosure provides a method of treating cancer, comprising administering to a subject in need thereof an IFN receptor agonist or pharmaceutical composition as described herein. In some embodiments, when an IFN receptor agonist comprises one or more protease-cleavable linkers, an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases expressed by the cancer tissue. Accordingly, in some embodiments, the IFN receptor agonist is selectively activated in the cancer tissue.

[0293] In some embodiments, the disclosure provides a method of treating cancer with an IFN protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IFN receptor agonist or pharmaceutical composition comprising one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by cancer tissue to which the IFN protein is intended. Thus, an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the cancer tissue.

[0294] The present disclosure further provides a method of localized delivery of an IFN protein, comprising administering to a subject an IFN receptor agonist or pharmaceutical composition as described herein, where the IFN receptor agonist has one or more targeting moieties and/or protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue to which the IFN protein is to be locally delivered. As used herein, the term “locally delivered” does not require local administration but rather indicates that the active component of the IFN receptor agonist refers to selective targeting with a targeting moiety that recognize a target molecule expressed in the intended site and/or activation of the protein by a protease active at the intended site.

[0295] The present disclosure further provides a method of administering to the subject IFN therapy with reduced systemic exposure and/or reduced systemic toxicity, comprising administering to a subject the IFN therapy in the form of an IFN receptor agonist or pharmaceutical composition as described herein, where the IFN receptor agonist has one or more targeting moieties that bind to a target molecule expressed by a tissue for which IFN

therapy is desirable and/or intended, and/or protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IFN therapy is desirable and/or intended.

[0296] Accordingly, the foregoing methods permit IFN therapy with reduced off-target side effects by virtue of preferential delivery and/or activation of an IFN receptor agonist at a locale intended for IFN treatment.

[0297] Accordingly, the present disclosure provides a method of targeted delivery of an activated IFN protein to a locale intended for treatment, *e.g.*, cancer tissue, comprising administering to a subject an IFN receptor agonist or pharmaceutical composition as described herein, wherein the IFN comprises one or more targeting moieties that recognize a target molecule expressed in the locale or by the tissue intended for treatment (*e.g.*, cancer tissue) and which optionally has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IFN therapy is desirable and/or intended.

[0298] The present disclosure further provides method of locally inducing an immune response in a target tissue, comprising administering to a subject IFN receptor agonist or pharmaceutical composition as described herein which has one or more targeting moieties capable of binding a target molecule expressed in the target tissue and optionally one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in the target tissue. When a protease-activated linker is present, an activated IFN protein comprising the IFN moiety can then be produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the target tissue. The resulting activated IFN protein can then induce the immune response against at least one cell type in the target tissue. In the absence of activation, an IFN protein with attenuated activity (*e.g.*, by virtue of masking) can induce the immune response against at least one cell type in the target tissue.

[0299] In some embodiments, the administration is not local to the tissue. For example, when the target tissue is cancer tissue, the administration can be systemic or subcutaneous.

[0300] The IFN receptor agonists of the disclosure can be used in the treatment of any proliferative disorder (*e.g.*, cancer) that expresses a target molecule (either on the tumor cells or in the tumor microenvironment, *e.g.*, the extracellular matrix or the tumor

lymphocytes). In particular embodiments, the cancer is acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), adrenocortical carcinoma, anal cancer, appendix cancer, astrocytoma, basal cell carcinoma, brain tumor, bile duct cancer, bladder cancer, bone cancer, breast cancer, bronchial tumor, Burkitt Lymphoma, carcinoma of unknown primary origin, cardiac tumor, cervical cancer, chordoma, chronic lymphocytic leukemia (CLL), chronic myelogenous leukemia (CML), chronic myeloproliferative neoplasm, colon cancer, colorectal cancer, craniopharyngioma, cutaneous T-cell lymphoma, ductal carcinoma, embryonal tumor, endometrial cancer, ependymoma, esophageal cancer, esthesioneuroblastoma, fibrous histiocytoma, Ewing sarcoma, eye cancer, germ cell tumor, gallbladder cancer, gastric cancer, gastrointestinal carcinoid tumor, gastrointestinal stromal tumor, gestational trophoblastic disease, glioma, head and neck cancer, hairy cell leukemia, hepatocellular cancer, histiocytosis, Hodgkin lymphoma, hypopharyngeal cancer, intraocular melanoma, islet cell tumor, Kaposi sarcoma, kidney cancer, Langerhans cell histiocytosis, laryngeal cancer, leukemia, lip and oral cavity cancer, liver cancer, lobular carcinoma in situ, lung cancer, lymphoma, macroglobulinemia, malignant fibrous histiocytoma, melanoma, Merkel cell carcinoma, mesothelioma, metastatic squamous neck cancer with occult primary, midline tract carcinoma involving NUT gene, mouth cancer, multiple endocrine neoplasia syndrome, multiple myeloma, mycosis fungoides, myelodysplastic syndrome, myelodysplastic/myeloproliferative neoplasm, nasal cavity and para-nasal sinus cancer, nasopharyngeal cancer, neuroblastoma, non-Hodgkin lymphoma, non-small cell lung cancer, oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, papillomatosis, paraganglioma, parathyroid cancer, penile cancer, pharyngeal cancer, pheochromocytomas, pituitary tumor, pleuropulmonary blastoma, primary central nervous system lymphoma, prostate cancer, rectal cancer, renal cell cancer, renal pelvis and ureter cancer, retinoblastoma, rhabdoid tumor, salivary gland cancer, Sezary syndrome, skin cancer, small cell lung cancer, small intestine cancer, soft tissue sarcoma, spinal cord tumor, stomach cancer, T-cell lymphoma, teratoid tumor, testicular cancer, throat cancer, thymoma and thymic carcinoma, thyroid cancer, urethral cancer, uterine cancer, vaginal cancer, vulvar cancer, or Wilms tumor.

[0301] Table I below shows exemplary indications for which IFN receptor agonists targeting particular target molecules can be used.

TABLE I Examples of Target Molecule Indications	
Target	**Exemplary Indication(s)**
ADRB3	Ewing sarcoma
ALK	NSCLC, ALCL, IMT, neuroblastoma
B7H3	melanoma, osteosarcoma, leukemia, breast, prostate, ovarian, pancreatic, colorectal cancers
BCMA	multiple myeloma, leukemia (*e.g.*, acute lymphoblastic leukemia ("ALL"), acute myeloid leukemia ("AML"), chronic lymphocytic leukemia ("CLL"), chronic myeloid leukemia ("CML") and hairy cell leukemia ("HCL")); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, including diffuse large B-cell lymphoma ("DLBCL"))
Cadherin 17	gastric, pancreatic, and colorectal adenocarcinomas
CAIX	clear-cell renal cell carcinoma, hypoxic solid tumors, head and neck squamous carcinoma
CD123	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma. In a preferred embodiment, the indication is AML.
CD171	neuroblastoma, paraganglioma
CD179a	B cell malignancies
CD19	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma.
CD20	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma.
CD22	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma; lung cancer
CD24	ovarian, breast, prostate, bladder, renal, non-small cell carcinomas
CD30	anaplastic large cell lymphoma, embryonal carcinoma, Hodgkin Lymphoma
CD32b	B cell malignancies, gastric, pancreatic, esophageal, glioblastoma, breast, colorectal
CD33	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma. In a preferred embodiment, the indication is AML.
CD38	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma
CD44v6	colon cancer, head and neck small cell carcinoma
CD97	B cell malignancies, gastric, pancreatic, esophageal, glioblastoma, breast, colorectal
CEA (CEACAM5)	colorectal carcinoma, gastric carcinoma, pancreatic carcinoma, lung cancer, breast cancer, medullary thyroid carcinoma
CLDN6	ovarian, breast, lung cancer
CLL-1	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL); lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL); multiple myeloma. In a preferred embodiment, the indication is AML.
CS1 (SLAMF7)	multiple myeloma

TABLE I Examples of Target Molecule Indications	
Target	**Exemplary Indication(s)**
EGFR	squamous cell carcinoma of lung, anal cancer, glioblastoma, epithelial tumors of head and neck, colon cancer
EGFRvIII	Glioblastoma
EPCAM	gastrointestestinal carcinoma, colorectal cancer
EphA2	kaposi's sarcoma, glioblastoma, solid tumors, glioma
Ephrin B2	thyroid cancer, breast cancer, malignant melanoma
ERBB2 (Her2/neu)	breast, ovarian, gastric cancers, lung adenocarcinoma, non-small cell lung cancer, uterine cancer, uterine serous endometrial carcinoma, salivary duct carcinoma
FAP	pancreatic cancer, colorectal cancer, metastasis, epithelial cancers, soft tissue sarcomas
FCRL5	multiple myeloma
FLT3	leukemia (*e.g.*, ALL, CLL, AML, CML, HCL), lymphoma (*e.g.*, Hodgkin's lymphoma, non-Hodgkin's lymphoma, *e.g.*, DLBCL), multiple myeloma
Folate receptor alpha	ovarian, breast, renal, lung, colorectal, brain cancers
Folate receptor beta	ovarian cancer
Fucosyl GM1	AML, myeloma
GD2	malignant melanoma, neuroblastoma
GD3	Melanoma
GloboH	ovarian, gastric, prostate, lung, breast, and pancreatic cancers
gp100	Melanoma
GPNMB	breast cancer, head and neck cancers
GPR20	GIST
GPR64	Ewing sarcoma, prostate, kidney and lung sarcomas
GPRC5D	multiple myeloma
HAVCR1	renal cancer
HER2	HER-2 (+) adenocarcinoma of gastroesophageal junction, HER-2 positive gastric adenocarcinoma, HER2 positive carcinoma of breast
HER3	colon and gastric cancers
HMWMAA	melanoma, glioblastoma, breast cancer
IGF-I receptor	breast, prostate, lung cancers
IL11Rα	papillary thyroid cancer, osteosarcoma, colorectal adenocarcinoma, lymphocytic leukemia
IL13Rα2	renal cell carcinoma, prostate cancer, gliomas, head and neck cancer, astrocytoma
KIT	myeloid leukemia, Kaposi's sarcoma, erythroleukemia, gastrointestinal stromal tumors
KLRG2	breast cancers, lung cancers and ovarian cancers.
LewisY	squamous cell lung carcinoma, lung adenocarcinoma, ovarian carcinoma, and colorectal adenocarcinoma
LMP2	prostate cancer, Hodgkin's lymphoma, nasopharyngeal carcinoma
LRP6	breast cancer

TABLE I Examples of Target Molecule Indications	
Target	**Exemplary Indication(s)**
LY6K	breast, lung, ovarian, and cervical cancer
LYPD8	colorectal and gastric cancers
Mesothelin	mesothelioma, pancreatic cancer, ovarian cancer, stomach cancer, lung cancer, endometrial cancer
MUC1	breast and ovarian cancers, lung, stomach, pancreatic, prostate cancers
NCAM	melanoma, Wilms' tumor, small cell lung cancer, neuroblastoma, myeloma, paraganglioma, pancreatic acinar cell carcinoma, myeloid leukemia
NY-BR-1	breast cancer
o-acetyl GD2	neuroblastoma, melanoma
OR51E2	prostate cancer
PANX3	Osteosarcoma
PLAC1	hepatocellular carcinoma
Polysialic acid	small cell lung cancer
PDGFR-beta	myelomonocytic leukemia, chronic myeloid leukemia, acute myelogenous leukemia, acute lymphoblastic leukemia
PRSS21	colon cancer, testicular cancer, ovarian cancer
PSCA	prostate cancer, gastric and bladder cancers
PSMA	prostate cancer
ROR1	metastatic cancers, chronic lymphocytic leukemia, solid tumors in lung, breast, ovarian, colon, pancreatic, sarcoma
SLC34A2	bladder cancer
SLC39A6	breast cancer, esophageal cancer
SLITRK6	breast cancer, urothelial cancer, lung cancer
SSEA-4	breast cancer, cancer stem cells, epithelial ovarian carcinoma
STEAP1	prostate cancer
STEAP2	prostate cancer (including castrate-resistant prostate cancer), bladder cancer, cervical cancer, lung cancer, colon cancer, kidney cancer, breast cancer, pancreatic cancer, stomach cancer, uterine cancer, ovarian cancer, preferably prostate cancer
TACSTD2	carcinomas, *e.g.*, non-small-cell lung cancer
TAG72	ovarian, breast, colon, lung, pancreatic cancers, gastric cancer
TEM1/CD248	colorectal cancer
TEM7R	colorectal cancer
Tn	colorectal, breast cancers, cervical, lung, stomach cancers
TSHR	thyroid cancer, multiple myeloma
Tyrosinase	prostate cancer, melanoma
UPK2	bladder cancer
VEGFR2	ovarian and pancreatic cancers, renal cell carcinoma, colorectal cancer, medullary thyroid carcinoma

[0302] Additional target molecules and corresponding indications are disclosed in, *e.g.*, Hafeez *et al.*, 2020, Molecules 25:4764, doi:10.3390/molecules25204764, particularly in Table 1. Table 1 is incorporated by reference in its entirety here.

[0303] In further embodiments, the IFN receptor agonists can be used to enhance an immune response elicited by another agent. Thus, in some embodiments an IFN receptor agonist of the disclosure is administered as an adjunct therapy with an immunogenic agent. In some embodiments, the immunogenic agent is an adjuvanted or unadjuvanted vaccine. The IFN receptor agonists can thus enhance an antigen-specific immune response elicited by the vaccine. In various embodiments, the vaccine is a prophylactic or therapeutic cancer vaccine or a prophylactic or therapeutic vaccine against an infectious agent, *e.g.*, a virus, bacteria, or parasite.

7. NUMBERED EMBODIMENTS

[0304] While various specific embodiments have been illustrated and described, it will be appreciated that various changes can be made without departing from the spirit and scope of the disclosure(s). The present disclosure is exemplified by the numbered embodiments set forth below.

[0305] In the numbered embodiments that follow, the targeting moiety preferably binds to a mammalian target molecule, the IFN moiety is preferably derived from a mammalian IFN, the Fc domains are preferably derived from a mammalian antibody, and the subjects are preferably mammals. More preferably, the mammal is human.

1. A Type I interferon (IFN) receptor agonist, comprising a

(a) a first polypeptide chain comprising a first Fc domain;

(b) a second polypeptide chain comprising a second Fc domain associated with the first Fc domain;

wherein the first polypeptide chain and/or the second polypeptide chain comprises a Type I interferon (IFN) moiety that is attenuated, optionally by (i) masking by a Type I interferon receptor (IFNR) moiety; (ii) one or more mutations in the IFN moiety as compared to wild-type interferon, *e.g.*, by one or more amino acid substitutions and/or truncations; (iii)

use of native IFN sequences with a lower receptor affinity than IFNα2b and/or IFNβ or (iv) any combination of two or all three of (i), (ii) and (iii).

2. The IFN receptor agonist of embodiment 1, which comprises an IFNR moiety on the same polypeptide chain as the IFN moiety.

3. The IFN receptor agonist of embodiment 1, which comprises an IFNR moiety on a different polypeptide chain than the IFN moiety.

4. The IFN receptor agonist of any one of embodiments 1 to 3, wherein the first polypeptide chain comprises a first IFN moiety and the second polypeptide chain comprises a second IFN moiety, each masked by an IFNR moiety.

5. The IFN receptor agonist of any one of embodiments 1 to 4, which is bivalent for the IFN moiety.

6. The IFN receptor agonist of embodiment 5, wherein each IFN moiety is singly masked.

7. The IFN receptor agonist of embodiment 5, wherein each IFN moiety is doubly masked.

8. The IFN receptor agonist of any one of embodiments 4 to 7, wherein the first IFN moiety is N-terminal to the first Fc domain and the second IFN moiety is N-terminal to the second Fc domain.

9. The IFN receptor agonist of any one of embodiments 4 to 7, wherein the first IFN moiety is C-terminal to the first Fc domain and the second IFN moiety is C-terminal to the second Fc domain.

10. The IFN receptor agonist of any one of embodiments 1 to 3, wherein either the first polypeptide chain or the second polypeptide chain comprises an IFN moiety, masked by an IFNR moiety.

11. The IFN receptor agonist of any one of embodiments 1 to 3 and 10, which is monovalent for the IFN moiety.

12. The IFN receptor agonist of embodiment 11, wherein the IFN moiety is singly masked.

13. The IFN receptor agonist of embodiment 11, wherein the IFN moiety is doubly masked.

14. The IFN receptor agonist of any one of embodiments 10 to 13, wherein the IFN moiety is N-terminal to the first Fc domain or N-terminal to the second Fc domain.

15. The IFN receptor agonist of any one of embodiments 10 to 13, wherein the IFN moiety is C-terminal to the first Fc domain or C-terminal to the second Fc domain.

16. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 15, which comprises any of half antibody pairs designated 1-23 as set forth in Table 2.

17. The IFN receptor agonist of any one of embodiments 1 to 16, which does not comprise a protease-cleavable linker (PCL).

18. The IFN receptor agonist of any one of embodiments 1 to 16, which comprises at least one protease-cleavable linker (PCL).

19. The IFN receptor agonist of embodiment 18, wherein the PCL comprises a substrate sequence cleavable by any protease set forth in Table A.

20. The IFN receptor agonist of embodiment 18 or embodiment 19, wherein the PCL comprises one or more substrate sequences selected from the substrate sequences set forth in Table B.

21. The IFN receptor agonist of any one of embodiments 18 to 20, wherein the PCL comprises one or more spacer sequences selected from the substrate sequences set forth in Table C.

22. The IFN receptor agonist of any one of embodiments 18 to 21, wherein the PCL comprises the amino acid sequence of any of the PCL sequences set forth in Table D or a variant thereof with up to 5 amino acid substitutions, *e.g.*, a variant thereof with 1 amino acid substitution, 2 amino acid substitutions, 3 amino acid substitutions, 4 amino acid substitutions, or 5 amino acid substitutions.

23. The IFN receptor agonist of any one of embodiments 18 to 22, which is configured such that cleavage of the protease-cleavable linker (PCL) unmasks the IFN moiety.

24. The IFN receptor agonist of embodiment 23, wherein the unmasked IFN moiety is in the form of an IFN polypeptide comprising an Fc domain.

25. The IFN receptor agonist of embodiment 23, wherein the unmasked IFN moiety is in the form of an IFN polypeptide lacking an Fc domain.

26. The IFN receptor agonist of any one of embodiments 1 to 25, wherein each IFN moiety comprises an amino acid sequence having at least about 90% sequence identity to (a) full length mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ or (b) a mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ having up to a 15-amino acid truncation at its N-terminus and/or its C-terminus.

27. The IFN receptor agonist of any one of embodiments 1 to 25, wherein each IFN moiety comprises an amino acid sequence having about 95% sequence identity to (a) full length mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ or (b) a mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ having up to a 15-amino acid truncation at its N-terminus and/or its C-terminus.

28. The IFN receptor agonist of any one of embodiments 1 to 25, wherein each IFN moiety comprises an amino acid sequence having about 98% sequence identity to (a) full length mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ or (b) a mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ having up to a 15-amino acid truncation at its N-terminus and/or its C-terminus.

29. The IFN receptor agonist of any one of embodiments 1 to 28, wherein each IFN moiety comprises an amino acid sequence having one or more attenuating mutations as compared to mature human IFNα1 or IFNα2b.

30. The IFN receptor agonist of any one of embodiments 1 to 28, which has one or more mutations selected from L26A, F27A, R33A, R33K, L30A, D35E, H57Y, E58N, Q61S, H57S, E58S, H57A, E58A, Q61A, Q90A, E96A, R120A, L135A, R144A, R144S, R144T, R144Y, R144I, R144L, A145D, A145H, A145K, A145M, A145V, A145Y, R149A, R149K, S152A, R162A, and E165D.

31. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution R33A.

32. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution R33K.

33. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution Q90A.

34. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution E96A.

35. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution R120A.

36. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution A145M.

37. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution R149A.

38. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution R149K.

39. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitution S152A.

40. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitutions R33A, H57Y, E58N and Q61S.

41. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitutions H57Y, E58N, Q61S and R144A.

42. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitutions A145M and R149K.

43. The IFN receptor agonist of any one of embodiments 1 to 29, which comprises the amino acid substitutions Q90A and R120A.

44. The IFN receptor agonist of any one of embodiments 1 to 43, wherein the IFNR moiety is an interferon alpha receptor (IFNAR) moiety.

45. The IFN receptor agonist of embodiment 44, wherein the IFNAR moiety is an IFNAR1 moiety.

46. The IFN receptor agonist of embodiment 45, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to (i) the SD2 and

SD3 domains of human IFNAR1, (ii) the SD1, SD2 and SD3 domains of human IFNAR1, or (iii) the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

47. The IFN receptor agonist of embodiment 46, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to the SD2 and SD3 domains of human IFNAR1.

48. The IFN receptor agonist of embodiment 46, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to the SD1, SD2 and SD3 domains of human IFNAR1.

49. The IFN receptor agonist of embodiment 46, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

50. The IFN receptor agonist of embodiment 45, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to (i) the SD2 and SD3 domains of human IFNAR1, (ii) the SD1, SD2 and SD3 domains of human IFNAR1, or (iii) the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

51. The IFN receptor agonist of embodiment 50, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to the SD2 and SD3 domains of human IFNAR1.

52. The IFN receptor agonist of embodiment 50, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to the SD1, SD2 and SD3 domains of human IFNAR1.

53. The IFN receptor agonist of embodiment 50, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

54. The IFN receptor agonist of embodiment 45, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to (i) the SD2 and SD3 domains of human IFNAR1, (ii) the SD1, SD2 and SD3 domains of human IFNAR1, or (iii) the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

55. The IFN receptor agonist of embodiment 54, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to the SD2 and SD3 domains of human IFNAR1.

56. The IFN receptor agonist of embodiment 54, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to the SD1, SD2 and SD3 domains of human IFNAR1.

57. The IFN receptor agonist of embodiment 54, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

58. The IFN receptor agonist of embodiment 44, wherein the IFNAR moiety is an IFNAR2 moiety.

59. The IFN receptor agonist of embodiment 58, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90% sequence identity to (i) the D1 domain of human IFNAR2 or (ii) the D1 and D2 domains of human IFNAR2.

60. The IFN receptor agonist of embodiment 59, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90% sequence identity to the D1 domain of human IFNAR2.

61. The IFN receptor agonist of embodiment 59, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90% sequence identity to the D1 and D2 domains of human IFNAR2.

62. The IFN receptor agonist of embodiment 58, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 95% sequence identity to (i) the D1 domain of human IFNAR2 or (ii) the D1 and D2 domains of human IFNAR2.

63. The IFN receptor agonist of embodiment 62, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 95% sequence identity to the D1 domain of human IFNAR2.

64. The IFN receptor agonist of embodiment 62, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 95% sequence identity to the D1 and D2 domains of human IFNAR2.

65. The IFN receptor agonist of embodiment 58, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 98% sequence identity to (i) the D1 domain of human IFNAR2 or (ii) the D1 and D2 domains of human IFNAR2.

66. The IFN receptor agonist of embodiment 65, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 98% sequence identity to the D1 domain of human IFNAR2.

67. The IFN receptor agonist of embodiment 65, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 98% sequence identity to the D1 and D2 domains of human IFNAR2.

68. The IFN receptor agonist of any one of embodiments 1 to 67, wherein the IFN moiety is attenuated by masking by an IFNAR1 moiety and an IFNAR2 moiety.

69. The IFN receptor agonist of claim 68, which is monovalent for the IFN moiety, the IFNAR1 moiety, and the IFNAR2 moiety.

70. The IFN receptor agonist of claim 68, which is bivalent for the IFN moiety, the IFNAR1 moiety, and the IFNAR2 moiety.

71. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to the SD2 and SD3 domains of human IFNAR1.

72. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to the SD2 and SD3 domains of human IFNAR1.

73. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to the SD2 and SD3 domains of human IFNAR1.

74. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises the amino acid sequence of the SD2 and SD3 domains of human IFNAR1.

75. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to the SD1, SD2, and SD3 domains of human IFNAR1.

76. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to the SD1, SD2, and SD3 domains of human IFNAR1.

77. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to the SD1, SD2, and SD3 domains of human IFNAR1.

78. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises the amino acid sequence of the SD1, SD2, and SD3 domains of human IFNAR1.

79. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90% sequence identity to the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

80. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 95% sequence identity to the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

81. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 98% sequence identity to the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

82. The IFN receptor agonist of any one of embodiments 68 to 70, wherein the IFNAR1 moiety comprises the amino acid sequence of the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

83. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90% sequence identity to the D1 domain of human IFNAR2.

84. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 95% sequence identity to the D1 domain of human IFNAR2.

85. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 98% sequence identity to the D1 domain of human IFNAR2.

86. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises the amino acid sequence of the D1 domain of human IFNAR2.

87. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90% sequence identity to the D1 and D2 domains of human IFNAR2.

88. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 95% sequence identity to the D1 and D2 domains of human IFNAR2.

89. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 98% sequence identity to the D1 and D2 domains of human IFNAR2.

90. The IFN receptor agonist of any one of embodiments 68 to 82, wherein the IFNAR2 moiety comprises the amino acid sequence of the D1 and D2 domains of human IFNAR2.

91. The IFN receptor agonist of any one of embodiments 68 to 90, wherein the IFN moiety has one or more mutations selected from L26A, F27A, R33A, R33K, L30A, D35E, H57Y, E58N, Q61S, H57S, E58S, H57A, E58A, Q61A, Q90A, E96A, R120A, L135A, R144A, R144S, R144T, R144Y, R144I, R144L, A145D, A145H, A145K, A145M, A145V, A145Y, R149A, R149K, S152A, R162A, and E165D, optionally wherein:

(a) the IFN moiety comprises the amino acid substitution R33A;

(b) the IFN moiety comprises the amino acid substitution R33K;

(c) the IFN moiety comprises the amino acid substitution Q90A;

(d) the IFN moiety comprises the amino acid substitution E96A;

(e) the IFN moiety comprises the amino acid substitution R120A;

(f) the IFN moiety comprises the amino acid substitution A145M;

(g) the IFN moiety comprises the amino acid substitution R149A;

(h) the IFN moiety comprises the amino acid substitution R149K;

(i) the IFN moiety comprises the amino acid substitution S152A;

(j) the IFN moiety comprises the amino acid substitutions R33A, H57Y, E58N and Q61S;

(k) the IFN moiety comprises the amino acid substitutions H57Y, E58N, Q61S and R144A;

(l) the IFN moiety comprises the amino acid substitutions A145M and R149K; or

(m) the IFN moiety comprises the amino acid substitutions Q90A and R120A.

92. The IFN receptor agonist of any one of embodiments 68 to 91, wherein the IFN moiety and the IFNAR1 moiety are on the same polypeptide chain.

93. The IFN receptor agonist of any one of embodiments 68 to 91, wherein the IFN moiety and the IFNAR2 moiety are on the same polypeptide chain.

94. The IFN receptor agonist of any one of embodiments 68 to 91, wherein the IFN moiety and the IFNAR1 moiety are on different polypeptide chains.

95. The IFN receptor agonist of any one of embodiments 68 to 91, wherein the IFN moiety and the IFNAR2 moiety are on different polypeptide chains.

96. The IFN receptor agonist of any one of embodiments 68 to 91, wherein the IFN moiety, the IFNAR1 moiety, and the IFNAR2 moiety are on the same polypeptide chain.

97. The IFN receptor agonist of any one of embodiments 68 to 96, wherein the first polypeptide chain comprises the IFN moiety.

98. The IFN receptor agonist of embodiment 97, wherein the IFN moiety is N-terminal to the first Fc domain.

99. The IFN receptor agonist of embodiment 97, wherein the IFN moiety is C-terminal to the first Fc domain.

100. The IFN receptor agonist of any one of embodiments 97 to 99, wherein the first polypeptide chain comprises the IFNAR1 moiety.

101. The IFN receptor agonist of embodiment 100, wherein the IFNAR1 moiety is N-terminal to the IFN moiety.

102. The IFN receptor agonist of embodiment 100, wherein the IFNAR1 moiety is C-terminal to the IFN moiety.

103. The IFN receptor agonist of any one of embodiments 97 to 102, wherein the first polypeptide chain comprises the IFNAR2 moiety.

104. The IFN receptor agonist of embodiment 103, wherein the IFNAR2 moiety is N-terminal to the IFN moiety.

105. The IFN receptor agonist of embodiment 103, wherein the IFNAR2 moiety is C-terminal to the IFN moiety.

106. The IFN receptor agonist of any one of embodiments 97 to 105, further comprising one or more linkers connecting two or more of the first Fc domain, the IFN moiety, the IFNAR1 moiety, and the IFNAR2 moiety.

107. The IFN receptor agonist of embodiment 97, wherein the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR1 moiety, the IFN moiety, and the IFNAR2 moiety.

108. The IFN receptor agonist of embodiment 107, further comprising a first linker connecting the first Fc domain and the first IFNAR1 moiety, a second linker connecting the IFNAR1 moiety and the IFN moiety, and a third linker connecting the IFN moiety and the IFNAR2 moiety.

109. The IFN receptor agonist of embodiment 97, wherein the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, a first linker, the IFNAR2 moiety, a second linker, the IFN moiety, a third linker, and the IFNAR1 moiety.

110. The IFN receptor agonist of embodiment 109, further comprising a first linker connecting the first Fc domain and the first IFNAR2 moiety, a second linker connecting the IFNAR2 moiety and the IFN moiety, and a third linker connecting the IFN moiety and the IFNAR1 moiety.

111. The IFN receptor agonist of embodiment 97, wherein (i) the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR2 moiety, and the IFN moiety, and (ii) the second polypeptide comprises, in N- to C-terminal orientation, the second Fc domain, and the IFNAR1 moiety.

112. The IFN receptor agonist of embodiment 111, further comprising a first linker connecting the first Fc domain and the IFNAR2 moiety, a second linker connecting the IFNAR2 moiety and the IFN moiety, and a third linker connecting the second Fc domain and the IFNAR1 moiety.

113. The IFN receptor agonist of embodiment 97, wherein (i) the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR1 moiety, and the IFN moiety, and (ii) the second polypeptide comprises, in N- to C-terminal orientation, the second Fc domain and the IFNAR2 moiety.

114. The IFN receptor agonist of embodiment 113, further comprising a first linker connecting the first Fc domain and the IFNAR1 moiety, a second linker connecting the IFNAR1 moiety and the IFN moiety, and a third linker connecting the second Fc domain and the IFNAR2 moiety.

115. The IFN receptor agonist according to any one of embodiments 1 to 114, wherein the first Fc domain and/or the second Fc domain comprises a hinge domain.

116. The IFN receptor agonist of any one of embodiments 1 to 115, wherein the Fc region is homodimeric.

117. The IFN receptor agonist of any one of embodiments 1 to 115, wherein the Fc region is heterodimeric.

118. The IFN receptor agonist of any one of embodiments 1 to 117, which comprises any pair of half-antibodies delineated in Table 2.

119. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2B.

120. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2C.

121. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2D.

122. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2E.

123. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2F.

124. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2G.

125. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2H.

126. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2I.

127. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2J.

128. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2K.

129. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2L.

130. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2M.

131. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2N.

132. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2O.

133. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2P.

134. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2Q.

135. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2R.

136. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2S.

137. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2T.

138. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2U.

139. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2V.

140. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2W.

141. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 117, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2X.

142. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 1 to 141 which comprises any of half antibody pairs designated 1-23 as set forth in Table 2.

143. The IFN receptor agonist of any one of embodiments 1 to 142, which further comprises one or more targeting moieties that bind to one or more target molecules.

144. The IFN receptor agonist of embodiment 143, which comprises a first targeting moiety that binds to a first target molecule and optionally a second targeting moiety that binds to a second target molecule.

145. The IFN receptor agonist of embodiment 144, wherein the first targeting moiety and optional second targeting moiety are antibodies or antigen-binding fragments thereof.

146. The IFN receptor agonist of embodiment 145, wherein the first targeting moiety and optional second targeting moiety are Fabs or scFvs.

147. The IFN receptor agonist of any one of embodiments 144 to 146, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

148. The IFN receptor agonist of any one of embodiments 144 to 147, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to any target molecule identified in Section 6.7.

149. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety (a) comprises the (i) CDR or (ii) VH and VL sequences of antibody set forth in Table F or (b) competes with the antibody set forth in Table F for binding to the target molecule.

150. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to an ECM antigen which is optionally selected from syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (*e.g.*, nectin-4), tenascin, collagen (*e.g.*, collagen type X) and matrixin.

151. The IFN receptor agonist of embodiment 150, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a nectin, *e.g.*, nectin 4.

152. The IFN receptor agonist of embodiment 150, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a collagen, *e.g.*, collagen X.

153. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a cell surface molecule of tumor or viral lymphocytes.

154. The IFN receptor agonist of embodiment 153, wherein the antigen is a T-cell co-stimulatory protein.

155. The IFN receptor agonist of embodiment 154, wherein the T-cell co-stimulatory protein is CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, or B7-H3.

156. The IFN receptor agonist of embodiment 155, wherein the T-cell co-stimulatory protein is B7-H3.

157. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a checkpoint inhibitor.

158. The IFN receptor agonist of embodiment 157, wherein the checkpoint inhibitor is CTLA-4, PD1, PDL1, PDL2, B7-H3, B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK1, or CHK2.

159. The IFN receptor agonist of embodiment 158, wherein the checkpoint inhibitor is PDL1.

160. The IFN receptor agonist of embodiment 158, wherein the checkpoint inhibitor is PD1.

161. The IFN receptor agonist of embodiment 158, wherein the checkpoint inhibitor is LAG3.

162. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a tumor-associated antigen (TAA).

163. The IFN receptor agonist of embodiment 162, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to AFP, ALK, a BAGE protein, BIRC5 (survivin), BIRC7, β-catenin, brc-abl, BRCA1, BORIS, CA9, carbonic anhydrase IX, caspase-8, CALR, CEACAM5 (also known as carcinoembryonic antigen or

CEA), CCR5, CD19, CD20 (MS4A1), CD22, CD30, CD40, CDK4, CEA, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, ErbB2/Her2, ErbB3, ErbB4, ETV6-AML, EpCAM, EphA2, Fra-1, FOLR1, a GAGE protein (*e.g.*, GAGE-1 or -2), GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/k-ras, HLA/MAGE-A3, hTERT, LMP2, MAGE proteins (*e.g.*, MAGE-1, -2, -3, -4, -6, and -12), MART-1, mesothelin, ML-IAP, Muc1, Muc2, Muc3, Muc4, Muc5, Muc16 (CA-125), MUM1, NA17, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PLAC1, PRLR, PRAME, PSMA (FOLH1), RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, STEAP1, STEAP2, TAG-72, TGF-β, TMPRSS2, Thompson-nouvelle antigen (Tn), TRP-1, TRP-2, tyrosinase, or uroplakin-3.

164. The IFN receptor agonist of embodiment 163, wherein the TAA is EGFR.

165. The IFN receptor agonist of embodiment 163, wherein the TAA is HER2.

166. The IFN receptor agonist of embodiment 163, wherein the TAA is EPCAM.

167. The IFN receptor agonist of embodiment 163, wherein the TAA is CEACAM5.

168. The IFN receptor agonist of embodiment 163, wherein the TAA is CD20.

169. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a dendritic cell (DC) or other antigen-presenting cell (APC) antigen which is optionally selected from XCR1, Clec9a, CD1c, CD11c, CD14, PDL1, macrophage mannose receptor (CD206), and DEC-205.

170. The IFN receptor agonist of embodiment 169, wherein the dendritic cell antigen is XCR1.

171. The IFN receptor agonist of embodiment 169, wherein the dendritic cell antigen is Clec9a.

172. The IFN receptor agonist of embodiment 169, wherein the dendritic cell antigen is DEC-205.

173. The IFN receptor agonist of any one of embodiments 144 to 148, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a natural killer (NK) cell antigen.

174. The IFN receptor agonist of any one of embodiments 1 to 142, which further comprises one or more targeting moieties each comprising means for binding to a target molecule.

175. The IFN receptor agonist of embodiment 174, which comprises a first targeting moiety comprising means for binding to a first target molecule and optionally a second targeting moiety comprising means for binding to a second target molecule.

176. The IFN receptor agonist of embodiment 175, wherein the first targeting moiety and optional second targeting moiety are antibodies or antigen-binding fragments thereof.

177. The IFN receptor agonist of embodiment 176, wherein the first antibody or antigen-binding fragment thereof and optional second antibody or antigen-binding fragment thereof are Fabs or scFvs.

178. The IFN receptor agonist of any one of embodiments 175 to 177, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

179. The IFN receptor agonist of any one of embodiments 175 to 178, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to any target molecule identified in Section 6.7.

180. The IFN receptor agonist of any one of embodiments 175 to 179, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to an ECM antigen which is optionally selected from syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (*e.g.*, nectin-4), tenascin, collagen (*e.g.*, collagen type X) and matrixin.

181. The IFN receptor agonist of embodiment 180, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a nectin, *e.g.*, nectin 4.

182. The IFN receptor agonist of embodiment 180, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a collagen, *e.g.*, collagen X.

183. The IFN receptor agonist of any one of embodiments 175 to 177, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a cell surface molecule of tumor or viral lymphocytes.

184. The IFN receptor agonist of embodiment 183, wherein the cell surface molecule is a T-cell co-stimulatory protein.

185. The IFN receptor agonist of embodiment 184, wherein the T-cell co-stimulatory protein is CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, or B7-H3.

186. The IFN receptor agonist of embodiment 185, wherein the T-cell co-stimulatory protein is B7-H3.

187. The IFN receptor agonist of any one of embodiments 175 to 177, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a checkpoint inhibitor.

188. The IFN receptor agonist of embodiment 187, wherein the checkpoint inhibitor is CTLA-4, PD1, PDL1, PDL2, B7-H3, B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK1, or CHK2.

189. The IFN receptor agonist of embodiment 188, wherein the checkpoint inhibitor is PDL1.

190. The IFN receptor agonist of embodiment 188, wherein the checkpoint inhibitor is PD1.

191. The IFN receptor agonist of embodiment 188, wherein the checkpoint inhibitor is LAG3.

192. The IFN receptor agonist of any one of embodiments 175 to 177, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a tumor-associated antigen (TAA).

193. The IFN receptor agonist of embodiment 192, wherein the TAA is AFP, ALK, a BAGE protein, BIRC5 (survivin), BIRC7, β-catenin, brc-abl, BRCA1, BORIS, CA9, carbonic anhydrase IX, caspase-8, CALR, CEACAM5 (also known as carcinoembryonic antigen or CEA), CCR5, CD19, CD20 (MS4A1), CD22, CD30, CD40, CDK4, CEA, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, ErbB2/Her2, ErbB3, ErbB4, ETV6-AML, EpCAM, EphA2, Fra-1, FOLR1, a GAGE protein (*e.g.*, GAGE-1 or -2), GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/k-ras, HLA/MAGE-A3, hTERT, LMP2, MAGE proteins (*e.g.*, MAGE-1, -2, -3, -4, -6, and -12), MART-1, mesothelin, ML-IAP, Muc1, Muc2, Muc3, Muc4, Muc5, Muc16 (CA-125), MUM1, NA17, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PLAC1, PRLR, PRAME, PSMA (FOLH1),

RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, STEAP1, STEAP2, TAG-72, TGF-β, TMPRSS2, Thompson-nouvelle antigen (Tn), TRP-1, TRP-2, tyrosinase, or uroplakin-3.

194. The IFN receptor agonist of embodiment 193, wherein the TAA is EGFR.

195. The IFN receptor agonist of embodiment 193, wherein the TAA is HER2.

196. The IFN receptor agonist of embodiment 193, wherein the TAA is EPCAM.

197. The IFN receptor agonist of embodiment 193, wherein the TAA is CEACAM5.

198. The IFN receptor agonist of embodiment 193, wherein the TAA is CD20.

199. The IFN receptor agonist of any one of embodiments 175 to 177, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a dendritic cell (DC) or other antigen-presenting cell (APC) antigen which is optionally selected from XCR1, Clec9a, CD1c, CD11c, CD14, PDL1, macrophage mannose receptor (CD206), and DEC-205.

200. The IFN receptor agonist of embodiment 199, wherein the dendritic cell antigen is XCR1.

201. The IFN receptor agonist of embodiment 199, wherein the dendritic cell antigen is Clec9a.

202. The IFN receptor agonist of embodiment 199, wherein the dendritic cell antigen is DEC-205.

203. The IFN receptor agonist of any one of embodiments 175 to 177, wherein the first targeting moiety and/or optional second targeting moiety comprises means for binding to a natural killer (NK) cell antigen.

204. The IFN receptor agonist of any one of embodiments 143 to 203, which comprises any pair of half-antibodies delineated in Table 2.

205. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3B.

206. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3C.

207. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3D.

208. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3E.

209. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3F.

210. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3G.

211. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3H.

212. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3I.

213. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3J.

214. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3K.

215. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3L.

216. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3M.

217. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3N.

218. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3O.

219. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3P.

220. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3Q.

221. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3R.

222. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3S.

223. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3T.

224. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3U.

225. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3V.

226. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3W.

227. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 143 to 203, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3X.

228. A nucleic acid or plurality of nucleic acids encoding the IFN receptor agonist of any one of embodiments 1 to 227.

229. A host cell engineered to express the IFN receptor agonist of any one of embodiments 1 to 227 or the nucleic acid(s) of embodiment 228.

230. A method of producing the IFN receptor agonist of any one of embodiments 1 to 227, comprising culturing the host cell of embodiment 229 and recovering the IFN receptor agonist expressed thereby.

231. A pharmaceutical composition comprising the IFN receptor agonist of any one of embodiments 1 to 227 and an excipient.

232. A method of treating cancer, comprising administering to a subject in need thereof the IFN receptor agonist of any one of embodiments 1 to 227 or the pharmaceutical composition of embodiment 231.

233. The method of embodiment 232, wherein the IFN receptor agonist comprises at least one targeting moiety that is capable of binding to a target molecule.

234. The method of embodiment 232, wherein the IFN receptor agonist comprises at least one targeting moiety comprising means for binding to a target molecule.

235. The method of embodiment 233 or 234, wherein the cancer is associated with expression of the target molecule, *e.g.*, a TAA and associated cancer as set forth in Table I.

236. The method of any one of embodiments 232 to 235, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases expressed by the cancer tissue.

237. The method of embodiment 236, wherein the IFN protein is selectively activated in the cancer tissue.

238. A method of localized delivery of an IFN protein, comprising administering to a subject an IFN receptor agonist according to any one of embodiments 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue to which the IFN protein is to be locally delivered.

239. The method of embodiment 238, wherein the IFN receptor agonist comprises one or more targeting moieties that recognize a target molecule expressed by the tissue.

240. The method of embodiment 239, wherein the IFN receptor agonist comprises two targeting moieties that each recognize a target molecule expressed by the tissue.

241. The method of embodiment 238, wherein the IFN receptor agonist comprises one or more targeting moieties each comprising means for binding to a target molecule expressed by the tissue.

242. The method of embodiment 241, wherein the IFN receptor agonist comprises two targeting moieties each comprising means for binding to a target molecule expressed by the tissue.

243. The method of any one of embodiments 238 to 242, wherein the tissue is cancer tissue.

244. The method of embodiment 243, wherein the target molecule expressed by the tissue is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

245. The method of any one of embodiments 238 to 244, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the tissue.

246. A method of treating cancer with an IFN protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IFN receptor agonist according to any one of embodiments 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by cancer tissue, *e.g.*, a cancer tissue to which the IFN receptor agonist is targeted.

247. The method of embodiment 246, wherein the IFN receptor agonist comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue or associated immune cells.

248. The method of embodiment 247, wherein the IFN receptor agonist comprises two targeting moieties that each recognize a target molecule expressed by the cancer tissue or associated immune cells.

249. The method of embodiment 246, wherein the IFN receptor agonist comprises one or more targeting moieties each comprising means for binding to a target molecule expressed by the cancer tissue or associated immune cells.

250. The method of embodiment 249, wherein the IFN receptor agonist comprises two targeting moieties each comprising means for binding to a target molecule expressed by the cancer tissue or associated immune cells.

251. The method of any one of embodiments 246 to 250, wherein the target molecule expressed by the cancer tissue or associated immune cells is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated

antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

252. The method of any one of embodiments 246 to 251, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the cancer tissue.

253. A method of administering to the subject IFN therapy with reduced systemic exposure and/or reduced systemic toxicity, comprising administering to a subject the IFN therapy in the form of an IFN receptor agonist according to any one of embodiments 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IFN therapy is desirable and/or intended.

254. The method of embodiment 253, wherein the IFN receptor agonist comprises one or more targeting moieties that recognize a target molecule expressed by the tissue.

255. The method of embodiment 254, wherein the IFN receptor agonist comprises two targeting moieties that each recognize a target molecule expressed by the tissue.

256. The method of embodiment 253, wherein the IFN receptor agonist comprises one or more targeting moieites comprising means for binding to a target molecule expressed by the tissue.

257. The method of embodiment 256, wherein the IFN receptor agonist comprises two targeting moieties comprising means for binding to a target molecule expressed by the tissue.

258. The method of any one of embodiments 253 to 257, wherein the tissue is cancer tissue or associated immune cells.

259. The method of embodiment 258, wherein the target molecule expressed by the tissue is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

260. The method of any one of embodiments 253 to 259, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the tissue.

261. A method of treating cancer with an IFN protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IFN receptor agonist according to any one of embodiments 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by the cancer tissue.

262. The method of embodiment 261, wherein the IFN receptor agonist comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue or associated immune cells.

263. The method of embodiment 262, wherein the IFN receptor agonist comprises two targeting moieties that each recognize a target molecule expressed by the cancer tissue or associated immune cells.

264. The method of embodiment 261, wherein the IFN receptor agonist comprises one or more targeting moieties each comprising means for binding to a target molecule expressed by the cancer tissue or associated immune cells.

265. The method of embodiment 264, wherein the IFN receptor agonist comprises two targeting moieties each comprising means for binding to a target molecule expressed by the cancer tissue or associated immune cells.

266. The method of any one of embodiments 261 to 265, wherein the target molecule expressed by the cancer tissue or associated immune cells is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

267. The method of any one of embodiments 261 to 266, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the cancer tissue.

268. A method of targeted delivery of an activated IFN protein to cancer tissue, comprising administering to a subject an IFN receptor agonist according to any one of embodiments 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient), wherein the IFN receptor agonist:

(a) comprises (i) one or more targeting moieties that recognize a target molecule expressed by the cancer tissue or associated immune cells or (ii) one or more targeting moieties each comprising means for binding to a target molecule expressed by the cancer tissue or associated immune cells; and

(b) has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in a tissue for which IFN therapy is desirable and/or intended.

269. The method of embodiment 268, wherein the IFN receptor agonist comprises (i) two targeting moieties that each recognize a target molecule expressed by the cancer tissue or associated immune cells or (ii) two targeting moieties each comprising means for binding to a target molecule expressed by the cancer tissue or associated immune cells.

270. The method of embodiment 268 or embodiment 269, wherein the target molecule expressed by the cancer tissue or associated immune cells is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor

or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

271. The method of any one of embodiments 268 to 270, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the cancer tissue.

272. A method of locally inducing an immune response in a target tissue, comprising administering to a subject an IFN receptor agonist according to any one of embodiments 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has (i) one or more targeting moieties capable of binding a target molecule expressed in the target tissue or (ii) one or more targeting moieties each comprising means for binding to a target molecule expressed in the target tissue and one or more protease-cleavable linkers, each protease-cleavable linker comprising one or more substrates for one or more proteases expressed in the target tissue.

273. The method of embodiment 272, wherein the IFN receptor agonist comprises (i) two targeting moieties that each recognize a target molecule expressed in the target tissue or associated immune cells or (ii) two targeting moieties each comprising means for binding to a target molecule expressed in the target tissue or associated immune cells.

274. The method of embodiment 272 or embodiment 273, wherein the target tissue is cancer tissue.

275. The method of any one of embodiments 272 to 274, wherein the target molecule expressed in the target tissue or associated immune cells is an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

276. The method of any one of embodiments 272 to 275, wherein an activated IFN protein comprising the IFN moiety is produced by cleavage of one or more protease-cleavable linkers in the IFN receptor agonist by one or more proteases in the target tissue.

277. The method of embodiment 276, wherein the IFN protein induces the immune response against at least one cell type in the target tissue.

278. A method of enhancing an immune response against an antigen, comprising administering to a subject an immunogenic agent that elicits an immune response against the antigen together with an IFN receptor agonist according to any one of claims 1 to 227 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) or a nucleic acid encoding such IFN receptor agonist, *e.g.*, as described in Section 6.11.1).

279. The method of embodiment 278, wherein the administration of the immunogenic agent and the IFN receptor agonist are concurrent, separate but simultaneous, or sequential.

280. The method of embodiment 278 or embodiment 279, wherein the immunogenic agent is a vaccine, optionally wherein the vaccine is a cancer vaccine or a vaccine against an infectious agent.

281. The method of any one of embodiments 232 to 280, wherein the administration is non-local.

282. The method of embodiment 281, wherein the administration is systemic.

283. The method of embodiment 281, wherein the administration is subcutaneous.

284. A Type I interferon (IFN) receptor agonist, comprising:

(a) a first polypeptide chain comprising a first Fc domain and a Type I interferon (IFN) moiety attenuated by masking by an interferon alpha receptor 1 (IFNAR) moiety and an interferon alpha receptor 2 (IFNAR2) moiety; and

(b) a second polypeptide chain comprising a second Fc domain associated with the first Fc domain.

285. The IFN receptor agonist of embodiment 284, wherein the IFN moiety is N-terminal to the first Fc domain.

286. The IFN receptor agonist of embodiment 284, wherein the IFN moiety is C-terminal to the first Fc domain.

287. The IFN receptor agonist of any one of embodiments 284 to 286, wherein the first polypeptide chain comprises the IFNAR1 moiety.

288. The IFN receptor agonist of embodiment 287, wherein the IFNAR1 moiety is N-terminal to the IFN moiety.

289. The IFN receptor agonist of embodiment 287, wherein the IFNAR1 moiety is C-terminal to the IFN moiety.

290. The IFN receptor agonist of any one of embodiments 284 to 289, wherein the first polypeptide chain comprises the IFNAR2 moiety.

291. The IFN receptor agonist of embodiment 290, wherein the IFNAR2 moiety is N-terminal to the IFN moiety.

292. The IFN receptor agonist of embodiment 290, wherein the IFNAR2 moiety is C-terminal to the IFN moiety.

293. The IFN receptor agonist of any one of embodiments 284 to 292, further comprising one or more linkers connecting two or more of the first Fc domain, the IFN moiety, the IFNAR1 moiety, and the IFNAR2 moiety.

294. The IFN receptor agonist of embodiment 284, wherein the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR1 moiety, the IFN moiety, and the IFNAR2 moiety.

295. The IFN receptor agonist of embodiment 294, further comprising a first linker connecting the first Fc domain and the first IFNAR1 moiety, a second linker connecting the IFNAR1 moiety and the IFN moiety, and a third linker connecting the IFN moiety and the IFNAR2 moiety.

296. The IFN receptor agonist of embodiment 284, wherein the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, a first linker, the IFNAR2 moiety, a second linker, the IFN moiety, a third linker, and the IFNAR1 moiety.

297. The IFN receptor agonist of embodiment 296, further comprising a first linker connecting the first Fc domain and the first IFNAR2 moiety, a second linker connecting the IFNAR2 moiety and the IFN moiety, and a third linker connecting the IFN moiety and the IFNAR1 moiety.

298. The IFN receptor agonist of embodiment 284, wherein (i) the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR2 moiety, and the IFN moiety, and (ii) the second polypeptide comprises, in N- to C-terminal orientation, the second Fc domain, and the IFNAR1 moiety.

299. The IFN receptor agonist of embodiment 298, further comprising a first linker connecting the first Fc domain and the IFNAR2 moiety, a second linker connecting the IFNAR2 moiety and the IFN moiety, and a third linker connecting the second Fc domain and the IFNAR1 moiety.

300. The IFN receptor agonist of embodiment 284, wherein (i) the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR1 moiety, and the IFN moiety, and (ii) the second polypeptide comprises, in N- to C-terminal orientation, the second Fc domain and the IFNAR2 moiety.

301. The IFN receptor agonist of embodiment 300, further comprising a first linker connecting the first Fc domain and the IFNAR1 moiety, a second linker connecting the IFNAR1 moiety and the IFN moiety, and a third linker connecting the second Fc domain and the IFNAR2 moiety.

302. The IFN receptor agonist of any one of embodiments 295, 297, 299, or 301, wherein one or more of the first linker, the second linker, and the third linker is a protease-cleavable linker (PCL).

303. The IFN receptor agonist of embodiment 302, wherein the second linker is a PCL.

304. The IFN receptor agonist of embodiment 302 or 303, wherein the PCL comprises a substrate sequence cleavable by any protease set forth in Table A.

305. The IFN receptor agonist of any one of embodiments 302 to 304, wherein the PCL comprises one or more substrate sequences selected from the substrate sequences set forth in Table B.

306. The IFN receptor agonist of any one of embodiments 302 to 305, wherein the PCL comprises one or more spacer sequences selected from the spacer sequences set forth in Table C.

307. The IFN receptor agonist of any one of embodiments 302 to 306, wherein the PCL comprises the amino acid sequence of any of the PCL sequences set forth in Table D or a variant thereof with up to 5 amino acid substitutions.

308. The IFN receptor agonist of any one of embodiments 302 to 307, which is configured such that cleavage of the protease-cleavable linker (PCL) unmasks the IFN moiety.

309. The IFN receptor agonist of any one of embodiments 284 to 308, wherein the second polypeptide chain comprises an additional IFN moiety masked by an additional IFNAR1 moiety and an additional IFNAR2 moiety.

310. The IFN receptor agonist of any one of embodiments 284 to 309, wherein the IFN moiety comprises an amino acid sequence having at least about 90%, at least about 95%, or at least about 98% sequence identity to (a) full length mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ or (b) a mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ having up to a 15-amino acid truncation at its N-terminus and/or its C-terminus.

311. The IFN receptor agonist of any one of embodiments 284 to 310, wherein the IFN moiety comprises an amino acid sequence having one or more attenuating mutations as compared to mature human IFNα1 or IFNα2b.

312. The IFN receptor agonist of any one of embodiments 284 to 311, wherein the IFN moiety has one or more mutations selected from L26A, F27A, R33A, R33K, L30A, D35E, H57Y, E58N, Q61S, H57S, E58S, H57A, E58A, Q61A, Q90A, E96A, R120A, L135A, R144A, R144S, R144T, R144Y, R144I, R144L, A145D, A145H, A145K, A145M, A145V, A145Y, R149A, R149K, S152A, R162A, and E165D.

313. The IFN receptor agonist of any one of embodiments 284 to 312, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90%, at least 95%, or at least 98% sequence identity to (i) the SD2 and SD3 domains of human IFNAR1, (ii) the SD1, SD2 and SD3 domains of human IFNAR1, or (iii) the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

314. The IFN receptor agonist of any one of embodiments 284 to 313, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90%, at least 95%, or at least 98% sequence identity to (i) the D1 domain of human IFNAR2 or (ii) the D1 and D2 domains of human IFNAR2.

315. The IFN receptor agonist according to any one of embodiments 284 to 314, wherein the first Fc domain and/or the second Fc domain comprises a hinge domain.

316. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2N.

317. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2L.

318. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2M.

319. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2O.

320. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2P.

321. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2Q.

322. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2U.

323. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 284 to 315, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2V.

324. The IFN receptor agonist of any one of embodiments 284 to 323, which comprises a first targeting moiety that binds to a first target molecule and optionally a second targeting moiety that binds to a second target molecule.

325. The IFN receptor agonist of embodiment 324, wherein the first targeting moiety and optional second targeting moiety are antibodies or antigen-binding fragments thereof.

326. The IFN receptor agonist of embodiment 324 or 325, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

327. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety (a) comprises the (i) CDR or (ii) VH and VL sequences of antibody set forth in Table F or (b) competes with the antibody set forth in Table F for binding to the target molecule.

328. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to an ECM antigen which is optionally selected from syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (*e.g.*, nectin-4), tenascin, collagen (*e.g.*, collagen type X) and matrixin.

329. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a cell surface molecule of tumor or viral lymphocytes.

330. The IFN receptor agonist of embodiment 329, wherein the cell surface molecule is a T-cell co-stimulatory protein, optionally selected from CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, and B7-H3.

331. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a checkpoint inhibitor.

332. The IFN receptor agonist of embodiment 331, wherein the checkpoint inhibitor is PDL1.

333. The IFN receptor agonist of embodiment 331, wherein the checkpoint inhibitor is PD1.

334. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a tumor-associated antigen (TAA), optionally selected from AFP, ALK, a BAGE protein, BIRC5 (survivin), BIRC7, β-catenin, brc-abl, BRCA1, BORIS, CA9, carbonic anhydrase IX, caspase-8, CALR, CEACAM5 (also known as carcinoembryonic antigen or CEA), CCR5, CD19, CD20 (MS4A1), CD22, CD30, CD40, CDK4, CEA, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, ErbB2/Her2, ErbB3, ErbB4, ETV6-AML, EpCAM, EphA2, Fra-1, FOLR1, a GAGE protein (*e.g.*, GAGE-1 or -2), GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/k-ras, HLA/MAGE-A3, hTERT, LMP2, MAGE proteins (*e.g.*, MAGE-1, -2, -3, -4, -6, and -12), MART-1, mesothelin, ML-IAP, Muc1, Muc2, Muc3, Muc4, Muc5, Muc16 (CA-125), MUM1, NA17, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PLAC1, PRLR, PRAME, PSMA (FOLH1), RAGE proteins, Ras,

RGS5, Rho, SART-1, SART-3, STEAP1, STEAP2, TAG-72, TGF-β, TMPRSS2, Thompson-nouvelle antigen (Tn), TRP-1, TRP-2, tyrosinase, and uroplakin-3.

335. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a dendritic cell (DC) or other antigen-presenting cell (APC) antigen which is optionally selected from XCR1, Clec9a, CD1c, CD11c, CD14, macrophage mannose receptor (CD206), and DEC-205.

336. The IFN receptor agonist of any one of embodiments 324 to 326, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a natural killer (NK) cell antigen.

337. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3N.

338. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3L.

339. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3M.

340. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3O.

341. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3P.

342. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3Q.

343. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3U.

344. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of embodiments 324 to 336, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3V.

345. A nucleic acid or plurality of nucleic acids encoding the IFN receptor agonist of any one of embodiments 284 to 344.

346. A host cell engineered to express the IFN receptor agonist of any one of embodiments 284 to 344 or the nucleic acid(s) of embodiment 345.

347. A method of producing the IFN receptor agonist of any one of embodiments 284 to 344, comprising culturing the host cell of embodiment 346 and recovering the IFN receptor agonist expressed thereby.

348. A pharmaceutical composition comprising the IFN receptor agonist of any one of embodiments 284 to 344 and an excipient.

349. A method of treating cancer, comprising administering to a subject in need thereof the IFN receptor agonist of any one of embodiments 284 to 344 or the pharmaceutical composition of embodiment 348.

350. A method of localized delivery of an IFN protein, comprising administering to a subject an IFN receptor agonist according to any one of embodiments 284 to 344 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which

has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue to which the IFN protein is to be locally delivered.

351. A method of treating cancer with an IFN protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IFN receptor agonist according to any one of embodiments 284 to 344 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by cancer tissue.

352. A method of administering to the subject IFN therapy with reduced systemic exposure and/or reduced systemic toxicity, comprising administering to a subject the IFN therapy in the form of an IFN receptor agonist according to any one of embodiments 284 to 344 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IFN therapy is desirable and/or intended.

353. A method of targeted delivery of an activated IFN protein to cancer tissue, comprising administering to a subject an IFN receptor agonist according to any one of embodiments 284 to 344 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient), wherein the IFN receptor agonist:

(a) comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue or associated immune cells; and

(b) has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in a tissue for which IFN therapy is desirable and/or intended.

354. A method of locally inducing an immune response in a target tissue, comprising administering to a subject an IFN receptor agonist according to any one of

embodiments 284 to 344 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more targeting moieties capable of binding a target molecule expressed in the target tissue and one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in the target tissue.

355. A method of enhancing an immune response against an antigen, comprising administering to a subject an immunogenic agent that elicits an immune response against the antigen together with an IFN receptor agonist according to any one of embodiments 284 to 344 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) or a nucleic acid encoding such IFN receptor agonist.

356. The method of any one of embodiments 349 to 355, wherein the administration is non-local.

8. EXAMPLES

8.1. IFN Receptor Agonists Construct Sequences

[0306] Table 6 below provides sequences of IFN receptor agonist and control constructs utilized in the studies described herein. Targeting moieties may be included in all of these as specified above.

Table 6

Construct	Sequence
Fc-IFN (homodimeric, both chains identical) Chain 1: Fc-linker-IFN Chain 2: Fc-Linker IFN	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHSL GSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAE TIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDL EACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 385) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV

Table 6	
Construct	**Sequence**
	SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHSL GSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAE TIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDL EACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 385)
IFN-Fc (homodimeric, both chains identical) Chain 1: IFN-linker-Fc Chain 2: IFN-linker-Fc	Chain 1: CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEE FGNQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYT ELYQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYL KEKKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGSGGGGS GGGGSGESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQF NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKA KGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVF SCSVMHEALHNHYTQKSLSLSLGK (SEQ ID NO: 386) Chain 2: CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEE FGNQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYT ELYQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYL KEKKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGSGGGGS GGGGSGESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISR TPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQF NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKA KGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVF SCSVMHEALHNHYTQKSLSLSLGK (SEQ ID NO: 386)
Fc-IFNAR1(SD1-3)-IFN (homodimeric, both chains identical) Chain 1: Fc-linker-IFNAR1(SD1-3)-linker-IFN Chain 2: Fc-linker-IFNAR1(SD1-3)-linker-IFN	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQKV EVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSGC QNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTPF RKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSL VIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKI GVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYAN MTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFPQ

Table 6	
Construct	**Sequence**
	NVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCDLP QTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQ FQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQ QLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKK YSPCAWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 387) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQKV EVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSGC QNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTPF RKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSL VIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKI GVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYAN MTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFPQ NVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCDLP QTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQ FQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQ QLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKK YSPCAWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 387)
Fc-IFNAR1(SD2-3)-IFN (homodimeric, both chains identical) Chain 1: Fc-linker-IFNAR1(SD2-3)-linker-IFN Chain 2: Fc-linker-IFNAR1(SD2-3)-linker-IFN	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSIGPPEVHLE AEDKAIVIHISPGTKDSVMWALDGLSFTYSLVIWKNSSGVEERI ENIYSRHKIYKLSPETTYCLKVKAALLTSWKIGVYSPVHCIKTTV ENELPPPENIEVSVQNQNYVLKWDYTYANMTFQVQWLHAFLK RNPGNHLYKWKQIPDCENVKTTQCVFPQNVFQKGIYLLRVQA SDGNNTSFWSEEIKFDTEIQGGGGSCDLPQTHSLGSRRTLML LAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHEMI QQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQGV GVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRAEI MRSFSLSTNLQESLRSKE (SEQ ID NO: 388) Chain 2:

Table 6	
Construct	**Sequence**
	ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSIGPPEVHLE AEDKAIVIHISPGTKDSVMWALDGLSFTYSLVIWKNSSGVEERI ENIYSRHKIYKLSPETTYCLKVKAALLTSWKIGVYSPVHCIKTTV ENELPPPENIEVSVQNQNYVLKWDYTYANMTFQVQWLHAFLK RNPGNHLYKWKQIPDCENVKTTQCVFPQNVFQKGIYLLRVQA SDGNNTSFWSEEIKFDTEIQGGGGSCDLPQTHSLGSRRTLML LAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHEMI QQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQGV GVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRAEI MRSFSLSTNLQESLRSKE (SEQ ID NO: 388)
Fc-IFN-IFNAR2(D1-2) (homodimeric, both chains identical) Chain 1: Fc-linker-IFN-linker-IFNAR2(D1-2) Chain 2: Fc-linker-IFN-linker-IFNAR2(D1-2)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHSL GSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAE TIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDL EACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSISYDSP DYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPE DLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTL FSCSHNFWLAIDMSFEPPEFEIVGFTNHINVMVKFPSIVEEELQ FDLSLVIEEQSEGIVKKHKPEIKGNMSGNFTYIIDKLIPNTNYCV SVYLEHSDEQAVIKSPLKCTLLPP (SEQ ID NO: 389) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHSL GSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAE TIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDL EACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSISYDSP DYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPE DLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTL

Table 6	
Construct	**Sequence**
	FSCSHNFWLAIDMSFEPPEFEIVGFTNHINVMVKFPSIVEEELQ FDLSLVIEEQSEGIVKKHKPEIKGNMSGNFTYIIDKLIPNTNYCV SVYLEHSDEQAVIKSPLKCTLLPP (SEQ ID NO: 389)
Fc-IFN-IFNAR2(D1) (homodimeric, both chains identical) Chain 1: Fc-linker-IFN-linker-IFNAR2(D1) Chain 2: Fc-linker-IFN-linker-IFNAR2(D1)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHSL GSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAE TIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDL EACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSISYDSP DYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPE DLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTL FSCSHNFWLAIDMSFEP (SEQ ID NO: 390) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHSL GSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAE TIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDL EACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCA WEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSISYDSP DYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPE DLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTL FSCSHNFWLAIDMSFEP (SEQ ID NO: 390)
Fc-IFNAR2(D1)-IFN (homodimeric, both chains identical) Chain 1: Fc-linker-IFNAR2(D1)-linker-IFN	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSISYDSPDYT DESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLK VVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSC SHNFWLAIDMSFEPGGGGSCDLPQTHSLGSRRTLMLLAQMR RISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHEMIQQIFNL FSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQGVGVTETP

Table 6	
Construct	**Sequence**
Chain 2: Fc-linker-IFNAR2(D1)-linker-IFN	LMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRAEIMRSFSL STNLQESLRSKE (SEQ ID NO: 391) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSISYDSPDYT DESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLK VVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSC SHNFWLAIDMSFEPGGGGSCDLPQTHSLGSRRTLMLLAQMR RISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHEMIQQIFNL FSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQGVGVTETP LMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRAEIMRSFSL STNLQESLRSKE (SEQ ID NO: 391)
Fc-IFN x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFN Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHS LGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLND LEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPC AWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 392) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK* (SEQ ID NO: 393)
Fc-IFN x Fc-R1(SD1-3) (heterodimeric) Chain 1: Fc_{knob}-linker-IFN Chain 2: Fc_{hole}-linker-IFNAR1(SD1-3)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHS LGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA

Table 6	
Construct	**Sequence**
	ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLND LEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPC AWEVVRAEIMRSFSLSTNLQESLRSKE(SEQ ID NO: 392) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGKGGGGSGGGGSGGGGSKNLKSPQKV EVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSGC QNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTPF RKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSL VIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKI GVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYAN MTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFPQ NVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQ (SEQ ID NO: 394)
Fc-IFN x Fc-R2(D1-2) (heterodimeric) Chain 1: Fc_{knob}-linker-IFN Chain 2: Fc_{hole} -linker-IFNAR2(D1-2)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHS LGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLND LEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPC AWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 392) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGKGGGGSGGGGSGGGGSISYDSPDYT DESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLK VVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSC SHNFWLAIDMSFEPPEFEIVGFTNHINVMVKFPSIVEEELQFDL SLVIEEQSEGIVKKHKPEIKGNMSGNFTYIIDKLIPNTNYCVSVY LEHSDEQAVIKSPLKCTLLPP (SEQ ID NO: 395)

Table 6	
Construct	**Sequence**
Fc-R1-IFN x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR1-linker-IFN Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQK VEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSG CQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTP FRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYS LVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSW KIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYA NMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFP QNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCD LPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFG NQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTEL YQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKE KKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 396) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK (SEQ ID NO: 397)
Fc-IFN-R2(D1-2) x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFN-IFNAR2(D1-2) Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHS LGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLND LEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPC AWEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSISYDS PDYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKP EDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTT LFSCSHNFWLAIDMSFEPPEFEIVGFTNHINVMVKFPSIVEEEL QFDLSLVIEEQSEGIVKKHKPEIKGNMSGNFTYIIDKLIPNTNYC VSVYLEHSDEQAVIKSPLKCTLLPP (SEQ ID NO: 398) Chain 2:

Table 6	
Construct	**Sequence**
	ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK (SEQ ID NO: 397)
Fc-IFN x Fc-R2(D1) (heterodimeric) Chain 1: Fc_{knob}-linker-IFN Chain 2: Fc_{hole}-linker - IFNAR2(D1)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHS LGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLND LEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPC AWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 392) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGKGGGGSGGGGSGGGGSISYDSPDYT DESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLK VVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSC SHNFWLAIDMSFEP (SEQ ID NO: 399)
Fc-R2(D1)-IFN x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR2(D1)-linker-IFN Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSISYDSPDY TDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDL KVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFS CSHNFWLAIDMSFEPGGGGSCDLPQTHSLGSRRTLMLLAQM RRISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHEMIQQIFN LFSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQGVGVTET PLMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRAEIMRSFS LSTNLQESLRSKE (SEQ ID NO: 400)

Table 6	
Construct	**Sequence**
	Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK (SEQ ID NO: 397)
Fc-R1(SD1-3)-IFN x Fc-R2(D1-2) (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR1(SD1-3)-linker-IFN Chain 2: Fc_{hole}-linker-IFNAR2(D1-2)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQK VEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSG CQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTP FRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYS LVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSW KIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYA NMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFP QNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCD LPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFG NQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTEL YQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKE KKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 396) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGKGGGGSGGGGSGGGGSISYDSPDYT DESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLK VVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSC SHNFWLAIDMSFEPPEFEIVGFTNHINVMVKFPSIVEEELQFDL SLVIEEQSEGIVKKHKPEIKGNMSGNFTYIIDKLIPNTNYCVSVY LEHSDEQAVIKSPLKCTLLPP (SEQ ID NO: 395)
Fc-R2(D1)-IFN x Fc-R1(SD1-3) (heterodimeric)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP

Table 6	
Construct	**Sequence**
Chain 1: Fc_{knob}-linker-IFNAR2(D1)-linker-IFN Chain 2: Fc_{hole}-linker-IFNAR1(SD1-3)	ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSISYDSPDY TDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDL KVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFS CSHNFWLAIDMSFEPGGGGSCDLPQTHSLGSRRTLMLLAQM RRISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHEMIQQIFN LFSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQGVGVTET PLMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRAEIMRSFS LSTNLQESLRSKE (SEQ ID NO: 400) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGKGGGGSGGGGSGGGGSKNLKSPQKV EVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSGC QNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTPF RKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSL VIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKI GVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYAN MTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFPQ NVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQ (SEQ ID NO: 394)
Fc-IFNAR1(SD1-3)-IFN-IFNAR2(D1) (homodimeric, both chains identical) Chain 1: Fc-linker-IFNAR1(SD1-3)-linker-IFN-linker-IFNAR2(D1) Chain 2: Fc-linker-IFNAR1(SD1-3)-linker-IFN-linker-IFNAR2(D1	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQKV EVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSGC QNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTPF RKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSL VIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKI GVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYAN MTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFPQ NVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCDLP QTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQ FQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQ QLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKK YSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSI SYDSPDYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTI

Table 6	
Construct	**Sequence**
	MSKPEDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFS GNTTLFSCSHNFWLAIDMSFEP (SEQ ID NO: 401) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHE ALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQKV EVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSGC QNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTPF RKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYSL VIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSWKI GVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYAN MTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFPQ NVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCDLP QTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQ FQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQ QLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKK YSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSI SYDSPDYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTI MSKPEDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFS GNTTLFSCSHNFWLAIDMSFEP (SEQ ID NO: 401)
Fc-R1(SD1-3)-IFN x Fc-R2(D1) (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR1(SD1-3)-linker-IFN Chain 2: Fc_{hole}-linker-IFNAR2(D1)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQK VEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSG CQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTP FRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYS LVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSW KIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYA NMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFP QNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCD LPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFG NQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTEL YQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKE KKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKE (SEQ ID NO: 396)

Table 6	
Construct	**Sequence**
	Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGKGGGGSGGGGSGGGGSISYDSPDYT DESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLK VVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSC SHNFWLAIDMSFEP (SEQ ID NO: 399)
Fc-IFNAR1(SD1-3)-IFN-IFNAR2(D1) x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR1(SD1-3)-linker-IFN-linker-IFNAR2(D1) Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQK VEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSG CQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTP FRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYS LVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSW KIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYA NMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFP QNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSCD LPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFG NQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTEL YQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKE KKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGG GSISYDSPDYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLL YTIMSKPEDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEG FSGNTTLFSCSHNFWLAIDMSFEP (SEQ ID NO: 402) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK (SEQ ID NO: 397)
Fc-IFN-R2(D1) x Fc (heterodimeric)	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP

Table 6	
Construct	**Sequence**
Chain 1: Fc_{knob}-linker-IFN-linker-IFNAR2(D1) Chain 2: Fc_{hole}	ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSCDLPQTHS LGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKA ETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLND LEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPC AWEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGGGGSISYDS PDYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKP EDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTT LFSCSHNFWLAIDMSFEP (SEQ ID NO: 403) Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK (SEQ ID NO: 397)
Fc x IFN-Fc (heterodimeric) Chain 1: Fc_{knob} Chain 2: IFN-linker-Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGK (SEQ ID NO: 404) Chain 2: CDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFPQEE FGNQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLDKFYT ELYQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRITLYL KEKKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGGGGSGG GGSGGGGSESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREE QFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIS KAKGQPREPQVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAV EWESNGQPENNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEG NVFSCSVMHEALHNRFTQKSLSLSPGK (SEQ ID NO: 405)
Fc x R1(SD1-3)-IFN-Fc (heterodimeric) Chain 1: Fc_{knob} Chain 2: IFNAR1(SD1-3)-linker-IFN-linker-Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGK (SEQ ID NO: 404)

Table 6	
Construct	**Sequence**
	Chain 2: KNLKSPQKVEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGM DNWIKLSGCQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSW YEVDSFTPFRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWAL DGLSFTYSLVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVK AALLTSWKIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLK WDYTYANMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKT TQCVFPQNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGG GGSCDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFP QEEFGNQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLD KFYTELYQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRI TLYLKEKKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGGG GSGGGGSGGGGSESKYGPPCPPCPAPPVAGPSVFLFPPKPK DTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTK PREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSI EKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLSCAVKGFYP SDIAVEWESNGQPENNYKTTPPVLDSDGSFFLVSRLTVDKSR WQEGNVFSCSVMHEALHNRFTQKSLSLSPGK (SEQ ID NO: 406)
Fc x R2(D1)-IFN-Fc (heterodimeric) Chain 1: Fc_{knob} Chain 2: IFNAR2D1-linker-IFN-linker-Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGK (SEQ ID NO: 404) Chain 2: ISYDSPDYTDESCTFKISLRNFRSILSWELKNHSIVPTHYTLLYTI MSKPEDLKVVKNCANTTRSFCDLTDEWRSTHEAYVTVLEGFS GNTTLFSCSHNFWLAIDMSFEPGGGGSCDLPQTHSLGSRRTL MLLAQMRRISLFSCLKDRHDFGFPQEEFGNQFQKAETIPVLHE MIQQIFNLFSTKDSSAAWDETLLDKFYTELYQQLNDLEACVIQG VGVTETPLMKEDSILAVRKYFQRITLYLKEKKYSPCAWEVVRA EIMRSFSLSTNLQESLRSKEGGGGSGGGGSGGGGSESKYGP PCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQ EDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH QDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPENNYKTTP PVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHEALHNRFT QKSLSLSPGK (SEQ ID NO: 407)

Table 6

Construct	Sequence
Fc x R1(SD1-3)-IFN-R2(D1)-Fc (heterodimeric) Chain 1: Fc_{knob} Chain 2: IFNAR1(SD1-3)-linker-IFN-linker-IFNAR2(D1)-linker-Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGK (SEQ ID NO: 404) Chain 2: KNLKSPQKVEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGM DNWIKLSGCQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSW YEVDSFTPFRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWAL DGLSFTYSLVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVK AALLTSWKIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLK WDYTYANMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKT TQCVFPQNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGG GGSCDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRHDFGFP QEEFGNQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWDETLLD KFYTELYQQLNDLEACVIQGVGVTETPLMKEDSILAVRKYFQRI TLYLKEKKYSPCAWEVVRAEIMRSFSLSTNLQESLRSKEGGG GSGGGGSISYDSPDYTDESCTFKISLRNFRSILSWELKNHSIVP THYTLLYTIMSKPEDLKVVKNCANTTRSFCDLTDEWRSTHEAY VTVLEGFSGNTTLFSCSHNFWLAIDMSFEPGGGGSGGGGSG GGGSESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTP EVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNS TYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKG QPREPQVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWES NGQPENNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFS CSVMHEALHNRFTQKSLSLSPGK (SEQ ID NO: 408)
Fc-R1(SD1-3)-3xG4S-IFN-3xG4S-R2(D1) x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR1(SD1-3)-3xG4S-IFN-3xG4S-IFNAR2(D1) Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQK VEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSG CQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTP FRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYS LVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSW KIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYA NMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFP QNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSGG GGSGGGGSCDLPQTHSLGSRRTLMLLAQMRRISLFSCLKDRH

Table 6	
Construct	**Sequence**
	DFGFPQEEFGNQFQKAETIPVLHEMIQQIFNLFSTKDSSAAWD ETLLDKFYTELYQQLNDLEACVIQGVGVTETPLMKEDSILAVRK YFQRITLYLKEKKYSPCAWEVVRAEIMRSFSLSTNLQESLRSK EGGGGSGGGGSGGGGSISYDSPDYTDESCTFKISLRNFRSIL SWELKNHSIVPTHYTLLYTIMSKPEDLKVVKNCANTTRSFCDLT DEWRSTHEAYVTVLEGFSGNTTLFSCSHNFWLAIDMSFEP Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK
Fc-R1(SD1-3)-4xG4S-IFN-4xG4S-R2(D1) x Fc (heterodimeric) Chain 1: Fc_{knob}-linker-IFNAR1(SD1-3)-4xG4S-IFN-4xG4S-IFNAR2(D1) Chain 2: Fc_{hole}	Chain 1: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLWCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMH EALHNHYTQKSLSLSLGKGGGGSGGGGSGGGGSKNLKSPQK VEVDIIDDNFILRWNRSDESVGNVTFSFDYQKTGMDNWIKLSG CQNITSTKCNFSSLKLNVYEEIKLRIRAEKENTSSWYEVDSFTP FRKAQIGPPEVHLEAEDKAIVIHISPGTKDSVMWALDGLSFTYS LVIWKNSSGVEERIENIYSRHKIYKLSPETTYCLKVKAALLTSW KIGVYSPVHCIKTTVENELPPPENIEVSVQNQNYVLKWDYTYA NMTFQVQWLHAFLKRNPGNHLYKWKQIPDCENVKTTQCVFP QNVFQKGIYLLRVQASDGNNTSFWSEEIKFDTEIQGGGGSGG GGSGGGGSGGGGSCDLPQTHSLGSRRTLMLLAQMRRISLFS CLKDRHDFGFPQEEFGNQFQKAETIPVLHEMIQQIFNLFSTKD SSAAWDETLLDKFYTELYQQLNDLEACVIQGVGVTETPLMKED SILAVRKYFQRITLYLKEKKYSPCAWEVVRAEIMRSFSLSTNLQ ESLRSKEGGGGSGGGGSGGGGSGGGGSISYDSPDYTDESC TFKISLRNFRSILSWELKNHSIVPTHYTLLYTIMSKPEDLKVVKN CANTTRSFCDLTDEWRSTHEAYVTVLEGFSGNTTLFSCSHNF WLAIDMSFEP Chain 2: ESKYGPPCPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCV VVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREP QVYTLPPSQEEMTKNQVSLSCAVKGFYPSDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLVSRLTVDKSRWQEGNVFSCSVMHE ALHNRFTQKSLSLSPGK

8.2. Materials and Methods

8.2.1. Production of IFN Receptor Agonist Constructs

[0307] Constructs encoding IFN receptor agonists were generated in standard mammalian protein expression DNA vectors (pcDNA3.4 or similar) suitable for high yield protein production and containing standard elements such as promoter sequence, polyA sequence, regulatory elements, and resistance genes. Where applicable, sequences were codon optimized. A 29-amino acid signal sequence from murine inactive tyrosine-protein kinase transmembrane receptor ROR1 (mROR1) was added to the N-termini of the constructs to serve as a signal for secretion. All IFN receptor agonists were expressed as preproteins containing the signal sequence which is cleaved by intracellular processing to produce a mature protein. The constructs were expressed in Expi293FTM cells by transient transfection (Thermo Fisher Scientific). Proteins in Expi293F supernatant were purified using the ProteinMaker system (Protein BioSolutions, Gaithersburg, MD) with either HiTrapTM Protein G HP or MabSelect SuRe pcc columns (Cytiva). After single step elution, the proteins were neutralized, dialyzed into a final buffer of phosphate buffered saline (PBS) with 5% glycerol, aliquoted and stored at -80 °C. Samples were further analyzed by SE-UPLC to determine the presence of high or low molecular weight species relative to the species of interest.

8.2.2. Engineering of reporter KG-1a cells

[0308] The promyeloblast macrophage cell line KG-1a was transduced with an Interferon-Stimulated Response Element (ISRE)-driven luciferase reporter construct and maintained in Iscove's modified Dulbecco's medium supplemented with 2mM L-Glutamine/Penicillin/Streptomycin + 20% FBS + 1μg/mL puromycin. A single cell clone, having high responsiveness to IFNα2b, was isolated. PDL1 expression was knocked out in this clone using CRISPR-Cas9 technology, and the resulting cell line, KG-1a/ISRE-Luc/PDL1 KO (also referred to as PDL1 KO KG-1a cells), was validated by flow cytometry. KG-1a/ISRE-Luc cells were engineered to overexpress PDL1 (amino acids M1-T290 of accession #NP_054862.1), followed by flow sorting for high PDL1 expressing cells, resulting in the cell line KG-1a/ISRE-Luc/hPDL1 (also referred to as PDL1 OE KG-1a cells).

8.2.3. Luciferase assay setup

[0309] RPMI1640 media supplemented with 2mM L-Glutamine/Penicillin/Streptomycin + 10% FBS was used as the assay medium to prepare cell suspensions and fusion protein dilutions.

[0310] The day of the assay, cells were centrifuged and resuspended in assay medium at a density of 5 x 10^5/mL. Recombinant IFNα2b (sometimes referred to as "recombinant IFN" or simply "IFN"), IFNα1, IFNβ, or IFN fusion proteins were diluted 1:5 following a 11-point dilution range (100nM to 10.2fM range or 500nM to 51.2fM range), with the 12^{th} point containing no recombinant protein. 2.5 x 10^4 reporter cells were added to 96-well white flat bottom plates and incubated with serially diluted recombinant IFN or IFN fusion protein. Plates were incubated for 5 hours at 37°C and 5% CO_2, before the addition of 100µL ONE-Glo™ (Promega) reagent to lyse cells and detect luciferase activity. The emitted light was captured in relative light units (RLU) on a multilabel plate reader Envision (PerkinElmer). All serial dilutions were tested in duplicates.

8.2.4. Splenocyte Cultures

[0311] Mice expressing the human IFNAR1 and IFNAR2 receptor were generated in house. Spleens were excised and homogenized. Cell suspensions were lysed with RBC lysis buffer for 5 min, then washed in RPMI1640 supplemented with 10% FBS. Cells were plated at a density of 2.5 x 10^5 cells/ well in a 96-well U bottom plate.

8.2.5. PBMC Cultures

[0312] Human PBMC's were thawed and allowed to recover overnight in RPMI1640 supplemented with 10% FBS. On the day of stimulation, cells were collected and plated at density of 7.5 x 10^4 cells per well in a 96 well U bottom plate.

8.2.6. Detection of pSTAT1

[0313] On the day of stimulation, molecules were diluted to a 11-point dilution range (100nM to 10.2fM) and added to plated splenocytes and PBMCs. Plates were incubated for 20 minutes at 37°C and 5% CO_2. Following stimulation, cells were washed and fixed with BD Cytofix buffer (cat 554655) and incubated at 37°C and 5% CO_2 for 12 minutes. Cells were spun and permeabilized with BD Perm Buffer III (cat 558050) for 10 minutes on ice. Cells were washed twice and stained with cell surface and intracellular antibodies (BD: CD4, B220, CD11b, CD44, CD3, CD8a, NK1.1, pSTAT1) made in BD Horizon Brilliant Buffer (cat

566349) with 2% mouse serum for 60 minutes at room temperature. Cells were washed twice and acquired on a BD Fortessa flow cytometer.

8.3. Example 1: SE-UPLC Profiles of Fc-linked Interferon Molecules

[0314] SE-UPLC was conducted to assess IFN molecules that can be incorporated into the IFN agonists of the disclosure. The four exemplary constructs analyzed with SE-UPLC, Fc-IFNα1 (FIG. 5A), Fc-IFNα2b (FIG. 5B), IFNα2b-Fc (FIG. 5C), and Fc-IFN x Fc (FIG. 5D), displayed discrete main peaks with varying levels of high molecular weight species. The main peak percent area of Fc-IFNα1 was calculated to be 37.43, whereas these percentage values were larger for the Fc-IFNα2b and IFNα2b-Fc, which were calculated to be 57.66 and 56.4, respectively. The largest main peak area percentage value 85, which was observed with Fc-IFN x Fc.

8.4. Example 2: Activity of Interferon Molecules

[0315] An Interferon-Stimulated Response Element (ISRE)-driven luciferase reporter was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate ability of IFN molecules to drive an ISRE-dependent transcriptional response in KG-1a cells.

[0316] The results, shown in FIG. 6, indicated that recombinant protein and Fc fusions of IFN variants show varying degrees of attenuation in an *in vitro* luciferase assay of Interferon-Stimulated Response Element (ISRE). Relative to IFNα2b, all three IFN molecule structures shown in FIG. 6A displayed weaker interferon signaling (FIG. 6B). However, the level of attenuation was more similar for Fc-IFNα2b and IFNα2b-Fc, both of which were associated with somewhat higher attenuation than Fc-IFNα2b x Fc. Next, *in vitro* activity of two Fc-IFN constructs, Fc-IFNα2b and Fc-IFNα1, were compared to the activity of three IFN variants, IFNα2b, IFNα1, and IFNβ (FIG. 6C). Among the IFN variants, the highest level of activity was observed with IFNβ and IFNα2b, whereas the activity of IFNα1 was relatively weaker. Recombinant proteins with Fc fusions displayed an attenuated level of activity compared to the IFN variants. In conclusion, Fc fusion leads to attenuation of interferon signaling relative to free interferon.

8.5. Example 3: SE-UPLC Profiles of Mutant IFN Constructs

[0317] SE-UPLC was conducted to assess mutant IFN molecules that are linked to Fc domains on the C-terminus. The four exemplary constructs analyzed with SE-UPLC, Fc-IFNα2bR33A (FIG. 7A), Fc-IFNα2bR149A (FIG. 7B), Fc-IFNα2bR120A (FIG. 7C), and Fc-IFNα2bS152A (FIG. 7D), displayed discrete main peaks with varying levels of high molecular weight species.

8.6. Example 4: Activity of Mutant IFN Constructs

[0318] The ISRE-driven luciferase reporter was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate ability of mutant IFN constructs to drive an ISRE-dependent transcriptional response in KG-1a cells.

[0319] Activity of IFN variants correlates with their affinity to IFNAR. Hence, mutations that affect the IFN-IFNAR binding can influence the activity of Fc-IFN constructs. A series of mutations were introduced to IFNα2b either on its IFNAR1 or IFNAR2 interface (FIG. 8A and 8B). Relative to wild-type Fc-IFNα2b, most mutations that interfere with IFNAR1 or IFNAR2 binding of Fc-IFNα2b attenuated the ISRE-luciferase activity. Moreover, the degree of this attenuation varied; whereas some mutations caused only a slight attenuation of activity, others led to very high levels of attenuation (FIG. 8B).

8.7. Example 5: SEC Profiles of Exemplary Interferon Receptor Agonist Constructs

[0320] SE-UPLC was conducted to assess the presence of high or low molecular weight species in samples of exemplary IFN receptor agonist constructs as described in Section 8.2.1. FIG. 9 illustrates the profiles of six exemplary IFN receptor agonist constructs described in FIG. 4: Fc-IFNAR1(SD1-3)-IFNα2b (FIG. 9A), Fc-R1(SD1-3)-IFNα2b x Fc (FIG. 9B), Fc-IFNα2b-IFNAR2(D1) (FIG. 9C), Fc-IFNAR2(D1)-IFNα2b (FIG. 9D), Fc-IFNα2b x Fc-R2(D1-2) (FIG. 9E), and Fc-IFNα2b-R2(D1-2) x Fc (FIG. 9F). In general, all six receptor agonist constructs showed discrete main peaks with varying levels smaller peaks that correspond to high molecular weight species.

8.8. Example 6: Activity of Exemplary IFN Receptor Agonist Constructs

[0321] The ISRE-driven luciferase reporter was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate ability of IFN receptor agonist constructs to drive an ISRE-dependent transcriptional response in KG-1a cells.

[0322] The results, shown in FIG. 10, indicated that IFN receptor agonists showed varying degrees of attenuation of ISRE-luciferase activity relative to recombinant, free interferon (IFNα2b; "recombinant IFN" in FIG. 10). Receptor masking attenuated the activity of the wild-type interferon to varying degrees depending on the receptor mask used.

8.9. Example 7: Effect of Masking on the Interferon Receptor Agonist Activity

[0323] An Interferon-Stimulated Response Element (ISRE)-driven luciferase reporter was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate the effect of masking on the ability of IFN fusion proteins to drive an ISRE-dependent transcriptional response in KG-1a cells.

[0324] The results shown in FIG. 11A indicate that masked Fc-IFN constructs attenuated the reporter activity, but with varying potencies. Relative to the reporter response obtained with unfused IFN (IFNα2b; "recombinant IFN" in FIGS. 11A and 11B), both N-terminus and C-terminus Fc fusions of IFN attenuated the reporter activity. Similar potencies were obtained regardless of the N- or C-terminus location of Fc fusion. Fc-IFNAR1(SD1-3)-IFN and Fc-IFNAR1(SD2-3)-IFN masking constructs displayed relatively higher attenuation of the reporter response than Fc-IFN and IFN-Fc, similar to the results observed in FIG. 10. Fc-IFNAR2(D1)-IFN attenuated the reporter response even further than Fc-IFNAR1(SD2-3)-IFN; however, Fc-IFN-IFNAR2(D1) was the most effective construct in attenuating the reporter response, which was approximately 4.7 times less potent than Fc-IFNAR2(D1)-IFN.

[0325] Next, the ability of heterodimeric IFN fusion protein constructs to drive an ISRE-dependent transcriptional response in KG-1a cells was evaluated. The results shown in FIG. 11B reveal that KiH monomeric IFN fusion constructs differ from dimeric Fc-IFN in attenuating the reporter response. As seen in FIG. 11A, IFN-Fc and Fc-IFN attenuated the reporter response with nearly identical potencies. Fc-IFN x Fc was associated with less attenuation than Fc-IFN since the former was associated with nearly 2-fold higher potency in

inducing reporter response than the latter construct. Fc-IFN-R2(D1-2) x Fc and Fc-IFN x Fc-R2(D1-2) displayed similar potencies, indicating that the placement of IFNAR2 mask on the same versus different Fc chain did not have a detectable effect. In contrast, Fc-R1(SD1-3)-IFN x Fc was significantly more effective in attenuating the reporter response than Fc-IFN x Fc-R1(SD1-3), indicating that placement of the IFNAR1(SD1-3) mask on the same Fc chain was associated with a better attenuation.

8.10. Example 8: Activity of IFN Receptor Agonists in hIFNAR-expressing Mouse Cells

[0326] To assess the activity of exemplary IFN receptor agonists, mouse splenocytes were isolated as described in Section 8.2.4. Distinct cell types were evaluated for pSTAT1 presence as described in Section 8.2.6., reported as a percentage of cells positive for pSTAT1.

[0327] In hIFNAR $CD8^+$ T cells, both Fc-IFN and IFN-Fc were associated with an attenuation of the % of pSTAT1 response relative to the % of pSTAT1 response obtained with the increasing concentrations of unfused IFN (IFNα2b). These attenuations displayed a similar potency regardless of the N- or C-terminus location of Fc fusion (FIG. 12A). Fc-IFNAR2(D1)-IFN and Fc-IFN-IFNAR2(D1) attenuated the response further than Fc-IFN and IFN-Fc; however, Fc-IFNAR1(SD1-3)-IFN was the most effective construct in attenuating the response (FIG. 12A). The assessment of KiH fusion constructs revealed that Fc-IFN-R2(D1-2) x Fc and Fc-IFN x Fc-R2(D1-2) displayed similar potencies, indicating that the placement of IFNAR2 mask on the same versus different Fc chain did not have a detectable effect (FIG. 12B). In contrast, Fc-R1(SD1-3)-IFN x Fc was significantly more effective in attenuating the pSTAT1 response than Fc-IFN x Fc-R1(SD1-3), indicating that placement of the IFNAR1(SD1-3) mask on the same Fc chain was associated with a better attenuation (FIG. 12B).

[0328] In hIFNAR $CD11b^+$ cells, the responses obtained with both Fc-IFN and IFN-Fc were associated with an attenuation of the % of pSTAT1 response relative to the % of pSTAT1 response obtained with the increasing concentrations of unfused IFN (IFNα2b), in which the attenuation associated with IFN-Fc was more pronounced (FIG. 13A). Fc-IFNAR2(D1)-IFN, Fc-IFN-IFNAR2(D1), and Fc-IFNAR1(SD1-3)-IFN attenuated the response even further (FIG. 13A). The assessment of heterodimeric KiH fusion constructs revealed that Fc-IFN-

R2(D1-2) x Fc was associated with a slightly less pronounced attenuation than Fc-IFN x Fc-R2(D1-2), indicating that the placement of IFNAR2 mask on different Fc chains may be associated with a better attenuation (FIG. 13B). In contrast, Fc-R1(SD1-3)-IFN x Fc was significantly more effective in attenuating the pSTAT1 response than Fc-IFN x Fc-R1(SD1-3), indicating that placement of the IFNAR1(SD1-3) mask on the same Fc chain was associated with a better attenuation (FIG. 13B).

[0329] In hIFNAR $CD4^+$ T cells, both Fc-IFN and IFN-Fc were associated with an attenuation of the % of pSTAT1 response relative to the % of pSTAT1 response obtained with the increasing concentrations of unfused IFN (IFNα2b). These attenuations displayed a similar potency regardless of the N- or C-terminus location of Fc fusion (FIG. 14A). Fc-IFNAR2(D1)-IFN and Fc-IFN-IFNAR2(D1) attenuated the response further than Fc-IFN and IFN-Fc; however, Fc-IFNAR1(SD1-3)-IFN was the most effective construct in attenuating the response (FIG. 14A). The assessment of heterodimeric KiH fusion constructs revealed that Fc-IFN-R2(D1-2) x Fc was associated with a slightly less pronounced attenuation than Fc-IFN x Fc-R2(D1-2), indicating that the placement of IFNAR2 mask on different Fc chains may be associated with a better attenuation (FIG. 14B). In contrast, Fc-R1(SD1-3)-IFN x Fc was significantly more effective in attenuating the pSTAT1 response than Fc-IFN x Fc-R1(SD1-3), indicating that placement of the IFNAR1(SD1-3) mask on the same Fc chain was associated with a better attenuation (FIG. 14B).

[0330] Lastly, in hIFNAR NK cells, both Fc-IFN and IFN-Fc were associated with an attenuation of the % of pSTAT1 response relative to the % of pSTAT1 response obtained with the increasing concentrations of unfused IFN (IFNα2b). These attenuations displayed a similar potency regardless of the N- or C-terminus location of Fc fusion (FIG. 15A). Fc-IFNAR2(D1)-IFN and Fc-IFN-IFNAR2(D1) attenuated the response further than Fc-IFN and IFN-Fc; however, Fc-IFNAR1(SD1-3)-IFN was the most effective construct in attenuating the response (FIG. 15A). The assessment of heterodimeric KiH fusion constructs revealed that Fc-IFN-R2(D1-2) x Fc was associated with a less pronounced attenuation than Fc-IFN x Fc-R2(D1-2), indicating that the placement of IFNAR2 mask on different Fc chains was associated with a better attenuation (FIG. 14B). In contrast, Fc-R1(SD1-3)-IFN x Fc was significantly more effective in attenuating the pSTAT1 response than Fc-IFN x Fc-R1(SD1-

3), indicating that placement of the IFNAR1(SD1-3) mask on the same Fc chain was associated with a better attenuation (FIG. 15B).

8.11. Example 9: Activity of IFN Receptor Agonists in Human Peripheral Blood Mononuclear Cells

[0331] To assess the activity of exemplary IFN receptor agonists, human PBMC were isolated as described in Section 8.2.5. Distinct cell types were evaluated for pSTAT1 presence as described in Section 8.2.6, reported as a percentage of cells positive for pSTAT1.

[0332] Both Fc-IFN and Fc-IFN x Fc were associated with similar levels of attenuation of the % of pSTAT1 response relative to the % of pSTAT1 response obtained with the increasing concentrations of unfused IFN in PBMC $CD8^+$ T cells (FIG. 16A) and PBMC NK cells (FIG. 16B). Responses associated with Fc-IFNAR1(SD1-3)-IFN and Fc-R1(SD1-3)-IFN x Fc were even more severely attenuated in both cell types (FIGS. 16A and 16B).

8.12. Example 10: Activity of Exemplary Monovalent and Bivalent Masked IFN Receptor Agonists

[0333] The ISRE-driven luciferase reporter was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate the ability of IFN receptor agonists to drive an ISRE-dependent transcriptional response in KG-1a cells. The monovalent and bivalent IFN receptor agonists used in this evaluation that are single- or dual-masked are set forth in Table 7 below.

Table 7

Construct	Mask Type	Construct Format	Visual Representation
Fc-IFNα2b x Fc	Unmasked	Monovalent	FIG. 2G
Fc-IFNα2b x Fc-R1(SD1-3)	Single-masked	Monovalent	FIG. 2H
Fc-R1(SD1-3)-IFNα2b x Fc	Single-masked	Monovalent	FIG. 2J
Fc-IFNα2b x Fc-R2(D1-2)	Single-masked	Monovalent	FIG. 2I
Fc-IFNα2b-R2(D1-2) x Fc	Single-masked	Monovalent	FIG. 2K
Fc-IFNα2b x Fc-R2(D1)	Single-masked (short R2 mask)	Monovalent	FIG. 2I
Fc-R2(D1)-IFNα2b x Fc	Single-masked (short R2 mask)	Monovalent	FIG. 2W
Fc-IFNα2b-R2(D1) x Fc	Single-masked (short R2 mask)	Monovalent	FIG. 2K
Fc-R2(D1)-IFNα2b x Fc-R1(SD1-3)	Dual-masked (short R2 mask)	Monovalent	FIG. 2Q

Table 7			
Construct	**Mask Type**	**Construct Format**	**Visual Representation**
Fc-R1(SD1-3)-IFNα2b x Fc-R2(D1)	Dual-masked (short R2 mask)	Monovalent	FIG. 2P
Fc-R1(SD1-3)-IFNα2b-R2(D1) x Fc	Dual-masked (short R2 mask)	Monovalent	FIG. 2N
Fc-R1(SD1-3)-IFNα2b x Fc-R2(D1-2)	Dual-masked	Monovalent	FIG. 2P
Fc-R1(SD1-3)-IFNα2b	Single-masked	Bivalent	FIG. 2C
Fc-R2(D1)-IFNα2b	Single-masked (short R2 mask)	Bivalent	FIG. 2E
Fc-IFNα2b-R2(D1)	Single-masked (short R2 mask)	Bivalent	FIG. 2F
Fc-R1(SD1-3)-IFNα2b-R2(D1)	Dual-masked (short R2 mask)	Bivalent	FIG. 2L

[0334] The results, shown in FIG. 17, indicated that monovalent IFN receptor agonists showed varying degrees of attenuation of ISRE activation relative to IFNα2b, depending on the type and orientation of the receptor mask. Dual-masked construct Fc-R1(SD1-3)-IFNα2b-R2(D1-2) x Fc showed the strongest signal attenuation.

[0335] Similarly, bivalent IFN receptor agonists also showed varying degrees of attenuation of ISRE activation (FIG. 18). In this evaluation, attenuation of ISRE activation was assessed using a dual-masked bivalent and three single-masked bivalent IFN receptor agonists (Table 7), wherein the dual masked construct displayed increased attenuation relative to the single-masked constructs (FIG. 18).

8.13. Example 11: Effect of PDL1 Targeting on the Potency of IFN Receptor Agonists

[0336] The ISRE-driven luciferase reporter assay was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate the ability of PDL1 targeted IFN receptor agonists to drive an ISRE-dependent transcriptional response in KG-1a cells. Isotype ("Iso") or PDL1 targeted, monovalent and bivalent IFN receptor agonists used in this evaluation are set forth in Table 8 below.

Table 8			
Construct	**Mask Type**	**Construct Format**	**Visual Representation**
Iso-IFNα2b x Iso	Unmasked	Monovalent	FIG. 3G
aPDL1-IFNα2b x aPDL1	Unmasked	Monovalent	FIG. 3G
Iso-IFNα2b-R2(D1-2) x Iso	Single-masked	Monovalent	FIG. 3K
aPDL1-IFNα2b-R2(D1-2) x aPDL1	Single-masked	Monovalent	FIG. 3K
Iso-IFNα2b x Iso-R2(D1-2)	Single-masked	Monovalent	FIG. 3I
aPDL1-IFNα2b x aPDL1-R2(D1-2)	Single-masked	Monovalent	FIG. 3I
Iso-R2(D1)-IFNα2b	Single-masked (short R2 mask)	Bivalent	FIG. 3E
aPDL1-R2(D1)-IFNα2b	Single-masked (short R2 mask)	Bivalent	FIG. 3E

[0337] In PDL1 OE KG-1a cells, PDL1 targeted monovalent IFN receptor agonist constructs displayed enhanced potency relative to their non-targeted (isotype) counterparts (FIG. 19A). This difference in potency between PDL1 targeted and non-targeted constructs was absent in PDL1 KO KG-1a cells (FIG. 19B). Similar results were observed with a PDL1 targeted bivalent IFN receptor agonist construct relative to its isotype counterpart (FIGS. 19C and 19D).

8.14. Example 12: Effect of Linker Length on the Potency of IFN Receptor Agonists

[0338] The ISRE-driven luciferase reporter assay was incorporated into the promyeloblast macrophage cell line KG-1a as described in Section 8.2.2 and was used as described in Section 8.2.3 to evaluate the effect of linker length on the ability of a PDL1 targeted IFN receptor agonists to drive an ISRE-dependent transcriptional response in KG-1a cells.

[0339] FIGS. 20A-20D show the structure of IFN receptor agonist constructs evaluated in this assessment, where the linkers between the IFNα2b and masking moieties in a dual-masked monovalent PDL1 targeted or isotype constructs varied between 5 and 20 amino acids. PDL1 targeted constructs with various linker lengths resulted in similar levels of ISRE-luciferase activity in PDL1 OE KG-1a cells (FIG. 20E), suggesting that the potency of the IFN receptor-masked construct was not affected by increasing the lengths of the linkers.

9. CITATION OF REFERENCES

[0340] All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes. In the event that there is an inconsistency between the teachings of one or more of the references incorporated herein and the present disclosure, the teachings of the present specification are intended.

CLAIMS

1. A Type I interferon (IFN) receptor agonist, comprising:

(a) a first polypeptide chain comprising a first Fc domain and a Type I interferon (IFN) moiety attenuated by masking by an interferon alpha receptor 1 (IFNAR1) moiety and an interferon alpha receptor 2 (IFNAR2) moiety; and

(b) a second polypeptide chain comprising a second Fc domain associated with the first Fc domain.

2. The IFN receptor agonist of claim 1, wherein the IFN moiety is N-terminal to the first Fc domain.

3. The IFN receptor agonist of claim 1, wherein the IFN moiety is C-terminal to the first Fc domain.

4. The IFN receptor agonist of any one of claims 1 to 3, wherein the first polypeptide chain comprises the IFNAR1 moiety.

5. The IFN receptor agonist of claim 4, wherein the IFNAR1 moiety is N-terminal to the IFN moiety.

6. The IFN receptor agonist of claim 4, wherein the IFNAR1 moiety is C-terminal to the IFN moiety.

7. The IFN receptor agonist of any one of claims 1 to 6, wherein the first polypeptide chain comprises the IFNAR2 moiety.

8. The IFN receptor agonist of claim 7, wherein the IFNAR2 moiety is N-terminal to the IFN moiety.

9. The IFN receptor agonist of claim 7, wherein the IFNAR2 moiety is C-terminal to the IFN moiety.

10. The IFN receptor agonist of any one of claims 1 to 6, further comprising one or more linkers connecting two or more of the first Fc domain, the IFN moiety, the IFNAR1 moiety, and the IFNAR2 moiety.

11. The IFN receptor agonist of claim 1, wherein the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR1 moiety, the IFN moiety, and the IFNAR2 moiety.

12. The IFN receptor agonist of claim 11, further comprising a first linker connecting the first Fc domain and the first IFNAR1 moiety, a second linker connecting the IFNAR1 moiety and the IFN moiety, and a third linker connecting the IFN moiety and the IFNAR2 moiety.

13. The IFN receptor agonist of claim 1, wherein the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, a first linker, the IFNAR2 moiety, a second linker, the IFN moiety, a third linker, and the IFNAR1 moiety.

14. The IFN receptor agonist of claim 13, further comprising a first linker connecting the first Fc domain and the first IFNAR2 moiety, a second linker connecting the IFNAR2 moiety and the IFN moiety, and a third linker connecting the IFN moiety and the IFNAR1 moiety.

15. The IFN receptor agonist of claim 1, wherein (i) the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR2 moiety, and the IFN moiety, and (ii) the second polypeptide comprises, in N- to C-terminal orientation, the second Fc domain, and the IFNAR1 moiety.

16. The IFN receptor agonist of claim 15, further comprising a first linker connecting the first Fc domain and the IFNAR2 moiety, a second linker connecting the IFNAR2 moiety and the IFN moiety, and a third linker connecting the second Fc domain and the IFNAR1 moiety.

17. The IFN receptor agonist of claim 1, wherein (i) the first polypeptide comprises, in N- to C-terminal orientation, the first Fc domain, the IFNAR1 moiety, and the IFN moiety, and (ii) the second polypeptide comprises, in N- to C-terminal orientation, the second Fc domain and the IFNAR2 moiety.

18. The IFN receptor agonist of claim 17, further comprising a first linker connecting the first Fc domain and the IFNAR1 moiety, a second linker connecting the IFNAR1 moiety and the IFN moiety, and a third linker connecting the second Fc domain and the IFNAR2 moiety.

19. The IFN receptor agonist of any one of claims 12, 14, 16, or 18, wherein one or more of the first linker, the second linker, and the third linker is a protease-cleavable linker (PCL).

20. The IFN receptor agonist of claim 19, wherein the second linker is a PCL.

21. The IFN receptor agonist of claim 19 or 20, wherein the PCL comprises a substrate sequence cleavable by any protease set forth in Table A.

22. The IFN receptor agonist of any one of claims 19 to 21, wherein the PCL comprises one or more substrate sequences selected from the substrate sequences set forth in Table B.

23. The IFN receptor agonist of any one of claims 19 to 22, wherein the PCL comprises one or more spacer sequences selected from the spacer sequences set forth in Table C.

24. The IFN receptor agonist of any one of claims 19 to 23, wherein the PCL comprises the amino acid sequence of any of the PCL sequences set forth in Table D or a variant thereof with up to 5 amino acid substitutions.

25. The IFN receptor agonist of any one of claims 19 to 24, which is configured such that cleavage of the protease-cleavable linker (PCL) unmasks the IFN moiety.

26. The IFN receptor agonist of any one of claims 1 to 25, wherein the second polypeptide chain comprises an additional IFN moiety masked by an additional IFNAR1 moiety and an additional IFNAR2 moiety.

27. The IFN receptor agonist of any one of claims 1 to 26, wherein the IFN moiety comprises an amino acid sequence having at least about 90%, at least about 95%, or at least about 98% sequence identity to (a) full length mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ or (b) a mature human IFNα1, IFNα2b, IFNβ, IFNω, IFNε or IFNκ having up to a 15-amino acid truncation at its N-terminus and/or its C-terminus.

28. The IFN receptor agonist of any one of claims 1 to 27, wherein the IFN moiety comprises an amino acid sequence having one or more attenuating mutations as compared to mature human IFNα1 or IFNα2b.

29. The IFN receptor agonist of any one of claims 1 to 28, wherein the IFN moiety has one or more mutations selected from L26A, F27A, R33A, R33K, L30A, D35E, H57Y, E58N, Q61S, H57S, E58S, H57A, E58A, Q61A, Q90A, E96A, R120A, L135A, R144A, R144S, R144T, R144Y, R144I, R144L, A145D, A145H, A145K, A145M, A145V, A145Y, R149A, R149K, S152A, R162A, and E165D.

30. The IFN receptor agonist of any one of claims 1 to 29, wherein the IFNAR1 moiety comprises an amino acid sequence having at least 90%, at least 95%, or at least 98% sequence identity to (i) the SD2 and SD3 domains of human IFNAR1, (ii) the SD1, SD2 and SD3 domains of human IFNAR1, or (iii) the SD1, SD2, SD3 and SD4 domains of human IFNAR1.

31. The IFN receptor agonist of any one of claims 1 to 30, wherein the IFNAR2 moiety comprises an amino acid sequence having at least 90%, at least 95%, or at least

98% sequence identity to (i) the D1 domain of human IFNAR2 or (ii) the D1 and D2 domains of human IFNAR2.

32. The IFN receptor agonist according to any one of claims 1 to 31, wherein the first Fc domain and/or the second Fc domain comprises a hinge domain.

33. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2N.

34. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2L.

35. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2M.

36. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2O.

37. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2P.

38. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2Q.

39. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2U.

40. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 1 to 32, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 2V.

41. The IFN receptor agonist of any one of claims 1 to 40, which comprises a first targeting moiety that binds to a first target molecule and optionally a second targeting moiety that binds to a second target molecule.

42. The IFN receptor agonist of claim 41, wherein the first targeting moiety and optional second targeting moiety are antibodies or antigen-binding fragments thereof.

43. The IFN receptor agonist of claim 41 or 42, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to an extracellular matrix (ECM) antigen, a tumor reactive lymphocyte antigen, a cell surface molecule of tumor or viral lymphocytes, a T-cell antigen (TCA), a checkpoint inhibitor, a tumor-associated antigen (TAA), a dendritic cell (DC) or other antigen-presenting cell (APC) antigen, or a natural killer (NK) cell antigen.

44. The IFN receptor agonist of any one of claims 41 to 43, wherein the first targeting moiety and/or optional second targeting moiety (a) comprises the (i) CDR or (ii) VH and VL sequences of antibody set forth in Table F or (b) competes with the antibody set forth in Table F for binding to the target molecule.

45. The IFN receptor agonist of any one of claims 41 to 44, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to an ECM antigen which is optionally selected from syndecan, heparanase, integrins, osteopontin, link, cadherins, laminin, laminin type EGF, lectin, fibronectin, notch, nectin (*e.g.*, nectin-4), tenascin, collagen (*e.g.*, collagen type X) and matrixin.

46. The IFN receptor agonist of any one of claims 41 to 44, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a cell surface molecule of tumor or viral lymphocytes.

47. The IFN receptor agonist of claim 46, wherein the cell surface molecule is a T-cell co-stimulatory protein, optionally selected from CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, and B7-H3.

48. The IFN receptor agonist of any one of claims 41 to 44, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a checkpoint inhibitor.

49. The IFN receptor agonist of claim 48, wherein the checkpoint inhibitor is PDL1.

50. The IFN receptor agonist of claim 48, wherein the checkpoint inhibitor is PD1.

51. The IFN receptor agonist of any one of claims 41 to 44, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a tumor-associated antigen (TAA), optionally selected from AFP, ALK, a BAGE protein, BIRC5 (survivin), BIRC7, β-catenin, brc-abl, BRCA1, BORIS, CA9, carbonic anhydrase IX, caspase-8, CALR, CEACAM5 (also known as carcinoembryonic antigen or CEA), CCR5, CD19, CD20 (MS4A1), CD22, CD30, CD40, CDK4, CEA, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, ErbB2/Her2, ErbB3, ErbB4, ETV6-AML, EpCAM, EphA2, Fra-1, FOLR1, a GAGE protein (*e.g.*, GAGE-1 or -2), GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/k-ras, HLA/MAGE-A3, hTERT, LMP2, MAGE proteins (*e.g.*, MAGE-1, -2, -3, -4, -6, and -12), MART-1, mesothelin, ML-IAP, Muc1, Muc2, Muc3, Muc4, Muc5, Muc16 (CA-125), MUM1, NA17, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PLAC1, PRLR, PRAME, PSMA (FOLH1), RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, STEAP1, STEAP2, TAG-72, TGF-β, TMPRSS2, Thompson-nouvelle antigen (Tn), TRP-1, TRP-2, tyrosinase, and uroplakin-3.

52. The IFN receptor agonist of any one of claims 41 to 44, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a dendritic cell (DC) or other antigen-presenting cell (APC) antigen which is optionally selected from XCR1, Clec9a, CD1c, CD11c, CD14, macrophage mannose receptor (CD206), and DEC-205.

53. The IFN receptor agonist of any one of claims 41 to 44, wherein the first targeting moiety and/or optional second targeting moiety is capable of binding to a natural killer (NK) cell antigen.

54. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3N.

55. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3L.

56. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3M.

57. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3O.

58. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3P.

59. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3Q.

60. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3U.

61. An IFN receptor agonist, which is optionally an IFN receptor agonist according to any one of claims 41 to 53, which comprises polypeptide chains having the configuration of the two half-antibodies illustrated in FIG. 3V.

62. A nucleic acid or plurality of nucleic acids encoding the IFN receptor agonist of any one of claims 1 to 61.

63. A host cell engineered to express the IFN receptor agonist of any one of claims 1 to 61 or the nucleic acid(s) of claim 62.

64. A method of producing the IFN receptor agonist of any one of claims 1 to 61, comprising culturing the host cell of claim 63 and recovering the IFN receptor agonist expressed thereby.

65. A pharmaceutical composition comprising the IFN receptor agonist of any one of claims 1 to 61 and an excipient.

66. A method of treating cancer, comprising administering to a subject in need thereof the IFN receptor agonist of any one of claims 1 to 61 or the pharmaceutical composition of claim 65.

67. A method of localized delivery of an IFN protein, comprising administering to a subject an IFN receptor agonist according to any one of claims 1 to 61 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which

has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue to which the IFN protein is to be locally delivered.

68. A method of treating cancer with an IFN protein that is selectively activated in cancer tissue, comprising administering to a subject in need thereof an IFN receptor agonist according to any one of claims 1 to 61 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by cancer tissue.

69. A method of administering to the subject IFN therapy with reduced systemic exposure and/or reduced systemic toxicity, comprising administering to a subject the IFN therapy in the form of an IFN receptor agonist according to any one of claims 1 to 61 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) which has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed by a tissue for which IFN therapy is desirable and/or intended.

70. A method of targeted delivery of an activated IFN protein to cancer tissue, comprising administering to a subject an IFN receptor agonist according to any one of claims 1 to 61 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient), wherein the IFN receptor agonist:

(a) comprises one or more targeting moieties that recognize a target molecule expressed by the cancer tissue or associated immune cells; and

(b) has one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in a tissue for which IFN therapy is desirable and/or intended.

71. A method of locally inducing an immune response in a target tissue, comprising administering to a subject an IFN receptor agonist according to any one of claims 1 to 61 (or a pharmaceutical composition comprising the IFN receptor agonist and an

excipient) which has one or more targeting moieties capable of binding a target molecule expressed in the target tissue and one or more protease-cleavable linkers, each comprising one or more substrates for one or more proteases expressed in the target tissue.

72. A method of enhancing an immune response against an antigen, comprising administering to a subject an immunogenic agent that elicits an immune response against the antigen together with an IFN receptor agonist according to any one of claims 1 to 61 (or a pharmaceutical composition comprising the IFN receptor agonist and an excipient) or a nucleic acid encoding such IFN receptor agonist.

73. The method of any one of claims 66 to 72, wherein the administration is non-local.




FIG. 1




FIG. 2A




FIG. 2B



CH2
CH2
CH3
CH3
CH2
CH2
CH3
CH3
cis
trans

FIG. 2C



CH2
CH2
CH3
CH3
CH2
CH2
CH3
CH3
cis
trans

FIG. 2D




FIG. 2E




FIG. 2F




FIG. 2G




FIG. 2H




FIG. 2I




FIG. 2J




FIG. 2K




FIG. 2L




FIG. 2M




FIG. 2N




FIG. 2O




FIG. 2P




FIG. 2Q




FIG. 2R




FIG. 2S




FIG. 2T




FIG. 2U




FIG. 2V




FIG. 2W




FIG. 2X




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E




FIG. 3F




FIG. 3G




FIG. 3H




FIG. 3I




FIG. 3J




FIG. 3K




FIG. 3L




FIG. 3M




FIG. 3N




FIG. 3O




FIG. 3P




FIG. 3Q




FIG. 3R




FIG. 3S




FIG. 3T




FIG. 3U




FIG. 3V




FIG. 3W




FIG. 3X

Molecule	Chain 1	Chain 2
Fc-IFN	Fc-IFN	Fc-IFN
Fc-IFNAR1(SD1-3)-IFN	Fc-IFNAR1(SD1-3)-IFN	Fc-IFNAR1(SD1-3)-IFN
Fc-IFNAR1(SD2-3)-IFN	Fc-IFNAR1(SD2-3)-IFN	Fc-IFNAR1(SD2-3)-IFN
Fc-IFN-IFNAR2(D1-2)	Fc-IFN-IFNAR2(D1-2)	Fc-IFN-IFNAR2(D1-2)
Fc-IFN-IFNAR2(D1)	Fc-IFN-IFNAR2(D1)	Fc-IFN-IFNAR2(D1)
Fc-IFNAR2(D1)-IFN	Fc-IFNAR2(D1)-IFN	Fc-IFNAR2(D1)-IFN
Fc-IFN x Fc	Fc(knob)-IFN	Fc(Hole*)
Fc-IFN x Fc-R1(SD1-3)	Fc(knob)-IFN	Fc(Hole*)-R1(SD1-3)
Fc-IFN x Fc-R2(D1-2)	Fc(knob)-IFN	Fc(Hole*)-R2(D1-2)
Fc-R1-IFN x Fc	Fc(knob)-R1-IFN	Fc(Hole*)
Fc-IFN-R2(D1-2) x Fc	Fc(knob)-IFN-R2(D1-2)	Fc(Hole*)
Fc-IFN x Fc-R2(D1)	Fc(knob)-IFN	Fc(Hole*)-IFNAR2(D1)
Fc-R2(D1)-IFN x Fc	Fc(knob)-IFNAR2(D1)-IFN	Fc(Hole*)

FIG. 4A

Molecule	Chain 1	Chain 2
Fc-R1(SD1-3)-IFN x Fc-R2(D1-2)	Fc(knob)-IFNAR1(SD1-3)-IFN	Fc(hole*)-IFNAR2(D1-2)
Fc-R2(D1)-IFN x Fc-R1(SD1-3)	Fc(knob)-IFNAR2(D1)-IFN	Fc(hole*)-IFNAR1(SD1-3)
Fc-IFNAR1(SD1-3)-IFN-IFNAR2(D1)	Fc-IFNAR1(SD1-3)-IFN-IFNAR2(D1)	Fc-IFNAR1(SD1-3)-IFN-IFNAR2(D1)
Fc-R1(SD1-3)-IFN x Fc-R2(D1)	Fc(knob)-IFNAR1(SD1-3)-IFN	Fc(hole*)-IFNAR2(D1)
Fc-R1(SD1-3)-IFN-R2(D1) x Fc	Fc(knob)-IFNAR1(SD1-3)-IFN-IFNAR2(D1)	Fc(hole*)
Fc-IFN-R2(D1) x Fc	Fc(knob)-IFN-IFNAR2(D1)	Fc(hole*)
Fc x IFN-Fc	Fc(knob)	IFN-Fc(hole*)
Fc x R1(SD1-3)-IFN-Fc	Fc(knob)	IFNAR1(SD1-3)-IFN-Fc(hole*)
Fc x R2(D1)-IFN-Fc	Fc(knob)	IFNAR2(D1)-IFN-Fc(hole*)
Fc x R1(SD1-3)-IFN-R2(D1)-Fc	Fc(knob)	IFNAR1(SD1-3)-IFN-IFNAR2(D1)-Fc(hole*)

FIG. 4B



Fc-IFNα1
8.312
7.431
7.139
6.719
5.975
AU
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
Minutes
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00

FIG. 5A



Fc-IFNα2b
8.270
7.134
6.739
AU
0.00
0.10
0.20
0.30
0.40
Minutes
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00

FIG. 5B



IFNα2b-Fc
8.231
7.280
6.716
6.397
6.225
AU
0.10
0.08
0.06
0.04
0.02
0.00
Minutes
3.00
3.50
4.00
4.50
5.00
5.50
6.00
6.50
7.00
7.50
8.00
8.50
9.00
9.50

FIG. 5C



Fc-IFN x Fc
8.695
7.727
AU
0.10
0.08
0.06
0.04
0.02
0.00
Minutes
3.00
3.50
4.00
4.50
5.00
5.50
6.00
6.50
7.00
7.50
8.00
8.50
9.00
9.50

FIG. 5D



Fc-IFN
IFN-Fc
Fc-IFN x Fc

FIG. 6A



IFNα2b
Fc-IFNα2b
IFNα2b-Fc
Fc-IFNα2b x Fc
4x10^5
3x10^5
2x10^5
1x10^5
0
ISRE-Luc activity (RLU)
-15
-14
-13
-12
-11
-10
-9
-8
-7
-6
rProtein concentration Log10[M]

FIG. 6B



IFNα2b
Fc-IFNα2b
IFNα1
Fc-IFNα1
IFNβ
ISRE-Luc activity (RLU)
5x10^5
4x10^5
3x10^5
2x10^5
1x10^5
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10[M]

FIG. 6C



Fc-IFNα2bR33A
8.286
7.267
7.122
6.702
AU
0.50
0.40
0.30
0.20
0.10
0.00
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
Minutes

FIG. 7A



Fc-IFNα2bR149A
8.266
7.303
7.103
6.706
AU
0.40
0.30
0.20
0.10
0.00
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
Minutes

FIG. 7B



Fc-IFNα2bR120A
8.263
7.276
7.118
6.717
AU
0.35
0.30
0.25
0.20
0.15
0.10
0.05
0.00
Minutes
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00

FIG. 7C



Fc-IFNα2bS152A
8.296
7.298
7.130
6.730
AU
0.10
0.08
0.06
0.04
0.02
0.00
Minutes
3.00
3.50
4.00
4.50
5.00
5.50
6.00
6.50
7.00
7.50
8.00
8.50
9.00
9.50
10.00

FIG. 7D



Fc-IFN (WT)
IFN-Fc (Mut)

FIG. 8A



ISRE-Luc activity (RLU)
5x10^5
4x10^5
3x10^5
2x10^5
1x10^5
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10[M]
IFNα2b
Fc-IFNα2b
Fc-IFNα2bR33A
Fc-IFNα2bR149A
Fc-IFNα2bA145M
Fc-IFNα2bR149K
Fc-IFNα2bR33K
Fc-IFNα2bS152A
Fc-IFNα2bR120A
Fc-IFNα2bQ90A
Fc-IFNα2bE96A
Fc-IFNα2bQ90AR120A
IFNAR2 interface
IFNAR1 interface

FIG. 8B



Fc-IFNAR1(SD1-3)IFNα2b
6.677
6.084
AU
Minutes
0.00
0.10
0.20
0.30
0.40
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00
14.00
15.00

FIG. 9A



Fc-R1(SD1-3)IFNα2b x Fc
7.324
6.433
6.140
AU
Minutes
0.00
0.20
0.40
0.60
0.80
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00
14.00
15.00

FIG. 9B



Fc-IFNα2b-IFNAR2(D1)
7.269
6.337
6.061
AU
Minutes
0.00
0.05
0.10
0.15
0.20
0.25
0.30
0.35
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00
14.00
15.00

FIG. 9C



Fc-IFNAR2(D1)-IFNα2b
7.323
6.466
6.089
AU
Minutes
0.00
0.20
0.40
0.60
0.80
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00
14.00
15.00

FIG. 9D



Fc-IFNα2b x Fc-R2(D1-2)
7.638
6.663
6.272
AU
0.00
0.20
0.40
0.60
0.80
Minutes
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00
14.00
15.00

FIG. 9E



Fc-IFNα2b-R2(D1-2) x Fc
7.588
6.596
6.245
AU
0.00
0.20
0.40
0.60
0.80
Minutes
0.00
1.00
2.00
3.00
4.00
5.00
6.00
7.00
8.00
9.00
10.00
11.00
12.00
13.00
14.00
15.00

FIG. 9F



Recombinant IFN
Fc-IFN
Fc-IFNAR1(SD1-3)-IFN
Fc-IFNAR1(SD2-3)-IFN
Fc-IFN-IFNAR2(D1)
Fc-IFN x Fc-R1(SD1-3)
Fc-IFN x Fc-R2(D1-2)
Fc-R1-IFN x Fc
Fc-IFN-R2(D1-2) x Fc
ISRE-Luc activity (RLU)
5x10^5
4x10^5
3x10^5
2x10^5
1x10^5
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10[M]

FIG. 10



Recombinant IFN
Fc-IFN
IFN-Fc
Fc-IFNAR2(D1)-IFN
Fc-IFN-IFNAR2(D1)
Fc-IFNAR1(SD1-3)-IFN
Fc-IFNAR1(SD2-3)-IFN
Relative ISRE-Luc activity (RLU)
0
20
40
60
80
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10[M]

FIG. 11A



Recombinant IFN
Fc-IFN
IFN-Fc
Fc-IFN x Fc
Fc-IFN x Fc-R1(SD1-3)
Fc-IFN x Fc-R2(D1-2)
Fc-R1-IFN x Fc
Fc-IFN-R2(D1-2) x Fc
ISRE-Luc activity (RLU)
4x10^5
3x10^5
2x10^5
1x10^5
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
-6
rProtein concentration Log10[M]

FIG. 11B



IFN
Fc-IFN
IFN-Fc
Fc-IFNAR2(D1)-IFN
Fc-IFN-IFNAR2(D1)
Fc-IFNAR1(SD1-3)-IFN
hIFNAR CD8+ T Cell
% of pSTAT1+
0
20
40
60
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
Conc [M]

FIG. 12A



hIFNAR CD8+ T Cell
% of pSTAT1+
60
40
20
0
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
Conc [M]
IFN
Fc-IFN
Fc-IFN x Fc
Fc-IFN x Fc-R1(SD1-3)
Fc-R1(SD1-3)-IFN x Fc
Fc-IFN x Fc R2(D1-2)
Fc-IFN-R2(D1-2) x Fc

FIG. 12B



hIFNAR CD11b+ Cells
% of pSTAT1+
0
20
40
60
80
Conc [M]
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
IFN
Fc-IFN
IFN-Fc
Fc-IFNAR2(D1)-IFN
Fc-IFN-IFNAR2(D1)
Fc-IFNAR1(SD1-3)-IFN

FIG. 13A



IFN
Fc-IFN
Fc-IFN x Fc
Fc-IFN x Fc-R1(SD1-3)
Fc-R1(SD1-3)-IFN x Fc
Fc-IFN x Fc R2(D1-2)
Fc-IFN-R2(D1-2) x Fc
hIFNAR CD11b+ Cells
% of pSTAT1+
0
20
40
60
80
Conc [M]
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6

FIG. 13B



hIFNAR CD4+ T Cells
% of pSTAT1+
0
20
40
60
80
Conc [M]
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
IFN
Fc-IFN
IFN-Fc
Fc-IFNAR2(D1)-IFN
Fc-IFN-IFNAR2(D1)
Fc-IFNAR1(SD1-3)-IFN

FIG. 14A



hIFNAR CD4+ T Cells
% of pSTAT1+
0
20
40
60
80
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
Conc [M]
IFN
Fc-IFN
Fc-IFN x Fc
Fc-IFN x Fc-R1(SD1-3)
Fc-R1(SD1-3)-IFN x Fc
Fc-IFN x Fc R2(D1-2)
Fc-IFN-R2(D1-2) x Fc

FIG. 14B



hIFNAR NK Cells
% of pSTAT1+
0
20
40
60
80
Conc [M]
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
IFN
Fc-IFN
IFN-Fc
Fc-IFNAR2(D1)-IFN
Fc-IFN-IFNAR2(D1)
Fc-IFNAR1(SD1-3)-IFN

FIG. 15A



hIFNAR NK Cells
% of pSTAT1+
Conc [M]
IFN
Fc-IFN
Fc-IFN x Fc
Fc-IFN x Fc-R1(SD1-3)
Fc-R1(SD1-3)-IFN x Fc
Fc-IFN x Fc R2(D1-2)
Fc-IFN-R2(D1-2) x Fc

FIG. 15B



IFN
Fc-IFN
Fc-IFNAR1(SD1-3)-IFN
Fc-IFN x Fc
Fc-R1(SD1-3)-IFN x Fc
PBMC CD8+ T Cell
% of pSTAT1+
100
80
60
40
20
0
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
Conc [M]

FIG. 16A



PBMC NK Cell
% of pSTAT1+
100
80
60
40
20
0
Conc [M]
10-15
10-14
10-13
10-12
10-11
10-10
10-9
10-8
10-7
10-6
IFN
Fc-IFN
Fc-IFNAR1(SD1-3)-IFN
Fc-IFN x Fc
Fc-R1(SD1-3)-IFN x Fc

FIG. 16B



IFNa2b
Fc-IFNa2b
Fc-IFNa2b x Fc
Fc-IFNa2b x Fc-R1(SD1-3)
Fc-R1(SD1-3)-IFNa2b x Fc
Fc-IFNa2b x Fc-R2(D1-2)
Fc-IFNa2b-R2(D1-2) x Fc
Fc-IFNa2b x Fc-R2(D1)
Fc-R2(D1)-IFNa2b x Fc
Fc-IFNa2b-R2(D1) x Fc
Fc-R2(D1)-IFNa2b x Fc-R1(SD1-3)
Fc-R1(SD1-3)-IFNa2b x Fc-R2(D1)
Fc-R1(SD1-3)-IFNa2b-R2(D1) x Fc
Fc-R1(SD1-3)-IFNa2b x Fc-R2(D1-2)
ISRE-Luc activity (RLU)
6x10^5
4x10^5
2x10^5
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)

FIG. 17



IFNα2b
Fc-IFNα2b
Fc-R1(SD1-3)-IFNα2b
Fc-R2(D1)-IFNα2b
Fc-IFNα2b-R2(D1)
Fc-R1(SD1-3)-IFNα2b-R2(D1)
ISRE-Luc activity (RLU)
6x10^5
4x10^5
2x10^5
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)

FIG. 18



PD-L1 OE
Relative ISRE-Luc activity (RLU)
60
40
20
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)
IFNα2b
Iso-IFNα2b x Iso
Iso-IFNα2b-R2(D1-2) x Iso
Iso-IFNα2b x Iso-R2(D1-2)
aPDL1-IFNα2b x aPDL1
aPDL1-IFNα2b-R2(D1-2) x aPDL1
aPDL1-IFNα2b x aPDL1-R2(D1-2)

FIG. 19A



PD-L1 KO
Relative ISRE-Luc activity (RLU)
0
20
40
60
80
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)
IFNα2b
Iso-IFNα2b x Iso
Iso-IFNα2b-R2(D1-2) x Iso
Iso-IFNα2b x Iso-R2(D1-2)
aPDL1-IFNα2b x aPDL1
aPDL1-IFNα2b-R2(D1-2) x aPDL1
aPDL1-IFNα2b x aPDL1-R2(D1-2)

FIG. 19B



IFNα2b
Iso-IFNα2b x Iso
Iso-R2(D1)-IFNα2b
aPDL1-IFNα2b x aPDL1
aPDL1-R2(D1)-IFNα2b
PD-L1 OE
Relative ISRE-Luc activity (RLU)
60
40
20
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)

FIG. 19C



PD-L1 KO
IFNα2b
Iso-IFNα2b x Iso
Iso-R2(D1)-IFNα2b
aPDL1-IFNα2b x aPDL1
aPDL1-R2(D1)-IFNα2b
Relative ISRE-Luc activity (RLU)
0
20
40
60
80
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)

FIG. 19D




FIG. 20A




FIG. 20B




FIG. 20C




FIG. 20D



PD-L1 OE
IFNα2b
Iso-IFNα2b x Iso
aPDL1-IFNα2b x aPDL1
Iso-R1-20aa-IFN-20aa-R2
aPDL1-R1-5aa-IFN-10aa-R2 x aPDL1
aPDL1-R1-15aa-IFN-15aa-R2 x aPDL1
aPDL1-R1-20aa-IFN-20aa-R2 x aPDL1
Relative ISRE-Luc activity (RLU)
120
100
80
60
40
20
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)

FIG. 20E



PD-L1 KO
IFNα2b
Iso-IFNα2b x Iso
aPDL1-IFNα2b x aPDL1
Iso-R1-20aa-IFN-20aa-R2
aPDL1-R1-5aa-IFN-10aa-R2 x aPDL1
aPDL1-R1-15aa-IFN-15aa-R2 x aPDL1
aPDL1-R1-20aa-IFN-20aa-R2 x aPDL1
Relative ISRE-Luc activity (RLU)
60
40
20
0
-15
-14
-13
-12
-11
-10
-9
-8
-7
rProtein concentration Log10 (M)

FIG. 20F

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2023/072516

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07K14/555 C07K14/715
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2022/155541 A1 (ASKGENE PHARMA INC [US]) 21 July 2022 (2022-07-21) paragraph [0071]; claims 1-21; figure 1	1-73
A	WO 2021/207669 A1 (CYTOMX THERAPEUTICS INC [US]) 14 October 2021 (2021-10-14) figure 5	1-73
A	US 2019/194293 A1 (POSADA JAMES ARTHUR [US] ET AL) 27 June 2019 (2019-06-27) the whole document	1-73
A	US 2020/123227 A1 (FU YANG-XIN [US] ET AL) 23 April 2020 (2020-04-23) claims 1-40	1-73

-/--

[x] Further documents are listed in the continuation of Box C.

[x] See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance;; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance;; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
3 November 2023	21/11/2023

Name and mailing address of the ISA/	Authorized officer
European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Seroz, Thierry

1

Form PCT/ISA/210 (second sheet) (April 2005)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2023/072516

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:

   a. [X] forming part of the international application as filed.

   b. [ ] furnished subsequent to the international filing date for the purposes of international search (Rule 13*ter*.1(a)).

      [ ] accompanied by a statement to the effect that the sequence listing does not go beyond the disclosure in the international application as filed.

2. [ ] With regard to any nucleotide and/or amino acid sequence disclosed in the international application, this report has been established to the extent that a meaningful search could be carried out without a WIPO Standard ST.26 compliant sequence listing.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2023/072516

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HOLDER PATRICK G. ET AL: "Engineering interferons and interleukins for cancer immunotherapy", ADVANCED DRUG DELIVERY REVIEWS, vol. 182, 1 March 2022 (2022-03-01), page 114112, XP093022598, Amsterdam , NL ISSN: 0169-409X, DOI: 10.1016/j.addr.2022.114112 Retrieved from the Internet: URL:https://www.sciencedirect.com/science/article/pii/S0169409X22000023/pdfft?md5=08d56ac50085ae8ec1ca4700081ca389&pid=1-s2.0-S0169409X22000023-main.pdf> the whole document	1-73
A	WO 2022/015711 A1 (IMMUNE TARGETING INC [US]) 20 January 2022 (2022-01-20) paragraphs [0103], [0114], [0125]; claims 1-40; figure 1	1-73
A	CAO XUEZHI ET AL: "Next generation of tumor-activating type I IFN enhances anti-tumor immune responses to overcome therapy resistance", NATURE COMMUNICATIONS, vol. 12, no. 1, 1 December 2021 (2021-12-01), XP055970199, DOI: 10.1038/s41467-021-26112-2 Retrieved from the Internet: URL:https://www.nature.com/articles/s41467-021-26112-2.pdf> the whole document	1-73
A	WO 2022/165443 A1 (ASKGENE PHARMA INC [US]) 4 August 2022 (2022-08-04) paragraph [0103]; claims 1-59; figures 1, 2, 4b	1-73

1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No
PCT/US2023/072516

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2022155541 A1	21-07-2022	EP 4277707 A1	22-11-2023
		WO 2022155541 A1	21-07-2022
WO 2021207669 A1	14-10-2021	AU 2021254283 A1	10-11-2022
		BR 112022020440 A2	27-12-2022
		CA 3174786 A1	14-10-2021
		CN 115667523 A	31-01-2023
		EP 4133085 A1	15-02-2023
		IL 297225 A	01-12-2022
		JP 2023520922 A	22-05-2023
		KR 20220167295 A	20-12-2022
		US 2021317177 A1	14-10-2021
		US 2022402990 A1	22-12-2022
		WO 2021207669 A1	14-10-2021
US 2019194293 A1	27-06-2019	AU 2018321359 A1	05-03-2020
		BR 112020003670 A2	01-09-2020
		CA 3073537 A1	28-02-2019
		CN 111315767 A	19-06-2020
		EP 3672986 A1	01-07-2020
		IL 272749 A	30-04-2020
		JP 2020533020 A	19-11-2020
		JP 2023060304 A	27-04-2023
		KR 20200039778 A	16-04-2020
		SG 11202001311V A	30-03-2020
		US 2019194293 A1	27-06-2019
		US 2021155673 A1	27-05-2021
		WO 2019040674 A1	28-02-2019
US 2020123227 A1	23-04-2020	CA 3067539 A1	27-12-2018
		CN 111050803 A	21-04-2020
		EP 3641829 A1	29-04-2020
		JP 2020528878 A	01-10-2020
		KR 20200015742 A	12-02-2020
		RU 2020101640 A	20-07-2021
		US 2020123227 A1	23-04-2020
		WO 2018236701 A1	27-12-2018
WO 2022015711 A1	20-01-2022	NONE	
WO 2022165443 A1	04-08-2022	NONE	