



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/00		A2	(11) International Publication Number: WO 98/50424
			(43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/US98/09096 (22) International Filing Date: 6 May 1998 (06.05.98) (30) Priority Data: 08/851,974 7 May 1997 (07.05.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/851,974 (CIP) Filed on 7 May 1997 (07.05.97) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US). SHAH, Purvi [IN/US]; 1608 Queen Charlotte Drive #5, Sunnyvale, CA 94087 (US).		(74) Agent: BILLINGS, Lucy, J.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AT, AU, BR, CA, CH, CN, DE, DK, ES, FI, GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, US, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	

(54) Title: HUMAN SERINE PROTEASE PRECURSOR

```

      9      18      27      36      45      54
NAC ACC AAC AGC TCT GAC CTG GGC AGC CTT CCT GAG AAA ATG CAG CCA TTC CTC
                               M  Q  P  F  L

      63      72      81      90      99      108
CTC CTG TTG GCC TTT CTT CTG ACC CCT GGG GCT GGG ACA GAG GAG ATC ATC GGG
L  L  L  A  F  L  L  T  P  G  A  G  T  E  E  I  I  G

      117      126      135      144      153      162
GGC CAT GAG GCC AAG CCC CAC TCC CGC CCC TAC ATG GCC TTT GTT CAG TTT CTC
G  H  E  A  K  P  H  S  R  P  Y  M  A  F  V  Q  F  L

      171      180      189      198      207      216
CAA GAG AAG AGT CGG AAG AGG TGT GGC GGC ATC CTA GTG AGA AAG GAC TTT GTG
Q  E  K  S  R  K  R  C  G  G  I  L  V  R  K  D  F  V

      225      234      243      252      261      270
CTG ACA GCT GCT CAC TGC CAG GGA AGC TCC ATA AAT GTC ACC TTG GGG GCC CAC
L  T  A  A  H  C  Q  G  S  S  I  N  V  T  L  G  A  H

      279      288      297      306      315      324
AAT ATC AAA GAA CAG GAG CCG ACC CAG CAG TTT ATC CCT GTG AAA AGA CCC ATC
N  I  K  E  Q  E  P  T  Q  Q  F  I  P  V  K  R  P  I

      333      342      351      360      369      378
CCC CAT CCA GCC TAT AAT CCT AAG AAC TTC TCC AAC GAC ATC ATG CTA CTG CAG
P  H  P  A  Y  N  P  K  N  F  S  N  D  I  M  L  L  Q

```

(57) Abstract

The present invention provides a human serine protease precursor (HSPP) and polynucleotides which encode HSPP. The invention also provides expression vectors, host cells, agonists, antisense molecules, antibodies, or antagonists. The invention also provides methods for treating disorders associated with expression of HSPP.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN SERINE PROTEASE PRECURSOR

TECHNCIAL FIELD

This invention relates to nucleic acid and amino acid sequences of a new serine protease precursor protein and to the use of these sequences in the diagnosis, prevention, and treatment of
5 immune disorders and cancer.

BACKGROUND ART

The serine proteases (SP) are a large family of proteolytic enzymes that include the digestive enzymes, trypsin and chymotrypsin, components of the complement cascade and of the blood-clotting cascade, and enzymes that control the degradation and turnover of macromolecules
10 of the extracellular matrix. SP are so named because of the presence of a serine residue in the active catalytic site for protein cleavage. SP have a wide range of substrate specificities and can be subdivided into subfamilies on the basis of these specificities. The main sub-families are trypases (cleavage after arginine or lysine), aspases (cleavage after aspartate), chymases (cleavage after phenylalanine or leucine), metases (cleavage after methionine), and serases (cleavage after
15 serine).

A series of six SP have been identified in murine cytotoxic T-lymphocytes (CTL) and natural killer (NK) cells. These SP are involved with CTL and NK cells in the destruction of virally transformed cells and tumor cells and in organ and tissue transplant rejection (Zunino, S.J. et al. (1990) J. Immunol. 144:2001-9; Sayers, T.J. et al. (1994) J. Immunol. 152:2289-97).
20 Human homologs of most of these enzymes have been identified (Trapani, J.A. et al. (1988) Proc. Nalt. Acad. Sci. 85:6924-28; Caputo, A. et al. (1990) J. Immunol. 145:737-44). Like all SP, the CTL-SP share three distinguishing features: 1) the presence of a catalytic triad of histidine, serine, and aspartate residues which comprise the active site; 2) the sequence GDSGGP which contains the active site serine; and 3) an N-terminal IIGG sequence which characterizes the
25 mature SP.

The SP are secretory proteins which contain N-terminal signal peptides that serve to export the immature protein across the endoplasmic reticulum and are then cleaved (von Heijne (1986) Nuc. Acid. Res. 14:5683-90). Differences in these signal sequences provide one means of distinguishing individual SP. Some SP, particularly the digestive enzymes, exist as inactive
30 precursors or preproenzymes, and contain a leader or activation peptide sequence 3' of the signal peptide. This activation peptide may be 2-12 amino acids in length, and it extends from the cleavage site of the signal peptide to the N-terminal IIGG sequence of the active, mature protein. Cleavage of this sequence activates the enzyme. This sequence varies in different SP according

to the biochemical pathway and/or its substrate (Zunino et al, supra; Sayers et al, supra). Other features that distinguish various SP are the presence or absence of N-linked glycosylation sites that provide membrane anchors, the number and distribution of cysteine residues that determine the secondary structure of the SP, and the sequence of a substrate binding sites such as S'. The S' substrate binding region is defined by residues extending from approximately +17 to +29 relative to the N-terminal I (+1). Differences in this region of the molecule are believed to determine SP substrate specificities (Zunino et al, supra).

The discovery of a new serine protease precursor and the polynucleotides encoding it satisfies a need in the art by providing new diagnostic or therapeutic compositions useful in the treatment or prevention of immune disorders and cancer.

DISCLOSURE OF THE INVENTION

The present invention features a new human serine protease precursor hereinafter designated HSPP and characterized as having similarity to other serine protease precursors.

Accordingly, the invention features a substantially purified HSPP having the amino acid sequence shown in SEQ ID NO:1.

One aspect of the invention features isolated and substantially purified polynucleotides that encode HSPP. In a particular aspect, the polynucleotide is the nucleotide sequence of SEQ ID NO:2.

The invention also relates to a polynucleotide sequence comprising the complement of SEQ ID NO:2 or variants thereof. In addition, the invention features polynucleotide sequences which hybridize under stringent conditions to SEQ ID NO:2.

The invention additionally features fragments of the polynucleotides encoding HSPP, expression vectors and host cells comprising polynucleotides that encode HSPP and a method for producing HSPP using the vectors and host cells. The present invention also features antibodies which bind specifically to HSPP, and pharmaceutical compositions comprising substantially purified HSPP. The invention also features antagonists of HSPP. The invention also provides methods for treating disorders associated with expression of HSPP by administration of antagonists of HSPP and methods for detection of polynucleotides encoding a serine protease precursor protein in a biological sample.

BRIEF DESCRIPTION OF DRAWINGS

Figures 1A, 1B, and 1C, show the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of HSPP. The alignment was produced using MacDNASIS PRO™ software (Hitachi Software Engineering Co., Ltd., San Bruno, CA).

Figures 2A, and 2B, show the amino acid sequence alignments between HSPP (SEQ ID NO:1), the rat natural killer cell protease-1 precursor, RNKP-1 (GI 206690; SEQ ID NO:3), and a human serine esterase from cytotoxic T lymphocytes, SECT (GI 306682; SEQ ID NO:4). The alignment was produced using the multisequence alignment program of DNASTAR™ software
5 (DNASTAR Inc, Madison WI).

Figures 3A, 3B, and 3C show the hydrophobicity plots (MacDNASIS PRO™) for HSPP, SEQ ID NO:1; rat RNKP-1, SEQ ID NO:3; and human SECT (SEQ ID NO:4). The positive X axis reflects amino acid position, and the negative Y axis, hydrophobicity.

MODES FOR CARRYING OUT THE INVENTION

10 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

15 It must be noted that as used herein and in the appended claims, the singular forms “a”, “an”, and “the” include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to “a host cell” includes a plurality of such host cells, reference to the “antibody” is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

20 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are incorporated herein by
25 reference for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

30 “Nucleic acid sequence” as used herein refers to an oligonucleotide, nucleotide, or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand. Similarly, “amino acid sequence” as used herein refers to an oligopeptide, peptide, polypeptide,

or protein sequence, and fragments or portions thereof, and to naturally occurring or synthetic molecules.

Where “amino acid sequence” is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms, such as

5 “polypeptide” or “protein” are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

“Peptide nucleic acid”, as used herein, refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their
10 complementary strand of nucleic acid (Nielsen, P.E. et al. (1993) Anticancer Drug Des. 8:53-63).

HSPP, as used herein, refers to the amino acid sequences of substantially purified HSPP obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic, or recombinant.

15 “Consensus”, as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, or which has been extended using XL-PCR™ (Perkin Elmer, Norwalk, CT) in the 5' and/or the 3' direction and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte clone using the GELVIEW™ Fragment Assembly system (GCG, Madison, WI), or which has been both extended and
20 assembled.

A “variant” of HSPP, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties, e.g., replacement of leucine with isoleucine. More rarely, a variant may have “nonconservative” changes, e.g., replacement of a
25 glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTAR software.

A “deletion”, as used herein, refers to a change in either amino acid or nucleotide
30 sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

An “insertion” or “addition”, as used herein, refers to a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring molecule.

A “substitution”, as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

The term “biologically active”, as used herein, refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise,

5 “immunologically active” refers to the capability of the natural, recombinant, or synthetic HSPP, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term “agonist”, as used herein, refers to a molecule which, when bound to HSPP, causes a change in HSPP which modulates the activity of HSPP. Agonists may include proteins,
10 nucleic acids, carbohydrates, or any other molecules which bind to HSPP.

The terms “antagonist” or “inhibitor”, as used herein, refer to a molecule which, when bound to HSPP, blocks or modulates the biological or immunological activity of HSPP. Antagonists and inhibitors may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to HSPP.

15 The term “modulate”, as used herein, refers to a change or an alteration in the biological activity of HSPP. Modulation may be an increase or a decrease in protein activity, a change in binding characteristics, or any other change in the biological, functional or immunological properties of HSPP.

The term “mimetic”, as used herein, refers to a molecule, the structure of which is
20 developed from knowledge of the structure of HSPP or portions thereof and, as such, is able to effect some or all of the actions of serine protease-like molecules.

The term “derivative”, as used herein, refers to the chemical modification of a nucleic acid encoding HSPP or the encoded HSPP. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would
25 encode a polypeptide which retains essential biological characteristics of the natural molecule.

The term “substantially purified”, as used herein, refers to nucleic or amino acid sequences that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated.

30 “Amplification” as used herein refers to the production of additional copies of a nucleic acid sequence and is generally carried out using polymerase chain reaction (PCR) technologies well known in the art (Dieffenbach, C.W. and G.S. Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY).

The term “hybridization”, as used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term “hybridization complex”, as used herein, refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen binds between complementary
5 G and C bases and between complementary A and T bases; these hydrogen bonds may be further stabilized by base stacking interactions. The two complementary nucleic acid sequences hydrogen bond in an antiparallel configuration. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., membranes, filters, chips,
10 pins or glass slides to which cells have been fixed for in situ hybridization).

The terms “complementary” or “complementarity”, as used herein, refer to the natural binding of polynucleotides under permissive salt and temperature conditions by base-pairing. For example, for the sequence “A-G-T” binds to the complementary sequence “T-C-A”.

Complementarity between two single-stranded molecules may be “partial”, in which only some
15 of the nucleic acids bind, or it may be complete when total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands.

20 The term “homology”, as used herein, refers to a degree of complementarity. There may be partial homology or complete homology (i.e., identity). A partially complementary sequence is one that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid; it is referred to using the functional term “substantially homologous.” The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined
25 using a hybridization assay (Southern or northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (i.e., the hybridization) of a completely homologous sequence or probe to the target sequence under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that
30 the binding of two sequences to one another be a specific (i.e., selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% identity); in the absence of non-specific binding, the probe will not hybridize to the second non-complementary target

sequence.

As known in the art, numerous equivalent conditions may be employed to comprise either low or high stringency conditions. Factors such as the length and nature (DNA, RNA, base composition) of the sequence, nature of the target (DNA, RNA, base composition, presence in
5 solution or immobilization, etc.), and the concentration of the salts and other components (e.g., the presence or absence of formamide, dextran sulfate and/or polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of either low or high stringency different from, but equivalent to, the above listed conditions.

The term "stringent conditions", as used herein, is the "stringency" which occurs within a
10 range from about $T_m - 5^\circ\text{C}$ (5°C below the melting temperature (T_m) of the probe) to about 20°C to 25°C below T_m . As will be understood by those of skill in the art, the stringency of hybridization may be altered in order to identify or detect identical or related polynucleotide sequences.

The term "antisense", as used herein, refers to nucleotide sequences which are
15 complementary to a specific DNA or RNA sequence. The term "antisense strand" is used in reference to a nucleic acid strand that is complementary to the "sense" strand. Antisense molecules may be produced by any method, including synthesis by ligating the gene(s) of interest in a reverse orientation to a viral promoter which permits the synthesis of a complementary strand. Once introduced into a cell, this transcribed strand combines with natural sequences
20 produced by the cell to form duplexes. These duplexes then block either the further transcription or translation. In this manner, mutant phenotypes may be generated. The designation "negative" is sometimes used in reference to the antisense strand, and "positive" is sometimes used in reference to the sense strand.

The term "portion", as used herein, with regard to a protein (as in "a portion of a given
25 protein") refers to fragments of that protein. The fragments may range in size from four amino acid residues to the entire amino acid sequence minus one amino acid. Thus, a protein "comprising at least a portion of the amino acid sequence of SEQ ID NO:1" encompasses the full-length human HSPP and fragments thereof.

"Transformation", as defined herein, describes a process by which exogenous DNA enters
30 and changes a recipient cell. It may occur under natural or artificial conditions using various methods well known in the art. Transformation may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method is selected based on the host cell being transformed and may include, but is not limited to, viral

infection, electroporation, lipofection, and particle bombardment. Such "transformed" cells include stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome. They also include cells which transiently express the inserted DNA or RNA for limited periods of time.

5 The term "antigenic determinant", as used herein, refers to that portion of a molecule that makes contact with a particular antibody (i.e., an epitope). When a protein or fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to a given region or three-dimensional structure on the protein; these regions or structures are referred to as antigenic determinants. An antigenic
10 determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The terms "specific binding" or "specifically binding", as used herein, in reference to the interaction of an antibody and a protein or peptide, mean that the interaction is dependent upon the presence of a particular structure (i.e., the antigenic determinant or epitope) on the protein; in
15 other words, the antibody is recognizing and binding to a specific protein structure rather than to proteins in general. For example, if an antibody is specific for epitope "A", the presence of a protein containing epitope A (or free, unlabeled A) in a reaction containing labeled "A" and the antibody will reduce the amount of labeled A bound to the antibody.

The term "sample", as used herein, is used in its broadest sense. A biological sample
20 suspected of containing nucleic acid encoding HSPP or fragments thereof may comprise a cell, chromosomes isolated from a cell (e.g., a spread of metaphase chromosomes), genomic DNA (in solution or bound to a solid support such as for Southern analysis), RNA (in solution or bound to a solid support such as for northern analysis), cDNA (in solution or bound to a solid support), an extract from cells or a tissue, and the like.

25 The term "correlates with expression of a polynucleotide", as used herein, indicates that the detection of the presence of ribonucleic acid that is similar to SEQ ID NO:2 by northern analysis is indicative of the presence of mRNA encoding HSPP in a sample and thereby correlates with expression of the transcript from the polynucleotide encoding the protein.

"Alterations" in the polynucleotide of SEQ ID NO: 2, as used herein, comprise any
30 alteration in the sequence of polynucleotides encoding HSPP including deletions, insertions, and point mutations that may be detected using hybridization assays. Included within this definition is the detection of alterations to the genomic DNA sequence which encodes HSPP (e.g., by alterations in the pattern of restriction fragment length polymorphisms capable of hybridizing to

SEQ ID NO:2), the inability of a selected fragment of SEQ ID NO: 2 to hybridize to a sample of genomic DNA (e.g., using allele-specific oligonucleotide probes), and improper or unexpected hybridization, such as hybridization to a locus other than the normal chromosomal locus for the polynucleotide sequence encoding HSPP (e.g., using fluorescent *in situ* hybridization [FISH] to
5 metaphase chromosomes spreads).

As used herein, the term “antibody” refers to intact molecules as well as fragments thereof, such as Fa, F(ab')₂, and Fv, which are capable of binding the epitopic determinant. Antibodies that bind HSPP polypeptides can be prepared using intact polypeptides or fragments containing small peptides of interest as the immunizing antigen. The polypeptide or peptide used
10 to immunize an animal can be derived from the transition of RNA or synthesized chemically, and can be conjugated to a carrier protein, if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin and thyroglobulin. The coupled peptide is then used to immunize the animal (e.g., a mouse, a rat, or a rabbit).

The term “humanized antibody”, as used herein, refers to antibody molecules in which
15 amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody, while still retaining the original binding ability.

THE INVENTION

The invention is based on the discovery of a new human serine protease precursor protein (HSPP), the polynucleotides encoding HSPP, and the use of these compositions for the diagnosis,
20 prevention, or treatment of immune disorders and cancer.

Nucleic acids encoding the human HSPP of the present invention were first identified in Incyte Clone 854243 from the ganglioneuroma tissue cDNA library (NGANNOT01) through a computer search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones
25 854243/ NGANNOT01, 1271243/ TESTTUT02, and 1876511/ LEUKNOT03

In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1 as shown in Figures 1A, 1B, and 1C. HSPP is 247 amino acids in length and has potential N-linked glycosylation sites at N71 and N104. Cysteine residues at C49, C65, C142, C173, C188, C209, and C230 represent potential disulfide bridging sites. As shown
30 in Figures 2A and 2B, HSPP has chemical and structural homology with rat RNKP-1 (GI 206690; SEQ ID NO:3) and human SECT (GI 306682; SEQ ID NO:4). In particular, HSPP shares 83% and 93% identity with RNKP-1 and SECT, respectively. The IIGG N-terminal sequence of the mature serine protease is located in HSPP beginning at I121. The three residues comprising the

active catalytic site are located at H64, D108, and S203 in HSPP. The sequence GDSGGP surrounding the active serine residue is conserved in HSPP. The N-linked glycosylation site at N71 in HSPP is shared by RNKP-1 and SECT. The glycosylation site at N104 is shared by SECT. The seven cysteine residues in HSPP are shared by SECT, and all but C230 are also
5 shared by RNKP-1. On the other hand, differences in the N-terminal region of HSPP concerning the potential signal and leader peptides and the S' substrate binding region distinguish HSPP from RNKP-1 and SECT. The N-terminal 20 amino acid segment from M1 to E20 in HSPP precedes the beginning of the active serine protease at IIGG. This segment represents the potential signal and proenzyme regions of HSPP and has only 70% homology with SECT and
10 45% homology with RNKP-1. In particular, differences in the eight amino acid segment immediately prior to the IIGG sequence in HSPP (TPGAGTEE) indicates a significantly different specificity. The amino acid sequence defining the potential S' substrate binding region extending from F37 to R48 in HSPP is also different from that of either RNKP-1 or SECT. As illustrated by Figures 3A, 3B, and 3C, HSPP, RNKP-1, and SECT have rather similar hydrophobicity plots.
15 A notable difference, however, is found in a peak of hydrophilicity centered at approximately residues 40 to 45 in HSPP, and reflects a notable difference in the S' substrate binding sequence of the two proteins. Northern analysis shows the expression of HSPP in various libraries, 47% of which are associated with the immune system and 33% of which are associated with cancer.

The invention also encompasses HSPP variants. A preferred HSPP variant is one having
20 at least 80%, and more preferably 90%, amino acid sequence identity to the HSPP amino acid sequence (SEQ ID NO:1). A most preferred HSPP variant is one having at least 95% amino acid sequence identity to SEQ ID NO:1.

The invention also encompasses polynucleotide which encode HSPP. Accordingly, any nucleic acid sequence which encodes the amino acid sequence of HSPP can be used to generate
25 recombinant molecules which express HSPP. In a particular embodiment, the invention encompasses the polynucleotide comprising the nucleic acid sequence of SEQ ID NO:2 as shown in Figures 1A, 1B and 1c.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences encoding HSPP, some bearing minimal
30 homology to the nucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the

nucleotide sequence of naturally occurring HSPP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode HSPP and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring HSPP under
5 appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding HSPP or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence
10 encoding HSPP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences, or portions thereof, which encode HSPP and its derivatives, entirely by synthetic chemistry. After production, the
15 synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents that are well known in the art at the time of the filing of this application. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding HSPP or any portion thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of
20 hybridizing to the claimed nucleotide sequences, and in particular, those shown in SEQ ID NO:2, under various conditions of stringency. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe, as taught in Wahl, G.M. and S.L. Berger (1987; Methods Enzymol. 152:399-407) and Kimmel, A.R. (1987; Methods Enzymol. 152:507-511), and may be used at a defined stringency.

25 Altered nucleic acid sequences encoding HSPP which are encompassed by the invention include deletions, insertions, or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent HSPP. The encoded protein may also contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent HSPP. Deliberate amino acid substitutions may be made on
30 the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological activity of HSPP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid; positively charged amino acids may include lysine and arginine; and amino acids with uncharged polar head

groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; phenylalanine and tyrosine.

Also included within the scope of the present invention are alleles of the genes encoding HSPP. As used herein, an "allele" or "allelic sequence" is an alternative form of the gene which
5 may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each
10 of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

Methods for DNA sequencing which are well known and generally available in the art may be used to practice any embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham,
15 Chicago, IL), or combinations of recombinant polymerases and proofreading exonucleases such as the ELONGASE Amplification System marketed by Gibco BRL (Gaithersburg, MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI 377 DNA sequencers (Perkin Elmer).

20 The nucleic acid sequences encoding HSPP may be extended utilizing a partial nucleotide sequence and employing various methods known in the art to detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed, "restriction-site" PCR, uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). In particular, genomic DNA is first
25 amplified in the presence of primer to linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR may also be used to amplify or extend sequences using divergent primers
30 based on a known region (Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186). The primers may be designed using OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. The method

uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which may be used is capture PCR which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA
5 (Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119). In this method, multiple restriction enzyme digestions and ligations may also be used to place an engineered double-stranded sequence into an unknown portion of the DNA molecule before performing PCR.

Another method which may be used to retrieve unknown sequences is that of Parker, J.D.
10 et al. (1991; Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PromoterFinder™ libraries to walk in genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that they
15 will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into the 5' and 3' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to
20 analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity may be converted to electrical signal using appropriate software (e.g. Genotyper™ and Sequence
25 Navigator™, Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof
30 which encode HSPP, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of HSPP in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences

may be used to clone and express HSPP.

As will be understood by those of skill in the art, it may be advantageous to produce HSPP-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate
5 of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter HSPP encoding sequences for a variety of reasons,
10 including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

15 In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding HSPP may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of HSPP activity, it may be useful to encode a chimeric HSPP protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the HSPP
20 encoding sequence and the heterologous protein sequence, so that HSPP may be cleaved and purified away from the heterologous moiety.

In another embodiment, sequences encoding HSPP may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232).
25 Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of HSPP, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J.Y. et al. (1995) Science 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer).

30 The newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) Proteins, Structures and Molecular Principles, WH Freeman and Co., New York, NY). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation

procedure; Creighton, *supra*). Additionally, the amino acid sequence of HSPP, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

In order to express a biologically active HSPP, the nucleotide sequences encoding HSPP
5 or functional equivalents, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence.

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding HSPP and appropriate transcriptional and
10 translational control elements. These methods include *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. Such techniques are described in Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, and Ausubel, F.M. et al. (1989) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY.

15 A variety of expression vector/host systems may be utilized to contain and express sequences encoding HSPP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression
20 vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

The "control elements" or "regulatory sequences" are those non-translated regions of the vector--enhancers, promoters, 5' and 3' untranslated regions--which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and
25 specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, LaJolla, CA) or pSport1™ plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in
30 insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO; and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell

line that contains multiple copies of the sequence encoding HSPP, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for HSPP. For example, when large quantities of HSPP are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as Bluescript® (Stratagene), in which the sequence encoding HSPP may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Promega, Madison, WI) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) Methods Enzymol. 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding HSPP may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.

An insect system may also be used to express HSPP. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding HSPP

may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of HSPP will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which HSPP may be expressed (Engelhard, E.K. et al. (1994) Proc. Nat. Acad. Sci. 91:3224-3227).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding HSPP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing HSPP in infected host cells (Logan, J. and Shenk, T. (1984) Proc. Natl. Acad. Sci. 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding HSPP. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding HSPP, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational

activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express HSPP may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression
5 elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using
10 tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1980) *Cell* 22:817-23) genes which can be employed in *tk⁻* or *aprt⁻* cells, respectively. Also, antimetabolite,
15 antibiotic or herbicide resistance can be used as the basis for selection; for example, *dhfr* which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); *npt*, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14) and *als* or *pat*, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have
20 been described, for example, *trpB*, which allows cells to utilize indole in place of tryptophan, or *hisD*, which allows cells to utilize histinol in place of histidine (Hartman, S.C. and R.C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but
25 also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C.A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding HSPP is inserted within a marker gene sequence, recombinant cells
30 containing sequences encoding HSPP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding HSPP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells which contain the nucleic acid sequence encoding HSPP and express HSPP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include membrane, solution, or chip based
5 technologies for the detection and/or quantification of nucleic acid or protein.

The presence of polynucleotide sequences encoding HSPP can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or portions or fragments of polynucleotides encoding HSPP. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the sequences encoding HSPP to detect transformants
10 containing DNA or RNA encoding HSPP. As used herein "oligonucleotides" or "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides, which can be used as a probe or amplimer.

A variety of protocols for detecting and measuring the expression of HSPP, using either
15 polyclonal or monoclonal antibodies specific for the protein are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on HSPP is preferred, but a competitive binding assay may be employed. These and other assays are described, among other
20 places, in Hampton, R. et al. (1990; Serological Methods, a Laboratory Manual, APS Press, St Paul, MN) and Maddox, D.E. et al. (1983; J. Exp. Med. 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding HSPP
25 include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding HSPP, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures
30 may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical Corp., Cleveland, OH). Suitable reporter molecules or labels, which may be used, include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors,

magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding HSPP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode HSPP may be designed to contain signal sequences which direct secretion of HSPP through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding HSPP to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and HSPP may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing HSPP and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography as described in Porath, J. et al. (1992, Prot. Exp. Purif. 3:263-281) while the enterokinase cleavage site provides a means for purifying HSPP from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. et al. (1993; DNA Cell Biol. 12:441-453).

In addition to recombinant production, fragments of HSPP may be produced by direct peptide synthesis using solid-phase techniques Merrifield J. (1963) J. Am. Chem. Soc. 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Various fragments of HSPP may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

THERAPEUTICS

Chemical and structural homology exists among HSPP, rat RNKP-1 and human SECT. In addition, northern analysis shows the expression of HSPP in tissues associated with the immune system and cancer. HSPP therefore appears to be associated with the development of immune disorders and cancer.

An increase in the level or activity of HSPP appears to be associated with the development of immune disorders. Therefore in one embodiment, an antagonist or inhibitor of HSPP may be administered to a subject to treat or prevent an immune disorder. Immune disorders may include, but are not limited to, Addison's disease, adult respiratory distress syndrome, allergies, anemia, asthma, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, ulcerative colitis, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, atrophic gastritis, glomerulonephritis, gout, Graves' disease, hypereosinophilia, irritable bowel syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, rheumatoid arthritis, scleroderma, Sjögren's syndrome, and autoimmune thyroiditis; complications of cancer, hemodialysis, extracorporeal circulation; viral, bacterial, fungal, parasitic, protozoal, and helminthic infections; and trauma.

An increase in the level or activity of HSPP also appears to be associated with the development of cancer. Therefore in another embodiment, an antagonist or inhibitor of HSPP may be administered to a subject to treat or prevent cancer, including astrocytoma, glioma, ganglioneuroma, neurocytoma, neuroblastoma, adenocarcinoma, sarcoma, melanoma, lymphoma, leukemia, and myeloma. In particular, types of cancer may include, but are not limited to, cancer of the adrenal gland, bladder, bone, brain, breast, gastrointestinal tract, heart, kidney, liver, lung, ovary, pancreas, paraganglia, parathyroid, prostate, salivary glands, skin, spleen, testis, thyroid, and uterus.

In any of the above embodiments, antibodies which are specific for HSPP may be used directly as an antagonist, or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express HSPP.

In another embodiment, the complement of the polynucleotide encoding HSPP or an antisense molecule may be administered to a subject to treat or prevent any of the immune disorders listed above.

In another embodiment, the complement of the polynucleotide encoding HSPP or an antisense molecule may be administered to a subject to treat or prevent any of the types of cancer listed above.

In other embodiments, the antagonists, antibodies, antisense or complementary sequences or vectors described above may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles.

The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

5 Antagonists or inhibitors of HSPP may be produced using methods which are generally known in the art. In particular, purified HSPP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which bind HSPP.

The antibodies may be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain,
10 Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies, (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others, may be immunized by injection with HSPP or any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants
15 may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

20 It is preferred that the peptides, fragments, or oligopeptides used to induce antibodies to HSPP have an amino acid sequence consisting of at least five amino acids, and more preferably at least 10 amino acids. It is also preferable that they are identical to a portion of the amino acid sequence of the natural protein, and they may contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of HSPP amino acids may be fused with those of
25 another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule.

Monoclonal antibodies to HSPP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the
30 EBV-hybridoma technique (Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. 80:2026-2030; Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120).

In addition, techniques developed for the production of "chimeric antibodies", the splicing

of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; Takeda, S. et al. (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies
5 may be adapted, using methods known in the art, to produce HSPP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton D.R. (1991) Proc. Natl. Acad. Sci. 88:11120-3).

Antibodies may also be produced by inducing in vivo production in the lymphocyte
10 population or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. 86: 3833-3837; Winter, G. et al. (1991) Nature 349:293-299).

Antibody fragments which contain specific binding sites for HSPP may also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be
15 produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse, W.D. et al. (1989) Science 254:1275-1281).

Various immunoassays may be used for screening to identify antibodies having the
20 desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between HSPP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering HSPP epitopes is preferred, but a
25 competitive binding assay may also be employed (Maddox, supra).

In another embodiment of the invention, the polynucleotides encoding HSPP, or any fragment thereof, or antisense molecules, may be used for therapeutic purposes. In one aspect, antisense to the polynucleotide encoding HSPP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with
30 sequences complementary to polynucleotides encoding HSPP. Thus, antisense molecules may be used to modulate HSPP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligomers or larger fragments, can be designed from various locations along the coding or control regions of sequences encoding HSPP.

Expression vectors derived from retro viruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can be used to construct recombinant vectors which will express antisense molecules complementary to the polynucleotides of the gene encoding HSPP. These techniques are described both in Sambrook et al. (supra) and in Ausubel et al. (supra).

Genes encoding HSPP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide or fragment thereof which encodes HSPP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector and even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA, or PNA, to the control regions of the gene encoding HSPP, i.e., the promoters, enhancers, and introns. Oligonucleotides derived from the transcription initiation site, e.g., between positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee, J.E. et al. (1994) In: Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY). The antisense molecules may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples which may be used include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding HSPP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be

evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Antisense molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding HSPP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection and by liposome injections may be achieved using methods which are well known in the art.

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of HSPP, antibodies to HSPP, mimetics, agonists, antagonists, or inhibitors of HSPP. The compositions may be administered alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical

carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any
5 number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which
10 facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using
15 pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active
20 compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums
25 including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel,
30 polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of

gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty
5 oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl
10 cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly
15 concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating,
20 dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized
25 powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of HSPP, such labeling would include amount, frequency, and method of administration.

30 Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes
5 for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example HSPP or fragments thereof, antibodies of HSPP, agonists, antagonists or inhibitors of HSPP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g.,
10 ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD₅₀/ED₅₀. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in
15 such compositions is preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient
20 levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life
25 and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or
30 their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind HSPP may be used for the

diagnosis of conditions or diseases characterized by expression of HSPP, or in assays to monitor patients being treated with HSPP, agonists, antagonists or inhibitors. The antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for HSPP include methods which utilize the antibody and a label
5 to detect HSPP in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules which are known in the art may be used, several of which are described above.

A variety of protocols including ELISA, RIA, and FACS for measuring HSPP are known
10 in the art and provide a basis for diagnosing altered or abnormal levels of HSPP expression. Normal or standard values for HSPP expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to HSPP under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, but preferably by photometric, means. Quantities of HSPP
15 expressed in subject, control and disease, samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding HSPP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide
20 sequences, antisense RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of HSPP may be correlated with disease. The diagnostic assay may be used to distinguish between absence, presence, and excess expression of HSPP, and to monitor regulation of HSPP levels during therapeutic intervention.

25 In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding HSPP or closely related molecules, may be used to identify nucleic acid sequences which encode HSPP. The specificity of the probe, whether it is made from a highly specific region, e.g., 10 unique nucleotides in the 5' regulatory region, or a less specific region, e.g., especially in the 3' coding region, and the
30 stringency of the hybridization or amplification (maximal, high, intermediate, or low) will determine whether the probe identifies only naturally occurring sequences encoding HSPP, alleles, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably

contain at least 50% of the nucleotides from any of the HSPP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements, and introns of the naturally occurring HSPP.

5 Means for producing specific hybridization probes for DNAs encoding HSPP include the cloning of nucleic acid sequences encoding HSPP or HSPP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a
10 variety of reporter groups, for example, radionuclides such as ³²P or ³⁵S, or enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding HSPP may be used for the diagnosis of conditions or diseases which are associated with expression of HSPP. Examples of such conditions or diseases
15 include immune disorders such as Addison's disease, adult respiratory distress syndrome, allergies, anemia, asthma, atherosclerosis, bronchitis, cholecystitis, Crohn's disease, ulcerative colitis, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, atrophic gastritis, glomerulonephritis, gout, Graves' disease, hypereosinophilia, irritable bowel syndrome, lupus erythematosus, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation,
20 osteoarthritis, osteoporosis, pancreatitis, polymyositis, rheumatoid arthritis, scleroderma, Sjögren's syndrome, and autoimmune thyroiditis; complications of cancer, hemodialysis, extracorporeal circulation; viral, bacterial, fungal, parasitic, protozoal, and helminthic infections; and trauma; and cancer, such as cancer of the blood, thyroid, spleen, brain, lung, prostate, testes, ovaries, and pancreas. The polynucleotide sequences encoding HSPP may be used in Southern or
25 northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; or in dip stick, pin, ELISA or chip assays utilizing fluids or tissues from patient biopsies to detect altered HSPP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding HSPP may be useful in assays that detect activation or induction of various cancers, particularly those mentioned above. The
30 nucleotide sequences encoding HSPP may be labeled by standard methods, and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the biopsied or extracted sample

is significantly altered from that of a comparable control sample, the nucleotide sequences have hybridized with nucleotide sequences in the sample, and the presence of altered levels of nucleotide sequences encoding HSPP in the sample indicates the presence of the associated disease. Such assays may also be used to evaluate the efficacy of a particular therapeutic
5 treatment regimen in animal studies, in clinical trials, or in monitoring the treatment of an individual patient.

In order to provide a basis for the diagnosis of disease associated with expression of HSPP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a
10 sequence, or a fragment thereof, which encodes HSPP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with those from an experiment where a known amount of a substantially purified polynucleotide is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients who are symptomatic for disease.
15 Deviation between standard and subject values is used to establish the presence of disease.

Once disease is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to evaluate whether the level of expression in the patient begins to approximate that which is observed in the normal patient. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to
20 months.

With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ
25 preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding HSPP may involve the use of PCR. Such oligomers may be chemically synthesized, generated enzymatically, or produced from a recombinant source. Oligomers will preferably consist of two
30 nucleotide sequences, one with sense orientation (5'→3') and another with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for detection and/or quantitation of closely related DNA

or RNA sequences.

Methods which may also be used to quantitate the expression of HSPP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated (Melby, P.C. et al. (1993) J.

- 5 Immunol. Methods, 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 229-236). The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

- In another embodiment of the invention, the nucleic acid sequences which encode HSPP
10 may also be used to generate hybridization probes which are useful for mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. Such techniques include FISH, FACS, or artificial chromosome constructions, such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial P1 constructions or single chromosome cDNA libraries as
15 reviewed in Price, C.M. (1993) Blood Rev. 7:127-134, and Trask, B.J. (1991) Trends Genet. 7:149-154.

- FISH (as described in Verma et al. (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York, NY) may be correlated with other physical chromosome mapping techniques and genetic map data. Examples of genetic map data can be found in the
20 1994 Genome Issue of Science (265:1981f). Correlation between the location of the gene encoding HSPP on a physical chromosomal map and a specific disease, or predisposition to a specific disease, may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier, or affected individuals.

- 25 In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts
30 thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti, R.A. et al. (1988) Nature 336:577-580), any sequences mapping

to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

In another embodiment of the invention, HSPP, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between HSPP and the agent being tested, may be measured.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the protein of interest as described in published PCT application WO84/03564. In this method, as applied to HSPP large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with HSPP, or fragments thereof, and washed. Bound HSPP is then detected by methods well known in the art. Purified HSPP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding HSPP specifically compete with a test compound for binding HSPP. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with HSPP.

In additional embodiments, the nucleotide sequences which encode HSPP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

INDUSTRIAL APPLICABILITY

I NGANNOT01 cDNA Library Construction

The NGANNOT01 cDNA library was constructed from ganglioneuroma which was excised from a 9-year-old Causcasian male. This ganglioneuroma formed an encapsulated lobulated mass. The pathology report also noted that excision included the left chest wall and tissue from pleura surrounding the tumor and containing fibrotic tissue with chronic

inflammation which extended into the patient's overlying adipose tissue.

The frozen tissue was homogenized and lysed using a Brinkmann Homogenizer Polytron PT-3000 (Brinkmann Instruments, Westbury NJ). The lysate was centrifuged over a 5.7 M CsCl cushion using a Beckman SW28 rotor in a Beckman L8-70M Ultracentrifuge (Beckman Instruments) for 18 hours at 25,000 rpm at ambient temperature. The RNA was extracted with phenol chloroform pH 4.0, precipitated using 0.3 M sodium acetate and 2.5 volumes of ethanol, resuspended in DEPC-treated water and DNase treated at 37°C. RNA extraction and precipitation were repeated as before. The RNA was isolated with the Qiagen Oligotex kit (QIAGEN Inc,) and used to construct the cDNA library.

The RNA was handled according to the recommended protocols in the SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Cat. #18248-013; Gibco/BRL, Gaithersburg, MD). cDNAs were fractionated on a Sepharose CL4B column (Cat. #275105, Pharmacia), and those cDNAs exceeding 400 bp were ligated into pSport I. The plasmid pSport I was subsequently transformed into DH5a™ competent cells (Cat. #18258-012, Gibco/BRL).

II Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 Plasmid Kit (Catalog #26173; QIAGEN, Inc.). This kit enables the simultaneous purification of 96 samples in a 96-well block using multi-channel reagent dispensers. The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth (Catalog #22711, Gibco/BRL) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) the cultures were incubated for 19 hours after the wells were inoculated and then lysed with 0.3 ml of lysis buffer; 3) following isopropanol precipitation, the plasmid DNA pellet was resuspended in 0.1 ml of distilled water. After the last step in the protocol, samples were transferred to a Beckman 96-well block for storage.

The cDNAs were sequenced by the method of Sanger F and AR Coulson (1975; J Mol Biol 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno NV) in combination with Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown MA) and Applied Biosystems 377 DNA Sequencing Systems; and the reading frame was determined.

III Homology Searching of cDNA Clones and Their Deduced Proteins

The nucleotide sequences of the Sequence Listing or amino acid sequences deduced from them were used as query sequences against databases such as GenBank, SwissProt, BLOCKS, and Pima II. These databases which contain previously identified and annotated sequences were searched for regions of homology (similarity) using BLAST, which stands for Basic Local

Alignment Search Tool (Altschul, S.F. (1993) J. Mol. Evol. 36:290-300; Altschul et al. (1990) J. Mol. Biol. 215:403-410).

BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in
 5 determining exact matches or in identifying homologs which may be of prokaryotic (bacterial) or eukaryotic (animal, fungal or plant) origin. Other algorithms such as the one described in Smith RF and TF Smith (1992; Protein Engineering 5:35-51), incorporated herein by reference, can be used when dealing with primary sequence patterns and secondary structure gap penalties. As disclosed in this application, the sequences have lengths of at least 49 nucleotides, and no more
 10 than 12% uncalled bases (where N is recorded rather than A, C, G, or T).

The BLAST approach, as detailed in Karlin, S. and S.F. Atschul (1993; Proc. Nat. Acad. Sci. 90:5873-7) and incorporated herein by reference, searches for matches between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. In this
 15 application, threshold was set at 10^{-25} for nucleotides and 10^{-14} for peptides.

Incyte nucleotide sequences were searched against the GenBank databases for primate (pri), rodent (rod), and mammalian sequences (mam), and deduced amino acid sequences from the same clones are searched against GenBank functional protein databases, mammalian (mamp), vertebrate (vrtp) and eukaryote (eukp), for homology. The relevant database for a particular
 20 match were reported as a GIXxx±p (where xxx is pri, rod, etc and if present, p = peptide).

IV Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al., supra).

25 Analogous computer techniques using BLAST (Altschul, S.F. 1993 and 1990, supra) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte Pharmaceuticals). This analysis is much faster than multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

30 The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the

length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

- 5 The results of northern analysis are reported as a list of libraries in which the transcript encoding HSPP occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

10 **V Extension of HSPP-Encoding Polynucleotides**

- Nucleic acid sequence of Incyte Clone 854243 or SEQ ID NO:2 is used to design oligonucleotide primers for extending a partial nucleotide sequence to full length or for obtaining 5' or 3', intron or other control sequences from genomic libraries. One primer is synthesized to initiate extension in the antisense direction (XLR) and the other is synthesized to extend sequence
15 in the sense direction (XLF). Primers are used to facilitate the extension of the known sequence "outward" generating amplicons containing new, unknown nucleotide sequence for the region of interest. The initial primers are designed from the cDNA using OLIGO 4.06 (National Biosciences), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C.
20 Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations is avoided.

The original, selected cDNA libraries, or a human genomic library are used to extend the sequence; the latter is most useful to obtain 5' upstream regions. If more extension is necessary or desired, additional sets of primers are designed to further extend the known region.

- 25 By following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix, high fidelity amplification is obtained. Beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR is performed using the Peltier Thermal Cycler (PTC200; M.J. Research, Watertown, MA) and the following parameters:

- | | | |
|----|--------|--|
| 30 | Step 1 | 94° C for 1 min (initial denaturation) |
| | Step 2 | 65° C for 1 min |
| | Step 3 | 68° C for 6 min |
| | Step 4 | 94° C for 15 sec |
| | Step 5 | 65° C for 1 min |

- | | | |
|---|---------|--|
| | Step 6 | 68° C for 7 min |
| | Step 7 | Repeat step 4-6 for 15 additional cycles |
| | Step 8 | 94° C for 15 sec |
| | Step 9 | 65° C for 1 min |
| 5 | Step 10 | 68° C for 7:15 min |
| | Step 11 | Repeat step 8-10 for 12 cycles |
| | Step 12 | 72° C for 8 min |
| | Step 13 | 4° C (and holding) |

10 A 5-10 μ l aliquot of the reaction mixture is analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products are selected and removed from the gel. Further purification involves using a commercial gel extraction method such as QIAQuick™ (QIAGEN Inc., Chatsworth, CA). After recovery of the DNA, Klenow enzyme is
15 used to trim single-stranded, nucleotide overhangs creating blunt ends which facilitate religation and cloning.

After ethanol precipitation, the products are redissolved in 13 μ l of ligation buffer, 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase are added, and the mixture is incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in
20 40 μ l of appropriate media) are transformed with 3 μ l of ligation mixture and cultured in 80 μ l of SOC medium (Sambrook et al., supra). After incubation for one hour at 37° C, the whole transformation mixture is plated on Luria Bertani (LB)-agar (Sambrook et al., supra) containing 2x Carb. The following day, several colonies are randomly picked from each plate and cultured in 150 μ l of liquid LB/2x Carb medium placed in an individual well of an appropriate,
25 commercially-available, sterile 96-well microtiter plate. The following day, 5 μ l of each overnight culture is transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5 μ l of each sample is transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for
30 the extension reaction are added to each well. Amplification is performed using the following conditions:

- | | | |
|----|--------|--|
| | Step 1 | 94° C for 60 sec |
| | Step 2 | 94° C for 20 sec |
| | Step 3 | 55° C for 30 sec |
| 35 | Step 4 | 72° C for 90 sec |
| | Step 5 | Repeat steps 2-4 for an additional 29 cycles |
| | Step 6 | 72° C for 180 sec |
| | Step 7 | 4° C (and holding) |

Aliquots of the PCR reactions are run on agarose gels together with molecular weight markers. The sizes of the PCR products are compared to the original partial cDNAs, and appropriate clones are selected, ligated into plasmid, and sequenced.

VI Labeling and Use of Hybridization Probes

5 Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250 μ Ci of [γ - ^{32} P] adenosine triphosphate (Amersham) and T4 polynucleotide kinase (DuPont NEN[®], Boston, MA).
10 The labeled oligonucleotides are substantially purified with Sephadex G-25 superfine resin column (Pharmacia & Upjohn). A portion containing 10^7 counts per minute of each of the sense and antisense oligonucleotides is used in a typical membrane based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN[®]).
15

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR[™] film (Kodak, Rochester, NY) is exposed to the
20 blots in a Phosphorimager cassette (Molecular Dynamics, Sunnyvale, CA) for several hours, hybridization patterns are compared visually.

VII Complementary Polynucleotide, Antisense Molecules

Polynucleotide complementary to the HSPP-encoding sequence, or any part thereof, or
25 an antisense molecule is used to inhibit in vivo expression of naturally occurring HSPP. Although use of antisense oligonucleotides, comprising about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. An oligonucleotide based on the coding sequences of HSPP, as shown in Figures 1A, 1B and 1C, is used to inhibit expression of naturally occurring HSPP. The complementary oligonucleotide is
30 designed from the most unique 5' sequence as shown in Figures 1A, 1B and 1C and used either to inhibit transcription by preventing promoter binding to the upstream nontranslated sequence or translation of an HSPP-encoding transcript by preventing the ribosome from binding. Using an appropriate portion of the signal and 5' sequence of SEQ ID NO:2, an effective antisense

oligonucleotide includes any 15-20 nucleotides spanning the region which translates into the signal or 5' coding sequence of the polypeptide as shown in Figures 1A, 1B and 1C.

VIII Expression of HSPP

Expression of HSPP is accomplished by subcloning the cDNAs into appropriate vectors and transforming the vectors into host cells. In this case, the cloning vector is used to express HSPP in *E. coli*. Upstream of the cloning site, this vector contains a promoter for β -galactosidase, followed by sequence containing the amino-terminal Met, and the subsequent seven residues of β -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

10 Induction of an isolated, transformed bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first eight residues of β -galactosidase, about 5 to 15 residues of linker, and the full length protein. The signal residues direct the secretion of HSPP into the bacterial growth media which can be used directly in the following assay for activity.

IX Demonstration of HSPP Activity

15 Serine protease activity of HSPP is measured by the hydrolysis of various peptide thiobenzyl ester substrates. The substrates are chosen to represent the different SP types (chymase, trypase, aspase, etc.). Assays are performed at room temperature (~25°C) and contain an aliquot of HSPP and the appropriate substrate in HEPES buffer, pH 7.5 containing 0.01M CaCl_2 and 8% dimethylsulfoxide. The reaction also contains 0.34 mM dithiopyridine, which
20 reacts with the thiobenzyl group that is released during hydrolysis and converts it to thiopyridone. The reaction is carried out in an optical cuvette and the generation of thiopyridone is measured in a spectrophotometer by the absorption produced at 324 nm. The amount of thiopyridone produced in the reaction is proportional to the activity of HSPP.

X Production of HSPP Specific Antibodies

25 HSPP that is substantially purified using PAGE electrophoresis (Sambrook, supra), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence deduced from SEQ ID NO:2 is analyzed using DNASTAR software (DNASTAR Inc) to determine regions of high immunogenicity and a corresponding oligopolypeptide is synthesized and used to raise antibodies by means known to
30 those of skill in the art. Selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions, is described by Ausubel et al. (supra), and others.

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole

limpet hemocyanin (KLH, Sigma, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel et al., supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radioiodinated, goat anti-rabbit IgG.

XI Purification of Naturally Occurring HSPP Using Specific Antibodies

Naturally occurring or recombinant HSPP is substantially purified by immunoaffinity chromatography using antibodies specific for HSPP. An immunoaffinity column is constructed by covalently coupling HSPP antibody to an activated chromatographic resin, such as CnBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing HSPP is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of HSPP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/HSPP binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope, such as urea or thiocyanate ion), and HSPP is collected.

XII Identification of Molecules Which Interact with HPRS

HSPP or biologically active fragments thereof are labeled with ¹²⁵I Bolton-Hunter reagent (Bolton et al. (1973) Biochem. J. 133:529). Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled HSPP, washed and any wells with labeled HSPP complex are assayed. Data obtained using different concentrations of HSPP are used to calculate values for the number, affinity, and association of HSPP with the candidate molecules.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: INCYTE PHARMACEUTICALS, INC.
- (ii) TITLE OF THE INVENTION: HUMAN SERINE PROTEASE PRECURSOR
- (iii) NUMBER OF SEQUENCES: 4
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Incyte Pharmaceuticals, Inc.
 - (B) STREET: 3174 Porter Drive
 - (C) CITY: Palo Alto
 - (D) STATE: CA
 - (E) COUNTRY: USA
 - (F) ZIP: 94304
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ for Windows Version 2.0
- (vi) CURRENT APPLICATION DATA:
 - (A) PCT APPLICATION NUMBER: To Be Assigned
 - (B) FILING DATE: Herewith
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: US 08/851,974
 - (B) FILING DATE: 02-MAY-1997
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Billings, Lucy J.
 - (B) REGISTRATION NUMBER: 36,749
 - (C) REFERENCE/DOCKET NUMBER: PF-0288 PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 650-855-0555
 - (B) TELEFAX: 650-845-4166

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 247 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (vii) IMMEDIATE SOURCE:
 - (A) LIBRARY: NGANNOT01
 - (B) CLONE: 854243

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met	Gln	Pro	Phe	Leu	Leu	Leu	Leu	Ala	Phe	Leu	Leu	Thr	Pro	Gly	Ala
1				5				10						15	
Gly	Thr	Glu	Glu	Ile	Ile	Gly	Gly	His	Glu	Ala	Lys	Pro	His	Ser	Arg
			20					25						30	
Pro	Tyr	Met	Ala	Phe	Val	Gln	Phe	Leu	Gln	Glu	Lys	Ser	Arg	Lys	Arg
			35					40						45	
Cys	Gly	Gly	Ile	Leu	Val	Arg	Lys	Asp	Phe	Val	Leu	Thr	Ala	Ala	His
			50					55						60	
Cys	Gln	Gly	Ser	Ser	Ile	Asn	Val	Thr	Leu	Gly	Ala	His	Asn	Ile	Lys
65					70					75					80

```

Glu Gln Glu Pro Thr Gln Gln Phe Ile Pro Val Lys Arg Pro Ile Pro
      85                      90                      95
His Pro Ala Tyr Asn Pro Lys Asn Phe Ser Asn Asp Ile Met Leu Leu
      100                    105                    110
Gln Leu Glu Arg Lys Ala Lys Arg Thr Arg Ala Val Gln Pro Leu Arg
      115                    120                    125
Leu Pro Ser Asn Lys Ala Gln Val Lys Pro Gly Gln Thr Cys Ser Val
      130                    135                    140
Ala Gly Trp Gly Gln Thr Ala Pro Leu Gly Lys His Ser His Thr Leu
      145                    150                    155                    160
Gln Glu Val Lys Met Thr Val Gln Glu Asp Arg Lys Cys Glu Ser Asp
      165                    170                    175
Leu Arg His Tyr Tyr Asp Ser Thr Ile Glu Leu Cys Val Gly Asp Pro
      180                    185                    190
Glu Ile Lys Lys Thr Ser Phe Lys Gly Asp Ser Gly Gly Pro Leu Val
      195                    200                    205
Cys Asn Lys Val Ala Gln Gly Ile Val Ser Tyr Gly Arg Asn Asn Gly
      210                    215                    220
Met Pro Pro Arg Ala Cys Thr Lys Val Ser Ser Phe Val His Trp Ile
      225                    230                    235                    240
Lys Lys Thr Met Lys Arg Tyr
      245

```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 884 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: NGANNOT01
- (B) CLONE: 854243

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

ACACCAACAG CTCTGACCTG GGCAGCCTTC CTGAGAAAAT GCAGCCATTC CTCCTCCTGT      60
TGGCCTTTCT TCTGACCCCT GGGGCTGGGA CAGAGGAGAT CATCGGGGGC CATGAGGCCA      120
AGCCCCACTC CCGCCCCTAC ATGGCCTTTG TTCAGTTTCT GCAAGAGAAG AGTCGGAAGA      180
GGTGTGGCGG CATCCTAGTG AGAAAGGACT TTGTGCTGAC AGCTGCTCAC TGCCAGGGAA      240
GCTCCATAAA TGTCACCTTG GGGGCCCCACA ATATCAAAGA ACAGGAGCCG ACCCAGCAGT      300
TTATCCCTGT GAAAAGACCC ATCCCCCATC CAGCCTATAA TCCTAAGAAC TTCTCCAACG      360
ACATCATGCT ACTGCAGCTG GAGAGAAAAG CCAAGCGGAC CAGAGCTGTG CAGCCCCCTCA      420
GGCTACCTAG CAACAAGGCC CAGGTGAAGC CAGGGCAGAC ATGCAGTGTG GCCGGCTGGG      480
GGCAGACGGC CCCCTGGGA AAACACTCAC ACACACTACA AGAGGTGAAG ATGACAGTGC      540
AGGAAGATCG AAAGTGCGAA TCTGACTTAC GCCATTATTA CGACAGTACC ATTGAGTTGT      600
GCGTGGGGGA CCCAGAGATT AAAAAGACTT CTTTAAGGG GGA CTCTGGA GGCCCTCTTG      660
TGTGTAACAA GGTGGCCAG GGCATTGTCT CCTATGGACG AAACAATGGC ATGCCTCCAC      720
GAGCCTGCAC CAAAGTCTCA AGCTTTGTAC ACTGGATAAA GAAAACCATG AAACGCTACT      780
AACTACAGGA AGCAAATAA GCCCCGCTG TAATGAAACA CCTTCTCTGG AGCCAAGTCC      840
AGATTACAC TGGGAGAGGT GCCAGCAACT GAATAAATCC TCTT      884

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 248 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
- (B) CLONE: 206690

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Met Lys Leu Leu Leu Leu Leu Leu Ser Phe Ser Leu Ala Pro Lys Thr
 1          5          10          15
Glu Ala Gly Glu Ile Ile Gly Gly His Glu Ala Lys Pro His Ser Arg
          20          25          30
Pro Tyr Met Ala Tyr Leu Gln Ile Met Asp Glu Tyr Ser Gly Ser Lys
          35          40          45
Lys Cys Gly Gly Phe Leu Ile Arg Glu Asp Phe Val Leu Thr Ala Ala
          50          55          60
His Cys Ser Gly Ser Lys Ile Asn Val Thr Leu Gly Ala His Asn Ile
65          70          75          80
Lys Glu Gln Glu Lys Met Gln Gln Ile Ile Pro Val Val Lys Ile Ile
          85          90          95
Pro His Pro Ala Tyr Asn Ser Lys Thr Ile Ser Asn Asp Ile Met Leu
          100          105          110
Leu Lys Leu Lys Ser Lys Ala Lys Arg Ser Ser Ala Val Lys Pro Leu
          115          120          125
Asn Leu Pro Arg Arg Asn Val Lys Val Lys Pro Gly Asp Val Cys Tyr
130          135          140
Val Ala Gly Trp Gly Lys Leu Gly Pro Met Gly Lys Tyr Ser Asp Thr
145          150          155          160
Leu Gln Glu Val Glu Leu Thr Val Gln Glu Asp Gln Lys Cys Glu Ser
          165          170          175
Tyr Leu Lys Asn Tyr Phe Asp Lys Ala Asn Glu Ile Cys Ala Gly Asp
          180          185          190
Pro Lys Ile Lys Arg Ala Ser Phe Arg Gly Asp Ser Gly Gly Pro Leu
          195          200          205
Val Cys Lys Lys Val Ala Ala Gly Ile Val Ser Tyr Gly Gln Asn Asp
210          215          220
Gly Ser Thr Pro Arg Ala Phe Thr Lys Val Ser Thr Phe Leu Ser Trp
225          230          235          240
Ile Lys Lys Thr Met Lys Lys Ser
          245

```

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 247 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
- (B) CLONE: 306682

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Gln Pro Ile Leu Leu Leu Leu Ala Phe Leu Leu Leu Pro Arg Ala
 1          5          10          15
Asp Ala Gly Glu Ile Ile Gly Gly His Glu Ala Lys Pro His Ser Arg
          20          25          30
Pro Tyr Met Ala Tyr Leu Met Ile Trp Asp Gln Lys Ser Leu Lys Arg
          35          40          45
Cys Gly Gly Phe Leu Ile Gln Asp Asp Phe Val Leu Thr Ala Ala His
          50          55          60
Cys Trp Gly Ser Ser Ile Asn Val Thr Leu Gly Ala His Asn Ile Lys
65          70          75          80
Glu Gln Glu Pro Thr Gln Gln Phe Ile Pro Val Lys Arg Pro Ile Pro
          85          90          95
His Pro Ala Tyr Asn Pro Lys Asn Phe Ser Asn Asp Ile Met Leu Leu
          100          105          110
Gln Leu Glu Arg Lys Ala Lys Arg Thr Arg Ala Val Gln Pro Leu Arg
          115          120          125
Leu Pro Ser Asn Lys Ala Gln Val Lys Pro Gly Gln Thr Cys Ser Val
130          135          140
Ala Gly Trp Gly Gln Thr Ala Pro Leu Gly Lys His Ser His Thr Leu
145          150          155          160

```

Gln	Glu	Val	Lys	Met	Thr	Val	Gln	Glu	Asp	Arg	Lys	Cys	Glu	Ser	Asp
				165					170					175	
Leu	Arg	His	Tyr	Tyr	Asp	Ser	Thr	Ile	Glu	Leu	Cys	Val	Gly	Asp	Pro
			180					185					190		
Glu	Ile	Lys	Lys	Thr	Ser	Phe	Lys	Gly	Asp	Ser	Gly	Gly	Pro	Leu	Val
		195					200					205			
Cys	Asn	Lys	Val	Ala	Gln	Gly	Ile	Val	Ser	Tyr	Gly	Arg	Asn	Asn	Gly
	210					215					220				
Met	Pro	Pro	Arg	Ala	Cys	Thr	Lys	Val	Ser	Ser	Phe	Val	His	Trp	Ile
225					230					235					240
Lys	Lys	Thr	Met	Lys	Arg	Tyr									
				245											

What is claimed is:

1. A substantially purified human serine protease precursor protein comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof.
2. An isolated and purified polynucleotide sequence encoding the serine protease precursor protein of claim 1.
3. A polynucleotide sequence which hybridizes under stringent conditions to the polynucleotide sequence of claim 2.
4. A hybridization probe comprising the polynucleotide sequence of claim 2.
5. An isolated and purified polynucleotide sequence comprising SEQ ID NO:2 or variants thereof.
6. A polynucleotide sequence which is complementary to the polynucleotide sequence of claim 2 or variants thereof.
7. A hybridization probe comprising the polynucleotide sequence of claim 6.
8. An expression vector containing the polynucleotide sequence of claim 2.
9. A host cell containing the vector of claim 8.
10. A method for producing a polypeptide comprising the amino acid sequence of SEQ ID NO:1 the method comprising the steps of:
 - a) culturing the host cell of claim 9 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
11. A pharmaceutical composition comprising a substantially purified serine protease precursor protein having an amino acid sequence of claim 1 in conjunction with a suitable pharmaceutical carrier.
12. A purified antibody which binds specifically to the polypeptide of claim 1.
13. A purified agonist which modulates the activity of the polypeptide of claim 1.
14. A purified antagonist which decreases the activity of the polypeptide of claim 1.
15. A method for treating or preventing an immune disorder comprising administering to a subject in need of such treatment and effective amount of the purified antagonist of claim 14.
16. A method for treating or preventing cancer comprising administering to a subject in need of such treatment an effective amount of the purified antagonist of claim 14.
17. A method for detecting a polynucleotide which encodes a serine protease precursor protein in a biological sample comprising the steps of:
 - a) hybridizing the polynucleotide of claim 6 to nucleic acid material of a

biological sample, thereby forming a hybridization complex; and

b) detecting said hybridization complex, wherein the presence of said complex correlates with the presence of a polynucleotide encoding a serine protease precursor protein in said biological sample.

5

1/8

NAC	ACC	AAC	AGC	TCT	GAC	CTG	GGC	AGC	CCT	CTT	CCT	GAG	AAA	ATG	CAG	CCA	TTC	CTC
9	18	27	36	45	54													
L	L	F	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L	L
63	72	81	90	99	108													
CTC	CTG	TTG	GCC	TTT	CTT	CTG	ACC	CCT	GGG	GCT	GGG	GAG	ACA	GAG	GAG	ATC	ATC	GGG
L	L	A	A	F	L	L	T	P	G	A	G	T	E	E	I	I	I	G
117	126	135	144	153	162													
GGC	CAT	GAG	GCC	AAG	CCC	CAC	TCC	CGC	CCC	TAC	ATG	GCC	TTT	TTT	GTT	CAG	TTT	CTG
G	H	E	A	K	P	H	S	R	P	Y	M	A	F	V	Q	Q	F	L
171	180	189	198	207	216													
CAA	GAG	AAG	AGT	CGG	AAG	AGG	TGT	GGC	GGC	ATC	CTA	GTG	AGA	AAG	GAC	GAC	TTT	GTG
Q	E	K	S	R	K	R	C	G	G	I	L	V	R	K	D	F	F	V
225	234	243	252	261	270													
CTG	ACA	GCT	GCT	CAC	TGC	CAG	GGA	AGC	TCC	ATA	AAT	GTC	ACC	TTG	GGG	GCC	CAC	
L	T	A	A	H	C	Q	G	S	S	I	N	V	T	L	G	A	H	
279	288	297	306	315	324													
AAT	ATC	AAA	GAA	CAG	GAG	CCG	ACC	CAG	CAG	TTT	ATC	CCT	GTG	AAA	AGA	CCC	ATC	
N	I	K	E	Q	E	P	T	Q	Q	F	I	P	V	K	R	P	I	
333	342	351	360	369	378													
CCC	CAT	CCA	GCC	TAT	AAT	CCT	AAG	AAC	TTC	TCC	AAC	GAC	ATC	ATG	CTA	CTG	CAG	
P	H	P	A	Y	N	P	K	N	F	S	N	D	I	M	L	L	L	Q

FIGURE 1A

387	CTG GAG AGA AAG GCC AAG CGG ACC AGA GCT GTG CAG CCC CTC AGG CTA CCT AGC	405	414	423	432
	L E R K A K A AAG GCC AAG CGG ACC AGA GCT GTG CAG CCC CTC AGG CTA CCT AGC				
441	AAC AAG GCC CAG GTG AAG CCA GGG CAG ACA TGC AGT GTG GCC TGG GGG CAG	459	468	477	486
	N K A Q V K K AAG CCA GGG CAG ACA TGC AGT GTG GCC TGG GGG CAG				
495	ACG GCC CCC CTG GGA AAA CAC TCA CAC ACA CTA CAA GAG GTG AAG ATG ACA GTG	513	522	531	540
	T A P L G K K AAG CCA GGG CAG ACA TGC AGT GTG GCC TGG GGG CAG				
549	CAG GAA GAT CGA AAG TGC GAA TCT GAC TTTA CGC CAT TAT TAC TAC AGT ACC ATT	567	576	585	594
	Q E D R K C E AAG TGC GAA TCT GAC TTTA CGC CAT TAT TAC TAC AGT ACC ATT				
603	GAG TTG TGC GTG GGG GAC CCA GAG ATT AAA AAG ACT TCC TTTT AAG GGG GAC TCT	621	630	639	648
	E L C V G D D AAG CCA GAG ATT AAA AAG ACT TCC TTTT AAG GGG GAC TCT				
657	GGA GGC CCT CTT GTG TGT AAC AAG GTG GCC CAG GGC ACT TCC TTTT AAG GGG GAC TCT	675	684	693	702
	G G P L V C N AAG GTG GCC CAG GGC ACT TCC TTTT AAG GGG GAC TCT				
711	AAC AAT GGC ATG CCT CCA CGA GCC TGC ACC AAA GTC TCA AGC TTTT GTA CAC TGG	729	738	747	756
	N N G M P P CCA CGA GCC TGC ACC AAA GTC TCA AGC TTTT GTA CAC TGG				

FIGURE 1B

3 / 8

765	774	783	792	801	810
ATA AAG AAA ACC ATG AAA CGC TAC TAA CTA CAG GAA GCA AAC TAA GCC CCC GCT					
I K K T M K R Y					
819	828	837	846	855	864
GTA ATG AAA CAC CTT CTC TGG AGC CAA GTC CAG ATT TAC ACT GGG AGA GGT GCC					
873	882				
AGC AAC TGA ATA AAT CCT CTT					

FIGURE 1C

1	M	Q	P	F	L	L	L	A	F	L	L	T	P	G	A	G	T	E	E	I	I	G	G	H	E	A	K	P	H	HSPP	
1	M	K	L	L	L	L	L	S	F	S	L	A	P	K	T	E	A	G	E	I	I	G	G	H	E	A	K	P	H	g206690	
1	M	Q	P	I	L	L	L	A	F	L	L	L	P	R	A	D	A	G	E	I	I	G	G	H	E	A	K	P	H	g306682	
31	S	R	P	Y	M	A	F	V	Q	F	L	Q	E	-	K	S	R	K	R	C	G	G	I	L	V	R	K	D	F	V	HSPP
31	S	R	P	Y	M	A	Y	L	Q	I	M	D	E	Y	S	G	S	K	K	C	G	G	F	L	I	R	E	D	F	V	g206690
31	S	R	P	Y	M	A	Y	L	M	I	W	D	Q	-	K	S	L	K	R	C	G	G	F	L	I	Q	D	D	F	V	g306682
60	L	T	A	A	H	C	Q	G	S	S	I	N	V	T	L	G	A	H	N	I	K	E	Q	E	P	T	Q	Q	F	I	HSPP
61	L	T	A	A	H	C	S	G	S	K	I	N	V	T	L	G	A	H	N	I	K	E	Q	E	K	M	Q	Q	I	I	g206690
60	L	T	A	A	H	C	W	G	S	S	I	N	V	T	L	G	A	H	N	I	K	E	Q	E	P	T	Q	Q	F	I	g306682
90	P	V	K	R	P	I	P	H	P	A	Y	N	P	K	N	F	S	N	D	I	M	L	L	Q	L	E	R	K	A	K	HSPP
91	P	V	K	I	I	P	H	P	A	Y	N	S	K	T	I	S	N	D	I	M	L	L	K	L	K	S	K	A	K	g206690	
90	P	V	K	R	P	I	P	H	P	A	Y	N	P	K	N	F	S	N	D	I	M	L	L	Q	L	E	R	K	A	K	g306682
120	R	T	R	A	V	Q	P	L	R	L	P	S	N	K	A	Q	V	K	P	G	Q	T	C	S	V	A	G	W	G	Q	HSPP
121	R	S	S	A	V	K	P	L	N	L	P	R	R	N	V	K	V	K	P	G	D	V	C	Y	V	A	G	W	G	K	g206690
120	R	T	R	A	V	Q	P	L	R	L	P	S	N	K	A	Q	V	K	P	G	Q	T	C	S	V	A	G	W	G	Q	g306682

FIGURE 2A

5/8

150	T	A	P	L	G	K	H	S	H	T	L	Q	E	V	K	M	T	V	Q	E	D	R	K	C	E	S	D	L	R	H	HSPP
151	L	G	P	M	G	K	Y	S	D	T	L	Q	E	V	E	L	T	V	Q	E	D	Q	K	C	E	S	Y	L	K	N	g206690
150	T	A	P	L	G	K	H	S	H	T	L	Q	E	V	K	M	T	V	Q	E	D	R	K	C	E	S	D	L	R	H	g306682
180	Y	Y	D	S	T	I	E	L	C	V	G	D	P	E	I	K	K	T	S	F	K	G	D	S	G	G	P	L	V	C	HSPP
181	Y	F	D	K	A	N	E	I	C	A	G	D	P	K	I	K	R	A	S	F	R	G	D	S	G	G	P	L	V	C	g206690
180	Y	Y	D	S	T	I	E	L	C	V	G	D	P	E	I	K	K	T	S	F	K	G	D	S	G	G	P	L	V	C	g306682
210	N	K	V	A	Q	G	I	V	S	Y	G	R	N	N	G	M	P	P	R	A	C	T	K	V	S	S	F	V	H	W	HSPP
211	K	K	V	A	A	G	I	V	S	Y	G	Q	N	D	G	S	T	P	R	A	F	T	K	V	S	T	F	L	S	W	g206690
210	N	K	V	A	Q	G	I	V	S	Y	G	R	N	N	G	M	P	P	R	A	C	T	K	V	S	S	F	V	H	W	g306682
240	I	K	K	T	M	K	R	Y																							HSPP
241	I	K	K	T	M	K	K	S																							g206690
240	I	K	K	T	M	K	R	Y																							g306682

FIGURE 2B




FIGURE3A

7/8




FIGURE3B




FIGURE 3C