



US00PP10424P

# United States Patent [19]

[11] Patent Number: Plant 10,424

Bolwijn

[45] Date of Patent: Jun. 2, 1998

[54] EUONYMUS PLANT NAMED 'INTERBOLWI'

[56] References Cited

[75] Inventor: **Gustaaf Emiel Ferdinand Bolwijn**,  
Putten, Netherlands

U.S. PATENT DOCUMENTS

P.P. 4,340 12/1978 Nielsen ..... Plt./63  
P.P. 6,127 3/1988 Dugan ..... Plt./63

[73] Assignee: **Spring Meadow Nursery, Inc.**, Grand  
Haven, Mich.

Primary Examiner—James R. Feyrer  
Attorney, Agent, or Firm—C. A. Whealy

[21] Appl. No.: **802,396**

[57] ABSTRACT

[22] Filed: **Feb. 18, 1997**

A distinct cultivar of *Euonymus* plant named 'Interbolwi', characterized by its showy green and yellow variegated foliage; compact, spreading and rounded plant form; moderate growth rate; flat leaves; and excellent resistance to leaf spot diseases. It is intended that the new *Euonymus* will be marketed under the trade name *Blondy*.

[51] Int. Cl.<sup>6</sup> ..... **A01H 5/00**

[52] U.S. Cl. .... **Plt./63**

[58] Field of Search ..... **Plt./63**

**2 Drawing Sheets**

1

2

The present invention relates to a new and distinct cultivar of *Euonymus* plant, botanically known as *Euonymus fortunei*, and hereinafter referred to by the cultivar name *Interbolwi*. It is intended that the new *Euonymus* will be marketed under the trade name *Blondy*.

The new cultivar is a naturally-occurring leaf color mutation of the *Euonymus fortunei* cultivar *Sunspot* (disclosed in U.S. Plant Pat. No. 4,340). In 1988, the new cultivar was discovered by the inventor in a controlled environment in Putten, The Netherlands, within a population of plants of the cultivar *Sunspot*.

Asexual reproduction of the new cultivar by hardwood or softwood cuttings taken at Putten, The Netherlands, has shown that the unique features of this new *Euonymus* plant are stable and reproduced true to type in successive generations of asexual reproduction.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Interbolwi'. These characteristics in combination distinguish 'Interbolwi' as a new and distinct cultivar:

1. Showy green and yellow variegated foliage.
2. Compact, spreading and rounded plant form.
3. Moderate growth rate.
4. Flat leaves.
5. Excellent resistance to leaf spot diseases.

In side-by-side comparisons in Grand Haven, Mich., under commercial practice, plants of the new *Euonymus* differed from plants of the parent cultivar *Sunspot* in the following characteristics:

1. Plants of the new *Euonymus* are more compact than plants of the cultivar *Sunspot*.
2. Plants of the new *Euonymus* are more spreading than plants of the cultivar *Sunspot*.
3. Stems of plants of the new *Euonymus* have more green streaking than stems of plants of the cultivar *Sunspot*.
4. Upper foliage surfaces of plants of the new *Euonymus* have a dull appearance whereas upper foliage surfaces of plants of the cultivar *Sunspot* are glossy.
5. Leaves of the new *Euonymus* are flatter than leaves of the cultivar *Sunspot*.
6. Foliage variegation is more pronounced on plants of the new *Euonymus* compared to plants of the cultivar *Sunspot*. Plants of the new *Euonymus* are considered more showy than plants of the cultivar *Sunspot*.
7. Foliage variegation color on plants of the new *Euonymus* is more golden than foliage variegation color on plants of the cultivar *Sunspot*.

8. Plants of the new *Euonymus* are more resistant to leaf spot diseases than plants of the cultivar *Sunspot*.

Plants of the new *Euonymus* can be compared to plants of the cultivar *Moonshadow* (disclosed in U.S. Plant Pat. No. 6,127), an *Euonymus* cultivar that is also a naturally-occurring mutation of the cultivar *Sunspot*. In side-by-side comparisons in Grand Haven, Mich., under commercial practice, plants of the new *Euonymus* differed from plants of the cultivar *Moonshadow* in the following characteristics:

1. Plants of the new *Euonymus* are more compact than plants of the cultivar *Moonshadow*.
2. Plants of the new *Euonymus* are more vigorous than plants of the cultivar *Moonshadow*.
3. Stems of plants of the new *Euonymus* have more green streaking than stems of plants of the cultivar *Moonshadow*.
4. Upper foliage surfaces of plants of the new *Euonymus* have a dull appearance whereas upper foliage surfaces of plants of the cultivar *Moonshadow* are glossy.
5. Leaves of the new *Euonymus* are flatter and wider than plants of the cultivar *Moonshadow*.
6. Foliage variegation is more pronounced on plants of the new *Euonymus* compared to plants of the cultivar *Moonshadow*. Plants of the new *Euonymus* are considered more showy than plants of the cultivar *Moonshadow*.
7. Foliage variegation color on plants of the new *Euonymus* is more golden than foliage variegation color on plants of the cultivar *Moonshadow*.

8. Plants of the new *Euonymus* are more resistant to leaf spot diseases than plants of the cultivar *Moonshadow*.

The cultivar *Interbolwi* has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The following observations, measurements and comparisons describe plants grown in Grand Haven, Mich., in a double polyethylene-covered greenhouse with day temperatures ranging from 5 to 26.5C and night temperatures ranging from 5 to 15.5C.

The accompanying colored photographs illustrate the overall appearance of the new *Euonymus*, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type.

The first photograph comprises a top perspective view of typical plants of the cultivar *Interbolwi*.

The second photograph comprises a close-up view of individual stems of the cultivar Interbolwi showing the green and yellow variegation pattern. Foliage colors in the photographs may appear different than the actual colors due to light reflectance.

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

*Botanical classification:* *Euonymus fortunei* cultivar Interbolwi.

*Parentage:* Naturally-occurring leaf color mutation of *Euonymus fortunei* cultivar Sunspot (U.S. Plant Pat. No. 4,340).

*Propagation:*

*Type cutting.*—Hardwood or softwood cuttings.

*Time to initiate and develop roots.*—Summer: About 21 days at 22C. Winter: About 40 days at 12C.

*Rooting habit.*—Fleshy and well-branched. Grey-brown aerial roots are occasionally observed.

*Plant description:*

*Form.*—Compact, rounded, spreading perennial plant.

*Branching habit.*—Basally branching, pruning increases branching.

*Plant height.*—38 to 50 cm from soil level to top of plant plane.

*Plant diameter or area of spread.*—60 to 90 cm.

*Vigor.*—Moderate.

*Stem description.*—Lateral branch diameter: About 2 mm. Internode length: 1.5 to 2.5 cm. Color: 14C/14D with green, 143C, streaks. Texture: Glabrous.

*Foliage description.*—Leaves single and generally symmetrical, opposite arrangement. Usually 10 to 20 leaves per lateral branch. Size (largest leaves): Length: 2 to 4 cm. Width: 1 to 2.2 cm. Shape: Ovate to elliptic. Apex: Obtuse. Base: Cuneate. Margin: Crenate/serrate. Texture: Smooth and leathery with a dull finish. Aspect: Flat. Color: Variegated with yellow and green patterns that are variable. Usually at least 60% of the leaf surface area is yellow. Young and fully expanded leaves: Upper surface: Center: 14C/14D. Margin: 147A. Upper surface: Center: 14D Margin: 147A. Petiole: Length: 3 to 5 mm. Diameter: 1 mm or less. Color: 14C/14D.

*Flower description:* Flowers have not been observed on plants of the new *Euonymus* grown to date.

*Disease resistance:* Plants of the new *Euonymus* have shown excellent resistance to leaf spot diseases.

It is claimed:

1. A new and distinct cultivar of *Euonymus* plant named 'Interbolwi', as illustrated and described.

\* \* \* \* \*



