



(51) International Patent Classification:

G01N 33/50 (2006.01)

(21) International Application Number:

PCT/IB2013/060458

(22) International Filing Date:

27 November 2013 (27.11.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/732,251 30 November 2012 (30.11.2012) US

13/802,117 13 March 2013 (13.03.2013) US

(71) Applicant: NESTEC S.A. [CH/CH]; Avenue Nestlé 55, CH-1800 Vevey (CH).

(72) Inventors: HAUENSTEIN, Scott; 3565 Caminito El Rincon #211, San Diego, California 92130 (US). SINGH, Sharat; PMB 69, P.O. Box 5000, Rancho Santa Fe, California 92067 (US).

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

(54) Title: ASSAYS FOR DETECTING NEUTRALIZING AUTOANTIBODIES TO BIOLOGIC THERAPY

Binding (non-NAB) Antibody



Neutralizing Antibody (NAB)



(57) Abstract: The present invention provides assays for detecting and measuring the presence or level of neutralizing and non-neutralizing autoantibodies to biologics such as anti-TNF α drug therapeutics in a sample. The present invention is useful for monitoring the formation of neutralizing and/or non-neutralizing anti-drug antibodies over time while a subject is on biologic therapy. The present invention is also useful for predicting and/or determining the cross-reactivity of neutralizing anti-drug antibodies in a subject's sample with alternative biologic therapies. As such, the present invention provides information for guiding treatment decisions for those subjects receiving therapy with a biologic agent and improves the accuracy of optimizing therapy, reducing toxicity, and/or monitoring the efficacy of therapeutic treatment to biologic therapy.

FIG. 22

ASSAYS FOR DETECTING NEUTRALIZING AUTOANTIBODIES TO BIOLOGIC THERAPY

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Application No. 13/802,117, filed March 13, 2013, which is a continuation-in-part of PCT Application No. PCT/US2012/045794, filed July 6, 2012, which claims priority to U.S. Provisional Application No. 61/505,031, filed July 6, 2011, U.S. Provisional Application No. 61/528,072, filed August 26, 2011, and U.S. Provisional Application No. 61/535,884, filed September 16, 2011. This application also claims priority to U.S. Provisional Application No. 61/732,251, filed November 30, 2012. The disclosures of each of these applications are hereby incorporated by reference in their entirety for all purposes.

BACKGROUND OF THE INVENTION

[0002] Autoimmune disorders are a significant and widespread medical problem. For example, rheumatoid arthritis (RA) is an autoimmune disease affecting more than two million people in the United States. RA causes chronic inflammation of the joints and typically is a progressive illness that has the potential to cause joint destruction and functional disability. The cause of rheumatoid arthritis is unknown, although genetic predisposition, infectious agents and environmental factors have all been implicated in the etiology of the disease. In active RA, symptoms can include fatigue, lack of appetite, low grade fever, muscle and joint aches and stiffness. Also during disease flare ups, joints frequently become red, swollen, painful and tender, due to inflammation of the synovium. Furthermore, since RA is a systemic disease, inflammation can affect organs and areas of the body other than the joints, including glands of the eyes and mouth, the lung lining, the pericardium, and blood vessels.

[0003] Traditional treatments for the management of RA and other autoimmune disorders include fast acting "first line drugs" and slower acting "second line drugs." The first line drugs reduce pain and inflammation. Example of such first line drugs include aspirin, naproxen, ibuprofen, etodolac and other non-steroidal anti-inflammatory drugs (NSAIDs), as well as corticosteroids, given orally or injected directly into tissues and joints. The second line drugs promote disease remission and prevent progressive joint destruction and are also referred to as disease-modifying anti-rheumatic drugs or DMARDs. Examples of second line drugs include gold, hydrochloroquine, azulfidine and immunosuppressive agents, such as

methotrexate, azathioprine, cyclophosphamide, chlorambucil and cyclosporine. Many of these drugs, however, can have detrimental side-effects. Thus, additional therapies for rheumatoid arthritis and other autoimmune disorders have been sought.

[0004] Tumor necrosis factor alpha (TNF- α) is a cytokine produced by numerous cell types, including monocytes and macrophages, that was originally identified based on its ability to induce the necrosis of certain mouse tumors. Subsequently, a factor termed cachectin, associated with cachexia, was shown to be identical to TNF- α . TNF- α has been implicated in the pathophysiology of a variety of other human diseases and disorders, including shock, sepsis, infections, autoimmune diseases, RA, Crohn's disease, transplant rejection and graft-versus-host disease.

[0005] Because of the harmful role of human TNF- α (hTNF- α) in a variety of human disorders, therapeutic strategies have been designed to inhibit or counteract hTNF- α activity. In particular, antibodies that bind to, and neutralize, hTNF- α have been sought as a means to inhibit hTNF- α activity. Some of the earliest of such antibodies were mouse monoclonal antibodies (mAbs), secreted by hybridomas prepared from lymphocytes of mice immunized with hTNF- α (*see, e.g.*, U.S. Pat. No. 5,231,024 to Moeller *et al.*). While these mouse anti-hTNF- α antibodies often displayed high affinity for hTNF- α and were able to neutralize hTNF- α activity, their use *in vivo* has been limited by problems associated with the administration of mouse antibodies to humans, such as a short serum half-life, an inability to trigger certain human effector functions, and elicitation of an unwanted immune response against the mouse antibody in a human (the "human anti-mouse antibody" (HAMA) reaction).

[0006] More recently, biological therapies have been applied to the treatment of autoimmune disorders such as rheumatoid arthritis. For example, four TNF α inhibitors, REMICADE™ (infliximab), a chimeric anti-TNF α mAb, ENBREL™ (etanercept), a TNFR-Ig Fc fusion protein, HUMIRA™ (adalimumab), a human anti-TNF α mAb, and CIMZIA® (certolizumab pegol), a PEGylated Fab fragment, have been approved by the FDA for treatment of rheumatoid arthritis. CIMZIA® is also used for the treatment of moderate to severe Crohn's disease (CD). While such biologic therapies have demonstrated success in the treatment of rheumatoid arthritis and other autoimmune disorders such as CD, not all subjects treated respond, or respond well, to such therapy. Moreover, administration of TNF α inhibitors can induce an immune response to the drug and lead to the production of autoantibodies such as human anti-chimeric antibodies (HACA), human anti-humanized

antibodies (HAHA), and human anti-mouse antibodies (HAMA). Such HACA, HAHA, or HAMA immune responses can be associated with hypersensitive reactions and dramatic changes in pharmacokinetics and biodistribution of the immunotherapeutic TNF α inhibitor that preclude further treatment with the drug. Thus, there is a need in the art for assays to
5 detect the presence of autoantibodies to biologic agents such as anti-TNF α drugs in a patient sample to monitor biologic therapy and to guide treatment decisions. The present invention satisfies this need and provides related advantages as well.

BRIEF SUMMARY OF THE INVENTION

[0007] The present invention provides assays for detecting and measuring the presence or
10 level of neutralizing and non-neutralizing autoantibodies to biologics such as anti-TNF α drug therapeutics in a sample. The present invention is useful for monitoring the formation of neutralizing and/or non-neutralizing anti-drug antibodies over time while a subject is on biologic therapy (*e.g.*, anti-TNF α drug therapy). The present invention is also useful for predicting and/or determining the cross-reactivity of neutralizing anti-drug antibodies in a
15 subject's sample with alternative biologic therapies (*e.g.*, alternative anti-TNF α therapies). As such, the present invention provides information for guiding treatment decisions for those subjects receiving therapy with a biologic agent and improves the accuracy of optimizing therapy, reducing toxicity, and/or monitoring the efficacy of therapeutic treatment to biologic therapy.

20 [0008] In one aspect, the present invention provides a method for detecting the presence of a neutralizing and/or non-neutralizing form of an autoantibody to a biologic in a sample, the method comprising:

- (a) contacting the sample with a labeled biologic and a labeled biologic binding moiety to form:
 - 25 (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
 - (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the
30 labeled biologic, the labeled biologic binding moiety, and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled

biologic binding moiety, free labeled biologic, and/or a complex of labeled biologic and labeled biologic binding moiety;

(c) measuring the level of free labeled biologic binding moiety after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled biologic binding moiety peak following size exclusion chromatography (SEC)); and

(d) comparing the level of the free labeled biologic binding moiety measured in step (c) to the level of free labeled biologic binding moiety in a control sample (*e.g.*, by measuring the AUC of the free labeled biologic binding moiety peak following SEC of a reference sample containing only free labeled biologic binding moiety), thereby detecting the presence of a neutralizing and/or non-neutralizing form of the autoantibody.

[0009] In certain embodiments, a neutralizing form of the autoantibody is detected when the level of the free labeled biologic binding moiety measured in step (c) is the same or substantially the same as the level of the free labeled biologic binding moiety in the control sample. In certain other embodiments, a non-neutralizing form of the autoantibody is detected when the level of the free labeled biologic binding moiety measured in step (c) is decreased (*e.g.*, substantially decreased) or absent (*e.g.*, undetectable) compared to the level of the free labeled biologic binding moiety in the control sample.

[0010] In another aspect, the present invention provides a method for measuring the level or percent of a neutralizing form of an autoantibody to a biologic in a sample, the method comprising:

(a) contacting the sample with a labeled biologic and a labeled biologic binding moiety to form:

(i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or

(ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic, the labeled biologic binding moiety, and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);

(b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled

biologic binding moiety, free labeled biologic, and/or a complex of labeled biologic and labeled biologic binding moiety;

(c) measuring the level of free labeled biologic binding moiety after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled biologic binding moiety peak following size exclusion chromatography (SEC)); and

(d) comparing the level of free labeled biologic binding moiety measured in step (c) to a normalized level or percent of free labeled biologic binding moiety in a control sample (*e.g.*, by measuring and normalizing the AUC of the free labeled biologic binding moiety peak following SEC of a reference sample containing only free labeled biologic binding moiety to calculate the level or percent of free labeled biologic binding moiety), wherein the normalized level or percent of the free labeled biologic binding moiety in the control sample corresponds to the level or percent of a neutralizing form of the autoantibody.

[0011] In some embodiments, the difference between the normalized level or percent of the free labeled biologic binding moiety in the control sample and the level of free labeled biologic binding moiety measured in step (c) corresponds to the level or percent of a non-neutralizing form of the autoantibody.

[0012] In yet another aspect, the present invention provides a method for measuring the percent or the level of a neutralizing form of an autoantibody to a biologic in a sample, the method comprising:

(a) contacting the sample with a labeled biologic and a labeled biologic binding moiety to form:

(i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or

(ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic, the labeled biologic binding moiety, and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);

(b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled

biologic binding moiety, free labeled biologic, and/or a complex of labeled biologic and labeled biologic binding moiety;

(c) measuring the level of free labeled biologic binding moiety after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled biologic binding moiety peak following size exclusion chromatography (SEC)); and

(d) comparing the level of free labeled biologic binding moiety measured in step (c) to the level of free labeled biologic binding moiety in a control sample (*e.g.*, by measuring the AUC of the free labeled biologic binding moiety peak following SEC of a reference sample), thereby measuring the percent or level of a neutralizing form of the autoantibody.

[0013] In another aspect, the present invention provides a method for determining whether a neutralizing form of an autoantibody to a first biologic is cross-reactive with a second (*i.e.*, different) biologic, the method comprising:

(a) detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample in accordance with an assay described herein to determine whether the sample is positive or negative for the neutralizing form of the autoantibody; and

if the sample is positive for the neutralizing form of the autoantibody, then:

(b) contacting the sample with a labeled second biologic to form a labeled complex of the labeled second biologic and the neutralizing form of the autoantibody (*i.e.*, wherein the components of the labeled complex are not covalently attached to each other);

(c) subjecting the labeled complex to size exclusion chromatography to separate the labeled complex (*e.g.*, from free labeled second biologic); and

(d) detecting the labeled complex, thereby determining whether a neutralizing form of an autoantibody to a first biologic is cross-reactive with a second biologic.

[0014] In certain embodiments, the presence of the labeled complex is an indication that the neutralizing autoantibody against the first biologic is cross-reactive with the second biologic, *i.e.*, the neutralizing autoantibody will inhibit the activity of both the first and second biological drugs.

[0015] In certain other embodiments, the absence of the labeled complex is an indication that the neutralizing autoantibody against the first biologic is not cross-reactive with the

second biologic, *i.e.*, the neutralizing autoantibody will not inhibit the activity of the second biological drug.

[0016] In some embodiments, the biologic includes antibodies (*e.g.*, anti-TNF α monoclonal antibodies), antibody fragments, proteins (*e.g.*, cytokines such as interleukins), polypeptides, peptides, fusion proteins, multivalent binding proteins, antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and combinations thereof.

[0017] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving biologic therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis), an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)), or cancer.

[0018] In certain embodiments, the sample has or is suspected of having an autoantibody to the biologic. In other embodiments, the biologic autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0019] In certain aspects, the assay methods of the present invention further comprise an acid dissociation step comprising contacting a sample with an acid prior to, during, and/or after contacting the sample with a labeled biologic and a labeled biologic binding moiety.

[0020] In certain other aspects, the assay methods of the present invention comprise detecting the presence or level of one or more isotypes of a neutralizing and/or non-neutralizing form of an autoantibody to a biologic in a sample.

[0021] In one particular aspect, the present invention provides a method for detecting the presence of a neutralizing and/or non-neutralizing form of an autoantibody to an anti-TNF α drug in a sample, the method comprising:

(a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to form:

(i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or

(ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug, the labeled TNF α , and the autoantibody (*i.e.*,

wherein the components of the second labeled complex are not covalently attached to each other);

- 5 (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled TNF α , free labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and labeled TNF α ;
- (c) measuring the level of free labeled TNF α after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled TNF α peak following size exclusion chromatography (SEC)); and
- 10 (d) comparing the level of the free labeled TNF α measured in step (c) to the level of free labeled TNF α in a control sample (*e.g.*, by measuring the AUC of the free labeled TNF α peak following SEC of a reference sample containing only free labeled TNF α), thereby detecting the presence of a neutralizing and/or non-neutralizing form of the autoantibody.

15 [0022] In certain embodiments, a neutralizing form of the autoantibody is detected when the level of the free labeled TNF α measured in step (c) is the same or substantially the same as the level of the free labeled TNF α in the control sample. In certain other embodiments, a non-neutralizing form of the autoantibody is detected when the level of the free labeled TNF α measured in step (c) is decreased (*e.g.*, substantially decreased) or absent (*e.g.*, undetectable)

20 compared to the level of the free labeled TNF α in the control sample.

[0023] In another particular aspect, the present invention provides a method for measuring the level or percent of a neutralizing form of an autoantibody to an anti-TNF α drug in a sample, the method comprising:

- 25 (a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to form:
- (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
- 30 (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug, the labeled TNF α , and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);

- (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled TNF α , free labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and labeled TNF α ;
- 5 (c) measuring the level of free labeled TNF α after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled TNF α peak following size exclusion chromatography (SEC)); and
- (d) comparing the level of free labeled TNF α measured in step (c) to a normalized level or percent of free labeled TNF α in a control sample (*e.g.*, by measuring and normalizing the AUC of the free labeled TNF α peak following SEC of a reference sample containing only free labeled TNF α to calculate the level or percent of free labeled TNF α), wherein the normalized level or percent of the free labeled TNF α in the control sample corresponds to the level or percent of a neutralizing form of the autoantibody.
- 10

15 [0024] In some embodiments, the difference between the normalized level or percent of the free labeled TNF α in the control sample and the level of free labeled TNF α measured in step (c) corresponds to the level or percent of a non-neutralizing form of the autoantibody.

[0025] In yet another particular aspect, the present invention provides a method for measuring the percent or the level of a neutralizing form of an autoantibody to an anti-TNF α drug in a sample, the method comprising:

20

- (a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to form:
- (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
- 25 (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug, the labeled TNF α , and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- 30 (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled TNF α , free labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and labeled TNF α ;

- (c) measuring the level of free labeled TNF α after size exclusion chromatography (e.g., by measuring the area under the curve (AUC) of the free labeled TNF α peak following size exclusion chromatography (SEC)); and
- (d) comparing the level of free labeled TNF α measured in step (c) to the level of free labeled TNF α in a control sample (e.g., by measuring the AUC of the free labeled TNF α following SEC of a reference sample), thereby measuring the percent or level of a neutralizing form of the autoantibody.

- [0026] In still yet another particular aspect, the present invention provides a method for determining whether a neutralizing form of an autoantibody to a first anti-TNF α drug is cross-reactive with a second (*i.e.*, different) anti-TNF α drug, the method comprising:
- (a) detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample in accordance with an assay described herein to determine whether the sample is positive or negative for the neutralizing form of the autoantibody; and
- if the sample is positive for the neutralizing form of the autoantibody, then:
- (b) contacting the sample with a labeled second anti-TNF α drug to form a labeled complex of the labeled second anti-TNF α drug and the neutralizing form of the autoantibody (*i.e.*, wherein the components of the labeled complex are not covalently attached to each other);
- (c) subjecting the labeled complex to size exclusion chromatography to separate the labeled complex (e.g., from free labeled second anti-TNF α drug); and
- (d) detecting the labeled complex, thereby determining whether a neutralizing form of an autoantibody to a first anti-TNF α drug is cross-reactive with a second anti-TNF α drug.

- [0027] In certain embodiments, the presence of the labeled complex is an indication that the neutralizing autoantibody against the first anti-TNF α drug is cross-reactive with the second anti-TNF α drug, *i.e.*, the neutralizing autoantibody will inhibit the activity of both the first and second anti-TNF α drugs.

- [0028] In certain other embodiments, the absence of the labeled complex is an indication that the neutralizing autoantibody against the first anti-TNF α drug is not cross-reactive with the second anti-TNF α drug, *i.e.*, the neutralizing autoantibody will not inhibit the activity of the second anti-TNF α drug.

[0029] In some embodiments, the anti-TNF α drug is selected from the group consisting of REMICADE™ (infliximab), ENBREL™ (etanercept), HUMIRA™ (adalimumab), CIMZIA® (certolizumab pegol), SIMPONI® (golimumab; CNTO 148), and combinations thereof.

5 [0030] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving anti-TNF α drug therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a TNF α -mediated disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis) or an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)).

10 [0031] In certain embodiments, the sample has or is suspected of having an autoantibody to the anti-TNF α drug. In other embodiments, the anti-TNF α drug autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

15 [0032] In certain aspects, the assay methods of the present invention further comprise an acid dissociation step comprising contacting a sample with an acid prior to, during, and/or after contacting the sample with a labeled anti-TNF α drug and a labeled TNF α .

[0033] In certain other aspects, the assay methods of the present invention comprise detecting the presence or level of one or more isotypes of a neutralizing and/or non-neutralizing form of an autoantibody to an anti-TNF α drug in a sample.

20 [0034] In a further aspect, the present invention provides a method for monitoring and/or optimizing therapy to a biologic in a subject receiving a course of therapy with the biologic, the method comprising:

- (a) detecting or measuring the presence, level, or percent of a neutralizing form of an autoantibody to the biologic in accordance with the assay described herein at a plurality of time points over the course of therapy;
- 25 (b) detecting a change in the presence, level, or percent of the neutralizing form of the autoantibody over time; and
- (c) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the presence, level, or percent of the neutralizing
- 30 form of the autoantibody over time.

[0035] In one particular aspect, the present invention provides a method for monitoring and/or optimizing therapy to a biologic in a subject receiving a course of therapy with the biologic, the method comprising:

- 5 (a) measuring the level or percent of a neutralizing form of an autoantibody to the biologic in a first sample from the subject as described herein at time point t_0 ;
- (b) measuring the level or percent of the neutralizing form of the autoantibody in a second sample from the subject as described herein at time point t_1 ;
- (c) optionally repeating step (b) with n additional samples from the subject at time points t_{n+1} , wherein n is an integer from 1 to about 25 (*e.g.*, n is 1, 2, 3, 4, 5, 6,
10 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or any range therein);
- (d) detecting a change in the level or percent of the neutralizing form of the autoantibody from time points t_0 to t_1 or from time points t_0 to t_{n+1} ; and
- 15 (e) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the level or percent of the neutralizing form of the autoantibody over time.

[0036] In an additional aspect, the present invention provides a method for optimizing therapy and/or reducing toxicity in a subject receiving a course of therapy with a first
20 biologic, the method comprising:

- (a) determining whether a neutralizing form of an autoantibody to the first biologic is cross-reactive with a second (*i.e.*, different) biologic by detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample from the subject in accordance with an assay
25 described herein; and
- (b) determining that a different course of therapy should be administered to the subject if the neutralizing form of the autoantibody is cross-reactive with the second biologic.

[0037] In one particular aspect, the present invention provides a method for monitoring and/or optimizing therapy to an anti-TNF α drug in a subject receiving a course of therapy
30 with the anti-TNF α drug, the method comprising:

- (a) detecting or measuring the presence, level, or percent of a neutralizing form of an autoantibody to the anti-TNF α drug in accordance with the assay described herein at a plurality of time points over the course of therapy;

- (b) detecting a change in the presence, level, or percent of the neutralizing form of the autoantibody over time; and
- (c) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the presence, level, or percent of the neutralizing form of the autoantibody over time.

[0038] In another particular aspect, the present invention provides a method for monitoring and/or optimizing therapy to an anti-TNF α drug in a subject receiving a course of therapy with the anti-TNF α drug, the method comprising:

- (a) measuring the level or percent of a neutralizing form of an autoantibody to the anti-TNF α drug in a first sample from the subject as described herein at time point t_0 ;
- (b) measuring the level or percent of the neutralizing form of the autoantibody in a second sample from the subject as described herein at time point t_1 ;
- (c) optionally repeating step (b) with n additional samples from the subject at time points t_{n+1} , wherein n is an integer from 1 to about 25 (*e.g.*, n is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or any range therein);
- (d) detecting a change in the level or percent of the neutralizing form of the autoantibody from time points t_0 to t_1 or from time points t_0 to t_{n+1} ; and
- (e) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the level or percent of the neutralizing form of the autoantibody over time.

[0039] In yet another particular aspect, the present invention provides a method for optimizing therapy and/or reducing toxicity in a subject receiving a course of therapy with a first anti-TNF α drug, the method comprising:

- (a) determining whether a neutralizing form of an autoantibody to the first anti-TNF α drug is cross-reactive with a second (*i.e.*, different) anti-TNF α drug by detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample from the subject in accordance with an assay described herein; and

- (b) determining that a different course of therapy should be administered to the subject if the neutralizing form of the autoantibody is cross-reactive with the second anti-TNF α drug.

[0040] Other objects, features, and advantages of the present invention will be apparent to one of skill in the art from the following detailed description and figures.

BRIEF DESCRIPTION OF THE DRAWINGS

- [0041] Figure 1 illustrates that there was a clear relationship between NAb percent (y-axis) and ATI levels.
- [0042] Figure 2 illustrates that an ATI concentration ≥ 60 U/ml is predictive of NAb+.
- 10 [0043] Figure 3 illustrates that ATI predicts NAb with a ROC AUC of 0.931.
- [0044] Figure 4 illustrates detection of ATI by the fluid phase mobility shift assay of the present invention.
- [0045] Figure 5 illustrates an exemplary ATI/IFX fluid phase mobility shift assay of the present invention.
- 15 [0046] Figure 6 illustrates a non-neutralizing anti-drug antibody (ADA) assay of the present invention.
- [0047] Figure 7 illustrates a neutralizing ADA assay of the present invention.
- [0048] Figure 8 illustrates the levels of IFX and ATI over a time course of 5 samples from a UC patient taken 1, 2, or 3 months apart.
- 20 [0049] Figure 9 shows peak analysis to determine the percentage of free TNF α over time in a UC patient.
- [0050] Figure 10 illustrates a shift from the presence of non-neutralizing autoantibodies to neutralizing autoantibodies over time as exemplified in 3 samples from a UC patient taken 2 or 3 months apart and spiked with IFX.
- 25 [0051] Figure 11 shows peak analysis to determine the percentage of free TNF α over time in samples from a UC patient that were spiked with IFX.
- [0052] Figure 12 shows the use of rabbit anti-human IgG1 Fc as a non-neutralizing antibody (Ab) control.

- [0053] Figure 13 shows the use of ATI positive serum as a mixed neutralizing antibody (NAb)/non-neutralizing antibody (Ab) control.
- [0054] Figure 14 shows that purification of ATI from ATI positive serum results in loss of weaker affinity NAb.
- 5 [0055] Figure 15 illustrates peak analysis from a UC patient case study to determine the percentage of free TNF α in these various controls.
- [0056] Figure 16 shows a peak analysis from a CD patient case study to determine the percentage of free TNF α over a time course of 4 samples taken 7 or 8 weeks apart during a 30-week period.
- 10 [0057] Figure 17 shows a peak analysis from another CD patient case study to determine the percentage of free TNF α over a time course of 3 samples taken during a 50-week period.
- [0058] Figure 18 shows a peak analysis from 4 additional CD patient case studies to determine the percentage of free TNF α in a sample at a particular week during or after induction or maintenance of therapy.
- 15 [0059] Figure 19 shows detection of non-neutralizing antibody activity via the mobility shift assay.
- [0060] Figure 20 depicts the cross-reactivity of ADA against both IFX and ADL, wherein the binding site of ADA mimics the binding site of TNF α and can therefore bind to multiple anti-TNF drugs.
- 20 [0061] Figure 21 shows two patient examples (Patients 1 and 2) in which cross-reactivity of NAb produced in response to one anti-TNF drug was determined for other anti-TNF drugs. In particular, NAb which developed when the patient was on Remicade (IFX) were tested against Humira (ADL).
- [0062] Figure 22 shows exemplary embodiments of the assays of the present invention to
25 detect the presence of non-neutralizing antibodies (non-NAb) (top) or neutralizing antibodies (NAb) (bottom) against a drug such as IFX or ADL.
- [0063] Figure 23 shows the generation and use of a NAb standard curve.
- [0064] Figure 24 provides the results of a case study for Patient 3, who was treated with IFX but lost response to IFX, to determine the cross-reactivity of NAb generated against IFX
30 to ADL.

[0065] Figure 25 provides the results of a case study for Patient 4, who was treated with IFX but lost response to IFX, to determine the cross-reactivity of NAb generated against IFX to ADL.

5 [0066] Figure 26 shows non-limiting examples of patient studies which demonstrate ATI affinity maturation and the development of cross-reactive ATI.

[0067] Figure 27 illustrates one embodiment of the competitive ligand binding NAb assay of the invention. Free TNF-532 represents the maximum assay signal. TNF-532/IFX-488 negative control containing no ATI represents minimum assay signal. (A) Binding ATI that does not affect assay signal. (B) Neutralizing ATI patient serum that restores assay signal.

10 [0068] Figure 28 shows the clinical utility of the neutralizing assay of the invention.

[0069] Figure 29 shows the performance characteristics of the neutralizing assay of the invention.

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

15 [0070] The present invention is based in part on the discovery that a homogeneous mobility shift assay using size exclusion chromatography and optionally acid dissociation to enable equilibration of immune complexes is particularly advantageous for measuring the presence or level of neutralizing and non-neutralizing forms of autoantibodies (*e.g.*, HACA, HAHA, *etc.*) that are generated against biologics such as anti-TNF α drugs. Such autoantibodies are
20 also known as anti-drug antibodies or ADA, and neutralizing and non-neutralizing forms thereof are also known as NAb and non-NAb, respectively.

[0071] In particular embodiments, the homogeneous mobility shift assays of the invention are performed by contacting a subject's sample with (*e.g.*, fluorescently) labeled biologic (*e.g.*, anti-TNF α drug) and (*e.g.*, fluorescently) labeled biologic binding moiety (*e.g.*, TNF α).

25 The assays described herein are advantageous for at least the following reasons: they obviate the need for wash steps which remove low affinity ADA; they use distinct labels such as fluorophores that allow for detection on the visible, IR, and/or near IR (NIR) spectra which decreases background and serum interference issues; they increase the ability to detect neutralizing and/or non-neutralizing ADA in subjects with a low titer due to the high
30 sensitivity of fluorescent label detection; and they occur as a liquid phase reaction, thereby reducing the chance of any changes in the epitope by attachment to a solid surface such as an ELISA plate.

- [0072] In exemplary embodiments, the assays of the present invention are advantageous because they enable time course case studies of IBD subjects on anti-TNF α drug therapy for monitoring the formation of neutralizing and/or non-neutralizing anti-drug antibodies in multiple samples at different time points. The assays of the present invention are also
5 advantageous because they enable the determination of whether there is a shift from non-neutralizing to neutralizing anti-drug antibodies over time while a subject is on anti-TNF α drug therapy. Without being bound to any particular theory, neutralizing anti-drug antibodies may have significant negative clinical consequences because they interfere with the binding between the anti-TNF α drug and TNF α , thereby inducing a loss of efficacy.
- 10 [0073] In additional exemplary embodiments, the assays of the present invention find utility in predicting and/or determining the cross-reactivity of neutralizing anti-drug antibodies in a subject's sample with alternative biological drugs such as other anti-TNF drugs. For illustration purposes only, if the sample contains neutralizing ADA to one anti-TNF α drug, these neutralizing ADA will likely cross-react and be neutralizing to other anti-TNF α drugs, such that the recommended treatment adjustment for the subject would be to
15 switch to a drug with a different mechanism of action (*e.g.*, a non-anti-TNF agent). However, if the sample contains non-neutralizing ADA to one anti-TNF α drug, the recommended treatment adjustment for the subject could be to switch to another anti-TNF α drug.
- [0074] Accordingly, the present invention addresses and overcomes current limitations
20 associated with the administration of anti-TNF α drugs, such as infliximab and adalimumab, in part, by providing information useful for guiding treatment decisions for those subjects receiving anti-TNF α drug therapy. The methods of the present invention are particularly useful for monitoring those subjects receiving an anti-TNF α drug to detect or measure the formation and/or development of neutralizing ADA (*e.g.*, over time during a course of anti-TNF α drug therapy) and are also useful to detect or measure a change in (*e.g.*, increase) the
25 amount, percent, or ratio of neutralizing ADA compared to non-neutralizing ADA over time while a subject is on anti-TNF α drug therapy.
- [0075] As such, the present invention provides methods for determining when and/or how
30 (1) to adjust or modify (*e.g.*, increase or decrease) the subsequent dose of an anti-TNF α drug to optimize therapeutic efficacy and/or to reduce toxicity in view of the presence, level, or percent of neutralizing ADA, (2) to combine an anti-TNF α drug (*e.g.*, at an initial, increased, decreased, or same dose) with one or more immunosuppressive agents such as methotrexate (MTX) or azathioprine (AZA) in view of the presence, level, or percent of neutralizing ADA, and/or (3) to change the current course of therapy (*e.g.*, switch to a different anti-TNF α drug

or to a drug that targets a different mechanism) in view of the presence, level, or percent of neutralizing ADA. Such methods are useful for ensuring that subjects receiving anti-TNF α drugs are getting the right dose, that they are not developing an immune response against the drug, and that they should be switched to a different drug due to failure with the initial drug
5 (e.g., development of cross-reactive neutralizing ADA against the initial anti-TNF α drug).

II. Definitions

[0076] As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

[0077] The terms “biologic” or “biologic agent” or “biological drug” as used herein
10 encompass products and substances produced from or extracted from a biological system (e.g., a living organism). Non-limiting examples of biologics include antibodies, antibody fragments, proteins, polypeptides, peptides, fusion proteins (e.g., Ig fusion proteins or Fc fusion proteins), multivalent binding proteins (e.g., DVD Ig), antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and
15 combinations thereof.

[0078] The term “biologic binding moiety” includes any molecule, agent, or substance that (e.g., specifically) binds to or interacts with a biologic. In certain instances, a neutralizing form of the autoantibody interferes with the binding between the biologic binding moiety and the biologic. In certain other instances, a non-neutralizing form of the autoantibody does not
20 interfere with the binding between the biologic binding moiety and the biologic. As one non-limiting example, the biologic binding moiety comprises TNF α when the biologic comprises an anti-TNF α drug. As another non-limiting example, the biologic binding moiety comprises an interleukin receptor (e.g., a soluble extracellular fragment of an interleukin receptor) when the biologic comprises an interleukin such as IL-2.

[0079] The terms “anti-TNF α drug” or “TNF α inhibitor” as used herein are intended to encompass agents including proteins, antibodies, antibody fragments, fusion proteins (e.g., Ig fusion proteins or Fc fusion proteins), multivalent binding proteins (e.g., DVD Ig), small molecule TNF α antagonists and similar naturally- or nonnaturally-occurring molecules, and/or recombinant and/or engineered forms thereof, that, directly or indirectly, inhibit TNF α
25 activity, such as by inhibiting interaction of TNF α with a cell surface receptor for TNF α , inhibiting TNF α protein production, inhibiting TNF α gene expression, inhibiting TNF α secretion from cells, inhibiting TNF α receptor signaling or any other means resulting in decreased TNF α activity in a subject. The term “anti-TNF α drug” or “TNF α inhibitor”
30

preferably includes agents which interfere with TNF α activity. Examples of anti-TNF α drugs include, without limitation, infliximab (REMICADE™, Johnson and Johnson), human anti-TNF monoclonal antibody adalimumab (D2E7/HUMIRA™, Abbott Laboratories), etanercept (ENBREL™, Amgen), certolizumab pegol (CIMZIA®, UCB, Inc.), golimumab (SIMPONI®; 5 CANTO 148), CDP 571 (Celltech), CDP 870 (Celltech), as well as other compounds which inhibit TNF α activity, such that when administered to a subject suffering from or at risk of suffering from a disorder in which TNF α activity is detrimental (e.g., RA), the disorder is treated.

[0080] The term “TNF α ” is intended to include a human cytokine that exists as a 17 kDa secreted form and a 26 kDa membrane associated form, the biologically active form of which 10 is composed of a trimer of noncovalently bound 17 kDa molecules. The structure of TNF α is described further in, for example, Jones *et al.*, *Nature*, 338:225-228 (1989). The term TNF α is intended to include human TNF α , a recombinant human TNF α (rhTNF- α), or TNF α that is at least about 80% identity to the human TNF α protein. Human TNF α consists of a 35 amino 15 acid (aa) cytoplasmic domain, a 21 aa transmembrane segment, and a 177 aa extracellular domain (ECD) (Pennica, D. *et al.* (1984) *Nature* 312:724). Within the ECD, human TNF α shares 97% aa sequence identity with rhesus TNF α , and 71% to 92% aa sequence identity with bovine, canine, cotton rat, equine, feline, mouse, porcine, and rat TNF α . TNF α can be prepared by standard recombinant expression methods or purchased commercially (R & D 20 Systems, Catalog No. 210-TA, Minneapolis, Minn.).

[0081] In certain embodiments, “TNF α ” is an “antigen,” which includes a molecule or a portion of the molecule capable of being bound by an anti-TNF- α drug. TNF α can have one or more than one epitope. In certain instances, TNF α will react, in a highly selective manner, 25 with an anti-TNF α antibody. Preferred antigens that bind antibodies, fragments, and regions of anti-TNF α antibodies include at least 5 amino acids of human TNF α . In certain instances, TNF α is a sufficient length having an epitope of TNF α that is capable of binding anti-TNF α antibodies, fragments, and regions thereof.

[0082] The term “size exclusion chromatography” or “SEC” includes a chromatographic method in which molecules in solution are separated based on their size and/or hydrodynamic 30 volume. It is applied to large molecules or macromolecular complexes such as proteins and their conjugates. Typically, when an aqueous solution is used to transport the sample through the column, the technique is known as gel filtration chromatography.

[0083] The terms “complex,” “immuno-complex,” “conjugate,” and “immunoconjugate” include, but are not limited to, TNF α bound (*e.g.*, by non-covalent means) to an anti-TNF α drug, an anti-TNF α drug bound (*e.g.*, by non-covalent means) to an autoantibody against the anti-TNF α drug (*e.g.*, a neutralizing or non-neutralizing anti-drug antibody), and an anti-TNF α drug bound (*e.g.*, by non-covalent means) to both TNF α and an autoantibody against the anti-TNF α drug (*e.g.*, a neutralizing or non-neutralizing anti-drug antibody).

[0084] As used herein, an entity that is modified by the term “labeled” includes any entity, molecule, protein, enzyme, antibody, antibody fragment, cytokine, or related species that is conjugated with another molecule or chemical entity that is empirically detectable. Chemical species suitable as labels for labeled-entities include, but are not limited to, fluorescent dyes, *e.g.* Alexa Fluor[®] dyes such as Alexa Fluor[®] 647, quantum dots, optical dyes, luminescent dyes, and radionuclides, *e.g.* ¹²⁵I.

[0085] The phrase “fluorescence label detection” includes a means for detecting a fluorescent label. Means for detection include, but are not limited to, a spectrometer, a fluorimeter, a photometer, and a detection device commonly incorporated with a chromatography instrument such as, but not limited to, size exclusion-high performance liquid chromatography, such as, but not limited to, an Agilent-1200 HPLC System.

[0086] The phrase “optimize therapy” includes optimizing the dose (*e.g.*, the effective amount or level) and/or the type of a particular therapy. For example, optimizing the dose of an anti-TNF α drug includes increasing or decreasing the amount of the anti-TNF α drug subsequently administered to a subject. In certain instances, optimizing the type of an anti-TNF α drug includes changing the administered anti-TNF α drug from one drug to a different drug (*e.g.*, a different anti-TNF α drug or a drug that targets a different mechanism). In other instances, optimizing therapy includes co-administering a dose of an anti-TNF α drug (*e.g.*, at an increased, decreased, or same dose as the previous dose) in combination with one or more immunosuppressive drugs.

[0087] The term “co-administer” includes to administer more than one active agent, such that the duration of physiological effect of one active agent overlaps with the physiological effect of a second active agent.

[0088] The term “subject,” “patient,” or “individual” typically includes humans, but also includes other animals such as, *e.g.*, other primates, rodents, canines, felines, equines, ovines, porcines, and the like.

[0089] The term “course of therapy” includes any therapeutic approach taken to relieve or prevent one or more symptoms associated with a disease or disorder. The term encompasses administering any compound, drug, procedure, and/or regimen useful for improving the health of an individual with a disease or disorder and includes any of the therapeutic agents described herein. As a non-limiting example, the course of therapy or the dose of the current course of therapy can be changed (*e.g.*, increased or decreased) based upon the presence or concentration level of TNF α , anti-TNF α drug, and/or anti-drug antibody (*e.g.*, the presence, level, or percent of neutralizing and/or non-neutralizing anti-drug antibody determined using the methods of the invention).

[0090] The term “immunosuppressive drug” or “immunosuppressive agent” includes any substance capable of producing an immunosuppressive effect, *e.g.*, the prevention or diminution of the immune response, as by irradiation or by administration of drugs such as anti-metabolites, anti-lymphocyte sera, antibodies, *etc.* Examples of immunosuppressive drugs include, without limitation, thiopurine drugs such as azathioprine (AZA) and metabolites thereof; anti-metabolites such as methotrexate (MTX); sirolimus (rapamycin); temsirolimus; everolimus; tacrolimus (FK-506); FK-778; anti-lymphocyte globulin antibodies, anti-thymocyte globulin antibodies, anti-CD3 antibodies, anti-CD4 antibodies, and antibody-toxin conjugates; cyclosporine; mycophenolate; mizoribine monophosphate; scoparone; glatiramer acetate; metabolites thereof; pharmaceutically acceptable salts thereof; derivatives thereof; prodrugs thereof; and combinations thereof.

[0091] The term “thiopurine drug” includes azathioprine (AZA), 6-mercaptopurine (6-MP), or any metabolite thereof that has therapeutic efficacy and includes, without limitation, 6-thioguanine (6-TG), 6-methylmercaptopurine riboside, 6-thioinosine nucleotides (*e.g.*, 6-thioinosine monophosphate, 6-thioinosine diphosphate, 6-thioinosine triphosphate), 6-thioguanine nucleotides (*e.g.*, 6-thioguanosine monophosphate, 6-thioguanosine diphosphate, 6-thioguanosine triphosphate), 6-thioxanthosine nucleotides (*e.g.*, 6-thioxanthosine monophosphate, 6-thioxanthosine diphosphate, 6-thioxanthosine triphosphate), derivatives thereof, analogues thereof, and combinations thereof.

[0092] The term “sample” includes any biological specimen obtained from an individual. Samples include, without limitation, whole blood, plasma, serum, red blood cells, white blood cells (*e.g.*, peripheral blood mononuclear cells (PBMC), polymorphonuclear (PMN) cells), ductal lavage fluid, nipple aspirate, lymph (*e.g.*, disseminated tumor cells of the lymph node), bone marrow aspirate, saliva, urine, stool (*i.e.*, feces), sputum, bronchial lavage fluid, tears, fine needle aspirate (*e.g.*, harvested by random periareolar fine needle aspiration), any

other bodily fluid, a tissue sample such as a biopsy of a site of inflammation (*e.g.*, needle biopsy), cellular extracts thereof, and an immunoglobulin enriched fraction derived from one or more of these bodily fluids or tissues. In some embodiments, the sample is whole blood, a fractional component thereof such as plasma, serum, or a cell pellet, or an immunoglobulin enriched fraction thereof. One skilled in the art will appreciate that samples such as serum samples can be diluted prior to the analysis. In certain embodiments, the sample is obtained by isolating PBMCs and/or PMN cells using any technique known in the art. In certain other embodiments, the sample is a tissue biopsy such as, *e.g.*, from a site of inflammation such as a portion of the gastrointestinal tract or synovial tissue.

10 [0093] The steps of the methods of the present invention do not necessarily have to be performed in the particular order in which they are presented. A person of ordinary skill in the art would understand that other orderings of the steps of the methods of the invention are encompassed within the scope of the present invention.

15 [0094] Brackets, “[]” indicate that the species within the brackets are referred to by their concentration.

III. Description of the Embodiments

[0095] The present invention provides assays for detecting and measuring the presence or level of neutralizing and non-neutralizing autoantibodies to biologics such as anti-TNF α drug therapeutics in a sample. The present invention is useful for monitoring the formation of neutralizing and/or non-neutralizing anti-drug antibodies over time while a subject is on biologic therapy (*e.g.*, anti-TNF α drug therapy). The present invention is also useful for predicting and/or determining the cross-reactivity of neutralizing anti-drug antibodies in a subject's sample with alternative biologic therapies (*e.g.*, alternative anti-TNF α therapies). As such, the present invention provides information for guiding treatment decisions for those subjects receiving therapy with a biologic agent and improves the accuracy of optimizing therapy, reducing toxicity, and/or monitoring the efficacy of therapeutic treatment to biologic therapy.

25 [0096] In one aspect, the present invention provides a method for detecting the presence of a neutralizing and/or non-neutralizing form of an autoantibody to a biologic in a sample, the method comprising:

- 30 (a) contacting the sample with a labeled biologic and a labeled biologic binding moiety to form:

- (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
- 5 (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic, the labeled biologic binding moiety, and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- (b) 10 subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled biologic binding moiety, free labeled biologic, and/or a complex of labeled biologic and labeled biologic binding moiety;
- (c) 15 measuring the level of free labeled biologic binding moiety after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled biologic binding moiety peak following size exclusion chromatography (SEC)); and
- (d) 20 comparing the level of the free labeled biologic binding moiety measured in step (c) to the level of free labeled biologic binding moiety in a control sample (*e.g.*, by measuring the AUC of the free labeled biologic binding moiety peak following SEC of a reference sample containing only free labeled biologic binding moiety), thereby detecting the presence of a neutralizing and/or non-neutralizing form of the autoantibody.

[0097] In some embodiments, a neutralizing form of the autoantibody interferes with the binding between the biologic and biologic binding moiety. In other embodiments, a non-
 25 neutralizing form of the autoantibody does not interfere with the binding between the biologic and biologic binding moiety.

[0098] In some instances, free labeled biologic binding moiety consists of labeled biologic binding moiety that is substantially free of bound biologic (*e.g.*, labeled and/or unlabeled biologic).

30 [0099] In certain embodiments, a neutralizing form of the autoantibody is detected when the level of the free labeled biologic binding moiety measured in step (c) is the same or substantially the same as the level of the free labeled biologic binding moiety in the control sample. In certain other embodiments, a non-neutralizing form of the autoantibody is detected when the level of the free labeled biologic binding moiety measured in step (c) is

decreased (*e.g.*, substantially decreased) or absent (*e.g.*, undetectable) compared to the level of the free labeled biologic binding moiety in the control sample.

[0100] In particular embodiments, the level of the free labeled biologic binding moiety measured in step (c) is considered to be substantially the same as the level of the free labeled biologic binding moiety in the control sample when it is at least about 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% the level of the free labeled biologic binding moiety measured in the control sample. In particular embodiments, the level of the free labeled biologic binding moiety measured in step (c) is considered to be substantially decreased compared to the level of the free labeled biologic binding moiety in the control sample when it is at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% less than the level of the free labeled biologic binding moiety measured in the control sample.

[0101] In some embodiments, the level of free labeled biologic binding moiety is measured by integrating the area under the curve (AUC) of the free labeled biologic binding moiety peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC).

[0102] In some embodiments, the biologic includes antibodies (*e.g.*, anti-TNF α monoclonal antibodies), antibody fragments, proteins (*e.g.*, cytokines such as interleukins), polypeptides, peptides, fusion proteins, multivalent binding proteins, antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and combinations thereof.

[0103] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving biologic therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis), an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)), or cancer.

[0104] In certain embodiments, the sample has or is suspected of having an autoantibody to the biologic. In other embodiments, the biologic autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0105] In another aspect, the present invention provides a method for measuring the level or percent of a neutralizing form of an autoantibody to a biologic in a sample, the method comprising:

- (a) contacting the sample with a labeled biologic and a labeled biologic binding moiety to form:
- (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
- (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic, the labeled biologic binding moiety, and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled biologic binding moiety, free labeled biologic, and/or a complex of labeled biologic and labeled biologic binding moiety;
- (c) measuring the level of free labeled biologic binding moiety after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled biologic binding moiety peak following size exclusion chromatography (SEC)); and
- (d) comparing the level of free labeled biologic binding moiety measured in step (c) to a normalized level or percent of free labeled biologic binding moiety in a control sample (*e.g.*, by measuring and normalizing the AUC of the free labeled biologic binding moiety peak following SEC of a reference sample containing only free labeled biologic binding moiety to calculate the level or percent of free labeled biologic binding moiety), wherein the normalized level or percent of the free labeled biologic binding moiety in the control sample corresponds to the level or percent of a neutralizing form of the autoantibody.

[0106] In some embodiments, the difference between the normalized level or percent of the free labeled biologic binding moiety in the control sample and the level of free labeled biologic binding moiety measured in step (c) corresponds to the level or percent of a non-neutralizing form of the autoantibody.

[0107] In some instances, free labeled biologic binding moiety consists of labeled biologic binding moiety that is substantially free of bound biologic (*e.g.*, labeled and/or unlabeled biologic).

- [0108] In particular embodiments, the level or percent of the free labeled biologic binding moiety in a control sample is normalized by measuring the peak area (*e.g.*, by measuring the AUC) of a complex formed between the labeled biologic and the labeled biologic binding moiety (*e.g.*, “labeled complex”), and then subtracting the measured peak area of the labeled complex from the peak area of the free labeled biologic binding moiety (*e.g.*, by measuring the AUC of the free labeled biologic binding moiety peak).
- [0109] In certain embodiments, the level of the free labeled biologic binding moiety is measured by integrating the area under the curve (AUC) of the free labeled biologic binding moiety peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC). In other embodiments, the level of a complex formed between the labeled biologic and labeled biologic binding moiety is measured by integrating the AUC of the free labeled biologic binding moiety peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC).
- [0110] In certain embodiments, a subpopulation of the autoantibody to a biologic (*e.g.*, ADA) is a neutralizing form of the autoantibody (*e.g.*, NAb). In some embodiments, the total level of an autoantibody to a biologic in a sample can be calculated by adding the levels of both neutralizing and non-neutralizing forms of the autoantibody measured in accordance with the methods of the invention.
- [0111] In some embodiments, the level of the free labeled biologic binding moiety measured in step (c) is further compared to a negative control, a positive control, or a combination thereof. In further embodiments, the percent of the neutralizing form of the autoantibody (*e.g.*, NAb) determined in step (d) is compared to a cutoff value or reference range established from a healthy control (*e.g.*, normal human serum). In some embodiments, the cutoff value or reference range is expressed as a threshold percent of NAb that the sample must have in order to be considered positive for NAb. In such embodiments, the sample is positive for NAb when the percent of NAb determined in step (d) is greater than or equal to the cutoff value or reference range established from the healthy control. In other embodiments, the sample is negative for NAb when the percent of NAb determined in step (d) is less than the cutoff value or reference range established from the healthy control. Non-limiting examples of cutoff values or reference ranges include, *e.g.*, at least about 0.25%, 0.50%, 0.75%, 1.00%, 1.50%, 2.00%, 2.50%, 2.60%, 2.70%, 2.80%, 2.90%, 3.00%, 3.01%, 3.02%, 3.03%, 3.04%, 3.05%, 3.06%, 3.07%, 3.08%, 3.09%, 3.10%, 3.20%, 3.30%, 3.40%,

3.50%, 4.00%, 4.50%, 5.00%, 5.50%, 6.00%, 6.50%, 7.00%, 7.50%, 8.00%, 8.50%, 9.00%, 9.50%, 10.00% NAb, or any range therein.

[0112] In some embodiments, all of the autoantibodies to the biologic are neutralizing antibodies and the sample is defined as having 100% neutralizing anti-drug antibodies (NAb) and/or 0% non-neutralizing anti-drug antibodies (non-NAb). In these embodiments, the level of the free labeled biologic binding moiety measured in step (c) is generally the same as the level of the free labeled biologic binding moiety in the control sample, and the autoantibodies are predicted to completely block or interfere with the binding between the biologic and the biologic binding moiety.

[0113] In other embodiments, none of the autoantibodies to the biologic are neutralizing antibodies and the sample is defined as having 100% non-NAb and/or 0% NAb. In these embodiments, the level of the free labeled biologic binding moiety measured in step (c) is generally absent (*e.g.*, undetectable) compared to the level of the free labeled biologic binding moiety in the control sample, and the autoantibodies are predicted to not completely block or interfere with the binding between the biologic and the biologic binding moiety.

[0114] In further embodiments, when both neutralizing and non-neutralizing forms of the autoantibody are present in a sample, the percent of each species can be expressed on their own (*e.g.*, 50% NAb or 50% non-NAb is defined as an equal proportion of NAb and non-NAb in a sample) or as a ratio. In certain instances, the ratio is calculated by dividing the percent of NAb by the percent of non-NAb, or vice versa. In other instances, the ratio is calculated by dividing the level of NAb by the level of non-NAb, or vice versa.

[0115] In some embodiments, the biologic includes antibodies (*e.g.*, anti-TNF α monoclonal antibodies), antibody fragments, proteins (*e.g.*, cytokines such as interleukins), polypeptides, peptides, fusion proteins, multivalent binding proteins, antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and combinations thereof.

[0116] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving biologic therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis), an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)), or cancer.

[0117] In certain embodiments, the sample has or is suspected of having an autoantibody to the biologic. In other embodiments, the biologic autoantibody includes, but is not limited to,

human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0118] In yet another aspect, the present invention provides a method for measuring the percent or the level of a neutralizing form of an autoantibody to a biologic in a sample, the method comprising:

- (a) contacting the sample with a labeled biologic and a labeled biologic binding moiety to form:
 - (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
 - (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled biologic, the labeled biologic binding moiety, and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled biologic binding moiety, free labeled biologic, and/or a complex of labeled biologic and labeled biologic binding moiety;
- (c) measuring the level of free labeled biologic binding moiety after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled biologic binding moiety peak following size exclusion chromatography (SEC)); and
- (d) comparing the level of free labeled biologic binding moiety measured in step (c) to the level of free labeled biologic binding moiety in a control sample (*e.g.*, by measuring the AUC of the free labeled biologic binding moiety peak following SEC of a reference sample), thereby measuring the percent or level of a neutralizing form of the autoantibody.

[0119] In some embodiments, the area under the curve (AUC) of the free labeled biologic binding moiety peak is calculated for the sample (*e.g.*, from a patient) and the control sample. In certain embodiments, the labeled biologic binds the labeled biologic binding moiety and reduces the assay signal (*e.g.*, reduces the AUC of the free labeled biologic binding moiety peak of the sample when compared to the AUC of the free labeled biologic binding moiety peak of the control sample). In other embodiments, the neutralizing form of the autoantibody

(NAb) neutralizes the labeled biologic and recovers the assay signal (*e.g.*, restores or recovers the AUC of the free labeled biologic binding moiety peak of the sample to a level comparable to that of the AUC of the free labeled biologic binding moiety peak of the control sample). In some embodiments, the NAb activity is directly proportional to the measured assay signal. In particular embodiments, the percent NAb in the sample is equivalent to the percent recovery of the assay signal. In certain instances, the percent NAb is calculated as a ratio of the AUC of the free labeled biologic binding moiety peak of the sample to the AUC of the free labeled biologic binding moiety peak of the control sample. As a non-limiting example, the percent NAb is calculated according to the following formula: % Recovery = (Free labeled biologic binding moiety_{AUC} of the Sample – BKGD)/(Free labeled biologic binding moiety_{AUC} of the Control Sample – BKGD)*100.

[0120] In some instances, free labeled biologic binding moiety consists of labeled biologic binding moiety that is substantially free of bound biologic (*e.g.*, labeled and/or unlabeled biologic).

[0121] In certain embodiments, the level of the free labeled biologic binding moiety is measured by integrating the area under the curve (AUC) of the free labeled biologic binding moiety peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC). In other embodiments, the level of a complex formed between the labeled biologic and labeled biologic binding moiety is measured by integrating the AUC of the free labeled biologic binding moiety peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC).

[0122] In certain embodiments, a subpopulation of the autoantibody to a biologic (*e.g.*, ADA) is a neutralizing form of the autoantibody (*e.g.*, NAb). In some embodiments, the total level of an autoantibody to a biologic in a sample can be calculated by adding the levels of both neutralizing and non-neutralizing forms of the autoantibody measured in accordance with the methods of the invention. In particular embodiments, the neutralizing form of the autoantibody is present in a population of autoantibodies comprising both neutralizing autoantibodies and non-neutralizing autoantibodies.

[0123] In some embodiments, the level of the free labeled biologic binding moiety measured in step (c) is further compared to a negative control, a positive control, or a combination thereof. In further embodiments, the percent of the neutralizing form of the autoantibody (*e.g.*, NAb) determined in step (d) is compared to a cutoff value or reference

range that is established, *e.g.*, from samples with low positivity for the autoantibody. In some embodiments, the cutoff value or reference range is expressed as a threshold percent of NAb that the sample must have in order to be considered positive for NAb. In such embodiments, the sample is positive for NAb when the percent of NAb determined in step (d) is greater than
5 (or equal to) the established cutoff value or reference range. In other embodiments, the sample is negative for NAb when the percent of NAb determined in step (d) is less than (or equal to) the established cutoff value or reference range. Non-limiting examples of cutoff values or reference ranges include, *e.g.*, at least about 0.10%, 0.20%, 0.30%, 0.40%, 0.50%, 0.60%, 0.70%, 0.80%, 0.90%, 1.00%, 1.10%, 1.15%, 1.20%, 1.21%, 1.22%, 1.23%, 1.24%,
10 1.25%, 1.26%, 1.27%, 1.28%, 1.29%, 1.30%, 1.40%, 1.50%, 2.00%, 2.50%, 2.60%, 2.70%, 2.80%, 2.90%, 3.00%, 3.01%, 3.02%, 3.03%, 3.04%, 3.05%, 3.06%, 3.07%, 3.08%, 3.09%, 3.10%, 3.20%, 3.30%, 3.40%, 3.50%, 4.00%, 4.50%, 5.00%, 5.50%, 6.00%, 6.50%, 7.00%, 7.50%, 8.00%, 8.50%, 9.00%, 9.50%, 10.00% NAb, or any range therein.

[0124] In some embodiments, all of the autoantibodies to the biologic are neutralizing
15 antibodies and the sample is defined as having 100% neutralizing anti-drug antibodies (NAb) and/or 0% non-neutralizing anti-drug antibodies (non-NAb). In these embodiments, the level of the free labeled biologic binding moiety measured in step (c) is generally the same as the level of the free labeled biologic binding moiety in the control sample, and the autoantibodies are predicted to completely block or interfere with the binding between the biologic and the
20 biologic binding moiety.

[0125] In other embodiments, none of the autoantibodies to the biologic are neutralizing
antibodies and the sample is defined as having 100% non-NAb and/or 0% NAb. In these
embodiments, the level of the free labeled biologic binding moiety measured in step (c) is
generally absent (*e.g.*, undetectable) compared to the level of the free labeled biologic
25 binding moiety in the control sample, and the autoantibodies are predicted to not completely
block or interfere with the binding between the biologic and the biologic binding moiety.

[0126] In further embodiments, when both neutralizing and non-neutralizing forms of the
autoantibody are present in a sample, the percent of each species can be expressed on their
own (*e.g.*, 50% NAb or 50% non-NAb is defined as an equal proportion of NAb and non-
30 NAb in a sample) or as a ratio. In certain instances, the ratio is calculated by dividing the
percent of NAb by the percent of non-NAb, or vice versa. In other instances, the ratio is
calculated by dividing the level of NAb by the level of non-NAb, or vice versa.

[0127] In some embodiments, the biologic includes antibodies (*e.g.*, anti-TNF α monoclonal antibodies), antibody fragments, proteins (*e.g.*, cytokines such as interleukins), polypeptides, peptides, fusion proteins, multivalent binding proteins, antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and combinations thereof.

[0128] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving biologic therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis), an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)), or cancer.

[0129] In certain embodiments, the sample has or is suspected of having an autoantibody to the biologic. In other embodiments, the biologic autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0130] In another aspect, the present invention provides a method for determining whether a neutralizing form of an autoantibody to a first biologic is cross-reactive with a second (*i.e.*, different) biologic, the method comprising:

- (a) detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample in accordance with an assay described herein to determine whether the sample is positive or negative for the neutralizing form of the autoantibody; and

if the sample is positive for the neutralizing form of the autoantibody, then:

- (b) contacting the sample with a labeled second biologic to form a labeled complex of the labeled second biologic and the neutralizing form of the autoantibody (*i.e.*, wherein the components of the labeled complex are not covalently attached to each other);
- (c) subjecting the labeled complex to size exclusion chromatography to separate the labeled complex (*e.g.*, from free labeled second biologic); and
- (d) detecting the labeled complex, thereby determining whether a neutralizing form of an autoantibody to a first biologic is cross-reactive with a second biologic.

[0131] In certain embodiments, the presence of the labeled complex is an indication that the neutralizing autoantibody against the first biologic is cross-reactive with the second

biologic, *i.e.*, the neutralizing autoantibody will inhibit the activity of both the first and second biological drugs.

[0132] In certain other embodiments, the absence of the labeled complex is an indication that the neutralizing autoantibody against the first biologic is not cross-reactive with the second biologic, *i.e.*, the neutralizing autoantibody will not inhibit the activity of the second biological drug.

[0133] In some embodiments, the first and second biologics are independently selected from the group consisting of antibodies (*e.g.*, anti-TNF α monoclonal antibodies), antibody fragments, proteins (*e.g.*, cytokines such as interleukins), polypeptides, peptides, fusion proteins, multivalent binding proteins, antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and combinations thereof.

[0134] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving biologic therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis), an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)), or cancer.

[0135] In certain embodiments, the sample has or is suspected of having an autoantibody to the biologic. In other embodiments, the biologic autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0136] In certain aspects, the assay methods of the present invention further comprise an acid dissociation step comprising contacting a sample with an acid prior to, during, and/or after contacting the sample with a labeled biologic and a labeled biologic binding moiety.

[0137] In certain other aspects, the assay methods of the present invention comprise detecting the presence or level of one or more isotypes of a neutralizing and/or non-neutralizing form of an autoantibody to a biologic in a sample.

[0138] In one particular aspect, the present invention provides a method for detecting the presence of a neutralizing and/or non-neutralizing form of an autoantibody to an anti-TNF α drug in a sample, the method comprising:

- (a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to form:

- (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or
- 5 (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug, the labeled TNF α , and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- (b) 10 subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled TNF α , free labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and labeled TNF α ;
- (c) measuring the level of free labeled TNF α after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled TNF α peak following size exclusion chromatography (SEC)); and
- 15 (d) comparing the level of the free labeled TNF α measured in step (c) to the level of free labeled TNF α in a control sample (*e.g.*, by measuring the AUC of the free labeled TNF α peak following SEC of a reference sample containing only free labeled TNF α), thereby detecting the presence of a neutralizing and/or
- 20 non-neutralizing form of the autoantibody.

[0139] In some embodiments, a neutralizing form of the autoantibody interferes with the binding between the anti-TNF α drug and TNF α . In other embodiments, a non-neutralizing form of the autoantibody does not interfere with the binding between the anti-TNF α drug and TNF α .

25 [0140] In some instances, free labeled TNF α consists of labeled TNF α that is substantially free of bound anti-TNF α drug (*e.g.*, labeled and/or unlabeled anti-TNF α drug).

[0141] In certain embodiments, a neutralizing form of the autoantibody is detected when the level of the free labeled TNF α measured in step (c) is the same or substantially the same as the level of the free labeled TNF α in the control sample. In certain other embodiments, a

30 non-neutralizing form of the autoantibody is detected when the level of the free labeled TNF α measured in step (c) is decreased (*e.g.*, substantially decreased) or absent (*e.g.*, undetectable) compared to the level of the free labeled TNF α in the control sample.

[0142] In particular embodiments, the level of the free labeled TNF α measured in step (c) is considered to be substantially the same as the level of the free labeled TNF α in the control

sample when it is at least about 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% the level of the free labeled TNF α measured in the control sample. In particular embodiments, the level of the free labeled TNF α measured in step (c) is considered to be substantially decreased compared to the level of the free labeled TNF α in the control sample when it is at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95% less than the level of the free labeled TNF α measured in the control sample.

[0143] In certain embodiments, the level of free labeled TNF α is measured by integrating the area under the curve (AUC) of the free labeled TNF α peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC).

[0144] In some embodiments, the anti-TNF α drug is selected from the group consisting of REMICADE™ (infliximab), ENBREL™ (etanercept), HUMIRA™ (adalimumab), CIMZIA® (certolizumab pegol), SIMPONI® (golimumab; CNTO 148), and combinations thereof.

[0145] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving anti-TNF α drug therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a TNF α -mediated disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis) or an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)).

[0146] In certain embodiments, the sample has or is suspected of having an autoantibody to the anti-TNF α drug. In other embodiments, the anti-TNF α drug autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0147] In another particular aspect, the present invention provides a method for measuring the level or percent of a neutralizing form of an autoantibody to an anti-TNF α drug in a sample, the method comprising:

(a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to form:

(i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or

(ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug, the labeled TNF α , and the autoantibody (*i.e.*,

wherein the components of the second labeled complex are not covalently attached to each other);

- 5 (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled TNF α , free labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and labeled TNF α ;
- (c) measuring the level of free labeled TNF α after size exclusion chromatography (*e.g.*, by measuring the area under the curve (AUC) of the free labeled TNF α peak following size exclusion chromatography (SEC)); and
- 10 (d) comparing the level of free labeled TNF α measured in step (c) to a normalized level or percent of free labeled TNF α in a control sample (*e.g.*, by measuring and normalizing the AUC of the free labeled TNF α peak following SEC of a reference sample containing only free labeled TNF α to calculate the level or percent of free labeled TNF α), wherein the normalized level or percent of the
- 15 free labeled TNF α in the control sample corresponds to the level or percent of a neutralizing form of the autoantibody.

[0148] In some embodiments, the difference between the normalized level or percent of the free labeled TNF α in the control sample and the level of free labeled TNF α measured in step (c) corresponds to the level or percent of a non-neutralizing form of the autoantibody.

- 20 [0149] In some instances, free labeled TNF α consists of labeled TNF α that is substantially free of bound anti-TNF α drug (*e.g.*, labeled and/or unlabeled anti-TNF α drug).

- [0150] In particular embodiments, the level or percent of the free labeled TNF α in a control sample is normalized by measuring the peak area (*e.g.*, by measuring the AUC) of a complex formed between the labeled anti-TNF α drug and labeled TNF α (*e.g.*, "labeled complex"), and
- 25 then subtracting the measured peak area of the labeled complex from the peak area of the free labeled TNF α (*e.g.*, by measuring the AUC of the free labeled TNF α peak).

- [0151] In certain embodiments, the level of free labeled TNF α is measured by integrating the area under the curve (AUC) of the free labeled TNF α peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC). In
- 30 other embodiments, the level of a complex formed between the labeled anti-TNF α drug and labeled TNF α is measured by integrating the AUC of the free labeled TNF α peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC).

[0152] In certain embodiments, a subpopulation of the autoantibody to an anti-TNF α drug (e.g., ADA) is a neutralizing form of the autoantibody (e.g., NAb). In some embodiments, the total level of an autoantibody to an anti-TNF α drug in a sample can be calculated by adding the levels of both neutralizing and non-neutralizing forms of the autoantibody measured in accordance with the methods of the invention.

[0153] In some embodiments, the level of the free labeled TNF α measured in step (c) is further compared to a negative control, a positive control, or a combination thereof. Non-limiting examples of negative controls include a mouse monoclonal anti-human IgG₁ Fc sample and/or a rabbit monoclonal anti-human IgG₁ Fc sample. Non-limiting examples of positive controls include a pooled ADA-positive patient serum sample and/or a sample of rabbit polyclonal antibodies against the F(ab')₂ fragment of an anti-TNF α drug.

[0154] In further embodiments, the percent of the neutralizing form of the autoantibody (e.g., NAb) determined in step (d) is compared to a cutoff value or reference range established from a healthy control (e.g., normal human serum). In particular embodiments, the cutoff value or reference range is expressed as a threshold percent of NAb that the sample must have in order to be considered positive for NAb. In such embodiments, the sample is positive for NAb when the percent of NAb determined in step (d) is greater than or equal to the cutoff value or reference range established from the healthy control. In other embodiments, the sample is negative for NAb when the percent of NAb determined in step (d) is less than the cutoff value or reference range established from the healthy control. Non-limiting examples of cutoff values or reference ranges include, e.g., at least about 0.25%, 0.50%, 0.75%, 1.00%, 1.50%, 2.00%, 2.50%, 2.60%, 2.70%, 2.80%, 2.90%, 3.00%, 3.01%, 3.02%, 3.03%, 3.04%, 3.05%, 3.06%, 3.07%, 3.08%, 3.09%, 3.10%, 3.20%, 3.30%, 3.40%, 3.50%, 4.00%, 4.50%, 5.00%, 5.50%, 6.00%, 6.50%, 7.00%, 7.50%, 8.00%, 8.50%, 9.00%, 9.50%, 10.00% NAb, or any range therein. In some instances, the cutoff value or reference range is about 3.00% NAb or about 3.06% NAb or between about 3.00%-3.10% NAb.

[0155] In some embodiments, all the autoantibodies to the anti-TNF α drug are neutralizing antibodies and the sample is defined as having 100% neutralizing anti-drug antibodies (NAb) and/or 0% non-neutralizing anti-drug antibodies (non-NAb). In these embodiments, the level of the free labeled TNF α measured in step (c) is generally the same as the level of the free labeled TNF α in the control sample, and the autoantibodies are predicted to completely block or interfere with the binding between the anti-TNF α drug and TNF α .

[0156] In certain other embodiments, none of the autoantibodies to the anti-TNF α drug are neutralizing antibodies and the sample is defined as having 100% non-NAb and/or 0% NAb. In these embodiments, the level of the free labeled TNF α measured in step (c) is generally absent (*e.g.*, undetectable) compared to the level of the free labeled TNF α in the control
5 sample, and the autoantibodies are predicted to not completely block or interfere with the binding between the anti-TNF α drug and TNF α .

[0157] In further embodiments, when both neutralizing and non-neutralizing forms of the autoantibody are present in a sample, the percent of each species can be expressed on their own (*e.g.*, 50% NAb or 50% non-NAb is defined as an equal proportion of NAb and non-
10 NAb in a sample) or as a ratio. In certain instances, the ratio is calculated by dividing the percent of NAb by the percent of non-NAb, or vice versa. In other instances, the ratio is calculated by dividing the level of NAb by the level of non-NAb, or vice versa.

[0158] In some embodiments, the anti-TNF α drug is selected from the group consisting of REMICADE™ (infliximab), ENBREL™ (etanercept), HUMIRA™ (adalimumab), CIMZIA®
15 (certolizumab pegol), SIMPONI® (golimumab; CNTO 148), and combinations thereof.

[0159] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving anti-TNF α drug therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a TNF α -mediated disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis) or an inflammatory disease (*e.g.*,
20 inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)).

[0160] In certain embodiments, the sample has or is suspected of having an autoantibody to the anti-TNF α drug. In other embodiments, the anti-TNF α drug autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

25 [0161] In yet another particular aspect, the present invention provides a method for measuring the percent or the level of a neutralizing form of an autoantibody to an anti-TNF α drug in a sample, the method comprising:

- (a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to form:
30 (i) a first labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug and the autoantibody (*i.e.*, wherein the components of the first labeled complex are not covalently attached to each other); and/or

- (ii) a second labeled complex (*i.e.*, immuno-complex or conjugate) of the labeled anti-TNF α drug, the labeled TNF α , and the autoantibody (*i.e.*, wherein the components of the second labeled complex are not covalently attached to each other);
- 5 (b) subjecting the first labeled complex and/or the second labeled complex to size exclusion chromatography to separate them from free (*i.e.*, unbound) labeled TNF α , free labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and labeled TNF α ;
- (c) measuring the level of free labeled TNF α after size exclusion chromatography (e.g., by measuring the area under the curve (AUC) of the free labeled TNF α peak following size exclusion chromatography (SEC)); and
- 10 (d) comparing the level of free labeled TNF α measured in step (c) to the level of free labeled TNF α in a control sample (e.g., by measuring the AUC of the free labeled TNF α following SEC of a reference sample), thereby measuring the
- 15 percent or level of a neutralizing form of the autoantibody.

[0162] In some embodiments, the area under the curve (AUC) of the free labeled TNF α peak is calculated for the sample (e.g., from a patient) and the control sample. In certain embodiments, the labeled anti-TNF α drug binds the labeled TNF α and reduces the assay signal (e.g., reduces the AUC of the free labeled TNF α peak of the sample when compared to the AUC of the free labeled TNF α peak of the control sample). In other embodiments, the neutralizing form of the autoantibody (NAb) neutralizes the labeled anti-TNF α drug and recovers the assay signal (e.g., restores or recovers the AUC of the free labeled TNF α peak of the sample to a level comparable to that of the AUC of the free labeled TNF α peak of the control sample). In some embodiments, the NAb activity is directly proportional to the measured assay signal. In particular embodiments, the percent NAb in the sample is equivalent to the percent recovery of the assay signal. In certain instances, the percent NAb is calculated as a ratio of the AUC of the free labeled TNF α peak of the sample to the AUC of the free labeled TNF α peak of the control sample. As a non-limiting example, the percent NAb is calculated according to the following formula: % Recovery = (Free labeled TNF α _{AUC} of the Sample - BKGD)/(Free labeled TNF α _{AUC} of the Control Sample - BKGD)*100.

[0163] In some instances, free labeled TNF α consists of labeled TNF α that is substantially free of bound anti-TNF α drug (e.g., labeled and/or unlabeled anti-TNF α drug).

[0164] In certain embodiments, the level of free labeled TNF α is measured by integrating the area under the curve (AUC) of the free labeled TNF α peak from a plot of signal intensity

as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC). In other embodiments, the level of a complex formed between the labeled anti-TNF α drug and labeled TNF α is measured by integrating the AUC of the free labeled TNF α peak from a plot of signal intensity as a function of elution time from the size exclusion chromatography (*e.g.*, SEC-HPLC).

[0165] In certain embodiments, a subpopulation of the autoantibody to an anti-TNF α drug (*e.g.*, ADA) is a neutralizing form of the autoantibody (*e.g.*, NAb). In some embodiments, the total level of an autoantibody to an anti-TNF α drug in a sample can be calculated by adding the levels of both neutralizing and non-neutralizing forms of the autoantibody measured in accordance with the methods of the invention. In particular embodiments, the neutralizing form of the autoantibody is present in a population of autoantibodies comprising both neutralizing autoantibodies and non-neutralizing autoantibodies.

[0166] In some embodiments, the level of the free labeled TNF α measured in step (c) is further compared to a negative control, a positive control, or a combination thereof. Non-limiting examples of negative controls include a mouse monoclonal anti-human IgG₁ Fc sample and/or a rabbit monoclonal anti-human IgG₁ Fc sample. Non-limiting examples of positive controls include a pooled ADA-positive patient serum sample and/or a sample of rabbit polyclonal antibodies against the F(ab')₂ fragment of an anti-TNF α drug.

[0167] In further embodiments, the percent of the neutralizing form of the autoantibody (*e.g.*, NAb) determined in step (d) is compared to a cutoff value or reference range that is established, *e.g.*, from samples with low positivity for the autoantibody. In particular embodiments, the cutoff value or reference range is expressed as a threshold percent of NAb that the sample must have in order to be considered positive for NAb. In such embodiments, the sample is positive for NAb when the percent of NAb determined in step (d) is greater than (or equal to) the established cutoff value or reference range. In other embodiments, the sample is negative for NAb when the percent of NAb determined in step (d) is less than (or equal to) the established cutoff value or reference range. Non-limiting examples of cutoff values or reference ranges include, *e.g.*, at least about 0.10%, 0.20%, 0.30%, 0.40%, 0.50%, 0.60%, 0.70%, 0.80%, 0.90%, 1.00%, 1.10%, 1.15%, 1.20%, 1.21%, 1.22%, 1.23%, 1.24%, 1.25%, 1.26%, 1.27%, 1.28%, 1.29%, 1.30%, 1.40%, 1.50%, 2.00%, 2.50%, 2.60%, 2.70%, 2.80%, 2.90%, 3.00%, 3.01%, 3.02%, 3.03%, 3.04%, 3.05%, 3.06%, 3.07%, 3.08%, 3.09%, 3.10%, 3.20%, 3.30%, 3.40%, 3.50%, 4.00%, 4.50%, 5.00%, 5.50%, 6.00%, 6.50%, 7.00%, 7.50%, 8.00%, 8.50%, 9.00%, 9.50%, 10.00% NAb, or any range therein. In some instances, the cutoff value or reference range is about 1.28% NAb or between about 1.25%-1.30% NAb.

[0168] In some embodiments, all the autoantibodies to the anti-TNF α drug are neutralizing antibodies and the sample is defined as having 100% neutralizing anti-drug antibodies (NAb) and/or 0% non-neutralizing anti-drug antibodies (non-NAb). In these embodiments, the level of the free labeled TNF α measured in step (c) is generally the same as the level of the free labeled TNF α in the control sample, and the autoantibodies are predicted to completely block or interfere with the binding between the anti-TNF α drug and TNF α .

[0169] In certain other embodiments, none of the autoantibodies to the anti-TNF α drug are neutralizing antibodies and the sample is defined as having 100% non-NAb and/or 0% NAb. In these embodiments, the level of the free labeled TNF α measured in step (c) is generally absent (*e.g.*, undetectable) compared to the level of the free labeled TNF α in the control sample, and the autoantibodies are predicted to not completely block or interfere with the binding between the anti-TNF α drug and TNF α .

[0170] In further embodiments, when both neutralizing and non-neutralizing forms of the autoantibody are present in a sample, the percent of each species can be expressed on their own (*e.g.*, 50% NAb or 50% non-NAb is defined as an equal proportion of NAb and non-NAb in a sample) or as a ratio. In certain instances, the ratio is calculated by dividing the percent of NAb by the percent of non-NAb, or vice versa. In other instances, the ratio is calculated by dividing the level of NAb by the level of non-NAb, or vice versa.

[0171] In some embodiments, the anti-TNF α drug is selected from the group consisting of REMICADE[™] (infliximab), ENBREL[™] (etanercept), HUMIRA[™] (adalimumab), CIMZIA[®] (certolizumab pegol), SIMPONI[®] (golimumab; CNTO 148), and combinations thereof.

[0172] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving anti-TNF α drug therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a TNF α -mediated disease or disorder such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis) or an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)).

[0173] In certain embodiments, the sample has or is suspected of having an autoantibody to the anti-TNF α drug. In other embodiments, the anti-TNF α drug autoantibody includes, but is not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0174] In still yet another particular aspect, the present invention provides a method for determining whether a neutralizing form of an autoantibody to a first anti-TNF α drug is cross-reactive with a second (*i.e.*, different) anti-TNF α drug, the method comprising:

- (a) detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample in accordance with an assay described herein to determine whether the sample is positive or negative for the neutralizing form of the autoantibody; and
- 5 if the sample is positive for the neutralizing form of the autoantibody, then:
 - (b) contacting the sample with a labeled second anti-TNF α drug to form a labeled complex of the labeled second anti-TNF α drug and the neutralizing form of the autoantibody (*i.e.*, wherein the components of the labeled complex are not covalently attached to each other);
 - 10 (c) subjecting the labeled complex to size exclusion chromatography to separate the labeled complex (*e.g.*, from free labeled second anti-TNF α drug); and
 - (d) detecting the labeled complex, thereby determining whether a neutralizing form of an autoantibody to a first anti-TNF α drug is cross-reactive with a second anti-TNF α drug.

15 [0175] In certain embodiments, the presence of the labeled complex is an indication that the neutralizing autoantibody against the first anti-TNF α drug is cross-reactive with the second anti-TNF α drug, *i.e.*, the neutralizing autoantibody will inhibit the activity of both the first and second anti-TNF α drugs.

20 [0176] In certain other embodiments, the absence of the labeled complex is an indication that the neutralizing autoantibody against the first anti-TNF α drug is not cross-reactive with the second anti-TNF α drug, *i.e.*, the neutralizing autoantibody will not inhibit the activity of the second anti-TNF α drug.

[0177] In particular embodiments, the first and second anti-TNF α drugs are independently selected from the group consisting of REMICADE™ (infliximab), ENBREL™ (etanercept),
 25 HUMIRA™ (adalimumab), CIMZIA® (certolizumab pegol), SIMPONI® (golimumab; CNTO 148), and combinations thereof.

[0178] In other embodiments, the sample is a whole blood, serum, or plasma sample, *e.g.*, from a subject receiving anti-TNF α drug therapy. In preferred embodiments, the sample is serum. In particular embodiments, the subject has a TNF α -mediated disease or disorder such
 30 as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis) or an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)).

[0179] In certain embodiments, the sample has or is suspected of having an autoantibody to the anti-TNF α drug. In other embodiments, the anti-TNF α drug autoantibody includes, but is

not limited to, human anti-chimeric antibodies (HACA), human anti-humanized antibodies (HAHA), and human anti-mouse antibodies (HAMA), as well as combinations thereof.

[0180] In certain aspects, the assay methods of the present invention further comprise an acid dissociation step comprising contacting a sample with an acid prior to, during, and/or after contacting the sample with a labeled anti-TNF α drug and a labeled TNF α .

[0181] Methods for detecting anti-drug antibodies using acid dissociation are described herein and in PCT Application No. PCT/US2012/025437, filed February 16, 2012, the disclosure of which is hereby incorporated by reference in its entirety for all purposes.

[0182] In certain other aspects, the assay methods of the present invention comprise detecting the presence or level of one or more isotypes of a neutralizing and/or non-neutralizing form of an autoantibody to an anti-TNF α drug in a sample. As a non-limiting example, the assays of the present invention can be used to determine different neutralizing and/or non-neutralizing ADA isotypes in samples from ADA-positive patients receiving an anti-TNF α drug such as REMICADE™ (infliximab) or HUMIRA™ (adalimumab). In certain embodiments, the one or more isotypes comprises a plurality of at least two, three, four, five, or more isotypes. In other embodiments, the one or more isotypes is selected from the group consisting of IgA, IgD, IgE, IgG, and IgM isotypes, subclasses thereof, and combinations thereof. In certain embodiments, each autoantibody isotype is characterized, identified, and/or detected by its retention time. In other embodiments, each autoantibody isotype is characterized, identified, and/or detected upon a signal that is generated by the proximity binding of detector moieties such as labeled anti-TNF α drug and labeled anti-Ig antibodies specific for different antibody isotypes. In certain instances, the signal comprises a fluorescent signal that can be detected by fluorescence resonance energy transfer (FRET).

[0183] Methods for detecting anti-drug antibody (ADA) isotypes are further described in PCT Publication No. WO 2012/054532, the disclosure of which is hereby incorporated by reference in its entirety for all purposes.

[0184] A biologic (e.g., anti-TNF α drug) or biologic binding moiety (e.g., TNF α) can be labeled with any of a variety of detectable group(s). In preferred embodiments, the biologic (e.g., anti-TNF α drug) and the biologic binding moiety (e.g., TNF α) comprise different labels. In certain embodiments, a biologic (e.g., anti-TNF α drug) or biologic binding moiety (e.g., TNF α) is labeled with a fluorophore or a fluorescent dye. Non-limiting examples of fluorophores or fluorescent dyes include those listed in the Molecular Probes Catalogue, which is herein incorporated by reference (*see*, R. Haugland, *The Handbook-A Guide to*

Fluorescent Probes and Labeling Technologies, 10th Edition, Molecular probes, Inc. (2005)). Such exemplary fluorophores or fluorescent dyes include, but are not limited to, Alexa Fluor[®] dyes such as Alexa Fluor[®] 350, Alexa Fluor[®] 405, Alexa Fluor[®] 430, Alexa Fluor[®] 488, Alexa Fluor[®] 514, Alexa Fluor[®] 532, Alexa Fluor[®] 546, Alexa Fluor[®] 555, Alexa Fluor[®] 568, Alexa Fluor[®] 594, Alexa Fluor[®] 610, Alexa Fluor[®] 633, Alexa Fluor[®] 635, Alexa Fluor[®] 647, Alexa Fluor[®] 660, Alexa Fluor[®] 680, Alexa Fluor[®] 700, Alexa Fluor[®] 750, and/or Alexa Fluor[®] 790, as well as other fluorophores including, but not limited to, Dansyl Chloride (DNS-Cl), 5-(iodoacetamido)fluorescein (5-IAF), fluorescein 5-isothiocyanate (FITC), tetramethylrhodamine 5- (and 6-)isothiocyanate (TRITC), 6-acryloyl-2-dimethylaminonaphthalene (acrylodan), 7-nitrobenzo-2-oxa-1,3,-diazol-4-yl chloride (NBD-Cl), ethidium bromide, Lucifer Yellow, 5-carboxyrhodamine 6G hydrochloride, Lissamine rhodamine B sulfonyl chloride, Texas Red[™] sulfonyl chloride, BODIPY[™], naphthalamine sulfonic acids (*e.g.*, 1-anilinonaphthalene-8-sulfonic acid (ANS), 6-(p-toluidinyl)naphthalene-2-sulfonic acid (TNS), and the like), Anthroyl fatty acid, DPH, Parinaric acid, TMA-DPH, Fluorenyl fatty acid, fluorescein-phosphatidylethanolamine, Texas Red-phosphatidylethanolamine, Pyrenyl-phosphatidylcholine, Fluorenyl-phosphatidylcholine, Merocyanine 540, 1-(3-sulfonatopropyl)-4-[β-[2[(di-n-butylamino)-6-naphthyl]vinyl]pyridinium betaine (Naphtyl Styryl), 3,3'-dipropylthiadicarbocyanine (diS-C₃-(5)), 4-(p-dipentyl aminostyryl)-1-methylpyridinium (di-5-ASP), Cy-3 Iodo Acetamide, Cy-5-N-Hydroxysuccinimide, Cy-7-Isothiocyanate, rhodamine 800, IR-125, Thiazole Orange, Azure B, Nile Blue, Al Phthalocyanine, Oxazine 1, 4', 6-diamidino-2-phenylindole (DAPI), Hoechst 33342, TOTO, Acridine Orange, Ethidium Homodimer, N(ethoxycarbonylmethyl)-6-methoxyquinolinium (MQAE), Fura-2, Calcium Green, Carboxy SNARF-6, BAPTA, coumarin, phytofluors, Coronene, metal-ligand complexes, IRDye[®] 700DX, IRDye[®] 700, IRDye[®] 800RS, IRDye[®] 800CW, IRDye[®] 800, Cy5, Cy5.5, Cy7, DY 676, DY680, DY682, DY780, and mixtures thereof. Additional suitable fluorophores include enzyme-cofactors; lanthanide, green fluorescent protein, yellow fluorescent protein, red fluorescent protein, or mutants and derivatives thereof. In one embodiment of the invention, the second member of the specific binding pair has a detectable group attached thereto.

[0185] Typically, the fluorescent group is a fluorophore selected from the category of dyes comprising polymethines, phthalocyanines, cyanines, xanthenes, fluorenes, rhodamines, coumarins, fluoresceins and BODIPY[™].

[0186] In one embodiment, the fluorescent group is a near-infrared (NIR) fluorophore that emits in the range of between about 650 to about 900 nm. Use of near infrared fluorescence

technology is advantageous in biological assays as it substantially eliminates or reduces background from auto fluorescence of biosubstrates. Another benefit to the near-IR fluorescent technology is that the scattered light from the excitation source is greatly reduced since the scattering intensity is proportional to the inverse fourth power of the wavelength.

5 Low background fluorescence and low scattering result in a high signal to noise ratio, which is essential for highly sensitive detection. Furthermore, the optically transparent window in the near-IR region (650 nm to 900 nm) in biological tissue makes NIR fluorescence a valuable technology for *in vivo* imaging and subcellular detection applications that require the transmission of light through biological components. Within aspects of this embodiment, the
10 fluorescent group is preferably selected from the group consisting of IRDye[®] 700DX, IRDye[®] 700, IRDye[®] 800RS, IRDye[®] 800CW, IRDye[®] 800, Alexa Fluor[®] 660, Alexa Fluor[®] 680, Alexa Fluor[®] 700, Alexa Fluor[®] 750, Alexa Fluor[®] 790, Cy5, Cy5.5, Cy7, DY 676, DY680, DY682, and DY780. In certain embodiments, the near infrared group is IRDye[®] 800CW, IRDye[®] 800, IRDye[®] 700DX, IRDye[®] 700, or Dynamic DY676.

15 [0187] Fluorescent labeling is accomplished using a chemically reactive derivative of a fluorophore. Common reactive groups include amine reactive isothiocyanate derivatives such as FITC and TRITC (derivatives of fluorescein and rhodamine), amine reactive succinimidyl esters such as NHS-fluorescein, and sulfhydryl reactive maleimide activated fluors such as fluorescein-5-maleimide, many of which are commercially available. Reaction
20 of any of these reactive dyes with a biologic (*e.g.*, anti-TNF α drug) or biologic binding moiety (*e.g.*, TNF α) results in a stable covalent bond formed between a fluorophore and a biologic (*e.g.*, anti-TNF α drug) or biologic binding moiety (*e.g.*, TNF α).

[0188] In certain instances, following a fluorescent labeling reaction, it is often necessary to remove any nonreacted fluorophore from the labeled target molecule. This is often
25 accomplished by size exclusion chromatography, taking advantage of the size difference between fluorophore and labeled protein.

[0189] Reactive fluorescent dyes are available from many sources. They can be obtained with different reactive groups for attachment to various functional groups within the target molecule. They are also available in labeling kits that contain all the components to carry out
30 a labeling reaction. In one preferred aspect, Alexa Fluor[®] 647 C2 maleimide is used from Invitrogen (Cat. No. A-20347).

[0190] Specific immunological binding of a neutralizing and/or non-neutralizing anti-drug antibody (*e.g.*, NAb and/or non-NAb) to a biologic (*e.g.*, anti-TNF α drug) and/or biologic

binding moiety (*e.g.*, TNF α) can be detected directly or indirectly. Direct labels include fluorescent or luminescent tags, metals, dyes, radionuclides, and the like, attached to the antibody. In certain instances, a biologic (*e.g.*, anti-TNF α drug) or biologic binding moiety (*e.g.*, TNF α) labeled with different radionuclides can be used for determining the presence or level of NAb and/or non-NAb in a sample. In other instances, a chemiluminescence assay using chemiluminescent biologic (*e.g.*, anti-TNF α drug) and biologic binding moiety (*e.g.*, TNF α) is suitable for sensitive, non-radioactive detection of the presence or level of NAb and/or non-NAb in a sample. In particular instances, a biologic (*e.g.*, anti-TNF α drug) and biologic binding moiety (*e.g.*, TNF α) labeled with different fluorochromes is suitable for detection of the presence or level of NAb and/or non-NAb in a sample. Examples of fluorochromes include, without limitation, Alexa Fluor[®] dyes, DAPI, fluorescein, Hoechst 33258, R-phycoerythrin, B-phycoerythrin, R-phycoerythrin, rhodamine, Texas red, and lissamine. Secondary antibodies linked to fluorochromes can be obtained commercially, *e.g.*, goat F(ab')₂ anti-human IgG-FITC is available from Tago Immunologicals (Burlingame, CA).

[0191] Indirect labels include various enzymes well-known in the art, such as horseradish peroxidase (HRP), alkaline phosphatase (AP), β -galactosidase, urease, and the like. A horseradish-peroxidase detection system can be used, for example, with the chromogenic substrate tetramethylbenzidine (TMB), which yields a soluble product in the presence of hydrogen peroxide that is detectable at 450 nm. An alkaline phosphatase detection system can be used with the chromogenic substrate p-nitrophenyl phosphate, for example, which yields a soluble product readily detectable at 405 nm. Similarly, a β -galactosidase detection system can be used with the chromogenic substrate o-nitrophenyl- β -D-galactopyranoside (ONPG), which yields a soluble product detectable at 410 nm. An urease detection system can be used with a substrate such as urea-bromocresol purple (Sigma Immunochemicals; St. Louis, MO). A useful secondary antibody linked to an enzyme can be obtained from a number of commercial sources, *e.g.*, goat F(ab')₂ anti-human IgG-alkaline phosphatase can be purchased from Jackson ImmunoResearch (West Grove, PA.).

[0192] A signal from the direct or indirect label can be analyzed, for example, using a spectrophotometer to detect color from a chromogenic substrate; a radiation counter to detect radiation such as a gamma counter for detection of ¹²⁵I; or a fluorometer to detect fluorescence in the presence of light of a certain wavelength. For detection of enzyme-linked antibodies, a quantitative analysis of NAb and/or non-NAb levels can be made using a spectrophotometer such as an EMAX Microplate Reader (Molecular Devices; Menlo Park,

CA) in accordance with the manufacturer's instructions. If desired, the assays of the present invention can be automated or performed robotically, and the signal from multiple samples can be detected simultaneously.

5 [0193] In certain embodiments, size exclusion chromatography is used. The underlying principle of SEC is that particles of different sizes will elute (filter) through a stationary phase at different rates. This results in the separation of a solution of particles based on size. Provided that all the particles are loaded simultaneously or near simultaneously, particles of the same size elute together. Each size exclusion column has a range of molecular weights that can be separated. The exclusion limit defines the molecular weight at the upper end of
10 this range and is where molecules are too large to be trapped in the stationary phase. The permeation limit defines the molecular weight at the lower end of the range of separation and is where molecules of a small enough size can penetrate into the pores of the stationary phase completely and all molecules below this molecular mass are so small that they elute as a single band.

15 [0194] In certain aspects, the eluent is collected in constant volumes, or fractions. The more similar the particles are in size, the more likely they will be in the same fraction and not detected separately. Preferably, the collected fractions are examined by spectroscopic techniques to determine the concentration of the particles eluted. Typically, the spectroscopy detection techniques useful in the present invention include, but are not limited to,
20 fluorometry, refractive index (RI), and ultraviolet (UV). In certain instances, the elution volume decreases roughly linearly with the logarithm of the molecular hydrodynamic volume (*i.e.*, heavier moieties come off first).

[0195] In a further aspect, the present invention provides a method for monitoring and/or optimizing therapy to a biologic in a subject receiving a course of therapy with the biologic,
25 the method comprising:

- (a) detecting or measuring the presence, level, or percent of a neutralizing form of an autoantibody to the biologic in accordance with the assay described herein at a plurality of time points over the course of therapy;
- (b) detecting a change in the presence, level, or percent of the neutralizing form of
30 the autoantibody over time; and
- (c) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the presence, level, or percent of the neutralizing form of the autoantibody over time.

[0196] In certain embodiments, the plurality of time points comprises at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, or more time points.

[0197] In one particular aspect, the present invention provides a method for monitoring and/or optimizing therapy to a biologic in a subject receiving a course of therapy with the

5 biologic, the method comprising:

- (a) measuring the level or percent of a neutralizing form of an autoantibody to the biologic in a first sample from the subject as described herein at time point t_0 ;
- (b) measuring the level or percent of the neutralizing form of the autoantibody in a second sample from the subject as described herein at time point t_1 ;
- 10 (c) optionally repeating step (b) with n additional samples from the subject at time points t_{n+1} , wherein n is an integer from 1 to about 25 (e.g., n is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or any range therein);
- (d) detecting a change in the level or percent of the neutralizing form of the autoantibody from time points t_0 to t_1 or from time points t_0 to t_{n+1} ; and
- 15 (e) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the level or percent of the neutralizing form of the autoantibody over time.

20 [0198] In certain other embodiments, the level or percent of the neutralizing form of the autoantibody (e.g., NAb) is measured during the course of biologic drug therapy at one or more (e.g., a plurality) of the following weeks: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 80, 90, 100, etc.

25 [0199] In some embodiments, determining a subsequent dose of the course of therapy for the subject comprises maintaining, increasing, or decreasing a subsequent dose of the course of therapy for the subject. In other embodiments, determining a different course of therapy for the subject comprises treatment with a different biologic drug. In other embodiments, determining a different course of therapy for the subject comprises treatment with the current
30 course of therapy along with another therapeutic agent. In further embodiments, determining a different course of therapy for the subject comprises changing the current course of therapy (e.g., switching to a different biologic or to a drug that targets a different mechanism).

[0200] In particular embodiments, an increase in the level or percent of the neutralizing form of the autoantibody (*e.g.*, NAb) over time is an indication that treatment adjustment should be recommended for the subject. In certain other embodiments, a change from an absence of the neutralizing form of the autoantibody (*e.g.*, NAb) to the presence thereof over time is an indication that treatment adjustment should be recommended for the subject. In these embodiments, the subject can be treated with the current course of therapy (*e.g.*, taking the existing biologic) along with one or more other therapeutic agents. In certain alternative embodiments, the subject can be switched to a different biologic. In certain other alternative embodiments, the subject can be switched to a drug (*e.g.*, biologic and/or non-biologic) that targets a different mechanism.

[0201] In an additional aspect, the present invention provides a method for optimizing therapy and/or reducing toxicity in a subject receiving a course of therapy with a first biologic, the method comprising:

- (a) determining whether a neutralizing form of an autoantibody to the first biologic is cross-reactive with a second (*i.e.*, different) biologic by detecting or measuring the presence, level, or percent of a neutralizing form of the autoantibody in a sample from the subject in accordance with an assay described herein; and
- (b) determining that a different course of therapy should be administered to the subject if the neutralizing form of the autoantibody is cross-reactive with the second biologic.

[0202] In certain embodiments, determining that a different course of therapy should be administered comprises switching to a drug (*e.g.*, biologic and/or non-biologic) that targets a different mechanism.

[0203] In some embodiments, the method further comprises determining that a subsequent dose of the current course of therapy be increased or decreased, or that a different course of therapy should be administered to the subject if the neutralizing form of the autoantibody is not cross-reactive with the second biologic. In certain instances, the different course of therapy comprises treatment with the second biologic. In certain other instances, the different course of therapy comprises treatment with the first or second biologic along with one or more other therapeutic agents.

[0204] In one particular aspect, the present invention provides a method for monitoring and/or optimizing therapy to an anti-TNF α drug in a subject receiving a course of therapy with the anti-TNF α drug, the method comprising:

- 5 (a) detecting or measuring the presence, level, or percent of a neutralizing form of an autoantibody to the anti-TNF α drug in accordance with the assay described herein at a plurality of time points over the course of therapy;
- (b) detecting a change in the presence, level, or percent of the neutralizing form of the autoantibody over time; and
- 10 (c) determining a subsequent dose of the course of therapy for the subject or whether a different course of therapy should be administered to the subject based upon the change in the presence, level, or percent of the neutralizing form of the autoantibody over time.

[0205] In certain embodiments, the plurality of time points comprises at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, or more time points.

15 [0206] In another particular aspect, the present invention provides a method for monitoring and/or optimizing therapy to an anti-TNF α drug in a subject receiving a course of therapy with the anti-TNF α drug, the method comprising:

- 20 (a) measuring the level or percent of a neutralizing form of an autoantibody to the anti-TNF α drug in a first sample from the subject as described herein at time point t_0 ;
- (b) measuring the level or percent of the neutralizing form of the autoantibody in a second sample from the subject as described herein at time point t_1 ;
- (c) optionally repeating step (b) with n additional samples from the subject at time points t_{n+1} , wherein n is an integer from 1 to about 25 (*e.g.*, n is 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or any range therein);
- 25 (d) detecting a change in the level or percent of the neutralizing form of the autoantibody from time points t_0 to t_1 or from time points t_0 to t_{n+1} ; and
- (e) determining a subsequent dose of the course of therapy for the subject or
- 30 whether a different course of therapy should be administered to the subject based upon the change in the level or percent of the neutralizing form of the autoantibody over time.

[0207] In certain other embodiments, the level or percent of the neutralizing form of the autoantibody (*e.g.*, NAb) is measured during the course of anti-TNF α drug therapy at one or

more (*e.g.*, a plurality) of the following weeks: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 80, 90, 100, *etc.*

[0208] In some embodiments, determining a subsequent dose of the course of therapy for the subject comprises maintaining, increasing, or decreasing a subsequent dose of the course of therapy for the subject. In other embodiments, determining a different course of therapy for the subject comprises treatment with a different anti-TNF α drug. In other embodiments, determining a different course of therapy for the subject comprises treatment with the current course of therapy along with another therapeutic agent including, but not limited to, an anti-TNF therapy, an immunosuppressive agent, a corticosteroid, a drug that targets a different mechanism, a nutrition therapy, and other combination treatments. In further embodiments, determining a different course of therapy for the subject comprises changing the current course of therapy (*e.g.*, switching to a different anti-TNF drug or to a drug that targets a different mechanism such as an IL-6 receptor-inhibiting monoclonal antibody, anti-integrin molecule (*e.g.*, Tysabri, Vedaluzamab), JAK-2 inhibitor, and tyrosine kinase inhibitor, or to a nutrition therapy (*e.g.*, special carbohydrate diet)).

[0209] In particular embodiments, an increase in the level or percent of the neutralizing form of the autoantibody (*e.g.*, NAb) over time is an indication that treatment adjustment should be recommended for the subject. In certain other embodiments, a change from an absence of the neutralizing form of the autoantibody (*e.g.*, NAb) to the presence thereof over time is an indication that treatment adjustment should be recommended for the subject. In these embodiments, the subject can be treated with the current course of therapy (*e.g.*, taking the existing anti-TNF α drug) along with one or more immunosuppressive agents such as, *e.g.*, methotrexate (MTX) or azathioprine (AZA). In certain alternative embodiments, the subject can be switched to a different anti-TNF α drug. In certain other alternative embodiments, the subject can be switched to a drug that targets a different mechanism (*e.g.*, a non-anti-TNF α drug).

[0210] In yet another particular aspect, the present invention provides a method for optimizing therapy and/or reducing toxicity in a subject receiving a course of therapy with a first anti-TNF α drug, the method comprising:

- (a) determining whether a neutralizing form of an autoantibody to the first anti-TNF α drug is cross-reactive with a second (*i.e.*, different) anti-TNF α drug by detecting or measuring the presence, level, or percent of a neutralizing form of

the autoantibody in a sample from the subject in accordance with an assay described herein; and

- (b) determining that a different course of therapy should be administered to the subject if the neutralizing form of the autoantibody is cross-reactive with the second anti-TNF α drug.

[0211] In certain embodiments, determining that a different course of therapy should be administered comprises switching to a drug that targets a different mechanism (*e.g.*, a non-anti-TNF α drug). Non-limiting examples of such drugs include an IL-6 receptor-inhibiting monoclonal antibody, anti-integrin molecule (*e.g.*, Tysabri, Vedaluzamab), JAK-2 inhibitor, tyrosine kinase inhibitor, a nutrition therapy (*e.g.*, special carbohydrate diet), and mixtures thereof.

[0212] In some embodiments, the method further comprises determining that a subsequent dose of the current course of therapy be increased or decreased, or that a different course of therapy should be administered to the subject if the neutralizing form of the autoantibody is not cross-reactive with the second anti-TNF α drug. In certain instances, the different course of therapy comprises treatment with the second anti-TNF α drug. In certain other instances, the different course of therapy comprises treatment with the first or second anti-TNF α drug along with one or more immunosuppressive agents such as MTX or AZA.

[0213] Methods for detecting anti-TNF α drugs and anti-drug antibodies are further described in PCT Publication No. WO 2011/056590, the disclosure of which is hereby incorporated by reference in its entirety for all purposes.

[0214] In certain instances, the present invention may further comprise administering to a subject a therapeutically effective amount of a course of therapy such as an anti-TNF α drug or a drug that targets a different mechanism (*e.g.*, a non-anti-TNF α drug) useful for treating one or more symptoms associated with a TNF α -mediated disease or disorder (*e.g.*, IBD such as CD or UC). For therapeutic applications, the course of therapy can be administered alone or co-administered in combination with one or more additional agents as described herein. As such, the present invention advantageously enables a clinician to practice "personalized medicine" by guiding treatment decisions and informing therapy selection and optimization for anti-TNF α drugs such that the right drug is given to the right patient at the right time.

IV. Acid Dissociation

[0215] In certain aspects, the assay methods of the present invention further comprise an acid dissociation step, *e.g.*, to enable equilibration of immune complexes for measuring the

presence or level of neutralizing autoantibodies (NAb), non-neutralizing autoantibodies (non-NAb), and/or isotypes thereof that are generated against biologics such as anti-TNF α drugs. As a result, the presence or level of NAb and/or non-NAb to a biologic (*e.g.*, anti-TNF α drug) administered to a subject in need thereof can be measured without substantial interference from the administered biologic that is also present in the subject's sample. In particular, a subject's sample can be incubated with an amount of acid that is sufficient to provide for the measurement of the presence or level of NAb and/or non-NAb in the presence of the biologic (*e.g.*, anti-TNF α drug) but without substantial interference from high biologic drug levels.

[0216] In some embodiments, step (a) of the assay methods of the present invention may comprise:

- (a') contacting the sample with an acid to dissociate preformed complexes of the autoantibody (*e.g.*, including neutralizing and/or non-neutralizing forms thereof) and the biologic (*e.g.*, anti-TNF α drug);
- (b') contacting the sample with a labeled biologic (*e.g.*, anti-TNF α drug) and a labeled biologic binding moiety (*e.g.*, TNF α) following dissociation of the preformed complexes; and
- (c') neutralizing the acid in the sample to form:
 - (i) a first labeled complex of the labeled biologic (*e.g.*, anti-TNF α drug) and the autoantibody; and/or
 - (ii) a second labeled complex of the labeled biologic (*e.g.*, anti-TNF α drug), the labeled biologic binding moiety (*e.g.*, TNF α), and the autoantibody.

[0217] In some alternative embodiments, steps (a') and (b') are performed simultaneously, *e.g.*, the sample is contacted with an acid, a labeled biologic (*e.g.*, anti-TNF α drug), and a labeled biologic binding moiety (*e.g.*, TNF α) at the same time. In other alternative embodiments, step (b') is performed prior to step (a'), *e.g.*, the sample is first contacted with a labeled biologic (*e.g.*, anti-TNF α drug) and a labeled biologic binding moiety (*e.g.*, TNF α), and then contacted with an acid. In further embodiments, steps (b') and (c') are performed simultaneously, *e.g.*, the sample is contacted with a labeled biologic (*e.g.*, anti-TNF α drug) and a labeled biologic binding moiety (*e.g.*, TNF α) and neutralized (*e.g.*, by contacting the sample with one or more neutralizing agents) at the same time.

[0218] In particular embodiments, the sample is contacted with an amount of an acid that is sufficient to dissociate preformed complexes of the autoantibody and the biologic (*e.g.*, anti-TNF α drug), such that the labeled biologic binding moiety (*e.g.*, TNF α), the labeled biologic

(*e.g.*, anti-TNF α drug), the unlabeled biologic (*e.g.*, anti-TNF α drug), and the autoantibody to the biologic (*e.g.*, anti-TNF α drug) can equilibrate and form complexes therebetween. In certain embodiments, the sample can be contacted with an amount of an acid that is sufficient to allow for the detection and/or measurement of the autoantibody in the presence of a high level of the biologic (*e.g.*, anti-TNF α drug).

[0219] In some embodiments, the phrase “high level of a biologic” such as a high level of an anti-TNF α drug includes drug levels of from about 10 to about 100 $\mu\text{g/mL}$, about 20 to about 80 $\mu\text{g/mL}$, about 30 to about 70 $\mu\text{g/mL}$, or about 40 to about 80 $\mu\text{g/mL}$. In other embodiments, the phrase “high level of a biologic” such as a high level of an anti-TNF α drug includes drug levels greater than or equal to about 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 $\mu\text{g/mL}$.

[0220] In some embodiments, the acid comprises an organic acid. In other embodiments, the acid comprises an inorganic acid. In further embodiments, the acid comprises a mixture of an organic acid and an inorganic acid. Non-limiting examples of organic acids include citric acid, isocitric acid, glutamic acid, acetic acid, lactic acid, formic acid, oxalic acid, uric acid, trifluoroacetic acid, benzene sulfonic acid, aminomethanesulfonic acid, camphor-10-sulfonic acid, chloroacetic acid, bromoacetic acid, iodoacetic acid, propanoic acid, butanoic acid, glyceric acid, succinic acid, malic acid, aspartic acid, and combinations thereof. Non-limiting examples of inorganic acids include hydrochloric acid, nitric acid, phosphoric acid, sulfuric acid, boric acid, hydrofluoric acid, hydrobromic acid, and combinations thereof.

[0221] In certain embodiments, the amount of an acid corresponds to a concentration of from about 0.01M to about 10M, about 0.1M to about 5M, about 0.1M to about 2M, about 0.2M to about 1M, or about 0.25M to about 0.75M of an acid or a mixture of acids. In other embodiments, the amount of an acid corresponds to a concentration of greater than or equal to about 0.01M, 0.05M, 0.1M, 0.2M, 0.3M, 0.4M, 0.5M, 0.6M, 0.7M, 0.8M, 0.9M, 1M, 2M, 3M, 4M, 5M, 6M, 7M, 8M, 9M, or 10M of an acid or a mixture of acids. The pH of the acid can be, for example, about 0.1, 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, or 6.5.

[0222] In some embodiments, the sample is contacted with an acid an amount of time that is sufficient to dissociate preformed complexes of the autoantibody and the biologic (*e.g.*, anti-TNF α drug). In certain instances, the sample is contacted (*e.g.*, incubated) with an acid for a period of time ranging from about 0.1 hours to about 24 hours, about 0.2 hours to about 16 hours, about 0.5 hours to about 10 hours, about 0.5 hours to about 5 hours, or about 0.5 hours to about 2 hours. In other instances, the sample is contacted (*e.g.*, incubated) with an

acid for a period of time that is greater than or equal to about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 6, 7, 8, 9, or 10 hours. The sample can be contacted with an acid at 4°C, room temperature (RT), or 37°C.

[0223] In certain embodiments, the step of neutralizing the acid comprises raising the pH of the sample to allow the formation of first and/or second labeled complexes described herein. In some embodiments, the acid is neutralized by the addition of one or more neutralizing agents such as, for example, strong bases, weak bases, buffer solutions, and combinations thereof. One skilled in the art will appreciate that neutralization reactions do not necessarily imply a resultant pH of 7. In some instances, acid neutralization results in a sample that is basic. In other instances, acid neutralization results in a sample that is acidic (but higher than the pH of the sample prior to adding the neutralizing agent). In particular embodiments, the neutralizing agent comprises a buffer such as phosphate buffered saline (*e.g.*, 10x PBS) at a pH of about 7.3.

[0224] In some embodiments, step (b') further comprises contacting an internal control with the sample together with a labeled biologic (*e.g.*, anti-TNF α drug) and a labeled biologic binding moiety (*e.g.*, TNF α) (*e.g.*, before, during, or after dissociation of the preformed complexes). In certain instances, the internal control comprises a labeled internal control such as, *e.g.*, Biocytin-Alexa 488. In certain other instances, the amount of the labeled internal control ranges from about 1 ng to about 25 ng, about 5 ng to about 25 ng, about 5 ng to about 20 ng, about 1 ng to about 20 ng, about 1 ng to about 10 ng, or about 1 ng to about 5 ng per 100 μ L of sample analyzed. In further instances, the amount of the labeled internal control is greater than or equal to about 1 ng, 5 ng, 10 ng, 15 ng, 20 ng, or 25 ng per 100 μ L of sample analyzed.

[0225] As one non-limiting example of the methods of the present invention, samples such as serum samples (*e.g.*, serum from subjects receiving therapy with an anti-TNF α drug such as Remicade (IFX)) can be incubated with 0.5M citric acid, pH 3.0 for one hour at room temperature. Following the dissociation of preformed complexes between (unlabeled) anti-TNF α drug and autoantibodies to the anti-TNF α drug (*e.g.*, anti-drug antibodies such as anti-IFX antibodies (ATI)), labeled anti-TNF α drug (*e.g.*, IFX-Alexa 488), labeled TNF α (*e.g.*, TNF α -Alexa 532), and optionally an internal control can be added and the reaction mixture (*e.g.*, immediately) neutralized with a neutralizing agent such as 10x PBS, pH 7.3. After neutralization, the reaction mixture can be incubated for another hour at room temperature (*e.g.*, on a plate shaker) to allow equilibration and to complete the reformation of immune complexes between the labeled TNF α , the labeled anti-TNF α drug, the unlabeled anti-TNF α

drug, and/or the autoantibody to the anti-TNF α drug. The samples can then be filtered and analyzed by SEC-HPLC as described herein.

[0226] In particular embodiments, the methods of the present invention (*e.g.*, comprising acid dissociation followed by homogeneous solution phase binding kinetics) significantly increases the IFX drug tolerance such that NAb and/or non-NAb ATI can be measured in the presence of IFX up to about 60 $\mu\text{g/mL}$. In other words, the methods of the present invention can detect the presence or level of NAb and/or non-NAb to anti-TNF α drugs such as ATI as well as autoantibodies to other anti-TNF α drugs in the presence of high levels of anti-TNF α drugs (*e.g.*, IFX), but without substantial interference therefrom.

[0227] Methods for detecting anti-drug antibodies using acid dissociation are further described in PCT Application No. PCT/US2012/025437, filed February 16, 2012, the disclosure of which is hereby incorporated by reference in its entirety for all purposes.

V. Biologic Therapy

[0228] The assays of the present invention are suitable for detecting and/or measuring the presence or absence (*e.g.*, whether positive or negative), level, or percent of neutralizing and/or non-neutralizing autoantibodies to any biologic in a sample from a subject (*e.g.*, a subject receiving biologic therapy). Non-limiting examples of biologics include antibodies, antibody fragments, proteins, polypeptides, peptides, fusion proteins (*e.g.*, Ig fusion proteins or Fc fusion proteins), multivalent binding proteins (*e.g.*, DVD Ig), antibody-drug conjugates, vaccines, nucleic acids, sugars, recombinant forms thereof, engineered forms thereof, and combinations thereof.

[0229] Examples of antibody-based biologics include, but are not limited to, therapeutic monoclonal antibodies and antigen-binding fragments thereof. In particular embodiments, the antibody comprises an anti-TNF α drug such as REMICADE[™] (infliximab), HUMIRA[™] (adalimumab), CIMZIA[®] (certolizumab pegol), SIMPONI[®] (golimumab; CNTO 148), or combinations thereof. Additional examples of antibody-based biologics include antibody-drug conjugates such as Adcetris[™] (brentuximab vedotin). Table 1 provides an exemplary list of therapeutic monoclonal antibodies which have either been approved or are currently in development. An extensive list of monoclonal antibody therapeutics in clinical development and approved products is provided in the 2006 PhRMA Report entitled "418 Biotechnology Medicines in Testing Promise to Bolster the Arsenal Against Disease," the disclosure of which is hereby incorporated by reference in its entirety for all purposes.

TABLE 1		
<u>Therapeutic monoclonal antibodies</u>		
Product Name	Company	Indication(s)
Inflammatory Diseases		
Remicade™ (infliximab)	Janssen Biotech, Inc.	Crohn's disease
ABT 874	Abbott Laboratories	Crohn's disease
Stelara® (ustekinumab)	Janssen Biotech, Inc.	Crohn's disease
Humira™ (adalimumab)	Abbott Laboratories	Crohn's disease
MDX-1100	Millennium Pharmaceuticals	ulcerative colitis
Nuvion® (visilizumab)	PDL BioPharma	I.V. steroid-refractory ulcerative colitis and Crohn's disease
Tysabri® (natalizumab)	Biogen Idec	Crohn's disease
Simponi® (golimumab)	Janssen Biotech, Inc.	uveitis
Autoimmune disorders		
Humira™ (adalimumab)	Abbott Laboratories	rheumatoid arthritis, ankylosing spondylitis, juvenile rheumatoid arthritis, psoriasis
Remicade™ (infliximab)	Janssen Biotech, Inc.	rheumatoid arthritis
Simponi® (golimumab)	Janssen Biotech, Inc.	rheumatoid arthritis, ankylosing spondylitis, psoriatic arthritis
Rituxan® (rituximab)	Genentech Biogen Idec	rheumatoid arthritis, lupus, primary progressive multiple sclerosis, SLE, relapsing-remitting multiple sclerosis
Tysabri® (natalizumab)	Biogen Idec	multiple sclerosis
Stelara® (ustekinumab)	Janssen Biotech, Inc.	plaque psoriasis, multiple sclerosis
ART 874	Abbott Laboratories	multiple sclerosis
Acterra	Roche	rheumatoid arthritis
AME 527	Applied Molecular	rheumatoid arthritis
AMG 108	Amgen	rheumatoid arthritis
AMG 714	Amgen	rheumatoid arthritis
anti-CD16 MAb	MacroGenics	immune thrombocytopenic
daclizumab (anti-CD25 MAb)	PDL BioPharma Biogen Idec	multiple sclerosis
denosumab (AMG 162)	Amgen	rheumatoid arthritis
ETI-201	Elusys Therapeutics	SLE
HuMax-CD20 (ofatumumab)	Genmab	rheumatoid arthritis
HuZAF™ (fontolizumab)	PDL BioPharma Biogen Idec	rheumatoid arthritis
IMMU-106 (hCD20)	Immunomedics	autoimmune disease
LymphoStat-B™ (belimumab)	Human Genome Sciences	rheumatoid arthritis, SLE
MEDI-545 (MDX-1103)	Medarex MedImmune	lupus
siplizumab (MEDI-507)	MedImmune	psoriasis
MLN 1202	Millennium Pharmaceuticals	multiple sclerosis
ocrelizumab (anti-CD20) (R1594)	Genentech Biogen Idec Roche	multiple sclerosis, rheumatoid arthritis
OKT3-gamma-I	Johnson & Johnson	psoriatic arthritis
TRX 1 (anti-CD4)	TolerRx	cutaneous lupus erythematosus
TRX 4	TolerRx	psoriasis
Infectious diseases		
Synagis® (palivizumab)	MedImmune	prevention of respiratory syncytial virus (RSV)

TABLE 1		
Therapeutic monoclonal antibodies		
Product Name	Company	Indication(s)
MDX-066 (CDA-1)	Medarex	<i>C. difficile</i> disease
anti-HIV-1 MAb	Polymun Scientific	HIV infection
CCR5 MAb	Hunan Genome Sciences	HIV infection
Cytolin® (anti-CD8 MAb)	CytoDyn	HIV infection
NM01	SRD Pharmaceuticals	HIV infection
PRO 140	Progenics Pharmaceuticals	HIV infection
TNX-355	Tanox	HIV infection
ABthrax™ (raxibacumab)	Human Genome Sciences	anthrax
Anthim™ (ETI-204)	Elusys Therapeutics	anthrax
anti-hsp90 MAb	NeuTec Pharma	candidiasis
anti-staph MAb	MedImmune	prevention of staphylococcal infections
Aurexis (tefibazumab)	Inhibitex	prevention and treatment of <i>S. aureus</i> bacteremia
bavituximab	Peregrine Pharmaceuticals	hepatitis C
MDX-1303	Medarex PharmAthene	anthrax
Numax™ (motavizumab)	MedImmune	RSV
Tarvacin™	Peregrine Pharmaceuticals	hepatitis C
XTL 6865	XTL Biopharmaceuticals	hepatitis C
Cancer		
Avastin™ (bevacizumab)	Genentech	metastatic colorectal cancer
Bexxar® (tositumomab)	GlaxoSmithKline	non-Hodgkin's lymphoma
Campath® (alemtuzumab)	Berlex Laboratories Genzyme	B-cell chronic lymphocytic leukemia
Erbix™ (cetuximab)	Bristol-Myers Squibb Medarex	colorectal cancer, squamous cell cancer of the head and neck
Herceptin® (trastuzumab)	Genentech	HER2-overexpressing early stage or metastatic breast cancer
Mylotarg™ (gemtuzumab ozogarnicin)	Wyeth	acute myeloid leukemia
Rituxan® (rituximab)	Genentech Biogen Idec	B-cell non-Hodgkin's lymphoma, indolent non-Hodgkin's lymphoma induction therapy, relapsed or refractory CLL
Zevalin™ (ibritumomab tiuxetan)	Biogen Idec	Non-Hodgkin's lymphoma
1311-huA33	Life Science Pharmaceuticals	colorectal cancer
1D09C3	GPC Biotech	relapsed/refractory B-cell lymphomas
AGS PSCA MAb	Agensys Merck	prostate cancer
AMG 102	Amgen	cancer
AMG 479	Amgen	cancer
AMG 623	Amgen	B-cell chronic lymphocytic leukemia (CLL)
AMG 655	Amgen	cancer
AMG 706	Amgen	imatinib-resistant GIST, advanced thyroid cancer
AMG 706	Amgen	imatinib resistant GIST, advanced thyroid cancer
anti-CD23 MAb	Biogen Idec	CLL
anti-CD80 MAb	Biogen Idec	non-Hodgkin's B-cell lymphoma
anti-idiotypic cancer vaccine	Viventia Biotech	malignant melanoma

TABLE 1		
Therapeutic monoclonal antibodies		
Product Name	Company	Indication(s)
anti-lymphotoxin beta receptor MAb	Biogen Idec	solid tumors
anti-PEM MAb	Somanta Pharmaceuticals	cancer
anti-Tac(Fv)-E38 immunotoxin	National Cancer Institute	leukemia, lymphoma
Avastin® (bevacizumab)	Genentech	relapsed metastatic colorectal cancer, first line metastatic breast cancer, first-line non-squamous NSCLC cancers
AVE 9633 maytansin-loaded anti-CD33 MAb	Sanofi Aventis	AML
bavituximab	Peregrine Pharmaceuticals	solid cancers
CAT 3888	Cambridge Antibody Technology	hairy cell leukemia
chimeric MAb	National Cancer Institute	neuroblastoma
siltuximab (CNTO 328)	Janssen Biotech, Inc.	renal cancer, prostate cancer, multiple myeloma
Cotara™	Peregrine Pharmaceuticals	brain cancer
bivatuzumab	Boehringer Ingelheim Pharmaceuticals	cancer
CP-751871 (figitumumab)	Pfizer	adrenocortical carcinoma, non-small cell lung cancer
CS-1008 (tigatuzumab)	Daiichi Sankyo	pancreatic cancer, colorectal cancer, non-small cell lung cancer, ovarian cancer
BrevaRex™	ViRexx	breast cancer, multiple myeloma
denosumab	Amgen	bone loss induced by hormone ablation therapy for breast or prostate cancer, prolonging bone metastases-free survival, bone metastases in breast cancer
ecromeximab	Kyowa Hakko USA	malignant melanoma
EMD 273063	EMD Lexigen	solid tumors, malignant melanoma, neuroblastoma, SCLC
Erbitux™	Bristol Myers Squibb	head/neck cancer, first-line pancreatic, first-line NSCLC, second-line NSCLC, first line colorectal cancer, second-line colorectal cancer
GMK	Progenies Pharmaceuticals	prevention of recurrence following surgery to remove primary melanoma in high-risk patients
Campath® (alemtuzumab)	National Cancer Institute Berlex Laboratories	leukemia, lymphoma
HGS-ETR1	Human Genome Sciences	hematologic and solid tumors
HGS ETR2 (mapatumumab)	Human Genome Sciences	hematologic and solid tumors
HGS-TR2J	Human Genome Sciences	advanced solid tumors
HuC242-DM4	ImmunoGen	colorectal, gastrointestinal, NSCLC, pancreatic cancers
HuMax-CD4 (zanolimumab)	Genmab Serono	cutaneous T-cell lymphoma, non-cutaneous T-cell lymphoma
HuMax CD20 (ofatumumab)	Genmab	CLL, non-Hodgkin's lymphoma
HuMax-EGFr	Genmab	head and neck cancer
huN901-DM1	ImmunoGen	SCLC multiple myeloma
ipilimumab	Bristol-Myers Squibb Medarex	melanoma monotherapy, leukemia, lymphoma, ovarian, prostate, renal

TABLE 1		
<u>Therapeutic monoclonal antibodies</u>		
Product Name	Company	Indication(s)
		cell cancers, melanoma (MCX-010 +/- DTIC), second-line metastatic melanoma (MDX-010 disomotide/overmotide MDX-1379)
M195-bismuth 213 conjugate	Actinium Pharmaceuticals	AML
M200 (volociximab)	PDL BioPharma Fremont, CA Biogen Idec Cambridge, MA	advanced solid tumors
MAB HeFi-1	National Cancer Institute Bethesda, MD	lymphoma, non-Hodgkin's lymphoma
MDX-060 (iratimumab)	Medarex	Hodgkin's disease, anaplastic large-cell-lymphoma
MDX-070	Medarex	prostate cancer
MDX-214	Medarex	ECFR-expressing cancers
MEDI-522	MedImmune	T-cell lymphoma, melanoma, prostate cancer, solid tumors
MORAb 003	Morphotek	ovarian cancer
MORAb 009	Morphotek	mesothelin-expressing tumors
neuradiab	Bradner Pharmaceuticals	glioblastoma
nimotuzumab	YM Biosciences	squamous cell carcinomas of the head and neck, recurrent or refractory high grade malignant glioma, anaplastic astrocytomas, glioblastomas and diffuse intrinsic pontine glioma
Omnitarg™ (pertuzumab)	Genentech	ovarian cancer
OvaRex® (oregovomab)	ViRexx MAb	ovarian cancer
PAM 4	Merck	pancreatic cancer
panitumumab (rHuMAb EGFr)	Abgenix	colorectal cancer
PSMA-ADC	Progenics Pharmaceuticals	prostate cancer
R1550 RadioTheraCIM	Roche YM BioSciences	metastatic breast cancer, glioma
RAV 12	Raven Biotechnologies	cancer
Rencarex® G250	Wilex AG	renal cancer
SGN30	Seattle Genetics	cutaneous anaplastic large-cell MAb lymphoma, systemic anaplastic large-cell lymphoma, Hodgkin's disease
SGN-33 (lintuzumab)	Seattle Genetics	AML, myelodysplastic syndromes CLL multiple myeloma, non Hodgkin's lymphoma
SGN-40	Seattle Genetics	AML, myelodysplastic syndromes CLL multiple myeloma, non Hodgkin's lymphoma
sibroturtumab	Life Science Pharmaceuticals	colorectal, head and neck, lung cancers
Tarvacin™ (bavituximab)	Peregrine Pharmaceuticals	solid tumors
tremelimumab	Pfizer	metastatic melanoma, prostate cancer
TNX-650	Tanox	refractory Hodgkin's lymphoma
Zevalin™ (ibritumomab tiuxetan)	Spectrum Pharmaceuticals	non-Hodgkin's lymphoma
Blood disorders		
ReoPro® (abciximab)	Eli Lilly	adjunct to percutaneous coronary intervention for the prevention of cardiac ischemic complications

TABLE 1		
Therapeutic monoclonal antibodies		
Product Name	Company	Indication(s)
urtoxazumab	Teijin Pharma	hemolytic uremic
afelimomab	Abbot Laboratories	sepsis, septic shock
eculizumab	Alexion Pharmaceuticals	paroxysmal nocturnal hemoglobinuria
Cardiovascular disease		
MLN 1202	Millennium Pharmaceuticals	atherosclerosis
pexelizumab	Alexion Pharmaceuticals Procter & Gamble Pharmaceuticals	acute myocardial infarction, cardiopulmonary bypass
Diabetes and Related Conditions		
anti-CD3 MAb	MacroGenics	type-1 diabetes mellitus
OKT3-gamma-1	Johnson & Johnson	type-1 diabetes mellitus
TRX 4 (anti-CD3)	TolerRx	type-1 diabetes mellitus
Genetic Disorders		
Soliris™ (eculizumab)	Alexion Pharmaceuticals	paroxysmal nocturnal hemoglobinuria (PNH)
Neurological Disorders		
RN624	Rinat Neuroscience	osteoarthritis pain
RN1219	Rinat Neuroscience	Alzheimer's disease
Respiratory Disorders		
ABN 912	Novartis Pharmaceuticals	asthma, chronic obstructive pulmonary disorders (COPD)
ABX-IL8	Amgen	COPD
AMG 317	Amgen	asthma
daclizumab (anti-CD25 MAb)	Protein Design Labs Roche	asthma
MEDI-528 (anti-TL-9 MAb)	MedImmune	asthma
mepolizumab (anti-TL5 MAb)	GlaxoSmithKline	asthma and nasal polyposis
TNX-832	Tanox Houston, TX	respiratory diseases
Xolair® (omalizumab)	Genentech Novartis Pharmaceuticals	pediatric asthma
Transplantation		
ORTHOCLONE OKT® 3 (muromomab-CD3)	Ortho Biotech	acute kidney transplant rejection, reversal of heart and liver transplant rejection
Simulect® (basiliximab)	Novartis Pharmaceuticals	prevention of renal transplant rejection
Zenapax® (daclizumab)	Roche	prophylaxis of acute kidney transplant rejection
OKT3-gamma-1	Protein Design Labs Johnson & Johnson	renal transplant rejection
Other		
CR 0002	CuraGen	kidney inflammation
denosumab (AMG 162)	Amgen	postmenopausal osteoporosis
mepolizumab (anti-IL5 MAb)	GlaxoSmithKline	hypereosinophilic syndrome, eosinophilic esophagitis
Xolair® (omalizumab)	Genentech Tanox	peanut allergy

- [0230] Non-limiting examples of protein-based or polypeptide-based biologics include cytokines (*e.g.*, interleukins), chemokines, growth factors, blood-production stimulating proteins (*e.g.*, erythropoietin), hormones (*e.g.*, Elonva[®] (follicle stimulating hormone), growth hormone), enzymes (*e.g.*, Pulmozyme[®] (dornase alfa)), clotting factors, insulin, albumin, fragments thereof, conservatively modified variants thereof, analogs thereof, and combinations thereof.
- [0231] Examples of cytokines include, but are not limited to, TNF α , TNF-related weak inducer of apoptosis (TWEAK), osteoprotegerin (OPG), IFN- α , IFN- β , IFN- γ , interleukins (*e.g.*, IL-1 α , IL-1 β , IL-1 receptor antagonist (IL-1ra), IL-2, IL-4, IL-5, IL-6, soluble IL-6 receptor (sIL-6R), IL-7, IL-8, IL-9, IL-10, IL-12, IL-13, IL-15, IL-17, IL-23, and IL-27), adipocytokines (*e.g.*, leptin, adiponectin, resistin, active or total plasminogen activator inhibitor-1 (PAI-1), visfatin, and retinol binding protein 4 (RBP4)), and combinations thereof. In particular embodiments, the interleukin comprises IL-2 such as Proleukin[®] (aldesleukin; recombinant IL-2).
- [0232] Examples of chemokines include, but are not limited to, CXCL1/GRO1/GRO α , CXCL2/GRO2, CXCL3/GRO3, CXCL4/PF-4, CXCL5/ENA-78, CXCL6/GCP-2, CXCL7/NAP-2, CXCL9/MIG, CXCL10/IP-10, CXCL11/I-TAC, CXCL12/SDF-1, CXCL13/BCA-1, CXCL14/BRAK, CXCL15, CXCL16, CXCL17/DMC, CCL1, CCL2/MCP-1, CCL3/MIP-1 α , CCL4/MIP-1 β , CCL5/RANTES, CCL6/C10, CCL7/MCP-3, CCL8/MCP-2, CCL9/CCL10, CCL11/Eotaxin, CCL12/MCP-5, CCL13/MCP-4, CCL14/HCC-1, CCL15/MIP-5, CCL16/LEC, CCL17/TARC, CCL18/MIP-4, CCL19/MIP-3 β , CCL20/MIP-3 α , CCL21/SLC, CCL22/MDC, CCL23/MPIF1, CCL24/Eotaxin-2, CCL25/TECK, CCL26/Eotaxin-3, CCL27/CTACK, CCL28/MEC, CL1, CL2, CX₃CL1, and combinations thereof.
- [0233] Non-limiting examples of growth factors include epidermal growth factor (EGF), heparin-binding epidermal growth factor (HB-EGF), vascular endothelial growth factor (VEGF), pigment epithelium-derived factor (PEDF; also known as SERPINF1), amphiregulin (AREG; also known as schwannoma-derived growth factor (SDGF)), basic fibroblast growth factor (bFGF), hepatocyte growth factor (HGF), transforming growth factor- α (TGF- α), transforming growth factor- β (TGF- β 1, TGF- β 2, TGF- β 3, *etc.*), endothelin-1 (ET-1), keratinocyte growth factor (KGF; also known as FGF7), bone morphogenetic proteins (*e.g.*, BMP1-BMP15), platelet-derived growth factor (PDGF), nerve growth factor (NGF), β -nerve growth factor (β -NGF), neurotrophic factors (*e.g.*, brain-derived neurotrophic factor (BDNF), neurotrophin 3 (NT3), neurotrophin 4 (NT4), *etc.*), growth differentiation

factor-9 (GDF-9), granulocyte-colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), myostatin (GDF-8), erythropoietin (EPO), thrombopoietin (TPO), and combinations thereof.

[0234] Examples of receptor construct-based or fusion protein-based biologics include, but are not limited to, naturally-occurring receptors linked to an immunoglobulin frame (*e.g.*, Orencia® (abatacept; immunoglobulin CTLA-4 fusion protein), Amevive® (alefacept; IgG1 fusion protein), ENBREL™ (etanercept; recombinant human TNF-receptor fusion protein), engineered proteins combining two different polypeptide species (*e.g.*, Ontak® (denileukin diftitox; engineered protein comprising interleukin-2 and diphtheria toxin), and combinations thereof.

[0235] The present invention can therefore be used in methods for detecting and measuring the presence or level of neutralizing and non-neutralizing autoantibodies to biologics such as anti-TNFα drug therapeutics in a sample from a subject receiving biologic therapy for one or more of the diseases or disorders referred to herein and Table 1, including one or more of the following:

[0236] Inflammatory diseases, such as inflammatory bowel disease (IBD) (*e.g.*, Crohn's disease (CD) and ulcerative colitis (UC)), uveitis, sarcoidosis, Wegener's granulomatosis, and other diseases with inflammation as a central feature;

[0237] Autoimmune diseases, such as rheumatoid arthritis (RA), multiple sclerosis (MS), systemic lupus erythematosus (SLE), ankylosing spondylitis (Bechterew's disease), lupus, psoriatic arthritis, juvenile idiopathic arthritis, psoriasis, and erythematosus;

[0238] Cancer, such as digestive and gastrointestinal cancers (*e.g.*, colorectal cancer, small intestine (small bowel) cancer; gastrointestinal stromal tumors, gastrointestinal carcinoid tumors, colon cancer, rectal cancer, anal cancer, bile duct cancer, gastric (stomach) cancer; esophageal cancer; appendix cancer; and the like); gallbladder cancer; liver cancer; pancreatic cancer; breast cancer; lung cancer (*e.g.*, non-small cell lung cancer); prostate cancer; ovarian cancer; renal cancer (*e.g.*, renal cell carcinoma); cancer of the central nervous system; skin cancer; choriocarcinomas; head and neck cancers; hematological malignancies (*e.g.*, leukemia, lymphoma such as B-cell non-Hodgkin's lymphoma); osteogenic sarcomas (*e.g.*, Ewing sarcoma); soft tissue sarcomas (*e.g.*, Dermatofibrosarcoma Protuberans (DFSP), rhabdomyosarcoma); other soft tissue malignancies, and papillary thyroid carcinomas;

- [0239] Infectious diseases, such as *C. difficile* disease, respiratory syncytial virus (RSV), HIV, anthrax, candidiasis, staphylococcal infections, and hepatitis C;
- [0240] Blood disorders, such as sepsis, septic shock, paroxysmal nocturnal hemoglobinuria, and hemolytic uremic syndrome;
- 5 [0241] Cardiovascular disease, such as atherosclerosis, acute myocardial infarction, cardiopulmonary bypass, and angina;
- [0242] Metabolic disorders, such as diabetes, *e.g.*, type-I diabetes mellitus;
- [0243] Genetic disorders, such as paroxysmal nocturnal hemoglobinuria (PNH);
- [0244] Neurological disorders, such as osteoarthritis pain and Alzheimer's disease;
- 10 [0245] Respiratory disorders, such as asthma, chronic obstructive pulmonary disorders (COPD), nasal polyposis, and pediatric asthma;
- [0246] Skin diseases, such as psoriasis, including chronic moderate to severe plaque psoriasis;
- [0247] Transplant rejection, such as acute kidney transplant rejection, reversal of heart and
15 liver transplant rejection, prevention of renal transplant rejection, prophylaxis of acute kidney transplant rejection, and renal transplant rejection; and/or
- [0248] Other disorders, such as kidney inflammation, postmenopausal osteoporosis (bone disorders), hypereosinophilic syndrome, eosinophilic esophagitis and peanut allergy.
- [0249] In particular embodiments, the subject has a TNF α -mediated disease or disorder
20 such as, *e.g.*, an autoimmune disease (*e.g.*, rheumatoid arthritis) or an inflammatory disease (*e.g.*, inflammatory bowel disease (IBD) such as Crohn's disease (CD) or ulcerative colitis (UC)).

VI. Examples

- [0250] The present invention will be described in greater detail by way of specific
25 examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of noncritical parameters which can be changed or modified to yield essentially the same results.

- [0251] The examples from PCT Application No. PCT/US2012/025437, filed February 16,
30 2012, are hereby incorporated by reference in their entirety for all purposes.

Example 1. Development of a Novel Assay to Monitor Neutralizing Anti-Drug Antibody Formation in IBD Patients.

[0252] This example illustrates a novel homogeneous assay for detecting or measuring the presence or level of neutralizing and/or non-neutralizing anti-drug autoantibodies (ADA) in a patient sample (*e.g.*, serum) using size exclusion chromatography in the presence of labeled (*e.g.*, fluorescently labeled) anti-TNF α drug and labeled TNF α . In particular embodiments, this assay is advantageous because it obviates the need for wash steps which remove low affinity ADA, uses distinct labels (*e.g.*, fluorophores) that allow for detection on the visible and/or IR spectra which decreases background and serum interference issues, increases the ability to detect neutralizing and/or non-neutralizing ADA in patients with a low titer due to the high sensitivity of fluorescent label detection, and occurs as a liquid phase reaction, thereby reducing the chance of any changes in the epitope by attachment to a solid surface such as an ELISA plate.

[0253] Infliximab (IFX) and adalimumab (ADL) are anti-TNF monoclonal antibodies prescribed for the treatment of inflammatory bowel disease (IBD). Anti-drug antibodies (ADA) often develop during the course of therapy. A proportion of these ADA are neutralizing antibodies (NAb). While ADA will negatively impact drug pharmacokinetics, the presence of NAb will additionally cause loss of drug efficacy through blockage of the drug's binding site. This example describes an assay to monitor the development of NAb in IBD patients receiving IFX treatment based on a homogenous mobility shift assay (HMSA) platform and shows the correlation between antibody-to-infliximab (ATI) maturation and NAb formation.

[0254] **Methods:** Serum concentrations of IFX and ATI were measured by HMSA as described in, *e.g.*, PCT Application No. PCT/US2012/025437, filed February 16, 2012, and PCT Publication No. WO 2011/056590, the disclosures of which are hereby incorporated by reference in their entirety for all purposes. For the NAb assay, patient serum containing ATI was first acid dissociated, then two labeled proteins (*e.g.*, IFX-Alexa488 and TNF alpha-Alexa532) were added, followed by neutralization. The solution was diluted to 2% serum, injected by HPLC on a size exclusion column and complexes monitored by fluorescence. The area under the curve (AUC) of the free TNF-Alexa532 peak in each spectrum (*e.g.*, plot or chromatogram) was calculated for controls and patient samples and then a percent NAb calculated. ATI that completely block antigen binding are defined as 100% NAb, 50% means that an equal proportion of ATI in the sample is non-NAb, and 0% means that all ATI is non-NAb. A reference range was established using serum from 75 healthy volunteers. ATI

positive serum samples (>3.13 U/mL) from 132 residual IBD patient serum screened for IFX and ATI levels were analyzed for NAb. Positive controls were created using pooled ATI positive patient serum.

[0255] For data analysis, a peak detection algorithm is used to find all of the peaks and troughs in each spectrum per experiment. A cubic smoothing spline is fit to each spectrum, and peaks and troughs are defined as a change in the first derivative of the signal. A peak is a sign change of the spectrum's slope from positive to negative. Conversely, troughs are defined as a change in sign from negative to positive. The tallest peak within a window at the expected location of the free TNF-Alexa532 peak (e.g., 11.5 to 13 minutes) is taken to be the free peak itself. The troughs directly above and below the detected free peak define the upper and lower limits of the peak itself. Areas under the bound, free (TNF and IFX) and negative control peaks are found by integrating the peak area within the limits described above using the trapezoid rule. The % of the TNF-Alexa532 peak area is then calculated for each sample by using the formula:

$$\% = [(a-b)/c]*100$$

wherein a = AUC of the TNF-Alexa532 peak in an unknown sample, b = AUC of the TNF-Alexa532 peak from a NAb negative control (e.g., IFX-Alexa488 + TNF-Alexa532 in normal human serum), and c = AUC of the free TNF-Alexa532 in normal human serum. For the calculation, "c" is set to 100% and "b" is as close to 0% as possible, although it may vary based on reaction conditions. The range between "b" and "c" defines the maximum window of sensitivity.

[0256] **Results:** The NAb assay of the invention has demonstrated high reproducibility, accuracy, and precision. The intra- and inter-assay precision is less than 20% of CV, and the accuracy of the assay is within 25%. The precision and accuracy obtained with the NAb assay of the invention is substantially better than cell-based assays or ELISAs. IFX drug tolerance is ~6 µg/mL, while TNFα interferes at greater than 1.0 ng/mL. Positive controls from pooled ATI positive patient serum dilute linearly from 40-5% NAb. Analysis of healthy controls shows that samples that return a value of ≥3% (e.g., 3.06%) are considered NAb positive. More than 30 ATI positive patient serum samples (3.12-199.43 U/mL) were screened for NAb, and 26 out of 132 (19.7%) of the ATI positive patient serum samples were NAb positive (mean 22.47%, range 3.29-51.63%). ATI levels greater than 60 U/mL corresponded to highly neutralizing Ab. Further analysis of NAb positive samples reveals a linear correlation between ATI titer and NAb positivity. In particular, Figure 1 illustrates that there was a clear relationship between NAb percent (y-axis) and ATI levels (Spearman Rank

Correlation, $\rho=0.564$, $p < 0.0001$). Figure 2 illustrates that an ATI concentration ≥ 60 U/ml is predictive of NAb positivity (NAb+). Sensitivity = 77.8%; Specificity = 98.1%; Odds ratio = 63.6, $p < 0.0001$, Fisher's Exact Test. Figure 3 illustrates an ATI cutoff analysis and demonstrates that ATI predicts NAb with a ROC AUC of 0.931. True Positive Rate (TPR) = Sensitivity; False Positive Rate (FPR) = $1 - \text{Specificity}$.

[0257] **Conclusion:** Monitoring of NAb, in addition to drug and ADA levels, provides necessary information on the ADA response and helps guide early therapeutic intervention. This method can be applied to the characterization of ADA against any biologic therapy.

Example 2. Patient Case Studies for Monitoring the Formation of Neutralizing Anti-Drug Antibodies Over Time.

[0258] This example illustrates additional embodiments of a novel homogeneous assay for detecting or measuring the presence or level of neutralizing and/or non-neutralizing anti-drug autoantibodies (ADA) in a patient sample (e.g., serum) using size exclusion chromatography in the presence of labeled (e.g., fluorescently labeled) anti-TNF α drug and labeled TNF α . In addition, this example demonstrates time course case studies of IBD patients on anti-TNF α drug therapy for monitoring the formation of neutralizing and/or non-neutralizing anti-drug antibodies and/or a shift from non-neutralizing to neutralizing anti-drug antibodies while the patient is on therapy.

1. Drug and anti-drug antibody assays

[0259] Figure 4 illustrates detection of ATI (i.e., antibody to IFX; "HACA") by the fluid phase mobility shift assay described herein. For example, 444 ng of Alexa488 labeled IFX (18.8 $\mu\text{g/ml}$ in 100% serum) was spiked into a sample to outcompete free IFX. In particular embodiments, patient serum samples containing complexes of IFX and ATI can be subjected to acid dissociation, wherein equilibration with acid dissociation and label addition followed by neutralization is performed.

[0260] Figure 5 illustrates an exemplary ATI/IFX fluid phase mobility shift assay of the present invention. For example, samples containing various concentrations of ATI (standards or unknowns) equilibrated with fluorescently labeled Infliximab (IFX-488) were injected on size exclusion columns in 2% serum. Figure 5 shows that large IFX-488/ATI complexes eluted first, followed by smaller complexes and then unbound IFX-488 and the Alexa488 loading control. Unknown concentrations were determined by interpolation from a standard curve. Detection of IFX followed a similar methodology.

2. Neutralizing and non-neutralizing anti-drug antibody assays

[0261] Figures 6 and 7 illustrate assays of the present invention for determining whether anti-drug antibodies such as ATI are neutralizing or non-neutralizing autoantibodies using size exclusion chromatography to detect the binding of these autoantibodies to fluorescently labeled anti-TNF α drug in the presence of fluorescently labeled TNF α . In one exemplary embodiment, an anti-TNF α drug such as IFX is labeled with a fluorophore "F1", wherein the fluorophore can be detected on either or both the visible and IR spectra. Similarly, TNF α is labeled with a fluorophore "F2", wherein the fluorophore can also be detected on either or both the visible and IR spectra, and wherein "F1" and "F2" are different fluorophores. The labeled anti-TNF α drug and the labeled TNF α are incubated with human serum in a liquid phase reaction to allow the formation of complexes (*i.e.*, immune complexes) between the labeled anti-TNF α drug (*e.g.*, IFX), labeled TNF α , and/or anti-drug antibodies (*e.g.*, ATI) present in the serum.

[0262] Following incubation, the samples are loaded directly onto a size exclusion column and subjected to the HPLC mobility shift assay. Figure 6 illustrates a non-neutralizing anti-drug antibody (ADA) assay of the present invention in which binding of both the anti-drug antibody (*e.g.*, ATI) and the labeled TNF α (*e.g.*, Alexa532 labeled TNF α ; "TNF-532") to the labeled anti-TNF α drug (*e.g.*, Alexa488 labeled IFX; "IFX-488") results in a decrease in free TNF-532 levels. Figure 7 illustrates a neutralizing ADA assay of the present invention in which binding of anti-drug antibody (*e.g.*, ATI) to the labeled anti-TNF α drug (*e.g.*, IFX-488) without binding of the labeled TNF α (*e.g.*, TNF-532) results in substantially the same amount of free TNF-532 levels as the TNF-532 control.

3. Time course studies for monitoring neutralizing and non-neutralizing anti-drug antibodies

[0263] Figures 8-11 illustrate data from a UC patient case study for determining whether anti-drug antibodies such as ATI are neutralizing or non-neutralizing autoantibodies using the mobility shift assays of the present invention. For example, Figure 8 illustrates the levels of IFX and ATI over a time course of 5 samples taken 1, 2, or 3 months apart. Figure 9 shows peak analysis to determine the percentage of free TNF α over time. In particular, the peak area of TNF-532/IFX-488 complexes was subtracted from the free labeled TNF α area of all samples and then % of free TNF α was calculated. Notably, Figure 9 demonstrates an increase in the level of free TNF α over the time course of 5 samples taken 1, 2, or 3 months apart, indicating an increase in neutralizing autoantibody levels. Figure 10 illustrates a shift from the presence of non-neutralizing autoantibodies to neutralizing autoantibodies over time

as exemplified in 3 samples taken 2 or 3 months apart and spiked with IFX. For the "Nov Year 1" sample, non-neutralizing antibody binds to spiked-in IFX and shows a decrease in the TNF-532 peak. For the "Jan Year 2" sample, a mixture of neutralizing antibody (NAb)/non-neutralizing antibody (Ab) shows a small decrease in the TNF-532 peak relative to the level of the initial complex. As ATI becomes almost completely neutralizing ("April Year 2" sample), high IFX levels cannot overcome ATI binding to IFX, preventing any TNF α binding. As such, Figure 10 demonstrates a UC patient ATI profile in which the ATI profile shifts from a non-neutralizing ATI profile to a profile containing a mixture of neutralizing ATI and non-neutralizing ATI to a neutralizing ATI profile over the course of IFX therapy. Figure 11 shows peak analysis to determine the percentage of free TNF α over time in samples that were spiked with IFX. In particular, the peak area of TNF-532/IFX-488 complexes was subtracted from the free TNF α area of all samples and then the percent (%) of free TNF α was calculated. Notably, Figure 11 demonstrates an increase in the level of free TNF α over the time course of samples taken from the UC patient, indicating an increase in neutralizing autoantibody levels and a shift from non-neutralizing ATI to neutralizing ATI while the patient is on IFX therapy.

[0264] Figures 12-14 illustrate various controls performed using the mobility shift assays of the present invention. In particular, Figure 12 shows the use of rabbit anti-human IgG1 Fc as a non-neutralizing antibody (Ab) control. Figure 13 shows the use of ATI positive serum as a mixed neutralizing antibody (NAb)/non-neutralizing antibody (Ab) control. Figure 14 shows that purification of ATI from ATI positive serum results in loss of weaker affinity NAb. Figure 15 illustrates peak analysis from a UC patient case study to determine the percentage of free TNF α in these various controls. In particular, the peak area of the TNF-532/IFX-488 complex was subtracted from the free TNF α area of all samples and then the percent (%) of free TNF α was calculated.

[0265] Figures 16-18 illustrate data from CD patient case studies for determining whether anti-drug antibodies such as ATI are neutralizing or non-neutralizing autoantibodies using the mobility shift assays of the present invention. For example, Figure 16 shows a peak analysis from a CD patient case study to determine the percentage of free TNF α over a time course of 4 samples taken 7 or 8 weeks apart during a 30-week period. Moreover, Figure 17 shows a peak analysis from another CD patient case study to determine the percentage of free TNF α over a time course of 3 samples taken during a 50-week period. In addition, Figure 18 shows a peak analysis from 4 additional CD patient case studies to determine the percentage of free TNF α in a sample at a particular week during or after induction or maintenance of therapy.

Example 3: Detection of Neutralizing Antibody (NAb) Activity via an HPLC Mobility Shift Competitive Ligand-Binding Assay.

- [0266] This example illustrates yet additional embodiments of a novel homogeneous assay for detecting or measuring the presence or level of neutralizing and/or non-neutralizing anti-drug autoantibodies (ADA) in a patient sample (*e.g.*, serum) using an HPLC size exclusion chromatography assay. In addition, this example demonstrates methods for predicting and/or determining the cross-reactivity of NAb with alternative biological drugs such as other anti-TNF drugs.
- [0267] In some embodiments, a multi-tiered approach to immunogenicity testing comprises first screening both drug and anti-drug antibodies by a rapid, sensitive screening assay. This approach is recommended by both the FDA and the EMEA and is a useful management tool for large clinical trials and multiple time points per patient. After confirming the presence of ADA such as ATI, patient samples are then further examined for the presence of neutralizing antibodies that may have significant negative clinical consequences. Neutralizing antibodies interfere with the biological activity by binding to or near the active site, or by induction of conformational changes, inducing a loss of efficacy. Samples containing ATI may also be screened for isotype and epitope specificity. Comparison of patients' clinical responses to product before and following ADA development can provide information on the correlation between ADA development (and antibody characteristics) and clinical responses.
- [0268] A NAb assay has been developed as disclosed herein that utilizes an HPLC mobility shift assay. In certain embodiments, the multi-tiered approach or test comprises or consists of any one, two, or all three of the following tiers: (1) screening to qualitatively determine if a sample is NAb positive (yes/no based on cutpoint established from analysis of normal human serum); (2) confirming that the sample is NAb positive using, *e.g.*, immunocompetition and/or immunodepletion; and/or (3) predicting and/or determining the cross-reactivity of NAb with alternative biological drugs.

I. Screening Tier

- [0269] After a patient sample has been confirmed as positive for ADA, it can be screened for NAb. In certain aspects, a subpopulation of ADA is NAb. In certain embodiments, patient serum containing ADA (*e.g.*, antibody to IFX, also known as "ATI" or "HACA") is first acid dissociated with 0.5M citric acid in HPLC water for 1 hr at room temperature (RT). Samples are prepared in a 96 well plate and incubation is conducted in the dark on a plate shaker. Next, two labeled proteins (*e.g.*, drug-Alexa488 (*e.g.*, IFX-Alexa488) and TNF α -

Alexa532 in HPLC water containing 0.1% BSA) are added. The samples are neutralized by the addition of 10X PBS, pH 7.3, and incubation for 1 hour at RT in the dark on a plate shaker. The samples are diluted to 2% serum with additional 10X buffer and HPLC water. The samples are then injected by HPLC on a size exclusion column. Complexes or species of differing sizes are separated and monitored by fluorescence, *e.g.*, Free TNF α -Alexa532 ("TNF532"), Free IFX-Alexa488 ("IFX488"), TNF532/IFX488 complexes, TNF532/IFX488/ATI complexes (non-neutralizing Ab), and ATI/IFX488 complexes (NAb). After comparing the results to negative (*see, e.g.*, Figures 12, 19) and positive (*see, e.g.*, Figure 13) controls along with a cutoff established from normal human sera (*e.g.*, reference range of 3.06% NAb), the sample can be designated as positive or negative for NAb and a titer can be determined.

[0270] Figure 19 demonstrates detection of non-neutralizing antibody activity via the mobility shift assay. Upon combination of TNF532 with IFX488, there is a shift to the retention time of approximately 8 minutes, indicating the formation of a higher molecular weight complex. The Free IFX-488 peak (around 10.5 minutes) completely disappears and the Free TNF-532 peak (around 12 minutes) almost completely disappears as well (indicating the formation of an ATI/IFX/TNF ternary complex). A non-neutralizing Ab that binds away from the active site of IFX follows a similar pattern. The mouse monoclonal antibody (*e.g.*, around 7 minutes) performs as desired.

[0271] Figure 13 demonstrates detection of neutralizing antibody activity via the mobility shift assay. A completely neutralizing Ab prevents the ability of IFX to bind to TNF (*e.g.*, due to blockage of the active site). This is seen in the chromatogram as a disappearance of the IFX-488 peak with the formation of a higher molecular weight species. The TNF-532 peak will not change. In reality, most patients experience a combination of non-neutralizing/non-neutralizing Ab as seen in the pooled patient serum in Figure 13 (ATI Pos. Serum, solid black line). Rabbit polyclonal antibodies against the F(ab')₂ fragment of IFX/Humira as an improved NAb positive control are also useful.

[0272] Figure 8 illustrates the development of a NAb response over time in a patient during the course of IFX treatment. While they are positive for ATI at all time points, it is not until the Jan Year 2 (light grey arrow, third from top at ~12 min) time point that NAb develops. The ATI/IFX-488 complexes shift to a slightly different retention time (~7.8 minutes) that indicates a different sized complex as compared to complexes of TNF532/IFX488/ATI (~8.2 and 8.8 mins). Confirmation of neutralizing activity in the presence of additional IFX versus

an irrelevant protein (immunocompetition) may be performed as well. Patients such as this would be ideal candidates for treatment adjustment.

[0273] Figure 9 plots the data as a bar graph of the AUC of the % free TNF peak remaining, clearly demonstrating that over time the patient is developing NAb. Even low levels of NAb development observed at early time points are predictive of disease relapse; treatment adjustment for patients displaying this activity is recommended. For example, the patient should be placed on one or more immunosuppressive agents such as methotrexate (MTX) or azathioprine (AZA) while taking the existing anti-TNF drug and/or switched to a different anti-TNF drug.

10 II. Confirmatory Tier

[0274] In the confirmatory tier, drug (*e.g.*, anti-TNF α antibody) is spiked into the sample at a variety of concentrations (*e.g.*, 1-50 $\mu\text{g/mL}$) to determine the neutralizing capability of the sample. In parallel, non-specific IgG is spiked in at similar levels. The samples spiked with drug should show a dose response to the drug and an EC50 of the NAb can be calculated.

15 Non-specific IgG should have no effect. Immunodepletion can also be performed to rule out the effect of the matrix, if necessary.

[0275] Figure 10 illustrates a shift from the presence of non-neutralizing autoantibodies to neutralizing autoantibodies over time as exemplified in 3 samples taken 2 or 3 months apart and spiked with IFX. Patient serum from each time point responds to spiked-in IFX, showing specificity of response. Over time, the NAb becomes more neutralizing and eventually can neutralize >20 $\mu\text{g/mL}$ IFX (the April Year 2 sample does not decrease when IFX is spiked-in). A complete titration can be performed to determine the EC50 of the NAb at each time point.

III. Cross-Reactivity Tier

25 [0276] The cross-reactivity tier is particularly useful for predicting whether a patient will respond to a drug or therapy such as, *e.g.*, an anti-TNF α drug or therapy.

[0277] In some embodiments, the present invention provides methods to rapidly determine which therapeutic drugs will or will not work in a patient (*e.g.*, a Crohn's disease, ulcerative colitis, or rheumatoid arthritis patient) based on the ability of an anti-drug antibody (ADA) to cross-react with a series of different anti-TNF therapeutics. As a non-limiting example, one or more of the following drugs may be tested in patients (*e.g.*, Crohn's disease, ulcerative colitis, and/or rheumatoid arthritis patients) that have NAb to Remicade (infliximab): Enbrel

(etanercept); Humira (adalimumab); Cimzia (certolizumab pegol); and Simponi (golimumab). After testing positive for NAb with a specific drug (e.g., IFX), the NAb assay can then be performed with a series of other drugs (e.g., fluorescently-labeled drugs) using the method of the initial NAb test described above.

5 [0278] The predictive test of the present invention is useful in the management of patient treatment by preventing the use of a drug (e.g., an anti-TNF α drug) that will be neutralized by a patient's antibodies. Without being bound by any particular theory, the sequence of the binding site of the neutralizing ADA has likely developed in such a way to resemble that of TNF α (see, Figure 20). If the NAb neutralizes any of the other anti-TNF drugs, then those
10 other anti-TNF drugs would likely be a poor alternative to the drug that is already being administered as the patient will likely have an immune response. In some embodiments, a cutoff established from normal human serum can be used to determine if a test sample from a patient is positive or negative. The test can be run in a rapid, cost-effective manner in an *in vitro* setting.

15 [0279] The following non-limiting case studies included Patients 1 and 2, who were treated with Remicade (infliximab), but who subsequently lost response to Remicade. Patient 1 had UC and Patient 2 had CD. The mobility shift assay described herein clearly demonstrated that Patients 1 and 2 lost response to Remicade as they developed anti-Remicade antibodies (e.g., ATI). These anti-Remicade antibodies were then shown to be neutralizing antibodies
20 (e.g., NAb).

[0280] Figure 21 illustrates that Patients 1 and 2 developed neutralizing antibodies (NAb). These NAb compete with TNF α for the Remicade binding site. Importantly, these NAb might cross-react with other anti-TNF therapeutics. If the NAb cross-react with other anti-TNF therapeutics, changing to another anti-TNF therapeutic will not help these patients. As
25 such, the predictive assays of the present invention provide advantages over current methods of managing patients who lose response to Remicade, in which positive HACA (detectable antibody) is managed by changing to another anti-TNF agent (see, e.g., Afif *et al.*, *Am. J. Gastroenterol.*, 105(5):1133-9 (2010)).

[0281] To determine the cross-reactivity of NAb produced in response to one anti-TNF
30 drug with other anti-TNF drugs, NAb which developed when the patient was on Remicade (IFX) were tested against Humira (adalimumab). The data shown in Figure 21 clearly demonstrated that NAb generated against IFX cross-react with Humira. Figure 21 illustrates that the free Humira peak (between 10 and 11 minutes, bottom panel of each patient study) is

completely shifted to a higher molecular weight when the patient serum containing NAb is added (~12 minutes, patient study #1; ~12 minutes, patient study #2; bottom panel of each patient study). These results indicate that the NAb binds to Humira in such a way that, to an extent, the NAb prevents TNF α from accessing the antigen-binding site of Humira. Figure 5 22 depicts this schematically for both NAb and non-NAb determinations.

[0282] In certain embodiments, the assay methods of the present invention predict that these patients will not respond to Humira or any other anti-TNF therapeutics. The patient should not be treated with anti-TNF therapy and should be switched to alternative therapy options, including, but not limited to, Actemra, Kineret, Orencia, Rituxan, and/or Arzerra for 10 rheumatoid arthritis (RA), or Tysabri and/or steroids for Crohn's disease (CD).

[0283] Accordingly, the methods of the present invention are particularly advantageous for predicting whether a patient will respond to anti-TNF α therapy by determining or measuring the presence and/or concentration level of neutralizing antibodies (NAb) and/or non-NAb in a sample from the patient. In one embodiment, if the sample contains NAb to one anti-TNF α 15 drug, these NAb will likely cross-react and be neutralizing to other anti-TNF α drugs, such that the recommended treatment adjustment for the patient would be to switch to a drug with a different mechanism of action (e.g., a non-anti-TNF agent). In another embodiment, if the sample contains non-neutralizing ADA to one anti-TNF α drug, then the recommended treatment adjustment for the patient would be to switch to another anti-TNF α drug.

20 **Example 4: Assays for Detecting the Presence and Cross-Reactivity of Neutralizing Anti-Drug Antibodies (NAb).**

[0284] This example illustrates additional embodiments related to the assay methods of the present invention for screening to determine if a sample is NAb positive and predicting and/or determining the cross-reactivity of NAb with alternative biological drugs (*see, e.g.,* 25 Example 3). In particular embodiments, the assay methods described herein are useful for predicting whether a subject receiving a first anti-TNF α drug will respond to alternative anti-TNF α therapy by determining whether a sample obtained from the subject is either positive or negative for NAb. If the sample is positive for NAb, the methods comprise determining whether the NAb will cross-react with a second anti-TNF α drug and recommending that the 30 subject be switched to a non-anti-TNF α drug when the NAb cross-react with the second anti-TNF α drug. If the sample is negative for NAb, the methods comprise recommending that the subject be switched to a second anti-TNF α drug.

[0285] Figure 23 shows the generation and use of an exemplary NAb standard curve of the invention. Samples containing various concentrations of rabbit (Rb) anti-IFX antibody (ATI) serum (*i.e.*, standards or unknowns) equilibrated with fluorescently labeled TNF-532/IFX-488 were injected onto size exclusion columns in 2% serum. Large immune complexes eluted first, followed by smaller complexes and then unbound IFX-488 and TNF-532. Unknown concentrations can be determined by interpolation from the standard curve. Rabbit serum containing different mixtures of NAb and non-NAb can be combined to make controls. The NAb assay described herein has an improved cut-off of 2.72% compared to an old cut-off of 11.63% (N = 50 normal samples). Table 2 provides a summary of NAb clinical studies by patient.

Table 2. NAb Clinical Summary – By Patient

	Study 1 n=154 (290 samples)	Study 2 n=328 (952 samples)	Study 3 n=64 (812 samples)	Study 4
ATI+ (%total)	43 (28%)	73 (22%)	58 (91%)	30
NAB + (% ATI + tested)	12 (28%)	9 (64%)	3 (23 samples) (60%)	5 (17%)
High Nab activity (20 ug/mL) (% total) (% NAB+)	4 (2.6%) (33%)	4 (1.2%) (44%)	2 (3.1%) (66%)	4 (80%)

[0286] The cross-reactivity assay methods of the present invention are particularly useful for predicting whether switching to another biological treatment will be beneficial. After finding that a patient is NAb positive to one drug, fluorescently-labeled alternative drugs can be used in the assay. If patient serum still shows neutralizing capability, the new drug will be unlikely to succeed. Such methods are advantageous because they can be used to screen a panel of drugs in a cost-effective and timely manner to enable a suggestion or indication of the best treatment options.

[0287] Figures 24 and 25 provide additional case studies to the patient studies described in Example 3 and set forth in Figure 21. In particular, Patients 3 and 4, who were treated with Remicade (infliximab, IFX), but who subsequently lost response to IFX, were identified as being patients who will likely not respond to Humira (adalimumab, ADL) because NAb which developed when the patient was on IFX were determined to be cross-reactive with ADL.

[0288] Figure 26 shows non-limiting examples of patient studies which demonstrate ATI affinity maturation and the development of cross-reactive ATI.

Example 5: Development of a Novel Assay to Monitor Neutralizing Anti-Drug Antibody Formation in IBD Patients.

BACKGROUND AND AIMS

[0289] Anti-TNF monoclonal antibodies, such as infliximab (IFX), adalimumab (ADL), and others are prescribed for the treatment of inflammatory bowel disease (IBD). Certain patients will generate anti-drug antibodies (ADA) that can cause loss of drug efficacy and adverse reactions. Neutralizing ADA (NAb) bind to the antigen binding site, preventing access to TNF and directly reducing or preventing drug efficacy. Binding ADA interact with structural components other than the antigen binding site and do not directly affect drug efficacy, though they may alter drug pharmacokinetics. We have developed an assay to monitor the development of NAb in IBD patients receiving IFX or ADL treatment based on our homogenous mobility shift assay (HMSA) platform and show the correlation between ATI/ATA maturation and NAb formation.

METHODS

I. Principle of the Competitive Ligand Binding NAb Assay for ATI

- ATI positive patient serum is acid dissociated.
- IFX-Alexa488 and TNF-Alexa532 added and solution neutralized.
- The solution is diluted to 2% serum, injected by HPLC on a size exclusion column and complexes monitored by fluorescence.
- The area under the curve (AUC) of the free TNF-Alexa532 peak is calculated for controls and patient samples.
- IFX-Alexa488 binds TNF-Alexa532, reducing the assay signal.
- NAb neutralizes IFX-Alexa488 and recovers the assay signal.
- NAb activity is directly proportional to the measured assay signal.
- $\% \text{ Recovery} = (\text{TNF}_{\text{AUC_sample}} - \text{BKGD}) / (\text{TNF}_{\text{AUC_Free Label}} - \text{BKGD}) * 100$, *e.g.*, $(\text{Free labeled TNF}_{\alpha\text{AUC}} \text{ of the Sample} - \text{BKGD (i.e., "background")}) / (\text{Free labeled TNF}_{\alpha\text{AUC}} \text{ of the Control Sample} - \text{BKGD}) * 100$.
- Figure 27 provides an illustration of the competitive ligand binding assay of the invention.

II. Drug and ADA Level Measurement

- Serum concentrations of infliximab, adalimumab and associated ADA were measured by the HMSA as described previously (*see, Wang et al., J. Immunol. Methods, 2012, 382:177*).

III. Samples

- 5 • Serum was from residual samples or an IRB approved study of IBD patients and healthy controls were purchased from Golden West Biologics, Inc.
- Serum isolated from rabbits immunized with infliximab or adalimumab was used to generate polyclonal antibodies against infliximab and adalimumab. The immunoglobulin fraction was purified by affinity chromatography and concentration
10 determined by UV/VIS spectroscopy. Purified ATI and ATA was spiked into normal human serum for the sensitivity and specificity experiments.

RESULTS

[0290] Figure 28 illustrates the clinical utility of the neutralizing assay. (Left) 130 ATI positive residual serum samples from patients undergoing routine testing were analyzed for neutralizing capability (0.0- 430 U/mL, mean = 49.22 U/mL, median = 19.10 U/mL). The
15 assay cut-off was determined from measuring NAb levels in a subset of 33 patients with low ATI positivity. The cut-off with a 99% one-side confidence limit was calculated to be 1.28% by the formula of mean + 2.33 x SD. Of the remaining patients, 37/97 (38.1%) were NAb positive (mean = 7.43% Recovery). (Right) 11 ATA positive serum samples from a cross-
20 sectional study of patients experiencing loss of response to adalimumab therapy were tested for neutralizing capability (7.93-147.9 U/ml, mean=36.97 U/mL, median = 16.39 U/mL). The assay cut-off was set to 1.28% as above. 5/11 (45.5%) of the serum samples were NAb positive (mean = 8.19% Recovery). The higher percentage of NAb in this population may reflect the loss of response seen in these patients.

25 [0291] Figure 29 illustrates the performance characteristics of the neutralizing assay. ATI and ATA was purified from rabbit serum as described in the methods section.

(A) Sensitivity and specificity of the NAb assay. Purified ATI (19.3 µg/mL, left), purified ATA (49.1 µg/mL, right) and monoclonal mouse anti-human Fc (36.8 µg/mL) were spiked into 50% normal human serum, 2-fold serial dilutions performed, and tested in the
30 neutralizing assay. The assay sensitivity is defined as the lowest concentration of NAb in neat serum that generates a positive result in the NAb assay (*i.e.*, greater than 1.28%, horizontal dashed line). The assay sensitivity of the ATI and ATA neutralizing assays were

determined to be 0.15 and 0.914 $\mu\text{g/mL}$, respectively (neat serum, vertical dashed lines). Both assays are specific for ATI (left) or ATA (right).

(B) Interference of infliximab in the NAb assay. In this experiment, the high control was tested in the presence of 2-fold serial dilutions of infliximab, starting at 50 $\mu\text{g/mL}$. The data was normalized to the control maximum signal and plotted as %inhibition of signal (%inhibition = $100 - \% \text{Recovery}_{\text{Nom}}$). The leftmost vertical dashed line indicates the concentration of infliximab (4 $\mu\text{g/mL}$) that increases the CV to greater than 25%. The rightmost vertical dashed line indicates the IC_{50} of 9.9 $\mu\text{g/mL}$.

(C) Precision of NAb positive controls. High, Medium, and Low positive controls were made by diluting pooled ATI positive patient serum (200 U/mL ATI as determined by HMSA) 2-fold with normal human serum. Each control was tested across multiple different days ($n=20$). The mean (CV%) for each control were 51.35 (10.4%), 28.50 (15.03%) and 10.38 (24.22%) for the High, Medium, and Low controls, respectively.

CONCLUSIONS

[0292] The homogenous, fluid phase NAb assay can detect both ATI and ATA with neutralizing capacity. As little as 0.15 and 0.914 $\mu\text{g/mL}$ of neutralizing ATI or ATA, respectively, can be detected with high specificity. The assay can detect NAb in the presence of higher levels of drug (up to 4.0 $\mu\text{g/mL}$) than most cell based or other competitive ligand binding assays. The NAb assay has high accuracy and precision ($\text{CV} < 25\%$) across an extended dynamic range. The presence of neutralizing antibodies may predict loss of response in patients undergoing infliximab or adalimumab therapy. Monitoring of NAb, in addition to drug and ADA levels, will provide necessary information on the ADA response and help guide early therapeutic intervention.

Example 6: Characterization of Neutralizing Anti-Drug Antibody Response in Patients with Loss of Response to Anti-TNF Therapy.

[0293] Anti-TNF monoclonal antibodies, such as infliximab (IFX) and adalimumab (ADL), are prescribed for the treatment of inflammatory bowel disease. Certain patients will generate anti-drug antibodies (ADA) that can cause loss of drug efficacy and adverse reactions. Neutralizing ADA (NAb) bind to the antigen binding site, preventing access to TNF and directly preventing drug efficacy. Binding ADA interact with structural components other than the antigen binding site and do not directly affect drug efficacy, though they may alter drug pharmacokinetics. We have developed an assay to monitor the development of NAb in IBD patients receiving IFX or ADL treatment based on our homogenous mobility shift assay

(HMSA) platform and show the correlation between ATI/ATA maturation and NAb formation.

[0294] **Methods:** 141 ATI or ATA positive patients were collected from an IRB approved study of IBD patients. Serum concentrations of IFX, ADL and associated ADA were measured by HMSA as described previously (Wang *et al.*, *J. Immunol. Methods*, 2012, 382:177). Healthy controls were purchased from Golden West Biologics, Inc. Serum isolated from rabbits immunized with IFX or ADL was used to generate polyclonal antibodies against IFX and ADL. Purified rabbit ATI and ATA was spiked into normal human serum for the sensitivity and specificity experiments. For the NAb Assay, ATI positive patient serum is first acid dissociated. IFX-Alexa488 and TNF-Alexa532 is added and the solution neutralized. The solution is diluted to 2% serum, injected by HPLC on a size exclusion column and complexes monitored by fluorescence. The area under the curve (AUC) of the free TNF-Alexa532 peak is calculated for controls and patient samples. IFX-Alexa488 binds TNF-Alexa532, reducing the assay signal. NAb neutralizes IFX-Alexa488 and recovers the assay signal. NAb activity is directly proportional to the measured assay signal.

[0295] **Results:** Validation of the NAB assay demonstrated high accuracy and precision (CV <25%) across an extended dynamic range. 130 ATI positive patients were analyzed for neutralizing capability (0.0- 430 U/mL, mean = 49.22 U/mL, median = 19.10 U/mL). The assay cut-off was determined from measuring NAb levels in a subset of 33 patients with low ATI positivity. The cut-off with a 99% one-side confidence limit was calculated to be 1.28% by the formula of mean + 2.33 x SD. Of the remaining patients, 37/97 (38.1%) were NAb positive (mean = 7.43% Recovery). 11 ATA positive patients experiencing loss of response to adalimumab therapy were tested for neutralizing capability (7.93-147.9 U/ml, mean=36.97 U/mL, median = 16.39 U/mL). The assay cut-off was set to 1.28% as above. 5/11 (45.5%) of the serum samples were NAb positive (mean = 8.19% Recovery).

[0296] **Conclusions:** The presence of neutralizing antibodies may predict loss of response in patients undergoing infliximab or adalimumab therapy. Monitoring of NAb, in addition to drug and ADA levels, will provide necessary information on the ADA response and may help guide early therapeutic intervention.

Example 7: Improved Homogeneous Mobility Shift Assay (HMSA) for the Detection of Neutralizing Antibodies (Nab) in IBD Patients Treated with Infliximab or Adalimumab.

[0297] **Background:** When treated with infliximab (IFX) and adalimumab (ADA), certain IBD patients will generate antibodies-to-infliximab or adalimumab (ATI/ATA) that can cause

loss of drug efficacy and adverse reactions. ATI and ATA can be neutralizing antibodies (NAb) that bind to the drug's antigen binding site, preventing access to TNF and directly preventing drug efficacy. This example describes the NAb HMSA of the present invention optimized for improved sensitivity and specificity, and shows a correlation with CRP and drug levels in patient samples.

[0298] **Methods:** Serum concentrations of IFX, ADA and associated anti-drug antibodies were measured by HMSA (*see, e.g., Wang et al., J. Immunol. Methods*, 382:177-88 (2012)) in healthy controls and residual patient samples from routine testing. Serum isolated from rabbits immunized with IFX or ADA was used to generate polyclonal antibodies against IFX and ADA, purified and spiked into normal human serum to establish positive controls. For the NAb Assay, ATI positive patient serum is first acid dissociated, IFX-Alexa488 is added and the solution is neutralized, followed by TNF-Alexa488 addition. The solution is diluted to 4% serum, injected by HPLC on a size exclusion column and complexes are monitored by fluorescence. The area under the curve of the free TNF-Alexa488 peak is calculated for controls and patient samples. IFX-Alexa488 binds TNF-Alexa488, reducing the assay signal. NAb neutralizes IFX-Alexa488 and recovers the assay signal. NAb activity is directly proportional to the measured assay signal.

[0299] **Results:** The NAb assay described in Example 6 demonstrated high accuracy and precision (CV <25%) across an extended dynamic range with a positive cutoff of 1.28% recovery (29.8 ng/mL). Optimization of the assay led to the ability to detect less than 10 ng/mL NAb in patient serum and improved accuracy (CV <15%). Assay optimization included the following changes to the NAb assay described in Example 6: (1) switched to TNF-Alexa488 (kept IFX-Alexa488); (2) changed fluorescence detector wavelengths from Ex510_Em540 to Ex494_Em525 (Ex=excitation, Em=emission); (3) changed PMT (*i.e.*, photomultiplier tube) gain from 14 to 18; (4) adjusted the molar ratio of TNF:IFX and decreased the amount of IFX/TNF in the assay; and (5) adjusted incubation conditions, *i.e.*, added IFX, neutralized for 30 minutes, followed by the addition of TNF for 30 minutes. ATI positive patients were analyzed for neutralizing capability (3.1- 200 U/mL) using the optimized assay described herein compared to the assay described in Example 6. The two assay formats correlated well; however, the optimized assay detected a broader range of neutralizing antibodies ($R^2=0.998$). The presence of detectable neutralizing antibodies was associated with both elevated CRP and decreased drug levels ($p<0.001$).

[0300] **Conclusions:** The presence of neutralizing antibodies (NAb) predicts loss of response, elevated CRP, and decreased drug levels in patients undergoing infliximab or

adalimumab therapy. Monitoring of NAb, in addition to drug and anti-drug antibody levels, provides necessary information on the immune response and helps guide early therapeutic intervention.

- [0301] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, one of skill in the art will appreciate that certain changes and modifications may be practiced within the scope of the appended claims. In addition, each reference provided herein is incorporated by reference in its entirety to the same extent as if each reference was individually incorporated by reference.
- 5

WHAT IS CLAIMED IS:

- 1 1. A method for measuring the percent or the level of a neutralizing form
2 of an autoantibody to an anti-TNF α drug in a sample, the method comprising:
 - 3 (a) contacting the sample with a labeled anti-TNF α drug and a labeled TNF α to
4 form:
 - 5 (i) a first labeled complex of the labeled anti-TNF α drug and the
6 autoantibody; and/or
 - 7 (ii) a second labeled complex of the labeled anti-TNF α drug, the labeled
8 TNF α , and the autoantibody;
 - 9 (b) subjecting the first labeled complex and/or the second labeled complex to size
10 exclusion chromatography to separate them from free labeled TNF α , free
11 labeled anti-TNF α drug, and/or a complex of labeled anti-TNF α drug and
12 labeled TNF α ;
 - 13 (c) measuring the level of free labeled TNF α after size exclusion chromatography;
14 and
 - 15 (d) comparing the level of free labeled TNF α measured in step (c) to the level of
16 free labeled TNF α in a control sample, thereby measuring the percent or level
17 of a neutralizing form of the autoantibody.
- 1 2. The method of claim 1, wherein the anti-TNF α drug is selected from
2 the group consisting of REMICADE™ (infliximab), ENBREL™ (etanercept), HUMIRA™
3 (adalimumab), CIMZIA® (certolizumab pegol), SIMPONI® (golimumab; CNTO 148), and
4 combinations thereof.
- 1 3. The method of claim 1 or 2, wherein the autoantibody to the anti-TNF α
2 drug is selected from the group consisting of a human anti-chimeric antibody (HACA), a
3 human anti-humanized antibody (HAHA), a human anti-mouse antibody (HAMA), and
4 combinations thereof.
- 1 4. The method of any one of claims 1 to 3, wherein step (c) comprises
2 measuring the area under the curve (AUC) of the free labeled TNF α peak of the sample after
3 size exclusion chromatography.

1 5. The method of claim 4, wherein the level of the free labeled TNF α in
2 the control sample is calculated by measuring the AUC of the free labeled TNF α peak in the
3 control sample after size exclusion chromatography.

1 6. The method of claim 5, wherein step (d) comprises calculating a ratio
2 of the AUC of the free labeled TNF α peak of the sample to the AUC of the free labeled
3 TNF α peak of the control sample.

1 7. The method of any one of claims 1 to 6, wherein the sample is positive
2 for the neutralizing form of the autoantibody when the sample has greater than an established
3 threshold value of the neutralizing form of the autoantibody.

1 8. The method of claim 7, wherein the threshold value is between about
2 1.25% to about 1.30% of the neutralizing form of the autoantibody.

1 9. The method of any one of claims 1 to 8, wherein the control sample is
2 a reference sample containing only free labeled TNF α .

1 10. The method of any one of claims 1 to 9, wherein the sample is serum.

1 11. The method of any one of claims 1 to 10, wherein the sample is
2 obtained from a subject receiving therapy with the anti-TNF α drug.

1 12. The method of any one of claims 1 to 11, wherein the labeled anti-
2 TNF α drug and the labeled TNF α comprise different fluorophores or fluorescent dyes.

1 13. The method of any one of claims 1 to 12, wherein the neutralizing
2 form of the autoantibody is present in a population of autoantibodies comprising neutralizing
3 autoantibodies and non-neutralizing autoantibodies.

1 14. A method for determining whether a neutralizing form of an
2 autoantibody to a first anti-TNF α drug is cross-reactive with a second anti-TNF α drug, the
3 method comprising:

- 4 (a) measuring the percent or the level of a neutralizing form of the autoantibody
5 in a sample according to a method of any one of claims 1 to 13 to determine
6 whether the sample is positive or negative for the neutralizing form of the
7 autoantibody; and

8 if the sample is positive for the neutralizing form of the autoantibody, then:

- 9 (b) contacting the sample with a labeled second anti-TNF α drug to form a labeled
10 complex of the labeled second anti-TNF α drug and the neutralizing form of
11 the autoantibody;
- 12 (c) subjecting the labeled complex to size exclusion chromatography to separate
13 the labeled complex; and
- 14 (d) detecting the labeled complex, thereby determining whether a neutralizing
15 form of an autoantibody to a first anti-TNF α drug is cross-reactive with a
16 second anti-TNF α drug.

1 15. The method of claim 14, wherein the first and second anti-TNF α drugs
2 are independently selected from the group consisting of REMICADE™ (infliximab),
3 ENBREL™ (etanercept), HUMIRA™ (adalimumab), CIMZIA® (certolizumab pegol),
4 SIMPONI® (golimumab; CNTO 148), and combinations thereof.

1 16. The method of claim 14 or 15, wherein the autoantibody to the first
2 anti-TNF α drug is selected from the group consisting of a human anti-chimeric antibody
3 (HACA), a human anti-humanized antibody (HAHA), a human anti-mouse antibody
4 (HAMA), and combinations thereof.

1 17. The method of any one of claims 14 to 16, wherein the presence of the
2 labeled complex is an indication that the neutralizing autoantibody against the first anti-TNF α
3 drug is cross-reactive with the second anti-TNF α drug.

1 18. The method of any one of claims 14 to 16, wherein the absence of the
2 labeled complex is an indication that the neutralizing autoantibody against the first anti-TNF α
3 drug is not cross-reactive with the second anti-TNF α drug.

1 19. The method of any one of claims 14 to 18, wherein the sample is
2 serum.

1 20. The method of any one of claims 14 to 19, wherein the sample is
2 obtained from a subject receiving therapy with the anti-TNF α drug.

1 21. The method of any one of claims 14 to 20, wherein the labeled second
2 anti-TNF α drug comprises a fluorophore or fluorescent dye.

1 22. A method for monitoring or optimizing therapy to an anti-TNF α drug
2 in a subject receiving a course of therapy with the anti-TNF α drug, the method comprising:

- 3 (a) measuring the percent or the level of a neutralizing form of an autoantibody to
4 the anti-TNF α drug according to a method of any one of claims 1 to 13 at a
5 plurality of time points over the course of therapy;
- 6 (b) detecting a change in the percent or the level of the neutralizing form of the
7 autoantibody over time; and
- 8 (c) determining a subsequent dose of the course of therapy for the subject or
9 whether a different course of therapy should be administered to the subject
10 based upon the change in the percent or the level of the neutralizing form of
11 the autoantibody over time.

1 23. The method of claim 22, wherein the anti-TNF α drug is selected from
2 the group consisting of REMICADE™ (infliximab), ENBREL™ (etanercept), HUMIRA™
3 (adalimumab), CIMZIA® (certolizumab pegol), SIMPONI® (golimumab; CNTO 148), and
4 combinations thereof.

1 24. The method of claim 22 or 23, wherein the autoantibody to the anti-
2 TNF α drug is selected from the group consisting of a human anti-chimeric antibody (HACA),
3 a human anti-humanized antibody (HAHA), a human anti-mouse antibody (HAMA), and
4 combinations thereof.

1 25. The method of any one of claims 22 to 24, wherein the subsequent
2 dose of the course of therapy is increased, decreased, or maintained based upon the change in
3 the percent or the level of the neutralizing form of the autoantibody over time.

1 26. The method of any one of claims 22 to 24, wherein the different course
2 of therapy comprises a different anti-TNF α drug, the current course of therapy along with an
3 immunosuppressive agent, or switching to a course of therapy that is not an anti-TNF α drug.

1 27. The method of claim 26, wherein the different course of therapy is
2 administered when the percent or the level of the neutralizing form of the autoantibody
3 increases over time.

1 28. A method for optimizing therapy and/or reducing toxicity in a subject
2 receiving a course of therapy with a first anti-TNF α drug, the method comprising:

- 3 (a) determining whether a neutralizing form of an autoantibody to the first anti-
4 TNF α drug is cross-reactive with a second anti-TNF α drug by measuring the

5 percent or the level of a neutralizing form of the autoantibody in a sample
6 from the subject according to a method of any one of claims 1 to 13; and
7 (b) determining that a different course of therapy should be administered to the
8 subject if the neutralizing form of the autoantibody is cross-reactive with the
9 second anti-TNF α drug.

1 29. The method of claim 28, wherein the first and second anti-TNF α drugs
2 are independently selected from the group consisting of REMICADE™ (infliximab),
3 ENBREL™ (etanercept), HUMIRA™ (adalimumab), CIMZIA® (certolizumab pegol),
4 SIMPONI® (golimumab; CNTO 148), and combinations thereof.


1 30. The method of claim 28 or 29, wherein the autoantibody to the first
2 anti-TNF α drug is selected from the group consisting of a human anti-chimeric antibody
3 (HACA), a human anti-humanized antibody (HAHA), a human anti-mouse antibody
4 (HAMA), and combinations thereof.

1 31. The method of any one of claims 28 to 30, wherein the different course
2 of therapy comprises switching to a course of therapy that is not an anti-TNF α drug.

1 32. The method of claim 31, wherein the non-anti-TNF α drug is selected
2 from the group consisting of an IL-6 receptor-inhibiting monoclonal antibody, anti-integrin
3 molecule, JAK-2 inhibitor, tyrosine kinase inhibitor, nutrition therapy, and mixtures thereof.

1 33. The method of any one of claims 28 to 32, wherein the method further
2 comprises determining that a subsequent dose of the current course of therapy should be
3 increased or decreased, or that a different course of therapy should be administered to the
4 subject, if the neutralizing form of the autoantibody is not cross-reactive with the second anti-
5 TNF α drug.

1/29



2/29




FIG. 2

3/29






FIG. 3

4/29

Serum
ATI IFX

Equilibration with
Acid dissociation



Label addition followed
by neutralization



Alexa488

444 ng (18.8 $\mu\text{g/mL}$ in
100% serum) of FI-IFX
spiked into sample to
outcompete free
Infiximab.






FIG. 4




- Samples containing various concentrations of ATI (standards or unknowns) equilibrated with fluorescently labeled Infliximab-(IFX-488) injected on size exclusion column in 2% serum
- Large IFX-488/ATI complexes elute first, followed by smaller complexes and then unbound IFX-488 and the Alexa488 loading control
- Unknown concentrations determined by interpolation from standard curve
- Detection of IFX follows similar methodology



Parameter	IFX [µg/mL]	ATI (U/mL)
LOD	0.10	6.22
LLOQ (= cutoff)	0.98	6.75
ULOQ	34.00	150

FIG. 5

6/29



6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
84

7/29





FIG. 7

8/29



Collection Date	IFX [µg/mL]	ATI [U/mL]	Collection Date	IFX [µg/mL]	ATI [U/mL]
July Year 1	<0.98	12.07	Jan Year 2	<0.98	>150 U/mL
Oct Year 1	<0.98	>150 U/mL	April Year 2	<0.98	>150 U/mL
Nov Year 1	<0.98	>150 U/mL			

FIG. 8

9/29





FIG. 9

Collection Date	IFX [µg/mL]	ATI [U/mL]
July Year 1	<0.98	12.07
Oct Year 1	<0.98	>150 U/mL
Nov Year 1	<0.98	>150 U/mL
Jan Year 2	<0.98	>150 U/mL
April Year 2	<0.98	>150 U/mL

Peak area of TNF532/IFX488 complex subtracted from Free TNF area of all samples and then % of free TNF calculated


10/29




1. Non-neutralizing Ab (Nov Year 1) binds to spiked-in IFX and shows decrease in TNF-532 peak
2. Mixed Nab/non-neutralizing Ab (Jan Year 2), shows small decrease in TNF-532 peak to level of initial complex
3. As ATI becomes almost completely neutralizing (April Year 2 sample), high IFX levels cannot overcome ATI binding to IFX, preventing any TNF binding

FIG. 10

11/29




12/29




Rabbit Anti-Human IgG₁ F_c: Non-neutralizing
Ab Control

FIG. 12



ATI Pos. Serum: Mixed Nab/Non-neutralizing
Ab Control

FIG. 13



Purification of ATI from ATI Positive Serum
results in loss of weaker affinity Nab

FIG. 14





FIG. 15

Sample ID	Week #	IFX [µg/mL]	ATI [U/mL]
10113A	7	26.14	17.09
10113B	14	19.26	14.37
10113C	22	17.09	15.12
10113D	30	7.22	10.16




Induction: Weeks 1-14
Maintenance: Weeks 22-50

Peak area of TNF532/IFX488 complex subtracted from all samples and then % of free TNF calculated

FIG. 16

Sample ID	Week #	IFX [µg/mL]	ATI [U/mL]
10509A	22	12.77	11.42
10509B	46	5.63	10.15
10509C	50	27.24	17.57




Induction: Weeks 1-14
Maintenance: Weeks 22-50

Peak area of TNF532/IFX488 complex subtracted from Free TNF area of all samples and then % of free TNF calculated

FIG. 17

Sample ID	Week #	IFX [µg/mL]	ATI [U/mL]
10111	46	8.76	13.59
11505	14	11.42	12.66
11903	60	20.95	11.79
10308	14	1.09	54.64



Induction: Weeks 1-14
Maintenance: Weeks 22-50

Peak area of TNF532/IFX488 complex subtracted from Free TNF area of all samples and then % of free TNF calculated

FIG. 18

19/29




FIG. 19

20/29






FIG. 20

21/29

Patient Study #1 – UC Patient




Collection Date	IFX [μg/mL]	ATI [U/mL]	Confirmed positive (>20 μg/mL, lab activity)
GRD0125	<0.05	>150 U/mL	Yes




Patient will likely not respond to adalimumab

Patient Study #2 – CD Patient





Collection Date	IFX [μg/mL]	ATI [U/mL]	Confirmed positive (>20 μg/mL, lab activity)
PL_CL01956.7	<0.05	356.82 U/mL	Yes



Patient will likely not respond to adalimumab

FIG. 21

22/29

Binding (non-NAB) Antibody**Neutralizing Antibody (NAB)****FIG. 22**





FIG. 23



Patient will likely not respond to adalimumab





FIG. 24


Collection Date	IFX [µg/mL]	ATI [U/mL]
S106080537	<0.98	77.85 U/mL




Patient will
likely not
respond to
adalimumab

FIG. 25

26/29



Note:
All IFX samples with result of NA plotted as 0
All Total ATI samples <3.13 U/mL plotted as 0



Note:
All IFX samples with result of NA plotted as 0
All Total ATI samples <3.13 U/mL plotted as 0

FIG. 26

27/29




FIG. 27




FIG. 28




FIG. 29



(51) International Patent Classification:

G01N 33/50 (2006.01)

(21) International Application Number:

PCT/IB2013/060458

(22) International Filing Date:

27 November 2013 (27.11.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/732,251 30 November 2012 (30.11.2012) US

13/802,117 13 March 2013 (13.03.2013) US

(71) Applicant: NESTEC S.A. [CH/CH]; Avenue Nestlé 55, CH-1800 Vevey (CH).

(72) Inventors: HAUENSTEIN, Scott; 3565 Caminito El Rincon #211, San Diego, California 92130 (US). SINGH, Sharat; PMB 69, P.O. Box 5000, Rancho Santa Fe, California 92067 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:


— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))


[Continued on next page]

(54) Title: ASSAYS FOR DETECTING NEUTRALIZING AUTOANTIBODIES TO BIOLOGIC THERAPY

Binding (non-NAB) Antibody



Neutralizing Antibody (NAB)



(57) Abstract: The present invention provides assays for detecting and measuring the presence or level of neutralizing and non-neutralizing autoantibodies to biologics such as anti-TNF α drug therapeutics in a sample. The present invention is useful for monitoring the formation of neutralizing and/or non-neutralizing anti-drug antibodies over time while a subject is on biologic therapy. The present invention is also useful for predicting and/or determining the cross-reactivity of neutralizing anti-drug antibodies in a subject's sample with alternative biologic therapies. As such, the present invention provides information for guiding treatment decisions for those subjects receiving therapy with a biologic agent and improves the accuracy of optimizing therapy, reducing toxicity, and/or monitoring the efficacy of therapeutic treatment to biologic therapy.

FIG. 22



(88) Date of publication of the international search report:
31 July 2014

INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2013/060458

A. CLASSIFICATION OF SUBJECT MATTER

INV. G01N33/50

ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, COMPENDEX, EMBASE, INSPEC, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2011/056590 A1 (PROMETHEUS LAB INC [US]; SINGH SHARAT [US]; WANG SHUI LONG [US]; OHRMU) 12 May 2011 (2011-05-12) paragraph [0007] - paragraph [0023]; claims 1-79 -----	1-13
X	WO 2012/154253 A1 (PROMETHEUS LAB INC [US]; SINGH SHARAT [US]; WANG SHUI LONG [US]; OHRMU) 15 November 2012 (2012-11-15) paragraph [0007] - paragraph [0022]; claims 1-28 -----	1-13
X	WO 2012/054532 A1 (PROMETHEUS LAB INC [US]; WANG SHUI LONG [US]; OHRMUND LINDA [US]; HAUE) 26 April 2012 (2012-04-26) paragraph [0009] - paragraph [0027]; claims 1-26 -----	1-13



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 March 2014

Date of mailing of the international search report

27/05/2014

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Stachowiak, Olaf

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IB2013/060458

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2011056590 A1	12-05-2011	AU 2010315547 A1	10-05-2012
		CA 2778454 A1	12-05-2011
		CN 102695955 A	26-09-2012
		EP 2494352 A1	05-09-2012
		JP 2013508739 A	07-03-2013
		KR 20120108978 A	05-10-2012
		RU 2012121710 A	10-12-2013
		US 2012329172 A1	27-12-2012
		US 2014057367 A1	27-02-2014
		WO 2011056590 A1	12-05-2011
WO 2012154253 A1	15-11-2012	AU 2012254150 A1	02-05-2013
		CA 2827609 A1	15-11-2012
		CN 103502815 A	08-01-2014
		EP 2676137 A1	25-12-2013
		JP 2014508930 A	10-04-2014
		KR 20140008397 A	21-01-2014
		SG 192770 A1	30-09-2013
		US 2014045276 A1	13-02-2014
		WO 2012154253 A1	15-11-2012
WO 2012054532 A1	26-04-2012	AU 2011317149 A1	02-05-2013
		CA 2815026 A1	26-04-2012
		CN 103299190 A	11-09-2013
		EP 2630495 A1	28-08-2013
		JP 2014500482 A	09-01-2014
		KR 20130139289 A	20-12-2013
		SG 189391 A1	31-05-2013
		US 2013344621 A1	26-12-2013
		WO 2012054532 A1	26-04-2012

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB2013/060458

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-13

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-13

Methods for measuring the level of a neutralizing form of an autoantibody to an anti-TNFalpha drug in a sample.

2. claims: 14-21

Methods for determining whether a neutralizing form of an autoantibody to a first anti-TNFalpha drug is cross-reactive with a second anti-TNFalpha drug.

3. claims: 22-33

Methods for monitoring or optimizing therapy to an anti-TNFalpha drug in a subject receiving a therapy with the anti-TNFalpha drug. Methods for reducing toxicity.

摘要

本发明提供用于检测和测量样品中抗生物制品（如抗 TNF α 药物治疗剂）的中和性自身抗体和非中和性自身抗体的存在或水平的测定。本发明用于监测个体进行生物制品治疗时，随着时间推移，中和性抗药物抗体和/或非中和性抗药物抗体的形成。本发明还用于预测和/或测定个体样品中的中和性抗药物抗体与备选生物制品治疗的交叉反应性。因此，本发明提供用于指导接受生物制剂治疗的那些个体的治疗决定的信息，并改善优化治疗、降低毒性和/或监测生物制品治疗的治疗处理的功效的准确性。