




(86) **Date de dépôt PCT/PCT Filing Date:** 2006/10/05
 (87) **Date publication PCT/PCT Publication Date:** 2007/04/19
 (45) **Date de délivrance/Issue Date:** 2015/01/20
 (85) **Entrée phase nationale/National Entry:** 2008/04/07
 (86) **N° demande PCT/PCT Application No.:** EP 2006/067061
 (87) **N° publication PCT/PCT Publication No.:** 2007/042447
 (30) **Priorité/Priority:** 2005/10/12 (EP05022222.3)

(51) **Cl.Int./Int.Cl. A01N 43/54** (2006.01),
A01N 25/32 (2006.01), **A01P 13/02** (2006.01),
A01N 41/06 (2006.01)
 (72) **Inventeurs/Inventors:**
 ZAGAR, CYRILL, CN;
 SIEVERNICH, BERND, DE
 (73) **Propriétaire/Owner:**
 BASF SE, DE
 (74) **Agent:** ROBIC

(54) **Titre : COMPOSITIONS HERBICIDES BASEES SUR 3-PHENYLURACILES ET N-[[4-[(CYCLOPROPYLAMINO)-
 CARBONYL]PHENYL]SULFONYL]-2-METHOXYBENZAMIDE**
 (54) **Title: HERBICIDAL COMPOSITIONS BASED ON 3-PHENYLURACILS AND N-[[4-[(CYCLOPROPYLAMINO)-
 CARBONYL]PHENYL]SULFONYL]-2-METHOXYBENZAMIDE**



(57) **Abrégé/Abstract:**

The present invention is related to compositions comprising 3-phenyluracils I, wherein the variables R1 to R7 are as defined in the specification, including their agriculturally acceptable salts; and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II including its agriculturally acceptable salts; and optionally at least one herbicide III selected from the classes III.1) to III.15) defined as in the specification including their agriculturally acceptable salts and, provided they have a carboxyl group, their agriculturally acceptable derivatives.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 April 2007 (19.04.2007)

PCT

(10) International Publication Number
WO 2007/042447 A3

(51) International Patent Classification:

A01N 43/54 (2006.01) A01P 13/02 (2006.01)
A01N 25/32 (2006.01) A01N 41/06 (2006.01)

(21) International Application Number:

PCT/EP2006/067061

(22) International Filing Date: 5 October 2006 (05.10.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

05022222.3 12 October 2005 (12.10.2005) EP

(71) Applicant (for all designated States except US): BASF AKTIENGESELLSCHAFT [DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): ZAGAR, Cyrill [DE/CN]; No. 1 Austin Road West, Kowloon, Hong Kong (CN). SIEVERNICH, Bernd [DE/DE]; Bertolt-Brecht-str. 18a, 67454 Hassloch (DE).

(74) Common Representative: BASF AKTIENGESELLSCHAFT; 67056 Ludwigshafen (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

13 December 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HERBICIDAL COMPOSITIONS BASED ON 3-PHENYLURACILS AND N-[[4-[(CYCLOPROPYLAMINO)-CARBONYL]PHENYL]SULFONYL]-2-METHOXYBENZAMIDE

(57) Abstract: The present invention is related to compositions comprising 3-phenyluracils I, wherein the variables R1 to R7 are as defined in the specification, including their agriculturally acceptable salts; and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxy-benzamide II including its agriculturally acceptable salts; and optionally at least one herbicide III selected from the classes III.1) to III.15) defined as in the specification including their agriculturally acceptable salts and, provided they have a carboxyl group, their agriculturally acceptable derivatives.



WO 2007/042447 A3

Herbicidal compositions based on 3-phenyluracils and N-[[4-[(cyclopropylamino)-carbonyl]phenyl]sulfonyl]-2-methoxybenzamide

The present invention relates to herbicidally active compositions comprising 3-phenyluracils I, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and optionally at least one further herbicide III.


In crop protection products, it is desirable in principle to increase the specificity and the reliability of the action of active compounds. In particular, it is desirable for the crop protection product to control the harmful plants effectively and, at the same time, to be tolerated by the useful plants in question. It is known that in some cases better crop plant compatibility can be achieved by joint application of specifically acting herbicides with organic active compounds, which act as antidotes or antagonists. Owing to the fact that they can reduce or even prevent damage to the crop plants, they are also referred to as safeners.

3-phenyluracils I and compositions comprising 3-phenyluracils I are known as highly effective herbicides (e.g. WO 01/83459, WO 03/24221, WO 04/80183). However, their compatibility with dicotyledonous crop plants such as cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers and soybeans and some graminaceous plants such as oat, barley, millet, corn, rice, wheat and sugar cane is not always satisfactory, i. e. in addition to the harmful plants, the crop plants are also damaged to an extent which is not acceptable. It is possible to spare the useful plants by lowering the application rates; however the extent of the control of harmful plants is naturally also reduced.

Therefore it is an object of the present invention to improve the crop-plant compatibility of 3-phenyluracils I, optionally in the presence of further herbicides III.

Surprisingly it has now been found that compositions comprising at least one 3-phenyluracil I or its agriculturally acceptable salts, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II or its agriculturally acceptable salts; and optionally at least one further herbicide III, improve the crop-plant compatibility of 3-phenyluracils I.


N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II (CAS no. 221667-31-8)



is a safener, which is disclosed, for example, in WO 99/16744 and WO 05/00797.

The herbicides III of groups III.1) to III.15) are known herbicides, see the quoted literature references and, for example, Farm Chemicals Handbook 2000, Vol. 86, Meister Publishing Company, 2000; B. Hock, C. Fedtke, R.R. Schmidt, Herbizide, Georg Thieme Verlag, Stuttgart 1995; W. H. Ahrens, Herbicide Handbook, 7th Edition, Weed Science Society of America, 1004; and K.K. Hatzios, Herbicide Handbook, Supplement I to 7th Edition, Weed Science Society of America, 1998.

The present invention is therefore related to compositions comprising a 3-phenyluracil I



I,


wherein the variables R¹ to R⁷ are as defined below:

- R¹ is methyl or NH₂;
- R² is C₁-C₂-haloalkyl;
- R³ is hydrogen or halogen;
- R⁴ is halogen or cyano;
- R⁵ is hydrogen or C₁-C₆-alkyl;
- R⁶, R⁷ independently of one another are hydrogen, C₁-C₆-alkyl, C₁-C₆-alkoxy, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkenyl, phenyl or benzyl;

including its agriculturally acceptable salts; and


N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II

2a



including its agriculturally acceptable salts; and

and optionally at least one herbicide III selected from the following classes III.1) to III.15):



- 5 III.1) lipid biosynthesis inhibitors;
III.2) acetolactate synthase inhibitors (ALS inhibitors);
III.3) photosynthesis inhibitors;
III.4) protoporphyrinogen-IX oxidase inhibitors;
5 III.5) bleacher herbicides;
III.6) enolpyruvyl shikimate 3-phosphate synthase inhibitors (EPSP inhibitors);
III.7) glutamine synthetase inhibitors;
III.8) 7,8-dihydropteroate synthase inhibitors (DHP inhibitors);
III.9) mitose inhibitors;
10 III.10) inhibitors of the synthesis of long chain fatty acids (VLCFA inhibitors);
III.11) cellulose biosynthesis inhibitors;
III.12) decoupler herbicides;
III.13) auxin herbicides;
III.14) auxin transport inhibitors;
15 III.15) other herbicides selected from the group consisting of benzoylprop, flamprop, flamprop-M, bromobutide, chlorflurenol, cinmethylin, methyldymuron, etobenzanid, fosamine, metam, pyributicarb, oxaziclomefone, dazomet, triaziflam and methyl bromide;
- 20 including their agriculturally acceptable salts and, provided they have a carboxyl group, their agriculturally acceptable derivatives.

The invention relates in particular to compositions in the form of safened herbicidally active crop protection compositions comprising a herbicidally effective amount of at least one composition of 3-phenyluracil I with N-[[4-[(cyclopropylamino)carbonyl]-phenyl]sulfonyl]-2-methoxybenzamide II and optionally at least one herbicide III, as defined above, and at least one inert liquid and/or solid carrier, if desired at least one or more surfactants and, if appropriate, at least one or more further auxiliaries customary for crop protection compositions.

30 The invention also relates to compositions in the form of a crop protection composition formulated as a 2-component composition comprising a first component, which comprises the 3-phenyluracil I and the N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, a solid or liquid carrier and, if appropriate, one or more surfactants, and
35 a second component, which comprises at least one further herbicide III and optionally the N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, a solid or liquid carrier and, if appropriate, one or more surfactants, where both components may additionally comprise further auxiliaries customary for
40 crop protection compositions.

The invention also relates to compositions in the form of a crop protection composition formulated as a 2-component composition comprising a first component, which comprises the 3-phenyluracil I and optionally the N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, a solid or liquid carrier and, if appropriate, one or more surfactants, and a second component, which comprises at least one further herbicide III and the N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, a solid or liquid carrier and, if appropriate, one or more surfactants, where both components may additionally comprise further auxiliaries customary for crop protection compositions.

The invention furthermore relates to a method for controlling undesired vegetation, which comprises allowing a herbicidally effective amount of a composition according to the present invention to act on plants, their habitat or on seed.

The invention furthermore relates to a method for controlling undesired vegetation, which comprises applying a herbicidal composition according to the present invention before, during and/or after the emergence of the undesirable plants, the components I, II and optionally III being applied simultaneously or in succession.

The invention furthermore relates to a method for controlling undesirable vegetation in crops, in particular in crops of oat, barley, millet, corn, rice, wheat, sugar cane, cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers and soybeans or in perennial crops.

The invention furthermore relates to a method for controlling undesirable vegetation in crops which, by genetic engineering or by breeding, are resistant to one or more herbicides and/ or fungicides and/or or to attack by insects.

The invention furthermore relates to a method of safening the phytotoxic activity of a 3-phenyluracil I, or an agriculturally acceptable salt thereof, according to the present invention on crops;

preferably crops selected from oat, barley, millet, corn, rice, wheat, sugar cane, cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers, soybeans, rye, triticale, potato, peas, beans, sorghum, grapes, citrus, apple and almond; especially preferred crops selected from oat, barley, millet, corn, rice, wheat, sugar cane, cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers and soybeans;

which comprises applying said 3-phenyluracil I, or an agriculturally acceptable salt thereof, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide of formula II or an agriculturally acceptable salt thereof;

and optionally at least one herbicide III selected from the classes III.1) to III.15) as defined herein,

in an amount effective to reduce or eliminate the phytotoxic activity of said 3-phenyluracil I.

5

In these methods it is immaterial whether the herbicide I, safener II and optionally III are formulated and applied jointly or separately, and, in the case of separate application, in which order the application takes place.

10 The organic moieties mentioned in the definition of the substituents R², R⁵, R⁶, R⁷ in formula I are - like the term halogen - collective terms for individual enumerations of the individual group members. All hydrocarbon chains, i.e. all alkyl, haloalkyl, cycloalkyl, alkoxy, alkenyl, cycloalkenyl and alkynyl groups, can be straight-chain or branched, the prefix C_n-C_m denoting in each case the possible number of carbon atoms in the group.

15 Halogenated substituents preferably carry one, two, three, four or five identical or different halogen atoms. The term halogen denotes in each case fluorine, chlorine, bromine or iodine.

Examples of other meanings are:

20

- C₁-C₄-alkyl: CH₃, C₂H₅, n-propyl, CH(CH₃)₂, n-butyl, CH(CH₃)-C₂H₅, CH₂-CH(CH₃)₂ and C(CH₃)₃;

25

- C₁-C₆-alkyl: C₁-C₄-alkyl as mentioned above, and also, for example, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl or 1-ethyl-2-methylpropyl, preferably methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1,1-dimethylethyl, n-pentyl or n-hexyl;

30

35

- C₁-C₂-haloalkyl: a methyl or ethyl radical, which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, for example CH₂F, CHF₂, CF₃, CH₂Cl, dichloromethyl, trichloromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 2-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-iodoethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-tri-chloroethyl, C₂F₅;

40

- C₁-C₄-alkoxy: OCH₃, OC₂H₅, n-propoxy, OCH(CH₃)₂, n-butoxy, OCH(CH₃)-C₂H₅, OCH₂-CH(CH₃)₂ or OC(CH₃)₃, preferably OCH₃, OC₂H₅ or OCH(CH₃)₂;

- C₁-C₆-alkoxy: a C₁-C₄-alkoxy radical as mentioned above, and also, for example pentoxy, 1-methylbutoxy, 2-methylbutoxy, 3-methoxybutoxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy, 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy;

5

10
- C₃-C₆-alkenyl: prop-1-en-1-yl, allyl, 1-methylethenyl, 1-buten-1-yl, 1-buten-2-yl, 1-buten-3-yl, 2-buten-1-yl, 1-methylprop-1-en-1-yl, 2-methylprop-1-en-1-yl, 1-methylprop-2-en-1-yl, 2-methylprop-2-en-1-yl, n-penten-1-yl, n-penten-2-yl, n-penten-3-yl, n-penten-4-yl, 1-methylbut-1-en-1-yl, 2-methylbut-1-en-1-yl, 3-methylbut-1-en-1-yl, 1-methylbut-2-en-1-yl, 2-methylbut-2-en-1-yl, 3-methylbut-2-en-1-yl, 1-methylbut-3-en-1-yl, 2-methylbut-3-en-1-yl, 3-methylbut-3-en-1-yl, 1,1-dimethylprop-2-en-1-yl, 1,2-dimethylprop-1-en-1-yl, 1,2-dimethylprop-2-en-1-yl, 1-ethylprop-1-en-2-yl, 1-ethylprop-2-en-1-yl, n-hex-1-en-1-yl, n-hex-2-en-1-yl, n-hex-3-en-1-yl, n-hex-4-en-1-yl, n-hex-5-en-1-yl, 1-methylpent-1-en-1-yl, 2-methylpent-1-en-1-yl, 3-methylpent-1-en-1-yl, 4-methylpent-1-en-1-yl, 1-methylpent-2-en-1-yl, 2-methylpent-2-en-1-yl, 3-methylpent-2-en-1-yl, 4-methylpent-2-en-1-yl, 1-methylpent-3-en-1-yl, 2-methylpent-3-en-1-yl, 3-methylpent-3-en-1-yl, 4-methylpent-3-en-1-yl, 1-methylpent-4-en-1-yl, 2-methylpent-4-en-1-yl, 3-methylpent-4-en-1-yl, 4-methylpent-4-en-1-yl, 1,1-dimethylbut-2-en-1-yl, 1,1-dimethylbut-3-en-1-yl, 1,2-dimethylbut-1-en-1-yl, 1,2-dimethylbut-2-en-1-yl, 1,2-dimethylbut-3-en-1-yl, 1,3-dimethylbut-1-en-1-yl, 1,3-dimethylbut-2-en-1-yl, 1,3-dimethylbut-3-en-1-yl, 2,2-dimethylbut-3-en-1-yl, 2,3-dimethylbut-1-en-1-yl, 2,3-dimethylbut-2-en-1-yl, 2,3-dimethylbut-3-en-1-yl, 3,3-dimethylbut-1-en-1-yl, 3,3-dimethylbut-2-en-1-yl, 1-ethylbut-1-en-1-yl, 1-ethylbut-2-en-1-yl, 1-ethylbut-3-en-1-yl, 2-ethylbut-1-en-1-yl, 2-ethylbut-2-en-1-yl, 2-ethylbut-3-en-1-yl, 1,1,2-trimethylprop-2-en-1-yl, 1-ethyl-1-methylprop-2-en-1-yl, 1-ethyl-2-methylprop-1-en-1-yl or 1-ethyl-2-methylprop-2-en-1-yl;

15

20

25

30
- C₃-C₆-alkynyl: prop-1-yn-1-yl, prop-2-yn-1-yl, n-but-1-yn-1-yl, n-but-1-yn-3-yl, n-but-1-yn-4-yl, n-but-2-yn-1-yl, n-pent-1-yn-1-yl, n-pent-1-yn-3-yl, n-pent-1-yn-4-yl, n-pent-1-yn-5-yl, n-pent-2-yn-1-yl, n-pent-2-yn-4-yl, n-pent-2-yn-5-yl, 3-methylbut-1-yn-3-yl, 3-methylbut-1-yn-4-yl, n-hex-1-yn-1-yl, n-hex-1-yn-3-yl, n-hex-1-yn-4-yl, n-hex-1-yn-5-yl, n-hex-1-yn-6-yl, n-hex-2-yn-1-yl, n-hex-2-yn-4-yl, n-hex-2-yn-5-yl, n-hex-2-yn-6-yl, n-hex-3-yn-1-yl, n-hex-3-yn-2-yl, 3-methylpent-1-yn-1-yl, 3-methylpent-1-yn-3-yl, 3-methylpent-1-yn-4-yl, 3-methylpent-1-yn-5-yl, 4-methylpent-1-yn-1-yl, 4-methylpent-2-yn-4-yl or 4-methylpent-2-yn-5-yl, preferably prop-2-yn-1-yl;

35

40

- C₃-C₇-cycloalkyl: a monocyclic saturated hydrocarbon ring having 3 to 7 ring members, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl;
- 5 - C₃-C₇-cycloalkenyl: monocyclic unsaturated hydrocarbon ring having 3 to 7 ring members, such as cycloprop-1-enyl, cycloprop-2-enyl, cyclobut-1-enyl, cyclobut-2-enyl, cyclobut-1,3-dienyl, cyclopent-1-enyl, cyclopent-2-enyl, cyclopent-3-enyl, cyclopent-2,4-dienyl, cyclohex-1-enyl, cyclohex-2-enyl, cyclohex-3-enyl; cyclohex-1,3-dienyl, cyclohex-1,5-dienyl, cyclohex-2,4-dienyl, or cyclohex-2,5-
- 10 dienyl.

If the phenyluracils I and/or the herbicides III are capable of forming geometrical isomers, for example E/Z isomers, it is possible to use both the pure isomers and compositions thereof in the compositions according to the invention. If the phenyluracils I and/or the herbicides III have one or more centers of chirality and, as a consequence, are present as enantiomers or diastereomers, it is possible to use both the pure enantiomers and diastereomers and their compositions in the compositions according to the invention.

20 If the phenyluracils and/or the herbicides III have functional groups, which can be ionized, they can also be used in the form of their agriculturally acceptable salts. Also, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide can be used in the form of its agriculturally acceptable salts. In general, the salts of those cations or anions are suitable whose cations or anions have no adverse effect on the action of the

25 active compounds ("agricultural acceptable").

Preferred cations are the ions of the alkali metals, preferably of lithium, sodium and potassium, of the alkaline earth metals, preferably of calcium and magnesium, and of the transition metals, preferably of manganese, copper, zinc and iron, furthermore ammonium and substituted ammonium in which one to four hydrogen atoms are replaced by C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkoxy-C₁-C₄-alkyl, phenyl or benzyl, preferably ammonium, methylammonium, isopropylammonium, dimethylammonium, diisopropylammonium, trimethylammonium, tetramethylammonium, tetraethylammonium, tetrabutylammonium, 2-

35 hydroxyethylammonium, 2-(2-hydroxyethoxy)eth-1-ylammonium, di(2-hydroxyeth-1-yl)ammonium, benzyltrimethylammonium, benzyltriethylammonium, furthermore phosphonium ions, sulfonium ions, preferably tri(C₁-C₄-alkyl)sulfonium such as trimethylsulfonium, and sulfoxonium ions, preferably tri(C₁-C₄-alkyl)sulfoxonium.

40 It is possible to use the phenyluracils of the formula I, the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and optionally at least one herbicide III selected from chlorazifop, clodinafop, clofop, cyhalofop, diclofop,

fenoxaprop, fenoxaprop-P, fenthiaprop, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, isoxapyrifop, propaquizafop, quizalofop, quizalofop-P, trifop, alloxymid, butoxydim, clethodim, cloproxydim, cycloxydim, profoxydim, sethoxydim, tepraloxymid, tralkoxydim, amidosulfuron, azimsulfuron, bensulfuron, chlorimuron, chloresulfuron, 5 cinosulfuron, cyclosulfamuron, ethametsulfuron, ethoxysulfuron, flazasulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, propoxycarbazon, flucarbazon, 10 imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriithiobac, flucetosulfuron, orthosulfamuron, pyrimisulfam, [N-(5,7-dimethoxy[1,2,4]triazolo[1,5-a]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3-pyridinsulfonamide, pyriminobac, bentazon, acifluorfen, ethoxyfen, fluoroglycofen, 15 fomesafen, halosafen, lactofen, pyraflufen, flumiclorac, fluthiacet, carfentrazone, flufenpyr, mesotrione, sulcotrione, topramezone, 4-hydroxy-3-[[2-methyl-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo-[3.2.1]oct-3-en-2-one, 4-hydroxy-3-[[2-(2-methoxyethoxy)methyl-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, 4-hydroxy-3-[4-(methylsulfonyl)-2-nitrobenzoyl]-bicyclo[3.2.1]oct-3-en-2-one, 2-[2-chloro-4-(methylsulfonyl)-3-[(2,2,2-trifluoroethoxy)methyl]benzoyl]-3-hydroxy-2- 20 cyclohexen-1-one, pyrasulfotole, glyphosate, glufosinate, bilanaphos, clomeprop, 2,4-D, 2,4-DB, dichlorprop, dichlorprop-P, MCPA, MCPB, mecoprop, mecoprop-P, 2,4,5-T, chloramben, dicamba, 2,3,6-TBA, tricamba, quinclorac, quinmerac, clopyralid, fluroxypyr, picloram, triclopyr, aminopyralid, naptalam, diflufenzopyr, cloquintocet, 25 fenchlorazole, isoxadifen and mefenpyr, in the form of salts with the agriculturally useful cations mentioned above.

Anions of useful acid addition salts are primarily chloride, bromide, fluoride, iodide, hydrogen sulfate, methyl sulfate, sulfate, dihydrogen phosphate, hydrogen phosphate, 30 nitrate, dicarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate and the anions of C₁-C₄-alkanoic acids, preferably formate, acetate, propionate and butyrate.

According to the invention, the herbicides cyperquat, diethamquat, difenzoquat, diquat, 35 morfamquat and paraquat are usually employed in the form of salts with the agriculturally useful anions mentioned above.

According to the invention, the active compounds which carry a carboxyl group can, instead of the active compounds mentioned above, also be employed in the form of an 40 agriculturally acceptable derivative, for example as amides such as mono- or di-C₁-C₆-alkylamides or arylamides, as esters, for example as allyl esters, propargyl esters, C₁-C₁₀-alkyl esters or alkoxyalkyl esters, and also as thioesters, for example as C₁-C₁₀-

alkyl thioesters. Examples of active compounds having a COOH group which can also be employed as derivatives are: chlorazifop, clodinafop, clofop, cyhalofop, diclofop, fenoxaprop, fenoxaprop-P, fenthiaprop, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, isoxapyrifop, propaquizafop, quizalofop, quizalofop-P, trifop, bensulfuron, chlorimuron, 5 ethametsulfuron, flupyrsulfuron, halosulfuron, iodosulfuron, mesosulfuron, metsulfuron, primisulfuron, pyrazosulfuron, sulfometuron, thifensulfuron, tribenuron, triflusulfuron, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, bispyribac, pyriithiobac, pyriminobac, acifluorfen, ethoxyfen, fluoroglycofen, lactofen, pyraflufen, flumiclorac, fluthiacet, carfentrazone, flufenpyr, 10 clomeprop, 2,4-D, 2,4-DB, dichlorprop, dichlorprop-P, MCPA, MCPB, mecoprop, mecoprop-P, 2,4,5-T, chloramben, dicamba, 2,3,6-TBA, tricamba, quinclorac, quinmerac, clopyralid, fluroxypyr, picloram, triclopyr, aminopyralid, naptalam, diflufenzopyr, cloquintocet, fenchlorazole, isoxadifen and mefenpyr.

Preferred mono- and di-C₁-C₆-alkylamides are the methyl- and the dimethylamides. 15 Preferred arylamides are, for example, the anilidines and the 2-chloroanilides. Preferred alkyl esters are, for example, the methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, mexyl (1-methylhexyl) or isooctyl (2-ethylhexyl) esters. Preferred C₁-C₄-alkoxy-C₁-C₄-alkyl esters are the straight-chain or branched C₁-C₄-alkoxyethyl esters, for example the methoxyethyl, ethoxyethyl or butoxyethyl esters. An example of 20 the straight-chain or branched C₁-C₁₀-alkyl thioesters is the ethyl thioester.

Among the 3-phenyluracils of formula I, preference is given to those wherein the variables R¹ to R⁷ independently of one another, but preferably combined, have the 25 meanings given below:

R¹ is methyl or NH₂;
 R² is trifluoromethyl;
 R³ is hydrogen, fluorine or chlorine,
 30 in particular fluorine;
 R⁴ is halogen or cyano,
 in particular chlorine or cyano;
 R⁵ is hydrogen;
 R⁶, R⁷ independently of one another are hydrogen, C₁-C₆-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkenyl, phenyl or benzyl;
 35 in particular hydrogen or C₁-C₆-alkyl.

R⁶ and R⁷ are in particular identical or different C₁-C₆-alkyl radicals.

40 In a particularly preferred embodiment of the invention, the compositions comprise at least one 3-phenyluracil I in which the variables R¹ to R⁷ in formula I have the following meanings (hereinbelow also referred to as 3-phenyluracils Ia):


- R¹ is methyl;
 R² is trifluoromethyl;
 R³ is fluorine;
 5 R⁴ is chlorine;
 R⁵ is hydrogen;
 R⁶, R⁷ independently of one another are C₁-C₆-alkyl.

10 In another particularly preferred embodiment of the invention, the compositions comprise at least one 3-phenyluracil I in which the variables R¹ to R⁷ in formula I have the meanings below (hereinbelow also referred to as 3-phenyluracils Ib):

- R¹ is NH₂;
 R² is trifluoromethyl;
 15 R³ is fluorine;
 R⁴ is chlorine;
 R⁵ is hydrogen;
 R⁶, R⁷ independently of one another are C₁-C₆-alkyl.

20

Particularly preferred herbicides I are 3-phenyluracils I.1, especially preferred 3-phenyluracils I.1.1 to I.1.74, listed below in table 1, wherein R¹, R⁶ and R⁷ have the meanings given in table 1:



25 Table 1

3-phenyluracil	R ¹	R ⁶	R ⁷
I.1.1	methyl	methyl	methyl
I.1.2	amino	methyl	methyl
I.1.3	methyl	methyl	ethyl
I.1.4	amino	methyl	ethyl
I.1.5	methyl	methyl	propyl
I.1.6	amino	methyl	propyl
I.1.7	methyl	methyl	isopropyl
I.1.8	amino	methyl	isopropyl
I.1.9	methyl	methyl	butyl
I.1.10	amino	methyl	butyl
I.1.11	methyl	methyl	s-butyl

3-phenyluracil	R ¹	R ⁶	R ⁷
I.1.12	amino	methyl	s-butyl
I.1.13	methyl	methyl	isobutyl
I.1.14	amino	methyl	isobutyl
I.1.15	methyl	methyl	t-butyl
I.1.16	amino	methyl	t-butyl
I.1.17	methyl	methyl	n-pentyl
I.1.18	amino	methyl	n-pentyl
I.1.19	methyl	methyl	n-hexyl
I.1.20	amino	methyl	n-hexyl
I.1.21	methyl	methyl	allyl
I.1.22	amino	methyl	allyl
I.1.23	methyl	methyl	propargyl
I.1.24	amino	methyl	propargyl
I.1.25	methyl	methyl	phenyl
I.1.26	amino	methyl	phenyl
I.1.27	methyl	methyl	benzyl
I.1.28	amino	methyl	benzyl
I.1.29	methyl	ethyl	ethyl
I.1.30	amino	ethyl	ethyl
I.1.31	methyl	ethyl	propyl
I.1.32	amino	ethyl	propyl
I.1.33	methyl	ethyl	isopropyl
I.1.34	amino	ethyl	isopropyl
I.1.35	methyl	ethyl	butyl
I.1.36	amino	ethyl	butyl
I.1.37	methyl	ethyl	n-pentyl
I.1.38	amino	ethyl	n-pentyl
I.1.39	methyl	ethyl	n-hexyl
I.1.40	amino	ethyl	n-hexyl
I.1.41	methyl	propyl	propyl
I.1.42	amino	propyl	propyl
I.1.43	methyl	propyl	isopropyl
I.1.44	amino	propyl	isopropyl
I.1.45	methyl	propyl	butyl
I.1.46	amino	propyl	butyl
I.1.47	methyl	propyl	n-pentyl
I.1.48	amino	propyl	n-pentyl
I.1.49	methyl	propyl	n-hexyl
I.1.50	amino	propyl	n-hexyl
I.1.51	methyl	isopropyl	isopropyl
I.1.52	amino	isopropyl	isopropyl

3-phenyluracil	R ¹	R ⁶	R ⁷
I.1.53	methyl	isopropyl	butyl
I.1.54	amino	isopropyl	butyl
I.1.55	methyl	isopropyl	n-pentyl
I.1.56	amino	isopropyl	n-pentyl
I.1.57	methyl	isopropyl	n-hexyl
I.1.58	amino	isopropyl	n-hexyl
I.1.59	methyl	butyl	butyl
I.1.60	amino	butyl	butyl
I.1.61	methyl	butyl	n-pentyl
I.1.62	amino	butyl	n-pentyl
I.1.63	methyl	butyl	n-hexyl
I.1.64	amino	butyl	n-hexyl
I.1.65	methyl	n-pentyl	n-pentyl
I.1.66	amino	n-pentyl	n-pentyl
I.1.67	methyl	n-pentyl	n-hexyl
I.1.68	amino	n-pentyl	n-hexyl
I.1.69	methyl	n-hexyl	n-hexyl
I.1.70	amino	n-hexyl	n-hexyl
I.1.71	methyl	-(CH ₂) ₄ -	
I.1.72	amino	-(CH ₂) ₄ -	
I.1.73	methyl	-(CH ₂) ₂ -O-(CH ₂) ₂ -	
I.1.74	amino	-(CH ₂) ₂ -O-(CH ₂) ₂ -	

In an especially preferred embodiment of the invention, the compositions comprise 3-phenyluracil I.1.7.

5

Among those compositions according to the present invention, preference is given to those which comprise at least one herbicide III selected from groups III.1 to III. 7, III.9 to III.11, III.13, III.14 and III.15; preferably in combination with a 3-phenyluracil Ia or Ib.

10

Among those compositions according to the present invention, preference is given to those which comprise at least one herbicide III selected from groups III.1, III.2, III.3, III.5, III.6, III.7, III.9, III.10, III.13, III.14 and III.15; in particular selected from groups III.1, III.2, III.3, III.5, III.13 and III.15; preferably in combination with a 3-phenyluracil Ia or Ib.

15

Preferred herbicides of groups III.1) to III.15) are the compounds listed below:

III.1) from the group of the lipid biosynthesis inhibitors:

- chlorazifop, clodinafop, clofop, cyhalofop, diclofop, fenoxaprop, fenoxaprop-p, fenthiaprop, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, isoxapyrifop, metamifop, propaquizafop, quizalofop, quizalofop-P, trifop, alloxydim, butroxydim, clethodim, cloproxydim, cycloxydim, profoxydim, sethoxydim, 5 tepraloxym, tralkoxydim, butylate, cycloate, diallate, dimepiperate, EPTC, esprocarb, ethiolate, isopolinate, methiobencarb, molinate, orbencarb, pebulate, prosulfocarb, sulfallate, thiobencarb, tiocarbazil, triallate, vernolate, benfuresate, ethofumesate, bensulide and pinoxaden;
- 10 III.2) from the group of the ALS inhibitors:
 amidosulfuron, azimsulfuron, bensulfuron, chlorimuron, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron, ethoxysulfuron, flazasulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, 15 pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, 20 pyriftalid, pyrithiobac, flucetosulfuron, orthosulfamuron, pyrimisulfan and [N-(5,7-dimethoxy[1,2,4]triazolo[1,5- α]pyrimidin-2-yl)-2-methoxy-4-(trifluormethyl)-3-pyridinesulfonamide, known from WO 02/36595; especially preferred amidosulfuron, azimsulfuron, bensulfuron, chlorimuron, chlorsulfuron, cinosulfuron, cyclosulfamuron, ethametsulfuron, ethoxysulfuron, 25 flazasulfuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamethabenz, imazamox, imazapic, imazapyr, 30 imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid, pyrithiobac, flucetosulfuron, orthosulfamuron and pyrimisulfan;
- 35 III.3) from the group of the photosynthesis inhibitors:
 atraton, atrazine, ametryne, aziprotryne, cyanazine, cyanatryn, chlorazine, cyprazine, desmetryne, dimethametryne, dipropetryn, eglazine, ipazine, mesoprazine, methometon, methoprotryne, procyazine, proglazine, prometon, prometryne, propazine, sebuthylazine, secbumeton, simazine, simeton, 40 simetryne, terbumeton, terbuthylazine, terbutryne, trietazine, ametrudione, amibuzin, hexazinone, isomethiozin, metamitron, metribuzin, bromacil, isocil, lenacil, terbacil, brompyrazon, chloridazon, dimidazon, desmedipham,

phenisopham, phenmedipham, phenmedipham-ethyl, benzthiazuron, buthiuron,
 ethidimuron, isouron, methabenzthiazuron, monoisouron, tebuthiuron,
 thiazafluron, anisuron, buturon, chlorbromuron, chloreturon, chlorotoluron,
 chloroxuron, difenoxuron, dimefuron, diuron, fenuron, fluometuron, fluothiuron,
 5 isoproturon, linuron, methiuron, metobenzuron, metobromuron, metoxuron,
 monolinuron, monuron, neburon, parafluron, phenobenzuron, siduron,
 tetrafluron, thidiazuron, cyperquat, diethamquat, difenzoquat, diquat,
 morfamquat, paraquat, bromobonil, bromoxynil, chloroxynil, iodobonil, ioxynil,
 amicarbazone, bromofenoxim, flumezin, methazole, bentazone, propanil,
 10 pentanochlor, pyridate, and pyridafol;

III.4) from the group of the protoporphyrinogen-IX oxidase inhibitors:
 acifluorfen, bifenox, chlomethoxyfen, chlornitrofen, ethoxyfen, fluorodifen,
 fluoroglycofen, fluoronitrofen, fomesafen, furyloxyfen, halosafen, lactofen,
 15 nitrofen, nitrofluorfen, oxyfluorfen, fluazolate, pyraflufen, cinidon-ethyl,
 flumiclorac, flumioxazin, flumipropyn, fluthiacet, thidiazimin, oxadiazon,
 oxadiargyl, azafenidin, carfentrazone, sulfentrazone, pentoxazone,
 benzfendizone, butafenacil, pyraclonil, profluazol, flufenpyr, flupropacil,
 nipyraclufen, etnipromid, bencarbazon and 3-{2-chloro-4-fluoro-5-[3-methyl-2,6-
 20 dioxo-4-(trifluoromethyl)-1,2,3,6-tetrahydropyrimidin-1-yl]phenoxy}-2-
 (ethoxycarbonyl)methoxypyridine (known from WO 06/61562);
 especially preferred acifluorfen, bifenox, chlomethoxyfen, chlornitrofen,
 ethoxyfen, fluorodifen, fluoroglycofen, fluoronitrofen, fomesafen, furyloxyfen,
 halosafen, lactofen, nitrofen, nitrofluorfen, oxyfluorfen, fluazolate, pyraflufen,
 25 cinidon-ethyl, flumiclorac, flumioxazin, flumipropyn, fluthiacet, thidiazimin,
 oxadiazon, oxadiargyl, azafenidin, carfentrazone, sulfentrazone, pentoxazone,
 benzfendizone, butafenacil, pyraclonil, profluazol, flufenpyr, flupropacil,
 nipyraclufen, etnipromid, and bencarbazon;

III.5) from the group of the bleacher herbicides:
 metflurazon, norflurazon, flufenican, diflufenican, picolinafen, beflubutamid,
 fluridone, flurochloridone, flurtamone, mesotrione, sulcotrione, isoxachlortole,
 isoxaflutole, benzofenap, pyrazolynate, pyrazoxyfen, benzobicyclon, amitrole,
 clomazone, aclonifen, 4-(3-trifluoromethyl- phenoxy)-2-(4-
 35 trifluoromethylphenyl)pyrimidine, known from EP 723960, topramezone, 4-
 hydroxy-3-[[2-methyl-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-
 en-2-one, known from WO 00/15615, 4-hydroxy-3-[[2-(2-methoxyethoxy)methyl-
 6-(trifluoro-methyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, known from
 WO 01/94339, 4-hydroxy-3-[4-(methylsulfonyl)-2-nitrobenzoyl]bicyclo[3.2.1]-oct-
 40 3-en-2-one, known from EP 338992, 2-[2-chloro-4-(methylsulfonyl)-3-[(2,2,2-
 trifluoroethoxy)methyl]benzoyl]-3-hydroxy-2-cyclohexen-1-one (known from DE
 19846792), and pyrasulfotole;

- III.6) from the group of the EPSP synthase inhibitors:
glyphosate;
- 5 III.7) from the group of the glutamine synthase inhibitors:
glufosinate and bilanaphos;
- III.8) from the group of the DHP synthase inhibitors:
asulam;
- 10 III.9) from the group of the mitose inhibitors:
benfluralin, butralin, dinitramine, ethalfluralin, fluchloralin, isopropalin,
methalpropalin, nitralin, oryzalin, pendimethalin, prodiamine, profluralin, trifluralin,
amiprofos-methyl, butamifos, dithiopyr, thiazopyr, propyzamide, tebutam,
15 chlorthal, carbetamide, chlorbufam, chlorpropham and propham;
- III.10) from the group of the VLCFA inhibitors:
acetochlor,alachlor, butachlor, butenachlor, delachlor, diethatyl, dimethachlor,
dimethenamid, dimethenamid-P, metazachlor, metolachlor, S-metolachlor,
20 pretilachlor, propachlor, propisochlor, prynachlor, terbuchlor, thenylchlor,
xylachlor, allidochlor, CDEA, epronaz, diphenamid, napropamide, naproanilide,
pethoxamid, flufenacet, mefenacet, fentrazamide, anilofos, piperophos,
cafenstrole, indanofan and tridiphane;
- 25 III.11) from the group of the cellulose biosynthesis inhibitors:
dichlobenil, chlorthiamid, isoxaben and flupoxam;
- III.12) from the group of the decoupler herbicides:
dinofenate, dinoprop, dinosam, dinoseb, dinoterb, DNOC, etinofen and
30 medinoterb;
- III.13) from the group of the auxin herbicides:
clomeprop, 2,4-D, 2,4,5-T, MCPA, MCPA thioethyl, dichlorprop, dichlorprop-P,
mecoprop, mecoprop-P, 2,4-DB, MCPB, chloramben, dicamba, 2,3,6-TBA,
35 tricamba, quinclorac, quinmerac, clopyralid, fluroxypyr, picloram, triclopyr,
benazolin and aminopyralid;
- III.14) from the group of the auxin transport inhibitors:
naptalam, diflufenzopyr;
- 40 III.15) benzoylprop, flamprop, flamprop-M, bromobutide, chlorflurenol, cinmethylin,
methyldymron, etobenzanid, fosamine, metam, pyributicarb, oxaziclomefone,

dazomet, triaziflam, methyl bromide;

and the agriculturally acceptable salts and the agriculturally acceptable derivatives of these herbicides, provided they have a carboxyl group.

5

Particularly preferred herbicides of groups III.1) to III.15) are the compounds listed below:

III.1) from the group of the lipid biosynthesis inhibitors:

10 clodinafop, cyhalofop, diclofop, fenoxaprop, fenoxaprop-p, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, metamifop, quizalofop, quizalofop-P, alloxydim, butoxydim, clethodim, cloproxydim, cycloxydim, profoxydim, sethoxydim, tepraloxym, tralkoxydim and pinoxaden;

15 III.2) from the group of the ALS inhibitors:

amidosulfuron, bensulfuron, chlorimuron, chlorsulfuron, cyclosulfamuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid, pyriothiobac, flucetosulfuron, orthosulfamuron, pyrimisulfan and [N-(5,7-

20 dimethoxy[1,2,4]triazolo[1,5- α]pyrimidin-2-yl-2-methoxy-4-(trifluoromethyl)-3-pyridinesulfonamide;

especially preferred amidosulfuron, bensulfuron, chlorimuron, chlorsulfuron, cyclosulfamuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid, pyriothiobac, flucetosulfuron, orthosulfamuron and pyrimisulfan;

30

35

III.3) from the group of the photosynthesis inhibitors:

40 atrazine, cyanazine, simazine, terbuthylazine, metamitron, metribuzin, chloridazon, amicarbazone, chlorotoluron, diuron, isoproturon, methabenzthiazuron, bentazone, propanil, bromoxynil, ioxynil and paraquat;

- III.5) from the group of the bleacher herbicides:
norflurazon, diflufenican, picolinafen, beflubutamid, mesotrione, sulcotrione,
isoxaflutole, benzofenap, pyrazolynate, pyrazoxyfen, benzobicyclon, 4-(3-
trifluoromethylphenoxy)-2-(4-trifluoromethylphenyl)pyrimidine, topramezone, 4-
5 hydroxy-3-[[2-methyl-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-
en-2-one, 4-hydroxy-3-[[2-(2-methoxyethoxy)methyl-6-(trifluoro-methyl)-3-
pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, 4-hydroxy-3-[4-(methylsulfonyl)-2-
nitrobenzoyl]bicyclo[3.2.1]-oct-3-en-2-one, 2-[2-chloro-4-(methylsulfonyl)-3-
10 [(2,2,2-trifluoroethoxy)methyl]benzoyl]-3-hydroxy-2-cyclohexen-1-one and
pyrasulfotole;
- III.6) from the group of the EPSP synthase inhibitors:
glyphosate;
- 15 III.7) from the group of the glutamine synthase inhibitors:
glufosinate;
- III.9) from the group of the mitose inhibitors:
oryzalin, pendimethalin and trifluralin;
- 20 III.10) from the group of the VLCFA inhibitors:
acetochlor, butachlor, dimethenamid, dimethenamid-P, metazachlor,
metolachlor, S-metolachlor, pethoxamid, pretilachlor, flufenacet, mefenacet,
fentrazamide, cafenstrole and indanofan;
- 25 III.13) from the group of the auxin herbicides:
2,4-D, dichlorprop, dichlorprop-P, mecoprop, MCPA, mecoprop-P, dicamba,
quinclorac, quinmerac, clopyralid, fluroxypyr, picloram, triclopyr and
aminopyralid;
- 30 III.14) from the group of the auxin transport inhibitors:
diflufenzopyr;
- III.15) cinmethylin, oxaziclomefone and triaziflam;
- 35 and the agriculturally acceptable salts and the agriculturally acceptable derivatives of
these herbicides, provided they have a carboxyl group.

The categorization of the herbicides III according to their mode of action is based on
40 current understanding. If a herbicide acts by more than one mode of action, this
substance was assigned to only one mode of action.

Particular preference is given to those binary and ternary compositions which comprise at least one 3-phenyluracil I and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and, if appropriate, one or more herbicides III.

5

Here and below, the term "binary compositions" includes compositions which comprise one or more (for example 2 or 3) 3-phenyluracils I and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II.

10 Correspondingly, the term "ternary compositions" includes compositions which comprise one or more (for example 2 or 3) 3-phenyluracils I, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and one or more (for example 2 or 3) herbicides III.

15 In binary compositions the weight ratio of the active compounds I : II is usually in the range from 1:10 to 10:1, preferably in the range from 1:5 to 5:1, in particular in the range from 1:3 to 3:1.

20 In ternary compositions which comprise both a 3-phenyluracil I, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and at least one herbicide III, the relative weight ratios of the components I : II : III are usually in the range from 10:1:1 to 1:10:20, preferably from 5:1:1 to 1:5:10, in particular from 3:1:1 to 1:3:5.

25 In these ternary compositions, the weight ratio of N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II to herbicide III is preferably in the range from 10:1 to 1:20.

30 In a particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II.

35 Preference is also given to those compositions of the invention comprising as the only herbicidal active compound a 3-phenyluracil I,
preferably phenyluracil Ia or Ib,
particularly preferred phenyluracil I selected from the group consisting of 3-phenyluracils I.1.1 to I.1.74,
especially preferred 3-phenyluracil I.1.7,
40 in combination with the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II.

Preference is also given to those compositions of the invention comprising as the only herbicidal active compound phenyluracil Ia in combination with the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II.

5

Furthermore preference is given to those compositions of the invention comprising a 3-phenyluracil I,

preferably phenyluracil Ia or Ib,

particularly preferred phenyluracil I selected from the group consisting of 3-

10

phenyluracils I.1.1 to I.1.74,

especially preferred 3-phenyluracil I.1.7,

in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and

at least one, especially exactly one, herbicide III selected from the classes III.1) to

15

III.15).

Preference is also given to those compositions of the invention comprising 3-phenyluracil Ia, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and at least one, especially exactly one, herbicide III selected from the classes III.1) to III.15).

20

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.1), in particular selected from the group consisting of clodinafop, cyhalofop, diclofop, fenoxaprop, fenoxaprop-p, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, metamifop, quizalofop, quizalofop-P, alloxydim, butoxydim, clethodim, cloproxydim, cycloxydim, profoxydim, sethoxydim, tepraloxydim, tralkoxydim and pinoxaden.

25

30

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.2), in particular selected from the group consisting of amidosulfuron, bensulfuron, chlorimuron, chlorsulfuron, cyclosulfamuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin,

35

40

imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid, pyriothiobac, flucetosulfuron, orthosulfamuron, pyrimisulfan and [N-(5,7-dimethoxy[1,2,4]triazolo[1,5- α]pyrimidin-2-yl-2-methoxy-4-(trifluoromethyl)-3-pyridinesulfonamide.

In a further particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.2), in particular selected from the group consisting of amidosulfuron, bensulfuron, chlorimuron, chlorsulfuron, cyclosulfamuron, flupyrsulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron, tritosulfuron, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid, pyriothiobac, flucetosulfuron, orthosulfamuron and pyrimisulfan.

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.3), in particular selected from the group consisting of atrazine, cyanazine, simazine, terbuthylazine, metamitron, metribuzin, chloridazon, amicarbazone, chlorotoluron, diuron, isoproturon, methabenzthiazuron, bentazone, propanil, pyridate, bromoxynil, ioxynil and paraquat.

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.5), in particular selected from the group consisting of diflufenican, picolinafen, mesotrione, sulcotrione, isoxaflutole, benzofenap, pyrazolynate, pyrazoxyfen, benzobicyclon, 4-(3-trifluoromethylphenoxy)-2-(4-trifluoromethylphenyl)pyrimidine, topramezone, 4-hydroxy-3-[[2-methyl-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, 4-hydroxy-3-[[2-(2-methoxyethoxy)methyl-6-(trifluoro-methyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, 4-hydroxy-3-[4-(methylsulfonyl)-2-

nitrobenzoyl]bicyclo[3.2.1]-oct-3-en-2-one, 2-[2-chloro-4-(methylsulfonyl)-3-[(2,2,2-trifluoroethoxy)methyl]benzoyl]-3-hydroxy-2-cyclohexen-1-one and pyrasulfotole.

5 In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.6), in particular glyphosate.

10

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.7), in particular glufosinate.

15

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.9), in particular oryzalin, pendimethalin and trifluralin.

20

25 In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.10), in particular selected from the group consisting of acetochlor, butachlor, dimethenamid, dimethenamid-P, metazachlor, metolachlor, S-metolachlor, pethoxamid, pretilachlor, flufenacet, mefenacet, fentrazamide, cafenstrole and indanofan.

30

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.13), in particular selected from the group consisting of 2,4-D, dichlorprop, dichlorprop-P, mecoprop, MCPA, mecoprop-P, dicamba, quinclorac, quinmerac, clopyralid, fluroxypyr, picloram, triclopyr and aminopyralid.

35

40

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.14), in particular diflufenzopyr.

In another particularly preferred embodiment of the invention, preference is given to those compositions of the invention which comprise a 3-phenyluracil I, especially Ia or Ib, preferably phenyluracil Ia, particularly preferred phenyluracil I.1.7, in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and at least one, especially exactly one, herbicidally active compound of the group III.15), in particular selected from the group consisting of cinmethylin, oxaziclomefone and triaziflam.

In the preferred or especially preferred compositions described above the 3-phenyluracils I, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and the herbicides III can be used in the form of their agriculturally acceptable salts or in the form of an agriculturally acceptable derivative thereof as described above.

The weight ratios of the individual components in the compositions are within the limits stated above.

Among the especially preferred compositions, particular preference is given to those compositions of the invention wherein the variables R¹ to R⁷ have the preferred meanings, especially the particularly preferred meanings. Particular preference is given to 3-phenyluracils Ia and Ib, very particular to 3-phenyluracils I.1.1 to I.1.74, as defined above.

Also particular preference is given to 3-phenyluracils Ia, especially to 3-phenyluracil I.1.7, as defined above.

Preference is also given, for example, to those compositions which comprise, as active compound I, the phenyluracil I.1.1 and, as further active compound, the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and, as further active compound, the substances listed in one row of table 2 (compositions 1.1 to 1.173).

The weight ratios of the individual components in the compositions 1.1 to 1.173 are within the stated limits, in the case of binary compositions of phenyluracil I.1.1 and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II for example 1:1, and in the case of ternary compositions of phenyluracil I.1.1, N-[[4-

[(cyclopropylamino)carbonyl]phenyl)sulfonyl]-2-methoxybenzamide II and herbicide III for example 1:1:1, 2:1:1, 1:2:1, 1:5:1 or 1:5:2.

5 Table 2

composition no.	herbicide III
1.1	clodinafop
1.2	cyhalofop
1.3	diclofop
1.4	fenoxaprop
1.5	fenoxaprop-P
1.6	fluazifop
1.7	fluazifop-P
1.8	haloxyfop
1.9	haloxyfop-P
1.10	metamifop
1.11	quizalofop
1.12	quizalofop-P
1.13	alloxydim
1.14	butroxydim
1.15	clethodim
1.16	cloproxydim
1.17	cycloxydim
1.18	profoxydim
1.19	sethoxydim
1.20	tepraloxym
1.21	tralkoxydim
1.22	pinoxaden
1.23	amidosulfuron
1.24	azimsulfuron
1.25	bensulfuron
1.26	chlorimuron
1.27	chlorsulfuron
1.28	cinosulfuron
1.29	cyclosulfamuron
1.30	ethametsulfuron
1.31	ethoxysulfuron
1.32	flazasulfuron
1.33	flupyrsulfuron
1.34	foramsulfuron
1.35	halosulfuron

composition no.	herbicide III
1.36	imazosulfuron
1.37	iodosulfuron
1.38	mesosulfuron
1.39	metsulfuron
1.40	nicosulfuron
1.41	oxasulfuron
1.42	primisulfuron
1.43	prosulfuron
1.44	pyrazosulfuron
1.45	rimsulfuron
1.46	sulfometuron
1.47	sulfosulfuron
1.48	thifensulfuron
1.49	triasulfuron
1.50	tribenuron
1.51	trifloxysulfuron
1.52	triflusulfuron
1.53	tritosulfuron
1.54	imazamethabenz
1.55	imazamox
1.56	imazapic
1.57	imazapyr
1.58	imazaquin
1.59	imazethapyr
1.60	cloransulam
1.61	diclosulam
1.62	florasulam
1.63	flumetsulam
1.64	metosulam
1.65	penoxsulam
1.66	bispyribac
1.67	pyriminobac
1.68	propoxycarbazone
1.69	flucarbazone
1.70	pyribenzoxim
1.71	pyriftalid
1.72	pyrithiobac
1.73	flucetosulfuron
1.74	orthosulfamuron
1.75	pyrimisulfan

composition no.	herbicide III
1.76	[N-(5,7-dimethoxy[1,2,4]triazolo[1,5- α]pyrimidin-2-yl)-2-methoxy-4-(trifluormethyl)-3-pyridinesulfonamide
1.77	atrazine
1.78	cyanazine
1.79	simazine
1.80	terbuthylazine
1.81	metamitron
1.82	metribuzin
1.83	amicarbazone
1.84	chloridazon
1.85	chlorotoluron
1.86	diuron
1.87	isoproturon
1.88	methabenzthiazuron
1.89	propanil
1.90	bromoxynil
1.91	ioxynil
1.92	bentazone
1.93	pyridate
1.94	paraquat
1.95	acifluorfen
1.96	fluoroglycofen
1.97	halosafen
1.98	lactofen
1.99	oxyfluorfen
1.100	fluazolate
1.101	pyraflufen
1.102	cinidon-ethyl
1.103	flumiclorac
1.104	flumioxazin
1.105	fluthiacet
1.106	oxadiazon
1.107	oxadiargyl
1.108	azafenidin
1.109	carfentrazone
1.110	sulfentrazone
1.111	pentoxazone
1.112	benzfendizone
1.113	butafenacil
1.114	pyraclonil

composition no.	herbicide III
1.115	profluazol
1.116	flufenpyr
1.117	nipyraclofen
1.118	bencarbazone
1.119	norflurazon
1.120	diflufenican
1.121	picolinafen
1.122	beflubutamid
1.123	mesotrione
1.124	sulcotrione
1.125	isoxaflutole
1.126	benzofenap
1.127	pyrazolynate
1.128	pyrazoxyfen
1.129	benzobicyclon
1.130	4-(3-trifluoromethylphenoxy)-2-(4-trifluoromethylphenyl)pyrimidine
1.131	topramezone
1.132	4-hydroxy-3-[[2-methyl-6-(trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one
1.133	4-hydroxy-3-[[2-(2-methoxyethoxy)methyl-6-(trifluoro-methyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one
1.134	4-hydroxy-3-[4-(methylsulfonyl)-2-nitrobenzoyl]bicyclo[3.2.1]-oct-3-en-2-one
1.135	2-[2-chloro-4-(methylsulfonyl)-3-[(2,2,2-trifluoroethoxy)methyl]benzoyl]-3-hydroxy-2-cyclohexen-1-one
1.136	pyrasulfotole
1.137	glyphosate
1.138	glufosinate
1.139	oryzalin
1.140	pendimethalin
1.141	trifluralin
1.142	acetochlor
1.143	butachlor
1.144	dimethenamid
1.145	dimethenamid-P
1.146	metazachlor
1.147	metolachlor
1.148	S-metolachlor
1.149	pethoxamid
1.150	pretilachlor

composition no.	herbicide III
1.151	flufenacet
1.152	mefenacet
1.153	fentrazamide
1.154	cafenstrole
1.155	indanofan
1.156	2,4-D
1.157	dichlorprop
1.158	dichlorprop-P
1.159	MCPA
1.160	mecoprop
1.161	mecoprop-P
1.162	dicamba
1.163	quinclorac
1.164	quinmerac
1.165	clopyralid
1.166	fluroxypyr
1.167	picloram
1.168	triclopyr
1.169	aminopyralid
1.170	diflufenzopyr
1.171	cinmethylin
1.172	oxaziclomefone
1.173	triaziflam

Preference is also given to the compositions 2.1 – 2.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.2.

5

Preference is also given to the compositions 3.1 – 3.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.3.

10

Preference is also given to the compositions 4.1 – 4.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.4.

15

Preference is also given to the compositions 5.1 – 5.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.5.

Preference is also given to the compositions 6.1 – 6.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.6.

- 5 Preference is also given to the compositions 7.1 – 7.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.7.

- 10 Preference is also given to the compositions 8.1 – 8.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.8.

- 15 Preference is also given to the compositions 9.1 – 9.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.9.

- 20 Preference is also given to the compositions 10.1 – 10.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.10.

- 25 Preference is also given to the compositions 11.1 – 11.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.11.

- 30 Preference is also given to the compositions 12.1 – 12.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.12.

- 35 Preference is also given to the compositions 13.1 – 13.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.13.

- 40 Preference is also given to the compositions 14.1 – 14.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I. 1.14.

Preference is also given to the compositions 15.1 – 15.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.15.

Preference is also given to the compositions 16.1 – 16.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.16.

- 5 Preference is also given to the compositions 17.1 – 17.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.17.

- 10 Preference is also given to the compositions 18.1 – 18.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.18.

- 15 Preference is also given to the compositions 19.1 – 19.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.19.

- 20 Preference is also given to the compositions 20.1 – 20.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.20.

- 25 Preference is also given to the compositions 21.1 – 21.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.21.

- 30 Preference is also given to the compositions 22.1 – 22.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.22.

- 35 Preference is also given to the compositions 23.1 – 23.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.23.

- 40 Preference is also given to the compositions 24.1 – 24.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.24.

Preference is also given to the compositions 25.1 – 25.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.25.

Preference is also given to the compositions 26.1 – 26.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.26.

- 5 Preference is also given to the compositions 27.1 – 27.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.27.

- 10 Preference is also given to the compositions 28.1 – 28.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.28.

- 15 Preference is also given to the compositions 29.1 – 29.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.29.

- 20 Preference is also given to the compositions 30.1 – 30.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.30.

- 25 Preference is also given to the compositions 31.1 – 31.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.31.

- 30 Preference is also given to the compositions 32.1 – 32.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.32.

- 35 Preference is also given to the compositions 33.1 – 33.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.33.

- 40 Preference is also given to the compositions 34.1 – 34.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.34.

Preference is also given to the compositions 35.1 – 35.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.35.

Preference is also given to the compositions 36.1 – 36.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.36.

- 5 Preference is also given to the compositions 37.1 – 37.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.37.

- 10 Preference is also given to the compositions 38.1 – 38.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.38.

- 15 Preference is also given to the compositions 39.1 – 39.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.39.

- Preference is also given to the compositions 40.1 – 40.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.40.

20

Preference is also given to the compositions 41.1 – 41.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.41.

- 25 Preference is also given to the compositions 42.1 – 42.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.42.

- 30 Preference is also given to the compositions 43.1 – 43.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.43.

- 35 Preference is also given to the compositions 44.1 – 44.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.44.

- Preference is also given to the compositions 45.1 – 45.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.45.

40

Preference is also given to the compositions 46.1 – 46.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.46.

- 5 Preference is also given to the compositions 47.1 – 47.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.47.

- 10 Preference is also given to the compositions 48.1 – 48.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.48.

- 15 Preference is also given to the compositions 49.1 – 49.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.49.

- 20 Preference is also given to the compositions 50.1 – 50.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.50.

- 25 Preference is also given to the compositions 51.1 – 51.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.51.

- 30 Preference is also given to the compositions 52.1 – 52.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.52.

- 35 Preference is also given to the compositions 53.1 – 53.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.53.

- 40 Preference is also given to the compositions 54.1 – 54.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.54.

Preference is also given to the compositions 55.1 – 55.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.55.

Preference is also given to the compositions 56.1 – 56.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.56.

- 5 Preference is also given to the compositions 57.1 – 57.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.57.

- 10 Preference is also given to the compositions 58.1 – 58.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.58.

- 15 Preference is also given to the compositions 59.1 – 59.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.59.

- Preference is also given to the compositions 60.1 – 60.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.60.

20

Preference is also given to the compositions 61.1 – 61.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.61.

- 25 Preference is also given to the compositions 62.1 – 62.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.62.

- 30 Preference is also given to the compositions 63.1 – 63.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.63.

- 35 Preference is also given to the compositions 64.1 – 64.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.64.

- Preference is also given to the compositions 65.1 – 65.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.65.

40

Preference is also given to the compositions 66.1 – 66.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.66.

- 5 Preference is also given to the compositions 67.1 – 67.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.67.

- 10 Preference is also given to the compositions 68.1 – 68.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.68.

- 15 Preference is also given to the compositions 69.1 – 69.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.69.

- 20 Preference is also given to the compositions 70.1 – 70.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.70.

- 25 Preference is also given to the compositions 71.1 – 71.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.71.

- 30 Preference is also given to the compositions 72.1 – 72.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.72.

- 35 Preference is also given to the compositions 73.1 – 73.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.73.

- 40 Preference is also given to the compositions 74.1 – 74.173, which differ from the corresponding compositions 1.1 – 1.173 only in that the phenyluracil I.1.1 is replaced by the phenyluracil I.1.74.

- 45 Preference is also given to those compositions, which comprise, as active compound, the substances listed in one row of table 1 (3-phenyluracil I.1.1 – I.1.74), and, as further active compound, the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, and, as further active compound, 3-{2-chloro-4-fluoro-5-[3-methyl-2,6-dioxo-4-(trifluoromethyl)-1,2,3,6-tetrahydropyrimidin-1-yl]phenoxy}-2-(ethoxycarbonyl)methoxypyridine.

The weight ratios of the individual components in the compositions 1.1 to 74.173 are within the limits stated above, in the case of binary compositions of 3-phenyluracil I and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II for example 5 1:1, 1:2 or 1:5, and in the case of ternary compositions of 3-phenyluracil I, N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and herbicide III for example 1:1:1, 2:1:1, 1:2:1, 1:5:1 or 1:5:2.

10 In the ready-to-use preparations, i.e. in the compositions according to the invention in the form of crop protection products, the components I and II and optionally III, in suspended, emulsified or dissolved form, can be present formulated jointly or separately. The use forms depend entirely on the intended use.

15 The compositions according to the invention can be applied, for example, in the form of directly sprayable aqueous solutions, powders, suspensions, also highly-concentrated aqueous, oily or other suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading or granules, by means of spraying, atomizing, dusting, broadcasting or watering. The use forms depend on the intended use; in any case, they should ensure the finest possible distribution of the active compounds.

20 Depending on the form in which the ready-to-use preparations are present in the compositions according to the invention, they comprise one or more liquid or solid carriers, if appropriate surfactants and if appropriate further auxiliaries which are customary for formulating crop protection products. The person skilled in the art is 25 sufficiently familiar with the recipes for such formulations.

The ready-to-use preparations comprise the components I and II and optionally III and auxiliaries which are customary for formulating crop protection products, which auxiliaries may also comprise a liquid carrier.

30 Suitable inert additives with carrier function are essentially: mineral oil fractions of medium to high boiling point, such as kerosene and diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, e.g. paraffins, tetrahydronaphthalene, alkylated naphthalenes and their derivatives, 35 alkylated benzenes and their derivatives, alcohols such as methanol, ethanol, propanol, butanol and cyclohexanol, ketones such as cyclohexanone, strongly polar solvents, e.g. amines such as N-methylpyrrolidone, and water.

40 Aqueous use forms can be prepared from emulsion concentrates, suspensions, pastes, wettable powders or water-dispersible granules by adding water. To prepare emulsions, pastes or oil dispersions, the active compounds I, II or III, as such or dissolved in an oil or solvent, can be homogenized in water by means of wetting agent,

tackifier, dispersant or emulsifier. Alternatively, it is possible to prepare concentrates consisting of active substance, wetting agent, tackifier, dispersant or emulsifier and, if desired, solvent or oil, and these concentrates are suitable for dilution with water.

5 Suitable surfactants are the alkali metal salts, alkaline earth metal salts and ammonium salts of aromatic sulfonic acids, e.g. ligno-, phenol-, naphthalene- and
dibutyl-naphthalenesulfonic acid, and of fatty acids, of alkyl- and alkylarylsulfonates, of
alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates, and salts of sulfated hexa-
10 , hepta- and octadecanols and of fatty alcohol glycol ethers, condensates of sulfonated
naphthalene and its derivatives with formaldehyde, condensates of naphthalene or of
the naphthalenesulfonic acids with phenol and formaldehyde, polyoxyethylene
octylphenol ether, ethoxylated isooctyl-, octyl- or nonylphenol, alkylphenyl polyglycol
ethers, tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol,
fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl
15 ether or polyoxypropylene alkyl ether, lauryl alcohol polyglycol ether acetate, sorbitol
esters, liginosulfite waste liquors or methylcellulose.

Powders, materials for spreading and dusts can be prepared by mixing or concomitant grinding of the active substances with a solid carrier.

20 Granules, e.g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients to solid carriers. Solid carriers are mineral earths such as silicas, silica gels, silicates, talc, kaolin, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate,
25 magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders, or other solid carriers.

30 The concentrations of the active compounds in the ready-to-use preparations can be varied within wide ranges. In general, the formulations comprise from 0.001 to 98% by weight, preferably 0.01 to 95% by weight, of active ingredients. The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

35 The compounds according to the invention can, for example, be formulated as follows:

I 20 parts by weight of the active compound composition in question are dissolved in a composition composed of 80 parts by weight of alkylated benzene, 10 parts
40 by weight of the adduct of 8 to 10 mol of ethylene oxide to 1 mol of oleic acid N-monoethanolamide, 5 parts by weight of calcium dodecylbenzenesulfonate and 5 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil.

Pouring the solution into 100 000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.

- 5 II 20 parts by weight of the active compound composition in question are dissolved in a composition composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 mol of ethylene oxide to 1 mol of isooctylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring the solution into 100 000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.
- 10
- III 20 parts by weight of the active compound composition in question are dissolved in a composition composed of 25 parts by weight of cyclohexanone, 65 parts by weight of a mineral oil fraction of boiling point 210 to 280°C and 10 parts by weight of the adduct of 40 mol of ethylene oxide to 1 mol of castor oil. Pouring the solution into 100 000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.
- 15
- IV 20 parts by weight of the active compound composition in question are mixed thoroughly with 3 parts by weight of sodium diisobutyl naphthalenesulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel, and the composition is ground in a hammer mill. Finely distributing the composition in 20 000 parts by weight of water gives a spray composition which comprises 0.1% by weight of the active ingredient.
- 20
- 25
- V 3 parts by weight of the active compound composition in question are mixed with 97 parts by weight of finely divided kaolin. This gives a dust which comprises 3% by weight of the active ingredient.
- 30
- VI 20 parts by weight of the active compound composition in question are mixed intimately with 2 parts by weight of calcium dodecylbenzenesulfonate, 8 parts by weight of fatty alcohol polyglycol ether, 2 parts by weight of the sodium salt of a phenol-urea-formaldehyde condensate and 68 parts by weight of a paraffinic mineral oil. This gives a stable oily dispersion.
- 35
- VII 1 part by weight of the active compound composition in question is dissolved in a composition composed of 70 parts by weight of cyclohexanone, 20 parts by weight of ethoxylated isooctylphenol and 10 parts by weight of ethoxylated castor oil. This gives a stable emulsion concentrate.
- 40

VIII 1 part by weight of the active compound composition in question is dissolved in a composition composed of 80 parts by weight of cyclohexanone and 20 parts by weight of Wettol® EM 31 (nonionic emulsifier based on ethoxylated castor oil).
5 This gives a stable emulsion concentrate.

The components I and II and optionally III can be formulated jointly or separately.

10 The components I and II and optionally III can be applied jointly or separately, simultaneously or successively, before, during or after emergence of the plants.

If the active compounds I and II and optionally III are less well tolerated by certain crop plants, it is possible to use application methods in which the herbicidal compositions are sprayed with the aid of sprayers in such a way that the leaves of the sensitive crop
15 plants are as far as possible unaffected, whereas the active compounds reach the leaves of the undesirable plants growing underneath or the uncovered soil surface (post-directed, lay-by).

The required application rate of the composition of the pure active compounds, i.e. of I
20 and II and optionally III without formulation auxiliary, depends on the density of the undesired vegetation, on the development stage of the plants, on the climatic conditions of the location where the composition is used and on the application method. In general, the application rate of I and II and optionally III is from 0.001 to 3 kg/ha, preferably from 0.005 to 2 kg/ha and in particular from 0.01 to 1 kg/ha of active
25 substance.

The required application rates of the 3-phenyluracils I and N-[[4-[(cyclopropylamino)-
30 carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II are generally in the range from 0.1 g/ha to 1 kg/ha and preferably in the range from 1 g/ha to 500 g/ha or from 5 g/ha to 500 g/ha of active substance.

The compositions are applied to the plants mainly by spraying, in particular foliar
35 spraying. Application can be carried out by customary spraying techniques using, for example, water as carrier and spray liquor rates of from about 50 to 1000 l/ha (for example from 300 to 400 l/ha). Application of the herbicidal compositions by the low-volume and the ultra-low-volume method is possible, as is their application in the form of microgranules.

40 The compositions according to the present invention are suitable for controlling common harmful plants in useful plants, in particular in crops such as oat, barley, millet, corn, rice, wheat, sugar cane, cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers and soybeans or in perennial crops.

Moreover, it may be useful to apply the compositions according to the invention jointly as a composition with other crop protection products, for example with pesticides or agents for controlling phytopathogenic fungi or bacteria. Also of interest is the miscibility with mineral salt solutions which are employed for treating nutritional and trace element deficiencies. Non-phytotoxic oils and oil concentrates may also be added.

The compositions according to the invention can also be used in crop plants which, by genetic engineering or by breeding, are resistant to one or more herbicides and/ or fungicides and/or or to attack by insects.

Suitable are for example crop plants preferably corn, wheat, barley, sunflower, rice, canola, soybeans which are resistant to herbicidal EPSP synthase inhibitors, such as, for example, glyphosate, to herbicidal glutamine synthase inhibitors, such as, for example, glufosinate, to herbicidal protoporphyrinogen-IX oxidase inhibitors, such as, for example, butafenacil, or to herbicidal ALS inhibitors, such as, for example, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, or crop plants which, owing to introduction of the gene for Bt toxin by genetic modification, are resistant to attack by certain insects.

Surprisingly, the compositions according to the invention which, in addition to the 3-phenyluracil I and optionally a herbicide III, comprise the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II, are better tolerated by useful plants than the respective compositions comprising a 3-phenyluracil I and optionally a herbicide III without safener II.

The 3-phenyluracils of formula I can be prepared by the preparation processes disclosed in WO 01/83459. With respect to the preparation of individual compounds, reference is made to the examples of WO 01/83459. Compounds, which are not explicitly disclosed in this document, can be prepared in an analogous manner.

N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II can be prepared by the preparation processes disclosed by the earlier applications WO 99/16744 and WO 05/00797.

Use Examples

The effect of the herbicidal compositions according to the invention of components I and II and, if appropriate, III on the growth of undesirable plants compared to the herbicidally active compounds alone can be demonstrated by the following greenhouse experiments.

For the pre-emergence treatment, directly after sowing the active compounds, which had been suspended or emulsified in water, were applied by means of finely distributed nozzles. The containers were irrigated gently to promote germination and growth and subsequently covered with transparent plastic hoods until plant had rooted. This cover
 5 caused uniform germination of the tests plants, unless this was adversely affected by active compounds.

For the post-emergence treatment, the test plants were first grown to a height of 3 to 20 cm, depending on the plant habit, and only then treated. Here, the herbicidal
 10 compositions were suspended or emulsified in water as distribution medium and sprayed using finely distributing nozzles.

The respective component I was formulated as a 12% by weight strength emulsion concentrate, and component II as a 10% by weight suspension concentrate. Optionally
 15 components III can be used in their appropriate available formulation types.

The components I, II and optionally III were introduced to the spray liquid with the amount of solvent system used for applying the active compound.

20 The test period extended over 21 days. During this time, the plants were tended, and their response to the treatments with active compound was evaluated.

The evaluation for the damage caused by the chemical compositions was carried out using a scale from 0 to 100%, compared to the untreated control plants. Here, 0 means
 25 no damage and 100 means complete destruction of the plants.

A safener action is present if the damage to the crop plant caused by a mixture according to the present invention which contains N-[[4-
 30 [(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide is less than the damage caused when the phenyluracil I optionally in mixture with a herbicide III according to the present invention is used without N-[[4-
 [(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide.

35 The plants used in these greenhouse experiments belong to the following species:

BAYER-Code	Common name
TRZAS	spring wheat
HORVW	winter barley
ZEAMX	corn
POLCO	wild buckwheat
CHEAL	lambsquaters

ABUTH AMARE	velvetleaf common amaranth
----------------	-------------------------------


- Example 1: Herbicidal action of the post-emergence applied mixture of 3-phenyluracil I.1.7 and N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II against POLCO, CHEAL, ABUTH and AMARE and safener action with regard to TRZAS, HORVW and ZEAMX

Application rate in g/ha		Damage to crop plant			Herbicidal action against			
I.1.7	II	TRZAS	HORVW	ZEAMX	POLCO	CHEAL	ABUTH	AMARE
6.25	--	35	40	50	100	100	100	100
6.25	150	20	35	35	100	100	100	100
3.125	--	20	35	35	100	100	100	100
3.125	150	10	20	30	100	100	100	100

- 10 The safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II has not shown damaging effect on TRZAS, HORVW and ZEAMX and has not shown herbicidal activity on POLCO, CHEAL, ABUTH and AMARE at application rates of 150 g/ha applied post-emergence.

CLAIMS

1. A composition comprising a 3-phenyluracil I




I

wherein the variables R^1 to R^7 are as defined below:

- R^1 is methyl or NH_2 ;
 R^2 is C_1 - C_2 -haloalkyl;
 R^3 is hydrogen or halogen;
 R^4 is halogen or cyano;
 R^5 is hydrogen or C_1 - C_6 -alkyl;
 R^6, R^7 independently of one another are hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -cycloalkenyl, phenyl or benzyl;

including its agriculturally acceptable salts; and

N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II



II,

including its agriculturally acceptable salts; and

optionally at least one herbicide III selected from the group consisting of the following classes III.1) to

III.15):

III.1) lipid biosynthesis inhibitors;

5 III.2) acetolactate synthase inhibitors (ALS inhibitors);

III.3) photosynthesis inhibitors;

III.4) protoporphyrinogen-IX oxidase inhibitors;

III.5) bleacher herbicides;

III.6) enolpyruvyl shikimate 3-phosphate synthase inhibitors (EPSP inhibitors);

10 III.7) glutamine synthetase inhibitors;

III.8) 7,8-dihydropteroate synthase inhibitors (DHP inhibitors);

III.9) mitose inhibitors;

III.10) inhibitors of the synthesis of long chain fatty acids (VLCFA inhibitors);

III.11) cellulose biosynthesis inhibitors;

15 III.12) decoupler herbicides;

III.13) auxin herbicides;

III.14) auxin transport inhibitors; and

III.15) other herbicides selected from the group consisting of benzoylprop,

flamprop, flamprop-M, bromobutide, chlorflurenol, cinmethylin,

20 methyldymuron, etobenzanid, fosamine, metam, pyributicarb,

oxaziclomefone, dazomet, triaziflam and methyl bromide;

including their agriculturally acceptable salts and, provided they have a carboxyl group, their agriculturally acceptable derivatives.

2. A composition as claimed in claim 1, wherein the variables R^1 to R^7 in formula I

25 are as defined below:

R^1 is methyl or NH_2 ;

R^2 is trifluoromethyl;

R^3 is hydrogen, fluorine or chlorine;

R⁴ is halogen or cyano;

R⁵ is hydrogen;

R⁶, R⁷ independently of one another are hydrogen, C₁-C₆-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkenyl, phenyl or benzyl.

3. A composition as claimed in claims 1 or 2, wherein R⁶ and R⁷ in formula I are identical or different C₁-C₆-alkyl radicals.

4. A composition as claimed in claim 1, wherein the variables R¹ to R⁷ in formula I have the following meanings:

10 R¹ is methyl;

R² is trifluoromethyl;


R³ is fluorine;

R⁴ is chlorine;

R⁵ is hydrogen;

15 R⁶, R⁷ independently of one another are C₁-C₆-alkyl.

5. A composition as claimed in claim 1, wherein the 3-phenyluracil I is the compound of formula:



wherein R¹ is methyl, R⁶ is methyl and R⁷ is isopropyl.

6. A composition as claimed in any one of claims 1 to 5 comprising as the only herbicidal active compound a 3-phenyluracil I in combination with the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II.

7. A composition as claimed in any one of claims 1 to 5 comprising a 3-phenyluracil I, N[[4-cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide II and at least one herbicide III selected from the classes III.1) to III.15.

8. A composition as claimed in any one of claims 1 to 5 and 7 wherein the herbicides III are selected from the group consisting of:

III.1) lipid biosynthesis inhibitors selected from the group consisting of:

10 clodinafop, cyhalofop, diclofop, fenoxaprop, fenoxaprop-p, fluazifop, fluazifop-P, haloxyfop, haloxyfop-P, metamifop, quizalofop, quizalofop-P, alloxydim, butroxydim, clethodim, cloproxydim, cycloxydim, profoxydim, sethoxydim, tepraloxym, tralkoxydim and pinoxaden;

III.2) ALS inhibitors selected from the group consisting of :

15 amidosulfuron, bensulfuron, chlorimuron, chlorsulfuron, cyclosulfamuron, flupyr-sulfuron, foramsulfuron, halosulfuron, imazosulfuron, iodosulfuron, mesosulfuron, metsulfuron, nicosulfuron, oxasulfuron, primisulfuron, prosulfuron, pyrazosulfuron, rimsulfuron, sulfometuron, sulfosulfuron, thifensulfuron, triasulfuron, tribenuron, trifloxysulfuron, triflusulfuron,
 20 tritosulfuron, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, cloransulam, diclosulam, florasulam, flumetsulam, metosulam, penoxsulam, bispyribac, pyriminobac, propoxycarbazone, flucarbazone, pyribenzoxim, pyriftalid, pyri-thiobac, flucetosulfuron, orthosulfamuron, pyrimisulfan and [N-(5,7-dimethoxy[1,2,4]triazolo[1,5- α]pyrimidin-2-yl)-2-methoxy-4-(trifluoromethyl)-3-pyridinesulfonamide;
 25

III.3) photosynthesis inhibitors selected from the group consisting of:
 atrazine, cyanazine, simazine, terbuthylazine, metamiltron, metribuzin,
 chloridazon, amicarbazone, chlorotoluron, diuron, isoproturon,
 methabenzthiazuron, bentazone, propanil, bromoxynil, ioxynil and
 5 paraquat;

III.5) bleacher herbicides selected from the group consisting of:
 norflurazon, diflufenican, picolinafen, beflubutamid, mesotrione,
 sulcotrione, isoxaflutole, benzofenap, pyrazolynate, pyrazoxyfen,
 benzobicyclon, 4-(3-trifluoromethylphenoxy)-2-(4-
 10 trifluoromethylphenyl)pyrimidine, topramezone, 4-hydroxy-3-[[2-methyl-6-
 (trifluoromethyl)-3-pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, 4-hydroxy-3-
 {[2-(2-methoxyethoxy)methyl-6-(trifluoro-methyl)-3-
 pyridinyl]carbonyl]bicyclo[3.2.1]oct-3-en-2-one, 4-hydroxy-3-[4-
 (methylsulfonyl)2-nitrobenzoyl]bicyclo[3.2.1]-oct-3-en-2-one, 2-[2-chloro-4-
 15 (methylsulfonyl)-3-[(2,2,2-trifluoroethoxy)methyl]benzoyl]-3-hydroxy-2-
 cyclohexen-1-one and pyrasulfotole;

III.6) a EPSP synthase inhibitor consisting of:
 glyphosate;

III.7) a glutamine synthase inhibitor consisting of:
 20 glufosinate;

III.9) mitose inhibitors selected from the group consisting of:
 oryzalin, pendimethalin and trifluralin;

III.10) VLCFA inhibitors selected from the group consisting of :
 acetochlor, butachlor, dimethenamid, dimethenamid-P, metazachlor,
 25 metolachlor, S-metolachlor, pethoxamid, pretilachlor, flufenacet,
 mefenacet, fentrazamide, cafenstrole and indanofan;

III.13) auxin herbicides selected from the group consisting of:

2,4-D, dichlorprop, dichlorprop-P, mecoprop, MCPA, mecoprop-P, dicamba, quinclorac, quinmerac, clopyralid, fluroxypyr, picloram, triclopyr and aminopyralid;

5 III.14) a auxin transport inhibitor consisting of:

diflufenzopyr; and

III.15) cinmethylin, oxaziclomefone and triaziflam;

their agriculturally acceptable salts and,

provided they have a carboxyl group, the agriculturally acceptable derivatives

10 thereof.

9. A composition as claimed in any one of claims 1 to 8 in form of a crop protection composition comprising additionally at least one inert liquid and/or solid carrier, optionally at least one or more surfactants and optionally at least one or more further auxiliaries.

15 10. A method for controlling undesired vegetation, which comprises allowing a herbicidally effective amount of a composition as claimed in any one of claims 1 to 9 to act on plants, their habitat or on seed.


11. A method as claimed in claim 10 wherein a herbicidal composition as claimed in any one of claims 1 to 9 is applied before, during and/or after the emergence of the
20 undesirable plants, the components I, II and optionally III being applied simultaneously or in succession.

12. The use of compositions as claimed in any one of claims 1 to 9 for controlling undesirable vegetation in crops.

13. The use as claimed in claim 12 wherein the crops are crops of oat, barley, millet, corn, rice, wheat, sugar cane, cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers and soybeans.

14. The use of compositions as claimed in any one of claims 1 to 9 for controlling undesirable vegetation in crops of plants, wherein the crop is a herbicide resistant crop, an insect resistant crop, a fungus resistant crop or two or all of these crops.

15. A method of safening the phytotoxic activity of a 3-phenyluracil I



I,


wherein the variables R^1 to R^7 are as defined below:

- R^1 is methyl or NH_2 ;
 R^2 is C_1 - C_2 -haloalkyl;
 R^3 is hydrogen or halogen;
 R^4 is halogen or cyano;
 R^5 is hydrogen or C_1 - C_6 -alkyl;
 R^6 , R^7 independently of one another are hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, C_3 - C_6 -alkenyl, C_3 - C_6 -alkynyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -cycloalkenyl, phenyl or benzyl;

or an agriculturally acceptable salt thereof;

on crops, which comprises applying said 3-phenyluracil I, or an agriculturally

acceptable salt thereof, in combination with N-[[4-[(cyclopropylamino)carbonyl]-phenyl]sulfonyl]-2-methoxybenzamide of formula II



or an agriculturally acceptable salt thereof; and

5 optionally at least one herbicide III selected from the group consisting of:

- III.1) lipid biosynthesis inhibitors;
- III.2) acetolactate synthase inhibitors (ALS inhibitors);
- III.3) photosynthesis inhibitors;
- III.4) protoporphyrinogen-IX oxidase inhibitors;
- 10 III.5) bleacher herbicides;
- III.6) enolpyruvyl shikimate 3-phosphate synthase inhibitors (EPSP inhibitors);
- III.7) glutamine synthetase inhibitors;
- III.8) 7,8-dihydropteroate synthase inhibitors (DHP inhibitors);
- 15 III.9) mitose inhibitors;
- III.10) inhibitors of the synthesis of long chain fatty acids (VLCFA inhibitors);
- III.11) cellulose biosynthesis inhibitors;
- III.12) decoupler herbicides;
- III.13) auxin herbicides;
- 20 III.14) auxin transport inhibitors; and
- III.15) other herbicides selected from the group consisting of benzoylprop, flamprop, flamprop-M, bromobutide, chlorflurenol, cinmethylin, methyldymuron, etobenzanid, fosamine, metam, pyributicarb, oxaziclomefone, dazomet, triaziflam and methyl bromide;

including their agriculturally acceptable salts and, provided they have a carboxyl group, their agriculturally acceptable derivatives;

in order to reduce or eliminate the phytotoxic activity of said 3- phenyluracil I.

16. A method of safening as claimed in claim 15 wherein the crops are selected
5 from the group consisting of oat, barley, millet, corn, rice, wheat, sugar cane, cotton, oilseed rape, flax, lentil, sugar beet, tobacco, sunflowers and soybeans.

17. A method of safening as claimed in claim 15 or 16 which comprises applying as only herbicidal active compound said 3-phenyluracil I in combination with the safener N-[[4-[(cyclopropylamino)carbonyl]phenyl]sulfonyl]-2-methoxybenzamide.

10 18. A method of safening as claimed in claim 15 or 16 which comprises applying said phenyluracil I in combination with N-[[4-[(cyclopropylamino)carbonyl]phenyl]-sulfonyl]-2-methoxybenzamide II, and at least one herbicide III selected from the group consisting of the herbicides III.1) to III.15) recited in claim 15.

15 19. A method of safening as claimed in any one of claims 15 to 18 wherein the variables R^1 to R^7 in formula I have the following meanings:

R^1 is methyl;

R^2 is trifluoromethyl;

R^3 is fluorine;

R^4 is chlorine;

20 R^5 is hydrogen;

R^6, R^7 independently of one another are C_1-C_6 -alkyl.

