



- (51) **International Patent Classification:** Not classified
- (21) **International Application Number:**
PCT/US2016/067430
- (22) **International Filing Date:**
19 December 2016 (19.12.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/269,044 17 December 2015 (17.12.2015) US
62/291,651 5 February 2016 (05.02.2016) US
- (71) **Applicant (for all designated States except US):** **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, 4056 Basel (CH).
- (72) **Inventors; and**
- (71) **Applicants (for US only):** **BILIC, Sanela** [US/US]; c/o Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, New Jersey 07936 (US). **HOWARD, JR., Danny Roland** [US/US]; c/o Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, New Jersey 07936 (US). **CAMERON, John Scott** [US/US]; c/o Novartis Institutes for Biomedical Research, Inc., 220 Massachusetts Avenue, Cambridge, Massachusetts 021399 (US).
- (74) **Agent:** **CHAPMAN, Jennifer**; Novartis Pharmaceuticals Corporation, One Health Plaza, East Hanover, New Jersey 07936 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(54) **Title:** COMBINATION OF C-MET INHIBITOR WITH ANTIBODY MOLECULE TO PD-1 AND USES THEREOF

Heavy Chain (murine IgG1)

```

FWH1          CDRH1  FWH2          CDRH2
QVQLQQSGSS LVRPGASVKL SCKASGYTFT TYWHWVRQR PQQLEWIGN IYPGTGGSNF DEKFKNRTSL
QVQLQQSGSS LVRPGASVKL SCKASGYTFT TYWHWVRQR PQQLEWIGN IYPGTGGSNF DEKFKNRTSL

FWH3          CDRH3  FWH4
TVDTSSITAY MHLASLTSED SAVYYCTRWT TGTGAYWQQG TLVTVSA
TVDTSSITAY MHLASLTSED SAVYYCTRWT TGTGAYWQQG TLVTVSAKT TTPSVYELAP GSAA

```

Light Chain (murine κ)

```

FWL1          CDRL1  FWL2          CDRL2
DIVMTQSPSS LTVTAGEKVT MSCKSSQSLL DSGNQKNFLT WYQKPGQPP KILIFWASTR ESGVDFRFTG
DIVMTQSPSS LTVTAGEKVT MSCKSSQSLL DSGNQKNFLT WYQKPGQPP KILIFWASTR ESGVDFRFTG

FWL3          CDRL3  FWL4
SGSVDFTLT ISSVQAEDLA VYYCQNDYSY FCTFGGKTL EIK
SGSVDFTLT ISSVQAEDLA VYYCQNDYSY FCTFGSGTKL EIKRAD

```

FIGURE 1

(57) **Abstract:** The present invention relates to a pharmaceutical combination which comprises (a) at least one antibody molecule (e.g., humanized antibody molecules) that bind to Programmed Death 1 (PD-1), and (b) at least one c-Met receptor tyrosine kinase inhibitor or pharmaceutically acceptable salt thereof, for simultaneous, separate or sequential administration for the treatment of a proliferative disease, particularly a c-Met dependent proliferative disease; a pharmaceutical composition comprising such combination; a method of treating a subject having a proliferative disease comprising administration of said combination to a subject in need thereof; use of such combination for the treatment of proliferative disease; and a commercial package comprising such combination.



**COMBINATION OF C-MET INHIBITOR WITH ANTIBODY MOLECULE TO PD-1
AND USES THEREOF**

FIELD OF THE INVENTION

5 The present invention relates to a pharmaceutical combination which comprises (a) at least one antibody molecule (*e.g.*, humanized antibody molecules) that bind to Programmed Death 1 (PD-1) , and (b) at least one c-Met receptor tyrosine kinase inhibitor or pharmaceutically acceptable salt thereof, for simultaneous, separate or sequential administration for the treatment of a proliferative disease, particularly a c-Met dependent proliferative disease; a pharmaceutical
10 composition comprising such combination; a method of treating a subject having a proliferative disease comprising administration of said combination to a subject in need thereof; use of such combination for the treatment of proliferative disease; and a commercial package comprising such combination.

BACKGROUND

20 The ability of T cells to mediate an immune response against an antigen requires two distinct signaling interactions (Viglietta, V. *et al.* (2007) *Neurotherapeutics* 4:666-675; Korman, A. J. *et al.* (2007) *Adv. Immunol.* 90:297-339). First, an antigen that has been arrayed on the surface of antigen-presenting cells (APC) is presented to an antigen-specific naive CD4⁺ T cell.
25 Such presentation delivers a signal via the T cell receptor (TCR) that directs the T cell to initiate an immune response specific to the presented antigen. Second, various co-stimulatory and inhibitory signals mediated through interactions between the APC and distinct T cell surface molecules trigger the activation and proliferation of the T cells and ultimately their inhibition.

30 The immune system is tightly controlled by a network of costimulatory and co-inhibitory ligands and receptors. These molecules provide the second signal for T cell activation and provide a balanced network of positive and negative signals to maximize immune responses

against infection, while limiting immunity to self (Wang, L. *et al.* (Epub Mar. 7, 2011) *J. Exp. Med.* 208(3):577-92; Lepenies, B. *et al.* (2008) *Endocrine, Metabolic & Immune Disorders--Drug Targets* 8:279-288). Examples of costimulatory signals include the binding between the B7.1 (CD80) and B7.2 (CD86) ligands of the APC and the CD28 and CTLA-4 receptors of the CD4⁺ T-lymphocyte (Sharpe, A. H. *et al.* (2002) *Nature Rev. Immunol.* 2:116-126; Lindley, P. S. *et al.* (2009) *Immunol. Rev.* 229:307-321). Binding of B7.1 or B7.2 to CD28 stimulates T cell activation, whereas binding of B7.1 or B7.2 to CTLA-4 inhibits such activation (Dong, C. *et al.* (2003) *Immunolog. Res.* 28(1):39-48; Greenwald, R. J. *et al.* (2005) *Ann. Rev. Immunol.* 23:515-548). CD28 is constitutively expressed on the surface of T cells (Gross, J., *et al.* (1992) *J. Immunol.* 149:380-388), whereas CTLA-4 expression is rapidly up-regulated following T-cell activation (Linsley, P. *et al.* (1996) *Immunity* 4:535-543).

Other ligands of the CD28 receptor include a group of related B7 molecules, also known as the "B7 Superfamily" (Coyle, A. J. *et al.* (2001) *Nature Immunol.* 2(3):203-209; Sharpe, A. H. *et al.* (2002) *Nature Rev. Immunol.* 2:116-126; Collins, M. *et al.* (2005) *Genome Biol.* 6:223.1-223.7; Korman, A. J. *et al.* (2007) *Adv. Immunol.* 90:297-339). Several members of the B7 Superfamily are known, including B7.1 (CD80), B7.2 (CD86), the inducible co-stimulator ligand (ICOS-L), the programmed death-1 ligand (PD-L1; B7-H1), the programmed death-2 ligand (PD-L2; B7-DC), B7-H3, B7-H4 and B7-H6 (Collins, M. *et al.* (2005) *Genome Biol.* 6:223.1-223.7).

The Programmed Death 1 (PD-1) protein is an inhibitory member of the extended CD28/CTLA-4 family of T cell regulators (Okazaki *et al.* (2002) *Curr Opin Immunol* 14:391779-82; Bennett *et al.* (2003) *J. Immunol.* 170:711-8). Other members of the CD28 family include CD28, CTLA-4, ICOS and BTLA. PD-1 is suggested to exist as a monomer, lacking the unpaired cysteine residue characteristic of other CD28 family members. PD-1 is expressed on activated B cells, T cells, and monocytes.

The PD-1 gene encodes a 55 kDa type I transmembrane protein (Agata *et al.* (1996) *Int Immunol.* 8:765-72). Although structurally similar to CTLA-4, PD-1 lacks the MYPPY motif (SEQ ID NO: 236) that is important for B7-1 and B7-2 binding. Two ligands for PD-1 have been identified, PD-L1 (B7-H1) and PD-L2 (B7-DC), that have been shown to downregulate T cell activation upon binding to PD-1 (Freeman *et al.* (2000) *J. Exp. Med.* 192:1027-34; Carter *et al.* (2002) *Eur. J. Immunol.* 32:634-43). Both PD-L1 and PD-L2 are B7 homologs that bind to

PD-1, but do not bind to other CD28 family members. PD-L1 is abundant in a variety of human cancers (Dong *et al.* (2002) *Nat. Med.* 8:787-9).

PD-1 is known as an immunoinhibitory protein that negatively regulates TCR signals (Ishida, Y. *et al.* (1992) *EMBO J.* 11:3887-3895; Blank, C. *et al.* (Epub 2006 Dec. 29) *Immunol. Immunother.* 56(5):739-745). The interaction between PD-1 and PD-L1 can act as an immune checkpoint, which can lead to, *e.g.*, a decrease in tumor infiltrating lymphocytes, a decrease in T-cell receptor mediated proliferation, and/or immune evasion by cancerous cells (Dong *et al.* (2003) *J. Mol. Med.* 81:281-7; Blank *et al.* (2005) *Cancer Immunol. Immunother.* 54:307-314; Konishi *et al.* (2004) *Clin. Cancer Res.* 10:5094-100). Immune suppression can be reversed by inhibiting the local interaction of PD-1 with PD-L1 or PD-L2; the effect is additive when the interaction of PD-1 with PD-L2 is blocked as well (Iwai *et al.* (2002) *Proc. Nat'l. Acad. Sci. USA* 99:12293-7; Brown *et al.* (2003) *J. Immunol.* 170:1257-66).

Given the importance of immune checkpoint pathways in regulating an immune response, the need exists for developing novel combination therapies that modulate the activity of immunoinhibitory proteins, such as PD-1, thus leading to activation of the immune system. Such agents can be used, *e.g.*, for cancer immunotherapy and treatment of other conditions, such as chronic infection.

c-Met, a proto-oncogene, is a member of a distinct subfamily of heterodimeric receptor tyrosine kinases which include Met, Ron, and Sea (Birchmeier, C. *et al.*, *Nat. Rev. Mol. Cell Biol.* 2003, 4(12):915-925; Christensen, J. G. *et al.*, *Cancer Lett.* 2005, 225(1):1-26). The only high affinity ligand for c-Met is the hepatocyte growth factor (HGF), also known as scatter factor (SF). Binding of HGF to c-Met induces activation of the receptor via autophosphorylation resulting in an increase of receptor dependent signaling. Both c-Met and HGF are widely expressed in a variety of organs, but their expression is normally confined to the cells of epithelial and mesenchymal origin, respectively. The biological functions of c-Met (or c-Met signaling pathway) in normal tissues and human malignancies such as cancer have been well documented (Christensen, J.G. *et al.*, *Cancer Lett.* 2005, 225(1):1-26; Corso, S. *et al.*, *Trends in Mol. Med.* 2005, 11(6):284-292).

HGF and c-Met are each required for normal mammalian development, and abnormalities reported in both HGF- and c-Met-null mice are consistent with proximity of embryonic expression and epithelial-mesenchymal transition defects during organ morphogenesis

(Christensen, J.G. et al., Cancer Lett. 2005, 225(1):1-26). Consistent with these findings, the transduction of signaling and subsequent biological effects of HGF/c-Met pathway have been shown to be important for epithelial-mesenchymal interaction and regulation of cell migration, invasion, cell proliferation and survival, angiogenesis, morphogenesis and organization of three-dimensional tubular structures (e.g. renal tubular cells, gland formation) during development. The specific consequences of c-Met pathway activation in a given cell/tissue are highly context-dependent.

Evidence shows that dysregulated c-Met pathway plays important and sometimes causative (in the case of genetic alterations) roles in tumor formation, growth, maintenance and progression (Birchmeier, C. et al., Nat. Rev. Mol. Cell. Biol. 2003, 4(12):915-925; Boccaccio, C. et al., Nat. Rev. Cancer 2006, 6(8):637-645; Christensen, J.G. et al., Cancer Lett. 2005, 225(1):1-26). HGF and/or c-Met are overexpressed in significant portions of most human cancers, and are often associated with poor clinical outcomes such as more aggressive disease, disease progression, tumor metastasis and shortened patient survival. Further, patients with high levels of HGF/c-Met proteins are more resistance to chemotherapy and radiotherapy. In addition to the abnormal HGF/c-Met expression, c-Met receptor can also be activated in cancer patients through genetic mutations (both germline and somatic) and gene amplification. Although gene amplification and mutations are the most common genetic alterations that have been reported in patients, the receptor can also be activated by deletions, truncations, gene rearrangement, as well as abnormal receptor processing and defective negative regulatory mechanisms.

SUMMARY

Disclosed herein is pharmaceutical combination which comprises (a) at least one antibody molecule (*e.g.*, humanized antibody molecules) that bind to Programmed Death 1 (PD-1), and (b) at least one c-Met receptor tyrosine kinase inhibitor or pharmaceutically acceptable salt thereof, for simultaneous, separate or sequential administration for the treatment of a proliferative disease, particularly a c-Met dependent proliferative disease; a pharmaceutical composition comprising such combination; a method of treating a subject having a proliferative disease comprising administration of said combination to a subject in need thereof; use of such

combination for the treatment of proliferative disease; and a commercial package comprising such combination.

In one embodiment, the invention features a method of treating (*e.g.*, inhibiting, reducing, ameliorating, or preventing) a disorder, *e.g.*, a hyperproliferative condition or disorder (*e.g.*, a cancer) in a subject. The method includes administering, in combination with a c-Met receptor tyrosine kinase inhibitor, to the subject an anti-PD-1 antibody molecule, *e.g.*, an anti-PD-1 antibody molecule described herein, at a dose of about 300 mg to 400 mg once every three weeks or once every four weeks. In certain embodiments, the anti-PD-1 antibody molecule is administered at a dose of about 300 mg once every three weeks. In other embodiments, the anti-PD-1 antibody molecule is administered at a dose of about 400 mg once every four weeks. In some embodiments, the disorder is a cancer, *e.g.*, a cancer described herein.

In some embodiments, the anti-PD-1 antibody molecule is administered by injection (*e.g.*, subcutaneously or intravenously) at a dose (*e.g.*, a flat dose) of about 200 mg to 500 mg, *e.g.*, about 250 mg to 450 mg, about 300 mg to 400 mg, about 250 mg to 350 mg, about 350 mg to 450 mg, or about 300 mg or about 400 mg. The dosing schedule (*e.g.*, flat dosing schedule) can vary from *e.g.*, once a week to once every 2, 3, 4, 5, or 6 weeks. In one embodiment, the anti-PD-1 antibody molecule is administered at a dose from about 300 mg to 400 mg once every three weeks or once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 300 mg once every three weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 400 mg once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 300 mg once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 400 mg once every three weeks.

In another aspect, the invention features a method of reducing an activity (*e.g.*, growth, survival, or viability, or all), of a hyperproliferative (*e.g.*, a cancer) cell. The method includes contacting the cell with an anti-PD-1 antibody molecule, *e.g.*, an anti-PD-1 antibody molecule described herein. The method can be performed in a subject, *e.g.*, as part of a therapeutic protocol in combination with a c-Met receptor tyrosine kinase inhibitor, *e.g.*, at a dose of about 300 mg to 400 mg of an anti-PD-1 antibody molecule once every three weeks or once every four weeks. In certain embodiments, the dose is about 300 mg of an anti-PD-1 antibody

molecule once every three weeks. In other embodiments, the dose is about 400 mg of an anti-PD-1 antibody molecule once every four weeks. The cancer cell can be, *e.g.*, a cell from a cancer described herein, such as lung cancer (squamous), lung cancer (adenocarcinoma), head and neck cancer, cervical cancer (squamous), stomach cancer, thyroid cancer, melanoma, nasopharyngeal cancer (*e.g.*, differentiated or undifferentiated metastatic or locally recurrent nasopharyngeal carcinoma), or breast cancer.

In another aspect, the invention features a composition (*e.g.*, one or more compositions or dosage forms), that includes an anti-PD-1 antibody molecule (*e.g.*, an anti-PD-1 antibody molecule as described herein). Formulations, *e.g.*, dosage formulations, and kits, *e.g.*, therapeutic kits, that include an anti-PD-1 antibody molecule (*e.g.*, an anti-PD-1 antibody molecule as described herein), are also described herein. In certain embodiments, the composition or formulation comprises 300 mg or 400 mg of an anti-PD-1 antibody molecule (*e.g.*, an anti-PD-1 antibody molecule as described herein). In some embodiments, the composition of formulation is administered or used once every three weeks or once every four weeks. Such composition is used in combination with at least one c-Met receptor tyrosine kinase inhibitor or pharmaceutically acceptable salt thereof, for simultaneous, separate or sequential administration.

The combinations disclosed herein can be administered together in a single composition or administered separately in two or more different compositions, *e.g.*, compositions or dosage forms as described herein. The administration of the therapeutic agents can be in any order. The first agent and the additional agents (*e.g.*, second, third agents) can be administered via the same administration route or via different administration routes.

Antibody Molecules to PD-1

In one embodiment, the PD-1 inhibitor is an anti-PD-1 antibody molecule as described in USSN 14/604,415, entitled "Antibody Molecules to PD-1 and Uses Thereof," incorporated by reference in its entirety. In one embodiment, the anti-PD-1 antibody molecule comprises at least one antigen-binding region, *e.g.*, a variable region or an antigen-binding fragment thereof, from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-

hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (e.g., at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one, two, three or four variable regions from an antibody described herein, e.g., an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (e.g., at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one or two heavy chain variable regions from an antibody described herein, e.g., an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (e.g., at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one or two light chain variable regions from an antibody described herein, e.g., an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or

BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule includes a heavy chain
5 constant region for an IgG4, *e.g.*, a human IgG4. In one embodiment, the human IgG4 includes a substitution at position 228 according to EU numbering (*e.g.*, a Ser to Pro substitution). In still another embodiment, the anti-PD-1 antibody molecule includes a heavy chain constant region for an IgG1, *e.g.*, a human IgG1. In one embodiment, the human IgG1 includes a substitution at position 297 according to EU numbering (*e.g.*, an Asn to Ala substitution). In one embodiment,
10 the human IgG1 includes a substitution at position 265 according to EU numbering, a substitution at position 329 according to EU numbering, or both (*e.g.*, an Asp to Ala substitution at position 265 and/or a Pro to Ala substitution at position 329). In one embodiment, the human IgG1 includes a substitution at position 234 according to EU numbering, a substitution at position 235 according to EU numbering, or both (*e.g.*, a Leu to Ala substitution at position 234
15 and/or a Leu to Ala substitution at position 235). In one embodiment, the heavy chain constant region comprises an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto.

In yet another embodiment, the anti-PD-1 antibody molecule includes a kappa light chain constant region, *e.g.*, a human kappa light chain constant region. In one embodiment, the light
20 chain constant region comprises an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto.

In another embodiment, the anti-PD-1 antibody molecule includes a heavy chain constant region for an IgG4, *e.g.*, a human IgG4, and a kappa light chain constant region, *e.g.*, a human
25 kappa light chain constant region, *e.g.*, a heavy and light chain constant region comprising an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto. In one embodiment, the human IgG4 includes a substitution at position 228 according to EU numbering (*e.g.*, a Ser to Pro substitution). In yet another embodiment, the anti-PD-1 antibody molecule includes a heavy
30 chain constant region for an IgG1, *e.g.*, a human IgG1, and a kappa light chain constant region, *e.g.*, a human kappa light chain constant region, *e.g.*, a heavy and light chain constant region

comprising an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto. In one embodiment, the human IgG1 includes a substitution at position 297 according to EU numbering (*e.g.*, an Asn to Ala substitution). In one embodiment, the human IgG1 includes a substitution at position 265 according to EU numbering, a substitution at position 329 according to EU numbering, or both (*e.g.*, an Asp to Ala substitution at position 265 and/or a Pro to Ala substitution at position 329). In one embodiment, the human IgG1 includes a substitution at position 234 according to EU numbering, a substitution at position 235 according to EU numbering, or both (*e.g.*, a Leu to Ala substitution at position 234 and/or a Leu to Ala substitution at position 235).

In another embodiment, the anti-PD-1 antibody molecule includes a heavy chain variable domain and a constant region, a light chain variable domain and a constant region, or both, comprising the amino acid sequence of BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences. The anti-PD-1 antibody molecule, optionally, comprises a leader sequence from a heavy chain, a light chain, or both, as shown in Table 4; or a sequence substantially identical thereto.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three complementarity determining regions (CDRs) from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs (or collectively all of the CDRs) from a heavy chain variable region comprising an amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1.

In one embodiment, one or more of the CDRs (or collectively all of the CDRs) have one, two, three, four, five, six or more changes, *e.g.*, amino acid substitutions or deletions, relative to the amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, 5 or three CDRs from a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, 10 BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequence.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs (or collectively all of the CDRs) from a light chain variable region comprising an 15 amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1. In one embodiment, one or more of the CDRs (or collectively all of the CDRs) have one, two, three, four, five, six or more changes, *e.g.*, amino acid substitutions or deletions, relative to the amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1. In certain embodiments, the anti-PD-1 antibody molecule includes a substitution in a light chain 20 CDR, *e.g.*, one or more substitutions in a CDR1, CDR2 and/or CDR3 of the light chain. In one embodiment, the anti-PD-1 antibody molecule includes a substitution in the light chain CDR3 at position 102 of the light variable region, *e.g.*, a substitution of a cysteine to tyrosine, or a cysteine to serine residue, at position 102 of the light variable region according to Table 1 (*e.g.*, SEQ ID NO: 16 or 24 for murine or chimeric, unmodified; or any of SEQ ID NOs: 34, 42, 46, 25 54, 58, 62, 66, 70, 74, or 78 for a modified sequence).

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, three, four, five or six CDRs (or collectively all of the CDRs) from a heavy and light chain variable region comprising an amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1. In one embodiment, one or more of the CDRs (or collectively all of 30 the CDRs) have one, two, three, four, five, six or more changes, *e.g.*, amino acid substitutions or

deletions, relative to the amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1.

In one embodiment, the anti-PD-1 antibody molecule includes all six CDRs from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1, or closely related CDRs, *e.g.*, CDRs which are identical or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions). In one embodiment, the anti-PD-1 antibody molecule may include any CDR described herein. In certain embodiments, the anti-PD-1 antibody molecule includes a substitution in a light chain CDR, *e.g.*, one or more substitutions in a CDR1, CDR2 and/or CDR3 of the light chain. In one embodiment, the anti-PD-1 antibody molecule includes a substitution in the light chain CDR3 at position 102 of the light variable region, *e.g.*, a substitution of a cysteine to tyrosine, or a cysteine to serine residue, at position 102 of the light variable region according to Table 1 (*e.g.*, SEQ ID NO: 16 or 24 for murine or chimeric, unmodified; or any of SEQ ID NOs: 34, 42, 46, 54, 58, 62, 66, 70, 74, or 78 for a modified sequence).

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs according to Kabat *et al.* (*e.g.*, at least one, two, or three CDRs according to the Kabat definition as set out in Table 1) from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two,

three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three CDRs according to Kabat *et al.* shown in Table 1.

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs according to Kabat *et al.* (*e.g.*, at least one, two, or three CDRs according to the Kabat definition as set out in Table 1) from a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three CDRs according to Kabat *et al.* shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, three, four, five, or six CDRs according to Kabat *et al.* (*e.g.*, at least one, two, three, four, five, or six CDRs according to the Kabat definition as set out in Table 1) from the heavy and light chain variable regions of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, three, four, five, or six CDRs according to Kabat *et al.* shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes all six CDRs according to Kabat *et al.* (*e.g.*, all six CDRs according to the Kabat definition as set out in Table

1) from the heavy and light chain variable regions of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to all six CDRs according to Kabat *et al.* shown in Table 1. In one embodiment, the anti-PD-1 antibody molecule may include any CDR described herein.

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three Chothia hypervariable loops (*e.g.*, at least one, two, or three hypervariable loops according to the Chothia definition as set out in Table 1) from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or at least the amino acids from those hypervariable loops that contact PD-1; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three hypervariable loops according to Chothia *et al.* shown in Table 1.

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three Chothia hypervariable loops (*e.g.*, at least one, two, or three hypervariable loops according to the Chothia definition as set out in Table 1) of a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12,

BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or at least the amino acids from those hypervariable loops that contact PD-1; or which have at least one amino acid alteration, but not
 5 more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three hypervariable loops according to Chothia *et al.* shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, three, four, five, or six hypervariable loops (*e.g.*, at least one, two, three, four, five, or six
 10 hypervariable loops according to the Chothia definition as set out in Table 1) from the heavy and light chain variable regions of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15,
 15 BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or at least the amino acids from those hypervariable loops that contact PD-1; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, three,
 20 four, five or six hypervariable loops according to Chothia *et al.* shown in Table 1.

In one embodiment, the anti-PD-1 antibody molecule includes all six hypervariable loops (*e.g.*, all six hypervariable loops according to the Chothia definition as set out in Table 1) of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07,
 25 BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, or closely related hypervariable loops, *e.g.*, hypervariable loops which are identical or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions); or which have at least one amino acid
 30 alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or

insertions, *e.g.*, conservative substitutions) relative to all six hypervariable loops according to Chothia *et al.* shown in Table 1. In one embodiment, the anti-PD-1 antibody molecule may include any hypervariable loop described herein.

In still another embodiment, the anti-PD-1 antibody molecule includes at least one, two,
 5 or three hypervariable loops that have the same canonical structures as the corresponding hypervariable loop of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15,
 10 BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, *e.g.*, the same canonical structures as at least loop 1 and/or loop 2 of the heavy and/or light chain variable domains of an antibody described herein. *See, e.g.*, Chothia *et al.*, (1992) *J. Mol. Biol.* 227:799-817; Tomlinson *et al.*, (1992) *J. Mol. Biol.* 227:776-798 for descriptions of hypervariable loop canonical structures. These structures can be determined by
 15 inspection of the tables described in these references.

In certain embodiments, the anti-PD-1 antibody molecule includes a combination of CDRs or hypervariable loops defined according to the Kabat *et al.* and Chothia *et al.*

In one embodiment, the anti-PD-1 antibody molecule includes at least one, two or three CDRs or hypervariable loops from a heavy chain variable region of an antibody described
 20 herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, according to the Kabat and Chothia
 25 definition (*e.g.*, at least one, two, or three CDRs or hypervariable loops according to the Kabat and Chothia definition as set out in Table 1); or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or
 30 insertions, *e.g.*, conservative substitutions) relative to one, two, or three CDRs or hypervariable loops according to Kabat and/or Chothia shown in Table 1.

For example, the anti-PD-1 antibody molecule can include VH CDR1 according to Kabat *et al.* or VH hypervariable loop 1 according to Chothia *et al.*, or a combination thereof, *e.g.*, as shown in Table 1. In one embodiment, the combination of Kabat and Chothia CDR of VH CDR1 comprises the amino acid sequence GYTFTTYWMH (SEQ ID NO: 224), or an amino acid sequence substantially identical thereto (*e.g.*, having at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions)). The anti-PD-1 antibody molecule can further include, *e.g.*, VH CDRs 2-3 according to Kabat *et al.* and VL CDRs 1-3 according to Kabat *et al.*, *e.g.*, as shown in Table 1. Accordingly, in some embodiments, framework regions are defined based on a combination of CDRs defined according to Kabat *et al.* and hypervariable loops defined according to Chothia *et al.* For example, the anti-PD-1 antibody molecule can include VH FR1 defined based on VH hypervariable loop 1 according to Chothia *et al.* and VH FR2 defined based on VH CDRs 1-2 according to Kabat *et al.*, *e.g.*, as shown in Table 1. The anti-PD-1 antibody molecule can further include, *e.g.*, VH FRs 3-4 defined based on VH CDRs 2-3 according to Kabat *et al.* and VL FRs 1-4 defined based on VL CDRs 1-3 according to Kabat *et al.*

The anti-PD-1 antibody molecule can contain any combination of CDRs or hypervariable loops according to the Kabat and Chothia definitions. In one embodiment, the anti-PD-1 antibody molecule includes at least one, two or three CDRs from a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, according to the Kabat and Chothia definition (*e.g.*, at least one, two, or three CDRs according to the Kabat and Chothia definition as set out in Table 1).

In one embodiment, the anti-PD-1 antibody molecule includes:

(a) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a light chain variable region (VL) comprising a VLCDR1 amino

acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33;

(b) a VH comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32;

(c) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33; or

(d) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

In the combinations herein, in another embodiment, the anti-PD-1 antibody molecule comprises (i) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1, SEQ ID NO: 4, or SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 5; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and (ii) a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10 or SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 11 or SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 32 or SEQ ID NO: 33.

In an embodiment, *e.g.*, an embodiment comprising a variable region, a CDR (*e.g.*, Chothia CDR or Kabat CDR), or other sequence referred to herein, *e.g.*, in Table 1, the antibody molecule is a monospecific antibody molecule, a bispecific antibody molecule, or is an antibody molecule that comprises an antigen binding fragment of an antibody, *e.g.*, a half antibody or antigen binding fragment of a half antibody. In certain embodiments the antibody molecule is a bispecific antibody molecule having a first binding specificity for PD-1 and a second binding specificity for TIM-3, LAG-3, CEACAM (*e.g.*, CEACAM-1, CEACAM-3, and/or CEACAM-5), PD-L1 or PD-L2. In one embodiment, the bispecific antibody molecule binds to PD-1 and TIM-3. In another embodiment, the bispecific antibody molecule binds to PD-1 and LAG-3. In

another embodiment, the bispecific antibody molecule binds to PD-1 and CEACAM (*e.g.*, CEACAM-1, CEACAM-3, and/or CEACAM-5). In another embodiment, the bispecific antibody molecule binds to PD-1 and CEACAM-1. In yet another embodiment, the bispecific antibody molecule binds to PD-1 and CEACAM-5. In another embodiment, the bispecific antibody molecule binds to PD-1 and PD-L1. In yet another embodiment, the bispecific antibody molecule binds to PD-1 and PD-L2. Any combination of the aforesaid molecules can be made in a multispecific antibody molecule, *e.g.*, a trispecific antibody that includes a first binding specificity to PD-1, and a second and third binding specificity to one or more of: TIM-3, LAG-3, CEACAM (*e.g.*, CEACAM-1, CEACAM-3, or CEACAM-5), PD-L1 or PD-L2.

C-Met Receptor Tyrosine Kinase Inhibitor

c-Met Receptor Tyrosine Kinase Inhibitor of the present invention is disclosed, for example, in US Patent 7,767,675, incorporated herein by reference in its entirety.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is 2-fluoro-N-methyl-4-[7-quinolin-6-yl-methyl)-imidazo[1,2-b][1,2,4]triazin-2yl]benzamide or pharmaceutically acceptable salt thereof.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is 2-fluoro-N-methyl-4-[7-quinolin-6-yl-methyl)-imidazo[1,2-b][1,2,4]triazin-2yl]benzamide dihydrochloric acid salt.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is capmatinib.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is capmatinib dihydrochloric acid salt. In one embodiment, capmatinib is administered at a dose of about 400-600 mg (*e.g.*, per day), *e.g.*, about 400, 500, or 600 mg, or about 400-500 or 500-600 mg.

In one embodiment, capmatinib is administered orally.

Uses of the Combination Therapies

The combinations disclosed herein can result in one or more of: an increase in antigen presentation, an increase in effector cell function (*e.g.*, one or more of T cell proliferation, IFN- γ secretion or cytolytic function), inhibition of regulatory T cell function, an effect on the activity of multiple cell types, such as regulatory T cell, effector T cells and NK cells), an increase in

tumor infiltrating lymphocytes, an increase in T-cell receptor mediated proliferation, and a decrease in immune evasion by cancerous cells. In one embodiment, the use of a PD-1 inhibitor in the combinations inhibits, reduces or neutralizes one or more activities of PD-1, resulting in blockade or reduction of an immune checkpoint. Thus, such combinations can be used to treat or prevent disorders where enhancing an immune response in a subject is desired.

Accordingly, in another aspect, a method of modulating an immune response in a subject is provided. The method comprises administering to the subject a combination disclosed herein (*e.g.*, a combination comprising a therapeutically effective amount of an anti-PD-1 antibody molecule), alone or in combination with one or more agents or procedures, such that the immune response in the subject is modulated. In one embodiment, the antibody molecule enhances, stimulates or increases the immune response in the subject. The subject can be a mammal, *e.g.*, a primate, preferably a higher primate, *e.g.*, a human (*e.g.*, a patient having, or at risk of having, a disorder described herein). In one embodiment, the subject is in need of enhancing an immune response. In one embodiment, the subject has, or is at risk of, having a disorder described herein, *e.g.*, a cancer or an infectious disorder as described herein. In certain embodiments, the subject is, or is at risk of being, immunocompromised. For example, the subject is undergoing or has undergone a chemotherapeutic treatment and/or radiation therapy. Alternatively, or in combination, the subject is, or is at risk of being, immunocompromised as a result of an infection.

In one aspect, a method of treating (*e.g.*, one or more of reducing, inhibiting, or delaying progression) a cancer or a tumor in a subject is provided. The method comprises administering to the subject a combination disclosed herein (*e.g.*, a combination comprising a therapeutically effective amount of an anti-PD-1 antibody molecule).

In certain embodiments, the cancer treated with the combination, includes but is not limited to, a solid tumor, a hematological cancer (*e.g.*, leukemia, lymphoma, myeloma, *e.g.*, multiple myeloma), and a metastatic lesion. In one embodiment, the cancer is a solid tumor. Examples of solid tumors include malignancies, *e.g.*, sarcomas and carcinomas, *e.g.*, adenocarcinomas of the various organ systems, such as those affecting the lung, breast, ovarian, lymphoid, gastrointestinal (*e.g.*, colon), anal, genitals and genitourinary tract (*e.g.*, renal, urothelial, bladder cells, prostate), pharynx, CNS (*e.g.*, brain, neural or glial cells), head and neck, skin (*e.g.*, melanoma), and pancreas, as well as adenocarcinomas which include

malignancies such as colon cancers, rectal cancer, renal-cell carcinoma, liver cancer (*e.g.*, hepatocellular carcinoma), non-small cell lung cancer, cancer of the small intestine and cancer of the esophagus. The cancer may be at an early, intermediate, late stage or metastatic cancer.

In one embodiment, the cancer is chosen from a lung cancer (*e.g.*, a non-small cell lung cancer (NSCLC) (*e.g.*, a NSCLC with squamous and/or non-squamous histology, or a NSCLC adenocarcinoma)), a melanoma (*e.g.*, an advanced melanoma), a renal cancer (*e.g.*, a renal cell carcinoma), a liver cancer (*e.g.*, hepatocellular carcinoma), a myeloma (*e.g.*, a multiple myeloma), a prostate cancer, a breast cancer (*e.g.*, a breast cancer that does not express one, two or all of estrogen receptor, progesterone receptor, or Her2/neu, *e.g.*, a triple negative breast cancer), a colorectal cancer, a pancreatic cancer, a head and neck cancer (*e.g.*, head and neck squamous cell carcinoma (HNSCC), anal cancer, gastro-esophageal cancer, thyroid cancer, cervical cancer, a lymphoproliferative disease (*e.g.*, a post-transplant lymphoproliferative disease) or a hematological cancer, T-cell lymphoma, B-cell lymphoma, a non-Hodgkin lymphoma, or a leukemia (*e.g.*, a myeloid leukemia or a lymphoid leukemia).

In another embodiment, the cancer is chosen from a carcinoma (*e.g.*, advanced or metastatic carcinoma), melanoma or a lung carcinoma, *e.g.*, a non-small cell lung carcinoma.

In one embodiment, the cancer is a lung cancer, *e.g.*, a non-small cell lung cancer or small cell lung cancer.

In one embodiment, the cancer is a melanoma, *e.g.*, an advanced melanoma. In one embodiment, the cancer is an advanced or unresectable melanoma that does not respond to other therapies. In other embodiments, the cancer is a melanoma with a BRAF mutation (*e.g.*, a BRAF V600 mutation). In yet other embodiments, the combination disclosed herein (*e.g.*, the combination comprising the anti-PD-1 antibody molecule) is administered after treatment with an anti-CTLA4 antibody (*e.g.*, ipilimumab) with or without a BRAF inhibitor (*e.g.*, vemurafenib or dabrafenib).

In another embodiment, the cancer is a hepatocarcinoma, *e.g.*, an advanced hepatocarcinoma, with or without a viral infection, *e.g.*, a chronic viral hepatitis.

In another embodiment, the cancer is a prostate cancer, *e.g.*, an advanced prostate cancer.

In yet another embodiment, the cancer is a myeloma, *e.g.*, multiple myeloma.

In yet another embodiment, the cancer is a renal cancer, *e.g.*, a renal cell carcinoma (RCC) (*e.g.*, a metastatic RCC or clear cell renal cell carcinoma (CCRCC)).

In one embodiment, the cancer microenvironment has an elevated level of PD-L1 expression. Alternatively, or in combination, the cancer microenvironment can have increased IFN γ and/or CD8 expression.

5 In some embodiments, the subject has, or is identified as having, a tumor that has one or more of high PD-L1 level or expression, or as being Tumor Infiltrating Lymphocyte (TIL)+ (*e.g.*, as having an increased number of TILs), or both. In certain embodiments, the subject has, or is identified as having, a tumor that has high PD-L1 level or expression and that is TIL+. In some embodiments, the methods described herein further include identifying a subject based on having a tumor that has one or more of high PD-L1 level or expression, or as being TIL+, or both. In
10 certain embodiments, the methods described herein further include identifying a subject based on having a tumor that has high PD-L1 level or expression and as being TIL+. In some embodiments, tumors that are TIL+ are positive for CD8 and IFN γ . In some embodiments, the subject has, or is identified as having, a high percentage of cells that are positive for one, two or more of PD-L1, CD8, and/or IFN γ . In certain embodiments, the subject has or is identified as
15 having a high percentage of cells that are positive for all of PD-L1, CD8, and IFN γ .

In some embodiments, the methods described herein further include identifying a subject based on having a high percentage of cells that are positive for one, two or more of PD-L1, CD8, and/or IFN γ . In certain embodiments, the methods described herein further include identifying a subject based on having a high percentage of cells that are positive for all of PD-L1, CD8, and
20 IFN γ . In some embodiments, the subject has, or is identified as having, one, two or more of PD-L1, CD8, and/or IFN γ , and one or more of a lung cancer, *e.g.*, squamous cell lung cancer or lung adenocarcinoma; a head and neck cancer; a squamous cell cervical cancer; a stomach cancer; an esophageal cancer; a thyroid cancer; a melanoma, and/or a nasopharyngeal cancer (NPC). In certain embodiments, the methods described herein further describe identifying a subject based
25 on having one, two or more of PD-L1, CD8, and/or IFN γ , and one or more of a lung cancer, *e.g.*, squamous cell lung cancer or lung adenocarcinoma; a head and neck cancer; a squamous cell cervical cancer; a stomach cancer; a thyroid cancer; a melanoma, and or a nasopharyngeal cancer.

30 Methods and compositions disclosed herein are useful for treating metastatic lesions associated with the aforementioned cancers.

In a further aspect, the invention provides a method of treating an infectious disease in a subject, comprising administering to a subject a combination as described herein, *e.g.*, a combination comprising a therapeutically effective amount of an anti-PD-1 antibody molecule described herein. In one embodiment, the infection disease is chosen from hepatitis (*e.g.*, hepatitis C infection), or sepsis.

Still further, the invention provides a method of enhancing an immune response to an antigen in a subject, comprising administering to the subject: (i) the antigen; and (ii) a combination as described herein, *e.g.*, a combination comprising a therapeutically effective amount of an anti-PD-1 antibody molecule described herein, such that an immune response to the antigen in the subject is enhanced. The antigen can be, for example, a tumor antigen, a viral antigen, a bacterial antigen or an antigen from a pathogen.

The combinations as described herein can be administered to the subject systemically (*e.g.*, orally, parenterally, subcutaneously, intravenously, rectally, intramuscularly, intraperitoneally, intranasally, transdermally, or by inhalation or intracavitary installation), topically, or by application to mucous membranes, such as the nose, throat and bronchial tubes.

Dosages and therapeutic regimens of the therapeutic agents disclosed herein can be determined by a skilled artisan. In certain embodiments, the anti-PD-1 antibody molecule is administered by injection (*e.g.*, subcutaneously or intravenously) at a dose of about 1 to 30 mg/kg, *e.g.*, about 5 to 25 mg/kg, about 10 to 20 mg/kg, about 1 to 5 mg/kg, or about 3 mg/kg. The dosing schedule can vary from *e.g.*, once a week to once every 2, 3, or 4 weeks. In one embodiment, the anti-PD-1 antibody molecule is administered at a dose from about 10 to 20 mg/kg every other week.

In some embodiments, the anti-PD-1 antibody molecule is administered by injection (*e.g.*, subcutaneously or intravenously) at a dose (*e.g.*, a flat dose) of about 200 mg to 500 mg, *e.g.*, about 250 mg to 450 mg, about 300 mg to 400 mg, about 250 mg to 350 mg, about 350 mg to 450 mg, or about 300 mg or about 400 mg. The dosing schedule (*e.g.*, flat dosing schedule) can vary from *e.g.*, once a week to once every 2, 3, 4, 5, or 6 weeks. In one embodiment, the anti-PD-1 antibody molecule is administered at a dose from about 300 mg to 400 mg once every three weeks or once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 300 mg once every three weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 400 mg once every four

weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 300 mg once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 400 mg once every three weeks.

In one embodiment, the anti-PD-1 antibody molecule is administered, alone or in
5 combination (*e.g.*, in combination with an anti-LAG-3 antibody molecule), at a dose of less than, or about, 5 mg/kg; less than, or about, 4 mg/kg; less than, or about, 3 mg/kg; less than, or about, 2 mg/kg; less than, or about, 1 mg/kg, every other week. In one embodiment, the anti-PD-1 antibody molecule is administered at a dose of 1 to 5 mg/kg every other week; 1 to 4 mg/kg every other week, 1 to 3 mg/kg every other week, or 1 to 2 mg/kg every other week. In one
10 embodiment, the anti-LAG-3 antibody molecule is administered, alone or in combination (*e.g.*, in combination with an anti-PD-1 antibody molecule) at a dose of 1 to 5 mg/kg every other week; 1 to 4 mg/kg every other week, 1 to 3 mg/kg every other week, or 1 to 2 mg/kg every other week.

The antibody molecules described herein are preferred for use in the methods described
15 herein, although other anti-PD-1 antibodies can be used instead, or in combination with an anti-PD-1 antibody molecule of the invention.

Further Combination Therapies

The methods and combinations described herein can be used in combination with other
20 agents or therapeutic modalities. In one embodiment, the methods described herein include administering to the subject a combination comprising an anti-PD-1 antibody molecule as described herein, in combination with an agent or therapeutic procedure or modality, in an amount effective to treat or prevent a disorder. The anti-PD-1 antibody molecule and the agent or therapeutic procedure or modality can be administered simultaneously or sequentially in any
25 order. Any combination and sequence of the anti-PD-1 antibody molecules and other therapeutic agents, procedures or modalities (*e.g.*, as described herein) can be used. The antibody molecule and/or other therapeutic agents, procedures or modalities can be administered during periods of active disorder, or during a period of remission or less active disease. The antibody molecule can be administered before the other treatment, concurrently with the treatment, post-treatment,
30 or during remission of the disorder.

In certain embodiments, the methods and compositions described herein are administered in combination with one or more of other antibody molecules, chemotherapy, other anti-cancer therapy (*e.g.*, targeted anti-cancer therapies, gene therapy, viral therapy, RNA therapy bone marrow transplantation, nanotherapy, or oncolytic drugs), cytotoxic agents, immune-based therapies (*e.g.*, cytokines or cell-based immune therapies), surgical procedures (*e.g.*, lumpectomy or mastectomy) or radiation procedures, or a combination of any of the foregoing. The additional therapy may be in the form of adjuvant or neoadjuvant therapy. In some embodiments, the additional therapy is an enzymatic inhibitor (*e.g.*, a small molecule enzymatic inhibitor) or a metastatic inhibitor. Exemplary cytotoxic agents that can be administered in combination with include antimicrotubule agents, topoisomerase inhibitors, anti-metabolites, mitotic inhibitors, alkylating agents, anthracyclines, vinca alkaloids, intercalating agents, agents capable of interfering with a signal transduction pathway, agents that promote apoptosis, proteasome inhibitors, and radiation (*e.g.*, local or whole body irradiation (*e.g.*, gamma irradiation)). In other embodiments, the additional therapy is surgery or radiation, or a combination thereof. In other embodiments, the additional therapy is a therapy targeting one or more of PI3K/AKT/mTOR pathway, an HSP90 inhibitor, or a tubulin inhibitor.

Alternatively, or in combination with the aforesaid combinations, the methods and compositions described herein can be administered in combination with one or more of: an immunomodulator (*e.g.*, an activator of a costimulatory molecule or an inhibitor of an inhibitory molecule, *e.g.*, an immune checkpoint molecule); a vaccine, *e.g.*, a therapeutic cancer vaccine; or other forms of cellular immunotherapy.

Exemplary non-limiting combinations and uses of the combinations disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, include the following.

In certain embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with a modulator of a costimulatory molecule or an inhibitory molecule, *e.g.*, a co-inhibitory ligand or receptor.

In one embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with a modulator, *e.g.*, agonist, of a costimulatory molecule. In one embodiment, the agonist of the costimulatory molecule is chosen from an agonist (*e.g.*, an agonistic antibody or antigen-binding fragment thereof, or a soluble fusion) of OX40, CD2, CD27, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), 4-1BB

(CD137), GITR, CD30, CD40, BAFFR, HVEM, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3 or CD83 ligand.

In one embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with an inhibitor of an inhibitory (or immune checkpoint) molecule chosen from PD-L1, PD-L2, CTLA-4, TIM-3, LAG-3, CEACAM (*e.g.*, CEACAM-1, CEACAM-3, and/or CEACAM-5), VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and/or TGFR beta. In one embodiment, the inhibitor is a soluble ligand (*e.g.*, a CTLA-4-Ig), or an antibody or antibody fragment that binds to PD-L1, PD-L2 or CTLA-4. For example, the anti-PD-1 antibody molecule can be administered in combination with an anti-CTLA-4 antibody, *e.g.*, ipilimumab, for example, to treat a cancer (*e.g.*, a cancer chosen from: a melanoma, *e.g.*, a metastatic melanoma; a lung cancer, *e.g.*, a non-small cell lung carcinoma; or a prostate cancer). In one embodiment, the anti-PD-1 antibody molecule is administered after treatment with an anti-CTLA-4 antibody (*e.g.*, ipilimumab) with or without a BRAF inhibitor (*e.g.*, vemurafenib or dabrafenib).

In another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with an anti-LAG-3 antibody or antigen-binding fragment thereof.

In another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with an anti-TIM-3 antibody or antigen-binding fragment thereof.

In yet other embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with an anti-LAG-3 antibody and an anti-TIM-3 antibody (or antigen-binding fragments thereof).

In another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with a CEACAM inhibitor (*e.g.*, CEACAM-1 and/or CEACAM-5 inhibitor), *e.g.*, an anti-CEACAM antibody molecule. In another embodiment, the anti-PD-1 antibody molecule is administered in combination with a CEACAM-1 inhibitor, *e.g.*, an anti-CEACAM-1 antibody molecule. In another embodiment, the anti-PD-1 antibody molecule is administered in combination with a CEACAM-5 inhibitor, *e.g.*, an anti-CEACAM-5 antibody molecule.

The combination of antibodies recited herein can be administered separately, *e.g.*, as separate antibodies or antigen-binding fragments thereof, or linked, *e.g.*, as a bispecific or trispecific antibody molecule. In one embodiment, a bispecific antibody that includes an anti-PD-1 antibody molecule and an anti-TIM-3, anti-CEACAM (*e.g.*, anti-CEACAM-1, CEACAM-3, and/or anti-CEACAM-5), or anti-LAG-3 antibody, or an antigen-binding fragment thereof, is administered. In certain embodiments, the combination of antibodies recited herein is used to treat a cancer, *e.g.*, a cancer as described herein (*e.g.*, a solid tumor or a hematologic malignancy).

In other embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with a cytokine. The cytokine can be administered as a fusion molecule to the anti-PD-1 antibody molecule, or as separate compositions. In one embodiment, the anti-PD-1 antibody is administered in combination with one, two, three or more cytokines, *e.g.*, as a fusion molecule or as separate compositions. In one embodiment, the cytokine is an interleukin (IL) chosen from one, two, three or more of IL-1, IL-2, IL-12, IL-15 or IL-21. In one embodiment, a bispecific antibody molecule has a first binding specificity to a first target (*e.g.*, to PD-1), a second binding specificity to a second target (*e.g.*, LAG-3 or TIM-3), and is optionally linked to an interleukin (*e.g.*, IL-12) domain *e.g.*, full length IL-12 or a portion thereof. In certain embodiments, the combination of anti-PD-1 antibody molecule and the cytokine described herein is used to treat a cancer, *e.g.*, a cancer as described herein (*e.g.*, a solid tumor).

In certain embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with an antibody specific against an HLA C, *e.g.*, an antibody specific to Killer-cell Immunoglobulin-like Receptors (also referred to herein as an “anti-KIR antibody”). In certain embodiments, the combination of anti-PD-1 antibody molecule and anti-KIR antibody is used to treat a cancer, *e.g.*, a cancer as described herein (*e.g.*, a solid tumor, *e.g.*, an advanced solid tumor).

In one embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with a cellular immunotherapy (*e.g.*, Provenge® (*e.g.*, Sipuleucel-T)), and optionally in combination with cyclophosphamide.

In certain embodiments, the combination of anti-PD-1 antibody molecule, Provenge® and/or

cyclophosphamide is used to treat a cancer, *e.g.*, a cancer as described herein (*e.g.*, a prostate cancer, *e.g.*, an advanced prostate cancer).

In another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with a vaccine, *e.g.*, a cancer vaccine, (*e.g.*, a dendritic cell renal carcinoma (DC-RCC) vaccine). In one embodiment, the vaccine is peptide-based, DNA-based, RNA-based, or antigen-based, or a combination thereof. In embodiments, the vaccine comprises one or more peptides, nucleic acids (*e.g.*, DNA or RNA), antigens, or a combination thereof. In certain embodiments, the combination of anti-PD-1 antibody molecule and the DC-RCC vaccine is used to treat a cancer, *e.g.*, a cancer as described herein (*e.g.*, a renal carcinoma, *e.g.*, metastatic renal cell carcinoma (RCC) or clear cell renal cell carcinoma (CCRCC)).

In another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with an adjuvant.

In yet another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered in combination with chemotherapy, and/or immunotherapy. For example, the anti-PD-1 antibody molecule can be used to treat a myeloma, alone or in combination with one or more of: chemotherapy or other anti-cancer agents (*e.g.*, thalidomide analogs, *e.g.*, lenalidomide), an anti-TIM-3 antibody, tumor antigen-pulsed dendritic cells, fusions (*e.g.*, electrofusions) of tumor cells and dendritic cells, or vaccination with immunoglobulin idiotype produced by malignant plasma cells. In one embodiment, the anti-PD-1 antibody molecule is used in combination with an anti-TIM-3 antibody to treat a myeloma, *e.g.*, a multiple myeloma.

In one embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is used in combination with chemotherapy to treat a lung cancer, *e.g.*, non-small cell lung cancer. In one embodiment, the anti-PD-1 antibody molecule is used with standard lung, *e.g.*, NSCLC, chemotherapy, *e.g.*, platinum doublet therapy, to treat lung cancer. In yet other embodiments, the anti-PD-1 antibody molecule is used in combination with an indoleamine-pyrrole 2,3-dioxygenase (IDO) inhibitor (*e.g.*, (4E)-4-[(3-chloro-4-fluoroanilino)-nitrosomethylidene]-1,2,5-oxadiazol-3-amine (also known as INCB24360), indoximod (1-methyl-D-tryptophan), α -cyclohexyl-5H-Imidazo[5,1-a]isoindole-5-ethanol (also

known as NLG919), etc.) in a subject with advanced or metastatic cancer (*e.g.*, a patient with metastatic and recurrent NSCL cancer).

In yet other embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is used in combination with one or more of: an
5 immune-based strategy (*e.g.*, interleukin-2 or interferon- α), a targeting agent (*e.g.*, a VEGF inhibitor such as a monoclonal antibody to VEGF); a VEGF tyrosine kinase inhibitor such as sunitinib, sorafenib, axitinib and pazopanib; an RNAi inhibitor; or an inhibitor of a downstream mediator of VEGF signaling, *e.g.*, an inhibitor of the mammalian target of rapamycin (mTOR),
10 *e.g.*, everolimus and temsirolimus. Any of such combinations can be used to treat a renal cancer, *e.g.*, renal cell carcinoma (RCC) (*e.g.*, clear cell renal cell carcinoma (CCRCC)) or metastatic RCC.

In some embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is used in combination with a MEK inhibitor (*e.g.*, a MEK inhibitor as described herein). In some embodiments, the combination of the anti-PD-1 antibody
15 and the MEK inhibitor is used to treat a cancer (*e.g.*, a cancer described herein). In some embodiments, the cancer treated with the combination is chosen from a melanoma, a colorectal cancer, a non-small cell lung cancer, an ovarian cancer, a breast cancer, a prostate cancer, a pancreatic cancer, a hematological malignancy or a renal cell carcinoma. In certain
20 embodiments, the cancer includes a BRAF mutation (*e.g.*, a BRAF V600E mutation), a BRAF wildtype, a KRAS wildtype or an activating KRAS mutation. The cancer may be at an early, intermediate or late stage.

In another embodiment, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is used in combination with one, two or all of oxaliplatin, leucovorin or 5-FU (*e.g.*, a FOLFOX co-treatment). Alternatively or in combination,
25 combination further includes a VEGF inhibitor (*e.g.*, a VEGF inhibitor as disclosed herein). In some embodiments, the combination of the anti-PD-1 antibody, the FOLFOX co-treatment, and the VEGF inhibitor is used to treat a cancer (*e.g.*, a cancer described herein). In some
30 embodiments, the cancer treated with the combination is chosen from a melanoma, a colorectal cancer, a non-small cell lung cancer, an ovarian cancer, a breast cancer, a prostate cancer, a pancreatic cancer, a hematological malignancy or a renal cell carcinoma. The cancer may be at an early, intermediate or late stage.

In other embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered with a tyrosine kinase inhibitor (*e.g.*, axitinib) to treat renal cell carcinoma and other solid tumors.

5 In other embodiments, the combination disclosed herein, *e.g.*, a combination comprising an anti-PD-1 antibody molecule, is administered with a 4-1BB receptor targeting agent (*e.g.*, an antibody that stimulates signaling through 4-1BB (CD-137), *e.g.*, PF-2566). In one embodiment, the anti-PD-1 antibody molecule is administered in combination with a tyrosine kinase inhibitor (*e.g.*, axitinib) and a 4-1BB receptor targeting agent.

10 The anti-PD-1 antibody molecule can be bound to a substance, *e.g.*, a cytotoxic agent or moiety (*e.g.*, a therapeutic drug; a compound emitting radiation; molecules of plant, fungal, or bacterial origin; or a biological protein (*e.g.*, a protein toxin) or particle (*e.g.*, a recombinant viral particle, *e.g.*, via a viral coat protein). For example, the antibody can be coupled to a radioactive isotope such as an α -, β -, or γ -emitter, or a β - and γ -emitter.

15 Any combination and sequence of the anti-PD-1 antibody molecules and other therapeutic agents, procedures or modalities (*e.g.*, as described herein) can be used. The antibody molecule and/or other therapeutic agents, procedures or modalities can be administered during periods of active disorder, or during a period of remission or less active disease. The antibody molecule can be administered before the other treatment, concurrently with the treatment, post-treatment, or during remission of the disorder.

20

Additional Combination Therapies

In certain embodiments, any of the combinations disclosed herein further includes one or more of the agents described in Table 7.

25 In some embodiments, the additional therapeutic agent is chosen from one or more of: 1) a protein kinase C (PKC) inhibitor; 2) a heat shock protein 90 (HSP90) inhibitor; 3) an inhibitor of a phosphoinositide 3-kinase (PI3K) and/or target of rapamycin (mTOR); 4) an inhibitor of cytochrome P450 (*e.g.*, a CYP17 inhibitor or a 17 α -Hydroxylase/C17-20 Lyase inhibitor); 5) an iron chelating agent; 6) an aromatase inhibitor; 7) an inhibitor of p53, *e.g.*, an inhibitor of a p53/Mdm2 interaction; 8) an apoptosis inducer; 9) an angiogenesis inhibitor; 10) an aldosterone synthase inhibitor; 11) a smoothened (SMO) receptor inhibitor; 12) a prolactin receptor (PRLR) inhibitor; 13) a Wnt signaling inhibitor; 14) a CDK4/6 inhibitor; 15) a fibroblast growth factor

30

receptor 2 (FGFR2)/fibroblast growth factor receptor 4 (FGFR4) inhibitor; 16) an inhibitor of macrophage colony-stimulating factor (M-CSF); 17) an inhibitor of one or more of c-KIT, histamine release, Flt3 (*e.g.*, FLK2/STK1) or PKC; 18) an inhibitor of one or more of VEGFR-2 (*e.g.*, FLK-1/KDR), PDGFRbeta, c-KIT or Raf kinase C; 19) a somatostatin agonist and/or a growth hormone release inhibitor; 20) an anaplastic lymphoma kinase (ALK) inhibitor; 21) an insulin-like growth factor 1 receptor (IGF-1R) inhibitor; 22) a P-Glycoprotein 1 inhibitor; 23) a vascular endothelial growth factor receptor (VEGFR) inhibitor; 24) a BCR-ABL kinase inhibitor; 25) an FGFR inhibitor; 26) an inhibitor of CYP11B2; 27) a HDM2 inhibitor, *e.g.*, an inhibitor of the HDM2-p53 interaction; 28) an inhibitor of a tyrosine kinase; 29) an inhibitor of c-MET; 30) an inhibitor of JAK; 31) an inhibitor of DAC; 32) an inhibitor of 11 β -hydroxylase; 33) an inhibitor of IAP; 34) an inhibitor of PIM kinase; 35) an inhibitor of Porcupine; 36) an inhibitor of BRAF, *e.g.*, BRAF V600E or wild-type BRAF; 37) an inhibitor of HER3; 38) an inhibitor of MEK; or 39) an inhibitor of a lipid kinase, *e.g.*, as described herein and in Table 7.

In one embodiment, the cancer is chosen from a lung cancer (*e.g.*, a non-small cell lung cancer (NSCLC) (*e.g.*, a NSCLC with squamous and/or non-squamous histology, or a NSCLC adenocarcinoma), or disclosed in a publication listed in Table 7.

Additional embodiments

Additional embodiments provide a method of treating a cancer, comprising: identifying in a subject or a sample (*e.g.*, a subject's sample comprising cancer cells and optionally immune cells such as TILs) the presence of one, two or all of PD-L1, CD8, or IFN- γ , thereby providing a value for one, two or all of PD-L1, CD8, and IFN- γ . The method can further include comparing the PD-L1, CD8, and/or IFN- γ values to a reference value, *e.g.*, a control value. If the PD-L1, CD8, and/or IFN- γ values are greater than the reference value, *e.g.*, the control values, administering a therapeutically effective amount of a combination as described herein (*e.g.*, a combination that includes an anti-PD-1 antibody described herein) to the subject, optionally in combination with one or more other agents, thereby treating the cancer. The cancer may be, *e.g.*, a cancer described herein, such as lung cancer (squamous), lung cancer (adenocarcinoma), head and neck cancer, cervical cancer (squamous), stomach cancer, thyroid cancer, melanoma, nasopharyngeal cancer, or breast cancer, *e.g.*, TN breast cancer, *e.g.*, IM-TN breast cancer. In some embodiments, the cancer is ER+ breast cancer or pancreatic cancer.

Also provided is a method of treating a cancer, comprising: testing a subject or a sample (*e.g.*, a subject's sample comprising cancer cells) for the presence of PD-L1, thereby identifying a PD-L1 value, comparing the PD-L1 value to a control value, and if the PD-L1 value is greater than the control value, administering a therapeutically effective amount of a combination as described herein (*e.g.*, a combination that includes an anti-PD-1 antibody described herein) to the subject, optionally in combination with one or more other agents, thereby treating the cancer. The cancer may be, *e.g.*, a cancer as described herein, such as cancer is non-small cell lung (NSCLC) adenocarcinoma (ACA), NSCLC squamous cell carcinoma (SCC), or hepatocellular carcinoma (HCC).

In another aspect, the invention features diagnostic or therapeutic kits that include the antibody molecules described herein and instructions for use.

All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety.

Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the amino acid sequences of the light and heavy chain variable regions of murine anti-PD-1 mAb BAP049. The upper and lower sequences were from two independent analyses. The light and heavy chain CDR sequences based on Kabat numbering are underlined. The light heavy chain CDR sequences based on Chothia numbering are shown in bold italics. The unpaired Cys residue at position 102 of the light chain sequence is boxed. Sequences are disclosed as SEQ ID NOs: 8, 228, 16 and 229, respectively, in order of appearance.

Figure 2A depicts the amino acid sequences of the light and heavy chain variable regions of murine anti-PD-1 mAb BAP049 aligned with the germline sequences. The upper and lower sequences are the germline (GL) and BAP049 (Mu mAb) sequences, respectively. The light and heavy chain CDR sequences based on Kabat numbering are underlined. The light heavy chain CDR sequences based on Chothia numbering are shown in bold italics. "-" means identical amino acid residue. Sequences disclosed as SEQ ID NOs: 230, 8, 231 and 16, respectively, in order of appearance.

Figure 2B depicts the sequence of murine κ J2 gene and the corresponding mutation in

murine anti-PD-1 mAb BAP049. “-” means identical nucleotide residue. Sequences disclosed as SEQ ID NOs: 233, 232, 234 and 235, respectively, in order of appearance.

Figures 3A-3B depict the competition binding between fluorescently labeled murine anti-PD-1 mAb BAP049 (Mu mAb) and three chimeric versions of BAP049 (Chi mAb).

Experiment was performed twice, and the results are shown in Figures 3A and 3B, respectively. The three chimeric BAP049 antibodies (Chi mAb (Cys), Chi mAb (Tyr) and Chi mAb (Ser)) have Cys, Tyr and Ser residue at position 102 of the light chain variable region, respectively. Chi mAb (Cys), Chi mAb (Tyr) and Chi mAb (Ser) are also known as BAP049-chi, BAP049-chi-Y, and BAP049-chi-S, respectively.

Figure 4 is a bar graph showing the results of FACS binding analysis for the sixteen humanized BAP049 clones (BAP049-hum01 to BAP049-hum16). The antibody concentrations are 200, 100, 50, 25 and 12.5 ng/ml from the leftmost bar to the rightmost bar for each tested mAb.

Figure 5 depicts the structural analysis of the humanized BAP049 clones (a, b, c, d and e represent various types of framework region sequences). The concentrations of the mAbs in the samples are also shown.

Figure 6A-6B depicts the binding affinity and specificity of humanized BAP049 mAbs measured in a competition binding assay using a constant concentration of Alexa 488-labeled murine mAb BAP049, serial dilutions of the test antibodies, and PD-1-expressing 300.19 cells.

Experiment was performed twice, and the results are shown in Figures 6A and 6B, respectively.

Figure 7 depicts the ranking of humanized BAP049 clones based on FACS data, competition binding and structural analysis. The concentrations of the mAbs in the samples are also shown.

Figures 8A-8B depict blocking of ligand binding to PD-1 by selected humanized BAP049 clones. Blocking of PD-L1-Ig and PD-L2-Ig binding to PD-1 is shown in Figure 8A. Blocking of PD-L2-Ig binding to PD-1 is shown in Figure 8B. BAP049-hum01, BAP049-hum05, BAP049-hum08, BAP049-hum09, BAP049-hum10, and BAP049-hum11 were evaluated. Murine mAb BAP049 and chimeric mAb having Tyr at position 102 of the light chain variable region were also included in the analyses.

Figures 9A-9B depict the alignment of heavy chain variable domain sequences for the sixteen humanized BAP049 clones and BAP049 chimera (BAP049-chi). In Figure 9A, all of the

sequences are shown (SEQ ID NOs: 22, 38, 38, 38, 38, 38, 38, 38, 38, 38, 50, 50, 50, 50, 82, 82 and 86, respectively, in order of appearance). In Figure 9B, only amino acid sequences that are different from mouse sequence are shown (SEQ ID NOs: 22, 38, 38, 38, 38, 38, 38, 38, 38, 38, 50, 50, 50, 50, 82, 82 and 86, respectively, in order of appearance).

Figures 10A-10B depict the alignment of light chain variable domain sequences for the sixteen humanized BAP049 clones and BAP049 chimera (BAP049-chi). In Figure 10A, all of the sequences are shown (SEQ ID NOs: 24, 66, 66, 66, 66, 70, 70, 70, 58, 62, 78, 74, 46, 46, 42, 54 and 54, respectively, in order of appearance). In Figure 10B, only amino acid sequences that are different from mouse sequence are shown (SEQ ID NOs: 24, 66, 66, 66, 66, 70, 70, 70, 58, 62, 78, 74, 46, 46, 42, 54 and 54, respectively, in order of appearance).

Figure 11 is a schematic diagram that outlines the antigen processing and presentation, effector cell responses and immunosuppression pathways targeted by the combination therapies disclosed herein.

Figure 12 depicts the predicted Ctrough (Cmin) concentrations across the different weights for patients while receiving the same dose of an exemplary anti-PD-1 antibody molecule.

Figure 13 depicts observed versus model predicted (population or individual based) Cmin concentrations.

Figure 14 depicts the accumulation, time course and within subject variability of the model used to analyze pharmacokinetics.

Figure 15 depicts the amino acid sequence of the light chain of the Antibody Molecule A.

Figure 16 depicts the amino acid sequence of the heavy chain of the Antibody Molecule A.

BRIEF DESCRIPTION OF THE TABLES

Table 1 is a summary of the amino acid and nucleotide sequences for the murine, chimeric and humanized anti-PD-1 antibody molecules. The antibody molecules include murine mAb BAP049, chimeric mAbs BAP049-chi and BAP049-chi-Y, and humanized mAbs BAP049-hum01 to BAP049-hum16 and BAP049-Clone-A to BAP049-Clone-E. The amino acid and nucleotide sequences of the heavy and light chain CDRs, the amino acid and nucleotide

sequences of the heavy and light chain variable regions, and the amino acid and nucleotide sequences of the heavy and light chains are shown in this Table.

Table 2 depicts the amino acid and nucleotide sequences of the heavy and light chain framework regions for humanized mAbs BAP049-hum01 to BAP049-hum16 and BAP049-Clone-A to BAP049-Clone-E.

Table 3 depicts the constant region amino acid sequences of human IgG heavy chains and human kappa light chain.

Table 4 shows the amino acid sequences of the heavy and light chain leader sequences for humanized mAbs BAP049-Clone-A to BAP049-Clone-E.

Table 5 depicts exemplary PK parameters based on flat dosing schedules.

Table 6 depicts the expected disulfide linkages in the Antibody Molecule A.

DETAILED DESCRIPTION

Disclosed herein, at least in part, are antibody molecules (*e.g.*, humanized antibody molecules) that bind to Programmed Death 1 (PD-1) with high affinity and specificity. Nucleic acid molecules encoding the antibody molecules, expression vectors, host cells and methods for making the antibody molecules are also provided. Pharmaceutical compositions and dose formulations comprising the antibody molecules are also provided. The anti-PD-1 antibody molecules disclosed herein can be used (alone or in combination with other agents or therapeutic modalities) to treat, prevent and/or diagnose disorders, such as cancerous disorders (*e.g.*, solid and soft-tissue tumors), as well as infectious diseases (*e.g.*, chronic infectious disorders or sepsis). Thus, compositions and methods for detecting PD-1, as well as methods for treating various disorders including cancer and/or infectious diseases, using the anti-PD-1 antibody molecules are disclosed herein. In certain embodiments, the anti-PD-1 antibody molecule is administered or used at a flat or fixed dose.

Also disclosed herein are methods and compositions comprising a combination of two, three or more therapeutic agents chosen from one, two, or all of the following categories (i)-(iii): (i) an agent that enhances antigen presentation (*e.g.*, tumor antigen presentation) (*e.g.*, by enhancing one or more of dendritic cell activity or maturation, antigen uptake, or antigen processing); (ii) an agent that enhances an effector cell response (*e.g.*, an immune effector cell

response, *e.g.*, B cell and/or T cell activation and/or mobilization, *e.g.*, in the lymph node); or (iii) an agent that decreases tumor immunosuppression (*e.g.*, increasing T cell infiltration and tumor cell killing). In some embodiments, the combination includes a PD-1 inhibitor (*e.g.*, an anti-PD-1 antibody molecule as described herein). Without wishing to be bound by theory, it is believed that therapeutic approaches that enhance anti-tumor immunity work more effectively when the immune response is optimized via multiple targets at different stages of the immune response. Each of these stages is depicted in schematic form in Figure 21. For example, approaches that result in activation of dendritic cells combined with approaches that enhance cellular and humoral immune can result in a more effective and/or prolonged therapeutic response.

Additional terms are defined below and throughout the application.

As used herein, the articles "a" and "an" refer to one or to more than one (*e.g.*, to at least one) of the grammatical object of the article.

The term "or" is used herein to mean, and is used interchangeably with, the term "and/or", unless context clearly indicates otherwise.

"About" and "approximately" shall generally mean an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Exemplary degrees of error are within 20 percent (%), typically, within 10%, and more typically, within 5% of a given value or range of values.

By "a combination" or "in combination with," it is not intended to imply that the therapy or the therapeutic agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope described herein. The therapeutic agents in the combination can be administered concurrently with, prior to, or subsequent to, one or more other additional therapies or therapeutic agents. The therapeutic agents or therapeutic protocol can be administered in any order. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. It will further be appreciated that the additional therapeutic agent utilized in this combination may be administered together in a single composition or administered separately in different compositions. In general, it is expected that additional therapeutic agents utilized in combination be utilized at levels that

do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination will be lower than those utilized individually.

In embodiments, the additional therapeutic agent is administered at a therapeutic or lower-than therapeutic dose. In certain embodiments, the concentration of the second therapeutic agent that is required to achieve inhibition, *e.g.*, growth inhibition, is lower when the second therapeutic agent is administered in combination with the first therapeutic agent, *e.g.*, the anti-PD-1 antibody molecule, than when the second therapeutic agent is administered individually. In certain embodiments, the concentration of the first therapeutic agent that is required to achieve inhibition, *e.g.*, growth inhibition, is lower when the first therapeutic agent is administered in combination with the second therapeutic agent than when the first therapeutic agent is administered individually. In certain embodiments, in a combination therapy, the concentration of the second therapeutic agent that is required to achieve inhibition, *e.g.*, growth inhibition, is lower than the therapeutic dose of the second therapeutic agent as a monotherapy, *e.g.*, 10-20%, 20-30%, 30-40%, 40-50%, 50-60%, 60-70%, 70-80%, or 80-90% lower. In certain embodiments, in a combination therapy, the concentration of the first therapeutic agent that is required to achieve inhibition, *e.g.*, growth inhibition, is lower than the therapeutic dose of the first therapeutic agent as a monotherapy, *e.g.*, 10-20%, 20-30%, 30-40%, 40-50%, 50-60%, 60-70%, 70-80%, or 80-90% lower.

The term “inhibition,” “inhibitor,” or “antagonist” includes a reduction in a certain parameter, *e.g.*, an activity, of a given molecule, *e.g.*, an immune checkpoint inhibitor. For example, inhibition of an activity, *e.g.*, a PD-1 or PD-L1 activity, of at least 5%, 10%, 20%, 30%, 40% or more is included by this term. Thus, inhibition need not be 100%.

The term “activation,” “activator,” or “agonist” includes an increase in a certain parameter, *e.g.*, an activity, of a given molecule, *e.g.*, a costimulatory molecule. For example, increase of an activity, *e.g.*, a costimulatory activity, of at least 5%, 10%, 25%, 50%, 75% or more is included by this term.

The term “anti-cancer effect” refers to a biological effect which can be manifested by various means, including but not limited to, *e.g.*, a decrease in tumor volume, a decrease in the number of cancer cells, a decrease in the number of metastases, an increase in life expectancy, decrease in cancer cell proliferation, decrease in cancer cell survival, or amelioration of various physiological symptoms associated with the cancerous condition. An “anti-cancer effect” can

also be manifested by the ability of the peptides, polynucleotides, cells and antibodies in prevention of the occurrence of cancer in the first place.

The term “anti-tumor effect” refers to a biological effect which can be manifested by various means, including but not limited to, *e.g.*, a decrease in tumor volume, a decrease in the
 5 number of tumor cells, a decrease in tumor cell proliferation, or a decrease in tumor cell survival.

The term “cancer” refers to a disease characterized by the rapid and uncontrolled growth of aberrant cells. Cancer cells can spread locally or through the bloodstream and lymphatic system to other parts of the body. Examples of various cancers are described herein and include but are not limited to, breast cancer, prostate cancer, ovarian cancer, cervical cancer, skin cancer,
 10 pancreatic cancer, colorectal cancer, renal cancer, liver cancer, brain cancer, lymphoma, leukemia, lung cancer and the like. The terms “tumor” and “cancer” are used interchangeably herein, *e.g.*, both terms encompass solid and liquid, *e.g.*, diffuse or circulating, tumors. As used herein, the term “cancer” or “tumor” includes premalignant, as well as malignant cancers and tumors.

15 The term “antigen presenting cell” or “APC” refers to an immune system cell such as an accessory cell (*e.g.*, a B-cell, a dendritic cell, and the like) that displays a foreign antigen complexed with major histocompatibility complexes (MHC's) on its surface. T-cells may recognize these complexes using their T-cell receptors (TCRs). APCs process antigens and present them to T-cells.

20 The term “costimulatory molecule” refers to the cognate binding partner on a T cell that specifically binds with a costimulatory ligand, thereby mediating a costimulatory response by the T cell, such as, but not limited to, proliferation. Costimulatory molecules are cell surface molecules other than antigen receptors or their ligands that are required for an efficient immune response. Costimulatory molecules include, but are not limited to, an MHC class I molecule,
 25 TNF receptor proteins, Immunoglobulin-like proteins, cytokine receptors, integrins, signaling lymphocytic activation molecules (SLAM proteins), activating NK cell receptors, BTLA, a Toll ligand receptor, OX40, CD2, CD7, CD27, CD28, CD30, CD40, CDS, ICAM-1, LFA-1 (CD11a/CD18), 4-1BB (CD137), B7-H3, CDS, ICAM-1, ICOS (CD278), GITR, BAFR, LIGHT, HVEM (LIGHTR), KIRDS2, SLAMF7, NKp80 (KLRF1), NKp44, NKp30, NKp46,
 30 CD19, CD4, CD8alpha, CD8beta, IL2R beta, IL2R gamma, IL7R alpha, ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD11d, ITGAE, CD103, ITGAL,

CD11a, LFA-1, ITGAM, CD11b, ITGAX, CD11c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, NKG2D, NKG2C, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRTAM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3),
5 BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, PAG/Cbp, CD19a, and a ligand that specifically binds with CD83.

“Immune effector cell,” or “effector cell” as that term is used herein, refers to a cell that is involved in an immune response, *e.g.*, in the promotion of an immune effector response. Examples of immune effector cells include T cells, *e.g.*, alpha/beta T cells and gamma/delta T
10 cells, B cells, natural killer (NK) cells, natural killer T (NKT) cells, mast cells, and myeloid-derived phagocytes.

“Immune effector” or “effector” “function” or “response,” as that term is used herein, refers to function or response, *e.g.*, of an immune effector cell, that enhances or promotes an immune attack of a target cell. *E.g.*, an immune effector function or response refers a property of
15 a T or NK cell that promotes killing or the inhibition of growth or proliferation, of a target cell. In the case of a T cell, primary stimulation and co-stimulation are examples of immune effector function or response.

The term “effector function” refers to a specialized function of a cell. Effector function of a T cell, for example, may be cytolytic activity or helper activity including the
20 secretion of cytokines.

As used herein, the terms “treat”, “treatment” and “treating” refer to the reduction or amelioration of the progression, severity and/or duration of a disorder, *e.g.*, a proliferative disorder, or the amelioration of one or more symptoms (preferably, one or more discernible symptoms) of the disorder resulting from the administration of one or more therapies. In specific
25 embodiments, the terms “treat,” “treatment” and “treating” refer to the amelioration of at least one measurable physical parameter of a proliferative disorder, such as growth of a tumor, not necessarily discernible by the patient. In other embodiments the terms “treat”, “treatment” and “treating” -refer to the inhibition of the progression of a proliferative disorder, either physically by, *e.g.*, stabilization of a discernible symptom, physiologically by, *e.g.*, stabilization of a
30 physical parameter, or both. In other embodiments the terms “treat”, “treatment” and “treating” refer to the reduction or stabilization of tumor size or cancerous cell count.

The compositions and methods of the present invention encompass polypeptides and nucleic acids having the sequences specified, or sequences substantially identical or similar thereto, *e.g.*, sequences at least 85%, 90%, 95% identical or higher to the sequence specified. In the context of an amino acid sequence, the term "substantially identical" is used herein to refer to a first amino acid that contains a sufficient or minimum number of amino acid residues that are i) identical to, or ii) conservative substitutions of aligned amino acid residues in a second amino acid sequence such that the first and second amino acid sequences can have a common structural domain and/or common functional activity. For example, amino acid sequences that contain a common structural domain having at least about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a reference sequence, *e.g.*, a sequence provided herein.

In the context of nucleotide sequence, the term "substantially identical" is used herein to refer to a first nucleic acid sequence that contains a sufficient or minimum number of nucleotides that are identical to aligned nucleotides in a second nucleic acid sequence such that the first and second nucleotide sequences encode a polypeptide having common functional activity, or encode a common structural polypeptide domain or a common functional polypeptide activity. For example, nucleotide sequences having at least about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identity to a reference sequence, *e.g.*, a sequence provided herein.

The term "functional variant" refers to polypeptides that have a substantially identical amino acid sequence to the naturally-occurring sequence, or are encoded by a substantially identical nucleotide sequence, and are capable of having one or more activities of the naturally-occurring sequence.

Calculations of homology or sequence identity between sequences (the terms are used interchangeably herein) are performed as follows.

To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues

or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or
5 nucleic acid "homology").

The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two
10 sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch ((1970) *J. Mol. Biol.* 48:444-453) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a
15 length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A particularly preferred set of parameters (and the one that should be used unless otherwise specified) are a Blossum 62 scoring
20 matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller ((1989) *CABIOS*, 4:11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

25 The nucleic acid and protein sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12
30 to obtain nucleotide sequences homologous to a nucleic acid (SEQ ID NO: 1) molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50,

wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

As used herein, the term "hybridizes under low stringency, medium stringency, high stringency, or very high stringency conditions" describes conditions for hybridization and washing. Guidance for performing hybridization reactions can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6, which is incorporated by reference. Aqueous and nonaqueous methods are described in that reference and either can be used. Specific hybridization conditions referred to herein are as follows: 1) low stringency hybridization conditions in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by two washes in 0.2X SSC, 0.1% SDS at least at 50°C (the temperature of the washes can be increased to 55°C for low stringency conditions); 2) medium stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C; 3) high stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65°C; and preferably 4) very high stringency hybridization conditions are 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C. Very high stringency conditions (4) are the preferred conditions and the ones that should be used unless otherwise specified.

It is understood that the molecules of the present invention may have additional conservative or non-essential amino acid substitutions, which do not have a substantial effect on their functions.

The term "amino acid" is intended to embrace all molecules, whether natural or synthetic, which include both an amino functionality and an acid functionality and capable of being included in a polymer of naturally-occurring amino acids. Exemplary amino acids include naturally-occurring amino acids; analogs, derivatives and congeners thereof; amino acid analogs having variant side chains; and all stereoisomers of any of any of the foregoing. As used herein the term "amino acid" includes both the D- or L- optical isomers and peptidomimetics.

A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues

having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

The terms "polypeptide", "peptide" and "protein" (if single chain) are used interchangeably herein to refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation, such as conjugation with a labeling component. The polypeptide can be isolated from natural sources, can be produced by recombinant techniques from a eukaryotic or prokaryotic host, or can be a product of synthetic procedures.

The terms "nucleic acid," "nucleic acid sequence," "nucleotide sequence," or "polynucleotide sequence," and "polynucleotide" are used interchangeably. They refer to a polymeric form of nucleotides of any length, either deoxyribonucleotides or ribonucleotides, or analogs thereof. The polynucleotide may be either single-stranded or double-stranded, and if single-stranded may be the coding strand or non-coding (antisense) strand. A polynucleotide may comprise modified nucleotides, such as methylated nucleotides and nucleotide analogs. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. The nucleic acid may be a recombinant polynucleotide, or a polynucleotide of genomic, cDNA, semisynthetic, or synthetic origin which either does not occur in nature or is linked to another polynucleotide in a nonnatural arrangement.

The term "isolated," as used herein, refers to material that is removed from its original or native environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated by human intervention from some or all of the co-existing materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be

isolated in that such vector or composition is not part of the environment in which it is found in nature.

Various aspects of the invention are described in further detail below. Additional definitions are set out throughout the specification.

5

Antibody Molecules

In one embodiment, the antibody molecule binds to a mammalian, *e.g.*, human, PD-1. For example, the antibody molecule binds specifically to an epitope, *e.g.*, linear or conformational epitope, (*e.g.*, an epitope as described herein) on PD-1.

10 As used herein, the term "antibody molecule" refers to a protein, *e.g.*, an immunoglobulin chain or fragment thereof, comprising at least one immunoglobulin variable domain sequence. The term "antibody molecule" includes, for example, a monoclonal antibody (including a full length antibody which has an immunoglobulin Fc region). In an embodiment, an antibody molecule comprises a full length antibody, or a full length immunoglobulin chain. In an
15 embodiment, an antibody molecule comprises an antigen binding or functional fragment of a full length antibody, or a full length immunoglobulin chain. In an embodiment, an antibody molecule is a multispecific antibody molecule, *e.g.*, it comprises a plurality of immunoglobulin variable domain sequences, wherein a first immunoglobulin variable domain sequence of the plurality has binding specificity for a first epitope and a second immunoglobulin variable domain
20 sequence of the plurality has binding specificity for a second epitope. In an embodiment, a multispecific antibody molecule is a bispecific antibody molecule. A bispecific antibody has specificity for no more than two antigens. A bispecific antibody molecule is characterized by a first immunoglobulin variable domain sequence which has binding specificity for a first epitope and a second immunoglobulin variable domain sequence that has binding specificity for a second
25 epitope.

In an embodiment, an antibody molecule is a monospecific antibody molecule and binds a single epitope. *E.g.*, a monospecific antibody molecule having a plurality of immunoglobulin variable domain sequences, each of which binds the same epitope.

In an embodiment an antibody molecule is a multispecific antibody molecule, *e.g.*, it
30 comprises a plurality of immunoglobulin variable domains sequences, wherein a first immunoglobulin variable domain sequence of the plurality has binding specificity for a first

epitope and a second immunoglobulin variable domain sequence of the plurality has binding specificity for a second epitope. In an embodiment the first and second epitopes are on the same antigen, *e.g.*, the same protein (or subunit of a multimeric protein). In an embodiment the first and second epitopes overlap. In an embodiment the first and second epitopes do not overlap. In an embodiment the first and second epitopes are on different antigens, *e.g.*, the different proteins (or different subunits of a multimeric protein). In an embodiment a multispecific antibody molecule comprises a third, fourth or fifth immunoglobulin variable domain. In an embodiment, a multispecific antibody molecule is a bispecific antibody molecule, a trispecific antibody molecule, or tetraspecific antibody molecule,

In an embodiment a multispecific antibody molecule is a bispecific antibody molecule. A bispecific antibody has specificity for no more than two antigens. A bispecific antibody molecule is characterized by a first immunoglobulin variable domain sequence which has binding specificity for a first epitope and a second immunoglobulin variable domain sequence that has binding specificity for a second epitope. In an embodiment the first and second epitopes are on the same antigen, *e.g.*, the same protein (or subunit of a multimeric protein). In an embodiment the first and second epitopes overlap. In an embodiment the first and second epitopes do not overlap. In an embodiment the first and second epitopes are on different antigens, *e.g.*, the different proteins (or different subunits of a multimeric protein). In an embodiment a bispecific antibody molecule comprises a heavy chain variable domain sequence and a light chain variable domain sequence which have binding specificity for a first epitope and a heavy chain variable domain sequence and a light chain variable domain sequence which have binding specificity for a second epitope. In an embodiment a bispecific antibody molecule comprises a half antibody having binding specificity for a first epitope and a half antibody having binding specificity for a second epitope. In an embodiment a bispecific antibody molecule comprises a half antibody, or fragment thereof, having binding specificity for a first epitope and a half antibody, or fragment thereof, having binding specificity for a second epitope. In an embodiment a bispecific antibody molecule comprises a scFv, or fragment thereof, have binding specificity for a first epitope and a scFv, or fragment thereof, have binding specificity for a second epitope. In an embodiment the first epitope is located on PD-1 and the second epitope is located on a TIM-3, LAG-3, CEACAM (*e.g.*, CEACAM-1 and/or CEACAM-5), PD-L1, or PD-L2.

In an embodiment, an antibody molecule comprises a diabody, and a single-chain molecule, as well as an antigen-binding fragment of an antibody (*e.g.*, Fab, F(ab')₂, and Fv). For example, an antibody molecule can include a heavy (H) chain variable domain sequence (abbreviated herein as VH), and a light (L) chain variable domain sequence (abbreviated herein as VL). In an embodiment an antibody molecule comprises or consists of a heavy chain and a light chain (referred to herein as a half antibody. In another example, an antibody molecule includes two heavy (H) chain variable domain sequences and two light (L) chain variable domain sequence, thereby forming two antigen binding sites, such as Fab, Fab', F(ab')₂, Fc, Fd, Fd', Fv, single chain antibodies (scFv for example), single variable domain antibodies, diabodies (Dab) (bivalent and bispecific), and chimeric (*e.g.*, humanized) antibodies, which may be produced by the modification of whole antibodies or those synthesized *de novo* using recombinant DNA technologies. These functional antibody fragments retain the ability to selectively bind with their respective antigen or receptor. Antibodies and antibody fragments can be from any class of antibodies including, but not limited to, IgG, IgA, IgM, IgD, and IgE, and from any subclass (*e.g.*, IgG1, IgG2, IgG3, and IgG4) of antibodies. The preparation of antibody molecules can be monoclonal or polyclonal. An antibody molecule can also be a human, humanized, CDR-grafted, or *in vitro* generated antibody. The antibody can have a heavy chain constant region chosen from, *e.g.*, IgG1, IgG2, IgG3, or IgG4. The antibody can also have a light chain chosen from, *e.g.*, kappa or lambda. The term "immunoglobulin" (Ig) is used interchangeably with the term "antibody" herein.

Examples of antigen-binding fragments of an antibody molecule include: (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a diabody (dAb) fragment, which consists of a VH domain; (vi) a camelid or camelized variable domain; (vii) a single chain Fv (scFv), *see e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883); (viii) a single domain antibody. These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies.

The term "antibody" includes intact molecules as well as functional fragments thereof. Constant regions of the antibodies can be altered, *e.g.*, mutated, to modify the properties of the antibody (*e.g.*, to increase or decrease one or more of: Fc receptor binding, antibody glycosylation, the number of cysteine residues, effector cell function, or complement function).

5 Antibody molecules can also be single domain antibodies. Single domain antibodies can include antibodies whose complementary determining regions are part of a single domain polypeptide. Examples include, but are not limited to, heavy chain antibodies, antibodies naturally devoid of light chains, single domain antibodies derived from conventional 4-chain antibodies, engineered antibodies and single domain scaffolds other than those derived from
10 antibodies. Single domain antibodies may be any of the art, or any future single domain antibodies. Single domain antibodies may be derived from any species including, but not limited to mouse, human, camel, llama, fish, shark, goat, rabbit, and bovine. According to another aspect of the invention, a single domain antibody is a naturally occurring single domain antibody known as heavy chain antibody devoid of light chains. Such single domain antibodies are
15 disclosed in WO 9404678, for example. For clarity reasons, this variable domain derived from a heavy chain antibody naturally devoid of light chain is known herein as a VHH or nanobody to distinguish it from the conventional VH of four chain immunoglobulins. Such a VHH molecule can be derived from antibodies raised in *Camelidae* species, for example in camel, llama, dromedary, alpaca and guanaco. Other species besides *Camelidae* may produce heavy chain
20 antibodies naturally devoid of light chain; such VHHs are within the scope of the invention.

The VH and VL regions can be subdivided into regions of hypervariability, termed "complementarity determining regions" (CDR), interspersed with regions that are more conserved, termed "framework regions" (FR or FW).

The extent of the framework region and CDRs has been precisely defined by a number of
25 methods (*see*, Kabat, E. A., *et al.* (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Chothia, C. *et al.* (1987) *J. Mol. Biol.* 196:901-917; and the AbM definition used by Oxford Molecular's AbM antibody modeling software. *See*, generally, *e.g.*, *Protein Sequence and Structure Analysis of Antibody Variable Domains*. In: Antibody Engineering Lab Manual (Ed.:
30 Duebel, S. and Kontermann, R., Springer-Verlag, Heidelberg).

The terms “complementarity determining region,” and “CDR,” as used herein refer to the sequences of amino acids within antibody variable regions which confer antigen specificity and binding affinity. In general, there are three CDRs in each heavy chain variable region (HCDR1, HCDR2, HCDR3) and three CDRs in each light chain variable region (LCDR1, LCDR2,
5 LCDR3).

The precise amino acid sequence boundaries of a given CDR can be determined using any of a number of well-known schemes, including those described by Kabat *et al.* (1991), “Sequences of Proteins of Immunological Interest,” 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (“Kabat” numbering scheme), Al-Lazikani *et al.*, (1997) *JMB*
10 273,927-948 (“Chothia” numbering scheme). As used herein, the CDRs defined according the “Chothia” number scheme are also sometimes referred to as “hypervariable loops.”

For example, under Kabat, the CDR amino acid residues in the heavy chain variable domain (VH) are numbered 31-35 (HCDR1), 50-65 (HCDR2), and 95-102 (HCDR3); and the CDR amino acid residues in the light chain variable domain (VL) are numbered 24-34 (LCDR1),
15 50-56 (LCDR2), and 89-97 (LCDR3). Under Chothia the CDR amino acids in the VH are numbered 26-32 (HCDR1), 52-56 (HCDR2), and 95-102 (HCDR3); and the amino acid residues in VL are numbered 26-32 (LCDR1), 50-52 (LCDR2), and 91-96 (LCDR3). By combining the CDR definitions of both Kabat and Chothia, the CDRs consist of amino acid residues 26-35 (HCDR1), 50-65 (HCDR2), and 95-102 (HCDR3) in human VH and amino acid residues 24-34
20 (LCDR1), 50-56 (LCDR2), and 89-97 (LCDR3) in human VL.

Generally, unless specifically indicated, the anti-PD-1 antibody molecules can include any combination of one or more Kabat CDRs and/or Chothia hypervariable loops, *e.g.*, described in Table 1. In one embodiment, the following definitions are used for the anti-PD-1 antibody molecules described in Table 1: HCDR1 according to the combined CDR definitions of both
25 Kabat and Chothia, and HCCDRs 2-3 and LCCDRs 1-3 according the CDR definition of Kabat. Under all definitions, each VH and VL typically includes three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

As used herein, an “immunoglobulin variable domain sequence” refers to an amino acid
30 sequence which can form the structure of an immunoglobulin variable domain. For example, the sequence may include all or part of the amino acid sequence of a naturally-occurring variable

domain. For example, the sequence may or may not include one, two, or more N- or C-terminal amino acids, or may include other alterations that are compatible with formation of the protein structure.

The term "antigen-binding site" refers to the part of an antibody molecule that comprises determinants that form an interface that binds to the PD-1 polypeptide, or an epitope thereof. With respect to proteins (or protein mimetics), the antigen-binding site typically includes one or more loops (of at least four amino acids or amino acid mimics) that form an interface that binds to the PD-1 polypeptide. Typically, the antigen-binding site of an antibody molecule includes at least one or two CDRs and/or hypervariable loops, or more typically at least three, four, five or six CDRs and/or hypervariable loops.

The terms "compete" or "cross-compete" are used interchangeably herein to refer to the ability of an antibody molecule to interfere with binding of an anti-PD-1 antibody molecule, *e.g.*, an anti-PD-1 antibody molecule provided herein, to a target, *e.g.*, human PD-1. The interference with binding can be direct or indirect (*e.g.*, through an allosteric modulation of the antibody molecule or the target). The extent to which an antibody molecule is able to interfere with the binding of another antibody molecule to the target, and therefore whether it can be said to compete, can be determined using a competition binding assay, for example, a FACS assay, an ELISA or BIACORE assay. In some embodiments, a competition binding assay is a quantitative competition assay. In some embodiments, a first anti-PD-1 antibody molecule is said to compete for binding to the target with a second anti-PD-1 antibody molecule when the binding of the first antibody molecule to the target is reduced by 10% or more, *e.g.*, 20% or more, 30% or more, 40% or more, 50% or more, 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more, 98% or more, 99% or more in a competition binding assay (*e.g.*, a competition assay described herein).

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope. A monoclonal antibody can be made by hybridoma technology or by methods that do not use hybridoma technology (*e.g.*, recombinant methods).

An "effectively human" protein is a protein that does not evoke a neutralizing antibody response, *e.g.*, the human anti-murine antibody (HAMA) response. HAMA can be problematic

in a number of circumstances, *e.g.*, if the antibody molecule is administered repeatedly, *e.g.*, in treatment of a chronic or recurrent disease condition. A HAMA response can make repeated antibody administration potentially ineffective because of an increased antibody clearance from the serum (*see, e.g.*, Saleh *et al.*, *Cancer Immunol. Immunother.*, 32:180-190 (1990)) and also
5 because of potential allergic reactions (*see, e.g.*, LoBuglio *et al.*, *Hybridoma*, 5:5117-5123 (1986)).

The antibody molecule can be a polyclonal or a monoclonal antibody. In other embodiments, the antibody can be recombinantly produced, *e.g.*, produced by phage display or by combinatorial methods.

10 Phage display and combinatorial methods for generating antibodies are known in the art (as described in, *e.g.*, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* International Publication No. WO 92/18619; Dower *et al.* International Publication No. WO 91/17271; Winter *et al.* International Publication WO 92/20791; Markland *et al.* International Publication No. WO 92/15679; Breitling *et al.* International Publication WO 93/01288; McCafferty *et al.*
15 International Publication No. WO 92/01047; Garrard *et al.* International Publication No. WO 92/09690; Ladner *et al.* International Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982, the contents of all of which are incorporated by reference herein).

In one embodiment, the antibody is a fully human antibody (*e.g.*, an antibody made in a mouse which has been genetically engineered to produce an antibody from a human
25 immunoglobulin sequence), or a non-human antibody, *e.g.*, a rodent (mouse or rat), goat, primate (*e.g.*, monkey), camel antibody. Preferably, the non-human antibody is a rodent (mouse or rat antibody). Methods of producing rodent antibodies are known in the art.

Human monoclonal antibodies can be generated using transgenic mice carrying the human immunoglobulin genes rather than the mouse system. Splenocytes from these transgenic
30 mice immunized with the antigen of interest are used to produce hybridomas that secrete human mAbs with specific affinities for epitopes from a human protein (*see, e.g.*, Wood *et al.*

International Application WO 91/00906, Kucherlapati *et al.* PCT publication WO 91/10741; Lonberg *et al.* International Application WO 92/03918; Kay *et al.* International Application 92/03917; Lonberg, N. *et al.* 1994 *Nature* 368:856-859; Green, L.L. *et al.* 1994 *Nature Genet.* 7:13-21; Morrison, S.L. *et al.* 1994 *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Bruggeman *et al.* 5 1993 *Year Immunol* 7:33-40; Tuailon *et al.* 1993 *PNAS* 90:3720-3724; Bruggeman *et al.* 1991 *Eur J Immunol* 21:1323-1326).

An antibody can be one in which the variable region, or a portion thereof, *e.g.*, the CDRs, are generated in a non-human organism, *e.g.*, a rat or mouse. Chimeric, CDR-grafted, and humanized antibodies are within the invention. Antibodies generated in a non-human organism, 10 *e.g.*, a rat or mouse, and then modified, *e.g.*, in the variable framework or constant region, to decrease antigenicity in a human are within the invention.

Chimeric antibodies can be produced by recombinant DNA techniques known in the art (see Robinson *et al.*, International Patent Publication PCT/US86/02269; Akira, *et al.*, European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison *et al.*, European Patent Application 173,494; Neuberger *et al.*, International Application WO 15 86/01533; Cabilly *et al.* U.S. Patent No. 4,816,567; Cabilly *et al.*, European Patent Application 125,023; Better *et al.* (1988 *Science* 240:1041-1043); Liu *et al.* (1987) *PNAS* 84:3439-3443; Liu *et al.*, 1987, *J. Immunol.* 139:3521-3526; Sun *et al.* (1987) *PNAS* 84:214-218; Nishimura *et al.*, 1987, *Canc. Res.* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; and Shaw *et al.*, 1988, 20 *J. Natl Cancer Inst.* 80:1553-1559).

A humanized or CDR-grafted antibody will have at least one or two but generally all three recipient CDRs (of heavy and or light immunoglobulin chains) replaced with a donor CDR. The antibody may be replaced with at least a portion of a non-human CDR or only some of the CDRs may be replaced with non-human CDRs. It is only necessary to replace the number of 25 CDRs required for binding of the humanized antibody to PD-1. Preferably, the donor will be a rodent antibody, *e.g.*, a rat or mouse antibody, and the recipient will be a human framework or a human consensus framework. Typically, the immunoglobulin providing the CDRs is called the "donor" and the immunoglobulin providing the framework is called the "acceptor." In one embodiment, the donor immunoglobulin is a non-human (*e.g.*, rodent). The acceptor framework 30 is a naturally-occurring (*e.g.*, a human) framework or a consensus framework, or a sequence about 85% or higher, preferably 90%, 95%, 99% or higher identical thereto.

As used herein, the term "consensus sequence" refers to the sequence formed from the most frequently occurring amino acids (or nucleotides) in a family of related sequences (See *e.g.*, Winnaker, *From Genes to Clones* (Verlagsgesellschaft, Weinheim, Germany 1987). In a family of proteins, each position in the consensus sequence is occupied by the amino acid occurring most frequently at that position in the family. If two amino acids occur equally frequently, either can be included in the consensus sequence. A "consensus framework" refers to the framework region in the consensus immunoglobulin sequence.

An antibody can be humanized by methods known in the art (*see e.g.*, Morrison, S. L., 1985, *Science* 229:1202-1207, by Oi *et al.*, 1986, *BioTechniques* 4:214, and by Queen *et al.* US 5,585,089, US 5,693,761 and US 5,693,762, the contents of all of which are hereby incorporated by reference).

Humanized or CDR-grafted antibodies can be produced by CDR-grafting or CDR substitution, wherein one, two, or all CDRs of an immunoglobulin chain can be replaced. *See e.g.*, U.S. Patent 5,225,539; Jones *et al.* 1986 *Nature* 321:552-525; Verhoeyan *et al.* 1988 *Science* 239:1534; Beidler *et al.* 1988 *J. Immunol.* 141:4053-4060; Winter US 5,225,539, the contents of all of which are hereby expressly incorporated by reference. Winter describes a CDR-grafting method which may be used to prepare the humanized antibodies of the present invention (UK Patent Application GB 2188638A, filed on March 26, 1987; Winter US 5,225,539), the contents of which is expressly incorporated by reference.

Also within the scope of the invention are humanized antibodies in which specific amino acids have been substituted, deleted or added. Criteria for selecting amino acids from the donor are described in US 5,585,089, *e.g.*, columns 12-16 of US 5,585,089, *e.g.*, columns 12-16 of US 5,585,089, the contents of which are hereby incorporated by reference. Other techniques for humanizing antibodies are described in Padlan *et al.* EP 519596 A1, published on December 23, 1992.

The antibody molecule can be a single chain antibody. A single-chain antibody (scFV) may be engineered (*see, for example*, Colcher, D. *et al.* (1999) *Ann N Y Acad Sci* 880:263-80; and Reiter, Y. (1996) *Clin Cancer Res* 2:245-52). The single chain antibody can be dimerized or multimerized to generate multivalent antibodies having specificities for different epitopes of the same target protein.

In yet other embodiments, the antibody molecule has a heavy chain constant region chosen from, *e.g.*, the heavy chain constant regions of IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgD, and IgE; particularly, chosen from, *e.g.*, the (*e.g.*, human) heavy chain constant regions of IgG1, IgG2, IgG3, and IgG4. In another embodiment, the antibody molecule has a light chain constant region chosen from, *e.g.*, the (*e.g.*, human) light chain constant regions of kappa or lambda. The constant region can be altered, *e.g.*, mutated, to modify the properties of the antibody (*e.g.*, to increase or decrease one or more of: Fc receptor binding, antibody glycosylation, the number of cysteine residues, effector cell function, and/or complement function). In one embodiment the antibody has: effector function; and can fix complement. In other embodiments the antibody does not; recruit effector cells; or fix complement. In another embodiment, the antibody has reduced or no ability to bind an Fc receptor. For example, it is a isotype or subtype, fragment or other mutant, which does not support binding to an Fc receptor, *e.g.*, it has a mutagenized or deleted Fc receptor binding region.

Methods for altering an antibody constant region are known in the art. Antibodies with altered function, *e.g.* altered affinity for an effector ligand, such as FcR on a cell, or the C1 component of complement can be produced by replacing at least one amino acid residue in the constant portion of the antibody with a different residue (*see e.g.*, EP 388,151 A1, U.S. Pat. No. 5,624,821 and U.S. Pat. No. 5,648,260, the contents of all of which are hereby incorporated by reference). Similar type of alterations could be described which if applied to the murine, or other species immunoglobulin would reduce or eliminate these functions.

An antibody molecule can be derivatized or linked to another functional molecule (*e.g.*, another peptide or protein). As used herein, a "derivatized" antibody molecule is one that has been modified. Methods of derivatization include but are not limited to the addition of a fluorescent moiety, a radionucleotide, a toxin, an enzyme or an affinity ligand such as biotin. Accordingly, the antibody molecules of the invention are intended to include derivatized and otherwise modified forms of the antibodies described herein, including immunoadhesion molecules. For example, an antibody molecule can be functionally linked (by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other molecular entities, such as another antibody (*e.g.*, a bispecific antibody or a diabody), a detectable agent, a cytotoxic agent, a pharmaceutical agent, and/or a protein or peptide that can mediate association of the

antibody or antibody portion with another molecule (such as a streptavidin core region or a polyhistidine tag).

One type of derivatized antibody molecule is produced by crosslinking two or more antibodies (of the same type or of different types, *e.g.*, to create bispecific antibodies). Suitable crosslinkers include those that are heterobifunctional, having two distinctly reactive groups separated by an appropriate spacer (*e.g.*, *m*-maleimidobenzoyl-*N*-hydroxysuccinimide ester) or homobifunctional (*e.g.*, disuccinimidyl suberate). Such linkers are available from Pierce Chemical Company, Rockford, Ill.

Useful detectable agents with which an antibody molecule of the invention may be derivatized (or labeled) to include fluorescent compounds, various enzymes, prosthetic groups, luminescent materials, bioluminescent materials, fluorescent emitting metal atoms, *e.g.*, europium (Eu), and other anthanides, and radioactive materials (described below). Exemplary fluorescent detectable agents include fluorescein, fluorescein isothiocyanate, rhodamine, 5dimethylamine-1-naphthalenesulfonyl chloride, phycoerythrin and the like. An antibody may also be derivatized with detectable enzymes, such as alkaline phosphatase, horseradish peroxidase, β -galactosidase, acetylcholinesterase, glucose oxidase and the like. When an antibody is derivatized with a detectable enzyme, it is detected by adding additional reagents that the enzyme uses to produce a detectable reaction product. For example, when the detectable agent horseradish peroxidase is present, the addition of hydrogen peroxide and diaminobenzidine leads to a colored reaction product, which is detectable. An antibody molecule may also be derivatized with a prosthetic group (*e.g.*, streptavidin/biotin and avidin/biotin). For example, an antibody may be derivatized with biotin, and detected through indirect measurement of avidin or streptavidin binding. Examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of bioluminescent materials include luciferase, luciferin, and aequorin.

Labeled antibody molecule can be used, for example, diagnostically and/or experimentally in a number of contexts, including (i) to isolate a predetermined antigen by standard techniques, such as affinity chromatography or immunoprecipitation; (ii) to detect a predetermined antigen (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the

abundance and pattern of expression of the protein; (iii) to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to determine the efficacy of a given treatment regimen.

An antibody molecules may be conjugated to another molecular entity, typically a label or a therapeutic (*e.g.*, a cytotoxic or cytostatic) agent or moiety. Radioactive isotopes can be used in diagnostic or therapeutic applications.

The invention provides radiolabeled antibody molecules and methods of labeling the same. In one embodiment, a method of labeling an antibody molecule is disclosed. The method includes contacting an antibody molecule, with a chelating agent, to thereby produce a conjugated antibody.

As is discussed above, the antibody molecule can be conjugated to a therapeutic agent. Therapeutically active radioisotopes have already been mentioned. Examples of other therapeutic agents include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, maytansinoids, *e.g.*, maytansinol (*see* U.S. Pat. No. 5,208,020), CC-1065 (*see* U.S. Pat. Nos. 5,475,092, 5,585,499, 5,846, 545) and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (*e.g.*, methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (*e.g.*, mechlorethamine, thioepa chlorambucil, CC-1065, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (*e.g.*, daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (*e.g.*, dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (*e.g.*, vincristine, vinblastine, taxol and maytansinoids).

In one aspect, the invention features a method of providing a target binding molecule that specifically binds to a target disclosed herein, *e.g.*, PD-1 receptor. For example, the target binding molecule is an antibody molecule. The method includes: providing a target protein that comprises at least a portion of non-human protein, the portion being homologous to (at least 70, 75, 80, 85, 87, 90, 92, 94, 95, 96, 97, 98% identical to) a corresponding portion of a human target protein, but differing by at least one amino acid (*e.g.*, at least one, two, three, four, five, six,

seven, eight, or nine amino acids); obtaining an antibody molecule that specifically binds to the antigen; and evaluating efficacy of the binding agent in modulating activity of the target protein. The method can further include administering the binding agent (*e.g.*, antibody molecule) or a derivative (*e.g.*, a humanized antibody molecule) to a human subject.

5

Multispecific Antibody Molecules

In certain embodiments, the antibody molecule is a multi-specific (*e.g.*, a bispecific or a trispecific) antibody molecule. Protocols for generating bispecific or heterodimeric antibody molecules are known in the art; including but not limited to, for example, the “knob in a hole”
10 approach described in, *e.g.*, US 5731168; the electrostatic steering Fc pairing as described in, *e.g.*, WO 09/089004, WO 06/106905 and WO 2010/129304; Strand Exchange Engineered Domains (SEED) heterodimer formation as described in, *e.g.*, WO 07/110205; Fab arm exchange as described in, *e.g.*, WO 08/119353, WO 2011/131746, and WO 2013/060867; double antibody conjugate, *e.g.*, by antibody cross-linking to generate a bi-specific structure using a
15 heterobifunctional reagent having an amine-reactive group and a sulfhydryl reactive group as described in, *e.g.*, US 4433059; bispecific antibody determinants generated by recombining half antibodies (heavy-light chain pairs or Fabs) from different antibodies through cycle of reduction and oxidation of disulfide bonds between the two heavy chains, as described in, *e.g.*, US 4444878; trifunctional antibodies, *e.g.*, three Fab' fragments cross-linked through sulfhydryl
20 reactive groups, as described in, *e.g.*, US5273743; biosynthetic binding proteins, *e.g.*, pair of scFvs cross-linked through C-terminal tails preferably through disulfide or amine-reactive chemical cross-linking, as described in, *e.g.*, US5534254; bifunctional antibodies, *e.g.*, Fab fragments with different binding specificities dimerized through leucine zippers (*e.g.*, c-fos and c-jun) that have replaced the constant domain, as described in, *e.g.*, US5582996; bispecific and
25 oligospecific mono- and oligovalent receptors, *e.g.*, VH-CH1 regions of two antibodies (two Fab fragments) linked through a polypeptide spacer between the CH1 region of one antibody and the VH region of the other antibody typically with associated light chains, as described in, *e.g.*, US5591828; bispecific DNA-antibody conjugates, *e.g.*, crosslinking of antibodies or Fab fragments through a double stranded piece of DNA, as described in, *e.g.*, US5635602; bispecific
30 fusion proteins, *e.g.*, an expression construct containing two scFvs with a hydrophilic helical peptide linker between them and a full constant region, as described in, *e.g.*, US5637481;

multivalent and multispecific binding proteins, *e.g.*, dimer of polypeptides having first domain with binding region of Ig heavy chain variable region, and second domain with binding region of Ig light chain variable region, generally termed diabodies (higher order structures are also disclosed creating bispecific, trispecific, or tetraspecific molecules, as described in, *e.g.*,
5 US5837242; minibody constructs with linked VL and VH chains further connected with peptide spacers to an antibody hinge region and CH3 region, which can be dimerized to form bispecific/multivalent molecules, as described in, *e.g.*, US5837821; VH and VL domains linked with a short peptide linker (*e.g.*, 5 or 10 amino acids) or no linker at all in either orientation, which can form dimers to form bispecific diabodies; trimers and tetramers, as described in, *e.g.*,
10 US5844094; String of VH domains (or VL domains in family members) connected by peptide linkages with crosslinkable groups at the C-terminus further associated with VL domains to form a series of FVs (or scFvs), as described in, *e.g.*, US5864019; and single chain binding polypeptides with both a VH and a VL domain linked through a peptide linker are combined into multivalent structures through non-covalent or chemical crosslinking to form, *e.g.*,
15 homobivalent, heterobivalent, trivalent, and tetravalent structures using both scFV or diabody type format, as described in, *e.g.*, US5869620. Additional exemplary multispecific and bispecific molecules and methods of making the same are found, for example, in US5910573, US5932448, US5959083, US5989830, US6005079, US6239259, US6294353, US6333396, US6476198, US6511663, US6670453, US6743896, US6809185, US6833441, US7129330,
20 US7183076, US7521056, US7527787, US7534866, US7612181, US2002004587A1, US2002076406A1, US2002103345A1, US2003207346A1, US2003211078A1, US2004219643A1, US2004220388A1, US2004242847A1, US2005003403A1, US2005004352A1, US2005069552A1, US2005079170A1, US2005100543A1, US2005136049A1, US2005136051A1, US2005163782A1, US2005266425A1,
25 US2006083747A1, US2006120960A1, US2006204493A1, US2006263367A1, US2007004909A1, US2007087381A1, US2007128150A1, US2007141049A1, US2007154901A1, US2007274985A1, US2008050370A1, US2008069820A1, US2008152645A1, US2008171855A1, US2008241884A1, US2008254512A1, US2008260738A1, US2009130106A1, US2009148905A1, US2009155275A1,
30 US2009162359A1, US2009162360A1, US2009175851A1, US2009175867A1, US2009232811A1, US2009234105A1, US2009263392A1, US2009274649A1, EP346087A2,

WO0006605A2, WO02072635A2, WO04081051A1, WO06020258A2, WO2007044887A2, WO2007095338A2, WO2007137760A2, WO2008119353A1, WO2009021754A2, WO2009068630A1, WO9103493A1, WO9323537A1, WO9409131A1, WO9412625A2, WO9509917A1, WO9637621A2, WO9964460A1. The contents of the above-referenced applications are incorporated herein by reference in their entireties.

In other embodiments, the anti-PD-1 antibody molecule (*e.g.*, a monospecific, bispecific, or multispecific antibody molecule) is covalently linked, *e.g.*, fused, to another partner *e.g.*, a protein *e.g.*, one, two or more cytokines, *e.g.*, as a fusion molecule for example a fusion protein. In other embodiments, the fusion molecule comprises one or more proteins, *e.g.*, one, two or more cytokines. In one embodiment, the cytokine is an interleukin (IL) chosen from one, two, three or more of IL-1, IL-2, IL-12, IL-15 or IL-21. In one embodiment, a bispecific antibody molecule has a first binding specificity to a first target (*e.g.*, to PD-1), a second binding specificity to a second target (*e.g.*, LAG-3 or TIM-3), and is optionally linked to an interleukin (*e.g.*, IL-12) domain *e.g.*, full length IL-12 or a portion thereof.

A “fusion protein” and a “fusion polypeptide” refer to a polypeptide having at least two portions covalently linked together, where each of the portions is a polypeptide having a different property. The property may be a biological property, such as activity *in vitro* or *in vivo*. The property can also be simple chemical or physical property, such as binding to a target molecule, catalysis of a reaction, etc. The two portions can be linked directly by a single peptide bond or through a peptide linker, but are in reading frame with each other.

This invention provides an isolated nucleic acid molecule encoding the above antibody molecule, vectors and host cells thereof. The nucleic acid molecule includes but is not limited to RNA, genomic DNA and cDNA.

Exemplary PD-1 Inhibitors

PD-1 is a CD28/CTLA-4 family member expressed, *e.g.*, on activated CD4⁺ and CD8⁺ T cells, T_{regs}, and B cells. It negatively regulates effector T cell signaling and function. PD-1 is induced on tumor-infiltrating T cells, and can result in functional exhaustion or dysfunction (Keir *et al.* (2008) *Annu. Rev. Immunol.* 26:677-704; Pardoll *et al.* (2012) *Nat Rev Cancer* 12(4):252-64). PD-1 delivers a coinhibitory signal upon binding to either of its two ligands, Programmed Death-Ligand 1 (PD-L1) or Programmed Death-Ligand 2 (PD-L2). PD-L1 is expressed on a

number of cell types, including T cells, natural killer (NK) cells, macrophages, dendritic cells (DCs), B cells, epithelial cells, vascular endothelial cells, as well as many types of tumors. High expression of PD-L1 on murine and human tumors has been linked to poor clinical outcomes in a variety of cancers (Keir *et al.* (2008) *Annu. Rev. Immunol.* 26:677-704; Pardoll *et al.* (2012) *Nat Rev Cancer* 12(4):252-64). PD-L2 is expressed on dendritic cells, macrophages, and some tumors. Blockade of the PD-1 pathway has been pre-clinically and clinically validated for cancer immunotherapy. Both preclinical and clinical studies have demonstrated that anti-PD-1 blockade can restore activity of effector T cells and results in robust anti-tumor response. For example, blockade of PD-1 pathway can restore exhausted/dysfunctional effector T cell function (e.g., proliferation, IFN- γ secretion, or cytolytic function) and/or inhibit T_{reg} cell function (Keir *et al.* (2008) *Annu. Rev. Immunol.* 26:677-704; Pardoll *et al.* (2012) *Nat Rev Cancer* 12(4):252-64). Blockade of the PD-1 pathway can be effected with an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide of PD-1, PD-L1 and/or PD-L2.

As used herein, the term “Programmed Death 1” or “PD-1” include isoforms, mammalian, e.g., human PD-1, species homologs of human PD-1, and analogs comprising at least one common epitope with PD-1. The amino acid sequence of PD-1, e.g., human PD-1, is known in the art, e.g., Shinohara T *et al.* (1994) *Genomics* 23(3):704-6; Finger LR, *et al.* *Gene* (1997) 197(1-2):177-87.

The anti-PD-1 antibody molecules described herein can be used alone or in combination with one or more additional agents described herein in accordance with a method described herein. In certain embodiments, the combinations described herein include a PD-1 inhibitor, e.g., an anti-PD-1 antibody molecule (e.g., humanized antibody molecules) as described herein.

In some embodiments, the anti-PD-1 antibody molecule (e.g., an isolated or recombinant antibody molecule) has one or more of the following properties:

(i) binds to PD-1, e.g., human PD-1, with high affinity, e.g., with an affinity constant of at least about 10^7 M^{-1} , typically about 10^8 M^{-1} , and more typically, about 10^9 M^{-1} to 10^{10} M^{-1} or stronger;

(ii) does not substantially bind to CD28, CTLA-4, ICOS or BTLA;

(iii) inhibits or reduces binding of PD-1 to a PD-1 ligand, e.g., PD-L1 or PD-L2, or both;

(iv) binds specifically to an epitope on PD-1, *e.g.*, the same or similar epitope as the epitope recognized by murine monoclonal antibody BAP049 or a chimeric antibody BAP049, *e.g.*, BAP049-chi or BAP049-chi-Y;

(v) shows the same or similar binding affinity or specificity, or both, as any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E;

(vi) shows the same or similar binding affinity or specificity, or both, as an antibody molecule (*e.g.*, an heavy chain variable region and light chain variable region) described in Table 1;

(vii) shows the same or similar binding affinity or specificity, or both, as an antibody molecule (*e.g.*, an heavy chain variable region and light chain variable region) having an amino acid sequence shown in Table 1;

(viii) shows the same or similar binding affinity or specificity, or both, as an antibody molecule (*e.g.*, an heavy chain variable region and light chain variable region) encoded by the nucleotide sequence shown in Table 1;

(ix) inhibits, *e.g.*, competitively inhibits, the binding of a second antibody molecule to PD-1, wherein the second antibody molecule is an antibody molecule described herein, *e.g.*, an antibody molecule chosen from, *e.g.*, any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E;

(x) binds the same or an overlapping epitope with a second antibody molecule to PD-1, wherein the second antibody molecule is an antibody molecule described herein, *e.g.*, an antibody molecule chosen from, *e.g.*, any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13,

BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E;

(xi) competes for binding, and/or binds the same epitope, with a second antibody molecule to PD-1, wherein the second antibody molecule is an antibody molecule described herein, *e.g.*, an antibody molecule chosen from, *e.g.*, any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E;

(xii) has one or more biological properties of an antibody molecule described herein, *e.g.*, an antibody molecule chosen from, *e.g.*, any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E;

(xiii) has one or more pharmacokinetic properties of an antibody molecule described herein, *e.g.*, an antibody molecule chosen from, *e.g.*, any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E;

(xiv) inhibits one or more activities of PD-1, *e.g.*, results in one or more of: an increase in tumor infiltrating lymphocytes, an increase in T-cell receptor mediated proliferation, or a decrease in immune evasion by cancerous cells;

(xv) binds human PD-1 and is cross-reactive with cynomolgus PD-1;

(xvi) binds to one or more residues within the C strand, CC' loop, C' strand, or FG loop of PD-1, or a combination two, three or all of the C strand, CC' loop, C' strand or FG loop of PD-1, *e.g.*, wherein the binding is assayed using ELISA or Biacore; or

(xvii) has a VL region that contributes more to binding to PD-1 than a VH region.

In some embodiments, the antibody molecule binds to PD-1 with high affinity, *e.g.*, with a K_D that is about the same, or at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or

90% higher or lower than the K_D of a murine or chimeric anti-PD-1 antibody molecule, *e.g.*, a murine or chimeric anti-PD-1 antibody molecule described herein. In some embodiments, the K_D of the murine or chimeric anti-PD-1 antibody molecule is less than about 0.4, 0.3, 0.2, 0.1, or 0.05 nM, *e.g.*, measured by a Biacore method. In some embodiments, the K_D of the murine or chimeric anti-PD-1 antibody molecule is less than about 0.2 nM, *e.g.*, about 0.135 nM. In other embodiments, the K_D of the murine or chimeric anti PD-1 antibody molecule is less than about 10, 5, 3, 2, or 1 nM, *e.g.*, measured by binding on cells expressing PD-1 (*e.g.*, 300.19 cells). In some embodiments, the K_D of the murine or chimeric anti PD-1 antibody molecule is less than about 5 nM, *e.g.*, about 4.60 nM (or about 0.69 $\mu\text{g/mL}$).

In some embodiments, the anti-PD-1 antibody molecule binds to PD-1 with a K_{off} slower than 1×10^{-4} , 5×10^{-5} , or $1 \times 10^{-5} \text{ s}^{-1}$, *e.g.*, about $1.65 \times 10^{-5} \text{ s}^{-1}$. In some embodiments, the anti-PD-1 antibody molecule binds to PD-1 with a K_{on} faster than 1×10^4 , 5×10^4 , 1×10^5 , or $5 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, *e.g.*, about $1.23 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$.

In some embodiments, the expression level of the antibody molecule is higher, *e.g.*, at least about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10-fold higher, than the expression level of a murine or chimeric antibody molecule, *e.g.*, a murine or chimeric anti-PD-1 antibody molecule described herein. In some embodiments, the antibody molecule is expressed in CHO cells.

In some embodiments, the anti-PD-1 antibody molecule reduces one or more PD-1-associated activities with an IC_{50} (concentration at 50% inhibition) that is about the same or lower, *e.g.*, at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% lower, than the IC_{50} of a murine or chimeric anti-PD-1 antibody molecule, *e.g.*, a murine or chimeric anti-PD-1 antibody molecule described herein. In some embodiments, the IC_{50} of the murine or chimeric anti-PD-1 antibody molecule is less than about 6, 5, 4, 3, 2, or 1 nM, *e.g.*, measured by binding on cells expressing PD-1 (*e.g.*, 300.19 cells). In some embodiments, the IC_{50} of the murine or chimeric anti-PD-1 antibody molecule is less than about 4 nM, *e.g.*, about 3.40 nM (or about 0.51 $\mu\text{g/mL}$). In some embodiments, the PD-1-associated activity reduced is the binding of PD-L1 and/or PD-L2 to PD-1. In some embodiments, the anti-PD-1 antibody molecule binds to peripheral blood mononucleated cells (PBMCs) activated by Staphylococcal enterotoxin B (SEB). In other embodiments, the anti-PD-1 antibody molecule increases the expression of IL-2 on whole blood activated by SEB. For example, the anti-PD-1 antibody increases the expression

of IL-2 by at least about 2, 3, 4, or 5-fold, compared to the expression of IL-2 when an isotype control (*e.g.*, IgG4) is used.

In some embodiments, the anti-PD-1 antibody molecule has improved stability, *e.g.*, at least about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10-fold more stable *in vivo* or *in vitro*, than a murine or chimeric anti-PD-1 antibody molecule, *e.g.*, a murine or chimeric anti-PD-1 antibody molecule described herein.

In one embodiment, the anti PD-1 antibody molecule is a humanized antibody molecule and has a risk score based on T cell epitope analysis of 300 to 700, 400 to 650, 450 to 600, or a risk score as described herein.

In another embodiment, the anti-PD-1 antibody molecule comprises at least one antigen-binding region, *e.g.*, a variable region or an antigen-binding fragment thereof, from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one, two, three or four variable regions from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one or two heavy chain variable regions from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-

hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one or two light chain variable regions from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule includes a heavy chain constant region for an IgG4, *e.g.*, a human IgG4. In one embodiment, the human IgG4 includes a substitution at position 228 according to EU numbering (*e.g.*, a Ser to Pro substitution). In still another embodiment, the anti-PD-1 antibody molecule includes a heavy chain constant region for an IgG1, *e.g.*, a human IgG1. In one embodiment, the human IgG1 includes a substitution at position 297 according to EU numbering (*e.g.*, an Asn to Ala substitution). In one embodiment, the human IgG1 includes a substitution at position 265 according to EU numbering, a substitution at position 329 according to EU numbering, or both (*e.g.*, an Asp to Ala substitution at position 265 and/or a Pro to Ala substitution at position 329). In one embodiment, the human IgG1 includes a substitution at position 234 according to EU numbering, a substitution at position 235 according to EU numbering, or both (*e.g.*, a Leu to Ala substitution at position 234 and/or a Leu to Ala substitution at position 235). In one embodiment, the heavy chain constant region comprises an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto.

In yet another embodiment, the anti-PD-1 antibody molecule includes a kappa light chain constant region, *e.g.*, a human kappa light chain constant region. In one embodiment, the light

chain constant region comprises an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto.

In another embodiment, the anti-PD-1 antibody molecule includes a heavy chain constant
5 region for an IgG4, *e.g.*, a human IgG4, and a kappa light chain constant region, *e.g.*, a human kappa light chain constant region, *e.g.*, a heavy and light chain constant region comprising an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto. In one embodiment, the human IgG4 includes a substitution at position 228 according to EU numbering (*e.g.*, a Ser to
10 Pro substitution). In yet another embodiment, the anti-PD-1 antibody molecule includes a heavy chain constant region for an IgG1, *e.g.*, a human IgG1, and a kappa light chain constant region, *e.g.*, a human kappa light chain constant region, *e.g.*, a heavy and light chain constant region comprising an amino sequence set forth in Table 3, or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) thereto. In one
15 embodiment, the human IgG1 includes a substitution at position 297 according to EU numbering (*e.g.*, an Asn to Ala substitution). In one embodiment, the human IgG1 includes a substitution at position 265 according to EU numbering, a substitution at position 329 according to EU numbering, or both (*e.g.*, an Asp to Ala substitution at position 265 and/or a Pro to Ala substitution at position 329). In one embodiment, the human IgG1 includes a substitution at
20 position 234 according to EU numbering, a substitution at position 235 according to EU numbering, or both (*e.g.*, a Leu to Ala substitution at position 234 and/or a Leu to Ala substitution at position 235).

In another embodiment, the anti-PD-1 antibody molecule includes a heavy chain variable domain and a constant region, a light chain variable domain and a constant region, or both,
25 comprising the amino acid sequence of BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences. The anti-PD-1 antibody molecule, optionally, comprises a leader sequence from a heavy chain, a light chain, or
30 both, as shown in Table 4; or a sequence substantially identical thereto.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three complementarity determining regions (CDRs) from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs (or collectively all of the CDRs) from a heavy chain variable region comprising an amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1. In one embodiment, one or more of the CDRs (or collectively all of the CDRs) have one, two, three, four, five, six or more changes, *e.g.*, amino acid substitutions or deletions, relative to the amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs from a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequence.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs (or collectively all of the CDRs) from a light chain variable region comprising an amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1. In one embodiment, one or more of the CDRs (or collectively all of the CDRs) have one, two, three, four, five, six or more changes, *e.g.*, amino acid substitutions or deletions, relative to the amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1.

In certain embodiments, the anti-PD-1 antibody molecule includes a substitution in a light chain CDR, *e.g.*, one or more substitutions in a CDR1, CDR2 and/or CDR3 of the light chain. In one embodiment, the anti-PD-1 antibody molecule includes a substitution in the light chain CDR3 at position 102 of the light variable region, *e.g.*, a substitution of a cysteine to tyrosine, or a cysteine to serine residue, at position 102 of the light variable region according to Table 1 (*e.g.*, SEQ ID NO: 16 or 24 for murine or chimeric, unmodified; or any of SEQ ID NOs: 34, 42, 46, 54, 58, 62, 66, 70, 74, or 78 for a modified sequence).

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, three, four, five or six CDRs (or collectively all of the CDRs) from a heavy and light chain variable region comprising an amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1. In one embodiment, one or more of the CDRs (or collectively all of the CDRs) have one, two, three, four, five, six or more changes, *e.g.*, amino acid substitutions or deletions, relative to the amino acid sequence shown in Table 1, or encoded by a nucleotide sequence shown in Table 1.

In one embodiment, the anti-PD-1 antibody molecule includes all six CDRs from an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1, or closely related CDRs, *e.g.*, CDRs which are identical or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions). In one embodiment, the anti-PD-1 antibody molecule may include any CDR described herein. In certain embodiments, the anti-PD-1 antibody molecule includes a substitution in a light chain CDR, *e.g.*, one or more substitutions in a CDR1, CDR2 and/or CDR3 of the light chain. In one embodiment, the anti-PD-1 antibody molecule includes a substitution in the light chain CDR3 at position 102 of the light variable region, *e.g.*, a substitution of a cysteine to tyrosine, or a cysteine to serine residue, at position 102 of the light variable region according to Table 1 (*e.g.*, SEQ ID NO: 16 or 24 for murine or chimeric, unmodified; or any of SEQ ID NOs: 34, 42, 46, 54, 58, 62, 66, 70, 74, or 78 for a modified sequence).

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs according to Kabat *et al.* (*e.g.*, at least one, two, or three CDRs according to the Kabat definition as set out in Table 1) from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three CDRs according to Kabat *et al.* shown in Table 1.

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three CDRs according to Kabat *et al.* (*e.g.*, at least one, two, or three CDRs according to the Kabat definition as set out in Table 1) from a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three CDRs according to Kabat *et al.* shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, three, four, five, or six CDRs according to Kabat *et al.* (*e.g.*, at least one, two, three, four, five, or six CDRs according to the Kabat definition as set out in Table 1) from the heavy and light chain variable regions of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06,

BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence
 5 substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, three, four, five, or six CDRs according to Kabat *et al.* shown in Table 1.

10 In yet another embodiment, the anti-PD-1 antibody molecule includes all six CDRs according to Kabat *et al.* (*e.g.*, all six CDRs according to the Kabat definition as set out in Table 1) from the heavy and light chain variable regions of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09,
 15 BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at
 20 least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to all six CDRs according to Kabat *et al.* shown in Table 1. In one embodiment, the anti-PD-1 antibody molecule may include any CDR described herein.

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or
 25 three Chothia hypervariable loops (*e.g.*, at least one, two, or three hypervariable loops according to the Chothia definition as set out in Table 1) from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12,
 30 BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in

Table 1, or encoded by the nucleotide sequence in Table 1; or at least the amino acids from those hypervariable loops that contact PD-1; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three hypervariable loops according to Chothia *et al.* shown in Table 1.

In another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three Chothia hypervariable loops (*e.g.*, at least one, two, or three hypervariable loops according to the Chothia definition as set out in Table 1) of a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or at least the amino acids from those hypervariable loops that contact PD-1; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three hypervariable loops according to Chothia *et al.* shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule includes at least one, two, three, four, five, or six hypervariable loops (*e.g.*, at least one, two, three, four, five, or six hypervariable loops according to the Chothia definition as set out in Table 1) from the heavy and light chain variable regions of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or at least the amino acids from those hypervariable loops that contact PD-1; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, three, four, five or six hypervariable loops according to Chothia *et al.* shown in Table 1.

In one embodiment, the anti-PD-1 antibody molecule includes all six hypervariable loops (*e.g.*, all six hypervariable loops according to the Chothia definition as set out in Table 1) of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, or closely related hypervariable loops, *e.g.*, hypervariable loops which are identical or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions); or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to all six hypervariable loops according to Chothia *et al.* shown in Table 1. In one embodiment, the anti-PD-1 antibody molecule may include any hypervariable loop described herein.

In still another embodiment, the anti-PD-1 antibody molecule includes at least one, two, or three hypervariable loops that have the same canonical structures as the corresponding hypervariable loop of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, *e.g.*, the same canonical structures as at least loop 1 and/or loop 2 of the heavy and/or light chain variable domains of an antibody described herein. *See, e.g.*, Chothia *et al.*, (1992) *J. Mol. Biol.* 227:799-817; Tomlinson *et al.*, (1992) *J. Mol. Biol.* 227:776-798 for descriptions of hypervariable loop canonical structures. These structures can be determined by inspection of the tables described in these references.

In certain embodiments, the anti-PD-1 antibody molecule includes a combination of CDRs or hypervariable loops defined according to the Kabat *et al.* and Chothia *et al.*

In one embodiment, the anti-PD-1 antibody molecule includes at least one, two or three CDRs or hypervariable loops from a heavy chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03,

BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, according to the Kabat and Chothia definition (*e.g.*, at least one, two, or three CDRs or hypervariable loops according to the Kabat and Chothia definition as set out in Table 1); or encoded by the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical) to any of the aforesaid sequences; or which have at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions) relative to one, two, or three CDRs or hypervariable loops according to Kabat and/or Chothia shown in Table 1.

For example, the anti-PD-1 antibody molecule can include VH CDR1 according to Kabat *et al.* or VH hypervariable loop 1 according to Chothia *et al.*, or a combination thereof, *e.g.*, as shown in Table 1. In one embodiment, the combination of Kabat and Chothia CDR of VH CDR1 comprises the amino acid sequence GYTFTTYWMH (SEQ ID NO: 224), or an amino acid sequence substantially identical thereto (*e.g.*, having at least one amino acid alteration, but not more than two, three or four alterations (*e.g.*, substitutions, deletions, or insertions, *e.g.*, conservative substitutions)). The anti-PD-1 antibody molecule can further include, *e.g.*, VH CDRs 2-3 according to Kabat *et al.* and VL CDRs 1-3 according to Kabat *et al.*, *e.g.*, as shown in Table 1. Accordingly, in some embodiments, framework regions are defined based on a combination of CDRs defined according to Kabat *et al.* and hypervariable loops defined according to Chothia *et al.* For example, the anti-PD-1 antibody molecule can include VH FR1 defined based on VH hypervariable loop 1 according to Chothia *et al.* and VH FR2 defined based on VH CDRs 1-2 according to Kabat *et al.*, *e.g.*, as shown in Table 1. The anti-PD-1 antibody molecule can further include, *e.g.*, VH FRs 3-4 defined based on VH CDRs 2-3 according to Kabat *et al.* and VL FRs 1-4 defined based on VL CDRs 1-3 according to Kabat *et al.*

The anti-PD-1 antibody molecule can contain any combination of CDRs or hypervariable loops according to the Kabat and Chothia definitions. In one embodiment, the anti-PD-1 antibody molecule includes at least one, two or three CDRs from a light chain variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-

hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, according to the
 5 Kabat and Chothia definition (*e.g.*, at least one, two, or three CDRs according to the Kabat and Chothia definition as set out in Table 1).

In an embodiment, *e.g.*, an embodiment comprising a variable region, a CDR (*e.g.*, Chothia CDR or Kabat CDR), or other sequence referred to herein, *e.g.*, in Table 1, the antibody molecule is a monospecific antibody molecule, a bispecific antibody molecule, or is an antibody
 10 molecule that comprises an antigen binding fragment of an antibody, *e.g.*, a half antibody or antigen binding fragment of a half antibody. In certain embodiments the antibody molecule is a bispecific antibody molecule having a first binding specificity for PD-1 and a second binding specificity for TIM-3, LAG-3, CEACAM (*e.g.*, CEACAM-1 and/or CEACAM-5), PD-L1 or PD-L2.

15 In one embodiment, the anti-PD-1 antibody molecule includes:

(a) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a
 20 VLCDR3 amino acid sequence of SEQ ID NO: 33;

(b) a VH comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ
 25 ID NO: 32;

(c) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33; or

30 (d) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3;

and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

In one embodiment, the anti-PD-1 antibody molecule comprises a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33.

In one embodiment, the anti-PD-1 antibody molecule comprises a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

In one embodiment, the anti-PD-1 antibody molecule comprises a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33.

In one embodiment, the anti-PD-1 antibody molecule comprises a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

In one embodiment, the antibody molecule is a humanized antibody molecule. In another embodiment, the antibody molecule is a monospecific antibody molecule. In yet another embodiment, the antibody molecule is a bispecific antibody molecule.

In one embodiment, the anti-PD-1 antibody molecule includes:

(i) a heavy chain variable region (VH) including a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1, SEQ ID NO: 4 or SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and

(ii) a light chain variable region (VL) including a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

In another embodiment, the anti-PD-1 antibody molecule includes:

5 (i) a heavy chain variable region (VH) including a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1, SEQ ID NO: 4 or SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and

(ii) a light chain variable region (VL) including a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid
10 sequence of SEQ ID NO: 33.

In one embodiment, the anti-PD-1 antibody molecule comprises the VHCDR1 amino acid sequence of SEQ ID NO: 1. In another embodiment, the anti-PD-1 antibody molecule comprises the VHCDR1 amino acid sequence of SEQ ID NO: 4. In yet another embodiment, the anti-PD-1 antibody molecule comprises the VHCDR1 amino acid sequence of SEQ ID NO: 224.

15 In one embodiment, the light or the heavy chain variable framework (*e.g.*, the region encompassing at least FR1, FR2, FR3, and optionally FR4) of the anti-PD-1 antibody molecule can be chosen from: (a) a light or heavy chain variable framework including at least 80%, 85%, 87%, 90%, 92%, 93%, 95%, 97%, 98%, or preferably 100% of the amino acid residues from a human light or heavy chain variable framework, *e.g.*, a light or heavy chain variable framework
20 residue from a human mature antibody, a human germline sequence, or a human consensus sequence; (b) a light or heavy chain variable framework including from 20% to 80%, 40% to 60%, 60% to 90%, or 70% to 95% of the amino acid residues from a human light or heavy chain variable framework, *e.g.*, a light or heavy chain variable framework residue from a human mature antibody, a human germline sequence, or a human consensus sequence; (c) a non-human
25 framework (*e.g.*, a rodent framework); or (d) a non-human framework that has been modified, *e.g.*, to remove antigenic or cytotoxic determinants, *e.g.*, deimmunized, or partially humanized. In one embodiment, the light or heavy chain variable framework region (particularly FR1, FR2 and/or FR3) includes a light or heavy chain variable framework sequence at least 70, 75, 80, 85, 87, 88, 90, 92, 94, 95, 96, 97, 98, 99% identical or identical to the frameworks of a VL or VH
30 segment of a human germline gene.

In certain embodiments, the anti-PD-1 antibody molecule comprises a heavy chain variable domain having at least one, two, three, four, five, six, seven, ten, fifteen, twenty or more changes, *e.g.*, amino acid substitutions or deletions, from an amino acid sequence of BAP049-chi-HC, *e.g.*, the amino acid sequence of the FR region in the entire variable region, *e.g.*, shown in FIGs. 9A-9B, or SEQ ID NO: 18, 20, 22 or 30. In one embodiment, the anti-PD-1 antibody molecule comprises a heavy chain variable domain having one or more of: E at position 1, V at position 5, A at position 9, V at position 11, K at position 12, K at position 13, E at position 16, L at position 18, R at position 19, I or V at position 20, G at position 24, I at position 37, A or S at position 40, T at position 41, S at position 42, R at position 43, M or L at position 48, V or F at position 68, T at position 69, I at position 70, S at position 71, A or R at position 72, K or N at position 74, T or K at position 76, S or N at position 77, L at position 79, L at position 81, E or Q at position 82, M at position 83, S or N at position 84, R at position 87, A at position 88, or T at position 91 of amino acid sequence of BAP049-chi-HC, *e.g.*, the amino acid sequence of the FR in the entire variable region, *e.g.*, shown in FIGs. 9A-9B, or SEQ ID NO: 18, 20, 22 or 30.

Alternatively, or in combination with the heavy chain substitutions of BAP049-chi-HC described herein, the anti-PD-1 antibody molecule comprises a light chain variable domain having at least one, two, three, four, five, six, seven, ten, fifteen, twenty or more amino acid changes, *e.g.*, amino acid substitutions or deletions, from an amino acid sequence of BAP049-chi-LC, *e.g.*, the amino acid sequence shown in FIGs. 10A-10B, or SEQ ID NO: 24 or 26. In one embodiment, the anti-PD-1 antibody molecule comprises a heavy chain variable domain having one or more of: E at position 1, V at position 2, Q at position 3, L at position 4, T at position 7, D or L or A at position 9, F or T at position 10, Q at position 11, S or P at position 12, L or A at position 13, S at position 14, P or L or V at position 15, K at position 16, Q or D at position 17, R at position 18, A at position 19, S at position 20, I or L at position 21, T at position 22, L at position 43, K at position 48, A or S at position 49, R or Q at position 51, Y at position 55, I at position 64, S or P at position 66, S at position 69, Y at position 73, G at position 74, E at position 76, F at position 79, N at position 82, N at position 83, L or I at position 84, E at position 85, S or P at position 86, D at position 87, A or F or I at position 89, T or Y at position 91, F at position 93, or Y at position 102 of the amino acid sequence of BAP049-chi-LC, *e.g.*, the amino acid sequence shown in FIGs. 10A-10B, or SEQ ID NO: 24 or 26.

In other embodiments, the anti-PD-1 antibody molecule includes one, two, three, or four heavy chain framework regions (*e.g.*, a VHFW amino acid sequence shown in Table 2, or encoded by the nucleotide sequence shown in Table 2), or a sequence substantially identical thereto.

5 In yet other embodiments, the anti-PD-1 antibody molecule includes one, two, three, or four light chain framework regions (*e.g.*, a VLFW amino acid sequence shown in Table 2, or encoded by the nucleotide sequence shown in Table 2), or a sequence substantially identical thereto.

10 In other embodiments, the anti-PD-1 antibody molecule includes one, two, three, or four heavy chain framework regions (*e.g.*, a VHFW amino acid sequence shown in Table 2, or encoded by the nucleotide sequence shown in Table 2), or a sequence substantially identical thereto; and one, two, three, or four light chain framework regions (*e.g.*, a VLFW amino acid sequence shown in Table 2, or encoded by the nucleotide sequence shown in Table 2), or a sequence substantially identical thereto.

15 In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework region 1 (VHFW1) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or
20 BAP049-Clone-E (*e.g.*, SEQ ID NO: 147). In some embodiments, the antibody molecule comprises the heavy chain framework region 1 (VHFW1) of BAP049-hum14 or BAP049-hum15 (*e.g.*, SEQ ID NO: 151).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework region 2 (VHFW2) of BAP049-hum01, BAP049-hum02, BAP049-hum05, BAP049-
25 hum06, BAP049-hum07, BAP049-hum09, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 153). In some embodiments, the antibody molecule comprises the heavy chain framework region 2 (VHFW2) of BAP049-hum03, BAP049-hum04, BAP049-hum08, BAP049-hum10, BAP049-hum14, BAP049-hum15, or BAP049-Clone-D (*e.g.*, SEQ ID NO: 157). In some
30 embodiments, the antibody molecule comprises the heavy chain framework region 2 (VHFW2) of BAP049-hum16 (*e.g.*, SEQ ID NO: 160).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework region 3 (VHFW3) of BAP049-hum01, BAP049-hum02, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum09, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 162). In some embodiments, the antibody molecule comprises the heavy chain framework region 3 (VHFW3) of BAP049-hum03, BAP049-hum04, BAP049-hum08, BAP049-hum10, BAP049-hum14, BAP049-hum15, BAP049-hum16, or BAP049-Clone-D (*e.g.*, SEQ ID NO: 166).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework region 4 (VHFW4) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 169).

In some embodiments, the anti-PD-1 antibody molecule comprises the light chain framework region 1 (VLFW1) of BAP049-hum08, BAP049-hum09, BAP049-hum15, BAP049-hum16, or BAP049-Clone-C (*e.g.*, SEQ ID NO: 174). In some embodiments, the antibody molecule comprises the light chain framework region 1 (VLFW1) of BAP049-hum01, BAP049-hum04, BAP049-hum05, BAP049-hum07, BAP049-hum10, BAP049-hum11, BAP049-hum14, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 177). In some embodiments, the antibody molecule comprises the light chain framework region 1 (VLFW1) of BAP049-hum06 (*e.g.*, SEQ ID NO: 181). In some embodiments, the antibody molecule comprises the light chain framework region 1 (VLFW1) of BAP049-hum13 (*e.g.*, SEQ ID NO: 183). In some embodiments, the antibody molecule comprises the light chain framework region 1 (VLFW1) of BAP049-hum02, BAP049-hum03, or BAP049-hum12 (*e.g.*, SEQ ID NO: 185).

In some embodiments, the anti-PD-1 antibody molecule comprises the light chain framework region 2 (VLFW2) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum06, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 187). In some embodiments, the antibody molecule

comprises the light chain framework region 2 (VLFW2) of BAP049-hum04, BAP049-hum05, BAP049-hum07, BAP049-hum13, or BAP049-Clone-C (*e.g.*, SEQ ID NO: 191). In some embodiments, the antibody molecule comprises the light chain framework region 2 (VLFW2) of BAP049-hum12 (*e.g.*, SEQ ID NO: 194).

5 In some embodiments, the anti-PD-1 antibody molecule comprises the light chain framework region 3 (VLFW3) of BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 196). In some embodiments, the antibody molecule comprises the light chain
10 framework region 3 (VLFW3) of BAP049-hum02 or BAP049-hum03 (*e.g.*, SEQ ID NO: 200). In some embodiments, the antibody molecule comprises the light chain framework region 3 (VLFW3) of BAP049-hum01 or BAP049-Clone-A (*e.g.*, SEQ ID NO: 202). In some embodiments, the antibody molecule comprises the light chain framework region 3 (VLFW3) of BAP049-hum04, BAP049-hum05, or BAP049-Clone-B (*e.g.*, SEQ ID NO: 205).

15 In some embodiments, the anti-PD-1 antibody molecule comprises the light chain framework region 4 (VLFW4) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C,
20 BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 208).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum01, BAP049-hum02, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum09, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 147
25 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)). In some embodiments, the antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum03, BAP049-hum04, BAP049-hum08, BAP049-hum10, or BAP049-Clone-D (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)). In some embodiments, the antibody molecule comprises the heavy chain framework regions 1-3 of
30 BAP049-hum14 or BAP049-hum15 (*e.g.*, SEQ ID NO: 151 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)). In some embodiments, the antibody molecule

comprises the heavy chain framework regions 1-3 of BAP049-hum16 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 160 (VHFW2), and SEQ ID NO: 166 (VHFW3)). In some embodiments, the antibody molecule further comprises the heavy chain framework region 4 (VHFW4) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 169).

In some embodiments, the anti-PD-1 antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum01 or BAP049-Clone-A (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 202 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum02 or BAP049-hum03 (*e.g.*, SEQ ID NO: 185 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 200 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum04, BAP049-hum05, or BAP049-Clone-B (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 191 (VLFW2), and SEQ ID NO: 205 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum06 (*e.g.*, SEQ ID NO: 181 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum07 (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 191 (VLFW2), and SEQ ID NO: 196 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum08, BAP049-hum09, BAP049-hum15, BAP049-hum16, or BAP049-Clone-C (*e.g.*, SEQ ID NO: 174 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum10, BAP049-hum11, BAP049-hum14, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum12 (*e.g.*, SEQ ID NO: 185 (VLFW1), SEQ ID NO: 194 (VLFW2), and SEQ ID NO: 196 (VLFW3)). In some embodiments, the antibody molecule comprises the light chain framework regions 1-3 of BAP049-hum13 (*e.g.*, SEQ ID NO: 183 (VLFW1), SEQ ID

NO: 191 (VLFW2), and SEQ ID NO: 196 (VLFW3)). In some embodiments, the antibody molecule further comprises the light chain framework region 4 (VLFW4) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 208).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum01 or BAP049-Clone-A (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum01 or BAP049-Clone-A (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 202 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum02 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum02 (*e.g.*, SEQ ID NO: 185 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 200 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum03 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum03 (*e.g.*, SEQ ID NO: 185 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 200 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum04 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum04 (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 191 (VLFW2), and SEQ ID NO: 205 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum05 or BAP049-Clone-B (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain

framework regions 1-3 of BAP049-hum05 or BAP049-Clone-B (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 191 (VLFW2), and SEQ ID NO: 205 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum06 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum06 (*e.g.*, SEQ ID NO: 181 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum07 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum07 (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 191 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum08 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum08 (*e.g.*, SEQ ID NO: 174 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum09 or BAP049-Clone-C (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum09 or BAP049-Clone-C (*e.g.*, SEQ ID NO: 174 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum10 or BAP049-Clone-D (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum10 or BAP049-Clone-D (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum11 or BAP049-Clone-E (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain

framework regions 1-3 of BAP049-hum11 or BAP049-Clone-E (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum12 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum12 (*e.g.*, SEQ ID NO: 185 (VLFW1), SEQ ID NO: 194 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum13 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 153 (VHFW2), and SEQ ID NO: 162 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum13 (*e.g.*, SEQ ID NO: 183 (VLFW1), SEQ ID NO: 191 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum14 (*e.g.*, SEQ ID NO: 151 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum14 (*e.g.*, SEQ ID NO: 177 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum15 (*e.g.*, SEQ ID NO: 151 (VHFW1), SEQ ID NO: 157 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum15 (*e.g.*, SEQ ID NO: 174 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule comprises the heavy chain framework regions 1-3 of BAP049-hum16 (*e.g.*, SEQ ID NO: 147 (VHFW1), SEQ ID NO: 160 (VHFW2), and SEQ ID NO: 166 (VHFW3)) and the light chain framework regions 1-3 of BAP049-hum16 (*e.g.*, SEQ ID NO: 174 (VLFW1), SEQ ID NO: 187 (VLFW2), and SEQ ID NO: 196 (VLFW3)).

In some embodiments, the anti-PD-1 antibody molecule further comprises the heavy chain framework region 4 (VHFW4) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13,

BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 169) and the light chain framework region 4 (VLFW4) of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E (*e.g.*, SEQ ID NO: 208).

In some embodiments, the anti-PD-1 antibody molecule comprises a heavy chain framework region having a combination of framework regions FW1, FW2 and FW3 as shown in FIGs. 5 or 7. In other embodiment, the antibody molecule comprises a light chain framework region having a combination of framework regions FW1, FW2 and FW3 as shown in FIGs. 5 or 7. In yet other embodiments, the antibody molecule comprises a heavy chain framework region having a combination of framework regions FW1, FW2 and FW3 as shown in FIGs. 5 or 7, and a light chain framework region having a combination of framework regions FW1, FW2 and FW3 as shown in FIGs. 5 or 7.

In one embodiment, the heavy or light chain variable domain, or both, of the anti-PD-1 antibody molecule includes an amino acid sequence, which is substantially identical to an amino acid disclosed herein, *e.g.*, at least 80%, 85%, 90%, 92%, 95%, 97%, 98%, 99% or higher identical to a variable region of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1, or encoded by the nucleotide sequence in Table 1; or which differs at least 1 or 5 residues, but less than 40, 30, 20, or 10 residues, from a variable region of an antibody described herein.

In one embodiment, the heavy or light chain variable region, or both, of the anti-PD-1 antibody molecule includes an amino acid sequence encoded by a nucleic acid sequence described herein or a nucleic acid that hybridizes to a nucleic acid sequence described herein (*e.g.*, a nucleic acid sequence as shown in Tables 1 and 2) or its complement, *e.g.*, under low

stringency, medium stringency, or high stringency, or other hybridization condition described herein.

In another embodiment, the anti-PD-1 antibody molecule comprises at least one, two, three, or four antigen-binding regions, *e.g.*, variable regions, having an amino acid sequence as set forth in Table 1, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 1, 2, 5, 10, or 15 amino acid residues from the sequences shown in Table 1. In another embodiment, the anti-PD-1 antibody molecule includes a VH and/or VL domain encoded by a nucleic acid having a nucleotide sequence as set forth in Table 1, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 3, 6, 15, 30, or 45 nucleotides from the sequences shown in Table 1.

In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one, two, or three CDRs from a heavy chain variable region having an amino acid sequence as set forth in Table 1, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions). In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one, two, or three CDRs from a light chain variable region having an amino acid sequence as set forth in Table 1, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions). In yet another embodiment, the anti-PD-1 antibody molecule comprises at least one, two, three, four, five or six CDRs from heavy and light chain variable regions having an amino acid sequence as set forth in Table 1), or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions).

In one embodiment, the anti-PD-1 antibody molecule comprises at least one, two, or three CDRs and/or hypervariable loops from a heavy chain variable region having an amino acid sequence of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16,

BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, as summarized in Table 1, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions). In another

5 embodiment, the anti-PD-1 antibody molecule comprises at least one, two, or three CDRs and/or hypervariable loops from a light chain variable region having an amino acid sequence of of an antibody described herein, *e.g.*, an antibody chosen from any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12,
10 BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, as summarized in Table 1, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions). In one embodiment, the anti-PD-1
15 antibody molecule comprises all six CDRs and/or hypervariable loops described herein, *e.g.*, described in Table 1.

In one embodiment, the anti-PD-1 antibody molecule has a variable region that is identical in sequence, or which differs by 1, 2, 3, or 4 amino acids from a variable region described herein (*e.g.*, an FR region disclosed herein).

20 In one embodiment, the anti-PD-1 antibody molecule is a full antibody or fragment thereof (*e.g.*, a Fab, F(ab')₂, Fv, or a single chain Fv fragment (scFv)). In certain embodiments, the anti-PD-1 antibody molecule is a monoclonal antibody or an antibody with single specificity. The anti-PD-1 antibody molecule can also be a humanized, chimeric, camelid, shark, or an *in vitro*-generated antibody molecule. In one embodiment, the anti-PD-1 antibody molecule thereof
25 is a humanized antibody molecule. The heavy and light chains of the anti-PD-1 antibody molecule can be full-length (*e.g.*, an antibody can include at least one, and preferably two, complete heavy chains, and at least one, and preferably two, complete light chains) or can include an antigen-binding fragment (*e.g.*, a Fab, F(ab')₂, Fv, a single chain Fv fragment, a single domain antibody, a diabody (dAb), a bivalent antibody, or bispecific antibody or fragment
30 thereof, a single domain variant thereof, or a camelid antibody).

In yet other embodiments, the anti-PD-1 antibody molecule has a heavy chain constant region (Fc) chosen from, *e.g.*, the heavy chain constant regions of IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgD, and IgE; particularly, chosen from, *e.g.*, the heavy chain constant regions of IgG1, IgG2, IgG3, and IgG4, more particularly, the heavy chain constant region of IgG1 or IgG2 (*e.g.*, human IgG1, IgG2 or IgG4). In one embodiment, the heavy chain constant region is human IgG1. In another embodiment, the anti-PD-1 antibody molecule has a light chain constant region chosen from, *e.g.*, the light chain constant regions of kappa or lambda, preferably kappa (*e.g.*, human kappa). In one embodiment, the constant region is altered, *e.g.*, mutated, to modify the properties of the anti-PD-1 antibody molecule (*e.g.*, to increase or decrease one or more of: Fc receptor binding, antibody glycosylation, the number of cysteine residues, effector cell function, or complement function). For example, the constant region is mutated at positions 296 (M to Y), 298 (S to T), 300 (T to E), 477 (H to K) and 478 (N to F) to alter Fc receptor binding (*e.g.*, the mutated positions correspond to positions 132 (M to Y), 134 (S to T), 136 (T to E), 313 (H to K) and 314 (N to F) of SEQ ID NOs: 212 or 214; or positions 135 (M to Y), 137 (S to T), 139 (T to E), 316 (H to K) and 317 (N to F) of SEQ ID NOs: 215, 216, 217 or 218). In another embodiment, the heavy chain constant region of an IgG4, *e.g.*, a human IgG4, is mutated at position 228 according to EU numbering (*e.g.*, S to P), *e.g.*, as shown in Table 3. In certain embodiments, the anti-PD-1 antibody molecules comprises a human IgG4 mutated at position 228 according to EU numbering (*e.g.*, S to P), *e.g.*, as shown in Table 3; and a kappa light chain constant region, *e.g.*, as shown in Table 3. In still another embodiment, the heavy chain constant region of an IgG1, *e.g.*, a human IgG1, is mutated at one or more of position 297 according to EU numbering (*e.g.*, N to A), position 265 according to EU numbering (*e.g.*, D to A), position 329 according to EU numbering (*e.g.*, P to A), position 234 according to EU numbering (*e.g.*, L to A), or position 235 according to EU numbering (*e.g.*, L to A), *e.g.*, as shown in Table 3. In certain embodiments, the anti-PD-1 antibody molecules comprises a human IgG1 mutated at one or more of the aforesaid positions, *e.g.*, as shown in Table 3; and a kappa light chain constant region, *e.g.*, as shown in Table 3.

In one embodiment, the anti-PD-1 antibody molecule is isolated or recombinant.

In one embodiment, the anti-PD-1 antibody molecule is a humanized antibody molecule.

In one embodiment, the anti-PD-1 antibody molecule has a risk score based on T cell epitope analysis of less than 700, 600, 500, 400 or less.

In one embodiment, the anti-PD-1 antibody molecule is a humanized antibody molecule and has a risk score based on T cell epitope analysis of 300 to 700, 400 to 650, 450 to 600, or a risk score as described herein.

In one embodiment, the anti-PD-1 antibody molecule includes:

- 5 (a) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33;
- 10 (b) a VH comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32;
- 15 (c) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33; or
- (d) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.
- 20

In certain embodiments, the anti-PD-1 antibody molecule comprises:

- 25 (i) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1, SEQ ID NO: 4 or SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and
- (ii) a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

- 30 In other embodiments, the anti-PD-1 antibody molecule comprises:

(i) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1, SEQ ID NO: 4 or SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and

5 (ii) a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33.

In embodiments of the aforesaid antibody molecules, the VHCDR1 comprises the amino acid sequence of SEQ ID NO: 1. In other embodiments, the VHCDR1 comprises the amino acid sequence of SEQ ID NO: 4. In yet other embodiments, the VHCDR1 amino acid sequence of
10 SEQ ID NO: 224.

In embodiments, the aforesaid antibody molecules have a heavy chain variable region comprising at least one framework (FW) region comprising the amino acid sequence of any of SEQ ID NOs: 147, 151, 153, 157, 160, 162, 166, or 169, or an amino acid sequence at least 90% identical thereto, or having no more than two amino acid substitutions, insertions or deletions
15 compared to the amino acid sequence of any of SEQ ID NOs: 147, 151, 153, 157, 160, 162, 166, or 169.

In other embodiments, the aforesaid antibody molecules have a heavy chain variable region comprising at least one framework region comprising the amino acid sequence of any of SEQ ID NOs: 147, 151, 153, 157, 160, 162, 166, or 169.

20 In yet other embodiments, the aforesaid antibody molecules have a heavy chain variable region comprising at least two, three, or four framework regions comprising the amino acid sequences of any of SEQ ID NOs: 147, 151, 153, 157, 160, 162, 166, or 169.

In other embodiments, the aforesaid antibody molecules comprise a VHFW1 amino acid sequence of SEQ ID NO: 147 or 151, a VHFW2 amino acid sequence of SEQ ID NO: 153, 157, or 160, and a VHFW3 amino acid sequence of SEQ ID NO: 162 or 166, and, optionally, further comprising a VHFW4 amino acid sequence of SEQ ID NO: 169.

In other embodiments, the aforesaid antibody molecules have a light chain variable region comprising at least one framework region comprising the amino acid sequence of any of SEQ ID NOs: 174, 177, 181, 183, 185, 187, 191, 194, 196, 200, 202, 205, or 208, or an amino
30 acid sequence at least 90% identical thereto, or having no more than two amino acid

substitutions, insertions or deletions compared to the amino acid sequence of any of 174, 177, 181, 183, 185, 187, 191, 194, 196, 200, 202, 205, or 208.

In other embodiments, the aforesaid antibody molecules have a light chain variable region comprising at least one framework region comprising the amino acid sequence of any of
5 SEQ ID NOs: 174, 177, 181, 183, 185, 187, 191, 194, 196, 200, 202, 205, or 208.

In other embodiments, the aforesaid antibody molecules have a light chain variable region comprising at least two, three, or four framework regions comprising the amino acid sequences of any of SEQ ID NOs: 174, 177, 181, 183, 185, 187, 191, 194, 196, 200, 202, 205, or 208.

10 In other embodiments, the aforesaid antibody molecules comprise a VLFW1 amino acid sequence of SEQ ID NO: 174, 177, 181, 183, or 185, a VLFW2 amino acid sequence of SEQ ID NO: 187, 191, or 194, and a VLFW3 amino acid sequence of SEQ ID NO: 196, 200, 202, or 205, and, optionally, further comprising a VLFW4 amino acid sequence of SEQ ID NO: 208.

In other embodiments, the aforesaid antibodies comprise a heavy chain variable domain
15 comprising an amino acid sequence at least 85% identical to any of SEQ ID NOs: 38, 50, 82, or 86.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38, 50, 82, or 86.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable
20 domain comprising an amino acid sequence at least 85% identical to any of SEQ ID NOs: 42, 46, 54, 58, 62, 66, 70, 74, or 78.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 42, 46, 54, 58, 62, 66, 70, 74, or 78.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable
25 domain comprising the amino acid sequence of SEQ ID NO: 38.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 91.

30 In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 52 or SEQ ID NO: 102.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82.

5 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 84.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 86.

10 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 88.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 42.

In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 44.

15 In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46.

In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 48.

20 In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54.

In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 56.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 58.

25 In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 60.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 62.

30 In other embodiments, the aforesaid antibodies comprise a light chain comprising the amino acid sequence of SEQ ID NO: 64.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 68.

5 In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70.

In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 72.

10 In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 74.

In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 76.

In other embodiments, the aforesaid antibody molecules comprise a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 78.

15 In other embodiments, the aforesaid antibody molecules comprise a light chain comprising the amino acid sequence of SEQ ID NO: 80.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 42.

20 In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable
25 domain comprising the amino acid sequence of SEQ ID NO: 70.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable
30 domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46.

5 In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54.

10 In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 58.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 62.

15 In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 74.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 78.

25 In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

30

In other embodiments, the aforesaid antibody molecules comprise a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 86 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

5 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of SEQ ID NO: 44.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of SEQ ID NO: 56.

10 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of SEQ ID NO: 68.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 91 and a light chain comprising the amino acid sequence of SEQ ID NO: 72.

15 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 102 and a light chain comprising the amino acid sequence of SEQ ID NO: 72.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 44.

20 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 48.

25 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 52 and a light chain comprising the amino acid sequence of SEQ ID NO: 48.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 52 and a light chain comprising the amino acid sequence of SEQ ID NO: 56.

30

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 56.

5 In other embodiments, the aforesaid antibodies comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 60.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 64.

10 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 52 and a light chain comprising the amino acid sequence of SEQ ID NO: 68.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 68.

15 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 52 and a light chain comprising the amino acid sequence of SEQ ID NO: 72.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 72.

20 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 76.

25 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 40 and a light chain comprising the amino acid sequence of SEQ ID NO: 80.

In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 84 and a light chain comprising the amino acid sequence of SEQ ID NO: 72.

30

In other embodiments, the aforesaid antibodies comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 84 and a light chain comprising the amino acid sequence of SEQ ID NO: 68.

5 In other embodiments, the aforesaid antibody molecules comprise a heavy chain comprising the amino acid sequence of SEQ ID NO: 88 and a light chain comprising the amino acid sequence of SEQ ID NO: 68.

In other embodiments, the aforesaid antibody molecules are chosen from a Fab, F(ab')₂, Fv, or a single chain Fv fragment (scFv).

10 In other embodiments, the aforesaid antibody molecules comprise a heavy chain constant region selected from IgG1, IgG2, IgG3, and IgG4.

In other embodiments, the aforesaid antibody molecules comprise a light chain constant region chosen from the light chain constant regions of kappa or lambda.

15 In other embodiments, the aforesaid antibody molecules comprise a human IgG4 heavy chain constant region with a mutation at position 228 according to EU numbering or position 108 of SEQ ID NO: 212 or 214 and a kappa light chain constant region.

In other embodiments, the aforesaid antibody molecules comprise a human IgG4 heavy chain constant region with a Serine to Proline mutation at position 228 according to EU numbering or position 108 of SEQ ID NO: 212 or 214 and a kappa light chain constant region.

20 In other embodiments, the aforesaid antibody molecules comprise a human IgG1 heavy chain constant region with an Asparagine to Alanine mutation at position 297 according to EU numbering or position 180 of SEQ ID NO: 216 and a kappa light chain constant region.

25 In other embodiments, the aforesaid antibody molecules comprise a human IgG1 heavy chain constant region with an Aspartate to Alanine mutation at position 265 according to EU numbering or position 148 of SEQ ID NO: 217, and Proline to Alanine mutation at position 329 according to EU numbering or position 212 of SEQ ID NO: 217 and a kappa light chain constant region.

30 In other embodiments, the aforesaid antibody molecules comprise a human IgG1 heavy chain constant region with a Leucine to Alanine mutation at position 234 according to EU numbering or position 117 of SEQ ID NO: 218, and Leucine to Alanine mutation at position 235 according to EU numbering or position 118 of SEQ ID NO: 218 and a kappa light chain constant region.

In other embodiments, the aforesaid antibody molecules are capable of binding to human PD-1 with a dissociation constant (K_D) of less than about 0.2 nM.

In some embodiments, the aforesaid antibody molecules bind to human PD-1 with a K_D of less than about 0.2 nM, 0.15 nM, 0.1 nM, 0.05 nM, or 0.02 nM, *e.g.*, about 0.13 nM to 0.03 nM, *e.g.*, about 0.077 nM to 0.088 nM, *e.g.*, about 0.083 nM, *e.g.*, as measured by a Biacore method.

In other embodiments, the aforesaid antibody molecules bind to cynomolgus PD-1 with a K_D of less than about 0.2 nM, 0.15 nM, 0.1 nM, 0.05 nM, or 0.02 nM, *e.g.*, about 0.11 nM to 0.08 nM, *e.g.*, about 0.093 nM, *e.g.*, as measured by a Biacore method.

10 In certain embodiments, the aforesaid antibody molecules bind to both human PD-1 and cynomolgus PD-1 with similar K_D , *e.g.*, in the nM range, *e.g.*, as measured by a Biacore method. In some embodiments, the aforesaid antibody molecules bind to a human PD-1-Ig fusion protein with a K_D of less than about 0.1 nM, 0.075 nM, 0.05 nM, 0.025 nM, or 0.01 nM, *e.g.*, about 0.04 nM, *e.g.*, as measured by ELISA.

15 In some embodiments, the aforesaid antibody molecules bind to Jurkat cells that express human PD-1 (*e.g.*, human PD-1-transfected Jurkat cells) with a K_D of less than about 0.1 nM, 0.075 nM, 0.05 nM, 0.025 nM, or 0.01 nM, *e.g.*, about 0.06 nM, *e.g.*, as measured by FACS analysis.

20 In some embodiments, the aforesaid antibody molecules bind to cynomolgus T cells with a K_D of less than about 1nM, 0.75 nM, 0.5 nM, 0.25 nM, or 0.1 nM, *e.g.*, about 0.4 nM, *e.g.*, as measured by FACS analysis.

In some embodiments, the aforesaid antibody molecules bind to cells that express cynomolgus PD-1 (*e.g.*, cells transfected with cynomolgus PD-1) with a K_D of less than about 1nM, 0.75 nM, 0.5 nM, 0.25 nM, or 0.01 nM, *e.g.*, about 0.6 nM, *e.g.*, as measured by FACS analysis.

25 In certain embodiments, the aforesaid antibody molecules are not cross-reactive with mouse or rat PD-1. In other embodiments, the aforesaid antibodies are cross-reactive with rhesus PD-1. For example, the cross-reactivity can be measured by a Biacore method or a binding assay using cells that expresses PD-1 (*e.g.*, human PD-1-expressing 300.19 cells). In other
30 embodiments, the aforesaid antibody molecules bind an extracellular Ig-like domain of PD-1.

In other embodiments, the aforesaid antibody molecules are capable of reducing binding of PD-1 to PD-L1, PD-L2, or both, or a cell that expresses PD-L1, PD-L2, or both. In some embodiments, the aforesaid antibody molecules reduce (*e.g.*, block) PD-L1 binding to a cell that expresses PD-1 (*e.g.*, human PD-1-expressing 300.19 cells) with an IC₅₀ of less than about 1.5 nM, 1 nM, 0.8 nM, 0.6 nM, 0.4 nM, 0.2 nM, or 0.1 nM, *e.g.*, between about 0.79 nM and about 1.09 nM, *e.g.*, about 0.94 nM, or about 0.78 nM or less, *e.g.*, about 0.3 nM. In some embodiments, the aforesaid antibodies reduce (*e.g.*, block) PD-L2 binding to a cell that expresses PD-1 (*e.g.*, human PD-1-expressing 300.19 cells) with an IC₅₀ of less than about 2 nM, 1.5 nM, 1 nM, 0.5 nM, or 0.2 nM, *e.g.*, between about 1.05 nM and about 1.55 nM, or about 1.3 nM or less, *e.g.*, about 0.9 nM.

In other embodiments, the aforesaid antibody molecules are capable of enhancing an antigen-specific T cell response.

In embodiments, the antibody molecule is a monospecific antibody molecule or a bispecific antibody molecule. In embodiments, the antibody molecule has a first binding specificity for PD-1 and a second binding specificity for TIM-3, LAG-3, CEACAM (*e.g.*, CEACAM-1, CEACAM-3, and/or CEACAM-5), PD-L1 or PD-L2. In embodiments, the antibody molecule comprises an antigen binding fragment of an antibody, *e.g.*, a half antibody or antigen binding fragment of a half antibody.

In some embodiments, the aforesaid antibody molecules increase the expression of IL-2 from cells activated by Staphylococcal enterotoxin B (SEB) (*e.g.*, at 25 µg/mL) by at least about 2, 3, 4, 5-fold, *e.g.*, about 2 to 3-fold, *e.g.*, about 2 to 2.6-fold, *e.g.*, about 2.3-fold, compared to the expression of IL-2 when an isotype control (*e.g.*, IgG4) is used, *e.g.*, as measured in a SEB T cell activation assay or a human whole blood *ex vivo* assay.

In some embodiments, the aforesaid antibody molecules increase the expression of IFN-γ from T cells stimulated by anti-CD3 (*e.g.*, at 0.1 µg/mL) by at least about 2, 3, 4, 5-fold, *e.g.*, about 1.2 to 3.4-fold, *e.g.*, about 2.3-fold, compared to the expression of IFN-γ when an isotype control (*e.g.*, IgG4) is used, *e.g.*, as measured in an IFN-γ activity assay.

In some embodiments, the aforesaid antibody molecules increase the expression of IFN-γ from T cells activated by SEB (*e.g.*, at 3 pg/mL) by at least about 2, 3, 4, 5-fold, *e.g.*, about 0.5 to 4.5-fold, *e.g.*, about 2.5-fold, compared to the expression of IFN-γ when an isotype control (*e.g.*, IgG4) is used, *e.g.*, as measured in an IFN-γ activity assay.

In some embodiments, the aforesaid antibody molecules increase the expression of IFN- γ from T cells activated with an CMV peptide by at least about 2, 3, 4, 5-fold, *e.g.*, about 2 to 3.6-fold, *e.g.*, about 2.8-fold, compared to the expression of IFN- γ when an isotype control (*e.g.*, IgG4) is used, *e.g.*, as measured in an IFN- γ activity assay.

5 In some embodiments, the aforesaid antibody molecules increase the proliferation of CD8⁺ T cells activated with an CMV peptide by at least about 1, 2, 3, 4, 5-fold, *e.g.*, about 1.5-fold, compared to the proliferation of CD8⁺ T cells when an isotype control (*e.g.*, IgG4) is used, *e.g.*, as measured by the percentage of CD8⁺ T cells that passed through at least *n* (*e.g.*, *n* = 2 or 4) cell divisions.

10 In certain embodiments, the aforesaid antibody molecules has a C_{max} between about 100 $\mu\text{g/mL}$ and about 500 $\mu\text{g/mL}$, between about 150 $\mu\text{g/mL}$ and about 450 $\mu\text{g/mL}$, between about 250 $\mu\text{g/mL}$ and about 350 $\mu\text{g/mL}$, or between about 200 $\mu\text{g/mL}$ and about 400 $\mu\text{g/mL}$, *e.g.*, about 292.5 $\mu\text{g/mL}$, *e.g.*, as measured in monkey.

In certain embodiments, the aforesaid antibody molecules has a $T_{1/2}$ between about 250
15 hours and about 650 hours, between about 300 hours and about 600 hours, between about 350 hours and about 550 hours, or between about 400 hours and about 500 hours, *e.g.*, about 465.5 hours, *e.g.*, as measured in monkey.

In some embodiments, the aforesaid antibody molecules bind to PD-1 with a K_d slower than 5×10^{-4} , 1×10^{-4} , 5×10^{-5} , or $1 \times 10^{-5} \text{ s}^{-1}$, *e.g.*, about $2.13 \times 10^{-4} \text{ s}^{-1}$, *e.g.*, as measured by a
20 Biacore method. In some embodiments, the aforesaid antibody molecules bind to PD-1 with a K_a faster than 1×10^4 , 5×10^4 , 1×10^5 , or $5 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, *e.g.*, about $2.78 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, *e.g.*, as measured by a Biacore method.

In some embodiments, the aforesaid anti-PD-1 antibody molecules bind to one or more residues within the C strand, CC' loop, C' strand and FG loop of PD-1. The domain structure of
25 PD-1 is described, *e.g.*, in Cheng et al., "Structure and Interactions of the Human Programmed Cell Death 1 Receptor" *J. Biol. Chem.* 2013, 288:11771-11785. As described in Cheng *et. al.*, the C strand comprises residues F43-M50, the CC' loop comprises S51-N54, the C' strand comprises residues Q55-F62, and the FG loop comprises residues L108-I114 (amino acid numbering according to Chang *et al. supra*). Accordingly, in some embodiments, an anti-PD-1
30 antibody as described herein binds to at least one residue in one or more of the ranges F43-M50, S51-N54, Q55-F62, and L108-I114 of PD-1. In some embodiments, an anti-PD-1 antibody as

described herein binds to at least one residue in two, three, or all four of the ranges F43-M50, S51-N54, Q55-F62, and L108-I114 of PD-1. In some embodiments, the anti-PD-1 antibody binds to a residue in PD-1 that is also part of a binding site for one or both of PD-L1 and PD-L2.

In another aspect, the invention provides an isolated nucleic acid molecule encoding any
5 of the aforesaid antibody molecules, vectors and host cells thereof.

An isolated nucleic acid encoding the antibody heavy chain variable region or light chain variable region, or both, of any the aforesaid antibody molecules is also provided.

In one embodiment, the isolated nucleic acid encodes heavy chain CDRs 1-3, wherein said nucleic acid comprises a nucleotide sequence of SEQ ID NO: 108-112, 223, 122-126, 133-
10 137, or 144-146.

In another embodiment, the isolated nucleic acid encodes light chain CDRs 1-3, wherein said nucleic acid comprises a nucleotide sequence of SEQ ID NO: 113-120, 127-132, or 138-143.

In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence
15 encoding a heavy chain variable domain, wherein said nucleotide sequence is at least 85% identical to any of SEQ ID NO: 39, 51, 83, 87, 90, 95, or 101.

In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a heavy chain variable domain, wherein said nucleotide sequence comprises any of
SEQ ID NO: 39, 51, 83, 87, 90, 95, or 101.

20 In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a heavy chain, wherein said nucleotide sequence is at least 85% identical to any of SEQ ID NO: 41, 53, 85, 89, 92, 96, or 103.

In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a heavy chain, wherein said nucleotide sequence comprises any of SEQ ID NO: 41, 53,
25 85, 89, 92, 96, or 103.

In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a light chain variable domain, wherein said nucleotide sequence is at least 85% identical to any of SEQ ID NO: 45, 49, 57, 61, 65, 69, 73, 77, 81, 94, 98, 100, 105, or 107.

30 In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a light chain variable domain, wherein said nucleotide sequence comprises any of SEQ ID NO: 45, 49, 57, 61, 65, 69, 73, 77, 81, 94, 98, 100, 105, or 107.

In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a light chain, wherein said nucleotide sequence is at least 85% identical to any of SEQ ID NO: 45, 49, 57, 61, 65, 69, 73, 77, 81, 94, 98, 100, 105 or 107.

5 In other embodiments, the aforesaid nucleic acid further comprises a nucleotide sequence encoding a light chain, wherein said nucleotide sequence comprises any of SEQ ID NO: 45, 49, 57, 61, 65, 69, 73, 77, 81, 94, 98, 100, 105 or 107.

In certain embodiments, one or more expression vectors and host cells comprising the aforesaid nucleic acids are provided.

10 A method of producing an antibody molecule or fragment thereof, comprising culturing the host cell as described herein under conditions suitable for gene expression is also provided.

In one aspect, the invention features a method of providing an antibody molecule described herein. The method includes: providing a PD-1 antigen (*e.g.*, an antigen comprising at least a portion of a PD-1 epitope); obtaining an antibody molecule that specifically binds to the PD-1 polypeptide; and evaluating if the antibody molecule specifically binds to the PD-1
15 polypeptide, or evaluating efficacy of the antibody molecule in modulating, *e.g.*, inhibiting, the activity of the PD-1. The method can further include administering the antibody molecule to a subject, *e.g.*, a human or non-human animal.

In another aspect, the invention provides, compositions, *e.g.*, pharmaceutical compositions, which include a pharmaceutically acceptable carrier, excipient or stabilizer, and at least one of
20 the therapeutic agents, *e.g.*, anti-PD-1 antibody molecules described herein. In one embodiment, the composition, *e.g.*, the pharmaceutical composition, includes a combination of the antibody molecule and one or more agents, *e.g.*, a therapeutic agent or other antibody molecule, as described herein. In one embodiment, the antibody molecule is conjugated to a label or a therapeutic agent. *C-Met Receptor Tyrosine Kinase Inhibitor*

25 c-Met Receptor Tyrosine Kinase Inhibitor of the present invention is disclosed, for example, in US Patent 7,767,675, incorporated herein by reference in its entirety.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is 2-fluoro-N-methyl-4-[7-quinolin-6-yl-methyl]-imidazo[1,2-b][1,2,4]triazin-2-yl]benzamide or pharmaceutically acceptable salt thereof.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is 2-fluoro-N-methyl-4-[7-quinolin-6-yl-methyl)-imidazo[1,2-b][1,2,4]triazin-2yl]benzamide dihydrochloric acid salt.

In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is capmatinib.

5 In a preferred embodiment, the c-Met receptor tyrosine kinase inhibitor is capmatinib dihydrochloric acid salt. In one embodiment, capmatinib is administered at a dose of about 400-600 mg (*e.g.*, per day), *e.g.*, about 400, 500, or 600 mg, or about 400-500 or 500-600 mg.

In one embodiment, capmatinib is administered orally.

10 *Pharmaceutical Compositions and Kits*

In another aspect, the present invention provides compositions, *e.g.*, pharmaceutically acceptable compositions, which include an antibody molecule described herein, formulated together with a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, isotonic and absorption
15 delaying agents, and the like that are physiologically compatible. The carrier can be suitable for intravenous, intramuscular, subcutaneous, parenteral, rectal, spinal or epidermal administration (*e.g.* by injection or infusion).

The compositions of this invention may be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (*e.g.*, injectable and
20 infusible solutions), dispersions or suspensions, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions. The preferred mode of administration is parenteral (*e.g.*, intravenous, subcutaneous, intraperitoneal, intramuscular). In a preferred embodiment, the antibody is administered by intravenous infusion or injection. In
25 another preferred embodiment, the antibody is administered by intramuscular or subcutaneous injection.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal,
30 intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous,

subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

Therapeutic compositions typically should be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high antibody concentration. Sterile injectable solutions can be prepared by incorporating the active compound (*i.e.*, antibody or antibody portion) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

The antibody molecules can be administered by a variety of methods known in the art, although for many therapeutic applications, the preferred route/mode of administration is intravenous injection or infusion. For example, the antibody molecules can be administered by intravenous infusion at a rate of more than 20 mg/min, *e.g.*, 20-40 mg/min, and typically greater than or equal to 40 mg/min to reach a dose of about 35 to 440 mg/m², typically about 70 to 310 mg/m², and more typically, about 110 to 130 mg/m². In embodiments, the antibody molecules can be administered by intravenous infusion at a rate of less than 10mg/min; preferably less than or equal to 5 mg/min to reach a dose of about 1 to 100 mg/m², preferably about 5 to 50 mg/m², about 7 to 25 mg/m² and more preferably, about 10 mg/m². As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible

polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. *See, e.g., Sustained and Controlled Release Drug Delivery Systems*, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

In certain embodiments, an antibody molecule can be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the invention by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. Therapeutic compositions can also be administered with medical devices known in the art.

Dosage regimens are adjusted to provide the optimum desired response (*e.g.*, a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of an antibody molecule is 0.1-30 mg/kg, more preferably 1-25 mg/kg. Dosages and therapeutic regimens of the anti-PD-1 antibody molecule can be determined by a skilled artisan.

In certain embodiments, the anti-PD-1 antibody molecule is administered by injection (*e.g.*, subcutaneously or intravenously) at a dose of about 1 to 40 mg/kg, *e.g.*, 1 to 30 mg/kg, *e.g.*,

about 5 to 25 mg/kg, about 10 to 20 mg/kg, about 1 to 5 mg/kg, 1 to 10 mg/kg, 5 to 15 mg/kg, 10 to 20 mg/kg, 15 to 25 mg/kg, or about 3 mg/kg. The dosing schedule can vary from *e.g.*, once a week to once every 2, 3, or 4 weeks. In one embodiment, the anti-PD-1 antibody molecule is administered at a dose from about 10 to 20 mg/kg every other week.

5 As another example, non-limiting range for a therapeutically or prophylactically effective amount of an antibody molecule is 200-500 mg, more preferably 300-400 mg/kg. Dosages and therapeutic regimens of the anti-PD-1 antibody molecule can be determined by a skilled artisan. In certain embodiments, the anti-PD-1 antibody molecule is administered by injection (*e.g.*, subcutaneously or intravenously) at a dose (*e.g.*, a flat dose) of about 200 mg to 500 mg, *e.g.*,
10 about 250 mg to 450 mg, about 300 mg to 400 mg, about 250 mg to 350 mg, about 350 mg to 450 mg, or about 300 mg or about 400 mg. The dosing schedule (*e.g.*, flat dosing schedule) can vary from *e.g.*, once a week to once every 2, 3, 4, 5, or 6 weeks. In one embodiment the anti-PD-1 antibody molecule is administered at a dose from about 300 mg to 400 mg once every three or once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is
15 administered at a dose from about 300 mg once every three weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 400 mg once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 300 mg once every four weeks. In one embodiment, the the anti-PD-1 antibody molecule is administered at a dose from about 400 mg once every three weeks. While not wishing to be
20 bound by theory, in some embodiments, flat or fixed dosing can be beneficial to patients, for example, to save drug supply and to reduce pharmacy errors.

In some embodiments, the clearance (CL) of the anti-PD-1 antibody molecule is from about 6 to 16 mL/h, *e.g.*, about 7 to 15 mL/h, about 8 to 14 mL/h, about 9 to 12 mL/h, or about 10 to 11 mL/h, *e.g.*, about 8.9 mL/h, 10.9 mL/h, or 13.2 mL/h.

25 In some embodiments, the exponent of weight on CL of the anti-PD-1 antibody molecule is from about 0.4 to 0.7, about 0.5 to 0.6, or 0.7 or less, *e.g.*, 0.6 or less, or about 0.54.

In some embodiments, the volume of distribution at steady state (V_{ss}) of the anti-PD-1 antibody molecule is from about 5 to 10 V, *e.g.*, about 6 to 9 V, about 7 to 8 V, or about 6.5 to 7.5 V, *e.g.*, about 7.2 V.

In some embodiments, the half-life of the anti-PD-1 antibody molecule is from about 10 to 30 days, *e.g.*, about 15 to 25 days, about 17 to 22 days, about 19 to 24 days, or about 18 to 22 days, *e.g.*, about 20 days.

In some embodiments, the Cmin (*e.g.*, for a 80 kg patient) of the anti-PD-1 antibody molecule is at least about 0.4 µg/mL, *e.g.*, at least about 3.6 µg/mL, *e.g.*, from about 20 to 50 µg/mL, *e.g.*, about 22 to 42 µg/mL, about 26 to 47 µg/mL, about 22 to 26 µg/mL, about 42 to 47 µg/mL, about 25 to 35 µg/mL, about 32 to 38 µg/mL, *e.g.*, about 31 µg/mL or about 35 µg/mL. In one embodiment, the Cmin is determined in a patient receiving the anti-PD-1 antibody molecule at a dose of about 400 mg once every four weeks. In another embodiment, the Cmin is determined in a patient receiving the anti-PD-1 antibody molecule at a dose of about 300 mg once every three weeks. In some embodiments, In certain embodiments, the Cmin is at least about 50-fold higher, *e.g.*, at least about 60-fold, 65-fold, 70-fold, 75-fold, 80-fold, 85-fold, 90-fold, 95-fold, or 100-fold, *e.g.*, at least about 77-fold, higher than the EC50 of the anti-PD-1 antibody molecule, *e.g.*, as determined based on IL-2 change in an SEB *ex-vivo* assay. In other embodiments, the Cmin is at least 5-fold higher, *e.g.*, at least 6-fold, 7-fold, 8-fold, 9-fold, or 10-fold, *e.g.*, at least about 8.6-fold, higher than the EC90 of the anti-PD-1 antibody molecule, *e.g.*, as determined based on IL-2 change in an SEB *ex-vivo* assay.

The antibody molecule can be administered by intravenous infusion at a rate of more than 20 mg/min, *e.g.*, 20-40 mg/min, and typically greater than or equal to 40 mg/min to reach a dose of about 35 to 440 mg/m², typically about 70 to 310 mg/m², and more typically, about 110 to 130 mg/m². In embodiments, the infusion rate of about 110 to 130 mg/m² achieves a level of about 3 mg/kg. In other embodiments, the antibody molecule can be administered by intravenous infusion at a rate of less than 10 mg/min, *e.g.*, less than or equal to 5 mg/min to reach a dose of about 1 to 100 mg/m², *e.g.*, about 5 to 50 mg/m², about 7 to 25 mg/m², or, about 10 mg/m². In some embodiments, the antibody is infused over a period of about 30 min. It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

The pharmaceutical compositions of the invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of an antibody or antibody portion of the invention. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the modified antibody or antibody fragment may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the modified antibody or antibody fragment is outweighed by the therapeutically beneficial effects. A "therapeutically effective dosage" preferably inhibits a measurable parameter, *e.g.*, tumor growth rate by at least about 20%, more preferably by at least about 40%, even more preferably by at least about 60%, and still more preferably by at least about 80% relative to untreated subjects. The ability of a compound to inhibit a measurable parameter, *e.g.*, cancer, can be evaluated in an animal model system predictive of efficacy in human tumors. Alternatively, this property of a composition can be evaluated by examining the ability of the compound to inhibit, such inhibition *in vitro* by assays known to the skilled practitioner.

A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

Also within the scope of the invention is a kit comprising an antibody molecule described herein. The kit can include one or more other elements including: instructions for use; other reagents, *e.g.*, a label, a therapeutic agent, or an agent useful for chelating, or otherwise coupling, an antibody to a label or therapeutic agent, or a radioprotective composition; devices or other materials for preparing the antibody for administration; pharmaceutically acceptable carriers; and devices or other materials for administration to a subject.

Uses of the Combination Therapies

The combinations, *e.g.*, the anti-PD-1 antibody molecules disclosed herein, have *in vitro* and *in vivo* diagnostic, as well as therapeutic and prophylactic utilities. For example, these molecules can be administered to cells in culture, *in vitro* or *ex vivo*, or to a subject, *e.g.*, a

human subject, to treat, prevent, and/or diagnose a variety of disorders, such as cancers and infectious disorders.

Accordingly, in one aspect, the invention provides a method of modifying an immune response in a subject comprising administering to the subject the combination described herein, such that the immune response in the subject is modified. In one embodiment, the immune response is enhanced, stimulated or up-regulated.

As used herein, the term "subject" is intended to include human and non-human animals. In one embodiment, the subject is a human subject, *e.g.*, a human patient having a disorder or condition characterized by abnormal PD-1 functioning. The term "non-human animals" includes mammals and non-mammals, such as non-human primates. In one embodiment, the subject is a human. In one embodiment, the subject is a human patient in need of enhancement of an immune response. In one embodiment, the subject is immunocompromised, *e.g.*, the subject is undergoing, or has undergone a chemotherapeutic or radiation therapy. Alternatively, or in combination, the subject is, or is at risk of being, immunocompromised as a result of an infection. The methods and compositions described herein are suitable for treating human patients having a disorder that can be treated by augmenting the T-cell mediated immune response. For example, the methods and compositions described herein can enhance a number of immune activities. In one embodiment, the subject has increased number or activity of tumour-infiltrating T lymphocytes (TILs). In another embodiment, the subject has increased expression or activity of interferon-gamma (IFN- γ). In yet another embodiment, the subject has decreased PD-L1 expression or activity.

Therapeutic Uses

Blockade of PD-1 can enhance an immune response to cancerous cells in a subject. The ligand for PD-1, PD-L1, is not expressed in normal human cells, but is abundant in a variety of human cancers (Dong *et al.* (2002) *Nat Med* 8:787-9). The interaction between PD-1 and PD-L1 can result in a decrease in tumor infiltrating lymphocytes, a decrease in T-cell receptor mediated proliferation, and/or immune evasion by the cancerous cells (Dong *et al.* (2003) *J Mol Med* 81:281-7; Blank *et al.* (2005) *Cancer Immunol. Immunother.* 54:307-314; Konishi *et al.* (2004) *Clin. Cancer Res.* 10:5094-100). Immune suppression can be reversed by inhibiting the local interaction of PD-1 to PD-L1; the effect is additive when the interaction of PD-1 to PD-L2 is

blocked as well (Iwai *et al.* (2002) *PNAS* 99:12293-7; Brown *et al.* (2003) *J. Immunol.* 170:1257-66). Thus, inhibition of PD-1 can result in augmenting an immune response.

In one aspect, the invention relates to treatment of a subject *in vivo* using an anti-PD-1 antibody molecule such that growth of cancerous tumors is inhibited or reduced. An anti-PD-1
5 antibody may be used alone to inhibit the growth of cancerous tumors. Alternatively, an anti-PD-1 antibody may be used in combination with one or more of: a standard of care treatment (*e.g.*, for cancers or infectious disorders), another antibody or antigen-binding fragment thereof, an immunomodulator (*e.g.*, an activator of a costimulatory molecule or an inhibitor of an inhibitory molecule); a vaccine, *e.g.*, a therapeutic cancer vaccine; or other forms of cellular
10 immunotherapy, as described below.

Accordingly, in one embodiment, the invention provides a method of inhibiting growth of tumor cells in a subject, comprising administering to the subject a therapeutically effective amount of an anti-PD-1 antibody molecule described herein.

In one embodiment, the methods are suitable for the treatment of cancer *in vivo*. To
15 achieve antigen-specific enhancement of immunity, the anti-PD-1 antibody molecule can be administered together with an antigen of interest. When antibodies to PD-1 are administered in combination with one or more agents, the combination can be administered in either order or simultaneously.

In another aspect, a method of treating a subject, *e.g.*, reducing or ameliorating, a
20 hyperproliferative condition or disorder (*e.g.*, a cancer), *e.g.*, solid tumor, a hematological cancer, soft tissue tumor, or a metastatic lesion, in a subject is provided. The method includes administering to the subject one or more of the combinations disclosed herein.

As used herein, the term "cancer" is meant to include all types of cancerous growths or oncogenic processes, metastatic tissues or malignantly transformed cells, tissues, or organs,
25 irrespective of histopathologic type or stage of invasiveness. Examples of cancerous disorders include, but are not limited to, solid tumors, hematological cancers, soft tissue tumors, and metastatic lesions. Examples of solid tumors include malignancies, *e.g.*, sarcomas, and carcinomas (including adenocarcinomas and squamous cell carcinomas), of the various organ systems, such as those affecting liver, lung, breast, lymphoid, gastrointestinal (*e.g.*, colon),
30 genitourinary tract (*e.g.*, renal, urothelial cells), prostate and pharynx. Adenocarcinomas include malignancies such as most colon cancers, rectal cancer, renal-cell carcinoma, liver cancer, non-

small cell carcinoma of the lung, cancer of the small intestine and cancer of the esophagus. Squamous cell carcinomas include malignancies, *e.g.*, in the lung, esophagus, skin, head and neck region, oral cavity, anus, and cervix. In one embodiment, the cancer is a melanoma, *e.g.*, an advanced stage melanoma. Metastatic lesions of the aforementioned cancers can also be treated or prevented using the methods and compositions of the invention.

Exemplary cancers whose growth can be inhibited using the antibodies molecules disclosed herein include cancers typically responsive to immunotherapy. Non-limiting examples of preferred cancers for treatment include melanoma (*e.g.*, metastatic malignant melanoma), renal cancer (*e.g.*, clear cell carcinoma), prostate cancer (*e.g.*, hormone refractory prostate adenocarcinoma), breast cancer, colon cancer and lung cancer (*e.g.*, non-small cell lung cancer). Additionally, refractory or recurrent malignancies can be treated using the antibody molecules described herein.

Examples of other cancers that can be treated include bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular malignant melanoma, uterine cancer, ovarian cancer, rectal cancer, anal cancer, gastro-esophageal, stomach cancer, testicular cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Merkel cell cancer, Hodgkin lymphoma, non-Hodgkin lymphoma, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, chronic or acute leukemias including acute myeloid leukemia, chronic myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, solid tumors of childhood, lymphocytic lymphoma, cancer of the bladder, multiple myeloma, myelodysplastic syndromes, cancer of the kidney or ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally induced cancers including those induced by asbestos (*e.g.*, mesothelioma), and combinations of said cancers.

In some embodiments, the therapies here can be used to treat a patient that has (or is identified as having) a cancer associated with an infection, *e.g.*, a viral or bacterial infection. Exemplary cancers include cervical cancer, anal cancer, HPV-associated head and neck

squamous cell cancer, HPV-associated esophageal papillomas, HHV6-associated lymphomas, EBV-associated lymphomas (including Burkitt lymphoma), Gastric MALT lymphoma, other infection-associated MALT lymphomas, HCC, and Kaposi's sarcoma.

In other embodiments, the cancer is a hematological malignancy or cancer including but is not limited to a leukemia or a lymphoma. For example, the anti-PD-1 antibody molecule can be used to treat cancers and malignancies including, but not limited to, *e.g.*, acute leukemias including but not limited to, *e.g.*, B-cell acute lymphoid leukemia ("BALL"), T-cell acute lymphoid leukemia ("TALL"), acute lymphoid leukemia (ALL); one or more chronic leukemias including but not limited to, *e.g.*, chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL); additional hematologic cancers or hematologic conditions including, but not limited to, *e.g.*, B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, diffuse large B cell lymphoma, Follicular lymphoma, Hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and "preleukemia" which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and the like.

In one embodiment, the cancer is chosen from a lung cancer (*e.g.*, a non-small cell lung cancer (NSCLC) (*e.g.*, a NSCLC with squamous and/or non-squamous histology, or a NSCLC adenocarcinoma)), a melanoma (*e.g.*, an advanced melanoma), a renal cancer (*e.g.*, a renal cell carcinoma, *e.g.*, clear cell renal cell carcinoma), a liver cancer (*e.g.*, hepatocellular carcinoma), a myeloma (*e.g.*, a multiple myeloma), a prostate cancer, a breast cancer (*e.g.*, a breast cancer that does not express one, two or all of estrogen receptor, progesterone receptor, or Her2/neu, *e.g.*, a triple negative breast cancer), a colorectal cancer, a pancreatic cancer, a head and neck cancer (*e.g.*, head and neck squamous cell carcinoma (HNSCC), anal cancer, gastro-esophageal cancer, thyroid cancer, cervical cancer, a lymphoproliferative disease (*e.g.*, a post-transplant lymphoproliferative disease) or a hematological cancer, T-cell lymphoma, a non-Hodgkin's lymphoma, or a leukemia (*e.g.*, a myeloid leukemia).

In another embodiment, the cancer is chosen from a carcinoma (*e.g.*, advanced or metastatic carcinoma), melanoma or a lung carcinoma, *e.g.*, a non-small cell lung carcinoma.

In one embodiment, the cancer is a lung cancer, *e.g.*, a non-small cell lung cancer.

In another embodiment, the cancer is a hepatocarcinoma, *e.g.*, an advanced hepatocarcinoma, with or without a viral infection, *e.g.*, a chronic viral hepatitis.

In another embodiment, the cancer is a prostate cancer, *e.g.*, an advanced prostate cancer.

5 In yet another embodiment, the cancer is a myeloma, *e.g.*, multiple myeloma.

In yet another embodiment, the cancer is a renal cancer, *e.g.*, a renal cell carcinoma (RCC) (*e.g.*, a metastatic RCC or clear cell renal cell carcinoma).

Methods and compositions disclosed herein are useful for treating metastatic lesions associated with the aforementioned cancers.

10

Infectious Diseases

Other methods of the invention are used to treat patients that have been exposed to particular toxins or pathogens. Accordingly, another aspect of the invention provides a method of treating an infectious disease in a subject comprising administering to the subject a
15 combination as disclosed herein, *e.g.*, a combination including an anti-PD-1 antibody molecule, such that the subject is treated for the infectious disease.

In the treatment of infection (*e.g.*, acute and/or chronic), administration of the anti-PD-1 antibody molecules can be combined with conventional treatments in addition to or in lieu of stimulating natural host immune defenses to infection. Natural host immune defenses to
20 infection include, but are not limited to inflammation, fever, antibody-mediated host defense, T-lymphocyte-mediated host defenses, including lymphokine secretion and cytotoxic T-cells (especially during viral infection), complement mediated lysis and opsonization (facilitated phagocytosis), and phagocytosis. The ability of the anti-PD-1 antibody molecules to reactivate dysfunctional T-cells would be useful to treat chronic infections, in particular those in which
25 cell-mediated immunity is important for complete recovery.

Similar to its application to tumors as discussed above, antibody mediated PD-1 blockade can be used alone, or as an adjuvant, in combination with vaccines, to stimulate the immune response to pathogens, toxins, and self-antigens. Examples of pathogens for which this therapeutic approach may be particularly useful, include pathogens for which there is currently
30 no effective vaccine, or pathogens for which conventional vaccines are less than completely effective. These include, but are not limited to HIV, Hepatitis (A, B, & C), Influenza, Herpes,

Giardia, Malaria, Leishmania, *Staphylococcus aureus*, *Pseudomonas Aeruginosa*. PD-1 blockade is particularly useful against established infections by agents such as HIV that present altered antigens over the course of the infections. These novel epitopes are recognized as foreign at the time of anti-human PD-1 administration, thus provoking a strong T cell response that is not dampened by negative signals through PD-1.

Combination Therapies

Combinations disclosed herein, e.g., combination of PD-1 antibody molecules, with one or more further therapeutics are provided herein. Many of the combinations in this section are useful in treating cancer, but other indications are also described. This section focuses on combinations of anti-PD-1 antibody molecules. In the combinations herein below, in one embodiment, the anti-PD-1 antibody molecule includes:

(a) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33;

(b) a VH comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32;

(c) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33; or

(d) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

In the combinations herein below, in another embodiment, the anti-PD-1 antibody molecule comprises (i) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence chosen from SEQ ID NO: 1, SEQ ID NO: 4, or SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2 or SEQ ID NO: 5; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and (ii) a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10 or SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 11 or SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 32 or SEQ ID NO: 33.

In another embodiment, the combination, *e.g.*, a combination comprising an anti-PD-1 antibody molecule as described herein, is used in combination with a c-Met receptor tyrosine kinase inhibitor inhibitor, 2-fluoro-N-methyl-4-(7-(quinolin-6-ylmethyl)imidazo[1,2-b][1,2,4]triazin-2-yl)benzamide (Capmatinib), or a dihydrochloric salt thereof, or a compound disclosed in PCT Publication No. WO 2007/070514, to treat a disorder, *e.g.*, a disorder described herein. In one embodiment, the c-Met receptor tyrosine kinase inhibitor inhibitor is 2-fluoro-N-methyl-4-(7-(quinolin-6-ylmethyl)imidazo[1,2-b][1,2,4]triazin-2-yl)benzamide (Capmatinib), or a dihydrochloric salt thereof, or a compound disclosed in PCT Publication No. WO 2007/070514. In one embodiment, a PD-1 antibody molecule is used in combination with 2-fluoro-N-methyl-4-(7-(quinolin-6-ylmethyl)imidazo[1,2-b][1,2,4]triazin-2-yl)benzamide (Capmatinib), or a dihydrochloric salt thereof, or a compound disclosed in PCT Publication No. WO 2007/070514, to treat a disorder such as colorectal cancer, myeloid leukemia, liver cancer, lung cancer, hematological cancer, autoimmune disease, non-Hodgkin lymphoma, or thrombocytopenia.

In one embodiment, the c-Met receptor tyrosine kinase inhibitor inhibitor or a 2-fluoro-N-methyl-4-(7-(quinolin-6-ylmethyl)imidazo[1,2-b][1,2,4]triazin-2-yl)benzamide (Capmatinib), or a dihydrochloric salt thereof is administered at a dose of about, 200-600mg, preferably 400-600 mg (*e.g.*, per day), *e.g.*, about 400, 500, or 600 mg, or about 400-500 or 500-600 mg.

In one embodiment, Capmatinib is administered orally.

In one embodiment, Capmatinib is administered orally, twice daily (BID).

In one embodiment, Capmatinib is administered orally, 100mg, twice daily (BID).

In one embodiment, Capmatinib is administered orally, 150mg, twice daily (BID).

In one embodiment, Capmatinib is administered orally, 200mg, twice daily (BID).

In one embodiment, Capmatinib is administered orally, 300mg, twice daily (BID).

In one embodiment, Capmatinib is administered orally, twice daily (BID) continuously.

In one embodiment, Capmatinib is administered orally, 100mg, twice daily (BID)

5 continuously.

In one embodiment, Capmatinib is administered orally, 150mg, twice daily (BID) continuously.

In one embodiment, Capmatinib is administered orally, 200mg, twice daily (BID) continuously.

10 In one embodiment, Capmatinib is administered orally, 300mg, twice daily (BID) continuously.

Exemplary huMLR assay and B or T cell proliferation assays are provided below.

Human mixed lymphocyte reaction

The Mixed Lymphocyte Reaction (MLR) is a functional assay which measures the
15 proliferative response of lymphocytes from one individual (the responder) to lymphocytes from another individual (the stimulator). To perform an allogeneic MLR, peripheral blood mononuclear cells (PBMC) from three donors were isolated from buffy-coats of unknown HLA type (Kantonspital Blutspendezentrum from Bern and Aarau, Switzerland). The cells were prepared at 2.10⁵ in 0.2mL of culture medium containing RPMI 1640 GlutaMAX™ with 10%
20 fetal calf serum (FCS), 100U penicillin/ 100µg streptomycin, 50µM 2-Mercaptoethanol. Individual 2-way reactions were set up by mixing PBMC from two different donors at a 1:1 ratio and co-cultures were done in triplicates in flat-bottomed 96-well tissue culture plates for 6 days at 37°C, 5% CO₂, in presence or not of an 8-point concentration range of test compounds. Cells were pulsed with 3H-TdR (1 µCi/0.2mL) for the last 16h of culture and incorporated
25 radioactivity was used as a measure of cell proliferation. The concentration that inhibited 50% of the maximal huMLR response (IC₅₀) was calculated for each compound. Cyclosporine was used as a positive control of huMLR inhibition.

Human B cell proliferation assay

PBMC were freshly isolated by Ficoll-Paque density gradient from human blood and subjected to negative B-cell isolation. B cells were resuspended in culture medium (RPMI 1640, HEPES, 10% FCS, 50µg/mL gentamicine, 50µM 2-Mercaptoethanol, 1x ITS (Insulin, Transferrin and Sodium Selenite), 1x Non-Essential Amino-Acids) at a concentration of 9.104 per well in a flat-bottom 96-well culture plate. B cell stimulation was performed by human anti-IgM antibody molecule (30ug/mL) and IL-4 (75ng/mL) or by CD40 ligand (3ug/mL) and IL-4 (75ng/mL) in presence or not of a 7-point concentration range of test compounds. After 72h of culture at 37°C, 10% CO₂, cells were pulsed with 3H-TdR (1 µCi/well) for the last 6h of culture. B cells were then harvested and the incorporation of thymidine was measured using a scintillation counter. Of each duplicate treatment, the mean was calculated and these data were plotted in XLfit 4 to determine the respective IC₅₀ values.

Human T cell proliferation assay

PBMC were freshly isolated by Ficoll-Paque density gradient from human blood and subjected to negative isolation of T cells. T cells were prepared in culture medium (RPMI 1640, HEPES, 10% FCS, 50µg/mL gentamicine, 50µM 2-Mercaptoethanol, 1x ITS (Insulin, Transferrin and Sodium Selenite), 1x Non-Essential Amino-Acids) at a concentration of 8.104 per well in a flat-bottom 96-well culture plate. T cell stimulation was performed by human anti-CD3 antibody molecule (10ug/mL) or by human anti-CD3 antibody molecule (5µg/mL) and anti-CD28 antibody molecule (1µg/mL) in presence or not of a 7-point concentration range of test compounds. After 72h of culture at 37°C, 10% CO₂, cells were pulsed with 3H-TdR (1 µCi/well) for the last 6h of culture. Cell proliferation was measured by the incorporation of thymidine allowing IC₅₀ determination for each tested compound.

Nucleic Acids

The invention also features nucleic acids comprising nucleotide sequences that encode heavy and light chain variable regions and CDRs or hypervariable loops of the anti-PD-1 antibody molecules, as described herein. For example, the invention features a first and second nucleic acid encoding heavy and light chain variable regions, respectively, of an anti-PD-1 antibody molecule chosen from one or more of the antibody molecules disclosed herein. The

nucleic acid can comprise a nucleotide sequence as set forth in the tables herein, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 3, 6, 15, 30, or 45 nucleotides from the sequences shown in the tables herein).

5 In certain embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs or hypervariable loops from a heavy chain variable region having an amino acid sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one or more substitutions, *e.g.*, conserved substitutions). In other embodiments, the
10 nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs or hypervariable loops from a light chain variable region having an amino acid sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one or more substitutions, *e.g.*, conserved substitutions). In yet another embodiment, the nucleic acid can comprise a
15 nucleotide sequence encoding at least one, two, three, four, five, or six CDRs or hypervariable loops from heavy and light chain variable regions having an amino acid sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one or more substitutions, *e.g.*, conserved substitutions).

20 In certain embodiments, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs or hypervariable loops from a heavy chain variable region having the nucleotide sequence as set forth in the tables herein, a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein). In another
25 embodiment, the nucleic acid can comprise a nucleotide sequence encoding at least one, two, or three CDRs or hypervariable loops from a light chain variable region having the nucleotide sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein). In yet another embodiment, the
30 nucleic acid can comprise a nucleotide sequence encoding at least one, two, three, four, five, or six CDRs or hypervariable loops from heavy and light chain variable regions having the

nucleotide sequence as set forth in the tables herein, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or capable of hybridizing under the stringency conditions described herein).

In another aspect, the application features host cells and vectors containing the nucleic acids described herein. The nucleic acids may be present in a single vector or separate vectors present in the same host cell or separate host cell, as described in more detail hereinbelow.

In certain embodiments, one or more nucleic acid molecule that comprises one or both nucleotide sequences that encode heavy and light chain variable regions, CDRs, hypervariable loops, framework regions of the anti-PD-1 antibody molecules is provided. In certain
10 embodiments, the nucleotide sequence that encodes the anti-PD-1 antibody molecule is codon optimized. For example, the invention features a first and second nucleic acid encoding heavy and light chain variable regions, respectively, of an anti-PD-1 antibody molecule chosen from one or more of, *e.g.*, any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09,
15 BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, as summarized in Table 1, or a sequence substantially identical thereto. For example, the nucleic acid can comprise a nucleotide sequence as set forth in Tables 1 and 2, or a sequence substantially identical thereto (*e.g.*, a sequence at least about
20 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 3, 6, 15, 30, or 45 nucleotides from the sequences shown in Tables 1 and 2).

In other embodiments, the nucleic acid molecule comprises a nucleotide sequence that encodes a heavy chain variable domain and/or a heavy chain constant region comprising the amino acid sequence of BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-
25 Clone-D, or BAP049-Clone-E; or as described in Table 1; or the nucleotide sequence in Table 1; or a sequence substantially identical (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical) to any of the aforesaid sequences.

In other embodiments, the nucleic acid molecule comprises a nucleotide sequence that encodes a light chain variable domain and/or a light chain constant region comprising the amino
30 acid sequence of BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E; or as described in Table 1; or the nucleotide sequence in Table 1; or a

sequence substantially identical (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical) to any of the aforesaid sequences.

The aforesaid nucleotide sequences encoding the anti-PD-1 heavy and light chain variable domain and constant regions can be present in a separate nucleic acid molecule, or in the same nucleic acid molecule. In certain embodiments, the nucleic acid molecules comprise a nucleotide sequence encoding a leader sequence, *e.g.*, a leader sequence as shown in Table 4, or a sequence substantially identical thereto.

In certain embodiments, the nucleic acid molecule comprises a nucleotide sequence encoding at least one, two, or three CDRs, or hypervariable loops, from a heavy chain variable region having an amino acid sequence as set forth in Table 1, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions).

In another embodiment, the nucleic acid molecule comprises a nucleotide sequence encoding at least one, two, or three CDRs, or hypervariable loops, from a light chain variable region having an amino acid sequence as set forth in Table 1, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions).

In yet another embodiment, the nucleic acid molecule comprises a nucleotide sequence encoding at least one, two, three, four, five, or six CDRs, or hypervariable loops, from heavy and light chain variable regions having an amino acid sequence as set forth in Table 1, or a sequence substantially homologous thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, and/or having one, two, three or more substitutions, insertions or deletions, *e.g.*, conserved substitutions).

In one embodiment, the nucleic acid molecule includes a nucleotide sequence encoding an anti-PD-1 antibody molecule that includes a substitution in the light chain CDR3 at position 102 of the light variable region, *e.g.*, a substitution of a cysteine to tyrosine, or a cysteine to serine residue, at position 102 of the light variable region according to Table 1 (*e.g.*, SEQ ID NO: 16 or 24 for murine or chimeric, unmodified; or any of SEQ ID NOs: 34, 42, 46, 54, 58, 62, 66, 70, 74, or 78 for a modified sequence).

In another embodiment, the nucleic acid molecule includes one or more heavy chain framework region (*e.g.*, any of VHFW1 (type a), VHFW1 (type b), VHFW2 (type a), VHFW2 (type b), VHFW2 (type c), VHFW3 (type a), VHFW3 (type b), or VHFW4, or any combination thereof, *e.g.*, a framework combination as described herein) for any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, as summarized in Table 1 and 2, or a sequence substantially identical thereto. For example, the nucleic acid molecule can comprise a nucleotide sequence as set forth in Tables 1 and 2, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 3, 6, 15, 30, or 45 nucleotides from the sequences shown in Tables 1 and 2).

In another embodiment, the nucleic acid molecule includes one or more light chain framework region (*e.g.*, any of VLFW1 (type a), VLFW1 (type b), VLFW1 (type c), VLFW1 (type d), VLFW1 (type e), VLFW2 (type a), VLFW2 (type b), VLFW2 (type c), VLFW3 (type a), VLFW3 (type b), VLFW3 (type c), VLFW3 (type d), or VLFW4, or any combination thereof, *e.g.*, a framework combination as described herein) for any of BAP049-hum01, BAP049-hum02, BAP049-hum03, BAP049-hum04, BAP049-hum05, BAP049-hum06, BAP049-hum07, BAP049-hum08, BAP049-hum09, BAP049-hum10, BAP049-hum11, BAP049-hum12, BAP049-hum13, BAP049-hum14, BAP049-hum15, BAP049-hum16, BAP049-Clone-A, BAP049-Clone-B, BAP049-Clone-C, BAP049-Clone-D, or BAP049-Clone-E, as summarized in Table 1 and 2, or a sequence substantially identical thereto. For example, the nucleic acid molecule can comprise a nucleotide sequence as set forth in Tables 1 and 2, or a sequence substantially identical thereto (*e.g.*, a sequence at least about 85%, 90%, 95%, 99% or more identical thereto, or which differs by no more than 3, 6, 15, 30, or 45 nucleotides from the sequences shown in Tables 1 and 2).

In another embodiment, the nucleic acid molecule includes one or more heavy chain framework region and one or more light chain framework region as described herein. The heavy and light chain framework regions may be present in the same vector or separate vectors.

Vectors and Host Cells

In another aspect, the application features host cells and vectors containing the nucleic acids described herein. The nucleic acids may be present in a single vector or separate vectors present in the same host cell or separate host cell.

5 In one embodiment, the vectors comprise nucleotides encoding an antibody molecule described herein. In one embodiment, the vectors comprise the nucleotide sequences described herein. The vectors include, but are not limited to, a virus, plasmid, cosmid, lambda phage or a yeast artificial chromosome (YAC).

10 Numerous vector systems can be employed. For example, one class of vectors utilizes DNA elements which are derived from animal viruses such as, for example, bovine papilloma virus, polyoma virus, adenovirus, vaccinia virus, baculovirus, retroviruses (Rous Sarcoma Virus, MMTV or MOMLV) or SV40 virus. Another class of vectors utilizes RNA elements derived from RNA viruses such as Semliki Forest virus, Eastern Equine Encephalitis virus and Flaviviruses.

15 Additionally, cells which have stably integrated the DNA into their chromosomes may be selected by introducing one or more markers which allow for the selection of transfected host cells. The marker may provide, for example, prototrophy to an auxotrophic host, biocide resistance (*e.g.*, antibiotics), or resistance to heavy metals such as copper, or the like. The selectable marker gene can be either directly linked to the DNA sequences to be expressed, or
20 introduced into the same cell by cotransformation. Additional elements may also be needed for optimal synthesis of mRNA. These elements may include splice signals, as well as transcriptional promoters, enhancers, and termination signals.

Once the expression vector or DNA sequence containing the constructs has been prepared for expression, the expression vectors may be transfected or introduced into an appropriate host
25 cell. Various techniques may be employed to achieve this, such as, for example, protoplast fusion, calcium phosphate precipitation, electroporation, retroviral transduction, viral transfection, gene gun, lipid based transfection or other conventional techniques. In the case of protoplast fusion, the cells are grown in media and screened for the appropriate activity.

Methods and conditions for culturing the resulting transfected cells and for recovering the
30 antibody molecule produced are known to those skilled in the art, and may be varied or

optimized depending upon the specific expression vector and mammalian host cell employed, based upon the present description.

The invention also provides host cells comprising a nucleic acid encoding an antibody molecule as described herein.

In one embodiment, the host cells are genetically engineered to comprise nucleic acids encoding the antibody molecule.

In one embodiment, the host cells are genetically engineered by using an expression cassette. The phrase "expression cassette," refers to nucleotide sequences, which are capable of affecting expression of a gene in hosts compatible with such sequences. Such cassettes may include a promoter, an open reading frame with or without introns, and a termination signal. Additional factors necessary or helpful in effecting expression may also be used, such as, for example, an inducible promoter.

The invention also provides host cells comprising the vectors described herein.

The cell can be, but is not limited to, a eukaryotic cell, a bacterial cell, an insect cell, or a human cell. Suitable eukaryotic cells include, but are not limited to, Vero cells, HeLa cells, COS cells, CHO cells, HEK293 cells, BHK cells and MDCKII cells. Suitable insect cells include, but are not limited to, Sf9 cells.

In some embodiments, the host cell is an eukaryotic cell, *e.g.*, a mammalian cell, an insect cell, a yeast cell, or a prokaryotic cell, *e.g.*, *E. coli*. For example, the mammalian cell can be a cultured cell or a cell line. Exemplary mammalian cells include lymphocytic cell lines (*e.g.*, NSO), Chinese hamster ovary cells (CHO), COS cells, oocyte cells, and cells from a transgenic animal, *e.g.*, mammary epithelial cell.

Table 1. Amino acid and nucleotide sequences for murine, chimeric and humanized antibody molecules. The antibody molecules include murine mAb BAP049, chimeric mAbs BAP049-chi and BAP049-chi-Y, and humanized mAbs BAP049-hum01 to BAP049-hum16 and BAP049-Clone-A to BAP049-Clone-E. The amino acid and nucleotide sequences of the heavy and light chain CDRs, the heavy and light chain variable regions, and the heavy and light chains are shown.

BAP049 HC		
-----------	--	--

SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 6	VH	QVQLQQPGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSTTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGLVTVSA
SEQ ID NO: 7	DNA VH	CAGGTCCAGCTGCAGCAACCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAAGGG ACTCTGGTCACTGTCTCTGCA
SEQ ID NO: 8	VH	QVQLQQSGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSTTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGLVTVSA
SEQ ID NO: 9	DNA VH	CAGGTCCAGCTGCAGCAGTCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAAGGG ACTCTGGTCACTGTCTCTGCA
BAP049 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 12 (Kabat)	LCDR3	QNDYSYPCT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 15 (Chothia)	LCDR3	DYSYPC
SEQ ID NO: 16	VL	DIVMTQSPSSLTVTAGEKVTMSCKSSQSLLDSG NQKNFLTWYQQKPGQPPLLIIFWASTRESGVPD RFTGSGSVTDFTLTISVQAEDLAVYYCQNDYS YPCTFGGGTKLEIK
SEQ ID NO: 17	DNA VL	GACATTGTGATGACCCAGTCTCCATCCTCCCTG ACTGTGACAGCAGGAGAGAAGGTCACTATGAGC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCAGGGCAGCCTCCTAACTGTTGATCTTC

		TGGGCATCCACTAGGGAATCTGGGGTCCCTGAT CGCTTCACAGGCAGTGGATCTGTAACAGATTTC ACTCTCACCATCAGCAGTGTGCAGGCTGAAGAC CTGGCAGTTTATTACTGTCAGAATGATTATAGT TATCCGTGCACGTTCCGAGGGGGGACCAAGCTG GAAATAAAA
BAP049-chi HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 18	VH	QVQLQQPGSELVVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGTTVTVSS
SEQ ID NO: 19	DNA VH	CAGGTCCAGCTGCAGCAGCCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 20	HC	QVQLQQPGSELVVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVTPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 21	DNA HC	CAGGTCCAGCTGCAGCAGCCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC

		AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTGTTCC AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCCTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTACAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACCTACAAGACCACGCCTCCCGTGTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
SEQ ID NO: 22	VH	QVQLQQSGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 23	DNA VH	CAGGTCCAGCTGCAGCAGTCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 30	HC	QVQLQQSGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVTPSSSLGKITY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL

		HNHYTQKSLSLSLGK
		CAGGTCCAGCTGCAGCAGTCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTCACTACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGGAAGTCAAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTCGTGGTGGTGGACGTGAGC CAGGAAGACCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCCTCACCGTCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAGGCGCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACCTACAAGACCACGCCTCCCGTGTCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
SEQ ID NO: 31	DNA HC	
BAP049-chi LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 12 (Kabat)	LCDR3	QNDYSYPCT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 15 (Chothia)	LCDR3	DYSYPC
		DIVMTQSPSSLTVTAGEKVTMSCKSSQSLLDSG NQKNFLTWYQQKPGQPPKLLIFWASTRESGVPD RFTGSGSVTDFTLTISVQAEDLAVYYCQNDYS YPCTFGQGTKVEIK
SEQ ID NO: 24	VL	

SEQ ID NO: 25	DNA VL	GACATTGTGATGACCCAGTCTCCATCCTCCCTG ACTGTGACAGCAGGAGAGAAGGTCACTATGAGC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCAGGGCAGCCTCCTAAACTGTTGATCTTC TGGGCATCCACTAGGGAATCTGGGGTCCCTGAT CGCTTCACAGGCAGTGGATCTGTAACAGATTTT ACTCTCACCATCAGCAGTGTGCAGGCTGAAGAC CTGGCAGTTTATTACTGTGAGAATGATTATAGT TATCCGTGCACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 26	LC	DIVMTQSPSSLTVTAGEKVTMSCKSSQSLDSG NQKNFLTWYQQKPGQPPKLLIFWASTRESGVDP RFTGSGSVTDFTLTISVQAEDLAVYYCQNDYS YPCTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 27	DNA LC	GACATTGTGATGACCCAGTCTCCATCCTCCCTG ACTGTGACAGCAGGAGAGAAGGTCACTATGAGC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCAGGGCAGCCTCCTAAACTGTTGATCTTC TGGGCATCCACTAGGGAATCTGGGGTCCCTGAT CGCTTCACAGGCAGTGGATCTGTAACAGATTTT ACTCTCACCATCAGCAGTGTGCAGGCTGAAGAC CTGGCAGTTTATTACTGTGAGAATGATTATAGT TATCCGTGCACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-chi-Y HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 18	VH	QVQLQQPGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSTTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQTTVTVSS
SEQ ID NO: 19	DNA VH	CAGGTCCAGCTGCAGCAGCCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT

		GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 20	HC	QVQLQQPGSELVVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQGLEWIGNIYPGTGGSNFDEKFN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 21	DNA HC	CAGGTCCAGCTGCAGCAGCCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTCTGTTCCCCCA AAACCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA

		ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
SEQ ID NO: 22	VH	QVQLQQSGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGTTVTVSS
SEQ ID NO: 23	DNA VH	CAGGTCCAGCTGCAGCAGTCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 30	HC	QVQLQQSGSELVRPGASVKLSCKASGYTFTTYW MHWVRQRPQGQLEWIGNIYPGTGGSNFDEKFKN RTSLTVDTSSSTAYMHLASLTSEDSAVYYCTRW TTGTGAYWGQGTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 31	DNA HC	CAGGTCCAGCTGCAGCAGTCTGGGTCTGAGCTG GTGAGGCCTGGAGCTTCAGTGAAGCTGTCCTGC AAGGCGTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGAGGCAGAGGCCTGGACAAGGC CTTGAGTGGATTGGAAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAAAAC AGGACCTCACTGACTGTAGACACATCCTCCACC ACAGCCTACATGCACCTCGCCAGCCTGACATCT GAGGACTCTGCGGTCTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCCTGGTGGTGGACGTGAGC

		CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCCTCACCGTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-chi-Y LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 34	VL	DIVMTQSPSSLTVTAGEKVTMSCKSSQSLLDSG NQKNFLTWYQQKPGQPPKLLIFWASTRESGVPD RFTGSGSVTDFTLTISSVQAEDLAVYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 35	DNA VL	GACATTGTGATGACCCAGTCTCCATCCTCCCTG ACTGTGACAGCAGGAGAGAAGGTCACTATGAGC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCAGGGCAGCCTCCTAAACTGTTGATCTTC TGGGCATCCACTAGGGAATCTGGGGTCCCTGAT CGCTTCACAGGCAGTGGATCTGTAACAGATTTC ACTCTCACCATCAGCAGTGTGCAGGCTGAAGAC CTGGCAGTTTATTACTGTGAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 36	LC	DIVMTQSPSSLTVTAGEKVTMSCKSSQSLLDSG NQKNFLTWYQQKPGQPPKLLIFWASTRESGVPD RFTGSGSVTDFTLTISSVQAEDLAVYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 37	DNA LC	GACATTGTGATGACCCAGTCTCCATCCTCCCTG ACTGTGACAGCAGGAGAGAAGGTCACTATGAGC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCAGGGCAGCCTCCTAAACTGTTGATCTTC TGGGCATCCACTAGGGAATCTGGGGTCCCTGAT CGCTTCACAGGCAGTGGATCTGTAACAGATTTC ACTCTCACCATCAGCAGTGTGCAGGCTGAAGAC

		CTGGCAGTTTATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAACCTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACCTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum01 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDVAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDVAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVTPSSSLGKITY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLGLK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC

		AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCCTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAGGCCCTCCCGTCTCCATCGAG AAAACCATCTCCAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGTCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum01 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 42	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTEFTLTITISLQPDDEFATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 43	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCATCA AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTC ACTCTCACCATCAGCAGCCTGTCAGCCTGATGAT

		TTTGCAACTTATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 44	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTEFTLTISLQPDFFATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 45	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCATCA AGGTTTCAGCGGCAGTGGATCTGGGACAGAATTC ACTCTCACCATCAGCAGCCTGCAGCCTGATGAT TTTGCAACTTATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCAAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum02 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDYAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC

SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTKY TCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFIYPSDIAVEWESNGQPENNYKTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLGLK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGTCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum02 LC		

SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 46	VL	DIQMTQSPSSLSASVGDRVITITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGIPP RFSGSGYGTDFTLTINNIESEDAAYYFCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 47	DNA VL	GACATCCAGATGACCCAGTCTCCATCCTCCCTG TCTGCATCTGTAGGAGACAGAGTCACCATCACT TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGATCCCACCT CGATTCACTGGCAGCGGGTATGGAACAGATTTT ACCCTCACAATTAATAACATAGAATCTGAGGAT GCTGCATATTACTTCTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 48	LC	DIQMTQSPSSLSASVGDRVITITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGIPP RFSGSGYGTDFTLTINNIESEDAAYYFCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKSTYLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 49	DNA LC	GACATCCAGATGACCCAGTCTCCATCCTCCCTG TCTGCATCTGTAGGAGACAGAGTCACCATCACT TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGATCCCACCT CGATTCACTGGCAGCGGGTATGGAACAGATTTT ACCCTCACAATTAATAACATAGAATCTGAGGAT GCTGCATATTACTTCTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum03 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY

SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 50	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSS
SEQ ID NO: 51	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 52	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHLEAL HNHYTQKSLSLSLGK
SEQ ID NO: 53	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGGAAGTCAAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTGTTCCCTG AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC

		GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCCTCACCGTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum03 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 46	VL	DIQMTQSPSSLSASVGDRTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGIPP RFGSGYGTDFTLTINNIESEDAAYYFCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 47	DNA VL	GACATCCAGATGACCCAGTCTCCATCCTCCCTG TCTGCATCTGTAGGAGACAGAGTCACCATCACT TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGATCCCACCT CGATTCAAGTGGCAGCGGGTATGGAACAGATTTT ACCCTCACAATTAATAACATAGAATCTGAGGAT GCTGCATATTACTTCTGTGAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 48	LC	DIQMTQSPSSLSASVGDRTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGIPP RFGSGYGTDFTLTINNIESEDAAYYFCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 49	DNA LC	GACATCCAGATGACCCAGTCTCCATCCTCCCTG TCTGCATCTGTAGGAGACAGAGTCACCATCACT TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGATCCCACCT CGATTCAAGTGGCAGCGGGTATGGAACAGATTTT ACCCTCACAATTAATAACATAGAATCTGAGGAT GCTGCATATTACTTCTGTGAGAATGATTATAGT

		TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCAAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum04 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 50	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSS
SEQ ID NO: 51	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 52	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 53	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC

		ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGACACCTTCCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTCACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCCTCACCGTCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACCTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum04 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPY
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 54	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLPEDIATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 55	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCATCA AGGTTTCAGTGGAAAGTGGATCTGGGACAGATTTT ACTTTTACCATCAGCAGCCTGCAGCCTGAAGAT ATTGCAACATATTACTGTGAGAATGATTATAGT

		TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 56	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLPEDIATYYCQNDYS YPYTFGGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 57	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCATCA AGGTTTCAGTGGAAGTGGATCTGGGACAGATTTT ACTTTTCACCATCAGCAGCCTGCAGCCTGAAGAT ATTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCTGCTGAAT AACTTCTATCCCAGAGAGGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum05 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYFTFTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTTVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYFTFTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN

		RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVT VSWNSGALTS GVHTFPAVLQSSGLYSLSSVTV PSSSLGKTY TCNVDHKPSNTKVDKRVESKYGP PCPPAPEF LGGPSVFLFPPKPKDTLMISRT PEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTK PREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKV SNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSV MHEAL HNHYTQKSLSLGLK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTACTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCGAACC GGTTG ACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAAC TACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum05 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLD SGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES

SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 54	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLPEDATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 55	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTGTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCATCA AGGTTTCAGTGGAAGTGGATCTGGGACAGATTTT ACTTTTCACCATCAGCAGCCTGCAGCCTGAAGAT ATTGCAACATATTACTGTGAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 56	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLPEDATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFPYPAKGVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 57	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTGTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCATCA AGGTTTCAGTGGAAGTGGATCTGGGACAGATTTT ACTTTTCACCATCAGCAGCCTGCAGCCTGAAGAT ATTGCAACATATTACTGTGAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum06 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY

SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACACAGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTCCCA AAACCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTACAGCGTCTCACCCTGCTGACCAG

		GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTCAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum06 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 58	VL	DIVMTQTPLSLPVTGPGEPAISCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 59	DNA VL	GATATTGTGATGACCCAGACTCCACTCTCCCTG CCCGTCACCCCTGGAGAGCCGGCCTCCATCTCC TGCAAGTCCAGTCAGAGTCTGTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTCCGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 60	LC	DIVMTQTPLSLPVTGPGEPAISCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFPYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 61	DNA LC	GATATTGTGATGACCCAGACTCCACTCTCCCTG CCCGTCACCCCTGGAGAGCCGGCCTCCATCTCC TGCAAGTCCAGTCAGAGTCTGTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTCCGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA

		TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCCCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum07 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVTPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPPCPPPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLGLK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC

		ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCCTCACCGTCTGCAACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTCAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAATAACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum07 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 62	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLDSDG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 63	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA

SEQ ID NO: 64	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 65	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCC GCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCCCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum08 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 50	VH	EVQLVQSGAEVKKPGESLRISCKGSGYFTTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQTTTVTVSS
SEQ ID NO: 51	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCCGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 52	HC	EVQLVQSGAEVKKPGESLRISCKGSGYFTTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQTTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS

		GVHTFPAVLQSSGLYSLSSVVTVPSSSLGKTY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEAL HNHYTQKSLSLGLK
		GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAG GGCGTGACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAAGCGTCTCACCCTGCTGACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTACGCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
SEQ ID NO: 53	DNA HC	
BAP049-hum08 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF

SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 66	VL	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 67	DNA VL	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAGAAGTCTTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 68	LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFPYAPKRVQWQVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 69	DNA LC	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAGAAGTCTTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum09 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIIYPGTGGSNFDEKFKN

		RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TTVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLGLK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG

		GTGTCCAACAAAGGCCTCCCGTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTCAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum09 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 66	VL	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 67	DNA VL	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 68	LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNMFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 69	DNA LC	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT

		AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCCCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum10 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 50	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSS
SEQ ID NO: 51	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTCACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 52	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPPCPPPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 53	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTCACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG

		GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCCTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTACAGCTCCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCTTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGTCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum10 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 70	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLDSDG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 71	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTT ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA

SEQ ID NO: 72	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 73	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCCCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum11 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYFTTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQTTTVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCCGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYFTTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQTTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS

		GVHTFPAVLQSSGLYSLSSVVTVPSSSLGKTY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEAL HNHYTQKSLSLGLK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACACAGCGCCCTGACCAGC GGCGTGACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAAGCGTCTCACCCTGCTGACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTACGCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum11 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF

SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 70	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 71	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAGAAGTCTTGTACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTGAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 72	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 73	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAGAAGTCTTGTACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTGAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCAACCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum12 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIIYPGTGGSNFDEKFKN

		RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TTVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTKY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG

		GTGTCCAACAAAGGCCTCCCGTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTCAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum12 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 74	VL	DIQMTQSPSSLSASVGDRTITCKSSQSLLDSG NQKNFLTWYLQKPGQSPQLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 75	DNA VL	GACATCCAGATGACCCAGTCTCCATCCTCCCTG TCTGCATCTGTAGGAGACAGAGTCACCATCACT TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCTGCAG AAGCCAGGGCAGTCTCCACAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 76	LC	DIQMTQSPSSLSASVGDRTITCKSSQSLLDSG NQKNFLTWYLQKPGQSPQLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 77	DNA LC	GACATCCAGATGACCCAGTCTCCATCCTCCCTG TCTGCATCTGTAGGAGACAGAGTCACCATCACT TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCTGCAG AAGCCAGGGCAGTCTCCACAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAC TGCTCTGTTGTGTGCTGCTGAAT

		AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum13 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 39	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 40	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 41	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTACCACTTACTGG ATGCACTGGGTGCGACAGGCCACTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGAGTCACGATTACCGCGGACAAATCCACGAGC ACAGCCTACATGGAGCTGAGCAGCCTGAGATCT GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG

		GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTGTTCC AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCCTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAAGCGTCCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGTCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum13 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 78	VL	DVVMTQSPLSLPVTLGQPASISCKSSQSLDSDG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 79	DNA VL	GATGTTGTGATGACTCAGTCTCCACTCTCCCTG CCCGTCACCCCTTGGACAGCCGGCCTCCATCTCC TGCAAGTCCAGTCAGAGTCTGTAGACAGTGGA AATCAAAAGAACTTCTTAACCTGGTATCAGCAG AAACCAGGGAAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA

SEQ ID NO: 80	LC	DVVMTQSPSLPVLTLGQPASISCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 81	DNA LC	GATGTTGTGATGACTCAGTCTCCACTCTCCCTG CCCGTCACCCTTGGACAGCCGGCCTCCATCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTAACCTGGTATCAGCAG AAACCAGGGAAGCTCCTAAGCTCCTGATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTCCGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCC GCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACAAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum14 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 82	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRW TTGTGAYWGQTTTVTVSS
SEQ ID NO: 83	DNA VH	CAGGTT CAGCTGGTGCAGTCTGGAGCTGAGGTG AAGAAGCCTGGGGCCTCAGTGAAGGTCTCCTGC AAGGCTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTACTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 84	HC	QVQLVQSGAEVKKPGASVKVSCKASGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRW TTGTGAYWGQTTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS

		GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTKTY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDLMISRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEAL HNHYTQKSLSLGLK
		CAGGTTTCAGCTGGTGCAGTCTGGAGCTGAGGTG AAGAAGCCTGGGGCCTCAGTGAAGGTCTCCTGC AAGGCTTCTGGCTACACATTACCACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTACCATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTACTGGGGCCAGGGC ACCACCGTGACCGTGTCTCCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAG GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCCCCA AAACCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAAGCGTCTCACCCTGCTGACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAATAACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
SEQ ID NO: 85	DNA HC	
BAP049-hum14 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF

SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 70	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 71	DNA VL	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAGAAGTCTTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 72	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 73	DNA LC	GAAATTGTGTTGACACAGTCTCCAGCCACCCTG TCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAGAAGTCTTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum15 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 82	VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN

		RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSS
SEQ ID NO: 83	DNA VH	CAGGTT CAGCTGGTGCAGTCTGGAGCTGAGGTG AAGAAGCCTGGGGCCTCAGTGAAGGTCTCCTGC AAGGCTTCTGGCTACACATTCACTACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTCACTATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTACTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 84	HC	QVQLVQSGAEVKKPGASVKV SCKASGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTKY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVDS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSV MHEAL HNHYTQKSLSLSLGK
SEQ ID NO: 85	DNA HC	CAGGTT CAGCTGGTGCAGTCTGGAGCTGAGGTG AAGAAGCCTGGGGCCTCAGTGAAGGTCTCCTGC AAGGCTTCTGGCTACACATTCACTACTTACTGG ATGCACTGGATCAGGCAGTCCCCATCGAGAGGC CTTGAGTGGCTGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTCACTATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTACTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTTCCCCCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACGTGCGTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG

		GTGTCCAACAAAGGCCTCCCGTCCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTCAGCCTGACC TGCCTGGTCAAAGGCTTCTACCCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum15 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 66	VL	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 67	DNA VL	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA
SEQ ID NO: 68	LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 69	DNA LC	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT

		AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCCCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-hum16 HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 86	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQAPGQGLEWMGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TTVTVSS
SEQ ID NO: 87	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTACTACTGG ATGCACTGGGTGCGACAGGCCCTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTCACTATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCC
SEQ ID NO: 88	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQAPGQGLEWMGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVDVVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLGK
SEQ ID NO: 89	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTG AAAAAGCCCGGGGAGTCTCTGAGGATCTCCTGT AAGGGTTCTGGCTACACATTCACTACTACTGG ATGCACTGGGTGCGACAGGCCCTGGACAAGGG CTTGAGTGGATGGGTAATATTTATCCTGGTACT GGTGGTTCTAACTTCGATGAGAAGTTCAAGAAC AGATTCACTATCTCCAGAGACAATTCCAAGAAC ACGCTGTATCTTCAAATGAACAGCCTGAGAGCC GAGGACACGGCCGTGTATTACTGTACAAGATGG ACTACTGGGACGGGAGCTTATTGGGGCCAGGGC ACCACCGTGACCGTGTCTCTCCGCTTCCACCAAG

		GGCCCATCCGTCTTCCCCCTGGCGCCCTGCTCC AGGAGCACCTCCGAGAGCACAGCCGCCCTGGGC TGCCTGGTCAAGGACTACTTCCCCGAACCGGTG ACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGC GGCGTGCACACCTTCCCGGCTGTCTTACAGTCC TCAGGACTCTACTCCCTCAGCAGCGTGGTGACC GTGCCCTCCAGCAGCTTGGGCACGAAGACCTAC ACCTGCAACGTAGATCACAAGCCCAGCAACACC AAGGTGGACAAGAGAGTTGAGTCCAAATATGGT CCCCCATGCCCACCGTGCCAGCACCTGAGTTC CTGGGGGGACCATCAGTCTTCTGTTCCTGACCA AAACCCAAGGACACTCTCATGATCTCCCGGACC CCTGAGGTACAGTGCCTGGTGGTGGACGTGAGC CAGGAAGACCCCGAGGTCCAGTTCAACTGGTAC GTGGATGGCGTGGAGGTGCATAATGCCAAGACA AAGCCGCGGGAGGAGCAGTTCAACAGCACGTAC CGTGTGGTACAGCTCCTCACCCTCCTGCACCAG GACTGGCTGAACGGCAAGGAGTACAAGTGCAAG GTGTCCAACAAAGGCCTCCCGTCTCCATCGAG AAAACCATCTCCAAAGCCAAAGGGCAGCCCCGA GAGCCACAGGTGTACACCCTGCCCCCATCCCAG GAGGAGATGACCAAGAACCAGGTGAGCCTGACC TGCTTGGTCAAAGGCTTCTACCCAGCGACATC GCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACCTACAAGACCACGCCTCCCGTGTCTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAGGCTA ACCGTGGACAAGAGCAGGTGGCAGGAGGGGAAT GTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACACAGAAGAGCCTCTCCCTG TCTCTGGGTAAA
BAP049-hum16 LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 66	VL	EIVLTQSPDFQSVTPKEKVTITCKSSQSLDSDG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 67	DNA VL	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTGTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCCTCG AGGTTTCAGTGGCAGTGGATCTGGGACAGATTTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAA

SEQ ID NO: 68	LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 69	DNA LC	GAAATTGTGCTGACTCAGTCTCCAGACTTTCAG TCTGTGACTCCAAAGGAGAAAGTCACCATCACC TGCAAGTCCAGTCAGAGTCTGTTAGACAGTGGA AATCAAAAGAACTTCTTGACCTGGTACCAGCAG AAACCTGGCCAGGCTCCCAGGCTCCTCATCTAT TGGGCATCCACTAGGGAATCTGGGGTCCCCTCG AGGTTCAGTGGCAGTGGATCTGGGACAGATTTTC ACCTTTACCATCAGTAGCCTGGAAGCTGAAGAT GCTGCAACATATTACTGTCAGAATGATTATAGT TATCCGTACACGTTTCGGCCAAGGGACCAAGGTG GAAATCAAACGTACGGTGGCTGCACCATCTGTC TTCATCTTCCC GCCATCTGATGAGCAGTTGAAA TCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAAT AACTTCTATCCCAGAGAGGCCAAAGTACAGTGG AAGGTGGATAACGCCCTCCAATCGGGTAACTCC CAGGAGAGTGTACAGAGCAGGACAGCAAGGAC AGCACCTACAGCCTCAGCAGCACCTGACGCTG AGCAAAGCAGACTACGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT
BAP049-Clone-A HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQTTTVTVSS
SEQ ID NO: 90	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTG AAGAAGCCTGGCGAGTCCCTGCGGATCTCCTGC AAGGGCTCTGGCTACACCTTACCACCTACTGG ATGCACTGGGTGCGACAGGCTACCGGCCAGGGC CTGGAATGGATGGGCAACATCTATCCTGGCACC GGCGGCTCCAACCTTCGACGAGAAGTTCAAGAAC AGAGTGACCATCACCGCCGACAAGTCCACCTCC ACCGCCTACATGGAAGTGTCTCCCTGAGATCC GAGGACACCGCCGTGTACTACTGCACCCGGTGG ACAACCGGCACAGGCGCTTATTGGGGCCAGGGC ACCACAGTGACCGTGTCTCT
SEQ ID NO: 91	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQTTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS

		GVHTFPAVLQSSGLYSLSSVVTVPSSSLGTKTY TCNVDPKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEAL HNHYTQKSLSLGLG
SEQ ID NO: 92	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTG AAGAAGCCTGGCGAGTCCCTGCGGATCTCCTGC AAGGGCTCTGGCTACACCTTACCACCTACTGG ATGCACTGGGTGCGACAGGCTACCGGCCAGGGC CTGGAATGGATGGGCAACATCTATCCTGGCACC GGCGGCTCCAACCTTCGACGAGAAGTTCAAGAAC AGAGTGACCATCACCGCCGACAAGTCCACCTCC ACCGCCTACATGGAAGTGTCTCCTGAGATCC GAGGACACCGCCGTGTACTACTGCACCCGGTGG ACAACCGGCACAGGCGCTTATTGGGGCCAGGGC ACCACAGTGACCGTGTCTCTGCTTCTACCAAG GGGCCCAGCGTGTTCCTTGGCCCCCTGCTCC AGAAGCACCAGCGAGAGCACAGCCGCCCTGGGC TGCCTGGTGAAGGACTACTTCCCCGAGCCCGTG ACCGTGTCTTGAACAGCGGAGCCCTGACCAGC GGCGTGCACACCTTCCCCGCCGTGCTGCAGAGC AGCGGCCTGTACAGCCTGAGCAGCGTGGTGACC GTGCCCAGCAGCAGCCTGGGCACCAAGACCTAC ACCTGTAACGTGGACCACAAGCCCAGCAACACC AAGGTGGACAAGAGGGTGGAGAGCAAGTACGGC CCACCCTGCCCCCCTGCCAGCCCCCGAGTTC CTGGGCGGACCCAGCGTGTTCCTGTTCCCCC AAGCCCAAGGACACCCTGATGATCAGCAGAACC CCCGAGGTGACCTGTGTGGTGGTGGACGTGTCC CAGGAGGACCCGAGGTCCAGTTCAACTGGTAC GTGGACGGCGTGGAGGTGCACAACGCCAAGACC AAGCCCAGAGAGGAGCAGTTTAACAGCACCTAC CGGGTGGTGTCCGTGCTGACCGTGTGACCAG GACTGGCTGAACGGCAAAGAGTACAAGTGAAG GTCTCCAACAAGGGCCTGCCAAGCAGCATCGAA AAGACCATCAGCAAGGCCAAGGGCCAGCCTAGA GAGCCCCAGGTCTACACCCTGCCACCCAGCCAA GAGGAGATGACCAAGAACCAGGTGTCCCTGACC TGTCTGGTGAAGGGCTTACCCCAAGCGACATC GCCGTGGAGTGGGAGAGCAACGCCAGCCCCGAG AACAATAACAAGACCACCCCCCAGTGCTGGAC AGCGACGGCAGCTTCTTCTGTACAGCAGGCTG ACCGTGGACAAGTCCAGATGGCAGGAGGGCAAC GTCTTTAGCTGCTCCGTGATGCACGAGGCCCTG CACAACCACTACACCCAGAAGAGCCTGAGCCTG TCCCTGGGC
BAP049-Clone-A LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF

SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 42	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTEFTLTISLQPDDEFATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 93	DNA VL	GAGATCGTGCTGACCCAGTCCCCTGCCACCCTG TCACTGTCTCCAGGCGAGAGAGCTACCCTGTCC TGCAAGTCCTCCCAGTCCCTGCTGGACTCCGGC AACCAGAAGAACTTCCTGACCTGGTATCAGCAG AAGCCCGGCCAGGCCCCCAGACTGCTGATCTAC TGGGCCTCCACCCGGGAATCTGGCGTGCCCTCT AGATTCTCCGGCTCCGGCTCTGGCACCAGATTT ACCCTGACCATCTCCAGCCTGCAGCCCCGACGAC TTCGCCACCTACTACTGCCAGAACGACTACTCC TACCCCTACACCTTCGGCCAGGGCACCAAGGTG GAAATCAAG
SEQ ID NO: 44	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTEFTLTISLQPDDEFATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFPYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 94	DNA LC	GAGATCGTGCTGACCCAGTCCCCTGCCACCCTG TCACTGTCTCCAGGCGAGAGAGCTACCCTGTCC TGCAAGTCCTCCCAGTCCCTGCTGGACTCCGGC AACCAGAAGAACTTCCTGACCTGGTATCAGCAG AAGCCCGGCCAGGCCCCCAGACTGCTGATCTAC TGGGCCTCCACCCGGGAATCTGGCGTGCCCTCT AGATTCTCCGGCTCCGGCTCTGGCACCAGATTT ACCCTGACCATCTCCAGCCTGCAGCCCCGACGAC TTCGCCACCTACTACTGCCAGAACGACTACTCC TACCCCTACACCTTCGGCCAGGGCACCAAGGTG GAAATCAAGCGTACGGTGGCCGCTCCCAGCGTG TTCATCTTCCCCCAAGCGACGAGCAGCTGAAG AGCGGCACCGCCAGCGTGGTGTGTCTGCTGAAC AACTTCTACCCAGGGAGGCCAAGGTGCAGTGG AAGGTGGACAACGCCCTGCAGAGCGGCAACAGC CAGGAGAGCGTCACCGAGCAGGACAGCAAGGAC TCCACCTACAGCCTGAGCAGCACCCCTGACCCCTG AGCAAGGCCGACTACGAGAAGCACAAGGTGTAC GCCTGTGAGGTGACCCACCAGGGCCTGTCCAGC CCCGTGACCAAGAGCTTCAACAGGGGCGAGTGC
BAP049-Clone-B HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN

		RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TVTVSS
SEQ ID NO: 95	DNA VH	GAGGTGCAGCTGGTGCAGTCAGGCGCCGAAGTG AAGAAGCCCGGCGAGTCACTGAGAATTAGCTGT AAAGGTTTCAGGCTACACCTTCACTACCTACTGG ATGCACTGGGTCCGCCAGGCTACCGGTCAAGGC CTCGAGTGGATGGGTAATATCTACCCCGGCACC GGCGGCTCTAACTTCGACGAGAAGTTTAAGAAT AGAGTGACTATCACCGCCGATAAGTCTACTAGC ACCGCCTATATGGAAGTGTCTAGCCTGAGATCA GAGGACACCGCCGTCTACTACTGCACTAGGTGG ACTACCGGCACAGGCGCCTACTGGGGTCAAGGC ACTACCGTGACCGTGTCTAGC
SEQ ID NO: 91	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSED TAVYYCTRW TTGTGAYWGQGT TVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGTKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMHREAL HNHYTQKSLSLSLG
SEQ ID NO: 96	DNA HC	GAGGTGCAGCTGGTGCAGTCAGGCGCCGAAGTG AAGAAGCCCGGCGAGTCACTGAGAATTAGCTGT AAAGGTTTCAGGCTACACCTTCACTACCTACTGG ATGCACTGGGTCCGCCAGGCTACCGGTCAAGGC CTCGAGTGGATGGGTAATATCTACCCCGGCACC GGCGGCTCTAACTTCGACGAGAAGTTTAAGAAT AGAGTGACTATCACCGCCGATAAGTCTACTAGC ACCGCCTATATGGAAGTGTCTAGCCTGAGATCA GAGGACACCGCCGTCTACTACTGCACTAGGTGG ACTACCGGCACAGGCGCCTACTGGGGTCAAGGC ACTACCGTGACCGTGTCTAGCGCTAGCACTAAG GGCCCGTCCGTGTTCCCCCTGGCACCTTGTAGC CGGAGCACTAGCGAATCCACCGCTGCCCTCGGC TGCCTGGTCAAGGATTACTTCCCGGAGCCCGTG ACCGTGTCCTGGAACAGCGGAGCCCTGACCTCC GGAGTGACACCTTCCCCGCTGTGCTGCAGAGC TCCGGCTGTACTCGCTGTGCTCGGTGGTCACG GTGCCTTCATCTAGCCTGGGTACCAAGACCTAC ACTTGCAACGTGGACCACAAGCCTTCCAACACT AAGGTGGACAAGCGCGTCAATCGAAGTACGGC CCACCGTGCCCGCCTTGTCCCGCGCCGGAGTTC CTCGGCGGTCCCTCGGTCTTTCTGTTCCACCG AAGCCCAAGGACACTTTGATGATTTCCCGCACC CCTGAAGTGACATGCGTGGTGGTGGACGTGTCA CAGGAAGATCCGGAGGTGCAGTTCAATTGGTAC GTGGATGGCGTCGAGGTGCACAACGCCAAAACC AAGCCGAGGGAGGAGCAGTTCAACTCCACTTAC CGCGTCGTGTCCGTGCTGACGGTGTGTCATCAG GACTGGCTGAACGGGAAGGAGTACAAGTGCAAA GTGTCCAACAAGGGACTTCTAGCTCAATCGAA

		AAGACCATCTCGAAAGCCAAGGGACAGCCCCGG GAACCCCAAGTGTATACCCTGCCACCGAGCCAG GAAGAAATGACTAAGAACCAAGTCTCATTGACT TGCCTTGTGAAGGGCTTCTACCCATCGGATATC GCCGTGGAATGGGAGTCCAACGGCCAGCCGGAA ACAACACTACAAGACCACCCCTCCGGTGCTGGAC TCAGACGGATCCTTCTTCTCTACTCGCGGCTG ACCGTGGATAAGAGCAGATGGCAGGAGGAAAT GTGTTTCAGCTGTTCTGTGATGCATGAAGCCCTG CACAACCACTACACTCAGAAGTCCCTGTCCCTC TCCCTGGGA
BAP049-Clone-B LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 54	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLPEDIATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 97	DNA VL	GAGATCGTCCTGACTCAGTCACCCGCTACCCCTG AGCCTGAGCCCTGGCGAGCGGGCTACACTGAGC TGTAATCTAGTCAGTCACTGCTGGATAGCGGT AATCAGAAGAACTTCCTGACCTGGTATCAGCAG AAGCCCGGTAAAGCCCCTAAGCTGCTGATCTAC TGGGCCTCTACTAGAGAATCAGGCGTGCCCTCT AGGTTTAGCGGTAGCGGTAGTGGCACCGACTTC ACCTTCACTATCTCTAGCCTGCAGCCCGAGGAT ATCGCTACCTACTACTGTGAGAACGACTATAGC TACCCCTACACCTTCGGTCAAGGCACTAAGGTC GAGATTAAG
SEQ ID NO: 56	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGKAPKLLIYWASTRESGVPS RFGSGSGTDFTFTISSLPEDIATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFPYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSPVTKSFNRGEC
SEQ ID NO: 98	DNA LC	GAGATCGTCCTGACTCAGTCACCCGCTACCCCTG AGCCTGAGCCCTGGCGAGCGGGCTACACTGAGC TGTAATCTAGTCAGTCACTGCTGGATAGCGGT AATCAGAAGAACTTCCTGACCTGGTATCAGCAG AAGCCCGGTAAAGCCCCTAAGCTGCTGATCTAC TGGGCCTCTACTAGAGAATCAGGCGTGCCCTCT AGGTTTAGCGGTAGCGGTAGTGGCACCGACTTC ACCTTCACTATCTCTAGCCTGCAGCCCGAGGAT ATCGCTACCTACTACTGTGAGAACGACTATAGC TACCCCTACACCTTCGGTCAAGGCACTAAGGTC GAGATTAAGCGTACGGTGGCCGCTCCCAGCGTG TTCATCTTCCCCCCCAGCGACGAGCAGCTGAAG AGCGGCACCGCCAGCGTGGTGTGCCTGCTGAAC AACTTCTACCCCGGGAGGCCAAGGTGCAGTGG

		AAGGTGGACAACGCCCTGCAGAGCGGCAACAGC CAGGAGAGCGTCACCGAGCAGGACAGCAAGGAC TCCACCTACAGCCTGAGCAGCACCTGACCCTG AGCAAGGCCGACTACGAGAAGCATAAGGTGTAC GCCTGCGAGGTGACCCACCAGGGCCTGTCCAGC CCCGTGACCAAGAGCTTCAACAGGGGCGAGTGC
BAP049-Clone-C HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSS
SEQ ID NO: 90	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTG AAGAAGCCTGGCGAGTCCCTGCGGATCTCCTGC AAGGGCTCTGGCTACACCTTCACCACCTACTGG ATGCACTGGGTGCGACAGGCTACCGGCCAGGGC CTGGAATGGATGGGCAACATCTATCCTGGCACC GGCGGCTCCAACCTTCGACGAGAAGTTCAAGAAC AGAGTGACCATCACCGCCGACAAGTCCACCTCC ACCGCCTACATGGAAGTGTCTCCCTGAGATCC GAGGACACCGCCGTGTACTACTGCACCCGGTGG ACAACCGGCACAGGCGCTTATTGGGGCCAGGGC ACCACAGTGACCGTGTCTCT
SEQ ID NO: 91	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTITVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVTPSSSLGKTY TCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEF LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLG
SEQ ID NO: 92	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTG AAGAAGCCTGGCGAGTCCCTGCGGATCTCCTGC AAGGGCTCTGGCTACACCTTCACCACCTACTGG ATGCACTGGGTGCGACAGGCTACCGGCCAGGGC CTGGAATGGATGGGCAACATCTATCCTGGCACC GGCGGCTCCAACCTTCGACGAGAAGTTCAAGAAC AGAGTGACCATCACCGCCGACAAGTCCACCTCC ACCGCCTACATGGAAGTGTCTCCCTGAGATCC GAGGACACCGCCGTGTACTACTGCACCCGGTGG ACAACCGGCACAGGCGCTTATTGGGGCCAGGGC ACCACAGTGACCGTGTCTCTGCTTCTACCAAG GGGCCCAGCGTGTTCCTCCCTGGCCCCCTGCTCC

		AGAAGCACCAGCGAGAGCACAGCCGCCCTGGGC TGCCTGGTGAAGGACTACTTCCCCGAGCCCGTG ACCGTGTCCTGGAACAGCGGAGCCCTGACCAGC GGCGTGCACACCTTCCCCGCCGTGCTGCAGAGC AGCGGCCTGTACAGCCTGAGCAGCGTGGTGACC GTGCCCAGCAGCAGCCTGGGCACCAAGACCTAC ACCTGTAACGTGGACCACAAGCCCAGCAACACC AAGGTGGACAAGAGGGTGGAGAGCAAGTACGGC CCACCCTGCCCCCCTGCCAGCCCCCGAGTTC CTGGGCGGACCCAGCGTGTTCTGTTCCCCCC AAGCCCAAGGACACCCTGATGATCAGCAGAACC CCCGAGGTGACCTGTGTGGTGGTGGACGTGTCC CAGGAGGACCCCGAGGTCCAGTTCAACTGGTAC GTGGACGGCGTGGAGGTGCACAACGCCAAGACC AAGCCCAGAGAGGAGCAGTTTAACAGCACCTAC CGGGTGGTGTCCGTGCTGACCGTGCTGCACCAG GACTGGCTGAACGGCAAAGAGTACAAGTGTAA GTCTCCAACAAGGGCCTGCCAAGCAGCATCGAA AAGACCATCAGCAAGGCCAAGGGCCAGCCTAGA GAGCCCCAGGTCTACACCCTGCCACCCAGCCAA GAGGAGATGACCAAGAACCAGGTGTCCCTGACC TGTCTGGTGAAGGGCTTCTACCCAAGCGACATC GCCGTGGAGTGGGAGAGCAACGGCCAGCCCGAG AACAACTACAAGACCACCCCCCAGTGCTGGAC AGCGACGGCAGCTTCTTCTGTACAGCAGGCTG ACCGTGGACAAGTCCAGATGGCAGGAGGGCAAC GTCTTTAGCTGCTCCGTGATGCACGAGGCCCTG CACAACCACTACACCCAGAAGAGCCTGAGCCTG TCCCTGGGC
BAP049-Clone-C LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 66	VL	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 99	DNA VL	GAGATCGTGCTGACCCAGTCCCCCGACTTCCAG TCCGTGACCCCCAAAGAAAAAGTGACCATCACA TGCAAGTCCTCCAGTCCCTGCTGGACTCCGGC AACCAGAAGAACTTCTGACCTGGTATCAGCAG AAGCCCGGCCAGGCCCCAGACTGCTGATCTAC TGGGCCTCCACCCGGAATCTGGCGTGCCCTCT AGATTCTCCGGCTCCGGCTCTGGCACCGACTTT ACCTTCACCATCTCCAGCCTGGAAGCCGAGGAC GCCGCCACCTACTACTGCCAGAACGACTACTCC TACCCCTACACCTTCGGCCAGGGCACCAAGGTG GAAATCAAG
SEQ ID NO: 68	LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS

		YPYTFGQGTKEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSTYSLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 100	DNA LC	GAGATCGTGCTGACCCAGTCCCCCGACTTCCAG TCCGTGACCCCCAAAGAAAAAGTGACCATCACA TGCAAGTCCTCCAGTCCCTGCTGGACTCCGGC AACCAGAAGAACTTCTGACCTGGTATCAGCAG AAGCCCGGCCAGGCCCCCAGACTGCTGATCTAC TGGGCCTCCACCCGGAATCTGGCGTGCCCTCT AGATTCTCCGGCTCCGGCTCTGGCACCGACTTT ACCTTCACCATCTCCAGCCTGGAAGCCGAGGAC GCCGCCACCTACTACTGCCAGAACGACTACTCC TACCCCTACACCTTCGGCCAGGGCACCAGGTG GAAATCAAGCGTACGGTGGCCGCTCCCAGCGTG TTCATCTTCCCCCAAGCGACGAGCAGCTGAAG AGCGGCACCGCCAGCGTGGTGTGTCTGCTGAAC AACTTCTACCCAGGGAGGCCAAGGTGCAGTGG AAGGTGGACAACGCCCTGCAGAGCGGCAACAGC CAGGAGAGCGTCACCGAGCAGGACAGCAAGGAC TCCACCTACAGCCTGAGCAGCACCTGACCCTG AGCAAGGCCGACTACGAGAAGCACAAGGTGTAC GCCTGTGAGGTGACCCACCAGGGCCTGTCCAGC CCCGTGACCAAGAGCTTCAACAGGGCGAGTGC
BAP049-Clone-D HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 50	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TTVTVSS
SEQ ID NO: 101	DNA VH	GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTG AAGAAGCCTGGCGAGTCCCTGCGGATCTCCTGC AAGGGCTCTGGCTACACCTTACCACCTACTGG ATGCACTGGATCCGGCAGTCCCCCTCTAGGGGC CTGGAATGGCTGGGCAACATCTACCCTGGCACC GGCGGCTCCAACCTTCGACGAGAAGTTCAAGAAC AGGTTCACCATCTCCCGGGACAACCTCAAGAAC ACCCTGTACCTGCAGATGAACTCCCTGCGGGCC GAGGACACCGCCGTGTACTACTGTACCAGATGG ACCACCGGAACCGGCGCCTATTGGGGCCAGGGC ACAACAGTGACCGTGTCTCTCC
SEQ ID NO: 102	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWIRQSPSRGLEWLGNIYPGTGGSNFDEKFKN RFTISRDN SKNTLYLQMN SLRAEDTAVYYCTRW TTGTGAYWGQGT TTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVTPSSSLGTKTY TCNV D HKPSNTKVDKRVESKYGP P CP P P A P E F LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS

		QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLGLG
SEQ ID NO: 103	DNA HC	GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTG AAGAAGCCTGGCGAGTCCCTGCGGATCTCCTGC AAGGGCTCTGGCTACACCTTACCACCTACTGG ATGCACTGGATCCGGCAGTCCCCCTCTAGGGGC CTGGAATGGCTGGGCAACATCTACCCTGGCACC GGCGGCTCCAACCTTCGACGAGAAGTTCAAGAAC AGGTTCACCATCTCCCGGGACAACCTCCAAGAAC ACCCTGTACCTGCAGATGAACTCCCTGCGGGCC GAGGACACCGCCGTGTACTACTGTACCAGATGG ACCACCGGAACCGGCGCCTATTGGGGCCAGGGC ACAACAGTGACCGTGTCTCTCCGCTTCTACCAAG GGGCCAGCGTGTTCCTCCCTGGCCCCCTGCTCC AGAAGCACCAGCGAGAGCACAGCCGCCCTGGGC TGCCTGGTGAAGGACTACTTCCCCGAGCCCGTG ACCGTGTCTCTGGAACAGCGGAGCCCTGACCAGC GGCGTGCACACCTTCCCCGCCGTGCTGCAGAGC AGCGGCCTGTACAGCCTGAGCAGCGTGGTGACC GTGCCCAGCAGCAGCCTGGGCACCAAGACCTAC ACCTGTAACGTGGACCACAAGCCCAGCAACACC AAGGTGGACAAGAGGGTGGAGAGCAAGTACGGC CCACCCTGCCCCCCTGCCCAGCCCCCGAGTTC CTGGGCGGACCCAGCGTGTTCCTGTTCCCCCCC AAGCCCAAGGACACCCTGATGATCAGCAGAACC CCCGAGGTGACCTGTGTGGTGGTGGACGTGTCC CAGGAGGACCCCGAGGTCCAGTTCAACTGGTAC GTGGACGGCGTGGAGGTGCACAACGCCAAGACC AAGCCCAGAGAGGAGCAGTTTAACAGCACCTAC CGGGTGGTGTCCGTGCTGACCGTGTGACACCAG GACTGGCTGAACGGCAAAGAGTACAAGTGTAAAG GTCTCCAACAAGGGCCTGCCAAGCAGCATCGAA AAGACCATCAGCAAGGCCAAGGGCCAGCCTAGA GAGCCCCAGGTCTACACCCTGCCACCCAGCCAA GAGGAGATGACCAAGAACCAGGTGTCCCTGACC TGTCTGGTGAAGGGCTTCTACCCAAGCGACATC GCCGTGGAGTGGGAGAGCAACGGCCAGCCCGAG AACAACATAAGACCACCCCCCAGTGCTGGAC AGCGACGGCAGCTTCTTCTGTACAGCAGGCTG ACCGTGGACAAGTCCAGATGGCAGGAGGGCAAC GTCTTTAGCTGCTCCGTGATGCACGAGGCCCTG CACAACCACTACACCCAGAAGAGCCTGAGCCTG TCCCTGGGC
BAP049-Clone-D LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLDSDGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPY
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLDSDGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY

SEQ ID NO: 70	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 104	DNA VL	GAGATCGTGCTGACCCAGTCCCCTGCCACCCTG TCACTGTCTCCAGGCGAGAGAGCTACCCTGTCC TGCAAGTCCTCCAGTCCCTGCTGGACTCCGGC AACCAGAAGAACTTCTGACCTGGTATCAGCAG AAGCCCGGCCAGGCCCCCAGACTGCTGATCTAC TGGGCCTCCACCCGGAATCTGGCGTGCCCTCT AGATTCTCCGGCTCCGGCTCTGGCACCGACTTT ACCTTCACCATCTCCAGCCTGGAAGCCGAGGAC GCCGCCACCTACTACTGCCAGAACGACTACTCC TACCCCTACACCTTCGGCCAGGGCACCAAGGTG GAAATCAAG
SEQ ID NO: 72	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFSGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKSTYSLSSLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 105	DNA LC	GAGATCGTGCTGACCCAGTCCCCTGCCACCCTG TCACTGTCTCCAGGCGAGAGAGCTACCCTGTCC TGCAAGTCCTCCAGTCCCTGCTGGACTCCGGC AACCAGAAGAACTTCTGACCTGGTATCAGCAG AAGCCCGGCCAGGCCCCCAGACTGCTGATCTAC TGGGCCTCCACCCGGAATCTGGCGTGCCCTCT AGATTCTCCGGCTCCGGCTCTGGCACCGACTTT ACCTTCACCATCTCCAGCCTGGAAGCCGAGGAC GCCGCCACCTACTACTGCCAGAACGACTACTCC TACCCCTACACCTTCGGCCAGGGCACCAAGGTG GAAATCAAGCGTACGGTGGCCGCTCCCAGCGTG TTCATCTTCCCCCAAGCGACGAGCAGCTGAAG AGCGGCACCGCCAGCGTGGTGTGTCTGCTGAAC AACTTCTACCCAGGGAGGCCAAGGTGCAGTGG AAGGTGGACAACGCCCTGCAGAGCGGCAACAGC CAGGAGAGCGTCACCGAGCAGGACAGCAAGGAC TCCACCTACAGCCTGAGCAGCACCCCTGACCCTG AGCAAGGCCGACTACGAGAAGCACAAGGTGTAC GCCTGTGAGGTGACCCACCAGGGCCTGTCCAGC CCCGTGACCAAGAGCTTCAACAGGGGCGAGTGC
BAP049-Clone-E HC		
SEQ ID NO: 1 (Kabat)	HCDR1	TYWMH
SEQ ID NO: 2 (Kabat)	HCDR2	NIYPGTGGSNFDEKFKN
SEQ ID NO: 3 (Kabat)	HCDR3	WTTGTGAY
SEQ ID NO: 4 (Chothia)	HCDR1	GYTFTTY
SEQ ID NO: 5 (Chothia)	HCDR2	YPGTGG
SEQ ID NO: 3 (Chothia)	HCDR3	WTTGTGAY
SEQ ID NO: 38	VH	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQGTTVTVSS
SEQ ID NO: 95	DNA VH	GAGGTGCAGCTGGTGCAGTCAGGCGCCGAAGTG

		AAGAAGCCCGGCGAGTCACTGAGAATTAGCTGT AAAGGTTTCAGGCTACACCTTCACTACCTACTGG ATGCACTGGGTCCGCCAGGCTACCGGTCAAGGC CTCGAGTGGATGGGTAATATCTACCCCGGCACC GGCGGCTCTAACTTCGACGAGAAGTTTAAGAAT AGAGTGACTATCACCGCCGATAAGTCTACTAGC ACCGCCTATATGGAAGTGTCTAGCCTGAGATCA GAGGACACCGCCGTCTACTACTGCACTAGGTGG ACTACCGGCACAGGCGCCTACTGGGGTCAAGGC ACTACCGTGACCGTGTCTAGC
SEQ ID NO: 91	HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYW MHWVRQATGQGLEWMGNIYPGTGGSNFDEKFKN RVTITADKSTSTAYMELSSLRSEDTAVYYCTRW TTGTGAYWGQTTVTVSSASTKGPSVFPLAPCS RSTSESTAALGCLVKDYFPEPVTVSWNSGALTS GVHTFPAVLQSSGLYSLSSVTVPSSSLGTKTY TCNVDHKPSNTKVDKRVESKYGPPCPPCPAPEF LGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS QEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIE KTISKAKGQPREPQVYTLPPSQEEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSRLTVDKSRWQEGNVFSCSVMEAL HNHYTQKSLSLSLG
SEQ ID NO: 96	DNA HC	GAGGTGCAGCTGGTGCAGTCAGGCGCCGAAGTG AAGAAGCCCGGCGAGTCACTGAGAATTAGCTGT AAAGGTTTCAGGCTACACCTTCACTACCTACTGG ATGCACTGGGTCCGCCAGGCTACCGGTCAAGGC CTCGAGTGGATGGGTAATATCTACCCCGGCACC GGCGGCTCTAACTTCGACGAGAAGTTTAAGAAT AGAGTGACTATCACCGCCGATAAGTCTACTAGC ACCGCCTATATGGAAGTGTCTAGCCTGAGATCA GAGGACACCGCCGTCTACTACTGCACTAGGTGG ACTACCGGCACAGGCGCCTACTGGGGTCAAGGC ACTACCGTGACCGTGTCTAGCGCTAGCACTAAG GGCCCGTCCGTGTTCCCCCTGGCACCTTGTAGC CGGAGCACTAGCGAATCCACCGCTGCCCTCGGC TGCCTGGTCAAGGATTACTTCCCGGAGCCCGTG ACCGTGTCCTGGAACAGCGGAGCCCTGACCTCC GGAGTGACACCTTCCCCGCTGTGCTGCAGAGC TCCGGGCTGTACTCGCTGTCTGCTGGTGCAG GTGCCTTCATCTAGCCTGGGTACCAAGACCTAC ACTTGCAACGTGGACCACAAGCCTTCCAACACT AAGGTGGACAAGCGCGTCGAATCGAAGTACGGC CCACCGTGCCCGCCTTGTCCCGCGCCGGAGTTC CTCGGCGGTCCCTCGGTCTTTCTGTTCCCAACG AAGCCCAAGGACACTTTGATGATTTCCCGCACC CCTGAAGTGACATGCGTGGTGGTGGACGTGTCA CAGGAAGATCCGGAGGTGCAGTTCAATTGGTAC GTGGATGGCGTTCGAGGTGCACAACGCCAAAACC AAGCCGAGGGAGGAGCAGTTCAACTCCACTTAC CGCGTCTGTCCGTGCTGACGGTGTGCTGCATCAG GACTGGCTGAACGGGAAGGAGTACAAGTGCAAA GTGTCCAACAAGGGACTTCCTAGCTCAATCGAA AAGACCATCTCGAAAGCCAAGGGACAGCCCCGG GAACCCCAAGTGTATACCCTGCCACCGAGCCAG GAAGAAATGACTAAGAACCAAGTCTCATTGACT

		TGCCTTGTGAAGGGCTTCTACCCATCGGATATC GCCGTGGAATGGGAGTCCAACGGCCAGCCGGAA AACAACTACAAGACCACCCCTCCGGTGCTGGAC TCAGACGGATCCTTCTTCTCTACTCGCGGCTG ACCGTGGATAAGAGCAGATGGCAGGAGGGAAAT GTGTTTCAGCTGTTCTGTGATGCATGAAGCCCTG CACAACCACTACACTCAGAAGTCCCTGTCCCTC TCCCTGGGA
BAP049-Clone-E LC		
SEQ ID NO: 10 (Kabat)	LCDR1	KSSQSLLDSGNQKNFLT
SEQ ID NO: 11 (Kabat)	LCDR2	WASTRES
SEQ ID NO: 32 (Kabat)	LCDR3	QNDYSYPYT
SEQ ID NO: 13 (Chothia)	LCDR1	SQSLLDSGNQKNF
SEQ ID NO: 14 (Chothia)	LCDR2	WAS
SEQ ID NO: 33 (Chothia)	LCDR3	DYSYPY
SEQ ID NO: 70	VL	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIK
SEQ ID NO: 106	DNA VL	GAGATCGTCCTGACTCAGTCACCCGCTACCCCTG AGCCTGAGCCCTGGCGAGCGGGCTACACTGAGC TGTAATCTAGTCAGTCACTGCTGGATAGCGGT AATCAGAAGAACTTCCTGACCTGGTATCAGCAG AAGCCCGGTCAAGCCCCTAGACTGCTGATCTAC TGGGCCTCTACTAGAGAATCAGGCGTGCCCTCT AGGTTTAGCGGTAGCGGTAGTGGCACCGACTTC ACCTTCACTATCTCTAGCCTGGAAGCCGAGGAC GCCGCTACCTACTACTGTGAGAACGACTATAGC TACCCCTACACCTTCGGTCAAGGCACTAAGGTC GAGATTAAG
SEQ ID NO: 72	LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSG NQKNFLTWYQQKPGQAPRLLIYWASTRESGVPS RFGSGSGTDFTFTISSLEAEDAATYYCQNDYS YPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLK SGTASVCLLNNFYFPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTLSSTLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC
SEQ ID NO: 107	DNA LC	GAGATCGTCCTGACTCAGTCACCCGCTACCCCTG AGCCTGAGCCCTGGCGAGCGGGCTACACTGAGC TGTAATCTAGTCAGTCACTGCTGGATAGCGGT AATCAGAAGAACTTCCTGACCTGGTATCAGCAG AAGCCCGGTCAAGCCCCTAGACTGCTGATCTAC TGGGCCTCTACTAGAGAATCAGGCGTGCCCTCT AGGTTTAGCGGTAGCGGTAGTGGCACCGACTTC ACCTTCACTATCTCTAGCCTGGAAGCCGAGGAC GCCGCTACCTACTACTGTGAGAACGACTATAGC TACCCCTACACCTTCGGTCAAGGCACTAAGGTC GAGATTAAGCGTACGGTGGCCGCTCCCAGCGTG TTCATCTTCCCCCCCAGCGACGAGCAGCTGAAG AGCGGCACCGCCAGCGTGGTGTGCCTGCTGAAC AACTTCTACCCCCGGGAGGCCAAGGTGCACTGG AAGGTGGACAACGCCCTGCAGAGCGGCAACAGC CAGGAGAGCGTCACCGAGCAGGACAGCAAGGAC TCCACCTACAGCCTGAGCAGCACCTGACCCTG

		AGCAAGGCCGACTACGAGAAGCATAAGGTGTAC GCCTGCGAGGTGACCCACCAGGGCCTGTCCAGC CCCGTGACCAAGAGCTTCAACAGGGGCGAGTGC
BAP049 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 115 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTGCACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 118 (Chothia)	LCDR3	GATTATAGTTATCCGTGC
BAP049-chi HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-chi LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 115 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTGCACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 118 (Chothia)	LCDR3	GATTATAGTTATCCGTGC
BAP049-chi Y HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT

SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-chi Y LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum01 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
		AATATTTATCCTGGTACTGGTGGTTCTAACTTC
SEQ ID NO: 109 (Kabat)	HCDR2	GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum01 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum02 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
		AATATTTATCCTGGTACTGGTGGTTCTAACTTC
SEQ ID NO: 109 (Kabat)	HCDR2	GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum02 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum03 HC		

SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum03 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum04 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum04 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum05 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum05 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT

SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum06 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum06 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum07 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum07 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum08 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC

SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum08 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum09 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum09 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum10 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum10 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC

BAP049-hum11 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum11 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum12 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum12 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum13 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum13 LC		
SEQ ID NO: 121 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTAACC

SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum14 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 223 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAC
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 223 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAC
BAP049-hum14 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum15 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 223 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAC
SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 223 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAC
BAP049-hum15 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAAGAACTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG
SEQ ID NO: 116 (Chothia)	LCDR1	AGTCAGAGTCTGTTAGACAGTGGAAATCAAAAG AACTTC
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-hum16 HC		
SEQ ID NO: 108 (Kabat)	HCDR1	ACTTACTGGATGCAC
SEQ ID NO: 109 (Kabat)	HCDR2	AATATTTATCCTGGTACTGGTGGTTCTAACTTC GATGAGAAGTTCAAGAAC
SEQ ID NO: 110 (Kabat)	HCDR3	TGGACTACTGGGACGGGAGCTTAT

SEQ ID NO: 111 (Chothia)	HCDR1	GGCTACACATTACCACTTAC
SEQ ID NO: 112 (Chothia)	HCDR2	TATCCTGGTACTGGTGGT
SEQ ID NO: 110 (Chothia)	HCDR3	TGGACTACTGGGACGGGAGCTTAT
BAP049-hum16 LC		
SEQ ID NO: 113 (Kabat)	LCDR1	AAGTCCAGTCAGAGTCTGTTAGACAGTGGAAAT CAAAGAAGTTCTTGACC
SEQ ID NO: 114 (Kabat)	LCDR2	TGGGCATCCACTAGGGAATCT
SEQ ID NO: 119 (Kabat)	LCDR3	CAGAATGATTATAGTTATCCGTACACG AGTCAGAGTCTGTTAGACAGTGGAAATCAAAG AACTTC
SEQ ID NO: 116 (Chothia)	LCDR1	
SEQ ID NO: 117 (Chothia)	LCDR2	TGGGCATCC
SEQ ID NO: 120 (Chothia)	LCDR3	GATTATAGTTATCCGTAC
BAP049-Clone-A HC		
SEQ ID NO: 122 (Kabat)	HCDR1	ACCTACTGGATGCAC
		AACATCTATCCTGGCACC GGCGGCTCCAAGTTTC
SEQ ID NO: 123 (Kabat)	HCDR2	GACGAGAAGTTCAAGAAC
SEQ ID NO: 124 (Kabat)	HCDR3	TGGACAACCGGCACAGGCGCTTAT
SEQ ID NO: 125 (Chothia)	HCDR1	GGCTACACCTTCACCACCTAC
SEQ ID NO: 126 (Chothia)	HCDR2	TATCCTGGCACC GGCGGCG
SEQ ID NO: 124 (Chothia)	HCDR3	TGGACAACCGGCACAGGCGCTTAT
BAP049-Clone-A LC		
		AAGTCCTCCAGTCCCTGCTGGACTCCGGCAAC
SEQ ID NO: 127 (Kabat)	LCDR1	CAGAAGAAGTTCTTGACC
SEQ ID NO: 128 (Kabat)	LCDR2	TGGGCCTCCACCCGGGAATCT
SEQ ID NO: 129 (Kabat)	LCDR3	CAGAACGACTACTCCTACCCCTACACC TCCAGTCCCTGCTGGACTCCGGCAACCAGAAG AACTTC
SEQ ID NO: 130 (Chothia)	LCDR1	
SEQ ID NO: 131 (Chothia)	LCDR2	TGGGCCTCC
SEQ ID NO: 132 (Chothia)	LCDR3	GACTACTCCTACCCCTAC
BAP049-Clone-B HC		
SEQ ID NO: 133 (Kabat)	HCDR1	ACCTACTGGATGCAC
		AATATCTACCCGGCACC GGCGGCTCTAAGTTTC
SEQ ID NO: 134 (Kabat)	HCDR2	GACGAGAAGTTTAAGAAT
SEQ ID NO: 135 (Kabat)	HCDR3	TGGACTACCGGCACAGGCGCCTAC
SEQ ID NO: 136 (Chothia)	HCDR1	GGCTACACCTTCACTACCTAC
SEQ ID NO: 137 (Chothia)	HCDR2	TACCCCGGCACC GGCGGCG
SEQ ID NO: 135 (Chothia)	HCDR3	TGGACTACCGGCACAGGCGCCTAC
BAP049-Clone-B LC		
		AAATCTAGTCAGTCACTGCTGGATAGCGGTAAT
SEQ ID NO: 138 (Kabat)	LCDR1	CAGAAGAAGTTCTTGACC
SEQ ID NO: 139 (Kabat)	LCDR2	TGGGCCTCTACTAGAGAATCA
SEQ ID NO: 140 (Kabat)	LCDR3	CAGAACGACTATAGCTACCCCTACACC AGTCAGTCACTGCTGGATAGCGGTAATCAGAAG AACTTC
SEQ ID NO: 141 (Chothia)	LCDR1	
SEQ ID NO: 142 (Chothia)	LCDR2	TGGGCCTCT

SEQ ID NO: 143 (Chothia)	LCDR3	GACTATAGCTACCCCTAC
BAP049-Clone-C HC		
SEQ ID NO: 122 (Kabat)	HCDR1	ACCTACTGGATGCAC
SEQ ID NO: 123 (Kabat)	HCDR2	AACATCTATCCTGGCACC GGCGGCTCCAACTTC GACGAGAAGTTCAAGAAC
SEQ ID NO: 124 (Kabat)	HCDR3	TGGACAACCGGCACAGGCGCTTAT
SEQ ID NO: 125 (Chothia)	HCDR1	GGCTACACCTTCACCACCTAC
SEQ ID NO: 126 (Chothia)	HCDR2	TATCCTGGCACC GGCGGC
SEQ ID NO: 124 (Chothia)	HCDR3	TGGACAACCGGCACAGGCGCTTAT
BAP049-Clone-C LC		
SEQ ID NO: 127 (Kabat)	LCDR1	AAGTCCTCCCAGTCCCTGCTGGACTCCGGCAAC CAGAAGAACTTCCTGACC
SEQ ID NO: 128 (Kabat)	LCDR2	TGGGCCTCCACCCGGAATCT
SEQ ID NO: 129 (Kabat)	LCDR3	CAGAACGACTACTCCTACCCCTACACC TCCCAGTCCCTGCTGGACTCCGGCAACCAGAAG AACTTC
SEQ ID NO: 130 (Chothia)	LCDR1	TGGGCCTCC
SEQ ID NO: 131 (Chothia)	LCDR2	TGGGCCTCC
SEQ ID NO: 132 (Chothia)	LCDR3	GACTACTCCTACCCCTAC
BAP049-Clone-D HC		
SEQ ID NO: 122 (Kabat)	HCDR1	ACCTACTGGATGCAC
SEQ ID NO: 144 (Kabat)	HCDR2	AACATCTACCCTGGCACC GGCGGCTCCAACTTC GACGAGAAGTTCAAGAAC
SEQ ID NO: 145 (Kabat)	HCDR3	TGGACCACCGGAACCGGCGCCTAT
SEQ ID NO: 125 (Chothia)	HCDR1	GGCTACACCTTCACCACCTAC
SEQ ID NO: 146 (Chothia)	HCDR2	TACCCTGGCACC GGCGGC
SEQ ID NO: 145 (Chothia)	HCDR3	TGGACCACCGGAACCGGCGCCTAT
BAP049-Clone-D LC		
SEQ ID NO: 127 (Kabat)	LCDR1	AAGTCCTCCCAGTCCCTGCTGGACTCCGGCAAC CAGAAGAACTTCCTGACC
SEQ ID NO: 128 (Kabat)	LCDR2	TGGGCCTCCACCCGGAATCT
SEQ ID NO: 129 (Kabat)	LCDR3	CAGAACGACTACTCCTACCCCTACACC TCCCAGTCCCTGCTGGACTCCGGCAACCAGAAG AACTTC
SEQ ID NO: 130 (Chothia)	LCDR1	TGGGCCTCC
SEQ ID NO: 131 (Chothia)	LCDR2	TGGGCCTCC
SEQ ID NO: 132 (Chothia)	LCDR3	GACTACTCCTACCCCTAC
BAP049-Clone-E HC		
SEQ ID NO: 133 (Kabat)	HCDR1	ACCTACTGGATGCAC
SEQ ID NO: 134 (Kabat)	HCDR2	AATATCTACCCGGCACC GGCGGCTCTAACTTC GACGAGAAGTTTAAGAAT
SEQ ID NO: 135 (Kabat)	HCDR3	TGGACTACCGGCACAGGCGCCTAC
SEQ ID NO: 136 (Chothia)	HCDR1	GGCTACACCTTCACTACCTAC
SEQ ID NO: 137 (Chothia)	HCDR2	TACCCCGGCACC GGCGGC
SEQ ID NO: 135 (Chothia)	HCDR3	TGGACTACCGGCACAGGCGCCTAC
BAP049-Clone-E LC		

SEQ ID NO: 138 (Kabat)	LCDR1	AAATCTAGTCAGTCACTGCTGGATAGCGGTAAT CAGAAGAACTTCCTGACC
SEQ ID NO: 139 (Kabat)	LCDR2	TGGGCCTCTACTAGAGAATCA
SEQ ID NO: 140 (Kabat)	LCDR3	CAGAACGACTATAGCTACCCCTACACC
SEQ ID NO: 141 (Chothia)	LCDR1	AGTCAGTCACTGCTGGATAGCGGTAATCAGAAG AACTTC
SEQ ID NO: 142 (Chothia)	LCDR2	TGGGCCTCT
SEQ ID NO: 143 (Chothia)	LCDR3	GACTATAGCTACCCCTAC

Table 2. Amino acid and nucleotide sequences of the heavy and light chain framework regions for humanized mAbs BAP049-hum01 to BAP049-hum16 and BAP049-Clone-A to BAP049-Clone-E

	Amino Acid Sequence	Nucleotide Sequence
VHFW1 (type a)	EVQLVQSGAEVKKPGESLRISCKGS (SEQ ID NO: 147)	GAAGTGCAGCTGGTGCAGTCTGGAGCAGAGGTGAAAAA GCCCCGGGGAGTCTCTGAGGATCTCCTGTAAGGGTTCT (SEQ ID NO: 148) GAAGTGCAGCTGGTGCAGTCTGGCGCCGAAGTGAAGAA GCCTGGCGAGTCCCTGCGGATCTCCTGCAAGGGCTCT (SEQ ID NO: 149) GAGGTGCAGCTGGTGCAGTCAGGCGCCGAAGTGAAGAA GCCCCGGCGAGTCACTGAGAATTAGCTGTAAAGGTTCA (SEQ ID NO: 150)
VHFW1 (type b)	QVQLVQSGAEVKKPGASVKVSKAS (SEQ ID NO: 151)	CAGGTTCACTGGTGCAGTCTGGAGCTGAGGTGAAGAA GCCTGGGGCCTCAGTGAAGGTCTCCTGCAAGGCTTCT (SEQ ID NO: 152)
VHFW2 (type a)	WVRQATGQGLEWMG (SEQ ID NO: 153)	TGGGTGCGACAGGCCACTGGACAAGGGCTTGAGTGGAT GGGT (SEQ ID NO: 154) TGGGTGCGACAGGCTACCGGCCAGGGCCTGGAATGGAT GGGC (SEQ ID NO: 155) TGGGTCCGCCAGGCTACCGGTCAAGGCCTCGAGTGGAT GGGT (SEQ ID NO: 156)
VHFW2 (type b)	WIRQSPSRGLEWLG (SEQ ID NO: 157)	TGGATCAGGCAGTCCCCATCGAGAGGCCTTGAGTGGCT GGGT (SEQ ID NO: 158) TGGATCCGGCAGTCCCCCTCTAGGGGCCTGGAATGGCT GGGC (SEQ ID NO: 159)
VHFW2 (type c)	WVRQAPGQGLEWMG (SEQ ID NO: 160)	TGGGTGCGACAGGCCCTGGACAAGGGCTTGAGTGGAT GGGT (SEQ ID NO: 161)
VHFW3 (type a)	RVTITADKSTSTAYMELSSLRSEDVAVY YCTR (SEQ ID NO: 162)	AGAGTCACGATTACCGCGGACAAATCCACGAGCACAGC CTACATGGAGCTGAGCAGCCTGAGATCTGAGGACACGG CCGTGTATTACTGTACAAGA (SEQ ID NO: 163) AGAGTGACCATCACCGCCGACAAGTCCACCTCCACCGC CTACATGGAAGTGTCTCCTGAGATCCGAGGACACCG

		CCGTGTACTACTGCACCCGG (SEQ ID NO: 164) AGAGTGACTATCACCGCCGATAAGTCTACTAGCACCCGCTATATGGAAGTGTCTAGCCTGAGATCAGAGGACACCGCCGTCTACTACTGCACTAGG (SEQ ID NO: 165)
VHFW3 (type b)	RFTISRDN SKNTLYLQMN SLRAEDTAVY YCTR (SEQ ID NO: 166)	AGATTACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTTCAAATGAACAGCCTGAGAGCCGAGGACACGCCGTGTATTACTGTACAAGA (SEQ ID NO: 167) AGGTTACCATCTCCCGGGACAACCTCCAAGAACACCCGTGTACCTGCAGATGAACTCCCTGCGGGCCGAGGACACCGCCGTGTACTACTGTACCAGA (SEQ ID NO: 168)
VHFW4	WGQGT TVTVSS (SEQ ID NO: 169)	TGGGGCCAGGGCACCACCGTGACCGTGTCTCTCC (SEQ ID NO: 170) TGGGGCCAGGGCACCACAGTGACCGTGTCTCTCT (SEQ ID NO: 171) TGGGGTCAAGGCACTACCGTGACCGTGTCTAGC (SEQ ID NO: 172) TGGGGCCAGGGCACAACAGTGACCGTGTCTCTCC (SEQ ID NO: 173)
VLFW1 (type a)	EIVLTQSPDFQSVTPKEKVTITC (SEQ ID NO: 174)	GAAATTGTGCTGACTCAGTCTCCAGACTTTTCACTCTGTGACTCCAAAGGAGAAAGTCACCATCACCTGC (SEQ ID NO: 175) GAGATCGTGCTGACCCAGTCCCCCGACTTCCAGTCCGTGACCCCCAAAGAAAAAGTGACCATCACATGC (SEQ ID NO: 176)
VLFW1 (type b)	EIVLTQSPATLSLSPGERATLSC (SEQ ID NO: 177)	GAAATTGTGTTGACACAGTCTCCAGCCACCCTGTCTTTGTCTCCAGGGGAAAGAGCCACCCTCTCTCTGC (SEQ ID NO: 178) GAGATCGTGCTGACCCAGTCCCCCTGCCACCCTGTCACTGTCTCCAGGCGAGAGAGCTACCCTGTCTCTGC (SEQ ID NO: 179) GAGATCGTCTCTGACTCAGTCACCCGCTACCCTGAGCCTGAGCCCTGGCGAGCGGGCTACACTGAGCTGT (SEQ ID NO: 180)
VLFW1 (type c)	DIVMTQTPLSLPVTGPGEPAISIC (SEQ ID NO: 181)	GATATTGTGATGACCCAGACTCCACTCTCCCTGCCCGTCACCCCTGGAGAGCCGGCCTCCATCTCTCTGC (SEQ ID NO: 182)
VLFW1 (type d)	DVVMTQSPLSLPVTLGQPASISIC (SEQ ID NO: 183)	GATGTTGTGATGACTCAGTCTCCACTCTCCCTGCCCGTCACCCCTGGACAGCCGGCCTCCATCTCTCTGC (SEQ ID NO: 184)
VLFW1 (type e)	DIQMTQSPSSLSASVGRVTITC (SEQ ID NO: 185)	GACATCCAGATGACCCAGTCTCCATCCTCCCTGTCTGCTATCTGTAGGAGACAGAGTCACCATCACTTGC (SEQ ID NO: 186)

VLFW2 (type a)	WYQQKPGQAPRLLIY (SEQ ID NO: 187)	TGGTACCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCT CATCTAT (SEQ ID NO: 188) TGGTATCAGCAGAAGCCCGGCCAGGCCCCCAGACTGCT GATCTAC (SEQ ID NO: 189) TGGTATCAGCAGAAGCCCGGTCAAGCCCCTAGACTGCT GATCTAC (SEQ ID NO: 190)
VLFW2 (type b)	WYQQKPGKAPKLLIY (SEQ ID NO: 191)	TGGTATCAGCAGAAACCAGGGAAAGCTCCTAAGCTCCT GATCTAT (SEQ ID NO: 192) TGGTATCAGCAGAAGCCCGGTAAAGCCCCTAAGCTGCT GATCTAC (SEQ ID NO: 193)
VLFW2 (type c)	WYLQKPGQSPQLLIY (SEQ ID NO: 194)	TGGTACCTGCAGAAGCCAGGGCAGTCTCCACAGCTCCT GATCTAT (SEQ ID NO: 195)
VLFW3 (type a)	GVPSRFSGSGSGTDFTFTISSLEAEDAA TYYC (SEQ ID NO: 196)	GGGGTCCCCTCGAGGTTTCAGTGGCAGTGGATCTGGGAC AGATTTTACCTTTACCATCAGTAGCCTGGAAGCTGAAG ATGCTGCAACATATTACTGT (SEQ ID NO: 197) GGCGTGCCCTCTAGATTCTCCGGCTCCGGCTCTGGCAC CGACTTTACCTTCACCATCTCCAGCCTGGAAGCCGAGG ACGCCGCCACCTACTACTGC (SEQ ID NO: 198) GGCGTGCCCTCTAGGTTTTCAGCGGTAGCGGTAGTGGCAC CGACTTCACCTTCACCTATCTCTAGCCTGGAAGCCGAGG ACGCCGCTACCTACTACTGT (SEQ ID NO: 199)
VLFW3 (type b)	GIPPRFSGSGYGTDFLTINNIESEDAA YYFC (SEQ ID NO: 200)	GGGATCCCACCTCGATTTCAGTGGCAGCGGGTATGGAAC AGATTTTACCCTCACAATTAATAACATAGAATCTGAGG ATGCTGCATATTACTTCTGT (SEQ ID NO: 201)
VLFW3 (type c)	GVPSRFSGSGSGTEFTLTISLQPDFA TYYC (SEQ ID NO: 202)	GGGGTCCCATCAAGGTTTCAGCGGCAGTGGATCTGGGAC AGAATTCACCTCTCACCATCAGCAGCCTGCAGCCTGATG ATTTTGCAACTTATTACTGT (SEQ ID NO: 203) GGCGTGCCCTCTAGATTCTCCGGCTCCGGCTCTGGCAC CGAGTTTACCCTGACCATCTCCAGCCTGCAGCCCGACG ACTTCGCCACCTACTACTGC (SEQ ID NO: 204)
VLFW3 (type d)	GVPSRFSGSGSGTDFTFTISSLQPDIA TYYC (SEQ ID NO: 205)	GGGGTCCCATCAAGGTTTCAGTGGAAAGTGGATCTGGGAC AGATTTTACTTTTACCATCAGCAGCCTGCAGCCTGAAG ATATTGCAACATATTACTGT (SEQ ID NO: 206) GGCGTGCCCTCTAGGTTTTCAGCGGTAGCGGTAGTGGCAC CGACTTCACCTTCACCTATCTCTAGCCTGCAGCCCGAGG ATATCGCTACCTACTACTGT (SEQ ID NO: 207)
VLFW4	FGQGTKVEIK (SEQ ID NO: 208)	TTCGGCCAAGGGACCAAGGTGGAAATCAAA (SEQ ID NO: 209) TTCGGCCAGGGCACCAAGGTGGAAATCAAG (SEQ ID NO: 210) TTCGGTCAAGGCACTAAGGTCGAGATTAAG (SEQ ID NO: 211)

Table 3. Constant region amino acid sequences of human IgG heavy chains and human kappa light chain

HC	IgG4 (S228P) mutant constant region amino acid sequence (EU Numbering) ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTKT YTCNVDPKPS NTKVDRVES KYGPPCPPCP APEFLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG NVFSCSVME ALHNHYTQKS LSLSLGK (SEQ ID NO: 212)
LC	Human kappa constant region amino acid sequence RTVAAPSVFI FPPSDEQLKS GTASVCLLN NFYPREAKVQ WKVDNALQSG NSQESVTEQD SKDSTYSLSS TLTLSKADYE KHKVYACEVT HQGLSSPVTK SFNRGEC (SEQ ID NO: 213)
HC	IgG4 (S228P) mutant constant region amino acid sequence lacking C-terminal lysine (K) (EU Numbering) ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTKT YTCNVDPKPS NTKVDRVES KYGPPCPPCP APEFLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG NVFSCSVME ALHNHYTQKS LSLSLG (SEQ ID NO: 214)
HC	IgG1 wild type ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDRVEP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTPEVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTCLVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK (SEQ ID NO: 215)
HC	IgG1 (N297A) mutant constant region amino acid sequence (EU Numbering) ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDRVEP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTPEVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYA STYRVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTCLVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK (SEQ ID NO: 216)
HC	IgG1 (D265A, P329A) mutant constant region amino acid sequence (EU Numbering) ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDRVEP KSCDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMISRTPEVTCVVVAVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVSVLT VLHQDWLNGK EYKCKVSNKA LAAPIEKTIS KAKGQPREPQ VYTLPPSREE

	MTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK (SEQ ID NO: 217)
HC	IgG1 (L234A, L235A) mutant constant region amino acid sequence (EU Numbering) ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKRVEP KSCDKHTTCP PCPAPEAAGG PSVFLFPPKP KDTLMISRTPEVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSREE MTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV MHEALHNHYT QKSLSLSPGK (SEQ ID NO: 218)

Table 4. Amino acid sequences of the heavy and light chain leader sequences for humanized mAbs BAP049-Clone-A to BAP049-Clone-E

BAP049-Clone-A	HC	MEWSWVFLFFLSVTTGVHS (SEQ ID NO: 219)
	LC	MSVPTQVLGLLLLLWLT DARC (SEQ ID NO: 220)
BAP049-Clone-B	HC	MAVWVTL PFLMAAAQSVQA (SEQ ID NO: 221)
	LC	MSVLTQVLALLLLLWLTGTRC (SEQ ID NO: 222)
BAP049-Clone-C	HC	MEWSWVFLFFLSVTTGVHS (SEQ ID NO: 219)
	LC	MSVPTQVLGLLLLLWLT DARC (SEQ ID NO: 220)
BAP049-Clone-D	HC	MEWSWVFLFFLSVTTGVHS (SEQ ID NO: 219)
	LC	MSVPTQVLGLLLLLWLT DARC (SEQ ID NO: 220)
BAP049-Clone-E	HC	MAVWVTL PFLMAAAQSVQA (SEQ ID NO: 221)
	LC	MSVLTQVLALLLLLWLTGTRC (SEQ ID NO: 222)

EXAMPLES

The Examples below are set forth to aid in the understanding of the inventions but are not intended to, and should not be construed to, limit its scope in any way.

Example 1: Pharmacokinetics Analysis of Flat Dosing Schedules

Based on pharmacokinetic (PK) modeling, utilizing flat dose is expected provide the exposure to patients at the appropriate C_{min} concentrations. Over 99.5% of patients will be above EC₅₀ and over 93% of patients will be above EC₉₀. Predicted steady state mean C_{min} for the exemplary anti-PD-1 antibody molecule utilizing either 300mg once every three weeks (Q3W) or 400 mg once every four weeks (Q4W) is expected to be above 20ug/mL (with highest weight, 150 kg) on average.

Table 5. Exemplary PK parameters based on flat dosing schedules

Number of patients in PK dataset	46
CL (mL/h)	10.9 [8.9, 13.2]; IIV: 62%
Exponent of Weight on CL	0.54 [0.021, 1.06]
Volume of distribution at SS (L)	7.2 [6.5, 7.9]; IIV: 22%
Half-Life (days)	20 [17, 23]; IIV: 64%
Predicted C_{min} (ug/mL) for 80 kg patient	31 [22, 42] (400mg q4w) 35 [26, 47] (300mg q3w)

The expected mean steady state C_{min} concentrations for the exemplary anti-PD-1 antibody molecule observed with either doses/regimens (300 mg q3w or 400 mg q4w) will be at least 77 fold higher than the EC₅₀ (0.42ug/mL) and about 8.6 fold higher than the EC₉₀. The *ex vivo* potency is based on IL-2 change in SEB *ex-vivo* assay.

Less than 10% of patients are expected to achieve C_{min} concentrations below 3.6 µg/mL for either 300 mg Q3W or 400 mg Q4W. Less than 0.5% of patients are expected to achieve C_{min} concentrations below 0.4 µg/mL for either 300 mg Q3W or 400 mg Q4W.

Predicted C_{trough} (C_{min}) concentrations across the different weights for patients while receiving the same dose of the exemplary anti-PD-1 antibody molecule are shown in Figure 12. Body weight based dosing is compared to fixed dose (3.75 mg/kg Q3W vs. 300 mg Q3W and 5 mg/kg Q4W vs. 400 mg Q4W). Figure 12 supports flat dosing of the exemplary anti-PD-1 antibody molecule.

The PK model further is validated. As shown in Figure 13, the observed versus model predicted concentrations lie on the line of unity. Figure 14 shows that the model captures accumulation, time course, and within subject variability.

Example 2: A phase Ib/II, open-label, multi-center study of Capmatinib in combination with an anti-PD-1 antibody molecule (“Antibody Molecule A”, detailed below) or Antibody Molecule A single agent in advanced hepatocellular carcinoma

Antibody Molecule A:

Antibody Molecule A is a high-affinity fully humanized anti-human-PD-1 monoclonal antibody that belongs to the IgG4/κ isotype subclass. It is expressed in a Chinese hamster ovary cell line (CHO-C8TD) and consists of two heavy chains and two light chains. Both heavy chains of Antibody Molecule A contain oligosaccharide chains linked to the protein backbone at Asn294.

The amino acid sequences of the light chain (220 amino acids) and the heavy chain (443 amino acids) respectively, as deduced from the DNA sequence, are shown in Figure 15 and Figure 16. The expected disulfide linkages derived from primary sequence are listed in Table 6.

Table 6. Expected disulfide linkages

Intra-light chain	Cys ₂₃ -Cys ₉₄ , Cys ₁₄₀ -Cys ₂₀₀ (x2)
Intra-heavy chain	Cys ₂₂ -Cys ₉₆ , Cys ₁₄₄ -Cys ₂₀₀ , Cys ₂₅₈ -Cys ₃₁₈ , Cys ₃₆₄ -Cys ₄₂₂ (x2)
Inter-chain	light chain Cys ₂₂₀ - heavy chain Cys ₁₃₁ (x2)

	heavy chain Cys ₂₂₃ - heavy chain Cys ₂₂₃ heavy chain Cys ₂₂₆ - heavy chain Cys ₂₂₆
--	--

The theoretical average molecular mass of Antibody Molecule A based on the amino acid composition as deduced from DNA sequence is 145759 Da. This mass takes into account the expected peptide

- 5 bonds and formation of disulfide bonds. Other post-translational modifications are disregarded, e.g. N-terminal glutamine is not considered as pyroglutamate, asparagine glycosylation sites are considered unchanged (Asn instead of Asp) and no other chemical modification is considered in the calculation.

Antibody Molecule A in the form of lyophilisate in vial for i.v. infusion. The starting
10 dose is 300mg, and it is administered every 3 weeks (Q3W).

Antibody Molecule A will be administered via i.v. infusion over 30 minutes (up to 2 hours, if clinically indicated) once every 3 weeks. The next scheduled dose may be delayed by up to 7 days to recover from previous AEs. If the next dose cannot be administered within the above mentioned 7-days delay, then the assessments should be shifted accordingly.

15

Capmatinib

Capmatinib tablet will be administered orally on a continuous twice daily (BID) dosing schedule, on a flat scale of mg/day and not individually adjusted by weight or body surface area.

Except on days of PK sampling, patients should take Capmatinib tablets twice daily (BID)
20 at approximately the same time each day starting at Cycle 1 Day 1.

Each dose of Capmatinib is to be taken with a glass of water (at least 8 ounces – approximately 250 mL) and consumed over as short a time as possible (i.e., not slower than 1 tablet every 2 minutes).

The dose of Capmatinib is 200mg BID.

25 Patients should be instructed to swallow the tablets whole and not to chew them.

- Capmatinib should be administered in the fasted state, at least 1 hour before or 2 hours after a meal. The morning and the evening doses should be taken 12 ($\pm$ 4) hours apart, although 12-hour interval is highly recommended. If a dose is not taken within 4 hours of the planned dosing time, the missed dose should not be replaced.

On days when PK blood samples are to be collected, patients will be instructed to hold their dose until arrival at the study center. Capmatinib will be administered at the site in the morning. The exact time of drug administration should be recorded in the appropriate eCRF. The PK blood draws will be supervised by a member of the research team. If a patient vomits
5 within 4 hours of Capmatinib dosing, the time of vomiting should be recorded on the eCRF.

Patients should be instructed not to make up for missed doses or partial doses (i.e., when the entire dose is not taken as instructed). A missed or partial dose will be defined as a case when the full dose is not taken within 4 hours of the scheduled twice daily dosing. If that occurs, then the dose (or part remaining dose) should not be taken and dosing should restart with
10 the next scheduled dose. If vomiting occurs, no attempt should be made to replace the vomited dose before the next scheduled dose.

During the whole duration of treatment with Capmatinib, the patient is recommended to use precautionary measures against ultraviolet exposure (e.g., use of sunscreen, protective clothing, avoid sunbathing or using a solarium).

15 *Objective of the trial:*

Main objective:

1. Phase Ib part:

To characterize the safety and tolerability of Capmatinib in combination with Antibody
20 Molecule A and identify the MTD and/or RP2D

2. Phase II part:

(a) For cMET low HCC patients, to compare the efficacy of Capmatinib in combination with Antibody Molecule A vs. Antibody Molecule A single agent.

25 (b) For cMET high HCC patients, to estimate the efficacy of Capmatinib in combination with Antibody Molecule A and Antibody Molecule A single agent, respectively.

Secondary objectives:

1. Phase Ib/II part:

To characterize the efficacy of Capmatinib in combination with Antibody Molecule A and Antibody Molecule A single agent in cMET high and low HCC.

2. Phase II part:

To characterize the safety and tolerability of Capmatinib in combination with Antibody Molecule A and Antibody Molecule A single agent.

3. Phase II part:

To investigate the association between Antibody Molecule A single agent efficacy and the cMET status in HCC patients.

4. Phase Ib/II parts:

To characterize the pharmacokinetic profile of Capmatinib and Antibody Molecule A.

5. Phase Ib/II parts:

To assess the pharmacodynamic effect of Capmatinib in combination with Antibody Molecule A and Antibody Molecule A single agent in tumor biopsy and peripheral blood in cMET high and low HCC.

6. Phase II part:

To compare the efficacy of Capmatinib in combination with Antibody Molecule A vs. Capmatinib single agent in cMET high HCC patients by using the historical data from a Capmatinib single agent study in 1st line Asian HCC patients.

PRINCIPAL INCLUSION CRITERIA

1. Histologically documented locally advanced recurrent or metastatic HCC. Current cirrhotic status of Child Pugh Class A (5-6 points), with no encephalopathy and/or ascites. Child Pugh status must be calculated based on clinical and laboratory results during the screening period.

2. Baseline tumor tissue (newly obtained) must be available at screening. Patient must have a site of disease amenable to biopsy, and be a candidate for tumor biopsy according to the treating institution's guidelines and requirements for such procedure.

3. Phase II: Documented evidence of cMet amplification (gene copy number) by FISH and cMet expression by IHC by a designated laboratory.

The cMET status will be classified as high or low according to the cMET expression assessed by IHC and the gene amplification assessed by FISH:

cMET high: if any one of the following criteria is satisfied

- 5 • IHC = 3+ in at least 50% of tumor cells and any gene (regardless of gene copy number (GCN))
- IHC = 2+ in at least 50% of tumor cells and $GCN \geq 5$

cMET low: if any one of the following criteria is satisfied

- 10 • IHC = 2+ in at least 50% of tumor cells and $GCN < 5$
 - IHC = 2+ in less than 50% of tumor cells and any GCN
 - IHC = 0 or 1+ (regardless of and any GCN)
4. Patients must be willing to undergo a new tumor biopsy during the study (6-9 weeks after start of study treatment, if medically feasible).
 - 15 For patients in the phase II part of the study, exceptions may be granted after documented discussion with Novartis. After a sufficient number of paired biopsies are collected, the decision may be taken to stop collecting the biopsies.
 5. Patients must have received prior systemic sorafenib treatment for HCC with documented progression during or after discontinuation of sorafenib treatment (for France only: patients must have received at least 8 weeks of prior sorafenib treatment), are intolerant to sorafenib (defined as documented Grade 3 or 4 adverse events that led to sorafenib discontinuation) or refused sorafenib treatment.
 - 20
 6. Patients must be tested during screening for Hepatitis-B-Virus surface antigen (HbsAg) status. Patients are included in the study if they have adequately controlled hepatitis B, defined by:
 - 25 • receiving a nucleoside analog anti-viral drug for 3 or more months, and
 - serum hepatitis B virus (HBV) deoxyribonucleic acid (DNA) level of less than 100 IU/ml via polymerase chain reaction quantification assays prior to enrollment.

7. Patients must be tested during study screening for hepatitis C virus ribonucleic acid (HCV RNA) status, patients are included in the study if they have adequately controlled hepatitis C; defined by having undetectable level of serum HCV RNA level prior to enrollment.

5 PRINCIPAL EXCLUSION CRITERIA

1. Patient has received the following therapies prior to the first dose of study treatment:

- Previous systemic anti-cancer therapy (including therapeutic cancer vaccines and immunotherapeutics) other than sorafenib (sorafenib must be completed within > 1 week prior to the first dose of study treatment) or Capmatinib.

10 • Previous locoregional therapy (e.g. hepatic arterial embolization, radio-frequency ablation, radiation therapy) if:

- administered after sorafenib treatment with the exception of palliative radiotherapy to a limited field, such as for the treatment of bone pain. Loco regional therapy for the focally painful liver tumor mass will be discussed on a case by case with trial sponsor.

15 - completed within 4 weeks prior to the dosing and, if present any related acute toxicity > grade 1.

- Use of any vaccines (except inactivated seasonal influenza vaccines) within 4 weeks of initiation of study treatment.

20 • Major surgery within 2 weeks of the first dose of study treatment (mediastinoscopy, insertion of a central venous access device, and insertion of a feeding tube are not considered major surgery).

- Participation in an interventional, investigational study within 2 weeks of the first dose of study treatment, unless agreed otherwise with trial sponsor.

25 • Presence of CTCAE grade ≥ 1 toxicity (except alopecia, peripheral neuropathy and ototoxicity, which are excluded if CTCAE grade ≥ 3) due to prior cancer therapy, unless agreed otherwise with trial sponsor.

30 • Use of hematopoietic colony-stimulating growth factors (e.g. G-CSF, GM-CSF, M-CSF) ≤ 2 weeks prior start or study drug. An erythroid stimulating agent is allowed as long as it was initiated at least 2 weeks prior to the first dose of study treatment and the patient is on a stable dose.

2. History of severe hypersensitivity reactions to other mAbs.

3. Known history of testing positive for human immunodeficiency virus (HIV) or known acquired immunodeficiency syndrome (testing is not required).

4. Clinically significant pleural effusion that either required pleurocentesis or is associated with shortness of breath.

5. Patients receiving treatment with medications that are either strong inducers or inhibitors of CYP3A, or CYP3A or CYP1A2 substrates with narrow therapeutic index, and cannot be discontinued at least 1 week prior to the start of treatment with Capmatinib and for the duration of the study.

6. Unable to stop herbal/food supplements or treatments which are considered to be capable of significantly causing either PK or PD herb/food-drug interactions.

7. Active autoimmune disease or a documented history of autoimmune disease, including ulcerative colitis and Crohn's disease or any condition that requires systemic steroids or any immunosuppressive therapy, except vitiligo or resolved asthma/atopy that is treated with bronchodilators (e.g., albuterol).

8. Clinically significant, uncontrolled heart diseases.

- Unstable angina within 6 months prior to screening
- Myocardial infarction within 6 months prior to screening
- History of documented congestive heart failure (New York Heart Association functional

classification III-IV)

• Uncontrolled hypertension defined by a Systolic Blood Pressure (SBP) ≥ 160 mm Hg and/or Diastolic Blood Pressure (DBP) ≥ 100 mm Hg, with or without antihypertensive medication. Initiation or adjustment of antihypertensive medication(s) is allowed prior to screening

- Ventricular arrhythmias
- Supraventricular and nodal arrhythmias not controlled with medication
- Other cardiac arrhythmia not controlled with medication
- QTcF ≥ 450 ms (male patients), ≥ 460 ms (female patients) on the screening ECG (as mean of triplicate ECG)

END POINTS:

Primary End Point

Phase Ib part:

5 Safety: Incidence and severity of AEs and SAEs, including changes in laboratory values, vital signs and ECGs. Incidence of DLT during the first 2 cycles of treatment.

Tolerability: Dose interruptions, reductions, and dose intensity

Phase II part:

10 Overall response rate (ORR) per Response Evaluation Criteria in Solid Tumors (RECIST v1.1)

Secondary End Point:

- 15 1. Best overall response (BOR), duration of overall response (DOR), time to response (TTR), progression-free survival (PFS), time to progression (TTP), overall survival (OS), Overall response rate (ORR)
2. Safety: Incidence and severity of adverse events (AEs) and serious adverse events, including changes in laboratory parameters, vital signs and electrocardiograms (ECGs). Tolerability: Dose interruptions, reductions and dose intensity
- 20 3. Best overall response (BOR), time to progression (TTP), cMET IHC score and GCN
4. Plasma/serum PK parameters (e.g., AUC, Cmax, Tmax) Plasma/serum concentration vs. time profiles.
5. H&E TIL & TIL characterization (CD8, CD3, CD4), TReg (FoxP3), PDL1, p-cMet on tumor biopsy (IHC)
- 25 6. Best overall response (BOR) and TTP

Preliminary Clinical Finding

As of mid December 2016, 4 patients have been enrolled and treated on this study and 1 tumor shrinkage that meets criteria for a confirmed partial response has been observed. The clinical trial is still on-going.

30

INCORPORATION BY REFERENCE

Other embodiments and examples including figures and tables are disclosed in International Patent Application Publication No. WO 2015/112900 and U.S. Patent Application
5 Publication No. US 2015/0210769, entitled “Antibody Molecules to PD-1 and Uses Thereof,”
which are incorporated by reference in its entirety.

All publications, patents, and Accession numbers mentioned herein are hereby
incorporated by reference in their entirety as if each individual publication or patent was
10 specifically and individually indicated to be incorporated by reference.

EQUIVALENTS

While specific embodiments of the subject invention have been discussed, the above
specification is illustrative and not restrictive. Many variations of the invention will become
15 apparent to those skilled in the art upon review of this specification and the claims below. The
full scope of the invention should be determined by reference to the claims, along with their full
scope of equivalents, and the specification, along with such variations.

CLAIMS

What is claimed is:

1. A method of treating a cancer in a subject, comprising

(A) administering to the subject a c-Met receptor tyrosine kinase inhibitor which is 2-fluoro-N-methyl-4-[7-quinolin-6-yl-methyl)-imidazo[1,2-b][1,2,4]triazin-2yl]benzamide or pharmaceutically acceptable salt thereof; and

(B) administering to the subject an anti-PD-1 antibody molecule at a dose of about 300 mg to 400 mg once every three weeks or once every four weeks,

wherein the anti- PD-1 antibody molecule comprises:

10 (a) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a light chain variable region (VL) comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33;

15 (b) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32;

20 (c) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33; or

25 (d) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

30

2. The method of claim 1 wherein the c-Met receptor tyrosine kinase inhibitor and the anti-PD-1 antibody molecule are administered separately, simultaneously or sequentially.

3. The method of claim 1 or 2, wherein the anti-PD-1 antibody molecule is
5 administered at a dose of about 300 mg once every three weeks.

4. The method of any of claims 1-2, wherein the anti-PD-1 antibody molecule is administered at a dose of about 400 mg once every four weeks.

10 5. The method of any of claims 1-3, wherein the c-Met receptor tyrosine kinase inhibitor is administered at 200 mg twice daily on a continuous schedule.

6. The method of any of claims 1-5, wherein the anti-PD-1 antibody molecule is administered intravenously.

15 7. The method of any of claims 1-6, wherein the c-Met receptor tyrosine kinase inhibitor is administered orally.

8. The method of any of claims 1-7, wherein the anti-PD-1 antibody molecule
20 comprises:

(a) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 42;

25 (b) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66;

(c) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70;

(d) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70;

5 (e) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46;

(f) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46;

10 (g) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54;

(h) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54;

(i) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 58;

20 (j) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 62;

(k) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66;

25 (l) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 74;

(m) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 78;

30

(n) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70;

(o) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66; or

(p) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 86 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

9. The method of any of claims 1-8, wherein the cancer is chosen from a lung cancer, a squamous cell lung cancer, a melanoma, a renal cancer, a liver cancer, a myeloma, a prostate cancer, a breast cancer, an ER+ breast cancer, an IM-TN breast cancer, a colorectal cancer, a colorectal cancer with high microsatellite instability, an EBV+ gastric cancer, a pancreatic cancer, a thyroid cancer, a hematological cancer, a non-Hodgkin's lymphoma, or a leukemia, or a metastatic lesion of the cancer.

10. The method of any of claims 1-9, wherein the cancer is chosen from a non-small cell lung cancer (NSCLC), a NSCLC adenocarcinoma, a NSCLC squamous cell carcinoma, a hepatocellular carcinoma.

11. A pharmaceutical combination comprising
(A) a c-Met receptor tyrosine kinase inhibitor which is 2-fluoro-N-methyl-4-[7-quinolin-6-yl-methyl)-imidazo[1,2-b][1,2,4]triazin-2-yl]benzamide or pharmaceutically acceptable salt thereof; and

(B) an anti-PD-1 antibody molecule for use in a dose of about 300 mg to 400 mg once every three weeks or once every four weeks,

wherein the anti- PD-1 antibody molecule comprises:

(a) a heavy chain variable region (VH) comprising a VHCDR1 amino acid sequence of SEQ ID NO: 4, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a light chain variable region (VL) comprising a

VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33;

(b) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 1; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32;

(c) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224, a VHCDR2 amino acid sequence of SEQ ID NO: 5, and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 13, a VLCDR2 amino acid sequence of SEQ ID NO: 14, and a VLCDR3 amino acid sequence of SEQ ID NO: 33; or

(d) a VH comprising a VHCDR1 amino acid sequence of SEQ ID NO: 224; a VHCDR2 amino acid sequence of SEQ ID NO: 2; and a VHCDR3 amino acid sequence of SEQ ID NO: 3; and a VL comprising a VLCDR1 amino acid sequence of SEQ ID NO: 10, a VLCDR2 amino acid sequence of SEQ ID NO: 11, and a VLCDR3 amino acid sequence of SEQ ID NO: 32.

12. The pharmaceutical combination of claim 11 wherein the c-Met receptor tyrosine kinase inhibitor is in oral dosage form.

13. The pharmaceutical combination of claim 11 wherein the anti-PD-1 antibody molecule is injectable dosage form.

14. The pharmaceutical combination of any of claims 11-13 wherein the anti-PD-1 antibody molecule is for use in a dose of about 300 mg to 400 mg once every three weeks or once every four weeks.

15. The pharmaceutical combination of any of claims 11-14 wherein c-Met receptor tyrosine kinase inhibitor is for use in a oral dosage form for continuously dosing at 200mg BID.

16. The pharmaceutical combination of any of claim 11-15, wherein the dose of the anti-PD-1 antibody molecule is about 300 mg once every three weeks.

5

17. The pharmaceutical combination of any of claim 11-15, wherein the dose is about 400 mg once every four weeks.

18. The pharmaceutical composition or dose formulation of any of claims 11-17,
10 wherein the anti-PD-1 antibody molecule comprises:

(a) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 42;

(b) a heavy chain variable domain comprising the amino acid sequence of SEQ ID
15 NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66;

(c) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70;

(d) a heavy chain variable domain comprising the amino acid sequence of SEQ ID
20 NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70;

(e) a heavy chain variable domain comprising the amino acid sequence of SEQ ID
25 NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46;

(f) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 46;

(g) a heavy chain variable domain comprising the amino acid sequence of SEQ ID
30 NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54;

(h) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 54;

(i) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 58;

(j) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 62;

(k) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 50 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66;

(l) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 74;

(m) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 38 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 78;

(n) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 70;

(o) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 82 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66; or

(p) a heavy chain variable domain comprising the amino acid sequence of SEQ ID NO: 86 and a light chain variable domain comprising the amino acid sequence of SEQ ID NO: 66.

19. The pharmaceutical combination of any of claims 11-18, for use in the treatment of a cancer.

20. The pharmaceutical combination of claim 19, wherein the cancer is chosen from a lung cancer, a squamous cell lung cancer, a melanoma, a renal cancer, a liver cancer, a myeloma, a prostate cancer, a breast cancer, an ER+ breast cancer, an IM-TN breast cancer, a colorectal cancer, a colorectal cancer with high microsatellite instability, an EBV+ gastric cancer, a
5 pancreatic cancer, a thyroid cancer, a hematological cancer, a non-Hodgkin's lymphoma, or a leukemia, or a metastatic lesion of the cancer.

21. The pharmaceutical combination of claim 19, wherein the cancer is chosen from a non-small cell lung cancer (NSCLC), a NSCLC adenocarcinoma, a NSCLC squamous cell
10 carcinoma, a hepatocellular carcinoma.

Heavy Chain (murine IgG1)

FWH1	CDRH1	FWH2	CDRH2
QVQLQQSGSE LVRPGASVKL SCKASGYTFT	TYMMHWVRQR	PQGLEWIGN	<u>IYPGTGGSNF</u> DEKFNRTSL
QVQLQQSGSE LVRPGASVKL SCKASGYTFT	TYMMHWVRQR	PQGLEWIGN	<u>IYPGTGGSNF</u> DEKFNRTSL
FWH3	CDRH3	FWH4	
TVDTSSSTAY MHLASLTSED SAVYYCTRMT	TGTGAYWGQG	TLVTVSA	
TVDTSSSTAY MHLASLTSED SAVYYCTRMT	TGTGAYWGQG	TLVTVSA	AKT TTPSVYPLAP GSAA

Light Chain (murine K)

FWL1	CDRL1	FWL2	CDRL2
DIVMTQSPSS LTVTAGEKVT MSCKSSQSLL	DSGNQKNFLT	WYQKPGQPP	KLLIFWASTR
DIVMTQSPSS LTVTAGEKVT MSCKSSQSLL	DSGNQKNFLT	WYQKPGQPP	KLLIFWASTR
FWL3	CDRL3	FWL4	
SGSVTDFTLT ISSVQAEDLA VYYCONDYSY	PCIFGGGTKL	EIK	
SGSVTDFTLT ISSVQAEDLA VYYCONDYSY	PCIFGGGTKL	EIKRAD	

FIGURE 1

<u>Heavy Chain</u>	
GL	QVQLQQPGSE LVRPGASVKL SCKASGYTFT <u>SYWMHWVKQR</u> HQGGLGWIGN <u>IYPGSGSTNY</u>
Mu mAb	-----S-----T-----R--P-----T- <u>GS-F</u>
GL	DEKFKSKGTL TVDTSSSTAY MHLSSLTSED SAVYYCTR
Mu mAb	-----NRTS-----T-----A-----WT <u>TGTGAYWGQG</u> TLVTVSA
<u>Light Chain</u>	
GL	DIVMTQSPSS LTVTAGEKVT MSCKSSQSLL NSGNQKNYLT WYQKPGQPP <u>KLLIYWASTR</u>
Mu mAb	-----D-----F-----F-----
GL	ESGVPRFTG SCSTDETLT ISSVQAECLA VYYQNDYSY <u>P</u>
Mu mAb	-----V-----CTECGGTKL EIK

FIGURE 2A

mAb	C	T	F	G	G	G	T	K	L	E	I	K	
mAb	g	tgc	acg	ttc	gga	ggg	ggg	acc	aag	ctg	gaa	ata	aaa
J2	-	-a-	-	-	-	-	-	-	-	-	-	-	-c
J2		Y											

FIGURE 2B




FIGURE 3A




FIGURE 3B




FIGURE 4

Clone No.	Concentration $\mu\text{g/mL}$	Sequence					
		HC			LC		
		FW1	FW2	FW3	FW1	FW2	FW3
		4 unique HC			9 unique LC		
1	23.3	a	a	a	b	a	c
2	45.5	a	a	a	e	a	b
3	58.4	a	b	b	e	a	b
4	52.9	a	b	b	b	b	d
5	30	a	a	a	b	b	d
6	7.9	a	a	a	c	a	a
7	24.9	a	a	a	b	b	a
8	32.8	a	b	b	a	a	a
9	16.3	a	a	a	a	a	a
10	61.5	a	b	b	b	a	a
11	31.4	a	a	a	b	a	a
12	34.8	a	a	a	e	c	a
13	8.6	a	a	a	d	b	a
14	48.4	b	b	b	b	a	a
15	20.7	b	b	b	a	a	a
16	32.8	a	c	b	a	a	a

FIGURE 5




FIGURE 6A




FIGURE 6B

Clone No.	Conc. $\mu\text{g/mL}$	Sequence						Ranking	Competition Binding		Ranking
		HC			LC			FACS data	1st exp.	2nd exp.*	
		FW1	FW2	FW3	FW1	FW2	FW3				
Chimeric	20.6	4 unique HC			9 unique LC						
1	23.3	a	a	a	b	a	c	2	7	2	A
2	45.5	a	a	a	e	a	b	6	3	2	D
3	58.4	a	b	b	e	a	b	7	8	14	E
4	52.9	a	b	b	b	b	d	14	15	15	B
5	30	a	a	a	b	b	d	5	5		A
6	7.9	a	a	a	c	a	a	1	7	3	D
7	24.9	a	a	a	b	b	a	4	7		D
8	32.8	a	b	b	a	a	a	7	7	4	C
9	16.3	a	a	a	a	a	a	7	2	4	B
10	61.5	a	b	b	b	a	a	7	6		C
11	31.4	a	a	a	b	a	a	6	4		B
12	34.8	a	a	a	e	c	a	3	8	16	D
13	8.6	a	a	a	d	b	a	6	1	1	D
14	48.4	b	b	b	b	a	a	16	7	15	C
15	20.7	b	b	b	a	a	a	6	7	15	C
16	32.8	a	c	b	a	a	a	15	16	15	C

*empty boxes means worse than 4

FIGURE 7




FIGURE 8A




FIGURE 8B

	10	20	30	40	50	60
						
BAP049-chi-HC	QVQLQQSGSELVRPGASVKLSCKASGYTFTTYWMHWVRQRPQGQGLEWIGNIYPGTGGSNF					
BAP049-hum01-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum02-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum05-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum06-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum07-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum09-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum11-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum12-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum13-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQATGQGLEWMGNIPGTGGSNF					
BAP049-hum03-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWIRQSPSRGLEWLGNIYPGTGGSNF					
BAP049-hum04-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWIRQSPSRGLEWLGNIYPGTGGSNF					
BAP049-hum08-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWIRQSPSRGLEWLGNIYPGTGGSNF					
BAP049-hum10-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWIRQSPSRGLEWLGNIYPGTGGSNF					
BAP049-hum14-HC	QVQLVQSGAEVKKPGASVKVSKASGYTFTTYWMHWIRQSPSRGLEWLGNIYPGTGGSNF					
BAP049-hum15-HC	QVQLVQSGAEVKKPGASVKVSKASGYTFTTYWMHWIRQSPSRGLEWLGNIYPGTGGSNF					
BAP049-hum16-HC	EVQLVQSGAEVKKPGESLRISCKGSGYTFPTYWMHWVRQAPQGQGLEWMGNIPGTGGSNF					

	70	80	90	100	110
					
BAP049-chi-HC	DEKFKNRTSLTVDTSSTAYMHLASLTSEDSAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum01-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum02-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum05-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum06-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum07-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum09-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum11-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum12-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum13-HC	DEKFKNRVTITADKSTSTAYMELSSLRSEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum03-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum04-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum08-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum10-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum14-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum15-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				
BAP049-hum16-HC	DEKFKNRFTISRDN SKNTLYLQMNSLRAEDTAVYYCTRWTGTGAYWQGQTTVTVSS				

FIGURE 9A

	10	20	30	40	50	60
BAP049-chi-HC	QVQLQQSGSELVRPGASVKLSCKASGYTF	FTTYWMHWVRQRPQG	LEWIGNIYPGTGGSNF			
BAP049-hum01-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum02-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum05-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum06-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum07-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum09-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum11-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum12-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum13-HC	E...V...A.VKK..E.LRI...G.....	AT.....M.....				
BAP049-hum03-HC	E...V...A.VKK..E.LRI...G.....	I..S.SR...L.....				
BAP049-hum04-HC	E...V...A.VKK..E.LRI...G.....	I..S.SR...L.....				
BAP049-hum08-HC	E...V...A.VKK..E.LRI...G.....	I..S.SR...L.....				
BAP049-hum10-HC	E...V...A.VKK..E.LRI...G.....	I..S.SR...L.....				
BAP049-hum14-HC	...V...A.VKK.....V.....	I..S.SR...L.....				
BAP049-hum15-HC	...V...A.VKK.....V.....	I..S.SR...L.....				
BAP049-hum16-HC	E...V...A.VKK..E.LRI...G.....	A.....M.....				
	70	80	90	100	110	
BAP049-chi-HC	DEKFKNRTSLTVDTSSTTAYMHLASLTSEDS	AVVYCTRWTTGTGAYWGQGT	TVT	VSS		
BAP049-hum01-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum02-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum05-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum06-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum07-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum09-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum11-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum12-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum13-HC	VTI.A.K.TS....E.S..R...T.....					
BAP049-hum03-HC	FTISR.N.KN.L.LQMN..RA..T.....					
BAP049-hum04-HC	FTISR.N.KN.L.LQMN..RA..T.....					
BAP049-hum08-HC	FTISR.N.KN.L.LQMN..RA..T.....					
BAP049-hum10-HC	FTISR.N.KN.L.LQMN..RA..T.....					
BAP049-hum14-HC	FTISR.N.KN.L.LQMN..RA..T.....					
BAP049-hum15-HC	FTISR.N.KN.L.LQMN..RA..T.....					
BAP049-hum16-HC	FTISR.N.KN.L.LQMN..RA..T.....					

FIGURE 9B

		10	20	30	40	50	60
							
BAP049-chi-LC	DIVMTQSPSSLTVTAGEKVTMSCKSSQSLLDSGNQKNFLT	WYQQKPGQPPKLLIFW	ASTR				
BAP049-hum08-LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum09-LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum15-LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum16-LC	EIVLTQSPDFQSVTPKEKVTITCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum10-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum11-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum14-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum06-LC	DIVMTQTPLSLPVTLPGEPAISCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum07-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGKAPKLLIYW	ASTR				
BAP049-hum13-LC	DVVMTQSPSLPVTLPQPASISCKSSQSLLDSGNQKNFLT	WYQQKPGKAPKLLIYW	ASTR				
BAP049-hum12-LC	DIQMTQSPSSLSASVGDRTVITCKSSQSLLDSGNQKNFLT	WYLOKPGQSPQLLIYW	ASTR				
BAP049-hum02-LC	DIQMTQSPSSLSASVGDRTVITCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum03-LC	DIQMTQSPSSLSASVGDRTVITCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum01-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGQAPRLLIYW	ASTR				
BAP049-hum04-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGKAPKLLIYW	ASTR				
BAP049-hum05-LC	EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLT	WYQQKPGKAPKLLIYW	ASTR				

		70	80	90	100	110
						
BAP049-chi-LC	ESGVFDRFTGSGSVTDFTLT	ISSVQAEDLAVYYCQNDYSYPCTFGQGT	KVEIK			
BAP049-hum08-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum09-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum15-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum16-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum10-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum11-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum14-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum06-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum07-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum13-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum12-LC	ESGVPSRFGSGSGTDFTFT	ISSLEAEDAATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum02-LC	ESGIPPRFSGSGYGTDFLT	INNIESDAAYYFCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum03-LC	ESGIPPRFSGSGYGTDFLT	INNIESDAAYYFCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum01-LC	ESGVPSRFGSGSGTEFTLT	ISSLPDDFATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum04-LC	ESGVPSRFGSGSGTDFTFT	ISSLPEDIATYYCQNDYSYPYTFGQGT	KVEIK			
BAP049-hum05-LC	ESGVPSRFGSGSGTDFTFT	ISSLPEDIATYYCQNDYSYPYTFGQGT	KVEIK			

FIGURE 10A

	10	20	30	40	50	60
BAP049-chi-LC	DI	VT	QSPSS	LT	VT	AGEKVT
BAP049-hum08-LC	E..L...	DFQS..PK...	IT.....		A.R...	Y....
BAP049-hum09-LC	E..L...	DFQS..PK...	IT.....		A.R...	Y....
BAP049-hum15-LC	E..L...	DFQS..PK...	IT.....		A.R...	Y....
BAP049-hum16-LC	E..L...	DFQS..PK...	IT.....		A.R...	Y....
BAP049-hum10-LC	E..L...	AT.SLSP..RA.L			A.R...	Y....
BAP049-hum11-LC	E..L...	AT.SLSP..RA.L			A.R...	Y....
BAP049-hum14-LC	E..L...	AT.SLSP..RA.L			A.R...	Y....
BAP049-hum06-LC		T.L..P..P..PASI			A.R...	Y....
BAP049-hum07-LC	E..L...	AT.SLSP..RA.L			KA....	Y....
BAP049-hum13-LC	V.....	L..P..L.QPASI			KA....	Y....
BAP049-hum12-LC	..Q.....	SASV.DR..IT			L....S.Q...	Y....
BAP049-hum02-LC	..Q.....	SASV.DR..IT			A.R...	Y....
BAP049-hum03-LC	..Q.....	SASV.DR..IT			A.R...	Y....
BAP049-hum01-LC	E..L...	AT.SLSP..RA.L			A.R...	Y....
BAP049-hum04-LC	E..L...	AT.SLSP..RA.L			KA....	Y....
BAP049-hum05-LC	E..L...	AT.SLSP..RA.L			KA....	Y....
	70	80	90	100	110	
BAP049-chi-LC	ES	GV	PDR	FT	GS	SVTDFTLT
BAP049-hum08-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum09-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum15-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum16-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum10-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum11-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum14-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum06-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum07-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum13-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum12-LC	S..S...	G...F...LE...	A.T.....	Y.....		
BAP049-hum02-LC	...I.P..S...	YG.....NNIES	A.Y.F.....	Y.....		
BAP049-hum03-LC	...I.P..S...	YG.....NNIES	A.Y.F.....	Y.....		
BAP049-hum01-LC	S..S...	G.E.....L.PD.F.T		Y.....		
BAP049-hum04-LC	S..S...	G...F...L.P..I.T		Y.....		
BAP049-hum05-LC	S..S...	G...F...L.P..I.T		Y.....		

FIGURE 10B




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14

EIVLTQSPATLSLSPGERATLSCKSSQSLLDSGNQKNFLTWYQQKPGQ
ESGVPSRFSGSGSGTDFTFTISSLEAEDAATYYCQNDYSYPYTFGQGTK
VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQES
LSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

NST N-Glycosylation site
CDR regions are shown in **bold**

Figure 15. Amino acid sequence of the Antibody Molecule A light chain

EVQLVQSGAEVKKPGESLRISCKGSGYTFTTYWMHWVRQATGQGLEWM
DEKFKNRVTITADKSTSTAYMELSSLRSEDNAVYYCTRWTTGTGAYWGC
KGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFP
SLSSVVTVPSSSLGTQTYTCNVDPHKPSNTKVDKRVESKYGPPCPPCPAPEFI
PPKPKDTLMISRTPEVTCVVVDVSDQEDPEVQFNWYVDGVEVHNAKTKPRI
SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPS
SLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDK
SCSVMHREALHNHYTQKSLSLGLG

NST N-Glycosylation site
CDR regions are shown in **bold**

Figure 16. Amino acid sequence of the Antibody Molecule A heavy