



US00PP36094P2

(12) **United States Plant Patent**
Koot

(10) **Patent No.:** **US PP36,094 P2**

(45) **Date of Patent:** **Sep. 3, 2024**

- (54) *Petunia* PLANT NAMED ‘Dopetpotbluvei’
- (50) Latin Name: *Petunia X hybrida*
Varietal Denomination: **Dopetpotbluvei**
- (71) Applicant: **DUMMEN GROUP B.V.**, De Lier (NL)
- (72) Inventor: **Arjan Koot**, Oeffelt (NL)
- (73) Assignee: **DÜMMEN GROUP B.V.**, De Lier (NL)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **18/384,865**
- (22) Filed: **Oct. 29, 2023**
- (51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/82 (2018.01)
- (52) **U.S. Cl.**
USPC **Plt./356.19**

(58) **Field of Classification Search**
USPC Plt./356.19
CPC . A01H 5/02; A01H 5/00; A01H 6/824; A01H 6/82
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS
PP33,746 P2 * 12/2021 Koot Plt./356.19
* cited by examiner
Primary Examiner — June Hwu
(74) *Attorney, Agent, or Firm* — C. Anne Whealy

(57) **ABSTRACT**
A new and distinct cultivar of *Petunia* plant named ‘Dopetpotbluvei’, characterized by its compact and uniformly mounding plant habit; moderately vigorous growth habit and moderate growth rate; freely branching habit; early and freely flowering habit; large bright purple-colored flowers with darker purple-colored venation; and good container and garden performance.

2 Drawing Sheets

1

Botanical designation: *Petunia X hybrida*.
Cultivar denomination: ‘DOPETPOTBLUVEI’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Petunia* plant, botanically known as *Petunia X hybrida* and hereinafter referred to by the name ‘Dopetpotbluvei’.

The new *Petunia* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program is to create new compact, freely branching and early-flowering *Petunia* plants with numerous attractive flowers.

The new *Petunia* plant originated from a cross-pollination made by the Inventor in July, 2020 in Rheinberg, Germany of a proprietary selection of *Petunia X hybrida* identified as code number TT19-K0130, not patented, as the female, or seed, parent with a proprietary selection of *Petunia X hybrida* identified as code number TT20-K0805, not patented, as the male, or pollen, parent. The new *Petunia* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2021.

Asexual reproduction of the new *Petunia* plant by terminal vegetative cuttings in a controlled greenhouse environment in Rheinberg, Germany since May, 2021 has shown that the unique features of this new *Petunia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Petunia* have not been observed under all possible combinations of environmental conditions and

2

cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Dopetpotbluvei’. These characteristics in combination distinguish ‘Dopetpotbluvei’ as a new and distinct *Petunia* plant:

1. Compact and uniformly mounding plant habit.
2. Moderately vigorous growth habit and moderate growth rate.
3. Freely branching habit.
4. Early and freely flowering habit.
5. Large bright purple-colored flowers with darker purple-colored venation.
6. Good container and garden performance.

Plants of the new *Petunia* can be compared to plants of the female parent selection. In side-by-side comparisons, plants of the new *Petunia* differ primarily from plants of the female parent selection in flower shape as flowers of plants of the new *Petunia* are more rounded than flowers of plants of the female parent selection. In addition, flowers of plants of the new *Petunia* are bright purple in color with darker purple-colored venation whereas flowers of plants of the female parent selection are solid purple in color that tend to fade to light purple with subsequent development.

Plants of the new *Petunia* can be compared to plants of the male parent selection. In side-by-side comparisons, plants of the new *Petunia* differ primarily from plants of the male parent selection in flower shape and substance as flowers of plants of the new *Petunia* are more rounded and sturdier than flowers of plants of the male parent selection. In addition, flowers of plants of the new *Petunia* are bright purple in

color with darker purple-colored venation whereas flowers of plants of the male parent selection are solid purple in color.

Plants of the new *Petunia* can be compared to plants of *Petunia X hybrida* 'Sunsurfbv', disclosed in U.S. Plant Pat. No. 17,735. In side-by-side comparisons, plants of the new *Petunia* and 'Sunsurfbv' differ primarily in the following characteristics:

1. Plants of the new *Petunia* are more compact than and not as vigorous as plants of 'Sunsurfbv'.
2. Plants of the new *Petunia* are more freely flowering than plants of 'Sunsurfbv'.
3. Flowers of plants of the new *Petunia* are larger than and not as ruffled as flowers of plants of 'Sunsurfbv'.
4. Flowers of plants of the new *Petunia* are bright purple in color with darker purple-colored venation whereas flowers of plants of 'Sunsurfbv' are light lavender in color with purple-colored venation.
5. Plants of the new *Petunia* are resistant to Tobacco Mosaic Virus whereas plants of 'Sunsurfbv' are not resistant to Tobacco Mosaic Virus.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new *Petunia* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Petunia* plant.

The photograph on the first sheet (FIG. 1) is a side perspective view of a typical flowering plant of 'Dopetpotbluvei' grown in a container.

The photograph on the second sheet (FIG. 2) is a close-up view of typical flowers of 'Dopetpotbluvei'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown during the spring and summer in 22-cm containers in a glass-covered greenhouse in Rheinberg, Germany and under cultural practices typical of commercial *Petunia* production. During the production of the plants, day and night temperatures averaged 18 C and light levels averaged 4,500 lux. Plants were twelve weeks old when the photographs were taken and 25 weeks old when the description was taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, Fifth Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Petunia X hybrida* 'Dopetpotbluvei'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Petunia X hybrida* identified as code number TT19-K0130, not patented.

Male, or pollen, parent.—Proprietary selection of *Petunia X hybrida* identified as code number TT20-K0805, not patented.

Propagation:

Type.—By terminal vegetative cuttings.

Time to initiate roots, summer.—About five days at temperatures about 20 C.

Time to initiate roots, winter.—About seven days at temperatures about 20 C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures about 20 C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures about 20 C.

Root description.—Fine, fibrous; close to 155B in color, actual color of the roots is dependent on substrate composition, water quality, fertilizers, substrate temperature and age of roots.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Compact and uniformly mounding plant habit; freely branching habit with about six primary lateral branches each with about eight secondary branches developing after pinching; moderately vigorous growth habit and moderate growth rate.

Plant height, soil level to top of foliar plane.—About 15 cm.

Plant height, soil level to top of floral plane.—About 18 cm.

Plant diameter.—About 57.5 cm.

Lateral branch description:

Length.—About 33 cm.

Diameter.—About 4 mm.

Internode length.—About 1.5 cm.

Strength.—Moderately strong.

Aspect.—Initially upright to somewhat outwardly spreading.

Texture and luster.—Pubescent; semi-glossy.

Color, developing and developed.—Close to 144C.

Leaf description:

Arrangement.—Before flowering, alternate; after flowering, opposite; simple.

Length.—About 4.5 cm.

Width.—About 2.2 cm.

Shape.—Spatulate.

Apex.—Obtuse.

Base.—Attenuate.

Margin.—Entire.

Texture and luster, upper and lower surfaces.—Pubescent; coriaceous; semi-glossy.

Venation pattern.—Pinnate; arcuate.

Color.—Developing leaves, upper surface: Close to 143A. Developing leaves, lower surface: Close to 143C. Fully expanded leaves, upper surface: Close to 146B; venation, close to 144B. Fully expanded leaves, lower surface: Close to 146C; venation, close to 146D.

Petioles.—Length: About 7.5 mm. Diameter: About 4 mm. Strength: Moderately strong; firm. Texture and luster, upper and lower surfaces: Smooth, glabrous; matte. Color, upper surface: Close to 144C. Color, lower surface: Close to 144D.

Flower description:

Flower type and flowering habit.—Single salverform flowers arising from leaf axils; freely flowering habit with usually about 300 flowers and flower buds developing per plant during the flowering season; flowers face mostly upright to outwardly.

Fragrance.—None detected.

Natural flowering season.—Plants flower continuously during the spring and summer in Germany; early

flowering habit, plants typically beginning flowering about eleven weeks after planting.

Flower longevity.—Individual flowers last about two to three days on the plant; flowers persistent.

Flower buds.—Length: About 4 cm. Diameter: About 7.5 mm. Shape: Ovoid. Texture and luster: Rippled; semi-glossy. Color: Close to 4B and N77C.

Flower diameter.—About 5.6 cm by 5.7 cm.

Flower depth (height).—About 5.5 cm.

Flower throat diameter.—About 1.1 cm.

Flower tube length.—About 2.5 cm.

Flower tube diameter, proximally.—About 4 mm.

Corolla.—Arrangement: Five petals fused at the base and opening into a flared trumpet. Petal lobe length (from throat): About 2.9 cm. Petal lobe width: About 2.8 cm. Petal shape: Roughly spatulate. Petal apex: Obtuse, rounded. Petal margin: Entire; slightly undulate. Petal texture and luster, upper and lower surfaces: Rippled, glabrous; semi-glossy. Throat texture and luster: Rippled; semi-glossy. Tube texture and luster: Rippled; semi-glossy. Color: Petal lobe, when opening, upper surface: Close to N80A. Petal lobe, when opening, lower surface: Close to N82C. Petal lobe, fully opened, upper surface: Close to N80B; venation, close to N92A; color becoming closer to 84B with subsequent development. Petal lobe, fully opened, lower surface: Close to 76A; venation, close to N92A; color becoming closer to 84B with subsequent development. Flower throat: Close to 86B; venation, close to N92A. Flower tube: Close to 85A; venation, close to N92A.

Sepals.—Arrangement: Five sepals fused at the base forming a tubular star-shaped calyx. Length: About 2.4 cm. Diameter: About 3.9 mm. Shape: Oblong. Apex: Rounded. Base: Decurrent. Margin: Entire. Texture and luster, upper and lower surfaces: Smooth, glabrous; semi-glossy. Color: When open-

ing and fully opened, upper surface: Close to 137A. When opening and fully opened, lower surface: Close to 137B.

Peduncles.—Length: About 4 cm. Diameter: About 2 mm. Strength: Moderately strong. Texture and luster: Smooth, glabrous; semi-glossy. Color: Close to 145A.

Reproductive organs.—Stamens: Quantity per flower: Five. Filament length: About 2 cm. Filament color: Close to 155A and N82D. Anther length: About 1.6 mm. Anther shape: Ovate. Anther color: Close to 103A. Pollen amount: Abundant. Pollen color: Close to 94B. Pistils: Quantity per flower: One. Pistil length: About 2.3 cm. Style length: About 2 cm. Style color: Close to N77A, N77C and 145C. Stigma diameter: About 2 mm. Stigma shape: Rounded. Stigma color: Close to 147A. Ovary color: Close to 142A. Fruits: Quantity produced per plant: About 200 during the flowering season. Length: About 5 mm. Diameter: About 4 mm. Texture: Smooth, glabrous. Color: Close to 199B. Seeds: Quantity per flower: About 50. Length: About 0.7 mm. Diameter: About 0.7 mm. Texture: Smooth, glabrous. Color: Close to 200A.

Garden performance: Plants of the new *Petunia* have been observed to have good garden performance and tolerate wind, rain, temperatures ranging from about 5 C to about 40 C and to be hardy to USDA Hardiness Zone 11.

Pathogen & pest resistance: Plants of the new *Petunia* have been observed to be resistant to Tobacco Mosaic Virus. To date, plants of the new *Petunia* have not been observed to be resistant to pests and other pathogens common to *Petunia* plants.

It is claimed:

1. A new and distinct *Petunia* plant named ‘Dopetpotbluvei’ as illustrated and described.

* * * * *



FIG. 1



FIG. 2