



(12) **Patent Application Publication**
Eaton et al.

(43) **Pub. Date:** **Sep. 25, 2003**

The present invention is directed to novel polypeptides and to nucleic acid molecules encoding those polypeptides. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

FIGURE 1

GGGGCTTCGGCGCCAGCGGCCAGCGCTAGTCGGTCTGGTAAGGATTTACAAAAGGTGCAGGTATG
AGCAGGTCTGAAGACTAACATTTTGTGAAGTTGTAAAACAGAAAACCTGTTAGAAATGTGGTGGT
TTCAGCAAGGCCTCAGTTTCCTTCCTTCAGCCCTTGTAATTTGGACATCTGCTGCTTTCATATTT
TCATACATTACTGCAGTAACACTCCACCATATAGACCCGGCTTTACCTTATATCAGTGACACTGG
TACAGTAGCTCCAGAAAAATGCTTATTTGGGGCAATGCTAAATATTGCGGCAGTTTTATGCATTG
CTACCATTTATGTTTCGTTATAAGCAAGTTCATGCTCTGAGTCCTGAAGAGAACGTTATCATCAA
TTAAACAAGGCTGGCCTTGTAAGTTCGTAAGTTCGTAAGTTCGTAAGTTCGTAAGTTCGTAAGTTC
CCAGAAAACAACCTTTTTTGCTGCACATGTAAGTGGAGCTGTGCTTACCTTTGGTATGGGCTCAT
TATATATGTTTGTTCAGACCATCCTTTCCTACCAAATGCAGCCCCAAAATCCATGGCAAACAAGTC
TTCTGGATCAGACTGTTGTTGGTTATCTGGTGTGGAGTAAGTGCACCTAGCATGCTGACTTGCTC
ATCAGTTTTGCACAGTGGCAATTTTGGGACTGATTTAGAACAGAACTCCATTGGAACCCCGAGG
ACAAAGGTTATGTGCTTCACATGATCACTACTGCAGCAGAATGGTCTATGTCATTTTCCTTCTTT
GGTTTTTTCCTGACTTACATTCGTGATTTTCAGAAAATTTCTTTACGGGTGGAAGCCAATTTACA
TGGATTAACCTCTATGACACTGCACCTTGCCCTATTAACAATGAACGAACACGGCTACTTTCCA
GAGATATTTGATGAAAGGATAAAATATTTCTGTAATGATTATGATTCTCAGGGATTGGGGAAAGG
TTCACAGAAGTTGCTTATTCTTCTCTGAAATTTTCAACCACTTAATCAAGGCTGACAGTAACACT
GATGAATGCTGATAATCAGGAAACATGAAAGAAGCCATTTGATAGATTATTCTAAAGGATATCAT
CAAGAAGACTATTAAAAACACCTATGCCTATACTTTTTTATCTCAGAAAATAAAGTCAAAGACT
ATG

FIGURE 2

<subunit 1 of 1, 266 aa, 1 stop

<MW: 29766, pI: 8.39, NX(S/T): 0

MWWFQQGLSFLPSALVIWTSAAFIIFS YITAVTLHHIDPALPYISDTGTVAPEKCLFGAMLNIAAV
LCIATIYVRYKQVHALSPEENVIIKLNKAGLVLGILSCLGLSIVANFQKTTLFAAHVSGAVLTFG
MGSLYMFVQTILSYQMOPKIHGKQVFWIRLLLVWCGVSALSMLTCSSVLHSGNFGTDLEQKLHW
NPEDKGYVLHMITTAAEWSMSFSFFGFFLT YIRDFQKISLRVEANLHGLTLYDTAPCPINNERTR
LLSRDI

Important features:

Type II transmembrane domain:

amino acids 13-33

Other Transmembrane domains:

amino acids 54-73, 94-113, 160-180, 122-141

N-myristoylation sites.

amino acids 57-63, 95-101, 99-105, 124-130, 183-189

FIGURE 3

CGGACGCGTGGGCGGACGCGTGGGGGAGAGCCGCGAGTCCCGGCTGCAGCACCTGGGAGAAGGCAGACC
 GTGTGAGGGGGCCTGTGGCCCCAGCGTGCTGTGGCCTCGGGGAGTGGGAAGTGGAGGCAGGAGCCTTC
 CTTACACTTCGCCATGAGTTTCCTCATCGACTCCAGCATCATGATTACCTCCCAGATACTATTTTTTG
 GATTTGGGTGGCTTTTCTTCATGCGCCAATTGTTTAAAGACTATGAGATACGTCAGTATGTTGTACAG
 GTGATCTTCTCCGTGACGTTTGCATTTTCTTGACCATGTTTGAGCTCATCATCTTTGAAATCTTAGG
 AGTATTGAATAGCAGCTCCCGTTATTTTCACTGGAAAATGAACCTGTGTGTAATTCTGCTGATCCTGG
 TTTTCATGGTGCCTTTTACATTGGCTATTTTATTGTGAGCAATATCCGACTACTGCATAAACAACGA
 CTGCTTTTTTCTGTCTCTTATGGCTGACCTTTATGTATTTCTTCTGGAACTAGGAGATCCCTTTCC
 CATTCCTCAGCCCCAAACATGGGATCTTATCCATAGAACAGCTCATCAGCCGGGTGGTGTGATTGGAG
 TGACTCTCATGGCTCTTCTTTCTGGATTTGGTGCTGTCAACTGCCCATACACTTACATGTCTTACTTC
 CTCAGGAATGTGACTGACACGGATATTCTAGCCCTGGAACGGCGACTGCTGCAAACCATGGATATGAT
 CATAAGCAAAAAGAAAAGGATGGCAATGGCACGGAGAACAATGTTCCAGAAGGGGGGAAGTGCATAACA
 AACCATCAGGTTTTCTGGGGAATGATAAAAAGTGTTACCACTTCAGCATCAGGAAGTGAAATCTTACT
 CTTATTCAACAGGAAGTGGATGCTTTGGAAGAATTAAGCAGGCAGCTTTTTCTGGAAACAGCTGATCT
 ATATGCTACCAAGGAGAGAATAGAATACTCCAAAACCTTCAAGGGGAAATATTTTAATTTTCTTGTT
 ACTTTTTCTCTATTTACTGTGTTTGGAAAATTTTCATGGCTACCATCAATATTGTTTTTGATCGAGTT
 GGGAAAACGGATCCTGTCACAAGAGGCATTGAGATCACTGTGAATTATCTGGGAATCCAATTTGATGT
 GAAGTTTTGGTCCCAACACATTTCTTCATTCTTGTGTTGGAATAATCATCGTCACATCCATCAGAGGAT
 TGCTGATCACTCTTACCAAGTTCTTTTATGCCATCTCTAGCAGTAAGTCCTCCAATGTCATTGTCCTG
 CTATTAGCACAGATAATGGGCATGTACTTTGTCTCCTCTGTGCTGCTGATCCGAATGAGTATGCCTTT
 AGAATACCGCACCATAATCACTGAAGTCCTTGGAGAACTGCAGTTCAACTTCTATCACCGTTGGTTTG
 ATGTGATCTTCCTGGTCAGCGCTCTCTCTAGCATACTCTTCCTCTATTTGGCTCACAAACAGGCACCA
 GAGAAGCAAATGGCACCTTGAACTTAAGCCTACTACAGACTGTTAGAGGCCAGTGGTTTCAAATTTA
 GATATAAGAGGGGGGAAAAATGGAACCAGGGCCTGACATTTTATAAACAAACAAAATGCTATGGTAGC
 ATTTTTACCTTCATAGCATACTCCTTCCCCGTCAGGTGATACTATGACCATGAGTAGCATCAGCCAG
 AACATGAGAGGGAGAACTAACTCAAGACAATACTCAGCAGAGAGCATCCCGTGTGGATATGAGGCTGG
 TGTAGAGGCGGAGAGGAGCAAGAACTAAAGGTGAAAAATACACTGGAACCTCTGGGGCAAGACATGT
 CTATGGTAGCTGAGCCAAACACGTAGGATTTCCGTTTTAAGGTTACATGGAAAAGGTTATAGCTTTG
 CCTTGAGATTGACTCATTAAATCAGAGACTGTAACAAAAAAAAAAAAAAAAAAGGGCGGCCGCG
 ACTCTAGAGTCGACCTGCAGAAGCTTGGCCGCCATGGCCCACTTGTTTATTGCAGCTTATAATG

FIGURE 4

MSFLIDSSIMITSQILFFGFGWLFFMRQLFKDYEIRQYVVQVIFSVTFAFSCTMFELIIFEILGV
LNSSSRYPFHWMNLCVILLILVFMVPFYIGYFIVSNIRLLHKQRLLFSCLLWLTFMYFFWKLGBP
FPILSPKHGILSIEQLISRVGIVGVTLMALLSGFGAVNCPYTYMSYFLRNVTDTDILALERLLQ
TMDMIISKKKRMAMARRTMFQKGEVHNKPSGFWGMIKSVTTSASGSENLTLIQQEVDALAEELSRQ
LFLETADLYATKERIEYSKTFKGKYFNFLGYFFSIYCVWKIFMATINIVFDRVGKTDVPVTRGIEI
TVNYLGIQFDVKFWSQHISFILVGIIIVTSIRGLLITLTKFFYAISSSKSSNVIVLLLAQIMGY
FVSSVLLIRMSMPLEYRTIITEVLGELQFNFYHRWFDVIFLVSALSSILFLYLAKQAPEKQMAP

Important features:

Signal peptide:

amino acids 1-23

Potential transmembrane domains:

amino acids 37-55, 81-102, 150-168, 288-311, 338-356, 375-398,
425-444

N-glycosylation sites.

amino acids 67-70, 180-183 and 243-246

Eukaryotic cobalamin-binding proteins

amino acids 151-160

FIGURE 5

AGCAGGGAAATCCGGATGTCTCGGTTATGAAGTGGAGCAGTGAGTGTGAGCCTCAACATAGTTCC
 AGAACTCTCCATCCGGACTAGTTATTGAGCATCTGCCTCTCATATCACCAGTGGCCATCTGAGGT
 GTTTCCTTGGCTCTGAAGGGGTAGGCACGATGGCCAGGTGCTTCAGCCTGGTGTGCTTCTCACT
 TCCATCTGGACCACGAGGCTCCTGGTCCAAGGCTCTTTGCGTGCAGAAGAGCTTTCCATCCAGGT
 GTCATGCAGAATTATGGGGATCACCTTGTGAGCAAAAAGGCGAACCAGCAGCTGAATTTACAG
 AAGCTAAGGAGGCCTGTAGGCTGCTGGGACTAAGTTTGGCCGGCAAGGACCAAGTTGAAACAGCC
 TTGAAAGCTAGCTTTGAAACTTGCAGCTATGGCTGGGTGGAGATGGATTTCGTGGTCATCTCTAG
 GATTAGCCCCAAACCCCAAGTGTGGGAAAAATGGGGTGGGTGTCCTGATTTGGAAGGTTCCAGTGA
 GCCGACAGTTTGCAGCCTATTGTTACAACCTCATCTGATACTTGGACTAACTCGTGCATTCCAGAA
 ATTATCACCACCAAGATCCCATATTCAACACTCAAACGCAACACAAACAAGAAATTTATTGT
 CAGTGACAGTACCTACTCGGTGGCATCCCCTTACTCTACAATACTGCCCCCTACTACTCTCTC
 CTGCTCCAGCTTCCACTTCTATTCCACGGAGAAAAAATTGATTTGTGTACAGAAGTTTTTATG
 GAACTAGCACCATGTCTACAGAACTGAACCATTTGTTGAAAATAAAGCAGCATTCAGAATGA
 AGCTGCTGGGTTTGGAGGTGTCCCCACGGCTCTGCTAGTGCTTGCTCTCTCTTCTTGGTGCTG
 CAGCTGGTCTTGGATTTTGCTATGTCAAAAGGTATGTGAAGGCCTTCCCTTTTACAAACAAGAAT
 CAGCAGAAGGAAATGATCGAAACCAAGTAGTAAAGGAGGAGAAGGCCAATGATAGCAACCCTAA
 TGAGGAATCAAAGAAAAGTGAATAAAACCCAGAAGAGTCCAAGAGTCCAAGCAAAAAGTACCCTGC
 GATGCCTGGAAGCTGAAGTTTAGATGAGACAGAAATGAGGAGACACACCTGAGGCTGGTTTCTTT
 CATGCTCCTTACCCCTGCCCCAGCTGGGGAAATCAAAGGGCCAAAGAACCAAGAAGAAAAGTCCA
 CCCTTGGTTCTTAAGTGAATCAGCTCAGGACTGCCATTGGACTATGGAGTGCACCAAAGAGAAT
 GCCCTTCTCCTTATTGTAACCCTGTCTGGATCCTATCCTCCTACCTCCAAAGCTTCCACGGCCT
 TTCTAGCCTGGCTATGTCCTAATAATATCCCAGTGGGAGAAAGGAGTTTTTGCAAAGTGCAAGGAC
 CTAAACATCTCATCAGTATCCAGTGGTAAAAAGGCCTCCTGGCTGTCTGAGGCTAGGTGGGTTG
 AAAGCCAAGGAGTCACTGAGACCAAGGCTTCTCTACTGATTCCGCAGCTCAGACCCTTTCTTCA
 GCTCTGAAAGAGAAACACGTATCCACCTGACATGTCCTTCTGAGCCCGGTAAGAGCAAAAGAAT
 GGCAGAAAAGTTTAGCCCCCTGAAAGCCATGGAGATTCTCATAACTTGAGACCTAATCTCTGTAAA
 GCTAAAATAAAGAAATAGAACAAGGCTGAGGATACGACAGTACACTGTCAGCAGGGACTGTAAAC
 ACAGACAGGGTCAAAGTGTCTTCTCTGAACACATTGAGTTGGAATCACTGTTTAGAACACACACA
 CTTACTTTTTTCTGGTCTCTACCACTGCTGATATTTTCTCTAGGAAATATACTTTTACAAGTAACA
 AAAATAAAAAGTCTTATAAATTTCTATTTTTATCTGAGTTACAGAAATGATTACTAAGGAAGATT
 ACTCAGTAATTTGTTTTAAAAGTAATAAAATTCACAAACATTTGCTGAATAGCTACTATATGTC
 AAGTGCTGTGCAAGGTATTACACTCTGTAATTGAATATTATTCCTCAAAAAATTGCACATAGTAG
 AACGCTATCTGGGAAGCTATTTTTTTTCAGTTTTGATATTTCTAGCTTATCTACTTCCAACTAAT
 TTTTATTTTTTCTGAGACTAATCTTATTCATTTTCTCTAATATGGCAACCATTATAACCTTAATT
 TATTATTAACATACCTAAGAAGTACATTGTTACCTCTATATACCAAAGCACATTTTAAAAGTGCC
 ATTAACAAATGTATCACTAGCCCTCCTTTTTTCCAACAAGAAGGACTGAGAGATGCAGAAATATT
 TGTGACAAAAAATTAAAGCATTTAGAAAACCT

FIGURE 6

MARCFSLVLLLLTSIWTRLLVQGSRLAEELSIQVSCRIMGITLVSKKANQQLNFTAEAKEACRLLG
LSLAGKDQVETALKASFETCSYGWVGDFVVISRISPNPKCGKNGVGVLIWKVPVSRQFAAYCYN
SSDTWTNSCIPEIIITTKDPIFNTQTATQTTEFIVSDSTYSVASPYSTIPAPTTTTPPAPASTSIPR
RKKLICVTEVFMETSTMSTETEPFVENKAAFKNEAAGFGGVPTALLVLALLFFGAAAGLGFCYVK
RYVKAFPFTNKNQQKEMIETKVVKEEKANDSNPNEEKKTDKNPEESKSPSKTTVRCLEAEV

Signal sequence:

amino acids 1-16

Transmembrane domain:

amino acids 235-254

N-glycosylation site.

amino acids 53-57, 130-134, 289-293

Casein kinase II phosphorylation site.

amino acids 145-149, 214-218

Tyrosine kinase phosphorylation site.

amino acids 79-88

N-myristoylation site.

amino acids 23-29, 65-71, 234-240, 235-239, 249-255, 253-259

FIGURE 7

CGCCGCGCTCCCGCACCCGCGGGCCCGCCACCGCGCCGCTCCCGCATCTGCACCCGCGAGCCCGGC
GGCCTCCCGGCGGGAGCGAGCAGATCCAGTCCGGCCCCGAGCGCAACTCGGTCCAGTCCGGGGCGG
CGGCTGCGGGCGCAGAGCGGAGATGCGAGCGGCTTGGGGCCACCCTGCTGTGCCTGCTGCTGGCGG
CGGCGGTCCCCACGGCCCCCGCGCCCGCTCCGACGGCGACCTCGGTCCAGTCAAGCCCGGCCCG
GCTCTCAGCTACCCGCAGGAGGAGGCCACCCTCAATGAGATGTTCCGCGAGGTTGAGGAACTGAT
GGAGGACACGCAGCACAAATTGCGCAGCGCGGTGGAAGAGATGGAGGCAGAAGAAGCTGCTGCTA
AAGCATCATCAGAAGTGAACCTGGCAAACCTTACCTCCCAGCTATCACAATGAGACCAACACAGAC
ACGAAGGTTGGAAATAATACCATCCATGTGCACCGAGAAATTCACAAGATAACCAACAACCAGAC
TGGACAAATGGTCTTTTCAGAGACAGTTATCACATCTGTGGGAGACGAAGAAGGCAGAAGGAGCC
ACGAGTGCATCATCGACGAGGACTGTGGGCCCAGCATGTACTGCCAGTTTGCCAGCTTCCAGTAC
ACCTGCCAGCCATGCCGGGGCCAGAGGATGCTCTGCACCCGGGACAGTGAGTGCTGTGGAGACCA
GCTGTGTGTCTGGGGTCACTGCACCAAAATGGCCACCAGGGGCGAGCAATGGGACCATCTGTGACA
ACCAGAGGGACTGCCAGCCGGGGCTGTGCTGTGCCTTCCAGAGAGGCCTGCTGTTCCCTGTGTGC
ACACCCCTGCCCGTGGAGGGCGAGCTTTGCCATGACCCCGCCAGCCGGCTTCTGGACCTCATCAC
CTGGGAGCTAGAGCCTGATGGAGCCTTGGACCGATGCCCTTGTGCCAGTGGCCTCCTCTGCCAGC
CCCACAGCCACAGCCTGGTGTATGTGTGCAAGCCGACCTTCGTGGGGAGCCGTGACCAAGATGGG
GAGATCCTGCTGCCCAGAGAGGTCCCCGATGAGTATGAAGTTGGCAGCTTCATGGAGGAGGTGCG
CCAGGAGCTGGAGGACCTGGAGAGGAGCCTGACTGAAGAGATGGCGCTGGGGGAGCCTGCGGCTG
CCGCCGCTGCACTGCTGGGAGGGGAAGAGATTAGATCTGGACCAGGCTGTGGGTAGATGTGCAA
TAGAAATAGCTAATTTATTTCCCCAGGTGTGTGCTTTAGGCGTGGGCTGACCAGGCTTCTTCCTA
CATCTTCTTCCCAGTAAGTTTCCCCTCTGGCTTGACAGCATGAGGTGTTGTGCATTTGTTTCAGCT
CCCCCAGGCTGTTCTCCAGGCTTCACAGTCTGGTGCTTGGGAGAGTCAGGCAGGGTTAAACTGCA
GGAGCAGTTTGCCACCCCTGTCCAGATTATTGGCTGCTTTGCCTCTACCAGTTGGCAGACAGCCG
TTTGTCTACATGGCTTTGATAATTGTTTGAGGGGAGGAGATGGAACAATGTGGAGTCTCCCTC
TGATTGGTTTGGGGAAATGTGGAGAAGAGTGCCCTGCTTTGCAAACATCAACCTGGCAAAAATG
CAACAAATGAATTTCCACGCAGTTCTTTCCATGGGCATAGGTAAGCTGTGCCTTCAGCTGTTGC
AGATGAAATGTTCTGTTACCCCTGCATTACATGTGTTTATTCATCCAGCAGTGTGCTCAGCTCC
TACCTCTGTGCCAGGGCAGCATTTTCATATCCAAGATCAATTCCTCTCTCAGCACAGCCTGGGG
AGGGGGTCATTGTTCTCCTCGTCCATCAGGGATCTCAGAGGCTCAGAGACTGCAAGCTGCTTGCC
CAAGTCACACAGCTAGTGAAGACCAGAGCAGTTTCATCTGGTTGTGACTCTAAGCTCAGTGCTCT
CTCCACTACCCACACCAGCCTTGGTGCCACCAAAAGTGCTCCCCAAAAGGAAGGAGAATGGGAT
TTTTCTTGAGGCATGCACATCTGGAATTAAGGTCAAACCTAATTCTCACATCCCTCTAAAAGTAAA
CTACTGTTAGGAACAGCAGTGTCTCACAGTGTGGGGCAGCCGTCTTCTAATGAAGACAATGAT
ATTGACACTGTCCCTCTTTGGCAGTTGCATTAGTAACTTTGAAAGGTATATGACTGAGCGTAGCA
TACAGGTTAACCTGCAGAAACAGTACTTAGGTAATTGTAGGGCGAGGATTATAAATGAAATTTGC
AAAATCACTTAGCAGCAACTGAAGACAATTATCAACCACGTGGAGAAAATCAAACCGAGCAGGGC
TGTGTGAAACATGGTTGTAATATGCGACTGCGAACACTGAACTCTACGCCACTCCACAAATGATG
TTTTCAGGTGTGATGGACTGTTGCCACCATGTATTCATCCAGAGTTCTTAAAGTTTAAAGTTGCA
CATGATTGTATAAGCATGCTTTCTTTGAGTTTTAAATTATGTATAAACATAAGTTGCATTTAGAA
ATCAAGCATAAATCACTTCAACTGCAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 8

MQRLGATLLCLLLAAVPTAPAPAPTATSAPVKPGPALSYPOEEATLNEMFREVEELMEDTQHKL
RSAVEEMEAEAEAAKASSEVNLANLPPTYHNETNTDTKVGNNTIHVHREIHKITNNQTGQMVFSE
TVITSVGDEEGRRSHECIIDEDCGPSMYCQFASFQYTCQPCRGQRMCLTRDSECCGDQLCVWGHC
TKMATRGSNGTICDNQRDCQPGGCCAFQRGLLFPVCTPLPVEGELCHDPASRLLDLITWELEPDG
ALDRPCASGLLCQPHSHSLVYVCKPTFVGSRDQDGEILLPREVPDEYEVGSFMEEVRQLELDLE
RSLTEEMALGEPAAAAAALLGGEI

Signal sequence:

amino acids 1-19

N-glycosylation site.

amino acids 96-100, 106-110, 121-125, 204-208

Casein kinase II phosphorylation site.

amino acids 46-50, 67-71, 98-102, 135-139, 206-210, 312-316,
327-331

N-myristoylation site.

amino acids 202-208, 217-223

Amidation site.

amino acids 140-144

FIGURE 9

CGGACGCGTGGGCGGACGCGTGGGGGCTGTGAGAAAGTGCCAATAAATACATCATGCAACCCAC
GGCCACCTTGTGAACTCCTCGTGCCCAGGGCTGATGTGCGTCTTCCAGGGCTACTCATCCAAAG
GCCTAATCCAACGTTCTGTCTTCAATCTGCAAATCTATGGGGTCCTGGGGCTCTTCTGGACCCTT
AACTGGGTACTGGCCCTGGGCCAATGCGTCCTCGCTGGAGCCTTTGCCTCCTTCTACTGGGCCTT
CCACAAGCCCCAGGACATCCCTACCTTCCCCTTAATCTCTGCCTTCATCCGCACACTCCGTTACC
AACTGGGTCAATTGGCATTGGAGCCCTCATCCTGACCCTTGTGCAGATAGCCCGGGTCATCTTG
GAGTATATTGACCACAAGCTCAGAGGAGTGCAGAACCTGTAGCCCGCTGCATCATGTGCTGTTT
CAAGTGCTGCCTCTGGTGTCTGGAAAAATTTATCAAGTTCCTAAACCGCAATGCATACATCATGA
TCGCCATCTACGGGAAGAATTTCTGTGTCTCAGCCAAAAATGCGTTCATGCTACTCATGCGAAAC
ATTGTCAGGGTGGTCTGCTGACAAAGTCACAGACCTGCTGCTGTTCTTTGGGAAGCTGCTGGT
GGTCGGAGGCGTGGGGGTCTGTCTTCTTTTTTTCTCCGGTCGCATCCCGGGGTGGGTAAAG
ACTTTAAGAGCCCCACCTCAACTATTACTGGCTGCCCATCATGACCTCCATCCTGGGGGCCTAT
GTCATCGCCAGCGGCTTCTTCAGCGTTTTTCGGCATGTGTGTGGACACGCTCTTCCTCTGCTTCCT
GGAAGACCTGGAGCGGAACAACGGCTCCCTGGACCGGCCCTACTACATGTCCAAGAGCCTTCTAA
AGATTCTGGGCAAGAAGAACGAGGCGCCCCCGGACAACAAGAAGAGGAAGAAGTGACAGCTCCGG
CCCTGATCCAGGACTGCACCCCCACCCCAACCGTCCAGCCATCCAACCTCACTTCGCCTTACAGGT
CTCCATTTTGTGGTAAAAAAGGTTTTAGGCCAGGCGCCGTGGCTCACGCCTGTAATCCAACACT
TTGAGAGGCTGAGGCGGGCGGATCACCTGAGTCAGGAGTTCGAGACCAGCCTGGCCAACATGGTG
AAACCTCCGTCTCTATTAAAAATACAAAAATTAGCCGAGAGTGGTGGCATGCACCTGTCTATCCCA
GCTACTCGGGAGGCTGAGGCAGGAGAATCGCTTGAACCCGGGAGGCAGAGGTTGCAGTGAGCCGA
GATCGCGCCACTGCACTCCAACCTGGGTGACAGACTCTGTCTCCAAAACAAAACAAAACAAA
AAGATTTTATTAAAGATATTTTGTTAAGTC

FIGURE 10

RTRGRTRGGCEKVPINTSCNPTAHLVNSSCPGLMCVFQGYSSKGLIQRSVFNLQIYGVLGLEFWTL
NWVLALGQCVLGAFASFYWAFHKPQDIPTFPLISAFIRTLRYHTGSLAFGALILTLVQIARVIL
EYIDHKLRGVQNPVARCIMCCFKCCLWCLEKFIKFLNRNAYIMIAIYGKNFCVSAKNAFMLLMRN
IVRVVVLDKVTDLLLFEGKLLVVGVGVLSEFFFSGRIPGLGKDFKSPHLNYYWLPIMTSILGAY
VIASGFFSVFGMCDTLFLCFLEDLERNNGSLDRPYMSKSLKILGKKNEAPPDNKKRKK

Important features:

Transmembrane domains:

amino acids 57-80 (type II), 110-126, 215-231, 254-274

N-glycosylation sites.

amino acids 16-20, 27-31, 289-293

Hypothetical YBR002c family proteins.

amino acids 276-288

Ammonium transporters proteins.

amino acids 204-231

N-myristoylation sites.

amino acids 60-66, 78-84

Amidation site.

amino acids 306-310

FIGURE 11

GCCCCGCGCCCGGCCCGCCCGAAGCCGGAGCCACCGCCATGGGGGGGCTGCCTGGGAGGCTGC
 TCCCTGCTCAGCTGCGCGTCTCTGCCTCTGCGGCTCTGCCCCCTGCATCCTGTGCAGCTGCTGCCCCG
 CAGCCGCAACTCCACCGTGAGCCGCCTCATCTTCACGTTCTTCCTCTTCCTGGGGGTGCTGGTGTCCA
 TCATTATGCTGAGCCCGGGCGTGGAGAGTCAGCTCTACAAGCTGCCCTGGGTGTGTGAGGAGGGGGCC
 GGGATCCCCACCGTCTTGACAGGGCCACATCGACTGTGGCTCCCTGCTTGGCTACCGCGCTGTCTACCG
 CATGTGCTTCGCCACGGCGGCCTTCTTCTTCTTCTTTTACCCCTGCTCATGCTCTGCGTGAGCAGCA
 GCCGGGACCCCCGGGCTGCCATCCAGAATGGGTTTTGGTTCTTTAAGTTCTTGATCCTGGTGGGCCTC
 ACCGTGGGTGCCTTCTACATCCCTGACGGCTCCTTCACCAACATCTGGTTCTACTTCGGCGTCTGTGG
 CTCCTTCCTCTTCATCCTCATCCAGCTGGTGCTGCTCATCGACTTTGCGCACTCCTGGAACCAGCGGT
 GGCTGGGCAAGGCCGAGGAGTGCATTCCCGTGCTGGTACGCAGGCCTCTTCTTCTTCACTCTCCTC
 TTCTACTTGCTGTGATCGCGCCGTGGCGCTGATGTTTCATGTACTACACTGAGCCCAGCGGCTGCCA
 CGAGGGCAAGGTCTTCATCAGCCTCAACCTCACCTTCTGTGTCTGCGTGTCCATCGCTGCTGTCTGCTG
 CCAAGGTCCAGGACGCCCAGCCCAACTCGGGTCTGCTGCAGGCCTCGGTATCACCTCTACACCATG
 TTTGTACCTGGTCAGCCCTATCCAGTATCCCTGAACAGAAATGCAACCCCCATTTGCCAACCAGCT
 GGGCAACGAGACAGTTGTGGCAGGCCCGAGGGCTATGAGACCCAGTGGTGGGATGCCCCGAGCATTG
 TGGGCCTCATCATCTTCCTCCTGTGCACCTCTTCATCAGTCTGCGCTCCTCAGACCACCGGCAGGTG
 AACAGCCTGATGCAGACCGAGGAGTGCCACCTATGCTAGACGCCACACAGCAGCAGCAGCAGCAGGT
 GGCAGCCTGTGAGGGCCGGGCCTTTGACAACGAGCAGGACGGCGCTCACCTACAGCTACTCCTTCTTCC
 ACTTCTGCCTGGTGTGTCCTCACTGCACGTATGATGACGCTCACCAACTGGTACAAGCCCGGTGAG
 ACCCGGAAGATGATCAGCACGTGGACCGCCGTGTGGGTGAAGATCTGTGCCAGCTGGGCAGGGCTGCT
 CCTCTACCTGTGGACCCTGGTAGCCCCACTCCTCCTGCGCAACCGCGACTTCAGCTTGAGGCAGCCTCA
 CAGCCTGCCATCTGGTGCCTCCTGCCACCTGGTGCCTCTCGGCTCGGTGACAGCCAACCTGCCCCCTC
 CCCACACCAATCAGCCAGGCTGAGCCCCCACCCTGCCCCAGCTCCAGGACCTGCCCCTGAGCCGGGC
 CTTCTAGTCGTAGTGCCTTCAGGGTCCGAGGAGCATCAGGCTCCTGCAGAGCCCCATCCCCCGCCAC
 ACCCACACGGTGGAGCTGCCTCTTCCTTCCCCTCCTCCCTGTTGCCATACTCAGCATCTCGGATGAA
 AGGGCTCCCTTGTCTCAGGCTCCACGGGAGCGGGGCTGCTGGAGAGAGCGGGGAACCTCCCACCACAG
 TGGGGCATCCGGCACTGAAGCCCTGGTGTTCCTGGTACGTCCCCCAGGGGACCCTGCCCCCTTCTTG
 GACTTCGTGCCTTACTGAGTCTCTAAGACTTTTTCTAATAAAACAAGCCAGTGCGTGTAAAAAAA

FIGURE 12

MGACLGACSLSCASCLCGSAPCILCSCCPASRNSTVSRLI FTFFFLGVLVSI IMLSPGVESQL
YKLPWVCEEGAGIPTVLQGHIDCGSLLGYRAVYRMCFATAA FFFFFFFTLLMLCVSSSRDPRAAIQ
NGFWFFKFLILVGLTVGAFYIPDGSFTNIWFFYFGVVGSFLFILIQLVLLIDFAHSWNQRWL GKAE
ECDSRAWYAGLFFFTLLFYLLSIAAVALMFMYYTEPSGCHEGKVFISLNLTFVCVVSIAAVLPKV
QDAQPN SGLLQASVITLYTMFVTWSALSSIPEQKCNPHLPTQLGNETVVAGPEGYETQWW DAPSI
VGLIIFLLCTLFISLRSSDHRQVNSLMQTEECPPMLDATQQQQQQVAACEGRAFDNEQDGV TYSY
SEFFHFCVLVSLHVMMLTNWYKPGETRKMISTWTAVWVKICASWAGLLLYLWTLVAPLLLRNRD
FS

Signal sequence:

amino acids 1-20

Transmembrane domains:

amino acids 40-58, 101-116, 134-150, 162-178, 206-223, 240-257,
272-283, 324-340, 391-406, 428-444

FIGURE 13

CGGGCCAGCCTGGGGCGGGCCGGCCAGGAACCAACCCGTTAAGGTGTCTTCTCTTTAGGGATGGTGA
GGTTGGAAAAAGACTCCTGTAACCCCTCCTCCAGGATGAACCACCTGCCAGAAGACATGGAGAACG
CTCTCACCGGGAGCCAGAGCTCCCATGCTTCTCTGCGCAATATCCATTCCATCAACCCACACAA
CTCATGGCCAGGATTGAGTCCTATGAAGGAAGGGAAAAGAAAGGCATATCTGATGTCAGGAGGAC
TTTCTGTTTGTGTTGTACCTTTGACCTCTTATTCGTAACATTACTGTGGATAATAGAGTTAAATG
TGAATGGAGGCATTGAGAACACATTAGAGAAGGAGGTGATGCAGTATGACTACTATTCTTCATAT
TTTGATATATTTCTTCTGGCAGTTTTTCGATTTAAAGTGTTAATACTTGCATATGCTGTGTGCAG
ACTGCGCCATTGGTGGGCAATAGCGTTGACAACGGCAGTGACCAGTGCCTTTTTACTAGCAAAAG
TGATCCTTTCGAAGCTTTTCTCTCAAGGGGCTTTTGGCTATGTGCTGCCCATCATTTTCATTCATC
CTTGCCTGGATTGAGACGTGGTTCCTGGATTTCAAAGTGTTACCTCAAGAAGCAGAAGAAGAAAA
CAGACTCCTGATAGTTCAGGATGCTTCAGAGAGGGCAGCACTTATACCTGGTGGTCTTTCTGATG
GTCAGTTTTATTCCCCTCCTGAATCCGAAGCAGGATCTGAAGAAGCTGAAGAAAAACAGGACAGT
GAGAAACCACTTTTAGAACTATGAGTACTACTTTTGTAAATGTGAAAAACCCTCACAGAAAGTC
ATCGAGGCCAAAAAGAGGCAGGCAGTGGAGTCTCCCTGTCGACAGTAAAGTTGAAATGGTGACGTC
CACTGCTGGCTTTATTGAACAGCTAATAAAGATTTATTTATTGTAATACCTCACAAACGTTGTAC
CATATCCATGCACATTTAGTTGCCTGCCTGTGGCTGGTAAGGTAATGTCATGATTCATCCTCTCT
TCAGTGAGACTGAGCCTGATGTGTTAACAAATAGGTGAAGAAAGTCTTGTGCTGTATTCCCTAATC
AAAAGACTTAATATATTGAAGTAACACTTTTTTAGTAAGCAAGATACCTTTTTATTTCATTCAC
AGAATGGAATTTTTTTGTTTCATGTCTCAGATTTATTTTGTATTTCTTTTTTAACACTCTACATT
TCCCTTGTTTTTTAACTCATGCACATGTGCTCTTTGTACAGTTTTTAAAAGTGTAATAAAATCTG
ACATGTCAATGTGGCTAGTTTTATTTTTCTTGTGTTTGCATTATGTGTATGGCCTGAAGTGTTGGA
CTTGCAAAAAGGGGAAGAAAGGAATTGCGAATACATGTAAATGTCACCAGACATTTGTATTATTT
TTATCATGAAATCATGTTTTTCTCTGATTGTTCTGAAATGTTCTAAATACTCTTATTTTGAATGC
ACAAAATGACTTAAACCATTATCATATCATGTTTCCTTTGCGTTCAGCCAATTTCAATTAAATGAA
CTAAATTAAAAA

FIGURE 14

MNHLPEDMENALTGSQSSHASLRNIHSINPTQLMARIESYEGREKKGISDVRRTFCLFVTFDLLF
VTLLWIIELNVNGGIENLEKEVMQYDYYSSYFDIFLLAVFRFKVLILAYAVCRLRHWWAIALTT
AVTSAFLLAKVILSKLFSQGAFGYVLPPIISFILAWIETWFLDFKVLQPQAEENRLLIVQDASER
AALIPGGLSDGQFYSPPESEAGSEEAEEKQDSEKPLLEL

Important features of the protein:

Signal peptide:

amino acids 1-20

Transmembrane domains:

amino acids 54-72, 100-118, 130-144, 146-166

N-myristoylation sites.

amino acids 14-20, 78-84, 79-85, 202-208, 217-223

FIGURE 15

ACTCGAACGCAGTTGCTTCGGGACCCAGGACCCCTCGGGCCCGACCCGCCAGGAAAGACTGAGG
 CCGCGGCCTGCCCCGCGCGCTCCCTGCGCCGCGCGCGCTCCCGGGACAGAAGATGTGCTCCAG
 GGTCCCTCTGCTGCTGCCGCTGCTCCTGCTACTGGCCCTGGGGCTGGGGTGCAGGGCTGCCCAT
 CCGGCTGCCAGTGCCAGCCAGCCACAGACAGTCTTCTGCACTGCCCGCCAGGGGACCACGGTGCCCC
 CGAGACGTGCCACCCGACACGGTGGGGCTGTACGTCTTTGAGAACGGCATCACCATGCTCGACGC
 AGGCAGCTTTGCCGCGCTGCCGGGCTGCAGCTCCTGGACCTGTCACAGAACCAGATCGCCAGCC
 TGCCCAGCGGGGTCTTCCAGCCACTCGCCAACCTCAGCAACCTGGACCTGACGGCCAACAGGCTG
 CATGAAATCACCAATGAGACCTTCCGTGGCCTGCGGCGCTCGAGCGCTCTACCTGGGCAAGAA
 CCGCATCCGCCACATCCAGCCTGGTGCCTTCGACACGCTCGACCGCTCCTGGAGCTCAAGCTGC
 AGGACAACGAGCTGCGGGCACTGCCCCGCTGCGCCTGCCCCGCTGCTGCTGCTGGACCTCAGC
 CACAACAGCCTCCTGGCCCTGGAGCCCGGCATCCTGGACACTGCCAACCTGGAGGCGCTGCGGCT
 GGCTGGTCTGGGGCTGCAGCAGCTGGACGAGGGGCTCTTCAGCCGCTTGCGAACCTCCACGACC
 TGGATGTGTCCGACAACCAGCTGGAGCGAGTGCCACCTGTGATCCGAGGCCTCCGGGGCTGACG
 CGCTGCGGCTGGCGGCAACACCCGCTATTGCCAGCTGCGGCCGAGGACCTGGCCGGCTGGC
 TGCCCTGCAGGAGCTGGATGTGAGCAACCTAAGCCTGCAGGCCCTGCCTGGCGACCTCTCGGGCC
 TCTTCCCCCGCTGCGGCTGCTGGCAGCTGCCCGCAACCCCTTCAACTGCGTGTGCCCCCTGAGC
 TGGTTTGGCCCCCTGGGTGCGCGAGAGCCACGTACACTGGCCAGCCCTGAGGAGACGCGCTGCCA
 CTTCCCGCCCAAGAACGCTGGCCGGCTGCTCCTGGAGCTTGACTACGCCGACTTTGGCTGCCAG
 CCACCACCACACAGCCACAGTGCCACCACGAGGCCCGTGGTGCGGGAGCCACAGCCTTGTCT
 TCTAGCTTGGCTCCTACCTGGCTTAGCCCCACAGCGCCGGCCACTGAGGCCCCAGCCCCGCTC
 CACTGCCCCACCGACTGTAGGGCCTGTCCCCAGCCCCAGGACTGCCACCCTCCACCTGCCTCA
 ATGGGGGCACATGCCACCTGGGGACACGGCACCACTGGCGTGCTTGTGCCCCGAAGGCTTCACG
 GGCTGTACTGTGAGAGCCAGATGGGGCAGGGGACACGGCCAGCCCTACACCAGTCACGCCGAG
 GCCACCACGGTCCCTGACCCTGGGCATCGAGCCGGTGAGCCCCACCTCCCTGCGCGTGGGGCTGC
 AGCGCTACCTCCAGGGGAGCTCCGTGCAGCTCAGGAGCCTCCGTCTACCTATCGCAACCTATCG
 GGCCCTGATAAGCGGCTGGTGACGCTGCGACTGCCTGCCTCGCTCGCTGAGTACACGGTCACCCA
 GCTGCGGGCCCAACGCCACTTACTCCGTCTGTGTCATGCCTTTGGGGCCCGGGCGGGTGGCGGAGG
 GCGAGGAGGCCTGCGGGGAGGCCATACACCCCGAGCCGTCCACTCCAACCACGCCCCAGTCACC
 CAGGCCCGCGAGGGCAACCTGCCGCTCCTCATTGCGCCCGCCCTGGCCGCGGTGCTCCTGGCCGC
 GCTGGCTGCGGTGGGGGCAGCCTACTGTGTGCGGCGGGGCGGGCCATGGCAGCAGCGGCTCAGG
 ACAAAGGGCAGGTGGGGCCAGGGGCTGGGCCCTGGAACCTGGAGGGAGTGAAGGTCCCTTGGAG
 CCAGGCCCGAAGGCAACAGAGGGCGGTGGAGAGGCCCTGCCAGCGGGTCTGAGTGTGAGGTGCC
 ACTCATGGGCTTCCCAGGGCCTGGCCTCCAGTCACCCCTCCACGCAAAGCCCTACATCTAAGCCA
 GAGAGAGACAGGGCAGCTGGGGCCGGGCTCTCAGCCAGTGAGATGGCCAGCCCCCTCCTGCTGCC
 ACACCACGTAAGTTCTCAGTCCCAACCTCGGGGATGTGTGCAGACAGGGCTGTGTGACCACAGCT
 GGGCCCTGTTCCCTCTGGACCTCGGTCTCCTCATCTGTGAGATGCTGTGGCCCAGCTGACGAGCC
 CTAACGTCCCCAGAACCAGTGCCATGAGGACAGTGTCCGCCCTGCCCTCCGCAACGTGCAGTC
 CCTGGGCACGGCGGGCCCTGCCATGTGCTGGTAACGCATGCCTGGGTCTGCTGGGCTCTCCAC
 TCCAGGCGGACCCTGGGGGCCAGTGAAGGAAGCTCCCGGAAAGAGCAGAGGGAGAGCGGGTAGGC
 GGCTGTGTGACTCTAGTCTTGGCCCCAGGAAGCGAAGGAACAAAAGAACTGGAAAGGAAGATGC
 TTTAGGAACATGTTTTGCTTTTTTAAATATATATATTTATAAGAGATCCTTTCCCATTTATTCT
 GGGAAAGATGTTTTTCAAACCTCAGAGACAAGGACTTTGGTTTTTGTAAAGACAAACGATGATATGAA
 GGCTTTTTGTAAAGAAAAATAAAGATGAAGTGTGAAA

FIGURE 16

MCSRVP LLLPL LLL LLL LALGPGVQGCPSGCQCSQPQT V FCTARQGTTPRDPVPD TVGLYVFENGIT
MLDAGSFAGLPGLQLLDLSQNQIASLPSGVFQPLANLSNLDLTANRLHEITNETFRGLRRLERLY
LGKNRIRHIQPGAFDTLDRLLELKLQDNELRALPPLRLPRL LLLDL SHNSLLALEPGILDTANVE
ALRLAGLGLQQLD EGLFSRLRNLDLDVSDNQLERVPPVIRGLRGLTRLRLAGNTRIAQLRPEDL
AGLAALQELDVSNLSLQALPGDLSGLFPRLRL LAAARNPFNCV CPLSWFGPWWRESHVTIASPEE
TRCHFPPKNAGRLLLELDYADFGCPATTTTATVPTTRPVVREPTALSSSLAPT WLSPTAPATEAP
SPPSTAPPTVGPVPQPD CPPSTCLNGGTCHLGTRHHLACLCEGFTGLYCESQMGQGTRPSPTP
VTPRPPRSLT LGIEPVSP TSLRVGLQRYLQGS SVQLRSLRLTYRNLSGPDKRLVT LRLPASLA EY
TVTQLRPNATYSVCMPLGPGRVPEGEEACGEAHTPPAVHSNHAPVTQAREGNLPLLIAPALAAV
LLAALAAVGAAYCVR RGRAMAAAAQDKGQVGP GAGPLELEGVKVPLEPGPKATEGGGEALPSGSE
CEVPLMGFP GPGLQSPLHAKPYI

Important features:

Signal peptide:

amino acids 1-23

Transmembrane domain:

amino acids 579-599

EGF-like domain cysteine pattern signature.

amino acids 430-442

Leucine zipper pattern.

amino acids 197-219, 269-291

N-glycosylation sites.

amino acids 101-105, 117-121, 273-277, 500-504, 528-532

Tyrosine kinase phosphorylation sites.

amino acids 124-131, 337-345

N-myristoylation sites.

amino acids 23-29, 27-33, 70-76, 142-148, 187-193, 348-354,
594-600, 640-646

FIGURE 17

GCAGCGGCGAGGCGGCGGTGGTGGCTGAGTCCGTGGTGGCAGAGGCGAAGGCGACAGCTCATGCG
GGTCCGGATAGGGCTGACGCTGCTGCTGTGTGCGGTGCTGCTGAGCTTGGCCTCGGCGTCCTCGG
ATGAAGAAGGCAGCCAGGATGAATCCTTAGATTCCAAGACTACTTTGACATCAGATGAGTCAGTA
AAGGACCATACTACTGCAGGCAGAGTAGTTGCTGGTCAAATATTTCTTGATTGAGAAGAATCTGA
ATTAGAATCCTCTATTCAAGAAGAGGAAGACAGCCTCAAGAGCCAAGAGGGGGAAAGTGTCACAG
AAGATATCAGCTTTCTAGAGTCTCCAAATCCAGAAAACAAGGACTATGAAGAGCCAAAGAAAAGTA
CGGAAACCAGCTTTGACCGCCATTGAAGGCACAGCACATGGGGAGCCCTGCCACTTCCCTTTTCT
TTTCCTAGATAAGGAGTATGATGAATGTACATCAGATGGGAGGGAAGATGGCAGACTGTGGTGTG
CTACAACCTATGACTACAAAGCAGATGAAAAGTGGGGCTTTTGTGAACTGAAGAAGAGGCTGCT
AAGAGACGGCAGATGCAGGAAGCAGAAATGATGTATCAAACCTGGAATGAAAATCCTTAATGGAAG
CAATAAGAAAAGCCAAAAAAGAGAAGCATATCGGTATCTCCAAAAGGCAGCAAGCATGAACCATA
CCAAAGCCCTGGAGAGAGTGTTCATATGCTCTTTTATTTGGTGATTACTTGCCACAGAATATCCAG
GCAGCGAGAGAGATGTTTGAGAAGCTGACTGAGGAAGGCTCTCCCAAGGGACAGACTGCTCTTGG
CTTTCTGTATGCCTCTGGACTTGGTGTTAATTCAAGTCAGGCAAAGGCTCTTGATATATTACAT
TTGGAGCTCTTGGGGGCAATCTAATAGCCACATGGTTTTTGGTAAGTAGACTTTAGTGGAAGGCT
AATAATATTAACATCAGAAGAATTTGTGGTTTATAGCGGCCACAACCTTTTTCAGCTTTCATGATC
CAGATTTGCTTGTATTAAGACCAAATATTCAGTTGAACTTCCTTCAAATTCTTGTTAATGGATAT
AACACATGGAATCTACATGTAAATGAAAGTTGGTGGAGTCCACAATTTTTCTTTAAATGATTAG
TTTGGCTGATTGCCCCATAAAAAGAGAGATCTGATAAATGGCTCTTTTTAAATTTTCTCTGAGTTG
GAATTGTCAGAATCATTTTTTACATTAGATTATCATAATTTTAAAAATTTTCTTTAGTTTTTCA
AAATTTTGTAATGGTGGCTATAGAAAAACAACATGAAATATTATACAATATTTTGCAACAATGC
CCTAAGAATTGTTAAATTCATGGAGTTATTTGTGCAGAATGACTCCAGAGAGCTCTACTTTCTG
TTTTTTACTTTTCATGATTGGCTGTCTTCCCATTATTCTGGTCATTTATTGCTAGTGACACTGT
GCCTGCTTCCAGTAGTCTCATTTTCCCTATTTTGCTAATTTGTTACTTTTTCTTTGCTAATTTGG
AAGATTAACTCATTTTAAATAAAATTATGTCTAAGATTAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AA

FIGURE 18

MRVRIGLTLLLCVLLSLASASSDEEGSQDES LDSKTTLTSDSVKDHTTAGRVVAGQIFLDSESESEL
ESSIQEEEDSLKSQEGESVTEDISFLESPNPENKDYEEPKKVRKPALTAIEGTAHGEPCHFPPFLFLDK
EYDECTSDGREDGRLWCATTYDYKADEKWGFCETEEEEAAKRRQMQEAEEMMYQTGMKILNGSNKKSQKR
EAYRYLQKAASMNHTKALERVSYALLFGDYLPQNIQAAREMFEEKLTEEGSPKGQTALGFLYASGLGVN
SSQAKALVYYTFGALGGNLI AHMVLVSRL

Important features:

Signal peptide:

amino acids 1-21

N-glycosylation sites.

amino acids 195-199, 217-221, 272-276

Tyrosine kinase phosphorylation site.

amino acids 220-228

N-myristoylation sites.

amino acids 120-126, 253-259, 268-274, 270-274, 285-291, 289-295

Glycosaminoglycan attachment site.

amino acids 267-271

Microbodies C-terminal targeting signal.

amino acids 299-303

Type II fibronectin collagen-binding domain protein.

amino acids 127-169

Fructose-bisphosphate aldolase class-II protein.

amino acids 101-119

FIGURE 19

AATTCAGATTTTAAGCCCATTCTGCAGTGGAATTTTCATGAACTAGCAAGAGGACACCATCTTCTT
GTATTATACAAGAAAGGAGTGTACCTATCACACACAGGGGGAAAAAATGCTCTTTTGGGTGCTAGG
CCTCCTAATCCTCTGTGGTTTTCTGTGGACTCGTAAAGGAAAACTAAAGATTGAAGACATCACTG
ATAAGTACATTTTTATCACTGGATGTGACTCGGGCTTTGGAACTTGGCAGCCAGAAGCTTTTGAT
AAAAAGGGATTTTCATGTAATCGCTGCCTGTCTGACTGAATCAGGATCAACAGCTTTAAAGGCAGA
AACCTCAGAGAGACTTCGTACTGTGCTTCTGGATGTGACCGACCCAGAGAATGTCAAGAGGACTG
CCCAGTGGGTGAAGAACCAAGTTGGGGAGAAAGGTCTCTGGGGTCTGATCAATAATGCTGGTGTT
CCCGGCGTGCTGGCTCCCACTGACTGGCTGACACTAGAGGACTACAGAGAACCTATTGAAGTGAA
CCTGTTTGGACTCATCAGTGTGACACTAAATATGCTTCCTTTGGTCAAGAAAGCTCAAGGGAGAG
TTATTAATGTCTCCAGTGTGGAGGTCGCCTTGCAATCGTTGGAGGGGGCTATACTCCATCCAAA
TATGCAGTGGAAGGTTTCAATGACAGCTTAAGACGGGACATGAAAGCTTTTGGTGTGCACGTCTC
ATGCATTGAACCAGGATTGTTCAAAACAACTTGGCAGATCCAGTAAAGGTAATTGAAAAAAAC
TCGCCATTTGGGAGCAGCTGTCTCCAGACATCAAACAACAATATGGAGAAGGTTACATTGAAAAA
AGTCTAGACAACTGAAAGGCAATAAATCCTATGTGAACATGGACCTCTCTCCGGTGGTAGAGTG
CATGGACCACGCTCTAACAAGTCTCTTCCCTAAGACTCATTATGCCGCTGGAAAAGATGCCAAAA
TTTTCTGGATACCTCTGTCTCACATGCCAGCAGCTTTGCAAGACTTTTTATTGTTGAAACAGAAA
GCAGAGCTGGCTAATCCCAAGGCAGTGTGACTCAGCTAACCACAAATGTCTCCTCCAGGCTATGA
AATTGGCCGATTTCAAGAACACATCTCCTTTTCAACCCCATTCCTTATCTGCTCCAACCTGGACT
CATTTAGATCGTGCTTATTTGGATTGCAAAGGGAGTCCCACCATCGCTGGTGGTATCCCAGGGT
CCCTGCTCAAGTTTTCTTTGAAAAGGAGGGCTGGAATGGTACATCACATAGGCAAGTCCTGCCCT
GTATTTAGGCTTTGCCTGCTTGGTGTGATGTAAGGGAAATTGAAAGACTTGCCCATTCAAAATGA
TCTTTACCGTGGCCTGCCCCATGCTTATGGTCCCCAGCATTTACAGTAACTTGTGAATGTTAAGT
ATCATCTCTTATCTAAATATTAAAAGATAAGTCAACCCAAAAAAAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAA

FIGURE 20

MLFWVLGLLILCGFLWTRKGLKIEDITDKYIFITGCDSGFGNLAARTFDKKG FHVIAACLTESG
STALKAETSERLRTVLLDVPENVKRTAQWVKNOVGEKGLWGLINNAGVPGVLAPTDWLTLEDY
REPIEVNLFGLISVTLNMLPLVKKAQGRVINVSSVGGRLAIVGGGYTPSKYAVEGFND SLRRDMK
AFGVHVS CIEPGLFKTNLADPVKVIEKKLAIWEQLSPDIKQQYGE GYIEKSLDKLKGNKSYVNMD
LSPVVECMDHALTSLEFPKTHYAAGKDAKIFWIPLSHMPAALQDFLLLKQKAELANPKAV

Important features of the protein:

Signal peptide:

amino acids 1-17

Transmembrane domain:

amino acids 136-152

N-glycosylation sites.

amino acids 161-163, 187-190 and 253-256

Glycosaminoglycan attachment site.

amino acids 39-42

N-myristoylation sites.

amino acids 36-41, 42-47, 108-113, 166-171, 198-203 and 207-212

FIGURE 21

CTGAGGCGGCGGTAGCATGGAGGGGGAGAGTACGTCGGCGGTGCTCTCGGGCTTTGTGCTCGGCG
 CACTCGCTTTCCAGCACCTCAACACGGACTCGGACACGGAAGGTTTTCTTCTTGGGGAAGTAAAA
 GGTGAAGCCAAGAACAGCATTACTGATTCCCAAATGGATGATGTTGAAGTTGTTTATACAATTGA
 CATTTCAGAAATATATTCCATGCTATCAGCTTTTTAGCTTTTATAATTCTTCAGGCGAAGTAAATG
 AGCAAGCACTGAAGAAAATATTATCAAATGTCAAAAAGAATGTGGTAGGTTGGTACAAATTCGGT
 CGTCATTTCAGATCAGATCATGACGTTTAGAGAGAGGCTGCTTCACAAAACTTGCAGGAGCATTT
 TTCAAACCAAGACCTTGTTTTTCTGCTATTAAACACCAAGTATAATAACAGAAAGCTGCTCTACTC
 ATCGACTGGAACATTCCTTATATAAACCTCAAAAAGGACTTTTTTCACAGGGTACCTTTAGTGGTT
 GCCAATCTGGGCATGTCTGAACAACCTGGGTTATAAACTGTATCAGGTTCCCTGTATGTCCACTGG
 TTTTAGCCGAGCAGTACAAACACACAGCTCTAAATTTTTTGAAGAAGATGGATCCTTAAAGGAGG
 TACATAAGATAAATGAAATGTATGCTTCATTACAAGAGGAATTAAAGAGTATATGCAAAAAAGTG
 GAAGACAGTGAACAAGCAGTAGATAAACTAGTAAAGGATGTAAACAGATTAAAACGAGAAATTGA
 GAAAAGGAGAGGAGCACAGATTCAGGCAGCAAGAGAGAAGAACATCCAAAAAGACCCTCAGGAGA
 ACATTTTTCTTTGTGTCAGGCATTACGGACCTTTTTTCCAAATCTGAATTTCTTCATTCATGTGTT
 ATGTCTTTAAAAAATAGACATGTTTCTAAAAGTAGCTGTAACATAACCACCATCTCGATGTAGT
 AGACAATCTGACCTTAATGGTAGAACACACTGACATTCCTGAAGCTAGTCCAGCTAGTACACCAC
 AAATCATTAAGCATAAAGCCTTAGACTTAGATGACAGATGGCAATTCAAGAGATCTCGGTTGTTA
 GATACACAAGACAAACGATCTAAAGCAAATACTGGTAGTAGTAACCAAGATAAAGCATCCAAAAT
 GAGCAGCCCAGAAACAGATGAAGAAATTGAAAAGATGAAGGGTTTTGGTGAATATTCACGGTCTC
 CTACATTTTTGATCCTTTTAACCTTACAAGGAGATTTTTTTATTTGGCTGATGGGTAAAGCCAAAC
 ATTTCTATTGTTTTTACTATGTTGAGCTACTTGCAGTAAGTTCATTTGTTTTTACTATGTTTACC
 TGTTTGCAGTAATACACAGATAACTCTTAGTGCAATTTACTTCACAAAGTACTTTTTTCAAACATCA
 GATGCTTTTTATTTCCAAACCTTTTTTTTACCTTTCACATAAGTTGTTGAGGGGAAGGCTTACACAG
 ACACATTCTTTAGAATTGGAAAAGTGAGACCAGGCACAGTGGCTCACACCTGTAATCCCAGCACT
 TAGGGAAGACAAGTCAGGAGGATTGATTGAAGCTAGGAGTTAGAGACCAGCCTGGGCAACGTATT
 GAGACCATGTCTATTAAAAAATAAAATGGAAAAGCAAGAATAGCCTTATTTTCAAATATGGAAA
 GAAATTTATATGAAAATTTATCTGAGTCATTAAATTTCTCCTTAAGTGATACTTTTTTGAAGTA
 CATTATGGCTAGAGTTGCCAGATAAAATGCTGGATATCATGCAATAAATTTGCAAAACATCATCT
 AAAATTTAAAAAAAAAAAAAAAAAAAAA

FIGURE 22

MEGESTSAVLSG FVLGALAFQHLNTDS DTEGFLLG EVKGEAKNSITDSQMDDVEVVYTIDIQKYI
PCYQLFSFYNSSGEVNEQALKKILSNVKKNVVGWYKFRRHSDQIMTFRERLLHKNLQEHFSNQDL
VFLLLTPSIITESCSTHRLEHS LYKPKGLFHRVPLVVANLGMSEQLGYKTVSGSCMSTGFSRAV
QTHSSKFFEE DGSLKEVHKINEMYASLQEELKSICKKVEDSEQAVDKLVKDVNRLKREIEKRRGA
QIQAAAREKNIQKDPQENIFLCQALRTFFPNSEFLHSCVMSLKNRHVSKSSCNYNHHLDVVDNLTL
MVEHTDIPEAS PASTPQIIKHKALDLD DRWQFKRSRLD TQDKRSKANTGSSNQDKASKMSSPET
DEEIEKMKGFGEYSRSPTE

Important features:

Signal peptide:

amino acids 1-19

N-glycosylation sites.

amino acids 75-79, 322-326

N-myristoylation site.

amino acids 184-154

Growth factor and cytokines receptors family.

amino acids 134-150

FIGURE 23

GGCACAGCCGCGCGGCGGAGGGCAGAGTCAGCCGAGCCGAGTCCAGCCGGACGAGCGGACCAGCGCAGGGCAGCCCAA
GCAGCGCGCAGCGAACGCCCGCCGCCGCCACACCCCTCTGCGGTCCCCGCGGCGCCTGCCACCCCTCCCTCCTTCCCC
GCGTCCCCGCTCGCCGCGCCAGTCAGCTTGCCGGGTTCGCTGCCCGCGAAACCCCGAGGTACCAGCCCGCGCCTCT
GCTTCCCTGGGCGCGCGCCGCTCCACGCCCTCCTTCTCCCTGGCCCGCGCCTGGCACCGGGACCGTTGCCTGA
CGCGAGGGCCAGCTCTACTTTTCGCCCCGCGTCTCCTCCGCTGCTCGCCTTCCACCAACTCCAACCTCCTTCTCC
TCCAGCTCCACTCGCTAGTCCCCGACTCCGCCAGCCCTCGGCCGCTGCCGTAGCGCCGCTTCCCGTCCGGTCCCAAA
GGTGGGAACGCGTCCGCCCGCGCCGACCAATGGCACGGTTCGGCTTGCCCGCGCTTCTCTGCACCCTGGCAGTGCTC
AGCGCCGCGCTGCTGGCTGCCGAGCTCAAGTCGAAAAGTTGCTCGGAAGTGGGACGTCTTTACGTGTCCAAAGGCTTC
AACAAGAACGATGCCCCCTCCACGAGATCAACGGTGATCATTTGAAGATCTGTCCCGAGGTTCTACCTGCTGCTCT
CAAGAGATGGAGGAGAAGTACAGCTGCAAAGTAAAGATGATTTCAAAGTGTGGTCAGCGAACAGTGCAATCATTTG
CAAGCTGTCTTTGCTTCACGTTACAAGAAGTTTGATGAATCTTCAAAGAAGTACTTGAAAATGCAGAGAAATCCCTG
AATGATATGTTTGTGAAGACATATGGCCATTTATACATGCAAATTTCTGAGCTATTTAAAGATCTCTTCGTAGAGTTG
AAACGTTACTACGTGGTGGGAAATGTGAACCTGGAAGAAATGCTAAATGACTTCTGGGCTCGCCTCCTGGAGCGGATG
TTCCGCCTGGTGAACCTCCAGTACCACTTTACAGATGAGTATCTGGAATGTGTGAGCAAGTATACGGAGCAGCTGAAG
CCCTTCGGAGATGTCCCTCGCAAATTGAAGCTCCAGGTTACTCGTGCTTTTGTAGCAGCCCGTACTTTGCTCAAGGC
TTAGCGGTTGCGGGAGATGTCTGAGCAAGGTCTCCGTGGTAAACCCACAGCCAGTGATCCCATGCCCTGTTGAAG
ATGATCTACTGCTCCCACTGCCGGGTCTCGTGACTGTGAAGCCATGTTACAACTACTGCTCAAACATCATGAGAGGC
TGTTTGCCAAACCAAGGGGATCTCGATTTTGAATGGAACAATTTCATAGATGCTATGCTGATGGTGGCAGAGAGGCTA
GAGGGTCTTTCAACATTGAATCGGTATGGATCCCATCGATGTGAAGATTTCTGATGCTATTATGAACATGCAGGAT
AATAGTGTTCAGTGTCTCAGAAGGTTTCCAGGGATGTGGACCCCCAAGCCCCCTCCAGCTGGACGAATTTCTCGT
TCCATCTCTGAAAGTGCCCTTCAGTGCTCGCTTCAGACCACATCACCCCGAGGAACGCCCAACCACAGCAGCTGGCACT
AGTTTGACCGACTGGTTACTGATGTCAAGGAGAAACTGAAACAGGCCAAGAAATCTGGTCTCCCTTCCGAGCAAC
GTTTGCAACGATGAGAGGATGGCTGCAGGAAACGGCAATGAGGATGACTGTTGGAATGGGAAAGGCAAAAGCAGGTAC
CTGTTTGCACTGACAGGAAATGGATTAGCCAACCAGGGCAACAACCCAGAGGTCCAGGTTGACACCAGCAAACCAGAC
ATACTGATCCTTCGTCAAATCATGGCTCTTCGAGTGATGACCAGCAAGATGAAGAATGCATACAATGGGAACGACGTG
GACTTCTTTGATATCAGTGATGAAAGTAGTGGAGAAGGAAGTGAAGTGGCTGTGAGTATCAGCAGTGCCCTTCAGAG
TTTGACTACAATGCCACTGACCATGCTGGGAAGAGTGCCAATGAGAAAGCCGACAGTGCTGGTGTCCGTCTGGGGCA
CAGGCCTACCTCCTCACTGTCTTCTGCATCTTGTTCCTGGTTATGCAGAGAGAGTGAGATTAATTCTCAAACCTGAG
AAAAAGTGTTCATCAAAAAGTTAAAGGCACCAAGTTATCACTTTTCTACCATCCTAGTGACTTTGCTTTTTAAATGAA
TGGACAACAATGTACAGTTTTTACTATGTGGCCACTGTTTTAAGAAGTGTGACTTTGTTTTCTCATTGAGTTTTGGG
AGGAAAAGGGACTGTGCATTGAGTTGGTTCCTGCTCCCCCAAACCATGTTAAACGTGGCTAACAGTGAGGTACAGAA
CTATAGTTAGTTGTGCATTTGTGATTTTCACTCTATTATTGTTTGTATGTTTTTCTCATTTCGTTTGTGGGTT
TTTTTTTCCAACTGTGATCTCGCTTGTTCCTTACAAGCAAACAGGGTCCCTTCTTGGCACGTACATGTACGTATT
TCTGAAATATTAAATAGCTGTACAGAAGCAGGTTTTATTATCATGTTATCTTATTAAGAAAAAGCCCCAAAAGC

FIGURE 24

MARFGLPALLCTLAVLSAALLAAELKSKSCSEVRRRLYVSKGFNKNDAPLHEINGDHLKICPQGST
CCSQEMEEKYSLQSKDDFKSVVSEQCNHLQAVFASRYKKFDEFFKELLENAEKSLNDMFVKTYGH
LYMQNSELFKDLFVELKRYVVGNNLEMLNDFWARLLERMFRLVNSQYHFTDEYLECVSKYTE
QLKPFQDVPRKLKLQVTRAFVAARTFAQGLAVAGDVVSKVSVVNPTAQCTHALLKMIYCSHCRGL
VTVKPCYNYCSNIMRGCLANQGDLD FEWNNFIDAMLMVAERLEGPFNIESVMDPIDVKISDAIMN
MQDNSVQVSQKVFQCGPPKPLPAGRISRISSESAFSARFRPHHPEERPTTAAGTSLDRLVTDVK
EKLKQAKKFWSSLPSNVCNDERMAAGNGNEDDCWNGKKGKSRYLFAVTGNGLANQGNNPEVQVDT
KPDILILRQIMALRVMTSKMKNAYNGNDVDFDISDESSGEGSGSGCEYQQCPSEFDYNATDHAG
KSANEKADSAGVRPGAQAYLLTVFCILFLVMQREWR

Important features:

Signal peptide:

amino acids 1-22

ATP/GTP-binding site motif A (P-loop).

amino acids 515-524

N-glycosylation site.

amino acids 514-518

Glycosaminoglycan attachment sites.

amino acids 494-498, 498-502

N-myristoylation sites.

amino acids 63-69, 224-230, 276-282, 438-444, 497-503, 531-537

Glypicans proteins.

amino acids 54-75, 105-157, 238-280, 309-346, 423-460, 468-506

FIGURE 25

CTCGCCCTCAAATGGGAACGCTGGCCTGGGACTAAAGCATAGACCACCAGGCTGAGTATCCTGAC
CTGAGTCATCCCCAGGGATCAGGAGCCTCCAGCAGGGAACCTTCCATTATATTCTTCAAGCAACT
TACAGCTGCACCGACAGTTGCGATGAAAGTTCTAATCTCTTCCCTCCTCCTGTTGCTGCCACTAA
TGCTGATGTCCATGGTCTCTAGCAGCCTGAATCCAGGGGTGCCAGAGGCCACAGGGACCGAGGC
CAGGCTTCTAGGAGATGGCTCCAGGAAGGCGGCCAAGAATGTGAGTGCAAAGATTGGTTCCTGAG
AGCCCCGAGAAGAAAATTCATGACAGTGTCTGGGCTGCCAAAGAAGCAGTGCCCCCTGTGATCATT
TCAAGGGCAATGTGAAGAAAACAAGACACCAAAGGCACCACAGAAAGCCAAACAAGCATTCCAGA
GCCTGCCAGCAATTTCTCAAACAATGTCAGCTAAGAAGCTTTGCTCTGCCTTTGTAGGAGCTCTG
AGCGCCCCTCTTCCAATTAAACATTCTCAGCCAAGAAGACAGTGAGCACACCTACCAGACACTC
TTCTTCTCCACCTCACTCTCCCACTGTACCCACCCCTAAATCATTCAGTGCTCTCAAAAAGCA
TGTTTTTCAAGATCATTTTGTTGTTGCTCTCTCTAGTGTCTTCTTCTCTCGTCAGTCTTAGCCT
GTGCCCTCCCCTTACCCAGGCTTAGGCTTAATTACCTGAAAGATTCCAGGAAACTGTAGCTTCCT
AGCTAGTGTCATTTAACCTTAAATGCAATCAGGAAAGTAGCAAACAGAAGTCAATAAATATTTTT
AAATGTCAAAAAAAAAAAAAAAAAA

FIGURE 26

MKVLISLLLLLLPLMLMSMVSSSLNPGVARGHRDRGQASRRWLQEGGQECECKDWFLRAPRRKFM
TVSGLPKKQCPCDHFKGNVKKTRHQRHHRKPNKHSRACQQFLKQCQLRSFALPL

Important features:

Signal peptide:

amino acids 1-22

N-myristoylation sites.

amino acids 27-33, 46-52

FIGURE 27

GGACGCCAGCGCCTGCAGAGGCTGAGCAGGGAAAAAGCCAGTGCCCCAGCGGAAGCACAGCTCAG
AGCTGGTCTGCCATGGACATCCTGGTCCCACCTCCTGCAGCTGCTGGTGCTGCTTCTTACCCTGCC
CCTGCACCTCATGGCTCTGCTGGGCTGCTGGCAGCCCCTGTGCAAAAGCTACTTCCCCCTACCTGA
TGGCCGTGCTGACTCCCAAGAGCAACCGCAAGATGGAGAGCAAGAAACGGGAGCTCTTCAGCCAG
ATAAAGGGGCTTACAGGAGCCTCCGGGAAAGTGGCCCTACTGGAGCTGGGCTGCGGAACCGGAGC
CAACTTTTCAGTTCTACCCACCGGGCTGCAGGGTCACCTGCCTAGACCCAAATCCCCACTTTGAGA
AGTTCCTGACAAAGAGCATGGCTGAGAACAGGCACCTCCAATATGAGCGGTTTGTGGTGGCTCCT
GGAGAGGACATGAGACAGCTGGCTGATGGCTCCATGGATGTGGTGGTCTGCACTCTGGTGCTGTG
CTCTGTGCAGAGCCCAAGGAAGGTCTGCAGGAGGTCCGGAGAGTACTGAGACCGGGAGGTGTGC
TCTTTTTCTGGGAGCATGTGGCAGAACCATATGGAAGCTGGGCCTTCATGTGGCAGCAAGTTTTC
GAGCCCACCTGGAAACACATTGGGGATGGCTGCTGCCTCACCAGAGAGACCTGGAAGGATCTTGA
GAACGCCCAGTTCTCCGAAATCCAAATGGAACGACAGCCCCCTCCCTTGAAGTGGCTACCTGTTG
GGCCCCACATCATGGGAAAGGCTGTCAAACAATCTTTCCCAAGCTCCAAGGCACTCATTTGCTCC
TTCCCCAGCCTCCAATTAGAACAAGCCACCCACCAGCCTATCTATCTTCCACTGAGAGGGACCTTA
GCAGAATGAGAGAAGACATTGATGTACCACCTACTAGTCCCTCTCTCCCCAACCTCTGCCAGGGC
AATCTCTAACTTCAATCCCGCCTTCGACAGTGAAAAAGCTCTACTTCTACGCTGACCCAGGGAGG
AAACACTAGGACCCTGTTGTATCCTCAACTGCAAGTTTCTGGACTAGTCTCCCAACGTTTGCCTC
CCAATGTTGTCCCTTTCCCTTCGTTCCCATGGTAAAGCTCCTCTCGCTTTCTCCTGAGGCTACAC
CCATGCGTCTCTAGGAAGTGGTCACAAAAGTCATGGTGCCTGCATCCCTGCCAAGCCCCCCTGAC
CCTCTCTCCCCACTACCACCTTCTTCTGAGCTGGGGGCACCAGGGAGAATCAGAGATGCTGGGG
ATGCCAGAGCAAGACTCAAAGAGGCAGAGGTTTTGTTCTCAAATATTTTTTAATAAATAGACGAA
ACCACG

FIGURE 28

MDILVPLLQLLVLLLTLP LHL MALLGCWQPLCKSYFPYLM AVLTPKSNRKMESKKRELFSQIKGL
TGASGKVALLELGCGTGANFQFYPPGCRVTCLDPNPHFEKFLTKSMAENRHLQYERFVVAPGEDM
RQLADGSMDVVVCTLVLC SVQSPRKVLQEVRRLRPGGVLEFFWEHVAEPYGSWAFMWQQVFEPTW
KHIGDGCCLTRETWKDLENAQFSEIQMERQPPPLKWLPVGP HIMGKAVKQSFPS SKALICSFPSL
QLEQATHQPIYLP LRG T

Important features:

Signal peptide:

amino acids 1-23

Leucine zipper pattern.

amino acids 10-32

N-myristoylation sites.

amino acids 64-70, 78-84, 80-86, 91-97, 201-207

FIGURE 29

CAATGTTTGCCTATCCACCTCCCCAAGCCCCTTTACCTATGCTGCTGCTAACGCTGCTGCTGCT
GCTGCTGCTGCTTAAAGGCTCATGCTTGGAGTGGGGACTGGTCGGTGCCCAGAAAGTCTCTTCTG
CCACTGACGCCCCCATCAGGGATTGGGCCTTCTTTCCCCCTTCCTTTCTGTGTCTCCTGCCTCAT
CGGCCTGCCATGACCTGCAGCCAAGCCCAGCCCCGTGGGGAAGGGGAGAAAGTGGGGGATGGCTA
AGAAAGCTGGGAGATAGGGAACAGAAGAGGGTAGTGGGTGGGCTAGGGGGGCTGCCTTATTTAAA
GTGGTTGTTTATGATTCTTATACTAATTTATACAAAGATATTAAGGCCCTGTTCAATTAAGAAATT
GTTCCCTTCCCCTGTGTTCAATGTTTGTAAGATTGTTCTGTGTAAATATGTCTTTATAATAAAC
AGTTAAAAGCTGAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 30

MLLTLLLLLLLLLKGSCLWGLVGAQKVSSATDAPIRDWAFFPPSFLCLLPHRPAMTCSQAQPRG
EGEKVGDG

Important features:

Signal peptide:

amino acids 1-15

Growth factor and cytokines receptors family:

amino acids 3-18

FIGURE 31

GTTTGAATTCCTTCAACTATACCCACAGTCCAAAAGCAGACTCACTGTGTCCCAGGCTACCAAGTT
CCTCCAAGCAAGTCATTTCCCTTATTTAACCGATGTGTCCCTCAAACACCTGAGTGCTACTCCCT
ATTTGCATCTGTTTTGATAAATGATGTTGACACCCTCCACCGAATTCTAAGTGGAATCATGTCGG
GAAGAGATACAATCCTTGGCCTGTGTATCCTCGCATTAGCCTTGTCTTTGGCCATGATGTTTACC
TTCAGATTCATCACCACCCTTCTGGTTTACATTTTCATTTTATTGGTTATTTTGGGATTGTTGTT
TGTCTGCGGTGTTTTATGGTGGCTGTATTATGACTATACCAACGACCTCAGCATAGAATTGGACA
CAGAAAGGGAAAATATGAAGTGCCTGCTGGGGTTTGCTATCGTATCCACAGGCATCACGGCAGTG
CTGCTCGTCTTGATTTTTGTTCTCAGAAAGAGAATAAAATTGACAGTTGAGCTTTTCCAAATCAC
AAATAAAGCCATCAGCAGTGCTCCCTTCCTGCTGTTCCAGCCACTGTGGACATTTGCCATCCTCA
TTTTCTTCTGGGTCCCTCTGGGTGGCTGTGCTGCTGAGCCTGGGAACTGCAGGAGCTGCCCAGGTT
ATGGAAGGCGGCCAAGTGGAAATATAAGCCCCCTTCGGGCATTTCGGTACATGTGGTCTGACCATTT
AATTGGCCTCATCTGGACTAGTGAATTCATCCTTGCGTGCCAGCAAATGACTATAGCTGGGGCAG
TGGTTACTTGTTATTTCAACAGAAGTAAAAATGATCCTCCTGATCATCCCATCCTTTCGTCTCTC
TCCATTCTCTTCTTCTACCATCAAGGAACCGTTGTGAAAGGGTCATTTTTAATCTCTGTGGTGAG
GATTCCGAGAATCATTGTCATGTACATGCAAAACGCACTGAAAGAACAGCAGCATGGTGCATTGT
CCAGGTACCTGTTCCGATGCTGCTACTGCTGTTTCTGGTGTCTTGACAAATACCTGCTCCATCTC
AACCAGAATGCATATACTACAACCTGCTATTAATGGGACAGATTTCTGTACATCAGCAAAAGATGC
ATTCAAAATCTTGTCCAAGAACTCAAGTCACTTTACATCTATTAAGTCTTTGGAGACTTCATAA
TTTTTCTAGGAAAGGTGTTAGTGGTGTGTTTCACTGTTTTTGGAGGACTCATGGCTTTTAACTAC
AATCGGGCATTCCAGGTGTGGGCAGTCCCTCTGTTATTGGTAGCTTTTTTGCCTACTTAGTAGC
CCATAGTTTTTTATCTGTGTTTGAAACTGTGCTGGATGCACTTTTCTGTGTTTTGCTGTTGATC
TGGAACAAATGATGGATCGTCAGAAAAGCCCTACTTTATGGATCAAGAATTTCTGAGTTTCGTA
AAAAGGAGCAACAAATTAAACAATGCAAGGGCACAGCAGGACAAGCACTCATTAAGGAATGAGGA
GGGAACAGAACTCCAGGCCATTGTGAGATAGATACCCATTTAGGTATCTGTACCTGGAAAACATT
TCCTTCTAAGAGCCATTTACAGAATAGAAGATGAGACCCTAGAGAAAAGTTAGTGAATTTTTTT
TTAAAAGACCTAATAAACCTATTCTTCCTCAAAA

FIGURE 32

MSGRDTILGLCILALALSLAMMFTFRFITLLVHIFISLVILGLLFVCGVLWWLYDYDTNDSIE
LDTERENMKCVLGFAIVSTGITAVLLVLI FVLRKRIKLTVELEQITNKAISSAPFLLFQPLWTEA
ILIFFWVLWVAVLLSLGTAGAAQVMEGGQVEYKPLSGIRYMWSYHLIGLIWTSEFILACQQMTIA
GAVVTCYFNRSKNDPPDHPILSSLSILFFYHQGTVVKGSFLISVVRIPRIIVMYMQNALKEQQHG
ALSRYLFRCCYCCFWCLDKYLLHLNQAYTTTAINGTDECTSAKDAFKILSKNSSHFTSINCFGD
FIIFLGKVLVVCFTVFGGLMAFNYNRAFQVWAVPLLLVAFFAYLVAHSFLSVFETVLDALFLCFA
VDLETNDGSSEKPYFMDQEFLSFVKRSNKLNNARAQQDKHSLRNEEGTELQAIVR

Important features:

Signal peptide:

amino acids 1-20

Putative transmembrane domains:

amino acids 35-54, 75-97, 126-146, 185-204, 333-350, 352-371

N-glycosylation sites.

amino acids 204-208, 295-299, 313-317

N-myristoylation sites.

amino acids 147-153, 178-184, 196-202, 296-275, 342-348

FIGURE 33

[illegible]

FIGURE 34

MRTVVLTMKASVIEMFLVLLVTGVHSNKETAKKIKRPKFTVPQINCDVKAGKIIDPEFIVKCPAG
CQDPKYHVGTDVYASYSSVCGAAVHSGVLDNSGGKILVRKVAGQSGYKGSYSNGVQSLSLPRWR
ESFIVLESKPKKGVITYPSALTYSSSKSPAAQAGETTKAYQRPPIPGTTAQPVTLMQLLAVTVAVA
TPTTLPRPSPSAASTTIPRPQSVGHRSEQEMDLWSTATYTSSQNRPRADPGIQRQDPSGAAFQKP
VGADVSLGLVPKEELSTQSLEPVSLGDPNCKIDLSFLIDGSTSIGKRRFRIQKQLLADVAQALDI
GPAGPLMGVVQYGDNPATHTNLKTHNSRDLKTAIEKITQRGGLSNVGRAISFVTKNFFSKANGN
RSGAPNVVVVMVDGWPTDKVEEASRLARESGINIFFITIEGAAENKQYVVEPNFANKAVCRTNG
FYSLHVQSWFGLHKTLOPLVKRVCDTDRILACSKTCLNSADIGFVIDGSSSVGTGNFRTVLQFVTN
LTKEFEISDTRIGAVQYTYEQRLFEFGFDKYSSKPDILNAIKRVGYWGGTSTGAAINFALQQL
FKKSKPNKRKLMILITDGRSYDDVRIPAMAAHLKGVITYAIGVAAQEELEVIATHPARDHSFF
VDEFDNLHQYVPRIIQNICTEFNQPRN

Important features:

Signal peptide:

amino acids 1-26

Transmembrane domain:

amino acids 181-200

N-glycosylation sites.

amino acids 390-394, 520-524

N-myristoylation sites.

amino acids 23-29, 93-99, 115-121, 262-268, 367-373, 389-395,
431-437, 466-472, 509-515, 570-576, 571-577, 575-581, 627-633

Amidation site.

amino acids 304-308

FIGURE 35

CCGAGCACAGGAGATTGCCTGCGTTTAGGAGGTGGCTGCGTTGTGGGAAAAGCTATCAAGGAAGAAATTGC
CAAACCATGTCTTTTTTCTGTTTTTCAGAGTAGTTCACAACAGATCTGAGTGTTTTAATTAAGCATGGAAT
ACAGAAAACAACAAAAAACTTAAGCTTTAATTTTCATCTGGAATTCCACAGTTTTCTTAGCTCCCTGGACCC
GGTTGACCTGTTGGCTCTTCCCGCTGGCTGCTCTATCACGTGGTGCTCTCCGACTACTCACCCCGAGTGTA
AAGAACCTTCGGCTCGCGTGCTTCTGAGCTGCTGTGGATGGCCCTCGGCTCTCTGGACTGTCTTCCGAGTA
GGATGTCACCTGAGATCCCTCAAATGGAGCCTCCTGCTGCTGTCACCTCCTGAGTTTTCTTTGTGATGTGGTAC
CTCAGCCTTCCCCACTACAATGTGATAGAACGCGTGAACCTGGATGTACTTCTATGAGTATGAGCCGATTTA
CAGACAAGACTTTCACTTCACACTTCGAGAGCATTCAAACCTGCTCTCATCAAAATCCATTTCTGGTCATTCT
TGGTGACCTCCCACCCCTTCAGATGTGAAAGCCAGGCAGGCCATTAGAGTTACTTGGGGTGAAAAAAGTCT
TGGTGGGGATATGAGETTCTTACATTTTTCTTATTAGGCCAAGAGGCTGAAAAGGAAGACAAAAATGTTGGC
ATTGTCCTTAGAGGATGAACACCTTCTTTATGGTGACATAATCCGACAAGATTTTTTAGACACATATAATA
ACCTGACCTTGAAAACCATTTATGGCATTTCAGGTGGGTAACCTGAGTTTTGCCCAATGCCAAGTACGTAATG
AAGACAGACACTGATGTTTTTCATCAATACTGGCAATTAGTGAAGTATCTTTTAAACCTAAACCACTCAGA
GAAGTTTTTCACAGGTTATCCTCTAATTGATAATTATTCCTATAGAGGATTTTACCAAAAAACCCATATTT
CTTACCAGGAGTATCCTTTCAAGGTGTTCCCTCCATACTGCAGTGGGTGGGTATATAATGTCCAGAGAT
TTGGTGCCAAGGATCTATGAAATGATGGGTACGTAAACCCATCAAGTTTGAAGATGTTTATGTCCGGAT
CTGTTTGAATTTATTAAAAGTGAACATTCATATTCCAGAAGACACAAATCTTTCTTTCTATATAGAATCC
ATTTGGATGTCTGTCAACTGAGACGTGTGATTGCAGCCCATGGCTTTTCTTCCAAGGAGATCATCACTTTT
TGGCAGGTCTAGCTAAGGAACACCACATGCCATTATTAACCTTCACATTCTACAAAAAGCCTAGAAGGACAG
GATACCTTGTGGAAAGTGTTAAATAAAGTAGGTACTGTGAAAAATTCATGGGGAGGTCAGTGTGCTGGCTT
ACACTGAACTGAAACTCATGAAAAACCCAGACTGGAGACTGGAGGGTTACACTTGTGATTTATTAGTCAGG
CCCTTCAAAGATGATATGTGGAGGAATTAATATAAAGGAATTGGAGGTTTTTGCTAAAGAAATTAATAGG
ACCAACAATTTGGACATGTCATTCTGTAGACTAGAAATTTCTTAAAAGGGTGTTACTGAGTTATAAGCTCA
CTAGGCTGTAAAAACAAAACAATGTAGAGTTTTATTATTGAACAATGTAGTCACTTGAAGGTTTTGTGTA
TATCTTATGTGGATTACCAATTTAAAAATATATGTAGTTCTGTGTCAAAAACTTCTTCACTGAAGTTATA
CTGAACAAAATTTTACCTGTTTTTGGTCATTTATAAAGTACTTCAAGATGTTGCAGTATTTACAGTTATT
ATTATTTAAAATTACTTCAACTTTGTGTTTTTAAATGTTTTGACGATTTCAATACAAGATAAAAAGGATAG
TGAATCATTCTTTACATGCAAACATTTTCCAGTTACTTAACTGATCAGTTTATTATTGATACATCACTCCA
TTAATGTAAAGTCATAGGTCAATTATTGCATATCAGTAATCTCTTGGACTTTGTAAATATTTTACTGTGGT
AATATAGAGAAGAATTAAAGCAAGAAATCTGAAAA

FIGURE 36

MASALWTVLPSRMSLRSLKWSLLLLSLLSFFVMWYLSLPHYNVIERVNWMYFYEYEPYRQDFHF
TLREHSNCSEQNPFLVILVTSHPSDVKARQAIRVTWGEKKSWWGYEVLTFLLGQEAEKEDKMLA
LSLEDEHLLYGDIIRQDFLDTYNNLTLKTIMAFRWVTEFCPNAKYVMKTDTDVFINTGNLVKYL
NLNHSEKFFTGYPLIDNYSYRGFYQKTHISYQYEPFKVFPPYCSGLGYIMSRDLVPRIYEMMGHV
KPIKFEDVYVGICLNLLKVNHIPEDTNLFYRIHLDVCQLRRVIAAHGFSSKEIITFWQVMLR
NTTCHY

Important features:

Type II transmembrane domain:

amino acids 20-39

N-glycosylation sites.

amino acids 72-76, 154-158, 198-202, 212-216, 326-330

Glycosaminoglycan attachment site.

amino acids 239-243

Ly-6 / u-PAR domain proteins.

amino acids 23-37

N-myristoylation site.

amino acids 271-277

FIGURE 37

CGCTCGGGCACCAGCCGCGGCAAGGATGGAGCTGGGTTGCTGGACGAGTTGGGGCTCACTTTTCTTCAGCTCCTTCATC
TCGTCCTTGCCAAGAGAGTACACAGTCATTAATGAAGCCTGCCCTGGAGCAGAGTGAATATCATGTGTCGGGAGTGCTGTG
AATATGATCAGATTGAGTGCCTCTGCCCCGAAAGAGGGAAGTCGTGGGTTATACCATCCCTTGCTGCAGGAATGAGGAGAA
TGAGTGTGACTCCTGCCTGATCCACCCAGGTTGTACCATCTTTGAAAACGCAAGAGCTGCCGAAATGGCTCATGGGGGGGT
ACCTTGGATGACTTCTATGTGAAGGGGTTCTACTGTGCAGAGTGCCGAGCAGGCTGGTACGGAGGAGACTGCATGCGATGTG
GCCAGGTTCTGCGAGCCCCAAAGGTCAGATTTTGTGGAAAGCTATCCCTAAATGCTCACTGTGAATGGACCATTTCATGC
TAAACCTGGGTTTGTATCCAACCTAAGATTTGTATGTTGAGTCTGGAGTTTGACTACATGTGCCAGTATGACTATGTTGAG
GTTTCGTGATGGAGACAACCGCGATGGCCAGATCATCAAGCGTGTCTGTGGCAACGAGCGGCCAGCTCCTATCCAGAGCATAG
GATCCTCACTCCACGTCCTCTTCCACTCCGATGGCTCCAAGAATTTGACGGTTTCCATGCCATTTATGAGGAGATCACAGC
ATGCTCCTCATCCCTTGTTCATGACGGCACGTGCGTCTTGTACAAGGCTGGATCTTACAAGTGTGCCTGCTTGGCAGGC
TATAC TGGGCAGCGCTGTGAAAATCTCCTTGAAGAAAGAACTGCTCAGACCC TGGGGGCCAGTCAATGGGTACCAGAAAA
TAACAGGGGGCCCTGGGCTTATCAACGGACGCCATGCTAAATTTGGCACCCTGGTGTCTTTCTTTTGTAAACAACCTCCTATGT
TCTTAGTGGCAATGAGAAAAGAACTTGCCAGCAGAATGGAGAGTGGTCAGGGAAACAGCCCCATCTGCATAAAAGCCTGCCGA
GAACCAAAGATTTAGACCTGGTGAGAAGGAGAGTTCTTCCGATGCAGGTTCAAGGGAGACACCATTACACCAGCTAT
ACTCAGCGGCCCTCAGCAAGCAGAACTGCAGAGTGCCCCATCCAAGAAGCCAGCCCTTCCCTTTGGAGATCTGCCCATGGG
ATACCAACATCTGCATACCCAGCTCCAGTATGAGTGCATCTCACCTTCTACCGCGCCTGGGCAGCAGCAGGAGGACATGT
CTGAGGACTGGGAAGTGGAGTGGGCGGGCACCATCTGCATCCCTATCTGCGGGAATTTGAGAATCACTGTCTCAAAGA
CCCAAGGGTTGCGCTGGCCGTGGCAGGCAGCCATCTACAGGAGGACCAGCGGGGTGCATGACGGCAGCTACACAAGGGAGC
GTGGTTCTTAGTCTGCAGCGGTGCCCTGGTGAATGAGCGCACTGTGGTGGTGGCTGCCACTGTGTTACTGACCTGGGGAG
GTCACCATGATCAAGACAGCAGACCTGAAGTTGTTTGGGGAAATCTACCGGGATGATGACCGGGATGAGAAGACCATCC
AGAGCCTACAGATTTCTGCTATCATCTCTGCATCCCACTATGACCCCATCTGCTTGATGCTGACATCGCCATCCTGAAGCT
CCTAGACAAGGCCCGTATCAGCACCCGAGTCCAGCCCATCTGCCTCGCTGCCAGTCCGGGATCTCAGCACTTCTTCCAGGAG
TCCCACATCACTGTGGCTGGCTGGAATGTCTGGCAGACGTGAGGAGCCCTGGCTTCAAGAACGACACACTGCGCTCTGGGG
TGGTCAGTGTGGTGGACTCGCTGCTGTGTGAGGAGCAGCATGAGGACCATGGCATCCAGTGAGTGTCACTGATAACATGTT
CTGTGCCAGCTGGGAACCCACTGCCCCCTTCTGATATCTGCATGCAGAGACAGGAGGCATCGCGGCTGTGTCTTCCCGGA
CGAGCATCTCCTGAGCCACGCTGGCATCTGATGGGACTGGTCAGCTGGAGCTATGATAAAACATGCAGCCACAGGCTCTCCA
CTGCCCTTACCAAGGTGCTGCCTTTTAAAGACTGGATTGAAAGAAATATGAAATGAACCATGCTCATGCACTCCTTGAGAAG
TGTTTCTGTATATCCGTCTGTACGTGTGTATTGCGTGAAGCAGTGTGGGCTGAAGTGTGATTGGCTGTGAACTTGCT
GTGCCAGGGCTTCTGACTTCAGGGACAAACTCAGTGAAGGGTGAGTAGACCTCCATTGCTGGTAGGCTGATGCCGCTCCA
CTACTAGGACAGCCAATTGGAAGATGCCAGGGCTTGCAAGAAGTAAGTTTCTTCAAAGAAGACCATATACAAAACCTCTCCA
CTCCACTGACCTGGTGGTCTTCCCCAACTTTTCAGTTATACGAATGCCATCAGCTTGACCAGGAAGATCTGGGCTTCATGAG
GCCCCTTTGGAGCTCTCAAGTTCTAGAGAGCTGCCGTGGGACAGCCAGGGCAGCAGAGCTGGGATGTGGTGCATGCCTT
TGTGTACATGGCCACAGTACAGTCTGGTCCTTTTCCCTTCCCCATCTCTGTACACATTTTAAATAAAATAAGGGTTGGCTTCT
GAACTACAAAAA
AAAAA

FIGURE 38

MELGCWTQLGLTFLQLLLISSLPREYTVINEACPGAENIMCRECCEYDQIECVCPGKREVVGYT
IPCCRNEENECDSCLIHPGCTIFENCKSCRNGSWGGLDDFYVKGIFYCAECRAGWYGGDCMRCGQ
VLRAPKGQILLESYPLNAHCEWTIHAKPGFVIQLRFVMLSLEFDYMCQYDYVEVRDGDNRDGQII
KRVCGNERPAPIQSIGSSSLHVLFHSDGSKNFDGFHAIYEEITACSSSPCFHDGTCVLDKAGSYKC
ACLAGYTGQRCENLLEERNCSDFGGPVNGYQKITGGPGLINGRHAKIGTVVSFFCNNSYVLSGNE
KRTCQONGEWSGKQPICIKACREPKISDLVRRRVLPQVQSRETPLHQLYSAAFSKQKLQSAPTK
KPALPFGDLPMGYQHLHTQLQYECISPFYRRLGSSRRTCLRTGKWSGRAPSCIPICGKIENITAP
KTQGLRWPWQAAIYRRTSGVHDGSLHKGAWFLVCSGALVNERTVVVAACHCVTDLGKVTMIKTADL
KVVLGKFYRDDDRDEKTIQSLQISAILHPNYDPILLDADIAILKLLDKARISTRVQPICLAASR
DLSTSFQESHITVAGWNVLADVRSFGFKNDTLRSGVSVVDSLLCEEQHEDHGIPVSVTDNMFCA
SWEPTAPSDICTAETGGIAAVSFPGRASPEPRWHLMLGLVSWSYDKTCSHRELTAFKVLFPFKDWI
ERNMK

Important features of the protein:

Signal peptide:

amino acids 1-23

EGF-like domain cysteine pattern signature.

amino acids 260-272

N-glycosylation sites.

amino acids 96-100, 279-283, 316-320, 451-455, 614-618

N-myristoylation sites.

amino acids 35-41, 97-103, 256-262, 284-290, 298-304, 308-314,
474-480, 491-497, 638-644, 666-672

Amidation site.

amino acids 56-60

Serine proteases, trypsin family.

amino acids 489-506

CUB domain proteins profile.

amino acids 150-167

FIGURE 39

GGTTCCTACATCCTCTCATCTGAGAATCAGAGAGCATAATCTTCTTACGGGCCCCGTGATTTATTAACGTGGCTTAATC
TGAAGGTTCTCAGTCAAATTTCTTTGTGATCTACTGATTGTGGGGGCATGGCAAGGTTTGCTTAAAGGAGCTTGGCTGG
TTTGGGCCCCCTGTAGCTGACAGAAGGTGGCCAGGGAGAATGCAGCACACTGCTCGGAGAATGAAGGCGCTTCTGTTGC
TGGTCTTGCCCTGGCTCAGTCTGTAACTACATTGACAATGTGGGCAACCTGCACTTCCTGTATTGAGAACTCTGTA
AAGGTGCCTCCCACTACGGCCTGACCAAAGATAGGAAGAGGCGCTCACAAGATGGCTGTCCAGACGGCTGTGCGAGCC
TCACAGCCACGGCTCCCTCCCCAGAGGTTTCTGCAGCTGCCACCATCTCCTTAATGACAGACGAGCCTGGCCCTAGACA
ACCTTGCTACGTGTCTCGGCAGAGGACGGGCAGCCAGCAATCAGCCCAGTGGACTCTGGCCGGAGCAACCGAACTA
GGGCACGGCCCTTTGAGAGATCCACTATTAGAAGCAGATCATTTAAAAAAATAAATCGAGCTTTGAGTGTTCTTCGAA
GGACAAAGAGCGGGAGTGCASTTGCCAACCATGCCGACCAGGGCAGGGAAAAATTCTGAAAAACCACTGCCCTGAAG
TCTTTCCAAGGTTGTACCACCTGATTCCAGATGGTGAAATTACCAGCATCAAGATCAATCGAGTAGATCCCACTGAAA
GCCTCTCTATTAGGCTGGTGGGAGGTAGCGAAACCCCACTGGTCCATATCATTATCCAACACATTTATCGTGATGGGG
TGATCGCCAGAGACGGCCGGCTACTGCCAGGAGACATCATTTCTAAAGGTCACGGGATGGACATCAGCAATGTCCCTC
ACAACCTACGCTGTGCGTCTCCTGCGGCAGCCCTGCCAGGTGCTGTGGCTGACTGTGATGCGTGAACAGAAGTTCCGCA
GCAGGAACAATGGACAGGCCCCGGATGCCTACAGACCCCGAGATGACAGCTTTCATGTGATTCTCAACAAAAGTAGCC
CCGAGGAGCAGCTTGGAATAAACTGGTGCGCAAGGTGGATGAGCCTGGGGTTTTTCATCTTCAATGTGCTGGATGGCG
GTGTGGCATATCGACATGCTCAGCTTGAGGAGAATGACCGTGTGTTAGCCATCAATGGACATGATCTTCGATATGGCA
GGCCAGAAAGTGCGGCTCATCTGATTGAGGCCAGTGAAAGACGTGTTACCTCGTGTGTCGCCAGGTTCCGCAGC
GGAGCCCTGACATCTTTGAGGAAGCCGGCTGGAACAGCAATGGCAGCTGGTCCCCAGGGCCAGGGGAGAGGAGCAACA
CTCCCAAGCCCTCCATCTACAATTAATTTGTGATGAGAAGGTGGTAAATATCCAAAAGACCCCGGTGAATCTCTCG
GCATGACCGTGCAGGGGGAGCATCACATAGAGAATGGGATTTGCCTATCTATGTGATCAGTGTTGAGCCCGGAGGAG
TCATAAGCAGAGATGGAAGAATAAAACAGGTGACATTTTGTGTAATGTGGATGGGGTCGAACTGACAGAGGTGAGCC
GGAGTGAGGCAGTGGCATTATTGAAAAGAACATCATCTCGATAGTACTCAAAGCTTTGGAAGTCAAAGAGTATGAGC
CCCAGGAAGACTGCAGCAGCCAGCAGCCCTGGACTCCAACCACAACATGGCCCCACCCAGTGACTGGTCCCCATCCT
GGGTCATGTGGCTGGAATTACCACGGTGCTTGTATAACTGTAAAGATATTGTATTACGAAGAAACACAGCTGGAAGTC
TGGGCTTCTGCATTGTAGGAGGTTATGAAGAATACAATGGAACAAACCTTTTTTTCATCAAATCCATTGTTGAAGGAA
CACCAGCATACAATGATGGAAGAATTAGATGTGGTGATATTCTTCTTGCTGTCAATGGTAGAAGTACATCAGGAATGA
TACATGCTTGCTTGGCAAGACTGCTGAAAGAATTAAGGAAGAATTACTCTAACTATTGTTTCTTGGCCTGGCACTT
TTTTATAGAAATCAATGATGGGTGAGAGGAAAACAGAAAAATCACAAATAGGCTAAGAAGTTGAAACACTATATTTATC
TTGTCAGTTTTTATATTTAAAGAAAGAATACATTGTAAAAATGTCAGGAAAAGTATGATCATCTAATGAAAGCCAGTT
ACACCTCAGAAAAATATGATTCCAAAAAATTAACCTACTAGTTTTTTTTTCAGTGTGGAGGATTTCTCATTACTCTAC
AACATTGTTTATATTTTTCTATTCAATAAAAAGCCCTAAACAACCTAAATGATTGATTGTATACCCCACTGAATT
CAAGCTGATTAAATTTAAATTTGGTATATGCTGAAGTCTGCCAAGGTACATTATGGCCATTTTTAATTTACAGCT
AAAATATTTTTTAAATGCATTGCTGAGAAACGTTGCTTTTCATCAAACAAGAATAAATATTTTTTCAGAAGTTAAA

FIGURE 40

MKALLLLVLPWLSPANYIDNVGNLHFLYSELCKGASHYGLTKDRKRRSQDGCPCDGCASLTATAPS
PEVSAAATISLMTDEPGLDNPAYVSSAEDGQPAISPVDSGRSNRTRARPFERSTIRSRSEFKKINR
ALSVLRRTKSGSAVANHADQGRESENTTAPEVFPRLYHLIPDGEITSIKINRVDPSESLSIRLV
GGSETPLVHIIIQHIYRDGVIARDGRLLPGDIILKVNMGDISNVPHNYAVRLLRQPCQVLWLTVM
REQKFRSRNNGQAPDAYRPRDDSFHVILNKSSPEEQLGIKLVKRVDEPGVFIENVLDGGVAYRHG
QLEENDRVLAINGHDLRYGSPESAHLIQASERRVHLVVSQRQVRQSPDIFQEAGWNSNGSWSPG
PGERSENTPKPLHPTITCHEKVNIQKDPGESLGMTVAGGASHREWDLPYVISVEPGGVISRDGR
IKTG DILLNVDGVELTEVSRSEAVALLKRTSSSIVLKALEVKEYEPQEDCSSPAALDSNHNMAP
SDWSPSWVMWLELPRCLYNCKDIVLRRNTAGSLGFCIVGGYEEYNGNKPFFIKSIVEGTPAYNDG
RIRCGDILLAVNGRSTSGMIHACLARLLKELKGRITLTIVSWPGTFL

Important features:

Signal peptide:

amino acids 1-15

N-glycosylation sites.

amino acids 108-112, 157-161, 289-293, 384-388

Tyrosine kinase phosphorylation sites.

amino acids 433-441, 492-500

N-myristoylation sites.

amino acids 51-57, 141-147, 233-239, 344-350, 423-429, 447-453,
467-473, 603-609

FIGURE 41

ACCAGGCATTGTATCTTCAGTTGTCATCAAGTTCGCAATCAGATTGGAAAAGCTCAACTTGAAGCTTT
 CTTGCCCTGCAGTGAAGCAGAGAGATAGATATTATTACGTAATAAAAAACATGGGCTTCAACCTGACT
 TTCCACCTTTCTTACAAATCCGATTACTGTTGCTGTTGACTTTGTGCCTGACAGTGGTTGGGTGGGC
 CACCAGTAACTACTTCGTGGGTGCCAATCAAGAGATTCCCTAAAGCAAAGGAGTTCATGGCTAATTTCC
 ATAAGACCCTCATTTTGGGGAAAGGGAAAACTCTGACTAATGAAGCATCCACGAAGAAGGTAGAACTT
 GACAACTGTCCTTCTGTGTCTCCTTACCTCAGAGGCCAGAGCAAGCTCATTTTCAAACCAGATCTCAC
 TTTGGAAGAGGTACAGGCAGAAAATCCCAAAGTGTCCAGAGGCCGGTATCGCCCTCAGGAATGTAAAG
 CTTTACAGAGGGTCGCCATCCTCGTTCCCCACCGGAACAGAGAGAAACACCTGATGTACCTGCTGGAA
 CATCTGCATCCCTTCTCTGCAGAGGCAGCAGCTGGATTATGGCATCTACGTCATCCACCAGGCTGAAGG
 TAAAAAGTTTAATCGAGCCAACTCTTGAATGTGGGCTATCTAGAAGCCCTCAAGGAAGAAAATTGGG
 ACTGCTTTTATATTCCACGATGTGGACCTGGTACCCGAGAATGACTTTAACCTTTACAAGTGTGAGGAG
 CATCCCAAGCATCTGGTGGTTGGCAGGAACAGCACTGGGTACAGGTTACGTTACAGTGGATATTTTGG
 GGGTGTACTGCCCTAAGCAGAGAGCAGTTTTTCAAGGTGAATGGATTCTCTAACAACCTACTGGGGAT
 GGGGAGGCCAAGACGATGACCTCAGACTCAGGGTTGAGCTCCAAAGAATGAAAATTTCCCGGCCCTG
 CCTGAAGTGGGTAAATATACAATGGTCTTCCACACTAGAGACAAAGGCAATGAGGTGAACGCAGAACC
 GATGAAGCTCTTACACCAAGTGTACGAGTCTGGAGAACAGATGGGTTGAGTAGTTGTTCTTATAAAT
 TAGTATCTGTGGAACACAATCCTTTATATATCAACATCACAGTGGATTTCTGGTTTGGTGCATGACCC
 TGGATCTTTTGGTGATGTTTGGAGAAGTGAATCTTTGTTTGAATAATTTTGGCCTAGAGACTTCAA
 ATAGTAGCACACATTAAAGAACCTGTTACAGCTCATTGTTGAGCTGAATTTTTCCTTTTGTATTTCT
 TAGCAGAGCTCCTGGTGATGTAGAGTATAAAACAGTTGTAACAAGACAGCTTTCTTAGTCATTTTGAT
 CATGAGGGTTAAATATTGTAATATGGATACTTGAAGGACTTTATATAAAAGGATGACTCAAAGGATAA
 AATGAACGCTATTTGAGGACTCTGGTTGAAGGAGATTTATTTAAATTTGAAGTAATATATTATGGGAT
 AAAAGGCCACAGGAAATAAGACTGCTGAATGTCTGAGAGAACCAGAGTTGTTCTCGTCCAAGGTAGAA
 AGGTACGAAGATACAATACTGTTATTCAATTTATCCTGTACAATCATCTGTGAAGTGGTGGTGTGAGGT
 GAGAAGGCGTCCACAAAAGAGGGGAGAAAAGGCGACGAATCAGGACACAGTGAACCTTGGGAATGAAGA
 GGTAGCAGGAGGGTGGAGTGTGGCTGCAAAGGCAGCAGTAGCTGAGCTGGTTGCAGGTGCTGATAGC
 CTTACAGGGGAGGACCTGCCCAGGTATGCCTTCCAGTGTATGCCACCAGAGAATACATTCTCTATTAGT
 TTTTAAAGAGTTTTTGTAAAATGATTTTGTACAAGTAGGATATGAATTAGCAGTTTACAAGTTTACAT
 ATTAATAATAATAATATGTCTATCAAATACCTCTGTAGTAAAATGTGAAAAAGCAAAA

FIGURE 42

MGFNLT FHL SYKFRLLLLLTLC LTVVGWATS NYFVGAIQEIPKAKEFMANFHKTLILGKGKTLTN
EASTKKVELDNCPSVSPYLRGQSKLIFKPDLTLEEVQAENPKVSRGRYRPQECKALQRVAILVPH
RNREKHLMYLLEHLHPFLQRQQLDYGIYVIHQAEKGKFNRAKLLNVGYLEALKEENWDCFI FHDV
DLVPENDEFNLYKCEEHPKHLVVGRNSTGYRLRYSGYFGGVTALSREQFFKVNGFSNNYWGWWGGED
DDLRLRVELQRMKISRPLPEVGKYTMVFHTRDKGNEVNAERMKLLHQVSRVWRDGLSSCSYKLV
SVEHNPLYINITVDFWFGA

Important features:

Signal peptide:

amino acids 1-27

N-glycosylation sites.

amino acids 4-8, 220-224, 335-339

Xylose isomerase proteins.

amino acids 191-202

FIGURE 43

[illegible]

FIGURE 44

MALSSQIWAACLLLLLLLASLTSGSVFPQQTGQLAELQPQDRAGARASWMPMFQRRRRRDTHFPI
CIFCCGCCHRSKCGMCCKT

Important features:

Signal peptide:

amino acids 1-24

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 58-59

N-myristoylation site.

amino acids 44-50

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 1-12

FIGURE 45

GTGGCTTCATTTTCAGTGGCTGACTTCCAGAGAGCAATATGGCTGGTTCCCCAACATGCCTCACCC
TCATCTATATCCTTTGGCAGCTCACAGGGTCAGCAGCCTCTGGACCCGTGAAAGAGCTGGTCGGT
TCCGTTGGTGGGGCCGTGACTTTCCCCCTGAAGTCCAAAGTAAAGCAAGTTGACTCTATTGTCTG
GACCTTCAACACAACCCCTCTTGTCACCATACAGCCAGAAGGGGGCACTATCATAGTGACCCAAA
ATCGTAATAGGGAGAGAGTAGACTTCCCAGATGGAGGCTACTCCCTGAAGCTCAGCAAAGTGAAG
AAGAATGACTCAGGGATCTACTATGTGGGGATATACAGCTCATCACTCCAGCAGCCCTCCACCCA
GGAGTACGTGCTGCATGTCTACGAGCACCTGTCAAAGCCTAAAGTCACCATGGGTCTGCAGAGCA
ATAAGAATGGCACCTGTGTGACCAATCTGACATGCTGCATGGAACATGGGGAAGAGGATGTGATT
TATACCTGGAAGGCCCTGGGGCAAGCAGCCAATGAGTCCCATAATGGGTCCATCCTCCCCATCTC
CTGGAGATGGGGAGAAAGTGATATGACCTTCATCTGCGTTGCCAGGAACCCTGTCAGCAGAACT
TCTCAAGCCCCATCCTTGCCAGGAAGCTCTGTGAAGGTGCTGCTGATGACCAGATTCCCTCCATG
GTCCTCCTGTGTCTCCTGTTGGTGGCCCTCCTGCTCAGTCTCTTTGTACTGGGGCTATTTCTTTG
GTTTCTGAAGAGAGAGAGACAAGAAGAGTACATTGAAGAGAAGAAGAGAGTGGACATTTGTCGGG
AACTCCTAACATATGCCCCATTCTGGAGAGAACACAGAGTACGACACAATCCCTCACACTAAT
AGAACAATCCTAAAGGAAGATCCAGCAAATACGGTTTACTCCACTGTGGAAATACCGAAAAAGAT
GGAAAATCCCCTCACTGCTCACGATGCCAGACACACCAAGGCTATTTGCCTATGAGAATGTTA
TCTAGACAGCAGTGCCTCCCCTAAGTCTCTGCTCA

FIGURE 46

MAGSPTCLTLIYILWQLTGSAASGPVKELVGSVGGAVTFPLKSKVKQVDSIVWTFNTTPLVTIQP
EGGTIIIVTQNRNRERVDFFPDGGYSLKLSKLKKNDSGIYYVGIYSSSLQQPSTQEYVLHVYEHLSK
PKVTMGLQSNKNGTCVTNLTCCEHGEEDVIYTWKALGQAANESHNGSILPISWRWGESDMTFIC
VARNPVSRNFSSPILARKLCEGAADDPDSSMVLLCLLLVPLLLSLFVLGLFLWFLKRERQEEYIE
EKKRVDICRETPNICPHSGENTEYDTIPHTNRTILKEDPANTVYSTVEIPKKMENPHSLLTMPDT
PRLFAYENVI

Important features:

Signal peptide:

amino acids 1-22

Transmembrane domain:

amino acids 224-250

Leucine zipper pattern.

amino acids 229-251

N-glycosylation sites.

amino acids 98-102, 142-146, 148-152, 172-176, 176-180, 204-208,
291-295

FIGURE 47

GGCTCGAGCGTTTCTGAGCCAGGGGTGACCAATGACCTGCTGCGAAGGATGGACATCCTGCAATGG
ATTGAGCCTGCTGGTTCTACTGCTGTTAGGAGTAGTTCTCAATGCGATACCTCTAATTGTCAGCT
TAGTTGAGGAAGACCAATTTTCTCAAACCCCATCTCTTGCTTTGAGTGGTGGTTCCCAGGAATT
ATAGGAGCAGGTCTGATGGCCATTCCAGCAACAACAATGTCCTTGACAGCAAGAAAAAGAGCGTG
CTGCAACAACAGAACTGGAATGTTTCTTTTCATCATTTTTTCAGTGTGATCACAGTCATTGGTGCTC
TGTATTGCATGCTGATATCCATCCAGGCTCTCTTAAAAGGTCCTCTCATGTGTAATTCTCCAAGC
AACAGTAATGCCAATTGTGAATTTTCATTGAAAAACATCAGTGACATTCATCCAGAATCCTTCAA
CTTGCAAGTGGTTTTTCAATGACTCTTGTGCACCTCCTACTGGTTTCAATAAACCCACCAGTAACG
ACACCATGGCGAGTGGCTGGAGAGCATCTAGTTTCCACTTCGATTCTGAAGAAAACAAACATAGG
CTTATCCACTTCTCAGTATTTTTAGGTCTATTGCTTGTGGAATTCTGGAGGTCCTGTTTGGGCT
CAGTCAGATAGTCATCGGTTTCCTTGGCTGTCTGTGTGGAGTCTCTAAGCGAAGAAGTCAAATTG
TGTAGTTTAATGGGAATAAAATGTAAGTATCAGTAGTTTGAAAAAAAAAAAA

FIGURE 48

MTCCEGWTS CNGFSLLVLLLLGVVLNAIPLIVSLVEEDQFSQNPISCFEWWFPGIIGAGLMAIPA
TTMSLTARKRACCNRTGMFLSSFFSVITVIGALYCM LISIQALLKGPLMCNSPSNSNANCEFSL
KNISDIHPESFNLQWFFNDSCAPPTGFNKPTSNDTMASGWRASSFHFDSEENKHRLIHFSVFLGL
LLVGILEVLFGLSQIVIGFLGCLCGVSKRRSQIV

Important features:

Transmembrane domains:

amino acids 10-31 (type II), 50-72, 87-110, 191-213

N-glycosylation sites.

amino acids 80-84, 132-136, 148-152, 163-167

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 223-227

N-myristoylation sites.

amino acids 22-28, 54-60, 83-89, 97-103, 216-222

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 207-218

TNFR/NGFR family cysteine-rich region protein.

amino acids 4-12

FIGURE 49

ATCCGTTCTCTGCGCTGCCAGCTCAGGTGAGCCCTCGCCAAGGTGACCTCGCAGGACACTGGTGA
AGGAGCAGTGAGGAACCTGCAGAGTCACACAGTTGCTGACCAATTGAGCTGTGAGCCTGGAGCAG
ATCCGTGGGCTGCAGACCCCCGCCCCAGTGCCTCTCCCCCTGCAGCCCTGCCCCCTCGAACTGTGA
CATGGGAGAGAGTGACCCTGGCCCTTCTCCTACTGGCAGGCCTGACTGCCTTGGAAGCCAATGACC
CATTTGCCAATAAAGACGATCCCTTCTACTATGACTGGAAAAACCTGCAGCTGAGCGGACTGATC
TGCGGAGGGCTCCTGGCCATTGCTGGGATCGCGGCAGTTCTGAGTGGCAAATGCAAATACAAGAG
CAGCCAGAAGCAGCACAGTCCTGTACCTGAGAAGGCCATCCCACTCATCACTCCAGGCTCTGCCA
CTACTTGCTTGAGCACAGGACTGGCCTCCAGGGATGGCCTGAAGCCTAACACTGGCCCCCAGCACC
TCCTCCCCTGGGAGGCCTTATCCTCAAGGAAGGACTTCTCTCCAAGGGCAGGCTGTTAGGCCCCT
TTCTGATCAGGAGGCTTCTTTATGAATTAACTCGCCCCACCACCCCCTCA

FIGURE 50

MERVTLALLLLAGLTALEANDPFANKDDPFYYDWKNLQLSGLICGGLLAIAAGIAAVLSGKCKYKS
SQKQHSPVPEKAIPPLITPGSATTC

Important features:

Signal peptide:

amino acids 1-16

Transmembrane domain:

amino acids 36-59

N-myristoylation sites.

amino acids 41-47, 45-51, 84-90

Extracellular proteins SCP/Tpx-1/Ag5/PR-1/Sc7.

amino acids 54-67

FIGURE 51

[illegible]

FIGURE 52

MKFQGPLACLLLLALCLGSGEAGPLQSGEESTGTNIGEALGHGLGDALSEG VGKAI GKEAGGAAGSKVS
EALGQGTREAVGTGVRQVPGFGAADALGNRVGEAAHALGNTGHEIGRQAEDVIRHGADAVRGSWQGV
GHSGAWETSGGHGIFGSQGGGLGGQGQGNPGGLGTPWVHGYPGNSAGSFGMNPQGAPWQGGNGGPPNF
GTNTQGAVAQPGYGSVRASNQNEGCTNPPPSGSGGGSSNSGGGSGSQSGSSGSGSNGDNNNGSSSSGGS
SSGSSSSGSSSGSSSGSSSGSSGSGSGSRGDSGSESSWGSSTGSSSGNHGGSGGGNGHKPGCEKPGNE
ARGSGESGIQGFRRQGVSSNMREISKEGNRLLGGSGDNYRGQSSWGSGGGDAVGGVNTVNSETSPGM
FNFDTFWKNFKSKLGFINWDAINKDQRSSRIP

Signal peptide:

amino acids 1-21

N-glycosylation site.

amino acids 265-269

Glycosaminoglycan attachment site.

amino acids 235-239, 237-241, 244-248, 255-259, 324-328, 388-392

Casein kinase II phosphorylation site.

amino acids 26-30, 109-113, 259-263, 300-304, 304-308

N-myristoylation site.

amino acids 17-23, 32-38, 42-48, 50-56, 60-66, 61-67, 64-70, 74-80,
90-96, 96-102, 130-136, 140-146, 149-155, 152-158, 155-161,
159-165, 163-169, 178-184, 190-196, 194-200, 199-205, 218-224,
236-242, 238-244, 239-245, 240-246, 245-251, 246-252, 249-252,
253-259, 256-262, 266-272, 270-276, 271-277, 275-281, 279-285,
283-289, 284-290, 287-293, 288-294, 291-297, 292-298, 295-301,
298-304, 305-311, 311-317, 315-321, 319-325, 322-328, 323-329,
325-331, 343-349, 354-360, 356-362, 374-380, 381-387, 383-389,
387-393, 389-395, 395-401

Cell attachment sequence.

amino acids 301-304

FIGURE 53

GGAGAAGAGGTTGTGTGGGACAAGCTGCTCCCGACAGAAGGATGTCGCTGCTGAGCCTGCCCTGG
CTGGGCCTCAGACCGGTGGCAATGTCCCATGGCTACTCCTGCTGCTGGTTGTGGGCTCCTGGCT
ACTCGCCCGCATCCTGGCTTGGACCTATGCCTTCTATAACAACCTGCCGCCGGCTCCAGTGTTTCC
CACAGCCCCCAAACGGAACCTGGTTTTGGGGTCACCTGGGCCTGATCACTCCTACAGAGGAGGGC
TTGAAGGACTCGACCCAGATGTCGGCCACCTATTCCCAGGGCTTTACGGTATGGCTGGGTCCCAT
CATCCCCTTCATCGTTTTATGCCACCCTGACACCATCCGGTCTATCACC AATGCCTCAGCTGCCA
TTGCACCCAAGGATAATCTCTTCATCAGGTTCTGAAGCCCTGGCTGGGAGAAGGGATACTGCTG
AGTGGCGGTGACAAGTGGAGCCGCCACCGTCGGATGCTGACGCCCGCCTTCCATTTCAACATCCT
GAAGTCCTATATAACGATCTTCAACAAGAGTGCAAACATCATGCTTGACAAGTGGCAGCACCTGG
CCTCAGAGGGCAGCAGTCGTCTGGACATGTTTGAGCACATCAGCCTCATGACCTTGGACAGTCTA
CAGAAATGCATCTTCAGCTTTGACAGCCATTGTCAGGAGAGGCCAGTGAATATATTGCCACCAT
CTTGAGCTCAGTGCCCTTGTAGAGAAAAGAAGCCAGCATATCCTCCAGCACATGGACTTTCTGT
ATTACCTCTCCCATGACGGGCGGCGCTTCCACAGGGCCTGCCGCCTGGTGCATGACTTCACAGAC
GCTGTCTATCCGGGAGCGGCGTCGCACCCTCCCCACTCAGGGTATTGATGATTTTTTCAAAGACAA
AGCCAAGTCCAAGACTTTGGATTTCAATTGATGTGCTTCTGCTGAGCAAGGATGAAGATGGGAAGG
CATTGTCAGATGAGGATATAAGAGCAGAGGCTGACACCTTCATGTTTGAGAGGCCATGACACCACG
GCCAGTGGCCTCTCCTGGGTCTGTACAACCTTGCGAGGCACCCAGAATACCAGGAGCGCTGCCG
ACAGGAGGTGCAAGAGCTTCTGAAGGACCGGATCCTAAAGAGATTGAATGGGACGACCTGGCCC
AGCTGCCCTTCCTGACCATGTGCGTGAAGGAGAGCCTGAGGTTACATCCCCAGCTCCCTTCATC
TCCCGATGCTGCACCCAGGACATTGTTCTCCAGATGGCCGAGTCATCCCCAAAGGCATTACCTG
CCTCATCGATATTATAGGGGTCCATCACAACCCAACTGTGTGGCCGGATCCTGAGGTCTACGACC
CCTTCCGCTTTGACCCAGAGAACAGCAAGGGGAGGTACCTCTGGCTTTTATTCTTTCTCCGCA
GGGCCCAGGAACCTGCATCGGGCAGGCGTTGCGCATGGCCGAGATGAAAGTGGTCTGGCGTTGAT
GCTGCTGCACTTCCGGTTCCTGCCAGACCACACTGAGCCCCGCAGGAAGCTGGAATTGATCATGC
GCGCCGAGGGCGGGCTTTGGCTGCGGGTGGAGCCCCTGAATGTAGGCTTGCATGACTTTCTGAC
CCATCCACCTGTTTTTTTTGCAGATTGTCATGAATAAAACGGTGCTGTCAAA

FIGURE 54

MSLLSLPWLGLRPVAMSPWLLLLLVGSWLLARILAWTYAFYNNCRRLQCFPQPPKRNWFWGHLG
LITPTEEGLKDSTQMSATYSQGFTVWLGPIIPFIVLCHPDTIRSITNASAAIAPKDNLFIRFLKP
WLGEGILLSGGDKWSRHRRLTPAFHFNILKSYITIFNKSANIMLDKWQHLASEGSSRLDMFEHI
SLMTLDSLQKCIFSFDSHCQERPSEYIATILELSALVEKRSQHILQHMDFLYYLSHDGRRFHRAC
RLVHDFTDVIRERRRTLPTQGIDDFKDKAKSKTLD FIDVLLLSKDEDGKALSDEDIRAEADTF
MFGGHDTTASGLSWVLYNLARHPYQERCQEVQELLKDRDPKEIEWDDLAQLPFLTMCVKESLR
LHPPAPFISRCCTQDIVLPDGRVIPKGITCLIDIIGVHHNPTVWPDPEVYDPFRFDPENSKGRSP
LAFIPFSAGPRNCIGQAFAMAEMKVVLALMLLHFRFLPDHTEPRRKLELIMRAEGGLWLRVEPLN
VGLQ

Important features:

Transmembrane domains:

amino acids 13-32 (type II), 77-102

Cytochrome P450 cysteine heme-iron ligand signature.

amino acids 461-471

N-glycosylation sites.

amino acids 112-116, 168-172

FIGURE 55

[illegible]

FIGURE 56

MGFVKQLKRMFEPTRLIATIMVLLCFALTLCSAFWWHNKGLALIFCILQSLALTWYSLSEIPFAR
DAVKKCFVCLA

Important features:

Signal peptide:

amino acids 1-33

Type II fibronectin collagen-binding domain protein.

amino acids 30-72

FIGURE 57

[illegible]

FIGURE 58

MLCLCLYVPVIGEAEQTEFQYFESKGLPAELKSIFKLSVFIPSQEFSTYRQWKQKIVQAGDKDLDG
QLDFEEFVHYLQDHEKKLRILVFKILDKKNDGRIDAQEIMQSLRDLGVKISEQQAEEKILKSMDKNG
TMTIDWNEWRDYHLLHPVENIPEIILYWKHSTIFDVGENLTVPDEFTVEERQTGMWWRHLVAGGG
AGAVSRTCTAPLDRLKVLQMQLVHASRSNNMGIVGGFTQMIREGGARSLWRGNGINVLKIAPESAIK
FMAYEQIKRLVGSDQETLRIHERLVAGSLAGAIAQSSIYPMEVLKTRMALRKTGQYSGMLDCARR
ILAREGVAAFYKGYVPNMLGIIPYAGIDLAVYETLKNAWLQHYAVNSADPGVFVLLACGTMSSTC
GQLASYPLALVRTRMQAQASIEGAPEVTMSSSLFKHILRTEGAFLYRGLAPNFMKVIPAVSISYV
VYENLKITLGVQSR

Important features:

Signal peptide:

amino acids 1-16

Putative transmembrane domains:

amino acids 284-304, 339-360, 376-394

Mitochondrial energy transfer proteins signature.

amino acids 206-215, 300-309

N-glycosylation sites.

amino acids 129-133, 169-173

Elongation Factor-hand calcium-binding protein.

amino acids 54-73, 85-104, 121-140

FIGURE 59

GGAAGGCAGCGGCAGCTCCACTCAGCCAGTACCCAGATACGCTGGGAACCTTCCCCAGCCATGGC
TTCCCTGGGGCAGATCCTCTTCTGGAGCATAATTAGCATCATCATTATTCTGGCTGGAGCAATTG
CACTCATCATTTGGCTTTGGTATTTTCAAGGAGACACTCCATCACAGTCACTACTGTGCGCTCAGCT
GGGAACATTGGGGAGGATGGAATCCTGAGCTGCACTTTTGAACCTGACATCAAACCTTTCTGATAT
CGTGATACAATGGCTGAAGGAAGGTGTTTTAGGCTTGGTCCATGAGTTCAAAGAAGGCAAAGATG
AGCTGTGCGGAGCAGGATGAAATGTTTCAAGGCCGGACAGCAGTGTGTTGCTGATCAAGTGATAGTT
GGCAATGCCTCTTTGCGGCTGAAAAACGTGCAACTCACAGATGCTGGCACCTACAAATGTTATAT
CATCACTTCTAAAGGCAAGGGGAATGCTAACCTTGAGTATAAACTGGAGCCTTCAGCATGCCGG
AAGTGAATGTGGACTATAATGCCAGCTCAGAGACCTTGCGGTGTGAGGCTCCCCGATGGTTCCCC
CAGCCCACAGTGGTCTGGGCATCCCAAGTTGACCAGGGAGCCAACTTCTCGGAAGTCTCCAATAC
CAGCTTTGAGCTGAACTCTGAGAATGTGACCATGAAGGTTGTGTCTGTGCTCTACAATGTTACGA
TCAACAACACATACTCCTGTATGATTGAAAATGACATTGCCAAAGCAACAGGGGATATCAAAGTG
ACAGAATCGGAGATCAAAGGCGGAGTCACCTACAGCTGCTAAACTCAAAGGCTTCTCTGTGTGT
CTCTTCTTTCTTTGCCATCAGCTGGGCACCTTCTGCCTCTCAGCCCTTACCTGATGCTAAAATAA
GTGCCTTGGCCACAAAAAGCATGCAAAGTCATTGTTACAACAGGGATCTACAGAACTATTTTAC
CACCAGATATGACCTAGTTTTATATTTCTGGGAGGAAATGAATTCATATCTAGAAGTCTGGAGTG
AGCAAACAAGAGCAAGAAACAAAAAGAAGCCAAAGCAGAAGGCTCCAATATGAACAAGATAAAT
CTATCTTCAAAGACATATTAGAAGTTGGGAAAATAATTCATGTGAACTAGACAAGTGTGTTAAGA
GTGATAAGTAAAATGCACGTGGAGACAAGTGCATCCCCAGATCTCAGGGACCTCCCCCTGCCTGT
CACCTGGGGAGTGAGAGGACAGGATAGTGCATGTTCTTTGTCTCTGAATTTTTAGTTATATGTGC
TGTAATGTTGCTCTGAGGAAGCCCCCTGGAAAGTCTATCCCAACATATCCACATCTTATATTCAC
AAATTAAGCTGTAGTATGTACCCTAAGACGCTGCTAATTGACTGCCACTTCGCAACTCAGGGGCG
GCTGCATTTTAGTAATGGGTCAAATGATTCACTTTTTATGATGCTTCCAAAGGTGCCTTGGCTTC
TCTTCCCAACTGACAAATGCCAAAGTTGAGAAAAATGATCATAATTTTAGCATAAACAGAGCAGT
CGGGGACACCGATTTTATAAATAAACTGAGCACCTTCTTTTTAAACAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 6o

MASLGQILFWSIISIIIIILAGAIALIIIGFGISGRHSITVTTVASAGNIGEDGILSCTFEPDIKLS
DIVIQWLKEGVLGLVHEFKEGKDELSEQDEMFRGRTAVFADQVIVGNASRLKKNVQLTDAGTYKC
YIITSKGGKGNANLEYKTAGFSMPEVNVVDYNASSETLRCEAPRWFPQPTVVWASQVDQGANFSEVS
NTSFELNSENVTMKVVSVLYNVTINNTYSCMIENDIAKATGDIKVTSESEIKRRSHLQLLNSKASL
CVSSFFAISWALLPLSPYLMLK

Important features:

Signal peptide:

amino acids 1-28

Transmembrane domain:

amino acids 258-281

N-glycosylation sites.

amino acids 112-116, 160-164, 190-194, 196-200, 205-209, 216-220,
220-224

N-myristoylation sites.

amino acids 52-58, 126-132, 188-194

FIGURE 61

TGACGTCAGAATCACCATGGCCAGCTATCCTTACCGGCAGGGCTGCCCAGGAGCTGCAGGACAAG
CACCAGGAGCCCCCTCCGGGTAGCTACTACCCTGGACCCCCCAATAGTGGAGGGCAGTATGGTAGT
GGGCTACCCCCCTGGTGGTGGTTATGGGGGTCTTGCCCCCTGGAGGGCCTTATGGACCACCAGCTGG
TGGAGGGCCCTATGGACACCCCAATCCTGGGATGTTCCCCTCTGGAACCTCCAGGAGGACCATATG
GCGGTGCAGCTCCCGGGGGCCCCCTATGGTCAGCCACCTCCAAGTTCCTACGGTGCCAGCAGCCT
GGGCTTTATGGACAGGGTGGCGCCCCCTCCCAATGTGGATCCTGAGGCCTACTCCTGGTTCCAGTC
GGTGGACTCAGATCACAGTGGCTATATCTCCATGAAGGAGCTAAAGCAGGCCCTGGTCAACTGCA
ATTGGTCTTCATTCAATGATGAGACCTGCCTCATGATGATAAACATGTTTGACAAGACCAAGTCA
GGCCGCATCGATGTCTACGGCTTCTCAGCCCTGTGGAAATTCATCCAGCAGTGAAGAACCTCTT
CCAGCAGTATGACCGGGACCGCTCGGGCTCCATTAGCTACACAGAGCTGCAGCAAGCTCTGTCCC
AAATGGGCTACAACCTGAGCCCCCAGTTCACCCAGCTTCTGGTCTCCCGCTACTGCCACGCTCT
GCCAATCCTGCCATGCAGCTTGACCGCTTCATCCAGGTGTGCACCCAGCTGCAGGTGCTGACAGA
GGCCTTCCGGGAGAAGGACACAGCTGTACAAGGCAACATCCGGCTCAGCTTCGAGGACTTCGTCA
CCATGACAGCTTCTCGGATGCTATTGACCCAACCATCTGTGGAGAGTGGAGTGCACCAGGGACCTT
TCCTGGCTTCTTAGAGTGAGAGAAGTATGTGGACATCTCTTCTTTTCCTGTCCCTCTAGAAGAAC
ATTCTCCCTTGCTTGATGCAACACTGTTCCAAAAGAGGGTGGAGAGTCTGCATCATAGCCACCA
AATAGTGAGGACCGGGGCTGAGGCCACACAGATAGGGGCCTGATGGAGGAGAGGATAGAAGTTGA
ATGTCCTGATGGCCATGAGCAGTTGAGTGGCACAGCCTGGCACCAGGAGCAGGTCTTGTAATGG
AGTTAGTGTCCAGTCAGCTGAGCTCCACCCTGATGCCAGTGGTGAGTGTTTCATCGGCCTGTTACC
GTTAGTACCTGTGTTCCCTCACCAGGCCATCCTGTCAAACGAGCCCATTTTCTCCAAAGTGGAAAT
CTGACCAAGCATGAGAGAGATCTGTCTATGGGACCAGTGGCTTGGATTCTGCCACACCCATAAAT
CCTTGTTGTGTTAACTTCTAGCTGCCTGGGGCTGGCCCTGCTCAGACAAATCTGCTCCCTGGGCAT
CTTTGGCCAGGCTTCTGCCCCCTGCAGCTGGGACCCCTCACTTGCCCTGCCATGCTCTGCTCGGCT
TCAGTCTCCAGGAGACAGTGGTCACCTCTCCCTGCCAATACTTTTTTTAATTTGCATTTTTTTTC
ATTTGGGGCCAAAAGTCCAGTGAAATTGTAAGCTTCAATAAAAGGATGAACTCTGA

FIGURE 62

MASYPYRQGCPGAAGQAPGAPPGSYYPGPPNSGGQYGSGLPPGGGYGGPAPGGPYGPPAGGGPYG
HPNPGMFPSGTPGGPYGGAAPGGPYGQPPSSYGAQQPGLYGQGGAPPNVDPEAYSWFQSVDS DH
SGYISMKELKQALVNCNWSSFNDETCLMMINMFDKTKSGRIDVYGFSALWKFIQQWKNLFQQYDR
DRSGSISYTELQQALSQMGYNLSPQFTQLLVSRYCPRSANPAMQLDRFIQVCTQLQVLTEAFREK
DTAVQGNIRLSFEDFVTMTASRML

Important features of the protein:

Signal peptide:

amino acids 1-19

N-glycosylation site.

amino acids 147-150

Casein kinase II phosphorylation sites.

amino acids 135-138, 150-153, 202-205, 271-274

N-myristoylation sites.

amino acids 9-14, 15-20, 19-24, 33-38, 34-39, 39-44, 43-48, 61-
66, 70-75, 78-83, 83-88, 87-92, 110-115

FIGURE 63

[illegible]

FIGURE 64

MQGRVAGSCAPLGLLLVLHLPGLFARSIGVVEEKVSNFGTNLPQLGQPSSTGSPNSEHPQPAL
DPRSNDLARVPLKLSVPPSDGFPPAGGSQVQRWPPSWGLPAMDSWPPEDPWQMAAAAEDRLGEA
LPEELSYLSSAAALAPGSGPLPGESSPDATGLSPEASLLHQDSESRRLPRSNLSGAGGKILSQRP
PWSLIHRVLPDHPWGTLNPSVSWGGGGPGTGWGTRPMHPHEGIWGINNQPPGTSGWNINRYPGGS
WGNINRYPGGSWGNINRYPGGSWGNIHLYPGINNPFPFPGVLRPPGSSWNIPAGFPNPPSPRLQWG

Important features of the protein:

Signal peptide:

amino acids 1-26

Casein kinase II phosphorylation sites.

amino acids 56-59, 155-158

N-myristoylation sites.

amino acids 48-53, 220-225, 221-226, 224-229, 247-252, 258-263,
259-264, 269-274, 270-275, 280-285, 281-286, 305-310

FIGURE 65

AAGGAGAGGCCACCGGGACTTCAGTGTCTCCTCCATCCCAGGAGCGCAGTGGCCACTATGGGGTC
TGGGCTGCCCCCTTGTCTCCTCTTGACCCTCCTTGGCAGCTCACATGGAACAGGGCCGGGTATGA
CTTTGCAACTGAAGCTGAAGGAGTCTTTTCTGACAAATTCTCCTATGAGTCCAGCTTCCTGGAA
TTGCTTGAAAAGCTCTGCCTCCTCCTCCATCTCCCTTCAGGGACCAGCGTCACCCTCCACCATGC
AAGATCTCAACACCATGTTGTCTGCAACACATTGACAGCCATTGAAGCCTGTGTCCTTCTTGGCCC
GGGCTTTTGGGCCGGGGATGCAGGAGGCAGGCCCCGACCCTGTCTTTCAGCAGGCCCCCACCCTC
CTGAGTGGCAATAAATAAAATTCGGTATGCTG

FIGURE 66

MGSGLPLVLLLTLLGSSHGTGPGMTLQKLKESFLTNSSYESSFLELLEKLCLLLHLPSGTSVTL
HHARSQHHVVCNT

Important features:

Signal peptide:

amino acids 1-19

N-glycosylation site.

amino acids 37-41

N-myristoylation sites.

amino acids 15-21, 19-25, 60-66

FIGURE 67

ACGGACCGAGGGTTCGAGGGAGGGACACGGACCAGGAACCTGAGCTAGGTCAAAGACGCCCCGGGC
CAGGTGCCCCGTCGCAGGTGCCCCCTGGCCGGAGATGCGGTAGGAGGGGCGAGCGCGAGAAGCCCC
TTCCTCGGCGCTGCCAACCCGCCACCCAGCCCATGGCGAACCCCGGGCTGGGGCTGCTTCTGGCG
CTGGGCCCTGCCGTTCTTGCTGGCCCGCTGGGGCCGAGCCTGGGGGCAAATACAGACCACTTCTGC
AAATGAGAATAGCACTGTTTTGCCTTCATCCACCAGCTCCAGCTCCGATGGCAACCTGCGTCCGG
AAGCCATCACTGCTATCATCGTGCTTCTTCTCCCTCTTGGCTGCCTTGCTCCTGGCTGTGGGGCTG
GCACTGTTGGTGCGGAAGCTTCGGGAGAAGCGGCAGACGGAGGGCACCTACCGGCCCAGTAGCGA
GGAGCAGTTCTCCCATGCAGCCGAGGCCCGGGCCCCCTCAGGACTCCAAGGAGACGGTGCAAGGCT
GCCTGCCCATCTAGGTCCCCCTCTCCTGCATCTGTCTCCCTTCATTGCTGTGTGACCTTGGGGAAA
GGCAGTGCCCTCTCTGGGCAGTCAGATCCACCCAGTGCTTAATAGCAGGGAAGAAGGTACTTCAA
AGACTCTGCCCCCTGAGGTCAAGAGAGGATGGGGCTATTCACTTTTATATATTTATATAAAATTAG
TAGTGAGATGTAAAAAAAAAAAAAAAAAAAAA

FIGURE 68

MANPGLGLLLALGLPFLRLRWGRAWGQIQTTSANENSTVLPSTSSSSSDGNLRPEAITAIIVFS
LLAALLLAVGLALLVRKLREKRQTEGTYRPSSEEQFSHAAEARAPQDSKETVQGCLPI

Important features:

Signal peptide:

amino acids 1-19

Transmembrane domain:

amino acids 56-80

N-glycosylation site.

amino acids 36-40

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 86-90

Tyrosine kinase phosphorylation site.

amino acids 86-94

N-myristoylation sites.

amino acids 7-13, 26-32

FIGURE 69

[illegible]

FIGURE 70

MGLFRGFVFLVLCLLHQSNSTFIKLNNGFEDIVIVIDPSVPEDEKIIIEQIEDMVTASTYLFE
ATEKRFFFKNVSILIPENWKENPQYKRPKHENHKHADVIVAPPTLPGRDEPYTKQFTECGEKGEY
IHFTPDLLLGGKQNEYGPPGKLFVHEWAHLRWGVFDEYNEDQPFYRAKSKKIEATRCISAGISGRN
RVYKCQGGSCLSRACRIDSTTKLYGKDCQFFPDKVQTEKASIMFMQSIDSVVEFCNEKTHNQEAP
SLQNIKCNFRSTWEVISNSEDFKNTIPMVTPPPPPVFSLKISQRIVCLVLDKSGSMGGKDRLNR
MNQAAKHFLLOTVENGSWGMVHFDSTATIVNKLIQIKSSDERNTLMAGLPTYPLGGTSICSGIK
YAFQVIGELHSQLDGSEVLLLTGDEDNTASSCIDEVKQSGAIVHFIALGRAADEAVIEMSKITGG
SHFYVSDEAQNGGLIDAFGALTSGNTDLSQKSLQLESKGLTLNSNAWMNDTVIIDSTVGKDTFFL
ITWNSLPPSISLWDPSGTIMENFTVDATSKMAYLSIPGTAKVGTWAYNLQAKANPETLTITVTSR
AANSSVPPITVNAKMNDVNSFPSPMIVYAEILQGYVPVLGANVTAFIESQNGHTEVLELLDNGA
GADSEKNDGVYSRYFTAYTENGRYSLKVRAGGANTARLKLAPPLNRAAYIPGWVVGGEIEANPP
RPEIDEDTQTTLEDFSRASGGAFVVSQVPSLPLPDQYPPSQITDLDATVHEDKIILTWTPAGDN
FDVGKVQRYIIRISASILDRLDSFDDALQVNTTDLSPKEANSKESFAFKPENISEENATHIFIAI
KSIDKSNLTSKVSNIAQVTLFIPQANPDDIDPTPTPTPTPTPDKSHNSGVNISTLVLSVIGSVVI
VNFILSTTI

Signal peptide:

amino acids 1-21

Putative transmembrane domains:

amino acids 284-300, 617-633

Leucine zipper pattern.

amino acids 469-491, 476-498

N-glycosylation site.

amino acids 20-24, 75-79, 340-344, 504-508, 542-546, 588-592,
628-632, 811-815, 832-836, 837-841, 852-856, 896-900

FIGURE 71

CTCCTTAGGTGGAACCCCTGGGAGTAGAGTACTGACAGCAAAGACCGGGAAGACCATACGTCCCCGGGCAGGGGTGA
CAACAGGTGTCTCTTTTGTCTCGTGTGTGGCTGCCCTTCTATTTCAAGGAAAGACGCCAAGGTAATTTTGACCCA
GAGGAGCAATGATGTAGCCACCTCCTAACCTTCCCTTCTTGAACCCCCAGTTATGCCAGGATTTACTAGAGAGTGTCA
ACTCAACCAGCAAGCGGCTCCTTCGGCTTAACCTTGTGGTTGGAGGAGAGAACCTTTGTGGGGCTGCCCTCTCTTAGCA
GTGCTCAGAAGTGACTTGCCTGAGGGTGGACCAGAGAAGAAAGGTAAGGTCCTCTTGTGTGGCTGCACATCAGGAA
GGCTGTGATGGGAATGAAGGTGAAAACCTTGGAGATTTCACTTCAGTCATTGCTTCTGCCCTGCAAGATCATCCTTTAAA
AGTAGAGAAGCTGCTCTGTGTGGTGGTTAACTCCAAGAGGCAGAACTCGTCTAGAAGGAAATGGATGCAAGCAGCTC
CGGGGGCCCCAAACGCATGCTTCTGTGGTCTAGCCAGGGAAGCCCTTCCGTGGGGGGCCCCGGCTTTGAGGGATGCC
ACCGGTTCTGGACGCATGGCTGATTCTGAAATGATGATGGTTGCGCGGGGGCTGCTTGCCTGGATTTCGGGGTGGTG
GTTTTGCTGGTGTCTCTCTGTGTGTCTATCTCTGTCTGTACATGTTGGCCTGCACCCCAAAGGTGACGAGGAGCAG
CTGGCACTGCCAGGGCCAACAGCCCCACGGGAAGGAGGGGTACCAGGCCGTCTTTCAGGAGTGGGAGGAGCAGCAC
CGCAACTACGTGAGCAGCCTGAAGCGGCAGATCGCACAGCTCAAGGAGGAGCTGCAGGAGAGGAGTGCAGCAGCTCAGG
AATGGGCAGTACCAAGCAGCGATGCTGCTGGCCTGGGTCTGGACAGGAGCCCCCAGAGAAAACCCAGGCCGACCTC
CTGGCCTTCTCGACTCGCAGGTGGACAAGGCAGAGGTGAATGCTGGCCTCAAGCTGGCCACAGAGTATGCAGCAGTG
CCTTTTCGATAGCTTTTACTCTACAGAAGGTGTACCAGCTGGAGACTGGCCTTACCGCCACCCCGAGGAGAAGCCTGTG
AGGAAGGACAAGCGGGATGAGTTGGTGGAGCCATTGAATCAGCCTTGGAGACCCTGAACAATCCTGCAGAGAACAGC
CCCAATCACCGTCTTACACGGCCTCTGATTTCATAGAAGGGATCTACCGAACAGAAAGGGACAAGGGACATTGTAT
GAGCTCACCTTCAAAGGGGACCACAAACACGAATTCAAACGGCTCATCTTATTTGACCATTCAGCCCCATCATGAAA
GTGAAAAATGAAAAGCTCAACATGGCCAACACGCTTATCAATGTTATCGTGCCTCTAGCAAAAAGGGTGGACAAGTTC
CGSCAGTTTATGCGAATTTTCAGGGAGATGTGCATTGAGCAGGATGGGAGAGTCCATCTCACTGTTGTTTACTTTGGG
AAAGAAGAAATAAATGAAGTCAAAGGAATACTTGAAAACACTTCCAAAGCTGCCAATTCAGGAACCTTACCTTCATC
CAGCTGAATGGAGAATTTTCTCGGGGAAAGGGACTTGATGTTGGAGCCCGCTTCTGGAAGGGAAGCAACGTCCTTCTC
TTTTCTGTGATGTGGACATCTACTTCACATCTGAATTCCTCAATACGTGTAGGCTGAATACACAGCCAGGGAAGAAG
GTATTTTATCCAGTTCTTTTCAGTCAGTACAATCCTGGCATAATATACGGCCACCATGATGCAGTCCCTCCCTTGGAA
CAGCAGCTGGTCATAAAGAAGGAACTGGATTTTGGAGAGACTTTGGATTGTTGGGATGACGTGTGAGTATCGGTCAGAC
TTCATCAATATAGGTGGGTTTGTCTGGACATCAAAGGCTGGGGCGGAGAGGATGTGCACCTTTATCGCAAGTATCTC
CAGACCAACCTCATAGTGGTACGGACGCCGTGTGCGAGGACTCTTCCACCTCTGGCATGAGAAGCGCTGCATGGACGAG
CTGACCCCCGAGCAGTACAAGATGTGCATGCACTCAAGGCCATGAACGAGGCATCCCACGGCCAGCTGGGCATGCTG
GTGTTTCAGGCACGAGATAGAGGCTCACCTTCGCAACAGAAACAGAAAGACAAGTAGCAAAAAACATGAAGTCCCAGA
GAAGGATTGTGGGAGACACTTTTCTTTCTTTTGCATTTACTGAAAGTGGCTGCAACAGAGAAAAGACTTCCATAAA
GGACGACAAAAGATTGGACTGATGGGTGAGAGATGAGAAGCCTCCGATTTCTCTCTGTTGGGCTTTTACAAACAGA
AATCAAAATCTCCGCTTTGCCCTGCAAAAGTAACCCAGTTGCACCTGTGAAGTGTCTGACAAAGGCAGAAATGCTTGTG
AGATTATAAGCCTAATGGTGTGAGGTTTTGATGGTGTTTACAATACACTGAGACCTGTTGTTTTGTGTGCTCATTGA
AATATTCTGATTTAAGAGCAGTTTTGTAAAAAATTCATTAGCATGAAAGGCAAGCATATTTCTCCTCATATGAATGA
GCCTATCAGCAGGGCTCTAGTTTCTAGGAATGCTAAAATATCAGAAGGCAGGAGAGGAGATAGGCTTATTATGATACT
AGTGASTACATTAAGTAAAATAAAATGGACCAGAAAAGAAAAGAAACCATAAATATCGTGTCTATTTTCCCCAAGAT
TAACCAAAAATAATCTGCTTATCTTTTTGGTTGTCTTTTAACTGTCTCCGTTTTTTTCTTTTATTTAAAAATGCACT
TTTTTTCCCTTGTGAGTTATAGTCTGCTTATTTAATTACCACTTTGCAAGCCTTACAAGAGAGCACAGTTGGCCTAC
ATTTTTATATTTTTTAAGAAGATACTTTGAGATGCATTATGAGAACTTTCAGTTCAAAGCATCAAATGATGCCATAT
CCAAGGACATGCCAAATGCTGATTCTGTGAGGCACTGAATGTGAGGCATTGAGACATAGGGAAGGAATGGTTTGTACT
AATACAGACGTACAGATACTTTCTCTGAAGAGTATTTTCGAAGAGGAGCAACTGAACACTGGAGGAAAAGAAAATGAC
ACTTTCTGCTTTACAGAAAAGGAACTCATTGAGACTGGTGATATCGTGATGTACCTAAAAGTCAGAAAACCACATTTT
CTCCTCAGAAGTAGGGACCGCTTTCTTACCTGTTTAAATAAACCAGTATACCGTGTGAACCAACAATCTCTTTTC
AAAACAGGGTGTCTCCTCCTGGCTTCTGGCTCCATAAGAAGAAATGGAGAAAAATATATATATATATATATATATGT
GAAAGATCAATCCATCTGCCAGAACTAGTGGGATGGAAGTTTTTGCTACATGTTATCCACCCAGGCCAGGTGGAAG
TAACCTGAATTATTTTTTAAATTAAGCAGTTCTACTCAATCACCAGATGCTTCTGAAAATTGCATTTTATTACCATTT
CAAACTATTTTTTAAAAATAAATACAGTTAAATAGAGTGGTTTTCTTCAATTCATGTGAAAATATTAGCCAGCACCAG
ATGCATGAGCTAATTATCTCTTTGAGTCCCTGCTTCTGTTTGTCTCAGGTAAACTCATTGTTTAAAGCTTCAAGAAC
ATTCAAGCTGTTGGTGTGTTAAAAAATGCATTGTATTGATTGTACTGGTAGTTTATGAAATTTAATTAAACACAGG
CCATGAATGGAAGGTGGTATTGCACAGCTAATAAATATGATTTGTGGATATGAA

FIGURE 72

MMVRRGLLAWISRVVLLVLLCCAISVLYMLACTPKGDEEQALALPRANSPTGKEGYQAVLQEW
EQHRNYVSSLKRQIAQLKEELQERSEQLRNGQYQASDAAGLGLDRSPPEKTQADLLAFLHSQVDK
AEVNAGVKLATEYAAVPFDSFTLQKVYQLETGLTRHPEEKPVKDKRDELVEAIESALETNNPA
ENSPNHRPYTASDFIEGIYRTERDKGTLYELTFKGDHKHEFKRLILFRPFSPIMKVNEKLNMAN
TLINVIVPLAKRVDKFRQFMQNFREMCIEQDGRVHLTVVYFGKEEINEVKGILENTSKAANFRNF
TFIQLNGEFSRGKGLDVGARFWKGSNVLLFFCDVDIYFTSEFLNTCRLNTQPGKKVFYPVLESQY
NPGIYGHHDVPPLEQQLVIKKETGFWRDFGFGMTCQYRSDFINIGGFDLDIKGWGGEDVHLYR
KYLHSNLIVVRTPVRLFHLWHEKRCMDLTPEQYKCMQSKAMNEASHGQLGMLVFRHEIEAHL
RKQKQKTSSKKT

Important features:

Signal peptide:

amino acids 1-27

N-glycosylation sites.

amino acids 315-319, 324-328

N-myristoylation sites.

amino acids 96-102, 136-142, 212-218, 311-317, 339-345, 393-399

Amidation site.

amino acids 377-381

FIGURE 73

GAGACTGCAGAGGGAGATAAAGAGAGAGGGCAAAGAGGCAGCAAGAGATTTGTCCTGGGGATCCA
 GAAACCCATGATACCCTACTGAACACCGAATCCCCTGGAAGCCACAGAGACAGAGACAGCAAGA
 GAAGCAGAGATAAATACACTCAGGCCAGGAGCTCGCTCGCTCTCTCTCTCTCTCTCTCACTCCTC
 CCTCCCTCTCTCTCTGCCTGTCCTAGTCCTCTAGTCCTCAAATTTCCAGTCCCCTGCACCCCTTC
 CTGGGACACTATGTTGTTCTCCGCCCTCCTGCTGGAGGTGATTTGGATCCTGGCTGCAGATGGGG
 GTCAACACTGGACGTATGAGGGCCACATGGTCAGGACCATTGGCCAGCCTCTTACCCTGAGTGT
 GGAAACAATGCCCAGTCGCCCATCGATATTTCAGACAGACAGTGTGACATTTGACCCTGATTTGCC
 TGCTCTGCAGCCCCACGGATATGACCAGCCTGGCACCGAGCCTTTGGACCTGCACAACAATGGCC
 ACACAGTGCAACTCTCTCTGCCCTCTACCCTGTATCTGGGTGGACTTCCCCGAAAATATGTAGCT
 GCCCAGCTCCACCTGCACTGGGGTCAGAAAGGATCCCCAGGGGGGTGAGAACACCAGATCAACAG
 TGAAGCCACATTTGCAGAGCTCCACATTGTACATTATGACTCTGATTCCATGACAGCTTGAGTG
 AGGCTGCTGAGAGGCCTCAGGGCCTGGCTGTCCTGGGCATCCTAATTGAGGTGGGTGAGACTAAG
 AATATAGCTTATGAACACATTCTGAGTCACTTGCATGAAGTCAGGCATAAAGATCAGAAGACCTC
 AGTGCCTCCCTTCAACCTAAGAGAGCTGCTCCCCAACAGCTGGGGCAGTACTTCCGCTACAATG
 GCTCGCTCACAACCTCCCCCTTGCTACCAGAGTGTGCTCTGGACAGTTTTTTATAGAAGGTCCCAG
 ATTTCAATGGAACAGCTGGAAAAGCTTCAGGGGACATTGTTCTCCACAGAAGAGGAGCCCTCTAA
 GCTTCTGGTACAGAACTACCGAGCCCTTCAGCCTCTCAATCAGCGCATGGTCTTTGCTTCTTTCA
 TCCAAGCAGGATCCTCGTATACCACAGGTGAAATGCTGAGTCTAGGTGTAGGAATCTTGGTTGGC
 TGTCTCTGCCTTCTCCTGGCTGTTTATTTTCATTGCTAGAAAGATTGGAAGAAGAGGCTGGAAAA
 CCGAAAGAGTGTGGTCTTCACCTCAGCACAAGCCACGACTGAGGCATTAAATTCCTTCTCAGATAC
 CATGGATGTGGATGACTTCCCTTCATGCCTATCAGGAAGCCTCTAAAATGGGGTGTAGGATCTGG
 CCAGAAACACTGTAGGAGTAGTAAGCAGATGTCTCCTTCCCCTGGACATCTCTTAGAGAGGAAT
 GGACCCAGGCTGTCATTCCAGGAAGAACTGCAGAGCCTTCAGCCTCTCCAAACATGTAGGAGGAA
 ATGAGGAAATCGCTGTGTTGTTAATGCAGAGANCAAACTCTGTTTAGTTGCAGGGGAAGTTTGGG
 ATATACCCCAAAGTCCTCTACCCCTCACTTTTATGGCCCTTCCCTAGATATACTGCGGGATCT
 CTCCTTAGGATAAAGAGTTGCTGTTGAAGTTGTATATTTTGATCAATATATTTGGAAATTAAAG
 TTTCTGACTTT

FIGURE 74

MLFSALLLEVIWILAADGGQHWTYEGPHGQDHPASYPECGNNAQSPIDIQTDSVTFDPDLPALQ
PHGYDQPGTEPLDLHNNGHTVQLSLPSTLYLGGLPRKYVAAQLHLHWGQKGSPPGGSEHQINSEAT
FAELHIVHYDSYDSLSEAAERPQGLAVLGILIEVGETKNIAYEHILSHLHEVRHKDQKTSVPP
FNLRELLPKQLGQYFRYNGSLTTPPCYQSVLWTVFYRRSQISMEQLEKLQGTLFSTEEEPSKLLV
QNYRALQPLNQRMVFASFIQAGSSYTTGEMLSLGVGILVGCLCLLLAVYFIARKIRKKRLENRKS
VVFTSAQATTEA

Important features of the protein:

Signal peptide:

amino acids 1-15

Transmembrane domain:

amino acids 291-310

N-glycosylation site.

amino acids 213-216

Eukaryotic-type carbonic anhydrases proteins

amino acids 197-245, 104-140, 22-69

FIGURE 75

TGCCGCTGCCGCCGCTGCTGCTGTTGCTCCTGGCGGCGCCTTGGGGACGGGCAGTTCCCTGTGTC
TCTGGTGGTTTGCCTAAACCTGCAAACATCACCTTCTTATCCATCAACATGAAGAATGTCCTACA
ATGGACTCCACCAGAGGGTCTTCAAGGAGTTAAAGTTACTTACACTGTGCAGTATTTTCATCACAA
ATTGGCCCCACCAGAGGTGGCACTGACTACAGATGAGAAGTCCATTTCTGTTGCTCTGACAGCTCC
AGAGAAGTGGAAGAGAAATCCAGAAGACCTTCCTGTTTCCATGCAACAAATATACTCCAATCTGA
AGTATAACGTGTCTGTGTTGAATACTAAATCAAACAGAACGTGGTCCCAGTGTGTGACCAACCAC
ACGCTGGTGCTCACCTGGCTGGAGCCGAACACTCTTTACTGCGTACACGTGGAGTCCTTCGTCCC
AGGGCCCCCTCGCCGTGCTCAGCCTTCTGAGAAGCAGTGTGCCAGGACTTTGAAAGATCAATCAT
CAGAGTTCAAGGCTAAAATCATCTTCTGGTATGTTTTGCCCATATCTATTACCGTGTTTCTTTTT
TCTGTGATGGGCTATTCCATCTACCGATATATCCACGTTGGCAAAGAGAAACACCCAGCAAATTT
GATTTTGATTTATGGAAATGAATTTGACAAAAGATTCTTTGTGCCTGCTGAAAAAATCGTGATTA
ACTTTATCACCCCTCAATATCTCGGATGATTCTAAAATTTCTCATCAGGATATGAGTTTACTGGGA
AAAAGCAGTGATGTATCCAGCCTTAATGATCCTCAGCCCAGCGGGAACCTGAGGCCCCCTCAGGA
GGAAGAGGAGGTGAAACATTTAGGGTATGCTTCGCATTTGATGGAAATTTTTTGTGACTCTGAAG
AAAACACGGAAGGTACTTCTCTACCCAGCAAGAGTCCCTCAGCAGAACAAATACCCCCGGATAAA
ACAGTCATTGAATATGAATATGATGTCAGAACCACTGACATTTGTGCGGGGCTGAAGAGCAGGA
GCTCAGTTTGCAGGAGGAGGTGTCCACACAAGGAACATTATTGGAGTCGCAGGCAGCGTTGGCAG
TCTTGGGCCCCGAAACGTTACAGTACTCATACACCCCTCAGCTCCAAGACTTAGACCCCCCTGGCG
CAGGAGCACACAGACTCGGAGGAGGGGCCGGAGGAAGAGCCATCGACGACCCTGGTCTGACTGGGA
TCCCCAAACTGGCAGGCTGTGTATTCCTTCGCTGTCCAGCTTCGACCAGGATTAGAGGGCTGCG
AGCCTTCTGAGGGGGATGGGCTCGGAGAGGAGGGTCTTCTATCTAGACTCTATGAGGAGCCGGCT
CCAGACAGGCCACCAGGAGAAAAATGAAACCTATCTCATGCAATTCATGGAGGAATGGGGTTATA
TGTGCAGATGGAAAACTGATGCCAAACACTTCCTTTTGCCCTTTGTTTCCTGTGCAACAAGTGAG
TCACCCCTTTGATCCAGCCATAAAGTACCTGGGATGAAAGAAGTTTTTCCAGTTTGTGAGTGT
CTGTGAGAATTACTTATTTCTTTTCTCTATTCTCATAGCACGTGTGTGATTGGTTTCATGCATGTA
GGTCTCTTAACAATGATGGTGGGCCCTCTGGAGTCCAGGGGCTGGCCGTTGTTCTATGCAGAGAA
AGCAGTCAATAAATGTTTGCCAGACTGGGTGCAGAATTTATTCAGGTGGGTGT

FIGURE 76

MSYNGLHQRVFKELKLLTLCSSIQIGPPEVALTTDEKSI SVVLTAPEKWKRN PEDLPVSMQQIY
SNLKYNVSVLNTKSNRTWSQCVTNHTLVLTWLEPNTLYCVHVESFVPGPPRAQ PSEKQCARTLK
DQSSEFKAKIIFWYVLPISITVFLFSVMGYSIYRYIHVGKEKH PANLILYIGNEFDKRFFVPAEK
IVINFITLNISSDDSKISHQDMSLLGKSSDVSSLNDPQPSGNLRPPQEEEEVKHLGYASHLMEIFC
DSEENTEGTSLTQQESLSRTIPDPKTVIEYDYDVRTTDICAGPEEQELSLQEEVSTQGTLLESQA
ALAVLGPQTLQYSYTPQLQDL DPLAQEHTDSEEGPEEEPSTTLVDWDPQTGR LCIPLSSFDQDS
EGCEPSEGDGLGEEGLLSRLYEEPAPDRPPGENETYLMQFMEEWGLYVQ MEN

Important features:

Signal peptide:

amino acids 1-28

Transmembrane domain:

amino acids 140-163

N-glycosylation sites.

amino acids 71-74, 80-83, 89-92, 204-207, 423-426

FIGURE 77

GAGGAGCGGGCCGAGGACTCCAGCGTGCCCGAGGTCTGGCATCCTGCACTTGCTGCCCTCTGACAC
 CTGGGAAGATGGCCCGGCCCGTGACCTTCACCCTTCTCTGTGGTTTGCTGGCAGCCACCTTGATC
 CAAGCCACCCTCAGTCCCCTGCACTGAGTTCTCATCCTCGGCCCAAAGTCATCAAAGAAAAGCTGAC
 ACAGGAGCTGAAGGACCACAACGCCACCAGCATCCTGCAGCAGCTGCCGCTGCTCAGTGCCATGC
 GGGAAAAGCCAGCCGGAGGCATCCCTGTGCTGGGCAGCCTGGTGAACACCGTCCTGAAGCACATC
 ATCTGGCTGAAGGTCATCACAGCTAACATCCTCCAGCTGCAGGTGAAGCCCTCGGCCAATGACCA
 GGAGCTGCTAGTCAAGATCCCCCTGGACATGGTGGCTGGATTCAACACGCCCTGGTCAAGACCA
 TCGTGGAGTTCCACATGACGACTGAGGCCCCAAGCCACCATCCGCATGGACACCAGTGCAAGTGGC
 CCCACCCGCCTGGTCCTCAGTGACTGTGCCACCAGCCATGGGAGCCTGCGCATCCAAGTCTGTGA
 TAAGCTCTCCTTCCTGGTGAACGCCTTAGCTAAGCAGGTCATGAACCTCCTAGTGCCATCCCTGC
 CCAATCTAGTGAAAAACCAGCTGTGTCCCGTGATCGAGGCTTCCTTCAATGGCATGTATGCAGAC
 CTCCTGCAGCTGGTGAAGGTGCCCATTTCCTCAGCATTGACCGTCTGGAGTTTGACCTTCTGTGA
 TCCTGCCATCAAGGGTGACACCATTAGCTCTACCTGGGGGCCAAGTTGTTGGACTCACAGGGAA
 AGGTGACCAAGTGGTTCAATAACTCTGCAGCTTCCCTGACAATGCCACCCTGGACAACATCCCCG
 TTCAGCCTCATCGTGAGTCAGGACGTGGTGAAAGCTGCAGTGGCTGCTGTGCTCTCTCCAGAAGA
 ATTCATGGTCCTGTTGGACTCTGTGCTTCCTGAGAGTGCCCATCGGCTGAAGTCAAGCATCGGGC
 TGATCAATGAAAAGGCTGCAGATAAGCTGGGATCTACCCAGATCGTGAAGATCCTAACTCAGGAC
 ACTCCCGAGTTTTTTATAGACCAAGGCCATGCCAAGGTGGCCCAACTGATCGTGCTGGAAGTGTT
 TCCCTCCAGTGAAGCCCTCCGCCCTTTGTTACCCCTGGGCATCGAAGCCAGCTCGGAAGCTCAGT
 TTTACACCAAAGGTGACCAACTTATACTCAACTTGAATAACATCAGCTCTGATCGGATCCAGCTG
 ATGAACTCTGGGATTGGCTGGTTCCAACCTGATGTTCTGAAAAACATCATCACTGAGATCATCCA
 CTCCATCCTGCTGCCGAACCAGAATGGCAAATTAAGATCTGGGGTCCCAGTGTGATTGGTGAAGG
 CCTTGGGATTTCGAGGCAGCTGAGTCCTCACTGACCAAGGATGCCCTTGTGCTTACTCCAGCCTCC
 TTGTGGAAACCCAGCTCTCCTGTCTCCCAGTGAAGACTTGGATGGCAGCCATCAGGGAAGGCTGG
 GTCCCAGCTGGGAGTATGGGTGTGAGCTCTATAGACCATCCCTCTCTGCAATCAATAAACACTTG
 CCTGTGAAAAA

FIGURE 78

MAGPWTFTLLCGLLAATLIQATLSPTAVLILGPKVIKEKLTQELKDNATSILQQLPLLSAMREK
PAGGIPVLGSLVNTVLKHIIWLKVITANILQLQVKPSANDQELLVKIPLDMVAGENTPLVKTIIVE
FHMTTEAQATIRMDTSASGPTRLVLSDCATSHGSLRIQLLYKLSFLVNALAKQVMNLLVPSLPNL
VKNQLCPVIEASFNGMYADLLQLVKVPISLSIDRLEFDLLYPAIKGDITQLYLGAKLLDSQGKVT
KWFNNSAASLTMPITLDNIPFSLIVSQDVVKAAVAVALSPEEFMVLLDSVLPESAHRLKSSIGLIN
EKAADKLGSTQIVKILTQDTPEFFIDQGHAKVAQLIVLEVFPSSSEALRPLFTLGIEASSEAQFYT
KGDQLILNLNLISSDRIQLMNSGIGWFQPDVLKNIITEIIHSILLPNQNGKLRSGVPVSLVKALG
FEAAESSLTKDALVLTPASLWKPPSPVSQ

Important features of the protein:

Signal peptide:

amino acids 1-21

N-glycosylation sites.

amino acids 48-51, 264-267, 401-404

Glycosaminoglycan attachment site.

amino acids 412-415

LBP / BPI / CETP family proteins.

amino acids 407-457

FIGURE 79

GAGAGAAGTCAGCCTGGCAGAGAGACTCTGAAATGAGGGATTAGAGGTGTTCAAGGAGCAAGAGC
TTCAGCCTGAAGACAAGGGAGCAGTCCCTGAAGACGCTTCTACTGAGAGGTCTGCCATGGCCTCT
CTTGGCCTCCAACTTGTGGGCTACATCCTAGGCCTTCTGGGGCTTTTGGGCACACTGGTTGCCAT
GCTGCTCCCCAGCTGGAAAAACAAGTTCTTATGTGGGTGCCAGCATTGTGACAGCAGTTGGCTTCT
CCAAGGGCCTCTGGATGGAATGTGCCACACACAGCACAGGCATCACCCAGTGTGACATCTATAGC
ACCCTTCTGGGCCTGCCCCGCTGACATCCAGGCTGCCCAGGCCATGATGGTGACATCCAGTGCAAT
CTCCTCCCTGGCCTGCATTATCTCTGTGGTGGGCATGAGATGCACAGTCTTCTGCCAGGAATCCC
GAGCCAAAGACAGAGTGGCGGTAGCAGGTGGAGTCTTTTTCATCCTTGGAGGCCTCCTGGGATTCT
ATTCTGTTGCCTGGAATCTTCATGGGATCCTACGGGACTTCTACTCACCCTGGTGCCCTGACAG
CATGAAATTTGAGATTGGAGAGGCTCTTTACTTGGGCATTATTTCTTCCCTGTTCTCCCTGATAG
CTGGAATCATCCTCTGCTTTTCCCTGCTCATCCCAGAGAAATCGCTCCAACCTACTACGATGCCTAC
CAAGCCCAACCTCTTGCCACAAGGAGCTCTCCAAGGCCTGGTCAACCTCCCAAAGTCAAGAGTGA
GTTCAATTCTACAGCCTGACAGGGTATGTGTGAAAGAACCAGGGGCCAGAGCTGGGGGGTGGCTG
GGTCTGTGAAAAACAGTGGACAGCACCCCGAGGGCCACAGGTGAGGGACACTACCACTGGATCGT
GTCAGAAGGTGCTGCTGAGGATAGACTGACTTTGGCCATTGGATTGAGCAAAGGCAGAAATGGGG
GCTAGTGTAACAGCATGCAGGTTGAATTGCCAAGGATGCTCGCCATGCCAGCCTTTCTGTTTTCC
TCACCTTGCTGCTCCCCTGCCCTAAGTCCCCAACCTCAACTTGAAACCCCATTCCTTAAGCCA
GGACTCAGAGGATCCCTTTGCCCTCTGGTTTACCTGGGACTCCATCCCCAAACCCACTAATCACA
TCCCCTGACTGACCCTCTGTGATCAAAGACCCTCTCTCTGGCTGAGGTTGGCTCTTAGCTCATT
GCTGGGGATGGGAAGGAGAAGCAGTGGCTTTTGTGGGCATTGCTCTAACCTACTTCTCAAGCTTC
CCTCCAAAGAACTGATTGGCCCTGGAACCTCCATCCCCTCTTGTTATGACTCCACAGTGTCCA
GACTAATTTGTGCATGAACTGAAATAAAACCATCCTACGGTATCCAGGGAACAGAAAGCAGGATG
CAGGATGGGAGGACAGGAAGGCAGCCTGGGACATTTAAAAAATA

FIGURE 8o

MASLGLQLVGYILGLLGGLTLVAMLLPSWKTSSYVGASIVTAVGFSGKLWMECATHSTGITQCD
IYSTLLGLPADIQAAQAMMVTSSAIISSLACIISVVGMRCTVFCQESRAKDRVAVAGGVFFILGGL
LGFIPVAWNLHGILRDFYSPLVPDSMKFEIGEALYLGIISSLSLIAGIILCFSCSSQRNRSNYY
DAYQAQPLATRSSPRPGQPPKVKSEFNSYSLTGYV

Important features of the protein:

Signal peptide:

amino acids 1-24

Transmembrane domains:

amino acids 82-102, 117-140, 163-182

N-glycosylation site.

amino acids 190-193

PMP-22 / EMP / MP20 family proteins.

amino acids 46-59

FIGURE 81

CCCACGCGTCCGCGCCTCTCCCTTCTGCTGGACCTTCCTTCGTCTCTCCATCTCTCCCTCCTTTT
CCCGCGTTCTCTTTCCACCTTTCTCTTCTTCCACCTTAGACCTCCCTTCCTGCCCTCCTTTCCCT
GCCCACCGCTGCTTCCTGGCCCTTCTCCGACCCCGCTCTAGCAGCAGACCTCCTGGGGTCTGTGG
GTTGATCTGTGGCCCCTGTGCCTCCGTGTCTTTTCGTCTCCCTTCCTCCCGACTCCGCTCCCGG
ACCAGCGGCCTGACCCTGGGGAAAGGATGGTTCCCGAGGTGAGGGTCCTCTCCTCCTTGCTGGGA
CTCGCGCTGCTCTGGTTCCCCCTGGACTCCCACGCTCGAGCCCGCCCAGACATGTTCTGCCTTTT
CCATGGGAAGAGATACTCCCCGGCGAGAGCTGGCACCCCTACTTGAGCCACAAGGCCTGATGT
ACTGCCTGCGCTGTACCTGCTCAGAGGGCGCCCATGTGAGTTGTTACCGCCTCCACTGTCCGCCT
GTCCACTGCCCCCAGCCTGTGACGGAGCCACAGCAATGCTGTCCCAAGTGTGTGGAACCTCACAC
TCCCTCTGGACTCCGGGCCCCACCAAAGTCCTGCCAGCACAAACGGGACCATGTACCAACACGGAG
AGATCTTCAGTGCCCATGAGCTGTTCCCCTCCCGCCTGCCCAACCAGTGTGTCTCTGCAGCTGC
ACAGAGGGCCAGATCTACTGCGGCCTCACAACCTGCCCCGAACCAGGCTGCCCAGCACCCCTCCC
ACTGCCAGACTCCTGCTGCCAAGCCTGCAAAGATGAGGCAAGTGAGCAATCGGATGAAGAGGACA
GTGTGCAGTCGCTCCATGGGGTGAGACATCCTCAGGATCCATGTTCCAGTGATGCTGGGAGAAAG
AGAGGCCCCGGGCACCCACGCCCCACTGGCCTCAGCGCCCTCTGAGCTTCATCCCTCGCCACTT
CAGACCCAAGGGAGCAGGCAGCACAACTGTCAAGATCGTCCTGAAGGAGAAACATAAGAAAGCCT
GTGTGCATGGCGGGAAGACGTACTCCACGGGGAGGTGTGGCACCCGGCCTTCCGTGCCTTCGGC
CCCTTGCCCTGCATCCTATGCACCTGTGAGGATGGCCGCCAGGACTGCCAGCGTGTGACCTGTCC
CACCGAGTACCCCTGCCGTACCCCGAGAAAGTGGCTGGGAAGTGCTGCAAGATTTGCCCAGAGG
ACAAAGCAGACCCTGGCCACAGTGAGATCAGTTCTACCAGGTGTCCCAAGGCACCGGGCCGGGTC
CTCGTCCACACATCGGTATCCCCAAGCCAGACAACCTGCGTCGCTTTGCCCTGGAACACGAGGC
CTCGGACTTGGTGGAGATCTACCTCTGGAAGCTGGTAAAAGATGAGGAACTGAGGCTCAGAGAG
GTGAAGTACCTGGCCCAAGGCCACACAGCCAGAATCTTCCACTTGACTCAGATCAAGAAAGTCAG
GAAGCAAGACTTCCAGAAAGAGGCCACAGCACTTCCGACTGCTCGCTGGCCCCCACGAAGGTCCT
GGAACGTCTTCTAGCCCAGACCCTGGAGCTGAAGGTCACGGCCAGTCCAGACAAAGTGACCAAG
ACATAACAAAGACCTAACAGTTGCAGATATGAGCTGTATAATTGTTGTTATTATATTAATAAA
TAAGAAGTTGCATTACCCTCAAAAAAAAAAAAAAAAAAAAAA

FIGURE 82

MVPEVRVLSSLLGLALLWFPLDSHARARPD MFCLFHGKRYS PGESWHPYLEPQGLMYCLRCTCSE
GAHVSCYRLHCPPVHCPQPVTEPQQCCPKC VEPHTPSGLRAPPKSCQHNGTMYQHGEIFSAHELF
PSRLPNQCVLCSCTEGQIYCGLTTCPEPGCPAPLPLPDSCCQACKDEASEQSDEEDSVQSLHGVR
HPQDPCSSDAGRKRGP GTPAPTGLSAPLSFIPRHFRPKGAGSTTVKIVLKEKHKKACVHGGKTYS
HGEVWHPAFRAFGPLPCILCTCEDGRQDCQRVTCPT EYPCRHPK VAGKCKICPEDKADPGHSE
ISSTRCPKAPGRVLVHTSVSPSPDNLRRFALEHEASDLVEIYLWKLVKDEETEAQRGEVPGPRPH
SQNLPLDSDQESQEARLPERGTALPTARWPPRRSLERLPSDPGAEGHGQSRQSDQDITKT

Signal peptide:

amino acids 1-25

FIGURE 83

GACAGCTGTGTCTCGATGGAGTAGACTCTCAGAACAGCGCAGTTTGCCCTCCGCTCACGCAGAGCCTCTCC
 GTGGCTTCCGCACCTTGAGCATTAGGCCAGTTCTCCTCTTCTCTCTAATCCATCCGTACCTCTCCTGTCA
 TCCGTTTCCATGCCGTGAGGTCCATTACAGAACACATCCATGGCTCTCATGCTCAGTTTGGTTCTGAGTC
 TCCTCAAGCTGGGATCAGGGCAGTGGCAGGTGTTTGGGCCAGACAAGCCTGTCCAGGCCTTGGTGGGGGAG
 GACGCAGCATTCTCCTGTTTCTGTCTCCTAAGACCAATGCAGAGGCCATGGAAGTGC GGTTCTTCAGGGG
 CCAGTTCTCTAGCGTGGTCCACCTCTACAGGGACGGGAAGGACCAGCCATTTATGCAGATGCCACAGTATC
 AAGGCAGGACAAAACCTGGTGAAGGATTCTATTGCGGAGGGGCGCATCTCTCTGAGGCTGGAAAACATTACT
 GTGTTGGATGCTGGCCTCTATGGGTGCAGGATTAGTTCCAGTCTTACTACCAGAAGGCCATCTGGGAGCT
 ACAGGTGTCAGCACTGGGCTCAGTTCCCTCTCATTTCCATCACGGGATATGTTGATAGAGACATCCAGCTAC
 TCTGTCAGTCCTCGGGCTGGTTCCCCCGGCCACAGCGAAGTGGAAGGTCCACAAGGACAGGATTTGTCC
 ACAGACTCCAGGACAAAACAGAGACATGCATGGCCTGTTTGTATGTGGAGATCTCTCTGACCGTCCAAGAGAA
 CGCCGGGAGCATATCCTGTTCCATGCGGCATGCTCATCTGAGCCGAGAGGTGGAATCCAGGGTACAGATAG
 GAGATACCTTTTTTCGAGCCTATATCGTGGCACCTGGCTACCAAAGTACTGGGAATACTCTGCTGTGGCCTA
 TTTTTTGGCATTGTTGGACTGAAGATTTTCTTCTCCAAATTCAGTGGAATCCAGGCGGAACCTGGACTG
 GAGAAGAAAGCACCGACAGGCAGAATTGAGAGACGCCCGAAACACGCAGTGGAGGTGACTCTGGATCCAG
 AGACGGCTCACCCGAAGCTCTGCGTTTCTGATCTGAAAACGTAAACCCATAGAAAAGCTCCCCAGGAGGTG
 CCTCACTCTGAGAAGAGATTTACAAGGAAGAGTGTGGTGGCTTCTCAGAGTTTCCAAGCAGGGAAACATTA
 CTGGGAGGTGGACGGAGGACACAATAAAAGGTGGCGCGTGGGAGTGTGCCGGGATGATGTGGACAGGAGGA
 AGGAGTACGTGACTTTGTCTCCCGATCATGGGTACTGGGTCTCAGACTGAATGGAGAACATTTGTATTTT
 ACATTAAATCCCCGTTTTATCAGCGTCTTCCCCAGGACCCACCTACAAAAATAGGGGTCTTCTTGACTA
 TGAGTGTGGGACCATCTCCTTCTTCAACATAAATGACCAGTCCCTTATTTATACCTGACATGTGGTTTG
 AAGGCTTATTGAGGCCCTACATTGAGTATCCGTCTTATAATGAGCAAAATGGAACCTCCATAGTCATCTGC
 CCAGTCAACCAGGAATCAGAGAAAGAGGCCTCTTGGCAAAGGGCCTCTGCAATCCCAGAGACAAGCAACAG
 TGAGTCCTCCTCACAGGCAACCACGCCCTTCTCCCCAGGGGTGAAATGTAGGATGAATCACATCCCACAT
 TCTTCTTTAGGGATATTAAGGTCTCTCTCCAGATCCAAAGTCCCGCAGCAGCCGGCCAAGGTGGCTTCCA
 GATGAAGGGGGACTGGCCTGTCCACATGGGAGTCAGGTGTCATGGCTGCCCTGAGCTGGGAGGGAAGAAGG
 CTGACATTACATTTAGTTTGCTCTCACTCCATCTGGCTAAGTGATCTTGAAATACCACCTCTCAGGTGAAG
 AACCGTCAGGAATTCCCATCTCACAGGCTGTGGTGTAGATTAAGTAGACAAGGAATGTGAATAATGCTTAG
 ATCTTATTGATGACAGAGTGTATCCTAATGGTTTGTTTCATTATATTACACTTTCAGTAAAAAAA

FIGURE 84

MALMLSLVLSLLKLGSGQWQVFGDPKPVQALVGEDAAFSCFLSPKTNAEAMEVRFFRGQFSSVVH
LYRDGKDQPFMQMPQYQGRTKLVKDSIAEGRISLRLENITVLDAGLYGCRISQSYQKAIWELQ
VSALGSVPLISITGYVDRDIQLLCQSSGWFFRPTAKWKGPQGQDLSTDSRTNRDMHGLFDVEISL
TVQENAGSISCSMRHAHLSREVESRVQIGDTFFEPISWHLATKVLGILCCGLFFGIVGLKIFFSK
FQWKIQAELDWRRKHGQAEIRDARKHAVEVTLDPETAHPKLCVSDLKTVTHRKAQEVPHSEKRF
TRKSVVASQSFQAGKHYWEVDGGHNKRWRVGVCRDDVDRRKEYVTLSPDHGYWVLRNLNGEHLFT
LNPRFISVFPRTPTTKIGVFLDYECGTISFFNINDQSLIYTLTCRFEGLLRPYIEYPSYNEQNGT
PIVICPVTQESEKEASWQRASALPETSNSSESSSQATTFFLPRGEM

Signal peptide:

amino acids 1-17

Transmembrane domain:

amino acids 239-255

FIGURE 85

AACAGACGTTCCCTCGCGGCCCTGGCACCTCTAACCCAGACATGCTGCTGCTGCTGCTGCCCCCT
GCTCTGGGGGAGGGAGAGGGCGGAAGGACAGACAAGTAACTGCTGACGATGCAGAGTTCCGTGA
CGGTGCAGGAAGGCCTGTGTGTCCATGTGCCCTGCTCCTTCTCCTACCCCTCGCATGGCTGGATT
TACCCCTGGCCCAGTAGTTTCATGGCTACTGGTTCCGGGAAGGGGCCAATACAGACCAGGATGCTCC
AGTGGCCACAAACAACCCAGCTCGGGCAGTGTGGGAGGAGACTCGGGACCGATTCCACCTCCTTG
GGGACCCACATACCAAGAATTGCACCCTGAGCATCAGAGATGCCAGAAGAAGTGATGCGGGGAGA
TACTTCTTTCGTATGGAGAAAGGAAGTATAAAATGGAATTATAAACATCACCGGCTCTCTGTGAA
TGTGACAGCCTTGACCCACAGGCCCAACATCCTCATCCCAGGCACCCTGGAGTCCGGCTGCCCCC
AGAATCTGACCTGCTCTGTGCCCTGGGCCTGTGAGCAGGGGACACCCCTATGATCTCCTGGATA
GGGACCTCCGTGTCCCCCTGGACCCCTCCACCACCCGCTCCTCGGTGCTCACCCCTATCCCACA
GCCCCAGGACCATGGCACCAGCCTCACCTGTCAGGTGACCTTCCCTGGGGCCAGCGTGACCACGA
ACAAGACCGTCCATCTCAACGTGTCTTACCCGCTCAGAACTTGACCATGACTGTCTTCCAAGGA
GACGGCACAGTATCCACAGTCTTGGGAAATGGCTCATCTCTGTCACTCCCAGAGGGCCAGTCTCT
GCGCCTGGTCTGTGCAGTTGATGCAGTTGACAGCAATCCCCCTGCCAGGCTGAGCCTGAGCTGGA
GAGGCCTGACCCTGTGCCCTCACAGCCCTCAAACCCGGGGGTGCTGGAGCTGCCTTGGGTGCAC
CTGAGGGATGCAGCTGAATTCACCTGCAGAGCTCAGAACCCCTCTCGGCTCTCAGCAGGTCTACCT
GAACGTCTCCCTGCAGAGCAAAGCCACATCAGGAGTGAATCAGGGGGTGGTCCGGGGGAGCTGGAG
CCACAGCCCTGGTCTTCTGTCTTCTGCGTCATCTTCGTTGTAGTGAGGTCTTGCAGGAAGAAA
TCGGCAAGGCCAGCAGCGGGCGTGGGAGATACGGGCATAGAGGATGCAAACGCTGTCAGGGGTTC
AGCCTCTCAGGGGCCCCCTGACTGAACCTTGGGCAGAAGACAGTCCCCCAGACCAGCTCCCCCAG
CTTCTGCCCCGCTCCTCAGTGGGGGAAGGAGAGCTCCAGTATGCATCCCTCAGCTTCCAGATGGTG
AAGCCTTGGGACTCGCGGGGACAGGAGGCCACTGACACCGAGTACTCGGAGATCAAGATCCACAG
ATGAGAACTGCAGAGACTCACCTGATTGAGGGATCACAGCCCCTCCAGGCAAGGGAGAAGTCA
GAGGCTGATTCTTGTAGAATTAACAGCCCTCAACGTGATGAGCTATGATAACACTATGAATTATG
TGCAGAGTGAAAAGCACACAGGCTTTAGAGTCAAAGTATCTCAAACCTGAATCCACACTGTGCCC
TCCCTTTTATTTTTTAACTAAAAGACAGACAAATTCCTA

FIGURE 86

MLLLLLLPLLWGRERAEGQTSKLLTMQSSVTVQEGLCVHVPCSFSPSHGWIYPGPVVHGYWFREG
ANTDQDAPVATNNPARAVWEETRDRFHLLGDPHTKNCTLSIRDARRSDAGRYFFRMEKGSIKWNY
KHHRLSVNVTALTHRPNILIPGTLESGCPQNLTCVWPWACEQGTPPMISWIGTSVSPLDPSTTRS
SVLTLLIPQPQDHGTS LTCQVTFFPGASVTTNKT VHLNVSYPQNLMTVFQGDGTVSTVLGNGSSL
SLPEGQSLRLVCAVDAVDSNPPARLSLSWRGLTLCPSQPSNPGVLELPWWHLRDAAEFTCRAQNP
LGSQQVYLNVS LQSKATSGVTQGVVGGAGATALVFLSFCVIFVVRSCRKKSARPAAGVGD TGIE
DANAVRGSASQGPLEPWAEDSPPDQPPPASARSSVGE GELQYASLSFQMKPWDSRGQEATDTE
YSEIKIHR

Signal peptide:

amino acids 1-15

Transmembrane domain:

amino acids 351-370

FIGURE 87

AGAAAGCTGCACTCTGTTGAGCTCCAGGGCGCAGTGGAGGGAGGGAGTGAAGGAGCTCTCTGTAC
CCAAGGAAAGTGCAGCTGAGACTCAGACAAGATTACAATGAACCAACTCAGCTTCCTGCTGTTTC
TCATAGCGACCACCAGAGGATGGAGTACAGATGAGGCTAATACTTACTTCAAGGAATGGACCTGT
TCTTCGTCTCCATCTCTGCCCAGAAGCTGCAAGGAAATCAAAGACGAATGTCCTAGTGCATTTGA
TGGCCTGTATTTTCTCCGCACTGAGAATGGTGTATCTACCAGACCTTCTGTGACATGACCTCTG
GGGGTGGCGGCTGGACCCTGGTGGCCAGCGTGCATGAGAATGACATGCGTGGGAAGTGCACGGTG
GGCGATCGCTGGTCCAGTCAGCAGGGCAGCAAAGCAGACTACCCAGAGGGGGACGGCAACTGGGC
CAACTACAACACCTTTGGATCTGCAGAGGCGGCCACGAGCGATGACTACAAGAACCCTGGCTACT
ACGACATCCAGGCCAAGGACCTGGGCATCTGGCACGTGCCCAATAAGTCCCCCATGCAGCACTGG
AGAAACAGCTCCCTGCTGAGGTACCGCACGGACACTGGCTTCCTCCAGACACTGGGACATAATCT
GTTTGGCATCTACCAGAAATATCCAGTGAATATGGAGAAGGAAAGTGTGGACTGACAACGGCC
CGGTGATCCCTGTGGTCTATGATTTTGGCGACGCCCAGAAAACAGCATCTTATTACTCACCTAT
GGCCAGCGGGAATTCAGTGCGGGATTTGTTCA GTTCAGGGTATTTAATAACGAGAGAGCAGCCAA
CGCCTTGTGTGCTGGAATGAGGGTCACCGGATGTAACACTGAGCATCACTGCATTGGTGGAGGAG
GATACTTTCCAGAGGCCAGTCCCCAGCAGTGTGGAGATTTTCTGGTTTTGATTGGAGTGGATAT
GGA²ACTCATGTTGGTTACAGCAGCAGCCGTGAGATAACTGAGGCAGCTGTGCTTCTATTCTATCG
TTGAGAGTTTTGTGGGAGGGAACCCAGACCTCTCCTCCCAACCATGAGATCCCAAGGATGGAGAA
CAACTACCCAGTAGCTAGAATGTTAATGGCAGAAGAGAAAACAATAAATCATATTGACTCAAGA
AAAAAA

FIGURE 88

MNQLSFLLFLIATTRGWSTDEANTYFKEWTCSSSPSLPRSCKEIKDECPSAFDGLYFLRTENGVI
YQTFCDMTSGGGGWTLVASVHENDMRGKCTVGDRWSSQOGSKADYPEGDGNWANYNTFGSAEAAT
SDDYKNPGYYDIQAKDLGIWHVPNKSPMQHWRNSSLLRYRTDTGFLQTLGHNLFGIYQKYPVKYG
EGKCWTDNGPVI PVVYDFGDAQKTASYYSPIYGQREFTAGFVQFRVFNNERAANALCAGMRVTGCN
TEHHCIGGGGYFPEASPOQCGDFSGFDWSGYGTHVGYSSSREITEAAVLLFYR

Important features:

Signal peptide:

amino acids 1-16

N-glycosylation site.

amino acids 163-167

Glycosaminoglycan attachment sites.

amino acids 74-78, 289-293

N-myristoylation sites.

amino acids 76-82, 115-121, 124-130, 253-259, 292-298

FIGURE 89

CTAGATTTGTGGCTTGCGGGGAGACTTCAGGAGTCGCTGTCTCTGAACTTCCAGCCTCAGAGAC
CGCCGCCCTTGTCCCCGAGGGCCATGGGCCGGGTCTCAGGGCTTGTGCCCTCTCGCTTCCTGACG
CTCCTGGCGCATCTGGTGGTCGTCATCACCTTATTCTGGTCCCGGGACAGCAACATACAGGCCTG
CCTGCCTCTCACGTTCACCCCGAGGAGTATGACAAGCAGGACATTACAGCTGGTGGCCGCGCTCT
CTGTCACCCTGGGCCTCTTTGCAGTGGAGCTGGCCGGTTTCCTCTCAGGAGTCTCCATGTTCAAC
AGCACCCAGAGCCTCATCTCCATTGGGGCTCACTGTAGTGCATCCGTGGCCCTGTCCTTCTTCAT
ATTCGAGCGTTGGGAGTGCACTACGTATTGGTACATTTTTGTCTTCTGCAGTGCCCTTCCAGCTG
TCACTGAAATGGCTTTATTTCGTCACCGTCTTTGGGCTGAAAAGAAACCTTCTTGATTACCTTCA
TGACGGGAACCTAAGGACGAAGCCTACAGGGGCAAGGGCCGCTTCGTATTCCTGGAAGAAGGAAG
GCATAGGCTTCGGTTTTCCCCTCGGAACTGCTTCTGCTGGAGGATATGTGTTGGAATAATTACG
TCTTGAGTCTGGGATTATCCGCATTGTATTTAGTGCTTTGTAATAAAATATGTTTTGTAGTAACA
TTAAGACTTATATACAGTTTTAGGGGACAATTAAAAAAAAAAAA

FIGURE 90

MGRVSGLVPSRFLTLLAHLVVVITLFWSRDSNIQACLPLTFTPEEYDKQDIQLVAALSVTLGLFA
VELAGFLSGVSMFNSTQSLISIGAHCSASVALSFFIFERWECTTYWYIFVFCSALPAVTEMALFV
TVFGLKKKPF

Transmembrane domain:

amino acids 12-28 (type II), 51-66, 107-124

FIGURE 91

CTGGGACCCCGAAAAGAGAAGGGGAGAGCGAGGGGACGAGAGCGGAGGAGGAAGATGCAACTGAC
TCGCTGCTGCTTCGTGTTCCCTGGTGCAGGGTAGCCTCTATCTGGTCATCTGTGGCCAGGATGATG
GTCCTCCCGGCTCAGAGGACCCTGAGCGTGATGACCACGAGGGCCAGCCCCGGCCCCGGGTGCCT
CGGAAGCGGGGCCACATCTCACCTAAGTCCCGCCCCATGGCCAATTCCACTCTCCTAGGGCTGCT
GGCCCCGCCTGGGGAGGCTTGGGGCATTCTTGGGCAGCCCCCAACCGCCCCGAACCACAGCCCCC
CACCCTCAGCCAAGGTGAAGAAAATCTTTGGCTGGGGCGACTTCTACTCCAACATCAAGACGGTG
GCCCTGAACCTGCTCGTCAAGGGAAGATTGTGGACCATGGCAATGGGACCTTCAGCGTCCACTT
CCAACACAATGCCACAGGCCAGGGAACATCTCCATCAGCCTCGTGCCCCCAGTAAAGCTGTAG
AGTTCCACCAGGAACAGCAGATCTTCATCGAAGCCAAGGCCTCCAAAATCTTCAACTGCCGGATG
GAGTGGGAGAAGGTAGAACGGGGCCGCCGACCTCGCTTTCACCCACGACCCAGCCAAGATCTG
CTCCCGAGACCACGCTCAGAGCTCAGCCACCTGGAGCTGCTCCAGCCCTTCAAAGTCGTCTGTG
TCTACATCGCCTTCTACAGCACGGACTATCGGCTGGTCCAGAAGGTGTGCCCAGATTACAACCTAC
CATAGTGATACCCCCTACTACCCATCTGGGTGAACCCGGGGCAGGCCACAGAGGCCAGGCCAGGGC
TGGGAAGGACAGGCCTGCCCATGCAGGAGACCATCTGGACACCGGGCAGGGAAGGGGTTGGGCCTC
AGGCAGGGAGGGGGGTGGAGACGAGGAGATGCCAAGTGGGGCCAGGGCCAAGTCTCAAGTGGCAG
AGAAAGGGTCCCAAGTGCTGGTCCCAACCTGAAGCTGTGGAGTGACTAGATCACAGGAGCACTGG
AGGAGGAGTGGGCTCTCTGTGCAGCCTCACAGGGCTTTGCCACGGAGCCACAGAGAGATGCTGGG
TCCCCGAGGCCTGTGGGCAGGCCGATCAGTGTGGCCCCAGATCAAGTCATGGGAGGAAGCTAAGC
CCTTGGTTCTTGCCATCCTGAGGAAAGATAGCAACAGGGAGGGGGAGATTTTCATCAGTGTGGACA
GCCTGTCAACTTAGGATGGATGGCTGAGAGGGCTTCCTAGGAGCCAGTCAGCAGGGTGGGGTGGG
GCCAGAGGAGCTCTCCAGCCCTGCCTAGTGGGCGCCCTGAGCCCTTGTCGTGTGCTGAGCATGG
CATGAGGCTGAAGTGGCAACCCTGGGGTCTTTGATGTCTTGACAGATTGACCATCTGTCTCCAGC
CAGGCCACCCCTTTCCAAAATTCCTCTTCTGCCAGTACTCCCCCTGTACCACCCATTGCTGATG
GCACACCCATCCTTAAGCTAAGACAGGACGATTGTGGTCCTCCACACTAAGGCCACAGCCCATC
CGCGTGCTGTGTGTCCCTCTTCCACCCCAACCCCTGCTGGCTCCTCTGGGAGCATCCATGTCCCG
GAGAGGGGTCCCTCAACAGTCAGCCTCACCTGTCAGACCGGGGTCTCCCGGATCTGGATGGCGC
CGCCCTCTCAGCAGCGGGCACGGGTGGGGCGGGGCCGGCCGAGAGCATGTGCTGGATCTGTTC
TGTGTGTCTGTCTGTGGGTGGGGGAGGGGAGGGAAGTCTTGTGAAACCGCTGATTGCTGACTTT
TGTGTGAAGAATCGTGTTCTTGGAGCAGGAATAAAGCTTGCCCCGGGGCA

FIGURE 92

MQLTRCCFVFLVQGSLYLVICGQDDGPPGSEDPERDDHEGQPRPRVPRKRGHISPKSRPMANSTL
LGLLAPPGEAWGILGQPPNRPNHSPPPSAKVKKIFGWGDFYSNIKTVALNLLVTGKIVDHGNGTF
SVHFQHNATGQGNISISLVPPSKAVEFHQEQQIFIEAKASKIFNCRMWEKVERGRRTSLCTHDP
AKICSRDHAQSSATWSCSQPFKVVCVYIAFYSTDYRLVQKVCPDINYHSDTPYYPSG

Important features of the protein:

Signal peptide:

amino acids 1-14

N-glycosylation sites.

amino acids 62-65, 127-130, 137-140, 143-146

2-oxo acid dehydrogenases acyltransferase

amino acids 61-71

FIGURE 93

CGGTGGCCATGACTGCGGCCGTGTTCTTCGGCTGCGCCTTCATTGCCTTCGGGCCTGCGCTCGCC
CTTTATGTCTTCACCATCGCCATCGAGCCGTTGCGTATCATCTTCCTCATCGCCGGAGCTTTCTT
CTGGTTGGTGTCTCTACTGATTTTCGTCCCTTGTTTGGTTTCATGGCAAGAGTCATTATTGACAACA
AAGATGGACCAACACAGAAATATCTGCTGATCTTTGGAGCGTTTGTCTCTGTCTATATCCAAGAA
ATGTTCCGATTTGCATATTATAAACTCTTAAAAAAGCCAGTGAAGGTTTGAAGAGTATAAACCC
AGGTGAGACAGCACCCCTCTATGCGACTGCTGGCCTATGTTTCTGGCTTGGGCTTTGGAATCATGA
GTGGAGTATTTTCCTTTGTGAATACCCTATCTGACTCCTTGGGGCCAGGCACAGTGGGCATTTCAT
GGAGATTCTCCTCAATTCTTCCTTTATTCAGCTTTCATGACGCTGGTCATTATCTTGCTGCATGT
ATTCTGGGGCATTGTATTTTTTGATGGCTGTGAGAAGAAAAAGTGGGGCATCCTCCTTATCGTTC
TCCTGACCCACCTGCTGGTGTGAGCCAGACCTTCATAAGTTCTTATTATGGAATAAACCTGGCG
TCAGCATTTATAATCCTGGTGCTCATGGGCACCTGGGCATTCTTAGCTGCGGGAGGCAGCTGCCG
AAGCCTGAAACTCTGCCTGCTCTGCCAAGACAAGAACTTTCTTCTTTACAACCAGCGCTCCAGAT
AACCTCAGGGAACCAGCACTTCCCAAACCGCAGACTACATCTTTAGAGGAAGCACAACTGTGCCT
TTTTCTGAAAATCCCTTTTTCTGGTGGAATTGAGAAAGAAATAAACTATGCAGATA

FIGURE 94

MTAAVFFGCAFIAGFPALALYVFTIAIEPLRIIFLIAGAFFWLVSLLISSLVWFEMARVIIDNKDG
PTQKYLLIFGAFVSVYIQEMFRFAYYKLLKKASEGLKSINPGETAPSMRLLAYVSGLGFGIMSGV
FSFVNTLSDSLGPSTVGIHGDSPQFFLYSAFMTLVIILLHVFWGIVFFDGCEKKKWGILLIVLLT
HLLVSAQTFISSYYGINLASAFIILVLMGTWAFILAAGGSCRSCLKCLLCQDKNFLLYNQRSR

Important features of the protein:

Signal peptide:

amino acids 1-19

Transmembrane domains:

amino acids 32-51, 119-138, 152-169, 216-235

Glycosaminoglycan attachment site.

amino acids 120-123

Sodium:neurotransmitter symporter family protein

amino acids 31-65

FIGURE 95

AATTTTTCACCAGAGTAAACTTGAGAAACCAACTGGACCTTGAGTATTGTACATTTTGCCTCGTG
GACCCAAAGGTAGCAATCTGAAACATGAGGAGTACGATTCTACTGTTTTGTCTTCTAGGATCAAC
TCGGTCATTACCACAGCTCAAACCTGCTTTGGGACTCCCTCCCACAAAAGTGGCTCCGGATCAGG
GAACACTACCAAACCAACAGCAGTCAAATCAGGTCTTTCCTTCTTTAAGTCTGATACCATTAAACA
CAGATGCTCACACTGGGGCCAGATCTGCATCTGTTAAATCCTGCTGCAGGAATGACACCTGGTAC
CCAGACCCACCCATTGACCCTGGGAGGGTTGAATGTACAACAGCAACTGCACCCACATGTGTTAC
CAATTTTTTGTACACAACCTTGAGGCCAGGGCACTATCCTAAGCTCAGAGGAATTGCCACAAATC
TTCACGAGCCTCATCATCCATTTCCTTGTTCCCGGGAGGCATCCTGCCCACCAGTCAGGCAGGGGC
TAATCCAGATGTCCAGGATGGAAGCCTTCCAGCAGGAGGAGCAGGTGTAAATCCTGCCACCCAGG
GAACCCAGCAGGCCGCCTCCCAACTCCCAGTGGCACAGATGACGACTTTGCAGTGACCACCCCT
GCAGGCATCCAAAGGAGCACACATGCCATCGAGGAAGCCACCACAGAATCAGCAAATGGAATTCA
GTAAGCTGTTTCAAATTTTTTCAACTAAGCTGCCTCGAATTTGGTGATACATGTGAATCTTTATC
ATTGATTATATTATGGAATAGATTGAGACACATTGGATAGTCTTAGAAGAAATTAATTCTTAATT
TACCTGAAAATATTCTTGAAATTTCAGAAAATATGTTCTATGTAGAGAATCCCAACTTTTAAAAA
CAATAATTCAATGGATAAATCTGTCTTTGAAATATAACATTATGCTGCCTGGATGATATGCATAT
TAAAACATATTTGAAAACTGAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 96

MRSTILLEFCLLGSTRSLPQLKPALGLPPTKLAPDQGTLPNQQQSNQVFPSLSLIPLTQM
LTLGPDHLHLLNPAAGMTPGTQTHPLTLGGLNVQQQLHPHVLPIFVTQLGAQGTLISSEE
LPQIFTSLLIHSLEFPGGILPTSQAGANPDVQDGLPAGGAGVNPATQGTPAGRLPTPSG
TDDDFAVTTPAGIQRSTHAIEEATTESANGIQ

Signal peptide:

amino acids 1-16

FIGURE 97

GCTCAAGTGCCCTGCCTTGCCCCACCCAGCCCAGCCTGGCCAGAGCCCCCTGGAGAAGGAGCTCT
 CTTCTTGCTTGGCAGCTGGACCAAGGGAGCCAGTCTTGGGCGCTGGAGGGCCTGTCTGACCATG
 GTCCCTGCCTGGCTGTGGCTGCTTTGTGTCTCCGTCCCCCAGGCTCTCCCCAAGGCCAGCCTGC
 AGAGCTGTCTGTGGAAGTTCCAGAAAACCTATGGTGGAAATTTCCCTTTATACCTGACCAAGTTGC
 CGCTGCCCCGTGAGGGGGCTGAAGGCCAGATCGTGCTGTCAGGGGACTCAGGCAAGGCAACTGAG
 GGCCCATTTGCTATGGATCCAGATTCTGGCTTCCTGCTGCTGACAGGGCCCTGGACCGAGAGGA
 GCAGGCAGAGTACCAGCTACAGGTCACCCTGGAGATGCAGGATGGACATGTCTTGTGGGGTCCAC
 AGCCTGTGCTTGTGCACGTGAAGGATGAGAATGACCAGGTGCCCCATTTCTCTCAAGCCATCTAC
 AGAGCTCGGCTGAGCCGGGGTACCAGGCCTGGCATCCCCCTCCTCTTCCTTGAGGCTTCAGACCG
 GGATGAGCCAGGCACAGCCAACTCGGATCTTCGATTCCACATCCTGAGCCAGGCTCCAGCCCAGC
 CTTCCCCAGACATGTTCCAGCTGGAGCCTCGGCTGGGGGCTCTGGCCCTCAGCCCCAAGGGGAGC
 ACCAGCCTTGACCACGCCCTGGAGAGGACCTACCAGCTGTTGGTACAGGTCAAGGACATGGGTGA
 CCAGGCCTCAGGCCACCAGGCCACTGCCACCGTGGAAAGTCTCCATCATAGAGAGCACCTGGGTGT
 CCCTAGAGCCTATCCACCTGGCAGAGAATCTCAAAGTCTATACCCGCACCATGAGCCAGGTA
 CACTGGAGTGGGGGTGATGTGCACTATCACCTGGAGAGCCATCCCCGGGACCCCTTTGAAGTGAA
 TGCAGAGGGAAACCTCTACGTGACCAGAGAGCTGGACAGAGAAGCCCAGGCTGAGTACCTGCTCC
 AGGTGCGGGCTCAGAATTCCCATGGCGAGGACTATGCGGCCCTCTGGAGCTGCACGTGCTGGTG
 ATGGATGAGAATGACAACGTGCCTATCTGCCCTCCCCGTGACCCACAGTCAGCATCCCTGAGCT
 CAGTCCACCAGGTACTGAAGTGACTAGACTGTGAGCAGAGGATGCAGATGCCCCCGGCTCCCCCA
 ATTTCCACGTTGTGTATCAGCTCCTGAGCCCTGAGCCTGAGGATGGGGTAGAGGGGAGAGCCTTC
 CAGGTGGACCCCACTTCAGGCAGTGTGACGCTGGGGGTGCTCCCACTCCGAGCAGGCCAGAACAT
 CCTGCTTCTGGTGTGGCCATGGACCTGGCAGGCGCAGAGGGTGGCTTCAGCAGCACGTGTGAAG
 TCGAAGTCGCAGTCACAGATATCAATGATCACGCCCTGAGTTCATCACTTCCAGATTGGGCCT
 ATAAGCCTCCCTGAGGATGTGGAGCCCCGGGACTCTGGTGGCCATGCTAACAGCCATTGATGCTGA
 CCTCGAGCCCGCCTTCGCGCTCATGGATTTTGCCATTGAGAGGGGAGACACAGAAGGGACTTTTG
 GCCTGGATTGGGAGCCAGACTCTGGGCATGTTAGACTCAGACTCTGCAAGAACCTCAGTTATGAG
 GCAGCTCCAAGTCATGAGGTGGTGGTGGTGGTGCAGAGTGTGGCGAAGCTGGTGGGGCCAGGCCC
 AGGCCCTGGAGCCACCGCCACGGTGACTGTGCTAGTGGAGAGAGTGATGCCACCCCCCAAGTTGG
 ACCAGGAGAGCTACGAGGCCAGTGTCCCCATCAGTGCCCCAGCCGGCTCTTTCTGTGCTGACCATC
 CAGCCCTCCGACCCCATCAGCCGAACCTCAGGTTCTCCCTAGTCAATGACTCAGAGGGCTGGCT
 CTGCATTGAGAAATTCTCCGGGGAGGTGCACACCGCCCAGTCCCTGCAGGGCGCCAGCCTGGGG
 ACACCTACACGGTGCTTGTGGAGGCCCAGGATACAGCCCTGACTCTTGCCCCCTGTGCCCTCCCAA
 TACCTCTGCACACCCCGCCAAGACCATGGCTTGATCGTGAGTGGACCCAGCAAGGACCCCGATCT
 GGCCAGTGGGCACGGTCCCTACAGCTTACCCTTGGTCCCAACCCACGGTGCAACGGGATTGGC
 GCCTCCAGACTCTCAATGGTTCCCATGCCTACCTCACCTTGGCCCTGCATTGGGTGGAGCCACGT
 GAACACATAATCCCGTGGTGGTTCAGCCACAATGCCAGATGTGGCAGCTCCTGGTTTCAGATGAT
 CGTGTGTCGCTGCAACGTGGAGGGGAGTGCATGCGCAAGGTGGGCCGCATGAAGGGCATGCCCA
 CGAAGCTGTGCGCAGTGGGCATCCTTGTAGGCACCTGGTAGCAATAGGAATCTTCCTCATCCTC
 ATTTTCACCCACTGGACCATGTCAAGGAAGAAGGACCCGGATCAACCAGCAGACAGCGTGCCCT
 GAAGGCGACTGTCTGAATGGCCCAGGCAGCTCTAGCTGGGAGCTTGGCCTCTGGCTCCATCTGAG
 TCCCCCTGGGAGAGAGCCAGCACCCAAAGATCCAGCAGGGGACAGGACAGAGTAGAAGCCCCCTCCA
 TCTGCCCTGGGGTGGAGGCACCATCACCATCACCAGGCATGTCTGCAGAGCCTGGACACCAACTT
 TATGGACTGCCCATGGGAGTGCTCCAAATGTGAGGGTGTTTGCCCAATAATAAAGCCCCAGAGAA
 CTGGGCTGGGCCCTATGGGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAG

FIGURE 98

MVPAWLWLLCVSVPQALPKAQPAELSVEVPENYGGNFPLYLTKLPLPREGAEGQIVLSGDSGKAT
EGPFAMDPDSGFLLVTRALDREEQAEYQLQVTLEMQDGHVLWGPQPVLVHVKDENDQVPHFSQAI
YRARLSRGTRPGIPFLFLEASDRDEPGTANSDLRFHILSQAPAQSPDMFQLEPRLGALALSPKG
STSLDHALERTYQLLVQVKMDGDQASGHQATATVEVSI IESTWVSLEPIHLAENLKVLYPHHMAQ
VHWSGGDVHYHLESHPPGPFEEVNAEGNLYVTRELDREAQAEYLLQVRAQNSHGEDIAAPLELHVL
VMDENDNVPICPPRDPTVSIPELSPPGTEVTRLAEDADAPGSPNSHVYQLLSPEPEDGVEGRA
FQVDPTSGSVTLGVLPLRAGQNILLVLAMDLAGAEGGFSSTCEVEVAVTDINDHAPEFITSQIG
PISLPEDVEPGTLVAMLT AIDADLEPAFRMLDFAIERGDTEGTFGLDWEPDSGHVRLRLCKNLSY
EAAPSHEVVVVVQSVAKLVGPGPGPGATATVTVLVERVMPPPKLDQESYEASVPISAPAGSFLLT
IQPSDPISRTLRFSLVNDSEGWLCIEKFSGEVHTAQSLQGAQPGDTYTVLVEAQDTALT LAPVPS
QYLCTPRQDHGLIVSGPSKDPDLASGHGPYSFTLGPNPTVQRDWRLQTLNGSHAYLTLALHWVEP
REHIIPVVVSHNAQMWQLLVRVIVCRCNVEGQCMRKVGRMKGMPTKLSAVGILVGTLVAIGIFLI
LIFTHWTMSRKKDPDQPADSVPLKATV

Signal peptide:

amino acids 1-18

Transmembrane domain:

amino acids 762-784

FIGURE 99

GGCTGACCGTGCTACATTGCCTGGAGGAAGCCTAAGGAACCCAGGCATCCAGCTGCCCACGCCTG
AGTCCAAGATTCTTCCCAGGAACACAAACGTAGGAGACCCACGCTCCTGGAAGCACCAGCCTTTA
TCTCTTCACCTTCAAGTCCCCCTTCTCAAGAATCCTCTGTTCTTTGCCCTCTAAAGTCTTGGTAC
ATCTAGGACCCAGGCATCTTGCTTTCCAGCCACAAAGAGACAGATGAAGATGCAGAAAGGAAATG
TTCTCCTTATGTTTGGTCTACTATTGCATTTAGAAGCTGCAACAAATTCCAATGAGACTAGCACC
TCTGCCAACACTGGATCCAGTGTGATCTCCAGTGGAGCCAGCACAGCCACCAACTCTGGGTCCAG
TGTGACCTCCAGTGGGGTCAGCACAGCCACCATCTCAGGGTCCAGCGTGACCTCCAATGGGGTCA
GCATAGTCACCAACTCTGAGTTCATACAACCTCCAGTGGGATCAGCACAGCCACCAACTCTGAG
TTCAGCACAGCGTCCAGTGGGATCAGCATAGCCACCAACTCTGAGTCCAGCACAACTCCAGTGG
GGCCAGCACAGCCACCAACTCTGAGTCCAGCACACCCTCCAGTGGGGCCAGCACAGTCCACCAACT
CTGGGTCCAGTGTGACCTCCAGTGGAGCCAGCACTGCCACCAACTCTGAGTCCAGCACAGTGTCC
AGTAGGGCCAGCACTGCCACCAACTCTGAGTCTAGCACACTCTCCAGTGGGGCCAGCACAGCCAC
CAACTCTGACTCCAGCACAACTCCAGTGGGGCTAGCACAGCCACCAACTCTGAGTCCAGCACAA
CCTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTCCAGCACAGTGTCCAGTAGGGCCAGCACT
GCCACCAACTCTGAGTCCAGCACAACTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTCCAG
AACGACCTCCAATGGGGCTGGCACAGCCACCAACTCTGAGTCCAGCACGACCTCCAGTGGGGCCA
GCACAGCCACCAACTCTGACTCCAGCACAGTGTCCAGTGGGGCCAGCACTGCCACCAACTCTGAG
TCCAGCACGACCTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTCCAGCACGACCTCCAGTGG
GGCTAGCACAGCCACCAACTCTGACTCCAGCACAACTCCAGTGGGGCCGGCACAGCCACCAACT
CTGAGTCCAGCACAGTGTCCAGTGGGATCAGCACAGTCCACCAATTCTGAGTCCAGCACACCCTCC
AGTGGGGCCAACACAGCCACCAACTCTGAGTCCAGTACGACCTCCAGTGGGGCCAACACAGCCAC
CAACTCTGAGTCCAGCACAGTGTCCAGTGGGGCCAGCACTGCCACCAACTCTGAGTCCAGCACAA
CCTCCAGTGGGGTCCAGCACAGCCACCAACTCTGAGTCCAGCACAACTCCAGTGGGGCTAGCACA
GCCACCAACTCTGACTCCAGCACAACTCCAGTGGGGCCAGCACAGCCACCAACTCTGAGTCTAG
CACAGTGTCCAGTGGGATCAGCACAGTCCACCAATTCTGAGTCCAGCACAACTCCAGTGGGGCCA
ACACAGCCACCAACTCTGGGTCCAGTGTGACCTCTGCAGGCTCTGGAACAGCAGCTCTGACTGGA
ATGCACACAACTTCCCATAGTGCATCTACTGCAGTGAGTGAGGCAAAGCCTGGTGGGTCCCTGGT
GCCGTGGGAAATCTTCCTCATCACCTGGTCTCGGTTGTGGCGGCCGTGGGGCTCTTTGCTGGGC
TCTTCTTCTGTGTGAGAAACAGCCTGTCCCTGAGAAACACCTTTAACACAGCTGTCTACCACCT
CATGGCCTCAACCATGGCCTTGGTCCAGGCCCTGGAGGGAATCATGGAGCCCCCAGAGGCCAG
GTGGAGTCCCTAACTGGTTCTGGAGGAGACCAAGTATCATCGATAGCCATGGAGATGAGCGGGAGGA
ACAGCGGGCCCTGAGCAGCCCCGGAAGCAAGTGCCGCATTCTTCAGGAAGGAAGAGACCTGGGCA
CCCAAGACCTGGTTTCCTTTCATTCATCCCAGGAGACCCCTCCAGCTTTGTTTGAGATCCTGAA
AATCTTGAAGAAGGTATTCCTCACCTTTCTTGCTTTACCAGACACTGGAAAGAGAATACTATAT
TGCTCATTTAGCTAAGAAATAAATACATCTCATCTAACACACACGACAAAGAGAAGCTGTGCTTG
CCCCGGGGTGGGTATCTAGCTCTGAGATGAACTCAGTTATAGGAGAAAACCTCCATGCTGGACTC
CATCTGGCATTCAAAATCTCCACAGTAAATCCAAAGACCTCAAAAAAAAAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 100

MKMQKGNVLLMFGLLLHLEAATNSNETSTSANTGSSVSSGASTATNSGSSVTSSGVSTATISGS
SVTSNGVSIIVTNSEFHTTSSGISTATNSEFSTASSGISIATNSESSTTSSGASTATNSESSTPSS
GASTVTNSGSSVTSSGASTATNSESSTVSSRASTATNSESSTLSSGASTATNSDSSTTSSGASTA
TNSESSTTSSGASTATNSESSTVSSRASTATNSESSTTSSGASTATNSESRTTNGAGTATNSES
STTSSGASTATNSDSSTVSSGASTATNSESSTTSSGASTATNSESSTTSSGASTATNSDSSTTSS
GAGTATNSESSTVSSGISTVTNSESSTPSSGANTATNSESSTTSSGANTATNSESSTVSSGASTA
TNSESSTTSSGVSTATNSESSTTSSGASTATNSDSSTTSSEASTATNSESSTVSSGISTVTNSES
STTSSGANTATNSGSSVTSAGSGTAALTGMHTTSHSASTAVSEAKPGGSLVPWEIFLITLVSVA
AVGLFAGLFFCVRNSLSLRNTFNTAVYHPHGLNHGLGPGPGGNHGAPHRPRWSPNWFWRPVSII
AMEMSGRNSGP

Signal peptide:

amino acids 1-20

Transmembrane domain:

amino acids 510-532

FIGURE 101

GGCCGGACGCCTCCGCGTTACGGGATGAATTAACGGCGGGTTCCGCACGGAGGTTGTGACCCCTA
CGGAGCCCCAGCTTGCCACGCACCCCACTCGGCGTCGCGCGGCGTGCCCTGCTTGTACAGGTG
GGAGGCTGGAAGTATCAGGCTGAAAAACAGAGTGGGTACTCTCTTCTGGGAAGCTGGCAACAAAT
GGATGATGTGATATATGCATTCCAGGGGAAGGGAAATTGTGGTGCTTCTGAACCCATGGTCAATT
AACGAGGCAGTTTCTAGCTACTGCACGTACTTCATAAAGCAGGACTCTAAAAGCTTTGGAATCAT
GGTGTGATGGAAGGGATTTACTTTTATACTGACTCTGTTTTGGGGAAGCTTTTTTGAAGCATTT
TCATGCTGAGTCCCTTTTTTACCTTTGATGTTTGTAACCCATCTTGGTATCGCTGGATCAACAAC
CGCCTTGTGGCAACATGGCTCACCTACCTGTGCCATTATTGGAGACCATGTTTGGTGTAAAAGT
GATTATAACTGGGGATGCATTTGTTCTTGAGAAAGAAGTGTGATTATCATGAACCATCGGACAA
GAATGGACTGGATGTTCTGTGGAATTGCCTGATGCGATATAGCTACCTCAGATTGGAGAAAAAT
TGCCTCAAAGCGAGTCTCAAAGGTGTTCTTGATTGTGGTTGGGCCATGCAGGCTGCTGCCTATAT
CTTCATTTCATAGGAAATGGAAGGATGACAAGAGCCATTCGAAGACATGATTGATTACTTTTGTG
ATATTCACGAACCACTTCAACTCCTCATATTCCCAGAAGGGACTGATCTCACAGAAAACAGCAAG
TCTCGAAGTAATGCATTTGCTGAAAAAATGGACTTCAGAAATATGAATATGTTTTACATCCAAG
AACTACAGGCTTTACTTTTGTGGTAGACCGTCTAAGAGAAGGTAAGAACCTTGATGCTGTCCATG
ATATCACTGTGGCGTATCCTCACAACATTCTCAATCAGAGAAGCACCTCCTCCAAGGAGACTTT
CCCAGGGAAATCCACTTTCACGTCCACCGGTATCCAATAGACACCCTCCCCACATCCAAGGAGGA
CCTTCAACTCTGGTGCCACAAACGGTGGGAAGAGAAAGAAGAGAGGCTGCGTTCCTTCTATCAAG
GGGAGAAGAATTTTTATTTTACCGGACAGAGTGTGATTCCACCTTGCAAGTCTGAACTCAGGGTC
CTTGTGGTCAAATTGCTCTCTATACTGTATTGGACCCTGTTTCAGCCCTGCAATGTGCCTACTCAT
ATATTTGTACAGTCTTGTTAAGTGGTATTTTATAATCACCATTGTAATCTTTGTGCTGCAAGAGA
GAATATTTGGTGGACTGGAGATCATAGAACTTGCATGTTACCGACTTTTACACAAACAGCCACAT
TTAAATTCAAAGAAAAATGAGTAAGATTATAAGGTTGCCATGTGAAAACCTAGAGCATATTTTG
GAAATGTTCTAAACCTTTCTAAGCTCAGATGCATTTTGCATGACTATGTGCAATATTTCTTACT
GCCATCATTATTTGTTAAAGATATTTTGCACTTAATTTTGTGGGAAAAATATTGCTACAATTTT
TTTAATCTCTGAATGTAATTTTGATACTGTGTACATAGCAGGGAGTGATCGGGGTGAAATAACTT
GGGCCAGAATATTATTAAACAATCATCAGGCTTTTAAA

FIGURE 102

MHSRGREIVVLLNPWSINEAVSSYCTYFIKQDSKSGIMVSWKGIYFILTLFWGSFFGSIFMLSP
FLPLMFVNPSWYRWINNRLVATWLTLPVALLETMFGVKVIITGDAFVPGERSVIIMNHRTRMDWM
FLWNCLMRYSYLRLEKICLKASLKGVPGEFGWAMQAAAYIFIHRKWKDDKSHFEDMIDYFCDIHEP
LQLLIFPEGTDLTENSKSRSNFAEKNGLQKYEYVLHPRTTGFTFVVDRLREGKNLDAVHDITVA
YPHNIPQSEKHLQGDFFPREIHFHVHRYPIDTLPTSKEDLQLWCHKRWEEKEERLRSFYQGEKNF
YFTGQSVIPPCKSELRVLVVKLLSILYWTLEFSPAMCLLIYLYSLVKWYFIITIVIFVLQERIFGG
LEIIEIACYRLLHKQPHLNSKKNE

Important features of the protein:

Signal peptide:

amino acids 1-22

Transmembrane domains:

amino acids 44-63, 90-108, 354-377

FIGURE 103

CGGCTCGAGCGGCTCGAGTGAAGAGCCTCTCCACGGCTCCTGCGCCTGAGACAGCTGGCCTGACC
TCCAAATCATCCATCCACCCCTGCTGTCATCTGTTTTCATAGTGTGAGATCAACCCACAGGAATA
TCCATGGCTTTTGTGCTCATTTTGGTTCTCAGTTTCTACGAGCTGGTGTGAGGACAGTGGCAAGT
CACTGGACCGGGCAAGTTTGTCCAGGCCTTGGTGGGGGAGGACGCCGTGTTCTCCTGCTCCCTCT
TTCTGAGACCAGTGCAGAGGCTATGGAAGTGCGGTTCTTCAGGAATCAGTTCCATGCTGTGGTC
CACCTCTACAGAGATGGGGAAGACTGGGAATCTAAGCAGATGCCACAGTATCGAGGGAGAAGTGA
GTTTGTGAAGGACTCCATTGCAGGGGGGCGTGTCTCTAAGGCTAAAAACATCACTCCCTCGG
ACATCGGCCTGTATGGGTGCTGGTTCAGTTCCAGATTACGATGAGGAGGCCACCTGGGAGCTG
CGGGTGGCAGCACTGGGCTCACTTCCTCTCATTTCCATCGTGGGATATGTTGACGGAGGTATCCA
GTTACTCTGCCTGTCTCAGGCTGGTTCCCCCAGCCACAGCCAAGTGGAAAGGTCCACAAGGAC
AGGATTTGTCTTCAGACTCCAGAGCAAATGCAGATGGGTACAGCCTGTATGATGTGGAGATCTCC
ATTATAGTCCAGGAAAATGCTGGGAGCATATTGTGTTCCATCCACCTTGCTGAGCAGAGTCATGA
GGTGGAATCCAAGGTATTGATAGGAGAGACGTTTTTCCAGCCCTCACCTTGGCGCCTGGCTTCTA
TTTTACTCGGGTTACTCTGTGGTGGCCTGTGTGGTGTGTCATGGGGATGATAATTGTTTTCTTC
AAATCCAAAGGGAAAATCCAGGCGGAAGTGGACTGGAGAAGAAAGCACGGACAGGCAGAATTGAG
AGACGCCCGGAAACACGCAGTGGAGGTGACTCTGGATCCAGAGACGGCTCACCCGAAGCTCTGCG
TTTCTGATCTGAAAATGTAACCCATAGAAAAGCTCCCCAGGAGGTGCCTCACTCTGAGAAGAGA
TTTACAAGGAAGAGTGTGGTGGCTTCTCAGGGTTTCCAAGCAGGGAGACATTACTGGGAGGTGGA
CGTGGGACAAAATGTAGGGTGGTATGTGGGAGTGTGTGCGGATGACGTAGACAGGGGGAAGAACA
ATGTGACTTTGTCTCCCAACAATGGGTATTGGGTCCCTCAGACTGACAACAGAACATTGTATTTC
ACATTCAATCCCCATTTTATCAGCCTCCCCCCCCAGCACCCCTCCTACACGAGTAGGGGTCTTCCT
GGACTATGAGGGTGGGACCATCTCCTTCTTCAATACAAATGACCAGTCCCTTATTTATACCTGC
TGACATGTCAGTTTGAAGGCTTGTGAGACCTATATCCAGCATGCGATGTATGACGAGGAAAAG
GGGACTCCCATATTATATGTCCAGTGTCTGGGGATGAGACAGAGAAGACCCGTCTTAAAGGGC
CCCACACCACAGACCCAGACACAGCCAAGGGAGAGTGTCTCCGACAGGTGGCCCCAGCTTCCTCT
CCGGAGCCTGCGCACAGAGAGTCACGCCCCCACTCTCCTTTAGGGAGCTGAGGTTCTTCTGCCC
TGAGCCCTGCAGCAGCGGCAGTCACAGCTTCCAGATGAGGGGGGATTGGCCTGACCCTGTGGGAG
TCAGAAGCCATGGCTGCCCTGAAGTGGGGACGGAATAGACTCACATTAGGTTTAGTTTGTGAAAA
CTCCATCCAGCTAAGCGATCTTGAACAAGTCACAACCTCCCAGGCTCCTCATTTGCTAGTCACGG
ACAGTGATTCTCTGCCTCACAGGTGAAGATTAAAGAGACAACGAATGTGAATCATGCTTGCAGGTT
TGAGGGCACAGTGTGCTAATGATGTGTTTTTATATTATACATTTTCCACCATAAACTCTGTT
TGCTTATTCCACATTAATTTACTTTTTCTCTATACCAAAATCACCCATGGAATAGTTATTGAACACC
TGCTTTGTGAGGCTCAAAGAATAAAGAGGAGGTAGGATTTTTCACTGATTCTATAAGCCCAGCAT
TACCTGATACCAAAACCAGGCAAAGAAAACAGAAGAAGAGGAAGGAAAACCTACAGGTCCATATCC
CTCATTAACACAGACACAAAAATTCTAAATAAAATTTTAACAAATTAACTAAACAATATATTTA
AAGATGATATATAACTACTCAGTGTGGTTTGTCCACAAATGCAGAGTTGGTTTAATATTTAAAT
ATCAACCAGTGTAATTCAGCACATTAATAAAGTAAAAAAGAAAACCATAAAAAAAAAAAAAA

FIGURE 104

MAFVLILVLSFYELVSGQWQVTGPGKFVQALVGEDAVFSCSLFPETSAEAMEVRFFRNQFHAVVH
LYRDGEDWESKQMPQYRGRTEFVKDSIAGGRVSLRLKNITPSDIGLYGCWFSSQIYDEEATWELR
VAALGSLPLISIVGYVDGGIQLLCLSSGWFPQPTAKWKGPQGQDLSSDSRANADGYSLYDVEISI
IVQENAGSILCSIHLAEQSHEVESKVLIGETFFQPSPWRLASILLGLLCGALCGVVMGMIIVFFK
SKGKIQAEILDWRRKHGQAEIRDARKHAVEVTLDPETAHPKLCVSDLKTVTHRKAQEVPHSEKRF
TRKSVVASQGFQAGRHYWEVDVGQNVGWYVGVCRRDDVDRGKNNVTLSPNNGYWVLRLTTEHLYFT
FNPHFISLPPSTPPTRVGVFLDYEGGTISFFNTNDQSLIYTLLTCQFEGLLRPYIQHAMYDEEKG
TPIFICPVSWG

Signal peptide:

amino acids 1-17

Transmembrane domains:

amino acids 131-150, 235-259

FIGURE 105

CCTTCACAGGACTCTTCATTGCTGGTTGGCAATGATGTATCGGCCAGATGTGGTGAGGGCTAGGAAAAGAG
 TTTGTTGGGAACCCCTGGGTTATCGGCCTCGTCATCTTCATATCCCTGATTGTCTGGCAGTGTGCATTGGA
 CTCACTGTTTCATTATGTGAGATATAATCAAAAGAAGACCTACAATTACTATAGCACATTGTCAATTTACAAC
 TGACAAACTATATGCTGAGTTTGGCAGAGAGGCTTCTAACAATTTTACAGAAATGAGCCAGAGACTTGAAT
 CAATGGTGAAAAATGCATTTTATAAATCTCCATTAAGGGAAGAATTTGTCAAGTCTCAGGTTATCAAGTTC
 AGTCAACAGAAGCATGGAGTGTGGCTCATATGCTGTTGATTTGTAGATTTCACTCTACTGAGGATCCTGA
 AACTGTAGATAAAATTGTTCAACTTGTTTTACATGAAAAGCTGCAAGATGCTGTAGGACCCCTAAAGTAG
 ATCCTCACTCAGTTAAAATTAAAAAATCAACAAGACAGAAACAGACAGCTATCTAAACCATTGCTGCGGA
 ACACGAAGAAGTAAACTCTAGGTCAGAGTCTCAGGATCGTTGGTGGGACAGAAGTAGAAGAGGGTGAATG
 GCCCTGGCAGGCTAGCCTGCAGTGGGATGGGAGTCATCGCTGTGGAGCAACCTTAATTAATGCCACATGGC
 TTGTGAGTGCTGCTCACTGTTTTACAACATATAAGAACCCTGCCAGATGGACTGCTTCCTTTGGAGTAACA
 ATAAACCTTCGAAAATGAAACGGGGTCTCCGGAGAATAATTGTCCATGAAAATACAAACACCCATCACA
 TGACTATGATATTTCTCTTGAGAGCTTTCTAGCCCTGTTCCTACACAAATGCAGTACATAGAGTTTGTCT
 TCCCTGATGCATCCTATGAGTTTCAACCAGGTGATGTGATGTTTGTGACAGGATTTGGAGCACTGAAAAAT
 GATGGTTACAGTCAAAATCATCTTCGACAAGCACAGGTGACTCTCATAGACGCTACAACCTGCAATGAACC
 TCAAGCTTACAATGACGCCATAACTCCTAGAATGTTATGTGCTGGCTCCTTAGAAGGAAAAACAGATGCAT
 GCCAGGGTGAAGTCTGGAGGACCACTGGTTAGTTCAGATGCTAGAGATATCTGGTACCTTGGTGAATAGTG
 AGCTGGGGAGATGAATGTGCGAAACCCAACAAGCCTGGTGTCTTATACTAGAGTTACGGCCTTGCGGGACTG
 GATTACTTCAAAACTGGTATCTAAGAGACAAAAGCCTCATGGAACAGATAACATTTTTTTTTTGTTTTTTG
 GGTGTGGAGGCCATTTTTAGAGATACAGAATTGGAGAAGACTTGCAAAACAGCTAGATTTGACTGATCTCA
 ATAAACTGTTTGCTTGATGCATGTATTTCTTCCCAGCTCTGTTCCGCACGTAAGCATCCTGCTTCTGCCA
 GATCAACTCTGTCTATCTGTGAGCAATAGTTGAACTTTATGTACATAGAGAAATAGATAATACAATATTAC
 ATTACAGCCTGTATTCAATTTGTTCTCTAGAAGTTTGTGAGAATTTTGACTTGTGACATAAATTTGTAAT
 GCATATATACAATTTGAAGCACTCCTTTTCTTCAGTTCTCTCAGCTCCTCTCATTTTCAGCAAATATCCATTT
 TCAAGGTGCAGAACAAAGGAGTGAAAGAAAATATAAGAAGAAAAAATCCCCTACATTTTATTGGCACAGAA
 AAGTATTAGGTGTTTTTCTTAGTGGAATATTAGAAATGATCATATTCATTATGAAAGGTCAAGCAAAGACA
 GCAGAATACCAATCACTTCATCATTTAGGAAGTATGGGAAGTAAGTTAAGGAAGTCCAGAAAGAAGCCAAG
 ATATATCCTTATTTTCATTTCCAAACAACACTACTATGATAAATGTGAAGAAGATTCTGTTTTTTTGTGACCT
 ATAATAATTATACAACTTCATGCAATGTACTTGTCTAAGCAAATTAAAGCAAATATTTATTTAACATTG
 TTACTGAGGATGTCAACATATAACAATAAAATATAAATCACCCA

FIGURE 106

MMYRPDVVRARKRVCWEPWVIGLVIFISLIVLAVCIGLTVHYVRYNQKKTYNYSTLSFTTDKLY
AEFGREASNNFTEMSQRLESMVKNAFYKSPLREEFVKSQVIKFSQQKHGVLAHMLLICRFHSTED
PETVDKIVQLVLHEKLQDAVGPPKVDPHSVKIKKINKTETDSYLNHCCGTRRSKTLGQSLRIVGG
TEVEEGEWPWQASLQWDGSHRCGATLINATWLVSAAHCFTTYKNPARWTASFGVTIKPSKMKRGL
RRIIVHEKYKHPSHDYDISLAELSSPVYPYTNVHRVCLPDASYEFQPGDVMFVTGFGALKNDGYS
QNHLRQAQVTLIDATTCNEPQAYNDAITPRMLCAGSLEGKTDACQGDSGGPLVSSDARDIWYLAG
IVSWGDECAKPNKPGVYTRVTALRDWITSKTGI

Transmembrane domain:

amino acids 21-40 (type II)

FIGURE 107

AGAGAAAGAAGCGTCTCCAGCTGAAGCCAATGCAGCCCTCCGGCTCTCCGCGAAGAAGTTCCTG
 CCCCAGATGAGCCCCCGCGTGCCTCCCGACTATCCCCAGGCGGGCGTGGGGCACCGGGCCCAGC
 GCCGACGATCGCTGCCGTTTTGCCCCTGGGAGTAGGATGTGGTGAAAGGATGGGGCTTCTCCCTT
 ACGGGGCTCACAAATGCCAGAGAAGATTCCGTGAAGTGTCTGCGCTGCCTGCTCTACGCCCTCAA
 TCTGCTCTTTTGGTTAATGTCCATCAGTGTGTGGCAGTTTCTGCTTGGATGAGGGACTACCTAA
 ATAATGTTCTCACTTTAACTGCAGAAACGAGGGTAGAGGAAGCAGTCATTTGACTTACTTTCTT
 GTGGTTCATCCGGTCATGATTGCTGTTTGTCTGTTTCTTATCATTGTGGGGATGTTAGGATATTG
 TGGAACGGTGAAAAGAAATCTGTTGCTTCTTGCATGGTACTTTGGAAGTTTGTCTGTCATTTTCT
 GTGTAGAACTGGCTTGTGGCGTTTGGACATATGAACAGGAACCTTATGGTTCAGTACAATGGTCA
 GATATGGTCACTTTGAAAGCCAGGATGACAAATTATGGATTACCTAGATATCGGTGGCTTACTCA
 TGCTTGAATTTTTTTCAGAGAGAGTTTAAAGTGTGTGGAGTAGTATATTTCACTGACTGGTTGG
 AAATGACAGAGATGGACTGGCCCCCAGATTCTGCTGTGTGTAGAGAATTTCCAGGATGTTCCAAA
 CAGGCCCCACAGGAAGATCTCAGTGACCTTTATCAAGAGGGTGTGGGAAGAAAATGTATTCCTT
 TTTGAGAGGAACCAACAACCTGCAGGTGCTGAGGTTTCTGGGAATCTCCATTGGGGTGACACAAA
 TCCTGGCCATGATTCTCACCATTACTCTGCTCTGGGCTCTGTATTATGATAGAAGGGAGCCTGGG
 ACAGACCAAATGATGTCCTTGAAGAATGACAACTCTCAGCACCTGTCATGTCCCTCAGTAGAACT
 GTTGAAACCAAGCCTGTCAAGAATCTTTGAACACACATCCATGGCAAACAGCTTTAATACACACT
 TTGAGATGGAGGAGTTATAAAAAAGAAATGTCACAGAAGAAAACCACAACTTGTTTTATTGGACT
 TGTGAATTTTGTAGTACATACTATGTGTTTCAGAAATATGTAGAAAATAAAATGTTGCCATAAAA
 TAACACCTAAGCATATACTATTCTATGCTTTAAATGAGGATGGAAAAGTTTCATGTCATAAGTC
 ACCACCTGGACAATAATTGATGCCCTTAAATGCTGAAGACAGATGTCATACCCACTGTGTAGCC
 TGTGTATGACTTTTACTGAACACAGTTATGTTTTGAGGCAGCATGGTTTGATTAGCATTTCCGCA
 TCCATGCAAACGAGTCACATATGGTGGGACTGGAGCCATAGTAAAGGTTGATTTACTTCTACCAA
 CTAGTATATAAAGTACTAATTAAATGCTAACATAGGAAGTTAGAAAATACTAATAACTTTTATTA
 CTCAGCGATCTATTCTTCTGATGCTAAATAAATTATATATCAGAAAACCTTTCAATATTGGTGACT
 ACCTAAATGTGATTTTTGCTGGTTACTAAAATATTCTTACCCTTAAAAGAGCAAGCTAACACAT
 TGTCTTAAGCTGATCAGGGATTTTTTGTATATAAGTCTGTGTTAAATCTGTATAATTCACTCGAT
 TTCAGTTCTGATAATGTTAAGAATAACCATATGAAAAGGAAAATTTGTCTGTATAGCATCATT
 ATTTTGTAGCCTTTCTGTAAATAAAGCTTTACTATTCTGTCTGGGCTTATATTACACATATAAC
 TGTTATTTAAATACTTAACCACTAATTTGAAAATTACCAGTGTGATACATAGGAATCATTATTC
 AGAATGTAGTCTGGTCTTTAGGAAGTATTAATAAGAAAATTTGCACATAACTTAGTTGATTGAGA
 AAGGACTTGTATGCTGTTTTTCTCCCAAATGAAGACTCTTTTGGACACTAAACACTTTTTAAAAA
 GCTTATCTTTGCCCTTCTCCAAACAAGAAGCAATAGTCTCCAAGTCAATATAAATTTACAGAAAA
 TAGTGTCTTTTTTCTCCAGAAAAATGCTGTGAGAATCATTAAACATGTGACAATTTAGAGATT
 CTTTGTTTTATTTCACTGATTAAATATACTGTGGCAAATTACACAGATTATTAAATTTTTTTACAA
 GAGTATAGTATATTTATTTGAAATGGGAAAAGTGCATTTTACTGTATTTTGTGTATTTTGTATTAT
 TTCTCAGAAATATGGAAGAAAATTAATATGTGTCAATAAATATTTCTAGAGAGTAA

FIGURE 108

MAREDSVKCLRCLLYALNLLFWLMSISVLAVSAWMRDYLNNVLTTLTAETRVEEAVILTYFPVVHP
VMIAVCCFLIIVGMLGYCGTVKRNLLLLAWYFGSLLVIFCVELACGVWTYEQELMVPVQWSDMVT
LKARMTNYGLPRYRWLTHAWNFFQREFKCCGVVYFTDWLEMTMDWPPDSCCVREFPGCSKQAHQ
EDLSDLYQEGCGKKMYSFLRGTKQLQVLRFLGISIGVTQILAMILTITLLWALYYDRREPSTDQM
MSLKNDNSQHLSCPSVELLKPSLSRIFEHTSMANSFNTHFEMEEL

Signal peptide:

amino acids 1-33

Transmembrane domains:

amino acids 12-35, 57-86, 94-114, 226-248

FIGURE 109

CCAAGGCCAGAGCTGTGGACACCTTATCCCACTCATCCTCATCCTCTTCCTCTGATAAAGCCCTACCAGTGCT
GATAAAGTCTTTCTCGTGAGAGCCTAGAGGCCCTAAAAAAGTGCTTGAAAGAGAAGGGGACAAAGGAACA
CCAGTATTAAGAGGATTTTCCAGTGTTTCTGGCAGTTGGTCCAGAAGCATGCTCCATTCTGCTTCTCACCTG
CCTCTTCATCACAGGCACCTCCGTGTACCCGTGGCCCTAGATCCTTGTTCTGCTTACATCAGCCTGAATGAGC
CCTGGAGGAACACTGACCACCAGTTGGATGAGTCTCAAGGTCCTCCTCTATGTGACAACCATGTGAATGGGGAG
TGGTACCACCTTCACGGGCATGGCGGGAGATGCCATGCCCTACCTTCTGCATACCAGAAAACCACTGTGGAACCCA
CGCACCTGTCTGGCTCAATGGCAGCCACCCCTAGAAGGCGACGGCATTGTGCAACGCCAGGCTTGTGCCAGCT
TCAATGGGAACCTGCTGTCTCTGGAACACCACGGTGGAAAGTCAAGGCTTGCCCTGGAGGCTACTATGTGTATCGT
CTGACCAAGCCCAGCGTCTGCTTCCACGTCTACTGTGGTCATTTTATGACATCTGCGACGAGGACTGCCATGG
CAGCTGCTCAGATACCAGCGAGTGCACATGCCCTCCAGGAAGTGTGCTAGGCCCTGACAGGCAGACATGCTTTG
ATGAAATGAATGTGAGCAAAACAACGGTGGCTGCAGTGAGATCTGTGTGAACCTCAAAAACCTCCTACCGCTGT
GAGTGTGGGGTTGGCCGTGTGCTAAGAAGTGTGCAAGACTTGTGAAGACGTTGAAGGATGCCACAATAACAA
TGGTGGCTGCAGCCACTCTTGCCCTGGATCTGAGAAAGGCTACCAGTGTGAATGTCCCGGGGCTGGTGCTGT
CTGAGGATAACCACTTGGCAAGTCCCTGTGTGTGCAAATCAAATGCCATTGAAGTGAACATCCCCAGGGAG
CTGGTTGGTGGCCTGGAGCTCTTCTGACCAACCTCCTGCCGAGGAGTGTCCAACGGCACCCATGTCAACAT
CCTCTTCTCTCAAGACATGTGGTACAGTGGTCCATGTGGTGAATGACAAGATTGTGGCCAGCAACCTCGTGA
CAGGTCTACCCAAGCAGACCCCGGGGAGCAGCGGGGACTTCATCATCCGAACCAGCAAGCTGCTGATCCCGGTG
ACCTGCGAGTTTCCACGCCTGTACACCATTCTGAAGGATACGTTCCCAACCTTCGAAACTCCCCACTGGAAAT
CATGAGCCGAAATCATGGGATCTTCCCATTCCTCTGGAGATCTTCAAGGACAATGAGTTTGAAGAGCCTTACC
GGGAAGCTCTGCCACCCCTCAAGCTTCGTACTCCCTCTACTTTGGCATTGAGCCCGTGGTGCACGTGAGCGGC
TTGGAAGCTTGGTGGAGAGCTGCTTTGCCACCCCCACCTCCAAGATCGACGAGGTCCTGAAATACTACCTCAT
CCGGGATGGCTGTGTTTTCAGATGACTCGGTAAAGCAGTACACATCCCGGGATCACCTAGCAAAGCACTTCCAGG
TCCCTGTCTTCAAGTTTGTGGGCAAAGACCACAAGGAAGTGTCTGCACTGCCGGGTCTTGTCTGTGGAGTG
TTGGACGAGCGTTCCCGCTGTGCCAGGGTTGCCACCGGCGAATGCGTCGTGGGGCAGGAGGAGAGGACTCAGC
CGGTCTACAGGGCCAGACGCTAACAGGCGGCCGATCCGCATCGACTGGGAGGACTAGTTTCGTAGCCATACCTC
GAGTCCCTGCATTGGACGGCTCTGCTCTTTGGAGCTTCTCCCCCACCGCCCTCTAAGAACATCTGCCAACAGC
TGGGTTTCAGACTTCACACTGTGAGTTTCAGACTCCAGCACCAACTCACTCTGATTCTGGTCCATTCACTGGGCA
CAGGTCACAGCACTGCTGAACAATGTGGCCTGGGTGGGGTTTCATCTTTCTAGGGTTGAAAATAAAGTGTCCA
CCCAGAAAGACACTCACCCCATTTCCCTCATTTCTTCTTACACTTAAATACCTCGTGTATGGTGAATCAGAC
CACAAATCAGAAGCTGGGTATAATATTTCAAGTTACAAACCCTAGAAAAATTAAACAGTTACTGAAATTATGA
CTTAAATACCAATGACTCCTTAAATATGTAAATTATAGTTATACCTTGAAATTTCAATTCAAATGCAGACTAA
TTATAGGGAATTTGGAAGTGTATCAATAAAACAGTATATAATTTT

FIGURE 110

MPPFLLLTCLFITGTSVSPVALDPCSAYISLNEPWRNTDHQLDESQGPPLCDNHVNGEWYHFTGMAGDAMP
TFCIPENHCGTHAPVWLNGSHPLEGDGIVQRQACASFNGNCCLWNTTVEVKACPGGYVYRLTKPSVCFHV
YCGHFYDIDCEDCHGSCSDTSECTCAPGTVLGPDRQTCFDENECEQNNGGCSEICVNLKNSYRCECGVGRV
LRSDGKTCEDVEGCHNNNGGCSHSCLGSEKGYQCECPRGLVLSEDNHTCQVPVLCKSNAIEVNI PRELVGG
LELFLTNTSCRGVSNNGTHVNILFSLKTCGTVVDVNDKIVASNLVTGLPKQTPGSSGDFIIRTSKLLIPVT
CEFPRLYTISEGYVPNLRNSPLEIMSRNHGIFPFTLEIFKDNEFEOPYREALPTLKLRLDSLYFGIEPVVHV
SGLESIVESCFAPTSKIDEVLKYILIRDGCVSDSVKQYTSRDHLAKHFQVPVFKFVGKDHKEVELHCRV
LVCGVLDERSRCAQGCCHRRMRRGAGGEDSAGLQGQTLTGGPIDWED

Important features of the protein:

Signal peptide:

amino acids 1-16

N-glycosylation sites.

amino acids 89-93, 116-120, 259-263, 291-295, 299-303

Tyrosine kinase phosphorylation sites.

amino acids 411-418, 443-451

N-myristoylation sites.

amino acids 226-232, 233-239, 240-246, 252-258, 296-302, 300-306,
522-528, 531-537

Aspartic acid and asparagine hydroxylation site.

amino acids 197-209

ZP domain proteins.

amino acids 431-457

Calcium-binding EGF-like proteins.

amino acids 191-212, 232-253

FIGURE 111

GAGAGAGGCAGCAGCTTGCTCAGCGGACAAGGATGCTGGGCGTGAGGGACCAAGGCCTGCCCTGCACTCGG
 GCCTCCTCCAGCCAGTGCTGACCAGGGACTTCTGACCTGCTGGCCAGCCAGGACCTGTGTGGGGAGGCCCT
 CCTGCTGCCCTTGGGGTGACAATCTCAGCTCCAGGCTACAGGGAGACCGGGAGGATCACAGAGCCAGCATGT
 TACAGGATCCTGACAGTGATCAACCTCTGAACAGCCTCGATGTCAAACCCCTGCGCAAACCCCGTATCCCC
 ATGGAGACCTTCAGAAAGGTGGGGATCCCCATCATCATAGCACTACTGAGCCTGGCGAGTATCATCATTTGT
 GGTGTCTCATCAAGGTGATTCTGGATAAATACTACTTCTCTGCGGGCAGCCTCTCCACTTCATCCCGA
 GGAAGCAGCTGTGTGACGGAGAGCTGGACTGTCCCTTGGGGGAGGACGAGGAGCACTGTGTCAAGAGCTTC
 CCCGAAGGGCCTGCAGTGGCAGTCCGCCCTCTCCAAGGACCGATCCACACTGCAGGTGCTGGACTCGGCCAC
 AGGGAAGTGGTTCTCTGCCTGTTTCGAGAACTTCACAGAAGCTCTCGCTGAGACAGCCTGTAGGCAGATGG
 GCTACAGCAGAGCTGTGGAGATTGGCCCAGACCAGGATCTGGATGTTGTTGAAATCACAGAAAACAGCCAG
 GAGCTTCGCATGCGGAACTCAAGTGGGCCCTGTCTCTCAGGCTCCCTGGTCTCCCTGCACTGTCTTGCCCTG
 TGGGAAGAGCCTGAAGACCCCCCGTGTGGTGGGTGGGAGGAGCCCTCTGTGGATTCTTGGCCTTGGCAGG
 TCAGCATCCAGTACGACAAACAGCACGTCTGTGGAGGGAGCATCCTGGACCCCCACTGGGTCTCACGGCA
 GCCCCTGCTTCAGGAAACATACCGATGTGTTCAACTGGAAGGTGCGGGCAGGCTCAGACAACTGGGCAG
 CTTCCCATCCCTGGCTGTGGCCAAGATCATCATATTGAATTCAACCCCATGTACCCCAAAGACAATGACA
 TCGCCCTCATGAAGCTGCAGTTCCTCACTTTCTCAGGCACAGTCAGGCCCATCTGTCTGCCCTTCTTT
 GATGAGGAGCTCACTCCAGCCACCCCACTCTGGATCATTGGATGGGGCTTTACGAAGCAGAATGGAGGAA
 GATGTCTGACATACTGCTGCAGGCGTCAGTCCAGGTCATTGACAGCACACGGTGCAATGCAGACGATGCGT
 ACCAGGGGGAAGTCACCGAGAAGATGATGTGTGCAGGCATCCCGGAAGGGGGTGTGGACACCTGCCAGGGT
 GACAGTGGTGGGCCCTGATGTACCAATCTGACCAGTGGCATGTGGTGGGCATCGTTAGCTGGGGCTATGG
 CTGCGGGGGCCCGAGCACCCAGGAGTATACACCAAGGTCTCAGCCTATCTCAACTGGATCTACAATGTCT
 GGAAGGCTGAGCTGTAATGCTGCTGCCCCTTGCAGTGTGGGAGCCGCTTCCTTCCTGCCCTGCCACCT
 GGGGATCCCCCAAAGTCAGACACAGAGCAAGAGTCCCCTTGGGTACACCCCTCTGCCACAGCCTCAGCAT
 TTCTTGAGAGCAGCAAAGGGCCTCAATTCCTGTAAGAGACCCTCGCAGCCCAGAGGCGCCAGAGGAAGTCA
 GCAGCCCTAGCTCGGCCACACTTGGTGCTCCCAGCATCCCAGGGAGAGACACAGCCCACTGAACAAGGTCT
 CAGGGGTATTGCTAAGCCAAGAAGGAACCTTCCACACTACTGAATGGAAGCAGGCTGTCTTGTAAGCC
 CAGATCACTGTGGCTGGAGAGGAGAAGGAAAGGGTCTGCGCCAGCCCTGTCCGTCTTACCCATCCCCAA
 GCCTACTAGAGCAAGAAACCAGTTGTAATATAAAATGCACTGCCCTACTGTTGGTATGACTACCGTTACCT
 ACTGTTGTCAATTGTTATTACAGCTATGGCCACTATTATTAAAGAGCTGTGTAACATCTCTGGCAAAAAAA
 AAAA

FIGURE 112

MLQDPDSQPLNSLDVKPLRKPRIPMETFRKVGIPII IALLSLASII IIVVVLIKVILDKYYFLCG
QPLHFIPRKQLCDGELDCPLGEDEEHCVKSFPEGPAVAVRLSKDRSTLQVLDSATGNWFSACFDN
FTEALAETACRQMGYSRAVEIGPDQDL DVVEITENSQELMRNSSGPCLSGSLVSLHCLACGKSL
KTPRVVGEEASVDSWPWQVSIQYDKQHVC GGSILDPHWVLTAAHCFRKHTDVFNWKVRAGSDKL
GSFPSLAVAKIIIIIEFNPMYPKNDIALMKLQFPLTFSGTVRPICLPFFDEELTPATPLWIIGWG
FTKQNGGKMSDILLQASVQVIDSTRCNADDAYQGEVTEKMMCAGIPEGGVDTCCQDSSGGPLMYQS
DQWHVVGIVSWGYGCGGPSTPGVYTKVSAYLNWIYNVWKAEL

Transmembrane domain:

amino acids 32-53 (typeII)

FIGURE 113

GGCTGGACTGGAACCTCCTGGTCCCAAGTGATCCACCCGCCTCAGCCTCCCAAGGTGCTGTGATTA
TAGGTGTAAGCCACCGTGTCTGGCCTCTGAACAACTTTTTTCAGCAACTAAAAAGCCACAGGAGT
TGAAGTGTAGGATTCTGACTATGCTGTGGTGGCTAGTGCTCCTACTCCTACCTACATTAAAATC
TGTTTTTTGTTCTCTTGTAAGTACCTTTACCTTCCTAACACAGAGGATCTGTCACTGTGGCTCT
GGCCCAAACCTGACCTTCACTCTGGAACGAGAACAGAGGTTTCTACCCACACCGTCCCCTCGAAG
CCGGGGACAGCCTCACCTTGCTGGCCTCTCGCTGGAGCAGTGCCCTCACCAACTGTCTCACGTCT
GGAGGCACTGACTCGGGCAGTGACAGGTAGCTGAGCCTCTTGGTAGCTGCGGCTTTCAAGGTGGGC
CTTGCCCTGGCCGTAGAAGGGATTGACAAGCCCGAAGATTTTCATAGGCGATGGCTCCCCTGCCC
AGGCATCAGCCTTGCTGTAGTCAATCACTGCCCTGGGGCCAGGACGGGCGTGGAACACCTGCTCA
GAAGCAGTGGGTGAGACATCACGTGCCCCGCCATCTAACCTTTTCATGTCTGCACATCACCTG
ATCCATGGGCTAATCTGAACTCTGTCCCAAGGAACCCAGAGCTTGAGTGAGCTGTGGCTCAGACC
CAGAAGGGGTCTGCTTAGACCACCTGGTTTATGTGACAGGACTTGCATTCTCCTGGAACATGAGG
GAACGCCGGAGGAAAGCAAAGTGGCAGGGAAGGAACCTGTGCCAAATTATGGGTGAGAAAAGATG
GAGGTGTTGGGTATCACAAAGGCATCGAGTCTCCTGCATTCACTGGACATGTGGGGGAAGGGCTG
CCGATGGCGCATGACACACTCGGGACTCACCTCTGGGGCCATCAGACAGCCGTTTCCGCCCCGAT
CCACGTACCAGCTGCTGAAGGGCAACTGCAGGCCGATGCTCTCATCAGCCAGGCAGCAGCCAAAA
TCTGCGATCACCAGCCAGGGGCAGCCGTCTGGGAAGGAGCAAGCAAAGTGACCATTTCTCCTCCC
CTCCTTCCCTCTGAGAGGCCCTCCTATGTCCCTACTAAAGCCACCAGCAAGACATAGCTGACAGG
GGCTAATGGCTCAGTGTGGCCAGGAGGTCAGCAAGGCCTGAGAGCTGATCAGAAGGGCCTGCT
GTGCGAACACGGAAATGCCTCCAGTAAGCACAGGCTGCAAAATCCCCAGGCAAAGGACTGTGTGG
CTCAATTTAAATCATGTTCTAGTAATTGGAGCTGTCCCCAAGACCAAAGGAGCTAGAGCTTGTT
CAAATGATCTCCAAGGGCCCTTATACCCCAGGAGACTTTGATTTGAATTTGAAACCCCAAATCCA
AACCTAAGAACCAGGTGCATTAAGAATCAGTTATTGCCGGGTGTGGTGGCCTGTAATGCCAACAT
TTTGGGAGGCCGAGGCGGGTAGATCACCTGAGGTCAGGAGTTCAAGACCAGCCTGGCCAACATGG
TGAAACCCCTGTCTCTACTAAAAATACAAAAAACTAGCCAGGCATGGTGGTGTGTGCCTGTATC
CCAGCTACTCGGGAGGCTGAGACAGGAGAATTACTTGAACCTGGGAGGTGAAGGAGGCTGAGACA
GGAGAATCACTTCAGCCTGAGCAACACAGCGAGACTCTGTCTCAGAAAAAATAAAAAAGAATTA
TGTTTATTTGTAA

FIGURE 114

MLWWLVLLLLPTLKSVFCSLVTSLYLPNTEDLSLWLWPKPDLHSGTRTEVSTHTVPSKPGTASPC
WPLAGAVPSPTVSRLEALTRAVQVAEPLGSCGFQGGPCPGRRRD

Signal peptide:

amino acids 1-15

FIGURE 115

CAGCAGTGGTCTCTCAGTCCTCTCAAAGCAAGGAAAGAGTACTGTGTGCTGAGAGACCATGGCAA
AGAATCCTCCAGAGAATTGTGAAGACTGTCACATTCTAAATGCAGAAGCTTTTAAATCCAAGAAA
ATATGTAAATCACTTAAGATTTGTGGACTGGTGTGTTGGTATCCTGGCCCTAACTCTAATTGTCCT
GTTTTGGGGGAGCAAGCACTTCTGGCCGGAGGTACCCAAAAAGCCTATGACATGGAGCACACTT
TCTACAGCAATGGAGAGAAGAAGAAGATTTACATGGAAATTGATCCTGTGACCAGAACTGAAATA
TTCAGAAGCGGAAATGGCACTGATGAAACATTGGAAGTGCACGACTTTAAAAACGGATACACTGG
CATCTACTTCGTGGGTCTTCAAAAATGTTTTATCAAACTCAGATTAAAGTGATTCCTGAATTTT
CTGAACCAGAAGAGGAAATAGATGAGAATGAAGAAATTACCACAACCTTTCTTTGAACAGTCAGTG
ATTTGGGTCCCAGCAGAAAAGCCTATTGAAAACCGAGATTTTCTTAAAAATTCCAAAATTCTGGA
GATTTGTGATAACGTGACCATGTATTGGATCAATCCCACTCTAATATCAGTTTCTGAGTTACAAG
ACTTTGAGGAGGAGGGAGAAGATCTTCACTTTCTGCCAACGAAAAAAAGGGATTGAACAAAAT
GAACAGTGGGTGGTCCCTCAAGTGAAAGTAGAGAAGACCCGTCACGCCAGACAAGCAAGTGAGGA
AGAACTTCCAATAAATGACTATACTGAAAATGGAATAGAATTTGATCCCATGCTGGATGAGAGAG
GTTATTGTTGTATTTACTGCCGTCGAGGCAACCGCTATTGCCGCCGCGTCTGTGAACCTTTACTA
GGCTACTACCCATATCCATACTGCTACCAAGGAGGACGAGTCATCTGTGTCATCATGCCTTG
TAACTGGTGGGTGGCCCGCATGCTGGGGAGGGTCTTAATAGGAGGTTTGAGCTCAAATGCTTAAAC
TGCTGGCAACATATAATAAATGCATGCTATTCAATGAATTTCTGCCTATGAGGCATCTGGCCCCCT
GGTAGCCAGCTCTCCAGAATTACTTGTAGGTAATTCCTCTCTTCATGTTCTAATAAACTTCTACA
TTATCACCAAAAAAAAAAAAAAAAAAAAA

FIGURE 116

MAKNPPENCEDCHILNAEAFKSKKICKSLKICGLVFGILALTILVLFWGSKHFWPEVPPKAYDME
HTFYSNGEKKKIYMEIDPVTRTEIFRSGNGTDETLVHDFKNGYTGIIYFVGLQKCFIKTQIKVIP
EFSEPEEEIDENEEITTTFFEQSVIWWPAEKPIENRDFLKNKILEICDNVTMYWINPTLISVSE
LQDFEEEGEDLHFPANEKKGIEQNEQWVVPQVKVEKTRHARQASEEELPINDYTENGIEFDPMLD
ERGYCCIIYCRGNRYCRRVCEPLLGYYPYPYCYQGGRVICRVIMPCNWWVARMLGRV

Important features of the protein:

Signal peptide:

amino acids 1-40

Transmembrane domain:

amino acids 25-47 (type II)

N-glycosylation sites.

amino acids 94-97, 180-183

Glycosaminoglycan attachment sites.

amino acids 92-95, 70-73, 85-88, 133-136, 148-151, 192-195, 239-242

N-myristoylation sites.

amino acids 33-38, 95-100, 116-121, 215-220, 272-277

Microbodies C-terminal targeting signal.

amino acids 315-317

Cytochrome c family heme-binding site signature.

amino acids 9-14

FIGURE 117

GAGCTCCCCTCAGGAGCGCGTTAGCTTCACACCTTCGGCAGCAGGAGGGCGGCAGCTTCTCGCAGGCGGCA
GGGCGGGCGGCCAGGATCATGTCCACCACCACATGCCAAGTGGTGGCGTTCTCTCTGTCCATCCTGGGGCT
GGCCGGCTGCATCGCGGCCACCGGGATGGACATGTGGAGCACCCAGGACCTGTACGACAACCCCGTCACCT
CCGTGTTCCAGTACGAAGGGCTCTGGAGGAGCTGCGTGAGGCAGAGTTTACGGCTTCACCGAATGCAGGCCC
TATTTACCATCCTGGGACTTCCAGCCATGCTGCAGGCAGTGGGAGCCCTGATGATCGTAGGCATCGTCTCT
GGGTGCCATTGGCCTCCTGGTATCCATCTTTGCCCTGAAATGCATCCGCATTGGCAGCATGGAGGACTCTG
CCAAAGCCAACATGACACTGACCTCCGGGATCATGTTTATTGTCTCAGGTCTTTGTGCAATTGCTGGAGTG
TCTGTGTTTGCCAACATGCTGGTGACTAACTTCTGGATGTCCACAGCTAACATGTACACCGGCATGGGTGG
GATGGTGCAGACTGTTTACAGCCAGGTACACATTTGGTGCGGCTCTGTTTCGTGGGCTGGGTCTGAGGCC
TCACACTAATTGGGGGTGTGATGATGTGCATCGCCTGCCGGGGCTGGCACCAGAAGAAACCAACTACAAA
GCCGTTTCTTATCATGCCTCAGGCCACAGTGTGCTTACAAGCCTGGAGGCTTCAAGGCCAGCACTGGCTT
TGGGTCCAACACCAAAAACAAGAAGATATACGATGGAGGTGCCCGCACAGAGGACGAGGTACAATCTTATC
CTTCCAAGCAGCACTATGTGTAATGCTCTAAGACCTCTCAGCACGGGCGGAAGAACTCCCGGAGAGCTCA
CCCAAAAAACAAGGAGATCCCATCTAGATTTCTTCTTGCTTTTGACTCACAGCTGGAAGTTAGAAAAGCCT
CGATTTTCATCTTTGGAGAGGCCAAATGGTCTTAGCCTCAGTCTCTGTCTCTAAATATTCCACCATAAAACA
GCTGAGTTATTTATGAATTAGAGGCTATAGCTCACATTTTCAATCCTCTATTTCTTTTTTAAATATAACT
TTCTACTCTGATGAGAGAATGTGGTTTTAATCTCTCTCTCACATTTTGATGATTTAGACAGACTCCCCCTC
TTCCTCCTAGTCAATAAACCCATTGATGATCTATTTCCAGCTTATCCCCAAGAAACTTTTGAAGGAAA
GAGTAGACCCAAAGATGTTATTTTCTGCTGTTTGAATTTTGTCTCCCCACCCCAACTTGGCTAGTAATAA
ACACTTACTGAAGAAGAAGCAATAAGAGAAAGATATTTGTAATCTCTCCAGCCCATGATCTCGGTTTTCTT
ACACTGTGATCTTAAAGTTACCAAACCAAAGTCATTTTTCAGTTTGAGGCAACCAAACCTTTCTACTGCTG
TTGACATCTTCTTATTACAGCAACACCATTCTAGGAGTTTCTTGAGCTCTCCACTGGAGTCTCTTCTGT
CGCGGGTCAGAAATTGTCCCTAGATGAATGAGAAAATTATTTTTTTAATTTAAGTCTAAATATAGTTAA
AATAAATAATGTTTTAGTAAATGATACACTATCTCTGTGAAATAGCCTCACCCCTACATGTGGATAGAAG
GAAATGAAAAAATAATTGCTTTGACATTGTCTATATGGTACTTTGTAAAGTCATGCTTAAGTACAAATTCC
ATGAAAAGCTCACACCTGTAATCCTAGCACTTTGGGAGGCTGAGGAGGAAGGATCACTTGAGCCCAGAAGT
TCGAGACTAGCCTGGGCAACATGGAGAAGCCCTGTCTCTACAAAATACAGAGAGAAAAAATCAGCCAGTCA
TGGTGGCATAACCTGTAGTCCCAGCATTCGGGAGGCTGAGGTGGGAGGATCACTTGAGCCCAGGGAGGT
TGGGGCTGCAGTGAGCCATGATCACACCACTGCACTCCAGCCAGGTGACATAGCGAGATCCTGTCTAAAAA
AATAAAAAATAAATAATGGAACACAGCAAGTCCTAGGAAGTAGGTTAAACTAATTCTTTAA

FIGURE 118

MSTTTCQVVAFLLSILGLAGCIAATGMDMWSTQDLYDNPVTSVFQYEGLRSCVRQSSGFTECRP
YFTILGLPAMLQAVRALMIVGIVLGAIGLLVSI FALKCIRIGSMEDSAKANMTLTSGIMFIVSGL
CAIAGVSVFANMLVTNFWMSTANMYTGMGGMVQTVQTRYTFGAALFVGWVAGGLTLIGGVMMCIA
CRGLAPEETNYKAVSYHASGHSVAYKPGGFKASTGFGSNTKNKKIYDGGARTEDEVQSYPSKHDY
V

Signal peptide:

amino acids 1-23

Transmembrane domains:

amino acids 81-100, 121-141, 173-194

FIGURE 119

GGAAAACTGTTCTCTTCTGTGGCACAGAGAACCCTGCTTCAAAGCAGAAGTAGCAGTTCCGGAGTCC
 AGCTGGCTAAAACTCATCCCAGAGGATAATGGCAACCCATGCCTTAGAAATCGCTGGGCTGTTTCTTG
 GTGGTGTGGAAATGGTGGGCACAGTGGCTGTCACTGTCACTGCCTCAGTGGAGAGTGTGGCCTTCATT
 GAAAACAACATCGTGGTTTTTGAAACTTCTGGGAAGGACTGTGGATGAATTGCGTGAGGCAGGCTAA
 CATCAGGATGCAGTGCAAAATCTATGATTCCTGCTGGCTCTTTCTCCGGACCTACAGGCAGCCAGAG
 GACTGATGTGTGCTGCTTCCGTGATGTCCTTCTTGGCTTTCATGATGGCCATCCTTGGCATGAAATGC
 ACCAGGTGCACGGGGGACAATGAGAAGGTGAAGGCTCACATTCTGCTGACGGCTGGAATCATCTTCAT
 CATCACGGGCATGGTGGTGCTCATCCCTGTGAGCTGGGTTGCCAATGCCATCATCAGAGATTTCTATA
 ACTCAATAGTGAATGTTGCCCAAAACGTGAGCTTGGAGAAGCTCTCTACTTAGGATGGACCACGGCA
 CTGGTGTGATTGTTGGAGGAGCTCTGTTCTGCTGCGTTTTTTGTTGCAACGAAAAGAGCAGTAGCTA
 CAGATACTCGATACCTTCCCATCGCACAAACCAAAAAAGTTATCACACCGGAAAGAAGTCACCGAGCG
 TCTACTCCAGAAGTCAGTATGTGTAGTTGTGTATGTTTTTTAACTTTACTATAAAGCCATGCAAATG
 ACAAAAATCTATATTACTTTCTCAAAATGGACCCCAAGAACTTTGATTTACTGTTCTTAACTGCCT
 AATCTTAATTACAGGAACGTGTCATCAGCTATTTATGATTCTATAAGCTATTTACGCAGAATGAGATA
 TTAAACCAATGCTTTGATTGTTCTAGAAAGTATAGTAATTTGTTTTCTAAGGTGGTTCAAGCATCTA
 CTCTTTTTATCATTTACTTCAAAATGACATTGCTAAAGACTGCATTATTTTACTACTGTAATTTCTCC
 ACGACATAGCATTATGTACATAGATGAGTGTAACATTTATATCTCACATAGAGACATGCTTATATGGT
 TTTATTTAAATGAAATGCCAGTCCATTACACTGAATAAATAGAACTCAACTATTGCTTTTCAGGGAA
 ATCATGGATAGGGTTGAAGAAGTTACTATTAATTGTTTAAAAACAGCTTAGGGATTAATGTCCTCCA
 TTTATAATGAAGATTAAATGAAGGCTTTAATCAGCATTGTAAAGGAAATTGAATGGCTTTCTGATAT
 GCTGTTTTTTAGCCTAGGAGTTAGAAATCCTAACTTCTTTATCCTCTTCTCCAGAGGCTTTTTTTTT
 CTTGTGTATTAAATTAACATTTTTAAACGCAGATATTTGTCAAGGGGCTTTGCATTCAAACCTGCTT
 TTCCAGGGCTATACTCAGAAGAAAGATAAAAGTGTGATCTAAGAAAAAGTGATGGTTTTAGGAAAGTG
 AAAATATTTTTGTTTTGTATTTGAAGAAGAATGATGCATTTTGACAAGAAATCATATATGTATGGAT
 ATATTTTAATAAGTATTTGAGTACAGACTTTGAGGTTTCATCAATATAAATAAAAGAGCAGAAAAATA
 TGTCTTGGTTTTTCATTTGCTTACCAAAAAACAACAACAAAAAAGTTGTCCTTTGAGAACTTCACCT
 GCTCCTATGTGGGTACCTGAGTCAAAATGTCAATTTTGTCTGTGAAAAATAAATTTCTTCTTGTA
 CCATTTCTGTTTAGTTTTACTAAAATCTGTAAATACTGTATTTTTCTGTTTATTCCAAATTTGATGAA
 ACTGACAATCCAATTTGAAAGTTTGTGTCGACGTCTGTCTAGCTTAAATGAATGTGTTCTATTTGCTT
 TATACATTTATATTAATAAATTTGTACATTTTCTAATT

FIGURE 120

MATHALEIAGLFLGGVGMVGTVAVTVMPOWRVSAFIENNIIVVFENFWEGLWMNCVRQANIRMQCK
IYDSELLALSPDLQAARGLMCAASVMSFLAFMMAILGMKCTRCTGDNEKVKAHILLTAGIIFIITG
MVLIPVSWVANAIIRDFYNSIVNVAQKRELGEALYLGWTTALVLIVGGALFCCVFCCNEKSSSY
RYSIPSHRTTQKSYHTGKKSPSVYSRSQYV

Signal peptide:

amino acids 1-17

Transmembrane domains:

amino acids 82-101, 118-145, 164-188

FIGURE 121

GGAGAGAGGCGCGCGGGTGAAAGGCGCATTGATGCAGCCTGCGGCGGCCTCGGAGCGCGGCGGAG
CCAGACGCTGACCACGTTCTCTCCTCGGTCTCCTCCGCCTCCAGCTCCGCGCTGCCCGGCAGCC
GGGAGCCATGCGACCCCAGGGCCCCGCGCCTCCCCGCAGCGGCTCCGCGGCCTCCTGCTGCTCC
TGCTGCTGCAGCTGCCCGCGCCGTCGAGCGCCTCTGAGATCCCCAAGGGGAAGCAAAAGGCGCAG
CTCCGGCAGAGGGAGGTGGTGGACCTGTATAATGGAATGTGCTTACAAGGGCCAGCAGGAGTGCC
TGGTCGAGACGGGAGCCCTGGGGCCAATGTTATTCCGGGTACACCTGGGATCCCAGGTCCGGGATG
GATTCAAAGGAGAAAAGGGGGAATGTCTGAGGGAAAGCTTTGAGGAGTCCTGGACACCCAACTAC
AAGCAGTGTTTCATGGAGTTCATTGAATTATGGCATAGATCTTGGGAAAATTGCCGAGTGTTACATT
TACAAAGATGCGTTCAAATAGTGCTCTAAGAGTTTTGTTTCAGTGGCTCACTTCGGCTAAAATGCA
GAAATGCATGCTGTGACGTTGGTATTTTACATTCAATGGAGCTGAATGTTCAGGACCTCTTCCC
ATTGAAGCTATAATTTATTTGGACCAAGGAAGCCCTGAAATGAATTCAACAATTAATATTCATCG
CACTTCTTCTGTGGAAGGACTTTGTGAAGGAATTGGTGTGATTAGTGGATGTTGCTATCTGGG
TTGGCACTTGTTTCAGATTACCCAAAAGGAGATGCTTCTACTGGATGGAATTCAGTTTCTCGCATC
ATTATTGAAGAACTACCAAAATAAATGCTTTAATTTTCATTTGCTACCTCTTTTTTTATTATGCC
TTGGAATGGTTCACTTAAATGACATTTTAAATAAGTTTATGTATACATCTGAATGAAAAGCAAAG
CTAAATATGTTTACAGACCAAAGTGTGATTTACACTGTTTTTAAATCTAGCATTATTCATTTTG
CTTCAATCAAAAGTGGTTTCAATATTTTTTTTAGTTGGTTAGAATACTTTCTTCATAGTCACATT
CTCTCAACCTATAATTTGGAATATTGTTGTGGTCTTTTGTTTTTCTCTTAGTATAGCATTTTTTA
AAAAATATAAAAGCTACCAATCTTTGTACAATTTGTAAATGTTAAGAATTTTTTTTATATCTGT
TAAATAAAAATTATTTCCAACA

FIGURE 122

MRPQGPAASPQRLRGLLLLLLLQLPAPSSASEIPKGGKQKAQLRQREVVDLYNGMCLQGPAAGVPGR
DGSPGANVIPGTPGIPGRDGFKGEGECLRESFEESWTPNYKQCSWSSLNYGIDLGKIAECTFTK
MRSNSALRVLFSGSLRLKCRNACCQRWYFTFNGAECSGPLPIEAIYLDQGSPEMNSTINIHRTS
SVEGLCEGIGAGLVDVAIWVGTCSDYPKGDASTGWNSVSRIIIIEELPK

Signal peptide:

amino acids 1-30

Transmembrane domain:

amino acids 195-217

FIGURE 123

GCTGAGCGTGTGCGCGGTACGGGGCTCTCCTGCCTTCTGGGCTCCAACGCAGCTCTGTGGCTGAA
 CTGGGTGCTCATCACGGGAAGTCTGGGCTATGGAATACAGATGTGGCAGCTCAGGTAGCCCCAA
 ATTGCCTGGAAGAATACATCATGTTTTTCGATAAGAAGAAATTGTAGGATCCAGTTTTTTTTTTA
 ACCGCCCCCTCCCCACCCCCCAAAAAAAGTGTAAAGATGCAAAAACGTAATATCCATGAAGATCC
 TATTACCTAGGAAGATTTTGATGTTTTGCTGCGAATGCGGTGTTGGGATTTATTTGTTCTTGAG
 TGTTCTGCGTGGCTGGCAAAGAATAATGTTCCAAAATCGGTCCATCTCCCAAGGGGTCCAATTTT
 TCTTCCTGGGTGTGAGCGAGCCCTGACTCACTACAGTGCAGCTGACAGGGGCTGTGATGCAACTG
 GCCCCTAAGCCAAAGCAAAAGACCTAAGGACGACCTTTGAACAATACAAAGGATGGGTTTCAATG
 TAATTAGGCTACTGAGCGGATCAGCTGTAGCACTGGTTATAGCCCCACTGTCTTACTGACAATG
 CTTTCTTCTGCCGAACGAGGATGCCCTAAGGGCTGTAGGTGTGAAGGCAAAATGGTATATTGTGA
 ATCTCAGAAATTACAGGAGATACCCCTCAAGTATATCTGCTGGTTGCTTAGGTTTGTCCCTTCGCT
 ATAACAGCCTTCAAAAAGTAAAGTATAATCAATTTAAAGGGCTCAACCAGCTCACCTGGCTATAC
 CTTGACCATAACCATCAGCAATATTGACGAAAAATGCTTTTAATGGAATACGCAGACTCAAAGA
 GCTGATTCTTAGTTCCAATAGAATCTCCTATTTTCTTAACAATACCTTCAGACCTGTGACAAATT
 TACGGAACCTTGGATCTGTCTATAATCAGCTGCATTCTCTGGGATCTGAACAGTTTCGGGGCTTG
 CGGAAGCTGCTGAGTTTACATTTACGGTCTAACCCTGAGAACCATGCCCTGTGCGAATATTCCA
 AGACTGCCGCAACCTGGAACCTTTGGACCTGGGATATAACCGGATCCGAAGTTTAGCCAGGAATG
 TCTTTGCTGGCATGATCAGACTCAAAGAAGTTCACCTGGAGCACAATCAATTTTCCAAGCTCAAC
 CTGGCCCTTTTTTCCAAGGTTGGTCAGCCTTCAGAACCTTTACTTGCAAGTGAATAAAATCAGTGT
 CATAGGACAGACCATGTCCTGGACCTGGAGCTCCTTACAAAGGCTTGATTTATCAGGCAATGAGA
 TCGAAGCTTTTCAGTGGACCCAGTGTTTTCCAGTGTGTCCCGAATCTGCAGCGCCTCAACCTGGAT
 TCCAACAAGCTCACATTTATTGGTCAAGAGATTTTGGATTCTTGATATCCCTCAATGACATCAG
 TCTTGCTGGGAATATATGGGAATGCAGCAGAAATATTTGCTCCCTTGTAAGTGGCTGAAAAGTT
 TTAAAGGTCTAAGGGAGAATACAATTATCTGTGCCAGTCCCAAAGAGCTGCAAGGAGTAAATGTG
 ATCGATGCAGTGAAGAACTACAGCATCTGTGGCAAAAGTACTACAGAGAGGTTTGATCTGGCCAG
 GGCTCTCCCAAAGCCGACGTTTAAAGCCCAAGCTCCCCAGGCCGAAGCATGAGAGCAAACCCCTT
 TGCCCCCGACGGTGGGAGCCACAGAGCCCGGCCAGAGACCGATGCTGACGCCGAGCACATCTCT
 TTCCATAAAATCATCGCGGGCAGCGTGGCGCTTTTCTGTCCGTGCTCGTCATCCTGCTGGTTAT
 CTACGTGTGATGGAAGCGGTACCCTGCGAGCATGAAGCAGCTGCAGCAGCGCTCCCTCATGCCAA
 GGCACAGGAAAAAGAAAAGACAGTCCCTAAAGCAAATGACTCCAGCACCCAGGAATTTTATGTA
 GATTATAAACCACCAACACGGAGACCAGCGAGATGCTGCTGAATGGGACGGGACCCCTGCACCTA
 TAACAAATCGGGCTCCAGGGAGTGTGAGGTATGAACCATTTGTGATAAAAAGAGCTCTTAAAGCT
 GGGAAATAAGTGGTGTCTTTATTGAACTCTGGTGACTATCAAGGGAACGCGATGCCCCCTCCCC
 TTCCCTCTCCCTCTCACTTTGGTGGCAAGATCCTTCCTTGTCGGTTTTAGTGCATTCATAATACT
 GGTCATTTTCTCTCATACATAATCAACCCATTGAAATTTAAATACCACAATCAATGTGAAGCTT
 GAACTCCGTTTTAATATAATACCTATTGTATAAGACCCTTTACTGATTCCATTAATGTGCGATTT
 GTTTTAAGATAAACTTCTTTCATAGGTAAAAA

FIGURE 124

MGFNVIRLLSGSAVALVIAPTVLLTMLSSAERGCPKGCRCEGKMVYCESQKLQEIPSSISAGCLG
LSLRYNLSLQKLKYNQFKGLNQLTWLYLDHNNHISNIDENAFNGIRRLKELILSSNRISYFLNNTFR
PVTNLRNLDSLQNQLHSLGSEQFRGLRKLLSLHLRSNSLRTIPVRI FQDCRNLELLDLGYNRIRS
LARNVFAGMIRLKEHLEHNQFSKLNALFPRLVSLQNLQWNKISVIGQTMSTWSSLQRLDL
SGNEIEAFSGPSVFQCVPNLQRLNLDNKLTFIGQEILDSWISLNDISLAGNIWECSRNICSLVN
WLKSEFKGLRENTIICASPKELOGVNVDAVKNYSICGKSTTERFDLARALPKPTFKPKLPRPKHE
SKPPLPPTVGATEPGPETDADAEHISFHKIIAGSVALFLSVLVILLVIYVSWKRYPASMKQLQQR
SLMRRHRKKKRQSLKQMPSTQEFYVDYKPTNTETSEMLLNGTGPCTYNKSGSRECEV

Important features of the protein:

Signal peptide:

amino acids 1-33

Transmembrane domain:

amino acids 420-442

N-glycosylation sites.

amino acids 126-129, 357-360, 496-499, 504-507

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 465-468

Tyrosine kinase phosphorylation site.

amino acids 136-142

N-myristoylation sites.

amino acids 11-16, 33-38, 245-250, 332-337, 497-502, 507-512

FIGURE 125

CCGTTATCGTCTTGCGCTACTGCTGAATGTCCGTCCCGGAGGAGGAGAGGCTTTTGCCGCTG
ACCCAGAGATGGCCCCGAGCGAGCAAATTCCTACTGTCCGGCTGCGCGGCTACCGTGCCGAGCT
AGCAACCTTTCCCCTGGATCTCACAAAACTCGACTCCAAATGCAAGGAGAAGCAGCTCTTGCTC
GGTTGGGAGACGGTGCAAGAGAATCTGCCCCCTATAGGGGAATGGTGCGCACAGCCCTAGGGATC
ATTGAAGAGGAAGGCTTTCTAAAGCTTTGGCAAGGAGTGACACCCGCCATTTACAGACACGTAGT
GTATTCTGGAGGTCGAATGGTCACATATGAACATCTCCGAGAGGTTGTGTTTGGCAAAAGTGAAG
ATGAGCATTATCCCCCTTGGAATCAGTCATTGGAGGGATGATGGCTGGTGTTATTGGCCAGTTT
TTAGCCAATCCAACCTGACCTAGTGAAGGTTGAGATGCAAATGGAAGGAAAAGGAACTGGAAGG
AAAACCATTTGCGATTTCTGTGGTGTACATCATGCATTTGCAAAAATCTTAGCTGAAGGAGGAATAC
GAGGGCTTTGGGCAGGCTGGGTACCCAATATACAAAGAGCAGCACTGGTGAATATGGGAGATTTA
ACCACTTATGATACAGTGAAACACTACTTGGTATTGAATACACCACTTGAGGACAATATCATGAC
TCACGGTTTATCAAGTTTATGTTCTGGACTGGTAGCTTCTATTCTGGGAACACCAGCCGATGTCA
TCAAAAGCAGAATAATGAATCAACCACGAGATAAACAAGGAAGGGGACTTTTGTATAAATCATCG
ACTGACTGCTTGATTGAGGCTGTTCAAGGTGAAGGATTCATGAGTCTATATAAAGGCTTTTTACC
ATCTTGGCTGAGAATGACCCCTTGGTCAATGGTGTCTGGCTTACTTATGAAAAATCAGAGAGA
TGAGTGGAGTCAGTCCATTTTAA

FIGURE 126

MSVPEEEEEERLLPLTQRWPRASKFLLSGCAATVAELATFFPLDLTKTRLQMQGEAALARLGDGARES
APYRGMVRTALGIIEEEGFLKLWQGVTPAIYRHVVYSGGRMVITYEHLREVVFVGKSEDEHYPLWKS
VIGGMMAGVIGQFLANPTDLVKVQMMEGKRKLEGKPLRFRGVHHAFAKILAEGGIRGLWAGWVP
NIQRAALVNMGDLTTYDITVKHYLVNTPLEDNIMTHGLSSLCGLVASILGTPADVIKSRIMNQP
RDKQGRGLLYKSSTDCLIQAVQGEFMSLYKGFLPSWLRMTPWSMVFWLTYEKIREMSGVSPF

Transmembrane domains:

amino acids 25-38, 130-147, 233-248

FIGURE 127

CGCGGATCGGACCCAAGCAGGTCGGCGGCGGGCGGCAGGAGAGCGGCCGGGCGTCAGCTCCTCGAC
CCCCGTGTGGGCTAGTCCAGCGAGGCGGACGGGCGGCGTGGGCCCCATGGCCAGGCCCGGCATGG
AGCGGTGGCGCGACCGGCTGGCGCTGGTGACGGGGGCCCTCGGGGGGCATCGGCGCGGGCCGTGGCC
CGGGCCCTGGTCCAGCAGGGACTGAAGGTGGTGGGCTGCGCCCGCACTGTGGGCAACATCGAGGA
GCTGGCTGCTGAATGTAAGAGTGCAGGCTACCCCGGGACTTTGATCCCCTACAGATGTGACCTAT
CAAATGAAGAGGACATCCTCTCCATGTTCTCAGCTATCCGTTCTCAGCACAGCGGTGTAGACATC
TGCATCAACAATGCTGGCTTGGCCCGGCCCTGACACCCTGCTCTCAGGCAGCACCAGTGGTTGGAA
GGACATGTTCAATGTGAACGTGCTGGCCCTCAGCATCTGCACACGGGAAGCCTACCAGTCCATGA
AGGAGCGGAATGTGGACGATGGGCACATCATTAAACATCAATAGCATGTCTGGCCACCGAGTGTTA
CCCCTGTCTGTGACCCACTTCTATAGTGCCACCAAGTATGCCGTCCTGCGCTGACAGAGGGACT
GAGGCAAGAGCTTCGGGAGGCCCAGACCCACATCCGAGCCACGTGCATCTCTCCAGGTGTGGTGG
AGACACAATTCGCCTTCAAACCTCCACGACAAGGACCCTGAGAAGGCAGCTGCCACCTATGAGCAA
ATGAAGTGTCTCAAACCCGAGGATGTGGCCGAGGCTGTTATCTACGTCCTCAGCACCCCCGCACA
CATCCAGATTGGAGACATCCAGATGAGGCCCACGGAGCAGGTGACCTAGTGACTGTGGGAGCTCC
TCCTTCCCTCCCCACCCTTCATGGCTTGCCTCCTGCCTCTGGATTTTAGGTGTTGATTTCTGGAT
CACGGGATACCACTTCCTGTCCACACCCCGACCAGGGGCTAGAAAATTTGTTTGAGATTTTATA
TCATCTTGTCAAATTGCTTCAGTTGTAAATGTGAAAAATGGGCTGGGGAAAGGAGGTGGTGTCCC
TAATTGTTTTACTTGTTAACTTGTTCTTGTTGCCCTGGGCACTTGGCCTTTGTCTGCTCTCAGTG
TCTTCCCTTTGACATGGGAAAGGAGTTGTGGCCAAAATCCCCATCTTCTTGACCTCAACGTCTG
TGGCTCAGGGCTGGGGTGGCAGAGGGAGGCCTTACCTTATATCTGTGTTGTTATCCAGGGCTCC
AGACTTCCTCCTCTGCCTGCCCCACTGCACCCTCTCCCCCTTATCTATCTCCTTCTCGGCTCCCC
AGCCCAGTCTTGGCTTCTTGTCCCTCCTGGGGTCATCCCTCCACTCTGACTCTGACTATGGCAG
CAGAACACCAGGGCCTGGCCCAGTGGATTTTCATGGTGATCATTAAGAAAGAAAAATCGCAACCAA
AAAAAAAAA

FIGURE 128

MARPGMERWRDRLALVTGASGGIGA AAVARALVQQGLKVVG CARTVGNIEELAAECKSAGYPGTLI
PYRCDLSNEEDILSMFSAIRSQHSGVDICINNAGLARPD TLLSGSTSGWKDMFNVNVLALSICTR
EAYQSMKERNVDDGHIININSMGHRVLP LSVTHFY SATKYAVTALTEGLRQELREAQTHIRATC
ISPGVVETQFAFKLHDKDPEKAAATYEQMKCLKPEDVAEAVIYVLSTPAHIQIGDIQMRPTEQVT

Important features of the protein:

Signal peptide:

amino acids 1-17

N-myristoylation sites.

amino acids 18-24, 21-27, 22-28, 24-30, 40-46, 90-96, 109-115,
199-205

Short-chain alcohol dehydrogenase.

amino acids 30-42, 104-114

FIGURE 129

AACTTCTACATGGGCCTCCTGCTGCTGGTGCTCTTCCTCAGCCTCCTGCCGGTGGCCTACACCAT
CATGTCCCTCCCACCCTCCTTTGACTGCGGGCCGTTCAGGTGCAGAGTCTCAGTTGCCCGGGAGC
ACCTCCCCTCCCGAGGCAGTCTGCTCAGAGGGCCTCGGCCCAGAATTCCAGTTCTGGTTTCATGC
CAGCCTGTAAAAGGCCATGGAACTTTGGGTGAATCACCGATGCCATTTAAGAGGGTTTTCTGCCA
GGATGGAAATGTTAGGTCGTTCTGTGTCTGCGCTGTTTCATTTTCAGTAGCCACCAGCCACCTGTGG
CCGTTGAGTGCTTGAAATGAGGAACTGAGAAAATTAATTTCTCATGTATTTTTCTCATTTATTTA
TTAATTTTTAACTGATAGTTGTACATATTTGGGGGTACATGTGATATTTGGATACATGTATACAA
TATATAATGATCAAATCAGGGTAACTGGGATATCCATCACATCAAACATTTATTTTTATTCTTT
TTAGACAGAGTCTCACTCTGTCACCCAGGCTGGAGTGCAGTGGTGCCATCTCAGCTTACTGCAAC
CTCTGCCTGCCAGGTTCAAGCGATTCTCATGCCTCCACCTCCCAAGTAGCTGGGACTACAGGCAT
GCACCACAATGCCCAACTAATTTTTGTATTTTTAGTAGAGACGGGGTTTTGCCATGTTGCCCAGG
CTGGCCTTGAACCTCCTGGCCTCAAACAATCCACTTGCTCGGCCTCCCAAAGTGTTATGATTACA
GGCGTGAGCCACCGTGCTGGCCTAAACATTTATCTTTCTTTGTGTTGGGAACTTTGAAATTAT
ACAATGAATTATTGTTAACTGTCTCTCCCTGCTGTGCTATGGAACACTGGGACTTCTTCCCTCT
ATCTAACTGTATATTTGTACCAGTTAACCAACCGTACTTCATCCCCACTCCTCTCTATCCTTCCC
AACCTCTGATCACCTCATTCTACTCTCTACCTCCATGAGATCCACTTTTTAGCTCCCATGTG
AGTAAGAAAATGCAATATTTGTCTTTCTGTGCCTGGCTTATTTCACTTAACATAATGACTTCCTG
TTCCATCCATGTTGCTGCAAATGACAGGATTCGTTCTTAATTTCAATTAAATAACACACATG
GCAAAAA

FIGURE 130

MGLLLLVLFLSLLPVAYTIMSLPPSFDCGPFRCRVSVAREHLPSRGSLLRGPRPRIPVLVSCQPV
KGHGTLGESPMFVKRVFCQDGNVRSFCVCAVHFSSHQPPVAVECLK

Important features of the protein:

Signal peptide:

amino acids 1-18

N-myristoylation site.

amino acids 86-92

Zinc carboxypeptidases, zinc-binding region 2 signature.

amino acids 68-79

FIGURE 131

TTCTGAAGTAACGGAAGCTACCTTGTATAAAGACCTCAACACTGCTGACCATGATCAGCGCAGCCTGGAGC
ATCTTCCTCATCGGGACTAAAAATTGGGCTGTTCCCTTCAAGTAGCACCTCTATCAGTTATGGCTAAATCCTG
TCCATCTGTGTGTCGCTGCGATGCGGGTTTCATTTACTGTAATGATCGCTTCTGACATCCATTCCAACAG
GAATACCAGAGGATGCTACAACCTCTTACCTTCAGAACAACCAAATAAATAATGCTGGGATTCCCTTCAGAT
TTGAAAAACTTGCTGAAAGTAGAAAGAATATACCTTATACCACAACAGTTTAGATGAATTTCTTACCAACCT
CCCAAAGTATGTAAAAGAGTTACATTTGCAAGAAAATAACATAAGGACTATCACTTATGATTCACTTTCAA
AAATTCCTTATCTGGAAGAATTACATTTAGATGACAACCTCTGTCTCTGCAGTTAGCATAGAAGAGGGAGCA
TTCCGAGACAGCAACTATCTCCGACTGCTTTTCCCTGTCCCGTAATCACCTTAGCACAAATCCCTGGGGTTT
GCCCAGGACTATAGAAGAACTACGCTTGGATGATAATCGCATATCCACTATTTATCACCATCTCTTCAAG
GTCTCACTAGTCTAAAACGCCTGGTTCTAGATGGAAACCTGTTGAACAATCATGGTTTAGGTGACAAAGTT
TTCTTCAACCTAGTTAATTTGACAGAGCTGTCCCTGGTGCGGAATTCCTGACTGCTGCACCAGTAAACCT
TCCAGGCACAAACCTGAGGAAGCTTTATCTTCAAGATAACCACATCAATCGGGTGCCCCAAATGCTTTTT
CTTATCTAAGGCAGCTCTATCGACTGGATATGTCCAATAATAACCTAAGTAATTTACCTCAGGGTATCTTT
GATGATTTGGACAATATAACACAACCTGATTCTTCGCAACAATCCCTGGTATTGCGGGTGCAAGATGAAATG
GGTACGTGACTGGTTACAATCACTACCTGTGAAGGTCAACGTGCGTGGGCTCATGTGCCAAGCCCCAGAAA
AGGTTCTGTTGGGATGGCTATTAAGGATCTCAATGCAGAACTGTTTGATTGTAAGGACAGTGGGATTGTAAGC
ACCATTCAAGATAACCACTGCAATACCCAACACAGTGTATCCTGCCAAGGACAGTGGCCAGCTCCAGTGAC
CAAACAGCCAGATATTAAGAACCCCAAGCTCACTAAGGATCAACAAACCACAGGGAGTCCCTCAAGAAAAA
CAATTACAATTACTGTGAAGTCTGTCACCTCTGATACCATTCAATATCTCTTGAAACTTGCTCTACCTATG
ACTGCTTTGAGACTCAGCTGGCTTAAACTGGGCCATAGCCCGGCATTTGGATCTATAACAGAAACAATTGT
AACAGGGGAACGCAGTGAGTACTTGGTCACAGCCCTGGAGCCTGATTCACCCTATAAAGTATGCATGGTTT
CCATGGAAACCAGCAACCTCTACCTATTTGATGAAACTCCTGTTTGTATTGAGACTGAAACTGCACCCCTT
CGAATGTACAACCCTACAACCACCTCAATCGAGAGCAAGAGAAAGAACCCTTACAAAACCCCAATTTACC
TTTGGCTGCCATCATTGGTGGGGCTGTGGCCCTGGTTACCATTGCCCTTCTTGCTTTAGTGTGTTGGTATG
TTCATAGGAATGGATCGCTCTTCTCAAGGAACCTGTGCATATAGCAAAGGGAGGAGAAGAAAGGATGACTAT
GCAGAAGCTGGCACTAAGAAGGACAACCTCTATCCTGGAAATCAGGGAACTTCTTTTCAGATGTTACCAAT
AAGCAATGAACCCATCTCGAAGGAGGAGTTTGTAATACACACCATATTTCTCCTAATGGAATGAATCTGT
ACAAAACAATCACAGTGAAAGCAGTAGTAACCGAAGCTACAGAGACAGTGGTATTCCAGACTCAGATCAC
TCACACTCATGATGTGAAGGACTCACAGCAGACTTGTGTTTTGGGTTTTTTAAACCTAAGGGAGGTGATG
GT

FIGURE 132

MISAAWSIFLIGTKIGLFLQVAPLSVMAKSCPSVCRC DAGFIYCND RFLTSIPTGIPEDATTLYL
QNNQINNAGIPSDLKNLLKVERIYLYHNSLDEFPTNLPKYVKELHLQENNI RTITYDSLSKIPYL
EELHLLDDNSVSAVSIEEGA FRDSNYLRLLFLSRNHLSTIPWGLPRTIEELRLDDNRISTISSPSL
QGLTSLKRLVLDGNLLNNHGLGDKVFFNLVNLTELSLVRNSLTAAPVNLPGTNLRKLYLQDNHIN
RVPPNAFSYLRQLYRLDMSNNLSNLPQGIFDDLDNITQLILRN NPWYCGCKMKWVRDWLQSLPV
KVNVRGLMCQAPEKVRGMAIKDLNAELFDCKDSGIVSTIQITTAIPNTVYPAQGQWPAPVTKQPD
IKNPKLTKDQQTGSPSRKTITITVKSVTSDTIHISWKLALPMTALRLSWLKLGHSPAFGSITET
IVTGERSEYLVTALEPDSPYKVCMPMETSNLYLFDETPVC IETETAPLRMYNPPTTTLNREQEKE
PYKNPNLPLAAIIGGAVALVTIALLLALVCWYVHRNGSLFSRNCAYS KGRRRKDDYAEAGTKKDNS
ILEIRETSFQMLPISNEPISKEEFVIHTIFPPNGMNL YKNNHSESSSNRSYRDSGIPDSDHSHS

Important features of the protein:

Signal peptide:

amino acids 1-28

Transmembrane domain:

amino acids 531-552

N-glycosylation sites.

amino acids 226-229, 282-285, 296-299, 555-558, 626-629, 633-636

Tyrosine kinase phosphorylation site.

amino acids 515-522

N-myristoylation sites.

amino acids 12-17, 172-177, 208-213, 359-364, 534-539, 556-561,
640-645

Amidation site.

amino acids 567-570

Leucine zipper pattern.

amino acids 159-180

Phospholipase A2 aspartic acid active site.

amino acids 34-44

FIGURE 133

CCGTCATCCCCCTGCAGCCACCCTTCCCAGAGTCCTTTGCCCAGGCCACCCCAGGCTTCTTGCCA
GCCCTGCCGGGGCCACTTGTCTTCAATGTCTGCCAGGGGGAGGTGGGAAGGAGGTGGGAGGAGGGCG
TGCAGAGGCAGTCTGGGCTTGGCCAGAGCTCAGGGTGCTGAGCGTGTGACCAGCAGTGAGCAGAG
GCCGGCCATGGCCAGCCTGGGGCTGCTGCTCCTGCTCTTACTGACAGCACTGCCACCGCTGTGGT
CCTCCTCACTGCCTGGGCTGGACACTGCTGAAAGTAAAGCCACCATTGCAGACCTGATCCTGTCT
GCGCTGGAGAGAGCCACCGTCTTCCCTAGAACAGAGGCTGCCTGAAATCAACCTGGATGGCATGGT
GGGGGTCCGAGTGCTGGAAGAGCAGCTAAAAAGTGTCCGGGAGAAGTGGGCCCAGGAGCCCCCTGC
TGCAGCCGCTGAGCCTGCGCGTGGGGATGCTGGGGGAGAAGCTGGAGGCTGCCATCCAGAGATCC
CTCCACTACCTCAAGCTGAGTGATCCCAAGTACCTAAGAGAGTTCCAGCTGACCCTCCAGCCCCG
GTTTTGGAAGCTCCCACATGCCTGGATCCACACTGATGCCTCCTTGGTGTACCCACGTTCCGGGC
CCCAGGACTCATTCTCAGAGGAGAGAAGTGACGTGTGCCTGGTGCAGCTGCTGGGAACCGGGACG
GACAGCAGCGAGCCCCTGCGGCCTCTCAGACCTCTGCAGGAGCCTCATGACCAAGCCCGGCTGCTC
AGGCTACTGCCTGTCCCACTGCTCTTCTTCTCTGCGCCAGAATGAGGGGATGCACACAGG
GACCACTCCAACAGAGCCAGGACTATATCAACCTCTTCTGCGCCAACATGATGGACTTGAACCGC
AGAGCTGAGGCCATCGGATACGCTACCCCTACCCGGGACATCTTCATGGAAAACATCATGTTCTG
TGGAATGGGCGGCTTCTCCGACTTCTACAAGCTCCGGTGGCTGGAGGCCATTCTCAGCTGGCAGA
AACAGCAGGAAGGATGCTTCGGGGAGCCTGATGCTGAAGATGAAGAATTATCTAAAGCTATTCAA
TATCAGCAGCATTTTTTCGAGGAGAGTGAAGAGGCGAGAAAAACAATTTCCAGATTCTCGCTCTGT
TGCTCAGGCTGGAGTACAGTGGCGCAATCTCGGCTCACTGCAACCTTTGCCTCCTGGGTTCAGC
AATTCTCTTGCTCATCTCCCGAGTAGCTGGGACTACAGGAGCGTGCCACCATACTGGCTAAT
TTTTATATTTTTTTAGTAGAGACAGGGTTTCATCATGTGCTCATGCTGGTCTCGAACTCCTGAT
CTCAAGAGATCCGCCCACCTCAGGCTCCCAAAGTGTGGGATTATAGGTGTGAGCCACCGTGTCTG
GCTGAAAAGCACTTTCAAAGAGACTGTGTTGAATAAAGGGCCAAGTTCTTGCCACCCAGCACTC
ATGGGGGCTCTCTCCCCTAGATGGCTGCTCCTCCCAACACAGCCACAGCAGTGGCAGCCCTGG
GTGGCTTCTTATACATCCTGGCAGAATACCCCCCAGCAAACAGAGAGCCACCCCATCCACACCG
CCACCACCAAGCAGCCGCTGAGACGGACGGTTCCATGCCAGCTGCCTGGAGGAGGAACAGACCCC
TTTAGTCCTCATCCCTTAGATCCTGGAGGGCACGGATCACATCCTGGGAAGAAGGCATCTGGAGG
ATAAGCAAAGCCACCCCGACACCCAATCTTGGAAGCCCTGAGTAGGCAGGGCCAGGGTAGGTGGG
GGCCGGGAGGGACCCAGGTGTGAACGGATGAATAAAGTTCAACTGCAACTGAAAAAAAAA

FIGURE 134

MSARGRWEGGGRRACRGSGLGLARAQGAERVTSSEQRPAMASLGLLLLLLLLLTALPPLWSSSLPGLD
TAESKATIADLILSALERATVFLEQRLPEINLDGMVGVRVLEEQLKSVREKWAQEPLLQPLSLRV
GMLGEKLEAAIQRSYLHYLKLSDPKYLREFQLTLQPGFWKLPHAWIHTDASLVYPTFGPQDSFSEE
RSDVCLVQLLGTGTDSEPCGLSDLCRSLMTKPGCSGYCLSHQLLFFLWARMRGCTQGPLQQSQD
YINLFCANMMDLNRRAEAIGYAYPTRDI FMENIMFCGMGGFSDFYKLRWLEAILSQKQEGCFG
EPDAEDEELSKAIQYQQHFSRRVKRREKQFPDSRSVAQAGVQWRNLGSLQPLPPGFKQFSLILP
SSWDYRSVPPYLANFYIFLVETGFHHVAHAGLELLISRDPTSGSQSVGL

Important features of the protein:

Signal peptide:

amino acids 1-26

Transmembrane domain:

amino acids 39-56

Tyrosine kinase phosphorylation sites.

amino acids 149-156, 274-282

N-myristoylation sites.

amino acids 10-16, 20-26, 63-69, 208-214

Amidation site.

amino acids 10-14

Glycoprotein hormones beta chain signature 1.

amino acids 230-237

FIGURE 135

GGTCTGAGTGCAGAGCTGCTGTCCATGGCGGCCGCTCTGTGGGGCTTCTTTCCCGTCCTGCTGCTG
CTGCTGCTATCGGGGGATGTCCAGAGCTCGGAGGTGCCCGGGGCTGCTGCTGAGGGATCGGGAGG
GAGTGGGGTCGGCATAGGAGATCGCTTCAAGATTGAGGGGCGTGCAGTTGTTCCAGGGGTGAAGC
CTCAGGACTGGATCTCGGCGGCCCGAGTGCTGGTAGACGGAGAAGAGCACGTCGGTTTCTCTTAAG
ACAGATGGGAGTTTTGTGGTTCATGATATACCTTCTGGATCTTATGTAGTGGAAGTTGTATCTCC
AGCTTACAGATTTGATCCCGTTCGAGTGGATATCACTTCGAAAGGAAAAATGAGAGCAAGATATG
TGAATTACATCAAAACATCAGAGGTTGTCAGACTGCCCTATCCTCTCCAAATGAAATCTTCAGGT
CCACCTTCTTACTTTATTAAAAGGGAATCGTGGGGCTGGACAGACTTTCCTAATGAACCCAATGGT
TATGATGATGGTTCTTCCTTTATTGATATTTGTGCTTCTGCCTAAAGTGGTCAACACAAGTGATC
CTGACATGAGACGGGAAATGGAGCAGTCAATGAATATGCTGAATTCCAACCATGAGTTGCCTGAT
GTTTCTGAGTTCATGACAAGACTCTTCTCTTCAAAATCATCTGGCAAATCTAGCAGCGGCAGCAG
TAAACAGGCAAAAGTGGGGCTGGCAAAGGAGGTAGTCAGGCCGTCCAGAGCTGGCATTTCAC
AAACACGGCAACACTGGGTGGCATCCAAGTCTTGAAAACCGTGTGAAGCAACTACTATAAACTT
GAGTCATCCCGACGTTGATCTCTTACAACCTGTGTATGTT
AACTTTTTAGCACATGTTTTGTACTTGGTACACGAGAAAACCCAGCTTTCATCTTTTGTCTGTAT
GAGGTCAATATTGATGTCACTGAATTAATTACAGTGTCCTATAGAAAATGCCATTAATAAATTAT
ATGAACTACTATACATTATGTATATTAATTAAAACATCTTAATCCAGAAATCAAAAAAAAAAAAAA
AAAAAAAAAAAAAAAA

FIGURE 136

MAAALWGFFPVLLLLLLSGDVQSSEVPGAAAEGSGSGVGIGDRFKIEGRAVVPGVKPDWISAA
RVLVDGEEHVGFLLKTDGSFVVHDIPSGSYVVEVVSAPAYRFDVVRVDITSGKGMRRARYVNYIKTSE
VVRLPYPLQMKSSGPPSYFIKRESWGWTDFLMNPMVMMMLVLPLLIFVLLPKVVNTSDPDMRREME
QSMNMLNSNHELPDVSEFMTRLFSSKSSGKSSSGSSSKTGKSGAGKRR

Important features of the protein:

Signal sequence:

amino acids 1-23

Transmembrane domain:

amino acids 161-182

N-glycosylation site.

amino acids 184-187

Glycosaminoglycan attachment sites.

amino acids 37-40, 236-239

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 151-154

N-myristoylation sites.

amino acids 33-38, 36-41, 38-44, 229-234

Amidation site.

amino acids 238-241

ATP/GTP-binding site motif A (P-loop).

amino acids 229-236

FIGURE 137

GATGGCGCAGCCACAGCTTCTGTGAGATTTCGATTCTCCCCAGTTCCCCTGTGGGTCTGAGGGGA
CCAGAAGGGTGAGCTACGTTGGCTTTCTGGAAGGGGAGGCTATATGCGTCAATTCCCCAAAACAA
GTTTTGACATTTCCCCTGAAATGTCATTCTCTATCTATTCACTGCAAGTGCCTGCTGTTCCAGGC
CTTACCTGCTGGGCACTAACGGCGGAGCCAGGATGGGGACAGAATAAAGGAGCCACGACCTGTGC
CACCAACTCGCACTCAGACTCTGAACTCAGACCTGAAATCTTCTCTTCACGGGAGGCTTGGCAGT
TTTTCTTACTCCTGTGGTCTCCAGATTTCAGGCCTAAGATGAAAGCCTCTAGTCTTGCCTTCAGC
CTTCTCTCTGCTGCGTTTTATCTCCTATGGACTCCTTCCACTGGACTGAAGACACTCAATTTGGG
AAGCTGTGTGATCGCCACAAACCTTCAGGAAATACGAAATGGATTTTCTGAGATACGGGGCAGTG
TGCAAGCCAAAGATGGAAACATTGACATCAGAATCTTAAGGAGGACTGAGTCTTGAAGACACA
AAGCCTGCGAATCGATGCTGCCTCCTGCGCCATTTGCTAAGACTCTATCTGGACAGGGTATTTAA
AACTACCAGACCCCTGACCATTATACTCTCCGGAAGATCAGCAGCCTCGCCAATTCCTTTCTTA
CCATCAAGAAGGACCTCCGGCTCTCTCATGCCACATGACATGCCATTGTGGGGAGGAAGCAATG
AAGAAATACAGCCAGATTCTGAGTCACCTTTGAAAAGCTGGAACCTCAGGCAGCAGTTGTGAAGGC
TTTGGGGGAACTAGACATTCTTCTGCAATGGATGGAGGAGACAGAATAGGAGGAAAGTGATGCTG
CTGCTAAGAATATTTCGAGGTCAAGAGCTCCAGTCTTCAATACCTGCAGAGGAGGCATGACCCCAA
ACCACCATCTCTTTACTGTACTAGTCTTGTGCTGGTCACAGTGTATCTTATTTATGCATTACTTG
CTTCCTTGCATGATTGTCTTTATGCATCCCCAATCTTAATTGAGACCATACTTGTATAAGATTTT
TGTAATATCTTTCTGCTATTGGATATATTTATTAGTTAATATATTTATTTATTTTTTGCTATTTA
ATGTATTTATTTTTTTTACTTGGACATGAACTTTAAAAAAATTACAGATTATATTTATAACCTG
ACTAGAGCAGGTGATGTATTTTTATACAGTAAAAAAAAAAAAACCTTGTAATTCTAGAAGAGTGG
CTAGGGGGGTTATTCATTTGTATTCAACTAAGGACATATTTACTCATGCTGATGCTCTGTGAGAT
ATTTGAAATTGAACCAATGACTACTTAGGATGGGTTGTGGAATAAGTTTTGATGTGGAATTGCAC
ATCTACCTTACAATTACTGACCATCCCCAGTAGACTCCCCAGTCCCATAATTGTGTATCTTCCAG
CCAGGAATCCTACACGGCCAGCATGTATTTCTACAAATAAAGTTTTCTTTGCATACCAAAAAAAA
AAAAAAAAAAAA

FIGURE 138

MRQFPKTSFDISPEMSFSIYSLQVPAVPG LTCWALTAEPGWGQNGATT CATNSHSDSEL RPEIF
SSREAWQFFLLLWSPDFRPKMKASSLAFSLLSAAFYLLWTPSTGLKTLNLGSCVIATNLQEIRNG
FSEIRGSVQAKDGNIDIRILRRTESLQDTKPANRCCLLRHLLRLYLDRVFKNYQTPDHYTLRKIS
SLANSFLT IKKDLRLSHAHTCHCGEEAMKKYSQILSHFEKLEPQAAVVKALGELDILLQWMEET
E

Important features of the protein:

Signal peptide:

amino acids 1-42

cAMP- and cGMP-dependent protein kinase phosphorylation sites.

amino acids 192-195, 225-228

N-myristoylation sites.

amino acids 42-47, 46-51, 136-141

FIGURE 139

CCTGGAGCCGGAAGCGCGGCTGCAGCAGGGCGAGGCTCCAGGTGGGGTTCGGTTCCGCATCCAGCC
 TAGCGTGTCCACGATGCGGCTGGGCTCCGGGACTTTTCGCTACCTGTTGCGTAGCGATCGAGGTGC
 TAGGGATCGCGGTCTTCCTTCGGGGATTCTTCCCGGCTCCCGTTCGTTCTCTGCCAGAGCGGAA
 CACGGAGCGGAGCCCCAGCGCCCGAACCCCTCGGCTGGAGCCAGTTCTAACTGGACCACGCTGCC
 ACCACCTCTCTTCAGTAAAGTTGTTATTGTTCTGATAGATGCCTTGAGAGATGATTTTGTGTTTG
 GGTCAAAGGGTGTGAAATTTATGCCCTACACAACCTTACCTTGTGGAAAAAGGAGCATCTCACAGT
 TTTGTGGCTGAAGCAAAGCCACCTACAGTTACTATGCCTCGAATCAAGGCATTGATGACGGGGAG
 CCTTCCTGGCTTTGTTCGACGTCATCAGGAACCTCAATTCTCCTGCACTGCTGGAAGACAGTGTGA
 TAAGACAAGCAAAAGCAGCTGGAAAAAGAATAGTCTTTTATGGAGATGAAACCTGGGTAAATTA
 TTCCCAAAGCATTTTGTGGAATATGATGGAACAACCTCATTTTTCGTGTGAGATTACACAGAGGT
 GGATAATAATGTCACGAGGCATTTGGATAAAGTATTAAAAAGAGGAGATTGGGACATATTAATCC
 TCCACTACCTGGGGCTGGACCACATTGGCCACATTTTCAGGGCCCAACAGCCCCCTGATTGGGCGAG
 AAGCTGAGCGAGATGGACAGCGTGTGATGAAGATCCACACCTCACTGCAGTGAAGGAGAGAGA
 GACGCCTTTACCCAATTTGCTGGTTCTTTGTGGTGACCATGGCATGTCTGAAACAGGAAGTCACG
 GGGCCTCCTCCACCGAGGAGGTGAATACACCTCTGATTTTAATCAGTTCTGCGTTTGAAAGGAAA
 CCGGCTGATATCCGACATCCAAAGCACGTCCAATAGACGGATGTGGCTGCGACACTGGCGATAGC
 ACTTGGCTTACCGATTCCAAAAGACAGTGTAGGGAGCCTCCTATTCCCAGTTGTGGAAGGAAGAC
 CAATGAGAGAGCAGTTGAGATTTTACATTTGAATACAGTGCAGCTTAGTAAACTGTTGCAAGAG
 AATGTGCCGTCAATATGAAAAAGATCCTGGGTTTGAGCAGTTTAAATGTGAGAAAGATTGCATGG
 GAACTGGATCAGACTGTACTTGGAGGAAAAGCATTCAGAAAGTCCTATTCAACCTGGGCTCCAAGG
 TTCTCAGGCAGTACCTGGATGCTCTGAAGACGCTGAGCTTGTCCTGAGTGCACAAGTGGCCAG
 TTCTCACCTGCTCCTGCTCAGCGTCCCACAGGCACTGCACAGAAAGGCTGAGCTGGAAGTCCCA
 CTGTCACTCCTGGGTTTTCTCTGCTCTTTTATTTGGTGATCCTGGTTCTTTTCGGCCGTTACAGT
 CATTGTGTGCACCTCAGCTGAAAGTTCGTGCTACTTCTGTGSCCTCTCGTGGCTGGCGGCAGGCT
 GCCTTTCGTTTACCAGACTCTGGTTGAACACCTGGTGTGTGCCAAGTGTGTCAGTGCCCTGGAC
 AGGGGGCCTCAGGGAAGGACGTGGAGCAGCCTTATCCCAGGCCTCTGGGTGTCCCGACACAGGTG
 TTCACATCTGTGCTGTGAGGTGAGATGCCTCAGTTCTTGGAAAGCTAGGTTCTGCGACTGTTAC
 CAAGGTGATTGTAAAGAGCTGGCGGTACAGAGGAACAAGCCCCCAGCTGAGGGGGTGTGTGAA
 TCGGACAGCCTCCCAGCAGAGGTGTGGGAGCTGCAGCTGAGGGAAGAAGAGACAATCGGCCTGGA
 CACTCAGGAGGGTCAAAGGAGACTTGGTTCGACCACTCATCCTGCCACCCCCAGAATGCATCCT
 GCCTCATCAGGTCCAGATTTCTTTCCAAGGCGGACGTTTTCTGTTGGAATTCTTAGTCCTTGGCC
 TCGGACACCTTCATTTCGTTAGCTGGGGAGTGGTGGTGAGGCAGTGAAGAAGAGGCGGATGGTCAC
 ACTCAGATCCACAGAGCCCAGGATCAAGGGACCACTGCAGTGGCAGCAGGACTGTTGGGCCCCC
 ACCCAACCCTGCACAGCCCTCATCCCTCTTGGCTTGAGCCGTGAGAGGCCCTGTGCTGAGTGT
 CTGACCGAGACACTCACAGCTTTGTGATCAGGGCACAGGCTTCCTCGGAGCCAGGATGATCTGTG
 CCACGCTTGCACCTCGGGCCCATCTGGGCTCATGCTCTCTCCTGCTATTGAATTAGTACCTAG
 CTGCACACAGTATGTAGTTACCAAAAGAATAAACGGCAATAATTGAGAAAAAAA

FIGURE 140

MRLGSGTFATCCVAIEVLGIAVFLRGFFPAPVRSSARAEHGAEPPEPSAGASSNWTTLPPPLF
SKVVIVLIDALRDDFVFGSKGVKEMPYTTYLVEKGASHSFVAEAKPPTVTMPRIKALMTGSLPGF
VDVIRNLNSPALLEDVIRQAKAAGKRIVFYGDETWVKLFPKHFVEYDGTTSFFVSDYTEVDNNV
TRHLDKVLKRGDWDILILHYLGLDGHIGHISGPN SPLIGQKLSEMDSVLMKIHTSLQSKERETPLP
NLLVLCGDHGMSETGSHGASSTEEVNTPLILISSAFERKPGDIRHPKHVQ

Important features of the protein:

Signal peptide:

amino acids 1-34

Transmembrane domain:

amino acids 58-76

N-glycosylation sites.

amino acids 56-60, 194-198

N-myristoylation sites.

amino acids 6-12, 52-58, 100-106, 125-131, 233-239, 270-276,
275-281, 278-284

Amidation site.

amino acids 154-158

Cell attachment sequence.

amino acids 205-208

FIGURE 141

GGCACGAGGCAAGCCTTCCAGGTTATCGTGACGCACCTTGAAAGTCTGAGAGCTACTGCCCTACA
GAAAGTTACTAGTGCCCTAAAGCTGGCGCTGGCACTGATGTTACTGCTGCTGTTGGAGTACAAC
TCCCTATAGAAAACAACCTGCCAGCACCTTAAGACCACTCACACCTTCAGAGTGAAGAACTTAAAC
CCGAAGAAATTCAGCATTCATGACCAGGATCACAAAGTACTGGTCCTGGACTCTGGGAATCTCAT
AGCAGTTCAGATAAAAACTACATACGCCCAGAGATCTTCTTTGCATTAGCCTCATCCTTGAGCT
CAGCCTCTGCGGAGAAAGGAAGTCCGATTCTCCTGGGGGTCTCTAAAGGGGAGTTTTGTCTCTAC
TGTGACAAGGATAAAGGACAAAGTCATCCATCCCTTCAGCTGAAGAAGGAGAACTGATGAAGCT
GGCTGCCCAAAGGAATCAGCACGCCGGCCCTTCATCTTTTATAGGGCTCAGGTGGGCTCCTGGA
ACATGCTGGAGTCGGCGGCTCACCCCGGATGGTTCATCTGCACCTCCTGCAATTGTAATGAGCCT
GTTGGGGTGACAGATAAATTTGAGAACAGGAAACACATTGAATTTTCATTTCAACCAGTTTGCAA
AGCTGAAATGAGCCCCAGTGAGGTCAGCGATTAGGAACTGCCCCATTGAACGCCTTCCTCGCTA
ATTTGAACTAATTGTATAAAAACACCAAACCTGCTCACT

FIGURE 142

MLLLLLLEYNFPPIENNCQHLKTTHTFRVKNLNPKKFSIHDQDHKVLVLDSGNLIAVPDKNYIRPEI
FFALASSLSSASAEKGSPIILLGVSKGEFCLYCDKDKGQSHPSLQLKKEKLMKLAQKESARRPFI
FYRAQVGSWNMLESAAHPGWFICTSCNCNEPVGVTDKFENRKHIEFSFQPVCKAEMSPSEVSD

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 33-36

N-myristoylation site.

amino acids 50-55, 87-92

Interleukin-1

amino acids 37-182

FIGURE 144

MLGLPWKGGLSWALLLLLLLGSQILLIYAWHFHEQRDCDEHNVMARYLPATVEFAVHTFNQQSKDY
YAYRLGHILNSWKEQVESKTVFSMELLGRTRCGKFEDDIDNCHFQESTELNNTFTCFFTISTRP
WMTQFSLLNKTCLEGFH

Important features of the protein:

Signal peptide:

amino acids 1-25

N-glycosylation sites.

amino acids 117-121, 139-143

N-myristoylation site.

amino acids 9-15

FIGURE 145

CTGTGCAGCTCGAGGCTCCAGAGGCACACTCCAGAGAGAGCCAAGGTTCTGACGCGATGAGGAAG
CACCTGAGCTGGTGGTGGCTGGCCACTGTCTGCATGCTGCTCTTCAGCCACCTCTCTGCGGTCCA
GACGAGGGGCATCAAGCACAGAATCAAGTGAACCGGAAGGCCCTGCCCAGCACTGCCCAGATCA
CTGAGGCCCAGGTGGCTGAGAACCGCCCGGGAGCCTTCATCAAGCAAGGCCGCAAGCTCGACATT
GACTTCGGAGCCGAGGGCAACAGGTACTACGAGGCCAACTACTGGCAGTTCCTCCGATGGCATCCA
CTACAACGGCTGCTCTGAGGCTAATGTGACCAAGGAGGCATTTGTCACCGGCTGCATCAATGCCA
CCCAGGCGGCGAACCAGGGGGAGTTCCAGAAGCCAGACAACAAGCTCCACCAGCAGGTGCTCTGG
CGGCTGGTCCAGGAGCTCTGCTCCCTCAAGCATTGCGAGTTTTGGTTGGAGAGGGGCGCAGGACT
TCGGGTCACCATGCACCAGCCAGTGCTCCTCTGCCTTCTGGCTTTGATCTGGCTCATGGTGAAAT
AAGCTTGCCAGGAGGCTGGCAGTACAGAGCGCAGCAGCGAGCAAATCCTGGCAAGTGACCCAGCT
CTTCTCCCCCAAACCCACGCGTGTTCTGAAGGTGCCAGGAGCGGCGATGCACTCGCACTGCAAA
TGCCGCTCCCACGTATGCGCCCTGGTATGTGCCTGCGTTCTGATAGATGGGGGACTGTGGCTTCT
CCGTCACTCCATTCTCAGCCCCTAGCAGAGCGTCTGGCACACTAGATTAGTAGTAAATGCTTGAT
GAGAAGAACACATCAGGCACTGCGCCACCTGCTTCACAGTACTTCCCAACAACCTCTTAGAGGTAG
GTGTATTCCCGTTTTACAGATAAGGAACTGAGGCCCAGAGAGCTGAAGTACTGCACCCAGCATC
ACCAGCTAGAAAGTGGCAGAGCCAGGATTCAACCCTGGCTTGTCTAACCCAGGTTTTCTGCTCT
GTCCAATTCCAGAGCTGTCTGGTGATCACTTTATGTCTCACAGGGACCCACATCCAAACATGTAT
CTCTAATGAAATTGTGAAAGCTCCATGTTTAGAAATAAATGAAAACACCTGA

FIGURE 146

MRKHLSSWWLATVCMLLFSHL SAVQTRGIKHRIKWNRKALPSTAQITEAQVAENRPGAFIKQGRK
LDIDFGAEGNRYYEANYWQFPDGIHYNGCSEANVTKEAFVTGCINATQAANQGEFQKPDNKLHQQ
VLWRLVQELCSLKHCEFWLERGAGLRVTMHQPVLLCLLALIWL MVK

Important features of the protein:

Signal peptide:

amino acids 1-26

Transmembrane domain:

amino acids 157-171

N-glycosylation sites.

amino acids 98-102, 110-114

Tyrosine kinase phosphorylation site.

amino acids 76-83

N-myristoylation sites.

amino acids 71-77, 88-94, 93-99, 107-113, 154-160

Amidation site.

amino acids 62-66

FIGURE 147

GCCTTGGCCTCCCAAAGGGCTGGGATTATAGGCGTGACCACCATGTCTGGTCCAGAGTCTCATTT
CCTGATGATTTATAGACTCAAAGAAAACTCATGTTCAGAAGCTCTCTTCTCTTCTGGCCTCCTCT
CTGTCTTCTTTCCCTCTTTCTTCTTATTTTAATTAGTAGCATCTACTCAGAGTCATGCAAGCTGG
AAATCTTTCATTTTGCTTGTCAGTGGGGTAGGTCACTGAGTCTTAGTTTTTATTTTTTGAAATTT
CAACTTTCAGATTCAGGGGGTACATGTGAAGGTTTGTTTTATGAGTATATTGCATTGATGCTGAGG
TTTGGGGT

FIGURE 148

MFRSSLLFWPPLCLLSLFLILISSIYSESKLEIFHFACQWGRSLSLSFYFLKFQLSDSGGTCE
GLFYEYIA

Important features of the protein:

Signal peptide:

amino acids 1-25

N-myristoylation site.

amino acids 62-68

FIGURE 149

GTCTCCGCGTCACAGGAACCTTCAGCACCCACAGGGCGGACAGCGCTCCCCTCTACCTGGAGACTTGAC
TCCCGCGCGCCCCAACCCCTGCTTATCCCTTGACCGTCGAGTGTGAGAGATCCTGCAGCCGCCCAGTCC
CGGCCCCCTCTCCCGCCCCACACCCACCCCTCCTGGCTCTTCCTGTTTTTACTCCTCCTTTTCATTCAIA
ACAAAAGCTACAGCTCCAGGAGCCCAGCGCCGGGCTGTGACCCAAGCCGAGCGTGGAAGAATGGGGTT
CCTCGGGACCGGCACTTGGAATTCTGGTGTAGTGCTCCCGATTCAAGCTTTCCCCAACCTGGAGGAA
GCCAAGACAAATCTCTACATAATAGAGAATTAAGTGCAGAAAGACCTTTGAATGAACAGATTGCTGAA
GCAGAAGAAGACAAGATTAAAAAACATATCCTCCAGAAAACAAGCCAGGTCAGAGCAACTATTCTTT
TGTTGATAACTTGAACCTGCTAAAGGCAATAACAGAAAAGGAAAAAATTGAGAAAGAAAGACAATCTA
TAAGAAGCTCCCCACTTGATAATAAGTTGAATGTGGAAGATGTTGATTCAACCAAGAATCGAAAACCTG
ATCGATGATTATGACTCTACTAAGAGTGGATTGGATCATAAATTTCAAGATGATCCAGATGGTCTTCA
TCAACTAGACGGGACTCCTTTAACCGCTGAAGACATTGTCCATAAAATCGCTGCCAGGATTTATGAAG
AAAATGACAGAGCCGTGTTTGACAAGATTGTTTCTAACTACTTAATCTCGGCCTTATCACAGAAAGC
CAAGCACATACACTGGAAGATGAAGTAGCAGAGGTTTTACAAAAATTAATCTCAAAGGAAGCCAAACA
TTATGAGGAGGATCCCAATAAGCCCACAAGCTGGACTGAGAATCAGGCTGGAAAAATACCAGAGAAAG
TGACTCCAATGGCAGCAATTCAAGATGGTCTTGCTAAGGGAGAAAACGATGAAACAGTATCTAACACA
TTAACCTTGACAAATGGCTTGGAAGGAGAACTAAACCTACAGTGAAGACAACCTTTGAGGAACTCCA
ATATTTCCCAAATTTCTATGCGCTACTGAAAAGTATTGATTGAGAAAAGAAGCAAAAGAGAAAGAAA
CACTGATTACTATCATGAAAACACTGATTGACTTTGTGAAGATGATGGTGAAATATGGAACAATATCT
CCAGAAGAAGGTGTTTCCTACCTTGAAAACCTGGATGAAATGATTGCTCTTCAGACCAAAAAACAAGCT
AGAAAAAATGCTACTGACAATATAAGCAAGCTTTTCCCAGCACCATCAGAGAAGAGTCATGAAGAAA
CAGACAGTACCAAGGAAGAAGCAGCTAAGATGGAAGGAATATGGAAGCTTGAAGGATTCCACAAAA
GATGATAACTCCAACCCAGGAGGAAAGACAGATGAACCCAAAGGAAAAACAGAAGCCTATTTGGAAGC
CATCAGAAAAAATATTGAATGGTTGAAGAAACATGACAAAAAGGGAAATAAAGAAGATTATGACCTTT
CAAAGATGAGAGACTTCATCAATAAACAAGCTGATGCTTATGTGGAGAAAGGCATCCTTGACAAGGAA
GAAGCCGAGGCCATCAAGCGCATTTATAGCAGCCTGTAAAAATGGCAAAAGATCCAGGAGTCTTTCAA
CTGTTTCAGAAAACATAATATAGCTTAAAACACTTCTAATTCTGTGATTAAAAATTTTTTGACCCAAGG
GTTATTAGAAAGTGCTGAATTTACAGTAGTTAACCTTTTACAAGTGGTTAAACATAGCTTTCTTCCC
GTAAAAACTATCTGAAAGTAAAGTTGTATGTAAGCTGAAAAAAAAAAAAAAAAAAAAA

FIGURE 150

MGFLGTGTWILVVLVLP IQAFPKPGGSQDKSLHNRELSAERPLNEQIAEAEEDKIKKTYPPENKPG
QSNYSFVDNLNLLKAITEKEKIEKERQSIRSSPLDNKLNVEDVDSTKNRKLIDDYDSTKSGLDHK
FQDDPDGLHQLDGTPLTAEDIVHKIAARIYEENDRAVFDKIVSKLLNLGLITESQAHTLEDEVAE
VLQKLISKEANNYEEDPNKPTSWTENQAGKIKEKVTMAAIQDGLAKGENDETVSNTLTLTNGLE
RRTKTYSEDNFEELQYFPNFYALLKSIDSEKEAKEKETLITIMKTLIDFVKMMVKYGTISPEEGV
SYLENLDEMIALQTKNKLEKNATDNISKLFAPSEKSHEETDSTKEEAAKMEKEYGSLKDSTKDD
NSNPGGKTDEPKGKTEAYLEAIRKNIEWLKKHDKKGNKEDYDL SKMRDFINKQADAYVEKGILDK
EEAEAIKRIYSSL

N-glycosylation sites:

amino acids 68-71, 346-349, 350-353

Casein kinase II phosphorylation site:

amino acids 70-73, 82-85, 97-100, 125-128, 147-150, 188-191, 217-
220, 265-268, 289-292, 305-308, 320-323, 326-329, 362-365, 368-
341, 369-372, 382-385, 386-389, 387-390

N-myristoylation sites:

amino acids 143-148, 239-244

FIGURE 151

CGGCTCGAGGCTCCCGCCAGGAGAAAGGAACATTCTGAGGGGAGTCTACACCCTGTGGAGCTCAA
GATGGTCCTGAGTGGGGCGCTGTGCTTCCGAATGAAGGACTCGGCATTGAAGGTGCTTTATCTGC
ATAATAACCAGCTTCTAGCTGGAGGGCTGCATGCAGGGAAGGTCATTAAAGGTGAAGAGATCAGC
GTGGTCCCCAATCGGTGGCTGGATGCCAGCCTGTCCCCCGTCATCCTGGGTGTCCAGGGTGAAG
CCAGTGCCTGTGATGTGGGTGGGGCAGGAGCCGACTCTAACACTAGAGCCAGTGAACATCATGG
AGCTCTATCTTGGTGCCAAGGAATCCAAGAGCTTACCTTCTACCGGCGGGACATGGGGCTCACC
TCCAGCTTCGAGTCGGCTGCCTACCGGGCTGGTTCCTGTGCACGGTGCCTGAAGCCGATCAGCC
TGTCAGACTCACCCAGCTTCCCGAGAATGGTGGCTGGAATGCCCCCATCACAGACTTCTACTTCC
AGCAGTGTGACTTAGGGCAACGTGCCCCCAGAACTCCCTGGGCAGAGCCAGCTCGGGTGAGGGGT
GAGTGGAGGAGACCCATGGCGGACAATCACTCTCTGCTCTCAGGACCCCCACGTCTGACTTAG
TGGGCACCTGACCACTTTGTCTTCTGGTTCCAGTTTGGATAAATTCTGAGATTTGGAGCTCAGT
CCACGGTCTCCCCACTGGATGGTGTCTACTGCTGTGGAACCTTGTA AAAACCATGTGGGGTAAA
CTGGGAATAACATGAAAAGATTTCTGTGGGGGTGGGGTGGGGGAGTGGTGGGAATCATTCCCTGCT
TAATGGTAACTGACAAGTGTTACCCTGAGCCCCGAGGCCAACCCTCCCTTTAATCCTGCCACTGTCTATA
GGGTGAGTAGCTCTCCACATGAAGTCTGTCACTCACCCTGTGCAGGAGAGGGAGGTGGTCATA
GAGTCAGGGATCTATGGCCCTTGGCCCAGCCCCACCCCTTCCCTTTAATCCTGCCACTGTCTATA
TGCTACCTTTCCCTATCTCTTCCCTCATCATCTTGTTGTGGGCATGAGGAGGTGGTGATGTCAGAA
GAAATGGCTCGAGCTCAGAAGATAAAAGATAAGTAGGGTATGCTGATCCTCTTTTAAAAACCCAA
GATACAATCAAAATCCCAGATGCTGGTCTCTATTCCCATGAAAAAGTGCTCATGACATATTGAGA
AGACCTACTTACAAAGTGGCATATATTGCAATTTATTTTAAATTAAAAGATACCTATTTATATATT
TCTTTATAGAAAAAAGTCTGGAAGAGTTTACTTCAATTGTAGCAATGTGAGGGTGGTGGCAGTAT
AGGTGATTTTTCTTTTAAATTCTGTAAATTTATCTGTATTTCTTAATTTTTCTACAATGAAGATGA
ATTCTTGTATAAAAAATAAGAAAAGAAATTAATCTTGAGGTAAGCAGAGCAGACATCATCTCTGA
TTGTCCTCAGCCTCCACTTCCCCAGAGTAAATTCAAATTGAATCGAGCTCTGCTGCTCTGGTTGG
TTGTAGTAGTGATCAGGAAACAGATCTCAGCAAAGCCACTGAGGAGGAGGCTGTGCTGAGTTTGT
GTGGCTGGAATCTCTGGGTAAAGGAACCTAAAGAACA AAAATCATCTGGTAATTCTTTCCCTAGAAG
GATCACAGCCCCCTGGGATTCCAAGGCATTGGATCCAGTCTCTAAGAAGGCTGCTGTACTGGTTGA
ATTGTGTCCCCCTCAAATTCACATCCTTCTTGGAATCTCAGTCTGTGAGTTTATTTGGAGATAAG
GTCTCTGCAGATGTAGTTAGTTAAGACAAGGTGATGCTGGATGAAGGTAGACCTAAATTCATAT
GACTGGTTTTCTTGTATGAAAAGGAGAGGACACAGAGACAGAGGAGACGCGGGGAAGACTATGTA
AAGATGAAGGCAGAGATCGGAGTTTTGCAGCCACAAGCTAAGAAACACCAAGGATTGTGGCAACC
ATCAGAAGCTTGGAAGAGGCAAAGAAGAATTCTTCCCTAGAGGCTTTAGAGGGATAACGGCTCTG
CTGAAACCTTAATCTCAGACTTCCAGCCTCCTGAACGAAGAAAGAATAAATTTCCGCTGTTTTAA
GCCACCAAGGATAATTGGTTACAGCAGCTCTAGGAACTAATAACAGCTGCTAAAATGATCCCTGT
CTCCTCGTGTTTACATTTCTGTGTGTGTCCCCCTCCACAATGTACCAAAGTTGTCTTTGTGACCAA
TAGAATATGGCAGAAGTGATGGCATGCCACTTCCAAGATTAGGTTATAAAAGACACTGCAGCTTC
TACTTGAGCCCTCTCTCTCTGCCACCCACCGCCCCCAATCTATCTTGGCTCACTCGCTCTGGGGG
AAGCTAGCTGCCATGCTATGAGCAGGCCATATAAAGAGACTTACGTGGTAAAAAATGAAGTCTCCT
GCCCACAGCCACATTAGTGAACCTAGAAGCAGAGACTCTGTGAGATAATCGATGTTTGTGTTTT
AAGTTGCTCAGTTTTGGTCTAACTTGTTATGCAGCAATAGATAAATAATATGCAGAGAAAGAG

FIGURE 152

MVLSGALCFRMKDSALKVLYLHNNQLLAGGLHAGKVIKGEESISVVPNRWLDASLSPVILGVQGG
QCLSCGVGQEP TLTLEPVNIMELYLGAKESKSFTFYRRDMGLTSSFESAAYPGWFLCTVPEADQP
VRLTQLPENGGWNAPITDFYFQQCD

N-myristoylation sites.

amino acids 29-34, 30-35, 60-65, 63-68, 73-78, 91-96, 106-111

Interleukin-1 signature.

amino acids 111-131

Interleukin-1 proteins.

amino acids 8-29, 83-120, 95-134, 64-103

FIGURE 153

CTTCAGAACAGGTTCTCCTTCCCCAGTCACCAGTTGCTCGAGTTAGAATTGTCTGCAATGGCCGC
CCTGCAGAAATCTGTGAGCTCTTTCCTTATGGGGACCCTGGCCACCAGCTGCCTCCTTCTCTTGG
CCCTCTTGGTACAGGGAGGAGCAGCTGCGCCCATCAGCTCCCACTGCAGGCTTGACAAGTCCAAC
TTCCAGCAGCCCTATATCACCAACCGCACCTTCATGCTGGCTAAGGAGGCTAGCTTGGCTGATAA
CAACACAGACGTTCTGCTCATTGGGGAGAACTGTTCCACGGAGTCAGTATGAGTGAGCGCTGCT
ATCTGATGAAGCAGGTGCTGAACTTCACCCTTGAAGAAGTGCTGTTCCCTCAATCTGATAGGTTT
CAGCCTTATATGCAGGAGGTGGTGCCCTTCCTGGCCAGGCTCAGCAACAGGCTAAGCACATGTCA
TATTGAAGGTGATGACCTGCATATCCAGAGGAATGTGCAAAAGCTGAAGGACACAGTGAAAAAGC
TTGGAGAGAGTGGAGAGATCAAAGCAATTGGAGAACTGGATTGCTGTTTATGTCTCTGAGAAAT
GCCTGCATTTGACCAGAGCAAAGCTGAAAAATGAATAACTAACCCCTTTCCTGCTAGAAATAA
CAATTAGATGCCCCAAAGCGATTTTTTTTAAACCAAAGGAAGATGGGAAGCCAACTCCATCATG
ATGGGTGGATTCCAAATGAACCCCTGCGTTAGTTACAAAGGAAACCAATGCCACTTTTGTATATA
AGACCAGAAGGTAGACTTTCTAAGCATAGATATTTATTGATAACATTCATTGTAAGTGGTGTTT
TATACACAGAAAACAATTTATTTTTTAAATAATTGTCTTTTCCATAAAAAAGATTACTTTCCAT
TCCTTTAGGGGAAAAAACCCCTAAATAGCTTCATGTTTCCATAATCAGTACTTTATATTTATAAA
TGTATTTATTATTATTATAAGACTGCATTTTATTTATATCATTTTATTAATATGGATTTATTTAT
AGAAACATCATTCGATATTGCTACTTGAGTGTAAGGCTAATATTGATATTTATGACAATAATTAT
AGAGCTATAACATGTTTATTTGACCTCAATAAACACTTGGATATCCC

FIGURE 154

MAALQKSVSSFLMGTLATSCLLLLALLVQGGAAAPISSHCRLDKSNFQQPYITNRTFMLAKEASL
ADNNTDVRLIGEKLFHGVSMSERCYLMKQVLNFTLEEVLFQSDRFQPYMQEVVPFLARLSNRLS
TCHIEGDDLHIQRNVQKLKDTVKKLGESGEIKAIGELDLLFMSLRNACI

Important features of the protein:

Signal peptide:

amino acids 1-33

N-glycosylation sites.

amino acids 54-58, 68-72, 97-101

N-myristoylation sites.

amino acids 14-20, 82-88

Prokaryotic membrane lipoprotein lipid attachment site.

amino acids 10-21

FIGURE 155

GGCTTGCTGAAAATAAAATCAGGACTCCTAACCTGCTCCAGTCAGCCTGCTTCCACGAGGCCTGT
CAGTCAGTGCCCGACTTGTGACTGAGTGTGCAGTGCCCGAGCATGTACCAGGTTCAGTGCAGAGGGC
TGCCTGAGGGCTGTGCTGAGAGGGAGAGGAGCAGAGATGCTGCTGAGGGTGGAGGGAGGCCAAGC
TGCCAGGTTTGGGGCTGGGGGCCAAGTGGAGTGAGAACTGGGATCCCAGGGGGAGGGTGCAGAT
GAGGGAGCGACCCAGATTAGGTGAGGACAGTTCTCTCATTAGCCTTTTCCTACAGGTGGTTGCAT
TCTTGGAATGGTCATGGGAACCCACACCTACAGCCACTGGCCCAGCTGCTGCCCCAGCAAAGGG
CAGGACACCTCTGAGGAGCTGCTGAGGTGGAGCACTGTGCCTGTGCCTCCCCTAGAGCCTGCTAG
GCCCCAACCGCCACCCAGAGTCCTGTAGGGCCAGTGAAGATGGACCCCTCAACAGCAGGGCCATCT
CCCCCTGGAGATATGAGTTGGACAGAGACTTGAACCGGCTCCCCCAGGACCTGTACCACGCCCCGT
TGCCTGTGCCCCGCACTGCGTCAGCCTACAGACAGGCTCCACATGGACCCCCGGGGCAACTCGGA
GCTGCTCTACCACAACCAGACTGTCTTCTACAGGCGGCCATGCCATGGCGAGAAGGGCACCCACA
AGGGCTACTGCCTGGAGCGCAGGCTGTACCGTGTTTCCTTAGCTTGTGTGTGTGTGCGGCCCCGT
GTGATGGGCTAGCCGGACCTGCTGAGGCTGGTCCCTTTTGGGAAACCTGGAGCCAGGTGTACA
ACCACTTGCCATGAAGGGCCAGGATGCCAGATGCTTGGCCCCCTGTGAAGTGCTGTCTGGAGCAG
CAGGATCCCGGGACAGGATGGGGGGCTTTGGGGAAAACCTGCACTTCTGCACATTTTGAAAAGAG
CAGCTGCTGCTTAGGGCCGCCGGAAGCTGGTGTCTGTGTCATTTTCTCTCAGGAAAGGTTTTCAA
GTTCTGCCATTTCTGGAGGCCACCACTCCTGTCTCTTCTCTTTTCCCATCCCCTGCTACCCCTG
GCCCAGCACAGGCACTTTCTAGATATTTCCCCCTTGCTGGAGAAGAAAGAGCCCCCTGGTTTTATT
TGTTTGTTTACTCATCACTCAGTGAGCATCTACTTTGGGTGCATTCTAGTGTAGTTACTAGTCTT
TTGACATGGATGATTCTGAGGAGGAAGCTGTTATTGAATGTATAGAGATTTATCCAAATAAATAT
CTTTATTTAAAAATGAAAAA

FIGURE 156

MRERPRLGEDSSLISLFLQVVAFLAMVMGTHTYSHWPSCCPSKGQDTSEELLRWSTVPVPPLEPA
RPNRHPESCRASEDGPLNSRAISPWRYELDRDLNRLPQDLYHARCLCPHCVSLQTGSHMDPRGNS
ELLYHNQTVFYRRPCHGEKGTKGYCLERRLYRVSLACVCVRPRVMG

Important features of the protein:

Signal peptide:

amino acids 1-32

N-glycosylation site.

amino acids 136-140

Tyrosine kinase phosphorylation site.

amino acids 127-135

N-myristoylation sites.

amino acids 44-50, 150-156

FIGURE 157

CCGGCGATGTCGCTCGTGCTGCTAAGCCTGGCCGCGCTGTGCAGGAGCGCCGTACCCCGAGAGCC
GACCGTTCAATGTGGCTCTGAAACTGGGCCATCTCCAGAGTGGATGCTACAACATGATCTAATCC
CCGGAGACTTGAGGGACCTCCGAGTAGAACCTGTTACAACACTAGTGTTGCAACAGGGGACTATTCA
ATTTTGATGAATGTAAGCTGGGTACTCCGGGCAGATGCCAGCATCCGCTTGTTGAAGGCCACCAA
GATTTGTGTGACGGGCAAAAGCAACTTCCAGTCCTACAGCTGTGTGAGGTGCAATTACACAGAGG
CCTTCCAGACTCAGACCAGACCCTCTGGTGGTAAATGGACATTTTCCTACATCGGCTTCCCTGTA
GAGCTGAACACAGTCTATTTTCATTGGGGCCCATATATTCCTAATGCAAATATGAATGAAGATGG
CCCTTCCATGTCTGTGAATTTACCTCACCAGGCTGCCTAGACCACATAATGAAATATAAAAAAA
AGTGTGTCAAGGCCGGAAGCCTGTGGGATCCGAACATCACTGCTTGTAAGAAGAATGAGGAGACA
GTAGAAGTGAAC TTCACAACCACTCCCTGGGAAACAGATACATGGCTCTTATCCAACACAGCAC
TATCATCGGGTTTTCTCAGGTGTTTGAGCCACACCAGAAGAAACAAACGCGAGCTTCAGTGGTGA
TTCCAGTGA CTGGGGATAGTGAAGGTGCTACGGTGCAGCTGACTCCATATTTTCCTACTTGTGGC
AGCGACTGCATCCGACATAAAGGAACAGTTGTGCTCTGCCCACAAACAGGCGTCCCTTTCCCTCT
GGATAACAACAAAAGCAAGCCGGGAGGCTGGCTGCCTCTCCTCCTGCTGTCTCTGCTGGTGGCCA
CATGGGTGCTGGTGGCAGGGATCTATCTAATGTGGAGGCACGAAAGGATCAAGAAGACTTCCTTT
TCTACCACCACACTACTGCCCCCATTAAGGTTCTTGTGGTTTACCCATCTGAAATATGTTTCCA
TCACACAATTTGTTACTTCACTGAATTTCTTCAAACCATTCGAGAAGTGAGGTGTCCTTGAAA
AGTGGCAGAAAAAGAAAATAGCAGAGATGGGTCCAGTGCAGTGGCTTGCCACTCAAAGAGGCA
GCAGACAAAGTCGTCTTCCTTCTTTCCAATGACGTCAACAGTGTGTGCGATGGTACCTGTGGCAA
GAGCGAGGGCAGTCCCAGTGAGAACTCTCAAGACCTCTTCCCCCTTGCTTTAACCTTTTCTGCA
GTGATCTAAGAAGCCAGATTCATCTGCACAAATACGTGGTGGTCTACTTTAGAGAGATTGATACA
AAAGACGATTACAATGCTCTCAGTGTCTGCCCCAAGTACCACCTCATGAAGGATGCCACTGCTTT
CTGTGCAGAACTTCTCCATGTCAAGCAGCAGGTGTCAGCAGGAAAAAGATCACAAGCCTGCCACG
ATGGCTGCTGCTCCTTGTAG

FIGURE 158

MSLVLLSLAALCRSAVPREPTVQCGSETGPSPEWMLQHDLPGLRDLRVEPVTTSVATGDYSILMNVS
WVLRADASIRLLKATKICVTGKSNFQSYSCVRCNYTEAFQTQTRPSGGKWTFSYIGFPVELNTVYFIGAHNIP
NANMNEDGPSMSVNFTSPGCGLDHIMKYKKKCVKAGSLWDPNITACKKNEETVEVNETTTPLGNRYMALIQH
STIIGFSQVFEHPHQKQTRASVVIPVTGDSEGATVQLTPYFPTCGSDCIRHKGTVVLCPTGVPFPLDNNK
SKPGGWLPLLLLSLLVATWVLVAGIYLMWRHERIKKTSFSTTTLLPPIKVLVVYPSEICFHHTICYFTEFL
QNHCRSEVILEKWQKKKIAEMGPVQWLATQKKAADKVVFLSNDVNSVCDGTCGKSEGSPSENSQDLFPLA
ENLFCSDLRSQIHLHKYVVVYFREIDTKDDYNALSVC PKYHLMKDATAFCAELLHV KQVVSAGKRSQACHD
GCCSL

Important features of the protein:

Signal peptide:

amino acids 1-14

Transmembrane domain:

amino acids 290-309

N-glycosylation sites.

amino acids 67 - 71, 103 - 107, 156 - 160, 183 - 187, 197 - 201 and 283
- 287

cAMP- and cGMP-dependent protein kinase phosphorylation sites.

amino acids 228 - 232 and 319 - 323

Casein kinase II phosphorylation sites.

amino acids 178 - 182, 402 - 406, 414 - 418 and 453 - 457

N-myristoylation site.

amino acids 116-122

Amidation site.

amino acids 488-452

FIGURE 159

AGCCACCAGCGCAACATGACAGTGAAGACCCTGCATGGCCCAGCCATGGTCAAGTACTTGCTGCT
GTCGATATTGGGGCTTGCCTTTCTGAGTGAGGCGGCAGCTCGGAAAATCCCCAAAGTAGGACATA
CTTTTTTCCAAAAGCCTGAGAGTTGCCCCGCTGTGCCAGGAGGTAGTATGAAGCTTGACATTGGC
ATCATCAATGAAAACCAGCGCGTTTCCATGTCACGTAACATCGAGAGCCGCTCCACCTCCCCCTG
GAATTACACTGTCACTTGGGACCCCAACCGGTACCCCTCGGAAGTTGTACAGGCCCAGTGTAGGA
ACTTGGGCTGCATCAATGCTCAAGGAAAGGAAGACATCTCCATGAATTCCGTTCCCATCCAGCAA
GAGACCCTGGTCGTCCGGAGGAAGCACCAAGGCTGCTCTGTTTCTTTCCAGTTGGAGAAGGTGCT
GGTGACTGTTGGCTGCACCTGCGTCACCCCTGTCATCCACCATGTGCAGTAAGAGGTGCATATCC
ACTCAGCTGAAGAAG

FIGURE 16o

MTVKTLHGPMVKYLLLSILGLAFLSEAAARKIPKVGHTFFQKPESCPPVPGGSMKLDIGIINEN
QRVMSRNIERSRSTSPWNYTWTWDPNRYPSEVVQAQCRNLGCINAQGKEDISMNSVPIQQETLVV
RRKHQGC SVSFQLEKVLVTVGCTCVTPVIHHVQ

Signal sequence:

amino acids 1-30

N-glycosylation site.

amino acids 83-87

N-myristoylation sites.

amino acids 106-111, 136-141

FIGURE 161

ACACTGGCCAAACAAAAACGAAAGCACTCCGTGCTGGAAGTAGGAGGAGAGTCAGGACTCCCAGG
 ACAGAGAGTGCACAAACTACCCAGCACAGCCCCCTCCGCCCCCTCTGGAGGCTGAAGAGGGATT
 CAGCCCCCTGCCACCCACAGACACGGGCTGACTGGGGTGTCTGCCCCCTTGGGGGGGGGCAGCAC
 AGGGCCTCAGGCCTGGGTGCCACCTGGCACCTAGAAGATGCCTGTGCCCTGGTTCTTGCTGTCCT
 TGGCACTGGGCCGAAGCCCAGTGGTCTTTCTCTGGAGAGGCTTGTGGGGCCTCAGGACGCTACC
 CACTGCTCTCCGGGCTCTCCTGCCGCTCTGGGACAGTGACATACTCTGCCTGCCTGGGGACAT
 CGTGCTGCTCCGGGCCCCGTGCTGGCGCTACGCACCTGCAGACAGAGCTGGTGCTGAGGTGCC
 AGAAGGAGACCGACTGTGACCTCTGTCTGCGTGTGGCTGTCCACTTGGCCGTGCATGGGCACTGG
 GAAGAGCCTGAAGATGAGGAAAAGTTTGGAGGAGCAGCTGACTCAGGGGTGGAGGAGCCTAGGAA
 TGCCTCTCTCCAGGCCCAAGTCGTGCTCTCCTTCCAGGCCTACCCTACTGCCCCGTGCGTCTCTGC
 TGGAGGTGCAAGTGCTGCTGCCCTTGTGCASTTTGGTCAGTCTGTGGGCTCTGTGGTATATGAC
 TGCTTCGAGGCTGCCCTAGGGAGTGAGGTACGAATCTGGTCTTATACTCAGCCCAGGTACGAGAA
 GGAACCTCAACCACACACAGCAGCTGCCTGCCCTGCCCTGGCTCAACGTGTCAGCAGATGGTGACA
 ACGTGCTCTGGTTCTGAATGTCTCTGAGGAGCAGCACTTCGGCCTCTCCCTGTACTGGAATCAG
 GTCCAGGGCCCCCAAAACCCCGGTGGCACAAAACCTGACTGGACCGCAGATCATTACCTTGAA
 CCACACAGACCTGGTTCCCTGCCTCTGTATTACAGGTGTGGCCTCTGGAACCTGACTCCGTTAGGA
 CGAACATCTGCCCCCTCAGGGAGGACCCCCGCGCACACCAGAACCTCTGGCAAGCCGCCCGACTG
 CGACTGCTGACCTTGACAGCTGGCTGCTGGACGCACCGTGCTCGCTGCCCCGAGAAGCGGCACT
 GTGCTGGCGGGCTCCGGGTGGGGACCCCTGCCAGCCACTGGTCCCACCGCTTCTCTGGGAGAAGC
 TCACTGTGGACAAGGTTCTCGAGTTCCCATTTGCTGAAAGGCCACCCTAACCTCTGTGTTACAGGTG
 AACAGCTCGGAGAAGCTGCAGCTGCAGGAGTGCTTGTGGGCTGACTCCCTGGGGCCTCTCAAAGA
 CGATGTGCTACTGTTGGAGACACGAGGCCCCAGGACAACAGATCCCTCTGTGCCTTGGAAACCCA
 GTGGCTGTACTTCACTACCCAGCAAAGCCTCCACGAGGGCAGCTCGCCTTGGAGAGTACTTACTA
 CAAGACCTGCAGTCAGGCCAGTGTCTGCAGCTATGGGACGATGACTTGGGAGCGCTATGGGCCTG
 CCCCATGGACAAATACATCCACAAGCGCTGGGGCCCTCGTGTGGCTGGCCTGCCTACTCTTTGCCG
 CTGCGCTTTCCCTCATCTCCTTCTCAAAAAGGATCACGCGAAAGGGTGGCTGAGGCTCTTGAA
 CAGGACGTCCGCTCGGGGGCGGGCCGCCAGGGGGCCGCGCGGCTCTGCTCCTCTACTCAGCCGATGA
 CTCGGGTTTCGAGCGCCTGGTGGGCGCCCTGGCGTCGGCCCTGTGCCAGCTGCCGCTGCGCGTGG
 CCGTAGACCTGTGGAGCCGTCTGTGAACCTGAGCGCGCAGGGGCCCGTGGCTTGGTTTCACGCGCAG
 CGGCGCCAGACCTTGACAGGAGGGCGGCGTGGTGGTCTTGCTCTTCTCTCCCGGTGCGGTGGCGCT
 GTGCAGCGAGTGGCTACAGGATGGGGTGTCCGGGCCCCGGGCGCACGGCCCCGCACGACGCCTTCC
 GCGCCTCGCTCAGCTGCGTGCTGCCCGACTTCTTGCAGGGCCGGGCGCCCGGCAGCTACGTGGGG
 GCCTGCTTCGACAGGCTGCTCCACCCGGACGCGGTACCCGCCCTTTTCGCGACCGTGCCCGTCTT
 CACACTGCCCTCCCAACTGCCAGACTTCTTGGGGGCCCTGCAGCAGCCTCGCGCCCCGCGTTCG
 GGCGGCTCCAAGAGAGAGCGGAGCAAGTGTCCCGGGCCCTTCAGCCAGCCCTGGATAGCTACTTC
 CATCCCCGGGGACTCCCGCGCCGGGACGCGGGTGGGACCAGGGGCGGGACCTGGGGCGGGGA
 CGGGACTTAAATAAAGGCAGACGCTGTTTTCTAAAAAAA

FIGURE 162

MPVPWFLLSLALGRSPVVLSELERLVGPQDATHCSPGLSCRLWDS DILCLPGDIVPAPGPFVLAPTHLQTELV
LRCQKETDCDLCLRVAVHLAVHGHWEPEDEEKFGGAADSGVEEPRNASLQAQVVLSFQAYPTARCVLLEV
QVPAALVQFGQSVGSSVYDCFEAALGSEVRIWSYTPRYEKELNHTQQLPALPWLNV SADGDNVHLVLNV
EEQHFGLSLYWNQVQGP KPRWHKNLTGPQIITLNHTDLVPCLCIQVWPLEPDSVRTNICPFREDPRAHQN
LWQAARLRLTLQSWLLDAPCSLPAEAALCWRAPGGDPCQPLVPPLSWENVTVDKVLEFP LLKGHPNLCVQ
VNSSEKLQLQECLWADSLGPLKDDVLLLETRGPQDNRS LCALEPSGCTSLPSKASTRAARLGEYLLQDLQS
GQCLQLWDDDLGALWACPMKYIHKRWALVWLACLLFAAALSLILLKKDHAKGWLRLLLKQDVRS GAAARG
RAALLLYSADD SGFERLVGALASALCQLPLRVAVDLWSRRELSAQGPVAFHAQRRQTLQEGGVVLLFSP
GAVALCSEWLQDGVSGPGAHGPHDAFRASLSCVLPDFLQGRAPGSYVGACFDRLLHPDAVPALFRTVPVFT
LPSQLPDEF LGALQQPRAPRSGR LQERAEQVSRA LQPALDSYFHPPGTPAPGRGVGPGAGPGAGDGT

Signal sequence:

amino acids 1-20

Transmembrane domain.

amino acids 453-475

N-glycosylation sites.

amino acids 118-121, 186-189, 198-201, 211-214, 238-241, 248-251,
334-337, 357-360, 391-394

Glycosaminoglycan attachment site.

amino acids 583-586

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 552-555

N-myristoylation sites.

amino acids 107-112, 152-157, 319-324, 438-443, 516-521, 612-617,
692-697, 696-701, 700-705

FIGURE 163

GGGAGGGCTCTGTGCCAGCCCCGATGAGGACGCTGCTGACCATCTTGACTGTGGGATCCCTGGCT
GCTCACGCCCCCTGAGGACCCCTCGGATCTGCTCCAGCACGTGAAATTCCAGTCCAGCAACTTTGA
AAACATCCTGACGTGGGACAGCGGGCCAGAGGGCACCCAGACACGGTCTACAGCATCGAGTATA
AGACGTACGGAGAGAGGGACTGGGTGGCAAAGAAGGGCTGTCAGCGGATCACCCGGAAGTCCTGC
AACCTGACGGTGGAGACGGGCAACCTCACGGAGCTCTACTATGCCAGGGTCACCGCT
GTCAGTGCAGGGAGGCCGGTCAGCCACCAAGATGACTGACAGGTTGAGCTCTCTGCAGCACACTAC
CCTCAAGCCACCTGATGTGACCTGTATCTCCAAAGTGAGATCGATTGAGATGATTGTTTCATCCTA
CCCCACGCCAATCCGTGCAGGCGATGGCCACCGGCTAACCCCTGGAAGACATCTTCCATGACCTG
TTCTACCACTTAGAGCTCCAGGTCAACCGCACCTACCAAATGCACCTTGAGAGGGAAGCAGAGAGA
ATATGAGTTCTTCGGCCTGACCCCTGACACAGAGTTCCTTGGCACCATCATGATTTGCGTTCCCA
CCTGGGCCAAGGAGAGTGCCCCCTACATGTGCCGAGTGAAGACACTGCCAGACCGGACATGGACC
TACTCCTTCTCCGGAGCCTTCTGTCTCCATGGGCTTCTCTCGTCCGAGTACTCTGCTACCTGAG
CTACAGATATGTCACCAAGCCGCTGCACCTCCCAACTCCCTGAACGTCCAGCGAGTCCCTGACTT
TCCAGCCGCTGCGCTTCATCCAGGAGCACGTCTGATCCCTGTCTTTGACCTCAGCGGCCCCAGC
AGTCTGGCCCAGCCTGTCCAGTACTCCAGATCAGGGTGTCTGGACCCAGGGAGCCCGCAGGAGC
TCCACAGCGGCATAGCCTGTCCGAGATCACCTACTTAGGGCAGCCAGACATCTCCATCCTCCAGC
CCTCCAACGTGCCACCTCCCCAGATCCTCTCCCCACTGTCTATGCCCCAAACGCTGCCCCCTGAG
GTCGGGCCCCCATCTATGCACCTCAGGTGACCCCCGAAGCTCAATTCCCATTCTACGCCCCACA
GGCCATCTCTAAGGTCCAGCCTTCTCTATGCCCCCTCAAGCCACTCCGGACAGCTGGCCTCCCT
CCTATGGGGTATGCATGGAAGTTCTGGCAAAGACTCCCCCACTGGGACACTTCTAGTCTTAAA
CACCTTAGGCCCTAAAGGTGAGCTTCAGAAAGAGCCACCAGCTGGAAGCTGCATGTTAGGTGGCCT
TTCTCTGCAGGAGGTGACCTCCTTGGCTATGGAGGAATCCCAAGAAGCAAAATCATTTGCCACAGC
CCCTGGGGATTTGCACAGACAGAACATCTGACCCAAATGTGCTACACAGTGGGGAGGAAGGGACA
CCACAGTACCTAAAGGGCCAGCTCCCCCTCTCTCTCAGTCCAGATCGAGGGCCACCCCATGTC
CCTCCCTTTGCAACCTCCTTCCGGTCCATGTTCCCCCTCGGACCAAGGTCCAAGTCCCTGGGGCC
TGCTGGAGTCCCTTGTGTGTCCCAAGGATGAAGCCAAGAGCCCAGCCCCCTGAGACCTCAGACCTG
GAGCAGCCACAGAACTGGATTCTCTTTTCAGAGGCCTGGCCCTGACTGTGCAGTGGGAGTCTTG
AGGGGAATGGGAAAGGCTTGGTGCTTCTCCCTGTCCCTACCCAGTGTACATCCTTGGCTGTCA
ATCCCATGCCTGCCATGCCACACACTCTGCGATCTGGCCTCAGACGGGTGCCCTTGAGAGAAGC
AGAGGGAGTGGCATGCAGGGCCCCCTGCCATGGGTGCGCTCCTCACCGGAACAAAGCAGCATGATA
AGGACTGCAGCGGGGAGCTCTGGGGAGCAGCTTGTGTAGACAAGCGCGTGCTCGCTGAGCCCTG
CAAGGCAGAAATGACAGTGAAGGAGGAAATGCAGGGAACTCCCGAGGTCCAGAGCCCCACCTC
CTAACACCATGGATTCAAAGTGCTCAGGGAATTTGCCTCTCCTTGCCCCATTCCTGGCCAGTTTC
ACAATCTAGCTCGACAGAGCATGAGGCCCCCTGCCTCTTCTGTCAATTGTTCAAAGGTGGGAAGAGA
GCCTGGAAAAGAACCAGGCCTGGAAAAGAACCAGAAGGAGGCTGGGCAGAACCAGAACAACCTGC
ACTTCTGCCAAGGCCAGGGCCAGCAGGACGGCAGGACTCTAGGGAGGGGTGTGGCCTGCAGCTCA
TTCCAGCCAGGGCAACTGCCTGACGTTGCACGATTTAGCTTCATTCTCTGATAGAACAAGC
GAAATGCAGGTCCACCAGGGAGGGAGACACACAAGCCTTTTCTGCAGGCAGGAGTTTCAGACCCCT
ATCCTGAGAAATGGGGTTTGAAAGGAAGGTGAGGGCTGTGGCCCCCTGGACGGGTACAATAACACAC
TGTACTGATGTCACAACTTTGCAAGCTCTGCCTTGGGTTAGCCCATCTGGGCTCAAATTCAGC
CTCACCACTCACAAGCTGTGTGACTTCAAACAAATGAAATCAGTGCCAGAACCTCGGTTTCTCTC
ATCTGTAATGTGGGGATCATAACACCTACCTCATGGAGTTGTGGTGAAGATGAAATGAAGTCATG
TCTTTAAAGTGCTTAATAGTGCTTGGTACATGGGCAGTGCCCAATAAACGGTAGCTATTTAAAAA
AAAAAAA

FIGURE 164

MRTLLTILTVGSLAAHAPEDPSDLLQHVKFQSSNFENILTWDSGPEGTPDTVYSIEYKTYGERDW
VAKKGCQRITRKSCNLTVETGNLTelyyARVTAVSAGGRSATKMTDRFSSLQHTTLKPPDVTCIS
KVRSIQMIVHPTPTPIRAGDGHRLTLEDIFHDLFYHLELQVNRTYQMHLGGKQREYEFFGLTPDT
EFLGTIMICVPTWAKESAPYMCrvKTLpDRTWTYSFSGAFLESMGFLVAVLCYLSYRYVTKPPAP
PNSLNVQRVLTFQPLRFIQEHVLIPVFDLSGPSSLAQPvQYSQIRVSGPREPAGAPQRHSLSEIT
YLGQPDISILQPSNVPPPQILSPLSYAPNAAPEVGPPSYAPQVTPEAQFPFYAPQAISKVQPSSY
APQATPDSWPPSYGVCMEGSGKDSPTGTLSSPKHLRPKGQLQKEPPAGSCMLGGLSLQEVTSLAM
EESQEAKSLHQPLGICTDRTSDPNVLHSGEEGTPQYLKGQLPLLSSVQIEGHPMSLPLQPPSGPC
SPSDQGPSPWGLLLESVCPKDEAKSPAPETSDLEQPTELDSLFRGLALTvQWES

Signal sequence.

amino acids 1-17

Transmembrane domain.

amino acids 233-250

N-glycosylation sites.

amino acids 80-83, 87-90, 172-175

N-myristoylation sites.

amino acids 11-16, 47-52, 102-107, 531-536, 565-570

FIGURE 165

TGGCCTACTGGAAAAAAAAAAAAAAAAAAAAAGTCACCCGGGCCCCGGGTGGCCACAACATGG
CTGCGGCGCCGGGGCTGCTCTTCTGGCTGTTTCGTGCTGGGGGCGCTCTGGTGGGTCCC GGCCAG
TCGGATCTCAGCCACGGACGGCGTTTCTCGGACCTCAAAGTGTGCGGGGACGAAGAGTGCAGCAT
GTTAATGTACCGTGGGAAAGCTCTTGAAGACTTCACGGGGCCCTGATTGTCGTTTTGTGAATTTTA
AAAAAGGTGACGATGTATATGTCTACTACAACTGGCAGGGGGATCCCTTGAAC TTTGGGCTGGA
AGTGTTGAACACAGTTTTGGATATTTTCCAAAAGATTTGATCAAGGTACTTCATAAATACACGGA
AGAAGAGCTACATATTCCAGCAGATGAGACAGACTTTGTCTGCTTTGAAGGAGGAAGAGATGATT
TTAATAGTTATAATGTAGAAGAGCTTTTAGGATCTTTGGA ACTGGAGGACTCTGTACCTGAAGAG
TCGAAGAAAGCTGAAGAAGTTTCTCAGCACAGAGAGAAATCTCCTGAGGAGTCTCGGGGGCGTGA
ACTTGACCCTGTGCCTGAGCCCAGGCATTCAGAGCTGATT CAGAGGATGGAGAAGGTGCTTTCT
CAGAGAGCACCGAGGGGGCTGCAGGGACAGCCCTCAGCTCAGGAGAGCCACCCTCACACCAGCGGT
CCTGCGGCTAACGCTCAGGGAGTGCAGTCTTCGTTGGACACTTTTGAAGAAATTCTGCACGATAA
ATTGAAAGTGCCGGGAAGCGAAAGCAGAACTGGCAATAGTTCTCCTGCCTCGGTGGAGCGGGAGA
AGACAGATGCTTACAAAGTCCTGAAAACAGAAATGAGTCAGAGAGGAAGTGGACAGTGC GTTATT
CATTACAGCAAAGGATTTTCGTTGGCATCAAAATCTAAGTTTGTTTTACAAAGATTGTTTTTAGTA
CTAAGCTGCCTTGGCAGTTTGCATTTTTGAGCCAAACAAAAATATATTATTTTCCCTTCTAAGTA
AAAAAAAAAAAAAAAAAAAAA

FIGURE 166

MAAAPGLLFWLFLVGLALWWVPGQSDLSHGRRFSDLKVCGDEEC SMLMYRGKALEDFTGPDCRFVN
FKKGDDVYVYYKLAGGSLELWAGSVEHSFGYFPKDLIKVLHKYTEELHIPADETDFVCFEGGRD
DENSYNVEELLGSLELED SVPEESKKAEEVSQHREKSP EESRGRELDPVPEPEAFRA DSEEDGEGA
FSESTEGLQGQPSAQESH PHTSGPAANAQGVQSSLDTFEEILHDKLKVP GSESRTGNSSPASVER
EKTDAYKVLKTEMSQRGSGQC VIHYSKGFRWHQNL SLFYKDCF

Important features of the protein:

Signal peptide:

amino acids 1-22

N-glycosylation site.

amino acids 294-298

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 30-34

Tyrosine kinase phosphorylation site.

amino acids 67-76

N-myristoylation sites.

amino acids 205-211, 225-231, 277-283

Amidation site.

amino acids 28-32

FIGURE 167

CCAGGACCAGGGCGCACC GGCTCAGCCTCTCACTTGTCAGAGGCCGGGGAAGAGAAGCAAAGCGC
AACGGTGTGGTCCAAGCCGGGGCTTCTGCTTCGCCTCTAGGACATACACGGGACCCCTAACTTC
AGTCCCCCAAACGCGCACCCCTCGAAGTCTTGAAGTCCAGCCCCGCACATCCACGCGCGGCACAGG
CGCGGCAGGCGGCAGGTCCCGGCCGAAGGCGATGCGCGCAGGGGGTTCGGGCAGCTGGGCTCGGGC
GGCGGGAGTAGGGCCCGGCAGGGAGGCAGGGAGGCTGCATATTCAGAGTCGCGGGGCTGCGCCCTG
GGCAGAGGCCCGCCCTCGCTCCACGCAACACCTGCTGCTGCCACCGCGCCGCGATGAGCCGCGTGG
TCTCGCTGCTGCTGGGCGCCGCGCTGCTCTGCGGCCACGGAGCCTTCTGCCGCCGCGTGGTCAGC
GGCCAAAAGGTGTGTTTTGCTGACTTCAAGCATCCCTGCTACAAAATGGCCTACTTCCATGAAGT
GTCCAGCCGAGTGAGCTTTCAGGAGGCACGCCTGGCTTGTGAGAGTGAGGGAGGAGTCTCTCTCA
GCCTTGAGAATGAAGCAGAACAGAAAGTTAATAGAGAGCATGTTGCAAAACCTGACAAAACCCGGG
ACAGGGATTTCTGATGGTGATTTCTGGATAGGGCTTTGGAGGAATGGAGATGGGCAAACATCTGG
TGCCTGCCCAGATCTCTACCAGTGGTCTGATGGAAGCAATTCAGTACCGAAACTGGTACACAG
ATGAACCTTCCTGCGGAAGTGAAGAGTGTGTTGTGATGTATCACCACCAACTGCCAATCCTGGC
CTTGGGGGTCCCTACCTTTACCAGTGGAAATGATGACAGGTGTAACATGAAGCACAATTATATTTG
CAAGTATGAACCAGAGATTAATCCAACAGCCCCTGTAGAAAAGCCTTATCTTACAAATCAACCAG
GAGACACCCATCAGAATGTGGTTGTTACTGAAGCAGGTATAATTCCCAATCTAATTTATGTTGTT
ATACCAACAATACCCCTGCTCTTACTGATACTGGTTGCTTTTGGAACTGTTGTTTCCAGATGCT
GCATAAAAGTAAAGGAAGAACAAAACACTAGTCCAAACCAGTCTACACTGTGGATTTCAAAGAGTA
CCAGAAAAGAAAGTGGCATGGAAGTATAATAACTCATTGACTTGGTTCCAGAATTTTGTAATCT
GGATCTGTATAAGGAATGGCATCAGAACAAATAGCTTGGAAATGGCTTGAAATCACAAGGATCTGC
AAGATGAACTGTAAGCTCCCCCTTGAGGCAATATTAAGTAATTTTTATATGTCTATTATTTCA
TTTAAAGAATATGCTGTGCTAATAATGGAGTGAGACATGCTTATTTTGCTAAAGGATGCACCCAA
ACTTCAAACCTCAAGCAAATGAAATGGACAATGCAGATAAAGTTGTTATCAACACGTCGGGAGTA
TGTGTGTTAGAAGCAATTCCTTTTATTTCTTTACCTTTCATAAGTTGTTATCTAGTCAATGTAA
TGTATATTGTATTGAAATTTACAGTGTGCAAAAGTATTTTACCTTTGCATAAGTGTGTTGATAAAA
ATGAAGTGTCTAATATTTATTTTATGGCATCTCATTTTTCAATACATGCTCTTTTGATTAAAG
AACTTATTACTGTTGTCAACTGAATTCACACACACACAAATATAGTACCATAGAAAAAGTTTGT
TTTCTCGAAATAATTCATCTTTTACGCTTCTCTGCTTTTGGTCAATGTCTAGGAAATCTCTTCAGA
AATAAGAAGCTATTTCAATTAAGTGTGATATAAACCTCCTCAAACATTTTACTTAGAGGCAAGGAT
TGTCTAATTTCAATTGTGCAAGACATGTGCCTTATAATTATTTTTAGCTTAAAATTAAACAGATT
TTGTAATAATGTAACCTTTGTTAATAGGTGCATAAACACTAATGCAGTCAATTTGAACAAAAGAAG
TGACATACACAATATAAATCATATGTCTTCACACGTTGCCTATATAATGAGAAGCAGCTCTCTGA
GGGTTCTGAAATCAATGTGGTCCCTCTCTTGCCCACTAAACAAAGATGGTTGTTTCGGGGTTTGGG
ATTGACACTGGAGGCAGATAGTTGCAAAGTTAGTCTAAGGTTTCCCTAGCTGTATTTAGCCTCTG
ACTATATTAGTATACAAAGAGGTCATGTGGTTGAGACCAGGTGAATAGTCACTATCAGTGTGGAG
ACAAGCACAGCACACAGACATTTTAGGAAGGAAAGGAACTACGAAATCGTGTGAAAATGGGTTGG
AACCCATCAGTGATCGCATATTCATTGATGAGGGTTTGCTTGAGATAGAAAATGGTGGCTCCTTT
CTGTCCTTATCTCCTAGTTTCTTCAATGCTTACGCCTTGTCTTCTCAAGAGAAAGTTGTAAGTCT
CTGGTCTTCATATGTCCTGTGCTCCTTTTAACCAAATAAAGAGTTCTTGTCTTCTGGGGGAAAAA
AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 168

MSRVVSLLLGAALLCGHGAFRRVVSQGKVCFADFKHPCYKMAYFHELSSRVSFQEARLACESE
GGVLLSLENEAEQKLIESMLQNLTKPGTGISDGDFWIGLWRNGDGQTSGACPDLYQWSDGSNSQ
YRNWYTDEPSCGSEKCVVMYHQPTANPGLGGPYLYQWNDDRCNMKHNYICKYEPEINPTAPVEK
PYLTNQPGDTHQNVVTEAGIIPNLIYVVIPTIPLLLLILVAFGTCCFQMLHKSCKGRTKTSPNQ
STLWISKSTRKESGMEV

Important features of the protein:

Signal peptide:

amino acids 1-21

Transmembrane domain:

amino acids 214-235

N-glycosylation sites.

amino acids 86-89, 255-258

cAMP- and cGMP-dependent protein kinase phosphorylation site.

amino acids 266-269

N-myristoylation sites.

amino acids 27-32, 66-71, 91-96, 93-98, 102-107, 109-114, 140-145, 212-217

**SECRETED AND TRANSMEMBRANE
POLYPEPTIDES AND NUCLEIC ACIDS
ENCODING THE SAME**

**CROSS REFERENCE TO RELATED
APPLICATIONS**

[0001] This is a continuation application claiming priority under 35 USC §120 to U.S. Ser. No. 10/006867 Filed Dec. 6, 2001, and which claims priority under 35 USC §119 to U.S. provisional serial Nos. 60/063,435 Filed Oct. 29, 1997; 60/064,215 Filed Oct. 29, 1997; 60/082,797 Filed Apr. 22, 1998; 60/083,495 Filed Apr. 29, 1998; 60/085,579 Filed May 15, 1998; 60/087,759 Filed Jun. 2, 1998; 60/088,021 Filed Jun. 4, 1998; 60/088,029 Filed Jun. 4, 1998; 60/088,030 Filed Jun. 4, 1998; 60/088,734 Filed Jun. 10, 1998; 60/088,740 Filed Jun. 10, 1998; 60/088,811 Filed Jun. 10, 1998; 60/088,824 Filed Jun. 10, 1998; 60/088,825 Filed Jun. 10, 1998; 60/088,863 Filed Jun. 11, 1998; 60/089,102 Filed Jun. 12, 1998; 60/089,514 Filed Jun. 16, 1998; 60/089,653 Filed Jun. 17, 1998; 60/089,952 Filed Jun. 19, 1998; 60/090,246 Filed Jun. 22, 1998; 60/090,444 Filed Jun. 24, 1998; 60/090,688 Filed Jun. 25, 1998; 60/090,696 Filed Jun. 25, 1998; 60/090,862 Filed Jun. 26, 1998; 60/091,628 Filed Jul. 2, 1998; 60/096,012 Filed Aug. 10, 1998; 60/096,757 Filed Aug. 17, 1998; 60/096,949 Filed Aug. 18, 1998; 60/096,959 Filed Aug. 18, 1998; 60/097,954 Filed Aug. 26, 1998; 60/097,971 Filed Aug. 26, 1998; 60/097,979 Filed Aug. 26, 1998; 60/098,749 Filed Sep. 10, 1998; 60/099,741 Filed Sep. 10, 1998; 60/099,763 Filed Sep. 10, 1998; 60/099,792 Filed Sep. 10, 1998; 60/099,812 Filed Sep. 10, 1998; 60/099,815 Filed Sep. 10, 1998; 60/100,627 Filed Sep. 16, 1998; 60/100,662 Filed Sep. 16, 1998; 60/100,683 Filed Sep. 17, 1998; 60/100,684 Filed Sep. 17, 1998; 60/100,930 Filed Sep. 17, 1998; 60/101,279 Filed Sep. 22, 1998; 60/101,475 Filed Sep. 23, 1998; 60/101,738 Filed Sep. 24, 1998; 60/101,743 Filed Sep. 24, 1998; 60/101,916 Filed Sep. 24, 1998; 60/102,570 Filed Sep. 30, 1998; 60/103,449 Filed Oct. 6, 1998; 60/103,678 Filed Oct. 8, 1998; 60/103,679 Filed Oct. 8, 1998; 60/103,711 Filed Oct. 8, 1998; 60/105,000 Filed Oct. 20, 1998; 60/105,002 Filed Oct. 20, 1998; 60/105,881 Filed Oct. 27, 1998; 60/106,030 Filed Oct. 28, 1998; 60/106,464 Filed Oct. 30, 1998; 60/106,856 Filed Nov. 3, 1998; 60/108,807 Filed Nov. 17, 1998; 60/112,419 Filed Dec. 15, 1998; 60/112,422 Filed Dec. 15, 1998; 60/112,853 Filed Dec. 16, 1998; 60/113,011 Filed Dec. 16, 1998; 60/112,854 Filed Dec. 16, 1998; 60/113,300 Filed Dec. 22, 1998; 60/113,408 Filed Dec. 22, 1998; 60/113,430 Filed Dec. 23, 1998; 60/113,621 Filed Dec. 23, 1998; 60/114,223 Filed Dec. 30, 1998; 60/115,614 Filed Jan. 12, 1999; 60/116,527 Filed Jan. 20, 1999; 60/116,843 Filed Jan. 22, 1999; 60/119,285 Filed Feb. 9, 1999; 60/119,287 Filed Feb. 9, 1999; 60/119,525 Filed Feb. 10, 1999; 60/119,549 Filed Feb. 10, 1999; 60/120,014 Filed Feb. 11, 1999; 60/129,122 Filed Apr. 13, 1999; 60/129,674 Filed Apr. 16, 1999; 60/131,291 Filed Apr. 27, 1999; 60/138,387 Filed Jun. 9, 1999; 60/144,791 Filed Jul. 20, 1999; 60/169,495 Filed Dec. 7, 1999; 60/175,481 Filed Jan. 11, 2000; 60/191,007 Filed Mar. 21, 2000; 60/199,397 Filed Apr. 25, 2000 and which claims priority under 35 USC §120 to U.S. patent applications Ser. No. and PCT International Patent Application Serial Nos. 09/380,139 Filed Aug. 25, 1999; 09/311,832 Filed May 14, 1999; 09/380,137 Filed Aug. 25, 1999, now abandoned; 09/380,138 Filed Aug. 25, 1999, now abandoned; 09/380,142 Filed Aug. 25, 1999, now abandoned;

09/397,342 Filed Sep. 15, 1999; 09/403,297 Filed Oct. 18, 1999, now abandoned; 09/423,844 Filed Nov. 12, 1999, now abandoned; 09/644,848 Filed Aug. 22, 2000; 09/665,350 Filed Sep. 18, 2000; 09/664,610 Filed Sep. 18, 2000, now abandoned; 09/709,238 Filed Nov. 8, 2000; 09/747,259 Filed Dec. 20, 2000; 09/816,744 Filed Mar. 22, 2001; 09/854,208 Filed May 10, 2001; 09/854,280 Filed May 10, 2001; 09/870,574 Filed May 30, 2001; 09/874,503 Filed Jun. 10, 2001; 09/908,827 Filed Jul. 18, 2001; 09/869,566 Filed Feb. 19, 2002; PCT/US98/19330 Filed Sep. 16, 1998; PCT/US99/05028 Filed Mar. 8, 1999; PCT/US99/10733 Filed May 14, 1999; PCT/US99/12252 Filed Jun. 2, 1999; PCT/US99/20111 Filed Sep. 1, 1999; PCT/US99/21090 Filed Sep. 15, 1999; PCT/US99/21194 Filed Sep. 15, 1999; PCT/US99/30720 Filed Dec. 22, 1999; PCT/US00/04341 Filed Feb. 18, 2000; PCT/US00/04342 Filed Feb. 18, 2000; PCT/US00/04414 Filed Feb. 22, 2000; PCT/US00/05601 Filed Mar. 1, 2000; PCT/US00/08439 Filed Mar. 30, 2000; PCT/US00/14042 Filed May 22, 2000; PCT/US00/15264 Filed Jun. 2, 2000; PCT/US00/23522 Filed Aug. 23, 2000; PCT/US00/23328 Filed Aug. 24, 2000; PCT/US00/30873 Filed Nov. 10, 2000; PCT/US00/32678 Filed Dec. 1, 2000; PCT/US00/34956 Filed Dec. 20, 2000; PCT/US01/06520 Filed Feb. 28, 2001; PCT/US01/06666 Filed Mar. 1, 2001; PCT/US01/17443 Filed May 30, 2001; PCT/US01/17800 Filed Jun. 01, 2000; PCT/US01/19692 Filed Jun. 20, 2001; PCT/US01/21066 Filed Jun. 29, 2001; PCT/US01/21735 Filed Jul. 9, 2001, the entire disclosures of which are hereby incorporated by reference.

BACKGROUND OF THE INVENTION

[0002] The present invention relates generally to the identification and isolation of novel DNA and to the recombinant production of novel polypeptides.

[0003] Extracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment.

[0004] Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., Proc. Natl. Acad. Sci. 93:7108-7113 (1996); U.S. Pat. No. 5,536,637].

[0005] Membrane-bound proteins and receptors can play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and nerve growth factor receptor.

[0006] Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

[0007] Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

SUMMARY OF INVENTION

[0008] In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO polypeptide.

[0009] In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic

acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0010] In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule comprising the coding sequence of a full-length PRO polypeptide cDNA as disclosed herein, the coding sequence of a PRO polypeptide lacking the signal peptide as disclosed herein, the coding sequence of an extracellular domain of a transmembrane PRO polypeptide, with or without the signal peptide, as disclosed herein or the coding sequence of any other specifically defined fragment of the full-length amino acid sequence as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0011] In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively

at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid-sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein, or (b) the complement of the DNA molecule of (a).

[0012] Another aspect the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence encoding a PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated, or is complementary to such encoding nucleotide sequence, wherein the transmembrane domain(s) of such polypeptide are disclosed herein. Therefore, soluble extracellular domains of the herein described PRO polypeptides are contemplated.

[0013] Another embodiment is directed to fragments of a PRO polypeptide coding sequence, or the complement thereof, that may find use as, for example, hybridization probes, for encoding fragments of a PRO polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO antibody or as antisense oligonucleotide probes. Such nucleic acid fragments are usually at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 160 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700 nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. It is noted that novel fragments of a PRO polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO polypeptide-encoding nucleotide sequences are contemplated herein. Also contemplated are the PRO polypeptide fragments encoded by these nucleotide molecule frag-

ments, preferably those PRO polypeptide fragments that comprise a binding site for an anti-PRO antibody.

[0014] In another embodiment, the invention provides isolated PRO polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

[0015] In a certain aspect, the invention concerns an isolated PRO polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a PRO polypeptide having a full-length amino acid sequence as disclosed herein, an amino acid sequence lacking the signal peptide as disclosed herein, an extracellular domain of a transmembrane protein, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of the full-length amino acid sequence as disclosed herein.

[0016] In a further aspect, the invention concerns an isolated PRO polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to an amino acid sequence encoded by any of the human protein cDNAs deposited with the ATCC as disclosed herein.

[0017] In a specific aspect, the invention provides an isolated PRO polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0018] Another aspect the invention provides an isolated PRO polypeptide which is either transmembrane domain-deleted or transmembrane domain-inactivated. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO polypeptide and recovering the PRO polypeptide from the cell culture.

[0019] In yet another embodiment, the invention concerns agonists and antagonists of a native PRO polypeptide as defined herein. In a particular embodiment, the agonist or antagonist is an anti-PRO antibody or a small molecule.

[0020] In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO polypeptide which comprise contacting the PRO polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO polypeptide. Preferably, the PRO polypeptide is a native PRO polypeptide.

[0021] In a still further embodiment, the invention concerns a composition of matter comprising a PRO polypeptide, or an agonist or antagonist of a PRO polypeptide as herein described, or an anti-PRO antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

[0022] Another embodiment of the present invention is directed to the use of a PRO polypeptide, or an agonist or antagonist thereof as hereinbefore described, or an anti-PRO antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO polypeptide, an agonist or antagonist thereof or an anti-PRO antibody.

[0023] In other embodiments of the present invention, the invention provides vectors comprising DNA encoding any of the herein described polypeptides. Host cell comprising any such vector are also provided. By way of example, the host cells may be CHO cells, *E. coli*, or yeast. A process for producing any of the herein described polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of the desired polypeptide and recovering the desired polypeptide from the cell culture.

[0024] In other embodiments, the invention provides chimeric molecules comprising any of the herein described polypeptides fused to a heterologous polypeptide or amino acid sequence. Example of such chimeric molecules comprise any of the herein described polypeptides fused to an epitope tag sequence or a Fc region of an immunoglobulin.

[0025] In another embodiment, the invention provides an antibody which binds, preferably specifically, to any of the

above or below described polypeptides. Optionally, the antibody is a monoclonal antibody, humanized antibody, antibody fragment or single-chain antibody.

[0026] In yet other embodiments, the invention provides oligonucleotide probes useful for isolating genomic and cDNA nucleotide sequences or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences.

[0027] In yet other embodiments, the present invention is directed to methods of using the PRO polypeptides of the present invention for a variety of uses based upon the functional biological assay data presented in the Examples below.

BRIEF DESCRIPTION OF DRAWINGS

[0028] FIG. 1 shows a nucleotide sequence (SEQ ID NO: 1) of a native sequence PRO180 cDNA, wherein SEQ ID NO: 1 is a clone designated herein as "DNA26843-1389".

[0029] FIG. 2 shows the amino acid sequence (SEQ ID NO: 2) derived from the coding sequence of SEQ ID NO: 1 shown in FIG. 1.

[0030] FIG. 3 shows a nucleotide sequence (SEQ ID NO: 3) of a native sequence PRO218 cDNA, wherein SEQ ID NO: 3 is a clone designated herein as "DNA30867-1335".

[0031] FIG. 4 shows the amino acid sequence (SEQ ID NO: 4) derived from the coding sequence of SEQ ID NO: 3 shown in FIG. 3.

[0032] FIG. 5 shows a nucleotide sequence (SEQ ID NO: 5) of a native sequence PRO263 cDNA, wherein SEQ ID NO: 5 is a clone designated herein as "DNA34431-1177".

[0033] FIG. 6 shows the amino acid sequence (SEQ ID NO: 6) derived from the coding sequence of SEQ ID NO: 5 shown in FIG. 5.

[0034] FIG. 7 shows a nucleotide sequence (SEQ ID NO: 7) of a native sequence PRO295 cDNA, wherein SEQ ID NO: 7 is a clone designated herein as "DNA38268-1188".

[0035] FIG. 8 shows the amino acid sequence (SEQ ID NO: 8) derived from the coding sequence of SEQ ID NO: 7 shown in FIG. 7.

[0036] FIG. 9 shows a nucleotide sequence (SEQ ID NO: 9) of a native sequence PRO874 cDNA, wherein SEQ ID NO: 9 is a clone designated herein as "DNA40621-1440".

[0037] FIG. 10 shows the amino acid sequence (SEQ ID NO: 10) derived from the coding sequence of SEQ ID NO: 9 shown in FIG. 9.

[0038] FIG. 11 shows a nucleotide sequence (SEQ ID NO: 11) of a native sequence PRO300 cDNA, wherein SEQ ID NO: 11 is a clone designated herein as "DNA40625-1189".

[0039] FIG. 12 shows the amino acid sequence (SEQ ID NO: 12) derived from the coding sequence of SEQ ID NO: 11 shown in FIG. 11.

[0040] FIG. 13 shows a nucleotide sequence (SEQ ID NO: 13) of a native sequence PRO1864 cDNA, wherein SEQ ID NO: 13 is a clone designated herein as "DNA45409-2511".

[0041] FIG. 14 shows the amino acid sequence (SEQ ID NO: 14) derived from the coding sequence of SEQ ID NO: 13 shown in FIG. 13.

[0042] FIG. 15 shows a nucleotide sequence (SEQ ID NO: 15) of a native sequence PRO1282 cDNA, wherein SEQ ID NO: 15 is a clone designated herein as "DNA45495-1550".

[0043] FIG. 16 shows the amino acid sequence (SEQ ID NO: 16) derived from the coding sequence of SEQ ID NO: 15 shown in FIG. 15.

[0044] FIG. 17 shows a nucleotide sequence (SEQ ID NO: 17) of a native sequence PRO1063 cDNA, wherein SEQ ID NO: 17 is a clone designated herein as "DNA49820-1427".

[0045] FIG. 18 shows the amino acid sequence (SEQ ID NO: 18) derived from the coding sequence of SEQ ID NO: 17 shown in FIG. 17.

[0046] FIG. 19 shows a nucleotide sequence (SEQ ID NO: 19) of a native sequence PRO1773 cDNA, wherein SEQ ID NO: 19 is a clone designated herein as "DNA56406-1704".

[0047] FIG. 20 shows the amino acid sequence (SEQ ID NO: 20) derived from the coding sequence of SEQ ID NO: 19 shown in FIG. 19.

[0048] FIG. 21 shows a nucleotide sequence (SEQ ID NO: 21) of a native sequence PRO1013 cDNA, wherein SEQ ID NO: 21 is a clone designated herein as "DNA56410-1414".

[0049] FIG. 22 shows the amino acid sequence (SEQ ID NO: 22) derived from the coding sequence of SEQ ID NO: 21 shown in FIG. 21.

[0050] FIG. 23 shows a nucleotide sequence (SEQ ID NO: 23) of a native sequence PRO937 cDNA, wherein SEQ ID NO: 23 is a clone designated herein as "DNA56436-1448".

[0051] FIG. 24 shows the amino acid sequence (SEQ ID NO: 24) derived from the coding sequence of SEQ ID NO: 23 shown in FIG. 23.

[0052] FIG. 25 shows a nucleotide sequence (SEQ ID NO: 25) of a native sequence PRO842 cDNA, wherein SEQ ID NO: 25 is a clone designated herein as "DNA56855-1447".

[0053] FIG. 26 shows the amino acid sequence (SEQ ID NO: 26) derived from the coding sequence of SEQ ID NO: 25 shown in FIG. 25.

[0054] FIG. 27 shows a nucleotide sequence (SEQ ID NO: 27) of a native sequence PRO1180 cDNA, wherein SEQ ID NO: 27 is a clone designated herein as "DNA56860-1510".

[0055] FIG. 28 shows the amino acid sequence (SEQ ID NO: 28) derived from the coding sequence of SEQ ID NO: 27 shown in FIG. 27.

[0056] FIG. 29 shows a nucleotide sequence (SEQ ID NO: 29) of a native sequence PRO831 cDNA, wherein SEQ ID NO: 29 is a clone designated herein as "DNA56862-1343".

[0057] FIG. 30 shows the amino acid sequence (SEQ ID NO: 30) derived from the coding sequence of SEQ ID NO: 29 shown in FIG. 29.

[0058] FIG. 31 shows a nucleotide sequence (SEQ ID NO: 31) of a native sequence PRO1115 cDNA, wherein SEQ ID NO: 31 is a clone designated herein as "DNA56868-1478".

[0059] FIG. 32 shows the amino acid sequence (SEQ ID NO: 32) derived from the coding sequence of SEQ ID NO: 31 shown in FIG. 31.

[0060] FIG. 33 shows a nucleotide sequence (SEQ ID NO: 33) of a native sequence PRO1277 cDNA, wherein SEQ ID NO: 33 is a clone designated herein as "DNA56869-1545".

[0061] FIG. 34 shows the amino acid sequence (SEQ ID NO: 34) derived from the coding sequence of SEQ ID NO: 33 shown in FIG. 33.

[0062] FIG. 35 shows a nucleotide sequence (SEQ ID NO: 35) of a native sequence PRO1074 cDNA, wherein SEQ ID NO: 35 is a clone designated herein as "DNA57704-1452".

[0063] FIG. 36 shows the amino acid sequence (SEQ ID NO: 36) derived from the coding sequence of SEQ ID NO: 35 shown in FIG. 35.

[0064] FIG. 37 shows a nucleotide sequence (SEQ ID NO: 37) of a native sequence PRO1344 cDNA, wherein SEQ ID NO: 37 is a clone designated herein as "DNA58723-1588".

[0065] FIG. 38 shows the amino acid sequence (SEQ ID NO: 38) derived from the coding sequence of SEQ ID NO: 37 shown in FIG. 37.

[0066] FIG. 39 shows a nucleotide sequence (SEQ ID NO: 39) of a native sequence PRO1136 cDNA, wherein SEQ ID NO: 39 is a clone designated herein as "DNA57827-1493".

[0067] FIG. 40 shows the amino acid sequence (SEQ ID NO: 40) derived from the coding sequence of SEQ ID NO: 39 shown in FIG. 39.

[0068] FIG. 41 shows a nucleotide sequence (SEQ ID NO: 41) of a native sequence PRO1109 cDNA, wherein SEQ ID NO: 41 is a clone designated herein as "DNA58737-1473".

[0069] FIG. 42 shows the amino acid sequence (SEQ ID NO: 42) derived from the coding sequence of SEQ ID NO: 41 shown in FIG. 41.

[0070] FIG. 43 shows a nucleotide sequence (SEQ ID NO: 43) of a native sequence PRO1003 cDNA, wherein SEQ ID NO: 43 is a clone designated herein as "DNA58846-1409".

[0071] FIG. 44 shows the amino acid sequence (SEQ ID NO: 44) derived from the coding sequence of SEQ ID NO: 43 shown in FIG. 43.

[0072] FIG. 45 shows a nucleotide sequence (SEQ ID NO: 45) of a native sequence PRO1138 cDNA, wherein SEQ ID NO: 45 is a clone designated herein as "DNA58850-1495".

[0073] FIG. 46 shows the amino acid sequence (SEQ ID NO: 46) derived from the coding sequence of SEQ ID NO: 45 shown in FIG. 45.

[0074] FIG. 47 shows a nucleotide sequence (SEQ ID NO: 47) of a native sequence PRO994 cDNA, wherein SEQ ID NO: 47 is a clone designated herein as "DNA58855-1422".

[0075] FIG. 48 shows the amino acid sequence (SEQ ID NO: 48) derived from the coding sequence of SEQ ID NO: 47 shown in FIG. 47.

[0076] FIG. 49 shows a nucleotide sequence (SEQ ID NO: 49) of a native sequence PRO1069 cDNA, wherein SEQ ID NO: 49 is a clone designated herein as "DNA59211-1450".

[0077] FIG. 50 shows the amino acid sequence (SEQ ID NO: 50) derived from the coding sequence of SEQ ID NO: 49 shown in FIG. 49.

[0078] FIG. 51 shows a nucleotide sequence (SEQ ID NO: 51) of a native sequence PRO1411 cDNA, wherein SEQ ID NO: 51 is a clone designated herein as "DNA59212-1627".

[0079] FIG. 52 shows the amino acid sequence (SEQ ID NO: 52) derived from the coding sequence of SEQ ID NO: 51 shown in FIG. 51.

[0080] FIG. 53 shows a nucleotide sequence (SEQ ID NO: 53) of a native sequence PRO1129 cDNA, wherein SEQ ID NO: 53 is a clone designated herein as "DNA59213-1487".

[0081] FIG. 54 shows the amino acid sequence (SEQ ID NO: 54) derived from the coding sequence of SEQ ID NO: 53 shown in FIG. 53.

[0082] FIG. 55 shows a nucleotide sequence (SEQ ID NO: 55) of a native sequence PRO1027 cDNA, wherein SEQ ID NO: 55 is a clone designated herein as "DNA59605-1418".

[0083] FIG. 56 shows the amino acid sequence (SEQ ID NO: 56) derived from the coding sequence of SEQ ID NO: 55 shown in FIG. 55.

[0084] FIG. 57 shows a nucleotide sequence (SEQ ID NO: 57) of a native sequence PRO1106 cDNA, wherein SEQ ID NO: 57 is a clone designated herein as "DNA59609-1470".

[0085] FIG. 58 shows the amino acid sequence (SEQ ID NO: 58) derived from the coding sequence of SEQ ID NO: 57 shown in FIG. 57.

[0086] FIG. 59 shows a nucleotide sequence (SEQ ID NO: 59) of a native sequence PRO1291 cDNA, wherein SEQ ID NO: 59 is a clone designated herein as "DNA59610-1556".

[0087] FIG. 60 shows the amino acid sequence (SEQ ID NO: 60) derived from the coding sequence of SEQ ID NO: 59 shown in FIG. 59.

[0088] FIG. 61 shows a nucleotide sequence (SEQ ID NO: 61) of a native sequence PRO3573 cDNA, wherein SEQ ID NO: 61 is a clone designated herein as "DNA59837-2545".

[0089] FIG. 62 shows the amino acid sequence (SEQ ID NO: 62) derived from the coding sequence of SEQ ID NO: 61 shown in FIG. 61.

[0090] FIG. 63 shows a nucleotide sequence (SEQ ID NO: 63) of a native sequence PRO3566 cDNA, wherein SEQ ID NO: 63 is a clone designated herein as "DNA59844-2542".

[0091] FIG. 64 shows the amino acid sequence (SEQ ID NO: 64) derived from the coding sequence of SEQ ID NO: 63 shown in FIG. 63.

[0092] FIG. 65 shows a nucleotide sequence (SEQ ID NO: 65) of a native sequence PRO1098 cDNA, wherein SEQ ID NO: 65 is a clone designated herein as "DNA59854-1459".

[0093] FIG. 66 shows the amino acid sequence (SEQ ID NO: 66) derived from the coding sequence of SEQ ID NO: 65 shown in FIG. 65.

[0094] FIG. 67 shows a nucleotide sequence (SEQ ID NO: 67) of a native sequence PRO1158 cDNA, wherein SEQ ID NO: 67 is a clone designated herein as "DNA60625-1507".

[0095] FIG. 68 shows the amino acid sequence (SEQ ID NO: 68) derived from the coding sequence of SEQ ID NO: 67 shown in FIG. 67.

[0096] FIG. 69 shows a nucleotide sequence (SEQ ID NO: 69) of a native sequence PRO1124 cDNA, wherein SEQ ID NO: 69 is a clone designated herein as "DNA60629-1481".

[0097] FIG. 70 shows the amino acid sequence (SEQ ID NO: 70) derived from the coding sequence of SEQ ID NO: 69 shown in FIG. 69.

[0098] FIG. 71 shows a nucleotide sequence (SEQ ID NO: 71) of a native sequence PRO1287 cDNA, wherein SEQ ID NO: 71 is a clone designated herein as "DNA61755-1554".

[0099] FIG. 72 shows the amino acid sequence (SEQ ID NO: 72) derived from the coding sequence of SEQ ID NO: 71 shown in FIG. 71.

[0100] FIG. 73 shows a nucleotide sequence (SEQ ID NO: 73) of a native sequence PRO1335 cDNA, wherein SEQ ID NO: 73 is a clone designated herein as "DNA62812-1594".

[0101] FIG. 74 shows the amino acid sequence (SEQ ID NO: 74) derived from the coding sequence of SEQ ID NO: 73 shown in FIG. 73.

[0102] FIG. 75 shows a nucleotide sequence (SEQ ID NO: 75) of a native sequence PRO1315 cDNA, wherein SEQ ID NO: 75 is a clone designated herein as "DNA62815-1576".

[0103] FIG. 76 shows the amino acid sequence (SEQ ID NO: 76) derived from the coding sequence of SEQ ID NO: 75 shown in FIG. 75.

[0104] FIG. 77 shows a nucleotide sequence (SEQ ID NO: 77) of a native sequence PRO1357 cDNA, wherein SEQ ID NO: 77 is a clone designated herein as "DNA64881-1602".

[0105] FIG. 78 shows the amino acid sequence (SEQ ID NO: 78) derived from the coding sequence of SEQ ID NO: 77 shown in FIG. 77.

[0106] FIG. 79 shows a nucleotide sequence (SEQ ID NO: 79) of a native sequence PRO1356 cDNA, wherein SEQ ID NO: 79 is a clone designated herein as "DNA64886-1601".

[0107] FIG. 80 shows the amino acid sequence (SEQ ID NO: 80) derived from the coding sequence of SEQ ID NO: 79 shown in FIG. 79.

[0108] FIG. 81 shows a nucleotide sequence (SEQ ID NO: 81) of a native sequence PRO1557 cDNA, wherein SEQ ID NO: 81 is a clone designated herein as "DNA64902-1667".

[0109] FIG. 82 shows the amino acid sequence (SEQ ID NO: 82) derived from the coding sequence of SEQ ID NO: 81 shown in FIG. 81.

[0110] FIG. 83 shows a nucleotide sequence (SEQ ID NO: 83) of a native sequence PRO1347 cDNA, wherein SEQ ID NO: 83 is a clone designated herein as "DNA64950-1590".

[0111] FIG. 84 shows the amino acid sequence (SEQ ID NO: 84) derived from the coding sequence of SEQ ID NO: 83 shown in FIG. 83.

[0112] FIG. 85 shows a nucleotide sequence (SEQ ID NO: 85) of a native sequence PRO1302 cDNA, wherein SEQ ID NO: 85 is a clone designated herein as "DNA65403-1565".

[0113] FIG. 86 shows the amino acid sequence (SEQ ID NO: 86) derived from the coding sequence of SEQ ID NO: 85 shown in FIG. 85.

[0114] FIG. 87 shows a nucleotide sequence (SEQ ID NO: 87) of a native sequence PRO1270 cDNA, wherein SEQ ID NO: 87 is a clone designated herein as "DNA66308-1537".

[0115] FIG. 88 shows the amino acid sequence (SEQ ID NO: 88) derived from the coding sequence of SEQ ID NO: 87 shown in FIG. 87.

[0116] FIG. 89 shows a nucleotide sequence (SEQ ID NO: 89) of a native sequence PRO1268 cDNA, wherein SEQ ID NO: 89 is a clone designated herein as "DNA66519-1535".

[0117] FIG. 90 shows the amino acid sequence (SEQ ID NO: 90) derived from the coding sequence of SEQ ID NO: 89 shown in FIG. 89.

[0118] FIG. 91 shows a nucleotide sequence (SEQ ID NO: 91) of a native sequence PRO1327 cDNA, wherein SEQ ID NO: 91 is a clone designated herein as "DNA66521-1583".

[0119] FIG. 92 shows the amino acid sequence (SEQ ID NO: 92) derived from the coding sequence of SEQ ID NO: 91 shown in FIG. 91.

[0120] FIG. 93 shows a nucleotide sequence (SEQ ID NO: 93) of a native sequence PRO1328 cDNA, wherein SEQ ID NO: 93 is a clone designated herein as "DNA66658-1584".

[0121] FIG. 94 shows the amino acid sequence (SEQ ID NO: 94) derived from the coding sequence of SEQ ID NO: 93 shown in FIG. 93.

[0122] FIG. 95 shows a nucleotide sequence (SEQ ID NO: 95) of a native sequence PRO1329 cDNA, wherein SEQ ID NO: 95 is a clone designated herein as "DNA66660-1585".

[0123] FIG. 96 shows the amino acid sequence (SEQ ID NO: 96) derived from the coding sequence of SEQ ID NO: 95 shown in FIG. 95.

[0124] FIG. 97 shows a nucleotide sequence (SEQ ID NO: 97) of a native sequence PRO1340 cDNA, wherein SEQ ID NO: 97 is a clone designated herein as "DNA66663-1598".

[0125] FIG. 98 shows the amino acid sequence (SEQ ID NO: 98) derived from the coding sequence of SEQ ID NO: 97 shown in FIG. 97.

[0126] FIG. 99 shows a nucleotide sequence (SEQ ID NO: 99) of a native sequence PRO1342 cDNA, wherein SEQ ID NO: 99 is a clone designated herein as "DNA66674-1599".

[0127] FIG. 100 shows the amino acid sequence (SEQ ID NO: 100) derived from the coding sequence of SEQ ID NO: 99 shown in FIG. 99.

[0128] FIG. 101 shows a nucleotide sequence (SEQ ID NO: 101) of a native sequence PRO3579 cDNA, wherein SEQ ID NO: 101 is a clone designated herein as "DNA68862-2546".

[0129] FIG. 102 shows the amino acid sequence (SEQ ID NO: 102) derived from the coding sequence of SEQ ID NO: 101 shown in FIG. 101.

[0130] FIG. 103 shows a nucleotide sequence (SEQ ID NO: 103) of a native sequence PRO1472 cDNA, wherein SEQ ID NO: 103 is a clone designated herein as "DNA68866-1644".

[0131] FIG. 104 shows the amino acid sequence (SEQ ID NO: 104) derived from the coding sequence of SEQ ID NO: 103 shown in FIG. 103.

[0132] FIG. 105 shows a nucleotide sequence (SEQ ID NO: 105) of a native sequence PRO1461 cDNA, wherein SEQ ID NO: 105 is a clone designated herein as "DNA68871-1638".

[0133] FIG. 106 shows the amino acid sequence (SEQ ID NO: 106) derived from the coding sequence of SEQ ID NO: 105 shown in FIG. 105.

[0134] FIG. 107 shows a nucleotide sequence (SEQ ID NO: 107) of a native sequence PRO1568 cDNA, wherein SEQ ID NO: 107 is a clone designated herein as "DNA68880-1676".

[0135] FIG. 108 shows the amino acid sequence (SEQ ID NO: 108) derived from the coding sequence of SEQ ID NO: 107 shown in FIG. 107.

[0136] FIG. 109 shows a nucleotide sequence (SEQ ID NO: 109) of a native sequence PRO1753 cDNA, wherein SEQ ID NO: 109 is a clone designated herein as "DNA68883-1691".

[0137] **FIG. 110** shows the amino acid sequence (SEQ ID NO: 110) derived from the coding sequence of SEQ ID NO: 109 shown in **FIG. 109**.

[0138] **FIG. 111** shows a nucleotide sequence (SEQ ID NO: 111) of a native sequence PRO1570 cDNA, wherein SEQ ID NO: 111 is a clone designated herein as "DNA68885-1678".

[0139] **FIG. 112** shows the amino acid sequence (SEQ ID NO: 112) derived from the coding sequence of SEQ ID NO: 111 shown in **FIG. 111**.

[0140] **FIG. 113** shows a nucleotide sequence (SEQ ID NO: 113) of a native sequence PRO1446 cDNA, wherein SEQ ID NO: 113 is a clone designated herein as "DNA71277-1636".

[0141] **FIG. 114** shows the amino acid sequence (SEQ ID NO: 114) derived from the coding sequence of SEQ ID NO: 113 shown in **FIG. 113**.

[0142] **FIG. 115** shows a nucleotide sequence (SEQ ID NO: 115) of a native sequence PRO1565 cDNA, wherein SEQ ID NO: 115 is a clone designated herein as "DNA73727-1673".

[0143] **FIG. 116** shows the amino acid sequence (SEQ ID NO: 116) derived from the coding sequence of SEQ ID NO: 115 shown in **FIG. 115**.

[0144] **FIG. 117** shows a nucleotide sequence (SEQ ID NO: 117) of a native sequence PRO1572 cDNA, wherein SEQ ID NO: 117 is a clone designated herein as "DNA73734-1680".

[0145] **FIG. 118** shows the amino acid sequence (SEQ ID NO: 118) derived from the coding sequence of SEQ ID NO: 117 shown in **FIG. 117**.

[0146] **FIG. 119** shows a nucleotide sequence (SEQ ID NO: 119) of a native sequence PRO1573 cDNA, wherein SEQ ID NO: 119 is a clone designated herein as "DNA73735-1681".

[0147] **FIG. 120** shows the amino acid sequence (SEQ ID NO: 120) derived from the coding sequence of SEQ ID NO: 119 shown in **FIG. 119**.

[0148] **FIG. 121** shows a nucleotide sequence (SEQ ID NO: 121) of a native sequence PRO1550 cDNA, wherein SEQ ID NO: 121 is a clone designated herein as "DNA76393-1664".

[0149] **FIG. 122** shows the amino acid sequence (SEQ ID NO: 122) derived from the coding sequence of SEQ ID NO: 121 shown in **FIG. 121**.

[0150] **FIG. 123** shows a nucleotide sequence (SEQ ID NO: 123) of a native sequence PRO1693 cDNA, wherein SEQ ID NO: 123 is a clone designated herein as "DNA77301-1708".

[0151] **FIG. 124** shows the amino acid sequence (SEQ ID NO: 124) derived from the coding sequence of SEQ ID NO: 123 shown in **FIG. 123**.

[0152] **FIG. 125** shows a nucleotide sequence (SEQ ID NO: 125) of a native sequence PRO1566 cDNA, wherein SEQ ID NO: 125 is a clone designated herein as "DNA77568-1626".

[0153] **FIG. 126** shows the amino acid sequence (SEQ ID NO: 126) derived from the coding sequence of SEQ ID NO: 125 shown in **FIG. 125**.

[0154] **FIG. 127** shows a nucleotide sequence (SEQ ID NO: 127) of a native sequence PRO1774 cDNA, wherein SEQ ID NO: 127 is a clone designated herein as "DNA77626-1705".

[0155] **FIG. 128** shows the amino acid sequence (SEQ ID NO: 128) derived from the coding sequence of SEQ ID NO: 127 shown in **FIG. 127**.

[0156] **FIG. 129** shows a nucleotide sequence (SEQ ID NO: 129) of a native sequence PRO1928 cDNA, wherein SEQ ID NO: 129 is a clone designated herein as "DNA81754-2532".

[0157] **FIG. 130** shows the amino acid sequence (SEQ ID NO: 130) derived from the coding sequence of SEQ ID NO: 129 shown in **FIG. 129**.

[0158] **FIG. 131** shows a nucleotide sequence (SEQ ID NO: 131) of a native sequence PRO1865 cDNA, wherein SEQ ID NO: 131 is a clone designated herein as "DNA81757-2512".

[0159] **FIG. 132** shows the amino acid sequence (SEQ ID NO: 132) derived from the coding sequence of SEQ ID NO: 131 shown in **FIG. 131**.

[0160] **FIG. 133** shows a nucleotide sequence (SEQ ID NO: 133) of a native sequence PRO1925 cDNA, wherein SEQ ID NO: 133 is a clone designated herein as "DNA82302-2529".

[0161] **FIG. 134** shows the amino acid sequence (SEQ ID NO: 134) derived from the coding sequence of SEQ ID NO: 133 shown in **FIG. 133**.

[0162] **FIG. 135** shows a nucleotide sequence (SEQ ID NO: 135) of a native sequence PRO1926 cDNA, wherein SEQ ID NO: 135 is a clone designated herein as "DNA82340-2530".

[0163] **FIG. 136** shows the amino acid sequence (SEQ ID NO: 136) derived from the coding sequence of SEQ ID NO: 135 shown in **FIG. 135**.

[0164] **FIG. 137** shows a nucleotide sequence (SEQ ID NO: 137) of a native sequence PRO1801 cDNA, wherein SEQ ID NO: 137 is a clone designated herein as "DNA83500-2506".

[0165] **FIG. 138** shows the amino acid sequence (SEQ ID NO: 138) derived from the coding sequence of SEQ ID NO: 137 shown in **FIG. 137**.

[0166] **FIG. 139** shows a nucleotide sequence (SEQ ID NO: 139) of a native sequence PRO4405 cDNA, wherein SEQ ID NO: 139 is a clone designated herein as "DNA84920-2614".

[0167] **FIG. 140** shows the amino acid sequence (SEQ ID NO: 140) derived from the coding sequence of SEQ ID NO: 139 shown in **FIG. 139**.

[0168] **FIG. 141** shows a nucleotide sequence (SEQ ID NO: 141) of a native sequence PRO3435 cDNA, wherein SEQ ID NO: 141 is a clone designated herein as "DNA85066-2534".

[0169] FIG. 142 shows the amino acid sequence (SEQ ID NO: 142) derived from the coding sequence of SEQ ID NO: 141 shown in FIG. 141.

[0170] FIG. 143 shows a nucleotide sequence (SEQ ID NO: 143) of a native sequence PRO3543 cDNA, wherein SEQ ID NO: 143 is a clone designated herein as "DNA86571-2551".

[0171] FIG. 144 shows the amino acid sequence (SEQ ID NO: 144) derived from the coding sequence of SEQ ID NO: 143 shown in FIG. 143.

[0172] FIG. 145 shows a nucleotide sequence (SEQ ID NO: 145) of a native sequence PRO3443 cDNA, wherein SEQ ID NO: 145 is a clone designated herein as "DNA87991-2540".

[0173] FIG. 146 shows the amino acid sequence (SEQ ID NO: 146) derived from the coding sequence of SEQ ID NO: 145 shown in FIG. 145.

[0174] FIG. 147 shows a nucleotide sequence (SEQ ID NO: 147) of a native sequence PRO3442 cDNA, wherein SEQ ID NO: 147 is a clone designated herein as "DNA92238-2539".

[0175] FIG. 148 shows the amino acid sequence (SEQ ID NO: 148) derived from the coding sequence of SEQ ID NO: 147 shown in FIG. 147.

[0176] FIG. 149 shows a nucleotide sequence (SEQ ID NO: 149) of a native sequence PRO5990 cDNA, wherein SEQ ID NO: 149 is a clone designated herein as "DNA96042-2682".

[0177] FIG. 150 shows the amino acid sequence (SEQ ID NO: 150) derived from the coding sequence of SEQ ID NO: 149 shown in FIG. 149.

[0178] FIG. 151 shows a nucleotide sequence (SEQ ID NO: 151) of a native sequence PRO4342 cDNA, wherein SEQ ID NO: 151 is a clone designated herein as "DNA96787-2534".

[0179] FIG. 152 shows the amino acid sequence (SEQ ID NO: 152) derived from the coding sequence of SEQ ID NO: 151 shown in FIG. 151.

[0180] FIG. 153 shows a nucleotide sequence (SEQ ID NO: 153) of a native sequence PRO10096 cDNA, wherein SEQ ID NO: 153 is a clone designated herein as "DNA125185-2806".

[0181] FIG. 154 shows the amino acid sequence (SEQ ID NO: 154) derived from the coding sequence of SEQ ID NO: 153 shown in FIG. 153.

[0182] FIG. 155 shows a nucleotide sequence (SEQ ID NO: 155) of a native sequence PRO10272 cDNA, wherein SEQ ID NO: 155 is a clone designated herein as "DNA147531-2821".

[0183] FIG. 156 shows the amino acid sequence (SEQ ID NO: 156) derived from the coding sequence of SEQ ID NO: 155 shown in FIG. 155.

[0184] FIG. 157 shows a nucleotide sequence (SEQ ID NO: 157) of a native sequence PRO5801 cDNA, wherein SEQ ID NO: 157 is a clone designated herein as "DNA115291-2681".

[0185] FIG. 158 shows the amino acid sequence (SEQ ID NO: 158) derived from the coding sequence of SEQ ID NO: 157 shown in FIG. 157.

[0186] FIG. 159 shows a nucleotide sequence (SEQ ID NO: 159) of a native sequence PRO20110 cDNA, wherein SEQ ID NO: 159 is a clone designated herein as "DNA166819".

[0187] FIG. 160 shows the amino acid sequence (SEQ ID NO: 160) derived from the coding sequence of SEQ ID NO: 159 shown in FIG. 159.

[0188] FIG. 161 shows a nucleotide sequence (SEQ ID NO: 161) of a native sequence PRO20040 cDNA, wherein SEQ ID NO: 161 is a clone designated herein as "DNA64625-2890".

[0189] FIG. 162 shows the amino acid sequence (SEQ ID NO: 162) derived from the coding sequence of SEQ ID NO: 161 shown in FIG. 161.

[0190] FIG. 163 shows a nucleotide sequence (SEQ ID NO: 163) of a native sequence PRO20233 cDNA, wherein SEQ ID NO: 163 is a clone designated herein as "DNA165608".

[0191] FIG. 164 shows the amino acid sequence (SEQ ID NO: 164) derived from the coding sequence of SEQ ID NO: 163 shown in FIG. 163.

[0192] FIG. 165 shows a nucleotide sequence (SEQ ID NO: 165) of a native sequence PRO19670 cDNA, wherein SEQ ID NO: 165 is a clone designated herein as "DNA131639-2874".

[0193] FIG. 166 shows the amino acid sequence (SEQ ID NO: 166) derived from the coding sequence of SEQ ID NO: 165 shown in FIG. 165.

[0194] FIG. 167 shows a nucleotide sequence (SEQ ID NO: 167) of a native sequence PRO1890 cDNA, wherein SEQ ID NO: 167 is a clone designated herein as "DNA79230-2525".

[0195] FIG. 168 shows the amino acid sequence (SEQ ID NO: 168) derived from the coding sequence of SEQ ID NO: 167 shown in FIG. 167.

DETAILED DESCRIPTION

I. Definitions

[0196] The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The term "PRO polypeptide" refers to each individual PRO/number polypeptide disclosed herein. All disclosures in this specification which refer to the "PRO polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the prepa-

ration of, purification of, derivation of, formation of antibodies to or against, administration of, compositions containing, treatment of a disease with, etc., pertain to each polypeptide of the invention individually. The term "PRO polypeptide" also includes variants of the PRO/number polypeptides disclosed herein.

[0197] A "native sequence PRO polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO polypeptide derived from nature. Such native sequence PRO polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence PRO polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position 1 in the figures may be employed as the starting amino acid residue for the PRO polypeptides.

[0198] The PRO polypeptide "extracellular domain" or "ECD" refers to a form of the PRO polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

[0199] The approximate location of the "signal peptides" of the various PRO polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1-6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal

sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

[0200] "PRO polypeptide variant" means an active PRO polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Such PRO polypeptide variants include, for instance, PRO polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO polypeptide sequence as disclosed herein, a PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, PRO variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20 amino acids in length, alternatively at least about 30 amino acids in length, alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length, alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

[0201] "Percent (%) amino acid sequence identity" with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a

candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0202] In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

[0203] 100 times the fraction $\frac{X}{Y}$

[0204] where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X", "Y" and "Z" each represent different hypothetical amino acid residues.

[0205] Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as

described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an the amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

[0206] Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

[0207] In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

[0208] 100 times the fraction $\frac{X}{Y}$

[0209] where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

[0210] "PRO variant polynucleotide" or "PRO variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO polypeptide as defined below and which has at least about 80% nucleic acid sequence identity

with a nucleotide acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Ordinarily, a PRO variant polynucleotide will have at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO polypeptide sequence as disclosed herein, a full-length native sequence PRO polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

[0211] Ordinarily, PRO variant polynucleotides are at least about 30 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 210 nucleotides in length, alternatively at least about 240 nucleotides in length, alternatively at least about 270 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 900 nucleotides in length, or more.

[0212] "Percent (%) nucleic acid sequence identity" with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the

sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code: shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

[0213] In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

[0214] $100 \text{ times the fraction } \frac{W}{Z}$

[0215] where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA", wherein "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides.

[0216] Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., Methods in Enzymology 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid

molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement “an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B”, the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

[0217] Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

[0218] In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

[0219] $100 \text{ times the fraction } \frac{W}{Z}$

[0220] where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

[0221] In other embodiments, PRO variant polynucleotides are nucleic acid molecules that encode an active PRO polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO polypeptide as disclosed herein. PRO variant polypeptides may be those that are encoded by a PRO variant polynucleotide.

[0222] “Isolated,” when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other pro-

teinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide in situ within recombinant cells, since at least one component of the PRO polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

[0223] An “isolated” PRO polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

[0224] The term “control sequences” refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

[0225] Nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0226] The term “antibody” is used in the broadest sense and specifically covers, for example, single anti-PRO monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO antibody compositions with polypeptidic specificity, single chain anti-PRO antibodies, and fragments of anti-PRO antibodies (see below). The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

[0227] “Stringency” of hybridization reactions is readily determinable by one of ordinary skill in the art, and gener-

ally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

[0228] “Stringent conditions” or “high stringency conditions”, as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50° C.; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42° C.; or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt’s solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC (sodium chloride/sodium citrate) and 50% formamide at 55° C., followed by a high-stringency wash consisting of 0.1×SSC containing EDTA at 55° C.

[0229] “Moderately stringent conditions” may be identified as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and %SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37° C. in a solution comprising: 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt’s solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37-50° C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

[0230] The term “epitope tagged” when used herein refers to a chimeric polypeptide comprising a PRO polypeptide fused to a “tag polypeptide”. The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

[0231] As used herein, the term “immunoadhesin” designates antibody-like molecules which combine the binding

specificity of a heterologous protein (an “adhesin”) with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is “heterologous”), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

[0232] “Active” or “activity” for the purposes herein refers to form(s) of a PRO polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO, wherein “biological” activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO and an “immunological” activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO.

[0233] The term “antagonist” is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO polypeptide disclosed herein. In a similar manner, the term “agonist” is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO polypeptide may comprise contacting a PRO polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO polypeptide.

[0234] “Treatment” refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

[0235] “Chronic” administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. “Intermittent” administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

[0236] “Mammal” for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

[0237] Administration “in combination with” one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

[0238] "Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEENTM, polyethylene glycol (PEG), and PLURONICSTM.

[0239] "Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata et al., Protein Eng. 8(10): 1057-1062[1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

[0240] Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

[0241] "Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0242] The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0243] The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

[0244] Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins

can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

[0245] "Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 13, Rosenberg and Moore eds., Springer-Verlag, N.Y., pp. 269-315 (1994).

[0246] The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., Proc. Natl. Acad. Sci. USA, 90:6444-6448 (1993).

[0247] An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[0248] An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

[0249] The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

[0250] By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain

embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Pat. No. 4,275,149.

[0251] A “liposome” is a small vesicle composed of various types of lipids, phospholipids and/or surfactant

which is useful for delivery of a drug (such as a PRO polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

[0252] A “small molecule” is defined herein to have a molecular weight below about 500 Daltons

TABLE 1

```
/*
 *
 * C—C increased from 12 to 15
 * Z is average of EQ
 * B is average of ND
 * match with stop is _M; stop—stop = 0; J (joker) match = 0
 */
#define _M -8 /* value of a match with a stop */
int
_day[26][26] = {
/* A B C D E F G H I J K L M N O P Q R S T U V W X Y Z */
/* A */ {2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */ {0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */ {-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */ {0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */ {0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */ {-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */ {1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */ {-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */ {-1, -2, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */ {-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */ {-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2}
/* M */ {-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */ {0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */ {_M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */ {1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */ {0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */ {-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */ {1, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */ {1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */ {0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */ {-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */ {0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */ {-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */ {0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, 0, -2, -6, 0, -4, 4},
};
/*
 */
#include <stdio.h>
#include <ctype.h>
#define MAXJMP 16 /* max jumps in a diag */
#define MAXGAP 24 /* don't continue to penalize gaps larger than this */
#define JMPS 1024 /* max jmps in an path */
#define MX 4 /* save if there's at least MX-1 bases since last jmp */
#define DMAT 3 /* value of matching bases */
#define DMIS 0 /* penalty for mismatched bases */
#define DINS0 8 /* penalty for a gap */
#define DINS1 1 /* penalty per base */
#define PINS0 8 /* penalty for a gap */
#define PINS1 4 /* penalty per residue */
struct jmp {
    short n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
    /* limits seq to 2 16 -1 */
};
struct diag {
    int score; /* score at last jmp */
    long offset; /* offset of prev block */
    short ijmp; /* current jmp index */
    struct jmp jp; /* list of jmps */
};
struct path {
    int spc; /* number of leading spaces */
};
```

TABLE 1-continued

```
short      n[JMP$];          /* size of jmp (gap) */
int        x[JMP$];          /* loc of jmp (last elem before gap) */
};
char        *ofile;           /* output file name */
char        *namex[2];        /* seq names: getseqs() */
char        *prog;            /* prog name for err msgs */
char        *seqx[2];         /* seqs: getseqs() */
int         dmax;             /* best diag: nw() */
int         dmax0;            /* final diag */
int         dna;              /* set if dna: main() */
int         endgaps;          /* set if penalizing end gaps */
int         gapx, gapy;        /* total gaps in seqs */
int         len0, len1;        /* seq lens */
int         ngapx, ngapy;      /* total size of gaps */
int         smax;             /* max score: nw() */
int         *xbm;             /* bitmap for matching */
long        offset;           /* current offset in jmp file */
struct diag *dx;              /* holds diagonals */
struct path pp[2];            /* holds path for seqs */
char        *calloc(), *malloc(), *index(), *strcpy();
char        *getseq(), *g_calloc();
/* Needleman-Wunsch alignment program
 *
 * usage: progs file1 file2
 *   where file1 and file2 are two dna or two protein sequences.
 *   The sequences can be in upper- or lower-case and may contain ambiguity
 *   Any lines beginning with ';', '>' or '<' are ignored
 *   Max file length is 65535 (limited by unsigned short x in the jmp struct)
 *   A sequence with 1/3 or more of its elements ACGTU is assumed to be DNA
 *   Output is in the file "align.out"
 *
 * The program may create a tmp file in /tmp to hold info about traceback.
 * Original version developed under BSD 4.3 on a vax 8650
 */
#include "nw.h"
#include "day.h"
static  _dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};
static  _pbval[26] = {
    1, 2[(1<<('D'-'A'))|(1<<('N'-'A'))], 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25[(1<<('E'-'A'))|(1<<('Q'-'A'))]
};
main(ac, av)
int      ac;
char     *av[];
{
    prog = av[0];
    if(ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
        fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
        fprintf(stderr, "The sequences can be in upper- or lower-case\n");
        fprintf(stderr, "Any lines beginning with ';' or '<' are ignored\n");
        fprintf(stderr, "Output is in the file \"align.out\"\n");
        exit(1);
    }
    namex[0] = av[1];
    namex[1] = av[2];
    seqx[0] = getseq(namex[0], &len0);
    seqx[1] = getseq(namex[1], &len1);
    xbm = (dna)? _dbval : _pbval;
    endgaps = 0;          /* 1 to penalize endgaps */
    ofile = "align.out";  /* output file */
    nw();                 /* fill in the matrix, get the possible jmps */
    readjmps();           /* get the actual jmps */
    print();              /* print stats, alignment */
    cleanup(0);           /* unlink any tmp files */
}
/* do the alignment, return best score: main()
 * dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
 * pro: PAM 250 values
 * When scores are equal, we prefer mismatches to any gap, prefer
 * a new gap to extending an ongoing gap, and prefer a gap in seqx
 * to a gap in seq y.
```

TABLE 1-continued

```

*/
nw()
{
    char    *px, *py;        /* seqs and ptrs */
    int     *ndely, *dely;    /* keep track of dely */
    int     ndelx, delx;      /* keep track of delx */
    int     *tmp;            /* for swapping row0, row1 */
    int     mis;             /* score for each type */
    int     ins0, ins1;       /* insertion penalties */
    register id;             /* diagonal index */
    register ij;             /* jmp index */
    register *col0, *col1;    /* score for curr, last row */
    register xx, yy;         /* index into seqs */
    dx = (struct diag *)g_calloc("to get diag", len0+len1+1, sizeof(struct diag));
    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;
    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0;        /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;
    /* fill in match matrix
    */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
        */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0]-ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }

    for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
        mis = col0[yy-1];
        if (dna)
            mis += (xbm[*px-'A']&xbm[*py-'A'])? DMAT : DMIS;
        else
            mis += _day[*px-'A'][*py-'A'];
        /* update penalty for del in x seq;
        * favor new del over ongoing del
        * ignore MAXGAP if weighting endgaps
        */
        if (endgaps || ndely[yy] < MAXGAP) {
            if (col0[yy] - ins0 >= dely[yy]) {
                dely[yy] = col0[yy] - (ins0+ins1);
                ndely[yy] = 1;
            } else {
                dely[yy] -= ins1;
                ndely[yy]++;
            }
        } else {
            if (col0[yy] - (ins0+ins1) >= dely[yy]) {
                dely[yy] = col0[yy] - (ins0+ins1);
                ndely[yy] = 1;
            } else
                ndely[yy]++;
        }
        /* update penalty for del in y seq;
        * favor new del over ongoing del
    }
}

```

TABLE 1-continued

```

    */
    if (endgaps || ndelx < MAXGAP) {
        if (col1[yy-1] - ins0 >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else {
            delx -= ins1;
            ndelx++;
        }
    } else {
        if (col1[yy-1] - (ins0+ins1) >= delx) {
            delx = col1[yy-1] - (ins0+ins1);
            ndelx = 1;
        } else
            ndelx++;
    }
    /* pick the maximum score; we're favoring
    * mis over any del and delx over dely
    */

    id = xx - yy + len1 - 1;
    if (mis >= delx && mis >= dely[yy])
        col1[yy] = mis;
    else if (delx >= dely[yy]) {
        col1[yy] = delx;
        ij = dx[id].ijmp;
        if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
        && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
            dx[id].ijmp++;
            if (++ij >= MAXJMP) {
                writeimps(id);
                ij = dx[id].ijmp = 0;
                dx[id].offset = offset;
                offset += sizeof(struct jmp) + sizeof(offset);
            }
        }
        dx[id].jp.n[ij] = ndelx;
        dx[id].jp.x[ij] = xx;
        dx[id].score = delx;
    }
    else {
        col1[yy] = dely[yy];
        ij = dx[id].ijmp;
        if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
        && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
            dx[id].ijmp++;
            if (++ij >= MAXJMP) {
                writeimps(id);
                ij = dx[id].ijmp = 0;
                dx[id].offset = offset;
                offset += sizeof(struct jmp) + sizeof(offset);
            }
        }
        dx[id].jp.n[ij] = ndely[yy];
        dx[id].jp.x[ij] = xx;
        dx[id].score = dely[yy];
    }
    if (xx == len0 && yy < len1) {
        /* last col
        */
        if (endgaps)
            col1[yy] -= ins0+ins1*(len1-yy);
        if (col1[yy] > smax) {
            smax = col1[yy];
            dmax = id;
        }
    }
}
if (endgaps && xx < len0)
    col1[yy-1] -= ins0+ins1*(len0-xx);
if (col1[yy-1] > smax) {
    smax = col1[yy-1];
    dmax = id;
}
tmp = col0; col0 = col1; col1 = tmp;
}

```

...nw

TABLE 1-continued

```

        (void) free((char *)ndely);
        (void) free((char *)dely);
        (void) free((char *)col0);
        (void) free((char *)col1);
    }
/*
 *
 * print() -- only routine visible outside this module
 *
 * static:
 * getmat() -- trace back best path, count matches: print()
 * pr_align() -- print alignment of described in array p[]: print()
 * dumpblock() -- dump a block of lines with numbers, stars: pr_align()
 * nums() -- put out a number line: dumpblock()
 * putline() -- put out a line (name, [num], seq, [num]): dumpblock()
 * stars() -- put a line of stars: dumpblock()
 * stripname() -- strip any path and prefix from a seqname
 */
#include "nw.h"
#define SPC 3
#define P_LINE 256 /* maximum output line */
#define P_SPC 3 /* space between name or num and seq */
extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */
print()
{
    int lx, ly, firstgap, lastgap; /* overlap */
    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}
/*
 * trace back the best path, count matches
 */
static
getmat(lx, ly, firstgap, lastgap)
int lx, ly; /* "core" (minus endgaps) */
int firstgap, lastgap; /* leading trailing overlap */
{
    int nm, i0, i1, siz0, siz1;
    char outx[32];
    double pct;
    register n0, n1;
    register char *p0, *p1;
    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;

```

TABLE 1-continued

```

n1 = pp[0].spc + 1;
nm = 0;
while ( *p0 && *p1 ) {
    if (siz0) {
        p1++;
        n1++;
        siz0--;
    }
    else if (siz1) {
        p0++;
        n0++;
        siz1--;
    }
    else {
        if (xbm[*p0-'A']&xbm[*p1-'A'])
            nm++;
        if (n0++ == pp[0].x[i0])
            siz0 = pp[0].n[i0++];
        if (n1++ == pp[1].x[i1])
            siz1 = pp[1].n[i1++];
        p0++;
        p1++;
    }
}
/* pct homology:
 * if penalizing endgaps, base is the shorter seq
 * else, knock off overhangs and take shorter core
 */
if (endgaps)
    lx = (len0 < len1)? len0 : len1;
else
    lx = (lx < ly)? lx : ly;
pct = 100.*(double)nm/(double)lx;
fprintf(fx, "\n");
fprintf(fx, "<%d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);
fprintf(fx, "<gaps in first sequence: %d", gapx);
if (gapx) {
    (void) sprintf(outx, "(%d %s%s)",
        ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
    fprintf(fx, "%s", outx);
    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, "(%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
}
if (dna)
    fprintf(fx,
        "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
        smax, DMAT, DMIS, DINS0, DINS1);
else
    fprintf(fx,
        "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
        smax, PINS0, PINS1);
if (endgaps)
    fprintf(fx,
        "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
        firstgap, (dna)? "base" : "residue", (firstgap == 1)? "" : "s",
        lastgap, (dna)? "base" : "residue", (lastgap == 1)? "" : "s");
else
    fprintf(fx, "<endgaps not penalized\n");
}
static      nm;                /* matches in core -- for checking */
static      lmax;              /* lengths of stripped file names */
static      ij[2];             /* jmp index for a path */
static      nc[2];             /* number at start of current line */
static      ni[2];             /* current elem number -- for gapping */
static      siz[2];
static char  *ps[2];           /* ptr to current element */
static char  *po[2];           /* ptr to next output char slot */
static char  out[2][P_LINE];   /* output line */
static char  star[P_LINE];     /* set by stars() */
/*
 * print alignment of described in struct path pp[]
 */

```

TABLE 1-continued

```

static
pr_align()
{
    int      nn;          /* char count */
    int      more;
    register i;
    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(name[i]);
        if (nn > lmax)
            lmax = nn;
        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = out[i];
    }
    for (nn = nm = 0, more = 1; more;) {
        for (i = more = 0; i < 2; i++) {
            /*
             * do we have more of this sequence?
             */
            if (!*ps[i])
                continue;
            more++;
            if (pp[i].spc) { /* leading space */
                *po[i]++ = ' ';
                pp[i].spc--;
            }
            else if (siz[i]) { /* in a gap */
                *po[i]++ = '-';
                siz[i]--;
            }
            else { /* we're putting a seq element
             */
                *po[i] = *ps[i];
                if (islower(*ps[i]))
                    *ps[i] = toupper(*ps[i]);
                po[i]++;
                ps[i]++;
                /*
                 * are we at next gap for this seq?
                 */
                if (ni[i] == pp[i].x[ij[i]]) {
                    /*
                     * we need to merge all gaps
                     * at this location
                     */
                    siz[i] == pp[i].n[ij[i]++];
                    while (ni[i] == pp[i].x[ij[i]])
                        siz[i] += pp[i].n[ij[i]++];
                }
                ni[i]++;
            }
        }
        if (++nn == olen || !more && nn) {
            dumpblock();
            for (i = 0; i < 2; i++)
                po[i] = out[i];
            nn = 0;
        }
    }
}
/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()
{
    register i;
    for(i = 0; i < 2; i++)
        *po[i]-- = '\0';

    (void) puts("\n", fx);
    for (i = 0; i < 2; i++) {
        if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
            if (i == 0)
                nums(i);
        }
    }
}

```

pr_align

...pr_align

dumpblock

...dumpblock

TABLE 1-continued

<pre> if (i == 0 && *out[1]) stars(); putline(i); if (i == 0 && *out[1]) fprintf(fx, star); if (i == 1) nums(i); } } /* * put out a number line: dumpblock() */ static nums(ix) int ix; /* index in out[] holding seq line */ { char nline[P_LINE]; register i, j; register char *pn, *px, *py; for(pn = nline, i = 0; i < lmax+P_SPC; i++, pn++) *pn = ' '; for (i = nc[ix], py = out[ix]; *py; py++, pn++) { if (*py == ' ' *py == '-') *pn = ' '; else { if (i%10 == 0 (i == 1 && nc[ix] != 1)) { j = (i < 0)? -i : i; for (px = pn; j; j/= 10, px--) *px = j%10 + '0'; if (i < 0) *px = '-'; } else *pn = ' '; i++; } } *pn = '\0'; nc[ix] = i; for (pn = nline; *pn; pn++) (void) putc(*pn, fx); (void) putc('\n', fx); } /* * put out a line (name, [num], seq. [num]): dumpblock() */ static putline(ix) int ix; { int i; register char *px; for (px = namex[ix], i = 0; *px && *px != '.'; px++, i++) (void) putc(*px, fx); for (; i < lmax+P_SPC; i++) (void) putc(' ', fx); /* these count from 1: * ni[] is current element (from 1) * nc[] is number at start of current line */ for (px = out[ix]; *px; px++) (void) putc(*px&0x7F, fx); (void) putc('\n', fx); } } /* * put a line of stars (seqs always in out[0], out[1]): dumpblock() */ static stars() { int i; register char *p0, *p1, cx, *px; if (!(*out[0] (*out[0] == ' ' && *(p0[0]) == ' ') !*out[1] (*out[1] == ' ' && *(p0[1]) == ' ')) return; </pre>		
		nums
		putline
		...putline
		stars

TABLE 1-continued

```

    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';
    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A']&xbm[*p1-'A']) {
                cx = '*';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)
char    *pn;                /* file name (may be path) */
{
    register char    *px, *py;
    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}
/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_alloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>
char    *jname = "/tmp/homgXXXXXX";    /* tmp file for jumps */
FILE    *fj;
int    cleanup();                /* cleanup tmp file */
long    lseek();
/*
 * remove any tmp file if we blow
 */
cleanup(i)
int    i;
{
    if (fj)
        (void) unlink(jname);
    exit(i);
}
/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char    *
getseq(file, len)
char    *file;                /* file name */
int    *len;                /* seq len */
{
    char    line[1024], *pseq;
    register char    *px, *py;
    int    natgc, tlen;
    FILE    *fp;
    if ((fp = fopen(file, "r")) == 0) {
        fprintf(stderr, "%s: can't read %s\n", prog, file);
        exit(1);
    }

```

TABLE 1-continued

<pre> } tlen = natgc = 0; while (fgets(line, 1024, fp)) { if (*line == ';' *line == '<' *line == '>') continue; for (px = line; *px != '\n'; px++) if (isupper(*px) islower(*px)) tlen++; } if ((pseq = malloc((unsigned)(tlen+6))) == 0) { fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file); exit(1); } pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0'; py = pseq + 4; *len = tlen; rewind(fp); while (fgets(line, 1024, fp)) { if (*line == ';' *line == '<' *line == '>') continue; for (px = line; *px != '\n'; px++) { if (isupper(*px)) *py++ = *px; else if (islower(*px)) *py++ = toupper(*px); if (index("ATGCU", *(py-1))) natgc++; } } *py++ = '\0'; *py = '\0'; (void) fclose(fp); dna = natgc > (tlen/3); return(pseq+4); } char * g__alloc(msg, nx, sz) char *msg; int nx, sz; /* program, calling routine */ /* number and size of elements */ { char *px, *calloc(); if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) { if (*msg) { fprintf(stderr, "%s: g__alloc() failed %s (n= %d, sz= %d)\n", prog, msg, nx, sz); exit(1); } } return(px); } /* * get final jmps from dx[] or tmp file, set pp[], reset dmax: main() */ readjmps() { int fd = -1; int siz, i0, i1; register i, j, xx; if (fj) { (void) fclose(fj); if ((fd = open(jname, O_RDONLY, 0)) < 0) { fprintf(stderr, "%s: can't open() %s\n", prog, jname); cleanup(1); } } for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++) { while (1) { for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--) ; if (j < 0 && dx[dmax].offset && fj) { (void) lseek(fd, dx[dmax].offset, 0); (void) read(fd, (char *) &dx[dmax].jp, sizeof(struct jmp)); (void) read(fd, (char *) &dx[dmax].offset, sizeof(dx[dmax].offset)); dx[dmax].ijmp = MAXJMP-1;</pre>	...getseq
	g__alloc
	readjmps
	...readjmps

TABLE 1-continued

```

    }
    else
        break;
    }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
    }
    if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
        if (siz < 0) { /* gap in second seq */
            pp[1].n[i1] = -siz;
            xx += siz;
            /* id = xx - yy + len1 - 1
            */
            pp[1].x[i1] = xx - dmax + len1 - 1;
            gapy++;
            ngapy -= siz;
/* ignore MAXGAP when doing endgaps */
            siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
            i1++;
        }
        else if (siz > 0) { /* gap in first seq */
            pp[0].n[i0] = siz;
            pp[0].x[i0] = xx;
            gapx++;
            ngapx += siz;
/* ignore MAXGAP when doing endgaps */
            siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
            i0++;
        }
    }
    else
        break;
}
/* reverse the order of jmps
*/
for (j = 0, i0--; j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--; j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}

/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)
int ix;
{
    char *mktemp();
    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}

```

[0253]

TABLE 2

PRO	XXXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXXXYYYYYYY	(Length = 12 amino acids)

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 15 = 33.3%

[0254]

TABLE 3

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXXXYYYZZYZ	(Length = 15 amino acids)

% amino acid sequence identity = (the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide) = 5 divided by 10 = 50%

[0255]

TABLE 4

PRO-DNA	NNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 6 divided by 14 = 42.9%

[0256]

TABLE 5

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

% nucleic acid sequence identity = (the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence) = 4 divided by 12 = 33.3%

II. Compositions and Methods of the Invention

[0257] A. Fall-Length PRO Polypeptides

[0258] The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO polypeptides. In particular, cDNAs encoding various PRO polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

[0259] As disclosed in the Examples below, various cDNA clones have been deposited with the ATCC. The actual

nucleotide sequences of those clones can readily be determined by the skilled artisan by sequencing of the deposited clone using routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

[0260] 5. PRO Polypeptide Variants

[0261] In addition to the full-length native sequence PRO polypeptides described herein, it is contemplated that PRO variants can be prepared. PRO variants can be prepared by introducing appropriate nucleotide changes into the PRO DNA, and/or by synthesis of the desired PRO polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

[0262] Variations in the native full-length sequence PRO or in various domains of the PRO described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Pat. No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO that results in a change in the amino acid sequence of the PRO as compared with the native sequence PRO. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

[0263] PRO polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO polypeptide.

[0264] PRO fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by poly-

merase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO polypeptide fragments share at least one biological and/or immunological activity with the native PRO polypeptide disclosed herein.

[0265] In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

TABLE 6

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

[0266] Substantial modifications in function or immunological identity of the PRO polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- [0267] (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- [0268] (2) neutral hydrophilic: cys, ser, thr;
- [0269] (3) acidic: asp, glu;
- [0270] (4) basic: asn, gin, his, lys, arg;
- [0271] (5) residues that influence chain orientation: gly, pro; and
- [0272] (6) aromatic: trp, tyr, phe.

[0273] Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

[0274] The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO variant DNA.

[0275] Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W. H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

[0276] C. Modifications of PRO

[0277] Covalent modifications of PRO are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of the PRO. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis (diazocetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propioimide.

[0278] Other modifications include deamidation of glutamyl and asparaginy residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T. E. Creighton, Proteins: Structure and Molecular Properties, W. H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[0279] Another type of covalent modification of the PRO polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means),

and/or adding one or more glycosylation sites that are not present in the native sequence PRO. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

[0280] Addition of glycosylation sites to the PRO polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO (for O-linked glycosylation sites). The PRO amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

[0281] Another means of increasing the number of carbohydrate moieties on the PRO polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published 22 Sep. 1987, and in Aplin and Wriston, *CRC Crit. Rev. Biochem.*, pp. 259-306 (1981).

[0282] Removal of carbohydrate moieties present on the PRO polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., *Arch. Biochem. Biophys.*, 259:52 (1987) and by Edge et al., *Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., *Meth. Enzymol.*, 138:350 (1987).

[0283] Another type of covalent modification of PRO comprises linking the PRO polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

[0284] The PRO of the present invention may also be modified in a way to form a chimeric molecule comprising PRO fused to another, heterologous polypeptide or amino acid sequence.

[0285] In one embodiment, such a chimeric molecule comprises a fusion of the PRO with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl-terminus of the PRO. The presence of such epitope-tagged forms of the PRO can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547-553

(1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, 6:1204-1210(1988)]; the KT3 epitopepeptide [Martinet et al., *Science*, 255:192-194 (1992)]; an α -tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, 87:6393-6397 (1990)].

[0286] In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an "immunoadhesin"), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG molecule. For the production of immunoglobulin fusions see also U.S. Pat. No. 5,428,130 issued Jun. 27, 1995.

[0287] D. Preparation of PRO

[0288] The description below relates primarily to production of PRO by culturing cells transformed or transfected with a vector containing PRO nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO. For instance, the PRO sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., *Solid-Phase Peptide Synthesis*, W. H. Freeman Co., San Francisco, Calif. (1969); Merrifield, *J. Am. Chem. Soc.*, 85:2149-2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, Calif.) using manufacturer's instructions. Various portions of the PRO may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO.

[0289] 1. Isolation of DNA Encoding PRO

[0290] DNA encoding PRO may be obtained from a cDNA library prepared from tissue believed to possess the PRO mRNA and to express it at a detectable level. Accordingly, human PRO DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

[0291] Libraries can be screened with probes (such as antibodies to the PRO or oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO is to use PCR methodology [Sambrook et al., *supra*; Dieffenbach et al., *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)].

[0292] The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

[0293] Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

[0294] Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

[0295] 2. Selection and Transformation of Host Cells

[0296] Host cells are transfected or transformed with expression or cloning vectors described herein for PRO production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

[0297] Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, $\text{CaCl}_2\text{-CaPO}_4$, liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 Jun. 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene,

polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

[0298] Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Elwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *S. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 Apr. 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype tonA; *E. coli* W310 strain 9E4, which has the complete genotype tonA ptr3; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype tonA ptr3 phoA E15 (argF-lac)769 degP ompT kan^r; *E. coli* W3110 strain 37D6, which has the complete genotype tonA ptr3 phoA E15 (argF-lac)769 degP ompT rbs7 ilvG kan^r; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant degP deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Pat. No. 4,946,783 issued 7 Aug. 1990. Alternatively, in vitro methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

[0299] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, Nature, 290: 140[1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Pat. No. 4,943,529; Fleer et al., Bio/Technology, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., J. Bacteriol., 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilum* (ATCC 36,906; Van den Berg et al., Bio/Technology, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., J. Basic Microbiol., 28:265-278 [1988]); *Candida*; *Trichoderma reesia* (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 Oct. 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published 10 Jan. 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983];

Tilburn et al., *Gene*, 26:205-221 [1983]; Yelton et al., *Proc. Natl. Acad. Sci. USA*, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, *EMBO J.*, 4:475-479 [1985]). Methylo-tropic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylo-trophs*, 269 (1982).

[0300] Suitable host cells for the expression of glycosylated PRO are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); Chinese hamster ovary cells/DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

[0301] 3. Selection and use of a Replicable Vector

[0302] The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

[0303] The PRO may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* (α -factor leaders, the latter described in U.S. Pat. No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 Apr. 1990), or the signal described in WO 90/13646 published 15 Nov. 1990. In mammalian cell expression, mammalian signal

sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

[0304] Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

[0305] Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

[0306] An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980). A suitable selection gene for use in yeast is the trp 1 gene present in the yeast plasmid YRp7 [Stinchcomb et al., *Nature*, 282:39 (1979); Kingsman et al., *Gene*, 7:141 (1979); Tschemper et al., *Gene*, 10:157 (1980)]. The trp 1 gene provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, *Genetics*, 85:12 (1977)].

[0307] Expression and cloning vectors usually contain a promoter operably linked to the PRO-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., *Nature*, 275:615 (1978); Goeddel et al., *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (trp) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the tac promoter [deBoer et al., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO.

[0308] Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes Hess et al., *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

[0309] Other yeast promoters, which are inducible promoters having the additional advantage of transcription

controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

[0310] PRO transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowpox virus (UK 2,211,504 published 5 Jul. 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

[0311] Transcription of a DNA encoding the PRO by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO coding sequence, but is preferably located at a site 5' from the promoter.

[0312] Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO.

[0313] Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO in recombinant vertebrate cell culture are described in Gething et al., *Nature*, 293:620-625 (1981); Mantei et al., *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

[0314] 4. Detecting Gene Amplification/Expression

[0315] Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201(1980)], dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

[0316] Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO DNA and encoding a specific antibody epitope.

[0317] 5. Purification of Polypeptide

[0318] Forms of PRO may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

[0319] It may be desired to purify PRO from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, *Methods in Enzymology*, 182 (1990); Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, N.Y. (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO produced.

[0320] E. Uses for PRO

[0321] Nucleotide sequences (or their complement) encoding PRO have various applications in the art of molecular biology, including uses as hybridization probes, in chromosome and gene mapping and in the generation of anti-sense RNA and DNA. PRO nucleic acid will also be useful for the preparation of PRO polypeptides by the recombinant techniques described herein.

[0322] The full-length native sequence PRO gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length PRO cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO or PRO from other species) which have a desired sequence identity to the native PRO sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence PRO. By way of example, a screening method will comprise isolating the coding region of the PRO gene using the known DNA sequence to synthesize a selected probe of about 40 bases.

Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ^{32}P or ^{35}S , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

[0323] Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

[0324] Other useful fragments of the PRO nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target PRO mRNA (sense) or PRO DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of PRO DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (Cancer Res. 48:2659, 1988) and van der Krol et al. (BioTechniques 6:958, 1988).

[0325] Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of PRO proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable in vivo (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

[0326] Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increases affinity of the oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

[0327] Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO_4 mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either in vivo or ex vivo. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retro-

virus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCTSC (see WO 90/13641).

[0328] Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

[0329] Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

[0330] Antisense or sense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

[0331] The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO coding sequences.

[0332] Nucleotide sequences encoding a PRO can also be used to construct hybridization probes for mapping the gene which encodes that PRO and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as in situ hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

[0333] When the coding sequences for PRO encode a protein which binds to another protein (example, where the PRO is a receptor), the PRO can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO or a receptor for PRO. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein

binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

[0334] Nucleic acids which encode PRO or its modified forms can also be used to generate either transgenic animals or “knock out” animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

[0335] Alternatively, non-human homologues of PRO can be used to construct a PRO “knock out” animal which has a defective or altered gene encoding PRO as a result of homologous recombination between the endogenous gene encoding PRO and altered genomic DNA encoding PRO introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO can be used to clone genomic DNA encoding PRO in accordance with established techniques. A portion of the genomic DNA encoding PRO can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal (e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113-152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a “knock out” animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock-

out animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the PRO polypeptide.

[0336] Nucleic acid encoding the PRO polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve in vivo synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. “Gene therapy” includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes in vivo. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik et al., *Proc. Natl. Acad. Sci. USA* 83:4143-4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

[0337] There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells in vitro, or in vivo in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells in vitro include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred in vivo gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11, 205-210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429-4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410-3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., *Science* 256, 808-813 (1992).

[0338] The PRO polypeptides described herein may also be employed as molecular weight markers for protein electrophoresis purposes and the isolated nucleic acid sequences may be used for recombinantly expressing those markers.

[0339] The nucleic acid molecules encoding the PRO polypeptides or fragments thereof described herein are useful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each PRO nucleic acid molecule of the present invention can be used as a chromosome marker.

[0340] The PRO polypeptides and nucleic acid molecules of the present invention may also be used diagnostically for tissue typing, wherein the PRO polypeptides of the present invention may be differentially expressed in one tissue as compared to another, preferably in a diseased tissue as compared to a normal tissue of the same tissue type. PRO nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

[0341] The PRO polypeptides described herein may also be employed as therapeutic agents. The PRO polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the PRO product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEENTM, PLURONICSTM or PEG.

[0342] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

[0343] Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[0344] The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

[0345] Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. "The use of interspecies scaling in toxicokinetics" In Toxicokinetics and New Drug Development, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42-96.

[0346] When in vivo administration of a PRO polypeptide or agonist or antagonist thereof is employed, normal dosage

amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 μ g/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue.

[0347] Where sustained-release administration of a PRO polypeptide is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of the PRO polypeptide, microencapsulation of the PRO polypeptide is contemplated. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon-(rhIFN-), interleukin-2, and MN rgp120. Johnson et al., Nat. Med., 2:795-799 (1996); Yasuda, Biomed. Ther., 27:1221-1223 (1993); Hora et al., Bio/Technology, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Poly-lactide Polyglycolide Microsphere Systems," in Vaccine Design: The Subunit and Adjuvant Approach, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

[0348] The sustained-release formulations of these proteins were developed using poly-lactic-coglycolic acid (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), Biodegradable Polymers as Drug Delivery Systems (Marcel Dekker: New York, 1990), pp. 1-41.

[0349] This invention encompasses methods of screening compounds to identify those that mimic the PRO polypeptide (agonists) or prevent the effect of the PRO polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

[0350] The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

[0351] All assays for antagonists are common in that they call for contacting the drug candidate with a PRO polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

[0352] In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction

mixture. In a particular embodiment, the PRO polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments. Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

[0353] If the candidate compound interacts with but does not bind to a particular PRO polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, *Nature* (London), 340:245-246 (1989); Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA*, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-lacZ reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKER™) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

[0354] Compounds that interfere with the interaction of a gene encoding a PRO polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the

reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

[0355] To assay for antagonists, the PRO polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the PRO polypeptide indicates that the compound is an antagonist to the PRO polypeptide. Alternatively, antagonists may be detected by combining the PRO polypeptide and a potential antagonist with membrane-bound PRO polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO polypeptide can be labeled, such as by radioactivity, such that the number of PRO polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., *Current Protocols in Immun.*, 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO polypeptide. Transfected cells that are grown on glass slides are exposed to labeled PRO polypeptide. The PRO polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

[0356] As an alternative approach for receptor identification, labeled PRO polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

[0357] In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled PRO polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

[0358] More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with PRO polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain anti-

bodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO polypeptide.

[0359] Another potential PRO polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix—see Lee et al., *Nucl. Acids Res.*, 6:3073 (1979); Cooney et al., *Science*, 241: 456 (1988); Dervan et al., *Science*, 251:1360 (1991)), thereby preventing transcription and the production of the PRO polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into the PRO polypeptide (antisense—Okano, *Neurochem.*, 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression (CRC Press: Boca Raton, Fla., 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed in vivo to inhibit production of the PRO polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

[0360] Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO polypeptide, thereby blocking the normal biological activity of the PRO polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

[0361] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, *Current Biology*, 4:469-471(1994), and PCT publication No. WO 97/33551 (published Sep. 18, 1997).

[0362] Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, supra.

[0363] These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

[0364] Diagnostic and therapeutic uses of the herein disclosed molecules may also be based upon the positive functional assay hits disclosed and described below.

[0365] F. Ant-PRO Antibodies

[0366] The present invention further provides anti-PRO antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

[0367] 1. Polyclonal Antibodies

[0368] The anti-PRO antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

[0369] 2. Monoclonal Antibodies

[0370] The anti-PRO antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

[0371] The immunizing agent will typically include the PRO polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl

transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

[0372] Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, Calif. and the American Type Culture Collection, Manassas, Va. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

[0373] The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against PRO. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

[0374] After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown in vivo as ascites in a mammal.

[0375] The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

[0376] The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Pat. No. 4,816,567; Morrison et al., supra] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be sub-

stituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

[0377] The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

[0378] In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

[0379] 3. Human and Humanized Antibodies

[0380] The anti-PRO antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-329 (1988); and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)].

[0381] Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers Bones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeven et al., Science, 239:1534-1536 (1988)], by: substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corre-

sponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

[0382] Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)). Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *Bio/Technology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51, (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

[0383] The antibodies may also be affinity matured using known selection and/or mutagenesis methods as described above. Preferred affinity matured antibodies have an affinity which is five times, more preferably 10 times, even more preferably 20 or 30 times greater than the starting antibody (generally murine, humanized or human) from which the matured antibody is prepared.

[0384] 4. Bispecific Antibodies

[0385] Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

[0386] Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Trautnecker et al., *EMBO J.*, 10:3655-3659 (1991).

[0387] Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain

constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

[0388] According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

[0389] Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

[0390] Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217(1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

[0391] Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148 (5):1547-1553(1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the

hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., Proc. Natl. Acad. Sci. USA 90:6444-6448(1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., J. Immunol. 152:5368 (1994). Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al. J. Immunol. 147:60 (1991).

[0392] Exemplary bispecific antibodies may bind to two different epitopes on a given PRO polypeptide herein. Alternatively, an anti-PRO polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc γ R), such as Fc γ RI (CD64), Fc γ RII (CD32) and Fc γ RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO polypeptide. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular PRO polypeptide. These antibodies possess a PRO-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO polypeptide and further binds tissue factor (TF).

[0393] 5. Heteroconjugate Antibodies

[0394] Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Pat. No. 4,676,980], and for treatment of HIV infection [WO 91, 000360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimide and those disclosed, for example, in U.S. Pat. No. 4,676,980.

[0395] 6. Effector Function Engineering

[0396] It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., J. Exp. Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using

heterobifunctional cross-linkers as described in Wolff et al. Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., Anti-Cancer Drug Design, 3: 219-230 (1989).

[0397] 7. Immunoconjugates

[0398] The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

[0399] Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolaca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcumin, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ^{212}Bi , ^{131}I , ^{131}In , ^{90}Y , and ^{186}Re . Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimide HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionuclide to the antibody. See WO 94/11026.

[0400] In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionuclide).

[0401] 8. Immunoliposomes

[0402] The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., Proc. Natl. Acad. Sci. USA, 82: 3688 (1985); Hwang et al., Proc. Natl. Acad. Sci. USA, 77:4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556.

[0403] Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid compo-

sition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem., 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., J. National Cancer Inst., 81(19): 1484 (1989).

[0404] 9. Pharmaceutical Compositions of Antibodies

[0405] Antibodies specifically binding a PRO polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

[0406] If the PRO polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., Proc. Natl. Acad. Sci. USA, 90: 7889-7893 (1993). The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

[0407] The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nano-capsules) or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, supra.

[0408] The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

[0409] Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOTTM

(injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

[0410] G. Uses for Anti-PRO Antibodies

[0411] The anti-PRO antibodies of the invention have various utilities. For example, anti-PRO antibodies may be used in diagnostic assays for PRO, e.g., detecting its expression (and in some cases, differential expression) in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, Monoclonal Antibodies: A Manual of Techniques, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ³H, ¹⁴C, ³²P, ³⁵S, or ¹²⁵I, a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., Nature, 144:945 (1962); David et al., Biochemistry, 13:1014 (1974); Pain et al., J. Immunol. Meth., 40:219 (1981); and Nygren, J. Histochem. and Cytochem., 30:407 (1982).

[0412] Anti-PRO antibodies also are useful for the affinity purification of PRO from recombinant cell culture or natural sources. In this process, the antibodies against PRO are immobilized on a suitable support, such as Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO from the antibody.

[0413] The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

[0414] All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

[0415] Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, Va.

Example 1

[0416] Extracellular Domain Homology Screening to Identify Novel Polypeptides and cDNA Encoding Therefor

[0417] The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search EST databases. The EST databases included public databases (e.g., Dayhoff, GenBank), and proprietary databases (e.g., LIFESEQ™, Incyte Pharmaceuticals, Palo Alto, Calif.). The search was performed using the computer program BLAST or BLAST-2 (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)) as a comparison of the ECD protein sequences to a 6 frame translation of the EST sequences. Those comparisons with a BLAST score of 70 (or in some cases 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Wash.).

[0418] Using this extracellular domain homology screen, consensus DNA sequences were assembled relative to the other identified EST sequences using phrap. In addition, the consensus DNA sequences obtained were often (but not always) extended using repeated cycles of BLAST or BLAST-2 and phrap to extend the consensus sequence as far as possible using the sources of EST sequences discussed above.

[0419] Based upon the consensus sequences obtained as described above, oligonucleotides were then synthesized and used to identify by PCR a cDNA library that contained the sequence of interest and for use as probes to isolate a clone of the full-length coding sequence for a PRO polypeptide. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

[0420] The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, Calif. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

Example 2

[0421] Isolation of cDNA Clones by Amylase Screening

[0422] 1. Preparation of Oligo dT Primed cDNA Library

[0423] mRNA was isolated from a human tissue of interest using reagents and protocols from Invitrogen, San Diego, Calif. (Fast Track 2). This RNA was used to generate an oligo dT primed cDNA library in the vector pRK5D using reagents and protocols from Life Technologies, Gaithersburg, Md. (Super Script Plasmid System). In this procedure, the double stranded cDNA was sized to greater than 1000 bp and the SalI/NotI tinkered cDNA was cloned into XhoI/NotI cleaved vector. pRK5D is a cloning vector that has an sp6 transcription initiation site followed by an SfiI restriction enzyme site preceding the XhoI/NotI cDNA cloning sites.

[0424] 2. Preparation of Random Primed cDNA Library

[0425] A secondary cDNA library was generated in order to preferentially represent the 5' ends of the primary cDNA clones. Sp6 RNA was generated from the primary library (described above), and this RNA was used to generate a random primed cDNA library in the vector pSST-AMY.0 using reagents and protocols from Life Technologies (Super Script Plasmid System, referenced above). In this procedure the double stranded cDNA was sized to 500-1000 bp, tinkered with blunt to NotI adaptors, cleaved with SfiI, and cloned into SfiI/NotI cleaved vector. pSST-AMY.0 is a cloning vector that has a yeast alcohol dehydrogenase promoter preceding the cDNA cloning sites and the mouse amylase sequence (the mature sequence without the secretion signal) followed by the yeast alcohol dehydrogenase terminator, after the cloning sites. Thus, cDNAs cloned into this vector that are fused in frame with amylase sequence will lead to the secretion of amylase from appropriately transfected yeast colonies.

[0426] 3. Transformation and Detection

[0427] DNA from the library described in paragraph 2 above was chilled on ice to which was added electrocompetent DH10B bacteria (Life Technologies, 20 ml). The bacteria and vector mixture was then electroporated as recommended by the manufacturer. Subsequently, SOC media (Life Technologies, 1 ml) was added and the mixture was incubated at 37° C. for 30 minutes. The transformants were then plated onto 20 standard 150 mm LB plates containing ampicillin and incubated for 16 hours (37° C.). Positive colonies were scraped off the plates and the DNA was isolated from the bacterial pellet using standard protocols, e.g. CsCl-gradient. The purified DNA was then carried on to the yeast protocols below.

[0428] The yeast methods were divided into three categories: (1) Transformation of yeast with the plasmid/cDNA combined vector; (2) Detection and isolation of yeast clones secreting amylase; and (3) PCR amplification of the insert directly from the yeast colony and purification of the DNA for sequencing and further analysis.

[0429] The yeast strain used was HD56-5A (ATCC-90785). This strain has the following genotype: MAT alpha, ura3-52, leu2-3, leu2-112, his3-11, his3-15, MAL⁺, SUC⁺, GAL⁺. Preferably, yeast mutants can be employed that have deficient post-translational pathways. Such mutants may have translocation deficient alleles in sec 71, sec 72, sec 62,

with truncated sec 71 being most preferred. Alternatively, antagonists (including antisense nucleotides and/or ligands) which interfere with the normal operation of these genes, other proteins implicated in this post translation pathway (e.g., SEC61p, SEC72p, SEC62p, SEC63p, TDJ1p or SSA1p-4p) or the complex formation of these proteins may also be preferably employed in combination with the amylase-expressing yeast.

[0430] Transformation was performed based on the protocol outlined by Gietz et al., Nucl. Acid. Res., 20:1425 (1992). Transformed cells were then inoculated from agar into YEPD complex media broth (100 ml) and grown overnight at 30° C. The YEPD broth was prepared as described in Kaiser et al., Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., p. 207 (1994). The overnight culture was then diluted to about 2×10^7 cells/ml (approx. OD₆₀₀=0.1) into fresh YEPD broth (500 ml) and regrown to 1×10^7 cells/ml (approx. OD₆₀₀=0.4-0.5).

[0431] The cells were then harvested and prepared for transformation by transfer into GS3 rotor bottles in a Sorval GS3 rotor at 5,000 rpm for 5 minutes, the supernatant discarded, and then resuspended into sterile water, and centrifuged again in 50 ml falcon tubes at 3,500 rpm in a Beckman GS-6KR centrifuge. The supernatant was discarded and the cells were subsequently washed with LiAc/TE (10 ml, 10 mM Tris-HCl, 1 mM EDTA pH 7.5, 100 mM Li₂OOCCH₃), and resuspended into LiAc/TE (2.5 ml).

[0432] Transformation took place by mixing the prepared cells (100 μ l) with freshly denatured single stranded salmon testes DNA (Lofstrand Labs, Gaithersburg, Md.) and transforming DNA (1 μ g, vol. <10 μ l) in microfuge tubes. The mixture was mixed briefly by vortexing, then 40% PEG/TE (600 μ l, 40% polyethylene glycol-4000, 10 mM Tris-HCl, 1 mM EDTA, 100 mM Li₂OOCCH₃, pH 7.5) was added. This mixture was gently mixed and incubated at 30° C. while agitating for 30 minutes. The cells were then heat shocked at 42° C. for 15 minutes, and the reaction vessel centrifuged in a microfuge at 12,000 rpm for 5-10 seconds, decanted and resuspended into TE (500 μ l, 10 mM Tris-HCl, 1 mM EDTA pH 7.5) followed by recentrifugation. The cells were then diluted into TE (1 ml) and aliquots (200 μ l) were spread onto the selective media previously prepared in 150 mm growth plates (VWR).

[0433] Alternatively, instead of multiple small reactions, the transformation was performed using a single, large scale reaction, wherein reagent amounts were scaled up accordingly.

[0434] The selective media used was a synthetic complete dextrose agar lacking uracil (SCD-Ura) prepared as described in Kaiser et al., Methods in Yeast Genetics, Cold Spring Harbor Press, Cold Spring Harbor, N.Y., p. 208-210 (1994). Transformants were grown at 30° C. for 2-3 days.

[0435] The detection of colonies secreting amylase was performed by including red starch in the selective growth media. Starch was couple d to the red dye (Reactive Red-120, Sigma) as per the procedure described by Biely et al., Anal. Biochem., 172:176-179 (1988). The couple d starch was incorporated into the SCD-Ura agar plates at a final

concentration of 0.15% (w/v), and was buffered with potassium phosphate to a pH of 7.0 (50-100 mM final concentration).

[0436] The positive colonies were picked and streaked across fresh selective media (onto 150 mm plates) in order to obtain well isolated and identifiable single colonies. Well isolated single colonies positive for amylase secretion were detected by direct incorporation of red starch into buffered SCD-Ura agar. Positive colonies were determined by their ability to break down starch resulting in a clear halo around the positive colony visualized directly.

[0437] 4. Isolation of DNA by PCR Amplification

[0438] When a positive colony was isolated, a portion of it was picked by a toothpick and diluted into sterile water (30 μ l) in a 96 well plate. At this time, the positive colonies were either frozen and stored for subsequent analysis or immediately amplified. An aliquot of cells (5 μ l) was used as a template for the PCR reaction in a 25 μ l volume containing: 0.5 μ l KlenTaq (Clontech, Palo Alto, Calif.); 4.0 μ l 10 mM dNTP's (Perkin Elmer-Cetus); 2.5 μ l Kentaq buffer (Clontech); 0.25 μ l forward oligo 1; 0.25 μ l reverse oligo 2; 12.5 μ l distilled water. The sequence of the forward oligonucleotide 1 was:

[0439] 5'-TGTAMACGACGGCCAGTTAMTA-GACCTGCMTTATTAATCT-3' (SEQ ID NO: 169)

[0440] The sequence of reverse oligonucleotide 2 was:

[0441] 5'-CAGGAMCAGCTATGACCACCTGCA-CACCTGCMATCCATT-3' (SEQ ID NO: 170)

[0442] PCR was then performed as follows:

a.		Denature	92° C.,	5 minutes
b.	3 cycles of:	Denature	92° C.,	30 seconds
		Anneal	59° C.,	30 seconds
		Extend	72° C.,	60 seconds
c.	3 cycles of:	Denature	92° C.,	30 seconds
		Anneal	57° C.,	30 seconds
		Extend	72° C.,	60 seconds
d.	25 cycles of:	Denature	92° C.,	30 seconds
		Anneal	55° C.,	30 seconds
		Extend	72° C.,	60 seconds
e.		Hold	4° C.	

[0443] The underlined regions of the oligonucleotides annealed to the ADH promoter region and the amylase region, respectively, and amplified a 307 bp region from vector pSST-AMY.0 when no insert was present. Typically, the first 18 nucleotides of the 5' end of these oligonucleotides contained annealing sites for the sequencing primers. Thus, the total product of the PCR reaction from an empty vector was 343 bp. However, signal sequence-fused cDNA resulted in considerably longer nucleotide sequences.

[0444] Following the PCR, an aliquot of the reaction (5 μ l) was examined by agarose gel electrophoresis in a 1% agarose gel using a Tris-Borate-EDTA (TBE) buffering system as described by Sambrook et al., supra. Clones resulting in a single strong PCR product larger than 400 bp were further analyzed by DNA sequencing after purification with a 96 Qiaquick PCR clean-up column (Qiagen Inc., Chatsworth, Calif.).

Example 3

[0445] Isolation of cDNA Clones Using Signal Algorithm Analysis

[0446] Various polypeptide-encoding nucleic acid sequences were identified by applying a proprietary signal sequence finding algorithm developed by Genentech, Inc. (South San Francisco, Calif.) upon ESTs as well as clustered and assembled EST fragments from public (e.g., GenBank) and/or private (LIFESeq®, Incyte Pharmaceuticals, Inc., Palo Alto, Calif.) databases. The signal sequence algorithm computes a secretion signal score based on the character of the DNA nucleotides surrounding the first and optionally the second methionine codon(s) (ATG) at the 5'-end of the sequence or sequence fragment under consideration. The nucleotides following the first ATG must code for at least 35 unambiguous amino acids without any stop codons. If the first ATG has the required amino acids, the second is not examined. If neither meets the requirement, the candidate sequence is not scored. In order to determine whether the EST sequence contains an authentic signal sequence, the DNA and corresponding amino acid sequences surrounding the ATG codon are scored using a set of seven sensors (evaluation parameters) known to be associated with secretion signals. Use of this algorithm resulted in the identification of numerous polypeptide-encoding nucleic acid sequences.

Example 4

[0447] Isolation of cDNA Clones Encoding Human PRO Polypeptides

[0448] Using the techniques described in Examples 1 to 3 above, numerous full-length cDNA clones were identified as encoding PRO polypeptides as disclosed herein. These cDNAs were then deposited under the terms of the Budapest Treaty with the American Type Culture Collection, 10801 University Blvd., Manassas, Va. 20110-2209, USA (ATCC) as shown in Tabl 7 below.

TABLE 7

Material	ATCC Dep. No.	Deposit Date
DNA26843-1389	203099	Aug. 4, 1998
DNA30867-1335	209807	Apr. 28, 1998
DNA34431-1177	209399	Oct. 17, 1997
DNA38268-1188	209421	Oct. 28, 1997
DNA40621-1440	209922	Jun. 2, 1998
DNA40625-1189	209788	Apr. 21, 1998
DNA45409-2511	203579	Jan. 12, 1999
DNA45495-1550	203156	Aug. 25, 1998
DNA49820-1427	209932	Jun. 2, 1998
DNA56406-1704	203478	Nov. 17, 1998
DNA56410-1414	209923	Jun. 2, 1998
DNA56436-1448	209902	May 27, 1998
DNA56855-1447	203004	Jun. 23, 1998
DNA56860-1510	209952	Jun. 9, 1998
DNA56862-1343	203174	Sep. 1, 1998
DNA56868-1478	203024	Jun. 23, 1998
DNA56869-1545	203161	Aug. 25, 1998
DNA57704-1452	209953	Jun. 9, 1998
DNA58723-1588	203133	Aug. 18, 1998
DNA57827-1493	203045	Jul. 1, 1998
DNA58737-1473	203136	Aug. 18, 1998
DNA58846-1409	209957	Jun. 9, 1998
DNA58850-1495	209956	Jun. 9, 1998
DNA58855-1422	203018	Jun. 23, 1998
DNA59211-1450	209960	Jun. 9, 1998

TABLE 7-continued

Material	ATCC Dep. No.	Deposit Date
DNA59212-1627	203245	Sep. 9, 1998
DNA59213-1487	209959	Jun. 9, 1998
DNA59605-1418	203005	Jun. 23, 1998
DNA59609-1470	209963	Jun. 9, 1998
DNA59610-1556	209990	Jun. 16, 1998
DNA59837-2545	203658	Feb. 9, 1999
DNA59844-2542	203650	Feb. 9, 1999
DNA59854-1459	209974	Jun. 16, 1998
DNA60625-1507	209975	Jun. 16, 1998
DNA60629-1481	209979	Jun. 16, 1998
DNA61755-1554	203112	Aug. 11, 1998
DNA62812-1594	203248	Sep. 9, 1998
DNA62815-1576	203247	Sep. 9, 1998
DNA64881-1602	203240	Sep. 9, 1998
DNA64886-1601	203241	Sep. 9, 1998
DNA64902-1667	203317	Oct. 6, 1998
DNA64950-1590	203224	Sep. 15, 1998
DNA65403-1565	203230	Sep. 15, 1998
DNA66308-1537	203159	Aug. 25, 1998
DNA66519-1535	203236	Sep. 15, 1998
DNA66521-1583	203225	Sep. 15, 1998
DNA66658-1584	203229	Sep. 15, 1998
DNA66660-1585	203279	Sep. 22, 1998
DNA66663-1598	203268	Sep. 22, 1998
DNA66674-1599	203281	Sep. 22, 1998
DNA68862-2546	203652	Feb. 9, 1999
DNA68866-1644	203283	Sep. 22, 1998
DNA68871-1638	203280	Sep. 22, 1998
DNA68880-1676	203319	Oct. 6, 1998
DNA68883-1691	203535	Dec. 15, 1998
DNA68885-1678	203311	Oct. 6, 1998
DNA71277-1636	203285	Sep. 22, 1998
DNA73727-1673	203459	Nov. 3, 1998
DNA73734-1680	203363	Oct. 20, 1998
DNA73735-1681	203356	Oct. 20, 1998
DNA76393-1664	203323	Oct. 6, 1998
DNA77301-1708	203407	Oct. 27, 1998
DNA77568-1626	203134	Aug. 18, 1998
DNA77626-1705	203536	Dec. 15, 1998
DNA81754-2532	203542	Dec. 15, 1998
DNA81757-2512	203543	Dec. 15, 1998
DNA82302-2529	203534	Dec. 15, 1998
DNA82340-2530	203547	Dec. 22, 1998
DNA83500-2506	203391	Oct. 29, 1998
DNA84920-2614	203966	Apr. 27, 1999
DNA85066-2534	203588	Jan. 12, 1999
DNA86571-2551	203660	Feb. 9, 1999
DNA87991-2540	203656	Feb. 9, 1999
DNA92238-2539	203602	Jan. 20, 1999
DNA96042-2682	PTA-382	Jul. 20, 1999
DNA96787-2534	203589	Jan. 12, 1999
DNA125185-2806	PTA-1031	Dec. 7, 1999
DNA147531-2821	PTA-1185	Jan. 11, 2000
DNA115291-2681	PTA-202	Jun. 8, 1999
DNA164625-28890	PTA-1535	Mar. 21, 2000
DNA131639-2874	PTA-1784	Apr. 25, 2000
DNA79230-2525	203549	Dec. 22, 1998

[0449] These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent

application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to 35 USC §122 and the Commissioner's rules pursuant thereto (including 37 CFR §1.14 with particular reference to 886 OG 638).

[0450] The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

Example 5

[0451] Use of PRO as a Hybridization Probe

[0452] The following method describes use of a nucleotide sequence encoding PRO as a hybridization probe.

[0453] DNA comprising the coding sequence of full-length or mature PRO as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO) in human tissue cDNA libraries or human tissue genomic libraries.

[0454] Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO-derived probe to the filters is performed in a solution of 50% formamide, 5×SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8, 2×Denhardt's solution, and 10% dextran sulfate at 42° C. for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1×SSC and 0.1% SDS at 42° C.

[0455] DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO can then be identified using standard techniques known in the art.

Example 6

[0456] Expression of PRO in *E. coli*

[0457] This example illustrates preparation of an unglycosylated form of PRO by recombinant expression in *E. coli*.

[0458] The DNA sequence encoding PRO is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO coding region, lambda transcriptional terminator, and an argU gene.

[0459] The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., *supra*. Transformants are identified by their

ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

[0460] Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

[0461] After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

[0462] PRO may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) Ion galE rpoHts (htpRts) clpP(lacIq)). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30° C. with shaking until an O.D.600 of 3-5 is reached. Cultures are then diluted 50-100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate-2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield hycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20-30 hours at 30° C. with shaking. Samples are removed to verify expression by SDS-page analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

[0463] *E. coli* paste from 0.5 to 1 L fermentations (6-10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1 M and 0.02 M, respectively, and the solution is stirred overnight at 4° C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3-5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Ultrapure grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4° C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

[0464] The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM

cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4° C. for 12-36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2-10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

[0465] Fractions containing the desired folded PRO polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

[0466] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 7

[0467] Expression of PRO in Mammalian Cells

[0468] This example illustrates preparation of a potentially glycosylated form of PRO by recombinant expression in mammalian cells.

[0469] The vector, pRK5 (see EP 307,247, published Mar. 15, 1989), is employed as the expression vector. Optionally, the PRO DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO DNA using ligation methods such as described in Sambrook et al., *supra*. The resulting vector is called pRK5-PRO.

[0470] In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 μ g pRK5-PRO DNA is mixed with about 1 μ g DNA encoding the VA RNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 μ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 μ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25° C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37° C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

[0471] Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture

medium (alone) or culture medium containing 200 μ Ci/ml ³⁵S-cysteine and 200 μ Ci/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

[0472] In an alternative technique, PRO may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 μ g pRK5-PRO DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 μ g/ml bovine insulin and 0.1 μ g/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

[0473] In another embodiment, PRO can be expressed in CHO cells. The pRK5-PRO can be transfected into CHO cells using known reagents such as CaPO₄ or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ³⁵S-methionine. After determining the presence of PRO polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO can then be concentrated and purified by any selected method.

[0474] Epitope-tagged PRO may also be expressed in host CHO cells. The PRO may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the expressed poly-His tagged PRO can then be concentrated and purified by any selected method, such as by Ni²⁺-chelate affinity chromatography.

[0475] PRO may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

[0476] Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

[0477] Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using

standard techniques as described in Ausubel et al., *Current Protocols of Molecular Biology*, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., *Nucl. Acids Res.* 24:9 (1774-1779) (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

[0478] Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Quiagen), Dosper® or Eugene® (Boehringer Mannheim). The cells are grown as described in Lucas et al., *supra*. Approximately 3×10^7 cells are frozen in an ampule for further growth and production as described below.

[0479] The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mLs of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2 μ m filtered PS20 with 5% 0.2 μ m diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37° C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with 3×10^5 cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Pat. No. 5,122,469, issued Jun. 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2×10^6 cells/mL. On day 0, the cell number pH is determined. On day 1, the spinner is sample d and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33° C., and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane emulsion, Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate was either stored at 4° C. or immediately loaded onto columns for purification.

[0480] For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 mL Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4, buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4-5 mL/min. at 4° C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 mL G25 Superfine (Pharmacia) column and stored at -80° C.

[0481] Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The condi-

tioned medium is pumped onto a 5 mL Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 mL fractions into tubes containing 275 μ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

[0482] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 5

[0483] Expression of PRO in Yeast

[0484] The following method describes recombinant expression of PRO in yeast.

[0485] First, yeast expression vectors are constructed for intracellular production or secretion of PRO from the ADH2/GAPDH promoter. DNA encoding PRO and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO. For secretion, DNA encoding PRO can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO signal peptide or other mammalian signal peptide, or, for example, a yeast alpha-factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO.

[0486] Yeast cells, such as yeast strain AB 110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

[0487] Recombinant PRO can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO may further be purified using selected column chromatography resins.

[0488] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 9

[0489] Expression of PRO in Baculovirus-Infected Insect Cells

[0490] The following method describes recombinant expression of PRO in Baculovirus-infected insect cells.

[0491] The sequence coding for PRO is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO or the desired portion of the coding sequence of PRO such as the sequence encoding the extracellular domain of a transmem-

brane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5' primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

[0492] Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4-5 days of incubation at 28° C., the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilley et al., *Baculovirus expression vectors: A Laboratory Manual*, Oxford: Oxford University Press (1994).

[0493] Expressed poly-his tagged PRO can then be purified, for example, by Ni²⁺-chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., *Nature*, 362:175-179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mL Hepes, pH 7.9; 12.5 mM MgCl₂; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 μm filter. A Ni²⁺-NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A₂₈₀ with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A₂₈₀ baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni²⁺-NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His₁₀-tagged PRO are pooled and dialyzed against loading buffer.

[0494] Alternatively, purification of the IgG tagged (or Fc tagged) PRO can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

[0495] Many of the PRO polypeptides disclosed herein were successfully expressed as described above.

Example 10

[0496] Preparation of Antibodies that Bind PRO

[0497] This example illustrates preparation of monoclonal antibodies which can specifically bind PRO.

[0498] Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, *supra*. Immunogens that may be employed include purified PRO, fusion proteins containing PRO, and cells expressing recombinant PRO on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

[0499] Mice, such as Balb/c, are immunized with the PRO immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1-100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, Mont.) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO antibodies.

[0500] After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

[0501] The hybridoma cells will be screened in an ELISA for reactivity against PRO. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO is within the skill in the art.

[0502] The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

Example 11

[0503] Purification of PRO Polypeptides Using Specific Antibodies

[0504] Native or recombinant PRO polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO polypeptide, mature PRO polypeptide, or pre-PRO polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO polypeptide antibody to an activated chromatographic resin.

[0505] Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotech-

nology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

[0506] Such an immunoaffinity column is utilized in the purification of PRO polypeptide by preparing a fraction from cells containing PRO polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

[0507] A soluble PRO polypeptide-containing preparation is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PRO polypeptide (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO polypeptide binding (e.g., a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO polypeptide is collected.

Example 12

[0508] Drug Screening

[0509] This invention is particularly useful for screening compounds by using PRO polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO polypeptide and its target cell or target receptors caused by the agent being tested.

[0510] Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO polypeptide-associated disease or disorder. These methods comprise contacting such an agent with a PRO polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO polypeptide or fragment, or (ii) for the presence of a complex between the PRO polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO polypeptide or fragment is typically labeled. After suitable incubation, free PRO polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO polypeptide or to interfere with the PRO polypeptide/cell complex.

[0511] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on Sep. 13, 1984. Briefly stated,

large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO polypeptide, the peptide test compounds are reacted with PRO polypeptide and washed. Bound PRO polypeptide is detected by methods well known in the art. Purified PRO polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

[0512] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO polypeptide specifically compete with a test compound for binding to PRO polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO polypeptide.

Example 13

[0513] Rational Drug Design

[0514] The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (i.e., a PRO polypeptide) or of small molecules with which they interact, e.g., agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO polypeptide or which enhance or interfere with the function of the PRO polypeptide in vivo (cf, Hodgson, *Bio/Technology*, 9: 19-21(1991)).

[0515] In one approach, the three-dimensional structure of the PRO polypeptide, or of a PRO polypeptide-inhibitor complex, is determined by X-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, *Biochemistry*, 31:7796-7801(1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda et al., *J. Biochem.*, 113:742-746(1993).

[0516] It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

[0517] By virtue of the present invention, sufficient amounts of the PRO polypeptide may be made available to

perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to X-ray crystallography.

Example 14

[0518] Pericyte c-Fos Induction (Assay 93)

[0519] This assay shows that certain polypeptides of the invention act to induce the expression of c-fos in pericyte cells and, therefore, are useful not only as diagnostic markers for particular types of pericyte-associated tumors but also for giving rise to antagonists which would be expected to be useful for the therapeutic treatment of pericyte-associated tumors. Induction of c-fos expression in pericytes is also indicative of the induction of angiogenesis and, as such, PRO polypeptides capable of inducing the expression of c-fos would be expected to be useful for the treatment of conditions where induced angiogenesis would be beneficial including, for example, wound healing, and the like. Specifically, on day 1, pericytes are received from VEC Technologies and all but 5 ml of media is removed from flask. On day 2, the pericytes are trypsinized, washed, spun and then plated onto 96 well plates. On day 7, the media is removed and the pericytes are treated with 100 μ l of PRO polypeptide test samples and controls (positive control=DME+5% serum+/-PDGF at 500 ng/ml; negative control=protein 32). Replicates are averaged and SD/CV are determined. Fold increase over Protein 32 (buffer control) value indicated by chemiluminescence units (RLU) luminometer reading verses frequency is plotted on a histogram. Two-fold above Protein 32 value is considered positive for the assay. ASY Matrix: Growth media=low glucose DMEM=20% FBS+1 $\times$ pen strep+1 $\times$ fungizone. Assay Media=low glucose DMEM+5% FBS.

[0520] The following polypeptides tested positive in this assay: PRO1347 and PRO1340.

Example 15

[0521] Ability of PRO Polypeptides to Stimulate the Release of Proteoglycans from Cartilage (Assay 97)

[0522] The ability of various PRO polypeptides to stimulate the release of proteoglycans from cartilage tissue was tested as follows.

[0523] The metacarpophalangeal joint of 4-6 month old pigs was aseptically dissected, and articular cartilage was removed by free hand slicing being careful to avoid the underlying bone. The cartilage was minced and cultured in bulk for 24 hours in a humidified atmosphere of 95% air, 5% CO₂ in serum free (SF) media (DME/F12 1:1) with 0.1% BSA and 100 U/ml penicillin and 100 μ g/ml streptomycin. After washing three times, approximately 100 mg of articular cartilage was aliquoted into micronics tubes and incubated for an additional 24 hours in the above SF media. PRO polypeptides were then added at 1% either alone or in combination with 18 ng/ml interleukin-1 α , a known stimulator of proteoglycan release from cartilage tissue. The supernatant was then harvested and assayed for the amount of proteoglycans using the 1,9-dimethyl-methylene blue (DMB) colorimetric assay (Farndale and Buttle, Biochem. Biophys. Acta 883:173-177 (1985)). A positive result in this

assay indicates that the test polypeptide will find use, for example, in the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis.

[0524] When various PRO polypeptides were tested in the above assay, the polypeptides demonstrated a marked ability to stimulate release of proteoglycans from cartilage tissue both basally and after stimulation with interleukin-1 α and at 24 and 72 hours after treatment, thereby indicating that these PRO polypeptides are useful for stimulating proteoglycan release from cartilage tissue. As such, these PRO polypeptides are useful for the treatment of sports-related joint problems, articular cartilage defects, osteoarthritis or rheumatoid arthritis. The polypeptides testing positive in this assay are: PRO1565, PRO1693, PRO1801 and PRO10096.

Example 16

[0525] Detection of Polypeptides That Affect Glucose or FFA Uptake in Skeletal Muscle (Assay 106)

[0526] This assay is designed to determine whether PRO polypeptides show the ability to affect glucose or FFA uptake by skeletal muscle cells. PRO polypeptides testing positive in this assay would be expected to be useful for the therapeutic treatment of disorders where either the stimulation or inhibition of glucose uptake by skeletal muscle would be beneficial including, for example, diabetes or hyper- or hypo-insulinemia.

[0527] In a 96 well format, PRO polypeptides to be assayed are added to primary rat differentiated skeletal muscle, and allowed to incubate overnight. Then fresh media with the PRO polypeptide and +/- insulin are added to the wells. The sample media is then monitored to determine glucose and FFA uptake by the skeletal muscle cells. The insulin will stimulate glucose and FFA uptake by the skeletal muscle, and insulin in media without the PRO polypeptide is used as a positive control, and a limit for scoring. As the PRO polypeptide being tested may either stimulate or inhibit glucose and FFA uptake, results are scored as positive in the assay if greater than 1.5 times or less than 0.5 times the insulin control.

[0528] The following PRO polypeptides tested positive as either stimulators or inhibitors of glucose and/or FFA uptake in this assay: PRO4405.

Example 17

[0529] Identification of PRO Polypeptides that Stimulate TNF- α or Release in Human Blood (Assay 128)

[0530] This assay shows that certain PRO polypeptides of the present invention act to stimulate the release of TNF- α in human blood. PRO polypeptides testing positive in this assay are useful for, among other things, research purposes where stimulation of the release of TNF- α would be desired and for the therapeutic treatment of conditions wherein enhanced TNF- α release would be beneficial. Specifically, 200 μ l of human blood supplemented with 50 mM Hepes buffer (pH 7.2) is aliquotted per well in a 96 well test plate. To each well is then added 300 μ l of either the test PRO polypeptide in 50 mM Hepes buffer (at various concentrations) or 50 mM Hepes buffer alone (negative control) and the plates are incubated at 37° C. for 6 hours. The samples are then centrifuged and 50 μ l of plasma is collected from

each well and tested for the presence of TNF- α by ELISA assay. A positive in the assay is a higher amount of TNF- α in the PRO polypeptide treated samples as compared to the negative control samples.

[0531] The following PRO polypeptides tested positive in this assay: PRO263, PRO295, PRO1282, PRO1063, PRO1356, PRO3543, and PRO5990.

Example 18

[0532] Tumor Versus Normal Differential Tissue Expression Distribution

[0533] Oligonucleotide probes were constructed from some of the PRO polypeptide-encoding nucleotide sequences shown in the accompanying figures for use in quantitative PCR amplification reactions. The oiouecode probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human tumor and normal human tissue samples and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO polypeptide-encoding nucleic acid in the various tumor and normal tissues tested. β -actin was used as a control to assure that equivalent amounts of nucleic acid was used in each reaction. Identification of the differential expression of the PRO polypeptide-encoding nucleic acid in one or more tumor tissues as compared to one or more normal tissues of the same tissue type renders the molecule useful diagnostically for the determination of the presence or absence of tumor in a subject suspected of possessing a tumor as well as therapeutically as a target for the treatment of a tumor in a subject possessing such a tumor. These assays provided the following results.

Molecule	is more highly expressed in:	as compared to:
DNA26843-1389	normal lung	lung tumor
	rectum tumor	normal rectum
DNA30867-1335	normal kidney	kidney tumor
DNA40621-1440	normal lung	lung tumor
DNA40625-1189	normal lung	lung tumor
DNA45409-2511	melanoma tumor	normal akin
DNA56406-1704	kidney tumor	normal kidney
	normal skin	melanoma tumor
DNA56410-1414	normal stomach	stomach tumor
DNA56436-1448	normal skin	melanoma tumor
DNA56855-1447	normal esophagus	esophageal tumor
	rectum tumor	normal rectum
DNA56860-1510	normal kidney	kidney tumor
	rectum tumor	normal rectum
DNA56862-1343	kidney tumor	normal kidney
	normal lung	lung tumor
DNA56868-1478	normal stomach	stomach tumor
	normal lung	lung tumor
DNA56869-1545	normal esophagus	esophageal tumor
	normal skin	melanoma tumor
DNA57704-1452	normal stomach	stomach tumor
	rectum tumor	normal rectum
DNA58723-1588	normal stomach	stomach tumor
	kidney tumor	normal kidney
	normal akin	melanoma tumor
DNA57827-1493	normal stomach	stomach tumor
	normal skin	melanoma tumor
DNA58737-1473	esophageal tumor	normal esophagus
	normal stomach	stomach tumor

-continued

Molecule	is more highly expressed in:	as compared to:
DNA58846-1409	lung tumor	normal lung
DNA58850-1495	esophageal tumor	normal esophagus
	kidney tumor	normal kidney
DNA58855-1422	normal stomach	stomach tumor
	rectum tumor	normal rectum
DNA59211-1450	normal kidney	kidney tumor
DNA59212-1627	normal skin	melanoma tumor
DNA59213-1487	normal stomach	stomach tumor
	normal skin	melanoma tumor
DNA59605-1418	melanoma tumor	normal skin
DNA59609-1470	esophageal tumor	normal esophagus
DNA59610-1556	esophageal tumor	normal esophagus
	lung tumor	normal lung
	normal skin	melanoma tumor
DNA59837-2545	normal skin	melanoma tumor
DNA59844-2542	normal skin	melanoma tumor
	esophageal tumor	normal esophagus
DNA59854-1459	normal esophagus	esophageal tumor
	stomach tumor	normal stomach
	normal lung	lung tumor
DNA60625-1507	normal lung	lung tumor
DNA60629-1481	normal esophagus	esophageal tumor
	normal rectum	rectum tumor
DNA61755-1554	normal stomach	stomach tumor
	kidney tumor	normal kidney
DNA62812-1594	normal stomach	stomach tumor
	normal lung	lung tumor
	normal rectum	rectum tumor
	normal skin	melanoma tumor
DNA62815-1576	esophageal tumor	normal esophagus
DNA64881-1602	normal stomach	stomach tumor
	normal lung	lung tumor
DNA64902-1667	esophageal tumor	normal esophagus
	kidney tumor	normal kidney
DNA65403-1565	normal esophagus	esophageal tumor
DNA66308-1537	normal lung	lung tumor
DNA66519-1535	kidney tumor	normal kidney
DNA66521-1583	normal esophagus	esophageal tumor
	normal stomach	stomach tumor
	normal lung	lung tumor
	normal rectum	rectum tumor
	normal skin	melanoma tumor
DNA66658-1584	normal lung	lung tumor
	melanoma tumor	normal skin
DNA66660-1585	lung tumor	normal lung
DNA66674-1599	kidney tumor	normal kidney
	normal lung	lung tumor
DNA68862-2546	melanoma tumor	normal skin
DNA68866-1644	normal stomach	stomach tumor
DNA68871-1638	lung tumor	normal lung
	normal skin	melanoma tumor
DNA68880-1676	normal lung	lung tumor
	normal skin	melanoma tumor
DNA68883-1691	esophageal tumor	normal esophagus
DNA68885-1678	lung tumor	normal lung
DNA71277-1636	normal stomach	stomach tumor
DNA73734-1680	normal lung	lung tumor
DNA73735-1681	esophageal tumor	normal esophagus
	normal kidney	kidney tumor
	lung tumor	normal lung
	normal skin	melanoma tumor
DNA76393-1664	esophageal tumor	normal esophagus
	stomach tumor	normal stomach
	lung tumor	normal lung
	rectum tumor	normal rectum
DNA77568-1626	normal stomach	stomach tumor
	lung tumor	normal lung
DNA77626-1705	normal rectum	rectum tumor
DNA81754-2532	normal skin	melanoma tumor
DNA81757-2512	esophageal tumor	normal esophagus
	normal stomach	stomach tumor
	melanoma tumor	normal skin
DNA82302-2529	normal stomach	stomach tumor
	normal lung	lung tumor
DNA82340-2530	normal esophagus	esophageal tumor

-continued		
Molecule	is more highly expressed in:	as compared to:
DNA85066-2534	lung tumor normal skin	normal lung melanoma tumor
DNA87991-2540	esophageal tumor	normal esophagus
DNA92238-2539	normal skin	melanoma tumor
DNA96787-2534	normal kidney	kidney tumor

Example 19

[0534] Identification of Receptor/Ligand Interactions

[0535] In this assay, various PRO polypeptides are tested for ability to bind to a panel of potential receptor or ligand molecules for the purpose of identifying receptor/ligand interactions. The identification of a ligand for a known receptor, a receptor for a known ligand or a novel receptor/ligand pair is useful for a variety of indications including, for example, targeting bioactive molecules (linked to the ligand or receptor) to a cell known to express the receptor or ligand, use of the receptor or ligand as a reagent to detect the presence of the ligand or receptor in a composition suspected of containing the same, wherein the composition may comprise cells suspected of expressing the ligand or receptor, modulating the growth of or another biological or immunological activity of a cell known to express or respond to the receptor or ligand, modulating the immune response of cells or toward cells that express the receptor or ligand, allowing the preparation of agonists, antagonists and/or antibodies directed against the receptor or ligand which will modulate the growth of or a biological or immunological activity of a cell expressing the receptor or ligand, and various other indications which will be readily apparent to the ordinarily skilled artisan.

[0536] The assay is performed as follows. A PRO polypeptide of the present invention suspected of being a ligand for a receptor is expressed as a fusion protein containing the Fc domain of human IgG (an immunoadhesin). Receptor-ligand binding is detected by allowing interaction of the immunoadhesin polypeptide with cells (e.g. Cos cells) expressing candidate PRO polypeptide receptors and visualization of bound immunoadhesin with fluorescent reagents directed toward the Fc fusion domain and examination by microscope. Cells expressing candidate receptors are produced by transient transfection, in parallel, of defined subsets of a library of cDNA expression vectors encoding PRO polypeptides that may function as receptor molecules. Cells are then incubated for 1 hour in the presence of the PRO polypeptide immunoadhesin being tested for possible receptor binding. The cells are then washed and fixed with paraformaldehyde. The cells are then incubated with fluorescent conjugated antibody directed against the Fc portion of the PRO polypeptide immunoadhesin (e.g. FITC conju-

gated goat anti-human-Fc antibody). The cells are then washed again and examined by microscope. A positive interaction is judged by the presence of fluorescent labeling of cells transfected with cDNA encoding a particular PRO polypeptide receptor or pool of receptors and an absence of similar fluorescent labeling of similarly prepared cells that have been transfected with other cDNA or pools of cDNA. If a defined pool of cDNA expression vectors is judged to be positive for interaction with a PRO polypeptide immunoadhesin, the individual cDNA species that comprise the pool are tested individually (the pool is "broken down") to determine the specific cDNA that encodes a receptor able to interact with the PRO polypeptide immunoadhesin.

[0537] In another embodiment of this assay, an epitope-tagged potential ligand PRO polypeptide (e.g. 8 histidine "His" tag) is allowed to interact with a panel of potential receptor PRO polypeptide molecules that have been expressed as fusions with the Fc domain of human IgG (immunoadhesins). Following a 1 hour co-incubation with the epitope tagged PRO polypeptide, the candidate receptors are each immunoprecipitated with protein A beads and the beads are washed. Potential ligand interaction is determined by western blot analysis of the immunoprecipitated complexes with antibody directed towards the epitope tag. An interaction is judged to occur if a band of the anticipated molecular weight of the epitope tagged protein is observed in the western blot analysis with a candidate receptor, but is not observed to occur with the other members of the panel of potential receptors.

[0538] Using these assays, the following receptor/ligand interactions have been herein identified:

- [0539] (1) PRO10272 binds to PRO5801.
- [0540] (2) PRO20110 binds to the human IL-17 receptor (Yao et al., Cytokine 9(11):794-800 (1997); also herein designated as PRO1) and to PRO20040.
- [0541] (3) PRO10096 binds to PRO20233.
- [0542] (4) PRO19670 binds to PRO1890.

[0543] The foregoing written specification is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the construct deposited, since the deposited embodiment is intended as a single illustration of certain aspects of the invention and any constructs that are functionally equivalent are within the scope of this invention. The deposit of material herein does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustrations that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 170

<210> SEQ ID NO 1
<211> LENGTH: 1173
<212> TYPE: DNA

-continued

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 1

```

ggggcttcgg cgccagcggc cagcgctagt cggctctgta aggatttaca      50
aaaggtgcag gtagagcag gtctgaagac taacattttg tgaagttgta      100
aaacagaaaa cctgttagaa atgtgggtgtt ttcagcaagg cctcagtttc      150
cttccttcag cccttgtaat ttggacatct gctgctttca tattttcata      200
cattactgca gtaacactcc accatataga cccggcttta ccttatatca      250
gtgacactgg tacagtagct ccagaaaaat gcttatttgg ggcaatgcta      300
aatattgcgg cagttttatg cattgctacc atttatgttc gttataagca      350
agttcatgct ctgagtcctg aagagaacgt tatcatcaaa ttaacaagg      400
ctggccttgt acttgaata ctgagttgtt taggactttc tattgtggca      450
aacttccaga aaacaaccct ttttgctgca catgtaagtg gagctgtgct      500
taccttttgt atgggctcat tatatatgtt tgttcagacc atcctttcct      550
accaaagtca gcccaaaatc catggcaaac aagtcttctg gatcagactg      600
ttgttggtta tctggtgtgg agtaagtgca cttagcatgc tgacttgctc      650
atcagttttg cacagtggca attttgggac tgatttagaa cagaaactcc      700
attggaaccg cgaggacaaa ggttatgtgc ttcacatgat cactactgca      750
gcagaatggt ctatgtcatt ttccttcttt ggttttttcc tgactttacat      800
tcgtgatttt cagaaaattt ctttacgggt ggaagccaat ttacatggat      850
taacctctta tgacactgca ccttgccta ttaacaatga acgaacacgg      900
ctactttcca gagatatattg atgaaaggat aaaatatttc tgtaatgatt      950
atgattctca gggattgggg aaaggttcac agaagttgct tattcttctc     1000
tgaaattttc aaccacttaa tcaaggctga cagtaacact gatgaatgct     1050
gataatcagg aaacatgaaa gaagccattt gatagattat tctaaaggat     1100
atcatcaaga agactattaa aaacacctat gcctatactt ttttatctca     1150
gaaaataaag tcaaaagact atg                                     1173

```

<210> SEQ ID NO 2

<211> LENGTH: 266

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 2

```

Met Trp Trp Phe Gln Gln Gly Leu Ser Phe Leu Pro Ser Ala Leu
 1             5             10            15
Val Ile Trp Thr Ser Ala Ala Phe Ile Phe Ser Tyr Ile Thr Ala
 20            25            30
Val Thr Leu His His Ile Asp Pro Ala Leu Pro Tyr Ile Ser Asp
 35            40            45
Thr Gly Thr Val Ala Pro Glu Lys Cys Leu Phe Gly Ala Met Leu
 50            55            60
Asn Ile Ala Ala Val Leu Cys Ile Ala Thr Ile Tyr Val Arg Tyr
 65            70            75
Lys Gln Val His Ala Leu Ser Pro Glu Glu Asn Val Ile Ile Lys

```

-continued

80					85					90				
Leu	Asn	Lys	Ala	Gly	Leu	Val	Leu	Gly	Ile	Leu	Ser	Cys	Leu	Gly
				95					100					105
Leu	Ser	Ile	Val	Ala	Asn	Phe	Gln	Lys	Thr	Thr	Leu	Phe	Ala	Ala
				110					115					120
His	Val	Ser	Gly	Ala	Val	Leu	Thr	Phe	Gly	Met	Gly	Ser	Leu	Tyr
				125					130					135
Met	Phe	Val	Gln	Thr	Ile	Leu	Ser	Tyr	Gln	Met	Gln	Pro	Lys	Ile
				140					145					150
His	Gly	Lys	Gln	Val	Phe	Trp	Ile	Arg	Leu	Leu	Leu	Val	Ile	Trp
				155					160					165
Cys	Gly	Val	Ser	Ala	Leu	Ser	Met	Leu	Thr	Cys	Ser	Ser	Val	Leu
				170					175					180
His	Ser	Gly	Asn	Phe	Gly	Thr	Asp	Leu	Glu	Gln	Lys	Leu	His	Trp
				185					190					195
Asn	Pro	Glu	Asp	Lys	Gly	Tyr	Val	Leu	His	Met	Ile	Thr	Thr	Ala
				200					205					210
Ala	Glu	Trp	Ser	Met	Ser	Phe	Ser	Phe	Phe	Gly	Phe	Phe	Leu	Thr
				215					220					225
Tyr	Ile	Arg	Asp	Phe	Gln	Lys	Ile	Ser	Leu	Arg	Val	Glu	Ala	Asn
				230					235					240
Leu	His	Gly	Leu	Thr	Leu	Tyr	Asp	Thr	Ala	Pro	Cys	Pro	Ile	Asn
				245					250					255
Asn	Glu	Arg	Thr	Arg	Leu	Leu	Ser	Arg	Asp	Ile				
				260					265					

<210> SEQ ID NO 3
<211> LENGTH: 2037
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 3

cggaacgcgtg ggcggacgcg tgggggagag ccgcagtcgcc ggtgcagca	50
cctgggagaa ggcagaccgt gtgagggggc ctgtggcccc agcgtgctgt	100
ggcctcgggg agtgggaagt ggaggcagga gccttcctta cacttcgccca	150
tgagtttcct catcgactcc agcatcatga ttacctccca gatactattt	200
tttggaattg ggtggctttt cttcatgcgc caattgttta aagactatga	250
gatacgtcag tatgttgtac aggtgatctt ctccgtgacg ttgcatttt	300
cttgacccat gtttgagctc atcatctttg aaatcctagg agtattgaat	350
agcagctccc gttattttca ctggaaaatg aacctgtgtg taattctgct	400
gatcctggtt ttcattgtgc ctttttacat tggctatttt attgtgagca	450
atatccgact actgcataaa caacgactgc ttttttcctg tctcttatgg	500
ctgaccttta tgtatttcct ctggaaacta ggagatccct ttcccattct	550
cagcccaaaa catgggatct tatccataga acagctcatc agccgggttg	600
gtgtgattgg agtgactctc atggctcttc tttctggatt tgggtctgtc	650
aactgcccat acacttacat gtcttacttc ctccaggaatg tgactgacac	700
ggatattcta gccctggaac ggcgactgct gcaaaccatg gatatgatca	750

-continued

taagcaaaaa gaaaaggatg gcaatggcac ggagaacaat gttccagaag	800
ggggaagtgc ataacaacc atcaggtttc tggggaatga taaaaagtgt	850
taccacttca gcatcaggaa gtgaaaatct tactcttatt caacaggaag	900
tggaatgcttt ggaagaatta agcaggcagc tttttctgga aacagctgat	950
ctatatgcta ccaaggagag aatagaatac tccaaaacct tcaaggggaa	1000
atatttttaat tttcttggtt actttttctc tatttactgt gtttgaaaa	1050
ttttcatggc taccatcaat attgtttttg atcgagttgg gaaaacggat	1100
cctgtcacaa gaggcattga gatcactgtg aattatctgg gaatccaatt	1150
tgatgtgaag ttttgggtccc aacacatttc cttcattctt gttggaataa	1200
tcacgtcac atccatcaga ggattgtga tcaactctac caagttcttt	1250
tatgccatct ctacgagtaa gtctccaat gtcattgtcc tgctattagc	1300
acagataatg ggcattgtact ttgtctcttc tgtgctgctg atccgaatga	1350
gtatgccttt agaataccgc accataatca ctgaagtcct tggagaactg	1400
cagttcaact tctatcaccg ttggtttgat gtgatcttcc tggtcagcgc	1450
tctctctagc atactcttcc tctatttggc tcacaaacag gcaccagaga	1500
agcaaatggc accttgaact taagcctact acagactgtt agaggccagt	1550
ggtttcaaaa tttagatata agagggggga aaaatggaac cagggcctga	1600
cattttataa acaacaacaa tgctatggta gcatttttca cttcatagc	1650
atactccttc ccggtcaggt gatactatga coatgagtag catcagccag	1700
aacatgagag ggagaactaa ctcaagacaa tactcagcag agagcatccc	1750
gtgtggatat gaggctgggtg tagaggcgga gaggagccaa gaaactaaag	1800
gtgaaaaata cactggaact ctggggcaag acatgtctat ggtagctgag	1850
ccaaacacgt aggatttcog ttttaagggt cacatggaaa aggttatagc	1900
tttgccctga gattgactca ttaaaatcag agactgtaac aaaaaaaaaa	1950
aaaaaaaaa agggcgggccg cgactctaga gtcgacctgc agaagcttgg	2000
ccgccatggc ccaacttggt tattgcagct tataatg	2037

<210> SEQ ID NO 4
<211> LENGTH: 455
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 4

Met Ser Phe Leu Ile Asp Ser Ser Ile Met Ile Thr Ser Gln Ile	
1 5 10 15	
Leu Phe Phe Gly Phe Gly Trp Leu Phe Phe Met Arg Gln Leu Phe	
20 25 30	
Lys Asp Tyr Glu Ile Arg Gln Tyr Val Val Gln Val Ile Phe Ser	
35 40 45	
Val Thr Phe Ala Phe Ser Cys Thr Met Phe Glu Leu Ile Ile Phe	
50 55 60	
Glu Ile Leu Gly Val Leu Asn Ser Ser Ser Arg Tyr Phe His Trp	
65 70 75	
Lys Met Asn Leu Cys Val Ile Leu Leu Ile Leu Val Phe Met Val	

-continued

<210> SEQ ID NO 5
<211> LENGTH: 2372
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 5

agcagggaaa tccggatgtc tcggttatga agtggagcag tgagtgtgag	50
cctcaacata gttccagaac tctccatccg gactagttag tgagcatctg	100
cctctcatat caccagtggc catctgaggt gtttccctgg ctctgaagg	150
gtaggcacga tggccaggtg cttcagcctg gtgttgcttc tcacttccat	200
ctggaccacg aggtctctgg tccaaggctc tttgcgtgca gaagagcttt	250
ccatccaggt gtcatgcaga attatgggga tcacccttgt gagcaaaaag	300
gcgaaccagc agctgaattt cacagaagct aaggaggcct gtaggctgct	350
gggactaagt ttggccggca aggaccaagt tgaacagcc ttgaaagcta	400
gctttgaaac ttgcagctat ggctgggttg gagatggatt cgtggctcat	450
tctaggatta gcccaaaccc caagtgtggg aaaaatgggg tgggtgtcct	500
gatttgaag gttccagtga gccgacagtt tgcagcctat tgttacaact	550
catctgatac ttggactaac tcgtgcattc cagaaattat caccacaaa	600
gatcccatat tcaaacactc aactgcaaca caaacacag aattttattgt	650
cagtgcagct acctactcgg tggcatcccc ttactctaca atacctgccc	700
ctactactac tcctcctgct ccagcttcca cttctattcc acggagaaaa	750
aaattgattt gtgtcacaga agtttttatg gaaactagca ccatgtctac	800
agaaactgaa ccatttggtt aaaataaagc agcattcaag aatgaagctg	850
ctgggtttgg aggtgtcccc acggctctgc tagtgcttgc tctcctcttc	900
tttggtgctg cagctggtct tggattttgc tatgtcaaaa ggtatgtgaa	950
ggccttccct tttacaaaca agaatcagca gaaggaaatg atcgaaacca	1000
aagtagtaaa ggaggagaag gccaatgata gcaaccctaa tgaggaaatca	1050
aagaaaactg ataaaaaccc agaagagtcc aagagtccaa gcaaaactac	1100
cgtgcgatgc ctggaagctg aagtttagat gagacagaaa tgaggagaca	1150
cacctgaggc tggtttcttt catgctcctt accctgcccc agctggggaa	1200
atcaaaaggg ccaagaagcc aaagaagaaa gtccaccctt ggttcctaac	1250
tggaatcagc tcaggactgc cattggacta tggagtgcac caaagagaat	1300
gcccttctcc ttattgtaac cctgtctgga tcctatcctc ctacctccaa	1350
agcttccccg ggcctttcta gcctggctat gtcctaataa tatcccactg	1400
ggagaaagga gttttgcaaa gtgcaaggac ctaaaacatc tcatcagtat	1450
ccagtggtaa aaaggcctcc tggctgtctg aggctagggt ggttgaaagc	1500
caaggagtca ctgagaccaa ggctttctct actgattccg cagctcagac	1550
cctttcttca gctctgaaag agaaacagct atcccactg acatgtcctt	1600
ctgagcccg taagagcaaa agaatggcag aaaagttag cccctgaaag	1650
ccatggagat tctcataact tgagacctaa tctctgtaaa gctaaaataa	1700

-continued

agaaatagaa caaggctgag gatacgacag tacactgtca gcagggactg 1750
taaacacaga caggggtcaaa gtgttttctc tgaacacatt gagttggaat 1800
cactgttttag aacacacaca ctactttttt ctggtctcta ccactgctga 1850
tatttttctct aggaaatata cttttacaag taacaaaaat aaaaactctt 1900
ataaatttct atttttatct gagttacaga aatgattact aaggaagatt 1950
actcagtaat ttgttttaaa agtaataaaa ttcaacaaac atttgctgaa 2000
tagctactat atgtcaagtg ctgtgcaagg tattacactc tgtaattgaa 2050
tattattcct caaaaaattg cacatagtag aacgctatct gggaagctat 2100
ttttttcagt ttgataattt ctagcttato tacttccaaa ctaattttta 2150
tttttgctga gactaatctt attcattttc tctaatatgg caaccattat 2200
aaccttaatt tattattaac atacctaaga agtacattgt tacctctata 2250
taccaaagca catttttaaa gtgccattaa caaatgtatc actagccctc 2300
ctttttccaa caagaaggga ctgagagatg cagaaatatt tgtgacaaaa 2350
aattaaagca tttagaaaac tt 2372

<210> SEQ ID NO 6
<211> LENGTH: 322
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 6

Met Ala Arg Cys Phe Ser Leu Val Leu Leu Leu Thr Ser Ile Trp
1 5 10 15
Thr Thr Arg Leu Leu Val Gln Gly Ser Leu Arg Ala Glu Glu Leu
20 25 30
Ser Ile Gln Val Ser Cys Arg Ile Met Gly Ile Thr Leu Val Ser
35 40 45
Lys Lys Ala Asn Gln Gln Leu Asn Phe Thr Glu Ala Lys Glu Ala
50 55 60
Cys Arg Leu Leu Gly Leu Ser Leu Ala Gly Lys Asp Gln Val Glu
65 70 75
Thr Ala Leu Lys Ala Ser Phe Glu Thr Cys Ser Tyr Gly Trp Val
80 85 90
Gly Asp Gly Phe Val Val Ile Ser Arg Ile Ser Pro Asn Pro Lys
95 100 105
Cys Gly Lys Asn Gly Val Gly Val Leu Ile Trp Lys Val Pro Val
110 115 120
Ser Arg Gln Phe Ala Ala Tyr Cys Tyr Asn Ser Ser Asp Thr Trp
125 130 135
Thr Asn Ser Cys Ile Pro Glu Ile Ile Thr Thr Lys Asp Pro Ile
140 145 150
Phe Asn Thr Gln Thr Ala Thr Gln Thr Thr Glu Phe Ile Val Ser
155 160 165
Asp Ser Thr Tyr Ser Val Ala Ser Pro Tyr Ser Thr Ile Pro Ala
170 175 180
Pro Thr Thr Thr Pro Pro Ala Pro Ala Ser Thr Ser Ile Pro Arg
185 190 195

-continued

Arg	Lys	Lys	Leu	Ile	Cys	Val	Thr	Glu	Val	Phe	Met	Glu	Thr	Ser
			200						205					210
Thr	Met	Ser	Thr	Glu	Thr	Glu	Pro	Phe	Val	Glu	Asn	Lys	Ala	Ala
			215						220					225
Phe	Lys	Asn	Glu	Ala	Ala	Gly	Phe	Gly	Gly	Val	Pro	Thr	Ala	Leu
			230						235					240
Leu	Val	Leu	Ala	Leu	Leu	Phe	Phe	Gly	Ala	Ala	Ala	Gly	Leu	Gly
			245						250					255
Phe	Cys	Tyr	Val	Lys	Arg	Tyr	Val	Lys	Ala	Phe	Pro	Phe	Thr	Asn
			260						265					270
Lys	Asn	Gln	Gln	Lys	Glu	Met	Ile	Glu	Thr	Lys	Val	Val	Lys	Glu
			275						280					285
Glu	Lys	Ala	Asn	Asp	Ser	Asn	Pro	Asn	Glu	Glu	Ser	Lys	Lys	Thr
			290						295					300
Asp	Lys	Asn	Pro	Glu	Glu	Ser	Lys	Ser	Pro	Ser	Lys	Thr	Thr	Val
			305						310					315
Arg	Cys	Leu	Glu	Ala	Glu	Val								
			320											

<210> SEQ ID NO 7
<211> LENGTH: 2586
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 7

cgccgcgctc ccgcacccgc ggcccgccca cgcgcgcgct cccgcacatctg	50
caccgcgagc ccggcggcct ccggcgggga gcgagcagat ccagtccggc	100
ccgcagcgca actcgggtcca gtcggggcgg cggctgcggg cgcagagcgg	150
agatgcagcg gcttggggcc accctgctgt gcctgctgct ggcggcggcg	200
gtccccacgg cccccgcgcc cgctccgacg gcgacctcgg ctccagtcaa	250
gcccggcccg gctctcagct acccgcagga ggaggccacc ctcaatgaga	300
tgttccgcga ggttgaggaa ctgatggagg acacgcagca caaattgcgc	350
agcgcggtgg aagagatgga ggcagaagaa gctgctgcta aagcatcatc	400
agaagtgaac ctggcaaact tacctcccag ctatcacaat gagaccaaca	450
cagacacgaa ggttggaat aataccatcc atgtgcaccg agaaattcac	500
aagataacca acaaccagac tggacaaatg gtcttttcag agacagttaa	550
cacatctgtg ggagacgaag aaggcagaag gagccacgag tgcacatctg	600
acgaggactg tgggcccagc atgtactgcc agtttgccag ctccagttac	650
acctgccagc catgccgggg ccagaggatg ctctgcaccc gggacagtga	700
gtgctgtgga gaccagctgt gtgtctgggg tcaactgcacc aaaatggcca	750
ccaggggcag caatgggacc atctgtgaca accagaggga ctgccagccg	800
gggctgtgct gtgccttcca gagaggcctg ctgttccctg tgtgcacacc	850
ctgccccgtg gagggcgagc ttgtccatga ccccgccagc cggtctcttg	900
acctcatcac ctgggagcta gagcctgatg gagccttgga ccgatgccct	950
tgtgccagtg gcctcctctg ccagccccac agccacagcc tgggtgtatgt	1000
gtgcaagcgc accttcgtgg ggagccgtga ccaagatggg gagatcctgc	1050

-continued

tgcccagaga ggtcccgat gagtatgaag ttggcagctt catggaggag	1100
gtgcgccagg agctggagga cctggagagg agcctgactg aagagatggc	1150
gctgggggag cctgcggtg ccgccgtgc actgctggga ggggaagaga	1200
tttagatctg gaccaggctg tgggtagatg tgcaatagaa atagctaatt	1250
tatttcccca ggtgtgtgct ttaggcgtgg gctgaccagg cttcttccta	1300
catcttcttc ccagtaagtt tcccctctgg cttgacagca tgagggtgtg	1350
tgcatattgt cagctcccc aggctgttct ccaggcttca cagtctggtg	1400
cttgggagag tcaggcaggg ttaaactgca ggagcagttt gccaccctg	1450
tccagattat tggctgcttt gcctctacca gttggcagac agccgtttgt	1500
tctacatggc ttgataaatt gtttgagggg aggagatgga aacaatgtgg	1550
agtcctccct tgattggttt tggggaaatg tggagaagag tgccctgctt	1600
tgcaaacatc aacctggcaa aaatgcaaca aatgaatttt ccacgcagtt	1650
ctttccatgg gcataggtaa gctgtgcctt cagctgttgc agatgaaatg	1700
ttctgttcac cctgcattac atgtgtttat tcattccagca gtgttgctca	1750
gtcctacct ctgtgccagg gcagcatttt catatccaag atcaattccc	1800
tctctcagca cagcctgggg agggggtcat tgttctctc gtccatcagg	1850
gatctcagag gctcagagac tgcaagctgc ttgcccaagt cacacagcta	1900
gtgaagacca gagcagtttc atctggttgt gactctaagc tcagtgtctt	1950
ctccactacc ccacaccagc cttggtgcc ccaaaagtgc tccccaaaag	2000
gaaggagaat gggatttttc ttgaggcatg cacatctgga attaaggtca	2050
aactaattct cacatccctc taaaagttaa ctactgttag gaacagcagt	2100
gttctcacag tgtggggcag ccgtccttct aatgaagaca atgatattga	2150
cactgtccct ctttggcagt tgcattagta actttgaaag gtatatgact	2200
gagcgtagca tacaggttaa cctgcagaaa cagtacttag gtaattgtag	2250
ggcgaggatt ataaatgaaa ttgcaaaat cacttagcag caactgaaga	2300
caattatcaa ccacgtggag aaaatcaaac cgagcagggc tgtgtgaaac	2350
atggttgtaa tatgcgactg cgaacactga actctacgcc actccacaaa	2400
tgatgttttc aggtgtcatg gactgttgcc accatgtatt catccagagt	2450
ttttaagttt taaagttgca catgattgta taagcatgct ttctttgagt	2500
tttaaatatt gtataaacat aagttgcatt tagaaatcaa gcataaatca	2550
cttcaactgc aaaaaaaaaa aaaaaaaaaa aaaaaa	2586

<210> SEQ ID NO 8
<211> LENGTH: 350
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 8

Met	Gln	Arg	Leu	Gly	Ala	Thr	Leu	Leu	Cys	Leu	Leu	Ala	Ala
1				5					10				15
Ala	Val	Pro	Thr	Ala	Pro	Ala	Pro	Ala	Pro	Thr	Ala	Thr	Ser
				20					25				30

-continued

Pro	Val	Lys	Pro	Gly	Pro	Ala	Leu	Ser	Tyr	Pro	Gln	Glu	Glu	Ala	
				35					40					45	
Thr	Leu	Asn	Glu	Met	Phe	Arg	Glu	Val	Glu	Glu	Leu	Met	Glu	Asp	
				50					55					60	
Thr	Gln	His	Lys	Leu	Arg	Ser	Ala	Val	Glu	Glu	Met	Glu	Ala	Glu	
				65					70					75	
Glu	Ala	Ala	Ala	Lys	Ala	Ser	Ser	Glu	Val	Asn	Leu	Ala	Asn	Leu	
				80					85					90	
Pro	Pro	Ser	Tyr	His	Asn	Glu	Thr	Asn	Thr	Asp	Thr	Lys	Val	Gly	
				95					100					105	
Asn	Asn	Thr	Ile	His	Val	His	Arg	Glu	Ile	His	Lys	Ile	Thr	Asn	
				110					115					120	
Asn	Gln	Thr	Gly	Gln	Met	Val	Phe	Ser	Glu	Thr	Val	Ile	Thr	Ser	
				125					130					135	
Val	Gly	Asp	Glu	Glu	Gly	Arg	Arg	Ser	His	Glu	Cys	Ile	Ile	Asp	
				140					145					150	
Glu	Asp	Cys	Gly	Pro	Ser	Met	Tyr	Cys	Gln	Phe	Ala	Ser	Phe	Gln	
				155					160					165	
Tyr	Thr	Cys	Gln	Pro	Cys	Arg	Gly	Gln	Arg	Met	Leu	Cys	Thr	Arg	
				170					175					180	
Asp	Ser	Glu	Cys	Cys	Gly	Asp	Gln	Leu	Cys	Val	Trp	Gly	His	Cys	
				185					190					195	
Thr	Lys	Met	Ala	Thr	Arg	Gly	Ser	Asn	Gly	Thr	Ile	Cys	Asp	Asn	
				200					205					210	
Gln	Arg	Asp	Cys	Gln	Pro	Gly	Leu	Cys	Cys	Ala	Phe	Gln	Arg	Gly	
				215					220					225	
Leu	Leu	Phe	Pro	Val	Cys	Thr	Pro	Leu	Pro	Val	Glu	Gly	Glu	Leu	
				230					235					240	
Cys	His	Asp	Pro	Ala	Ser	Arg	Leu	Leu	Asp	Leu	Ile	Thr	Trp	Glu	
				245					250					255	
Leu	Glu	Pro	Asp	Gly	Ala	Leu	Asp	Arg	Cys	Pro	Cys	Ala	Ser	Gly	
				260					265					270	
Leu	Leu	Cys	Gln	Pro	His	Ser	His	Ser	Leu	Val	Tyr	Val	Cys	Lys	
				275					280					285	
Pro	Thr	Phe	Val	Gly	Ser	Arg	Asp	Gln	Asp	Gly	Glu	Ile	Leu	Leu	
				290					295					300	
Pro	Arg	Glu	Val	Pro	Asp	Glu	Tyr	Glu	Val	Gly	Ser	Phe	Met	Glu	
				305					310					315	
Glu	Val	Arg	Gln	Glu	Leu	Glu	Asp	Leu	Glu	Arg	Ser	Leu	Thr	Glu	
				320					325					330	
Glu	Met	Ala	Leu	Gly	Glu	Pro	Ala	Ala	Ala	Ala	Ala	Ala	Leu	Leu	
				335					340					345	
Gly	Gly	Glu	Glu	Ile											
				350											

<210> SEQ ID NO 9
<211> LENGTH: 1395
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 9

cggacgcgtg ggcggacgcg tgggggctgt gagaaagtgc caataaatac

-continued

atcatgcaac cccacggccc accttgtgaa ctctctgtgc ccagggctga	100
tgtgcgtctt ccagggctac tcatccaaag gcctaatacca acgttctgtc	150
ttcaatctgc aaatctatgg ggtcctgggg ctcttctgga cccttaactg	200
ggtactggcc ctgggccaat gcgtcctcgc tggagccttt gcctccttct	250
actgggcctt ccacaagccc caggacatcc ctaccttccc cttaatctct	300
gccttcaccc gcacactccg ttaccacact gggtcattgg catttggagc	350
cctcatcctg acccttgtgc agatagcccg ggtcatcttg gagtatattg	400
accacaagct cagaggagtg cagaaccctg tagcccgctg catcatgtgc	450
tgtttcaagt gctgcctctg gtgtctggaa aaatttatca agttcctaaa	500
ccgcaatgca tacatcatga tcgccatcta cggaagaat ttctgtgtct	550
cagccaaaaa tgcgttcatg ctactcatgc gaaacattgt cagggtggtc	600
gtcctggaca aagtcacaga cctgctgtg ttctttggga agctgctggt	650
ggtcggaggc gtgggggtcc tgtccttctt ttttttctcc ggtcgcatcc	700
cggggctggg taaagacttt aagagccccc acctcaacta ttactggctg	750
cccacatgca cctccatcct gggggcctat gtcacgcca gcggcttctt	800
cagcgttttc ggcattgtgt tggacacgct cttcctctgc ttcttggaag	850
acctggagcg gaacaacggc tccctggacc ggccctacta catgtccaag	900
agccttctaa agattctggg caagaagaac gaggcgcccc cggacaacaa	950
gaagagggaag aagtgcacgc tccggccctg atccaggact gacccccacc	1000
cccaccgtcc agccatccaa cctcacttcg ccttacaggt ctccattttg	1050
tggtaaaaaa aggttttagg ccaggcgccg tggctcacgc ctgtaatacca	1100
acacttttag aggctgaggc gggcggtatc cctgagtcag gagttcgaga	1150
ccagcctggc caacatggtg aaacctccgt ctctattaaa aatacaaaaa	1200
ttagccgaga gtggtggcat gcacctgtca tcccagctac tcgggaggct	1250
gaggcaggag aatcgcttga acccgggagg cagagggtgc agtgagccga	1300
gatcgcgcca ctgcactcca acctgggtga cagactctgt ctccaaaaca	1350
aaacaaacaa acaaaaagat ttattataag atattttggt aactc	1395

<210> SEQ ID NO 10
<211> LENGTH: 321
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 10

Arg Thr Arg Gly Arg Thr Arg Gly Gly Cys Glu Lys Val Pro Ile	
1 5 10 15	
Asn Thr Ser Cys Asn Pro Thr Ala His Leu Val Asn Ser Ser Cys	
20 25 30	
Pro Gly Leu Met Cys Val Phe Gln Gly Tyr Ser Ser Lys Gly Leu	
35 40 45	
Ile Gln Arg Ser Val Phe Asn Leu Gln Ile Tyr Gly Val Leu Gly	
50 55 60	
Leu Phe Trp Thr Leu Asn Trp Val Leu Ala Leu Gly Gln Cys Val	
65 70 75	

-continued

Leu Ala Gly Ala Phe Ala Ser Phe Tyr Trp Ala Phe His Lys Pro
80 85 90
Gln Asp Ile Pro Thr Phe Pro Leu Ile Ser Ala Phe Ile Arg Thr
95 100 105
Leu Arg Tyr His Thr Gly Ser Leu Ala Phe Gly Ala Leu Ile Leu
110 115 120
Thr Leu Val Gln Ile Ala Arg Val Ile Leu Glu Tyr Ile Asp His
125 130 135
Lys Leu Arg Gly Val Gln Asn Pro Val Ala Arg Cys Ile Met Cys
140 145 150
Cys Phe Lys Cys Cys Leu Trp Cys Leu Glu Lys Phe Ile Lys Phe
155 160 165
Leu Asn Arg Asn Ala Tyr Ile Met Ile Ala Ile Tyr Gly Lys Asn
170 175 180
Phe Cys Val Ser Ala Lys Asn Ala Phe Met Leu Leu Met Arg Asn
185 190 195
Ile Val Arg Val Val Val Leu Asp Lys Val Thr Asp Leu Leu Leu
200 205 210
Phe Phe Gly Lys Leu Leu Val Val Gly Gly Val Gly Val Leu Ser
215 220 225
Phe Phe Phe Phe Ser Gly Arg Ile Pro Gly Leu Gly Lys Asp Phe
230 235 240
Lys Ser Pro His Leu Asn Tyr Tyr Trp Leu Pro Ile Met Thr Ser
245 250 255
Ile Leu Gly Ala Tyr Val Ile Ala Ser Gly Phe Phe Ser Val Phe
260 265 270
Gly Met Cys Val Asp Thr Leu Phe Leu Cys Phe Leu Glu Asp Leu
275 280 285
Glu Arg Asn Asn Gly Ser Leu Asp Arg Pro Tyr Tyr Met Ser Lys
290 295 300
Ser Leu Leu Lys Ile Leu Gly Lys Lys Asn Glu Ala Pro Pro Asp
305 310 315
Asn Lys Lys Arg Lys Lys
320

<210> SEQ ID NO 11
<211> LENGTH: 1901
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 11

gccccgcgcc cggcgcggg cgccgaagc cgggagccac cgccatgggg 50
gcctgcctgg gagcctgctc cctgtcagc tgcgcgtcct gcctctgcgg 100
ctctgcccc tgcacctgt gcagctgtg ccccgccagc cgcaactcca 150
ccgtgagcgg cctcatcttc acgtttcttc tcttctctggg ggtgctggtg 200
tccatcatta tgctgagccc gggcgtggag agtcagctct acaagctgcc 250
ctgggtgtgt gaggaggggg ccgggatccc caccgtcctg cagggccaca 300
tcgactgtgg ctccctgctt ggctaccgcg ctgtctaccg catgtgcttc 350
gccacggcgg ccttcttctt cttctttttc accctgctca tgctctgcgt 400

-continued

gagcagcagc cgggaccccc gggtgccat ccagaatggg ttttggttct	450
ttaagttcct gatcctggtg ggctcaccg tgggtgcctt ctacatccct	500
gacggctcct tcaccaacat ctggttctac ttcggcgtcg tgggtccctt	550
cctcttcatc ctcatccagc tgggtgtgct catcgacttt gcgcactcct	600
ggaaccagcg gtggctgggc aaggccgagg agtgcgattc ccgtgcctgg	650
tacgcaggcc tcttcttctt cactctcctc ttctacttgc tgtcgatcgc	700
ggcgtggcg ctgatgttca tgtactacac tgagcccagc ggctgccacg	750
agggcaaggt cttcatcagc ctcaacctca ccttctgtgt ctgcggtgcc	800
atcgctgctg tcctgcccaa ggtccaggac gccagccca actcgggtct	850
gctgcaggcc tcggtcatca ccctctacac catgtttgtc acctggtcag	900
ccctatccag tatccctgaa cagaaatgca accccattt gccaacccag	950
ctgggcaacg agacagttgt ggcaggcccc gagggctatg agaccagtg	1000
gtgggatgcc ccgagcattg tgggcctcat catcttctc ctgtgcaccc	1050
tcttcatcag tctgcgtcc tcagaccacc ggcaggtgaa cagcctgatg	1100
cagaccgagg agtgcccacc tatgctagac gccacacagc agcagcagca	1150
gcaggtgga gctgtgagg gccgggcctt tgacaacgag caggacggcg	1200
tcacctagc ctactccttc ttccacttct gcctggtgct ggcctcactg	1250
cacgtcatga tgacgtcac caactggtac aagcccggtg agaccggaa	1300
gatgatcagc acgtggaccg ccgtgtgggt gaagatctgt gccagctggg	1350
cagggtctgt cctctacctg tggaccctgg tagccccact cctcctgcgc	1400
aaccgcgact tcagctgagg cagcctcaca gcctgccatc tggtgccctc	1450
tgccacctgg tgcctctcgg ctcggtgaca gccaacctgc cccctcccca	1500
caccaatcag ccaggctgag cccccacccc tgcccagct ccaggacctg	1550
cccctgagcc gggccttcta gtcgtagtgc cttcagggtc cgaggagcat	1600
caggctcctg cagagcccca tcccccgcc acaccacac ggtggagctg	1650
cctcttctct cccctcctcc ctggtgccca tactcagcat ctcggatgaa	1700
agggctccct tgtcctcagg ctccacggga gcggggctgc tggagagagc	1750
ggggaactcc caccacagtg gggcatccgg cactgaagcc ctggtgttcc	1800
tggtcacgtc cccagggga ccctgcccc ttcttggaact tcgtgcctta	1850
ctgagtctct aagacttttt ctaataaaca agccagtgcg tgtaaaaaaa	1900
a	1901

<210> SEQ ID NO 12
<211> LENGTH: 457
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 12

Met Gly Ala Cys Leu Gly Ala Cys Ser Leu Leu Ser Cys Ala Ser
1 5 10 15
Cys Leu Cys Gly Ser Ala Pro Cys Ile Leu Cys Ser Cys Cys Pro
20 25 30

-continued

Ala	Ser	Arg	Asn	Ser	Thr	Val	Ser	Arg	Leu	Ile	Phe	Thr	Phe	Phe		35	40	45
Leu	Phe	Leu	Gly	Val	Leu	Val	Ser	Ile	Ile	Met	Leu	Ser	Pro	Gly		50	55	60
Val	Glu	Ser	Gln	Leu	Tyr	Lys	Leu	Pro	Trp	Val	Cys	Glu	Glu	Gly		65	70	75
Ala	Gly	Ile	Pro	Thr	Val	Leu	Gln	Gly	His	Ile	Asp	Cys	Gly	Ser		80	85	90
Leu	Leu	Gly	Tyr	Arg	Ala	Val	Tyr	Arg	Met	Cys	Phe	Ala	Thr	Ala		95	100	105
Ala	Phe	Phe	Phe	Phe	Phe	Phe	Thr	Leu	Leu	Met	Leu	Cys	Val	Ser		110	115	120
Ser	Ser	Arg	Asp	Pro	Arg	Ala	Ala	Ile	Gln	Asn	Gly	Phe	Trp	Phe		125	130	135
Phe	Lys	Phe	Leu	Ile	Leu	Val	Gly	Leu	Thr	Val	Gly	Ala	Phe	Tyr		140	145	150
Ile	Pro	Asp	Gly	Ser	Phe	Thr	Asn	Ile	Trp	Phe	Tyr	Phe	Gly	Val		155	160	165
Val	Gly	Ser	Phe	Leu	Phe	Ile	Leu	Ile	Gln	Leu	Val	Leu	Leu	Ile		170	175	180
Asp	Phe	Ala	His	Ser	Trp	Asn	Gln	Arg	Trp	Leu	Gly	Lys	Ala	Glu		185	190	195
Glu	Cys	Asp	Ser	Arg	Ala	Trp	Tyr	Ala	Gly	Leu	Phe	Phe	Phe	Thr		200	205	210
Leu	Leu	Phe	Tyr	Leu	Leu	Ser	Ile	Ala	Ala	Val	Ala	Leu	Met	Phe		215	220	225
Met	Tyr	Tyr	Thr	Glu	Pro	Ser	Gly	Cys	His	Glu	Gly	Lys	Val	Phe		230	235	240
Ile	Ser	Leu	Asn	Leu	Thr	Phe	Cys	Val	Cys	Val	Ser	Ile	Ala	Ala		245	250	255
Val	Leu	Pro	Lys	Val	Gln	Asp	Ala	Gln	Pro	Asn	Ser	Gly	Leu	Leu		260	265	270
Gln	Ala	Ser	Val	Ile	Thr	Leu	Tyr	Thr	Met	Phe	Val	Thr	Trp	Ser		275	280	285
Ala	Leu	Ser	Ser	Ile	Pro	Glu	Gln	Lys	Cys	Asn	Pro	His	Leu	Pro		290	295	300
Thr	Gln	Leu	Gly	Asn	Glu	Thr	Val	Val	Ala	Gly	Pro	Glu	Gly	Tyr		305	310	315
Glu	Thr	Gln	Trp	Trp	Asp	Ala	Pro	Ser	Ile	Val	Gly	Leu	Ile	Ile		320	325	330
Phe	Leu	Leu	Cys	Thr	Leu	Phe	Ile	Ser	Leu	Arg	Ser	Ser	Asp	His		335	340	345
Arg	Gln	Val	Asn	Ser	Leu	Met	Gln	Thr	Glu	Glu	Cys	Pro	Pro	Met		350	355	360
Leu	Asp	Ala	Thr	Gln	Gln	Gln	Gln	Gln	Gln	Val	Ala	Ala	Cys	Glu		365	370	375
Gly	Arg	Ala	Phe	Asp	Asn	Glu	Gln	Asp	Gly	Val	Thr	Tyr	Ser	Tyr		380	385	390
Ser	Phe	Phe	His	Phe	Cys	Leu	Val	Leu	Ala	Ser	Leu	His	Val	Met		395	400	405
Met	Thr	Leu	Thr	Asn	Trp	Tyr	Lys	Pro	Gly	Glu	Thr	Arg	Lys	Met				

-continued

[illegible]

-continued

tctctgattg ttctgaaatg ttctaaatac tcttattttg aatgcacaaa 1500
atgacttaaa ccattcatat catgtttcct ttgcgttcag ccaatttcaa 1550
ttaaaatgaa ctaaattaaa aa 1572

<210> SEQ ID NO 14
<211> LENGTH: 234
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 14

Met Asn His Leu Pro Glu Asp Met Glu Asn Ala Leu Thr Gly Ser
1 5 10 15
Gln Ser Ser His Ala Ser Leu Arg Asn Ile His Ser Ile Asn Pro
20 25 30
Thr Gln Leu Met Ala Arg Ile Glu Ser Tyr Glu Gly Arg Glu Lys
35 40 45
Lys Gly Ile Ser Asp Val Arg Arg Thr Phe Cys Leu Phe Val Thr
50 55 60
Phe Asp Leu Leu Phe Val Thr Leu Leu Trp Ile Ile Glu Leu Asn
65 70 75
Val Asn Gly Gly Ile Glu Asn Thr Leu Glu Lys Glu Val Met Gln
80 85 90
Tyr Asp Tyr Tyr Ser Ser Tyr Phe Asp Ile Phe Leu Leu Ala Val
95 100 105
Phe Arg Phe Lys Val Leu Ile Leu Ala Tyr Ala Val Cys Arg Leu
110 115 120
Arg His Trp Trp Ala Ile Ala Leu Thr Thr Ala Val Thr Ser Ala
125 130 135
Phe Leu Leu Ala Lys Val Ile Leu Ser Lys Leu Phe Ser Gln Gly
140 145 150
Ala Phe Gly Tyr Val Leu Pro Ile Ile Ser Phe Ile Leu Ala Trp
155 160 165
Ile Glu Thr Trp Phe Leu Asp Phe Lys Val Leu Pro Gln Glu Ala
170 175 180
Glu Glu Glu Asn Arg Leu Leu Ile Val Gln Asp Ala Ser Glu Arg
185 190 195
Ala Ala Leu Ile Pro Gly Gly Leu Ser Asp Gly Gln Phe Tyr Ser
200 205 210
Pro Pro Glu Ser Glu Ala Gly Ser Glu Glu Ala Glu Glu Lys Gln
215 220 225
Asp Ser Glu Lys Pro Leu Leu Glu Leu
230

<210> SEQ ID NO 15
<211> LENGTH: 2768
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 15

actgaacgc agttgcttcg ggacccagga cccctcggg cccgacccgc 50
caggaaagac tgaggccgcg gcctgccccg cccggtccc tgcgcgcgcg 100
ccgcctcccg ggacagaaga tgtgtccag ggtccctctg ctgctgccgc 150

-continued

tgctcctgct actggccctg gggcctgggg tgcagggctg cccatccggc	200
tgccagtgea gccagccaca gacagtcttc tgactgccc gccaggggac	250
cacggtgccc cgagacgtgc caccgacac ggtggggctg tacgtctttg	300
agaacggcat caccatgctc gacgcaggca gctttgccgg cctgccgggc	350
ctgcagctcc tggacctgtc acagaaccag atcgccagcc tgcccagcgg	400
ggtcttccag ccactcgcca acctcagcaa cctggacctg acggccaaca	450
ggctgcata aatcaccaat gagaccttcc gtggcctgcg gcgcctcgag	500
cgctcttacc tgggcaagaa ccgcctccgc cacatccagc ctggtgcctt	550
cgacacgtc gaccgcctcc tggagctcaa gctgcaggac aacgagctgc	600
gggcactgcc cccgctgcg cctgcccgc tgctgctgct ggacctcagc	650
cacaacagcc tcctggccct ggagcccgcc atcctggaca ctgccaacgt	700
ggaggcgctg cggctggctg gtctggggct gcagcagctg gacgaggggc	750
tcttcagcgg cttgcgcaac ctccacgacc tggatgtgtc cgacaaccag	800
ctggagcgag tgccacctgt gatccgaggc ctccggggcc tgacgcgcct	850
gcggctggcc ggcaacaccc gcattgccc gctgcggccc gaggacctgg	900
ccggcctggc tgccctgcag gagctggatg tgagcaacct aagcctgcag	950
gccctgcctg gcgacctctc gggcctcttc cccgcctgc ggctgctggc	1000
agctgcccgc aacccttca actgcgtgtg cccctgagc tggtttgcc	1050
cctgggtgcy cgagagccac gtcacactgg ccagccctga ggagacgcgc	1100
tgccacttcc cgccaagaa cgtggccgg ctgctcctgg agcttgacta	1150
cgccgacttt ggctgcccag ccaccaccac cacagccaca gtgcccacca	1200
cgagcccggt ggtgcgggag cccacagcct tgtcttctag cttggctcct	1250
acctggctta gcccacagc gcgggccact gaggccccca gccgcctc	1300
cactgcccga ccgactgtag ggcctgtccc ccagccccag gactgcccac	1350
cgctccacctg cctcaatggg ggcacatgcc acctggggac acggcaccac	1400
ctggcgctgt tgtgcccga aggtctcacg ggcctgtact gtgagagcca	1450
gatggggcag gggacacggc ccagccctac accagtacag ccgagggcac	1500
cacggtccct gaccctgggc atcgagccgg tgagccccac ctccctgcgc	1550
gtggggctgc agcgctacct ccaggggagc tccgtgcagc tcaggagcct	1600
ccgtctcacc tatcgcaacc tatcgggccc tgataagcgg ctggtgacgc	1650
tgcgactgcc tgctcgtctc gctgagtaca cggtcaccca gctgcggccc	1700
aacgccactt actccgtctg tgtcatgcct ttggggcccg ggcgggtgcc	1750
ggagggcgag gaggcctgcy gggaggccca tacaccccca gccgtccact	1800
ccaaccacgc ccagtcacc caggcccgcy agggcaacct gccgtcctc	1850
attgcgccc ccctggccgc ggtgtcctg gccgcgctgg ctgcggtggg	1900
ggcagcctac tgtgtgcggc gggggcgggc catggcagca gcggctcagg	1950
acaaagggca ggtggggcca ggggctgggc ccctggaact ggagggagtg	2000
aaggctccct tggagccagg ccgaaggca acagaggcg gtggagaggc	2050

-continued

```

cctgccagc gggctctgagt gtgaggtgcc actcatgggc tccccagggc      2100
ctggcctcca gtcacccctc cacgcaaagc cctacatcta agccagagag      2150
agacagggca gctggggccg ggctctcagc cagtgagatg gccagccccc      2200
tcctgctgcc acaccacgta agttctcagt cccaacctcg gggatgtgtg      2250
cagacagggc tgtgtgacca cagctggggc ctgttccctc tggacctcgg      2300
tctcctcatc tgtgagatgc tgtggcccag ctgacgagcc ctaacgtccc      2350
cagaaccgag tgcctatgag gacagtgtcc gccctgccct ccgcaacgtg      2400
cagtccctgg gcacggcggg ccctgccatg tgctggtaac gcatgcctgg      2450
gtcctgctgg gctctccac tccaggcggg ccctgggggc cagtgaagga      2500
agctcccgga aagagcagag ggagagcggg taggcggctg tgtgactcta      2550
gtcttggccc caggaagcga aggaacaaaa gaaactggaa aggaagatgc      2600
tttaggaaca tgttttgctt ttttaaaata tataatatta taagagatcc      2650
tttcccatth attctgggaa gatgtttttc aaactcagag acaaggactt      2700
tggtttttgt aagacaaacg atgatatgaa ggccttttgt aagaaaaaat      2750
aaaagatgaa gtgtgaaa      2768

```

<210> SEQ ID NO 16

<211> LENGTH: 673

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 16

```

Met Cys Ser Arg Val Pro Leu Leu Leu Pro Leu Leu Leu Leu Leu
  1             5             10             15

Ala Leu Gly Pro Gly Val Gln Gly Cys Pro Ser Gly Cys Gln Cys
  20             25             30

Ser Gln Pro Gln Thr Val Phe Cys Thr Ala Arg Gln Gly Thr Thr
  35             40             45

Val Pro Arg Asp Val Pro Pro Asp Thr Val Gly Leu Tyr Val Phe
  50             55             60

Glu Asn Gly Ile Thr Met Leu Asp Ala Gly Ser Phe Ala Gly Leu
  65             70             75

Pro Gly Leu Gln Leu Leu Asp Leu Ser Gln Asn Gln Ile Ala Ser
  80             85             90

Leu Pro Ser Gly Val Phe Gln Pro Leu Ala Asn Leu Ser Asn Leu
  95            100            105

Asp Leu Thr Ala Asn Arg Leu His Glu Ile Thr Asn Glu Thr Phe
 110            115            120

Arg Gly Leu Arg Arg Leu Glu Arg Leu Tyr Leu Gly Lys Asn Arg
 125            130            135

Ile Arg His Ile Gln Pro Gly Ala Phe Asp Thr Leu Asp Arg Leu
 140            145            150

Leu Glu Leu Lys Leu Gln Asp Asn Glu Leu Arg Ala Leu Pro Pro
 155            160            165

Leu Arg Leu Pro Arg Leu Leu Leu Leu Asp Leu Ser His Asn Ser
 170            175            180

Leu Leu Ala Leu Glu Pro Gly Ile Leu Asp Thr Ala Asn Val Glu

```

-continued

185	190	195
Ala Leu Arg Leu	Ala Gly Leu Gly Leu	Gln Gln Leu Asp Glu Gly
200	205	210
Leu Phe Ser Arg	Leu Arg Asn Leu His	Asp Leu Asp Val Ser Asp
215	220	225
Asn Gln Leu Glu	Arg Val Pro Pro Val	Ile Arg Gly Leu Arg Gly
230	235	240
Leu Thr Arg Leu	Arg Leu Ala Gly Asn	Thr Arg Ile Ala Gln Leu
245	250	255
Arg Pro Glu Asp	Leu Ala Gly Leu Ala	Ala Leu Gln Glu Leu Asp
260	265	270
Val Ser Asn Leu	Ser Leu Gln Ala Leu	Pro Gly Asp Leu Ser Gly
275	280	285
Leu Phe Pro Arg	Leu Arg Leu Leu Ala	Ala Ala Arg Asn Pro Phe
290	295	300
Asn Cys Val Cys	Pro Leu Ser Trp Phe	Gly Pro Trp Val Arg Glu
305	310	315
Ser His Val Thr	Leu Ala Ser Pro Glu	Glu Thr Arg Cys His Phe
320	325	330
Pro Pro Lys Asn	Ala Gly Arg Leu Leu	Leu Glu Leu Asp Tyr Ala
335	340	345
Asp Phe Gly Cys	Pro Ala Thr Thr Thr	Thr Ala Thr Val Pro Thr
350	355	360
Thr Arg Pro Val	Val Arg Glu Pro Thr	Ala Leu Ser Ser Ser Leu
365	370	375
Ala Pro Thr Trp	Leu Ser Pro Thr Ala	Pro Ala Thr Glu Ala Pro
380	385	390
Ser Pro Pro Ser	Thr Ala Pro Pro Thr	Val Gly Pro Val Pro Gln
395	400	405
Pro Gln Asp Cys	Pro Pro Ser Thr Cys	Leu Asn Gly Gly Thr Cys
410	415	420
His Leu Gly Thr	Arg His His Leu Ala	Cys Leu Cys Pro Glu Gly
425	430	435
Phe Thr Gly Leu	Tyr Cys Glu Ser Gln	Met Gly Gln Gly Thr Arg
440	445	450
Pro Ser Pro Thr	Pro Val Thr Pro Arg	Pro Pro Arg Ser Leu Thr
455	460	465
Leu Gly Ile Glu	Pro Val Ser Pro Thr	Ser Leu Arg Val Gly Leu
470	475	480
Gln Arg Tyr Leu	Gln Gly Ser Ser Val	Gln Leu Arg Ser Leu Arg
485	490	495
Leu Thr Tyr Arg	Asn Leu Ser Gly Pro	Asp Lys Arg Leu Val Thr
500	505	510
Leu Arg Leu Pro	Ala Ser Leu Ala Glu	Tyr Thr Val Thr Gln Leu
515	520	525
Arg Pro Asn Ala	Thr Tyr Ser Val Cys	Val Met Pro Leu Gly Pro
530	535	540
Gly Arg Val Pro	Glu Gly Glu Glu Ala	Cys Gly Glu Ala His Thr
545	550	555
Pro Pro Ala Val	His Ser Asn His Ala	Pro Val Thr Gln Ala Arg
560	565	570

-continued

Glu Gly Asn Leu Pro Leu Leu Ile Ala Pro Ala Leu Ala Ala Val	575	580	585
Leu Leu Ala Ala Leu Ala Ala Val Gly Ala Ala Tyr Cys Val Arg	590	595	600
Arg Gly Arg Ala Met Ala Ala Ala Ala Gln Asp Lys Gly Gln Val	605	610	615
Gly Pro Gly Ala Gly Pro Leu Glu Leu Glu Gly Val Lys Val Pro	620	625	630
Leu Glu Pro Gly Pro Lys Ala Thr Glu Gly Gly Gly Glu Ala Leu	635	640	645
Pro Ser Gly Ser Glu Cys Glu Val Pro Leu Met Gly Phe Pro Gly	650	655	660
Pro Gly Leu Gln Ser Pro Leu His Ala Lys Pro Tyr Ile	665	670	

<210> SEQ ID NO 17
<211> LENGTH: 1672
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 17

gcagcggcga ggcggcggtg gtggctgagt ccgtggtggc agaggcgaag	50
gcgacagctc atgcgggtcc ggatagggct gacgctgctg ctgtgtgcgg	100
tgctgctgag cttggcctcg gcgtcctcgg atgaagaagg cagccaggat	150
gaatccttag attccaagac tactttgaca tcagatgagt cagtaaagga	200
ccatactact gcaggcagag tagttgctgg tcaaataatt cttgattcag	250
aagaatctga attagaatcc tctattcaag aagaggaaga cagcctcaag	300
agccaagagg gggaaagtgt cacagaagat atcagctttc tagagtctcc	350
aaatccagaa aacaaggact atgaagagcc aaagaaagta cggaaaccag	400
ctttgaccgc cattgaaggc acagcacatg gggagccctg ccacttcctt	450
tttcttttcc tagataagga gtatgatgaa tgtacatcag atgggaggga	500
agatggcaga ctgtggtgtg ctacaaccta tgactacaaa gcagatgaaa	550
agtggggctt ttgtgaaact gaagaagagg ctgctaagag acggcagatg	600
caggaagcag aaatgatgta tcaaactgga atgaaatcc ttaatggaag	650
caataagaaa agccaaaaaa gagaagcata tcggtatctc caaaaggcag	700
caagcatgaa ccataccaaa gccctggaga gagtgtcata tgctctttta	750
tttggtgatt acttgccaca gaatatccag gcagcgagag agatgtttga	800
gaagctgact gaggaaggct ctccaaggg acagactgct cttggctttc	850
tgtatgcctc tggacttggg gttaattcaa gtcaggcaaa ggctcttgta	900
tattatacat ttggagctct tgggggcaat ctaatagccc acatggtttt	950
ggtaagtaga ctttagtgga aggctaataa tattaacatc agaagaattt	1000
gtggtttata gcggccacaa ctttttcagc tttcatgata cagatttgct	1050
tgtattaaga ccaaataatt agttgaactt cttcaaat cttgttaaat	1100
gatataacac atggaatcta catgaaatg aaagtgtgtg gagtccacaa	1150

-continued

tttttcttta aaatgattag ttgggtgat tgcccctaaa aagagagatc	1200
tgataaatgg ctctttttaa attttctctg agttggaatt gtcagaatca	1250
ttttttacat tagattatca taattttaaa aatttttctt tagtttttca	1300
aaattttgta aatggtggct atagaaaaac aacatgaaat attatacaat	1350
attttgcaac aatgccctaa gaattgttaa aattcatgga gttatttgtg	1400
cagaatgact ccagagagct ctactttctg ttttttactt ttcatgattg	1450
gctgtcttcc catttattct ggctatttat tgctagtgc actgtgcctg	1500
cttccagtag tctcattttc cctattttgc taatttgta ctttttcttt	1550
gctaatttgg aagattaact catttttaat aaaattatgt ctaagattaa	1600
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1650
aaaaaaaaa aaaaaaaaaa aa	1672

<210> SEQ ID NO 18

<211> LENGTH: 301

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 18

Met Arg Val Arg Ile Gly Leu Thr Leu Leu Leu Cys Ala Val Leu	1	5	10	15
Leu Ser Leu Ala Ser Ala Ser Ser Asp Glu Glu Gly Ser Gln Asp	20	25	30	
Glu Ser Leu Asp Ser Lys Thr Thr Leu Thr Ser Asp Glu Ser Val	35	40	45	
Lys Asp His Thr Thr Ala Gly Arg Val Val Ala Gly Gln Ile Phe	50	55	60	
Leu Asp Ser Glu Glu Ser Glu Leu Glu Ser Ser Ile Gln Glu Glu	65	70	75	
Glu Asp Ser Leu Lys Ser Gln Glu Gly Glu Ser Val Thr Glu Asp	80	85	90	
Ile Ser Phe Leu Glu Ser Pro Asn Pro Glu Asn Lys Asp Tyr Glu	95	100	105	
Glu Pro Lys Lys Val Arg Lys Pro Ala Leu Thr Ala Ile Glu Gly	110	115	120	
Thr Ala His Gly Glu Pro Cys His Phe Pro Phe Leu Phe Leu Asp	125	130	135	
Lys Glu Tyr Asp Glu Cys Thr Ser Asp Gly Arg Glu Asp Gly Arg	140	145	150	
Leu Trp Cys Ala Thr Thr Tyr Asp Tyr Lys Ala Asp Glu Lys Trp	155	160	165	
Gly Phe Cys Glu Thr Glu Glu Glu Ala Ala Lys Arg Arg Gln Met	170	175	180	
Gln Glu Ala Glu Met Met Tyr Gln Thr Gly Met Lys Ile Leu Asn	185	190	195	
Gly Ser Asn Lys Lys Ser Gln Lys Arg Glu Ala Tyr Arg Tyr Leu	200	205	210	
Gln Lys Ala Ala Ser Met Asn His Thr Lys Ala Leu Glu Arg Val	215	220	225	
Ser Tyr Ala Leu Leu Phe Gly Asp Tyr Leu Pro Gln Asn Ile Gln				

-continued

230										235										240									
Ala	Ala	Arg	Glu	Met	Phe	Glu	Lys	Leu	Thr	Glu	Glu	Gly	Ser	Pro															
				245						250				255															
Lys	Gly	Gln	Thr	Ala	Leu	Gly	Phe	Leu	Tyr	Ala	Ser	Gly	Leu	Gly															
				260						265				270															
Val	Asn	Ser	Ser	Gln	Ala	Lys	Ala	Leu	Val	Tyr	Tyr	Thr	Phe	Gly															
				275						280				285															
Ala	Leu	Gly	Gly	Asn	Leu	Ile	Ala	His	Met	Val	Leu	Val	Ser	Arg															
				290						295				300															

Leu

<210> SEQ ID NO 19
<211> LENGTH: 1508
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 19

aattcagatt ttaagcccat tctgcagtgg aatttcatga actagcaaga	50
ggacaccatc ttcttgtatt atacaagaaa ggagtgtacc tatcacacac	100
agggggaaaa atgctctttt ggggtgctagg cctcctaatac ctctgtggtt	150
ttctgtggac tcgtaaagga aaactaaaga ttgaagacat cactgataag	200
tacattttta tcaactggatg tgactcgggc tttggaaact tggcagccag	250
aacttttgat aaaaagggat ttcattgtaat cgctgcctgt ctgactgaat	300
caggatcaac agctttaaag gcagaaacct cagagagact tcgtactgtg	350
cttctggatg tgaccgaccc agagaatgac aagaggactg cccagtgggt	400
gaagaaccaa gttggggaga aaggtctctg gggctctgac aataatgctg	450
gtgttcccgg cgtgctggct cccactgact ggctgacact agaggactac	500
agagaacctt ttgaagtga cctgtttgga ctcatcagtg tgacactaaa	550
tatgcttctt ttggtcaaga aagctcaagg gagagttatt aatgtctcca	600
gtgttgagg tcgccttgca atcgttgagg ggggctatac tccatccaaa	650
tatgcagtgg aaggtttcaa tgacagctta agacgggaca tgaaagcttt	700
tggtgtgcac gtctcatgca ttgaaccagg attgttcaaa aaaaacttgg	750
cagatccagt aaaggttaatt gaaaaaaac tcgccatttg ggagcagctg	800
tctccagaca tcaacaaca atatggagaa ggttacattg aaaaaagtct	850
agacaaactg aaaggcaata aatcctatgt gaacatggac ctctctccgg	900
tggtagagtg catggaccac gctctaacaa gtctcttccc taagactcat	950
tatgccgctg gaaaagatgc caaaattttc tggatacctc tgtctcacat	1000
gccagcagct ttgcaagact ttttattgtt gaaacagaaa gcagagctgg	1050
ctaattccaa ggcagtgtga ctacagtaac cacaatgctc tcctccaggc	1100
tatgaaattg gccgatttca agaacacatc tccttttcaa cccatttcct	1150
tatctgctcc aacctggact catttagatc gtgcttattt ggattgcaaa	1200
agggagtccc accatcgctg gtggtatccc agggtccttg ctcaagtttt	1250
ctttgaaaag gagggctgga atggtacatc acataggcaa gtcctgcctc	1300

-continued

gtatttaggc tttgctgct tgggtgatg taagggaat tgaagactt	1350
gccattcaa aatgatcttt accgtggcct gcccctgct tatggcccc	1400
agcatttaca gtaacttggt aatgttaagt atcatctctt atctaaatat	1450
taaaagataa gtcaacccaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1500
aaaaaaaa	1508

<210> SEQ ID NO 20

<211> LENGTH: 319

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 20

Met	Leu	Phe	Trp	Val	Leu	Gly	Leu	Leu	Ile	Leu	Cys	Gly	Phe	Leu	
1				5					10					15	
Trp	Thr	Arg	Lys	Gly	Lys	Leu	Lys	Ile	Glu	Asp	Ile	Thr	Asp	Lys	
			20						25					30	
Tyr	Ile	Phe	Ile	Thr	Gly	Cys	Asp	Ser	Gly	Phe	Gly	Asn	Leu	Ala	
			35						40					45	
Ala	Arg	Thr	Phe	Asp	Lys	Lys	Gly	Phe	His	Val	Ile	Ala	Ala	Cys	
			50						55					60	
Leu	Thr	Glu	Ser	Gly	Ser	Thr	Ala	Leu	Lys	Ala	Glu	Thr	Ser	Glu	
			65						70					75	
Arg	Leu	Arg	Thr	Val	Leu	Leu	Asp	Val	Thr	Asp	Pro	Glu	Asn	Val	
			80						85					90	
Lys	Arg	Thr	Ala	Gln	Trp	Val	Lys	Asn	Gln	Val	Gly	Glu	Lys	Gly	
			95						100					105	
Leu	Trp	Gly	Leu	Ile	Asn	Asn	Ala	Gly	Val	Pro	Gly	Val	Leu	Ala	
			110						115					120	
Pro	Thr	Asp	Trp	Leu	Thr	Leu	Glu	Asp	Tyr	Arg	Glu	Pro	Ile	Glu	
			125						130					135	
Val	Asn	Leu	Phe	Gly	Leu	Ile	Ser	Val	Thr	Leu	Asn	Met	Leu	Pro	
			140						145					150	
Leu	Val	Lys	Lys	Ala	Gln	Gly	Arg	Val	Ile	Asn	Val	Ser	Ser	Val	
			155						160					165	
Gly	Gly	Arg	Leu	Ala	Ile	Val	Gly	Gly	Gly	Tyr	Thr	Pro	Ser	Lys	
			170						175					180	
Tyr	Ala	Val	Glu	Gly	Phe	Asn	Asp	Ser	Leu	Arg	Arg	Asp	Met	Lys	
			185						190					195	
Ala	Phe	Gly	Val	His	Val	Ser	Cys	Ile	Glu	Pro	Gly	Leu	Phe	Lys	
			200						205					210	
Thr	Asn	Leu	Ala	Asp	Pro	Val	Lys	Val	Ile	Glu	Lys	Lys	Leu	Ala	
			215						220					225	
Ile	Trp	Glu	Gln	Leu	Ser	Pro	Asp	Ile	Lys	Gln	Gln	Tyr	Gly	Glu	
			230						235					240	
Gly	Tyr	Ile	Glu	Lys	Ser	Leu	Asp	Lys	Leu	Lys	Gly	Asn	Lys	Ser	
			245						250					255	
Tyr	Val	Asn	Met	Asp	Leu	Ser	Pro	Val	Val	Glu	Cys	Met	Asp	His	
			260						265					270	
Ala	Leu	Thr	Ser	Leu	Phe	Pro	Lys	Thr	His	Tyr	Ala	Ala	Gly	Lys	
			275						280					285	
Asp	Ala	Lys	Ile	Phe	Trp	Ile	Pro	Leu	Ser	His	Met	Pro	Ala	Ala	

-continued

290													295			300		
Leu	Gln	Asp	Phe	Leu	Leu	Leu	Lys	Gln	Lys	Ala	Glu	Leu	Ala	Asn				
				305						310					315			
Pro Lys Ala Val																		
<210> SEQ ID NO 21																		
<211> LENGTH: 1849																		
<212> TYPE: DNA																		
<213> ORGANISM: Homo Sapien																		
<400> SEQUENCE: 21																		
ctgagggcggg ggtagcatgg aggggggagag tacgtcggcg gtgctctcgg																50		
gctttgtgct cggcgcactc gctttccagc acctcaacac ggactcggac																100		
acggaaggtt ttcttcttgg ggaagtaaaa ggtgaagcca agaacagcat																150		
tactgattcc caaatggatg atgttgaagt tgtttataca attgacattc																200		
agaaatatat tccatgctat cagcttttta gcttttataa ttcttcaggc																250		
gaagtaaatg agcaagcact gaagaaaata ttatcaaatg tcaaaaagaa																300		
tgtgttaggt tggtaacaaat tccgtcgtca ttcagatcag atcatgacgt																350		
ttagagagag gctgcttcac aaaaacttgc aggagcattt ttcaaaccaa																400		
gacctgtgtt ttctgctatt aacaccaagt ataataacag aaagctgctc																450		
tactcatcga ctggaacatt ccttatataa acctcaaaaa ggaacttttc																500		
acagggatcc tttagtgggt gccaatctgg gcatgtctga acaactgggt																550		
tataaaactg tatcaggttc ctgtatgtcc actggtttta gccgagcagt																600		
acaaacacac agctctaaat tttttgaaga agatggatcc ttaaaggagg																650		
tacataagat aaatgaaatg tatgcttcat tacaagagga attaaagagt																700		
atatgcaaaa aagtgaaga cagtgaacaa gcagtagata aactagtaaa																750		
ggatgtaaac agattaaaac gagaaattga gaaaaggaga ggagcacaga																800		
ttcaggcagc aagagagaag aacatccaaa aagaccctca ggagaacatt																850		
tttctttgtc aggcattacg gacctttttt ccaaattctg aatttcttca																900		
ttcatgtgtt atgtctttaa aaaatagaca tgtttctaaa agtagctgta																950		
actacaacca ccatctcgat gtagtagaca atctgacctt aatggtagaa																1000		
cacactgaca ttcttgaagc tagtccagct agtacaccac aaatcattaa																1050		
gcataaagcc ttagacttag atgacagatg gcaattcaag agatctcggg																1100		
tgttagatac acaagacaaa cgatctaaag caaatactgg tagtagtaac																1150		
caagataaag catccaaaat gagcagccca gaaacagatg aagaaattga																1200		
aaagatgaag ggttttggtg aatattcacg gtctcctaca ttttgatcct																1250		
tttaacctta caaggagatt tttttatttg gctgatgggt aaagccaaac																1300		
atctctattg tttttactat gttgagctac ttgcagtaag ttcatttgtt																1350		
tttactatgt tcacctgttt gcagtaatac acagataact cttagtgcac																1400		
ttacttcaca aagtactttt tcaaacatca gatgctttta ttccaaacc																1450		
tttttttcac ctttcactaa gttgttgagg ggaaggctta cacagacaca																1500		
ttcttttagaa ttggaaaagt gagaccaggc acagtggctc acacctgtaa																1550		

-continued

tcccagcact tagggaagac aagtcaggag gattgattga agctaggagt	1600
tagagaccag cctgggcaac gtattgagac catgtctatt aaaaaataaa	1650
atggaaaagc aagaatagcc ttattttcaa aatatggaaa gaaatttata	1700
tgaaaaatta tctgagtcac taaaattctc cttaagtgat acttttttag	1750
aagtacatta tggctagagt tgccagataa aatgctggat atcatgcaat	1800
aaatttgcaa aacatcatct aaaattttaa aaaaaaaaaa aaaaaaaaaa	1849

<210> SEQ ID NO 22

<211> LENGTH: 409

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 22

Met	Glu	Gly	Glu	Ser	Thr	Ser	Ala	Val	Leu	Ser	Gly	Phe	Val	Leu	1	5	10	15
Gly	Ala	Leu	Ala	Phe	Gln	His	Leu	Asn	Thr	Asp	Ser	Asp	Thr	Glu	20	25	30	
Gly	Phe	Leu	Leu	Gly	Glu	Val	Lys	Gly	Glu	Ala	Lys	Asn	Ser	Ile	35	40	45	
Thr	Asp	Ser	Gln	Met	Asp	Asp	Val	Glu	Val	Val	Tyr	Thr	Ile	Asp	50	55	60	
Ile	Gln	Lys	Tyr	Ile	Pro	Cys	Tyr	Gln	Leu	Phe	Ser	Phe	Tyr	Asn	65	70	75	
Ser	Ser	Gly	Glu	Val	Asn	Glu	Gln	Ala	Leu	Lys	Lys	Ile	Leu	Ser	80	85	90	
Asn	Val	Lys	Lys	Asn	Val	Val	Gly	Trp	Tyr	Lys	Phe	Arg	Arg	His	95	100	105	
Ser	Asp	Gln	Ile	Met	Thr	Phe	Arg	Glu	Arg	Leu	Leu	His	Lys	Asn	110	115	120	
Leu	Gln	Glu	His	Phe	Ser	Asn	Gln	Asp	Leu	Val	Phe	Leu	Leu	Leu	125	130	135	
Thr	Pro	Ser	Ile	Ile	Thr	Glu	Ser	Cys	Ser	Thr	His	Arg	Leu	Glu	140	145	150	
His	Ser	Leu	Tyr	Lys	Pro	Gln	Lys	Gly	Leu	Phe	His	Arg	Val	Pro	155	160	165	
Leu	Val	Val	Ala	Asn	Leu	Gly	Met	Ser	Glu	Gln	Leu	Gly	Tyr	Lys	170	175	180	
Thr	Val	Ser	Gly	Ser	Cys	Met	Ser	Thr	Gly	Phe	Ser	Arg	Ala	Val	185	190	195	
Gln	Thr	His	Ser	Ser	Lys	Phe	Phe	Glu	Glu	Asp	Gly	Ser	Leu	Lys	200	205	210	
Glu	Val	His	Lys	Ile	Asn	Glu	Met	Tyr	Ala	Ser	Leu	Gln	Glu	Glu	215	220	225	
Leu	Lys	Ser	Ile	Cys	Lys	Lys	Val	Glu	Asp	Ser	Glu	Gln	Ala	Val	230	235	240	
Asp	Lys	Leu	Val	Lys	Asp	Val	Asn	Arg	Leu	Lys	Arg	Glu	Ile	Glu	245	250	255	
Lys	Arg	Arg	Gly	Ala	Gln	Ile	Gln	Ala	Ala	Arg	Glu	Lys	Asn	Ile	260	265	270	
Gln	Lys	Asp	Pro	Gln	Glu	Asn	Ile	Phe	Leu	Cys	Gln	Ala	Leu	Arg				

-continued

275					280					285				
Thr	Phe	Phe	Pro	Asn	Ser	Glu	Phe	Leu	His	Ser	Cys	Val	Met	Ser
290					295					300				
Leu	Lys	Asn	Arg	His	Val	Ser	Lys	Ser	Ser	Cys	Asn	Tyr	Asn	His
305					310					315				
His	Leu	Asp	Val	Val	Asp	Asn	Leu	Thr	Leu	Met	Val	Glu	His	Thr
320					325					330				
Asp	Ile	Pro	Glu	Ala	Ser	Pro	Ala	Ser	Thr	Pro	Gln	Ile	Ile	Lys
335					340					345				
His	Lys	Ala	Leu	Asp	Leu	Asp	Asp	Arg	Trp	Gln	Phe	Lys	Arg	Ser
350					355					360				
Arg	Leu	Leu	Asp	Thr	Gln	Asp	Lys	Arg	Ser	Lys	Ala	Asn	Thr	Gly
365					370					375				
Ser	Ser	Asn	Gln	Asp	Lys	Ala	Ser	Lys	Met	Ser	Ser	Pro	Glu	Thr
380					385					390				
Asp	Glu	Glu	Ile	Glu	Lys	Met	Lys	Gly	Phe	Gly	Glu	Tyr	Ser	Arg
395					400					405				

Ser Pro Thr Phe

<210> SEQ ID NO 23
<211> LENGTH: 2651
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 23

ggcacagcgcg	cgcgggcggag	ggcagagtca	gccgagccga	gtccagcccg	50
acgagcggac	cagcgcaggg	cagcccaagc	agcgcgcagc	gaacgcccgc	100
cgccgccac	accctctgcg	gtccccgcgg	cgcttgccac	ccttcctcc	150
ttccccgcgt	ccccgcctcg	ccggccagtc	agcttgccgg	gttcgtgccc	200
ccgcgaaacc	ccgaggtcac	cagccgcgc	ctctgcttcc	ctgggcccgcg	250
cgccgcctcc	acgcccctct	tctcccctgg	cccgggcgct	ggcaccgggg	300
accgttgct	gacgcgaggc	ccagctctac	ttttcgcccc	gcgtctctc	350
cgctgctcg	cctcttccac	caactccaac	tccttctccc	tccagctcca	400
ctcgctagtc	ccgactccg	ccagccctcg	gcccgctgcc	gtagcgccgc	450
ttcccgctcg	gtcccaaagg	tgggaacgcg	tccgccccgg	cccgcaaccat	500
ggcacgggtc	ggcttgcccc	cgcttctctg	cacctggca	gtgctcagcg	550
ccgcgctgct	ggctgccgag	ctcaagtcga	aaagttgctc	ggaagtgcga	600
cgtcttttacg	tgtccaaagg	cttcaacaag	aacgatgccc	ccctccacga	650
gatcaacggt	gatcatttga	agatctgtcc	ccagggttct	acctgctgct	700
ctcaagagat	ggaggagaag	tacagcctgc	aaagtaaaga	tgatttcaaa	750
agtgtgtgta	gcgaacagtg	caatcatttg	caagctgtct	ttgcttcacg	800
ttacaagaag	tttgatgaat	tcttcaaaga	actacttgaa	aatgcagaga	850
aatccctgaa	tgatatgttt	gtgaagacat	atggccatth	atacatgcaa	900
aattctgagc	tatttaaaga	tctcttcgta	gagttgaaac	gttactacgt	950
ggtgggaaat	gtgaacctgg	aagaatgct	aatgacttc	tgggctcgcc	1000

-continued

tcctggagcg gatgttccgc ctggtgaact cccagtacca ctttacagat	1050
gagtatctgg aatgtgtgag caagtatacg gagcagctga agcccttcgg	1100
agatgtccct cgcaaattga agctccaggt tactcgtgct tttgtagcag	1150
cccgtacttt cgctcaaggc ttagcggttg cgggagatgt cgtgagcaag	1200
gtctccgtgg taaacccac agcccagtg acccatgccc tgttgaagat	1250
gatctactgc tcccactgcc ggggtctcgt gactgtgaag ccatgttaca	1300
actactgctc aaacatcatg agaggctgtt tggccaacca aggggatctc	1350
gattttgaat ggaacaattt catagatgct atgctgatgg tggcagagag	1400
gctagagggg cctttcaaca ttgaatcggg catggatccc atcgaagtga	1450
agattttctga tgctattatg aacatgcagg ataatagtgt tcaagtgtct	1500
cagaaggttt tccagggatg tggaccccc aagcccctcc cagctggacg	1550
aatttctcgt tccatctctg aaagtgcctt cagtgtctgc ttcagaccac	1600
atcaccccca ggaacgcca accacagcag ctggcactag tttggaccga	1650
ctggttactg atgtcaagga gaaactgaaa caggccaaga aattctggtc	1700
ctcccttcgg agcaacgttt gcaacgatga gaggatggct gcaggaaacg	1750
gcaatgagga tgactgttgg aatgggaaa gcaaaagcag gtacctgttt	1800
gcagtgacag gaaatggatt agccaaccag ggcaacaacc cagagggtcca	1850
ggttgacacc agcaaaccag acatactgat ccttcgtcaa atcatggctc	1900
ttcgaagtga gaccagcaag atgaagaatg catacaatgg gaacgacgtg	1950
gacttctttg atatcagtga tgaaagtagt ggagaaggaa gtggaagtgg	2000
ctgtgagtat cagcagtgcc cttcagagtt tgactacaat gccactgacc	2050
atgtcgggaa gagtgccaat gagaaagccg acagtgtctg tgtccgtcct	2100
ggggcacagg cctacctcct cactgtcttc tgcactctgt tcttggttat	2150
gcagagagag tggagataat tctcaaactc tgagaaaaag tgttcacaa	2200
aaagttaaaa ggcaccagtt atcaactttc taccatccta gtgactttgc	2250
tttttaaatg aatggacaac aatgtacagt ttttactatg tggccactgg	2300
tttaagaagt gctgactttg ttttctcatt cagttttggg aggaaaaggg	2350
actgtgcatt gagtgggtc ctgctcccc aaaccatgtt aaacgtggct	2400
aacagtgtag gtacagaact atagttagtt gtgcatttgt gattttatca	2450
ctctattatt tgtttgtatg ttttttctc atttcgtttg tgggtttttt	2500
tttccaaactg tgatctcgcc ttgtttctta caagcaaacc agggtcctt	2550
cttggcacgt aacatgtacg tatttctgaa atattaaata gctgtacaga	2600
agcagggtttt atttatcatg ttatcttatt aaaagaaaaa gcccaaaaag	2650
c	2651

<210> SEQ ID NO 24

<211> LENGTH: 556

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 24

-continued

Met	Ala	Arg	Phe	Gly	Leu	Pro	Ala	Leu	Leu	Cys	Thr	Leu	Ala	Val
1				5					10					15
Leu	Ser	Ala	Ala	Leu	Leu	Ala	Ala	Glu	Leu	Lys	Ser	Lys	Ser	Cys
				20					25					30
Ser	Glu	Val	Arg	Arg	Leu	Tyr	Val	Ser	Lys	Gly	Phe	Asn	Lys	Asn
				35					40					45
Asp	Ala	Pro	Leu	His	Glu	Ile	Asn	Gly	Asp	His	Leu	Lys	Ile	Cys
				50					55					60
Pro	Gln	Gly	Ser	Thr	Cys	Cys	Ser	Gln	Glu	Met	Glu	Glu	Lys	Tyr
				65					70					75
Ser	Leu	Gln	Ser	Lys	Asp	Asp	Phe	Lys	Ser	Val	Val	Ser	Glu	Gln
				80					85					90
Cys	Asn	His	Leu	Gln	Ala	Val	Phe	Ala	Ser	Arg	Tyr	Lys	Lys	Phe
				95					100					105
Asp	Glu	Phe	Phe	Lys	Glu	Leu	Leu	Glu	Asn	Ala	Glu	Lys	Ser	Leu
				110					115					120
Asn	Asp	Met	Phe	Val	Lys	Thr	Tyr	Gly	His	Leu	Tyr	Met	Gln	Asn
				125					130					135
Ser	Glu	Leu	Phe	Lys	Asp	Leu	Phe	Val	Glu	Leu	Lys	Arg	Tyr	Tyr
				140					145					150
Val	Val	Gly	Asn	Val	Asn	Leu	Glu	Glu	Met	Leu	Asn	Asp	Phe	Trp
				155					160					165
Ala	Arg	Leu	Leu	Glu	Arg	Met	Phe	Arg	Leu	Val	Asn	Ser	Gln	Tyr
				170					175					180
His	Phe	Thr	Asp	Glu	Tyr	Leu	Glu	Cys	Val	Ser	Lys	Tyr	Thr	Glu
				185					190					195
Gln	Leu	Lys	Pro	Phe	Gly	Asp	Val	Pro	Arg	Lys	Leu	Lys	Leu	Gln
				200					205					210
Val	Thr	Arg	Ala	Phe	Val	Ala	Ala	Arg	Thr	Phe	Ala	Gln	Gly	Leu
				215					220					225
Ala	Val	Ala	Gly	Asp	Val	Val	Ser	Lys	Val	Ser	Val	Val	Asn	Pro
				230					235					240
Thr	Ala	Gln	Cys	Thr	His	Ala	Leu	Leu	Lys	Met	Ile	Tyr	Cys	Ser
				245					250					255
His	Cys	Arg	Gly	Leu	Val	Thr	Val	Lys	Pro	Cys	Tyr	Asn	Tyr	Cys
				260					265					270
Ser	Asn	Ile	Met	Arg	Gly	Cys	Leu	Ala	Asn	Gln	Gly	Asp	Leu	Asp
				275					280					285
Phe	Glu	Trp	Asn	Asn	Phe	Ile	Asp	Ala	Met	Leu	Met	Val	Ala	Glu
				290					295					300
Arg	Leu	Glu	Gly	Pro	Phe	Asn	Ile	Glu	Ser	Val	Met	Asp	Pro	Ile
				305					310					315
Asp	Val	Lys	Ile	Ser	Asp	Ala	Ile	Met	Asn	Met	Gln	Asp	Asn	Ser
				320					325					330
Val	Gln	Val	Ser	Gln	Lys	Val	Phe	Gln	Gly	Cys	Gly	Pro	Pro	Lys
				335					340					345
Pro	Leu	Pro	Ala	Gly	Arg	Ile	Ser	Arg	Ser	Ile	Ser	Glu	Ser	Ala
				350					355					360
Phe	Ser	Ala	Arg	Phe	Arg	Pro	His	His	Pro	Glu	Glu	Arg	Pro	Thr
				365					370					375
Thr	Ala	Ala	Gly	Thr	Ser	Leu	Asp	Arg	Leu	Val	Thr	Asp	Val	Lys

-continued

380										385										390									
Glu	Lys	Leu	Lys	Gln	Ala	Lys	Lys	Phe	Trp	Ser	Ser	Leu	Pro	Ser															
				395					400					405															
Asn	Val	Cys	Asn	Asp	Glu	Arg	Met	Ala	Ala	Gly	Asn	Gly	Asn	Glu															
				410					415					420															
Asp	Asp	Cys	Trp	Asn	Gly	Lys	Gly	Lys	Ser	Arg	Tyr	Leu	Phe	Ala															
				425					430					435															
Val	Thr	Gly	Asn	Gly	Leu	Ala	Asn	Gln	Gly	Asn	Asn	Pro	Glu	Val															
				440					445					450															
Gln	Val	Asp	Thr	Ser	Lys	Pro	Asp	Ile	Leu	Ile	Leu	Arg	Gln	Ile															
				455					460					465															
Met	Ala	Leu	Arg	Val	Met	Thr	Ser	Lys	Met	Lys	Asn	Ala	Tyr	Asn															
				470					475					480															
Gly	Asn	Asp	Val	Asp	Phe	Phe	Asp	Ile	Ser	Asp	Glu	Ser	Ser	Gly															
				485					490					495															
Glu	Gly	Ser	Gly	Ser	Gly	Cys	Glu	Tyr	Gln	Gln	Cys	Pro	Ser	Glu															
				500					505					510															
Phe	Asp	Tyr	Asn	Ala	Thr	Asp	His	Ala	Gly	Lys	Ser	Ala	Asn	Glu															
				515					520					525															
Lys	Ala	Asp	Ser	Ala	Gly	Val	Arg	Pro	Gly	Ala	Gln	Ala	Tyr	Leu															
				530					535					540															
Leu	Thr	Val	Phe	Cys	Ile	Leu	Phe	Leu	Val	Met	Gln	Arg	Glu	Trp															
				545					550					555															
Arg																													

<210> SEQ ID NO 25
<211> LENGTH: 870
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 25

ctcgccctca aatgggaacg ctggcctggg actaaagcat agaccaccag	50
gctgagtatc ctgacctgag tcatccccag ggatcaggag cctccagcag	100
ggaaccttcc attatattct tcaagcaact tacagctgca ccgacagttg	150
cgatgaaagt tctaattctt tccctcctcc tgttgctgcc actaatgctg	200
atgtccatgg tctctagcag cctgaatcca ggggtcgcca gaggccacag	250
ggaccgaggc caggcttcta ggagatggct ccaggaaggc ggccaagaat	300
gtgagtgcaa agattgggtc ctgagagccc cgagaagaaa attcatgaca	350
gtgtctgggc tgccaaagaa gcagtgcccc tgtgatcatt tcaagggcaa	400
tgtgaagaaa acaagacacc aaaggcacca cagaaagcca aacaagcatt	450
ccagagcctg ccagcaattt ctcaaacaat gtcagctaag aagctttgct	500
ctgcctttgt aggagctctg agcgcccact cttccaatta aacattctca	550
gccaaagaag cagtgagcac acctaccaga cactcttctt ctcccacctc	600
actctcccac tgtaccaccc cctaaatcat tccagtgtct tcaaaaagca	650
tgtttttcaa gatcattttg ttgttgctc tctctagtgt cttcttctct	700
cgtcagtcct agcctgtgcc ctccccttac ccaggcttag gcttaattac	750
ctgaaagatt ccaggaaact gtagcttcct agctagtgtc atttaacctt	800

-continued

aaatgcaatc aggaaagtag caaacagaag tcaataaata tttttaaatg	850
tcaaaaaaaaa aaaaaaaaaa	870

<210> SEQ ID NO 26
<211> LENGTH: 119
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 26

Met Lys Val Leu Ile Ser Ser Leu Leu Leu Leu Leu Pro Leu Met	1 5 10 15
Leu Met Ser Met Val Ser Ser Ser Leu Asn Pro Gly Val Ala Arg	20 25 30
Gly His Arg Asp Arg Gly Gln Ala Ser Arg Arg Trp Leu Gln Glu	35 40 45
Gly Gly Gln Glu Cys Glu Cys Lys Asp Trp Phe Leu Arg Ala Pro	50 55 60
Arg Arg Lys Phe Met Thr Val Ser Gly Leu Pro Lys Lys Gln Cys	65 70 75
Pro Cys Asp His Phe Lys Gly Asn Val Lys Lys Thr Arg His Gln	80 85 90
Arg His His Arg Lys Pro Asn Lys His Ser Arg Ala Cys Gln Gln	95 100 105
Phe Leu Lys Gln Cys Gln Leu Arg Ser Phe Ala Leu Pro Leu	110 115

<210> SEQ ID NO 27
<211> LENGTH: 1371
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 27

ggacgccagc gcctgcagag gctgagcagg gaaaaagcca gtgccccagc	50
ggaagcacag ctgagagctg gtctgccatg gacatcctgg tcccactcct	100
gcagctgctg gtgctgcttc ttacctgccc cctgcacctc atggctctgc	150
tgggctgctg gcagcccctg tgcaaaagct acttccccta cctgatggcc	200
gtgctgactc ccaagagcaa ccgcaagatg gagagcaaga aacgggagct	250
cttcagccag ataaaggggc ttacaggagc ctccgggaaa gtggccctac	300
tggagctggg ctgcggaacc ggagccaact ttcagttcta cccaccgggc	350
tgcagggtca cctgcctaga cccaaatccc cactttgaga agttcctgac	400
aaagagcatg gctgagaaca ggcacctcca atatgagcgg tttgtggtgg	450
ctcctggaga ggacatgaga cagctggctg atggctccat ggatgtggtg	500
gtctgcactc tgggtgctgt ctctgtgcag agcccaagga aggtcctgca	550
ggaggtccgg agagtactga gaccgggagg tgtgctcttt ttctgggagc	600
atgtggcaga accatatgga agctgggcct tcatgtggca gcaagttttc	650
gagcccacct ggaacacat tggggatggc tgctgcctca ccagagagac	700
ctggaaggat cttgagaacg cccagttcto cgaatatcaa atggaacgac	750
agccccctcc cttgaagtgg ctacctgttg ggccccacat catgggaaag	800

-continued

gctgtcaaac aatctttccc aagctccaag gcaactcattt gtccttccc	850
cagcctccaa ttagaacaag ccaccacca gcctatctat ctccactga	900
gagggaccta gcagaatgag agaagacatt catgtaccac ctactagtcc	950
ctctctcccc aacctctgcc agggcaatct ctaacttcaa tccgccttc	1000
gacagtga aaagctctact tctacgctga cccagggagg aaacactagg	1050
accctgttgt atcctcaact gcaagtttct ggactagtct cccaacgttt	1100
gcctcccaat gttgtccctt tccttcgttc ccatggtaaa gtcctctctg	1150
ctttctcctt gaggctacac ccatgctgtct cttaggaactg gtcacaaaag	1200
tcatggtgcc tgcacccctg ccaagccccc ctgaccctct ctccccacta	1250
ccaccttctt cctgagctgg gggcaccagg gagaatcaga gatgctgggg	1300
atgccagagc aagactcaaa gaggcagagg ttttgttctc aaatatTTTT	1350
taataaatag acgaaaccac g	1371

<210> SEQ ID NO 28

<211> LENGTH: 277

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 28

Met Asp Ile Leu Val Pro Leu Leu Gln Leu Leu Val Leu Leu Leu	1	5	10	15
Thr Leu Pro Leu His Leu Met Ala Leu Leu Gly Cys Trp Gln Pro	20	25	30	
Leu Cys Lys Ser Tyr Phe Pro Tyr Leu Met Ala Val Leu Thr Pro	35	40	45	
Lys Ser Asn Arg Lys Met Glu Ser Lys Lys Arg Glu Leu Phe Ser	50	55	60	
Gln Ile Lys Gly Leu Thr Gly Ala Ser Gly Lys Val Ala Leu Leu	65	70	75	
Glu Leu Gly Cys Gly Thr Gly Ala Asn Phe Gln Phe Tyr Pro Pro	80	85	90	
Gly Cys Arg Val Thr Cys Leu Asp Pro Asn Pro His Phe Glu Lys	95	100	105	
Phe Leu Thr Lys Ser Met Ala Glu Asn Arg His Leu Gln Tyr Glu	110	115	120	
Arg Phe Val Val Ala Pro Gly Glu Asp Met Arg Gln Leu Ala Asp	125	130	135	
Gly Ser Met Asp Val Val Val Cys Thr Leu Val Leu Cys Ser Val	140	145	150	
Gln Ser Pro Arg Lys Val Leu Gln Glu Val Arg Arg Val Leu Arg	155	160	165	
Pro Gly Gly Val Leu Phe Phe Trp Glu His Val Ala Glu Pro Tyr	170	175	180	
Gly Ser Trp Ala Phe Met Trp Gln Gln Val Phe Glu Pro Thr Trp	185	190	195	
Lys His Ile Gly Asp Gly Cys Cys Leu Thr Arg Glu Thr Trp Lys	200	205	210	
Asp Leu Glu Asn Ala Gln Phe Ser Glu Ile Gln Met Glu Arg Gln	215	220	225	

-continued

Pro Pro Pro Leu Lys Trp Leu Pro Val Gly Pro His Ile Met Gly
230 235 240
Lys Ala Val Lys Gln Ser Phe Pro Ser Ser Lys Ala Leu Ile Cys
245 250 255
Ser Phe Pro Ser Leu Gln Leu Glu Gln Ala Thr His Gln Pro Ile
260 265 270
Tyr Leu Pro Leu Arg Gly Thr
275

<210> SEQ ID NO 29
<211> LENGTH: 494
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 29
caatgtttgc cstatccacct cccccaagcc cctttaccta tgctgctgct 50
aacgctgctg ctgctgctgc tgctgcttaa aggctcatgc ttggagtggg 100
gactggtcgg tgcccagaaa gtctcttctg ccaactgacgc ccccatcagg 150
gattgggcct tctttccccc ttcctttctg tgtctcctgc ctcatcggcc 200
tgccatgacc tgcagccaag ccagccccg tggggaaggg gagaaagtgg 250
gggatggcta agaaagctgg gagataggga acagaagagg gtagtgggtg 300
ggctaggggg gctgccttat ttaaagtggg tgtttatgat tcttatacta 350
atttatacaa agatattaag gccctgttca ttaagaaatt gttcccttcc 400
cctgtgttca atgtttgtaa agattgttct gtgtaaatat gtctttataa 450
taaacagtta aaagctgaaa aaaaaaaaaa aaaaaaaaaa aaaa 494

<210> SEQ ID NO 30
<211> LENGTH: 73
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 30
Met Leu Leu Leu Thr Leu Leu Leu Leu Leu Leu Lys Gly
1 5 10 15
Ser Cys Leu Glu Trp Gly Leu Val Gly Ala Gln Lys Val Ser Ser
20 25 30
Ala Thr Asp Ala Pro Ile Arg Asp Trp Ala Phe Phe Pro Pro Ser
35 40 45
Phe Leu Cys Leu Leu Pro His Arg Pro Ala Met Thr Cys Ser Gln
50 55 60
Ala Gln Pro Arg Gly Glu Gly Glu Lys Val Gly Asp Gly
65 70

<210> SEQ ID NO 31
<211> LENGTH: 1660
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 31
gtttgaattc cttcaactat acccacagtc caaaagcaga ctactgtgt 50
cccaggctac cagttcctcc aagcaagtca tttcccttat ttaaccgatg 100

-continued

tgtccctcaa acacctgagt gctactccct atttgcattct gttttgataa	150
atgatgttga caccctccac cgaattctaa gtggaatcat gtcgggaaga	200
gatacaatcc ttggcctgtg tatcctcgca ttagccttgt ctttggccat	250
gatgtttacc ttcagattca tcaccaccct tctgggtcac attttcattt	300
cattggttat tttgggattg ttgtttgtct gcggtgtttt atggtggctg	350
tattatgact ataccaacga cctcagcata gaattggaca cagaaagga	400
aaatatgaag tgcgtgctgg ggtttgcctat cgtatccaca ggcattcacg	450
cagtgcctgct cgtcttgatt tttgttctca gaaagagaat aaaattgaca	500
gttgagcttt tccaaatcac aaataaagcc atcagcagtg ctcccttcct	550
gctgttccag ccactgtgga catttgccat cctcattttc ttctgggtcc	600
ttctgggtggc tgtgctgctg agcctgggaa ctgcaggagc tgcccaggtt	650
atggaaggcg gccaaagtga atataagccc ctttcgggca ttcggtacat	700
gtggtcgtac catttaattg gcctcatctg gactagttaa ttcattcctg	750
cgtgccagca aatgactata gctggggcag tggttacttg ttatttcaac	800
agaagtaaaa atgatcctcc tgatcatccc atcctttcgt ctctctccat	850
ttctctcttc taccatcaag gaaccgttgt gaaaggtca tttttaatct	900
ctgtggtgag gattccgaga atcattgtca tgtacatgca aaacgcactg	950
aaagaacagc agcatggtgc attgtccagg tacctgttcc gatgctgcta	1000
ctgctgtttc tgggtgtcttg acaaatacct gctccatctc aaccagaatg	1050
catatactac aactgctatt aatgggacag atttctgtac atcagcaaaa	1100
gatgcattca aaatcttgtc caagaactca agtcacttta catctattaa	1150
ctgctttgga gacttcataa tttttctagg aaagggtta gtggtgtgtt	1200
tcactgtttt tggaggactc atggctttta actacaatcg ggcattccag	1250
gtgtgggcag tccctctgtt attggtagct tttttgcct acttagtagc	1300
ccatagtttt ttatctgtgt ttgaaactgt gctggatgca cttttcctgt	1350
gttttgcgtg tgatctggaa acaaatgatg gatcgtcaga aaagccctac	1400
tttatggatc aagaatttct gagtttcgta aaaaggagca acaaattaaa	1450
caatgcaagg gcacagcagg acaagcactc attaaggaat gaggagggaa	1500
cagaactcca ggccattgtg agatagatac ccatttaggt atctgtacct	1550
ggaaaacatt tccttctaag agccatttac agaatagaag atgagaccac	1600
tagagaaaaa ttagtgaatt tttttttaa agacctata aacctattc	1650
ttcctcaaaa	1660

<210> SEQ ID NO 32
<211> LENGTH: 445
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 32
Met Ser Gly Arg Asp Thr Ile Leu Gly Leu Cys Ile Leu Ala Leu
1 5 10 15
Ala Leu Ser Leu Ala Met Met Phe Thr Phe Arg Phe Ile Thr Thr

-continued

20					25					30				
Leu	Leu	Val	His	Ile	Phe	Ile	Ser	Leu	Val	Ile	Leu	Gly	Leu	Leu
				35					40					45
Phe	Val	Cys	Gly	Val	Leu	Trp	Trp	Leu	Tyr	Tyr	Asp	Tyr	Thr	Asn
				50					55					60
Asp	Leu	Ser	Ile	Glu	Leu	Asp	Thr	Glu	Arg	Glu	Asn	Met	Lys	Cys
				65					70					75
Val	Leu	Gly	Phe	Ala	Ile	Val	Ser	Thr	Gly	Ile	Thr	Ala	Val	Leu
				80					85					90
Leu	Val	Leu	Ile	Phe	Val	Leu	Arg	Lys	Arg	Ile	Lys	Leu	Thr	Val
				95					100					105
Glu	Leu	Phe	Gln	Ile	Thr	Asn	Lys	Ala	Ile	Ser	Ser	Ala	Pro	Phe
				110					115					120
Leu	Leu	Phe	Gln	Pro	Leu	Trp	Thr	Phe	Ala	Ile	Leu	Ile	Phe	Phe
				125					130					135
Trp	Val	Leu	Trp	Val	Ala	Val	Leu	Leu	Ser	Leu	Gly	Thr	Ala	Gly
				140					145					150
Ala	Ala	Gln	Val	Met	Glu	Gly	Gly	Gln	Val	Glu	Tyr	Lys	Pro	Leu
				155					160					165
Ser	Gly	Ile	Arg	Tyr	Met	Trp	Ser	Tyr	His	Leu	Ile	Gly	Leu	Ile
				170					175					180
Trp	Thr	Ser	Glu	Phe	Ile	Leu	Ala	Cys	Gln	Gln	Met	Thr	Ile	Ala
				185					190					195
Gly	Ala	Val	Val	Thr	Cys	Tyr	Phe	Asn	Arg	Ser	Lys	Asn	Asp	Pro
				200					205					210
Pro	Asp	His	Pro	Ile	Leu	Ser	Ser	Leu	Ser	Ile	Leu	Phe	Phe	Tyr
				215					220					225
His	Gln	Gly	Thr	Val	Val	Lys	Gly	Ser	Phe	Leu	Ile	Ser	Val	Val
				230					235					240
Arg	Ile	Pro	Arg	Ile	Ile	Val	Met	Tyr	Met	Gln	Asn	Ala	Leu	Lys
				245					250					255
Glu	Gln	Gln	His	Gly	Ala	Leu	Ser	Arg	Tyr	Leu	Phe	Arg	Cys	Cys
				260					265					270
Tyr	Cys	Cys	Phe	Trp	Cys	Leu	Asp	Lys	Tyr	Leu	Leu	His	Leu	Asn
				275					280					285
Gln	Asn	Ala	Tyr	Thr	Thr	Thr	Ala	Ile	Asn	Gly	Thr	Asp	Phe	Cys
				290					295					300
Thr	Ser	Ala	Lys	Asp	Ala	Phe	Lys	Ile	Leu	Ser	Lys	Asn	Ser	Ser
				305					310					315
His	Phe	Thr	Ser	Ile	Asn	Cys	Phe	Gly	Asp	Phe	Ile	Ile	Phe	Leu
				320					325					330
Gly	Lys	Val	Leu	Val	Val	Cys	Phe	Thr	Val	Phe	Gly	Gly	Leu	Met
				335					340					345
Ala	Phe	Asn	Tyr	Asn	Arg	Ala	Phe	Gln	Val	Trp	Ala	Val	Pro	Leu
				350					355					360
Leu	Leu	Val	Ala	Phe	Phe	Ala	Tyr	Leu	Val	Ala	His	Ser	Phe	Leu
				365					370					375
Ser	Val	Phe	Glu	Thr	Val	Leu	Asp	Ala	Leu	Phe	Leu	Cys	Phe	Ala
				380					385					390
Val	Asp	Leu	Glu	Thr	Asn	Asp	Gly	Ser	Ser	Glu	Lys	Pro	Tyr	Phe
				395					400					405

-continued

Met Asp Gln Glu Phe Leu Ser Phe Val Lys Arg Ser Asn Lys Leu
410 415 420
Asn Asn Ala Arg Ala Gln Gln Asp Lys His Ser Leu Arg Asn Glu
425 430 435
Glu Gly Thr Glu Leu Gln Ala Ile Val Arg
440 445

<210> SEQ ID NO 33
<211> LENGTH: 2773
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 33

gttcgattag ctcctctgag aagaagagaa aaggttcttg gacctctccc	50
tgtttcttcc ttagaataat ttgtatggga tttgtgatgc aggaaagcct	100
aagggaaaaa gaatattcat tctgtgtggt gaaaaatttt tgaaaaaaaa	150
attgccttct tcaaacaagg gtgtcattct gatatttatg aggactgttg	200
ttctcactat gaaggcatct gttattgaaa tgttccttgt tttgctggtg	250
actggagtac attcaaacaa agaaacggca aagaagatta aaaggcccaa	300
gttcactgtg cctcagatca actgcgatgt caaagccgga aagatcatcg	350
atcctgagtt cattgtgaaa tgtccagcag gatccaaga ccccaaatac	400
catgtttatg gcaactgacgt gtatgcatcc tactccagtg tgtgtggcgc	450
tgccgtacac agtggtgtgc ttgataattc aggagggaaa atacttgttc	500
ggaaggttgc tggacagtct gggtacaaag ggagttattc caacggtgtc	550
caatcgttat ccctaccacg atggagagaa tcctttatcg tcttagaaaag	600
taaacccaaa aagggtgtaa cctaccatc agctcttaca tactcatcat	650
cgaaaagtcc agctgcccaa gcaggtgaga ccacaaaagc ctatcagagg	700
ccacctattc cagggacaac tgcacagcgg gtcactctga tgcagcttct	750
ggctgtcact gtagctgtgg ccacccccac caccttgcca aggccatccc	800
cttctgtctg ttctaccacc agcatcccca gaccacaatc agtgggccac	850
aggagccagg agatggatct ctggtccact gccacctaca caagcagcca	900
aaacaggccc agagctgac caggtatcca aaggcaagat ccttcaggag	950
ctgccttcca gaaacctggt ggagcggatg tcagcctggg acttgttcca	1000
aaagaagaat tgagcacaca gtctttggag ccagtatccc tgggagatcc	1050
aaactgcaaa attgacttgt cgtttttaat tgatgggagc accagcattg	1100
gcaaacggcg attccgaatc cagaagcagc tcctggctga tgttgcccaa	1150
gctcttgaca ttggccctgc cggtcactg atgggtgttg tccagtatgg	1200
agacaacctt gctactcact ttaacctcaa gacacacacg aattctcgag	1250
atctgaagac agccatagag aaaattactc agagaggagg actttcta	1300
gtaggtcggg ccatctccct tgtgaccaag aacttctttt ccaaagccaa	1350
tggaacaga agcggggctc ccaatgtggt ggtggtgatg gtggatggct	1400
ggccccagga caaagtggag gaggcttcaa gacttgcgag agagtcagga	1450

-continued

atcaacattt tttcatcac cattgaaggt gctgctgaaa atgagaagca	1500
gtatgtggtg gagcccaact ttgcaaaaca ggccgtgtgc agaacaacg	1550
gcttctactc gctccacgtg cagagctggt ttggcctcca caagaccctg	1600
cagcctctgg tgaagcgggt ctgcgacact gaccgcctgg cctgcagcaa	1650
gacctgcttg aactcggctg acattggctt cgtcatcgac ggctccagca	1700
gtgtggggac gggcaacttc cgcaccgtcc tccagtttgt gaccaacctc	1750
accaaagagt ttgagatttc cgacacggac acgcgcatcg gggccgtgca	1800
gtacacctac gaacagcggc tggagtttgg gttcgacaag tacagcagca	1850
agcctgacat cctcaacgcc atcaagaggg tgggctactg gagtgtggc	1900
accagcacgg gggctgccat caacttcgcc ctggagcagc tttcaagaa	1950
gtccaagccc aacaagagga agttaatgat cctcatcacc gacgggaggt	2000
cctacgacga cgtccggatc ccagccatgg ctgccatct gaagggagt	2050
atcacctatg cgataggcgt tgctgggct gcccaagagg agctagaagt	2100
cattgccact caccgccca gagaccactc cttctttgtg gacgagttt	2150
acaacctcca tcagtatgtc cccaggatca tccagaacat ttgtacagag	2200
ttcaactcac agcctcggaa ctgaattcag agcaggcaga gcaccagcaa	2250
gtgctgcttt actaactgac gtgttgacc accccaccgc ttaatggggc	2300
acgcacgggt catcaagtct tgggcagggc atggagaaac aaatgtctt	2350
ttattattct ttgccatcat gctttttcat attccaaaac ttggagttac	2400
aaagatgac acaaacgtat agaatgagcc aaaaggctac atcatgttga	2450
gggtgctgga gattttacat ttgacaatt gttttcaaaa taaatgttcg	2500
gaatacagtg cagcccttac gacaggctta cgtagagctt ttgtgagatt	2550
tttaagttgt tatttctgat ttgaactctg taaccctcag caagtttcat	2600
ttttgtcatg acaatgtagg aattgctgaa ttaaatgttt agaaggatga	2650
aaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa	2700
aaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa	2750
aaaaaaaa aaaaaaaaa aag	2773

<210> SEQ ID NO 34
<211> LENGTH: 678
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 34

Met Arg Thr Val Val Leu Thr Met Lys Ala Ser Val Ile Glu Met	
1 5 10 15	
Phe Leu Val Leu Leu Val Thr Gly Val His Ser Asn Lys Glu Thr	
20 25 30	
Ala Lys Lys Ile Lys Arg Pro Lys Phe Thr Val Pro Gln Ile Asn	
35 40 45	
Cys Asp Val Lys Ala Gly Lys Ile Ile Asp Pro Glu Phe Ile Val	
50 55 60	
Lys Cys Pro Ala Gly Cys Gln Asp Pro Lys Tyr His Val Tyr Gly	
65 70 75	

-continued

Thr	Asp	Val	Tyr	Ala	Ser	Tyr	Ser	Ser	Val	Cys	Gly	Ala	Ala	Val	80	85	90
His	Ser	Gly	Val	Leu	Asp	Asn	Ser	Gly	Gly	Lys	Ile	Leu	Val	Arg	95	100	105
Lys	Val	Ala	Gly	Gln	Ser	Gly	Tyr	Lys	Gly	Ser	Tyr	Ser	Asn	Gly	110	115	120
Val	Gln	Ser	Leu	Ser	Leu	Pro	Arg	Trp	Arg	Glu	Ser	Phe	Ile	Val	125	130	135
Leu	Glu	Ser	Lys	Pro	Lys	Lys	Gly	Val	Thr	Tyr	Pro	Ser	Ala	Leu	140	145	150
Thr	Tyr	Ser	Ser	Ser	Lys	Ser	Pro	Ala	Ala	Gln	Ala	Gly	Glu	Thr	155	160	165
Thr	Lys	Ala	Tyr	Gln	Arg	Pro	Pro	Ile	Pro	Gly	Thr	Thr	Ala	Gln	170	175	180
Pro	Val	Thr	Leu	Met	Gln	Leu	Leu	Ala	Val	Thr	Val	Ala	Val	Ala	185	190	195
Thr	Pro	Thr	Thr	Leu	Pro	Arg	Pro	Ser	Pro	Ser	Ala	Ala	Ser	Thr	200	205	210
Thr	Ser	Ile	Pro	Arg	Pro	Gln	Ser	Val	Gly	His	Arg	Ser	Gln	Glu	215	220	225
Met	Asp	Leu	Trp	Ser	Thr	Ala	Thr	Tyr	Thr	Ser	Ser	Gln	Asn	Arg	230	235	240
Pro	Arg	Ala	Asp	Pro	Gly	Ile	Gln	Arg	Gln	Asp	Pro	Ser	Gly	Ala	245	250	255
Ala	Phe	Gln	Lys	Pro	Val	Gly	Ala	Asp	Val	Ser	Leu	Gly	Leu	Val	260	265	270
Pro	Lys	Glu	Glu	Leu	Ser	Thr	Gln	Ser	Leu	Glu	Pro	Val	Ser	Leu	275	280	285
Gly	Asp	Pro	Asn	Cys	Lys	Ile	Asp	Leu	Ser	Phe	Leu	Ile	Asp	Gly	290	295	300
Ser	Thr	Ser	Ile	Gly	Lys	Arg	Arg	Phe	Arg	Ile	Gln	Lys	Gln	Leu	305	310	315
Leu	Ala	Asp	Val	Ala	Gln	Ala	Leu	Asp	Ile	Gly	Pro	Ala	Gly	Pro	320	325	330
Leu	Met	Gly	Val	Val	Gln	Tyr	Gly	Asp	Asn	Pro	Ala	Thr	His	Phe	335	340	345
Asn	Leu	Lys	Thr	His	Thr	Asn	Ser	Arg	Asp	Leu	Lys	Thr	Ala	Ile	350	355	360
Glu	Lys	Ile	Thr	Gln	Arg	Gly	Gly	Leu	Ser	Asn	Val	Gly	Arg	Ala	365	370	375
Ile	Ser	Phe	Val	Thr	Lys	Asn	Phe	Phe	Ser	Lys	Ala	Asn	Gly	Asn	380	385	390
Arg	Ser	Gly	Ala	Pro	Asn	Val	Val	Val	Val	Met	Val	Asp	Gly	Trp	395	400	405
Pro	Thr	Asp	Lys	Val	Glu	Glu	Ala	Ser	Arg	Leu	Ala	Arg	Glu	Ser	410	415	420
Gly	Ile	Asn	Ile	Phe	Phe	Ile	Thr	Ile	Glu	Gly	Ala	Ala	Glu	Asn	425	430	435
Glu	Lys	Gln	Tyr	Val	Val	Glu	Pro	Asn	Phe	Ala	Asn	Lys	Ala	Val	440	445	450

-continued

Cys	Arg	Thr	Asn	Gly	Phe	Tyr	Ser	Leu	His	Val	Gln	Ser	Trp	Phe
			455						460					465
Gly	Leu	His	Lys	Thr	Leu	Gln	Pro	Leu	Val	Lys	Arg	Val	Cys	Asp
			470						475					480
Thr	Asp	Arg	Leu	Ala	Cys	Ser	Lys	Thr	Cys	Leu	Asn	Ser	Ala	Asp
			485						490					495
Ile	Gly	Phe	Val	Ile	Asp	Gly	Ser	Ser	Ser	Val	Gly	Thr	Gly	Asn
			500						505					510
Phe	Arg	Thr	Val	Leu	Gln	Phe	Val	Thr	Asn	Leu	Thr	Lys	Glu	Phe
			515						520					525
Glu	Ile	Ser	Asp	Thr	Asp	Thr	Arg	Ile	Gly	Ala	Val	Gln	Tyr	Thr
			530						535					540
Tyr	Glu	Gln	Arg	Leu	Glu	Phe	Gly	Phe	Asp	Lys	Tyr	Ser	Ser	Lys
			545						550					555
Pro	Asp	Ile	Leu	Asn	Ala	Ile	Lys	Arg	Val	Gly	Tyr	Trp	Ser	Gly
			560						565					570
Gly	Thr	Ser	Thr	Gly	Ala	Ala	Ile	Asn	Phe	Ala	Leu	Glu	Gln	Leu
			575						580					585
Phe	Lys	Lys	Ser	Lys	Pro	Asn	Lys	Arg	Lys	Leu	Met	Ile	Leu	Ile
			590						595					600
Thr	Asp	Gly	Arg	Ser	Tyr	Asp	Asp	Val	Arg	Ile	Pro	Ala	Met	Ala
			605						610					615
Ala	His	Leu	Lys	Gly	Val	Ile	Thr	Tyr	Ala	Ile	Gly	Val	Ala	Trp
			620						625					630
Ala	Ala	Gln	Glu	Glu	Leu	Glu	Val	Ile	Ala	Thr	His	Pro	Ala	Arg
			635						640					645
Asp	His	Ser	Phe	Phe	Val	Asp	Glu	Phe	Asp	Asn	Leu	His	Gln	Tyr
			650						655					660
Val	Pro	Arg	Ile	Ile	Gln	Asn	Ile	Cys	Thr	Glu	Phe	Asn	Ser	Gln
			665						670					675
Pro	Arg	Asn												

<210> SEQ ID NO 35
<211> LENGTH: 2095
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 35

ccgagcacag gagattgcct gcgtttagga ggtggctgcg ttgtgggaaa	50
agctatcaag gaagaaattg ccaaaccatg tctttttttc tgttttcaga	100
gtagttcaca acagatctga gtgttttaat taagcatgga atacagaaaa	150
caacaaaaaa cttaagcttt aatttcattc ggaattccac agttttctta	200
gtccctcgga cccgggtgac ctgtttggctc ttcccgctgg ctgctctatc	250
acgtgggtgct ctccgactac tcaccccgag tgtaaagaac cttcggtctg	300
cggtgcttctg agctgctgtg gatggcctcg gctctctgga ctgtccttcc	350
gagtaggatg tcaactagat ccctcaaatg gagcctcctg ctgctgtcac	400
tcctgagttt ctttgtgatg tggtaacctca gccttcccca ctacaatgtg	450
atagaacgcg tgaactggat gtacttctat gagtatgagc cgatttacag	500
acaagacttt cacttcacac ttcgagagca ttcaaactgc tctcatcaaa	550

-continued

```
atccatttct ggtcattctg gtgacctccc acccttcaga tgtgaaagcc      600
aggcaggcca ttagagttac ttggggtgaa aaaaagtctt ggtggggata      650
tgaggttctt acatttttct tattaggcca agaggctgaa aaggaagaca      700
aatgtttggc attgtcctta gaggatgaac accttcttta tggtgacata      750
atccgacaag atttttttaga cacatataat aacctgacct tgaaaaccat      800
tatggcattc aggtgggtaa ctgagttttg cccaatgcc aagtacgtaa      850
tgaagacaga cactgatgtt ttcataata ctggcaattt agtgaagtat      900
cttttaaac taaaccactc agagaagttt ttcacagggt atcctctaatt      950
tgataattat tcctatagag gattttacca aaaaacccat atttcttacc     1000
aggagtatcc tttcaagggt ttccctccat actgcagtgg gttgggttat     1050
ataatgtcca gagatttggg gccaaagatc tatgaaatga tgggtcacgt     1100
aaaaccatc aagtttgaag atgtttatgt cgggatctgt ttgaatttat     1150
taaaagtga cttcatatt ccagaagaca caaatctttt ctttctatat     1200
agaatccatt tggatgtctg tcaactgaga cgtgtgattg cagcccatgg     1250
cttttcttcc aaggagatca tcactttttg gcaggtcatg ctaaggaaca     1300
ccacatgcca ttattaactt cacattctac aaaaagccta gaaggacagg     1350
ataccttgtg gaaagtgtta aataaagtag gtactgtgga aaattcatgg     1400
ggaggtcagt gtgctggcct acactgaact gaaactcatg aaaaaccag      1450
actggagact ggagggttac acttgtgatt tattagtcag gcccttcaaa     1500
gatgatattg ggaggaatta aatataaagg aattggaggt ttttgctaaa     1550
gaaattaata ggaccaaaca atttgacat gtcattctgt agactagaat     1600
ttcttaaaa ggtgttactg agttataagc tcactaggct gtaaaaacaa     1650
aacaatgtag agttttattt attgaacaat gtagtcaatt gaaggttttg     1700
tgtatatctt atgtggatta ccaatttaaa aatatatgta gttctgtgtc     1750
aaaaaacttc ttactgaag ttatactgaa caaaatttta cctgtttttg     1800
gtcatttata aagtacttca agatgttgca gtatttcaca gttattatta     1850
tttaaaatta cttcaacttt gtgtttttta atgttttgac gatttcaata     1900
caagataaaa aggatagtga atcattcttt acatgcaaac attttccagt     1950
tacttaactg atcagtttat tattgataca tcaactccatt aatgtaaagt     2000
cataggtcatt tattgcatat cagtaatctc ttggactttg ttaaatattt     2050
tactgtggta atatagagaa gaattaaagc aagaaaatct gaaaa      2095
```

<210> SEQ ID NO 36
<211> LENGTH: 331
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 36

```
Met Ala Ser Ala Leu Trp Thr Val Leu Pro Ser Arg Met Ser Leu
 1             5             10             15
Arg Ser Leu Lys Trp Ser Leu Leu Leu Ser Leu Leu Ser Phe
                20             25             30
```

-continued

Phe	Val	Met	Trp	Tyr	Leu	Ser	Leu	Pro	His	Tyr	Asn	Val	Ile	Glu
				35					40					45
Arg	Val	Asn	Trp	Met	Tyr	Phe	Tyr	Glu	Tyr	Glu	Pro	Ile	Tyr	Arg
				50					55					60
Gln	Asp	Phe	His	Phe	Thr	Leu	Arg	Glu	His	Ser	Asn	Cys	Ser	His
				65					70					75
Gln	Asn	Pro	Phe	Leu	Val	Ile	Leu	Val	Thr	Ser	His	Pro	Ser	Asp
				80					85					90
Val	Lys	Ala	Arg	Gln	Ala	Ile	Arg	Val	Thr	Trp	Gly	Glu	Lys	Lys
				95					100					105
Ser	Trp	Trp	Gly	Tyr	Glu	Val	Leu	Thr	Phe	Phe	Leu	Leu	Gly	Gln
				110					115					120
Glu	Ala	Glu	Lys	Glu	Asp	Lys	Met	Leu	Ala	Leu	Ser	Leu	Glu	Asp
				125					130					135
Glu	His	Leu	Leu	Tyr	Gly	Asp	Ile	Ile	Arg	Gln	Asp	Phe	Leu	Asp
				140					145					150
Thr	Tyr	Asn	Asn	Leu	Thr	Leu	Lys	Thr	Ile	Met	Ala	Phe	Arg	Trp
				155					160					165
Val	Thr	Glu	Phe	Cys	Pro	Asn	Ala	Lys	Tyr	Val	Met	Lys	Thr	Asp
				170					175					180
Thr	Asp	Val	Phe	Ile	Asn	Thr	Gly	Asn	Leu	Val	Lys	Tyr	Leu	Leu
				185					190					195
Asn	Leu	Asn	His	Ser	Glu	Lys	Phe	Phe	Thr	Gly	Tyr	Pro	Leu	Ile
				200					205					210
Asp	Asn	Tyr	Ser	Tyr	Arg	Gly	Phe	Tyr	Gln	Lys	Thr	His	Ile	Ser
				215					220					225
Tyr	Gln	Glu	Tyr	Pro	Phe	Lys	Val	Phe	Pro	Pro	Tyr	Cys	Ser	Gly
				230					235					240
Leu	Gly	Tyr	Ile	Met	Ser	Arg	Asp	Leu	Val	Pro	Arg	Ile	Tyr	Glu
				245					250					255
Met	Met	Gly	His	Val	Lys	Pro	Ile	Lys	Phe	Glu	Asp	Val	Tyr	Val
				260					265					270
Gly	Ile	Cys	Leu	Asn	Leu	Leu	Lys	Val	Asn	Ile	His	Ile	Pro	Glu
				275					280					285
Asp	Thr	Asn	Leu	Phe	Phe	Leu	Tyr	Arg	Ile	His	Leu	Asp	Val	Cys
				290					295					300
Gln	Leu	Arg	Arg	Val	Ile	Ala	Ala	His	Gly	Phe	Ser	Ser	Lys	Glu
				305					310					315
Ile	Ile	Thr	Phe	Trp	Gln	Val	Met	Leu	Arg	Asn	Thr	Thr	Cys	His
				320					325					330

Tyr

<210> SEQ ID NO 37

<211> LENGTH: 2846

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 37

cgctcgggca ccagccgcgg caaggatgga gctgggttgc tggacgcagt	50
tggggctcac ttttcttcag ctcttctca tctcgtcctt gccaaagagag	100

-continued

tacacagtc	taa	gaagc	ctgccctgga	gcagagtga	atatcatgtg	150
tcgggagtg	c	tgtgaatatg	atcagattga	gtgcgtctgc	cccgaaaga	200
gggaagtcgt	g	gggtataacc	atcccttgct	gcaggaaatga	ggagaatgag	250
tgtgactcct	g	cctgatcca	cccagggttg	accatctttg	aaaactgcaa	300
gagctgccga	a	atggctcat	gggggggtac	cttgatgac	ttctatgtga	350
aggggttcta	ct	gtgcagag	tgccgagcag	gctggtacgg	aggagactgc	400
atgcgatgtg	gcc	aggttct	gcgagcccca	aagggtcaga	ttttgttgga	450
aagctatccc	ctaa	atgctc	actgtgaatg	gaccattcat	gctaaacctg	500
ggtttgtcat	cca	actaaga	tttgtcatgt	tgagtctgga	gtttgactac	550
atgtgccagt	at	gactatgt	tgaggttcgt	gatggagaca	accgcgatgg	600
ccagatcacc	aag	cgtgtct	gtggcaacga	gcggccagct	cctatccaga	650
gcataaggatc	ctc	actccac	gtcctcttcc	actccgatgg	ctccaagaat	700
tttgacgggt	tcc	atgccat	ttatgaggag	atcacagcat	gctcctcatc	750
ccctgttttc	cat	gacggca	cgtgcgtcct	tgacaaggct	ggatcttaca	800
agtggtcctg	ctt	ggcaggc	tatactgggc	agcgtgtga	aaatctcctt	850
gaagaaagaa	act	gctcaga	ccctgggggc	ccagtcaatg	ggtaccagaa	900
aataacaggg	ggc	cctgggc	ttatcaacgg	acgccatgct	aaaattggca	950
ccgtggtgtc	ttt	cttttgt	aacaactcct	atgttcttag	tggaatgag	1000
aaaagaactt	gcc	agcagaa	tgagagtg	tcagggaac	agcccatctg	1050
cataaaagcc	tgcc	gagaac	caaagatttc	agacctggtg	agaaggagag	1100
ttcttccgat	gcag	gttcag	tcaagggaga	caccattaca	ccagctatac	1150
tcagcggcct	tcag	caagca	gaaactgcag	agtgccctca	ccaagaagcc	1200
agcccttccc	tttg	gagatc	tgcccatggg	ataccaacat	ctgcataccc	1250
agctccagta	tgag	tgcatc	tcacccttct	accgccgcct	gggcagcagc	1300
aggaggacat	gtct	gaggac	tggaagtgg	agtgggcggg	caccatcctg	1350
catccctatc	tg	gggaaaa	ttgagaacat	cactgctcca	aagacccaag	1400
ggttgcgctg	gcc	gtggcag	gcagccatct	acaggaggac	cagcggggtg	1450
catgacggca	gcct	acacaa	gggagcgtgg	ttcctagtct	gcagcgggtg	1500
cctggtgaat	gag	cgcactg	tggtggtggc	tgcccactgt	gttactgacc	1550
tggggaaggt	cacc	atgac	gcagcagc	acctgaaagt	tgttttgggg	1600
aaattctacc	ggg	atgatga	ccgggatgag	aagaccatcc	agagcctaca	1650
gatttctgct	at	cattctgc	atcccaacta	tgaccccatc	ctgcttgatg	1700
ctgacatcgc	cat	cctgaag	ctcctagaca	agggccgtat	cagcaccgga	1750
gtccagccca	tct	gcctcgc	tgccagtcgg	gatctcagca	cttccctcca	1800
ggagtccac	at	cactgtgg	ctggctggaa	tgtcctggca	gacgtgagga	1850
gccctggctt	ca	agaacgac	acactgcgct	ctgggggtgt	cagtgtgtgtg	1900
gactcgctgc	tgt	gtgagga	gcagcatgag	gaccatggca	tcccagtga	1950
tgctcactgat	aac	atgttct	gtgccagctg	ggaaccact	gccccttctg	2000

-continued

atatctgcac tgcagagaca ggaggcatcg cggtctgtgc cttcccggga	2050
cgagcatctc ctgagccacg ctggcatctg atgggactgg tcagctggag	2100
ctatgataaa acatgcagcc acaggctctc cactgccttc accaagggtg	2150
tgctttttta agactggatt gaaagaaata tgaaatgaac catgctcatg	2200
cactccttga gaagtgtttc tgtatatccg tctgtacgtg tgtcattgcg	2250
tgaagcagtg tgggcctgaa gtgtgatttg gcctgtgaac ttggctgtgc	2300
cagggtcttc gacttcaggg acaaaactca gtgaagggtg agtagacctc	2350
cattgctggt aggctgatgc cgcgtccact actaggacag ccaattggaa	2400
gatgccaggg cttgcaagaa gtaagtttct tcaaagaaga ccatatacaa	2450
aacctctcca ctccactgac ctggtggtct tccccaactt tcagttatac	2500
gaatgccatc agcttgacca gggaagatct gggcttcatg aggccccttt	2550
tgaggctctc aagttctaga gagctgcctg tgggacagcc cagggcagca	2600
gagctgggat gtggtgcatg cttttgtgta catggccaca gtacagtctg	2650
gtccttttcc ttccccatct cttgtacaca ttttaataaa ataagggttg	2700
gcttctgaac tacaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	2750
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	2800
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa	2846

<210> SEQ ID NO 38

<211> LENGTH: 720

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 38

Met	Glu	Leu	Gly	Cys	Trp	Thr	Gln	Leu	Gly	Leu	Thr	Phe	Leu	Gln
1				5					10					15
Leu	Leu	Leu	Ile	Ser	Ser	Leu	Pro	Arg	Glu	Tyr	Thr	Val	Ile	Asn
				20					25					30
Glu	Ala	Cys	Pro	Gly	Ala	Glu	Trp	Asn	Ile	Met	Cys	Arg	Glu	Cys
				35					40					45
Cys	Glu	Tyr	Asp	Gln	Ile	Glu	Cys	Val	Cys	Pro	Gly	Lys	Arg	Glu
				50					55					60
Val	Val	Gly	Tyr	Thr	Ile	Pro	Cys	Cys	Arg	Asn	Glu	Glu	Asn	Glu
				65					70					75
Cys	Asp	Ser	Cys	Leu	Ile	His	Pro	Gly	Cys	Thr	Ile	Phe	Glu	Asn
				80					85					90
Cys	Lys	Ser	Cys	Arg	Asn	Gly	Ser	Trp	Gly	Gly	Thr	Leu	Asp	Asp
				95					100					105
Phe	Tyr	Val	Lys	Gly	Phe	Tyr	Cys	Ala	Glu	Cys	Arg	Ala	Gly	Trp
				110					115					120
Tyr	Gly	Gly	Asp	Cys	Met	Arg	Cys	Gly	Gln	Val	Leu	Arg	Ala	Pro
				125					130					135
Lys	Gly	Gln	Ile	Leu	Leu	Glu	Ser	Tyr	Pro	Leu	Asn	Ala	His	Cys
				140					145					150
Glu	Trp	Thr	Ile	His	Ala	Lys	Pro	Gly	Phe	Val	Ile	Gln	Leu	Arg
				155					160					165
Phe	Val	Met	Leu	Ser	Leu	Glu	Phe	Asp	Tyr	Met	Cys	Gln	Tyr	Asp

-continued

170					175					180				
Tyr	Val	Glu	Val	Arg	Asp	Gly	Asp	Asn	Arg	Asp	Gly	Gln	Ile	Ile
				185					190					195
Lys	Arg	Val	Cys	Gly	Asn	Glu	Arg	Pro	Ala	Pro	Ile	Gln	Ser	Ile
				200					205					210
Gly	Ser	Ser	Leu	His	Val	Leu	Phe	His	Ser	Asp	Gly	Ser	Lys	Asn
				215					220					225
Phe	Asp	Gly	Phe	His	Ala	Ile	Tyr	Glu	Glu	Ile	Thr	Ala	Cys	Ser
				230					235					240
Ser	Ser	Pro	Cys	Phe	His	Asp	Gly	Thr	Cys	Val	Leu	Asp	Lys	Ala
				245					250					255
Gly	Ser	Tyr	Lys	Cys	Ala	Cys	Leu	Ala	Gly	Tyr	Thr	Gly	Gln	Arg
				260					265					270
Cys	Glu	Asn	Leu	Leu	Glu	Glu	Arg	Asn	Cys	Ser	Asp	Pro	Gly	Gly
				275					280					285
Pro	Val	Asn	Gly	Tyr	Gln	Lys	Ile	Thr	Gly	Gly	Pro	Gly	Leu	Ile
				290					295					300
Asn	Gly	Arg	His	Ala	Lys	Ile	Gly	Thr	Val	Val	Ser	Phe	Phe	Cys
				305					310					315
Asn	Asn	Ser	Tyr	Val	Leu	Ser	Gly	Asn	Glu	Lys	Arg	Thr	Cys	Gln
				320					325					330
Gln	Asn	Gly	Glu	Trp	Ser	Gly	Lys	Gln	Pro	Ile	Cys	Ile	Lys	Ala
				335					340					345
Cys	Arg	Glu	Pro	Lys	Ile	Ser	Asp	Leu	Val	Arg	Arg	Arg	Val	Leu
				350					355					360
Pro	Met	Gln	Val	Gln	Ser	Arg	Glu	Thr	Pro	Leu	His	Gln	Leu	Tyr
				365					370					375
Ser	Ala	Ala	Phe	Ser	Lys	Gln	Lys	Leu	Gln	Ser	Ala	Pro	Thr	Lys
				380					385					390
Lys	Pro	Ala	Leu	Pro	Phe	Gly	Asp	Leu	Pro	Met	Gly	Tyr	Gln	His
				395					400					405
Leu	His	Thr	Gln	Leu	Gln	Tyr	Glu	Cys	Ile	Ser	Pro	Phe	Tyr	Arg
				410					415					420
Arg	Leu	Gly	Ser	Ser	Arg	Arg	Thr	Cys	Leu	Arg	Thr	Gly	Lys	Trp
				425					430					435
Ser	Gly	Arg	Ala	Pro	Ser	Cys	Ile	Pro	Ile	Cys	Gly	Lys	Ile	Glu
				440					445					450
Asn	Ile	Thr	Ala	Pro	Lys	Thr	Gln	Gly	Leu	Arg	Trp	Pro	Trp	Gln
				455					460					465
Ala	Ala	Ile	Tyr	Arg	Arg	Thr	Ser	Gly	Val	His	Asp	Gly	Ser	Leu
				470					475					480
His	Lys	Gly	Ala	Trp	Phe	Leu	Val	Cys	Ser	Gly	Ala	Leu	Val	Asn
				485					490					495
Glu	Arg	Thr	Val	Val	Val	Ala	Ala	His	Cys	Val	Thr	Asp	Leu	Gly
				500					505					510
Lys	Val	Thr	Met	Ile	Lys	Thr	Ala	Asp	Leu	Lys	Val	Val	Leu	Gly
				515					520					525
Lys	Phe	Tyr	Arg	Asp	Asp	Asp	Arg	Asp	Glu	Lys	Thr	Ile	Gln	Ser
				530					535					540
Leu	Gln	Ile	Ser	Ala	Ile	Ile	Leu	His	Pro	Asn	Tyr	Asp	Pro	Ile
				545					550					555

-continued

Leu	Leu	Asp	Ala	Asp	Ile	Ala	Ile	Leu	Lys	Leu	Leu	Asp	Lys	Ala
			560						565					570
Arg	Ile	Ser	Thr	Arg	Val	Gln	Pro	Ile	Cys	Leu	Ala	Ala	Ser	Arg
			575						580					585
Asp	Leu	Ser	Thr	Ser	Phe	Gln	Glu	Ser	His	Ile	Thr	Val	Ala	Gly
			590						595					600
Trp	Asn	Val	Leu	Ala	Asp	Val	Arg	Ser	Pro	Gly	Phe	Lys	Asn	Asp
			605						610					615
Thr	Leu	Arg	Ser	Gly	Val	Val	Ser	Val	Val	Asp	Ser	Leu	Leu	Cys
			620						625					630
Glu	Glu	Gln	His	Glu	Asp	His	Gly	Ile	Pro	Val	Ser	Val	Thr	Asp
			635						640					645
Asn	Met	Phe	Cys	Ala	Ser	Trp	Glu	Pro	Thr	Ala	Pro	Ser	Asp	Ile
			650						655					660
Cys	Thr	Ala	Glu	Thr	Gly	Gly	Ile	Ala	Ala	Val	Ser	Phe	Pro	Gly
			665						670					675
Arg	Ala	Ser	Pro	Glu	Pro	Arg	Trp	His	Leu	Met	Gly	Leu	Val	Ser
			680						685					690
Trp	Ser	Tyr	Asp	Lys	Thr	Cys	Ser	His	Arg	Leu	Ser	Thr	Ala	Phe
			695						700					705
Thr	Lys	Val	Leu	Pro	Phe	Lys	Asp	Trp	Ile	Glu	Arg	Asn	Met	Lys
			710						715					720

<210> SEQ ID NO 39
<211> LENGTH: 2571
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 39

ggttcctaca tcctctcatc tgagaatcag agagcataat cttottacgg	50
gcccgtgatt tattaacgtg gcttaatctg aaggttctca gtcaaattct	100
ttgtgatcta ctgattgtgg gggcatggca aggtttgctt aaaggagctt	150
ggctggtttg ggcccttgta gctgacagaa ggtggccagg gagaatgcag	200
cacactgctc ggagaatgaa ggcgcttctg ttgctgtctc tgccttggtc	250
cagtcctgct aactacattg acaatgtggg caacctgcac ttcctgtatt	300
cagaactctg taaaggtgcc tcccactacg gcctgaccaa agataggaag	350
agggcctcac aagatggctg tccagacggc tgtgcgagcc tcacagccac	400
ggctccctcc ccagaggttt ctgcagctgc caccatctcc ttaatgacag	450
acgagcctgg cctagacaac cctgcctacg tgtcctcggc agaggacggg	500
cagccagcaa tcagcccagt ggactctggc cggagcaacc gaactagggc	550
acggcccttt gagagatcca ctattagaag cagatcattt aaaaaataa	600
atcgagcttt gagtgttctt cgaaggacaa agagcgggag tgcagttgcc	650
aaccatgccg accagggcag ggaaaattct gaaaacacca ctgccctga	700
agtctttcca aggttgtacc acctgattcc agatggtgaa attaccagca	750
tcaagatcaa tcgagtagat ccagtgaaa gcctctctat taggctggtg	800
ggaggtagcg aaacccact ggtccatcctc attatccaac acatttatcg	850

-continued

tgatggggtg atcgccagag acggccggct actgccagga gacatcattc	900
taaaggtcaa cgggatggac atcagcaatg tccctcacia ctacgtgtg	950
cgtctcctcg ggcagccctg ccaggtgctg tggctgactg tgatgcgtga	1000
acagaagtgc cgcagcagga acaatggaca ggccccggat gcctacagac	1050
cccagatga cagctttcat gtgattctca acaaaagtag ccccgaggag	1100
cagcttgaa taaaactggt gcgcaagggt gatgagcctg gggttttcat	1150
cttcaatgtg ctggatggcg gtgtggcata tcgacatggt cagcttgagg	1200
agaatgaccg tgtgttagcc atcaatggac atgatcttcg atatggcagc	1250
ccagaaagtg cggtcatctt gattcaggcc agtgaaagac gtgttcacct	1300
cgtcgtgtcc cgccagggtc ggcagcggag cctgacatc ttccaggaag	1350
ccggctggaa cagcaatggc agctggtccc cagggccagg ggagaggagc	1400
aacactccca agcccccca tcctacaatt acttgtcatg agaaggtggt	1450
aaatatccaa aaagaccccc gtgaatctct cggcattgacc gtcgcagggg	1500
gagcatcaca tagagaatgg gatttgccta tctatgtcat cagtgttgag	1550
cccggaggag tcataagcag agatggaaga ataaaaacag gtgacatttt	1600
gttgaatgtg gatggggctg aactgacaga ggtcagccgg agtgaggcag	1650
tggcattatt gaaaagaaca tcatcctcga tagtactcaa agctttggaa	1700
gtcaaaagt atgagcccca ggaagactgc agcagcccag cagccctgga	1750
ctccaaccac aacatggccc caccagtgat ctggtcccca tcctgggtca	1800
tgtgctgga attaccacgg tgcttgata actgtaaaga tattgtatta	1850
cgaagaaaca cagctggaag tctgggcttc tgcattgtag gaggttatga	1900
agaatacaat ggaacaaac cttttttcat caaatccatt gttgaaggaa	1950
caccagcata caatgatgga agaattagat gtggtgatat tcttcttgct	2000
gtcaatggta gaagtacatc aggaatgata catgcttgct tggcaagact	2050
gctgaaagaa cttaaaggaa gaattactct aactattgtt tcttgccctg	2100
gcactttttt atagaatcaa tgatgggtca gagaaaaca gaaaaatcac	2150
aaataggcta agaagttgaa acactatatt tatcttgtca gttttatat	2200
ttaaagaag aatacattgt aaaaatgtca ggaaaagtat gatcatctaa	2250
tgaaagccag ttacacctca gaaaatatga ttccaaaaaa attaaaacta	2300
ctagtttttt ttcagtgtgg aggatttctc attactctac aacattgttt	2350
atattttttt tattcaataa aaagccctaa aacaactaaa atgattgatt	2400
tgtatacccc actgaattca agctgattta aatttaaaat ttggtatatg	2450
ctgaagtctg ccaagggtac attatggcca tttttaattt acagctaaaa	2500
tatttttttaaatgcatgtc tgagaaacgt tgctttcatc aaacaagaat	2550
aaatattttt cagaagttaa a	2571

<210> SEQ ID NO 40

<211> LENGTH: 632

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 40

Met	Lys	Ala	Leu	Leu	Leu	Val	Leu	Pro	Trp	Leu	Ser	Pro	Ala	
1			5					10					15	
Asn	Tyr	Ile	Asp	Asn	Val	Gly	Asn	Leu	His	Phe	Leu	Tyr	Ser	Glu
			20					25						30
Leu	Cys	Lys	Gly	Ala	Ser	His	Tyr	Gly	Leu	Thr	Lys	Asp	Arg	Lys
			35					40						45
Arg	Arg	Ser	Gln	Asp	Gly	Cys	Pro	Asp	Gly	Cys	Ala	Ser	Leu	Thr
			50					55						60
Ala	Thr	Ala	Pro	Ser	Pro	Glu	Val	Ser	Ala	Ala	Ala	Thr	Ile	Ser
			65					70						75
Leu	Met	Thr	Asp	Glu	Pro	Gly	Leu	Asp	Asn	Pro	Ala	Tyr	Val	Ser
			80					85						90
Ser	Ala	Glu	Asp	Gly	Gln	Pro	Ala	Ile	Ser	Pro	Val	Asp	Ser	Gly
			95					100						105
Arg	Ser	Asn	Arg	Thr	Arg	Ala	Arg	Pro	Phe	Glu	Arg	Ser	Thr	Ile
			110					115						120
Arg	Ser	Arg	Ser	Phe	Lys	Lys	Ile	Asn	Arg	Ala	Leu	Ser	Val	Leu
			125					130						135
Arg	Arg	Thr	Lys	Ser	Gly	Ser	Ala	Val	Ala	Asn	His	Ala	Asp	Gln
			140					145						150
Gly	Arg	Glu	Asn	Ser	Glu	Asn	Thr	Thr	Ala	Pro	Glu	Val	Phe	Pro
			155					160						165
Arg	Leu	Tyr	His	Leu	Ile	Pro	Asp	Gly	Glu	Ile	Thr	Ser	Ile	Lys
			170					175						180
Ile	Asn	Arg	Val	Asp	Pro	Ser	Glu	Ser	Leu	Ser	Ile	Arg	Leu	Val
			185					190						195
Gly	Gly	Ser	Glu	Thr	Pro	Leu	Val	His	Ile	Ile	Ile	Gln	His	Ile
			200					205						210
Tyr	Arg	Asp	Gly	Val	Ile	Ala	Arg	Asp	Gly	Arg	Leu	Leu	Pro	Gly
			215					220						225
Asp	Ile	Ile	Leu	Lys	Val	Asn	Gly	Met	Asp	Ile	Ser	Asn	Val	Pro
			230					235						240
His	Asn	Tyr	Ala	Val	Arg	Leu	Leu	Arg	Gln	Pro	Cys	Gln	Val	Leu
			245					250						255
Trp	Leu	Thr	Val	Met	Arg	Glu	Gln	Lys	Phe	Arg	Ser	Arg	Asn	Asn
			260					265						270
Gly	Gln	Ala	Pro	Asp	Ala	Tyr	Arg	Pro	Arg	Asp	Asp	Ser	Phe	His
			275					280						285
Val	Ile	Leu	Asn	Lys	Ser	Ser	Pro	Glu	Glu	Gln	Leu	Gly	Ile	Lys
			290					295						300
Leu	Val	Arg	Lys	Val	Asp	Glu	Pro	Gly	Val	Phe	Ile	Phe	Asn	Val
			305					310						315
Leu	Asp	Gly	Gly	Val	Ala	Tyr	Arg	His	Gly	Gln	Leu	Glu	Glu	Asn
			320					325						330
Asp	Arg	Val	Leu	Ala	Ile	Asn	Gly	His	Asp	Leu	Arg	Tyr	Gly	Ser
			335					340						345
Pro	Glu	Ser	Ala	Ala	His	Leu	Ile	Gln	Ala	Ser	Glu	Arg	Arg	Val
			350					355						360
His	Leu	Val	Val	Ser	Arg	Gln	Val	Arg	Gln	Arg	Ser	Pro	Asp	Ile
			365					370						375

-continued

Phe Gln Glu Ala Gly Trp Asn Ser Asn Gly Ser Trp Ser Pro Gly
380 385 390

Pro Gly Glu Arg Ser Asn Thr Pro Lys Pro Leu His Pro Thr Ile
395 400 405

Thr Cys His Glu Lys Val Val Asn Ile Gln Lys Asp Pro Gly Glu
410 415 420

Ser Leu Gly Met Thr Val Ala Gly Gly Ala Ser His Arg Glu Trp
425 430 435

Asp Leu Pro Ile Tyr Val Ile Ser Val Glu Pro Gly Gly Val Ile
440 445 450

Ser Arg Asp Gly Arg Ile Lys Thr Gly Asp Ile Leu Leu Asn Val
455 460 465

Asp Gly Val Glu Leu Thr Glu Val Ser Arg Ser Glu Ala Val Ala
470 475 480

Leu Leu Lys Arg Thr Ser Ser Ser Ile Val Leu Lys Ala Leu Glu
485 490 495

Val Lys Glu Tyr Glu Pro Gln Glu Asp Cys Ser Ser Pro Ala Ala
500 505 510

Leu Asp Ser Asn His Asn Met Ala Pro Pro Ser Asp Trp Ser Pro
515 520 525

Ser Trp Val Met Trp Leu Glu Leu Pro Arg Cys Leu Tyr Asn Cys
530 535 540

Lys Asp Ile Val Leu Arg Arg Asn Thr Ala Gly Ser Leu Gly Phe
545 550 555

Cys Ile Val Gly Gly Tyr Glu Glu Tyr Asn Gly Asn Lys Pro Phe
560 565 570

Phe Ile Lys Ser Ile Val Glu Gly Thr Pro Ala Tyr Asn Asp Gly
575 580 585

Arg Ile Arg Cys Gly Asp Ile Leu Leu Ala Val Asn Gly Arg Ser
590 595 600

Thr Ser Gly Met Ile His Ala Cys Leu Ala Arg Leu Leu Lys Glu
605 610 615

Leu Lys Gly Arg Ile Thr Leu Thr Ile Val Ser Trp Pro Gly Thr
620 625 630

Phe Leu

<210> SEQ ID NO 41
<211> LENGTH: 1964
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 41

accaggcatt gtatcttcag ttgtcatcaa gttcgcaatc agattggaaa	50
agctcaactt gaagctttct tgctgcagt gaagcagaga gatagatatt	100
attcacgtaa taaaaaacat gggcttcaac ctgactttcc acctttccta	150
caaattccga ttactgttgc tgttgacttt gtgcctgaca gtggttgggt	200
gggccaccag taactacttc gtgggtgcc ttcaagagat tcctaaagca	250
aaggagtcca tggctaattt ccataagacc ctcattttgg ggaagggaaa	300
aactctgact aatgaagcat ccacgaagaa ggtagaactt gacaactgtc	350

-continued

cttctgtgtc tccttacctc agaggccaga gcaagctcat tttcaaacca	400
gatctcactt tggaagaggt acaggcagaa aatcccaaag tgtccagagg	450
ccggtatcgc cctcaggaat gtaaagcttt acagagggtc gccatcctcg	500
ttccccaccg gaacagagag aaacacctga tgtacctgct ggaacatctg	550
catcccttcc tgcagaggca gcagctggat tatggcatct acgtcatcca	600
ccaggctgaa ggtaaaaagt ttaatcgagc caaactcttg aatgtgggct	650
atctagaagc cctcaaggaa gaaaattggg actgctttat attccacgat	700
gtggacctgg tacccgagaa tgactttaac ctttacaagt gtgaggagca	750
tcccagcat ctggtggttg gcaggaaacag cactgggtac aggttacgtt	800
acagtggata ttttgggggt gttactgccc taagcagaga gcagtttttc	850
aagggtgaatg gattctctaa caactactgg ggatggggag gcgaagacga	900
tgacctcaga ctcagggttg agctccaaag aatgaaaatt tcccggtccc	950
tgctgaagt gggtaaatat acaatgggtc tccacactag agacaaaggc	1000
aatgaggtda acgcagaacg gatgaagctc ttacaccaag tgtcacgagt	1050
ctggagaaca gatgggttga gtagttgttc ttataaatta gtatctgttg	1100
aacacaatcc tttatatatc aacatcacag tggatttctg gtttggtgca	1150
tgaccttgga tcttttggtg atgtttggaa gaactgattc tttgtttgca	1200
ataattttgg cctagagact tcaaatagta gcacacatta agaacctgtt	1250
acagctcatt gttgagctga atttttcctt tttgtatttt cttagcagag	1300
ctcctggtda tgtagagtat aaaacagttg taacaagaca gctttcttag	1350
tcattttgat catgaggggt aaatatgtga atatggatac ttgaaggact	1400
ttatataaaa ggatgactca aaggataaaa tgaacgctat ttgaggactc	1450
tggttgaagg agattttatt aaatttgaag taatatatta tgggataaaa	1500
ggccacagga aataagactg ctgaatgtct gagagaacca gagttgttct	1550
cgtccaaggt agaaaggtag gaagatacaa tactgttatt cattttacct	1600
gtacaatcat ctgtgaagtg gtggtgtcag gtgagaaggc gtccacaaaa	1650
gaggggagaa aaggcgacga atcaggacac agtgaacttg ggaatgaaga	1700
ggtagcagga ggggtgagtg tcggctgcaa aggcagcagt agctgagctg	1750
gttgacagtg ctgatagcct tcaggggagg acctgcccag gtatgccttc	1800
cagtgatgcc caccagagaa tacattctct attagttttt aaagagtttt	1850
tgtaaaatga tttgtacaa gtaggatatg aattagcagt ttacaagttt	1900
acatatatac taataataaa tatgtctatc aaatacctct gtagtaaaat	1950
gtgaaaaagc aaaa	1964

<210> SEQ ID NO 42

<211> LENGTH: 344

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 42

Met	Gly	Phe	Asn	Leu	Thr	Phe	His	Leu	Ser	Tyr	Lys	Phe	Arg	Leu
1					5					10				15

-continued

Leu	Leu	Leu	Leu	Thr	Leu	Cys	Leu	Thr	Val	Val	Gly	Trp	Ala	Thr	
				20					25					30	
Ser	Asn	Tyr	Phe	Val	Gly	Ala	Ile	Gln	Glu	Ile	Pro	Lys	Ala	Lys	
				35					40					45	
Glu	Phe	Met	Ala	Asn	Phe	His	Lys	Thr	Leu	Ile	Leu	Gly	Lys	Gly	
				50					55					60	
Lys	Thr	Leu	Thr	Asn	Glu	Ala	Ser	Thr	Lys	Lys	Val	Glu	Leu	Asp	
				65					70					75	
Asn	Cys	Pro	Ser	Val	Ser	Pro	Tyr	Leu	Arg	Gly	Gln	Ser	Lys	Leu	
				80					85					90	
Ile	Phe	Lys	Pro	Asp	Leu	Thr	Leu	Glu	Glu	Val	Gln	Ala	Glu	Asn	
				95					100					105	
Pro	Lys	Val	Ser	Arg	Gly	Arg	Tyr	Arg	Pro	Gln	Glu	Cys	Lys	Ala	
				110					115					120	
Leu	Gln	Arg	Val	Ala	Ile	Leu	Val	Pro	His	Arg	Asn	Arg	Glu	Lys	
				125					130					135	
His	Leu	Met	Tyr	Leu	Leu	Glu	His	Leu	His	Pro	Phe	Leu	Gln	Arg	
				140					145					150	
Gln	Gln	Leu	Asp	Tyr	Gly	Ile	Tyr	Val	Ile	His	Gln	Ala	Glu	Gly	
				155					160					165	
Lys	Lys	Phe	Asn	Arg	Ala	Lys	Leu	Leu	Asn	Val	Gly	Tyr	Leu	Glu	
				170					175					180	
Ala	Leu	Lys	Glu	Glu	Asn	Trp	Asp	Cys	Phe	Ile	Phe	His	Asp	Val	
				185					190					195	
Asp	Leu	Val	Pro	Glu	Asn	Asp	Phe	Asn	Leu	Tyr	Lys	Cys	Glu	Glu	
				200					205					210	
His	Pro	Lys	His	Leu	Val	Val	Gly	Arg	Asn	Ser	Thr	Gly	Tyr	Arg	
				215					220					225	
Leu	Arg	Tyr	Ser	Gly	Tyr	Phe	Gly	Gly	Val	Thr	Ala	Leu	Ser	Arg	
				230					235					240	
Glu	Gln	Phe	Phe	Lys	Val	Asn	Gly	Phe	Ser	Asn	Asn	Tyr	Trp	Gly	
				245					250					255	
Trp	Gly	Gly	Glu	Asp	Asp	Asp	Leu	Arg	Leu	Arg	Val	Glu	Leu	Gln	
				260					265					270	
Arg	Met	Lys	Ile	Ser	Arg	Pro	Leu	Pro	Glu	Val	Gly	Lys	Tyr	Thr	
				275					280					285	
Met	Val	Phe	His	Thr	Arg	Asp	Lys	Gly	Asn	Glu	Val	Asn	Ala	Glu	
				290					295					300	
Arg	Met	Lys	Leu	Leu	His	Gln	Val	Ser	Arg	Val	Trp	Arg	Thr	Asp	
				305					310					315	
Gly	Leu	Ser	Ser	Cys	Ser	Tyr	Lys	Leu	Val	Ser	Val	Glu	His	Asn	
				320					325					330	
Pro	Leu	Tyr	Ile	Asn	Ile	Thr	Val	Asp	Phe	Trp	Phe	Gly	Ala		
				335					340						

<210> SEQ ID NO 43
<211> LENGTH: 485
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 43

gctcaagacc cagcagtggg acagccagac agacggcacg atggcactga

-continued

gctcccagat ctgggccgct tgccctcctgc tccctcctcct cctcgccagc	100
ctgaccagtg gctctgtttt cccacaacag acgggacaac ttgcagagct	150
gcaaccccag gacagagctg gagccagggc cagctggatg cccatgttcc	200
agaggcgaag gaggcgagac acccacttcc ccatctgcat tttctgctgc	250
ggctgctgtc atcgatcaaa gtgtgggatg tgctgcaaga cgtagaacct	300
acctgccctg cccccgtccc ctcccttcct tatttattcc tgctgcccc	350
gaacataggt cttggaataa aatggctggt tcttttgttt tccaaaaaaa	400
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	450
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa	485

<210> SEQ ID NO 44
<211> LENGTH: 84
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 44

Met	Ala	Leu	Ser	Ser	Gln	Ile	Trp	Ala	Ala	Cys	Leu	Leu	Leu	Leu
1				5					10					15
Leu	Leu	Leu	Ala	Ser	Leu	Thr	Ser	Gly	Ser	Val	Phe	Pro	Gln	Gln
			20						25					30
Thr	Gly	Gln	Leu	Ala	Glu	Leu	Gln	Pro	Gln	Asp	Arg	Ala	Gly	Ala
			35						40					45
Arg	Ala	Ser	Trp	Met	Pro	Met	Phe	Gln	Arg	Arg	Arg	Arg	Arg	Asp
			50						55					60
Thr	His	Phe	Pro	Ile	Cys	Ile	Phe	Cys	Cys	Gly	Cys	Cys	His	Arg
			65						70					75
Ser	Lys	Cys	Gly	Met	Cys	Cys	Lys	Thr						
			80											

<210> SEQ ID NO 45
<211> LENGTH: 1076
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 45

gtggcttcat ttcagtggct gacttccaga gagcaatatg gctggttccc	50
caacatgcct caccctcatc tatatccttt ggcagctcac agggtcagca	100
gcctctggag ccgtgaaaga gctggtcggt tccgttggtg gggccgtgac	150
tttccccctg aagtccaaag taaagcaagt tgactctatt gtctggacct	200
tcaacacaac ccctcttgtc accatacagc cagaaggggg cactatcata	250
gtgacccaaa atcgtaatat ggagagagta gacttcccag atggaggcta	300
ctccctgaag ctacgaaac tgaagaagaa tgactcaggg atctactatg	350
tggggatata cagctcatca ctccagcagc cctccacca ggagtacgtg	400
ctgcatgtct acgagcacct gtcaaagcct aaagtcacca tgggtctgca	450
gagcaataag aatggcacct gtgtgaccaa tctgacatgc tgcattgaac	500
atggggaaga ggatgtgatt tatacctgga aggcctggg gcaagcagcc	550
aatgagtccc ataatgggtc catcctcccc atctcctgga gatggggaga	600

-continued

aagtgatatg accttcatct gcgttgccag gaaccctgtc agcagaaact 650
tctcaagccc catccttgcc aggaagctct gtgaaggtgc tgctgatgac 700
ccagattcct ccatggctct cctgtgtctc ctgttggtgc ccctcctgct 750
cagtctcttt gtactggggc tatttctttg gtttctgaag agagagagac 800
aagaagagta cattgaagag aagaagagag tggacatttg tcgggaaact 850
cctaacatat gccccattc tggagagaac acagagtacg acacaatccc 900
tcacactaat agaacaatcc taaaggaaga tccagcaaat acggtttact 950
ccactgtgga aataccgaaa aagatggaaa atccccactc actgctcacg 1000
atgccagaca caccaaggct atttgctat gagaatgtta tctagacagc 1050
agtgcactcc cctaagtctc tgctca 1076

<210> SEQ ID NO 46
<211> LENGTH: 335
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 46

Met Ala Gly Ser Pro Thr Cys Leu Thr Leu Ile Tyr Ile Leu Trp
1 5 10 15
Gln Leu Thr Gly Ser Ala Ala Ser Gly Pro Val Lys Glu Leu Val
20 25 30
Gly Ser Val Gly Gly Ala Val Thr Phe Pro Leu Lys Ser Lys Val
35 40 45
Lys Gln Val Asp Ser Ile Val Trp Thr Phe Asn Thr Thr Pro Leu
50 55 60
Val Thr Ile Gln Pro Glu Gly Gly Thr Ile Ile Val Thr Gln Asn
65 70 75
Arg Asn Arg Glu Arg Val Asp Phe Pro Asp Gly Gly Tyr Ser Leu
80 85 90
Lys Leu Ser Lys Leu Lys Lys Asn Asp Ser Gly Ile Tyr Tyr Val
95 100 105
Gly Ile Tyr Ser Ser Ser Leu Gln Gln Pro Ser Thr Gln Glu Tyr
110 115 120
Val Leu His Val Tyr Glu His Leu Ser Lys Pro Lys Val Thr Met
125 130 135
Gly Leu Gln Ser Asn Lys Asn Gly Thr Cys Val Thr Asn Leu Thr
140 145 150
Cys Cys Met Glu His Gly Glu Glu Asp Val Ile Tyr Thr Trp Lys
155 160 165
Ala Leu Gly Gln Ala Ala Asn Glu Ser His Asn Gly Ser Ile Leu
170 175 180
Pro Ile Ser Trp Arg Trp Gly Glu Ser Asp Met Thr Phe Ile Cys
185 190 195
Val Ala Arg Asn Pro Val Ser Arg Asn Phe Ser Ser Pro Ile Leu
200 205 210
Ala Arg Lys Leu Cys Glu Gly Ala Ala Asp Asp Pro Asp Ser Ser
215 220 225
Met Val Leu Leu Cys Leu Leu Leu Val Pro Leu Leu Leu Ser Leu
230 235 240

-continued

Phe Val Leu Gly Leu Phe Leu Trp Phe Leu Lys Arg Glu Arg Gln
245 250 255
Glu Glu Tyr Ile Glu Glu Lys Lys Arg Val Asp Ile Cys Arg Glu
260 265 270
Thr Pro Asn Ile Cys Pro His Ser Gly Glu Asn Thr Glu Tyr Asp
275 280 285
Thr Ile Pro His Thr Asn Arg Thr Ile Leu Lys Glu Asp Pro Ala
290 295 300
Asn Thr Val Tyr Ser Thr Val Glu Ile Pro Lys Lys Met Glu Asn
305 310 315
Pro His Ser Leu Leu Thr Met Pro Asp Thr Pro Arg Leu Phe Ala
320 325 330
Tyr Glu Asn Val Ile
335

<210> SEQ ID NO 47
<211> LENGTH: 766
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 47

ggctcgagcg tttctgagcc aggggtgacc atgacctgct gcgaaggatg 50
gacatcctgc aatggattca gcctgctggt tctactgctg ttaggagtag 100
ttctcaatgc gatacctcta attgtcagct tagttgagga agaccaattt 150
tctcaaaacc ccatctcttg ctttgagtgg tggttcccag gaattatagg 200
agcagggtctg atggccattc cagcaacaac aatgtccttg acagcaagaa 250
aaagagcgtg ctgcaacaac agaactggaa tgtttctttc atcatttttc 300
agtgtgatca cagtcattgg tgctctgtat tgcattgctga tatccatcca 350
ggctctctta aaaggtcctc tcatgtgtaa ttctccaagc aacagtaatg 400
ccaattgtga attttcattg aaaaacatca gtgacattca tccagaatcc 450
ttcaacttgc agtgggtttt caatgactct tgtgcacctc ctactggttt 500
caataaaacc accagtaacg acaccatggc gagtggctgg agagcatcta 550
gtttccactt cgattctgaa gaaaacaaac ataggcttat ccacttctca 600
gtatttttag gtctattgct tgttggaatt ctggaggtcc tgtttgggct 650
cagtcagata gtcacgggtt tccttggtg tctgtgtgga gtctctaagc 700
gaagaagtca aattgtgtag tttaatggga ataaaatgta agtatcagta 750
gtttgaaaaa aaaaaa 766

<210> SEQ ID NO 48
<211> LENGTH: 229
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 48

Met Thr Cys Cys Glu Gly Trp Thr Ser Cys Asn Gly Phe Ser Leu
1 5 10 15
Leu Val Leu Leu Leu Gly Val Val Leu Asn Ala Ile Pro Leu
20 25 30

-continued

Ile	Val	Ser	Leu	Val	Glu	Glu	Asp	Gln	Phe	Ser	Gln	Asn	Pro	Ile	
	35								40					45	
Ser	Cys	Phe	Glu	Trp	Trp	Phe	Pro	Gly	Ile	Ile	Gly	Ala	Gly	Leu	
	50								55					60	
Met	Ala	Ile	Pro	Ala	Thr	Thr	Met	Ser	Leu	Thr	Ala	Arg	Lys	Arg	
	65								70					75	
Ala	Cys	Cys	Asn	Asn	Arg	Thr	Gly	Met	Phe	Leu	Ser	Ser	Phe	Phe	
	80								85					90	
Ser	Val	Ile	Thr	Val	Ile	Gly	Ala	Leu	Tyr	Cys	Met	Leu	Ile	Ser	
	95								100					105	
Ile	Gln	Ala	Leu	Leu	Lys	Gly	Pro	Leu	Met	Cys	Asn	Ser	Pro	Ser	
	110								115					120	
Asn	Ser	Asn	Ala	Asn	Cys	Glu	Phe	Ser	Leu	Lys	Asn	Ile	Ser	Asp	
	125								130					135	
Ile	His	Pro	Glu	Ser	Phe	Asn	Leu	Gln	Trp	Phe	Phe	Asn	Asp	Ser	
	140								145					150	
Cys	Ala	Pro	Pro	Thr	Gly	Phe	Asn	Lys	Pro	Thr	Ser	Asn	Asp	Thr	
	155								160					165	
Met	Ala	Ser	Gly	Trp	Arg	Ala	Ser	Ser	Phe	His	Phe	Asp	Ser	Glu	
	170								175					180	
Glu	Asn	Lys	His	Arg	Leu	Ile	His	Phe	Ser	Val	Phe	Leu	Gly	Leu	
	185								190					195	
Leu	Leu	Val	Gly	Ile	Leu	Glu	Val	Leu	Phe	Gly	Leu	Ser	Gln	Ile	
	200								205					210	
Val	Ile	Gly	Phe	Leu	Gly	Cys	Leu	Cys	Gly	Val	Ser	Lys	Arg	Arg	
	215								220					225	
Ser	Gln	Ile	Val												

<210> SEQ ID NO 49
<211> LENGTH: 636
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 49

atccgttctc tgcgctgcc a gctcaggtga gccctcgcca aggtgacctc	50
gcaggacact ggtgaaggag cagtgaggaa cctgcagagt cacacagttg	100
ctgaccaatt gagctgtgag cctggagcag atccgtgggc tgcagacccc	150
cgccccagtg cctctcccc tgcagccctg cccctcgaac tgtgacatgg	200
agagagtgc cctggccctt ctctactgg caggcctgac tgccttgga	250
gccaatgacc catttgcaa taaagacgat cccttctact atgactggaa	300
aaacctgcag ctgagcggac tgatctgcgg agggctcctg gccattgctg	350
ggatcgcggc agttctgagt ggcaaatgca aatacaagag cagccagaag	400
cagcacagtc ctgtacctga gaaggccatc cactcatca ctccaggctc	450
tgccactact tgctgagcac aggactggcc tccagggatg gctgaagcc	500
taacactggc cccagcacc tcctcccctg ggaggcctta tcctcaagga	550
aggacttctc tccaagggca ggctgttagg cccctttctg atcaggaggc	600
ttctttatga attaaactcg cccaccacc cctca	636

-continued

<210> SEQ ID NO 50
<211> LENGTH: 89
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 50

Met	Glu	Arg	Val	Thr	Leu	Ala	Leu	Leu	Leu	Leu	Ala	Gly	Leu	Thr
1				5					10				15	
Ala	Leu	Glu	Ala	Asn	Asp	Pro	Phe	Ala	Asn	Lys	Asp	Asp	Pro	Phe
			20						25				30	
Tyr	Tyr	Asp	Trp	Lys	Asn	Leu	Gln	Leu	Ser	Gly	Leu	Ile	Cys	Gly
			35					40					45	
Gly	Leu	Leu	Ala	Ile	Ala	Gly	Ile	Ala	Ala	Val	Leu	Ser	Gly	Lys
			50					55					60	
Cys	Lys	Tyr	Lys	Ser	Ser	Gln	Lys	Gln	His	Ser	Pro	Val	Pro	Glu
			65					70					75	
Lys	Ala	Ile	Pro	Leu	Ile	Thr	Pro	Gly	Ser	Ala	Thr	Thr	Cys	
			80					85						

<210> SEQ ID NO 51
<211> LENGTH: 1734
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 51

gtggactctg agaagccag gcagttgagg acaggagaga gaaggctgca	50
gacccagagg gagggaggac agggagtcgg aaggaggagg acagaggagg	100
gcacagagac gcagagcaag ggcggcaagg aggagaccct ggtgggagga	150
agacactctg gagagagagg gggctgggca gagatgaagt tccaggggcc	200
cctggcctgc ctctgtctgg ccctctgcct gggcagtgga gaggtggcc	250
ccctgcagag cggagaggaa agcactggga caaatattgg ggaggccctt	300
ggacatggcc tgggagacgc cctgagcgaa ggggtgggaa aggccattgg	350
caaaagagcc ggaggggcag ctggctctaa agtcagttag gcccttgcc	400
aagggaccag agaagcagtt ggcactggag tcaggcaggt tccaggcttt	450
ggcgacgacg atgctttggg caacagggtc ggggaagcag cccatgctct	500
gggaaacact gggcacgaga ttggcagaca ggcagaagat gtcattcgac	550
acggagcaga tgctgtccgc ggctcctggc aggggggtgcc tggccacagt	600
ggtgcttggg aaacttctgg aggccatggc atctttggct ctcaaggtgg	650
ccttgaggag cagggccagg gcaatcctgg aggtctgggg actccgtggg	700
tccacggata ccccggaac tcagcaggca gctttggaat gaatcctcag	750
ggagctccct ggggtcaagg aggcaatgga gggccaccaa actttgggac	800
caacactcag ggagctgtgg cccagcctgg ctatggttca gtgagagcca	850
gcaaccagaa tgaaggtgac acgaatcccc caccatctgg ctcaagtgga	900
ggctccagca actctggggg aggcagcggc tcacagtcgg gcagcagtg	950
cagtggcagc aatggtgaca acaacaatgg cagcagcagt ggtggcagca	1000
gcagtggcag cagcagtggc agcagcagtg gggcagcag tggcggcagc	1050
agtggtggca gcagtggaac cagtggtggc agcagaggtg acagcggcag	1100

-continued

tgagtctctcc tggggatcca gcaccggctc ctcctccggc aaccacggtg	1150
ggagcggcgg aggaaatgga cataaacccg ggtgtgaaaa gccagggaat	1200
gaagcccgcg ggagcgggga atctgggatt cagggcttca gaggacaggg	1250
agtttccagc aacatgaggg aaataagcaa agagggcaat cgcctccttg	1300
gaggctctgg agacaattat cgggggcaag ggtcgagctg gggcagtgga	1350
ggaggtgacg ctgttggtgg agtcaatact gtgaactctg agacgtctcc	1400
tgggatgttt aactttgaca ctttctggaa gaattttaaa tccaagctgg	1450
gtttcatcaa ctgggatgcc ataaacaagg accagagaag ctctcgcac	1500
ccgtgacctc cagacaagga gccaccagat tggatgggag ccccccacact	1550
ccctccttaa aacaccaccc tctcatcact aatctcagcc cttgcccttg	1600
aaataaacct tagctgcccc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1650
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1700
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa	1734

<210> SEQ ID NO 52
<211> LENGTH: 440
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 52

Met Lys Phe Gln Gly Pro Leu Ala Cys Leu Leu Leu Ala Leu Cys	
1 5 10 15	
Leu Gly Ser Gly Glu Ala Gly Pro Leu Gln Ser Gly Glu Glu Ser	
20 25 30	
Thr Gly Thr Asn Ile Gly Glu Ala Leu Gly His Gly Leu Gly Asp	
35 40 45	
Ala Leu Ser Glu Gly Val Gly Lys Ala Ile Gly Lys Glu Ala Gly	
50 55 60	
Gly Ala Ala Gly Ser Lys Val Ser Glu Ala Leu Gly Gln Gly Thr	
65 70 75	
Arg Glu Ala Val Gly Thr Gly Val Arg Gln Val Pro Gly Phe Gly	
80 85 90	
Ala Ala Asp Ala Leu Gly Asn Arg Val Gly Glu Ala Ala His Ala	
95 100 105	
Leu Gly Asn Thr Gly His Glu Ile Gly Arg Gln Ala Glu Asp Val	
110 115 120	
Ile Arg His Gly Ala Asp Ala Val Arg Gly Ser Trp Gln Gly Val	
125 130 135	
Pro Gly His Ser Gly Ala Trp Glu Thr Ser Gly Gly His Gly Ile	
140 145 150	
Phe Gly Ser Gln Gly Gly Leu Gly Gly Gln Gly Gln Gly Asn Pro	
155 160 165	
Gly Gly Leu Gly Thr Pro Trp Val His Gly Tyr Pro Gly Asn Ser	
170 175 180	
Ala Gly Ser Phe Gly Met Asn Pro Gln Gly Ala Pro Trp Gly Gln	
185 190 195	
Gly Gly Asn Gly Gly Pro Pro Asn Phe Gly Thr Asn Thr Gln Gly	
200 205 210	

-continued

Ala Val Ala Gln Pro Gly Tyr Gly Ser Val Arg Ala Ser Asn Gln
215 220 225
Asn Glu Gly Cys Thr Asn Pro Pro Pro Ser Gly Ser Gly Gly Gly
230 235 240
Ser Ser Asn Ser Gly Gly Gly Ser Gly Ser Gln Ser Gly Ser Ser
245 250 255
Gly Ser Gly Ser Asn Gly Asp Asn Asn Asn Gly Ser Ser Ser Gly
260 265 270
Gly Ser Ser Ser Gly Ser Ser Ser Gly Ser Ser Ser Gly Gly Ser
275 280 285
Ser Gly Gly Ser Ser Gly Gly Ser Ser Gly Asn Ser Gly Gly Ser
290 295 300
Arg Gly Asp Ser Gly Ser Glu Ser Ser Trp Gly Ser Ser Thr Gly
305 310 315
Ser Ser Ser Gly Asn His Gly Gly Ser Gly Gly Gly Asn Gly His
320 325 330
Lys Pro Gly Cys Glu Lys Pro Gly Asn Glu Ala Arg Gly Ser Gly
335 340 345
Glu Ser Gly Ile Gln Gly Phe Arg Gly Gln Gly Val Ser Ser Asn
350 355 360
Met Arg Glu Ile Ser Lys Glu Gly Asn Arg Leu Leu Gly Gly Ser
365 370 375
Gly Asp Asn Tyr Arg Gly Gln Gly Ser Ser Trp Gly Ser Gly Gly
380 385 390
Gly Asp Ala Val Gly Gly Val Asn Thr Val Asn Ser Glu Thr Ser
395 400 405
Pro Gly Met Phe Asn Phe Asp Thr Phe Trp Lys Asn Phe Lys Ser
410 415 420
Lys Leu Gly Phe Ile Asn Trp Asp Ala Ile Asn Lys Asp Gln Arg
425 430 435
Ser Ser Arg Ile Pro
440

<210> SEQ ID NO 53
<211> LENGTH: 1676
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 53

ggagaagagg ttgtgtggga caagctgctc ccgacagaag gatgtcgctg 50
ctgagcctgc cctggctggg cctcagaccg gtggcaatgt ccccatggct 100
actcctgctg ctggttgtgg gctcctggct actcgccgc atcctggctt 150
ggacctatgc cttctataac aactgccgc ggctccagtg tttccacag 200
ccccaaaaa ggaactgggt ttgggtcac ctgggcctga tcaactctac 250
agaggagggc ttgaaggact cgaccagat gtcggccacc tattcccagg 300
gctttacggt atggctgggt cccatcatcc cttcatcgt tttatgccac 350
cctgacacca tccggtctat caccaatgcc tcagctgcca ttgcaccaa 400
ggataatctc ttcacaggt tcctgaagcc ctggctggga gaagggatac 450
tgctgagtgg cggtgacaag tggagccgc accgtcggat gctgacgcc 500

-continued

gccttcatt tcaacatcct gaagtcctat ataacgatct tcaacaagag 550
tgcaaacatc atgcttgaca agtggcagca cctggcctca gagggcagca 600
gtcgtctgga catgtttgag cacatcagcc tcatgacctt ggacagtcta 650
cagaaatgca tcttcagctt tgacagccat tgtcaggaga ggcccagtga 700
atatattgcc accatcttgg agctcagtgc ccttgtagag aaaagaagcc 750
agcatatcct ccagcacatg gactttctgt attacctctc ccatgacggg 800
cggcgcttcc acagggcctg ccgcctggtg catgacttca cagacgctgt 850
catccgggag cggcgtcgca ccctcccccac tcagggtatt gatgattttt 900
tcaaagacaa agccaagtcc aagactttgg atttcattga tgtgcttctg 950
ctgagcaagg atgaagatgg gaaggcattg tcagatgagg atataagagc 1000
agaggctgac accttcatgt ttggaggcca tgacaccacg gccagtggcc 1050
ttctctgggt cctgtacaac cttgcgaggc acccagaata ccaggagcgc 1100
tgccgacagg aggtgcaaga gcttctgaag gaccgcgatc ctaaagagat 1150
tgaatgggac gacctggccc agctgccctt cctgaccatg tgcgtgaagg 1200
agagcctgag gttacatccc ccagctccct tcatctcccg atgctgcacc 1250
caggacattg ttctcccaga tggccgagtc atcccccagg gcattacctg 1300
cctcatcgat attatagggg tccatcacia cccaactgtg tggccggatc 1350
ctgagggtcta cgacccttc cgctttgacc cagagaacag caaggggagg 1400
tcacctctgg cttttattcc tttctccgca gggcccagga actgcatcgg 1450
gcaggcgctt gccatggcgg agatgaaagt ggtcctggcg ttgatgctgc 1500
tgcaacttccg gttcctgcca gaccacactg agccccgcag gaagctggaa 1550
ttgatcatgc gcgccgaggg cgggctttgg ctgcgggtgg agcccctgaa 1600
tgtaggcttg cagtgacttt ctgaccatc cactgtttt tttgcagatt 1650
gtcatgaata aaacggtgct gtcaaa 1676

<210> SEQ ID NO 54
<211> LENGTH: 524
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 54

Met Ser Leu Leu Ser Leu Pro Trp Leu Gly Leu Arg Pro Val Ala
1 5 10 15
Met Ser Pro Trp Leu Leu Leu Leu Val Val Gly Ser Trp Leu
20 25 30
Leu Ala Arg Ile Leu Ala Trp Thr Tyr Ala Phe Tyr Asn Asn Cys
35 40 45
Arg Arg Leu Gln Cys Phe Pro Gln Pro Pro Lys Arg Asn Trp Phe
50 55 60
Trp Gly His Leu Gly Leu Ile Thr Pro Thr Glu Glu Gly Leu Lys
65 70 75
Asp Ser Thr Gln Met Ser Ala Thr Tyr Ser Gln Gly Phe Thr Val
80 85 90
Trp Leu Gly Pro Ile Ile Pro Phe Ile Val Leu Cys His Pro Asp

-continued

95					100					105				
Thr	Ile	Arg	Ser	Ile	Thr	Asn	Ala	Ser	Ala	Ala	Ile	Ala	Pro	Lys
				110					115					120
Asp	Asn	Leu	Phe	Ile	Arg	Phe	Leu	Lys	Pro	Trp	Leu	Gly	Glu	Gly
				125					130					135
Ile	Leu	Leu	Ser	Gly	Gly	Asp	Lys	Trp	Ser	Arg	His	Arg	Arg	Met
				140					145					150
Leu	Thr	Pro	Ala	Phe	His	Phe	Asn	Ile	Leu	Lys	Ser	Tyr	Ile	Thr
				155					160					165
Ile	Phe	Asn	Lys	Ser	Ala	Asn	Ile	Met	Leu	Asp	Lys	Trp	Gln	His
				170					175					180
Leu	Ala	Ser	Glu	Gly	Ser	Ser	Arg	Leu	Asp	Met	Phe	Glu	His	Ile
				185					190					195
Ser	Leu	Met	Thr	Leu	Asp	Ser	Leu	Gln	Lys	Cys	Ile	Phe	Ser	Phe
				200					205					210
Asp	Ser	His	Cys	Gln	Glu	Arg	Pro	Ser	Glu	Tyr	Ile	Ala	Thr	Ile
				215					220					225
Leu	Glu	Leu	Ser	Ala	Leu	Val	Glu	Lys	Arg	Ser	Gln	His	Ile	Leu
				230					235					240
Gln	His	Met	Asp	Phe	Leu	Tyr	Tyr	Leu	Ser	His	Asp	Gly	Arg	Arg
				245					250					255
Phe	His	Arg	Ala	Cys	Arg	Leu	Val	His	Asp	Phe	Thr	Asp	Ala	Val
				260					265					270
Ile	Arg	Glu	Arg	Arg	Arg	Thr	Leu	Pro	Thr	Gln	Gly	Ile	Asp	Asp
				275					280					285
Phe	Phe	Lys	Asp	Lys	Ala	Lys	Ser	Lys	Thr	Leu	Asp	Phe	Ile	Asp
				290					295					300
Val	Leu	Leu	Leu	Ser	Lys	Asp	Glu	Asp	Gly	Lys	Ala	Leu	Ser	Asp
				305					310					315
Glu	Asp	Ile	Arg	Ala	Glu	Ala	Asp	Thr	Phe	Met	Phe	Gly	Gly	His
				320					325					330
Asp	Thr	Thr	Ala	Ser	Gly	Leu	Ser	Trp	Val	Leu	Tyr	Asn	Leu	Ala
				335					340					345
Arg	His	Pro	Glu	Tyr	Gln	Glu	Arg	Cys	Arg	Gln	Glu	Val	Gln	Glu
				350					355					360
Leu	Leu	Lys	Asp	Arg	Asp	Pro	Lys	Glu	Ile	Glu	Trp	Asp	Asp	Leu
				365					370					375
Ala	Gln	Leu	Pro	Phe	Leu	Thr	Met	Cys	Val	Lys	Glu	Ser	Leu	Arg
				380					385					390
Leu	His	Pro	Pro	Ala	Pro	Phe	Ile	Ser	Arg	Cys	Cys	Thr	Gln	Asp
				395					400					405
Ile	Val	Leu	Pro	Asp	Gly	Arg	Val	Ile	Pro	Lys	Gly	Ile	Thr	Cys
				410					415					420
Leu	Ile	Asp	Ile	Ile	Gly	Val	His	His	Asn	Pro	Thr	Val	Trp	Pro
				425					430					435
Asp	Pro	Glu	Val	Tyr	Asp	Pro	Phe	Arg	Phe	Asp	Pro	Glu	Asn	Ser
				440					445					450
Lys	Gly	Arg	Ser	Pro	Leu	Ala	Phe	Ile	Pro	Phe	Ser	Ala	Gly	Pro
				455					460					465
Arg	Asn	Cys	Ile	Gly	Gln	Ala	Phe	Ala	Met	Ala	Glu	Met	Lys	Val
				470					475					480

-continued

Val Leu Ala Leu Met Leu Leu His Phe Arg Phe Leu Pro Asp His
485 490 495
Thr Glu Pro Arg Arg Lys Leu Glu Leu Ile Met Arg Ala Glu Gly
500 505 510
Gly Leu Trp Leu Arg Val Glu Pro Leu Asn Val Gly Leu Gln
515 520

<210> SEQ ID NO 55
<211> LENGTH: 644
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 55
atcgcatcaa ttgggagtac catcttcctc atgggaccag tgaacacgct 50
gaagcgaatg tttgagccta ctcgtttgat tgcaactatc atggtgctgt 100
tgtgttttgc acttacccctg tgttctgcct tttggtggca taacaaggga 150
cttgcaactta tcttctgcat ttgcagctct ttggcattga cgtggtacag 200
cctttccttc ataccatttg caagggatgc tgtgaagaag tgttttgccg 250
tgtgtcttgc ataattcatg gccagtttta tgaagctttg gaaggcacta 300
tggacagaag ctggtggaca gttttgtaac tatcttcgaa acctctgtct 350
tacagacatg tgccttttat ctgcagcaa tgtgttgctt gtgattcgaa 400
catttgaggg ttacttttgg aagcaacaat acattctcga acctgaatgt 450
cagtagcaca ggatgagaag tgggttctgt atcttgtgga gtggaatctt 500
cctcatgtac ctgtttcctc tctggatgtt gtccactga attcccatga 550
atacaaacct attcagcaac agcaaaaaaa aaaaaaaaaa aaaaaaaaaa 600
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 644

<210> SEQ ID NO 56
<211> LENGTH: 77
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 56
Met Gly Pro Val Lys Gln Leu Lys Arg Met Phe Glu Pro Thr Arg
1 5 10 15
Leu Ile Ala Thr Ile Met Val Leu Leu Cys Phe Ala Leu Thr Leu
20 25 30
Cys Ser Ala Phe Trp Trp His Asn Lys Gly Leu Ala Leu Ile Phe
35 40 45
Cys Ile Leu Gln Ser Leu Ala Leu Thr Trp Tyr Ser Leu Ser Phe
50 55 60
Ile Pro Phe Ala Arg Asp Ala Val Lys Lys Cys Phe Ala Val Cys
65 70 75
Leu Ala

<210> SEQ ID NO 57
<211> LENGTH: 3334
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 57

-continued

cggtctcgagc tcgagccgaa tcggctcgag gggcagtgga gcacccagca	50
ggccgccaac atgctctgtc tgtgcctgta cgtgccggtc atcggggaag	100
cccagaccga gttccagtac tttagtcga aggggctccc tgccgagctg	150
aagtccattt tcaagctcag tgtcttcac cctcccagg aattctccac	200
ctaccgccag tggaagcaga aaattgtaca agctggagat aaggaccttg	250
atgggcagct agactttgaa gaatttgtcc attatctcca agatcatgag	300
aagaagctga ggctggtgtt taagattttg gacaaaaaga atgatggacg	350
cattgacgag caggagatca tgcagtcctt gcgggacttg ggagtcaaga	400
tatctgaaca gcaggcagaa aaaattctca agagcatgga taaaaacggc	450
acgatgacca tcgactggaa cgagtggaga gactaccacc tcctccaccc	500
cgtgaaaaac atccccgaga tcctcctcta ctggaagcat tccacgatct	550
ttgatgtggg tgagaatcta acggctcccg atgagttcac agtggaggag	600
aggcagacgg ggatgtggtg gagacacctg gtggcaggag gtggggcagg	650
ggcgtatcc agaacctgca cggccccctt ggacaggctc aagggtctca	700
tgcagggtcca tgcctccgc agcaacaaca tgggcatcgt tggtggttc	750
actcagatga ttcgagaagg aggggccagg tcaactctggc ggggcaatgg	800
catcaacgtc ctcaaaattg cccccaatc agccatcaaa ttcattggct	850
atgagcagat caagcgctt gttgtagtg accaggagac tctgaggatt	900
cacgagaggg ttgtggcagg gtccttggca ggggccatcg cccagagcag	950
catctaccca atggaggtcc tgaagaccg gatggcgctg cgaagacag	1000
gccagtactc aggaatgctg gactgcgcca ggaggatcct ggccagagag	1050
gggggtggcg ccttctacaa aggtatgtc cccaacatgc tgggcatcat	1100
cccctatgcc ggcacgacc ttgcagtcta cgagacgctc aagaatgcct	1150
ggctgcagca ctatgcagtg aacagcgcg accccggcgt gtttgtgctc	1200
ctggcctgtg gcaccatgtc cagtacctgt ggccagctgg ccagctaccc	1250
cctggcccta gtcaggacct ggatgcaggc gcaagcctct attgagggcg	1300
ctccggaggt gaccatgagc agcctcttca aacatatcct gcggaccgag	1350
ggggccttcg ggctgtacag ggggctggcc cccaacttca tgaaggatcat	1400
cccagctgtg agcatcagct acgtgttcta cgagaacctg aagatcacc	1450
tgggcgtgca gtcgcgtga cggggggagg gcccccggc agtggactcg	1500
ctgatcctgg gccgcagcct ggggtgtgca gccatctcat tctgtgaatg	1550
tgccaacact aagctgtctc gagccaagct gtgaaaacc tagacgcacc	1600
cgcagggagg gtggggagag ctggcaggcc cagggttgt cctgctgacc	1650
ccagcagacc ctctgttgg ttccagcgaa gaccacaggc attccttagg	1700
gtccagggtc agcaggctcc gggctcacat gtgtaaggac aggacattt	1750
ctgcagtgcc tgccaatagt gagcttggag cctggaggcc ggcttagttc	1800
ttccatttca cccttgacg cagctgttgg ccacggcccc tgccctctgg	1850
tctgcgctgc atctccctgt gccctcttgc tgccctgctg tctgctgag	1900

-continued

taaggtggga ggagggctac agcccacatc ccaccccctc gtccaatccc	1950
ataatccatg atgaaaggtg aggtcacgtg gcctcccagg cctgacttcc	2000
caacctacag cattgacgcc aacttggtctg tgaaggaaga ggaaaggatc	2050
tggccttggtg gtcactggca tctgagccct gctgatggct ggggctctcg	2100
ggcatgcttg ggagtgacag gggctcgggc tgctggcct ggctgcacag	2150
aaggcaagtg ctggggctca tgggtgctctg agctggcctg gaccctgtca	2200
ggatggggccc cacctcagaa ccaaactcac tgtcccact gtggcatgag	2250
ggcagtgag caccatgttt gagggcgaag ggcagagcgt ttgtgtgttc	2300
tggggaggga aggaaaaggt gttggaggcc ttaattatgg actgttggga	2350
aaagggtttt gtccagaagg acaagccgga caaatgagcg acttctgtgc	2400
ttccagagga agacgagga gcaggagctt ggctgactgc tcagagtctg	2450
ttctgacgcc ctgggggttc ctgtccaacc ccagcagggg cgcagcggga	2500
ccagccccac attccacttg tgctactgct tggaacctat ttattttgta	2550
tttatttgaa cagagttagt tcctaactat ttttatagat ttgtttaatt	2600
aatagcttgt catTTTcaag ttcatTTTT attcataatt atgttcatgg	2650
ttgattgtac cttcccaagc ccgccagtg ggatgggagg aggaggagaa	2700
ggggggcctt gggccgctgc agtcacatct gtccagagaa attccttttg	2750
ggactggagg cagaaaagcg gccagaagcg agcagccctg gctcctttcc	2800
tttggcaggt tggggaaggg ctgccccca gccttaggat ttcagggttt	2850
gactgggggg gtggagagag agggaggaac ctcaataacc ttgaagggtg	2900
aatccagtta tttcctgcgc tgcgagggtt tctttatttc actcttttct	2950
gaatgtcaag gcagtgaggt gcctctcact gtgaatttgt ggtgggcggg	3000
ggctggagga gaggggtggg ggctggctcc gtccctccca gccttctgct	3050
gcccttgctt aacaatgccg gccaaactggc gacctcacgg ttgcacttcc	3100
attccaccag aatgacctga tgaggaaatc ttcaatagga tgcaaagatc	3150
aatgcaaaaa ttgttatata tgaacatata actggagtcg tcaaaaagca	3200
aattaagaaa gaattggacg ttagaagttg tcattttaaag cagccttcta	3250
ataaagtgtg ttcaaagctg aaaaaaaaa aaaaaaaaa aaaaaaaaa	3300
aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaa	3334

<210> SEQ ID NO 58
<211> LENGTH: 469
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 58

Met	Leu	Cys	Leu	Cys	Leu	Tyr	Val	Pro	Val	Ile	Gly	Glu	Ala	Gln
1				5					10					15
Thr	Glu	Phe	Gln	Tyr	Phe	Glu	Ser	Lys	Gly	Leu	Pro	Ala	Glu	Leu
			20						25					30
Lys	Ser	Ile	Phe	Lys	Leu	Ser	Val	Phe	Ile	Pro	Ser	Gln	Glu	Phe
			35						40					45

-continued

Ser	Thr	Tyr	Arg	Gln	Trp	Lys	Gln	Lys	Ile	Val	Gln	Ala	Gly	Asp	50	55	60
Lys	Asp	Leu	Asp	Gly	Gln	Leu	Asp	Phe	Glu	Glu	Phe	Val	His	Tyr	65	70	75
Leu	Gln	Asp	His	Glu	Lys	Lys	Leu	Arg	Leu	Val	Phe	Lys	Ile	Leu	80	85	90
Asp	Lys	Lys	Asn	Asp	Gly	Arg	Ile	Asp	Ala	Gln	Glu	Ile	Met	Gln	95	100	105
Ser	Leu	Arg	Asp	Leu	Gly	Val	Lys	Ile	Ser	Glu	Gln	Gln	Ala	Glu	110	115	120
Lys	Ile	Leu	Lys	Ser	Met	Asp	Lys	Asn	Gly	Thr	Met	Thr	Ile	Asp	125	130	135
Trp	Asn	Glu	Trp	Arg	Asp	Tyr	His	Leu	Leu	His	Pro	Val	Glu	Asn	140	145	150
Ile	Pro	Glu	Ile	Ile	Leu	Tyr	Trp	Lys	His	Ser	Thr	Ile	Phe	Asp	155	160	165
Val	Gly	Glu	Asn	Leu	Thr	Val	Pro	Asp	Glu	Phe	Thr	Val	Glu	Glu	170	175	180
Arg	Gln	Thr	Gly	Met	Trp	Trp	Arg	His	Leu	Val	Ala	Gly	Gly	Gly	185	190	195
Ala	Gly	Ala	Val	Ser	Arg	Thr	Cys	Thr	Ala	Pro	Leu	Asp	Arg	Leu	200	205	210
Lys	Val	Leu	Met	Gln	Val	His	Ala	Ser	Arg	Ser	Asn	Asn	Met	Gly	215	220	225
Ile	Val	Gly	Gly	Phe	Thr	Gln	Met	Ile	Arg	Glu	Gly	Gly	Ala	Arg	230	235	240
Ser	Leu	Trp	Arg	Gly	Asn	Gly	Ile	Asn	Val	Leu	Lys	Ile	Ala	Pro	245	250	255
Glu	Ser	Ala	Ile	Lys	Phe	Met	Ala	Tyr	Glu	Gln	Ile	Lys	Arg	Leu	260	265	270
Val	Gly	Ser	Asp	Gln	Glu	Thr	Leu	Arg	Ile	His	Glu	Arg	Leu	Val	275	280	285
Ala	Gly	Ser	Leu	Ala	Gly	Ala	Ile	Ala	Gln	Ser	Ser	Ile	Tyr	Pro	290	295	300
Met	Glu	Val	Leu	Lys	Thr	Arg	Met	Ala	Leu	Arg	Lys	Thr	Gly	Gln	305	310	315
Tyr	Ser	Gly	Met	Leu	Asp	Cys	Ala	Arg	Arg	Ile	Leu	Ala	Arg	Glu	320	325	330
Gly	Val	Ala	Ala	Phe	Tyr	Lys	Gly	Tyr	Val	Pro	Asn	Met	Leu	Gly	335	340	345
Ile	Ile	Pro	Tyr	Ala	Gly	Ile	Asp	Leu	Ala	Val	Tyr	Glu	Thr	Leu	350	355	360
Lys	Asn	Ala	Trp	Leu	Gln	His	Tyr	Ala	Val	Asn	Ser	Ala	Asp	Pro	365	370	375
Gly	Val	Phe	Val	Leu	Leu	Ala	Cys	Gly	Thr	Met	Ser	Ser	Thr	Cys	380	385	390
Gly	Gln	Leu	Ala	Ser	Tyr	Pro	Leu	Ala	Leu	Val	Arg	Thr	Arg	Met	395	400	405
Gln	Ala	Gln	Ala	Ser	Ile	Glu	Gly	Ala	Pro	Glu	Val	Thr	Met	Ser	410	415	420
Ser	Leu	Phe	Lys	His	Ile	Leu	Arg	Thr	Glu	Gly	Ala	Phe	Gly	Leu			

-continued

425										430										435									
Tyr	Arg	Gly	Leu	Ala	Pro	Asn	Phe	Met	Lys	Val	Ile	Pro	Ala	Val															
				440						445				450															
Ser	Ile	Ser	Tyr	Val	Val	Tyr	Glu	Asn	Leu	Lys	Ile	Thr	Leu	Gly															
				455						460				465															
Val	Gln	Ser	Arg																										
<210> SEQ ID NO 59																													
<211> LENGTH: 1658																													
<212> TYPE: DNA																													
<213> ORGANISM: Homo Sapien																													
<400> SEQUENCE: 59																													
ggaaggcagc ggcagctcca ctcagccagt acccagatac gctgggaacc																									50				
ttccccagcc atggcttccc tggggcagat cctcttctgg agcataatta																									100				
gcatcatcat tattctggct ggagcaattg cactcatcat tggctttggt																									150				
atctcagga gacactccat cacagtcact actgtcgct cagctgggaa																									200				
cattggggag gatggaatcc tgagctgcac ttttgaacct gacatcaaac																									250				
ttttctgatat cgtgatacaa tggctgaagg aaggtgtttt aggcttggtc																									300				
catgagttca aagaaggcaa agatgagctg tcggagcagg atgaaatgtt																									350				
cagaggccgg acagcagtgt ttgctgatca agtgatagtt ggcaatgcct																									400				
ctttcgggct gaaaaacgtg caactcacag atgctggcac ctacaaatgt																									450				
tatatcatca cttctaaagg caaggggaat gctaaccttg agtataaaac																									500				
tggagccttc agcatgccgg aagtgaatgt ggactataat gccagctcag																									550				
agacottgag gtgtgaggct ccccgatggt tccccagcc cacagtggtc																									600				
tgggcatccc aagttgacca gggagccaac ttctcggaag tctccaatac																									650				
cagcttttag ctgaactctg agaatgtgac catgaaggtt gtgtctgtgc																									700				
tctacaatgt tacgatcaac aacacatact cctgtatgat tgaaaatgac																									750				
attgccaaag caacagggga tatcaaagt acagaatcgg agatcaaaag																									800				
gcggagtcat ctacagctgc taaactcaaa ggcttctctg tgtgtctctt																									850				
ctttctttgc catcagctgg gcaactctgc ctctcagccc ttacctgatg																									900				
ctaaaaaat gtgccttggc cacaaaaaag catgcaaagt cattgttaca																									950				
acagggatct acagaactat ttcaccacca gatatgacct agttttatat																									1000				
ttctgggagg aaatgaattc atatctagaa gtctggagt agcaaacaag																									1050				
agcaagaaac aaaaagaagc caaaagcaga aggctccaat atgaacaaga																									1100				
taaactctatc ttcaaagaca tattagaagt tgggaaaata attcatgtga																									1150				
actagacaag tgtgttaaga gtgataagta aaatgcacgt ggagacaagt																									1200				
gcatccccag atctcagga cctccccctg cctgtcacct ggggagtga																									1250				
aggacaggat agtgcagtgt ctttgtctct gaatttttag ttatatgtgc																									1300				
tgtaatgttg ctctgaggaa gccctggaa agtctatccc aacatatcca																									1350				
catcttatat tccacaaatt aagctgtagt atgtacccta agacgctgct																									1400				
aattgactgc cacttcgcaa ctcagggcg gctgcatttt agtaatgggt																									1450				

-continued

caaatgattc actttttatg atgcttccaa aggtgccttg gcttctcttc	1500
ccaactgaca aatgccaaag ttgagaaaaa tgatcataat tttagcataa	1550
acagagcagt cggggacacc gattttataa ataaactgag caccttcttt	1600
ttaaacaataa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1650
aaaaaaaaa	1658

<210> SEQ ID NO 60

<211> LENGTH: 282

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 60

Met	Ala	Ser	Leu	Gly	Gln	Ile	Leu	Phe	Trp	Ser	Ile	Ile	Ser	Ile	1	5	10	15
Ile	Ile	Ile	Leu	Ala	Gly	Ala	Ile	Ala	Leu	Ile	Ile	Gly	Phe	Gly	20	25	30	
Ile	Ser	Gly	Arg	His	Ser	Ile	Thr	Val	Thr	Thr	Val	Ala	Ser	Ala	35	40	45	
Gly	Asn	Ile	Gly	Glu	Asp	Gly	Ile	Leu	Ser	Cys	Thr	Phe	Glu	Pro	50	55	60	
Asp	Ile	Lys	Leu	Ser	Asp	Ile	Val	Ile	Gln	Trp	Leu	Lys	Glu	Gly	65	70	75	
Val	Leu	Gly	Leu	Val	His	Glu	Phe	Lys	Glu	Gly	Lys	Asp	Glu	Leu	80	85	90	
Ser	Glu	Gln	Asp	Glu	Met	Phe	Arg	Gly	Arg	Thr	Ala	Val	Phe	Ala	95	100	105	
Asp	Gln	Val	Ile	Val	Gly	Asn	Ala	Ser	Leu	Arg	Leu	Lys	Asn	Val	110	115	120	
Gln	Leu	Thr	Asp	Ala	Gly	Thr	Tyr	Lys	Cys	Tyr	Ile	Ile	Thr	Ser	125	130	135	
Lys	Gly	Lys	Gly	Asn	Ala	Asn	Leu	Glu	Tyr	Lys	Thr	Gly	Ala	Phe	140	145	150	
Ser	Met	Pro	Glu	Val	Asn	Val	Asp	Tyr	Asn	Ala	Ser	Ser	Glu	Thr	155	160	165	
Leu	Arg	Cys	Glu	Ala	Pro	Arg	Trp	Phe	Pro	Gln	Pro	Thr	Val	Val	170	175	180	
Trp	Ala	Ser	Gln	Val	Asp	Gln	Gly	Ala	Asn	Phe	Ser	Glu	Val	Ser	185	190	195	
Asn	Thr	Ser	Phe	Glu	Leu	Asn	Ser	Glu	Asn	Val	Thr	Met	Lys	Val	200	205	210	
Val	Ser	Val	Leu	Tyr	Asn	Val	Thr	Ile	Asn	Asn	Thr	Tyr	Ser	Cys	215	220	225	
Met	Ile	Glu	Asn	Asp	Ile	Ala	Lys	Ala	Thr	Gly	Asp	Ile	Lys	Val	230	235	240	
Thr	Glu	Ser	Glu	Ile	Lys	Arg	Arg	Ser	His	Leu	Gln	Leu	Leu	Asn	245	250	255	
Ser	Lys	Ala	Ser	Leu	Cys	Val	Ser	Ser	Phe	Phe	Ala	Ile	Ser	Trp	260	265	270	
Ala	Leu	Leu	Pro	Leu	Ser	Pro	Tyr	Leu	Met	Leu	Lys	275	280					

-continued

<210> SEQ ID NO 61

<211> LENGTH: 1617

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 61

tgacgtcaga atcaccatgg ccagctatcc ttaccggcag ggctgcccag	50
gagctgcagg acaagcacca ggagcccctc cgggtagcta ctaccctgga	100
cccccaata gtggagggca gtatggtagt gggctacccc ctggtggtgg	150
ttatgggggt cctgccctg gagggcctta tggaccacca gctggtgagg	200
ggccctatgg acacccaat cctgggatgt tcccctctgg aactccagga	250
ggaccatatg gcggtgcagc tcccgggggc ccctatggtc agccacctcc	300
aagttcctac ggtgcccagc agcctgggct ttatggacag ggtggcgccc	350
ctcccaatgt ggatcctgag gcctactcct ggttcagtc ggtggactca	400
gatcacagtg gctatatctc catgaaggag cttaaagcagg ccctgggtcaa	450
ctgcaattgg tcttcattca atgatgagac ctgcctcatg atgataaaca	500
tgtttgacaa gaccaagtca ggccgcctcg atgtctacgg cttctcagcc	550
ctgtggaaat tcatccagca gtggaagaac ctcttcagc agtatgaccg	600
ggaccgctcg ggctccatta gctacacaga gctgcagcaa gctctgtccc	650
aaatgggcta caacctgagc cccagttca ccagcttct ggtctccgc	700
tactgccac gctctgcaa tcctgccatg cagcttgacc gcttcatcca	750
gggtgtgcacc cagctgcagg tgctgacaga ggccttcgg gagaaggaca	800
cagctgtaca aggcaacatc cggtcagct tcgaggactt cgtaacctg	850
acagcttctc ggatgctatg acccaacct ctgtggagag tggagtgcac	900
cagggacctt tcctggcttc ttagagttag agaagtatgt ggacatctct	950
tcttttctcg tccctctaga agaacattct cccttgcttg atgcaacact	1000
gttccaaaag agggtgagaga gtccctgcac atagccacca aatagttagg	1050
accggggctg aggccacaca gataggggcc tgatggagga gaggatagaa	1100
gttgaatgtc ctgatggcca tgagcagttg agtggcacag cctggcacca	1150
ggagcaggtc cttgtaatgg agttagtgtc cagtcagctg agctccaccc	1200
tgatgccagt ggtgagtgtt catcggcctg ttaccgttag tacctgtgtt	1250
ccctcaccag gccatcctgt caaacgagcc cattttctcc aaagtggaat	1300
ctgaccaagc atgagagaga tctgtctatg ggaccagtgg cttggattct	1350
gccacaccca taaatccttg tgtgttaact tctagctgcc tggggctggc	1400
cctgctcaga caaatctgct ccctgggcat ctttggccag gcttctgcc	1450
cctgcagctg ggaccctca cttgcctgcc atgctctgct cggcttcagt	1500
ctccaggaga cagtggtcac ctctccctgc caatactttt tttaatttgc	1550
atttttttct atttggggcc aaaagtccag tgaattgta agcttcaata	1600
aaaggatgaa actctga	1617

<210> SEQ ID NO 62

<211> LENGTH: 284

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 62

Met Ala Ser Tyr Pro Tyr Arg Gln Gly Cys Pro Gly Ala Ala Gly
1 5 10 15
Gln Ala Pro Gly Ala Pro Pro Gly Ser Tyr Tyr Pro Gly Pro Pro
20 25 30
Asn Ser Gly Gly Gln Tyr Gly Ser Gly Leu Pro Pro Gly Gly Gly
35 40 45
Tyr Gly Gly Pro Ala Pro Gly Gly Pro Tyr Gly Pro Pro Ala Gly
50 55 60
Gly Gly Pro Tyr Gly His Pro Asn Pro Gly Met Phe Pro Ser Gly
65 70 75
Thr Pro Gly Gly Pro Tyr Gly Gly Ala Ala Pro Gly Gly Pro Tyr
80 85 90
Gly Gln Pro Pro Pro Ser Ser Tyr Gly Ala Gln Gln Pro Gly Leu
95 100 105
Tyr Gly Gln Gly Gly Ala Pro Pro Asn Val Asp Pro Glu Ala Tyr
110 115 120
Ser Trp Phe Gln Ser Val Asp Ser Asp His Ser Gly Tyr Ile Ser
125 130 135
Met Lys Glu Leu Lys Gln Ala Leu Val Asn Cys Asn Trp Ser Ser
140 145 150
Phe Asn Asp Glu Thr Cys Leu Met Met Ile Asn Met Phe Asp Lys
155 160 165
Thr Lys Ser Gly Arg Ile Asp Val Tyr Gly Phe Ser Ala Leu Trp
170 175 180
Lys Phe Ile Gln Gln Trp Lys Asn Leu Phe Gln Gln Tyr Asp Arg
185 190 195
Asp Arg Ser Gly Ser Ile Ser Tyr Thr Glu Leu Gln Gln Ala Leu
200 205 210
Ser Gln Met Gly Tyr Asn Leu Ser Pro Gln Phe Thr Gln Leu Leu
215 220 225
Val Ser Arg Tyr Cys Pro Arg Ser Ala Asn Pro Ala Met Gln Leu
230 235 240
Asp Arg Phe Ile Gln Val Cys Thr Gln Leu Gln Val Leu Thr Glu
245 250 255
Ala Phe Arg Glu Lys Asp Thr Ala Val Gln Gly Asn Ile Arg Leu
260 265 270
Ser Phe Glu Asp Phe Val Thr Met Thr Ala Ser Arg Met Leu
275 280

<210> SEQ ID NO 63
<211> LENGTH: 1234
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 63

caggatgcag ggccgcgtgg caggagactg cgctcctctg ggctgctcc 50
tggtctgtct tcatctccca ggctcttttg cccggagcat cgggtgtgtg 100
gaggagaaa tttcccaaaa cttcgggacc aacttgctc agctcggaca 150

```

<210> SEQ ID NO 64
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 64

Met Gln Gly Arg Val Ala Gly Ser Cys Ala Pro Leu Gly Leu Leu
  1             5             10             15

Leu Val Cys Leu His Leu Pro Gly Leu Phe Ala Arg Ser Ile Gly
          20             25             30

Val Val Glu Glu Lys Val Ser Gln Asn Phe Gly Thr Asn Leu Pro
          35             40             45

Gln Leu Gly Gln Pro Ser Ser Thr Gly Pro Ser Asn Ser Glu His
          50             55             60

Pro Gln Pro Ala Leu Asp Pro Arg Ser Asn Asp Leu Ala Arg Val
          65             70             75

Pro Leu Lys Leu Ser Val Pro Pro Ser Asp Gly Phe Pro Pro Ala
          80             85             90

Gly Gly Ser Ala Val Gln Arg Trp Pro Pro Ser Trp Gly Leu Pro
          95             100            105

Ala Met Asp Ser Trp Pro Pro Glu Asp Pro Trp Gln Met Met Ala
          110            115            120

```

-continued

Ala	Ala	Ala	Glu	Asp	Arg	Leu	Gly	Glu	Ala	Leu	Pro	Glu	Glu	Leu
			125						130					135
Ser	Tyr	Leu	Ser	Ser	Ala	Ala	Ala	Leu	Ala	Pro	Gly	Ser	Gly	Pro
			140						145					150
Leu	Pro	Gly	Glu	Ser	Ser	Pro	Asp	Ala	Thr	Gly	Leu	Ser	Pro	Glu
			155						160					165
Ala	Ser	Leu	Leu	His	Gln	Asp	Ser	Glu	Ser	Arg	Arg	Leu	Pro	Arg
			170						175					180
Ser	Asn	Ser	Leu	Gly	Ala	Gly	Gly	Lys	Ile	Leu	Ser	Gln	Arg	Pro
			185						190					195
Pro	Trp	Ser	Leu	Ile	His	Arg	Val	Leu	Pro	Asp	His	Pro	Trp	Gly
			200						205					210
Thr	Leu	Asn	Pro	Ser	Val	Ser	Trp	Gly	Gly	Gly	Gly	Pro	Gly	Thr
			215						220					225
Gly	Trp	Gly	Thr	Arg	Pro	Met	Pro	His	Pro	Glu	Gly	Ile	Trp	Gly
			230						235					240
Ile	Asn	Asn	Gln	Pro	Pro	Gly	Thr	Ser	Trp	Gly	Asn	Ile	Asn	Arg
			245						250					255
Tyr	Pro	Gly	Gly	Ser	Trp	Gly	Asn	Ile	Asn	Arg	Tyr	Pro	Gly	Gly
			260						265					270
Ser	Trp	Gly	Asn	Ile	Asn	Arg	Tyr	Pro	Gly	Gly	Ser	Trp	Gly	Asn
			275						280					285
Ile	His	Leu	Tyr	Pro	Gly	Ile	Asn	Asn	Pro	Phe	Pro	Pro	Gly	Val
			290						295					300
Leu	Arg	Pro	Pro	Gly	Ser	Ser	Trp	Asn	Ile	Pro	Ala	Gly	Phe	Pro
			305						310					315
Asn	Pro	Pro	Ser	Pro	Arg	Leu	Gln	Trp	Gly					
			320						325					

<210> SEQ ID NO 65
<211> LENGTH: 422
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

aaggagaggc caccgggact tcagtgtctc ctccatccca ggagcgcagt	50
ggccactatg gggctctgggc tgccccttgt cctcctcttg accctccttg	100
gcagctcaca tggaacaggg ccgggtatga ctttgcaact gaagctgaag	150
gagtcttttc tgacaaattc ctccatgag tccagcttcc tggaattgct	200
tgaaaagctc tgcctcctcc tccatctccc ttcaggggacc agcgtcacc	250
tccaccatgc aagatctcaa caccatgttg tctgcaacac atgacagcca	300
ttgaagcctg tgtccttctt ggcccgggct tttgggccgg ggatgcagga	350
ggcaggcccc gaccctgtct ttcagcaggc cccaccctc ctgagtggca	400
ataaataaaa ttcggtatgc tg	422

<210> SEQ ID NO 66
<211> LENGTH: 78
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 66

-continued

Met	Gly	Ser	Gly	Leu	Pro	Leu	Val	Leu	Leu	Leu	Thr	Leu	Leu	Gly
1				5					10					15
Ser	Ser	His	Gly	Thr	Gly	Pro	Gly	Met	Thr	Leu	Gln	Leu	Lys	Leu
			20						25					30
Lys	Glu	Ser	Phe	Leu	Thr	Asn	Ser	Ser	Tyr	Glu	Ser	Ser	Phe	Leu
			35						40					45
Glu	Leu	Leu	Glu	Lys	Leu	Cys	Leu	Leu	Leu	His	Leu	Pro	Ser	Gly
			50						55					60
Thr	Ser	Val	Thr	Leu	His	His	Ala	Arg	Ser	Gln	His	His	Val	Val
			65						70					75
Cys	Asn	Thr												

<210> SEQ ID NO 67
<211> LENGTH: 744
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 67

acggaccgag ggttcgaggg agggacacgg accaggaacc tgagctaggt	50
caaagacgcc cgggccaggt gccccgtcgc aggtgccctt ggccggagat	100
gcggtaggag gggcgagcgc gagaagcccc ttcctcggcg ctgccaaacc	150
gccaccacgc ccatggcgaa ccccgggctg gggctgcttc tggcgctggg	200
cctgccgttc ctgctggccc gctggggccg agcctggggg caaatacaga	250
ccacttctgc aaatgagaat agcactgttt tgccttcata caccagctcc	300
agctccgatg gcaacctgcg tccggaagcc atcactgcta tcatcgtagt	350
cttctccctc ttggctgcct tgetcctggc tgtggggctg gcactgttgg	400
tgcggaagct tcgggagaag cggcagacgg agggcaccta ccggcccagt	450
agcgaggagc agttctccca tgcagccgag gcccgggccc ctcaggactc	500
caaggagacg gtgcagggct gcctgcccat ctaggctccc tctcctgcat	550
ctgtctccct tcattgctgt gtgaccttgg ggaaaggcag tgccctctct	600
gggcagtcag atccaccacg tgcttaatag cagggaagaa ggtacttcaa	650
agactctgcc cctgaggtca agagaggatg gggctattca cttttatata	700
tttatataaa attagtagtg agatgtaaaa aaaaaaaaaa aaaa	744

<210> SEQ ID NO 68
<211> LENGTH: 123
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 68

Met	Ala	Asn	Pro	Gly	Leu	Gly	Leu	Leu	Leu	Ala	Leu	Gly	Leu	Pro
1				5					10					15
Phe	Leu	Leu	Ala	Arg	Trp	Gly	Arg	Ala	Trp	Gly	Gln	Ile	Gln	Thr
			20						25					30
Thr	Ser	Ala	Asn	Glu	Asn	Ser	Thr	Val	Leu	Pro	Ser	Ser	Thr	Ser
			35						40					45
Ser	Ser	Ser	Asp	Gly	Asn	Leu	Arg	Pro	Glu	Ala	Ile	Thr	Ala	Ile
			50						55					60
Ile	Val	Val	Phe	Ser	Leu	Leu	Ala	Ala	Leu	Leu	Leu	Ala	Val	Gly

-continued

												65	70	75
Leu	Ala	Leu	Leu	Val	Arg	Lys	Leu	Arg	Glu	Lys	Arg	Gln	Thr	Glu
												80	85	90
Gly	Thr	Tyr	Arg	Pro	Ser	Ser	Glu	Glu	Gln	Phe	Ser	His	Ala	Ala
												95	100	105
Glu	Ala	Arg	Ala	Pro	Gln	Asp	Ser	Lys	Glu	Thr	Val	Gln	Gly	Cys
												110	115	120
Leu Pro Ile														
<210> SEQ ID NO 69														
<211> LENGTH: 3265														
<212> TYPE: DNA														
<213> ORGANISM: Homo Sapien														
<400> SEQUENCE: 69														
gccaggaata actagagagg aacaatgggg ttattcagag gttttgtttt													50	
cctcttagtt ctgtgcctgc tgcaccagtc aaatacttcc ttcattaagc													100	
tgaataataa tggctttgaa gatattgtca ttgttataga tcctagtgtg													150	
ccagaagatg aaaaaataat tgaacaaata gaggatatgg tgactacagc													200	
ttctacgtac ctgtttgaag ccacagaaaa aagatttttt ttcaaaaatg													250	
tatctatatt aattcctgag aattggaagg aaaatcctca gtacaaaagg													300	
ccaaaacatg aaaaccataa acatgctgat gttatagttg caccacctac													350	
actcccaggt agagatgaac catacaccaa gcagttcaca gaatgtggag													400	
agaaaggcga atacattcac ttcaccccctg accttctact tggaaaaaaa													450	
caaaatgaat atggaccacc aggcaaacctg tttgtccatg agtgggctca													500	
cctccggtgg ggagtgtttg atgagtacaa tgaagatcag cctttctacc													550	
gtgctaagtc aaaaaaaatc gaagcaacaa ggtgttccgc aggtatctct													600	
ggtagaaata gagtttataa gtgtcaagga ggcagctgtc ttagtagagc													650	
atgcagaatt gattctacaa caaaactgta tggaaaagat tgtcaattct													700	
ttcctgataa agtacaaaca gaaaagcat ccataatggt tatgcaaagt													750	
attgattctg ttgttgaatt ttgtaacgaa aaaaccata atcaagaagc													800	
tccaagccta caaaacataa agtgcaattt tagaagtaca tgggagggtga													850	
ttagcaattc tgaggatttt aaaaacacca taccatggt gacaccacct													900	
cctccacctg tcttctcatt gctgaagatc agtcaaagaa ttgtgtgctt													950	
agttcttgat aagtctggaa gcatgggggg taaggaccgc ctaaatcgaa													1000	
tgaatcaagc agcaaaacat ttcctgctgc agactgttga aaatggatcc													1050	
tgggtgggga tggttcactt tgatagtact gccactattg taaataagct													1100	
aatccaaata aaaagcagtg atgaaagaaa cacactcatg gcaggattac													1150	
ctacatatcc tctgggagga acttccatct gctctggaat taaatatgca													1200	
tttcagtgta ttggagagct acattcccaa ctcgatggat ccgaagtact													1250	
gctgctgact gatggggagg ataacactgc aagttcttgt attgatgaag													1300	
tgaaacaaag tggggccatt gttcatttta ttgctttggg aagagctgct													1350	
gatgaagcag taatagagat gagcaagata acaggaggaa gtcattttta													1400	

-continued

tgtttcagat gaagctcaga acaatggcct cattgatgct tttggggctc	1450
ttacatcagg aaatactgat ctctcccaga agtcccttca gtcgaaaagt	1500
aagggattaa cactgaatag taatgcctgg atgaacgaca ctgtcataat	1550
tgatagtaca gtgggaaagg acacgttctt tctcatcaca tggaacagtc	1600
tgctctccag tatttctctc tgggatccca gtggaacaat aatggaaaat	1650
ttcacagtgg atgcaacttc caaaatggcc tatctcagta ttccaggaac	1700
tgcaaagggtg ggcacttggg catacaatct tcaagccaaa gcgaaccag	1750
aaacattaac tattacagta acttctcgag cagcaaattc ttctgtgcct	1800
ccaatcacag tgaatgctaa aatgaataag gacgtaaaca gtttcccag	1850
cccaatgatt gtttacgcag aaattctaca aggatatgta cctgttcttg	1900
gagccaatgt gactgctttc attgaatcac agaatggaca tacagaagtt	1950
ttggaacttt tggataatgg tgcaggcgct gattctttca agaatgatgg	2000
agtctactcc aggtatttta cagcatatac agaaaatggc agatatagct	2050
taaaagtctg ggctcatgga ggagcaaaca ctgccaggct aaaattacgg	2100
cctccactga atagagccgc gtacatacca ggctgggtag tgaacgggga	2150
aattgaagca aaccgcgcaa gacctgaaat tgatgaggat actcagacca	2200
ccttgaggga tttcagccga acagcatccg gaggtgcatt tgtggtatca	2250
caagtcccaa gccttccctt gcctgaccaa taccaccaa gtcaaatcac	2300
agacottgat gccacagttc atgaggataa gattattctt acatggacag	2350
caccaggaga taattttgat gttggaaaag ttcaacgtta tatcataaga	2400
ataagtgcaa gtattcttga tctaagagac agttttgatg atgctcttca	2450
agtaaaact actgatctgt caccaaagga ggccaactcc aaggaaaagt	2500
ttgcatttaa accagaaaat atctcagaag aaaatgcaac ccacatattt	2550
attgccatta aaagtataga taaaagcaat ttgacatcaa aagtatccaa	2600
cattgcacaa gtaactttgt ttatccctca agcaaatcct gatgacattg	2650
atcctacacc tactcctact cctactccta ctctgataa aagtcataat	2700
tctggaggta atattttctac gctggtattg tctgtgattg ggtctgttgt	2750
aattgttaac tttattttaa gtaccacat ttgaacctta acgaagaaaa	2800
aaatcttcaa gtagacctag aagagagttt taaaaacaa aacaatgtaa	2850
gtaaaggata tttctgaatc ttaaaattca tcccatgtgt gatcataaac	2900
tcataaaaaat aattttaaga tgtcggaaaa ggatactttg attaaataaa	2950
aacactcatg gatagttaa aactgtcaag attaaaattt aatagtttca	3000
tttattttgt attttatttg taagaaatag tgatgaacaa agatcctttt	3050
tcatactgat acctggttgt atattatttg atgcaacagt tttctgaaat	3100
gatattttcaa attgcatcaa gaaattaaaa tcatctatct gagtagtcaa	3150
aatacaagta aagagagca aataaacaac atttgaaaa aaaaaaaaaa	3200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	3250
aaaaaaaaa aaaaa	3265

-continued

<210> SEQ ID NO 70

<211> LENGTH: 919

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 70

```

Met Gly Leu Phe Arg Gly Phe Val Phe Leu Leu Val Leu Cys Leu
 1          5          10          15

Leu His Gln Ser Asn Thr Ser Phe Ile Lys Leu Asn Asn Asn Gly
 20          25          30

Phe Glu Asp Ile Val Ile Val Ile Asp Pro Ser Val Pro Glu Asp
 35          40          45

Glu Lys Ile Ile Glu Gln Ile Glu Asp Met Val Thr Thr Ala Ser
 50          55          60

Thr Tyr Leu Phe Glu Ala Thr Glu Lys Arg Phe Phe Phe Lys Asn
 65          70          75

Val Ser Ile Leu Ile Pro Glu Asn Trp Lys Glu Asn Pro Gln Tyr
 80          85          90

Lys Arg Pro Lys His Glu Asn His Lys His Ala Asp Val Ile Val
 95          100         105

Ala Pro Pro Thr Leu Pro Gly Arg Asp Glu Pro Tyr Thr Lys Gln
110          115         120

Phe Thr Glu Cys Gly Glu Lys Gly Glu Tyr Ile His Phe Thr Pro
125          130         135

Asp Leu Leu Leu Gly Lys Lys Gln Asn Glu Tyr Gly Pro Pro Gly
140          145         150

Lys Leu Phe Val His Glu Trp Ala His Leu Arg Trp Gly Val Phe
155          160         165

Asp Glu Tyr Asn Glu Asp Gln Pro Phe Tyr Arg Ala Lys Ser Lys
170          175         180

Lys Ile Glu Ala Thr Arg Cys Ser Ala Gly Ile Ser Gly Arg Asn
185          190         195

Arg Val Tyr Lys Cys Gln Gly Gly Ser Cys Leu Ser Arg Ala Cys
200          205         210

Arg Ile Asp Ser Thr Thr Lys Leu Tyr Gly Lys Asp Cys Gln Phe
215          220         225

Phe Pro Asp Lys Val Gln Thr Glu Lys Ala Ser Ile Met Phe Met
230          235         240

Gln Ser Ile Asp Ser Val Val Glu Phe Cys Asn Glu Lys Thr His
245          250         255

Asn Gln Glu Ala Pro Ser Leu Gln Asn Ile Lys Cys Asn Phe Arg
260          265         270

Ser Thr Trp Glu Val Ile Ser Asn Ser Glu Asp Phe Lys Asn Thr
275          280         285

Ile Pro Met Val Thr Pro Pro Pro Pro Pro Val Phe Ser Leu Leu
290          295         300

Lys Ile Ser Gln Arg Ile Val Cys Leu Val Leu Asp Lys Ser Gly
305          310         315

Ser Met Gly Gly Lys Asp Arg Leu Asn Arg Met Asn Gln Ala Ala
320          325         330

Lys His Phe Leu Leu Gln Thr Val Glu Asn Gly Ser Trp Val Gly

```


-continued

335										340					345				
Met	Val	His	Phe	Asp	Ser	Thr	Ala	Thr	Ile	Val	Asn	Lys	Leu	Ile					
				350					355					360					
Gln	Ile	Lys	Ser	Ser	Asp	Glu	Arg	Asn	Thr	Leu	Met	Ala	Gly	Leu					
				365					370					375					
Pro	Thr	Tyr	Pro	Leu	Gly	Gly	Thr	Ser	Ile	Cys	Ser	Gly	Ile	Lys					
				380					385					390					
Tyr	Ala	Phe	Gln	Val	Ile	Gly	Glu	Leu	His	Ser	Gln	Leu	Asp	Gly					
				395					400					405					
Ser	Glu	Val	Leu	Leu	Leu	Thr	Asp	Gly	Glu	Asp	Asn	Thr	Ala	Ser					
				410					415					420					
Ser	Cys	Ile	Asp	Glu	Val	Lys	Gln	Ser	Gly	Ala	Ile	Val	His	Phe					
				425					430					435					
Ile	Ala	Leu	Gly	Arg	Ala	Ala	Asp	Glu	Ala	Val	Ile	Glu	Met	Ser					
				440					445					450					
Lys	Ile	Thr	Gly	Gly	Ser	His	Phe	Tyr	Val	Ser	Asp	Glu	Ala	Gln					
				455					460					465					
Asn	Asn	Gly	Leu	Ile	Asp	Ala	Phe	Gly	Ala	Leu	Thr	Ser	Gly	Asn					
				470					475					480					
Thr	Asp	Leu	Ser	Gln	Lys	Ser	Leu	Gln	Leu	Glu	Ser	Lys	Gly	Leu					
				485					490					495					
Thr	Leu	Asn	Ser	Asn	Ala	Trp	Met	Asn	Asp	Thr	Val	Ile	Ile	Asp					
				500					505					510					
Ser	Thr	Val	Gly	Lys	Asp	Thr	Phe	Phe	Leu	Ile	Thr	Trp	Asn	Ser					
				515					520					525					
Leu	Pro	Pro	Ser	Ile	Ser	Leu	Trp	Asp	Pro	Ser	Gly	Thr	Ile	Met					
				530					535					540					
Glu	Asn	Phe	Thr	Val	Asp	Ala	Thr	Ser	Lys	Met	Ala	Tyr	Leu	Ser					
				545					550					555					
Ile	Pro	Gly	Thr	Ala	Lys	Val	Gly	Thr	Trp	Ala	Tyr	Asn	Leu	Gln					
				560					565					570					
Ala	Lys	Ala	Asn	Pro	Glu	Thr	Leu	Thr	Ile	Thr	Val	Thr	Ser	Arg					
				575					580					585					
Ala	Ala	Asn	Ser	Ser	Val	Pro	Pro	Ile	Thr	Val	Asn	Ala	Lys	Met					
				590					595					600					
Asn	Lys	Asp	Val	Asn	Ser	Phe	Pro	Ser	Pro	Met	Ile	Val	Tyr	Ala					
				605					610					615					
Glu	Ile	Leu	Gln	Gly	Tyr	Val	Pro	Val	Leu	Gly	Ala	Asn	Val	Thr					
				620					625					630					
Ala	Phe	Ile	Glu	Ser	Gln	Asn	Gly	His	Thr	Glu	Val	Leu	Glu	Leu					
				635					640					645					
Leu	Asp	Asn	Gly	Ala	Gly	Ala	Asp	Ser	Phe	Lys	Asn	Asp	Gly	Val					
				650					655					660					
Tyr	Ser	Arg	Tyr	Phe	Thr	Ala	Tyr	Thr	Glu	Asn	Gly	Arg	Tyr	Ser					
				665					670					675					
Leu	Lys	Val	Arg	Ala	His	Gly	Gly	Ala	Asn	Thr	Ala	Arg	Leu	Lys					
				680					685					690					
Leu	Arg	Pro	Pro	Leu	Asn	Arg	Ala	Ala	Tyr	Ile	Pro	Gly	Trp	Val					
				695					700					705					
Val	Asn	Gly	Glu	Ile	Glu	Ala	Asn	Pro	Pro	Arg	Pro	Glu	Ile	Asp					
				710					715					720					

-continued

Glu Asp Thr Gln Thr Thr Leu Glu Asp Phe Ser Arg Thr Ala Ser
725 730 735
Gly Gly Ala Phe Val Val Ser Gln Val Pro Ser Leu Pro Leu Pro
740 745 750
Asp Gln Tyr Pro Pro Ser Gln Ile Thr Asp Leu Asp Ala Thr Val
755 760 765
His Glu Asp Lys Ile Ile Leu Thr Trp Thr Ala Pro Gly Asp Asn
770 775 780
Phe Asp Val Gly Lys Val Gln Arg Tyr Ile Ile Arg Ile Ser Ala
785 790 795
Ser Ile Leu Asp Leu Arg Asp Ser Phe Asp Asp Ala Leu Gln Val
800 805 810
Asn Thr Thr Asp Leu Ser Pro Lys Glu Ala Asn Ser Lys Glu Ser
815 820 825
Phe Ala Phe Lys Pro Glu Asn Ile Ser Glu Glu Asn Ala Thr His
830 835 840
Ile Phe Ile Ala Ile Lys Ser Ile Asp Lys Ser Asn Leu Thr Ser
845 850 855
Lys Val Ser Asn Ile Ala Gln Val Thr Leu Phe Ile Pro Gln Ala
860 865 870
Asn Pro Asp Asp Ile Asp Pro Thr Pro Thr Pro Thr Pro Thr Pro
875 880 885
Thr Pro Asp Lys Ser His Asn Ser Gly Val Asn Ile Ser Thr Leu
890 895 900
Val Leu Ser Val Ile Gly Ser Val Val Ile Val Asn Phe Ile Leu
905 910 915
Ser Thr Thr Ile

<210> SEQ ID NO 71
<211> LENGTH: 3877
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 71

ctccttaggt ggaaaccctg ggagtagagt actgacagca aagaccggga 50
aagaccatac gtccccgggc aggggtgaca acaggtgtca tctttttgat 100
ctcgtgtgtg gctgccttcc tatttcaagg aaagacgcca aggtaatttt 150
gacccagagg agcaatgatg tagccacctc ctaaccttcc cttcttgaac 200
ccccagttat gccaggattt actagagagt gtcaactcaa ccagcaagcg 250
gtccttcgg cttaacttgt ggttgaggga gagaaccttt gtggggctgc 300
gttctcttag cagtgtcag aagtgacttg cctgagggtg gaccagaaga 350
aaggaaaggt cccctcttgc tgttggtgc acatcaggaa ggctgtgatg 400
ggaatgaagg tgaaaacttg gagatttcac ttcagtcatt gcttctgcct 450
gcaagatcat cctttaaag tagagaagct gctctgtgtg gtggttaact 500
ccaagaggca gaactcgttc tagaaggaaa tggatgcaag cagctccggg 550
ggccccaac gcattgcttc tgtgtctag ccagggaag cccttccgtg 600
ggggccccgg ctttgaggga tgccaccggt tctggacgca tggctgattc 650

-continued

ctgaatgatg atgggttcgcc gggggctgct tgcgtggatt tcccgggtgg	700
tggttttgct ggtgctcctc tgctgtgcta tctctgtcct gtacatgttg	750
gcctgcaccc caaaagtgta cgaggagcag ctggcactgc ccagggccaa	800
cagccccacg gggaaggagg ggtaccaggc cgtccttcag gagtgggagg	850
agcagcaccg caactacgtg agcagcctga agcggcagat cgcacagctc	900
aaggaggagc tgcaggagag gagtgagcag ctcagggaatg ggcagtacca	950
agccagcgat gctgctggcc tgggtctgga caggagcccc ccagagaaaa	1000
cccaggccga cctcctggcc ttcctgcact cgcagggtga caaggcagag	1050
gtgaatgctg gcgtcaagct ggccacagag tatgcagcag tgcctttcga	1100
tagctttact ctacagaagg tgtaccagct ggagactggc cttaccgcc	1150
accccagga gaagcctgtg aggaaggaca agcgggatga gttggtgaa	1200
gccattgaat cagccttgga gaccctgaac aatcctgcag agaacagccc	1250
caatcacctg ccttacacgg cctctgattt catagaaggg atctaccgaa	1300
cagaaaagga caaagggaca ttgtatgagc tcaccttcaa aggggaccac	1350
aaacacgaat tcaaacggct catcttattt cgaccattca gccccatcat	1400
gaaagtgaat aatgaaaagc tcaacatggc caacacgctt atcaatgtta	1450
tcgtgcctct agcaaaaagg gtggacaagt tccggcagtt catgcagaat	1500
ttcagggaga tgtgcattga gcaggatggg agagtccatc tcaactgtgt	1550
ttactttggg aaagaagaaa taaatgaagt caaaggaata cttgaaaaa	1600
cttccaaagc tgccaacttc aggaacttta ccttcattca gctgaatgga	1650
gaattttctc ggggaaaggg acttgatgtt ggagcccgtc tctggaaggg	1700
aagcaacgct cttctctttt tctgtgatgt ggacatctac ttcacatctg	1750
aattcctcaa tacgtgtagg ctgaatacac agccaggga gaaggatttt	1800
tatccagttc ttttcagtca gtacaatcct ggcataatat acggccacca	1850
tgatgcagtc cctcccttg aacagcagct ggtcataaag aaggaaactg	1900
gattttggag agactttgga tttgggatga cgtgtcagta tcggtcagac	1950
ttcatcaata taggtggggt tgatctggac atcaaaggct ggggcggaga	2000
ggatgtgcac ctttatcgca agtatctcca cagcaacctc atagtggtag	2050
ggacgcctgt gcgaggactc ttccacctct ggcatgagaa gcgctgcatg	2100
gacgagctga cccccgagca gtacaagatg tgcatgcagt ccaaggccat	2150
gaacgagga tcccacggcc agctgggcat gctgggtgtc aggcacgaga	2200
tagaggctca ctttcgcaa cagaacaga agacaagtag caaaaaaaca	2250
tgaactccca gagaaggatt gtgggagaca ctttttcttt ctttttgcaa	2300
ttactgaaag tggctgcaac agagaaaaga cttccataaa ggacgacaaa	2350
agaattggac tgatgggtca gagatgagaa agcctccgat ttctctctgt	2400
tgggcttttt acaacagaaa tcaaaatctc cgctttgcct gcaaaagtaa	2450
cccagttgca ccctgtgaag tgtctgacaa aggcagaatg cttgtgagat	2500
tataagccta atggtgtgga ggttttgatg gtgtttacaa tacactgaga	2550

-continued

```

cctgttgttt tgtgtgctca ttgaaatatt catgatttaa gagcagtttt      2600
gtaaaaaatt cattagcatg aaaggcaagc atatttctcc tcatatgaat      2650
gagcctatca gcagggctct agtttctagg aatgctaaaa tatcagaag      2700
caggagagga gataggctta ttatgatact agtgagtaca ttaagtaaaa      2750
taaaatggac cagaaaagaa aagaaaccat aaatatcgtg tcatattttc      2800
cccaagatta accaaaaata atctgcttat ctttttggtt gtccttttaa      2850
ctgtctccgt ttttttcttt tatttaaaaa tgcacttttt ttcccttggt      2900
agttatagtc tgcttattta attaccactt tgcaagcctt acaagagagc      2950
acaagttggc ctacattttt atatttttta agaagatact ttgagatgca      3000
ttatgagaac tttcagttca aagcatcaaa ttgatgccat atccaaggac      3050
atgccaaatg ctgattctgt caggcactga atgtcaggca ttgagacata      3100
gggaaggaat ggtttgtact aatacagacg tacagatact ttctctgaag      3150
agtattttcg aagaggagca actgaacact ggaggaaaag aaaatgacac      3200
tttctgcttt acagaaaagg aaactcatc agactggtga tatcgtgatg      3250
tacctaaaag tcagaaacca cattttctcc tcagaagtag ggaccgcttt      3300
cttacctggt taaataaacc aaagtatacc gtgtgaacca aacaatctct      3350
tttcaaaaca ggggtgctcct cctggcttct ggcttcata agaagaaatg      3400
gagaaaaata tatatatata tatatatatt gtgaaagatc aatccatctg      3450
ccagaatcta gtgggatgga agtttttgct acatgttatc caccocaggc      3500
caggtggaag taactgaatt attttttaaa ttaagcagtt ctactcaatc      3550
accaagatgc ttctgaaaat tgcattttat taccatttca aactattttt      3600
taaaaaataa tacagttaac atagagtggg ttcttcattc atgtgaaaat      3650
tattagccag caccagatgc atgagctaat tatctctttg agtccttgct      3700
tctgtttgct cacagtaaac tcattgttta aaagcttcaa gaacattcaa      3750
gctgttggtg tgtaaaaaa tgcattgtat tgatttgtac tggtagttta      3800
tgaaatttaa ttaaaacaca ggccatgaat ggaagggtgg attgcacagc      3850
taataaaaata tgatttgtgg atatgaa      3877

```

<210> SEQ ID NO 72

<211> LENGTH: 532

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 72

```

Met Met Met Val Arg Arg Gly Leu Leu Ala Trp Ile Ser Arg Val
 1             5             10            15
Val Val Leu Leu Val Leu Leu Cys Cys Ala Ile Ser Val Leu Tyr
                20            25            30
Met Leu Ala Cys Thr Pro Lys Gly Asp Glu Glu Gln Leu Ala Leu
                35            40            45
Pro Arg Ala Asn Ser Pro Thr Gly Lys Glu Gly Tyr Gln Ala Val
                50            55            60
Leu Gln Glu Trp Glu Glu Gln His Arg Asn Tyr Val Ser Ser Leu
                65            70            75

```

-continued

Lys	Arg	Gln	Ile	Ala	Gln	Leu	Lys	Glu	Glu	Leu	Gln	Glu	Arg	Ser	80	85	90
Glu	Gln	Leu	Arg	Asn	Gly	Gln	Tyr	Gln	Ala	Ser	Asp	Ala	Ala	Gly	95	100	105
Leu	Gly	Leu	Asp	Arg	Ser	Pro	Pro	Glu	Lys	Thr	Gln	Ala	Asp	Leu	110	115	120
Leu	Ala	Phe	Leu	His	Ser	Gln	Val	Asp	Lys	Ala	Glu	Val	Asn	Ala	125	130	135
Gly	Val	Lys	Leu	Ala	Thr	Glu	Tyr	Ala	Ala	Val	Pro	Phe	Asp	Ser	140	145	150
Phe	Thr	Leu	Gln	Lys	Val	Tyr	Gln	Leu	Glu	Thr	Gly	Leu	Thr	Arg	155	160	165
His	Pro	Glu	Glu	Lys	Pro	Val	Arg	Lys	Asp	Lys	Arg	Asp	Glu	Leu	170	175	180
Val	Glu	Ala	Ile	Glu	Ser	Ala	Leu	Glu	Thr	Leu	Asn	Asn	Pro	Ala	185	190	195
Glu	Asn	Ser	Pro	Asn	His	Arg	Pro	Tyr	Thr	Ala	Ser	Asp	Phe	Ile	200	205	210
Glu	Gly	Ile	Tyr	Arg	Thr	Glu	Arg	Asp	Lys	Gly	Thr	Leu	Tyr	Glu	215	220	225
Leu	Thr	Phe	Lys	Gly	Asp	His	Lys	His	Glu	Phe	Lys	Arg	Leu	Ile	230	235	240
Leu	Phe	Arg	Pro	Phe	Ser	Pro	Ile	Met	Lys	Val	Lys	Asn	Glu	Lys	245	250	255
Leu	Asn	Met	Ala	Asn	Thr	Leu	Ile	Asn	Val	Ile	Val	Pro	Leu	Ala	260	265	270
Lys	Arg	Val	Asp	Lys	Phe	Arg	Gln	Phe	Met	Gln	Asn	Phe	Arg	Glu	275	280	285
Met	Cys	Ile	Glu	Gln	Asp	Gly	Arg	Val	His	Leu	Thr	Val	Val	Tyr	290	295	300
Phe	Gly	Lys	Glu	Glu	Ile	Asn	Glu	Val	Lys	Gly	Ile	Leu	Glu	Asn	305	310	315
Thr	Ser	Lys	Ala	Ala	Asn	Phe	Arg	Asn	Phe	Thr	Phe	Ile	Gln	Leu	320	325	330
Asn	Gly	Glu	Phe	Ser	Arg	Gly	Lys	Gly	Leu	Asp	Val	Gly	Ala	Arg	335	340	345
Phe	Trp	Lys	Gly	Ser	Asn	Val	Leu	Leu	Phe	Phe	Cys	Asp	Val	Asp	350	355	360
Ile	Tyr	Phe	Thr	Ser	Glu	Phe	Leu	Asn	Thr	Cys	Arg	Leu	Asn	Thr	365	370	375
Gln	Pro	Gly	Lys	Lys	Val	Phe	Tyr	Pro	Val	Leu	Phe	Ser	Gln	Tyr	380	385	390
Asn	Pro	Gly	Ile	Ile	Tyr	Gly	His	His	Asp	Ala	Val	Pro	Pro	Leu	395	400	405
Glu	Gln	Gln	Leu	Val	Ile	Lys	Lys	Glu	Thr	Gly	Phe	Trp	Arg	Asp	410	415	420
Phe	Gly	Phe	Gly	Met	Thr	Cys	Gln	Tyr	Arg	Ser	Asp	Phe	Ile	Asn	425	430	435
Ile	Gly	Gly	Phe	Asp	Leu	Asp	Ile	Lys	Gly	Trp	Gly	Gly	Glu	Asp	440	445	450

-continued

Val	His	Leu	Tyr	Arg	Lys	Tyr	Leu	His	Ser	Asn	Leu	Ile	Val	Val
				455					460					465
Arg	Thr	Pro	Val	Arg	Gly	Leu	Phe	His	Leu	Trp	His	Glu	Lys	Arg
				470					475					480
Cys	Met	Asp	Glu	Leu	Thr	Pro	Glu	Gln	Tyr	Lys	Met	Cys	Met	Gln
				485					490					495
Ser	Lys	Ala	Met	Asn	Glu	Ala	Ser	His	Gly	Gln	Leu	Gly	Met	Leu
				500					505					510
Val	Phe	Arg	His	Glu	Ile	Glu	Ala	His	Leu	Arg	Lys	Gln	Lys	Gln
				515					520					525
Lys	Thr	Ser	Ser	Lys	Lys	Thr								
				530										

<210> SEQ ID NO 73
<211> LENGTH: 1701
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien
<220> FEATURE:
<221> NAME/KEY: unsure
<222> LOCATION: 1528
<223> OTHER INFORMATION: unknown base

<400> SEQUENCE: 73	
gagactgcag agggagataa agagagaggg caaagaggca gcaagagatt	50
tgctctgggg atccagaaac ccatgatacc ctactgaaca ccgaatcccc	100
tggaagccca cagagacaga gacagcaaga gaagcagaga taaatacact	150
cacgccagga gctcgctcgc tctctctctc tctctctcac tcctccctcc	200
ctctctctct gcctgtccta gtctcttagt cctcaaattc ccagtccctc	250
gcaccccttc ctgggacact atgttggttct ccgccctcct gctggagggtg	300
atttggatcc tggctgcaga tgggggtcaa cactggacgt atgagggccc	350
acatggtcag gaccattggc cagcctctta ccctgagtgt ggaacaatg	400
cccagtcgcc catcgatatt cagacagaca gtgtgacatt tgaccctgat	450
ttgcctgctc tgcagcccca cggatatgac cagcctggca ccgagccttt	500
ggacctgcac aacaatggcc acacagtgca actctctctg ccctctaccc	550
tgtatctggg tggacttccc cgaaaatatg tagctgcca gctccacctg	600
cactggggtc agaaaggatc ccagggggg tcagaacacc agatcaacag	650
tgaagccaca tttgcagagc tccacattgt acattatgac tctgattcct	700
atgacagctt gagtgaggct gctgagaggc ctcagggcct ggctgtcctg	750
ggcatccctaa ttgaggtggg tgagactaag aatatagctt atgaacacat	800
tctgagtcat ttgcatgaag tcaggcataa agatcagaag acctcagtgc	850
ctcccttcaa cctaagagag ctgctcccca aacagctggg gcagtacttc	900
cgctacaatg gctcgctcac aactccccct tgctaccaga gtgtgctctg	950
gacagttttt tatagaaggt cccagatttc aatggaacag ctggaaaagc	1000
ttcaggggac attgttctcc acagaagagg agccctctaa gcttctggta	1050
cagaactacc gagcccttca gcctctcaat cagcgcatgg tctttgcttc	1100
tttcatccaa gcaggatcct cgtataccac aggtgaaatg ctgagtctag	1150

-continued

gtgtaggaat cttggttggc tgtctctgcc ttctcctggc tgtttatttc	1200
attgctagaa agattcggaa gaagaggctg gaaaaccgaa agagtgtggt	1250
cttcacctca gcacaagcca cgactgaggc ataaattcct tctcagatac	1300
catggatgtg gatgacttcc cttcatgcct atcaggaagc ctctaaaatg	1350
gggtgtagga tctggccaga aacactgtag gagtagtaag cagatgtcct	1400
ccttccctg gacatctctt agagaggaat ggaccaggc tgtcattcca	1450
ggaagaactg cagagccttc agcctctcca aacatgtagg aggaaatgag	1500
gaaatcgctg tgttggttaat gcagaganca aactctgttt agttgcaggg	1550
gaagtttggg atatacccca aagtcctcta cccctcact tttatggccc	1600
tttccctaga tatactgcgg gatctctcct taggataaag agttgtgttt	1650
gaagttgtat atttttgatc aatatatttg gaaattaaag tttctgactt	1700
t	1701

<210> SEQ ID NO 74

<211> LENGTH: 337

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 74

Met Leu Phe Ser Ala Leu Leu Leu Glu Val Ile Trp Ile Leu Ala	
1 5 10 15	
Ala Asp Gly Gly Gln His Trp Thr Tyr Glu Gly Pro His Gly Gln	
20 25 30	
Asp His Trp Pro Ala Ser Tyr Pro Glu Cys Gly Asn Asn Ala Gln	
35 40 45	
Ser Pro Ile Asp Ile Gln Thr Asp Ser Val Thr Phe Asp Pro Asp	
50 55 60	
Leu Pro Ala Leu Gln Pro His Gly Tyr Asp Gln Pro Gly Thr Glu	
65 70 75	
Pro Leu Asp Leu His Asn Asn Gly His Thr Val Gln Leu Ser Leu	
80 85 90	
Pro Ser Thr Leu Tyr Leu Gly Gly Leu Pro Arg Lys Tyr Val Ala	
95 100 105	
Ala Gln Leu His Leu His Trp Gly Gln Lys Gly Ser Pro Gly Gly	
110 115 120	
Ser Glu His Gln Ile Asn Ser Glu Ala Thr Phe Ala Glu Leu His	
125 130 135	
Ile Val His Tyr Asp Ser Asp Ser Tyr Asp Ser Leu Ser Glu Ala	
140 145 150	
Ala Glu Arg Pro Gln Gly Leu Ala Val Leu Gly Ile Leu Ile Glu	
155 160 165	
Val Gly Glu Thr Lys Asn Ile Ala Tyr Glu His Ile Leu Ser His	
170 175 180	
Leu His Glu Val Arg His Lys Asp Gln Lys Thr Ser Val Pro Pro	
185 190 195	
Phe Asn Leu Arg Glu Leu Leu Pro Lys Gln Leu Gly Gln Tyr Phe	
200 205 210	
Arg Tyr Asn Gly Ser Leu Thr Thr Pro Pro Cys Tyr Gln Ser Val	
215 220 225	

-continued

Leu	Trp	Thr	Val	Phe	Tyr	Arg	Arg	Ser	Gln	Ile	Ser	Met	Glu	Gln
			230						235					240
Leu	Glu	Lys	Leu	Gln	Gly	Thr	Leu	Phe	Ser	Thr	Glu	Glu	Glu	Pro
			245						250					255
Ser	Lys	Leu	Leu	Val	Gln	Asn	Tyr	Arg	Ala	Leu	Gln	Pro	Leu	Asn
			260						265					270
Gln	Arg	Met	Val	Phe	Ala	Ser	Phe	Ile	Gln	Ala	Gly	Ser	Ser	Tyr
			275						280					285
Thr	Thr	Gly	Glu	Met	Leu	Ser	Leu	Gly	Val	Gly	Ile	Leu	Val	Gly
			290						295					300
Cys	Leu	Cys	Leu	Leu	Leu	Ala	Val	Tyr	Phe	Ile	Ala	Arg	Lys	Ile
			305						310					315
Arg	Lys	Lys	Arg	Leu	Glu	Asn	Arg	Lys	Ser	Val	Val	Phe	Thr	Ser
			320						325					330
Ala	Gln	Ala	Thr	Thr	Glu	Ala								
			335											

<210> SEQ ID NO 75
<211> LENGTH: 1743
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 75

tgccgctgccc gccgctgctg ctgttgctcc tggcggcgccc ttggggacgg	50
gcagttccct gtgtctctgg tggtttgcc aaacctgcaa acatcacctt	100
cttatccatc aacatgaaga atgtcctaca atggactcca ccagagggtc	150
ttcaaggagt taaagttact tacactgtgc agtatttcat cacaaattgg	200
cccaccagag gtggcactga ctacagatga gaagtccatt tctgttgtcc	250
tgacagctcc agagaagtgg aagagaaatc cagaagacct tcctgtttcc	300
atgcaacaaa tatactcaa tctgaagtat aacgtgtctg tgtgaatac	350
taaatcaaac agaactgggt cccagtgtgt gaccaaccac acgctgggtc	400
tcacctggct ggagccgaac actctttact gcgtacacgt ggagtccttc	450
gtcccagggc cccctcgccg tgctcagcct tctgagaagc agtgtgccag	500
gactttgaaa gatcaatcat cagagttcaa ggctaaaatc atcttctggt	550
atgttttgcc catatctatt accgtgtttc ttttttctgt gatgggctat	600
tccatctacc gatatatcca cgttggcaca gagaaacacc cagcaaattt	650
gattttgatt tatggaaatg aatttgacaa aagattcttt gtgcctgctg	700
aaaaaatcgt gattaacttt atcacctca atatctcgga tgattctaaa	750
atttctcatc aggatatgag ttactggga aaaagcagtg atgtatccag	800
ccttaatgat cctcagccca gcgggaacct gagggccct caggaggaag	850
aggaggtgaa acatttaggg tatgcttcgc atttgatgga aattttttgt	900
gactctgaag aaaacacgga aggtacttct ctcaccacgc aagagtcctt	950
cagcagaaca ataccccgga ataaaacagt cattgaatat gaatatgat	1000
tcagaaccac tgacatttgt gcggggcctg aagagcagga gctcagtttg	1050
caggaggagg tgtccacaca aggaacatta ttggagtcgc aggcagcgtt	1100

-continued

```
ggcagtcttg ggcccgcaaa cgttacagta ctcatacacc cctcagctcc      1150
aagacttaga ccccttgccg caggagcaca cagactcgga ggaggggccg      1200
gaggaagagc catcgacgac cctggtcgac tgggatcccc aaactggcag      1250
gctgtgtatt ccttcgctgt ccagcttcga ccaggattca gagggctgcg      1300
agccttctga gggggatggg ctcggagagg agggctcttct atctagactc      1350
tatgaggagc cggctccaga caggccacca ggagaaaatg aaacctatct      1400
catgcaattc atggaggaat gggggttata tgtgcagatg gaaaactgat      1450
gccaacactt ccttttgccct tttgtttcct gtgcaaacia gtgagtcacc      1500
cctttgatcc cagccataaa gtacctggga tgaaagaagt tttttccagt      1550
ttgtcagtgt ctgtgagaat tacttatttc ttttctctat tctcatagca      1600
cgtgtgtgat tggttcatgc atgtaggctc cttacaatg atggtgggcc      1650
tctggagtcc aggggctggc cggttgttct atgcagagaa agcagtcaat      1700
aaatgtttgc cagactgggt gcagaattta ttcaggtggg tgt          1743
```

<210> SEQ ID NO 76

<211> LENGTH: 442

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 76

```
Met Ser Tyr Asn Gly Leu His Gln Arg Val Phe Lys Glu Leu Lys
 1              5              10             15
Leu Leu Thr Leu Cys Ser Ile Ser Ser Gln Ile Gly Pro Pro Glu
              20              25             30
Val Ala Leu Thr Thr Asp Glu Lys Ser Ile Ser Val Val Leu Thr
              35              40             45
Ala Pro Glu Lys Trp Lys Arg Asn Pro Glu Asp Leu Pro Val Ser
              50              55             60
Met Gln Gln Ile Tyr Ser Asn Leu Lys Tyr Asn Val Ser Val Leu
              65              70             75
Asn Thr Lys Ser Asn Arg Thr Trp Ser Gln Cys Val Thr Asn His
              80              85             90
Thr Leu Val Leu Thr Trp Leu Glu Pro Asn Thr Leu Tyr Cys Val
              95             100            105
His Val Glu Ser Phe Val Pro Gly Pro Pro Arg Arg Ala Gln Pro
             110             115            120
Ser Glu Lys Gln Cys Ala Arg Thr Leu Lys Asp Gln Ser Ser Glu
             125             130            135
Phe Lys Ala Lys Ile Ile Phe Trp Tyr Val Leu Pro Ile Ser Ile
             140             145            150
Thr Val Phe Leu Phe Ser Val Met Gly Tyr Ser Ile Tyr Arg Tyr
             155             160            165
Ile His Val Gly Lys Glu Lys His Pro Ala Asn Leu Ile Leu Ile
             170             175            180
Tyr Gly Asn Glu Phe Asp Lys Arg Phe Phe Val Pro Ala Glu Lys
             185             190            195
Ile Val Ile Asn Phe Ile Thr Leu Asn Ile Ser Asp Asp Ser Lys
             200             205            210
```

-continued

Ile Ser His Gln Asp Met Ser Leu Leu Gly Lys Ser Ser Asp Val
215 220 225

Ser Ser Leu Asn Asp Pro Gln Pro Ser Gly Asn Leu Arg Pro Pro
230 235 240

Gln Glu Glu Glu Glu Val Lys His Leu Gly Tyr Ala Ser His Leu
245 250 255

Met Glu Ile Phe Cys Asp Ser Glu Glu Asn Thr Glu Gly Thr Ser
260 265 270

Leu Thr Gln Gln Glu Ser Leu Ser Arg Thr Ile Pro Pro Asp Lys
275 280 285

Thr Val Ile Glu Tyr Glu Tyr Asp Val Arg Thr Thr Asp Ile Cys
290 295 300

Ala Gly Pro Glu Glu Gln Glu Leu Ser Leu Gln Glu Glu Val Ser
305 310 315

Thr Gln Gly Thr Leu Leu Glu Ser Gln Ala Ala Leu Ala Val Leu
320 325 330

Gly Pro Gln Thr Leu Gln Tyr Ser Tyr Thr Pro Gln Leu Gln Asp
335 340 345

Leu Asp Pro Leu Ala Gln Glu His Thr Asp Ser Glu Glu Gly Pro
350 355 360

Glu Glu Glu Pro Ser Thr Thr Leu Val Asp Trp Asp Pro Gln Thr
365 370 375

Gly Arg Leu Cys Ile Pro Ser Leu Ser Ser Phe Asp Gln Asp Ser
380 385 390

Glu Gly Cys Glu Pro Ser Glu Gly Asp Gly Leu Gly Glu Glu Gly
395 400 405

Leu Leu Ser Arg Leu Tyr Glu Glu Pro Ala Pro Asp Arg Pro Pro
410 415 420

Gly Glu Asn Glu Thr Tyr Leu Met Gln Phe Met Glu Glu Trp Gly
425 430 435

Leu Tyr Val Gln Met Glu Asn
440

<210> SEQ ID NO 77
<211> LENGTH: 1636
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 77

gaggagcggg ccgaggactc cagcgtgccc aggtctggca tcctgcactt 50

gctgccctct gacacctggg aagatggccg gcccgtagac cttcacccctt 100

ctctgtggtt tgctggcagc caccttgatc caagccaccc tcagtcccac 150

tgcagttctc atcctcggcc caaaagtcac caaagaaaag ctgacacagg 200

agctgaagga ccacaacgcc accagcatcc tgcagcagct gccgctgctc 250

agtgccatgc gggaaaagcc agccggaggc atccctgtgc tgggcagcct 300

ggtgaacacc gtccatgaagc acatcatctg gctgaaggtc atcacagcta 350

acatcctcca gctgcaggtg aagccctcgg ccaatgacca ggagctgcta 400

gtcaagatcc ccctggacat ggtggctgga ttcaaacacgc ccctggtcaa 450

gaccatcgtg gagtccaca tgacgactga ggcccaagcc accatccgca 500

-continued

```

tggacaccag tgcaagtggc cccacccgcc tggtcctcag tgactgtgcc      550
accagccatg ggagcctgcg catccaactg ctgtataagc tctccttcct      600
ggatgaacgcc ttagctaagc aggtcatgaa cctcctagtgc ccatccctgc      650
ccaatctagt gaaaaaccag ctgtgtcccg tgatcgaggc ttccttcaat      700
ggcatgtatg cagacctcct gcagctgggtg aaggtgcccc tttccctcag      750
cattgaccgt ctggagtttg accttctgta tcctgccatc aagggtgaca      800
ccattcagct ctacctgggg gccaatgtgt tggactcaca gggaaagggtg      850
accaagtggg tcaataactc tgcagcttcc ctgacaatgc ccaccctgga      900
caacatcccg ttcagcctca tcgtgagtca ggacgtgggtg aaagctgcag      950
tggctgtgtg gctctctcca gaagaattca tggtcctggt ggactctgtg     1000
cttcctgaga gtgcccatcg gctgaagtca agcatcgggc tgatcaatga     1050
aaaggctgca gataagctgg gatctacca gatcgtgaag atcctaactc     1100
aggacactcc cgagtttttt atagaccaag gccatgccaa ggtggcccaa     1150
ctgatcgtgc tggaagtgtt tcctccagtg gaagccctcc gcccttgtgt     1200
caccttgggc atcgaagcca gtcggaagc tcagttttac accaaagggtg     1250
accaacttat actcaacttg aataacatca gctctgatcg gatccagctg     1300
atgaactctg ggattggctg gttccaacct gatgttctga aaaacatcat     1350
cactgagatc atccactcca tcctgtctgc gaaccagaat ggcaaattaa     1400
gatctggggg cccagtgtca ttggtgaagg ccttgggatt cgaggcagct     1450
gagtcctcac tgaccaagga tgccttctgt cttactccag cctccttctg     1500
gaaaccagc tctcctgtct cccagtgaag acttggatgg cagccatcag     1550
ggaaggctgg gtcccagctg ggagtatggg tgtgagctct atagaccatc     1600
cctctctgca atcaataaac acttgcctgt gaaaaa                    1636

```

<210> SEQ ID NO 78

<211> LENGTH: 484

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 78

```

Met Ala Gly Pro Trp Thr Phe Thr Leu Leu Cys Gly Leu Leu Ala
 1             5             10            15
Ala Thr Leu Ile Gln Ala Thr Leu Ser Pro Thr Ala Val Leu Ile
          20            25            30
Leu Gly Pro Lys Val Ile Lys Glu Lys Leu Thr Gln Glu Leu Lys
          35            40            45
Asp His Asn Ala Thr Ser Ile Leu Gln Gln Leu Pro Leu Leu Ser
          50            55            60
Ala Met Arg Glu Lys Pro Ala Gly Gly Ile Pro Val Leu Gly Ser
          65            70            75
Leu Val Asn Thr Val Leu Lys His Ile Ile Trp Leu Lys Val Ile
          80            85            90
Thr Ala Asn Ile Leu Gln Leu Gln Val Lys Pro Ser Ala Asn Asp
          95           100           105

```

-continued

Gln	Glu	Leu	Leu	Val	Lys	Ile	Pro	Leu	Asp	Met	Val	Ala	Gly	Phe	110	115	120
Asn	Thr	Pro	Leu	Val	Lys	Thr	Ile	Val	Glu	Phe	His	Met	Thr	Thr	125	130	135
Glu	Ala	Gln	Ala	Thr	Ile	Arg	Met	Asp	Thr	Ser	Ala	Ser	Gly	Pro	140	145	150
Thr	Arg	Leu	Val	Leu	Ser	Asp	Cys	Ala	Thr	Ser	His	Gly	Ser	Leu	155	160	165
Arg	Ile	Gln	Leu	Leu	Tyr	Lys	Leu	Ser	Phe	Leu	Val	Asn	Ala	Leu	170	175	180
Ala	Lys	Gln	Val	Met	Asn	Leu	Leu	Val	Pro	Ser	Leu	Pro	Asn	Leu	185	190	195
Val	Lys	Asn	Gln	Leu	Cys	Pro	Val	Ile	Glu	Ala	Ser	Phe	Asn	Gly	200	205	210
Met	Tyr	Ala	Asp	Leu	Leu	Gln	Leu	Val	Lys	Val	Pro	Ile	Ser	Leu	215	220	225
Ser	Ile	Asp	Arg	Leu	Glu	Phe	Asp	Leu	Leu	Tyr	Pro	Ala	Ile	Lys	230	235	240
Gly	Asp	Thr	Ile	Gln	Leu	Tyr	Leu	Gly	Ala	Lys	Leu	Leu	Asp	Ser	245	250	255
Gln	Gly	Lys	Val	Thr	Lys	Trp	Phe	Asn	Asn	Ser	Ala	Ala	Ser	Leu	260	265	270
Thr	Met	Pro	Thr	Leu	Asp	Asn	Ile	Pro	Phe	Ser	Leu	Ile	Val	Ser	275	280	285
Gln	Asp	Val	Val	Lys	Ala	Ala	Val	Ala	Ala	Val	Leu	Ser	Pro	Glu	290	295	300
Glu	Phe	Met	Val	Leu	Leu	Asp	Ser	Val	Leu	Pro	Glu	Ser	Ala	His	305	310	315
Arg	Leu	Lys	Ser	Ser	Ile	Gly	Leu	Ile	Asn	Glu	Lys	Ala	Ala	Asp	320	325	330
Lys	Leu	Gly	Ser	Thr	Gln	Ile	Val	Lys	Ile	Leu	Thr	Gln	Asp	Thr	335	340	345
Pro	Glu	Phe	Phe	Ile	Asp	Gln	Gly	His	Ala	Lys	Val	Ala	Gln	Leu	350	355	360
Ile	Val	Leu	Glu	Val	Phe	Pro	Ser	Ser	Glu	Ala	Leu	Arg	Pro	Leu	365	370	375
Phe	Thr	Leu	Gly	Ile	Glu	Ala	Ser	Ser	Glu	Ala	Gln	Phe	Tyr	Thr	380	385	390
Lys	Gly	Asp	Gln	Leu	Ile	Leu	Asn	Leu	Asn	Asn	Ile	Ser	Ser	Asp	395	400	405
Arg	Ile	Gln	Leu	Met	Asn	Ser	Gly	Ile	Gly	Trp	Phe	Gln	Pro	Asp	410	415	420
Val	Leu	Lys	Asn	Ile	Ile	Thr	Glu	Ile	Ile	His	Ser	Ile	Leu	Leu	425	430	435
Pro	Asn	Gln	Asn	Gly	Lys	Leu	Arg	Ser	Gly	Val	Pro	Val	Ser	Leu	440	445	450
Val	Lys	Ala	Leu	Gly	Phe	Glu	Ala	Ala	Glu	Ser	Ser	Leu	Thr	Lys	455	460	465
Asp	Ala	Leu	Val	Leu	Thr	Pro	Ala	Ser	Leu	Trp	Lys	Pro	Ser	Ser	470	475	480
Pro	Val	Ser	Gln														

-continued

<210> SEQ ID NO 79
<211> LENGTH: 1475
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 79

gagagaagtc agcctggcag agagactctg aaatgaggga ttagaggtgt 50
tcaaggagca agagcttcag cctgaagaca agggagcagt ccctgaagac 100
gcttctactg agaggtctgc catggcctct cttggcctcc aacttgtggg 150
ctacatccta ggcttctctg ggcttttgga cactactggt gccatgctgc 200
tccccagctg gaaaacaagt tcttatgtcg gtgccagcat tgtgacagca 250
gttggtctct ccaagggcct ctggatggaa tgtgccacac acagcacagg 300
catcaccagc tgtgacatct atagaccctc tctgggcctg cccgtgaca 350
tccaggctgc ccaggccatg atggtgacat ccagtgaat ctctccctg 400
gcctgcatta tctctgtggt ggcatgaga tgcacagtct tctgccagga 450
atcccagacc aaagacagag tggcggtagc aggtggagtc ttttcatcc 500
ttggaggcct cctgggattc attcctgttg cctggaatct tcatgggatc 550
ctacgggact tctactcacc actggtgcct gacagcatga aatttgagat 600
tggagaggct ctttacttgg gcattatttc ttccctgttc tccctgatag 650
ctggaatcat cctctgcttt tctgtctcat cccagagaaa tcgctccaac 700
tactacgatg cctaccaagc ccaacctctt gccacaagga gctctccaag 750
gcctggtcaa cctcccaaag tcaagagtga gttcaattcc tacagcctga 800
cagggtatgt gtgaagaacc aggggccaga gctggggggt ggctgggtct 850
gtgaaaaaca gtggacagca ccccgagggc cacaggtgag ggacactacc 900
actggatcgt gtcagaaggt gctgtgagg atagactgac tttggccatt 950
ggattgagca aaggcagaaa tgggggctag tgtaacagca tgcaggttga 1000
attgccaaag atgctcgcca tgccagcctt tctgttttcc tcaccttget 1050
gtccccctgc cctaagtccc caacctcaa cttgaaaccc cattccctta 1100
agccaggact cagaggatcc ctttgccctc tggtttacct gggactccat 1150
ccccaaaccc actaatcaca tcccactgac tgaccctctg tgatcaaaga 1200
ccctctctct ggctgaggtt ggctcttagc tcattgctgg gcatgggaag 1250
gagaagcagt ggcttttgtg ggcattgctc taacctactt ctcaagcttc 1300
cctccaaaga aactgattgg ccctggaacc tccatccac tctgttatg 1350
actccacagt gtccagacta atttgtgcat gaactgaaat aaaaccatcc 1400
tacggtatcc aggaacaga aagcaggatg caggatggga ggacaggaag 1450
gcagcctggg acatttaaaa aaata 1475

<210> SEQ ID NO 80
<211> LENGTH: 230
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 80

-continued

Met Ala Ser Leu Gly Leu Gln Leu Val Gly Tyr Ile Leu Gly Leu
1 5 10 15
Leu Gly Leu Leu Gly Thr Leu Val Ala Met Leu Leu Pro Ser Trp
20 25 30
Lys Thr Ser Ser Tyr Val Gly Ala Ser Ile Val Thr Ala Val Gly
35 40 45
Phe Ser Lys Gly Leu Trp Met Glu Cys Ala Thr His Ser Thr Gly
50 55 60
Ile Thr Gln Cys Asp Ile Tyr Ser Thr Leu Leu Gly Leu Pro Ala
65 70 75
Asp Ile Gln Ala Ala Gln Ala Met Met Val Thr Ser Ser Ala Ile
80 85 90
Ser Ser Leu Ala Cys Ile Ile Ser Val Val Gly Met Arg Cys Thr
95 100 105
Val Phe Cys Gln Glu Ser Arg Ala Lys Asp Arg Val Ala Val Ala
110 115 120
Gly Gly Val Phe Phe Ile Leu Gly Gly Leu Leu Gly Phe Ile Pro
125 130 135
Val Ala Trp Asn Leu His Gly Ile Leu Arg Asp Phe Tyr Ser Pro
140 145 150
Leu Val Pro Asp Ser Met Lys Phe Glu Ile Gly Glu Ala Leu Tyr
155 160 165
Leu Gly Ile Ile Ser Ser Leu Phe Ser Leu Ile Ala Gly Ile Ile
170 175 180
Leu Cys Phe Ser Cys Ser Ser Gln Arg Asn Arg Ser Asn Tyr Tyr
185 190 195
Asp Ala Tyr Gln Ala Gln Pro Leu Ala Thr Arg Ser Ser Pro Arg
200 205 210
Pro Gly Gln Pro Pro Lys Val Lys Ser Glu Phe Asn Ser Tyr Ser
215 220 225
Leu Thr Gly Tyr Val
230

<210> SEQ ID NO 81
<211> LENGTH: 1732
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 81

cccacgcgtc cgcgccctctc ccttctgctg gaccttcctt cgtctctcca 50
tctctccctc ctttccccgc gttctctttc cacctttctc ttcttccac 100
cttagacctc ccttctctgcc ctcccttcct gccaccgct gcttctctggc 150
ccttctccga ccccgctcta gcagcagacc tcctggggtc tgtgggttga 200
tctgtggccc ctgtgcctcc gtgtcctttt cgtctccctt cctcccgact 250
ccgctcccg accagcgcc tgacctggg gaaaggatgg ttcccagggt 300
gagggctctc tcctccttgc tgggactcgc gctgctctgg ttccccctgg 350
actcccacgc tcgagccgc ccagacatgt tctgcctttt ccatgggaag 400
agatactccc ccggcgagag ctggcacccc tacttggagc cacaaggcct 450
gatgtactgc ctgcgctgta cctgctcaga gggcgcccat gtgagttgtt 500

-continued

```
accgcctcca ctgtccgcct gtccactgcc cccagcctgt gacggagcca      550
cagcaatgct gtcccaagtg tgtggaacct cacactccct ctggactccg      600
ggccccacca aagtcctgcc agcacaacgg gaccatgtac caacacggag      650
agatcttcag tgcccatgag ctgttccctc cccgcctgcc caaccagtgt      700
gtcctctgca gctgcacaga gggccagatc tactgcggcc tcacaacctg      750
ccccgaacca ggctgcccag caccctccc actgccagac tcctgctgcc      800
aagcctgcaa agatgaggca agtgagcaat cggatgaaga ggacagtgtg      850
cagtcgctcc atggggtgag acatcctcag gatccatgtt ccagtgatgc      900
tgaggagaaag agaggcccg gacccccag cccactggc ctcagcgccc      950
ctctgagctt catccctgc cacttcagac ccaagggagc aggcagcaca     1000
actgtcaaga tcgtcctgaa ggagaaacat aagaaagcct gtgtgcatgg     1050
cggaagacg tactcccacg gggagggtgtg gcacccggcc ttccgtgcct     1100
tcggccccct gccctgcac ctatgcacct gtgaggatgg ccgccaggac     1150
tgccagcgtg tgacctgtcc caccgagtac cctgcccgtc accccgagaa     1200
agtggctggg aagtgtctga agatttgccc agaggacaaa gcagaccctg     1250
gccacagtga gatcagttct accaggtgtc ccaaggcacc gggccgggtc     1300
ctcgtccaca catcggtatc cccaagccca gacaacctgc gtcgctttgc     1350
cctggaacac gaggcctcgg acttggtgga gatctacctc tggaagctgg     1400
taaaagatga ggaaactgag gctcagagag gtgaagtacc tggcccaagg     1450
ccacacagcc agaatcttcc acttgactca gatcaagaaa gtcaggaagc     1500
aagacttcca gaaagaggca cagcacttcc gactgctcgc tggcccccac     1550
gaaggtcact ggaacgtctt cctagcccag accctggagc tgaaggtcac     1600
ggccagtcca gaaaagtga ccaagacata acaaagacct aacagttgca     1650
gatatgagct gtataattgt tgttattata tattaataaa taagaagttg     1700
cattaccctc aaaaaaaaaa aaaaaaaaaa aa                          1732
```

<210> SEQ ID NO 82
<211> LENGTH: 451
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 82

```
Met Val Pro Glu Val Arg Val Leu Ser Ser Leu Leu Gly Leu Ala
 1             5             10            15
Leu Leu Trp Phe Pro Leu Asp Ser His Ala Arg Ala Arg Pro Asp
                20             25            30
Met Phe Cys Leu Phe His Gly Lys Arg Tyr Ser Pro Gly Glu Ser
                35             40            45
Trp His Pro Tyr Leu Glu Pro Gln Gly Leu Met Tyr Cys Leu Arg
                50             55            60
Cys Thr Cys Ser Glu Gly Ala His Val Ser Cys Tyr Arg Leu His
                65             70            75
Cys Pro Pro Val His Cys Pro Gln Pro Val Thr Glu Pro Gln Gln
                80             85            90
```

-continued

Cys	Cys	Pro	Lys	Cys	Val	Glu	Pro	His	Thr	Pro	Ser	Gly	Leu	Arg	95	100	105
Ala	Pro	Pro	Lys	Ser	Cys	Gln	His	Asn	Gly	Thr	Met	Tyr	Gln	His	110	115	120
Gly	Glu	Ile	Phe	Ser	Ala	His	Glu	Leu	Phe	Pro	Ser	Arg	Leu	Pro	125	130	135
Asn	Gln	Cys	Val	Leu	Cys	Ser	Cys	Thr	Glu	Gly	Gln	Ile	Tyr	Cys	140	145	150
Gly	Leu	Thr	Thr	Cys	Pro	Glu	Pro	Gly	Cys	Pro	Ala	Pro	Leu	Pro	155	160	165
Leu	Pro	Asp	Ser	Cys	Cys	Gln	Ala	Cys	Lys	Asp	Glu	Ala	Ser	Glu	170	175	180
Gln	Ser	Asp	Glu	Glu	Asp	Ser	Val	Gln	Ser	Leu	His	Gly	Val	Arg	185	190	195
His	Pro	Gln	Asp	Pro	Cys	Ser	Ser	Asp	Ala	Gly	Arg	Lys	Arg	Gly	200	205	210
Pro	Gly	Thr	Pro	Ala	Pro	Thr	Gly	Leu	Ser	Ala	Pro	Leu	Ser	Phe	215	220	225
Ile	Pro	Arg	His	Phe	Arg	Pro	Lys	Gly	Ala	Gly	Ser	Thr	Thr	Val	230	235	240
Lys	Ile	Val	Leu	Lys	Glu	Lys	His	Lys	Lys	Ala	Cys	Val	His	Gly	245	250	255
Gly	Lys	Thr	Tyr	Ser	His	Gly	Glu	Val	Trp	His	Pro	Ala	Phe	Arg	260	265	270
Ala	Phe	Gly	Pro	Leu	Pro	Cys	Ile	Leu	Cys	Thr	Cys	Glu	Asp	Gly	275	280	285
Arg	Gln	Asp	Cys	Gln	Arg	Val	Thr	Cys	Pro	Thr	Glu	Tyr	Pro	Cys	290	295	300
Arg	His	Pro	Glu	Lys	Val	Ala	Gly	Lys	Cys	Cys	Lys	Ile	Cys	Pro	305	310	315
Glu	Asp	Lys	Ala	Asp	Pro	Gly	His	Ser	Glu	Ile	Ser	Ser	Thr	Arg	320	325	330
Cys	Pro	Lys	Ala	Pro	Gly	Arg	Val	Leu	Val	His	Thr	Ser	Val	Ser	335	340	345
Pro	Ser	Pro	Asp	Asn	Leu	Arg	Arg	Phe	Ala	Leu	Glu	His	Glu	Ala	350	355	360
Ser	Asp	Leu	Val	Glu	Ile	Tyr	Leu	Trp	Lys	Leu	Val	Lys	Asp	Glu	365	370	375
Glu	Thr	Glu	Ala	Gln	Arg	Gly	Glu	Val	Pro	Gly	Pro	Arg	Pro	His	380	385	390
Ser	Gln	Asn	Leu	Pro	Leu	Asp	Ser	Asp	Gln	Glu	Ser	Gln	Glu	Ala	395	400	405
Arg	Leu	Pro	Glu	Arg	Gly	Thr	Ala	Leu	Pro	Thr	Ala	Arg	Trp	Pro	410	415	420
Pro	Arg	Arg	Ser	Leu	Glu	Arg	Leu	Pro	Ser	Pro	Asp	Pro	Gly	Ala	425	430	435
Glu	Gly	His	Gly	Gln	Ser	Arg	Gln	Ser	Asp	Gln	Asp	Ile	Thr	Lys	440	445	450

Thr

-continued

<210> SEQ ID NO 83

<211> LENGTH: 2052

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 83

gacagctgtg tctcgatgga gtagactctc agaacagcgc agtttgccct	50
ccgctcacgc agagcctctc cgtggcttcc gcaccttgag cattaggcca	100
gttctcctct tctctctaata ccatccgtca cctctcctgt catccgtttc	150
catgccgtga ggtccattca cagaacacat ccatggctct catgctcagt	200
ttggttctga gtctcctcaa gctgggatca gggcagtggc aggtgtttgg	250
gccagacaag cctgtccagg ccttggtggg ggaggacgca gcattctcct	300
gtttcctgtc tcctaagacc aatgcagagg ccatggaagt gcggttcttc	350
aggggccagt tctctagcgt ggtccacctc tacagggacg ggaaggacca	400
gccattttatg cagatgccac agtatcaagg caggacaaaa ctggtgaagg	450
attctattgc ggaggggcgc atctctctga ggctggaaaa cattactgtg	500
ttggatgctg gcctctatgg gtgcaggatt agttccagct cttactacca	550
gaaggccatc tgggagctac aggtgtcagc actgggctca gttcctctca	600
tttccatcac gggatatgtt gatagagaca tccagctact ctgtcagctc	650
tcgggctggt tccccggcc cacagcgaag tggaaaggtc cacaaggaca	700
ggatttgtcc acagactcca ggacaaacag agacatgcat ggcctgtttg	750
atgtggagat ctctctgacc gtccaagaga acgccgggag catatcctgt	800
tccatgcggc atgctcatct gagccgagag gtggaatcca ggtacagat	850
aggagatacc tttttcgagc ctatatcgtg gcacctggct accaaagtac	900
tgggaatact ctgctgtggc ctattttttg gcattgtttg actgaagatt	950
ttcttctcca aattccagtg gaaaatccag gcggaactgg actggagaag	1000
aaagcacgga caggcagaat tgagagacgc ccgaaacac gcagtggagg	1050
tgactctgga tccagagacg gctcaccgga agctctgcgt ttctgatctg	1100
aaaactgtaa cccatagaaa agctccccag gagtgccctc actctgagaa	1150
gagattttaca aggaagagtg tgggtgcttc tcagagtttc caagcaggga	1200
aacattactg ggaggtggac ggaggacaca ataaaagggt gcgcgtggga	1250
gtgtgccggg atgatgtgga caggaggaag gagtacgtga ctttgtctcc	1300
cgatcatggg tactgggtcc tcagactgaa tggagaacat ttgtatttca	1350
cattaaatcc ccgttttatc agcgtcttcc ccaggacccc acctacaaaa	1400
ataggggtct tcctggacta tgagtgtggg accatctcct tcttcaacat	1450
aatgaccag tcctttatct ataccctgac atgtcggttt gaaggcttat	1500
tgaggcccta cattgagtat ccgtcctata atgagcaaaa tggaactccc	1550
atagtcatct gccagtcac ccaggaatca gagaaagggt cctcttgcca	1600
aagggcctct gcaatcccag agacaagcaa cagtgagtcc tcctcacagg	1650
caaccacgcc cttcctcccc aggggtgaaa tgtaggatga atcacatccc	1700
acattcttct ttagggatat taaggtctct ctcccagatc caaagtcccg	1750

-continued

cagcagccgg ccaaggtggc ttccagatga agggggactg gcctgtccac 1800
atgggagatca ggtgtcatgg ctgccctgag ctgggagggga agaaggctga 1850
cattacattt agtttgctct cactccatct ggctaagtga tcttgaaata 1900
ccacctctca ggtgaagaac cgtcaggaat tcccatctca caggctgtgg 1950
tgtagattaa gtagacaagg aatgtgaata atgcttagat cttattgatg 2000
acagagtgtg tcctaattgt ttgttcatta tattacactt tcagtaaaaa 2050
aa 2052

<210> SEQ ID NO 84
<211> LENGTH: 500
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 84

Met Ala Leu Met Leu Ser Leu Val Leu Ser Leu Leu Lys Leu Gly
1 5 10 15
Ser Gly Gln Trp Gln Val Phe Gly Pro Asp Lys Pro Val Gln Ala
20 25 30
Leu Val Gly Glu Asp Ala Ala Phe Ser Cys Phe Leu Ser Pro Lys
35 40 45
Thr Asn Ala Glu Ala Met Glu Val Arg Phe Phe Arg Gly Gln Phe
50 55 60
Ser Ser Val Val His Leu Tyr Arg Asp Gly Lys Asp Gln Pro Phe
65 70 75
Met Gln Met Pro Gln Tyr Gln Gly Arg Thr Lys Leu Val Lys Asp
80 85 90
Ser Ile Ala Glu Gly Arg Ile Ser Leu Arg Leu Glu Asn Ile Thr
95 100 105
Val Leu Asp Ala Gly Leu Tyr Gly Cys Arg Ile Ser Ser Gln Ser
110 115 120
Tyr Tyr Gln Lys Ala Ile Trp Glu Leu Gln Val Ser Ala Leu Gly
125 130 135
Ser Val Pro Leu Ile Ser Ile Thr Gly Tyr Val Asp Arg Asp Ile
140 145 150
Gln Leu Leu Cys Gln Ser Ser Gly Trp Phe Pro Arg Pro Thr Ala
155 160 165
Lys Trp Lys Gly Pro Gln Gly Gln Asp Leu Ser Thr Asp Ser Arg
170 175 180
Thr Asn Arg Asp Met His Gly Leu Phe Asp Val Glu Ile Ser Leu
185 190 195
Thr Val Gln Glu Asn Ala Gly Ser Ile Ser Cys Ser Met Arg His
200 205 210
Ala His Leu Ser Arg Glu Val Glu Ser Arg Val Gln Ile Gly Asp
215 220 225
Thr Phe Phe Glu Pro Ile Ser Trp His Leu Ala Thr Lys Val Leu
230 235 240
Gly Ile Leu Cys Cys Gly Leu Phe Phe Gly Ile Val Gly Leu Lys
245 250 255
Ile Phe Phe Ser Lys Phe Gln Trp Lys Ile Gln Ala Glu Leu Asp
260 265 270

-continued

Trp Arg Arg Lys His Gly Gln Ala Glu Leu Arg Asp Ala Arg Lys
275 280 285

His Ala Val Glu Val Thr Leu Asp Pro Glu Thr Ala His Pro Lys
290 295 300

Leu Cys Val Ser Asp Leu Lys Thr Val Thr His Arg Lys Ala Pro
305 310 315

Gln Glu Val Pro His Ser Glu Lys Arg Phe Thr Arg Lys Ser Val
320 325 330

Val Ala Ser Gln Ser Phe Gln Ala Gly Lys His Tyr Trp Glu Val
335 340 345

Asp Gly Gly His Asn Lys Arg Trp Arg Val Gly Val Cys Arg Asp
350 355 360

Asp Val Asp Arg Arg Lys Glu Tyr Val Thr Leu Ser Pro Asp His
365 370 375

Gly Tyr Trp Val Leu Arg Leu Asn Gly Glu His Leu Tyr Phe Thr
380 385 390

Leu Asn Pro Arg Phe Ile Ser Val Phe Pro Arg Thr Pro Pro Thr
395 400 405

Lys Ile Gly Val Phe Leu Asp Tyr Glu Cys Gly Thr Ile Ser Phe
410 415 420

Phe Asn Ile Asn Asp Gln Ser Leu Ile Tyr Thr Leu Thr Cys Arg
425 430 435

Phe Glu Gly Leu Leu Arg Pro Tyr Ile Glu Tyr Pro Ser Tyr Asn
440 445 450

Glu Gln Asn Gly Thr Pro Ile Val Ile Cys Pro Val Thr Gln Glu
455 460 465

Ser Glu Lys Glu Ala Ser Trp Gln Arg Ala Ser Ala Ile Pro Glu
470 475 480

Thr Ser Asn Ser Glu Ser Ser Ser Gln Ala Thr Thr Pro Phe Leu
485 490 495

Pro Arg Gly Glu Met
500

<210> SEQ ID NO 85
<211> LENGTH: 1665
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 85

aacagacgtt ccctcgcggc cctggcacct ctaaccccag acatgctgct 50

gctgctgctg cccctgctct gggggaggga gagggcggaa ggacagacaa 100

gtaaactgct gacgatgcag agttccgtga cggtcgagga aggcctgtgt 150

gtccatgtgc cctgctcctt ctccctacccc tcgcatggct ggatttaccc 200

tggcccagta gttcatggct actggttccg ggaaggggcc aatacagacc 250

aggatgctcc agtggccaca aacaaccag ctcgggcagt gtgggaggag 300

actcgggacc gattccacct ccttggggac ccacatacca agaattgcac 350

cctgagcatc agagatgcc aagaagtga tgcggggaga tacttctttc 400

gtatggagaa aggaagtata aaatggaatt ataaacatca ccggctctct 450

gtgaatgtga cagccttgac ccacaggccc aacatcctca tcccaggcac 500

-continued

cctggagtc	ggctgcccc	agaatctgac	ctgctctgtg	ccctgggcct	550
gtgagcaggg	gacacccct	atgatctcct	ggataggac	ctccgtgtcc	600
ccccggacc	cctccaccac	ccgctcctcg	gtgctcacc	tcacccaca	650
gccccaggac	catggcacca	gcctcacctg	tcaggtgacc	ttccctggg	700
ccagcgtgac	cacgaacaag	accgtccatc	tcaacgtgtc	ctacccgcct	750
cagaacttga	ccatgactgt	cttccaagga	gacggcacag	tatccacagt	800
cttgggaaat	ggctcatctc	tgtcactccc	agagggccag	tctctgcgcc	850
tggctctgtc	agttgatgca	gttgacagca	atccccctgc	caggctgagc	900
ctgagcttga	gaggcctgac	cctgtgcccc	tcacagccct	caaaccggg	950
ggtgctggag	ctgccttggg	tgacactgag	ggatgcagct	gaattcacct	1000
gcagagctca	gaaccctctc	ggctctcagc	aggtctacct	gaacgtctcc	1050
ctgcagagca	aagccacatc	aggagtgact	cagggggtgg	tcgggggagc	1100
tggagccaca	gccctggtct	tcctgtcctt	ctgcgtcatc	ttcgtttag	1150
tgaggtcctg	caggaagaaa	tcggcaaggc	cagcagcggg	cgtgggagat	1200
acgggcatag	aggatgcaaa	cgtgtcagg	ggttcagcct	ctcaggggcc	1250
cctgactgaa	ccttgggcag	aagacagtcc	cccagaccag	cctccccag	1300
cttctgccc	ctcctcagtg	ggggaaggag	agctccagta	tgcatccctc	1350
agcttcacga	tggatgaagc	ttgggactcg	cggggacagg	aggccactga	1400
caccgagtac	tcggagatca	agatccacag	atgagaaact	gcagagactc	1450
accctgattg	agggatcaca	gcccctccag	gcaagggaga	agtcagaggc	1500
tgattcttgt	agaattaaca	gccctcaacg	tgatgagcta	tgataacact	1550
atgaattatg	tcagagtgta	aaagcacaca	ggcttttagag	tcaaagtatc	1600
tcaaacctga	atccacactg	tgccctccct	tttatTTTTT	taactaaaag	1650
acagacaaat	tccta				1665

<210> SEQ ID NO 86

<211> LENGTH: 463

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 86

Met	Leu	Leu	Leu	Leu	Leu	Pro	Leu	Leu	Trp	Gly	Arg	Glu	Arg	Ala
1				5						10				15
Glu	Gly	Gln	Thr	Ser	Lys	Leu	Leu	Thr	Met	Gln	Ser	Ser	Val	Thr
				20					25					30
Val	Gln	Glu	Gly	Leu	Cys	Val	His	Val	Pro	Cys	Ser	Phe	Ser	Tyr
				35					40					45
Pro	Ser	His	Gly	Trp	Ile	Tyr	Pro	Gly	Pro	Val	Val	His	Gly	Tyr
				50					55					60
Trp	Phe	Arg	Glu	Gly	Ala	Asn	Thr	Asp	Gln	Asp	Ala	Pro	Val	Ala
				65					70					75
Thr	Asn	Asn	Pro	Ala	Arg	Ala	Val	Trp	Glu	Glu	Thr	Arg	Asp	Arg
				80					85					90
Phe	His	Leu	Leu	Gly	Asp	Pro	His	Thr	Lys	Asn	Cys	Thr	Leu	Ser
				95					100					105

-continued

Ile Arg Asp Ala	Arg Arg Ser Asp Ala	Gly Arg Tyr Phe Phe Arg
110		115 120
Met Glu Lys Gly	Ser Ile Lys Trp Asn Tyr Lys His His Arg Leu	
125		130 135
Ser Val Asn Val	Thr Ala Leu Thr His Arg Pro Asn Ile Leu Ile	
140		145 150
Pro Gly Thr Leu	Glu Ser Gly Cys Pro Gln Asn Leu Thr Cys Ser	
155		160 165
Val Pro Trp Ala	Cys Glu Gln Gly Thr Pro Pro Met Ile Ser Trp	
170		175 180
Ile Gly Thr Ser	Val Ser Pro Leu Asp Pro Ser Thr Thr Arg Ser	
185		190 195
Ser Val Leu Thr	Leu Ile Pro Gln Pro Gln Asp His Gly Thr Ser	
200		205 210
Leu Thr Cys Gln	Val Thr Phe Pro Gly Ala Ser Val Thr Thr Asn	
215		220 225
Lys Thr Val His	Leu Asn Val Ser Tyr Pro Pro Gln Asn Leu Thr	
230		235 240
Met Thr Val Phe	Gln Gly Asp Gly Thr Val Ser Thr Val Leu Gly	
245		250 255
Asn Gly Ser Ser	Leu Ser Leu Pro Glu Gly Gln Ser Leu Arg Leu	
260		265 270
Val Cys Ala Val	Asp Ala Val Asp Ser Asn Pro Pro Ala Arg Leu	
275		280 285
Ser Leu Ser Trp	Arg Gly Leu Thr Leu Cys Pro Ser Gln Pro Ser	
290		295 300
Asn Pro Gly Val	Leu Glu Leu Pro Trp Val His Leu Arg Asp Ala	
305		310 315
Ala Glu Phe Thr	Cys Arg Ala Gln Asn Pro Leu Gly Ser Gln Gln	
320		325 330
Val Tyr Leu Asn	Val Ser Leu Gln Ser Lys Ala Thr Ser Gly Val	
335		340 345
Thr Gln Gly Val	Val Gly Gly Ala Gly Ala Thr Ala Leu Val Phe	
350		355 360
Leu Ser Phe Cys	Val Ile Phe Val Val Val Arg Ser Cys Arg Lys	
365		370 375
Lys Ser Ala Arg	Pro Ala Ala Gly Val Gly Asp Thr Gly Ile Glu	
380		385 390
Asp Ala Asn Ala	Val Arg Gly Ser Ala Ser Gln Gly Pro Leu Thr	
395		400 405
Glu Pro Trp Ala	Glu Asp Ser Pro Pro Asp Gln Pro Pro Pro Ala	
410		415 420
Ser Ala Arg Ser	Ser Val Gly Glu Gly Glu Leu Gln Tyr Ala Ser	
425		430 435
Leu Ser Phe Gln	Met Val Lys Pro Trp Asp Ser Arg Gly Gln Glu	
440		445 450
Ala Thr Asp Thr	Glu Tyr Ser Glu Ile Lys Ile His Arg	
455		460

<210> SEQ ID NO 87

<211> LENGTH: 1176

-continued

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 87

```
agaaagctgc actctgttga gctccagggc gcagtggagg gagggagtga      50
aggagctctc tgtaccaag gaaagtgcag ctgagactca gacaagatta      100
caatgaacca actcagcttc ctgctgtttc tcatagcgac caccagagga      150
tggagtagac atgaggctaa tacttacttc aaggaatgga cctgttcttc      200
gtctccatct ctgcccagaa gctgcaagga aatcaaagac gaatgtccta      250
gtgcatttga tggcctgtat ttctccgca ctgagaatgg tgttatctac      300
cagaccttct gtgacatgac ctctgggggt ggcggctgga ccctggtggc      350
cagcgtgcac gagaatgaca tgcgtgggaa gtgcacggtg ggcgatcgct      400
ggtccagcta gcagggcagc aaagcagact acccagaggg ggacggcaac      450
tgggccaact acaacacctt tggatctgca gaggcggcca cgagcgatga      500
ctacaagaac cctggctact acgacatcca ggccaaggac ctgggcatct      550
ggcacgtgcc caataagtcc cccatgcagc actggagaaa cagctccctg      600
ctgaggtacc gcacggacac tggcttcctc cagacactgg gacataatct      650
gtttggcatc taccagaaat atccagtga atattggagaa ggaaagtgtt      700
ggactgacaa cggcccgggtg atccctgtgg tctatgattt tggcgacgcc      750
cagaaaacag catcttatta ctacacctat ggccagcggg aattcactgc      800
gggatttgtt cagttcaggg tatttaataa cgagagagca gccaacgcct      850
tgtgtgctgg aatgagggtc accggatgta aactgagca tcaactgcatt      900
ggtggaggag gatactttcc agaggccagt cccagcagt gtggagattt      950
ttctggtttt gattggagtg gatatggaac tcatgttggg tacagcagca     1000
gccgtgagat aactgaggca gctgtgcttc tattctatcg ttgagagttt     1050
tgtgggaggg aaccagagacc tctcctccca accatgagat cccaaggatg     1100
gagaacaact taccagtag ctagaatgtt aatggcagaa gagaaaacaa     1150
taaatacatat tgactcaaga aaaaaa                                1176
```

<210> SEQ ID NO 88

<211> LENGTH: 313

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 88

```
Met Asn Gln Leu Ser Phe Leu Leu Phe Leu Ile Ala Thr Thr Arg
 1             5             10            15
Gly Trp Ser Thr Asp Glu Ala Asn Thr Tyr Phe Lys Glu Trp Thr
          20            25            30
Cys Ser Ser Ser Pro Ser Leu Pro Arg Ser Cys Lys Glu Ile Lys
          35            40            45
Asp Glu Cys Pro Ser Ala Phe Asp Gly Leu Tyr Phe Leu Arg Thr
          50            55            60
Glu Asn Gly Val Ile Tyr Gln Thr Phe Cys Asp Met Thr Ser Gly
          65            70            75
```

-continued

Gly	Gly	Gly	Trp	Thr	Leu	Val	Ala	Ser	Val	His	Glu	Asn	Asp	Met	
				80					85					90	
Arg	Gly	Lys	Cys	Thr	Val	Gly	Asp	Arg	Trp	Ser	Ser	Gln	Gln	Gly	
				95					100					105	
Ser	Lys	Ala	Asp	Tyr	Pro	Glu	Gly	Asp	Gly	Asn	Trp	Ala	Asn	Tyr	
				110					115					120	
Asn	Thr	Phe	Gly	Ser	Ala	Glu	Ala	Ala	Thr	Ser	Asp	Asp	Tyr	Lys	
				125					130					135	
Asn	Pro	Gly	Tyr	Tyr	Asp	Ile	Gln	Ala	Lys	Asp	Leu	Gly	Ile	Trp	
				140					145					150	
His	Val	Pro	Asn	Lys	Ser	Pro	Met	Gln	His	Trp	Arg	Asn	Ser	Ser	
				155					160					165	
Leu	Leu	Arg	Tyr	Arg	Thr	Asp	Thr	Gly	Phe	Leu	Gln	Thr	Leu	Gly	
				170					175					180	
His	Asn	Leu	Phe	Gly	Ile	Tyr	Gln	Lys	Tyr	Pro	Val	Lys	Tyr	Gly	
				185					190					195	
Glu	Gly	Lys	Cys	Trp	Thr	Asp	Asn	Gly	Pro	Val	Ile	Pro	Val	Val	
				200					205					210	
Tyr	Asp	Phe	Gly	Asp	Ala	Gln	Lys	Thr	Ala	Ser	Tyr	Tyr	Ser	Pro	
				215					220					225	
Tyr	Gly	Gln	Arg	Glu	Phe	Thr	Ala	Gly	Phe	Val	Gln	Phe	Arg	Val	
				230					235					240	
Phe	Asn	Asn	Glu	Arg	Ala	Ala	Asn	Ala	Leu	Cys	Ala	Gly	Met	Arg	
				245					250					255	
Val	Thr	Gly	Cys	Asn	Thr	Glu	His	His	Cys	Ile	Gly	Gly	Gly	Gly	
				260					265					270	
Tyr	Phe	Pro	Glu	Ala	Ser	Pro	Gln	Gln	Cys	Gly	Asp	Phe	Ser	Gly	
				275					280					285	
Phe	Asp	Trp	Ser	Gly	Tyr	Gly	Thr	His	Val	Gly	Tyr	Ser	Ser	Ser	
				290					295					300	
Arg	Glu	Ile	Thr	Glu	Ala	Ala	Val	Leu	Leu	Phe	Tyr	Arg			
				305					310						

<210> SEQ ID NO 89
<211> LENGTH: 759
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 89

ctagatttgt	cggtctgcgg	ggagacttca	ggagtcgctg	tctctgaact	50
tccagcctca	gagaccgccg	cccttgcccc	cgagggccat	gggccgggtc	100
tcagggcttg	tgccctctcg	cttcctgacg	ctcctggcgc	atctggtggt	150
cgtcatcacc	ttattctggt	cccgggacag	caacatacag	gcctgcctgc	200
ctctcacggt	cacccccgag	gagtatgaca	agcaggacat	tcagctggtg	250
gccgcgctct	ctgtcaccct	gggcctcttt	gcagtggagc	tggccggttt	300
cctctcagga	gtctccatgt	tcaacagcac	ccagagcctc	atctccattg	350
gggctcactg	tagtgcatcc	gtggccctgt	ccttcttcat	attcgagcgt	400
tgggagtgca	ctacgtattg	gtacattttt	gtcttctgca	gtgcccttcc	450
agctgtcact	gaaatggctt	tattcgtcac	cgtctttggg	ctgaaaaaga	500

-continued

aacccttctg attaccttca tgacgggaac ctaaggacga agcctacagg	550
ggcaagggcc gcttcgtatt cctggaagaa ggaaggcata ggcttcggtt	600
ttcccctcgg aaactgcttc tgctggagga tatgtgttg aataattacg	650
tcttgagtct gggattatcc gcattgtatt tagtgctttg taataaaata	700
tgttttgtag taacattaag acttatatac agttttaggg gacaattaa	750
aaaaaaaa	759

<210> SEQ ID NO 90
<211> LENGTH: 140
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 90

Met Gly Arg Val Ser Gly Leu Val Pro Ser Arg Phe Leu Thr Leu	1	5	10	15
Leu Ala His Leu Val Val Val Ile Thr Leu Phe Trp Ser Arg Asp	20	25	30	
Ser Asn Ile Gln Ala Cys Leu Pro Leu Thr Phe Thr Pro Glu Glu	35	40	45	
Tyr Asp Lys Gln Asp Ile Gln Leu Val Ala Ala Leu Ser Val Thr	50	55	60	
Leu Gly Leu Phe Ala Val Glu Leu Ala Gly Phe Leu Ser Gly Val	65	70	75	
Ser Met Phe Asn Ser Thr Gln Ser Leu Ile Ser Ile Gly Ala His	80	85	90	
Cys Ser Ala Ser Val Ala Leu Ser Phe Phe Ile Phe Glu Arg Trp	95	100	105	
Glu Cys Thr Thr Tyr Trp Tyr Ile Phe Val Phe Cys Ser Ala Leu	110	115	120	
Pro Ala Val Thr Glu Met Ala Leu Phe Val Thr Val Phe Gly Leu	125	130	135	
Lys Lys Lys Pro Phe	140			

<210> SEQ ID NO 91
<211> LENGTH: 1871
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 91

ctgggacccc gaaaagagaa ggggagagcg aggggacgag agcggaggag	50
gaagatgcaa ctgactcgct gctgcttcgt gttcctgggt cagggtagcc	100
tctatctgggt catctgtggc caggatgatg gtcctcccg ctcagaggac	150
cctgagcgtg atgaccacga gggccagccc cgccccggg tgcctcggaa	200
gcggggccac atctcaccta agtcccggcc catggccaat tccactctcc	250
tagggctgct ggccccgcct ggggaggctt ggggcattct tgggcagccc	300
cccaaccgcc cgaaccacag cccccacccc tcagccaagg tgaagaaaat	350
ctttggctgg ggcgacttct actccaacat caagacggtg gccctgaacc	400
tgctcgtcac agggaagatt gtggaccatg gcaatgggac cttcagcgtc	450

-continued

cacttccaac acaatgccac aggccaggga aacatctcca tcagcctcgt	500
gccccccagt aaagctgtag agttccacca ggaacagcag atcttcatcg	550
aagccaaggc ctccaaaatc ttcaactgcc ggatggagtg ggagaaggta	600
gaacggggcc gccggacctc gctttgcacc cacgacccag ccaagatctg	650
ctcccagagac cacgctcaga gctcagccac ctggagctgc tcccagccct	700
tcaaagtctg ctgtgtctac atcgctttct acagcacgga ctatcggctg	750
gtccagaagg tgtgcccaga ttacaactac catagtgata ccccctacta	800
cccctctggg tgacccgggg caggccacag aggccaggcc agggctggaa	850
ggacaggcct gcccatgcag gagaccatct ggacaccggg caggggaaggg	900
gttgggcctc aggcagggag gggggtggag acgaggagat gccaagtggg	950
gccaggggcca agtctcaagt ggcagagaaa gggtoccaag tgcgtgtccc	1000
aacctgaagc tgtggagtga ctagatcaca ggagcactgg aggaggagtg	1050
ggctctctgt gcagcctcac agggctttgc cagggagcca cagagagatg	1100
ctgggtcccc gaggcctgtg ggcaggccga tcagtgtggc cccagatcaa	1150
gtcatgggag gaagctaagc ccttggttct tgccatcctg aggaaagata	1200
gcaacaggga gggggagatt tcatcagtgt ggacagcctg tcaacttagg	1250
atggatggct gagagggctt cctaggagcc agtcagcagg gtggggtggg	1300
gccagaggag ctctccagcc ctgcctagtg ggcgccctga gcccttgtc	1350
gtgtgctgag catggcatga ggctgaagtg gcaaccctgg ggtctttgat	1400
gtcttgacag attgaccatc tgtctccagc caggccaccc ctttccaaaa	1450
ttccctcttc tgccagtact ccccctgtac caccatttgc tgatggcaca	1500
cccctcctta agctaagaca ggacgattgt ggtcctccca cactaaggcc	1550
acagcccac cgcgtgctgt gtgtccctct tccaccccaa cccctgctgg	1600
ctcctctggg agcatccatg tcccggagag gggtcctca acagtcagcc	1650
tcacctgtca gaccgggggt ctcccggatc tggatggcgc cgcctctca	1700
gcagcgggca cgggtggggc ggggcccggc cgcagagcat gtgctggatc	1750
tgttctgtgt gtctgtctgt ggggtggggg aggggaggga agtcttgtga	1800
aaccgctgat tgctgacttt tgtgtgaaga atcgtgttct tggagcagga	1850
aataaagctt gccccggggc a	1871

<210> SEQ ID NO 92
<211> LENGTH: 252
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 92

Met	Gln	Leu	Thr	Arg	Cys	Cys	Phe	Val	Phe	Leu	Val	Gln	Gly	Ser
1				5					10					15
Leu	Tyr	Leu	Val	Ile	Cys	Gly	Gln	Asp	Asp	Gly	Pro	Pro	Gly	Ser
				20					25					30
Glu	Asp	Pro	Glu	Arg	Asp	Asp	His	Glu	Gly	Gln	Pro	Arg	Pro	Arg
				35					40					45
Val	Pro	Arg	Lys	Arg	Gly	His	Ile	Ser	Pro	Lys	Ser	Arg	Pro	Met

-continued

50										55										60									
Ala	Asn	Ser	Thr	Leu	Leu	Gly	Leu	Leu	Ala	Pro	Pro	Gly	Glu	Ala															
				65						70																			
Trp	Gly	Ile	Leu	Gly	Gln	Pro	Pro	Asn	Arg	Pro	Asn	His	Ser	Pro															
				80					85																				
Pro	Pro	Ser	Ala	Lys	Val	Lys	Lys	Ile	Phe	Gly	Trp	Gly	Asp	Phe															
				95					100																				
Tyr	Ser	Asn	Ile	Lys	Thr	Val	Ala	Leu	Asn	Leu	Leu	Val	Thr	Gly															
				110					115																				
Lys	Ile	Val	Asp	His	Gly	Asn	Gly	Thr	Phe	Ser	Val	His	Phe	Gln															
				125					130																				
His	Asn	Ala	Thr	Gly	Gln	Gly	Asn	Ile	Ser	Ile	Ser	Leu	Val	Pro															
				140					145																				
Pro	Ser	Lys	Ala	Val	Glu	Phe	His	Gln	Glu	Gln	Gln	Ile	Phe	Ile															
				155					160																				
Glu	Ala	Lys	Ala	Ser	Lys	Ile	Phe	Asn	Cys	Arg	Met	Glu	Trp	Glu															
				170					175																				
Lys	Val	Glu	Arg	Gly	Arg	Arg	Thr	Ser	Leu	Cys	Thr	His	Asp	Pro															
				185					190																				
Ala	Lys	Ile	Cys	Ser	Arg	Asp	His	Ala	Gln	Ser	Ser	Ala	Thr	Trp															
				200					205																				
Ser	Cys	Ser	Gln	Pro	Phe	Lys	Val	Val	Cys	Val	Tyr	Ile	Ala	Phe															
				215					220																				
Tyr	Ser	Thr	Asp	Tyr	Arg	Leu	Val	Gln	Lys	Val	Cys	Pro	Asp	Tyr															
				230					235																				
Asn	Tyr	His	Ser	Asp	Thr	Pro	Tyr	Tyr	Pro	Ser	Gly																		
				245					250																				

<210> SEQ ID NO 93
<211> LENGTH: 902
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 93

cggtggccat gactgcggcc gtgttcttcg gctgcgcctt cattgccttc	50
gggcctgcgc tcgcccttta tgtcttcacc atcgccatcg agcogttgcg	100
tatcatcttc ctcatcgccg gagctttctt ctggttggtg tctctactga	150
tttcgtccct tgtttggttc atggcaagag tcattattga caacaaagat	200
ggaccaacac agaaatatct gctgatcttt ggagcgtttg tctctgtcta	250
tatccaagaa atgttccgat ttgcatatta taaactctta aaaaaagcca	300
gtgaaggttt gaagagtata aaccaggtg agacagcacc ctctatgcga	350
ctgtgcgcct atgtttcttg cttgggcttt ggaatcatga gtggagtatt	400
ttcctttgtg aataccctat ctgactcctt ggggccaggc acagtgggca	450
ttcatggaga ttctcctcaa ttcttccttt attcagcttt catgacgtg	500
gtcattatct tgctgcatgt attctggggc attgtatttt ttgatggctg	550
tgagaagaaa aagtggggca tcctccttat cgttctcctg acccacctgc	600
tggtgtcagc ccagaccttc ataagttctt attatggaat aaacctggcg	650
tcagcattta taatcctggt gctcatgggc acctgggcat tcttagctgc	700

-continued

gggaggcagc tgccgaagcc tgaaactctg cctgctctgc caagacaaga 750
actttcttct ttacaaccag cgctccagat aacctcaggg aaccagcact 800
tcccaaaccg cagactacat ctttagagga agcacaactg tgcctttttc 850
tgaaaatccc tttttctggt ggaattgaga aagaaataaa actatgcaga 900
ta 902

<210> SEQ ID NO 94
<211> LENGTH: 257
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 94

Met Thr Ala Ala Val Phe Phe Gly Cys Ala Phe Ile Ala Phe Gly
1 5 10 15
Pro Ala Leu Ala Leu Tyr Val Phe Thr Ile Ala Ile Glu Pro Leu
20 25 30
Arg Ile Ile Phe Leu Ile Ala Gly Ala Phe Phe Trp Leu Val Ser
35 40 45
Leu Leu Ile Ser Ser Leu Val Trp Phe Met Ala Arg Val Ile Ile
50 55 60
Asp Asn Lys Asp Gly Pro Thr Gln Lys Tyr Leu Leu Ile Phe Gly
65 70 75
Ala Phe Val Ser Val Tyr Ile Gln Glu Met Phe Arg Phe Ala Tyr
80 85 90
Tyr Lys Leu Leu Lys Lys Ala Ser Glu Gly Leu Lys Ser Ile Asn
95 100 105
Pro Gly Glu Thr Ala Pro Ser Met Arg Leu Leu Ala Tyr Val Ser
110 115 120
Gly Leu Gly Phe Gly Ile Met Ser Gly Val Phe Ser Phe Val Asn
125 130 135
Thr Leu Ser Asp Ser Leu Gly Pro Gly Thr Val Gly Ile His Gly
140 145 150
Asp Ser Pro Gln Phe Phe Leu Tyr Ser Ala Phe Met Thr Leu Val
155 160 165
Ile Ile Leu Leu His Val Phe Trp Gly Ile Val Phe Phe Asp Gly
170 175 180
Cys Glu Lys Lys Lys Trp Gly Ile Leu Leu Ile Val Leu Leu Thr
185 190 195
His Leu Leu Val Ser Ala Gln Thr Phe Ile Ser Ser Tyr Tyr Gly
200 205 210

Ile Asn Leu Ala Ser Ala Phe Ile Ile Leu Val Leu Met Gly Thr
215 220 225
Trp Ala Phe Leu Ala Ala Gly Gly Ser Cys Arg Ser Leu Lys Leu
230 235 240
Cys Leu Leu Cys Gln Asp Lys Asn Phe Leu Leu Tyr Asn Gln Arg
245 250 255

Ser Arg

<210> SEQ ID NO 95

-continued

<211> LENGTH: 1073

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 95

aatttttcac cagagtaaac ttgagaaacc aactggacct tgagtattgt	50
acattttgcc tcgtggaccc aaaggtagca atctgaaaca tgaggagtac	100
gattctactg ttttgtcttc taggatcaac tcggtcatta ccacagctca	150
aacctgcttt gggactccct cccacaaaac tggctccgga tcagggaaca	200
ctaccaaacc aacagcagtc aaatcaggtc tttccttctt taagtctgat	250
accattaaca cagatgctca cactggggcc agatctgcat ctgttaaact	300
ctgtgcagag aatgacacct ggtaccaga cccaccatt gaccctggga	350
gggttgaatg tacaacagca actgcacca catgtgttac caatttttgt	400
cacacaactt ggagcccagg gcactatcct aagctcagag gaattgccac	450
aaatcttcac gagcctcatc atccattcct tgttcccggg aggcactctg	500
cccaccagtc aggcaggggc taatccagat gtccaggatg gaagccttcc	550
agcaggagga gcaggtgtaa atcctgccac ccagggaacc ccagcaggcc	600
gcctcccaac tcccagtggc acagatgacg actttgcagt gaccaccct	650
gcaggcatcc aaaggagcac acatgccatc gaggaagcca ccacagaatc	700
agcaaatgga attcagtaag ctgtttcaaa tttttcaac taagctgcct	750
cgaatttggg gatacatgtg aatctttatc attgattata ttatggaata	800
gattgagaca cattggatag tcttagaaga aattaattct taatttacct	850
gaaaatatcc ttgaaatttc agaaaatatg ttctatgtag agaatcccaa	900
cttttaaaaa caataattca atggataaat ctgtctttga aatataacat	950
tatgctgcct ggatgatatg catattaaaa catatttggg aaactggaaa	1000
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	1050
aaaaaaaaa aaaaaaaaaa aaa	1073

<210> SEQ ID NO 96

<211> LENGTH: 209

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 96

Met Arg Ser Thr Ile Leu Leu Phe Cys Leu Leu Gly Ser Thr Arg	
1 5 10 15	
Ser Leu Pro Gln Leu Lys Pro Ala Leu Gly Leu Pro Pro Thr Lys	
20 25 30	
Leu Ala Pro Asp Gln Gly Thr Leu Pro Asn Gln Gln Ser Asn	
35 40 45	
Gln Val Phe Pro Ser Leu Ser Leu Ile Pro Leu Thr Gln Met Leu	
50 55 60	
Thr Leu Gly Pro Asp Leu His Leu Leu Asn Pro Ala Ala Gly Met	
65 70 75	
Thr Pro Gly Thr Gln Thr His Pro Leu Thr Leu Gly Gly Leu Asn	
80 85 90	

-continued

Val	Gln	Gln	Gln	Leu	His	Pro	His	Val	Leu	Pro	Ile	Phe	Val	Thr
				95					100					105
Gln	Leu	Gly	Ala	Gln	Gly	Thr	Ile	Leu	Ser	Ser	Glu	Glu	Leu	Pro
				110					115					120
Gln	Ile	Phe	Thr	Ser	Leu	Ile	Ile	His	Ser	Leu	Phe	Pro	Gly	Gly
				125					130					135
Ile	Leu	Pro	Thr	Ser	Gln	Ala	Gly	Ala	Asn	Pro	Asp	Val	Gln	Asp
				140					145					150
Gly	Ser	Leu	Pro	Ala	Gly	Gly	Ala	Gly	Val	Asn	Pro	Ala	Thr	Gln
				155					160					165
Gly	Thr	Pro	Ala	Gly	Arg	Leu	Pro	Thr	Pro	Ser	Gly	Thr	Asp	Asp
				170					175					180
Asp	Phe	Ala	Val	Thr	Thr	Pro	Ala	Gly	Ile	Gln	Arg	Ser	Thr	His
				185					190					195
Ala	Ile	Glu	Glu	Ala	Thr	Thr	Glu	Ser	Ala	Asn	Gly	Ile	Gln	
				200					205					

<210> SEQ ID NO 97
<211> LENGTH: 2848
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 97

gctcaagtgc	cctgccttgc	cccaccacgc	ccagcctggc	cagagccccc	50
tggagaagga	gctctcttct	tgcttggcag	ctggaccaag	ggagccagtc	100
ttgggcgctg	gagggcctgt	cctgaccatg	gtccctgcct	ggctgtggct	150
gctttgtgtc	tccgtccccc	aggctctccc	caaggcccag	cctgcagagc	200
tgtctgtgga	agttccagaa	aactatgggt	gaaatttccc	tttatacctg	250
accaagttgc	cgctgccccg	tgagggggct	gaaggccaga	tcgtgctgtc	300
aggggactca	ggcaaggcaa	ctgagggccc	atttgctatg	gatccagatt	350
ctggcttcct	gctggtgacc	agggccctgg	accgagagga	gcaggcagag	400
taccagctac	aggtcaccct	ggagatgcag	gatggacatg	tcttgtgggg	450
tccacagcct	gtgcttgtgc	acgtgaagga	tgagaatgac	caggtgcccc	500
atttctctca	agccatctac	agagctcggc	tgagccgggg	taccaggcct	550
ggcatccccc	tcctcttcct	tgaggcttca	gaccgggatg	agccaggcac	600
agccaactcg	gatcttcgat	tccacatcct	gagccaggct	ccagcccagc	650
cttccccaga	catgttccag	ctggagcctc	ggctgggggc	tctggccctc	700
agccccaag	ggagcaccag	ccttgaccac	gccctggaga	ggacctacca	750
gctgttggtg	caggtcaagg	acatgggtga	ccaggcctca	ggccaccagg	800
ccactgccac	cgtggaagtc	tccatcatag	agagcacctg	ggtgtcccta	850
gagcctatcc	acctggcaga	gaatctcaaa	gtcctatacc	cgcaccacat	900
ggcccaggta	cactggagtg	gggggtgatg	gcactatcac	ctggagagcc	950
atcccccg	accctttgaa	gtgaatgcag	agggaaacct	ctacgtgacc	1000
agagagctgg	acagagaagc	ccaggctgag	tacctgtctc	aggtgcgggc	1050
tcagaattcc	catggcgagg	actatgcggc	ccctctggag	ctgcacgtgc	1100

-continued

tggtgatgga tgagaatgac aacgtgccta tctgccctcc ccgtgacccc	1150
acagtcagca tccctgagct cagtccacca ggtactgaag tgactagact	1200
gtcagcagag gatgcagatg cccccggctc ccccaattcc cacgttgtgt	1250
atcagctcct gagccctgag cctgaggatg gggtagaggg gagagccttc	1300
caggtggacc ccacttcagg cagtgtgacg ctgggggtgc tccactccg	1350
agcaggccag aacatcctgc ttctggtgct ggccatggac ctggcaggcg	1400
cagagggtgg cttcagcagc acgtgtgaag tcgaagtcgc agtcacagat	1450
atcaatgac acgcccctga gttcatcact tcccagattg ggcctataag	1500
cctccctgag gatgtggagc cggggactct ggtggccatg ctaacagcca	1550
ttgatgctga cctcagagcc gccttcgcc tcattggattt tgccattgag	1600
aggggagaca cagaagggac ttttggcctg gattgggagc cagactctgg	1650
gcatgttaga ctcagactct gcaagaacct cagttatgag gcagctocaa	1700
gtcatgaggt ggtggtggtg gtgcagagtg tggcgaagct ggtggggcca	1750
ggcccaggcc ctggagccac cgccacggtg actgtgctag tgagagagat	1800
gatgccaccc cccaagttgg accaggagag ctacgaggcc agtgtcccca	1850
tcagtgcctc agccggctct ttctgctga ccctcagcc ctccgacccc	1900
atcagccgaa ccctcagggt ctccctagtc aatgactcag agggctggct	1950
ctgcattgag aaattctccg gggaggtgca caccgccag tcctgcagag	2000
gcgcccagcc tggggacacc tacacggtgc ttgtggaggc ccaggataca	2050
gccctgactc ttgcccctgt gccctcccaa tacctctgca caccocgcca	2100
agaccatggc ttgatcgtga gtggacccag caaggacccc gatctggcca	2150
gtgggcacgg tccctacagc ttcaccttg gtcccaaccc cacggtgcaa	2200
cgggattggc gcctccagac tctcaatggt tccatgcct acctcacctt	2250
ggccctgcat tgggtggagc cacgtgaaca cataatcccc gtggtggtca	2300
gccacaatgc ccagatgtgg cagctcctgg ttcgagtgat cgtgtgtcgc	2350
tgcaacgtgg aggggcagtg catgcgcaag gtgggccgca tgaaggcat	2400
gcccacgaag ctgtcggcag tgggcatcct tgtaggcacc ctggtagcaa	2450
taggaatctt cctcatcctc attttcaccc actggacat gtcaagggaag	2500
aaggacccgg atcaaccagc agacagcgtg cccctgaagg cgactgtctg	2550
aatggcccag gcagctctag ctgggagctt ggctctggc tccatctgag	2600
tcccctggga gagagcccag cacccaagat ccagcagggg acaggacaga	2650
gtagaagccc ctccatctgc cctggggtg aggcaccatc accatcacca	2700
ggcatgtctg cagagcctgg acaccaactt tatggactgc ccatgggagt	2750
gtccaaatg tcaggggtgt tgcccaataa taaagcccca gagaactggg	2800
ctgggcccta tgggaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaag	2848

<210> SEQ ID NO 98

<211> LENGTH: 807

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 98

Met	Val	Pro	Ala	Trp	Leu	Trp	Leu	Leu	Cys	Val	Ser	Val	Pro	Gln	1	5	10	15
Ala	Leu	Pro	Lys	Ala	Gln	Pro	Ala	Glu	Leu	Ser	Val	Glu	Val	Pro	20	25	30	
Glu	Asn	Tyr	Gly	Gly	Asn	Phe	Pro	Leu	Tyr	Leu	Thr	Lys	Leu	Pro	35	40	45	
Leu	Pro	Arg	Glu	Gly	Ala	Glu	Gly	Gln	Ile	Val	Leu	Ser	Gly	Asp	50	55	60	
Ser	Gly	Lys	Ala	Thr	Glu	Gly	Pro	Phe	Ala	Met	Asp	Pro	Asp	Ser	65	70	75	
Gly	Phe	Leu	Leu	Val	Thr	Arg	Ala	Leu	Asp	Arg	Glu	Glu	Gln	Ala	80	85	90	
Glu	Tyr	Gln	Leu	Gln	Val	Thr	Leu	Glu	Met	Gln	Asp	Gly	His	Val	95	100	105	
Leu	Trp	Gly	Pro	Gln	Pro	Val	Leu	Val	His	Val	Lys	Asp	Glu	Asn	110	115	120	
Asp	Gln	Val	Pro	His	Phe	Ser	Gln	Ala	Ile	Tyr	Arg	Ala	Arg	Leu	125	130	135	
Ser	Arg	Gly	Thr	Arg	Pro	Gly	Ile	Pro	Phe	Leu	Phe	Leu	Glu	Ala	140	145	150	
Ser	Asp	Arg	Asp	Glu	Pro	Gly	Thr	Ala	Asn	Ser	Asp	Leu	Arg	Phe	155	160	165	
His	Ile	Leu	Ser	Gln	Ala	Pro	Ala	Gln	Pro	Ser	Pro	Asp	Met	Phe	170	175	180	
Gln	Leu	Glu	Pro	Arg	Leu	Gly	Ala	Leu	Ala	Leu	Ser	Pro	Lys	Gly	185	190	195	
Ser	Thr	Ser	Leu	Asp	His	Ala	Leu	Glu	Arg	Thr	Tyr	Gln	Leu	Leu	200	205	210	
Val	Gln	Val	Lys	Asp	Met	Gly	Asp	Gln	Ala	Ser	Gly	His	Gln	Ala	215	220	225	
Thr	Ala	Thr	Val	Glu	Val	Ser	Ile	Ile	Glu	Ser	Thr	Trp	Val	Ser	230	235	240	
Leu	Glu	Pro	Ile	His	Leu	Ala	Glu	Asn	Leu	Lys	Val	Leu	Tyr	Pro	245	250	255	
His	His	Met	Ala	Gln	Val	His	Trp	Ser	Gly	Gly	Asp	Val	His	Tyr	260	265	270	
His	Leu	Glu	Ser	His	Pro	Pro	Gly	Pro	Phe	Glu	Val	Asn	Ala	Glu	275	280	285	
Gly	Asn	Leu	Tyr	Val	Thr	Arg	Glu	Leu	Asp	Arg	Glu	Ala	Gln	Ala	290	295	300	
Glu	Tyr	Leu	Leu	Gln	Val	Arg	Ala	Gln	Asn	Ser	His	Gly	Glu	Asp	305	310	315	
Tyr	Ala	Ala	Pro	Leu	Glu	Leu	His	Val	Leu	Val	Met	Asp	Glu	Asn	320	325	330	
Asp	Asn	Val	Pro	Ile	Cys	Pro	Pro	Arg	Asp	Pro	Thr	Val	Ser	Ile	335	340	345	
Pro	Glu	Leu	Ser	Pro	Pro	Gly	Thr	Glu	Val	Thr	Arg	Leu	Ser	Ala	350	355	360	
Glu	Asp	Ala	Asp	Ala	Pro	Gly	Ser	Pro	Asn	Ser	His	Val	Val	Tyr	365	370	375	

-continued

Gln Leu Leu Ser	Pro Glu Pro Glu Asp	Gly Val Glu Gly Arg Ala	
380		385	390
Phe Gln Val Asp	Pro Thr Ser Gly Ser	Val Thr Leu Gly Val Leu	
395		400	405
Pro Leu Arg Ala	Gly Gln Asn Ile Leu	Leu Leu Val Leu Ala Met	
410		415	420
Asp Leu Ala Gly	Ala Glu Gly Gly Phe	Ser Ser Thr Cys Glu Val	
425		430	435
Glu Val Ala Val	Thr Asp Ile Asn Asp	His Ala Pro Glu Phe Ile	
440		445	450
Thr Ser Gln Ile	Gly Pro Ile Ser Leu	Pro Glu Asp Val Glu Pro	
455		460	465
Gly Thr Leu Val	Ala Met Leu Thr Ala	Ile Asp Ala Asp Leu Glu	
470		475	480
Pro Ala Phe Arg	Leu Met Asp Phe Ala	Ile Glu Arg Gly Asp Thr	
485		490	495
Glu Gly Thr Phe	Gly Leu Asp Trp Glu	Pro Asp Ser Gly His Val	
500		505	510
Arg Leu Arg Leu	Cys Lys Asn Leu Ser	Tyr Glu Ala Ala Pro Ser	
515		520	525
His Glu Val Val	Val Val Val Gln Ser	Val Ala Lys Leu Val Gly	
530		535	540
Pro Gly Pro Gly	Pro Gly Ala Thr Ala	Thr Val Thr Val Leu Val	
545		550	555
Glu Arg Val Met	Pro Pro Pro Lys Leu	Asp Gln Glu Ser Tyr Glu	
560		565	570
Ala Ser Val Pro	Ile Ser Ala Pro Ala	Gly Ser Phe Leu Leu Thr	
575		580	585
Ile Gln Pro Ser	Asp Pro Ile Ser Arg	Thr Leu Arg Phe Ser Leu	
590		595	600
Val Asn Asp Ser	Glu Gly Trp Leu Cys	Ile Glu Lys Phe Ser Gly	
605		610	615
Glu Val His Thr	Ala Gln Ser Leu Gln	Gly Ala Gln Pro Gly Asp	
620		625	630
Thr Tyr Thr Val	Leu Val Glu Ala Gln	Asp Thr Ala Leu Thr Leu	
635		640	645
Ala Pro Val Pro	Ser Gln Tyr Leu Cys	Thr Pro Arg Gln Asp His	
650		655	660
Gly Leu Ile Val	Ser Gly Pro Ser Lys	Asp Pro Asp Leu Ala Ser	
665		670	675
Gly His Gly Pro	Tyr Ser Phe Thr Leu	Gly Pro Asn Pro Thr Val	
680		685	690
Gln Arg Asp Trp	Arg Leu Gln Thr Leu	Asn Gly Ser His Ala Tyr	
695		700	705
Leu Thr Leu Ala	Leu His Trp Val Glu	Pro Arg Glu His Ile Ile	
710		715	720
Pro Val Val Val	Ser His Asn Ala Gln	Met Trp Gln Leu Leu Val	
725		730	735
Arg Val Ile Val	Cys Arg Cys Asn Val	Glu Gly Gln Cys Met Arg	
740		745	750

-continued

Lys	Val	Gly	Arg	Met	Lys	Gly	Met	Pro	Thr	Lys	Leu	Ser	Ala	Val
				755					760					765
Gly	Ile	Leu	Val	Gly	Thr	Leu	Val	Ala	Ile	Gly	Ile	Phe	Leu	Ile
				770					775					780
Leu	Ile	Phe	Thr	His	Trp	Thr	Met	Ser	Arg	Lys	Lys	Asp	Pro	Asp
				785					790					795
Gln	Pro	Ala	Asp	Ser	Val	Pro	Leu	Lys	Ala	Thr	Val			
				800					805					

<210> SEQ ID NO 99
<211> LENGTH: 2436
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 99

ggctgaccgt gctacattgc ctggaggaag cctaaggaac ccaggcatcc	50
agctgcccc gcctgagtcc aagattcttc ccaggaacac aaacgtagga	100
gacccacgct cctggaagca ccagccttta tctcttcacc ttcaagtccc	150
ctttctcaag aatcctctgt tctttgccct ctaaagtctt ggtacatcta	200
ggacccaggc atcttgcttt ccagccacaa agagacagat gaagatgcag	250
aaaggaaatg ttctccttat gtttggctta ctattgcatt tagaagctgc	300
aacaaattcc aatgagacta gcacctctgc caacactgga tccagtgtga	350
tctccagtgg agccagcaca gccaccaact ctgggtccag tgtgacctcc	400
agtggtggta gcacagccac catctcaggg tccagcgtga cctccaatgg	450
ggtcagcata gtcaccaact ctgagttcca tacaacctcc agtgggatca	500
gcacagccac caactctgag ttcagcacag cgtccagtgg gatcagcata	550
gccaccaact ctgagtcag cacaacctcc agtggggcca gcacagccac	600
caactctgag tccagcacac cctccagtgg ggccagcaca gtcaccaact	650
ctgggtccag tgtgacctcc agtggagcca gcaactgccc caactctgag	700
tccagcacag tgtccagtag ggccagcact gccaccaact ctgagtctag	750
cacactctcc agtggggcca gcacagccac caactctgac tccagcaca	800
cctccagtgg ggctagcaca gccaccaact ctgagtcag cacaacctcc	850
agtggggcca gcacagccac caactctgag tccagcacag tgtccagtag	900
ggccagcact gccaccaact ctgagtcag cacaacctcc agtggggcca	950
gcacagccac caactctgag tccagaacga cctccaatgg ggtgggcaca	1000
gccaccaact ctgagtcag cagcacctcc agtggggcca gcacagccac	1050
caactctgac tccagcacag tgtccagtgg ggccagcact gccaccaact	1100
ctgagtcag cagcacctcc agtggggcca gcacagccac caactctgag	1150
tccagcacga cctccagtgg ggctagcaca gccaccaact ctgactccag	1200
cacaacctcc agtggggccg gcacagccac caactctgag tccagcacag	1250
tgtccagtgg gatcagcaca gtcaccaatt ctgagtcag cacacctcc	1300
agtggggcca acacagccac caactctgag tccagtacga cctccagtgg	1350
ggccaacaca gccaccaact ctgagtcag cacagtgtcc agtggggcca	1400

-continued

gcactgccac caactctgag tccagcacaa cctccagtgg ggtcagcaca	1450
gccaccaact ctgagtcag cacaacctcc agtggggcta gcacagccac	1500
caactctgac tccagcacaa cctccagtga ggccagcaca gccaccaact	1550
ctgagtctag cacagtgtcc agtgggatca gcacagtcac caattctgag	1600
tccagcacaa cctccagtgg ggccaacaca gccaccaact ctgggtccag	1650
tgtgacctct gcaggctctg gaacagcagc tctgactgga atgcacacaa	1700
cttcccatag tgcattctact gcagtgagtg aggcaaagcc tgggtgggtcc	1750
ctgggtgccgt gggaaatctt cctcatcacc ctggtctcgg ttgtggcggc	1800
cgtggggctc tttgctgggc tcttcttctg tgtgagaaac agcctgtccc	1850
tgagaaacac ctttaacaca gctgtctacc accctcatgg cctcaaccat	1900
ggccttggtc caggccctgg agggaatcat ggagccccc acaggccag	1950
gtggagtctt aactggttct ggaggagacc agtatcatcg atagccatgg	2000
agatgagcgg gaggaacagc gggccctgag cagccccgga agcaagtgcc	2050
gcattcttca ggaaggaaga gacctgggca cccaagacct ggtttccttt	2100
cattcatccc aggagacccc tcccagcttt gtttgagatc ctgaaaatct	2150
tgaagaaggt attcctcacc tttcttgctt ttaccagaca ctggaagag	2200
aatactatat tgctcattta gctaagaaat aaatacatct catctaacac	2250
acacgacaaa gagaagctgt gcttgccccg gggtggttat ctgctctga	2300
gatgaactca gttataggag aaaacctcca tgctggactc catctggcat	2350
tcaaaatctc cacagtaaaa tccaagacc tcaaaaaaaaa aaaaaaaaaa	2400
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa	2436

<210> SEQ ID NO 100

<211> LENGTH: 596

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 100

Met	Lys	Met	Gln	Lys	Gly	Asn	Val	Leu	Leu	Met	Phe	Gly	Leu	Leu
1				5					10					15
Leu	His	Leu	Glu	Ala	Ala	Thr	Asn	Ser	Asn	Glu	Thr	Ser	Thr	Ser
				20					25					30
Ala	Asn	Thr	Gly	Ser	Ser	Val	Ile	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				35					40					45
Thr	Asn	Ser	Gly	Ser	Ser	Val	Thr	Ser	Ser	Gly	Val	Ser	Thr	Ala
				50					55					60
Thr	Ile	Ser	Gly	Ser	Ser	Val	Thr	Ser	Asn	Gly	Val	Ser	Ile	Val
				65					70					75
Thr	Asn	Ser	Glu	Phe	His	Thr	Thr	Ser	Ser	Gly	Ile	Ser	Thr	Ala
				80					85					90
Thr	Asn	Ser	Glu	Phe	Ser	Thr	Ala	Ser	Ser	Gly	Ile	Ser	Ile	Ala
				95					100					105
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				110					115					120
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Pro	Ser	Ser	Gly	Ala	Ser	Thr	Val
				125					130					135

-continued

Thr	Asn	Ser	Gly	Ser	Ser	Val	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				140						145				150
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Val	Ser	Ser	Arg	Ala	Ser	Thr	Ala
				155						160				165
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Leu	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				170						175				180
Thr	Asn	Ser	Asp	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				185						190				195
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				200						205				210
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Val	Ser	Ser	Arg	Ala	Ser	Thr	Ala
				215						220				225
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				230						235				240
Thr	Asn	Ser	Glu	Ser	Arg	Thr	Thr	Ser	Asn	Gly	Ala	Gly	Thr	Ala
				245						250				255
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				260						265				270
Thr	Asn	Ser	Asp	Ser	Ser	Thr	Val	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				275						280				285
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				290						295				300
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				305						310				315
Thr	Asn	Ser	Asp	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Gly	Thr	Ala
				320						325				330
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Val	Ser	Ser	Gly	Ile	Ser	Thr	Val
				335						340				345
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Pro	Ser	Ser	Gly	Ala	Asn	Thr	Ala
				350						355				360
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Asn	Thr	Ala
				365						370				375
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Val	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				380						385				390
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Val	Ser	Thr	Ala
				395						400				405
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Ser	Thr	Ala
				410						415				420
Thr	Asn	Ser	Asp	Ser	Ser	Thr	Thr	Ser	Ser	Glu	Ala	Ser	Thr	Ala
				425						430				435
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Val	Ser	Ser	Gly	Ile	Ser	Thr	Val
				440						445				450
Thr	Asn	Ser	Glu	Ser	Ser	Thr	Thr	Ser	Ser	Gly	Ala	Asn	Thr	Ala
				455						460				465
Thr	Asn	Ser	Gly	Ser	Ser	Val	Thr	Ser	Ala	Gly	Ser	Gly	Thr	Ala
				470						475				480
Ala	Leu	Thr	Gly	Met	His	Thr	Thr	Ser	His	Ser	Ala	Ser	Thr	Ala
				485						490				495
Val	Ser	Glu	Ala	Lys	Pro	Gly	Gly	Ser	Leu	Val	Pro	Trp	Glu	Ile
				500						505				510

-continued

Phe	Leu	Ile	Thr	Leu	Val	Ser	Val	Val	Ala	Ala	Val	Gly	Leu	Phe
			515						520				525	
Ala	Gly	Leu	Phe	Phe	Cys	Val	Arg	Asn	Ser	Leu	Ser	Leu	Arg	Asn
			530						535				540	
Thr	Phe	Asn	Thr	Ala	Val	Tyr	His	Pro	His	Gly	Leu	Asn	His	Gly
			545						550				555	
Leu	Gly	Pro	Gly	Pro	Gly	Gly	Asn	His	Gly	Ala	Pro	His	Arg	Pro
			560						565				570	
Arg	Trp	Ser	Pro	Asn	Trp	Phe	Trp	Arg	Arg	Pro	Val	Ser	Ser	Ile
			575						580				585	
Ala	Met	Glu	Met	Ser	Gly	Arg	Asn	Ser	Gly	Pro				
			590						595					

<210> SEQ ID NO 101
<211> LENGTH: 1728
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 101

ggccggacgc	ctccgcgtta	cgggatgaat	taacggcggg	ttccgcacgg	50
aggttgtagc	ccctacggag	ccccagcttg	cccacgcacc	ccactcggcg	100
tcgcgcggcg	tgccttgctt	gtcacagggtg	ggaggctgga	actatcaggc	150
tgaaaaacag	agtggttact	ctcttctggg	aagctggcaa	caaattggatg	200
atgtgatata	tgcattccag	gggaagggaa	attgtggtgc	ttctgaaccc	250
atggtaatt	aacgaggcag	tttctagcta	ctgcacgtac	ttcataaagc	300
aggactctaa	aagctttgga	atcatgggtg	catggaaagg	gatttacttt	350
atactgactc	tgttttgggg	aagctttttt	ggaagcattt	tcattgctgag	400
tcccttttta	cctttgatgt	ttgtaaaccc	atcttggtat	cgctggatca	450
acaaccgcct	tgtggcaaca	tggctcaccc	tacctgtggc	attattggag	500
accatgtttg	gtgtaaaagt	gattataact	ggggatgcat	ttgttcctgg	550
agaaagaagt	gtcattatca	tgaaccatcg	gacaagaatg	gactggatgt	600
tcctgtggaa	ttgcctgatg	cgatatagct	acctcagatt	ggagaaaatt	650
tgctcaaag	cgagtctcaa	aggtgttcct	ggatttggtt	gggccatgca	700
ggctgctgcc	tatatcttca	ttcataggaa	atggaaggat	gacaagagcc	750
atttcgaaga	catgattgat	tactttttgtg	atattcacga	accacttcaa	800
ctcctcatat	tcccagaagg	gactgatctc	acagaaaaca	gcaagtctcg	850
aagtaatgca	tttgctgaaa	aaaatggact	tcagaaatat	gaatatgttt	900
tacatccaag	aactacaggc	tttacttttg	tggtagaccg	tctaagagaa	950
ggtaagaacc	ttgatgctgt	ccatgatatc	actgtggcgt	atcctcacia	1000
cattcctcaa	tcagagaagc	acctcctcca	aggagacttt	cccagggaaa	1050
tccactttca	cgtccaccgg	tatccaatag	acaccctccc	cacatccaag	1100
gaggaccttc	aactctggtg	ccacaaacgg	tgggaagaga	aagaagagag	1150
gctgcgttcc	ttctatcaag	gggagaagaa	tttttatatt	accggacaga	1200
gtgtcattcc	accttgcaag	tctgaactca	gggtccttgt	ggtcaaatg	1250

-continued

ctctctatac tgtattggac cctgttcagc cctgcaatgt gcctactcat	1300
atatttgtac agtcttgta agtggtattt tataatcacc attgtaatct	1350
ttgtgctgca agagagaata ttgtgtggac tggagatcat agaacttgca	1400
tgttaccgac ttttacacaa acagccacat ttaaattcaa agaaaaatga	1450
gtaagattat aaggtttgcc atgtgaaaac ctagagcata ttttggaat	1500
gttctaaacc tttctaagct cagatgcatt tttgcatgac tatgtcgaat	1550
atttcttact gccatcatta ttgtttaaag atattttgca cttaattttg	1600
tgggaaaaat attgctacaa ttttttttaa tctctgaatg taatttcgat	1650
actgtgtaca tagcagggag tgatcgggt gaaataactt gggccagaat	1700
attattaaac aatcatcagg cttttaaa	1728

<210> SEQ ID NO 102

<211> LENGTH: 414

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 102

Met His Ser Arg Gly Arg Glu Ile Val Val Leu Leu Asn Pro Trp	
1 5 10 15	
Ser Ile Asn Glu Ala Val Ser Ser Tyr Cys Thr Tyr Phe Ile Lys	
20 25 30	
Gln Asp Ser Lys Ser Phe Gly Ile Met Val Ser Trp Lys Gly Ile	
35 40 45	
Tyr Phe Ile Leu Thr Leu Phe Trp Gly Ser Phe Phe Gly Ser Ile	
50 55 60	
Phe Met Leu Ser Pro Phe Leu Pro Leu Met Phe Val Asn Pro Ser	
65 70 75	
Trp Tyr Arg Trp Ile Asn Asn Arg Leu Val Ala Thr Trp Leu Thr	
80 85 90	
Leu Pro Val Ala Leu Leu Glu Thr Met Phe Gly Val Lys Val Ile	
95 100 105	
Ile Thr Gly Asp Ala Phe Val Pro Gly Glu Arg Ser Val Ile Ile	
110 115 120	
Met Asn His Arg Thr Arg Met Asp Trp Met Phe Leu Trp Asn Cys	
125 130 135	
Leu Met Arg Tyr Ser Tyr Leu Arg Leu Glu Lys Ile Cys Leu Lys	
140 145 150	
Ala Ser Leu Lys Gly Val Pro Gly Phe Gly Trp Ala Met Gln Ala	
155 160 165	
Ala Ala Tyr Ile Phe Ile His Arg Lys Trp Lys Asp Asp Lys Ser	
170 175 180	
His Phe Glu Asp Met Ile Asp Tyr Phe Cys Asp Ile His Glu Pro	
185 190 195	
Leu Gln Leu Leu Ile Phe Pro Glu Gly Thr Asp Leu Thr Glu Asn	
200 205 210	
Ser Lys Ser Arg Ser Asn Ala Phe Ala Glu Lys Asn Gly Leu Gln	
215 220 225	
Lys Tyr Glu Tyr Val Leu His Pro Arg Thr Thr Gly Phe Thr Phe	
230 235 240	

-continued

Val	Val	Asp	Arg	Leu	Arg	Glu	Gly	Lys	Asn	Leu	Asp	Ala	Val	His
				245					250					255
Asp	Ile	Thr	Val	Ala	Tyr	Pro	His	Asn	Ile	Pro	Gln	Ser	Glu	Lys
				260					265					270
His	Leu	Leu	Gln	Gly	Asp	Phe	Pro	Arg	Glu	Ile	His	Phe	His	Val
				275					280					285
His	Arg	Tyr	Pro	Ile	Asp	Thr	Leu	Pro	Thr	Ser	Lys	Glu	Asp	Leu
				290					295					300
Gln	Leu	Trp	Cys	His	Lys	Arg	Trp	Glu	Glu	Lys	Glu	Glu	Arg	Leu
				305					310					315
Arg	Ser	Phe	Tyr	Gln	Gly	Glu	Lys	Asn	Phe	Tyr	Phe	Thr	Gly	Gln
				320					325					330
Ser	Val	Ile	Pro	Pro	Cys	Lys	Ser	Glu	Leu	Arg	Val	Leu	Val	Val
				335					340					345
Lys	Leu	Leu	Ser	Ile	Leu	Tyr	Trp	Thr	Leu	Phe	Ser	Pro	Ala	Met
				350					355					360
Cys	Leu	Leu	Ile	Tyr	Leu	Tyr	Ser	Leu	Val	Lys	Trp	Tyr	Phe	Ile
				365					370					375
Ile	Thr	Ile	Val	Ile	Phe	Val	Leu	Gln	Glu	Arg	Ile	Phe	Gly	Gly
				380					385					390
Leu	Glu	Ile	Ile	Glu	Leu	Ala	Cys	Tyr	Arg	Leu	Leu	His	Lys	Gln
				395					400					405
Pro	His	Leu	Asn	Ser	Lys	Lys	Asn	Glu						
				410										

<210> SEQ ID NO 103
<211> LENGTH: 2403
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 103

cggctcgagc	ggctcgagt	aagagcctct	ccacggctcc	tgcgcctgag	50
acagctggcc	tgacctccaa	atcatccatc	cacctctgct	gtcatctgtt	100
ttcatagtgt	gagatcaacc	cacaggaata	tccatggctt	ttgtgctcat	150
tttggttctc	agtttctacg	agctggtgtc	aggacagtgg	caagtcactg	200
gaccgggcaa	gtttgtccag	gccttggtgg	gggaggacgc	cgtgttctcc	250
tgctccctct	ttcctgagac	cagtgcagag	gctatggaag	tgcggttctt	300
caggaatcag	ttccatgctg	tggtccacct	ctacagagat	ggggaagact	350
gggaatctaa	gcagatgcca	cagtatcgag	ggagaactga	gtttgtgaag	400
gactccattg	caggggggcg	tgtctctcta	aggctaaaaa	acatcactcc	450
ctcggacatc	ggcctgtatg	ggtgctggtt	cagttcccag	atttacgatg	500
aggaggccac	ctgggagctg	cgggtggcag	cactgggctc	acttcctctc	550
atttccatcg	tgggatatgt	tgacggagg	atccagttac	tctgcctgtc	600
ctcaggctgg	ttccccagc	ccacagccaa	gtgaaaggt	ccacaaggac	650
aggatttgtc	ttcagactcc	agagcaaagt	cagatgggta	cagcctgtat	700
gatgtggaga	tctccattat	agtccaggaa	aatgctggga	gcatattgtg	750
ttccatccac	cttgctgagc	agagtcatga	ggtggaatcc	aaggtattga	800

```
<210> SEQ ID NO 104
<211> LENGTH: 466
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 104
```

Met Ala Phe Val Leu Ile Leu Val Leu Ser Phe Tyr Glu Leu Val
1 5 10 15

-continued

Ser Gly Gln Trp	Gln Val Thr Gly Pro	Gly Lys Phe Val Gln Ala	20	25	30
Leu Val Gly Glu	Asp Ala Val Phe Ser	Cys Ser Leu Phe Pro Glu	35	40	45
Thr Ser Ala Glu	Ala Met Glu Val Arg	Phe Phe Arg Asn Gln Phe	50	55	60
His Ala Val Val	His Leu Tyr Arg Asp	Gly Glu Asp Trp Glu Ser	65	70	75
Lys Gln Met Pro	Gln Tyr Arg Gly Arg	Thr Glu Phe Val Lys Asp	80	85	90
Ser Ile Ala Gly	Gly Arg Val Ser Leu	Arg Leu Lys Asn Ile Thr	95	100	105
Pro Ser Asp Ile	Gly Leu Tyr Gly Cys	Trp Phe Ser Ser Gln Ile	110	115	120
Tyr Asp Glu Glu	Ala Thr Trp Glu Leu	Arg Val Ala Ala Leu Gly	125	130	135
Ser Leu Pro Leu	Ile Ser Ile Val Gly	Tyr Val Asp Gly Gly Ile	140	145	150
Gln Leu Leu Cys	Leu Ser Ser Gly Trp	Phe Pro Gln Pro Thr Ala	155	160	165
Lys Trp Lys Gly	Pro Gln Gly Gln Asp	Leu Ser Ser Asp Ser Arg	170	175	180
Ala Asn Ala Asp	Gly Tyr Ser Leu Tyr	Asp Val Glu Ile Ser Ile	185	190	195
Ile Val Gln Glu	Asn Ala Gly Ser Ile	Leu Cys Ser Ile His Leu	200	205	210
Ala Glu Gln Ser	His Glu Val Glu Ser	Lys Val Leu Ile Gly Glu	215	220	225
Thr Phe Phe Gln	Pro Ser Pro Trp Arg	Leu Ala Ser Ile Leu Leu	230	235	240
Gly Leu Leu Cys	Gly Ala Leu Cys Gly	Val Val Met Gly Met Ile	245	250	255
Ile Val Phe Phe	Lys Ser Lys Gly Lys	Ile Gln Ala Glu Leu Asp	260	265	270
Trp Arg Arg Lys	His Gly Gln Ala Glu	Leu Arg Asp Ala Arg Lys	275	280	285
His Ala Val Glu	Val Thr Leu Asp Pro	Glu Thr Ala His Pro Lys	290	295	300
Leu Cys Val Ser	Asp Leu Lys Thr Val	Thr His Arg Lys Ala Pro	305	310	315
Gln Glu Val Pro	His Ser Glu Lys Arg	Phe Thr Arg Lys Ser Val	320	325	330
Val Ala Ser Gln	Gly Phe Gln Ala Gly	Arg His Tyr Trp Glu Val	335	340	345
Asp Val Gly Gln	Asn Val Gly Trp Tyr	Val Gly Val Cys Arg Asp	350	355	360
Asp Val Asp Arg	Gly Lys Asn Asn Val	Thr Leu Ser Pro Asn Asn	365	370	375
Gly Tyr Trp Val	Leu Arg Leu Thr Thr	Glu His Leu Tyr Phe Thr	380	385	390

-continued

Phe	Asn	Pro	His	Phe	Ile	Ser	Leu	Pro	Pro	Ser	Thr	Pro	Pro	Thr
			395					400						405
Arg	Val	Gly	Val	Phe	Leu	Asp	Tyr	Glu	Gly	Gly	Thr	Ile	Ser	Phe
			410					415						420
Phe	Asn	Thr	Asn	Asp	Gln	Ser	Leu	Ile	Tyr	Thr	Leu	Leu	Thr	Cys
			425					430						435
Gln	Phe	Glu	Gly	Leu	Leu	Arg	Pro	Tyr	Ile	Gln	His	Ala	Met	Tyr
			440					445						450
Asp	Glu	Glu	Lys	Gly	Thr	Pro	Ile	Phe	Ile	Cys	Pro	Val	Ser	Trp
			455					460						465

Gly

<210> SEQ ID NO 105
<211> LENGTH: 2103
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 105

ccttcacagg	actcttcatt	gctggttggc	aatgatgtat	cggccagatg	50
tggtgagggc	taggaaaaga	gtttgttggg	aaccctgggt	tatcggcctc	100
gtcatcttca	tatccctgat	tgtcctggca	gtgtgcattg	gactcactgt	150
tcattatgtg	agatataatc	aaaagaagac	ctacaattac	tatagcacat	200
tgtcatttac	aactgacaaa	ctatatgctg	agtttggcag	agaggcttct	250
aacaatttta	cagaaatgag	ccagagactt	gaatcaatgg	tgaaaaatgc	300
attttataaa	tctccattaa	gggaagaatt	tgtcaagtct	caggttatca	350
agttcagtc	acagaagcat	ggagtgttgg	ctcatatgct	gttgatttgt	400
agatttcact	ctactgagga	tcctgaaact	gtagataaaa	ttgttcaact	450
tgttttacat	gaaaagctgc	aagatgctgt	aggaccccct	aaagtagatc	500
ctcactcagt	taaaattaaa	aaaatcaaca	agacagaaac	agacagctat	550
ctaaaccatt	gctgcggaac	acgaagaagt	aaaactctag	gtcagagtct	600
caggatcggt	ggtgggacag	aagtagaaga	gggtgaatgg	ccctggcagg	650
ctagcctgca	gtgggatggg	agtcatcgct	gtggagcaac	cttaattaat	700
gccacatggc	ttgtgagtgc	tgctcactgt	tttacaacat	ataagaaccc	750
tgccagatgg	actgcttcct	ttggagtaac	aataaaacct	tcgaaaatga	800
aacggggtct	ccggagaata	attgtccatg	aaaaatacaa	acacccatca	850
catgactatg	atatttctct	tcagagctt	tctagccctg	ttccctacac	900
aaatgcagta	catagagttt	gtctccctga	tgcatcctat	gagtttcaac	950
cagggtgatg	gatgtttgtg	acaggatttg	gagcactgaa	aaatgatgg	1000
tacagtcaaa	atcatcttcg	acaagcacag	gtgactctca	tagacgctac	1050
aacttgcaat	gaacctcaag	cttacaatga	cgccataact	cctagaatgt	1100
tatgtgctgg	ctccttagaa	ggaaaacag	atgcatgcc	gggtgactct	1150
ggaggaccac	tggttagttc	agatgctaga	gatatctggt	accttgctgg	1200
aatagtgagc	tggggagatg	aatgtgcgaa	acccaacaag	cctggtgttt	1250
atactagagt	tacggccttg	cgggactgga	ttacttcaaa	aactggtatc	1300

-continued

taagagacaa aagcctcatg gaacagataa cttttttttt tgttttttg	1350
gtgtggaggc cttttttaga gatacagaat tggagaagac ttgcaaaaca	1400
gctagatttg actgatctca ataaactgtt tgcttgatgc atgtattttc	1450
ttcccagctc tgttccgcac gtaagcatcc tgcttctgcc agatcaactc	1500
tgatcatctgt gagcaatagt tgaaacttta tgtacataga gaaatagata	1550
atacaatatt acattacagc ctgtattcat ttgttctcta gaagttttgt	1600
cagaattttg acttggtgac ataaatttgt aatgcatata tacaatttga	1650
agcactcctt ttcttcagtt cctcagctcc tctcatttca gcaaatatcc	1700
attttcaagg tgcagaacaa ggagtgaag aaaatataag aagaaaaaaa	1750
tcccctacat tttattggca cagaaaagta ttaggtgttt ttcttagtgg	1800
aatattagaa atgatcatat tcattatgaa aggtcaagca aagacagcag	1850
aataccaatc acttcatcat ttaggaagta tgggaactaa gttaaggaag	1900
tccagaaaga agccaagata tatccttatt ttcatttcca aacaactact	1950
atgataaatg tgaagaagat tctgtttttt tgtgacctat aataattata	2000
caaacttcat gcaatgtact tgttctaagc aaattaaagc aaatatttat	2050
ttaacattgt tactgaggat gtcaacatat aacaataaaa tataaatcac	2100
cca	2103

<210> SEQ ID NO 106

<211> LENGTH: 423

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 106

Met Met Tyr Arg Pro Asp Val Val Arg Ala Arg Lys Arg Val Cys	
1 5 10 15	
Trp Glu Pro Trp Val Ile Gly Leu Val Ile Phe Ile Ser Leu Ile	
20 25 30	
Val Leu Ala Val Cys Ile Gly Leu Thr Val His Tyr Val Arg Tyr	
35 40 45	
Asn Gln Lys Lys Thr Tyr Asn Tyr Tyr Ser Thr Leu Ser Phe Thr	
50 55 60	
Thr Asp Lys Leu Tyr Ala Glu Phe Gly Arg Glu Ala Ser Asn Asn	
65 70 75	
Phe Thr Glu Met Ser Gln Arg Leu Glu Ser Met Val Lys Asn Ala	
80 85 90	
Phe Tyr Lys Ser Pro Leu Arg Glu Glu Phe Val Lys Ser Gln Val	
95 100 105	
Ile Lys Phe Ser Gln Gln Lys His Gly Val Leu Ala His Met Leu	
110 115 120	
Leu Ile Cys Arg Phe His Ser Thr Glu Asp Pro Glu Thr Val Asp	
125 130 135	
Lys Ile Val Gln Leu Val Leu His Glu Lys Leu Gln Asp Ala Val	
140 145 150	
Gly Pro Pro Lys Val Asp Pro His Ser Val Lys Ile Lys Lys Ile	
155 160 165	

-continued

Asn	Lys	Thr	Glu	Thr	Asp	Ser	Tyr	Leu	Asn	His	Cys	Cys	Gly	Thr
			170						175					180
Arg	Arg	Ser	Lys	Thr	Leu	Gly	Gln	Ser	Leu	Arg	Ile	Val	Gly	Gly
			185						190					195
Thr	Glu	Val	Glu	Glu	Gly	Glu	Trp	Pro	Trp	Gln	Ala	Ser	Leu	Gln
			200						205					210
Trp	Asp	Gly	Ser	His	Arg	Cys	Gly	Ala	Thr	Leu	Ile	Asn	Ala	Thr
			215						220					225
Trp	Leu	Val	Ser	Ala	Ala	His	Cys	Phe	Thr	Thr	Tyr	Lys	Asn	Pro
			230						235					240
Ala	Arg	Trp	Thr	Ala	Ser	Phe	Gly	Val	Thr	Ile	Lys	Pro	Ser	Lys
			245						250					255
Met	Lys	Arg	Gly	Leu	Arg	Arg	Ile	Ile	Val	His	Glu	Lys	Tyr	Lys
			260						265					270
His	Pro	Ser	His	Asp	Tyr	Asp	Ile	Ser	Leu	Ala	Glu	Leu	Ser	Ser
			275						280					285
Pro	Val	Pro	Tyr	Thr	Asn	Ala	Val	His	Arg	Val	Cys	Leu	Pro	Asp
			290						295					300
Ala	Ser	Tyr	Glu	Phe	Gln	Pro	Gly	Asp	Val	Met	Phe	Val	Thr	Gly
			305						310					315
Phe	Gly	Ala	Leu	Lys	Asn	Asp	Gly	Tyr	Ser	Gln	Asn	His	Leu	Arg
			320						325					330
Gln	Ala	Gln	Val	Thr	Leu	Ile	Asp	Ala	Thr	Thr	Cys	Asn	Glu	Pro
			335						340					345
Gln	Ala	Tyr	Asn	Asp	Ala	Ile	Thr	Pro	Arg	Met	Leu	Cys	Ala	Gly
			350						355					360
Ser	Leu	Glu	Gly	Lys	Thr	Asp	Ala	Cys	Gln	Gly	Asp	Ser	Gly	Gly
			365						370					375
Pro	Leu	Val	Ser	Ser	Asp	Ala	Arg	Asp	Ile	Trp	Tyr	Leu	Ala	Gly
			380						385					390
Ile	Val	Ser	Trp	Gly	Asp	Glu	Cys	Ala	Lys	Pro	Asn	Lys	Pro	Gly
			395						400					405
Val	Tyr	Thr	Arg	Val	Thr	Ala	Leu	Arg	Asp	Trp	Ile	Thr	Ser	Lys
			410						415					420
Thr	Gly	Ile												

<210> SEQ ID NO 107
<211> LENGTH: 2397
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 107

agagaaagaa gcgctccag ctgaagccaa tgcagccctc cggtctccg	50
cgaagaagtt ccctgcccc atgagcccc gccgtgcgtc cccgactatc	100
cccaggcggg cgtggggcac cgggccacg gccgacgac gctgccgttt	150
tgcccttggg agtaggatgt ggtgaaagga tggggcttct cccttacggg	200
gtcacaaatg gccagagaag attccgtgaa gtgtctgcgc tgctgtctct	250
acgccctcaa tctgtctttt tggttaatgt ccatcagtgt gttggcagtt	300
tctgtctgga tgagggacta cctaaataat gttctcactt taactgcaga	350
aacgagggta gaggaagcag tcattttgac ttactttcct gtggttcac	400

-continued

cggtcatgat tgctgtttgc tgtttcctta tcattgtggg gatgttagga	450
tattgtggaa cggtgaaaag aaatctgttg cttcttgcat ggtactttgg	500
aagtttgctt gtcattttct gtgtagaact ggcttgtggc gtttgacat	550
atgaacagga acttatgggt ccagtacaat ggtcagatat ggtcactttg	600
aaagccagga tgacaaatta tggattacct agatatcggg ggcttactca	650
tgcttggaaat ttttttcaga gagagttaa gtgctgtgga gtagtatatt	700
tcactgactg gttggaaatg acagagatgg actggccccc agattcctgc	750
tgtgttagag aattcccagg atgttccaaa caggcccacc aggaagatct	800
cagtgcctt tatcaagagg gttgtgggaa gaaaatgtat tcctttttga	850
gaggaaccaa acaactgcag gtgctgaggt ttctgggaat ctccattggg	900
gtgacacaaa tcctggccat gattctcacc attactctgc tctgggctct	950
gtattatgat agaagggagc ctgggacaga ccaaatgatg tcottgaaga	1000
atgacaactc tcagcacctg tcatgtccct cagtagaact gttgaaacca	1050
agcctgtcaa gaatctttga acacacatcc atggcaaaca gctttaatac	1100
acactttgag atggaggagt tataaaaaga aatgtcacag aagaaaacca	1150
caaaacttgt ttattggact tgtgaatttt tgagtacata ctatgtgttt	1200
cagaaatatg tagaaataaa aatgttgcca taaaataaca cctaagcata	1250
tactattcta tgctttaaaa tgaggatgga aaagtttcat gtcataagtc	1300
accacctgga caataattga tgcccttaaa atgctgaaga cagatgtcat	1350
accactgtg tagcctgtgt atgactttta ctgaacacag ttatgttttg	1400
aggcagcatg gtttgattag catttccgca tccatgcaa cgagtcacat	1450
atgggtgggac tggagccata gtaaaggttg atttacttct accaactagt	1500
atataaagta ctaattaaat gtaacatag gaagttagaa aatactaata	1550
acttttatta ctacgcgac tattcttctg atgctaaata aattatatat	1600
cagaaaactt tcaatatagg tgactaccta aatgtgattt ttgctgggta	1650
ctaaaaatatt cttaccactt aaaagagcaa gctaacacat tgtcttaagc	1700
tgatcaggga ttttttgat ataagtctgt gttaaactcg tataattcag	1750
tcgatttcag ttctgataat gttaagaata accattatga aaaggaaaat	1800
ttgtcctgta tagcatcatt atttttagcc ttctctgtta ataaagcttt	1850
actattctgt cctgggctta tattacacat ataactgtta tttaaatact	1900
taaccactaa ttttgaaaat taccagtgtg atacatagga atcattattc	1950
agaatgtagt ctggtcttta ggaagtatta ataagaaaat ttgcacataa	2000
cttagttgat tcagaaagga cttgtatgct gtttttctcc caaatgaaga	2050
ctctttttga cactaaacac tttttaaaaa gcttatcttt gccttctcca	2100
aacaagaagc aatagtctcc aagtcaatat aaattctaca gaaaatagtg	2150
ttctttttct ccagaaaaat gcttgtgaga atcattaaaa catgtgacaa	2200
tttagagatt ctttgtttta tttcactgat taatatactg tggcaaatga	2250
cacagattat taaatttttt tacaagagta tagtatattt atttgaaatg	2300

-continued

ggaaaagtgc attttactgt attttgtgta ttttgtttat ttctcagaat 2350

atggaaagaa aattaaaatg tgtcaataaa tattttctag agagtaa 2397

<210> SEQ ID NO 108

<211> LENGTH: 305

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 108

Met Ala Arg Glu Asp Ser Val Lys Cys Leu Arg Cys Leu Leu Tyr
1 5 10 15Ala Leu Asn Leu Leu Phe Trp Leu Met Ser Ile Ser Val Leu Ala
20 25 30Val Ser Ala Trp Met Arg Asp Tyr Leu Asn Asn Val Leu Thr Leu
35 40 45Thr Ala Glu Thr Arg Val Glu Glu Ala Val Ile Leu Thr Tyr Phe
50 55 60Pro Val Val His Pro Val Met Ile Ala Val Cys Cys Phe Leu Ile
65 70 75Ile Val Gly Met Leu Gly Tyr Cys Gly Thr Val Lys Arg Asn Leu
80 85 90Leu Leu Leu Ala Trp Tyr Phe Gly Ser Leu Leu Val Ile Phe Cys
95 100 105Val Glu Leu Ala Cys Gly Val Trp Thr Tyr Glu Gln Glu Leu Met
110 115 120Val Pro Val Gln Trp Ser Asp Met Val Thr Leu Lys Ala Arg Met
125 130 135Thr Asn Tyr Gly Leu Pro Arg Tyr Arg Trp Leu Thr His Ala Trp
140 145 150Asn Phe Phe Gln Arg Glu Phe Lys Cys Cys Gly Val Val Tyr Phe
155 160 165Thr Asp Trp Leu Glu Met Thr Glu Met Asp Trp Pro Pro Asp Ser
170 175 180Cys Cys Val Arg Glu Phe Pro Gly Cys Ser Lys Gln Ala His Gln
185 190 195Glu Asp Leu Ser Asp Leu Tyr Gln Glu Gly Cys Gly Lys Lys Met
200 205 210Tyr Ser Phe Leu Arg Gly Thr Lys Gln Leu Gln Val Leu Arg Phe
215 220 225Leu Gly Ile Ser Ile Gly Val Thr Gln Ile Leu Ala Met Ile Leu
230 235 240Thr Ile Thr Leu Leu Trp Ala Leu Tyr Tyr Asp Arg Arg Glu Pro
245 250 255Gly Thr Asp Gln Met Met Ser Leu Lys Asn Asp Asn Ser Gln His
260 265 270Leu Ser Cys Pro Ser Val Glu Leu Leu Lys Pro Ser Leu Ser Arg
275 280 285Ile Phe Glu His Thr Ser Met Ala Asn Ser Phe Asn Thr His Phe
290 295 300Glu Met Glu Glu Leu
305

-continued

<210> SEQ ID NO 109

<211> LENGTH: 2339

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 109

ccaaggccag agctgtggac accttatccc actcatcctc atcctcttcc	50
tctgataaag cccctaccag tgctgataaa gtctttctcg tgagagccta	100
gaggcccttaa aaaaaaaagt gcttgaaaga gaaggggaca aaggaacacc	150
agtattaaga ggattttcca gtgtttctgg cagttggtcc agaaggatgc	200
ctccattcct gcttctcacc tgccctcttca tcacaggcac ctccgtgtca	250
cccgtggccc tagatccttg ttctgcttac atcagcctga atgagcctg	300
gaggaacact gaccaccagt tggatgagtc tcaaggtcct cctctatgtg	350
acaaccatgt gaatggggag tggtagcact tcacgggcat ggcgggagat	400
gccatgccta cttctgcac accagaaaac cactgtggaa cccacgcacc	450
tgtctggctc aatggcagcc accccctaga aggcgacggc attgtgcaac	500
gccaggcttg tgccagcttc aatgggaact gctgtctctg gaacaccacg	550
gtggaagtca aggcttgccc tggaggctac tatgtgtatc gtctgaccaa	600
gccagcgctc tgcttccacg tctactgtgg tcatttttat gacatctgcg	650
acgaggactg ccatggcagc tgctcagata ccagcgagtg cacatcgctc	700
ccaggaactg tgctaggccc tgacaggcag acatgctttg atgaaaatga	750
atgtgagcaa aacaacgggtg gctgcagtga gatctgtgtg aacctcaaaa	800
actcctaccg ctgtgagtgt ggggttggcc gtgtgctaag aagtgatggc	850
aagacttgtg aagacgttga aggatgccac aataacaatg gtggctgcag	900
ccactcttgc cttggatctg agaaaggcta ccagtgtgaa tgtccccggg	950
gcctggtgct gtctgaggat aaccacactt gccaaagccc tgtgttgtgc	1000
aaatcaaatg ccattgaagt gaacatcccc agggagctgg ttggtggcct	1050
ggagctcttc ctgaccaaca cctcctgcgg aggagtgtcc aacggcacc	1100
atgtcaacat cctcttctct ctcaagacat gtggtacagt ggtcgatgtg	1150
gtgaatgaca agattgtggc cagcaacctc gtgacaggtc tacccaagca	1200
gaccccgggg agcagcggg acttcatcat ccgaaccagc aagctgctga	1250
tcccggtgac ctgcgagttt ccacgcctgt acaccatttc tgaaggatac	1300
gttcccaacc ttcgaaactc cccactggaa atcatgagcc gaaatcatgg	1350
gatcttccca ttactctgg agatcttcaa ggacaatgag tttgaagagc	1400
cttaccggga agctctgccc accctcaagc ttcgtgactc cctctacttt	1450
ggcattgagc ccgtggtgca cgtgagcggc ttgaaaagct tggtgagag	1500
ctgctttgcc accccacct ccaagatcga cgaggctctg aaatactacc	1550
tcacccggga tggctgtgtt tcagatgact cggtaaagca gtacacatcc	1600
cggtatcacc tagcaaagca cttccaggto cctgtcttca agtttgtggg	1650
caaagaccac aaggaagtgt ttctgcactg ccgggttctt gtctgtggag	1700

-continued

tggtggacga gcgttccgc tgtgccagg gttgccaccg gcgaatgcgt	1750
cgtggggcag gaggagagga ctacagccgt ctacagggcc agacgctaac	1800
aggcgccccc atccgcacgc actgggagga ctagtctgta gccatacctc	1850
gagtcctgc attggacggc tctgctcttt ggagcttctc cccccaccgc	1900
cctctaagaa catctgcaa cagctgggtt cagacttcac actgtgagtt	1950
cagactccca gcaccaactc actctgattc tggtcattc agtgggcaca	2000
ggtcacagca ctgctgaaca atgtggcctg ggtggggttt catctttcta	2050
gggttgaaaa ctaaactgtc caccagaaa gacactcacc ccatttcct	2100
catttctttc ctacacttaa atacctcgtg tatggtgcaa tcagaccaca	2150
aaatcagaag ctgggtataa tatttcaagt tacaaccct agaaaaatta	2200
aacagttact gaaattatga cttaaatacc caatgactcc ttaaatatgt	2250
aaattatagt tataccttga aatttcaatt caaatgcaga ctaattatag	2300
ggaatttgga agtgtatcaa taaaacagta tataatttt	2339

<210> SEQ ID NO 110

<211> LENGTH: 545

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 110

Met	Pro	Pro	Phe	Leu	Leu	Leu	Thr	Cys	Leu	Phe	Ile	Thr	Gly	Thr
1				5					10					15
Ser	Val	Ser	Pro	Val	Ala	Leu	Asp	Pro	Cys	Ser	Ala	Tyr	Ile	Ser
				20					25					30
Leu	Asn	Glu	Pro	Trp	Arg	Asn	Thr	Asp	His	Gln	Leu	Asp	Glu	Ser
				35					40					45
Gln	Gly	Pro	Pro	Leu	Cys	Asp	Asn	His	Val	Asn	Gly	Glu	Trp	Tyr
				50					55					60
His	Phe	Thr	Gly	Met	Ala	Gly	Asp	Ala	Met	Pro	Thr	Phe	Cys	Ile
				65					70					75
Pro	Glu	Asn	His	Cys	Gly	Thr	His	Ala	Pro	Val	Trp	Leu	Asn	Gly
				80					85					90
Ser	His	Pro	Leu	Glu	Gly	Asp	Gly	Ile	Val	Gln	Arg	Gln	Ala	Cys
				95					100					105
Ala	Ser	Phe	Asn	Gly	Asn	Cys	Cys	Leu	Trp	Asn	Thr	Thr	Val	Glu
				110					115					120
Val	Lys	Ala	Cys	Pro	Gly	Gly	Tyr	Tyr	Val	Tyr	Arg	Leu	Thr	Lys
				125					130					135
Pro	Ser	Val	Cys	Phe	His	Val	Tyr	Cys	Gly	His	Phe	Tyr	Asp	Ile
				140					145					150
Cys	Asp	Glu	Asp	Cys	His	Gly	Ser	Cys	Ser	Asp	Thr	Ser	Glu	Cys
				155					160					165
Thr	Cys	Ala	Pro	Gly	Thr	Val	Leu	Gly	Pro	Asp	Arg	Gln	Thr	Cys
				170					175					180
Phe	Asp	Glu	Asn	Glu	Cys	Glu	Gln	Asn	Asn	Gly	Gly	Cys	Ser	Glu
				185					190					195
Ile	Cys	Val	Asn	Leu	Lys	Asn	Ser	Tyr	Arg	Cys	Glu	Cys	Gly	Val
				200					205					210

-continued

Gly	Arg	Val	Leu	Arg	Ser	Asp	Gly	Lys	Thr	Cys	Glu	Asp	Val	Glu	
			215						220					225	
Gly	Cys	His	Asn	Asn	Asn	Gly	Gly	Cys	Ser	His	Ser	Cys	Leu	Gly	
			230						235					240	
Ser	Glu	Lys	Gly	Tyr	Gln	Cys	Glu	Cys	Pro	Arg	Gly	Leu	Val	Leu	
			245						250					255	
Ser	Glu	Asp	Asn	His	Thr	Cys	Gln	Val	Pro	Val	Leu	Cys	Lys	Ser	
			260						265					270	
Asn	Ala	Ile	Glu	Val	Asn	Ile	Pro	Arg	Glu	Leu	Val	Gly	Gly	Leu	
			275						280					285	
Glu	Leu	Phe	Leu	Thr	Asn	Thr	Ser	Cys	Arg	Gly	Val	Ser	Asn	Gly	
			290						295					300	
Thr	His	Val	Asn	Ile	Leu	Phe	Ser	Leu	Lys	Thr	Cys	Gly	Thr	Val	
			305						310					315	
Val	Asp	Val	Val	Asn	Asp	Lys	Ile	Val	Ala	Ser	Asn	Leu	Val	Thr	
			320						325					330	
Gly	Leu	Pro	Lys	Gln	Thr	Pro	Gly	Ser	Ser	Gly	Asp	Phe	Ile	Ile	
			335						340					345	
Arg	Thr	Ser	Lys	Leu	Leu	Ile	Pro	Val	Thr	Cys	Glu	Phe	Pro	Arg	
			350						355					360	
Leu	Tyr	Thr	Ile	Ser	Glu	Gly	Tyr	Val	Pro	Asn	Leu	Arg	Asn	Ser	
			365						370					375	
Pro	Leu	Glu	Ile	Met	Ser	Arg	Asn	His	Gly	Ile	Phe	Pro	Phe	Thr	
			380						385					390	
Leu	Glu	Ile	Phe	Lys	Asp	Asn	Glu	Phe	Glu	Glu	Pro	Tyr	Arg	Glu	
			395						400					405	
Ala	Leu	Pro	Thr	Leu	Lys	Leu	Arg	Asp	Ser	Leu	Tyr	Phe	Gly	Ile	
			410						415					420	
Glu	Pro	Val	Val	His	Val	Ser	Gly	Leu	Glu	Ser	Leu	Val	Glu	Ser	
			425						430					435	
Cys	Phe	Ala	Thr	Pro	Thr	Ser	Lys	Ile	Asp	Glu	Val	Leu	Lys	Tyr	
			440						445					450	
Tyr	Leu	Ile	Arg	Asp	Gly	Cys	Val	Ser	Asp	Asp	Ser	Val	Lys	Gln	
			455						460					465	
Tyr	Thr	Ser	Arg	Asp	His	Leu	Ala	Lys	His	Phe	Gln	Val	Pro	Val	
			470						475					480	
Phe	Lys	Phe	Val	Gly	Lys	Asp	His	Lys	Glu	Val	Phe	Leu	His	Cys	
			485						490					495	
Arg	Val	Leu	Val	Cys	Gly	Val	Leu	Asp	Glu	Arg	Ser	Arg	Cys	Ala	
			500						505					510	
Gln	Gly	Cys	His	Arg	Arg	Met	Arg	Arg	Gly	Ala	Gly	Gly	Glu	Asp	
			515						520					525	
Ser	Ala	Gly	Leu	Gln	Gly	Gln	Thr	Leu	Thr	Gly	Gly	Pro	Ile	Arg	
			530						535					540	
Ile	Asp	Trp	Glu	Asp											
			545												

<210> SEQ ID NO 111

<211> LENGTH: 2063

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 111

-continued

gagagaggca gcagcttgct cagcggacaa ggatgctggg cgtgaggga	50
caaggcctgc cctgcactcg ggctcctcc agccagtgt gaccagggac	100
ttctgacctg ctggccagcc aggacctgtg tggggaggcc ctctgtctgc	150
cttgggggtga caatctcagc tccaggctac agggagaccg ggaggatcac	200
agagccagca tgttacagga tcctgacagt gatcaacctc tgaacagcct	250
cgatgtcaaa cccctgcgca aaccccgat ccccatggag accttcagaa	300
aggtggggat ccccatcatc atagcactac tgagcctggc gagtatcatc	350
attgtggttg tcctcatcaa ggtgattctg gataaatact acttcctctg	400
cgggcagcct ctccacttca tcccaggaa gcagctgtgt gacggagagc	450
tggactgtcc cttgggggag gacgaggagc actgtgtcaa gagcttcccc	500
gaagggcctg cagtggcagt ccgcctctcc aaggaccgat ccacactgca	550
ggtgctggac tcggccacag ggaactggtt ctctgcctgt ttcgacaact	600
tcacagaagc tctcgtgag acagcctgta ggcagatggg ctacagcaga	650
gctgtggaga ttggcccaga ccaggatctg gatgttgtt aaatcacaga	700
aaacagccag gagcttcgca tgcggaactc aagtggggcc tgtctctcag	750
gtcccttggt ctccctgcac tgtcttgct gtgggaagag cctgaagacc	800
ccccgtgtgg tgggtgggga ggaggcctct gtggattctt gcccttggca	850
ggtcagcatc cagtacgaca aacagcacgt ctgtggaggg agcatcctg	900
acccccactg ggtcctcacg gcagccact gcttcaggaa acataccgat	950
gtgttcaact ggaaggtgcg ggcaggctca gacaaactgg gcagcttccc	1000
atccctggct gtggccaaga tcatcatcat tgaattcaac cccatgtacc	1050
ccaaagacaa tgacatcgcc ctcatgaagc tgcagttccc actcactttc	1100
tcaggcacag tcaggcccat ctgtctgccc ttctttgatg aggagctcac	1150
tccagccacc ccactctgga tcattggatg gggctttacg aagcagaatg	1200
gagggaagat gtctgacata ctgtgcagg cgtcagttcca ggtcattgac	1250
agcacacggt gcaatgcaga cgatgcgtac cagggggaag tcaccgagaa	1300
gatgatgtgt gcaggcatcc cggaagggg tgtggacacc tgccaggggtg	1350
acagtgggtg gccctgatg taccaatctg accagtggca tgtgggtggc	1400
atcgttagct ggggctatgg ctgcgggggc ccgagacccc caggagtata	1450
caccaaggtc tcagcctatc tcaactggat ctacaatgtc tggaaggctg	1500
agctgtaatg ctgctgcccc ttgtcagtgc tgggagccgc ttccttctctg	1550
ccctgccac ctggggatcc cccaaagtca gacacagagc aagagtcccc	1600
ttgggtacac ccctctgccc acagcctcag catttcttgg agcagcaaag	1650
ggcctcaatt cctgtaagag accctcgag cccagaggcg cccagaggaa	1700
gtcagcagcc ctagctcggc cacacttggt gctcccagca tcccaggag	1750
agacacagcc cactgaacaa ggtctcagg gtattgctaa gccagaagg	1800
aactttccca cactactgaa tggaagcagg ctgtcttgta aaagcccaga	1850
tcactgtggg ctggagagga gaaggaaagg gtctgcgcca gccctgtccg	1900

-continued

tcttcaccca tccccaagcc tactagagca agaaaccagt tgtaatataa	1950
aatgcactgc cctactgttg gtatgactac cgttacctac tgttgtcatt	2000
gttattacag ctatggccac tattattaaa gagctgtgta acatctctgg	2050
caaaaaaaaaaaa aaa	2063

<210> SEQ ID NO 112

<211> LENGTH: 432

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 112

Met	Leu	Gln	Asp	Pro	Asp	Ser	Asp	Gln	Pro	Leu	Asn	Ser	Leu	Asp	1	5	10	15
Val	Lys	Pro	Leu	Arg	Lys	Pro	Arg	Ile	Pro	Met	Glu	Thr	Phe	Arg	20	25	30	
Lys	Val	Gly	Ile	Pro	Ile	Ile	Ile	Ala	Leu	Leu	Ser	Leu	Ala	Ser	35	40	45	
Ile	Ile	Ile	Val	Val	Leu	Ile	Lys	Val	Ile	Leu	Asp	Lys	Tyr	50	55	60		
Tyr	Phe	Leu	Cys	Gly	Gln	Pro	Leu	His	Phe	Ile	Pro	Arg	Lys	Gln	65	70	75	
Leu	Cys	Asp	Gly	Glu	Leu	Asp	Cys	Pro	Leu	Gly	Glu	Asp	Glu	Glu	80	85	90	
His	Cys	Val	Lys	Ser	Phe	Pro	Glu	Gly	Pro	Ala	Val	Ala	Val	Arg	95	100	105	
Leu	Ser	Lys	Asp	Arg	Ser	Thr	Leu	Gln	Val	Leu	Asp	Ser	Ala	Thr	110	115	120	
Gly	Asn	Trp	Phe	Ser	Ala	Cys	Phe	Asp	Asn	Phe	Thr	Glu	Ala	Leu	125	130	135	
Ala	Glu	Thr	Ala	Cys	Arg	Gln	Met	Gly	Tyr	Ser	Arg	Ala	Val	Glu	140	145	150	
Ile	Gly	Pro	Asp	Gln	Asp	Leu	Asp	Val	Val	Glu	Ile	Thr	Glu	Asn	155	160	165	
Ser	Gln	Glu	Leu	Arg	Met	Arg	Asn	Ser	Ser	Gly	Pro	Cys	Leu	Ser	170	175	180	
Gly	Ser	Leu	Val	Ser	Leu	His	Cys	Leu	Ala	Cys	Gly	Lys	Ser	Leu	185	190	195	
Lys	Thr	Pro	Arg	Val	Val	Gly	Gly	Glu	Glu	Ala	Ser	Val	Asp	Ser	200	205	210	
Trp	Pro	Trp	Gln	Val	Ser	Ile	Gln	Tyr	Asp	Lys	Gln	His	Val	Cys	215	220	225	
Gly	Gly	Ser	Ile	Leu	Asp	Pro	His	Trp	Val	Leu	Thr	Ala	Ala	His	230	235	240	
Cys	Phe	Arg	Lys	His	Thr	Asp	Val	Phe	Asn	Trp	Lys	Val	Arg	Ala	245	250	255	
Gly	Ser	Asp	Lys	Leu	Gly	Ser	Phe	Pro	Ser	Leu	Ala	Val	Ala	Lys	260	265	270	
Ile	Ile	Ile	Ile	Glu	Phe	Asn	Pro	Met	Tyr	Pro	Lys	Asp	Asn	Asp	275	280	285	
Ile	Ala	Leu	Met	Lys	Leu	Gln	Phe	Pro	Leu	Thr	Phe	Ser	Gly	Thr	290	295	300	

-continued

Val Arg Pro Ile Cys Leu Pro Phe Phe Asp Glu Glu Leu Thr Pro
305 310 315
Ala Thr Pro Leu Trp Ile Ile Gly Trp Gly Phe Thr Lys Gln Asn
320 325 330
Gly Gly Lys Met Ser Asp Ile Leu Leu Gln Ala Ser Val Gln Val
335 340 345
Ile Asp Ser Thr Arg Cys Asn Ala Asp Asp Ala Tyr Gln Gly Glu
350 355 360
Val Thr Glu Lys Met Met Cys Ala Gly Ile Pro Glu Gly Gly Val
365 370 375
Asp Thr Cys Gln Gly Asp Ser Gly Gly Pro Leu Met Tyr Gln Ser
380 385 390
Asp Gln Trp His Val Val Gly Ile Val Ser Trp Gly Tyr Gly Cys
395 400 405
Gly Gly Pro Ser Thr Pro Gly Val Tyr Thr Lys Val Ser Ala Tyr
410 415 420
Leu Asn Trp Ile Tyr Asn Val Trp Lys Ala Glu Leu
425 430

<210> SEQ ID NO 113
<211> LENGTH: 1768
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 113

ggctggactg gaactcctgg tcccaagtga tccacccgcc tcagcctccc 50
aagggtgctgt gattataggt gtaagccacc gtgtctggcc tctgaacaac 100
tttttcagca actaaaaaag ccacaggagt tgaactgcta ggattctgac 150
tatgtctgtg tggctagtgc tcctactcct acctacatta aaatctgttt 200
tttgtttctt tgtaactagc ctttaccttc ctaacacaga ggatctgtca 250
ctgtggctct ggcccaaacc tgaccttcac tctggaacga gaacagaggt 300
ttctacccac accgtcccct cgaagccggg gacagcctca ccttgtctggc 350
ctctcgctgg agcagtgcc tcaccaactg tctcacgtct ggaggcactg 400
actcgggcag tgcaggtagc tgagcctctt ggtagctgcg gctttcaagg 450
tgggccttgc cctggccgta gaagggattg acaagcccga agatttcata 500
ggcgatggct ccactgccc aggcatacag cttgctgtag tcaatcactg 550
ccctggggcc aggacgggcc gtggacacct gctcagaagc agtgggtgag 600
acatcacgct gccgcgccat ctaacctttt catgtcctgc acatcacctg 650
atccatgggc taatctgaac tctgtcccaa ggaaccaga gcttgagtga 700
gctgtggctc agaccagaa ggggtctgct tagaccacct ggtttatgtg 750
acaggacttg cattctcctg gaacatgagg gaacgccgga ggaagcaaa 800
gtggcagggg aggaacttgt gccaaattat gggtcagaaa agatggaggt 850
gttgggttat cacaaggcat cgagtctcct gcattcagtg gacatgtggg 900
ggaagggtcg ccgatggcg atgacacact cgggactcac ctctggggcc 950
atcagacagc cgtttccgcc ccgatccag taccagctgc tgaagggcaa 1000

-continued

ctgcaggccg atgctctcat cagccaggca gcagccaaaa tctgcatca	1050
ccagccaggg gcagccgtct gggaaggagc aagcaaagt accatttctc	1100
ctccctcct tccctctgag aggcctcct atgtccctac taaagccacc	1150
agcaagacat agctgacagg ggctaattggc tcagtgttg cccaggaggt	1200
cagcaaggcc tgagagctga tcagaagggc ctgctgtgcg aacacggaaa	1250
tgctccagt aagcacaggc tgcaaatcc ccaggcaaag gactgtgtgg	1300
ctcaatttaa atcatgttct agtaattgga gctgtcccca agaccaaagg	1350
agctagagct tggttcaaat gatctccaag ggcccttata cccaggaga	1400
ctttgatttg aatttgaaac cccaaatcca aacctaagaa ccagggtgat	1450
taagaatcag ttattgccgg gtgtggtggc ctgtaatgcc aacattttgg	1500
gaggccgagg cgggtagatc acctgaggtc aggagttaa gaccagcctg	1550
gccaacatgg tgaaccctt gtcttacta aaaatacaaa aaaactagcc	1600
aggcatggtg gtgtgtgcct gtatcccagc tactcgggag gctgagacag	1650
gagaattact tgaacctggg aggtgaagga ggctgagaca ggagaatcac	1700
ttcagcctga gcaacacagc gagactctgt ctcagaaaaa ataaaaaaag	1750
aattatggtt attttaa	1768

<210> SEQ ID NO 114

<211> LENGTH: 109

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 114

Met Leu Trp Trp Leu Val Leu Leu Leu Pro Thr Leu Lys Ser	
1 5 10 15	
Val Phe Cys Ser Leu Val Thr Ser Leu Tyr Leu Pro Asn Thr Glu	
20 25 30	
Asp Leu Ser Leu Trp Leu Trp Pro Lys Pro Asp Leu His Ser Gly	
35 40 45	
Thr Arg Thr Glu Val Ser Thr His Thr Val Pro Ser Lys Pro Gly	
50 55 60	
Thr Ala Ser Pro Cys Trp Pro Leu Ala Gly Ala Val Pro Ser Pro	
65 70 75	
Thr Val Ser Arg Leu Glu Ala Leu Thr Arg Ala Val Gln Val Ala	
80 85 90	
Glu Pro Leu Gly Ser Cys Gly Phe Gln Gly Gly Pro Cys Pro Gly	
95 100 105	
Arg Arg Arg Asp	

<210> SEQ ID NO 115

<211> LENGTH: 1197

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 115

cagcagtggc ctctcagtc tctcaaagca aggaaagagt actgtgtgct	50
gagagaccat ggcaaagaat cctccagaga attgtgaaga ctgtcacatt	100
ctaaatgcag aagcttttaa atccaagaaa atatgtaaat cacttaagat	150

-continued

```
ttgtggactg gtgtttggta tcctggccct aactctaatt gtccctgttt      200
gggggagcaa gcacttcttg ccggaggtag ccaaaaaagc ctatgacatg      250
gagcacactt tctacagcaa tggagagaag aagaagattt acatggaaat      300
tgatcctgtg accagaactg aaatattcag aagcggaat ggactgatg      350
aaacattgga agtgcacgac tttaaaaacg gatacactgg catctacttc      400
gtgggtcttc aaaaatgttt tatcaaaact cagattaaag tgattcctga      450
attttctgaa ccagaagagg aaatagatga gaatgaagaa attaccacaa      500
ctttctttga acagtcagtg atttgggtcc cagcagaaaa gcctattgaa      550
aaccgagatt ttcttaaaaa ttccaaaatt ctggagattt gtgataacgt      600
gaccatgtat tggatcaatc cactctaat atcagtttct gagttacaag      650
actttgagga ggaggagaa gatcttctact ttctgcca cgaaaaaaaaa      700
gggattgaac aaatgaaca gtgggtggtc cctcaagtga aagtagagaa      750
gacccgtagc gccagacaag caagtgagga agaacttcca ataatgact      800
atactgaaaa tggaatagaa ttgatccca tgctggatga gagaggttat      850
tgttgtattt actgccgtcg aggcaaccgc tattgccgcc gcgtctgtga      900
acctttacta ggctactacc catatccata ctgctaccaa ggaggacgag      950
tcactgtctg tgtcatcatg ccttgtaact ggtgggtggc ccgatgctg     1000
gggagggtct aataggaggt ttgagctcaa atgcttaaac tgctggcaac     1050
atataataaa tgcatgctat tcaatgaatt tctgcctatg aggcactctg     1100
ccccgtgtag ccagctctcc agaattactt gtaggtaatt cctctcttca     1150
tgttctaata aacttctaca ttatcaccaa aaaaaaaaaa aaaaaaa      1197
```

<210> SEQ ID NO 116
<211> LENGTH: 317
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 116

```
Met Ala Lys Asn Pro Pro Glu Asn Cys Glu Asp Cys His Ile Leu
 1          5          10         15
Asn Ala Glu Ala Phe Lys Ser Lys Lys Ile Cys Lys Ser Leu Lys
 20         25         30
Ile Cys Gly Leu Val Phe Gly Ile Leu Ala Leu Thr Leu Ile Val
 35         40         45
Leu Phe Trp Gly Ser Lys His Phe Trp Pro Glu Val Pro Lys Lys
 50         55         60
Ala Tyr Asp Met Glu His Thr Phe Tyr Ser Asn Gly Glu Lys Lys
 65         70         75
Lys Ile Tyr Met Glu Ile Asp Pro Val Thr Arg Thr Glu Ile Phe
 80         85         90
Arg Ser Gly Asn Gly Thr Asp Glu Thr Leu Glu Val His Asp Phe
 95        100       105
Lys Asn Gly Tyr Thr Gly Ile Tyr Phe Val Gly Leu Gln Lys Cys
110       115       120
Phe Ile Lys Thr Gln Ile Lys Val Ile Pro Glu Phe Ser Glu Pro
```

-continued

	125		130		135
Glu Glu Glu Ile Asp Glu Asn Glu Glu Ile Thr Thr Thr Phe Phe	140		145		150
Glu Gln Ser Val Ile Trp Val Pro Ala Glu Lys Pro Ile Glu Asn	155		160		165
Arg Asp Phe Leu Lys Asn Ser Lys Ile Leu Glu Ile Cys Asp Asn	170		175		180
Val Thr Met Tyr Trp Ile Asn Pro Thr Leu Ile Ser Val Ser Glu	185		190		195
Leu Gln Asp Phe Glu Glu Glu Gly Glu Asp Leu His Phe Pro Ala	200		205		210
Asn Glu Lys Lys Gly Ile Glu Gln Asn Glu Gln Trp Val Val Pro	215		220		225
Gln Val Lys Val Glu Lys Thr Arg His Ala Arg Gln Ala Ser Glu	230		235		240
Glu Glu Leu Pro Ile Asn Asp Tyr Thr Glu Asn Gly Ile Glu Phe	245		250		255
Asp Pro Met Leu Asp Glu Arg Gly Tyr Cys Cys Ile Tyr Cys Arg	260		265		270
Arg Gly Asn Arg Tyr Cys Arg Arg Val Cys Glu Pro Leu Leu Gly	275		280		285
Tyr Tyr Pro Tyr Pro Tyr Cys Tyr Gln Gly Gly Arg Val Ile Cys	290		295		300
Arg Val Ile Met Pro Cys Asn Trp Trp Val Ala Arg Met Leu Gly	305		310		315

Arg Val

<210> SEQ ID NO 117
<211> LENGTH: 2121
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 117

gagctccct caggagcgcg ttagcttcac accttcggca gcaggagggc	50
ggcagcttct cgcaggcggc agggcggcg gccaggatca tgtccaccac	100
cacatgccaa gtggtggcgt tcctcctgtc catcctgggg ctggccggct	150
gcacgcggc caccgggatg gacatgtgga gcaccagga cctgtacgac	200
aaccccgta cctccgtgtt ccagtacgaa gggctctgga ggagctgcgt	250
gaggcagagt tcaggcttca ccgaatgcag gccctatttc accatcctgg	300
gacttcacg catgctgcag gcagtgcgag ccctgatgat cgtaggcatc	350
gtcctgggtg ccattggcct cctggtatcc atctttgccc tgaatgcata	400
ccgcattggc agcatggagg actctgccaa agccaacatg aactgacct	450
ccgggatcat gttcattgtc tcaggctctt gtgcaattgc tggagtgtct	500
gtgtttgcc aatgctggt gactaacttc tggatgtcca cagctaacat	550
gtacaccggc atgggtggga tggcgcagac tgttcagacc aggtacacat	600
ttggtgcggc tctgttcgtg ggctgggtcg ctggaggcct cacactaatt	650
gggggtgtga tgatgtgcat cgctgccgg ggctggcac cagaagaaac	700

-continued

```

caactacaaa gccgtttctt atcatgcctc aggccacagt gttgcctaca      750
agcctggagg cttcaaggcc agcactggct ttgggtccaa caccaaaaac      800
aagaagatat acgatggagg tgcccgcaca gaggacgagg tacaatctta      850
tccttccaag cagcactatg tgtaatgctc taagacctct cagcacgggc      900
ggaagaaact cccggagagc tcacccaaaa aacaaggaga tcccatctag      950
atcttctctt gcttttgact cacagctgga agttagaaaa gcctcgattt     1000
catctttgga gaggccaaat ggtcttagcc tcagtctctg tctctaaata     1050
ttccaccata aaacagctga gttatttatg aattagaggc tatagctcac     1100
attttcaatc ctctatttct ttttttaaataaatactttct actctgatga     1150
gagaatgtgg ttttaatctc tctctcacat tttgatgatt tagacagact     1200
ccccctcttc ctctagtca ataaacccat tgatgatcta tttccagct      1250
tatccccaag aaaacttttg aaaggaaaga gtagacccaa agatgttatt     1300
ttctgctgtt tgaattttgt ctccccacc ccaacttggc tagtaataaa     1350
cacttactga agaagaagca ataagagaaa gatatttgta atctctccag     1400
cccatgatct cggttttctt aactgtgat cttaaaagtt accaaaccaa     1450
agtcattttc agtttgaggc aaccaaacct ttctactgct gttgacatct     1500
tcttattaca gcaacaccat tctaggagtt tcctgagctc tccactggag     1550
tcctctttct gtcgcggggtc agaaattgtc cctagatgaa tgagaaaatt     1600
atttttttta atttaagtcc taaatatagt taaaataaat aatgttttag     1650
taaaatgata cactatctct gtgaaatagc ctcacccta catgtggata     1700
gaaggaaatg aaaaaataat tgctttgaca ttgtctatat ggtactttgt     1750
aaagtcatgc ttaagtacaa attccatgaa aagctcacac ctgtaatcct     1800
agcactttgg gaggtgagg aggaaggatc acttgagccc agaagttcga     1850
gactagcctg ggcaacatgg agaagccctg tctctacaaa atacagagag     1900
aaaaaatcag ccagtcatgg tggcatcacac ctgtagtccc agcattccgg     1950
gaggctgagg tgggaggatc acttgagccc agggaggttg gggctgcagt     2000
gagccatgat cacaccactg cactccagcc aggtgacata gcgagatcct     2050
gtctaaaaaa ataaaaaata aataatggaa cacagcaagt cctaggaagt     2100
aggttaaaac taattcttta a                                     2121

```

<210> SEQ ID NO 118

<211> LENGTH: 261

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 118

```

Met Ser Thr Thr Thr Cys Gln Val Val Ala Phe Leu Leu Ser Ile
 1           5           10          15
Leu Gly Leu Ala Gly Cys Ile Ala Ala Thr Gly Met Asp Met Trp
          20          25          30
Ser Thr Gln Asp Leu Tyr Asp Asn Pro Val Thr Ser Val Phe Gln
          35          40          45
Tyr Glu Gly Leu Trp Arg Ser Cys Val Arg Gln Ser Ser Gly Phe

```

-continued

50					55					60				
Thr	Glu	Cys	Arg	Pro 65	Tyr	Phe	Thr	Ile	Leu 70	Gly	Leu	Pro	Ala	Met 75
Leu	Gln	Ala	Val	Arg 80	Ala	Leu	Met	Ile	Val 85	Gly	Ile	Val	Leu	Gly 90
Ala	Ile	Gly	Leu	Leu 95	Val	Ser	Ile	Phe	Ala 100	Leu	Lys	Cys	Ile	Arg 105
Ile	Gly	Ser	Met	Glu 110	Asp	Ser	Ala	Lys	Ala 115	Asn	Met	Thr	Leu	Thr 120
Ser	Gly	Ile	Met	Phe 125	Ile	Val	Ser	Gly	Leu 130	Cys	Ala	Ile	Ala	Gly 135
Val	Ser	Val	Phe	Ala 140	Asn	Met	Leu	Val	Thr 145	Asn	Phe	Trp	Met	Ser 150
Thr	Ala	Asn	Met	Tyr 155	Thr	Gly	Met	Gly	Gly 160	Met	Val	Gln	Thr	Val 165
Gln	Thr	Arg	Tyr	Thr 170	Phe	Gly	Ala	Ala	Leu 175	Phe	Val	Gly	Trp	Val 180
Ala	Gly	Gly	Leu	Thr 185	Leu	Ile	Gly	Gly	Val 190	Met	Met	Cys	Ile	Ala 195
Cys	Arg	Gly	Leu	Ala 200	Pro	Glu	Glu	Thr	Asn 205	Tyr	Lys	Ala	Val	Ser 210
Tyr	His	Ala	Ser	Gly 215	His	Ser	Val	Ala	Tyr 220	Lys	Pro	Gly	Gly	Phe 225
Lys	Ala	Ser	Thr	Gly 230	Phe	Gly	Ser	Asn	Thr 235	Lys	Asn	Lys	Lys	Ile 240
Tyr	Asp	Gly	Gly	Ala 245	Arg	Thr	Glu	Asp	Glu 250	Val	Gln	Ser	Tyr	Pro 255
Ser	Lys	His	Asp	Tyr 260	Val									

<210> SEQ ID NO 119
<211> LENGTH: 2010
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 119

ggaaaaactg ttctcttctg tggcacagag aaccctgctt caaagcagaa	50
gtagcagttc cggagtccag ctggctaaaa ctcatcccag aggataatgg	100
caacccatgc cttagaaatc gctgggctgt ttcttggtgg tgttggaatg	150
gtgggcacag tggctgtcac tgtcatgcct cagtggagag tgcggcctt	200
cattgaaaac aacatcgtgg tttttgaaaa cttctgggaa ggactgtgga	250
tgaattgcgt gaggcaggct aacatcagga tgcagtgcaa aatctatgat	300
tccctgctgg ctctttctcc ggacctacag gcagccagag gactgatgtg	350
tgctgcttcc gtgatgtcct tcttggtttt catgatggcc atccttgcca	400
tgaaatgcac cagtgacag ggggacaatg agaaggtgaa ggctcacatt	450
ctgctgacgg ctggaatcat cttcatcatc acgggcatgg tgggtgctcat	500
ccctgtgagc tgggttgcca atgccatcat cagagatttc tataactcaa	550
tagtgaatgt tgcccaaaaa cgtgagcttg gagaagctct ctacttagga	600

-continued

tggaccacgg cactggtgct gattgttgga ggagctctgt tctgctgcgt	650
tttttgttgc aacgaaaaga gcagtagcta cagatactcg ataccttccc	700
atcgcacaac ccaaaaaagt tatcacaccg gaaagaagtc accgagcgtc	750
tactccagaa gtcagtatgt gtagttgtgt atgttttttt aactttacta	800
taaagccatg caaatgacaa aaatctatat tactttctca aaatggaccc	850
caaagaaact ttgatttact gttcttaact gcctaactctt aattacagga	900
actgtgcacg agctatttat gattctataa gctatttcag cagaatgaga	950
tattaaaccg aatgctttga ttgttctaga aagtatagta atttgttttc	1000
taagtggtgt caagcatcta ctctttttat catttacttc aaaatgacat	1050
tgctaaagac tgcattatgt tactactgta atttctccac gacatagcat	1100
tatgtacata gatgagtgtg acatttatat ctcacataga gacatgctta	1150
tatgggttta tttaaaatga aatgccagtc cattacactg aataaataga	1200
actcaactat tgcttttcag ggaaatcatg gatagggttg aagaaggtta	1250
ctattaattg tttaaaaaca gcttagggat taatgtcctc cattttataat	1300
gaagattaaa atgaaggctt taatcagcat tgtaaaggaa attgaatggc	1350
tttctgatat gctgtttttt agcctaggag ttagaaatcc taacttcttt	1400
atcctcttct cccagaggct ttttttttct tgtgtattaa attaacattt	1450
ttaaaacgca gatattttgt caaggggctt tgcattcaaa ctgcttttcc	1500
agggctatata tcagaagaaa gataaaagtg tgatctaaga aaaagtgatg	1550
gttttaggaa agtgaaaata tttttgtttt tgtatttgaa gaagaatgat	1600
gcattttgac aagaaatcat atatgtatgg atatatttta ataagtattt	1650
gagtacagac tttgaggttt catcaatata aataaaagag cagaaaaata	1700
tgtcttggtt ttcatttgct taccaaaaaa acaacaacaa aaaaagttgt	1750
cccttgagaa cttcacctgc tcctatgtgg gtacctgagt caaaattgtc	1800
atttttgttc tgtgaaaaat aaatttcctt cttgtacat ttctgtttag	1850
ttttactaaa atctgtaaat actgtatttt tctgtttatt ccaaatttga	1900
tgaaactgac aatccaattt gaaagtttgt gtcgacgtct gtctagctta	1950
aatgaatgtg ttctatttgc tttatacatt tatattaata aattgtacat	2000
ttttctaatt	2010

<210> SEQ ID NO 120
<211> LENGTH: 225
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 120

Met Ala Thr His Ala Leu Glu Ile Ala Gly Leu Phe Leu Gly Gly
1 5 10 15

Val Gly Met Val Gly Thr Val Ala Val Thr Val Met Pro Gln Trp
20 25 30

Arg Val Ser Ala Phe Ile Glu Asn Asn Ile Val Val Phe Glu Asn
35 40 45

Phe Trp Glu Gly Leu Trp Met Asn Cys Val Arg Gln Ala Asn Ile

-continued

50					55					60				
Arg	Met	Gln	Cys	Lys	Ile	Tyr	Asp	Ser	Leu	Leu	Ala	Leu	Ser	Pro
65					70					75				
Asp	Leu	Gln	Ala	Ala	Arg	Gly	Leu	Met	Cys	Ala	Ala	Ser	Val	Met
80					85					90				
Ser	Phe	Leu	Ala	Phe	Met	Met	Ala	Ile	Leu	Gly	Met	Lys	Cys	Thr
95					100					105				
Arg	Cys	Thr	Gly	Asp	Asn	Glu	Lys	Val	Lys	Ala	His	Ile	Leu	Leu
110					115					120				
Thr	Ala	Gly	Ile	Ile	Phe	Ile	Ile	Thr	Gly	Met	Val	Val	Leu	Ile
125					130					135				
Pro	Val	Ser	Trp	Val	Ala	Asn	Ala	Ile	Ile	Arg	Asp	Phe	Tyr	Asn
140					145					150				
Ser	Ile	Val	Asn	Val	Ala	Gln	Lys	Arg	Glu	Leu	Gly	Glu	Ala	Leu
155					160					165				
Tyr	Leu	Gly	Trp	Thr	Thr	Ala	Leu	Val	Leu	Ile	Val	Gly	Gly	Ala
170					175					180				
Leu	Phe	Cys	Cys	Val	Phe	Cys	Cys	Asn	Glu	Lys	Ser	Ser	Ser	Tyr
185					190					195				
Arg	Tyr	Ser	Ile	Pro	Ser	His	Arg	Thr	Thr	Gln	Lys	Ser	Tyr	His
200					205					210				
Thr	Gly	Lys	Lys	Ser	Pro	Ser	Val	Tyr	Ser	Arg	Ser	Gln	Tyr	Val
215					220					225				

<210> SEQ ID NO 121
<211> LENGTH: 1257
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 121

ggagagagggc ggcgcgggtga aaggcgcatc gatgcagcct gcggcggcct	50
cggagcgcgcg cggagccaga cgtcgaccac gttcctctcc tcggtctcct	100
ccgcctccag ctccgcgctg cccggcagcc gggagccatg cgacccagc	150
gccccgcgcg cccccgcag cggtccgcg gcctcctgct gctcctgctg	200
ctgcagctgc ccgcgcgctc gagcgctct gagatcccca aggggaagca	250
aaagcgcgag ctccggcaga gggaggtggt ggacctgtat aatggaatgt	300
gcttacaagg gccagcagga gtgcctggtc gagacgggag ccctggggcc	350
aatgttatcc cgggtacacc tgggatccca ggtcgggatg gattcaaagg	400
agaaaagggg gaatgtctga gggaaagctt tgaggagtcc tggacaccca	450
actacaagca gtgttcattg agttcattga attatggcat agatcttggg	500
aaaattgcgg agtgtacatt tacaaagatg cgttcaaata gtgctctaag	550
agttttgttc agtggctcac ttcggctaaa atgcagaaat gcatgctgtc	600
agcgttggtg ttccacattc aatggagctg aatgttcagg acctcttccc	650
attgaagcta taatttatcc ggaccaagga agccctgaaa tgaattcaac	700
aattaatatc catcgcaact cttctgtgga aggactttgt gaaggaattg	750
gtgctggatt agtggatggt gctatctggg ttggcacttg ttcagattac	800
ccaaaaggag atgcttctac tggatggaat tcagtttctc gcatcattat	850

-continued

tgaagaacta ccaaaataaa tgctttaatt ttcatttgct acctcttttt	900
ttattatgcc ttggaatggt tcacttaaat gacattttta ataagtttat	950
gtatacatct gaatgaaaag caaagctaaa tatgtttaca gaccaaagtg	1000
tgatttcaca ctgtttttta atctagcatt attcattttg cttcaatcaa	1050
aagtgggttc aatatttttt ttagttgggtt agaatacttt cttcatagtc	1100
acattctctc aacctataat ttggaatatt gttgtggtct tttgtttttt	1150
ctcttagtat agcattttta aaaaaatata aaagctacca atctttgtac	1200
aatttgtaaa tgtaagaat tttttttata tctgttaaataaaaaattatt	1250
tccaaca	1257

<210> SEQ ID NO 122

<211> LENGTH: 243

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 122

Met	Arg	Pro	Gln	Gly	Pro	Ala	Ala	Ser	Pro	Gln	Arg	Leu	Arg	Gly	1	5	10	15
Leu	Leu	Leu	Leu	Leu	Leu	Leu	Gln	Leu	Pro	Ala	Pro	Ser	Ser	Ala	20	25	30	
Ser	Glu	Ile	Pro	Lys	Gly	Lys	Gln	Lys	Ala	Gln	Leu	Arg	Gln	Arg	35	40	45	
Glu	Val	Val	Asp	Leu	Tyr	Asn	Gly	Met	Cys	Leu	Gln	Gly	Pro	Ala	50	55	60	
Gly	Val	Pro	Gly	Arg	Asp	Gly	Ser	Pro	Gly	Ala	Asn	Val	Ile	Pro	65	70	75	
Gly	Thr	Pro	Gly	Ile	Pro	Gly	Arg	Asp	Gly	Phe	Lys	Gly	Glu	Lys	80	85	90	
Gly	Glu	Cys	Leu	Arg	Glu	Ser	Phe	Glu	Glu	Ser	Trp	Thr	Pro	Asn	95	100	105	
Tyr	Lys	Gln	Cys	Ser	Trp	Ser	Ser	Leu	Asn	Tyr	Gly	Ile	Asp	Leu	110	115	120	
Gly	Lys	Ile	Ala	Glu	Cys	Thr	Phe	Thr	Lys	Met	Arg	Ser	Asn	Ser	125	130	135	
Ala	Leu	Arg	Val	Leu	Phe	Ser	Gly	Ser	Leu	Arg	Leu	Lys	Cys	Arg	140	145	150	
Asn	Ala	Cys	Cys	Gln	Arg	Trp	Tyr	Phe	Thr	Phe	Asn	Gly	Ala	Glu	155	160	165	
Cys	Ser	Gly	Pro	Leu	Pro	Ile	Glu	Ala	Ile	Ile	Tyr	Leu	Asp	Gln	170	175	180	
Gly	Ser	Pro	Glu	Met	Asn	Ser	Thr	Ile	Asn	Ile	His	Arg	Thr	Ser	185	190	195	
Ser	Val	Glu	Gly	Leu	Cys	Glu	Gly	Ile	Gly	Ala	Gly	Leu	Val	Asp	200	205	210	
Val	Ala	Ile	Trp	Val	Gly	Thr	Cys	Ser	Asp	Tyr	Pro	Lys	Gly	Asp	215	220	225	
Ala	Ser	Thr	Gly	Trp	Asn	Ser	Val	Ser	Arg	Ile	Ile	Ile	Glu	Glu	230	235	240	
Leu	Pro	Lys																

-continued

<210> SEQ ID NO 123

<211> LENGTH: 2379

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 123

gctgagcgtg tgcgcggtac ggggctctcc tgccttctgg gtcceaacgc	50
agctctgtgg ctgaactggg tgctcatcac gggaactgct gggctatgga	100
atacagatgt ggcagctcag gtagcccaa attgcctgga agaatacatc	150
atgtttttcg ataagaagaa attgtaggat ccagtttttt ttttaaccgc	200
cccccccca cccccaaaa aaactgtaaa gatgcaaaaa cgtaatatcc	250
atgaagatcc tattacntag gaagattttg atgttttgct gcgaatgcgg	300
tggtgggatt tatttgttct tggagtgttc tgcgtggctg gcaaagaata	350
atgttccaaa atcgggtccat ctccaagggt gtccaatttt tcttcctggg	400
tgtcagcgag ccctgactca ctacagtgca gctgacaggg gctgtcatgc	450
aactggcccc taagccaaag caaaagacct aaggacgacc tttgaacaat	500
acaaaggatg ggtttcaatg taattaggct actgagcgga tcagctgtag	550
cactggttat agccccact gtcttactga caatgctttc tctgccgaa	600
cgaggatgoc ctaagggctg taggtgtgaa ggcaaatgg tatattgtga	650
atctcagaaa ttacaggaga taccctcaag tatactctgct ggttgcttag	700
gtttgtccct tcgctataac agccttcaaa aacttaagta taatcaattt	750
aaagggctca accagctcac ctggctatac cttgaccata accatatcag	800
caatattgac gaaaatgctt ttaatggaat acgcagactc aaagagctga	850
ttcttagttc caatagaatc tcctattttc ttaacaatac cttcagacct	900
gtgacaaatt tacggaactt ggaatgtgcc tataatcagc tgcattctct	950
gggatctgaa cagtttcggg gcttgcgga gctgctgagt ttacatttac	1000
ggcttaactc cctgagaacc atccctgtgc gaatattcca agactgccgc	1050
aacctggaac ttttgacctt gggatataac cggatccgaa gtttagccag	1100
gaatgtcttt gctggcatga tcagactcaa agaacttcac ctggagcaca	1150
atcaattttc caagctcaac ctggcccttt ttccaagggt ggtcagcctt	1200
cagaaccttt acttgcagtg gaataaaatc agtgtcatag gacagacct	1250
gtcctggacc tggagctcct tacaaaggct tgatttatca ggcaatgaga	1300
tcgaagcttt cagtggacct agtgttttcc agtgtgtccc gaatctgcag	1350
cgcctcaacc tggattccaa caagctcaca tttattggtc aagagatttt	1400
ggattcttgg atatccctca atgacatcag tcttgctggg aatatatggg	1450
aatgcagcag aaatatttgc tcccttgtaa actggctgaa aagttttaaa	1500
ggctaaaggg agaatacaat tatctgtgcc agtcccaaag agctgcaagg	1550
agtaaatgtg atcgatgcag tgaagaacta cagcatctgt ggcaaaagta	1600
ctacagagag gtttgatctg gccagggtcc tcccaaagcc gacgtttaag	1650
cccaagctcc ccaggccgaa gcatgagagc aaacccccctt tgcccccgac	1700

-continued

```

ggtgggagcc acagagcccg gcccagagac cgatgctgac gccgagcaca      1750
tctctttcca taaaatcatc gcgggcagcg tggcgctttt cctgtccgtg      1800
ctcgtcatcc tgctgggttat ctacgtgtca tggaagcggg accctgcgag      1850
catgaagcag ctgcagcagc gctccctcat gcgaaggcac aggaaaaaga      1900
aaagacagtc cctaaagcaa atgactccca gcaccagga attttatgta      1950
gattataaac ccaccaacac ggagaccagc gagatgctgc tgaatgggac      2000
gggaccctgc acctataaca aatcgggctc cagggagtgt gaggtatgaa      2050
ccattgtgat aaaaagagct cttaaagctc gggaaataag tgggtgcttta      2100
ttgaactctg gtgactatca agggaacgcg atgccccccc tccccctccc      2150
tctccctctc actttggtgg caagatcctt ccttgccgtt tttagtgcatt      2200
tcataatact ggtcattttc ctctcataca taatcaaccc attgaaattt      2250
aaataccaca atcaatgtga agcttgaact cgggtttaat ataataccta      2300
ttgtataaga ccctttactg attccattaa tgcgcattt gttttaagat      2350
aaaacttctt tcataggtaa aaaaaaaaaa      2379

```

<210> SEQ ID NO 124

<211> LENGTH: 513

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 124

```

Met Gly Phe Asn Val Ile Arg Leu Leu Ser Gly Ser Ala Val Ala
 1             5             10            15
Leu Val Ile Ala Pro Thr Val Leu Leu Thr Met Leu Ser Ser Ala
          20            25            30
Glu Arg Gly Cys Pro Lys Gly Cys Arg Cys Glu Gly Lys Met Val
          35            40            45
Tyr Cys Glu Ser Gln Lys Leu Gln Glu Ile Pro Ser Ser Ile Ser
          50            55            60
Ala Gly Cys Leu Gly Leu Ser Leu Arg Tyr Asn Ser Leu Gln Lys
          65            70            75
Leu Lys Tyr Asn Gln Phe Lys Gly Leu Asn Gln Leu Thr Trp Leu
          80            85            90
Tyr Leu Asp His Asn His Ile Ser Asn Ile Asp Glu Asn Ala Phe
          95          100          105
Asn Gly Ile Arg Arg Leu Lys Glu Leu Ile Leu Ser Ser Asn Arg
          110          115          120
Ile Ser Tyr Phe Leu Asn Asn Thr Phe Arg Pro Val Thr Asn Leu
          125          130          135
Arg Asn Leu Asp Leu Ser Tyr Asn Gln Leu His Ser Leu Gly Ser
          140          145          150
Glu Gln Phe Arg Gly Leu Arg Lys Leu Leu Ser Leu His Leu Arg
          155          160          165
Ser Asn Ser Leu Arg Thr Ile Pro Val Arg Ile Phe Gln Asp Cys
          170          175          180
Arg Asn Leu Glu Leu Leu Asp Leu Gly Tyr Asn Arg Ile Arg Ser
          185          190          195

```

-continued

Leu	Ala	Arg	Asn	Val	Phe	Ala	Gly	Met	Ile	Arg	Leu	Lys	Glu	Leu
			200						205					210
His	Leu	Glu	His	Asn	Gln	Phe	Ser	Lys	Leu	Asn	Leu	Ala	Leu	Phe
			215						220					225
Pro	Arg	Leu	Val	Ser	Leu	Gln	Asn	Leu	Tyr	Leu	Gln	Trp	Asn	Lys
			230						235					240
Ile	Ser	Val	Ile	Gly	Gln	Thr	Met	Ser	Trp	Thr	Trp	Ser	Ser	Leu
			245						250					255
Gln	Arg	Leu	Asp	Leu	Ser	Gly	Asn	Glu	Ile	Glu	Ala	Phe	Ser	Gly
			260						265					270
Pro	Ser	Val	Phe	Gln	Cys	Val	Pro	Asn	Leu	Gln	Arg	Leu	Asn	Leu
			275						280					285
Asp	Ser	Asn	Lys	Leu	Thr	Phe	Ile	Gly	Gln	Glu	Ile	Leu	Asp	Ser
			290						295					300
Trp	Ile	Ser	Leu	Asn	Asp	Ile	Ser	Leu	Ala	Gly	Asn	Ile	Trp	Glu
			305						310					315
Cys	Ser	Arg	Asn	Ile	Cys	Ser	Leu	Val	Asn	Trp	Leu	Lys	Ser	Phe
			320						325					330
Lys	Gly	Leu	Arg	Glu	Asn	Thr	Ile	Ile	Cys	Ala	Ser	Pro	Lys	Glu
			335						340					345
Leu	Gln	Gly	Val	Asn	Val	Ile	Asp	Ala	Val	Lys	Asn	Tyr	Ser	Ile
			350						355					360
Cys	Gly	Lys	Ser	Thr	Thr	Glu	Arg	Phe	Asp	Leu	Ala	Arg	Ala	Leu
			365						370					375
Pro	Lys	Pro	Thr	Phe	Lys	Pro	Lys	Leu	Pro	Arg	Pro	Lys	His	Glu
			380						385					390
Ser	Lys	Pro	Pro	Leu	Pro	Pro	Thr	Val	Gly	Ala	Thr	Glu	Pro	Gly
			395						400					405
Pro	Glu	Thr	Asp	Ala	Asp	Ala	Glu	His	Ile	Ser	Phe	His	Lys	Ile
			410						415					420
Ile	Ala	Gly	Ser	Val	Ala	Leu	Phe	Leu	Ser	Val	Leu	Val	Ile	Leu
			425						430					435
Leu	Val	Ile	Tyr	Val	Ser	Trp	Lys	Arg	Tyr	Pro	Ala	Ser	Met	Lys
			440						445					450
Gln	Leu	Gln	Gln	Arg	Ser	Leu	Met	Arg	Arg	His	Arg	Lys	Lys	Lys
			455						460					465
Arg	Gln	Ser	Leu	Lys	Gln	Met	Thr	Pro	Ser	Thr	Gln	Glu	Phe	Tyr
			470						475					480
Val	Asp	Tyr	Lys	Pro	Thr	Asn	Thr	Glu	Thr	Ser	Glu	Met	Leu	Leu
			485						490					495
Asn	Gly	Thr	Gly	Pro	Cys	Thr	Tyr	Asn	Lys	Ser	Gly	Ser	Arg	Glu
			500						505					510
Cys	Glu	Val												

<210> SEQ ID NO 125
<211> LENGTH: 998
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 125

ccgttatcgt cttgcgctac tgctgaatgt ccgtcccgga ggaggaggag	50
aggccttttcg cgctgaccga gagatggccc cgagcgagca aattcctact	100

-continued

gtccggctgc gcggtaccg tggccgagct agcaaccttt cccctggatc	150
tcacaaaaac tcgactccaa atgcaaggag aagcagctct tgctcggttg	200
ggagacggtg caagagaatc tgccccctat aggggaatgg tgcgcacagc	250
cctagggatc attgaagagg aaggctttct aaagctttgg caaggagtga	300
caccgcgatc ttacagacac gtagtgatt ctggaggtcg aatggtcaca	350
tatgaacatc tccgagaggt tgtgtttggc aaaagtgaag atgagcatta	400
ttccctttgg aaatcagtca ttggagggat gatggctggt gttattggcc	450
agtttttagc caatccaact gacctagtga aggttcagat gcaaatggaa	500
ggaaaaagga aactggaagg aaaaccattg cgatttcgtg gtgtacatca	550
tgcatttgca aaaatcttag ctgaaggagg aatacggagg ctttgggcag	600
gctgggtacc caatatacaa agagcagcac tggatgaatat gggagattta	650
accacttatg atacagtga acactacttg gtattgaata caccacttga	700
ggacaatatc atgactcacg gtttatcaag tttatgttct ggactggtag	750
cttctattct gggaacacca gccgatgtca tcaaagcag aataatgaat	800
caaccacgag ataacaagg aaggggactt ttgtataaat catcgactga	850
ctgcttgatt caggctgttc aaggtgaagg attcatgagt ctatataaag	900
gctttttacc atcttggctg agaatgacct ctgtgtcaat ggtgttctgg	950
cttacttatg aaaaaatcag agagatgagt ggagtcagtc cattttaa	998

<210> SEQ ID NO 126
<211> LENGTH: 323
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 126

Met Ser Val Pro Glu Glu Glu Arg Leu Leu Pro Leu Thr Gln	1 5 10 15
Arg Trp Pro Arg Ala Ser Lys Phe Leu Leu Ser Gly Cys Ala Ala	20 25 30
Thr Val Ala Glu Leu Ala Thr Phe Pro Leu Asp Leu Thr Lys Thr	35 40 45
Arg Leu Gln Met Gln Gly Glu Ala Ala Leu Ala Arg Leu Gly Asp	50 55 60
Gly Ala Arg Glu Ser Ala Pro Tyr Arg Gly Met Val Arg Thr Ala	65 70 75
Leu Gly Ile Ile Glu Glu Glu Gly Phe Leu Lys Leu Trp Gln Gly	80 85 90
Val Thr Pro Ala Ile Tyr Arg His Val Val Tyr Ser Gly Gly Arg	95 100 105
Met Val Thr Tyr Glu His Leu Arg Glu Val Val Phe Gly Lys Ser	110 115 120
Glu Asp Glu His Tyr Pro Leu Trp Lys Ser Val Ile Gly Gly Met	125 130 135
Met Ala Gly Val Ile Gly Gln Phe Leu Ala Asn Pro Thr Asp Leu	140 145 150
Val Lys Val Gln Met Gln Met Glu Gly Lys Arg Lys Leu Glu Gly	

-continued

				155					160					165
Lys	Pro	Leu	Arg	Phe 170	Arg	Gly	Val	His	His 175	Ala	Phe	Ala	Lys	Ile 180
Leu	Ala	Glu	Gly	Gly 185	Ile	Arg	Gly	Leu	Trp 190	Ala	Gly	Trp	Val	Pro 195
Asn	Ile	Gln	Arg	Ala 200	Ala	Leu	Val	Asn	Met 205	Gly	Asp	Leu	Thr	Thr 210
Tyr	Asp	Thr	Val	Lys 215	His	Tyr	Leu	Val	Leu 220	Asn	Thr	Pro	Leu	Glu 225
Asp	Asn	Ile	Met	Thr 230	His	Gly	Leu	Ser	Ser 235	Leu	Cys	Ser	Gly	Leu 240
Val	Ala	Ser	Ile	Leu 245	Gly	Thr	Pro	Ala	Asp 250	Val	Ile	Lys	Ser	Arg 255
Ile	Met	Asn	Gln	Pro 260	Arg	Asp	Lys	Gln	Gly 265	Arg	Gly	Leu	Leu	Tyr 270
Lys	Ser	Ser	Thr	Asp 275	Cys	Leu	Ile	Gln	Ala 280	Val	Gln	Gly	Glu	Gly 285
Phe	Met	Ser	Leu	Tyr 290	Lys	Gly	Phe	Leu	Pro 295	Ser	Trp	Leu	Arg	Met 300
Thr	Pro	Trp	Ser	Met 305	Val	Phe	Trp	Leu	Thr 310	Tyr	Glu	Lys	Ile	Arg 315
Glu	Met	Ser	Gly	Val 320	Ser	Pro	Phe							

<210> SEQ ID NO 127
<211> LENGTH: 1505
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 127

cgcggtatcgg	acccaagcag	gtcggcggcg	gcggcaggag	agcggccggg	50
cgtcagctcc	tcgacccccg	tgtcgggcta	gtccagcgag	gcggacgggc	100
ggcgtgggcc	catggccagg	cccggcatgg	agcggtggcg	cgaccggctg	150
gcgctggtga	cgggggcctc	ggggggcatc	ggcgcggccg	tggcccgggc	200
cctggtccag	cagggactga	aggtggtggg	ctgcgcccg	actgtgggca	250
acatcgagga	gctggctgct	gaatgtaaga	gtgcaggcta	ccccgggact	300
ttgatcccct	acagatgtga	cctatcaaat	gaagaggaca	tcctctccat	350
gttctcagct	atccgttctc	agcacagcgg	tgtagacatc	tgcatcaaca	400
atgctggctt	ggccccgcct	gacaccctgc	tctcaggcag	caccagtggg	450
tggaaggaca	tgttcaatgt	gaacgtgctg	gccctcagca	tctgcacacg	500
ggaagcctac	cagtccatga	aggagcggaa	tgtggacgat	gggcacatca	550
ttaacatcaa	tagcatgtct	ggccaccgag	tgttaccctt	gtctgtgacc	600
cactttctata	gtgccaccaa	gtatgccgtc	actgcgctga	cagaggggact	650
gaggcaagag	cttcggggagg	cccagaccga	catccgagcc	acgtgcatct	700
ctccagggtg	ggtggagaca	caattgcctt	tcaaactcca	cgacaaggac	750
cctgagaagg	cagctgccac	ctatgagcaa	atgaagtgtc	tcaaaccgga	800
ggatgtggcc	gaggctgtta	tctacgtcct	cagcaccctt	gcacacatcc	850

-continued

```
agattggaga catccagatg aggcccacgg agcaggtgac ctagtgactg      900
tgggagctcc tccttccctc cccacccttc atggcttgcc tcctgcctct      950
ggatttttagg tgttgatttc tggatcacgg gataccactt cctgtccaca     1000
ccccgaccag gggtagaaa atttgtttga gatttttata tcatcttgtc     1050
aaattgcttc agttgtaaat gtgaaaaatg ggctggggaa aggaggtggt     1100
gtccctaatt gttttacttg ttaacttggt cttgtgcccc tgggcacttg     1150
gcctttgtct gctctcagtg tcttcccttt gacatgggaa aggagttgtg     1200
gccaaaatcc ccatcttctt gcacctcaac gtctgtggct cagggctggg     1250
gtggcagagg gaggccttca ccttatatct gtgttgttat ccagggtccc     1300
agacttcctc ctctgcctgc cccactgcac cctctcccc ttatctatct     1350
ccttctcggc tcccagccc agtcttggct tcttgctccc tcctggggtc     1400
atccctccac tctgactctg actatggcag cagaacacca gggcctggcc     1450
cagtggaatt catggtgatc attaaaaaag aaaaatcgca accaaaaaaa     1500
aaaaa                                                    1505
```

<210> SEQ ID NO 128
<211> LENGTH: 260
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 128

```
Met Ala Arg Pro Gly Met Glu Arg Trp Arg Asp Arg Leu Ala Leu
 1             5             10            15
Val Thr Gly Ala Ser Gly Gly Ile Gly Ala Ala Val Ala Arg Ala
20            25            30
Leu Val Gln Gln Gly Leu Lys Val Val Gly Cys Ala Arg Thr Val
35            40            45
Gly Asn Ile Glu Glu Leu Ala Ala Glu Cys Lys Ser Ala Gly Tyr
50            55            60
Pro Gly Thr Leu Ile Pro Tyr Arg Cys Asp Leu Ser Asn Glu Glu
65            70            75
Asp Ile Leu Ser Met Phe Ser Ala Ile Arg Ser Gln His Ser Gly
80            85            90
Val Asp Ile Cys Ile Asn Asn Ala Gly Leu Ala Arg Pro Asp Thr
95            100           105
Leu Leu Ser Gly Ser Thr Ser Gly Trp Lys Asp Met Phe Asn Val
110           115           120
Asn Val Leu Ala Leu Ser Ile Cys Thr Arg Glu Ala Tyr Gln Ser
125           130           135
Met Lys Glu Arg Asn Val Asp Asp Gly His Ile Ile Asn Ile Asn
140           145           150
Ser Met Ser Gly His Arg Val Leu Pro Leu Ser Val Thr His Phe
155           160           165
Tyr Ser Ala Thr Lys Tyr Ala Val Thr Ala Leu Thr Glu Gly Leu
170           175           180
Arg Gln Glu Leu Arg Glu Ala Gln Thr His Ile Arg Ala Thr Cys
185           190           195
```

-continued

Ile	Ser	Pro	Gly	Val	Val	Glu	Thr	Gln	Phe	Ala	Phe	Lys	Leu	His
			200						205					210
Asp	Lys	Asp	Pro	Glu	Lys	Ala	Ala	Ala	Thr	Tyr	Glu	Gln	Met	Lys
			215						220					225
Cys	Leu	Lys	Pro	Glu	Asp	Val	Ala	Glu	Ala	Val	Ile	Tyr	Val	Leu
			230						235					240
Ser	Thr	Pro	Ala	His	Ile	Gln	Ile	Gly	Asp	Ile	Gln	Met	Arg	Pro
			245						250					255
Thr	Glu	Gln	Val	Thr										
			260											

<210> SEQ ID NO 129
<211> LENGTH: 1177
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 129

aacttctaca tgggcctcct gctgctggtg ctcttcctca gcctcctgcc	50
ggtggcctac accatcatgt ccctcccacc ctctttgac tgcggggcgt	100
tcaggtgcag agtctcagtt gcccgggagc acctcccctc ccgaggcagt	150
ctgctcagag ggcctcggcc cagaattcca gttctggttt catgccagcc	200
tgtaaaaggc catggaactt tgggtgaatc accgatgcca tttaagaggg	250
ttttctgcca ggatggaaat gttaggtcgt tctgtgtctg cgctgttcat	300
ttcagtagcc accagccacc tgtggccgtt gagtgctga aatgaggaac	350
tgagaaaatt aatttctcat gtatttttct catttattta ttaattttta	400
actgatagtt gtacatatat gggggtacat gtgatatttg gatacatgta	450
tacaatatat aatgatcaaa tcagggtaac tgggatatcc atcacatcaa	500
acatttattt tttattcttt ttagacagag tctcactctg tcaccagggc	550
tggagtgcag tgggtgccatc tcagettact gcaacctctg cctgccaggt	600
tcaagcgatt ctcatgcctc cacctcccaa gtagctggga ctacaggcat	650
gcaccacaat gcccaactaa tttttgtatt tttagtagag acgggggtttt	700
gccatgttgc ccaggctggc ctggaactcc tggcctcaaa caatccactt	750
gcctcggcct cccaaagtgt tatgattaca ggcgtgagcc accgtgcctg	800
gcctaaacat ttatcttttc tttgtgttgg gaactttgaa attatacaat	850
gaattattgt taactgtcat ctccctgctg tgctatggaa cactgggact	900
tcttccctct atctaactgt atatttgtac cagttaacca accgtacttc	950
atccccactc ctctctatcc ttcccaacct ctgatcacct cattctactc	1000
tctacctcca tgagatccac ttttttagct cccacatgtg agtaagaaaa	1050
tgcaatatat gtctttctgt gcctggctta tttcacttaa cataatgact	1100
tcctgtttcca tccatgttgc tgcaaatgac aggatttcgt tcttaatttc	1150
aattaaaata accacacatg gcaaaaa	1177

<210> SEQ ID NO 130
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 130

Met	Gly	Leu	Leu	Leu	Leu	Val	Leu	Phe	Leu	Ser	Leu	Leu	Pro	Val
1				5					10				15	
Ala	Tyr	Thr	Ile	Met	Ser	Leu	Pro	Pro	Ser	Phe	Asp	Cys	Gly	Pro
				20					25				30	
Phe	Arg	Cys	Arg	Val	Ser	Val	Ala	Arg	Glu	His	Leu	Pro	Ser	Arg
				35					40				45	
Gly	Ser	Leu	Leu	Arg	Gly	Pro	Arg	Pro	Arg	Ile	Pro	Val	Leu	Val
				50					55				60	
Ser	Cys	Gln	Pro	Val	Lys	Gly	His	Gly	Thr	Leu	Gly	Glu	Ser	Pro
				65					70				75	
Met	Pro	Phe	Lys	Arg	Val	Phe	Cys	Gln	Asp	Gly	Asn	Val	Arg	Ser
				80					85				90	
Phe	Cys	Val	Cys	Ala	Val	His	Phe	Ser	Ser	His	Gln	Pro	Pro	Val
				95					100				105	
Ala	Val	Glu	Cys	Leu	Lys									
				110										

<210> SEQ ID NO 131
<211> LENGTH: 2061
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 131

ttctgaagta	acggaagcta	ccttgtataa	agacctcaac	actgctgacc	50
atgatcagcg	cagcctggag	catcttcctc	atcgggacta	aaattgggct	100
gttccttcaa	gtagcacctc	tatcagttat	ggctaaatcc	tgtccatctg	150
tgtgtcgctg	cgatgcgggt	ttcattttact	gtaatgatcg	ctttctgaca	200
tccattccaa	caggaatacc	agaggatgct	acaactctct	accttcagaa	250
caaccaaata	aataatgctg	ggattccttc	agatttgaaa	aacttgctga	300
aagtagaaag	aatataccta	taccacaaca	gtttagatga	atttcctacc	350
aacctcccaa	agtatgtaaa	agagttacat	ttgcaagaaa	ataacataag	400
gactatcact	tatgattcac	tttcaaaaat	tcctatctcg	gaagaattac	450
atttagatga	caactctgtc	tctgcagtta	gcatagaaga	gggagcattc	500
cgagacagca	actatctccg	actgcttttc	ctgtcccgta	atcaccttag	550
cacaattccc	tggggtttgc	ccaggactat	agaagaacta	cgcttggtatg	600
ataatcgcat	atccactatt	tcatcaccat	ctcttcaagg	tctcactagt	650
ctaaaacgcc	tggttctaga	tggaaacctg	ttgaacaatc	atggtttagg	700
tgacaaagtt	ttcttcaacc	tagttaattt	gacagagctg	tccttggtgc	750
ggaattccct	gactgctgca	ccagtaaacc	ttccaggcac	aaacctgagg	800
aagctttatc	ttcaagataa	ccacatcaat	cgggtgcccc	caaatgcttt	850
ttcttatcta	aggcagctct	atcgactgga	tatgtccaat	aataacctaa	900
gtaattttacc	tcagggtatc	tttgatgatt	tggacaatat	aacacaactg	950
attcttcgoa	acaatccctg	gtattgctgg	tgcaagatga	aatgggtacg	1000
tgactgggta	caatcactac	ctgtgaaggt	caacgtgcgt	gggctcatgt	1050

-continued

```
gccaagcccc agaaaagggt cgtgggatgg ctattaagga tctcaatgca      1100
gaactgtttg attgtaagga cagtgggatt gtaagcacca ttcagataac      1150
cactgcaata cccaacacag tgtatcctgc ccaaggacag tggccagctc      1200
cagtgaccaa acagccagat attaagaacc ccaagctcac taaggatcaa      1250
caaaccacag ggagtccctc aagaaaaaca attacaatta ctgtgaagtc      1300
tgtcacctct gataccattc atatctcttg gaaacttgct ctacctatga      1350
ctgcttttag actcagctgg cttaaactgg gccatagccc ggcatttgga      1400
tctataacag aaacaattgt aacaggggaa cgcagtgagt acttggtcac      1450
agccctggag cctgattcac cctataaagt atgcatgggt cccatggaaa      1500
ccagcaacct ctacctattt gatgaaactc ctgtttgtat tgagactgaa      1550
actgcacccc ttcgaatgta caaccctaca accaccctca atcgagagca      1600
agagaaagaa ccttacaaaa accccaattt acctttggct gccatcattg      1650
gtggggctgt ggccctgggt accattgccc ttcttgcttt agtgtgttg      1700
tatgttcata ggaatggatc gctcttctca aggaactgtg catatagcaa      1750
agggaggaga agaaaggatg actatgcaga agctggcact aagaaggaca      1800
actctatcct ggaaatcagg gaaacttctt ttcagatggt accaataagc      1850
aatgaaccca tctcgaagga ggagtttgta atacacacca tatttcctcc      1900
taatggaatg aatctgtaca aaaacaatca cagtgaaagc agtagtaacc      1950
gaagctacag agacagtggg attccagact cagatcactc aactcatga      2000
tgctgaagga ctcacagcag acttgtgttt tgggtttttt aaacctaagg      2050
gaggtgatgg t                                     2061
```

<210> SEQ ID NO 132

<211> LENGTH: 649

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 132

```
Met Ile Ser Ala Ala Trp Ser Ile Phe Leu Ile Gly Thr Lys Ile
  1             5             10             15
Gly Leu Phe Leu Gln Val Ala Pro Leu Ser Val Met Ala Lys Ser
  20             25             30
Cys Pro Ser Val Cys Arg Cys Asp Ala Gly Phe Ile Tyr Cys Asn
  35             40             45
Asp Arg Phe Leu Thr Ser Ile Pro Thr Gly Ile Pro Glu Asp Ala
  50             55             60
Thr Thr Leu Tyr Leu Gln Asn Asn Gln Ile Asn Asn Ala Gly Ile
  65             70             75
Pro Ser Asp Leu Lys Asn Leu Leu Lys Val Glu Arg Ile Tyr Leu
  80             85             90
Tyr His Asn Ser Leu Asp Glu Phe Pro Thr Asn Leu Pro Lys Tyr
  95             100            105
Val Lys Glu Leu His Leu Gln Glu Asn Asn Ile Arg Thr Ile Thr
  110            115            120
Tyr Asp Ser Leu Ser Lys Ile Pro Tyr Leu Glu Glu Leu His Leu
```

-continued

125					130					135				
Asp	Asp	Asn	Ser	Val	Ser	Ala	Val	Ser	Ile	Glu	Glu	Gly	Ala	Phe
				140						145				150
Arg	Asp	Ser	Asn	Tyr	Leu	Arg	Leu	Leu	Phe	Leu	Ser	Arg	Asn	His
				155						160				165
Leu	Ser	Thr	Ile	Pro	Trp	Gly	Leu	Pro	Arg	Thr	Ile	Glu	Glu	Leu
				170						175				180
Arg	Leu	Asp	Asp	Asn	Arg	Ile	Ser	Thr	Ile	Ser	Ser	Pro	Ser	Leu
				185						190				195
Gln	Gly	Leu	Thr	Ser	Leu	Lys	Arg	Leu	Val	Leu	Asp	Gly	Asn	Leu
				200						205				210
Leu	Asn	Asn	His	Gly	Leu	Gly	Asp	Lys	Val	Phe	Phe	Asn	Leu	Val
				215						220				225
Asn	Leu	Thr	Glu	Leu	Ser	Leu	Val	Arg	Asn	Ser	Leu	Thr	Ala	Ala
				230						235				240
Pro	Val	Asn	Leu	Pro	Gly	Thr	Asn	Leu	Arg	Lys	Leu	Tyr	Leu	Gln
				245						250				255
Asp	Asn	His	Ile	Asn	Arg	Val	Pro	Pro	Asn	Ala	Phe	Ser	Tyr	Leu
				260						265				270
Arg	Gln	Leu	Tyr	Arg	Leu	Asp	Met	Ser	Asn	Asn	Asn	Leu	Ser	Asn
				275						280				285
Leu	Pro	Gln	Gly	Ile	Phe	Asp	Asp	Leu	Asp	Asn	Ile	Thr	Gln	Leu
				290						295				300
Ile	Leu	Arg	Asn	Asn	Pro	Trp	Tyr	Cys	Gly	Cys	Lys	Met	Lys	Trp
				305						310				315
Val	Arg	Asp	Trp	Leu	Gln	Ser	Leu	Pro	Val	Lys	Val	Asn	Val	Arg
				320						325				330
Gly	Leu	Met	Cys	Gln	Ala	Pro	Glu	Lys	Val	Arg	Gly	Met	Ala	Ile
				335						340				345
Lys	Asp	Leu	Asn	Ala	Glu	Leu	Phe	Asp	Cys	Lys	Asp	Ser	Gly	Ile
				350						355				360
Val	Ser	Thr	Ile	Gln	Ile	Thr	Thr	Ala	Ile	Pro	Asn	Thr	Val	Tyr
				365						370				375
Pro	Ala	Gln	Gly	Gln	Trp	Pro	Ala	Pro	Val	Thr	Lys	Gln	Pro	Asp
				380						385				390
Ile	Lys	Asn	Pro	Lys	Leu	Thr	Lys	Asp	Gln	Gln	Thr	Thr	Gly	Ser
				395						400				405
Pro	Ser	Arg	Lys	Thr	Ile	Thr	Ile	Thr	Val	Lys	Ser	Val	Thr	Ser
				410						415				420
Asp	Thr	Ile	His	Ile	Ser	Trp	Lys	Leu	Ala	Leu	Pro	Met	Thr	Ala
				425						430				435
Leu	Arg	Leu	Ser	Trp	Leu	Lys	Leu	Gly	His	Ser	Pro	Ala	Phe	Gly
				440						445				450
Ser	Ile	Thr	Glu	Thr	Ile	Val	Thr	Gly	Glu	Arg	Ser	Glu	Tyr	Leu
				455						460				465
Val	Thr	Ala	Leu	Glu	Pro	Asp	Ser	Pro	Tyr	Lys	Val	Cys	Met	Val
				470						475				480
Pro	Met	Glu	Thr	Ser	Asn	Leu	Tyr	Leu	Phe	Asp	Glu	Thr	Pro	Val
				485						490				495
Cys	Ile	Glu	Thr	Glu	Thr	Ala	Pro	Leu	Arg	Met	Tyr	Asn	Pro	Thr
				500						505				510

-continued

Thr	Thr	Leu	Asn	Arg	Glu	Gln	Glu	Lys	Glu	Pro	Tyr	Lys	Asn	Pro
			515						520				525	
Asn	Leu	Pro	Leu	Ala	Ala	Ile	Ile	Gly	Gly	Ala	Val	Ala	Leu	Val
			530						535				540	
Thr	Ile	Ala	Leu	Ala	Ala	Leu	Val	Cys	Trp	Tyr	Val	His	Arg	Asn
			545						550				555	
Gly	Ser	Leu	Phe	Ser	Arg	Asn	Cys	Ala	Tyr	Ser	Lys	Gly	Arg	Arg
			560						565				570	
Arg	Lys	Asp	Asp	Tyr	Ala	Glu	Ala	Gly	Thr	Lys	Lys	Asp	Asn	Ser
			575						580				585	
Ile	Leu	Glu	Ile	Arg	Glu	Thr	Ser	Phe	Gln	Met	Leu	Pro	Ile	Ser
			590						595				600	
Asn	Glu	Pro	Ile	Ser	Lys	Glu	Glu	Phe	Val	Ile	His	Thr	Ile	Phe
			605						610				615	
Pro	Pro	Asn	Gly	Met	Asn	Leu	Tyr	Lys	Asn	Asn	His	Ser	Glu	Ser
			620						625				630	
Ser	Ser	Asn	Arg	Ser	Tyr	Arg	Asp	Ser	Gly	Ile	Pro	Asp	Ser	Asp
			635						640				645	
His	Ser	His	Ser											

<210> SEQ ID NO 133
<211> LENGTH: 1882
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 133

ccgtcatccc cctgcagcca cccttcccag agtcctttgc ccaggccacc	50
ccaggcttct tggcagccct gccggggccac ttgtcttcat gtctgccagg	100
gggagggtgg aaggagggtg gaggagggcg tgcagaggca gtctgggctt	150
ggccagagct cagggtgctg agcgtgtgac cagcagtga cagaggccgg	200
ccatggccag cctggggctg ctgctcctgc tcttactgac agcactgcca	250
ccgctgtggt cctcctcact gcctgggctg gacactgctg aaagtaaacg	300
caccattgca gacctgatcc tgtctgcgct ggagagagcc accgtcttcc	350
tagaacagag gctgcctgaa atcaacctgg atggcatggt gggggtcoga	400
gtgctggaag agcagctaaa aagtgtccgg gagaagtggg ccaggagacc	450
cctgtgtcag ccgctgagcc tgcgcgtggg gatgctggg gagaagctgg	500
aggctgccat ccagagatcc ctccactacc tcaagctgag tgatcccaag	550
tacctaagag agttccagct gaccctccag cccgggtttt ggaagtccc	600
acatgcctgg atccacactg atgcctcctt ggtgtacccc acgttcgggc	650
cccaggactc attctcagag gagagaagtg acgtgtgcct ggtgcagctg	700
ctgggaacgg ggacggacag cagcgagccc tgcggcctct cagacctctg	750
caggagcctc atgaccaagc ccggctgctc aggtactgac ctgtcccacc	800
aactgctctt cttcctctgg gccagaatga ggggatgcac acagggacca	850
ctccaacaga gccaggacta tatcaacctc ttctgcgcca acatgatgga	900
cttgaaccgc agagctgagg ccacgagata cgcctaccct acccgggaca	950

-continued

tcttcacatgga aaacatcatg ttctgtggaa tgggcgggctt ctccgacttc	1000
tacaagctcc ggtggctgga ggccattctc agctggcaga aacagcagga	1050
aggatgcttc ggggagcctg atgctgaaga tgaagaatta tctaaagcta	1100
ttcaatatca gcagcatctt tcgaggagag tgaagaggcg agaaaaacaa	1150
ttccagatt ctgctctgt tgctcaggct ggagtacagt ggcgcaatct	1200
cggctcactg caacctttgc ctctgggtt caagcaattc tcttgctca	1250
tcctcccgag tagctgggac tacaggagcg tgccaccata cctggcta	1300
ttttatat ttttagtaga gacagggtt catcatgttg ctcatgctgg	1350
tctcgaactc ctgatctcaa gagatccgcc cacctcaggc tcccaaagt	1400
tgggattata ggtgtgagcc accgtgtctg gctgaaaagc actttcaaag	1450
agactgtgtt gaataaagg ccaaggttct tgccaccag cactcatggg	1500
ggctctctcc cctagatggc tgctctctcc acaacacagc cacagcagtg	1550
gcagccctgg gtggcttct atacatcctg gcagaatacc cccagcaaa	1600
cagagagcca caccatcca caccgccacc accaagcagc cgctgagacg	1650
gacggttcca tgccagctgc ctggaggagg aacagacccc ttagtctctc	1700
atcccttaga tcctggaggg caccgatcac atcctgggaa gaaggcatct	1750
ggaggataag caaagccacc ccgacacca atcttgaag ccctgagtag	1800
gcagggccag ggtaggtggg ggccgggagg gaccaggtg tgaacggatg	1850
aataaagttc aactgcaact gaaaaaaaa aa	1882

<210> SEQ ID NO 134

<211> LENGTH: 440

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 134

Met Ser Ala Arg Gly Arg Trp Glu Gly Gly Gly Arg Arg Ala Cys	
1 5 10 15	
Arg Gly Ser Leu Gly Leu Ala Arg Ala Gln Gly Ala Glu Arg Val	
20 25 30	
Thr Ser Ser Glu Gln Arg Pro Ala Met Ala Ser Leu Gly Leu Leu	
35 40 45	
Leu Leu Leu Leu Leu Thr Ala Leu Pro Pro Leu Trp Ser Ser Ser	
50 55 60	
Leu Pro Gly Leu Asp Thr Ala Glu Ser Lys Ala Thr Ile Ala Asp	
65 70 75	
Leu Ile Leu Ser Ala Leu Glu Arg Ala Thr Val Phe Leu Glu Gln	
80 85 90	
Arg Leu Pro Glu Ile Asn Leu Asp Gly Met Val Gly Val Arg Val	
95 100 105	
Leu Glu Glu Gln Leu Lys Ser Val Arg Glu Lys Trp Ala Gln Glu	
110 115 120	
Pro Leu Leu Gln Pro Leu Ser Leu Arg Val Gly Met Leu Gly Glu	
125 130 135	
Lys Leu Glu Ala Ala Ile Gln Arg Ser Leu His Tyr Leu Lys Leu	
140 145 150	

-continued

Ser	Asp	Pro	Lys	Tyr	Leu	Arg	Glu	Phe	Gln	Leu	Thr	Leu	Gln	Pro
			155						160					165
Gly	Phe	Trp	Lys	Leu	Pro	His	Ala	Trp	Ile	His	Thr	Asp	Ala	Ser
			170						175					180
Leu	Val	Tyr	Pro	Thr	Phe	Gly	Pro	Gln	Asp	Ser	Phe	Ser	Glu	Glu
			185						190					195
Arg	Ser	Asp	Val	Cys	Leu	Val	Gln	Leu	Leu	Gly	Thr	Gly	Thr	Asp
			200						205					210
Ser	Ser	Glu	Pro	Cys	Gly	Leu	Ser	Asp	Leu	Cys	Arg	Ser	Leu	Met
			215						220					225
Thr	Lys	Pro	Gly	Cys	Ser	Gly	Tyr	Cys	Leu	Ser	His	Gln	Leu	Leu
			230						235					240
Phe	Phe	Leu	Trp	Ala	Arg	Met	Arg	Gly	Cys	Thr	Gln	Gly	Pro	Leu
			245						250					255
Gln	Gln	Ser	Gln	Asp	Tyr	Ile	Asn	Leu	Phe	Cys	Ala	Asn	Met	Met
			260						265					270
Asp	Leu	Asn	Arg	Arg	Ala	Glu	Ala	Ile	Gly	Tyr	Ala	Tyr	Pro	Thr
			275						280					285
Arg	Asp	Ile	Phe	Met	Glu	Asn	Ile	Met	Phe	Cys	Gly	Met	Gly	Gly
			290						295					300
Phe	Ser	Asp	Phe	Tyr	Lys	Leu	Arg	Trp	Leu	Glu	Ala	Ile	Leu	Ser
			305						310					315
Trp	Gln	Lys	Gln	Gln	Glu	Gly	Cys	Phe	Gly	Glu	Pro	Asp	Ala	Glu
			320						325					330
Asp	Glu	Glu	Leu	Ser	Lys	Ala	Ile	Gln	Tyr	Gln	Gln	His	Phe	Ser
			335						340					345
Arg	Arg	Val	Lys	Arg	Arg	Glu	Lys	Gln	Phe	Pro	Asp	Ser	Arg	Ser
			350						355					360
Val	Ala	Gln	Ala	Gly	Val	Gln	Trp	Arg	Asn	Leu	Gly	Ser	Leu	Gln
			365						370					375
Pro	Leu	Pro	Pro	Gly	Phe	Lys	Gln	Phe	Ser	Cys	Leu	Ile	Leu	Pro
			380						385					390
Ser	Ser	Trp	Asp	Tyr	Arg	Ser	Val	Pro	Pro	Tyr	Leu	Ala	Asn	Phe
			395						400					405
Tyr	Ile	Phe	Leu	Val	Glu	Thr	Gly	Phe	His	His	Val	Ala	His	Ala
			410						415					420
Gly	Leu	Glu	Leu	Leu	Ile	Ser	Arg	Asp	Pro	Pro	Thr	Ser	Gly	Ser
			425						430					435
Gln	Ser	Val	Gly	Leu										
			440											

<210> SEQ ID NO 135
<211> LENGTH: 884
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 135

ggctctgagtgcagagctgctgtcatggcggcgctctgtg	50
gggcttctttcccgtcctgc	
tgctgctgctgctatcggggatgtccagagtcgagggt	100
gccccgggctgctgctgaggatcgggaggagtggtggggtc	
ggcataggagatcgttcaa	150
gattgagggcggtgcagttgtccagggtgaagcctcag	200

-continued

gactggatct cggcggcccc agtgctggta gacggagaag agcacgtcgg	250
tttccttaag acagatggga gttttgtggt tcatgatata ccttctggat	300
cttatgtagt ggaagttgta tctccagctt acagatttga tcccgttcga	350
gtggatatac cttcgaaagg aaaaatgaga gcaagatatg tgaattacat	400
caaaacatca gaggttgta gactgcccta tcctctccaa atgaaatctt	450
caggtccacc ttcttacttt attaaaagg aatcgtggg ctggacagac	500
tttctaatag acccaatggt tatgatgatg gttcttcctt tattgatatt	550
tgtgcttctg cctaaagtgg tcaacacaag tgatcctgac atgagacggg	600
aaatggagca gtcaatgaat atgctgaatt ccaaccatga gttgcctgat	650
gtttctgagt tcatgacaag actcttctct tcaaaatcat ctggcaaate	700
tagcagcggc agcagtaaaa caggcaaaa tggggctggc aaaaggaggt	750
agtcaggcgc tccagagctg gcatttgcac aaacacggca aactgggtg	800
gcatccaagt cttggaaaac cgtgtgaagc aactactata aacttgagtc	850
atcccgcagt tgatctctta caactgtgta tggt	884

<210> SEQ ID NO 136
<211> LENGTH: 242
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 136

Met Ala Ala Ala Leu Trp Gly Phe Phe Pro Val Leu Leu Leu Leu	
1 5 10 15	
Leu Leu Ser Gly Asp Val Gln Ser Ser Glu Val Pro Gly Ala Ala	
20 25 30	
Ala Glu Gly Ser Gly Gly Ser Gly Val Gly Ile Gly Asp Arg Phe	
35 40 45	
Lys Ile Glu Gly Arg Ala Val Val Pro Gly Val Lys Pro Gln Asp	
50 55 60	
Trp Ile Ser Ala Ala Arg Val Leu Val Asp Gly Glu Glu His Val	
65 70 75	
Gly Phe Leu Lys Thr Asp Gly Ser Phe Val Val His Asp Ile Pro	
80 85 90	
Ser Gly Ser Tyr Val Val Glu Val Val Ser Pro Ala Tyr Arg Phe	
95 100 105	
Asp Pro Val Arg Val Asp Ile Thr Ser Lys Gly Lys Met Arg Ala	
110 115 120	
Arg Tyr Val Asn Tyr Ile Lys Thr Ser Glu Val Val Arg Leu Pro	
125 130 135	
Tyr Pro Leu Gln Met Lys Ser Ser Gly Pro Pro Ser Tyr Phe Ile	
140 145 150	
Lys Arg Glu Ser Trp Gly Trp Thr Asp Phe Leu Met Asn Pro Met	
155 160 165	
Val Met Met Met Val Leu Pro Leu Leu Ile Phe Val Leu Leu Pro	
170 175 180	
Lys Val Val Asn Thr Ser Asp Pro Asp Met Arg Arg Glu Met Glu	
185 190 195	
Gln Ser Met Asn Met Leu Asn Ser Asn His Glu Leu Pro Asp Val	

-continued

ccatccccag tagactcccc agtcccataa ttgtgtatct tccagccagg	1500
aatcctacac ggccagcatg tattttctaca aataaagttt tctttgcata	1550
ccaaaaaaaa aaaaaaaaaa a	1571

<210> SEQ ID NO 138
<211> LENGTH: 261
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 138

Met Arg Gln Phe Pro Lys Thr Ser Phe Asp Ile Ser Pro Glu Met	
1 5 10 15	
Ser Phe Ser Ile Tyr Ser Leu Gln Val Pro Ala Val Pro Gly Leu	
20 25 30	
Thr Cys Trp Ala Leu Thr Ala Glu Pro Gly Trp Gly Gln Asn Lys	
35 40 45	
Gly Ala Thr Thr Cys Ala Thr Asn Ser His Ser Asp Ser Glu Leu	
50 55 60	
Arg Pro Glu Ile Phe Ser Ser Arg Glu Ala Trp Gln Phe Phe Leu	
65 70 75	
Leu Leu Trp Ser Pro Asp Phe Arg Pro Lys Met Lys Ala Ser Ser	
80 85 90	
Leu Ala Phe Ser Leu Leu Ser Ala Ala Phe Tyr Leu Leu Trp Thr	
95 100 105	
Pro Ser Thr Gly Leu Lys Thr Leu Asn Leu Gly Ser Cys Val Ile	
110 115 120	
Ala Thr Asn Leu Gln Glu Ile Arg Asn Gly Phe Ser Glu Ile Arg	
125 130 135	
Gly Ser Val Gln Ala Lys Asp Gly Asn Ile Asp Ile Arg Ile Leu	
140 145 150	
Arg Arg Thr Glu Ser Leu Gln Asp Thr Lys Pro Ala Asn Arg Cys	
155 160 165	
Cys Leu Leu Arg His Leu Leu Arg Leu Tyr Leu Asp Arg Val Phe	
170 175 180	
Lys Asn Tyr Gln Thr Pro Asp His Tyr Thr Leu Arg Lys Ile Ser	
185 190 195	
Ser Leu Ala Asn Ser Phe Leu Thr Ile Lys Lys Asp Leu Arg Leu	
200 205 210	
Ser His Ala His Met Thr Cys His Cys Gly Glu Glu Ala Met Lys	
215 220 225	
Lys Tyr Ser Gln Ile Leu Ser His Phe Glu Lys Leu Glu Pro Gln	
230 235 240	
Ala Ala Val Val Lys Ala Leu Gly Glu Leu Asp Ile Leu Leu Gln	
245 250 255	
Trp Met Glu Glu Thr Glu	
260	

<210> SEQ ID NO 139
<211> LENGTH: 2395
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 139

-continued

cctggagccg gaagcgcggc tgcagcaggg cgaggctcca ggtggggtcg	50
gttcgcatac cagcctagcg tgtccacgat gcggctgggc tccgggactt	100
tcgctacctg ttgcgtagcg atcgaggtgc tagggatcgc ggtcttcctt	150
cggggattct tcccggctcc cgttcgttcc tctgccagag cggaacacgg	200
agcggagccc ccagcgccc aaccctcggc tggagccagt tctaactgga	250
ccacgctgcc accacctctc ttcagtaaag ttgttattgt tctgatagat	300
gccttgagag atgattttgt gtttgggtca aaggggtgta aatttatgcc	350
ctacacaact taccttgtagg aaaaaggagc atctcacagt tttgtggctg	400
aagcaaagcc acctacagtt actatgcctc gaatcaaggc attgatgacg	450
gggagccttc ctggctttgt cgacgtcatc aggaacctca attctcctgc	500
actgctggaa gacagtgtga taagacaagc aaaagcagct ggaaaaagaa	550
tagtctttta tggagatgaa acctgggtta aattattccc aaagcatttt	600
gtggaatatg atggaacaac ctcatTTTTc gtgtcagatt acacagaggt	650
ggataataat gtcacgaggc atttgataaa agtattaaaa agaggagatt	700
gggacataatt aatcctccac tacctggggc tggaccacat tggccacatt	750
tcagggccca acagccccct gattgggcag aagctgagcg agatggacag	800
cgtgctgatg aagatccaca cctactgca gtcgaaggag agagagacgc	850
ctttaccocaa tttgctgggt ctttggtgtg accatggcat gtctgaaaca	900
ggaaagtcaog gggcctcctc caccgaggag gtgaatacac ctctgatTTT	950
aatcagttct gcgtttgaaa ggaaacccgg tgatatccga catccaaagc	1000
acgtccaata gacggatgtg gctgcgacac tggcgatagc acttggttta	1050
ccgattccaa aagacagtgt agggagcctc ctattcccag ttgtggaagg	1100
aagaccaatg agagagcagt tgagatTTTT acatttgaat acagtgcagc	1150
ttagtaaaact gttgcaagag aatgtgccgt catatgaaaa agatcctggg	1200
tttgagcagt ttaaaatgtc agaaagattg catgggaact ggatcagact	1250
gtacttgtag gaaaagcatt cagaagtcct attcaacctg ggctccaagg	1300
ttctcaggca gtacctggat gctctgaaga cgctgagctt gtccctgagt	1350
gcacaagtgg ccagttctc accctgctcc tgctcagcgt ccacagggca	1400
ctgcacagaa aggctgagct ggaagtccca ctgtcatctc ctgggttttc	1450
tctgtctctt tatttggtga tcctggttct ttcggccggt cacgtcattg	1500
tggtgcacct agctgaaagt tcgtgtact tctgtggcct ctcgtggctg	1550
gcggcaggct gcctttcgtt taccagactc tgggtgaaca cctgggtgtg	1600
gccaaagtgt ggcagtgcc tggacagggg gcctcaggga aggacgtgga	1650
gcagccttat ccagggcctc tgggtgtccc gacacagggt ttcacatctg	1700
tgctgtcagg tcagatgcct cagttcttgg aaagctaggt tcctgcgact	1750
gttaccaaagg tgattgtaaa gagctggcgg tcacagagga acaagcccc	1800
cagctgaggg ggtgtgtgaa tcggacagcc toccagcaga ggtgtgggag	1850
ctgcagctga gggaagaaga gacaatcggc ctggacactc aggagggtca	1900

-continued

aaaggagact tggtcgcacc actcatcctg ccacccccag aatgcatacct	1950
gcctcatcag gtccagattt ctttccaagg cggacgtttt ctgttggaat	2000
tcttagtcct tggcctcgga caccttcatt cgtagctgg ggagtgggtg	2050
tgaggcagtg aagaagaggc ggatgggtcac actcagatcc acagagccca	2100
ggatcaaggg acccactgca gtggcagcag gactgttggg cccccacccc	2150
aacctgcac agccctcatc ccctcttggc ttgagccgtc agaggccctg	2200
tgctgagtg ctgaccgaga cactcacagc tttgtcatca gggcacaggc	2250
ttcctcggag ccaggatgat ctgtgccacg cttgcacctc gggcccatct	2300
gggtcatgc tctctctcct gctattgaat tagtacctag ctgcacacag	2350
tatgtagtta ccaaagaat aaacggcaat aattgagaaa aaaaa	2395

<210> SEQ ID NO 140
<211> LENGTH: 310
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 140

Met Arg Leu Gly Ser Gly Thr Phe Ala Thr Cys Cys Val Ala Ile	1	5	10	15
Glu Val Leu Gly Ile Ala Val Phe Leu Arg Gly Phe Phe Pro Ala	20	25	30	
Pro Val Arg Ser Ser Ala Arg Ala Glu His Gly Ala Glu Pro Pro	35	40	45	
Ala Pro Glu Pro Ser Ala Gly Ala Ser Ser Asn Trp Thr Thr Leu	50	55	60	
Pro Pro Pro Leu Phe Ser Lys Val Val Ile Val Leu Ile Asp Ala	65	70	75	
Leu Arg Asp Asp Phe Val Phe Gly Ser Lys Gly Val Lys Phe Met	80	85	90	
Pro Tyr Thr Thr Tyr Leu Val Glu Lys Gly Ala Ser His Ser Phe	95	100	105	
Val Ala Glu Ala Lys Pro Pro Thr Val Thr Met Pro Arg Ile Lys	110	115	120	
Ala Leu Met Thr Gly Ser Leu Pro Gly Phe Val Asp Val Ile Arg	125	130	135	
Asn Leu Asn Ser Pro Ala Leu Leu Glu Asp Ser Val Ile Arg Gln	140	145	150	
Ala Lys Ala Ala Gly Lys Arg Ile Val Phe Tyr Gly Asp Glu Thr	155	160	165	
Trp Val Lys Leu Phe Pro Lys His Phe Val Glu Tyr Asp Gly Thr	170	175	180	
Thr Ser Phe Phe Val Ser Asp Tyr Thr Glu Val Asp Asn Asn Val	185	190	195	
Thr Arg His Leu Asp Lys Val Leu Lys Arg Gly Asp Trp Asp Ile	200	205	210	
Leu Ile Leu His Tyr Leu Gly Leu Asp His Ile Gly His Ile Ser	215	220	225	
Gly Pro Asn Ser Pro Leu Ile Gly Gln Lys Leu Ser Glu Met Asp	230	235	240	

-continued

Ser	Val	Leu	Met	Lys	Ile	His	Thr	Ser	Leu	Gln	Ser	Lys	Glu	Arg
			245						250					255
Glu	Thr	Pro	Leu	Pro	Asn	Leu	Leu	Val	Leu	Cys	Gly	Asp	His	Gly
			260						265					270
Met	Ser	Glu	Thr	Gly	Ser	His	Gly	Ala	Ser	Ser	Thr	Glu	Glu	Val
			275						280					285
Asn	Thr	Pro	Leu	Ile	Leu	Ile	Ser	Ser	Ala	Phe	Glu	Arg	Lys	Pro
			290						295					300
Gly	Asp	Ile	Arg	His	Pro	Lys	His	Val	Gln					
			305						310					

<210> SEQ ID NO 141
<211> LENGTH: 754
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 141

ggcacgagggc aagccttcca ggttatcgtg acgcaccttg aaagtctgag	50
agctactgcc ctacagaaag ttactagtgc cctaaagctg gcgctggcac	100
tgatgttact gctgctgttg gagtacaact tccctataga aaacaactgc	150
cagcacctta agaccactca caccttcaga gtgaagaact taaacccgaa	200
gaaattcagc attcatgacc aggatcacaa agtactggtc ctggactctg	250
ggaatctcat agcagttcca gataaaaact acatacgccc agagatcttc	300
tttgattag cctcatcctt gagctcagcc tctgcggaga aaggaagtcc	350
gattctcctg ggggtctcta aaggggagtt ttgtctctac tgtgacaagg	400
ataaaggaca aagtcattcca tcccttcagc tgaagaagga gaaactgatg	450
aagctggctg cccaaaagga atcagcacgc cggcccttca tcttttatag	500
ggctcaggtg ggctcctgga acatgctgga gtcggcggct caccctcgat	550
ggttcatctg cacctcctgc aattgtaatg agcctgttgg ggtgacagat	600
aaatttgaga acaggaaaca cattgaattt tcatttcaac cagtttgcaa	650
agctgaaatg agccccagtg aggtcagcga ttaggaaact gccccattga	700
acgccttctc cgctaatttg aactaattgt ataaaaacac caaacctgct	750
cact	754

<210> SEQ ID NO 142
<211> LENGTH: 193
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 142

Met	Leu	Leu	Leu	Leu	Leu	Glu	Tyr	Asn	Phe	Pro	Ile	Glu	Asn	Asn
1				5					10					15
Cys	Gln	His	Leu	Lys	Thr	Thr	His	Thr	Phe	Arg	Val	Lys	Asn	Leu
			20						25					30
Asn	Pro	Lys	Lys	Phe	Ser	Ile	His	Asp	Gln	Asp	His	Lys	Val	Leu
			35						40					45
Val	Leu	Asp	Ser	Gly	Asn	Leu	Ile	Ala	Val	Pro	Asp	Lys	Asn	Tyr
			50						55					60
Ile	Arg	Pro	Glu	Ile	Phe	Phe	Ala	Leu	Ala	Ser	Ser	Leu	Ser	Ser

-continued

	65		70		75
Ala Ser Ala Glu Lys Gly Ser Pro Ile Leu Leu Gly Val Ser Lys	80		85		90
Gly Glu Phe Cys Leu Tyr Cys Asp Lys Asp Lys Gly Gln Ser His	95		100		105
Pro Ser Leu Gln Leu Lys Lys Glu Lys Leu Met Lys Leu Ala Ala	110		115		120
Gln Lys Glu Ser Ala Arg Arg Pro Phe Ile Phe Tyr Arg Ala Gln	125		130		135
Val Gly Ser Trp Asn Met Leu Glu Ser Ala Ala His Pro Gly Trp	140		145		150
Phe Ile Cys Thr Ser Cys Asn Cys Asn Glu Pro Val Gly Val Thr	155		160		165
Asp Lys Phe Glu Asn Arg Lys His Ile Glu Phe Ser Phe Gln Pro	170		175		180
Val Cys Lys Ala Glu Met Ser Pro Ser Glu Val Ser Asp	185		190		

<210> SEQ ID NO 143
<211> LENGTH: 961
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 143

ctagagagta tagggcagaa ggatggcaga tgagtgactc cacatccaga	50
gctgcctccc tttaatccag gatcctgtcc ttcctgtcct gtaggagtgc	100
ctgttgccag tgtggggtga gacaagtttg tcccacaggg ctgtctgagc	150
agataagatt aagggtctgg tctgtgctca attaacctct gtgggcacgg	200
gggtctggaa gagcaaagtc agcgggtgcct acagtcagca ccatgctggg	250
cctgccgtgg aagggaagtc tgtcctgggc gctgctgctg cttctcttag	300
gctcccagat cctgctgata tatgcctggc atttcacga gcaaaggac	350
tgtagatgaac acaatgtcat ggctcgttac ctccctgcc aagtggagtt	400
tgctgtccac acattcaacc aacagagcaa ggactactat gcctacagac	450
tggggcacat cttgaattcc tggaaggagc aggtggagtc caagactgta	500
ttctcaatgg agctactgct ggggagaact aggtgtggga aatttgaaga	550
cgacattgac aactgccatt tccaagaaag cacagagctg aacaatactt	600
tcacctgctt cttcaccatc agcaccaggc cctggatgac tcagttcagc	650
ctcctgaaca agacctgctt ggagggattc cactgagtga aaccactca	700
caggcttgtc catgtgtctc tcccacattc cgtggacatc agcactactc	750
tcctgaggac tcttcagtgg ctgagcagct ttggacttgt ttgttatcct	800
attttgcatg tgtttgagat ctcatatcag tgttttagaa aatccacaca	850
tcttgagcct aatcatgtag ttagatcat taaacatcag cattttaaga	900
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	950
aaaaaaaaa a	961

<210> SEQ ID NO 144

-continued

<211> LENGTH: 147
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 144
Met Leu Gly Leu Pro Trp Lys Gly Gly Leu Ser Trp Ala Leu Leu
1 5 10 15
Leu Leu Leu Leu Gly Ser Gln Ile Leu Leu Ile Tyr Ala Trp His
20 25 30
Phe His Glu Gln Arg Asp Cys Asp Glu His Asn Val Met Ala Arg
35 40 45
Tyr Leu Pro Ala Thr Val Glu Phe Ala Val His Thr Phe Asn Gln
50 55 60
Gln Ser Lys Asp Tyr Tyr Ala Tyr Arg Leu Gly His Ile Leu Asn
65 70 75
Ser Trp Lys Glu Gln Val Glu Ser Lys Thr Val Phe Ser Met Glu
80 85 90
Leu Leu Leu Gly Arg Thr Arg Cys Gly Lys Phe Glu Asp Asp Ile
95 100 105
Asp Asn Cys His Phe Gln Glu Ser Thr Glu Leu Asn Asn Thr Phe
110 115 120
Thr Cys Phe Phe Thr Ile Ser Thr Arg Pro Trp Met Thr Gln Phe
125 130 135
Ser Leu Leu Asn Lys Thr Cys Leu Glu Gly Phe His
140 145

<210> SEQ ID NO 145
<211> LENGTH: 1157
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 145
ctgtgcagct cgaggctcca gaggcacact ccagagagag ccaaggttct 50
gacgcgatga ggaagcacct gagctggtgg tggctggcca ctgtctgcat 100
gctgtctctc agccacctct ctgcggtcca gacgaggggc atcaagcaca 150
gaatcaagtg gaaccggaag gccctgccca gcaactgccca gatcactgag 200
gcccaggtgg ctgagaaccg cccgggagcc ttcataaagc aaggccgcaa 250
gctcgacatt gacttcggag ccgagggcaa caggtactac gaggccaact 300
actggcagtt ccccgatggc atccactaca acggctgctc tgaggctaata 350
gtgaccaagg aggcatttgt caccggctgc atcaatgccca cccaggcggc 400
gaaccagggg gagttccaga agccagacaa caagctccac cagcaggtgc 450
tctggcggct ggtccaggag ctctgctccc tcaagcattg cgagttttgg 500
ttggagaggg gcgcaggact tcgggtcacc atgcaccagc cagtgtctct 550
ctgccttctg gctttgatct ggctcatggt gaaataagct tgccaggagg 600
ctggcagtag agagcgagc agcgagcaaa tcctggcaag tgaccagct 650
cttctcccc aaaccacgc gtgttctgaa ggtgcccagg agcggcgatg 700
cactgcgact gcaaatgccg ctcccacgta tgcgccctgg tatgtgcttg 750
cgttctgata gatgggggac tgtggcttct ccgtcactcc attctcagcc 800

-continued

cctagcagag cgtctggcac actagattag tagtaaatgc ttgatgagaa	850
gaacacatca ggcactgcg cacctgcttc acagtacttc ccaacaactc	900
ttagaggtag gtgtattccc gttttacaga taaggaaact gaggcccaga	950
gagctgaagt actgcaccca gcatcaccag ctagaaagtg gcagagccag	1000
gattcaaccc tggcttgtct aaccccaggt tttctgtct gtccaattcc	1050
agagctgtct ggtgatcact ttatgtctca cagggacca catccaaaca	1100
tgtatctcta atgaaattgt gaaagctcca tgtttagaaa taaatgaaaa	1150
cacctga	1157

<210> SEQ ID NO 146
<211> LENGTH: 176
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 146

Met Arg Lys His Leu Ser Trp Trp Trp Leu Ala Thr Val Cys Met	
1 5 10 15	
Leu Leu Phe Ser His Leu Ser Ala Val Gln Thr Arg Gly Ile Lys	
20 25 30	
His Arg Ile Lys Trp Asn Arg Lys Ala Leu Pro Ser Thr Ala Gln	
35 40 45	
Ile Thr Glu Ala Gln Val Ala Glu Asn Arg Pro Gly Ala Phe Ile	
50 55 60	
Lys Gln Gly Arg Lys Leu Asp Ile Asp Phe Gly Ala Glu Gly Asn	
65 70 75	
Arg Tyr Tyr Glu Ala Asn Tyr Trp Gln Phe Pro Asp Gly Ile His	
80 85 90	
Tyr Asn Gly Cys Ser Glu Ala Asn Val Thr Lys Glu Ala Phe Val	
95 100 105	
Thr Gly Cys Ile Asn Ala Thr Gln Ala Ala Asn Gln Gly Glu Phe	
110 115 120	
Gln Lys Pro Asp Asn Lys Leu His Gln Gln Val Leu Trp Arg Leu	
125 130 135	
Val Gln Glu Leu Cys Ser Leu Lys His Cys Glu Phe Trp Leu Glu	
140 145 150	
Arg Gly Ala Gly Leu Arg Val Thr Met His Gln Pro Val Leu Leu	
155 160 165	
Cys Leu Leu Ala Leu Ile Trp Leu Met Val Lys	
170 175	

<210> SEQ ID NO 147
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 147

gccttggcct cccaaagggc tgggattata ggcgtgacca ccatgtctgg	50
tccagagtct catttcctga tgatttatag actcaaagaa aactcatgtt	100
cagaagctct cttctcttct ggctctctct ctgtcttctt tccctctttc	150
ttcttatttt aattagtagc atctactcag agtcatgcaa gctggaaatc	200

-continued

tttcattttg cttgtcagtg gggtaggtca ctgagtcctta gtttttattt	250
tttgaaattht caactttcag attcaggggg tacatgtgaa ggthttgttt	300
atgagtatat tgcattgatgc tgaggthttgg ggt	333
<210> SEQ ID NO 148	
<211> LENGTH: 73	
<212> TYPE: PRT	
<213> ORGANISM: Homo Sapien	
<400> SEQUENCE: 148	
Met Phe Arg Ser Ser Leu Leu Phe Trp Pro Pro Leu Cys Leu Leu	
1 5 10 15	
Ser Leu Phe Leu Leu Ile Leu Ile Ser Ser Ile Tyr Ser Glu Ser	
20 25 30	
Cys Lys Leu Glu Ile Phe His Phe Ala Cys Gln Trp Gly Arg Ser	
35 40 45	
Leu Ser Leu Ser Phe Tyr Phe Leu Lys Phe Gln Leu Ser Asp Ser	
50 55 60	
Gly Gly Thr Cys Glu Gly Leu Phe Tyr Glu Tyr Ile Ala	
65 70	
<210> SEQ ID NO 149	
<211> LENGTH: 1893	
<212> TYPE: DNA	
<213> ORGANISM: Homo Sapien	
<400> SEQUENCE: 149	
gtctccgcgt cacaggaact tcagcaccct cagggcggac agcgcctccc	50
tctacctgga gacttgactc ccgcgcgccc caacctgct tatcccttga	100
ccgtcgagtg tcagagatcc tgcagccgcc cagtcccggc ccctctccc	150
ccccacaccc accctcctgg ctcttcctgt ttttactcct ccttttcatt	200
cataacaaaa gctacagctc caggagccca gcgcgggct gtgacccaag	250
ccgagcgtgg aagaatgggg ttctctggga ccggcacttg gattctggtg	300
ttagtgctcc cgattcaagc tttcccaaaa cctggaggaa gccaaagaaa	350
atctctacat aatagagaat taagtgcaga aagaccttg aatgaacaga	400
ttgctgaagc agaagaagac aagattaaaa aaacatatcc tccagaaaa	450
aagccaggtc agagcaacta ttcttttgtt gataacttga acctgctaaa	500
ggcaataaca gaaaaggaaa aaattgagaa agaaagaaa tctataagaa	550
gtccccact tgataataag ttgaatgtg aagatgttga ttcaaccaag	600
aatcgaaaa tgatcgatga ttatgactct actaagagtg gattggatca	650
taaatthtcaa gatgatccag atgggtcttca tcaactagac gggactcctt	700
taaccgtga agacattgtc cataaaatcg ctgccaggat ttatgaagaa	750
aatgacagag ccgtgtttga caagattgtt tctaaactac ttaatctcgg	800
ccttatcaca gaaagccaag cacatacact ggaagatgaa gtagcagagg	850
ttttacaaaa attaatctca aaggaagcca acaattatga ggaggatccc	900
aataagccca caagctggac tgagaatcag gctggaaaaa taccagagaa	950
agtactcca atggcagcaa ttcaagatgg tottgctaag ggagaaaacg	1000

-continued

atgaaacagt atctaacaca ttaaccttga caaatggctt ggaaaggaga	1050
actaaaacct acagtgaaga caactttgag gaactccaat atttcccaaa	1100
tttctatgcg ctactgaaaa gtattgattc agaaaaagaa gcaaaagaga	1150
aagaaacact gattactatc atgaaaacac tgattgactt tgtgaagatg	1200
atggtgaaat atggaacaat atctccagaa gaaggtgttt cctaccttga	1250
aaacttggat gaaatgattg ctcttcagac caaaaacaag ctagaaaaaa	1300
atgctactga caatataagc aagcttttcc cagcaccatc agagaagagt	1350
catgaagaaa cagacagtac caaggaagaa gcagctaaga tggaaaagga	1400
atatggaagc ttgaaggatt ccacaaaaga tgataactcc aaccaggag	1450
gaaagacaga tgaacccaaa ggaaaaacag aagcctattht ggaagccatc	1500
agaaaaata ttgaatggtt gaagaacat gacaaaaagg gaaataaaga	1550
agattatgac ctttcaaaga tgagagactt catcaataaa caagctgatg	1600
cttatgtgga gaaaggcatc ctgacaagg aagaagccga ggccatcaag	1650
cgcatttata gcagcctgta aaaatggcaa aagatccagg agtctttcaa	1700
ctgtttcaga aaacataata tagcttaaaa cacttctaatt tctgtgatta	1750
aaatthtttg acccaagggt tattagaaag tgctgaattht acagtagtta	1800
accttttaca agtgggttaa acatagcttht cttcccgtaa aaactatctg	1850
aaagtaaagt tgtatgtaag ctgaaaaaaa aaaaaaaaaa aaa	1893

<210> SEQ ID NO 150

<211> LENGTH: 468

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 150

Met Gly Phe Leu Gly Thr Gly Thr Trp Ile Leu Val Leu Val Leu	
1 5 10 15	
Pro Ile Gln Ala Phe Pro Lys Pro Gly Gly Ser Gln Asp Lys Ser	
20 25 30	
Leu His Asn Arg Glu Leu Ser Ala Glu Arg Pro Leu Asn Glu Gln	
35 40 45	
Ile Ala Glu Ala Glu Glu Asp Lys Ile Lys Lys Thr Tyr Pro Pro	
50 55 60	
Glu Asn Lys Pro Gly Gln Ser Asn Tyr Ser Phe Val Asp Asn Leu	
65 70 75	
Asn Leu Leu Lys Ala Ile Thr Glu Lys Glu Lys Ile Glu Lys Glu	
80 85 90	
Arg Gln Ser Ile Arg Ser Ser Pro Leu Asp Asn Lys Leu Asn Val	
95 100 105	
Glu Asp Val Asp Ser Thr Lys Asn Arg Lys Leu Ile Asp Asp Tyr	
110 115 120	
Asp Ser Thr Lys Ser Gly Leu Asp His Lys Phe Gln Asp Asp Pro	
125 130 135	
Asp Gly Leu His Gln Leu Asp Gly Thr Pro Leu Thr Ala Glu Asp	
140 145 150	
Ile Val His Lys Ile Ala Ala Arg Ile Tyr Glu Glu Asn Asp Arg	

-continued

				155								160								165			
Ala	Val	Phe	Asp	Lys	Ile	Val	Ser	Lys	Leu	Leu	Asn	Leu	Gly	Leu									
				170					175								180						
Ile	Thr	Glu	Ser	Gln	Ala	His	Thr	Leu	Glu	Asp	Glu	Val	Ala	Glu									
				185					190								195						
Val	Leu	Gln	Lys	Leu	Ile	Ser	Lys	Glu	Ala	Asn	Asn	Tyr	Glu	Glu									
				200					205								210						
Asp	Pro	Asn	Lys	Pro	Thr	Ser	Trp	Thr	Glu	Asn	Gln	Ala	Gly	Lys									
				215					220								225						
Ile	Pro	Glu	Lys	Val	Thr	Pro	Met	Ala	Ala	Ile	Gln	Asp	Gly	Leu									
				230					235								240						
Ala	Lys	Gly	Glu	Asn	Asp	Glu	Thr	Val	Ser	Asn	Thr	Leu	Thr	Leu									
				245					250								255						
Thr	Asn	Gly	Leu	Glu	Arg	Arg	Thr	Lys	Thr	Tyr	Ser	Glu	Asp	Asn									
				260					265								270						
Phe	Glu	Glu	Leu	Gln	Tyr	Phe	Pro	Asn	Phe	Tyr	Ala	Leu	Leu	Lys									
				275					280								285						
Ser	Ile	Asp	Ser	Glu	Lys	Glu	Ala	Lys	Glu	Lys	Glu	Thr	Leu	Ile									
				290					295								300						
Thr	Ile	Met	Lys	Thr	Leu	Ile	Asp	Phe	Val	Lys	Met	Met	Val	Lys									
				305					310								315						
Tyr	Gly	Thr	Ile	Ser	Pro	Glu	Glu	Gly	Val	Ser	Tyr	Leu	Glu	Asn									
				320					325								330						
Leu	Asp	Glu	Met	Ile	Ala	Leu	Gln	Thr	Lys	Asn	Lys	Leu	Glu	Lys									
				335					340								345						
Asn	Ala	Thr	Asp	Asn	Ile	Ser	Lys	Leu	Phe	Pro	Ala	Pro	Ser	Glu									
				350					355								360						
Lys	Ser	His	Glu	Glu	Thr	Asp	Ser	Thr	Lys	Glu	Glu	Ala	Ala	Lys									
				365					370								375						
Met	Glu	Lys	Glu	Tyr	Gly	Ser	Leu	Lys	Asp	Ser	Thr	Lys	Asp	Asp									
				380					385								390						
Asn	Ser	Asn	Pro	Gly	Gly	Lys	Thr	Asp	Glu	Pro	Lys	Gly	Lys	Thr									
				395					400								405						
Glu	Ala	Tyr	Leu	Glu	Ala	Ile	Arg	Lys	Asn	Ile	Glu	Trp	Leu	Lys									
				410					415								420						
Lys	His	Asp	Lys	Lys	Gly	Asn	Lys	Glu	Asp	Tyr	Asp	Leu	Ser	Lys									
				425					430								435						
Met	Arg	Asp	Phe	Ile	Asn	Lys	Gln	Ala	Asp	Ala	Tyr	Val	Glu	Lys									
				440					445								450						
Gly	Ile	Leu	Asp	Lys	Glu	Glu	Ala	Glu	Ala	Ile	Lys	Arg	Ile	Tyr									
				455					460								465						
Ser Ser Leu																							

<210> SEQ ID NO 151
<211> LENGTH: 2598
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 151

cggctcgagg ctccgccag gagaaaggaa cattctgagg ggagtctaca	50
cctgtggag ctcaagatgg tcctgagtgg ggcgtgtgc ttccgaatga	100

-continued

aggactcggc attgaaggtg ctttatctgc ataataacca gcttctagct	150
ggagggctgc atgcaggga ggtcattaaa ggtgaagaga tcagcgtggt	200
ccccaatcgg tggctggatg ccagcctgtc ccccgtcac ctaggtgtcc	250
agggtggaag ccagtgcctg tcatgtgggg tggggcagga gccgactcta	300
acactagagc cagtgaacat catggagctc tatcttggtg ccaaggaatc	350
caagagcttc accttctacc ggccggacat ggggctcacc tccagcttcg	400
agtcggctgc ctacccgggc tggttcctgt gcacggtgcc tgaagccgat	450
cagcctgtca gactcaccca gcttcccag aatggtggct ggaatgcccc	500
catcacagac ttctacttcc agcagtgtga ctagggaac gtgccccca	550
gaactccctg ggcagagcca gctcgggtga ggggtgagtg gaggagaccc	600
atggcggaca atcactctct ctgctctcag gacccccacg tctgacttag	650
tgggcacctg accactttgt cttctggttc ccagtttga taaattotga	700
gatttgagc tcagtcacg gtctccccc actggatggt gctactgctg	750
tggaaccttg taaaaacat gtggggtaaa ctgggaataa catgaaaaga	800
tttctgtggg ggtgggggtg gggagtggg ggaatcattc ctgcttaatg	850
gtaactgaca agtgttacc tgagccccg aggccaccc atccccagtt	900
gagccttata gggtcagtag ctctccacat gaagtcctgt cactcaccac	950
tgtgcaggag agggaggtg tcatagagtc agggatctat ggcccttggc	1000
ccagccccac ccccttccct ttaatcctgc cactgtcata tgctacctt	1050
cctatctctt ccctcatcat cttgttgtg gcataggag gtggtgatgt	1100
cagaagaaat ggctcgagct cagaagataa aagataagta gggtagctg	1150
atcctctttt aaaaaccaa gataaatca aaatccaga tgctggtctc	1200
tattcccatg aaaaagtgt catgacatat tgagaagacc tacttacaaa	1250
gtggcatata ttgcaattta ttttaattaa aagataccta tttatatatt	1300
tctttataga aaaaagtctg gaagagtta cttcaattgt agcaatgtca	1350
gggtggtggc agtatagggt atttttcttt taattctgtt aatttatctg	1400
tatttcctaa tttttctaca atgaagatga attccttgta taaaaataag	1450
aaaagaaatt aatcttgagg taagcagagc agacatcatc tctgattgtc	1500
ctcagcctcc acttccccag agtaaatca aattgaatcg agctctgctg	1550
ctctggttgg ttgtagtagt gatcaggaaa cagatctcag caaagccact	1600
gaggaggagg ctgtgctgag tttgtgtggc tggaatctct gggtaaggaa	1650
cttaagaac aaaaatcatc tggtaattct ttcctagaag gatcacagcc	1700
cctgggattc caaggcattg gatccagtct ctaagaaggc tgctgtactg	1750
gttgaattgt gtccccctca aattcacatc cttcttgga tctcagtctg	1800
tgagtttatt tggagataag gtctctgcag atgtagttag ttaagacaag	1850
gtcatgctgg atgaagtag acctaaatc aatatgactg gtttccttgt	1900
atgaaaagga gaggacacag agacagagga gacgcgggga agactatgta	1950
aagatgaagg cagagatcgg agttttgcag ccacaagcta agaaacacca	2000

-continued

aggattgtgg caaccatcag aagcttgga gaggcaaaga agaattcttc	2050
cctagaggct ttagagggat aacggctctg ctgaaacctt aatctcagac	2100
ttccagcctc ctgaacgaag aaagaataaa tttcggctgt tttaagccac	2150
caaggataat tggttacagc agctctagga aactaataca gctgctaaaa	2200
tgatccctgt ctctcgtgt ttacattctg tgtgtgtccc ctcccacaat	2250
gtaccaaagt tgtctttgtg accaatagaa tatggcagaa gtgatggcat	2300
gccacttcca agattaggtt ataaaagaca ctgcagcttc tacttgagcc	2350
ctctctctct gccacccacc gcccacaatc tatcttggt cactcgtct	2400
gggggaagct agctgccatg ctatgagcag gcctataaag agacttacgt	2450
ggtaaaaaat gaagtctcct gccacagcc acattagtga acctagaagc	2500
agagactctg tgagataatc gatgtttgtt gttttaagtt gtcagtttt	2550
ggtctaactt gttatgcagc aatagataaa taatatgcag agaaagag	2598
<210> SEQ ID NO 152	
<211> LENGTH: 155	
<212> TYPE: PRT	
<213> ORGANISM: Homo Sapien	
<400> SEQUENCE: 152	
Met Val Leu Ser Gly Ala Leu Cys Phe Arg Met Lys Asp Ser Ala	
1 5 10 15	
Leu Lys Val Leu Tyr Leu His Asn Asn Gln Leu Leu Ala Gly Gly	
20 25 30	
Leu His Ala Gly Lys Val Ile Lys Gly Glu Glu Ile Ser Val Val	
35 40 45	
Pro Asn Arg Trp Leu Asp Ala Ser Leu Ser Pro Val Ile Leu Gly	
50 55 60	
Val Gln Gly Gly Ser Gln Cys Leu Ser Cys Gly Val Gly Gln Glu	
65 70 75	
Pro Thr Leu Thr Leu Glu Pro Val Asn Ile Met Glu Leu Tyr Leu	
80 85 90	
Gly Ala Lys Glu Ser Lys Ser Phe Thr Phe Tyr Arg Arg Asp Met	
95 100 105	
Gly Leu Thr Ser Ser Phe Glu Ser Ala Ala Tyr Pro Gly Trp Phe	
110 115 120	
Leu Cys Thr Val Pro Glu Ala Asp Gln Pro Val Arg Leu Thr Gln	
125 130 135	
Leu Pro Glu Asn Gly Gly Trp Asn Ala Pro Ile Thr Asp Phe Tyr	
140 145 150	
Phe Gln Gln Cys Asp	
155	
<210> SEQ ID NO 153	
<211> LENGTH: 1152	
<212> TYPE: DNA	
<213> ORGANISM: Homo Sapien	
<400> SEQUENCE: 153	
cttcagaaca ggttctcctt cccagctcac cagttgctcg agttagaatt	50
gtctgcaatg gccgccctgc agaaatctgt gagctctttc cttatgggga	100

-continued

```
ccctggccac cagctgcctc cttctcttgg ccctcttggg acagggagga      150
gcagctgcgc ccatcagctc ccactgcagg cttgacaagt ccaacttcca      200
gcagccctat atcaccaacc gcaccttcat gctggctaag gaggctagct      250
tggtcgataa caacacagac gttcgtctca ttggggagaa actgttccac      300
ggagtcagta tgagtgcgct ctgctatctg atgaagcagg tgctgaactt      350
cacccttgaa gaagtgcctg tccctcaatc tgatagggtc cagccttata      400
tgcaggaggt ggtgcccttc ctggccaggc tcagcaacag gctaagcaca      450
tgtcatattg aaggtgatga cctgcataac cagaggaatg tgcaaaagct      500
gaaggacaca gtgaaaaagc ttggagagag tggagagatc aaagcaattg      550
gagaactgga tttgctgttt atgtctctga gaaatgcctg catttgacca      600
gagcaaaagt gaaaaatgaa taactaaccc cctttccctg ctagaataaa      650
caattagatg ccccaaagcg atttttttta accaaaagga agatgggaag      700
ccaaactcca tcatgatggg tggattccaa atgaaccctt gcgttagtta      750
caaaggaaac caatgccact tttgtttata agaccagaag gtagactttc      800
taagcataga tatttattga taacatttca ttgtaactgg tgttctatac      850
acagaaaaca atttattttt taaataattg tctttttcca taaaaaagat      900
tactttccat tccttttagg gaaaaaaccc ctaaatagct tcatgtttcc      950
ataatcagta ctttatattt ataaatgtat ttattattat tataagactg     1000
cattttattt atatcatttt attaatatgg atttatttat agaaacatca     1050
ttcgatattg ctacttgagt gtaaggctaa tattgatatt tatgacaata     1100
attatagagc tataacatgt ttatttgacc tcaataaaca cttggatatc     1150
cc                                                                1152
```

<210> SEQ ID NO 154

<211> LENGTH: 179

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 154

```
Met Ala Ala Leu Gln Lys Ser Val Ser Ser Phe Leu Met Gly Thr
 1             5             10             15
Leu Ala Thr Ser Cys Leu Leu Leu Leu Ala Leu Leu Val Gln Gly
          20             25             30
Gly Ala Ala Ala Pro Ile Ser Ser His Cys Arg Leu Asp Lys Ser
          35             40             45
Asn Phe Gln Gln Pro Tyr Ile Thr Asn Arg Thr Phe Met Leu Ala
          50             55             60
Lys Glu Ala Ser Leu Ala Asp Asn Asn Thr Asp Val Arg Leu Ile
          65             70             75
Gly Glu Lys Leu Phe His Gly Val Ser Met Ser Glu Arg Cys Tyr
          80             85             90
Leu Met Lys Gln Val Leu Asn Phe Thr Leu Glu Glu Val Leu Phe
          95             100            105
Pro Gln Ser Asp Arg Phe Gln Pro Tyr Met Gln Glu Val Val Pro
          110            115            120
```

-continued

Phe	Leu	Ala	Arg	Leu	Ser	Asn	Arg	Leu	Ser	Thr	Cys	His	Ile	Glu
				125					130					135
Gly	Asp	Asp	Leu	His	Ile	Gln	Arg	Asn	Val	Gln	Lys	Leu	Lys	Asp
				140					145					150
Thr	Val	Lys	Lys	Leu	Gly	Glu	Ser	Gly	Glu	Ile	Lys	Ala	Ile	Gly
				155					160					165
Glu	Leu	Asp	Leu	Leu	Phe	Met	Ser	Leu	Arg	Asn	Ala	Cys	Ile	
				170					175					

<210> SEQ ID NO 155
<211> LENGTH: 1320
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 155

ggcttgctga aaataaaatc aggactccta acctgctcca gtcagcctgc	50
ttccacgagg cctgtcagtc agtgcccgac ttgtgactga gtgtgcagtg	100
cccagcatgt accaggtcag tgcagagggc tgcctgaggg ctgtgctgag	150
agggagagga gcagagatgc tgctgagggt ggagggaggc caagctgcc	200
ggtttggggc tgggggccaa gtggagttag aaactgggat cccaggggga	250
gggtgcagat gagggagcga cccagattag gtgaggacag ttctctcatt	300
agccttttcc tacaggtggt tgcattcttg gcaatggtca tgggaacca	350
cacctacagc cactggccca gctgctgccc cagcaaaggg caggacacct	400
ctgaggagct gctgaggtgg agcactgtgc ctgtgcctcc cctagagcct	450
gctaggccca accgccccc agagtctgtg agggccagtg aagatggacc	500
cctcaacagc agggccatct ccccctggag atatgagttg gacagagact	550
tgaaccggct cccccaggac ctgtaccacg ccggttgctt gtgcccgcac	600
tgcgctcagc tacagacagg ctcccacatg gacccccggg gcaactcgga	650
gctgctctac cacaaccaga ctgtcttcta caggcggccca tgccatggcg	700
agaagggcac ccacaagggc tactgccttg agcgcaggct gtaccgtgtt	750
tccttagctt gtgtgtgtgt gcggccccgt gtgatgggct agccggacct	800
gctggaggct ggtccctttt tgggaaacct ggagccaggt gtacaaccac	850
ttgccatgaa gggccaggat gccagatgc ttggcccctg tgaagtgtctg	900
tctggagcag caggatcccc ggacaggatg gggggctttg gggaaaacct	950
gcacttctgc acattttgaa aagagcagct gctgcttagg gccgcgggaa	1000
gctggtgtcc tgtcattttc tctcaggaaa ggttttcaaa gttctgccca	1050
tttctggagg ccaccactcc tgtctcttcc tcttttccca tcccctgcta	1100
ccctggccca gcacaggcac tttctagata tttccccctt gctggagaag	1150
aaagagcccc tggttttatt tgtttgttta ctcatcactc agtgagcatc	1200
tactttgggt gcattctagt gtagttacta gtcttttgac atggatgatt	1250
ctgaggagga agctgttatt gaatgtatag agatttatcc aaataaatat	1300
ctttatttaa aatgaaaaa	1320

-continued

<210> SEQ ID NO 156
<211> LENGTH: 177
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 156

Met	Arg	Glu	Arg	Pro	Arg	Leu	Gly	Glu	Asp	Ser	Ser	Leu	Ile	Ser
1				5					10					15
Leu	Phe	Leu	Gln	Val	Val	Ala	Phe	Leu	Ala	Met	Val	Met	Gly	Thr
			20						25					30
His	Thr	Tyr	Ser	His	Trp	Pro	Ser	Cys	Cys	Pro	Ser	Lys	Gly	Gln
			35						40					45
Asp	Thr	Ser	Glu	Glu	Leu	Leu	Arg	Trp	Ser	Thr	Val	Pro	Val	Pro
			50						55					60
Pro	Leu	Glu	Pro	Ala	Arg	Pro	Asn	Arg	His	Pro	Glu	Ser	Cys	Arg
			65						70					75
Ala	Ser	Glu	Asp	Gly	Pro	Leu	Asn	Ser	Arg	Ala	Ile	Ser	Pro	Trp
			80						85					90
Arg	Tyr	Glu	Leu	Asp	Arg	Asp	Leu	Asn	Arg	Leu	Pro	Gln	Asp	Leu
			95						100					105
Tyr	His	Ala	Arg	Cys	Leu	Cys	Pro	His	Cys	Val	Ser	Leu	Gln	Thr
			110						115					120
Gly	Ser	His	Met	Asp	Pro	Arg	Gly	Asn	Ser	Glu	Leu	Leu	Tyr	His
			125						130					135
Asn	Gln	Thr	Val	Phe	Tyr	Arg	Arg	Pro	Cys	His	Gly	Glu	Lys	Gly
			140						145					150
Thr	His	Lys	Gly	Tyr	Cys	Leu	Glu	Arg	Arg	Leu	Tyr	Arg	Val	Ser
			155						160					165
Leu	Ala	Cys	Val	Cys	Val	Arg	Pro	Arg	Val	Met	Gly			
			170						175					

<210> SEQ ID NO 157
<211> LENGTH: 1515
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 157

ccggcgatgt cgctcgtgct gctaagcctg gccgcgctgt gcaggagcgc	50
cgtaacccca gagccgaccg ttcaatgtgg ctctgaaact gggccatctc	100
cagagtggat gctacaacat gatctaattcc ccggagactt gagggacctc	150
cgagtagaac ctgttacaac tagtgttgca acaggggact attcaatttt	200
gatgaatgta agctgggtac tccgggcaga tgccagcatc cgcttgttga	250
aggccaccaa gatttgtgtg acgggcaaaa gcaacttcca gtcctacagc	300
tgtgtgaggt gcaattacac agaggccttc cagactcaga ccagaccctc	350
tggttggtaaa tggacatttt cctacatcgg cttccctgta gagctgaaca	400
cagtctattt cattggggcc cataatatcc ctaatgcaaa tatgaatgaa	450
gatggccctt ccatgtctgt gaatttcacc tcaccaggct gcctagacca	500
cataatgaaa tataaaaaaa agtgtgtcaa ggccggaagc ctgtgggatc	550
cgaacatcac tgcttgtaag aagaatgagg agacagtaga agtgaacttc	600
acaaccactc ccctgggaaa cagatacatg gctcttatcc aacacagcac	650

-continued

```
tatcatcggg ttttctcagg tgtttgagcc acaccagaag aaacaaacgc      700
gagcttcagt ggtgattcca gtgactgggg atagtgaagg tgctacggtg      750
cagctgactc catattttcc tacttggtgc agcgactgca tccgacataa      800
aggaacagtt gtgctctgcc cacaacacag cgtccctttc cctctggata      850
acaacaaaaa caagccggga ggctggctgc ctctcctcct gctgtctctg      900
ctggtgggcca catgggtgct ggtggcaggg atctatctaa tgtggaggca      950
cgaaaggatc aagaagactt ccttttctac caccacacta ctgcccccca     1000
ttaaggttct tgtggtttac ccattcgaaa tatgtttcca tcacacaatt     1050
tgttacttca ctgaatttct tcaaaaccat tgcagaagtg aggtcatcct     1100
tgaaaagtgg cagaaaaaga aaatagcaga gatgggtcca gtgcagtggc     1150
ttgccactca aaagaaggca gcagacaaag tcgtcttcct tctttccaat     1200
gacgtcaaca gtgtgtgcga tggtagctgt ggcaagagcg agggcagtc     1250
cagtgagaac tctcaagacc tcttccccct tgcctttaac cttttctgca     1300
gtgatctaag aagccagatt catctgcaca aatacgtggt ggtctacttt     1350
agagagattg atacaaaaga cgattacaat gctctcagtg tctgccccaa     1400
gtaccacctc atgaaggatg ccaactgcttt ctgtgcagaa cttctccatg     1450
tcaagcagca ggtgtcagca ggaaaagat cacaagcctg ccacgatggc     1500
tgctgctcct tgtag                                           1515
```

<210> SEQ ID NO 158

<211> LENGTH: 502

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 158

```
Met Ser Leu Val Leu Leu Ser Leu Ala Ala Leu Cys Arg Ser Ala
  1             5             10             15
Val Pro Arg Glu Pro Thr Val Gln Cys Gly Ser Glu Thr Gly Pro
 20             25             30
Ser Pro Glu Trp Met Leu Gln His Asp Leu Ile Pro Gly Asp Leu
 35             40             45
Arg Asp Leu Arg Val Glu Pro Val Thr Thr Ser Val Ala Thr Gly
 50             55             60
Asp Tyr Ser Ile Leu Met Asn Val Ser Trp Val Leu Arg Ala Asp
 65             70             75
Ala Ser Ile Arg Leu Leu Lys Ala Thr Lys Ile Cys Val Thr Gly
 80             85             90
Lys Ser Asn Phe Gln Ser Tyr Ser Cys Val Arg Cys Asn Tyr Thr
 95            100            105
Glu Ala Phe Gln Thr Gln Thr Arg Pro Ser Gly Gly Lys Trp Thr
110            115            120
Phe Ser Tyr Ile Gly Phe Pro Val Glu Leu Asn Thr Val Tyr Phe
125            130            135
Ile Gly Ala His Asn Ile Pro Asn Ala Asn Met Asn Glu Asp Gly
140            145            150
Pro Ser Met Ser Val Asn Phe Thr Ser Pro Gly Cys Leu Asp His
```

-continued

155	160	165
Ile Met Lys Tyr Lys Lys Lys Cys Val Lys Ala Gly Ser Leu Trp		
170	175	180
Asp Pro Asn Ile Thr Ala Cys Lys Lys Asn Glu Glu Thr Val Glu		
185	190	195
Val Asn Phe Thr Thr Thr Pro Leu Gly Asn Arg Tyr Met Ala Leu		
200	205	210
Ile Gln His Ser Thr Ile Ile Gly Phe Ser Gln Val Phe Glu Pro		
215	220	225
His Gln Lys Lys Gln Thr Arg Ala Ser Val Val Ile Pro Val Thr		
230	235	240
Gly Asp Ser Glu Gly Ala Thr Val Gln Leu Thr Pro Tyr Phe Pro		
245	250	255
Thr Cys Gly Ser Asp Cys Ile Arg His Lys Gly Thr Val Val Leu		
260	265	270
Cys Pro Gln Thr Gly Val Pro Phe Pro Leu Asp Asn Asn Lys Ser		
275	280	285
Lys Pro Gly Gly Trp Leu Pro Leu Leu Leu Leu Ser Leu Leu Val		
290	295	300
Ala Thr Trp Val Leu Val Ala Gly Ile Tyr Leu Met Trp Arg His		
305	310	315
Glu Arg Ile Lys Lys Thr Ser Phe Ser Thr Thr Thr Leu Leu Pro		
320	325	330
Pro Ile Lys Val Leu Val Val Tyr Pro Ser Glu Ile Cys Phe His		
335	340	345
His Thr Ile Cys Tyr Phe Thr Glu Phe Leu Gln Asn His Cys Arg		
350	355	360
Ser Glu Val Ile Leu Glu Lys Trp Gln Lys Lys Lys Ile Ala Glu		
365	370	375
Met Gly Pro Val Gln Trp Leu Ala Thr Gln Lys Lys Ala Ala Asp		
380	385	390
Lys Val Val Phe Leu Leu Ser Asn Asp Val Asn Ser Val Cys Asp		
395	400	405
Gly Thr Cys Gly Lys Ser Glu Gly Ser Pro Ser Glu Asn Ser Gln		
410	415	420
Asp Leu Phe Pro Leu Ala Phe Asn Leu Phe Cys Ser Asp Leu Arg		
425	430	435
Ser Gln Ile His Leu His Lys Tyr Val Val Val Tyr Phe Arg Glu		
440	445	450
Ile Asp Thr Lys Asp Asp Tyr Asn Ala Leu Ser Val Cys Pro Lys		
455	460	465
Tyr His Leu Met Lys Asp Ala Thr Ala Phe Cys Ala Glu Leu Leu		
470	475	480
His Val Lys Gln Gln Val Ser Ala Gly Lys Arg Ser Gln Ala Cys		
485	490	495
His Asp Gly Cys Cys Ser Leu		
500		

<210> SEQ ID NO 159

<211> LENGTH: 535

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 159

gccaccagc gcaacatgac agtgaagacc ctgcatggcc cagccatggt	50
caagtacttg ctgctgtcga tattggggct tgcctttctg agtgaggcgg	100
cagctcgaa aatcccaaa gtaggacata cttttttcca aaagcctgag	150
agttgccgc ctgtgccagg aggtagtatg aagcttgaca ttggcatcat	200
caatgaaaac cagcgcgttt ccatgtcacg taacatcgag agccgctcca	250
cctcccccctg gaattacact gtcacttggg accccaaccg gtaccctctg	300
gaagttgtac agggccagtg taggaacttg ggctgcatca atgctcaagg	350
aaaggaagac atctccatga attccgttcc catccagcaa gagaccctgg	400
tcgtccggag gaagcaccaa ggctgctctg tttctttcca gttggagaag	450
gtgctggtga ctgttggtg cactgcgctc acccctgtca tccaccatgt	500
gcagtaagag gtgcatatcc actcagctga agaag	535

<210> SEQ ID NO 160

<211> LENGTH: 163

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 160

Met Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu	
1 5 10 15	
Leu Leu Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Ala	
20 25 30	
Arg Lys Ile Pro Lys Val Gly His Thr Phe Phe Gln Lys Pro Glu	
35 40 45	
Ser Cys Pro Pro Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly	
50 55 60	
Ile Ile Asn Glu Asn Gln Arg Val Ser Met Ser Arg Asn Ile Glu	
65 70 75	
Ser Arg Ser Thr Ser Pro Trp Asn Tyr Thr Val Thr Trp Asp Pro	
80 85 90	
Asn Arg Tyr Pro Ser Glu Val Val Gln Ala Gln Cys Arg Asn Leu	
95 100 105	
Gly Cys Ile Asn Ala Gln Gly Lys Glu Asp Ile Ser Met Asn Ser	
110 115 120	
Val Pro Ile Gln Gln Glu Thr Leu Val Val Arg Arg Lys His Gln	
125 130 135	
Gly Cys Ser Val Ser Phe Gln Leu Glu Lys Val Leu Val Thr Val	
140 145 150	
Gly Cys Thr Cys Val Thr Pro Val Ile His His Val Gln	
155 160	

<210> SEQ ID NO 161

<211> LENGTH: 2380

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 161

acactggcca aacaaaaacg aaagcactcc gtgctggaag taggaggaga	50
--	----

-continued

gtcaggactc ccaggacaga gagtgcacaa actaccacgc acagccccct	100
ccgccccctc tggaggctga agagggatc cagcccctgc caccacaga	150
cacgggctga ctggggtgtc tgccccctt gggggggggc agcacaggc	200
ctcaggcctg ggtgccacct ggcacctaga agatgcctgt gccctggttc	250
ttgctgtcct tggcactggg ccgaagccca gtggtccttt ctctggagag	300
gcttgtgggg cctcaggacg ctaccactg ctctccgggc ctctcctgcc	350
gcctctggga cagtgcata ctctgcctgc ctggggacat cgtgcctgct	400
ccgggccccg tgctggcgcc tacgcacctg cagacagagc tggtgctgag	450
gtgcagaag gagaccgact gtgacctctg tctgcgtgtg gctgtccact	500
tggcgtgca tgggactgg gaagagcctg aagatgagga aaagtttgga	550
ggagcagctg actcaggggt ggaggagcct aggaatgcct ctctccaggc	600
caaagtcgtg ctctccttcc aggcctaccc tactgccgcg tgcgtcctgc	650
tggaggtgca agtgcctgct gcccttgtgc agtttggtca gtctgtgggc	700
tctgtggtat atgactgctt cgaggctgcc ctagggagtg aggtacgaat	750
ctggtcctat actcagccca ggtacgagaa ggaactcaac cacacacagc	800
agctgcctgc cctgccctgg ctcaacgtgt cagcagatgg tgacaacgtg	850
catctgggtc tgaatgtctc tgaggagcag cacttcggcc tctccctgta	900
ctggaatcag gtccaggggc ccccaaaacc ccggtggcac aaaaacctga	950
ctggaccgca gatcattacc ttgaaccaca cagacctggt tccctgctc	1000
tgtattcagg tgtggcctct ggaacctgac tccgttagga cgaacatctg	1050
ccccttcagg gaggaccccc gcgcacacca gaacctctgg caagccgccc	1100
gactgcgaat gctgaccctg cagagctggc tgctggagcg accgtgctcg	1150
ctgcccgcag aagcggaact gtgctggcgg gctccgggtg gggacccctg	1200
ccagccactg gtcccaccgc ttctctggga gaacgtcact gtggacaagg	1250
ttctcgagtt ccattgctg aaaggccacc ctaacctctg tgttcagggt	1300
aacagctcgg agaagctgca gctgcaggag tgcttggtgg ctgactccct	1350
ggggcctctc aaagacgatg tgctactgtt ggagacacga gggccccagg	1400
acaacagatc cctctgtgcc ttggaaccca gtgctgttac ttcactacc	1450
agcaaaacct ccacgagggc agctcgcctt ggagagtact tactacaaga	1500
cctgcagtca ggccagtgtc tgcagctatg ggacgatgac ttgggagcgc	1550
tatgggcctg ccccatggac aaatacatcc acaagcgctg ggcctcgtg	1600
tggctggcct gcctactctt tgccgctgcg ctttccctca tcctccttct	1650
caaaaaggat cagcgaaaag ggtggctgag gctcttgaaa caggacgtcc	1700
gctcgggggc ggccgccagg ggccgcgcgg ctctgctcct ctactcagcc	1750
gatgactcgg gtttcgagcg cctgggtggc gccctggcgt cggccctgtg	1800
ccagctgcog ctgcgcgtgg ccgtagacct gtggagccgt cgtgaactga	1850
gcgcgcaggg gcccgtagct tggtttcacg cgcagcgcg ccagacctg	1900
caggaggggc gcgtggtggt ctgtctcttc tctcccggtg cggtggcgct	1950

-continued

gtgcagcgag tggctacagg atgggggtgtc cgggcccggg gcgcacggcc	2000
cgcacgacgc cttccgcgcc tcgctcagct gcgtgctgcc cgactttctg	2050
cagggccggg cgcccggcag ctacgtgggg gcctgcttcg acaggctgct	2100
ccaccgggac gccgtaccgg cccttttccg caccgtgccc gtcttcacac	2150
tgccctccca actgccagac ttcttggggg ccctgcagca gcctcgcgcc	2200
ccgcgttcgg ggcgggtcca agagagagcg gagcaagtgt cccggggccct	2250
tcagccagcc ctggatagct acttccatcc cccggggact cccgcgccgg	2300
gacgcgggggt gggaccaggg gcggggacctg gggcggggga cgggacttaa	2350
ataaaggcag acgctgtttt tctaaaaaaa	2380

<210> SEQ ID NO 162

<211> LENGTH: 705

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 162

Met	Pro	Val	Pro	Trp	Phe	Leu	Leu	Ser	Leu	Ala	Leu	Gly	Arg	Ser	1	5	10	15
Pro	Val	Val	Leu	Ser	Leu	Glu	Arg	Leu	Val	Gly	Pro	Gln	Asp	Ala	20	25	30	
Thr	His	Cys	Ser	Pro	Gly	Leu	Ser	Cys	Arg	Leu	Trp	Asp	Ser	Asp	35	40	45	
Ile	Leu	Cys	Leu	Pro	Gly	Asp	Ile	Val	Pro	Ala	Pro	Gly	Pro	Val	50	55	60	
Leu	Ala	Pro	Thr	His	Leu	Gln	Thr	Glu	Leu	Val	Leu	Arg	Cys	Gln	65	70	75	
Lys	Glu	Thr	Asp	Cys	Asp	Leu	Cys	Leu	Arg	Val	Ala	Val	His	Leu	80	85	90	
Ala	Val	His	Gly	His	Trp	Glu	Glu	Pro	Glu	Asp	Glu	Glu	Lys	Phe	95	100	105	
Gly	Gly	Ala	Ala	Asp	Ser	Gly	Val	Glu	Glu	Pro	Arg	Asn	Ala	Ser	110	115	120	
Leu	Gln	Ala	Gln	Val	Val	Leu	Ser	Phe	Gln	Ala	Tyr	Pro	Thr	Ala	125	130	135	
Arg	Cys	Val	Leu	Leu	Glu	Val	Gln	Val	Pro	Ala	Ala	Leu	Val	Gln	140	145	150	
Phe	Gly	Gln	Ser	Val	Gly	Ser	Val	Val	Tyr	Asp	Cys	Phe	Glu	Ala	155	160	165	
Ala	Leu	Gly	Ser	Glu	Val	Arg	Ile	Trp	Ser	Tyr	Thr	Gln	Pro	Arg	170	175	180	
Tyr	Glu	Lys	Glu	Leu	Asn	His	Thr	Gln	Gln	Leu	Pro	Ala	Leu	Pro	185	190	195	
Trp	Leu	Asn	Val	Ser	Ala	Asp	Gly	Asp	Asn	Val	His	Leu	Val	Leu	200	205	210	
Asn	Val	Ser	Glu	Glu	Gln	His	Phe	Gly	Leu	Ser	Leu	Tyr	Trp	Asn	215	220	225	
Gln	Val	Gln	Gly	Pro	Pro	Lys	Pro	Arg	Trp	His	Lys	Asn	Leu	Thr	230	235	240	
Gly	Pro	Gln	Ile	Ile	Thr	Leu	Asn	His	Thr	Asp	Leu	Val	Pro	Cys	245	250	255	

-continued

Leu Cys Ile Gln Val Trp Pro Leu Glu	Pro Asp Ser Val Arg Thr
260	265 270
Asn Ile Cys Pro Phe Arg Glu Asp Pro	Arg Ala His Gln Asn Leu
275	280 285
Trp Gln Ala Ala Arg Leu Arg Leu Leu	Thr Leu Gln Ser Trp Leu
290	295 300
Leu Asp Ala Pro Cys Ser Leu Pro Ala	Glu Ala Ala Leu Cys Trp
305	310 315
Arg Ala Pro Gly Gly Asp Pro Cys Gln	Pro Leu Val Pro Pro Leu
320	325 330
Ser Trp Glu Asn Val Thr Val Asp Lys	Val Leu Glu Phe Pro Leu
335	340 345
Leu Lys Gly His Pro Asn Leu Cys Val	Gln Val Asn Ser Ser Glu
350	355 360
Lys Leu Gln Leu Gln Glu Cys Leu Trp	Ala Asp Ser Leu Gly Pro
365	370 375
Leu Lys Asp Asp Val Leu Leu Leu Glu	Thr Arg Gly Pro Gln Asp
380	385 390
Asn Arg Ser Leu Cys Ala Leu Glu Pro	Ser Gly Cys Thr Ser Leu
395	400 405
Pro Ser Lys Ala Ser Thr Arg Ala Ala	Arg Leu Gly Glu Tyr Leu
410	415 420
Leu Gln Asp Leu Gln Ser Gly Gln Cys	Leu Gln Leu Trp Asp Asp
425	430 435
Asp Leu Gly Ala Leu Trp Ala Cys Pro	Met Asp Lys Tyr Ile His
440	445 450
Lys Arg Trp Ala Leu Val Trp Leu Ala	Cys Leu Leu Phe Ala Ala
455	460 465
Ala Leu Ser Leu Ile Leu Leu Leu Lys	Lys Asp His Ala Lys Gly
470	475 480
Trp Leu Arg Leu Leu Lys Gln Asp Val	Arg Ser Gly Ala Ala Ala
485	490 495
Arg Gly Arg Ala Ala Leu Leu Leu Tyr	Ser Ala Asp Asp Ser Gly
500	505 510
Phe Glu Arg Leu Val Gly Ala Leu Ala	Ser Ala Leu Cys Gln Leu
515	520 525
Pro Leu Arg Val Ala Val Asp Leu Trp	Ser Arg Arg Glu Leu Ser
530	535 540
Ala Gln Gly Pro Val Ala Trp Phe His	Ala Gln Arg Arg Gln Thr
545	550 555
Leu Gln Glu Gly Gly Val Val Val Leu	Leu Phe Ser Pro Gly Ala
560	565 570
Val Ala Leu Cys Ser Glu Trp Leu Gln	Asp Gly Val Ser Gly Pro
575	580 585
Gly Ala His Gly Pro His Asp Ala Phe	Arg Ala Ser Leu Ser Cys
590	595 600
Val Leu Pro Asp Phe Leu Gln Gly Arg	Ala Pro Gly Ser Tyr Val
605	610 615
Gly Ala Cys Phe Asp Arg Leu Leu His	Pro Asp Ala Val Pro Ala
620	625 630

-continued

Leu	Phe	Arg	Thr	Val	Pro	Val	Phe	Thr	Leu	Pro	Ser	Gln	Leu	Pro
				635					640					645
Asp	Phe	Leu	Gly	Ala	Leu	Gln	Gln	Pro	Arg	Ala	Pro	Arg	Ser	Gly
				650					655					660
Arg	Leu	Gln	Glu	Arg	Ala	Glu	Gln	Val	Ser	Arg	Ala	Leu	Gln	Pro
				665					670					675
Ala	Leu	Asp	Ser	Tyr	Phe	His	Pro	Pro	Gly	Thr	Pro	Ala	Pro	Gly
				680					685					690
Arg	Gly	Val	Gly	Pro	Gly	Ala	Gly	Pro	Gly	Ala	Gly	Asp	Gly	Thr
				695					700					705

<210> SEQ ID NO 163
<211> LENGTH: 2478
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 163

gtcagtgctgaggccgggtc agccaccaag atgactgaca ggttcagctc	50
tctgcagcac actaccctca agccacctga tgtgacctgt atctccaaag	100
tgagatcgat tcagatgatt gttcatccta cccccacgcc aatccgtgca	150
ggcgtatggcc accggctaac cctggaagac atcttccatg acctgttcta	200
ccacttagag ctccaggtca accgcacctc ccaaatgcac cttggaggga	250
agcagagaga atatgagttc ttccggcctga cccctgacac agagttcctt	300
ggcaccatca tgatttgctg tcccacctgg gcccaaggaga gtgcccccta	350
catgtgccga gtgaagacac tgccagaccg gacatggacc tactccttct	400
ccggagcctt cctgttctcc atgggcttcc tcgtgcagct actctgctac	450
ctgagctaca gatatgtcac caagccgcct gcacctccca actccctgaa	500
cgtccagcga gtcctgactt tccagccgct gcgcttcac caggagcacg	550
tcctgatccc tgtctttgac ctccagcgcc ccagcagtct ggcccagcct	600
gtccagtact ccagatcag ggtgtctgga ccagggagc ccgcaggagc	650
tccacagcgg catagcctgt ccgagatcac ctacttaggg cagccagaca	700
tctccatcct ccagccctcc aacgtgccac ctccccagat cctctcccca	750
ctgtcctatg ccccaaacgc tgcccctgag gtcgggcccc catcctatgc	800
acctcaggtg acccccgaag ctcaattccc attctacgcc ccacaggcca	850
tctctaaggt ccagccttcc tcctatgccc ctcaagccac tccggacagc	900
tggcctccct cctatggggt atgcatggaa ggttctggca aagactcccc	950
cactgggaca ctttctagtc ctaaacacct taggcctaaa ggtcagcttc	1000
agaaagagcc accagctgga agctgcatgt taggtggcct ttctctgcag	1050
gaggtgacct ccttggtat ggaggaatcc caagaagcaa aatcattgca	1100
ccagcccctg gggatttgca cagacagaac atctgaccca aatgtgctac	1150
acagtgggga ggaagggaca ccacagtacc taaagggccca gctccccctc	1200
ctctcctcag tccagatcga gggccacccc atgtccctcc ctttgcaacc	1250
tccttccggt ccatgttccc cctcggaacca aggtccaagt ccttggggcc	1300
tgctggagtc ccttgtgtgt cccaaggatg aagccaagag cccagcccct	1350

-continued

```
gagacctcag acctggagca gcccacagaa ctggattctc ttttcagagg      1400
cctggccctg actgtgcagt gggagtcctg aggggaatgg gaaaggcttg      1450
gtgcttcctc cctgtcccta cccagtgtca catccttggc tgtcaatccc      1500
atgcctgccc atgccacaca ctctgcgacg tggcctcaga cgggtgccct      1550
tgagagaagc agagggagtg gcatgcaggg cccctgccat gggtgcgctc      1600
ctcaccggaa caaagcagca tgataaggac tgcagcgggg gagctctggg      1650
gagcagcttg tgtagacaag cgcgtgctcg ctgagccctg caaggcagaa      1700
atgacagtgc aaggaggaaa tgcagggaaa ctcccagagt ccagagcccc      1750
acctcctaac accatggatt caaagtgctc agggaatttg cctctccttg      1800
ccccattcct ggccagtttc acaatctagc tcgacagagc atgaggcccc      1850
tgccctcttc gtcattgttc aaagtgggga agagagcctg gaaaagaacc      1900
aggcctggaa aagaaccaga aggaggctgg gcagaaccag aacaacctgc      1950
acttctgcc aagccagggc cagcaggacg gcaggactct agggaggggt      2000
gtggcctgca gctcattccc agccagggca actgcctgac gttgcacgat      2050
ttcagcttca ttcctctgat agaacaaagc gaaatgcagg tccaccaggg      2100
agggagacac acaagccttt tctgcaggca ggagtttcag accctatcct      2150
gagaatgggg tttgaaagga aggtgagggc tgtggcccct ggacgggtac      2200
aataacacac tgtactgatg tcacaacttt gcaagctctg ccttgggttc      2250
agcccatctg ggctcaaatt ccagcctcac cactcacaag ctgtgtgact      2300
tcaaacaaat gaaatcagtg cccgaacctc cggtttcctc atctgtaatg      2350
tggggatcat aacacctacc tcatggagtt gtggtgaaga tgaaatgaag      2400
tcatgtcttt aaagtgttta atagtgcctg gtacatgggc agtgcccaat      2450
aaacggtagc tatttaaaaa aaaaaaaa      2478
```

<210> SEQ ID NO 164
<211> LENGTH: 574
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 164

```
Met Arg Thr Leu Leu Thr Ile Leu Thr Val Gly Ser Leu Ala Ala      1 5 10 15
His Ala Pro Glu Asp Pro Ser Asp Leu Leu Gln His Val Lys Phe      20 25 30
Gln Ser Ser Asn Phe Glu Asn Ile Leu Thr Trp Asp Ser Gly Pro      35 40 45
Glu Gly Thr Pro Asp Thr Val Tyr Ser Ile Glu Tyr Lys Thr Tyr      50 55 60
Gly Glu Arg Asp Trp Val Ala Lys Lys Gly Cys Gln Arg Ile Thr      65 70 75
Arg Lys Ser Cys Asn Leu Thr Val Glu Thr Gly Asn Leu Thr Glu      80 85 90
Leu Tyr Tyr Ala Arg Val Thr Ala Val Ser Ala Gly Gly Arg Ser      95 100 105
```

-continued

Ala Thr Lys Met Thr Asp Arg Phe Ser Ser Leu Gln His Thr Thr	110	115	120
Leu Lys Pro Pro Asp Val Thr Cys Ile Ser Lys Val Arg Ser Ile	125	130	135
Gln Met Ile Val His Pro Thr Pro Thr Pro Ile Arg Ala Gly Asp	140	145	150
Gly His Arg Leu Thr Leu Glu Asp Ile Phe His Asp Leu Phe Tyr	155	160	165
His Leu Glu Leu Gln Val Asn Arg Thr Tyr Gln Met His Leu Gly	170	175	180
Gly Lys Gln Arg Glu Tyr Glu Phe Phe Gly Leu Thr Pro Asp Thr	185	190	195
Glu Phe Leu Gly Thr Ile Met Ile Cys Val Pro Thr Trp Ala Lys	200	205	210
Glu Ser Ala Pro Tyr Met Cys Arg Val Lys Thr Leu Pro Asp Arg	215	220	225
Thr Trp Thr Tyr Ser Phe Ser Gly Ala Phe Leu Phe Ser Met Gly	230	235	240
Phe Leu Val Ala Val Leu Cys Tyr Leu Ser Tyr Arg Tyr Val Thr	245	250	255
Lys Pro Pro Ala Pro Pro Asn Ser Leu Asn Val Gln Arg Val Leu	260	265	270
Thr Phe Gln Pro Leu Arg Phe Ile Gln Glu His Val Leu Ile Pro	275	280	285
Val Phe Asp Leu Ser Gly Pro Ser Ser Leu Ala Gln Pro Val Gln	290	295	300
Tyr Ser Gln Ile Arg Val Ser Gly Pro Arg Glu Pro Ala Gly Ala	305	310	315
Pro Gln Arg His Ser Leu Ser Glu Ile Thr Tyr Leu Gly Gln Pro	320	325	330
Asp Ile Ser Ile Leu Gln Pro Ser Asn Val Pro Pro Pro Gln Ile	335	340	345
Leu Ser Pro Leu Ser Tyr Ala Pro Asn Ala Ala Pro Glu Val Gly	350	355	360
Pro Pro Ser Tyr Ala Pro Gln Val Thr Pro Glu Ala Gln Phe Pro	365	370	375
Phe Tyr Ala Pro Gln Ala Ile Ser Lys Val Gln Pro Ser Ser Tyr	380	385	390
Ala Pro Gln Ala Thr Pro Asp Ser Trp Pro Pro Ser Tyr Gly Val	395	400	405
Cys Met Glu Gly Ser Gly Lys Asp Ser Pro Thr Gly Thr Leu Ser	410	415	420
Ser Pro Lys His Leu Arg Pro Lys Gly Gln Leu Gln Lys Glu Pro	425	430	435
Pro Ala Gly Ser Cys Met Leu Gly Gly Leu Ser Leu Gln Glu Val	440	445	450
Thr Ser Leu Ala Met Glu Glu Ser Gln Glu Ala Lys Ser Leu His	455	460	465
Gln Pro Leu Gly Ile Cys Thr Asp Arg Thr Ser Asp Pro Asn Val	470	475	480
Leu His Ser Gly Glu Glu Gly Thr Pro Gln Tyr Leu Lys Gly Gln			

-continued

				485					490					495
Leu	Pro	Leu	Leu	Ser	Ser	Val	Gln	Ile	Glu	Gly	His	Pro	Met	Ser
				500					505					510
Leu	Pro	Leu	Gln	Pro	Pro	Ser	Gly	Pro	Cys	Ser	Pro	Ser	Asp	Gln
				515					520					525
Gly	Pro	Ser	Pro	Trp	Gly	Leu	Leu	Glu	Ser	Leu	Val	Cys	Pro	Lys
				530					535					540
Asp	Glu	Ala	Lys	Ser	Pro	Ala	Pro	Glu	Thr	Ser	Asp	Leu	Glu	Gln
				545					550					555
Pro	Thr	Glu	Leu	Asp	Ser	Leu	Phe	Arg	Gly	Leu	Ala	Leu	Thr	Val
				560					565					570
Gln	Trp	Glu	Ser											

<210> SEQ ID NO 165
<211> LENGTH: 1060
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 165

tgccctactg	gaaaaaaaa	aaaaaaaaa	aaaagtcacc	cgggcccgcg	50
gtggccacaa	catggctgcg	gcgccggggc	tgctcttctg	gctgttcgtg	100
ctggggggcg	tctggtgggt	cccgggccag	tcggatctca	gccacggacg	150
gcgtttctcg	gacctcaaag	tgtgcgggga	cgaagagtgc	agcatgttaa	200
tgtaccgtgg	gaaagctctt	gaagacttca	cgggccctga	ttgtcgtttt	250
gtgaatttta	aaaaagtgta	cgatgtatat	gtctactaca	aactggcagg	300
gggatccctt	gaactttggg	ctggaagtgt	tgaacacagt	tttggatatt	350
ttccaaaaga	tttgatcaag	gtacttcata	aatacacgga	agaagagcta	400
catattccag	cagatgagac	agactttgtc	tgctttgaag	gaggaagaga	450
tgattttaat	agttataatg	tagaagagct	tttaggatct	ttggaactgg	500
aggactctgt	acctgaagag	tcgaagaaag	ctgaagaagt	ttctcagcac	550
agagagaaat	ctcctgagga	gtctcggggg	cgtgaacttg	accctgtgcc	600
tgagcccag	gcattcagag	ctgattcaga	ggatggagaa	ggtgctttct	650
cagagagcac	cgaggggctg	cagggacagc	cctcagctca	ggagagccac	700
cctcacacca	gcggtcctgc	ggctaacgct	cagggagtgc	agtcttcggt	750
ggacactttt	gaagaaattc	tgcacgataa	attgaaagtg	ccgggaagcg	800
aaagcagaac	tggcaatagt	tctcctgcct	cggtggagcg	ggagaagaca	850
gatgcttaca	aagtcctgaa	aacagaaatg	agtcagagag	gaagtggaca	900
gtgcgttatt	cattacagca	aaggatttcg	ttggcatcaa	aatctaagtt	950
tgttttacaa	agattgtttt	tagtactaag	ctgccttggc	agtttgcatt	1000
tttgagccaa	acaaaaatat	attattttcc	cttctaagta	aaaaaaaaaa	1050
aaaaaaaaaa					1060

<210> SEQ ID NO 166
<211> LENGTH: 303
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

-continued

<400> SEQUENCE: 166

Met	Ala	Ala	Ala	Pro	Gly	Leu	Leu	Phe	Trp	Leu	Phe	Val	Leu	Gly
1				5					10					15
Ala	Leu	Trp	Trp	Val	Pro	Gly	Gln	Ser	Asp	Leu	Ser	His	Gly	Arg
				20					25					30
Arg	Phe	Ser	Asp	Leu	Lys	Val	Cys	Gly	Asp	Glu	Glu	Cys	Ser	Met
				35					40					45
Leu	Met	Tyr	Arg	Gly	Lys	Ala	Leu	Glu	Asp	Phe	Thr	Gly	Pro	Asp
				50					55					60
Cys	Arg	Phe	Val	Asn	Phe	Lys	Lys	Gly	Asp	Asp	Val	Tyr	Val	Tyr
				65					70					75
Tyr	Lys	Leu	Ala	Gly	Gly	Ser	Leu	Glu	Leu	Trp	Ala	Gly	Ser	Val
				80					85					90
Glu	His	Ser	Phe	Gly	Tyr	Phe	Pro	Lys	Asp	Leu	Ile	Lys	Val	Leu
				95					100					105
His	Lys	Tyr	Thr	Glu	Glu	Glu	Leu	His	Ile	Pro	Ala	Asp	Glu	Thr
				110					115					120
Asp	Phe	Val	Cys	Phe	Glu	Gly	Gly	Arg	Asp	Asp	Phe	Asn	Ser	Tyr
				125					130					135
Asn	Val	Glu	Glu	Leu	Leu	Gly	Ser	Leu	Glu	Leu	Glu	Asp	Ser	Val
				140					145					150
Pro	Glu	Glu	Ser	Lys	Lys	Ala	Glu	Glu	Val	Ser	Gln	His	Arg	Glu
				155					160					165
Lys	Ser	Pro	Glu	Glu	Ser	Arg	Gly	Arg	Glu	Leu	Asp	Pro	Val	Pro
				170					175					180
Glu	Pro	Glu	Ala	Phe	Arg	Ala	Asp	Ser	Glu	Asp	Gly	Glu	Gly	Ala
				185					190					195
Phe	Ser	Glu	Ser	Thr	Glu	Gly	Leu	Gln	Gly	Gln	Pro	Ser	Ala	Gln
				200					205					210
Glu	Ser	His	Pro	His	Thr	Ser	Gly	Pro	Ala	Ala	Asn	Ala	Gln	Gly
				215					220					225
Val	Gln	Ser	Ser	Leu	Asp	Thr	Phe	Glu	Glu	Ile	Leu	His	Asp	Lys
				230					235					240
Leu	Lys	Val	Pro	Gly	Ser	Glu	Ser	Arg	Thr	Gly	Asn	Ser	Ser	Pro
				245					250					255
Ala	Ser	Val	Glu	Arg	Glu	Lys	Thr	Asp	Ala	Tyr	Lys	Val	Leu	Lys
				260					265					270
Thr	Glu	Met	Ser	Gln	Arg	Gly	Ser	Gly	Gln	Cys	Val	Ile	His	Tyr
				275					280					285
Ser	Lys	Gly	Phe	Arg	Trp	His	Gln	Asn	Leu	Ser	Leu	Phe	Tyr	Lys
				290					295					300

Asp Cys Phe

<210> SEQ ID NO 167
<211> LENGTH: 2570
<212> TYPE: DNA
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 167

ccaggaccag	ggcgaccgg	ctcagcctct	cacttgtcag	aggccgggga	50
agagaagcaa	agcgcaacgg	tgtggtccaa	gccggggcctt	ctgcttcgcc	100

-continued

tctaggacat acacgggacc ccctaacttc agtcccccaa acgcgcaccc	150
tcgaagtctt gaactccagc cccgcacatc cacgcgcggc acaggcgcgg	200
caggcggcag gtcccggccg aaggcgatgc gcgcaggggg tcgggcagct	250
gggctcgggc ggcgggagta gggcccggca gggaggcagg gaggctgcat	300
attcagagtc gcgggctgcg ccctggggcag aggccgccct cgctccacgc	350
aacacctgct gctgccaccg cgccgcgatg agccgcgtgg tctcgctgct	400
gctggggccc gcgctgctct gcggccacgg agccttctgc cgccgcgtgg	450
tcagcggcca aaaggtgtgt tttgctgact tcaagcatcc ctgctacaaa	500
atggcctact tccatgaact gtccagccga gtgagctttc aggaggcacg	550
cctggcttgt gagagtgagg gaggagtctt cctcagcctt gagaatgaag	600
cagaacagaa gttaatagag agcatgttgc aaaacctgac aaaaccggg	650
acagggattt ctgatggtga tttctggata gggctttgga ggaatggaga	700
tgggcaaaaa tctggtgcct gccagatct ctaccagtgg tctgatggaa	750
gcaattccca gtaccgaaac tggtagacag atgaaccttc ctgcggaagt	800
gaaaagtgtg ttgtgatgta tcaccaacca actgccaatc ctggccttgg	850
gggtccctac ctttaccagt ggaatgatga cagggtgaac atgaagcaca	900
attatatttt caagtatgaa ccagagatta atccaacagc ccctgtagaa	950
aagccttato ttacaaatca accaggagac acccatcaga atgtggttgt	1000
tactgaagca ggtataattc ccaatccta tttatgttgtt ataccaacaa	1050
taccctgct cttactgata ctggttgctt ttggaacctg ttgtttccag	1100
atgtctgata aaagtaaagg aagaacaaaa actagtccaa accagtctac	1150
actgtggatt tcaaagagta ccagaaaaga aagtggcatg gaagtataat	1200
aactcattga cttggttcca gaattttgta attctggatc tgtataagga	1250
atggcatcag aacaatagct tggaatggct tgaaatcaca aaggatctgc	1300
aagatgaact gtaagctccc ccttgaggca aatattaaag taatttttat	1350
atgtctatta tttcatttaa agaatatgct gtgctaataa tggagtgaga	1400
catgcttatt ttgctaagg atgcacccaa acttcaaact tcaagcaaat	1450
gaaatggaca atgcagataa agttgttatc aacacgtcgg gagtatgtgt	1500
gttagaagca attcctttta tttctttcac ctttcataag ttgttatcta	1550
gtcaatgtaa tgtatatgtt attgaaattt acagtgtgca aaagtatttt	1600
acctttgcat aagtgtttga taaaaatgaa ctgttctaata atttattttt	1650
atggcatctc atttttcaat acatgctctt ttgattaaag aaacttatta	1700
ctgttgtcaa ctgaattcac acacacacaa atatagtacc atagaaaaag	1750
tttgttttct cgaaataatt catctttcag cttctctgct tttggccaat	1800
gtctaggaaa tctcttcaga aataagaagc tatttcatta agtgtgatat	1850
aaacctctc aaacatttta cttagaggca aggattgtct aatttcaatt	1900
gtgcaagaca tgtgccttat aattattttt agcttaaaat taaacagatt	1950
ttgtaataat gtaactttgt taataggtgc ataaacacta atgcagtcga	2000

-continued

```
tttgaacaaa agaagtgaca tacacaatat aaatcatatg tcttcacacg      2050
ttgcctatat atagagaagc agctctctga gggttctgaa atcaatgtgg      2100
tcctctcttt gccactaaa caaagatggg tggtcggggg ttgggattga      2150
cactggaggc agatagtgtc aaagttagtc taaggtttcc ctagctgtat      2200
ttagcctctg actatattag tatacaaaga ggtcatgtgg ttgagaccag      2250
gtgaatagtc actatcagtg tggagacaag cacagcacac agacatttta      2300
ggaaggaaa gaactacgaa atcgtgtgaa aatgggttgg aacccatcag      2350
tgatcgcata ttcattgatg agggtttgct tgagatagaa aatggtggct      2400
cctttctgtc ttatctccta gtttcttcaa tgcttacgcc ttgttcttct      2450
caagagaaa ttgtaactct ctggtcttca tatgtccctg tgctcctttt      2500
aaccataata agagttcttg tttctggggg aaaaaaaaaa aaaaaaaaaa      2550
aaaaaaaaa aaaaaaaaaa      2570
```

<210> SEQ ID NO 168

<211> LENGTH: 273

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 168

```
Met Ser Arg Val Val Ser Leu Leu Leu Gly Ala Ala Leu Leu Cys      15
  1          5          10
Gly His Gly Ala Phe Cys Arg Arg Val Val Ser Gly Gln Lys Val      30
  20          25          30
Cys Phe Ala Asp Phe Lys His Pro Cys Tyr Lys Met Ala Tyr Phe      45
  35          40          45
His Glu Leu Ser Ser Arg Val Ser Phe Gln Glu Ala Arg Leu Ala      60
  50          55          60
Cys Glu Ser Glu Gly Gly Val Leu Leu Ser Leu Glu Asn Glu Ala      75
  65          70          75
Glu Gln Lys Leu Ile Glu Ser Met Leu Gln Asn Leu Thr Lys Pro      90
  80          85          90
Gly Thr Gly Ile Ser Asp Gly Asp Phe Trp Ile Gly Leu Trp Arg     105
  95          100         105
Asn Gly Asp Gly Gln Thr Ser Gly Ala Cys Pro Asp Leu Tyr Gln     120
  110         115         120
Trp Ser Asp Gly Ser Asn Ser Gln Tyr Arg Asn Trp Tyr Thr Asp     135
  125         130         135
Glu Pro Ser Cys Gly Ser Glu Lys Cys Val Val Met Tyr His Gln     150
  140         145         150
Pro Thr Ala Asn Pro Gly Leu Gly Gly Pro Tyr Leu Tyr Gln Trp     165
  155         160         165
Asn Asp Asp Arg Cys Asn Met Lys His Asn Tyr Ile Cys Lys Tyr     180
  170         175         180
Glu Pro Glu Ile Asn Pro Thr Ala Pro Val Glu Lys Pro Tyr Leu     195
  185         190         195
Thr Asn Gln Pro Gly Asp Thr His Gln Asn Val Val Val Thr Glu     210
  200         205         210
Ala Gly Ile Ile Pro Asn Leu Ile Tyr Val Val Ile Pro Thr Ile
```

-continued																	
215						220						225					
Pro	Leu	Leu	Leu	Leu	Ile	Leu	Val	Ala	Phe	Gly	Thr	Cys	Cys	Phe			
230						235						240					
Gln	Met	Leu	His	Lys	Ser	Lys	Gly	Arg	Thr	Lys	Thr	Ser	Pro	Asn			
245						250						255					
Gln	Ser	Thr	Leu	Trp	Ile	Ser	Lys	Ser	Thr	Arg	Lys	Glu	Ser	Gly			
260						265						270					
Met Glu Val																	
<210> SEQ ID NO 169																	
<211> LENGTH: 43																	
<212> TYPE: DNA																	
<213> ORGANISM: Artificial Sequence																	
<220> FEATURE:																	
<223> OTHER INFORMATION: Synthetic oligonucleotide probe																	
<400> SEQUENCE: 169																	
tgtaaaacga cggccagtta aatagacctg caattattaa tct														43			
<210> SEQ ID NO 170																	
<211> LENGTH: 41																	
<212> TYPE: DNA																	
<213> ORGANISM: Artificial Sequence																	
<220> FEATURE:																	
<223> OTHER INFORMATION: Synthetic oligonucleotide probe																	
<400> SEQUENCE: 170																	
caggaaacag ctatgaccac ctgcacacct gcaaatccat t														41			

1. An isolated nucleic acid having at least 80% nucleic acid sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);

(b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;

(c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);

(d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;

(e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);

(f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or

(g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.

2. The isolated nucleic acid of claim 1 having at least 85% nucleic acid sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);

(b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
- (c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);

(d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;

(e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);

(f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or

(g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.

3. The isolated nucleic acid of claim 1 having at least 90% nucleic acid sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);

(b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;

(c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);

(d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;

- (e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);
 - (f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or
 - (g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.
4. The isolated nucleic acid of claim 1 having at least 95% nucleic acid sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);
 - (f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or
 - (g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.
5. The isolated nucleic acid of claim 1 having at least 99% nucleic acid sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);
 - (f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or
 - (g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.
6. An isolated nucleic acid comprising:
- (a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);
 - (f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or
 - (g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.
7. The isolated nucleic acid of claim 6 comprising a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96).
8. The isolated nucleic acid of claim 6 comprising a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide.
9. The isolated nucleic acid of claim 6 comprising a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96).
10. The isolated nucleic acid of claim 6 comprising a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide.
11. The isolated nucleic acid of claim 6 comprising the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95).
12. The isolated nucleic acid of claim 6 comprising the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95).
13. The isolated nucleic acid of claim 6 comprising the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.
14. An isolated nucleic acid that hybridizes to:
- (a) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (b) a nucleic acid sequence encoding the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (c) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96);
 - (d) a nucleic acid sequence encoding the extracellular domain of the polypeptide shown in **FIG. 96** (SEQ ID NO: 96), lacking its associated signal peptide;
 - (e) the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95);
 - (f) the full-length coding sequence of the nucleic acid sequence shown in **FIG. 95** (SEQ ID NO: 95); or
 - (g) the full-length coding sequence of the cDNA deposited under ATCC accession number 203279.
15. The isolated nucleic acid of claim 14, wherein said hybridization occurs under stringent conditions.
16. The isolated nucleic acid of claim 14 which is at least 10 nucleotides in length.
17. A vector comprising the nucleic acid of claim 1.
18. The vector of claim 17, wherein said nucleic acid is operably linked to control sequences recognized by a host cell transformed with the vector.
19. A host cell comprising the vector of claim 17.
20. The host cell of claim 19, wherein said cell is a CHO cell, an *E. coli* or a yeast cell.

* * * * *