




(86) Date de dépôt PCT/PCT Filing Date: 2008/03/12
(87) Date publication PCT/PCT Publication Date: 2008/09/25
(85) Entrée phase nationale/National Entry: 2009/09/03
(86) N° demande PCT/PCT Application No.: JP 2008/055028
(87) N° publication PCT/PCT Publication No.: 2008/114807
(30) Priorité/Priority: 2007/03/13 (US60/894,624)

(51) Cl.Int./Int.Cl. *A61K 31/513* (2006.01),
A61P 3/10 (2006.01), *A61P 31/18* (2006.01),
A61P 35/00 (2006.01)
(71) Demandeur/Applicant:
TAKEDA PHARMACEUTICAL COMPANY LIMITED, JP
(72) Inventeur/Inventor:
OGAWA, ATSUSHI, JP
(74) Agent: SMART & BIGGAR

(54) Titre : ADMINISTRATION HEBDOMADAIRE D'INHIBITEURS DE LA DIPEPTIDYLE PEPTIDASE
(54) Title: WEEKLY ADMINISTRATION OF DIPEPTIDYL PEPTIDASE INHIBITORS

FIG. 2



(57) **Abrégé/Abstract:**

Pharmaceutical compositions comprising 2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile and pharmaceutically acceptable salts thereof are provided as well as kits and articles of manufacture comprising the pharmaceutical compositions as well as methods of using the pharmaceutical compositions.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 September 2008 (25.09.2008)

PCT

(10) International Publication Number
WO 2008/114807 A1

(51) International Patent Classification:

A61K 31/513 (2006.01) A61P 35/00 (2006.01)
A61P 3/10 (2006.01) A61P 31/18 (2006.01)

(74) Agent: **TAKASHIMA, Hajime**; Meiji Yasuda Seimei Osaka Midosuji Bldg., 1-1, Fushimimachi 4-chome, Chuo-ku, Osaka-shi, Osaka 5410044 (JP).

(21) International Application Number:

PCT/JP2008/055028

(22) International Filing Date: 12 March 2008 (12.03.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/894,624 13 March 2007 (13.03.2007) US

(71) Applicant (for all designated States except US): **TAKEDA PHARMACEUTICAL COMPANY LIMITED** [JP/JP]; 1-1, Doshomachi 4-chome, Chuo-ku, Osaka-shi, Osaka, 5410045 (JP).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **OGAWA, Atsushi** [JP/JP]; c/o TAKEDA PHARMACEUTICAL COMPANY LIMITED, 17-85Jusohonmachi 2-chome, Yodogawa-ku, Osaka-shi, Osaka, 5328686 (JP).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(54) Title: WEEKLY ADMINISTRATION OF DIPEPTIDYL PEPTIDASE INHIBITORS

FIG. 2



(57) Abstract: Pharmaceutical compositions comprising 2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile and pharmaceutically acceptable salts thereof are provided as well as kits and articles of manufacture comprising the pharmaceutical compositions as well as methods of using the pharmaceutical compositions.

WO 2008/114807 A1

DESCRIPTION**WEEKLY ADMINISTRATION OF DIPEPTIDYL PEPTIDASE INHIBITORS****TECHNICAL FIELD OF THE INVENTION**

[0001] The invention relates to the method of administering
5 compounds and pharmaceutical compositions used to inhibit
dipeptidyl peptidase IV as well as treatment methods based
on such administration.

BACKGROUND OF THE INVENTION**(DESCRIPTION OF RELATED ART)**

10 [0002] Dipeptidyl Peptidase IV (IUBMB Enzyme Nomenclature
EC.3.4.14.5) is a type II membrane protein that has been
referred to in the literature by a wide a variety of names
including DPP4, DP4, DAP-IV, FAP β , adenosine deaminase
complexing protein 2, adenosine deaminase binding protein
15 (ADAbp), dipeptidyl aminopeptidase IV; Xaa-Pro-dipeptidyl-
aminopeptidase; Gly-Pro naphthylamidase; postproline
dipeptidyl aminopeptidase IV; lymphocyte antigen CD26;
glycoprotein GP110; dipeptidyl peptidase IV; glycylproline
aminopeptidase; glycylproline aminopeptidase; X-prolyl
20 dipeptidyl aminopeptidase; pep X; leukocyte antigen CD26;
glycylprolyl dipeptidylaminopeptidase; dipeptidyl-peptide
hydrolase; glycylprolyl aminopeptidase; dipeptidyl-
aminopeptidase IV; DPP IV/CD26; amino acyl-prolyl dipeptidyl
aminopeptidase; T cell triggering molecule Tp103; X-PDAP.
25 Dipeptidyl Peptidase IV is referred to herein as "DPP-IV."
[0003] DPP-IV is a non-classical serine aminodipeptidase
that removes Xaa-Pro dipeptides from the amino terminus (N-
terminus) of polypeptides and proteins. DPP-IV dependent
slow release of dipeptides of the type X-Gly or X-Ser has
30 also been reported for some naturally occurring peptides.
[0004] DPP-IV is constitutively expressed on epithelial and
endothelial cells of a variety of different tissues
(intestine, liver, lung, kidney and placenta), and is also

found in body fluids. DPP-IV is also expressed on circulating T-lymphocytes and has been shown to be synonymous with the cell-surface antigen, CD-26.

5 [0005] DPP-IV is responsible for the metabolic cleavage of certain endogenous peptides (GLP-1 (7-36), glucagon) *in vivo* and has demonstrated proteolytic activity against a variety of other peptides (GHRH, NPY, GLP-2, VIP) *in vitro*.

10 [0006] GLP-1 (7-36) is a 29 amino-acid peptide derived by post-translational processing of proglucagon in the small intestine. GLP-1 (7-36) has multiple actions *in vivo* including the stimulation of insulin secretion, inhibition of glucagon secretion, the promotion of satiety, and the slowing of gastric emptying. Based on its physiological profile, the actions of GLP-1 (7-36) are believed to be
15 beneficial in the prevention and treatment of type II diabetes and potentially obesity. For example, exogenous administration of GLP-1 (7-36) (continuous infusion) in diabetic patients has been found to be efficacious in this patient population. Unfortunately, GLP-1 (7-36) is degraded
20 rapidly *in vivo* and has been shown to have a short half-life *in vivo* ($t_{1/2}$ =1.5 minutes).

[0007] Based on a study of genetically bred DPP-IV knock out mice and on *in vivo* / *in vitro* studies with selective DPP-IV inhibitors, DPP-IV has been shown to be the primary
25 degrading enzyme of GLP-1 (7-36) *in vivo*. GLP-1 (7-36) is degraded by DPP-IV efficiently to GLP-1 (9-36), which has been speculated to act as a physiological antagonist to GLP-1 (7-36). Inhibiting DPP-IV *in vivo* is therefore believed to be useful for potentiating endogenous levels of GLP-1 (7-
30 36) and attenuating the formation of its antagonist GLP-1 (9-36). Thus, DPP-IV inhibitors are believed to be useful agents for the prevention, delay of progression, and/or treatment of conditions mediated by DPP-IV, in particular

diabetes and more particularly, type 2 diabetes mellitus, diabetic dislipidemia, conditions of impaired glucose tolerance (IGT), conditions of impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation and obesity.

[0008] Several compounds have been shown to inhibit DPP-IV.

Nonetheless, a need still exists for new DPP-IV inhibitors and methods of administering such inhibitors for the treatment of disease.

DISCLOSURE OF THE INVENTION

(SUMMARY OF THE INVENTION)

[0009] A method is provided comprising: administering a weekly dose of between 1 mg/week and 500 mg/week of Compound I to a patient, optionally between 12.5 mg/week and 400

mg/week of Compound I, optionally between 20 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and optionally between 100 mg/week and 400 mg/week of Compound I. In one variation, a weekly dose of 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg, 400 mg or 500mg of Compound I is administered, optionally a weekly dose of 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg or 400 mg of Compound I is administered.

[0010] A method is also provided comprising administering a weekly dose of more than 250 mg of Compound I, optionally at least 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at least 350 mg of Compound I, and optionally at least 400 mg of Compound I to a patient. In one variation, the weekly dose of Compound I administered to the patient is not more than 500 mg. In another variation, the weekly dose of Compound I administered to the patient is not more than 400 mg. In still another variation, the

weekly dose of Compound I administered to the patient is not more than 350 mg. In yet another variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 500 mg. In a further variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 400 mg. In other variations, a weekly dose of 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I is administered to a patient.

[0011] In still a further variation, administering is performed 1 time per week and may optionally be performed 1 time per week as a single dosage. Optionally, administering is performed 1 time per week for a period of at least 1 month, optionally for a period of at least 2 months and optionally for a period of at least 3 months. In one variation, administering is performed in the morning and optionally is performed in the morning prior to a first meal of the day for the patient.

[0012] Administering may be performed by a wide range of routes of administration including, but not limited to a route selected from the group consisting of orally, parenterally, intraperitoneally, intravenously, intraarterially, transdermally, sublingually, intramuscularly, rectally, transbuccally, intranasally, liposomally, via inhalation, vaginally, intraocularly, via local delivery, subcutaneously, intraadiposally, intraarticularly, intraperitoneally and intrathecally. In one particular variation, administering is performed orally.

[0013] A method is also provided for administering Compound I in combination with one or more antidiabetic or incretin compounds other than Compound I. In one variation, such combination therapy method is performed where a weekly dose of between 1 mg/week and 500 mg/week of Compound I, optionally between 12.5 mg/week and 400 mg/week of Compound

I, optionally between 20 mg/week and 400 mg/ of Compound I, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and optionally between 100 mg/week and 400 mg/week of

5 Compound I. In one variation, a weekly dose of 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg, 400mg or 500 mg of Compound I is administered to a patient in combination with one or more antidiabetic compounds other than Compound I.

10 [0014] In a further variation, such combination therapy method is performed where a weekly dose of more than 250 mg of Compound I, optionally at least 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at least 350 mg of Compound I, and optionally at least 400 mg

15 of Compound I is administered to a patient. In still a further variation, the weekly dose of Compound I administered to the patient is not more than 500 mg. In another variation, the weekly dose of Compound I administered to the patient is not more than 400 mg. In

20 still another variation, the weekly dose of Compound I administered to the patient is not more than 350 mg. In yet another variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 500 mg. In a further variation, the weekly dose of

25 Compound I administered to the patient is more than 250 mg and not more than 400 mg. In other variations, a weekly dose of 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I is administered to a patient in combination with one or more antidiabetic compounds other than Compound I.

30 [0015] In regard to each of the above embodiments and variations thereof, Compound I may be administered as a free base or as a pharmaceutically acceptable salt thereof. In particular variations, Compound I is administered as a HCl,

methanesulfonate, succinate, benzoate, toluenesulfonate, R-
(-)-mandelate or benzenesulfonate salt of Compound I.

[0016] Pharmaceutical compositions are also provided. In
one embodiment, a pharmaceutical composition is provided
5 that is formulated in a single dose form wherein such single
dose form comprises between 1 mg/week and 500 mg/week of
Compound I, optionally between 12.5 mg/week and 400 mg/week
of Compound I, optionally between 20 mg/week and 400 mg/ of
Compound I, optionally between 20 mg/week and 200 mg/week of
10 Compound I, optionally between 50 mg/week and 400 mg/week of
Compound I, and optionally between 100 mg/week and 400
mg/week of Compound I. In particular variations, the
pharmaceutical composition comprises 3.125 mg, 12.5 mg, 20
mg, 25mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200
15 mg, 250 mg, 400 mg or 500mg of Compound I.

[0017] In another embodiment, a pharmaceutical composition
is provided that is formulated in a single dose form wherein
such single dose form comprises more than 250 mg of Compound
I, optionally at least 275 mg of Compound I, optionally at
20 least 300 mg of Compound I, optionally at least 350 mg of
Compound I, and optionally at least 400 mg of Compound I.
In one variation, the pharmaceutical composition comprises
not more than 500 mg of Compound I. In another variation,
the pharmaceutical composition comprises not more than 400
25 mg of Compound I. In still another variation, the
pharmaceutical composition comprises not more than 350 mg of
Compound I. In yet another variation, the pharmaceutical
composition comprises more than 250 mg and not more than 500
mg of Compound I. In a further variation, the
30 pharmaceutical composition comprises more than 250 mg and
not more than 400 mg of Compound I. In other variations,
the pharmaceutical composition comprises 275 mg, 300 mg, 350
mg, 400 mg or 500 mg of Compound I.

[0018] In another embodiment, a pharmaceutical composition is provided that comprises Compound I and one or more antidiabetic or incretin compounds other than Compound I in a single dose form. In one variation, Compound I is present in the single dose form in a dosage amount between 1 mg/week and 500 mg/week of Compound I, optionally between 12.5 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 400 mg/ of Compound I, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and optionally between 100 mg/week and 400 mg/week of Compound I. In particular variations, the pharmaceutical composition comprises 3.125 mg, 12.5 mg, 20 mg, 25mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg, 400mg or 500mg of Compound I.

[0019] In another variation, Compound I is present in the single dose form in a dosage amount of more than 250 mg of Compound I, optionally at least 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at least 350 mg of Compound I, and optionally at least 400 mg of Compound I. In a further variation, the pharmaceutical composition comprises not more than 500 mg of Compound I. In another variation, the pharmaceutical composition comprises not more than 400 mg of Compound I. In still another variation, the pharmaceutical composition comprises not more than 350 mg of Compound I. In yet another variation, the pharmaceutical composition comprises more than 250 mg and not more than 500 mg of Compound I. In a further variation, the pharmaceutical composition comprises more than 250 mg and not more than 400 mg of Compound I. In other variations, the pharmaceutical composition comprises 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I.

[0020] In regard to each of the above embodiments and variations thereof regarding pharmaceutical compositions, Compound I may be present in the pharmaceutical composition as a free base or as a pharmaceutically acceptable salt thereof. In particular variations, Compound I is present as a HCl, methanesulfonate, succinate, benzoate, toluenesulfonate, R-(-)mandelate or benzenesulfonate salt of Compound I.

[0021] Also in regard to each of the above embodiments and variations thereof regarding pharmaceutical compositions, the pharmaceutical composition may optionally be a single dose form adapted for oral administration, optionally a solid formulation adapted for oral administration, and optionally a tablet or capsule adapted for oral administration. Further, in regard to each of the above embodiments and variations thereof regarding pharmaceutical compositions, the pharmaceutical composition may optionally be a single dose form adapted for parenteral (subcutaneous, intravenous, subdermal or intramuscular) administration, optionally a solution formulation adapted for parenteral administration, and optionally a suspension formulation adapted for parenteral administration. The pharmaceutical formulation may also be an extended release formulation adapted for oral administration.

[0022] Also in regard to each of the above embodiments and variations thereof regarding pharmaceutical compositions, the pharmaceutical composition may be employed to prevent or treat conditions mediated by DPP-IV such as diabetes and more particularly, type 2 diabetes mellitus; diabetic dislipidemia; impaired glucose tolerance (IGT); impaired fasting plasma glucose (IFG); metabolic acidosis; ketosis; appetite regulation; obesity; complications associated with diabetes including diabetic neuropathy, diabetic retinopathy

and kidney disease; hyperlipidemia including
hypertriglyceridemia, hypercholesteremia, hypoHDLemia and
postprandial hyperlipidemia; arteriosclerosis; hypertension;
myocardial infarction, angina pectoris, cerebral infarction,
5 cerebral apoplexy and metabolic syndrome.

[0023] Combinations of Compound I with one or more
antidiabetic or incretin compounds other than Compound I
provide excellent effects such as 1) enhancement in
therapeutic effects of Compound I and/or the antidiabetic or
10 incretin compounds; 2) reduction in side effects of Compound
I and/or the antidiabetic or incretin compounds; and 3)
reduction in a dose of Compound I and/or the antidiabetic or
incretin compounds. Accordingly, the present invention
comprises methods of administering Compound I in combination
15 with one or more other antidiabetic or incretin compounds
and pharmaceutical compositions comprising Compound I
together with one or more other antidiabetic or incretin
compounds. It is noted that several different dosage ranges
for particular antidiabetic and incretin compounds are
20 provided herein. It is intended for the scope of the
present invention to include drug combinations covering any
of the disclosed ranges for Compound I in combination with
any of the dosage ranges described herein for other
antidiabetic or incretin compounds.

25 [0024] With respect to each of the above embodiments and
variations thereof regarding methods and pharmaceutical
compositions comprising one or more antidiabetic or incretin
compounds other than Compound I, the one or more
antidiabetic or incretin compounds may be selected from any
30 of a variety of known antidiabetic and incretin compounds.
In one variation, the one or more antidiabetic or incretin
compounds used in combination with Compound I may optionally
be selected from the group consisting of insulin signaling

pathway modulators, compounds influencing a dysregulated hepatic glucose production, insulin sensitivity enhancers, and insulin secretion enhancers.

[0025] The one or more antidiabetic or incretin compounds
5 used in combination with Compound I may also optionally be selected from the group consisting of protein tyrosine phosphatase inhibitors, glutamine-fructose-6-phosphate amidotransferase inhibitors, glucose-6-phosphatase inhibitors, fructose-1,6-bisphosphatase inhibitors, glycogen
10 phosphorylase inhibitors, glucagon receptor antagonists, phosphoenolpyruvate carboxykinase inhibitors, pyruvate dehydrogenase kinase inhibitors, alpha-glucosidase inhibitors, inhibitors of gastric emptying, glucokinase activators, GLP-1 receptor agonists, GLP-2 receptor agonists,
15 UCP modulators, RXR modulators, GSK-3 inhibitors, PPAR modulators, metformin, insulin, α_2 -adrenergic antagonists, deacetylases (e.g., resveratrol, sirtuin agonist, polyphenols), and Sodium dependent glucose transport (SGLT2) inhibitors.

20 [0026] The one or more antidiabetic or incretin compounds used in combination with Compound I may also optionally be selected from the group consisting of GSK-3 inhibitors, retinoid X receptor agonists, Beta-3 AR agonists, UCP
25 modulators, antidiabetic thiazolidinediones, non-glitazone type PPAR gamma agonists, dual PPAR gamma/PPAR alpha agonists, antidiabetic vanadium containing compounds and biguanides.

[0027] The one or more antidiabetic or incretin compounds
used in combination with Compound I may also optionally be
30 thiazolidinediones selected from the group consisting of (S)-((3,4-dihydro-2-(phenyl-methyl)-2H-1-benzopyran-6-yl)methyl-thiazolidine-2,4-dione, 5-{[4-(3-(5-methyl-2-phenyl-4-oxazolyl)-1-oxo-propyl)-phenyl]-methyl}-

thiazolidine-2,4-dione, 5-([4-(1-methyl-cyclohexyl)methoxy)-phenyl)methyl]-thiazolidine-2,4-dione, 5-([4-(2-(1-indolyl)ethoxy)phenyl)methyl]-thiazolidine-2,4-dione, 5-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)-ethoxy]benzyl]-thiazolidine-2,4-dione, 5-(2-naphthylsulfonyl)-thiazolidine-2,4-dione, bis[4-[(2,4-dioxo-5-thiazolidinyl)-methyl]phenyl]methane, 5-[4-[2-(5-methyl-2-phenyl-4-oxazolyl)-2-hydroxyethoxy]-benzyl]-thiazolidine-2,4-dione, 5-[4-(1-phenyl-1-cyclopropanecarbonylamino)-benzyl]-thiazolidine-2,4-dione, 5-([4-(2-(2,3-dihydroindol-1-yl)ethoxy)phenyl)methyl]-thiazolidine-2,4-dione, 5-[3-(4-chloro-phenyl)]-2-propynyl]-5-phenylsulfonyl)thiazolidine-2,4-dione, 5-[3-(4-chlorophenyl)]-2-propynyl]-5-(4-fluorophenyl-sulfonyl)thiazolidine-2,4-dione, 5-([4-(2-(methyl-2-pyridinyl-amino)-ethoxy)phenyl)methyl]-thiazolidine-2,4-dione, 5-([4-(2-(5-ethyl-2-pyridyl)ethoxy)phenyl]-methyl)-thiazolidine-2,4-dione, 5-([4-((3,4-dihydro-6-hydroxy-2,5,7,8-tetramethyl-2H-1-benzopyran-2-yl)methoxy)-phenyl]-methyl)-thiazolidine-2,4-dione, 5-[6-(2-fluoro-benzyloxy)-naphthalen-2-ylmethyl]-thiazolidine-2,4-dione, 5-([2-(2-naphthyl)-benzoxazol-5-yl]-methyl)thiazolidine-2,4-dione and 5-(2,4-dioxothiazolidin-5-ylmethyl)-2-methoxy-N-(4-trifluoromethyl-benzyl)benzamide, including any pharmaceutically acceptable salts thereof.

[0028] In one variation, the one or more antidiabetic compounds used in combination with Compound I includes metformin. In one particular variation, the metformin in this combination comprises one or more pharmaceutically acceptable salts thereof. In another particular variation, the metformin in this combination comprises a metformin HCl salt. In still another particular variation, the metformin in this combination is administered in a daily dose of between 125 and 2550 mg. In yet another variation, the

metformin in this combination is administered in a daily dose of between 250 and 2550 mg. In other variations, the metformin in this combination is administered in an immediate release or an extended release formulation.

5 [0029] In another variation, the one or more antidiabetic compounds used in combination with Compound I includes one or more sulphonyl urea derivatives.

[0030] The one or more antidiabetic compounds used in combination with Compound I may also optionally be selected
10 from the group consisting of glisoxepid, glyburide, glibenclamide, acetohexamide, chloropropamide, glibornuride, tolbutamide, tolazamide, glipizide, carbutamide, gliquidone, glyhexamide, phenbutamide, tolcyclamide, glimepiride and gliclazide, including any pharmaceutically acceptable salts
15 thereof. In one variation, the one or more antidiabetic compounds administered in combination with Compound I includes glimepiride.

[0031] The one or more antidiabetic compounds used in combination with Compound I may also optionally be selected
20 from the group consisting of incretin hormones or mimics thereof, beta-cell imidazoline receptor antagonists, and short-acting insulin secretagogues.

[0032] In another variation, the one or more antidiabetic compounds used in combination with Compound I includes
25 insulin.

[0033] The one or more antidiabetic compounds used in combination with Compound I may also optionally be one or more GLP-1 agonists including, for example, extendatide.

[0034] The one or more antidiabetic compounds used in
30 combination with Compound I may also optionally be one or more GLP-2 agonists including, for example, human recombinant GLP-2.

[0035] The one or more antidiabetic compounds used in combination with Compound I may also optionally be one or more antidiabetic D-phenylalanine derivatives.

5 [0036] The one or more antidiabetic compounds used in combination with Compound I may also optionally be selected from the group consisting of repaglinide, mitiglinide and nateglinide, including any pharmaceutically acceptable salts thereof. In one variation, the one or more antidiabetic compounds administered in combination with Compound I
10 includes mitiglinide calcium salt hydrate.

[0037] The one or more antidiabetic compounds used in combination with Compound I may also optionally be one or more alpha-Glucosidase inhibitors.

15 [0038] The one or more antidiabetic compounds used in combination with Compound I may also optionally be selected from the group consisting of acarbose, voglibose and miglitol, including any pharmaceutically acceptable salts thereof. In one variation, the one or more antidiabetic compounds administered in combination with Compound I
20 includes voglibose. In another variation, the voglibose in this combination is administered in a daily dose of between 0.1 and 1 mg.

[0039] The one or more antidiabetic compounds used in combination with Compound I may also optionally be
25 rosiglitazone, including any pharmaceutically acceptable salts thereof. In one variation, the rosiglitazone in this combination comprises a rosiglitazone maleate salt

[0040] The one or more antidiabetic compounds used in combination with Compound I may also optionally be
30 tesaglitazar, muraglitazar or naveglitazar, including any pharmaceutically acceptable salts thereof.

[0041] The one or more antidiabetic compounds used in combination with Compound I may also optionally be

pioglitazone, including any pharmaceutically acceptable salts thereof. In one variation, the pioglitazone in this combination comprises a pioglitazone HCl salt. In another variation, the pioglitazone in this combination is
5 administered in a daily dose of between 7.5 and 60 mg. In still another variation, the pioglitazone in this combination is administered in a daily dose of between 15 and 45 mg.

[0042] The one or more antidiabetic compounds used in
10 combination with Compound I may also optionally comprise metformin and pioglitazone. In one variation, the pioglitazone in this combination comprises one or more pharmaceutically acceptable salts thereof. In another variation, the pioglitazone in this combination comprises a
15 pioglitazone HCl salt. In still another variation, the pioglitazone in this combination is administered in a daily dose of between 7.5 and 60 mg. In yet another variation, the pioglitazone in this combination is administered in a daily dose of between 15 and 45 mg. In another variation of
20 each of the above variations, the metformin in this combination comprises one or more pharmaceutically acceptable salts thereof. In one particular variation, the metformin in this combination comprises a metformin HCl salt. In another particular variation, the metformin in this
25 combination is administered in a daily dose of between 125 and 2550 mg. In still another variation, the metformin in this combination is administered in a daily dose of between 250 and 2550 mg.

[0043] Compound I and pharmaceutical compositions
30 comprising Compound I may be used to treat a range of diseases. In one variation, administering Compound I or a pharmaceutical composition comprising Compound I is performed to treat type I or type II diabetes disease state

of the patient. In another variation, administering Compound I or a pharmaceutical composition comprising Compound I is performed to treat a pre-diabetic patient. In still another variation, administering Compound I or a pharmaceutical composition comprising Compound I is performed to treat an inflammatory bowel disease, Crohn's disease, chemotherapy-induced enteritis, oral mucositis or Shortened Bowel syndrome. In yet another variation, administering Compound I or a pharmaceutical composition comprising Compound I is performed to increase engraftment efficiency after bone marrow transplantation. In another variation, administering Compound I or a pharmaceutical composition comprising Compound I is performed to treat a patient suffering from conditions mediated by DPP-IV such as diabetes and more particularly, type 2 diabetes mellitus; diabetic dyslipidemia; impaired glucose tolerance (IGT); impaired fasting plasma glucose (IFG); metabolic acidosis; ketosis; appetite regulation; obesity; complications associated with diabetes including diabetic neuropathy, diabetic retinopathy and kidney disease; hyperlipidemia including hypertriglyceridemia, hypercholesteremia, hypoHDLemia and postprandial hyperlipidemia; arteriosclerosis; hypertension; myocardial infarction, angina pectoris, cerebral infarction, cerebral apoplexy and metabolic syndrome.

[0044] In addition, Compound I and pharmaceutical compositions comprising Compound I may be employed to prevent or treat conditions mediated by DPP-IV such as diabetes and more particularly, type 2 diabetes mellitus; diabetic dyslipidemia; impaired glucose tolerance (IGT); impaired fasting plasma glucose (IFG); metabolic acidosis; ketosis; appetite regulation; obesity; complications associated with diabetes including diabetic neuropathy,

diabetic retinopathy and kidney disease; hyperlipidemia including hypertriglyceridemia, hypercholesteremia, hypoHDLemia and postprandial hyperlipidemia; arteriosclerosis; hypertension; myocardial infarction, angina pectoris, cerebral infarction, cerebral apoplexy and metabolic syndrome.

[0045] Also provided are kits comprising multiple doses of pharmaceutical compositions according to the present invention. In one variation, the kits further comprise instructions which comprise one or more forms of information selected from the group consisting of indicating a disease state for which the pharmaceutical composition is to be administered, storage information for the pharmaceutical composition, dosing information and instructions regarding how to administer the pharmaceutical composition.

[0046] Also provided are articles of manufacture comprising multiple doses of pharmaceutical composition according to the present invention. In one variation, the articles of manufacture further comprise packaging materials such as a container for housing the multiple doses of the pharmaceutical composition and or a label indicating one or more members of the group consisting of a disease state for which the compound is to be administered, storage information, dosing information and/or instructions regarding how to administer the composition.

[0047] Accordingly, the present invention relates to

[1] a method comprising: administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week,

[2] the method of the above-mentioned [1], wherein the weekly dose of Compound I administered to the patient is at least 275 mg,

[3] the method of the above-mentioned [1], wherein the weekly dose of Compound I administered to the patient is at least 300 mg,

5 [4] the method of the above-mentioned [1], wherein the weekly dose of Compound I administered to the patient is at least 350 mg,

[5] the method of the above-mentioned [1], wherein the weekly dose of Compound I administered to the patient is at least 400 mg,

10 [6] the method of any one of the above-mentioned [1]-[5], wherein the weekly dose of Compound I administered to the patient is not more than 500 mg,

[7] the method of any one of the above-mentioned [1]-[4], wherein the weekly dose of Compound I administered to the
15 patient is not more than 400 mg,

[8] the method of any one of the above-mentioned [1]-[3], wherein the weekly dose of Compound I administered to the patient is not more than 350 mg,

20 [9] the method of the above-mentioned [1], wherein the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 500 mg,

[10] the method of the above-mentioned [1], wherein the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 400 mg,

25 [11] the method of any one of the above-mentioned [1]-[10], wherein Compound I is administered as a free base,

[12] the method of any one of the above-mentioned [1]-[10],
wherein Compound I is administered as a pharmaceutically
acceptable salt,

[13] the method of any one of the above-mentioned [1]-[10],
5 wherein Compound I is administered as a succinate salt,

[14] a pharmaceutical composition formulated in a single
dose form comprising more than 250 mg of Compound I,

[15] the pharmaceutical composition of the above-mentioned
[14], wherein the single dose form comprises at least 275 mg
10 of Compound I,

[16] the pharmaceutical composition of the above-mentioned
[14], wherein the single dose form comprises at least 300 mg
of Compound I,

[17] the pharmaceutical composition of the above-mentioned
15 [14], wherein the single dose form comprises at least 350 mg
of Compound I,

[18] the pharmaceutical composition of the above-mentioned
[14], wherein the single dose form comprises not more than
500 mg of Compound I,

20 [19] the pharmaceutical composition of the above-mentioned
[14], wherein the single dose form comprises not more than
400 mg of Compound I,

[20] the pharmaceutical composition of the above-mentioned
[14], wherein the single dose form comprises not more than
25 350 mg of Compound I,

[21] the pharmaceutical composition of the above-mentioned [14], wherein the single dose form comprises more than 250 mg and not more than 500 mg of Compound I,

5 [22] the pharmaceutical composition of the above-mentioned [14], wherein the single dose form comprises more than 250 mg and not more than 400 mg of Compound I,

[23] the pharmaceutical composition of any one of the above-mentioned [14]-[22], wherein Compound I is present in the pharmaceutical composition as a free base,

10 [24] the pharmaceutical composition of any one of the above-mentioned [14]-[22], wherein Compound I is present in the pharmaceutical composition in a pharmaceutically acceptable salt,

15 [25] the pharmaceutical composition of any one of the above-mentioned [14]-[22], wherein Compound I is present in the pharmaceutical composition in a succinate salt,

[26] use of Compound I for the manufacture of a pharmaceutical composition of any one of the above-mentioned [14]-[25],

20 [27] a method of treating diabetes comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week,

25 [28] a method of treating cancer comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week,

[29] a method of treating autoimmune disorders comprising administering a weekly dose of more than 250 mg of Compound

I to a patient wherein the weekly dose is administered once per week,

[30] a method of treating HIV infection comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week;

and the like.

[0048] It is noted in regard to all of the above embodiments that the embodiments should be interpreted as being open ended in the sense that the methods may comprise further actions beyond those specified including the administration of other pharmaceutically active materials to a patient. Similarly, unless otherwise specified, the pharmaceutical compositions, kits and articles of manufacture may further include other materials including other pharmaceutically active materials.

BRIEF DESCRIPTION OF THE DRAWINGS

[0049] Figure 1 illustrates DPP IV inhibition in plasma after a single oral administration of Compound I in human.

[0050] Figure 2 illustrates DPP IV inhibition in plasma after a single oral administration of Compound I in human.

DETAILED DESCRIPTION OF THE INVENTION

(DEFINITIONS)

[0051] Unless otherwise stated, the following terms used in the specification and claims shall have the following meanings for the purposes of this Application.

[0052] "Disease" specifically includes any unhealthy condition of an animal or part thereof and includes an unhealthy condition that may be caused by, or incident to,

medical or veterinary therapy applied to that animal, i.e., the "side effects" of such therapy.

[0053] "Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary use as well as human pharmaceutical use.

[0054] "Pharmaceutically acceptable salts" means salts which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity. Such salts include, but are not limited to, acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or with organic acids such as acetic acid, trifluoroacetic acid, propionic acid, hexanoic acid, heptanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, o-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, p-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, p-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid and the like.

[0055] Pharmaceutically acceptable salts also include, but are not limited to, base addition salts which may be formed when acidic protons present are capable of reacting with

inorganic or organic bases. Acceptable inorganic bases include, but are not limited to, sodium hydroxide, sodium carbonate, potassium hydroxide, aluminum hydroxide and calcium hydroxide. Acceptable organic bases include, but
5 are not limited to, ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine and the like.

[0056] "Subject" and "patient" includes humans, non-human mammals (e.g., dogs, cats, rabbits, cattle, horses, sheep,
10 goats, swine, deer, and the like) and non-mammals (e.g., birds, and the like).

[0057] "Therapeutically effective amount" means that amount of a compound which, when administered to an animal for treating a disease, is sufficient to effect such treatment
15 for the disease.

[0058] "Treatment" or "treating" means any administration of a therapeutically effective amount of a compound and includes:

(1) preventing the disease from occurring in an animal
20 which may be predisposed to the disease but does not yet experience or display the pathology or symptomatology of the disease,


(2) inhibiting the disease in an animal that is experiencing or displaying the pathology or symptomatology
25 of the disease (i.e., arresting further development of the pathology and/or symptomatology), or

(3) ameliorating the disease in an animal that is experiencing or displaying the pathology or symptomatology
30 of the disease (i.e., reversing the pathology and/or symptomatology).

(DETAILED DESCRIPTION OF THE INVENTION)

1. 2-[6-(3-AMINO-PIPERIDIN-1-YL)-3-METHYL-2,4-DIOXO-3,4-DIHYDRO-2H-PYRIMIDIN-1-YLMETHYL]-4-FLUORO-BENZONITRILE AND COMPOSITIONS THEREOF

[0059] The present invention relates generally to the administration of 2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluorobenzonitrile (referred to herein as "Compound I") whose structure is provided below.



[0060] Example 1 describes one method for synthesizing Compound I. It is noted that other methods for synthesizing Compound I may be used as would be appreciated to one of ordinary skill in the art. As described in detail below, Compound I has long acting DPP-IV inhibitory affects.

[0061] Compound I may be administered in its free base form and may also be administered in the form of salts, hydrates and prodrugs that are converted *in vivo* into the free base form of Compound I. For example, it is within the scope of the present invention to administer Compound I as a pharmaceutically acceptable salt derived from various organic and inorganic acids and bases in accordance with procedures well known in the art. As used herein, Compound I is intended to encompass salts, hydrates and prodrugs of Compound I unless otherwise specified.

[0062] A pharmaceutically acceptable salt of Compound I preferably confers improved pharmacokinetic properties as

compared to the free base form of Compound I.

Pharmaceutically acceptable salts may also confer desirable pharmacokinetic properties on Compound I that it did not previously possess, and may even positively affect the pharmacodynamics of the compound with respect to its therapeutic activity in the body.

[0063] Particular examples of salts, hydrates and prodrugs of Compound I include, but are not limited to salt forms formed by inorganic or organic acids, e.g., hydrohalides such as hydrochloride, hydrobromide and hydroiodide; other mineral acids and their corresponding salts such as sulfate, nitrate, phosphate, etc.; alkyl and monoarylsulfonates such as ethanesulfonate, toluenesulfonate and benzenesulfonate; and other organic acids and their corresponding salts such as acetate, trifluoroacetate, tartrate, maleate, succinate, citrate, benzoate, salicylate and ascorbate. Further acid addition salts include, but are not limited to: adipate, alginate, arginate, aspartate, bisulfate, bisulfite, bromide, butyrate, camphorate, camphorsulfonate, caprylate, chloride, chlorobenzoate, cyclopentanepropionate, digluconate, dihydrogenphosphate, dinitrobenzoate, dodecylsulfate, fumarate, galacterate (from mucic acid), galacturonate, glucoheptaate, gluconate, glutamate, glycerophosphate, hemisuccinate, hemisulfate, heptanoate, hexanoate, hippurate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, iodide, isethionate, iso-butyrate, lactate, lactobionate, malate, malonate, mandelate, metaphosphate, methanesulfonate, methylbenzoate, monohydrogenphosphate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, oleate, pamoate, pectinate, persulfate, phenylacetate, 3-phenylpropionate, phosphate, phosphonate and phthalate.

[0064] In particular variations, Compound I is administered as a HCl, methanesulfonate, succinate, benzoate, toluenesulfonate, R-(-)mandelate or benzenesulfonate salt of Compound I. Example 1 describes the preparation of various salt forms of Compound I including TFA, HCl, benzoic, p-toluenesulfonic, succinic, R-(-)-mandelic and benzenesulfonic acid salts.

2. ADMINISTRATION AND USE OF COMPOUND I

[0065] The present invention relates generally to a method comprising administering Compound I to a patient at a weekly dose of between 1 mg/week and 500 mg/week of Compound I to a patient, optionally between 12.5 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 400 mg/week of Compound I, optionally between 20mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and optionally between 100 mg/week and 400 mg/week of Compound. Specific dosage amounts that may be used include, but are not limited to 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg, 400 mg and 500 mg of Compound I per week. It is noted that unless otherwise specifically specified, Compound I may be administered in its free base form or as a pharmaceutically acceptable salt. However, the dosage amounts and ranges provided herein are always based on the molecular weight of the free base form of Compound I.

[0066] The present invention also relates generally to a method comprising administering Compound I to a patient at a weekly dose of more than 250 mg of Compound I, optionally at least 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at least 350 mg of Compound I, and optionally at least 400 mg of Compound I. In one variation, the weekly dose of Compound I administered to the patient is not more than 500 mg. In another variation, the weekly dose

of Compound I administered to the patient is not more than 400 mg. In still another variation, the weekly dose of Compound I administered to the patient is not more than 350 mg. In yet another variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 500 mg. In a further variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 400 mg. In other variations, a weekly dose of 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I is administered to the patient. It is noted that unless otherwise specifically specified, Compound I may be administered in its free base form or as a pharmaceutically acceptable salt. However, the dosage amounts and ranges provided herein are based on the molecular weight of the free base form of Compound I.

[0067] Compound I may be administered by any route of administration. In particular embodiments, however, the method of the present invention is practiced by administering Compound I orally. This type of administration is advantageous in that it is easy and may be self-administered by the patient.

[0068] Compound I may be administered one or more times per week. An advantage of the present invention, however, is that Compound I can be effectively administered at the dosage levels specified herein one time per week and may also be administered as a single dosage form one time a week.

By being able to administer Compound I at the dosage levels specified herein only one time per week and orally, it is easier for patients to self-administer Compound I, thus improving the compliance of usage among patients requiring *in vivo* inhibition of DPP-IV activity.

[0069] Advantageously, Compound I is suitable for prolonged continuous use and may be administered to patients for an

extended period of time. Accordingly, the method may be performed where Compound I is administered to a patient each week (optionally 1 time weekly) for a period of at least 1 month, optionally for at least 2 months, optionally for at least 3 months, and, if necessary, optionally for the duration of the patients disease profile.

[0070] Advantageously, Compound I may be administered at any time during the day. Optionally, Compound I is administered one time per week where administration occurs in the morning before meals. Because Compound I can stimulate insulin secretion when blood glucose levels reach 100 mg/dl and above, it may be beneficial to have Compound I in systemic circulation before an elevation in blood glucose levels occurs postprandially.

[0071] Compound I may be administered to any patient who would benefit from a course of treatment leading to the reduction of *in vivo* DPP-IV activity. As described in detail below, Figures 1 and 2 illustrate the observed effect that administering Compound I has on human-plasma DPP-IV activity after a single oral administration. As can be seen from the data shown in Figure 1, by administering Compound I at the dosage levels specified herein, Compound I can be effectively used relative to disease states where it is desired to reduce plasma DPP-IV activity. In view of the data presented, it is believed that when at least 12.5 mg of Compound I is administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 10% relative to baseline for a period of at least 168 hours following administration; when at least 50 mg of Compound I is administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 35% relative to baseline for a period of at least 168 hours following administration; and when 200 mg or 400 mg of Compound I is

administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 70% relative to baseline for a period of at least 168 hours following administration.

5 [0072] Examples of particular applications for administering Compound I include, but are not limited to the prevention, delay of progression, and/or treatment of conditions mediated by DPP-IV, in particular diabetes and more particularly, type 2 diabetes mellitus, diabetic
10 dislipidemia, impaired glucose tolerance (IGT), impaired fasting plasma glucose (IFG), metabolic acidosis, ketosis, appetite regulation, obesity and complications associated with diabetes including diabetic neuropathy, diabetic retinopathy, inflammatory bowel disease, Crohn's disease,
15 chemotherapy-induced enteritis, oral mucositis, Shortened Bowel Syndrome and kidney disease. The conditions mediated by DPP-IV further includes hyperlipidemia such as hypertriglyceridemia, hypercholesteremia, hypoHDLemia and postprandial hyperlipidemia; arteriosclerosis; hypertension;
20 myocardial infarction, angina pectoris, cerebral infarction, cerebral apoplexy and metabolic syndrome.

[0073] It is believed that administration of Compound I to type I or type II diabetic patients following a minimum treatment of at least 30 days will improve one or more
25 cardiovascular measurements. It is also believed that administration of Compound I in combination with one or more antidiabetic or incretin compounds to type I or type II diabetic patients following a minimum treatment of at least 30 days will improve one or more cardiovascular measurements.
30 Examples of cardiac measurements that may be improved include, but are not limited to a decrease in mean systolic blood pressure, an increase in HDL cholesterol, improvement in LDL/HDL ratio and a reduction in triglycerides.

[0074] It is also believed that administration of Compound I in combination with one or more antidiabetic or incretin compounds to patients with gastrointestinal inflammatory disorders (including, but not be limited to inflammatory bowel disease, Crohn's disease, chemotherapy-induced enteritis, oral mucositis and Shortened Bowel Syndrome) following a minimum treatment of at least 30 days will improve the health of the mucosal lining of the gastrointestinal tract. Improvement in the health of the mucosal lining of the gastrointestinal tract may be demonstrated by, but is not limited to, an increase in the intestinal surface area, reduced inflammation, and/or increases in absorption of nutrients.

[0075] Administering Compound I, one time per week at the dosage levels specified herein, to a patient with type 2 diabetes may also be beneficial. Patients receiving Compound I may also have a malfunction in insulin secretion from pancreatic islets rather than patients who have developed insulin resistance in peripheral insulin sensitive tissues/organs.

[0076] Advantageously, administering Compound I one time per week, at the dosage levels specified herein may also be used to treat patients who are prediabetic. It is believed that administering Compound I in a patient who is prediabetic serves to delay development of type II diabetes in that patient. Sustained increase in blood glucose desensitizes pancreatic islet function and impairs insulin secretion. By improving cyclic AMP levels and the calcium dynamics in beta cells, the cells activate genes repairing damaged cell components and are less vulnerable to glucose toxicity.

[0077] Administering Compound I one time per week, at the dosage levels specified herein is expected to have a range

of desirous biological effects *in vivo*. For example, administering Compound I one time per week, at the dosage levels specified herein reduces the patient's blood glucose level when compared with placebo control. Such a decrease
5 in postprandial blood glucose levels helps diabetic patients to maintain lower glucose levels.

[0078] Administering Compound I one time per week, at the dosage levels specified herein is also expected to have the affect of increasing the patient's insulin level or insulin
10 sensitivity. Insulin facilitates entry of glucose into muscle, adipose and several other tissues. The mechanism by which cells can take up glucose is by facilitated diffusion through stimulation of insulin receptor. C-peptide and insulin are protein chains created by the activation and
15 division of proinsulin (an inactive precursor to insulin). C-peptide and insulin are created and stored in the beta cells of the pancreas. When insulin is released into the bloodstream, equal amounts of C-peptide also are released. This makes C-peptide useful as a marker of
20 insulin production. Administering Compound I according to the present invention is expected to increase the patient's C-peptide level.

[0079] Administering Compound I one time per week at the dosage levels specified herein is also expected to have the
25 affect of decreasing the patient's hemoglobin A1c level by greater than 0.5% when compared to placebo control after extended treatment with Compound I. Further, administering Compound I one time per week at the dosage levels specified herein is also expected to have the affect of decreasing the
30 patient's hemoglobin A1c level by greater than 0.2% when compared to placebo control after extended treatment with Compound I. Hb-A1c values are known to be directly proportional to the concentration of glucose in the blood

over the life span of the red blood cells. Hb-A1c thus gives an indication of a patient's blood glucose levels over the previous last 90 days, skewed to the most recent 30 days.

The observed reduction in the patient's hemoglobin A1c level thus verifies the sustained reduction in the patient's blood glucose levels as a result of administering Compound I one time per day at the dosage levels specified herein.

3. COMBINATION THERAPY INCLUDING COMPOUND I

[0080] The present invention also relates to the use of Compound I in combination with one or more other antidiabetic and/or incretin compounds. Examples of such other antidiabetic compounds include, but are not limited to insulin signaling pathway modulators, like protein tyrosine phosphatase (PTPase) inhibitors, and glutamine-fructose-6-phosphate amidotransferase (GFAT) inhibitors; compounds influencing a dysregulated hepatic glucose production, like glucose-6-phosphatase (G6Pase) inhibitors, fructose-1,6-bisphosphatase (F-1,6-BPase) inhibitors, glycogen phosphorylase (GP) inhibitors, glucagon receptor antagonists and phosphoenolpyruvate carboxykinase (PEPCK) inhibitors; pyruvate dehydrogenase kinase (PDHK) inhibitors; insulin sensitivity enhancers (insulin sensitizers); insulin secretion enhancers (insulin secretagogues); alpha-glucosidase inhibitors; inhibitors of gastric emptying; glucokinase activators, GLP-1 receptor agonists, GLP-2 receptor agonists, UCP modulators, RXR modulators, GSK-3 inhibitors, PPAR modulators, metformin, insulin; and α_2 -adrenergic antagonists. Compound I may be administered with such at least one other antidiabetic compound either simultaneously as a single dose, at the same time as separate doses, or sequentially (i.e., where one is administered before or after the other is administered).

[0081] Examples of PTPase inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in U.S. Patent. Nos. 6,057,316, 6,001,867, and PCT Publication Nos. WO 99/58518, WO 99/58522, WO 99/46268, WO 99/46267, WO 99/46244, WO 99/46237, WO 99/46236, and WO 99/15529.

[0082] Examples of GFAT inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in Mol. Cell. Endocrinol. 1997, 135(1), 67-77.

[0083] Examples of G6Pase inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in PCT Publication Nos. WO 00/14090, WO 99/40062 and WO 98/40385, European Patent Publication No. EP682024 and Diabetes 1998, 47, 1630-1636.

[0084] Examples of F-1,6-BPase inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in PCT Publication Nos. WO 00/14095, WO 99/47549, WO 98/39344, WO 98/39343 and WO 98/39342.

[0085] Examples of GP inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in U.S. Patent No. 5,998,463, PCT Publication Nos. WO 99/26659, WO 97/31901, WO 96/39384 and WO9639385 and European Patent Publication Nos. EP 978279 and EP 846464.

[0086] Examples of glucagon receptor antagonists that may be used in combination with Compound I include, but are not limited to those disclosed in U.S. Patent Nos. 5,880,139 and 5,776,954, PCT Publication Nos. WO 99/01423, WO 98/22109, WO 98/22108, WO 98/21957, WO 97/16442 and WO 98/04528 and those described in Bioorg Med. Chem. Lett 1992, 2, 915-918, J. Med. Chem. 1998, 41, 5150-5157, and J. Biol Chem. 1999, 274; 8694-8697.

[0087] Examples of PEPCK inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in U.S. Patent No. 6,030,837 and Mol. Biol. Diabetes 1994,2, 283-99.

5 [0088] Examples of PDHK inhibitors that may be used in combination with Compound I include, but are not limited to those disclosed in J. Med. Chem. 42 (1999) 2741-2746.

[0089] Examples of insulin sensitivity enhancers that may be used in combination with Compound I include, but are not
10 limited to GSK-3 inhibitors, retinoid X receptor (RXR) agonists, Beta-3 AR agonists, UCP modulators, antidiabetic thiazolidinediones (glitazones), non-glitazone type PPAR gamma agonists, dual PPAR gamma/PPAR alpha agonists, antidiabetic vanadium containing compounds and biguanides
15 such as metformin.

[0090] Examples of GSK-3 inhibitors include, but are not limited to those disclosed in PCT Publication Nos. WO 00/21927 and WO 97/41854.

[0091] Examples of RXR modulators include, but are not
20 limited to those disclosed in U.S. Patent Nos. 4,981,784, 5,071,773, 5,298,429 and 5,506,102 and PCT Publication Nos. WO89/05355, WO91/06677, WO92/05447, WO93/11235, WO95/18380, WO94/23068, and WO93/23431.

[0092] Examples of Beta-3 AR agonists include, but are not
25 limited to CL-316,243 (Lederle Laboratories) and those disclosed in U.S. Patent No. 5,705,515 and PCT Publication Nos. WO 99/29672, WO 98/32753, WO 98/20005, WO 98/09625, WO 97/46556, and WO 97/37646.

[0093] Examples of UCP modulators include agonists of UCP-1,
30 UCP-2 and UCP-3. Examples of UCP modulators include, but are not limited to those disclosed in Vidal-Puig et al., Biochem. Biophys. Res. Commun., Vol. 235(1) pp. 79-82 (1997).

[0094] Examples of antidiabetic, PPAR modulating thiazolidinediones (glitazones) include, but are not limited to, (S)-((3,4-dihydro-2-(phenyl-methyl)-2H-1-benzopyran-6-yl)methyl-thiazolidine-2,4-dione (englitazone), 5-{{4-(3-(5-methyl-2-phenyl-4-oxazolyl)-1-oxo-propyl)-phenyl]-methyl}-thiazolidine-2,4-dione (darglitazone), 5-{{4-(1-methyl-cyclohexyl)methoxy)-phenyl]methyl}-thiazolidine-2,4-dione (ciglitazone), 5-{{4-(2-(1-indolyl)ethoxy)phenyl]methyl}-thiazolidine-2,4-dione (DRF2189), 5-{{4-[2-(5-methyl-2-phenyl-4-oxazolyl)-ethoxy]benzyl}-thiazolidine-2,4-dione (BM-13.1246), 5-(2-naphthylsulfonyl)-thiazolidine-2,4-dione (AY-31637), bis{4-[(2,4-dioxo-5-thiazolidinyl)-methyl]phenyl}methane (YM268), 5-{{4-[2-(5-methyl-2-phenyl-4-oxazolyl)-2-hydroxyethoxy]-benzyl}-thiazolidine-2,4-dione (AD-5075), 5-[4-(1-phenyl-1-cyclopropanecarbonylamino)-benzyl]-thiazolidine-2,4-dione (DN-108) 5-{{4-(2-(2,3-dihydroindol-1-yl)ethoxy)phenylmethyl}-thiazolidine-2,4-dione, 5-[3-(4-chloro-phenyl)]-2-propynyl]-5-phenylsulfonyl)thiazolidine-2,4-dione, 5-[3-(4-chlorophenyl)]-2-propynyl]-5-(4-fluorophenyl-sulfonyl)thiazolidine-2,4-dione, 5-{{4-(2-(methyl-2-pyridinyl-amino)-ethoxy)phenyl]methyl}-thiazolidine-2,4-dione (rosiglitazone), 5-{{4-(2-(5-ethyl-2-pyridyl)ethoxy)phenyl]-methyl}-thiazolidine-2,4-dione (pioglitazone; marketed under the trademark ACTOSTM), 5-[6-(2-fluoro-benzyloxy)-naphthalen-2-ylmethyl]-thiazolidine-2,4-dione (MCC555), 5-([2-(2-naphthyl)-benzoxazol-5-yl]-methyl)thiazolidine-2,4-dione (T-174), edaglitazone (BM-13-1258), rivoglitazone (CS-011), and 5-(2,4-dioxothiazolidin-5-ylmethyl)-2-methoxy-N-(4-trifluoromethyl-benzyl)benzamide (KRP297).

[0095] Examples of non-glitazone type PPAR gamma agonists include, but are not limited to N-(2-benzoylphenyl)-L-

tyrosine analogues, such as GI-262570, reglixane (JTT501) and FK-614 and metaglidase (MBX-102).

[0096] Examples of dual PPAR gamma/PPAR alpha agonists include, but are not limited to omega.-

5 [(oxoquinazolinylalkoxy)phenyl]alkanoates and analogs thereof including those described in PCT Publication No. WO 99/08501 and Diabetes 2000, 49(5), 759-767; tesaglitazar, muraglitazar and naveglitazar.

[0097] Examples of antidiabetic vanadium containing
10 compounds include, but are not limited to those disclosed in the U.S. Patent No. 5,866,563.

[0098] Metformin (dimethyldiguanide) and its hydrochloride salt is marketed under the trademark GLUCOPHAGE™.

[0099] Examples of insulin secretion enhancers include but
15 are not limited to glucagon receptor antagonists (as described above), sulphonyl urea derivatives, incretin hormones or mimics thereof, especially glucagon-like peptide-1 (GLP-1) or GLP-1 agonists, beta-cell imidazoline receptor antagonists, and short-acting insulin secretagogues,
20 like antidiabetic phenylacetic acid derivatives, antidiabetic D-phenylalanine derivatives, and mitiglinide and pharmaceutical acceptable salts thereof.

[00100] Examples of sulphonyl urea derivatives include, but are not limited to, glisoxepid, glyburide, glibenclamide,
25 acetohexamide, chloropropamide, glibornuride, tolbutamide, tolazamide, glipizide, carbutamide, gliquidone, glyhexamide, phenbutamide, tolcyclamide; glimepiride and gliclazide. Tolbutamide, glibenclamide, gliclazide, glibornuride, gliquidone, glisoxepid and glimepiride can be administered
30 in the form that they are marketed under the trademarks RASTINON HOECHST™, AZUGLUCON™, DIAMICRONT™, GLUBORID™, GLURENORM™, PRO-DIABAN™ and AMARYL™, respectively.

[0101] Examples of GLP-1 agonists include, but are not limited to those disclosed in U.S. Patent Nos. 5,120,712, 5,118,666 and 5,512,549, and PCT Publication No. WO 91/11457.

In particular, GLP-1 agonists include those compounds like
5 GLP-1 (7-37) in which compound the carboxy-terminal amide functionality of Arg³⁶ is displaced with Gly at the 37th position of the GLP-1 (7-36)NH₂ molecule and variants and analogs thereof including GLN⁹-GLP-1 (7-37), D-GLN⁹-GLP-1 (7-37), acetyl LYS⁹-GLP-1 (7-37), LYS¹⁸-GLP-1 (7-37) and, in
10 particular, GLP-1 (7-37)OH, VAL⁸-GLP-1 (7-37), GLY⁸-GLP-1 (7-37), THR⁸-GLP-1 (7-37), GLP-1 (7-37) and 4-imidazopropionyl-GLP-1.

[0102] One particular example of a GLP-1 agonist is
Extendatide, a 39-amino acid peptide amide, which is
15 marketed under the trademark BYETTA™. Exenatide has the empirical formula C₁₈₄H₂₈₂N₅₀O₆₀S and molecular weight of 4186.6 Daltons. The amino acid sequence for Exenatide is as follows: H-His-Gly-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-
20 Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-NH₂

[0103] Examples of glucagon-like peptide-2 (GLP-2) or GLP-2 agonists include, but are not limited to those disclosed in U.S. Patent No. 7,056,886 and PCT Publication Nos. WO 00/53208, WO 01/49314 and WO 03/099854. One particular
25 example of a GLP-2 agonist is TEDUGLUTIDE™, a 39-amino acid peptide amide (NPS Pharmaceuticals, Inc.).

[0104] Examples of beta-cell imidazoline receptor antagonists include, but are not limited to those described in PCT Publication No. WO 00/78726 and J. Pharmacol. Exp.
30 Ther. 1996; 278; 82-89.

[0105] An example of an antidiabetic phenylacetic acid derivative is repaglinide and pharmaceutically acceptable salts thereof.

[0106] Examples of antidiabetic D-phenylalanine derivatives include, but are not limited to nateglinide (N-[(trans4-isopropylcyclohexyl)-carbonyl]-D-phenylalanine, EP 196222 and EP 526171) and repaglinide ((S)-2-ethoxy-4-{2-[[3-methyl-1-1-[2-(1-piperidinyl)phenyl]butyl]-amino]-2-oxoethyl}benzoic acid, EP 0 147 850 A2 and EP 0 207 331 A1).

Nateglinide is intended to include the particular crystal forms (polymorphs) disclosed in U.S. Patent No. 5,488,510 and European Patent Publication No. EP 0526171 B1.

Repaglinide and nateglinide may be administered in the form as they are marketed under the trademarks NOVONORM™ and STARLIX™, respectively.

[0107] Examples of alpha-Glucosidase inhibitors include, but are not limited to, acarbose, N-(1,3-dihydroxy-2-propyl)valiolamine (voglibose) and the 1-deoxynojirimycin derivative miglitol. Acarbose is 4",6"-dideoxy-4'-[(1S)-(1,4,6/5)-4,5,6-trihydroxy-3-hydroxymethyl-2-cyclohexenylamino)maltotriose. The structure of acarbose can as well be described as O-4,6-dideoxy-4-{[1S,4R,5S,6S]-4,5,6-trihydroxy-3-(hydroxymethyl)-2-cyclohexen-1-yl]-amino)-alpha-D-glucopyranosyl-(1-4)-O-alpha-D-glucopyranosyl-(1-4)-D-glucopyranose. (U.S. Patent No. 4,062,950 and European Patent Publication No. EP 0 226 121). Acarbose and miglitol may be administered in the forms that they are marketed under the trademarks GLUCOBAY™ and DIASTABOL 50™ respectively.

[0108] Examples of inhibitors of gastric emptying other than GLP-1 include, but are not limited to those disclosed in J. Clin. Endocrinol. Metab. 2000, 85(3), 1043-1048, and

Diabetes Care 1998; 21; 897-893, especially Amylin and analogs thereof such as pramlintide. Amylin is described in Diabetologia 39, 1996, 492-499.

[0109] Examples of α_2 -adrenergic antagonists include, but
5 are not limited to midaglizole which is described in Diabetes 36, 1987, 216-220. The insulin that may be used in combination with Compound I include, but are not limited to animal insulin preparations extracted from the pancreas of bovine and pig; human insulin preparations genetically
10 synthesized using *Escherichia coli* or yeast; zinc insulin; protamine zinc insulin; fragment or derivative of insulin (e.g., INS-1) and an oral insulin preparation.

[0110] In one particular embodiment, the antidiabetic compound administered in combination with Compound I is
15 selected from the group consisting of nateglinide, mitiglinide, repaglinide, metformin, extendatide, rosiglitazone, tesaglitazar, pioglitazone, glisoxepid, glyburide, glibenclamide, acetohexamide, chloropropamide, glibornuride, tolbutamide, tolazamide, glipizide,
20 carbutamide, gliquidone, glyhexamide, phenbutamide, tolcyclamide, glimepiride and gliclazide, including any pharmaceutically acceptable salts thereof.

[0111] Examples of the preparation and formulation of PTPase inhibitors, GSK-3 inhibitors, non-small molecule
25 mimetic compounds, GFAT inhibitors, G6Pase inhibitors, glucagon receptor antagonists, PEPCK inhibitors, F-1,6-BPase inhibitors, GP inhibitors, RXR modulators, Beta-3 AR agonists, PDHK inhibitors, inhibitors of gastric emptying and UCP modulators are disclosed in the patents,
30 applications and references provided herein.

[0112] In the case of combination therapy with Compound I, the other antidiabetic compound may be administered (e.g., route and dosage form) in a manner known per se for such

compound. Compound I and the other antidiabetic compound may be administered sequentially (*i.e.*, at separate times) or at the same time, either one after the other separately in two separate dose forms or in one combined, single dose form. In one particular embodiment, the other antidiabetic compound is administered with Compound I as a single, combined dosage form. The dose of the antidiabetic compound may be selected from the range known to be clinically employed for such compound. Any of therapeutic compounds of diabetic complications, antihyperlipemic compounds, antiobestic compounds or antihypertensive compounds can be used in combination with Compound I in the same manner as the above antidiabetic compounds. Examples of therapeutic compounds of diabetic complications include, but are not limited to, aldose reductase inhibitors such as tolrestat, epalrestat, zenarestat, zopolrestat, minalrestat, fidarestat, CT-112 and ranirestat; neurotrophic factors and increasing compounds thereof such as NGF, NT-3, BDNF and neurotrophin production-secretion promoters described in WO01/14372 (*e.g.*, 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(2-methylphenoxy)propyl]oxazole); neuranagenesis stimulators such as Y-128; PKC inhibitors such as ruboxistaurin mesylate; AGE inhibitors such as ALT946, pimagedine, N-phenacylthiazolium bromide (ALT766), ALT-711, EXO-226, pyridorin and pyridoxamine; reactive oxygen scavengers such as thiocctic acid; cerebral vasodilators such as tiapride and mexiletine; somatostatin receptor agonists such as BIM23190; and apoptosis signal regulating kinase-1 (ASK-1) inhibitors.

Examples of antihyperlipemic compounds include, but are not limited to, HMG-CoA reductase inhibitors such as pravastatin, simvastatin, lovastatin, atorvastatin, fluvastatin, rosuvastatin and pitavastatin; squalene synthase inhibitors such as compounds described in WO97/10224 (*e.g.*, N-

[[(3R, 5S) -1- (3-acetoxy-2,2-dimethylpropyl) -7-chloro-5- (2,3-dimethoxyphenyl) -2-oxo-1,2,3,5-tetrahydro-4,1-benzoxazepin-3-yl]acetyl]piperidine-4-acetic acid); fibrate compounds such as bezafibrate, clofibrate, simfibrate and
5 clinofibrate; ACAT inhibitors such as avasimibe and eflucimibe; anion exchange resins such as colestyramine; probucol; nicotinic acid drugs such as nicomol and niceritrol; ethyl icosapentate; and plant sterols such as soysterol and γ -oryzanol. Examples of antiobestic compounds
10 include, but are not limited to, dexfenfluramine, fenfluramine, phentermine, sibutramine, amfepramone, dexamphetamine, mazindol, phenylpropanolamine, clobenzorex; MCH receptor antagonists such as SB-568849 and SNAP-7941; neuropeptide Y antagonists such as CP-422935; cannabinoid
15 receptor antagonists such as SR-141716 and SR-147778; ghrelin antagonist; 11β -hydroxysteroid dehydrogenase inhibitors such as BVT-3498; pancreatic lipase inhibitors such as orlistat and ATL-962; Beta-3 AR agonists such as AJ-9677; peptidic anorexiant such as leptin and CNTF (Ciliary
20 Neurotropic Factor); cholecystokinin agonists such as lintitript and FPL-15849; and feeding deterrent such as P-57.

Examples of the antihypertensive compounds include angiotensin converting enzyme inhibitors such as captopril, enalapril and delapril; angiotensin II antagonists such as
25 candesartan cilexetil, losartan, eprosartan, valsartan, telmisartan, irbesartan, olmesartan medoxomil, tasosartan and 1-[[2'-(2,5-dihydro-5-oxo-4H-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]methyl]-2-ethoxy-1H-benzimidazole-7-carboxylic acid; calcium channel blockers such as manidipine,
30 nifedipine, nicardipine, amlodipine and efonidipine; potassium channel openers such as levcromakalim, L-27152, AL0671 and NIP-121; clonidine; deacetylases such as

reservatrol, sirtuin agonist; polyphenols; MCR4 agonist;
Sodium dependent glucose transport (SGLT2) inhibitors.

[0113] The structure of the active agents identified herein
by code nos., generic or trade names may be taken from the
5 actual edition of the standard compendium "The Merck Index"
or from databases, e.g. Patents International (e.g. IMS
World Publications). The corresponding content thereof is
hereby incorporated by reference. Any person skilled in the
art is fully enabled to identify the active agents and,
10 based on these references, likewise enabled to manufacture
and test the pharmaceutical indications and properties in
standard test models, both in vitro and in vivo.

4. COMPOSITIONS COMPRISING COMPOUND I

[0114] Compound I may be comprised within a pharmaceutical
15 composition adapted for a variety of routes of
administration. For example, Compound I may be comprised
within a pharmaceutical composition adapted to be
administered by a route selected from the group consisting
of orally, parenterally, intraperitoneally, intravenously,
20 intraarterially, transdermally, sublingually,
intramuscularly, rectally, transbuccally, intranasally,
liposomally, via inhalation, vaginally, intraocularly, via
local delivery (for example by catheter or stent),
subcutaneously, intraadiposally, intraarticularly,
25 intraperitoneally and intrathecally. As such, Compound I
may be formulated in a variety of pharmaceutically
acceptable compositions including injectable forms (e.g.,
subcutaneous, intravenous, intramuscular and intraperitoneal
injections), drip infusions, external application forms
30 (e.g., nasal spray preparations, transdermal preparations;
ointments, etc.), and suppositories (e.g., rectal and
vaginal suppositories). These different pharmaceutically
acceptable compositions can be manufactured by known

techniques conventionally used in the pharmaceutical industry with a pharmaceutically acceptable carrier conventionally used in the pharmaceutical industry.

[0115] As used herein, a composition comprising Compound I is intended to encompass the free base form of Compound I, salts, hydrates and prodrugs of Compound I, as well as other materials that may be included in such composition for its intended purpose, including other active ingredients, unless otherwise specified. Particular salt forms of Compound I that may be employed include, but are not limited to, the HCl, methanesulfonate, succinate, benzoate, toluenesulfonate, R-(-)mandelate or benzenesulfonate salt forms of Compound I.

[0116] As noted above, Compound I may advantageously be used when administered to a patient at a weekly dose of between 1 mg/week and 500 mg/week of Compound, optionally between 12.5 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 400 mg/week, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and optionally between 100 mg/week and 400 mg/week of Compound. Specific dosage amounts that may be used include, but are not limited to 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg 250 mg, 400 mg and 500 mg of Compound I per week. As also noted above, it is desirable for Compound I to be administered one time per week. Accordingly, pharmaceutical compositions of the present invention may be in the form of a single dose form comprising between 1 mg/week and 500 mg/week of Compound I, optionally between 12.5 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 400 mg/week, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and

optionally between 100 mg/week and 400 mg/week of Compound I.

In specific embodiments, the pharmaceutical composition comprises 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg 250 mg, 400 mg and 500 mg of Compound I. In each instance, the dosage amounts and ranges of Compound I provided is based on the molecular weight of the free base form of Compound I.

[0117] Compound I also may advantageously be used when administered to a patient at a weekly dose of more than 250 mg of Compound I, optionally at least 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at least 350 mg of Compound I, and optionally at least 400 mg of Compound I. In one variation, the weekly dose of Compound I administered to the patient is not more than 500 mg. In another variation, the weekly dose of Compound I administered to the patient is not more than 400 mg. In still another variation, the weekly dose of Compound I administered to the patient is not more than 350 mg. In yet another variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 500 mg. In a further variation, the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 400 mg. Specific dosage amounts that may be used include, but are not limited to 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I per week. In each instance, the dosage amounts and ranges of Compound I provided is based on the molecular weight of the free base form of Compound I.

[0118] As also noted above, it is desirable for Compound I to be administered one time per week. Accordingly, pharmaceutical compositions of the present invention may be in the form of a single dose form comprising a dose of between 1 mg/week and 500 mg/week of Compound I to a patient,

optionally between 12.5 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 400 mg/week, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and
5 optionally between 100 mg/week and 400 mg/week of Compound. In other variations, the pharmaceutical composition comprises 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg, 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I. In each instance,
10 the dosage amounts and ranges of Compound I provided is based on the molecular weight of the free base form of Compound I. [0119] Pharmaceutical compositions of the present invention also may be in the form of a single dose form comprising a dose of more than 250 mg of Compound I, optionally at least
15 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at least 350 mg of Compound I, and optionally at least 400 mg of Compound I. In one variation, the pharmaceutical composition comprises not more than 500 mg of Compound I. In another variation, the pharmaceutical
20 composition comprises not more than 400 mg of Compound I. In still another variation, the pharmaceutical composition comprises not more than 350 mg of Compound I. In yet another variation, the pharmaceutical composition comprises more than 250 mg and not more than 500 mg of Compound I. In
25 a further variation, the pharmaceutical composition comprises more than 250 mg and not more than 400 mg of Compound I. In other variations, the pharmaceutical composition comprises 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I. In each instance, the dosage amounts and
30 ranges of Compound I provided is based on the molecular weight of the free base form of Compound I.

[0120] As also noted above, Compound I may advantageously be used when administered orally. Accordingly, the

compositions of the present invention may optionally be adapted for oral administration. In one variation, such pharmaceutical composition is a solid formulation adapted for oral administration. In this regard, the composition, for example, may be in the form of a tablet or capsule. In another variation, such pharmaceutical composition is a liquid formulation adapted for oral administration.

[0121] As also noted above, Compound I may advantageously be used when administered parenterally. Accordingly, the compositions of the present invention may optionally be adapted for parenteral administration. In one variation, such pharmaceutical composition is a solution formulation adapted for parenteral administration. In another variation, such pharmaceutical composition is a suspension formulation adapted for parenteral administration.

[0122] As noted above, Compound I may advantageously be used in combination with one or more other antidiabetic and/or incretin compounds. Accordingly, the compositions of the present invention may optionally comprises Compound I in combination with one or more other antidiabetic or incretin compounds in a combined, single dose form. Optionally, such combined, single dose form comprising Compound I in combination with one or more other antidiabetic and/or incretin compounds is adapted for oral administration and optionally is a solid oral dose form. Alternatively, such combined, single dose form comprising Compound I in combination with one or more other antidiabetic and/or incretin compounds can be adapted for parenteral administration and optionally is a solution dose form.

[0123] In one variation, such combined, single dose form comprising Compound I in combination with one or more other antidiabetic compounds comprises a dose of between 1 mg/week and 500 mg/week of Compound I to a patient, optionally

between 12.5 mg/week and 400 mg/week of Compound I, optionally between 20 mg/week and 400 mg/week, optionally between 20 mg/week and 200 mg/week of Compound I, optionally between 50 mg/week and 400 mg/week of Compound I, and
5 optionally between 100 mg/week and 400 mg/week of Compound. In other variations, the pharmaceutical composition comprises 3.125 mg, 12.5 mg, 20 mg, 25 mg, 50 mg, 75 mg, 100 mg, 125 mg, 150 mg, 175 mg, 200 mg, 250 mg, 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I.
10 [0124] In another variation, such combined, single dose form comprising Compound I in combination with one or more other antidiabetic compounds comprises a dose of more than 250 mg of Compound I, optionally at least 275 mg of Compound I, optionally at least 300 mg of Compound I, optionally at
15 least 350 mg of Compound I, and optionally at least 400 mg of Compound I. In one variation, the pharmaceutical composition comprises not more than 500 mg of Compound I. In another variation, the pharmaceutical composition comprises not more than 400 mg of Compound I. In still
20 another variation, the pharmaceutical composition comprises not more than 350 mg of Compound I. In yet another variation, the pharmaceutical composition comprises more than 250 mg and not more than 500 mg of Compound I. In a further variation, the pharmaceutical composition comprises
25 more than 250 mg and not more than 400 mg of Compound I. In other variations, the pharmaceutical composition comprises 275 mg, 300 mg, 350 mg, 400 mg or 500 mg of Compound I.
[0125] Any antidiabetic compound, or set of antidiabetic compounds may be combined with Compound I to form such
30 combined, single dose form. In particular embodiments, such combined, single dose form includes Compound I and one or more members of the group consisting of insulin signaling pathway modulators, like protein tyrosine phosphatase

(PTPase) inhibitors, and glutamine-fructose-6-phosphate
 amidotransferase (GFAT) inhibitors, compounds influencing a
 dysregulated hepatic glucose production, like glucose-6-
 phosphatase (G6Pase) inhibitors, fructose-1,6-bisphosphatase
 5 (F-1,6-BPase) inhibitors, glycogen phosphorylase (GP)
 inhibitors, glucagon receptor antagonists and
 phosphoenolpyruvate carboxykinase (PEPCK) inhibitors,
 pyruvate dehydrogenase kinase (PDHK) inhibitors, insulin
 sensitivity enhancers (insulin sensitizers), insulin
 10 secretion enhancers (insulin secretagogues), alpha-
 glucosidase inhibitors, inhibitors of gastric emptying,
 glucokinase activators, GLP-1 receptor agonists, GLP-2
 receptor agonists, UCP modulators, RXR modulators, GSK-3
 inhibitors, PPAR modulators, metformin, insulin, and α_2 -
 15 adrenergic antagonists. Compound I may be administered with
 such at least one other antidiabetic compound either
 simultaneously as a single dose, at the same time as
 separate doses, or sequentially (i.e., where one is
 administered before or after the other is administered).
 20 [0126] In one variation, such combined, single dose form
 comprises Compound I and an antidiabetic thiazolidinedione.
 Particular examples of thiazolidinediones that may be used
 in this variation include, but are not limited to (S)-((3,4-
 dihydro-2-(phenyl-methyl)-2H-1-benzopyran-6-yl)methyl-
 25 thiazolidine-2,4-dione (englitazone), 5-{{[4-(3-(5-methyl-2-
 phenyl-4-oxazolyl)-1-oxo-propyl)-phenyl]-methyl}-
 thiazolidine-2,4-dione (darglitazone), 5-{{[4-(1-methyl-
 cyclohexyl)methoxy)-phenyl]methyl}-thiazolidine-2,4-dione
 (ciglitazone), 5-{{[4-(2-(1-indolyl)ethoxy)phenyl]methyl}-
 30 thiazolidine-2,4-dione (DRF2189), 5-{4-[2-(5-methyl-2-
 phenyl-4-oxazolyl)-ethoxy]benzyl}-thiazolidine-2,4-dione
 (BM-13.1246), 5-(2-naphthylsulfonyl)-thiazolidine-2,4-dione
 (AY-31637), bis{4-[(2,4-dioxo-5-thiazolidinyl)-

methyl]phenyl}methane (YM268), 5-{4-[2-(5-methyl-2-phenyl-4-oxazolyl)-2-hydroxyethoxy]-benzyl}-thiazolidine-2,4-dione (AD-5075), 5-[4-(1-phenyl-1-cyclopropanecarbonylamino)-benzyl]-thiazolidine-2,4-dione (DN-108) 5-{[4-(2-(2,3-dihydroindol-1-yl)ethoxy)phenylmethyl]-thiazolidine-2,4-dione, 5-[3-(4-chloro-phenyl))-2--propynyl]-5-phenylsulfonyl}thiazolidine-2,4-dione, 5-[3-(4-chlorophenyl))-2-propynyl]-5-(4-fluorophenyl-sulfonyl)thiazolidine-2,4-dione, 5-{[4-(2-(methyl-2-pyridinyl-amino)-ethoxy)phenyl]methyl}-thiazolidine-2,4-dione (rosiglitazone), 5-{[4-(2-(5-ethyl-2-pyridyl)ethoxy)phenyl]-methyl}-thiazolidine-2,4-dione (pioglitazone), 5-[6-(2-fluoro-benzyloxy)-naphthalen-2-ylmethyl]-thiazolidine-2,4-dione (MCC555), 5-([2-(2-naphthyl)-benzoxazol-5-yl]-methyl)thiazolidine-2,4-dione (T-174), edaglitazone (BM-13-1258), rivoglitazone (CS-011) and 5-(2,4-dioxothiazolidin-5-ylmethyl)-2-methoxy-N-(4-trifluoromethyl-benzyl)benzamide (KRP297).

[0127] In one particular variation, the thiazolidinedione in such combined, single dose form is 5-{[4-(2-(5-ethyl-2-pyridyl)ethoxy)phenyl]-methyl}-thiazolidine-2,4-dione (pioglitazone) and its hydrochloride salt which is marketed under the trademark ACTOS™.

[0128] In another particular variation, the thiazolidinedione is 5-{[4-(2-(methyl-2-pyridinyl-amino)-ethoxy)phenyl]methyl}-thiazolidine-2,4-dione (rosiglitazone) and its maleate salt.

[0129] In another variation, such combined, single dose form comprises Compound I and a non-glitazone type PPAR gamma agonist.

[0130] In another variation, such combined, single dose form comprises Compound I and a biguanide. A particular example of a biguanide that may be used in this variation is

Metformin (dimethyldiguanide) and its hydrochloride salt which is marketed under the trademark GLUCOPHAGE™.

[0131] In another variation, such combined, single dose form comprises Compound I and a sulphonyl urea derivative. Particular examples of sulphonyl urea derivatives that may be used in this variation include, but are not limited to glisoxepid, glyburide, glibenclamide, acetohexamide, chloropropamide, glibornuride, tolbutamide, tolazamide, glipizide, carbutamide, gliquidone, glyhexamide, phenbutamide, tolcyclamide, glimepiride and gliclazide. Tolbutamide, glibenclamide, gliclazide, glibornuride, gliquidone, glisoxepid and glimepiride can be administered in the form as they are marketed under the trademarks RASTINON HOECHST™, AZUGLUCON™, DIAMICRONT™, GLUBORID™, GLURENORM™, PRO-DIABAN™ and AMARYL™, respectively.

[0132] In another variation, such combined, single dose form comprises Compound I and an antidiabetic D-phenylalanine derivative. Particular examples of antidiabetic D-phenylalanine derivatives that may be used in this variation include, but are not limited to repaglinide and nateglinide which may be administered in the form as they are marketed under the trademarks NOVONORM™ and STARLIX™, respectively.

[0133] In another variation, such combined, single dose form comprises Compound I and an alpha-Glucosidase inhibitor. Particular examples of alpha-Glucosidase inhibitors that may be used in this variation include, but are not limited to acarbose, miglitol and voglibose which may be administered in the form as they are marketed under the trademarks GLUCOBAY™, DIASTABOL 50™ and BASEN™, respectively.

[0134] In one particular embodiment, the antidiabetic compound administered in combination with Compound I in such combined, single dose form is selected from the group consisting of nateglinide, repaglinide, metformin, extendatide, rosiglitazone, pioglitazone, glisoxepid, glyburide, glibenclamide, acetohexamide, chloropropamide, glibornuride, tolbutamide, tolazamide, glipizide, carbutamide, gliquidone, glyhexamide, phenbutamide, tolcyclamide, glimepiride and gliclazide, including any pharmaceutically acceptable salts thereof.

[0135] In regard to each of the above embodiments and variations regarding a combined, single dose form comprising the combination of Compound I and one or more other antidiabetic compounds, the pharmaceutical composition may optionally be adapted for oral administration and in this regard may optionally be a solid formulation such as a tablet or capsule or may alternatively be in a liquid formulation adapted for oral administration. The dose of the antidiabetic compound may be selected from the range known to be clinically employed for such compound. Any of therapeutic compounds of diabetic complications, antihyperlipemic compounds, antiobestic compounds or antihypertensive compounds can be used in combination with Compound I in the same manner as the above antidiabetic compounds. Examples of therapeutic compounds of diabetic complications include, but are not limited to, aldose reductase inhibitors such as tolrestat, epalrestat, zenarestat, zopolrestat, minalrestat, fidarestat, CT-112 and ranirestat; neurotrophic factors and increasing compounds thereof such as NGF, NT-3, BDNF and neurotrophin production-secretion promoters described in WO01/14372 (e.g., 4-(4-chlorophenyl)-2-(2-methyl-1-imidazolyl)-5-[3-(2-methylphenoxy)propyl]oxazole); neuranogenesis stimulators

such as Y-128; PKC inhibitors such as ruboxistaurin mesylate; AGE inhibitors such as ALT946, pimagedine, N-phenacylthiazolium bromide (ALT766), ALT-711, EXO-226, pyridorin and pyridoxamine; reactive oxygen scavengers such
5 as thiocctic acid; cerebral vasodilators such as tiapride and mexiletine; somatostatin receptor agonists such as BIM23190; and apoptosis signal regulating kinase-1 (ASK-1) inhibitors.

Examples of antihyperlipemic compounds include, but are not limited to, HMG-CoA reductase inhibitors such as pravastatin,
10 simvastatin, lovastatin, atorvastatin, fluvastatin, rosuvastatin and pitavastatin; squalene synthase inhibitors such as compounds described in WO97/10224 (e.g., N-[[[(3R,5S)-1-(3-acetoxy-2,2-dimethylpropyl)-7-chloro-5-(2,3-dimethoxyphenyl)-2-oxo-1,2,3,5-tetrahydro-4,1-benzoxazepin-
15 3-yl]acetyl]piperidine-4-acetic acid); fibrate compounds such as bezafibrate, clofibrate, simfibrate and clinofibrate; ACAT inhibitors such as avasimibe and eflucimibe; anion exchange resins such as colestyramine; probucol; nicotinic acid drugs such as nicomol and
20 niceritrol; ethyl icosapentate; and plant sterols such as soysterol and γ -oryzanol. Examples of antiobestic compounds include, but are not limited to, dexfenfluramine, fenfluramine, phentermine, sibutramine, amfepramone, dexamphetamine, mazindol, phenylpropanolamine, clobenzorex;
25 MCH receptor antagonists such as SB-568849 and SNAP-7941; neuropeptide Y antagonists such as CP-422935; cannabinoid receptor antagonists such as SR-141716 and SR-147778; ghrelin antagonist; 11 β -hydroxysteroid dehydrogenase inhibitors such as BVT-3498; pancreatic lipase inhibitors
30 such as orlistat and ATL-962; Beta-3 AR agonists such as AJ-9677; peptidic anorexiant such as leptin and CNTF (Ciliary Neurotropic Factor); cholecystokinin agonists such as

linitript and FPL-15849; and feeding deterrent such as P-57.

Examples of the antihypertensive compounds include angiotensin converting enzyme inhibitors such as captopril, enalapril and delapril; angiotensin II antagonists such as
5 candesartan cilexetil, losartan, eprosartan, valsartan, telmisartan, irbesartan, olmesartan medoxomil, tasosartan and 1-[[2'-(2,5-dihydro-5-oxo-4H-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]methyl]-2-ethoxy-1H-benzimidazole-7-carboxylic acid; calcium channel blockers such as manidipine,
10 nifedipine, nicardipine, amlodipine and efonidipine; potassium channel openers such as levcromakalim, L-27152, AL0671 and NIP-121; and clonidine.

5. KITS AND ARTICLES OF MANUFACTURE COMPRISING COMPOUND I

[0136] The present invention also relates to kits
15 comprising a pharmaceutical composition according to the present invention comprising Compound I (and optionally one or more other antidiabetic or incretin compounds) where such kit further comprises instructions that include one or more forms of information selected from the group consisting of
20 indicating a disease state for which the pharmaceutical composition is to be administered, storage information for the pharmaceutical composition, dosing information and instructions regarding how to administer the pharmaceutical composition. The kit may also comprise packaging materials.
25 The packaging material may also comprise a container for housing the pharmaceutical composition. The container may optionally comprise a label indicating the disease state for which the pharmaceutical composition is to be administered, storage information, dosing information and/or instructions
30 regarding how to administer the composition. The kit may also comprise additional components for storage or administration of the composition. The kit may also comprise the composition in single or multiple dose forms.

[0137] In one embodiment, the pharmaceutical composition in the kit comprises multiple doses of a pharmaceutical composition according to the present invention wherein such pharmaceutical composition is a single dose form that
5 comprises Compound I in one of the dosage ranges specified herein.

[0138] In another embodiment, the pharmaceutical composition in the kit comprises multiple doses of a pharmaceutical composition according to the present
10 invention wherein such pharmaceutical composition is a single dose form that comprises Compound I and one or more of the other antidiabetic or incretin compounds specified herein.

[0139] The present invention also relates to articles of
15 manufacture comprising a pharmaceutical composition according to the present invention comprising Compound I (and optionally one or more other antidiabetic or incretin compounds) where such articles of manufacture further comprise packaging materials. In one variation, the
20 packaging material comprises a container for housing the composition. In another variation, the invention provides an article of manufacture where the container comprises a label indicating one or more members of the group consisting of a disease state for which the composition is to be
25 administered, storage information, dosing information and/or instructions regarding how to administer the composition.

[0140] In one embodiment, the pharmaceutical composition in the article of manufacture comprises multiple doses of a pharmaceutical composition according to the present
30 invention wherein such pharmaceutical composition is a single dose form that comprises Compound I in one of the dosage ranges specified herein.

[0141] In another embodiment, the pharmaceutical composition in the article of manufacture comprises multiple doses of a pharmaceutical composition according to the present invention wherein such pharmaceutical composition is a single dose form that comprises Compound I and one or more of the other antidiabetic or incretin compounds specified herein.


[0142] It is noted that the packaging material used in kits and articles of manufacture according to the present invention may form a plurality of divided containers such as a divided bottle or a divided foil packet. The container can be in any conventional shape or form as known in the art which is made of a pharmaceutically acceptable material, for example a paper or cardboard box, a glass or plastic bottle or jar, a re-sealable bag (for example, to hold a "refill" of tablets for placement into a different container), or a blister pack with individual doses for pressing out of the pack according to a therapeutic schedule. The container that is employed will depend on the exact dosage form involved. It is feasible that more than one container can be used together in a single package to market a single dosage form. For example, tablets may be contained in a bottle that is in turn contained within a box.

[0143] One particular example of a kit according to the present invention is a so-called blister pack. Blister packs are well known in the packaging industry and are being widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material (preferably stiff transparent plastic material) covered with a foil. During the packaging process recesses are formed in the stiff material. The recesses have the size and shape of individual tablets or capsules to be packed or may have the

size and shape to accommodate multiple tablets and/or capsules to be packed. Next, the tablets or capsules are placed in the recesses accordingly and the sheet of relatively stiff material is sealed against the plastic foil at the face of the foil which is opposite from the direction in which the recesses were formed. As a result, the tablets or capsules are individually sealed or collectively sealed, as desired, in the recesses between the foil and the sheet. The strength of the sheet is preferably such that the tablets or capsules can be removed from the blister pack by manually applying pressure on the recesses whereby an opening is formed in the foil at the place of the recess. The tablet or capsule can then be removed via said opening.

EXAMPLES

1. Preparation of 2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluorobenzonitrile and pharmaceutically acceptable salts



4-Fluoro-2-methylbenzonitrile (3)

- [0144] A mixture of 2-bromo-5-fluorotoluene (2) (3.5 g, 18.5 mmol) and CuCN (2 g, 22 mmol) in DMF (100 mL) was

refluxed for 24 hours. The reaction was diluted with water and extracted with hexane. The organics were dried over MgSO_4 and the solvent removed to give product 3 (yield 60%).

^1H -NMR (400 MHz, CDCl_3): δ 7.60 (dd, $J=5.6, 8.8$ Hz, 1H),
5 6.93-7.06 (m, 2H), 2.55 (s, 3H).


2-Bromomethyl-4-fluorobenzonitrile (4)

[0145] A mixture of 4-fluoro-2-methylbenzonitrile (3) (2 g, 14.8 mmol), NBS (2.64 g, 15 mmol) and AIBN (100 mg) in CCl_4 was refluxed under nitrogen for 2 hours. The reaction was
10 cooled to room temperature. The solid was removed by filtration. The organic solution was concentrated to give crude product as an oil, which was used in the next step without further purification. ^1H -NMR (400 MHz, CDCl_3): δ
7.68 (dd, $J=5.2, 8.4$ Hz, 1H), 7.28 (dd, $J=2.4, 8.8$ Hz, 1H),
15 7.12 (m, 1H), 4.6 (s, 2H).

2-(6-Chloro-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl)-4-fluoro-benzonitrile (6)

[0146] A mixture of crude 3-methyl-6-chlorouracil (5) (0.6 g, 3.8 mmol), 2-Bromomethyl-4-fluorobenzonitrile (0.86 g, 4 mmol) and K_2CO_3 (0.5 g, 4 mmol) in DMSO (10 mL) was stirred
20 at 60 °C for 2 hours. The reaction was diluted with water and extracted with EtOAc. The organics were dried over MgSO_4 and the solvent removed. The residue was purified by column
25 chromatography. 0.66 g of the product was obtained (yield: 60%). ^1H -NMR (400 MHz, CDCl_3): δ 7.73 (dd, $J=7.2, 8.4$ Hz, 1H), 7.26 (d, $J=4.0$ Hz, 1H), 7.11-7.17 (m, 1H), 6.94 (dd, $J=2.0, 9.0$ Hz, 1H), 6.034 (s, 2H), 3.39 (s, 3H). MS (ES) [m+H] calc'd for $\text{C}_{13}\text{H}_9\text{ClFN}_3\text{O}_2$, 293.68; found 293.68.

2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile, TFA salt (1) (TFA salt of Compound I)




• TFA

- 5 [0147] 2-(6-Chloro-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl)-4-fluoro-benzonitrile (5) (300 mg, 1.0 mmol), (R)-3-amino-piperidine dihydrochloride (266 mg, 1.5 mmol) and sodium bicarbonate (500 mg, 5.4 mmol) were stirred in a sealed tube in EtOH (3 mL) at 100 °C for 2 hrs.
- 10 The final compound was obtained as a TFA salt after HPLC purification. ¹H-NMR (400 MHz, CD₃OD): δ. 7.77-7.84 (m, 1H), 7.16-7.27 (m, 2H), 5.46 (s, 1H), 5.17-5.34 (ABq, 2H, J = 35.2, 15.6 Hz), 3.33-3.47 (m, 2H), 3.22 (s, 3H), 2.98-3.08 (m, 1H), 2.67-2.92 (m, 2H), 2.07-2.17 (m, 1H), 1.82-1.92 (m, 1H), 1.51-1.79 (m, 2H). MS (ES) [m+H] calc'd for C₁₈H₂₀FN₅O₂, 357.38; found, 357.38.
- 15

2-[6-(3-Amino-piperidin-1-yl)-3-methyl-2,4-dioxo-3,4-dihydro-2H-pyrimidin-1-ylmethyl]-4-fluoro-benzonitrile, HCl salt

20



• HCl

[0148] The TFA salt of Compound I was suspended in DCM, and then washed with saturated Na₂CO₃. The organic layer was dried and removed in vacuo. The residue was dissolved in

acetonitrile and HCl in dioxane (1.5 eq.) was added at 0 °C.

The HCl salt was obtained after removing the solvent. ¹H-NMR (400 MHz, CD₃OD): δ. 7.77-7.84 (m, 1H), 7.12-7.26 (m, 2H), 5.47 (s, 1H), 5.21-5.32 (ABq, 2H, J = 32.0, 16.0 Hz), 3.35-3.5 (m, 2H), 3.22 (s, 3H), 3.01-3.1 (m, 1H), 2.69-2.93 (m, 2H), 2.07-2.17 (m, 1H), 1.83-1.93 (m, 1H), 1.55-1.80 (m, 2H). MS (ES) [m+H] calc'd for C₁₈H₂₀FN₅O₂, 357.38; found, 357.38.

General procedure for the preparation of salts of Compound I [0149] The benzonitrile product may be isolated as the free base if desired, but preferably, the product may be further converted to a corresponding acid addition salt.

Specifically, the benzonitrile product (approximately 10 mg) in a solution of MeOH (1 mL) was treated with various acids (1.05 equivalents). The solutions were allowed to stand for three days open to the air. If a precipitate formed, the mixture was filtered and the salt dried. If no solid formed, the mixture was concentrated in vacuo and the residue isolated. In this way, salts of Compound I were prepared from the following acids: benzoic, p-toluenesulfonic, succinic, R-(-)-Mandelic and benzenesulfonic.

[0150] The isolation and/or purification steps of the intermediate compounds in the above described process may optionally be avoided if the intermediates from the reaction mixture are obtained as relatively pure compounds and the by-products or impurities of the reaction mixture do not interfere with the subsequent reaction steps. Where feasible, one or more isolation steps may be eliminated to provide shorter processing times, and the elimination of further processing may also afford higher overall reaction yields.

2. EFFECT OF ADMINISTRATION ON PLASMA DPP-IV ACTIVITY

[0151] A single dose of Compound I was administered orally to 6 humans at a dosage of 12.5 mg, 25 mg, 50 mg, 100 mg, 200 mg and 400 mg, respectively (total 36 humans). Figure 1 illustrates the observed effect that administering Compound I has on the human plasma DPP-IV activity post dosing. As can be seen, Compound I reduced DPP-IV activity in human plasma by greater than 10% relative to baseline at 168 hours post dosing. Thus, as can be seen from the data shown in Figure 1, by administering Compound I one time per week at the dosage levels specified herein, Compound I can be effectively used relative to disease states where it is desired to reduce plasma DPP-IV activity. In view of the data presented, it is believed that when at least 50 mg of Compound I is administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 35% relative to baseline for a period of at least 168 hours following administration, when at least 100 mg of Compound I is administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 60% relative to baseline for a period of at least 168 hours following administration, and , when at least 200 mg of Compound I is administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 70% relative to baseline for a period of at least 168 hours following administration.

3. EFFECT OF ADMINISTRATION ON PLASMA DPP-IV ACTIVITY

[0152] A single dose of Compound I was administered orally to humans at a dosage of 3.125 mg (to 9 humans), 12.5 mg (to 8 humans), 50 mg (to 7 humans) and 100 mg (to 8 human), respectively. Figure 2 illustrates the observed effect that administering Compound I has on the human plasma DPP-IV

activity post dosing. As can be seen, Compound I reduced DPP-IV activity in human plasma by greater than 20% relative to baseline at 168 hours post dosing. Thus, as can be seen from the data shown in Figure 2, by administering Compound I one time per week at the dosage levels specified herein, Compound I can be effectively used relative to disease states where it is desired to reduce plasma DPP-IV activity.

In view of the data presented, it is believed that when at least 50 mg of Compound I is administered to a patient, the patient's plasma DPP-IV activity may be reduced by greater than 65% relative to baseline for a period of at least 168 hours following administration.

[0153] It will be apparent to those skilled in the art that various modifications and variations can be made to the compounds, compositions, kits, and methods of the present invention without departing from the spirit or scope of the invention. Thus, it is intended that the present invention cover the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents.

[0154] This application is based on US provisional application No. 60/894,624, the contents of which are incorporated hereinto by reference.

CLAIMS

1. A method comprising:

administering a weekly dose of more than 250 mg of
Compound I to a patient wherein the weekly dose is
5 administered once per week.

2. The method according to claim 1, wherein the weekly dose
of Compound I administered to the patient is at least 275 mg.

3. The method according to claim 1, wherein the weekly dose
of Compound I administered to the patient is at least 300 mg.

10 4. The method according to claim 1, wherein the weekly dose
of Compound I administered to the patient is at least 350 mg.

5. The method according to claim 1, wherein the weekly dose
of Compound I administered to the patient is at least 400 mg.

15 6. The method according to any one of claims 1-5, wherein the
weekly dose of Compound I administered to the patient is not
more than 500 mg.

7. The method according to any one of claims 1-4, wherein the
weekly dose of Compound I administered to the patient is not
more than 400 mg.

20 8. The method according to any one of claims 1-3, wherein the
weekly dose of Compound I administered to the patient is not
more than 350 mg.

25 9. The method according to claim 1, wherein the weekly dose
of Compound I administered to the patient is more than 250
mg and not more than 500 mg.

10. The method according to claim 1, wherein the weekly dose of Compound I administered to the patient is more than 250 mg and not more than 400 mg.
11. The method according to any one of claims 1-10, wherein
5 Compound I is administered as a free base.
12. The method according to any one of claims 1-10, wherein Compound I is administered as a pharmaceutically acceptable salt.
13. The method according to any one of claims 1-10, wherein
10 Compound I is administered as a succinate salt.
14. A pharmaceutical composition formulated in a single dose form comprising more than 250 mg of Compound I.
15. The pharmaceutical composition according to claim 14, wherein the single dose form comprises at least 275 mg of
15 Compound I.
16. The pharmaceutical composition according to claim 14, wherein the single dose form comprises at least 300 mg of Compound I.
17. The pharmaceutical composition according to claim 14,
20 wherein the single dose form comprises at least 350 mg of Compound I.
18. The pharmaceutical composition according to claim 14, wherein the single dose form comprises not more than 500 mg of Compound I.
- 25 19. The pharmaceutical composition according to claim 14, wherein the single dose form comprises not more than 400 mg of Compound I.

20. The pharmaceutical composition according to claim 14, wherein the single dose form comprises not more than 350 mg of Compound I.
21. The pharmaceutical composition according to claim 14, wherein the single dose form comprises more than 250 mg and not more than 500 mg of Compound I.
22. The pharmaceutical composition according to claim 14, wherein the single dose form comprises more than 250 mg and not more than 400 mg of Compound I.
23. The pharmaceutical composition according to any one of claims 14-22, wherein Compound I is present in the pharmaceutical composition as a free base.
24. The pharmaceutical composition according to any one of claims 14-22, wherein Compound I is present in the pharmaceutical composition in a pharmaceutically acceptable salt.
25. The pharmaceutical composition according to any one of claims 14-22, wherein Compound I is present in the pharmaceutical composition in a succinate salt.
26. Use of Compound I for the manufacture of a pharmaceutical composition according to any one of claims 14-25.
27. A method of treating diabetes comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week.
28. A method of treating cancer comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week.

29. A method of treating autoimmune disorders comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week.
- 5 30. A method of treating HIV infection comprising administering a weekly dose of more than 250 mg of Compound I to a patient wherein the weekly dose is administered once per week.

FIG. 1




FIG. 2




FIG. 2

