



(51) International Patent Classification:

C07D 401/12 (2006.01) *A61P 35/00* (2006.01)
C07D 401/14 (2006.01) *A61P 35/02* (2006.01)
A61K 31/438 (2006.01) *A61P 35/04* (2006.01)
A61K 31/444 (2006.01)

(21) International Application Number:

PCT/CN2013/000216

(22) International Filing Date:

1 March 2013 (01.03.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/606,291 2 March 2012 (02.03.2012) US

(71) Applicants: **GENENTECH, INC.** [US/US]; 1 DNA Way, South San Francisco, California 94080-4990 (US). **FORMA TM, LLC** [US/US]; 500 Arsenal Street, Suite 100, Watertown Massachusetts 02472 (US).

(72) Inventor; and

(71) Applicant (*for US only*): **YUEN, Po-Wai** [US/CN]; 6 Tai-He Road BDA, Beijing 100176 (CN).

(72) Inventors: **BAIR, Kenneth, W.**; 1 DNA Way, South San Francisco, California 94080-4990 (US). **BAUMEISTER, Timm, R.**; 500 Arsenal Street, Suite 100, Watertown, Massachusetts 02472 (US). **DRAGOVICH, Peter**; 1 DNA Way, South San Francisco, California 94080-4990 (US). **LIU, Xiongcai**; 6 Tai-He Road, BDA, Beijing 100176 (CN). **PATEL, Snahel**; 1 DNA Way, South San Francisco, California 94080-4990 (US). **ZAK, Mark**; 1 DNA Way, South San Francisco, California 94080-4990 (US). **ZHAO, Guiling**; 1 DNA Way, South San Francisco, California 94080-4990 (US). **ZHANG, Yamin**; 6 Tai-He Road, BDA, Beijing 100176 (CN). **ZHENG, Xiaozhang**; 500

Arsenal Street, Suite 100, Watertown, Massachusetts 02472 (US).

(74) Agent: **LIU, SHEN & ASSOCIATES**; A0601, Huibin Building, No. 8, Beichen Dong Street, Chaoyang District, Beijing 100101 (CN).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

— *with international search report (Art. 21(3))*

(54) Title: AMIDO SPIROCYCLIC AMIDE AND SULFONAMIDE DERIVATIVES

(57) Abstract: Provided are amido spirocyclic amide and sulfonamide compounds, pharmaceutical compositions comprising such compounds, and methods of treatment using such compounds.



AMIDO SPIROCYCLIC AMIDE AND SULFONAMIDE DERIVATIVES

FIELD OF THE INVENTION

5 The present invention relates to certain amido spirocyclic amide and sulfonamide compounds, pharmaceutical compositions comprising such compounds, and methods of treating cancer, including leukemias and solid tumors, inflammatory diseases, osteoporosis, atherosclerosis, irritable bowel syndrome, and other diseases and medical conditions, with such compounds and pharmaceutical compositions. The present invention also relates to certain amido spirocyclic amide and sulfonamide
10 compounds for use in inhibiting nicotinamide phosphoribosyltransferase ("NAMPT").

BACKGROUND OF THE INVENTION

 Nicotinamide adenine dinucleotide (NAD) plays a fundamental role in both
15 cellular energy metabolism and cellular signaling. NAD plays an important role in energy metabolism, as the pyridine ring in the NAD molecule readily accepts and donates electrons in hydride transfer reactions catalyzed by numerous dehydrogenases. The enzyme nicotinamide phosphoribosyltransferase (NAMPT, NMPRT, NMPRTase, or NAmPRTase, International nomenclature: E.C. 2.4.2.12),
20 promotes the condensation of nicotinamide with 5-phosphoribosyl-1-pyrophosphate to generate nicotinamide mononucleotide, which is a precursor in the biosynthesis of NAD.

 NAMPT is implicated in a variety of functions, including the promotion of
25 vascular smooth muscle cell maturation, inhibition of neutrophil apoptosis, activation of insulin receptors, development of T and B lymphocytes, and reduction of blood glucose. Thus, small molecule NAMPT inhibitors have potential uses as therapies in a variety of diseases or conditions, including cancers involving solid and liquid tumors, non-small cell lung cancer, leukemia, lymphoma, ovarian cancer, glioma, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate
30 cancer, skin cancer, rhino-gastric tumors, colorectal cancer, CNS cancer, bladder cancer, pancreatic cancer and Hodgkin's disease. NAMPT inhibitors also have

potential uses as therapies for diseases or conditions such as cancer, rheumatoid arthritis, diabetes, atherosclerosis, sepsis, or aging.

Rongvaux et al. have demonstrated that NAMPT is implicated in the regulation of cell viability during genotoxic or oxidative stress, and NAMPT inhibitors may therefore be useful as treatments for inflammation. Rongvaux, A., et al. *J. Immunol.* 2008, 181, 4685-4695. NAMPT may also have effects on the reaction of endothelial cells to high glucose levels, oxidative stress, and aging. Thus, NAMPT inhibitors may enable proliferating endothelial cells to resist the oxidative stress of aging and of high glucose, and to productively use excess glucose to support replicative longevity and angiogenic activity.

In particular, NAMPT inhibitors have been shown to interfere with NAD biosynthesis and to induce apoptotic cell death without any DNA damaging effects or primary effects on cellular energy metabolism, and thus have important anti-tumor effects. For example, the NAMPT inhibitor FK866 has these biochemical effects, and has also been shown to reduce NAD levels, induce a delay in tumor growth and enhance tumor radiosensitivity in a mouse mammary carcinoma model. *See, e.g.,* Hasmann M. and I. Schemainda, "FK866, a Highly Specific Noncompetitive Inhibitor of Nicotinamide Phosphoribosyltransferase, Represents a Novel Mechanism for Induction of Tumor Cell Apoptosis," *Cancer Res.* 2003, 63, 7436-7442; Dreys, J. et al., "Antiangiogenic potency of FK866/K22.175, a new inhibitor of intracellular NAD biosynthesis, in murine renal cell carcinoma," *Anticancer Res.* 2003, 23, 4853-4858.


More recently, another NAMPT inhibitor, CHS-828, has been shown to potently inhibit cell growth in a broad range of tumor cell lines. *See* Olesen, U.H. et al., "Anticancer agent CHS-828 inhibits cellular synthesis of NAD," *Biochem. Biophys. Res. Commun.* 2008, 367, 799-804; Ravaud, A. et al., "Phase I study and guanidine kinetics of CHS-828, a guanidine-containing compound, administered orally as a single dose every 3 weeks in solid tumors: an ECGS/EORTC study," *Eur. J. Cancer* 2005, 41, 702-707. Both FK866 and CHS-828 are currently in clinical trials as cancer treatments.

There remains a need for potent NAMPT inhibitors with desirable pharmaceutical properties. Certain amido spirocyclic amide and sulfonamide

derivatives have been found in the context of this invention to have NAMPT-modulating activity.

SUMMARY OF THE INVENTION

5 In one aspect, the invention is directed to compounds of Formula I:



wherein:

R is (a) an 8-, 9-, or 10-membered bicyclic heteroaryl comprising one heteroatom selected from N, S, and O, and one, two, or three additional N atoms, wherein said bicyclic heteroaryl is unsubstituted or is substituted with one or more substituents selected from the group consisting of deuterium, amino, alkylamino, dialkylamino, alkyl, halo, cyano, haloalkyl, hydroxy, hydroxyalkyl, and alkoxy, and wherein one or more N atoms of said bicyclic heteroaryl is optionally an N-oxide; or

15 (b) a five- or six-membered nitrogen-linked heterocycloalkyl ring fused to a phenyl or monocyclic five- or six-membered heteroaryl, wherein said phenyl or heteroaryl is unsubstituted or is substituted with one or more substituents selected from the group consisting of deuterium, amino, alkylamino, dialkylamino, alkyl, halo, cyano, haloalkyl, hydroxy, hydroxyalkyl, and alkoxy; and

20 R^1 is H, $-(C_{1-4}\text{alkylene})_{0-1}C(O)R^a$, $-(C_{1-4}\text{alkylene})_{0-1}CO_2R^a$, $-(C_{1-4}\text{alkylene})_{0-1}S(O)R^a$, $-(C_{1-4}\text{alkylene})_{0-1}SO_2R^a$, $-C(O)NH(R^a)$, $-C(O)N(R^a)_2$, or $-C(O)C(O)NH(R^a)$;

wherein each R^a is independently

25 (1) alkyl, unsubstituted or substituted with one or more R^m substituents, wherein each R^m is independently selected from the group consisting of deuterium, hydroxy, $-NR^bR^c$, alkoxy, cyano, halo, $-C(O)$ alkyl, $-CO_2$ alkyl, $-CONR^bR^c$, $-S(O)$ alkyl, $-SO_2$ alkyl, $-SO_2NR^bR^c$, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, phenoxy, and $-O$ -alkyl-OH;

wherein R^b is H or alkyl;

R^c is H, alkyl, alkoxyalkyl, haloalkyl, $-C(O)alkyl$, $-CO_2alkyl$,
 $-SO_2alkyl$, $-C(O)NH_2$, or $C(O)H$; and

each aryl, heteroaryl, cycloalkyl, and heterocycloalkyl group within

5 R^m is unsubstituted or substituted with one or more substituents
independently selected from the group consisting of alkyl,
haloalkyl, hydroxy, $-NR^bR^c$, alkoxy, haloalkoxy, cyano, halo, oxo,
 $-C(O)alkyl$, $-CO_2alkyl$, $-C(O)-heterocycloalkyl$, $-CONR^bR^c$,
 $-S(O)alkyl$, $-SO_2alkyl$, $-SO_2-haloalkyl$, $-SO_2NR^bR^c$, aryl, heteroaryl,
10 cycloalkyl, and heterocycloalkyl; or two substituents taken
together form a fused phenyl, heteroaryl, cycloalkyl, or
heterocycloalkyl ring;

wherein each alkyl or alkoxy is unsubstituted or substituted with

$-NR^bR^c$, heterocycloalkyl, heteroaryl, or $-C(O)alkyl$; and

15 each aryl, heteroaryl, cycloalkyl, and heterocycloalkyl is
unsubstituted or substituted with alkyl, halo, oxo, $-C(O)alkyl$;

(2) phenyl, cycloalkyl, heteroaryl, or heterocycloalkyl, each unsubstituted or
substituted with one or more R^s substituents;

wherein each R^s is independently selected from the group consisting of

20 alkyl, haloalkyl, hydroxy, $-NR^bR^c$, alkoxy, haloalkoxy, cyano, halo,
oxo, $-C(O)alkyl$, $-CO_2alkyl$, $-C(O)-heterocycloalkyl$, $-CONR^bR^c$,
 $-S(O)alkyl$, $-SO_2alkyl$, $-SO_2-haloalkyl$, $-SO_2NR^bR^c$, aryl, heteroaryl,
cycloalkyl, and heterocycloalkyl; or two R^s substituents taken together
form a fused phenyl, heteroaryl, cycloalkyl, or heterocycloalkyl ring;

25 wherein each alkyl or alkoxy is unsubstituted or substituted with

phenyl, $-NR^bR^c$, heterocycloalkyl, heteroaryl, or $-C(O)alkyl$; and

each aryl, heteroaryl, cycloalkyl, and heterocycloalkyl is unsubstituted
or substituted with alkyl, halo, $-CO_2alkyl$, or $-C(O)alkyl$; or

(3) $-NR^xR^y$,

30 where R^x is H or alkyl; and

R^y is H, alkyl, alkoxyalkyl, haloalkyl, $-C(O)alkyl$, $-CO_2alkyl$, or

-SO₂alkyl;

R² and R³ are each independently H or deuterium, and

n is 1 or 2;

and stereoisomers thereof, and pharmaceutically acceptable salts of such compounds

5 and stereoisomers.

In a further aspect, the invention relates to pharmaceutical compositions each comprising an effective amount of at least one compound of Formula I or a pharmaceutically acceptable salt of a compound of Formula I. Pharmaceutical compositions according to the invention may further comprise at least one

10 pharmaceutically acceptable excipient.

In another aspect, the invention is directed to a method of treating a subject suffering from a disease or medical condition mediated by NAMPT activity, comprising administering to the subject in need of such treatment an effective amount of at least one compound of Formula I or a pharmaceutically acceptable salt of a

15 compound of Formula I, or comprising administering to the subject in need of such treatment an effective amount of a pharmaceutical composition comprising an effective amount of at least one compound of Formula I or a pharmaceutically acceptable salt of a compound of Formula I.

An aspect of the present invention concerns the use of compound of Formula I for the preparation of a medicament used in the treatment, prevention, inhibition or

20 elimination of cancer.

An aspect of the present invention concerns the use of a compound of Formula I for the preparation of a medicament used in the treatment, prevention, inhibition or elimination of cancer, where the cancer can be selected from leukemia,

25 lymphoma, ovarian cancer, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, CNS cancer, bladder cancer, pancreatic cancer and Hodgkin's disease.

An aspect of the present invention concerns the use of a compound of Formula I for the preparation of a medicament used in the treatment, prevention,

30 inhibition or elimination of cancer, where the cancer can be selected from cancers with solid and liquid tumors, non-small cell lung cancer, leukemia, lymphoma,

ovarian cancer, glioma, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, rhino-gastric tumors, colorectal cancer, CNS cancer, bladder cancer, pancreatic cancer and Hodgkin's disease.

In another aspect, the compounds of Formula I and pharmaceutically acceptable salts thereof are useful as NAMPT modulators. Thus, the invention is directed to a method for modulating NAMPT activity, including when NAMPT is in a subject, comprising exposing NAMPT to an effective amount of at least one compound of Formula I or a pharmaceutically acceptable salt of a compound of Formula I.

In yet another aspect, the present invention is directed to methods of making compounds of Formula I and pharmaceutically acceptable salts thereof.

In certain embodiments of the compounds, pharmaceutical compositions, and methods of the invention, the compound of Formula I is a compound selected from those species described or exemplified in the detailed description below, or is a pharmaceutically acceptable salt of such a compound.

Additional embodiments, features, and advantages of the invention will be apparent from the following detailed description and through practice of the invention.

DETAILED DESCRIPTION AND PARTICULAR EMBODIMENTS

For the sake of brevity, the disclosures of the publications cited in this specification, including patents and patent applications, are herein incorporated by reference in their entirety.

Most chemical names were generated using IUPAC nomenclature herein. Some chemical names were generated using different nomenclatures or alternative or commercial names known in the art. In the case of conflict between names and structures, the structures prevail.

General Definitions

As used above, and throughout this disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings. If a definition is missing, the conventional definition as known to one skilled in the art

controls. If a definition provided herein conflicts or is different from a definition provided in any cited publication, the definition provided herein controls.

As used herein, the terms “including”, “containing” and “comprising” are used in their open, non-limiting sense.

5 As used herein, the singular forms “a”, “an” and “the” include plural referents unless the context clearly dictates otherwise.

To provide a more concise description, some of the quantitative expressions given herein are not qualified with the term “about”. It is understood that, whether the term “about” is used explicitly or not, every quantity given herein is meant to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including equivalents and approximations due to the experimental and/or measurement conditions for such given value. Whenever a yield is given as a percentage, such yield refers to a mass of the entity for which the yield is given with respect to the maximum amount of the same entity that could be obtained under the particular stoichiometric conditions. Concentrations that are given as percentages refer to mass ratios, unless indicated differently.

10
15

Definitions

20 As used herein, “alkyl” refers to a saturated, straight- or branched-chain hydrocarbon group having from 1 to 10 carbon atoms. Representative alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, and the like, and longer alkyl groups, such as heptyl, octyl, and the like. As used herein, “lower alkyl” means an alkyl having from 1 to 6 carbon atoms.

25

30 The term “alkoxy” as used herein includes -O-(alkyl), wherein alkyl is defined above.

As used herein, "alkoxyalkyl" means -(alkylenyl)-O-(alkyl), wherein each "alkyl" is independently an alkyl group defined above.

The term "amino" as used herein refers to an $-NH_2$ group.

The term "alkylamino" as used herein denotes an amino group as defined
5 above wherein one hydrogen atom of the amino group is replaced by an alkyl group as defined herein. Aminoalkyl groups can be defined by the following general formula $-NH$ -alkyl. This general formula includes groups of the following general formulae: $-NH-C_1-C_{10}$ -alkyl and $-NH-C_1-C_6$ -alkyl. Examples of aminoalkyl groups include, but are not limited to aminomethyl, aminoethyl, aminopropyl, aminobutyl.

10 The term "dialkylamino" as used herein denotes an amino group as defined above wherein two hydrogen atoms of the amino group are replaced by alkyl groups as defined herein. Diaminoalkyl groups can be defined by the following general formula $-N(alkyl)_2$, wherein the alkyl groups can be the same or can be different and can be selected from alkyls as defined herein, for example C_1-C_{10} -alkyl or C_1-C_6 -
15 alkyl.

"Aryl" means a mono-, bi-, or tricyclic aromatic group, wherein all rings of the group are aromatic. For bi- or tricyclic systems, the individual aromatic rings are fused to one another. Exemplary aryl groups include, but are not limited to, phenyl, naphthalene, and anthracene.

20 "Aryloxy" as used herein refers to an $-O$ -(aryl) group, wherein aryl is defined as above.

"Arylalkyl" as used herein refers to an $-(alkylenyl)$ -(aryl) group, wherein alkylenyl and aryl are as defined above. Non-limiting examples of arylalkyls comprise a lower alkyl group. Non-limiting examples of suitable arylalkyl groups
25 include benzyl, 2-phenethyl, and naphthalenylmethyl.

"Arylalkoxy" as used herein refers to an $-O$ -(alkylenyl)-aryl group wherein alkylenyl and aryl are as defined above.

The term "cyano" as used herein means a substituent having a carbon atom joined to a nitrogen atom by a triple bond.

30 The term "deuterium" as used herein means a stable isotope of hydrogen having one proton and one neutron.

The term "halogen" as used herein refers to fluorine, chlorine, bromine, or iodine. The term "halo" represents chloro, fluoro, bromo, or iodo. Halo can also denote chloro, fluoro, or bromo.

The term "haloalkyl" denotes an alkyl group as defined above wherein one or more, for example one, two, or three of the hydrogen atoms of the alkyl group are replaced by a halogen atom, for example fluoro, bromo, or chloro, in particular fluoro. Examples of haloalkyl include, but are not limited to, monofluoro-, difluoro-, or trifluoro-methyl, -ethyl or -propyl, for example, 3,3,3-trifluoropropyl, 2-fluoroethyl, 2,2,2-trifluoroethyl, fluoromethyl, difluoromethyl, or trifluoromethyl, or bromoethyl or chloroethyl. Similarly, the term "fluoroalkyl" refers to an alkyl group as defined above substituted with one or more, for example one, two, or three fluorine atoms.


The term "haloalkoxy" as used herein refers to an -O-(haloalkyl) group wherein haloalkyl is defined as above. Exemplary haloalkoxy groups are bromoethoxy, chloroethoxy, trifluoromethoxy and 2,2,2-trifluoroethoxy.

The term "hydroxy" means an -OH group.

The term "hydroxyalkyl" denotes an alkyl group that is substituted by at least one hydroxy group, for example, one, two or three hydroxy group(s). The alkyl portion of the hydroxyalkyl group provides the connection point to the remainder of a molecule. Examples of hydroxyalkyl groups include, but are not limited to, hydroxymethyl, hydroxyethyl, 1-hydroxypropyl, 2-hydroxyisopropyl, 1,4-dihydroxybutyl, and the like.

The term "oxo" means an =O group and may be attached to a carbon atom or a sulfur atom. The term "N-oxide" refers to the oxidized form of a nitrogen atom.

As used herein, the term "cycloalkyl" refers to a saturated or partially saturated, monocyclic, fused polycyclic, bridged polycyclic, or spiro polycyclic carbocycle having from 3 to 15 ring carbon atoms. A non limiting category of cycloalkyl groups are saturated or partially saturated, monocyclic carbocycles having from 3 to 6 carbon atoms. Illustrative examples of cycloalkyl groups include, but are not limited to, the following moieties:



ring systems that must be charged to be aromatic, such as pyrylium. Certain suitable 5-membered heteroaryl rings (as a monocyclic heteroaryl or as part of a polycyclic heteroaryl) have one oxygen, sulfur, or nitrogen atom, or one nitrogen plus one oxygen or sulfur, or 2, 3, or 4 nitrogen atoms. Certain suitable 6-membered

5 heteroaryl rings (as a monocyclic heteroaryl or as part of a polycyclic heteroaryl) have 1, 2, or 3 nitrogen atoms. Examples of heteroaryl groups include, but are not limited to, pyridinyl, imidazolyl, imidazopyridinyl, pyrimidinyl, pyrazolyl, triazolyl, pyrazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, thiazolyl, oxazolyl, isothiazolyl, pyrrolyl, quinolinyl, isoquinolinyl, indolyl, benzimidazolyl, benzofuranyl, cinnolinyl,


10 indazolyl, indoliziny, phthalazinyl, pyridazinyl, triazinyl, isoindolyl, pteridinyl, purinyl, oxadiazolyl, triazolyl, thiadiazolyl, furazanyl, benzofurazanyl, benzothiophenyl, benzothiazolyl, benzoxazolyl, quinazolinyl, quinoxaliny, naphthyridinyl, and furopyridinyl.


The term “bicyclic heteroaryl” refers to a heteroaryl as defined above, having

15 two constituent aromatic rings, wherein the two rings are fused to one another and at least one of the rings is a heteroaryl as defined above. Bicyclic heteroaryls include bicyclic heteroaryl groups comprising 1, 2, 3, or 4 heteroatom ring members and are unsubstituted or substituted with one or more substituents selected from the group consisting of amino and halo; and wherein one or more N atoms of said heteroaryl is


20 optionally an N-oxide. Bicyclic heteroaryls also include 8-, 9-, or 10-membered bicyclic heteroaryl groups. Bicyclic heteroaryls also include 8-, 9-, or 10-membered bicyclic heteroaryl groups that have 1, 2, 3 or 4 heteroatom ring members and that are unsubstituted or substituted with one or more substituents selected from the group consisting of amino and halo; and wherein one or more N atoms of said heteroaryl is

25 optionally an N-oxide. Illustrative examples of bicyclic heteroaryls include, but are not limited to:





The term “five- or six-membered nitrogen-linked heterocycloalkyl ring fused to a phenyl or monocyclic heteroaryl, wherein said phenyl or heteroaryl is unsubstituted or is substituted with amino” include, but are not limited to, the following groups:



Those skilled in the art will recognize that the species of heteroaryl, cycloalkyl, and heterocycloalkyl groups listed or illustrated above are not exhaustive, and that additional species within the scope of these defined terms may also be selected.

As used herein, the term “substituted” means that the specified group or moiety bears one or more suitable substituents. As used herein, the term “unsubstituted” means that the specified group bears no substituents. As used herein, the term “optionally substituted” means that the specified group is unsubstituted or substituted by the specified number of substituents. Where the term “substituted” is used to describe a structural system, the substitution is meant to occur at any valency-allowed position on the system.

As used herein, the expression “one or more substituents” denotes one to the maximum possible number of substitution(s) that can occur at any valency-allowed position on the system. In a certain embodiment, one or more substituents means 1, 2, 3, 4, or 5 substituents. In another embodiment, one or more substituent means 1, 2, or 3 substituents. Such substituents may be the same or different from one another.

Any atom that is represented herein with an unsatisfied valence is assumed to have the sufficient number of hydrogen atoms to satisfy the atom's valence.

When any variable (*e.g.*, alkyl, alkylenyl, heteroaryl, R¹, R², Rⁿ, etc.) appears in more than one place in any formula or description provided herein, the definition of that variable on each occurrence is independent of its definition at every other occurrence.

Numerical ranges, as used herein, are intended to include sequential whole numbers. For example, a range expressed as "from 0 to 4" or "0-4" includes 0, 1, 2, 3 and 4.

When a multifunctional moiety is shown, the point of attachment to the core is indicated by a line or hyphen. For example, aryloxy- refers to a moiety in which an oxygen atom is the point of attachment to the core molecule while aryl is attached to the oxygen atom.

As used herein, the term "subject" encompasses mammals and non-mammals. Examples of mammals include, but are not limited to, any member of the Mammalian class: humans; non-human primates such as chimpanzees, and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, swine; domestic animals such as rabbits, dogs, and cats; and laboratory animals including rodents, such as rats, mice and guinea pigs, and the like. Examples of non-mammals include, but are not limited to, birds, fish and the like. In one embodiment of the present invention, the mammal is a human.

"Patient" includes both human and animals.

The term "inhibitor" refers to a molecule such as a compound, a drug, an enzyme activator, or a hormone that blocks or otherwise interferes with a particular biologic activity.

The term "modulator" refers to a molecule, such as a compound of the present invention, that increases or decreases, or otherwise affects the activity of a given enzyme or protein.

The terms "effective amount" or "therapeutically effective amount" refer to a sufficient amount of the agent to provide the desired biological result. That result can be reduction and/or alleviation of the signs, symptoms, or causes of a disease or

medical condition, or any other desired alteration of a biological system. For example, an “effective amount” for therapeutic use is the amount of a compound, or of a composition comprising the compound, that is required to provide a clinically relevant change in a disease state, symptom, or medical condition. An appropriate
5 “effective” amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation. Thus, the expression “effective amount” generally refers to the quantity for which the active substance has a therapeutically desired effect.

As used herein, the terms “treat” or “treatment” encompass both
10 “preventative” and “curative” treatment. “Preventative” treatment is meant to indicate a postponement of development of a disease, a symptom of a disease, or medical condition, suppressing symptoms that may appear, or reducing the risk of developing or recurrence of a disease or symptom. “Curative” treatment includes reducing the severity of or suppressing the worsening of an existing disease, symptom,
15 or condition. Thus, treatment includes ameliorating or preventing the worsening of existing disease symptoms, preventing additional symptoms from occurring, ameliorating or preventing the underlying metabolic causes of symptoms, inhibiting the disorder or disease, e.g., arresting the development of the disorder or disease, relieving the disorder or disease, causing regression of the disorder or disease,
20 relieving a condition caused by the disease or disorder, or stopping the symptoms of the disease or disorder.

Additional Chemical Descriptions

Any formula given herein is intended to represent compounds having
25 structures depicted by the structural formula as well as certain variations or forms. For example, compounds of any formula given herein may have asymmetric or chiral centers and therefore exist in different stereoisomeric forms. All stereoisomers, including optical isomers, enantiomers, and diastereomers, of the compounds of the general formula, and mixtures thereof, are considered to fall within the scope of the
30 formula. Furthermore, certain structures may exist as geometric isomers (i.e., *cis* and *trans* isomers), as tautomers, or as atropisomers. All such isomeric forms, and

mixtures thereof, are contemplated herein as part of the present invention. Thus, any formula given herein is intended to represent a racemate, one or more enantiomeric forms, one or more diastereomeric forms, one or more tautomeric or atropisomeric forms, and mixtures thereof.

5 Diastereomeric mixtures may be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers may be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active
10 compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride, or formation of a mixture of diastereomeric salts), separating the diastereomers and converting (e.g., hydrolyzing or de-salting) the individual diastereomers to the corresponding pure enantiomers. Enantiomers may also be separated by use of chiral HPLC column. The chiral centers of compounds of the present invention may be
15 designated as "R" or "S" as defined by the IUPAC 1974 Recommendations.

The compounds of the invention can form pharmaceutically acceptable salts, which are also within the scope of this invention. A "pharmaceutically acceptable salt" refers to a salt of a free acid or base of a compound of Formula I that is non-toxic, is physiologically tolerable, is compatible with the pharmaceutical composition
20 in which it is formulated, and is otherwise suitable for formulation and/or administration to a subject. Reference to a compound herein is understood to include reference to a pharmaceutically acceptable salt of said compound unless otherwise indicated.

Compound salts include acidic salts formed with inorganic and/or organic
25 acids, as well as basic salts formed with inorganic and/or organic bases. In addition, where a given compound contains both a basic moiety, such as, but not limited to, a pyridine or imidazole, and an acidic moiety, such as, but not limited to, a carboxylic acid, one of skill in the art will recognize that the compound may exist as a zwitterion ("inner salt"); such salts are included within the term "salt" as used herein. Salts of
30 the compounds of the invention may be prepared, for example, by reacting a compound with an amount of a suitable acid or base, such as an equivalent amount, in

a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

Exemplary salts include, but are not limited, to sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, 5 ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate ("mesylate"), ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, and pamoate (i.e., 1,1'-methylene-bis(2-hydroxy-3-naphthoate)) salts. A pharmaceutically acceptable salt 10 may involve the inclusion of another molecule such as an acetate ion, a succinate ion or other counterion. The counterion may be any organic or inorganic moiety that stabilizes the charge on the parent compound. Furthermore, a pharmaceutically acceptable salt may have more than one charged atom in its structure. Instances where multiple charged atoms are part of the pharmaceutically acceptable salt can 15 have multiple counterions. Hence, a pharmaceutically acceptable salt can have one or more charged atoms and/or one or more counter ion.

Exemplary acid addition salts include acetates, ascorbates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, fumarates, hydrochlorides, hydrobromides, hydroiodides, lactates, 20 maleates, methanesulfonates, naphthalenesulfonates, nitrates, oxalates, phosphates, propionates, salicylates, succinates, sulfates, tartarates, thiocyanates, toluenesulfonates (also known as tosylates,) and the like.

Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and 25 magnesium salts, salts with organic bases (for example, organic amines) such as dicyclohexylamines, *t*-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quarternized with agents such as lower alkyl halides (e.g. methyl, ethyl, and butyl chlorides, bromides and iodides), dialkyl sulfates (e.g. dimethyl, diethyl, and dibutyl sulfates), long chain 30 halides (e.g. decyl, lauryl, and stearyl chlorides, bromides and iodides), aralkyl halides (e.g. benzyl and phenethyl bromides), and others.

Additionally, acids and bases which are generally considered suitable for the formation of pharmaceutically useful salts from pharmaceutical compounds are discussed, for example, by P. Stahl et al, Camille G. (eds.) Handbook of Pharmaceutical Salts. Properties, Selection and Use. (2002) Zurich: Wiley-VCH; S. Berge et al, Journal of Pharmaceutical Sciences (1977) 66(1) 1-19; P. Gould, International J. of Pharmaceutics (1986) 33 201-217; Anderson et al, The Practice of Medicinal Chemistry (1996), Academic Press, New York; and in The Orange Book (Food & Drug Administration, MD, available from FDA). These disclosures are incorporated herein by reference thereto.

10 Additionally, any compound described herein is intended to refer also to any unsolvated form, or a hydrate, solvate, or polymorph of such a compound, and mixtures thereof, even if such forms are not listed explicitly. "Solvate" means a physical association of a compound of the invention with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent
15 bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of a crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Suitable solvates include those formed with pharmaceutically acceptable solvents such as water, ethanol, and the like. In some embodiments, the
20 solvent is water and the solvates are hydrates.

One or more compounds of the invention may optionally be converted to a solvate. Methods for the preparation of solvates are generally known. Thus, for example, M. Caira et al., J. Pharmaceutical Sci., 93(3), 601-611 (2004), describes the preparation of the solvates of the antifungal fluconazole in ethyl acetate as well as
25 from water. Similar preparations of solvates, hemisolvate, hydrates, and the like are described by E. C. van Tonder et al, AAPS PharmSciTech., 5(1), article 12 (2004); and A. L. Bingham et al, Chem. Commun., 603-604 (2001). A typical, non-limiting process involves dissolving the inventive compound in a suitable amounts of the solvent (organic solvent or water or a mixture thereof) at a higher than ambient
30 temperature, and cooling the solution at a rate sufficient to form crystals which are then isolated by standard methods. Analytical techniques such as, for example,

infrared spectroscopy, show the presence of the solvent (or water) in the crystals as a solvate (or hydrate).

The invention also relates to pharmaceutically acceptable prodrugs of the compounds of Formula I, and treatment methods employing such pharmaceutically acceptable prodrugs. The term "prodrug" means a precursor of a designated
5 compound that, following administration to a subject, yields the compound *in vivo* via a chemical or physiological process such as solvolysis or enzymatic cleavage, or under physiological conditions (e.g., a prodrug on being brought to physiological pH is converted to the compound of Formula I). A "pharmaceutically acceptable
10 prodrug" is a prodrug that is non-toxic, biologically tolerable, and otherwise suitable for formulation and/or administration to the subject. Illustrative procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs", ed. H. Bundgaard, Elsevier, 1985.

Examples of prodrugs include pharmaceutically acceptable esters of the
15 compounds of the invention, which are also considered to be part of the invention. Pharmaceutically acceptable esters of the present compounds include the following groups: (1) carboxylic acid esters obtained by esterification of the hydroxy groups, in which the non-carbonyl moiety of the carboxylic acid portion of the ester grouping is selected from straight or branched chain alkyl (for example, acetyl, n-propyl, t-butyl,
20 or n-butyl), alkoxyalkyl (for example, methoxymethyl), aralkyl (for example, benzyl), aryloxyalkyl (for example, phenoxymethyl), aryl (for example, phenyl optionally substituted with, for example, halogen, C₁₋₄alkyl, or C₁₋₄alkoxy or amino); (2) sulfonate esters, such as alkyl- or aralkylsulfonyl (for example, methanesulfonyl); (3) amino acid esters (for example, L-valyl or L-isoleucyl); (4) phosphonate esters and (5)
25 mono-, di- or triphosphate esters. The phosphate esters may be further esterified by, for example, a C₁₋₂₀ alcohol or reactive derivative thereof, or by a 2,3-di(C₆₋₂₄)acyl glycerol. Additional discussion of prodrugs is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems (1987) 14 of the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design, (1987) Edward B. Roche, ed., American
30 Pharmaceutical Association and Pergamon Press.

- For example, if a compound of Formula I contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as, for example, (C₁-C₈)alkyl, (C₂-C₁₂)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 5 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxy-carbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxy-carbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxy-carbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxy-carbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxy-carbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁-C₂)alkylamino(C₂-C₃)alkyl (such as β-dimethylaminoethyl), carbamoyl-(C₁-C₂)alkyl, N,N-di(C₁-C₂)alkylcarbamoyl-(C₁-C₂)alkyl and piperidino-, pyrrolidino- or morpholine (C₂-C₃)alkyl, and the like.
- 10 Similarly, if a compound of Formula I contains an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as, for example, (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl, (C₁-C₆)alkoxy-carbonyloxymethyl, N-(C₁-C₆)alkoxy-carbonylaminomethyl, succinoyl, (C₁-C₆)alkanoyl, α-amino(C₁-C₄)alkanyl, arylacyl and α-aminoacyl, or α-aminoacyl-α-aminoacyl, where each α-aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate), and the like.
- 15 If a compound of Formula I incorporates an amine functional group, a prodrug can be formed by the replacement of a hydrogen atom in the amine group with a group such as, for example, R^{''}-carbonyl, R^{''}O-carbonyl, NR^{''}R[']-carbonyl where R^{''} and R['] are each independently (C₁-C₁₀)alkyl, (C₃-C₇) cycloalkyl, benzyl, or R^{''}-carbonyl is a natural α-aminoacyl or natural α-aminoacyl, -C(OH)C(O)OY¹ wherein 20 Y¹ is H, (C₁-C₆)alkyl or benzyl, -C(OY²)Y³ wherein Y² is (C₁-C₄) alkyl and Y³ is (C₁-C₆)alkyl, carboxy(C₁-C₆)alkyl, amino(C₁-C₄)alkyl or mono-N- or di-N,N-(C₁-

C₆)alkylaminoalkyl, -C(Y⁴)Y⁵ wherein Y⁴ is H or methyl and Y⁵ is mono-N- or di-N,N-(C₁-C₆)alkylamino morpholino, piperidin-1-yl or pyrrolidin-1-yl, and the like.

The present invention also relates to pharmaceutically active metabolites of compounds of Formula I, and uses of such metabolites in the methods of the
5 invention. A “pharmaceutically active metabolite” means a pharmacologically active product of metabolism in the body of a compound of Formula I or salt thereof. Prodrugs and active metabolites of a compound may be determined using routine techniques known or available in the art. See, e.g., Bertolini et al., *J. Med. Chem.* 1997, *40*, 2011-2016; Shan et al., *J. Pharm. Sci.* 1997, *86* (7), 765-767; Bagshawe,
10 *Drug Dev. Res.* 1995, *34*, 220-230; Bodor, *Adv. Drug Res.* 1984, *13*, 255-331; Bundgaard, *Design of Prodrugs* (Elsevier Press, 1985); and Larsen, *Design and Application of Prodrugs, Drug Design and Development* (Krogsgaard-Larsen et al., eds., Harwood Academic Publishers, 1991).


Any formula given herein is also intended to represent unlabeled forms as
15 well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such
20 as ²H, ³H, ¹¹C, ¹³C, ¹⁴C, ¹⁵N, ¹⁸O, ¹⁷O, ³¹P, ³²P, ³⁵S, ¹⁸F, ³⁶Cl, and ¹²⁵I, respectively. Such isotopically labelled compounds are useful in metabolic studies (for example with ¹⁴C), reaction kinetic studies (with, for example ²H or ³H), detection or imaging techniques [such as positron emission tomography (PET) or single-photon emission
25 computed tomography (SPECT)] including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ¹⁸F or ¹¹C labeled compound may be particularly suitable for PET or SPECT studies. Further, substitution with heavier isotopes such as deuterium (i.e., ²H) may afford certain therapeutic advantages resulting from greater metabolic stability, for example increased *in vivo*
30 half-life or reduced dosage requirements. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described

below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

The use of the terms “salt”, “solvate”, “polymorph”, “prodrug”, and the like, with respect to the compounds described herein is intended to apply equally to the salt, solvate, polymorph, and prodrug forms of enantiomers, stereoisomers, rotamers, tautomers, atropisomers, and racemates of the inventive compounds.


Compounds of the Invention

In some embodiments of Formula I, compounds of the invention have the following Formula I-a:




wherein R, R¹, R², and R³ are as defined herein for Formula I.

In some embodiments, R is an unsubstituted or substituted bicyclic heteroaryl as defined for Formula I. In some embodiments, the bicyclic heteroaryl has 1, 2, or 3 nitrogen ring atoms. In other embodiments, the bicyclic heteroaryl is a 9- or 10-membered bicyclic heteroaryl, unsubstituted or substituted as described for Formula I. In other embodiments, the bicyclic heteroaryl is an 8- or 9-membered heteroaryl, unsubstituted or substituted as described for Formula I. In other embodiments, R is a bicyclic heteroaryl selected from the group consisting of:




each unsubstituted or substituted as described for Formula I. In further embodiments, R is selected from the group consisting of:




each unsubstituted or substituted as described for Formula I. In further embodiments,

5 R is selected from the group consisting of:




In further embodiments, R is



10 In other embodiments, R is a five- or six-membered nitrogen-linked heterocycloalkyl ring fused to an unsubstituted or substituted phenyl or monocyclic heteroaryl, for examples a 6 membered heteroaryl, as defined in Formula I. In further

embodiments, R is



15 In other embodiments, R is substituted with one or more substituents selected from the group consisting of amino and halo.

In some embodiments, R¹ is H. In other embodiments, R¹ is -C(O)R^a, -CO₂R^a, -S(O)R^a, or -SO₂R^a. In further embodiments, R¹ is -C(O)R^a, -CO₂R^a, or

$-\text{SO}_2\text{R}^a$. In other embodiments, R^1 is $-\text{C}(\text{O})\text{NH}(\text{R}^a)$ or $-\text{C}(\text{O})\text{C}(\text{O})\text{NH}(\text{R}^a)$. In other embodiments, R^a is alkyl, unsubstituted or substituted as described for Formula I. In other embodiments, R^a is $-\text{NR}^x\text{R}^y$. In further embodiments, R^a is $-\text{CH}_2\text{C}(\text{O})\text{R}^a$ or $-\text{CH}_2\text{SO}_2\text{R}^a$.

5 In some embodiments, R^a is methyl, ethyl, propyl, isopropyl, tert-butyl, isobutyl, isopentyl, phenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, isoindolinyl, azetidyl, oxetanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, or tetrahydrothiophenyl, each unsubstituted or substituted as
10 described for Formula I. In other embodiments, R^a is a bicyclic heteroaryl, unsubstituted or substituted with one or more substituents selected from the group consisting of alkyl, $-\text{CO}_2$ -tert-butyl, oxo, and halo. In other embodiments, R^a is alkyl, phenyl, benzyl, cycloalkyl, or heterocycloalkyl, each unsubstituted or substituted as described for Formula I. In some embodiments, R^a is methyl, ethyl, propyl, isopropyl,
15 butyl, isobutyl, sec-butyl, tert-butyl, 2,2-dimethylpropyl, hydroxyethyl, aminoethyl, cyanoethyl, ethoxy, tert-butoxy, phenyl, benzyl, cyclobutyl, cyclopentyl, cyclohexyl, oxetanyl, pyrrolidinyl, piperidinyl, tetrahydrofuranyl, or tetrahydropyranyl, each unsubstituted or substituted as described for Formula I. In other embodiments, R^a is
20 tert-butoxy, 2,2-dimethylpropyl, benzyl, cyclohexyl, tetrahydropyranyl, phenyl, methyl, ethyl, or isopropyl.

In some embodiments, R^a is phenyl, cycloalkyl, heteroaryl, or heterocycloalkyl, each unsubstituted or substituted as described for Formula I. In other embodiments, R^a is phenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl,
25 pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, isoindolinyl, azetidyl, oxetanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, or tetrahydrothiophenyl, each unsubstituted or substituted as in Formula I. In other embodiments, R^a is phenyl, cycloalkyl, heteroaryl, or
30 heterocycloalkyl, each unsubstituted or substituted with one or more substituents selected from the group consisting of fluoro, oxo, methyl, $-\text{CONH}_2$, acetyl, -

SO₂methyl, -C(O)-isopropyl, pyridazinyl, triazolyl, dimethylaminomethyl, cyano, methyl-triazolyl-methoxy, trifluoromethoxy, pyrrolidinylmethyl, acetylamino, tetrazolylmethyl, methyl-tetrazolyl-methyl, methyl-imidazolyl-methyl, -NHSO₂methyl, 1,1-dioxothiomorpholinyl, 4-methyl-piperazinylmethyl, -NHCONH₂,
 5 -SO₂CF₃, morpholinylmethyl, imidazolyl, -SO₂NH₂, methylpiperidinyl, methyl-piperazinyl, -C(O)(4-methyl-piperazinyl), morpholinyl, trifluoromethyl, cyclopropyl, ethyl, isoxazolyl, tetrazolyl, isopropyl, phenyl, fluoro-phenyl, tert-butyl, benzyl, N-methylpyrrolidinyl, N-acetyl-pyrrolidinyl, isobutyl, propyl, methylpyrazolyl, trifluoroethyl, pyrimidinyl, oxo, acetyl, cyano, -CO₂-tert-butyl, and amino.

10 In other embodiments, R¹ is alkyl, unsubstituted or substituted with one or more substituents selected from the group consisting of fluoro, tert-butoxy, -C(O)NMe₂, -NHCHO, methoxy, phenoxy, cyano, acetyl, hydroxy, -OCH₂C(CH₃)=OH, -NH(acetyl), and -N(Me)(acetyl).

In other embodiments, R¹ is -SO₂R^a, where R^a is methyl, ethyl, phenyl,
 15 benzyl, or 2,2-dimethylpropyl.








In other embodiments, R¹ is -C(O)NHR^a, wherein R^a is methyl, ethyl, propyl, isopropyl, tertobutyl, cyclohexyl, -CH₂-cyclohexyl, oxetanyl, or methyloxetanyl, or R^a is a phenyl or benzyl group, each optionally substituted with one or more substituents selected from the group consisting of cyano, methyl, fluoro, methoxy,
 20 and chloro.








In some embodiments, one of R² and R³ is deuterium and the other is H. In other embodiments, both R² and R³ are H.

In some embodiments, each alkyl or alkylene described above is independently a C₁₋₁₀alkyl. In other embodiments, each alkyl or alkylene in Formula
 25 I is independently a C₁₋₆alkyl. In still other embodiments, each alkyl or alkylene in Formula I is independently a C₁₋₄alkyl.

In certain embodiments, the compound of Formula I is chosen from the following table:






Ex.	Structure	Chemical Name
1		tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
2		tert-butyl 2-[(thieno[2,3-c]pyridine-2-carbonylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
3		tert-butyl 2-[(imidazo[1,2-a]pyrimidine-6-carbonylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
4		tert-butyl 2-[(furo[2,3-c]pyridine-2-carbonylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
5		tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carbonylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
6		N-[[8-(3,3-dimethylbutanoyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
7		N-[[8-(2-phenylacetyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

8		tert-butyl 2-[(1H-pyrrolo[3,2-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
9		N-[[8-(cyclohexanecarbonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
10		N-[[8-(tetrahydropyran-4-carbonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
11		N-[(8-benzoyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
12		N-[[8-(2-methylpropanoyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
13		N-[[8-(benzenesulfonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
14		N-[(8-ethylsulfonyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

15		tert-butyl 2-[(1H-pyrazolo[3,4-b]pyridine-5-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate
16		N-[(8-acetyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
17		N-[(8-methylsulfonyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
18		N-(8-azaspiro[2.5]octan-2-ylmethyl)-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
19		N-(8-azaspiro[2.5]octan-2-ylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide
20		N-[(8-propanoyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide or
21		tert-butyl 2-((imidazo[1,2-a]pyridine-6-carboxamido)methyl)-7-azaspiro[3.5]nonane-7-carboxylate

or a pharmaceutically acceptable salt thereof, or a stereoisomer or a pharmaceutically acceptable salt of said stereoisomer.

In certain other embodiments, the compound of Formula I is chosen from the following table:







Ex.	Structure	Chemical Name
22		tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
23		tert-butyl 2-[(furo[2,3-c]pyridine-2-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
24		N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
25		tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
26		N-[[6-(2-imidazol-1-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide







<p>27</p>		<p>N-[[6-(4,4,4-trifluorobutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>28</p>		<p>N-[[6-[2-(3-methylsulfonylphenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>29</p>		<p>N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>30</p>		<p>N-[[6-[2-(4-methylpiperazin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>31</p>		<p>N-[[6-[2-(3-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>32</p>		<p>N-[[6-(2-cyclopropylacetyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>33</p>		<p>N-[[6-(2-pyrrolidin-1-ylacetyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>34</p>		<p>N-[[6-[2-(3-methylisoxazol-5-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>35</p>		<p>N-[[6-[2-(1-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>




<p>36</p>		<p>N-[[6-(2-tetrahydropyran-4-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>37</p>		<p>N-[[6-(2-tetrahydrofuran-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>38</p>		<p>N-[[6-(2-tert-butoxyacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




39		N-[[6-[2-(3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
40		N-[[6-[2-[1-(2,2,2-trifluoroethyl)-4-piperidyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
41		tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 1)
42		tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 2)
43		tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 1)
44		tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 2)




45		N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
46		N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
47		N-[[6-(2-phenylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
48		tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 1)
49		tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 2)
50		N-[[6-(2-pyrazin-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>51</p>		<p>N-[[6-(2-morpholinoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>52</p>		<p>N-[[6-(2-tetrahydropyran-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>53</p>		<p>N-[[6-(2-pyrrolidin-1-ylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




<p>54</p>		<p>N-[[6-[2-(tert-butylamino)-2-oxo-acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>55</p>		<p>N-[[6-[4-(dimethylamino)-4-oxo-butanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>56</p>		<p>N-[[6-(1-methyl-5-oxo-pyrrolidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




57		N-[[6-[2-(2-oxopyrrolidin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
58		N-[[6-(2,2-difluorocyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
59		N-[[6-(tetrahydrofuran-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

60	 <p>The structure shows a furo[2,3-c]pyridine ring system with a methyl group at the 2-position and a carboxamide group at the 3-position. The methyl group is substituted with a 6-azaspiro[2.5]octan-2-yl group. The nitrogen of the azaspiro ring is substituted with a tetrahydropyran-4-carbonyl group.</p>	N-[[6-(tetrahydropyran-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
61	 <p>The structure is similar to compound 60, but the tetrahydropyran ring is replaced by a tetrahydrofuran ring, and the carbonyl group is at the 2-position of the tetrahydrofuran ring.</p>	N-[[6-(tetrahydrofuran-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
62	 <p>The structure is similar to compound 60, but the tetrahydropyran ring is replaced by a 2-formamidoacetyl group attached to the nitrogen of the azaspiro ring.</p>	N-[[6-(2-formamidoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide




<p>63</p>		<p>N-[[6-[3-(3,5-dimethylpyrazol-1-yl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>64</p>		<p>N-[[6-(1-carbamoylpyrrolidin-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>65</p>		<p>N-[[6-(3-morpholinopropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>66</p>		<p>N-[[6-(1,1-dioxathiolane-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>67</p>		<p>N-[[6-[2-(3,5-dimethylpyrazol-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>68</p>		<p>N-[[6-(1-acetylpyrrolidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>



69		<p>N-[[6-(2-methylsulfonylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
70		<p>N-[[6-[3-(1,2,4-triazol-4-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
71		<p>N-[[6-[4-((dimethylamino)methyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

72		<p>N-[[6-(4-acetamidobenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
73		<p>N-[[6-(1-methylsulfonylpiperidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
74		<p>N-[[6-(1-methylsulfonylpiperidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>75</p>		<p>N-[[6-(2-morpholinopropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>76</p>		<p>N-[[6-(1-methylsulfonylpyrrolidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>77</p>		<p>N-[[6-[2-(3-oxoisindolin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




78		N-[[6-(1-carbamoylpiperidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
79		N-[[6-(1-carbamoylpiperidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
80		N-[[6-[2-(1,1-dioxothiolan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide




<p>81</p>		<p>N-[[6-[2-(2,4-dioxypyrimidin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>82</p>		<p>N-[[6-(1-acetylpiperidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>83</p>		<p>N-[[6-[4-(tetrazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




84		N-[[6-[4-(methanesulfonamido)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
85		N-[[6-[4-(1,1-dioxo-1,4-thiazinan-4-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide




<p>86</p>		<p>N-[[6-[4-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>87</p>		<p>N-[[6-[2-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>88</p>		<p>N-[[6-[3-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




<p>89</p>		<p>N-[[6-[3-(4-methylpiperazin-1-yl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>90</p>		<p>N-[[6-(1-isopropyl-5-oxopyrrolidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>91</p>		<p>N-[[6-[2-(3-oxopiperazin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>92</p>		<p>N-[[6-(3-methylsulfonylbenzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>93</p>		<p>N-[[6-(4-methylsulfonylbenzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>94</p>		<p>N-[[6-(4-ureidobenzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>

95		<p>N-[[6-[2-(4-acetylpiperazin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
96		<p>N-[[6-[4-[(5-methyltetrazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
97		<p>N-[[6-(pyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>98</p>		<p>N-[[6-(pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>99</p>		<p>N-[[6-(oxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>100</p>		<p>N-[[6-(1H-1,2,4-triazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>101</p>	 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring system. The 6-position of the spiro system is substituted with a 1H-imidazole-4-carbonyl group.</p>	<p>N-[[6-(1H-imidazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>102</p>	 <p>The structure is similar to 101, but the imidazole ring is replaced by a pyrimidine ring.</p>	<p>N-[[6-(pyrimidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>103</p>	 <p>The structure is similar to 101, but the imidazole ring is replaced by a pyridine ring.</p>	<p>N-[[6-(pyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




104	 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the spiro ring is substituted with a 1H-pyrazole-3-carbonyl group.</p>	N-[[6-(1H-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
105	 <p>The structure is similar to 104, but the pyrazole ring is replaced by a 1-methylimidazole ring. The nitrogen of the imidazole ring is substituted with a methyl group.</p>	N-[[6-(1-methylimidazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
106	 <p>The structure is similar to 104, but the pyrazole ring is substituted with a methyl group at the 3-position. The carbonyl group is attached to the 4-position of the pyrazole ring.</p>	N-[[6-(3-methyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>107</p>		<p>N-[[6-(2-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>108</p>		<p>N-[[6-(5-methyl-1H-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>109</p>		<p>N-[[6-(1-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>




<p>110</p>		<p>N-[[6-(pyridazine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>111</p>		<p>N-[[6-(pyrimidine-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>112</p>		<p>N-[[6-(pyrazine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




<p>113</p>		<p>N-[[6-(pyridazine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>114</p>		<p>N-[[6-(1-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>115</p>		<p>N-[[6-(1,5-dimethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>




<p>116</p>		<p>N-[[6-(2,5-dimethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>117</p>		<p>N-[[6-(5-methylisoxazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>118</p>		<p>N-[[6-(4-methyl-1,2,5-oxadiazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>

119		N-[[6-(2-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
120		N-[[6-(4-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
121		N-[[6-(2-methylpyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>122</p>		<p>N-[[6-(5-methylpyrazine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>123</p>		<p>N-[[6-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>124</p>		<p>N-[[6-(1-ethyl-5-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

125		N-[[6-(3,5-dimethylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
126		N-[[6-(1-ethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
127		N-[[6-(6-cyanopyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide




128		N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
129		N-[[6-(2-morpholinopyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
130		N-[[6-(cyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>131</p>		<p>N-[[6-[1-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>132</p>		<p>N-[[6-(4-isoxazol-5-yl-1-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>133</p>		<p>N-[[6-(2-acetamidopyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>





<p>134</p>		<p>N-[[6-[3-(tetrazol-1-yl)-1H-pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>135</p>		<p>N-[[6-(6-acetamidopyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>136</p>		<p>N-[[6-(3-ethyl-5-methylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>





<p>137</p>		<p>N-[[6-(2-ethyl-5-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>138</p>		<p>N-[[6-(1-ethyl-3-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>139</p>		<p>N-[[6-(3-methylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>140</p>		<p>N-[[6-[2-methyl-3-(3-methylpyrazol-1-yl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>141</p>		<p>N-[[6-(1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>142</p>		<p>N-[[6-[2-(1-methylsulfonyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

143		N-[[6-(4-methylmorpholine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide
144		N-[[6-[1-(2-methylpropanoyl)azetidine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide
145		N-[[6-(1-acetylazetidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide

146		<p>N-[[6-(1-pyridazin-3-yl)pyrrolidine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
147		<p>N-[[6-[2-(1-pyrimidin-2-yl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
148		<p>N-[[6-[2-(1-isopropyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
149		<p>N-[[6-[2-(1-methyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
150		<p>N-[[6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide</p>

151		isopropyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
152		N-[[6-[2-(2-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
153		N-[[6-(3-methoxypropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
154		N-[[6-[2-[4-(trifluoromethylsulfonyl)phenyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

155		N-[[6-[3-[acetyl(methyl)amino]propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
156		N-[[6-[2-(2-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
157		N-[[6-[(4-cyanophenyl)carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
158		N-[[6-(m-tolylmethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

159		N-[[6-(p-tolylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
160		N-[[6-(benzylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
161		N-[[6-[(2-fluorophenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
162		N-[[6-(p-tolylmethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
163		N-[[6-[(4-fluorophenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
164		N-[[6-[(2-methoxyphenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

165		N-[[6-[(3-methoxyphenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
166		N-[[6-[(3-fluorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
167		N-[[6-[(4-methoxyphenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
168		N-[[6-[(3,4-dichlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
169		N-[[6-(1-adamantyl)carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
170		N-[[6-[(4-chlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
171		N-[[6-[(2-chlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

172		N-[[6-(ethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
173		N-[[6-[(2,4-dichlorophenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
174		N-[[6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
175		N-[[6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
176		N-[[6-[2-(2-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
177		N-[[6-[2-(3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
178		N-[[6-[2-(3-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide









<p>179</p>		<p>N-[[6-[4-(1,2,4-triazol-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>180</p>		<p>N-[[6-[2-(2-methylimidazol-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>181</p>		<p>N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

182		N-[[6-(tert-butylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
183		N-[[6-(isopropylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
184		N-[[6-(cyclohexylmethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
185		N-[[6-(propylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
186		N-[[6-(cyclohexylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

187		N-[[6-(phenylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
188		N-[[6-[2-(1-isopropyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
189		N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
190		N-[[6-[2-(4-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
191		N-[[6-(2-cyanoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
192		ethyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
193		tert-butyl 2-[(1H-pyrrolo[3,2-c]pyridine-2-carboxylamino)methyl]-7-azaspiro[3.5]nonane-7-carboxylate

194		tert-butyl 2-[(furo[2,3-c]pyridine-2-carbonylamino)methyl]-7-azaspiro[3.5]nonane-7-carboxylate
195		N-[[6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
196		N-[[6-(2-cyclohexylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
197		N-[[6-(3-cyclohexylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
198		N-[[6-(2-morpholinoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
199		N-[[6-(3-phenylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
200		tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino)methyl]-7-azaspiro[3.5]nonane-7-carboxylate
201		N-[[6-[2-(3,5-difluorophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide






202		N-[[6-[2-[3-(trifluoromethyl)phenyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
203		N-[[6-(tert-butylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
204		[2,2,2-trideuterio-1,1-bis(trideuteriomethyl)ethyl] 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
205		(2-methoxy-1,1-dimethyl-2-oxoethyl) 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
206		N-[[6-[2-(3-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
207		N-[[6-[2-(4-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
208		N-[[6-(phenylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide





209		N-[[6-[3-(3-pyridyl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
210		N-[[6-(benzylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
211		N-[(6-benzylsulfonyl-6-azaspiro[2.5]octan-2-yl)methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
212		N-[[6-(tert-butylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
213		N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide (isomer 1)
214		N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide (isomer 2)
215		(2-hydroxy-1,1-dimethyl-ethyl) 2-[(furo[2,3-c]pyridine-2-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
216		N-[[6-[2-(4-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide





218		(2-methoxy-1,1-dimethyl-ethyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
219		N-[[6-(2-tetrahydropyran-4-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
220		N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
221		N-[[6-(1-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
222		N-[[6-(1-isopropyl-3,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
223		N-[[6-[1-(4-fluorophenyl)-3,5-dimethylpyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
228		(1-methylcyclobutyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
229		N-[[6-(3-morpholinopropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide

230		N-[[6-(2,2-difluoro-2-phenylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
231		N-[[6-(3-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
232		N-[[6-(3-methylpyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
233		N-[[6-(6-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide





<p>234</p>		<p>N-[[6-(6-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>235</p>		<p>N-[[6-(4-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>236</p>		<p>N-[[6-(1H-benzimidazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>237</p>		<p>N-[[6-(pyrrolo[1,2-c]pyrimidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

238		N-[[6-(1H-benzotriazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
239		N-[[6-(1H-pyrrolo[2,3-b]pyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
240		N-[[6-(1H-indazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
241		N-[[6-(1H-benzimidazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
242		N-[[6-(4-cyanobenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

243		N-[[6-(3-cyanobenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
244		N-[[6-[4-[(5-methyl-1,2,4-oxadiazol-3-yl)methoxy]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
245		N-[[6-[4-(morpholinomethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
246		N-[[6-[4-[(2-methylimidazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

247		N-[[6-[4-(trifluoromethoxy)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
248		N-[[6-[4-(pyrrolidin-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
249		N-[[6-[3-(1,2,4-triazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
250		N-[[6-[2-(1,2,4-triazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide








251		N-[[6-[4-(1,2,4-triazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
252		N-[[6-[3-(imidazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
253		N-[[6-(4-imidazol-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
254		N-[[6-[4-(1H-1,2,4-triazol-5-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide







255		N-[[6-(6-imidazol-1-yl)pyridine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
256		N-[[6-[4-(trifluoromethyl)pyridine-2-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
257		N-[[6-[6-(trifluoromethyl)pyridine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
258		N-[[6-(3-oxo-4H-1,4-benzoxazine-6-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>259</p>		<p>N-[[6-(4-sulfamoylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>260</p>		<p>N-[[6-[2-(imidazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>261</p>		<p>N-[[6-(quinoxaline-6-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>262</p>		<p>N-[[6-(1,6-naphthyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

263		N-[[6-(2-methyl-1H-benzimidazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
264		N-[[6-(3-acetamidobenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
265		N-[[6-(3-acetamidopyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>266</p>		<p>N-[[6-(2,1,3-benzoxadiazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>267</p>		<p>N-[[6-(1,8-naphthyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>268</p>		<p>N-[[6-(isoquinoline-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>269</p>		<p>N-[[6-(quinoxaline-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>270</p>		<p>N-[[6-(1,5-naphthyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

271		N-[[6-(1,8-naphthyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
272		N-[[6-(isoxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
273		N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
274		tert-butyl 2-[(4,6-dihydro-1H-pyrrolo[3,4-c]pyrazole-5-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
275		N-[[6-[2-(1,4-dimethyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
276		N-[[6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
277		N-[[6-[2-(3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

278		N-[[6-[2-(4-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
279		N-[[6-(1,3-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
280		N-[[6-[4-(2-methyltetrazol-5-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
281		N-[[6-(4,6-dimethylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
282		N-[[6-(4-oxopentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
283		N-[[6-(4-hydroxy-4-methylpentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

284		N-[[6-(3-hydroxy-3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
285		N-[[6-(2-(2-pyridyl)acetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
286		N-[[6-(2-pyrimidin-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
287		N-[[6-(2-pyrazin-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
288		N-[[6-(3-thiazol-2-ylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
289		N-[[6-(2-phenoxyacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
290		N-[[6-(3-tetrahydropyran-4-ylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
291		N-[[6-(4-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

292		N-[[6-(3,5-dimethyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
293		N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
294		N-[[6-(1-tert-butyl-3,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
295		N-[[6-(1-benzyl-3,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
296		N-[[6-(5-tert-butyl-2-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
297		N-[[6-(2-ethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
298		N-[[6-(2-tert-butyl-4-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
299		N-[[6-(2,4-dimethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

300		N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide (isomer 1)
301		N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide (isomer 2)
302		N-[[6-(1H-indazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
303		N-[[6-[1-(1-isopropylpyrrolidin-3-yl)pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>304</p>		<p>N-[[6-[1-(1-acetylpyrrolidin-3-yl)pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>305</p>		<p>N-[[6-[5-(1,3-dimethylpyrazol-4-yl)isoxazole-3-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>306</p>		<p>tert-butyl 3-[2-((furo[2,3-c]pyridine-2-carboxylamino)methyl)-6-azaspiro[2.5]octane-6-carbonyl]-4,6-dihydro-1H-pyrrolo[3,4-c]pyrazole-5-carboxylate</p>
<p>307</p>		<p>N-[[6-[1-(1-methylpyrrolidin-3-yl)pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>





308		N-[[6-(2-tert-butyl-5-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
309		N-[[6-(2-methylimidazo[1,2-a]pyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
310		N-[[6-(6-methyl-1H-benzimidazole-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
311		N-[[6-(5-methylimidazo[1,2-a]pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
312		N-[[6-(6-fluoro-1H-indazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
313		N-[[6-(pyrazolo[1,5-a]pyrimidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

314		<p>N-[[6-(1-ethyl-3,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
315		<p>N-[[6-(6-methylimidazo[1,2-a]pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
316		<p>N-[[6-(pyrazolo[1,5-a]pyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
317		<p>N-[[6-(imidazo[1,2-a]pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
318		<p>N-[[6-(5-isopropylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

319		N-[[6-(3-isobutyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide
320		N-[[6-(imidazo[1,2-a]pyrimidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide
321		N-[[6-(pyrazolo[1,5-a]pyrimidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide
322		N-[[6-(3-tert-butyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide
323		N-[[6-(5-isopropyl-1H-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide

324		N-[[6-(3-isopropyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
325		N-[[6-(2-isopropylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
326		N-[[6-(5-ethyl-2-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
327		N-[[6-(5-cyclopropylisoxazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide

<p>328</p>		<p>N-[[6-(5-cyclopropylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>329</p>		<p>N-[[6-(3,5-dimethyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>330</p>		<p>N-[[6-(1,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>331</p>		<p>N-[[6-(4-methyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>332</p>		<p>N-[[6-(3-cyclopropyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>333</p>		<p>N-[[6-(1,4,5,6-tetrahydrocyclopenta[c]pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>334</p>		<p>N-[[6-(2,5-dimethyloxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>335</p>		<p>N-[[6-(1,3-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>

<p>336</p>		<p>N-[[6-(oxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>337</p>		<p>N-[[6-(isoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>338</p>		<p>N-[[6-(1H-imidazole-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>
<p>339</p>		<p>N-[[6-[4-(4-methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide</p>






340		N-[[6-(3H-imidazo[4,5-b]pyridine-6-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
341		N-[[6-(5-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
343		N-[[6-[4-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
344		N-[[6-[3-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
345		N-[[6-[4-[(dimethylamino)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
346		N-[[6-[4-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide

347		N-[[6-(1,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
348		N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
350		(3-methyloxetan-3-yl) 2-[[1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino]methyl]-6-azaspiro[2.5]octane-6-carboxylate
351		(2-hydroxy-1,1-dimethyl-ethyl) 2-[[1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino]methyl]-6-azaspiro[2.5]octane-6-carboxylate
352		N-[[6-(4-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
353		N-[[6-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
354		N-[[6-(4-isoxazol-5-yl-1-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
355		N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide

356		N-[[6-[2-(1,4-dimethyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
357		N-[[6-[2-(2-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
358		(2-acetamido-1,1-dimethylethyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
359		N-[[6-(2-thiazol-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
360		N-[[6-(4-hydroxy-4-methylpentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
361		N-[[6-[(3-methyloxetan-3-yl)carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
362		N-[[6-[4-(4-methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
363		N-[[6-[4-(4-methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

364		N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
365		N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide (isomer 1)
366		N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide (isomer 2)
367		tert-butyl 2-[[[(1,1,3,3-tetradeuteriopyrrolo[3,4-c]pyridine-2-carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate
368		N-[[6-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
369		N-[[6-(4-isoxazol-5-yl-1-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
370		N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
371		N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide






372		N-[[6-[4-(4-methylpiperazine-1-carbonyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
373		N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
375		(3-methyltetrahydrofuran-3-yl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
376		(2-hydroxy-2-methyl-propyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
377		N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
378		N-[[6-(2,2-dimethylpropylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
379		N-[[6-(2,2-dimethylpropylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
381		N-[[6-(2,2-dimethylpropylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide

382		N-[[6-(4,4,4-trifluorobutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
383		tert-butyl 2-[(pyrazolo[1,5-b]pyridazine-5-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
384		N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide (single isomer)
385		N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
387		N-[[6-[2-(2-hydroxy-2-methylpropoxy)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

388		N-[[6-[2-methyl-4-(trifluoromethyl)thiazole-5-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
389		N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
392		tert-butyl 2-[(imidazo[1,2-a]pyridin-6-yl)methylcarbamoylamino]methyl]-6-azaspiro[2.5]octane-6-carboxylate
393		N-[[6-[2-(2-hydroxy-2-methylpropoxy)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
394		N-[[6-[2-(6-amino-3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
395		N-[[6-(morpholine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
396		N-[[6-[5-(trifluoromethyl)-1H-pyrazole-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
397		N-[[6-(4-hydroxy-4-methylpentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide

398		N-[[6-[4-(trifluoromethyl)pyridine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
399		N-[[6-[2-(4-methyltetrahydropyran-4-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
400		N-[[6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
401		(2-hydroxy-2-methyl-propyl) 2-[(1H-pyrrolo[3,2-c]pyridine-2-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
402		tert-butyl 2-[(imidazo[1,2-b]pyridazine-6-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
403		N-[[6-[2-[4-methyl-1-(2,2,2-trifluoroethyl)-4-piperidyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
404		N-[[6-(2,4-dimethylthiazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
405		N-[[6-[4-(1-methyl-4-piperidyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide

406		tert-butyl 2-[[6-amino-1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate
407		N-[[6-(2-pyrimidin-5-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
408		1,1,3,3-tetradeuterio-N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]pyrrolo[3,4-c]pyridine-2-carboxamide
409		(3-methyloxetan-3-yl) 2-[[[(1,1,3,3-tetradeuteriopyrrolo[3,4-c]pyridine-2-carbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate
410		1,1,3,3-tetradeuterio-N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]pyrrolo[3,4-c]pyridine-2-carboxamide
411		N-[[6-(benzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide
413		N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
414		N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide

415		N-[[6-(4-methyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
416		isopropyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
417		(3-methyloxetan-3-yl) 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate
418		N-[(6-benzoyl-6-azaspiro[2.5]octan-2-yl)methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide
419		(3-methyloxetan-3-yl) (2S)-2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate

or a pharmaceutically acceptable salt thereof, or a stereoisomer or a pharmaceutically acceptable salt of said stereoisomer.

Pharmaceutical Description

- 5 The dosage forms of the present invention may contain a mixture of one or more compounds of this invention, and may include additional materials known to those skilled in the art as pharmaceutical excipients. "Excipient" includes any excipient commonly used in pharmaceuticals and should be selected on the basis of compatibility and the release profile properties of the desired dosage form.
- 10 Exemplary excipients include, e.g., binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like. Exemplary excipients include, e.g., acacia, gelatin, colloidal silicon dioxide, calcium glycerophosphate, calcium lactate, maltodextrin, glycerine,

magnesium silicate, sodium caseinate, soy lecithin, sodium chloride, tricalcium phosphate, dipotassium phosphate, sodium stearoyl lactylate, carrageenan, monoglyceride, diglyceride, pregelatinized starch, and the like. See, e.g., Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pa.

5 1975.

Exemplary excipients include: stabilizing additives such as gum acacia, gelatin, methyl cellulose, polyethylene glycol, carboxylic acids and salts thereof, and polylysine; acidifying agents (acetic acid, glacial acetic acid, citric acid, fumaric acid, hydrochloric acid, diluted hydrochloric acid, malic acid, nitric acid, phosphoric acid, 10 diluted phosphoric acid, sulfuric acid, tartaric acid); aerosol propellants (butane, dichlorodifluoro-methane, dichlorotetrafluoroethane, isobutane, propane, trichloromonofluoromethane); air displacements (carbon dioxide, nitrogen); alcohol denaturants (denatonium benzoate, methyl isobutyl ketone, sucrose octacetate); alkalinizing agents (strong ammonia solution, ammonium carbonate, diethanolamine, 15 diisopropanolamine, potassium hydroxide, sodium bicarbonate, sodium borate, sodium carbonate, sodium hydroxide, triethylamine); anticaking agents (see "glidant" below); antifoaming agents (dimethicone, simethicone); antimicrobial preservatives (benzalkonium chloride, benzalkonium chloride solution, benzethonium chloride, benzoic acid, benzyl alcohol, butylparaben, cetylpyridinium chloride, chlorobutanol, 20 chlorocresol, cresol, dehydroacetic acid, ethylparaben, methylparaben, methylparaben sodium, phenol, phenylethyl alcohol, phenylmercuric acetate, phenylmercuric nitrate, potassium benzoate, potassium sorbate, propylparaben, propylparaben sodium, sodium benzoate, sodium dehydroacetate, sodium propionate, sorbic acid, thimerosal, thymol); antioxidants (ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, 25 butylated hydroxytoluene, hypophosphorous acid, monothioglycerol, propyl gallate, sodium formaldehyde sulfoxylate, sodium metabisulfite, sodium thiosulfate, sulfur dioxide, tocopherol, tocopherols excipient); buffering agents (acetic acid, ammonium carbonate, ammonium phosphate, boric acid, citric acid, lactic acid, phosphoric acid, potassium citrate, potassium metaphosphate, potassium phosphate monobasic, sodium acetate, sodium citrate, sodium lactate solution, dibasic sodium phosphate, monobasic 30 sodium phosphate); capsule lubricants (see "tablet and capsule lubricant" below);

chelating agents (edetate disodium, ethylenediaminetetraacetic acid and salts, edetic acid); coating agents (sodium carboxymethylcellulose, cellulose acetate, cellulose acetate phthalate, ethylcellulose, gelatin, pharmaceutical glaze, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose phthalate, 5 methacrylic acid copolymer, methylcellulose, polyethylene glycol, polyvinyl acetate phthalate, shellac, sucrose, titanium dioxide, carnauba wax, microcrystalline wax, zein); colorants (caramel, red, yellow, black or blends, ferric oxide); complexing agents (ethylenediaminetetraacetic acid and salts (EDTA), edetic acid, gentisic acid ethanolmaide, oxyquinoline sulfate); desiccants (calcium chloride, calcium sulfate, 10 silicon dioxide); emulsifying and/or solubilizing agents (acacia, cholesterol, diethanolamine (adjunct), glyceryl monostearate, lanolin alcohols, lecithin, mono- and di-glycerides, monoethanolamine (adjunct), oleic acid (adjunct), oleyl alcohol (stabilizer), poloxamer, polyoxyethylene 50 stearate, polyoxyl 35 castor oil, polyoxyl 40 hydrogenated castor oil, polyoxyl 10 oleyl ether, polyoxyl 20 cetostearyl ether, 15 polyoxyl 40 stearate, polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80, propylene glycol diacetate, propylene glycol monostearate, sodium lauryl sulfate, sodium stearate, sorbitan monolaurate, sorbitan monooleate, sorbitan monopalmitate, sorbitan monostearate, stearic acid, trolamine, emulsifying wax); filtering aids (powdered cellulose, purified siliceous earth); flavors and perfumes (anethole, 20 benzaldehyde, ethyl vanillin, menthol, methyl salicylate, monosodium glutamate, orange flower oil, peppermint, peppermint oil, peppermint spirit, rose oil, stronger rose water, thymol, tolu balsam tincture, vanilla, vanilla tincture, vanillin); glidants and/or anticaking agents (calcium silicate, magnesium silicate, colloidal silicon dioxide, talc); humectants (glycerin, hexylene glycol, propylene glycol, sorbitol); 25 plasticizers (castor oil, diacetylated monoglycerides, diethyl phthalate, glycerin, mono- and di-acetylated monoglycerides, polyethylene glycol, propylene glycol, triacetin, triethyl citrate); polymers (e.g., cellulose acetate, alkyl celluloses, hydroxyalkylcelluloses, acrylic polymers and copolymers); solvents (acetone, alcohol, diluted alcohol, amylene hydrate, benzyl benzoate, butyl alcohol, carbon tetrachloride, 30 chloroform, corn oil, cottonseed oil, ethyl acetate, glycerin, hexylene glycol, isopropyl alcohol, methyl alcohol, methylene chloride, methyl isobutyl ketone,

mineral oil, peanut oil, polyethylene glycol, propylene carbonate, propylene glycol, sesame oil, water for injection, sterile water for injection, sterile water for irrigation, purified water); sorbents (powdered cellulose, charcoal, purified siliceous earth); carbon dioxide sorbents (barium hydroxide lime, soda lime); stiffening agents

5 (hydrogenated castor oil, cetostearyl alcohol, cetyl alcohol, cetyl esters wax, hard fat, paraffin, polyethylene excipient, stearyl alcohol, emulsifying wax, white wax, yellow wax); suspending and/or viscosity-increasing agents (acacia, agar, alginic acid, aluminum monostearate, bentonite, purified bentonite, magma bentonite, carbomer 934p, carboxymethylcellulose calcium, carboxymethylcellulose sodium,

10 carboxymethylcellulose sodium 12, carrageenan, microcrystalline and carboxymethylcellulose sodium cellulose, dextrin, gelatin, guar gum, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, magnesium aluminum silicate, methylcellulose, pectin, polyethylene oxide, polyvinyl alcohol, povidone, propylene glycol alginate, silicon dioxide, colloidal silicon dioxide, sodium

15 alginate, tragacanth, xanthan gum); sweetening agents (aspartame, dextrates, dextrose, excipient dextrose, fructose, mannitol, saccharin, calcium saccharin, sodium saccharin, sorbitol, solution sorbitol, sucrose, compressible sugar, confectioner's sugar, syrup); tablet binders (acacia, alginic acid, sodium carboxymethylcellulose, microcrystalline cellulose, dextrin, ethylcellulose, gelatin, liquid glucose, guar gum,

20 hydroxypropyl methylcellulose, methylcellulose, polyethylene oxide, povidone, pregelatinized starch, syrup); tablet and/or capsule diluents (calcium carbonate, dibasic calcium phosphate, tribasic calcium phosphate, calcium sulfate, microcrystalline cellulose, powdered cellulose, dextrates, dextrin, dextrose excipient, fructose, kaolin, lactose, mannitol, sorbitol, starch, pregelatinized starch, sucrose,

25 compressible sugar, confectioner's sugar); tablet disintegrants (alginic acid, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, starch, pregelatinized starch); tablet and/or capsule lubricants (calcium stearate, glyceryl behenate, magnesium stearate, light mineral oil, polyethylene glycol, sodium stearyl fumarate, stearic acid, purified stearic

30 acid, talc, hydrogenated vegetable oil, zinc stearate); tonicity agent (dextrose, glycerin, mannitol, potassium chloride, sodium chloride); vehicle: flavored and/or sweetened

(aromatic elixir, compound benzaldehyde elixir, iso-alcoholic elixir, peppermint water, sorbitol solution, syrup, tolu balsam syrup); vehicle: oleaginous (almond oil, corn oil, cottonseed oil, ethyl oleate, isopropyl myristate, isopropyl palmitate, mineral oil, light mineral oil, myristyl alcohol, octyldodecanol, olive oil, peanut oil, persic oil, 5 seame oil, soybean oil, squalane); vehicle: solid carrier (sugar spheres); vehicle: sterile (bacteriostatic water for injection, bacteriostatic sodium chloride injection); viscosity-increasing (see "suspending agent" below); water repelling agent (cyclomethicone, dimethicone, simethicone); and wetting and/or solubilizing agent (benzalkonium chloride, benzethonium chloride, cetylpyridinium chloride, docusate 10 sodium, nonoxynol 9, nonoxynol 10, octoxynol 9, poloxamer, polyoxyl 35 castor oil, polyoxyl 40, hydrogenated castor oil, polyoxyl 50 stearate, polyoxyl 10 oleyl ether, polyoxyl 20, cetostearyl ether, polyoxyl 40 stearate, polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80, sodium lauryl sulfate, sorbitan monolaureate, sorbitan monooleate, sorbitan monopalmitate, sorbitan monostearate, tyloxapol). This list is 15 not meant to be exclusive, but instead merely representative of the classes of excipients and the particular excipients which may be used in dosage forms of the present invention.

In certain aspects, the invention relates to methods of treating diseases or conditions mediated by elevated levels of NAMPT, or which are generally mediated 20 by NAMPT activity. Such disease or condition can be one or more selected from the group consisting of cancer, ovarian cancer, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, bladder cancer, pancreatic cancer, leukemia, lymphoma, Hodgkin's disease, viral infections, Human Immunodeficiency Virus, hepatitis virus, herpes virus, herpes simplex, inflammatory 25 disorders, irritable bowel syndrome, inflammatory bowel disease, rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, osteoarthritis, osteoporosis, dermatitis, atopic dermatitis, psoriasis, systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, graft-versus-host disease, Alzheimer's disease, cerebrovascular accident, atherosclerosis, diabetes, glomerulonephritis, 30 metabolic syndrome, non-small cell lung cancer, small cell lung cancer, multiple myeloma, leukemias, lymphomas, squamous cell cancers, kidney cancer, uretral and

bladder cancers, cancers of head and neck, and cancers of the brain and central nervous system (CNS).

The inventive compounds can be useful in the therapy of proliferative diseases such as, but not limited to cancer, autoimmune diseases, viral diseases, fungal
5 diseases, neurological/neurodegenerative disorders, arthritis, inflammation, anti-proliferative (e.g., ocular retinopathy), neuronal, alopecia and cardiovascular disease.

More specifically, the compounds can be useful in the treatment of a variety of cancers, including (but not limited to) the following: carcinoma, including that of the bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, non-
10 small cell lung cancer, head and neck, esophagus, gall bladder, ovary, pancreas, stomach, cervix, thyroid, prostate, and skin, including squamous cell carcinoma, hematopoietic tumors of lymphoid lineage, including leukemia, acute lymphocytic leukemia, acute lymphoblastic leukemia, B-cell lymphoma, T-cell lymphoma, Hodgkins lymphoma, non-Hodgkins lymphoma, hairy cell lymphoma, mantle cell
15 lymphoma, myeloma, and Burkett's lymphoma, hematopoietic tumors of myeloid lineage, including acute and chronic myelogenous leukemias, myelodysplastic syndrome and promyelocytic leukemia, tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma, tumors of the central and peripheral nervous system, including astrocytoma, neuroblastoma, glioma and schwannomas, and other
20 tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xenoderoma pigmentosum, keratoctanthoma, thyroid follicular cancer and Kaposi's sarcoma.

The compounds of the invention may induce or inhibit apoptosis.

The compounds of the invention may also be useful in the chemoprevention of cancer. Chemoprevention is defined as inhibiting the development of invasive cancer
25 by either blocking the initiating mutagenic event or by blocking the progression of pre-malignant cells that have already suffered an insult or inhibiting tumor relapse.

A further aspect of the invention is a method of inhibiting a NAMPT pathway in a subject, said method comprising administering to said subject a pharmaceutically acceptable amount of a compound of the invention to a subject in need thereof.

Another embodiment of the invention comprises a pharmaceutical formulation of the invention, wherein the pharmaceutical formulation, upon administration to a human, results in a decrease in tumor burden.

5 Still another embodiment of the invention is a pharmaceutical formulation comprising at least one compound of Formula I and a pharmaceutically acceptable excipient. Such formulations may further comprise one or more adjunctive active agent. The pharmaceutical formulations of the invention may further comprise a therapeutic effective amount of an adjunctive active agent.

10 The compounds of the present invention are also useful in combination therapies with at least one adjunctive active agent. Such methods include regimes in which the compound of the invention and the at least one adjunctive active agent are administered simultaneously or sequentially. Also useful are pharmaceutical compositions in which at least one compound of the present invention and at least one adjunctive active agent are combined in a single formulation.

15 The expression "adjunctive active agent" generally refers to agents which targets the same or a different disease, symptom, or medical condition as the primary therapeutic agent. Adjunctive active agents may treat, alleviate, relieve, or ameliorate side effects caused by administration of the primary therapeutic agents. Examples of adjunctive active agents include, but are not limited to, antineoplastic agents, 20 filgrastim, and erythropoietin. Such agents include those which modify blood cell growth and maturation. Non-limiting examples of adjunctive active agent are filgrastim, pegfilgrastim and erythropoietin. Other such adjunctive active agents include those which inhibit nausea associated with administration of chemotherapeutic agents, such as a 5-HT₃ receptor inhibitor (e.g., dolansetron, 25 granisetron, or ondansetron), with or without dexamethasone. The invention also describes one or more uses of the compounds of the present invention with an adjunctive active agent such as TNF, GCSF, or other chemotherapeutic agents. Additional adjunctive active agents include those that mediate cytotoxicity of NAMPT inhibitors, such as nicotinic acid rescue agents, or other compounds that play 30 a role in the NAMPT pathway, such as niacin (nicotinic acid), nicotinamide, or

related compounds, or modified release formulations of such compounds, for example, NIASPAN[®].

The terms "chemotherapeutic agent" and "antineoplastic agent" generally refer to agents, which treat, prevent, cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect malignancies and their metastasis. Examples of such agents include, but are not limited to, prednisone, fluorouracil (e.g., 5-fluorouracil (5-FU)), anastrozole, bicalutamide, carboplatin, cisplatin, chlorambucil, cisplatin, carboplatin, docetaxel, doxorubicin, flutamide, interferon-alpha, letrozole, leuprolide, megestrol, mitomycin, oxaliplatin, paclitaxel, plicamycin (Mithracin[™]), tamoxifen, thiotepa, topotecan, valrubicin, vinblastine, vincristine, and any combination of any of the foregoing.

The invention is also directed to a method of treating or preventing a disorder associated with excessive rate of growth of cells in a mammal comprising administering to the mammal an effective amount of the pharmaceutical formulation of the invention. Non-limiting examples of disorder include cancer or metastasis from malignant tumors.

Another aspect of the invention is a method of inhibiting tumor cell growth and rate of division in a mammal with cancer, or other disorder associated with abnormally dividing cells comprising administering to the mammal an effective amount of the pharmaceutical formulation of this invention.

Another embodiment of the invention is a method of treating bone pain due to excessive growth of a tumor or metastasis to bone in a mammal in need thereof comprising administering to the mammal an effective amount of the pharmaceutical formulation of this invention.

A further embodiment of the invention is a method of preparing a pharmaceutical formulation comprising mixing at least one compound of the present invention, and, optionally, one or more pharmaceutically acceptable excipients.

The invention is also directed to methods of synthesizing compounds of the present invention.

Still another aspect of this invention is to provide a method for treating, preventing, inhibiting or eliminating a disease or condition in a patient by inhibiting

NAMPT in said patient by administering a therapeutically effective amount of at least one compound of this disclosure, wherein said disease or condition is selected from the group consisting of cancer, ovarian cancer, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, bladder cancer, 5 pancreatic cancer, leukemia, lymphoma, Hodgkin's disease, viral infections, Human Immunodeficiency Virus, hepatitis virus, herpes virus, herpes simplex, inflammatory disorders, irritable bowel syndrome, inflammatory bowel disease, rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, osteoarthritis, osteoporosis, dermatitis, atopic dermatitis, psoriasis, systemic lupus erythematosus, multiple sclerosis, 10 psoriatic arthritis, ankylosing spondylitis, graft-versus-host disease, Alzheimer's disease, cerebrovascular accident, atherosclerosis, diabetes, glomerulonephritis, metabolic syndrome, non-small cell lung cancer, small cell lung cancer, multiple myeloma, leukemias, lymphomas, squamous cell cancers, kidney cancer, uretral and bladder cancers, cancers of head and neck, cancers of the brain and central nervous 15 system.

In a certain embodiment, the compounds of Formula I can be used in the treatment of solid and liquid tumors, non-small cell lung cancer, leukemia, lymphoma, ovarian cancer, glioma, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, rhino-gastric tumors, colorectal cancer, 20 CNS cancer, bladder cancer, pancreatic cancer and Hodgkin's disease.

In a certain embodiment, the compounds of Formula I can be used in the treatment of solid and liquid tumors, non-small cell lung cancer, leukemia, lymphoma, ovarian cancer, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, rhino-gastric tumors, colorectal cancer, bladder 25 cancer, pancreatic cancer and Hodgkin's disease.

Another embodiment is a pharmaceutical formulation comprising a pharmaceutically acceptable compound of the present invention, which provides, upon administration to a subject (e.g., a human), a decrease in tumor burden and/or metastases. The pharmaceutical formulation can be administered by oral means or 30 other suitable means.

Yet another embodiment is a method of treating ovarian cancer in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention.

5 Yet another embodiment is a method of treating non-small cell lung cancer (NSCLC) in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention.

10 Yet another embodiment is a method of treating colon cancer in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention.

Yet another embodiment is a method of treating breast cancer in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the pharmaceutical formulation of the present invention.

15 Yet another embodiment is a method of treating leukemia in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention.

20 Yet another embodiment is a method of treating colon cancer before or after surgical resection and/or radiation therapy, in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention.

25 Yet another embodiment is a method of treating cancer before or after surgical resection and/or radiation therapy, in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention, including adjunctive therapy to treat nausea, with or without dexamethasone.

30 Yet another embodiment is a method of treating cancer before or after surgical resection and or radiation therapy, in a subject (e.g., a human) in need thereof by administering to the subject an effective amount of the compound or the pharmaceutical formulation of the present invention, including adjunctive therapy with one or more additional therapeutic agents, or their pharmaceutically acceptable salts. Non-limiting examples of such additional therapeutic agents include cytotoxic

agents (such as for example, but not limited to, DNA interactive agents (such as cisplatin or doxorubicin)); taxanes (e.g. taxotere, taxol); topoisomerase II inhibitors (such as etoposide); topoisomerase I inhibitors (such as irinotecan (or CPT-11), camptostar, or topotecan); tubulin interacting agents (such as paclitaxel, docetaxel or the epothilones); hormonal agents (such as tamoxifen); thymidilate synthase inhibitors (such as 5-fluorouracil or 5-FU); anti-metabolites (such as methotrexate); alkylating agents (such as temozolomide, cyclophosphamide); Farnesyl protein transferase inhibitors (such as, SARASARTM, (4-[2-[4-[(11R)-3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,-6]cyclohepta[1,2-b]pyridin-11-yl]-1-piperidinyl]-2-oxoethyl]-1-piperidine- carboxamide, or SCH 66336), tipifarnib (Zamestra[®] or R115777 from Janssen Pharmaceuticals), L778,123 (a farnesyl protein transferase inhibitor from Merck & Company, Whitehouse Station, N.J.), BMS 214662 (a farnesyl protein transferase inhibitor from Bristol-Myers Squibb Pharmaceuticals, Princeton, N.J.); signal transduction inhibitors (such as, Iressa[®] (from Astra Zeneca Pharmaceuticals, England), Tarceva[®] (EGFR kinase inhibitors), antibodies to EGFR (e.g., C225), GLEEVEC[®] (C-abl kinase inhibitor from Novartis Pharmaceuticals, East Hanover, N.J.); interferons such as, for example, intron[®] (from Merck & Company), Peg-Intron[®] (from Merck & Company); hormonal therapy combinations; aromatase combinations; ara-C, adriamycin, cytoxan, and gemcitabine.

Other anti-cancer (also known as anti-neoplastic) agents include but are not limited to Uracil mustard, Chlormethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, oxaliplatin, leucovorin, oxaliplatin (ELOXATIN[®] from Sanofi-Synthelabo Pharmaceuticals, France), Pentostatine, Vinblastine, Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, Mithramycin, Deoxycoformycin, Mitomycin-C, L-Asparaginase, Teniposide 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrolacetate, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone,

Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, goserelin, Cisplatin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, Anastrozole, Letrazole, Capecitabine, Reloxafine, Droloxafine, Hexamethylmelamine, Avastin, herceptin, Bexxar, Velcade[®], Zevalin, Trisenox, Xeloda, Vinorelbine, Porfimer, 5 Erbitux, Liposomal, Thiotepa, Altretamine, Melphalan, Trastuzumab, Lerazole, Fulvestrant, Exemestane, Ifosfomide, Rituximab, C225, and Campath, 5-fluorouracil and leucovorin, with or without a 5-HT₃ receptor inhibitor (e.g., dolansetron, granisetron, ondansetron) with or without dexamethasone.

10 Additionally, according to the present invention, the compounds of the invention described herein may be administered and/or formulated in combination with an adjunctive active agent. In certain embodiments, the adjunctive active agent is niacin, nicotinamide, nicotinic acid, nicotinamide mononucleotide (NMN), or variations thereof, including modified release formulations of niacin, such as 15 NIASPAN[®]. Niacin, nicotinamide, nicotinic acid, nicotinamide mononucleotide (NMN), or variations thereof have also been described in the literature as “rescue agents” or “rescuing agents” and these terms have been used herein. The role of nicotinamide and/or nicotinic acid as a rescuing or rescue agent has for example been described by Beauparlant et al. in *Anti-Cancer Drugs* 2009, 20:346–354 and by 20 Rongvaux et al. in *The Journal of Immunology*, 2008, 181: 4685–4695. These two references describe the role of a rescuing or rescue agent with regards to ameliorating possible toxic effects of NAMPT inhibitors.

 If formulated as a fixed dose, such combination products employ the 25 compounds of this invention within the dosage range described herein (or as known to those skilled in the art) and the other pharmaceutically active agents or treatments within its dosage range. For example, the CDC2 inhibitor olomucine has been found to act synergistically with known cytotoxic agents in inducing apoptosis (*J. Cell Sci.*, (1995) 108, 2897). The compounds of the invention may also be administered 30 sequentially with known anticancer or cytotoxic agents when a combination formulation is inappropriate. In any combination treatment, the invention is not

limited in the sequence of administration; compounds of the disclosed Formulas may be administered either prior to or after administration of the known anticancer or cytotoxic agent. For example, the cytotoxic activity of the cyclin-dependent kinase inhibitor flavopiridol is affected by the sequence of administration with anticancer agents. *Cancer Research*, (1997) 57, 3375. Such techniques are within the skills of persons skilled in the art as well as attending physicians.

Any of the aforementioned methods may be augmented by administration of fluids (such as water), loop diuretics, one or more adjunctive active agents, such as a chemotherapeutic or antineoplastic agent, such as leucovorin and fluorouracil, or an adjunctive chemotherapeutic agent (such as filgrastim and erythropoietin), or any combination of the foregoing.

Yet another embodiment is a method for administering a compound of the instant invention to a subject (e.g., a human) in need thereof by administering to the subject the pharmaceutical formulation of the present invention.

Yet another embodiment is a method of preparing a pharmaceutical formulation of the present invention by mixing at least one pharmaceutically acceptable compound of the present invention, and, optionally, one or more pharmaceutically acceptable additives or excipients.

For preparing pharmaceutical compositions from the compounds described by this invention, inert, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules, capsules, cachets and suppositories. The powders and tablets may be comprised of from about 5 to about 95 percent active ingredient. Suitable solid carriers are known in the art, e.g., magnesium carbonate, magnesium stearate, talc, sugar or lactose. Tablets, powders, cachets and capsules can be used as solid dosage forms suitable for oral administration. Examples of pharmaceutically acceptable carriers and methods of manufacture for various compositions may be found in A. Gennaro (ed.), *Remington's Pharmaceutical Sciences*, 18th Edition, (1990), Mack Publishing Co., Easton, Pa.

The compositions and formulations of the invention can be administered as sterile compositions and sterile formulations. Sterile pharmaceutical formulations are

compounded or manufactured according to pharmaceutical-grade sterilization standards (e.g., United States Pharmacopeia Chapters 797, 1072, and 1211; California Business & Professions Code 4127.7; 16 California Code of Regulations 1751, 21 Code of Federal Regulations 21, or ex-U.S. counterparts to such regulations) known
5 to those of skill in the art.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injection or addition of sweeteners and opacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal
10 administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier, such as an inert compressed gas, e.g. nitrogen.

Also included are solid form preparations that are intended to be converted,
15 shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions can take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type
20 as are conventional in the art for this purpose.

The compounds of this invention may also be delivered subcutaneously.

The compound can be administered orally or intravenously.

The pharmaceutical preparation can be in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses containing appropriate
25 quantities of the active component, e.g., an effective amount to achieve the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 1 mg to about 1000 mg, for example from about 1 mg to about 500 mg, in particular from about 1 mg to about 250 mg, or from about 1 mg to about
30 25 mg, according to the particular application.

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage regimen for a particular situation is within the skill of the art. For convenience, the total daily dosage may be divided and administered in portions during the day as required.

The amount and frequency of administration of the compounds of the invention and/or the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended daily dosage regimen for oral administration can range from about 1 mg/day to about 500 mg/day, preferably 1 mg/day to 200 mg/day, in two to four divided doses.

Schemes and Examples

Exemplary, non-limiting, chemical entities and methods useful in preparing compounds of the invention will now be described by reference to illustrative synthetic schemes for their general preparation below and the specific examples that follow. Those skilled in the art will appreciate that other synthetic routes may be used to synthesize the compounds according to the invention. Although specific starting materials and reagents are depicted and discussed herein, other starting materials and reagents can be easily substituted to provide a variety of derivatives and/or reaction conditions. In addition, many of the exemplary compounds prepared by the described methods can be further modified in light of this disclosure using conventional chemistry well known to those skilled in the art.

Artisans will recognize that, to obtain the various compounds herein, starting materials may be suitably selected so that the ultimately desired substituents will be carried through the reaction scheme with or without protection as appropriate to yield the desired product. Alternatively, it may be necessary or desirable to employ, in the place of the ultimately desired substituent, a suitable group that may be carried through the reaction scheme and replaced as appropriate with the desired substituent. Each of the reactions depicted in the reaction schemes is preferably run at a

temperature from about 0 °C to the reflux temperature of the solvent used. Unless otherwise specified, the variables shown in the schemes below are as defined above in reference to Formula I.

Compounds according to the invention may be synthesized by synthetic routes that include processes analogous to those well-known in the chemical arts, particularly in light of the description contained herein, and those for other heterocycles described in: Comprehensive Heterocyclic Chemistry II, Editors Katritzky and Rees, Elsevier, 1997, e.g. Volume 3; Liebigs Annalen der Chemie, (9):1910-16, (1985); Helvetica Chimica Acta, 41:1052-60, (1958); Arzneimittel-
10 Forschung, 40(12):1328-31, (1990), each of which are expressly incorporated by reference. Starting materials are generally available from commercial sources such as Sigma-Aldrich Chemicals (Milwaukee, WI) or are readily prepared using methods well known to those skilled in the art (e.g., prepared by methods generally described in Louis F. Fieser and Mary Fieser, Reagents for Organic Synthesis, v. 1-23, Wiley,
15 N.Y. (1967-2006 ed.), or Beilsteins Handbuch der organischen Chemie, 4, Aufl. ed. Springer-Verlag, Berlin, including supplements (also available via the Beilstein online database).

Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing compounds according to the
20 invention and necessary reagents and intermediates are known in the art and include, for example, those described in R. Larock, Comprehensive Organic Transformations, VCH Publishers (1989); T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Synthesis, 3rd Ed., John Wiley and Sons (1999); and L. Paquette, ed., Encyclopedia of Reagents for Organic Synthesis, John Wiley and Sons (1995) and
25 subsequent editions thereof. The need for such protection will vary depending on the nature of the remote functionality and the conditions of the preparation methods. Suitable amino-protecting groups include acetyl, trifluoroacetyl, t-butoxycarbonyl (BOC), benzyloxycarbonyl (CBz) and 9-fluorenylmethyleneoxycarbonyl (Fmoc). The need for such protection is readily determined by one skilled in the art.

Additional particularly useful reactions in preparing compounds of the present invention include alkylation, reductive amination, oxidation, reduction, and hydrolysis reactions. Such transformations are well within the ordinary skill in the art.

Compounds according to the invention may be prepared singly or as
5 compound libraries comprising, for example, at least two, or 5 to 1,000 compounds, or 10 to 100 compounds. Libraries of compounds of Formula I may be prepared by a combinatorial "split and mix" approach or by multiple parallel syntheses using either solution phase or solid phase chemistry, by procedures known to those skilled in the art. Thus, according to a further aspect of the invention there is provided a compound
10 library comprising at least two compounds of Formula I, or pharmaceutically acceptable salts thereof.

In the methods of preparing compounds according to the invention, it may be advantageous to separate reaction products from one another and/or from starting materials. The desired products of each step or series of steps is separated and/or
15 purified to the desired degree of homogeneity by the techniques common in the art. Typically such separations involve multiphase extraction, crystallization from a solvent or solvent mixture, distillation, sublimation, or chromatography. Chromatography can involve any number of methods including, for example: reverse-phase and normal phase; size exclusion; ion exchange; high, medium and low
20 pressure liquid chromatography methods and apparatus; small scale analytical; simulated moving bed (SMB) and preparative thin or thick layer chromatography, as well as techniques of small scale thin layer and flash chromatography.

Another class of separation methods involves treatment of a mixture with a reagent selected to bind to or render otherwise separable a desired product, unreacted
25 starting material, reaction by product, or the like. Such reagents include adsorbents or absorbents such as activated carbon, molecular sieves, ion exchange media, or the like. Alternatively, the reagents can be acids in the case of a basic material, bases in the case of an acidic material, binding reagents such as antibodies, binding proteins, selective chelators such as crown ethers, liquid/liquid ion extraction reagents (LIX),
30 or the like. Selection of appropriate methods of separation depends on the nature of the materials involved, such as, boiling point and molecular weight in distillation and

sublimation, presence or absence of polar functional groups in chromatography, stability of materials in acidic and basic media in multiphase extraction, and the like.

A single stereoisomer, e.g., an enantiomer, substantially free of its stereoisomer may be obtained by resolution of the racemic mixture using a method such as formation of diastereomers using optically active resolving agents (Eliel, E. and Wilen, S. "Stereochemistry of Organic Compounds," John Wiley & Sons, Inc., New York, 1994; Lochmuller, C. H., (1975) *J. Chromatogr.*, 113(3):283-302). Racemic mixtures of chiral compounds of the invention can be separated and isolated by any suitable method, including: (1) formation of ionic, diastereomeric salts with chiral compounds and separation by fractional crystallization or other methods, (2) formation of diastereomeric compounds with chiral derivatizing reagents, separation of the diastereomers, and conversion to the pure stereoisomers, and (3) separation of the substantially pure or enriched stereoisomers directly under chiral conditions. See: "Drug Stereochemistry, Analytical Methods and Pharmacology," Irving W. Wainer, Ed., Marcel Dekker, Inc., New York (1993)

Under method (1), diastereomeric salts can be formed by reaction of enantiomerically pure chiral bases such as brucine, quinine, ephedrine, strychnine, *α*-methyl-*b*-phenylethylamine (amphetamine), and the like with asymmetric compounds bearing acidic functionality, such as carboxylic acid and sulfonic acid. The diastereomeric salts may be induced to separate by fractional crystallization or ionic chromatography. For separation of the optical isomers of amino compounds, addition of chiral carboxylic or sulfonic acids, such as camphorsulfonic acid, tartaric acid, mandelic acid, or lactic acid can result in formation of the diastereomeric salts.

Alternatively, by method (2), the substrate to be resolved is reacted with one enantiomer of a chiral compound to form a diastereomeric pair (E. and Wilen, S. "Stereochemistry of Organic Compounds", John Wiley & Sons, Inc., 1994, p. 322). Diastereomeric compounds can be formed by reacting asymmetric compounds with enantiomerically pure chiral derivatizing reagents, such as menthyl derivatives, followed by separation of the diastereomers and hydrolysis to yield the pure or enriched enantiomer. A method of determining optical purity involves making chiral esters, such as a menthyl ester, e.g., (-) menthyl chloroformate in the presence of base,

or Mosher ester, *a*-methoxy-*a*-(trifluoromethyl)phenyl acetate of the racemic mixture and analyzing the ¹H NMR spectrum for the presence of the two atropisomeric enantiomers or diastereomers (Jacob III. J. Org. Chem. (1982) 47:4165). Stable diastereomers of atropisomeric compounds can be separated and isolated by normal- and reverse-phase chromatography following methods for separation of atropisomeric naphthyl-isoquinolines (WO 96/15111). By method (3), a racemic mixture of two enantiomers can be separated by chromatography using a chiral stationary phase ("Chiral Liquid Chromatography" (1989) W. J. Lough, Ed., Chapman and Hall, New York; Okamoto, J. Chromatogr., (1990) 513:375-378). Enriched or purified enantiomers can be distinguished by methods used to distinguish other chiral molecules with asymmetric carbon atoms, such as optical rotation and circular dichroism.


Abbreviations and acronyms used in the following Schemes and elsewhere herein are defined as follows:

CDCl ₃	deuterated chloroform
CD ₃ OD	deuterated methanol
δ	chemical shift (ppm)
DCM	Dichloromethane
DIPEA	Diisopropylethylamine
DMF	N,N-dimethylformamide
DMSO	Dimethyl sulfoxide
EDCI	1-Ethyl-3-(3-dimethylaminopropyl) carbodiimide
ELSD	Evaporative light scattering detector
equiv	Molar equivalent
ESI	Electrospray ionization
h	Hour(s)
H ₂	Hydrogen gas
HATU	<i>O</i> -(7-Azabenzotriazol-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
¹ H NMR	proton nuclear magnetic resonance spectroscopy
HOBt	1-Hydroxybenzotriazole
HPLC	High performance liquid chromatography
LC/MS	Liquid chromatography - mass spectrometry
MeOH	Methanol
MHz	megahertz
min	Minute(s)
PDA	Photo diode array detector
psi	Pounds per square inch

rt	Room temperature
Raney-Ni	Raney Nickel
R _f	Retention factor
TFA	Trifluoroacetic acid
Tf ₂ O	Trifluoromethanesulfonic anhydride
THF	Tetrahydrofuran
TLC	Thin layer chromatography

Exemplary general reaction schemes that are useful in preparing compounds of the invention are described below.

General Scheme A




Compounds of Formula I may be prepared as shown above in Scheme A. Compounds of Formula A, in which X is, for example, OH, chloro, or bromo, are reacted with amines B to produce compounds of Formula I. Where X is OH, coupling reactions may occur in the presence of a coupling reagent such as EDCI, HATU, or HOBt, and a base (e.g., K₂CO₃, Cs₂CO₃, trialkylamine, sodium or potassium alkoxide) in an inert solvent such as dichloromethane, N,N-dialkylformamide, N,N-dialkylacetamide, dialkylethers, cyclic ethers, DMSO, or N-methyl-2-pyrrolidinone, or a mixture thereof, at temperatures ranging from -78 °C to 200 °C. Such coupling reactions between amines and acids are well-known in the art.

10

Alternatively, compounds A where X is bromo or chloro may be reacted with amines B in the presence of a suitable base, such as triethylamine, K₂CO₃, or Cs₂CO₃, to form compounds of Formula I.

15

General Scheme B



Methods of Chemical Analysis of Example Compounds

Unless otherwise indicated, ^1H NMR spectra were recorded at ambient temperature using one of the following machines: Varian Unity Inova (400 MHz) spectrometer with a triple resonance 5 mm probe, Bruker Avance DRX400 (400 MHz) spectrometer with a triple resonance 5 mm probe, a Bruker Avance DPX 300 (300 MHz) equipped with a standard 5 mm dual frequency probe for detection of ^1H and ^{13}C , a Bruker AVIII (400 MHz) using a BBI Broad Band Inverse 5 mm probe, or a Bruker AVIII (500 MHz) using a QNP (Quad Nucleus detect) 5 mm probe. Chemical shifts are expressed in ppm relative to an internal standard; tetramethylsilane (ppm = 0.00). The following abbreviations have been used: br = broad signal, s = singlet, d = doublet, dd = double doublet, t = triplet, q = quartet, m = multiplet.

High Performance Liquid Chromatography - Mass Spectrometry (LC/MS) and supercritical fluid chromatography (SFC) experiments to determine retention times (RT) and associated mass ions, *e.g.*, $[\text{M}+\text{H}]^+$, $[\text{M}+\text{Na}]^+$, $[\text{M}-\text{H}]^-$, were performed using one of the following methods:

Method A

Instrument: SHIMADZU LCMS-2010EV

LC Parameters: Column: Shim-pack XR-ODS, 2.2 μm , 3.0*50 mm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 5% to 100% B in 2.0 min, 100% B for 1.1 min, 100% to 5% B in 0.2 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 $^\circ\text{C}$; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 μL .

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 $^\circ\text{C}$; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.7 kv.

Method B

Instrument: SHIMADZU LCMS-2010EV

LC Parameters: Column: Waters XBridge C18, 3.0x50 mm, 3.5 μm ; Mobile Phase A: Water/5mM Ammonium Acetate; Mobile Phase B: Methanol; Gradient: 10% to

100% B in 1.8 min, 100% B for 1.3 min, 100% to 10% B in 0.1 min, then stop; Flow Rate: 0.9 mL/min; Column Temperature: 40 °C; Detector: PDA and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 µL.

MS Parameters: Interface: ESI (Positive & Negative); Interface Voltage: 4.0kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900(m/z); Detector voltage: 1.5 kv.

Method C

Instrument: SHIMADZU LCMS-2010EV

10 LC Parameters: Column: Shim-pack XR-ODS, 2.2 µm, 3.0*50 mm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile/0.05% TFA; Gradient: 5% to 100% B in 2.0 min, 100% B for 1.1 min, 100% to 5% B in 0.2 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 µL.

15 MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.5 kv.

Method D

20 Instrument: SHIMADZU LC/MS-2010EV

LC Parameters: Column: Waters Xselect C18, 3.0x50 mm, 3.5 µm; Mobile Phase A: Water/0.1% formic acid; Mobile Phase B: Acetonitrile/0.05% formic acid; Gradient: 5% to 100% B in 2.0 min, 100% B for 1.2 min, 100% to 5% B in 0.1 min, then stop; Flow Rate: 0.9 mL/min; Column Temperature: 35 °C; Detector: 254 nm and ELSD;

25 Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 µL.

MS Parameters: Interface: ESI (Positive & Negative); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.5 kv.

30

Method E

Instrument: SHIMADZU LCMS-2010EV

LC Parameters: Column: Shim-pack XR-ODS, 3.0x50 mm, 2.2 μ m; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 5% to 100% B in 2.0 min, 100% B for 1 min, 100% to 5% B in 0.3 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.3 kv.

10 Method F

Instrument: SHIMADZU LCMS-2010EV

LC Parameters: Column: Shim-pack XR-ODS, 3.0x50 mm, 2.2 μ m; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 5% to 100% B in 2.0 min, 100% B for 1.2 min, 100% to 5% B in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 70-900 (m/z); Detector voltage: 1.1 kv.

20 Method G

Instrument: SHIMADZU LC/MS-2020EV

LC Parameters: Column: Shim-pack XR-ODS, 50 mm*3.0 mm, 2.2 μ m; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 5% to 100% B in 2.1 min, 100% B for 0.8 min, 100% to 5% B in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Acetonitrile; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.05 kv.

Method H

Instrument: SHIMADZU LCMS-2020

LC Parameters: Column: Shim-pack XR-ODS, 2.2 μ m, 3.0*50 mm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile/0.05% TFA; Gradient: 5% to 100%

5 B in 2.0 min, 100% B for 1.2 min, 100% to 5% B in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage:

10 1.1 kv.

Method I

Instrument: SHIMADZU LCMS-2020

LC Parameters: Column: Shim-pack XR-ODS 50*3.0 mm, 2.2 μ m; Mobile Phase A:

15 Water/0.05% TFA; Mobile Phase B: Acetonitrile/0.05% TFA; Gradient: 5% B to 100% B for 2.0 min, 100% B for 1.2 min, 100% B to 5% in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD;

Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block:

20 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 70-900 (m/z); Detector voltage: 1.05 kv.

Method J

Instrument: SHIMADZU LCMS-2020

25 LC Parameters: Column: Shim-pack XR-ODS, 3.0x50 mm, 2.2 μ m; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 5% to 100% B in 2.0 min, 100% B for 1.2 min, 100% to 5% B in 0.2 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Acetonitrile; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 200 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.05 kv.

5

Method K

Instrument: SHIMADZU LCMS-2020

LC Parameters: Column: Gemini-NX 3u C18 110A; Mobile Phase A: Water/0.04% Ammonia; Mobile Phase B: Acetonitrile; Gradient: 5% to 100% B in 2.0 min, 100% B for 1.1 min, 100% to 5% B in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column
10 Temperature: 35 °C; Detector: 254 nm and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 µL.

MS Parameters: Interface: ESI (Positive & Negative); Interface Voltage: 4.5 kv; Heat Block: 200 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 0.75 kv.

15

Method I.

Instrument	Waters mass-directed
Mobile phase A	0.1% H ₂ O w/ NH ₄ OH
Mobile phase B	acetonitrile
Column	Phenomenex Gemini N-X C18, 10 um, 21.5x100 mm
Column temperature	25 °C
LC gradient	5 to 85% in 10 min.
LC Flowrate	35 mL/min
UV wavelength	254 nm
Mass Spectrometer	Waters 3100
Ionization	ES+

Method M

HPLC	Waters mass-directed
Mobile phase A	0.1% H ₂ O w/ NH ₄ OH
Mobile phase B	acetonitrile

Column	Phenomenex Gemini N-X C18, 10 μ m, 30x100 mm
Column temperature	25 °C
LC gradient	5 to 50% in 15 min.
LC Flowrate	60 mL/min
UV wavelength	254 nm
Mass Spectrometer	Waters 3100
Ionization	ES+

Method N

Instrument: SHIMADZU LC/MS-2020

LC Parameters: Column: Shim-pack XR-ODS, 2.2 μ m, 3.0*50 mm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 5% B to 100% B for 2.0 min, 100% B for 1.2 min, 100% B to 5% in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: UV and ELSD; Sample Preparation: 1 mg/mL in Methanol; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 70-900 (m/z); Detector voltage: 1.1 kv.

Method O

Instrument: SHIMADZU UHPLCMS-2020EV (LC-30AD pump, Binary solvent manager, SIL-AC Auto Samples, SPDM20A Detector, Alltech 3300 ELSD Detector

LC Parameters: Column: Shim-pack XR-ODS, 1.6 μ m, 2.0*50 mm; Mobile Phase A: Water/0.1% formic acid; Mobile Phase B: Acetonitrile/0.1% formic acid; Gradient: 5% B to 100% B for 2.0 min, 100% B for 1.1 min, 100% B to 5% in 0.1 min, then stop; Flow Rate: 0.7 mL/min; Column Temperature: 40 °C; Detector: Diode Array Detector (DAD) and ELSD; Injection Volume: 1 μ L.

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.0 kv; Heat Block: 200 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 0.9 kv.

Method P

Instrument: SHIMADZU LC/MS-2020

LC Parameters: Column: Shim-pack XR-ODS, 2.2 μm , 3.0*50 mm; Mobile Phase A:

Water/0.1% formic acid; Mobile Phase B: Acetonitrile/0.05% formic acid; Gradient:

- 5 5% B to 100% B for 2.0 min, 100% B for 1.1 min, 100% B to 5% in 0.1 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: PDA and ELSD; Sample Preparation: 1 mg/mL in acetonitrile; Injection Volume: 1 μL .

MS Parameters: Interface: ESI (Positive); Interface Voltage: tuning file; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage:

- 10 0.9 kv.

Method Q

Instrument: SHIMADZU LC/MS-2020

LC Parameters: Column: Shim-pack XR-ODS, 2.2 μm , 3.0*50 mm; Mobile Phase A:

- 15 Water/0.1% formic acid; Mobile Phase B: Acetonitrile/0.05% formic acid; Gradient: 5% B to 100% B for 2.0 min, 100% B for 1.2 min, 100% B to 5% in 0.2 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: UV and ELSD; Sample Preparation: 1 mg/mL in acetonitrile; Injection Volume: 1 μL .

MS Parameters: Interface: ESI (Positive); Interface Voltage: 4.5 kv; Heat Block:

- 20 200 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 0.95 kv.

Method R

Instrument: SHIMADZU LC/MS-2020

- 25 LC Parameters: Column: Shim-pack XR-ODS, 2.2 μm , 3.0*50 mm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile/0.05% TFA; Gradient: 5% B to 100% B for 1.2 min, 100% B for 0.9 min, 100% B to 5% in 0.2 min, then stop; Flow Rate: 1.0 mL/min; Column Temperature: 40 °C; Detector: PDA and ELSD; Sample Preparation: 1 mg/mL in acetonitrile; Injection Volume: 1 μL .

MS Parameters: Interface: ESI (Positive); Interface Voltage: tuning file; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 1.1 kv.

5 Method S

Instrument: SHIMADZU UHPLC/MS-2020

LC Parameters: Column: Shim-pack XR-ODS, 1.6 µm, 2.0*50 mm; Mobile Phase A: Water/0.1% formic acid; Mobile Phase B: Acetonitrile/0.05% formic acid; Gradient: 5% B to 100% B for 2.0 min, 100% B for 1.1 min, 100% B to 5% in 0.1 min, then
10 stop; Flow Rate: 0.7 mL/min; Column Temperature: 40 °C; Detector: PDA and ELSD; Sample Preparation: 1 mg/mL in acetonitrile; Injection Volume: 1 µL.

MS Parameters: Interface: ESI (Positive); Interface Voltage: tuning file; Heat Block: 250 °C; Nebulizing Gas: 1.50 L/min; Scan Range: 90-900 (m/z); Detector voltage: 0.85 kv.

15

Method T

Instrument: HPLC Agilent 1200

LC Parameters: Column: Agilent SB C18, 2.1*30 mm, 1.8 µm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 3% B for 0.3 min, 3% B
20 to 95% B in 6.5 min, 95% B for 1.5 min, 95% to 3% B in 0.1 min, then stop; Flow Rate: 0.4 mL/min; Column Temperature: 25 °C; Detector: 254 nm.

MS Parameters: Agilent 6140 Quadrupole LC/MS; Interface: ESI (Positive); Scan Range: 90-1300 amu.

25 Method U

Instrument: Waters Acquity UPLC


LC Parameters: Column: Acquity UPLC BEH C18, 1.7 µm, 2.1*50 mm; Mobile Phase A: Water/0.05% TFA; Mobile Phase B: Acetonitrile; Gradient: 2% to 98% B in
17.5 min, 98% B for 1.5 min, equilibrate for 1.5 min, then stop; Flow Rate: 0.6
30 mL/min; Column Temperature: 40 °C; Detector: 254 nm and 220 nm.

MS Parameters: Waters LCT Premier XE; Interface: ESI (Positive); Scan Range: 80-1300 amu; Detector: Time of flight.

The following examples illustrate the preparation of representative compounds of the invention. Unless otherwise specified, all reagents and solvents were of standard commercial grade and were used without further purification.

I. Preparation of Intermediates

Intermediate 1: Furo[2,3-c]pyridine-2-carboxylic acid



10

Step 1. Ethyl 3-hydroxyisonicotinate. A solution of 3-hydroxyisonicotinic acid (495 g, 3.56 mol) in ethanol (7 L) and concentrated H₂SO₄ (250 mL) was heated under reflux for 72 h and then cooled to rt and concentrated under reduced pressure to remove the solvent. The residue was dissolved in water (3 L) and the pH was adjusted to 4 by addition of saturated aqueous NaHCO₃ solution. The resulting precipitate was removed by filtration and the filtrate was extracted with DCM (2 L×3). The combined organic phase was washed with brine, dried over anhydrous Na₂SO₄, and then concentrated under reduced pressure to give ethyl 3-hydroxyisonicotinate (414 g, 70%) as yellow oil.

15

Step 2. Ethyl 3-(2-ethoxy-2-oxoethoxy)isonicotinate. To a solution of triphenylphosphine (780 g, 2.97 mol) in THF (6 L) at -10 °C was added dropwise diisopropyl azodicarboxylate (600 mL, 2.97 mol). The reaction mixture was stirred at -10 °C for 30 min and then ethyl 3-hydroxyisonicotinate (414 g, 2.48 mol) in THF (1 L) solution was added dropwise. The resulting mixture was stirred at rt for 16 h and then concentrated under reduced pressure. The residue was partitioned between ethyl acetate (4 L) and 1 N HCl (2 L). The aqueous layer was separated and the organic phase was extracted by 1 N HCl (1 L×2). The combined aqueous layers were slowly adjusted to pH 8 by addition of solid NaHCO₃ and then extracted with ethyl acetate (2 L×2). The combined organic layers were dried over anhydrous Na₂SO₄ and then

20

25

concentrated under reduced pressure to give the title compound (380 g, 61%) as a brown oil.


Step 3. Ethyl 3-hydroxyfuro[2,3-c]pyridine-2-carboxylate. To a suspension of NaH (72 g, 1.8 mol, 60% suspension in mineral oil) in anhydrous THF (2 L) at 0 °C was added dropwise a solution of ethyl 3-(2-ethoxy-2-oxoethoxy)isonicotinate (380 g, 1.5 mol) in THF (1 L) under argon. The reaction mixture was stirred at rt for 16 h and then carefully quenched with ice water (1 L). The resulting mixture was concentrated to a volume of 1.2 L and then diluted with saturated aqueous NaHCO₃ solution (2.5 L), and stirred for an additional 30 min. The precipitated solid was collected by filtration and washed with ethyl acetate (1 L). The filtrate was washed with ethyl acetate (1 L×2) and the aqueous layer was combined with the solid and carefully acidified to a pH of 5 with acetic acid. The resulting solid was collected by filtration and dried under vacuum to give the title compound (210 g, 68%) as a yellow solid.

Step 4. Ethyl 3-(((trifluoromethyl)sulfonyl)oxy)furo[2,3-c]pyridine-2-carboxylate. To a solution of ethyl 3-hydroxyfuro[2,3-c]pyridine-2-carboxylate (210 g, 1.01 mol) and pyridine (107 mL, 1.3 mol) in anhydrous DCM (3 L) at 0 °C was added dropwise triflic anhydride (203 g, 1.2 mol). The reaction mixture was stirred at rt for 16 h and then quenched with ice water (1 L). The aqueous layer was extracted with DCM (1 L×2) and the combined organic layer was dried over anhydrous Na₂SO₄ and then concentrated under reduced pressure. The residue was purified by silica gel column chromatography eluting with 10% ethyl acetate/petroleum ether to give the title compound (298 g, 87%) as a white solid.

Step 5. Ethyl furo[2,3-c]pyridine-2-carboxylate. To a solution of ethyl 3-(((trifluoromethyl)sulfonyl)oxy)furo[2,3-c]pyridine-2-carboxylate (298 g, 0.88 mol) in ethanol (3 L) was added 10% Pd/C (30 g) and triethylamine (281 mL, 2.02 mol). The reaction mixture was stirred under an atmosphere of hydrogen for 16 h and then filtered through a pad of diatomaceous earth. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography eluting with 20% ethyl acetate/petroleum ether to give the title compound (158 g, 94%) as a pale yellow solid.

Step 6. To a solution of ethyl furo[2,3-c]pyridine-2-carboxylate (158 g, 0.83 mol) in water:THF:MeOH (1:1:1, 2.4 L) was added KOH (139 g, 2.49 mol). The reaction mixture was stirred at rt for 16 h and then concentrated to a volume of 750 mL. To this residue was added acetic acid until pH ~ 4. The resulting solids were collected by filtration, washed with water (300 mL×2) and dried in a vacuum oven overnight to give the title compound (101 g, 75%) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.07 (s, 1H), 8.47 (d, *J* = 5.6 Hz, 1H), 7.80 (d, *J* = 5.2 Hz, 1H), 7.61 (s, 1H). MS (ESI+) *m/z*: 164 [M+H]⁺.

10 Intermediate 2: Imidazo[1,2-a]pyridine-6-carboxylic acid




Step 1. Imidazo[1,2-a]pyridine-6-carboxylic acid hydrochloride salt. A mixture of 2-chloroacetaldehyde (277 g, 40%) and 6-aminonicotinic acid (150 g) in ethanol (330 mL) was heated to reflux and stirred for 8 h. After cooling, a solid precipitated and was isolated by vacuum filtration, then washed with ethanol and dried under vacuum to give the title compound as a light yellow solid (1.78 g, 82%).

Step 2. Imidazo[1,2-a]pyridine-6-carboxylic acid hydrochloride salt (170 g) was diluted with water (600 mL) and heated until a clear solution resulted, then an aqueous solution of NaOH (2 M) was added slowly to adjust the pH = 5-6. The reaction mixture was cooled to 0 °C using an ice-H₂O bath. The resulting precipitate was collected by vacuum filtration, then washed with ethanol and dried under vacuum to give the title product (107.2 g, 77%) as a light yellow powder. ¹H NMR (400 MHz, DMSO-*d*₆) δ 13.76–12.82 (br, 1H), 9.28 (s, 1H), 8.10 (s, 1H), 7.68 (s, 1H), 7.64–7.56 (m, 2H). MS (ESI+) *m/z*: 163 [M+H]⁺.

25

Intermediate 3: Imidazo[1,2-a]pyrimidine-6-carboxylic acid




Step 1. Sodium (Z)-2-(dimethoxymethyl)-3-methoxy-3-oxoprop-1-en-1-olate. Methyl 3,3-dimethoxypropanoate (100 g, 675 mmol) and methyl formate (81 g, 1350 mmol) were dissolved in anhydrous THF (450 mL). Sodium hydride (60% dispersion; 32.4 g, 810 mmol, 1.2 eq.) was then added slowly in portions at 0 °C. The reaction mixture was stirred at rt for 1 h, then was heated at 50 °C for 3 h. During this period, H₂ evolution was observed. After cooling to rt, the solvent was then removed under reduced pressure to give the crude product which was directly used in the next step without further purification.

Step 2. Methyl 2-aminopyrimidine-5-carboxylate. The crude enolate from step 1 was dissolved in DMF (200 mL), and guanidine hydrochloride (64 g, 670 mmol) was added. The mixture was heated at 100 °C under N₂ for 3 h. After cooling to rt, water was added and the mixture was cooled with an ice-water bath. The resulting precipitate was collected by vacuum filtration and dried under vacuum to give the desired product (63 g, 61% yield for 2 steps).

Step 3. Methyl imidazo[1,2-a]pyrimidine-6-carboxylate. To a mixture of 2-bromo-1,1-diehoxyethane (100.6 g, 0.51 mol) and methyl 2-aminopyrimidine-5-carboxylate (63 g, 0.41 mol) in ethanol (300 mL) was added concentrated HBr (40%) (55 g). The reaction mixture was heated to reflux for 3 h under N₂. After cooling to rt, the mixture was further cooled with an ice-water bath. The resulting precipitate was collected by vacuum filtration and dried under vacuum overnight to give the desired product (92 g, 87%).

Step 4. Into a round bottom flask containing methyl imidazo[1,2-a]pyrimidine-6-carboxylate (92 g, 356.5 mmol), was added water (200 mL). NaOH (6 N in H₂O, 2.5 eq.) was then added dropwise with stirring at rt. After stirring at rt for 1 h, the mixture was cooled with an ice-water bath and concentrated HCl was added (pH = 5-6). The resulting mixture was concentrated under reduced pressure to approximately 150 mL (3/4 volume) and cooled with an ice-water bath. The resulting precipitate was collected by vacuum filtration, washed with cold water (50 mL) and dried to give the title compound as an off-white solid (46 g, 79%). ¹H NMR (DMSO-*d*₆, 400 MHz) δ 9.29 (d, *J* = 2.0 Hz, 1H), 8.89 (d, *J* = 2.0 Hz, 1H), 7.94 (s, 1H), 7.70 (s, 1H). MS (*m z*, ES⁺): 164.1 [M+H]⁺, 186.1 [M + Na]⁺.

Intermediate 4: 1H-Pyrazolo[3,4-b]pyridine-5-carboxylic acid

5 Step 1. 1-(4-Methoxybenzyl)-1H-pyrazol-5-amine. To a solution of acrylonitrile (30 mL, 455 mmol) in THF (250 mL), $\text{NH}_2\text{NH}_2\cdot\text{H}_2\text{O}$ (23.19 mL, 478 mmol) was added drop-wise at 0 °C. After addition was complete, the mixture was stirred at rt for 2 h, then 4-methoxybenzaldehyde (55.4 mL, 455 mmol) was added drop-wise. The mixture was stirred at rt overnight, then at reflux for 2 h. After cooling to rt the mixture was quenched by addition of 300 mL of ice water. The

10 mixture was extracted with ethyl acetate (3 x), then the combined organic layers were extracted with 1 N HCl. The aqueous layer was neutralized with aqueous 10 N NaOH solution, then extracted with ethyl acetate. The organic layer was washed with H_2O and brine, then dried over Na_2SO_4 . Filtration, concentration, and recrystallization with Et_2O gave the target compound as a white solid (50 g, 60%).

15 Step 2. Ethyl 4-hydroxy-1-(4-methoxybenzyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. 1-(4-Methoxybenzyl)-1H-pyrazol-5-amine (3.94 g, 19.39 mmol), followed by diethyl 2-(ethoxymethylene)malonate (4 mL, 20 mmol) was added to a 200 mL round bottom flask fitted with a distillation head to remove ethanol. The mixture was heated to 130 °C for 45 min, then 10 mL of diphenyl ether was added

20 and the temperature was raised to 240 °C for 2 h. The reaction mixture was then cooled to rt and diethyl ether (100 mL) was added. The resulting precipitate was collected by vacuum filtration and dried under vacuum to afford the target compound as a white solid (4 g, 62%).


25 Step 3. Ethyl 4-chloro-1-(4-methoxybenzyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate. POCl_3 (10 mL) was added to ethyl 4-hydroxy-1-(4-methoxybenzyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate (7.5 g, 19.39 mmol). The mixture was stirred at 60 °C for 3 h. The mixture was poured into ice water and the resulting precipitate was collected by vacuum filtration and dried under vacuum to afford the target compound a light yellow solid (6.4 g, 80%).

Step 4. Ethyl 1-(4-methoxybenzyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate.

To a solution of ethyl 4-chloro-1-(4-methoxybenzyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate (5.9 g, 17 mmol) in THF (50 mL), triethylamine (1.7 g, 17 mmol), followed by Pd(OH)₂/C (300 mg) was added. The mixture was stirred at rt for 3 h
5 under H₂. The mixture was filtered and concentrated. The residue was dissolved in ethyl acetate and washed with saturated aqueous NaHCO₃ solution and brine, then dried over Na₂SO₄. Filtration and concentration gave target compound as a light gray solid (5.3 g, 100%).

Step 5. Ethyl 1-(4-methoxybenzyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxylate
10 (4.4 g, 14 mmol) was dissolved in TFA (158 mL) and heated to 80 °C. The mixture was stirred at 80 °C for 4 h, then was concentrated to dryness. The residue was poured into ice water, then aqueous NaOH solution (2 M) was added until the pH was approximately 14. The solid formed was removed by filtration, and the aqueous layer was washed with ethyl acetate. To the aqueous layer was added concentrated HCl
15 was added until the pH was approximately 7. The resulting precipitate was collected by vacuum filtration and dried under vacuum to afford the title compound as a white solid (2.1 g, 80%). ¹H NMR (400 MHz, DMSO-*d*₆) δ 14.38-13.62 (br, 1H), 9.07 (d, *J* = 1.6 Hz, 1H), 8.81 (d, *J* = 1.6 Hz, 1H), 8.32 (s, 1H). MS (*m/z*, ESI⁺): 164 [M+H]⁺.

20 Intermediate 5: 1H-Pyrrolo[3,2-c]pyridine-2-carboxylic acid



Step 1. 3-Iodopyridin-4-amine. To a 2 L 3-necked flask was added a solution of 38 mL of concentrated sulfuric acid in 200 mL water. The solution was cooled with an ice-water bath, then 4-aminopyridine (200 g, 2.12 mol) and acetic acid (700
25 mL) were added in batches. The mixture was then heated to reflux. Iodine (189 g, 0.745 mol) and periodic acid dihydrate (97 g, 0.424 mol) were both equally divided into four parts. One batch of iodine was added and then one batch of periodic acid dihydrate was added 15 min later. After 30 min, a new batch of iodine and periodic acid dihydrate were added in the same way. When all four batches of iodine and
30 periodic acid dihydrate were added, the mixture was kept refluxing for an additional

3 h. After cooling to rt the reaction mixture was slowly poured into water while stirring, then a 40% solution of NaOH in water was added until pH > 9. Na₂SO₃ was added to destroy the unreacted iodine. After cooling to rt, a filtration was performed. The collected solid was further purified by recrystallization in chloroform to give the
5 desired product (184 g, 39%).


Step 2. To a 2 L 3-necked flask was added DMF (700 mL), triethylene diamine (168 g, 1.5 mol), and 4-amino-3-iodopyridine (24, 110 g, 0.5 mol). The mixture was cooled with an ice-water bath and pyruvic acid (132 g, 1.5 mol) was slowly added, followed by palladium acetate (4.49 g, 0.02 mol). Under nitrogen
10 atmosphere, the mixture was heated to 115 °C. The reaction generated effervescence. The reaction mixture was kept at 115-120 °C for 11 h. The mixture was concentrated under reduced pressure. The residue was poured into water (500 mL), and concentrated HCl was added to adjust pH to <1. The mixture was cooled by adding ice and a filtration was performed. The cake thus obtained was a brownish black
15 solid.

The above cake was added into 500 mL of water. Concentrated HCl was added (to ensure complete protonation) followed by 5 g of active carbon. The mixture was heated to reflux for 20 min and then filtration was performed while hot. The solid was discarded and the hot filtrate was placed in a refrigerator to allow the
20 HCl salt of the desired product to precipitate. Upon cooling, filtration was performed which afforded a dark brown solid with a wet weight of 48 g as the HCl salt of the desired product.

The solid was then added to 250 mL of water and the mixture was heated until a clear solution resulted. Solid NaOH was slowly added to adjust pH to 5-6, then
25 active carbon and an additional 500 mL of water was added. The mixture was heated to reflux for 30 min, then filtration was performed while hot. The resulting cake was added to 750 mL of water, heated to reflux, and filtered again. The cake thus obtained was discarded. The two batches of filtrate were combined and cooled in a refrigerator. The resulting precipitate was collected by vacuum filtration, then
30 washed with ethanol to give the title compound as a slightly yellow solid (25 g, 31%).
MS (*m z*, ES⁺): 161.1 [M-1], 323.1 [2M-1]. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 12.20

(br s, 1H), 8.97 (s, 1H), 8.27 (d, $J = 5.6$ Hz, 1H), 7.41 (d, $J = 6.0$ Hz, 1H), 7.23 (s, 1H).

Intermediate 6: Thieno[2,3-c]pyridine-2-carboxylic acid



5


Step 1. 3,5-Dibromoisonicotinaldehyde. Lithium diisopropylamide (507 mmol, 1.2 eq.) was added to 200 mL of dry THF at -78 °C under N_2 . A solution of 3,5-dibromopyridine (100 g, 424 mmol) in 537 mL of dry THF was then added drop-wise over 30 min. The reaction mixture was stirred at -78 °C for 1 h. Ethyl formate (34.4 g, 465 mmol) was added drop-wise and stirred at -78 °C for 30 min, then the reaction mixture was poured into ice-cold saturated aqueous $NaHCO_3$ solution. The mixture was extracted with 3 x 500 mL of ethyl acetate. The organic layer was concentrated to provide a brown solid, which was filtered through a pad of silica gel (eluted with dichloromethane) to give the title compound as a yellow powder (70 g, 15 63%).

Step 2: Methyl 4-bromothieno[2,3-c]pyridine-2-carboxylate. 3,5-Dibromoisonicotinaldehyde (80 g, 303 mmol), followed by cesium carbonate (98 g, 302 mmol) was added to a 2 L round bottom flask containing THF (1.3 L) under N_2 . Methyl mercaptoacetate (32 g, 302 mmol) was added and the mixture was heated at 20 60 °C overnight. After cooling to rt, ethyl acetate was added and the organic layer was washed with water, aqueous saturated $NaHCO_3$ solution, and brine, then dried over Na_2SO_4 and filtered to give a white solid. The crude product was purified by recrystallization from ethyl acetate to give the desired product (60 g, 73%).

Step 3. Methyl thieno[2,3-c]pyridine-2-carboxylate. Methyl 4-bromothieno[2,3-c]pyridine-2-carboxylate (115 g, 423 mmol), triethylamine (42.7 g, 423 mmol), THF (1.5 L), and MeOH (500 mL) were mixed and degassed. Under nitrogen, palladium on carbon (10%, 14.7 g, 13.9 mmol) was added. The mixture was hydrogenated with a Parr apparatus at 45 psi H_2 for 3 days. The catalyst was filtered off and the filtrate was concentrated to give the desired compound as a white 25 30 solid (65 g, 80%).

Step 4. A three necked 2 L round bottom flask equipped with an overhead stirrer and thermocouple was charged with methyl thieno[2,3-c]pyridine-2-carboxylate (130 g, 674 mmol) and water (650 mL). Aqueous sodium hydroxide solution (10 N) was added with stirring at 20 °C. Over the next 20 min, the temperature rose to 25 °C and the solid dissolved. After 1 h, concentrated HCl (1.5 eq.) was slowly added to the reaction mixture with rapid stirring, generating a thick slurry. After stirring for 1 h, the slurry was filtered and the solid was dried under vacuum to give the title compound as a white solid (105.5 g, 88%). MS (*m/z*, ES⁻): 178.0 [M-1]. ¹H-NMR (DMSO-*d*₆, 400 MHz) δ 12.24 (br s, 1H), 8.97 (s, 1H), 8.27 (d, *J* = 6.0 Hz, 1H), 7.40 (d, *J* = 5.6 Hz, 1H), 7.23 (s, 1H).

Intermediate 7: Imidazo[1,2-b]pyridazine-6-carboxylic acid




Step 1. 6-Chloro-imidazo[1,2-b]pyridazine. A solution of 6-chloro-1,2-diazinan-3-amine (10 g, 73.75 mmol, 1.00 equiv), 2-bromo-1,1-dimethoxyethane (50 g, 295.83 mmol, 4.01 equiv), and HBr (40%, 45 mL) in ethanol (100 mL) was stirred overnight at 90 °C. The majority of the ethanol was removed under reduced pressure then the pH value of the solution was adjusted to 10 with 5% aqueous potassium carbonate solution. The resulting mixture was extracted with 6x500 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1/2~1/1) to give 6.5 g (57%) of the title compound as a yellow solid. ¹H NMR (300 MHz, CDCl₃) δ 7.95 (s, 1H), 7.91 (s, 1H), 7.80 (s, 1H), 7.05 (d, *J* = 9.3 Hz, 1H).

Step 2. Imidazo[1,2-b]pyridazine-6-carboxylic acid methyl ester. A mixture of 6-chloro-imidazo[1,2-b]pyridazine (200 mg, 1.30 mmol, 1.00 equiv), bis(triphenylphosphine)palladium(II) dichloride (200 mg, 0.28 mmol, 0.22 equiv), and triethylamine (0.5 mL) in methanol (4 mL) was stirred under carbon monoxide (10 atm) in a 50-mL pressure reactor overnight at 110 °C. The solid material was

removed by filtration. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1/1) to give 100 mg (43%) of the title compound as a yellow solid. ¹H NMR (300 MHz, CDCl₃) δ 8.16 (s, 1H), 8.08 (d, *J* = 9.6 Hz, 1H), 7.94 (s, 1H), 7.77 (d, *J* = 9.6 Hz, 1H), 4.09 (s, 3H).

Step 3. A mixture of imidazo[1,2-*b*]pyridazine-6-carboxylic acid methyl ester (900 mg, 5.08 mmol, 1.00 equiv) and 5% aqueous sodium hydroxide solution (15 mL, 3.75 equiv) in THF (3 mL) was stirred overnight at rt. The pH value of the solution was adjusted to 2 with 1 M HCl. The resulting mixture was concentrated under vacuum to give 3 g of crude title product as a yellow solid. The crude product was used without further purification. LC/MS (Method A, ESI): RT= 0.43 min, *m/z* = 164.0 [M+H]⁺.

Intermediate 8: Pyrazolo[1,5-*a*]pyridine-5-carboxylic acid



Step 1. 1-Amino-4-methoxypyridinium iodide. A solution of aminooxysulfonic acid (11.4 g, 100.80 mmol, 0.50 equiv) and 4-methoxypyridine (22 g, 201.60 mmol, 1.00 equiv) in water (200 mL) was stirred under nitrogen for 0.5 h at 90 °C. Potassium carbonate (14 g, 101.30 mmol, 0.50 equiv) was added at rt. The resulting mixture was concentrated under vacuum then ethanol (150 mL) was added to dissolve the residue. The insoluble material was removed by filtration. The filtrate was cooled to -20 °C and then hydroiodic acid (16 g, 40%) was added. The resulting solution was stirred for 1 h at -20 °C. The precipitated product was collected by filtration and washed with cold ethanol to give 9.3 g (46%) of the title compound as a white solid. TLC: 1:5 MeOH/DCM, R_f = 0.02.

Step 2. 5-Methoxy-pyrazolo[1,5-*a*]pyridine-3-carboxylic acid methyl ester. A mixture of 1-amino-4-methoxypyridinium iodide (6 g, 23.80 mmol, 1.00 equiv), potassium carbonate (5 g, 36.18 mmol, 1.50 equiv), and methyl propiolate (2 g, 23.79 mmol, 1.00 equiv) in DMF (50 mL) was stirred under nitrogen for 4 h at rt. After the reaction completed, the mixture was concentrated under vacuum. The residue was

dissolved in 150 mL of dichloromethane and then washed with 1x20 mL of saturated aqueous sodium bicarbonate solution. The organic layer was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/hexane (1:3) to give 1.5 g (31%) of title product as a solid. LC/MS (Method D, ESI): RT= 1.30 min, $m/z = 207.0$ [M+H]⁺.

5 Step 3. Pyrazolo[1,5-a]pyridin-5-ol. A mixture of methyl 5-methoxypyrazolo[1,5-a]pyridine-3-carboxylate (100 mg, 0.48 mmol, 1.00 equiv) in 40% HBr (5 mL) was stirred for 16 h at 100 °C. The reaction mixture was cooled to rt and the pH value of the solution was adjusted to 8 with 5 M potassium hydroxide
10 solution. The resulting solution was extracted with 2x50 mL of ether. The organic layers were combined and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:3 to 1:1) to yield 20 mg (31%) of the title compound as a white solid. LC/MS (Method D, ESI): RT= 0.41 min, $m/z = 135.0$ [M+H]⁺.


15 Step 4. Trifluoro-methanesulfonic acid pyrazolo[1,5-a]pyridin-5-yl ester. A mixture of pyrazolo[1,5-a]pyridin-5-ol (300 mg, 2.24 mmol, 1.00 equiv) and trifluoromethanesulfonic anhydride (0.5 mL) in pyridine (5 mL) was stirred for 10 h at rt. The resulting mixture was concentrated under vacuum and the residue was dissolved in 100 mL of dichloromethane. The mixture was washed with 1x10 mL of
20 sodium bicarbonate solution. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:3) to yield 200 mg (34%) of the title compound as a solid. LC/MS (Method B, ESI): RT= 2.13 min, $m/z = 267.0$ [M+H]⁺.

25 Step 5. Pyrazolo[1,5-a]pyridine-5-carboxylic acid methyl ester. A mixture of trifluoro-methanesulfonic acid pyrazolo[1,5-a]pyridin-5-yl ester (200 mg, 0.75 mmol, 1.00 equiv), triethylamine (227 mg, 2.24 mmol, 3.00 equiv), DMSO (98 mg, 1.25 mmol, 1.67 equiv), and bis(triphenylphosphine)palladium(II) dichloride (53 mg, 0.08 mmol, 0.10 equiv) in methanol (20 mL) was stirred under carbon monoxide (10 atm)
30 for 16 h at 100 °C in a 50-mL pressure reactor. After the reaction completed, the reaction mixture was cooled to rt and the mixture was concentrated under vacuum.

The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:3) to afford 130 mg of the title compound as a solid. LC/MS (Method H, ESI): RT= 1.36 min, $m/z = 177.0$ [M+H]⁺.

Step 6. A mixture of pyrazolo[1,5-a]pyridine-5-carboxylic acid methyl ester (130 mg, 0.74 mmol, 1.00 equiv) and potassium hydroxide (1 g, 17.82 mmol, 24.15 equiv) in methanol (2 mL), THF (2 mL), and water (5 mL) was stirred for 12 h at rt. The reaction mixture was washed with 2x50 mL of ethyl acetate. The aqueous layer was collected and the pH value of the solution was adjusted to 6 with 1 N HCl. The solution was extracted with 5x50 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to give 100 mg (84%) the title compound as a yellow solid. LC/MS (Method G, ESI): RT= 1.32 min, $m/z = 163.0$ [M+H]⁺.

Intermediate 9: 1H-Pyrazolo[4,3-b]pyridine-6-carboxylic acid



Step 1. 5-Bromo-2-methyl-pyridin-3-ylamine. To a stirred mixture of iron filings (5 g, 89.29 mmol, 3.88 equiv) and ammonium chloride (1 g, 18.70 mmol, 0.81 equiv) in ethanol (66 mL) and water (33 mL) was added a solution of 5-bromo-2-methyl-3-nitropyridine (5 g, 23.04 mmol, 1.00 equiv) in ethanol (50 mL) dropwise at 90 °C. The reaction mixture was stirred for 10 min at 90 °C and then cooled to rt. The solid material was removed by filtration. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:2) to yield 1.6 g (37%) of the title compound as a yellow solid. LC/MS (Method I, ESI): RT= 0.81 min, $m/z = 187.0; 189.0$ [M+H]⁺.

Step 2. N-(5-Bromo-2-methyl-pyridin-3-yl)-acetamide. A solution of 5-bromo-2-methyl-pyridin-3-ylamine (3 g, 16.04 mmol, 1.00 equiv) in acetic anhydride (20 mL) and acetic acid (10 mL) was stirred overnight at rt. The resulting mixture was concentrated under vacuum to give 2.6 g (71%) of the title compound as a light yellow solid. LC/MS (Method I, ESI): RT= 1.05 min, $m/z = 229.0; 231.0$ [M+H]⁺.

Step 3. 1-(6-Bromo-pyrazolo[4,3-b]pyridin-1-yl)-ethanone. A mixture of N-(5-bromo-2-methyl-pyridin-3-yl)-acetamide (3.5 g, 15.28 mmol, 1.00 equiv), isopentyl nitrite (4 g, 34.73 mmol, 2.27 equiv), potassium acetate (20 g), and acetic anhydride (30 mL) in toluene (150 mL) was stirred under nitrogen overnight at 90 °C.

5 The reaction mixture was cooled to rt and the solid material was removed by filtration. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:5) to give in 2 g (55%) of the title compound as a light yellow solid. LC/MS (Method I, ESI): RT= 1.44 min, m/z = 240.0; 242.0 [M+H]⁺.


10 Step 4. 1H-Pyrazolo[4,3-b]pyridine-6-carboxylic acid methyl ester. A mixture of 1-(6-bromo-pyrazolo[4,3-b]pyridin-1-yl)-ethanone (2 g, 8.33 mmol, 1.00 equiv), bis(triphenylphosphine)palladium(II) dichloride (1 g, 1.42 mmol, 0.17 equiv), and triethylamine (2.5 mL) in methanol (70 mL) was stirred overnight under carbon monoxide (10 atmospheres) at 100 °C in a 100 mL pressure reactor. The reaction

15 mixture was cooled to rt and the solid material was removed by filtration. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:5) to afford 0.8 g (54%) of the title compound as a light yellow solid. TLC: 1:1 ethyl acetate/petroleum ether, R_f = 0.2.

Step 5. A solution 1H-pyrazolo[4,3-b]pyridine-6-carboxylic acid methyl ester (200 mg, 1.13 mmol, 1.00 equiv) and sodium hydroxide (200 mg, 5.00 mmol, 4.43 equiv) in water (10 mL) was stirred overnight at rt. After the reaction was complete, the pH value of the solution was adjusted to 3 with concentrated HCl. The resulting mixture was concentrated under vacuum to give 1 g of crude title product as a light yellow solid. LC/MS (Method I, ESI): RT= 0.91 min, m/z = 164.0; 242.0 [M+H]⁺.

25

Intermediate 10: [1,2,4]Triazolo[1,5-a]pyridine-6-carboxylic acid



Step 1. N'-(5-Bromo-pyridin-2-yl)-N,N-dimethyl-formamidine. A solution of 5-bromopyridin-2-amine (4 g, 23.12 mmol, 1.00 equiv) and N,N-dimethylformamide dimethyl acetal (9.6 mL, 3.00 equiv) in DMF (30 mL) was stirred under nitrogen for

30

12 h at 130 °C. The reaction mixture was cooled to rt and then concentrated under vacuum to give 4 g (76%) of the title compound as an oil. TLC: 1:5 MeOH/DCM, $R_f = 0.6$.


Step 2. 6-Bromo-[1,2,4]triazolo[1,5-a]pyridine. To a solution of N'-(5-bromo-pyridin-2-yl)-N,N-dimethyl-formamide (4 g, 17.54 mmol, 1.00 equiv) in methanol (40 mL) maintained under nitrogen at 0 °C was added pyridine (4 mL, 2.00 equiv) and (aminoxy)sulfonic acid (3.6 g, 31.83 mmol, 1.30 equiv). The resulting solution was stirred for 12 h at rt. After the reaction completed, the mixture was concentrated under vacuum. The residue was diluted with 150 mL of ethyl acetate then washed with 1x50 mL of saturated aqueous sodium carbonate solution and 2x50 mL of water. The organic layer was dried over anhydrous sodium sulfate then concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/hexane (1:1) to give 2.5 g (72%) title compound as a solid. LC/MS (Method D, ESI): RT= 1.15 min, $m/z = 198.0 [M+H]^+$.

Step 3. [1,2,4]Triazolo[1,5-a]pyridine-6-carboxylic acid methyl ester. A mixture of 6-bromo-[1,2,4]triazolo[1,5-a]pyridine (2.4 g, 12.12 mmol, 1.00 equiv), bis(triphenylphosphine)palladium(II) dichloride (800 mg, 1.14 mmol, 0.10 equiv) and triethylamine (4 g, 39.53 mmol, 3.00 equiv) in DMSO (1.6 g, 20.48 mmol, 1.67 equiv) and methanol (50 mL) was stirred under carbon monoxide (10 atm) for 20 h at 100 °C. The reaction mixture was cooled to rt and quenched with brine (50 mL). The resulting solution was extracted with ethyl acetate (3x40 mL). The combined organic layers were dried over anhydrous sodium sulfate then concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/hexane (1:1) to give 0.98 g (46%) of the title compound as a crude solid. LC/MS (Method C, ESI): RT= 1.04 min, $m/z = 178.0 [M+H]^+$.

Step 4. A solution of [1,2,4]triazolo[1,5-a]pyridine-6-carboxylic acid methyl ester (200 mg, 1.13 mmol, 1.00 equiv) in THF (2 mL) was added to a solution of potassium hydroxide (1 g, 17.82 mmol, 15.79 equiv) in water (10 mL). The resulting mixture was stirred for 10 h at rt. After the reaction completed, the pH value of the solution was adjusted to 5-6 with 1 N HCl. The mixture was extracted with 3x50 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium

sulfate and concentrated under vacuum to give 112 mg (61%) of the title compound as a solid. LC/MS (Method C, ESI): RT = 0.9 min, $m/z = 164.0$ [M+H]⁺.

Intermediate 11: Pyrazolo[1,5-a]pyrimidine-5-carboxylic acid



5

Step 1. 4H-Pyrazolo[1,5-a]pyrimidin-5-one. A solution of 1H-pyrazol-3-ylamine (7 g, 84.24 mmol, 1.00 equiv) and ethyl prop-2-ynoate (50 mL) in dioxane (10 g, 1.21 equiv) was stirred under nitrogen overnight at 110 °C. The reaction mixture was cooled to rt and the precipitated product was collected by filtration to give 4 g (36%) of the title compound as a light brown solid. ¹H NMR (300 MHz, DMSO-*d*₆) δ 12.04 (s, 1H), 8.41-8.44 (m, 1H), 7.71 (d, *J* = 1.8 Hz, 1H), 5.88 (d, *J* = 8.1 Hz, 1H), 5.77 (m, 1H).

10

Step 2. 5-Chloro-pyrazolo[1,5-a]pyrimidine. A solution of 4H-pyrazolo[1,5-a]pyrimidin-5-one (1 g, 7.40 mmol, 1.00 equiv) in phosphorus oxychloride (15 mL) was stirred under nitrogen for 2 h at 120 °C. The reaction mixture was cooled to rt then concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:2) to give 0.6 g (53%) of the title compound as a light yellow solid. LC/MS (Method I, ESI): RT = 1.21 min, $m/z = 154.0$ [M+H]⁺.


15

Step 3. Pyrazolo[1,5-a]pyrimidine-5-carboxylic acid methyl ester. A mixture of 5-chloro-pyrazolo[1,5-a]pyrimidine (2 g, 13.02 mmol, 1.00 equiv), triethylamine (4 mL), methanol (80 mL), and bis(triphenylphosphine)palladium(II) dichloride (1 g, 1.42 mmol, 0.11 equiv) was stirred in a 100-mL pressure reactor overnight at 100 °C under 10 atmospheres of carbon monoxide. The reaction mixture was cooled to rt then concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:5) to yield 1.2 g (52%) of the title compound as a light yellow solid. LC/MS (Method I, ESI): RT = 1.09 min, $m/z = 178.0$ [M+H]⁺.

20
25

Step 4. To a solution of methyl pyrazolo[1,5-a]pyrimidine-5-carboxylic acid methyl ester (100 mg, 0.56 mmol, 1.00 equiv) in acetic acid (5 mL) was added concentrated HCl (37%, 5 mL). The resulting solution was stirred for 3 h at 120 °C, then concentrated under vacuum. The residue was dissolved in 3 mL of water and then adjusted to pH 5 with saturated aqueous sodium carbonate solution. The precipitated product was collected by filtration then air-dried to give 0.08 g (87%) of pyrazolo[1,5-a]pyrimidine-5-carboxylic acid as a light yellow solid. LC/MS (Method 1, ESI): RT= 0.95 min, $m/z = 164.0$ [M+H]⁺.

10 Intermediate 12: 3-tert-Butylamino-imidazo[1,2-a]pyridine-6-carboxylic acid




Step 1. 3-tert-Butylamino-imidazo[1,2-a]pyridine-6-carboxylic acid methyl ester. To a solution of methyl 6-aminopyridine-3-carboxylate (3.8 g, 24.98 mmol, 1.00 equiv) and 2-oxoacetic acid hydrate (3.9 g, 42.39 mmol, 1.70 equiv) in methanol (120 mL) was added perchloric acid (250 mg, 2.50 mmol, 0.10 equiv). The reaction mixture was stirred for 30 min and 2-isocyno-2-methylpropane (2.08 g, 25.02 mmol, 1.00 equiv) was then added. The reaction mixture was stirred for 12 h at rt and then concentrated under vacuum. The residue was purified on a silica gel column eluted with dichloromethane/ethyl acetate (2:1) to give 850 mg (14%) of the title compound as a yellow solid. ¹H NMR (300 MHz, CDCl₃) δ 8.97-8.96 (dd, $J = 0.9, 1.5$ Hz, 1H), 7.69-7.65 (dd, $J = 4.2, 9.6$ Hz, 1H), 7.53-7.50 (dd, $J = 4.2, 9.6$ Hz, 1H), 7.39 (s, 1H), 3.96 (s, 3H), 1.23 (s, 9H).

Step 2. Sodium 3-tert-Butylamino-imidazo[1,2-a]pyridine-6-carboxylate. To a solution of 3-tert-butylamino-imidazo[1,2-a]pyridine-6-carboxylic acid methyl ester (300 mg, 1.21 mmol, 1.00 equiv) in methanol (5 mL) was added a solution of sodium hydroxide (97 mg, 2.42 mmol, 2.00 equiv) in water (5 mL). The resulting solution was stirred for 1.5 h at 46 °C. The reaction mixture was cooled to rt and then quenched by the addition of 0.15 mL of HCl. The resulting mixture was concentrated under vacuum to give 345.6 mg (crude) of the title product as a yellow solid. LC/MS

(Method I, ESI): RT = 1.02 min, $m/z = 234.0$ [M+H - 22].

Step 3. Sodium 3-tert-butylamino-imidazo[1,2-a]pyridine-6-carboxylate (300 mg, 1.17 mmol, 1.00 equiv) was dissolved in acetic acid (10 mL) and then concentrated under vacuum. The residue was purified on a silica gel column eluted with dichloromethane/methanol (20:1) to give 150 mg (54%) of the title compound as a yellow solid. LC/MS (Method N, ESI): RT = 0.94 min, $m/z = 234.0$ [M+H].

Intermediate 13: 2,3-Dihydro-1H-pyrrolo[3,4-c]pyridine.



Step 1. Ethyl N-(prop-2-yn-1-yl)carbamate. To a solution of prop-2-yn-1-amine (11.5 g, 208.79 mmol, 1.00 equiv) and sodium hydroxide (9.1 g, 227.50 mmol, 1.09 equiv) in water (40 mL) and toluene (110 mL) maintained under nitrogen was added ethyl chloroformate (23.9 g, 220.23 mmol, 1.05 equiv) dropwise in 20 min with stirring at 10 °C. The resulting solution was stirred overnight at rt then extracted with 3x100 mL of toluene. The combined organic layers were dried over anhydrous sodium sulfate then concentrated under vacuum to give 15 g (57%) of ethyl N-(prop-2-yn-1-yl)carbamate as a light yellow oil. TLC: ethyl acetate/petroleum ether (1:2), $R_f = 0.5$.

Step 2. Pyrimidine-5-carboxaldehyde. To a solution of 5-bromopyrimidine (2 g, 12.58 mmol, 1.00 equiv) in THF (20 mL) placed in a 50-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen was added n-butyllithium (1.1 mL) at -78 °C. The reaction mixture was stirred at -78 °C for another 2 h. Ethyl formate (5.2 mL) was then added and the resulting solution was stirred for 2 h at -78 °C. The resulting mixture was warmed to 0 °C and washed with 50 mL of brine. The organic layer was dried with anhydrous sodium carbonate and concentrated. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:1) to give 11 g of crude pyrimidine-5-carboxaldehyde as a yellow oil. TLC: ethyl acetate/petroleum ether (1/1), $R_f = 0.2$.

Step 3. Pyrimidin-5-ylmethanol. A mixture of pyrimidine-5-carboxaldehyde (2 g, 18.50 mmol, 1.00 equiv) and sodium borohydride (2 g) in methanol (100 mL)

was stirred at 0 - 10°C for 30 min. The reaction mixture was concentrated under vacuum and the residue was purified on a silica gel column eluted with dichloromethane/methanol (50:1) to yield 1.2 g (59%) of pyrimidin-5-ylmethanol as a light yellow solid. LC/MS (Method N, ESI): RT= 0.74 min, $m/z = 111.0$ [M+H]⁺.

5 Step 4. 5-(Chloromethyl)pyrimidine. To a solution of pyrimidin-5-ylmethanol (1.1 g, 10 mmol, 1.00 equiv) in dichloromethane (30 mL) was added thionyl chloride (2 mL) dropwise with stirring. The resulting solution was stirred at rt for 2 h then concentrated in vacuum to give 1.1 g of crude 5-(chloromethyl)pyrimidine as a yellow oil. TLC: ethyl acetate/petroleum ether (1:1),
10 $R_f = 0.4$.


Step 5. Ethyl N-(prop-2-yn-1-yl)-N-(pyrimidin-5-ylmethyl)carbamate. A mixture of ethyl N-(prop-2-yn-1-yl)carbamate (1.27 g, 9.99 mmol, 1.00 equiv) benzyltriethylammonium chloride (500 mg, 2.60 mmol, 0.26 equiv), 5-(chloromethyl)pyrimidine (1.28 g, 9.96 mmol, 1.00 equiv) and potassium hydroxide
15 (3 g, 53.47 mmol, 5.37 equiv) in toluene (30 mL) was stirred overnight under nitrogen at rt. The resulting mixture was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:1) to afford 0.3 g (14%) of ethyl N-(prop-2-yn-1-yl)-N-(pyrimidin-5-ylmethyl)carbamate as a light yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 9.16 (s, 1H), 8.73 (s, 2H), 4.59 (s,
20 2H), 4.11-4.26 (m, 4H), 2.28 (t, $J = 2.4$ Hz, 1H), 1.30 (t, $J = 7.2$ Hz, 3H).

Step 6. Ethyl 1H,2H,3H-Pyrrolo[3,4-c]pyridine-2-carboxylate. A mixture of ethyl N-(prop-2-yn-1-yl)-N-(pyrimidin-5-ylmethyl)carbamate (1 g, 4.56 mmol, 1.00 equiv) in xylene (30 mL) was stirred under nitrogen at 150 °C for 2 days. The resulting mixture was concentrated under vacuum and the residue was purified on a
25 silica gel column eluted with ethyl acetate/petroleum ether (1/2) to give 0.4 g (46%) of ethyl 1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxylate as a light brown crude solid. ¹H NMR (300 MHz, CDCl₃) δ 8.53-8.93 (m, 2H), 7.24 (d, $J = 5.1$ Hz, 1H), 4.73-4.80 (m, 4H), 4.22-4.33 (m, 2H), 1.33-1.49 (m, 3H).

Step 7. 2,3-Dihydro-1H-pyrrolo[3,4-c]pyridine. A mixture of ethyl
30 1H,2H,3H-pyrrolo[3,4-c]pyridine-2(3H)-carboxylate (400 mg, 2.4 mmol, 1.00 equiv) and barium hydroxide (0.8 g) in water (100 mL) was stirred overnight at 120 °C. The

reaction mixture was cooled to rt and the solid material was collected by filtration. The residue was stirred in hot ethyl acetate (150 mL) and then filtered to remove solid material. The filtrate was concentrated under vacuum to give 0.18 g (72%) of 2,3-dihydro-1H-pyrrolo[3,4-c]pyridine as a light yellow oil. ¹H NMR (300 MHz, CDCl₃) δ 8.51 (s, 1H), 8.41-8.45 (t, *J* = 4.8 Hz, 1H), 7.13-7.20 (m, 1H), 4.25 (s, 2H), 4.22 (s, 2H).

Intermediate 14. 2H,4H,5H,6H-Pyrrolo[3,4-c]pyrazole hydrochloride.




10 Step 1. tert-Butyl 2H,4H,5H,6H-pyrrolo[3,4-c]pyrazole-5-carboxylate. A solution of tert-butyl 3-[(dimethylamino)methylidene]-4-oxopyrrolidine-1-carboxylate (1 g, 4.16 mmol, 1.00 equiv) and hydrazine hydrate (340 mg, 6.79 mmol, 1.63 equiv) in ethanol (10 mL) was stirred at rt for 5 h. The resulting mixture was concentrated under vacuum and the residue was purified on a silica gel column eluted
15 with ethyl acetate/petroleum ether (1:5 to 1:2) to give 250 mg (29%) of tert-butyl 2H,4H,5H,6H-pyrrolo[3,4-c]pyrazole-5-carboxylate as a yellow solid. LC/MS (Method C, ESI): RT = 1.30 min, *m/z* = 210.0 [M + H]⁺.

20 Step 2. A solution of tert-butyl 2H,4H,5H,6H-pyrrolo[3,4-c]pyrazole-5-carboxylate (250 mg, 1.19 mmol, 1.00 equiv) in DCM (5 mL) and TFA (5 mL) was stirred overnight at rt. The resulting mixture was concentrated under vacuum and the residue was dissolved in 20 mL of concentrated HCl. The resulting mixture was concentrated under vacuum to give 200 mg of crude 2H,4H,5H,6H-pyrrolo[3,4-c]pyrazole hydrochloride as a dark red solid. LC/MS (Method C, ESI): RT = 0.46 min, *m/z* = 110.0 [M + H]⁺.

25

Intermediate 15. Lithium pyrazolo[1,5-b]pyridazine-5-carboxylate.



Step 1. 4-Methoxypyridazine. A mixture of 3,6-dichloro-4-methoxypyridazine (30 g, 167.59 mmol, 1.00 equiv), ammonium formate (31 g, 491.63 mmol, 2.93 equiv) and 10% palladium on carbon (3 g) catalyst in MeOH (500 mL) was stirred under 1 atmosphere of hydrogen at rt overnight. The catalyst was removed by filtration and the filtrate was concentrated under vacuum. The residue was dissolved in 500 mL of DCM/MeOH (10:1). The solid material was removed by filtration. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (2:1 to 1:1) to give 15 g (81%) of 4-methoxypyridazine as a brown oil. TLC: petroleum ether:ethyl acetate =2:1, $R_f = 0.1$.

Step 2. Methyl 5-methoxypyrazolo[1,5-b]pyridazine-3-carboxylate. To a stirred solution of hydroxylamine-O-sulfonic acid (30.8 g, 272.57 mmol, 1.50 equiv) in water (100 mL) maintained under nitrogen at 5°C was added a solution of potassium bicarbonate (29.1 g, 290.67 mmol, 1.60 equiv) in water (100 mL) dropwise. The resulting mixture was stirred for 10 min then a solution of 4-methoxypyridazine (20 g, 181.63 mmol, 1.00 equiv) in water (100 mL) was added. The reaction mixture was stirred at 70°C for 5 h and then cooled back to rt. A solution of methyl prop-2-ynoate (16.8 g, 199.83 mmol, 1.10 equiv) in DCM (500 mL) followed by a solution of potassium hydroxide (17.3 g, 308.32 mmol, 1.70 equiv) in water (100 mL) were added to the reaction mixture. The resulting solution was stirred overnight at rt then extracted with 4000 mL of DCM. The organic layer was washed with 3x500 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (6:1 to 2:1) to give 1 g (3%) of methyl 5-methoxypyrazolo[1,5-b]pyridazine-3-carboxylate as a brown solid. TLC: petroleum ether:ethyl acetate=1:1, $R_f=0.4$.

Step 3. Pyrazolo[1,5-b]pyridazin-3-ol. A solution of methyl 5-methoxypyrazolo[1,5-b]pyridazine-3-carboxylate (2 g, 9.65 mmol, 1.00 equiv) in 40% hydrobromic acid (50 mL) was refluxed overnight. The resulting mixture was cooled to rt then concentrated under vacuum. The residue was dissolved in 50 mL of H₂O and the pH value of the solution was adjusted to 6-7 with saturated aqueous KHCO₃ solution (50mL). The mixture was concentrated and the residue was purified


on a silica gel column eluted with DCM:MeOH (10:1) to give 500 mg (38%) of pyrazolo[1,5-b]pyridazin-3-ol as a brown solid. LC/MS (Method D, ESI): RT = 0.41 min, $m/z = 136.0$ [M + H]⁺.

Step 4. Pyrazolo[1,5-b]pyridazin-3-yl trifluoromethanesulfonate. To a solution of pyrazolo[1,5-b]pyridazin-3-ol (450 mg, 3.33 mmol, 1.00 equiv) and pyridine (1.1 g, 13.91 mmol, 3.00 equiv) in DCM (20 mL) maintained under nitrogen at -5°C was added trifluoromethanesulfonic anhydride (1.9 g, 6.73 mmol, 2.02 equiv) dropwise with stirring in 20 min. The resulting solution was stirred for another 60 min at 0°C and then diluted with 200 mL of DCM. The mixture was washed with 3x100 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:10 to 1:5) to give 570 mg (64%) of pyrazolo[1,5-b]pyridazin-3-yl trifluoromethanesulfonate as a white solid. TLC: petroleum ether:ethyl acetate = 1:1, $R_f=0.2$.

Step 5. Methyl pyrazolo[1,5-b]pyridazine-3-carboxylate. A mixture of 4aH,5H-pyrazolo[1,5-b]pyridazin-3-yl trifluoromethanesulfonate (600 mg, 2.23 mmol, 1.00 equiv), Pd(PPh₃)₂Cl₂ (300 mg, 0.43 mmol, 0.19 equiv) and pyridine (530 mg, 6.70 mmol, 3.01 equiv) in DMF (20 mL) and MeOH (5 mL) was stirred under 10 atmosphere of CO in a 50-mL pressure tank reactor overnight at 80°C. The reaction mixture was cooled to rt then diluted with 100 mL of H₂O. The resulting solution was extracted with 3x100 mL of ethyl acetate. The combined organic layers were washed with 3x100 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:5) to yield 200 mg (51%) of methyl pyrazolo[1,5-b]pyridazine-3-carboxylate as a yellow solid. TLC: petroleum ether:ethyl acetate = 2:1, $R_f=0.4$.

Step 6. To a solution of methyl pyrazolo[1,5-b]pyridazine-3-carboxylate (60 mg, 0.34 mmol, 1.00 equiv) in THF (5 mL) was added a solution of LiOH (40 mg, 1.667 mmol, 1.7 equiv) in water (1 mL). The reaction mixture was stirred overnight at 50°C and then concentrated under vacuum to give 150 mg of crude lithium pyrazolo[1,5-b]pyridazine-5-carboxylate as a dark red solid.

Intermediate 16: 2-(6-Aminopyridin-3-yl)acetic acid.



Step 1. 1,3-Diethyl 2-(6-nitropyridin-3-yl)propanedioate. A mixture of 5-
5 bromo-2-nitropyridine (2 g, 9.85 mmol, 1.00 equiv), 1,3-diethyl propanedioate (8 g,
49.95 mmol, 5.07 equiv), sodium hydride (1.2 g, 50.00 mmol, 5.07 equiv), L-proline
(0.1 g) and CuI (100 mg, 0.53 mmol, 0.05 equiv) in DMF (100 mL) was stirred under
nitrogen at 100°C for 20 h. The reaction was cooled to rt and then concentrated under
vacuum. The mixture was diluted with 300 mL of ethyl acetate and then washed with
10 3x50 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate, filtered
and concentrated to give 2 g of crude 1,3-diethyl 2-(6-nitropyridin-3-yl)propanedioate
as red oil. TLC: ethyl acetate/petroleum ether = 1/1, R_f = 0.4.

Step 2. 1,3-Diethyl 2-(6-aminopyridin-3-yl)propanedioate. A mixture of 1,3-
15 diethyl 2-(6-nitropyridin-3-yl)propanedioate (1.5 g, 5.31 mmol, 1.00 equiv) and
Raney Ni (1 g) in MeOH (30 mL) was stirred under 1 atmosphere of H₂ at rt for 2 h.
The catalyst was removed by filtration and the filtrate was concentrated under
vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH
(5:1) to afford 1.3 g of crude 1,3-diethyl 2-(6-aminopyridin-3-yl)propanedioate as a
yellow oil. TLC: dichloromethane/MeOH=5/1; R_f = 0.5.


20 Step 3. Ethyl 2-(6-aminopyridin-3-yl)acetate. A mixture of 1,3-diethyl 2-(6-
aminopyridin-3-yl)propanedioate (1.2 g, 4.76 mmol, 1.00 equiv) and lithium chloride
(600 mg, 14.15 mmol, 2.98 equiv) in DMSO (10 mL) and water (1 mL) was stirred
under nitrogen at 130°C for 18 h. The reaction was cooled to rt and then diluted with
200 mL of ethyl acetate. The mixture was washed with 3x50 mL of H₂O. The organic
25 layer was dried over anhydrous sodium sulfate, filtered and concentrated under
vacuum. The residue was purified on a silica gel column eluted with ethyl
acetate/hexane (1/1) to give 200 mg (23%) of ethyl 2-(6-aminopyridin-3-yl)acetate as
a light yellow oil. TLC: DCM/MeOH=5:1; R_f = 0.6.

Step 4. To a solution of ethyl 2-(6-aminopyridin-3-yl)acetate (200 mg, 1.11
30 mmol, 1.00 equiv) in ethanol (5 mL) was added a solution of potassium hydroxide

(200 mg, 3.56 mmol, 3.21 equiv) in water (5 mL). The reaction mixture was stirred under nitrogen at rt for 11 h. The pH value of the solution was adjusted to 6 with 2 M HCl. The resulting mixture was concentrated under vacuum to give 0.5 g of crude 2-(6-aminopyridin-3-yl)acetic acid as an off-white solid. TLC: DCM/MeOH = 5/1; $R_f =$

5 0.2.

Intermediate 17. 1H,2H,3H-Pyrrolo[3,4-c]pyridin-6-amine.



Step 1. 6-Chloropyridine-3-carbonyl chloride. A mixture of 6-chloropyridine-3-carboxylic acid (20 g, 126.94 mmol, 1.00 equiv), DMF (1 g, 13.68 mmol, 0.11 equiv) and thionyl chloride (20 mL) in toluene (200 mL) was stirred under nitrogen for 3 h at 80°C. The resulting solution was cooled to rt and concentrated under vacuum to give 25 g of crude 6-chloropyridine-3-carbonyl chloride as a light yellow solid.

15 Step 2. 6-Chloro-N,N-bis(propan-2-yl)pyridine-3-carboxamide. To a solution of 6-chloropyridine-3-carbonyl chloride (25 g, 142.05 mmol, 1.00 equiv) in DCM (500 mL) maintained under nitrogen at 0°C was added diisopropylamine (50 g, 494.12 mmol, 3.48 equiv) dropwise with stirring. The reaction mixture was stirred for 50 min at 25°C and then quenched with the addition of H₂O (300 mL). The organic layer was collected and the aqueous layer was extracted with 2x300 mL of DCM. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to give 25 g (73%) of 6-chloro-N,N-bis(propan-2-yl)pyridine-3-carboxamide as a light yellow solid. TLC: ethyl acetate:petroleum ether=1:2, $R_f = 0.4$.

25 Step 3. 6-Chloro-4-formyl-N,N-bis(propan-2-yl)pyridine-3-carboxamide. To a stirred solution of diisopropylamine (1 g, 9.88 mmol, 4.76 equiv) in ether (30 mL) at -50°C maintained under nitrogen was added a 2.5M solution of n-BuLi (5 mL) in hexanes dropwise. The reaction mixture was stirred for 30 min at -50°C then solid 6-chloro-N,N-bis(propan-2-yl)pyridine-3-carboxamide (500 mg, 2.08 mmol, 1.00 equiv) was added in a single portion. The resulting solution was stirred for 30 min at -50°C.

30

DMF (1 mL) was then added dropwise with stirring. The reaction mixture was stirred at -50°C for 3 h and then warmed to rt and stirred overnight. The reaction was quenched by the addition of 10% aqueous citric acid solution (30 mL) and then extracted with 2x50 mL of ether. The combined organic layers was dried over anhydrous sodium sulfate and concentrated under vacuum to give 0.5 g of crude 6-chloro-4-formyl-N,N-bis(propan-2-yl)pyridine-3-carboxamide as a yellow solid. LC/MS (Method G, ESI): RT= 1.40 min, $m/z = 269.0$ [M+H]⁺.

Step 4. 6-Chloro-4-(hydroxymethyl)-N,N-bis(propan-2-yl)pyridine-3-carboxamide. A mixture of 6-chloro-4-formyl-N,N-bis(propan-2-yl)pyridine-3-carboxamide (500 mg, 1.86 mmol, 1.00 equiv) and NaBH₄ (500 mg, 13.22 mmol, 7.10 equiv) in ethanol (50 mL) was stirred for 50 min at 30°C. The reaction was then quenched by the addition of 1M HCl. The solid was removed by filtration and the filtrate was concentrated to provide 0.5 g of crude 6-chloro-4-(hydroxymethyl)-N,N-bis(propan-2-yl)pyridine-3-carboxamide as a light yellow solid. LC/MS (Method F, ESI): RT= 1.25 min, $m/z = 271.0$ [M+H]⁺.

Step 5. 6-Chloro-1H,3H-furo[3,4-c]pyridin-3-one. A mixture of 6-chloro-4-(hydroxymethyl)-N,N-bis(propan-2-yl)pyridine-3-carboxamide (2 g, 7.39 mmol, 1.00 equiv) in 6M HCl (40 mL) was stirred for 30 min at 100°C. The reaction mixture was cooled to rt and the pH of the solution was adjusted to 8 with sodium carbonate. The mixture was extracted with 200 mL of DCM. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum to yield 1 g of crude 6-chloro-1H,3H-furo[3,4-c]pyridin-3-one as a light yellow solid. LC/MS (Method R, ESI): RT = 1.13 min, $m/z = 170.0$ [M+H]⁺.

Step 6. [6-Chloro-4-(hydroxymethyl)pyridin-3-yl]methanol. A mixture of 6-chloro-1H,3H-furo[3,4-c]pyridin-3-one (1 g, 5.90 mmol, 1.00 equiv) and NaBH₄ (0.5 g) in ethanol (50 mL) was stirred for 60 min at 25°C. The pH of the solution was adjusted to 1 with 6M HCl. The solid was removed by filtration and the filtrate was concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (20:1) to give 0.4 g (39%) of [6-chloro-4-(hydroxymethyl)pyridin-3-yl]methanol as a light yellow solid. LCMS (Method R, ESI): RT = 0.95min, $m/z = 174.0$ [M+H]⁺.

Step 7. 2-Chloro-4,5-bis(chloromethyl)pyridine hydrochloride. A mixture of [6-chloro-4-(hydroxymethyl)pyridin-3-yl]methanol (100 mg, 0.58 mmol, 1.00 equiv) and thionyl chloride (2 mL) in DCM (20 mL) was stirred under nitrogen at rt for 1 h. The resulting mixture was concentrated under vacuum to give 0.1 g of crude 2-chloro-4,5-bis(chloromethyl)pyridine hydrochloride as a dark red solid.

Step 8. 6-Chloro-2-[(2,4-dimethoxyphenyl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine. A mixture of 2-chloro-4,5-bis(chloromethyl)pyridine hydrochloride (1 g, 4.05 mmol, 1.00 equiv), (2,4-dimethoxyphenyl) methanamine (1 g, 5.98 mmol, 1.48 equiv) and DIPEA (1 g, 7.74 mmol, 1.91 equiv) in DCM (60 mL) was stirred under nitrogen overnight at rt. The resulting mixture was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (2:1) to give 0.9 g of crude 6-chloro-2-[(2,4-dimethoxyphenyl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine as a reddish oil. LC/MS (Method R, ESI): RT = 0.94 min, $m/z = 305.0$ [M+H]⁺.


Step 9. N,2-bis[(2,4-Dimethoxyphenyl) methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridin-6-amine. A mixture of 6-chloro-2-[(2,4-dimethoxyphenyl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine (200 mg, 0.66 mmol, 1.00 equiv), Pd₂(dba)₃•CHCl₃ (0.1 g), t-BuONa (200 mg, 2.08 mmol, 3.17 equiv), BINAP (100 mg, 0.16 mmol, 0.24 equiv) and (2,4-dimethoxyphenyl)methanamine (400 mg, 2.39 mmol, 3.65 equiv) in toluene (20 mL) was stirred under nitrogen overnight at 80°C. The resulting solution was diluted with 20 mL of H₂O and extracted with 2x50 mL of ethyl acetate. The combined organic layers was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:1) to give 0.2 g (70%) of N,2-bis[(2,4-dimethoxyphenyl) methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridin-6-amine as a dark red solid. LC/MS (Method R, ESI): RT = 0.92 min, $m/z = 436.0$ [M+H]⁺.

Step 10. A solution of N,2-bis[(2,4-dimethoxyphenyl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridin-6-amine (300 mg, 0.69 mmol, 1.00 equiv) in TFA (20 mL) was stirred under nitrogen overnight at 90°C. The resulting mixture was concentrated under vacuum to remove most of the TFA. The pH of the residue was adjusted to 8 with saturated sodium carbonate solution. The mixture was concentrated under

vacuum and the residue was dissolved in hot ethyl acetate and filtered. The filtrate was concentrated under reduced pressure to provide 0.15 g of crude 1H,2H,3H-pyrrolo[3,4-c]pyridin-6-amine as a red oil. LC/MS (Method R, ESI): RT = 0.18 min, $m/z = 136.0$ [M+H]⁺.

5

Intermediate 18. 2,3-Dihydro-1H-pyrrolo[3,4-c]pyridine-d₄ hydrochloride.



Step 1. 3,4-Dimethyl pyridine-3,4-dicarboxylate. To a solution of pyridine-3,4-dicarboxylic acid (35 g, 209.43 mmol, 1.00 equiv) in MeOH (250 mL) was added concentrated sulfuric acid (55 mL) dropwise with stirring at rt. The resulting solution was stirred at 100°C overnight. The reaction mixture was cooled to rt and then concentrated under vacuum. The residue was diluted with 500 mL of H₂O and the pH of the solution was adjusted to 8 with 2M aqueous sodium carbonate solution. The resulting mixture was extracted with 3x300 mL of DCM. The combined organic layers were washed with 3x500 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum to give 30 g (73%) of 3,4-dimethyl pyridine-3,4-dicarboxylate as a yellow oil. LC/MS (Method R, ESI): RT = 1.23 min, $m/z = 195.0$ [M+H]⁺.

Step 2. 3,4-bis(Hydroxymethyl)pyridine-d₄ hydrochloride. To a stirred solution of NaBD₄ (4.3 g, 102.38 mmol, 4.00 equiv) in EtOD (15 mL) maintained under nitrogen at 0°C was added dropwise a solution of 3,4-dimethyl pyridine-3,4-dicarboxylate (5.0 g, 25.62 mmol, 1.00 equiv) in EtOD (10 mL) followed by a solution of CaCl₂ (2.5 g, 22.73 mmol, 0.90 equiv) in EtOD (20 mL) dropwise at 0°C. The reaction mixture was stirred for 3 h while the reaction temperature was maintained between 0 to 10°C by an ice/water bath. The resulting mixture was concentrated under vacuum and the residue was dissolved in 200 mL of ethanol. The solid material was removed by filtration. The filtrate was cooled to 0°C then hydrogen chloride gas was bubbled into the solution while keeping the reaction temperature below 5°C with an ice/water bath. The resulting solution was stirred at 0-10°C for 2 h.


The reaction mixture was concentrated under vacuum and the residue was triturated in 100 mL of THF. The solid was collected by filtration and dried in vacuum to give 2.9 g of crude 3,4-bis(hydroxymethyl)pyridine-d₄ hydrochloride as a white solid. ¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 9.00 (d, *J* = 2.1 Hz, 1H), 8.70 (s, 1H), 8.07 (d, *J* = 6.0 Hz, 1H).

Step 3. 2-(2,4-Dimethoxybenzyl)-2,3-dihydro-1H-pyrrolo[3,4-c]pyridine-d₄.
To a solution of 3,4-bis(hydroxymethyl)pyridine-d₄ hydrochloride (3.0 g, 1.00 equiv) in DCM (80 mL) maintained under nitrogen at -5°C was added dropwise triethylamine (10 mL, 5.00 equiv) followed by methanesulfonyl chloride (4.5 mL, 3.00 equiv) in 5 min. The reaction mixture was stirred at 0°C for 1 h. The reaction was then quenched by the addition of 20 mL of water and the resulting mixture was extracted with 2x50 mL of DCM. The combined organic layers was washed with 1x200 mL of brine, dried over anhydrous sodium sulfate and filtered. The filtrate was cooled to 0°C then DIPEA (5 mL, 2.0 equiv) followed by (2,4-dimethoxyphenyl)methanamine (3 mL, 1.10 equiv) was added dropwise at 0°C within 5 min. The resulting solution was stirred at rt for 8 h. The reaction was then quenched by the addition of 20 mL of water. The resulting mixture was washed with 1x200 mL of water and 2x200 mL of brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (96/4) to give 1.4 g of crude as 2-(2,4-dimethoxybenzyl)-2,3-dihydro-1H-pyrrolo[3,4-c]pyridine-d₄ as a red oil.

Step 4. A solution of 2-(2,4-dimethoxybenzyl)-2,3-dihydro-1H-pyrrolo[3,4-c]pyridine-d₄ (700 mg, 2.04 mmol, 1.00 equiv, 80%) in CF₃COOD (5 mL) was stirred at 70°C for 10 h. The resulting mixture was concentrated under vacuum and the residue was dissolved in 50 mL of DCM. Hydrogen chloride gas was bubbled into the stirred reaction mixture at 0°C for 30 min. The precipitate was collected by filtration and dried in vacuum to give 600 mg of crude 2,3-dihydro-1H-pyrrolo[3,4-c]pyridine-d₄ hydrochloride as a yellow solid. ¹H-NMR (400 MHz, D₂O, ppm): δ 8.77 (s, 1H), 8.69 (d, *J* = 6.0 Hz, 1H), 8.00 (d, *J* = 6.0 Hz, 1H).

II. Preparation of Example Compounds

Example 1: tert-Butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate



5

Step 1. 1-[[[(4-Nitrophenoxycarbonyl)amino)methyl]-6-azaspiro[2.5]octane-6-carboxylate. A solution of 4-nitrophenyl chloroformate (42.5 g, 210.85 mmol, 1.00 equiv) and tert-butyl 1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate (50 mg, 0.21 mmol) in toluene (5 mL) was stirred at 120 °C for 1 h. The resulting mixture was concentrated under vacuum to give 50 mg of tert-butyl 1-[[[(4-nitrophenoxycarbonyl)amino)methyl]-6-azaspiro[2.5]octane-6-carboxylate as a yellow solid. LC/MS (Method K, ESI): RT = 2.02 min, $m/z = 407.0$ [M + H]⁺.


10

Step 2. tert-Butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate. A solution of tert-butyl 1-[[[(4-nitrophenoxycarbonyl)amino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (81 mg, 0.20 mmol, 1.00 equiv) and 2,3-dihydro-1H-pyrrolo[3,4-c]pyridine (25 mg, 0.21 mmol, 1.04 equiv) in ethanol (5 mL) was stirred for 1 h at 80 °C. The resulting mixture was concentrated under vacuum and the residue was purified on a silica gel column eluted with dichloromethane/methanol (10/1) to yield 8.9 mg (12%) of the title compound as an off-white solid. LC/MS (Method H, ESI): RT = 1.40 min, $m/z = 387.0$ [M + H]⁺. ¹H NMR (300MHz, CD₃OD) δ 8.55 (s, 1H), 8.47-8.46 (d, $J = 5.1$ Hz, 1H), 7.46-7.44 (d, $J = 5.1$ Hz, 1H), 4.76 (s, 2H), 3.64-3.54 (m, 2H), 3.37 (s, 2H), 3.27-3.20 (m, 2H), 1.70-1.63 (m, 1H), 1.46 (s, 1H), 1.25-1.19 (m, 1H), 1.07-1.05 (m, 1H), 0.62-0.53 (m, 1H), 0.32-0.21 (m, 1H).

20

25

Example 6: N-[[[8-(3,3-dimethylbutanoyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide




Step 1. tert-Butyl 2-[(1H-pyrrolo[3,2-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate. A solution of tert-butyl 1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate (1750 mg, 7.28 mmol, 1.00 equiv), 1H-pyrrolo[3,2-c]pyridine-2-carboxylic acid (1299 mg, 8.01 mmol, 1.10 equiv), EDCI (2786 mg, 14.53 mmol, 2.00 equiv), triethylamine (2946 mg, 29.11 mmol, 4.00 equiv), and HOBt (1181 mg, 8.74 mmol, 1.20 equiv) in DMF (30 mL) was stirred overnight at room temperature. The resulting solution was diluted with 100 mL of water and extracted with 3x100 mL of ethyl acetate. The combined organic layers were washed with 3x200 mL of brine, dried over anhydrous sodium sulfate, and concentrated under vacuum to afford 2.3 g (82%) of the title compound as a red oil. LC/MS (Method C, ESI): RT = 1.29 min, $m/z = 385.0 [M + H]^+$.

Step 2. N-(8-azaspiro[2.5]octan-2-ylmethyl)-1H-pyrrolo[3,2-c]pyridine-2-carboxamide trifluoroacetate salt. A solution of tert-butyl 2-[(1H-pyrrolo[3,2-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate (3.2 g, 8.32 mmol, 1.00 equiv) in TFA (10 mL) and dichloromethane (10 mL) was stirred for 3 h at room temperature. The resulting mixture was concentrated under vacuum to give 4 g of crude title product as a brown oil. LC/MS (Method E, ESI): RT = 0.96 min, $m/z = 285.0 [M + H]^+$.

Step 3. To a solution of N-(8-azaspiro[2.5]octan-2-ylmethyl)-1H-pyrrolo[3,2-c]pyridine-2-carboxamide trifluoroacetate salt (179 mg, 0.44 mmol, 1.00 equiv) and triethylamine (136 mg, 1.32 mmol, 3.01 equiv) in dichloromethane (5 mL) at 0 °C was added 3,3-dimethylbutanoyl chloride (60 mg, 0.44 mmol, 1.00 equiv) dropwise with stirring. The resulting solution was stirred for 1 h at room temperature. The reaction mixture was diluted with 20 mL of dichloromethane, washed with 2x10 mL of brine, then dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product (110 mg) was purified by preparative HPLC (Instrument, Waters 2767-2; Column, sunfire-C18; mobile phase, water with NH_4HCO_3 (1 g/L) and

CH₃CN (25% CH₃CN up to 57% in 10 min); Detector, UV 254 nm) to give 2.3 mg (1%) of the title compound as a white solid. LC/MS (Method H, ESI): RT = 1.95 min, *m/z* = 383.0 [M + H]⁺. ¹H NMR (400 Hz, CDCl₃) δ 9.02 (s, 1H), 8.42-8.41 (d, *J* = 4.8 Hz, 1H), 7.54 (m, 1H), 7.01 (s, 1H), 6.48 (m, 1H), 4.10-4.09 (m, 1H), 3.67-3.32 (m, 5H), 2.35-2.25 (m, 2H), 1.63-1.50 (m, 2H), 1.28-1.22 (m, 1H), 1.07 (s, 1H), 0.71 (s, 1H), 0.39 (s, 1H).

Example 150. N-([6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-1-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide.



10

Step 1. Ethyl 2-(oxetan-3-ylidene)acetate. A solution of oxetan-3-one (2.0 g, 27.75 mmol, 1.00 equiv) and (ethoxycarbonylmethylene)triphenylphosphorane (10.2 g, 29.28 mmol, 1.10 equiv) in DCM (40 mL) was stirred at 0°C for 30 min. The reaction mixture was concentrated under vacuum and the residue was triturated in 200 mL of petroleum ether. The solid material was removed by filtration. The filtrate was concentrated under vacuum to give 3.3 g of crude ethyl 2-(oxetan-3-ylidene)acetate as a colorless liquid. LC/MS (Method A, ESI): RT = 1.32 min, *m/z* = 143.0 [M-H]⁻.

15

Step 2. Ethyl 2-(3-methyloxetan-3-yl)acetate. To a solution of ethyl 2-(oxetan-3-ylidene)acetate (1.5 g, 8.55 mmol, 1.00 equiv, 81%) and CuI (163 mg, 0.86 mmol, 0.10 equiv) in THF (10 mL) maintained under nitrogen was added a solution of chlorotrimethylsilane (1.85 g, 17.03 mmol, 2.00 equiv) in THF (30 mL). The reaction mixture was stirred at rt for 15 min and then cooled to -15°C. A 1.4 N solution of methylmagnesium chloride (24.5 mL, 4.00 equiv) in THF was added dropwise with stirring in 10 min. The resulting solution was stirred at 25°C for 1 h and then quenched by the addition of 20 mL of saturated aqueous NH₄Cl solution. The mixture was concentrated under vacuum. The residue was redissolved in 100 mL of DCM then washed with 1x100 mL of water and 2x100 mL of brine. The organic

25


layer was dried over anhydrous sodium sulfate and concentrated under vacuum to give 600 mg of crude ethyl 2-(3-methyloxetan-3-yl)acetate as a red oil.

Step 3. 2-(3-Methyloxetan-3-yl)acetic acid. To a solution of ethyl 2-(3-methyloxetan-3-yl)acetate (500 mg, 3.16 mmol, 1.00 equiv) in MeOH (15 mL) was added a 2M aqueous sodium hydroxide (5 mL) solution. The mixture was stirred at 40°C for 12 h and then cooled to rt. The reaction mixture was concentrated under vacuum. The residue was dissolved in 50 mL of water and then washed with 3x50 mL of DCM. The aqueous layer was collected and the pH value of the solution was adjusted to 5 with 1M HCl. The resulting mixture was concentrated under vacuum and the residue was dissolved with 50 mL of MeOH. The solid material was removed by filtration. The filtrate was concentrated under vacuum to give 260 mg of crude 2-(3-methyloxetan-3-yl)acetic acid as a red solid. ¹H NMR (400 MHz, D₂O-*d*₆, ppm): δ 4.63 (s, 2H), 4.33 (s, 2H), 2.19 (s, 2H), 1.27 (s, 3H).

Step 4. A solution of 2-(3-methyloxetan-3-yl)acetic acid (66 mg, 0.51 mmol, 2.00 equiv), EDCI (72 mg, 0.38 mmol, 1.50 equiv), HOBt (51 mg, 0.38 mmol, 1.50 equiv) and DIPEA (162 mg, 1.25 mmol, 5.00 equiv) in DMF (10 mL) was stirred at rt for 1 h. N-[6-Azaspiro[2.5]octan-1-ylmethyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide dihydrochloride (90 mg, 0.25 mmol, 1.00 equiv) was then added and the resulting solution was stirred at 25°C for 12 h. The reaction mixture was concentrated under vacuum. The residue was dissolved in 100 mL of DCM and then washed with 2x100 mL of brine. The organic layer was concentrated under vacuum and the residue was purified by Prep-HPLC (SHIMADZU-PDA (LC-08): Column, XSelect CSH Prep C18 OBD Column, 5µm, 19*150mm, mobile phase, water with 0.025% ammonium carbonate and CH₃CN (8% CH₃CN up to 22.0% in 12 min, up to 95.0% in 1 min, hold 95.0% in 1 min, down to 8.0% in 2 min); Detector, UV 254/220 nm) to give 13.5 mg (13%) of N-([6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-1-yl]methyl)-1H-pyrrolo[3,2-c]pyridine-2-carboxamide as a white solid. LC/MS (Method K, ESI): RT = 1.88 min, *m/z* = 397.1 [M+H]⁺. ¹H-NMR (400 MHz, CDCl₃, ppm): δ 9.03 (s, 1H), 8.42 (d, *J* = 5.6 Hz, 1H), 7.39 (d, *J* = 6.0 Hz, 1H), 6.98 (s, 1H), 6.40 (s, 1H), 4.60-4.59 (m, 2H), 4.48-4.46 (m, 2H), 3.99-3.97 (m, 1H), 3.65-3.60 (m,

2H), 3.46~3.31 (m, 3H), 2.73 (s, 2H), 1.76-1.63 (m, 4H), 1.47 (s, 3H), 1.28-1.23 (m, 1H), 1.10-1.08 (m, 1H), 0.74-0.72 (m, 1H), 0.42-0.39 (m, 1H).


Example 159. N-[[6-(p-tolylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide.



A solution of N-(6-azaspiro[2.5]octan-2-ylmethyl)furo[2,3-c]pyridine-2-carboxamide; 2,2,2-trifluoroacetic acid (60 mg, 0.15 mmol, 1.00 equiv), 1-isocyanato-4-methyl-benzene (20 mg, 0.15 mmol, 1.00 equiv) and triethylamine (65
 10 uL, 0.45 mmol, 3.00 equiv) in 10% DMF in DCM (2.0 mL) was stirred at 50 °C for 2 h. The reaction mixture was concentrated under vacuum and the crude product was purified by Prep-HPLC (Column, Sunfire C18 19x150; mobile phase, CH₃CN:NH₄CO₃/H₂O (10 mmol/L) = 5%-85%, 10min; Detector, UV 254 nm) to give
 15 16 mg (25%) of N-[[6-(p-tolylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide as an off-white solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.05 (s, 1H), 9.05-9.00 (m, 1H), 8.47 (d, *J* = 5.3 Hz, 1H), 8.33 (s, 1H), 7.82 (d, *J* = 5.2, 1.0 Hz, 1H), 7.61 (s, 1H), 7.31 (d, 2H), 7.01 (d, *J* = 8.2 Hz, 2H), 3.67-3.53 (m, 2H), 3.46-3.25 (m, 2H), 2.21 (s, 3H), 1.70-1.58 (m, 1H), 1.50-1.37 (m, 2H), 1.24-1.14 (m, 1H), 1.12-0.98 (m, 1H), 0.60-0.52 (m, 1H), 0.31 (t, *J* = 4.8 Hz, 1H).

20

Example 174. N-([6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-1-yl]methyl)furo[2,3-c]pyridine-2-carboxamide.



Step 1. tert-Butyl 2-(2,2-dichloro-3-oxocyclobutyl)acetate. A mixture of tert-butyl but-3-enoate (10 g, 70.33 mmol, 1.00 equiv) in ether (250 mL), trichloroacetyl chloride (34 g, 186.98 mmol, 2.66 equiv) and Zn-Cu (13 g) in ethylene glycol dimethyl ether (40 mL) was stirred under nitrogen at rt for 48 h. The reaction mixture was filtered to remove the solid material. The filtrate was concentrated under vacuum and the residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:10) to yield 21 g of tert-butyl 2-(2,2-dichloro-3-oxocyclobutyl)acetate as a brown liquid. TLC: petroleum ether:ethyl acetate = 2:1, R_f = 0.6.

10 Step 2. tert-Butyl 2-(3-oxocyclobutyl)acetate. A mixture of tert-butyl 2-(2,2-dichloro-3-oxocyclobutyl)acetate (500 mg, 1.98 mmol, 1.00 equiv) and zinc (635 mg, 9.71 mmol, 4.91 equiv) in a saturated NH_4Cl solution in MeOH (20 mL) was stirred at rt overnight. The mixture was filtered to remove the solid material. The filtrate was concentrated under vacuum to give 300 mg (82%) of tert-butyl 2-(3-oxocyclobutyl)acetate as a light yellow oil. TLC: petroleum ether:ethyl acetate = 2:1, R_f = 0.5.

20 Step 3. tert-Butyl 2-(3-hydroxy-3-methylcyclobutyl)acetate. To a stirred solution of tert-butyl 2-(3-oxocyclobutyl)acetate (300 mg, 1.63 mmol, 1.00 equiv) in THF (50 mL) maintained under nitrogen at 0°C was added dropwise a 3M solution of CH_3MgBr (0.8 mL) in THF. The reaction mixture was stirred at rt for 1 h and then quenched by the addition of 20 mL of saturated NH_4Cl solution. The organic layer was collected and the aqueous layer was extracted with 50 mL of DCM. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 270 mg (83%) of tert-butyl 2-(3-hydroxy-3-methylcyclobutyl)acetate as a light yellow oil.


25 Step 4. 2-(3-Hydroxy-3-methylcyclobutyl)acetic acid. To a solution of tert-butyl 2-(3-hydroxy-3-methylcyclobutyl)acetate (270 mg, 1.35 mmol, 1.00 equiv) in DCM (50 mL) was added TFA (1 mL). The resulting solution was stirred overnight at room temperature then concentrated under vacuum to give 240 mg of crude 2-(3-hydroxy-3-methylcyclobutyl)acetic acid as brown oil.

30

Step 5. A solution of 2-(3-hydroxy-3-methylcyclobutyl)acetic acid (200 mg, 1.39 mmol, 5.58 equiv), EDCI (265 mg, 1.38 mmol, 5.56 equiv), HOBT (187 mg, 1.38 mmol, 5.57 equiv) and DIPEA (1 mL) in DMF (10 mL) was stirred for 10 min at rt. N-[6-Azaspiro[2.5]octan-1-ylmethyl]furo[2,3-c]pyridine-2-carboxamide hydrochloride (80 mg, 0.25 mmol, 1.00 equiv) was then added. The reaction mixture was stirred overnight at rt and then concentrated under vacuum. The residue was dissolved in 100 mL of DCM and the resulting solution was washed with 50 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (20:1). The partially purified product (110 mg) was repurified by Prep-HPLC (Column, Xbridge RP18 19*150; mobile phase, CH₃CN:NH₄CO₃/H₂O (10 mmol/L), 5%-19% over 15 min; Detector, UV 254 nm) to give 19.6 mg (19%) of N-([6-[2-(3-hydroxy-3-methylcyclobutyl) acetyl]-6-azaspiro[2.5]octan-1-yl]methyl)furo[2,3-c]pyridine-2-carboxamide as a white solid. LC/MS (Method I, ESI): RT = 1.66 min, *m/z* = 412.0 [M+H]⁺. ¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 9.06 (t, 2H), 8.48 (d, *J* = 5.2 Hz, 2H), 7.81 (d, *J* = 4.4 Hz, 1H), 7.61 (s, 1H), 4.75 (s, 1H), 3.73-3.53 (m, 1H), 3.49-3.32 (m, 1H), 2.50-2.21 (m, 3H), 2.03-2.02 (m, 3H), 1.64 (s, 3H), 1.44-1.33 (m, 2H), 1.19-1.13 (m, 3H), 1.10-1.03 (m, 1H), 0.56-0.53 (m, 1H), 0.32-0.29 (m, 1H).

20

Example 274. tert-Butyl 2-[(4,6-dihydro-1H-pyrrolo[3,4-c]pyrazole-5-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate.




Step 1. tert-Butyl 1-[[4-(4-nitrophenoxycarbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate. To a stirred mixture of tert-butyl 1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate oxalic acid salt (200 mg, 0.61 mmol, 1.00 equiv) and 4-nitrophenyl chloroformate (122 mg, 0.61 mmol, 1.00 equiv) in DCM (10 mL) at 0-5°C was added triethylamine (184 mg, 1.82 mmol, 3.00 equiv)

25

dropwise. The resulting solution was stirred at 0-5°C for another 2 h. The resulting mixture was concentrated under vacuum to give 400 mg of crude tert-butyl 1-[[4-nitrophenoxycarbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate as a yellow solid. TLC: petroleum ether:ethyl acetate =2:1, R_f = 0.3.

5 Step 2. A solution of tert-butyl 1-[[4-nitrophenoxycarbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate (250 mg, 0.62 mmol, 1.00 equiv), 1H,4H,5H,6H-pyrrolo[3,4-c]pyrazole hydrochloride (90 mg, 0.62 mmol, 1.00 equiv) and triethylamine (187 mg, 1.85 mmol, 3.00 equiv) in ethanol (5 mL) was stirred at 78°C for 2 h. The resulting mixture was warmed to rt and concentrated under vacuum. The residue was dissolved in 50 mL of DCM and the resulting mixture was washed with 3x50 mL of brine. The organic layer was concentrated under vacuum and the residue was purified by Prep-HPLC (Waters-1; Column, Sunfire C18 19*150; mobile phase A: 0.2% aqueous NH_4HCO_3 , phase B: CH_3CN , gradient: 10% to 43% B over 10 min, up to 100% B in 13 min; Detector, UV 254 nm) to give 33.7 mg (15%) of tert-butyl 1-[[[1H,4H,5H,6H-pyrrolo[3,4-c]pyrazol-5-yl]carbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate as a white solid. LC/MS (Method Q, ESI): RT = 1.49 min, m/z = 376.2 [M+H]⁺. ¹H-NMR (300 MHz, CDCl_3 , ppm): δ 7.36 (s, 1H), 4.51 (s, 4H), 4.31 (s, 1H), 3.61 (br s, 2H), 3.34-3.22 (m, 4H), 1.67-1.60 (m, 1H), 1.49 (s, 1H), 1.46 (s, 9H), 1.39 (s, 1H), 1.18-1.14 (m, 1H), 0.97-0.92 (m, 1H), 0.60-0.57 (m, 1H), 0.29-0.21 (m, 1H).

Example 275. N-([6-[2-(1,4-dimethylpiperidin-4-yl)acetyl]-6-azaspiro[2.5]octan-1-yl]methyl)furo[2,3-c]pyridine-2-carboxamide.



25 Step 1. Ethyl 2-(1-methylpiperidin-4-ylidene)acetate. A mixture of 1-methylpiperidin-4-one (5 g, 44.19 mmol, 1.00 equiv) and (ethoxycarbonylmethylene)triphenylphosphorane (17 g, 48.80 mmol, 1.00 equiv) in DCM (200 mL) was stirred at 0-5°C for 3 h. The resulting mixture was concentrated

under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:10) to give 1 g (12%) of ethyl 2-(1-methylpiperidin-4-ylidene)acetate as a colorless oil. LC/MS (Method Q, ESI): RT = 0.35 min, $m/z = 184.0[M+H]^+$.

5 Step 2. Ethyl 2-(1,4-dimethylpiperidin-4-yl)acetate. A solution of ethyl 2-(1-methylpiperidin-4-ylidene)acetate (1 g, 5.46 mmol, 1.00 equiv), CuI (1.2 g, 6.30 mmol, 1.15 equiv) and $(CH_3)_3SiCl$ (2.3 g, 21.3 mmol, 3.90 equiv) in THF (50 mL) was stirred under nitrogen at rt for 2 h. The reaction mixture was cooled to $-30^\circ C$ and a 3M solution of MeMgBr (11 mL, 33.0 mmol, 6.04 equiv) in THF was added
10 dropwise with stirring. The reaction mixture was stirred at $-30^\circ C$ for 30 min and then slowly warmed to rt and then stirred overnight. The reaction was quenched by the addition of 50 mL of water and the resulting solution was extracted with 4x50 mL of ethyl acetate. The combined organic layers was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 800 mg of crude ethyl 2-(1,4-
15 dimethylpiperidin-4-yl)acetate as a light yellow oil. LC/MS (Method Q, ESI): RT = 0.26 min, $m/z = 200.0 [M+H]^+$.

Step 3. 2-(1,4-Dimethylpiperidin-4-yl)acetic acid. To a solution of ethyl 2-(1,4-dimethylpiperidin-4-yl)acetate (800 mg, 4.01 mmol, 1.00 equiv) in THF (10 mL) was added a 10% aqueous sodium hydroxide solution (6 mL). The resulting solution
20 was stirred overnight at $65^\circ C$. The reaction mixture was cooled to rt and the pH value of the solution was adjusted to 7 with 5% HCl. The resulting mixture was extracted with 3x50 mL of DCM. The combined organic layers was dried over anhydrous sodium sulfate and concentrated under vacuum to give 1.6 g of crude 2-(1,4-dimethylpiperidin-4-yl)acetic acid as a light yellow solid. LC/MS (Method R, ESI):
25 RT = 0.26 min, $m/z = 172.0 [M+H]^+$.


Step 4. tert-Butyl 1-([furo[2,3-c]pyridin-2-yl]formamido)methyl)-6-azaspiro[2.5]octane-6-carboxylate. A solution of furo[2,3-c]pyridine-2-carboxylic acid (510 mg, 3.13 mmol, 1.50 equiv), EDCI (517 mg, 2.70 mmol, 1.30 equiv), HOBt (365 mg, 2.70 mmol, 1.30 equiv) and DIPEA (2 mL) in DMF (20 mL) was stirred at
30 rt for 10 min. tert-Butyl 1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate (500 mg, 2.08 mmol, 1.00 equiv) was then added and the resulting solution was stirred

overnight at rt. The reaction mixture was diluted with 200 mL of ethyl acetate and washed with 100 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:1) to give 600 mg (75%) of tert-butyl 1-([furo[2,3-c]pyridin-2-ylformamido]methyl)-6-azaspiro[2.5]octane-6-carboxylate as a light yellow oil. LC/MS (Method J, ESI): RT = 1.23 min, *m/z* = 386.0[M+H]⁺.

Step 5. N-[6-Azaspiro[2.5]octan-1-ylmethyl]furo[2,3-c]pyridine-2-carboxamide hydrochloride. A solution of tert-butyl 1-([furo[2,3-c]pyridin-2-ylformamido]methyl)-6-azaspiro[2.5]octane-6-carboxylate (1 g, 2.59 mmol, 1.00 equiv) in 1M solution of hydrogen chloride in MeOH (100 mL) was stirred at rt for 3 h. The reaction mixture was concentrated under vacuum to give 0.8 g (96%) of N-[6-azaspiro[2.5]octan-1-ylmethyl]furo[2,3-c]pyridine-2-carboxamide hydrochloride as a white solid. LC/MS (Method Q, ESI): RT = 0.38 min, *m/z* = 286.0[M+H]⁺.

Step 6. A solution of 2-(1,4-dimethylpiperidin-4-yl)acetic acid (1.6 g (crude solid), 9.34 mmol, 11.16 equiv), EDCI (300 mg, 1.56 mmol, 1.87 equiv), HOBt (250 mg, 1.85 mmol, 2.21 equiv) and DIPEA (2 mL) in DMF (30 mL) was stirred at rt for 10 min. N-[6-Azaspiro[2.5]octan-1-ylmethyl]furo[2,3-c]pyridine-2-carboxamide dihydrochloride (300 mg, 0.84 mmol, 1.00 equiv) was then added and the resulting solution was stirred overnight at rt. The solid material was removed by filtration. The filtrate was concentrated under reduced pressure and the residue was purified by Prep-HPLC (Column: Sunfire C18 19*150; mobile phase, CH₃CN:NH₄CO₃/H₂O (10 mmol/L), 15%-60%, 10 min; Detector, UV 254 nm) to give 43 mg (12%) of N-([6-[2-(1,4-dimethylpiperidin-4-yl)acetyl]-6-azaspiro[2.5]octan-1-yl]methyl)furo[2,3-c]pyridine-2-carboxamide as a light yellow solid. LC/MS (Method A, ESI): RT = 1.70 min, *m/z* = 439.0 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 9.04 (s, 2H), 8.48 (d, *J* = 5.1 Hz, 1H), 7.80 (d, *J* = 5.4 Hz, 1H), 7.60 (s, 1H), 3.67-3.56 (m, 2H), 3.39-3.32 (m, 2H), 2.29-2.24 (m, 5H), 2.11-2.09 (m, 6H), 1.59-1.52 (m, 3H), 1.50-1.22 (m, 4H), 1.21-1.16 (m, 1H), 1.06-1.01 (m, 1H), 0.94 (d, *J* = 5.4 Hz, 3H), 0.60-0.50 (m, 1H), 0.31-0.29 (m, 1H).

Example 339. N-[[6-[4-(4-Methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide.




Step 1. Ethyl 4-(4-methylpiperazin-1-yl)benzoate. A solution of ethyl 4-fluorobenzoate (2 g, 11.89 mmol, 1.00 equiv), 1-methylpiperazine (1.6 g, 15.97 mmol, 1.34 equiv) and potassium carbonate (2.2 g, 15.92 mmol, 1.34 equiv) in DMF (30 mL) was stirred at 80°C for 2 days. The resulting solution was diluted with 150 mL of H₂O and the pH value of the solution was adjusted to 1 with 10% HCl. The resulting solution was washed with 100 mL of ethyl acetate. The aqueous layer was collected and the pH value of the solution was adjusted to 10 with sodium carbonate. The resulting solution was extracted with 2x100 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under vacuum to give 1.3 g (44%) of ethyl 4-(4-methylpiperazin-1-yl)benzoate as a white solid. TLC: DCM:MeOH = 10:1, R_f = 0.5.

Step 2. 4-(4-Methylpiperazin-1-yl)benzoic acid. To a solution of ethyl 4-(4-methylpiperazin-1-yl)benzoate (1.3 g, 5.24 mmol, 1.00 equiv) in THF (10 mL) was added a 10% aqueous sodium hydroxide solution (8 mL). The resulting solution was stirred at 70°C overnight. After cooling to rt, the pH value of the solution was adjusted to 5 with 5% HCl. The resulting solution was extracted with 3x100 mL of DCM. The combined organic layers was and concentrated under vacuum to give 2.2 g of crude 4-(4-methylpiperazin-1-yl)benzoic acid as a white solid. LC/MS (Method C, ESI): RT = 0.54 min, m/z = 221 [M+H]⁺.

Step 3. N-[[6-[4-(4-Methylpiperazin-1-yl)phenyl] carbonyl]-6-azaspiro[2.5]octan-1-yl)methyl]furo[2,3-c]pyridine-2-carboxamide. A solution of 4-(4-methylpiperazin-1-yl)benzoic acid (200 mg, 0.91 mmol, 1.00 equiv), EDCI (200 mg, 1.04 mmol, 1.15 equiv), HOBt (150 mg, 1.11 mmol, 1.22 equiv) and DIPEA (2 mL) in DMF (10 mL) was stirred at rt for 10 min. N-[6-Azaspiro[2.5]octan-1-yl)methyl]furo[2,3-c]pyridine-2-carboxamide dihydrochloride (100 mg, 0.28 mmol,

0.31 equiv) was then added and the resulting solution was stirred at rt overnight. The reaction mixture was concentrated under vacuum and the crude product was purified by Prep-HPLC (Column, Sunfire C18 19x150; mobile phase, CH₃CN: NH₄CO₃/H₂O (10 mmol/L) = 15%-60%, 10min; Detector, UV 254 nm) to give 40 mg (9%) of N-
 5 [(6-[[4-(4-methylpiperazin-1-yl)phenyl] carbonyl]-6-azaspiro[2.5]octan-1-yl)methyl]furo[2,3-c]pyridine-2-carboxamide as an off-white solid. LC/MS (Method F, ESI): RT = 1.12 min, *m/z* = 488.3[M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm):
 δ 9.07-9.05 (m, 2H), 8.48 (d, *J* = 5.4 Hz, 1H), 7.82-7.80 (m, 1H), 7.61 (s, 1H), 7.26 (d,
J = 8.7 Hz, 2H), 6.94 (d, *J* = 8.7 Hz, 2H), 3.62-3.58 (m, 2H), 3.39-3.25 (m, 3H), 3.25-
 10 3.17 (m, 4H), 2.49-2.41 (m, 4H), 2.21 (s, 3H), 1.68-1.62 (m, 1H), 1.46-1.43 (m, 2H),
 1.23-1.18 (m, 1H), 1.04-1.00 (m, 1H), 0.58-0.56 (m, 1H), 0.35-0.29 (m, 1H).

Example 383. tert-Butyl 2-[(pyrazolo[1,5-b]pyridazine-5-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate.




15

A solution of lithium pyrazolo[1,5-b]pyridazine-5-carboxylate (90 mg, 0.53 mmol, 1.00 equiv), tert-butyl 1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate trifluoroacetic acid salt (193 mg, 0.54 mmol, 1.01 equiv), EDCI (372 mg, 1.94 mmol, 3.66 equiv), HOBt (99 mg, 0.73 mmol, 1.38 equiv) and triethylamine (246 mg, 2.43
 20 mmol, 4.6 equiv) in DMF (5 mL) was stirred at rt for 3 h. The reaction was then quenched by the addition of 100 mL of H₂O and the resulting solution was extracted with 3x100 mL of ethyl acetate. The combined organic layers were washed with 3x100 mL of brine, dried over sodium sulfate and concentrated under vacuum. The residue was first purified on a silica gel column eluted with DCM:MeOH (30:1). The
 25 partially purified product was repurified by Prep-HPLC (Waters-1; Column, XBridge C18 19*150; mobile phase A: 0.2% aqueous NH₄HCO₃; mobile phase B: CH₃CN; gradient, 10 to 43% CH₃CN in 10 min, up to 100% in 13 min; Detector, UV 254 nm) to give 17.2 mg (9%) of tert-butyl 1-[(pyrazolo[1,5-b]pyridazin-3-

ylformamido]methyl)-6-azaspiro[2.5]octane-6-carboxylate as a white solid. LC/MS (Method S, ESI): RT = 1.48 min, $m/z = 386.0$ [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.77 (t, *J* = 2.4 Hz, 1H), 8.74 (s, 1H), 8.73 (s, 1H), 8.18 (d, *J* = 2.4 Hz, 1H), 7.05 (d, *J* = 2.7 Hz, 1H), 3.59-3.43 (m, 2H), 3.41-3.35 (m, 2H), 3.25-3.17 (m, 2H), 1.67-1.54 (m, 1H), 1.46-1.31 (m, 11H), 1.19-1.09 (m, 1H), 1.05-0.95 (m, 1H), 0.59-0.50 (m, 1H), 0.31-0.25 (m, 11H).

Example 394. N-[[6-[2-(6-Amino-3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide.




10

A solution of 2-(6-aminopyridin-3-yl)acetic acid (150 mg, 0.99 mmol, 1.00 equiv), N-6-azaspiro[2.5]octan-1-ylmethyl-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide (350 mg, 1.22 mmol, 1.20 equiv), DIPEA (400 mg, 3.00 equiv), EDCI (230 g, 1.20 mol, 1.20 equiv) and HOBt (170 mg, 1.26 mmol, 1.20 equiv) in DMF (20mL) was stirred at rt for 20 h. The resulting mixture was concentrated under vacuum and the residue was first purified on a silica gel column eluted with DCM/MeOH (5:1). The partially purified product (300 mg) was repurified by Flash-Prep-HPLC (IntelFlash-1: Column, silica gel; mobile phase, 15 to 45% CH₃CN in within 30 min; Detector, UV 254 nm) to give 93 mg (22%) of N-[[6-[2-(6-aminopyridin-3-yl)acetyl]-6-azaspiro[2.5]octan-1-yl]methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide as an off-white solid. LC/MS (Method C, ESI): RT = 1.01 min, $m/z = 421.1$ [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.57 (s, 1H), 8.47 (d, *J* = 4.8 Hz, 1H), 7.74 (s, 1H), 7.39 (d, *J* = 4.8 Hz, 1H), 7.21 (dd, *J* = 6.0 Hz, 2.4 Hz, 1H), 6.44-6.36 (m, 2H), 5.72 (s, 2H), 4.62 (d, *J* = 2.7 Hz, 4H), 3.65-3.54 (m, 4H), 3.40-3.33 (m, 1H), 3.18-3.09 (m, 3H), 1.54-1.48 (m, 1H), 1.32-0.95 (m, 4H), 0.55-0.45 (m, 1H), 0.25-0.15 (m, 1H).

20

25

Example 399. N-[[6-[2-(4-Methyltetrahydropyran-4-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide.



Step 1. Ethyl 2-(oxan-4-ylidene)acetate. A solution of oxan-4-one (2 g, 19.98 mmol, 1.00 equiv) and (ethoxycarbonylmethylene)triphenylphosphorane (8 g, 22.96 mmol, 1.15 equiv) in acetonitrile (50 mL) was stirred at 70°C for 16 h. The resulting mixture was cooled to rt and concentrated. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:5) to give 1.9 g (56%) of ethyl 2-(oxan-4-ylidene)acetate as a colorless solid. TLC: petroleum ether:ethyl acetate = 1:2, $R_f = 0.5$.

Step 2. Ethyl 2-(4-methyloxan-4-yl)acetate. To a stirred solution of CuI (2.9 g, 15.23 mmol, 3.24 equiv) in ether (50 mL) maintained under nitrogen at 0°C was added a 1.6 M solution of methyllithium (20 mL, 32 mmol, 6.8 equiv) in ether dropwise. The resulting solution was stirred at 0°C for 10 min and the ether solvent was removed from the reaction vessel under vacuum at 0°C. DCM (50 mL) was then added to the residue and the reaction was cooled to -78°C. Chlorotrimethylsilane (1.9 g, 17.6 mmol, 3.7 equiv) was then added dropwise followed by a solution of ethyl 2-(oxan-4-ylidene)acetate (800 mg, 4.70 mmol, 1.00 equiv) in DCM (20 mL) to the reaction mixture at -78°C. The resulting solution was stirred for another 30 min at -78°C and then at 0°C for 2 h. The reaction quenched by the addition of 50 mL of water. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum to give 750 mg (86%) of ethyl 2-(4-methyloxan-4-yl)acetate as a colorless oil.


Step 3. 2-(4-Methyloxan-4-yl)acetic acid. To a solution of ethyl 2-(4-methyloxan-4-yl)acetate (750 mg, 4.03 mmol, 1.00 equiv) in ethanol (20 mL) at 0°C was added dropwise a solution of sodium hydroxide (817 mg, 5.00 equiv) in water (4 mL) with stirring. The resulting solution was stirred overnight at rt and then concentrated to remove ethanol. The resulting solution was diluted with 10 mL of H₂O and washed with 3x20 mL of ether. The aqueous layer was collected and the pH value of the solution was adjusted to 4-5 with 2N HCl. The aqueous layer was

extracted with 4x15 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated to give 600 mg (94%) of 2-(4-methyloxan-4-yl)acetic acid as yellow oil. LC/MS (Method S, ESI): RT = 0.5 min, $m/z = 159 [M+H]^+$.

5 Step 4. A solution of 2-(4-methyloxan-4-yl)acetic acid (100 mg, 0.63 mmol, 1.26 equiv), DIPEA (1 mL), HOBt (85 mg, 0.63 mmol, 1.26 equiv), and EDCI (120 mg, 0.63 mmol, 1.25 equiv) in DMF (4 mL) was stirred at rt for 10 min. N-[6-Azaspiro[2.5]octan-1-ylmethyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide dihydrochloride (180 mg, 0.50 mmol, 1.00 equiv) was then added and the reaction
10 mixture was stirred overnight at rt. The resulting solution was diluted with 100 mL of DCM and then washed with 60 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was purified on a silica gel column eluted with DCM/MeOH (20:1) to give 110 mg (51%) of N-([6-[2-(4-methyloxan-4-yl)acetyl]-6-azaspiro[2.5]octan-1-yl]methyl)-1H,2H,3H-pyrrolo[3,4-
15 c]pyridine-2-carboxamide as a light yellow solid. LC/MS (Method F, ESI): RT = 1.19 min, $m/z = 428.0 [M+H]^+$. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.55 (s, 1H), 8.46 (d, *J* = 5.1 Hz, 1H), 7.38 (d, *J* = 5.1 Hz, 1H), 6.42 (t, *J* = 4.9 Hz, 1H), 4.61 (d, 4H), 3.69-3.31 (m, 8H), 3.15-3.08 (m, 2H), 2.31 (s, 2H), 1.62-1.49 (m, 3H), 1.45-1.25 (m, 4H), 1.21-1.05 (m, 1H), 1.04-0.89 (m, 4H), 0.55-0.51 (m, 1H), 0.25-0.15 (m, 1H).

20

Example 403. N-([6-[2-[4-Methyl-1-(2,2,2-trifluoroethyl)-4-piperidyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl)-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide.



25 Step 1. tert-Butyl 4-methyl-4-[2-oxo-2-(1-[[[1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl]carbonyl)amino]methyl]-6-azaspiro[2.5]octan-6-yl)ethyl]piperidine-1-carboxylate. A solution of 2-[1-[(tert-butoxy)carbonyl]-4-methylpiperidin-4-yl]acetic acid (450 mg, 1.75 mmol, 2.09 equiv), EDCI (340 mg), HOBt (250 mg, 1.85 mmol, 2.22 equiv) and DIPEA (2 mL) in DMF (15 mL) was stirred at rt for 10 min. N-[6-Azaspiro[2.5]octan-1-ylmethyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide

dihydrochloride (300 mg, 0.83 mmol, 1.00 equiv) was then added and the reaction mixture was stirred overnight at rt. The resulting solution was diluted with 200 mL of ethyl acetate and washed with 100 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate and concentrated. The residue was purified on a silica gel column eluted with DCM/MeOH (20:1) to yield 210 mg (48%) of tert-butyl 4-methyl-4-[2-oxo-2-(1-[[[1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl]carbonyl)amino]methyl]-6-azaspiro[2.5]octan-6-yl)ethyl]piperidine-1-carboxylate as a light yellow oil. TLC: DCM:MeOH=10:1, R_f = 0.5.

Step 2. N-([6-[2-(4-Methylpiperidin-4-yl)acetyl]-6-azaspiro[2.5]octan-1-yl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide hydrochloride.


Hydrogen chloride gas was bubbled into a solution of tert-butyl 4-methyl-4-[2-oxo-2-(1-[[[1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl]carbonyl)amino]methyl]-6-azaspiro[2.5]octan-6-yl)ethyl]piperidine-1-carboxylate (210 mg, 0.40 mmol, 1.00 equiv) in 1,4-dioxane (20 mL) at 0°C for 20 min. The resulting solution was stirred at rt for 1 h and then concentrated under vacuum to give 220 mg of crude N-([6-[2-(4-methylpiperidin-4-yl)acetyl]-6-azaspiro[2.5]octan-1-yl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide hydrochloride as a light yellow solid. LC/MS (Method F, ESI): RT = 0.89 min, m/z = 426.0[M+H]⁺

Step 3. To a solution of N-([6-[2-(4-methylpiperidin-4-yl)acetyl]-6-azaspiro[2.5]octan-1-yl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide dihydrochloride (220 mg, 0.44 mmol, 1.00 equiv) in DMF (10 mL) at rt was added 2,2,2-trifluoroethyl trifluoromethanesulfonate (140 mg, 0.60 mmol, 1.37 equiv) followed by DIPEA (2 mL). The reaction mixture was stirred at room temperature for 2 h and then diluted with 200 mL of ethyl acetate. The resulting mixture was washed with 100 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate, filtered and concentrated under vacuum. The residue was purified on a silica gel column eluted with dichloromethane/methanol (20:1) to give 35.2 mg (16%) of N-([6-[2-[4-methyl-1-(2,2,2-trifluoroethyl)piperidin-4-yl]acetyl]-6-azaspiro[2.5]octan-1-yl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide as a white solid.

LC/MS (Method I, ESI): RT = 1.15 min, m/z = 508.0 [M+H]⁺. ¹H NMR (300MHz, DMSO-*d*₆, ppm): δ 8.56 (s, 1H), 8.46 (d, *J* = 4.5 Hz, 1H), 7.39 (d, *J* = 4.8 Hz, 1H),

6.43 (s, 1H), 4.62 (s, 4H), 3.65-3.58 (m, 2H), 3.48-3.33 (m, 3H), 3.20-3.01 (m, 4H), 2.59-2.50 (m, 3H), 2.27 (s, 2H), 1.59-1.56 (m, 3H), 1.39-1.30 (m, 3H), 1.01-0.96 (m, 4H), 0.49-0.45 (m, 1H), 0.25-0.23 (m, 1H).

- 5 Example 405. N-[[6-[4-(1-Methyl-4-piperidyl)benzoyl]-6-azaspiro[2.5]octan-2-yl)methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide.



Step 1. tert-Butyl 4-[4-(ethoxycarbonyl)phenyl]-1,2,3,6-tetrahydropyridine-1-carboxylate. A mixture of ethyl 4-bromobenzoate (1 g, 4.37 mmol, 1.00 equiv), tert-butyl 4-(tetramethyl-1,3,2-dioxaborolan-2-yl)-1,2,3,6-tetrahydropyridine-1-carboxylate (1.4 g, 4.53 mmol, 1.04 equiv), Pd(PPh₃)₄ (0.2 g), and potassium carbonate (800 mg, 5.79 mmol, 1.33 equiv) in 1,4-dioxane (10 mL) was stirred under nitrogen in a 20-mL sealed tube at 100°C overnight. The reaction mixture was cooled to rt then diluted with 120 mL of ethyl acetate. The solid material was removed by
10 filtration and the filtrate was washed with 100 mL of water. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:20) to give 1 g (69%) of tert-butyl 4-[4-(ethoxycarbonyl)phenyl]-1,2,3,6-tetrahydropyridine-1-carboxylate as a white solid. TLC: petroleum ether:ethyl acetate = 5:1, R_f = 0.3.

Step 2. tert-Butyl 4-[4-(ethoxycarbonyl)phenyl]piperidine-1-carboxylate. A mixture of tert-butyl 4-[4-(ethoxycarbonyl)phenyl]-1,2,3,6-tetrahydropyridine-1-carboxylate (1 g, 3.02 mmol, 1.00 equiv) and 10% palladium on carbon (0.5 g) catalyst in ethanol (50 mL) was stirred under 1 atmosphere of H₂ at rt for 2 h. The catalyst was removed by filtration. The filtrate was concentrated under vacuum to
25 give 1.2 g of crude tert-butyl 4-[4-(ethoxycarbonyl)phenyl]piperidine-1-carboxylate as a gray colored oil. TLC: petroleum ether:ethyl acetate = 5:1, R_f = 0.4.

Step 3. Ethyl 4-(piperidin-4-yl)benzoate hydrochloride. Thionyl chloride (2 mL) was added dropwise with stirring at rt to ethanol (50 mL). tert-Butyl 4-[4-

(ethoxycarbonyl)phenyl]piperidine-1-carboxylate (1 g, 3.00 mmol, 1.00 equiv) was then added to the reaction mixture and the resulting solution was stirred at rt for 1 h. The reaction mixture was concentrated under vacuum to give 0.8 g (99%) of ethyl 4-(piperidin-4-yl)benzoate hydrochloride as an off-white solid. LC/MS (Method Q, ESI): RT = 1.03min, $m/z = 233 [M+H]^+$.


Step 4. Methyl 4-(1-methylpiperidin-4-yl)benzoate. A solution of methyl 4-(piperidin-4-yl)benzoate hydrochloride (800 mg, 3.13 mmol, 1.00 equiv) and paraformaldehyde (180 mg) in ethanol (50 mL) and water(10 mL) was stirred at 60°C for 1 h. Sodium cyanoborohydride (400 mg) and acetic acid (0.1 mL) were then added. The resulting solution was stirred overnight at 60°C. The reaction mixture was cooled to rt and concentrated under vacuum. The resulting solution was diluted with 50 mL of H₂O and extracted with 3x50 mL of DCM. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (10:1) to give 700 mg (96%) of methyl 4-(1-methylpiperidin-4-yl)benzoate as a colorless oil. LC/MS (Method I, ESI): RT = 1.12 min, $m z = 247 [M+H]^+$.

Step 5. 4-(1-Methylpiperidin-4-yl)benzoic acid. To a solution of ethyl 4-(1-methylpiperidin-4-yl)benzoate (700 mg, 2.83 mmol, 1.00 equiv) in ethanol (30 mL) was added a solution of sodium hydroxide (340 mg, 8.50 mmol, 3.00 equiv) in water (5 mL). The reaction mixture was stirred overnight at 60°C and then cooled tort. The pH of the solution was adjusted to 7 with 5% HCl and then extracted with 3x50 mL of DCM. The combined organic layers were dried over anhydrous sodium sulfate then concentrated under vacuum to give 1 g of crude 4-(1-methylpiperidin-4-yl)benzoic acid as a white solid. LC/MS (Method I, ESI): RT = 0.93 min, $m z = 220 [M+H]^+$.

Step 6. A solution of 4-(1-methylpiperidin-4-yl)benzoic acid (131 mg, 0.60 mmol, 2.15 equiv), EDCI (80 mg, 0.42 mmol, 1.50 equiv), HOBt (57 mg, 0.42 mmol, 1.52 equiv) and DIPEA (1 mL) in DMF (6 mL) was stirred at rt for 10 min. N-[6-Azaspiro[2.5]octan-1-ylmethyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide dihydrochloride (100 mg, 0.28 mmol, 1.00 equiv) was then added and the reaction mixture was stirred overnight at rt. The resulting solution was diluted with 30 mL of

H₂O then extracted with 3x50 mL of DCM. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. The crude product was purified by Prep-HPLC (Column, Sunfire C18 19*150; mobile phase, CH₃CN:NH₃/H₂O (5 mL/10 L), 15%-70%, 10min; Detector, UV 254 nm) to give 23.7 mg (17%) of N-[(6-[[4-(1-methylpiperidin-4-yl)phenyl]carbonyl]-6-azaspiro[2.5]octan-1-yl)methyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide as a white solid. LC/MS (Method I, ESI): RT = 1.25 min, *m z* = 488.0 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.56 (s, 1H), 8.46 (d, *J* = 4.5 Hz, 1H), 7.38 (d, *J* = 3.6 Hz, 1H), 7.29 (s, 4H), 6.43 (s, 1H), 4.63 (s, 4H), 3.76-3.74 (m, 1H), 3.51-3.33 (m, 3H), 3.20-3.12 (m, 2H), 2.83 (d, *J* = 5.1 Hz, 2H), 2.44-2.39 (m, 1H), 2.18 (s, 3H), 1.98-1.91 (m, 2H), 1.70-1.62 (m, 5H), 1.54-1.24 (m, 3H), 1.01-0.96 (m, 1H), 0.54-0.49 (m, 1H), 0.24-0.21 (m, 1H).


Example 406. tert-Butyl 2-[[[(6-amino-1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate.



A solution of tert-butyl 1-[[[(4-nitrophenoxycarbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate (200 mg, 0.49 mmol, 1.00 equiv), 1H,2H,3H-pyrrolo[3,4-c]pyridin-6-amine (80 mg, 0.59 mmol, 1.20 equiv) and DIPEA (200 mg, 1.55 mmol, 3.14 equiv) in DCM (50 mL) was stirred at rt for 2 h. The reaction mixture was concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (5:1) to give 0.113 g (57%) of tert-butyl 1-[[[(6-amino-1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl)carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate as a light yellow solid. LC/MS (Method K, ESI): RT = 1.65 min, *m z* = 402.2 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 7.94 (s, 1H), 6.59 (s, 1H), 5.25 (s, 2H), 4.64-4.52 (m, 6H), 3.61 (s, 2H), 3.27 (d, *J* = 8.1 Hz,

5H), 1.65 (d, $J = 8.4$ Hz, 1H), 7.94 (s, 9H), 1.14 -1.09 (m, 1H), 0.95-0.83 (m, 1H), 0.59-0.55 (m, 1H), 0.26-0.24 (m, 1H).

5 Example 409. (3-Methyloxetan-3-yl) 2-[[[(1,1,3,3-tetradeuteriopyrrolo[3,4-c]pyridine-2-carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate.




Step 1. tert-Butyl 1-[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl) carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate-d₄. To a solution of tert-butyl 1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate (1.0 g, 4.16 mmol, 1.00 equiv) in DCM (20 mL) at 0°C was added 4-nitrophenyl carbonochloridate (830 mg, 4.12 mmol, 1.00 equiv) and triethylamine (0.6 mL, 4.6 mmol, 1.10 equiv). The resulting solution was stirred at rt for 2 h. A solution of triethylamine (1.5 mL) in DCM (80 mL) followed by 2,3-dihydro-1H-pyrrolo[3,4-c]pyridine-d₄ hydrochloride (700 mg, crude) were added. The reaction mixture was stirred at 25°C for 3 h then washed with 1x100 mL of sodium bicarbonate, 1x100 mL of water and 1x100 mL of brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (95/5) to give 500 mg of tert-butyl 1-[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl) carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate-d₄ as a yellow solid. LC/MS (Method I, ESI): RT = 1.17 min, $m/z = 391.0$ [M+H]⁺. ¹H NMR (300 MHz, DMSO-d₆, ppm): δ 8.47 (s, 1H), 8.46 (d, $J = 4.5$ Hz, 1H), 7.38 (d, $J = 4.2$ Hz, 1H), 6.34 (s, 1H), 3.51-3.41 (m, 2H), 3.22-3.02 (m, 4H), 1.62-1.55 (m, 1H), 1.42-1.35 (m, 11H), 1.22-1.08 (m, 1H), 0.99-0.92 (m, 1H), 0.51-0.39 (m, 1H), 0.22-0.17 (m, 1H).

25 Step 2. N-{6-Azaspiro[2.5]octan-1-ylmethyl}-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide-d₄. To a solution of tert-butyl 1-[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl) carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate-d₄ (500 mg, 1.00 equiv) in DCM (15 mL) was added CF₃COOD (2 mL) dropwise at rt. The

resulting solution was stirred at rt for 1 h then concentrated under vacuum. The residue was dissolved in 10 mL of MeOH. The pH of the solution was adjusted to 8 with saturated aqueous sodium carbonate solution. The resulting mixture was concentrated under vacuum to give 1.5 g of crude N-{6-azaspiro[2.5]octan-1-ylmethyl}-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide-d₄ as a yellow solid.
5 LC/MS (Method I, ESI): RT = 0.82 min, *m/z* = 291.0 [M+H]⁺.

Step 3. A solution of N-{6-azaspiro[2.5]octan-1-ylmethyl}-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide-d₄ (500 mg, crude), 3-methyloxetan-3-yl 4-nitrophenyl carbonate (136 mg, 0.54 mmol, 1.2 equiv) and triethylamine (91 mg, 0.90 mmol, 2.0 equiv) in DCM (15 mL) was stirred at rt for 5 h. The reaction mixture was diluted with 50 mL of DCM and then washed with 1x50 mL of sodium bicarbonate, 1x50 mL of water and 1x50 mL of brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was first purified on a silica gel column eluted with DCM/MeOH (95/5). The partially purified product (80 mg) was repurified by Prep-HPLC (1#-Pre-HPLC-005(Waters): Column, SunFire Prep C18 OBD Column, 5µm, 19*150 mm; mobile phase, water with 10 mmol NH₄HCO₃ and CH₃CN (15.0% CH₃CN up to 34.0% in 10 min, up to 95.0% in 2 min, down to 15.0% in 2 min); Detector, UV 254/220 nm) to give 47 mg of 3-methyloxetan-3-yl 1-[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl)carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate-d₄ as a white solid.
15 LC/MS (Method I, ESI): RT = 1.39 min, *m/z* = 405.0 [M+H]⁺. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.57 (s, 1H), 8.46 (d, *J* = 5.1 Hz, 1H), 7.38 (d, *J* = 5.1 Hz, 1H), 6.42 (t, *J* = 5.1 Hz, 1H), 4.64-4.58 (m, 2H), 4.42-4.36 (m, 2H), 3.50-3.47 (m, 2H), 3.31-3.28 (m, 2H), 3.16-3.06 (m, 2H), 1.62 (s, 3H), 1.64-1.56 (m, 1H), 1.40-1.36 (m, 2H), 1.20-1.17 (m, 1H), 1.00-0.96 (m, 1H), 0.49-0.47 (m, 1H), 0.24-0.20 (m, 1H).
20 25

Example 419. (3-Methyloxetan-3-yl) (2S)-2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate.




Step 1. tert-Butyl 4-(2-ethoxy-2-oxoethylidene)piperidine-1-carboxylate. A mixture of tert-butyl 4-oxopiperidine-1-carboxylate (300 g, 1.51 mol, 1.00 equiv), ethyl 2-(diethoxyphosphoryl)acetate (405 g, 1.81 mol, 1.20 equiv) and potassium carbonate (314 g, 2.26 mol, 1.50 equiv) in DMF (4.5 L) was stirred overnight at 80°C. The reaction was cooled to rt and then quenched by the addition of 5 L of water/ice. The precipitate was collected by filtration and air-dried to give 252 g (62%) of tert-butyl 4-(2-ethoxy-2-oxoethylidene)piperidine-1-carboxylate as a white solid. TLC: ethyl acetate/petroleum ether = 1:2, R_f = 0.6.

Step 2. 6-tert-Butyl 1-ethyl 6-azaspiro[2.5]octane-1,6-dicarboxylate. A solution of trimethylsulfoxonium iodide (618 g, 2.81 mol, 3.00 equiv) and t-BuOK (315 g, 2.81 mol, 3.00 equiv) in DMSO (5 L) was stirred at rt for 1 h. tert-Butyl 4-(2-ethoxy-2-oxoethylidene)piperidine-1-carboxylate (252 g, 935.63 mmol, 1.00 equiv) was then added in several portions. The reaction mixture was stirred at rt overnight and then quenched by the addition of 10 L of saturated NH_4Cl solution. The resulting mixture was extracted with 3x3 L of ethyl acetate. The organic combined layers were washed with 2x1.5 L of brine. The organic layer was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1/10) to give 147 g of 6-tert-butyl 1-ethyl 6-azaspiro[2.5]octane-1,6-dicarboxylate as yellow oil. TLC: ethyl acetate/petroleum ether = 1:2, R_f = 0.5.

Step 3. 6-[(tert-Butoxy)carbonyl]-6-azaspiro[2.5]octane-1-carboxylic acid. To a stirred solution of 6-tert-butyl 1-ethyl 6-azaspiro[2.5]octane-1,6-dicarboxylate (147 g, 1.00 equiv) in ethanol (1 L) at 0°C was added dropwise a solution of sodium hydroxide (104 g, 2.60 mol, 5.00 equiv) in water (200 mL). The resulting solution was stirred at rt overnight. The reaction mixture was concentrated to remove the excess ethanol. Water (2L) was added and the mixture was washed with 3x500 mL of ethyl acetate. The aqueous layer was collected and the pH of the solution was

adjusted to 5-6 with 1M HCl. The resulting mixture was extracted with 4x600 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum to give 129 g of crude 6-[(tert-butoxy)carbonyl]-6-azaspiro[2.5]octane-1-carboxylic acid as a white solid. TLC: DCM/MeOH = 5:1, R_f = 0.1.

Step 4. tert-Butyl (1S)-1-[[[(1S)-1-phenylethyl]carbamoyl]-6-azaspiro[2.5]octane-6-carboxylate.



To a stirred solution of 6-[(tert-butoxy)carbonyl]-6-azaspiro[2.5]octane-1-carboxylic acid (129 g, 505.27 mmol, 1.00 equiv), DIPEA (196 g, 1.52 mol, 6.02 equiv) and (S)-(-)-1-phenylethylamine (87 g, 717.94 mmol, 1.10 equiv) in DMF (2 L) 0°C was added *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (231 g, 607.89 mmol, 1.20 equiv) in several portions. The resulting solution was stirred at rt overnight and then was quenched by the addition of 4.5 L of water/ice. The resulting mixture was extracted with 3x2 L of ethyl acetate. The combined organic layers were washed with 2x800 mL of brine, dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1/4) to give 83.15 g (46%) of tert-butyl 1-[[[(1S)-1-phenylethyl]carbamoyl]-6-azaspiro[2.5]octane-6-carboxylate as a mixture of diastereomers as a white solid. LC/MS (Method 1, ESI): RT_1 = 4.63 min, RT_2 = 4.73 min, m/z = 359.1 [M+H]⁺. The resulting mixture of diastereomers was separated by chiral supercritical fluid chromatography (Column: 3 x 25 cm, 5 μm Chiralpak AD; Mobile phase A: CO₂; Mobile phase B: 0.1% NH₄OH in MeOH; Isocratic conditions: 87:13 A:B; Flow rate: 200 g/min; UV: 220 nm; Back pressure: 100 bar, Column temperature: 40 °C). The first eluting diastereomer was carried forward into the next step.

Step 5. (1S)-N-[(1S)-1-Phenylethyl]-6-azaspiro[2.5]octane-1-carboxamide.
Hydrogen chloride gas was bubbled into a solution of tert-butyl (1S)-1-[[[(1S)-1-

phenylethyl]carbamoyl]-6-azaspiro[2.5]octane-6-carboxylate (20 g, 55.79 mmol, 1.00 equiv) in MeOH (200 mL) at 0°C for 30 min. The resulting solution was stirred overnight at rt and then concentrated under vacuum. The residue was diluted with 100 mL of H₂O and the pH of the solution was adjusted to 12 with 2N sodium hydroxide solution. The resulting mixture was extracted with 3x100 mL of DCM. The combined organic layers was dried over anhydrous sodium sulfate and concentrated under vacuum to give 15 g of (1S)-N-[(1S)-1-phenylethyl]-6-azaspiro[2.5]octane-1-carboxamide as a light yellow solid. LC/MS (Method I, ESI): RT = 1.03 min, *m/z* = 259.0 [M-H]⁻.

10 Step 6. tert-Butyl (1S)-1-([(1S)-1-phenylethyl]amino)methyl)-6-azaspiro[2.5]octane-6-carboxylate. To a stirred solution of (1S)-N-[(1S)-1-phenylethyl]-6-azaspiro[2.5]octane-1-carboxamide (8 g, 30.96 mmol, 1.00 equiv) in THF (800 mL) maintained under nitrogen at rt was added a 1 M solution of borane in THF (100 mL, 100 mmol, 3.3 equiv) dropwise. The resulting solution was refluxed overnight. The resulting mixture was concentrated under vacuum then water (50 mL) was added to the residue. The pH of the solution was adjusted to 1 with 5% HCl (5%). The resulting solution was stirred at reflux for 4 h. The reaction mixture was cooled to rt and the pH of the solution was adjusted to 12 with a 10% sodium hydroxide solution. The solution was cooled to 0°C then a solution of di-tert-butyl dicarbonate (6.7 g, 30.70 mmol, 0.68 equiv) in 20 mL of THF was added dropwise with stirring. The reaction mixture was stirred overnight at rt. The resulting mixture was extracted with 2x200 mL of ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:3) to give 2.9 g (19%) of tert-butyl (1S)-1-([(1S)-1-phenylethyl]amino)methyl)-6-azaspiro[2.5]octane-6-carboxylate as light brown oil. T.L.C: DCM:MeOH=10:1, R_f = 0.3.

30 Step 7. tert-Butyl (1S)-1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate. A mixture of tert-butyl (1S)-1-([(1S)-1-phenylethyl]amino)methyl)-6-azaspiro[2.5]octane-6-carboxylate (700 mg, 2.03 mmol, 1.00 equiv) and Pd(OH)₂ (100 mg) in MeOH (30 mL) was stirred under 5 atmosphere of H₂ in a 50-mL

pressure reactor overnight at rt. The catalyst was removed by filtration. The filtrate was concentrated under vacuum to afford 400 mg (82%) of tert-butyl (1S)-1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate as a light yellow oil. LC/MS (Method F, ESI): RT = 1.25 min, $m/z = 241.0$ [M+H].

5 Step 8. tert-Butyl (1S)-1-[[[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl)carbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate. A solution of tert-butyl (1S)-1-(aminomethyl)-6-azaspiro[2.5]octane-6-carboxylate (5 g, 20.80 mmol, 1.00 equiv), 4-nitrophenyl chloroformate (4.4 g, 21.83 mmol, 1.05 equiv) and DIPEA (10 mL) in THF (200 mL) was stirred overnight at rt. 1H,2H,3H-Pyrrolo[3,4-c]pyridine dihydrochloride (4.4 g, 22.79 mmol, 9.24 equiv) was then added and the
10 reaction mixture was stirred overnight at rt. The resulting mixture was concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (20/1) to give 2.3 g of tert-butyl (1S)-1-[[[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl)carbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate as a light
15 yellow solid. TLC: DCM:MeOH = 10:1, $R_f = 0.4$.

Step 9. N-[(1S)-6-Azaspiro[2.5]octan-1-ylmethyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide hydrochloride. A solution of tert-butyl (1S)-1-[[[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl)carbonyl]amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate (2.3 g, 5.95 mmol, 1.00 equiv) in a saturated
20 hydrogen chloride in MeOH solution (100 mL) was stirred at rt for 2 h. The resulting mixture was concentrated under vacuum to yield 1.7 g of crude N-[(1S)-6-azaspiro[2.5]octan-1-ylmethyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide hydrochloride as a brown solid.

Step 10. 3-Methyloxetan-3-yl 4-nitrophenyl carbonate. To a stirred solution
25 of oxetan-3-one (9 g, 124.89 mmol, 1.00 equiv) in THF (200 mL) maintained under nitrogen at -50°C was added dropwise a 1.6M solution of methyl lithium in ether (150 mL, 240 mmol, 1.9 equiv). The reaction mixture was warmed to 0°C and stirred for 2 h. A solution of 4-nitrophenyl chloroformate (26 g, 128.99 mmol, 1.03 equiv) in THF (100 mL) was then added dropwise at 0°C. The resulting solution was stirred for
30 another 2 h at rt after the addition was completed. The reaction was then quenched by the addition of 200 mL of water. The resulting mixture was extracted with 2x200 mL

of ethyl acetate. The combined organic layers was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was purified on a silica gel column eluted with ethyl acetate/petroleum ether (1:5) to give 4.5 g (14%) of 3-methyloxetan-3-yl 4-nitrophenyl carbonate as a white solid. TLC: petroleum ether:ethyl acetate = 2:1, $R_f = 0.6$.

5 Step 11. A solution of N-[(1S)-6-azaspiro[2.5]octan-1-ylmethyl]-1H,2H,3H-pyrrolo[3,4-c]pyridine-2-carboxamide hydrochloride (1.7 g, 5.27 mmol, 1.00 equiv), 3-methyloxetan-3-yl 4-nitrophenyl carbonate (1.7 g, 6.71 mmol, 1.27 equiv) and DIPEA (2 mL) in ethanol (50 mL) was stirred overnight at rt. The resulting mixture
10 was concentrated under vacuum. The residue was dissolved in 150 mL of DCM and then washed with 100 mL of H₂O. The organic layer was dried over anhydrous sodium sulfate then concentrated under vacuum. The residue was purified on a silica gel column eluted with DCM/MeOH (20/1) to give 1.43 g (68%) of 3-methyloxetan-3-yl (1S)-1-[[[(1H,2H,3H-pyrrolo[3,4-c]pyridin-2-yl)carbonyl)amino]methyl]-6-
15 azaspiro[2.5]octane-6-carboxylate as a white solid. LC/MS (Method I, ESI): RT= 1.24 min, $m/z = 401.1$ [M+H]⁺. ¹H-NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.56 (s, 1H), 8.47 (d, $J = 5.1$ Hz, 1H), 7.39 (d, $J = 5.1$ Hz, 2H), 6.42 (t, $J = 5.2$ Hz, 1H), 4.63-4.59 (m, 6H), 4.38 (d, $J = 7.5$ Hz, 2H), 3.48 (s, 2H), 3.31 (s, 2H), 3.12 (t, $J = 6.0$ Hz, 2H), 1.62 (s, 3H), 1.59-1.55 (m, 1H), 1.42-1.39 (m, 2H), 1.37-1.34 (m, 1H), 1.21-1.17 (m, 1H),
20 1.01-0.96 (m, 1H), 0.54-0.48 (m, 1H), 0.27-0.21 (m, 1H).

Additional example compounds were prepared using methods analogous to those described above. Particular example compounds were prepared using methods analogous to those indicated for the Example numbers listed below:

Ex.	Method of Example No.
22	6
23	6
24	6
25	1
26	339
27	339
28	339
29	339
30	339

31	339
32	339
33	339
34	339
35	339
36	339
37	339
38	339
39	339
40	6
41	383
42	383
43	339
44	339
45	1 and 6
46	6
47	1 and 6
48	1
49	1
50	339
51	339
52	339
53	339
54	339
55	339
56	339
57	339
58	339
59	339
60	339
61	339
62	339
63	339
64	339
65	339
66	339
67	339
68	339
69	339
70	339
71	339
72	339
73	339
74	339
75	339

76	339
77	339
78	339
79	339
80	339
81	339
82	339
83	339
84	339
85	339
86	339
87	339
88	339
89	339
90	339
91	339
92	339
93	339
94	339
95	339
96	339
97	339
98	339
99	339
100	339
101	339
102	339
103	339
104	339
105	339
106	339
107	339
108	339
109	339
110	339
111	339
112	339
113	339
114	339
115	339
116	339
117	339
118	339
119	339
120	339

121	339
122	339
123	339
124	339
125	339
126	339
127	339
128	339
129	339
130	339
131	339
132	339
133	339
134	339
135	339
136	339
137	339
138	339
139	339
140	339
141	339
142	339
143	339
144	339
145	339
146	339
147	339
148	6
149	6
151	1
152	339
153	339
154	339
155	339
156	339
157	159
158	159
160	159
161	159
162	159
163	159
164	159
165	159
166	159
167	159

168	159
169	159
170	159
171	159
172	159
173	159
175	150
176	1 and 6
177	1 and 6
178	1 and 6
179	339
180	339
181	339
182	159
183	159
184	159
185	159
186	159
187	159
188	1 and 6
189	1 and 6
190	1 and 6
191	6
192	6
193	1
194	1
195	150
196	6
197	6
198	6
199	6
200	1
201	1 and 6
202	1 and 6
203	6
204	419
205	419
206	6
207	1 and 6
208	6
209	6
210	6
211	6
212	1
213	6

214	6
215	419
216	6
218	1
219	150
220	150
221	150
222	339
223	339
228	1
229	6
230	1 and 6
231	339
232	339
233	339
234	339
235	339
236	339
237	339
238	339
239	339
240	339
241	339
242	339
243	339
244	339
245	339
246	339
247	339
248	339
249	339
250	339
251	339
252	339
253	339
254	339
255	339
256	339
257	339
258	339
259	339
260	339
261	339
262	339
263	339

264	339
265	339
266	339
267	339
268	339
269	339
270	339
271	339
272	339
273	339
275	6
276	174
277	6
278	6
279	1 and 6
280	339
281	339
282	339
283	339
284	339
285	6
286	6
287	6
288	6
289	6
290	6
291	6
292	1 and 6
293	6
294	339
295	339
296	339
297	339
298	339
299	339
300	339
301	339
302	339
303	339
304	339
305	339
306	339
307	339
308	339
309	339

310	339
311	339
312	339
313	339
314	339
315	339
316	339
317	339
318	339
319	339
320	339
321	339
322	339
323	339
324	339
325	339
326	339
327	339
328	339
329	339
330	339
331	339
332	339
333	339
334	339
335	339
336	339
337	339
338	339
340	339
341	339
343	6
344	6
345	6
346	1 and 6
347	1 and 6
348	6
350	1
351	1
352	1 and 6
353	1 and 6
354	1 and 6
355	1 and 6
356	6
357	1 and 6

358	6
359	6
360	1 and 6
361	1
362	339
363	339
364	6
365	1 and 6
366	1 and 6
367	409
368	6
369	6
370	6
371	6
372	6
373	6
375	1
376	1 and 6
377	6
378	6
379	6
381	6
382	1
384	339
385	339
387	6
388	1
389	6
392	1
393	1 and 6
395	1
396	1 and 6
397	6
398	1 and 6
399	6
400	174
401	6
402	6
404	1 and 6
407	1 and 6
408	409
410	409
411	6
413	419
414	419

415	1 and 6
416	1 and 6
417	1 and 6
418	1 and 6

Additional examples were prepared using methods analogous to those described above.

5 Analytical Characterization:

Each of the specifically exemplified compounds described herein was prepared using the methods analogous to those described above, and were analyzed by LC/MS. Data for each compound, along with the LC/MS method used to generate the data, is provided in Table 1 (NA = not available).

10

Table 1. LC/MS Data for Example Compounds.

Ex.	Retention time (min)	<i>m z</i>	Method
2	1.63	402.0	E
3	3.43	386.1	B
4	1.60	386.0	E
5	3.60	385.1	B
7	3.44	403.0	C
8	3.19	385.2	L
9	1.40	395.0	H
10	1.16	397.0	H
11	1.43	398.2	C
12	1.24	355.0	H
13	2.45	425.2	C
14	1.21	377.0	C
15	4.25	386.0	L
16	2.39	327.3	J
17	1.27	363.0	A
18	2.23	285.1	M
19	2.41	286.1	M
20	1.71	341.2	C
21	2.62	399.0	A
22	3.59	385.1	B
23	2.63	386.1	I
24	1.37	383.2	C
25	1.07	387.3	O
26	2.59	394.2	T

27	3.60	410.2	T
28	3.80	482.2	T
29	4.19	384.3	T
30	2.83	426.3	T
31	4.06	429.2	T
32	3.79	368.2	T
33	2.99	397.2	T
34	3.66	409.2	T
35	3.18	411.3	T
36	3.66	412.3	T
37	3.61	398.2	T
38	3.93	400.2	T
39	3.02	405.2	T
40	1.04	482.3	Q
41	4.46	385.2	T
42	4.45	385.3	T
43	4.50	386.2	T
44	4.49	386.2	T
45	1.35	385	I
46	1.95	369	B
47	1.31	405	I
48	4.36	387.3	T
49	4.35	387.3	T
50	3.48	406.2	T
51	3.57	413.3	T
52	3.94	412.3	T
53	3.15	411.3	T
54	3.85	413.2	T
55	3.48	413.2	T
56	3.39	411.2	T
57	3.45	411.2	T
58	3.91	390.2	T
59	3.54	384.2	T
60	3.63	398.2	T
61	3.58	384.2	T
62	3.18	371.2	T
63	3.64	436.3	T
64	3.34	426.3	T
65	3.04	427.3	T
66	3.41	432.2	T
67	3.53	422.3	T
68	3.44	425.3	T
69	3.87	468.2	T
70	3.60	457.2	T
71	3.79	475.2	T

72	3.70	447.2	T
73	3.79	475.2	T
74	3.65	475.2	T
75	2.95	427.3	T
76	3.54	461.2	T
77	3.88	459.2	T
78	3.36	440.3	T
79	3.36	440.3	T
80	3.46	446.2	T
81	3.81	472.2	T
82	3.52	439.3	T
83	3.81	472.2	T
84	3.79	483.2	T
85	3.87	523.2	T
86	3.29	502.3	T
87	3.41	502.3	T
88	3.54	502.3	T
89	2.91	440.3	T
90	3.70	439.3	T
91	2.79	426.3	T
92	3.81	468.2	T
93	3.85	468.2	T
94	3.52	448.2	T
95	2.91	454.3	T
96	3.84	486.3	T
97	3.12	391.2	T
98	3.56	391.2	T
99	3.49	381.2	T
100	3.19	381.2	T
101	2.97	380.2	T
102	3.41	392.2	T
103	3.21	391.2	T
104	3.44	380.2	T
105	2.98	394.2	T
106	3.43	394.2	T
107	3.43	394.2	T
108	3.62	394.2	T
109	3.59	394.2	T
110	3.33	392.2	T
111	3.39	392.2	T
112	3.46	392.2	T
113	3.32	392.2	T
114	3.49	394.2	T
115	3.58	408.2	T
116	3.76	408.2	T

117	3.72	395.2	T
118	4.09	396.2	T
119	3.12	405.2	T
120	3.10	405.2	T
121	3.15	405.2	T
122	3.65	406.2	T
123	3.85	420.2	T
124	3.74	422.2	T
125	3.72	409.2	T
126	3.65	408.2	T
127	3.77	416.2	T
128	3.70	409.2	T
129	3.37	476.3	T
130	3.73	354.2	T
131	4.30	462.2	T
132	3.75	461.2	T
133	3.51	448.2	T
134	3.42	448.2	T
135	3.70	448.2	T
136	3.91	423.2	T
137	3.88	422.2	T
138	3.73	422.2	T
139	3.65	395.2	T
140	3.80	436.3	T
141	3.37	380.2	T
142	3.82	489.3	T
143	2.95	413.3	T
144	3.73	439.3	T
145	3.37	411.2	T
146	3.27	461.3	T
147	3.74	489.3	T
148	1.29	453	I
149	1.74	425	I
150	1.88	397.1	K
151	0.97	373.2	S
152	4.10	429.2	T
153	3.44	372.2	T
154	4.64	536.2	T
155	3.38	413.3	T
156	2.97	405.1	T
157	4.07	430.2	T
158	4.24	433.3	T
159	4.20	419.2	T
160	4.03	419.2	T
161	4.08	437.2	T

162	4.24	433.3	T
163	4.09	437.2	T
164	4.13	449.2	T
165	4.05	449.2	T
166	4.09	437.2	T
167	4.03	449.2	T
168	4.51	489.2	T
169	4.71	463.3	T
170	4.31	453.2	T
171	4.25	453.2	T
172	3.47	357.2	T
173	4.53	489.2	T
174	1.66	412	I
175	2.58	399	B
176	1.67	406.1	B
177	2.32	406.2	K
178	2.1	429.9	C
179	3.71	457.2	T
180	3.13	408.2	T
181	3.48	422.2	T
182	3.83	385.3	T
183	3.59	371.2	T
184	4.35	425.3	T
185	3.63	371.2	T
186	4.12	411.3	T
187	3.98	405.2	T
188	1.09	454.1	C
189	1.9	371.1	B
190	1.61	406.1	B
191	1.52	340.1	K
192	1.21	359.2	F
193	1.55	399.2	G
194	1.57	344.1	G
195	1.39	398.1	K
196	2.05	411.3	F
197	1.6	425	C
198	0.87	414.2	F
199	1.43	418.9	C
200	1.49	401.2	I
201	1.33	441.3	F
202	3.49	473.1	B
203	1.31	384.2	I
204	3.10	396.6	T
205	4.13	430.2	T
206	1.33	428	I

207	1.88	430	I
208	1.92	406.2	F
209	1.04	441.9	C
210	1.29	420	I
211	1.98	440.9	C
212	2.04	386.3	F
213	4.11	383.3	T
214	4.12	383.3	T
215	3.83	402.2	T
216	1.32	428	I
218	2.12	439.2	C
219	1.27	413.2	L
220	1.06	423.3	F
221	1.21	395.2	R
222	3.74	450.3	T
223	4.12	502.3	T
228	1.41	399.2	Q
229	3.24	428.2	B
230	2.24	441	I
231	3.42	405.2	T
232	3.10	405.2	T
233	2.98	405.2	T
234	3.35	405.2	T
235	3.42	405.2	T
236	3.18	430.2	T
237	3.75	430.2	T
238	3.43	431.2	T
239	3.32	430.2	T
240	3.59	430.2	T
241	2.93	430.2	T
242	3.78	415.2	T
243	3.76	415.2	T
244	3.90	502.2	T
245	3.12	489.3	T
246	3.24	484.3	T
247	4.36	474.2	T
248	3.26	473.3	T
249	3.56	471.2	T
250	3.59	471.2	T
251	3.52	471.3	T
252	3.21	470.3	T
253	3.09	456.2	T
254	3.47	458.2	T
255	3.14	458.2	T
256	3.96	459.2	T

257	3.96	459.2	T
258	3.64	461.2	T
259	3.41	469.2	T
260	3.36	470.2	T
261	3.61	442.2	T
262	3.21	442.2	T
263	3.07	444.2	T
264	3.55	447.2	T
265	3.00	448.1	T
266	3.94	432.2	T
267	3.28	442.2	T
268	3.00	442.2	T
269	3.88	442.2	T
270	3.50	442.2	T
271	3.36	442.2	T
272	3.44	381.2	T
273	3.85	370.2	T
274	1.49	376.2	Q
275	1.7	439	A
276	1.28	413.2	R
277	1.34	404.2	J
278	2.56	404.1	B
279	1.62	409.2	B
280	3.89	472.2	T
281	3.09	419.2	T
282	3.36	384.2	T
283	3.43	400.2	T
284	3.39	386.2	T
285	1.71	404.1	B
286	0.92	407.1	R
287	1.42	407.2	R
288	1.02	426.1	R
289	1.78	421.2	R
290	1.15	427.3	C
291	0.9	404.2	R
292	1.6	409.1	B
293	0.98	421.2	R
294	3.91	464.3	T
295	4.04	498.3	T
296	4.15	450.3	T
297	3.58	408.2	T
298	3.94	450.3	T
299	3.59	408.2	T
300	3.43	422.2	T
301	3.43	422.2	T

302	3.77	430.2	T
303	3.22	491.3	T
304	3.35	491.3	T
305	3.82	475.2	T
306	4.03	521.3	T
307	3.04	463.2	T
308	3.88	450.3	T
309	3.14	444.2	T
310	3.83	444.2	T
311	3.19	444.2	T
312	3.88	448.2	T
313	3.34	431.2	T
314	3.47	436.3	T
315	3.39	444.2	T
316	3.62	430.2	T
317	3.00	430.2	T
318	4.53	423.2	T
319	4.13	436.3	T
320	3.19	431.2	T
321	3.42	431.2	T
322	3.92	436.3	T
323	3.76	422.2	T
324	3.49	422.2	T
325	3.67	422.2	T
326	3.73	422.2	T
327	3.94	421.2	T
328	4.41	421.1	T
329	3.15	408.2	T
330	3.29	408.2	T
331	3.39	395.2	T
332	3.41	420.2	T
333	3.65	420.2	T
334	3.57	409.2	T
335	3.35	408.2	T
336	3.23	381.2	T
337	3.39	381.2	T
338	2.63	380.2	T
339	1.12	488.3	F
340	3.12	431.2	T
341	3.49	305.2	T
343	0.99	501.3	C
344	1.4	501.3	C
345	1.02	446.3	C
346	1.42	503.4	F
347	2.51	409.2	B

348	1.37	381	I
350	2.11	401.2	F
351	1.16	403.3	F
352	0.94	406.3	F
353	1.87	421.3	F
354	1.11	462.3	F
355	1.14	410.2	F
356	1.37	440.4	C
357	1.99	430.3	F
358	1.24	466	C
359	0.76	412.2	O
360	2.85	401.2	B
361	2.38	400.2	O
362	2.02	489.4	F
363	1.03	487.3	F
364	2.24	381.2	F
365	1.06	423.3	F
366	1.06	423.3	F
367	2	391.2	K
368	1.29	419	I
369	1.35	460.2	O
370	0.74	408.3	S
371	1.76	423.2	O
372	1.2	517.4	F
373	1.75	382.2	O
375	1.23	415.3	F
376	1.78	403.3	F
377	2.03	383.3	O
378	1.78	420.1	K
379	1.64	419	I
381	2.43	421.2	K
382	0.9	411.2	O
383	1.48	386.0	S
384	4.65	409.2	U
385	4.76	409.2	U
387	1.3	415.2	K
388	2.68	502.4	C
389	1.24	405.3	C
392	1.42	414.2	F
393	1.68	417.3	F
394	1.01	421.1	C
395	1.09	400.2	F
396	1.3	449.3	F
397	1.85	399.3	F
398	1.44	460.2	O

399	1.19	428	F
400	1.76	411	I
401	1.41	401	I
402	1.71	386.3	P
403	1.15	508	I
404	1.16	426	I
405	1.25	488	I
406	1.65	402.2	K
407	1.6	407	I
408	1.27	375	I
409	1.39	405	I
410	1.15	414	I
411	2.19	390	I
413	1.26	371.2	I
414	1.3	371.2	I
415	1.25	396.2	F
416	1.58	372.2	F
417	1.33	400.1	I
418	1.43	391.1	I
419	1.24	401.1	I

It is understood that the person skilled in the art will be able to prepare the compounds of the present invention using methods known in the art along with the general method of synthesis described herein.

5

Assay 1: Biochemical Inhibition Assay

NAMPT protein purification. Recombinant His-tagged NAMPT was produced in *E.coli* cells, purified over a Ni column, and further purified over a size-exclusion column by XTAL Biostructures.

10 The NAMPT enzymatic reaction. The NAMPT enzymatic reactions were carried out in Buffer A (50 mM Hepes pH 7.5, 50 mM NaCl, 5 mM MgCl₂, and 1 mM THP) in 96-well V-bottom plates. The compound titrations were performed in a separate dilution plate by serially diluting the compounds in DMSO to make a 100X stock. Buffer A (89 μ L) containing 33 nM of NAMPT protein was added to 1 μ L of
 15 100X compound plate containing controls (e.g. DMSO or blank). The compound and enzyme mixture was incubated for 15 min at room temperature, then 10 μ L of 10X substrate and co-factors in Buffer A were added to the test well to make a final

concentration of 1 μ M NAM, 100 μ M 5-Phospho-D-ribose 1-diphosphate (PRPP), and 2.5 mM Adenosine 5'-triphosphate (ATP). The reaction was allowed to proceed for 30 min at room temperature, then was quenched with the addition of 11 μ L of a solution of formic acid and L-Cystathionine to make a final concentration of 1% formic acid and 10 μ M L-Cystathionine. Background and signal strength was determined by addition (or non-addition) of a serial dilution of NMN to a pre-quenched enzyme and cofactor mix.

Quantification of NMN. A mass spectrometry-based assay was used to measure the NAMPT reaction product, β -nicotinamide mononucleotide (NMN), and the internal control (L-Cystathionine). NMN and L-Cystathionine were detected using the services of Biocius Lifesciences with the RapidFire system. In short, the NMN and L-Cystathionine were bound to a graphitic carbon cartridge in 0.1% formic acid, eluted in 30% acetonitrile buffer, and injected into a Sciex 4000 mass spectrometer. The components of the sample were ionized with electrospray ionization and the positive ions were detected. The Q1 (parent ion) and Q3 (fragment ion) masses of NMN were 334.2 and 123.2, respectively. The Q1 and Q3 for L-Cystathionine were 223.1 and 134.1, respectively. The fragments are quantified and the analyzed by the following method.

Determination of IC₅₀ Values. First, the NMN signal was normalized to the L-Cystathionine signal by dividing the NMN signal by the L-Cystathionine signal for each well. The signal from the background wells were averaged and subtracted from the test plates. The compound treated cells were then assayed for percent inhibition by using this formula:

$$\% \text{ Inh} = 100 - 100 * x/y$$

wherein x denotes the average signal of the compound treated wells and y denotes the average signal of the DMSO treated wells.

IC₅₀ values were then determined using the following formula:

$$IC_{50} = 10^{(\text{LOG}_{10}(X) + ((50 - \% \text{ Inh at Cmpd Concentration 1}) / (XX - YY)) * (\text{LOG}_{10}(X) - \text{LOG}_{10}(Y)))}$$

wherein X denotes the compound concentration 1, Y denotes the compound concentration 2, XX denotes the % inhibition at compound concentration 1 (X), and YY denotes the % inhibition at compound concentration 2 (Y).

The compounds of this invention have IC₅₀ values that are preferably under 1 μM, more preferably under 0.1 μM, and most preferably under 0.01 μM. Results for the compounds tested in this assay are provided in Table 2 below.

Assay 2: In-Vitro Cell Proliferation Assay

Assay Method. A2780 cells were seeded in 96-well plates at 1×10^3 cells/well in 180 μL of culture medium (10% FBS, 1% Pen/Strep Amphotericin B, RPMI-1640) with and without the addition of either NMN or nicotinamide (NAM). After overnight incubation at 37 °C and 5% CO₂, the compound titrations were performed in a separate dilution plate by serially diluting the compounds in DMSO to make a 1000X stock. The compounds were then further diluted to 10X final concentration in culture media, whereupon 20 μL of each dilution was added to the plated cells with controls (e.g. DMSO and blank) to make a final volume of 200 μL. The final DMSO concentration in each well was 0.1%. The plates were then incubated for 72 hours at 37 °C in a 5% CO₂ incubator. The number of viable cells was then assessed using sulforhodamine B (SRB) assay. Cells were fixed at 4 °C for 1 hour with the addition of 50 μL 30% trichloroacetic acid (TCA) to make a final concentration of 6 % TCA. The plates were washed four times with H₂O and allowed to dry for at least 1 hour, whereupon 100 μL of a 4% SRB in 1% acetic acid solution was added to each well and incubated at room temperature for at least 30 min. The plates were then washed three times with 1% acetic acid, dried, and treated with 100 μL of 10mM Tris-Base solution. The plates were then read in a microplate reader at an absorbance of 570 nm. Background was generated on a separate plate with media only.

Determination of IC₅₀ Values. First, the signals from the background plate were averaged, then the background was subtracted from the test plates. The compound-treated cells were then assayed for % inhibition by using the following formula:

$$\% \text{ Inh} = 100 - 100 * x/y$$

wherein x denotes the average signal of the compound-treated cells and y denotes the average signal of the DMSO-treated cells.

IC₅₀ values were then determined using the following formula:

$$5 \quad \text{IC}_{50} = 10^{(\text{LOG}_{10}(\text{X}) + (((50 - \% \text{ Inh at Cmpd Concentration 1}) / (\text{XX} - \text{YY})) * (\text{LOG}_{10}(\text{X}) - \text{LOG}_{10}(\text{Y}))))}$$

wherein X denotes the compound concentration 1, Y denotes the compound concentration 2, XX denotes the % inhibition at compound concentration 1 (X), and YY denotes the % inhibition at compound concentration 2 (Y).

10 Specificity of cytotoxicity. Inhibition of NAMPT could be reversed by the addition of NAM or NMN. The specificity of the compounds were determined *via* cell viability assay in the presence of the compound and either NAM or NMN. Percent inhibitions were determined using the method given above.

The compounds of this invention have IC₅₀ values that are preferably under 15 1 μM, more preferably under 0.1 μM, and most preferably under 0.01 μM. Most preferable compounds of this invention are compounds that have IC₅₀ values in the enzymatic assay and the cell proliferation assay that are both under 1 μM, more preferably both of the values are under 0.1 μM, and most preferably both of the values are under 0.01 μM. Results for the compounds tested in this assay are 20 provided in Table 2 (NT = not tested).

Table 2. Biochemical and Cell Proliferation Assay Results.

Ex.	Biochemical (IC ₅₀) [μM]	Cell Proliferation (IC ₅₀) [μM]
1	0.00432	0.000775
2	0.00719	0.202
3	0.00896	0.0697
4	0.00953	0.0489
5	0.0183	0.119
6	0.0454	0.00533
7	0.0648	0.0962
8	0.0735	0.208
9	0.108	0.221
10	0.208	0.791
11	0.255	2.000
12	0.398	2.000

13	0.423	2.000
14	0.988	2.000
15	2.000	NT
16	2.000	2.000
17	2.000	2.000
18	NT	NT
19	NT	NT
20	0.39832	2.000
21	0.175	2.000
22	0.0175	0.119
23	0.0100	0.0325
24	0.0511	0.00829
25	0.0068	0.0013
26	0.304	2.0
27	0.0504	0.842
28	0.0642	0.616
29	0.0111	0.00303
30	0.416	0.614
31	0.0373	0.0121
32	0.153	2.0
33	1.12	2.0
34	0.0542	0.379
35	0.563	2.0
36	0.0893	0.185
37	0.155	0.643
38	0.040	0.0188
39	0.047	0.232
40	0.13	0.0243
41	0.0143	0.0861
42	0.143	2.0
43	0.279	2.0
44	0.0199	0.0289
45	0.00623	0.00158
46	0.12	0.402
47	0.00459	0.0126
48	0.00996	0.000995
49	0.0479	0.0335
50	0.309	2.0
51	0.173	0.367
52	0.135	0.674
53	0.42	0.846
54	2.0	2.0
55	1.06	2.0
56	0.575	2.0
57	0.223	1.6

58	0.422	2.0
59	0.782	2.0
60	0.309	1.53
61	0.768	2.0
62	0.395	2.0
63	0.177	1.39
64	0.398	2.0
65	0.526	0.931
66	1.04	2.0
67	0.21	0.193
68	0.812	0.414
69	0.386	0.194
70	0.0681	0.492
71	0.0641	0.0137
72	0.0338	0.0481
73	0.0574	0.0258
74	0.231	0.169
75	0.109	0.0809
76	0.214	0.228
77	0.169	1.15
78	0.296	2.0
79	0.755	2.0
80	0.244	1.34
81	0.626	2.0
82	0.398	0.71
83	0.211	1.08
84	0.0767	0.185
85	0.0388	0.0142
86	0.030	0.00453
87	0.802	2.0
88	0.022	0.00977
89	0.228	0.116
90	0.149	0.183
91	0.474	2.0
92	0.189	0.152
93	0.475	0.878
94	0.0706	0.437
95	0.426	2.0
96	0.153	0.711
97	0.421	2.0
98	0.504	2.0
99	0.826	2.0
100	0.462	2.0
101	0.625	2.0
102	1.21	2.0

103	0.44	2.0
104	2.92	2.0
105	0.246	0.755
106	0.396	0.743
107	0.372	2.0
108	0.216	0.449
109	0.0993	0.299
110	0.861	2.0
111	0.579	2.0
112	0.417	2.0
113	0.877	2.0
114	0.325	2.0
115	0.0499	0.0376
116	0.365	0.37
117	0.422	0.752
118	0.449	1.07
119	0.188	0.446
120	0.207	0.0928
121	0.308	1.09
122	0.0974	2.0
123	0.0661	0.083
124	0.151	0.124
125	0.0955	0.221
126	0.273	0.833
127	0.269	2.0
128	0.0914	0.0292
129	0.10	0.0373
130	2.0	2.0
131	0.356	0.326
132	0.0432	0.0072
133	0.194	0.696
134	0.0477	0.514
135	0.0509	0.0364
136	0.104	0.169
137	0.252	0.116
138	0.0941	2.0
139	0.163	2.0
140	0.78	2.0
141	0.222	2.0
142	0.116	0.0971
143	1.22	2.0
144	0.116	0.375
145	0.494	2.0
146	0.0801	0.0745
147	0.0553	0.0744

148	0.288	0.102
149	0.768	0.496
150	0.31	0.265
151	0.0148	0.00423
152	0.0601	0.262
153	0.686	2.0
154	0.208	2.61
155	0.261	2.0
156	NT	NT
157	0.0752	2.0
158	0.41	2.0
159	0.0332	0.274
160	0.951	2.0
161	0.123	2.0
162	0.441	2.0
163	0.559	2.0
164	0.118	1.54
165	0.309	2.0
166	0.406	2.0
167	0.314	2.0
168	0.113	2.0
169	0.0558	0.0278
170	0.161	2.0
171	0.0604	2.0
172	0.422	2.0
173	0.146	2.0
174	0.238	0.197
175	0.0467	0.0111
176	0.0376	0.0341
177	0.0363	0.0344
178	0.0257	0.0323
179	0.034	0.0254
180	2.0	2.46
181	0.0173	0.00331
182	0.255	2.0
183	0.314	2.0
184	0.422	2.0
185	0.502	2.0
186	0.0972	0.732
187	0.0446	2.46
188	0.0301	0.0453
189	0.0128	0.00438
190	0.0312	0.0507
191	0.156	1.04
192	0.0239	0.0294

193	0.212	2.0
194	0.668	2.0
195	0.0785	0.218
196	0.0144	0.00367
197	0.0111	0.00961
198	0.131	0.0618
199	0.0144	0.184
200	0.00456	0.0247
201	0.0173	0.0279
202	0.012	0.0282
203	0.544	2.0
204	0.00636	0.00336
205	0.108	0.13
206	0.263	2.0
207	0.0491	0.136
208	0.0347	0.0261
209	0.0803	0.203
210	0.0588	0.316
211	0.0278	0.131
212	0.0528	0.0207
213	0.0318	0.00578
214	0.936	2.0
215	0.0751	0.0658
216	0.179	1.34
218	0.00997	0.00159
219	0.0151	0.00531
220	0.0034	0.00622
221	0.0178	0.0254
222	0.0504	0.0117
223	0.027	0.00192
228	0.0222	0.00474
229	0.0548	0.0184
230	0.0185	0.0299
231	0.201	0.516
232	0.0983	0.207
233	0.0458	0.288
234	0.161	0.164
235	0.288	1.56
236	0.0339	0.0473
237	0.0501	0.166
238	0.0317	0.91
239	0.0329	0.0297
240	0.0263	0.0187
241	0.0273	0.0702
242	0.134	0.713

243	0.0736	0.434
244	0.0264	0.0261
245	0.0525	0.0126
246	0.0211	0.0138
247	0.205	2.0
248	0.0221	0.00439
249	0.254	0.935
250	0.549	2.0
251	0.0776	0.215
252	0.0695	0.344
253	0.0417	0.0554
254	0.0191	0.013
255	0.0499	0.179
256	0.147	0.596
257	0.313	2.0
258	0.0314	0.0203
259	0.161	1.2
260	0.916	2.0
261	0.0245	0.0158
262	0.0695	0.0795
263	0.0265	0.0254
264	0.0683	0.245
265	2.23	2.0
266	0.127	1.47
267	0.321	2.0
268	0.0691	0.0719
269	0.0277	0.016
270	0.0192	0.0192
271	0.0964	0.139
272	0.335	2.0
273	0.0178	0.207
274	0.384	2.0
275	0.121	0.0177
276	0.0391	0.0132
277	0.445	0.755
278	0.479	2.0
279	0.0934	0.00745
280	0.0152	0.00683
281	0.056	0.0185
282	2.0	2.0
283	0.362	0.362
284	0.106	0.105
285	2.0	2.0
286	0.027	0.027
287	0.0245	0.0245

288	0.0606	0.0606
289	0.00999	0.00999
290	0.00454	0.00454
291	0.722	0.722
292	0.693	0.693
293	0.0593	0.0593
294	0.0486	0.0121
295	0.0145	0.000944
296	0.0871	0.0108
297	0.0925	0.278
298	0.213	0.103
299	0.165	0.249
300	1.13	0.335
301	0.0425	0.00421
302	0.0648	0.0903
303	0.478	2.0
304	0.0313	0.0329
305	0.0979	0.0612
306	0.0678	0.0388
307	0.0858	0.0267
308	0.116	2.0
309	0.0822	0.0247
310	0.119	0.0308
311	0.0158	0.0121
312	0.368	0.00961
313	0.0363	0.0159
314	0.0459	0.00676
315	0.0572	0.104
316	0.744	2.0
317	0.0443	0.0152
318	0.436	2.0
319	0.0776	0.055
320	0.13	0.549
321	0.523	2.0
322	0.0282	0.0151
323	0.613	2.0
324	0.137	0.109
325	0.0698	0.0247
326	0.0339	0.0109
327	0.159	0.0776
328	0.040	0.0834
329	0.050	0.0551
330	0.0806	0.159
331	0.0243	0.00704
332	0.0169	0.0338


333	0.297	0.00681
334	0.0599	0.0565
335	0.32	0.59
336	0.0378	0.0635
337	0.0275	0.0231
338	0.169	2.0
339	0.0138	0.0019
340	0.0663	0.429
341	0.162	0.717
343	0.0541	0.0425
344	0.0973	0.072
345	0.14	0.0808
346	0.0103	0.00601
347	0.0113	0.00393
348	0.102	0.0426
350	0.0273	0.00396
351	0.0288	0.00356
352	0.0267	0.00651
353	0.0238	0.00755
354	0.0288	0.00707
355	0.00441	0.00132
356	0.0060	0.00762
357	0.0236	0.0106
358	0.0202	0.00919
359	0.0208	0.0217
360	0.0957	0.0509
361	0.0357	0.0561
362	0.00969	0.0019
363	0.0505	0.0114
364	0.365	0.11
365	0.0845	0.0463
366	0.00993	0.00295
367	0.0183	0.00211
368	0.17	0.991
369	0.167	0.0933
370	0.169	0.115
371	0.0286	0.0302
372	0.0227	0.0275
373	0.0152	0.00703
375	0.00888	0.00111
376	0.0675	0.0161
377	0.0149	0.00159
378	0.241	2.0
379	1.47	2.0
381	0.0233	0.0554

382	0.0229	0.0248
383	0.0158	2.0
384	0.928	0.436
385	0.0824	0.0171
387	0.724	0.535
388	0.034	0.00126
389	0.0216	0.000575
392	0.699	2.0
393	0.0495	0.0185
394	0.0205	0.0145
395	0.193	0.0243
396	0.028	0.0175
397	0.412	0.888
398	0.0125	0.000722
399	0.00331	0.000381
400	0.136	0.222
401	0.29	2.0
402	0.557	2.0
403	0.011	0.0029
404	0.0136	0.00375
405	0.00637	0.00534
406	0.29	2
407	0.0131	0.0726
408	0.00376	0.00101
409	0.0119	0.00103
410	0.0121	0.00393
411	0.121	2.0
413	0.0053	0.00292
414	0.34	0.0535
415	0.0128	0.0142
416	0.082	0.461
417	0.0506	0.83
418	0.0056	0.00352
419	0.00149	0.000489

While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

CLAIMS

1. A compound of Formula I:



5

wherein:

R is (a) an 8-, 9-, or 10-membered bicyclic heteroaryl comprising one heteroatom selected from N, S, and O, and one, two, or three additional N atoms, wherein said bicyclic heteroaryl is unsubstituted or is substituted with one or more substituents selected from the group consisting of deuterium, amino, alkylamino, dialkylamino, alkyl, halo, cyano, haloalkyl, hydroxy, hydroxyalkyl, and alkoxy, and wherein one or more N atoms of said bicyclic heteroaryl is optionally an N-oxide; or

10

(b) a five- or six-membered nitrogen-linked heterocycloalkyl ring fused to a phenyl or monocyclic five- or six-membered heteroaryl, wherein said phenyl or heteroaryl is unsubstituted or is substituted with one or more substituents selected from the group consisting of deuterium, amino, alkylamino, dialkylamino, alkyl, halo, cyano, haloalkyl, hydroxy, hydroxyalkyl, and alkoxy; and

15

R¹ is H, -(C₁₋₄alkylene)₀₋₁C(O)R^a, -(C₁₋₄alkylene)₀₋₁CO₂R^a, -(C₁₋₄alkylene)₀₋₁S(O)R^a, -(C₁₋₄alkylene)₀₋₁SO₂R^a, -C(O)NH(R^a), -C(O)N(R^a)₂, or -C(O)C(O)NH(R^a);

20

wherein each R^a is independently

(1) alkyl, unsubstituted or substituted with one or more R^m substituents,

25

wherein each R^m is independently selected from the group consisting of deuterium, hydroxy, -NR^bR^c, alkoxy, cyano, halo, -C(O)alkyl, -CO₂alkyl, -CONR^bR^c, -S(O)alkyl, -SO₂alkyl, -SO₂NR^bR^c, aryl, heteroaryl, cycloalkyl, heterocycloalkyl, phenoxy, and -O-alkyl-OH; wherein R^b is H or alkyl;

R^c is H, alkyl, alkoxyalkyl, haloalkyl, -C(O)alkyl, -CO₂alkyl,
-SO₂alkyl, -C(O)NH₂, or C(O)H; and

each aryl, heteroaryl, cycloalkyl, and heterocycloalkyl group within

R^m is unsubstituted or substituted with one or more substituents
independently selected from the group consisting of alkyl,
haloalkyl, hydroxy, -NR^bR^c, alkoxy, haloalkoxy, cyano, halo, oxo,
-C(O)alkyl, -CO₂alkyl, -C(O)-heterocycloalkyl, -CONR^bR^c,
-S(O)alkyl, -SO₂alkyl, -SO₂-haloalkyl, -SO₂NR^bR^c, aryl, heteroaryl,
cycloalkyl, and heterocycloalkyl; or two substituents taken
together form a fused heteroaryl, cycloalkyl, or heterocycloalkyl
ring;

wherein each alkyl or alkoxy is unsubstituted or substituted with
phenyl, -NR^bR^c, heterocycloalkyl, heteroaryl, or -C(O)alkyl;
and

each aryl, heteroaryl, cycloalkyl, and heterocycloalkyl is
unsubstituted or substituted with alkyl, halo, or -C(O)alkyl,
(2) phenyl, cycloalkyl, heteroaryl, or heterocycloalkyl, each unsubstituted or
substituted with one or more R^g substituents;

wherein each R^g is independently selected from the group consisting of
alkyl, haloalkyl, hydroxy, -NR^bR^c, alkoxy, haloalkoxy, cyano, halo,
oxo, -C(O)alkyl, -CO₂alkyl, -C(O)-heterocycloalkyl, -CONR^bR^c,
-S(O)alkyl, -SO₂alkyl, -SO₂-haloalkyl, -SO₂NR^bR^c, aryl, heteroaryl,
cycloalkyl, and heterocycloalkyl; or two R^g substituents taken together
form a fused phenyl, heteroaryl, cycloalkyl, or heterocycloalkyl ring;

wherein each alkyl or alkoxy is unsubstituted or substituted with
-NR^bR^c, heterocycloalkyl, heteroaryl, or -C(O)alkyl; and
each aryl, heteroaryl, cycloalkyl, and heterocycloalkyl is unsubstituted
or substituted with alkyl, halo, -CO₂alkyl, or -C(O)alkyl; or

(3) -NR^xR^y,

where R^x is H or alkyl; and

R^y is H, alkyl, alkoxyalkyl, haloalkyl, -C(O)alkyl, -CO₂alkyl, or

-SO₂alkyl;

R² and R³ are each independently H or deuterium; and


n is 1 or 2;

or a stereoisomer thereof, or a pharmaceutically acceptable salt of such a compound

5 or stereoisomer.

2. The compound of claim 1, wherein R is an 8- or 9-membered heteroaryl, unsubstituted or substituted as described for claim 1.

10 3. The compound of claim 1, wherein R is:



each unsubstituted or substituted as described for claim 1.


15 4. The compound of claim 1, wherein, R is a five- or six-membered nitrogen-linked heterocycloalkyl ring fused to an unsubstituted or substituted phenyl or monocyclic heteroaryl, as defined in claim 1.



20 6. The compound of claim 1, wherein R¹ is H.









7. The compound of claim 1, wherein R¹ is -C(O)R^a, -CO₂R^a, -S(O)R^a, or -SO₂R^a.
8. The compound of claim 1, wherein R^a is alkyl, unsubstituted or substituted as described for claim 1.
9. The compound of claim 1, wherein R^a is methyl, ethyl, propyl, isopropyl, tert-butyl, isobutyl, isopentyl, phenyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, pyrrolyl, furanyl, thiophenyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, triazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, isoindolinyl, azetidinyl, oxetanyl, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, tetrahydrofuranyl, tetrahydropyranyl, or tetrahydrothiophenyl, each unsubstituted or substituted.
10. The compound of claim 1, wherein R^a is phenyl, cycloalkyl, heteroaryl, or heterocycloalkyl, each unsubstituted or substituted with one or more substituents selected from the group consisting of fluoro, oxo, methyl, -CONH₂, acetyl, -SO₂methyl, -C(O)-isopropyl, pyridazinyl, triazolyl, dimethylaminomethyl, cyano, methyl-triazolyl-methoxy, trifluoromethoxy, pyrrolidinylmethyl, acetylamino, tetrazolylmethyl, methyl-tetrazolyl-methyl, methyl-imidazolyl-methyl, -NHISO₂methyl, 1,1-dioxothiomorpholinyl, 4-methyl-piperazinylmethyl, -NHCONH₂, -SO₂CF₃, morpholinylmethyl, imidazolyl, -SO₂NH₂, methylpiperidinyl, methyl-piperazinyl, -C(O)(4-methyl-piperazinyl), morpholinyl, trifluoromethyl, cyclopropyl, ethyl, isoxazolyl, tetrazolyl, isopropyl, phenyl, fluoro-phenyl, tert-butyl, benzyl, N-methylpyrrolidinyl, N-acetyl-pyrrolidinyl, isobutyl, propyl, methylpyrazolyl, trifluoroethyl, pyrimidinyl, oxo, acetyl, cyano, -CO₂-tert-butyl, and amino.
11. The compound of claim 1, wherein R^a is alkyl, unsubstituted or substituted with one or more substituents selected from the group consisting of fluoro, tert-butoxy, -C(O)NMe₂, -NHCHO, methoxy, phenoxy, cyano, acetyl, hydroxy, -OCH₂C(CH₃)=OH, -NH(acetyl), and -N(Me)(acetyl).









12. The compound of claim 1, wherein R^1 is $-\text{SO}_2R^a$, where R^a is methyl, ethyl, phenyl, benzyl, or 2,2-dimethylpropyl.
13. The compound of claim 1, wherein R^1 is $-\text{C}(\text{O})\text{NHR}^a$, wherein R^a is methyl, ethyl, propyl, isopropyl, tertobutyl, cyclohexyl, $-\text{CH}_2$ -cyclohexyl, oxetanyl, or methyloxetanyl, or R^a is a phenyl or benzyl group, each optionally substituted with one or more substituents selected from the group consisting of cyano, methyl, fluoro, methoxy, and chloro.
14. The compound of claim 1, wherein both R^2 and R^3 are H.
15. The compound of claim 1, which is a compound of Formula I-a:



- wherein R , R^1 , R^2 , and R^3 are as defined for Formula I;
- 15 or a stereoisomer thereof, or a pharmaceutically acceptable salt of such a compound or stereoisomer.
16. A compound selected from the group consisting of:




	tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;
	tert-butyl 2-[(thieno[2,3-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;
	tert-butyl 2-[(imidazo[1,2-a]pyrimidine-6-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;

	tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;
	tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;
	N-[[8-(3,3-dimethylbutanoyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[8-(2-phenylacetyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
	tert-butyl 2-[(1H-pyrrolo[3,2-c]pyridine-2-carboxylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;
	N-[[8-(cyclohexanecarbonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[8-(tetrahydropyran-4-carbonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[(8-benzoyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;




	N-[[8-(2-methylpropanoyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[8-(benzenesulfonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide
	N-[(8-ethylsulfonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	tert-butyl 2-[(1H-pyrazolo[3,4-b]pyridine-5-carbonylamino)methyl]-8-azaspiro[2.5]octane-8-carboxylate;
	N-[(8-acetyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[(8-methylsulfonyl)-8-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-(8-azaspiro[2.5]octan-2-ylmethyl)-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-(8-azaspiro[2.5]octan-2-ylmethyl)-1H-pyrazolo[3,4-b]pyridine-5-carboxamide;

	<p>N-[(8-propanoyl-8-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>tert-butyl 2-((imidazo[1,2-a]pyridine-6-carboxamido)methyl)-7-azaspiro[3.5]nonane-7-carboxylate;</p>
	<p>tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>

	<p>N-[[6-(2-imidazol-1-ylacetyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4,4,4-trifluorobutanoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-methylsulfonylphenyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position and a methyl group at the 6-position. The methyl group is attached to a 6-azaspiro[2.5]octane ring, which is further substituted with a 3,3-dimethylbutanoyl group.</p>	<p>N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position and a methyl group at the 6-position. The methyl group is attached to a 6-azaspiro[2.5]octane ring, which is further substituted with a 2-(4-methylpiperazin-1-yl)acetyl group.</p>	<p>N-[[6-[2-(4-methylpiperazin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position and a methyl group at the 6-position. The methyl group is attached to a 6-azaspiro[2.5]octane ring, which is further substituted with a 2-(3-cyanophenyl)acetyl group.</p>	<p>N-[[6-[2-(3-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2-cyclopropylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-pyrrolidin-1-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-methylisoxazol-5-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position and a methyl group at the 3-position. The methyl group is attached to a 6-azaspiro[2.5]octane ring. The nitrogen of the 6-azaspiro[2.5]octane ring is substituted with a 2-(1-piperidyl)acetyl group.</p>	<p>N-[[6-[2-(1-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the piperidine ring is replaced by a tetrahydropyran ring.</p>	<p>N-[[6-(2-tetrahydropyran-4-ylacetyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the piperidine ring is replaced by a tetrahydrofuran ring.</p>	<p>N-[[6-(2-tetrahydrofuran-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2-tert-butoxyacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-[1-(2,2,2-trifluoroethyl)-4-piperidyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 1);</p>

	tert-butyl 2-[(imidazo[1,2-a]pyridine-6-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 2);
	tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 1);
	tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 2);
	N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(2-phenylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 1);




	<p>tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate (isomer 2);</p>
	<p>N-[[6-(2-pyrazin-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-morpholinoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-tetrahydropyran-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2-pyrrolidin-1-ylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(tert-butylamino)-2-oxoacetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(dimethylamino)-4-oxo-butanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-methyl-5-oxo-pyrrolidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is connected to a 6-azaspiro[2.5]octane system. The 6-position of the azaspiro system is substituted with a 2-(2-oxopyrrolidin-1-yl)acetyl group.</p>	<p>N-[[6-[2-(2-oxopyrrolidin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is similar to the first one, but the 2-(2-oxopyrrolidin-1-yl)acetyl group is replaced by a 2,2-difluorocyclopropanecarbonyl group.</p>	<p>N-[[6-(2,2-difluorocyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is similar to the first one, but the 2-(2-oxopyrrolidin-1-yl)acetyl group is replaced by a tetrahydrofuran-3-carbonyl group.</p>	<p>N-[[6-(tetrahydrofuran-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(tetrahydropyran-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(tetrahydrofuran-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-formamidoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[3-(3,5-dimethylpyrazol-1-yl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-carbamoylpyrrolidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-morpholinopropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>




 <p>The structure shows a furo[2,3-c]pyridine ring system with a methyl group at the 2-position and a carboxamide group at the 3-position. The nitrogen of the carboxamide group is attached to a 6-azaspiro[2.5]octane ring system. The nitrogen of the azaspiro ring is further substituted with a 1,1-dioxothiolane-3-carbonyl group.</p>	<p>N-[[6-(1,1-dioxothiolane-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the 1,1-dioxothiolane-3-carbonyl group is replaced by a 2-(3,5-dimethylpyrazol-1-yl)acetyl group.</p>	<p>N-[[6-[2-(3,5-dimethylpyrazol-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide,</p>
 <p>The structure is similar to the first one, but the 1,1-dioxothiolane-3-carbonyl group is replaced by a 1-acetylpyrrolidine-2-carbonyl group.</p>	<p>N-[[6-(1-acetylpyrrolidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide,</p>

	<p>N-[[6-(2-methylsulfonylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[3-(1,2,4-triazol-4-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-((dimethylamino)methyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position and a methyl group at the 6-position. The methyl group is attached to the 6-position of a 6-azaspiro[2.5]octane ring. The 6-position of the azaspiro ring is further substituted with a 4-acetamidobenzoyl group.</p>	<p>N-[[6-(4-acetamidobenzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is similar to the first one, but the 4-acetamidobenzoyl group is replaced by a 1-methylsulfonylpiperidine-3-carbonyl group.</p>	<p>N-[[6-(1-methylsulfonylpiperidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is similar to the second one, but the 1-methylsulfonylpiperidine-3-carbonyl group is replaced by a 1-methylsulfonylpiperidine-4-carbonyl group.</p>	<p>N-[[6-(1-methylsulfonylpiperidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2-morpholinopropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-methylsulfonylpyrrolidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-oxoisindolin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at position 2 and a methyl group at position 3. The methyl group is attached to the 6-position of a 6-azaspiro[2.5]octane system. The 6-position of the spiro system is also attached to a piperidine ring, which has a carbonyl group at its 4-position. This carbonyl group is further attached to another piperidine ring, which has a carbonyl group at its 1-position. This final carbonyl group is attached to an amino group (H₂N).</p>	<p>N-[[6-(1-carbamoylpiperidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is identical to the first one, except that the carbonyl group on the piperidine ring is at the 3-position instead of the 4-position.</p>	<p>N-[[6-(1-carbamoylpiperidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is identical to the first one, except that the piperidine ring is attached to an acetyl group at its 6-position. The acetyl group is further attached to a 1,1-dioxothiolan-3-yl group.</p>	<p>N-[[6-[2-(1,1-dioxothiolan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the spiro ring is substituted with a 2-(2,4-dioxypyrimidin-1-yl)acetyl group.</p>	<p>N-[[6-[2-(2,4-dioxypyrimidin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the spiro ring is substituted with a 1-acetylpiperidine-4-carbonyl group.</p>	<p>N-[[6-(1-acetylpiperidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the spiro ring is substituted with a 4-(tetrazol-1-ylmethyl)benzoyl group.</p>	<p>N-[[6-[4-(tetrazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>




	<p>N-[[6-[4-(methanesulfonamido)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(1,1-dioxo-1,4-thiazinan-4-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a cyclopropane ring, which is further connected to a 6-azaspiro[2.5]octane system. The nitrogen of the spiro system is attached to a benzoyl group, which is substituted at the para position with a (4-methylpiperazin-1-yl)methyl group.</p>	<p>N-[[6-[4-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is similar to the first one, but the benzoyl group is substituted at the ortho position with a (4-methylpiperazin-1-yl)methyl group.</p>	<p>N-[[6-[2-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure is similar to the first one, but the benzoyl group is substituted at the meta position with a (4-methylpiperazin-1-yl)methyl group.</p>	<p>N-[[6-[3-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>




<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a bicyclic azaspiro[2.5]octane system. The 6-position of the azaspiro system is substituted with a propanoyl chain, which is further substituted with a 4-methylpiperazin-1-yl group.</p>	<p>N-[[6-[3-(4-methylpiperazin-1-yl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a bicyclic azaspiro[2.5]octane system. The 6-position of the azaspiro system is substituted with a pyrrolidine-3-carbonyl chain, which is further substituted with an isopropyl group.</p>	<p>N-[[6-(1-isopropyl-5-oxopyrrolidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a bicyclic azaspiro[2.5]octane system. The 6-position of the azaspiro system is substituted with an acetyl chain, which is further substituted with a 3-oxopiperazin-1-yl group.</p>	<p>N-[[6-[2-(3-oxopiperazin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(3-methylsulfonylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-methylsulfonylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-ureidobenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[2-(4-acetylpiperazin-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-[(5-methyltetrazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>




 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring, which is further substituted with a pyridine-2-carbonyl group.</p>	N-[[6-(pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
 <p>The structure is similar to the first one, but the pyridine ring is replaced by an oxazole ring.</p>	N-[[6-(oxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
 <p>The structure is similar to the first one, but the pyridine ring is replaced by a 1H-1,2,4-triazole ring.</p>	N-[[6-(1H-1,2,4-triazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;

	<p>N-[[6-(1H-imidazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrimidine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring system. The 6-position of the azaspiro ring is substituted with a 1H-pyrazole-3-carbonyl group.</p>	<p>N-[[6-(1H-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the pyrazole ring is replaced by a 1-methylimidazole ring, with the carbonyl group at the 4-position.</p>	<p>N-[[6-(1-methylimidazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the pyrazole ring has a methyl group at the 3-position and the carbonyl group is at the 4-position.</p>	<p>N-[[6-(3-methyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-methyl-1H-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(pyridazine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrimidine-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrazine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>




 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of the amide is attached to a 6-azaspiro[2.5]octane ring system. The 6-position of the azaspiro ring is substituted with a pyridazine-4-carbonyl group.</p>	<p>N-[[6-(pyridazine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the pyridazine ring is replaced by a 1-methylpyrazole ring. The amide group is shown as a secondary amide (HN).</p>	<p>N-[[6-(1-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the second one, but the 1-methylpyrazole ring is replaced by a 1,5-dimethylpyrazole ring. The amide group is shown as a secondary amide (HN).</p>	<p>N-[[6-(1,5-dimethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2,5-dimethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-methylisoxazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-methyl-1,2,5-oxadiazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(2-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-methylpyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(5-methylpyrazine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-ethyl-5-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(3,5-dimethylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-ethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(6-cyanopyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>




	<p>N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-morpholinopyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(cyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>






	<p>N-[[6-[1-methyl-5-(trifluoromethyl)pyrazole-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-isoxazol-5-yl-1-methyl-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-acetamidopyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of this amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the azaspiro ring is substituted with a carbonyl group, which is further linked to a 1H-pyrazole ring at its 4-position. The 3-position of the pyrazole ring is substituted with a tetrazol-1-yl group.</p>	<p>N-[[6-[3-(tetrazol-1-yl)-1H-pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of this amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the azaspiro ring is substituted with a carbonyl group, which is further linked to a pyridine ring at its 2-position. The 6-position of the pyridine ring is substituted with an acetamido group (-NH-C(=O)-CH3).</p>	<p>N-[[6-(6-acetamidopyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
<p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of this amide is attached to a 6-azaspiro[2.5]octane ring. The 6-position of the azaspiro ring is substituted with a carbonyl group, which is further linked to an isoxazole ring at its 4-position. The 3-position of the isoxazole ring is substituted with an ethyl group, and the 5-position is substituted with a methyl group.</p>	<p>N-[[6-(3-ethyl-5-methylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>





	<p>N-[[6-(2-ethyl-5-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-ethyl-3-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-methylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[2-methyl-3-(3-methylpyrazol-1-yl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1-methylsulfonyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a furo[2,3-c]pyridine ring system with a carboxamide group at the 2-position. The nitrogen of this amide is attached to a 6-membered azaspiro[2.5]octane ring. The 6-position of the azaspiro ring is substituted with a 4-methylmorpholine-2-carbonyl group.</p>	<p>N-[[6-(4-methylmorpholine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the 4-methylmorpholine group is replaced by a 1-(2-methylpropanoyl)azetidine-3-carbonyl group attached to the 6-position of the azaspiro ring.</p>	<p>N-[[6-[1-(2-methylpropanoyl)azetidine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the second one, but the 2-methylpropanoyl group is replaced by an acetyl group attached to the 1-position of the azetidine ring.</p>	<p>N-[[6-(1-acetylazetidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>







	<p>N-[[6-(1-pyridazin-3-yl)pyrrolidine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1-pyrimidin-2-yl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1-isopropyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1-methyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide</p>
	<p>N-[[6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl)methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>

	<p>isopropyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>N-[[6-[2-(2-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-methoxypropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-[4-(trifluoromethylsulfonyl)phenyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a furo[2,3-c]pyridine ring system. At the 2-position, there is a carboxamide group (-NH-C(=O)-) attached to a propano chain. The 3-position of the propano chain has an acetyl(methyl)amino group (-NH-C(=O)-CH₂-CH₂-N(CH₃)-C(=O)-CH₃). At the 6-position, the propano chain is attached to a methyl group, which is further connected to a piperidine ring. The piperidine ring is spiro-fused to a cyclopropane ring.</p>	<p>N-[[6-[3-acetyl(methyl)amino]propano]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure features a furo[2,3-c]pyridine ring with a carboxamide group at the 2-position. The 6-position is part of a 6-azaspiro[2.5]octane system. The nitrogen of this spiro system is substituted with a 2-(2-pyridyl)acetyl group (-NH-C(=O)-CH₂-CH₂-pyridine-2-yl). A methyl group at the 6-position of the spiro system is attached to the furo[2,3-c]pyridine ring.</p>	<p>N-[[6-[2-(2-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure shows a furo[2,3-c]pyridine ring with a carboxamide group at the 2-position. The 6-position is part of a 6-azaspiro[2.5]octane system. The nitrogen of this spiro system is substituted with a (4-cyanophenyl)carbamoyl group (-NH-C(=O)-NH-phenyl-4-CN). A methyl group at the 6-position of the spiro system is attached to the furo[2,3-c]pyridine ring.</p>	<p>N-[[6-[(4-cyanophenyl)carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure features a furo[2,3-c]pyridine ring with a carboxamide group at the 2-position. The 6-position is part of a 6-azaspiro[2.5]octane system. The nitrogen of this spiro system is substituted with an (m-tolylmethyl)carbamoyl group (-NH-C(=O)-NH-CH₂-CH₂-phenyl-3-CH₃). A methyl group at the 6-position of the spiro system is attached to the furo[2,3-c]pyridine ring.</p>	<p>N-[[6-(m-tolylmethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(p-tolylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(benzylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(2-fluorophenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(p-tolylmethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(4-fluorophenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(2-methoxyphenyl)methylcarbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[(3-methoxyphenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(3-fluorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(4-methoxyphenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(3,4-dichlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-adamantyl)carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(4-chlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[(2-chlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(ethylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[(2,4-dichlorophenyl)methyl]carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(2-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>








	<p>N-[[6-[2-(3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(1,2,4-triazol-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(2-methylimidazol-1-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(tert-butylcarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(isopropylcarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(cyclohexylmethylcarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(propylcarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(cyclohexylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(phenylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1-isopropyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(4-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-cyanoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>

	<p>ethyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>tert-butyl 2-[(1H-pyrrolo[3,2-c]pyridine-2-carboxylamino)methyl]-7-azaspiro[3.5]nonane-7-carboxylate;</p>
	<p>tert-butyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-7-azaspiro[3.5]nonane-7-carboxylate;</p>
	<p>N-[[6-[2-(3-methyloxetan-3-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-cyclohexylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-cyclohexylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-morpholinoacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(3-phenylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>tert-butyl 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-7-azaspiro[3.5]nonane-7-carboxylate;</p>
	<p>N-[[6-[2-(3,5-difluorophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-(trifluoromethyl)phenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(tert-butylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>[2,2,2-trideuterio-1,1-bis(trideuteriomethyl)ethyl] 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>(2-methoxy-1,1-dimethyl-2-oxo-ethyl) 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>





	N-[[6-[2-(3-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-[2-(4-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(phenylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[3-(3-pyridyl)propanoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(benzylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(benzylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(tert-butylcarbamoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;





	N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide (isomer 1);
	N-[[6-(3,3-dimethylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide (isomer 2);
	(2-hydroxy-1,1-dimethyl-ethyl) 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-[2-(4-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	(2-methoxy-1,1-dimethyl-ethyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-(2-tetrahydropyran-4-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(1-methylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;

	<p>N-[[6-(1-isopropyl-3,5-dimethyl-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[1-(4-fluorophenyl)-3,5-dimethyl-pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>(1-methylcyclobutyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>N-[[6-(3-morpholinopropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2,2-difluoro-2-phenylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(3-methylpyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(6-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(6-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(1H-benzimidazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrrolo[1,2-c]pyrimidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1H-benzotriazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1H-pyrrolo[2,3-b]pyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1H-indazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

 <p>The structure shows a 1H-benzimidazole ring system with a carbonyl group at the 5-position. This carbonyl is attached to the nitrogen atom of a 6-azaspiro[2.5]octane system. The 2-position of the azaspiro system is substituted with a methyl group. This methyl group is further substituted with a furo[2,3-c]pyridine-2-carboxamide moiety.</p>	<p>N-[[6-(1H-benzimidazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the first one, but the benzimidazole ring is replaced by a benzene ring with a cyano group (-C≡N) at the 4-position. The carbonyl group at the 1-position of the benzene ring is attached to the nitrogen atom of the 6-azaspiro[2.5]octane system.</p>	<p>N-[[6-(4-cyanobenzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure is similar to the second one, but the cyano group (-C≡N) is located at the 3-position of the benzene ring.</p>	<p>N-[[6-(3-cyanobenzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
 <p>The structure features a 5-methyl-1,2,4-oxadiazole ring with a methoxy group (-OCH2-) at the 3-position. This methoxy group is attached to a benzoyl group at the 4-position. The carbonyl group of this benzoyl group is attached to the nitrogen atom of the 6-azaspiro[2.5]octane system. The 2-position of the azaspiro system is substituted with a methyl group, which is further substituted with a furo[2,3-c]pyridine-2-carboxamide moiety.</p>	<p>N-[[6-[4-[(5-methyl-1,2,4-oxadiazol-3-yl)methoxy]benzoyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	N-[[6-[4-(morpholinomethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-[4-[(2-methylimidazol-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-[4-(trifluoromethoxy)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-[4-(pyrrolidin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;






	<p>N-[[6-[3-(1,2,4-triazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1,2,4-triazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(1,2,4-triazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[3-(imidazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(4-imidazol-1-ylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(1H-1,2,4-triazol-5-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(6-imidazol-1-ylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(trifluoromethyl)pyridine-2-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[6-(trifluoromethyl)pyridine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-oxo-4H-1,4-benzoxazine-6-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-sulfamoylbenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(imidazol-1-ylmethyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(quinoxaline-6-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>









	<p>N-[[6-(1,6-naphthyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-methyl-1H-benzimidazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-acetamidobenzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(3-acetamidopyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2,1,3-benzoxadiazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1,8-naphthyridine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(isoquinoline-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(quinoxaline-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1,5-naphthyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1,8-naphthyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(isoxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>tert-butyl 2-[(4,6-dihydro-1H-pyrrolo[3,4-c]pyrazole-5-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>N-[[6-[2-(1,4-dimethyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(4-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1,3-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(2-methyltetrazol-5-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(4,6-dimethylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-oxopentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-hydroxy-4-methylpentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-hydroxy-3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-(2-pyridyl)acetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-pyrimidin-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>

	N-[[6-(2-pyrazin-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(3-thiazol-2-ylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(2-phenoxyacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(3-tetrahydropyran-4-ylpropanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(4-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(3,5-dimethyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(1-tert-butyl-3,5-dimethyl-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;

	<p>N-[[6-(1-benzyl-3,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-tert-butyl-2-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-ethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-tert-butyl-4-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2,4-dimethylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide (isomer 1);</p>

	<p>N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide (isomer 2);</p>
	<p>N-[[6-(1H-indazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[1-(1-isopropylpyrrolidin-3-yl)pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[1-(1-acetylpyrrolidin-3-yl)pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-[5-(1,3-dimethylpyrazol-4-yl)isoxazole-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>tert-butyl 3-[2-((furo[2,3-c]pyridine-2-carbonylamino)methyl)-6-azaspiro[2.5]octane-6-carbonyl]-4,6-dihydro-1H-pyrrolo[3,4-c]pyrazole-5-carboxylate;</p>
	<p>N-[[6-[1-(1-methylpyrrolidin-3-yl)pyrazole-4-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-tert-butyl-5-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-methylimidazo[1,2-a]pyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>





	<p>N-[[6-(6-methyl-1H-benzimidazole-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-methylimidazo[1,2-a]pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(6-fluoro-1H-indazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrazolo[1,5-a]pyrimidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1-ethyl-3,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(6-methylimidazo[1,2-a]pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrazolo[1,5-a]pyrimidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(imidazo[1,2-a]pyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-isopropylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-isobutyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(imidazo[1,2-a]pyrimidine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(pyrazolo[1,5-a]pyrimidine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>








	<p>N-[[6-(3-tert-butyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-isopropyl-1H-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-isopropyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2-isopropylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	<p>N-[[6-(5-ethyl-2-methylpyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-cyclopropylisoxazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-cyclopropylisoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3,5-dimethyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>









	<p>N-[[6-(1,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-methyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-cyclopropyl-1H-pyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(1,4,5,6-tetrahydrocyclopenta[c]pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>








	N-[[6-(2,5-dimethyloxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-(1,3-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-(oxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-(isoxazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;









	<p>N-[[6-(1H-imidazole-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[4-(4-methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3H-imidazo[4,5-b]pyridine-6-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(5-methylpyridine-2-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>

	N-[[6-[4-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-[3-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-[4-[(dimethylamino)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-[4-[(4-methylpiperazin-1-yl)methyl]benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(1,5-dimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	(3-methyloxetan-3-yl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;








	<p>(2-hydroxy-1,1-dimethyl-ethyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>N-[[6-(4-methylpyridine-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(4-isoxazol-5-yl-1-methyl-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(1,4-dimethyl-4-piperidyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide; N-[[6-[2-(2-cyanophenyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>

	(2-acetamido-1,1-dimethyl-ethyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-(2-thiazol-2-ylacetyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(4-hydroxy-4-methylpentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[(3-methyloxetan-3-yl)carbamoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[4-(4-methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[4-(4-methylpiperazin-1-yl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-








	2-carboxamide (isomer 1); N-[[6-(1,3,5-trimethylpyrazole-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide (isomer 2);
	tert-butyl 2-[[[(1,1,3,3-tetradeuteriopyrrolo[3,4-c]pyridine-2-carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(4-isoxazol-5-yl-1-methyl-pyrazole-3-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-[4-(4-methylpiperazine-1-carbonyl)benzoyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;








	N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	(3-methyltetrahydrofuran-3-yl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	(2-hydroxy-2-methyl-propyl) 2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(2,2-dimethylpropylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;
	N-[[6-(2,2-dimethylpropylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	N-[[6-(2,2-dimethylpropylsulfonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(4,4,4-trifluorobutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;

	<p>tert-butyl 2-[(pyrazolo[1,5-b]pyridazine-5-carbonylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;</p>
	<p>N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide (single isomer);</p>
	<p>N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]furo[2,3-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-(2-hydroxy-2-methylpropoxy)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;</p>
	<p>N-[[6-[2-methyl-4-(trifluoromethyl)thiazole-5-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;</p>

	N-[[6-(2,2-dimethylcyclopropanecarbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	tert-butyl 2-[(imidazo[1,2-a]pyridin-6-yl)methylcarbamoylamino]methyl-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-[2-(2-hydroxy-2-methylpropoxy)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[2-(6-amino-3-pyridyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(morpholine-4-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[5-(trifluoromethyl)-1H-pyrazole-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(4-hydroxy-4-methylpentanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;

	N-[[6-[4-(trifluoromethyl)pyridine-3-carbonyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[2-(4-methyltetrahydropyran-4-yl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-[2-(3-hydroxy-3-methylcyclobutyl)acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1H-pyrrolo[3,2-c]pyridine-2-carboxamide;
	(2-hydroxy-2-methyl-propyl) 2-[[1H-pyrrolo[3,2-c]pyridine-2-carboxylamino]methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	tert-butyl 2-[[imidazo[1,2-b]pyridazine-6-carboxylamino]methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-[2-[4-methyl-1-(2,2,2-trifluoroethyl)-4-piperidyl]acetyl]-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(2,4-dimethylthiazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;

	N-[[6-[4-(1-methyl-4-piperidyl)benzoyl]-6-azaspiro[2.5]octan-2-yl)methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	tert-butyl 2-[[[(6-amino-1,3-dihydropyrrolo[3,4-c]pyridine-2-carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[[6-(2-pyrimidin-5-ylacetyl)-6-azaspiro[2.5]octan-2-yl)methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	1,1,3,3-tetradeuterio-N-[[6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl)methyl]pyrrolo[3,4-c]pyridine-2-carboxamide;
	(3-methyloxetan-3-yl) 2-[[[(1,1,3,3-tetradeuteriopyrrolo[3,4-c]pyridine-2-carbonyl)amino]methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	1,1,3,3-tetradeuterio-N-[[6-(2,4-dimethyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl)methyl]pyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(benzoyl)-6-azaspiro[2.5]octan-2-yl)methyl]furo[2,3-c]pyridine-2-carboxamide;

	N-[[<i>(2S)</i> -6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[<i>(2R)</i> -6-(3-methylbutanoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	N-[[6-(4-methyloxazole-5-carbonyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	isopropyl 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	(3-methyloxetan-3-yl) 2-[(furo[2,3-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;
	N-[(6-benzoyl)-6-azaspiro[2.5]octan-2-yl]methyl]-1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxamide;
	(3-methyloxetan-3-yl) (<i>2S</i>)-2-[(1,3-dihydropyrrolo[3,4-c]pyridine-2-carboxylamino)methyl]-6-azaspiro[2.5]octane-6-carboxylate;

and stereoisomers thereof, and pharmaceutically acceptable salts of such compounds and stereoisomers.

17. A pharmaceutical composition comprising: (a) an effective amount of at least

one compound of claim 1; and (b) a pharmaceutically acceptable carrier.

18. The pharmaceutical composition of claim 17, further comprising therapeutically effective amounts of one or more additional adjunctive active agents.

5

19. The pharmaceutical composition of claim 18, wherein said one or more additional adjunctive active agents are selected from the group consisting of cytotoxic agent, cisplatin, doxorubicin, taxotere, taxol, etoposide, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, the epothilones, tamoxifen, 5-fluorouracil, methotrexate, temozolomide, cyclophosphamide, SCH 66336, tipifarnib (Zarnestra[®]), R115777, L778,123, BMS 214662, Iressa[®], Tarceva[®], C225, GLEEVEC[®], intron[®], Peg-Intron[®], aromatase combinations, ara-C, adriamycin, cytoxan, gemcitabine, Uracil mustard, Chlormethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, oxaliplatin, leucovorin, oxaliplatin (ELOXATIN[®]), Pentostatine, Vinblastine, Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, MithramycinTM, Deoxycoformycin, Mitomycin-C, L-Asparaginase, Teniposide 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrol acetate, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, goserelin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, Anastrozole, Letrazole, Capecitabine, Reloxafine, Droloxafine, Hexamethylmelamine, Avastin, herceptin, Bexxar, Velcade, Zevalin, Trisenox, Xeloda, Vinorelbine, Porfimer, Erbitux, Liposomal, Thiotepa, Altretamine, Melphalan, Trastuzumab, Lerozole, Fulvestrant, Exemestane, Ifosfomide, Rituximab, C225, Campath, leucovorin, and dexamethasone,

10
15
20
25
30

bicalutamide, carboplatin, chlorambucil, cisplatin, letrozole, megestrol, valrubicin, vinblastine and NIASPAN[®].

20. The pharmaceutical composition of claim 17 further comprising a rescuing agent.

21. The pharmaceutical composition of claim 20, wherein the rescuing agent is selected from the group consisting of nicotinamide, nicotinic acid, and nicotinamide mononucleotide (NMN).

22. A method of treating a subject suffering from or diagnosed with a disease or medical condition mediated by NAMPT activity, comprising administering to the subject in need of such treatment an effective amount of at least one compound of claim 1.

23. The method of claim 22, wherein the disease or medical condition is a solid or liquid tumor, non-small cell lung cancer, leukemia, lymphoma, ovarian cancer, glioma, breast cancer, uterine cancer, colon cancer, cervical cancer, lung cancer, prostate cancer, skin cancer, rhino-gastric tumors, colorectal cancer, CNS cancer, bladder cancer, pancreatic cancer, Hodgkin's disease, rheumatoid arthritis, diabetes, atherosclerosis, sepsis, aging or inflammation.

24. The method of claim 22, further comprising administering to the subject an effective amount of at least one compound selected from the group consisting of: a cytotoxic agent, cisplatin, doxorubicin, taxotere, taxol, etoposide, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, the epothilones, tamoxifen, 5-fluorouracil, methotrexate, temozolomide, cyclophosphamide, SCH 66336, tipifarnib (Zarnestra[®]), R115777, L778,123, BMS 214662, Iressa[®], Tarceva[®], C225, GLEEVEC[®], intron[®], Peg-Intron[®], aromatase combinations, ara-C, adriamycin, cytoxan, gemcitabine, Uracil mustard, Chlormethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine,

Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, leucovorin, oxaliplatin (ELOXATIN[®]), Pentostatine, vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, MithramycinTM,
5 Deoxycoformycin, Mitomycin-C, L-Asparaginase, Teniposide 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrol acetate, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide,
10 Flutamide, Toremifene, goserelin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, Anastrozole, Letrozole, Capecitabine, Reloxafine, Droloxafine, Hexamethylmelamine, Avastin, herceptin, Bexxar, Velcade, Zevalin, Trisenox, Xeloda, Vinorelbine, Porfimer, Erbitux, Liposomal, Thiotepa, Altretamine, Melphalan, Trastuzumab, Lerazole,
15 Fulvestrant, Exemestane, Ifosfomide, Rituximab, C225, Campath, leucovorin, dexamethasone, bicalutamide, chlorambucil, letrozole, megestrol, valrubicin, vinblastine, and NIASPAN[®].

25. The method of claim 22 further comprising administering an effective amount
20 of a rescuing agent.

26. The pharmaceutical composition of claim 25, wherein the rescuing agent is selected from the group consisting of nicotinamide, nicotinic acid, and nicotinamide mononucleotide (NMN).

25

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2013/000216

A. CLASSIFICATION OF SUBJECT MATTER

See extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: C07D 401/-; A61K 31/-; A61P 35/-

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CNKI, CNPAT, EPODOC, WPI, REGISTRY, CAPLUS: Genentech, sipiro, azaspiro, octan+, amide?, cancer, tumor?, tumour?, leukemia

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2007/0280943 A1 (FRIEDMAN, Steven M. et al.) 06 December 2007(06.12.2007) See the whole document	1-26
A	WO 2009/098448 A1 (ASTRAZENECA AB et al.) 13 August 2009 (13.08.2009) See the whole document	1-26
A	WO 2007/062093 A2 (INCYTE CORPORATION et al.) 31 May 2007(31.05.2007) See the whole document	1-26
A	EP 2364982 A1 (ONO PHARMACEUTICAL CO., LTD. et al.) 14 September 2011 (14.09.2011) See the whole document	1-26

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:	“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
“A” document defining the general state of the art which is not considered to be of particular relevance	“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
“E” earlier application or patent but published on or after the international filing date	“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
“L” document which may throw doubts on priority claim (S) or which is cited to establish the publication date of another citation or other special reason (as specified)	“&” document member of the same patent family
“O” document referring to an oral disclosure, use, exhibition or other means	
“P” document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 15 May 2013 (15.05.2013)	Date of mailing of the international search report 06 Jun. 2013 (06.06.2013)
---	--

Name and mailing address of the ISA/CN
The State Intellectual Property Office, the P.R.China
6 Xitucheng Rd., Jimen Bridge, Haidian District, Beijing, China
100088
Facsimile No. 86-10-62019451

Authorized officer

JIANG Shichao

Telephone No. (86-10)82246761

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2013/000216

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: 22-25
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 22-25 are directed to methods for treating the alleged diseases of human/animal body, the search has been carried out and based on the alleged effects of the compounds/compositions.
2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CN2013/000216

Patent Documents referred in the Report	Publication Date	Patent Family	Publication Date
US2007280943A1	06.12.2007	WO2007143600A2	13.12.2007
		WO2007143600A3	14.02.2008
		US7910108B2	22.03.2011
WO2009098448A1	13.08.2009	US2009298807A1	03.12.2009
		WO2009098448A8	26.08.2010
		AU2009211251A1	13.08.2009
		EP2242759A1	27.10.2010
		KR20100121637A	18.11.2010
		CA2713322A1	13.08.2009
		INMUMNP201001622E	26.11.2010
		TW200938549A	16.09.2009
		MXPA10008577A	30.11.2010
		JP2011512331A	21.04.2011
		CN102083839A	01.06.2011
		VN26090A	25.05.2011
		PH12010501811A	13.08.2009
		ZA201005100A	28.12.2011
		HK1149258A0	30.09.2011
		US8148373B2	03.04.2012
		NZ586872A	30.03.2012
WO2007062093A2	31.05.2007	AU2009211251B2	02.02.2012
		JP4934223B2	16.05.2012
		MX299866B	05.06.2012
		HK1158196A0	13.07.2012
		EP2242759B1	12.09.2012
		US2007117809A1	24.05.2007
		EP1951242A2	06.08.2008
WO2007062093A3	23.10.2008		
		JP2009516747A	23.04.2009

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CN2013/000216

Patent Documents referred in the Report	Publication Date	Patent Family	Publication Date
EP2364982A1	14.09.2011	WO2004092169A1	28.10.2004
		EP1619193A1	25.01.2006
		JP2005505481T2	06.07.2006
		US2006229301A1	12.10.2006
		US7498323B2	03.03.2009
		US2010261641A1	14.10.2010
		JP4710606B2B2	29.06.2011

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CN2013/000216

A. CLASSIFICATION OF SUBJECT MATTER

C07D 401/12 (2006.01) i

C07D 401/14 (2006.01) i

A61K 31/438 (2006.01) i

A61K 31/444 (2006.01) i

A61P 35/00 (2006.01) i

A61P 35/02 (2006.01) i

A61P 35/04 (2006.01) i