

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3198018 T3**

(12) **Oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **C 12 N 15/864 (2006.01)** **A 61 K 48/00 (2006.01)** **C 12 N 15/90 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2021-03-01**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2020-12-09**
- (86) Europæisk ansøgning nr.: **15775884.8**
- (86) Europæisk indleveringsdag: **2015-09-23**
- (87) Den europæiske ansøgnings publiceringsdag: **2017-08-02**
- (86) International ansøgning nr.: **US2015051785**
- (87) Internationalt publikationsnr.: **WO2016049230**
- (30) Prioritet: **2014-09-24 US 201462054899 P** **2014-10-14 US 201462063587 P**
2015-08-25 US 201562209862 P
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **City of Hope, 1500 East Duarte Road, Duarte, CA 91010-3000, USA**
- (72) Opfinder: **CHATTERJEE, Saswati, 614 Church Canyon Place, Altadena, California 91001, USA**
SMITH, Laura, 405 N. Madison Avenue, Apt. 21, Pasadena, California 91101, USA
WONG, Kamehameha, 614 Church Canyon Place, Altadena, California 91001, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **VEKTORVARIANTER AF ADENOASSOCIERET VIRUS TIL HØJEFFEKTIV GENOMREDIGERING OG FREMGANGSMÅDER DERTIL**
- (56) Fremdragne publikationer:
WO-A1-2012/051343
US-A1- 2011 294 218
US-A1- 2013 096 182
LAURA J. SMITH ET AL: "Gene Transfer Properties and Structural Modeling of Human Stem Cell-Derived AAV", MOLECULAR THERAPY, 13 June 2014 (2014-06-13), XP055232861, GB ISSN: 1525-0016, DOI: 10.1038/mt.2014.107
IRAM F KHAN ET AL: "AAV-mediated gene targeting methods for human cells", NATURE PROTOCOLS, vol. 6, no. 4, 1 March 2011 (2011-03-01), pages 482-501, XP055232661, GB ISSN: 1754-2189, DOI: 10.1038/nprot.2011.301
SMITH LAURA J ET AL: "Receptor Binding and Post-Entry Processing of Stem Cell-Derived AAV Isolates in CD34+Cells", MOLECULAR THERAPY, NATURE PUBLISHING GROUP, GB , vol. 22, no. Supplement 1 24 May 2014 (2014-05-24), pages S27-S28, XP008178251, ISSN: 1525-0016, DOI: 10.1038/MT.2014.64 Retrieved from the Internet: URL:http://www.nature.com/mt/index.html
UL-HASAN TAIHRA ET AL: "Novel AAV Vectors Mediate Efficient Intramuscular Gene Delivery", MOLECULAR

Fortsættes ...

THERAPY, NATURE PUBLISHING GROUP, GB , vol. 22, no. Supplement 1 24 May 2014 (2014-05-24), pages S115-S116, XP008178252, ISSN: 1525-0016, DOI: 10.1038/MT.2014.64 Retrieved from the Internet: URL:<http://www.nature.com/mt/journal/v22/n1s/pdf/mt201464a.pdf>

CHANDRA MANASA ET AL: "Genomic Integration Profile of AAVHSC Vectors in Human CD34+Long-Term Engrafted Hematopoietic Stem Cell Xenografts in Immune-Deficient Mice", MOLECULAR THERAPY, vol. 23, no. Suppl. 1, May 2015 (2015-05), pages S40-S41, XP002751826, & 18TH ANNUAL MEETING OF THE AMERICAN-SOCIETY-OF-GENE-AND-CELL-THERAPY (ASGCT); NEW ORLEANS, LA, USA; MAY 13 -16, 2015

MILLER DANIEL G ET AL: "Gene targeting in vivo by adeno-associated virus vectors", NATURE BIOTECHNOLOGY, GALE GROUP INC, US, vol. 24, no. 8, 1 August 2006 (2006-08-01) , pages 1022-1026, XP002459153, ISSN: 1087-0156, DOI: 10.1038/NBT1231

SMITH LAURA J ET AL: "Stem cell-derived clade F AAVs mediate high-efficiency homologous recombination-based genome editing", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 115, no. 31, 31 July 2018 (2018-07-31), pages E7379-E7388, XP55609078, ISSN: 0027-8424

DESCRIPTION

[0001] The present invention was made with government support under Grant No. HL087285 awarded by the National Institutes of Health. The Government has certain rights in the invention.

BACKGROUND

[0002] The adeno-associated virus (AAV) genome is built of single-stranded deoxyribonucleic acid (ssDNA), either positive- or negative-sensed, which is about 4.9 kilobase long. The genome comprises inverted terminal repeats (ITRs) at both ends of the DNA strand, and two open reading frames (ORFs): rep and cap. Rep is composed of four overlapping genes encoding rep proteins required for the AAV life cycle, and cap contains overlapping nucleotide sequences of capsid proteins: VP1, VP2 and VP3, which interact together to form a capsid of an icosahedral symmetry.

[0003] Recombinant adeno-associated virus (rAAV) vectors derived from the replication defective human parvovirus AAV2 are proving to be safe and effective gene transfer vehicles that have yet to be definitively identified as either pathogenic or oncogenic [3-4, 6, 18-19, 26, 31]. rAAV transduce non-dividing primary cells, are low in immunogenicity, and direct sustained transgene expression in vivo [6, 10, 20]. Infection with wild type AAV is associated with inhibition of oncogenic transformation and AAV inverted terminal repeats may actually confer oncoprotection [2, 28, 52-55]. A recent survey of panels of human tissues found that the marrow and liver were the two most common sites of naturally occurring AAV isolates in humans, suggesting that infection of marrow cells by AAV is not rare.

[0004] Use of viral vectors for gene therapy has been long considered. Due to its potential for long-lived correction and the ease of ex vivo manipulation, the hematopoietic system was one of the earliest targets of gene therapy. Despite significant effort, however, actual therapeutic success remains elusive [5]. This is due to the recognized inability of most viral vectors to efficiently transduce quiescent, non-dividing hematopoietic stem cells (HSC) [23] as well as safety concerns arising from insertional oncogenesis [15, 22]. However, stable gene transfer has been successfully demonstrated to both murine and human HSC by rAAV [8, 11-12, 24, 27, 29-30, 37].

[0005] It has been additionally difficult to effectively use viral vectors in gene therapy for treating neurological conditions, particularly central nervous system diseases or disorders due to the difficulty of crossing the blood-brain barrier, a cellular and metabolic separation of the circulating blood from the brain extracellular fluid created by tight junctions between endothelial cells that restrict the passage of solutes.

[0006] CD34 is cell surface glycoprotein and a cell-cell adhesion factor. CD34 protein is

expressed in early hematopoietic and vascular tissue and a cell expressing CD34 is designated CD34⁺. Chromosomal integration of rAAV in human CD34⁺ HSC [8, 12, 16, 29] and efficient transduction of primitive, pluripotent, self-renewing human HSC capable of supporting primary and secondary multi-lineage engraftment has been demonstrated in immune-deficient NOD-SCID mice [29]. Transduction of primitive HSC capable of supporting serial engraftment was shown to be attributable to the propensity of rAAV to efficiently transduce primitive, quiescent CD34⁺CD38⁻ cells residing in GO [24]. Despite several reports of successful rAAV-mediated gene transfer into human HSC in vitro and in murine and non-human primate HSC in vivo, controversy regarding the utility of rAAV for HSC transduction still persists. These discrepancies arose primarily from short-term in vitro studies that assessed transduction by expression profiling and are attributable to the identified restrictions to transgene expression from rAAV2, including viral uncoating [35], intracellular trafficking [33], nuclear transport and second strand synthesis [36].

[0007] While AAV2 remains the best-studied prototypic virus for AAV-based vectors [1, 13, 18, 21], the identification of a large number of new AAV serotypes significantly enhances the repertoire of potential gene transfer vectors [14]. AAV1, 3 and 4 were isolated as contaminants of adenovirus stocks, and AAV5 was isolated from a human condylomatous wart. AAV6 arose as a laboratory recombinant between AAV1 and AAV2. Recently, more than 100 distinct isolates of naturally occurring AAV in human and non-human primate tissues were identified. This led to the use of capsids derived from some of these isolates for pseudotyping, replacing the envelope proteins of AAV2 with the novel envelopes, whereby rAAV2 genomes are then packaged using AAV2 rep and novel capsid genes. The use of novel capsids, the proteins as part of the viral shell, resulted in the circumvention of many limitations in transgene expression associated with AAV2 [32, 35-36].

[0008] In an effort to circumvent these restrictions, recent research has shown that novel capsid sequences result in reduced proteasome-mediated capsid degradation, increased nuclear trafficking and retention. Novel capsids, many of which utilize novel receptors, broadens the tropism of rAAV allowing for efficient transduction of previously refractory tissues and provides a means of circumventing highly prevalent pre-existing serologic immunity to AAV2, which posed major clinical limitations in a recent trial. Notably, some novel capsids appear to alter the intracellular processing of rAAV. For example, uncoating and transgene expression is accelerated in the context of AAV8 as compared to native AAV2 capsids. Recently, transgene expression was shown to be based upon capsid proteins, regardless of the serotype origin of the inverted terminal repeats (ITRs).

[0009] Naturally occurring AAV is identifiable in cytokine-primed peripheral blood stem cells. Capsid sequences of these AAV are unique. These capsids are capable of pseudotyping recombinant AAV2 genomes. US Patent Publication Number 20130096182A1 describes capsids AAVF1-17, and use thereof for cell transduction and gene transfer. Any improvement in the area of gene therapy regarding both permanent and reversible gene transfer and expression for therapeutic purposes would be a significant improvement in the art. Moreover, safe and efficient gene delivery to stem cells remains a significant challenge in the field despite

decades of research. Therefore the ability to genetically modify stem cells safely would represent a significant advance.

[0010] Further, genome editing by gene targeting or correction at a specific site in the genome without leaving a footprint in the genome is attractive for the precise correction of inherited and acquired diseases. Current technology accomplishes this through the use of exogenous endonucleases such as zinc finger nucleases, TAL endonucleases or caspase 9/CRISPR systems. However, these "traditional" approaches are associated with toxicity and off target effects of endonuclease cleavage. Therefore, the ability to genetically modify stem cells safely and efficiently at high frequencies without the need for exogenous endonuclease cleavage would represent a significant advance.

[0011] Additionally, current methods of genetic transduction of human HSCs involve *ex vivo* transduction of purified donor stem cells followed by transplantation into usually "conditioned" recipients. The cell harvest procedures are invasive and involve either bone marrow harvest or multiple days of granulocyte-colony stimulating factor (G-CSF) priming of the donor followed by apheresis. The *ex vivo* transduction procedures can affect the hematopoietic potential of the stem cells. Additionally, *in vitro* transduced cells must be tested for sterility, toxicity, etc. before transplantation. Prior to transplanting into recipients, the stem cells often have to undergo conditioning with chemotherapy or radiation to ensure engraftment. The process usually requires hospitalization of patients for at least several days and sometimes longer. Overall, this is an arduous, expensive and high risk procedure that greatly limits the utility of stem cell gene therapy. A procedure is needed that offers a better alternative to current stem cell transduction methods without the need for purification and *ex vivo* transduction.

[0012] Khan et al. (2011; Nature Protocols; 6(4): 482-501) provides an overview of AAV-derived vectors for gene targeting of different human cell types. Miller et al. (2006; Nature Biotechnology; 24(8): 1022-1026) describes AAV-mediated gene targeting *in vivo* to correct the GusB gene mutation responsible for murine mucopolysaccharidosis type VII. US2013/096182 describes AAV F clade isolates from human CD34⁺ hematopoietic stem cells for use in high-efficiency transduction of human cord blood stem cells.

SUMMARY

[0013] The present invention provides an *ex vivo* method for editing a target locus of a mammalian chromosome, wherein the method comprises transducing a mammalian cell comprising the mammalian chromosome with a replication-defective adeno-associated virus (AAV) comprising

1. a) an AAV Clade F capsid; and
2. b) a correction genome comprising:
 1. (i) an editing element for insertion of one or more nucleotides, deletion of one or more nucleotides, or substitution of one or more nucleotides relative to the target

locus, wherein the editing element is selected from an internucleotide bond or a nucleotide sequence for integration into the target locus;

2. (ii) a 5' homologous arm nucleotide sequence 5' of the editing element, having homology to a 5' region of the mammalian chromosome relative to the target locus; and
3. (iii) a 3' homologous arm nucleotide sequence 3' of the editing element, having homology to a 3' region of the mammalian chromosome relative to the target locus,

wherein the cell is transduced without co-transducing an exogenous nuclease or a nucleotide sequence that encodes an exogenous nuclease, wherein the AAV Clade F capsid comprises:

1. (i) a capsid protein comprising an amino acid sequence having 100% sequence identity with amino acids 203 to 736 of any one of SEQ ID NOs: 2-17;
2. (ii) a capsid protein comprising an amino acid sequence having 100% sequence identity with amino acids 138 to 736 of any one of SEQ ID NOs: 2-17; and/or
3. (iii) a capsid protein comprising an amino acid sequence having 100% sequence identity with amino acids 1 to 736 of any one of SEQ ID NOs: 2-17.

[0014] The present invention also provides a replication-defective adeno-associated virus (AAV) for use in a method of treating a disease or disorder in a subject, the method comprising administering the replication-defective AAV to the subject without co-administering an exogenous nuclease or a nucleotide sequence that encodes an exogenous nuclease, wherein the replication-defective AAV comprises:

1. a) an AAV Clade F capsid; and
2. b) a correction genome comprising:
 1. (i) an editing element for insertion of one or more nucleotides, deletion of one or more nucleotides, or substitution of one or more nucleotides relative to the target locus, wherein the editing element is selected from an internucleotide bond or a nucleotide sequence for integration into a target locus of a mammalian chromosome;
 2. (ii) a 5' homologous arm nucleotide sequence 5' of the editing element, having homology to a 5' region of the mammalian chromosome relative to the target locus; and
 3. (iii) a 3' homologous arm nucleotide sequence 3' of the editing element, having homology to a 3' region of the mammalian chromosome relative to the target locus,

wherein the AAV Clade F capsid comprises:

1. (i) a capsid protein comprising an amino acid sequence having 100% sequence identity with amino acids 203 to 736 of any one of SEQ ID NOs: 2-17;
2. (ii) a capsid protein comprising an amino acid sequence having 100% sequence identity with amino acids 138 to 736 of any one of SEQ ID NOs: 2-17; and/or
3. (iii) a capsid protein comprising an amino acid sequence having 100% sequence

identity with amino acids 1 to 736 of any one of SEQ ID NOs: 2-17.

[0015] In some embodiments, the AAV has a chromosomal integration efficiency of at least about 1% in the absence of an exogenous nuclease for integrating the editing element into the target locus of the mammalian chromosome in the cell.

[0016] In some embodiments, the correction genome comprises a 5' inverted terminal repeat (5' ITR) nucleotide sequence 5' of the 5' homologous arm nucleotide sequence, and a 3' inverted terminal repeat (3' ITR) nucleotide sequence 3' of the 3' homologous arm nucleotide sequence. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially identical to an AAV2 virus 5' ITR and an AAV2 virus 3' ITR, respectively. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially mirror images of each other. In some embodiments, the 5' ITR nucleotide sequence has at least 95% sequence identity to SEQ ID NO:36, and the 3' ITR nucleotide sequence has at least 95% sequence identity to SEQ ID NO:37. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially identical to an AAV5 virus 5' ITR and an AAV5 virus 3' ITR, respectively. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially mirror images of each other. In some embodiments, the 5' ITR nucleotide sequence has at least 95% sequence identity to SEQ ID NO:38, and the 3' ITR nucleotide sequence has at least 95% sequence identity to SEQ ID NO:39.

[0017] In some embodiments, the correction genome has an essential absence of a promoter operatively linked to the editing element nucleotide sequence. In some embodiments, the correction genome further comprises an exogenous promoter operatively linked to the editing element.

[0018] In some embodiments, the replication-defective AAV genome comprises an essential absence of an AAV *rep* gene and an AAV *cap* gene.

[0019] In some embodiments, each of the 5' and 3' homologous arm nucleotide sequences independently has a nucleotide length of between about 500 to 1000 nucleotides or between about 600 to 1000 nucleotides. In some embodiments, the 5' and 3' homologous arm nucleotide sequences have substantially equal nucleotide lengths. In some embodiments, the 5' and 3' homologous arm nucleotide sequences have asymmetrical nucleotide lengths. In some embodiments, the 5' homologous arm nucleotide sequence has at least about 95% nucleotide sequence identity to the 5' region of the mammalian chromosome relative to the target locus. In some embodiments, the 3' homologous arm nucleotide sequence has at least about 95% nucleotide sequence identity to the 3' region of the mammalian chromosome relative to the target locus. In some embodiments, the 5' homologous arm nucleotide sequence has 100% sequence identity to the 5' region of the mammalian chromosome relative to the target locus and the 3' homologous arm nucleotide sequence has 100% sequence

identity to the 3' region of the mammalian chromosome relative to the target locus.

[0020] In some embodiments, the editing element consists of one nucleotide. In some embodiments, the target locus is a nucleotide sequence consisting of one nucleotide, and the target locus represents a point mutation of the mammalian chromosome.

[0021] In some embodiments, the target locus can comprise an intron of a mammalian chromosome. In some embodiments, the target locus can comprise an exon of a mammalian chromosome. In some embodiments, the target locus can comprise a non-coding region of a mammalian chromosome. In some embodiments, the target locus can comprise a regulatory region of a mammalian chromosome. In some embodiments, the target locus may be a locus associated with a disease state as described herein.

[0022] In some embodiments, the editing element comprises at least 1, 2, 10, 100, 200, 500, 1000, 1500, 2000, 3000, 4000, or 5000 nucleotides. In some embodiments, the editing element comprises 1 to 5500, 1 to 5000, 1 to 4500, 1 to 4000, 1 to 3000, 1 to 2000, 1 to 1000, 1 to 500, 1 to 200, or 1 to 100 nucleotides, or 2 to 5500, 2 to 5000, 2 to 4500, 2 to 4000, 2 to 3000, 2 to 2000, 2 to 1000, 2 to 500, 2 to 200, or 2 to 100 nucleotides, or 10 to 5500, 10 to 5000, 10 to 4500, 10 to 4000, 10 to 3000, 10 to 2000, 10 to 1000, 10 to 500, 10 to 200, or 10 to 100 nucleotides.

[0023] In some embodiments, the editing element comprises an exon, an intron, a 5' untranslated region (UTR), a 3' UTR, a promoter, a splice donor, a splice acceptor, a sequence encoding or non-coding RNA, an insulator, a gene, or a combination thereof. In some embodiments, the editing element is a fragment of a coding sequence of a gene within or spanning the target locus.

[0024] In some embodiments, the target locus is a nucleotide sequence comprising n nucleotides where n is an integer greater than or equal to one; the editing element comprises m nucleotides where m is an integer equal to n ; and the editing element represents a substitution for the target locus of the mammalian chromosome. In some embodiments, the target locus is a nucleotide sequence comprising n nucleotides where n is an integer greater than or equal to one; the editing element comprises m nucleotides where m is an integer greater than n ; and the editing element represents a substitutive addition for the target locus of the mammalian chromosome. In some embodiments, the target locus is a nucleotide sequence comprising n nucleotides where n is an integer greater than or equal to two; the editing element comprises m nucleotides where m is an integer less than n ; and the editing element represents a substitutive deletion for the target locus of the mammalian chromosome. In some embodiments, the target locus is an internucleotide bond; the editing element comprises m nucleotides where m is an integer greater than or equal to one; and the editing element represents an addition for the target locus of the mammalian chromosome.

[0025] In some embodiments, the editing element is an internucleotide bond. In some embodiments, the target locus is a nucleotide sequence comprising one or more nucleotides,

and the editing element comprises a deletion for the target locus of the mammalian chromosome.

[0026] In some embodiments, the target locus of the mammalian chromosome is a mutant target locus comprising one or more mutant nucleotides, relative to a corresponding wild type mammalian chromosome. In some embodiments, the mutant target locus comprises a point mutation, a missense mutation, a nonsense mutation, an insertion of one or more nucleotides, a deletion of one or more nucleotides, or combinations thereof. In some embodiments, the mutant target locus comprises an amorphic mutation, a neomorphic mutation, or an antimorphic mutation. In some embodiments, the mutant target locus comprises an autosomal dominant mutation, an autosomal recessive mutation, a heterozygous mutation, a homozygous mutation, or combinations thereof. In some embodiments, the mutant target locus is selected from a promoter, an enhancer, a signal sequence, an intron, an exon, a splice donor site, a splice acceptor site, an internal ribosome entry site, an inverted exon, an insulator, a gene, a chromosomal inversion, and a chromosomal translocation within the mammalian chromosome.

[0027] The AAV Clade F capsid comprises at least one Clade F VP3 as set forth in any one of SEQ ID Nos: 2-17. In some embodiments, the AAV Clade F capsid comprises Clade F VP2 and Clade F VP3 as set forth in any one of SEQ ID Nos: 2-17. In some embodiments, the AAV Clade F capsid comprises Clade F VP1, Clade F VP2 and Clade F VP3 proteins as set forth in any one of SEQ ID Nos: 2-17. In some embodiments, the AAV Clade F capsid comprises a VP1 capsid protein selected from a VP1 capsid protein of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which corresponds to amino acids 1 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively. In some embodiments, the AAV Clade F capsid comprises a VP1 and a VP2 capsid protein independently selected from a VP1 and VP2 capsid protein of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which correspond to amino acids 1 to 736 and amino acids 138 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively. In some embodiments, the AAV Clade F capsid comprises a VP1, a VP2 and a VP3 capsid protein independently selected from a VP1, VP2 and VP3 capsid protein of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which correspond to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively. In some embodiments, the AAV Clade F capsid comprises each of the VP1, VP2 and VP3 capsid proteins of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which correspond to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively.

[0028] As used herein, AAVF1, AAVF2, AAVF3, AAVF4, AAVF5, AAVF6, AAVF7, AAVF8, AAVF9, AAVF10, AAVF11, AAVF12, AAVF13, AAVF14, AAVF15, AAVF16, and AAVF17 are also referred to as AAVHSC1, AAVHSC2, AAVHSC3, AAVHSC4, AAVHSC5, AAVHSC6, AAVHSC7, AAVHSC8, AAVHSC9, AAVHSC10, AAVHSC11, AAVHSC12, AAVHSC13, AAVHSC14, AAVHSC15, AAVHSC16, and AAVHSC17, respectively. In other words, any recitation of AAVF1, AAVF2, AAVF3, AAVF4, AAVF5, AAVF6, AAVF7, AAVF8, AAVF9, AAVF10, AAVF11, AAVF12,

AAVF13, AAVF14, AAVF15, AAVF16, or AAVF17 is equivalent to and may be replaced with AAVHSC1, AAVHSC2, AAVHSC3, AAVHSC4, AAVHSC5, AAVHSC6, AAVHSC7, AAVHSC8, AAVHSC9, AAVHSC10, AAVHSC11, AAVHSC12, AAVHSC13, AAVHSC14, AAVHSC15, AAVHSC16, or AAVHSC17, respectively.

[0029] In some embodiments, the mammalian chromosome is selected from human chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, X and Y. In some embodiments, the mammalian chromosome is selected from mouse chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, X and Y. In some embodiments, the mammalian chromosome is not human chromosome 19.

[0030] In some embodiments, the mammalian chromosome is a somatic cell chromosome. In some embodiments, the somatic cell is from a tissue selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, and epithelial tissue. In some embodiments, the somatic cell is from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage. In some embodiments, the somatic cell is a CD34+ cell.

[0031] In some embodiments, the cell is a stem cell. In some embodiments, the stem cell is a hematopoietic stem cell, a cord blood stem cell, a bone marrow stem cell, a fetal liver stem cell, or a peripheral blood stem cell. In some embodiments, the cell is selected from the group consisting of a CD34+ Hematopoietic stem cell line (HSC), a K562 CD34+ leukemia cell line, a HepG2 human liver cell line, a peripheral blood stem cell, a cord blood stem cell, a CD34+ peripheral blood stem cell, a WI-38 human diploid fibroblast cell line, a MCF7 human breast cancer cell line, a Y79 human retinoblastoma cell line, a SCID-X1 LBL human EBV-immortalized B cell line, a primary human hepatocyte, a primary hepatic sinusoidal endothelial cell, and a primary skeletal muscle myoblast.

[0032] In some embodiments, the AAV has a chromosomal integration efficiency of at least about 5% for integrating the editing element into the target locus of the mammalian chromosome in the cell. In some embodiments, the AAV has a chromosomal integration efficiency of at least about 10% for integrating the editing element into the target locus of the mammalian chromosome in the cell.

[0033] Other aspects of the disclosure relate to a composition comprising an AAV as described herein, wherein the composition is in a pharmaceutically acceptable formulation. The formulation may be constituted for administration to a mammal. The formulation may be constituted for administration to a mammal via intravenous injection, subcutaneous injection, intramuscular injection, autologous cell transfer, or allogeneic cell transfer. The pharmaceutically acceptable formulation may comprise an excipient. The excipient may be selected from a carrier, an adjuvant and a vehicle, or combinations thereof.

[0034] Further described herein is a packaging system for recombinant preparation of an adeno-associated virus (AAV), wherein the packaging system comprises a Rep nucleotide

sequence encoding one or more AAV Rep proteins; a Cap nucleotide sequence encoding one or more AAV Cap proteins of an AAV Clade F capsid; and a correction genome as described herein; wherein the packaging system is operative in a cell for enclosing the correction genome in the capsid to form the adeno-associated virus. The packaging system may comprise a first vector comprising the Rep nucleotide sequence and the Cap nucleotide sequence, and a second vector comprising the correction genome. The AAV Clade F capsid may comprise at least one protein selected from Clade F VP1, Clade F VP2 and Clade F VP3. The AAV Clade F capsid may comprise at least two proteins selected from Clade F VP1, Clade F VP2 and Clade F VP3. The AAV Clade F capsid may comprise Clade F VP1, Clade F VP2 and Clade F VP3 proteins. The AAV Clade F capsid may be any AAV Clade F capsid as described herein. The Rep nucleotide sequence may encode an AAV2 Rep protein. The AAV2 Rep protein encoded may be at least one of Rep 78/68 or Rep 68/52. The nucleotide sequence encoding the AAV2 Rep protein may comprise a nucleotide sequence having a minimum percent sequence identity to the AAV2 Rep nucleotide sequence of SEQ ID NO:40, wherein the minimum percent sequence identity is at least 70% across the length of the nucleotide sequence encoding the AAV2 Rep protein. In any one of the packaging systems described, the packaging system may further comprise a third vector, wherein the third vector is a helper virus vector. The helper virus vector may be an independent third vector. The helper virus vector may be integral with the first vector. The helper virus vector may be integral with the second vector. The third vector may comprise genes encoding helper virus proteins. The helper virus may be selected from the group consisting of adenovirus, herpes virus (including herpes simplex virus (HSV)), vaccinia virus, and cytomegalovirus (CMV). The helper virus may be adenovirus. The adenovirus genome may comprise one or more adenovirus RNA genes selected from the group consisting of E1, E2, E4 and VA. The helper virus may be HSV. The HSV genome may comprise one or more of HSV genes selected from the group consisting of UL5/8/52, ICPO, ICP4, ICP22 and UL30/UL42. The first vector and the third vector may be contained within a first transfecting plasmid. The nucleotides of the second vector and the third vector may be contained within a second transfecting plasmid. The nucleotides of the first vector and the third vector may be cloned into a recombinant helper virus. The nucleotides of the second vector and the third vector may be cloned into a recombinant helper virus. The AAV capsid may be the capsid of a Clade F AAV selected from the group consisting of AAV9, AAVF1, AAVF2, AAVF3, AAVF4, AAVF5, AAVF6, AAVF7, AAVF8, AAVF9, AAVF11, AAVF12, AAVF13, AAVF14, AAVF15, AAVF16, AAVF17, AAVHU31, and AAVHU32. Any of the packaging systems described herein may be comprised within a kit.

[0035] Also described herein is a method for recombinant preparation of an adeno-associated virus (AAV), wherein the method comprises transfecting or transducing a cell with a packaging system as described herein under conditions operative for enclosing the correction genome in the capsid to form the AAV.

[0036] As mentioned above, the disclosure provides an ex vivo method for editing a target locus of a mammalian chromosome, wherein the method comprises transducing a mammalian cell comprising the mammalian chromosome with an adeno-associated virus (AAV) as described herein. In some embodiments, the cell is a mammalian stem cell. In some

embodiments, the mammalian cell is from a tissue selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, and epithelial tissue. In some embodiments, the mammalian cell is from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage. In some embodiments, the mammalian cell is a stem cell. In some embodiments, the stem cell is a hematopoietic stem cell, a cord blood stem cell, or peripheral blood stem cell. In some embodiments, the mammalian cell is a myoblast, an endothelial cell, a liver cell, a fibroblast, a breast cell, a lymphocyte, or a retinal cell. Other aspects of the disclosure relate to a cell obtainable by any method described herein.

[0037] Another aspect of the disclosure relates to a method for editing a target locus of a mammalian genome, wherein the method comprises: (a) obtaining mammalian cells from a mammal; (b) culturing the mammalian cells ex-vivo to form an ex-vivo culture; (c) transducing the mammalian cells with an adeno-associated virus (AAV) as described herein in the ex-vivo culture to form transduced mammalian cells; and (d) administering the transduced mammalian cells to the mammal.

[0038] In other aspects, the disclosure relates to a method for editing a target locus of a mammalian genome, wherein the method comprises: (a) obtaining mammalian cells from a first mammal; (b) culturing the mammalian cells ex-vivo to form an ex-vivo culture; (c) transducing the mammalian cells with an adeno-associated virus (AAV) as described herein in the ex-vivo culture to form transduced mammalian cells; and (d) administering the transduced mammalian cells to a second mammal. The first mammal and the second mammal may be the same species. In some embodiments, the mammalian cells are from a tissue selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, and epithelial tissue. In some embodiments, the mammalian cells are from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage. In some embodiments, the mammalian cells are stem cells. In some embodiments, the stem cells are hematopoietic stem cells, cord blood stem cells, or peripheral blood stem cells. In some embodiments, the mammalian cells are a CD34+ cells. In some embodiments, the mammalian cells are myoblasts, endothelial cells, liver cells, fibroblasts, breast cells, lymphocytes, or retinal cells.

[0039] Another aspect of the disclosure relates to a method for editing a target locus of a mammalian genome, wherein the method comprises administering an AAV as described herein or a composition as described herein to a mammal in an amount effective to transduce cells of the mammal with the AAV in-vivo.

[0040] In any one of the methods provided, the AAV is transduced or administered without co-transducing or co-administering an exogenous nuclease or a nucleotide sequence that encodes an exogenous nuclease.

[0041] In some embodiments of any one of the methods provided, the AAV has a chromosomal integration efficiency of at least about 1% for integrating the editing element into

the target locus of the mammalian chromosome. In some embodiments, the chromosomal integration efficiency of the AAV is at least about 2%, 3%, 4% or 5% for integrating the editing element into the target locus of the mammalian chromosome. In some embodiments, the editing element of the correction genome is integrated into the target locus of the mammalian chromosome with a chromosomal integration efficiency of at least 10%, 20%, 40%, or 50% of the mammalian cells. In some embodiments, the editing element of the correction genome is integrated into the target locus of the mammalian chromosome with a chromosomal integration efficiency ranging from 10% to 70%, 20% to 70%, 40% to 70%, or 50% to 70% of the mammalian cells.

[0042] In some embodiments of any one of the methods provided, the AAV has a chromosomal integration efficiency further characterized by an allele frequency in a population of cells of at least about 10% for the allele comprising the editing element integrated into the target locus of the mammalian chromosome. In some embodiments, the AAV has a chromosomal integration efficiency further characterized by an allele frequency in a population of cells of at least about 50% for the allele comprising the editing element integrated into the target locus of the mammalian chromosome. In some embodiments, the AAV has a chromosomal integration efficiency further characterized by an allele frequency in a population of cells of at least about 75% for the allele comprising the editing element integrated into the target locus of the mammalian chromosome. In some embodiments, the allele frequency in a population of cells is an allele frequency in a population of cells in vitro.

[0043] Also described herein is a method for generating a transgenic non-human animal, the method comprising administering an AAV as described herein or a composition as described herein to a non-human animal; or transducing a non-human animal cell with the AAV as described herein or the composition as described herein and implanting the cell into a host non-human animal under conditions sufficient to generate a transgenic non-human animal from the host non-human animal (e.g., by allowing the implanted cell to form or become part of an embryo, which then develops in the host into a transgenic non-human animal). The transgenic non-human animal may be crossed with another non-human animal to generate further transgenic non-human animals. The non-human animal cell may be derived from a zygote or an embryo of a non-human animal. The non-human animal may be a mouse, rat, rabbit, pig, bovine, sheep, goat, chicken, cat, dog, ferret, or primate.

[0044] Further described herein is a transgenic non-human animal obtainable by a method described herein, such as a method described above. The transgenic non-human animal may be a mouse, rat, rabbit, pig, bovine, sheep, goat, chicken, cat, dog, ferret, or primate.

[0045] Also described herein is a tissue derived from a transgenic non-human animal as described herein. The tissue may be selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, endothelial tissue and epithelial tissue. The tissue may be from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage.

[0046] Further described herein is a cell derived from a transgenic non-human animal as described herein. The cell may be a primary cell. The cell may be a CD34+ cell, a myoblast, an endothelial cell, a liver cell, a fibroblast, a breast cell, a lymphocyte, or a retinal cell. The cell may be an inducible pluripotent stem (iPS) cell. The cell may be from a tissue selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, endothelial tissue, and epithelial tissue. The cell may be from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage. The cell may be a stem cell. The stem cell may be a hematopoietic stem cell, a cord blood stem cell, or peripheral blood stem cell.

[0047] Adeno-associated virus (AAV) Clade F vectors (e.g., replication-defective AAVs comprising correction genomes enclosed in a Clade F capsid) or AAV vector variants (e.g., replication-defective AAVs comprising capsid variants relative to AAV9 capsids) for editing the genome of a cell are described. An AAV Clade F vector or AAV vector variant may comprise one or more Clade F capsids or one or more capsid variants (relative to an AAV9 capsid), an editing element (also referred to herein as a targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (also referred to herein as a target site) of the genome, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site). The editing element (target cassette) may be contained within a correction genome as described herein comprising inverted terminal repeats (ITRs) as described herein. The one or more Clade F capsids or capsid variants may be any of the Clade F capsids or capsid variants described herein. The one or more Clade F capsids or capsid variants may comprise a polypeptide sequence selected from the group of AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13), variants, fragments, mutants and any combination thereof. The one or more Clade F capsids or the one or more capsid variants may comprise a polypeptide sequence selected from the group of AAVF5 (SEQ ID NO: 11), AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13), AAVF9 (SEQ ID NO: 10), AAVF16 (SEQ ID NO: 17), variants, fragments, mutants and any combination thereof. The one or more Clade F capsids or capsid variants may comprise a polypeptide sequence having a percent sequence identity of at least 95% to a polypeptide sequence selected from the group of AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13), variants, fragments, mutants and any combination thereof. The one or more Clade F capsids or capsid variants may comprise a polypeptide sequence having a percent sequence identity of at least 95% to a polypeptide sequence selected from the group of AAVF5 (SEQ ID NO: 11), AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13), AAVF9 (SEQ ID NO: 10), AAVF16 (SEQ ID NO: 17), variants, fragments, mutants and any combination thereof. In certain embodiments, the target locus (target site) may be a safe harbor site. In certain embodiments, the safe harbor site may be the AAVS1 locus on chromosome 19. In certain embodiments, the target locus (target site) may be a locus associated with a disease state as described herein. In certain embodiments, the cell may be a

stem cell. In certain embodiments, the stem cell may be a hematopoietic stem cell, a pluripotent stem cell, a non-human embryonic stem cell, or a mesenchymal stem cell.

[0048] According to certain embodiments, ex vivo methods of editing the genome of a cell as defined by the claims are provided. In certain embodiments, the methods of editing the genome of a cell may comprise transducing the cell with one or more AAV Clade F vectors (e.g., replication-defective AAVs comprising correction genomes enclosed in a Clade F capsid) as described herein. The transduction is performed without additional exogenous nucleases. The AAV Clade F vectors comprise one or more Clade F capsids as defined in the claims, an editing element (targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) of the genome, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site). The editing element (targeting cassette) may be contained within a correction genome as described herein comprising ITRs as described herein. In certain embodiments, the Clade F capsids may comprise a polypeptide sequence selected from the group of AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13). In certain embodiments, the Clade F capsids may comprise a polypeptide sequence selected from the group of AAVF5 (SEQ ID NO: 11), AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13), AAVF9 (SEQ ID NO: 10), AAVF16 (SEQ ID NO: 17). In certain embodiments, the AAV Clade F vector does not contain a promoter for the one or more therapeutic nucleotide sequences. In certain embodiments, the target locus (target site) may be a safe harbor site. In certain embodiments, the safe harbor site may be the AAVS1 locus on chromosome 19. In certain embodiments, the target locus (target site) may be a locus associated with a disease state as described herein. In certain embodiments, the cell may be a stem cell. In certain embodiments, the stem cell may be a hematopoietic stem cell, a pluripotent stem cell, a non-human embryonic stem cell, or a mesenchymal stem cell.

[0049] Also provided herein is a replication-defective AAV for use in methods of treating a disease or disorder in a subject by editing a genome of a cell of the subject. Further described are methods of treating a disease or disorder in a subject by editing a genome of a cell of the subject, including the steps of transducing the cell of the subject with an AAV Clade F vector or AAV vector variant as described herein and transplanting the transduced cell into the subject, wherein the transduced cell treats the disease or disorder. Transduction of the cell is performed without additional exogenous nucleases. In certain embodiments, AAV Clade F vectors may comprise one or more Clade F capsids as defined in the claims, an editing element (targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) of the genome, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site). The editing element (targeting cassette) may

be contained within a correction genome as described herein comprising ITRs as described herein. In certain embodiments, the Clade F capsids may comprise a polypeptide sequence selected from the group of AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13). In certain embodiments, the one or more Clade F capsids may comprise a polypeptide sequence selected from the group of AAVF5 (SEQ ID NO: 11), AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), AAVF17 (SEQ ID NO: 13), AAVF9 (SEQ ID NO: 10), AAVF16 (SEQ ID NO: 17). In certain embodiments, the AAV Clade F vector does not contain a promoter for the one or more therapeutic nucleotide sequences. In certain embodiments, the target locus (target site) may be a safe harbor site. In certain embodiments, the safe harbor site may be the AAVS1 locus on chromosome 19. In certain embodiments, the target locus (target site) may be a locus associated with a disease state as described herein. In certain embodiments, the cell may be a stem cell. In certain embodiments, the stem cell may be a hematopoietic stem cell, a pluripotent stem cell, a non-human embryonic stem cell, or a mesenchymal stem cell. In certain embodiments, the disease or disorder may be caused by one or more mutations in the cell genome. In certain embodiments, the disease or disorder may be selected from an inherited metabolic disease, lysosomal storage disease, mucopolysaccharidosis, immunodeficiency disease, and hemoglobinopathy disease and infection.

[0050] Also disclosed herein is a replication-defective AAV for use in a method of treating a disease or disorder in a subject by in vivo genome editing by directly administering the AAV Clade F vector to the subject in accordance with the claims. In certain embodiments, the AAV Clade F vector may comprise one or more Clade F capsids as defined by the claims, an editing element (targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) of the genome, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site), wherein the vector transduces a cell of the subject and integrates the one or more therapeutic nucleotide sequences into the genome of the cell. The editing element (target cassette) may be contained within a correction genome as described herein comprising ITRs as described herein. In certain embodiments, the one or more Clade F capsids may comprise a polypeptide sequence selected from the group of AAVF1 (SEQ ID NO: 2), AAVF2 (SEQ ID NO: 3), AAVF11 (SEQ ID NO: 4), AAVF3 (SEQ ID NO: 5), AAVF4 (SEQ ID NO: 6), AAVF6 (SEQ ID NO: 7), AAVF7 (SEQ ID NO: 8), AAVF8 (SEQ ID NO: 9), AAVF9 (SEQ ID NO: 10), AAVF5 (SEQ ID NO: 11), AAVF12 (SEQ ID NO: 12), AAVF17 (SEQ ID NO: 13), AAVF13 (SEQ ID NO: 14), AAVF14 (SEQ ID NO: 15), AAVF15 (SEQ ID NO: 16), and AAVF16 (SEQ ID NO: 17). In certain embodiments, the AAV Clade F vector does not contain a promoter for the one or more therapeutic nucleotide sequences. In certain embodiments, the target locus (target site) may be a safe harbor site. In certain embodiments, the safe harbor site may be the AAVS1 locus on chromosome 19. In certain embodiments, the target locus (target site) may be a locus associated with a disease state as described herein. In certain embodiments, the cell may be a stem cell. In certain embodiments, the stem cell may be a hematopoietic stem cell, a pluripotent stem cell, a non-human embryonic stem cell, or a

mesenchymal stem cell. In certain embodiments, the disease or disorder may be caused by one or more mutations in the cell genome. In certain embodiments, the disease or disorder may be selected from an inherited metabolic disease, lysosomal storage disease, mucopolysaccharidosis, immunodeficiency disease, and hemoglobinopathy disease and infection. Also described herein are kits comprising one or more AAV Clade F vectors or AAV vector variants for editing the genome of a cell.

BRIEF DESCRIPTION OF THE DRAWINGS

[0051] This application contains at least one drawing executed in color. Copies of this application with color drawing(s) will be provided by the Office upon request and payment of the necessary fees.

Figure 1 shows the alignment of Clade F AAV capsid variant polypeptide sequences in comparison to AAV9. A corresponding alignment of Clade F AAV capsid variant polynucleotide sequences is provided in Figure 1 of US Patent Publication Number US20130096182A1.

Figure 2 is a chart listing some of the nucleotide mutations in the capsid of each sequence, including the base change, the amino acid change, and whether it is in VP1 or VP3.

Figure 3 shows a schematic of a portion of one set of donor ITR-AAVS 1-FP vector constructs that were used for genome editing. The AAV vector contained 5' homology and 3' homology arms, and regulatory elements, which included a 2A sequence, splice acceptor sequence, and polyadenylation sequence. Yellow fluorescent protein ("YFP" or "FP") was used as the transgene. AAV2 ITRs flanked the homologous arms and the vector genome was packaged in AAVF capsids to form the AAVF-AAVS 1-FP donor vectors. Importantly, the vector containing the FP gene does not contain a promoter to drive expression. The FP gene will only be expressed if it integrates correctly into AAVS1, downstream from an endogenous chromosomal promoter.

Figure 4 shows a schematic map of the targeted chromosomal AAVS1 locus and the edited AAVS1 locus that was the target site for transgene integration mediated by the AAVF vector. The top schematic (Figure 4A), "Wild type AAVS1 locus", illustrates the wild-type AAVS1 locus that contains a 5' homology arm and a 3' homology arm, but does not contain a transgene. Amplification with primers located outside of the homology region using an "OUT Forward Primer Region" primer and an "OUT Reverse Primer Region" primer results in a fragment ~1.9 kb long (see line labeled "Fragment 1"), which indicates that the fragment does not contain an integrated transgene. The bottom schematic (Figure 4B), "Edited AAVS1 locus", illustrates the edited AAVS1 locus which contains a 5' homology arm, regulatory elements, an integrated transgene, and the 3' homology arm. Amplification with primers located outside of the homology region using an "OUT Forward Primer Region" primer and an "OUT Reverse Primer Region" primer results in a fragment ~3.0 kb long (see line labeled "Fragment 2"), which indicates that the fragment contains a transgene. Amplification of the 5' junction region (the junction between the 5' homology arm and the transgene) using an "OUT Forward Primer

Region" primer and an "In Reverse Primer" results in a fragment ~1.7 kb long (see line labeled "Fragment 3"). Amplification of the 3' junction region (the junction between the transgene and the 3' homology arm) using an "OUT Reverse Primer Region" primer and an "In Forward Primer" results in a fragment ~1.2 kb long (see line labeled "Fragment 4"). If the transgene is not integrated, there is no resulting product upon amplification of the 5' junction region or the 3' junction region.

Figure 5 shows representative scatter plots from flow cytometric analyses of YFP expression in K562 cells 24 hours after transduction. Cells were transduced with the AAVF7 FP vector at a variety of multiplicity of infections (MOIs) (A) Cells not transduced with any vector (untransduced), (B) 50,000 MOI, (C) 100,000 MOI, (D) 200,000 MOI, and (E) 400,000 MOI. Data shown is from representative samples. Events above the line of demarcation within each scatter plot represent FP expressing cells, indicating that in these cells, the promoterless FP gene from the Donor ITR-AAVS 1-FP vector integrated correctly into AAVS1 in the human chromosome 19, downstream from the endogenous chromosomal promoter.

Figure 6 shows representative scatter plots from flow cytometric analyses of YFP expression in K562 cells 72 hours after transduction. Cells were transduced with the AAVF7 FP vector at variety of multiplicity of infections (MOIs) (A) Cells not transduced with any vector (untransduced), (B) 50,000 MOI, (C) 100,000 MOI, (D) 200,000 MOI, and (E) 400,000 MOI. Events above the line of demarcation represent cells with correctly targeted integration of the promoterless FP gene in the Donor ITR-AAVS 1-FP vector.

Figure 7 shows the average percentage of YFP expression following targeted integration of the promoterless YFP transgene in the AAVS1 locus in CD34+ K562 leukemic cells. (A) A bar graph showing YFP expression 24 hours post-transduction with AAVF7 vector in cells with MOIs of 50,000; 100,000; 150,000; 200,000; 300,000; and 400,000. (B) A bar graph showing YFP expression 72 hours post-transduction with AAVF7 vector in cells with MOIs of 50,000; 100,000; 150,000; 200,000; and 400,000. Each bar represents data compiled from up to 7 samples.

Figure 8 shows PCR confirmation of targeted integration of the YFP transgene into AAVS1 locus in K562 cells. A) Gel showing amplified DNA from representative samples from K562 cells with no template, untransduced, or transduced with AAVF7 FP vector at an MOI of 100,000. Lane 1: DNA ladder, lane 2: no template control, lane 3: untransduced control, and lane 4: AAVF7 FP transduced K562. Arrows point to either the FP integrated AAVS1 -3.1 kb fragment or the non-integrated AAVS1 ~1.9 kb fragment. B) Gel showing amplified DNA from representative samples from K562 cells with no template, untransduced, or transduced with AAVF7 FP vector with an MOI of 100,000. Lane 1: DNA ladder, lane 2: no template control, lane 3: untransduced control, lane 4: AAVF7 FP vector transduced K562. The arrows point to either the amplified FP integrated AAVS1 -3.1 kb fragment or the amplified non-integrated AAVS1 ~1.9 kb fragment.

Figure 9 shows representative scatter plots of YFP expression in primary CD34+ cells after targeted integration 1 day post-transduction with AAVF FP vectors. (A) Cells not transduced with any vector (untransduced), (B) cells transduced with AAVF7 FP vector, and (C) cells

transduced with AAVF17 FP vector. Cells transduced with either AAVF7 or AAVF17 vector showed a significant amount of YFP expression compared with the untransduced cells (compare B and C, respectively, with A). YFP expression in (B) and (C) indicates that the promoterless FP gene delivered by the AAVF vector accurately integrated into the chromosomal AAVS1 locus.

Figure 10 shows representative scatter plots of YFP expression in primary CD34+ cells after targeted integration 4 days post-transduction with AAVF FP vectors. (A) Cells not transduced with any vector (untransduced), (B) cells transduced with AAVF7 FP vector, and (C) cells transduced with AAVF17 FP vector. Cells transduced with either AAVF7 or AAVF17 FP vector showed a significant amount of YFP expression compared with the untransduced cells (compare B and C, respectively, with A), indicating accurate targeted integration of the gene delivered by the AAVF vector.

Figure 11 shows representative scatter plots of YFP expression in primary CD34+ cells after targeted integration 18 days post-transduction with AAVF FP vectors from representative samples. (A) Cells not transduced with any vector (untransduced), (B) cells transduced with AAVF7 FP vector, and (C) cells transduced with AAVF17 FP vector. Cells transduced with either AAVF7 or AAVF17 FP vector showed a significant amount of YFP expression compared with the untransduced cells (compare B and C, respectively, with A).

Figure 12 shows YFP expression in primary CD34+ cells after targeted integration. (A) A table showing the percentage of YFP positive cells for untransduced cells and cells transduced with either AAVF7 FP vector or AAVF17 FP vector at 4, 18, 20, and 39 days post-transduction. (B) A line graph showing the frequency of YFP expressing primary CD34+ cells at 4, 18, 20, and 39 days post AAVF FP transduction with an MOI of 100,000. The line with diamonds represents untransduced cells, the line with squares represents cells transduced with AAVF7 FP vector, and the line with triangles represents cells transduced with AAVF17 FP vector.

Figure 13 shows PCR confirmation of targeted integration into the AAVS 1 locus in primary CD34+ cells. A gel showing amplified DNA from representative samples of primary CD34+ cells with no template, untransduced, or transduced with AAVF7 FP vector with an MOI of 150,000. Lane 1: DNA ladder, lane 2: no template control, lane 3: untransduced control, lane 4: DNA marker, and lane 5: AAVF7 FP vector transduced. The arrow points to the FP integrated AAVS1 (~1.7 kb fragment) showing the amplification product of the 5' junction region. Inset to the left shows the DNA ladder that was loaded in lane 1.

Figure 14 shows sequence confirmation of targeted integration of YFP gene sequences in the AAVS 1 locus beginning at the OUT Forward Primer Region. Sequencing results indicate that the YFP gene was present and was integrated correctly into the AAVS1 locus.

Figure 15 shows sequence confirmation of targeted integration of YFP sequences in the AAVS1 locus beginning near the 5' homology arm. Sequencing results indicate that the YFP gene was present and was integrated into the AAVS1 locus.

Figure 16 shows sequence confirmation of targeted integration of YFP sequences in the

AAVS1 locus beginning near the 5' region of the regulatory elements. Sequencing results indicate that the YFP gene was present and was integrated into the AAVS1 locus.

Figure 17 shows sequence confirmation of targeted integration of YFP sequences in the AAVS1 locus beginning near the 3' region of the regulatory elements. Sequencing results indicate that the YFP gene was present and was integrated into the AAVS1 locus.

Figure 18 shows sequence confirmation of targeted integration of YFP sequences in the AAVS1 locus beginning near the 5' region of the transgene. Sequencing results indicate that the YFP gene was present and was integrated into the AAVS1 locus.

Figure 19 shows sequence confirmation of targeted integration of YFP sequences in the AAVS1 locus beginning near the "IN Reverse Primer" region. Sequencing results indicate that the YFP gene was present and was integrated into the AAVS1 locus.

Figure 20 shows a schematic of the steps performed in the experiments in Example 4. One million human cord blood CD34+ cells were obtained (see Step 1) and injected into sublethally-irradiated immune deficient NOD/SCID adult mice (see Step 2). Two hours after injection with CD34+ cells, the mice were injected with AAVF-Luciferase vector (i.e., AAVF7-Luciferase vector or AAVF17-Luciferase vector). Two to seven days later the mice were injected with AAVF-Venus vectors (i.e., AAVF7-Venus vector or AAVF17-Venus vector) (see Step 3). Finally, in vivo luciferase expression was measured 4 weeks post injection and Venus expression was quantitated 6 weeks post-injection (see Step 4).

Figure 21 shows in vivo specific luciferase expression in representative recipients. Figure 21A shows that adult immune-deficient mice previously xenografted with human cord blood CD34+ HSCs that received intravenous injections of AAVF-Luciferase vector displayed specific luciferase expression in vertebrae, spleen, hips, and long bones, all sites of hematopoiesis after transplantation. Arrows indicate luciferase expression in vertebrae, spleen, liver, hips, and long bones. Flux for the liver and spleen was $4.08e9$ and flux for the tail was $1.74e9$. Figure 21B shows that adult immune-deficient mice that were not previously xenografted with human cord blood CD34+ HSCs that received intravenous injections of AAVF-Luciferase vector did not display high levels of specific luciferase expression. Flux for the liver and spleen was $1.47e8$ and flux for the tail was $2.22e8$.

Figure 22 shows histograms illustrating flow cytometry data of Venus-expressing human CD34+ or CD45+ cells in adult immune-deficient mice previously xenografted with human cord blood CD34+ HSCs that received intravenous injections of either AAVF7-Venus or AAVF17-Venus vectors. Figure 22A shows flow cytometry data from femoral CD34+ cells of xenografted mice injected with AAVF7-Venus vector. 9.23% of engrafted human hematopoietic cells expressed Venus. Figure 22B shows flow cytometry data from femoral CD45+ cells of xenografted mice injected with AAVF7-Venus vector. 8.35% of engrafted human hematopoietic cells expressed Venus. Figure 22C shows flow cytometry data from femoral CD34+ cells of xenografted mice injected with AAVF17-Venus vector. 8.92% of engrafted human hematopoietic cells expressed Venus. Figure 22D shows flow cytometry data from femoral CD45+ cells of xenografted mice injected with AAVF17-Venus vector. 8.59% of engrafted

human hematopoietic cells expressed Venus. Figure 22E shows flow cytometry data from vertebral CD45+ cells of xenografted mice injected with AAVF7-Venus vector. 15.3% of engrafted human hematopoietic cells expressed Venus. Figure 22F shows flow cytometry data from vertebral CD45+ cells of xenografted mice injected with AAVF17-Venus vector. 70.2% of engrafted human hematopoietic cells expressed Venus. Figure 22G shows flow cytometry data from spleen CD45+ cells of xenografted mice injected with AAVF7-Venus vector. 10.3% of engrafted human hematopoietic cells expressed Venus. Figure 22H shows flow cytometry data from spleen CD45+ cells of xenografted mice injected with AAVF17-Venus vector. 9.90% of engrafted human hematopoietic cells expressed Venus. Results from Figure 22 are also provided in Table 5.

Figure 23 shows a phylogram of the relationship of AAV Clade F viruses relative to each other and other AAV strains. The phylogram is based on nucleotide sequence homology of the capsid genes of AAVF viruses (Smith et al, Mol Ther. 2014 Sep;22(9):1625-34).

Figure 24 shows a map of a single stranded AAV vector genome for the insertion of a large DNA insert. The single stranded AAV2 genome contained AAV2 ITRs, homology arms, regulatory sequences and the promoterless Venus open reading frame (ORF). Venus is a fluorescent reporter protein. The promoterless Venus containing the Venus ORF is downstream from a splice acceptor and 2A sequence. The Venus ORF is followed by a polyadenylation signal. Each homology arms is 800bp long and targets Intron 1 of PPP1R12C gene on Chromosome 19

Figure 25 shows a schematic of an insertion site of an editing moiety in AAVS1. The transgene cassette consisting of the Venus open reading frame and a splice acceptor site followed by 2A sequence is flanked on either side by homology arms. Homology arms are complementary to Intron 1 of the human PPP1R12C gene within the AAVS1 locus on chromosome 19 and mediate insertion of Venus into the site between the two homology arms.

Figures 26A-F show targeted genomic insertion of a large protein coding sequence by recombinant AAVF vectors in human cell lines and primary cells demonstrating that AAVF-mediated genome editing is robust and that there is efficient editing in Human CD34+ cells and cell lines. Figure 26A: CD34+ represents primary human CD34+ cytokine-primed peripheral blood stem cells. Figure 26B: K562 is a human CD34+ erythroleukemia cell line. Figure 26C: HepG2 is a human liver cell line. The percent of cells displaying Venus expression, indicative of precise insertion, is shown for Figures 26A-C. Figure 26D shows representative flow profiles showing a distinct Venus expressing population of CD34+ cells after transduction with recombinant AAVF viruses, as compared with untransduced cells. Figure 26E shows editing activity of AAVF7, AAVF12, AAVF15, AAVF17 and AAV9 as compared with AAV6 and AAV8 in a K562 erythroleukemia line. Figure 26F shows editing activity of the same virus in HepG2, a liver cell line. Data shows the percent of cells displaying editing and Venus expression.

Figure 27 shows a targeted integration assay for the detection of large and small inserts into the AAVS1 locus on the human chromosome 19. The schematic maps show the location of primers. The 5' primer is complementary to chromosomal sequences. The 3' primer is specific for the insert. The specific amplicon is predicted to be 1.7kb for the large insert and 1 kb for the

small insert. The split primer pair (chromosomal and insert specific) lends specificity to the targeted integration assay.

Figures 28A-E show that AAVF vectors mediate nucleotide substitution at specified genomic sites. Figure 28A shows maps of single stranded AAV vector genomes for the insertion of a 10 bp insert in intron 1 of the human PPP1R12C gene. This vector encodes a wild type left homology arm (HA-L) which contains an Nhe1 restriction enzyme recognition site (GCTAGC). The NS mut vector, was designed to change the TA sequence in the left homology arm on chromosome 19 to AT. This change results in the conversion of an Nhe1 site to an Sph1 site, changing the sequence from GCTAGC to GCATGC. Figure 28B shows that the left homology arm was amplified using a forward primer located in upstream chromosomal sequences and a reverse primer located in the 10bp insert in Intron 1 of the PPP1R12C gene on chromosome 19. The upper schematic designates the relative sizes of the expected fragments created when genomic DNA from K562 cells is edited using either the wild type or the NS Mut AAVF vectors. Figure 28C is a gel that shows the actual amplicons derived from genomic DNA of K562 cells edited with a wild type AAVF vector. Lanes show the uncut amplicon (Un), the amplicon cut with Nhe 1 (Nhe1) and with Sph 1 (Sph 1). Figure 28D shows gel electrophoresis of K562 DNA after editing with AAVF7 or an AAVF17 vectors encoding either wild type or NS Mut genomes. Figure 28E shows gel electrophoresis of a hepatocellular carcinoma cell line, HepG2 after editing with AAVF7 or an AAVF17 vectors encoding either wild type or NS Mut genomes.

Figure 29 shows a sequence analysis of DNA from cells edited with AAVF7 and AAVF17 Wild type or NS Mut vectors.

Figure 30 is a table that showing AAVF vectors mediate editing in both dividing and non-dividing cells and that AAVF-mediated gene editing does not require DNA synthesis. The figure shows frequency of edited cells expressing Venus in the dividing and non-dividing subsets of primary human CD34+ cells. The percentage of all CD34+ cells that were Venus positive and either BrdU positive or negative was determined by flow cytometry. BrdU positive cells represent dividing cells and BrdU negative cells represent non-dividing cells.

Figures 31A-C show efficient editing of engrafted human hematopoietic stem cells in vivo by systemically delivered AAVF vectors. Figure 31A shows a diagram of the experimental design. Immune deficient NOD/SCID mice were engrafted with human cord blood CD34+ hematopoietic stem cells. Cells were allowed to engraft for 7 weeks prior to intravenous injection of AAVF17-Venus. Hematopoietic cells were harvested from the vertebral and femoral marrow and spleen of xenografted mice 12.5 weeks after AAVF injection. Cells were analyzed by multi-color flow cytometry for Venus expression as well as the presence of human-specific surface markers. Specifically, Venus expression was analyzed in the primitive CD34+ human hematopoietic stem/progenitor cells, CD45+ human differentiated mononuclear hematopoietic cells and glycophorin A+ cells of the erythroid lineage. Figure 31B shows a schematic of the differentiation pathway of the human erythroid lineage, from CD34+ progenitor cells to the glycophorin A+ red blood cells. Figure 31C shows flow cytometric profiles of long term engrafted human cells in the marrow and spleen cells of xenografted mice, 20 weeks after transplantation. Cells were analyzed for both expression of Venus, a marker of editing as well

as specific human cell surface markers.

Figure 32A and B are a summary of *in vivo* data following intravenous injection of AAVF vectors into immune deficient mice xenografted with human cord blood CD34+ hematopoietic stem/progenitor cells. Venus expression reflects targeted insertion of the promoterless Venus cassette into Intron 1 of the human PPP1R12C gene in *in vivo* engrafted human hematopoietic stem cells and their progeny. Figure 32B is a summary of the data in Figure 32A. Figures 32A and B show that editing is long term, that editing is stably inherited (the insert is efficiently expressed in differentiated progeny cells), that *in vivo* editing may be much more efficient than *ex vivo* transduction followed by transplant, and that progeny of edited CD34+ cells retain Venus expression long term.

Figure 33 shows a sequence analysis of targeted chromosomal insertion of a promoterless SA/2A venus ORF in K562 erythroleukemia cell line, primary human cytokine-primed peripheral blood CD34+ cells (PBSC) and HepG2 human liver cell line. Site-specifically integrated sequences were amplified using a chromosome-specific primer and an insert-specific primer. The amplified product was cloned into a TOPO-TA vector and sequenced using Sanger sequencing.

Figure 34 shows a sequence analysis of targeted chromosomal insertion of a 10bp insert in primary human cytokine-primed peripheral blood CD34+ cells and the HepG2 human liver cell line. Site-specifically integrated sequences were amplified using a chromosome-specific primer and an insert-specific primer. The amplified product was cloned into a TOPO-TA vector and sequenced using Sanger sequencing.

Figure 35 shows that AAVF targets integration of small inserts into AAVS1 - CD34+ cells.

Figure 36 shows AAV targeting of promoterless Venus & RFLP in a HepG2 (Hepatoma) cell line.

Figure 37 shows representative femoral bone marrow flow cytometry graphs showing total population sorted, backgating of CD34 and glycoA positive cells onto Venus positive and Venus negative populations, CD34/GlycoA positive cells in Venus negative populations, and glycoA positive and CD34 positive populations.

Figure 38 shows representative spleen flow cytometry graphs showing total population sorted, backgating of CD34 and glycoA positive cells onto Venus positive and Venus negative populations, CD34/GlycoA positive cells in Venus negative populations, and glycoA positive and CD34 positive populations.

Figure 39 shows maps of CBA-mCherry and AAS 1-Venus vector genomes used to determine the relative transduction versus editing efficiencies of AAV vectors.

Figures 40A and 40B show flow cytometric profiles of mCherry (Figure 40A) and Venus (Figure 40B) expression in human CD34+ cord blood cells. Figure 40C shows quantitation of mCherry and Venus expression in CD34+ cells 48 hours after transduction. Figure 40D shows a comparison of the relative expression of Venus to mCherry (Editing Ratio). Bars denote the

ratio of the proportion of cells expressing Venus as a ratio of those expressing mCherry with the corresponding capsid. The black horizontal bar denotes a ratio of 1, which would indicate equal efficiencies of Venus:mCherry expression.

DETAILED DESCRIPTION

[0052] Certain embodiments of the invention are described in detail, using specific examples, sequences, and drawings. One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs.

[0053] The invention relates to adeno-associated virus (AAV) Clade F vectors (e.g., replication-defective AAVs comprising correction genomes enclosed in a Clade F capsid) or AAV vector variants (e.g., replication-defective AAVs comprising capsid variants relative to AAV9 capsids) and related methods thereof, as defined in the claims, that were developed for precise editing of the genome of a cell using homologous recombination without the need for addition of exogenous nucleases. In certain embodiments, genome editing may include, without limitation, introducing insertions, deletions, point mutations or any combination thereof into the genome sequence of a cell (e.g., a target locus of a mammalian chromosome). In certain embodiments, the AAV Clade F vectors and related methods thereof provided herein may be used to insert one or more nucleotide sequences into a specific location of a cell genome without the need for addition of exogenous nucleases prior to integration of the one or more nucleotide sequences. In certain embodiments, the Clade F vectors and related methods thereof provided herein may be used to insert an internucleotide bond into a specific location of a cell genome without the need for addition of exogenous nucleases prior to integration of the internucleotide bond. Also described is a method of treating a disease or disorder in a subject by ex-vivo editing the genome of a cell of the subject via transducing the cell with a Clade F vector or AAV vector variant as described herein and further transplanting the transduced cell into the subject to treat the disease or disorder of the subject. Also provided herein is a replication-defective AAV for use in a method of treating a disease or disorder in a subject by in vivo genome editing by directly administering the Clade F vector in accordance with the claims to the subject. Also described herein are kits for genome editing of a cell comprising one or more of the Clade F vectors or AAV vector variants described herein.

[0054] Homologous recombination using various AAV vectors (e.g., AAV2, AAV6, and AAV8) has been previously reported; however, the reported efficiencies were very low - approximately 1 in a million cells. As shown in Examples 1 and 2 below, AAV Clade F vectors (or AAV vector variants) were used to reproducibly target gene insertion to specified chromosomal locations at significantly greater frequencies than previously seen. For example, targeted genome editing

was achieved by transducing primary cells with AAV Clade F vectors (or AAV vector variants) resulting in the insertion of the transgene into the genome of the primary cells at surprisingly high frequencies, with approximately 10% of the primary cells displaying insertion of the transgene six weeks post-transduction. This frequency is 1,000 to 100,000 fold more efficient than previously reported (see, e.g., Khan, 2011). As shown in Examples 1 and 2 below, high level genome editing was achieved using primary human CD34+ hematopoietic stem cells (K562) and CD34+ primary peripheral blood-derived human hematopoietic stem cells (PBSCs). Targeted gene insertion was observed in both short term (one day) and long term (up to almost six weeks) CD34+ cultures and was verified by transgene expression and sequence analysis. Furthermore, the Clade F vector or AAV vector variant targeted recombination as described herein allows for specific genome engineering with no associated toxicity. As shown in Example 3 below, intravenous injection of AAV vectors pseudotyped with AAVF7 or AAVF17 resulted in transduction of human CD34+ hematopoietic stem and progenitor cells *in vivo*. As shown in Example 4 below, genome editing was achieved both in cell culture and *in vivo* for both small (~10 bps) and large inserts (~800 bps) and was shown by sequencing to be precisely integrated into the target locus. As shown in Example 5 below, genome editing of various human cell lines (e.g., fibroblasts, hepatocellular carcinoma cells, breast cancer cells, retinoblastoma cells, leukemia cells and B cells) was achieved, demonstrating that Clade F vectors could be used to edit genomes in several distinct cell types, such as fibroblasts, liver cells, breast cells, retinal cells, and B cells. As such, this technique has tremendous potential for targeted genome editing in cells *ex vivo* as well as *in vivo* in specific organs.

[0055] Provided herein are Clade F vectors as defined by the claims (e.g., replication-defective AAVs comprising correction genomes enclosed in a Clade F capsid) for editing a genome of a cell and methods thereof as defined in the claims (via recombination, and without the use of exogenous nucleases). In certain embodiments, genome editing includes correction or insertion of one or more mutations in the genome, deletion of one or more nucleotides in the genome, alteration of genomic sequences including regulatory sequences, insertion of one or more nucleotides including transgenes at safe harbor sites or other specific locations in the genome, or any combination thereof as defined in the claims. In certain embodiments, genome editing using the Clade F vectors or and methods thereof as described herein may result in the induction of precise alterations of one or more genomic sequences without inserting exogenous viral sequences or other footprints.

[0056] In some aspects, the disclosure provides an *ex vivo* method for editing a target locus of a mammalian chromosome involving a replication-defective adeno-associated virus (AAV) comprising a correction genome enclosed in a capsid as described herein, e.g., an AAV Clade F capsid. In some embodiments, a "correction genome" is a nucleic acid molecule as defined in the claims, i.e. that contains an editing element as described herein along with additional element(s) (e.g., a 5' inverted terminal repeat (5' ITR) nucleotide sequence, or a fragment thereof, and a 3' inverted terminal repeat (3' ITR) nucleotide sequence, or a fragment thereof) sufficient for encapsidation within a capsid as described herein. It is to be understood that the term "correction genome" does not necessarily require that an editing element contained within the correction genome will "correct" a target locus in a genome, once integrated into the target

locus (e.g., correction of target locus containing a mutation by replacement with a wild-type sequence). Accordingly, in some embodiments, a correction genome may contain an editing element which may comprise a nucleotide sequence that is additive to the target locus (e.g., the target locus is the 3' end of a first open reading frame and the editing element is a second open reading frame that, when integrated into the target locus, will create a gene that encodes a fusion protein).

[0057] In some embodiments, the replication-defective adeno-associated virus (AAV) comprises a correction genome, the correction genome comprising (a) an editing element selected from an internucleotide bond or a nucleotide sequence for integration into a target locus of a mammalian chromosome, (b) a 5' homologous arm nucleotide sequence 5' of the editing element, having homology to a 5' region of the mammalian chromosome relative to the target locus, and (c) a 3' homologous arm nucleotide sequence 3' of the editing element, having homology to a 3' region of the mammalian chromosome relative to the target locus. In some embodiments, the replication-defective AAV comprises a correction genome, the correction genome comprising an editing element nucleotide sequence for integration into a target locus of a mammalian chromosome, the correction genome having an essential absence of a promoter operatively linked to the editing element nucleotide sequence. In some embodiments, the replication-defective AAV comprises a correction genome, the correction genome comprising an editing element selected from an internucleotide bond or a nucleotide sequence for integration into a target locus of a mammalian chromosome in a cell; the AAV having a chromosomal integration efficiency of at least about 1% (e.g., at least about 2%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90%) for integrating the editing element into the target locus of the mammalian chromosome in the cell. In some embodiments, the replication-defective AAV comprises a correction genome, the correction genome comprising an editing element selected from an internucleotide bond or a nucleotide sequence for integration into a target locus of a mammalian chromosome in a cell; the AAV having a chromosomal integration efficiency of at least about 1% (e.g., at least about 2%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90%) in the absence of an exogenous nuclease for integrating the editing element into the target locus of the mammalian chromosome in the cell. In some embodiments of any one of the correction genomes, the correction genome has an essential absence of a promoter operatively linked to the editing element nucleotide sequence. In some embodiments of any one of the correction genomes, the correction genome further comprises an exogenous promoter operatively linked to the editing element. In some embodiments of any one of the replication-defective AAVs, the AAV has a chromosomal integration efficiency of at least about 1%, at least about 2%, at least about 3%, at least about 4%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90% for integrating an editing element into a target locus of a mammalian chromosome in a cell.

[0058] Other aspects of the disclosure relate to a gene editing vector comprising a replication-defective adeno-associated virus (AAV) comprising a correction genome enclosed in an AAV capsid, the correction genome as described herein (e.g., comprising an editing element selected from an internucleotide bond or a nucleotide sequence for integration into a target locus of a mammalian cell chromosome; a 5' homologous arm nucleotide sequence 5' of the editing element having homology to a 5' region of the chromosome relative to the target locus; a 3' homologous arm nucleotide sequence 3' of the editing element having homology to a 3' region of the chromosome relative to the target locus); wherein the AAV has a chromosomal integration efficiency of at least 10% (e.g., at least 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%) for integrating an editing element as described herein into a target locus as described herein. In some embodiments, the chromosomal integration efficiency is at least 10% (e.g., at least 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%) for integrating an editing element as described herein into a target locus as described herein in the absence of an exogenous nuclease.

[0059] A correction genome as described herein can comprise a 5' inverted terminal repeat (5' ITR) nucleotide sequence 5' of the 5' homologous arm nucleotide sequence, and a 3' inverted terminal repeat (3' ITR) nucleotide sequence 3' of the 3' homologous arm nucleotide sequence. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially identical (e.g., at least 90%, at least 95%, at least 98%, at least 99% identical or 100% identical) to an AAV2 virus 5' ITR and an AAV2 virus 3' ITR, respectively. In some embodiments, the 5' ITR nucleotide sequence has at least 95% (e.g., at least 96%, at least 97%, at least 98%, at least 99%, or 100%) sequence identity to SEQ ID NO:36, and the 3' ITR nucleotide sequence has at least 95% (e.g., at least 96%, at least 97%, at least 98%, at least 99%, or 100%) sequence identity to SEQ ID NO:37. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially identical (e.g., at least 90%, at least 95%, at least 98%, at least 99% identical or 100% identical) to an AAV5 virus 5' ITR and an AAV5 virus 3' ITR, respectively. In some embodiments, the 5' ITR nucleotide sequence has at least 95% (e.g., at least 96%, at least 97%, at least 98%, at least 99%, or 100%) sequence identity to SEQ ID NO:38, and the 3' ITR nucleotide sequence has at least 95% (e.g., at least 96%, at least 97%, at least 98%, at least 99%, or 100%) sequence identity to SEQ ID NO:39. In some embodiments, the 5' ITR nucleotide sequence and the 3' ITR nucleotide sequence are substantially mirror images of each other (e.g., are mirror images of each other except for at 1, 2, 3, 4 or 5 nucleotide positions in the 5' or 3' ITR).

[0060] Exemplary AAV2 5' ITR (SEQ ID NO: 36) -

ttggccactccctctctgcgcgctcgtcgtcactgaggccgggcgaccaaaggctgccccgacgcccgggctttgcccgggcgggc
ctcagtgagcgagcgagcgcgagagagggagtggccaactccatcactaggggttcct

[0061] Exemplary AAV2 3' ITR (SEQ ID NO: 37) -

aggaaccctagtgatggagtggccactccctctctgcgcgctcgtcgtcactgaggccgggcgaccaaaggctgccccgacgc
ccgggctttgcccgggcggcctcagtgagcgagcgagcgcgagagagggagtggccaa

[0062] Exemplary AAV5 5' ITR (SEQ ID NO: 38) -

ctctccccctgtcgcgttcgctcgctcgctggctcgtttgggggggtggcagctcaaagagctgccagacgacggccctctggccgt
cgcccccccaaacgagccagcgagcgagcgagcgaacgcgacaggggggagagtcccacactctcaagcaaggggggtttgta

[0063] Exemplary AAV5 3' ITR (SEQ ID NO: 39) -

tacaaaacctccttgcttgagagtgtggcactctccccctgtcgcgttcgctcgctcgctggctcgtttgggggggtggcagctcaa
gagctgccagacgacggccctctggccgtcgcccccccaaacgagccagcgagcgagcgaacgcgacaggggggagag

[0064] In some embodiments, a correction genome as described herein is no more than 7kb (kilobases), no more than 6kb, no more than 5kb, or no more than 4kb in size. In some embodiments, a correction genome as described herein is between 4kb and 7kb, 4kb and 6kb, 4kb and 5kb, or 4.1kb and 4.9kb.

[0065] In certain embodiments, AAV Clade F vectors for editing a genome of a cell comprise one or more Clade F capsids as defined in the claims. In certain embodiments, a donor vector may be packaged into the Clade F capsids described herein according to a standard AAV packaging method resulting in formation of the AAV Clade F vector (see e.g., Chatterjee, 1992). In certain embodiments, the one or more Clade F capsids as defined in the claims may influence the tropism of the AAV Clade F vector for a particular cell.

[0066] According to certain embodiments, the one or more Clade F capsids are derived from human stem cell-derived AAV. It has been previously shown that that cytokine-primed peripheral blood CD34⁺ stem cells from healthy donors harbor endogenous natural AAV sequences in their genome (see, e.g., US Patent Publication Number US20130096182A1 and US20110294218A1). The efficacy of the AAV isolate variants (variant relative to AAV9) has been previously demonstrated, including the efficacy of individual capsid nucleotides and proteins for use in cell transduction (see, e.g., US Patent Publication Number US20130096182A1 and US20110294218A1).

[0067] Full length AAV capsid variant genes (variant relative to AAV9) from the donors harboring endogenous natural AAV sequences in their genome were isolated and sequenced. The polynucleotide and polypeptide sequences of the capsid variants are provided in Figure 1 and in U.S. Patent Publication Number US20130096182A1, and U.S. Patent Publication No. US20110294218A1. The AAV Clade F vectors or AAV vector variants described herein may comprise one or more Clade F capsids or capsid variants comprising a polynucleotide sequence selected from the group of AAVF1 (SEQ ID NO: 20), AAVF2 (SEQ ID NO: 21), AAVF3 (SEQ ID NO: 22), AAVF4 (SEQ ID NO: 23), AAVF5 (SEQ ID NO: 25), AAVF11 (SEQ ID NO: 26), AAVF7 (SEQ ID NO: 27), AAVF8 (SEQ ID NO: 28), AAVF9 (SEQ ID NO: 29), AAVF12 (SEQ ID NO: 30), AAVF13 (SEQ ID NO: 31), AAVF14 (SEQ ID NO: 32), AAVF15 (SEQ ID NO: 33), AAVF16 (SEQ ID NO: 34), AAVF17 (SEQ ID NO: 35), variants, fragments, mutants, and any combination thereof. The AAV Clade F vectors or AAV vector variants may comprise one

or more Clade F capsids or capsid variants comprising a polypeptide sequence selected from the group of AAVF1 (SEQ ID NO: 2), AAVF2 (SEQ ID NO: 3), AAVF11 (SEQ ID NO: 4), AAVF3 (SEQ ID NO: 5), AAVF4 (SEQ ID NO: 6), AAVF6 (SEQ ID NO: 7), AAVF7 (SEQ ID NO: 8), AAVF8 (SEQ ID NO: 9), AAVF9 (SEQ ID NO: 10), AAVF5 (SEQ ID NO: 11), AAVF12 (SEQ ID NO: 12), AAVF17 (SEQ ID NO: 13), AAVF13 (SEQ ID NO: 14), AAVF14 (SEQ ID NO: 15), AAVF15 (SEQ ID NO: 16), AAVF16 (SEQ ID NO: 17), variants, fragments, mutants, and any combination thereof (see, e.g., Figure 1).

[0068] The polynucleotide or polypeptide sequences of the Clade F capsids or capsid variants described herein may have at least about 95%, 96%, 97%, more preferably about 98%, and most preferably about 99% sequence identity to the sequences taught in the present specification. Percentage identity may be calculated using any of a number of sequence comparison programs or methods such as the Pearson & Lipman, Proc. Natl. Acad. Sci. USA, 85:2444 (1988), and programs implementing comparison algorithms such as GAP, BESTFIT, FASTA, or TFASTA (from the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), or BLAST, available through the National Center for Biotechnology Information web site.

[0069] The Clade F capsids or capsid variant sequences may be modified at one or more positions in the V1 and/or V3 cap genes, these genes or functional portions of the genes can be used separately or together in any of the AAV Clade F vectors or AAV vector variants and methods described herein. Cap genes, V1, V2, and V3, may be substituted out from multiple mutated sequences, and are typically used in a colinear fashion V1-V2-V3. However the sequences may be truncated such as partial V1-V2-V3 or V1-V3 or V1-V1-V2-V3. For example, one sequence could be V1 of (AAVF8)-V2 of (AAVF4)-V3 of AAVF14. Preferably, the Clade F capsids or capsid variants transduce the target cells on a level at or higher than AAV2.

[0070] Also described are capsid variants comprising a combination of one or more V1, V2, and V3 polynucleotide sequences of capsid variants (e.g., SEQ ID NOs: 20-35, variant relative to AAV9 capsid), the AAV9 capsid (SEQ ID NO: 18), the AAV2 capsid (SEQ ID NO: 19), variants, fragments, or mutants thereof. Also described are capsid variants comprising a combination of one or more V1, V2, and V3 polynucleotide sequences of capsid variants (SEQ ID NOs: 20-35, variant relative to AAV9 capsid), any other known AAV capsids, variants, fragments, or mutants thereof.

[0071] In certain embodiments, the one or more capsids used in the claimed methods may comprise a combination of V1, V2, and/or V3 polypeptide sequences of the Clade F capsids (SEQ ID NOs: 2-17, variant relative to AAV9).

[0072] In some embodiments, an AAV Clade F vector for editing a genome of a cell comprises a AAV Clade F capsid in accordance with the claims. An "AAV Clade F capsid" described herein may refer to a capsid that has an AAV VP1, VP2, and/or VP3 sequence that has at least 86% (e.g., at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at

least 99%) sequence identity with the AAV VP1, VP2, and/or VP3 sequence of AAV9, respectively. Exemplary Clade F capsids include AAVF1-17 (also referred to herein as AAVHSC1-17), AAV9, AAVHU31, AAVHU32, and AAVAnc110 (see, e.g., Zinn et al. In Silico Reconstruction of the Viral Evolutionary Lineage Yields a Potent Gene Therapy Vector (2015) Cell Reports, Vol 12, pp. 1056-1068).

[0073] In some embodiments, an AAV Clade F capsid comprises at least one or at least two proteins selected from Clade F VP1, Clade F VP2 and Clade F VP3 as defined by the claims. In some embodiments, an AAV Clade F capsid comprises Clade F VP1, Clade F VP2 and Clade F VP3 proteins as defined by the claims.

[0074] Exemplary AAV VP1, VP2, and VP3 protein sequences of AAV Clade F capsids are provided in the below table.

AAV capsid	VP1	VP2	VP3
AAV9	Amino acids 1 to 736 of SEQ ID NO: 1	Amino acids 138 to 736 of SEQ ID NO: 1	Amino acids 203 to 736 of SEQ ID NO: 1
AAVF1	Amino acids 1 to 736 of SEQ ID NO: 2	Amino acids 138 to 736 of SEQ ID NO: 2	Amino acids 203 to 736 of SEQ ID NO: 2
AAVF2	Amino acids 1 to 736 of SEQ ID NO: 3	Amino acids 138 to 736 of SEQ ID NO: 3	Amino acids 203 to 736 of SEQ ID NO: 3
AAVF3	Amino acids 1 to 736 of SEQ ID NO: 5	Amino acids 138 to 736 of SEQ ID NO: 5	Amino acids 203 to 736 of SEQ ID NO: 5
AAVF4	Amino acids 1 to 736 of SEQ ID NO: 6	Amino acids 138 to 736 of SEQ ID NO: 6	Amino acids 203 to 736 of SEQ ID NO: 6
AAVF5	Amino acids 1 to 736 of SEQ ID NO: 11	Amino acids 138 to 736 of SEQ ID NO: 11	Amino acids 203 to 736 of SEQ ID NO: 11
AAVF6	Amino acids 1 to 736 of SEQ ID NO: 7	Amino acids 138 to 736 of SEQ ID NO: 7	Amino acids 203 to 736 of SEQ ID NO: 7
AAVF7	Amino acids 1 to 736 of SEQ ID NO: 8	Amino acids 138 to 736 of SEQ ID NO: 8	Amino acids 203 to 736 of SEQ ID NO: 8
AAVF8	Amino acids 1 to 736 of SEQ ID NO: 9	Amino acids 138 to 736 of SEQ ID NO: 9	Amino acids 203 to 736 of SEQ ID NO: 9
AAVF9	Amino acids 1 to 736 of SEQ ID NO: 10	Amino acids 138 to 736 of SEQ ID NO: 10	Amino acids 203 to 736 of SEQ ID NO: 10
AAVF10	Amino acids 1 to 736 of SEQ ID NO: 3	Amino acids 138 to 736 of SEQ ID NO: 3	Amino acids 203 to 736 of SEQ ID NO: 3
AAVF11	Amino acids 1 to 736 of SEQ ID NO: 4	Amino acids 138 to 736 of SEQ ID NO: 4	Amino acids 203 to 736 of SEQ ID NO: 4
AAVF12	Amino acids 1 to 736 of SEQ ID NO: 12	Amino acids 138 to 736 of SEQ ID NO: 12	Amino acids 203 to 736 of SEQ ID NO: 12
AAVF13	Amino acids 1 to 736 of SEQ ID NO: 14	Amino acids 138 to 736 of SEQ ID NO: 14	Amino acids 203 to 736 of SEQ ID NO: 14

AAV capsid	VP1	VP2	VP3
AAVF14	Amino acids 1 to 736 of SEQ ID NO: 15	Amino acids 138 to 736 of SEQ ID NO: 15	Amino acids 203 to 736 of SEQ ID NO: 15
AAVF15	Amino acids 1 to 736 of SEQ ID NO: 16	Amino acids 138 to 736 of SEQ ID NO: 16	Amino acids 203 to 736 of SEQ ID NO: 16
AAVF16	Amino acids 1 to 736 of SEQ ID NO: 17	Amino acids 138 to 736 of SEQ ID NO: 17	Amino acids 203 to 736 of SEQ ID NO: 17
AAVF17	Amino acids 1 to 736 of SEQ ID NO: 13	Amino acids 138 to 736 of SEQ ID NO: 13	Amino acids 203 to 736 of SEQ ID NO: 13

[0075] An AAV Clade F capsid described herein may comprise a VP1, VP2, or VP3 protein that has at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736, amino acids 138 to 736 or amino acids 203 to 736 of SEQ ID NO:1, respectively, which correspond to the amino acid sequences of AAV9 capsid proteins VP1, VP2 and VP3, respectively. An AAV Clade F capsid described herein may comprise VP1 and VP2 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736 and amino acids 138 to 736 of SEQ ID NO:1, respectively, which correspond to the amino acid sequences of AAV9 capsid proteins VP1 and VP2, respectively; VP1 and VP3 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736 and amino acids 203 to 736 of SEQ ID NO:1, respectively, which correspond to the amino acid sequences of AAV9 capsid proteins VP1 and VP3, respectively; or VP2 and VP3 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 138 to 736 and amino acids 203 to 736 of SEQ ID NO:1, respectively, which correspond to the amino acid sequences of AAV9 capsid proteins VP2 and VP3, respectively. An AAV Clade F capsid described herein may comprise VP1, VP2, and VP3 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 of SEQ ID NO:1, respectively, which correspond to the amino acid sequences of AAV9 capsid proteins VP1, VP2 and VP3, respectively.

[0076] In some aspects, an AAV Clade F capsid comprises a VP1, VP2, or VP3 protein that has at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736, amino acids 138 to 736 or amino acids 203 to 736 of any one of SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively, which correspond to the amino acid sequences of AAVF1 through AAVF9 and AAVF11 through AAVF17 capsid

proteins VP1, VP2 and VP3, respectively. In some aspects, an AAV Clade F capsid comprises VP1 and VP2 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736 and amino acids 138 to 736 of any one of SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively, which correspond to the amino acid sequences of AAVF1 through AAVF9 and AAVF11 through AAVF17 capsid proteins VP1 and VP2, respectively; VP1 and VP3 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736 and amino acids 203 to 736 of any one of SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively, which correspond to the amino acid sequences of AAVF1 through AAVF9 and AAVF11 through AAVF17 capsid proteins VP1 and VP3, respectively; or VP2 and VP3 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 138 to 736 and amino acids 203 to 736 of any one of SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively, which correspond to the amino acid sequences of AAVF1 through AAVF9 and AAVF11 through AAVF17 capsid proteins VP2 and VP3, respectively. In some aspects, an AAV Clade F capsid comprises VP1, VP2, and VP3 proteins that have at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) amino acid sequence identity to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 of any one of SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively, which correspond to the amino acid sequences of AAVF1 through AAVF9 and AAVF11 through AAVF17 capsid proteins VP1, VP2 and VP3, respectively.

[0077] In some aspects, an AAV Clade F capsid comprises a VP1, VP2, or VP3 protein that is encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to SEQ ID NO:18, which corresponds to the nucleotide sequence encoding AAV9 capsid proteins VP1, VP2 and VP3, respectively. In some aspects, an AAV Clade F capsid comprises VP1 and VP2 proteins that are encoded by nucleotide sequences comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to SEQ ID NO: 18, which corresponds to the nucleotide sequence encoding AAV9 capsid proteins VP1, VP2, and VP3; VP1 and VP3 proteins that are encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to SEQ ID NO: 18; or VP2 and VP3 proteins that are encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to SEQ ID NO: 18. In some aspects, an AAV Clade F capsid comprises VP1, VP2, and VP3 proteins that are encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to SEQ

ID NO: 18, which corresponds to a nucleotide sequence encoding AAV9 capsid proteins VP1, VP2 and VP3.

[0078] In some aspects, an AAV Clade F capsid comprises a VP1, VP2, or VP3 protein that is encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to any one of SEQ ID NOs: 20, 21, 22, 23, 25, 24, 27, 28, 29, 26, 30, 31, 32, 33, 34 or 35, which correspond to nucleotide sequences encoding AAVF1 through AAVF17 capsid proteins VP1, VP2 and VP3, respectively. In some aspects, an AAV Clade F capsid comprises VP1 and VP2 proteins that are encoded by nucleotide sequences comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to any one of SEQ ID NOs:20-35; VP1 and VP3 proteins that are encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to any one of SEQ ID NOs:20-35; or VP2 and VP3 proteins that are encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to any one of SEQ ID NOs:20-35. In some aspects, an AAV Clade F capsid comprises VP1, VP2, and VP3 proteins that are encoded by a nucleotide sequence comprising at least 85% (e.g., at least 86%, at least 88%, at least 90%, at least 92%, at least 94%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) nucleotide sequence identity to any one of SEQ ID NOs: 20, 21, 22, 23, 25, 24, 27, 28, 29, 26, 30, 31, 32, 33, 34 or 35, which correspond to nucleotide sequences encoding AAVF1 through AAVF17 capsid proteins VP1, VP2 and VP3, respectively.

[0079] An AAV Clade F capsid described herein may comprise an AAV9 VP1, VP2, or VP3 capsid protein, which corresponds to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NO:1, respectively. An AAV Clade F capsid described herein may comprise AAV9 VP1 and VP2 capsid proteins, which correspond to amino acids 1 to 736 and amino acids 138 to 736 as set forth in SEQ ID NO:1, respectively; AAV9 VP1 and VP3 capsid proteins, which correspond to amino acids 1 to 736 and amino acids 203 to 736 as set forth in SEQ ID NO:1, respectively; or AAV9 VP2 and VP3 capsid proteins, which correspond to amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NO:1, respectively. An AAV Clade F capsid described herein may comprise AAV9 VP1, VP2 and VP3 capsid proteins, which correspond to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NO:1, respectively.

[0080] In some embodiments, an AAV Clade F capsid comprises a VP1 capsid protein selected from a VP1 capsid protein of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which corresponds to amino acids 1 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively. In some embodiments, an AAV Clade F capsid comprises a VP1 and a VP2 capsid protein independently selected from a VP1 and VP2 capsid protein of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which correspond

to amino acids 1 to 736 and amino acids 138 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively. In some embodiments, an AAV Clade F capsid comprises a VP2 and a VP3 capsid protein independently selected from a VP2 and VP3 capsid protein of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which correspond to amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively. In some embodiments, an AAV Clade F capsid comprises each of the VP1, VP2 and VP3 capsid proteins of any one of AAVF1 through AAVF9 and AAVF11 through AAVF17, which correspond to amino acids 1 to 736, amino acids 138 to 736 and amino acids 203 to 736 as set forth in SEQ ID NOs: 2, 3, 5, 6, 11, 7, 8, 9, 10, 4, 12, 14, 15, 16, 17 or 13, respectively.

[0081] As used herein, a fragment of a polynucleotide sequence may be a portion of the polynucleotide that encodes a polypeptide which provides substantially the same function as the polypeptide encoded by the full length polynucleotide sequence. As used herein, mutants of a polynucleotide sequence may be obtained by deletion, substitution, addition, and/or insertion of one or more nucleotides to the specific polynucleotide sequence. It should be understood that such fragments, and/or mutants of a polynucleotide sequence encode a polypeptide having substantially the same function as the polypeptide encoded by the full length polynucleotide sequence.

[0082] As used herein, a polypeptide sequence may include fragments, and/or mutants of the polypeptide sequence, while still providing substantially the same function as the full length polypeptide sequence. A fragment of a polypeptide sequence means a part of the polypeptide sequence that provides substantially the same function as the full length polypeptide sequence. Examples of mutants of a polypeptide sequence include deletions, substitutions, additions, and/or insertions of one or more amino acids to the polypeptide sequence.

[0083] In certain aspects, a polynucleotide sequence may be a recombinant or non-naturally occurring polynucleotide. In certain aspects, a polynucleotide sequence may be cDNA.

[0084] The AAV Clade F vectors or AAV vector variants described herein may comprise any of the AAVF (or AAVHSC) or any other AAV Clade F vectors described herein. An AAV Clade F vector or AAV vector variant described herein may comprise any of the AAVF (or AAVHSC) vectors described in herein, such as AAVF1, AAVF2, AAVF3, AAVF4, AAVF5, AAVF6, AAVF7, AAVF8, AAVF9, AAVF10, AAVF11, AAVF12, AAVF13, AAVF14, AAVF15, AAVF16, AAVF17, variants, fragments, mutants, or any combination thereof. A Clade F vector or AAV vector variant described herein may comprise any of AAV9, AAVF1, AAVF2, AAVF3, AAVF4, AAVF5, AAVF6, AAVF7, AAVF8, AAVF9, AAVF10, AAVF11, AAVF12, AAVF13, AAVF14, AAVF15, AAVF16, AAVF17, AAVHU31, AAVHU32, variants, fragments, mutants, or any combination thereof.

[0085] In certain embodiments, the AAV Clade F vectors provided herein for use in the claimed methods comprise an editing element (also referred to herein as a targeting cassette, meaning the terms "editing element" and "targeting cassette" are used interchangeably herein) as

defined by the claims comprising one or more nucleotide sequences or an internucleotide bond to be integrated into a target locus (also referred to herein as a target site, meaning the terms are used interchangeably herein) of the genome, a 5' homologous arm nucleotide sequence 5' of the editing element, having homology to a 5' region of the mammalian chromosome relative to the target locus (e.g., a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site)), and a 3' homologous arm nucleotide sequence 3' of the editing element, having homology to a 3' region of the mammalian chromosome relative to the target locus (e.g., a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site)).

[0086] In certain embodiments, the one or more nucleotide sequences to be integrated into a target site of the genome may be one or more therapeutic nucleotide sequences. The term "therapeutic" as used herein refers to a substance or process that results in the treatment of a disease or disorder. "Therapeutic nucleotide sequence" is a nucleotide sequence that provides a therapeutic effect. The therapeutic effect can be direct (e.g., substitution of a nucleic acid of a gene expressed as a protein, or insertion of a cDNA into an intron for expression) or indirect (e.g., correction of a regulatory element such as a promoter). In certain embodiments, the therapeutic nucleotide sequence may include one or more nucleotides. In certain embodiments, the therapeutic nucleotide sequence may be a gene, variant, fragment, or mutant thereof. In certain embodiments, when gene therapy is desired, the therapeutic nucleotide sequence may be any nucleotide sequence that encodes a protein that is therapeutically effective, including therapeutic antibodies. The Clade F vectors comprising the therapeutic nucleotide sequences are preferably administered in a therapeutically effective amount via a suitable route of administration, such as injection, inhalation, absorption, ingestion or other methods.

[0087] In some embodiments, an editing element as described herein consists of one nucleotide. In some embodiments, an editing element as described herein consists of one nucleotide and a target locus as described herein is a nucleotide sequence consisting of one nucleotide, the target locus representing a point mutation. In some embodiments, an editing element as described herein comprises at least 1, 2, 10, 100, 200, 500, 1000, 1500, 2000, 3000, 4000, or 5000 nucleotides. In some embodiments, an editing element as described herein comprises or consists of 1 to 5500, 1 to 5000, 1 to 4500, 1 to 4000, 1 to 3000, 1 to 2000, 1 to 1000, 1 to 500, 1 to 200, or 1 to 100 nucleotides, or 2 to 5500, 2 to 5000, 2 to 4500, 2 to 4000, 2 to 3000, 2 to 2000, 2 to 1000, 2 to 500, 2 to 200, or 2 to 100 nucleotides, or 10 to 5500, 10 to 5000, 10 to 4500, 10 to 4000, 10 to 3000, 10 to 2000, 10 to 1000, 10 to 500, 10 to 200, or 10 to 100 nucleotides. In some embodiments, an editing element as described herein comprises or consists of an exon, an intron, a 5' untranslated region (UTR), a 3' UTR, a promoter, a splice donor, a splice acceptor, a sequence encoding or non-coding RNA, an insulator, a gene, or a combination thereof. In some embodiments, an editing element as described herein is a fragment (e.g., no more than 2kb, no more than 1kb, no more than 500bp, no more than 250bp, no more than 100bp, no more than 50bp, or no more than 25bp) of a coding sequence of a gene within or spanning a target locus as described herein. In some

embodiments, an editing element as described herein is an internucleotide bond (e.g., a phosphodiester bond connecting two adjacent nucleotides). In some embodiments, an editing element as described herein is an internucleotide bond, a target locus in a chromosome as described herein is a nucleotide sequence comprising one or more nucleotides, and the editing element comprises a deletion for the target locus in the chromosome.

[0088] In certain embodiments, the editing element (or targeting cassette) of the AAV Clade F vector may comprise one or more regulatory element polynucleotide sequences. For example, in certain embodiments, the one or more regulatory element polynucleotide sequences may be selected from a 2A sequence, splice acceptor sequence, polyadenylation sequence, and any combination thereof. In certain embodiments, the targeting cassette may comprise one or more AAV inverted terminal repeat (ITR) polynucleotide sequences flanking the 5' and 3' homologous arm polynucleotide sequences. In certain embodiments, the editing element (or targeting cassette) does not contain a promoter to drive expression of the one or more nucleotide sequences. In certain embodiments, if the editing element (or targeting cassette) does not contain a promoter, the expression of the one or more nucleotide sequences after integration into the cell genome may be controlled by one or more regulatory elements of the cell. In certain embodiments, expression of the promoterless one or more nucleotide sequences demonstrates that the one or more nucleotide sequences was correctly integrated into the cell.

[0089] The AAV Clade F vector comprises a 5' homologous arm nucleotide sequences and a 3' homologous arm nucleotide sequence as defined in the claims. The homologous arm polynucleotide sequences are homologous to a region of the target locus (target site) of the genome as defined in the claims. The homologous arm polynucleotide sequences includes a 5' homologous arm polynucleotide sequence as defined in the claims. The 5' homologous arm polynucleotide sequence flanks the 5' end of the editing element (or targeting cassette) as defined in the claims. The 5' homologous arm polynucleotide sequence flanking the editing element (or targeting cassette) is homologous to a region that is upstream of a target locus (target site) of the genome as defined in the claims. The homologous arm polynucleotide sequences includes a 3' homologous arm polynucleotide sequence as defined in the claims. The 3' homologous arm polynucleotide sequence flanks the 3' end of the editing element (or targeting cassette) as defined in the claims. The 3' homologous arm polynucleotide sequence flanking the editing element (or targeting cassette) is homologous to a region that is downstream of the target locus (target site) of the genome as defined in the claims. In certain embodiments, the homologous arm polynucleotide sequences may be approximately 500 to 1,000 nucleotides long. For example, in certain embodiments, the homologous arm polynucleotide sequences may be approximately 800 nucleotides long. In certain embodiments, the homologous arm polynucleotide sequence may be up to approximately 3,000 nucleotides long. In some embodiments, each of the 5' and 3' homologous arm nucleotide sequences independently has a nucleotide length of between about 50 to 2000 nucleotides, such as between about 500-1000, about 600-1000, or about 700-900 nucleotides. In some embodiments, each of the 5' and 3' homologous arm nucleotide sequences independently has a nucleotide length of between about 600, about 800, or about 1000

nucleotides.

[0090] In some embodiments, the 5' and 3' homologous arm nucleotide sequences have substantially equal nucleotide lengths. In some embodiments, the 5' and 3' homologous arm nucleotide sequences have asymmetrical nucleotide lengths. In some embodiments, the asymmetry in nucleotide length is defined by a difference between the 5' and 3' homologous arm nucleotide sequence lengths of up to 50% in the length, such as up to 40%, 30%, 20%, or 10% difference in the length. In some embodiments, the asymmetry in nucleotide length is defined by one arm of the 5' and 3' homologous arm having a length of about 600 nucleotides and the other arm of the 5' and 3' homologous arm having a length of about 800 or about 900 nucleotides.

[0091] In some embodiments, the 5' homologous arm nucleotide sequence has at least about 90% (e.g., at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or at least about 99.5%) nucleotide sequence identity to the 5' region of the mammalian chromosome relative to the target locus. In some embodiments, the 3' homologous arm nucleotide sequence has at least about 90% (e.g., at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or at least about 99.5%) nucleotide sequence identity to the 3' region of the mammalian chromosome relative to the target locus. In some embodiments, differences in nucleotide sequences of the 5' homologous arm or the 3' homologous arm and the corresponding 5' region or 3' region of the mammalian chromosome, respectively, can comprise, consist essentially of or consist of non-coding differences in nucleotide sequences. In some embodiments, differences in nucleotide sequences of the 5' homologous arm or the 3' homologous arm and the corresponding 5' region or 3' region of the mammalian chromosome, respectively, can comprise, consist essentially of or consist of differences in nucleotide sequences that result in conservative amino acid changes (e.g., a basic amino acid changed to a different basic amino acid). In some embodiments, the 5' homologous arm nucleotide sequence has 100% sequence identity to the 5' region of the mammalian chromosome relative to the target locus and the 3' homologous arm nucleotide sequence has 100% sequence identity to the 3' region of the mammalian chromosome relative to the target locus. In some embodiments, the 5' homologous arm nucleotide sequence and the 3' homologous arm nucleotide sequence are considered homologous with the 5' region and 3' of the mammalian chromosome relative to the target locus, respectively, even if the target locus contains one or more mutations, such as one or more naturally occurring SNPs, compared to the 5' or 3' homologous arm.

[0092] In certain embodiments, the target locus (target site) of the cell genome may be any region of the genome where it is desired that the editing of the cell genome occur. For example, the target locus (target site) of the cell genome may comprise a locus of a chromosome in the cell (e.g., a region of a mammalian chromosome). In certain embodiments, the locus of the chromosome may be a safe harbor site. A safe harbor site is a location in the genome where a nucleotide sequence may integrate and function in a predictable manner without perturbing endogenous gene activity. In certain embodiments, the safe harbor site may

be the AAVS1 locus in human chromosome 19 (also known as PPP1R12C locus). In certain embodiments, the safe harbor site may be the first intron of PPP1R12C in the AAVS1 locus in human chromosome 19. The AAVS1 locus on chromosome 19 qter13.3-13.4 was previously shown to be a "safe harbor" site for the insertion of transgenes since genes inserted here are expressed with no pathogenic consequences, which is similar to wild-type AAV that integrates at this locus with no pathogenic consequences (Giraud 1994; Linden, 1996A; Linden 1996B). In some embodiments, the target locus (target site) is a locus associated with a disease state as described herein.

[0093] In certain embodiments, the target locus is a mutant target locus in a mammalian chromosome comprising one or more mutant nucleotides, relative to a corresponding wild type mammalian chromosome. In some embodiments, the mutant target locus comprises a point mutation, a missense mutation, a nonsense mutation, an insertion of one or more nucleotides, a deletion of one or more nucleotides, or combinations thereof. In some embodiments, the mutant target locus comprises an amorphic mutation, a neomorphic mutation, or an antimorphic mutation. In some embodiments, the mutant target locus comprises an autosomal dominant mutation, an autosomal recessive mutation, a heterozygous mutation, a homozygous mutation, or combinations thereof. In some embodiments of any one of the mutant target loci described herein, the mutant target locus is selected from a promoter, an enhancer, a signal sequence, an intron, an exon, a splice donor site, a splice acceptor site, an internal ribosome entry site, an inverted exon, an insulator, a gene, a chromosomal inversion, and a chromosomal translocation within the mammalian chromosome.

[0094] In some embodiments, a target locus in a chromosome as described herein is a nucleotide sequence comprising n nucleotides where n is an integer greater than or equal to one (e.g., 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, 100, 500, 1000, 2000, 3000, 4000, 5000, or any integer therebetween), an editing element as described herein comprises m nucleotides where m is an integer equal to n , and the editing element represents a substitution for the target locus of the chromosome. In some embodiments, a target locus in a chromosome as described herein is a nucleotide sequence comprising n nucleotides where n is an integer greater than or equal to one (e.g., 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, 100, 500, 1000, 2000, 3000, 4000, 5000, or any integer therebetween), an editing element as described herein comprises m nucleotides where m is an integer greater than n , and the editing element represents a substitutive addition for the target locus of the chromosome. In some embodiments, a target locus in a chromosome as described herein is a nucleotide sequence comprising n nucleotides where n is an integer greater than or equal to two (e.g., 2, 3, 4, 5, 10, 20, 30, 40, 50, 100, 500, 1000, 2000, 3000, 4000, 5000, or any integer therebetween), an editing element as described herein comprises m nucleotides where m is an integer less than n ; and the editing element represents a substitutive deletion for the target locus of the chromosome. In some embodiments, a target locus in a chromosome as described herein is an internucleotide bond, an editing element as described herein comprises m nucleotides where m is an integer greater than or equal to one (e.g., 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, 100, 500, 1000, 2000, 3000, 4000, 5000, or any integer therebetween); and the editing element represents an addition for the target locus of the chromosome.

[0095] In accordance with the claims, a target locus in a chromosome is a target locus in a mammalian chromosome (e.g. a human, mouse, bovine, equine, canine, feline, rat, or rabbit chromosome). In some embodiments, the target locus can comprise an intron of a mammalian chromosome. In some embodiments, the target locus can comprise an exon of a mammalian chromosome. In some embodiments, the target locus can comprise a non-coding region of a mammalian chromosome. In some embodiments, the target locus can comprise a regulatory region of a mammalian chromosome. In some embodiments, the mammalian chromosome is selected from human chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, X and Y. In some embodiments, the mammalian chromosome is selected from mouse chromosome 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, X and Y. In some embodiments, the mammalian chromosome is not human chromosome 19. In some embodiments, the mammalian chromosome is a somatic cell chromosome. Exemplary somatic cells are further described herein.

[0096] In certain embodiments, the one or more nucleotide sequences or editing element may be integrated into the genome through homologous recombination without the need for DNA cleavage prior to integration. In certain embodiments, the one or more nucleotide sequences or editing element may be integrated into the genome through homologous recombination without the need for the addition of exogenous nucleases such as a zinc finger nuclease (ZFN), a transcription activator-like effector nuclease (TALEN), or an RNA guided nuclease (CRISPR/Cas).

[0097] The cell that is edited by the AAV Clade F vectors or AAV vector variants described herein may be any type of mammalian cell. In certain embodiments, cells are the ones defined by the claims. Contemplated herein is a wide variety of mammalian cells, for example, cells of the liver, lung, cartilage and other connective tissue, eye, central and peripheral nervous system, lymphatic system, bone, muscle, blood, brain, skin, heart, and digestive tract. When the cell to be edited by the AAV Clade F vectors or AAV vector variants is, for example, a liver cell, the inserted nucleotide sequence is directed to treating (improving or curing a disorder or stopping further progression of a disease or disorder) or preventing a condition. When the cell to be edited by the AAV Clade F vectors or AAV vector variants is a liver cell, the liver conditions treated or prevented comprise hemophilia, enzyme delivery, cirrhosis, cancer, or atherosclerosis, among other liver conditions. In certain embodiments, the cell may be a somatic cell (e.g., a mammalian somatic cell). In certain embodiments, the cell (e.g., somatic cell such as mammalian somatic cell) may be from a tissue selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, and epithelial tissue. In certain embodiments, the cell (e.g., somatic cell such as mammalian somatic cell) may be from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage. In some embodiments, the cell is a CD34+ cell (e.g., a CD34+ somatic cell). In some embodiments, the cell (e.g., somatic cell such as mammalian somatic cell) is a liver cell, a fibroblast, a breast cell, a lymphocyte, or a retinal cell.

[0098] As shown herein, AAV packaged with the Clade F capsids or capsid variants described

herein demonstrate specific tropism for certain target tissues, such as blood stem cells, liver, heart, eye, breast, and joint tissue, and may be used to transduce stem cells for introduction of genes of interest into the target tissues. Certain of the vectors are able to cross tightly controlled biological junctions, such as the blood-brain barrier, which open up additional novel uses and target organs for the vectors, providing for methods of gene therapy through genome editing. Thus, Clade F vectors or AAV vector variants may demonstrate a tropism for a particular cell based on their Clade F capsids or capsid variants. For example a) for muscle tissue or cells, the AAV Clade F vector or AAV vector variant may be selected from the group of AAVF5, AAVF7, AAVF13, AAVF15, and AAVF17; b) for heart or lung tissue or cells, the vector may be selected from the group of AAVF13, AAVF15, and AAVF17; c) for liver or CNS tissue or cells, the vector may be selected from AAVF5, AAVF13, AAVF17, AAVF7 or AAVF15; d) for stem cells, the vector may be AAVF17; e) for B cell progenitors, the vector may be AAVF5; f) for myeloid and erythroid progenitors, the vector may be AAVF12; and g) for lymph node, kidney, spleen, cartilage and bone tissues or cells, the vector may be selected from the group of the vector selected from the group of AAVF7, AAVF13, AAVF15, and AAVF17.

[0099] In addition, Clade F vectors or AAV vector variants may have a tropism for cells containing various tags, such as a six-His tag or an affinity tag, or for interferon responses, such as naturally occurring antibodies elicited or introduced monoclonal antibodies administered in response to a pathogen or tumor cell.

[0100] In certain embodiments, the cell may be a stem cell (e.g., a mammalian stem cell). In certain embodiments, the stem cell may be any type of stem cell including a hematopoietic stem cell, a pluripotent stem cell, a non-human embryonic stem cell or a mesenchymal stem cell. In certain embodiments, the stem cell (e.g., mammalian stem cell) may be a hematopoietic stem cell, a cord blood stem cell, a bone marrow stem cell, a fetal liver stem cell, or a peripheral blood stem cell. In some embodiments, the stem cell may be a CD34+ stem cell. In certain embodiments, the stem cell (e.g., mammalian stem cell) may be a hematopoietic stem cell or peripheral blood stem cell. Transduction of the stem cell may be either transient or permanent (also called persistent). If transient, one embodiment allows for the length of time the therapeutic nucleotide is used or expressed to be controlled either by the vector, by substance attached to the vector, or by external factors or forces.

[0101] In certain embodiments, the cell may be selected from the group consisting of a CD34+ Hematopoietic stem cell line (HSC), a K562 CD34+ leukemia cell line, a HepG2 human liver cell line, a peripheral blood stem cell, a cord blood stem cell, a CD34+ peripheral blood stem cell, a WI-38 human diploid fibroblast cell line, a MCF7 human breast cancer cell line, a Y79 human retinoblastoma cell line, a SCID-X1 LBL human EBV-immortalized B cell line, a primary human hepatocyte, a primary hepatic sinusoidal endothelial cell, and a primary skeletal muscle myoblast.

[0102] Also provided herein are methods in accordance with the claims of ex-vivo editing a genome of a cell of a subject comprising transducing the cell with a Clade F vector as defined by the claims. In accordance with the claims, transducing the cell with a Clade F vector occurs

without additional exogenous nucleases, such as a zinc finger nuclease (ZFN), a transcription activator-like effector nuclease (TALEN), or an RNA guided nuclease (CRISPR/Cas). In accordance with the claims, the cell may be any type of mammalian cell. In certain embodiments, the cell may be a stem cell as described herein. For example, in certain embodiments, methods of editing the genome of a stem cell may comprise transducing the stem cell with one or more Clade F vectors as defined by the claims. In certain embodiments, transduction of the stem cell may be performed without the need for additional exogenous nucleases. In certain embodiments, the cell may be a somatic cell as described herein. For example, in certain embodiments, methods of editing the genome of a somatic cell may comprise transducing the somatic cell with one or more Clade F vectors as defined by the claims. In accordance with the claims, transduction of the somatic cell is performed without the need for additional exogenous nucleases. In certain embodiments, the Clade F vector comprises one or more Clade F capsids as defined by the claims, an editing element (targeting cassette) selected from an internucleotide bond or a nucleotide sequence for integration into a target locus of a mammalian chromosome or comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) of the genome, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site). In certain embodiments, the internucleotide bond or nucleotide sequence or the one or more therapeutic nucleotide sequences may be integrated into the genome without the need for additional exogenous nucleases for DNA cleavage prior to integration.

[0103] Also described herein are methods of treating a disease or disorder in a subject by ex-vivo editing a genome of a cell of the subject including transducing the cell with a Clade F vector or an AAV vector variant and further transplanting the transduced cell into the subject to treat the disease or disorder. In certain aspects, the method may comprise transducing the cell of the subject with a Clade F vector or an AAV vector variant vector described herein. In certain aspects, the cell may be transduced without additional exogenous nucleases. Transduction of the cell with the Clade F vector or the AAV vector variant may be performed as provided herein or by any method of transduction known to one of ordinary skill in the art. The cell may be transduced with the Clade F vector or the AAV vector variant at a multiplicity of infection (MOI) of 50,000; 100,000; 150,000; 200,000; 250,000; 300,000; 350,000; 400,000; 450,000; or 500,000, or at any MOI that provides for optimal transduction of the cell. In certain aspects, the transduced cell is further transplanted into the subject, wherein the transduced cell treats the disease or disorder. The cell may be any type of mammalian cell described herein.

[0104] Also provided herein is an ex vivo method of editing a target locus of a mammalian genome in accordance with the claims. In some embodiments, the method comprises transducing a cell as defined by the claims (such as a human, mouse, bovine, equine, canine, feline, rat, or rabbit cell) comprising the mammalian genome with an AAV as defined by the claims (e.g., a replication-defective AAV comprising a correction genome enclosed in a capsid). In some aspects, the method comprises (a) obtaining mammalian cells from a mammal (such

as a human, mouse, bovine, equine, canine, feline, rat, or rabbit); (b) culturing the mammalian cells ex-vivo to form an ex-vivo culture; (c) transducing the mammalian cells with an AAV as described herein (e.g., a replication-defective AAV comprising a correction genome enclosed in a capsid) in the ex-vivo culture to form transduced mammalian cells; and (d) administering the transduced mammalian cells to the mammal. In some aspects, the method comprises (a) obtaining mammalian cells from a first mammal; (b) culturing the mammalian cells ex-vivo to form an ex-vivo culture; (c) transducing the mammalian cells with an AAV as described herein (e.g., a replication-defective AAV comprising a correction genome enclosed in a capsid) in the ex-vivo culture to form transduced mammalian cells; and (d) administering the transduced mammalian cells to a second mammal. In some aspects, the first mammal and the second mammal are different species (e.g., the first mammal is human, mouse, bovine, equine, canine, feline, rat, or rabbit and the second mammal is a different species). In some aspects, the first mammal and the second mammal are the same species (e.g., are both human, mouse, bovine, equine, canine, feline, rat, or rabbit). The method may comprise administering an AAV as described herein (e.g., a replication-defective AAV comprising a correction genome enclosed in a capsid) to a mammal (such as a human, mouse, bovine, equine, canine, feline, rat, or rabbit) in an amount effective to transduce cells of the mammal with the AAV in-vivo.

[0105] In some embodiments of any one of the methods, the mammalian cells are from a tissue selected from the group consisting of connective tissue (including blood), muscle tissue, nervous tissue, and epithelial tissue. In some embodiments of any one of the methods, the mammalian cells are from an organ selected from the group consisting of lung, heart, liver, kidney, muscle, brain, eye, breast, bone, and cartilage. In some embodiments of any one of the methods, the mammalian cells are stem cells. In some embodiments, the stem cells are hematopoietic stem cells or peripheral blood stem cells. In some embodiments of any one of the methods, the mammalian cells are a CD34+ cells.

[0106] In some embodiments of any one of the methods, the AAV (e.g., Clade F AAV) has a chromosomal integration efficiency of at least about 1% (e.g., at least about 2%, at least about 3%, at least about 4%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 98%, or about 100%) for integrating the editing element into the target locus of the mammalian chromosome. In some embodiments of any one of the methods, the AAV (e.g., Clade F AAV) has a chromosomal integration efficiency of at least about 1% (e.g., at least about 2%, at least about 3%, at least about 4%, at least about 5%, at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 98%, or about 100%) for integrating the editing element into the target locus of the mammalian chromosome in the absence of an exogenous nuclease. In some embodiments of any one of the methods, the editing element of the correction genome is integrated into the target locus of the mammalian chromosome with a chromosomal integration efficiency ranging from 10% to 70%, 20% to 70%, 40% to 70%, 50% to 70%, 10% to 80%, 20% to 80%, 40% to 80%, 50% to 80%, 10% to 90%, 20% to 90%, 40% to 90%, 50% to 90%, 10% to 100%, 20% to 100%, 40% to 100%, or 50% to

100% of the mammalian cells. In some embodiments of any one of the methods, the editing element of the correction genome is integrated into the target locus of the mammalian chromosome with a chromosomal integration efficiency ranging from 10% to 70%, 20% to 70%, 40% to 70%, 50% to 70%, 10% to 80%, 20% to 80%, 40% to 80%, 50% to 80%, 10% to 90%, 20% to 90%, 40% to 90%, 50% to 90%, 10% to 100%, 20% to 100%, 40% to 100%, or 50% to 100% of the mammalian cells in the absence of an exogenous nuclease.

[0107] In some embodiments of any one of the methods, the AAV (e.g., Clade F AAV) has a chromosomal integration efficiency further characterized by an allele frequency in a population of cells of at least about 10% (e.g., at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 75%, at least about 85%, at least about 90%, or at least about 95%) for the allele comprising the editing element integrated into the target locus of the mammalian chromosome. In some embodiments, the allele frequency in a population of cells is an allele frequency in a population of cells in vitro, such as population of a cell type provided herein in vitro (e.g., a CD34+ Hematopoietic stem cell line (HSC), a K562 CD34+ leukemia cell line, a HepG2 human liver cell line, a peripheral blood stem cell, a cord blood stem cell, a CD34+ peripheral blood stem cell, a WI-38 human diploid fibroblast cell line, a MCF7 human breast cancer cell line, a Y79 human retinoblastoma cell line, a SCID-X1 LBL human EBV-immortalized B cell line, a primary human hepatocyte, a primary hepatic sinusoidal endothelial cell, or a primary skeletal muscle myoblast).

[0108] Also described herein are methods of treating a disease or disorder in a subject by ex-vivo editing a genome of stem cell of the subject and further transplanting the edited cell into the subject to treat the disease or disorder. In certain aspects, the methods of treating a disease or disorder in a subject by editing a genome of a stem cell of the subject may comprise the steps of transducing the stem cell of the subject with a AAV Clade F vector or an AAV vector variant as described herein and transplanting the transduced stem cell into the subject, wherein the transduced stem cell treats the disease or disorder. In certain aspects, the AAV Clade F vector or the AAV vector variant may comprise one or more Clade F capsids or capsid variants, an editing element (targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) in the genome of the stem cell, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site). Transducing the stem cell is performed without additional exogenous nucleases. In certain aspects, the one or more therapeutic nucleotide sequences may be integrated into the genome without the need for additional exogenous nucleases for DNA cleavage prior to integration.

[0109] In certain embodiments, when the cell is a stem cell, the disease or disorder that is treated may be any disease or disorder that is caused by one or more mutations of the genome. In certain embodiments, the disease or disorder that is treated is selected from inherited metabolic diseases, lysosomal storage diseases, mucopolysaccharidosis, immunodeficiency diseases, and hemoglobinopathy diseases and infections. In certain

embodiments, when the cell to be edited is a stem cell, the AAV Clade F vector may be selected from the group of AAVF7, AAVF12, AAVF15, and AAVF17,. In certain embodiments, when the cell to be edited is a stem cell, the Clade F vector may be selected from the group of AAVF5, AAVF7, AAVF12, AAVF15, and AAVF17,. In certain aspects, the Clade F vector or AAV vector variant may comprise one or more Clade F capsids or capsid variants comprising a polynucleotide sequence selected from the group of AAVF7 (SEQ ID NO: 27), AAVF12 (SEQ ID NO: 30), AAVF15 (SEQ ID NO: 33), AAVF17 (SEQ ID NO: 35), variants, fragments, mutants and combinations thereof. In certain aspects, the Clade F vector or the AAV vector variant may comprise one or more Clade F capsids or capsid variants comprising a polynucleotide sequence selected from the group of AAVF5 (SEQ ID NO: 25), AAVF7 (SEQ ID NO: 27), AAVF12 (SEQ ID NO: 30), AAVF15 (SEQ ID NO: 33), AAVF17 (SEQ ID NO: 35), variants, fragments, mutants and combinations thereof. In certain embodiments, the AAV Clade F vector may comprise one or more Clade F capsids comprising a polypeptide sequence selected from the group of AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), and AAVF17 (SEQ ID NO: 13). In certain embodiments, the AAV Clade F vector may comprise one or more Clade F capsids comprising a polypeptide sequence selected from the group of AAVF5 (SEQ ID NO: 11), AAVF7 (SEQ ID NO: 8), AAVF12 (SEQ ID NO: 12), AAVF15 (SEQ ID NO: 16), and AAVF17 (SEQ ID NO: 13).

[0110] In another aspect, the AAV Clade F vectors or AAV vector variants capable of genome editing, from CD34⁺ HSC or from another source, may be used for high efficiency transduction of stem cells, including HSC and iPSC, and other cells, such as those of the heart, joint, central nervous system, including the brain, muscle, and liver. If the AAV Clade F vectors or AAV vector variants are used in vitro, they may be used for research and investigation purposes or to prepare cells or tissues that will later be implanted into a subject. Preferably, the subject is a mammal, such as a human, but may be any other animal that has tissues that can be transduced by the present vectors and methods of using those vectors. The present AAV Clade F vectors or the AAV vector variants are well suited for both human and veterinary use. The AAV Clade F vectors or AAV vector variants may also be used in vitro for the transient transduction of stem cells, such as HSC. The length of transduction may be controlled by culture conditions. If the AAV Clade F vectors or AAV vector variants are used in vivo, they may be directly administered to the subject receiving the therapy for uptake or use in the target cells, such as liver or cartilage cells. If the AAV Clade F vectors or AAV vector variants are used for transducing cells of the central nervous system, they are preferably able to traverse the blood-brain barrier and maintain their efficacy.

[0111] Also provided herein is a replication-defective AAV for use in a method of treating a disease or disorder in a subject by in vivo genome editing of a cell of the subject by directly administering an AAV Clade F vector as defined by the claims to the subject. The AAV Clade F vector may be any AAV Clade F vector in accordance with the claims. In certain embodiments, the AAV Clade F vector may comprise one or more Clade F capsids as defined in the claims, an editing element (targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) in the genome of the stem cell, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and

having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is downstream of the target locus (target site). In certain embodiments, the AAV Clade F vector that is administered treats the disease or disorder by genome editing of the cell of the subject. In accordance with the claims, the in vivo genome editing occurs without additional exogenous nucleases. In certain embodiments, the one or more Clade F capsids comprise a polypeptide sequence as defined by the claims. In certain aspects, the polypeptide sequence may be selected from the sequences provided in Figure 1 of US Patent Publication Number 20130096182A1 or in Figure 1 herein, variants, fragments, mutants, and combinations thereof. In certain embodiments, the AAV Clade F vectors or AAV vector variants are preferably administered in a therapeutically effective amount via a suitable route of administration, such as injection, inhalation, absorption, ingestion or other methods.

[0112] Previous studies, including Xu *et al.*, Wang *et al.*, and Carbonaro *et al.*, have shown transduction of HSCs following in vivo delivery of a viral vector (see Xu 2004; Wang 2014; and Carbonaro 2006). However, all three of these studies involved either retroviruses (Xu 2004) or lentiviruses (Wang 2014 and Carbonaro 2006). Additionally, the injections in Xu *et al.* and Carbonaro *et al.* were performed in neonatal mice, and rapamycin and intrafemoral injection was required for efficient transduction in Wang *et al.* None of these papers, however, report transduction of HSCs by in vivo transduction of Clade F vectors or AAV vector variants into adult mice. The novel results provided in Example 3 are the first to show AAV vector transduction on HSCs by intravenous injection.

[0113] As shown in Example 3 below, intravenous injection of Clade F vectors (or AAV vector variants) pseudotyped with AAVF7 or AAVF17 resulted in transduction of human CD34+ hematopoietic stem and progenitor cells in vivo. The intravenous injected Clade F vectors or AAV vectors trafficked to sites of human hematopoiesis and transduced human cells. Intravenous injection of Clade F vectors or AAV vector variants resulted in Venus expression in human CD34+ stem progenitor cells as well as their CD45+ progeny. These data show that intravenous injection of Clade F vectors or AAV vector variants can be used for in vivo genome engineering without the need for stem cell harvest, ex vivo transduction, conditioning of the recipient, and subsequent transplantation of transduced cells. This approach makes stem cell gene therapy significantly safer, more accessible to patients worldwide, less expensive, and obviates the need for hospitalization.

[0114] In accordance with the claims, a replication-defective AAV for use in a method of treating a disease or disorder in a subject by in vivo genome editing of a cell of the subject by directly administering an AAV Clade F vector to the subject is disclosed. In certain embodiments, the AAV Clade F vector may comprise one or more Clade F capsids as defined in the claims, an editing element (targeting cassette) comprising one or more therapeutic nucleotide sequences to be integrated into a target locus (target site) of the genome, a 5' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and having homology to a region that is upstream of the target locus (target site), and a 3' homologous arm polynucleotide sequence flanking the editing element (targeting cassette) and

having homology to a region that is downstream of the target locus (target site), wherein the vector transduces a cell of the subject and integrates the one or more therapeutic nucleotide sequences into the genome of the cell. In certain embodiments, the one or more Clade F capsids may comprise a polypeptide sequence selected from the group of AAVF1 (SEQ ID NO: 2), AAVF2 (SEQ ID NO: 3), AAVF11 (SEQ ID NO: 4), AAVF3 (SEQ ID NO: 5), AAVF4 (SEQ ID NO: 6), AAVF6 (SEQ ID NO: 7), AAVF7 (SEQ ID NO: 8), AAVF8 (SEQ ID NO: 9), AAVF9 (SEQ ID NO: 10), AAVF5 (SEQ ID NO: 11), AAVF12 (SEQ ID NO: 12), AAVF17 (SEQ ID NO: 13), AAVF13 (SEQ ID NO: 14), AAVF14 (SEQ ID NO: 15), AAVF15 (SEQ ID NO: 16), and AAVF16 (SEQ ID NO: 17). In certain embodiments, the one or more Clade F capsids may comprise a polypeptide sequence of AAVF7 (SEQ ID NO: 8) or AAVF17 (SEQ ID NO: 13). In certain embodiments, the one or more Clade F capsids may comprise a polypeptide sequence of AAVF5 (SEQ ID NO: 11), AAVF (SEQ ID NO: 8) or AAVF17 (SEQ ID NO: 13). In certain embodiments, the AAV Clade F vector does not contain a promoter for the one or more therapeutic nucleotide sequences. In certain embodiments, the target locus (target site) may be a safe harbor site. In certain embodiments, the safe harbor site may be the AAVS1 locus on chromosome 19. In certain embodiments, the cell may be a stem cell. In certain embodiments, the stem cell may be a hematopoietic stem cell, a pluripotent stem cell, a non-human embryonic stem cell, or a mesenchymal stem cell. In certain embodiments, the disease or disorder may be caused by one or more mutations in the cell genome. In certain embodiments, the disease or disorder may be selected from an inherited metabolic disease, lysosomal storage disease, mucopolysaccharidosis, immunodeficiency disease, and hemoglobinopathy disease and infection.

[0115] Further demonstrating the efficacy of vivo applications, transplantation of transduced cells to immune-deficient mice with the isolate variants (relative to AAV9) resulted in prolonged and sustained transgene expression and may be used for gene therapy. In certain aspects, when delivered systemically, these vectors display a tropism for the liver and cartilage, with implications for therapy of inherited, acquired, infectious and oncologic diseases. With respect to the liver transduction, the present AAV isolates have up to approximately 10-fold higher liver transduction levels than the current gold standard for systemic gene delivery to the liver, AAV8. This property can be exploited for gene-based enzyme replacement therapy from the liver for diseases such as hemophilia, enzyme deficiency diseases, and atherosclerosis. The additional tropism of the present AAV isolates for cartilaginous tissue in joints may be exploited for the treatment of bone disorders such as arthritis, osteoporosis or other cartilage/bone based diseases. The variant sequences and methods may accordingly be used for transient transduction where long term integration is not desirable.

[0116] Members of the AAV Clade F capsid family or AAV capsid variant family transduce HSC, e.g. AAVF 15 and AAVF 17, giving rise to long-term engraftment with sustained gene expression and are thus strong candidates for stem cell gene therapy vectors. AAVF17 and AAVF15 (also referred in abbreviated form as "HSC17" and "HSC15") supported the highest levels of long-term in vivo transduction, up to 22 weeks post-transplantation. Serial bioluminescent imaging following intravenous injection of the AAV variants revealed that AAVF15 generally supported the highest levels of long-term transgene expression in vivo.

Other AAV variants including AAVF13 and 17 also supported strong in vivo transduction.

[0117] AAVF15 was found to be highly liver tropic, about 5-10 fold higher than AAV9. AAVF13 and AAVF15 also transduced the heart and skeletal muscle at least 10-fold better than AAV9. In vitro neutralization titers revealed that the prevalence of antibodies to AAVF 1-9 capsids in pooled human IVIG were similar to AAV9, while antibodies to AAVF13, AAVF15, AAVF16 and AAVF17 were somewhat less prevalent. In vivo neutralization assays confirmed that over 100-fold higher vector genome copies/cell were found in liver and muscle following IVIG administration with AAVF15 compared to AAV9, suggesting that pre-existing antibodies did not completely neutralize AAVF15. Muscle diseases or disorders may comprise any cell, tissue, organ, or system containing muscle cells which have a disease or disorder, including the heart, such as coronary heart disease or cardiomyopathy.

[0118] In addition, site-specific mutagenesis experiments indicate that the R505G mutation in AAVF15 is responsible for the enhanced liver tropism. The AAV Clade F vectors or AAV vector variants may be used to treat a whole host of genetic diseases such as hemophilia, atherosclerosis and a variety of inborn errors of metabolism. In one instance, AAVF15 effectively treats hemophilia B. Some members of this family also target the joints after systemic injection, which may be used to treat joint and cartilage diseases such as arthritis. Other members of the family target the heart upon intravenous injection. Yet other members of the family target the brain. In some aspects, a vector comprising AAVF5 capsid proteins is provided as part of a method, kit or composition provided herein, as AAVF5 was shown to transduce multiple cell types (see Figure 4).

[0119] In certain aspects, methods of treating a neurological disease or disorder in a subject by genome editing may comprise administering an AAV Clade F vector or AAV vector variant capable of crossing the blood-brain barrier, blood-ocular barrier, or blood-nerve barrier. Certain of the AAV Clade F vectors or AAV vector variants disclosed herein have the unique ability to traverse the biological junctions that were previously unknown to be accessible to any vector for gene therapy or other diagnostic or therapeutic purposes using a modified viral vector. These junctions have common characteristics. The blood-brain barrier is a separation between blood circulating in the body and the brain extracellular fluid in the central nervous system and is created by tight junctions around capillaries. The blood-brain barrier generally allows only the passage of by diffusion of small hydrophobic molecules. The blood-ocular barrier is a separation made by between the local blood vessels and most parts of the eye and is made by endothelium of capillaries of the retina and iris. The blood-nerve barrier is the physiological space within which the axons, Schwann cells, and other associated cells of a peripheral nerve function and is made of endoneurial microvessels within the nerve fascicle and the investing perineurium. As with three of these barriers, there is restricted permeability to protect in the internal environment, here, the nerve, from drastic concentration changes in the vascular and other extracellular spaces. The vector that traverses any of these barriers has a unique ability to deliver one or more therapeutic nucleotide sequences for treating the neurological disease or disorder or to act as a labeled and or diagnostic agent. Certain of the AAV Clade F vectors or AAV vector variants that have been experimentally validated as being particularly well suited

for crossing these biological barriers include AAVF15, AAVF15 A346T, and AAVF15 R505G.

[0120] There are many neurological diseases or disorders that are well known to one of skill in the art, which may be generally classified by cell or organ-type such as a disease or disorder of the brain, spinal cord, ganglia, motor nerve, sensory nerve, autonomic nerve, optic nerve, retinal nerve, and auditory nerve. By way of example, brain diseases or disorders may include cancer or other brain tumor, inflammation, bacterial infections, viral infections, including rabies, amoeba or parasite infections, stroke, paralysis, neurodegenerative disorders such as Alzheimer's Disease, Parkinson's Disease, or other dementia or reduction in cognitive functioning, plaques, encephalopathy, Huntington's Disease, aneurysm, genetic or acquired malformations, acquired brain injury, Tourette Syndrome, narcolepsy, muscular dystrophy, tremors, cerebral palsy, autism, Down Syndrome, attention deficit and attention deficit hyperactivity disorder, chronic inflammation, epilepsy, coma, meningitis, multiple sclerosis, myasthenia gravis, various neuropathies, restless leg syndrome, and Tay-Sachs disease.

[0121] Muscle diseases or disorders include, by way of example only, myopathies, chronic fatigue syndrome, fibromyalgia, muscular dystrophy, multiple sclerosis, atrophy, spasms, cramping, rigidity, various inflammations, such as dermatomyositis, rhabdomyolysis, myofascial pain syndrome, swelling, compartment syndrome, eosinophilia-myalgia syndrome, mitochondrial myopathies, myotonic disorder, paralysis, tendinitis, polymyalgia rheumatic, cancer, and tendon disorders such as tendinitis and tenosynovitis.

[0122] Heart diseases or disorders include, by way of example only, coronary artery disease, coronary heart disease, congestive heart failure, cardiomyopathy, myocarditis, pericardial disease, congenital heart disease, cancer, endocarditis, and valve disease.

[0123] Lung diseases or disorders include, by way of example only, asthma, allergies, chronic obstructive pulmonary disease, bronchitis, emphysema, cystic fibrosis, pneumonia, tuberculosis, pulmonary edema, cancer, acute respiratory distress syndrome, pneumoconiosis, and interstitial lung disease.

[0124] Liver diseases or disorders include, by way of example only, cancer, hepatitis A, B, and C, cirrhosis, jaundice, and liver disease. Kidney diseases or disorders include, by way of example only, cancer, diabetes, nephrotic syndrome, kidney stones, acquired kidney disease, congenital disease, polycystic kidney disease, nephritis, primary hyperoxaluria, and cystinuria. Spleen diseases or disorders include, by way of example only, cancer, splenic infarction, sarcoidosis, and Gaucher's disease. Bone diseases or disorders include, by way of example only, osteoporosis, cancer, low bone density, Paget's disease, and infection.

[0125] With any of these diseases or disorders treated using therapeutic nucleotide sequences or even small molecules transported by or with the AAV Clade F vectors or AAV vector variants, the therapeutic nucleotide sequence may be, by way of example, a nucleic acid encoding a protein therapeutic, such as for cancer--an apoptotic protein, miRNA, shRNA, siRNA, other RNA-subtypes or a combination thereof. In some embodiments, the vectors are isolated and

purified as described herein. Isolation and purification are preferred in vivo administration to increase efficacy and reduce contamination. The vector may permanently or transiently transduce a transgene, which is a gene or other genetic material that has been isolated from one organism and introduced into another. Here, the other organism may be the subject receiving the vector.

[0126] In certain embodiments, the AAV Clade F vectors for genome editing may be selected based on experimental results of the highest efficacy in the given target cell or tissue for the given disease or disorder as shown herein. For example a) for muscle disease or disorders and for antibody genes or other vaccine treatments administered to the subject via the muscle, the AAV Clade F vector selected from the group of AAVF5, AAVF7, AAVF13, AAVF15, and AAVF17; b) for heart and lung disease or disorders, the vector selected from the group of AAVF13, AAVF15, and AAVF17; c) for liver or neurological diseases or disorders, the vector selected from AAVF5 and AAVF15; d) for conditions treated by engrafting stem cells, vector AAVF17; e) for conditions treated by transducing B cell progenitors, vector AAVF5; f) for conditions treated by transducing myeloid and erythroid progenitors, vector AAVF12; and g) for lymph node, kidney, spleen, cartilage and bone disease or disorders, the vector selected from the group of the vector selected from the group of AAVF7, AAVF13, AAVF15, and AAVF17; wherein the AAV Clade F vector as defined by the claims transduces the cell or tissue and the one or more therapeutic nucleotide sequences are integrated into the genome of the cell and treat the disease or disorder. In certain embodiments, the AAV Clade F vector may comprise one or more Clade F capsids as defined in the claims that demonstrates tropism for a cell as described herein.

[0127] The subject is any animal for which the method works, but is preferably a mammal, which may be a human. If the vector contains an antibody gene or other vaccine treatment it may be administered via injection in the muscle and may provide immunological protection against diseases including from HIV, influenza, malaria, tetanus, measles, mumps, rubella, HPV, pertussis, or any other vaccine. The vector may be packaged, isolated, and purified and may transduce a stem cell of any type with the at least one therapeutic nucleotide sequence. The vector may also transduce a transgene or carry corrective genes endogenous to the subject and/or to the other subjects of the same species.

[0128] "AAV" is an adeno-associated virus. The term may be used to refer to the virus or derivatives thereof, virus subtypes, and naturally occurring and recombinant forms, unless otherwise indicated. AAV has over 100 different subtypes, which are referred to as AAV-1, AAV-2, etc., and includes both human and non-human derived AAV. There are about a dozen AAV serotypes. The various subtypes of AAVs can be used as recombinant gene transfer viruses to transduce many different cell types.

[0129] "Recombinant," as applied to a polynucleotide means that the polynucleotide is the product of various combinations of cloning, restriction or ligation steps, and other procedures that result in a construct that is distinct from a naturally-occurring polynucleotide. A recombinant virus is a viral particle comprising a recombinant polynucleotide, including

replicates of the original polynucleotide construct and progeny of the original virus construct. A "rAAV vector" refers to a recombinant AAV vector comprising a polynucleotide sequence not of AAV origin (i.e., a polynucleotide heterologous to AAV), which is usually a sequence of interest for the genetic transformation of a cell.

[0130] A "helper virus" for AAV as used herein is virus that allows AAV to be replicated and packaged by a mammalian cell. Helper viruses for AAV are known in the art, and include, for example, adenoviruses (such as Adenovirus type 5 of subgroup C), herpes viruses (such as herpes simplex viruses, Epstein-Bar viruses, and cytomegaloviruses) and poxviruses.

[0131] "Joint tissue" is comprised of a number of tissues including cartilage, synovial fluid, and mature, progenitor and stem cells that give rise to, or are: (i) cartilage producing cells; (ii) Type I synoviocytes; (iii) Type II synoviocytes; (iv) resident or circulating leukocytes; (v) fibroblasts; (vi) vascular endothelial cells; and (vii) pericytes.

[0132] A "replication-competent" virus refers to a virus that is infectious and capable of being replicated in an infected cell. In the case of AAV, replication competence generally requires the presence of functional AAV packaging genes, as well as helper virus genes, such as adenovirus and herpes simplex virus. In general, rAAV vectors are replication-incompetent (also referred to herein as replication-defective) because they lack of one or more AAV packaging genes. In some embodiments, an AAV may be considered replication defective (or replication-incompetent) if the AAV has an essential absence of an AAV *rep* gene and/or an AAV *cap* gene. In some embodiments, an AAV may be considered replication defective (or replication-incompetent) if the AAV lacks an AAV *rep* gene and/or an AAV *cap* gene. In some aspects, a composition comprising AAV Clade F vectors or AAV variant isolates is a cell-free composition. The composition is generally free of cellular proteins and/or other contaminants and may comprise additional elements such as a buffer (e.g., a phosphate buffer, a Tris buffer), a salt (e.g., NaCl, MgCl₂), ions (e.g., magnesium ions, manganese ions, zinc ions), a preservative, a solubilizing agent, or a detergent, (e.g., a non-ionic detergent; dimethylsulfoxide).

[0133] Also described herein is an expression cassette comprising a polynucleotide sequence encoding a polypeptide comprising one or more of the Clade F capsids or AAV variant isolates, wherein the polynucleotide sequence encoding the polypeptide comprises a sequence having at least about 95%, 96%, 97%, more preferably about 98%, and most preferably about 99% sequence identity to the sequences taught in the present specification. Percentage identity may be calculated using any of a number of sequence comparison programs or methods such as the Pearson & Lipman, Proc. Natl. Acad. Sci. USA, 85:2444 (1988), and programs implementing comparison algorithms such as GAP, BESTFIT, FASTA, or TFASTA (from the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), or BLAST, available through the National Center for Biotechnology Information web site.

[0134] Also described is an expression cassette comprising a polynucleotide sequence

encoding a polypeptide comprising one or more of the Clade F capsids or AAV variant isolates, wherein the sequence is comprised of portions of the three genes comprising the capsid protein, V1-V3 (also referred to as VP1-VP3). For example, the cassette may comprise V1 from capsid AAVF1, a standard V2 as compared to AAV9 hu.14, and V3 from AAVF17 capsids. A capsid may comprise more than one of each of the capsid gene components. For example, Clade F capsids or capsid variants may be selected from any of the VP1-VP3 (V1-V3) for the capsid sequences set forth herein and may be combined in any order and in any combination so long as the desired property of increased transduction is achieved. For example, the capsid sequence could be VP1A-VP1B-VP2-VP3 (V1A-V1B-V2-V3), VP3-VP1-VP2 (V3-V1-V2), or VP1-VP2-VP3A-VP3B (V1-V2-V3A-V3B).

[0135] Also described herein are methods of immunization of a subject. Compositions comprising the Clade F capsids or capsid variants may be introduced into a subject in a manner that causes an immunological reaction resulting in immunity in the subject. The Clade F capsids or capsid variants may be in the composition alone or as part of an expression cassette. The expression cassettes (or polynucleotides) can be introduced using a gene delivery vector. The gene delivery vector can, for example, be a non-viral vector or a viral vector. Exemplary viral vectors include, but are not limited to Sindbis-virus derived vectors, retroviral vectors, and lentiviral vectors. Compositions useful for generating an immunological response can also be delivered using a particulate carrier. Further, such compositions can be coated on, for example, gold or tungsten particles and the coated particles delivered to the subject using, for example, a gene gun. The compositions can also be formulated as liposomes. The subject may be a mammal and can, for example, be a human.

[0136] The term "affinity tag" is used herein to denote a polypeptide segment that can be attached to a second polypeptide to provide for purification or detection of the second polypeptide or provide sites for attachment of the second polypeptide to a substrate. In principal, any peptide or protein for which an antibody or other specific binding agent is available can be used as an affinity tag. Affinity tags include a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al., Methods Enzymol. 198:3, 1991), glutathione S transferase (Smith and Johnson, Gene 67:31, 1988), Glu-Glu affinity tag (Grussenmeyer et al., Proc. Natl. Acad. Sci. USA 82:7952-4, 1985), substance P, Flag.TM. peptide (Hopp et al., Biotechnology 6:1204-10, 1988), streptavidin binding peptide, or other antigenic epitope or binding domain. See, in general, Ford et al., Protein Expression and Purification 2: 95-107, 1991, DNAs encoding affinity tags are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, N.J.).

[0137] Of the number of affinity tag purification systems available, the most frequently employed utilize polyhistidine (His) or glutathione S-transferase (GST) tags. His binds with good selectivity to matrices incorporating Ni²⁺ ions, typically immobilized with either iminodiacetic acid or nitrilotriacetic acid chelating groups. The technique is known as immobilized metal affinity chromatography. Absorption of the His-tagged protein is performed at neutral to slightly alkaline pH to prevent protonation and loss of binding capacity of the weakly basic histidine imidazole groups. Elution of the bound protein is caused by

displacement with imidazole or low pH conditions.

[0138] Methods of generating induced pluripotent stem cells from somatic cells without permanent introduction of foreign DNA are also described. The method involved transiently transducing stem cells with vectors comprising a Clade F capsid or capsid variant nucleotide sequence as described herein encoding a polypeptide sequence, or VP1 (V1) or VP3 (V3) portion thereof.

[0139] For these and other experiments, a person skilled in the art knows how to modify and propagate AAV. For example, AAV-2 can be propagated both as lytic virus and as a provirus. For lytic growth, AAV requires co-infection with a helper virus. Either adenovirus or herpes simplex can supply helper function. When no helper is available, AAV can persist as an integrated provirus, which involves recombination between AAV termini and host sequences and most of the AAV sequences remain intact in the provirus. The ability of AAV to integrate into host DNA allows propagation absent a helper virus. When cells carrying an AAV provirus are subsequently infected with a helper, the integrated AAV genome is rescued and a productive lytic cycle occurs. The construction of rAAV vectors carrying particular modifications and the production of rAAV particles, e.g., with modified capsids, is described, e.g., in Shi et al. (2001), *Human Gene Therapy* 12:1697-1711; Rabinowitz et al. (1999), *Virology* 265:274-285; Nicklin et al. (2001), *Molecular Therapy* 4:174-181; Wu et al. (2000), *J. Virology* 74:8635-8647; and Grifman et al. (2001), *Molecular Therapy* 3:964-974.

[0140] Also described herein is a pharmaceutical composition containing an AAV Clade F vector or AAV vector variant or AAV particle as described herein. The pharmaceutical composition containing an AAV Clade F vector or AAV vector variant or particle, preferably, contains a pharmaceutically acceptable excipient, adjuvant, diluent, vehicle or carrier, or a combination thereof. A "pharmaceutically acceptable carrier" includes any material which, when combined with an active ingredient of a composition, allows the ingredient to retain biological activity and without causing disruptive physiological reactions, such as an unintended immune reaction. Pharmaceutically acceptable carriers include water, phosphate buffered saline, emulsions such as oil/water emulsion, and wetting agents. Compositions comprising such carriers are formulated by well-known conventional methods such as those set forth in Remington's Pharmaceutical Sciences, current Ed., Mack Publishing Co., Easton Pa. 18042, USA; A. Gennaro (2000) "Remington: The Science and Practice of Pharmacy", 20th edition, Lippincott, Williams, & Wilkins; Pharmaceutical Dosage Forms and Drug Delivery Systems (1999) H. C. Ansel et al., 7th ed., Lippincott, Williams, & Wilkins; and Handbook of Pharmaceutical Excipients (2000) A. H. Kibbe et al., 3rd ed. Amer. Pharmaceutical Assoc. Such carriers can be formulated by conventional methods and can be administered to the subject at a suitable dose. Administration of the suitable compositions may be effected by different ways, e.g. by intravenous, intraperitoneal, subcutaneous, intramuscular, topical or intradermal administration. The composition may be formulated for administration to a mammal. The composition may be formulated for administration to a mammal via intravenous injection, subcutaneous injection, intramuscular injection, autologous cell transfer, or allogeneic cell transfer. The route of administration, of course, depends, inter alia, on the kind of vector

contained in the pharmaceutical composition. The dosage regimen will be determined by the attending physician and other clinical factors. As is well known in the medical arts, dosages for any one patient depends on many factors, including the patient's size, body surface area, age, sex, the particular compound to be administered, time and route of administration, the kind and stage of infection or disease, general health and other drugs being administered concurrently.

[0141] Some of the AAV Clade F vectors or capsid variants are capable of supporting long-term stable transgene expression in vivo after transplantation of transduced hematopoietic stem cells or after direct systemic delivery of rAAV.

[0142] In certain aspects, a nucleic acid comprising the Clade F capsids or AAV capsid isolate variants may be inserted into the genome of a new virus, where in the addition of the Clade F capsid or capsid isolate variant genes transmits the same or similar tissue or organ tropisms of the Clade F capsids or AAV capsid isolates to the new virus. Such gene therapy may be effected using in vivo and ex vivo gene therapy procedures; see, e.g., U.S. Pat. No. 5,474,935; Okada, Gene Ther. 3:957-964, 1996. Gene therapy using the AAV Clade F capsid or AAV capsid variant gene will typically involve introducing the target gene in vitro into the new virus, either alone or with another gene intended for therapeutic purposes. If the tropic gene is introduced with one or more additional genes, preferably the resulting polypeptides are administered for therapeutic purposes in the tissue for which the Clade F capsid or AAV isolate has a tropism. The virus may then be administered to patient in need of such therapy or may be administered ex vivo, such as to an organ awaiting transplant. The virus may be a retrovirus, an RNA virus, a DNA virus such as an adenovirus vector, an adeno-associated virus vector, a vaccinia virus vector, a herpes virus vector, and the like. A transfection method using a virus vector that uses a liposome for administration in which the new virus vector is encapsulated is also contemplated.

[0143] Kits are described herein that comprise one or more AAV Clade F vectors or AAV vector variants described herein or compositions or formulations thereof. The one or more AAV Clade F vectors or AAV vector variants in the kits may be used for genome editing of a cell. The kit may be used as a research tool to investigate the effect of genome editing by the one or more AAV Clade F vectors or AAV vector variants.

[0144] Other aspects of the disclosure relate to a packaging system for recombinant preparation of an AAV as described herein (e.g., an AAV Clade F vector or an AAV variant vector) and methods of use thereof. The packaging system comprises a Rep nucleotide sequence encoding one or more AAV Rep proteins; a Cap nucleotide sequence encoding one or more AAV Cap proteins of an AAV Clade F capsid as described herein; and a correction genome as described herein, wherein the packaging system is operative in a cell for enclosing the correction genome in the capsid to form an adeno-associated virus.

[0145] The packaging system comprises a first vector comprising the Rep nucleotide sequence and the Cap nucleotide sequence, and a second vector comprising the correction genome. As used in the context of a packaging system as described herein, a "vector" refers to

a nucleic acid molecule that is a vehicle for introducing nucleic acids into a cell (e.g., a plasmid, a virus, a cosmid, an artificial chromosome, etc.).

[0146] In the packaging system, the AAV Clade F capsid may comprise at least one or at least two proteins selected from Clade F VP1, Clade F VP2 and Clade F VP3. In the packaging system, the AAV Clade F capsid may comprise Clade F VP1, Clade F VP2 and Clade F VP3 proteins. In the packaging system, the AAV Clade F capsid may be selected from the group consisting of AAV9, AAVHSC1, AAVHSC2, AAVHSC3, AAVHSC4, AAVHSC5, AAVHSC6, AAVHSC7, AAVHSC8, AAVHSC9, AAVHSC10, AAVHSC11, AAVHSC12, AAVHSC13, AAVHSC14, AAVHSC15, AAVHSC16, AAVHSC17, AAVHU31, and AAVHU32.

[0147] In the packaging system, the Rep nucleotide sequence may encode an AAV2 Rep protein. In the packaging system, the AAV2 Rep protein encoded may be at least one of Rep 78/68 or Rep 68/52. In the packaging system, the nucleotide sequence encoding the AAV2 Rep protein may comprise a nucleotide sequence that encodes a protein having a minimum percent sequence identity to the AAV2 Rep amino acid sequence of SEQ ID NO:40, wherein the minimum percent sequence identity is at least 70% (e.g., at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or 100%) across the length of the amino acid sequence of the AAV2 Rep protein.

[0148] Exemplary AAV2 Rep amino acid sequence (SEQ ID NO: 40) -

mpgfyevikvpsdldehlpgisdsfvnwvaekewelppdsdmdlnlieqapltvaeklqrdfltewrrvskapealffv
 qfekgesyfhmhvlvettgvksmvlgrflsqirekliqriyrgieptlpnwfavtktrngagggnkvvdecyipnyllpktqpelq
 wawtnmeqylsaclnlterkrivaqlhthvsqtqeqnkenqnpnsdapvirsktsarymelvgwlvdkgitsekqwqedqas
 yisfnaasnsrsqikaaldnagkimsllktapdylvgqqpvedissnriykilelgydpqyaasvflgwatkkfgkrntiwlfgpa
 ttgktniaeaiahtvpfygcvnwtnefpndcvdkmviwweegkmtakvvesakailggskvrvdqkckssaqidptpvivt
 sntnmcaavidgnsttfhqqlqdrmfkfeltrldhdfgkvtkqevkdffrwakdhvvevehefyvkkggakkrpapsdadis
 epkrvresvaqpstsdaeasinyadryqnkcsrhvgmnlmlfpcrqcermnqnsnicfthgqkdclecfpvsesqpvsvvkka
 yqkleyihhimgkvpdactadlvnvdlddcifeq

[0149] In the packaging system, the packaging system may further comprise a third vector, e.g., a helper virus vector. The third vector may be an independent third vector, integral with the first vector, or integral with the second vector. In some embodiments, the third vector comprises genes encoding helper virus proteins.

[0150] In the packaging system, the helper virus may be selected from the group consisting of adenovirus, herpes virus (including herpes simplex virus (HSV)), poxvirus (such as vaccinia virus), cytomegalovirus (CMV), and baculovirus. In the packaging system, where the helper virus is adenovirus, the adenovirus genome may comprise one or more adenovirus RNA genes selected from the group consisting of E1, E2, E4 and VA. In the packaging system, where the helper virus is HSV, the HSV genome may comprise one or more of HSV genes selected from the group consisting of UL5/8/52, ICPO, ICP4, ICP22 and UL30/UL42.

[0151] In the packaging system, the first, second, and/or third vector may be contained within one or more transfecting plasmids. The first vector and the third vector may be contained within a first transfecting plasmid. The second vector and the third vector may be contained within a second transfecting plasmid.

[0152] In the packaging system, the first, second, and/or third vector may be contained within one or more recombinant helper viruses. The first vector and the third vector may be contained within a recombinant helper virus. The second vector and the third vector may be contained within a recombinant helper virus.

[0153] Also described herein is a method for recombinant preparation of an AAV as described herein (e.g., an AAV Clade F vector or AAV variant vector), wherein the method comprises transfecting or transducing a cell with a packaging system as described under conditions operative for enclosing the correction genome in the capsid to form the AAV as described herein (e.g., an AAV Clade F vector or AAV variant vector). Exemplary methods for recombinant preparation of an AAV include transient transfection (e.g., with one or more transfection plasmids containing a first, and a second, and optionally a third vector as described herein), viral infection (e.g. with one or more recombinant helper viruses, such as a adenovirus, poxvirus (such as vaccinia virus), herpes virus (including herpes simplex virus (HSV), cytomegalovirus, or baculovirus, containing a first, and a second, and optionally a third vector as described herein), and stable producer cell line transfection or infection (e.g., with a stable producer cell, such as a mammalian or insect cell, containing a Rep nucleotide sequence encoding one or more AAV Rep proteins and/or a Cap nucleotide sequence encoding one or more AAV Cap proteins of an AAV Clade F capsid as described herein, and with a correction genome as described herein being delivered in the form of a transfecting plasmid or a recombinant helper virus).

[0154] The following examples are intended to illustrate various embodiments of the disclosure.

EXAMPLES

Example 1: AAV Clade F Vector Variant Mediated Genome Editing in a CD34+ Human Hematopoietic Stem Cell Line or a K562 Cell Line

[0155] Genome editing through site-specific insertion or targeted integration of specific DNA sequences using AAVF (AAVHSC) vectors without the use of an exogenous nuclease was performed in human CD34+ hematopoietic cell lines from healthy donors or the K562 cell line, which is a CD34+ erythroleukemic cell line. One set of donor recombinant AAV vectors, ITR-AAVS1-FP vectors, was constructed and was used to integrate a transgene into the AAVS1 locus on chromosome 19, the natural wild-type AAV integration site (Kotin, 1992; Giraud,

1994). The AAVS1 locus on chromosome 19 qter13.3-13.4 was previously shown to be a "safe harbor" site for the insertion of transgenes since genes inserted here are expressed with no pathogenic consequences, which is similar to wild-type AAV that integrates at this locus with no pathogenic consequences (Giraud, 1994; Linden, 1996A; Linden 1996B). The transgene to be integrated was a Venus yellow fluorescent protein ("YFP" or "FP") gene, which was flanked on each side by approximately 800 nucleotides having homology with the AAVS1 locus on human chromosome 19 (see schematic in Figure 3). The donor AAV vector was designed such that the transgene was promoterless and would only be expressed if it was integrated at the correct locus, which would be downstream from chromosomally encoded regulatory sequences (see Figure 4). Thus, any Venus YFP transgene expression that occurred was under the control of a chromosomal promoter located in or near AAVS1.

[0156] The donor vector, ITR-AAVS 1-FP, was packaged into AAVHSC capsids according to the standard AAV packaging method described in Chatterjee et al, 1992. Specifically, ITR-AAVS 1-FP was packaged into AAVHSC7, AAVHSC15, or AAVHSC17 capsids forming the pseudotyped AAVHSC-AAVS 1-FP vector (i.e., a AAV vector variant). Human CD34+ hematopoietic stem cell lines or K562 cells were transduced with the pseudotyped AAVHSC-AAVS1-FP at different multiplicities of infection (MOI) (i.e., 50,000 MOI; 100,000 MOI; 200,000 MOI; 300,000 MOI; and 400,000 MOI).

[0157] Integration of the YFP transgene into the AAVS1 locus by homologous recombination was initially assayed by cytofluorometric analysis of YFP expression in transduced K562 cells. Targeted integration using the AAVHSC7 FP vector resulted in expression of the YFP transgene 24 hours post-transduction (Figures 5 and 7A) and 72 hours post-transduction (Figures 6 and 7B). Additionally, as the MOI of the AAVHSC7 FP vector was increased, the average percentage of YFP expression also increased (Figures 7A and B).

[0158] Targeted integration of the YFP transgene was further confirmed by PCR amplification of the edited genome using primers located outside of the homology regions. Briefly, DNA was extracted from K562 cells transduced at an MOI of 100,000 vector genomes/cell with AAVHSC7 FP vector. PCR amplification was performed using the "OUT Forward Primer Region" and "OUT Reverse Primer Region" primers (see Figure 4). Integration of the YFP transgene resulted in an increase in size of the AAVS1 locus from the wild type size of ~1.9kb to the YFP transgene containing -3.1 kb fragment (see Figure 4, compare line labeled "Fragment 1" with line labeled "Fragment 2"). Amplification of the -3.1 kb fragment containing the YFP transgene within the chromosome 19 AAVS1 locus indicated that the YFP transgene was effectively integrated into the AAVS1 locus in cells transduced with the AAVHSC7 FP vector (see Figures 8A and 8B, lane 4).

EXAMPLE 2: AAV Vector Variant Mediated Genome Editing in Primary Human CD34+ Peripheral Blood Stem Cells.

[0159] Genome editing through site-specific insertion or targeted integration of specific DNA

sequences using AAVHSC vectors without the use of an exogenous nuclease was also performed in human CD34+ primary peripheral blood-derived human hematopoietic stem cells (PBSCs). Briefly, the vector, ITR-AAVS 1-FP, was packaged in AAVHSC capsids including AAVHSC7, AAVHSC12, AAVHSC15, and AAVHSC17 (see Chatterjee, 1992 for the standard AAV packaging method). Primary CD34+ cells were transduced with the pseudotyped AAVHSC-AAVS 1-FP vector (i.e., a AAV vector variant) at MOIs of 100,000 and 150,000.

[0160] Integration of the YFP transgene into the AAVS1 locus by homologous recombination was assayed by cytofluorometric analysis of YFP expression. Targeted integration using the AAVHSC7 FP and AAVHSC17 FP vectors in primary CD34+ cells resulted in expression of the YFP transgene 1 day post-transduction at an MOI of 150,000 (Figure 9), 4 days post-transduction at an MOI of 100,000 (Figure 10), and 18 days post-transduction at an MOI of 100,000 (Figure 11). The percentage of positive cells expressing YFP at 5.5 weeks post transduction at an MOI of 100,000 (39 days) did not decline (see Figures 12A and B, compare 20 day results with 39 day results). This long term expression of a promoterless YFP transgene in a dividing cell population indicates accurate integration of the transgene.

[0161] Targeted integration of the YFP transgene was further confirmed by PCR analysis. The edited genome was amplified using primers that amplify the 5' junction region between the inserted transgene sequence and the native chromosomal 5' homology arm sequence (see Figure 4, see line labeled "Fragment 3"). Briefly, DNA was extracted from primary CD34+ cells transduced at an MOI of 150,000 vector genomes/cell with the AAVHSC7 FP vector. PCR amplification was performed using the "OUT Forward Primer Region" and the "In Reverse Primer" primers (see Figure 4). Amplification of a -1.7 kb fragment of the 5' junction region for transduced primary CD34+ cells indicated that the YFP transgene was successfully integrated into the AAVS1 locus (see Figure 13, lane 5). Whereas, there was no amplified product for those cells that were not transduced with the AAVHSC7 FP vector (see Figure 13, lane 3).

[0162] Targeted integration of the YFP transgene was further confirmed by sequence analysis of different portions of the edited AAVS1 locus. Sequencing was performed beginning near the "OUT Forward Primer Region" (see Figure 14), near the 5' homology arm (see Figure 15), near the 5' region of the regulatory elements (see Figure 16), near the 3' region of the regulatory elements (see Figure 17), near the 5' region of the transgene (see Figure 18), and near the "IN Reverse Primer" region (see Figure 19). Sequencing results indicated that the YFP gene was present and was successfully integrated into the AAVS1 locus.

[0163] As provided in Examples 1 and 2, the AAVHSC vectors successfully mediated efficient targeted genome editing in human CD34+ hematopoietic cell lines and CD34+ PBSCs without the need for addition of exogenous endonucleases. AAVHSC vectors were capable of directing integration of the YFP transgene to the AAVS1 locus on chromosome 19 based upon flanking homology arms corresponding to the AAVS1 locus. AAV-mediated transgenesis has previously been reported; however, reported frequencies have been low, usually on the order of 1 in 1e6 cells to 1 in 1e4 cells. As shown herein, targeted genome editing using AAVHSC vectors occurred at frequencies of approximately 10% of primary cells long term, which is 1,000 to

100,000 fold more efficient than previously reported (see, e.g., Khan, 2011). Expression of the YFP transgene in human CD34+ hematopoietic cell lines was observed as early as day one post-transduction and was confirmed on day three. Expression of the YFP transgene in PBSCs was observed starting from day one and continued long term (up to almost 6 weeks), which was the latest time point analyzed. No overt toxicity was observed as a result of AAVHSC vector transduction. Based upon the high frequency of insertion, ease of use, and lack of toxicity observed, therapies based upon targeted genome editing using AAVHSC vectors is practical and feasible.

EXAMPLE 3: In Vivo Genome Engineering with AAV Vector Variants.

[0164] AAVHSC vectors encoding luciferase and AAVHSC vectors encoding Venus were injected into adult immune-deficient mice previously xenografted with human cord blood CD34+ HSCs. As shown below, intravenous injection of AAVHSC vectors resulted in transduction of human CD34+ hematopoietic stem and progenitor cells in vivo, and the intravenous injected AAVHSC vectors trafficked to sites of human hematopoiesis and transduced human cells.

[0165] *Methods.* Briefly, immune-deficient NOD/SCID adult mice were first irradiated with a sublethal dose of 350cGy from a ^{137}Cs source. Second, one million human cord blood CD34+ cells were injected into the sublethally-irradiated immune-deficient NOD/SCID mice. Next, two hours after CD34+ HSC transplantation, the mice were injected intravenously with approximately $1\text{E}11$ - $5\text{E}11$ particles of AAVHSC-Luciferase vector (either AAVHSC7-Luciferase vector or AAVHSC17-Luciferase vector). These vectors were used in the absence of an exogenous nuclease. The AAVHSC-Luciferase vectors encode the single-stranded firefly luciferase gene (ssLuc) under the control of the ubiquitous CBA promoter to permit serial in vivo bioluminescent monitoring of transgene expression. These vectors are described specifically in US Publication Number 20130096182A1 and in Smith *et al.* (see Smith, 2014). The AAVHSC-Luciferase vector was pseudotyped in the HSC7 capsid variant (the polynucleotide sequence of HSC7 capsid is provided as SEQ ID NO: 27 and the polypeptide sequence of HSC7 capsid is provided as SEQ ID NO:8 (see Figure 1)) or the HSC17 capsid variant (the polynucleotide sequence of HSC17 capsid is provided as SEQ ID NO: 35 and the polypeptide sequence of HSC17 capsid is provided as SEQ ID NO: 13 (see Figure 1)) as described in US Patent Publication Number 20130096182A1 and Smith *et al.*, to form the AAVHSC7-Luciferase vector and the AAVHSC17-Luciferase vector (see Smith, 2014) (see Chatterjee, 1992 for the standard AAV packaging method). Note that the AAVHSC-Luciferase vectors can transduce both mouse and human cells. However, in contrast to the Venus encoded in the AAVHSC-Venus vector described below, the luciferase will not integrate into AAVS1. Instead, the luciferase may integrate randomly or stay episomal.

[0166] Two to seven days after injection with the AAVHSC-Luciferase vector, the mice were injected with approximately $1\text{E}11$ - $5\text{E}11$ particles of AAVHSC-Venus vectors. Specifically, mice

that were first injected with AAVHSC7-Luciferase vector were injected with AAVHSC7-Venus vector and mice that were first injected with AAVHSC17-Luciferase vector were injected with AAVHSC17-Venus vector. The Venus donor vector used is described specifically in Examples 1 and 2 above. The donor AAV vector was designed such that the Venus transgene was promoterless and would only be expressed if it was integrated at the correct AAVS1 locus, which would be downstream from chromosomally encoded regulatory sequences (see Figures 3 and 4). Thus, any Venus transgene expression that occurred was under the control of a chromosomal promoter located in or near AAVS1. Importantly, the vector containing the Venus gene does not contain a promoter to drive expression. The Venus gene will only be expressed if it integrates correctly into AAVS1 in human cells, downstream from an endogenous chromosomal promoter. The donor vector, ITR-AAVS1-Venus (see Figure 3), was packaged into AAVHSC capsids according to the standard AAV packaging method described in Chatterjee et al, 1992. The vector was pseudotyped in the HSC7 capsid variant (the polynucleotide sequence of HSC7 capsid is provided as SEQ ID NO: 27 and the polypeptide sequence of HSC7 capsid is provided as SEQ ID NO:8 (see Figure 1)) or the HSC17 capsid variant (the polynucleotide sequence of HSC17 capsid is provided as SEQ ID NO: 35 and the polypeptide sequence of HSC17 capsid is provided as SEQ ID NO:13 (see Figure 1)) to form the AAVHSC7-Venus vector and the AAVHSC17-Venus vector.

[0167] Finally, in vivo luciferase expression was measured 4 weeks post-injection. Six weeks post-injection, engraftment of human CD34+ and CD45+ cells was measured and Venus expression was quantitated. See Figure 20 for an overall schematic of the experiments performed in this Example.

[0168] *Results.* Four weeks post-injection, in vivo imaging was performed on xenotransplanted and non-xenotransplanted mice that received intravenous injections of AAVHSC-Luciferase vectors. Results showed specific luciferase expression in vertebrae, spleen, hips, and long bones, which are all sites of hematopoiesis after transplantation (see Figure 21A). However, no specific luciferase expression in hematopoietic organs was observed in mice that were not previously xenografted with human cord blood CD34+ HSCs (see Figure 21B). These results indicate that intravenously injected AAVHSC vectors traffic to in vivo sites of human hematopoiesis and preferentially transduce stem and progenitor cells.

[0169] Six weeks after injection with the AAVHSC-Venus vectors, human CD34+ and CD45+ cells were analyzed using flow cytometry. Results indicated that the injected human cord blood CD34+ cells engrafted into the mice and gave rise to more mature blood cells. Specifically, primitive human blood progenitor cells (i.e., CD34+ cells) were observed in the bone marrow (see Table 1, CD34+ cells and femoral marrow). Additionally, human mononuclear blood cells (i.e., CD45+ cells) were evident in the femoral marrow, vertebral marrow, and spleen as shown in Table 1.

Table 1: Engraftment of Human Blood Cells in Immune Deficient Mice

AAVHSC7			
Cell type	Femoral Marrow	Vertebral Marrow	Spleen
CD45+	68.4	30.4	24.3
CD34+	22	NT*	NT
AAVHSC 17			
Cell Type	Femoral Marrow	Vertebral Marrow	Spleen
CD45+	46.6	24	18.3
CD34+	13.1	NT	NT
*NT= Not Tested			
CD45+ cells: human mononuclear blood cells			
CD34+ cells: human hematopoietic progenitor cells			

[0170] Six weeks post injection, flow cytometry was used to analyze Venus expression from human HSCs of xenotransplanted mice that received intravenous injections of either AAVHSC7-Venus or AAVHSC17-Venus vectors. Results revealed that AAVHSC transduction was readily observed in the CD45+ human HSCs as well as CD34+ human HSCs from the femoral and vertebral marrow (see Table 2 and Figures 22A-F).

Table 2: Percentage of Engrafted Human Hematopoietic Cells Expressing Venus

AAVHSC7			
Cell Type	Femoral Marrow	Vertebral Marrow	Spleen
CD45+	8.35	15.3	10.3
CD34+	9.23	NT*	NT
AAVHSC 17			
Cell Type	Femoral Marrow	Vertebral Marrow	Spleen
CD45+	8.59	70.2	9.9
CD34+	8.92	NT	NT
*NT= Not Tested			
CD45+ cells: human mononuclear blood cells			
CD34+ cells: human hematopoietic progenitor cells			

[0171] Additionally, human CD45+ cells in the spleen readily showed evidence of transduction as Venus was expressed in these cells (see Table 2 and Figures 22G and H). This demonstrates that CD45+ cells arising from the transplanted human cord blood CD34+ cells express Venus. These results indicate that intravenous injection of AAVHSC vectors in vivo results in transduction of human hematopoietic cells.

EXAMPLE 4: Insertion of large and small editing elements into a genome using AAV

clade F vectors

[0172] *Methods.* rAAV Production, Purification, and Titration. All targeting genomes were cloned into an AAV2 backbone using New England Biolabs Gibson Assembly Cloning Kit with primers designed using NEBuilder v.1.6.2 (Ipswich, MA). All targeting genomes were sequenced and AAV2 ITR integrity was confirmed using restriction digest and sequencing. Single-stranded targeting genomes were packaged into the AAVF capsids in herpes simplex virus (HSV)-infected 293 cells. The resulting recombinant AAV vectors were purified through two rounds of CsCl₂ density centrifugation gradients and titers were determined using qPCR with transgene-specific primers and probe.

[0173] K562, HepG2 and PBSC Transductions. The chronic myelogenous leukemia (CML) cell line, K562 and the hepatocellular carcinoma cell line HepG2, were obtained from American Type Culture Collection (ATCC) (Manassas, VA) and cultured according to ATCC guidelines. Peripheral blood stem cells (PBSCs) were purified from mononuclear cells from cytokine primed PB of healthy donors using CD34+ Indirect isolation kits (Miltenyi Biotech) and transductions performed immediately after isolation. PBSCs were cultured in Iscove's Modified Dulbecco's Medium (IMDM) (Irvine Scientific) containing 20% FCS, 100 ug/mL streptomycin, 100 U/mL penicillin, 2mmol/L L-glutamine, IL-3 (10 ng/mL; R&D Systems), IL-6 (10 ng/mL; R&D Systems), and stem cell factor (1 ng/mL; R&D Systems). HepG2 cells were split and plated approximately 24 hours prior to transductions. K562 cells were plated immediately prior to transductions. K562s, HepG2s or PBSCs were transduced with AAVF targeting vectors at MOIs ranging from 5E4 to 4E5. The cells were transduced and homologous recombination was achieved in the absence of an exogenous nuclease. Cells were harvested for flow and molecular analysis at time points between 1 to 39 days post transduction. BrdU labeling of in vitro transductions were performed prior to harvesting using the APC BrdU Flow Kit (BD Biosciences) as instructed.

[0174] TI PCR and Sequencing. High molecular weight DNA was isolated from K562s, HepG2s or PBSCs transduced with AAVF targeting vectors. TI specific PCR was performed using a primer that anneal to the chromosomal region outside the homology arms, Sigma AAVS1 forward primer (5' - GGC CCT GGC CAT TGT CAC TT - 3') and a primer that anneal to the inserted cassette, either Venus reverse primer (5' - AAC GAG AAG CGC GAT CAC A - 3') or RFLP HindIII reverse primer (5' - CCAATCCTGTCCCTAGTAAAGCTT - 3'). Roche Expand Hifidelity PCR system (Indianapolis, IN) was used and cycling conditions as follows: 1 cycle, 5 minutes - 95°C; 15 cycles, 30 seconds - 95°C, 30 seconds - start at 62°C and decrease by 0.5°C per cycle, 2 minutes - 68°C; 20 cycles, 30 seconds - 95°C, 30 seconds - 53°C, 2 minutes - 68°C; 1 cycle, 5 minutes - 68°C. PCR products were PCR purified for direct sequencing using Qiaquick PCR Purification Kit (Qiagen) or cloned using TOPO TA Cloning Kit for Sequencing and clones sequenced by Sanger Sequencing (Life Technologies).

[0175] Transplantation of CD34+ Cells. All animal care and experiments were performed under

protocols approved by a Institutional Animal Care and Use Committee. 6-8-week old male NOD.CB17-Prkdcscid/NCrCrI (NOD/SCID) mice were maintained in a specific pathogen free facility. Mice were placed on sulfamethoxazole and trimethoprim oral pediatric antibiotic (Hi-Tech Pharmacal (Amityville, NY), 10ml/500ml H₂O) for at least 48 hours before transplant. Mice were irradiated with a sublethal dose of 350cGy from a ¹³⁷Cs source and allowed to recover for a minimum of 4 hours prior to transplantation. Umbilical cord blood (CB) CD34⁺ cells were isolated using CD34⁺ Indirect isolation kits (Miltenyi Biotech). 1 X 10⁶ CB CD34⁺ cells were resuspended in approximately 200 ul and transplanted by tail vein injection. 2.4E11 to 6.0E11 particles of AAVF targeting vectors were injected intravenously through the tail vein at 1 or 7 weeks post CB CD34⁺ transplantation. Femoral bone marrow (BM), vertebral BM and spleen were harvested 6, 7 or 19 week post transplantation.

[0176] Flow Cytometric Analysis. In vitro transductions were analyzed for AAVF targeting vector mediated integration, and BrdU and 7-AAD using a Cyan ADP Flow Cytometer (Dako). Specific fluorescence was quantified following the subtraction of autofluorescence. In vivo expression of integrated fluorescent cassette in CD34⁺ and erythroid cells was analyzed in harvested vertebral BM, femoral BM and spleen of xenografted mice by staining with human specific antibodies, APC-conjugated anti-CD34 and PE-conjugated anti-Glycophorin A, and PE- and APC- conjugated IgG controls (BD Biosciences) on an FACS Aria SORP (BD Biosciences). Flow cytometry data was analyzed using FlowJo software (Treestar).

[0177] *Results.* Stem cell-derived AAV were shown to map to AAV clade F based on nucleotide sequence homology of the capsid genes (Figure 23, Smith et al, Mol Ther. 2014 Sep;22(9):1625-34). These stem cell-derived AAV were named AAVHSC1-17. These AAV are also referred to herein as AAVF1-17, respectively.

[0178] A singled stranded AAV vector genome was used to design a correction genome containing homology arms and a large insert (Figure 24). The insert contained a promoterless Venus open reading frame (ORF) downstream from a splice acceptor (SA) and a 2A sequence (2A) to allow for independent protein expression. Venus is a variant of yellow fluorescent protein (see, e.g., Nagai et al. Nat Biotechnol, 2002, 20(1): 87-90). The left and right homology arms (HA) were each 800 bp long and were complementary to sequences in Intron 1 of the human PPP1R12C located in the AAVS1 locus on chromosome 19 (Figure 25). A similar single stranded AAV vector genome was designed with a 10bp insert between the two homology arms (Figure 35). The AAVS1 locus is considered a safe harbor site for the insertion of heterologous transgenes.

[0179] The homology arms, the open reading frame and regulatory sequences were cloned between AAV2 inverted terminal repeats (ITRs). This correction genome was then packaged (pseudotyped) in different AAV capsids, including AAVHSC, AAV8, 9, 6 and 2. Recombinant viruses were then used to deliver the editing genome to the nuclei of target cells. Target cells tested included CD34⁺ erythroleukemia cell lines, liver cell lines and primary human CD34⁺ hematopoietic stem/progenitor cells and as well as their hematopoietic progeny.

[0180] AAVF vectors containing the Venus ORF, preceded by and flanked by homology arms complementary to Intron 1 of the human PPP1R12C genes were used to deliver the editing genome to primary human CD34+ cytokine-primed peripheral blood stem cells (Figure 26A), K562, a human CD34+ erythroleukemia cell line (Figure 26B), and HepG2, a human liver cell line (Figure 26C). Primary CD34+ cells supported the highest level of editing, up to 60% (Figures 26A-F). Immortalized cell lines, including K562 and HepG2, also showed significant levels of editing. In all cases, the level of editing achieved was consistently significantly higher than that achieved with non-Clade F viruses, including AAV6 and AAV8 (Figure 26A-F and Figure 36).

[0181] In another experiment, DNA extracted from cytokine primed CD34+ peripheral blood stem cells (PBSC) transduced with AAVF7, AAVF15 or AAVF17 vectors was amplified with a chromosome -specific primer and an insert-specific primer. The vectors included either a large insert (Venus) or a short insert (10bp, RFLP). Gels showed that correctly sized amplicons were amplified from the edited CD34 cells (Figure 27, Figure 35, and Figure 36). The presence of the 1.7 kb and 1kb bands reflected correctly targeted integration of large and small inserts, respectively. Targeted integration was shown at both short and long-term time points after editing with AAVF vectors (Figure 27).

[0182] In another experiment, single stranded AAV vector genomes were designed for the insertion of a 10 bp insert in intron 1 of the human PPP1R12C gene (Figure 28A). These vectors included a wild type left homology arm (HA-L) which contained a NheI restriction enzyme recognition site (GCTAGC). The NS mut vector, was designed to change the TA sequence in the left homology arm on chromosome 19 to AT. This change results in the conversion of an NheI site to an SphI site, changing the sequence from GCTAGC to GCATGC. Figure 28B shows the relative sizes of the expected fragments created by cutting with NheI or SphI when genomic DNA from K562 cells was edited using either the wild type or the NS Mut AAVF vectors. Actual amplicons derived from genomic DNA of K562 cells edited with a wild type AAVF vector were digested with NheI, but not with SphI, as predicted (Figure 28C). Results with amplified K562 DNA after editing with AAVF7 or an AAVF17 vectors encoding either wild type or NS Mut genomes showed that digestion with NheI no longer resulted in cleavage of the amplicon, comparable to the amplicon from unedited cells. Digestion with SphI resulted in cleavage, demonstrating that the NheI site in the left homology arm of the chromosome was replaced by an SphI site (Figure 28D). Electrophoresis of amplified DNA from a hepatocellular carcinoma cell line, HepG2, after editing with AAVF7 or an AAVF17 vectors encoding either wild type or NS Mut genomes showed that digestion with NheI no longer resulted in cleavage of the amplicon, comparable to the amplicon from unedited cells (Figure 28E). Digestion with SphI resulted in cleavage, demonstrating that the NheI site in the left homology arm of the chromosome was replaced by an SphI site. Sequence analysis confirmed editing with AAVF7 and AAVF17 Wild type or NS Mut vectors (Figure 29). These results demonstrate that both AAVF7 as well as AAVF17 successfully mediated the 2 nucleotide substitution in the chromosomal sequences in two different cell lines. These results also demonstrate the ability of AAVF vectors to mediate a 2 base pair substitution in genomic DNA of human cells, suggesting their use for correction of disease-causing mutations or

induction of new mutations in the genome.

[0183] In another experiment, the potential requirement for cell division on the editing capacity of AAVF vectors was tested on healthy human CD34⁺ PBSC. BrdU was incorporated into transduced human CD34⁺ PBSC through pulsing with 10 μ M of BrdU for 2 hours. AAVF transduced CD34⁺ cells were harvested, permeabilized and fixed prior to DNase treatment. After DNase treatment, treated cells were stained with anti-BrdU APC antibody for 20 minutes. BrdU labeling of in vitro edited cells was performed prior to harvesting using the APC BrdU Flow Kit (BD Biosciences) as per instructions. Cells were then analyzed by flow cytometry for Venus expression as well as BrdU labeling. Results revealed the similar frequencies of Venus expression in both the BrdU positive and negative populations, suggesting that cell division was not required for AAVF-mediated editing (Figure 30).

[0184] In another experiment, editing of engrafted human hematopoietic stem cells in vivo was tested by systemically delivered AAVF vectors immune deficient NOD/SCID mice engrafted with human cord blood CD34⁺ hematopoietic stem cells (Figure 31A and B). In both the marrow as well the spleen, the majority of human cells were found to express Venus, while no Venus expression was observed in mouse cells (Figure 31C, Figure 37 and Figure 38). Since the mouse genome does not contain an AAVS1 locus complementary to the homology arms, these findings demonstrate the specificity of gene targeting. Of the human cells analyzed, Venus expression was observed in primitive CD34⁺ progenitor cells, as well as mature glycophorin A⁺ erythroid cells in both the marrow as well the spleen.

[0185] In another experiment, mice were engrafted with human cord blood CD34⁺ cells and AAVF-Venus was injected by an intravenous route either 1 or 7 weeks later. Vertebral or femoral marrow or spleen was harvested either 5, 6 or 13 weeks after intravenous injection of Venus. These represented cumulative times post-transplant of 6, 7 or 20 weeks. Results reveal that intravenous injection of AAVF-Venus results in editing of both the primitive (CD34⁺) as well as the more mature, differentiated (CD45⁺) in vivo engrafted human hematopoietic cells. Cells of the human erythroid lineage demonstrated very efficient editing long-term after transplantation and injection (Figure 32). AAVF-mediated editing was found to be stable long term, and was stably inherited by the differentiated progeny of in vivo engrafted human CD34⁺ cells (Figure 32). The differentiated progeny of edited CD34⁺ cells expressed Venus long term (Figure 32).

[0186] In another experiment, sequence analysis was performed for targeted chromosomal insertion of a promoterless SA/2A venus ORF in a K562 erythroleukemia cell line, primary human cytokine-primed peripheral blood CD34⁺ cells and a HepG2 human liver cell line (Figure 33). Site-specifically integrated sequences were amplified using a chromosome-specific primer and an insert-specific primer. Results revealed precise insertion of the SA/2A Venus at the junction between the left and right homology arms in every case (Figure 33).

[0187] In another experiment, sequence analysis was performed for targeted chromosomal insertion of a 10bp insert in primary human cytokine-primed peripheral blood CD34⁺ cells and

a HepG2 human liver cell line (Figure 34). Site-specifically integrated sequences were amplified using a chromosome-specific primer and an insert-specific primer. Results revealed precise insertion of the 10 bp insert at the junction between the left and right homology arms in every case (Figure 34).

[0188] These data show that both large and short inserts can be successfully edited into a genome using AAV clade F vectors and that the integration into the genome is precise. These data also show that AAV clade F vectors could be used for high efficiency genome editing in the absence of an exogenous nuclease.

EXAMPLE 5: Editing of the PPP1R12c Locus in Human Cell Lines

Methods

[0189] To assess the editing of human cell types by AAVF vectors, the following human cell lines were used:

Cell Line	Tissue type
WI-38	normal human diploid fibroblasts
MCF7	human breast cancer cell line
Hep-G2	human hepatocellular carcinoma cell line
K562	CD34+ erythroleukemic cell line
Y79	human retinoblastoma cell line
SCID-X1 LBL	human EBV-immortalized B cell line from a SCID-X1 patient

[0190] The AAVF vectors used each contained a vector genome containing an editing element encoding a promoterless Venus reporter. The promoterless Venus contained the open reading frame (ORF) of Venus downstream from a splice acceptor (SA) and a 2A sequence (2A) to allow for independent protein expression. The left and right homology arms (HA) were each 800 bp long and were complementary to sequences in Intron 1 of human PPP1R12C located in the AAVS1 locus on chromosome 19. The AAVS1 locus is considered a safe harbor site for the insertion of heterologous transgenes. The editing element containing the homology arms, the Venus ORF, and regulatory sequences were cloned between AAV2 inverted terminal repeats (ITRs).

[0191] Cell Culture. All cell lines were grown in a humidified atmosphere of 5% CO2 at 37°C and cultured as follows: SCID-X1 lymphoblasts (Coriell) were cultured in RPMI1640 (Gibco, cat# 21875) supplemented with 15% fetal bovine serum (FBS) (Gibco, cat# 26140); K562 cells (ATCC) were cultured in DMEM (Corning, cat# 15-017-CVR) supplemented with 10% FBS (Gibco, cat# 26140) and 1% L-glutamine (Gibco, cat# 25030); HepG2 cells (ATCC) were

cultured in EMEM (ATCC, cat# 30-2003) supplemented with 10% FBS (Gibco, cat# 26140); MCF-7 cells (ATCC) were cultured in MEM (Gibco, cat# 11095) supplemented with 10% FBS (Gibco, cat# 26140), 1% MEM non-essential amino acids (Gibco, cat# 11140), 1% Sodium pyruvate (Gibco, cat# 11360) and 10 µg/ml human recombinant insulin (Gibco, cat#12585-014); WI-38 fibroblasts (ATCC) and HEK293 cells (ATCC) were cultured in MEM (Gibco, cat# 11095) supplemented with 10% FBS (Gibco, cat# 26140), 1% MEM non-essential amino acids (Gibco, cat# 11140) and 1% Sodium pyruvate (Gibco, cat# 11360); and Y79 cells (ATCC) were cultured in RPMI1640 (Gibco, cat# A10491) supplemented with 20% FBS (Gibco, cat# 26140).

[0192] AAVF-mediated editing of human cell lines. Adherent cells were seeded on day 0 at 20,000 cells/0.1 mL for a 96-well format or 20,000 cells/0.5 mL for a 24-well format. Cell counts were measured 24-h later (on day 1) prior to addition of the AAVF vectors. Suspension cells were seeded on day 1 on 20,000 cells/0.1 mL in a 96-well format or 20,000 cells/0.5 mL in 24-well plate format. On day 1, the AAVFs were added to the cells using a multiplicity of infection (MOI) of 150,000 vector genomes (VG)/cell. Before AAVF addition, the pipet tips used for transfer were coated with protamine sulfate (10 mg/ml). The AAVFs were thoroughly suspended on a vortex mixer at full speed for 30 seconds immediately before transduction. The volume of AAVF added to the cells did not exceed 5% of the total volume in the well. On day 3, cells were harvested (adherent cells were mildly trypsinized to remove them from the tissue culture plates), washed, and analyzed for Venus expression by flow cytometry using an Intellicyt flow cytometer fitted with a Hypercyt Autosampler. Editing was expressed as the percent of the total cell population that was Venus positive minus the background fluorescence observed in untransduced cells (typically less than 1% Venus positive cells). The AAVF editing experiments were carried out without the use of an exogenous nuclease.

Results

[0193] Editing of the human PPP1R12c locus was observed for all the AAVF tested and showed cell type selectivity (Table 3). In general, AAVF5 produced the highest levels of editing in each of the cell lines, from 12-45% of the total cell population, after 48 hours of infection. AAVF9 also produced high levels of gene editing in the B lymphoblast cell lines (SCID-X1 LBL and K562). AAVF17 produced the highest level of gene editing in normal human diploid fibroblasts (WI-38 cells) under these conditions. The AAVF1, AAVF4, AAVF7 vectors produced levels of editing that were consistently greater than that seen in untransduced cells but that were generally lower than the maximal levels observed with AAVF5, AAVF9, and AAVF17. These data demonstrate that the AAVFs have a broad tropism for human tissues and may be useful for gene editing in the liver, CNS (e.g., retina), tissue fibroblasts, breast, and lymphocytes among others.

Table 3. Editing of Human Cell Lines by AAVF

	Human Cell Lines					
Vector	K562	Hep-G2	WI-38	Y79	MCF7	SCID-X1 LBL
	Percent Venus Positive Cells					
AAVF1	2.06	1.92	1.39	0.7	2.71	0.57
AAVF4	1.36	2.95	6.29	0.35	3.42	0.12
AAVF5	45.32	-	11.85	12.5	20.47	45.02
AAVF7	5.78	3.83	4.87	1.42	3.6	0.94
AAVF9	18.06	-	-	2.46	6.14	6.79
AAVF17	3.18	3.71	17.6	1.29	5.66	0.44

EXAMPLE 6: AAVF Editing in Primary Human Cells

[0194] To assess the editing of primary human cells by AAVF vectors, primary cultures of human hepatocytes, hepatic sinusoidal endothelial cells, and skeletal muscle myoblasts were used.

[0195] The AAVF and AAV vectors used each packaged a vector genome encoding a promoter-less Venus reporter. The insert comprised a Venus open reading frame (ORF) downstream from a splice acceptor (SA) and a 2A sequence (2A) to allow for independent protein expression. The left and right homology arms (HA) were each 800 bp long and were complementary to sequences in Intron 1 of human PPP1R12C located in the AAVS1 locus on chromosome 19. The AAVS1 locus is considered a safe harbor site for the insertion of heterologous transgenes. The correction genome consisting of the homology arms, the open reading frame and regulatory sequences were cloned between AAV2 inverted terminal repeats (ITRs).

Methods

[0196] Cell culture. All primary human cells were cultured at 37°C, under 5% humidified CO₂ in a tissue culture incubator. All materials and media components were obtained from Life Technologies unless specified otherwise.

[0197] Primary cultures of human hepatocytes were obtained from Invitrogen and were cultured on type I collagen coated plates as suggested by the manufacturer. Cells were recovered from storage in liquid nitrogen in Thawing/Plating Medium [32.5 mL William's E Medium (#A12176) 1.6 mL fetal bovine serum, 3.2 uL of 10 mM Dexamethazone and 0.9 mL of Plating Cocktail A per 35.0 mL final volume]. Plating cocktail A consisted of 0.5 mL Penicillin (10,000 U/mL)/Streptomycin (10,000 ug/mL) solution (Cat # 15140), 0.05 mL of 4.0 mg human

recombinant insulin/mL (Catalog # 12585-014), 0.5 mL of 200 mM GlutaMAX™ solution (Catalog # 35050) and 0.75 mL of 1.0M Hepes, pH7.4 (Catalog # 15630) per 1.8 mL final volume. Human hepatocytes were maintained in Maintenance media which contained 100 mL of Williams E Medium, 0.001 mL of dexamethasone, and 3.6 mL of Maintenance Cocktail B (Catalog # A13448) per 103.6 mL final volume.

[0198] Primary cultures of human skeletal muscle myoblasts were obtained from Lonza and were cultured in SkGM™ medium as described by the manufacturer.

[0199] SkGM™-2 Bullit™ Kit (Lonza, Catalog No. CC-3245) contained 0.5 mL human Epidermal Growth Factor [hEGF] (#0000482653), 0.5 mL Dexamethasone (#0000474738), 10 mL L-glutamine (#0000474740), 50 mL Fetal Bovine Serum (#0000474741), 0.5 mL Gentamicin/Amphotericin-B [GA] (#0000474736), in 500 mL SkGM-2 medium (#0000482653).

[0200] Primary cultures of human hepatic sinusoidal endothelial cells were purchased from Creative Bioarray and were cultured in SuperCult® Endothelial Cell Medium and grown on a gelatin-based coating as described by the manufacturer. SuperCult® Endothelial Cell Medium Supplement Kit (Catalog # ECM-500 Kit) contained 0.5 mL VEGF (#15206), 0.5 mL Heparin (#15250), 0.5 mL EGF (#15217), 0.5 mL FGF (#15204), 0.5 mL Hydrocortisone (#15318), 5.0 mL Antibiotic-Antimycotic Solution (#15179), 5.0 mL L-glutamine (#15409), 10.0 mL Endothelial Cell Supplement (#15604), 50.0 mL FBS (#15310), and 500.0 mL of SuperCult® Endothelial Cell Medium (#15517).

[0201] AAVF-mediated editing of primary human cells. Human primary hepatocytes were seeded on day 0 on 2×10^4 cells/0.1 mL for the 96-well format or 2×10^4 cells/0.5 mL for the 24-well format. Viral vectors were added 48 h later on day 2 as described below. Human skeletal muscle myoblasts, and human hepatic sinusoidal endothelial cells were seeded on day 1 at 2×10^4 cells/0.1 mL in a 96-well format or 2×10^4 cells/0.5 mL in a 24-well format. On day 2, the viral vectors were added to these cells at a multiplicity of infection (MOI) of 150,000 VG(Vector genomes)/cell (For AAVF5 an MOI of 5×10^4 VG/cell was used and for AAVF17 an MOI of 2.5×10^4 VG/cell was used). Prior to the addition of vector to the cells, all pipet tips used for vector transfer were coated with protamine sulfate (10 mg/mL) and the vectors were thoroughly mixed by vortexing for 30 seconds immediately before transduction. The volume of vector added to the cells did not exceed 5% of the total volume in the well.

[0202] The culture media for the human primary hepatocytes was refreshed on day 3. On day 4, cells were harvested (adherent cells were trypsinized) and analyzed for Venus expression by flow cytometry using an Intellicyt flow cytometer fitted with a Hypercyt Autosampler. Editing was expressed as the percent of the total cell population that was Venus positive minus the background fluorescence observed in un-transduced cells (typically less than 1% Venus positive cells).

Results

[0203] Editing of the human PPP1R12c locus was observed for all the AAVF tested and showed cell type selectivity (Table 4). In general, AAVF5 gave the highest levels of editing in each of the primary cell populations, from a low of 2% in human hepatocytes to 24% to 35% in primary skeletal myoblasts and hepatic sinusoidal endothelial cells, respectively, after 48-h of infection. AAVF7 and AAVF17 also produced high levels of gene editing in the primary hepatic sinusoidal endothelial cells and skeletal myoblasts. Levels of editing with the AAVF vectors in these cells were 10- to 50-fold higher than that observed with AAV2 or AAV6. The AAVF vectors produced levels of editing that were consistently greater than that seen in untransduced cells or cells transduced with a AAV2 or AAV6 packaging the promoter-less Venus vector genome. Little or no editing was observed for either AAV2 or AAV6 in these cells. These data in primary human cells demonstrate that AAVF5, AAVF7, and AAVF17 have a broad tropism for human tissues and may be useful for gene therapy applications directed towards the liver, skeletal muscle, and endothelial cell populations, among others.

[0204] For comparison, each of the primary human cells populations were also transduced with AAVF gene transfer vectors packaging mCherry under control of the chicken beta actin (CBA) promoter. The ratios of protein expression of Venus /mCherry were then used to estimate an editing ratio - reflecting the ratio of number of cells having detectable protein expression in the various cell types using gene editing vectors of the experiment (Venus) to number of cells having detectable protein expression using the aforementioned gene transfer vectors (mCherry) - for AAV6, AAVF5, and AAVF7. As shown in Table 5, AAVF-mediated gene editing is generally more effective for protein expression than the corresponding AAVF-mediated gene transfer approach in the studied primary human cells, whereas AAV6-mediated gene editing was substantially the same as or slightly less effective than AAV6-mediated gene transfer in such cells. Notably, as also shown in Table 5, AAVF-mediated gene editing was higher than that observed with AAV6-mediated gene editing for the studied primary human cells. These data demonstrate that AAVF-mediated gene editing is more efficient than AAV6 in a variety of primary human cells, and that AAVF-mediated gene editing compares favorably over corresponding gene-transfer approaches in such cells.

[0205] The AAVF vectors packaging the mCherry reporter also effectively transduced primary human cells and showed cell type specificity (Table 6). For human umbilical vein endothelial cells (HUVEC) and hepatic sinusoidal endothelial cells (HSEC), transduction with AAVF9 produced 38-50% mCherry positive cells after 48-h of infection. AAVF9 also efficiently transduced human skeletal muscle myoblasts and to a lesser extent, all of the AAVF tested effectively transduced the HSEC under these conditions (Table 6). Without being bound by theory, it is likely that most, if not all, of the mCherry expression in these studies represents episomal expression of the reporter as no homology arms were present in the mCherry vector genomes with expression was driven by the CBA promoter.

Table 4. AAVF-Mediated Editing of Primary Human Cells

Vector	HSEC	Hepatocytes	SMM
Percent Venus Positive Cells			
AAV2	0.34	0.06	0.52
AAV6	0.00	0.00	0.00
AAVF5	34.70	2.00	24.20
AAVF7	18.40	0.62	18.65
AAVF17	21.40	1.37	19.60

Table 5. Editing ratios of AAVF in primary human cells

Vector	Hepatocytes		SMM	
	Venus/mCherry	AAVF/AAV6	Venus/mCherry	AAVF/AAV6
AAV6	0.84		0.80	
AAVF5	9.42	11.21	14.32	17.90
AAVF7	0.95	1.10	13.05	16.30

Vector	HSEC	
	Venus/mCherry	AAVF/AAV6
AAV6	0.60	
AAVF5	5.30	8.80
AAVF7	2.70	4.50

HSEC= hepatic sinusoidal endothelial cells

SMM = skeletal muscle myoblasts

Table 6. Transduction of Primary Human Cells with AAVF Vectors Packaging mCherry

Vector	HSEC	Hepatocytes	SMM	HUVEC
Percent mCherry positive cells				
AAVF1	7.04	0.13	1.65	1.47
AAVF4	9.10	0.05	1.72	0.61
AAVF5	4.00	0.01	1.03	0.71
AAVF7	4.43	0.70	0.77	2.15
AAVF9	38.24	1.05	29.27	50.13

HSEC= hepatic sinusoidal endothelial cells
SMM = skeletal muscle myoblasts
HUVEC = human umbilical vein endothelial cells

EXAMPLE 7: Study of AAVF relative gene-editing (HDR) versus gene-transfer (transduction) efficiencies

[0206] Two types of AAV-based vectors (CBA-mCherry gene transfer vectors, and AAVS1-Venus gene editing vectors, Figure 39) were used to assess the relative gene-editing efficiency versus the gene-transfer transduction efficiency of AAVF vectors, as well as AAV2 and AAV6,

as measured by relative protein expression. For the gene-transfer vectors, CBA-mCherry construct included the mCherry gene under the control of the chicken beta actin (CBA) promoter, and utilized a polyadenylation signal, but did not contain any homology arms. For the gene-editing vectors, AAVS1-Venus construct included the promoterless Venus open reading frame (ORF), with (HA-L) and right (HA-R) homology arms targeting the Venus ORF to Intron 1 of the PPP1R12C gene within the AAVS1 region of chromosome 19. There was also a splice acceptor (SA) and a 2A sequence upstream of the Venus ORF, which allowed the Venus transcript to be spliced out and expressed independent of the PPP1R12C gene.

[0207] The ability of AAVF, AAV2 and AAV6 vectors to mediate gene editing versus gene transfer was compared in primary human cord blood CD34+ hematopoietic stem/progenitor cells. The cells used were pooled from multiple donors. Purified CD34+ cells were transduced with either gene transfer vectors (AAV*-CBA-mCherry) or gene editing vectors (AAV*-Venus) at a multiplicity of 150,000 vector genomes (VG) per cell. Forty eight hours later, cells were harvested and analyzed by flow cytometry (Figure 40A and 40B). The data shown includes subtraction of background untransduced cells.

[0208] It was hypothesized that mCherry expression would represent transduction efficiency, whereas Venus expression would represent gene editing efficiency. Without being bound by theory, the following is a summary of the potential mechanism of AAV transduction versus editing. Upon infection, AAV binds to cell surface receptors and is internalized prior to nuclear translocation and entry. In the nucleus, the AAV undergoes uncoating and vector genomes are released. These processes likely occur at the same rate for each given capsid in the same cell population, regardless of the vector genome. Following uncoating, the single stranded CBA-mCherry genome undergoes second strand synthesis prior to mCherry expression. On the other hand, the promoterless Venus editing vector is directed to the genomic region of complementarity on Chromosome 19. The SA/2A-Venus cassette which is bounded by the homology arms may then recombine into the chromosome at the internal junction of the homology arms via homology dependent repair (HDR) mechanisms. Following this recombination event, Venus is expressed in the successfully edited cells.

[0209] Venus was expressed in a much higher proportion of cells than mCherry for all vectors except AAVF9 (Figure 40A and 40C). Venus expression was especially high following transduction with AAVF1, AAVF5, AAVF7, AAVF16 and AAVF17. The same capsids led to much lower levels of mCherry expression (Figure 40A and 40C). The least amount of both mCherry and Venus expression was observed following transduction with AAV2 and AAV6. A comparison of the relative expression of Venus to mCherry was also performed. Specifically, an editing ratio - reflecting the ratio of number of cells having detectable protein expression in the various cell types using gene editing vectors of the experiment (Venus) to the number of cells having detectable protein expression using the aforementioned gene transfer vectors (mCherry) - was determined. The editing ratio provides an estimate of the relative efficiency of editing mediated by each AAV capsid after normalization for the processes of virus entry through uncoating within the nucleus. All AAV vectors tested exhibited more efficient gene editing (Venus) as compared with gene transfer / transgene expression (mCherry), except for

AAVF9 (Figure 40D and Table 7). AAVF5 and AAVF7 displayed the highest editing ratio (Figure 40D and Table 7). AAVF-mediated gene editing (Venus) was also compared relative to AAV 2- and AAV6-mediated gene editing (Venus) (Table 7, which shows the Venus:mCherry ratio of AAVF divided by the same ratio for either AAV2 or AAV6). The gene-editing effectiveness of AAVF gene-editing constructs compared favorably relative to AAV2- and AAV6-gene editing constructs. The editing ratios of AAVF5 and AAVF7 were the highest of the vectors compared, indicating that these vectors in particular mediate highly efficient editing.

Table 7. Editing-to-Transduction ratio in CD34+ CB Cells

Vector	Ratio editing :transduction (Venus:mCherry)	Ratio AAVF editing: AAV6 editing	Ratio AAVF editing:AAV2 editing
AAVF1	4.32	1.23	3.02
AAVF4	3.42	0.98	0.98
AAVF5	16.60	4.74	11.62
AAVF7	14.66	4.19	10.26
AAVF9	0.73	0.21	0.51
AAV2	1.43	0.41	1.00
AAV6	3.50	1.00	2.45

REFERENCES

[0210]

1. BAINBRIDGE, J.W., SMITH, A.J., BARKER, S.S., ROBBIE, S., HENDERSON, R., BALAGGAN, K., VISWANATHAN, A., HOLDER, G.E., STOCKMAN, A., TYLER, N., PETERSEN-JONES, S., BHATTACHARYA, S.S., THRASHER, A.J., FITZKE, F.W., CARTER, B.J., RUBIN, G.S., MOORE, A.T., and ALI, R.R. (2008). Effect of gene therapy on visual function in Leber's congenital amaurosis. *N Engl J Med* 358, 2231-2239.
2. BATCHU, R.B., SHAMMAS, M.A., WANG, J.Y., FREEMAN, J., ROSEN, N., and MUNSHI, N.C. (2002). Adeno-associated virus protects the retinoblastoma family of proteins from adenoviral-induced functional inactivation. *Cancer Res* 62, 2982-2985.
3. BELL, P., WANG, L., LEBHERZ, C., FLIEDER, D.B., BOVE, M.S., WU, D., GAO, G.P., WILSON, J.M., and WIVEL, N.A. (2005). No evidence for tumorigenesis of AAV vectors in a large-scale study in mice. *Mol Ther* 12, 299-306.
4. BERNS, K.I., and GIRAUD, C. (1996). Biology of adeno-associated virus. *Curr Top Microbiol Immunol* 218, 1-23.
5. BIFFI, A., and CESANI, M. (2008). Human hematopoietic stem cells in gene therapy: preclinical and clinical issues. *Curr Gene Ther* 8, 135-146.
6. BRANTLY, M.L., CHULAY, J.D., WANG, L., MUELLER, C., HUMPHRIES, M., SPENCER, L.T., ROUHANI, F., CONLON, T.J., CALCEDO, R., BETTS, M.R., SPENCER,

- C., BYRNE, B.J., WILSON, J.M., and FLOTTE, T.R. (2009). Sustained transgene expression despite T lymphocyte responses in a clinical trial of rAAV1-AAT gene therapy. *Proc Natl Acad Sci U S A*.
7. 7. CHATTERJEE, S., JOHNSON, P.R., and WONG, K.K., JR. (1992). Dual-target inhibition of HIV-1 in vitro by means of an adeno-associated virus antisense vector. *Science* 258, 1485-1488.
8. 8. CHATTERJEE, S., LI, W., WONG, C.A., FISHER-ADAMS, G., LU, D., GUHA, M., MACER, J.A., FORMAN, S.J., and WONG, K.K., JR. (1999). Transduction of primitive human marrow and cord blood-derived hematopoietic progenitor cells with adeno-associated virus vectors. *Blood* 93, 1882-1894.
9. 9. CHATTERJEE, S., WONG, K.K. (1993). Adeno-Associated Viral Vectors for the Delivery of Antisense RNA. *METHODS -LONDON- A COMPANION TO METHODS IN ENZYMOLOGY-* 5, 1.
10. 10. CIDECIYAN, A.V., HAUSWIRTH, W.W., ALEMAN, T.S., KAUSHAL, S., SCHWARTZ, S.B., BOYE, S.L., WINDSOR, E.A., CONLON, T.J., SUMAROKA, A., PANG, J.J., ROMAN, A.J., BYRNE, B.J., and JACOBSON, S.G. (2009). Human RPE65 gene therapy for Leber congenital amaurosis: persistence of early visual improvements and safety at 1 year. *Hum Gene Ther* 20, 999-1004.
11. 11. EINERHAND, M.P., ANTONIOU, M., ZOLOTUKHIN, S., MUZYCZKA, N., BERNS, K.I., GROSVELD, F., and VALERIO, D. (1995). Regulated high-level human beta-globin gene expression in erythroid cells following recombinant adeno-associated virus-mediated gene transfer. *Gene Ther* 2, 336-343.
12. 12. FISHER-ADAMS, G., WONG, K.K., JR., PODSAKOFF, G., FORMAN, S.J., and CHATTERJEE, S. (1996). Integration of adeno-associated virus vectors in CD34+ human hematopoietic progenitor cells after transduction. *Blood* 88, 492-504.
13. 13. FLOTTE, T.R., BRANTLY, M.L., SPENCER, L.T., BYRNE, B.J., SPENCER, C.T., BAKER, D.J., and HUMPHRIES, M. (2004). Phase I trial of intramuscular injection of a recombinant adeno-associated virus alpha 1-antitrypsin (rAAV2-CB-hAAT) gene vector to AAT-deficient adults. *Hum Gene Ther* 15, 93-128.
14. 14. GAO, G., VANDENBERGHE, L.H., ALVIRA, M.R., LU, Y., CALCEDO, R., ZHOU, X., and WILSON, J.M. (2004). Clades of Adeno-associated viruses are widely disseminated in human tissues. *J Virol* 78, 6381-6388.
15. 15. HACEIN-BEY-ABINA, S., VON KALLE, C., SCHMIDT, M., LE DEIST, F., WULFFRAAT, N., MCINTYRE, E., RADFORD, I., VILLEVAL, J.L., FRASER, C.C., CAVAZZANA-CALVO, M., and FISCHER, A. (2003). A serious adverse event after successful gene therapy for X-linked severe combined immunodeficiency. *N Engl J Med* 348, 255-256.
16. 16. HAN, Z., ZHONG, L., MAINA, N., HU, Z., LI, X., CHOUTHAI, N.S., BISCHOF, D., WEIGEL-VAN AKEN, K.A., SLAYTON, W.B., YODER, M.C., and SRIVASTAVA, A. (2008). Stable integration of recombinant adeno-associated virus vector genomes after transduction of murine hematopoietic stem cells. *Hum Gene Ther* 19, 267-278.
17. 17. JAYANDHARAN, G.R., ZHONG, L., LI, B., KACHNIARZ, B., and SRIVASTAVA, A. (2008). Strategies for improving the transduction efficiency of single-stranded adeno-associated virus vectors in vitro and in vivo. *Gene Ther* 15, 1287-1293.
18. 18. KAPLITT, M.G., FEIGIN, A., TANG, C., FITZSIMONS, H.L., MATTIS, P., LAWLOR,

- P.A., BLAND, R.J., YOUNG, D., STRYBING, K., EIDELBERG, D., and DURING, M.J. (2007). Safety and tolerability of gene therapy with an adeno-associated virus (AAV) borne GAD gene for Parkinson's disease: an open label, phase I trial. *Lancet* 369, 2097-2105.
19. **19.** KELLS, A.P., HADACZEK, P., YIN, D., BRINGAS, J., VARENIKA, V., FORSAYETH, J., and BANKIEWICZ, K.S. (2009). Efficient gene therapy-based method for the delivery of therapeutics to primate cortex. *Proc Natl Acad Sci U S A* 106, 2407-2411.
 20. **20.** KESSLER, P.D., PODSAKOFF, G.M., CHEN, X., MCQUISTON, S.A., COLOSI, P.C., MATELIS, L.A., KURTZMAN, G.J., and BYRNE, B.J. (1996). Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein. *Proc Natl Acad Sci U S A* 93, 14082-14087.
 21. **21.** MANNO, C.S., CHEW, A.J., HUTCHISON, S., LARSON, P.J., HERZOG, R.W., ARRUDA, V.R., TAI, S.J., RAGNI, M.V., THOMPSON, A., OZELO, M., COUTO, L.B., LEONARD, D.G., JOHNSON, F.A., MCCLELLAND, A., SCALLAN, C., SKARSGARD, E., FLAKE, A.W., KAY, M.A., HIGH, K.A., and GLADER, B. (2003). AAV-mediated factor IX gene transfer to skeletal muscle in patients with severe hemophilia B. *Blood* 101, 2963-2972.
 22. **22.** MCCORMACK, M.P., and RABBITS, T.H. (2004). Activation of the T-cell oncogene LMO2 after gene therapy for X-linked severe combined immunodeficiency. *N Engl J Med* 350, 913-922.
 23. **23.** MILLER, D.G., ADAM, M.A., and MILLER, A.D. (1990). Gene transfer by retrovirus vectors occurs only in cells that are actively replicating at the time of infection. *Mol Cell Biol* 10, 4239-4242.
 24. **24.** PAZ, H., WONG, C.A., LI, W., SANTAT, L., WONG, K.K., and CHATTERJEE, S. (2007). Quiescent subpopulations of human CD34-positive hematopoietic stem cells are preferred targets for stable recombinant adeno-associated virus type 2 transduction. *Hum Gene Ther* 18, 614-626.
 25. **25.** PETERS-SILVA, H., DINCULESCU, A., LI, Q., MIN, S.H., CHIODO, V., PANG, J.J., ZHONG, L., ZOLOTUKHIN, S., SRIVASTAVA, A., LEWIN, A.S., and HAUSWIRTH, W.W. (2009). High-efficiency transduction of the mouse retina by tyrosine-mutant AAV serotype vectors. *Mol Ther* 17, 463-471.
 26. **26.** PODSAKOFF, G., WONG, K.K., JR., and CHATTERJEE, S. (1994). Efficient gene transfer into nondividing cells by adeno-associated virus-based vectors. *J Virol* 68, 5656-5666.
 27. **27.** PONNAZHAGAN, S., YODER, M.C., and SRIVASTAVA, A. (1997). Adeno-associated virus type 2-mediated transduction of murine hematopoietic cells with long-term repopulating ability and sustained expression of a human globin gene in vivo. *J Virol* 71, 3098-3104.
 28. **28.** RAJ, K., OGSTON, P., and BEARD, P. (2001). Virus-mediated killing of cells that lack p53 activity. *Nature* 412, 914-917.
 29. **29.** SANTAT, L., PAZ, H., WONG, C., LI, L., MACER, J., FORMAN, S., WONG, K.K., and CHATTERJEE, S. (2005). Recombinant AAV2 transduction of primitive human hematopoietic stem cells capable of serial engraftment in immune-deficient mice. *Proc Natl Acad Sci U S A* 102, 11053-11058.

30. **30.** SRIVASTAVA, A. (2004). Gene delivery to human and murine primitive hematopoietic stem and progenitor cells by AAV2 vectors. *Methods Mol Biol* 246, 245-254.
31. **31.** TOWNE, C., SCHNEIDER, B.L., KIERAN, D., REDMOND, D.E., JR., and AEBISCHER, P. (2009). Efficient transduction of non-human primate motor neurons after intramuscular delivery of recombinant AAV serotype 6. *Gene Ther.*
32. **32.** ZHONG, L., CHEN, L., LI, Y., QING, K., WEIGEL-KELLEY, K.A., CHAN, R.J., YODER, M.C., and SRIVASTAVA, A. (2004a). Self-complementary adeno-associated virus 2 (AAV)-T cell protein tyrosine phosphatase vectors as helper viruses to improve transduction efficiency of conventional single-stranded AAV vectors in vitro and in vivo. *Mol Ther* 10, 950-957.
33. **33.** ZHONG, L., LI, B., JAYANDHARAN, G., MAH, C.S., GOVINDASAMY, L., AGBANDJEMCKENNA, M., HERZOG, R.W., WEIGEL-VAN AKEN, K.A., HOBBS, J.A., ZOLOTUKHIN, S., MUZYCZKA, N., and SRIVASTAVA, A. (2008a). Tyrosine phosphorylation of AAV2 vectors and its consequences on viral intracellular trafficking and transgene expression. *Virology* 381, 194-202.
34. **34.** ZHONG, L., LI, B., MAH, C.S., GOVINDASAMY, L., AGBANDJE-MCKENNA, M., COOPER, M., HERZOG, R.W., ZOLOTUKHIN, I., WARRINGTON, K.H., JR., WEIGEL-VAN AKEN, K.A., HOBBS, J.A., ZOLOTUKHIN, S., MUZYCZKA, N., and SRIVASTAVA, A. (2008b). Next generation of adeno-associated virus 2 vectors: point mutations in tyrosines lead to high-efficiency transduction at lower doses. *Proc Natl Acad Sci U S A* 105, 7827-7832.
35. **35.** ZHONG, L., LI, W., YANG, Z., QING, K., TAN, M., HANSEN, J., LI, Y., CHEN, L., CHAN, R.J., BISCHOF, D., MAINA, N., WEIGEL-KELLEY, K.A., ZHAO, W., LARSEN, S.H., YODER, M.C., SHOU, W., and SRIVASTAVA, A. (2004b). Impaired nuclear transport and uncoating limit recombinant adeno-associated virus 2 vector-mediated transduction of primary murine hematopoietic cells. *Hum Gene Ther* 15, 1207-1218.
36. **36.** ZHONG, L., ZHAO, W., WU, J., LI, B., ZOLOTUKHIN, S., GOVINDASAMY, L., AGBANDJEMCKENNA, M., and SRIVASTAVA, A. (2007). A dual role of EGFR protein tyrosine kinase signaling in ubiquitination of AAV2 capsids and viral second-strand DNA synthesis. *Mol Ther* 15, 1323-1330.
37. **37.** ZHOU, S.Z., BROXMEYER, H.E., COOPER, S., HARRINGTON, M.A., and SRIVASTAVA, A. (1993). Adeno-associated virus 2-mediated gene transfer in murine hematopoietic progenitor cells. *Exp Hematol* 21, 928-933.
38. **38.** KHAN, I.G., HIRATA, R.K., RUSSELL, D.W. (2011) AAV-mediated gene targeting methods for human cells. *Nat Protoc.* 4, 482-501.
39. **39.** KHOTIN, R.M., LINDEN, R.M., and BERNIS, K.I. (1992). Characterization of a preferred site on human chromosome 19q for integration of adeno-associated virus DNA by non-homologous recombination. *EMBO* 13, 5071-8.
40. **40.** GIRAUD, C., WINOCOUR, E., and BERNIS, K.I. (1994). Site-specific integration by adeno-associated virus is directed by a cellular DNA sequence. *Proc. Natl Acad Sci USA* 21, 10039-43.
41. **41.** LINDEN, R.M., WARD, P., GIRAUD, C., WINOCOUR, E., and BERNIS, K.I. (1996A). Site-specific integration by adeno-associated virus. *Proc. Natl Acad Sci USA* 21, 11288-94.

42. **42.** LINDEN, R.M., WINOCOUR, E., and BERNIS, K.I. (1996B). The recombination signals for adeno-associated virus site-specific integration. *Proc. Natl Acad Sci USA* 15, 7966-72.
43. **43.** XU, L., O'MALLEY, T., SANDS, M.S., WANG, B., MEYERROSE, T., HASKINS, M.E., and PONDER, K.P (2004). In Vivo Transduction of Hematopoietic Stem Cells After Neonatal Intravenous Injection of an Amphotrophic Retroviral Vector in Mice. *Mol. Ther.* Jul; 10(1):37-44.
44. **44.** WANG, C.X., SATHER, B.D., WANG, X., ADAIR, J., KHAN, I., SINGH, S., LANG, S., ADAMS, A., CURINGA, G., KIEM, H-P., MIAO, C.H., RAWLINGS, D.J., and TORBETT, B.E. (2014). Rapamycin relieves lentiviral vector transduction resistance in human and mouse hematopoietic stem cells. *Blood.* Aug; 124(6):913-23.
45. **45.** CARBONARO, D.A., JIN, X., PETERSEN, D., WANG, X., DOREY, F., KIL, K.S., ALDRICH, M., BLACKBURN, M.R., KELLEMS, R.E., and KOHN, D.B. (2006) In Vivo Transduction by Intravenous Injection of a Lentiviral Vector Expressing Human ADA into Neonatal ADA Gene Knockout Mice: A Novel Form of Enzyme Replacement Therapy for ADA Deficiency. *Mol. Ther.* Jun; 13(6): 1110-20.
46. **46.** SMITH, L.J., UL-HASAN, T., CARVAINES, S.K., VAN VLIET, K., YANG, E., WONG, JR, K.K., AGBANDJE-MCKENNA, M., and CHATTERJEE, S. (2014) Gene Transfer Properties and Structural Modeling of Human Stem Cell-derived AAV. *Mol. Ther.* Sept; 22(9):1625-1634.

SEQUENCE LISTING

[0211]

<110> CITY OF HOPE CHATTERJEE, Saswati SMITH, Laura WONG, Kamehameha

<120> ADENO-ASSOCIATED VIRUS VECTOR VARIANTS FOR HIGH EFFICIENCY GENOME EDITING AND METHODS THEREOF

<130> 054435-8138.WO00

<150> US 62/209,862

<151> 2015-08-25

<150> US 62/063,587

<151> 2014-10-14

<150> US 62/054,899

<151> 2014-09-24

<160> 40

<170> PatentIn version 3.5

<210> 1

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> adeno-associated AAV9

<400> 1

Met	Ala	Ala	Asp	Gly	Tyr	Leu	Pro	Asp	Trp	Leu	Glu	Asp	Asn	Leu	Ser
1				5					10					15	

Glu	Gly	Ile	Arg	Glu	Trp	Trp	Ala	Leu	Lys	Pro	Gly	Ala	Pro	Gln	Pro
			20					25					30		

Lys	Ala	Asn	Gln	Gln	His	Gln	Asp	Asn	Ala	Arg	Gly	Leu	Val	Leu	Pro
		35					40					45			

Gly	Tyr	Lys	Tyr	Leu	Gly	Pro	Gly	Asn	Gly	Leu	Asp	Lys	Gly	Glu	Pro
	50					55					60				

Val	Asn	Ala	Ala	Asp	Ala	Ala	Ala	Leu	Glu	His	Asp	Lys	Ala	Tyr	Asp
65					70					75					80

Gln	Gln	Leu	Lys	Ala	Gly	Asp	Asn	Pro	Tyr	Leu	Lys	Tyr	Asn	His	Ala
			85						90					95	

Asp	Ala	Glu	Phe	Gln	Glu	Arg	Leu	Lys	Glu	Asp	Thr	Ser	Phe	Gly	Gly
		100						105						110	

Asn	Leu	Gly	Arg	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Leu	Leu	Glu	Pro
		115					120					125			

Leu	Gly	Leu	Val	Glu	Glu	Ala	Ala	Lys	Thr	Ala	Pro	Gly	Lys	Lys	Arg
	130					135					140				

Pro	Val	Glu	Gln	Ser	Pro	Gln	Glu	Pro	Asp	Ser	Ser	Ala	Gly	Ile	Gly
145					150					155					160

Lys	Ser	Gly	Ala	Gln	Pro	Ala	Lys	Lys	Arg	Leu	Asn	Phe	Gly	Gln	Thr
			165						170					175	

Gly	Asp	Thr	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Ile	Gly	Glu	Pro	Pro
		180						185					190		

Ala	Ala	Pro	Ser	Gly	Val	Gly	Ser	Leu	Thr	Met	Ala	Ser	Gly	Gly	Gly
		195					200					205			

Ala	Pro	Val	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Ser	Ser
	210					215					220				

Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Gln	Trp	Leu	Gly	Asp	Arg	Val	Ile
225					230					235					240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255
 Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270
 Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285
 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300
 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320
 Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350
 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asn Asn Val Asn Ala Asn Lys Val Met Ile

Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
545 550 555 560

Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
565 570 575

Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
580 585 590

Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 2

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 2

Met Thr Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245 250 255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290 295 300

Asn Trp Gly Phe Arg Pro Lys Gln Leu Asn Phe Lys Leu Phe Asn Ile
305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
325 330 335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
355 360 365

Ala	Asp	Val	Phe	Met	Ile	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	Asp	370	375	380	
Gly	Ser	Gln	Ala	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	385	390	395	400
Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Gln	Phe	Ser	Tyr	Glu	405	410	415	
Phe	Glu	Asn	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu	420	425	430	
Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	Ser	435	440	445	
Lys	Thr	Ile	Asn	Gly	Ser	Gly	Gln	Asn	Gln	Gln	Thr	Leu	Lys	Phe	Ser	450	455	460	
Val	Ala	Gly	Pro	Ser	Asn	Met	Ala	Val	Gln	Gly	Arg	Asn	Tyr	Ile	Pro	465	470	475	480
Gly	Pro	Ser	Tyr	Arg	Gln	Gln	Arg	Val	Ser	Thr	Thr	Val	Thr	Gln	Asn	485	490	495	
Asn	Asn	Ser	Glu	Phe	Ala	Trp	Pro	Gly	Ala	Ser	Ser	Trp	Ala	Leu	Asn	500	505	510	
Gly	Arg	Asn	Ser	Leu	Met	Asn	Pro	Gly	Pro	Ala	Met	Ala	Ser	His	Lys	515	520	525	
Glu	Gly	Glu	Asp	Arg	Phe	Phe	Pro	Leu	Ser	Gly	Ser	Leu	Ile	Phe	Gly	530	535	540	
Lys	Gln	Gly	Thr	Gly	Arg	Asp	Asn	Val	Asp	Ala	Asp	Lys	Val	Met	Ile	545	550	555	560
Thr	Asn	Glu	Glu	Glu	Ile	Lys	Thr	Thr	Asn	Pro	Val	Ala	Thr	Glu	Ser	565	570	575	
Tyr	Gly	Gln	Val	Ala	Thr	Asn	His	Gln	Ser	Ala	Gln	Ala	Gln	Ala	Gln	580	585	590	
Thr	Gly	Trp	Val	Gln	Asn	Gln	Gly	Ile	Leu	Pro	Gly	Met	Val	Trp	Gln	595	600	605	
Asp	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp	Ala	Lys	Ile	Pro	His	610	615	620	
Thr	Asp	Gly	Asn	Phe	His	Pro	Ser	Pro	Leu	Met	Gly	Gly	Phe	Gly	Met	625	630	635	640
Lys	His	Pro	Pro	Pro	Gln	Ile	Leu	Ile	Lys	Asn	Thr	Pro	Val	Pro	Ala	645	650	655	

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 3

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 3

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr

165										170			175			
Gly	Asp	Thr	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Ile	Gly	Glu	Pro	Pro	
			180					185					190			
Ala	Ala	Pro	Ser	Gly	Val	Gly	Ser	Leu	Thr	Met	Ala	Ser	Gly	Gly	Gly	
		195					200					205				
Ala	Pro	Val	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Ser	Ser	
	210					215					220					
Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Gln	Trp	Leu	Gly	Asp	Arg	Val	Ile	
225					230					235					240	
Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu	
				245					250					255		
Tyr	Lys	Gln	Ile	Ser	Asn	Ser	Thr	Ser	Gly	Gly	Ser	Ser	Asn	Asp	Asn	
			260					265					270			
Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg	
		275					280					285				
Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn	
	290					295					300					
Asn	Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	Lys	Leu	Phe	Asn	Ile	
305					310					315					320	
Gln	Val	Lys	Glu	Val	Thr	Asp	Asn	Asn	Gly	Val	Lys	Thr	Ile	Ala	Asn	
					325					330					335	
Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr	Asp	Ser	Asp	Tyr	Gln	Leu	
			340					345					350			
Pro	Tyr	Val	Leu	Gly	Ser	Ala	His	Glu	Gly	Cys	Leu	Pro	Pro	Phe	Pro	
		355					360					365				
Ala	Asp	Val	Phe	Met	Ile	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	Asp	
	370					375					380					
Gly	Ser	Gln	Ala	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe	
385					390					395					400	
Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Gln	Phe	Ser	Tyr	Glu	
				405					410					415		
Phe	Glu	Asn	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu	
			420					425					430			
Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	Ser	
		435					440					445				
Lys	Thr	Ile	Asn	Gly	Ser	Gly	Gln	Asn	Gln	Gln	Thr	Leu	Lys	Phe	Ser	
	450					455					460					

Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Gly Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700
 Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Gly Gly Val
 705 710 715 720
 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

<210> 4

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 4

Met	Ala	Ala	Asp	Gly	Tyr	Leu	Pro	Asp	Trp	Leu	Glu	Asp	Asn	Leu	Ser
1				5					10					15	

Glu	Gly	Ile	Arg	Glu	Trp	Trp	Ala	Leu	Lys	Pro	Gly	Ala	Pro	Gln	Pro
			20					25					30		

Lys	Ala	Asn	Gln	Gln	His	Gln	Asp	Asn	Ala	Arg	Gly	Leu	Val	Leu	Pro
		35					40					45			

Gly	Tyr	Lys	Tyr	Leu	Gly	Pro	Gly	Asn	Gly	Leu	Asp	Lys	Gly	Glu	Pro
	50					55						60			

Ile	Asn	Ala	Ala	Asp	Ala	Ala	Ala	Leu	Glu	His	Asp	Lys	Ala	Tyr	Asp
65					70					75					80

Gln	Gln	Leu	Lys	Ala	Gly	Asp	Asn	Pro	Tyr	Leu	Lys	Tyr	Asn	His	Ala
				85					90					95	

Asp	Ala	Glu	Phe	Gln	Glu	Arg	Leu	Lys	Glu	Asp	Thr	Ser	Phe	Gly	Gly
			100					105					110		

Asn	Leu	Gly	Arg	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Leu	Leu	Glu	Pro
		115					120					125			

Leu	Gly	Leu	Val	Glu	Glu	Ala	Ala	Lys	Thr	Ala	Pro	Gly	Lys	Lys	Arg
	130					135					140				

Pro	Val	Glu	Gln	Ser	Pro	Gln	Glu	Pro	Asp	Ser	Ser	Ala	Gly	Ile	Gly
145					150					155					160

Lys	Ser	Gly	Ala	Gln	Pro	Ala	Lys	Lys	Arg	Leu	Asn	Phe	Gly	Gln	Thr
				165					170					175	

Gly	Asp	Thr	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Ile	Gly	Glu	Pro	Pro
			180					185					190		

Ala	Ala	Pro	Ser	Gly	Val	Gly	Ser	Leu	Thr	Met	Ala	Ser	Gly	Gly	Gly
		195					200					205			

Ala	Pro	Val	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Ser	Ser
	210					215					220				

Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Gln	Trp	Leu	Gly	Asp	Arg	Val	Ile
225					230					235					240

Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu
				245					250					255	

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290 295 300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
325 330 335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
355 360 365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
370 375 380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385 390 395 400

Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
405 410 415

Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
420 425 430

Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
435 440 445

Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
450 455 460

Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
465 470 475 480

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
485 490 495

Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
500 505 510

Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
515 520 525

Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
530 535 540

Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
545 550 555 560

Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser

```

Tyr Asn Glu Glu Glu Thr Lys Thr Thr Asn Thr Val Ala Thr Glu Ser
565                               570                               575

Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
580                               585                               590

Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
595                               600                               605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610                               615                               620

Thr Tyr Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
625                               630                               635                               640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645                               650                               655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660                               665                               670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675                               680                               685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690                               695                               700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705                               710                               715                               720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725                               730                               735

<210> 5
<211> 736
<212> PRT
<213> Artificial Sequence

<220>
<223> novel AAV isolate

<400> 5
Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1           5           10           15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20           25           30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35           40           45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50           55           60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65           70           75           80

```

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
 100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
 115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Asp
 145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
 180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
 195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
 210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 6

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 6

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Leu Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245 250 255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290 295 300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
325 330 335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
355 360 365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
370 375 380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385 390 395 400

Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
405 410 415

Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
420 425 430

Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
435 440 445

Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
450 455 460

Val Ala Gly Ser Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
465 470 475 480

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
485 490 495

Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510

Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525

Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540

Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560

Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575

Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590

Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620

Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
 705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

<210> 7

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 7

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
 20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
 100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
 115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
 145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
 180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
 195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
 210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu

245 250 255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300
 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320
 Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350
 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Arg Ala Gln
 580 585 590

Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 8

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 8

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Val Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
 100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
 115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
 145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
 180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
 195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
 210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400

Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 9

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 9

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Arg Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser

210						215						220					
Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Gln	Trp	Leu	Gly	Asp	Arg	Val	Ile		
225					230					235					240		
Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu		
				245					250					255			
Tyr	Lys	Gln	Ile	Ser	Asn	Ser	Thr	Ser	Gly	Gly	Ser	Ser	Asn	Asp	Asn		
			260					265					270				
Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg		
		275					280					285					
Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn		
290						295					300						
Asn	Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	Lys	Leu	Phe	Asn	Ile		
305					310					315					320		
Gln	Val	Lys	Glu	Val	Thr	Asp	Asn	Asn	Gly	Val	Lys	Thr	Ile	Ala	Asn		
				325					330					335			
Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr	Asp	Ser	Asp	Tyr	Gln	Leu		
			340					345					350				
Pro	Tyr	Val	Leu	Gly	Ser	Ala	His	Glu	Gly	Cys	Leu	Pro	Pro	Phe	Pro		
		355					360					365					
Ala	Asp	Val	Phe	Met	Ile	Pro	Gln	Tyr	Gly	Tyr	Leu	Thr	Leu	Asn	Asp		
	370					375					380						
Gly	Ser	Gln	Ala	Val	Gly	Arg	Ser	Ser	Phe	Tyr	Cys	Leu	Glu	Tyr	Phe		
385					390					395					400		
Pro	Ser	Gln	Met	Leu	Arg	Thr	Gly	Asn	Asn	Phe	Gln	Phe	Ser	Tyr	Glu		
			405						410					415			
Phe	Glu	Asn	Val	Pro	Phe	His	Ser	Ser	Tyr	Ala	His	Ser	Gln	Ser	Leu		
			420					425					430				
Asp	Arg	Leu	Met	Asn	Pro	Leu	Ile	Asp	Gln	Tyr	Leu	Tyr	Tyr	Leu	Ser		
		435					440					445					
Lys	Thr	Ile	Asn	Gly	Ser	Gly	Gln	Asn	Gln	Gln	Thr	Leu	Lys	Phe	Ser		
	450					455					460						
Val	Ala	Gly	Pro	Ser	Asn	Met	Ala	Val	Gln	Gly	Arg	Asn	Tyr	Ile	Pro		
465					470					475					480		
Gly	Pro	Ser	Tyr	Arg	Gln	Gln	Arg	Val	Ser	Thr	Thr	Val	Thr	Gln	Asn		
				485					490					495			
Asn	Asn	Ser	Glu	Phe	Ala	Trp	Pro	Gly	Ala	Ser	Ser	Trp	Ala	Leu	Asn		
			500					505					510				
Gly	Arg	Asn	Ser	Leu	Met	Asn	Pro	Gly	Pro	Ala	Met	Ala	Ser	His	Lys		

515		520		525
Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly				
530		535		540
Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile				
545		550		555
Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser				
		565		570
				575
Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln				
		580		585
				590
Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln				
		595		600
				605
Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His				
610		615		620
Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met				
625		630		635
				640
Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala				
		645		650
				655
Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr				
		660		665
				670
Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln				
		675		680
				685
Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn				
690		695		700
Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val				
705		710		715
				720
Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu				
		725		730
				735
<210> 10				
<211> 736				
<212> PRT				
<213> Artificial Sequence				
<220>				
<223> novel AAV isolate				
<400> 10				
Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser				
1		5		10
				15
Glu Glv Ile Arg Glu Trp Trp Ala Leu Lvs Pro Glv Ala Pro Gln Pro				

```

-----
                20                25                30
Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
      35                40                45
Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
      50                55                60
Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
      65                70                75                80
Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
      85                90                95
Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
      100                105                110
Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
      115                120                125
Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
      130                135                140
Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
      145                150                155                160
Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
      165                170                175
Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
      180                185                190
Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Cys Gly Gly
      195                200                205
Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
      210                215                220
Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
      225                230                235                240
Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
      245                250                255
Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
      260                265                270
Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
      275                280                285
Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
      290                295                300
Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
      305                310                315                320

```

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350
 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Thr Leu Gln Gly Pro Ile Thr Ala Lys Ile Pro His

Asp Arg Asp Val Tyr Leu Gln Gly Phe Ile Trp Ala Lys Ile Phe His
610 615 620

Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 11

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 11

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Arg Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

```

---
Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130                      135                      140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145                      150                      155                      160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
                      165                      170                      175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180                      185                      190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195                      200                      205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210                      215                      220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
225                      230                      235                      240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245                      250                      255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260                      265                      270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275                      280                      285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290                      295                      300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
305                      310                      315                      320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
325                      330                      335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
340                      345                      350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
355                      360                      365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
370                      375                      380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385                      390                      395                      400

Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
405                      410                      415

```

Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430

Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445

Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460

Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495

Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510

Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525

Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540

Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560

Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575

Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590

Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620

Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685

Lys Lys Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
 705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 12

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 12

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile

```

225      -      -      230      -      -      -      235      -      -      -      240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
      245      250      255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
      260      265      270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
      275      280      285

Phe His Cys His Phe Ser Pro His Asp Trp Gln Arg Leu Ile Asn Asn
      290      295      300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
305      310      315      320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
      325      330      335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
      340      345      350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
      355      360      365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
      370      375      380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385      390      395      400

Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
      405      410      415

Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
      420      425      430

Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
      435      440      445

Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Asn
      450      455      460

Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
465      470      475      480

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
      485      490      495

Asn Asn Ser Glu Phe Ala Trp Pro Arg Ala Ser Ser Trp Ala Leu Asn
      500      505      510

Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
      515      520      525

Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
      530      535

```

530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Met Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700
 Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
 705 710 715 720
 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

<210> 13

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 13

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
 20 25 30
 Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195 200 205

Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210 215 220

Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245 250 255

Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260 265 270

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290 295 300

Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
325 330 335

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Arg Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala

```

-
        645              650              655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
        660              665              670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
        675              680              685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
        690              695              700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705              710              715              720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
        725              730              735

<210> 14
<211> 736
<212> PRT
<213> Artificial Sequence

<220>
<223> novel AAV isolate

<400> 14
Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1              5              10              15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
        20              25              30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
        35              40              45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
        50              55              60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65              70              75              80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
        85              90              95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
        100              105              110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
        115              120              125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
        130              135              140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145              150              155              160

```

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175
 Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
 180 185 190
 Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
 195 200 205
 Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
 210 215 220
 Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225 230 235 240
 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255
 Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270
 Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285
 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300
 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320
 Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350
 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445

Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Arg Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700
 Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
 705 710 715 720
 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

<210> 15

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 15

Met	Ala	Ala	Asp	Gly	Tyr	Leu	Pro	Asp	Trp	Leu	Glu	Asp	Asn	Leu	Ser
1				5					10					15	

Glu	Gly	Ile	Arg	Glu	Trp	Trp	Ala	Leu	Lys	Pro	Gly	Ala	Pro	Gln	Pro
			20					25					30		

Lys	Ala	Asn	Gln	Gln	His	Gln	Asp	Asn	Ala	Arg	Gly	Leu	Val	Leu	Pro
		35					40					45			

Gly	Tyr	Lys	Tyr	Leu	Gly	Pro	Gly	Asn	Gly	Leu	Asp	Lys	Gly	Glu	Pro
	50					55					60				

Val	Asn	Ala	Ala	Asp	Ala	Ala	Ala	Leu	Glu	His	Asp	Lys	Ala	Tyr	Asp
65					70					75					80

Gln	Gln	Leu	Lys	Ala	Gly	Asp	Asn	Pro	Tyr	Leu	Lys	Tyr	Asn	His	Ala
					85					90					95

Asp	Ala	Glu	Phe	Gln	Glu	Arg	Leu	Lys	Glu	Asp	Thr	Ser	Phe	Gly	Gly
			100					105					110		

Asn	Leu	Gly	Arg	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Leu	Leu	Glu	Pro
		115					120					125			

Leu	Gly	Leu	Val	Glu	Glu	Ala	Ala	Lys	Thr	Ala	Pro	Gly	Lys	Lys	Arg
	130					135					140				

Pro	Val	Glu	Gln	Ser	Pro	Gln	Glu	Pro	Asp	Ser	Ser	Ala	Gly	Ile	Gly
145					150					155					160

Lys	Ser	Gly	Ala	Gln	Pro	Ala	Lys	Lys	Arg	Leu	Asn	Phe	Gly	Gln	Thr
				165					170					175	

Gly	Asp	Thr	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Ile	Gly	Glu	Pro	Pro
			180					185					190		

Ala	Ala	Pro	Ser	Gly	Val	Gly	Ser	Leu	Thr	Met	Ala	Ser	Gly	Gly	Gly
		195					200					205			

Ala	Pro	Val	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Ser	Ser
	210					215					220				

Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Gln	Trp	Leu	Gly	Asp	Arg	Val	Ile
225					230					235					240

Met	Met	Glu	Met	Asp	Met	Met	Ala	Tyr	Met	Met	Met	Met	Met	Met	Met
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255
 Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270
 Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285
 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300
 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320
 Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350
 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Arg Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Thr Gln Gln Phe Gln Arg Arg Arg Val Arg Ala Arg Thr Val Met Thr

Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
545 550 555 560

Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
565 570 575

Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
580 585 590

Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Arg Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 16

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 16

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val	Asn	Ala	Ala	Asp	Ala	Ala	Ala	Leu	Glu	His	Asp	Lys	Ala	Tyr	Asp		
65					70					75					80		
Gln	Gln	Leu	Lys	Ala	Gly	Asp	Asn	Pro	Tyr	Leu	Lys	Tyr	Asn	His	Ala		
				85					90					95			
Asp	Ala	Glu	Phe	Gln	Glu	Arg	Leu	Lys	Glu	Asp	Thr	Ser	Phe	Gly	Gly		
			100					105					110				
Asn	Leu	Gly	Arg	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Leu	Leu	Glu	Pro		
		115					120					125					
Leu	Gly	Leu	Val	Glu	Glu	Ala	Ala	Lys	Thr	Ala	Pro	Gly	Lys	Lys	Arg		
	130					135					140						
Pro	Val	Glu	Gln	Ser	Pro	Gln	Glu	Pro	Asp	Ser	Ser	Ala	Gly	Ile	Gly		
145					150				155						160		
Lys	Ser	Gly	Ala	Gln	Pro	Ala	Lys	Lys	Arg	Leu	Asn	Phe	Gly	Gln	Thr		
				165					170					175			
Gly	Asp	Thr	Glu	Ser	Val	Pro	Asp	Pro	Gln	Pro	Ile	Gly	Glu	Pro	Pro		
			180					185					190				
Ala	Ala	Pro	Ser	Gly	Val	Gly	Ser	Leu	Thr	Met	Ala	Ser	Gly	Gly	Gly		
		195					200					205					
Ala	Pro	Val	Ala	Asp	Asn	Asn	Glu	Gly	Ala	Asp	Gly	Val	Gly	Ser	Ser		
	210					215					220						
Ser	Gly	Asn	Trp	His	Cys	Asp	Ser	Gln	Trp	Leu	Gly	Asp	Arg	Val	Ile		
225					230					235					240		
Thr	Thr	Ser	Thr	Arg	Thr	Trp	Ala	Leu	Pro	Thr	Tyr	Asn	Asn	His	Leu		
				245					250					255			
Tyr	Lys	Gln	Ile	Ser	Asn	Ser	Thr	Ser	Gly	Gly	Ser	Ser	Asn	Asp	Asn		
			260					265					270				
Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	Phe	Asp	Phe	Asn	Arg		
		275					280					285					
Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	Arg	Leu	Ile	Asn	Asn		
	290					295					300						
Asn	Trp	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	Lys	Leu	Phe	Asn	Ile		
305					310					315					320		
Gln	Val	Lys	Glu	Val	Thr	Asp	Asn	Asn	Gly	Val	Lys	Thr	Ile	Ala	Asn		
				325					330					335			
Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Ala	Asp	Ser	Asp	Tyr	Gln	Leu		
			340					345					350				
Pro	Tyr	Val	Leu	Gly	Ser	Ala	His	Glu	Gly	Cys	Leu	Pro	Pro	Phe	Pro		
		355					360					365					

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Phe Ala Trp Pro Arg Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

<210> 17

<211> 736

<212> PRT

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 17

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

Glu Asn Thr Glu Ser Val Pro Asn Pro Gln Pro Ile Glu Glu Pro Pro

Gly Asp Thr Glu Ser Val Phe Asp Phe Gln Phe Ile Gly Glu Phe Phe
 180 185 190
 Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
 195 200 205
 Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
 210 215 220
 Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225 230 235 240
 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255
 Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
 260 265 270
 Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285
 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300
 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320
 Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
 325 330 335
 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350
 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365
 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380
 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400
 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415
 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430
 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445
 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460
 Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480
 - - - - - - - - - - - - - - - - -

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
 485 490 495
 Asn Asn Ser Glu Ile Ala Trp Pro Arg Ala Ser Ser Trp Ala Leu Asn
 500 505 510
 Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
 530 535 540
 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
 545 550 555 560
 Thr Asn Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
 565 570 575
 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln
 580 585 590
 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685
 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700
 Tyr Cys Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
 705 710 715 720
 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

<210> 18

<211> 2211

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 18

atggctgccg atggttatct tccagattgg ctcgaggaca accttagtga aggaattcgc	60
gagtgggtggg ctttgaaacc tggagcccct caacccaagg caaatcaaca acatcaagac	120
aacgctcgag gtcttgtgct tccgggttac aaataccttg gacccggcaa cggactcgac	180
aagggggagc cggccaacgc agcagacgcg gcggccctcg agcacgacaa ggcctacgac	240
cagcagctca aggccggaga caaccggtac ctcaagtaca accacgccga cgccgagttc	300
caggagcggc tcaaagaaga tacgtctttt gggggcaacc tcgggcgagc agtcttcag	360
gccaaaaaga ggcttcttga acctcttggg ctgggtgagg aagcggctaa gacggctcct	420
ggaaagaaga ggcctgtaga gcagtctcct caggaaccgg actcctccgc gggattggc	480
aaatcgggtg cacagcccgc taaaaagaga ctcaatttcg gtcagactgg cgacacagag	540
tcagtcccag accctcaacc aatcggagaa cctcccgag cccctcagg tgtgggatct	600
cttacaatgg cttcaggtgg tggcgacca gtggcagaca ataacgaagg tgccgatgga	660
gtgggtagtt cctcgggaaa ttggcattgc gattcccaat ggctggggga cagagtcac	720
accaccagca ccogaacctg ggcctgccc acctacaaca atcacctcta caagcaaata	780
tccaacagca catctggagg atcttcaaata gacaacgcct acttcggcta cagcaccccc	840
tgggggtatt ttgacttcaa cagattccac tgccacttct caccacgtga ctggcagcga	900
ctcatcaaca acaactgggg attccggcct aagcgactca acttcaagct cttcaacatt	960
caggtcaaag aggttacgga caacaatgga gtcaagacca tcgccataa ccttaccagc	1020
acggtccagg tcttcacgga ctcagactat cagctcccgt acgtgctcgg gtcggctcac	1080
gagggctgcc tcccgccgtt ccagcggac gttttcatga ttctcagta cgggtatctg	1140
acgcttaatg atggaagcca ggccgtgggt cgttcgtcct tttactgcct ggaatatttc	1200
ccgtcgcaaa tgctaagaac gggtaacaac ttccagttca gctacgagtt tgagaacgta	1260
cctttccata gcagctacgc tcacagccaa agcctggacc gactaatgaa tccactcatc	1320
gaccaatact tgtactatct ctcaaagact attaacgggt ctggacagaa tcaacaaacg	1380
ctaaaattca gtgtggccgg acccagcaac atggctgtcc aggaagaaa ctacatacct	1440
ggaccagct accgacaaca acgtgtctca accactgtga ctcaaaacaa caacagcgaa	1500
tttgcttggc ctggagcttc ttcttgggct ctcaatggac gtaatagctt gatgaatcct	1560
ggacctgcta tggccagcca caaagaagga gaggaccgtt tctttccttt gtctggatct	1620
ttaatttttg gcaaacaagg aactggaaga gacaacgtgg atgcggacaa agtcatgata	1680
accaacgaag aagaaattaa aactactaac ccggtagcaa cggagtccta tggacaagtg	1740
gccacaaacc accagagtgc ccaagcacag gcgcagaccg gctgggttca aaaccaagga	1800
atacttccgg gtatggtttg gcaggacaga gatgtgtacc tgcaaggacc catttgggcc	1860
aaaattcctc acacggacgg caactttcac ccttctccgc tgatgggagg gtttggaatg	1920
aagcaccgc ctctcagat cctcatcaaa aacacacctg tacctgcgga tcctccaacg	1980
gccttcaaca aggacaagct gaactctttc atcaccagat attctactgg ccaagtcagc	2040
gtggagatcg agtgggagct gcagaaggaa aacagcaagc gctggaacct ggagatccag	2100

tacacttcca actattacaa gtctaataat gttgaatttg ctgttaatac tgaaggtgta 2160
 tatagtgaac cccgccccat tggcaccaga tacctgactc gtaatctgta a 2211

<210> 19

<211> 4679

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 19

ttggccactc cctctctgcg cgctcgctcg ctactgagg ccgggcgacc aaaggtcgcc 60
 cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc gagcgcgag agagggagtg 120
 gccaaactcca tcactagggg ttcttgagg ggtggagtcg tgacgtgaat tacgtcatag 180
 ggttagggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacaccat 240
 gtggtcacgc tgggtattta agcccagtg agcacgcagg gtctccattt tgaagcggga 300
 ggtttgaacg cgcagccgcc atgccgggt tttacgagat tgtgattaag gtccccagcg 360
 accttgacga gcatctgcc ggcatctctg acagctttgt gaactgggtg gccgagaagg 420
 aatgggagtt gccgccagat tctgacatgg atctgaatct gattgagcag gcacccctga 480
 ccgtggccga gaagctgcag cgcgactttc tgacggaatg gcgccgtgtg agtaaggccc 540
 cggaggccct tttctttgtg caatttgaga agggagagag ctacttccac atgcacgtgc 600
 tcgtggaaac caccggggtg aaatccatgg ttttgggacg tttcctgagt cagattcgcg 660
 aaaaactgat tcagagaatt taccgcggga tcgagccgac tttgccaaac tggttcgcg 720
 tcacaaagac cagaaatggc gccggaggcg ggaacaagg ggtggatgag tgctacatcc 780
 ccaattactt gctccccaaa acccagcctg agctccagtg ggcgtggact aatatggaac 840
 agtatttaag cgctgtttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga 900
 cgcacgtgtc gcagacgcag gagcagaaca aagagaatca gaatcccaat tctgatgcgc 960
 cggtgatcag atcaaaaact tcagccaggt acatggagct ggtcgggtgg ctcggtggaca 1020
 aggggattac ctcggagaag cagtggatcc aggaggacca ggcctcatac atctccttca 1080
 atgcggcctc caactcgcg tcccaaatac aggctgcctt ggacaatgcg ggaaagatta 1140
 tgagcctgac taaaaccgcc ccgactacc tgggtggcca gcagcccggtg gaggacattt 1200
 ccagcaatcg gatattataa attttggaa taaacgggta cgatcccaa tatgcggctt 1260
 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg 1320
 ggcctgcaac taccgggaag accaacaatc cgagggccat agcccacact gtgcccttct 1380
 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg 1440
 tgatctggtg ggaggagggg aagatgaccg ccaaggtcgt ggagtcggcc aaagccattc 1500
 tcggaggaag caaggtgcgc gtggaccaga aatgcaagtc ctcggccag atagaccga 1560
 ctcccgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga 1620

ccttcgaaca ccagcagccg ttgcaagacc ggatgttcaa atttgaactc acccgccgtc	1680
tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cgggtgggcaa	1740
aggatcacgt ggttgaggtg gagcatgaat tctacgtcaa aaaggggtgga gccaaagaaa	1800
gacccgcccc cagtgacgca gatataagtg agcccaaacg ggtgcgcgag tcagttgcg	1860
agccatcgac gtcagacgcg gaagcttcga tcaactacgc agacaggtac caaaacaaat	1920
gttctcgtca cgtgggcatg aatctgatgc tgtttccctg cagacaatgc gagagaatga	1980
atcagaattc aaatatctgc ttcactcacg gacagaaaga ctgtttagag tgctttccc	2040
tgtcagaatc tcaaccggtt tctgtcgtca aaaaggcgta tcagaaactg tgctacattc	2100
atcatatcat gggaaaggtg ccagacgctt gcactgcctg cgatctgggtc aatgtggatt	2160
tggatgactg catctttgaa caataaatga tttaaatacag gtatggctgc cgatgggttat	2220
cttcagatt ggctcgagga cactctctct gaaggaataa gacagtgggtg gaagctcaaa	2280
cctggcccac caccaccaa gcccgagag cggcataagg acgacagcag gggctctgtg	2340
cttcctgggt acaagtacct cggacccttc aacggactcg acaagggaga gccggtcaac	2400
gaggcagacg ccgcgccct cgagcacgac aaagcctacg accggcagct cgacagcgga	2460
gacaaccggt acctcaagta caaccacgcc gacgcggagt ttcaggagcg ccttaaagaa	2520
gatacgtctt ttgggggcaa cctcgagca gcagtcttcc aggcgaaaaa gagggttctt	2580
gaacctctgg gcctggttga ggaacctgtt aagacggctc cgggaaaaaa gaggccggt	2640
gagcactctc ctgtggagcc agactcctcc tcgggaaccg gaaaggcggg ccagcagcct	2700
gcaagaaaaa gattgaattt tggtcagact ggagacgcag actcagtacc tgacccccag	2760
cctctcggac agccaccagc agccccctct ggtctgggaa ctaatacgat ggctacaggc	2820
agtggcgcac caatggcaga caataacgag ggcgccgacg gagtgggtaa ttcctcggga	2880
aattggcatt gcgattccac atggatgggc gacagagtca tcaccaccag caccogaacc	2940
tgggccctgc ccacctaca caaccacctc taaaaacaaa tttccagcca atcaggagcc	3000
tcgaacgaca atcactactt tggctacagc accccttggg ggtattttga cttcaacaga	3060
ttccactgcc acttttcacc acgtgactgg caaagactca tcaacaacaa ctggggattc	3120
cgaccaaga gactcaactt caagctcttt aacattcaag tcaaagaggt cacgcagaat	3180
gacggtacga cgacgattgc caataacctt accagcacgg ttcagggtgtt tactgactcg	3240
gagtaccagc tcccgtacgt cctcggctcg gcgcacaaag gatgcctccc gccgttcca	3300
gcagacgtct tcatggtgcc acagtatgga tacctcacc tgaacaacgg gagtacggca	3360
gtaggacgct cttcatthta ctgcctggag tactttcctt ctcagatgct gcgtaccgga	3420
aacaacttta ccttcagcta cacttttgag gacgttcctt tccacagcag ctacgctcac	3480
agccagagtc tggaccgtct catgaatcct ctcacgacc agtacctgta ttacttgagc	3540
agaacaaaca ctccaagtgg aaccaccacg cagtcaaggc ttcagttttc tcaggccgga	3600
gcgagtgaca ttcgggacca gtctaggaac tggcttcctg gaccctgtta ccgccagcag	3660
cgagtatcaa agacatctgc ggataaacac aacagtgaat actcgtggac tggagctacc	3720
aagtaccacc tcaatggcag agactctctg gtgaatccgg gcccggccat ggcaagccac	3780
aaggacgatg aagaaaagtt ttttcctcag agcgggggtc tcatctttgg gaagcaaggc	3840

tcagagaaaa caaatgtgga cattgaaaag gtcattgatta cagacgaaga ggaaatcagg	3900
acaaccaatc ccgtggctac ggagcagtat ggttctgtat ctaccaacct ccagagagggc	3960
aacagacaag cagctaccgc agatgtcaac acacaaggcg ttcttccagg catggtcttg	4020
caggacagag atgtgtacct tcagggggccc atctgggcaa agattccaca cacggacgga	4080
cattttcacc cctctccctt catgggtgga ttcggactta aacaccctcc tccacagatt	4140
ctcatcaaga acaccccggt acctgcgaat ccttcgacca ccttcagtgc ggcaaagttt	4200
gcttccttca tcacacagta ctccacggga caggtcagcg tggagatcga gtgggagctg	4260
cagaaggaaa acagcaaacg ctggaatccc gaaattcagt acacttccaa ctacaacaag	4320
tctgttaatg tggactttac tgtggacact aatggcgtgt attcagagcc tcgccccatt	4380
ggcaccagat acctgactcg taatctgtaa ttgcttgta atcaataaac cgtttaattc	4440
gtttcagttg aactttggtc tctgcgtatt tctttcttat ctagtttcca tggctacgta	4500
gataagtagc atggcggggt aatcattaac tacaaggaa ccctagtgat ggagttggcc	4560
actccctctc tgcgcgctcg ctgcctcact gaggcggggc gaccaaaggc cggccgacgc	4620
ccgggctttg cccggggggc ctcagtgagc gagcgagcgc gcagagaggg agtggccaa	4679

<210> 20

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 20

ccatcgacgt cagacgcgga agcttcgac aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggg gtcaaagact gtttagagtg ctttcccgctg	180
tcagaatctc aacccgtttc tgcgtcaaa aaggcgtatc agaaactgtg ctacattcat	240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataatgact taaaccaggt atgactgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtggtggg ctttgaaacc	420
tggagcccct caacccaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gacccggcaa cggactcgac aagggggagc cggccaacgc	540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggcgggaga	600
caaccogtac ctcaagtaca accacgcca cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga	720
acctcttggt ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggcctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgag cccctcagg tgtgggatct cttaaatgg cttcagggtg	960

tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcato accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaato tccaacagca catctggagg	1140
atcttcaaat gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg	1260
attccggcct aagcaactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtccggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttccctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct tttactgcct ggaatatctc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcato gaccaatact tgtactatct	1680
ctcaaagact attaacgggt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atggctgtcc agggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctggagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttcogg gtatggtttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct	2340
gaactcttcc atcaccagct attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 21

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 21

ccatcgacgt cagacgcgga agcttcgato aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcagg tggggtcaa tctcgtcgtc tttccctgca gagaatcga gacagtgaat	120

ccccgcaacg	cyggcatgaa	ccgacgctg	cccccccgca	gacaaatgca	gagacatgaa	120
cagaattcaa	atatctgctt	cactcacggt	gtcaaagact	gtttagagtg	ctttcccgtg	180
tcagaatctc	aaccggtttc	tgtcgtcaaa	aaggcgtatc	agaaactgtg	ctacattcat	240
cacatcatgg	gaaaggtgcc	agacgcttgc	actgcttgcc	acctgggtcaa	tgtggacttg	300
gatgactgtg	tttctgaaca	ataaatgact	taaaccaggt	atggctgccg	atggttatct	360
tccagattgg	ctcgaggaca	accttagtga	aggaattcgc	gagtgggtggg	ctttgaaacc	420
tggagcccct	caacccaagg	caaatacaaca	acatcaagac	aacgctcgag	gtcttgtgct	480
tccgggttac	aaataccttg	gacccggcaa	cggactcgac	aagggggagc	cgggtcaacgc	540
agcagacgcg	gcggccctcg	agcacgacaa	ggcctacgac	cagcagctca	aggccggaga	600
caaccggtac	ctcaagtaca	accacgcccga	cgcgagttc	caggagcggc	tcaaagaaga	660
tacgtctttt	gggggcaacc	tcgggcgagc	agtcttccag	gccaaaaaga	ggcttcttga	720
acctcttggt	ctggttgagg	aagcggctaa	gacggctcct	ggaaagaaga	ggcctgtaga	780
gcagtctcct	caggaaccgg	actcctccgc	gggtattggc	aaatcgggtg	cacagcccgc	840
taaaaagaga	ctcaatttcg	gtcagactgg	cgacacagag	tcagtcccag	acctcaacc	900
aatcggagaa	cctcccgcag	ccccctcagg	tgtgggatct	cttacaatgg	cttcaggtgg	960
tggcgcacca	gtggcagaca	ataacgaagg	tgccgatgga	gtgggtagtt	cctcgggaaa	1020
ttggcattgc	gattcccaat	ggctggggga	cagagtcatc	accaccagca	cccgaacctg	1080
ggccctgccc	acctacaaca	atcacctcta	caagcaaatc	tccaacagca	catctggagg	1140
atcttcaaat	gacaacgcct	acttcggcta	cagcaccccc	tgggggtatt	ttgacttcaa	1200
cagattccac	tgccacttct	caccacgtga	ctggcagcga	ctcatcaaca	acaactgggg	1260
attccggcct	aagcgactca	acttcaagct	cttcaacatt	caggtcaaag	aggttacgga	1320
caacaatgga	gtcaagacca	tcgccataaa	ccttaccagc	acggtccagg	tcttcacgga	1380
ctcagactat	cagctcccgt	acgtgctcgg	gtcggctcac	gagggctgcc	tcccgccttt	1440
cccagcggac	gttttcatga	ttcctcagta	cgggtatctg	acgcttaatg	atggaagcca	1500
ggccgtgggt	cgttcgtcct	tttactgcct	ggaatatttc	ccgtcgcaaa	tgctaagaac	1560
gggtaacaac	ttccagttca	gctacgagtt	tgagaacgta	cctttccata	gcagctacgc	1620
tcacagccaa	agcctggacc	gactaatgaa	tccactcatc	gaccaatact	tgtactatct	1680
ctcaaagact	attaacgggt	ctggacagaa	tcaacaaacg	ctaaaattca	gtgtggccgg	1740
accagcaac	atggctgtcc	agggagaaaa	ctacatacct	ggacccagct	accgacaaca	1800
acgtgtctca	accactgtga	ctcaaaaaca	caacagcgaa	tttgcttggc	ctggggcttc	1860
ttcttgggct	ctcaatggac	gtaatagctt	gatgaatcct	ggacctgcta	tggccagcca	1920
caaagaagga	gaggaccgtt	tctttccttt	gtctggatct	ttaatttttg	gcaaacaagg	1980
aactggaaga	gacaacgtgg	atgcggacaa	agtcattgata	accaacgaag	aagaaattaa	2040
aactactaac	ccggtagcaa	cggagtccta	tggacaagtg	gccacaaacc	accagagtgc	2100
ccaagcacag	gcgcagaccg	gctgggttca	aaaccaagga	atacttccgg	gtatggtttg	2160
gcaggacaga	gatgtgtacc	tgcaaggacc	catttgggcc	aaaattcctc	acacgggcgg	2220
caactttcac	ccttctccgc	tgatgggagg	gtttggaatg	aagcacccgc	ctcctcagat	2280

```

cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct      2340
gaactctttc atcaccacgt attctactgg ccaagtcagc gtggagatcg agtgggagct      2400
gcagaaggaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattacaa      2460
gtctaataat gttgaatttg ctgttaatac tggaggtgta tatagtgaac ccgccccat      2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt      2580
cgtttcagtt gaactgcggc c                                     2601

```

<210> 22

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 22

```

ccatcgacgt cagacgcgga agcttcgac cactacgcgg acaggtacca aaacaaatgt      60

tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat      120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccggtg      180
tcagaatctc aacccgtttc tgctgcaaaa aaggcgatc agaaactgtg ctacattcat      240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctgggtcaa tgtggactcg      300
gatgactgtg tttctgaaca ataatgact taaaccaggt atggctgccg atggttatct      360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc      420
tggagccctt caaccaagg caaatcaaca acatcaagac aacgctcgag gtcttggtgt      480
tccgggttac aaataccttg gaccggcaa cggactcgac aagggggagc cggccaacgc      540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga      600
caaccggtac ctcaagtaca accacgcga cgcgagttc caggagcggc tcaaagaaga      660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga      720
acctcttggt ctgggtgagg aagcgggtaa gacggctcct ggaaagaaga ggctgtaga      780
gcagtctcct caggaaccgg actcctccgc gggattgac aaatcgggtg cacagcccg      840
taaaaagaga ctcaatttcg gtcagaactg cgacacagag tcagtcccag acctcaacc      900
aatcggagaa cctcccgag cccctcagg tgtgggatct cttacaatgg cttcaggtgg      960
tggcgaccca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa     1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg     1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg     1140
atcttcaaata gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa     1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg     1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga     1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga     1380
ctcggactat cagctcccg acgtgctcgg gtcggctcac gagggctgcc tcccgcggtt     1440
ggggggggg cttttgataa ttactgacta ggggtatctg ggggttaata atggaacgga     1500

```

```

ccagcggac gtttccatga ttttccagca tgggtatctg acgttcaatg atggaagcca 1500
ggccgtgggt cgttcgtcct tttactgcct ggaatatttc ccgtcgcaaa tgctaagaac 1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cttttccata gcagctacgc 1620
tcacagccaa agcctggacc gactaatgaa tccactcacc gaccaatact tgtactatct 1680
ctcaaagact attaacgggt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg 1740
accagcaac atggctgtcc agggaagaaa ctacatacct ggacccagct accgacaaca 1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctggagcttc 1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca 1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaacaagg 1980

aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa 2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc 2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatggtttg 2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg 2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctctcagat 2280
cctcatcaaa aacacacctg tacctgcgga tctccaacg gccttcaaca aggacaagct 2340
gaactctttc atcaccagct attctactgg ccaagtcagc gtggagatcg agtgggagct 2400
gcagaaggaa aacagcaagc gctggaaccg ggagatccag tacacttcca actattacaa 2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac ccgccccat 2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt 2580
cgtttcagtt gaactgcggc c 2601

```

<210> 23

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 23

```

ccatcgacgt cagacgcgga agcttcgacc aactacgcgg acaggtacca aaacaaatgt 60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat 120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg 180
tcagaatctc aacccgtttc tgtcgtcaaa aaggcgtacc agaaactgtg ctacattcat 240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg 300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct 360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc 420
tggagccctt caaccgaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct 480
tccgggttac aaataccttg gaccgggcaa cggactcgac aagggggagc cgggtcaacgc 540
agcagacgcg gcggccctcg agcagacaa ggcctacgac cagcagctca aggccggaga 600

```

caacccgtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcctccag gccaaaaaga ggcttcttga	720
acctcttggt ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag accctcaacc	900
aatcggagaa cctcccgcag cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaato tccaacagca catctggagg	1140
atcttcaa at gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcacaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccataaa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccg t acgtgctcgg gtcggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct	1680
ctcaaagact attaacggtt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
atccagcaac atggctgtcc aggggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctggagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccag gtatggtttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct	2340
gaactctttc atcaccagct attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttgtt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 24

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 24

ccatcgacgt cagacgcgga agcttcgata aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgtcgtcaaa aaggcgtatc agaaactgtg ctacattcat	240
cacatcatgg gaaaggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccct caaccgaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gaccgggcaa cggactcgac aagggggagc cgggtcaacgc	540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga	600
caacccgtag ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga	720
acctcttggg ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgcag cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcata accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaat gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtcggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcata gaccaatact tgtactatct	1680
ctcaaagact attaacggtt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atggctgtcc agggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa ttgcttggc ctggagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920

caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacgg gcgcagaccg gctgggttca aaaccaagga atacttcogg gtatggtttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctcctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tctccaacg gccttcaaca aggacaagct	2340
gaactctttc atcaccagc attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaacc ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac ccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 25

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 25

ccatcgacgt cagacgcgga agcttcgac aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgcgtcaaa aaggcgtatc agaaactgtg ctacattcat	240
cacatcatgg gaaaggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccct caaccgaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gacccggcaa cggactcgac aagggggagc cgggtcaacgc	540
agcagacgcg gcggccctcg agcacgacag ggcctacgac cagcagctca aggccggaga	600
caaccggtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttcag gccaaaaa ggcttcttga	720
acctcttggc ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggcctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgcag cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttaccattac aattcccaat aactaaaaa caaatcctc accaccaaca cccaaacctc	1080

```

-----
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg 1140
atcttcaaata gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa 1200
cagattccac tgcacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg 1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga 1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga 1380
ctcagactat cagctcccgt acgtgctcgg gtccggtcac gagggctgcc tcccgccgtt 1440
cccagcggac gttttcatga ttcttcagta cgggtatctg acgcttaatg atggaagcca 1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgaaa tgctaagaac 1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc 1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct 1680
ctcaaagact attaacggtt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg 1740
acctagcaac atggctgtcc aggaagaaa ctacatacct ggaccagct accgacaaca 1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctggagcttc 1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca 1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg 1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa 2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc 2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttcggg gtatggtttg 2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg 2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctcctcagat 2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct 2340
gaactctttc atcaccagc attctactgg ccaagtcagc gtggagatcg agtgggagct 2400
gcagaagaaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattacaa 2460
gtctaataat gttgaatttg ctgttaacac tgaaggtgta tatagtgaac cccgccccat 2520
tggcaccaga tacctgactc gtaatctgta attgcttgtt aatcaataaa ccgtttaatt 2580

cgtttcagtt gaactgcggc c 2601

```

<210> 26

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 26

```

ccatcgacgt cagacgcgga agcttcgata aactacgcgg acaggtacca aaacaaatgt 60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat 120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg 180
tcagaatctc aacccgtttc tctcctcaaa aagccctatc agaaactctg ctacattcat 240

```

caacatcatgg	gaaaggtgcc	agacgcttgc	actgcttgcg	acctgggtcaa	tgtggacttg	300
gatgactgtg	tttctgaaca	ataaatgact	taaaccaggt	atggctgccg	atggttatct	360
tccagattgg	ctcgaggaca	accttagtga	aggaattcgc	gagtgggtggg	ctttgaaacc	420
tggagcccct	caacccaagg	caaatcaaca	acatcaagac	aacgctcgag	gtcttgtgct	480
tccgggttac	aaataccttg	gacccggcaa	cggactcgac	aagggggagc	cgatcaacgc	540
agcagacgcg	gcggccctcg	agcacgacaa	ggcctacgac	cagcagctca	aggccggaga	600
caaccctgtac	ctcaagtaca	accacgccga	cgccgagttc	caggagcggc	tcaaagaaga	660
tacgtctttt	gggggcaacc	tggggcgagc	agtcttccag	gccaaaaaga	ggcttcttga	720
acctcttggg	ctggttgagg	aagcggctaa	gacggctcct	ggaaagaaga	ggcctgtaga	780
gcagtctcct	caggaaccgg	actcctccgc	gggtattggc	aaatcgggtg	cacagcccgc	840
taaaaagaga	ctcaatttcg	gtcagactgg	cgacacagag	tcagtcccag	accctcaacc	900
aatcggagaa	cctcccgcag	ccccctcagg	tgtgggatct	cttacaatgg	cttcaggtgg	960
tggcgcacca	gtggcagaca	ataacgaagg	tgccgatgga	gtgggtagtt	cctcgggaaa	1020
ttggcattgc	gattcccaat	ggctggggga	cagagtcctc	accaccagca	cccgaacctg	1080
ggccctgccc	acctacaaca	atcacctcta	caagcaaatc	tccaacagca	catctggagg	1140
atcttcaa	gacaacgcct	acttcggcta	cagcaccctc	tgggggtatt	ttgacttcaa	1200
cagattccac	tgccacttct	caccacgtga	ctggcagcga	ctcatcaaca	acaactgggg	1260
attccggcct	aagcgactca	acttcaagct	cttcaacatt	caggtcaaag	aggttacgga	1320
caacaatgga	gtcaagacca	tgcccaataa	ccttaccagc	acggtccagg	tcttcacgga	1380
ctcagactat	cagctcccgt	acgtgctcgg	gtcggctcac	gagggctgcc	tcccgccgtt	1440
cccagcggac	gttttcatga	ttcctcagta	cgggtatctg	acgcttaatg	atggaagcca	1500
ggccgtgggt	cgttcgtcct	tttactgcct	ggaatatttc	ccgtcgcaaa	tgctaagaac	1560
gggtaacaac	ttccagttca	gctacgagtt	tgagaacgta	cctttccata	gcagctacgc	1620
tcacagccaa	agcctggacc	gactaatgaa	tccactcatc	gaccaatact	tgtactatct	1680
ctcaaagact	attaacgggt	ctggacagaa	tcaacaaacg	ctaaaattca	gtgtggccgg	1740
accagcaac	atggctgtcc	aggggaagaa	ctacatacct	ggaccagct	accgacaaca	1800
acgtgtctca	accactgtga	ctcaaaacaa	caacagcgaa	tttgcttggc	ctggagcttc	1860
ttcttgggct	ctcaatggac	gtaatagctt	gatgaatcct	ggacctgcta	tggccagcca	1920
caaagaagga	gaggaccgtt	tctttccttt	gtctggatct	ttaatttttg	gcaaacaagg	1980
aactggaaga	gacaacgtgg	atgcggacaa	agtcattgata	accaacgaag	aagaaattaa	2040
aactactaac	ccggtagcaa	cggagtccta	tggacaagtg	gccacaaacc	accagagtgc	2100
ccaagcacag	gcgcagaccg	gctgggttca	aaaccaagga	atacttcggg	gtatggtttg	2160
gcaggacaga	gatgtgtacc	tgcaaggacc	catttgggcc	aaaattcctc	acacgtacgg	2220
caactttcac	ccttctccgc	tgatgggagg	gtttggaatg	aagcaccgcg	ctcctcagat	2280
cctcatcaaa	aacacacctg	tacctgcgga	tcctccaacg	gccttcaaca	aggacaagct	2340
gaactctttc	atcaccacgt	attctactgg	ccaagtcagc	gtggagatcg	agtgggagct	2400

gcagaaggaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 27

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 27

ccatcgacgt cagacgcgga agcttcgac aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgtcgtcaaa aaggcgtatc agaaactgtg ctacattcat	240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctgggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccct caaccaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gaccgggcaa cggactcgac aagggggagc cggtaaacgc	540
agtagacgcg gcggccctcg agcagacaaa ggcctacgac cagcagctca aggccggaga	600
caaccgtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcggggcagc agtcttccag gccaaaaaga ggcttcttga	720
acctcttggt ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggcctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgcag cccctcagg tgtgggatct ctacaatgg cttcagggtg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg	1080
ggcctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaat gacaacgctt acttcggcta cagcaccccc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg	1260
atttcggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtcggctcac gagggctgcc tcccgcggtt	1440
cccagcggac gttttcatga ttctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgcaaa tgctaagaac	1560

```

gggtaacaac ttccagttca gctacgagtt tgagaacgta cttttccata gcagctacgc 1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct 1680
ctcaaagact attaacgggt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg 1740
accagcaac atggctgtcc agggaagaaa ctacatacct ggaccagct accgacaaca 1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctggagcttc 1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca 1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg 1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa 2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc 2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatggtttg 2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggccc aaaattcctc acacggacgg 2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgcc ctctcagat 2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct 2340
gaactctttc atcaccagct attctactgg ccaagtcagc gtggagatcg agtgggagct 2400

gcagaaggaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattacaa 2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat 2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt 2580
cgtttcagtt gaactgcggc c 2601

```

<210> 28

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 28

```

ccatcgacgt cagacgcgga agcttcgata aactacgcgg acaggtagca aaacaaatgt 60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat 120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg 180
tcagaatctc aaccgtttc tgctgtcaaa aaggcgtatc agaaactgtg ctacattcat 240
cacatcatgg gaaaggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg 300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct 360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtggtggg ctttgaaacc 420
tggagcccct caacccaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct 480
tccgggttac aaataccttg gacccggcaa cggactcgac aagggggagc cgggtcaacgc 540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga 600
caaccgtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga 660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga 720

```

acctcttggg	ctggttgagg	aagcggctaa	gacggctcct	ggaaagaaga	ggcctgtaga	780
gcagtctcct	cgggaaccgg	actcctccgc	gggtattggc	aaatcgggtg	cacagcccgc	840
taaaaagaga	ctcaatttcg	gtcagactgg	cgacacagag	tcagtcccag	accctcaacc	900
aatcggagaa	cctcccgcag	ccccctcagg	tgtgggatct	cttacaatgg	cttcaggtgg	960
tggcgcacca	gtggcagaca	ataacgaagg	tgccgatgga	gtgggtagtt	cctcgggaaa	1020
ttggcattgc	gattcccaat	ggctggggga	cagagtcato	accaccagca	cccgaacctg	1080
ggccctgccc	acctacaaca	atcacctcta	caagcaaato	tccaacagca	catctggagg	1140
atcttcaa	at	gacaacgcct	acttcggcta	cagcaccccc	tgggggtatt	1200
cttcaa	at	gacttcaa	at	tgacttcaa	at	1260
cagattccac	tgccacttct	caccacgtga	ctggcagcga	ctcatcaaca	acaactgggg	1320
attcgggcct	aagcgactca	acttcaagct	cttcaacatt	caggtcaaag	aggttacgga	1380
caacaatgga	gtcaagacca	tgcgaataa	ccttaccage	acggtccagg	tcttcacgga	1440
ctcagactat	cagctcccgt	acgtgctcgg	gtcggctcac	gagggctgcc	tcccgcctgt	1500
cccagcggac	gttttcatga	ttcctcagta	cgggtatctg	acgcttaatg	atggaagcca	1560
ggccgtgggt	cggttcgtcct	tttactgcct	ggaatatttc	ccgtcgcata	tgctaagaac	1620
gggtaacaac	ttccagttca	gctacgagtt	tgagaacgta	cctttccata	gcagctacgc	1680
tcacagccaa	agcctggacc	gactaatgaa	tccactcato	gaccaatact	tgtactatct	1740
ctcaaagact	attaacgggt	ctggacagaa	tcaacaaacg	ctaaaattca	gtgtggccgg	1800
accagcaac	atggctgtcc	agggaagaaa	ctacatacct	ggaccagct	accgacaaca	1860
acgtgtctca	accactgtga	ctcaaaacaa	caacagcgaa	tttgcttggc	ctggagcttc	1920
ttcttgggct	ctcaatggac	gtaatagctt	gatgaatcct	ggacctgcta	tggccagcca	1980
caaagaagga	gaggaccgtt	tctttccttt	gtctggatct	ttaatttttg	gcaaacaagg	2040
aactggaaga	gacaacgtgg	atgcggacaa	agtcacgata	accaacgaag	aagaaattaa	2100
aactactaac	ccggtagcaa	cggagtccta	tggacaagtg	gccacaaacc	accagagtgc	2160
ccaagcacag	gcgcagaccg	gctgggttca	aaaccaagga	atacttccgg	gtatggtttg	2220
gcaggacaga	gatgtgtacc	tgcaaggacc	catttgggcc	aaaattcctc	acacggacgg	2280
caactttcac	ccttctccgc	tgatgggagg	gtttggaatg	aagcaccgcg	ctcctcagat	2340
cctcatcaaa	aacacacctg	tacctgcgga	tcctccaacg	gccttcaaca	aggacaagct	2400
gaactctttc	atcaccagct	attctactgg	ccaagtcagc	gtggagatcg	agtgggaact	2460
gcagaaggaa	aacagcaagc	gctggaaccc	ggagatccag	tacacttcca	actattacaa	2520
gtctaataat	gttgaatttg	ctgttaatac	tgaagggtga	tatagtgaac	cccgcccat	2580
tggcaccaga	tacctgactc	gtaatctgta	attgcttggt	aatcaataaa	ccgtttaatt	2601
cgtttcagtt	gaactgcggc	c				

<210> 29

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 29

ccatcgacgt cagacgggga agcttcgata aactacggcg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggg gtcaaagact gtttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgcgtcaaa aaggcgatc agaaactgtg ctacattcat	240
cacatcatgg gaaaggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccct caaccaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gaccgggcaa cggactcgac aagggggagc cggccaacgc	540
agcagacgcg gggccctcg agcacgacaa ggcctacgac cagcagctca aggcgggaga	600
caaccggtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggagagc agtctttcag gccaaaaaga ggcttcttga	720
acctcttggg ctgggtgagg aagcggctaa gacggctcct ggaaagaaga ggcctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagteccag acctcaacc	900
aatcggagaa cctcccgag cccctcagg tgtgggatct cttacaatgg ctcatgtgg	960
tggcgacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcata accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaata gacaacgcct acttcggcta cagcaccctc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtcgggtcac gagggctgcc tcccggcgtt	1440
cccagcggac gttttcatga ttcttcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct tttactgcct ggaatatctc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcata gaccaatact tgtactatct	1680
ctcaaagact attaacgggt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atgggtgtcc agggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctggagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040

aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatgggttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct	2340
gaactctttc atcaccaggt attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaacc ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 30

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 30

ccatcgacgt cagacgcgga agcttcgatc aactacgcgg acaggtacca aaacaaatgt	60
tctcgccacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgtcgtcaaa aaggcgtatc agaaactgtg ctacattcat	240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctgggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtggtggg ctttgaaacc	420
tggagcccct caacccaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gacccggcaa cggactcgac aagggggagc cgggtcaacgc	540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga	600
caacccgtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga	720
acctcttggt ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgtg cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140

```

atcttcaa at gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa 1200
cagattccac tgccacttct caccacatga ctggcagcga ctcatcaaca acaactgggg 1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga 1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga 1380
ctcagactat cagctcccgt acgtgctcgg gtccggctcac gagggctgcc tcccgcggtt 1440
cccagcggac gttttcatga ttccctcagta cgggtatctg acgcttaatg atggaagcca 1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgcaaa tgctaagaac 1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc 1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct 1680
ctcaaagact attaacggtt ctggacagaa tcaacaaacg ctaaaattca atgtggccgg 1740
accagcaac atggctgtcc agggaagaaa ctacatacct ggaccagct accgacaaca 1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa ttgctcggc ctagagcttc 1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca 1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg 1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa 2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc 2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatggtttg 2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg 2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctctcagat 2280
ctcatcaaaa aacacacctg tacctgcgga tctccaacg gccttcaaca aggacaagct 2340
gaactctttc atcaccagct attctactgg ccaagtcagc atggagatcg agtgggagct 2400
gcagaaggaa aacagcaagc gctggaaccg ggagatccag tacacttcca actattacaa 2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac ccgccccat 2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt 2580
cgtttcagtt gaactgcggc c 2601

```

<210> 31

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 31

```

ccatcgacgt cagacgcgga agcttcgac aactacgcgg acaggtacca aaacaaatgt 60

tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat 120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg 180
tcagaatctc aaccggtttc tgctgtcaaa aaggcgtatc agaaactgtg ctacattcat 240
cacatcatgg gaaagggtgc agacgcttgc actgcttgcg acctgggtcaa tgtggacttg 300

```

gatgactgtg tttctgaaca ataatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccoct caacccaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gacccggcaa cggactcgac aagggggagc cggccaacgc	540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggcgggaga	600
caaccctgtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga	720
acctcttggg ctggttgagg aagcgggtaa gacggctcct ggaaagaaga ggctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag accctcaacc	900
aatcggagaa cctcccgag cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaat gacaacgct acttcggcta cagcaccccc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcacaaaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccg acgtgctcgg gtcggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct tttactgcct ggaatatttc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cttttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcacc gaccaatact tgtactatct	1680
ctcaaagact attaacgggt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atggctgtcc agggaagaaa ctacatacct ggacccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa ttgcttggc ctagagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatggtttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct	2340
gaactctttc atcaccagat attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaacct ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac ccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttgtt aatcaataaa ccgtttaatt	2580

-- - - - -
cgtttcagtt gaactgcggc c

2601

<210> 32

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 32

ccatcgacgt cagacgcgga agcttcgata aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgtcgtcaaa aaggcgatc agaaactgtg ctacattcat	240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctgggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataatgact taaaccagggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccct caaccgaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gaccgggcaa cggactcgac aagggggagc cgggtcaacgc	540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga	600
caacccttac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga	720
acctcttggt ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggctgtaga	780
gcagtctcct caggaaccgg actcctccgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgcag cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcata accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaat gacaacgcct acttcggcta cagcaccctc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt cagggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccataaa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtcggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cttttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcata gaccaatact tgtactatct	1680

ctcaaagact	attaacggtt	ctggacagaa	tcaacaaacg	ctaaaattca	gtgtggccgg	1740
acccagcaac	atggctgtcc	aggggaagaaa	ctacatacct	ggacccagct	accgacaaca	1800
acgtgtctca	accactgtga	ctcaaaacaa	caacagcgaa	tttgcttggc	ctagagcttc	1860
ttcttgggct	ctcaatggac	gtaatagctt	gatgaatcct	ggacctgcta	tggccagcca	1920
caaagaagga	gaggaccgtt	tctttccttt	gtctggatct	ttaatTTTTg	gcaaacaagg	1980
aactggaaga	gacaacgtgg	atgcggacaa	agtcattgata	accaacgaag	aagaaattaa	2040
aactactaac	ccggtagcaa	cggagtcccta	tggacaagtg	gccacaaacc	accagagtgc	2100
ccaagcacag	gcgcagaccg	gctgggttca	aaaccaagga	atacttccgg	gtatggtttg	2160
gcaggacaga	gatgtgtacc	tgcaaggacc	catttggggc	aaaattcctc	acacggacgg	2220
caactttcac	ccttctccgc	tgatgggagg	gtttggaatg	aagcaccgcg	ctcctcagat	2280
cctcatcaaa	aacacacctg	tacctgcgga	tcctccaacg	gccttcaaca	aggacaagct	2340
gaactctttc	atcaccagct	attctactgg	ccaagtcagc	gtggagatcg	agtgggagcg	2400
gcagaaggaa	aacagcaagc	gctggaaccc	ggagatccag	tacacttcca	actattacaa	2460
gtctaataat	gttgaatttg	ctgttaatac	tgaagggtga	tatagtgaac	cccgcccat	2520
tggcaccaga	tacctgactc	gtaatctgta	attgcttggt	aatcaataaa	ccgtttaatt	2580
cgtttcagtt	gaactgcggc	c				2601

<210> 33

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 33

ccatcgacgt	cagacgcgga	agcttcgata	aactacgcgg	acaggtacca	aaacaaatgt	60
tctcgtcacg	tgggcatgaa	tctgatgctg	tttccctgca	gacaatgcga	gagactgaat	120
cagaattcaa	atatctgctt	cactcacggt	gtcaaagact	gtttagagtg	ctttcccgtg	180
tcagaatctc	aaccgtttc	tgctgtcaaa	aaggcgtatc	agaaactgtg	ctacattcat	240
cacatcatgg	gaaagggtgcc	agacgcttgc	actgcttgcg	acctgggtcaa	tgtggacttg	300
gatgactgtg	tttctgaaca	ataaatgact	taaaccaggt	atggctgccg	atggttatct	360
tccagattgg	ctcgaggaca	accttagtga	aggaattcgc	gagtgggtggg	ctttgaaacc	420
tggagcccct	caaccacaag	caaatcaaca	acatcaagac	aacgctcgag	gtcttgtgct	480
tccgggttac	aaataccttg	gacccggcaa	cggactcgac	aagggggagc	cgggtcaacgc	540
agcagacgcg	gcggccctcg	agcacgacaa	ggcctacgac	cagcagctca	aggccggaga	600
caacccttac	ctcaagtaca	accacgccga	cgccgagttc	caggagcggc	tcaaagaaga	660
tacgtctttt	gggggcaacc	tcgggcgagc	agtcttccag	gccaaaaaga	ggcttcttga	720
acctcttggt	ctggttgagg	aagcggctaa	gacggctcct	ggaaagaaga	ggcctgtaga	780
gcagtctcct	caggaaccgg	actcctccgc	gggtattggc	aaatcgggtg	cacagcccgc	840

taaaaagaga ctcaattttcg gtcagactgg cgacacagag tcagtcccag accctcaacc	900
aatcggagaa cctcccgcag cccctcagg tgtgggatct cttacaatgg cttcaggtgg	960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaat gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcacaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccataa ccttaccagc acggtccagg tcttcgcgga	1380
ctcagactat cagctcccgat acgtgctcgg gtcggctcac gagggctgcc tcccgcggtt	1440
cccagcggac gttttcatga ttccctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatctt ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct	1680
ctcaaagact attaacgggt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atggctgtcc aggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa tttgcttggc ctagagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatggtttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctcctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct	2340
gaactctttc atcaccagat attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaacc ggagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 34

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 34

ccatcgacgt cagacgcgga agcttcgac aactacgcgg acaggtacca aaacaaatgt	60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat	120
cagaattcaa atatctgctt cactcacggt gtcaaagact gttagagtg ctttcccgtg	180
tcagaatctc aacccgtttc tgcgtcaaa aaggcgtatc agaaactgtg ctacattcat	240
cacatcatgg gaaaggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg	300
gatgactgtg tttctgaaca ataaatgact taaaccaggt atggctgccg atggttatct	360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc	420
tggagcccct caaccgaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct	480
tccgggttac aaataccttg gaccgggcaa cggactcgac aagggggagc cgggtcaacgc	540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga	600
caaccgtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga	660
tacgtctttt gggggcaacc tcgggcgagc agtcttcag gccaaaaaga ggcttcttga	720
acctcttggg ctggttgagg aagcgggctaa gacggctcct ggaaagaaga ggcctgtaga	780
gcagtctcct caggaaccgg actcctcgc gggatttggc aaatcgggtg cacagcccgc	840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc	900
aatcggagaa cctcccgag cccctcagc tgtgggatct cttacaatgg cttcaggtgg	960
tggcgacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa	1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg	1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg	1140
atcttcaaata gacaacgcct acttcggcta cagcaccctc tgggggtatt ttgacttcaa	1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg	1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga	1320
caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtcggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttccctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct tttactgcct ggaatatctc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct	1680
ctcaaagact attaacggtt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atggctgtcc aggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa attgcttggc ctagagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccagtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttcggg gtatggtttg	2160

```

gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg      2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctcctcagat      2280
cctcatcaaa aacacacctg tacctgcgga tcctccaacg gccttcaaca aggacaagct      2340
gaactctttc atcaccacgt attctactgg ccaagtcagc gtggagatcg agtgggagct      2400
gcagaaggaa aacagcaagc gctggaaccc ggagatccag tacacttcca actattgcaa      2460
gtctaataat gttgaatttg ctgttaatac tgaaggtgta tatagtgaac cccgccccat      2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt      2580

cgtttcagtt gaactgcggc c                                              2601

```

<210> 35

<211> 2601

<212> DNA

<213> Artificial Sequence

<220>

<223> novel AAV isolate

<400> 35

```

ccatcgacgt cagacgcgga agcttcgata aactacgcgg acaggtacca aaacaaatgt      60
tctcgtcacg tgggcatgaa tctgatgctg tttccctgca gacaatgcga gagactgaat      120
cagaattcaa atatctgctt cactcacggt gtcaaagact gtttagagtg ctttcccgtg      180
tcagaatctc aaccgcgttc tgcgtcaaa aaggcgtatc agaaactgtg ctacattcat      240
cacatcatgg gaaagggtgcc agacgcttgc actgcttgcg acctggtcaa tgtggacttg      300
gatgtctgtg tttctgaaca ataaatgact taaaccagggt atggctgccg atggttatct      360
tccagattgg ctcgaggaca accttagtga aggaattcgc gagtgggtggg ctttgaaacc      420
tggagcccct caaccgaagg caaatcaaca acatcaagac aacgctcgag gtcttgtgct      480
tccgggttac aaataccttg gaccgggcaa cggactcgat aagggggagc cgggtcaacgc      540
agcagacgcg gcggccctcg agcacgacaa ggcctacgac cagcagctca aggccggaga      600
caaccogtac ctcaagtaca accacgccga cgccgagttc caggagcggc tcaaagaaga      660
tacgtctttt gggggcaacc tcgggcgagc agtcttccag gccaaaaaga ggcttcttga      720
acctcttggt ctggttgagg aagcggctaa gacggctcct ggaaagaaga ggctgtaga      780
gcagtctcct caggaaccgg actcctccgc gggattggc aaatcgggtg cacagcccgc      840
taaaaagaga ctcaatttcg gtcagactgg cgacacagag tcagtcccag acctcaacc      900
aatcggagaa cctcccgcag cccctcagg tgtgggatct cttacaatgg cttcaggtgg      960
tggcgcacca gtggcagaca ataacgaagg tgccgatgga gtgggtagtt cctcgggaaa     1020
ttggcattgc gattcccaat ggctggggga cagagtcac accaccagca cccgaacctg     1080
ggccctgccc acctacaaca atcacctcta caagcaaata tccaacagca catctggagg     1140
atcttcaaat gacaacgcct acttcggcta cagcaccccc tgggggtatt ttgacttcaa     1200
cagattccac tgccacttct caccacgtga ctggcagcga ctcatcaaca acaactgggg     1260
attccggcct aagcgactca acttcaagct cttcaacatt caggtcaaag aggttacgga     1320

```

caacaatgga gtcaagacca tcgccaataa ccttaccagc acggtccagg tcttcacgga	1380
ctcagactat cagctcccgt acgtgctcgg gtccggctcac gagggctgcc tcccgccgtt	1440
cccagcggac gttttcatga ttccctcagta cgggtatctg acgcttaatg atggaagcca	1500
ggccgtgggt cgttcgtcct ttactgcct ggaatatttc ccgtcgcaaa tgctaagaac	1560
gggtaacaac ttccagttca gctacgagtt tgagaacgta cctttccata gcagctacgc	1620
tcacagccaa agcctggacc gactaatgaa tccactcatc gaccaatact tgtactatct	1680
ctcaaagact attaacggtt ctggacagaa tcaacaaacg ctaaaattca gtgtggccgg	1740
accagcaac atggctgtcc aggaagaaa ctacatacct ggaccagct accgacaaca	1800
acgtgtctca accactgtga ctcaaaacaa caacagcgaa ttgcttggc ctagagcttc	1860
ttcttgggct ctcaatggac gtaatagctt gatgaatcct ggacctgcta tggccagcca	1920
caaagaagga gaggaccgtt tctttccttt gtctggatct ttaatttttg gcaaacaagg	1980
aactggaaga gacaacgtgg atgcggacaa agtcatgata accaacgaag aagaaattaa	2040
aactactaac ccggtagcaa cggagtccta tggacaagtg gccacaaacc accagagtgc	2100
ccaagcacag gcgcagaccg gctgggttca aaaccaagga atacttccgg gtatggtttg	2160
gcaggacaga gatgtgtacc tgcaaggacc catttgggcc aaaattcctc acacggacgg	2220
caactttcac ccttctccgc tgatgggagg gtttggaatg aagcaccgc ctccctcagat	2280
cctcatcaaa aacacacctg tacctgcgga tccccaacg gccttcaaca aggacaagct	2340
gaactctttc atcaccagct attctactgg ccaagtcagc gtggagatcg agtgggagct	2400
gcagaaggaa aacagcaagc gctggaacc ccgagatccag tacacttcca actattacaa	2460
gtctaataat gttgaatttg ctgttaatac tgaagggtga tatagtgaac cccgccccat	2520
tggcaccaga tacctgactc gtaatctgta attgcttggt aatcaataaa ccgtttaatt	2580
cgtttcagtt gaactgcggc c	2601

<210> 36

<211> 145

<212> DNA

<213> Artificial Sequence

<220>

<223> AAV2 5' ITR

<400> 36

ttggccactc cctctctgcg cgctcgctcg ctactgagg ccgggcgacc aaaggctgcc	60
cgacgccccg gctttgcccg ggccgcctca gtgagcgagc gagcgcgag agagggagtg	120
gccaaactcca tcactagggg ttccct	145

<210> 37

<211> 145

<212> DNA

<213> Artificial Sequence

<220>

<223> AAV2 3' ITR

<400> 37

aggaacccct agtgatggag ttggccactc cctctctgcg cgctcgctcg ctcaactgagg	60
ccgggcgacc aaaggctcgcc cgacgcccgg gctttgcccg ggccggcctca gtgagcgagc	120
gagcgcgag agagggagtg gccaa	145

<210> 38

<211> 167

<212> DNA

<213> Artificial Sequence

<220>

<223> AAV5 5' ITR

<400> 38

ctctcccccc tgtcgcgttc gctcgctcgc tggctcgttt ggggggggtgg cagctcaaag	60
agctgccaga cgacggccct ctggccgctcg cccccccaaa cgagccagcg agcgagcgaa	120
cgcgacaggg gggagagtgc cacactctca agcaaggggg ttttgta	167

<210> 39

<211> 167

<212> DNA

<213> Artificial Sequence

<220>

<223> AAV5 3' ITR

<400> 39

tacaaaacct ccttgcttga gagtgtggca ctctcccccc tgtcgcgttc gctcgctcgc	60
tggctcgttt ggggggggtgg cagctcaaag agctgccaga cgacggccct ctggccgctcg	120
cccccccaaa cgagccagcg agcgagcgaa cgcgacaggg gggagag	167

<210> 40

<211> 621

<212> PRT

<213> Artificial Sequence

<220>

<223> AAV2 Rep

<400> 40

Met	Pro	Gly	Phe	Tyr	Glu	Ile	Val	Ile	Lys	Val	Pro	Ser	Asp	Leu	Asp
1				5					10					15	
Glu	His	Leu	Pro	Gly	Ile	Ser	Asp	Ser	Phe	Val	Asn	Trp	Val	Ala	Glu
			20					25					30		

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
 35 40 45
 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu
 50 55 60
 Thr Glu Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val
 65 70 75 80
 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Met His Val Leu Val Glu
 85 90 95
 Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile
 100 105 110
 Arg Glu Lys Leu Ile Gln Arg Ile Tyr Arg Gly Ile Glu Pro Thr Leu
 115 120 125
 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly
 130 135 140
 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys
 145 150 155 160
 Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Gln Tyr Leu
 165 170 175
 Ser Ala Cys Leu Asn Leu Thr Glu Arg Lys Arg Leu Val Ala Gln His
 180 185 190
 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Gln Asn
 195 200 205
 Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr
 210 215 220
 Met Glu Leu Val Gly Trp Leu Val Asp Lys Gly Ile Thr Ser Glu Lys
 225 230 235 240
 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala
 245 250 255
 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys
 260 265 270
 Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val Gly Gln Gln
 275 280 285
 Pro Val Glu Asp Ile Ser Ser Asn Arg Ile Tyr Lys Ile Leu Glu Leu
 290 295 300
 Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
 305 310 315 320
 Thr Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala

```

-----
325                               330                               335

Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Thr Val Pro
340                               345                               350

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
355                               360                               365

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
370                               375                               380

Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
385                               390                               395                               400

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
405                               410                               415

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
420                               425                               430

Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
435                               440                               445

Glu Leu Thr Arg Arg Leu Asp His Asp Phe Gly Lys Val Thr Lys Gln
450                               455                               460

Glu Val Lys Asp Phe Phe Arg Trp Ala Lys Asp His Val Val Glu Val
465                               470                               475                               480

Glu His Glu Phe Tyr Val Lys Lys Gly Gly Ala Lys Lys Arg Pro Ala
485                               490                               495

Pro Ser Asp Ala Asp Ile Ser Glu Pro Lys Arg Val Arg Glu Ser Val
500                               505                               510

Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Ser Ile Asn Tyr Ala Asp
515                               520                               525

Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn Leu Met Leu
530                               535                               540

Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Ser Asn Ile Cys
545                               550                               555                               560

Phe Thr His Gly Gln Lys Asp Cys Leu Glu Cys Phe Pro Val Ser Glu
565                               570                               575

Ser Gln Pro Val Ser Val Val Lys Lys Ala Tyr Gln Lys Leu Cys Tyr
580                               585                               590

Ile His His Ile Met Gly Lys Val Pro Asp Ala Cys Thr Ala Cys Asp
595                               600                               605

Leu Val Asn Val Asp Leu Asp Asp Cys Ile Phe Glu Gln
610                               615                               620

```

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US20130096182A1 \[0009\] \[0051\] \[0066\] \[0066\] \[0067\] \[0111\] \[0165\] \[0165\]](#)
- [US2013096182A \[0012\]](#)
- [US20110294218A1 \[0066\] \[0066\] \[0067\]](#)
- [US5474935A \[0142\]](#)
- [US62209862B \[0211\]](#)
- [US62063587B \[0211\]](#)
- [US62054899B \[0211\]](#)

Non-patent literature cited in the description

- KHAN et al.Nature Protocols, 2011, vol. 6, 4482-501 [\[0012\]](#)
- MILLER et al.Nature Biotechnology, 2006, vol. 24, 81022-1026 [\[0012\]](#)
- SMITH et al.Mol Ther., 2014, vol. 22, 91625-34 [\[0051\] \[0177\]](#)
- PEARSONLIPMANProc. Natl. Acad. Sci. USA, 1988, vol. 85, 2444- [\[0068\] \[0133\]](#)
- ZINN et al.In Silico Reconstruction of the Viral Evolutionary Lineage Yields a Potent Gene Therapy VectorCell Reports, 2015, vol. 12, 1056-1068 [\[0072\]](#)
- NILSSON et al.EMBO J., 1985, vol. 4, 1075- [\[0136\]](#)
- NILSSON et al.Methods Enzymol., 1991, vol. 198, 3- [\[0136\]](#)
- SMITHJOHNSONGene, 1988, vol. 67, 31- [\[0136\]](#)
- GRUSSENMEYER et al.Proc. Natl. Acad. Sci. USA, 1985, vol. 82, 7952-4 [\[0136\]](#)
- HOPP et al.Biotechnology, 1988, vol. 6, 1204-10 [\[0136\]](#)
- FORD et al.Protein Expression and Purification, 1991, vol. 2, 95-107 [\[0136\]](#)
- SHI et al.Human Gene Therapy, 2001, vol. 12, 1697-1711 [\[0139\]](#)
- RABINOWITZ et al.Virology, 1999, vol. 265, 274-285 [\[0139\]](#)
- NICKLIN et al.Molecular Therapy, 2001, vol. 4, 174-181 [\[0139\]](#)
- WU et al.J. Virology, 2000, vol. 74, 8635-8647 [\[0139\]](#)

- **GRIFMAN et al.** Molecular Therapy, 2001, vol. 3, 964-974 [0139]
- Remington's Pharmaceutical Sciences Mack Publishing Co. [0140]
- **A. GENNARO** Remington: The Science and Practice of Pharmacy Lippincott, Williams, & Wilkins 2000 0000 [0140]
- Pharmaceutical Dosage Forms and Drug Delivery Systems Lippincott, Williams, & Wilkins 1999 0000 [0140]
- **A. H. KIBBE et al.** Handbook of Pharmaceutical Excipients Amer. Pharmaceutical Assoc 2000 0000 [0140]
- **OKADA** Gene Ther., 1996, vol. 3, 957-964 [0142]
- **NAGAI et al.** Nat Biotechnol, 2002, vol. 20, 187-90 [0178]
- **BAINBRIDGE, J.W. SMITH, A.J. BARKER, S.S. ROBBIE, SHENDERSON, R. BALAGGAN, K. VISWANATHAN, A. HOLDER, G.E. STOCKMAN, A. TYLER, N.** Effect of gene therapy on visual function in Leber's congenital amaurosis N Engl J Med, 2008, vol. 358, 2231-2239 [0210]
- **BATCHU, R.B. SHAMMAS, M.A. WANG, J.Y. FREEMAN, J. ROSEN, N. MUNSHI, N.C.** Adeno-associated virus protects the retinoblastoma family of proteins from adenoviral-induced functional inactivation Cancer Res, 2002, vol. 62, 2982-2985 [0210]
- **BELL, P. WANG, L. LEBHERZ, C. FLIEDER, D.B. BOVE, M.S. WU, D. GAO, G.P. WILSON, J.M. WIVEL, N.A.** No evidence for tumorigenesis of AAV vectors in a large-scale study in mice Mol Ther, 2005, vol. 12, 299-306 [0210]
- **BERNS, K.I. GIRAUD, C.** Biology of adeno-associated virus Curr Top Microbiol Immunol, 1996, vol. 218, 1-23 [0210]
- **BIFFI, A. CESANI, M.** Human hematopoietic stem cells in gene therapy: preclinical and clinical issues Curr Gene Ther, 2008, vol. 8, 135-146 [0210]
- **BRANTLY, M.L. CHULAY, J.D. WANG, L. MUELLER, C. HUMPHRIES, M. SPENCER, L.T. ROUHANI, F. CONLON, T.J. CALCEDO, R. BETTS, M.R.** Sustained transgene expression despite T lymphocyte responses in a clinical trial of rAAV1-AAT gene therapy Proc Natl Acad Sci U S A., 2009, [0210]
- **CHATTERJEE, S. JOHNSON, P.R. WONG, K.K., JR.** Dual-target inhibition of HIV-1 in vitro by means of an adeno-associated virus antisense vector Science, 1992, vol. 258, 1485-1488 [0210]
- **CHATTERJEE, S. LI, W. WONG, C.A. FISHER-ADAMS, G. LU, D. GUHA, M. MACER, J.A. FORMAN, S.J. WONG, K.K., JR.** Transduction of primitive human marrow and cord blood-derived hematopoietic progenitor cells with adeno-associated virus vectors Blood, 1999, vol. 93, 1882-1894 [0210]
- **CHATTERJEE, S. WONG, K.K.** Adeno-Associated Viral Vectors for the Delivery of Antisense RNAs METHODS -LONDON- A COMPANION TO METHODS IN ENZYMOLOGY, 1993, vol. 5, 1- [0210]
- **CIDECIYAN, A.V. HAUSWIRTH, W.W. ALEMAN, T.S. KAUSHAL, S. SCHWARTZ, S.B. BOYE, S.L. WINDSOR, E.A. CONLON, T.J. SUMAROKA, A. PANG, J.J.** Human RPE65 gene therapy for Leber congenital amaurosis: persistence of early visual improvements and safety at 1 year Hum Gene Ther, 2009, vol. 20, 999-1004 [0210]
- **EINERHAND, M.P. ANTONIOU, M. ZOLOTUKHIN, S. MUZYCZKA, N. BERNES, K.I. GROSVELD, F. VALERIO, D.** Regulated high-level human beta-globin gene

expression in erythroid cells following recombinant adeno-associated virus-mediated gene transfer *Gene Ther*, 1995, vol. 2, 336-343 [\[0210\]](#)

- FISHER-ADAMS, G.WONG, K.K., JR.PODSAKOFF, G.FORMAN, S.J.CHATTERJEE, S.Integration of adeno-associated virus vectors in CD34+ human hematopoietic progenitor cells after transduction *Blood*, 1996, vol. 88, 492-504 [\[0210\]](#)
- FLOTTE, T.R.BRANTLY, M.L.SPENCER, L.T.BYRNE, B.J.SPENCER, C.T.BAKER, D.J.HUMPHRIES, M.Phase I trial of intramuscular injection of a recombinant adeno-associated virus alpha 1-antitrypsin (rAAV2-CB-hAAT) gene vector to AAT-deficient adults *Hum Gene Ther*, 2004, vol. 15, 93-128 [\[0210\]](#)
- GAO, G.VANDENBERGHE, L.H.ALVIRA, M.R.LU, Y.CALCEDO, R.ZHOU, XWILSON, J.M.Clades of Adeno-associated viruses are widely disseminated in human tissues *J Virol*, 2004, vol. 78, 6381-6388 [\[0210\]](#)
- HACEIN-BEY-ABINA, S.VON KALLE, C.SCHMIDT, M.LE DEIST, F.WULFFRAAT, N.MCINTYRE, E.RADFORD, I.VILLEVAL, J.LFRASER, C.C.CAVAZZANA-CALVO, M.A serious adverse event after successful gene therapy for X-linked severe combined immunodeficiency *N Engl J Med*, 2003, vol. 348, 255-256 [\[0210\]](#)
- HAN, Z.ZHONG, L.MAINA, N.HU, Z.LI, X.CHOUTHAI, N.S.BISCHOF, D.WEIGEL-VAN AKEN, K.A.SLAYTON, W.B.YODER, M.C.Stable integration of recombinant adeno-associated virus vector genomes after transduction of murine hematopoietic stem cells *Hum Gene Ther*, 2008, vol. 19, 267-278 [\[0210\]](#)
- JAYANDHARAN, G.R.ZHONG, L.LI, B.KACHNIARZ, B.SRIVASTAVA, A.Strategies for improving the transduction efficiency of single-stranded adeno-associated virus vectors in vitro and in vivo *Gene Ther*, 2008, vol. 15, 1287-1293 [\[0210\]](#)
- KAPLITT, M.G.FEIGIN, A.TANG, CFITZSIMONS, H.L.MATTIS, P.LAWLOR, P.A.BLAND, R.J.YOUNG, D.STRYBING, KEIDELBERG, D.Safety and tolerability of gene therapy with an adeno-associated virus (AAV) borne GAD gene for Parkinson's disease: an open label, phase I trial *Lancet*, 2007, vol. 369, 2097-2105 [\[0210\]](#)
- KELLS, A.PHADACZEK, P.YIN, D.BRINGAS, J.VARENIKA, VFORSAYETH, J.BANKIEWICZ, K.S.Efficient gene therapy-based method for the delivery of therapeutics to primate cortex *Proc Natl Acad Sci U S A*, 2009, vol. 106, 2407-2411 [\[0210\]](#)
- KESSLER, P.D.PODSAKOFF, G.M.CHEN, XMCQUISTON, S.A.COLOSI, P.C.MATELIS, L.A.KURTZMAN, G.J.BYRNE, B.J.Gene delivery to skeletal muscle results in sustained expression and systemic delivery of a therapeutic protein *Proc Natl Acad Sci U S A*, 1996, vol. 93, 14082-14087 [\[0210\]](#)
- MANNO, C.S.CHEW, A.JHUTCHISON, SLARSON, P.J.HERZOG, R.W.ARRUDA, V.R.TAI, S.JRAGNI, M.VTHOMPSON, AOZELO, M.AAV-mediated factor IX gene transfer to skeletal muscle in patients with severe hemophilia B *Blood*, 2003, vol. 101, 2963-2972 [\[0210\]](#)
- MCCORMACK, M.P.RABBITS, T.H.Activation of the T-cell oncogene LMO2 after gene therapy for X-linked severe combined immunodeficiency *N Engl J Med*, 2004, vol. 350, 913-922 [\[0210\]](#)
- MILLER, D.G.ADAM, M.A.MILLER, A.D.Gene transfer by retrovirus vectors occurs only in cells that are actively replicating at the time of infection *Mol Cell Biol*, 1990, vol. 10,

4239-4242 [0210]

- PAZ, H.WONG, C.A.LI, W.SANTAT, L.WONG, K.K.CHATTERJEE, S.Quiescent subpopulations of human CD34-positive hematopoietic stem cells are preferred targets for stable recombinant adeno-associated virus type 2 transductionHum Gene Ther, 2007, vol. 18, 614-626 [0210]
- PETRS-SILVA, H.DINCULESCU, A.LI, Q.MIN, S.H.CHiodo, V.PANG, J.J.ZHONG, L.ZOLOTUKHIN, S.SRIVASTAVA, A.LEWIN, A.S.High-efficiency transduction of the mouse retina by tyrosine-mutant AAV serotype vectorsMol Ther, 2009, vol. 17, 463-471 [0210]
- PODSAKOFF, G.WONG, K.K., JR.CHATTERJEE, S.Efficient gene transfer into nondividing cells by adeno-associated virus-based vectorsJ Virol, 1994, vol. 68, 5656-5666 [0210]
- PONNAZHAGAN, S.YODER, M.C.SRIVASTAVA, A.Adeno-associated virus type 2-mediated transduction of murine hematopoietic cells with long-term repopulating ability and sustained expression of a human globin gene in vivoJ Virol, 1997, vol. 71, 3098-3104 [0210]
- RAJ, K.OGSTON, P.BEARD, P.Virus-mediated killing of cells that lack p53 activityNature, 2001, vol. 412, 914-917 [0210]
- SANTAT, L.PAZ, HWONG, C.LI, L.MACER, J.FORMAN, SWONG, K.K.CHATTERJEE, S.Recombinant AAV2 transduction of primitive human hematopoietic stem cells capable of serial engraftment in immune-deficient miceProc Natl Acad Sci U S A, 2005, vol. 102, 11053-11058 [0210]
- SRIVASTAVA, A.Gene delivery to human and murine primitive hematopoietic stem and progenitor cells by AAV2 vectorsMethods Mol Biol, 2004, vol. 246, 245-254 [0210]
- TOWNE, C.SCHNEIDER, B.L.KIERAN, D.REDMOND, D.E., JR.AEBISCHER, P.Efficient transduction of non-human primate motor neurons after intramuscular delivery of recombinant AAV serotype 6Gene Ther., 2009, [0210]
- ZHONG, L.CHEN, L.LI, Y.QING, K.WEIGEL-KELLEY, K.ACHAN, R.J.YODER, M.C.SRIVASTAVA, A.Self-complementary adeno-associated virus 2 (AAV)-T cell protein tyrosine phosphatase vectors as helper viruses to improve transduction efficiency of conventional single-stranded AAV vectors in vitro and in vivoMol Ther, 2004, vol. 10, 950-957 [0210]
- ZHONG, L.LI, B.JAYANDHARAN, GMAH, C.SGOVINDASAMY, LAGBANDJEMCKENNA, M.HERZOG, R.WWEIGEL-VAN AKEN, K.A.HOBBS, J.A.ZOLOTUKHIN, S.Tyrosine phosphorylation of AAV2 vectors and its consequences on viral intracellular trafficking and transgene expressionVirology, 2008, vol. 381, 194-202 [0210]
- ZHONG, L.LI, B.MAH, C.S.GOVINDASAMY, L.AGBANDJE-MCKENNA, MCOOPER, M.HERZOG, R.W.ZOLOTUKHIN, I.WARRINGTON, K.H., JR.WEIGEL-VAN AKEN, K.A.Next generation of adeno-associated virus 2 vectors: point mutations in tyrosines lead to high-efficiency transduction at lower dosesProc Natl Acad Sci U S A, 2008, vol. 105, 7827-7832 [0210]
- ZHONG, L.LI, W.YANG, Z.QING, K.TAN, MHANSEN, J.LI, Y.CHEN, L.CHAN, R.J.BISCHOF, D.Impaired nuclear transport and uncoating limit recombinant adeno-

associated virus 2 vector-mediated transduction of primary murine hematopoietic cellsHum Gene Ther, 2004, vol. 15, 1207-1218 [\[0210\]](#)

- **ZHONG, L.ZHAO, W.WU, J.LI, B.ZOLOTUKHIN, S.GOVINDASAMY, L.AGBANDJEMCKENNA, M.SRIVASTAVA, A.**A dual role of EGFR protein tyrosine kinase signaling in ubiquitination of AAV2 capsids and viral second-strand DNA synthesisMol Ther, 2007, vol. 15, 1323-1330 [\[0210\]](#)
- **ZHOU, S.Z.BROXMEYER, H.E.COOPER, S.HARRINGTON, M.A.SRIVASTAVA, A.**Adeno-associated virus 2-mediated gene transfer in murine hematopoietic progenitor cellsExp Hematol, 1993, vol. 21, 928-933 [\[0210\]](#)
- **KHAN, I.GHIRATA, R.K.RUSSELL, D.W.**AAV-mediated gene targeting methods for human cellsNat Protoc., 2011, vol. 4, 482-501 [\[0210\]](#)
- **KHOTIN, R.M.LINDEN, R.M.BERNS, K.I.**Characterization of a preferred site on human chromosome 19q for integration of adeno-associated virus DNA by non-homologous recombinationEMBO, 1992, vol. 13, 5071-8 [\[0210\]](#)
- **GIRAUD, C.WINOCOUR, E.BERNS, K.I.**Site-specific integration by adeno-associated virus is directed by a cellular DNA sequenceProc. Natl Acad Sci USA, 1994, vol. 21, 10039-43 [\[0210\]](#)
- **LINDEN, R.M.WARD, P.GIRAUD, C.WINOCOUR, E.BERNS, K.I.**Site-specific integration by adeno-associated virusProc. Natl Acad Sci USA, 1996, vol. 21, 11288-94 [\[0210\]](#)
- **LINDEN, R.M.WINOCOUR, E.BERNS, K.I.**The recombination signals for adeno-associated virus site-specific integrationProc. Natl Acad Sci USA, 1996, vol. 15, 7966-72 [\[0210\]](#)
- **XU, L.O'MALLEY, T.SANDS, M.S.WANG, B.MEYERROSE, T.HASKINS, M.E.PONDER, K.P.**In Vivo Transduction of Hematopoietic Stem Cells After Neonatal Intravenous Injection of an Amphotrophic Retroviral Vector in MiceMol. Ther., 2004, vol. 10, 137-44 [\[0210\]](#)
- **WANG, C.X.SATHER, B.D.WANG, X.ADAIR, J.KHAN, I.SINGH, S.LANG, S.ADAMS, A.CURINGA, G.KIEM, H-P.**Rapamycin relieves lentiviral vector transduction resistance in human and mouse hematopoietic stem cellsBlood, 2014, vol. 124, 6913-23 [\[0210\]](#)
- **CARBONARO, D.A.JIN, X.PETERSEN, D.WANG, XDOREY, F.KIL, K.S.ALDRICH, M.BLACKBURN, M.R.KELLEMS, R.E.KOHN, D.B.**In Vivo Transduction by Intravenous Injection of a Lentiviral Vector Expressing Human ADA into Neonatal ADA Gene Knockout Mice: A Novel Form of Enzyme Replacement Therapy for ADA DeficiencyMol. Ther., 2006, vol. 13, 61110-20 [\[0210\]](#)
- **SMITH, L.J.UL-HASAN, T.CARVAINES, S.K.VAN VLIET, K.YANG, E.WONG, JR, K.K.AGBANDJE-MCKENNA, M.CHATTERJEE, S.**Gene Transfer Properties and Structural Modeling of Human Stem Cell-derived AAVMol. Ther., 2014, vol. 22, 91625-1634 [\[0210\]](#)

Patentkrav

1. Ex vivo-fremgangsmåde til redigering af et mållocus i et mammaliakromosom, hvilken fremgangsmåde omfatter transduktion af en mammaliacelle, der omfatter mammaliakromosomet, med et replikationsdefekt adenoassocieret virus (AAV), der omfatter
- a) et AAV-clade F-capsid; og
 - b) et korrektionsgenom, der omfatter:
 - (i) et redigeringselement til insertion af et eller flere nukleotider, deletion af et eller flere nukleotider eller substitution af et eller flere nukleotider i forhold til mållocuset, hvor redigeringselementet er valgt blandt en internukleotidbinding eller en nukleotidsekvens til integrering i mållocuset;
 - (ii) en 5'-homolog arm-nukleotidsekvens 5' for redigeringselementet, der har homologi til en 5'-region af mammaliakromosomet i forhold til mållocuset; og
 - (iii) en 3'-homolog arm-nukleotidsekvens 3' for redigeringselementet, der har homologi til en 3'-region af mammaliakromosomet i forhold til mållocuset,
- Hvor cellen transduceres uden co-transduktion eller co-administration af en exogen nuklease eller en nukleotidsekvens, der koder for en exogen nuklease, hvor AAV-clade F-capsidet omfatter:
- (i) et capsidprotein, der omfatter en aminosyresekvens, der har 100 % sekvensidentitet med aminosyrerne 203 til 736 af en hvilken som helst af SEQ ID NO: 2-17;
 - (ii) et capsidprotein, der omfatter en aminosyresekvens, der har 100 % sekvensidentitet med aminosyrerne 138 til 736 af en hvilken som helst af SEQ ID NO: 2-17; og/eller
 - (iii) et capsidprotein, der omfatter en aminosyresekvens, der har 100 % sekvensidentitet med aminosyrerne 1 til 736 af en hvilken som helst af SEQ ID NO: 2-17.
2. Ex vivo-fremgangsmåde ifølge krav 1, hvor
- (a) mammaliacellen er en myoblast, en endotelcelle, en levercelle, en fibroblast, en brystcelle, en lymfocyt eller en retinalcelle; og/eller

(b) mammaliacellen er en stamcelle, eventuelt hvor stamcellen er en hæmatopoietisk stamcelle, en stamcelle fra navlestrengsblod eller en stamcelle fra perifert blod, og yderligere eventuelt hvor stamcellen er en CD34⁺-hæmatopoietisk stamcelle.

3. Replikationsdefekt adenoassocieret virus (AAV) til anvendelse i en fremgangsmåde til behandling af en sygdom eller forstyrrelse hos et individ, hvilken fremgangsmåde omfatter administrering af det replikationsdefekte AAV til individet uden co-transduktion eller co-administration af en exogen nuklease eller en nukleotidsekvens, der koder for en exogen nuklease,

hvor det replikationsdefekte AAV omfatter:

a) et AAV-clade F-capsid; og

b) et korrektionsgenom, der omfatter:

(i) et redigeringselement til insertion af et eller flere nukleotider, deletion af et eller flere nukleotider eller substitution af et eller flere nukleotider i forhold til mållocuset, hvor redigeringselementet er valgt blandt en internukleotidbinding eller en nukleotidsekvens til integrering i et mållocus i et mammaliakromosom;

(ii) en 5'-homolog arm-nukleotidsekvens 5' for redigeringselementet, der har homologi til en 5'-region af mammaliakromosomet i forhold til mållocuset; og

(iii) en 3'-homolog arm-nukleotidsekvens 3' for redigeringselementet, der har homologi til en 3'-region af mammaliakromosomet i forhold til mållocuset,

hvor AAV-clade F-capsidet omfatter:

(i) et capsidprotein, der omfatter en aminosyresekvens, der har 100 % sekvensidentitet med aminosyrerne 203 til 736 af en hvilken som helst af SEQ ID NO: 2-17;

(ii) et capsidprotein, der omfatter en aminosyresekvens, der har 100 % sekvensidentitet med aminosyrerne 138 til 736 af en hvilken som helst af SEQ ID NO: 2-17; og/eller

(iii) et capsidprotein, der omfatter en aminosyresekvens, der har 100 % sekvensidentitet med aminosyrerne 1 til 736 af en hvilken som helst af SEQ ID NO: 2-17.

4. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af kravene 1-3, hvor
korrektionsgenomet mangler en promotor, der er operativt
5 koblet til redigeringselement-nukleotidsekvensen.

5. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af kravene 1-3, hvor
korrektionsgenomet yderligere omfatter en promotor, der er
10 operativt koblet til redigeringselementet.

6. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af ovennævnte krav,
hvor hver af den 5'- og 3'-homologe arm-nukleotidsekvens
15 uafhængigt har en nukleotidlængde på ca. 50 til
2000 nukleotider.

7. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af ovennævnte krav,
20 hvor redigeringselementet omfatter et exon, et intron, en 5'-
utranslateret region (UTR), en 3'-utranslateret region (UTR),
en promotor, en splejsningsdonor, en splejsningsacceptor, en
sekvens, der svarer til et ikke-kodende RNA, en isolator, et
gene eller et fragment deraf, en kodende sekvens af et gen
25 eller en kombination deraf.

8. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af ovennævnte krav,
hvor mållocuset i mammaliakromosomet er et mutant mållocus,
30 der omfatter et eller flere mutantnukleotider i forhold til et
tilsvarende vildtypemammaliakromosom, eventuelt hvor:

(a) det mutante mållocus omfatter en amorf mutation, en
neomorf mutation, en antimorf mutation, en autosomal dominant
mutation, en autosomal recessiv mutation eller en kombination
35 deraf; og/eller

(b) det mutante mållocus er valgt blandt en promotor, en
enhancer, en signalsekvens, et intron, et exon, et
splejsningsdonorsted, et splejsningsacceptorsted, et internt

ribosomindgangssted, et inverteret exon, en isolator, et gen eller et fragment deraf, en kromosom-inversion og en kromosom-translokation i mammaliakromosomet.

- 5 9. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af ovennævnte krav,
hvor korrektionsgenomet omfatter en 5'-inverteret terminal
repetition (5'-ITR)-nukleotidsekvens 5' for den 5'-homologe
arm-nukleotidsekvens og en 3'-inverteret terminal repetition
10 (3'-ITR)-nukleotidsekvens 3' for den 3'-homologe arm-
nukleotidsekvens, eventuelt hvor:
(a) 5'-ITR-nukleotidsekvensen har mindst 95 % sekvensidentitet
med SEQ ID NO: 36, og 3'-ITR-nukleotidsekvensen har mindst
95 % sekvensidentitet med SEQ ID NO: 37; eller
15 (b) 5'-ITR-nukleotidsekvensen har mindst 95 % sekvensidentitet
med SEQ ID NO: 38, og 3'-ITR-nukleotidsekvensen har mindst
95 % sekvensidentitet med SEQ ID NO: 39.

10. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
20 anvendelse ifølge et hvilket som helst af ovennævnte krav,
hvor internukleotidbindingen er en phosphodiesterbinding.

11. Ex vivo-fremgangsmåde eller replikationsdefekt AAV til
anvendelse ifølge et hvilket som helst af ovennævnte krav,
25 hvor mållocuset er et "safe harbor"-sted.

Figure 1

60	SEQ ID NO: 2	HSC1	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 3	HSC2	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 4	HSC11	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 5	HSC3	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 6	HSC4	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 7	HSC6	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 8	HSC7	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 9	HSC8	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 10	HSC9	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 11	HSC5	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 12	HSC12	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 13	HSC17	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 14	HSC13	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 15	HSC14	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 16	HSC15	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 17	HSC16	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
	SEQ ID NO: 1	AAV9	Amino Acid Sequence	(1)	NTADGYLPDWLEDNLSGIRENNAWKPGAPQPKANQOQHDNARGGLVLPQYKYLGGFNGSLD
				61	120
		HSC1	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC2	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC11	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC3	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC4	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC6	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC7	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC8	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC9	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC5	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC12	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC17	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC13	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC14	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC15	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		HSC16	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ
		AAV9	Amino Acid Sequence	(61)	KGEPTNADAAALEHDKAYDQQLKAGDNPYLKYNHADAFFQERIKEDTSEGGNLGRAVETQ

Figure 1 (cont.)

HSC1	Amino Acid	Sequence	(121)	121	180
HSC2	Amino Acid	Sequence	(121)		
HSC11	Amino Acid	Sequence	(121)		
HSC3	Amino Acid	Sequence	(121)		
HSC4	Amino Acid	Sequence	(121)		
HSC6	Amino Acid	Sequence	(121)		
HSC7	Amino Acid	Sequence	(121)		
HSC8	Amino Acid	Sequence	(121)		
HSC9	Amino Acid	Sequence	(121)		
HSC5	Amino Acid	Sequence	(121)		
HSC12	Amino Acid	Sequence	(121)		
HSC17	Amino Acid	Sequence	(121)		
HSC13	Amino Acid	Sequence	(121)		
HSC14	Amino Acid	Sequence	(121)		
HSC15	Amino Acid	Sequence	(121)		
HSC16	Amino Acid	Sequence	(121)		
AAV9	Amino Acid	Sequence (121)			
HSC1	Amino Acid	Sequence	(181)	181	240
HSC2	Amino Acid	Sequence	(181)		
HSC11	Amino Acid	Sequence	(181)		
HSC3	Amino Acid	Sequence	(181)		
HSC4	Amino Acid	Sequence	(181)		
HSC6	Amino Acid	Sequence	(181)		
HSC7	Amino Acid	Sequence	(181)		
HSC8	Amino Acid	Sequence	(181)		
HSC9	Amino Acid	Sequence	(181)		
HSC5	Amino Acid	Sequence	(181)		
HSC12	Amino Acid	Sequence	(181)		
HSC17	Amino Acid	Sequence	(181)		
HSC13	Amino Acid	Sequence	(181)		
HSC14	Amino Acid	Sequence	(181)		
HSC15	Amino Acid	Sequence	(181)		
HSC16	Amino Acid	Sequence	(181)		
AAV9	Amino Acid	Sequence (181)			

Figure 1 (cont.)

HSC1	Amino Acid Sequence	(241)	241	300
HSC2	Amino Acid Sequence	(241)		
HSC11	Amino Acid Sequence	(241)		
HSC3	Amino Acid Sequence	(241)		
HSC4	Amino Acid Sequence	(241)		
HSC6	Amino Acid Sequence	(241)		
HSC7	Amino Acid Sequence	(241)		
HSC8	Amino Acid Sequence	(241)		
HSC9	Amino Acid Sequence	(241)		
HSC5	Amino Acid Sequence	(241)		
HSC12	Amino Acid Sequence	(241)		
HSC17	Amino Acid Sequence	(241)		
HSC13	Amino Acid Sequence	(241)		
HSC14	Amino Acid Sequence	(241)		
HSC15	Amino Acid Sequence	(241)		
HSC16	Amino Acid Sequence	(241)		
AAV9	Amino Acid Sequence (241)			
HSC1	Amino Acid Sequence	(301)	301	360
HSC2	Amino Acid Sequence	(301)		
HSC11	Amino Acid Sequence	(301)		
HSC3	Amino Acid Sequence	(301)		
HSC4	Amino Acid Sequence	(301)		
HSC6	Amino Acid Sequence	(301)		
HSC7	Amino Acid Sequence	(301)		
HSC8	Amino Acid Sequence	(301)		
HSC9	Amino Acid Sequence	(301)		
HSC5	Amino Acid Sequence	(301)		
HSC12	Amino Acid Sequence	(301)		
HSC17	Amino Acid Sequence	(301)		
HSC13	Amino Acid Sequence	(301)		
HSC14	Amino Acid Sequence	(301)		
HSC15	Amino Acid Sequence	(301)		
HSC16	Amino Acid Sequence	(301)		
AAV9	Amino Acid Sequence (301)			

Figure 1 (cont.)

HSC1	Amino Acid Sequence	(361)	361	420
HSC2	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC11	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC3	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC4	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC6	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC7	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC8	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC9	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC5	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC12	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC17	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC13	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC14	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC15	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
HSC16	Amino Acid Sequence	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	
AAV9	Amino Acid Sequence (361)	(361)	EGCLPEFPADVPMIFQYGYLTINDGSOAVGRSSFYCLIEYFPSSQMLRTGNNTQFSYELENV	421
HSC1	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	480
HSC2	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC11	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC3	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC4	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC6	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC7	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC8	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC9	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC5	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC12	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC17	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC13	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC14	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC15	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
HSC16	Amino Acid Sequence	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	
AAV9	Amino Acid Sequence (421)	(421)	PFHSSYAHSQSLDRIMNELIDQYLYLSKTTINGSGONQOCTLKEFSVACFSNNMAVQGRNYIP	

Figure 1 (cont.)

HSC1	Amino Acid Sequence	(481)	481	540
HSC2	Amino Acid Sequence	(481)		
HSC11	Amino Acid Sequence	(481)		
HSC3	Amino Acid Sequence	(481)		
HSC4	Amino Acid Sequence	(481)		
HSC6	Amino Acid Sequence	(481)		
HSC7	Amino Acid Sequence	(481)		
HSC8	Amino Acid Sequence	(481)		
HSC9	Amino Acid Sequence	(481)		
HSC5	Amino Acid Sequence	(481)		
HSC12	Amino Acid Sequence	(481)		
HSC17	Amino Acid Sequence	(481)		
HSC13	Amino Acid Sequence	(481)		
HSC14	Amino Acid Sequence	(481)		
HSC15	Amino Acid Sequence	(481)		
HSC16	Amino Acid Sequence	(481)		
AAV9	Amino Acid Sequence (481)			
HSC1	Amino Acid Sequence	(541)	541	600
HSC2	Amino Acid Sequence	(541)		
HSC11	Amino Acid Sequence	(541)		
HSC3	Amino Acid Sequence	(541)		
HSC4	Amino Acid Sequence	(541)		
HSC6	Amino Acid Sequence	(541)		
HSC7	Amino Acid Sequence	(541)		
HSC8	Amino Acid Sequence	(541)		
HSC9	Amino Acid Sequence	(541)		
HSC5	Amino Acid Sequence	(541)		
HSC12	Amino Acid Sequence	(541)		
HSC17	Amino Acid Sequence	(541)		
HSC13	Amino Acid Sequence	(541)		
HSC14	Amino Acid Sequence	(541)		
HSC15	Amino Acid Sequence	(541)		
HSC16	Amino Acid Sequence	(541)		
AAV9	Amino Acid Sequence (541)			

Figure 1 (cont.)

HSC1	Amino Acid Sequence	(601)	601	660
HSC2	Amino Acid Sequence	(601)		
HSC11	Amino Acid Sequence	(601)		
HSC3	Amino Acid Sequence	(601)		
HSC4	Amino Acid Sequence	(601)		
HSC6	Amino Acid Sequence	(601)		
HSC7	Amino Acid Sequence	(601)		
HSC8	Amino Acid Sequence	(601)		
HSC9	Amino Acid Sequence	(601)		
HSC5	Amino Acid Sequence	(601)		
HSC12	Amino Acid Sequence	(601)		
HSC17	Amino Acid Sequence	(601)		
HSC13	Amino Acid Sequence	(601)		
HSC14	Amino Acid Sequence	(601)		
HSC15	Amino Acid Sequence	(601)		
HSC16	Amino Acid Sequence	(601)		
AAV9	Amino Acid Sequence (601)			
HSC1	Amino Acid Sequence	(661)	661	720
HSC2	Amino Acid Sequence	(661)		
HSC11	Amino Acid Sequence	(661)		
HSC3	Amino Acid Sequence	(661)		
HSC4	Amino Acid Sequence	(661)		
HSC6	Amino Acid Sequence	(661)		
HSC7	Amino Acid Sequence	(661)		
HSC8	Amino Acid Sequence	(661)		
HSC9	Amino Acid Sequence	(661)		
HSC5	Amino Acid Sequence	(661)		
HSC12	Amino Acid Sequence	(661)		
HSC17	Amino Acid Sequence	(661)		
HSC13	Amino Acid Sequence	(661)		
HSC14	Amino Acid Sequence	(661)		
HSC15	Amino Acid Sequence	(661)		
HSC16	Amino Acid Sequence	(661)		
AAV9	Amino Acid Sequence (661)			

Figure 1 (cont.)

HSC1	Amino Acid Sequence	(721)	721	736
HSC2	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC11	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC3	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC4	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC6	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC7	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC8	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC9	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC5	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC12	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC17	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC13	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC14	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC15	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
HSC16	Amino Acid Sequence	(721)	YSEPRFICTRYLTNNL	
AAV9	Amino Acid Sequence (721)		YSEPRFICTRYLTNNL	

Figure 2

Capsid	Base Change	AAV 9 hu. Isolate Amino Acid	Clone New Amino Acid	Difference in Amino Acids	Placement of Amino Acid (HVR = Hypervariable Region)
HSC1 SEQ ID NO: 2	G to A	GCT - Alanine	ACT - Threonine	Alanine is nonpolar & neutral; Threonine is polar & neutral	4 bases into capsid; 2nd amino acid; Not in HVR, VP1
HSC1 SEQ ID NO: 2	G to A	CGA - Arginine	CAA - Glutamine	Arginine is polar & strongly basic; Glutamine is polar & neutral	935 bases into the capsid; 312th amino acid; Not in HVR, VP3
HSC2 SEQ ID NO: 3	A to G	GGA - Glycine	GGG - Glycine	No Amino Acid Difference	1515 bases into the capsid; 505th amino acid; 1st amino acid after HVR 7, VP3
HSC2 SEQ ID NO: 3	A to G	GAC- Aspartic Acid	GGC - Glycine	Aspartic Acid is polar & acidic; Glycine is nonpolar & neutral	1877 bases into the capsid; 626th amino acid; Not in HVR, VP3
HSC2 SEQ ID NO: 3	A to G	GAA - Glutamic Acid	GGA - Glycine	Glutamic Acid is polar & acidic; Glycine is nonpolar & neutral	2153 bases into the capsid; 718th amino acid; In HVR 12 last amino acid, VP3
HSC3 SEQ ID NO: 5	G to A	GGC - Glycine	GAC - Aspartic Acid	Glycine is nonpolar & neutral; Aspartic Acid is polar and acidic	479 bases into the capsid; 160th amino acid; Not in HVR, VP1
HSC3 SEQ ID NO: 5	A to G	TCA - Serine	TCG - Serine	No Amino Acid Difference	1044 bases into the capsid, 348th amino acid; Not in HVR VP3
HSC4 SEQ ID NO: 6	T to C	TTC - Phenylalanine	CTC - Leucine	Phenylalanine is nonpolar & neutral; Leucine is nonpolar & neutral	355 bases into the capsid, 119th amino acid; Not in HVR, VP1
HSC4 SEQ ID NO: 6	C to T	CCC - Proline	TCC - Serine	Proline is nonpolar & neutral; Serine is polar & neutral	1402 bases into the capsid; 468th amino acid; In HVR 5, VP3
HSC4 SEQ ID NO: 6	G to A	CCG - Proline	CCA - Proline	No Amino Acid Difference	1809 bases into the capsid; 603rd amino acid; Not in HVR, VP3

Figure 2 (cont.)

Capsid	Base Change	AAV 9 hu. Isolate Amino Acid	Clone New Amino Acid	Difference in Amino Acids	Placement of Amino Acid (HVR = Hypervariable Region)
HSC6 SEQ ID NO: 8	A to G	CAG - Glutamine	CGG - Arginine	Glutamine is polar & neutral; Arginine is polar & strongly basic	1769 bases into the capsid; 590th amino acid; In HVR 10; VP3
HSC5 SEQ ID NO: 11	A to G	AAG - Lysine	AGG - Arginine	Lysine is polar & basic; Arginine is polar & strongly basic	230 bases into the capsid; 77th amino acid; Not in HVR, VP1
HSC5 SEQ ID NO: 11	C to T	CCC - Proline	CCT - Proline	No Amino Acid Difference	1404 bases into the capsid; 468th amino acid; In HVR 5, VP3
HSC5 SEQ ID NO: 11	G to A	GAA - Glutamic Acid	AAA - Lysine	Glutamic Acid is polar & acidic; Lysine is polar & basic	2068 bases into the capsid; 690th amino acid; Not in HVR, VP3
HSC5 SEQ ID NO: 11	T to C	AAT - Asparagine	AAC - Asparagine	No Amino Acid Difference	2148 bases into the capsid; 716th amino acid; In HVR 13, VP3
HSC11 SEQ ID NO: 5	G to A	GTC - Valine	ATC - Isoleucine	Isoleucine is nonpolar & neutral; valine is nonpolar & neutral	193 bases into the capsid; 65th amino acid; Not in HVR; VP1
HSC11 SEQ ID NO: 4	G to T	GAC- Aspartic Acid	TAC - Tyrosine	Aspartic Acid is polar & acidic; Tyrosine is polar & neutral	1876 bases into the capsid; 626th amino acid; Not in HVR; VP3
HSC7 SEQ ID NO: 8	C to T	GCA - Alanine	GTA - Valine	Alanine is nonpolar & neutral; Valine is nonpolar & neutral	203 bases into the capsid; 68th amino acid; Not in HVR; VP1
HSC7 SEQ ID NO: 8	C to T	TTC - Phenylalanine	TTT - Phenylalanine	No Amino Acid Difference	924 bases into the capsid; 308th amino acid; Not in HVR; VP3
HSC8 SEQ ID NO: 9	A to G	CAG - Glutamine	CGG - Arginine	Glutamine is polar & neutral; Arginine is polar & strongly basic	452 bases into the capsid; 151th amino acid; In HVR 1; VP1

Figure 2 (cont.)

Capsid	Base Change	AAV 9 hu. Isolate Amino Acid	Clone New Amino Acid	Difference in Amino Acids	Placement of Amino Acid (HVR = Hypervariable Region)
HSC8 SEQ ID NO: 9	G to A	GAG - Glutamine Acid	GAA - Glutamine Acid	No Amino Acid Difference	2058 bases into the capsid; 686th amino acid; Not in HVR; VP3
HSC9 SEQ ID NO: 10	T to C	TTT - Phenylalanine	TTC - Phenylalanine	No Amino Acid Difference	357 bases into the capsid; 119th amino acid; Not in HVR; VP1
HSC9 SEQ ID NO: 10	T to G	TGT - Cysteine	GGT - Glycine	Cysteine is polar & neutral; Glycine is nonpolar & neutral	616 bases into the capsid; 206th amino acid; Not in HVR; Very beginning of VP3
HSC12 SEQ ID NO: 12	A to T	GCA - Alanine	GCT - Alanine	No Amino Acid Difference	579 bases into capsid; 193rd amino acid into capsid, in the end of VP2 near HVR 2
HSC12 SEQ ID NO: 12	G to A	CGT - Arginine	CAT - Histidine	Arginine is polar and strongly basic; Histidine is polar and weakly basic	887 bases into capsid; 296th amino acid into capsid; in VP3, not in HVR
HSC12 SEQ ID NO: 12	G to A	AGT - Serine	AAT - Asparagine	Serine is polar and neutral; Asparagine is polar and neutral	1391 bases into capsid; 464th amino acid into capsid; in VP3 in HVR 5
HSC12 SEQ ID NO: 12	T to C	GCT - Alanine	GCC - Alanine	No Amino Acid Difference	1506 bases into capsid; 502nd amino acid into capsid; in VP3 in HVR 7
HSC12 SEQ ID NO: 12	G to A	GGA - Glycine	AGA - Arginine	Glycine is nonpolar and neutral; Arginine is polar and strongly basic	1513 bases into capsid; 505th amino acid into capsid; in VP3, right after HVR 7
HSC12 SEQ ID NO: 12	G to A	GTG - Valine	ATG - Methionine	Valine is nonpolar and neutral; Methionine is nonpolar and neutral	2041 bases into capsid; 681st amino acid into capsid; in VP3; not in HVR
HSC13 SEQ ID NO: 14	G to A	GGA - Glycine	AGA - Arginine	Glycine is nonpolar and neutral; Arginine is polar and strongly basic	1513 bases into capsid; 505th amino acid into capsid; in VP3, right after HVR 7

Figure 2 (cont.)

Capsid	Base Change	AAV 9 hu. Isolate Amino Acid	Clone New Amino Acid	Difference in Amino Acids	Placement of Amino Acid (HVR = Hypervariable Region)
HSC14 SEQ ID NO: 15	G to A	GGA - Glycine	AGA - Arginine	Glycine is nonpolar and neutral; Arginine is polar and strongly basic	1513 bases into capsid; 505th amino acid into capsid; in VP3, right after HVR 7
HSC14 SEQ ID NO: 15	T to G	CTG - Leucine	CGG - Arginine	Leucine is nonpolar and neutral; Arginine is polar and strongly basic	2060 bases into capsid; 687th amino acid into capsid; in VP3; Not in HVR.
HSC15 SEQ ID NO: 16	A to G	ACG - Threonine	GCG - Alanine	Threonine is polar and neutral; Arginine is polar and strongly basic	1036 bases into capsid; 346th amino acid into the capsid; In VP3; Not in HVR
HSC15 SEQ ID NO: 16	G to A	GGA - Glycine	AGA - Arginine	Glycine is nonpolar and neutral; Arginine is polar and strongly basic	1513 bases into capsid; 505th amino acid into capsid; in VP3, right after HVR 7
HSC16 SEQ ID NO: 17	T to A	TTT - Phenylalanine	ATT - Isoleucine	Phenylalanine is nonpolar and neutral; Isoleucine is nonpolar and neutral	1501 bases into capsid; 501st amino acid into capsid; In VP3, In HVR7
HSC16 SEQ ID NO: 17	G to A	GGA - Glycine	AGA - Arginine	Glycine is nonpolar and neutral; Arginine is polar and strongly basic	1513 bases into capsid; 505th amino acid into capsid; in VP3, right after HVR 7
HSC16 SEQ ID NO: 17	G to A	CCG - Proline	CCA - Proline	No Amino Acid Difference	1713 bases into capsid; 571st amino acid into capsid; In VP3 Not in HVR
HSC16 SEQ ID NO: 17	A to G	TAC - Tyrosine	TGC - Cysteine	Tyrosine is polar and neutral; Cysteine is polar and neutral	2117 bases into capsid; 706th amino acid into capsid; In VP3; In HVR 12 and is next to a YF mutation
HSC17 SEQ ID NO: 13	C to T	GAC - Aspartic Acid	GAT - Aspartic Acid	No Amino Acid Difference	180 bases into the capsid; 60th amino acid into capsid; In VP1; Not in HVR
HSC17 SEQ ID NO: 13	G to A	GGA - Glycine	AGA - Arginine	Glycine is nonpolar and neutral; Arginine is polar and strongly basic	1513 bases into capsid; 505th amino acid into capsid; in VP3, right after HVR 7

Figure 3




Figure 4A






Figure 4B

Figure 5


A.



B.



C.



D.




Figure 5 (cont.)

E.





Figure 6


A.

untransduced




B.

50,000 MOI



C.

100,000 MOI



D.

200,000 MOI




Figure 6 (cont.)

E.





Figure 7

A.



B.








Figure 8**A.****B.**

Figure 9

A.



B.



C.






Figure 10

A.



B.



C.






Figure 11

A.



B.



C.




Figure 12

A.

% FP Positive Cells				
	4 day	18 day	20 day	39 day
Untransduced	0	0	0	0
AAVHSC7 FP Donor	3	7.23	4.07	4.61
AAVHSC17 FP Donor	20.2	5.43	2.85	9.18

B.




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20





Figure 21

A.



B.







Figure 22


A.



B.



C.



D.






Figure 22 (cont.)


E.




F.




G.



H.



Stem cell-derived AAV map to Clade F



• Nucleotide homology
Figure 23




Figure 24

Schema: Targeted Integration of Promoterless Reporter into
PPP1r12C INTRON (AAVS1)




Figure 25

Figure 26A




Figure 26B







Figure 26C





- AAVF mediates efficient gene targeting in primary CD34+ cells
- High efficiency targeting is observed in human leukemic & liver cell lines
- AAVF editing does not require DNA synthesis

Figure 26D

**Figure 26E**




Figure 26F

AAVF VECTORS TARGET INTEGRATION OF LARGE ORFS INTO THE AAVS1 LOCUS IN CD34+ CELLS





Figure 27

HA-L: 5'-Homology Arm
 HA-R: 3'-Homology Arm
 NTC: No Template Control
 Untd: Untransduced Cells

AAVF VECTORS FOR NUCLEOTIDE SUBSTITUTION

Figure 28A



- Mutates TA to AT
- Changes an NheI site to a SphI site

NheI $\longrightarrow$ SphI
 GCTAGC $\longrightarrow$ GCATGC

Figure 28B




Figure 28C




Figure 28D





Figure 28E



AAVF MEDIATES PRECISE NUCLEOTIDE SUBSTITUTION





Figure 29

	GFP+ BrdU- (Non-dividing Cells)	GFP+BrdU+ (Dividing Cells)
AAVHSC7	12	14.6
AAVHSC17	6.15	6.9
AAV9	4.31	3.73
AAV8	1.37	0.839

Figure 30

1e6 Human Cord blood CD34+ Stem Cells



Analysis of Human Blood Lineages in Marrow & Spleen

- Flow analysis
- Flow sort
- Venus, CD34+, Glyco A+ (erythroid)

Figure 31A




Figure 31B




Figure 31C

	EXP	2A	2B	2C	2D	2E	2F	2G	2H	2I
		ASPT	ASPT	ASPT	ASPT	ASPT	ASPT	ASPT	ASPT	ASPT
		CO34+VEN+*	CO34+VEN+*	CO34+VEN+*	CO34+VEN+*	CO34+VEN+*	CO34+VEN+*	CO34+VEN+*	CO34+VEN+*	CO34+VEN+*
WKS POST TX TO VENUS		1	1	1	1	1	1	1	1	1
WKS POST VENUS TO HAZ/EST		5	5	5	5	5	5	5	5	5
WKS POST TX TO HAZ/EST		6	6	7	7	7	7	7	7	7
VEN		13.4	14.7	13.4	14.7	14.7	14.7	14.7	14.7	14.7
HEN		13.2	9.49	19.48	20.9	22.2	22.2	22.2	22.2	22.2
SPN		15.9	14.8	15.9	17.4	17.4	17.4	17.4	17.4	17.4

Figure 32A

% Venus Positive	Mid-Term	Mid-Term	Mid-Term	Mid-Term	Long-Term	Long-Term
	AAVF7	AAVF7	AAVF17	AAVF17	AAVF17	AAVF17
<i>Compartment</i>	CD34+	CD45+	CD34+	CD45+	CD34+	GlycoA+
Vertebral Marrow	54.9	32.4	49.9	14.7	ND*	ND*
Femoral Marrow	20.3	9.49	26.8	11.2	75.5	99.4
Spleen	71.1	15.9	82	24.8	58.1	99.2

* ND: Not Done

Figure 32B

SEQUENCE ANALYSIS SHOWING CHROMOSOMAL INSERTION OF VENUS INTO INTRON 1 OF PPP1R12C BY AAVF VECTORS

ALIGNMENT OF AAV-TARGETED SEQUENCES IN AAVS1

```

> K1462 AAVBSC7 #1      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> K1462 AAVBSC7 #2      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> K1462 AAVBSC7 #3      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> K1462 AAVBSC7 #4      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> K1462 AAVBSC7 #5      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC7 #1      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC7 #2      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC7 #3      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC7 #4      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC15 #1     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC15 #2     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC17 #1     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC17 #2     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC17 #3     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> PBSC AAVBSC17 #4     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> BspC2 AAVBSC7 #1     (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
> AC010327 AAVS1      (2)  GGGCTGAGCAATTGTCACTTTGCGCTGCGCTGCTCTGCGCGCGGAGTGGCTTGGCTGTGCGCGCG
>   > Venus Donor      (1)
> Homology Arm 5'      (1)

> K1462 AAVBSC7 #1     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> K1462 AAVBSC7 #2     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> K1462 AAVBSC7 #3     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> K1462 AAVBSC7 #4     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> K1462 AAVBSC7 #5     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC7 #1     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC7 #2     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC7 #3     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC7 #4     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC15 #1     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC15 #2     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC17 #1     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC17 #2     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC17 #3     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> PBSC AAVBSC17 #4     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> BspC2 AAVBSC7 #1     (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
> AC010327 AAVS1      (67)  GAAGTCGAGCCCTCTAAGCGCTGCGCTGCTCTGCTGAGTCCGGAGCACTTTGAGGCTCTACTGGCTT
>   > Venus Donor      (1)
> Homology Arm 5'      (1)

> K1462 AAVBSC7 #1     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> K1462 AAVBSC7 #2     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> K1462 AAVBSC7 #3     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> K1462 AAVBSC7 #4     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> K1462 AAVBSC7 #5     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC7 #1     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC7 #2     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC7 #3     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC7 #4     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC15 #1     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC15 #2     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC17 #1     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC17 #2     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC17 #3     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> PBSC AAVBSC17 #4     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> BspC2 AAVBSC7 #1     (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
> AC010327 AAVS1      (132) CTGCGCGCGCTCTGAGCCCACTGTTTCCCGTTCCAGGCGAGGTCCGCTTTCTCTGACCTGCATTTC
>   > Venus Donor      (1)
> Homology Arm 5'      (1)

```

Figure 33

[illegible]

Figure 33 (cont.)

[illegible]

Figure 33 (cont.)

[illegible]

Figure 33 (cont.)

```

» PBSC AAVHSC7 #1      (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC7 #2      (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC7 #3      (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC7 #4      (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC15 #1     (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC15 #2     (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC17 #1     (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC17 #2     (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC17 #3     (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» PBSC AAVHSC17 #4     (762) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» RepG2 AAVHSC7 #1     (608) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA
» AC010327 AAVS1
  » RepG2 Prom.
» Homology Arm 5'      (608) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCTCCAAACCCGGGCCCCCTATGTCCA

» PBSC AAVHSC7 #1      (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC7 #2      (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC7 #3      (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC7 #4      (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC15 #1     (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC15 #2     (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC17 #1     (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC17 #2     (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC17 #3     (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» PBSC AAVHSC17 #4     (847) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» RepG2 AAVHSC7 #1     (673) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT
» AC010327 AAVS1
  » RepG2 Prom.
» Homology Arm 5'      (673) CTTCAGGACAGCATGTTTGCTGCGCTCCAGGATCCCTGTGTGTCGCCAGCTGGGACCAACCTTATATT

» PBSC AAVHSC7 #1      (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC7 #2      (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC7 #3      (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC7 #4      (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC15 #1     (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC15 #2     (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC17 #1     (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC17 #2     (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC17 #3     (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» PBSC AAVHSC17 #4     (912) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGCCCTCCACCCCACAGTGG
» RepG2 AAVHSC7 #1     (738) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGTGCCCTCCACCCCACAGTGG
» AC010327 AAVS1
  » RepG2 Prom.
» Homology Arm 5'      (738) CCCAGGCCCCGGTTAATGTGGCTCTGGTTCCTGGTACTTTTATCTGTGTGCCCTCCACCCCACAGTGG

```

Figure 33 (cont.)

[illegible][illegible][illegible][illegible][illegible][illegible]

Figure 33 (cont.)

[illegible]

Figure 33 (cont.)

> E112	AAVH5C7 #1	(1367)	CTAAGCTCTACGCTCCAGTAAAGGCAACATCTTCTTTAAAGGATGACCTGATACACAGAGATGCGCGTGG
> E113	AAVH5C7 #2	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E114	AAVH5C7 #3	(1367)	CTAAGCTCTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E115	AAVH5C7 #4	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E150	AAVH5C7 #1	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E151	AAVH5C7 #2	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E152	AAVH5C7 #3	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E153	AAVH5C7 #4	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E154	AAVH5C15 #1	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E155	AAVH5C15 #2	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E156	AAVH5C17 #1	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E157	AAVH5C17 #2	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E158	AAVH5C17 #3	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E159	AAVH5C17 #4	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E160	AAVH5C17 #1	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E161	AAVH5C17 #2	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E162	AAVH5C17 #3	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E163	AAVH5C17 #4	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> Rep12	AAVH5C7 #1	(1367)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> AC010327	AAV51	(+174)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> > Venus-5'		(+174)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> Homology Arm 5'		(+801)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> > Venus-pA		(271)	CTAAGGCTTACGCTCCAGTAAAGGCAACATCTTCTTCAAGGCAATGACCGGATATACATATAGCGGCTGGAG
> E112	AAVH5C7 #1	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E113	AAVH5C7 #2	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E114	AAVH5C7 #3	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E115	AAVH5C7 #4	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E150	AAVH5C7 #1	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E151	AAVH5C7 #2	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E152	AAVH5C7 #3	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E153	AAVH5C7 #4	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E154	AAVH5C15 #1	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E155	AAVH5C15 #2	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E156	AAVH5C17 #1	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E157	AAVH5C17 #2	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E158	AAVH5C17 #3	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E159	AAVH5C17 #4	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E160	AAVH5C17 #1	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E161	AAVH5C17 #2	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E162	AAVH5C17 #3	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E163	AAVH5C17 #4	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> Rep01	AAVH5C7 #1	(1367)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> AC010327	AAV51	(+174)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> > Venus-5'		(+174)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> Homology Arm 5'		(+801)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> > Venus-pA		(336)	AGGTTGAAGTTTCGAGGCGGACACACCTGCTTAAAGCGATCTGATGCTAAGGCTCATCGACTTCAGGAG
> E112	AAVH5C7 #1	(1367)	GACGGCAACATCTCTGCGGACACAAAGCTGCAATACAACTACAAAGCGGCTATATATACCGG
> E113	AAVH5C7 #2	(1367)	GACGGCAACATCTCTGCGGACACAAAGCTGCAATACAACTACAAAGCGGCTATATATACCGG
> E114	AAVH5C7 #3	(1367)	GACGGCAACATCTCTGCGGACACAAAGCTGCAATACAACTACAAAGCGGCTATATATACCGG
&			

Figure 33 (cont.)

[illegible]

Figure 33 (cont.)

Sequence Analysis Showing Chromosomal Insertion of a Short Sequence Into Intron 1 of PPP1R12C by AAVF Vectors

	1	65
» AAVESC7 RFLP PBSC #1	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» AAVESC7 RFLP PBSC #2	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» AAVESC7 RFLP HepG2 #1	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» AAVESC7 RFLP HepG2 #2	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» AAVESC7 RFLP HepG2 #3	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» AAVESC7 RFLP HepG2 #4	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» RFLP Vector Sequence	(1)	
» AC010327 AAVS1	(2) GGGCCCTGGCCATTGTGACCTTGGGCTGGCCCTCCTCTGGCCCGGAGTGGCCCTTGGTGTGGCGCG	
» Homology Arm 5'	(1)	
» Homology Arm 3'	(1)	
	66	130
» AAVESC7 RFLP PBSC #1	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» AAVESC7 RFLP PBSC #2	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» AAVESC7 RFLP HepG2 #1	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» AAVESC7 RFLP HepG2 #2	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» AAVESC7 RFLP HepG2 #3	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» AAVESC7 RFLP HepG2 #4	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» RFLP Vector Sequence	(1)	
» AC010327 AAVS1	(67) GAACCTGTGGCCCTCTAAAGCTGGCGTCTCTCTCTGAGTCCGGACCACTTTGAGCTCTACTGGCTT	
» Homology Arm 5'	(1)	
» Homology Arm 3'	(1)	
	131	195
» AAVESC7 RFLP PBSC #1	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCTGCAATC	
» AAVESC7 RFLP PBSC #2	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCTGCAATC	
» AAVESC7 RFLP HepG2 #1	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCTGCAATC	
» AAVESC7 RFLP HepG2 #2	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCTGCAATC	
» AAVESC7 RFLP HepG2 #3	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCTGCAATC	
» AAVESC7 RFLP HepG2 #4	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCTGCAATC	
» RFLP Vector Sequence	(1) TGCTTTCTCTGACCAAGCAATC	
» AC010327 AAVS1	(132) CTGGCGCGCCCTCTGGCCCACTGTTTCCGCCCTTCCAGGCAAGGTCTGCTTTCTCTGACCAAGCAATC	
» Homology Arm 5'	(1) TGCTTTCTCTGACCAAGCAATC	
» Homology Arm 3'	(1)	
	196	260
» AAVESC7 RFLP PBSC #1	(197) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» AAVESC7 RFLP PBSC #2	(197) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» AAVESC7 RFLP HepG2 #1	(197) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» AAVESC7 RFLP HepG2 #2	(197) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» AAVESC7 RFLP HepG2 #3	(197) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» AAVESC7 RFLP HepG2 #4	(197) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» RFLP Vector Sequence	(23) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(23) TCTCCGCTGGGCTGTGCGCGCTTCTCTCTGAGCTTGTGGCTGGGTCACCTCTAGGGCTGGGC	
» Homology Arm 3'	(1)	

Figure 34

	261	325
» AAVHSC7 RFLP PBSC #1	(262) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» AAVHSC7 RFLP PBSC #2	(262) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» AAVHSC7 RFLP HepG2 #1	(262) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» AAVHSC7 RFLP HepG2 #2	(262) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» AAVHSC7 RFLP HepG2 #3	(262) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» AAVHSC7 RFLP HepG2 #4	(262) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» RFLP Vector Sequence	(88) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(88) CAGATCCCTTCCTTCGCGCGCTCCTTCAGGTTCCGTCCTTCCTCCACTCCCTCTTCGCCCTTGGCTCCT	
» Homology Arm 3'	(1)	
	326	390
» AAVHSC7 RFLP PBSC #1	(327) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» AAVHSC7 RFLP PBSC #2	(327) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» AAVHSC7 RFLP HepG2 #1	(327) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» AAVHSC7 RFLP HepG2 #2	(327) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» AAVHSC7 RFLP HepG2 #3	(327) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» AAVHSC7 RFLP HepG2 #4	(327) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» RFLP Vector Sequence	(153) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(153) GCTGTGTTCCTGCCCCAAGGATGGTCTTTCCGGAGCACTTCCTTCCTCGGCGCTGCACCAAGCTGATG	
» Homology Arm 3'	(1)	
	391	455
» AAVHSC7 RFLP PBSC #1	(392) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» AAVHSC7 RFLP PBSC #2	(392) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» AAVHSC7 RFLP HepG2 #1	(392) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» AAVHSC7 RFLP HepG2 #2	(392) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» AAVHSC7 RFLP HepG2 #3	(392) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» AAVHSC7 RFLP HepG2 #4	(392) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» RFLP Vector Sequence	(218) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(218) TCCTCTGAGCGGATCCTTCCCGTGTCTGAGTCTCTCCGGGATCTCTCTCTCCCTCACCACCAACCC	
» Homology Arm 3'	(1)	
	456	520
» AAVHSC7 RFLP PBSC #1	(457) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» AAVHSC7 RFLP PBSC #2	(457) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» AAVHSC7 RFLP HepG2 #1	(457) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» AAVHSC7 RFLP HepG2 #2	(457) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» AAVHSC7 RFLP HepG2 #3	(457) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» AAVHSC7 RFLP HepG2 #4	(457) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» RFLP Vector Sequence	(283) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(283) CATGCCCTCTTCACTCCCTGGGTTCCCTTTTCCCTTCCTCTCTGGGGCTCTGCGCATCTCTCGTT	
» Homology Arm 3'	(1)	

Figure 34 (cont.)

	521	585
» AAVHSC7 RFLP PBSC #1	(522) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» AAVHSC7 RFLP PBSC #2	(522) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» AAVHSC7 RFLP HepG2 #1	(522) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» AAVHSC7 RFLP HepG2 #2	(522) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» AAVHSC7 RFLP HepG2 #3	(522) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» AAVHSC7 RFLP HepG2 #4	(522) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» RFLP Vector Sequence	(348) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(348) TCTTAGGATGGGCTTCTCCGACGGATGCTCTCCCTTGCGTCCCGGCTCCGCTTCTTGTAGGCTTGC	
» Homology Arm 3'	(1)	
	586	650
» AAVHSC7 RFLP PBSC #1	(587) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» AAVHSC7 RFLP PBSC #2	(587) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» AAVHSC7 RFLP HepG2 #1	(587) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» AAVHSC7 RFLP HepG2 #2	(587) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» AAVHSC7 RFLP HepG2 #3	(587) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» AAVHSC7 RFLP HepG2 #4	(587) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» RFLP Vector Sequence	(413) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(413) ATCATCACCCTTTTCTGGACAAACCCAAAGTACCCCGTCTCCCTGGCTTTAGGCACTCTCCAT	
» Homology Arm 3'	(1)	
	651	715
» AAVHSC7 RFLP PBSC #1	(652) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» AAVHSC7 RFLP PBSC #2	(652) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» AAVHSC7 RFLP HepG2 #1	(652) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» AAVHSC7 RFLP HepG2 #2	(652) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» AAVHSC7 RFLP HepG2 #3	(652) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» AAVHSC7 RFLP HepG2 #4	(652) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» RFLP Vector Sequence	(478) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(478) CCTCTTGCCTTTCTTTTGGCTGGACACCCCGTTCTCTCTGTGGATTCGGCTCAGCTCTCACTCCCTTC	
» Homology Arm 3'	(1)	
	716	780
» AAVHSC7 RFLP PBSC #1	(717) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» AAVHSC7 RFLP PBSC #2	(717) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» AAVHSC7 RFLP HepG2 #1	(717) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» AAVHSC7 RFLP HepG2 #2	(717) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» AAVHSC7 RFLP HepG2 #3	(717) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» AAVHSC7 RFLP HepG2 #4	(717) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» RFLP Vector Sequence	(543) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(543) ATTTGGGCAGCTCCCTACCCCGCTTACCTCTCTAGTCTGTGCTAGCTCTTCCAGCCCGCTGCA	
» Homology Arm 3'	(1)	

Figure 34 (cont.)

	781	845
» AAVHSC7 RFLP PBSC #1	(782) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» AAVHSC7 RFLP PBSC #2	(782) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» AAVHSC7 RFLP HepG2 #1	(782) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» AAVHSC7 RFLP HepG2 #2	(782) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» AAVHSC7 RFLP HepG2 #3	(782) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» AAVHSC7 RFLP HepG2 #4	(782) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» RFLP Vector Sequence	(608) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(608) TGGCATCTTCCAGGGGTCCGAGAGCTCAGCTAGTCTTCTTCTCCAAACCCGGGCCCCATGTCCA	
» Homology Arm 3'	(1)	
	846	910
» AAVHSC7 RFLP PBSC #1	(847) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» AAVHSC7 RFLP PBSC #2	(847) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» AAVHSC7 RFLP HepG2 #1	(847) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» AAVHSC7 RFLP HepG2 #2	(847) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» AAVHSC7 RFLP HepG2 #3	(847) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» AAVHSC7 RFLP HepG2 #4	(847) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» RFLP Vector Sequence	(673) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(673) CTTCAGGACAGCATGTTTGCTGCTCCAGGGATCCTGTGTCCCGAGCTGGGACACCTTATATT	
» Homology Arm 3'	(1)	
	911	975
» AAVHSC7 RFLP PBSC #1	(912) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» AAVHSC7 RFLP PBSC #2	(912) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» AAVHSC7 RFLP HepG2 #1	(912) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» AAVHSC7 RFLP HepG2 #2	(912) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» AAVHSC7 RFLP HepG2 #3	(912) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» AAVHSC7 RFLP HepG2 #4	(912) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» RFLP Vector Sequence	(738) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(738) CCCAGGGCCGGTTAAAGTGGCTCTGGTTCTGGGTACCTTTATCTGTCCCTCCACCCACAGTGG	
» Homology Arm 3'	(1)	
	976	1040
» AAVHSC7 RFLP PBSC #1	(977) GGCAGGCTTTACAGGACAGGATGG	
» AAVHSC7 RFLP PBSC #2	(977) GGCAGGCTTTACAGGACAGGATGG	
» AAVHSC7 RFLP HepG2 #1	(977) GGCAGGCTTTACAGGACAGGATGG	
» AAVHSC7 RFLP HepG2 #2	(977) GGCAGGCTTTACAGGACAGGATGG	
» AAVHSC7 RFLP HepG2 #3	(977) GGCAGGCTTTACAGGACAGGATGG	
» AAVHSC7 RFLP HepG2 #4	(977) GGCAGGCTTTACAGGACAGGATGG	
» RFLP Vector Sequence	(803) GGCAGGCTTTACAGGACAGGATGGTGACAGAAAAGCCCCATCCTTAGGCTCTCTCTCTCTA	
» AC010327 AAVS1	(•174)	
» Homology Arm 5'	(•801)	
» Homology Arm 3'	(1) TAGGACAGGATGGTGACAGAAAAGCCCCATCCTTAGGCTCTCTCTCTCTA	
>> Hind III	AGGCTT	

Figure 34 (cont.)

AAVF TARGETS INTEGRATION OF SMALL INSERTS INTO
AAVS1 - CD34+ CELLS





Figure 35

Figure 36






Figure 37




Figure 38




Figure 39




Figure 40A




Figure 40B




Figure 40C




Figure 40D