

(51) International Patent Classification:
C12Q 1/42 (2006.01)(21) International Application Number:
PCT/US2015/061017(22) International Filing Date:
17 November 2015 (17.11.2015)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/101,751 9 January 2015 (09.01.2015) US

(72) Inventor; and

(71) Applicant : MALO, Madhu, S. [US/US]; c/o Atoxin Biotech LLC, 100 Barber Avenue, Worcester, Massachusetts 01606 (US).

(74) Agents: SUSAN, Janine, M. et al.; Burns & Levinson, LLP, 125 Summer Street, Boston, Massachusetts 02110 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: DIAGNOSIS AND TREATMENT OF INCIPIENT DIABETES

FIG. 1



(57) Abstract: A method is described for predicting incipient diabetes, metabolic disorders or the metabolic syndrome by developing a personal temporal phosphatase profile, which is generated by measuring phosphatase concentration in stool at a single time-point or multiple time-points. The phosphatase profile further can be used for diagnosing and determining prognosis of other incipient or overt diseases, such as the metabolic syndrome, coronary heart disease, nonalcoholic fatty liver disease, cancers, other chronic or acute diseases or infectious diseases. Also described is a specific dose of phosphatase for therapeutic use in incipient diabetes and other incipient or overt diseases.

DIAGNOSIS AND TREATMENT OF INCIPIENT DIABETES

BACKGROUND

Type 2 diabetes mellitus (T2DM) is characterized by hyperglycemia, and it is a major
5 global health problem that affects nearly 5.3% population of the world with devastating
consequences in the context of healthcare cost, morbidity and mortality (American Diabetes
Association. Diagnosis and classification of diabetes mellitus. Diabetes Care. 2014 Jan;37
Suppl 1:S81-90; Abdullah et al. Int J Endocrinol. 2014;59:3982; International Diabetes
Federation, IDF Diabetes Atlas, 2012, <http://www.idf.org/diabetesatlas>.

10 <http://www.idf.org/diabetesatlas/update-2014>; <http://www.idf.org/diabetesatlas>;
<http://www.idf.org/diabetesatlas/update-2014>; American Diabetes Association. Economic
costs of diabetes in the U.S. in 2012. Diabetes Care 2013;36:1033--1046).

T2DM is diagnosed by measuring plasma glucose levels, and a fasting plasma glucose
(FPG) level > 7.0 millimole/liter (126 milligram/deciliter) is considered as diagnostic for
15 T2DM. Hyperglycemia is associated with established T2DM or with the terminal stage of
development of T2DM. Therefore, measuring FPG cannot be used to predict far (years) in
advance whether a person have incipient diabetes, which means that the person would
develop diabetes in the future although the current FPG value is normal. Therefore, a
diagnostic protocol predicting incipient diabetes and/or other diseases is warranted. Also,
20 therapeutic intervention of these incipient or overt diseases as well as determining the
prognosis of a disease is critically important.

SUMMARY OF THE INVENTION

According to the disclosure, patients with type 2 diabetes mellitus (T2DM) have less
25 amount of the brush-border enzyme intestinal alkaline phosphatase (IAP) in their stools
compared to the control non-diabetic healthy subjects. Disclosed herein is a protocol for
diagnosing the incipient diabetes and other metabolic disorders (e.g., cardiovascular diseases
(coronary heart disease), cerebrovascular disease (stroke), peripheral vascular disease,
nonalcoholic fatty liver disease, etc.) and the metabolic syndrome (e.g., obesity,
30 hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia,
hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein

(LDL), hypertension, fatty liver, etc.) by developing a temporal profile of stool IAP levels for humans and other animals, such as cattle, pig, sheep, goats, cows, horses, dogs, cats, monkeys, rabbits, rats, mice, chickens, turkeys, etc. The temporal profile of IAP consists of IAP concentration determined at a single time-point or multiple time-points. A low IAP level or a persistent loss of IAP from a previously high level in a healthy subject is indicative of incipient diabetes, other metabolic disorders or the metabolic syndrome. Also disclosed is the use of temporal profiling of IAP for diagnosis or prognosis of the metabolic diseases, acute or chronic diseases, or infectious diseases, including but not limited to myocardial infarction, cancer, tuberculosis, salmonellosis, AIDS, hepatitis, acute or chronic kidney injury. It is also disclosed that a personal stool IAP profile of a normal healthy subject could be used as a reference IAP profile for future use.

It is also disclosed that specific doses of IAP, as an oral supplementation, can be administered to a mammal to decrease the risk of incipient diabetes, other metabolic disorders or the metabolic syndrome. The level of IAP in stool greatly varies from one individual to another individual. An individual with less than the average IAP level in control healthy humans (approx. 65 U/gm stool using the assay conditions disclosed herein) can be immediately provided with IAP supplementation, whereas an individual with a higher level of IAP can be further investigated allowing for the development of a temporal profile of IAP levels encompassing multiple time-points. It is further disclosed that IAP can be administered to a mammal for the treatment of metabolic diseases, acute or chronic diseases, or infectious diseases including but not limited to myocardial infarction, cancer, tuberculosis, salmonellosis, AIDS, hepatitis, acute or chronic kidney injury.

In an illustrative embodiment, the method for diagnosing or for determining the prognosis of incipient diabetes, an incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease or metabolic syndrome involves measuring the concentration of phosphatase in stool of a mammal and comparing the concentration to a normal range of concentrations.

In another illustrative embodiment, the methods further involve developing a temporal profile by measuring the concentration of phosphatase at a single time-point or multiple time-points.

In another illustrative embodiment, the phosphatase is intestinal alkaline phosphatase, placental alkaline phosphatase, tissue nonspecific alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity.

In yet a further illustrative embodiment, in addition to measuring the concentration of phosphatase in a stool sample, a sample of saliva, urine, blood, plasma, serum, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, or a body cavity fluid is also analyzed.

In another illustrative embodiment, the incipient diabetes or diabetes is type 2 diabetes mellitus, gestational diabetes, diabetes insipidus, or type 1 diabetes.

In another illustrative embodiment, the incipient metabolic disease or metabolic disease is a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

In another illustrative embodiment, the incipient metabolic syndrome or metabolic syndrome is obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension or fatty liver.

In an illustrative embodiment, the method for decreasing the risk of or for treating incipient diabetes, incipient metabolic diseases, incipient metabolic syndrome, diabetes, metabolic diseases or metabolic syndrome involves administering an effective amount of phosphatase to a mammal in need thereof.

In another illustrative embodiment, the phosphatase is intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity.

In yet another illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is a native or recombinant isoform.

In yet another illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is a mutant isoform.

5 In yet a further illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is a native human intestinal alkaline phosphatase.

In another illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is a recombinant human intestinal alkaline phosphatase.

10 In another illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is a native calf or bovine intestinal alkaline phosphatase.

In yet another illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is a recombinant calf or bovine intestinal alkaline phosphatase.

In yet a further illustrative embodiment, the phosphatase is administered orally, topically, subcutaneously, intradermally, intramuscularly, intravenously, vaginally or rectally.

15 In yet a further illustrative embodiment, the phosphatase is administered by an oral, nasal, dermal, vaginal or rectal spray.

In a still further illustrative embodiment, the phosphatase is administered before, during or after a meal.

20 In another illustrative embodiment, the phosphatase is administered in an amount effective to achieve a therapeutic goal of preventive (i.e. decreasing the risk of), palliative, or curative endpoint.

In a further illustrative embodiment, the phosphatase is administered in an amount effective to maintain a steady-state level of intestinal alkaline phosphatase in stool.

25 In yet a further illustrative embodiment, the phosphatase is administered in an amount effective to maintain a steady-state level between 1 to 650 units of intestinal alkaline phosphatase per gram stool, preferably in an amount effective to maintain a steady-state level of 65 units of intestinal alkaline phosphatase per gram stool measured using the alkaline phosphatase assay described herein.

30 In another illustrative embodiment, the phosphatase is in an amount of 1 to 6,500 units of phosphatase per dose.

In another illustrative embodiment, the phosphatase is dispensed in a tablet, gel, capsule, liquid or semi-solid chewing gum.

In another illustrative embodiment, the phosphatase is dispensed in milk, food or a beverage.

5 In another illustrative embodiment, the phosphatase is dispensed with other medication from a group consisting of anti-diabetes, anti-hypertension, anti-cholesterol, anti-depression or antibiotics.

In another illustrative embodiment, the phosphatase is dispensed with other medication from a group consisting of up-regulators of IAP including, but not limited to,
10 sodium butyrate, thyroid hormone, curcumin, omega-3 fatty acid or combinations thereof.

In another illustrative embodiment, the phosphatase is dispensed with other medication from a group consisting of anti-dysbiosis factors including, but not limited to, a protein or a peptide, preferably a beta-lactamase.

In another illustrative embodiment, the phosphatase is dispensed with other
15 medication from a group consisting of probiotic bacteria.

In another illustrative embodiment, the phosphatase is an intestinal alkaline phosphatase that is highly purified, partially purified or unpurified.

In yet another illustrative embodiment, the phosphatase is administered to a subject, wherein the subject is human.

20 In yet a further illustrative embodiment, the phosphatase is administered to a subject, wherein the subject is a cow, pig, sheep, dog, cat, horse, monkey, cattle, chicken, turkey, rat, mouse, bird or a mammal.

In an illustrative embodiment, the method for diagnosing or for determining the prognosis of a chronic disease, an acute disease, an infectious disease, a disorder or a
25 syndrome involves measuring the concentration of phosphatase in stool of a mammal and comparing the concentration to a normal range of concentrations.

In a further illustrative embodiment, the method for decreasing the risk of or for treating a chronic disease, an acute disease, an infectious disease, a disorder or a syndrome involves administering an effective amount of phosphatase to a mammal in need thereof.

30 In another illustrative embodiment, the chronic disease is Acne, Acquired Immunodeficiency Syndrome (AIDS), Ankylosing spondylitis, Asthma, Addison's disease,

Allergic rhinitis, Alzheimer's Disease, Amyotrophic lateral sclerosis (ALS, Lou Gehrig's Disease), Aplastic anemia, Arthritis, Atherosclerosis, Attention Deficit Hyperactivity Disorder (ADHD, ADD), Autism Spectrum Disorder, Benign Prostatic Hypertrophy, Bipolar Mood Disorder, Bronchiectasis, Cancer, Cardiac Failure, Cardiomyopathy, Chronic

5 Alcoholism, Chronic anemia, Chronic hepatitis, Chronic Amebiasis Chronic obstructive pulmonary disorder, Chronic gastritis, Chronic esophagitis, Chronic oral diseases, Chronic Poisoning, Chronic renal disease, Collagen diseases, Crohn's disease, Cushing's diseases, Cystic Fibrosis, Dementia, Dermatomyositis, Depression, Dysrhythmias, Eating Disorders (Anorexia Nervosa, Bulimia Nervosa, Binge Eating), Eczema, Endometriosis, Epilepsy,

10 Fibrosing Alveolitis, Gastro-Esophageal Reflex Disease (GERD), Glaucoma, Gout, Graves Disease, Hemophilia, Hyperlipidemia, Hyperpituitarism, Hypopituitarism, Hypophyseal Adenoma, Hypothyroidism, Idiopathic Thrombocytopenic Purpura, Menopause, Migraine, Motor Neuron Disease, Multiple Sclerosis, Muscular Dystrophy, Myasthenia Gravis, Myopathy, Neuropathy, Obesity, Obsessive Compulsive Disorder, Osteoarthritis,

15 Osteoporosis, Paralysis (Paraplegia, Quadriplegia), Parkinson's disease, Paget's disease, Polyarteritis Nodosa, Polycystic ovarian disease, Psoriasis, Psoriatic Arthritis, Pulmonary Interstitial Fibrosis, Reflex Sympathetic Dystrophy (RSD) Syndrome, Rheumatoid Arthritis, Scleroderma, Schizophrenia, Sjögren's syndrome, Smoking and Tobacco-related Diseases, Systemic Lupus Erythematosus, Trigeminal Neuralgia, Tuberculosis, Ulcerative Colitis or

20 Urinary Incontinence.

In another illustrative embodiment, the acute disease is Acute Alcohol Poisoning, Acute Amebiasis, Acute Auditory Diseases, Acute Brain Diseases, Acute Cancers, Acute Cardiovascular Diseases, Acute Dental Diseases, Acute Dermatological Diseases, Acute Esophageal Diseases, Acute Gastrointestinal Diseases, Acute Kidney Diseases, Acute Liver

25 Diseases, Acute Muscular Diseases, Acute Ocular Diseases, Acute Poisoning, Acute Oral Diseases, Acute Osteoarthritis, Acute Pharyngeal Diseases, Acute Prostatic Diseases, Acute Physical Injury (Trauma), Acute Peritoneal Diseases, Acute Respiratory Diseases or Acute Urogenital Diseases.

In another illustrative embodiment, the infectious disease is caused by Bacillus

30 anthracis, Bordetella pertussis, Borrelia burgdorferi, Brucella abortus, Brucella canis, Brucella melitensis, Brucella suis, Campylobacter jejuni, Chlamydomydia pneumoniae, Chlamydia

trachomatis, Chlamydia psittaci, Clostridium botulinum, Clostridium difficile, Clostridium perfringens, Clostridium tetani, Corynebacterium diphtheriae, Entamoeba histolytica, Enterococcus faecalis, Enterococcus faecium, Escherichia coli, Enterotoxigenic Escherichia coli, Enteropathogenic E. coli, Enterohemorrhagic Escherichia coli O157:H7, 5 Francisella tularensis, Fungus (Candida albicans), Haemophilus influenza, Helicobacter pylori, Leishmania (Kala Azar), Legionella pneumophila, Leptospira interrogans, Leptospira interrogans, Listeria monocytogenes, Mycobacterium leprae, Mycobacterium tuberculosis, Mycoplasma pneumoniae, Neisseria gonorrhoeae, Neisseria meningitidis, Parasites (protozoa, helminths, ectoparasites), Plasmodium (malaria), Pseudomonas aeruginosa, Rickettsia 10 rickettsia, Salmonella enterica subsp. Enterica, Salmonella typhimurium, Shigella sonnei, Staphylococcus aureus, Staphylococcus epidermidis, Staphylococcus saprophyticus, Streptococcus agalactiae, Streptococcus pneumoniae, Streptococcus pyogenes, Treponema pallidum, Vibrio cholerae or Yersinia pestis.

In yet another illustrative embodiment, the infectious disease is caused by Adenovirus, 15 Coxsackievirus, Dengue virus, Ebola virus, Epstein-Barr virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Herpes simplex virus, Cytomegalovirus, Human immunodeficiency virus (HIV), Influenza virus, Measles virus, Mumps virus, Human papillomavirus, Human parainfluenza virus, Poliovirus, Rabies virus, Human respiratory syncytial virus, Rubella virus, Severe acute respiratory syndrome (SARS) coronavirus, Varicella zoster virus or 20 Yellow fever virus.

In an illustrative embodiment, there is a kit for determining the concentration of phosphatase in stool, wherein the kit comprises of a tool for measuring a defined quantity of stool, a stool dilution buffer or water, a substrate for phosphatase and phosphatase assay 25 buffer. The tool for measuring a defined quantity of stool can be, for example, a spoon or a scoop with a predetermined volume. The concentration of phosphatase is measured using a spectrophotometer, a biochemistry analyzer, a high-performance liquid chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical, biochemical, electrical or electronic device.

In an illustrative embodiment, there is a kit for determining the concentration of 30 phosphatase in stool, wherein the kit comprises of a tool (e.g. a spoon or scoop with a predetermined volume) for measuring a defined quantity of stool, a substrate for phosphatase

and phosphatase assay buffer. The stool sample is dissolved in a stool dilution buffer or water. Stool suspension, filtered or unfiltered, is then mixed with a substrate of phosphatase, waiting a specified period of time to allow a color to develop, comparing the developed color with photographs of standards and quantifying the amount of alkaline phosphatase in the biological sample. As an alternative to quantifying the amount of alkaline phosphatase by comparing the developed color to photographs of standards, the quantifying can be done by counting pixels of photographs.

In another illustrative embodiment, the phosphatase is intestinal alkaline phosphatase, placental alkaline phosphatase, tissue nonspecific alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity.

In another illustrative embodiment, the substrate for phosphatase is provided in liquid or solid form.

In another illustrative embodiment, the substrate for phosphatase is attached to a solid medium comprising a membrane, paper, stick, glass, plastic, a bead or a slide.

In yet a further illustrative embodiment, there is an enzyme-linked immunosorbent assay (ELISA) system for quantifying intestinal alkaline phosphatase in stool.

In yet a further illustrative embodiment, there is a CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'-(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt)-based luminescent phosphatase assay system for quantifying intestinal alkaline phosphatase in stool.

In yet a further illustrative embodiment, there is a malachite green-based phosphate detection system for quantifying intestinal alkaline phosphatase in stool.

In another illustrative embodiment, the method for diagnosing or for determining the prognosis of incipient diabetes, an incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease or metabolic syndrome involves providing a solid or liquid medium containing a substrate for phosphatase, contacting a biological sample, preferably a stool sample, with the solid or liquid medium, waiting a specified period of time to allow a color to develop, comparing the developed color with photographs of standards and

quantifying the amount of alkaline phosphatase in the biological sample. As an alternative to quantifying the amount of alkaline phosphatase by comparing the developed color to photographs of standards, the quantifying can be done by counting pixels of photographs.

5 In another illustrative embodiment, the method for diagnosing or for determining the prognosis of incipient diabetes, an incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease or metabolic syndrome involves providing a solid medium for attaching a biological sample, preferably a stool sample, contacting a substrate for phosphatase with the solid medium, waiting a specified period of time to allow a color to develop, comparing the developed color with photographs of standards and quantifying the
10 amount of alkaline phosphatase in the biological sample. As an alternative to quantifying the amount of alkaline phosphatase by comparing the developed color to photographs of standards, the quantifying can be done by counting pixels of photographs.

In yet another illustrative embodiment, the method for diagnosing or for determining the prognosis of incipient diabetes, an incipient metabolic disease, incipient metabolic
15 syndrome, diabetes, metabolic disease or metabolic syndrome involves mixing a substrate for phosphatase with a biological sample, preferably a stool sample waiting a specified period of time to allow a color to develop, comparing the developed color with photographs of standards and quantifying the amount of alkaline phosphatase in the biological sample. As an alternative to quantifying the amount of alkaline phosphatase by comparing the developed
20 color to photographs of standards, the quantifying can be done by counting pixels of photographs.

In another illustrative embodiment, it is anticipated that deficiency of IAP might be associated with increase in serum endotoxins (LPS, LTA, etc.), proinflammatory nucleotides (UDP, ATP, etc.), cytokines (Interleukins IL-1 to IL-36; tumor necrosis factors TNF-alpha,
25 beta and gamma; interferons INF--alpha, beta and gamma; transforming growth factor TGF-beta; IFN-g inducing factor (IGIF), etc.), chemokines (CXCL-8, CCL2, CCL3, CCL4, CCL5, CCL11, CXCL10, granulocyte-macrophage colony-stimulating factor GM-CSF), granulocyte-colony stimulating factor (G-CSF), etc.), and antagonists of cytokine receptors. Further, it is anticipated that IAP deficiency might also be associated with decrease in anti-inflammatory
30 factors such as receptors of interleukins, TNFs and INFs, and some interleukins with dual pro- and anti-inflammatory functions (IL-4, IL-6, IL-10, IL-11, IL-13, etc.).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a flow chart showing the steps for determining the concentrations of intestinal alkaline phosphatase (IAP) in stool.

5 Figures 2A-C are graphs showing that the stool alkaline phosphatase activity is mostly due to intestinal alkaline phosphatase (IAP).

Figures 3A-C are graphs showing that patients with type 2 diabetes mellitus (T2DM) have low levels of intestinal alkaline phosphatase (IAP) in their stool.

10 Figure 4 is a graph showing that stool IAP levels are low in T2DM patients at all percentile points.

Figure 5 is a graph showing that high levels of stool IAP is protective against T2DM irrespective of body mass index (BMI).

DETAILED DESCRIPTION

15 Type 2 diabetes mellitus: Type 2 diabetes mellitus (T2DM) is characterized by hyperglycemia resulting from defects in insulin secretion, insulin action, or both. T2DM is diagnosed as having fasting plasma glucose (FPG) level > 7.0 mmol/l (126 mg/dl) or > 11.1 mmol/l (200 mg/dl) of plasma glucose during oral glucose tolerance test (OGTT) performed by ingestion of 75 gm glucose and measuring plasma glucose levels at specific time intervals
20 (min). T2DM causes long-term damage, dysfunction, and failure of different organs, especially the eyes, kidneys, nerves, heart, and blood vessels (American Diabetes Association. Diagnosis and classification of diabetes mellitus. Diabetes Care. 2014 Jan;37 Suppl 1:S81-90). It is a major global health problem that affected 387 million people worldwide in 2014 and cost 612 billion US dollars (Abdullah et al. Int J Endocrinol. 2014;593982; International
25 Diabetes Federation, IDF Diabetes Atlas, 2012, <http://www.idf.org/diabetesatlas>; <http://www.idf.org/diabetesatlas/update-2014>). In the United States, estimated 24.4 million people (9.2% population) had diabetes in 2013, costing 306 billion dollars (<http://www.idf.org/diabetesatlas>; <http://www.idf.org/diabetesatlas/update-2014>; Abdullah et al. Int J Endocrinol. 2014;593982). The American Diabetes Association estimated the total
30 cost of diagnosed diabetes in 2012 as \$245 billion, including \$176 billion in direct medical

costs and \$69 billion in reduced productivity (American Diabetes Association. Economic costs of diabetes in the U.S. in 2012. *Diabetes Care* 2013;36:1033–1046).

Etiologically, various factors have been postulated to be involved in the development of T2DM, such as autoimmunity, the metabolic syndrome, diets, obesity, infection, ethnicity, genetic polymorphism and predisposition, drugs, stress, sedentary lifestyle, pregnancy, etc. (Kahn et al. *Lancet* 2014;383(9922):1068-83; Stumvoll et al. *Lancet* 2005;365(9467):1333-46; Pietropaolo et al. *Diabetes* 2007;56(5):1189-97. Bakker et al. *Eur J Endocrinol* 2013;169(5):R99-R114; Lin and Sun. *J Endocrinol* 2009;204(1):1-11; Leahy. *Arch Med Res* 2005;36(3):197-209; DeFronzo. *Med Clin North Am* 2004;88(4):787-835; Holt. *Br J Psychiatry Suppl.* 2004 Apr;47:S55-63).

Recently, a low-grade systemic inflammation, induced by persistently increased levels of endotoxin lipopolysaccharides (LPS) in blood (metabolic endotoxemia), has been implicated as an etiological factor for the metabolic syndrome that eventually leads to T2DM (Cani et al., *Diabetes* 2007;56(7):1761-72). We have shown that mice deficient in the brush-border enzyme intestinal alkaline phosphatase (IAP) develop T2DM (Kaliannan et al. *Proc Natl Acad Sci U S A* 2013;110(17):7003-8). We have also shown that IAP detoxifies LPS and reduces metabolic endotoxemia, and oral supplementation with IAP not only prevents but also cures high-fat diet-induced metabolic syndrome and T2DM in wild-type mice T2DM (Kaliannan et al. *Proc Natl Acad Sci U S A* 2013;110(17):7003-8). We postulated that a low level of IAP in the intestinal lumen might be associated with T2DM in humans, and therefore, we decided to determine the levels of IAP in the stools of diabetic and healthy non-diabetic populations. In this invention, we report that, indeed, the diabetic patients have less amounts of IAP in their stools compared to the healthy non-diabetic population. Based on this observation, we propose to develop a personal temporal profile of IAP levels in stool, and if the profile of a healthy person shows that IAP levels are low and/or rapidly decreasing then we can predict that the person most probably have incipient diabetes. We anticipate that a personal temporal profile of IAP levels in stool would be highly valuable for diagnosis, treatment and assessing prognosis of any disease.

Persons having incipient diabetes currently have a FPG value in the normal range, but are at risk of developing diabetes based on a low concentration of phosphatase or a

concentration of phosphatase that is persistently decreasing over time as well as other known risk factors.

The Metabolic Syndrome: The metabolic syndrome consists of a group of related disorders that includes obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia (abnormal lipid profile), hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension and fatty liver (Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8; Grundy et al. Circulation. 2004 Jan 27;109(3):433-8; Hotamisligil. Nature 444(7121):860–867). The ‘overt’ metabolic syndrome has been defined as the co-existence of at least 3 out of 5 criteria encompassing obesity (body mass index (BMI) > 30.0 kg/m²), hyperglycemia (FPG > 5.5 mmol/l), hypertension (systolic blood pressure > 130 mmHg or diastolic blood pressure > 85 mmHg), hypertriglyceridemia (> 150 mg/dl) and low high-density lipoprotein (HDL) (< 40 mg/dl for males or < 50 mg/dl for females) (Huang PL. A comprehensive definition for metabolic syndrome. Dis Model Mech 2009;2(5-6):231-7). International Diabetic Federation estimates that approximately 25% of the world population have the metabolic syndrome (Kaur. Cardiol Res Pract. 2014;2014:943162; <http://www.idf.org/metabolic-syndrome>). Approximately 40% of the US population have the syndrome (Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8; Ford. Diabetes Care. 2005 Nov;28(11):2745-9). The metabolic syndrome leads to metabolic diseases such as T2DM, cardiovascular disease (coronary heart disease), cerebrovascular disease (stroke), nonalcoholic fatty liver disease, etc. The metabolic syndrome has become an epidemic in the USA with devastating consequences in terms of healthcare expenditure, morbidity and mortality (Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8; Malik et al. Circulation. 2004 Sep 7;110(10):1245-50).

Pathogenesis of the metabolic syndrome is poorly understood; however, recent work by Cani et al. (Diabetes 2007; 56:1761–1772) established that “metabolic endotoxemia” plays a critical role for the development of the metabolic syndrome. Endotoxin, also known as lipopolysaccharides (LPS), is the cell-wall component of Gram-negative bacteria. The metabolic endotoxemia is defined as a persistent increase of two- to three-fold endotoxin levels in the circulation compared to the normal levels (Cani et al. Diabetes 2007; 56:1761–1772; Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8).

Metabolic endotoxemia causes low-grade systemic inflammation as evidenced by increased serum levels of interleukin (IL)-1, IL-6 and tumor necrosis factor-alpha (TNF- α) (Cani et al. Diabetes 2007; 56:1761–1772; Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8). On the other hand, it is well-evidenced that chronic inflammation adversely affects the function of pancreatic beta cells (Hohmeier et al. Int J Obes Relat Metab Disord 27(Suppl 3):S12–S16), hepatocytes (Gieling et al. Am J Physiol Gastrointest Liver Physiol 296(6):G1324–G1331), and vascular endothelial cells (Hansson. N Engl J Med 352(16):1685–1695), and dysfunction of these cells is recognized to contribute to the metabolic syndrome.

Antibiotic-associated dysbiosis sometimes leads to *Clostridium difficile* infection as well as metabolic syndrome (Alam et al. Ann Surg. 2014 Apr;259(4):715-22; Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8; Parekh et al. Clin Transl Gastroenterol. 2015 Jun 18;6:e91). It has been shown that beta-lactamase prevents dysbiosis associated with a few specific antibiotics (Goldberg et al. Am J Health Syst Pharm. 2015 Jun 15;72(12):1007-12).

As described above, we have shown that IAP knockout mice develop the metabolic syndrome that ultimately leads to T2DM. We have also shown that oral IAP supplementation can prevent as well as cure the high fat diet-induced metabolic syndrome.

Persons having the incipient metabolic syndrome currently have normal or abnormal values for glucose, insulin, cholesterol, lipids, blood pressure, liver fat, BMI, etc., but are at risk of developing the metabolic syndrome based on a low concentration of phosphatase or a concentration of phosphatase that is persistently decreasing over time as well as other known risk factors.

Persons having the incipient metabolic syndrome also include persons with the pre-metabolic syndrome who have at least one criterion of the 'overt' metabolic syndrome as defined above.

Persons having incipient metabolic diseases currently have normal or abnormal metabolic function of different organs (e.g., liver, lung, kidney, pancreas, brain, muscle, adrenal glands, heart, skin, etc.), but are at risk of developing metabolic diseases based on a low concentration of phosphatase or a concentration of phosphatase that is persistently decreasing over time as well as other known risk factors.

Intestinal Alkaline Phosphatase: Intestinal alkaline phosphatase (IAP) is a membrane-bound glycoprotein that is exclusively expressed in villus-associated enterocytes of proximal small intestine and hence recognized as an enterocyte differentiation marker (Hodin et al. *Am J Physiol.* K1995 Aug;269(2 Pt 1):C385-91; Eliakim et al. *Biochim Biophys Acta.* 1991 Jan 10;1091(1):1-8). As the name suggests, IAP optimally functions at a high pH, and it hydrolyzes monophosphate esters suggesting its important role in fat and phosphate metabolism (Malo et al. *Am J Physiol Gastrointest Liver Physiol.* 2006 Apr;290(4):G737-46, and references therein). From the enterocytes the enzyme is bidirectionally secreted into the intestinal lumen as well as the systemic circulation (Eliakim et al. *Biochim Biophys Acta.* 1991 Jan 10;1091(1):1-8). IAP in the intestinal lumen travels downwards from the proximal small intestine to the distal large intestine and then excreted with stool (Malo et al., *Gut.* 2010;59:1476-1484). IAP is strictly conserved among species (Goldstein et al. *Proc Natl Acad Sci U S A.* 1980 May;77(5):2857-60) indicating an essential physiological role of IAP. Various isozymes of alkaline phosphatases (APs) exist, and they include IAP, placental AP, tissue nonspecific (liver/bone/kidney /neutrophils) AP, and germ cell AP. These various AP enzymes share significant structural homology as well as functional similarities (Kaliannan et al. *Proc Natl Acad Sci U S A* 2013;110(17):7003-8; Goldstein et al. *Proc Natl Acad Sci U S A.* 1980 May;77(5):2857-60). IAP protein is a component of the surfactant-like particle (SLP), a unilamellar secreted membrane associated with the process of fat absorption and secreted from the apical surface of enterocytes (Malo et al. *Am J Physiol Gastrointest Liver Physiol.* 2006 Apr;290(4):G737-46; Mahmood et al. *Am J Physiol Gastrointest Liver Physiol.* 2003 Aug;285(2):G433-41).

Physiologically, IAP exerts two very important functions with respect to its existence in the luminal bacterial environment; firstly, it maintains the normal homeostasis of intestinal microbiota, and secondly, it detoxifies bacterial toxins. We have shown that IAP knockout mice harbor fewer bacteria compared to its wild-type littermates (Malo et al., *Gut.* 2010;59:1476-1484) and IAP promotes the gut bacterial growth by reducing the concentrations of intestinal luminal nucleotide triphosphates that have toxic effect on bacterial growth (Malo et al. *Am J Physiol Gastrointest Liver Physiol.* 2014 May 15;306(10):G826-38). We and others have shown that IAP detoxifies various bacterial toxins, such as lipopolysaccharides (LPS), CpG DNA, flagellin and uridine diphosphate (UDP), and IAP

probably destroys these targets by dephosphorylation (phosphohydrolysis) (Poelstra et al. USPTO patent no. 6,290,952, September 2001; Bentala et al. Shock. 2002 Dec;18(6):561-6; Chen et al. Am J Physiol Gastrointest Liver Physiol. 2010 Aug;299(2):G467-75; Moss et al. Am J Physiol Gastrointest Liver Physiol. 2013 Mar 15;304(6):G597-604; Malo et al. Am J Physiol Gastrointest Liver Physiol. 2014 May 15;306(10):G826-38). Recently, using mice deficient in IAP (IAP knockout) Narisawa et al. (Mol Cell Biol. 2003 Nov;23(21):7525-30) showed that IAP limits fat absorption, and IAP knockout mice become obese when fed a high fat diet.

Pharmacologically, various therapeutical uses of IAP have been described in different animal models as well as in humans. In a mouse model, we have shown that oral supplementation of IAP prevents antibiotic-induced susceptibility to enteric pathogens such as *Salmonella Typhimurium*, and *Clostridium difficile* (Malo et al., Gut. 2010;59:1476-1484; Alam et al. Ann Surg. 2014 Apr;259(4):715-22). We have shown that oral IAP supplementation not only prevents but also cures the high fat diet-induced metabolic syndrome in mice (Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8). Kiss has recently claimed use of IAP to correct insulin resistance, hyperinsulinemia and dyslipidemia in a patent application with USPTO (Kiss, USPTO patent application no. US 2013/0251701 A1, Sep. 26, 2013). Kiss has also previously claimed use of IAP to treat obesity, diabetes and cancer (USPTO patent nos. 7,858,085; 8,603,464 & 7,943,606).

Lukas et al. showed efficacious value of IAP in treating ulcerative colitis in humans (Lucas et al. Inflamm Bowel Dis. 2010 Jul;16(7):1180-6). In mouse models we have also shown that IAP has beneficial effects to treat chronic colitis (Ramasamy et al. Inflamm Bowel Dis. 2011 Feb;17(2):532-42). Whitehouse et al. reported a protective role of intestinal alkaline phosphatase in necrotizing enterocolitis (Whitehouse et al. J Surg Res. 2010 Sep;163(1):79-85). Van Veen et al. showed that bovine intestinal alkaline phosphatase attenuates the inflammatory response in secondary peritonitis in mice (van Veen et al. Infect Immun. 2005 Jul;73(7):4309-14). We have demonstrated that local peritoneal irrigation with IAP is protective against peritonitis in mice (Ebrahimi et al. J Gastrointest Surg. 2011 May;15(5):860-9).

General Methodology

Participants: A case-control study was used to assess the difference in the concentrations of IAP in stools of diabetic and control non-diabetic healthy populations. Participants, aged 30-70 yr., were recruited from a suburban community of Dhaka, Bangladesh by advertisement through local dignitaries, hospitals, clinics and physicians' offices. Based on preliminary data (unpublished) the sample size of each group was calculated to achieve statistical power of 80% or more (continuous endpoint, $\alpha = 0.05$). The study included 202 diabetic cases (63 males and 139 females) and 445 healthy control subjects (114 males and 331 females) of the same ethnicity. The study included more females than males just because of more accessibility to female participants. People with T2DM were diagnosed by a physician for at least 6 months prior to their recruitment, and were on oral antihyperglycemic agents and/or insulin medication. Newly diagnosed persons with hyperglycemia (FPG > 7.0 mmol/l) were also included in the diabetic group (see below). All participants were on unrestricted diets. Any person suffering from an acute disease was excluded from the study. Pregnant women, patients with Type 1 diabetes and history of cancer were excluded. All participants were tested for renal, hepatic and cardiovascular diseases, and participants with clinically significant renal, hepatic and cardiovascular diseases were excluded. The participants did not have any history of chronic alcohol consumption. Underweight persons (body mass index, BMI < 18.5 kg/m²) were also excluded.

Study approval. An informed consent form to participate in the study was signed by each participant. The Institutional Review Board (IRB) of Harvard Biotech BD Ltd. (Dhaka, Bangladesh) reviewed and approved the study.

Laboratory tests, physical examination and socio-medical history: Laboratory tests were performed using commercially available biochemical assay kits (Linear Chemicals S.L., Barcelona, Spain) and an automatic biochemistry analyzer (Sinnova Medical Science & Technology Co., Ltd, Nanjing, Jiangsu, China; Model: Sinnolab MT 5000, Version 5.00). The diabetic status of each participant (diabetics and controls) was confirmed by measuring fasting (at least 10 hours) plasma glucose (FPG). An FPG level > 7.0 mmol/l (126 mg/dl, milligrams/deciliter), was considered diagnostic for diabetes (American Diabetes Association, 2014). Any newly diagnosed diabetic patient (FPG > 7.0 mmol/l) was included in the diabetic group. All participants were also subjected to serum biochemical tests for cholesterol, low-density lipoproteins (LDL), high-density lipoprotein (HDL), triglycerides, creatinine and

ALT. Height and weight of each participant were measured to calculate the body mass index (BMI) defined as weight in kg divided by the square of height in meter (kg/m^2). Participants were asked for the history of alcohol consumption. Medical history was used to discover participants with kidney, liver and heart diseases. The status of cardiovascular disease was also evaluated by measuring blood pressure.

Homogenization of stool: The supernatant of a homogenized stool suspension was used for IAP assay. A small amount of stool (milligrams) was measured and then the 'stool dilution buffer' (10 mM (millimolar) Tris-HCl, pH 8.0, 1 mM magnesium chloride, 10 μM (micromolar) zinc chloride) at a defined ratio was added. Usually, 50 μl (microliter) of stool dilution buffer was added to 1 mg of stool. The sample was vigorously vortexed to prepare a homogenized stool suspension, which was then centrifuged at 10,000 x g for 20 min, and the supernatant containing IAP was collected and assayed for IAP concentration. Alternatively, stool can be suspended in water, however, stool suspended in water shows a bit lower IAP activity as compared to stool suspended in stool dilution buffer.

Alkaline Phosphatase Assay. The stool supernatant was assayed for alkaline phosphatase (AP) following an established protocol using the automatic biochemistry analyzer mentioned above (Nanjing, Jiangsu, China). In brief, 20 μl of supernatant was added to 1 ml of enzyme assay buffer (1.25 M diethanolamine (DEA) buffer, pH 10.2, 0.6 mM magnesium chloride) containing 10 mM p-nitrophenyl phosphate (pNPP), and the reaction mixture was incubated for one min at 37°C followed by measuring the AP concentration by the analyzer pre-calibrated with AP standards. To determine the major isoform among stool APs, prior to assaying for AP activity an aliquot of stool sample was treated for 10 min with L-phenylalanine (L-Phe, 10 mM final conc.), a specific inhibitor of IAP, and another aliquot with L-homoarginine (L-Arg, 10 mM final conc.), an inhibitor of tissue non-specific alkaline phosphatase (TNAP). Each aliquot treated with an inhibitor was then mixed with the reaction buffer containing an equal concentration (10 mM) of the respective inhibitor, and assayed for AP activity using the analyzer. Because most of the AP activity in stool was due to IAP, the stool AP values were expressed as units of IAP/g stool. All AP assays were performed by a single laboratory technologist who was blinded to the diagnoses of participants.

Statistical Analysis: The SAS System (SAS Institute, Cary, North Carolina) was used for statistical analysis. Mean and standard errors were calculated for T2DM cases and non-

T2DM controls stratified by sex. The correlation between IAP levels and various risk factors for T2DM was assessed via Pearson correlation coefficient stratified by sex and T2DM status (T2DM and non-T2DM controls). Mean differences in IAP levels between T2DM cases and non-T2DM controls were assessed via linear regression models controlling for the effects of age, sex, FPG and BMI on IAP levels and T2DM status. The statistical significance of the variance associated with independent variables were assessed from sum of square III using GLM procedure in SAS. Multiple logistic regression using Proc Logist procedure in SAS assessed association between T2DM cases with independent risk factors including IAP. Regression coefficients and odds ratios were used to assess the independent risk contribution of IAP to T2DM status. The statistical significance of the difference between two groups was determined using unpaired two-tailed Student's t-Test. The difference between two groups was considered significant when the p value was < 0.05. Student's t-Test was performed using Microsoft Excel program. Post-hoc statistical power analysis of two independent groups was performed using an online program (<http://clincalc.com/Stats/Power.aspx>).

EXAMPLES

The following examples are provided to further describe the invention; however, these examples do not limit the scope of the invention defined in the claims.

Example 1: The flow chart showing the steps for determining the concentrations of intestinal alkaline phosphatase (IAP) in stool.

Mice deficient in intestinal alkaline phosphatase (IAP-knockout) have been shown to develop type 2 diabetes mellitus (T2DM). Based on this observation we hypothesized that the T2DM patients might have less amount of IAP in their stools. Therefore, we planned to measure the IAP concentration in the stools of T2DM patients and control healthy participants. A schematic presentation of the protocol to determine IAP in stool is provided in Figure 1.

Example 2: Stool alkaline phosphatase activity is mostly due to intestinal alkaline phosphatase.

IAP knockout ($Akp3^{-/-}$) mice develop T2DM, and based on this observation we hypothesized that humans with T2DM might have IAP deficiency. Accordingly, we decided to measure IAP concentrations in the stools of diabetic and control non-diabetic healthy people. However, taking in consideration of possible association of different isoforms of

alkaline phosphatases (APs) with T2DM in humans, we first decided to determine the nature of major AP isoform in the stools of both diabetic and control healthy populations. Stool samples of non-diabetic healthy controls and T2DM patients were homogenized in a stool dilution buffer followed by centrifuging and collection of supernatant. We performed AP assays on individual aliquots of a stool sample supernatant in presence of L-phenylalanine (L-Phe, a specific inhibitor of IAP) as well as L-homoarginine (L-Arg, a specific inhibitor of TNAP).

Figure 2 is a graph showing stool alkaline phosphatase activity is mostly due to intestinal alkaline phosphatase (IAP). L-Phe inhibited approx. 80% of the AP activity in the stool of total healthy population, and the inhibition was approx. 70% in total diabetic population, however, the difference was statistically significant ($p = 1.4E-04$) (Figure 2A). On the other hand, L-Arg had no inhibitory effect on the stool AP activity of either group. Sex-matched distribution also showed similar AP inhibitory effect of L-Phe in males (Figure 2B) and females (Figure 2C) of both control and diabetic groups. This result indicates that most of the stool AP activity is due to IAP, and accordingly, we decided to refer stool AP as IAP in the following sections as it is applicable.

Figure 2: Stool alkaline phosphatase activity is mostly due to intestinal alkaline phosphatase (IAP). Stool samples of non-diabetic healthy controls and T2DM patients were homogenized in a stool dilution buffer followed by centrifuging and collection of supernatant. The supernatant was assayed for alkaline phosphatase (AP) concentration in presence of L-phenylalanine (L-Phe), a specific inhibitor of IAP as well as in presence of L-homoarginine (L-Arg), a specific inhibitor of tissue nonspecific alkaline phosphatase (TNAP), using an automatic biochemistry analyzer. (A) Effects of AP inhibitors on the AP activity in the stools of total control and diabetic populations. (B) Effects of AP inhibitors on the AP activity in the stools of male control and diabetic populations. (C) Effects of AP inhibitors on the AP activity in the stools of female control and diabetic populations. Statistics: Values are expressed as mean $\pm$ SEM. Statistical significance of the difference between two respective groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. **, $p < 0.01$; ***, $p < 0.001$.

Example 3: The levels of intestinal alkaline phosphatase (IAP) is less in the stools of diabetic patients compared to healthy controls.

Table 1 shows a few important characteristics along with the concentrations of IAP in diabetic and healthy populations. The study included 202 diabetic patients (T2DM) and 445 control healthy subjects. The diabetic group included 63 men and 139 women, whereas the control group comprised of 114 men and 331 women. The ages of the participants ranged from 30 to 70 years. A comparative analysis of IAP values of diabetic and healthy participants is provided. It is evident that compared to the total healthy population the total diabetic population has 47.6% less IAP in their stools (Mean $\pm$ SEM: 67.4 $\pm$ 3.2 vs 35.3 $\pm$ 2.5 units/g stool, respectively; $p = 5.6E-10$). Compared to healthy males, diabetic males have 51.6% less IAP (57.1 $\pm$ 4.5 vs 27.7 $\pm$ 3.2 units/g stool, respectively; $p = 1.1E-05$). Similarly, in comparison to healthy females, diabetic females have 45.3% less IAP (71.0 $\pm$ 4.0 vs 38.8 $\pm$ 3.3 units/g stool, respectively; $p = 1.6E-06$). It appears that healthy control males have the IAP levels 19.5% lower than the healthy females, however, the difference is not significant ($p = 0.061$). On the other hand, the diabetic males have 28.7% lower IAP levels compared to diabetic females, and the difference is statistically significance ($p = 0.041$). It is noteworthy that the numerical value of IAP concentration of a sample varies depending on different alkaline phosphatase assays and/or variation in assay conditions.

Table 1. Characteristics of healthy control participants and T2DM patients.

Type of Participants	Total Participants		Males		Females	
	Healthy	T2DM	Healthy	T2DM	Healthy	T2DM
No. of Participants	445	202	114	63	331	139
Age of Participants (yr.)	47.1 $\pm$ 0.4	47.6 $\pm$ 0.7	48.7 $\pm$ 0.8	50.9 $\pm$ 1.2	46.7 $\pm$ 0.5	46.0 $\pm$ 0.8
Weight (kg)	59.3 $\pm$ 0.5	59.6 $\pm$ 0.7	65.0 $\pm$ 0.9	65.7 $\pm$ 1.3	57.3 $\pm$ 0.6	56.9 $\pm$ 0.8
Height (m)	1.52 $\pm$ 0.00	1.52 $\pm$ 0.01	1.62 $\pm$ 0.01	1.61 $\pm$ 0.01	1.48 $\pm$ 0.00	1.48 $\pm$ 0.01
BMI (kg/m ²)	25.7 $\pm$ 0.2	26.0 $\pm$ 0.3	24.7 $\pm$ 0.3	25.5 $\pm$ 0.5	26.1 $\pm$ 0.2	26.2 $\pm$ 0.4
Systolic Blood Pressure (mmHg)	134.2 $\pm$ 1.1	137.3 $\pm$ 1.5	134.4 $\pm$ 1.7	133.3 $\pm$ 2.5	134.2 $\pm$ 1.4	139.0 $\pm$ 1.8
Diastolic Blood Pressure (mmHg)	78.9 $\pm$ 0.6	80.1 $\pm$ 0.8	80.0 $\pm$ 1.1	80.8 $\pm$ 1.4	78.5 $\pm$ 0.7	79.8 $\pm$ 1.0
Creatinine (mg/dl)	0.98 $\pm$ 0.15	0.91 $\pm$ 0.02	0.89 $\pm$ 0.03	0.95 $\pm$ 0.03	1.01 $\pm$ 0.20	0.89 $\pm$ 0.03
Cholesterol (mg/dl)	162.7 $\pm$ 1.3	172.5 $\pm$ 2.2***	160.3 $\pm$ 2.6	164.4 $\pm$ 4.1	162.7 $\pm$ 1.3	176.2 $\pm$ 2.5***

HDL (mg/dl)	38.1 +/- 0.4	38.8 +/- 0.5	36.7 +/- 0.7	38.6 +/- 0.9	38.6 +/- 0.4	38.8 +/- 0.6
LDL (mg/dl)	94.6 +/- 1.0	102.2 +/- 2.0***	93.2 +/- 2.5	94.5 +/- 4.0	95.0 +/- 1.1	105.7 +/- 2.2***
Triglycerides (mg/dl)	152.2 +/- 2.3	159.4 +/- 3.8	153.5 +/- 5.9	162.3 +/- 7.6	151.7 +/- 2.4	158.1 +/- 4.3
ALT (U/l)	44.8 +/- 0.9	47.7 +/- 1.7	46.9 +/- 2.4	49.0 +/- 2.8	44.1 +/- 0.8	47.1 +/- 2.1
FPG (mmol/l)	4.4 +/- 0.0	8.2 +/- 0.2***	4.4 +/- 0.1	8.3 +/- 0.4***	4.4 +/- 0.0	8.2 +/- 0.3***
IAP (U/g stool)	67.4 +/- 3.2	35.3 +/- 2.5***	57.1 +/- 4.5	27.7 +/- 3.2***	71.0 +/- 4.0	38.8 +/- 3.3***

Note: The participants were recruited from a suburb of Dhaka, Bangladesh. All participants were on overnight (10 h) fasting and investigated for all the physical and biochemical tests described above. Statistics: Values are expressed as mean +/- SEM.

- 5 Statistical significance of the difference between two respective groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. ***, $p < 0.001$. Percentage loss of IAP in T2DM patients compared to healthy controls: Total, 47.6%; Male, 51.6%; Female, 45.3%.

- 10 Table 2 shows the age group- and gender-matched distribution of stool IAP levels in the healthy control and diabetic populations. It is apparent that compared to non-diabetic group, T2DM patients have approx. 35-60% less mean stool IAP, and the difference between two respective groups is statistically significant. The least reduction (36.8%) in IAP level is observed in 40-49 yr. old diabetic females, whereas the highest reduction (59.0%) is evidenced in 60-70 yr. old male diabetes patients. However, there is no statistically significant difference in IAP levels between the younger and older groups of the same population.
- 15

Table 2. Age group- and gender-matched distributions of IAP along with a few critical parameters of T2DM.

Group	Type of Participants	Number of Participants	Age of Participants (years)	Body Mass Index (BMI) (kg/m ²)	Fasting Plasma Glucose (FPG) (mmol/l)	IAP Conc. in stool (U/gm Stool)	Loss of IAP compared to Controls (%)
30-39 yr. old	Healthy Controls	Total : 109	35.7 +/- 0.1	27.0 +/- 0.5	4.5 +/- 0.1	78.1 +/- 7.6	NA
		Males : 22	36.6 +/- 0.3	24.8 +/- 0.6	4.5 +/- 0.1	69.9 +/- 10.2	NA
		Females : 87	35.5 +/- 0.1	27.5 +/- 0.6	4.4 +/- 0.1	80.1 +/- 9.2	NA
	Diabetic Patients	Total : 43	35.0 +/- 0.3	25.8 +/- 0.7	7.7 +/- 0.4	40.6 +/- 6.1	48.0
		Males : 8	35.4 +/- 1.1	24.3 +/- 1.2	8.9 +/- 1.1	29.8 +/- 5.6	57.4
40-49 yr. old	Healthy Controls	Females : 35	34.9 +/- 0.3	26.2 +/- 0.8	7.5 +/- 0.4	43.0 +/- 7.4	46.3
		Total	1.2E-02	1.9E-01	5.2E-25***	3.6E-03**	NA
		Males	1.3E-01	6.6E-01	5.4E-07***	2.9E-02**	NA
	Diabetic Patients	Females	4.5E-02	1.9E-01	8.2E-20***	1.6E-02**	NA
		Total : 138	43.3 +/- 0.2	25.9 +/- 0.3	4.5 +/- 0.1	70.1 +/- 4.3	NA
	Healthy Controls	Males : 32	43.9 +/- 0.5	24.8 +/- 0.5	4.4 +/- 0.2	62.5 +/- 6.8	NA

		Females : 106	43.1 +/- 0.3	26.2 +/- 0.4	4.5 +/- 0.1	72.3 +/- 5.2	NA
	Diabetic Patients	Total : 69 Males : 19 Females : 50	43.3 +/- 0.3 43.9 +/- 0.6 43.1 +/- 0.4	25.3 +/- 0.4 25.1 +/- 0.8 25.3 +/- 0.5	8.4 +/- 0.4 9.0 +/- 0.8 8.2 +/- 0.5	43.6 +/- 5.1 38.1 +/- 7.3 45.7 +/- 6.5	37.8 39.1 36.8
	p Value (Controls vs Diabetics)	Total Males Females	9.2E-01 9.8E-01 9.9E-01	2.3E-01 7.9E-01 1.8E-01	8.7E-27*** 1.1E-08*** 3.5E-19***	2.6E-04*** 2.4E-02* 3.1E-03**	NA NA NA
50-59 yr. old	Healthy Controls	Total : 131 Males : 48 Females : 83	52.3 +/- 0.2 53.1 +/- 0.4 51.8 +/- 0.3	25.2 +/- 0.3 24.8 +/- 0.5 25.4 +/- 0.4	4.4 +/- 0.1 4.5 +/- 0.1 4.4 +/- 0.1	60.2 +/- 6.9 49.3 +/- 7.8 66.5 +/- 9.9	NA NA NA
		Total : 52 Males : 21 Females : 31	52.7 +/- 0.4 54.1 +/- 0.6 51.8 +/- 0.4	27.4 +/- 0.7 25.8 +/- 0.6 28.4 +/- 1.1	8.3 +/- 0.5 8.3 +/- 0.7 8.3 +/- 0.8	26.8 +/- 3.2 22.5 +/- 3.9 29.7 +/- 4.6	55.5 54.5 55.3
		Total Males Females	3.1E-01 1.7E-01 9.8E-01	1.2E-03** 2.0E-01 1.9E-03**	5.6E-22*** 1.4E-11*** 1.8E-12***	3.2E-03** 3.0E-02* 2.8E-02*	NA NA NA
	Diabetic Patients	Total : 67 Males : 12 Females : 55	63.0 +/- 0.4 63.8 +/- 0.9 62.9 +/- 0.5	24.4 +/- 0.5 23.3 +/- 0.9 24.7 +/- 0.6	4.4 +/- 0.1 4.3 +/- 0.2 4.4 +/- 0.1	58.7 +/- 6.5 50.3 +/- 12.3 60.5 +/- 7.5	NA NA NA
		Total : 38 Males : 15 Females : 23	62.4 +/- 0.7 63.7 +/- 1.0 61.7 +/- 0.9	25.7 +/- 0.8 26.4 +/- 1.5 25.2 +/- 0.9	8.3 +/- 0.6 7.2 +/- 0.9 8.9 +/- 0.8	26.2 +/- 4.5 20.7 +/- 6.9 29.8 +/- 6.0	55.4 59.0 50.8
		Total Males Females	5.6E-01 9.6E-01 2.4E-01	1.5E-01 1.5E-01 5.3E-01	2.6E-12*** 1.5E-02* 4.0E-12***	1.2E-03** 1.8E-02* 2.6E-02*	NA NA NA

Statistics: Values are expressed as mean +/- SEM. Statistical significance of the difference between two groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. NA, not applicable.

Figure 3 is a graph showing that IAP levels are less in the stools of diabetic patients compared to the healthy controls.

Comparison of IAP levels between the control healthy population and diabetic patients showed that on average the entire (total) diabetic population has much less IAP in their stools (Figure 3A). Gender-matched distribution showed that both male and female diabetic patients have less stool IAP compared to their healthy counterparts (Figure 3A). Age group- and gender-matched distributions of IAP also showed that the enzyme apparently decreases with age both in males (Figure 3B) and females (Figure 3C) of non-diabetic and diabetic populations.

Figure 3: Patients with type 2 diabetes mellitus (T2DM) have low levels of intestinal alkaline phosphatase (IAP) in their stool. Stool samples of healthy participants and T2DM patients were assayed for IAP concentration using an automatic biochemistry analyzer (see Figure 2). (A) IAP concentrations in the stools of total non-diabetic healthy control and diabetic populations. (B) Age-dependent distribution of IAP concentrations in male control and diabetic populations. (C) Age-dependent distribution of IAP concentrations in female control and diabetic populations. Statistics: Values are expressed as mean +/- SEM. Statistical

significance of the difference between two groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. The post-hoc statistical power analyses revealed the powers for respective total, male and female groups to be 100%, 100% and 100%, respectively, validating the adequacy of power (conventionally, $> 80\%$ power at $\alpha = 0.05$) for respective sample sizes. The difference in IAP values between younger and older groups of the same population (diabetic or non-diabetic) is not statistically significant.

Figure 4 is a graph showing that IAP levels are low in T2DM patients at all percentile points.

Taking in consideration of any influence of 'outliers' (few extremely high or low individual values compared to most values) in the average (mean) values of IAP levels as shown above (Figure 3A), we further verified the difference in IAP levels of diabetic and non-diabetic groups at different percentile points. We evaluated the percentile distribution of IAP values in 202 T2DM patients and 445 healthy controls. We organized individual IAP values of each group (diabetic or non-diabetic) from the lowest to the highest, and then calculated the average IAP value within each 10 percentile divisions. It is apparent that at all percentile points IAP values are less in T2DM patients compared to the healthy population, and the difference is highly significant. The percentile distribution of IAP levels confirms that T2DM patients, indeed, have lower levels of IAP compared to their counterparts.

Figure 4: IAP levels are low in T2DM patients at all percentile points. Individual IAP values from each group (Healthy Controls or T2DM Patients) were arranged from the lowest to the highest, and then the average IAP value within each 10th percentile was calculated ($n = 20$ within each 10th percentile for T2DM Patients, and $n = 44$ within each 10th percentile for Healthy Controls). Average values for corresponding percentiles are plotted. Statistics: Values are expressed as mean $\pm$ SEM. Statistical significance of the difference between two respective groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. ***, $p < 0.001$.

Example 4: IAP deficiency is associated with T2DM.

Table 3 shows the Pearson correlation coefficient analysis between the IAP level and age, gender, FPG, BMI, ALT, serum creatinine, blood pressure and lipid parameters. It is evident that there is no correlation between IAP and the above-mentioned T2DM risk factors.

Table 3. Pearson correlation coefficients showing no correlation of IAP levels with different risk factors of T2DM.

Type of participants	Male		Female	
	Healthy	T2DM	Healthy	T2DM
Risk factors	IAP correlation	IAP correlation	IAP correlation	IAP correlation
Age	-0.192	-0.221	-0.08	-0.131
BMI	-0.024	0.088	0.085	0.008
Creatinine	-0.013	-0.069	0.07	0.055
Total cholesterol	0.267	-0.132	0.004	0.127
HDL-cholesterol	0.101	-0.007	0.005	0.206
LDL-cholesterol	0.176	-0.130	-0.014	0.080
Triglycerides	0.138	0.014	-0.007	0.074
ALT	0.156	0.107	-0.065	-0.013
FPG	-0.076	0.039	0.05	-0.014
Systolic blood pressure	0.169	-0.257	-0.031	-0.153
Diastolic blood pressure	-0.040	-0.059	0.039	-0.060

Note: A Pearson correlation coefficient close to +1 or -1 indicates that the two variables are highly correlated (positively or negatively, respectively). A correlation coefficient between 0 and +0.30 or between 0 and -0.30 was considered of having no correlation between the two variables.

Table 4 shows a generalized linear regression model predicting a strong association of IAP with T2DM. It also predicts a mild association of IAP with age validating the observation that IAP decreases with age (see Figure 3B & C).

Table 4. A generalized linear regression model predicts an association of IAP with T2DM.

Source	DF	Type III SS	F Value	Pr > F
Diabetes	1	81665.00	23.3	< 0.0001
Sex	1	12309.33	3.51	0.0614
FPG (mmol/l)	1	180.46	0.05	0.8206
BMI (kg/m ²)	1	5335.29	1.52	0.2177
Age	1	23707.51	6.76	0.0095

Note: The model predicts a strong association of IAP with T2DM. It also predicts a mild association of IAP with age (IAP level gradually decreases with age; see Figure 3B & C).

Table 5 shows multiple logistic regression analyses predicting an independent inverse relationship between the IAP level and diabetes status. We performed multiple logistic regression analyses controlling for age, FPG, BMI, ALT, serum creatinine, blood pressure and

lipid parameters. The data showed an independent inverse relationship between the IAP level and diabetes status (Table 5). With each 25 U/g decrease in stool IAP, there is 35% increased risk of diabetes. We conclude that IAP deficiency is associated with diabetes independent of other known diabetic risk factors.

5

Table 5. Multiple logistic regression analyses predict an association of IAP deficiency with T2DM.

Explanatory variables	Logistic coefficients per unit change	Odds ratio (95% CI)
Age (yr.)	0.0226	1.023 (0.993-1.054)
Body mass index (BMI, kg/m ²)	0.0466	1.048 (0.986-1.114)
Creatinine (mg/dl)	-0.0166	0.983 (0.847-1.142)
Total cholesterol (mg/dl)	0.00183	1.002 (0.984-1.020)
HDL-cholesterol (mg/dl)	-0.0599	0.942 (0.900-0.985)
LDL-cholesterol (mg/dl)	0.0206	1.021 (1.001-1.041)
Triglycerides (mg/dl)	0.00473	1.005 (0.998-1.011)
ALT (U/l)	0.00944	1.009 (0.998-1.021)
Systolic blood pressure (mmHg)	-0.00012	1.000 (0.984-1.016)
Diastolic blood pressure (mmHg)	-0.0139	0.986 (0.958-1.015)
FPG (mmol/l)	1.3563	3.882 (2.972-5.070)
IAP (U/g stool)	-0.0145	0.986 (0.978-0.993)

Note: With each U/g decrease in stool IAP level there is 1.4% increase in the odds of diabetes diagnosis. For example, if there is 25 U/g decrease in IAP level then there will be a 35% increased risk of diabetes. Statistics: Proc Logist procedure (SAS) was used for multiple logistic regression analyses determining association between T2DM with independent risk factors including IAP.

Example 5: Hyper- and normo-glycemic T2DM patients have similar levels of stool IAP.

T2DM patients on rigorous antidiabetic medications and life-style changes very often achieve normoglycemic status (FPG < 7.0 mmol/l). We were interested to know whether the glycemic status of diabetic patients has any effect on the stool IAP level.

Table 6 shows that both hyperglycemic (FPG > 7.0 mmol/l) and normoglycemic diabetics have similar levels of IAP in their stools, and the difference in values between two respective groups is statistically insignificant.

Table 6. Hyper- and normo-glycemic diabetes patients have similar levels of stool IAP.

Type of Participants	Number of Participants	Age of Participants (years)	Body Mass Index (BMI) (kg/m ²)	Fasting Plasma Glucose (FPG) (mmol/l)	IAP Conc. (U/gm Stool)
Hyperglycemic diabetes patients (FPG > 7.0 mmol/l)	Total : 119	47.0 +/- 0.9	25.9 +/- 0.4	10.3 +/- 0.3	37.7 +/- 3.4
	Males : 36	49.1 +/- 1.4	25.8 +/- 0.7	10.6 +/- 0.4	29.6 +/- 4.4
	Females : 83	46.1 +/- 1.1	26.0 +/- 0.5	10.1 +/- 0.3	41.3 +/- 4.4
Normoglycemic diabetes patients (FPG < 7.0 mmol/l)	Total : 83	48.3 +/- 1.1	26.1 +/- 0.5	5.2 +/- 0.1	32.0 +/- 3.8
	Males : 27	53.3 +/- 2.1	25.2 +/- 0.7	5.3 +/- 0.2	25.1 +/- 4.6
	Females : 56	45.9 +/- 1.2	26.5 +/- 0.7	5.2 +/- 0.1	35.2 +/- 5.2
p Value (Hyperglycemic vs Normoglycemic)	Total	3.6E-01	7.9E-01	1.8E-33***	2.6E-01
	Males	9.3E-02	6.0E-01	2.0E-14***	5.0E-01
	Females	8.9E-01	5.3E-01	5.0E-21***	3.8E-01

Note: Diabetic patients were categorized into hyperglycemic (FPG > 7.0 mmol/l) and normoglycemic (FPG < 7.0 mmol/l) groups, and respective stool IAP levels were compared.

5 Statistics: Values are expressed as mean +/- SEM. Statistical significance of the difference between two groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. ***, $p < 0.001$.

Example 6: High levels of IAP is protective against diabetes irrespective of body mass index (BMI).

10 It is well recognized that overweight (BMI > 25.0 kg/m²) and obese (BMI > 30.0 kg/m²) populations are more vulnerable for developing T2DM than the population with normal weights (BMI < 25.0 – 18.5 kg/m²). We categorized the diabetics as well as healthy controls of this study in two groups, one with BMI > 25.0 kg/m² and the other having BMI < 25.0 kg/m².

15 Table 7 shows that a high level of IAP is protective against diabetes irrespective of body mass index (BMI). It is evident that both groups of diabetics (BMI > 25.0 kg/m² and BMI < 25.0 kg/m²) have significantly lower amounts of IAP compared to their healthy counterparts. These data suggest that IAP plays a protective role against the development of T2DM. An obese or overweight person with high IAP does not develop T2DM. Both groups
20 of diabetes patients have similar levels of IAP, and the difference is statistically insignificant ($p = 0.77$). We also observed that the levels of IAP in both groups of healthy controls are similar and the difference is insignificant ($p = 0.29$).

Table 7. A high level of IAP is protective against diabetes irrespective of body mass index (BMI).

Group	Type of Participants	Number of Participants	Age of Participants (years)	Body Mass Index (BMI) (kg/m ²)	Fasting Plasma Glucose (FPG) (mmol/l)	IAP Conc. in stool (U/gm stool)	Loss of IAP compared to Controls (%)
High BMI (> 25.0 kg/m ²)	Healthy Controls	Total : 227	45.7 +/- 0.6	28.8 +/- 0.2	4.5 +/- 0.1	66.5 +/- 4.4	NA
		Males : 49	47.8 +/- 1.2	27.4 +/- 0.3	4.5 +/- 0.1	52.0 +/- 7.3	NA
		Females : 178	45.1 +/- 0.7	29.2 +/- 0.3	4.4 +/- 0.1	70.5 +/- 5.2	NA
	Diabetic Patients	Total : 101	48.0 +/- 1.0	29.3 +/- 0.4	8.1 +/- 0.3	36.5 +/- 3.5	45.1
		Males : 28	52.5 +/- 1.9	28.7 +/- 0.7	8.3 +/- 0.6	29.0 +/- 5.1	44.2
		Females : 73	46.3 +/- 1.0	29.5 +/- 0.5	8.0 +/- 0.4	39.4 +/- 4.4	44.1
	p Value (Controls vs Diabetics)	Total	3.5E-02*	2.7E-01	5.6E-41***	2.3E-05***	NA
Low BMI (< 25.0 kg/m ²)	Healthy Controls	Total : 217	48.5 +/- 0.7	22.4 +/- 0.1	4.4 +/- 0.1	63.0 +/- 4.0	NA
		Males : 66	49.0 +/- 1.1	22.5 +/- 0.2	4.4 +/- 0.1	61.3 +/- 5.7	NA
		Females : 151	48.2 +/- 0.8	22.4 +/- 0.2	4.4 +/- 0.1	63.8 +/- 5.2	NA
	Diabetic Patients	Total : 101	47.1 +/- 1.0	22.7 +/- 0.2	8.3 +/- 0.4	34.2 +/- 3.7	45.8
		Males : 35	49.6 +/- 1.6	23.1 +/- 0.3	8.3 +/- 0.6	26.6 +/- 4.1	56.6
		Females : 66	45.8 +/- 1.3	22.5 +/- 0.2	8.0 +/- 0.4	38.2 +/- 5.2	40.1
	p Value (Controls vs Diabetics)	Total	2.5E-01	1.9E-01	4.1E-39***	9.9E-06***	NA
		Males	7.6E-01	1.8E-01	5.0E-14***	6.9E-05***	NA
		Females	1.1E-01	5.4E-01	2.4E-26***	3.4E-03**	NA

Note: There was no significant difference in IAP levels between high- and low-BMI groups of healthy controls as well as no significant difference between high- and low-BMI groups of diabetic patients. Obese people with high stool IAP do not develop T2DM. Statistics: Values are expressed as mean +/- SEM. Statistical significance of the difference between two groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. NA, not applicable.

Figure 5 is a graph showing the distributions of IAP in diabetes patients and healthy control subjects with high and low body mass index (BMI) as described in Table 7.

Figure 5: High levels of IAP is protective against T2DM irrespective of body mass index (BMI). Stool samples of healthy participants and T2DM patients were assayed for IAP concentrations using an automatic biochemistry analyzer (see Figure 3). The healthy controls as well as T2DM patients were categorized in two groups, one with high BMI (> 25.0 kg/m²) and the other having low BMI (< 25.0 kg/m²). Statistics: Values are expressed as mean +/- SEM. Statistical significance of the difference between two groups was tested using the unpaired two-tailed Student's t-Test. $p < 0.05$ is considered significant. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. Note: There was no significant difference in IAP levels between high- and low-BMI groups of healthy controls as well as no significant difference in IAP levels

between high- and low-BMI groups of diabetic patients. Obese people with high stool IAP have no T2DM.

Example 7: The majority of population has the incipient metabolic syndrome.

IAP deficiency causes the metabolic syndrome in mice (Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8). Thus we predict that the association of IAP deficiency with T2DM in humans would be useful to isolate the people with 'the incipient metabolic syndrome' who are apparently healthy persons, however, vulnerable to develop the metabolic syndrome in the near future due to IAP deficiency. Victims of 'the incipient metabolic syndrome' should include the persons with 'incipient' diabetes, heart disease, nonalcoholic fatty liver, hypertension, and other metabolic disorders. We observed that the average IAP level in healthy population is approx. 67.4 +/- 3.2 U/g stool (Table 1), and hence we define that a healthy person having IAP level less than 65.0 U/g stool, using the current assay condition, should be considered as having 'the incipient metabolic syndrome' that should include 'incipient diabetes' and 'the pre-metabolic syndrome (exhibiting at least one criterion of the metabolic syndrome)'. It is noteworthy that the numerical value of concentration of 65.0 U phosphatase per gm stool would vary depending on different alkaline phosphatase (AP) assays as well as on assay conditions, for example change of pH, time, temperature, substrate concentration, etc.

Table 8 shows that approx. 65% of the healthy population have 'the incipient metabolic syndrome' having IAP level < 65.0 U/g stool. This prevalence rate is in concordance with the 71% prevalence rate of the pre-metabolic syndrome (exhibiting at least one criterion of the metabolic syndrome) (Al-Qahtani DA, Imtiaz ML. Prevalence of metabolic syndrome in Saudi adult soldiers. Saudi Med J 2005;26(9):1360-6). Also, it was reported that 68-81% of Pakistani people have low levels of HDL-cholesterol, one criterion of the metabolic syndrome (Basit A, Shera AS. 2008. Prevalence of metabolic syndrome in Pakistan. Metab Syndr Relat Disord. 6(3):171-5). We calculated that approx. 40% of healthy subjects have amounts of IAP that are less than the average IAP value of the diabetic group (35.0 U/g stool). It is predicted that these people have 'the severe incipient metabolic syndrome' and are extremely vulnerable to develop T2DM or other metabolic disorders within a few years. Interestingly, the prevalence rate of 'overt' metabolic syndrome (co-existence of 3 criteria of the metabolic syndrome; Huang PL. A comprehensive definition for

metabolic syndrome. Dis Model Mech 2009;2(5-6):231-7) is also 40% (Kaliannan et al. Proc Natl Acad Sci U S A 2013;110(17):7003-8; Ford. Diabetes Care. 2005 Nov;28(11):2745-9).

Also, a few T2DM patients (approx. 15%) have stool IAP level > 67.0 U/g (the average value of healthy controls), and this observation means that either IAP is not associated with the pathogenesis of T2DM in these specific patients or a persistent loss of IAP from a previously high level might also lead to T2DM. A loss of approx. 50% IAP activity might be significant to precipitate T2DM as diabetic patients on average have approx. 50% less IAP compared to their healthy counterparts (see Table 1 and Figure 3). Thus it appears that a distinct population of incipient diabetics might also have higher IAP levels.

An incipient diabetic person is an apparently healthy person having a stool IAP level less than 65.0 U/gm stool using the assay conditions disclosed herein or previously having a higher IAP level but presently experiencing persistent loss of 50% IAP activity.

Table 8. The majority of population has the incipient metabolic syndrome.

Group	Characteristics of Participants	
Healthy Controls	Average IAP level (U/g stool)	67.4 +/- 3.2
	Total number of Healthy Controls	445
	Number of Controls with IAP levels < 65.0 U/g stool (Incipient Metabolic Syndrome)	299
	Number of Controls with IAP levels > 65.0 U/g stool	146
	Percentage (%) of Incipient Metabolic Syndrome (299/445*100)	67.2

Note: Based on the average IAP value of 67.4 +/- 3.2 U/g stool in healthy controls, it is defined that a healthy person with 'the incipient metabolic syndrome' has stool IAP levels < 65.0 U/g stool. The incipient metabolic syndrome should also include 'incipient diabetes'.

What is claimed is:

CLAIMS

1. A method for developing a temporal profile of stool phosphatase comprising measuring concentration of phosphatase in a stool sample from a subject, wherein said stool sample is collected at a single time point or at multiple time points.

5

2. The method of claim 1, wherein said multiple time points comprise daily, weekly, monthly or yearly.

3. The method of claim 1 or 2, wherein said subject is a human, cattle, pig, sheep, goat, cow,
10 horse, dog, cat, monkey, rabbit, rat, mouse, chicken, or turkey, preferably human.

4. The method of any one of claims 1-3, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;

contacting said substrate for phosphatase with a stool sample;

15

waiting a specified period of time to allow a color to develop;

comparing the developed color with photographs of standards; and

quantifying the concentration of phosphatase in said stool sample.

5. The method of any one of claims 1-3, wherein said measuring comprises:

20

providing a substrate for phosphatase in a solid or liquid form;

contacting said substrate for phosphatase with a stool sample;

waiting a specified period of time to allow a color to develop;

counting the pixels of photographs;

comparing the number of pixels to standards; and

25

quantifying the concentration of phosphatase in said stool sample.

6. The method of any one of claims 1-3, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;

contacting said substrate for phosphatase with a stool sample;

30

waiting a specified period of time to allow an enzymatic reaction to occur; and

quantifying the concentration of phosphatase in said stool sample using a spectrophotometer, a biochemistry analyzer, a high-performance liquid chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical, biochemical, electrical or electronic device.

5

7. The method of any one of claims 1-3, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a stool sample;
waiting a specified period of time to allow a color to develop;
10 comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said stool sample.

8. The method of any one of claims 1-3, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
15 contacting said solid medium with a stool sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said stool sample.

20

9. The method of any one of claims 1-3, wherein said measuring comprises:

providing a solid medium for attaching a stool sample thereto;
attaching said stool sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
25 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said stool sample.

10. The method of any one of claims 1-3, wherein said measuring comprises:

30 providing a solid medium for attaching a stool sample thereto;
attaching said stool sample to said solid medium;

contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
5 quantifying the concentration of phosphatase in said stool sample.

11. The method of any one of claims 7-10, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

12. The method of any one of claims 4-11, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl
15 phosphate disodium salt) or combinations thereof.

13. The method of any one of claims 1-12, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline
20 phosphatase, neutrophil alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

14. The method of any one of claims 1-13, wherein said stool sample is substituted with a biological sample comprising saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric
25 fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid.

15. A method for diagnosing incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising
30 measuring concentration of phosphatase in a biological sample from a subject and comparing the concentration to an average normal value of concentration.

16. The method of claim 15, wherein a concentration of phosphatase that is 1% - 100% less than an average normal value indicates incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

5 17. The method of claim 15 or 16, further comprising developing a temporal phosphatase profile comprising measuring the concentration of phosphatase in one or more than one biological sample from a subject, wherein said one or more than one biological sample is collected at a single time point or at multiple time points.

10 18. The method of claim 17, wherein said multiple time points comprise daily, weekly, monthly or yearly.

19. The method of claim 17 or 18, wherein the temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time indicates incipient diabetes,
15 incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

20. The method of claim 19, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

20

21. The method of any one of claims 15-20, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

25 22. The method of any one of claims 15-21, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
30 quantifying the concentration of phosphatase in said biological sample.

23. The method of any one of claims 15-21, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
5 counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

24. The method of any one of claims 15-21, wherein said measuring comprises:

10 providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
spectrophotometer, a biochemistry analyzer, a high-performance liquid
15 chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
biochemical, electrical or electronic device.

25. The method of any one of claims 15-21, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
20 contacting said solid medium with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

26. The method of any one of claims 15-21, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
25 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

27. The method of any one of claims 15-21, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
5 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

28. The method of any one of claims 15-21, wherein said measuring comprises:

10 providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
15 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

29. The method of any one of claims 25-28, wherein said solid medium comprises a
membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab,
20 preferably nylon or nitrocellulose membrane.

30. The method of any one of claims 22-28, wherein said substrate for phosphatase comprises
p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium
substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-
25 methoxyspiro{1,2-dioxetane-3,2'-(5'-chloro)-tricyclo [3.3.1.1^{3.7}]decan}-4-yl)-1-phenyl
phosphate disodium salt) or combinations thereof..

31. The method of any one of claims 15-30, wherein said phosphatase comprises an alkaline
phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific
30 tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline

phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

32. The method of any one of claims 15-30, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

33. The method of any one of claims 15-32, further comprising measuring concentration of one or more of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

34. The method of any one of claims 15-33, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

35. The method of any one of claims 15-33, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

36. The method of any one of claims 15-33, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

37. Use of a substrate of phosphatase for developing a temporal phosphatase profile comprising measuring the concentration of phosphatase in one or more than one biological

samples from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

38. The use of claim 37, wherein said multiple time points comprise daily, weekly, monthly
5 or yearly.

39. The use of claim 37 or 38, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

10 40. The use of any one of claims 37-39, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
15 quantifying the concentration of phosphatase in said biological sample.

41. The use of any one of claims 37-39, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
20 waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

25 42. The use of any one of claims 37-39, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
30 spectrophotometer, a biochemistry analyzer, a high-performance liquid

chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical, biochemical, electrical or electronic device.

43. The use of any one of claims 37-39, wherein said measuring comprises:

5 providing a solid medium having a substrate for phosphatase attached thereto;
 contacting said solid medium with a biological sample;
 waiting a specified period of time to allow a color to develop;
 comparing the developed color with photographs of standards; and
 quantifying the concentration of phosphatase in said biological sample.

10

44. The use of any one of claims 37-39, wherein said measuring comprises:

 providing a solid medium having a substrate for phosphatase attached thereto;
 contacting said solid medium with a biological sample;
 waiting a specified period of time to allow a pixilated color pattern to develop;
15 counting the pixels of photographs;
 comparing the number of pixels to standards; and
 quantifying the concentration of phosphatase in said biological sample.

15

45. The use of any one of claims 37-39, wherein said measuring comprises:

20 providing a solid medium for attaching a biological sample thereto;
 attaching said biological sample to said solid medium;
 contacting said solid medium with a substrate for phosphatase;
 waiting a specified period of time to allow a color to develop;
 comparing the developed color with photographs of standards; and
25 quantifying the concentration of phosphatase in said biological sample.

20

46. The use of any one of claims 37-39, wherein said measuring comprises:

 providing a solid medium for attaching a biological sample thereto;
 attaching said biological sample to said solid medium;
30 contacting said solid medium with a substrate for phosphatase;
 waiting a specified period of time to allow a pixilated color pattern to develop;

30

counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

5 47. The use of any one of claims 43-46, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

10 48. The use of any one of claims 40-46, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

15 49. The use of any one of claims 37-48, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

20 50. The use of any one of claims 37-49, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

25 51. Use of a substrate of phosphatase for detecting incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising measuring the concentration of phosphatase in a biological sample from a subject and comparing the concentration to an average normal value of concentration.

30

52. The use of claim 51, wherein a concentration of phosphatase that is 1% - 100% less than an average normal value indicates incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

5 53. The use of claim 51 or 52, further comprising developing a temporal phosphatase profile comprising measuring the concentration of phosphatase in one or more than one biological sample from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

10 54. The use of claim 53, wherein said multiple time points comprise daily, weekly, monthly or yearly.

55. The use of claim 53 or 54, wherein the temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time indicates incipient diabetes,
15 incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

56. The use of claim 55, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

20

57. The use of any one of claims 51-56, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

58. The use of any one of claims 51-57, wherein said measuring comprises:

25 providing a substrate for phosphatase in a solid or liquid form;
 contacting said substrate for phosphatase with a biological sample;
 waiting a specified period of time to allow a color to develop;
 comparing the developed color with photographs of standards; and
 quantifying the concentration of phosphatase in said biological sample.

30

59. The use of any one of claims 51-57, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
5 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

60. The use of any one of claims 51-57, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
10 contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
spectrophotometer, a biochemistry analyzer, a high-performance liquid
chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
15 biochemical, electrical or electronic device.

61. The use of any one of claims 51-57, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
20 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

62. The use of any one of claims 51-57, wherein said measuring comprises:

25 providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
30 quantifying the concentration of phosphatase in said biological sample.

63. The use of any one of claims 51-57, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
5 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

64. The use of any one of claims 51-57, wherein said measuring comprises:

10 providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
15 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

65. The use of any one of claims 61-64, wherein said solid medium comprises a membrane,
paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or
20 nitrocellulose membrane.

66. The use of any one of claims 51-65, wherein said substrate for phosphatase comprises p-
nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate
system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-
25 methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl
phosphate disodium salt) or combinations thereof.

67. The use of any one of claims 51-66, wherein said phosphatase comprises an alkaline
phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific
30 tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline

phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

68. The use of any one of claims 51-67, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular
5 fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

69. The use of any one of claims 51-68, further comprising measuring concentration of one or more of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines,
10 antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

70. The use of any one of claims 51-69, wherein said incipient diabetes or diabetes comprises
15 type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

71. The use of any one of claims 51-69, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy,
20 asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

72. The use of any one of claims 51-69, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia,
25 dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

73. A method of decreasing the risk of or a method of treating incipient diabetes, incipient
30 metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic

syndrome comprising administering a pharmaceutically effective amount of a phosphatase to a subject in need thereof.

74. The method of claim 73, wherein said subject in need thereof has a concentration of phosphatase that is 1% - 100% less than an average normal value.

75. The method of claim 73, wherein said subject in need thereof has a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

76. The method of claim 75, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

77. The method of any one of claims 73-76, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

78. The method of any one of claims 73-77, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

79. The method of any one of claims 73-78, wherein said phosphatase comprises a native intestinal alkaline phosphatase, a recombinant intestinal alkaline phosphatase, a mutant intestinal alkaline phosphatase, a peptide with intestinal alkaline phosphatase activity, a recombinant peptide with intestinal alkaline phosphatase activity, native human intestinal alkaline phosphatase, recombinant human intestinal alkaline phosphatase, native calf intestinal alkaline phosphatase or recombinant calf intestinal alkaline phosphatase.

80. The method of any one of claims 73-79, wherein said phosphatase is administered orally, topically, subcutaneously, intradermally, intramuscularly, intravenously, vaginally or rectally.

81. The method of any one of claims 73-80, wherein said phosphatase is administered before,
5 during or after a meal.

82. The method of any one of claims 73-81, wherein said phosphatase is administered in an amount effective to maintain a steady-state level of phosphatase within a normal range of 65-650 U/gram stool.

10 83. The method of any one of claims 73-82, wherein said phosphatase is administered in an amount effective to maintain a steady-state level between 65 and 650 units of phosphatase per gram of stool.

15 84. The method of any one of claims 73-83, wherein said phosphatase is administered in an amount effective to maintain a steady-state level of about 65 units or more of phosphatase per gram of stool.

85. The method of any one of claims 73-84, wherein said phosphatase is administered in an amount of 1 to 6,500 units of phosphatase per dose.

20 86. The method of any one of claims 73-85, wherein said phosphatase is dispensed in a form of a tablet, gel, capsule, liquid or semi-solid chewing gum.

87. The method of any one of claims 73-86, wherein said phosphatase is dispensed in a food
25 or beverage, wherein said beverage is preferably milk.

88. The method of any one of claims 73-87, wherein said phosphatase is dispensed with one or more medications comprising anti-diabetes medications, anti-hypertension medications, anti-cholesterol medications, anti-depression medications, antibiotics, probiotics, sodium
30 butyrate, thyroid hormone, curcumin, omega-3 fatty acid, beta-lactamase, a protein, a peptide or combinations thereof.

89. The method of any one of claims 73-88, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

90. The method of any one of claims 73-88, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

91. The method of any one of claims 73-88, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

92. Use of a phosphatase to decrease the risk of or to treat incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

93. The use of claim 92, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

94. The use of claim 92, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

95. The use of claim 94, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

96. The use of any one of claims 93-95, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

97. The use of any one of claims 92-96, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

98. The use of any one of claims 92-97, wherein said phosphatase comprises a native intestinal alkaline phosphatase, a recombinant intestinal alkaline phosphatase, a mutant intestinal alkaline phosphatase, a peptide with intestinal alkaline phosphatase activity, a recombinant peptide with intestinal alkaline phosphatase activity, native human intestinal alkaline phosphatase, recombinant human intestinal alkaline phosphatase, native calf intestinal alkaline phosphatase or recombinant calf intestinal alkaline phosphatase.

99. The use of any one of claims 92-98, wherein said phosphatase is for oral, topical, subcutaneous, intradermal, intramuscular, intravenous, vaginal or rectal administration.

100. The use of claim 99, wherein said administration is before, during or after a meal.

101. The use of claim 99 or 100, wherein said administration is in an amount effective to maintain a steady-state level of phosphatase within a normal range of 65-650 U/gram stool.

102. The use of any one of claims 99-101, wherein said administration is in an amount effective to maintain a steady-state level between 65 and 650 units of phosphatase per gram of stool.

103. The use of any one of claims 99-102, wherein said administration is in an amount effective to maintain a steady-state level of about 65 units or more of phosphatase per gram of stool.

5 104. The use of any one of claims 99-103, wherein said administration is in an amount of 1 to 6,500 units of phosphatase per dose.

105. The use of any one of claims 92-104, wherein said phosphatase is dispensed in a form of a tablet, gel, capsule, liquid or semi-solid chewing gum.

10

106. The use of any one of claims 92-105, wherein said phosphatase is dispensed in a food or beverage, wherein said beverage is preferably milk.

107. The use of any one of claims 92-106, wherein said phosphatase is dispensed with one or
15 more medications comprising anti-diabetes medications, anti-hypertension medications, anti-cholesterol medications, anti-depression medications, antibiotics, probiotics, sodium butyrate, thyroid hormone, curcumin, omega-3 fatty acid, beta-lactamase, a protein, a peptide or combinations thereof.

20 108. The use of any one of claims 92-107, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

109. The use of any one of claims 92-107, wherein said incipient metabolic disease or
25 metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

110. The use of any one of claims 92-107, wherein said incipient metabolic syndrome or
30 metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia,

dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

111. A pharmaceutical composition comprising phosphatase for use in decreasing the risk of or for use in treating incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

112. The pharmaceutical composition of claim 111, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

113. The pharmaceutical composition of claim 111, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

114. The pharmaceutical composition of claim 113, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

115. The pharmaceutical composition of any one of claims 112-114, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

116. The pharmaceutical composition of any one of claims 111-115, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

117. The pharmaceutical composition of any one of claims 111-116, wherein said phosphatase comprises a native intestinal alkaline phosphatase, a recombinant intestinal alkaline phosphatase, a mutant intestinal alkaline phosphatase, a peptide with intestinal alkaline phosphatase activity, a recombinant peptide with intestinal alkaline phosphatase activity, native human intestinal alkaline phosphatase, recombinant human intestinal alkaline phosphatase, native calf intestinal alkaline phosphatase or recombinant calf intestinal alkaline phosphatase.

118. The pharmaceutical composition of any one of claims 111-117, wherein said phosphatase is for oral, topical, subcutaneous, intradermal, intramuscular, intravenous, vaginal or rectal administration.

119. The pharmaceutical composition of claim 118, wherein said administration is before, during or after a meal.

120. The pharmaceutical composition of claim 118 or 119, wherein said administration is in an amount effective to maintain a steady-state level of phosphatase within a normal range of 65-650 U/gram stool.

121. The pharmaceutical composition of any one of claims 118-120, wherein said administration is in an amount effective to maintain a steady-state level between 65 and 650 units of phosphatase per gram of stool.

122. The pharmaceutical composition of any one of claims 118-121, wherein said administration is in an amount effective to maintain a steady-state level of about 65 units or more of phosphatase per gram of stool.

123. The pharmaceutical composition of any one of claims 118-122, wherein said administration is in an amount of 1 to 6,500 units of phosphatase per dose.

124. The pharmaceutical composition of any one of claims 121-123, wherein said phosphatase is dispensed in a form of a tablet, gel, capsule, liquid or semi-solid chewing gum.

125. The pharmaceutical composition of any one of claims 121-124, wherein said
5 phosphatase is dispensed in a food or beverage, wherein said beverage is preferably milk.

126. The pharmaceutical composition of any one of claims 121-125, wherein said phosphatase is dispensed with one or more medications comprising anti-diabetes medications, anti-hypertension medications, anti-cholesterol medications, anti-depression medications,
10 antibiotics, probiotics, sodium butyrate, thyroid hormone, curcumin, omega-3 fatty acid, beta-lactamase, a protein, a peptide or combinations thereof.

127. The pharmaceutical composition of any one of claims 121-126, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes
15 insipidus.

128. The pharmaceutical composition of any one of claims 121-126, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver
20 disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

129. The pharmaceutical composition of any one of claims 121-126, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin
25 resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

130. Use of a phosphatase in the preparation of a medicament for decreasing the risk of or for
30 the treatment of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

131. The use of claim 130, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

5

132. The use of claim 130, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

10

133. The use of claim 132, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

15

134. The use of any one of claims 131-133, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

20

135. The use of claim 130-134, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, a mammalian alkaline phosphatase, a bacterial alkaline phosphatase, a fungal alkaline phosphatase, an acid phosphatase or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

25

136. The use of any one of claims 130-135, wherein said phosphatase comprises a native intestinal alkaline phosphatase, a recombinant intestinal alkaline phosphatase, a mutant intestinal alkaline phosphatase, a peptide with intestinal alkaline phosphatase activity, a recombinant peptide with intestinal alkaline phosphatase activity, native human intestinal alkaline phosphatase, recombinant human intestinal alkaline phosphatase, native calf

30

intestinal alkaline phosphatase or recombinant calf intestinal alkaline phosphatase.

137. The use of any one of claims 130-136, wherein said phosphatase is for oral, topical, subcutaneous, intradermal, intramuscular, intravenous, vaginal or rectal administration.

138. The use of claim 137, wherein said administration is before, during or after a meal.

5

139. The use of claim 137 or 138, wherein said administration is in an amount effective to maintain a steady-state level of phosphatase within a normal range of 65-650 U/gram stool.

140. The use of any one of claims 137-139, wherein said administration is in an amount effective to maintain a steady-state level between 65 and 650 units of phosphatase per gram of stool.

10

141. The use of any one of claims 137-140, wherein said administration is in an amount effective to maintain a steady-state level of about 65 units or more of phosphatase per gram of stool.

15

142. The use of any one of claims 137-141, wherein said administration is in an amount of 1 to 6,500 units of phosphatase per dose.

143. The use of any one of claims 130-142, wherein said phosphatase is dispensed in a form of a tablet, gel, capsule, liquid or semi-solid chewing gum.

20

144. The use of any one of claims 130-143, wherein said phosphatase is dispensed in a food or beverage, wherein said beverage is preferably milk.

25

145. The use of any one of claims 130-144, wherein said phosphatase is dispensed with one or more medications comprising anti-diabetes medications, anti-hypertension medications, anti-cholesterol medications, anti-depression medications, antibiotics, probiotics, sodium butyrate, thyroid hormone, curcumin, omega-3 fatty acid, beta-lactamase, a protein, a peptide or combinations thereof.

30

146. The use of any one of claims 130-145, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

147. The use of any one of claims 130-145, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

148. The use of any one of claims 130-145, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

149. A kit for determining the concentration of phosphatase in stool wherein the kit comprises a tool for measuring a defined quantity of stool, a stool dilution buffer or water, a substrate for phosphatase and a phosphatase assay buffer.

150. The kit of claim 149, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

151. The kit of claim 149 or 150 wherein said substrate for phosphatase is provided in solid or liquid form or attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably paper or a tube.

152. The kit of any one of claims 149-151, wherein said stool sample is attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

153. The kit of any one of claims 149-152, wherein the kit comprises a tool for measuring a defined quantity of stool, a bottle or a tube fitted with a filtering device, and a tube containing a substrate for phosphatase mixed with phosphatase assay buffer.

5 154. The kit of any one of claims 149-153, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

10

155. The kit of any one of claims 149-154, wherein stool is substituted with a biological sample comprising saliva, urine, blood, plasma, serum, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid or a body cavity fluid.

15 156. Use of a kit for determining the concentration of phosphatase in a stool sample, wherein the kit comprises a tool for measuring a defined quantity of stool, a stool sample dilution buffer, a substrate for phosphatase and a phosphatase assay buffer.

20 157. The use of claim 156, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo[3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

25 158. The use of claim 156 or 157, wherein said substrate for phosphatase is provided in solid or liquid form or attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably paper or a tube.

30 159. The use of any one of claims 156-158, wherein said stool sample is attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

160. The use of any one of claims 156-159, wherein said kit comprises a tool for measuring a defined quantity of stool, a bottle or a tube fitted with a filtering device, and a tube containing a substrate for phosphatase mixed with phosphatase assay buffer.

5 161. The use of any one of claims 156-160, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

10

162. The use of any one of claims 156-161, wherein said stool is substituted with a biological sample comprising saliva, urine, blood, plasma, serum, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid or a body cavity fluid.

15 163. A kit for detecting incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome, wherein the kit comprises a tool for measuring a defined quantity of stool, a stool dilution buffer, a substrate for phosphatase and a phosphatase assay buffer.

20 164. The kit of claim 163, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

25 165. The kit of claim 163, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

30 166. The kit of claim 165, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

167. The kit of any one of claims 163-166, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

5 168. The kit of any one of claims 163-167, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

10

169. The kit of any one of claims 163-168, wherein said substrate for phosphatase is provided in solid or liquid form or attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably paper or a tube.

15

170. The kit of any one of claims 163-169, wherein the stool sample is attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

20

171. The kit of any one of claims 163-170, wherein the kit comprises a tool for measuring a defined quantity of stool, a bottle or a tube fitted with a filtering device, and a tube containing a substrate for phosphatase mixed with phosphatase assay buffer.

25

172. The kit of any one of claims 163-171, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

30

173. The kit of any one of claims 163-172, wherein said subject provides a biological sample comprising stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal

fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

174. The kit of any one of claims 163-173, wherein said kit further comprises a means for measuring concentration of one or more of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

175. The kit of any one of claims 163-174, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

176. The kit of any one of claims 163-174, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

177. The kit of any one of claims 163-174, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

178. Use of a kit for detecting incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome, wherein the kit comprises a tool for measuring a defined quantity of stool, a stool dilution buffer, a substrate for phosphatase and a phosphatase assay buffer.

179. The use of claim 178, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is

characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

180. The use of claim 178, wherein said incipient diabetes, incipient metabolic disease,
5 incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

181. The use of claim 180, wherein said persistent decrease is about a 50% decrease from a
10 previously measured concentration.

182. The use of any one of claims 179-181, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken, turkey, preferably human.

183. The use of any one of claims 178-182, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium
15 substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl
20 phosphate disodium salt) or combinations thereof.

184. The use of any one of claims 178-183, wherein said substrate for phosphatase is provided in solid or liquid form or attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably paper or a
25 tube.

185. The use of any one of claims 178-184, wherein the stool sample is attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

186. The use of any one of claims 178-185, wherein the kit comprises a tool for measuring a defined quantity of stool, a bottle or a tube fitted with a filtering device, and a tube containing a substrate for phosphatase mixed with phosphatase assay buffer.

5 187. The use of any one of claims 178-186, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

10

188. The use of any one of claims 178-187, wherein said subject provides a biological sample comprising stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

15

189. The use of any one of claims 178-188, wherein said kit further comprises a means for measuring concentration of one or more of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample from a subject, wherein said one or more than one

20

biological samples is collected at a single time point or at multiple time points.

190. The use of any one of claims 178-189, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

25

191. The use of any one of claims 178-189, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

30

192. The use of any one of claims 178-189, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

5

193. A method for determining prognosis of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising developing a temporal phosphatase profile comprising measuring the concentration of phosphatase in one or more than one biological sample from a subject, wherein said one or
10 more than one biological samples is collected at a single time point or at multiple time points.

194. The method of claim 193, wherein said multiple time points comprise daily, weekly, monthly or yearly.

15

195. The method of claim 193 or 194, wherein the temporal phosphatase profile comprising an increase in phosphatase concentration over time indicates a good prognosis for lessening the symptoms of or decreasing the risk of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome and a decrease in phosphatase concentration over time indicates a poor prognosis for lessening the
20 symptoms of or decreasing the risk of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

196. The method of claim 195, wherein said increase or decrease is about a 50% change from a previously measured concentration.

25

197. The method of claim 195, wherein said increase is to a concentration of phosphatase that is about within a normal range of 65-650 U/gram stool.

198. The method of claim 195, wherein said decrease is to a concentration of phosphatase
30 that is 1% - 100% less than an average normal value.

199. The method of any one of claims 193-198, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

5 200. The method of any one of claims 193-199, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
10 quantifying the concentration of phosphatase in said biological sample.

201. The method of any one of claims 193-199, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
15 waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

20 202. The method of any one of claims 193-199, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
25 spectrophotometer, a biochemistry analyzer, a high-performance liquid
chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
biochemical, electrical or electronic device.

203. The method of any one of claims 193-199, wherein said measuring comprises:
30 providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;

waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

- 5 204. The method of any one of claims 193-199, wherein said measuring comprises:
providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
10 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

205. The method of any one of claims 193-199, wherein said measuring comprises:
providing a solid medium for attaching a biological sample thereto;
15 attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

- 20 206. The method of any one of claims 193-199, wherein said measuring comprises:
providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
25 waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

207. The method of any one of claims 203-206, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

208. The method of any one of claims 200-207, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

209. The method of any one of claims 193-208, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

210. The method of any one of claims 193-209, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

211. The method of any one of claims 193-210, further comprising measuring the concentration of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

212. The method of any one of claims 193-211, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

213. The method of any one of claims 193-211, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

214. The method of any one of claims 193-211, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

215. Use of a substrate of phosphatase for determining prognosis of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising developing a temporal phosphatase profile comprising measuring the concentration of phosphatase in one or more than one biological samples from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

216. The use of claim 215, wherein said multiple time points comprise daily, weekly, monthly or yearly.

217. The use of claim 215 or 216, wherein the temporal phosphatase profile comprising an increase in phosphatase concentration over time indicates a good prognosis for lessening the symptoms of or decreasing the risk of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome and a decrease in phosphatase concentration over time indicates a poor prognosis for lessening the symptoms of or decreasing the risk of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

218. The use of claim 217, wherein said increase or decrease is about a 50% change from a previously measured concentration.

219. The use of claim 217, wherein said increase is to a concentration of phosphatase that is about within a normal range of 65-650 U/gram stool.

220. The use of claim 217, wherein said decrease is to a concentration of phosphatase that is 1% - 100% less than an average normal value.

221. The use of any one of claims 215-220, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

222. The use of any one of claims 215-221, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

223. The use of any one of claims 215-221, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

224. The use of any one of claims 215-221, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;

waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
spectrophotometer, a biochemistry analyzer, a high-performance liquid
chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
5 biochemical, electrical or electronic device.

225. The use of any one of claims 215-221, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
10 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

226. The use of any one of claims 215-221, wherein said measuring comprises:

15 providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
20 quantifying the concentration of phosphatase in said biological sample.

227. The use of any one of claims 215-221, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
25 contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

30 228. The use of any one of claims 215-221, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;

attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
5 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

229. The use of claims 225-228, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or
10 nitrocellulose membrane.

230. The use of any one of claims 215-229, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-
15 methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

231. The use of any one of claims 215-230, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific
20 tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

232. The use of any one of claims 215-231, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular
25 fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

233. The use of any one of claims 215-232, further comprising measuring the concentration
30 of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample

from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

234. The use of any one of claims 215-233, wherein said incipient diabetes or diabetes
5 comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

235. The use of any one of claims 215-233, wherein said incipient metabolic disease or
metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular
disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy,
10 asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis,
eczema or psoriasis.

236. The use of any one of claims 215-233, wherein said incipient metabolic syndrome or
metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia,
15 dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein
(HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

237. Use of a temporal phosphatase profile for detecting incipient diabetes, incipient
metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic
20 syndrome comprising measuring the concentration of phosphatase in one or more than one
biological samples from a subject, wherein said one or more than one biological sample is
collected at a single time point or at multiple time points.

238. The use of claim 237, wherein said multiple time points comprise daily, weekly,
25 monthly or yearly.

239. The use of claim 237 or 238, wherein a concentration of phosphatase that is 1% - 100%
less than an average normal value indicates incipient diabetes, incipient metabolic disease,
incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

240. The use of claim 237 or 238, wherein the temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time indicates incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome.

5

241. The use of claim 240, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

242. The use of any one of claims 237-241, wherein said subject is a human, cattle, pig,
10 sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

243. The use of any one of claims 237-242, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
15 contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

20 244. The use of any one of claims 237-242, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
25 comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

245. The use of any one of claims 237-242, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
30 contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow an enzymatic reaction to occur; and

quantifying the concentration of phosphatase in said biological sample using a spectrophotometer, a biochemistry analyzer, a high-performance liquid chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical, biochemical, electrical or electronic device.

5

246. The use of any one of claims 237-242, wherein said measuring comprises:
providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a color to develop;
10 comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

247. The use of any one of claims 237-242, wherein said measuring comprises:
providing a solid medium having a substrate for phosphatase attached thereto;
15 contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

20

248. The use of any one of claims 237-242, wherein said measuring comprises:
providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
25 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

249. The use of any one of claims 237-242, wherein said measuring comprises:
30 providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;

contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
5 quantifying the concentration of phosphatase in said biological sample.

250. The use of claims 246-249, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

251. The use of any one of claims 243-250, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo[3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl
15 phosphate disodium salt) or combinations thereof.

252. The use of any one of claims 237-251, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline
20 phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

253. The use of any one of claims 237-252, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular
25 fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

254. The use of any one of claims 237-253, further comprising measuring the concentration of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of
30 cytokine receptors or anti-inflammatory factors in one or more than one biological sample

from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

255. The use of any one of claims 237-254, wherein said incipient diabetes or diabetes
5 comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

256. The use of any one of claims 237-254, wherein said incipient metabolic disease or
metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular
disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy,
10 asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis,
eczema or psoriasis.

257. The use of any one of claims 237-254, wherein said incipient metabolic syndrome or
metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia,
15 dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein
(HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

258. A method for diagnosing or for determining prognosis of a chronic disease, an acute
disease, an infectious disease, a disorder or a syndrome comprising developing a temporal
20 phosphatase profile comprising measuring the concentration of phosphatase in one or more
than one biological sample from a subject, wherein said one or more than one biological
samples is collected at a single time point or at multiple time points.

259. The method of claim 258, wherein said multiple time points comprise daily, weekly,
25 monthly or yearly.

260. The method of claim 258 or 259, wherein said subject is a human, cattle, pig, sheep,
goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

30 261. The method of any one of claims 258-260, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;

contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

5

262. The method of any one of claims 258-260, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
10 counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

263. The method of any one of claims 258-260, wherein said measuring comprises:

15 providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
spectrophotometer, a biochemistry analyzer, a high-performance liquid
20 chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
biochemical, electrical or electronic device.

264. The method of any one of claims 258-260, wherein said measuring comprises:

25 providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

30 265. The method of any one of claims 258-260, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;

contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
5 quantifying the concentration of phosphatase in said biological sample.

266. The method of any one of claims 258-260, wherein said measuring comprises:
providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
10 contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

15 267. The method of any one of claims 258-260, wherein said measuring comprises:
providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
20 counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

268. The method of claims 264-267, wherein said solid medium comprises a membrane,
25 paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or
nitrocellulose membrane.

269. The method of any one of claims 261-268, wherein said substrate for phosphatase
comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue
30 tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-

chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof..

270. The method of any one of claims 258-269, wherein said phosphatase comprises an
 5 alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.
- 10 271. The method of any one of claims 258-270, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.
- 15 272. The method of any one of claims 258-271, wherein said chronic disease comprises Acne, Acquired Immunodeficiency Syndrome (AIDS), Ankylosing spondylitis, Asthma, Addison's disease, Allergic rhinitis, Alzheimer's Disease, Amyotrophic lateral sclerosis (ALS, Lou Gehrig's Disease), Aplastic anemia, Arthritis, Atherosclerosis, Attention Deficit Hyperactivity Disorder (ADHD, ADD), Autism Spectrum Disorder, Benign Prostatic
 20 Hypertrophy, Bipolar Mood Disorder, Bronchiectasis, Cancer, Cardiac Failure, Cardiomyopathy, Chronic Alcoholism, Chronic anemia, Chronic hepatitis, Chronic Amebiasis, Chronic obstructive pulmonary disorder, Chronic gastritis, Chronic esophagitis, Chronic oral diseases, Chronic Poisoning, Chronic renal disease, Collagen diseases, Crohn's disease, Cushing's diseases, Cystic Fibrosis, Dementia, Dermatomyositis, Depression, Dysrhythmias,
 25 Eating Disorders (Anorexia Nervosa, Bulimia Nervosa, Binge Eating), Eczema, Endometriosis, Epilepsy, Fibrosing Alveolitis, Gastro-Esophageal Reflex Disease (GERD), Glaucoma, Gout, Graves Disease, Hemophilia, Hyperlipidemia, Hyperpituitarism, Hypopituitarism, Hypophyseal Adenoma, Hypothyroidism, Idiopathic Thrombocytopenic Purpura, Menopause, Migraine, Motor Neuron Disease, Multiple Sclerosis, Muscular
 30 Dystrophy, Myasthenia Gravis, Myopathy, Neuropathy, Obesity, Obsessive Compulsive Disorder, Osteoarthritis, Osteoporosis, Paralysis (Paraplegia, Quadriplegia), Parkinson's

disease, Paget's disease, Polyarteritis Nodosa, Polycystic ovarian disease, Psoriasis, Psoriatic Arthritis, Pulmonary Interstitial Fibrosis, Reflex Sympathetic Dystrophy (RSD) Syndrome, Rheumatoid Arthritis, Scleroderma, Schizophrenia, Sjögren's syndrome, Smoking and Tobacco-related Diseases, Systemic Lupus Erythematosus, Trigeminal Neuralgia,
 5 Tuberculosis, Ulcerative Colitis or Urinary Incontinence.

273. The method of any one of claims 258-271, wherein said acute disease comprises Acute Alcohol Poisoning, Acute Amebiasis, Acute Auditory Diseases, Acute Brain Diseases, Acute Cancers, Acute Cardiovascular Diseases, Acute Dental Diseases, Acute Dermatological
 10 Diseases, Acute Esophageal Diseases, Acute Gastrointestinal Diseases, Acute Kidney Diseases, Acute Liver Diseases, Acute Muscular Diseases, Acute Ocular Diseases, Acute Poisoning, Acute Oral Diseases, Acute Osteoarthritis, Acute Pharyngeal Diseases, Acute Prostatic Diseases, Acute Physical Injury (Trauma), Acute Peritoneal Diseases, Acute Respiratory Diseases or Acute Urogenital Diseases.

274. The method of any one of claims 258-271, wherein said infectious disease is caused by an organism comprising *Bacillus anthracis*, *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella abortus*, *Brucella canis*, *Brucella melitensis*, *Brucella suis*, *Campylobacter jejuni*, *Chlamydomphila pneumoniae*, *Chlamydia trachomatis*, *Chlamydomphila psittaci*, *Clostridium*
 20 *botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Entamoeba histolytica*, *Enterococcus faecalis*, *Enterococcus faecium*, *Escherichia coli*, *Enterotoxigenic Escherichia coli*, *Enteropathogenic E. coli*, *Enterohemorrhagic Escherichia coli* O157:H7, *Francisella tularensis*, Fungus (*Candida albicans*), *Haemophilus influenza*, *Helicobacter pylori*, *Leishmania* (Kala Azar), *Legionella*
 25 *pneumophila*, *Leptospira interrogans*, *Leptospira interrogans*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, Parasites (protozoa, helminths, ectoparasites), *Plasmodium* (malaria), *Pseudomonas aeruginosa*, *Rickettsia rickettsia*, *Salmonella enterica* subsp. *Enterica*, *Salmonella typhimurium*, *Shigella sonnei*, *Staphylococcus aureus*,
 30 *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Streptococcus agalactiae*,

Streptococcus pneumoniae, *Streptococcus pyogenes*, *Treponema pallidum*, *Vibrio cholerae* or *Yersinia pestis*.

275. The method of any one of claims 258-271, wherein said infectious disease is caused by an organism comprising Adenovirus, Coxsackievirus, Dengue virus, Ebola virus, Epstein-Barr virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Herpes simplex virus, Cytomegalovirus, Human immunodeficiency virus (HIV), Influenza virus, Measles virus, Mumps virus, Human papillomavirus, Human parainfluenza virus, Poliovirus, Rabies virus, Human respiratory syncytial virus, Rubella virus, Varicella zoster virus or Yellow fever virus.

276. Use of a substrate of phosphatase for detecting a chronic disease, an acute disease, an infectious disease, a disorder or a syndrome comprising developing a temporal phosphatase profile comprising measuring the concentration of phosphatase in one or more than one biological samples from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

277. The use of claim 276, wherein said multiple time points comprise daily, weekly, monthly or yearly.

278. The use of claim 276 or 277, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

279. The use of any one of claims 276-278, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

280. The use of any one of claims 276-278, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;

contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
5 quantifying the concentration of phosphatase in said biological sample.

281. The use of any one of claims 276-278, wherein said measuring comprises:
providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
10 waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
spectrophotometer, a biochemistry analyzer, a high-performance liquid
chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
biochemical, electrical or electronic device.

15 282. The use of any one of claims 276-278, wherein said measuring comprises:
providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a color to develop;
20 comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

283. The use of any one of claims 276-278, wherein said measuring comprises:
providing a solid medium having a substrate for phosphatase attached thereto;
25 contacting said solid medium with a biological sample;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

30 284. The use of any one of claims 276-278, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a color to develop;
5 comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

285. The use of any one of claims 276-278, wherein said measuring comprises:
providing a solid medium for attaching a biological sample thereto;
10 attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
15 quantifying the concentration of phosphatase in said biological sample.

286. The use of claims 282-285, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

287. The use of any one of claims 279-286, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl
25 phosphate disodium salt) or combinations thereof..

288. The use of any one of claims 276-287, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline
30 phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

289. The use of any one of claims 276-288, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

5

290. The use of any one of claims 276-289, wherein said chronic disease comprises Acne, Acquired Immunodeficiency Syndrome (AIDS), Ankylosing spondylitis, Asthma, Addison's disease, Allergic rhinitis, Alzheimer's Disease, Amyotrophic lateral sclerosis (ALS, Lou Gehrig's Disease), Aplastic anemia, Arthritis, Atherosclerosis, Attention Deficit Hyperactivity Disorder (ADHD, ADD), Autism Spectrum Disorder, Benign Prostatic Hypertrophy, Bipolar
10 Mood Disorder, Bronchiectasis, Cancer, Cardiac Failure, Cardiomyopathy, Chronic Alcoholism, Chronic anemia, Chronic hepatitis, Chronic Amebiasis Chronic obstructive pulmonary disorder, Chronic gastritis, Chronic esophagitis, Chronic oral diseases, Chronic Poisoning, Chronic renal disease, Collagen diseases, Crohn's disease, Cushing's diseases, Cystic Fibrosis, Dementia, Dermatomyositis, Depression, Dysrhythmias, Eating Disorders
15 (Anorexia Nervosa, Bulimia Nervosa, Binge Eating), Eczema, Endometriosis, Epilepsy, Fibrosing Alveolitis, Gastro-Esophageal Reflex Disease (GERD), Glaucoma, Gout, Graves Disease, Hemophilia, Hyperlipidemia, Hyperpituitarism, Hypopituitarism, Hypophyseal Adenoma, Hypothyroidism, Idiopathic Thrombocytopenic Purpura, Menopause, Migraine, Motor Neuron Disease, Multiple Sclerosis, Muscular Dystrophy, Myasthenia Gravis,
20 Myopathy, Neuropathy, Obesity, Obsessive Compulsive Disorder, Osteoarthritis, Osteoporosis, Paralysis (Paraplegia, Quadriplegia), Parkinson's disease, Paget's disease, Polyarteritis Nodosa, Polycystic ovarian disease, Psoriasis, Psoriatic Arthritis, Pulmonary Interstitial Fibrosis, Reflex Sympathetic Dystrophy (RSD) Syndrome, Rheumatoid Arthritis, Scleroderma, Schizophrenia, Sjögren's syndrome, Smoking and Tobacco-related Diseases,
25 Systemic Lupus Erythematosus, Trigeminal Neuralgia, Tuberculosis, Ulcerative Colitis or Urinary Incontinence.

291. The use of any one of claims 276-289, wherein said acute disease comprises Acute
30 Alcohol Poisoning, Acute Amebiasis, Acute Auditory Diseases, Acute Brain Diseases, Acute Cancers, Acute Cardiovascular Diseases, Acute Dental Diseases, Acute Dermatological

Diseases, Acute Esophageal Diseases, Acute Gastrointestinal Diseases, Acute Kidney Diseases, Acute Liver Diseases, Acute Muscular Diseases, Acute Ocular Diseases, Acute Poisoning, Acute Oral Diseases, Acute Osteoarthritis, Acute Pharyngeal Diseases, Acute Prostatic Diseases, Acute Physical Injury (Trauma), Acute Peritoneal Diseases, Acute
 5 Respiratory Diseases or Acute Urogenital Diseases.

292. The use of any one of claims 276-289, wherein said infectious disease is caused by an organism comprising *Bacillus anthracis*, *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella abortus*, *Brucella canis*, *Brucella melitensis*, *Brucella suis*, *Campylobacter jejuni*,

10 *Chlamydomphila pneumoniae*, *Chlamydia trachomatis*, *Chlamydomphila psittaci*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Entamoeba histolytica*, *Enterococcus faecalis*, *Enterococcus faecium*, *Escherichia coli*, *Enterotoxigenic Escherichia coli*, *Enteropathogenic E. coli*, *Enterohemorrhagic Escherichia coli* O157:H7, *Francisella tularensis*, Fungus (*Candida albicans*), *Haemophilus influenza*, *Helicobacter pylori*, *Leishmania* (Kala Azar), *Legionella pneumophila*, *Leptospira interrogans*, *Leptospira interrogans*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, Parasites (protozoa, helminths, ectoparasites), *Plasmodium* (malaria), *Pseudomonas aeruginosa*, *Rickettsia rickettsia*, *Salmonella enterica*
 15 subsp. *Enterica*, *Salmonella typhimurium*, *Shigella sonnei*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Treponema pallidum*, *Vibrio cholerae* or *Yersinia pestis*.

25 293. The use of any one of claims 276-289, wherein said infectious disease is caused by an organism comprising Adenovirus, SARS coronavirus, Coxsackievirus, Dengue virus, Ebola virus, Epstein-Barr virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Herpes simplex virus, Cytomegalovirus, Human immunodeficiency virus (HIV), Influenza virus, Measles virus, Mumps virus, Human papillomavirus, Human parainfluenza virus, Poliovirus,
 30 Rabies virus, Human respiratory syncytial virus, Rubella virus, Varicella zoster virus or Yellow fever virus.

294. Use of a kit for detecting a chronic disease, an acute disease, an infectious disease, a disorder or a syndrome, wherein the kit comprises a tool for measuring a defined quantity of stool, a biological sample dilution buffer, a substrate for phosphatase and a phosphatase assay buffer.

5

295. The use of claim 294, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

10

296. The use of claim 294 or 295, wherein said substrate for phosphatase is attached to a solid medium comprising a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle a slide or a swab, preferably nylon or nitrocellulose membrane.

15

297. The use of any one of claims 294-296, wherein said detecting comprises measuring concentration of phosphatase in a biological sample from a subject.

298. The use of claim 297, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

20

299. The use of claim 297 or 298, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken, turkey, preferably human.

25

300. The use of any one of claims 297-299, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

30

301. The use of any one of claims 294-300, wherein said chronic disease comprises Acne, Acquired Immunodeficiency Syndrome (AIDS), Ankylosing spondylitis, Asthma, Addison's disease, Allergic rhinitis, Alzheimer's Disease, Amyotrophic lateral sclerosis (ALS, Lou Gehrig's Disease), Aplastic anemia, Arthritis, Atherosclerosis, Attention Deficit Hyperactivity Disorder (ADHD, ADD), Autism Spectrum Disorder, Benign Prostatic Hypertrophy, Bipolar Mood Disorder, Bronchiectasis, Cancer, Cardiac Failure, Cardiomyopathy, Chronic Alcoholism, Chronic anemia, Chronic hepatitis, Chronic Amebiasis Chronic obstructive pulmonary disorder, Chronic gastritis, Chronic esophagitis, Chronic oral diseases, Chronic Poisoning, Chronic renal disease, Collagen diseases, Crohn's disease, Cushing's diseases, Cystic Fibrosis, Dementia, Dermatomyositis, Depression, Dysrhythmias, Eating Disorders (Anorexia Nervosa, Bulimia Nervosa, Binge Eating), Eczema, Endometriosis, Epilepsy, Fibrosing Alveolitis, Gastro-Esophageal Reflex Disease (GERD), Glaucoma, Gout, Graves Disease, Hemophilia, Hyperlipidemia, Hyperpituitarism, Hypopituitarism, Hypophyseal Adenoma, Hypothyroidism, Idiopathic Thrombocytopenic Purpura, Menopause, Migraine, Motor Neuron Disease, Multiple Sclerosis, Muscular Dystrophy, Myasthenia Gravis, Myopathy, Neuropathy, Obesity, Obsessive Compulsive Disorder, Osteoarthritis, Osteoporosis, Paralysis (Paraplegia, Quadriplegia), Parkinson's disease, Paget's disease, Polyarteritis Nodosa, Polycystic ovarian disease, Psoriasis, Psoriatic Arthritis, Pulmonary Interstitial Fibrosis, Reflex Sympathetic Dystrophy (RSD) Syndrome, Rheumatoid Arthritis, Scleroderma, Schizophrenia, Sjögren's syndrome, Smoking and Tobacco-related Diseases, Systemic Lupus Erythematosus, Trigeminal Neuralgia, Tuberculosis, Ulcerative Colitis or Urinary Incontinence.

302. The use of any one of claims 294-300, wherein said acute disease comprises Acute Alcohol Poisoning, Acute Amebiasis, Acute Auditory Diseases, Acute Brain Diseases, Acute Cancers, Acute Cardiovascular Diseases, Acute Dental Diseases, Acute Dermatological Diseases, Acute Esophageal Diseases, Acute Gastrointestinal Diseases, Acute Kidney Diseases, Acute Liver Diseases, Acute Muscular Diseases, Acute Ocular Diseases, Acute Poisoning, Acute Oral Diseases, Acute Osteoarthritis, Acute Pharyngeal Diseases, Acute Prostatic Diseases, Acute Physical Injury (Trauma), Acute Peritoneal Diseases, Acute Respiratory Diseases or Acute Urogenital Diseases.

303. The use of any one of claims 294-300, wherein said infectious disease is caused by an organism comprising *Bacillus anthracis*, *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella abortus*, *Brucella canis*, *Brucella melitensis*, *Brucella suis*, *Campylobacter jejuni*, *Chlamydomphila pneumoniae*, *Chlamydia trachomatis*, *Chlamydomphila psittaci*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Entamoeba histolytica*, *Enterococcus faecalis*, *Enterococcus faecium*, *Escherichia coli*, *Enterotoxigenic Escherichia coli*, *Enteropathogenic E. coli*, *Enterohemorrhagic Escherichia coli* O157:H7, *Francisella tularensis*, Fungus (*Candida albicans*), *Haemophilus influenza*, *Helicobacter pylori*, *Leishmania* (Kala Azar), *Legionella pneumophila*, *Leptospira interrogans*, *Leptospira interrogans*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, Parasites (protozoa, helminths, ectoparasites), *Plasmodium* (malaria), *Pseudomonas aeruginosa*, *Rickettsia rickettsia*, *Salmonella enterica* subsp. *Enterica*, *Salmonella typhimurium*, *Shigella sonnei*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Treponema pallidum*, *Vibrio cholerae* or *Yersinia pestis*.

304. The use of any one of claims 294-300, wherein said infectious disease is caused by an organism comprising *Adenovirus*, *SARS coronavirus*, *Coxsackievirus*, *Dengue virus*, *Ebola virus*, *Epstein-Barr virus*, *Hepatitis A virus*, *Hepatitis B virus*, *Hepatitis C virus*, *Herpes simplex virus*, *Cytomegalovirus*, *Human immunodeficiency virus (HIV)*, *Influenza virus*, *Measles virus*, *Mumps virus*, *Human papillomavirus*, *Human parainfluenza virus*, *Poliovirus*, *Rabies virus*, *Human respiratory syncytial virus*, *Rubella virus*, *Varicella zoster virus* or *Yellow fever virus*.

305. Use of a temporal phosphatase profile for detecting a chronic disease, an acute disease, an infectious disease, a disorder or a syndrome comprising measuring the concentration of phosphatase in one or more than one biological samples from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

306. The use of claim 305, wherein said multiple time points comprise daily, weekly, monthly or yearly.

307. The use of claim 305 or 306, wherein said subject is a human, cattle, pig, sheep, goat,
5 cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

308. The use of any one of claims 305-307, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
10 waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

309. The use of any one of claims 305-307, wherein said measuring comprises:

15 providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
waiting a specified period of time to allow a color to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
20 quantifying the concentration of phosphatase in said biological sample.

310. The use of any one of claims 305-307, wherein said measuring comprises:

providing a substrate for phosphatase in a solid or liquid form;
contacting said substrate for phosphatase with a biological sample;
25 waiting a specified period of time to allow an enzymatic reaction to occur; and
quantifying the concentration of phosphatase in said biological sample using a
spectrophotometer, a biochemistry analyzer, a high-performance liquid
chromatography (HPLC) machine, a colorimeter, a luminometer or a mechanical,
biochemical, electrical or electronic device.

311. The use of any one of claims 305-307, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
waiting a specified period of time to allow a color to develop;
comparing the developed color with photographs of standards; and
5 quantifying the concentration of phosphatase in said biological sample.

312. The use of any one of claims 305-307, wherein said measuring comprises:

providing a solid medium having a substrate for phosphatase attached thereto;
contacting said solid medium with a biological sample;
10 waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
quantifying the concentration of phosphatase in said biological sample.

15 313. The use of any one of claims 305-307, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;
attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a color to develop;
20 comparing the developed color with photographs of standards; and
quantifying the concentration of phosphatase in said biological sample.

314. The use of any one of claims 305-307, wherein said measuring comprises:

providing a solid medium for attaching a biological sample thereto;
25 attaching said biological sample to said solid medium;
contacting said solid medium with a substrate for phosphatase;
waiting a specified period of time to allow a pixilated color pattern to develop;
counting the pixels of photographs;
comparing the number of pixels to standards; and
30 quantifying the concentration of phosphatase in said biological sample.

315. The use of claims 311-314, wherein said solid medium comprises a membrane, paper, stick, glass, plastic, wood, a bead, a tube, a bottle, a slide or a swab, preferably nylon or nitrocellulose membrane.

316. The use of any one of claims 308-315, wherein said substrate for phosphatase comprises p-nitrophenyl phosphate, 5-Bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium substrate system, Fast Red TR/Naphthol substrate system, CDP-star substrate (2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'-(5'-chloro)-tricyclo [3.3.1.1^{3,7}]decan}-4-yl)-1-phenyl phosphate disodium salt) or combinations thereof.

317. The use of any one of claims 305-316, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

318. The use of any one of claims 305-317, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

319. The use of any one of claims 305-318, wherein said chronic disease comprises Acne, Acquired Immunodeficiency Syndrome (AIDS), Ankylosing spondylitis, Asthma, Addison's disease, Allergic rhinitis, Alzheimer's Disease, Amyotrophic lateral sclerosis (ALS, Lou Gehrig's Disease), Aplastic anemia, Arthritis, Atherosclerosis, Attention Deficit Hyperactivity Disorder (ADHD, ADD), Autism Spectrum Disorder, Benign Prostatic Hypertrophy, Bipolar Mood Disorder, Bronchiectasis, Cancer, Cardiac Failure, Cardiomyopathy, Chronic Alcoholism, Chronic anemia, Chronic hepatitis, Chronic Amebiasis Chronic obstructive pulmonary disorder, Chronic gastritis, Chronic esophagitis, Chronic oral diseases, Chronic Poisoning, Chronic renal disease, Collagen diseases, Crohn's disease, Cushing's diseases, Cystic Fibrosis, Dementia, Dermatomyositis, Depression, Dysrhythmias, Eating Disorders

(Anorexia Nervosa, Bulimia Nervosa, Binge Eating), Eczema, Endometriosis, Epilepsy, Fibrosing Alveolitis, Gastro-Esophageal Reflex Disease (GERD), Glaucoma, Gout, Graves Disease, Hemophilia, Hyperlipidemia, Hyperpituitarism, Hypopituitarism, Hypophyseal Adenoma, Hypothyroidism, Idiopathic Thrombocytopenic Purpura, Menopause, Migraine, 5 Motor Neuron Disease, Multiple Sclerosis, Muscular Dystrophy, Myasthenia Gravis, Myopathy, Neuropathy, Obesity, Obsessive Compulsive Disorder, Osteoarthritis, Osteoporosis, Paralysis (Paraplegia, Quadriplegia), Parkinson's disease, Paget's disease, Polyarteritis Nodosa, Polycystic ovarian disease, Psoriasis, Psoriatic Arthritis, Pulmonary Interstitial Fibrosis, Reflex Sympathetic Dystrophy (RSD) Syndrome, Rheumatoid Arthritis, 10 Scleroderma, Schizophrenia, Sjögren's syndrome, Smoking and Tobacco-related Diseases, Systemic Lupus Erythematosus, Trigeminal Neuralgia, Tuberculosis, Ulcerative Colitis or Urinary Incontinence.

320. The use of any one of claims 305-318, wherein said acute disease comprises Acute 15 Alcohol Poisoning, Acute Amebiasis, Acute Auditory Diseases, Acute Brain Diseases, Acute Cancers, Acute Cardiovascular Diseases, Acute Dental Diseases, Acute Dermatological Diseases, Acute Esophageal Diseases, Acute Gastrointestinal Diseases, Acute Kidney Diseases, Acute Liver Diseases, Acute Muscular Diseases, Acute Ocular Diseases, Acute Poisoning, Acute Oral Diseases, Acute Osteoarthritis, Acute Pharyngeal Diseases, Acute 20 Prostatic Diseases, Acute Physical Injury (Trauma), Acute Peritoneal Diseases, Acute Respiratory Diseases or Acute Urogenital Diseases.

321. The use of any one of claims 305-318, wherein said infectious disease is caused by an organism comprising *Bacillus anthracis*, *Bordetella pertussis*, *Borrelia burgdorferi*, *Brucella abortus*, *Brucella canis*, *Brucella melitensis*, *Brucella suis*, *Campylobacter jejuni*, 25 *Chlamydophila pneumoniae*, *Chlamydia trachomatis*, *Chlamydophila psittaci*, *Clostridium botulinum*, *Clostridium difficile*, *Clostridium perfringens*, *Clostridium tetani*, *Corynebacterium diphtheriae*, *Entamoeba histolytica*, *Enterococcus faecalis*, *Enterococcus faecium*, *Escherichia coli*, *Enterotoxigenic Escherichia coli*, *Enteropathogenic E. coli*, 30 *Enterohemorrhagic Escherichia coli* O157:H7, *Francisella tularensis*, Fungus (*Candida albicans*), *Haemophilus influenza*, *Helicobacter pylori*, *Leishmania* (Kala Azar), *Legionella*

pneumophila, *Leptospira interrogans*, *Leptospira interrogans*, *Listeria monocytogenes*, *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, Parasites (protozoa, helminths, ectoparasites), *Plasmodium* (malaria), *Pseudomonas aeruginosa*, *Rickettsia rickettsia*, *Salmonella enterica* subsp. *Enterica*, *Salmonella typhimurium*, *Shigella sonnei*, *Staphylococcus aureus*,
5 *Staphylococcus epidermidis*, *Staphylococcus saprophyticus*, *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Treponema pallidum*, *Vibrio cholerae* or *Yersinia pestis*.

- 10 322. The use of any one of claims 305-318, wherein said infectious disease is caused by an organism comprising Adenovirus, SARS coronavirus, Coxsackievirus, Dengue virus, Ebola virus, Epstein-Barr virus, Hepatitis A virus, Hepatitis B virus, Hepatitis C virus, Herpes simplex virus, Cytomegalovirus, Human immunodeficiency virus (HIV), Influenza virus, Measles virus, Mumps virus, Human papillomavirus, Human parainfluenza virus, Poliovirus,
15 Rabies virus, Human respiratory syncytial virus, Rubella virus, Varicella zoster virus or Yellow fever virus.

323. An enzyme-linked immunosorbent assay (ELISA) system for determining concentration of phosphatase in a biological sample comprising a solid-phase support system, a primary
20 and/or secondary antibody that specifically binds to phosphatase, a horseradish peroxidase (HRP)- or alkaline phosphatase (AP)-conjugated antibody that specifically binds to said primary and/or secondary antibody, and chromogenic substrates for HRP or AP.

324. The system of claim 323, wherein said biological sample comprises stool, saliva, urine,
25 blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

325. Use of an enzyme-linked immunosorbent assay (ELISA) system for determining concentration of phosphatase in a biological sample comprising a solid-phase support system,
30 a primary and/or secondary antibody that specifically binds to phosphatase, a horseradish

peroxidase (HRP)- or alkaline phosphatase (AP)-conjugated antibody that specifically binds to said primary and/or secondary antibody, and chromogenic substrates for HRP or AP.

326. The use of claim 325, wherein said biological sample comprises stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

327. An enzyme-linked immunosorbent assay (ELISA) system for detecting incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease or metabolic syndrome, wherein the system comprises a solid-phase support system, a primary and/or secondary antibody that specifically binds to phosphatase, a horseradish peroxidase (HRP)- or alkaline phosphatase (AP)-conjugated antibody that specifically binds to said primary and/or secondary antibody, and chromogenic substrates for HRP or AP.

328. The system of claim 327, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

329. The system of claim 327, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

330. The system of claim 329, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

331. The system of any one of claims 328-330, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken, turkey, preferably human.

332. The system of any one of claims 327-331, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with
5 phosphatase activity, preferably intestinal alkaline phosphatase.

333. The system of any one of claims 328-332, wherein said subject provides a biological sample comprising stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body
10 cavity fluid, preferably a stool sample.

334. The system of any one of claims 327-333, wherein said system further comprises a means for measuring concentration of one or more of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory
15 factors in one or more than one biological sample from a subject, wherein said one or more than one biological samples is collected at a single time point or at multiple time points.

335. The system of any one of claims 327-334, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

336. The system of any one of claims 327-334, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis,
25 eczema or psoriasis.

337. The system of any one of claims 327-334, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein
30 (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

338. Use of an enzyme-linked immunosorbent assay (ELISA) system for detecting incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome, wherein the system comprises a solid-phase support system, a primary and/or secondary antibody that specifically binds to phosphatase, a horseradish peroxidase (HRP)- or alkaline phosphatase (AP)-conjugated antibody that specifically binds to said primary and/or secondary antibody, and chromogenic substrates for HRP or AP.

339. The use of claim 338, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a concentration of phosphatase that is 1% - 100% less than an average normal value.

340. The use of claim 338, wherein said incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome is characterized by a subject having a temporal phosphatase profile comprising a persistent decrease in phosphatase concentration over time.

341. The use of claim 340, wherein said persistent decrease is about a 50% decrease from a previously measured concentration.

342. The use of any one of claims 339-341, wherein said subject is a human, cattle, pig, sheep, goat, cow, horse, dog, cat, monkey, rabbit, rat, mouse, chicken or turkey, preferably human.

343. The use of any one of claims 338-342, wherein said phosphatase comprises an alkaline phosphatase, intestinal alkaline phosphatase, placental alkaline phosphatase, nonspecific tissue alkaline phosphatase (liver/bone/kidney alkaline phosphatase), germ cell alkaline phosphatase, neutrophil alkaline phosphatase, an acid phosphatase, or a peptide with phosphatase activity, preferably intestinal alkaline phosphatase.

344. The use of any one of claims 339-343, wherein said subject provides a biological sample comprising stool, saliva, urine, blood, plasma, serum, tears, sweat, pus, gastric fluid, intestinal fluid, ocular fluid, peritoneal fluid, vaginal fluid, cerebrospinal fluid or a body cavity fluid, preferably a stool sample.

5

345. The use of any one of claims 338-344, wherein said system further comprises a means for measuring concentration of one or more of serum endotoxins, proinflammatory nucleotides, cytokines, chemokines, antagonists of cytokine receptors or anti-inflammatory factors in one or more than one biological sample from a subject, wherein said one or more
10 than one biological samples is collected at a single time point or at multiple time points.

346. The use of any one of claims 338-345, wherein said incipient diabetes or diabetes comprises type 2 diabetes mellitus, gestational diabetes or diabetes insipidus.

15

347. The use of any one of claims 338-345, wherein said incipient metabolic disease or metabolic disease comprises a cardiovascular disease, coronary heart disease, cerebrovascular disease, stroke, peripheral vascular disease, nonalcoholic fatty liver disease, retinopathy, asthma, chronic pulmonary obstructive disease, cystic fibrosis, Alzheimer's disease, arthritis, eczema or psoriasis.

20

348. The use of any one of claims 338-345, wherein said incipient metabolic syndrome or metabolic syndrome comprises obesity, hyperglycemia, insulin resistance, hyperinsulinemia, dyslipidemia, hypercholesterolemia, hypertriglyceridemia, low high-density lipoprotein (HDL), high low-density lipoprotein (LDL), hypertension, fatty liver or combinations thereof.

25

349. A method of quantifying alkaline phosphatase in a stool sample comprising:
 contacting said stool sample with a phosphatase substrate;
 allowing alkaline phosphatase contained within said stool sample to hydrolyze said phosphatase substrate to release inorganic phosphate;
30 treating said inorganic phosphate with malachite green molybdate to form green molybdophosphoric acid complex; and

quantifying concentration of said complex;
wherein concentration of said complex is proportional to the concentration of alkaline phosphatase in said stool sample.


5 350. The use of malachite green-based phosphate quantitation assay, wherein the assay comprises:

contacting a stool sample with a phosphatase substrate;
allowing alkaline phosphatase contained within said stool sample to hydrolyze said phosphatase substrate to release inorganic phosphate;
10 treating said inorganic phosphate with malachite green molybdate to form green molybdophosphoric acid complex; and
quantifying concentration of said complex;

wherein concentration of said complex is proportional to the concentration of alkaline phosphatase in said stool sample.

15

FIG. 1



2/5

FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4





FIG. 5



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/061017

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - C12Q 1/42 (2016.01)

CPC - C12Q 1/42 (2016.02)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - C12Q 1/00, 1/34, 1/42; G01N 33/50, 33/53, 33/573 (2016.01)

CPC - C12Q 1/00, 1/34, 1/42; G01N 33/50, 33/53, 33/573 (2016.02)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC - 435/4, 18, 21; IPC(8) - C12Q 1/00, 1/34, 1/42; G01N 33/50, 33/53, 33/573 (2016.01); CPC - C12Q 1/00, 1/34, 1/42; G01N 33/50, 33/53, 33/573 (2016.02) (keyword delimited)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Orbit, Google Patents, Google Scholar, Google

Search terms used: diabetes metabolic disease phosphatase monitor measure record assay stool feces poop intestinal gut alkaline diagnose concentration level activity liver leukemia hypothyroidism metabolic syndrome disease hyperglycemia fatty liver heart disease cardiovascular disease decrease previous comparative

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	EP 1 487 501 B1 (THE JOHNS HOPKINS UNIVERSITY) 05 December 2012 (05.12.2012) entire document	1, 3, 37, 39, 258, 260, 276, 278 ----- 2, 38, 51-54, 149-151, 156-158, 215, 216, 259, 277, 294-296
X --- Y	TALBOT et al. Serum Phosphatase as an Aid in the Diagnosis of Cretinism and Juvenile Hypothyroidism. The American Journal of Diseases of Children. Vol. 62, No. 2. August 1941. [retrieved on 25 February 2016]. Retrieved from the Internet. <URL: http://archpedi.jamanetwork.com/article.aspx?articleid=1179250 >. entire document	15-17, 193, 237, 239, 305, 307 ----- 18, 51-54, 194, 215, 216, 238, 306
Y	BOZZETTI et al. Metabolic Bone Disease in preterm newborn: an update on nutritional issues. Italian Journal of Pediatrics. Vol. 35. 2009. [retrieved on 25 February 2016]. Retrieved from the Internet. <URL: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2729305/ >. entire document	2, 18, 38, 54, 194, 216, 238, 259, 277, 306
Y	GALINDO. Alkaline Phosphate (ALP). 23 August 2010. [retrieved on 25 February 2016]. Retrieved from the Internet. <URL: http://www.isu.edu/~galisusa/alp_sop.html >. entire document	149-151, 156-158, 294-296



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

26 February 2016

Date of mailing of the international search report

11 MAR 2016

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, VA 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/061017

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	APPLEYARD et al. Pretreatment with the probiotic VSL#3 delays transition from inflammation to dysplasia in a rat model of colitis-associated cancer. American Journal of Physiology Gastrointestinal and Liver Physiology. Vol. 301. 2011. [retrieved on 17 December 2015]. Retrieved from the Internet. <URL: http://ajpgi.physiology.org/content/ajpgi/301/6/G1004.full.pdf >. pp. G1004-G1013	1-3, 15-18, 37-39, 51-54, 149-151, 156-158, 163-167, 178-182, 193-198, 215-220, 237-241, 258-260, 276-278, 294-296, 305-307
A	LAN et al. IA-2, a transmembrane protein of the protein tyrosine phosphatase family, is a major autoantigen in insulin-dependent diabetes mellitus. PNAS, Proceedings of the National Academy of Sciences USA. Vol. 93. June 1996. [retrieved on 17 December 2015]. Retrieved from the Internet. <URL: http://www.pnas.org/content/93/13/6367.full.pdf >. pp. 6367-6370	1-3, 15-18, 37-39, 51-54, 149-151, 156-158, 163-167, 178-182, 193-198, 215-220, 237-241, 258-260, 276-278, 294-296, 305-307

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/061017

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: see claims listed in Extra Sheet
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See Extra Sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-3, 15-18, 37-39, 51-54, 149-151, 156-158, 163-167, 178-182, 193-198, 215-220, 237-241, 258-260, 276-278, 294-296, 305-307

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2015/061017

Claims Nos. 4-14, 19-36, 40-50, 55-72, 78-91, 97-110, 116-129, 135-148, 152-155, 159-162, 168-177, 183-192, 199-214, 221-236, 242-257, 261-275, 279-293, 297-304, 308-322, 332-337, and 343-348 have been found to be unsearchable under Article 17(2)(b) because of defects under Article 17(2)(a) and therefore have not been included with any invention.

<Continued from Box No. III: Observations where unity of invention is lacking>

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees need to be paid.

Group I: Claims 1-3, 15-18, 37-39, 51-54, 149-151, 156-158, 163-167, 178-182, 193-198, 215-220, 237-241, 258-260, 276-278, 294-296, and 305-307 are drawn to methods for developing a temporal profile, methods thereof, uses thereof, and kits thereof.

Group II: Claims 73-77, 92-96, 111-115, and 130-134 are drawn to methods of decreasing risk of or a method of treatment, uses thereof, pharmaceutical compositions thereof.

Group III: Claims 323-331 and 338-342 are drawn to enzyme-linked immunosorbent assay (ELISA) systems and uses thereof.

Group IV: Claims 349 and 350 are drawn to methods of quantifying alkaline phosphatase.

The inventions listed in Groups I through IV do not relate to a single general inventive concept under PCT Rule 13.1, because under PCT Rule 13.2 they lack the same or corresponding special technical features for the following reasons:

The special technical features of Group I, methods for developing a temporal profile, methods thereof, uses thereof, and kits thereof, are not present in Groups II through IV; the special technical features of Group II, methods of decreasing risk of or a method of treatment, uses thereof, pharmaceutical compositions thereof, are not present in Groups I, III, and IV; the special technical features of Group III, enzyme-linked immunosorbent assay (ELISA) systems and uses thereof, are not present in Groups I, II, and IV; and the special technical features of Group IV, methods of quantifying alkaline phosphatase, are not present in Groups I through III.

The Groups I through IV share the technical features of a method comprising measuring concentration of phosphatase in a stool sample; a method for diagnosing incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising measuring concentration of phosphatase in a biological sample; use of a substrate of phosphatase for measuring the concentration of phosphatase in one or more than one biological samples; use of a substrate of phosphatase for detecting incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising measuring the concentration of phosphatase in a biological sample; and determining prognosis of incipient diabetes, incipient metabolic disease, incipient metabolic syndrome, diabetes, metabolic disease, or metabolic syndrome comprising measuring the concentration of phosphatase in one or more than one biological sample from a subject. However, these shared technical features do not represent a contribution over the prior art.

Specifically, "Pretreatment with the probiotic VSL#3 delays transition from inflammation to dysplasia in a rat model of colitis-associated cancer" to Appleyard et al. teach a method comprising measuring concentration of phosphatase in a stool sample (See Pg. G1006, Col. 2, Para. 2); and use of a substrate of phosphatase for measuring the concentration of phosphatase in one or more than one biological samples (See Pg. G1006, Col. 2, Para. 2).

Additionally, "IA-2, a transmembrane protein of the protein tyrosine phosphatase family, is a major autoantigen in insulin-dependent diabetes mellitus" to Lan et al. teach a method for diagnosing diabetes comprising measuring phosphatase in a biological sample (See Pg. 6367, Col. 2, Materials and Methods; Pg. 6368, Col. 1, Para. 5. The sera used in Fig. 2 also were tested by D-ELISA for antibodies that reacted with GAD65. As seen in Figs. 3 and 4, 52% of the sera from IDDM patients, but none of the sera from controls, reacted with GAD 65, up to 86% of the IDDM patients developed autoantibodies to IA-2 and/or GAD65. Some of the patients (34%) developed antibodies to both antigens, 12% to only GAD65, and 4% to only IA-2. The current data, together with data from other studies (15), suggest that the age of the onset of IDDM may influence the nature of the immune response. Of 60 patients diagnosed with IDDM before age 20, 68% had antibodies to IA-2 and 60% had antibodies to GAD65. However, of 15 patients diagnosed with IDDM after age 20, only 46% had antibodies to IA-2, whereas 86% had antibodies to GAD65.; Pg. 6369, Fig. 4); use of a substrate of phosphatase for detecting diabetes comprising measuring the concentration of phosphatase in a biological sample (See Pg. 6367, Col. 2, Materials and Methods; Pg. 6368, Col. 1, Para. 5; Pg. 6369, Fig. 4); and determining prognosis of diabetes comprising measuring phosphatase in one or more than one biological sample from a subject (See Pg. 6367, Col. 2, Materials and Methods; Pg. 6368, Col. 1, Para. 5; Pg. 6369, Fig. 4).

The inventions listed in Groups I through IV therefore lack unity under Rule 13 because they do not share a same or corresponding special technical feature.