



(51) International Patent Classification:

G01N 33/62 (2006.01) A61P 19/06 (2006.01)

A61K 31/436 (2006.01) G01N 33/68 (2006.01)

A61K 38/44 (2006.01)

(21) International Application Number:

PCT/US2020/036116

(22) International Filing Date:

04 June 2020 (04.06.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/856,844 04 June 2019 (04.06.2019) US

62/932,728 08 November 2019 (08.11.2019) US

(71) Applicant: SELECTA BIOSCIENCES, INC. [GB/GB];

65 Grove Street, Watertown MA 02472 (GB).

(72) Inventors: JOHNSTON, Lloyd; 24 Oakley Road, Bel-

mont, MA 02478 (US). KISHIMOTO, Takashi, Kei; 46

Coolidge Avenue, Lexington, MA 02420 (US). SANDS,

Earl; 22 Blackheath Terrace, Alpharetta, GA 30005 (US).

CAUTREELS, Werner; 221 Massachusetts Avenue, No. 903, Boston, MA 02115 (US).

(74) Agent: VON REYN, Jessica, A. et al.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: FORMULATIONS AND DOSES OF PEGYLATED URICASE




Fig. 10

(57) Abstract: Provided herein are methods and compositions and kits related to identification and/or treatment of subjects, including subjects with hyperuricemia, gout or a condition associated with gout, and for preventing gout flare.



Published:

— *with international search report (Art. 21(3))*

FORMULATIONS AND DOSES OF PEGYLATED URICASE

RELATED APPLICATIONS

This application claims the benefit of priority under 35 U.S.C. § 119(e) of U.S. Provisional Application Serial No. 62/856,844, filed on June 4, 2019; U.S. Provisional Application Serial No. 62/932,728, filed on November 8, 2019; the entire contents of each of which are incorporated herein by reference.

FIELD OF THE INVENTION

Provided herein are methods and compositions and kits related to uricase compositions and/or compositions comprising synthetic nanocarriers comprising an immunosuppressant. Also provided herein are methods and compositions and kits for the treatment of subjects, including subjects with hyperuricemia, gout or a condition associated with gout, and for preventing gout flare. Also provided herein are methods of identifying a subject for treatment provided herein as well as methods for monitoring the effectiveness of a treatment. In some embodiments, the methods comprise step(s) of obtaining or determining an anti-uricase titer in the subject and/or comparing the anti-uricase titer with a threshold value. In some embodiments of such embodiments, the methods can further comprise any one of the steps or methods provided herein for treating any one of the subjects provided herein.

SUMMARY OF THE INVENTION

The development of anti-drug antibodies (ADAs) is a common cause for biotherapeutic treatment failure and adverse hypersensitivity reactions. It has been demonstrated that synthetic nanocarriers comprising an immunosuppressant are capable of inducing immunological tolerance to a composition comprising uricase, resulting in improved efficacy of the uricase-comprising composition. The improved efficacy has been demonstrated at least with a significantly higher rate of reduction in serum uric acid levels over time as compared to other treatments. It has also been demonstrated that synthetic nanocarriers comprising an immunosuppressant, when administered concomitantly with a composition comprising uricase, are capable of significantly reducing the incidence of gout flare as compared to other treatments. The compositions comprising synthetic nanocarriers comprising an immunosuppressant and compositions comprising a uricase as provided herein

-2-

can be used to efficaciously and durably (e.g., for at least 30 days) reduce serum uric acid levels and/or reduce the incidence of gout flare.

It has also been discovered that the use of a measure of anti-uricase antibodies in a subject can be helpful in the treatment of the subject. The measure of anti-uricase antibodies can be an anti-uricase titer in the subject. The measure can be compared to a threshold. Thus provided herein are methods comprising determining or obtaining a level of anti-uricase antibodies in any one of the subjects provided herein, such as in a sample from the subject, and/or comparing a level of anti-uricase antibodies in the subject to a threshold. In any one of the methods provided herein, the level of anti-uricase antibodies can be given as a titer. In any one of the methods provided herein, the threshold is a titer of less than about 1080. In any one of the methods provided herein, the threshold is about 1080. Any one of the foregoing methods may also comprise any one of more steps or any one of the methods of treatment as provided herein.

Also provided herein are methods comprising administering to a subject having an anti-uricase titer below a threshold as provided herein any one of the compositions comprising uricase provided herein alone or in combination with any one of the compositions comprising synthetic nanocarriers comprising an immunosuppressant provided herein. Also provided herein are methods of preventing gout flare, comprising concomitantly administering to a subject having an anti-uricase titer below a threshold a composition comprising synthetic nanocarriers comprising an immunosuppressant and a composition comprising uricase, such as one that is not administered an additional therapeutic to prevent gout flare concomitantly with the concomitant administration. In some embodiments, the subject is identified as having had or as being expected to have gout flare from treatment with a gout therapy without concomitant administration of an additional therapeutic to prevent gout flare. The subject may be in need thereof. The subject may be any one of the subjects described herein.

Also provided herein are methods of treating a subject having an anti-uricase titer below a threshold with gout or a condition associated with gout comprising administering any one of the compositions comprising uricase provided herein alone or in combination with any one of the compositions comprising synthetic nanocarriers comprising an immunosuppressant provided herein. In one embodiment of any one of the methods provided herein, the compositions comprising uricase provided herein alone or in combination with any one of the compositions comprising synthetic nanocarriers comprising an immunosuppressant may be

-3-

repeatedly administered to the subject. The subject may be one in need thereof. The subject may be any one of the subjects described herein.

In one aspect, a method of treating a human subject with gout or a condition associated with gout and having an anti-uricase titer below a threshold, comprising
5 administering to the subject a composition comprising uricase and a pharmaceutically acceptable carrier is provided. In one embodiment, the administration is via a non-intramuscular mode of administration. In one embodiment, the composition comprising uricase and a pharmaceutically acceptable carrier is administered more than once to the subject. In one embodiment, the composition comprising uricase and a pharmaceutically
10 acceptable carrier is administered more than twice, more than thrice, or more than four times to the subject. In one embodiment, the composition comprising uricase and a pharmaceutically acceptable carrier is administered every two to four weeks. In one embodiment, the composition comprising uricase and a pharmaceutically acceptable carrier is administered monthly. In one embodiment, the composition comprising uricase and a
15 pharmaceutically acceptable carrier is administered concomitantly with a composition comprising an immunosuppressant.

In one aspect, a method of treating a subject with gout or a condition associated with gout and having an anti-uricase titer below a threshold, comprising concomitantly
20 administering to the subject a composition comprising synthetic nanocarriers comprising an immunosuppressant and a composition comprising uricase is provided.

Also provided herein are methods of treating a subject having an anti-uricase titer below a threshold that may experience gout flare comprising administering any one of the compositions comprising uricase provided herein in combination with any one of the compositions comprising synthetic nanocarriers comprising an immunosuppressant provided
25 herein. In one aspect, a method of preventing gout flare in a subject having an anti-uricase titer below a threshold, comprising concomitantly administering to the subject a composition comprising synthetic nanocarriers comprising an immunosuppressant and a composition comprising uricase. In one embodiment, the subject is not administered an additional therapeutic to prevent the gout flare, such as an anti-gout flare therapeutic, concomitantly
30 with the concomitant administration. In some embodiments, the subject is not administered colchicine or an NSAID concomitantly with the concomitant administration. In one embodiment, the subject is identified as having had or as being expected to have gout flare from treatment with a gout therapeutic, such as a uric acid lowering therapeutic. In one

-4-

embodiment, the subject is identified as having had or as being expected to have gout flare without concomitant administration of an additional therapeutic to prevent the gout flare.

In one embodiment of any one of the methods provided herein, the subject is identified or monitored to have or maintain an anti-uricase antibody level (such as an antibody titer) below the threshold. In one embodiment of any one of the methods provided herein, the treatment doses and/or frequency is adjusted based on the identification or monitoring. In one embodiment of any one of the methods provided herein, the method further comprises one or more steps for identifying or monitoring the subject to determine an anti-uricase antibody level (such as an antibody titer).

In one embodiment of any one of the methods provided herein, the concomitant administration occurs more than once in the subject. In one embodiment of any one of the methods provided herein, the concomitant administration occurs at least twice (*e.g.*, at least three, four, five, six, seven, eight, nine, or ten times) in the subject. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered concomitantly every two to four weeks. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered monthly concomitantly. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered monthly for at least three months (*e.g.*, 4, 5, 6, 7, 8, 9, 10 or more months) concomitantly.

In one embodiment of any one of the methods provided herein, the composition comprising uricase is administered at a label dose of 0.1 mg/kg – 1.2 mg/kg uricase with each administration, such as each concomitant administration. In one embodiment of any one of the methods provided herein, the composition comprising uricase is administered at a label dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1.0 mg/kg, 1.1 mg/kg, or 1.2 mg/kg uricase with each administration, such as each concomitant administration. In one embodiment of any one of the methods provided herein, the composition comprising uricase is administered at a label dose of 0.2 - 0.4 mg/kg uricase with each administration, such as each concomitant administration.

In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a

-5-

label dose of 0.05 mg/kg – 0.5 mg/kg immunosuppressant with each concomitant administration. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.05 mg/kg, 0.07 mg/kg, 0.075 mg/kg, 0.08 mg/kg, 0.1 mg/kg, 0.125 mg/kg, 0.15 mg/kg, 0.2 mg/kg, 0.25 mg/kg, 0.3 mg/kg, 0.35 mg/kg, 0.4 mg/kg, 0.45 mg/kg, or 0.5 mg/kg immunosuppressant with each concomitant administration. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.075 – 0.2 mg/kg or 0.08 – 0.125 mg/kg immunosuppressant with each concomitant administration.

In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.5 mg/kg – 6.5 mg/kg with each concomitant administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.55 mg/kg, 0.6 mg/kg, 0.65 mg/kg, 0.7 mg/kg, 0.75 mg/kg, 0.8 mg/kg, 0.85 mg/kg, 0.9 mg/kg, 0.95 mg/kg, 1.0 mg/kg, 1.10 mg/kg, 1.125 mg/kg, 1.5 mg/kg, 1.75 mg/kg, 2.0 mg/kg, 2.5 mg/kg, 3.0 mg/kg, 3.5 mg/kg, 4.0 mg/kg, 4.5 mg/kg, 5 mg/kg, 5.5 mg/kg, 6.0 mg/kg, or 6.5 mg/kg with each concomitant administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.6 – 2.5 mg/kg, 0.7 - 2.5 mg/kg, 0.8 – 2.5 mg/kg, 0.9 - 2.5 mg/kg, 1.0 - 2.5 mg/kg, 1.5 - 2.5 mg/kg, or 2.0 - 2.5 mg/kg with each concomitant administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.65 – 2.5 mg/kg, 0.65 – 2.0 mg/kg, 0.65 – 1.5 mg/kg, or 0.65 – 1.0 mg/kg with each concomitant administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.75 – 2.0 mg/kg, 0.8 – 1.5 mg/kg, 0.9 – 1.5 mg/kg or 1 - 2 mg/kg with each concomitant administration, wherein the

-6-

dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a label dose of 0.9 – 2 mg/kg or 1 – 1.5 mg/kg with each concomitant administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant.

In one embodiment of any one of the methods provided herein, the method further comprises administering a composition comprising uricase to the subject at least once (*e.g.*, at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more times) after the concomitant administration(s) without concomitant administration of an additional therapeutic, such as a composition comprising an immunosuppressant, such as a composition comprising synthetic nanocarriers comprising an immunosuppressant. In one embodiment of any one of the methods provided herein, the method further comprises administering the composition comprising uricase at least twice after the concomitant administration(s). In one embodiment of any one of the methods provided herein, the method further comprises administering the composition comprising uricase monthly for two months after the concomitant administration(s) each administration without concomitant administration of an additional therapeutic, such as a composition comprising an immunosuppressant, such as a composition comprising synthetic nanocarriers comprising an immunosuppressant. In some embodiments, the composition comprising uricase is administered at a label dose of 0.1 – 1.2 mg/kg uricase with each administration after the one or more concomitant administrations without an immunosuppressant. In some embodiments, the composition comprising uricase is administered at a label dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1.0 mg/kg, 1.1 mg/kg, 1.2 mg/kg uricase with each administration after the one or more concomitant administrations without an immunosuppressant.

In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered prior to the composition comprising uricase, such as with each concomitant administration. In one embodiment of any one of the methods provided herein, the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered within an hour of each other.

In one embodiment of any one of the methods provided herein, the subject is not administered an additional therapeutic, such as an additional gout therapeutic, such as one to prevent gout flare. In one embodiment of these embodiments, the additional therapeutic,

-7-

such as the additional gout therapeutic, such as one to prevent gout flare, is not administered concomitantly with each concomitant administration.

Any one of the methods, compositions or kits provided herein may be used to treat any one of the subjects provided herein.

5 In one embodiment of any one of the methods, compositions or kits provided herein, the subject has an elevated serum uric acid level. In one embodiment of any one of the methods, compositions or kits provided herein, the subject has a serum uric acid level of ≥ 5 mg/dL. In one embodiment of any one of the methods, compositions or kits provided herein, the subject has a serum uric acid level of ≥ 6 mg/dL. In one embodiment of any one of the
10 methods, compositions or kits provided herein, the subject has a serum uric acid level of ≥ 7 mg/dL. In one embodiment of any one of the methods, compositions or kits provided herein, the subject has or is at risk of having hyperuricemia; acute gout; chronic gout with or without tophi; idiopathic gout; refractory gout; secondary gout; unspecified gout; gout associated with a cardiovascular condition, renal condition, pulmonary condition, neurological condition,
15 ocular condition, dermatological condition or hepatic condition; or has had a gout attack or gout flare. In one embodiment of any one of the methods, compositions or kits provided herein, the subject is expected to have gout flare from treatment with a gout therapeutic, such as a uric acid lowering therapeutic, such as a composition comprising uricase. In one embodiment of any one of the methods, compositions or kits provided herein, the subject has
20 gout having at least one of a) tophi, b) gout flare within the last 6 months and c) chronic gouty arthropathy.

In one embodiment of any one of the methods or compositions or kits provided herein, the uricase is a pegylated uricase. In one embodiment of any one of the methods or compositions or kits provided herein, the pegylated uricase is pegsiticase or pegloticase. In
25 one embodiment of any one of the methods provided herein, the pegylated uricase is pegsiticase.

In one embodiment of any one of the methods or compositions or kits provided herein, the immunosuppressant is encapsulated in the synthetic nanocarriers.

In one embodiment of any one of the methods or compositions or kits provided
30 herein, the immunosuppressant is an mTOR inhibitor. In one embodiment of any one of the methods or compositions or kits provided herein, the mTOR inhibitor is a rapalog. In one embodiment of any one of the methods or compositions or kits provided herein, the rapalog is rapamycin.

-8-

In one embodiment of any one of the methods or compositions or kits provided herein, the synthetic nanocarriers are polymeric synthetic nanocarriers. In one embodiment of any one of the methods or compositions or kits provided herein, the polymeric synthetic nanocarriers comprise a hydrophobic polyester. In one embodiment of any one of the methods or compositions or kits provided herein, the hydrophobic polyester comprises PLA, PLG, PLGA or polycaprolactone. In one embodiment of any one of the methods or compositions or kits provided herein, the polymeric synthetic nanocarriers further comprise PEG. In one embodiment of any one of the methods or compositions or kits provided herein, the PEG is conjugated to the PLA, PLG, PLGA or polycaprolactone. In one embodiment of any one of the methods or compositions or kits provided herein, the polymeric synthetic nanocarriers comprise PLA, PLG, PLGA or polycaprolactone and PEG conjugated to PLA, PLG, PLGA or polycaprolactone. In one embodiment of any one of the methods or compositions or kits provided herein, the polymeric synthetic nanocarriers comprise PLA and PLA-PEG. In one embodiment of any one of the methods or compositions or kits provided herein, the synthetic nanocarriers are those as described according to or obtainable by any one of the exemplified methods provided herein.

In one embodiment of any one of the methods or compositions or kits provided herein, the mean of a particle size distribution obtained using dynamic light scattering of the synthetic nanocarriers is a diameter greater than 120nm. In one embodiment of any one of the methods or compositions or kits provided herein, the diameter is greater than 150nm. In one embodiment of any one of the methods or compositions or kits provided herein, the diameter is greater than 200nm. In one embodiment of any one of the methods or compositions or kits provided herein, the diameter is greater than 250nm. In one embodiment of any one of the methods or compositions or kits provided herein, the diameter is less than 300nm. In one embodiment of any one of the methods or compositions or kits provided herein, the diameter is less than 250nm. In one embodiment of any one of the methods or compositions or kits provided herein, the diameter is less than 200nm.

In one embodiment of any one of the methods or compositions or kits provided herein, the load of the immunosuppressant of the synthetic nanocarriers is 7-12% or 8-12% by weight. In one embodiment of any one of the methods or compositions or kits provided herein, the load of the immunosuppressant of the synthetic nanocarriers is 7-10% or 8-10% by weight. In one embodiment of any one of the methods or compositions or kits provided herein, the load of the immunosuppressant of the synthetic nanocarriers is 9-11% by weight.

-9-

In one embodiment of any one of the methods or compositions or kits provided herein, the load of the immunosuppressant of the synthetic nanocarriers is 7%, 8%, 9%, 10%, 11% or 12% by weight.

5 In one embodiment of any one of the methods provided herein, each administration is an intravenous administration. In one embodiment of any one of the methods provided herein, the intravenous administration is an intravenous infusion.

In one embodiment of any one of the methods provided herein, the method further comprises administering an additional therapeutic to the subject. In one embodiment of any one of the methods provided herein, the additional therapeutic is an anti-inflammatory
10 therapeutic, such as a corticosteroid. In one embodiment of any one of the methods provided herein, the additional therapeutic is a gout therapeutic, such as an oral gout therapeutic. In one embodiment of any one of the methods provided herein, the additional therapeutic is administered subsequently. In one embodiment of any one of the methods provided herein, the additional therapeutic is administered subsequent to the completion of treatment with the
15 concomitant administration of the uricase composition(s) and synthetic nanocarrier composition(s), such as according to any one of the regimens provided herein.

In one embodiment of any one of the methods provided herein, the additional therapeutic is an anti-gout flare treatment. In one embodiment of any one of the methods provided herein, the anti-gout flare treatment is a prophylactic treatment administered
20 concomitantly but prior to the administration of each uricase composition that is administered, such as according to any one of the regimens provided herein. In one embodiment of any one of the methods provided herein, the anti-gout flare treatment is colchicine or an NSAID.

In one embodiment of any one of the methods provided herein, the additional
25 therapeutic is a corticosteroid, and the corticosteroid is administered concomitantly, such as concomitantly prior to the administration of each uricase composition that is administered, such as according to any one of the regimens provided herein.

In another aspect, a method comprising administering to any of the subjects described herein a composition comprising uricase at any one of the doses, including label doses,
30 provided herein and a pharmaceutically acceptable carrier one or more times (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or 10 or more times). In some embodiments, the at least one administration or each administration is via a non-intramuscular mode of administration. In some examples, at least one administration or each administration is an intravenous administration, such as

-10-

intravenous infusion. In some embodiments, the composition comprising uricase and a pharmaceutically acceptable carrier is administered every two or four weeks. In some embodiments, the composition comprising uricase and a pharmaceutically acceptable carrier is administered monthly. In some embodiments, the composition comprising uricase and a pharmaceutically acceptable carrier is administered concomitantly with any one of the compositions comprising an immunosuppressant described herein.

In one aspect is a composition or kit comprising one or more compositions comprising uricase alone or in combination with one or more compositions comprising synthetic nanocarriers comprising an immunosuppressant. Each composition comprising uricase may be any one of the compositions comprising uricase as provided herein in any one of the compositions or kits. Each composition comprising uricase may be in an amount such that it provides any one or more doses, including label doses, of the uricase as provided herein in any one of the compositions or kits. Each composition comprising uricase may be in lyophilized form in any one of the compositions or kits. Each composition comprising synthetic nanocarriers comprising an immunosuppressant may be any one of the compositions comprising synthetic nanocarriers comprising an immunosuppressant as provided herein in any one of the compositions or kits. Each composition comprising synthetic nanocarriers comprising an immunosuppressant may be in an amount such that it provides any one or more doses, including label doses, of the synthetic nanocarriers comprising an immunosuppressant or immunosuppressant as provided herein in any one of the compositions or kits. Each composition comprising synthetic nanocarriers comprising an immunosuppressant may be in lyophilized form in any one of the compositions or kits. Each composition comprising synthetic nanocarriers comprising an immunosuppressant may be in a frozen suspension in any one of the compositions or kits. In one embodiment of any one of the compositions or kits, the frozen suspension further comprises phosphate-buffered saline (PBS). In one embodiment of any one of the compositions or kits, the lyophilized form further comprises PBS and/or mannitol. In one embodiment of any one of the compositions or kits, the composition or kit further comprises 0.9% sodium chloride, USP.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is an image showing tophi/uric acid deposits visualized using DECT.

Fig. 2 is a cartoon representation of the components of SEL-212.

-11-

Fig. 3 is a graph of ADA levels in non-human primates after treatment with empty nanocarriers + pegsiticase or pegsiticase + 0.1X or 1X synthetic nanocarriers comprising rapamycin (SVP-Rapamycin).

5 **Fig. 4** is a graph of mean serum uric acid (sUA) levels in the 5 cohorts of the phase 1a clinical trial following a single intravenous infusion of pegsiticase.

Fig. 5 is a graphical illustration showing the serum uric acid levels and uricase-specific ADA levels for each subject in Cohort #3 of the Phase 1a clinical trial and Cohort #9, Cohort # 4, and Cohort #6 in the Phase 1b clinical trial.

10 **Fig. 6** is a graph showing the serum uric acid levels of Cohort #3 from the Phase 1a clinical and Cohort #9, Cohort #1, Cohort #2, Cohort #3, Cohort #4, Cohort #5 and Cohort #6 from the Phase 1b clinical trial trial.

Fig. 7 from left to right shows data from two replicate Kystexxa® trials, in the middle is the data of SVP-Rapamycin alone vs. pegsiticase alone (Cohort #9) and then Rapamycin alone vs. Cohort #6 (a SEL-212 cohort).

15 **Fig. 8** is a graphical illustration showing the serum uric acid levels of subjects treated with pegsiticase alone, or in combination with synthetic nanocarriers comprising rapamycin (SVP-Rapamycin) (0.1 or 0.3 mg/kg).

Fig. 9 shows doses for the phase 2 clinical trial.

20 **Fig. 10** is a graph illustrating anti-uricase titers. In this instance, non-responding treatment periods had titers of 1080 or greater.

Fig. 11 is a graph illustrating anti-uricase titers. Those less than 1:1080 had increased sustained uricase activity.

25 **Figs. 12A-12B** shows results with five monthly doses of rapamycin-containing nanocarriers co-administered with 0.2 mg/kg of pegadricase. Fig. 12A shows different doses of rapamycin-containing nanocarriers, and Fig. 12B shows different combinations of rapamycin-containing nanocarrier concentration administrations.

30 **Figs. 13A-13D** show that increasing the anti-uricase titer results in lower uricase activity (Figs. 13A-13B), while increasing the rapamycin-containing nanocarrier concentration leads to higher sustained uricase activity (Figs. 13C-13D). In Figs. 13A and 13C, doses of 0.2 mg/kg of pegadricase were used; in Figs. 13B and 13D, doses of 0.4 mg/kg pegadricase were used.

-12-

DETAILED DESCRIPTION OF THE INVENTION**A. OVERVIEW**

Gout can be painful and disabling and is thought to result from excess uric acid.

5 Additionally, high concentrations of uric acid, such as serum uric acid, can increase the risk of co-morbidities, including cardiovascular, cardiometabolic, joint and kidney disease. There are approximately 8.3 million and 10 million gout sufferers in the United States and the European Union, respectively.

Based on studies and data, examples of which are provided above and elsewhere
10 herein, it has been demonstrated that the compositions and methods provided are substantially more efficacious than currently available treatments, can reduce undesired immune responses associated with the delivery of uricase, such as pegylated uricase, can provide strong and durable control of serum uric acid levels in patients, can provide for the removal of painful and damaging uric acid deposits for patients, such as with chronic
15 tophaceous gout, and/or can substantially reduce or eliminate the risk of gout flare that may occur with uric acid lowering therapies, such as uricase.

It has also been demonstrated that the use of a measure of anti-uricase antibodies in a subject can be helpful in identifying a subject for treatment or assessing the treatment of the subject. The measure of anti-uricase antibodies can be an anti-uricase titer in the subject.

20 The measure can be compared to a threshold and inform treatment decisions.

B. DEFINITIONS

“Additional therapeutic”, as used herein, refers to any therapeutic that is used in addition to another treatment. For example, when the method is one directed to treatment
25 with synthetic nanocarriers comprising an immunosuppressant, and the method comprises the use of an additional therapeutic, the additional therapeutic is in addition to synthetic nanocarriers comprising an immunosuppressant. As another example, when the method is one directed to treatment with a combination of a composition comprising a uricase and a composition comprising synthetic nanocarriers comprising an immunosuppressant, and the
30 method comprises the use of an additional therapeutic, the additional therapeutic is in addition to the uricase and synthetic nanocarrier composition combination. Generally, the additional therapeutic will be a different therapeutic. The additional therapeutic may be administered at the same time or at a different time and/or via the same mode of

-13-

administration or via a different mode of administration, as that of the other therapeutic. In preferred embodiments, the additional therapeutic will be given at a time and in a way that will provide a benefit to the subject during the effective treatment window of the other therapeutic. When two compositions are administered with a specific time period, generally the time period is measured from the start of the first composition to the start of the second composition. As used herein, when two compositions are given within an hour, for example, the time before the start of the administration of the first composition is about an hour before the start of the administration of the second composition.

In some embodiments, the additional therapeutic is another therapeutic for the treatment of gout or a condition associated with gout. As used herein, a “gout therapeutic” is any therapeutic that can be administered and from which a subject with gout may derive a benefit because of its administration. In some embodiments, the gout therapeutic is an oral gout therapeutic (i.e., a gout therapeutic that can be taken or given orally).

The additional therapeutic may be any one of the previously approved therapeutics described herein or otherwise known in the art. In some embodiments, the additional therapeutic is an uric acid lowering therapeutic. Such a therapeutic is any that results in a lower serum uric acid level in a subject as compared to a serum uric acid level in the subject without the administration of the therapeutic. Such uric acid lowering therapeutics include, uricases.

In some embodiments, the additional therapeutic is a therapeutic for preventing gout flare or also referred to herein as an anti-gout flare therapeutic. Any therapeutic that can be used to prevent a gout flare is included in this class of therapeutics. In some of these embodiments, the therapeutic for preventing gout flare is given prior to the administration of the other therapeutic. In some embodiments, the therapeutic for preventing gout flare is colchicine. In other embodiments, the therapeutic for preventing gout flare is an NSAID.

In an embodiment, any one of the methods for treating any one of the subjects or any one of the compositions or kits as provided herein can include the administration of an additional therapeutic or an additional therapeutic, respectively. In another embodiment, any one of the methods for treating any one of the subjects or any one of the compositions or kits as provided herein does not include the administration of an additional therapeutic, such as within the effective treatment window of the other therapeutic, or an additional therapeutic, respectively.

-14-

"Administering" or "administration" or "administer" means giving a material to a subject in a manner such that there is a pharmacological result in the subject. This may be direct or indirect administration, such as by inducing or directing another subject, including another clinician or the subject itself, to perform the administration.

5 "Amount effective" in the context of a composition or dose for administration to a subject refers to an amount of the composition or dose that produces one or more desired responses in the subject. In some embodiments, the amount effective is a pharmacodynamically effective amount. Therefore, in some embodiments, an amount effective is any amount of a composition or dose provided herein that produces one or more
10 of the desired therapeutic effects and/or immune responses as provided herein. This amount can be for in vitro or in vivo purposes. For in vivo purposes, the amount can be one that a clinician would believe may have a clinical benefit for a subject in need thereof. Any one of the compositions or doses, including label doses, as provided herein can be in an amount effective.

15 Amounts effective can involve reducing the level of an undesired response, although in some embodiments, it involves preventing an undesired response altogether. Amounts effective can also involve delaying the occurrence of an undesired response. An amount that is effective can also be an amount that produces a desired therapeutic endpoint or a desired therapeutic result. In other embodiments, the amounts effective can involve enhancing the
20 level of a desired response, such as a therapeutic endpoint or result. Amounts effective, preferably, result in a therapeutic result or endpoint and/or reduced or eliminated ADAs against the treatment and/or result in prevention of gout flare in any one of the subjects provided herein. The achievement of any of the foregoing can be monitored by routine methods.

25 Amounts effective will depend, of course, on the particular subject being treated; the severity of a condition, disease or disorder; the individual patient parameters including age, physical condition, size and weight; the duration of the treatment; the nature of concurrent therapy (if any); the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary
30 skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however,

-15-

that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reason.

Doses of the components in any one of the compositions of the invention or used in any one of the methods of the invention may refer to the amount of the components in the composition, the actual amounts of the respective components received by an administered
5 subject, or the amount that appears on a label (also referred to herein as label dose). The dose can be administered based on the number of synthetic nanocarriers that provide the desired amount of the component(s).

“Attach” or “Attached” or “Couple” or “Coupled” (and the like) means to chemically
10 associate one entity (for example a moiety) with another. In some embodiments, the attaching is covalent, meaning that the attachment occurs in the context of the presence of a covalent bond between the two entities. In non-covalent embodiments, the non-covalent attaching is mediated by non-covalent interactions including but not limited to charge interactions, affinity interactions, metal coordination, physical adsorption, host-guest
15 interactions, hydrophobic interactions, π - π stacking interactions, hydrogen bonding interactions, van der Waals interactions, magnetic interactions, electrostatic interactions, dipole-dipole interactions, and/or combinations thereof. In embodiments, encapsulation is a form of attaching.

“Average”, as used herein, refers to the arithmetic mean unless otherwise noted.

“Concomitantly” means administering two or more materials/agents to a subject in a
20 manner that is correlated in time, preferably sufficiently correlated in time so as to provide a modulation in a physiologic or immunologic response, and even more preferably the two or more materials/agents are administered in combination. In embodiments, concomitant administration may encompass administration of two or more materials/agents within a
25 specified period of time, preferably within 1 month, more preferably within 1 week, still more preferably within 1 day, and even more preferably within 1 hour. In embodiments, the two or more materials/agents are sequentially administered. In embodiments, the materials/agents may be repeatedly administered concomitantly; that is concomitant administration on more than one occasion.

“Dose” refers to a specific quantity of a pharmacologically active material for
30 administration to a subject for a given time. Unless otherwise specified, the doses recited for compositions comprising pegylated uricase refer to the weight of the uricase (i.e., the protein without the weight of the PEG or any other components of the composition comprising the

-16-

pegylated uricase). Also, unless otherwise specified, the doses recited for compositions comprising synthetic nanocarriers comprising an immunosuppressant refer to the weight of the immunosuppressant (i.e., without the weight of the synthetic nanocarrier material or any of the other components of the synthetic nanocarrier composition). When referring to a dose for administration, in an embodiment of any one of the methods, compositions or kits provided herein, any one of the doses provided herein is the dose as it appears on a label/label dose.

“Encapsulate” means to enclose at least a portion of a substance within a synthetic nanocarrier. In some embodiments, a substance is enclosed completely within a synthetic nanocarrier. In other embodiments, most or all of a substance that is encapsulated is not exposed to the local environment external to the synthetic nanocarrier. In other embodiments, no more than 50%, 40%, 30%, 20%, 10% or 5% (weight/weight) is exposed to the local environment. Encapsulation is distinct from absorption, which places most or all of a substance on a surface of a synthetic nanocarrier, and leaves the substance exposed to the local environment external to the synthetic nanocarrier. In embodiments of any one of the methods or compositions provided herein, the immunosuppressants are encapsulated within the synthetic nanocarriers.

“Elevated serum uric acid level” refers to any level of uric acid in a subject’s serum that may lead to an undesirable result or would be deemed by a clinician to be elevated. In an embodiment, the subject of any one of the methods provided herein can have a serum uric acid level of ≥ 5 mg/dL, ≥ 6 mg/dL, or ≥ 7 mg/dL. Such a subject may be a hyperuremic subject. Whether or not a subject has elevated blood uric acid levels can be determined by a clinician, and in some embodiments, the subject is one in which a clinician has identified or would identify as having elevated serum uric acid levels.

“Gout” generally refers to a disorder or condition associated with the buildup of uric acid, such as deposition of uric crystals in tissues and joints, and/or a clinically relevant elevated serum uric acid level. Accumulation of uric acid may be due to overproduction of uric acid or reduced excretion of uric acid. Gout may range from asymptomatic to severe and painful inflammatory conditions. A “condition associated with gout” refers to any condition in a subject where the subject experiences local and/or systemic effects of gout, including inflammation and immune responses, and in which the condition is caused or exacerbated by, or the condition can result in or exacerbate, gout. A gout flare is an “attack” or exacerbation

-17-

of gout symptoms, which can happen at any time. Gout flares can include gout flares that occur after the administration of a uric acid lowering therapy.

“Hydrophobic polyester” refers to any polymer that comprises one or more polyester polymers or units thereof and that has hydrophobic characteristics. Polyester polymers include, but are not limited to, PLA, PLGA, PLG and polycaprolactone. “Hydrophobic” refers to a material that does not substantially participate in hydrogen bonding to water. Such materials are generally non-polar, primarily non-polar, or neutral in charge. Synthetic nanocarriers may be completely comprised of hydrophobic polyesters or units thereof. In some embodiments, however, the synthetic nanocarriers comprise hydrophobic polyesters or units thereof in combination with other polymers or units thereof. These other polymers or units thereof may be hydrophobic but are not necessarily so. In some preferred embodiments, when synthetic nanocarriers include one or more other polymers or units thereof in addition to a hydrophobic polyester, the matrix of other polymers or units thereof with the hydrophobic polyester is hydrophobic overall. Examples of synthetic nanocarriers that can be used in the invention and that comprise hydrophobic polyesters can be found in U.S. Publication Nos. US 2016/0128986 and US 2016/0128987, and such synthetic nanocarriers and the disclosure of such synthetic nanocarriers is incorporated herein by reference.

“Immunosuppressant”, as used herein, means a compound that can cause a tolerogenic immune response specific to an antigen, also referred to herein as an “immunosuppressive effect”. An immunosuppressive effect generally refers to the production or expression of cytokines or other factors by an antigen-presenting cell (APC) that reduces, inhibits or prevents an undesired immune response or that promotes a desired immune response, such as a regulatory immune response, against a specific antigen. When the APC acquires an immunosuppressive function (under the immunosuppressive effect) on immune cells that recognize an antigen presented by this APC, the immunosuppressive effect is said to be specific to the presented antigen. Examples of immunosuppressants include “mTOR inhibitors”, a class of drugs that inhibit mTOR, a serine/threonine-specific protein kinase that belongs to the family of phosphatidylinositol-3 kinase (PI3K) related kinases (PIKKs). mTOR inhibitors include, but are not limited to, rapalogs, such as rapamycin, as well as ATP-competitive mTOR kinase inhibitors, such as mTORC1/mTORC2 dual inhibitors.

In embodiments of any one of the methods, compositions or kits provided herein, the immunosuppressants provided herein are attached to synthetic nanocarriers. In preferable embodiments, the immunosuppressant is an element that is in addition to the material that makes up the structure of the synthetic nanocarrier. For example, in one embodiment, where the synthetic nanocarrier is made up of one or more polymers, the immunosuppressant is a compound that is in addition and attached to the one or more polymers. In embodiments, such as where the material of the synthetic nanocarrier also results in an immunosuppressive effect, the immunosuppressant is an element present in addition to the material of the synthetic nanocarrier that results in an immunosuppressive effect.

“Load”, when comprise in a composition comprising a synthetic nanocarrier, such as coupled thereto, is the amount of the immunosuppressant in the composition based on the total dry recipe weight of materials in an entire synthetic nanocarrier (weight/weight). Generally, such a load is calculated as an average across a population of synthetic nanocarriers. In one embodiment, the load on average across the synthetic nanocarriers is between 0.1% and 15%. In another embodiment, the load is between 0.1% and 10%. In a further embodiment, the load is between 1% and 15%. In yet a further embodiment, the load is between 5% and 15%. In still a further embodiment, the load is between 7% and 12%. In still a further embodiment, the load is between 8% and 12%. In still another embodiment, the load is between 7% and 10%. In still another embodiment, the load is between 8% and 10%. In yet a further embodiment, the load is 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, or 15% on average across the population of synthetic nanocarriers. In any one of the methods, compositions or kits provided herein, the load of the immunosuppressant, such as rapamycin, may be any one of the loads provided herein.

The rapamycin load of the nanocarrier in suspension is calculated by dividing the rapamycin content of the nanocarrier as determined by HPLC analysis of the test article by the nanocarrier mass. The total polymer content is measured either by gravimetric yield of the dry nanocarrier mass or by the determination of the nanocarrier solution total organic content following pharmacopeia methods and corrected for PVA content.

“Maximum dimension of a synthetic nanocarrier” means the largest dimension of a nanocarrier measured along any axis of the synthetic nanocarrier. “Minimum dimension of a synthetic nanocarrier” means the smallest dimension of a synthetic nanocarrier measured along any axis of the synthetic nanocarrier. For example, for a spheroidal synthetic nanocarrier, the maximum and minimum dimension of a synthetic nanocarrier would be

-19-

substantially identical, and would be the size of its diameter. Similarly, for a cuboidal synthetic nanocarrier, the minimum dimension of a synthetic nanocarrier would be the smallest of its height, width or length, while the maximum dimension of a synthetic nanocarrier would be the largest of its height, width or length. In an embodiment, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or greater than 100 nm. In an embodiment, a maximum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or less than 5 μm . Preferably, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is greater than 110 nm, more preferably greater than 120 nm, more preferably greater than 130 nm, and more preferably still greater than 150 nm. Aspects ratios of the maximum and minimum dimensions of synthetic nanocarriers may vary depending on the embodiment. For instance, aspect ratios of the maximum to minimum dimensions of the synthetic nanocarriers may vary from 1:1 to 1,000,000:1, preferably from 1:1 to 100,000:1, more preferably from 1:1 to 10,000:1, more preferably from 1:1 to 1000:1, still more preferably from 1:1 to 100:1, and yet more preferably from 1:1 to 10:1.

Preferably, a maximum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample is equal to or less than 3 μm , more preferably equal to or less than 2 μm , more preferably equal to or less than 1 μm , more preferably equal to or less than 800 nm, more preferably equal to or less than 600 nm, and more preferably still equal to or less than 500 nm. In preferred embodiments, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or greater than 100 nm, more preferably equal to or greater than 120 nm, more preferably equal to or greater than 130 nm, more preferably equal to or greater than 140 nm, and more preferably still equal to or greater than 150 nm. Measurement of synthetic nanocarrier dimensions (*e.g.*, effective diameter) may be obtained, in some embodiments, by suspending the synthetic nanocarriers in a liquid (usually aqueous) media and using dynamic light scattering (DLS) (*e.g.*, using a Brookhaven ZetaPALS instrument). For example, a

-20-

suspension of synthetic nanocarriers can be diluted from an aqueous buffer into purified water to achieve a final synthetic nanocarrier suspension concentration of approximately 0.01 to 0.5 mg/mL. The diluted suspension may be prepared directly inside, or transferred to, a suitable cuvette for DLS analysis. The cuvette may then be placed in the DLS, allowed to
5 equilibrate to the controlled temperature, and then scanned for sufficient time to acquire a stable and reproducible distribution based on appropriate inputs for viscosity of the medium and refractive indices of the sample. The effective diameter, or mean of the distribution, is then reported. Determining the effective sizes of high aspect ratio, or non-spheroidal, synthetic nanocarriers may require augmentative techniques, such as electron microscopy, to
10 obtain more accurate measurements. “Dimension” or “size” or “diameter” of synthetic nanocarriers means the mean of a particle size distribution, for example, obtained using dynamic light scattering.

“Obtaining” as used herein refers to any method by which the respective information or materials can be acquired. Thus, the respective information can be acquired by
15 experimental methods. Respective materials can be created, designed, etc. with various experimental or laboratory methods, in some embodiments. The respective information or materials can also be acquired by being given or provided with the information, such as in a report, or materials.

“Pegylated uricase” refers to any uricase that is attached to one or more PEG
20 (poly(ethylene glycol), poly (ethylene oxide) or poly (oxyethylene)) molecules (i.e, a poly(ethylene glycol), poly (ethylene oxide) or poly (oxyethylene) polymer or unit thereof). Preferably in some embodiments, the one or more PEG molecules are poly(ethylene glycol) molecules. The terms “pegylated” or “pegylation” refer to the conjugated form or the act of conjugating to the uricase, respectively. Such a modified uricase is referred to as pegylated
25 uricase. Pegylated uricases include, but are not limited to pegsiticase and pegloticase (KRYSTEXXA®).

“Pharmaceutically acceptable excipient” or “pharmaceutically acceptable carrier” means a pharmacologically inactive material used together with a pharmacologically active material to formulate the compositions. Pharmaceutically acceptable excipients comprise a
30 variety of materials known in the art, including but not limited to saccharides (such as glucose, lactose, and the like), preservatives such as antimicrobial agents, reconstitution aids, colorants, saline (such as phosphate buffered saline), and buffers. Any one of the compositions provided herein may include a pharmaceutically acceptable excipient or carrier.

-21-

“Rapalog” refers to rapamycin and molecules that are structurally related to (an analog) of rapamycin (sirolimus), and are preferably hydrophobic. Examples of rapalogs include, without limitation, temsirolimus (CCI-779), deforolimus, everolimus (RAD001), ridaforolimus (AP-23573), zotarolimus (ABT-578). Additional examples of rapalogs may be found, for example, in WO Publication WO 1998/002441 and U.S. Patent No. 8,455,510, the disclosure of such rapalogs are incorporated herein by reference in its entirety. In any one of the methods or compositions or kits provided herein, the immunosuppressant may be a rapalog.

“Subject” means animals, including warm blooded mammals such as humans and primates; avians; domestic household or farm animals such as cats, dogs, sheep, goats, cattle, horses and pigs; laboratory animals such as mice, rats and guinea pigs; fish; reptiles; zoo and wild animals; and the like. In any one of the methods, compositions and kits provided herein, the subject is human. In any one of the methods, compositions and kits provided herein, the subject is any one of the subjects provided herein, such as one that has any one of the conditions provided herein, such as gout or other condition associated with gout.

“Synthetic nanocarrier(s)” means a discrete object that is not found in nature, and that possesses at least one dimension that is less than or equal to 5 microns in size. Synthetic nanocarriers may be a variety of different shapes, including but not limited to spheroidal, cuboidal, pyramidal, oblong, cylindrical, toroidal, and the like. Synthetic nanocarriers comprise one or more surfaces.

A synthetic nanocarrier can be, but is not limited to, one or a plurality of lipid-based nanoparticles (also referred to herein as lipid nanoparticles, i.e., nanoparticles where the majority of the material that makes up their structure are lipids), polymeric nanoparticles, metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles (i.e., particles that are primarily made up of viral structural proteins but that are not infectious or have low infectivity), peptide or protein-based particles (also referred to herein as protein particles, i.e., particles where the majority of the material that makes up their structure are peptides or proteins) (such as albumin nanoparticles) and/or nanoparticles that are developed using a combination of nanomaterials such as lipid-polymer nanoparticles. Synthetic nanocarriers may be a variety of different shapes, including but not limited to spheroidal, cuboidal, pyramidal, oblong, cylindrical, toroidal, and the like. Examples of synthetic nanocarriers include (1) the biodegradable nanoparticles disclosed in US Patent 5,543,158 to Gref et al., (2) the polymeric nanoparticles of Published US Patent Application

-22-

20060002852 to Saltzman et al., (3) the lithographically constructed nanoparticles of
Published US Patent Application 20090028910 to DeSimone et al., (4) the disclosure of WO
2009/051837 to von Andrian et al., (5) the nanoparticles disclosed in Published US Patent
Application 2008/0145441 to Penades et al., (6) the nanoprecipitated nanoparticles disclosed
5 in P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently
Associate and Deliver Virus-like Particles" *Nanomedicine*. 5(6):843-853 (2010), and (7)
those of Look et al., "Nanogel-based delivery of mycophenolic acid ameliorates systemic
lupus erythematosus in mice" *J. Clinical Investigation* 123(4):1741-1749(2013).

Synthetic nanocarriers may have a minimum dimension of equal to or less than about
10 100 nm, preferably equal to or less than 100 nm, do not comprise a surface with hydroxyl
groups that activate complement or alternatively comprise a surface that consists essentially
of moieties that are not hydroxyl groups that activate complement. In an embodiment,
synthetic nanocarriers that have a minimum dimension of equal to or less than about 100 nm,
preferably equal to or less than 100 nm, do not comprise a surface that substantially activates
15 complement or alternatively comprise a surface that consists essentially of moieties that do
not substantially activate complement. In a more preferred embodiment, synthetic
nanocarriers according to the invention that have a minimum dimension of equal to or less
than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface that
activates complement or alternatively comprise a surface that consists essentially of moieties
20 that do not activate complement. In embodiments, synthetic nanocarriers exclude virus-like
particles. In embodiments, synthetic nanocarriers may possess an aspect ratio greater than
1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7, or greater than 1:10.

"Treating" refers to the administration of one or more therapeutics with the
expectation that the subject may have a resulting benefit due to the administration. The
25 treating may also result in the prevention of a condition as provided herein and, therefore,
treating includes prophylactic treatment. When used prophylactically, the subject is one in
which a clinician expects that there is a likelihood for the development of a condition or other
undesired response as provided herein. In some embodiments, a subject that is expected to
have a gout flare is one in which a clinician believes there is a likelihood that a gout flare will
30 occur. Treating may be direct or indirect, such as by inducing or directing another subject,
including another clinician or the subject itself, to treat the subject.

"Weight%" or "% by weight" refers to the ratio of one weight to another weight
times 100. For example, the weight% can be the ratio of the weight of one component to

-23-

another times 100 or the ratio of the weight of one component to a total weight of more than one component times 100. Generally, the weight% is measured as an average across a population of synthetic nanocarriers or an average across the synthetic nanocarriers in a composition or suspension.

5

C. METHODS AND RELATED COMPOSITIONS

Uricase and Pegylated Uricase

The methods and compositions and kits described herein involve compositions comprising uricase. Uricase is generally thought to catalyze the conversion of uric acid to allantoin, which is soluble and may be excreted. Uricase is an enzyme endogenous to all mammals, except for humans and certain primates. The gene encoding the uricase enzyme may be obtained from any source known in the art, including mammalian and microbial sources as well as by recombinant and synthetic technologies. As will be evident to one of ordinary skill in the art, a gene may be obtained from a source and recombinantly (or transgenically) expressed and produced in another organism using standard methods. See Erlich, H A, (Ed.) (1989) PCR Technology. Principles and Applications for DNA Amplification. New York: Stockton Press; Sambrook, J, et al., (1989) Molecular Cloning. A Laboratory Manual, Second Edition. Cold Spring Harbor, N.Y.: Cold Spring Harbor Laboratory Press. For example, U.S. Patent No. 5,700,674 describes recombinant production of uricase in *E. coli* cells. In some embodiments, the enzyme is produced by fermentation in *E. coli*.

In some embodiments, the gene encoding the uricase, or a portion thereof, is obtained from a mammal, for example a pig, bovine, sheep, goat, baboon, monkey mouse, rabbit, or domestic animal. In some embodiments, the gene encoding the uricase, or a portion thereof, is obtained from a microorganism, such as a bacteria or fungi (including yeast). In some embodiments, the gene encoding the uricase is obtained from a bacterial source, such as bacterium belonging to *Streptomyces* spp., *Bacillus* spp., or *E. coli*. In some embodiments, the gene encoding the uricase is obtained from a fungal (including yeast) source, such as *Candida* (e.g., *Candida utilis*), *Anthrobacter* (e.g., *Anthrobacter globiformis*), *Saccharomyces*, *Schizosaccharomyces*, *Emericella*, *Aspergillus* (e.g., *Aspergillus flavus*), and *Neurospora* spp. In some embodiments, the uricase is derived from *Candida utilis*. In some embodiments, the uricase is that of pegsiticase (3SBio as described in U.S. Patent No. 6,913,915, and such

-24-

uricase and description thereof is incorporated herein by reference). In some embodiments, the uricase is derived from *Aspergillus flavus*. In some embodiments, the uricase is rasburicase (ELITEK®; FASTURTEC®, from Sanofi Genzyme).

In some embodiments, the uricase is chimeric uricase, in which portions of the gene encoding the uricase are obtained from different sources. For example, a portion of the gene encoding the chimeric uricase may be obtained from one organism and one or more other portions of the gene encoding the chimeric uricase may be obtained from another organism. In some embodiments, a portion of the gene encoding the chimeric uricase is obtained from a pig and another portion of the gene encoding the chimeric uricase is obtained from a baboon. In some embodiments, the chimeric uricase is that of pegloticase/KRYSTEXXA®.

Also within the scope of the present invention are variant uricases, which may include one or more mutations (substitutions, insertions, deletions). Mutations may be made in the nucleotide sequence encoding the uricase protein, which may or may not result in an amino acid mutation. In general, mutations may be made, for example, to enhance production of the protein, turnover/half-life of the protein or mRNA encoding the protein, modulate (enhance or reduce) the enzymatic activity of the uricase.

In other embodiments, the gene encoding the uricase is obtained from a plant or invertebrate source, such as *Drosophila* or *C. elegans*.

Any of the uricase proteins described herein may be pegylated. Uricase may be covalently bonded to PEG via a biocompatible linking group, using methods known in the art, as described, for example, by Park et al, *Anticancer Res.*, 1:373-376 (1981); and Zaplisky and Lee, *Polyethylene Glycol Chemistry: Biotechnical and Biomedical Applications*, J. M. Harris, ed., Plenum Press, New York, Chapter 21 (1992). The linking group used to covalently attach PEG to uricase may be any biocompatible linking group, meaning the linking group non-toxic and may be utilized in vitro or in vivo without causing adverse effects. Alternatively, the PEG may be directly conjugated to the uricase, such as directly to a lysine residue of uricase.

Uricase may be pegylated at many different amino acid residues of the uricase protein. The number of PEG molecules and/or residue to which the PEG is conjugated may affect the activity of the uricase. In some embodiments, the pegylated uricase comprises at least one PEG molecule. In some embodiments, the pegylated uricase comprises at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 45, 50, or more PEG molecules on average per uricase protein.

-25-

In some embodiments, the pegylated uricase comprises about 20-25 PEG molecules per uricase protein.

On average, PEG has a molecular weight between 5 kDa to 100 kDa. Both the molecular weight (size) of the PEG used as well as the number of PEG molecules used to pegylate the uricase may be varied. In some embodiment the average molecular weight of the PEG is between 5 kDa to 100 kDa, 5 kDa to 75 kDa, 5 kDa to 50 kDa, 5 kDa to 30 kDa, 5 kDa to 20 kDa, 5 kDa to 10 kDa, 10 kDa to 75 kDa, 10 kDa to 50 kDa, 10 kDa to 30 kDa, 5 kDa to 30 kDa, 15 kDa to 50 kDa, 15 kDa to 30 kDa, 15 kDa to 25 kDa, 20 kDa to 75 kDa, 30 kDa to 80 kDa, 30 kDa to 70 kDa, or 30 kDa to 50 kDa. In some embodiments, the molecular weight of the PEG is about 5 kDa, 6 kDa, 7 kDa, 8 kDa, 9 kDa, 10 kDa, 11 kDa, 12 kDa, 13 kDa, 14 kDa, 15 kDa, 16 kDa, 17 kDa, 18 kDa, 19 kDa, 20 kDa, 21 kDa, 22 kDa, 23 kDa, 24 kDa, 25 kDa, 30 kDa, 35 kDa, 40 kDa, 45 kDa, 50 kDa, 55 kDa, 60 kDa, 65 kDa, 70 kDa, 75 kDa, 80 kDa, 85 kDa, 90 kDa, 95 kDa, or 100 kDa. In general, the PEG is referred to based on the molecular weight of the PEG. For example, PEG-20 refers to PEG molecules with a molecular weight of 20 kDa, and PEG-5 refers to PEG molecules with a molecular weight of 5 kDa. In some embodiments, the uricase is pegylated with PEG molecules having a molecule weight of 20 kDa (PEG-20).

Pegylated uricases include, without limitation, pegsiticase (available from 3Sbio, and as described in U.S. Patent No. 6,913,915, and such pegylated uricase and description thereof is incorporated herein by reference) and pegloticase/KRYSTEXXA® (Horizon Pharmaceuticals).

Preferably, in some embodiments of any one of the methods or compositions or kits provided herein, the pegylated uricase is pegsiticase, a recombinant uricase conjugated to multiple 20 kDa molecular weight poly (ethylene glycol) molecules. The uricase component of pegsiticase can be cloned from the yeast *Candida utilis* and expressed in *E. coli* for production.

The uric acid catalysis activity of uricase, including pegylated uricase, can be assessed using methods known in the art or as otherwise provided herein.

Synthetic Nanocarriers

A variety of synthetic nanocarriers can be used. In some embodiments, synthetic nanocarriers are spheres or spheroids. In some embodiments, synthetic nanocarriers are flat or plate-shaped. In some embodiments, synthetic nanocarriers are cubes or cubic. In some

-26-

embodiments, synthetic nanocarriers are ovals or ellipses. In some embodiments, synthetic nanocarriers are cylinders, cones, or pyramids.

In some embodiments, it is desirable to use a population of synthetic nanocarriers that is relatively uniform in terms of size or shape so that each synthetic nanocarrier has similar properties. For example, at least 80%, at least 90%, or at least 95% of the synthetic nanocarriers, based on the total number of synthetic nanocarriers, may have a minimum dimension or maximum dimension that falls within 5%, 10%, or 20% of the average diameter or average dimension of the synthetic nanocarriers.

Synthetic nanocarriers can be solid or hollow and can comprise one or more layers. In some embodiments, each layer has a unique composition and unique properties relative to the other layer(s). To give but one example, synthetic nanocarriers may have a core/shell structure, wherein the core is one layer (e.g. a polymeric core) and the shell is a second layer (e.g. a lipid bilayer or monolayer). Synthetic nanocarriers may comprise a plurality of different layers.

In preferred embodiments, the synthetic nanocarriers comprise a polymer as provided herein. Polymers may be natural or unnatural (synthetic) polymers. Polymers may be homopolymers or copolymers comprising two or more monomers. In terms of sequence, copolymers may be random, block, or comprise a combination of random and block sequences. Typically, polymers in accordance with the present invention are organic polymers.

The synthetic nanocarriers as provided herein, preferably, comprise hydrophobic polyesters. Such polyesters can include copolymers comprising lactic acid and glycolic acid units, such as poly(lactic acid-co-glycolic acid) and poly(lactide-co-glycolide), collectively referred to herein as "PLGA"; and homopolymers comprising glycolic acid units, referred to herein as "PGA," and lactic acid units, such as poly-L-lactic acid, poly-D-lactic acid, poly-D,L-lactic acid, poly-L-lactide, poly-D-lactide, and poly-D,L-lactide, collectively referred to herein as "PLA." In some embodiments, exemplary polyesters include, for example, polyhydroxyacids; PEG copolymers and copolymers of lactide and glycolide (e.g., PLA-PEG copolymers, PGA-PEG copolymers, PLGA-PEG copolymers, and derivatives thereof. In some embodiments, polyesters include, for example, poly(caprolactone), poly(caprolactone)-PEG copolymers, poly(L-lactide-co-L-lysine), poly(serine ester), poly(4-hydroxy-L-proline ester), poly[α -(4-aminobutyl)-L-glycolic acid], and derivatives thereof.

-27-

In some embodiments, the polyester may be PLGA. PLGA is a biocompatible and biodegradable co-polymer of lactic acid and glycolic acid, and various forms of PLGA are characterized by the ratio of lactic acid:glycolic acid. Lactic acid can be L-lactic acid, D-lactic acid, or D,L-lactic acid. The degradation rate of PLGA can be adjusted by altering the lactic acid:glycolic acid ratio. In some embodiments, PLGA to be used in accordance with the present invention is characterized by a lactic acid:glycolic acid ratio of approximately 85:15, approximately 75:25, approximately 60:40, approximately 50:50, approximately 40:60, approximately 25:75, or approximately 15:85.

The synthetic nanocarriers may comprise one or more non-polyester polymers or units thereof that are also hydrophobic and/or polymers or units thereof that are not hydrophobic. In some embodiments, it is preferred that overall the synthetic nanocarrier comprises a hydrophobic polyester and, in some embodiments, is itself hydrophobic.

The synthetic nanocarriers may comprise one or more polymers that are a non-methoxy-terminated, pluronic polymer, or a unit thereof. "Non-methoxy-terminated polymer" means a polymer that has at least one terminus that ends with a moiety other than methoxy. In some embodiments, the polymer has at least two termini that ends with a moiety other than methoxy. In other embodiments, the polymer has no termini that ends with methoxy. "Non-methoxy-terminated, pluronic polymer" means a polymer other than a linear pluronic polymer with methoxy at both termini.

The synthetic nanocarriers may comprise, in some embodiments, polyhydroxyalkanoates, polyamides, polyethers, polyolefins, polyacrylates, polycarbonates, polystyrene, silicones, fluoropolymers, or a unit thereof. Further examples of polymers that may be comprised in the synthetic nanocarriers provided herein include polycarbonate, polyamide, or polyether, or unit thereof. In other embodiments, the polymers of the synthetic nanocarriers may comprise poly(ethylene glycol) (PEG), polypropylene glycol, or unit thereof.

In some embodiments, it is preferred that the synthetic nanocarriers comprise polymer that is biodegradable. Therefore, in such embodiments, the polymers of the synthetic nanocarriers may include a polyether, such as poly(ethylene glycol) or polypropylene glycol or unit thereof. Additionally, the polymer may comprise a block-co-polymer of a polyether and a biodegradable polymer such that the polymer is biodegradable. In other embodiments, the polymer does not solely comprise a polyether or unit thereof, such as poly(ethylene glycol) or polypropylene glycol or unit thereof.

-28-

In some embodiments, polymers in accordance with the present invention include polymers which have been approved for use in humans by the U.S. Food and Drug Administration (FDA) under 21 C.F.R. § 177.2600.

Other examples of polymers suitable for use in synthetic nanocarriers include, but are not limited to polyethylenes, polycarbonates (e.g. poly(1,3-dioxan-2-one)), polyanhydrides (e.g. poly(sebacic anhydride)), polypropylfumerates, polyamides (e.g. polycaprolactam), polyacetals, polyethers, polyesters (e.g., polylactide, polyglycolide, polylactide-co-glycolide, polycaprolactone, polyhydroxyacid (e.g. poly(β -hydroxyalkanoate))), poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polyureas, polystyrenes, and polyamines, polylysine, polylysine-PEG copolymers, and poly(ethyleneimine), poly(ethylene imine)-PEG copolymers.

Still other examples of polymers that may be included in the synthetic nanocarriers include acrylic polymers, for example, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, aminoalkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), methacrylic acid alkylamide copolymer, poly(methyl methacrylate), poly(methacrylic acid anhydride), methyl methacrylate, polymethacrylate, poly(methyl methacrylate) copolymer, polyacrylamide, aminoalkyl methacrylate copolymer, glycidyl methacrylate copolymers, polycyanoacrylates, and combinations comprising one or more of the foregoing polymers.

In some embodiments, the polymers of a synthetic nanocarrier associate to form a polymeric matrix. A wide variety of polymers and methods for forming polymeric matrices therefrom are known conventionally. In some embodiments, a synthetic nanocarrier comprising a hydrophobic polyester has a hydrophobic environment within the synthetic nanocarrier.

In some embodiments, polymers may be modified with one or more moieties and/or functional groups. A variety of moieties or functional groups can be used in accordance with the present invention. In some embodiments, polymers may be modified with polyethylene glycol (PEG), with a carbohydrate, and/or with acyclic polyacetals derived from polysaccharides (Papisov, 2001, ACS Symposium Series, 786:301). Certain embodiments may be made using the general teachings of US Patent No. 5543158 to Gref et al., or WO publication WO2009/051837 by Von Andrian et al.

In some embodiments, polymers may be modified with a lipid or fatty acid group. In some embodiments, a fatty acid group may be one or more of butyric, caproic, caprylic,

capric, lauric, myristic, palmitic, stearic, arachidic, behenic, or lignoceric acid. In some embodiments, a fatty acid group may be one or more of palmitoleic, oleic, vaccenic, linoleic, alpha-linoleic, gamma-linoleic, arachidonic, gadoleic, arachidonic, eicosapentaenoic, docosahexaenoic, or erucic acid.

5 In some embodiments, polymers can be linear or branched polymers. In some embodiments, polymers can be dendrimers. In some embodiments, polymers can be substantially cross-linked to one another. In some embodiments, polymers can be substantially free of cross-links. In some embodiments, polymers can be used in accordance with the present invention without undergoing a cross-linking step. It is further to be
10 understood that the synthetic nanocarriers may comprise block copolymers, graft copolymers, blends, mixtures, and/or adducts of any of the foregoing and other polymers. Those skilled in the art will recognize that the polymers listed herein represent an exemplary, not comprehensive, list of polymers that can be of use in accordance with the present invention provided they meet the desired criteria.

15 The properties of these and other polymers and methods for preparing them are well known in the art (see, for example, U.S. Patents 6,123,727; 5,804,178; 5,770,417; 5,736,372; 5,716,404; 6,095,148; 5,837,752; 5,902,599; 5,696,175; 5,514,378; 5,512,600; 5,399,665; 5,019,379; 5,010,167; 4,806,621; 4,638,045; and 4,946,929; Wang et al., 2001, J. Am. Chem. Soc., 123:9480; Lim et al., 2001, J. Am. Chem. Soc., 123:2460; Langer, 2000, Acc. Chem.
20 Res., 33:94; Langer, 1999, J. Control. Release, 62:7; and Uhrich et al., 1999, Chem. Rev., 99:3181). More generally, a variety of methods for synthesizing certain suitable polymers are described in Concise Encyclopedia of Polymer Science and Polymeric Amines and Ammonium Salts, Ed. by Goethals, Pergamon Press, 1980; Principles of Polymerization by Odian, John Wiley & Sons, Fourth Edition, 2004; Contemporary Polymer Chemistry by
25 Allcock et al., Prentice-Hall, 1981; Deming et al., 1997, Nature, 390:386; and in U.S. Patents 6,506,577, 6,632,922, 6,686,446, and 6,818,732.

Synthetic nanocarriers may be prepared using a wide variety of methods known in the art. For example, synthetic nanocarriers can be formed by methods such as nanoprecipitation, flow focusing using fluidic channels, spray drying, single and double
30 emulsion solvent evaporation, solvent extraction, phase separation, milling (including cryomilling), supercritical fluid (such as supercritical carbon dioxide) processing, microemulsion procedures, microfabrication, nanofabrication, sacrificial layers, simple and complex coacervation, and other methods well known to those of ordinary skill in the art.

-30-

Alternatively or additionally, aqueous and organic solvent syntheses for monodisperse semiconductor, conductive, magnetic, organic, and other nanomaterials have been described (Pellegrino et al., 2005, *Small*, 1:48; Murray et al., 2000, *Ann. Rev. Mat. Sci.*, 30:545; and Trindade et al., 2001, *Chem. Mat.*, 13:3843). Additional methods have been described in the literature (see, e.g., Doubrow, Ed., "Microcapsules and Nanoparticles in Medicine and Pharmacy," CRC Press, Boca Raton, 1992; Mathiowitz et al., 1987, *J. Control. Release*, 5:13; Mathiowitz et al., 1987, *Reactive Polymers*, 6:275; and Mathiowitz et al., 1988, *J. Appl. Polymer Sci.*, 35:755; US Patents 5578325 and 6007845; P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" *Nanomedicine*. 5(6):843-853 (2010)).

Immunosuppressants may be encapsulated into synthetic nanocarriers as desirable using a variety of methods including but not limited to C. Astete et al., "Synthesis and characterization of PLGA nanoparticles" *J. Biomater. Sci. Polymer Edn*, Vol. 17, No. 3, pp. 247-289 (2006); K. Avgoustakis "Pegylated Poly(Lactide) and Poly(Lactide-Co-Glycolide) Nanoparticles: Preparation, Properties and Possible Applications in Drug Delivery" *Current Drug Delivery* 1:321-333 (2004); C. Reis et al., "Nanoencapsulation I. Methods for preparation of drug-loaded polymeric nanoparticles" *Nanomedicine* 2:8- 21 (2006); P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" *Nanomedicine*. 5(6):843-853 (2010). Other methods suitable for encapsulating materials into synthetic nanocarriers may be used, including without limitation methods disclosed in United States Patent 6,632,671 to Unger issued October 14, 2003.

In certain embodiments, synthetic nanocarriers are prepared by a nanoprecipitation process or spray drying. Conditions used in preparing synthetic nanocarriers may be altered to yield particles of a desired size or property (e.g., hydrophobicity, hydrophilicity, external morphology, "stickiness," shape, etc.). The method of preparing the synthetic nanocarriers and the conditions (e.g., solvent, temperature, concentration, air flow rate, etc.) used may depend on the materials to be included in the synthetic nanocarriers and/or the composition of the carrier matrix.

If synthetic nanocarriers prepared by any of the above methods have a size range outside of the desired range, such synthetic nanocarriers can be sized, for example, using a sieve.

-31-

Preferably, in some embodiments of any one of the methods or compositions or kits provided herein, the synthetic nanocarriers are those that comprise synthetic nanocarriers composed of PLA and PLA-PEG. PLA is part of the broader poly(lactic co glycolic acid), or PLGA, family of biodegradable polymers that have more than 30 years of commercial use and are formulation components in a number of approved products. Polyethylene glycol, or PEG, has been widely studied in clinical trials and is also a formulation component in many approved biologic products.

As examples, the synthetic nanocarriers comprising rapamycin are those produced or obtainable by one of the following methods:

1) PLA with an inherent viscosity of 0.41 dL/g is purchased from Evonik Industries (Rellinghauser Straße 1—11 45128 Essen, Germany), product code Resomer Select 100 DL 4A. PLA-PEG-OMe block co-polymer with a methyl ether terminated PEG block of approximately 5,000 Da and an overall inherent viscosity of 0.50 DL/g is purchased from Evonik Industries (Rellinghauser Straße 1—11 45128 Essen, Germany), product code Resomer Select 100 DL mPEG 5000 (15 wt% PEG). Rapamycin is purchased from Concord Biotech Limited (1482-1486 Trasad Road, Dholka 382225, Ahmedabad India), product code SIROLIMUS. EMPROVE® Polyvinyl Alcohol 4-88, USP (85-89% hydrolyzed, viscosity of 3.4-4.6 mPa·s) is purchased from MilliporeSigma (EMD Millipore, 290 Concord Road Billerica, Massachusetts 01821), product code 1.41350. Dulbecco's phosphate buffered saline 1X (DPBS) is purchased from Lonza (Muenchensteinerstrasse 38, CH-4002 Basel, Switzerland), product code 17-512Q. Sorbitan monopalmitate is purchased from Croda International (300-A Columbus Circle, Edison, NJ 08837), product code SPAN 40. Solutions are prepared as follows. Solution 1 is prepared by dissolving PLA at 150 mg/mL and PLA-PEG-Ome at 50 mg/mL in dichloromethane. Solution 2 is prepared by dissolving rapamycin at 100 mg/mL in dichloromethane. Solution 3 is prepared by dissolving SPAN 40 at 50 mg/mL in dichloromethane. Solution 4 is prepared by dissolving PVA at 75 mg/mL in 100 mM phosphate buffer pH 8. O/W emulsions are prepared by adding Solution 1 (0.50 mL), Solution 2 (0.12 mL), Solution 3 (0.10 mL), and dichloromethane (0.28 mL), in a thick walled glass pressure tube. The combined organic phase solutions are then mixed by repeat pipetting. To this mixture, Solution 4 (3 mL), is added. The pressure tube is then vortex mixed for 10 seconds. Next, the crude emulsion is homogenized by sonication at 30% amplitude for 1 minute using a

-32-

Branson Digital Sonifier 250 with a 1/8" tapered tip, and the pressure tube immersed in an ice water bath. The emulsion is then added to a 50 mL beaker containing DPBS (30 mL). This is stirred at room temperature for 2 hours to allow the dichloromethane to evaporate and for the nanocarriers to form. A portion of the nanocarriers is washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,600×g at 4 °C for 50 minutes, removing the supernatant, and re-suspended the pellet in DPBS containing 0.25% w/v PVA. The wash procedure is repeated and the pellet is re-suspended in DPBS containing 0.25% w/v PVA to achieve a nanocarrier suspension having a nominal concentration of 10 mg/mL on a polymer basis. The nanocarrier suspension is then filtered using a 0.22 µm PES membrane syringe filter from MilliporeSigma (EMD Millipore, 290 Concord Rd. Billerica MA, product code SLGP033RB). The filtered nanocarrier suspension is stored at -20°C.

2) PLA with an inherent viscosity of 0.41 dL/g is purchased from Evonik Industries (Rellinghauser Straße 1—11 45128 Essen, Germany), product code Resomer Select 100 DL 4A. PLA-PEG-OMe block co-polymer with a methyl ether terminated PEG block of approximately 5,000 Da and an overall inherent viscosity of 0.50 DL/g is purchased from Evonik Industries (Rellinghauser Straße 1—11 45128 Essen, Germany), product code Resomer Select 100 DL mPEG 5000 (15 wt% PEG). Rapamycin is purchased from Concord Biotech Limited (1482-1486 Trasad Road, Dholka 382225, Ahmedabad India), product code SIROLIMUS. Sorbitan monopalmitate is purchased from Sigma-Aldrich (3050 Spruce St., St. Louis, MO 63103), product code 388920. EMPROVE® Polyvinyl Alcohol (PVA) 4-88, USP (85-89% hydrolyzed, viscosity of 3.4-4.6 mPa·s) is purchased from MilliporeSigma (EMD Millipore, 290 Concord Road Billerica, Massachusetts 01821), product code 1.41350. Dulbecco's phosphate buffered saline 1X (DPBS) is purchased from Lonza (Muenchensteinerstrasse 38, CH-4002 Basel, Switzerland), product code 17-512Q. Solutions are prepared as follows: Solution 1: A polymer, rapamycin, and sorbitan monopalmitate mixture is prepared by dissolving PLA at 37.5 mg/mL, PLA-PEG-Ome at 12.5 mg/mL, rapamycin at 8 mg/mL, and sorbitan monopalmitate at 2.5 in dichloromethane. Solution 2: Polyvinyl alcohol is prepared at 50 mg/mL in 100 mM pH 8 phosphate buffer. An O/W emulsion is prepared by combining Solution 1 (1.0 mL) and Solution 2 (3 mL) in a small glass pressure tube, and vortex mixed for 10 seconds. The formulation is then homogenized by sonication at 30% amplitude for 1

-33-

minute using a Branson Digital Sonifier 250 with a 1/8" tapered tip, with the pressure tube immersed in an ice water bath. The emulsion is then added to a 50 mL beaker containing DPBS (15 mL), and covered with aluminum foil. A second O/W emulsion is prepared using the same materials and method as above and then added to the same beaker using a fresh aliquot of DPBS (15 mL). The combined emulsion is then left uncovered and stirred at room temperature for 2 hours to allow the dichloromethane to evaporate and for the nanocarriers to form. A portion of the nanocarriers is washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,600×g and 4 °C for 50 minutes, removing the supernatant, and re-suspending the pellet in DPBS containing 0.25% w/v PVA. The wash procedure is repeated and then the pellet re-suspended in DPBS containing 0.25% w/v PVA to achieve a nanocarrier suspension having a nominal concentration of 10 mg/mL on a polymer basis. The nanocarrier suspension is then filtered using a 0.22 µm PES membrane syringe filter from MilliporeSigma (EMD Millipore, 290 Concord Rd. Billerica MA, product code SLGP033RB). The filtered nanocarrier suspension is then stored at -20°C.

Immunosuppressants

Any immunosuppressant as provided herein can be used in any one of the methods or compositions provided and can be, in some embodiments, attached to synthetic nanocarriers.

Immunosuppressants include, but are not limited to, mTOR inhibitors. Examples of mTOR inhibitors include rapamycin and rapalogs (e.g., CCL-779, RAD001, AP23573, C20-methylrapamycin (C20-Marap), C16-(S)-butylsulfonamidrapamycin (C16-BSrap), C16-(S)-3-methylindolerapamycin (C16-iRap) (Bayle et al. Chemistry & Biology 2006, 13:99-107)), AZD8055, BEZ235 (NVP-BEZ235), chrysophanic acid (chrysophanol), deforolimus (MK-8669), everolimus (RAD0001), KU-0063794, PI-103, PP242, temsirolimus, and WYE-354 (available from Selleck, Houston, TX, USA).

Preferably, in some embodiments of any one of the methods or compositions or kits provided herein, the immunosuppressant is rapamycin. In some of such embodiments, the rapamycin is preferably encapsulated in the synthetic nanocarriers. Rapamycin is the active ingredient of Rapamune, an immunosuppressant which has extensive prior use in humans and is currently FDA approved for prophylaxis of organ rejection in kidney transplant patients aged 13 or older.

-34-

When coupled to a synthetic nanocarrier, the amount of the immunosuppressant coupled to the synthetic nanocarrier based on the total dry recipe weight of materials in an entire synthetic nanocarrier (weight/weight), is as described elsewhere herein. Preferably, in some embodiments of any one of the methods or compositions or kits provided herein, the load of the immunosuppressant, such as rapamycin or rapalog, is between 7% and 12% or 8% and 12% by weight.

Dosing

Unless otherwise specified herein, the amount (by weight) of a dose of a composition comprising pegylated uricase as well as the concentrations per vial provided herein refers to the amount or concentration of the uricase protein, respectively, not including the PEG molecules conjugated thereto or any added excipients in the composition. The actual amount of the pegylated uricase, in such instances, will be higher than the dose described due to the higher weight of the pegylated protein form. In one example, a dose of 0.4 mg/kg of a composition comprising pegylated uricase refers to a dose of 0.4 mg/kg uricase protein.

Thus, a dose of a composition comprising pegylated uricase for administration to a subject may be calculated based on the dose provided herein and the weight of the subject, according to the following equation:

$$(\text{dose in mg/kg (this is of the uricase protein)}) \times (\text{subject weight (kg)}) / (\text{concentration per mL in vial (again this is of the uricase protein)}) = \text{volume to be administered}$$

As an example, the pegylated uricase may be reconstituted in sterile water to a concentration of 6 mg/mL. Thus, for this example, for a dose of 0.4 mg/kg to be administered to a subject weighing 90.7 kg (200 lbs), 6.048 mL of the reconstituted pegylated uricase composition should be administered to the subject:

$$(0.4 \text{ mg/kg}) \times (90.7 \text{ kg}) / (6 \text{ mg/mL}) = 6.048 \text{ mL}$$

In some embodiments, the appropriate volume of the composition comprising pegylated uricase is diluted in a pharmaceutically acceptable excipient (e.g., sterile saline solution) for, for example, intravenous infusion to a subject over a desired period of time (e.g., 60 minutes).

Similarly, unless otherwise specified herein, the amount (by weight) of a dose of a composition comprising synthetic nanocarriers comprising an immunosuppressant as well as the concentrations per vial as provided herein refers to the amount or concentration of the immunosuppressant, respectively, and not including the synthetic nanocarrier material or any

-35-

added excipients or other components in the composition. The actual amount of the synthetic nanocarrier composition comprising the immunosuppressant will be higher than the dose described due to the added weight of the synthetic nanocarrier material and any added excipients or other components in the composition. In one example, a dose of 0.08 mg/kg of a composition comprising synthetic nanocarriers comprising an immunosuppressant refers to a dose of 0.08 mg/kg immunosuppressant.

Thus, a dose of a composition comprising synthetic nanocarriers comprising an immunosuppressant for administration to a subject may be calculated based on the weight of the subject, according to the following equation:

$$(\text{dose in mg/kg (this is of the immunosuppressant)}) \times (\text{subject weight (kg)}) / (\text{concentration per mL in vial (again this is of the immunosuppressant)}) = \text{volume to be administered}$$

As an example, the composition comprising synthetic nanocarriers comprising an immunosuppressant is at a concentration of 2 mg/mL (again this is the concentration of the immunosuppressant). Thus, for this example, for a dose of 0.08 mg/kg to be administered to a subject weighing 90.7 kg (200 lbs), 3.6 mL of the composition should be administered to the subject:

$$(0.08 \text{ mg/kg}) \times (90.7 \text{ kg}) / (2 \text{ mg/mL}) = 3.6 \text{ mL}$$

The load of the immunosuppressant (e.g., rapamycin) of the synthetic nanocarriers comprising an immunosuppressant may be determined by extracting the immunosuppressant from the synthetic nanocarriers using liquid liquid extraction compatible with both the immunosuppressant and the synthetic nanocarriers (e.g., polymers comprising the synthetic nanocarriers) and analyzing the extract by reverse phase liquid chromatography with UV detection specific for the analyte. The immunosuppressant load (content of the synthetic nanocarriers) may be accurately and precisely calculated from a calibration standard curve of a qualified reference standard prepared in conditions compatible with the chromatography and the nanoparticle extraction procedure and analyzed concomitantly.

The amount (by weight) of a dose of a composition comprising synthetic nanocarriers comprising an immunosuppressant may be calculated based on the amount (by weight) of the immunosuppressant dose, according to the following equation:

$$(1/\text{load of immunosuppressant}) \times (\text{dose given based on the amount of immunosuppressant}) = \text{dose of immunosuppressant given as the amount of the synthetic nanocarriers comprising the immunosuppressant}$$

-36-

As an example, the load of immunosuppressant in the synthetic nanocarriers can be about 10% and if a dose of 0.08 mg/kg of the immunosuppressant is desired, the dose given as the amount of the synthetic nanocarriers comprising the immunosuppressant is 8 mg/kg.

The amount of uricase protein present in a pegylated uricase may be determined using methods known in the art, for example colorimetry, UV absorbance or amino acid analysis. The colorimetric approach relies on a standardized kit commercially available leveraging typical dye based reactions such as those described for Bradford or bicinchoninic acid (BCA) assays. The uricase protein quantity is accurately and precisely calculated from a calibration standard curve of a qualified protein reference standard, preferably purchased from compendial sources, and analyzed concomitantly using the same spectrophotometer. Single or multiple point calibration of a known protein of similar or different chemical properties may be run within the same assay to ensure consistency of the read out at the chosen UV absorbance. The amino acid mixture obtained from acid hydrolysis of the drug product may also be analyzed and generally provides a precise and accurate quantification. The amino acid mixture is analyzed by HPLC with either UV or fluorescence detection and using pre-chromatography or post-chromatography derivatization of the primary and secondary amines. Commercially available mixtures of common amino acids are analyzed within the same assay to build the individual amino acid calibration curves against which each amino acid is quantified. In some embodiments, the determination of the uricase protein quantity is supplemented by measuring the enzyme activity, which may be performed by measuring the decrease of an excess of uric acid monitored by UV absorbance at 595 nm. Alternatively or in addition, the uricase activity can be determined using a commercially available kit, which may involve, for example, labeling the enzymatic reaction product and measuring the response of the uricase against a calibration curve established by analyzing a known quantity of the enzyme.

Similar to the immediately above formula, the amount (by weight) of a dose of a composition comprising pegylated uricase can be calculated based on the amount (by weight) of the uricase dose, according to the following equation:

$$\left(\frac{1}{(\text{weight of uricase of a pegylated uricase} / \text{weight of the pegylated uricase})} \right) \times (\text{dose given based on the amount of uricase}) = \text{dose of pegylated uricase given as the amount of the pegylated uricase}$$

It should be understood that the amount provided herein can be an average amount based on a population of the respective molecules in a composition.

-37-

Exemplary doses of uricase for the compositions comprising uricase, such as pegsiticase, as provided herein can be 0.10 mg/kg, 0.11 mg/kg, 0.12 mg/kg, 0.13 mg/kg, 0.14 mg/kg, 0.15 mg/kg, 0.16 mg/kg, 0.17 mg/kg, 0.18 mg/kg, 0.19 mg/kg, 0.20 mg/kg, 0.21 mg/kg, 0.22 mg/kg, 0.23 mg/kg, 0.24 mg/kg, 0.25 mg/kg, 0.26 mg/kg, 0.27 mg/kg, 0.28 mg/kg, 0.29 mg/kg, 0.30 mg/kg, 0.31 mg/kg, 0.32 mg/kg, 0.34 mg/kg, 0.35 mg/kg, 0.36 mg/kg, 0.37 mg/kg, 0.38 mg/kg, 0.39 mg/kg, 0.40 mg/kg, 0.41 mg/kg, 0.42 mg/kg, 0.43 mg/kg, 0.44 mg/kg, 0.45 mg/kg, 0.46 mg/kg, 0.47 mg/kg, 0.48 mg/kg, 0.49 mg/kg, 0.50 mg/kg, 0.51 mg/kg, 0.52 mg/kg, 0.53 mg/kg, 0.54 mg/kg, 0.55 mg/kg, 0.56 mg/kg, 0.57 mg/kg, 0.58 mg/kg, 0.59 mg/kg, 0.60 mg/kg, 0.61 mg/kg, 0.62 mg/kg, 0.63 mg/kg, 0.64 mg/kg, 0.65 mg/kg, 0.66 mg/kg, 0.67 mg/kg, 0.68 mg/kg, 0.69 mg/kg, 0.70 mg/kg, 0.71 mg/kg, 0.72 mg/kg, 0.73 mg/kg, 0.74 mg/kg, 0.75 mg/kg, 0.76 mg/kg, 0.77 mg/kg, 0.78 mg/kg, 0.79 mg/kg, 0.80 mg/kg, 0.81 mg/kg, 0.82 mg/kg, 0.83 mg/kg, 0.84 mg/kg, 0.85 mg/kg, 0.86 mg/kg, 0.87 mg/kg, 0.88 mg/kg, 0.89 mg/kg, 0.90 mg/kg, 0.91 mg/kg, 0.92 mg/kg, 0.93 mg/kg, 0.94 mg/kg, 0.95 mg/kg, 0.96 mg/kg, 0.97 mg/kg, 0.98 mg/kg, 0.99 mg/kg, 1.0 mg/kg, 1.01 mg/kg, 1.02 mg/kg, 1.03 mg/kg, 1.04 mg/kg, 1.05 mg/kg, 1.06 mg/kg, 1.07 mg/kg, 1.08 mg/kg, 1.09 mg/kg, 1.10 mg/kg, 1.11 mg/kg, 1.12 mg/kg, 1.13 mg/kg, 1.14 mg/kg, 1.15 mg/kg, 1.16 mg/kg, 1.17 mg/kg, 1.18 mg/kg, 1.19 mg/kg, or 1.20 mg/kg uricase.

Exemplary doses of rapamycin for the compositions comprising synthetic nanocarriers comprising rapamycin can be 0.050 mg/kg, 0.055 mg/kg, 0.060 mg/kg, 0.065 mg/kg, 0.070 mg/kg, 0.075 mg/kg, 0.080 mg/kg, 0.085 mg/kg, 0.090 mg/kg, 0.095 mg/kg, 0.100 mg/kg, 0.105 mg/kg, 0.110 mg/kg, 0.115 mg/kg, 0.120 mg/kg, 0.125 mg/kg, 0.130 mg/kg, 0.135 mg/kg, 0.140 mg/kg, 0.145 mg/kg, 0.150 mg/kg, 0.155 mg/kg, 0.160 mg/kg, 0.165 mg/kg, 0.170 mg/kg, 0.175 mg/kg, 0.180 mg/kg, 0.185 mg/kg, 0.190 mg/kg, 0.195 mg/kg, 0.200 mg/kg, 0.205 mg/kg, 0.210 mg/kg, 0.215 mg/kg, 0.220 mg/kg, 0.225 mg/kg, 0.230 mg/kg, 0.235 mg/kg, 0.240 mg/kg, 0.245 mg/kg, 0.250 mg/kg, 0.255 mg/kg, 0.260 mg/kg, 0.265 mg/kg, 0.270 mg/kg, 0.275 mg/kg, 0.280 mg/kg, 0.285 mg/kg, 0.290 mg/kg, 0.295 mg/kg, 0.300 mg/kg, 0.305 mg/kg, 0.310 mg/kg, 0.315 mg/kg, 0.320 mg/kg, 0.325 mg/kg, 0.330 mg/kg, 0.335 mg/kg, 0.340 mg/kg, 0.345 mg/kg, 0.350 mg/kg, 0.355 mg/kg, 0.360 mg/kg, 0.365 mg/kg, 0.370 mg/kg, 0.375 mg/kg, 0.380 mg/kg, 0.385 mg/kg, 0.390 mg/kg, 0.395 mg/kg, 0.400 mg/kg, 0.405 mg/kg, 0.410 mg/kg, 0.415 mg/kg, 0.420 mg/kg, 0.425 mg/kg, 0.430 mg/kg, 0.435 mg/kg, 0.440 mg/kg, 0.445 mg/kg, 0.450 mg/kg, 0.455

-38-

mg/kg, 0.460 mg/kg, 0.465 mg/kg, 0.470 mg/kg, 0.475 mg/kg, 0.480 mg/kg, 0.485 mg/kg, 0.490 mg/kg, 0.495 mg/kg, 0.500 mg/kg rapamycin.

Exemplary doses of compositions comprising synthetic nanocarriers comprising rapamycin as provided herein can be 0.55 mg/kg, 0.56 mg/kg, 0.57 mg/kg, 0.58 mg/kg, 0.59

5 mg/kg, 0.60 mg/kg, 0.61 mg/kg, 0.62 mg/kg, 0.63 mg/kg, 0.64 mg/kg, 0.65 mg/kg, 0.66 mg/kg, 0.67 mg/kg, 0.68 mg/kg, 0.69 mg/kg, 0.70 mg/kg, 0.71 mg/kg, 0.72 mg/kg, 0.73 mg/kg, 0.74 mg/kg, 0.75 mg/kg, 0.76 mg/kg, 0.77 mg/kg, 0.78 mg/kg, 0.79 mg/kg, 0.80 mg/kg, 0.81 mg/kg, 0.82 mg/kg, 0.83 mg/kg, 0.84 mg/kg, 0.85 mg/kg, 0.86 mg/kg, 0.87 mg/kg, 0.88 mg/kg, 0.89 mg/kg, 0.90 mg/kg, 0.91 mg/kg, 0.92 mg/kg, 0.93 mg/kg, 0.94
10 mg/kg, 0.95 mg/kg, 0.96 mg/kg, 0.97 mg/kg, 0.98 mg/kg, 0.99 mg/kg, 1.0 mg/kg, 1.01 mg/kg, 1.02 mg/kg, 1.03 mg/kg, 1.04 mg/kg, 1.05 mg/kg, 1.06 mg/kg, 1.07 mg/kg, 1.08 mg/kg, 1.09 mg/kg, 1.10 mg/kg, 1.11 mg/kg, 1.12 mg/kg, 1.13 mg/kg, 1.14 mg/kg, 1.15 mg/kg, 1.16 mg/kg, 1.17 mg/kg, 1.18 mg/kg, 1.19 mg/kg, 1.20 mg/kg, 1.21 mg/kg, 1.22 mg/kg, 1.23 mg/kg, 1.24 mg/kg, 1.25 mg/kg, 1.26 mg/kg, 1.27 mg/kg, 1.28 mg/kg, 1.29
15 mg/kg, 1.30 mg/kg, 1.31 mg/kg, 1.32 mg/kg, 1.33 mg/kg, 1.34 mg/kg, 1.35 mg/kg, 1.36 mg/kg, 1.37 mg/kg, 1.38 mg/kg, 1.39 mg/kg, 1.40 mg/kg, 1.41 mg/kg, 1.42 mg/kg, 1.43 mg/kg, 1.44 mg/kg, 1.45 mg/kg, 1.46 mg/kg, 1.47 mg/kg, 1.48 mg/kg, 1.49 mg/kg, 1.50 mg/kg, 1.51 mg/kg, 1.52 mg/kg, 1.53 mg/kg, 1.54 mg/kg, 1.55 mg/kg, 1.56 mg/kg, 1.57 mg/kg, 1.58 mg/kg, 1.59 mg/kg, 1.60 mg/kg, 1.61 mg/kg, 1.62 mg/kg, 1.63 mg/kg, 1.64
20 mg/kg, 1.65 mg/kg, 1.66 mg/kg, 1.67 mg/kg, 1.68 mg/kg, 1.69 mg/kg, 1.70 mg/kg, 1.71 mg/kg, 1.72 mg/kg, 1.73 mg/kg, 1.74 mg/kg, 1.75 mg/kg, 1.76 mg/kg, 1.77 mg/kg, 1.78 mg/kg, 1.79 mg/kg, 1.80 mg/kg, 1.81 mg/kg, 1.82 mg/kg, 1.83 mg/kg, 1.84 mg/kg, 1.85 mg/kg, 1.86 mg/kg, 1.87 mg/kg, 1.88 mg/kg, 1.89 mg/kg, 1.90 mg/kg, 1.91 mg/kg, 1.92 mg/kg, 1.93 mg/kg, 1.94 mg/kg, 1.95 mg/kg, 1.96 mg/kg, 1.97 mg/kg, 1.98 mg/kg, 1.99
25 mg/kg, 2.00 mg/kg, 2.01 mg/kg, 2.02 mg/kg, 2.03 mg/kg, 2.04 mg/kg, 2.05 mg/kg, 2.06 mg/kg, 2.07 mg/kg, 2.08 mg/kg, 2.09 mg/kg, 2.10 mg/kg, 2.11 mg/kg, 2.12 mg/kg, 2.13 mg/kg, 2.14 mg/kg, 2.15 mg/kg, 2.16 mg/kg, 2.17 mg/kg, 2.18 mg/kg, 2.19 mg/kg, 2.20 mg/kg, 2.21 mg/kg, 2.22 mg/kg, 2.23 mg/kg, 2.24 mg/kg, 2.25 mg/kg, 2.26 mg/kg, 2.27 mg/kg, 2.28 mg/kg, 2.29 mg/kg, 2.30 mg/kg, 2.31 mg/kg, 2.32 mg/kg, 2.33 mg/kg, 2.34
30 mg/kg, 2.35 mg/kg, 2.36 mg/kg, 2.37 mg/kg, 2.38 mg/kg, 2.39 mg/kg, 2.40 mg/kg, 2.41 mg/kg, 2.42 mg/kg, 2.43 mg/kg, 2.44 mg/kg, 2.45 mg/kg, 2.46 mg/kg, 2.47 mg/kg, 2.48 mg/kg, 2.49 mg/kg, 2.50 mg/kg, 2.51 mg/kg, 2.52 mg/kg, 2.53 mg/kg, 2.54 mg/kg, 2.55 mg/kg, 2.56 mg/kg, 2.57 mg/kg, 2.58 mg/kg, 2.59 mg/kg, 2.60 mg/kg, 2.61 mg/kg, 2.62

-39-

mg/kg, 2.63 mg/kg, 2.64 mg/kg, 2.65 mg/kg, 2.66 mg/kg, 2.67 mg/kg, 2.68 mg/kg, 2.69
mg/kg, 2.70 mg/kg, 2.71 mg/kg, 2.72 mg/kg, 2.73 mg/kg, 2.74 mg/kg, 2.75 mg/kg, 2.76
mg/kg, 2.77 mg/kg, 2.78 mg/kg, 2.79 mg/kg, 2.80 mg/kg, 2.81 mg/kg, 2.82 mg/kg, 2.83
mg/kg, 2.84 mg/kg, 2.85 mg/kg, 2.86 mg/kg, 2.87 mg/kg, 2.88 mg/kg, 2.89 mg/kg, 2.90
5 mg/kg, 2.91 mg/kg, 2.92 mg/kg, 2.93 mg/kg, 2.94 mg/kg, 2.95 mg/kg, 2.96 mg/kg, 2.97
mg/kg, 2.98 mg/kg, 2.99 mg/kg, 3.00 mg/kg, 3.01 mg/kg, 3.02 mg/kg, 3.03 mg/kg, 3.04
mg/kg, 3.05 mg/kg, 3.06 mg/kg, 3.07 mg/kg, 3.08 mg/kg, 3.09 mg/kg, 3.10 mg/kg, 3.11
mg/kg, 3.12 mg/kg, 3.13 mg/kg, 3.14 mg/kg, 3.15 mg/kg, 3.16 mg/kg, 3.17 mg/kg, 3.18
mg/kg, 3.19 mg/kg, 3.20 mg/kg, 3.21 mg/kg, 3.22 mg/kg, 3.23 mg/kg, 3.24 mg/kg, 3.25
10 mg/kg, 3.26 mg/kg, 3.27 mg/kg, 3.28 mg/kg, 3.29 mg/kg, 3.30 mg/kg, 3.31 mg/kg, 3.32
mg/kg, 3.33 mg/kg, 3.34 mg/kg, 3.35 mg/kg, 3.36 mg/kg, 3.37 mg/kg, 3.38 mg/kg, 3.39
mg/kg, 3.40 mg/kg, 3.41 mg/kg, 3.42 mg/kg, 3.43 mg/kg, 3.44 mg/kg, 3.45 mg/kg, 3.46
mg/kg, 3.47 mg/kg, 3.48 mg/kg, 3.49 mg/kg, 3.50 mg/kg, 3.51 mg/kg, 3.52 mg/kg, 3.53
mg/kg, 3.54 mg/kg, 3.55 mg/kg, 3.56 mg/kg, 3.57 mg/kg, 3.58 mg/kg, 3.59 mg/kg, 3.60
15 mg/kg, 3.61 mg/kg, 3.62 mg/kg, 3.63 mg/kg, 3.64 mg/kg, 3.65 mg/kg, 3.66 mg/kg, 3.67
mg/kg, 3.68 mg/kg, 3.69 mg/kg, 3.70 mg/kg, 3.71 mg/kg, 3.72 mg/kg, 3.73 mg/kg, 3.74
mg/kg, 3.75 mg/kg, 3.76 mg/kg, 3.77 mg/kg, 3.78 mg/kg, 3.79 mg/kg, 3.80 mg/kg, 3.81
mg/kg, 3.82 mg/kg, 3.83 mg/kg, 3.84 mg/kg, 3.85 mg/kg, 3.86 mg/kg, 3.87 mg/kg, 3.88
mg/kg, 3.89 mg/kg, 3.90 mg/kg, 3.91 mg/kg, 3.92 mg/kg, 3.93 mg/kg, 3.94 mg/kg, 3.95
20 mg/kg, 3.96 mg/kg, 3.97 mg/kg, 3.98 mg/kg, 3.99 mg/kg, 4.00 mg/kg, 4.01 mg/kg, 4.02
mg/kg, 4.03 mg/kg, 4.04 mg/kg, 4.05 mg/kg, 4.06 mg/kg, 4.07 mg/kg, 4.08 mg/kg, 4.09
mg/kg, 4.10 mg/kg, 4.11 mg/kg, 4.12 mg/kg, 4.13 mg/kg, 4.14 mg/kg, 4.15 mg/kg, 4.16
mg/kg, 4.17 mg/kg, 4.18 mg/kg, 4.19 mg/kg, 4.20 mg/kg, 4.21 mg/kg, 4.22 mg/kg, 4.23
mg/kg, 4.24 mg/kg, 4.25 mg/kg, 4.26 mg/kg, 4.27 mg/kg, 4.28 mg/kg, 4.29 mg/kg, 4.30
25 mg/kg, 4.31 mg/kg, 4.32 mg/kg, 4.33 mg/kg, 4.34 mg/kg, 4.35 mg/kg, 4.36 mg/kg, 4.37
mg/kg, 4.38 mg/kg, 4.39 mg/kg, 4.40 mg/kg, 4.41 mg/kg, 4.42 mg/kg, 4.43 mg/kg, 4.44
mg/kg, 4.45 mg/kg, 4.46 mg/kg, 4.47 mg/kg, 4.48 mg/kg, 4.49 mg/kg, 4.50 mg/kg, 4.51
mg/kg, 4.52 mg/kg, 4.53 mg/kg, 4.54 mg/kg, 4.55 mg/kg, 4.56 mg/kg, 4.57 mg/kg, 4.58
mg/kg, 4.59 mg/kg, 4.60 mg/kg, 4.61 mg/kg, 4.62 mg/kg, 4.63 mg/kg, 4.64 mg/kg, 4.65
30 mg/kg, 4.66 mg/kg, 4.67 mg/kg, 4.68 mg/kg, 4.69 mg/kg, 4.70 mg/kg, 4.71 mg/kg, 4.72
mg/kg, 4.73 mg/kg, 4.74 mg/kg, 4.75 mg/kg, 4.76 mg/kg, 4.77 mg/kg, 4.78 mg/kg, 4.79
mg/kg, 4.80 mg/kg, 4.81 mg/kg, 4.82 mg/kg, 4.83 mg/kg, 4.84 mg/kg, 4.85 mg/kg, 4.86
mg/kg, 4.87 mg/kg, 4.88 mg/kg, 4.89 mg/kg, 4.90 mg/kg, 4.91 mg/kg, 4.92 mg/kg, 4.93

-40-

mg/kg, 4.94 mg/kg, 4.95 mg/kg, 4.96 mg/kg, 4.97 mg/kg, 4.98 mg/kg, 4.99 mg/kg, 5.00 mg/kg, 5.01 mg/kg, 5.02 mg/kg, 5.03 mg/kg, 5.04 mg/kg, 5.05 mg/kg, 5.06 mg/kg, 5.07 mg/kg, 5.08 mg/kg, 5.09 mg/kg, 5.10 mg/kg, 5.11 mg/kg, 5.12 mg/kg, 5.13 mg/kg, 5.14 mg/kg, 5.15 mg/kg, 5.16 mg/kg, 5.17 mg/kg, 5.18 mg/kg, 5.19 mg/kg, 5.20 mg/kg, 5.21 mg/kg, 5.22 mg/kg, 5.23 mg/kg, 5.24 mg/kg, 5.25 mg/kg, 5.26 mg/kg, 5.27 mg/kg, 5.28 mg/kg, 5.29 mg/kg, 5.30 mg/kg, 5.31 mg/kg, 5.32 mg/kg, 5.33 mg/kg, 5.34 mg/kg, 5.35 mg/kg, 5.36 mg/kg, 5.37 mg/kg, 5.38 mg/kg, 5.39 mg/kg, 5.40 mg/kg, 5.41 mg/kg, 5.42 mg/kg, 5.43 mg/kg, 5.44 mg/kg, 5.45 mg/kg, 5.46 mg/kg, 5.47 mg/kg, 5.48 mg/kg, 5.49 mg/kg, 5.50 mg/kg, 5.51 mg/kg, 5.52 mg/kg, 5.53 mg/kg, 5.54 mg/kg, 5.55 mg/kg, 5.56 mg/kg, 5.57 mg/kg, 5.58 mg/kg, 5.59 mg/kg, 5.60 mg/kg, 5.61 mg/kg, 5.62 mg/kg, 5.63 mg/kg, 5.64 mg/kg, 5.65 mg/kg, 5.66 mg/kg, 5.67 mg/kg, 5.68 mg/kg, 5.69 mg/kg, 5.70 mg/kg, 5.71 mg/kg, 5.72 mg/kg, 5.73 mg/kg, 5.74 mg/kg, 5.75 mg/kg, 5.76 mg/kg, 5.77 mg/kg, 5.78 mg/kg, 5.79 mg/kg, 5.80 mg/kg, 5.81 mg/kg, 5.82 mg/kg, 5.83 mg/kg, 5.84 mg/kg, 5.85 mg/kg, 5.86 mg/kg, 5.87 mg/kg, 5.88 mg/kg, 5.89 mg/kg, 5.90 mg/kg, 5.91 mg/kg, 5.92 mg/kg, 5.93 mg/kg, 5.94 mg/kg, 5.95 mg/kg, 5.96 mg/kg, 5.97 mg/kg, 5.98 mg/kg, 5.99 mg/kg, 6.00 mg/kg, 6.01 mg/kg, 6.02 mg/kg, 6.03 mg/kg, 6.04 mg/kg, 6.05 mg/kg, 6.06 mg/kg, 6.07 mg/kg, 6.08 mg/kg, 6.09 mg/kg, 6.10 mg/kg, 6.11 mg/kg, 6.12 mg/kg, 6.13 mg/kg, 6.14 mg/kg, 6.15 mg/kg, 6.16 mg/kg, 6.17 mg/kg, 6.18 mg/kg, 6.19 mg/kg, 6.20 mg/kg, 6.21 mg/kg, 6.22 mg/kg, 6.23 mg/kg, 6.24 mg/kg, 6.25 mg/kg, 6.26 mg/kg, 6.27 mg/kg, 6.28 mg/kg, 6.29 mg/kg, 6.30 mg/kg, 6.31 mg/kg, 6.32 mg/kg, 6.33 mg/kg, 6.34 mg/kg, 6.35 mg/kg, 6.36 mg/kg, 6.37 mg/kg, 6.38 mg/kg, 6.39 mg/kg, 6.40 mg/kg, 6.41 mg/kg, 6.42 mg/kg, 6.43 mg/kg, 6.44 mg/kg, 6.45 mg/kg, 6.46 mg/kg, 6.47 mg/kg, 6.48 mg/kg, 6.49 mg/kg, or 6.50 mg/kg, wherein the dose is given as the mg of the synthetic nanocarriers comprising the rapamycin.

Any one of the doses provided herein for the composition comprising uricase, such as pegsiticase, can be used in any one of the methods or compositions or kits provided herein. Generally, when referring to a dose to be administered to a subject the dose is a label dose. Any one of the doses provided herein for the composition comprising synthetic nanocarriers comprising an immunosuppressant, such as rapamycin, can be used in any one of the methods or compositions or kits provided herein. Generally, when referring to a dose to be administered to a subject the dose is a label dose. Thus, in any one of the methods provided herein the dose(s) are label dose(s).

-41-

In some embodiments of any one of the methods provided herein, an additional volume (prime volume) may be used to prime the infusion line for administering any of the compositions provided herein to the subject.

5 Provided herein are a number of possible dosing schedules. Accordingly, any one of the subjects provided herein may be treated according to any one of the dosing schedules provided herein. As an example, any one of the subject provided herein may be treated with a composition comprising uricase, such as pegylated uricase, and/or composition comprising synthetic nanocarriers comprising an immunosuppressant, such as rapamycin, according to any one of these dosage schedules. The mode of administration for the composition(s) of any
10 one of the treatment methods provided may be by intravenous administration, such as an intravenous infusion that, for example, may take place over about 1 hour. Additionally, any one of the methods of treatment provided herein may also include administration of an additional therapeutic, such as a uric acid lowering therapeutic, such as a uricase, or an anti-gout flare prophylactic treatment. The administration of the additional therapeutic may be
15 according to any one of the applicable treatment regimens provided herein.

Preferably, in some embodiments, the treatment with a combination of synthetic nanocarrier composition comprising immunosuppressant, such as rapamycin, with a composition comprising uricase, such as pegylated uricase, can comprise three doses of the synthetic nanocarrier composition concomitantly with the uricase-comprising composition
20 followed by two doses of uricase without the concomitant administration of a composition comprising an immunosuppressant, such as a synthetic nanocarrier composition comprising an immunosuppressant, or without the concomitant administration of an additional therapeutic. In such an embodiment, each dose may be administered every two to four weeks. In one embodiment, a method is provided whereby any one of the subjects provided
25 herein is concomitantly administered three doses of a synthetic nanocarrier composition with a uricase-comprising composition monthly for three months. In another embodiment, this method further comprises administering 2, 3, 4, 5, 6, 7, 8, 9 or 10 or more monthly doses of a uricase-comprising composition alone or without the concomitant administration of immunosuppressant, such as a synthetic nanocarrier composition comprising an
30 immunosuppressant, or an additional therapeutic. In some embodiments of any one of the methods provided herein, the level of uric acid is measured in the subject at one or more time points before, during and/or after the treatment period.

Additional Therapeutics

Additional therapeutics for elevated uric acid levels, gout, gout flare, or conditions associated with gout, may be administered to any one of the subjects provided herein, such as for the reduction of uric acid levels and/or gout treatment and/or gout flare prevention. Any one of the methods provided herein may include the administration of one or more of these additional therapeutics. In some embodiments, any one of the methods provided herein do not comprise the concomitant administration of an additional therapeutic. Examples of additional therapeutics include, but are not limited to, the following. Other examples will be known to those of skill in the art.

Additional therapeutics include anti-inflammatory therapeutics (i.e., any therapeutic that can act to reduce inflammation). Anti-inflammatory therapeutics include, but are not limited to, corticosteroids or derivatives of cortisol (hydrocortisone). Corticosteroids include, but are not limited to, glucocorticoids and mineralocorticoids. Still other examples of corticosteroids include, but are not limited to, those that are natural (e.g., 11-Dehydrocorticosterone (11-oxocorticosterone, 17-deoxycortisone) = 21-hydroxypregn-4-ene-3,11,20-trione; 11-Deoxycorticosterone (deoxycortone, desoxycortone; 21-hydroxyprogesterone) = 21-hydroxypregn-4-ene-3,20-dione; 11-Deoxycortisol (cortodoxone, cortexolone) = 17 α ,21-dihydroxypregn-4-ene-3,20-dione; 11-Ketoprogesterone (11-oxoprogesterone; Ketogestin) = pregn-4-ene-3,11,20-trione; 11 β -Hydroxypregnenolone = 3 β ,11 β -dihydroxypregn-5-en-20-one; 11 β -Hydroxyprogesterone (21-deoxycorticosterone) = 11 β -hydroxypregn-4-ene-3,20-dione; 11 β ,17 α ,21-Trihydroxypregnenolone = 3 β ,11 β ,17 α ,21-tetrahydroxypregn-5-en-20-one; 17 α ,21-Dihydroxypregnenolone = 3 β ,17 α ,21-trihydroxypregn-5-en-20-one; 17 α -Hydroxypregnenolone = 3 β ,17 α -dihydroxypregn-5-en-20-one; 17 α -Hydroxyprogesterone = 17 α -hydroxypregn-4-ene-3,11,20-trione; 18-Hydroxy-11-deoxycorticosterone = 18,21-dihydroxypregn-4-ene-3,20-dione; 18-Hydroxycorticosterone = 11 β ,18,21-trihydroxypregn-4-ene-3,20-dione; 18-Hydroxyprogesterone = 18-hydroxypregn-4-ene-3,20-dione; 21-Deoxycortisol = 11 β ,17 α -dihydroxypregn-4-ene-3,20-dione; 21-Deoxycortisone = 17 α -hydroxypregn-4-ene-3,11,20-trione; 21-Hydroxypregnenolone (prebediolone) = 3 β ,21-dihydroxypregn-5-en-20-one; Aldosterone = 11 β ,21-dihydroxypregn-4-ene-3,18,20-trione; Corticosterone (17-deoxycortisol) = 11 β ,21-dihydroxypregn-4-ene-3,20-dione; Cortisol (hydrocortisone) = 11 β ,17 α ,21-trihydroxypregn-4-ene-3,20-dione; Cortisone = 17 α ,21-dihydroxypregn-4-ene-3,11,20-trione; Pregnenolone = pregn-5-en-3 β -ol-20-one; and Progesterone = pregn-4-ene-3,20-dione); those that are synthetic, such as

-43-

progesterone-type (e.g., Flugestone (flurogestone) = 9 α -fluoro-11 β ,17 α -dihydroxypregn-4-ene-3,20-dione; Fluorometholone = 6 α -methyl-9 α -fluoro-11 β ,17 α -dihydroxypregna-1,4-diene-3,20-dione; Medrysone (hydroxymethylprogesterone) = 6 α -methyl-11 β -hydroxypregn-4-ene-3,20-dione; and Prebediolone acetate (21-acetoxypregnenolone) = 3 β ,21-dihydroxypregn-5-en-20-one 21-acetate) and progesterone derivative progestins (e.g., chlormadinone acetate, cyproterone acetate, medrogestone, medroxyprogesterone acetate, megestrol acetate, and segesterone acetate); hydrocortisone-type (e.g., Chloroprednisone = 6 α -chloro-17 α ,21-dihydroxypregna-1,4-diene-3,11,20-trione; Cloprednol = 6-chloro-11 β ,17 α ,21-trihydroxypregna-1,4,6-triene-3,20-dione; Difluprednate = 6 α ,9 α -difluoro-11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione 17 α -butyrate 21-acetate; Fludrocortisone = 9 α -fluoro-11 β ,17 α ,21-trihydroxypregn-4-ene-3,20-dione; Fluocinolone = 6 α ,9 α -difluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione; Fluperolone = 9 α -fluoro-11 β ,17 α ,21-trihydroxy-21-methylpregna-1,4-diene-3,20-dione; Fluprednisolone = 6 α -fluoro-11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione; Loteprednol = 11 β ,17 α ,dihydroxy-21-oxa-21-chloromethylpregna-1,4-diene-3,20-dione; Methylprednisolone = 6 α -methyl-11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione; Prednicarbate = 11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione 17 α -ethylcarbonate 21-propionate; Prednisolone = 11 β ,17 α ,21-trihydroxypregna-1,4-diene-3,20-dione; Prednisone = 17 α ,21-dihydroxypregna-1,4-diene-3,11,20-trione; Tixocortol = 11 β ,17 α -dihydroxy-21-sulfanylpregn-4-ene-3,20-dione; and Triamcinolone = 9 α -fluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione); methasone-type (16-methylated) (e.g., Methasone; Alclometasone = 7 α -chloro-11 β ,17 α ,21-trihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Beclometasone = 9 α -chloro-11 β ,17 α ,21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione; Betamethasone = 9 α -fluoro-11 β ,17 α ,21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione; Clobetasol = 9 α -fluoro-11 β ,17 α -dihydroxy-16 β -methyl-21-chloropregna-1,4-diene-3,20-dione; Clobetasone = 9 α -fluoro-16 β -methyl-17 α -hydroxy-21-chloropregna-1,4-diene-3,11,20-trione; Clocortolone = 6 α -fluoro-9 α -chloro-11 β ,21-dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Desoximetasone = 9 α -fluoro-11 β ,21-dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Dexamethasone = 9 α -fluoro-11 β ,17 α ,21-trihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Diflorasone = 6 α ,9 α -difluoro-11 β ,17 α ,21-trihydroxy-16 β -methylpregna-1,4-diene-3,20-dione; Difluocortolone = 6 α ,9 α -difluoro-11 β ,21-dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Fluclorolone = 6 α -fluoro-9 α ,11 β -dichloro-16 α ,17 α ,21-trihydroxypregna-1,4-dien-3,20-dione; Flumetasone = 6 α ,9 α -difluoro-11 β ,17 α ,21-trihydroxy-16 α -methylpregna-

-44-

1,4-diene-3,20-dione; Fluocortin = 6 α -fluoro-11 β ,21-dihydroxy-16 α -methylpregna-1,4-diene-3,20,21-trione; Fluocortolone = 6 α -fluoro-11 β ,21-dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Fluprednidene = 9 α -fluoro-11 β ,17 α ,21-trihydroxy-16-methylenepregna-1,4-diene-3,20-dione; Fluticasone = 6 α ,9 α -difluoro-11 β ,17 α -dihydroxy-16 α -methyl-21-thia-21-fluoromethylpregna-1,4-dien-3,20-dione; Fluticasone furoate = 6 α ,9 α -difluoro-11 β ,17 α -dihydroxy-16 α -methyl-21-thia-21-fluoromethylpregna-1,4-dien-3,20-dione 17 α -(2-furoate); Halometasone = 2-chloro-6 α ,9 α -difluoro-11 β ,17 α ,21-trihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Meprednisone = 16 β -methyl-17 α ,21-dihydroxypregna-1,4-diene-3,11,20-trione; Mometasone = 9 α ,21-dichloro-11 β ,17 α -dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Mometasone furoate = 9 α ,21-dichloro-11 β ,17 α -dihydroxy-16 α -methylpregna-1,4-diene-3,20-dione 17 α -(2-furoate); Paramethasone = 6 α -fluoro-11 β ,17 α ,21-trihydroxy-16 α -methylpregna-1,4-diene-3,20-dione; Prednylidene = 11 β ,17 α ,21-trihydroxy-16-methylenepregna-1,4-diene-3,20-dione; Rimexolone = 11 β -hydroxy-16 α ,17 α ,21-trimethylpregna-1,4-dien-3,20-dione; and Ulobetasol (halobetasol) = 6 α ,9 α -difluoro-11 β ,17 α -dihydroxy-16 β -methyl-21-chloropregna-1,4-diene-3,20-dione); Acetonides and related (e.g., Amcinonide = 9 α -fluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with cyclopentanone, 21-acetate; Budesonide = 11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with butyraldehyde; Ciclesonide = 11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with (R)-cyclohexanecarboxaldehyde, 21-isobutyrate; Deflazacort = 11 β ,21-dihydroxy-2'-methyl-5'H-pregna-1,4-dieno[17,16-d]oxazole-3,20-dione 21-acetate; Desonide = 11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with acetone; Formocortal (fluoroformylone) = 3-(2-chloroethoxy)-9 α -fluoro-11 β ,16 α ,17 α ,21-tetrahydroxy-20-oxopregna-3,5-diene-6-carboxaldehyde cyclic 16 α ,17 α -acetal with acetone, 21-acetate; Fluclorolone acetonide (flucloronide) = 6 α -fluoro-9 α ,11 β -dichloro-16 α ,17 α ,21-trihydroxypregna-1,4-dien-3,20-dione cyclic 16 α ,17 α -acetal with acetone; Fludroxycortide (flurandrenolone, flurandrenolide) = 6 α -fluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregn-4-ene-3,20-dione cyclic 16 α ,17 α -acetal with acetone; Flunisolide = 6 α -fluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with acetone; Fluocinolone acetonide = 6 α ,9 α -difluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with acetone; Fluocinonide = 6 α ,9 α -difluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with acetone, 21-acetate; Halcinonide = 9 α -fluoro-11 β ,16 α ,17 α -trihydroxy-21-chloropregn-4-ene-3,20-dione cyclic

-45-

16 α ,17 α -acetal with acetone; and Triamcinolone acetonide = 9 α -fluoro-11 β ,16 α ,17 α ,21-tetrahydroxypregna-1,4-diene-3,20-dione cyclic 16 α ,17 α -acetal with acetone); and still others (e.g., Cortivazol = 6,16 α -dimethyl-11 β ,17 α ,21-trihydroxy-2'-phenyl[3,2-c]pyrazolopregna-4,6-dien-20-one 21-acetate; and RU-28362 = 6-methyl-11 β ,17 β -dihydroxy-17 α -(1-propynyl)androsta-1,4,6-trien-3-one).

Corticosteroids, particularly glyocorticoids, have anti-inflammatory and immunosuppressive effects that may be effective in managing symptoms, including pain and inflammation associated with gout, gout flare, and/or conditions associated with gout. Administration of corticosteroids may also aid in reducing hypersensitivity reactions associated with one or more additional therapies, for example uricase replacement therapy. Still other non-limiting examples of corticosteroids, include prednisone, prednisolone, Medrol, and methylprednisolone.

Additional therapeutics include short term therapies for gout flare or pain and inflammation associated with any of the symptoms associated with gout or a condition associated with gout include nonsteroidal anti-inflammatory drugs (NSAIDS), colchicine, oral corticosteroids. Non-limiting examples of NSAIDS include both over-the-counter NSAIDS, such as ibuprofen, aspirin, and naproxen, as well as prescription NSAIDS, such as celecoxib, diclofenac, diflunisal, etodolac, indomethacin, ketoprofen, ketorolac, nabumetrone, oxaprozin, piroxiam salsalate, sulindac, and tolmetin.

Colchicine is an anti-inflammatory agent that is generally considered as an alternative for NSAIDs for managing the symptoms, including pain and inflammation associated with gout, gout flare, and/or conditions associated with gout.

Further examples of additional therapeutics include xanthine oxidase inhibitors, which are molecules that inhibit xanthine oxidase, reducing or preventing the oxidation of xanthine to uric acid, thereby reducing the production of uric acid. Xanthine oxidase inhibitors are generally classified as either purine analogues and other types of xanthine oxidase inhibitors. Examples of xanthine oxidase inhibitors include allopurinol, oxypurinol, tisopurine, febuxostat, topiroxostat, inositols (e.g., phytic acid and myo-inositol), flavonoids (e.g., kaempferol, myricetin, quercetin), caffeic acid, and 3,4-dihydrox-5-nitrobenzaldehyde (DHNB).

Still other examples of additional therapeutics include uricosuric agents. Uricosuric agents aim to increase excretion of uric acid in order to reduce serum levels of uric acid by modulating renal tubule reabsorption. For example, some uricosuric agents modulate activity

-46-

of renal transporters of uric acid (e.g., URAT1/SLC22A12 inhibitors). Non-limiting examples of uricosuric agents include probenecid, benzbromarone, lesinurad, sulfinpyrazone. Other additional therapeutics may also have uricosuric activity, such as aspirin.

Additional therapeutics also include other uricase-based therapies, which include
5 pegylated uricase. Such therapies, such as when infused into humans, have been shown to reduce blood uric acid levels and improve gout symptoms. Rasburicase (Elitek®), an unpegylated recombinant uricase cloned from *Aspergillus flavus*, is approved for management of uric acid levels in patients with tumor lysis syndrome (Elitek®).
KRYSTEXXA® (pegloticase) is a recombinant uricase (primarily porcine with a carboxyl-
10 terminus sequence from baboon) bound by multiple 10 kDa PEG molecules approved for the treatment of chronic refractory gout. As mentioned elsewhere, however, the clinical experience with KRYSTEXXA® has shown that a significant number of patients will develop anti-drug antibodies which limit the long term efficacy of the drug. Thus, prior administration of KRYSTEXXA® may be a contraindication for the use of the methods
15 provided herein.

The treatments provided herein may allow patients to switch to oral gout therapy, such as with xanthine oxidase inhibitors, unless and until such patients experience a subsequent manifestation of uric acid deposits at which time a new course of treatment as provided herein according to any one of the methods provided is then undertaken. Any one
20 of the methods provided herein, thus, can include the subsequent administration of an oral gout therapeutic as an additional therapeutic after the treatment regimen according to any one of the methods provided is performed. It is believed that oral therapy may not completely prevent the build up over time of uric acid crystals in patients with a history of chronic tophaceous gout. As a result, it is anticipated that treatment as provided herein is likely to be
25 required intermittently in such patients. Thus, in such subjects, the subject is also further administered one or more compositions according to any one of the methods provided herein.

The treatments provided herein may allow patients to subsequently be treated with a uric acid lowering therapeutic, such as a uricase. In some embodiments, without an immunosuppressant. In some embodiments, without synthetic nanocarriers comprising an
30 immunosuppressant.

Treatment according to any one of the methods provided herein may also include a pre-treatment with an anti-gout flare therapeutic, such as with colchicine or NSAIDS. Accordingly, any one of the methods provided herein may further comprise such an anti-gout

-47-

flare therapeutic whereby the anti-gout flare therapeutic is concomitantly administered with the composition comprising uricase and the composition comprising synthetic nanocarriers comprising an immunosuppressant.

Identifying or monitoring of a subject, such as with the measurement of serum uric acid levels and/or ADAs, may be a step comprised in any one of the methods provided
5 herein. The methods provided herein can include a step of determining or obtaining an anti-uricase antibody level (such as a titer) in a subject, such as in a sample from the subject, and the level can be used to identify a subject for treatment or monitor a subject that is being treated. In one embodiment, the treatment may be any one of the treatments provided herein.
10 For example, if the anti-uricase antibody level is less than a threshold (e.g., 1080), the treatment may be indicated or may not need to be adjusted. If the anti-uricase antibody level is greater than a threshold (e.g., 1080), an adjustment to the treatment or further monitoring or assessment may be indicated.

In some embodiments, should such subject develop an undesired immune response,
15 the subject is further administered one or more compositions according to any one of the methods provided herein. In some embodiments of any one of the methods provided herein, the subject is monitored with dual energy computed tomography (DECT), that can be used to visualize uric acid deposits in joints and tissues. Imaging, such as with DECT, can be used to assess the efficacy of treatment with any one of the methods or compositions provided herein.
20 As a result, any one of the methods provided herein can further include a step of imaging, such as with DECT. In some embodiments of any one of the methods provided herein, the subject is one in which the gout, such as chronic tophaceous gout, or condition associated with gout has been diagnosed with such imaging, such as with DECT.

25 *Subjects*

Subjects provided herein can be in need of treatment according to any one of the methods or compositions or kits provided herein. Such subjects include those with elevated serum uric acid levels or uric acid deposits. Such subjects include those with hyperuricemia. It is within the skill of a clinician to be able to determine subjects in need of a treatment as
30 provided herein.

In some embodiments, any one of the subjects for treatment as provided in any one of the methods provided has gout or a condition associated with gout or another condition as

provided herein. In some embodiments, any one of the subjects for treatment as provided in any one of the methods provided the subject has had or is expected to have gout flare.

In some embodiments, the subject has or is at risk of having erosive bone disease associated with gout, cirrhosis or steatohepatitis associated with gout, or visceral gout.

5 In some embodiments, the subject has or is at risk of having an elevated uric acid level, e.g., an elevated plasma or serum uric acid level. When blood levels of uric acid may exceed the physiologic limit of solubility, the uric acid may crystallize in the tissues, including the joints, and may cause gout and gout-associated conditions.

In some embodiments, serum uric acid levels ≥ 5 mg/dL, ≥ 6 mg/dL, or ≥ 7 mg/dL are
10 indicative that a subject may be a candidate for treatment with any one of the methods or compositions or kits described herein. In some embodiments, such a subject has a serum level of uric acid ≥ 6 mg/dL, for example, between 6.1 mg/dL - 15 mg/dL, between 6.1 mg/dL - 10 mg/dL, 7 mg/dL - 15 mg/dL, 7 mg/dL - 10 mg/dL, 8 mg/dL - 15 mg/dL, 8 mg/dL - 10 mg/dL, 9 mg/dL - 15 mg/dL, 9 mg/dL - 10 mg/dL, 10 mg/dL - 15 mg/dL, or 11
15 mg/dL - 14 mg/dL. In some embodiments, the subject has serum level of uric acid of about 6.1 mg/dL, 6.2 mg/dL, 6.3 mg/dL, 6.4 mg/dL, 6.5 mg/dL, 6.7 mg/dL, 6.8 mg/dL, 6.9 mg/dL, 7.0 mg/dL, 7.1 mg/dL, 7.2 mg/dL, 7.3 mg/dL, 7.4 mg/dL, 7.5 mg/dL, 7.6 mg/dL, 7.7 mg/dL, 7.8 mg/dL, 7.9 mg/dL, 8.0 mg/dL, 8.1 mg/dL, 8.2 mg/dL, 8.3 mg/dL, 8.4 mg/dL, 8.5 mg/dL, 8.6 mg/dL, 8.7 mg/dL, 8.8 mg/dL, 8.9 mg/dL, 9.0 mg/dL, 9.1 mg/dL, 9.2 mg/dL, 9.3 mg/dL,
20 9.4 mg/dL, 9.5 mg/dL, 9.6 mg/dL, 9.7 mg/dL, 9.8 mg/dL, 9.9 mg/dL, 10.0 mg/dL, 10.1 mg/dL, 10.2 mg/dL, 10.3 mg/dL, 10.4 mg/dL, 10.5 mg/dL, 10.6 mg/dL, 10.7 mg/dL, 10.8 mg/dL, 10.9 mg/dL, 11.0 mg/dL, 11.1 mg/dL, 11.2 mg/dL, 11.3 mg/dL, 11.4 mg/dL, 11.5 mg/dL, 11.6 mg/dL, 11.7 mg/dL, 11.8 mg/dL, 11.9 mg/dL, 12.0 mg/dL, 12.1 mg/dL, 12.2 mg/dL, 12.3 mg/dL, 12.4 mg/dL, 12.5 mg/dL, 12.6 mg/dL, 12.7 mg/dL, 12.8 mg/dL, 12.9
25 mg/dL, 13.0 mg/dL, 13.1 mg/dL, 13.2 mg/dL, 13.3 mg/dL, 13.4 mg/dL, 13.5 mg/dL, 13.6 mg/dL, 13.7 mg/dL, 13.8 mg/dL, 13.9 mg/dL, 14.0 mg/dL, 14.1 mg/dL, 14.2 mg/dL, 14.3 mg/dL, 14.4 mg/dL, 14.5 mg/dL, 14.6 mg/dL, 14.7 mg/dL, 14.8 mg/dL, 14.9 mg/dL, 15.0 mg/dL or higher. In some embodiments, the subject has a plasma or serum uric acid level of 5.0 mg/dL, 5.1 mg/dL, 5.2 mg/dL, 5.3 mg/dL, 5.4 mg/dL, 5.5 mg/dL, 5.6 mg/dL, 5.7 mg/dL,
30 5.8 mg/dL, 5.9 mg/dL, 6.0 mg/dL, 6.1 mg/dL, 6.2 mg/dL, 6.3 mg/dL, 6.4 mg/dL, 6.5 mg/dL, 6.6 mg/dL, 6.7 mg/dL, 6.8 mg/dL, 6.9 mg/dL, or 7.0 mg/dL. In some embodiments, the subject has a plasma or serum uric acid level of greater than or equal to 5.0 mg/dL, 5.1 mg/dL, 5.2 mg/dL, 5.3 mg/dL, 5.4 mg/dL, 5.5 mg/dL, 5.6 mg/dL, 5.7 mg/dL, 5.8 mg/dL, 5.9

-49-

mg/dL, 6.0 mg/dL, 6.1 mg/dL, 6.2 mg/dL, 6.3 mg/dL, 6.4 mg/dL, 6.5 mg/dL, 6.6 mg/dL, 6.7 mg/dL, 6.8 mg/dL, 6.9 mg/dL, or 7.0 mg/dL.

In some embodiments, the subject has, or is at risk of having, hyperuricemia. In some embodiments, the subject has, or is at risk of having, gout, acute gout, acute intermittent gout, 5 gouty arthritis, acute gouty arthritis, acute gouty arthropathy, acute polyarticular gout, recurrent gouty arthritis, chronic gout (with or without tophi), tophaceous gout, chronic tophaceous gout, chronic advanced gout (with or without tophi), chronic polyarticular gout (with or without tophi), chronic gouty arthropathy (with or without tophi), idiopathic gout, idiopathic chronic gout (with or without tophi), primary gout, chronic primary gout (with or 10 without tophi), refractory gout, such as chronic refractory gout, axial gouty arthropathy, a gout attack, a gout flare, podagra (i.e., monarticular arthritis of the great toe), chiragra (i.e., monarticular arthritis of the hand), gonagra (i.e., monarticular arthritis of the knee), gouty bursitis, gouty spondylitis, gouty synovitis, gouty tenosynovitis, gout that affects tendons and ligaments, lead-induced gout (i.e., saturnine gout), drug induced gout, gout due to renal 15 impairment, gout due to kidney disease, chronic gout due to renal impairment (with or without tophi), chronic gout due to kidney disease (with or without tophi), erosive bone disease associated with gout, stroke associated with gout, vascular plaque associated with gout, cirrhosis or steatohepatitis associated with gout, liver-associated gout, incident and recurrent gout, diabetes associated with damage to pancreas in gout, general inflammatory 20 diseases exacerbated by gout, other secondary gout, or unspecified gout.

In some embodiments, the subject has, or is at risk of having, a condition associated with the renal system, for example, calculus of urinary tract due to gout, uric acid urolithiasis, uric acid nephrolithiasis, uric acid kidney stones, gouty nephropathy, acute gouty nephropathy, chronic gouty nephropathy, urate nephropathy, uric acid nephropathy, and 25 gouty interstitial nephropathy.

In some embodiments, the subject has, or is at risk of having, a condition associated with the nervous system, for example, peripheral autonomic neuropathy due to gout, gouty neuropathy, gouty peripheral neuropathy, gouty entrapment neuropathy, or gouty neuritis.

In some embodiments, the subject has, or is at risk of having, a condition associated 30 with the cardiovascular system, for example, metabolic syndrome, hypertension, obesity, diabetes, myocardial infarction, stroke, dyslipidemia, hypertriglyceridemia, insulin resistance/hyperglycemia, coronary artery disease/coronary heart disease, coronary artery disease or blockage associated with gout or hyperuricemia, heart failure, peripheral arterial

-50-

disease, stroke/cerebrovascular disease, peripheral vascular disease, and cardiomyopathy due to gout.

In some embodiments, the subject has, or is at risk of having, a condition associated with the ocular system including, for example, gouty iritis, inflammatory disease in the eye caused by gout, dry eye syndrome, red eye, uveitis, intraocular hypertension, glaucoma, and cataracts.

In some embodiments, the subject has, or is at risk of having, a condition associated with the skin including, for example, gout of the external ear, gouty dermatitis, gouty eczema, gouty panniculitis, and miliarial gout.

Compositions and Kits

Compositions provided herein may comprise inorganic or organic buffers (e.g., sodium or potassium salts of phosphate, carbonate, acetate, or citrate) and pH adjustment agents (e.g., hydrochloric acid, sodium or potassium hydroxide, salts of citrate or acetate, amino acids and their salts) antioxidants (e.g., ascorbic acid, alpha-tocopherol), surfactants (e.g., polysorbate 20, polysorbate 80, polyoxyethylene9-10 nonyl phenol, sodium desoxycholate), solution and/or cryo/lyo stabilizers (e.g., sucrose, lactose, mannitol, trehalose), osmotic adjustment agents (e.g., salts or sugars), antibacterial agents (e.g., benzoic acid, phenol, gentamicin), antifoaming agents (e.g., polydimethylsiloxane), preservatives (e.g., thimerosal, 2-phenoxyethanol, EDTA), polymeric stabilizers and viscosity-adjustment agents (e.g., polyvinylpyrrolidone, poloxamer 488, carboxymethylcellulose) and co-solvents (e.g., glycerol, polyethylene glycol, ethanol).

Compositions according to the invention may comprise pharmaceutically acceptable excipients. The compositions may be made using conventional pharmaceutical manufacturing and compounding techniques to arrive at useful dosage forms. Techniques suitable for use in practicing the present invention may be found in Handbook of Industrial Mixing: Science and Practice, Edited by Edward L. Paul, Victor A. Atiemo-Obeng, and Suzanne M. Kresta, 2004 John Wiley & Sons, Inc.; and Pharmaceutics: The Science of Dosage Form Design, 2nd Ed. Edited by M. E. Auten, 2001, Churchill Livingstone. In an embodiment, compositions are suspended in a sterile saline solution for injection together with a preservative.

It is to be understood that the compositions of the invention can be made in any suitable manner, and the invention is in no way limited to compositions that can be produced

-51-

using the methods described herein. Selection of an appropriate method of manufacture may require attention to the properties of the particular elements being associated.

In some embodiments, compositions are manufactured under sterile conditions or are initially or terminally sterilized. This can ensure that resulting compositions are sterile and non-infectious, thus improving safety when compared to non-sterile compositions. This provides a valuable safety measure, especially when subjects receiving the compositions have immune defects, are suffering from infection, and/or are susceptible to infection. In some embodiments, the compositions may be lyophilized and stored in suspension or as lyophilized powder depending on the formulation strategy for extended periods without losing activity.

Administration according to the present invention may be by a variety of routes, including but not limited to an intravenous route. The compositions referred to herein may be manufactured and prepared for administration using conventional methods.

The compositions of the invention can be administered in effective amounts, such as the effective amounts described elsewhere herein. Doses of compositions as provided herein may contain varying amounts of elements according to the invention. The amount of elements present in the compositions for dosing can be varied according to their nature, the therapeutic benefit to be accomplished, and other such parameters. The compositions for dosing may be administered according to any one of the frequencies provided herein.

Another aspect of the disclosure relates to kits. In some embodiments, the kit comprises any one or more of the compositions provided herein. In some embodiments of any one of the kits provided, the kit comprises any one or more of the compositions comprising uricase as provided herein. Preferably, the uricase-comprising composition(s) is/are in an amount to provide any one or more doses as provided herein. The uricase-comprising composition(s) can be in one container or in more than one container in the kit.

In some embodiments of any one of the kits provided, the kit further comprises any one or more of the synthetic nanocarrier compositions provided herein. Preferably, the synthetic nanocarrier composition(s) is/are in an amount to provide one or more of the synthetic nanocarrier doses provided herein. The synthetic nanocarrier composition(s) can be in one container or in more than one container in the kit. In some embodiments of any one of the kits provided, the container is a vial or an ampoule. In some embodiments of any one of the kits provided, the composition(s) are in lyophilized form each in a separate container or in the same container, such that they may be reconstituted at a subsequent time. In some embodiments of any one of the kits, the lyophilized composition further comprises a sugar,

-52-

such as mannitol. In some embodiments of any one of the kits provided, the composition(s) are in the form of a frozen suspension each in a separate container or in the same container, such that they may be reconstituted at a subsequent time. In some embodiments of any one of the kits, the frozen suspension further comprises PBS. In some embodiments of any one of the kits, the kit further comprises PBS and/or 0.9% sodium chloride, USP. In some embodiments of any one of the kits provided, the kit further comprises instructions for reconstitution, mixing, administration, etc. In some embodiments of any one of the kits provided, the instructions include a description of any one of the methods described herein. Instructions can be in any suitable form, e.g., as a printed insert or a label. In some embodiments of any one of the kits provided herein, the kit further comprises one or more syringes or other device(s) that can deliver the composition(s) in vivo to a subject.

EXAMPLES

Example 1: SEL 212 clinical trial results, non-human

Preclinical development

SEL 212 was used to treat uricase deficient mice and wild type mice, rats and nonhuman primates to evaluate efficacy, dose regimens and safety.

Proof of concept study in uricase deficient mice

A pharmacology study in mice that were genetically deficient in endogenous uricase was conducted. The study evaluated the efficacy of a dose regimen consisting of three immunizations with SEL 212 followed by doses of pegsiticase alone in preventing the formation of ADAs to pegsiticase. The treatment period consisted of the first 14 days of the study. In the study, mice were separated into three treatment groups. During the treatment period:

- the first group, referred to as the Untreated Group, received no treatment;
- the second group, referred to as the Pegsiticase Group, was treated with pegsiticase alone; and
- the third group, referred to as the SVP Rapamycin + Pegsiticase Group, was treated with SVP Rapamycin co administered with pegsiticase.

-53-

The Pegsiticase Group and SVP Rapamycin + Pegsiticase Group were treated on days zero, seven and 14 of the treatment period. Each group was then treated with pegsiticase alone on days 35 and 42 of the study, or the challenge period. Uricase specific ADA levels were recorded to determine the formation of ADAs to pegsiticase. Uric acid levels were measured to determine effectiveness of SVP Rapamycin co administered with pegsiticase in lowering uric acid levels below 6 mg/dl, which is the treatment target for gout patients.

Antibody formation. The Pegsiticase Group developed uricase specific ADAs when exposed to pegsiticase during the treatment period. The Untreated Group also developed uricase specific ADAs as soon as they were challenged with pegsiticase. Despite exposure to pegsiticase during both the treatment and challenge periods, the SVP Rapamycin + Pegsiticase Group did not develop uricase specific ADAs during either period.

Uric acid levels. After initial exposure to pegsiticase, the Untreated Group maintained high uric acid levels of approximately 10 mg/dl. The Pegsiticase Group recorded uric acid levels below 6 mg/dl after the first dose in the treatment period. However, during subsequent doses in the treatment period and challenge period, uric acid levels returned to levels well in excess of 6 mg/dl. In contrast, the SVP Rapamycin + Pegsiticase Group maintained uric acid levels that were close to zero throughout the study.

Proof of concept study in nonhuman primates

A preclinical study to evaluate the ability of SVP Rapamycin to mitigate the formation of uricase specific ADAs in nonhuman primates was also conducted. As depicted in **Fig. 3**, during the study:

- pegsiticase was administered alone, referred to as the Empty Nanoparticle Group, or
- was co-administered with one of two dose levels of SVP Rapamycin, referred to as the SVP Rapamycin 0.1X and SVP Rapamycin 1X Groups, respectively. The SVP Rapamycin 0.1X Group received a dose level of SVP Rapamycin of 0.3 mg/kg and the SVP Rapamycin 1X Group received a dose level of SVP Rapamycin of 3 mg/kg.

The Empty Nanoparticle Group received three monthly doses of pegsiticase and each of the SVP Rapamycin 0.1X Group and SVP Rapamycin 1X Group received three monthly doses of pegsiticase co-administered with SVP Rapamycin. All groups then received two monthly doses of pegsiticase alone. The SVP Rapamycin 0.1X Group received one tenth of the dose administered in the SVP Rapamycin 1X Group.

-54-

Antibody formation. It was observed that the Empty Nanoparticle Group produced high levels of uricase specific ADAs by the end of the study. The SVP Rapamycin 0.1X Group and SVP Rapamycin 1X Group were able to reduce the levels of uricase specific ADAs significantly compared to the Empty Nanoparticle Group and, in the case of the SVP Rapamycin 1X Group, inhibited the formation of antibodies. The observations in this study confirmed in non-human primates the mitigation of uricase specific ADAs that was observed in mice.

Uric acid levels. As expected, the effect that pegsiticase alone or pegsiticase co-administered with SVP Rapamycin had on uric acid levels in nonhuman primates could not be determined due to the activity of naturally occurring uricase in these animals.

Based on these preclinical studies, as well as toxicology studies conducted to conform to regulatory guidelines, referred to as current good laboratory practice, or GLP, it is believed that SEL 212 demonstrated sufficient efficacy and safety in the preclinical animal models to justify movement into clinical development.

Example 2: SEL 212 clinical trial results, human

Phase 1a clinical trial

The Phase 1a clinical trial for SEL 212 was an ascending dose trial of pegsiticase alone in 22 subjects with elevated serum uric acid levels greater than 6 mg/dl who were separated into five cohorts. Each cohort received a single intravenous infusion of pegsiticase at the following dose levels of 0.1 mg/kg for Cohort #1, 0.2 mg/kg for Cohort #2, 0.4 mg/kg for Cohort #3, 0.8 mg/kg for Cohort #4 and 1.2 mg/kg for Cohort #5. Dosing began with the lowest dose and only after an entire cohort was safely dosed was the next cohort started. The subjects were monitored during a 30 day period post infusion with visits occurring on day 7, 14, 21 and the end of trial visit on day 30. Blood and serum of each patient was evaluated for serum uric acid, ADAs (specifically anti-peg, anti-uricase and anti-pegsiticase) and safety parameters. It was observed that pegsiticase demonstrated no serious adverse events and was well tolerated at the five dose levels tested. Additionally, it was observed that pegsiticase rapidly reduced (within hours) and sustained average serum uric acid levels below 6 mg/dl for each cohort for 14 to 30 days, depending on the dose level. Consistent with preclinical studies in animals, pegsiticase induced uricase specific ADAs in all subjects with varying levels in this Phase 1a trial.

-55-

Fig. 4 depicts average serum uric acid levels of the Phase 1a clinical trial's five cohorts tested at different measurement intervals (Day 7, 14, 21 and 30) during the course of the 30 day period following the single intravenous infusion of pegsiticase at the outset of the trial.

5 The serum uric acid levels were measured at baseline and days seven, 14, 21 and 30 and uricase specific ADA levels at baseline and days seven, 14 and 30 following a single intravenous injection of pegsiticase. Uricase specific ADA levels at day 21 in the Phase 1a clinical trial were not measured. Based on the results from the Phase 1a clinical trial, it was observed that pegsiticase at a tolerated dose is capable of achieving and maintaining a
10 reduction of serum uric acid below the target of 6 mg/dl for a 30 day period in the absence of inhibitory uricase specific ADAs.

Phase 1b clinical trial

The Phase 1b clinical trial enrolled 63 patients with serum uric acid levels greater than
15 6 mg/dl separated into 11 cohorts. A single intravenous infusion of SVP Rapamycin alone at the following ascending dose levels was administered to four cohorts in ascending order. Each cohort consisted of seven patients and were designated as follows: Cohort #1 (0.03 mg/kg), Cohort #3 (0.1 mg/kg), Cohort #5 (0.3 mg/kg) and Cohort #7 (0.5 mg/kg) collectively the SVP Rapamycin Cohorts. After a cohort of the SVP-Rapamycin alone had
20 successfully and safely been dosed the corresponding dose level of SVP Rapamycin was combined with a fixed dose of pegsiticase (0.4mg/kg). The combination was co-administered sequentially as a single intravenous infusion, with the SVP Rapamycin infusion preceding the pegsiticase infusion. The cohort designation is as follows for the six cohorts (5 patients per cohort), which were Cohort #2 (SVP Rapamycin 0.03 mg/kg + 0.4 mg/kg pegsiticase),
25 Cohort #4 (SVP Rapamycin 0.1mg/kg + 0.4 mg/kg pegsiticase), Cohort #6 (SVP Rapamycin 0.3 mg/kg + 0.4 mg/kg pegsiticase), Cohort #10 (0.4 mg/kg pegsiticase + 0.03 mg/kg SVP Rapamycin separated by 48 hours), Cohort #12 (SVP Rapamycin 0.15 mg/kg + 0.4 mg/kg pegsiticase) and Cohort #14 (SVP Rapamycin 0.1 mg/kg + 0.4 mg/kg pegsiticase) collectively the SEL 212 Cohorts. In Cohort #9 a fixed amount of pegsiticase alone at a dose
30 level of 0.4 mg/kg was administered to five patients, which is referred to as the Pegsiticase Cohort. Methods of such treatment are also provided. The subjects were monitored during a 30 day period post infusion with visits occurring on day 7, 14 ,21 and the end of trial visit on day 30. Blood and serum of each patient was evaluated for serum uric acid, ADAs

-56-

(specifically anti-PEG, anti-uricase and anti-pegositicase) and safety parameters. The primary objective of the Phase 1b clinical trial was to evaluate the safety and tolerability of SVP Rapamycin alone and in combination with a fixed dose of pegositicase. A secondary clinical objective was to evaluate the ability of SVP Rapamycin co-administered with pegositicase to reduce serum uric acid levels and mitigate the formation of uricase specific ADAs when compared to administration of pegositicase alone.

Fig. 5 indicates the serum uric acid levels of Cohort #3 from the Phase 1a clinical trial, in which subjects received a fixed amount of pegositicase alone (at the same 0.4 mg/kg pegositicase. Also in the first graph is the data from Cohort# 9 (pegositicase 0.4 mg/kg) of the Phase 1b clinical trial. This graph represents the reproducibility of the data across two separate studies. In both cohorts there is initial control of the serum uric acid (levels maintained below 6mg/dL) but past day 14, individuals loose the enzyme activity. Also in **Fig. 5**, the data from the SVP rapamycin alone cohorts is displayed. All values remain essentially the same throughout the 30 days of testing indicating that SVP Rapamycin alone has no effect on serum uric acid levels. For Cohort #2 from the Phase 1b clinical trial, which received the lowest dose of SVP Rapamycin co-administered with pegositicase, it was observed that four out of five subjects tested maintained serum uric acid levels below 6 mg/dl through day 21 of the trial. It was also observed that four out of five subjects in Cohort #4 from the Phase 1b clinical trial, which received the second lowest dose of SVP Rapamycin co-administered with pegositicase, maintained levels of serum uric acid of less than 0.1 mg/dl through day 30. For Cohort #6 (SEL 212 Cohort), it was observed that four (out of the projected five) subjects maintained levels of serum uric acid of less than 0.1 mg/dl through day 21 and two (out of the projected five) subjects maintained levels of serum uric acid of less than 0.1 mg/dl through day 30. By comparison, for Cohort #9 (Pegositicase Cohort), four of the five subjects returned to baseline serum uric acid levels by day 30.

Fig. 5 shows the serum uric acid levels and uricase specific ADA levels for each subject in Cohort #3 of the Phase 1a clinical trial and Cohort #9 (Pegositicase Cohort) of the Phase 1b clinical trial for comparison to the serum uric acid levels and uricase specific ADA levels for each subject in Cohort # 4 (SEL 212 Cohort) in the Phase 1b clinical trial. Cohort #3 from the Phase 1a clinical trial is depicted along with Cohort #9 from the Phase 1b clinical trial for purposes of comparison against Cohort #4 from the Phase 1b clinical trial because the subjects in these cohorts received the same fixed dose of pegositicase. In addition, Cohort #4 from the Phase 1b clinical trial is depicted in **Fig. 5** because the subjects in Cohort #4

-57-

from the Phase 1b clinical trial received a higher dose of SVP Rapamycin than did the subjects in Cohort #2 in the Phase 1b clinical trial, the other SEL 212 Cohort for which 30 day observation period data from the Phase 1b clinical trial was available.

As depicted in **Fig. 5**, in Cohort #3 from the Phase 1a clinical trial and Cohort #9 from the Phase 1b clinical trial, uricase specific ADA formation at day 14 resulting in a return to baseline levels of serum uric acid was observed. In comparison, for Cohort #4 from the Phase 1b clinical trial, it was observed that minimal uricase specific ADA formation in four of the five subjects tested with corresponding maintenance of control of serum uric acid levels through day 30. In the Phase 1a clinical trial, uricase specific ADA levels at day 21 was not measured. However, in the course of conducting the Phase 1a clinical trial, it was learned that it would be useful to measure uricase specific ADA levels at day 21 to more fully understand any variations in such levels between day 14 and day 30. As a result, for the Phase 1b clinical trial, uricase specific ADA levels at day 21 were monitored.

Additional serum uric acid and uricase specific ADA data after day 30 was collected for three of the subjects in Cohort #4 (SEL 212 Cohort) that had no or very low serum uric acid and uricase specific ADA levels at day 30. Data on day 37 was collected for all three of these subjects and again on day 42 or day 44 for two of the three subjects. Each of these three subjects had no or very low uricase specific ADA levels on day 37, day 42 or day 44, as applicable. Serum uric acid levels remained below baseline on day 37 in all three subjects. With respect to the two subjects for which day 42 or day 44 data was available, serum uric acid levels approached or exceeded baseline by the last time point measured. Based on the observations from the Phase 1b clinical trial data that SEL 212 was capable of controlling uric acid levels for at least 30 days in the majority of subjects in Cohort # 4.

On a combined basis, a total of 85 subjects have been dosed with either SEL 212 (SVP Rapamycin and pegsiticase), SVP Rapamycin alone or pegsiticase alone in connection with the Phase 1a and Phase 1b clinical trials. It has been generally observed that SEL 212 and its components, SVP Rapamycin and pegsiticase, have been well tolerated. There have been a total of four serious adverse events, or SAEs, in both Phase 1 clinical trials. All SAEs fully resolved.

Fig. 6 shows the serum uric acid levels and uricase-specific ADA levels for each subject in Cohort #3 of the Phase 1a clinical trial and Cohort #9 (Pegsiticase Cohort) of the Phase 1b clinical trial for comparison to the serum uric acid levels and uricase-specific ADA levels for each subject in Cohort # 4 (SEL-212 Cohort) and Cohort #6 (SEL-212 Cohort) in

-58-

the Phase 1b clinical trial. Cohort #3 from the Phase 1a clinical trial is also depicted along with Cohort #9 from the Phase 1b clinical trial for purposes of comparison against Cohort #4 and Cohort #6 from the Phase 1b clinical trial because the subjects in these cohorts received the same fixed dose of pegsiticase. In addition, Cohort #4 from the Phase 1b clinical trial is depicted because the subjects in Cohort #4 from the Phase 1b clinical trial received a higher dose of SVP-Rapamycin than did the subjects in Cohort #2 in the Phase 1b clinical trial. Also included is Cohort #6 from the Phase 1b clinical trial because these subjects received the highest dose of SVP-Rapamycin tested to date—higher than both Cohorts #2 and #4.

Fig. 7 presents a non-head-to-head comparison of the efficacy of SEL-212 in Cohort #6 of the Phase 1b clinical trial with Cohort #5 of the Phase 1b clinical trial and data from two replicate, randomized, double-blind, placebo-controlled clinical trials of KRYSTEXXA® as reported in the Journal of the American Medical Association in 2011. These two KRYSTEXXA® clinical trials included 85 patients who received biweekly doses of KRYSTEXXA®, 84 patients who received monthly doses of KRYSTEXXA® and 43 patients who received a placebo.

KRYSTEXXA® has been approved for the treatment of refractory gout on a biweekly dose regimen whereas the monthly dose regimen of KRYSTEXXA® has not been approved for marketing. The graph on the left below depicts the data for the four-week period after the first dose of Krystexxa® from the cohorts of subjects in the KRYSTEXXA® clinical trials who received monthly doses.

The placebo control subjects, indicated in open circles in **Fig. 7**, had uric acid levels above 6 mg/dl for the entire four weeks. The KRYSTEXXA®-treated subjects that went on to become responders, as defined by maintenance of uric acid levels below 6 mg/dl for 80% of the time at months three and six, are indicated in black circles. The KRYSTEXXA®-treated subjects that went on to become non-responders, as defined by the inability to maintain uric acid levels below 6 mg/dl for 80% of the time at months three and six, are indicated in black triangles. Only 35% of KRYSTEXXA®-treated subjects in the monthly dosing cohorts were classified as responders. It is notable that, even at four weeks, the mean uric acid levels were above 6 mg/dl in the non-responders, representing 65% of subjects, and were above 4 mg/dl in the responders. 89% of all KRYSTEXXA®-treated subjects developed ADAs. In comparison, the graph on the right in **Fig. 7** depicts data from Cohort #5 of the Phase 1b clinical trial, which received a single dose of SVP-Rapamycin alone, and Cohort #6 of the Phase 1b clinical trial, which received a single dose of SEL-212. All five subjects in

-59-

Cohort #6 of the Phase 1b clinical trial, treated with SEL-212, maintained levels of serum uric acid of less than 0.1 mg/dl through day 30. Subjects in Cohort #5 of the Phase 1b clinical trial, treated with SVP-Rapamycin alone, experienced no significant reduction in uric acid levels, as such levels remained relatively constant over the 30-day period. Also shown is a comparison of data from Cohort #5 of the Phase 1b clinical trial, which received a single dose of SVP-Rapamycin alone, with Cohort #9 of the Phase 1b clinical trial, which received pegsiticase alone.

While it is believed that the above comparison is useful in evaluating the results of Cohort #6 of the Phase 1b clinical trial, the Phase 1b clinical trial and the KRYSTEXXA® clinical trials were separate trials conducted by different investigators at different sites. In addition, there were substantial differences, including, for example, that the KRYSTEXXA® clinical trials were double-blind trials involving a substantial number of patients with refractory gout while the Phase 1b clinical trial evaluated SEL-212 in an unblinded manner in a small number of subjects with elevated uric acid levels. Moreover, only the efficacy of SEL-212 with the four-week period following the first injection of KRYSTEXXA® could be compared as SEL-212 had not yet been evaluated in a multi-dose clinical trial.

Additional serum uric acid and uricase-specific ADA data was collected after day 30 for three of the subjects in Cohort #4 (SEL-212 Cohort) that had no or very low serum uric acid and uricase-specific ADA levels at day 30. Data was collected on day 37 for all three of these subjects and again on day 42 or day 44 for two of the three subjects. Each of these three subjects had no or very low uricase-specific ADA levels on day 37, day 42 or day 44, as applicable. Serum uric acid levels remained below baseline on day 37 in all three subjects. With respect to the two subjects for which day 42 or day 44 data was available, serum uric acid levels approached or exceeded baseline by the last time point measured.

Example 3 - Phase 2 Clinical Trial

Presented herein is a phase 2 clinical trial of SEL-212. The study consists of multiple doses of SEL-212 concomitantly administered with doses of SEL-037. SEL-212 is a combination of SEL-037 and SEL-110. SEL-037 comprises pegsiticase (Recombinant Pegylated Candida Urate Oxidase). SEL-110 is a nanoparticle comprising PLA (poly(D,L-lactide)) and PLA-PEG (poly(D,L-lactide)-block-poly (ethylene-glycol)) encapsulating rapamycin.

SEL-037 can be provided with phosphate buffer and mannitol as excipients. Prior to

-60-

administration, 6 mg, measured as uricase protein, lyophilized SEL-037 can be reconstituted with 1.1 ml of sterile water for injection, USP (United States Pharmacopeia) which forms a 6 mg/mL concentrated solution. A sufficient volume of reconstituted SEL-037 at 0.2 mg/kg or 0.4 mg/kg, measured as uricase protein, is diluted in 100 mL of 0.9% sodium chloride for injection, USP and dosed as a single intravenous infusion with an infusion pump over 60 minutes.

SEL-110 is provided as a 2 mg/mL, based on rapamycin content, suspension in PBS. The appropriate amount of SEL-110 on a mg/kg basis is drawn into a syringe or syringes and administered as an IV infusion with a syringe infusion pump. If a subject is part of Cohorts 3, 4, 5, 6, 7 and 8 then SEL-110 is administered prior to SEL-037. SEL-110 is delivered by syringe infusion pump at a single steady rate sufficient to deliver the dose volume over a period of 55 minutes concurrently with a 60 minute infusion of 125 mL of normal saline and then the SEL-037 infusion (0.2 mg/kg for Cohorts 3, 5 and 7; 0.4 mg/kg for Cohorts 4, 6 and 8) are started at the 60 minute mark.

48 subjects were divided into 8 dosing cohorts. Each cohort consists of 6 patients. Cohort 3 receives SEL-212 (with 0.05 mg/kg of SEL-110 + 0.2 mg/kg pegsiticase), Cohort 4 receives SEL-212 (with 0.05 mg/kg of SEL-110 + 0.4 mg/kg pegsiticase), Cohort 5 receives SEL-212 (with 0.08 mg/kg of SEL-110 + 0.2 mg/kg pegsiticase), Cohort 6 receives SEL-212 (with 0.08 mg/kg of SEL-110 + 0.4 mg/kg pegsiticase), Cohort 7 receives SEL-212 (with 0.1 mg/kg of SEL-110 + 0.2 mg/kg pegsiticase) and Cohort 8 receives SEL-212 (with 0.1 mg/kg of SEL-110 + 0.4 mg/kg pegsiticase).

Distribution of Subjects

All enrolled subjects were randomized initially to 4 cohorts such that upon reaching 12 subjects total for all 4 cohorts, each cohort contains 3 subjects. After the completion of at least one treatment cycle the subject experience is evaluated before enrollment is opened to all cohorts. The future enrollment is randomized between all open cohorts.

Premedication for Study Drug Treatments

All subjects receive 180 mg fexofenadine orally the night before receiving study drug (12 h $\pm$ 2h) and again 2 $\pm$ 1 hours before receiving study drug (i.e., SEL-110 for Cohorts 3, 4, 5, 6, 7 and 8). In addition, they also receive methylprednisolone 40 mg (or equivalent drug, for example prednisone 50 mg IV or dexamethasone 8 mg IV) intravenously 1 $\pm$ 0.5 hour

-61-

before receiving study drug (i.e. prior SEL-110 for Cohorts 3, 4, 5, 6, 7 and 8). This occurs for every treatment dosing of study drug (Part A, Treatment Periods 1-3 and for Part B, Treatment Periods 4 and 5). Cohorts 3-6 have received first and second doses.

5 **Premedication for Gout Flare**

All subjects that meet all inclusion and exclusion criteria are given premedication for gout flare prevention. The regimen begins 1 week prior to the first dosing of study drug and continue for as long as the subject is enrolled in the clinical study. Subjects are given colchicine 1.2 mg as a single loading dose. Then they will continue with colchicine 0.6 mg
10 QD for the remainder of their participation in the trial. If there is a contraindication to colchicine, the subject receives ibuprofen 600 mg TID or equivalent dose of a NSAID. If there is a contraindication to colchicine and to NSAIDs the subject receives no premedication for gout flare. The gout flare prevention medication continues as long as the subject is enrolled in the clinical study. Subjects who began receiving a NSAID as gout flare prevention
15 medication due to a contraindication to colchicine continue to receive the NSAID as long as the subject is enrolled in the study.

Duration of treatment for Cohort 3, Cohort 4, Cohort 5, Cohort 6, Cohort 7, and Cohort 8

20

Treatment period 1 - Part A

Subjects were screened within 45 days of dosing. Once they met inclusion/exclusion criteria and all assessments were considered acceptable they were instructed on when to start their premedication (date and medication, Day -7) for the prevention of gout flares. The day
25 of initial dosing of study drug was designated Day 0. Eligible subjects who have been assigned to Cohorts 3, 4, 5, 6, 7 and 8 received a single IV infusion of SEL-110 (dose based on a mg/kg basis). SEL-110 was delivered by syringe infusion pump at a single steady rate sufficient to deliver the dose volume over a period of 55 minutes. Concurrently to the administration of SEL-110, the subject received a 125 mL of normal saline over 60 minutes.
30 This was followed ($\pm$ 3 minutes) by an infusion delivered by infusion pump of SEL-037 (0.2 mg/kg for Cohorts 3, 5, and 7; 0.4 mg/kg for Cohorts 4, 6 and 8) diluted into 100 mL of normal saline delivered over 60 minutes. Subjects remained in the clinic for 9 hours after the start of the infusion of SEL-110 for safety evaluations and PK blood draws. Subjects

-62-

returned for PK and PD blood draws on Treatment Period 1, Days 1, 7, 14, 21 and safety and Antibody blood draws on Treatment Period 1, Days 7, 14, 21.

Treatment period 2 - Part A

5 On the morning of Treatment Period 2, Day 0, subjects reported to the clinic for the dosing of study drug. Eligible subjects who had been assigned to Cohorts 3, 4, 5, 6, 7 and 8 received a single IV infusion of SEL-110 (dose based on a mg/kg basis). SEL-110 was delivered by syringe infusion pump at a single steady rate sufficient to deliver the dose volume over a period of 55 minutes. Concurrently to the administration of SEL-110, the
10 subject received a 125 mL of normal saline over 60 minutes. This was followed ($\pm$ 3 minutes) by an infusion delivered by infusion pump of SEL-037 (0.2 mg/kg for Cohorts 3, 5 and 7; 0.4 mg/kg for Cohorts 4, 6 and 8) diluted into 100 mL of normal saline delivered over 60 minutes. Subjects remained in the clinic for 9 hours after the start of the infusion of SEL-110 for safety evaluations and PK blood draws. Subjects returned for PK and PD on
15 Treatment Period 2, Days 1, 7, 14 and 21 and safety and Antibody blood draws on Treatment Period 2, Days 7, 14 and 21.

Treatment period 3 - Part A

20 On the morning of Treatment Period 3, Day 0 subjects will report to the clinic for the dosing of study drug. Eligible subjects who have been assigned to Cohorts 3, 4, 5, 6, 7 and 8 will receive a single IV infusion of SEL-110 (dose based on a mg/kg basis). SEL-110 will be delivered by syringe infusion pump at a single steady rate sufficient to deliver the dose volume over a period of 55 minutes. Concurrently to the administration of SEL-110, the subject will receive a 125 mL of normal saline over 60 minutes. This will be followed ($\pm$ 3
25 minutes) by an infusion delivered by infusion pump of SEL-037 (0.2 mg/kg for Cohorts 3, 5 and 7; 0.4 mg/kg for Cohorts 4, 6 and 8) diluted into 100 mL of normal saline delivered over 60 minutes. Subjects will remain in the clinic for 9 hours after the start of the infusion of SEL-110 for safety evaluations and PK blood draws. Subjects will return for PK and PD blood draws on Treatment Period 3, Days 1, 7, 14 and 21 and safety and Antibody blood
30 draws on Treatment Period 3, Days 7, 14 and 21.

Treatment period 4 - Part B

On the morning of Treatment Period 4, Day 0 subjects will report to the clinic for the

-63-

dosing of study drug. Subjects will receive a single IV infusion of SEL-037 (0.2 mg/kg for Cohorts 3, 5 and 7; 0.4 mg/kg for Cohorts 4, 6 and 8) diluted into 100 mL of normal saline over 60 minutes by infusion pump. Subjects will remain in the clinic for 9 hours after the start of the infusion of SEL-037 for safety evaluations and PK blood draws. Subjects will return for PK and PD blood draws on Treatment Period 4, Days 1, 7, 14 and 21 and safety and Antibody blood draws on Treatment Period 4, Days 7, 14 and 21.

Treatment period 5 - Part B

On the morning of Treatment Period 5, Day 0 subjects will report to the clinic for the dosing of study drug. Subjects will receive a single IV infusion of SEL-037 (0.2 mg/kg for Cohorts 3, 5 and 7; 0.4 mg/kg for Cohorts 4, 6 and 8) diluted into 100 ml of normal saline over 60 minutes by infusion pump. Subjects will remain in the clinic for 9 hours after the start of the infusion of SEL-037 for safety evaluations and PK blood draws. Subjects will return for PK and PD blood draws on Treatment Period 5, Days 1, 7, 14 and 21 and safety and Antibody blood draws on Treatment Period 5, Days 7, 14 and 21.

Results

When pegsiticase was administered alone in the Phase 1 described in **Example 2**, 57% (4 out of 7 patients) of those with a history of gout had signs of gout flare in the first month after receiving the study drug (**Table 1**). In contrast, however, when PLA/PLA-PEG synthetic nanocarriers comprising rapamycin were concomitantly administered with pegsiticase in a Phase 2 trial described in **Example 3**, only one gout flare was reported in the subjects who had a history of gout (16 out of 63 enrolled patients) (**Table 2**). This subject was in the cohort that received only the rapamycin-comprising nanocarrier (without uricase). Because this subject did not receive the uricase therapy, this subject's serum uric acid level did not drop significantly. The flare was, therefore, unrelated to a change in serum uric acid. One additional subject, who did not have a prior diagnosis of gout, reported a post-treatment flare. This patient's serum uric acid level dropped from 8.8 mg/dL to 0.1 mg/dL within 90 minutes following drug administration. So, although this subject had only been diagnosed with asymptomatic hyperuricemia before the study, a flare did seem to coincide with a drop in serum uric acid.

Table 1. Flares in subjects with history of gout

-64-

Subject	Flare in 1 st month	Dose of SEL-037
1	no	0.1 mg/kg
2	yes	0.1 mg/kg
3	yes	0.1 mg/kg
4	yes (positive for tenderness and swelling of right 5 th PIP)	0.1 mg/kg
5	no	0.2 mg/kg
6	yes (mild gout, right 3 rd toe)	0.2 mg/kg
7	no	0.8 mg/kg

Table 2. Flares in SEL-212 subjects

SEL-212 subjects with gout	Flare in 1 st month	Cohort/dose
1	no	Cohort 1/SEL-110 0.03 mg/kg
2	no	Cohort 3/SEL-110 0.1 mg/kg
3	no	Cohort 4/SEL-212 0.1 mg/kg
4	no	Cohort 5/SEL-110 0.3 mg/kg
5	no	Cohort 4/SEL-212 0.1 mg/kg
6	no	Cohort 3/SEL-110 0.1 mg/kg
7	no	Cohort 14/SEL-212 0.1 mg/kg
8	no	Cohort 6/SEL-212 0.3 mg/kg
9	no	Cohort 5/ SEL-110 0.3 mg/kg
10	yes	Cohort 7/ SEL-110 0.5 mg/kg
11	no	Cohort 14/ SEL-212 0.1 mg/kg
12	no	Cohort 14/ SEL-212 0.1 mg/kg
13	no	Cohort 14/ SEL-212 0.1 mg/kg
14	no	Cohort 12/ SEL-212 0.15 mg/kg
15	no	Cohort 12/ SEL-212 0.15 mg/kg
16	no	Cohort 12/ SEL-212 0.15 mg/kg

-65-

17	yes	Cohort 2/ SEL-212 0.03 mg/kg
----	-----	------------------------------

A phase 2 study has been undertaken (**Example 3**). This study involved the administration of multiple IV infusions of PLA/PLA-PEG synthetic nanocarriers comprising rapamycin together with pegsiticase in order to assess its safety and tolerability. Thirty-eight subjects were randomized and dosed, with 8 subjects reported as suffering from a gout flare (**Table 3**).

Table 3. Subjects who suffered from gout flare following treatment

SEL-212 subject	Cohort	Dose	Gout flare prophylaxis with colchicine/NSAID
1	1	SEL-037 0.2 mg/kg	no
2	1	SEL-037 0.2 mg/kg	no
3	1	SEL-037 0.2 mg/kg	no
4	3	SEL-110 0.05 mg/kg + SEL-037 0.2 mg/kg	yes
5	3	SEL-110 0.05 mg/kg + SEL-037 0.2 mg/kg	no
6	4	SEL-110 0.05 mg/kg + SEL-037 0.4 mg/kg	no
7	4	SEL-110 0.05 mg/kg + SEL-037 0.4 mg/kg	no
8	5	SEL-110 0.08 mg/kg + SEL-037 0.2 mg/kg	yes

Flare rates in the above subjects were compared to the flare rates in the pegloticase trials. Those subjects who received gout flare prophylaxis (with colchicine or NSAIDS) only were chosen to match the pegloticase subject conditions. Flare frequency (number of flares per patient month) was selected as a measure by which to compare flare rates. This measure was chosen based on the fact that the trial data covers 2 months, or 2 treatment cycles; while the pegloticase trials varied in length from 35 days (Sundy et al., Pharmacokinetics and pharmacodynamics of intravenous PEGylated recombinant mammalian urate oxidase in

-66-

patients with refractory gout. *Arthritis and Rheumatism*. Vol. 56, No. 3, March 2007, pp 1021-1028) to 6 months (John S. Sundy, MD, PhD; Herbert S. B. Baraf, MD; Robert A. Yood, MD; et al. Efficacy and Tolerability of Pegloticase for the Treatment of Chronic Gout in Patients Refractory to Conventional TreatmentTwo Randomized Controlled Trials. *JAMA*. 2011;306(7):711-720). Patient monthly rates were chosen to be able to compare between trials.

Cohorts 3 and 4 were grouped together for this analysis, as they were given the same dose of synthetic nanocarriers comprising rapamycin (0.05 mg/kg), and likewise cohorts 5 and 6 have been grouped together (with a synthetic nanocarrier comprising rapamycin dose of 0.08 mg/kg). In cohorts 3 and 4, 19 subjects have been dosed for a total of 24 treatment cycles. Not all subjects received all treatments, as certain subjects were discontinued following protocol changes. In cohorts 5 and 6, thus far, 13 subjects have been dosed with a total of 24 treatment cycles. This means that, for subjects who received gout flare prophylaxis, 2 flares in total have occurred over 48 treatment cycles. This can be equated to 0.04 flares per treatment cycle; in other words, a flare frequency of 0.04 flares per patient month.

In contrast, the Phase 3 pegloticase trials (John S. Sundy, MD, PhD; Herbert S. B. Baraf, MD; Robert A. Yood, MD; et al. Efficacy and Tolerability of Pegloticase for the Treatment of Chronic Gout in Patients Refractory to Conventional TreatmentTwo Randomized Controlled Trials. *JAMA*. 2011;306(7):711-720) reported the following: 2.3 flares per patient over the first 3 months for 85 patients who received biweekly pegloticase, and 2.7 flares per patient over the first 3 months for 84 patients who received monthly pegloticase. These numbers equate to a flare frequency of 0.77 and 0.9 flares per patient month, respectively.

Further comparisons can be made with the two primary branded oral uric acid lowering medication, febuxostat and lesinurad. In a phase 3, randomized, double-blind, multi-center trial, the safety and efficacy of febuxostat was studied over 52 weeks (Michael A. Becker, M.D., H. Ralph Schumacher, Jr., M.D., Robert L. Wortmann, M.D., Patricia A. MacDonald, B.S.N., N.P., Denise Eustace, B.A., William A. Palo, M.S., Janet Streit, M.S., and Nancy Joseph-Ridge, M.D. Febuxostat Compared with Allopurinol in Patients with Hyperuricemia and Gout. *N Engl J Med* 2005; 353:2450-2461 December 8, 2005). The comparison period for this analysis included only the first 8 weeks of that study, when gout flare prophylaxis was administered. At a dose of 80 mg/day, 55 out of 255 subjects required

-67-

treatment for at least one gout flare. This would be the equivalent to a flare frequency of at least 0.22 flares per patient month, and possibly more. At a dose of 120 mg/day, 90 out of 250 subjects required treatment for at least one gout flare, equating to at least a flare frequency of 0.36 flares per patient month, and possibly more.

In a phase 2, randomized, double-blind study to assess the efficacy and tolerability of lesinurad, subjects were given colchicine for gout flare prophylaxis and treated with lesinurad at different doses over 1 month (Perez -Ruiz F, Sundry JS, Miner JN for the RDEA594-203 Study Group, et al. Lesinurad in combination with allopurinol: results of a phase 2, randomised, double-blind study in patients with gout with an inadequate response to allopurinol, *Annals of the Rheumatic Diseases* 2016;75:1074-1080). During this treatment period, gout flares requiring treatment were reported in 10 out of 46 patients in a month in those dosed at 200 mg daily, 13 out of 42 patients in a month in those dosed at 400 mg daily, and 15 out of 48 patients in a month in those dosed at 600 mg daily. This equates to a flare frequency of 0.22, 0.31, and 0.31 flares per patient month, respectively.

The tabulated data outlining the comparison in flare frequency between the different medications alongside their efficacy in reducing serum uric acid (sUA) is compiled in **Table 4**.

Table 4. Flares per patient month compared with other uric acid lowering treatments

Medication and dosage	Flares per patient month	Time of efficacy (area under curve of mean sUA levels over time)*
SEL-212 monthly	0.04	25.66
Pegloticase biweekly	0.77	Responders- 26.6 Nonresponders- 84.35
Pegloticase monthly	0.90	Responders-53.55 Nonresponders-99.75
Febuxostat 80 mg/day	0.22	
Febuxostat 120 mg/day	0.36	
Lesinurad 200 mg/day	0.22	158.5
Lesinurad 400 mg/day	0.31	165.8
Lesinurad 600 mg/day	0.31	167.5

* Indicator of efficacy.

-68-

The flare frequency is clearly reduced for the subjects who received the rapamycin-containing nanocarrier concomitantly administered with pegsiticase as compared to all of the other medications. This unexpected outcome is significantly better than with other therapies. This also has the benefit for patient adherence to uric acid lowering therapies, such as uricase, as adherence is greatly reduced when rebound flares occur following initiation of therapy (Treatment of chronic gouty arthritis: it is not just about urate-lowering therapy. Schlesinger N - Semin. Arthritis Rheum. - October 1, 2012; 42 (2); 155-65).

Example 4- Monthly dosing of rapamycin-containing nanocarriers combined with pegylated uricase mitigates formation of anti-drug antibodies resulting in sustained uricase activity in symptomatic gout patients

Background: Gout is caused by the deposition of monosodium urate (MSU) crystals in joints due to chronic hyperuricemia. Long term treatment focuses on reducing sUA levels, thus allowing MSU crystals to dissolve.

Pegylated uricases are therapies for treatment of severe chronic gout. However, uricases are limited by induction of anti-drug antibodies (ADAs) that can compromise efficacy and safety. SEL-212 is a novel combination product consisting of a pegylated uricase co-administered with rapamycin-containing nanocarriers made up of PLA and PLA-PEG (SEL-110). SEL-110 has been designed to mitigate the formation of ADAs by inducing tolerogenic dendritic cells and antigen-specific regulatory T cells. Prolonged therapeutic activity of the uricase plays a role in maintaining a sustained control of serum uric acid (SUA) levels in patients.

Methods: Patients with symptomatic gout (≥ 1 tophus, gout flare within 6 months or gouty arthropathy) and elevated SUA ≥ 6 mg/dL were treated with fixed doses of the pegylated uricase (0.2 mg/kg) in combination with SEL-110 (0.1 and 0.15 mg/kg), or the pegylated uricase (0.2 or 0.4 mg/kg) alone as a control. Patients were infused in 28-day cycles up to 5 times. Safety, tolerability, sUA, ADAs, and uricase activity were monitored.

In this example, “responder” indicates an individual with serum uric acid values < 6 mg/dL and serum uric acid ≤ 1 mg/dL on day 21 (stopping rule). In this example, a “non-responder” is a subject having serum uric acid levels ≥ 6 mg/dL at any time during a 28 day dosing cycle. The “maximum titer” is the maximum titer observed during a 28 day dosing cycle.

-69-

Anti-uricase antibodies were detected using a sandwich ELISA. Uricase was coated onto a plate and anti-uricase IgG was detected using an anti-human IgG secondary antibody. Uricase activity was measured fluorometrically. As uricase converts uric acid into allantoin and H₂O₂, horseradish peroxidase was added and reacted with H₂O₂ and a fluorescent probe to yield a quantifiable signal.

Results: Demographics of 46 patients treated with 28-day cycles x5 combination doses of SEL-110 and the pegylated uricase as compared to 6 patients treated with the pegylated uricase alone was 23-70 years old vs. 41-64 years old (mean 53.6 vs. 51.8 years), male 97.8% vs. 100%, and white 73.9% vs. 33.3%. The mean BMI at baseline was 34.5 vs. 38.9 kg/m². 71.7% vs. 100% of patients were obese with mean duration of established or symptomatic gout as 12.5 vs. 12.8 years. 43 SEL-212 patients were evaluable for this analysis.

The majority of treatment periods with SEL-212 had maximum antibody titers <1080 (83.4%, 111/133) while 14.3% (1/7) treatment periods of the pegylated uricase alone had maximum antibody titers <1080.

For the patients enrolled in cohorts receiving 5 combination doses of rapamycin-containing nanocarriers and the pegylated uricase, the average maximum anti-uricase titer observed during a treatment period was plotted for responders and non-responders (**Fig. 10**). All non-responding treatment periods had titers ≥1080 (11/11). The average maximum anti-uricase titer for non-responders dropped in successive treatment periods and subsequently no patients had titers greater than 1080 in treatment periods 4 and 5.

The 7 dosing cycles with the pegylated uricase alone had an average uricase activity AUC of 3437 mU/dL*wk (**Fig. 11**). For the cohorts with 5 monthly doses of SEL-212, the uricase activity AUC was categorized into individual dosing cycles with maximum anti-uricase titers either above or below a titer of 1:1080. For dosing cycles that had maximum anti-uricase titers greater than 1:1080, the average uricase activity AUC was 6776 mU/dL*wk. For treatment periods with anti-uricase titers less than 1:1080 the average uricase activity AUC was significantly higher (p=0.0029) at 8924 mU/dL*wk. The data indicates that uricase activity AUC is correlated with anti-uricase titer and along with **Fig. 10**, supports that an anti-uricase titer of 1:1080 is clinically relevant.

Patients treated with the pegylated uricase alone developed high titer antibodies and were discontinued from the study within the first two months. Combining the pegylated uricase with increasingly higher doses of rapamycin-containing nanocarriers resulted in better

-70-

anti-uricase titers at 3 months, with 78% of patients having titers $\leq 1:1080$ at 0.15 mg/kg of rapamycin-containing nanocarriers. However, dosing with only with the pegylated uricase in months 4 and 5 led to a drop in patients with titers $\leq 1:1080$ at month 5 (**Fig. 12A**). Five monthly doses of the pegylated uricase and rapamycin-containing nanocarriers resulted in a higher percentage (~66%) of patients with titers $\leq 1:1080$ at month 5 (**Fig. 12B**).

Data from all cohorts was used to plot the maximum anti-uricase titer reached during a dosing cycle against uricase activity AUC (**Figs. 13A-13B**). **Fig. 13A** shows 0.2 mg/kg doses of the pegylated uricase and **Fig. 13B** shows 0.4 mg/kg doses of the pegylated uricase. The plots show that as the anti-uricase titer increases, the uricase activity AUC correspondingly decreases.

Using data from all cohorts, uricase activity AUC showed rapamycin-containing nanocarriers dose dependency, with higher AUC associated with higher rapamycin-containing nanocarrier dose. **Fig. 13C** shows 0.2 mg/kg doses of the pegylated uricase and **Fig. 13D** shows 0.4 mg/kg doses of the pegylated uricase. The relationship between uricase AUC and rapamycin-containing nanocarrier dose appeared to be stronger at a dose of 0.4 mg/kg of the pegylated uricase.

Conclusion: By mitigating the formation of ADAs, monthly dosing of rapamycin-containing nanocarriers combined with the pegylated uricase increases uricase activity in symptomatic gout patients relative to the pegylated uricase alone. When anti-uricase titers are < 1080 , patients show sustained uricase activity, enabling 28 day treatment intervals.

-71-

OTHER EMBODIMENTS

All of the features disclosed in this specification may be combined in any combination. Each feature disclosed in this specification may be replaced by an alternative feature serving the same, equivalent, or similar purpose. Thus, unless expressly stated
5 otherwise, each feature disclosed is only an example of a generic series of equivalent or similar features.

From the above description, one skilled in the art can easily ascertain the essential characteristics of the present invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various
10 usages and conditions. Thus, other embodiments are also within the claims.

-72-

What is claimed is:

1. A method, comprising:
obtaining an anti-uricase antibody level (e.g., a titer) in any one of the subjects
5 provided herein, such as in a sample from the subject, and
assessing the subject for treatment of any one of the conditions provided herein.
2. The method of claim 1, wherein the method further comprises comparing the anti-uricase antibody level in the subject with a threshold.
3. The method of claim 1 or 2, wherein the method further comprises any one of the
10 treatment methods provided herein.
4. The method of any one of claims 1-3, wherein the method further comprises obtaining an anti-uricase antibody level in the subject at a subsequent point in time and assessing a treatment of the subject (e.g., treatment according to any one of the methods provided herein).
- 15 5. A method, comprising:
concomitantly administering to a subject having an anti-uricase antibody level (e.g., a titer) below a threshold, 1) a composition comprising synthetic nanocarriers comprising an immunosuppressant and 2) a composition comprising uricase.
6. The method of claim 5, wherein the subject is identified or monitored to have or
20 maintain an anti-uricase antibody level below the threshold.
7. The method of any one of the preceding claims, wherein the treatment doses and/or frequency is adjusted based on the identification or monitoring.
8. A method of any one of the preceding claims, wherein the method further comprises one or more steps for identifying or monitoring the subject to obtain an anti-uricase antibody
25 level.
9. A method, comprising,
identifying a subject as having an anti-uricase antibody level (e.g., titer) below a threshold and treating the subject.

-73-

10. The method of claim 9, where the treating comprises administering 1) a composition comprising synthetic nanocarriers comprising an immunosuppressant and 2) a composition comprising uricase.

11. The method of claim 9 or 10, wherein the subject is monitored to have or maintain an anti-uricase antibody level below the threshold.

12. The method of any one of claims 9-11, wherein the treatment doses and/or frequency is adjusted based on the monitoring.

13. A method of any one of claims 9-12, wherein the method further comprises one or more steps for monitoring the subject to obtain an anti-uricase antibody level.

14. A method of preventing gout flare, comprising:
concomitantly administering to a subject having an anti-uricase antibody level (e.g., titer) below a threshold, 1) a composition comprising synthetic nanocarriers comprising an immunosuppressant and 2) a composition comprising uricase, wherein the subject is not administered an additional therapeutic to prevent gout flare concomitantly with the concomitant administration.

15. The method of claim 14, wherein the subject is identified or monitored to have or maintain an anti-uricase antibody level below the threshold.

16. The method of claim 14 or 15, wherein the treatment doses and/or frequency is adjusted based on the identification or monitoring.

17. A method of any one of claims 14-16, wherein the method further comprises one or more steps for identifying or monitoring the subject to obtain an anti-uricase antibody level.

18. The method of any one of claims 14-17, wherein the subject is identified as having had or as being expected to have gout flare from treatment with a gout therapy without concomitant administration of an additional therapeutic to prevent gout flare.

19. The method of any one of the preceding claims, wherein the subject is a subject in need thereof.

-74-

20. The method of claim 19, wherein the subject is a subject with an elevated serum uric acid level and/or undesired uric acid deposits.

21. The method of claim 19, wherein the subject has hyperuricemia.

22. The method of any one of the preceding claims, wherein the subject has gout or a
5 condition associated with gout.

23. The method of any one of the preceding claims, wherein the concomitant administration occurs once or more than once in the subject.

24. The method of claim 23, wherein the concomitant administration occurs at least twice in the subject.

10 25. The method of claim 24, wherein the concomitant administration occurs at least thrice in the subject.

26. The method of claim 25, wherein the concomitant administration occurs at least four times in the subject.

27. The method of claim 26, wherein the concomitant administration occurs at least five
15 times, at least six times, at least seven times, at least eight times, at least nine times, or at least ten times in the subject.

28. The method of any one of the preceding claims, wherein the subject is not administered an additional therapeutic to prevent gout flare concomitantly with each concomitant administration.

20 29. The method of any one of the preceding claims, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered concomitantly every two to four weeks.

30. The method of any one of the preceding claims, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising
25 uricase are administered monthly.

31. The method of claim 30, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered

-75-

monthly for at least three months, four months, five months, six months, seven months, eight months, nine months, ten months or more.

32. The method of any one of the preceding claims, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a dose of 0.05 –
5 0.5 mg/kg immunosuppressant with each administration.

33. The method of claim 32, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a dose of 0.05 mg/kg, 0.08 mg/kg, 0.1 mg/kg, 0.125 mg/kg, 0.15 mg/kg, 0.2 mg/kg, 0.25 mg/kg, 0.3 mg/kg, 0.35 mg/kg, 0.4 mg/kg, 0.45 mg/kg, or 0.5 mg/kg immunosuppressant with each administration.

10 34. The method of any one of the preceding claims, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered at a dose of 0.55 – 6.5 mg/kg with each administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant.

35. The method of claim 34, wherein the composition comprising synthetic nanocarriers
15 comprising an immunosuppressant is administered at a dose of 0.55 mg/kg, 0.65 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1.0 mg/kg, 1.1 mg/kg, 1.25 mg/kg, 1.5 mg/kg, 2.0 mg/kg, 2.5 mg/kg, 3.0 mg/kg, 3.5 mg/kg, 4.0 mg/kg, 4.5 mg/kg, 5.0 mg/kg, 5.5 mg/kg, 6.0 mg/kg, or 6.5 mg/kg with each administration, wherein the dose is given as the mg of the synthetic nanocarriers comprising the immunosuppressant.

20 36. The method of any one of the preceding claims, wherein the composition comprising uricase is administered at a dose of 0.1 – 1.2 mg/kg uricase with each administration.

37. The method of claim 36, wherein the composition comprising uricase is administered at a dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1.0 mg/kg, 1.1 mg/kg, 1.2 mg/kg uricase with each administration.

25 38. The method of any one of the preceding claims, wherein a composition comprising uricase is administered to the subject at least once after the one or more concomitant administrations without an immunosuppressant.

-76-

39. The method of any one of the preceding claims, wherein a composition comprising uricase is administered to the subject at least once after the one or more concomitant administrations without synthetic nanocarriers comprising an immunosuppressant.

40. The method of claim 38 or 39, wherein the composition comprising uricase is administered at least twice after the one or more concomitant administrations.

41. The method of claim 40, wherein the composition comprising uricase is administered at least thrice after the one or more concomitant administrations.

42. The method of claim 41, wherein the composition comprising uricase is administered four, five or more times after the one or more concomitant administrations.

43. The method of claim 39 or 40, wherein the composition comprising uricase is administered monthly for two months after the one or more concomitant administrations.

44. The method of any one of claims 38-43, wherein the composition comprising uricase is administered at a dose of 0.1 – 1.2 mg/kg uricase with each administration after the one or more concomitant administrations without an immunosuppressant.

45. The method of claim 44, wherein the composition comprising uricase is administered at a dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1.0 mg/kg, 1.1 mg/kg, 1.2 mg/kg uricase with each administration after the one or more concomitant administrations without an immunosuppressant.

46. The method of any one of the preceding claims, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant is administered prior to the composition comprising uricase with each concomitant administration.

47. The method of any one of the preceding claims, wherein the composition comprising synthetic nanocarriers comprising an immunosuppressant and the composition comprising uricase are administered within an hour of each other with each concomitant administration.

48. The method of any one of the preceding claims, wherein the subject has acute gout; chronic gout with or without tophi; idiopathic gout; refractory gout, such as chronic refractory gout; secondary gout; unspecified gout; gout associated with a cardiovascular

-77-

condition, renal condition, pulmonary condition, neurological condition, ocular condition, dermatological condition or hepatic condition; or has had a gout attack or gout flare.

49. The method of any one of the preceding claims, wherein the uricase is pegylated uricase.

5 50. The method of claim 49, wherein the pegylated uricase is pegsiticase or pegloticase.

51. The method of claim 50, wherein the pegylated uricase is pegsiticase.

52. The method of any one of the preceding claims, wherein the immunosuppressant is an mTOR inhibitor.

53. The method of claim 52, wherein the mTOR inhibitor is a rapalog.

10 54. The method of claim 53, wherein the rapalog is rapamycin.

55. The method of any one of the preceding claims, wherein the immunosuppressant is encapsulated in the synthetic nanocarriers.

56. The method of any one of the preceding claims, wherein the synthetic nanocarriers are polymeric synthetic nanocarriers.

15 57. The method of claim 56, wherein the polymeric synthetic nanocarriers comprise a hydrophobic polyester.

58. The method of claim 57, wherein the hydrophobic polyester comprises PLA, PLG, PLGA or polycaprolactone.

59. The method of claim 57 or 58, wherein the polymeric synthetic nanocarriers further
20 comprise PEG.

60. The method of claim 59, wherein the PEG is conjugated to the PLA, PLG, PLGA or polycaprolactone.

61. The method of claim 60, wherein the polymeric synthetic nanocarriers comprise PLA, PLG, PLGA or polycaprolactone and PEG conjugated to PLA, PLG, PLGA or
25 polycaprolactone.

-78-

62. The method of claim 61, wherein the polymeric synthetic nanocarriers comprise PLA and PLA-PEG.

63. The method of any one of claims 56-62, wherein the mean of a particle size distribution obtained using dynamic light scattering of the synthetic nanocarriers is a diameter
5 greater than 120nm.

64. The method of claim 63, wherein the diameter is greater than 150nm.

65. The method of claim 64, wherein the diameter is greater than 200nm.

66. The method of claim 65, wherein the diameter is greater than 250nm.

67. The method of any one of claims 63-66, wherein the diameter is less than 300nm.

10 68. The method of any one of claims 63-66, wherein the diameter is less than 250nm.

69. The method of claim 63 or 64, wherein the diameter is less than 200nm.

70. The method of any one of the preceding claims, wherein the load of the immunosuppressant of the synthetic nanocarriers is 7-12% or 8-12% by weight.

15 71. The method of claim 70, wherein the load of the immunosuppressant of the synthetic nanocarriers is 7-10% or 8-10% by weight.

72. The method of claim 70, wherein the load of the immunosuppressant of the synthetic nanocarriers is 7%, 8%, 9%, 10%, 11%, or 12% by weight.

73. The method of any one of the preceding claims, wherein each administration of each composition is an intravenous administration.

20 74. The method of claim 73, wherein the intravenous administration is an intravenous infusion.

75. The method of any one of the preceding claims, wherein the method further comprises administering an additional therapeutic to the subject.

25 76. The method of claim 75, wherein the additional therapeutic is an oral gout therapeutic.

-79-

77. The method of claim 75 or 76, wherein the additional therapeutic is administered at least once after the one or more concomitant administrations without an immunosuppressant.

78. The method of claim 75 or 76, wherein the additional therapeutic is administered at least once after the one or more concomitant administrations without synthetic nanocarriers
5 comprising an immunosuppressant.

79. The method of claim 78, wherein the additional therapeutic is administered at least twice, at least thrice, four times, five times or more.

80. The method of claim 75, wherein the additional therapeutic is an anti-gout flare treatment.

10 81. The method of claim 80, wherein the anti-gout flare treatment is a prophylactic treatment administered concomitantly but prior to the administration of each uricase composition.

82. The method of claim 80 or 81, wherein the anti-gout flare treatment is colchicine or an NSAID.

15 83. A method, comprising:
administering to any one of the subjects described herein having an anti-uricase antibody level (e.g., titer) below a threshold, a composition comprising uricase at any one of the doses provided herein and a pharmaceutically acceptable carrier one or more times.

84. The method of claim 83, wherein the subject is identified or monitored to have or
20 maintain an anti-uricase antibody level below the threshold.

85. The method of claim 83 or 84, wherein the treatment doses and/or frequency is adjusted based on the identification or monitoring.

86. A method of any one of claims 83-85, wherein the method further comprises one or more steps for identifying or monitoring the subject to obtain an anti-uricase antibody level.

25 87. The method of any one of claims 83-86, wherein at least one administration or each administration is via a non-intramuscular mode of administration.

88. The method of any one of claims 83-87, wherein the uricase is pegylated uricase.

-80-

89. The method of claim 88, wherein the pegylated uricase is pegsiticase or pegloticase.

90. The method of claim 89, wherein the pegylated uricase is pegsiticase.

91. The method of any one of claims 83-90, wherein at least one administration or each administration is an intravenous administration.

5 92. The method of claim 91, wherein the intravenous administration is an intravenous infusion.

93. The method of any one of claims 83-92, wherein the composition comprising a uricase and a pharmaceutically acceptable carrier is administered at least twice, at least thrice, four times, five times or more to the subject.

10 94. The method of any one of claims 83-93, wherein the composition comprising uricase and a pharmaceutically acceptable carrier is administered every two to four weeks.

95. The method of claim 94, wherein the composition comprising uricase and a pharmaceutically acceptable carrier is administered monthly.

15 96. The method of any one of claims 83-95, wherein the composition comprising uricase and a pharmaceutically acceptable carrier is administered concomitantly with a composition comprising an immunosuppressant.

97. A method, comprising
concomitantly administering to a subject having an anti-uricase antibody level (e.g., titer) below a threshold, 1) a composition comprising polymeric synthetic nanocarriers
20 comprising PLA, PLA-PEG, and rapamycin; and 2) a composition comprising uricase, wherein the composition comprising polymeric synthetic nanocarriers comprising PLA, PLA-PEG, and rapamycin is administered at a dose of 0.05 mg/kg – 0.3 mg/kg rapamycin and the dose of the composition comprising uricase is 0.1 mg/kg - 0.5 mg/kg.

25 98. The method of claim 97, wherein the subject is identified or monitored to have or maintain an anti-uricase antibody level below the threshold.

99. The method of claim 97 or 98, wherein the treatment doses and/or frequency is adjusted based on the identification or monitoring.

-81-

100. A method of any one of claims 97-99, wherein the method further comprises one or more steps for identifying or monitoring the subject to obtain an anti-uricase antibody level.

101. The method of any one of claims 97-100, wherein the concomitant administration occurs at least twice or thrice in the subject.

5 102. The method of any one of claims 97-101, wherein the composition comprising polymeric synthetic nanocarriers comprising PLA, PLA-PEG, and rapamycin and the composition comprising uricase are administered concomitantly every two to four weeks.

103. The method of claim 102, wherein the composition comprising polymeric synthetic nanocarriers comprising PLA, PLA-PEG, and rapamycin and the composition comprising
10 uricase are administered monthly.

104. The method of any one of claims 97-103, wherein the composition comprising polymeric synthetic nanocarriers comprising PLA, PLA-PEG, and rapamycin is administered at a dose of 0.05 mg/kg 0.08 mg/kg, 0.1 mg/kg, 0.125 mg/kg, 0.15 mg/kg, 0.2 mg/kg, 0.25 mg/kg, or 0.3 mg/kg rapamycin.

15 105. The method of any one of claims 97-104, wherein the composition comprising uricase is administered at a dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, or 0.5 mg/kg.

106. The method of any one of claims 97-105, wherein the uricase is pegylated uricase.

107. The method of claim 106, wherein the pegylated uricase is pegsiticase or pegloticase.

108. The method of claim 107, wherein the pegylated uricase is pegsiticase.

20 109. The method of any one of claims 97-108, wherein the subject is a subject in need thereof.

110. The method of claim 109, wherein the subject has an elevated serum uric acid level.

111. The method of claim 109, wherein the subject has gout or a condition associated with gout.

25 112. The method of claim 109, wherein the subject is expected to have gout flare from treatment with a gout therapy.

-82-

113. A method, comprising:

concomitantly administering to a subject having an anti-uricase antibody level (e.g., titer) below a threshold, 1) a composition comprising polymeric synthetic nanocarriers comprising rapamycin; and 2) a composition comprising pegsiticase, wherein the
5 composition comprising polymeric synthetic nanocarriers is administered at a dose of 0.05 mg/kg – 0.3 mg/kg rapamycin and the dose of the composition comprising pegsiticase is 0.1 mg/kg - 0.5 mg/kg pegsiticase.

114. The method of claim 113, wherein the subject is identified or monitored to have or maintain an anti-uricase titer below the threshold.

10 115. The method of claim 113 or 114, wherein the treatment doses and/or frequency is adjusted based on the identification or monitoring.

116. A method of any one of claims 113-115, wherein the method further comprises one or more steps for identifying or monitoring the subject to obtain an anti-uricase antibody level.

117. The method of claims 113-116, wherein the concomitant administration occurs at
15 least twice or thrice in the subject.

118. The method of any one of claims 113-117, wherein the composition comprising polymeric synthetic nanocarriers comprising rapamycin and the composition comprising pegsiticase are administered concomitantly every two to four weeks.

119. The method of claim 118, wherein the composition comprising polymeric synthetic
20 nanocarriers comprising rapamycin and the composition comprising pegsiticase are administered monthly.

120. The method of any one of claims 113-119, wherein the composition comprising polymeric synthetic nanocarriers comprising rapamycin is administered at a dose of 0.05 mg/kg 0.08 mg/kg, 0.1 mg/kg, 0.125 mg/kg, 0.15 mg/kg, 0.2 mg/kg, 0.25 mg/kg, or 0.3
25 mg/kg rapamycin.

121. The method of any one of claims 113-120, wherein the composition comprising pegsiticase is administered at a dose of 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, or 0.5 mg/kg pegsiticase.

-83-

122. The method of any one of claims 113-121, wherein the polymeric synthetic nanocarriers comprise a hydrophobic polyester.

123. The method of claim 122, wherein the hydrophobic polyester comprises PLA, PLG, PLGA or polycaprolactone.

5 124. The method of claim 122 or 123, wherein the polymeric synthetic nanocarriers further comprise PEG.

125. The method of claim 124, wherein the PEG is conjugated to the PLA, PLG, PLGA or polycaprolactone.

10 126. The method of claim 125, wherein the polymeric synthetic nanocarriers comprise PLA, PLG, PLGA or polycaprolactone and PEG conjugated to PLA, PLG, PLGA or polycaprolactone.

127. The method of claim 126, wherein the polymeric synthetic nanocarriers comprise PLA and PLA-PEG.

15 128. The method of any one of claims 113-127, wherein the subject is a subject in need thereof.

129. The method of claim 128, wherein the subject has an elevated serum uric acid level.

130. The method of claim 128, wherein the subject has gout or a condition associated with gout.

20 131. The method of claim 128, wherein the subject is expected to have gout flare from treatment with a gout therapy.

132. The method of any one of the preceding claims, wherein the subject is any one of the subjects provided herein.

133. The method of any one of the preceding claims, wherein the threshold is an anti-uricase titer of less than 1080.

25 134. The method of any one of the preceding claims, wherein the threshold is an anti-uricase titer of 1080.

1/12




Fig. 1




Fig. 2

2/12




Fig. 3

Serum Uric Acid Levels After Single IV Dose of Pegsiticase
Across Five Cohorts




Fig. 4

3/12

Uricase-Specific ADA Titer and Serum Uric Acid Levels										
Subject number	Baseline		Day 7		Day 14		Day 21		Day 30	
	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)
1	7.4	Neg	<0.1	Neg	5	9720	6	N.A.	6.9	3240
2	7.5	Neg	<0.1	40	<0.1	40	<0.1	N.A.	0.4	40
3	7.3	120	<0.1	120	6.9	9720	7.6	N.A.	7.6	3240
4	7.6	Neg	<0.1	Neg	6.1	3240	7.5	N.A.	7.6	1080
5	4.9	Neg	<0.1	Neg	<0.1	1080	0.3	N.A.	5.1	1080

Phase 1a Cohort #3

Uricase-Specific ADA Titer and Serum Uric Acid Levels										
Subject number	Baseline		Day 7		Day 14		Day 21		Day 30	
	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)
1	5.4	Neg	<0.1	N.A.	5.6	1080	5.8	1080	7	1080
2	6.3	Neg	<0.1	N.A.	5.8	29160	5.5	29160	6	9720
3	7.4	Neg	<0.1	N.A.	<0.1	3240	<0.1	1080	1.9	1080
4	7.2	Neg	<0.1	N.A.	3.2	3240	7	3240	6.3	1080
5	8.1	Neg	<0.1	N.A.	<0.1	29160	7.8	9720	8.8	9720

Phase 1b Cohort #9

Fig. 5

4/12

Uricase-Specific ADA Titer and Serum Uric Acid Levels										
Subject number	Baseline		Day 7		Day 14		Day 21		Day 30	
	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)
1	6.7	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	Neg
2	5.8	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	Neg
3	7.3	Neg	<0.1	N.A.	<0.1	1080	4.8	29160	6.1	29160
4	6.2	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	120
5	5.5	Neg	<0.1	N.A.	<0.1	40	<0.1	Neg	<0.1	Neg

Phase 1b Cohort #4

Uricase-Specific ADA Titer and Serum Uric Acid Levels										
Subject number	Baseline		Day 7		Day 14		Day 21		Day 30	
	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)	Uric acid (mg/dl)	ADA (Titer)
1	7	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	Neg
2	7.4	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	Neg
3	7.5	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	Neg
4	5.6	120	<0.1	N.A.	<0.1	120	<0.1	120	<0.1	120
5	5.9	Neg	<0.1	N.A.	<0.1	Neg	<0.1	Neg	<0.1	Neg

Phase 1b Cohort #6

(Neg = Negative; N.A. = Sample not available)

Fig. 5 (Continued)

5/12





Fig. 6


Phase 1a Cohort #3 and Phase 1b Cohort #9	Phase 1b Cohorts #1, #2, #3, #4, #5 and #6
---	--



Data from the Phase 1b single dose clinical study of SEL-212

Data excerpted from the monthly dosing cohorts of the Krystexxa Phase 3 clinical studies
Sundy et al., JAMA, 2011 Vol 306, p717

Fig. 7



Day 30 sUA levels and ADA levels

Pegsiticase alone (0.4 mg/kg)			SEL-212 0.1 mg/kg SVP-Rapamycin + 0.4 mg/kg Pegsiticase			SEL-212 0.3 mg/kg SVP-Rapamycin + 0.4 mg/kg Pegsiticase		
Subject number	Day 30		Subject number	Day 30		Subject number	Day 30	
	Uric acid (mg/dL)	ADA (Titer)		Uric acid (mg/dL)	ADA (Titer)		Uric acid (mg/dL)	ADA (Titer)
108-0010	7	1080	107-0018	<0.1	Neg	107-0027	<0.1	Neg
103-0015	6	9720	107-0021	<0.1	Neg	107-0028	<0.1	Neg
104-0032	1.9	1080	104-0027	6.1	29160	104-0050	<0.1	Neg
109-0012	6.3	1080	108-0008	<0.1	120	104-0060	<0.1	120
104-0036	8.8	9720	102-0005	<0.1	Neg	103-0019	<0.1	Neg

(Neg = Negative)

Fig. 8

8/12

Cohort	SEL-110	SEL-037
1	NA	0.2 mg/kg
2	NA	0.4 mg/kg
3	0.05mg/kg	0.2 mg/kg
4	0.05mg/kg	0.4 mg/kg
5	0.08mg/kg	0.2 mg/kg
6	0.08mg/kg	0.4 mg/kg

Fig. 9

9/12




Fig. 10




Fig. 11




Fig. 12A




Fig. 12B

11/12




Fig. 13A




Fig. 13B

12/12




Fig. 13C




Fig. 13D

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2020/036116

A. CLASSIFICATION OF SUBJECT MATTER

INV. G01N33/62 A61K31/436 A61K38/44 A61P19/06 G01N33/68
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

G01N A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Rehan Azeem ET AL: "Initial Phase 2 Clinical Data of SEL-212 in Symptomatic Gout Patients: Monthly Dosing of a Pegylated Uricase (Pegadricase) with Svp-Rapamycin Enables Sustained Reduction of Acute Gout Flares", Arthritis Rheumatol., 22 October 2018 (2018-10-22), XP055730329, Retrieved from the Internet: URL:https://acrabstracts.org/abstract/initial-phase-2-clinical-data-of-sel-212-in-symptomatic-gout-patients-monthly-dosing-of-a-pegylated-uricase-pegadricase-with-svp-rapamycin-enables-sustained-reduction-of-acute-gout-flares/ [retrieved on 2020-09-14] pg 1, para 3 ; fig 1 ----- -/--	1-33, 35-123, 125-134



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

16 September 2020

Date of mailing of the international search report

24/09/2020

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Vadot-Van Geldre, E

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2020/036116

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Earl Sands ET AL: "Update of SEL-212 Phase 2 Clinical Data in Symptomatic Gout Patients: Svp-Rapamycin Combined with Pegadricase Mitigates Immunogenicity and Enables Sustained Reduction of Serum Uric Acid Levels, Low Rate of Gout Flares and Monthly Dosing", 1 October 2018 (2018-10-01), XP055730509, Retrieved from the Internet: URL:https://acrabstracts.org/abstract/update-of-sel-212-phase-2-clinical-data-in-symptomatic-gout-patients-svp-rapamycin-combined-with-pegadricase-mitigates-immunogenicity-and-enables-sustained-reduction-of-serum-uric-acid-levels-low-ra/ [retrieved on 2020-09-14] pg 1, last para -----	1-33, 35-123, 125-134
X	US 2018/289776 A1 (JOHNSTON LLOYD [US]) 11 October 2018 (2018-10-11) Figures ; claims ; para 12-13, 117, 157, 159-160, 170, 178 -----	1-134
X	US 9 377 454 B2 (ROSARIO-JANSEN THERESA [US]; WRIGHT DAVID ERICK [US] ET AL.) 28 June 2016 (2016-06-28) col 7, l 1-5 -----	1-4
A	SUSHRUSHA NAYAK ET AL: "Prevention and Reversal of Antibody Responses Against Factor IX in Gene Therapy for Hemophilia B", FRONTIERS IN MICROBIOLOGY, vol. 2, 1 January 2011 (2011-01-01), XP055730478, Lausanne ISSN: 1664-302X, DOI: 10.3389/fmicb.2011.00244 abstract ; Fig 1-2 ----- -/--	1-134

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2020/036116

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	Wesley Dehaan ET AL: "Monthly Dosing of ImmTOR Tolerogenic Nanoparticles Combined with Pegylated Uricase (Pegadricase) Mitigates Formation of Anti-Drug Antibodies Resulting in Sustained Uricas", American College of Rheumatology (ACR) and Association for Rheumatology Professionals (ARP) Annual Meeting, 1 November 2019 (2019-11-01), XP055730470, Retrieved from the Internet: URL:https://acrabstracts.org/abstract/monthly-dosing-of-immTOR-tolerogenic-nanoparticles-combined-with-pegylated-uricase-pegadricase-mitigates-formation-of-anti-drug-antibodies-resulting-in-sustained-uricase-activity-in-symptomatic-gout/ [retrieved on 2020-09-14] pg 1, para bridging pg 2 ; Fig -----	1-134

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2020/036116

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2018289776 A1	11-10-2018	AU 2018236123 A1 BR 112019018748 A2 CA 3055936 A1 CN 110612122 A EP 3592389 A1 JP 2020510687 A KR 20190124295 A US 2018289776 A1 WO 2018169811 A1	03-10-2019 07-04-2020 20-09-2018 24-12-2019 15-01-2020 09-04-2020 04-11-2019 11-10-2018 20-09-2018
US 9377454 B2	28-06-2016	AU 2010265964 A1 BR PI1010069 A2 CA 2766309 A1 CN 102803954 A EP 2446260 A1 EP 3482768 A1 IL 216956 A JP 2012531596 A JP 2015111137 A KR 20120046125 A MX 347903 B NZ 597089 A RU 2012102399 A SG 176897 A1 SG 10201408480R A US 2012301454 A1 US 2016377604 A1 US 2018188242 A1 US 2019317083 A1 WO 2010151823 A1	02-02-2012 15-03-2016 29-12-2010 28-11-2012 02-05-2012 15-05-2019 31-10-2016 10-12-2012 18-06-2015 09-05-2012 18-05-2017 30-05-2014 27-07-2013 30-01-2012 27-02-2015 29-11-2012 29-12-2016 05-07-2018 17-10-2019 29-12-2010