

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



(10) International Publication Number

WO 2017/011820 A2

(43) International Publication Date  
19 January 2017 (19.01.2017)

(51) International Patent Classification:  
*C07K 7/08* (2006.01)      *C07K 7/06* (2006.01)

Lassen Avenue, Mountain View, California 94043 (US).  
**FREDERICK, Brian Troy**; 8470 Glen Arbor Road, Ben Lomond, California 98005 (US). **ZHANG, Jie**; 101 Ness Road, Salisbury, Queensland 4107 (AU). **PATEL, Dinesh V.**; 45109 Cougar Circle, Fremont, California 94539 (US). **LIU, David**; 201 Ferne Avenue, Palo Alto, California 94306 (US).

(21) International Application Number:  
PCT/US2016/042680

(22) International Filing Date:  
15 July 2016 (15.07.2016)

(25) Filing Language: English

(74) Agents: **LAHERTY, Carol** et al.; Cooley LLP, 1299 Pennsylvania Avenue, NW, Suite 700, Washington, District of Columbia 20004 (US).

(26) Publication Language: English

(30) Priority Data:  
*PCT/US15/40658* 15 July 2015 (15.07.2015) US  
14/800,627 15 July 2015 (15.07.2015) US  
62/264,820 8 December 2015 (08.12.2015) US  
62/281,123 20 January 2016 (20.01.2016) US

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant: **PROTAGONIST THERAPEUTICS, INC.** [US/US]; 521 Cottonwood Drive, Suite A, Milpitas, California 95035 (US).


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,

(72) Inventor; and  
(71) Applicant : **BHANDARI, Ashok** [US/US]; 2629 Lotus Street, Pleasanton, California 94588 (US).

(72) Inventors: **BOURNE, Gregory**; 29 Moolingal Street, Jindalee, Queensland 4074 (AU). **CHENG, XiaoLi**; 263

[Continued on next page]

(54) Title: PEPTIDE INHIBITORS OF INTERLEUKIN-23 RECEPTOR AND THEIR USE TO TREAT INFLAMMATORY DISEASES



(57) Abstract: The present invention provides novel peptide inhibitors of the interleukin-23 receptor, and related compositions and methods of using these peptide inhibitors to treat or prevent a variety of diseases and disorders, including inflammatory bowel diseases.

FIG. 13A

**Published:**

- TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).
- *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*
- *with sequence listing part of description (Rule 5.2(a))*

**PEPTIDE INHIBITORS OF INTERLEUKIN-23 RECEPTOR AND THEIR USE  
TO TREAT INFLAMMATORY DISEASES**

**CROSS REFERENCE TO RELATED APPLICATIONS**

[0001] This application claims priority to and is a continuation-in-part of U.S. Application No. 14/800,627 filed on July 15, 2015, and also claims priority to International Patent Application PCT/US2015/040658 filed on July 15, 2015, U.S. Provisional Application No. 62/264,820 filed on December 8, 2015, and U.S. Provisional Application No. 62/281,123 filed on January 20, 2016, all of which are incorporated by reference herein in theirentireties.

**SEQUENCE LISTING**

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on July 15, 2016, is named PRTH-002-03WO\_SL.txt and is 504 KB in size

**FIELD OF THE INVENTION**

[0003] The present invention relates to novel peptide inhibitors of the interleukin-23 receptor, and their use to treat or prevent a variety of diseases and disorders, including inflammatory bowel disease, Crohn's disease and psoriasis.

**BACKGROUND**

[0004] The interleukin-23 (IL-23) cytokine has been implicated as playing a crucial role in the pathogenesis of autoimmune inflammation and related diseases and disorders, such as multiple sclerosis, asthma, rheumatoid arthritis, psoriasis, and inflammatory bowel diseases (IBDs), e.g., ulcerative colitis and Crohn's disease. Studies in acute and chronic mouse models of IBD revealed a primary role of IL-23R and downstream effector cytokines in disease pathogenesis. IL-23R is expressed on various adaptive and innate immune cells including Th17 cells,  $\gamma\delta$  T cells, natural killer (NK) cells, dendritic cells, macrophages, and innate lymphoid cells, which are found abundantly in the intestine. At the intestine mucosal surface, the gene expression and protein levels of IL-23R are found to be elevated in IBD patients. It is believed that IL-23

mediates this effect by promoting the development of a pathogenic CD4<sup>+</sup> T cell population that produces IL-6, IL-17, and tumor necrosis factor (TNF).

**[0005]** Production of IL-23 is enriched in the intestine, where it is believed to play a key role in regulating the balance between tolerance and immunity through T-cell-dependent and T-cell-independent pathways of intestinal inflammation through effects on T-helper 1 (Th1) and Th17-associated cytokines, as well as restraining regulatory T-cell responses in the gut, favoring inflammation. In addition, polymorphisms in the IL-23 receptor (IL-23R) have been associated with susceptibility to IBDs, further establishing the critical role of the IL-23 pathway in intestinal homeostasis.

**[0006]** Psoriasis, a chronic skin disease affecting about 2%-3% of the general population has been shown to be mediated by the body's T cell inflammatory response mechanisms. IL-23 has one of several interleukins implicated as a key player in the pathogenesis of psoriasis, purportedly by maintaining chronic autoimmune inflammation via the induction of interleukin-17, regulation of T memory cells, and activation of macrophages. Expression of IL-23 and IL-23R has been shown to be increased in tissues of patients with psoriasis, and antibodies that neutralize IL-23 showed IL-23-dependent inhibition of psoriasis development in animal models of psoriasis.

**[0007]** IL-23 is a heterodimer composed of a unique p19 subunit and the p40 subunit of IL-12, which is a cytokine involved in the development of interferon- $\gamma$  (IFN- $\gamma$ )-producing T helper 1 (T<sub>H</sub>1) cells. Although IL-23 and IL-12 both contain the p40 subunit, they have different phenotypic properties. For example, animals deficient in IL-12 are susceptible to inflammatory autoimmune diseases, whereas IL-23 deficient animals are resistant, presumably due to a reduced number of CD4<sup>+</sup> T cells producing IL-6, IL-17, and TNF in the CNS of IL-23-deficient animals. IL-23 binds to IL-23R, which is a heterodimeric receptor composed of IL-12R $\beta$ 1 and IL-23R subunits. Binding of IL-23 to IL-23R activates the Jak-stat signaling molecules, Jak2, Tyk2, and Stat1, Stat 3, Stat 4, and Stat 5, although Stat4 activation is substantially weaker and different DNA-binding Stat complexes form in response to IL-23 as compared with IL-12. IL-23R associates constitutively with Jak2 and in a ligand-dependent manner with Stat3. In contrast to

IL-12, which acts mainly on naive CD4(+) T cells, IL-23 preferentially acts on memory CD4(+) T cells.

**[0008]** Efforts have been made to identify therapeutic moieties that inhibit the IL-23 pathway, for use in treating IL-23-related diseases and disorders. A number of antibodies that bind to IL-23 or IL-23R have been identified, including ustekinumab, a humanized antibody that binds IL-23, which has been approved for the treatment of psoriasis. More recently, polypeptide inhibitors that bind to IL-23R and inhibit the binding of IL-23 to IL-23R have been identified (see, e.g., US Patent Application Publication No. US2013/0029907). Clinical trials in Crohn's Disease or psoriasis with ustekinumab and briakinumab (which target the common p40 subunit) and tildrakizumab, guselkumab, MEDI2070, and BI-655066 (which target the unique p19 subunit of IL-23) highlight the potential of IL-23 signaling blockade in treatment of human inflammatory diseases. While these findings are promising, challenges remain with respect to identifying stable and selective agents that preferentially target the IL-23 pathway in the intestine, which can be used for the treatment of intestinal inflammation, such as intestinal bowel diseases, including Crohn's disease, ulcerative colitis and related disorders.

**[0009]** Clearly, there remains a need in the art for new therapeutics targeting the IL-23 pathway, which may be used to treat and prevent IL-23-associated diseases, including those associated with autoimmune inflammation in the intestinal tract. In addition, compounds and methods for specific targeting of IL-23R from the luminal side of the gut may provide therapeutic benefit to IBD patients suffering from local inflammation of the intestinal tissue. The present invention addresses these needs by providing novel peptide inhibitors that bind IL-23R to inhibit IL-23 binding and signaling and which are suitable for oral administration.

#### BRIEF SUMMARY OF THE INVENTION

**[0010]** The present invention provides *inter alia* novel peptide inhibitors of IL-23R and related methods of use.

In a first aspect, the present invention provides a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Xa): X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Xa),

wherein:

X1, X2 and X3 are any amino acid or absent

X4 is any amino acid or chemical moiety capable of forming a bond with X9;

X5, X6, X7 and X8 are any amino acid;

X9 is any amino acid or chemical moiety capable of forming a bond with X4;

X10, X11, X12, X13, X14 and X15 are any amino acid; and

X16, X17, X18, X19 and X20 are any amino acid or absent;

wherein the peptide inhibitor is cyclized via a bond between X4 and X9, and

wherein the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

**[0011]** In certain embodiments of Xa:

**[0012]** X1 is absent; X2 is absent; X3 is absent; X4 is Cys, Abu or Pen; X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn, Gln, Arg, Ser or Thr; X6 is Asp or Thr; X7 is Trp or 6-Chloro-Trp; X8 is Glu, Gln or Val; X9 is Cys, Abu or Pen; X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog; X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl<sub>2</sub>), Trp or Tyr(3-tBu); X12 is 3-Pal, Acpc, Acbc, Acvc, Achc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cav, Cha, Cit, Cpa, D-Asn, Glu, His, hLeu, hArg, Lys, Leu, Octgly, Orn, 4-amino-4-carboxy-piperidine, Arg, Ser, Thr or THP; X13 is Cit, Asp, Dab, Dap, Phe, His, Dap(Peg2-Ac), Dap(pyroglutaric acid), Glu, HomoArg, Lys, Lys(Ac), Lys(Benzoic acid), Lys(glutaric acid), Lys(IVA), Lys(Peg4-isoGlu-Palm), Lys(pyroglutaric acid), Lys(succinic acid), Asn, Orn, Gln, Arg, Thr or Val; X14 is Asp, Dab(Ac), Dap(Ac), Phe, His, Lys(Ac), Met, Asn(isobutyl), Gln, Arg, Tyr or Asp(1,4-diaminobutane); and X15 is Ala,  $\beta$ Ala, Glu, Gly, Asn, Gln, Arg or Ser.

**[0013]** In certain embodiments of Xa: X1 is absent; X2 is absent; X3 is absent; X4 is Cys, Abu or Pen; X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, Orn, Gln, Arg,

Ser or Thr; X6 is Asp or Thr; X7 is Trp or 6-Chloro-Trp; X8 is Gln or Val; X9 is Cys, Abu or Pen; X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog; X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl<sub>2</sub>), Trp or Tyr(3-tBu); X12 is 3-Pal, Acpc, Acbc, Acvc, Achc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cav, Cha, Cit, Cpa, D-Asn, His, hLeu, hArg, Lys, Leu, Octgly, Orn, 4-amino-4-carboxy-piperidine, or THP; X13 is Cit, Asp, Dab, Dap, Phe, His, Dap(Peg2-Ac), Dap(pyroglutaric acid), Glu, hArg, Lys, Lys(Ac), Lys(Benzoic acid), Lys(glutaric acid), Lys(IVA), Lys(Peg4-isoGlu-Palm), Lys(pyroglutaric acid), Lys-(succinic acid), Asn, Orn, Gln, Arg, Thr or Val; X14 is Dab(Ac), Dap(Ac), Phe, His, Lys(Ac), Met, Asn, Gln, Arg, or Tyr; and X15 is Ala, betaAla, Gly, Asn, Gln, or Ser.

**[0014]** In certain embodiments of Xa: X1 is absent; X2 is absent; X3 is absent; X4 is Cys, Abu or Pen; X5 is Dap, Dap(Ac), Gly, Lys, Gln, Arg, Ser, Thr or Asn; X6 is Thr; X7 is Trp or 6-Chloro-Trp; X8 is Gln; X9 is Cys, Abu or Pen; X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog; X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), Phe(3,4-Cl<sub>2</sub>), or Trp; X12 is Acpc, Acbc, Acvc, Achc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, hLeu, Lys, Leu, Arg or THP; X13 is Cit, Asp, Dap, Dap(Peg2-Ac), Dap(pyroglutaric acid), Glu, hArg, Lys, Lys(Ac), Lys(Benzoic acid), Lys(glutaric acid), Lys(IVA), Lys(Peg4-isoGlu-Palm), Lys(pyroglutaric acid), Lys-(succinic acid), Asn, Orn, Gln, Arg, or Val; X14 is Dab(Ac), Dap(Ac), His, Lys(Ac), Asn, Gln, or Tyr; and X15 is Ala, betaAla, Gly, Asn, Gln, or Ser.

**[0015]** In certain embodiments of Xa: X1 is absent; X2 is absent; X3 is absent; X4 is Cys, Abu or Pen; X5 is Dap, Dap(Ac), Gln, Ser, Thr or Asn; X6 is Thr; X7 is Trp; X8 is Gln; X9 is Cys, Abu or Pen; X10 is a Phe analog, Tyr, or a Tyr analog; X11 is 2-Nal or Trp; X12 is Acpc, Acbc, Acvc, Achc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, hLeu, Leu, or THP; X13 is Cit, Asp, Glu, Lys, Lys(Ac), Asn, or Gln; X14 is Dab(Ac), Asn, or His; and X15 is Ala, betaAla, Gly, Asn, or Gln.

**[0016]** In certain embodiments of Xa: X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Sec, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-chloropropanoic acid, 4-chlorobutyric acid, 3-chloroisobutyric acid, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-

(3'-butenyl)glycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, 2-(5'-hexenyl)glycine, or Abu; X7 is Trp, Glu, Gly, Ile, Asn, Pro, Arg, Thr or OctGly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Leu, Val, Phe, or Ser, Sec, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, Ala, hCys, Abu, Met, MeCys, (D)Tyr or 2-(5'-hexenyl)glycine; X10 is Tyr, Phe(4-OMe), 1-Nal, 2-Nal, Aic,  $\alpha$ -MePhe, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, His, hPhe(3,4-dimethoxy), hTyr, N-Me-Tyr, Trp, Phe(4-CONH<sub>2</sub>), Phe(4-phenoxy), Thr, Tic, Tyr(3-tBu), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5-F<sub>2</sub>), Phe(4-CH<sub>2</sub>CO<sub>2</sub>H), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>), Phe(4-OCH<sub>3</sub>), Bip, Cha, 4-PyridylAlanine,  $\beta$ hTyr, OctGly, Phe(4-N<sub>3</sub>), Phe(4-Br), Phe[4-(2-aminoethoxy)] or Phe, a Phe analog, a Tyr analog, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe (3,4-F<sub>2</sub>), Phe(4-CO<sub>2</sub>H),  $\beta$ hPhe(4-F),  $\alpha$ -Me-Trp, 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tri-tert-Butyl), Phe(4-Oallyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino, Phe(4-OBzl), Octgly, Glu(Bzl), 4-Phenylbenzylalanine, Phe[4-(2-aminoethoxy)], 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, 1,2,3,4-tetrahydro-norharman, Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine, Bip, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acvc, Acbc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Aib,, D-Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Tyr, Aib,  $\alpha$ -MeLeu, $\alpha$ -MeOrn,  $\beta$ -Aib,  $\beta$ -Ala,  $\beta$ hAla,  $\beta$ hArg,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Glu, hArg, Ile, Lys, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Ser, Thr, Tle, t-butyl-Gly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Arg, Orn, Val,  $\beta$ hAla, Lys(Ac), (D)Asn, (D)Leu, (D)Phe, (D)Thr, Ala,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, hLeu, Asn, Ogl, Pro, Gln, Ser,  $\beta$ -spiro-pip, Thr, Tba, Tle or Aib, Cit, hArg, Lys, Asn, Orn, Gln or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X14 is Phe, Tyr, Glu, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr,

Tic $\beta$ hPhe, Arg, Lys(Ac), His; Dap(Ac), Dab(Ac), Asp or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X15 is Gly, Ser, Thr, Gln, Ala, (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Aea, Asp, Asn, Glu, Phe, Gly, Lys, Leu, Pro, Arg,  $\beta$ -Ala, Sarc, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, Gly, Ser, Pro, Asn, Thr or absent, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; and X17 is Leu, Lys, Arg, Glu, Ser, Gly, Gln or absent, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing.

[0017] In certain embodiments of peptide inhibitors of Xa, the bond is a disulfide bond, a thioether bond, a lactam bond, a triazole ring, a selenoether bond, a diselenide bond, or an olefin bond.

[0018] In particular embodiments of peptide inhibitors of Xa, X4 is Cys and X9 is Cys, and the bond is a disulfide bond. In particular embodiments, X4 is Pen and X9 is Pen, and the bond is a disulfide bond. In certain embodiments: X7 is Trp; X10 is Phe, Tyr, a Phe analog, or a Tyr analog; X11 is Trp, 1-Nal or 2-Nal; and X12 is Aib,  $\alpha$ -Me-Lys, a-Me-Leu, Achc, Acvc, Acpc, Acbc or THP. In certain embodiments: X7 is Trp; X10 is Phe, Tyr, a Phe analog, or a Tyr analog; X11 is Trp, 1-Nal or 2-Nal; and X12 is Aib,  $\alpha$ -Me-Lys or a-Me-Leu. In particular embodiments, the peptide inhibitor comprises any of the following the amino acid sequences: Pen-Q-T-W-Q-Pen-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -Me-Lys]-E-N-G; Pen-N-T-W-Q-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-N-N; Pen-Q-T-W-Q-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-N-N; or Pen-Q-T-W-Q-[Pen]-[Phe(4-CONH<sub>2</sub>)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-N-N, wherein the peptide inhibitor comprises a disulfide bond between the two Pen amino acids.

[0019] In particular embodiments of peptide inhibitors of Xa, X4 is an amino acid, aliphatic acid, alicyclic acid or modified 2- methyl aromatic acid having a carbon side chain capable of forming a thioether bind with X9; X9 is a sulfur-containing amino acid capable of forming a thioether bond with X4, and the bond between X4 and X9 is a thioether bond. In certain embodiments, X4 is Abu, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-chloro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid; and X9 is Abu, Cys, Pen, hCys, D-Pen, D-Cys, or D-hCys. In certain

embodiments, X4 is Abu; and X9 is Cys. In certain embodiments, X7 is Trp; X10 is Phe, Tyr, a Phe analog, or a Tyr analog; X11 is Trp, 1-Nal or 2-Nal; and X12 is  $\alpha$ -Me-Lys,  $\alpha$ -Me-Leu,  $\alpha$ -Me-Ser,  $\alpha$ -Me-Val, Achc, Acvc, Acpc, Acbc, or [4-amino-4-carboxy-tetrahydropyran]. In certain embodiments, X7 is Trp; X10 is Phe, Tyr, a Phe analog, or a Tyr analog; X11 is Trp, 1-Nal or 2-Nal; and X12 is  $\alpha$ -Me-Lys or [4-amino-4-carboxy-tetrahydropyran]. In particular embodiments, the peptide inhibitor comprises any of the following amino acid sequences: [Abu]-Q-T-W-Q-C-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-E-N-G; [Abu]-Q-T-W-Q-C-[Phe(4-(2-aminoethoxy))]-W-[ $\alpha$ -MeLys]-E-N-G; or [Abu]-Q-T-W-Q-C-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-E-N-N, wherein the peptide inhibitor comprises a thioether bond between the Abu and the C.

**[0020]** In certain embodiments of peptide inhibitors of Xa: X4 is Pen, Cys or hCys; X5 is any amino acid; X6 is any amino acid; X7 is Trp, Bip, Gln, His, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -Me-Trp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>), Phe(4-tBu),  $\beta\beta$ -diPheAla, Glu, Gly, Ile, Asn, Pro, Arg, Thr or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X8 is any amino acid; X9 is Pen, Cys or hCys; X10 is 1-Nal, 2-Nal, Aic, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, Phe, His, Trp, Thr, Tic, Tyr, 4-pyridylAla, Octgly, a Phe analog or a Tyr analog (optionally, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), or Phe(4-OBzl)), or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe (3,4-F<sub>2</sub>), Phe(4-CO<sub>2</sub>H),  $\beta$ hPhe(4-F),  $\alpha$ -Me-Trp, 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tr-tert-Butyl), Phe(4-Oallyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl), Octgly, Glu(Bzl), 4-Phenylbenzylalanine, Phe[4-(2-aminoethoxy)], 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, 1,2,3,4-tetrahydro-norharman, Phe(4-CONH<sub>2</sub>), Phe(3,4-OMe<sub>2</sub>) Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine or Bip, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu,  $\alpha$ -MeVal, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp,

(D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Lys, Arg, Orn, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Leu, Asn, Ogl, Pro, Gln, Asp, Arg, Ser, spiro-pip, Thr, Tba, Tlc, Val or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr, Lys(Ac), Orn or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser,  $\beta$ -Ala, Arg or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X16 is absent, Gly, Ala, Asp, Ser, Pro, Asn or Thr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X17 is absent, Glu, Ser, Gly or Gln, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X18 is absent or any amino acid; X19 is absent or any amino acid; and X20 is absent or any amino acid. In particular embodiments, the bond between X4 and X9 is a disulfide bond. In certain embodiments, X1, X2, and X3 are absent. In certain embodiments, X17, X19 and X20 are absent. In certain embodiments, one or both of X4 or X9 is Pen. In certain embodiments, both X4 and X9 are Pen. In particular embodiments, X18 is (D)-Lys. In certain embodiments, the peptide inhibitors comprise one or more, two or more, three or more, or four of the following: X5 is Arg, Asn, Gln, Dap, Orn; X6 is Thr or Ser; X7 is Trp, 2-Nal, 1-Nal, Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp or 1,2,3,4 -tetrahydro-norharman; and X8 is Gln, Val, Phe, Glu, Lys. In certain embodiments, the peptide inhibitors comprise one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following: X10 is Tyr, Phe(4-OBzl), Phe(4-OMe), Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-CO<sub>2</sub>H), Phe(4-(2aminoethoxy)) or Phe(4-guanadino); X11 is Trp, 2-Nal, 1-Nal, Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp or 1,2,3,4 -tetrahydro-norharman; X12 is Arg,  $\alpha$ -MeLys  $\alpha$ -MeLeu, Aib or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly, Sr or Ala; and X16 is absent or

AEA. In certain embodiments, X4 and X9 are Pen; X5 is Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr, Phe(4-OMe) or 2-Nal; X11 is Trp, 2-Nal or 1-Nal; X12 is Arg,  $\alpha$ -MeLys or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly; and X16 is absent. In certain embodiments, one or more of X1, X2 and X3 are absent; and one or more, two or more, three or more, or four of X17, X18, X19 and X20 are absent.

**[0021]** In certain embodiments of peptide inhibitors of Xa: X4 is Abu, Pen, or Cys; X7 is Trp, Bip, Gln, His, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>),  $\beta\beta$ -diPheAla, Phe(4-tBu), Glu, Gly, Ile, Asn, Pro, Arg, Thr or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X9 is Abu, Pen, or Cys; X10 is 1-Nal, 2-Nal, Aic, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, Phe, His, Trp, Thr, Tic, Tyr, 4-pyridylAla, Octgly a Phe analog or a Tyr analog, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, 4-phenylcyclohexyl, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(3,4-F<sub>2</sub>),  $\beta$ hPhe(4-F), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>), Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine,  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Bip, Nva(5-phenyl), Phe, His, hPhe, Tqa, Trp, Tyr, Phe(4-Me), Trp(2,5,7-tri-tertButyl), Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl), or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu, MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu,  $\beta$ Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Arg, Orn, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Leu, Asn, Ogl, Pro, Gln, Asp, Arg, Ser, spiro-pip, Thr, Tba, Tlc, Val or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr, or a

corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing, or X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, Asn, Ser, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X16 is absent, Gly, Ala, Asp, Ser, Pro, Asn or Thr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; and X17 is absent, Glu, Ser, Gly or Gln, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing. In particular embodiments, the peptide inhibitor is cyclized via an intramolecular bond between X4 and X9. In certain embodiments, one or more of X1, X2, and X3 are absent. In certain embodiments, one or more of X17, X19 and X20 are absent. In certain embodiments, one of X4 or X9 is Abu, and the other of X4 or X9 is not Abu. In certain embodiments, the peptide inhibitors comprise one or more, two or more, three or more, or four of the following: X5 is Arg, Gln, Dap or Orn; X6 is Thr or Ser; X7 is Trp, 2-Nal, 1-Nal, Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl), Phe(4-Me), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, or  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman; and X8 is Gln, Val, Phe, Glu or Lys. In certain embodiments, the peptide inhibitors comprise one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following: X10 is Tyr, Phe(4-OBzl), Phe(4-OMe), Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-CO<sub>2</sub>H), Phe(4-(2aminoethoxy)) or Phe(4-guanadino); X11 is Trp, 2-Nal, 1-Nal, Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp or 1,2,3,4 -tetrahydro-norharman; X12 is Arg, hLeu, (D)Asn, Aib,  $\alpha$ -MeLys,  $\alpha$ -MeLeu or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly, Ser or Ala, or X15 is Asn, Gly, Ser,  $\beta$ Ala or Ala; and X16 is absent or AEA.

[0022] In particular embodiments of any of the peptide inhibitors, X4 and X9 are Pen. In particular embodiments, X4 and X9 form a disulfide bond.

[0023] In particular embodiments, X4 is Abu and X9 is Cys. In particular embodiments, X4 and X9 form a thioether bond.

In particular embodiments, the peptide inhibitor comprises an amino acid sequence of any one of SEQ ID NOS: 365-370, 857-1029. In particular embodiments, the peptide inhibitor is cyclized via a bond between X4 and X9, and the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

**[0024]** In certain embodiments of peptide inhibitors of Xa, the peptide inhibitor comprises an amino acid sequence set forth in any of Formulas (V), (Va), (Vb), (Vc), (Vd), (Ve), (Vf), (Vg) and (Vh).

**[0025]** In certain embodiments of peptide inhibitors of Xa, the peptide inhibitor comprises any of the following amino acid sequences:

[Palm]-[isoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NNNH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-isoGlu-Palm)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Octanyl]-[IsoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Octanyl]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Palm]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Octanyl)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Palm)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Palm)-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Lauryl)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-IsoGlu-Palm)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-IsoGlu-Lauryl)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(IV A)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(Biotin)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(Octanyl)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-[Lys(IV A)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(IV A)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Biotin)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Biotin)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Octanyl)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(octanyl)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Palm)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Lys(Palm)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(PEG8)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(PEG8)]-N-NH<sub>2</sub>;

Ac-[Pen]-K(Peg11-Palm)TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Peg11-palm)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Cit]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Ac)]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(3,4-OCH<sub>3</sub>)<sub>2</sub>]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(2,4-CH<sub>3</sub>)<sub>2</sub>]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(3-CH<sub>3</sub>)]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(4-CH<sub>3</sub>)]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH<sub>2</sub>;

Ac-[(D)Tyr]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-N-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Lys(Ac)]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QQ-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Q-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Cit]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-NNH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-Q-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-[Lys(Ac)]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-[Cit]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QN-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-E-[Cit]-Q-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Cit]-N-[Cit]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Cit]-Q-[Cit]-NH<sub>2</sub>;

Ac-[Pen]-[Cit]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QNN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-ENQ-NH<sub>2</sub>;

Ac-[Pen]-GPWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-PGWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWN-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NSWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-N-[Aib]-WQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTW-[Aib]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]N-[Aib]-NH<sub>2</sub>;

Ac-[Pen]-QTW-[Lys(Ac)]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Ac)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]NNNH<sub>2</sub>;

Ac-[Pen]-QVWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[2-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[1-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-LN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-GN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-SN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Aib]-N-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-FN-NH<sub>2</sub>;

Ac-[Pen]-NTW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Tic]-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[nLeu]-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-G-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-R-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-W-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-S-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-L-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[AIB]-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[N-MeAla]-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[2-Nap]-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-F-[βAla]-NH<sub>2</sub>;

Ac-[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]NNNH<sub>2</sub>;

Biotin-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-E-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Asp]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-R-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

inoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-F-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Phe]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[2-Nal]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-T-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-L-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Gln]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Asn]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(PEG4-Alexa488)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

[Alexa488]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

[Alexa647]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

[Alexa-647]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Alexa647]-[PEG12]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>; and

[Alexa488]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>,

**[0026]** wherein the peptide inhibitor is cyclized via a disulfide bond between two Pen residues or by a thioether bond between Abu and a Cys residue, and wherein the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

**[0027]** In particular embodiments, any of the peptide inhibitors described herein comprise one or more half-life extension moiety and/or one or more linker moiety conjugated to the peptide inhibitor. In particular embodiments, the half-life extension moiety is conjugated to the peptide inhibitor via one or more linker moieties.

**[0028]** In certain embodiments, any of the peptide inhibitors described herein further comprise a conjugated chemical substituent. In particular embodiments, the conjugated chemical substituent

is a lipophilic substituent or a polymeric moiety, e.g., Ac, Palm, gamaGlu-Palm, isoGlu-Palm, PEG2-Ac, PEG4-isoGlu-Palm, (PEG)<sub>5</sub>-Palm, succinic acid, glutaric acid, pyroglutaric acid, benzoic acid, IVA, octanoic acid, 1,4 diaminobutane, isobutyl, or biotin. In certain embodiments, the conjugated chemical substituent is a polyethylene glycol with a molecular mass of 400 Da to 40,000 Da.

**[0029]** In another aspect, the present invention includes peptide inhibitors comprising the structure of Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein

$R^1$  is a bond, hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl, a C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing;

$R^2$  is a bond, OH or NH<sub>2</sub>; and

X is any of the peptide sequences described herein, e.g., Xa, I, Ia-It, II, IIa-IIId, III, IIIa-IIIe, IV, IVa-IVb, V, or Va-Vh.

**[0030]** In a related aspect, the present invention includes a peptide dimer inhibitor of an interleukin-23 receptor, wherein the peptide dimer inhibitor comprises two peptide monomer subunits connected via one or more linker moieties, wherein each peptide monomer subunit has a sequence or structure set forth herein. In certain embodiments, one or both peptide monomer subunit is cyclized via an intramolecular bond between X4 and X9. In certain embodiments, one or both intramolecular bond is a disulfide bond, a thioether bond, a lactam bond, a selenoether, diselenide, or an olefin bond. In certain embodiments, the linker is any of those shown in Table 2 or described herein. In certain embodiments, the linker moiety is a diethylene glycol linker, an iminodiacetic acid (IDA) linker, a  $\beta$ -Ala-iminodiaceticacid ( $\beta$ -Ala-IDA) linker, or a PEG linker. In particular embodiments, the N-terminus of each peptide monomer subunit is connected by the linker moiety. In particular embodiments, the C-terminus of each peptide monomer subunit is connected by the linker moiety. In certain embodiments, the linker connects an internal amino

acid residue of at least one of the peptide monomer subunits to the N-terminus, C-terminus, or an internal amino acid residue of the other peptide monomer subunit.

[0031] In a further related aspect, the present invention includes a polynucleotide comprising a sequence encoding a peptide inhibitor of the present invention or one or both peptide monomer subunit of a peptide dimer inhibitor of the present invention. The present invention also includes a vector comprising the polynucleotide.

[0032] In another aspect, the present invention includes a pharmaceutical composition comprising a peptide inhibitor or a peptide dimer inhibitor of the present invention, and a pharmaceutically acceptable carrier, excipient, or diluent. In particular embodiments, the pharmaceutical composition comprises an enteric coating. In certain embodiments, the enteric coating protects and releases the pharmaceutical composition within a subject's lower gastrointestinal system.

[0033] In another aspect, the present invention includes a method for treating or preventing a disease associated with IL-23 signalling, including but not limited to an Inflammatory Bowel Disease (IBD), ulcerative colitis, Crohn's disease, Celiac disease (*nontropical Sprue*), enteropathy associated with seronegative arthropathies, microscopic colitis, collagenous colitis, eosinophilic gastroenteritis, colitis associated with radio- or chemo-therapy, colitis associated with disorders of innate immunity as in leukocyte adhesion deficiency-1, chronic granulomatous disease, glycogen storage disease type 1b, Hermansky-Pudlak syndrome, Chediak-Higashi syndrome, and Wiskott-Aldrich Syndrome, pouchitis resulting after proctocolectomy and ileoanal anastomosis, gastrointestinal cancer, pancreatitis, insulin-dependent diabetes mellitus, mastitis, cholecystitis, cholangitis, pericholangitis, chronic bronchitis, chronic sinusitis, asthma, psoriasis, or graft versus host disease in a subject, comprising providing to the subject an effective amount of a peptide inhibitor or pharmaceutical composition of the present invention. In certain embodiments, the inflammatory bowel disease is ulcerative colitis or Crohn's disease. In particular embodiments, the peptide inhibitor or the peptide dimer inhibitor inhibits binding of an interleukin-23 (IL-23) to the interleukin-23 receptor (IL-23R). In certain embodiments, the pharmaceutical composition is provided to the subject by an oral, intravenous, peritoneal, intradermal, subcutaneous, intramuscular, intrathecal, inhalation, vaporization, nebulization,

sublingual, buccal, parenteral, rectal, intraocular, inhalation, vaginal, or topical route of administration. In particular embodiments, the pharmaceutical composition is provided orally for treating Inflammatory Bowel Disease (IBD), ulcerative colitis, Crohn's disease. In certain embodiments, the pharmaceutical composition is provided to the subject topically, parenterally, intravenously, subcutaneously, peritoneally, or intravenously for treating psoriasis.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0034] **FIG. 1** **Figure 1** provides an example of a rat IL-23 dose-response curve as measured by levels of IL-17A in the rat splenocyte assay.

[0035] **Figure 2** is a graph showing IL-12-dependent production of IFN $\gamma$  from human PBMCs treated with the indicated amounts of Compound A or Compound B.

[0036] **Figure 3** shows results for DAI values from Day 7. Statistical analysis for significance was determined using Student's t-test (GraphPad Prism). Differences were noted as significant \*p<0.05, \*\*p<0.01, \*\*\*p<0.001, \*\*\*\*p<0.0001.

[0037] **Figure 4** shows an alignment of the amino acid sequences of human IL23R, mouse IL-23R, rat IL23R, chimp IL-23R, dog IL-23R and cow IL-23R, with highly conserved amino acid residues shaded. The region of mouse IL-23R lacking in the other IL-23R species shown is shown, and a region of IL23R that may be bound by certain peptide inhibitors of the present invention is indicated by a dashed line.

[0038] **Figure 5** is a table outlining the study design for TNBS induced colitis in rats.

[0039] **Figures 6A-6D** are graphs showing colon weight to length (Figure 6A), colon wall thickness (Table 6B, colon macroscopic score (Table 6C) or myeloperoxidase (MPO) abundance in proximal colon extracts quantified by ELISA, following sham treatment, vehicle treatment, or treatment with the indicated amounts of anti-IL23p19 antibody or Compound C. Values are shown as mean  $\pm$ SD. Statistical significance assessed by one-way ANOVA: \* $\leq$ 0.05; \*\* $\leq$ 0.01; \*\*\*p $\leq$ 0.001; \*\*\*\*p $\leq$ 0.0001; ns, not significant.

[0040] **Figure 7** provides micrographs of colon lesions found in animals following sham treatment (upper left panel), vehicle treatment (upper right panel) showing transmural inflammation, presence of necrotic tissue, and mucosa devoid of crypts, anti-IL23p19 antibody (lower left panel), or 160 mg/kg/d Compound C (lower right panel) showing restriction of lesions to the mucosa.

[0041] **Figures 8A-8E** are graphs showing inflammation (Figure 8A), mucosal necrosis (Figure 8B), grand loss (Figure 8C), colon wall thickness (Figure 8D) and histological score (Figure 8E) following vehicle treatment, treatment with anti-IL23p19 antibody, or treatment with the indicated amount of Compound C

[0042] **Figure 9** shows the concentration of Compound C in the plasma and proximal colon determined one hour post last PO dose (left panel), and fold above IC75 of its activity as determined by the rat splenocyte assay (middle panel) and the rat IL23R ELISA assay (right panel).

[0043] **Figure 10** provides a schematic diagram depicting the structure of certain peptide inhibitors and illustrating representative types of bonds between X4 and X9.

[0044] **FIGS. 11A-11E** show pharmacokinetic data for the IL-23R peptide inhibitor Peptide 993 (SEQ ID NO: 993). FIG. 11A shows the concentration of Peptide 993 in serum (nM) measured at different time points up to 24 hours after oral administration of Peptide 993. FIGS. 11B-11D show the concentration of Peptide 993 (in nM) in samples taken from the Peyer's Patch (FIG. 11B), small intestine (FIG. 11C), and the colon (FIG. 11D). The dashed line indicates 350 mM. FIG. 11E shows the amount of Peptide 993 detected in feces 24 hours after oral administration (% dose).

[0045] **FIGS. 12A-12D** summarize experiments comparing systemic treatments with prednisolone or anti-IL-23p19 neutralizing antibody with treatment with Peptide 993 by oral administration in the TNBS model of acute colitis. FIG. 12A shows the change in body weight (percentage) from day 0 to day 7 from sham, vehicle, and Peptide 993 treated rats. FIG. 12B shows the ratio of colon weight to colon length in mg/cm of colons harvested from rats at day 7. FIG. 12C shows the colon macroscopic score of colons harvested from rats at day 7. FIG. 12D.

shows the sum of histopathology scores for colons taken from sham, vehicle, and Peptide 993 treated rats. For all experiments, statistical comparisons between groups were performed with a 1-Way ANOVA followed by a post hoc test: \* p< 0.05; \*\* p< 0.01; \*\*\* p< 0.001; \*\*\*\* p< 0.0001; ns, not significant.

**[0046] FIGS. 13A-13C** summarizes experiments comparing systemic treatments with prodnisolone or anti-IL-23p19 neutralizing antibody with treatment with Peptide 1185 by oral administration in the TNBS model of acute colitis. FIG. 13A shows the change in body weight (percentage) from day 0 to day 7 from sham, vehicle, and Peptide 1185 treated rats. FIG. 13B shows the ratio of colon weight to colon length in mg/cm of colons harvested from rats at day 7. FIG. 13C shows the colon macroscopic score of colons harvested from rats at day 7. For all experiments, statistical comparisons between groups were performed with a 1-Way ANOVA followed by a post hoc test: \* p< 0.05; \*\* p< 0.01; \*\*\* p< 0.001; \*\*\*\* p< 0.0001; ns, not significant.

**[0047] FIGS. 14A-14D** summarizes experiments comparing systemic treatments with prodnisolone or anti-IL-23p19 neutralizing antibody with treatment with Peptide 980 by oral administration in the TNBS model of acute colitis. FIG. 14A shows the change in body weight (percentage) from day 0 to day 7 from sham, vehicle, and Peptide 980 treated rats. FIG. 14B shows the ratio of colon weight to colon length in mg/cm of colons harvested from rats at day 7. FIG. 13C shows the colon macroscopic score of colons harvested from rats at day 7. FIG. 14D. shows the sum of histopathology scores for colons taken from sham, vehicle, and Peptide 980 treated rats. For all experiments, statistical comparisons between groups were performed with a 1-Way ANOVA followed by a post hoc test: \* p< 0.05; \*\* p< 0.01; \*\*\* p< 0.001; \*\*\*\* p< 0.0001; ns, not significant.

**[0048] FIGS. 15A-15E** show levels of disease and IL-23 directed biomarkers measured in colons from rats in the sham (not TNBS-exposed) experimental group, or TNBS-exposed experimental groups that received treatment with vehicle or Peptide 993. Data is shown for MPO (FIG. 15A), IL-6 (FIG. 15B), IL-1 beta (FIG. 15C), IL-22 (FIG. 15D), and IL-17A (FIG. 15E). For all experiments, statistical comparisons between groups were performed with a

1-Way ANOVA followed by a post hoc test: \* p< 0.05; \*\* p< 0.01; \*\*\* p< 0.001; \*\*\*\* p< 0.0001; ns, not significant.

**[0049]** **FIGS. 16A-16B** show levels of disease and IL-23 directed biomarkers measured in colons from rats in the sham (not TNBS-exposed) experimental group, or TNBS-exposed experimental groups that received treatment with vehicle or Peptide 980. Data is shown for MPO (FIG. 16A) and IL-22 (FIG. 16B). For all experiments, statistical comparisons between groups were performed with a 1-Way ANOVA followed by a post hoc test: \* p< 0.05; \*\* p< 0.01; \*\*\* p< 0.001; \*\*\*\* p< 0.0001; ns, not significant

**[0050]** **FIGS. 17A-17D** show Schild analysis of inhibitor peptides. FIG. 17A shows a graph depicting the % Emax response as a function of increasing concentrations of IL-23 in the presence of Peptide 993 in concentrations of 0 nM (closed circles), 0.3 nM (closed squares), 1 nM (triangles), 3 nM (inverted triangles), 10 nM (diamonds), 30 nM (open circles), or 100 nM (open squares). Properties of the curves are listed below the graph. FIG. 17B depicts results from the same set of experiments, and shows a graph displaying Log(dose ratio<sup>-1</sup>) as a function of Peptide 993 concentration (M) on a logarithmic scale. Properties of the resulting linear function are displayed below the graph. FIG. 17C shows a graph depicting the % Emax response as a function of increasing concentrations of IL-23 in the presence of the peptide of SEQ ID NO: 1169 in concentrations of 0 nM (closed circles), 0.3 nM (closed squares), 1 nM (triangles), 3 nM (inverted triangles), 10 nM (diamonds), 30 nM (open circles), or 100 nM (open squares). Properties of the curves are listed below the graph. FIG. 17D shows a graph depicting the % Emax response as a function of increasing concentrations of IL-23 in the presence of the peptide of SEQ ID NO: 1211 in concentrations of 0 nM (closed circles), 0.3 nM (closed squares), 1 nM (triangles), 3 nM (inverted triangles), 10 nM (diamonds), 30 nM (open circles), or 100 nM (open squares). Properties of the curves are listed below the graph.

**[0051]** **FIGS. 18A-18B** show pharmacokinetic data for the IL-23R peptide inhibitor Peptide 1185. FIG. 18A shows the concentration of Peptide 1185 in serum and in samples taken from small intestine and the colon. Fig 18B shows the amount of Peptide 1185 detected in urine and feces 24 hours after oral administration (% dose).

[0052] FIGS. 19A and 19B show pharmacokinetic data for the IL-23R peptide inhibitor Peptide 980. FIG. 19A shows the concentration of Peptide 980 in serum and in samples taken from small intestine and the colon. Fig 19B shows the amount of Peptide 980 detected in urine and feces 24 hours after oral administration (% dose).

### **DETAILED DESCRIPTION OF THE INVENTION**

[0053] Unless otherwise defined herein, scientific and technical terms used in this application shall have the meanings that are commonly understood by those of ordinary skill in the art. Generally, nomenclature used in connection with, and techniques of, chemistry, molecular biology, cell and cancer biology, immunology, microbiology, pharmacology, and protein and nucleic acid chemistry, described herein, are those well-known and commonly used in the art.

[0054] As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

[0055] Throughout this specification, the word “comprise” or variations such as “comprises” or “comprising” will be understood to imply the inclusion of a stated integer (or components) or group of integers (or components), but not the exclusion of any other integer (or components) or group of integers (or components).

[0056] The singular forms “a,” “an,” and “the” include the plurals unless the context clearly dictates otherwise.

[0057] The term “including” is used to mean “including but not limited to.” “Including” and “including but not limited to” are used interchangeably.

[0058] The terms “patient,” “subject,” and “individual” may be used interchangeably and refer to either a human or a non-human animal. These terms include mammals such as humans, primates, livestock animals (e.g., bovines, porcines), companion animals (e.g., canines, felines) and rodents (e.g., mice and rats).

[0059] The term “peptide,” as used herein, refers broadly to a sequence of two or more amino acids joined together by peptide bonds. It should be understood that this term does not connote a

specific length of a polymer of amino acids, nor is it intended to imply or distinguish whether the polypeptide is produced using recombinant techniques, chemical or enzymatic synthesis, or is naturally occurring.

**[0060]** The recitations “sequence identity”, “percent identity”, “percent homology”, or, for example, comprising a “sequence 50% identical to,” as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity.

**[0061]** Calculations of sequence similarity or sequence identity between sequences (the terms are used interchangeably herein) can be performed as follows. To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences can be aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In certain embodiments, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position.

**[0062]** The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0063] The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In some embodiments, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch, (1970, *J. Mol. Biol.* 48: 444-453) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package, using an NW<sub>Sgapdna.CMP</sub> matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. Another exemplary set of parameters includes a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5. The percent identity between two amino acid or nucleotide sequences can also be determined using the algorithm of E. Meyers and W. Miller (1989, *Cabios*, 4: 11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0064] The peptide sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al., (1990, *J. Mol. Biol.*, 215: 403-10). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al. (*Nucleic Acids Res.* 25:3389-3402, 1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

[0065] The term "conservative substitution" as used herein denotes that one or more amino acids are replaced by another, biologically similar residue. Examples include substitution of amino acid residues with similar characteristics, e.g., small amino acids, acidic amino acids, polar amino acids, basic amino acids, hydrophobic amino acids and aromatic amino acids. See, for

example, the table below. In some embodiments of the invention, one or more Met residues are substituted with norleucine (Nle) which is a bioisostere for Met, but which, as opposed to Met, is not readily oxidized. Another example of a conservative substitution with a residue normally not found in endogenous, mammalian peptides and proteins is the conservative substitution of Arg or Lys with, for example, ornithine, canavanine, aminoethylcysteine or another basic amino acid. In some embodiments, one or more cysteines of a peptide analogue of the invention may be substituted with another residue, such as a serine. For further information concerning phenotypically silent substitutions in peptides and proteins, see, for example, Bowie et.al. *Science* 247, 1306-1310, 1990. In the scheme below, conservative substitutions of amino acids are grouped by physicochemical properties. I: neutral, hydrophilic, II: acids and amides, III: basic, IV: hydrophobic, V: aromatic, bulky amino acids.

<b>I</b>	<b>II</b>	<b>III</b>	<b>IV</b>	<b>V</b>
A	N	H	M	F
S	D	R	L	Y
T	E	K	I	W
P	Q		V	
G			C	

**[0066]** In the scheme below, conservative substitutions of amino acids are grouped by physicochemical properties. VI: neutral or hydrophobic, VII: acidic, VIII: basic, IX: polar, X: aromatic.

<b>VI</b>	<b>VII</b>	<b>VIII</b>	<b>IX</b>	<b>X</b>
A	E	H	M	F
L	D	R	S	Y
I		K	T	W
P			C	
G			N	
V			Q	

[0067] The term “amino acid” or “any amino acid” as used here refers to any and all amino acids, including naturally occurring amino acids (e.g.,  $\alpha$ -amino acids), unnatural amino acids, modified amino acids, and non-natural amino acids. It includes both D- and L-amino acids. Natural amino acids include those found in nature, such as, e.g., the 23 amino acids that combine into peptide chains to form the building-blocks of a vast array of proteins. These are primarily L stereoisomers, although a few D-amino acids occur in bacterial envelopes and some antibiotics. The 20 “standard,” natural amino acids are listed in the above tables. The “non-standard,” natural amino acids are pyrrolysine (found in methanogenic organisms and other eukaryotes), selenocysteine (present in many noneukaryotes as well as most eukaryotes), and N-formylmethionine (encoded by the start codon AUG in bacteria, mitochondria and chloroplasts). “Unnatural” or “non-natural” amino acids are non-proteinogenic amino acids (i.e., those not naturally encoded or found in the genetic code) that either occur naturally or are chemically synthesized. Over 140 unnatural amino acids are known and thousands of more combinations are possible. Examples of “unnatural” amino acids include  $\beta$ -amino acids ( $\beta^3$  and  $\beta^2$ ), homo-amino acids, proline and pyruvic acid derivatives, 3-substituted alanine derivatives, glycine derivatives, ring-substituted phenylalanine and tyrosine derivatives, linear core amino acids, diamino acids, D-amino acids, alpha-methyl amino acids and N-methyl amino acids. Unnatural or non-natural amino acids also include modified amino acids. “Modified” amino acids include amino acids (e.g., natural amino acids) that have been chemically modified to include a group, groups, or chemical moiety not naturally present on the amino acid. According to certain embodiments, a peptide inhibitor comprises an intramolecular bond between two amino acid residues present in the peptide inhibitor. It is understood that the amino acid residues that form the bond will be altered somewhat when bonded to each other as compared to when not bonded to each other. Reference to a particular amino acid is meant to encompass that amino acid in both its unbonded and bonded state. For example, the amino acid residue homoSerine (hSer) or homoSerine(Cl) in its unbonded form may take the form of 2-aminobutyric acid (Abu) when participating in an intramolecular bond according to the present invention. The present invention includes both peptide inhibitors containing cross-links between X4 and X9, as well as the peptide inhibitors that do not contain cross-links between X4 and X9, e.g., before cross-link formation. As such, the names hSer and Abu are intended to indicate the same amino acids and are used interchangeably.

[0068] For the most part, the names of naturally occurring and non-naturally occurring aminoacyl residues used herein follow the naming conventions suggested by the IUPAC Commission on the Nomenclature of Organic Chemistry and the IUPAC-IUB Commission on Biochemical Nomenclature as set out in “Nomenclature of  $\alpha$ -Amino Acids (Recommendations, 1974)” Biochemistry, 14(2), (1975). To the extent that the names and abbreviations of amino acids and aminoacyl residues employed in this specification and appended claims differ from those suggestions, they will be made clear to the reader. Some abbreviations useful in describing the invention are defined below in the following Table 1A.



[0069] Table 1A. Abbreviations of Non-Natural Amino Acids and Chemical Moieties (for amino acid derivatives, all L unless stated)






Abbreviation	Definition
Ac-	Acetyl
Hy	Hydrogen (Free N-terminal)
Dap	L-Diaminopropionic acid
Dab	L-Diaminobutyric acid
Orn	L-Ornathine
Pen	L-Penicillamine
Sarc	Sarcosine
Cit	L-Citrulline
Cav	L-Cavanine
Phe-(4-Guanidino)	4-Guanidine-L-Phenylalanine
N-MeArg	N-Methyl-L-Arginine
N-MeTrp	N-Methyl-L-Tryptophan
N-MeGln	N-Methyl-L-Glutamine

N-MeAla	N-Methyl-L-Alanine
N-MeLys	N-Methyl-Lysine
N-MeAsn	N-Methyl-L-Asparagine
6-ChloroTrp	6-Chloro-L-Tryptophan
5-HydroxyTrp	5-Hydroxy-L-Tryptophan
1,2,3,4-tetrahydro-norharman	L-1,2,3,4-tetrahydro-norharman
2-Nal (also referred to as 2-Nap)	L-2-Naphylalanine
1-Nal (also referred to as 1-Nap)	L-1-Naphylalanine
Phe(4-OMe)	4-Methoxy-L-phenylalanine
Abu	2-Aminobutyric acid
Bip	L-4,4'-Biphenylalanine
βAla	beta-Alanine
βhTyr	beta homo-L-Tyrosine
βhTrp	beta homo-L-Trptophan
βhAla	beta homo-L-Alanine
βhLeu,	beta homo-L-Leucine
βhVal	beta homo-L-Valine
Aib	2-aminoisobutyric acid
Azt	L-azetidine-2-carboxylic acid
Tic	(3S)-1,2,3,4-Tetrahydroisoquinoline-7-hydroxy-3-carboxylic Acid
Phe(4-OMe)	4-methoxy-L-phenylalanine

N-Me-Lys	N-Methyl-L-Lysine
N-Me-Lys(Ac)	N- $\epsilon$ -Acetyl-D-lysine
CONH <sub>2</sub>	Carboxamide
COOH	Acid
3-Pal	L-3-Pyridylalanine
Phe(4-F)	4-Fluoro-L-Phenylalanine
DMT	2,6-DimethylTyrosine
Phe(4-OMe)	4-Methoxyphenylalanine
hLeu	L-homoLeucine
hArg	L-homoArginine
$\alpha$ -MeLys	alpha-methyl-L-Lysine
$\alpha$ -MeOrn	alpha-methyl-L-Ornathine
$\alpha$ -MeLeu	alpha-methyl-L-Leucine
$\alpha$ -MeTrp	alpha-methyl-L-Tryptophan
$\alpha$ -MePhe	alpha-methyl-L-Phenylalanine
$\alpha$ -MeTyr	alpha-methyl-L-Tyrosine
$\alpha$ -DiethylGly	$\alpha$ -DiethylGlycine
Lys(Ac)	N- $\epsilon$ -acetyl-L-Lysine
DTT	Dithiothreitol
Nle	L-Norleucine
$\beta$ hTrp	L- $\beta$ -homoTryptophan
$\beta$ hPhe	L- $\beta$ -homophenylalanine
$\beta$ hPro	L- $\beta$ -homoproline

Phe(4-CF <sub>3</sub> )	4-Trifluoromethyl-L-Phenylalanine
β-Glu	L-β-Glutamic acid
βhGlu	L-β-homoglutamic acid
2-2-Indane	2-Aminoindane-2-carboxylic acid
1-1-Indane	1-Aminoindane-1-carboxylic acid
hCha	L-homocyclohexylalanine
Cyclobutyl	L-cyclobutylalanine
βhPhe	L-β-homo-phenylalanine
Gla	Gama-Carboxy-L-Glutamic acid
Cpa	Cyclopentyl-L-alanine
Cha	Cyclohexyl-L-alanine
Octgly	L-Octylglycine
<i>t</i> -butyl-Ala	3-( <i>tert</i> -butyl)-L-Ala-OH
<i>t</i> -butyl-Gly	<i>tert</i> -butyl-glycine
AEP	3-(2-aminoethoxy)propanoic acid
AEA	(2-aminoethoxy)acetic acid
Phe(4-Phenoxy)]	4-Phenoxy-L-phenylalanine
Phe(4-OBzl)	<i>O</i> -Benzyl-L-tyrosine
Phe(4-CONH <sub>2</sub> )	4-Carbamoyl-L-phenylalanine
Phe(4-CO <sub>2</sub> H)	4-Carboxy-L-phenylalanine
Phe(3,4-Cl <sub>2</sub> )	3,4 dichloro-L-phenylalanine
Tyr(3- <i>t</i> -Bu)	3- <i>t</i> -butyl-L-tyrosine

Phe(t-Bu)	<i>t</i> -butyl-L-phenylalanine
Phe[4-(2-aminoethoxy)]	 4-(2-aminoethoxy)-L-phenylalanine
Phe(4-CN)	4-cyano-L-phenylalanine
Phe(4-Br)	4-bromo-L-phenylalanine
Phe(4-NH <sub>2</sub> )	4-amino-L-phenylalanine
Phe(4-Me)	4-methyl-L-phenylalanine
4-Pyridylalanine	4-L-Pyridylalanine
4-amino-4-carboxy-piperidine	 4-amino-4-carboxy-piperidine
hPhe(3,4-dimethoxy)	3,4-dimethoxy-L-homophenylalanine
Phe(2,4-Me <sub>2</sub> )	2,4-dimethyl-L-phenylalanine
Phe(3,5-F <sub>2</sub> )	3,5-difluoro-L-phenylalanine
Phe(penta-F)	pentafluoro-L-phenylalanine

2,5,7-tert butyl Trp	2,5,7-Tris-tert-butyl-L-tryptophan
Tic	 <p>L-1,2,3,4-tetrahydro-isoquinoline-3-carboxylic acid</p>
Phe(4-OAllyl)	O- <i>Allyl</i> -L-Tyrosine
Phe(4-N <sub>3</sub> )	4-azidophenylalanine
Achc	 <p>1-aminocyclohexanecarboxylic acid</p>
Acvc	 <p>1-aminocyclopentanecarboxylic acid</p>
Acbc	 <p>1-aminocyclobutanecarboxylic acid</p>
Acpc	

	1-aminocyclopropylcarboxylic acid
4-amino-4-carboxy- tetrahydropyran (also referred as THP)	<p>4-amino-4-carboxy-tetrahydropyran</p>

[0070] Throughout the present specification, unless naturally occurring amino acids are referred to by their full name (e.g. alanine, arginine, etc.), they are designated by their conventional three-letter or single-letter abbreviations (e.g. Ala or A for alanine, Arg or R for arginine, etc.). Unless otherwise indicated, three-letter and single-letter abbreviations of amino acids refer to the L-isomeric form of the amino acid in question. The term “L-amino acid,” as used herein, refers to the “L” isomeric form of a peptide, and conversely the term “D-amino acid” refers to the “D” isomeric form of a peptide (e.g., Dasp, (D)Asp or D-Asp; Dphe, (D)Phe or D-Phe). Amino acid residues in the D isomeric form can be substituted for any L-amino acid residue, as long as the desired function is retained by the peptide. D-amino acids may be indicated as customary in lower case when referred to using single-letter abbreviations.

[0071] In the case of less common or non-naturally occurring amino acids, unless they are referred to by their full name (e.g. sarcosine, ornithine, etc.), frequently employed three- or four-character codes are employed for residues thereof, including, Sar or Sarc (sarcosine, i.e. N-methylglycine), Aib ( $\alpha$ -aminoisobutyric acid), Dab (2,4-diaminobutanoic acid), Dapa (2,3-diaminopropanoic acid),  $\gamma$ -Glu ( $\gamma$ -glutamic acid), Gaba ( $\gamma$ -aminobutanoic acid),  $\beta$ -Pro (pyrrolidine-3-carboxylic acid), and 8Ado (8-amino-3,6-dioxaoctanoic acid), Abu (2-amino butyric acid),  $\beta$ hPro ( $\beta$ -homoproline),  $\beta$ hPhe ( $\beta$ -homophenylalanine) and Bip ( $\beta$ , $\beta$  diphenylalanine), and Ida (Iminodiacetic acid).

[0072] As is clear to the skilled artisan, the peptide sequences disclosed herein are shown proceeding from left to right, with the left end of the sequence being the N-terminus of the

peptide and the right end of the sequence being the C-terminus of the peptide. Among sequences disclosed herein are sequences incorporating a “Hy-” moiety at the amino terminus (N-terminus) of the sequence, and either an “-OH” moiety or an “-NH<sub>2</sub>” moiety at the carboxy terminus (C-terminus) of the sequence. In such cases, and unless otherwise indicated, a “Hy-” moiety at the N-terminus of the sequence in question indicates a hydrogen atom, corresponding to the presence of a free primary or secondary amino group at the N-terminus, while an “-OH” or an “-NH<sub>2</sub>” moiety at the C-terminus of the sequence indicates a hydroxy group or an amino group, corresponding to the presence of an amido (CONH<sub>2</sub>) group at the C-terminus, respectively. In each sequence of the invention, a C-terminal “-OH” moiety may be substituted for a C-terminal “-NH<sub>2</sub>” moiety, and vice-versa.

[0073] The term “DRP,” as used herein, refers to disulfide rich peptides.

[0074] The term “dimer,” as used herein, refers broadly to a peptide comprising two or more monomer subunits. Certain dimers comprise two DRPs. Dimers of the present invention include homodimers and heterodimers. A monomer subunit of a dimer may be linked at its C- or N-terminus, or it may be linked via internal amino acid residues. Each monomer subunit of a dimer may be linked through the same site, or each may be linked through a different site (e.g., C-terminus, N-terminus, or internal site).

[0075] The term “NH<sub>2</sub>,” as used herein, can refer to a free amino group present at the amino terminus of a polypeptide. The term “OH,” as used herein, can refer to a free carboxy group present at the carboxy terminus of a peptide. Further, the term “Ac,” as used herein, refers to Acetyl protection through acylation of the C- or N-terminus of a polypeptide. In certain peptides shown herein, the NH<sub>2</sub> locates at the C-terminus of the peptide indicates an amino group.

[0076] The term “carboxy,” as used herein, refers to -CO<sub>2</sub>H.

[0077] The term “isostere replacement,” as used herein, refers to any amino acid or other analog moiety having chemical and/or structural properties similar to a specified amino acid.

[0078] The term “cyclized,” as used herein, refers to a reaction in which one part of a polypeptide molecule becomes linked to another part of the polypeptide molecule to form a closed ring, such as by forming a disulfide bridge or other similar bond.

[0079] The term “subunit,” as used herein, refers to one of a pair of polypeptide monomers that are joined to form a dimer peptide composition.

[0080] The term “linker moiety,” as used herein, refers broadly to a chemical structure that is capable of linking or joining together two peptide monomer subunits to form a dimer.

[0081] The term “pharmaceutically acceptable salt,” as used herein, represents salts or zwitterionic forms of the peptides or compounds of the present invention which are water or oil-soluble or dispersible, which are suitable for treatment of diseases without undue toxicity, irritation, and allergic response; which are commensurate with a reasonable benefit/risk ratio, and which are effective for their intended use. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting an amino group with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, formate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isethionate), lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthalenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate, and undecanoate. Also, amino groups in the compounds of the present invention can be quaternized with methyl, ethyl, propyl, and butyl chlorides, bromides, and iodides; dimethyl, diethyl, dibutyl, and diamyl sulfates; decyl, lauryl, myristyl, and steryl chlorides, bromides, and iodides; and benzyl and phenethyl bromides. Examples of acids which can be employed to form therapeutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric. A pharmaceutically acceptable salt may suitably be a salt chosen, e.g., among acid addition salts and basic salts. Examples of acid addition salts include chloride salts, citrate salts and acetate salts. Examples of basic salts include salts where the cation is selected among alkali metal cations, such as sodium or potassium ions, alkaline earth metal cations, such as calcium or magnesium ions, as well as substituted ammonium ions, such as ions of the type  $N(R1)(R2)(R3)(R4)^+$ , where R1, R2, R3 and R4 independently will typically designate hydrogen, optionally substituted C1-6-alkyl or optionally substituted C2-6-alkenyl.

Examples of relevant C1-6-alkyl groups include methyl, ethyl, 1-propyl and 2-propyl groups. Examples of C2-6-alkenyl groups of possible relevance include ethenyl, 1-propenyl and 2-propenyl. Other examples of pharmaceutically acceptable salts are described in “Remington’s Pharmaceutical Sciences”, 17th edition, Alfonso R. Gennaro (Ed.), Mark Publishing Company, Easton, PA, USA, 1985 (and more recent editions thereof), in the “Encyclopaedia of Pharmaceutical Technology”, 3rd edition, James Swarbrick (Ed.), Informa Healthcare USA (Inc.), NY, USA, 2007, and in J. Pharm. Sci. 66: 2 (1977). Also, for a review on suitable salts, see *Handbook of Pharmaceutical Salts: Properties, Selection, and Use* by Stahl and Wermuth (Wiley-VCH, 2002). Other suitable base salts are formed from bases which form non-toxic salts. Representative examples include the aluminum, arginine, benzathine, calcium, choline, diethylamine, diolamine, glycine, lysine, magnesium, meglumine, olamine, potassium, sodium, tromethamine, and zinc salts. Hemisalts of acids and bases may also be formed, e.g., hemisulphate and hemicalcium salts.

**[0082]** The term “N(alpha)Methylation”, as used herein, describes the methylation of the alpha amine of an amino acid, also generally termed as an N-methylation.

**[0083]** The term “sym methylation” or “Arg-Me-sym”, as used herein, describes the symmetrical methylation of the two nitrogens of the guanidine group of arginine. Further, the term “asym methylation” or “Arg-Me-asym” describes the methylation of a single nitrogen of the guanidine group of arginine.

**[0084]** The term “acylating organic compounds”, as used herein refers to various compounds with carboxylic acid functionality that are used to acylate the N-terminus of an amino acid or a monomer or dimer, e.g., a monomer subunit prior to forming a C-terminal dimer. Non-limiting examples of acylating organic compounds include cyclopropylacetic acid, 4-Fluorobenzoic acid, 4-fluorophenylacetic acid, 3-Phenylpropionic acid, Succinic acid, Glutaric acid, Cyclopentane carboxylic acid, 3,3,3-trifluoropropeonic acid, 3-Fluoromethylbutyric acid, Tetrahedro-2H-Pyran-4-carboxylic acid.

**[0085]** The term “alkyl” includes a straight chain or branched, noncyclic or cyclic, saturated aliphatic hydrocarbon containing from 1 to 24 carbon atoms. Representative saturated straight chain alkyls include, but are not limited to, methyl, ethyl, *n*-propyl, *n*-butyl, *n*-pentyl, *n*-hexyl,

and the like, while saturated branched alkyls include, without limitation, isopropyl, *sec*-butyl, isobutyl, *tert*-butyl, isopentyl, and the like. Representative saturated cyclic alkyls include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like, while unsaturated cyclic alkyls include, without limitation, cyclopentenyl, cyclohexenyl, and the like.

**[0086]** The term “mammal” refers to any mammalian species such as a human, mouse, rat, dog, cat, hamster, guinea pig, rabbit, livestock, and the like.

**[0087]** As used herein, a “therapeutically effective amount” of the peptide inhibitor of the invention is meant to describe a sufficient amount of the peptide inhibitor to treat an IL-23/IL-23R-related disease, including but not limited to any of the diseases and disorders described herein (for example, to reduce inflammation associated with IBD). In particular embodiments, the therapeutically effective amount will achieve a desired benefit/risk ratio applicable to any medical treatment.

**[0088]** An “analog” of an amino acid, e.g., a “Phe analog” or a “Tyr analog” means an analog of the referenced amino acid. A variety of amino acid analogs are known and available in the art, including Phe and Tyr analogs. In certain embodiments, an amino acid analog, e.g., a Phe analog or a Tyr analog comprises one, two, three, four or five substitutions as compared to Phe or Tyr, respectively. In certain embodiments, the substitutions are present in the side chains of the amino acids. In certain embodiments, a Phe analog has the structure Phe(R<sup>2</sup>), wherein R<sup>2</sup> is a Hy, OH, CH<sub>3</sub>, CO<sub>2</sub>H, CONH<sub>2</sub>, CONH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, *t*-Bu, OCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, phenoxy, OCH<sub>3</sub>, OAllyl, Br, Cl, F, NH<sub>2</sub>, N3, or guanadino. In certain embodiments, R<sup>2</sup> is CONH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>, OCH<sub>3</sub>, CONH<sub>2</sub>, OCH<sub>3</sub> or CO<sub>2</sub>H. Examples of Phe analogs include, but are not limited to: hPhe, Phe(4-OMe),  $\alpha$ -Me-Phe, hPhe(3,4-dimethoxy), Phe(4-CONH<sub>2</sub>), Phe(4-phenoxy), Phe(4-guanadino), Phe(4-*t*Bu), Phe(4-CN), Phe(4-Br), Phe(4-OBzl), Phe(4-NH<sub>2</sub>), BhPhe(4-F), Phe(4-F), Phe(3,5 DiF), Phe(CH<sub>2</sub>CO<sub>2</sub>H), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe (3,4-F<sub>2</sub>), Phe(4-CF<sub>3</sub>),  $\beta$  $\beta$ -diPheAla, Phe(4-N<sub>3</sub>), Phe[4-(2-aminoethoxy)], 4-Phenylbenzylalanine, Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>), Phe(2,3-Cl<sub>2</sub>), and Phe(2,3-F<sub>2</sub>). Examples of Tyr analogs include, but are not limited to: hTyr, N-Me-Tyr, Tyr(3-*t*Bu), Tyr(4-N<sub>3</sub>) and  $\beta$ hTyr.

### Peptide Inhibitors of IL-23R

[0089] Genome-wide association studies (GWAS) have demonstrated significant association of the IL-23 receptor (IL-23R) gene with inflammatory bowel disease (IBD), suggesting that perturbation of IL-23 signaling could be relevant to the pathogenesis of the disease. The present invention provides compositions and methods to modulate the IL-23 pathway through selective antagonism of IL-23R by oral treatment with peptides that are stable and restricted to the gastrointestinal (GI) tissue. Novel inhibitory peptides that are uniquely resistant to oxidative/reductive conditions and proteolytic degradation in a variety of assays that mimic the various compartments of the GI environment were identified. Functionally, these peptides potently neutralize IL-23-mediated signaling in a transformed human cell line and in human primary cells. The binding of IL-23R is selective, since the peptides do not block the interaction between IL-6 to IL-6R or antagonize the IL-12 signaling pathway. Furthermore, these orally delivered peptides are efficacious in attenuating colitis in a 2,4,6-trinitrobenzenesulfonic acid (TNBS)-induced acute rat model of IBD, as shown by a significant reduction in the ratio of colon weight to length, colon macroscopic score, neutrophil infiltration, and histopathology comparable to that of the control anti-IL-23p19 mAb.

[0090] The present invention relates generally to peptides that have IL-23R antagonist activity, including both peptide monomers and peptide dimers. In certain embodiments, this invention demonstrates a new paradigm for treatment of IBD and other diseases and disorders by oral delivery of antagonists of IL-23. IBD represents a local inflammation of the intestinal tissue; therefore, advantageous therapeutic agents would act from the luminal side of the intestine, yielding high drug concentrations in diseased tissue, minimizing systemic availability and resulting in improved efficacy and safety when compared to systemic approaches. Oral administration of the compounds of the present invention is expected to maximize drug levels in diseased intestinal tissues while limiting drug concentrations in circulation, thereby providing efficacious, safe, and durable delivery for life-long treatment of IBD and other diseases and disorders.

[0091] In certain embodiments, the present invention relates to various peptides, or peptide dimers comprising hetero- or homo-monomer subunits, that form cyclized structures through disulfide or other bonds. In certain embodiments, the disulfide or other bonds are intramolecular

bonds. The cyclized structure of the peptide monomer inhibitors and the monomer subunits of the peptide dimer inhibitors has been shown to increase potency and selectivity of the peptide inhibitors. In certain embodiments, a peptide dimer inhibitor may include one or more intermolecular bonds linking the two monomer peptide subunits within the peptide dimer inhibitor, e.g., an intermolecular bridge between two cysteine residues, one in each peptide monomer subunit.

**[0092]** The present invention provides peptide inhibitors that bind to IL-23R, which may be monomers or dimers. In particular embodiments, the peptide inhibitors inhibit the binding of IL-23 to IL-23R. In certain embodiments, the IL-23R is human IL-23R, and the IL-23 is human IL-23. In certain embodiments, a peptide inhibitor of the present invention reduces IL-23 binding to IL-23R by at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% as compared to a negative control peptide. Methods of determining binding are known in the art and include ELISA assays, as described in the accompanying Examples.

**[0093]** In certain embodiments, a peptide inhibitor of the present invention has an IC<sub>50</sub> of > 1 mM, < 1 mM, 500 nM to 1000 nM, < 500 nM, < 250 nM, < 100 nM, < 50 nM, < 25 nM, < 10 nM, < 5 nM, < 2 nM, < 1 nM, or < 5 mM, e.g., for inhibiting binding of IL-23 to IL-23R (e.g., human IL-23 and human IL-23R). Methods of determining activity are known in the art and include any of those described in the accompanying Examples.

**[0094]** In certain embodiments, a peptide inhibitor of the present invention has increased stability, increased gastrointestinal stability, or increased stability in stimulated intestinal fluid (SIF) or simulated gastric fluid (SGF), and/or under redox conditions (e.g., DTT) as compared to a control peptide. In certain embodiments, a control peptide is an unrelated peptide of the same or similar length. In particular embodiments, a control peptide is a peptide having the identical or a highly related amino acid sequence (e.g., > 90% sequence identity) as the peptide inhibitor. In particular embodiments, a control peptide is a peptide having the identical or a highly related amino acid sequence (e.g., > 90% sequence identity) as the peptide inhibitor, but which does not have a cyclized structure, e.g., through an intramolecular bond between two amino acid residues within the control peptide, or which is not dimerized, or which does not comprise a conjugate for

stabilization. In particular embodiments, the only difference between the peptide inhibitor and the control peptide is that the peptide inhibitor comprises one or more amino acid substitutions that introduce one or more amino acid residues into the peptide inhibitor, wherein the introduced amino residue(s) forms an intrasulfide disulfide or thioether bond with another amino acid residue in the peptide inhibitor. One example of a control for a peptide dimer inhibitor is a monomer having the same sequence as one of the monomer subunits present in the peptide dimer inhibitor. One example of a control for a peptide inhibitor comprising a conjugate is a peptide having the same sequence but not including the conjugated moiety. In certain embodiments, a control peptide is a peptide (e.g., a naturally-occurring peptide) corresponding to a region of IL-23 that binds to IL-23R.

**[0095]** Methods of determining the stability of a peptide are known in the art. In certain embodiments, the stability of a peptide inhibitor is determined using an SIF assay, e.g., as described in Example 3. In certain embodiments, the stability of a peptide inhibitor is determined using an SGF assay, e.g., as described in Example 3. In particular embodiments, a peptide inhibitor has a half-life (e.g., in SIF or SGF or DTT) under a given set of conditions (e.g., temperature) of greater than 1 minute, greater than 10 minutes, greater than 20 minutes, greater than 30 minutes, greater than 60 minutes, greater than 90 minutes, greater than 120 minutes, greater than 3 hours, or greater than four hours when exposed to SIF or SGF or DTT. In certain embodiments, the temperature is about 25 °C, about 4 °C, or about 37 °C, and the pH is a physiological pH, or a pH about 7.4.

**[0096]** In some embodiments, the half-life is measured in vitro using any suitable method known in the art, e.g., in some embodiments, the stability of a peptide of the present invention is determined by incubating the peptide with pre-warmed human serum (Sigma) at 37 °C. Samples are taken at various time points, typically up to 24 hours, and the stability of the sample is analyzed by separating the peptide or peptide dimer from the serum proteins and then analyzing for the presence of the peptide or peptide dimer of interest using LC-MS.

**[0097]** In some embodiments, a peptide inhibitor of the present invention exhibits improved solubility or improved aggregation characteristics as compared to a control peptide. Solubility may be determined via any suitable method known in the art. In some embodiments, suitable

methods known in the art for determining solubility include incubating peptides in various buffers (Acetate pH4.0, Acetate pH5.0, Phos/Citrate pH5.0, Phos Citrate pH6.0, Phos pH 6.0, Phos pH 7.0, Phos pH7.5, Strong PBS pH 7.5, Tris pH7.5, Tris pH 8.0, Glycine pH 9.0, Water, Acetic acid (pH 5.0 and other known in the art) and testing for aggregation or solubility using standard techniques. These include, but are not limited to, visual precipitation, dynamic light scattering, Circular Dichroism and fluorescent dyes to measure surface hydrophobicity, and detect aggregation or fibrillation, for example. In some embodiments, improved solubility means the peptide is more soluble in a given liquid than is a control peptide. In some embodiments, improved aggregation means the peptide has less aggregation in a given liquid under a given set of conditions than a control peptide.

[0098] In certain embodiments advantageous for achieving high compound concentrations in intestinal tissues when delivered orally, peptide inhibitors of the present invention are stable in the gastrointestinal (GI) environment. Proteolytic metabolism in the GI tract is driven by enzymes (including pepsins, trypsin, chymotrypsin, elastase, aminopeptidases, and carboxypeptidase A/B) that are secreted from the pancreas into the lumen or are produced as brush border enzymes. Proteases typically cleave peptides and proteins that are in an extended conformation. In the reducing environment of intestinal fluids, disulfide bonds may be broken, resulting in a linear peptide and rapid proteolysis. This luminal redox environment is largely determined by the Cys/CySS redox cycle. In enterocytes, relevant activities include numerous digestive enzymes such as CYP450 and UDP-glucuronyl-transferase. Finally, bacteria, present in the large intestine at concentration ranging from  $10^{10}$  to  $10^{12}$  CFU/ml, constitute another metabolic barrier. In certain embodiments, the peptide inhibitors are stable to various pHs that range from strongly acidic in the stomach (pH 1.5-1.9), trending towards basic in the small intestine (pH 6-7.5), and then weakly acidic in the colon (pH 5-7). Such peptide inhibitors are stable during their transit through the various GI compartments, a process that has been estimated to take 3-4 h in the intestine and 6-48 h in the colon.

[0099] In some embodiments, the peptide inhibitors of the present invention have less degradation, e.g., over a period of time (i.e., more degradation stability), e.g., greater than or about 10% less, greater than or about 20% less, greater than or about 30% less, greater than or about 40 less, or greater than or about 50% less degradation than a control peptide. In some

embodiments, degradation stability is determined via any suitable method known in the art. In some embodiments, the degradation is enzymatic degradation. For example, in certain embodiments, the peptide inhibitors have reduced susceptibility to degradation by trypsin, chhromotrypsin or elastase. In some embodiments, suitable methods known in the art for determining degradation stability include the method described in Hawe et al., *J Pharm Sci*, VOL. 101, No. 3, 2012, p 895-913, incorporated herein in its entirety. Such methods are in some embodiments used to select potent peptide sequences with enhanced shelf lifes. In particular embodiments, peptide stability is determined using a SIF assay or SGF assay as described herein.

**[00100]** In certain embodiments, peptide inhibitors of the present invention inhibit or reduce IL-23-mediated inflammation. In related embodiments, peptide inhibitors of the present invention inhibit or reduce IL-23-mediated secretion of one or more cytokines, e.g., by binding to IL-23R on the cell surface, thus inhibiting IL-23 binding to the cell. In particular embodiments, peptide inhibitors of the present invention inhibit or reduce IL-23-mediated activation of Jak2, Tyk2, Stat1, Stat3, Stat4, or Stat5. Methods of determining inhibition of cytokine secretion and inhibition of signaling molecules are known in the art. For example, inhibition of IL-23/IL-23R signaling may be determined by measuring inhibition of phospho-Stat3 levels in cell lysates, as described in the accompanying Examples, including Example 2.

**[00101]** In certain embodiments, peptide inhibitors of the present invention inhibit or reduce IL-23-mediated inflammation. In related embodiments, peptide inhibitors of the present invention inhibit or reduce IL-23-mediated secretion of one or more cytokines, e.g., by binding to IL-23R on the cell surface, thus inhibiting IL-23 binding to the cell. In particular embodiments, peptide inhibitors of the present invention inhibit or reduce IL-23-mediated activation of Jak2, Tyk2, Stat1, Stat3, Stat4, or Stat5. Methods of determining inhibition of cytokine secretion and inhibition of signaling molecules are known in the art. For example, inhibition of IL-23/IL-23R signaling may be determined by measuring inhibition of phospho-Stat3 levels in cell lysates, as described in the accompanying Examples, including Example 2.

**[00102]** In certain embodiments, peptide inhibitors have increased redox stability as compared to a control peptide. A variety of assays that may be used to determine redox stability are known

and available in the art. Any of these may be used to determine the redox stability of peptide inhibitors of the present invention.

[00103] In certain embodiments, the present invention provides various peptide inhibitors that bind or associate with the IL-23R, *in vitro* or *in vivo*, to disrupt or block binding between IL-23 and IL-23R. In certain embodiments, the peptide inhibitors bind and/or inhibit human IL-23R. In certain embodiments, the peptide inhibitors bind and/or inhibit both human and rodent IL-23R. In certain embodiments, the peptide inhibitors bind and/or inhibit both human and rat IL-23R. In particular embodiments, the peptide inhibitors inhibit rat IL-23R at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95% as well as they bind or inhibit human IL-23R, e.g., as determined by an assay described herein. In certain embodiments, the peptide inhibitors preferentially bind and/or inhibit human and/or rat IL-23R as compared to mouse IL-23R. In particular embodiments, the peptide inhibitors preferentially bind to rat IL-23R as compared to mouse IL-23R. In particular embodiments, the peptide inhibitors preferentially bind to human IL-23R as compared to mouse IL-23R. In certain embodiments, binding of a peptide inhibitor to mouse IL-23R is less than 75%, less than 50%, less than 40%, less than 30%, less than 20%, or less than 10% of binding of the same peptide inhibitor to human IL-23R and/or rat IL-23R. In certain embodiments of peptide inhibitors that preferentially bind and/or inhibit human IL-23R and/or rat IL-23R as compared to mouse IL-23R, the peptide inhibitor binds to a region of IL-23R that is disrupted by the presence of additional amino acids present in mouse IL-23R but not human IL-23R or rat IL-23. In one embodiment, the additional amino acids present in the mouse IL-23R are in the region corresponding to about amino acid residue 315 to about amino acid residue 340 of the mouse IL23R protein, e.g., amino acid region NWQPWSSPFVHQTSQETGKR (see, e.g., Figure 4). In particular embodiments, the peptide inhibitors bind to a region of human IL-23R from about amino acid 230 to about amino acid residue 370.

[00104] In certain embodiments, peptide inhibitors show GI-restricted localization following oral administration. In particular embodiments, greater than 50%, greater than 60%, greater than 70%, greater than 80%, or greater than 90% of orally administered peptide inhibitor is localized to gastrointestinal organs and tissues. In particular embodiments, blood plasma levels of orally administered peptide inhibitor are less than 20%, less than 10%, less than 5%, less than 2%, less

than 1% or less than 0.5% the levels of peptide inhibitor found in the small intestine mucosa, colon mucosa, or proximal colon.

**[00105]** The various peptide inhibitors of the invention may be constructed solely of natural amino acids. Alternatively, the peptide inhibitors may include non-natural amino acids including, but not limited to, modified amino acids. In certain embodiments, modified amino acids include natural amino acids that have been chemically modified to include a group, groups, or chemical moiety not naturally present on the amino acid. The peptide inhibitors of the invention may additionally include one or more D-amino acids. Still further, the peptide inhibitors of the invention may include amino acid analogs.

**[00106]** In certain embodiments, peptide inhibitors of the present invention include one or more modified or unnatural amino acids. For example, in certain embodiments, a peptide inhibitor includes one or more of Dab, Dap, Pen, Sarc, Cit, Cav, hLeu, 2-Nal, D-1-Nal, D-2-Nal, Phe(4-OMe),  $\beta$ hTrp,  $\alpha$ -MePhe,  $\alpha$ -MeTyr,  $\alpha$ -MeTrp,  $\beta$ -HPhe, Phe(4-CF<sub>3</sub>), 2-2-Indane, 1-1-Indane, Cyclobutyl,  $\beta$ -hPhe, Gla, Phe(4-NH<sub>2</sub>), hPhe, 1-Nal, Nle, homoamino acids, D-amino acids, 4,4'-Biphenylalanine (Bip), cyclobutyl-Ala, hCha,  $\beta$ hPhe,  $\beta$ Glu, Phe(4-Guanidino), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminooethoxy)], Phe(4-CONH<sub>2</sub>), Phe(4-Me), Tyr(Bzl), or Tyr(Me), Phe(3,4-diF<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminooethoxy)], Phe(Br), Phe(4-CONH<sub>2</sub>), Phe(Cl), Phe(4-CN), Phe(4-guadino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N3), Tyr, Tyr(Bzl), or Tyr(Me), Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl<sub>2</sub>), Tyr(3-tBu), and various N-methylated amino acids and alpha-methyl amino acids. In some embodiments of the present invention, a peptide inhibitor includes one or more non-natural amino acids shown in Table 1A. One having skill in the art will appreciate that other modified or unnatural amino acids, and various other substitutions of natural amino acids with modified or unnatural amino acids, may be made to achieve similar desired results, and such substitutions are within the teaching and spirit of the present invention. In certain embodiments, peptide inhibitors of the present invention include any of those described herein, including but not limited to any of those comprising an amino acid sequence or peptide inhibitor structure shown in any one of the tables herein, the accompanying sequence listing or the accompanying figures, wherein one or more residues is substituted with a modified or unnatural amino acid.

[00107] The present invention also includes any of the peptide inhibitors described herein in either a free or a salt form. Thus, embodiments of any of the peptide inhibitors described herein (and related methods of use thereof) include a pharmaceutically acceptable salt of the peptide inhibitor.

[00108] The present invention also includes variants of any of the peptide inhibitors described herein, including but not limited to any of those comprising a sequence shown in any one of the tables herein, the accompanying sequence listing or the accompanying figures, wherein one or more L-amino acid residue is substituted with the D isomeric form of the amino acid residue, e.g., an L-Ala is substituted with a D-Ala.

[00109] In particular embodiments of the peptide inhibitors described herein, they comprise one or more unnatural or non-natural amino acid residue.

[00110] The present invention also includes any of the peptide monomer inhibitors described herein linked to a linker moiety, including any of the specific linker moieties described herein. In particular embodiments, a linker is attached to an N-terminal or C-terminal amino acid, while in other embodiments, a linker is attached to an internal amino acid. In particular embodiments, a linker is attached to two internal amino acids, e.g., an internal amino acid in each of two monomer subunits that form a dimer. In some embodiments of the present invention, a peptide inhibitor is attached to one or more linker moieties shown.

[00111] The present invention also includes peptides and peptide dimers comprising a peptide having at least 90%, at least 95%, at least 98%, or at least 99% sequence identity to the peptide sequence of a peptide inhibitor described herein. In particular embodiments, peptide inhibitors of the present invention comprise a core peptide sequence and one or more N-terminal and/or C-terminal modification (e.g., Ac and NH<sub>2</sub>) and/or one or more conjugated linker moiety and/or half-life extension moiety. As used herein, the core peptide sequence is the amino acid sequence of the peptide absent such modifications and conjugates. For example, for the peptide inhibitor: [Palm]-[isoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>, the core peptide sequence is: [Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NN.

[00112] In certain embodiments, a peptide inhibitor or a monomer subunit of a peptide inhibitor of the present invention comprises, consists essentially of, or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 20 amino acid residues, 8 to 20 amino acid residues, 9 to 20 amino acid residues, 10 to 20 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues, and, optionally, one or more additional non-amino acid moieties, such as a conjugated chemical moiety, e.g., a PEG or linker moiety. In particular embodiments, a peptide inhibitor of the present invention (or a monomer subunit thereof), including but not limited to those of any embodiments of Formula X, Formula I, Formula II, Formula III, Formula IV, or Formula V is greater than 10, greater than 12, greater than 15, greater than 20, greater than 25, greater than 30 or greater than 35 amino acids, e.g., 35 to 50 amino acids. In certain embodiments, a peptide inhibitor (or a monomer subunit thereof) is less than 50, less than 35, less than 30, less than 25, less than 20, less than 15, less than 12, or less than 10 amino acids. In particular embodiments, a monomer subunit of a peptide inhibitor (or a peptide monomer inhibitor) comprises or consists of 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, or 35 amino acid residues. In particular embodiments, a monomer subunit of a peptide inhibitor of the present invention comprises or consists of 10 to 18 amino acid residues and, optionally, one or more additional non-amino acid moieties, such as a conjugated chemical moiety, e.g., a PEG or linker moiety. In various embodiments, the monomer subunit comprises or consists of 7 to 35 amino acid residues, 7 to 20 amino acid residues, 8 to 20 amino acid residues, 9 to 20 amino acid residues, 10 to 20 amino acid residues, 8 to 18 amino acid residues, 8 to 19 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues. In particular embodiments of any of the various Formulas described herein, X comprises or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues.

[00113] Certain illustrative peptide inhibitors described herein comprise 12 or more amino acid residues. However, the present invention also includes peptide inhibitors comprising a fragment

of any of the peptide sequences described herein, including peptide inhibitors having 7, 8, 9, 10, or 11 amino acid residues. For example, peptide inhibitors of the present invention include peptides comprising or consisting of X4-X9, X4-X10, X4-X11, X4-X12, X4-X13, X4-X14, X4-X15, or X4-X16. In particular embodiments, the present invention includes peptide inhibitors having any of the sequences described herein, including but not limited to, those shown in any of the formulas described herein, the sequence listing, or any of the tables provided herein, wherein one or more of X10, X11, X12, X13, X14, X15, or X16 is absent. In particular embodiments, one or more of X13, X14, X15 or X16 is absent.

[00114] In particular embodiments of the present invention, the peptide inhibitors, or X regions thereof, are not present within an antibody. In particular embodiments, the peptide inhibitors, or X regions thereof, are not present within a V<sub>H</sub> or V<sub>L</sub> region of an antibody.

[00115] In particular embodiments of the peptide inhibitors described herein, they comprise one or more unnatural or non-natural amino acid residue.

[00116] In particular embodiments, peptide inhibitors of the present invention are cyclized via a cyclic amide bond, a disulfide bond, or a thioether bond. In particular embodiments, the bond is an intramolecular bond between two amino acid residues within the peptide inhibitor or a monomer subunit thereof.

### Peptide Inhibitors

[00117] Peptide inhibitors of the present invention include peptides having any of the amino acid sequences described herein, compounds having any of the structures described herein, including compounds comprising any of the peptide sequences described herein, and dimers of any of such peptides and compounds. Peptide inhibitors on the present invention include both peptides not having and those having a bond between X4 and X9, e.g., before and after a cross-link is introduced between X4 and X9. Illustrative peptides of the invention comprise an amino acid sequence or structure described in any of the accompanying tables, Examples, figures and sequence listing.

[00118] In certain embodiments, the present invention includes a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Xa):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Xa)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is any amino acid or chemical moiety capable of forming a bond with X9;

X5 is any amino acid;

X6 is any amino acid;

X7 is any amino acid;

X8 is any amino acid;

X9 is any amino acid or chemical moiety capable of forming a bond with X4;

X10 is any amino acid;

X11 is any amino acid;

X12 is any amino acid;

X13 is any amino acid;

X14 is any amino acid;

X15 is any amino acid,

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00119]wherein X4 and X9 are capable of forming a bond with each other. In particular embodiments, the bond is a disulfide bond, a thioether bond, a lactam bond, a triazole ring, a selenoether bond, a diselenide bond, or an olefin bond. In particular embodiments, the bond is a disulfide bond or a thioether bond. In certain embodiments, the peptide inhibitor is cyclized via the bond between X4 and X9. In certain embodiments, the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor. In particular embodiments, when X4 is not an amino acid, then X1, X2, and X3 are absent. In certain embodiments, X1 is a D-amino acid or absent. In certain embodiments, X2 is a D-amino acid or absent. In certain embodiments, X3 is a D-amino acid or absent. In certain embodiments, X16 is a D-amino acid or absent. In certain embodiments, X17 is a D-amino acid or absent. In certain embodiments, X18 is a D-amino acid or absent. In certain embodiments, X19 is a D-amino acid or absent. In certain embodiments, X20 is a D-amino acid or absent.

[00120]In one embodiment of the peptide inhibitor of Formula Xa,

X1 is absent;

X2 is absent;

X3 is Glu, D-Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln or absent;

X4 is Cys, Abu or Pen;

X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn, Gln, Arg, Ser or Thr;

X6 is Asp or Thr;

X7 is Trp or 6-Chloro-Trp;

X8 is Glu, Gln or Val;

X9 is Cys, Abu or Pen;

X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog, wherein in particular embodiments, X10 is 2-Nal, Phe(3,4-diF<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminoethoxy)], Phe(Br), Phe(4-CONH<sub>2</sub>), Phe(Cl), Phe(4-CN), Phe(4-guadino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N3), Tyr, Tyr(Bzl), or Tyr(Me);

X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl<sub>2</sub>), Trp or Tyr(3-tBu);

X12 is 3-Pal, Acpc, Acbc, Acvc, Achc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu, a-  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cav, Cha, Cit, Cpa, D-Asn, Glu, His, hLeu, hArg, Lys, Leu, Octgly, Orn, piperidine, Arg, Ser, Thr or THP;

X13 is Cit, Asp, Dab, Dap, Phe, His, Dap(Peg2-Ac), Dap(pyroglutaric acid), Glu, hArg, Lys, Lys(Ac), Lys(Benzoic acid), Lys(glutaric acid), Lys(IVA), Lys(Peg4-isoGlu-Palm), Lys(pyroglutaric acid), Lys-succinic acid, Asn, Orn, Gln, Arg, Thr or Val;

X14 is Asp, Dab(Ac), Dap(Ac), Phe, His, Lys(Ac), Met, Asn(isobutyl), Gln, Arg, Tyr or Asp(1,4-diaminobutane);

X15 is Ala, betaAla, Glu, Gly, Asn, Gln, Arg or Ser,

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

[00121] IN certain embodiments, X3 is absent. In particular embodiments, X16, X17, X18, X19 and X20 are absent. In particular embodiments, X4 and X9 are Cys, and X4 and X9 are linked via a disulfide bond. In particular embodiments, X4 is Abu and X9 is Pen, and X4 and X9 are linked via a thioether bond. In particular embodiments, X4 is Abu and X9 is Cys, and X4 and X9 are linked via a thioether bond.

[00122] In another embodiment of the peptide inhibitor of Formula Xa,

X1 is absent;

X2 is absent;

X3 is Glu, D-Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln or absent;

X4 is Cys, Abu or Pen;

X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, Orn, Gln, Arg, Ser or Thr;

X6 is Asp or Thr;

X7 is Trp or 6-Chloro-Trp;

X8 is Gln or Val;

X9 is Cys, Abu or Pen;

X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog, wherein in particular embodiments, X10 is 2-Nal, Phe(3,4-diF<sub>2</sub>), Phe(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminoethoxy)], Phe(Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guadino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Tyr, Tyr(Bzl), or Tyr(Me);

X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl<sub>2</sub>), Trp or Tyr(3-tBu);

X12 is 3-Pal, Acpc, Acbc, Acvc, Achc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cav, Cha, Cit, Cpa, D-Asn, His, hLeu, hArg, Lys, Leu, Octgly, Orn, 4-amino-4-carboxy-piperidine, or THP;

X13 is Cit, Asp, Dab, Dap, Phe, His, Dap(Peg2-Ac), Dap(pyroglutaric acid), Glu, hArg, Lys, Lys(Ac), Lys(Benzoic acid), Lys(glutaric acid), Lys(IVA), Lys(Peg4-isoGlu-Palm), Lys(pyroglutaric acid), Lys-succinic acid, Asn, Orn, Gln, Arg, Thr or Val;

X14 is Dab(Ac), Dap(Ac), Phe, His, Lys(Ac), Met, Asn, Gln, Arg, or Tyr;

X15 is Ala,  $\beta$ Ala, Gly, Asn, Gln, or Ser,

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00123]** In some embodiments, X3 is absent. In particular embodiments, X16, X17, X18, X19 and X20 are absent. In particular embodiments, X4 and X9 are Cys, and X4 and X9 are linked via a disulfide bond. In particular embodiments, X4 is Abu and X9 is Pen, and X4 and X9 are linked via a thioether bond. In particular embodiments, X4 is Abu and X9 is Cys, and X4 and X9 are linked via a thioether bond.

**[00124]** In another embodiment of the peptide inhibitor of Formula Xa,

X1 is absent;

X2 is absent;

X3 is Glu, D-Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln or absent;

X4 is Cys, Abu or Pen;

X5 is Dap, Dap(Ac), Gly, Lys, Gln, Arg, Ser, Thr or Asn;

X6 is Thr;

X7 is Trp or 6-Chloro-Trp;

X8 is Gln;

X9 is Cys, Abu or Pen;

X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog, wherein in particular embodiments, X10 is 2-Nal, Phe(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminooethoxy)], Phe(4-CONH<sub>2</sub>), Phe(4-Me), Phe(4-NH<sub>2</sub>), Tyr, Tyr(Bzl), or Tyr(Me);

X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), Phe(3,4-Cl<sub>2</sub>), or Trp;

X12 is Acpc, Acbc, Acvc, Achc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, hLeu, Lys, Leu, Arg or THP;

X13 is Cit, Asp, Dap, Dap(Peg2-Ac), Dap(pyroglutaric acid), Glu, hArg, Lys, Lys(Ac), Lys(Benzoic acid), Lys(glutaric acid), Lys(IVA), Lys(Peg4-isoGlu-Palm), Lys(pyroglutaric acid), Lys(succinic acid), Asn, Orn, Gln, Arg, or Val;

X14 is Dab(Ac), Dap(Ac), His, Lys(Ac), Asn, Gln, or Tyr;

X15 is Ala, betaAla, Gly, Asn, Gln, or Ser,

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

[00125] In some embodiments, X3 is absent. In particular embodiments, X16, X17, X18, X19 and X20 are absent. In particular embodiments, X4 and X9 are Cys, and X4 and X9 are linked via a disulfide bond. In particular embodiments, X4 is Abu and X9 is Pen, and X4 and X9 are linked via a thioether bond. In particular embodiments, X4 is Abu and X9 is Cys, and X4 and X9 are linked via a thioether bond.

[00126] In another embodiment of the peptide inhibitor of Formula Xa,

X1 is absent;

X2 is absent;

X3 is Glu, D-Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln or absent;

X4 is Cys, Abu or Pen;

X5 is Dap, Dap(Ac), Gln, Ser, Thr or Asn;

X6 is Thr;

X7 is Trp;

X8 is Gln;

X9 is Cys, Abu or Pen;

X10 is a Phe analog, Tyr, or a Tyr analog, wherein in particular embodiments, X10 is Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminoethoxy)], Phe(4-CONH<sub>2</sub>), Phe(4-Me), Tyr, Tyr(Bzl), or Tyr(Me);

X11 is 2-Nal or Trp;

X12 is Acpc, Acbc, Acvc, Achc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, hLeu, Leu, or THP;

X13 is Cit, Asp, Glu, Lys, Lys(Ac), Asn, or Gln;

X14 is Dab(Ac), Asn, or His;

X15 is Ala, betaAla, Gly, Asn, or Gln;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

[00127] In some embodiments, X3 is absent. In particular embodiments, X16, X17, X18, X19 and X20 are absent. In particular embodiments, X4 and X9 are Cys, and X4 and X9 are linked via a disulfide bond. In particular embodiments, X4 is Abu and X9 is Pen, and X4 and X9 are linked via a thioether bond. In particular embodiments, X4 is Abu and X9 is Cys, and X4 and X9 are linked via a thioether bond.

[00128] In particular embodiments, the peptide inhibitor comprises the amino acid sequence set forth in any of the various formula described herein, e.g., Ia-It, IIa-IIId, IIIa-IIIe, or IV.

[00129] In certain embodiments, the present invention includes a peptide inhibitor of an interleukin-23 receptor, wherein the peptide inhibitor has the structure of Formula I:



[00130] or a pharmaceutically acceptable salt or solvate thereof,

[00131] wherein R<sup>1</sup> is a bond, hydrogen, an C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing;

[00132] R<sup>2</sup> is a bond, OH or NH<sub>2</sub>; and

[00133] X is an amino acid sequence, e.g., an amino acid comprising 7 to 35 amino acid residues. In certain embodiments, R<sup>2</sup> is OH or NH<sub>2</sub>.

[00134] In certain embodiments, X comprises a sequence of Formula Xa.

[00135] In particular embodiments of formula (I), X comprises the sequence of Formula Ia:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Ia)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Sec, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-(3'-butenyl)glycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, 2-(5'-hexenyl)glycine or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, CitDap, Dab, Dap (Ac), Gly, Lys, Asn, N-Me-Gln, N-Me-Arg, Orn or Gln,

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Leu, Val, Phe, Ser, Sec, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, or 2-(5'-hexenyl)glycine;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu)

X12 is His, Phe, Arg, N-Me-His, or Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac), or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

In particular embodiments of Ia: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal or 2-Nal; X12 is His, Phe, Arg, N-Me-His, or Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00136] In particular embodiments, X4 is present.

[00137] In certain embodiments, the peptide inhibitor is cyclized.

[00138] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00139] In certain embodiments, the peptide inhibitor is cyclized, or contains an intramolecular bond, between X4 and X9.

[00140] In certain embodiments of Formula I, X comprises the sequence of Formula Ib:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Ib),

[00141] wherein:

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Sec, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-(3'-butenyl)glycine, 2-2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, 2-(5'-hexenyl)glycine, or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, CitDap, Dab, Dap (Ac), Gly, Lys, Asn, N-Me-Gln, N-Me-Arg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Sec, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, or 2-(5'-hexenyl)glycine;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>), Tyr(3-t-Bu)

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAlaAib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln

X14 is Phe, Tyr, or  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, or Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid, or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00142]** In particular embodiments of Ib: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla or Aib; X14 is Phe, Tyr or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

**[00143]** In particular embodiments, X4 is present.

[00144] In certain embodiments, the peptide inhibitor is cyclized.

[00145] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00146] In certain embodiments, the peptide inhibitor is cyclized, or contains an intramolecular bond, between X4 and X9.

[00147] In certain embodiments of Formula I, X comprises the sequence of Formula Ic:

X1-X2-X3-X4-X5-X6-W-X8-X9-Y-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20

(Ic)

[00148] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Sec, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, 2-(5'-hexenyl)glycine, or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, CitDap, Dab, Dap (Ac), Gly, Lys, Asn, N-Me-Gln, N-Me-Arg, Orn or Gln

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Sec, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, or 2-(5'-hexenyl)glycine;

X11 is Trp, 1-Nal, 2-Nal Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu)

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala t-butyl-Gly; 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00149]** In particular embodiments of Ic, X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

**[00150]** In particular embodiments, X4 is present.

[00151] In certain embodiments, the peptide inhibitor is cyclized.

[00152] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00153] In certain embodiments, the peptide inhibitor is cyclized, or contains an intramolecular bond, between X4 and X9.

[00154] In certain embodiments of Formula I, X comprises the sequence of Formula Id:

X1-X2-X3-C-X5-X6-W-X8-C-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20

(Id)

[00155] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, CitDap, Dab, Dap (Ac), Gly, Lys, Asn, N-Me-Gln, N-Me-Arg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X10 is Tyr Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>), Tyr(3-t-Bu)

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -

MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His, Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00156] wherein X4 and X9 are optionally linked by an intramolecular disulphide bridge.

[00157] In certain embodiments of Id: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00158] In certain embodiments of Formula I, X comprises the sequence of Formula Ie:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20

(Ie)

[00159] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Pen, hCys, D-Pen, D-Cys, or D-hCys;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, CitDap, Dab, Dap (Ac), Gly, Lys, Asn, N-Me-Gln, N-Me-Arg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Pen, hCys, D-Pen, D-Cys, D-hCys;

X10 is Tyr, Phe Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00160] wherein X4 and X9 are optionally linked by an intramolecular disulphide bridge.

[00161] In certain embodiments of Ie: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00162] In particular embodiments, X4 is present.

[00163] In certain embodiments, the peptide inhibitor is cyclized.

[00164] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00165] In certain embodiments, the peptide inhibitor is cyclized, or contains an intramolecular bond, between X4 and X9.

[00166] In particular embodiments, X4 and X9 and both Pen.

[00167] In certain embodiments of Formula I, X comprises the sequence of Formula If:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20

(If)

[00168] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, or Asp;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap (Ac), Gly, Lys, Asn, N-Me-Gln, N-Me-Arg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, or Asp;

X10 is Tyr, Phe, Phe(3,4-F2), Phe(3,4-Cl2), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH2), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH2), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00169] wherein X4 and X9 are optionally cyclized through an intramolecular bond.

[00170] In certain embodiments of If: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe; X8 is Val, Gln, Glu, or Lys; X9 is Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, or Asp; X10 is Tyr or Phe;; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; ;X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00171] In certain embodiments, the intramolecular bond is a lactam bond.

[00172] In certain embodiments of Formula I, X comprises the sequence of Formula Ig:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20

(Ig)

[00173] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is  $\beta$ -azido-Ala-OH, or propargylglycine;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is  $\beta$ -azido-Ala-OH or propargylglycine, ;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu)

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, or Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00174] wherein X4 and X9 are optionally cyclized through an intramolecular triazole ring.

[00175] In particular embodiments of Ig: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe; X8 is Val, Gln, Glu, or Lys; X9 is  $\beta$ -azido-Ala-OH or propargylglycine; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00176] In certain embodiments of Formula I, X comprises the sequence of Formula Ih:

X1-X2-X3-C-X5-X6-W-X8-C-Y-X11-H-X13-F-X15-X16-X17-X18-X19-X20

(Ih)

[00177] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, or 2-(5'-hexenyl)glycine;

X5 is Ala, Arg, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, or Asn;

X8 is Val, Gln, or Glu;

X9 is 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, or 2-(5'-hexenyl)glycine;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu)

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00178] wherein X4 and X9 are optionally cyclized via an intramolecular ring closing methasis to give the corresponding olefins.

[00179] In particular embodiments of Ih: X5 is Ala, Arg, or Sarc; X6 is Asp, Thr, or Asn; X11 is Trp, 1-Nal, or 2-Nal; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, or Aib; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00180] In certain embodiments of Formula I, X comprises the sequence of Formula II:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (II),

[00181] wherein:

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-chloro-propanoic acid, 4-chloro-butyric acid, or 3-chloro-isobutyric acid;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, or Abu;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu)

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

**[00182]**wherein X4 and X9 are optionally cyclized via an intramolecular thioether bond.

**[00183]**In particular embodiments of Ii: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

**[00184]**In certain embodiments of Formula I, X comprises the sequence of Formula Ij:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Ij),

**[00185]**wherein:

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Sec, 2-chloromethylbenzoic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, or Abu;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Sec or Abu;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-aminoethoxy), Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

**[00186]**wherein X4 and X9 are optionally cyclized via an intramolecular thioseleno or diselenide bond.

**[00187]**In particular embodiments of Ij: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X10 is Tyr or Phe; X11 is

Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, or Aib; X14 is Phe, Tyr, or  $\beta$ hPhe; X15 is Gly, Ser, Thr, Gln, Ala, or Sarc; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00188] In certain embodiments of Formula I, X comprises the sequence of Formula I<sub>k</sub>:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (I<sub>k</sub>),

[00189] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Leu, Val, Phe, or Ser;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00190]** In particular embodiments of I<sub>k</sub>: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib or absent; X14 is Phe, Tyr,  $\beta$ hPhe or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, Leu, or absent; and X17 is Leu, Lys, Arg, or absent.

In certain embodiments of Formula I, X comprises or consists of the sequence of Formula II:

X1-X2-X3-X4-X5-X6-W-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (II),

**[00191]** wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, or Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Leu, Val, Phe, or Ser;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, , Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln or absent;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

[00192] In particular embodiments of II: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, or Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Val,  $\beta$ hAla, Aib or absent; X14 is Phe, Tyr,  $\beta$ hPhe or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, Leu, or absent; and X17 is Leu, Lys, Arg, or absent.

[00193] In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, or Aib.

[00194] In certain embodiments, X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp. In certain embodiments, X14 is Phe, Tyr, or  $\beta$ hPhe.

[00195] In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, or Sarc.

[00196] In certain embodiments, X12 is alpha amino acid, e.g., 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Aib,  $\alpha$ -MeGly(diethyl),  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal.

[00197] In certain embodiments, X13 is present.

[00198] In certain embodiments, X13 and X14 are present.

[00199] In certain embodiments, X13, X14 and X15 are present.

[00200] In particular embodiments, X4 is present.

[00201] In certain embodiments, the peptide inhibitor is cyclized.

[00202] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00203] In certain embodiments, the peptide inhibitor is cyclized, or contains an intramolecular bond, between X4 and X9.

[00204] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula Im:

X1-X2-X3-X4-X5-X6-W-X8-X9-Y-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Im),

[00205] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn, or Gln;

X6 is Asp, Thr, Asn, or Phe;

X8 is Val, Gln, Glu, or Lys;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Leu, Val, Phe, or Ser;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>); 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>), or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -

MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln or absent;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac), or Asp or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00206]** In certain embodiments of Im: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, or Sarc; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib or absent; X14 is Phe, Tyr,  $\beta$ hPhe or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

**[00207]** In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, or Aib. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln.

**[00208]** In certain embodiments, X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp. In certain embodiments, X14 is Phe, Tyr, or  $\beta$ hPhe.

**[00209]** In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, or Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, or Sarc.

[00210] In certain embodiments, X12 is alpha amino acid, e.g., 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Aib,  $\alpha$ -MeGly(diethyl),  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal.

[00211] In certain embodiments, X13 is present.

[00212] In certain embodiments, X13 and X14 are present.

[00213] In certain embodiments, X13, X14, and X15 are present.

[00214] In particular embodiments, X4 is present.

[00215] In certain embodiments, the peptide inhibitor is cyclized.

[00216] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00217] In certain embodiments, the peptide inhibitor is cyclized, or contains an intramolecular bond, between X4 and X9.

[00218] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula I in:

X1-X2-X3-C-X5-X6-W-X8-C-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (In)

[00219] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, or Phe;

X8 is Val, Gln, Glu, or Lys;

X10 is Tyr Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln or absent;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

**[00220]**wherein the Cys at position X4 and and the Cys at position X9 are optionally linked by a disulphide bridge.

**[00221]**In certain embodiments of In: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln; X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr; X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-

OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu); X12 is His, Phe, Arg, N-Me-His, Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln or absent; X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00222] In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, or Aib.

[00223] In certain embodiments, X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp. In certain embodiments, X14 is Phe, Tyr, or  $\beta$ hPhe.

[00224] In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, or Sarc.

[00225] In certain embodiments, X12 is an alpha amino acid, e.g., 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal.

[00226] In certain embodiments, X13 is present.

[00227] In certain embodiments, X13 and X14 are present.

[00228] In certain embodiments, X13, X14 and X15 are present.

[00229] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula Io:

X1-X2-X3-C-X5-X6-W-X8-C-Y-X11-H-X13-X14-X15-X16-X17-X18-X19-X20 (Io)

[00230] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, or Phe;

X8 is Val, Gln, Glu, or Lys;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln or absent;

X14 is Phe, Tyr, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00231] wherein the Cys at position X4 and the Cys at position X9 are optionally linked by a disulphide bridge.

[00232] In certain embodiments of Io: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln; X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>), 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu); X13 is

Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, Gln or absent; X14 is Phe, Tyr, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) Asp or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn or absent; X16 is Asp, Glu, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

[00233] In certain embodiments, X12 is an alpha amino acid, e.g., 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal.

[00234] In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla or Aib.

[00235] In certain embodiments, X14 is Phe, Tyr, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp. In certain embodiments, X14 is Phe or Tyr.

[00236] In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala or Sarc.

[00237] In certain embodiments, X13 is present.

[00238] In certain embodiments, X13 and X14 are present.

[00239] In certain embodiments, X13, X14 and X15 are present.

[00240] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula Ip:

X1-X2-X3-C-X5-X6-W-X8-C-Y-X11-H-X13-F-X15-X16-X17-X18-X19-X20 (Ip)

[00241] wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is Ala, Arg, Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, or Asn;

X8 is Val, Gln, or Glu;

X11 is Trp, 1-Nal, 2-Nal, Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, Gln or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

**[00242]**wherein the Cys at position X4 and and the Cys at position X9 are optionally linked by a disulphide bridge.

**[00243]**In certain embodiments of I<sub>p</sub>: X5 is Ala, Arg, or Sarc; X11 is Trp, 1-Nal, or 2-Nal; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, or absent; and X17 is Leu, Lys, Arg, or absent.

**[00244]**In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla or Aib.

[00245] In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala or Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala or Sarc.

[00246] In certain embodiments, X13 is present.

[00247] In certain embodiments, X13 and X14 are present.

[00248] In certain embodiments, X13, X14 and X15 are present.

[00249] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula Iq:

X1-X2-X3-C-X5-X6-W-X8-C-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Iq),

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, D-Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn or Gln;

X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, or D-Phe;

X8 is Val, Gln, Glu, or Lys;

X10 is Tyr, Phe, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, 2-Nal, , Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is His, Phe, Arg, N-Me-His, Val, or D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, Gln or absent;

X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac), Asp or absent;

X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg, Asn or absent;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

[00250] wherein the Cys at position X4 and and the Cys at position X9 are optionally linked.

[00251] In certain embodiments of Iq: X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, or D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib or absent; X14 is Phe, Tyr,  $\beta$ hPhe or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, Leu, or absent; and X17 is Leu, Lys, Arg, or absent.

[00252] In certain embodiments, X12 is alpha amino acid, e.g., 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal.

[00253] In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla or Aib.

[00254] In certain embodiments, X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac), or Asp. In certain embodiments, X14 is Phe, Tyr or  $\beta$ hPhe.

[00255] In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala or Sarc.

[00256] In certain embodiments, X13 is present.

[00257] In certain embodiments, X13 and X14 are present.

[00258] In certain embodiments, X13, X14 and X15 are present.

[00259] In certain embodiments, Iq comprises or consists of the sequence of Formula Iq':

X1-X2-X3-C-X5-X6-W-X8-C-X10-X11-X12-X13-X14-X15 (Iq'),

wherein X1-X14 have the definition provided for Iq, and

wherein the Cys at position X4 and the Cys at position X9 are optionally linked.

[00260] In certain embodiments of Iq': X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, or D-Sarc; X10 is Tyr or Phe; X11 is Trp, 1-Nal, or 2-Nal; X12 is His, Phe, Arg, N-Me-His, Val, or D-His, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala, or t-butyl-Gly; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Val, Aib or absent; X14 is Phe, Tyr,  $\beta$ hPhe or absent; X15 is Gly, Ser, Thr, Gln, Ala, Sarc or absent; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, Leu, or absent; and X17 is Leu, Lys, Arg, or absent.

[00261] In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, Aib, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, or Gln. In certain embodiments, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla or Aib.

[00262] In certain embodiments, X14 is Phe, Tyr,  $\beta$ hPhe, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp. In certain embodiments, X14 is Phe, Tyr or  $\beta$ hPhe.

[00263] In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala or Sarc,  $\beta$ -Ala, Glu, Arg or Asn. In certain embodiments, X14 is Phe, Tyr or  $\beta$ hPhe.

[00264] In certain embodiments, X13 is present.

[00265] In certain embodiments, X13 and X14 are present.

[00266] In certain embodiments, X13, X14 and X15 are present.

[00267] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula Ir:

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Ir)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Met, Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Sec, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-(3'-butenyl)glycine, 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, 2-(5'-hexenyl)glycine, Abu or absent;

X5 is any amino acid;

X6 is any amino acid;

X7 is Trp, Glu, Gly, Ile, Asn, Pro, Arg, Thr or OctGly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X8 is any amino acid;

X9 is Cys, Pen, hCys, D-Pen, D-Cys, D-hCys, Glu, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu, D-Lys, Asp, Leu, Val, Phe, or Ser, Sec, Abu,  $\beta$ -azido-Ala-OH, propargylglycine, 2-2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, Ala, hCys, Abu, Met, MeCys, (D)Tyr or 2-(5'-hexenyl)glycine;

X10 is Tyr, Phe(4-OMe), 1-Nal, 2-Nal, Aic,  $\alpha$ -MePhe, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, His, hPhe(3,4-dimethoxy), hTyr, N-Me-Tyr, Trp, Phe(4-CONH<sub>2</sub>), Phe(4-phenoxy), Thr, Tic, Tyr(3-tBu), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5-F<sub>2</sub>), Phe(4-CH<sub>2</sub>CO<sub>2</sub>H), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>), Bip, Cha, 4-PyridylAlanine,  $\beta$ hTyr, OctGly, Phe(4-N<sub>3</sub>), Phe(4-Br), Phe[4-(2-aminoethoxy)] or Phe, a Phe analog, a Tyr analog, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe (3,4-F<sub>2</sub>), Phe(4-CO<sub>2</sub>H),  $\beta$ hPhe(4-F),  $\alpha$ -Me-Trp, 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>), , Phe(3,4-OMe<sub>2</sub>),  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tri-tert-Butyl), Phe(4-Oallyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino, Phe(4-OBzl), Octgly, Glu(Bzl), 4-Phenylbenzylalanine, Phe[4-(2-aminoethoxy)], 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, 1,2,3,4-tetrahydro-norharman, Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine or Bip, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X12 is His, Phe, Arg, N-Me-His, or Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala,  $\alpha$ -MeLys, D-Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Tyr, Aib,  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\beta$ -Aib,  $\beta$ -Ala,  $\beta$ hAla,  $\beta$ hArg,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Glu, hArg, Ile, Lys, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Ser, Thr, Tle or t-butyl-Gly, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Asn, Cit, Lys, Arg, Orn, Val,  $\beta$ hAla, Lys(Ac), (D)Asn, (D)Leu, (D)Phe, (D)Thr, Ala,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, hLeu, Asn, Ogl, Pro, Gln, Ser,  $\beta$ -spiro-pip, Thr, Tba, Tle or Aib, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X14 is Phe, Tyr, Glu, Gly, His, Lys, Leu, Met, Asn, Lys(Ac), Dap(Ac), Asp, Pro, Gln, Arg, Ser, Thr, Tic or  $\beta$ hPhe, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X15 is Gly, Ser, Thr, Gln, Ala, (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Aea, Asp, Asn, Glu, Phe, Gly, Lys, Leu, Pro, Arg,  $\beta$ -Ala, or Sarc, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00268]** In particular embodiments, the peptide is cyclized via X4 and X9.

**[00269]** In particular embodiments, X3 is Glu, D-Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln.

**[00270]** In certain embodiments of Ir: X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe (3,4-F<sub>2</sub>), Phe(4-CO<sub>2</sub>H),  $\beta$ hPhe(4-F),  $\alpha$ -Me-Trp, 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tri-tert-Butyl), Phe(4-Oallyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino, Phe(4-OBzl), Octgly, Glu(Bzl), 4-Phenylbenzylalanine, Phe[4-(2-aminoethoxy)], 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, 1,2,3,4-tetrahydro-norharman, Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine or Bip, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X12 is His, Phe, Arg, N-Me-

His, or Val, Cav, Cpa, Leu, Cit, hLeu, 3-Pal, t-butyl-Ala,  $\alpha$ -MeLys, D-Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Tyr, Aib,  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\beta$ -Aib,  $\beta$ -Ala,  $\beta$ hAla,  $\beta$ hArg,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Glu, hArg, Ile, Lys, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Ser, Thr, Tle or t-butyl-Gly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Arg, Orn, Val,  $\beta$ hAla, Lys(Ac), (D)Asn, (D)Leu, (D)Phe, (D)Thr, Ala,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, hLeu, Asn, Ogl, Pro, Gln, Ser,  $\beta$ -spiro-pip, Thr, Tba, Tle or Aib, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X14 is Phe, Tyr, Glu, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or  $\beta$ hPhe, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X15 is Gly, Ser, Thr, Gln, Ala, (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Aea, Asp, Asn, Glu, Phe, Gly, Lys, Leu, Pro, Arg or Sarc, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X16 is Asp, Glu, Ala, AEA, AEP,  $\beta$ hAla, Gaba, Gly, Ser, Pro, Asn, Thr or absent, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; and X17 is Leu, Lys, Arg, Glu, Ser, Gly, Gln or absent, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing.

[00271] In certain embodiments, both X4 and X9 are Pen. In particular embodiments, X4 and X9 are cyclized via a disulfide bond.

[00272] In certain embodiments, X4 is Abu and X9 is Cys. In certain embodiments, X4 and X9 are cyclized via a thioether bond.

[00273] In particular embodiments, X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, Cys, Cit, Asp, Dab, Dap, Gly, His, hCys, Lys, Met, Asn, N-Me-Ala, N-Me-Asn, N-Me-Lys, N-Me-Gln, Orn, Pro, Pen, Gln, Val,  $\alpha$ Me-Lys,  $\alpha$ Me-Orn, or D-Sarc,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, or Gln. In certain embodiments, X5 is Gln or Asn. In particular embodiments, X5 is Ala, Arg, Glu, Phe, Leu, Thr, Ser, Aib, Sarc, D-Ala, D-Arg, D-Glu, D-Phe, D-Leu, D-Thr, D-Ser, D-Aib, Cys, Cit, Asp, Dab, Dap, Gly, His, hCys, Lys, Met, Asn, N-Me-Ala, N-Me-Asn, N-Me-Lys, N-Me-Gln, N-Me-Arg, Orn, Pro, Pen, Gln, Val,  $\alpha$ Me-Lys,  $\alpha$ Me-Orn, or D-Sarc. In certain embodiments, X5 is Gln.

[00274] In particular embodiments, X6 is Asp, Thr, Asn, Phe, D-Asp, D-Thr, D-Asn, Glu, Arg, Ser or D-Phe. In particular embodiments, X6 is Thr.

[00275] In particular embodiments, X7 is Trp.

[00276] In particular embodiments, X8 is Val, Gln, Glu, Phe, Asn, Pro, Arg, Thr, Trp or Lys. In particular embodiments, X8 is Gln.

[00277] In particular embodiments, X1, X2 and X3 are absent.

[00278] In certain embodiments, X11 is a Trp analog.

[00279] In particular embodiments, X10 is a Phe analog. In particular embodiments, X10 is Phe(4-OMe), Phe(4-CONH<sub>2</sub>), or Phe[4-(2-aminoethoxy)] (also referred to herein as Phe[4-2ae]). In particular embodiments, X10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)] (also referred to herein as Phe[4-2ae]).

[00280] In particular embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X11 is 2-Nal.

[00281] In certain embodiments, X12 is  $\alpha$ -MeLys, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, or  $\alpha$ -MeVal. In certain embodiments, X12 is  $\alpha$ -MeLys.

[00282] In certain embodiments, X13 is Glu or Lys(Ac). In certain embodiments, X13 is Glu.

[00283] In certain embodiments, X14 is Asn.

[00284] In certain embodiments, X15 is Gly or Asn. In certain embodiments, X15 is Gly.

[00285] In certain embodiments, one or more, two or more, three or more, or four or more of X16, X17, X18, X19 and X20 are absent. In particular embodiments, X16, X17, X18, X19 and X20 are absent.

[00286] In particular embodiments of Ir, X4 and X9 are Cys, X7 is Trp, and X18 is [(D)Lys]. In particular embodiments of Ir, X4 and X9 are Cys, X7 is Trp, X10 is Tyr, and X18 is [(D)Lys]. In particular embodiments of Ir, X4 and X9 are Cys, X7 is Trp, X1, X2 and X3 are absent, X17 is

absent, X18 is [(D)Lys], and X19 and X20 are absent. In particular embodiments of Ir, X4 and X9 are Cys, X7 and X11 are Trp, X10 is Tyr, and X18 is [(D)Lys]. In certain embodiments, X1, X2, and X3 are absent; and in certain embodiments, X17 is absent.

[00287] In particular embodiments of Ir, X4 and X9 are Pen, and X12 is  $\alpha$ -MeLys. In particular embodiments of Ir, X4 and X9 are Pen, X12 is  $\alpha$ -MeLys, and X16, X17, X18, X19 and X20 are absent. In particular embodiments of Ir, X4 and X9 are Pen, X12 is  $\alpha$ -MeLys, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, X16, X17, X18, X19 and X20 are absent, and X7 is Trp. In particular embodiments of Ir, X4 and X9 are Pen, X12 is  $\alpha$ -MeLys, X16, X17, X18, X19 and X20 are absent, and X7 is Trp. In particular embodiments of Ir, X4 and X9 are Pen, X7 is Trp, and X12 is  $\alpha$ -MeLys. In certain embodiments, X1, X2, and X3 are absent. In particular embodiments, there is a disulfide bond between X4 and X9.

[00288] In particular embodiments of Ir, X4 is Abu, X9 is Cys, and X12 is 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys, or  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, or  $\alpha$ -MeVal. In particular embodiments of Ir, X4 is Abu, X9 is Cys, and X12 is  $\alpha$ -MeLys. In particular embodiments of Ir, X4 is Abu, X9 is Cys, X12 is  $\alpha$ -MeLys, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys, or  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, or  $\alpha$ -MeVal and X16, X17, X18, X19 and X20 are absent. In particular embodiments of Ir, X4 is Abu, X9 is Cys, X12 is  $\alpha$ -MeLys, and X16, X17, X18, X19 and X20 are absent. In particular embodiments of Ir, X4 is Abu, X9 is Cys, X12 is  $\alpha$ -MeLys, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys, or  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, or  $\alpha$ -MeVal, X16, X17, X18, X19 and X20 are absent, and X7 is Trp. In particular embodiments of Ir, X4 is Abu, X9 is Cys, X12 is  $\alpha$ -MeLys, X16, X17, X18, X19 and X20 are absent, and X7 is Trp. In particular embodiments of Ir, X4 is Abu, X9 is Cys, X7 is Trp, and X12 is  $\alpha$ -MeLys. In certain embodiments, X1, X2, and X3 are absent. In particular embodiments, there is a thioether bond between X4 and X9.

[00289] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula Is:

X1-X2-X3-C-X5-X6-W-X8-C-X10-X11-X12-X13-X14-G-X16-X17-X18-X19-X20 (Is)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is any amino acid;

X6 is any amino acid;

X8 is any amino acid;

X10 is Tyr, 1-Nal, 2-Nal, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl) or Tyr;

X11 is Trp, 1-Nal, Phe(3,4-OMe<sub>2</sub>), 5-Hydroxy-Trp, Phe(3,4-Cl<sub>2</sub>) or Tyr(3-t-Bu);

X12 is Arg, Lys, His, hArg, Cit, Orn, 1-Nal, D-Ala, D-Leu, D-Phe, D-Asn, D-Asp, Agp, Leu, βhLeu, Aib, βhAla, βhVal, βhArg, hLeu, Dap, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib, α-DiethylGly, α-MeLys, α-MeLys(Ac), α-Me-Leu, α-MeOrn, α-MeSer, α-MeVal, Cha, Cit, Cpa, (D)Asn, Glu, hArg, or Lys;

X13 is Cha, Ogl, Aib, Leu, Val, Dab, Glu, Lys, βhLeu, βhAla, βhVal, βGlu, Lys(Ac), Cit, Asp, Dab, Dap, Glu, hArg, Lys, Asn, Orn, Lys(Ac), or Gln;

X14 is Phe, Tic, Asn, Tyr, Asn, Arg, Qln, Lys(Ac), His; Dap(Ac), Dab(Ac) or Asp;

X16 is any amino acid;

X17 is absent;

X18 is D-Lys;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

[00290] In particular embodiments of Is: X10 is Tyr, 1-Nal or 2-Nal; X11 is Trp or 1-Nal; X12 is Arg, Lys, His, hArg, Cit, Orn, 1-Nal, D-Ala, D-Leu, D-Phe, D-Asn, D-Asp, Agp, Leu,  $\beta$ hLeu, Aib,  $\beta$ hAla,  $\beta$ hVal,  $\beta$ hArg, hLeu or Dap; X13 is Cha, Ogl, Aib, Leu, Val, Dab, Glu, Lys,  $\beta$ hLeu,  $\beta$ hAla,  $\beta$ hVal or  $\beta$ GLu; X14 is Phe, Tic, Asn or Tyr; and X16 is AEA, Ala or  $\beta$ Ala.

[00291] In particular embodiments, X5 is Glu, Arg, Ala, N-Me-Arg, N-Me-Ala, N-Me-Gln, Orn, N-Me-Asn, N-Me-Lys, Ser, Gln, Orn, Asn or Dap. In particular embodiments, X5 is Glu, Arg, Ala, N-Me-Arg, N-Me-Ala, N-Me-Gln, Orn, N-Me-Asn, N-Me-Lys, Ser, Asn or Dap.

[00292] In particular embodiments, X6 is Asp or Thr.

[00293] In particular embodiments, X8 is Gln or Val.

[00294] In particular embodiments, the peptide of Is is cyclized via a disulfide bond between X4 and X9.

[00295] In certain embodiments of the peptide inhibitor of Formula I, X comprises or consists of the sequence of Formula It:

X1-X2-X3-C-X5-X6-W-X8-C-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (It)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X5 is any amino acid;

X6 is any amino acid;

X8 is any amino acid;

X10 is Tyr, 1-Nal, 2-Nal, Phe[4-(2-aminoethoxy)], Phe(4-CONH<sub>2</sub>), Phe(4-OMe);

X11 is Trp, 1-Nal, 2-Nal, Bip, , Phe(3,4-OMe<sub>2</sub>) 5-Hydroxy-Trp,;

X12 is Arg, His, 3-Pal, Leu, Thr, Gln, Asn, Glu, Ile, Phe, Ser, Lys, hLeu,  $\alpha$ -MeLeu, D-Leu, D-Asn, h-Leu, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer or  $\alpha$ -MeVal;

X13 is Thr, Glu, Tyr, Lys, Gln, Asn, Lys, Lys (Ac), Asp, Arg, Ala, Ser, Leu;

X14 is Phe, Tyr, Asn, Gly, Ser, Met, Arg, His, Lys, Leu or Gln;

X15 is Gly, Ser, Arg, Leu, Asp, Ala,  $\beta$ -Ala, Glu, Arg or Asn;

X16 is absent or any amino acid;

X17 is absent or any amino acid;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

**[00296]** In certain embodiments of It: X10 is Tyr, 1-Nal or 2-Nal; X11 is Trp, 1-Nal, 2-Nal or Bip; X12 is Arg, His, 3-Pal, Leu, Thr, Gln, Asn, Glu, Ile, Phe, Ser, Lys, hLeu,  $\alpha$ -MeLeu, D-Leu, D-Asn, or h-Leu; X13 is Thr, Glu, Tyr, Lys, Gln, Asn, Lys, Asp, Arg, Ala, Ser, Leu; X15 is Gly, Ser, Arg, Leu, Asp or Ala; X16 is absent or Asn, Glu, Phe, Ala, Gly, Pro, Asp, Gln, Ser, Thr, D-Glu or Lys; and X17 is absent or Pro, Arg, Glu, Asp, Ser, Gly or Gln.

**[00297]** In particular embodiments, X5 is Ser, Asp, Asn, Gln, Ala, Met, Arg, His or Gly. In particular embodiments, X5 is Ser, Asp, Gln, Ala, Met, Arg, His or Gly.

[00298] In particular embodiments, X6 is any Asp, Ser or Thr.

[00299] In particular embodiments, X8 is Gln, Glu or Thr.

[00300] In particular embodiments, the peptide of It is cyclized via a disulfide bond between X4 and X9.

[00301] In a further embodiment, the present invention includes a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Va):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Va)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any D-amino acid or absent;

X4 is Cys, hCys, Pen, hPen, Abu, Ser, hSer or chemical moiety capable of forming a bond with X9;

X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn, Gln, Arg, Ser or Thr;

X6 is Thr, Ser, Asp, Ile or any amino acid;

X7 is Trp, 6-Chloro-Trp, 1-Nap or 2-Nap;

X8 is Glu, Gln, Asn, Lys(Ac), Cit, Cav, Lys(N- $\epsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), or Lys(N- $\epsilon$ -Palmitoyl;

X9 is Cys, hCys, Pen, hPen Abu, or any amino acid or chemical moiety capable of forming a bond with X4;

X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog;

X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl<sub>2</sub>), Trp or Tyr(3-tBu);

X12 is Aib, 4-amino-4-carboxy-tetrahydropyran, any alpha-methylamino acid, alpha-ethyl-amino acid, Achc, Acvc, Acbc, Acpc, 4-amino-4-carboxy-piperidine, 3-Pal, Agp,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu, a-  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cav, Cha, Cit, Cpa, D-Asn, Glu, His, hLeu, hArg, Lys, Leu, Octgly, Orn, piperidine, Arg, Ser, Thr or THP;

X13 is Lys(Ac), Gln, Cit, Glu, or any amino acid;

X14 is Asn, Gln, Lys(Ac), Cit, Cav, Lys(N- $\varepsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), Lys(N- $\varepsilon$ -Palmitoyl), or any amino acid;

X15 is  $\beta$ -Ala, Asn, Gly, Gln, Ala, Ser, Aib or Cit;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

wherein X4 and X9 are capable of forming a bond with each other. In particular embodiments, the bond is a ether, disulfide bond or a thioether bond. In certain embodiments, the peptide inhibitor is cyclized via the bond between X4 and X9.

[00302] In certain embodiments, X1 is a D-amino acid or absent. In certain embodiments, X2 is a D-amino acid or absent.

[00303] In certain embodiments, X16 is a D-amino acid or absent. In certain embodiments, X17 is a D-amino acid or absent. In certain embodiments, X18 is a D-amino acid or absent. In

certain embodiments, X19 is a D-amino acid or absent. In certain embodiments, X20 is a D-amino acid or absent.

**[00304]** In particular embodiments of Formula (Ia), X10 is 2-Nal, Phe(3,4-diF<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminooethoxy)], Phe(Br), Phe(4-CONH<sub>2</sub>), Phe(Cl), Phe(4-CN), Phe(4-guadino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Tyr, Tyr(Bzl), or Tyr(Me). In certain embodiments of Formula (Ia), X10 is Phe(4-ZR), Phe(3-ZR), or Phe(2-ZR), where R= CH<sub>2</sub>(CH<sub>2</sub>)<sub>n</sub>Y and n=1-25, Z=NH, O, CO, CONH, or CH<sub>2</sub>, and Y=NH<sub>2</sub>, CO<sub>2</sub>H, OH, or CH<sub>3</sub>.

**[00305]** In a further related embodiments, the present invention includes a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Vb):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Vb)

wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is D-Arg, D-Phe, any D amino acid or absent;

X4 is Cys, hCys, Pen, hPen, Abu, or a chemical moiety capable of forming a bond with X9;

X5 is Gln, Asn, Lys(Ac), Cit, Cav, Lys(N-ε-(N-α-Palmitoyl-L-γ-glutamyl)), or Lys(N-ε-Palmitoyl);

X6 is Thr, Ser, Asp, Ile or any amino acid;

X7 is Trp, 1-Nap or 2-Nap;

X8 is Gln, Asn, Lys(Ac), Cit, Cav, Lys(N-ε-(N-α-Palmitoyl-L-γ-glutamyl)), or Lys(N-ε-Palmitoyl);

X9 is Cys, hCys, Pen, hPen, Abu, any amino acid or a chemical moiety capable of forming a bond with X4;

X10 is Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminoethoxy)], Phe(4-CONH<sub>2</sub>), Phe(4-guadino), Phe(4-NH<sub>2</sub>), Tyr(Me) or Phe(4-ZR), where R= CH<sub>2</sub>(CH<sub>2</sub>)<sub>n</sub>Y; n=1-25; Z= O, CO, NH, CONH, or CH<sub>2</sub>; and Y=NH<sub>2</sub>, CO<sub>2</sub>H, OH, or CH<sub>3</sub>;

X11 is 2-Nal or Trp;

X12 is Aib, 4-amino-4-carboxy-tetrahydropyran,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, acid, Achc, Acvc, Acbc, Acpc, or 4-amino-4-carboxy-piperidine;

X13 is Lys(Ac), Gln, Cit, Glu, or any amino acid;

X14 is Asn, Gln, Lys(Ac), Cit, Cav, Lys(N- $\epsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), Lys(N- $\epsilon$ -Palmitoyl), or any amino acid;

X15 is  $\beta$ -Ala, Asn, Gln, Ala, Ser, or Aib;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

wherein X4 and X9 are capable of forming a bond with each other. In particular embodiments, the bond is a disulfide bond or a thioether bond. In certain embodiments, the peptide inhibitor is cyclized via the bond between X4 and X9.

[00306] In certain embodiments, X1 is a D-amino acid or absent. In certain embodiments, X2 is a D-amino acid or absent.

[00307] In certain embodiments, X16 is a D-amino acid or absent. In certain embodiments, X17 is a D-amino acid or absent. In certain embodiments, X18 is a D-amino acid or absent. In certain embodiments, X19 is a D-amino acid or absent. In certain embodiments, X20 is a D-amino acid or absent.

[00308] In another related embodiment, the present invention includes a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Vc):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Vc)

wherein

X1 is absent;

X2 is absent;

X3 is D-Arg or absent;

X4 is Cys, Pen, Abu, or a chemical moiety capable of forming a bond with X9;

X5 is Gln, Asn, Lys(Ac), Cit, or Cav;

X6 is Thr or Ser;

X7 is Trp, 1-Nap or 2-Nap;

X8 is Gln, Asn, Lys(Ac), Cit, or Cav;

X9 is Cys, hCys, Pen, hPen, Abu, or any amino acid or chemical moiety capable of forming a bond with X4;

X10 is Phe[4-(2-aminoethoxy)], Phe(4-CONH<sub>2</sub>) or Phe(4-OR) where R= CH<sub>2</sub>(CH<sub>2</sub>)<sub>n</sub>Y; n=1-25; and Y=NH<sub>2</sub>, CO<sub>2</sub>H, OH, or CH<sub>3</sub>;

X11 is Trp or 2-Nal;

X12 is Aib, 4-amino-4-carboxy-tetrahydropyran,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu, Achc, Acvc, Acbc or Acpc;

X13 is Lys(Ac) or Glu;

X14 is Asn, Gln, Lys(Ac), Lys(N- $\varepsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), or Lys(N- $\varepsilon$ -Palmitoyl);

X15 is Gly,  $\beta$ -Ala, Asn, Gln, Ala, Ser, or Aib;

X16 is absent;

X17 is absent;

X18 is absent;

X19 is absent; and

X20 is absent,

wherein X4 and X9 are capable of forming a bond with each other. In particular embodiments, the bond is a disulfide bond or a thioether bond. In certain embodiments, the peptide inhibitor is cyclized via the bond between X4 and X9.

[00309] In certain embodiments, X1 is a D-amino acid or absent. In certain embodiments, X2 is a D-amino acid or absent.

[00310] In certain embodiments, X16 is a D-amino acid or absent. In certain embodiments, X17 is a D-amino acid or absent. In certain embodiments, X18 is a D-amino acid or absent. In certain embodiments, X19 is a D-amino acid or absent. In certain embodiments, X20 is a D-amino acid or absent.

[00311] In another related embodiment, the present invention includes a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Vd):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Vd)

wherein

X1 is absent;

X2 is absent;

X3 is absent;

X4 is Pen or Abu;

X5 is Gln or Asn;

X6 is Thr or Ser;

X7 is Trp;

X8 is Gln or Asn;

X9 is Pen or Cys;

X10 is Phe[4-(2-aminoethoxy)] or Phe(4-CONH<sub>2</sub>);

X11 is Trp or 2-Nal;

X12 is Aib, 4-amino-4-carboxy-tetrahydropyran,  $\alpha$ -MeLys,  $\alpha$ -MeLeu, or Achc;

X13 is Lys(Ac) or Glu;

X14 is Asn, Gln or Lys(Ac);

X15 is Gly, Ala, Ser,  $\beta$ -Ala, Asn, or Gln;

X16 is absent;

X17 is absent;

X18 is absent;

X19 is absent; and

X20 is absent,

wherein X4 and X9 are capable of forming a bond with each other. In particular embodiments, the bond is a disulfide bond or a thioether bond. In certain embodiments, the peptide inhibitor is cyclized via the bond between X4 and X9.

[00312] Any of the peptide inhibitors of the present invention (e.g., any of those of Formula I (e.g., Ix, Ia-It) may be further defined, e.g., as described below. It is understood that each of the further defining features described herein may be applied to any peptide inhibitors where the amino acids designated at particular positions allow the presence of the further defining feature.

[00313] In certain embodiments, any of the Phe[4-(2-aminoethoxy)] residues present in a peptide inhibitor may be substituted by Phe[4-(2-acetylaminooethoxy)].

[00314] In certain embodiments, X1-X20 are any of the amino acids shown in the corresponding position relative to the cyclized Pen-Pen or cyclized Abu-Cys residues of the illustrative peptide inhibitors set forth in Tables 2-5.

[00315] In certain embodiments, any of the peptides inhibitors described herein, including but not limited to those of Formulas (X), (Va), (Vb), (Vc), (Vd), (Ve), (Vf), (Vg) or (Vh), further comprises a linker or spacer moiety between any two amino acid residues of the peptide. In particular embodiments, the linker or spacer moiety is a PEG moiety.

[00316] In certain embodiments, the peptide inhibitor is cyclized by a disulphide bridge.

[00317] In certain embodiments, X10 is Tyr, Phe[4-(2-aminoethoxy)], Phe(4-CONH<sub>2</sub>) or Phe(4-OMe). In certain embodiments, X10 is Tyr.

[00318] In certain embodiments, X11 is 2-Nal, Trp, or 5-Hydroxy-Trp. In certain embodiments, X11 is Trp.

[00319] In certain embodiments, X10 is Tyr or Phe[4-(2-aminoethoxy)], and X11 is Trp or 2-Nal. In certain embodiments, X10 is Tyr and X11 is Trp.

[00320] In particular embodiments, X4 and X9 are both Cys.

[00321] In particular embodiments, X4 is Cys, Pen, hCys, or absent.

[00322] In particular embodiments, X7 and X11 are not both W.

[00323] In particular embodiments, X7 and X11 are both W.

[00324] In particular embodiments, X7 and X11 are both W, X10 is Y, and X4 and X9 are both Cys.

[00325] In particular embodiments, X15 is Gly, Asn,  $\beta$ -ala or Ser. In particular embodiments, X15 is Gly or Ser.

[00326] In particular embodiments, X16 is AEA or AEP.

[00327] In particular embodiments, X10 is Tyr, Phe or Phe[4-(2-aminoethoxy)]. In particular embodiments, X10 is Tyr or Phe.

[00328] In particular embodiments, X11 is Trp or 2-Nal. In particular embodiments, X11 is Trp.

[00329] In particular embodiments, X1, X2 and X3 are absent.

[00330] In particular embodiments, X18, X19 and X20 are absent.

[00331] In particular embodiments, X1, X2, X3, X18, X19 and X20 are absent.

[00332] In particular embodiments, one or more of X1, X2 or X3 are present.

[00333] In particular embodiments of any of Ix, Ia-Ir, one of X1, X2 and X3 is present and the other two are absent. In one embodiment, the X1, X2 or X3 present is Ala.

[00334] In certain embodiments, X3 is present. In particular embodiments, X3 is Glu, (D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln. In certain embodiments, X3 is (D)Arg or (D)Phe. In particular embodiments, X1 and X2 are absent and X3 is present.

[00335] In particular embodiments, two of X1, X2 and X3 are present and the other one is absent. In certain embodiments, the two present consist of SG, NK, DA, PE, QV or DR.

[00336] In particular embodiments, X1, X2 and X3 are present. In certain embodiments, X1, X2 and X3 consist of ADQ, KEN, VQE, GEE, DGF, NAD, ERN, RVG, KAN, or YED.

[00337] In certain embodiments, the peptide comprises an AEP residue. In particular embodiments, any of X15, X16, X17, X18, X19 or X20 is AEP.

[00338] In certain embodiments of any of the peptide inhibitors or peptide monomer subunits, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys, Lys(Ac),  $\beta$ hAla, or Aib. In certain embodiments of any of the peptide inhibitors or peptide monomer subunits, X13 is Thr, Sarc, Glu, Phe, Arg, Leu, Lys,  $\beta$ hAla, or Aib. In certain embodiments, X14 is Phe, Asn, Tyr, or  $\beta$ hPhe. In certain embodiments, X14 is Phe, Tyr, or  $\beta$ hPhe. In certain embodiments, X15 is Gly, Asn Ser, Thr, Gln, Ala, or Sarc. In certain embodiments, X15 is Gly, Ser, Thr, Gln, Ala, or Sarc. In certain embodiments, X12 is alpha amino acid, e.g., 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, or  $\alpha$ -MeVal.

[00339] In certain embodiments, X13 is present.

[00340] In certain embodiments, X13 and 14 are present.

[00341] In certain embodiments, X13, X14 and X15 are present.

[00342] In particular embodiments of any one of Ia-It, one or more of X16-X20 are present. In particular embodiments, two or more or three or more of X16-X20 are present. In particular embodiments, X18 is [(D)Lys]. In particular embodiments, X17 is absent, and X18 is [(D)Lys]. In certain embodiments wherein X4 and X9 are optionally Cys, X4 and X9 are Cys, X7 is Trp, and X18 is [(D)Lys]. In particular embodiments wherein X4 and X9 are optionally Cys, X4 and X9 are Cys, X7 is Trp, X10 is Tyr or Phe[4-(2-aminoethoxy)], and X18 is [(D)Lys]. In particular embodiments wherein X4 and X9 are optionally Cys, X4 and X9 are Cys, X7 is Trp, X10 is Tyr, and X18 is [(D)Lys]. In particular embodiments wherein X4 and X9 are optionally Cys, X4 and X9 are Cys, X7 is Trp, X1, X2 and X3 are absent, X17 is absent, X18 is [(D)Lys], and X19 and

X20 are absent. In particular embodiments of Ir, X4 and X9 are Cys, X7 and X11 are Trp, X10 is Tyr, and X18 is [(D)Lys. In certain embodiments, X1, X2, and X3 are absent; and in certain embodiments, X17 is absent.

**[00343]** In certain embodiments, any of the peptide inhibitors (or monomer subunits) described herein is cyclized. In particular embodiments, the peptide inhibitor is cyclized via a bond between two or more internal amino acids of the peptide inhibitor. In particular embodiments, cyclized peptide inhibitors are not cyclized via a bond between the N-terminal and C-terminal amino acids of the peptide inhibitor. In certain embodiments, one of the amino acid residues participating in the intramolecular bond cyclizing the peptide in the amino terminal amino acid residue. In certain embodiments, any of the peptide inhibitors in cyclized via a peptide bond between its N-terminal amino acid and its C-terminal amino acid.

**[00344]** In certain embodiments of any of the peptide inhibitors, or one or both monomer subunits thereof, the peptide inhibitor (or one or both monomer subunit thereof) is cyclized via an intramolecular bond between X4 and X9 or by a triazole ring. In particular embodiments, the intramolecular bond is any disulfide bond, a thioether bond, a lactam bond, a triazole, a selenoether bond, a diselendide bond, or an olefin bond.

**[00345]** In one embodiment, X4 and X9 of the peptide inhibitor (or one or both monomer subunits thereof) are Cys, Pen, hCys, D-Pen, D-Cys or D-hCys, and the intramolecular bond is a disulfide bond. In certain embodiments, both X4 and X9 are Cys, or both X4 and X9 are Pen, and the intramolecular bond is a disulfide bond.

**[00346]** In one embodiment, X4 and X9 of the peptide inhibitor (or one or both monomer subunits thereof) are Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu or D-Lys, and the intramolecular bond is a lactam bond.

**[00347]** In one embodiment, X4 is Abu, 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, or 3-chloro-isobutyric acid; X9 is Abu, Cys, Pen, hCys, D-Pen, D-Cys or D-hCys; and the intramolecular bond is a thioether bond. In certain embodiments, X4 is Abu and X9 is Pen, and the intramolecular bond is a thioether bond. In particular embodiments, X4 is a 2-methylbenzoyl

moiety capable of forming a thioether bond with X9, and X9 is selected from Cys, N-Me-Cys, D-Cys, hCys, Pen, and D-Pen. In particular embodiments, X4 is Abu and X9 is Cys, and the intramolecular bond is a thioether bond. In particular instances, a peptide monomer, dimer, or subunit thereof of any of the Formulas and peptides described herein, X4 is selected from the group consisting of modified Ser, modified hSer (e.g., Homo-Ser-Cl), a suitable isostere, and corresponding D-amino acids. In other instances, X4 is an aliphatic acid having from one to four carbons and forming a thioether bond with X9. In some instances, X4 is a five- or six-membered alicyclic acid having a modified 2-methyl group that forms a thioether bond with X9. In some embodiments, X4 is a 2-methylbenzoyl moiety. In certain embodiments, X4 is selected from Cys, hCys, Pen, and a 2-methylbenzoyl moiety. In certain embodiments, X4 is selected from the group consisting of a modified Ser, a modified hSer, a suitable isostere, and corresponding D-amino acids. In one embodiment, X4 is a hSerCl (before the thioether bond is formed with X9 whereby the Cl is removed) or a hSer precursor (e.g., homoSer(O-TBDMS). In other instances, X4 is an aliphatic acid having from one to four carbons and forming a thioether bond with X9. In some instances, X4 is a five- or six-membered alicyclic acid having a modified 2-methyl group that forms a thioether bond with X9. In some instances, X4 is a 2-methylbenzoyl moiety. In certain embodiments wherein X4 is not an amino acid but is a chemical moiety that binds to X9, X1, X2, and X3 are absent, and X4 is conjugated to or bound to X5. In some embodiments, the amino acid directly carboxyl to X9 is an aromatic amino acid. In certain embodiments, X4 is an amino acid, while in other embodiments, X4 is another chemical moiety capable of binding to X9, e.g., to form a thioether bond. In particular embodiments, X4 is another chemical moiety selected from any of the non-amino acid moieties described herein for X4. In particular embodiments wherein X4 is another chemical moiety, X1, X2 and X3 are absent, and the another chemical moiety is bound to or conjugated to X5. In certain embodiments, X4 is defined as a chemical moiety including a group such as a chloride, e.g., in 2-chloromethylbenzoic acid, 2-chloro-acetic acid, 3-chloropropanoic acid, 4-chlorobutyric acid, 3-chloroisobutyric acid. However, the skilled artisan will appreciate that once the peptide has undergone ring closing cyclization to form a thioether bond between X4 and X9, the chloride group is no longer present. The description of chemical moieties at X4 that include a reactant group such as chloride thus means both the group with the chloride and also the group without the chloride, i.e., after formation of the bond with X9. The present invention also includes peptides comprising the same

structure as shown in any of the other formulas or tables described herein, but where the thioether bond is in the reverse orientation. In such embodiments of the invention, it may generally be considered that the amino acid residues or other chemical moieties shown at X4 are instead present at X9, and the amino acid residues shown at X9 are instead present at X4, i.e., the amino acid residue comprising the sulfur of the resulting thioether bond is located at X4 instead of X9, and the amino acid residue or other moiety having a carbon side chain capable of forming a thioether bond with X4 is located at X9. In this reverse orientation, however, the amino acid or chemical moiety at position X9 is one that comprises a free amine. For example, in particular embodiments, the amino acid at X9 is a protected homoserine, such as, e.g., homoserine (OTBDMS). Thus, in particular reverse orientation embodiments of peptide inhibitors of any of the formulas described herein, X9 is an amino acid residue having a side chain with one or two carbons, and forming a thioether bond with X4, and X4 is selected from the group consisting of Cys, N-Me-Cys, D-Cys, HCys, Pen, and D-Pen. Specific examples of amino acid residues and other chemical moieties present at corresponding positions of other formulas and tables are described herein.

[00348] One of skill in the art will appreciate that certain amino acids and other chemical moieties are modified when bound to another molecule. For example, an amino acid side chain may be modified when it forms an intramolecular bridge with another amino acid side chain, e.g., one or more hydrogen may be removed or replaced by the bond. In addition, when hSer-Cl binds to an amino acid such as Cys or Pen via a thioether bond, the Cl moiety is released. Accordingly, as used herein, reference to an amino acid or modified amino acid, such as hSer-Cl, present in a peptide dimer of the present invention (e.g., at position X4 or position X9) is meant to include the form of such amino acid or modified amino acid present in the peptide both before and after forming the intramolecular bond.

[00349] In certain embodiments, the peptide inhibitor of the peptide inhibitor (or one or both monomer subunits thereof) is cyclized through a triazole ring. In certain embodiments, the peptide inhibitor of the peptide inhibitor (or one or both monomer subunits thereof) is linear or not cyclized. In certain embodiments of any of the peptide inhibitors described herein, including both monomer peptide inhibitors and dimer peptide inhibitors, one (or both) peptide monomer subunits comprise or consist of a cyclized peptide having a structure or sequence set forth in any

of Ix, Ia, Ib, Ic, Id, Ie, If, Ig, Ih, Ii, Ij, Ik, Il, Im, In, Io, Ip, Iq, Iq', Ir, Is or It, IIa-IIId, IIIa-IIIe, Iva, or IVb.

[00350] In certain embodiments of any of the peptide inhibitors or monomer subunits, X7 and X11 are both W.

[00351] In certain embodiments of any of the peptide inhibitors or monomer subunits, X7 and X11 are not both W. In particular embodiments, X7 is W and X11 is not W.

[00352] In certain embodiments of any of the peptide inhibitors or monomer subunits, X4 and X9 are amino acid residues capable of forming an intramolecular bond between each other that is a thioether bond, a lactam bond, a triazole, a selenoether, a diselenide bond, or an olefin bond.

[00353] In certain embodiments, X7 and X11 are both W, X10 is Y, Phe[4-(2-aminoethoxy) or Phe(CONH<sub>2</sub>), and X4 and X9 are amino acid residues capable of forming an intramolecular bond between each other that is a thioether bond, a lactam bond, a triazole, a selenoether, a diselenide bond, or an olefin bond. In certain embodiments, X7 and X11 are both W, X10 is Y, and X4 and X9 are amino acid residues capable of forming an intramolecular bond between each other that is a thioether bond, a lactam bond, a triazole, a selenoether, a diselenide bond, or an olefin bond.

[00354] In certain embodiments, X7 and X11 are both W, X10 is Y, and X4 and X9 are both C.

[00355] In certain embodiments, X4 and X9 are each Cys, Pen, hCys, D-Pen, D-Cys or D-hCys, and the intramolecular bond is a disulfide bond.

[00356] In certain embodiments, X4 and X9 are each Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu or D-Lys, and the intramolecular bond is a lactam bond.

[00357] In certain embodiments, X4 and X9 are each  $\beta$ -azido-Ala-OH or propargylglycine, and the peptide inhibitor (or monomer subunit) is cyclized through a triazole ring.

[00358] In certain embodiments, X4 and X9 are each 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, or 2-(5'-hexenyl)glycine and the peptide inhibitor (or monomer subunit) is cyclized via ring closing methasis to give the corresponding olefin / “stapled peptide.”

[00359] In certain embodiments, X4 is 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, or hSer(Cl); X9 is hSer(Cl), Cys, Pen, hCys, D-Pen, D-Cys or D-hCys; and the intramolecular bond is a thioether bond. In certain embodiments, X4 is 2-chloromethylbenzoic acid or hSer(Cl); X9 is Cys or Pen, and the intramolecular bond is a thioether bond. In certain embodiments, X4 is Abu, and X9 is Cys or Pen.

[00360] In certain embodiments, X4 is 2-chloromethylbenzoic acid, 2-chloro-acetic acid, 3-choro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, Abu or Sec; X9 is Abu or Sec; and the intramolecular bond is a selenoether bond.

[00361] In certain embodiments, the intramolecular bond between X4 and X9 is a diselenide bond.

[00362] In certain embodiments of any of the peptide inhibitors described herein that contain two amino acid residues, e.g., cysteine residues, joined by an intramolecular bond, e.g., disulphide bond, the two amino acid residues participating in the intramolecular bond are not both located at either the N-terminal or C-terminal position of the peptide inhibitor. In certain embodiments, neither of the two amino acid residues, e.g., cysteines, participating in the intramolecular bond is located at the N-terminal or C-terminal position of the peptide inhibitor. In other words, in certain embodiments, at least one, or both, of the two amino acid residues, e.g., cysteines, participating in the intramolecular bond are internal amino acid residues of the peptide inhibitor. In certain embodiments, neither of the two amino acid residues, e.g., cysteines, participating in the intramolecular bond is located at the C-terminal position of the peptide inhibitor. At certain embodiment, the two amino acid residues participating in the intramolecular bond are Cys, Pen, hCys, D-Pen, D-Cys or D-hCys residues. In certain embodiments, the two amino acid residues participating in the intramolecular bond are located at X4 and X9. In one embodiment, there is a disulfide bond between the amino acid residues, e.g., cysteines or Pen residues, at X4 and X9. In particular embodiments, both X4 and X9 are Pen. In certain embodiments, one or both peptide monomer subunits in the peptide inhibitor is cyclized via a disulfide bond between two Pen residues, e.g., at positions X4 and X9.

[00363] In particular embodiments of any of the peptide inhibitors described herein, one or both peptide monomer subunits present in the peptide inhibitor, whether it is a monomer or a dimer, is cyclic or cyclized, e.g., by an intramolecular bond, such as a disulfide bond, between two cysteine residues present in the peptide monomer or peptide monomer subunit. In certain embodiments, a peptide inhibitor comprises two or more cysteine residues. In some embodiments, the peptide inhibitor is cyclized via an intramolecular disulfide bond between the two cysteine residues. In particular embodiments of peptide inhibitors having any of the Formulas described herein, the two cysteines occur at positions X4 and X9. In other embodiments, one or both peptide monomer subunits in the peptide inhibitor is cyclized via a disulfide bond between two Pen residues, e.g., at positions X4 and X9.

[00364] In some embodiments, a peptide inhibitor has a structure of any of the Formulas described herein (e.g., Formula I) and comprises a disulfide bond, e.g., an intramolecular disulfide bond, or a thioether bond. Illustrative examples of such peptide inhibitors are shown in Tables 3A-3H and 4A, 4B, 9, 11 or 15. Such disulfide bonded peptides may have a particular advantage in that the disulfide bonds enhance structural stability and can improve biological activity of many bioactive peptides. However, in certain situations, these bonds are labile to reducing agents. One of skill in the art will appreciate that disulfide is amenable to simple isosteric replacement. Illustrative examples of such replacements include, but are not limited to, thioethers, dithioethers, selenoethers, diselenides, triazoles, lacatams, alkane and alkene groups. Accordingly, in certain embodiments of any of the peptide inhibitors described herein, one, two or more cysteine residues are substituted, e.g., with a thioether, dithioether, selenoether, diselenide, triazoles, lacatam, alkane or alkene group, including but not limited to any of those shown below or described herein. In particular embodiments, two of these substituted groups form a bond (e.g., an intramolecular bond), thus cyclizing the peptide inhibitor or one or both monomer subunits thereof.

[00365] In certain embodiments, a peptide inhibitor of the present invention comprises or consists of an amino acid sequence shown herein, e.g., in any one of Tables 3A-3H, 4A, 4B, 5A-5C, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15. In certain embodiments, a peptide inhibitor of the present invention has a structure shown herein, e.g., in any one of Tables 3A-3H, 4A, 4B, 5A-5C, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15.

[00366] In certain embodiments, the present invention includes a peptide inhibitor that comprises a core consensus sequence selected from one of the following (shown in N-terminal to C-terminal direction):

$X_1X_2X_2WX_2X_1X_2W$ ;

$X_1X_2X_2WX_2X_1X_2$  (1-Nal);

$X_1X_2X_2WX_2X_1X_2$  (2-Nal);

$X_1X_2X_2WX_2X_1YW$ ;

$X_1X_2X_2WX_2X_1Y$ (1-Nal);

$X_1X_2X_2WX_2X_1Y$ (2-Nal);

$X_1X_2X_2WX_2X_1X_2X_2$ ;

$X_1X_2X_2WX_2X_1X_2X_2X_2X_2X_2$ -[(D)Lys];

$X_1X_2X_2WX_2X_1X_3X_2$ ;

$X_1X_2X_2WX_2X_1X_3$ (1-Nal); and

$X_1X_2X_2WX_2X_1X_3$ (2-Nal).

[00367] wherein W is tryptophan, Y is tyrosine, each the two X1 residues are amino acids or other chemical moieties capable of forming an intramolecular bond with each other; each X2 is independently selected from all amino acids, which include, e.g., natural amino acids, L-amino acids, D-amino acids, non-natural amino acids, and unnatural amino acids; and X3 is any amino acid. In particular embodiments, X3 is Phe, a Phe analog (e.g., Phe[4-(2-aminoethoxy)] or Phe(4-CONH<sub>2</sub>)), Tyr, or a Tyr analog (e.g., Tyr(Me)). In particular embodiments, each X1 is selected from Cys, Pen and Abu. In particular embodiments, each X1 is Cys. In certain embodiments, each X1 is Pen. In certain embodiments, one X1 is Cys and the other X1 is Abu. In particular embodiments, the N-terminal X1 is Abu and the C-terminal X1 is Cys. In particular embodiments, the N-terminal X1 is Cys and the C-terminal X1 is Abu. In particular embodiments, the residues between the two X1 residues are Gln, Thr, Trp and Gln. In particular embodiments, each X1 is selected from Cys, Pen and Abu; and X3 is Phe, a Phe analog (e.g., Phe[4-(2-aminoethoxy)] or Phe(4-carbomide)), Tyr, or a Tyr analog (e.g., Tyr(Me)). In particular embodiments, X3 is a Phe analog.

[00368] In certain embodiments, peptide inhibitors of the present invention comprises any of the following consensus sequences, wherein X1, X2, X3, X4, X5, X6, X7, X8, X9, X10, X11, X12, X13, X14 and X15 are defined as shown in any of the various Formula or peptide inhibitors described herein:

X1-X2-X3-Pen-X5-X6-W-X8-Pen-X10-X11-X12-X13-X14-X15;

Pen-X5-X6-W-Q-Pen;

Pen-X5-X6-W-X8-Pen;

Pen-X5-X6-W-X8-Pen-[Phe(4-CONH2)];

Pen-X5-X6-W-X8-Pen-[Phe[4-(2-aminoethoxy)]];

X1-X2-X3-Abu-X5-X6-W-X8-C-X9-X10-X11-X12-X13-X14-X15;

Abu-X5-X6-W-Q-C;

Abu-X5-X6-W-X8-C;

Abu-X5-X6-W-X8-C-[Phe(4-CONH2)]; or

Abu-X5-X6-W-X8-C-[Phe[4-(2-aminoethoxy)]].

[00369] In certain embodiments of any of the peptide inhibitors or monomer subunits, X7 and X11 are both W. In certain embodiments of any of the peptide inhibitors, X7 and X11 are both W, and X10 is Y. In certain embodiments, X7 and X11 are both W and X10 is Phe[4-(2-aminoethoxy)] or Phe(4-OMe) .

[00370] In certain embodiments of any of the peptide inhibitors or monomer subunits, X7 and X11 are not both W.

[00371] In certain embodiments of peptide inhibitors of Formula I, X4 and X9 are each Pen, and the intramolecular bond is a disulfide bond.

[00372] In certain embodiments, a peptide inhibitor of the present invention comprises or consists of an amino acid sequence shown in any one of the tables, sequence listing or the accompanying figures herein.

[00373] In certain embodiments of any of the peptide inhibitors described herein that contain two amino acid residues, e.g., Pen residues, joined by an intramolecular bond, e.g., disulphide bond, one or both of the two amino acid residues participating in the intramolecular bond are not located at either the N-terminal or C-terminal position of the peptide inhibitor. In certain embodiments, neither of the two amino acid residues, e.g., Pen, participating in the intramolecular bond is located at the N-terminal or C-terminal position of the peptide inhibitor. In other words, in certain embodiments, at least one, or both, of the two amino acid residues, e.g., Pens, participating in the intramolecular bond are internal amino acid residues of the peptide inhibitor. In certain embodiments, neither of the two amino acid residues, e.g., Pens, participating in the intramolecular bond is located at the C-terminal position of the peptide inhibitor.

[00374] In some embodiments, wherein a peptide of the invention is conjugated to an acidic compound such as, e.g., isovaleric acid, isobutyric acid, valeric acid, and the like, the presence of such a conjugation is referenced in the acid form. So, for example, but not to be limited in any way, instead of indicating a conjugation of isovaleric acid to a peptide by referencing isovaleroyl (e.g., isovaleroyl-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[a-MeLys]-[Lys(Ac)]-NG-NH<sub>2</sub>) in some embodiments, the present application references such a conjugation as isovaleric acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[a-MeLys]-[Lys(Ac)]-NG-NH<sub>2</sub>.

[00375] The present invention further includes peptide inhibitors that selectively bind to an epitope or binding domain present within amino acid residues 230 – 349 of the human IL23R protein. In particular embodiments, the peptide inhibitor binds human IL23R and not mouse IL-23R. In certain embodiments, the peptide inhibitor also binds to rat IL-23R.

[00376] In certain embodiments of peptide inhibitors of Formula I, X4 is Abu; X9 is Cys, Pen, homocys, and the intramolecular bond is a thioether bond.

[00377] In certain embodiments, peptide inhibitors do not include compounds, disclosed in PCT Application No. PCT/US2014/030352 or PCT Application No. PCT/US2015/038370.

Illustrative Peptide Inhibitors Comprising Pen-Pen Disulfide Bonds

[00378] In certain embodiments, the present invention includes a peptide inhibitor of an interleukin-23 receptor, wherein the peptide inhibitor has the structure of Formula II:



[00379] or a pharmaceutically acceptable salt or solvate thereof,

wherein  $R^1$  is a bond, hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl, a C1-C6 alkyl, a C1-C20 alkanoyl, an alkylsulphonate, an acid,  $\gamma$ -Glu or pGlu, appended to the N-terminus, and including PEGylated versions (e.g., 200 Da to 60,000 Da), alone or as a spacer of any of the foregoing;

[00380]  $R^2$  is a bond, OH or  $NH_2$ ; and

[00381] X is an amino acid sequence of 8 to 20 amino acids or 8 to 35 amino acids.

[00382] In particular embodiments of peptide inhibitor of Formula II, X comprises or consists of the sequence of Formula IIa:



wherein

X1 is absent or any amino acid;

X2 is absent or any amino acid;

X3 is absent or any amino acid;

X4 is Pen, Cys or homo-Cys;

X5 is any amino acid;

X6 is any amino acid;

X7 is Trp, Bip, Gln, His, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -Me-Trp, 1,2,3,4 - tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>), Phe(4-tBu),  $\beta\beta$ -diPheAla, Glu, Gly, Ile, Asn, Pro, Arg, Thr or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X8 is any amino acid;

X9 is Pen, Cys or hCys;

X10 is 1-Nal, 2-Nal, Aic, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, Phe, His, Trp, Thr, Tic, Tyr, 4-pyridylAla, Octgly, a Phe analog or a Tyr analog (optionally, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me), Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), or Phe(4-OBzl)), or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe(3,4-F<sub>2</sub>), Phe(4-CO<sub>2</sub>H),  $\beta$ hPhe(4-F),  $\alpha$ -Me-Trp, 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tri-tert-Butyl), Phe(4-Oallyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino, Phe(4-OBzl), Octgly, Glu(Bzl), 4-Phenylbenzylalanine, Phe[4-(2-aminoethoxy)], 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, 1,2,3,4-tetrahydro-norharman, Phe(4-CONH<sub>2</sub>), Phe(3,4-OMe<sub>2</sub>) Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine or Bip, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu,  $\alpha$ -MeVal, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Lys, Arg, Orn, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Leu, Asn, Ogl, Pro, Gln, Asp, Arg, Ser, spiro-pip, Thr, Tba, Tlc, Val or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr, Lys(Ac), Orn or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser,  $\beta$ -Ala, Arg or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X16 is absent, Gly, Ala, Asp, Ser, Pro, Asn or Thr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X17 is absent, Glu, Ser, Gly or Gln, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X18 is absent or any amino acid;

X19 is absent or any amino acid; and

X20 is absent or any amino acid.

[00383] In certain embodiments of IIa: X10 is 1-Nal, 2-Nal, Aic, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, Phe, His, Trp, Thr, Tic, Tyr, 4-pyridylAla, Octgly, a Phe analog or a Tyr analog, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe (3,4-F<sub>2</sub>), Phe(4-CO<sub>2</sub>H),  $\beta$ hPhe(4-F),  $\alpha$ -Me-Trp, 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tri-tert-Butyl), Phe(4-Oallyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl), Octgly, Glu(Bzl), 4-Phenylbenzylalanine, Phe[4-(2-aminoethoxy)], 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp, 1,2,3,4-tetrahydro-norharman, Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine or Bip, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu, MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Lys, Arg, Orn, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Leu, Asn, Ogl, Pro, Gln, Asp, Arg, Ser, spiro-pip, Thr, Tba, Tlc, Val or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; and X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing.

[00384] In certain embodiments, X3 is present. In particular embodiments, X3 is Glu,(D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln. In certain embodiments, X3 is (D)Arg or (D)Phe. In particular embodiments, X1 and X2 are absent and X3 is present.

[00385] In certain embodiments, X5 is Gln, Ala, Cit, Asp, Dab, Dap, Cit Glu, Phe, Gly, His, hCys, Lys, Leu, Met, Asn, N-Me-Ala, N-Me-Asn, N-Me-Lys,  $\alpha$ -Me-Lys,  $\alpha$ -Me-Orn, N-Me-Gln, N-Me-Arg,  $\alpha$ -MeSer, Orn, Pro, Arg, Ser, Thr, or Val. In certain embodiments, X5 is Gln, Ala, Cit, Asp, Dab, Dap, Glu, Phe, Gly, His, hCys, Lys, Leu, Met, Asn, N-Me-Ala, N-Me-Asn, N-Me-Lys,  $\alpha$ Me-Lys,  $\alpha$ Me-Orn, N-Me-Gln, N-Me-Arg, Orn, Pro, Arg, Ser, Thr, or Val. In certain embodiments, X5 is Gln or Asn.

[00386] In certain embodiments, X6 is Thr, Asp, Glu, Phe, Asn, Pro, Arg, or Ser.

[00387] In certain embodiments, X7 is Trp.

[00388] In certain embodiments, X8 is Gln, Glu, Phe, Lys, Asn, Pro, Arg, Val, Thr, or Trp.

[00389] In certain embodiments, X10 is a Tyr analog or a Phe analog. In particular embodiments, X10 is a Phe analog.

[00390] In certain embodiments wherein X10 is a Phe analog, X10 is selected from hPhe, Phe(4-OMe),  $\alpha$ -Me-Phe, hPhe(3,4-dimethoxy), Phe(4-CONH<sub>2</sub>), Phe(4-phenoxy), Phe(4-guanadino), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-OBzl), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5 DiF), Phe(CH<sub>2</sub>CO<sub>2</sub>H), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>),  $\beta\beta$ -diPheAla, Phe(4-N<sub>3</sub>) and Phe[4-(2-aminoethoxy)]. In particular embodiments, X10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]. In particular embodiments, X10 is Phe(4-OMe), Phe(4-CONH<sub>2</sub>) or Phe[4-(2-aminoethoxy)]. In certain embodiments where X10 wherein X10 is a Phe analog, X10 is selected from hPhe, Phe(4-OMe),  $\alpha$ -Me-Phe, hPhe(3,4-dimethoxy), Phe(4-CONH<sub>2</sub>), Phe(4-phenoxy), Phe(4-guanadino), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-OBzl), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5 DiF), Phe(CH<sub>2</sub>CO<sub>2</sub>H), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>),  $\beta\beta$ -diPheAla, Phe(4-N<sub>3</sub>) and Phe[4-(2-aminoethoxy)]. In particular embodiments, X10 is Phe(4-OMe).

[00391] In certain embodiments where X10 is a Tyr analog, X10 is selected from hTyr,  $\alpha$ -MeTyr, N-Me-Tyr, Tyr(3-tBu), Phe(4-CONH<sub>2</sub>), Phe[4-(2-aminoethoxy)], and bhTyr. In certain

embodiments where X10 is a Tyr analog, X10 is selected from hTyr,  $\alpha$ -MeTyr, N-Me-Tyr, Tyr(3-tBu), and bhTyr.

[00392] In certain embodiments, X10 is Tyr, Phe(4-OMe), Phe[4-(2-aminoethoxy)], Phe(4-CONH<sub>2</sub>), or 2-Nal. In certain embodiments, X10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]. In certain embodiments, X10 is not Tyr.

In certain embodiments, X11 is a Trp analog. In particular embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X11 is 2-Nal.

[00393] In certain embodiments, X12 is Aib,  $\alpha$ -MeLys or  $\alpha$ -MeLeu.

[00394] In particular embodiments of a peptide inhibitor of Formula II, one or both of X4 or X9 is Pen. In particular embodiments, both X4 and X9 are Pen.

[00395] In certain embodiments, the peptide inhibitor of Formula II is cyclized. In particular embodiments, the peptide inhibitor of Formula II is cyclized via an intramolecular bond between X4 and X9. In particular embodiments, the intramolecular bond is a disulfide bond. In particular embodiments, X4 and X9 are both Pen.

[00396] In certain embodiments, the peptide inhibitor of Formula II is linear or not cyclized. In particular embodiments of the linear peptide inhibitor of Formula I, X4 and/or X9 are any amino acid.

[00397] In particular embodiments of a peptide inhibitor of Formula II, one or more, two or more, or all three of X1, X2, and X3 are absent. In certain embodiments, X1 is absent. In certain embodiments, X1 and X2 are absent. In certain embodiments, X1, X2 and X3 are absent.

[00398] In particular embodiments of a peptide inhibitor of Formula II, one or more, two or more, three or more, four or more, or all of X16, X17, X18, X19 and X20 are absent. In particular embodiments of a peptide inhibitor of Formula I, one or more, two or more, three or more, or all of X17, X18, X19 and X20 are absent. In certain embodiments, one or more, two or more, or all three of X17, X19 and X20 are absent. In certain embodiments, one or more of X1, X2 and X3

are absent; and one or more, two or more, three or more, or four of X17, X18, X19 and X20 are absent.

**[00399]** In particular embodiments of a peptide inhibitor of Formula II, X18 is (D)-Lys. In certain embodiments, X18 is (D)-Lys and X17 is absent.

**[00400]** In particular embodiments of a peptide inhibitor of Formula II, the peptide inhibitor comprises one or more, two or more, three or more, or four of the following features: X5 is Asn, Arg or Gln; X6 is Thr; X7 is Trp; and X8 is Gln. In particular embodiments of a peptide inhibitor of Formula I, X4 is Pen; X5 is Gln, Asn or Arg; X6 is Thr; X7 is Trp, 5-hydroxy-Trp, 6-chloro-Trp, N-MeTrp, alpha-Me-Trp, or 1,2,3,4-tetrahydro-norharman; X8 is Gln; and X9 is Pen. In particular embodiments, X5 is Gln. In certain embodiments, X1, X2 and X3 are absent. In particular embodiments, both X4 and X9 are Pen.

**[00401]** In particular embodiments of a peptide inhibitor of Formula II, the peptide inhibitor comprises one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following features: X10 is Tyr, a Phe analog, a Tyr analog or 2-Nal; X11 is Trp, 5-hydroxy-Trp, 6-chloro-Trp, N-MeTrp, alpha-Me-Trp, 1,2,3,4-tetrahydro-norharman, 2-Nal or 1-Nal; X12 is Aib,  $\alpha$ -MeLys,  $\alpha$ -MeOrn and  $\alpha$ -MeLeu; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly, Ser or Ala; and X16 is absent or AEA. In certain embodiments, X10 is Tyr, Phe(4-OMe), Phe[4-(2-aminoethoxy)], Phe(CONH<sub>2</sub>), or 2-Nal. In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X10 is not Tyr. In certain embodiments, X1, X2 and X3 are absent. In particular embodiments, both X4 and X9 are Pen.

**[00402]** In particular embodiments of a peptide inhibitor of Formula II, the peptide inhibitor comprises one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven of the following features: X5 is Arg or Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is a Phe analog; X11 is Trp, 2-Nal or 1-Nal; X12 is Aib,  $\alpha$ -MeLys or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Asn; X15 is Gly, Ser or Ala; and X16 is absent or AEA. In certain embodiments, X10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]. In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent. In particular embodiments, both X4 and X9 are Pen.


[00403] In particular embodiments of a peptide inhibitor of Formula II, the peptide is cyclized via X4 and X9; X4 and X9 are Pen; X5 is Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr, a Phe analog or 2-Nal; X11 is Trp, 2-Nal or 1-Nal; X12 is Arg,  $\alpha$ -MeLys,  $\alpha$ -MeOrn, or  $\alpha$ -MeLeu; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly, Ser or Ala; and X16 is absent. In certain embodiments, X10 is Tyr, Phe(4-OMe), Phe[4-(2-aminoethoxy)], Phe(4-OMe) or 2-Nal. In certain embodiments, X10 is Phe(4-OMe). In certain embodiments, X10 is not Tyr. In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent.

[00404] In particular embodiments of a peptide inhibitor of Formula II, the peptide is cyclized via X4 and X9; X4 and X9 are Pen; X5 is Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr, Phe(4-OMe) or 2-Nal; X11 is Trp, 2-Nal or 1-Nal; X12 is Arg,  $\alpha$ -MeLys or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly; and X16 is absent. In certain embodiments, X10 is Phe(4-OMe). In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent.

[00405] In particular embodiments of a peptide inhibitor of Formula II, the peptide is cyclized via X4 and X9; X4 and X9 are Pen; X5 is Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]; X11 is Trp, 2-Nal or 1-Nal; X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn, or  $\alpha$ -MeLeu; X13 is Lys, Glu or Lys(Ac); X14 is Asn; X15 is Gly, Ser or Ala; and X16 is absent. In certain embodiments, X10 is Phe(4-OMe). In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent.

[00406] In particular embodiments of a peptide inhibitor of Formula II, X10 is not Tyr.

[00407] In certain embodiments, the present invention includes a peptide, optionally 8 to 35, 8 to 20, 8 to 16 or 8 to 12 amino acids in length, optionally cyclized, comprising or consisting of having a core sequence of Formula IIb:



[00408] wherein Xaa5, Xaa6 and Xaa8 are any amino acid residue; and Xaa10 is a Phe analogue, wherein the peptide inhibits binding of IL-23 to IL-23R. In particular embodiments, X10 is a Phe analog selected from  $\alpha$ -Me-Phe, Phe(4-OMe), Phe(4-OBzl), Phe(4-OMe), Phe(4-CONH<sub>2</sub>),

Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-CO<sub>2</sub>H), Phe[4-(2-aminoethoxy)] or Phe(4-guanadino). In particular embodiments, Xaa10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]. In one embodiment, Xaa10 is Phe(4-OMe). In certain embodiments, the peptide is cyclized via an intramolecular bond between Pen at Xaa4 and Pen at Xaa9. In particular embodiments, the peptide is a peptide inhibitor of Formula II, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain embodiments, a peptide of Formula IIb further comprises an amino acid bound to the N-terminal Pen residue. In particular embodiments, the bound amino acid is Glu,(D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln. In certain embodiments, it is is (D)Arg or (D)Phe.

[00409] In certain embodiments, the present invention includes a peptide, optionally 8 to 35, 8 to 20, 8 to 16, or 8 to 12 amino acids in length, optionally cyclized, comprising or consisting of a core sequence of Formula IIc:

Pen-Xaa5-Xaa6-Trp-Xaa8-Pen-Xaa10-[(2-Nal)] (IIc)

[00410] wherein Xaa5, Xaa6 and Xaa8 are any amino acid residue; and Xaa10 is Tyr, a Phe analog,  $\alpha$ -Me-Tyr,  $\alpha$ -Me-Trp or 2-Nal, wherein the peptide inhibits binding of IL-23 to IL-23R. In certain embodiments, X10 is Tyr, Phe(4-OMe), Phe[4-(2-aminoethoxy)],  $\alpha$ -Me-Tyr,  $\alpha$ -Me-Phe,  $\alpha$ -Me-Trp or 2-Nal. In certain embodiments, Xaa10 is Tyr, Phe(4-OMe), Phe(CONH<sub>2</sub>), Phe[4-(2-aminoethoxy)] or 2-Nal. In certain embodiments, Xaa10 is Tyr, Phe(4-OMe), Phe[4-(2-aminoethoxy)] or 2-Nal. In particular embodiments, Xaa10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]. In one embodiment, Xaa10 is Phe[4-(2-aminoethoxy)] or Phe(CONH<sub>2</sub>). In particular embodiments, Xaa10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)]. In one embodiment, Xaa10 is Phe[4-(2-aminoethoxy)]. In certain embodiments, Xaa10 is not Tyr. In certain embodiments, the peptide is cyclized via an intramolecular bond between Pen at Xaa4 and Pen at Xaa9. In particular embodiments, the peptide is a peptide inhibitor of Formula II, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain embodiments, a peptide of Formula IIc further comprises an amino acid bound to the N-terminal Pen residue. In particular embodiments, the

bound amino acid is Glu,(D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln. In certain embodiments, it is is (D)Arg or (D)Phe.

[00411] In certain embodiments, the present invention includes a peptide, optionally 8 to 35, 8 to 20, 8 to 16 or 8 to 12 amino acids in length, optionally cyclized, comprising or consisting of a core sequence of Formula IIId:



[00412] wherein Xaa5, Xaa6 and Xaa8 are any amino acid residue. In certain embodiments, the peptide comprises a disulfide bond between Xaa4 and Xaa9. In certain embodiments, the peptide is a peptide inhibitor of Formula I, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain embodiments, a peptide of Formula IIId further comprises an amino acid bound to the N-terminal Pen residue. In particular embodiments, the bound amino acid is Glu,(D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln. In certain embodiments, it is is (D)Arg or (D)Phe.

[00413] In particular embodiments of a peptide inhibitor of Formula II, the peptide inhibitor has a structure shown in any of Tables 2, 3, 4A, 4B, 8, 11 or 15 or comprises an amino acid sequence set forth in Tables 2, 3, 4A, 4B, 8, 11 or 15.

Table 2. Illustrative Di-Pen Inhibitors

[Palm]-[isoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-isoGlu-Palm)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Octanyl]-[IsoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Octanyl]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Palm]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Octanyl)]-NN-NH <sub>2</sub>

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Palm)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Palm)-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Lauryl)]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(PEG4-IsoGlu-Palm)]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-a-Me-K(PEG4-IsoGlu-Lauryl)]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(IVA)]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(Biotin)]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(Octanyl)]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-[Lys(IVA)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(IVA)]-N-NH <sub>2</sub>
Ac-[Pen]-[Lys(Biotin)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Biotin)]-N-NH <sub>2</sub>
Ac-[Pen]-[Lys(Octanyl)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(octanyl)]-N-NH <sub>2</sub>
Ac-[Pen]-[Lys(Palm)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Lys(Palm)]-N-NH <sub>2</sub>
Ac-[Pen]-[Lys(PEG8)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(PEG8)]-N-NH <sub>2</sub>
Ac-[Pen]-K(Peg11-Palm)TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Peg11-palm)]-N-NH <sub>2</sub>
Ac-[Pen]-[Cit]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-[Lys(Ac)]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NT-[Phe(3,4-OCH <sub>3</sub> ) <sub>2</sub> ]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NT-[Phe(2,4-CH <sub>3</sub> ) <sub>2</sub> ]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NT-[Phe(3-CH <sub>3</sub> )]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NT-[Phe(4-CH <sub>3</sub> )]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>
Ac[(D)Tyr]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-N-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Lys(Ac)]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QQ-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Q-[βAla]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Cit]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-NNH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-Q-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-[Lys(Ac)]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-[Cit]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QN-[βAla]-NH <sub>2</sub>

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-E-[Cit]-Q-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-CitNCitNH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Cit]-Q-[Cit]-NH <sub>2</sub>
Ac-[Pen]-[Cit]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QNN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-ENQ-NH <sub>2</sub>
Ac-[Pen]-GPWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-PGWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWN-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NSWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-N-[Aib]-WQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTW-[Aib]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]N-[Aib]-NH <sub>2</sub>
Ac-[Pen]-QTW-[Lys(Ac)]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-[Lys(Ac)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]NNNH <sub>2</sub>
Ac-[Pen]-QVWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NT-[2-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NT-[1-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-LN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-GN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-SN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Aib]-N-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-FN-NH <sub>2</sub>
Ac-[Pen]-NTW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Tic]-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[nLeu]-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-G-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-R-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-W-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-S-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-L-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[AIB]-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[N-MeAla]-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[2-Nap]-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-F-[ $\beta$ Ala]-NH <sub>2</sub>
Ac-[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>

Table 3. Illustrative Peptides Containing the Ac-[Pen]-XXWX-[Pen]-XXXX Motif and Analogues Thereof

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNE-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNF-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNK-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNW-NH
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNG-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNPK-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNPG-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNEP-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNKG-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNPT-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNKG-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNGW-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNGQ-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNGGG-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNKKK-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNEEE-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNFFF-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNTTT-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NNGGGR-NH <sub>2</sub>



Ac-EG-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-QG-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-TG-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Palm)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(isoGlu-Palm)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(PEG11-Palm)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ahx-Palm)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(isoGlu-Ahx-Palm)]-NN-NH <sub>2</sub>
[Palm]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Palm-isoGlu]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Palm-PEG11]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Palm-Ahx]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Palm-Ahx-isoGlu]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-Lys[Palm]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-Lys[isoGlu-Palm]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-Lys[PEG11-Palm]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-Lys[Ahx-Palm]-NH <sub>2</sub>
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-Lys[isoGlu-Ahx-Palm]-NH <sub>2</sub>

#### Illustrative Peptide Inhibitors Comprising Thioether Bonds

[00414] In certain embodiments, the present invention includes a peptide inhibitor of an interleukin-23 receptor, wherein the peptide inhibitor has the structure of Formula III:

$$R^1-X-R^2 \text{ (III)}$$

[00415] or a pharmaceutically acceptable salt or solvate thereof,

[00416] wherein  $R^1$  is a bond, hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl, a C1-C6 alkyl, a C1-C20 alkanoyl, an alkylsulphonate, an acid,  $\gamma$ -Glu or pGlu, appended to the N-terminus, and including PEGylated versions (e.g., 200 Da to 60,000 Da), alone or as a spacer of any of the foregoing,

[00417]  $R^2$  is a bond, OH or  $NH_2$ ; and

[00418] X is an amino acid sequence of 8 to 20 amino acids or 8 to 35 amino acids,

[00419] In particular embodiments of peptide inhibitors of Formula III, X comprises or consists of the sequence of Formula IIIa:

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (IIIa)

wherein

X1 is absent or any amino acid;

X2 is absent or any amino acid;

X3 is absent or any amino acid;

X4 is Abu, Pen, or Cys;

X5 is any amino acid;

X6 is any amino acid;

X7 is Trp, Bip, Gln, His, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-(OCH<sub>3</sub>)<sub>2</sub>), Phe(4-CF<sub>3</sub>),  $\beta\beta$ -diPheAla, Phe(4-tBu), Glu, Gly, Ile, Asn, Pro, Arg, Thr or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X8 is any amino acid;

X9 is Abu, Pen, or Cys;

X10 is 1-Nal, 2-Nal, Aic, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, Phe, His, Trp, Thr, Tic, Tyr, 4-pyridylAla, Octgly a Phe analog or a Tyr analog (optionally, Phe(3,4-F<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), F(3-Me),

Phe[4-(2-aminoethoxy)], Phe[4-(2-(acetyl-aminoethoxy)], Phe(4-Br), Phe(4-CONH<sub>2</sub>), Phe(4-Cl), Phe(4-CN), Phe(4-guanidino), Phe(4-Me), Phe(4-NH<sub>2</sub>), Phe(4-N<sub>3</sub>), Phe(4-OMe), Phe(4-OBzl)), or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, 4-phenylcyclohexyl, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(3,4-F<sub>2</sub>),  $\beta$ hPhe(4-F), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>), Phe(2,3-Cl<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine,  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Bip, Nva(5-phenyl), Phe, His, hPhe, Tqa, Trp, Tyr, Phe(4-Me), Trp(2,5,7-tri-tertButyl), Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl), or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu, MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle, amino-4-carboxy-tetrahydropyran (THP), Achc Acpc, Acbc, Acvc, Aib, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X13 is Lys Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu,  $\beta$ Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Arg, Orn, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Leu, Asn, Ogl, Pro, Gln, Asp, Arg, Ser, spiro-pip, Thr, Tba, Tlc, Val or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X14 is Asn, Glu, Phe, Gly, His, Lys, Lys (Ac), Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic, Asp or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg,  $\beta$ -Ala, or Ser, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X16 is absent, Gly, Ala, Asp, Ser, Pro, Asn or Thr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X17 is absent, Glu, Ser, Gly or Gln, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing;

X18 is absent or any amino acid;

X19 is absent or any amino acid; and

X20 is absent or any amino acid.

[00420] In certain embodiments, X14 is Asn, Glu, Phe, Gly, His, Lys, Lys (Ac), Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic, or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing

[00421] In certain embodiments of IIIa: X7 is Trp, Bip, Gln, His, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>),  $\beta\beta$ -diPheAla, Phe(4-tBu), Glu, Gly, Ile, Asn, Pro, Arg, Thr or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X10 is 1-Nal, 2-Nal, Aic, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, Phe, His, Trp, Thr, Tic, Tyr, 4-pyridylAla, Octgly a Phe analog or a Tyr analog, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, 4-phenylcyclohexyl, Glu(Bzl), 4-Phenylbenzylalanine, Tic, Phe[4-(2-aminoethoxy)], Phe(3,4-Cl<sub>2</sub>), Phe(3,4-F<sub>2</sub>),  $\beta$ hPhe(4-F), Phe(4-OMe), 5-Hydroxy-Trp, 6-Chloro-Trp, N-MeTrp,  $\alpha$ -MeTrp, 1,2,3,4 -tetrahydro-norharman, Phe(4-CO<sub>2</sub>H), Phe(4-CONH<sub>2</sub>), Phe(3,4-Dimethoxy), Phe(4-CF<sub>3</sub>), Phe(2,3-Cl<sub>2</sub>), Phe(2,3-F<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine,  $\alpha$ -MePhe,  $\beta$ hNal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, Bip, Nva(5-phenyl), Phe, His, hPhe, Tqa, Trp, Tyr, Phe(4-Me), Trp(2,5,7-tri-tertButyl), Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl), or Octgly, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu, MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu,  $\beta$ Ala,  $\beta$ hGlu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Arg, Orn, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Leu, Asn, Ogl, Pro, Gln, Asp, Arg, Ser, spiro-pip, Thr, Tba, Tlc, Val or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing; X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr, or a corresponding  $\alpha$ -methyl amino acid form of any of the

foregoing; and X15 is Gly, (D)Ala, (D)Asn, (D)Asp, Asn, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser, or a corresponding  $\alpha$ -methyl amino acid form of any of the foregoing.

[00422] In certain embodiments, X3 is present. In particular embodiments, X3 is Glu,(D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln. In certain embodiments, it is (D)Arg or (D)Phe.

[00423] In particular embodiments, X5 is Gln, Ala, Cys, Cit, Asp, Dab, Dap, Glu, Phe, Gly, His, hCys, Lys, Leu, Met, Asn, N-Me-Ala, N-M-Asn, N-Me-Lys, N-Me-Gln, N-Me-Arg, Orn, Pro, Pen, Gln, Arg, Ser, Thr, or Val.

[00424] In particular emodiments, X6 is Thr, Asp, Glu, Phe, Asn, Pro, Arg, Ser, or Thr.

[00425] In particular embodiments, X8 is Gln, Glu, Phe, Lys, Asn, Pro, Arg, Val, Thr, or Trp.

[00426] In certain embodiments, X10 is a Tyr analog or a Phe analog. In particular embodiments, X10 is Phe(4-OMe), Phe(CONH<sub>2</sub>) or Phe[4-(2-aminoethoxy)]. In certain embodiments, X10 is a Tyr analog or a Phe analog. In particular embodiments, X10 is Phe(4-OMe) or Phe[4-(2-aminoethoxy)].

[00427] In certain embodiments where X10 is a the Phe analog, X10 is selected from hPhe, Phe(4-OMe),  $\alpha$ -MePhe, hPhe(3,4-dimethoxy), Phe(4-CONH<sub>2</sub>), Phe(4-O-Bzl), Phe(4-guanadino), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5 DiF), Phe(CH<sub>2</sub>CO<sub>2</sub>H), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>),  $\beta\beta$ -diPheAla, Phe(4-N<sub>3</sub>) and Phe[4-(2-aminoethoxy)]. In particular embodiments, X10 is Phe[4-(2-aminoethoxy)] or Phe(CONH<sub>2</sub>). In particular embodiments, X10 is Phe[4-(2-aminoethoxy)].

[00428] In certain embodiments where X10 is a Tyr analog, X10 is selected from hTyr, N-Me-Tyr, Tyr(3-tBu), Phe(4-OMe) and bhTyr. In particular embodiments, X10 is Phe(4-OMe).

[00429] In particular embodiments, X10 is Tyr, Phe(4-OMe), Phe(4-OBzl), Phe(4-OMe), Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-carboxy), Phe[4-(2aminoethoxy)] or Phe(4-guanadino). In particular embodiments, X10 is not Tyr.

[00430] In certain embodiments, X11 is Trp or a Trp analog. In particular embodiments, X11 is 2-Nal or 1-Nal.

[00431] In particular embodiments, the peptide inhibitor of Formula III is cyclized. In certain embodiments, the peptide inhibitor is cyclized via an intramolecular bond between X4 and X9. In certain embodiments, the intramolecular bond is a thioether bond.

[00432] In certain embodiments, the peptide inhibitor of Formula III is linear or not cyclized. In particular embodiments of the linear peptide inhibitor of Formula III, X4 and/or X9 are any amino acid.

[00433] In particular embodiments of a peptide inhibitor of Formula III, one or more, two or more, or all three of X1, X2, and X3 are absent. In certain embodiments, X1 is absent. In certain embodiments, X1 and X2 are absent. In certain embodiments, X1, X2 and X3 are absent.

[00434] In particular embodiments of a peptide inhibitor of Formula III, one or more, two or more, three or more, four or more, or all of X16, X17, X18, X19 and X20 are absent. In particular embodiments of a peptide inhibitor of Formula III, one or more, two or more, three or more, or all of X17, X18, X19 and X20 are absent. In certain embodiments, one or more, two or more, or all three of X17, X19 and X20 are absent. In certain embodiments, one or more of X1, X2 and X3 are absent; and one or more, two or more, three or more, or four of X17, X18, X19 and X20 are absent.

[00435] In particular embodiments of a peptide inhibitor of Formula III, one of X4 or X9 is Abu, and the other of X4 or X9 is not Abu. In certain embodiments, X4 is Abu and X9 is Cys.

[00436] In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, or four of the following features: X5 is Arg or Gln; X6 is Thr; X7 is Trp; and X8 is Gln. In particular embodiments, X5 is Gln, X6 is Thr, X7 is Trp, and X8 is Gln. In certain embodiments, X5 is Gln. In certain embodiments, X1, X2 and X3 are absent. In certain embodiments, X4 is Abu and X9 is Cys.

[00437] In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following

features: X10 is Tyr or a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl) or Phe(4-Me); X12 is Arg, hLeu, (D)Asn, or any alpha methyl amino acids including, Aib,  $\alpha$ -MeLys,  $\alpha$ -MeLeu or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is  $\beta$ -Ala, Gln, Gly, Ser, Ala; and X16 is absent or AEA. In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following features: X10 is Tyr or a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl) or Phe(4-Me); X12 is Arg, hLeu, (D)Asn, or any alpha methyl amino acids including, Aib,  $\alpha$ -MeLys,  $\alpha$ -MeLeu or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly, Ser, Ala; and X16 is absent or AEA. In certain embodiments, the Phe analog is Phe(4-OBzl), Phe(4-OMe), Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-carboxy), Phe[4-(2aminoethoxy)] or Phe(4-guanadino). In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent. In certain embodiments, X4 is Abu and X9 is Cys.

[00438] In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following features: X10 is Tyr or a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(4-OBzl) or Phe(4-Me); X12 is Arg, hLeu, (D)Asn, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal,; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly; and X16 is absent or AEA. In certain embodiments, the Phe analog is Phe(4-OBzl), Phe(4-OMe), Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-carboxy), Phe(4-(2aminoethoxy)) or Phe(4-guanadino). In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent. In certain embodiments, X4 is Abu and X9 is Cys.


[00439] In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven of the following features: X5 is Arg or Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me); X12 is Aib,  $\alpha$ -MeLys,  $\alpha$ -MeLeu, 4-

amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Agp, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeSer,  $\alpha$ -MeVal,  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is  $\beta$ -ala, Gly, Ser, Ala; and X16 is absent or AEA. In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven of the following features: X5 is Arg or Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me); X12 is Aib,  $\alpha$ -MeLys,  $\alpha$ -MeLeu or  $\alpha$ -MeOrn; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly, Ser, Ala; and X16 is absent or AEA. In certain embodiments, the Phe analog is Phe(4-OBzl), Phe(4-OMe), Phe[4-(2aminoethoxy)], Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-CO<sub>2</sub>H), or Phe(4-guanadino). In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent. In certain embodiments, X4 is Abu and X9 is Cys.

[00440] In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven of the following features: X5 is Arg or Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr or a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me); X12 is Arg, hLeu, (D)Asn, 4-amino-4-carboxy-tetrahydropyran, Achc, Acpc, Acbc, Acvc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeSer,  $\alpha$ -MeVal,; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is  $\beta$ -Ala, Asn or Gly; and X16 is absent or AEA. In particular embodiments, a peptide inhibitor of Formula III comprises one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, or eleven of the following features: X5 is Arg or Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr or a Phe analog; X11 is Trp, 2-Nal, 1-Nal, Phe(4-O-Allyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Phe(Bzl) or Phe(4-Me); X12 is Arg, hLeu, (D)Asn,  $\alpha$ -MeLys,  $\alpha$ -MeLeu or  $\alpha$ -MeOrn, Aib; X13 is Lys, Glu or Lys(Ac); X14 is Phe or Asn; X15 is Gly; and X16 is absent or AEA. In certain embodiments, the Phe analog is Phe(4-OBzl), Phe(4OMe), Phe(4-CONH<sub>2</sub>), Phe(3,4-Cl<sub>2</sub>), Phe(4-tBu), Phe(4-NH<sub>2</sub>), Phe(4-Br), Phe(4-CN), Phe(4-CO<sub>2</sub>H), Phe(4-(2-aminoethoxy)) or Phe(4-

guanidino). In certain embodiments, X11 is 2-Nal or 1-Nal. In certain embodiments, X1, X2 and X3 are absent. In certain embodiments, X4 is Abu and X9 is Cys.

**[00441]** In certain embodiments, the present invention includes a peptide of 8 to 20, 8 to 16 or 8 to 12 amino acids, optionally cyclized, comprising or consisting of a core sequence of Formula IIIb:



**[00442]** wherein Xaa4 and Xaa9 are each independently selected from Abu and Cys, wherein Xaa4 and Xaa9 are not both the same; Xaa5, Xaa6 and Xaa8 are any amino acid residue; Xaa10 is Tyr, a Phe analog or 2-Nal, and Xaa11 is 2-Nal or Trp, wherein the peptide inhibits binding of IL-23 to IL-23R. In particular embodiments, Xaa10 is Phe(4-OMe), 2-Nal, or Phe[4-(2-aminoethoxy)]. In one embodiment, Xaa10 is Phe(4-OMe). In one embodiment, Xaa7 is Phe[4-(2-aminoethoxy)]. In one embodiment, Xaa11 is 2-Nal. In certain embodiments, the peptide is cyclized via Xaa4 and Xaa9. In particular embodiments, the Phe analog is Phe[4-(2-aminoethoxy)] or Phe(4-OMe). In certain embodiments, Xaa4 is Abu and Xaa9 is Cys, and the peptide is cyclized via Xaa4 and Xaa9. In particular embodiments, the peptide is a peptide inhibitor of Formula III, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain embodiments, a peptide of Formula IIIb comprises a Glu,(D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln bound to Xaa4. In certain embodiments, it is (D)Arg or (D)Phe.

**[00443]** In certain embodiments, the present invention includes a peptide of 8 to 20, 8 to 16 or 8 to 12 amino acids, optionally cyclized, comprising or consisting of a core sequence of Formula IIIc:



**[00444]** wherein Xaa5, Xaa6 and Xaa8 are any amino acid residue; and wherein the peptide inhibits binding of IL-23 to IL-23R. In certain embodiments, the peptide is cyclized via Abu at Xaa4 and Cys at Xaa9. In certain embodiments, the peptide is a peptide inhibitor of Formula III, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain embodiments, a peptide of Formula

IIIc comprises a Glu, (D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln bound to Abu. In certain embodiments, it is (D)Arg or (D)Phe.

**[00445]** In certain embodiments, the present invention includes a peptide of 8 to 20, 8 to 16 or 8 to 12 amino acids, optionally cyclized, comprising or consisting of a core sequence of Formula IIId:



**[00446]** wherein Xaa5, Xaa6 and Xaa8 are any amino acid residue; Xaa10 is a modified Phe; and wherein the peptide inhibits binding of IL-23 to IL-23R. In particular embodiments, the modified Phe is Phe(4-tBu), Phe(4-guanidino), Phe[4-(2-aminoethoxy)], Phe(4-CO<sub>2</sub>H), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(CONH<sub>2</sub>) or Phe(4-Me). In particular embodiments, the modified Phe is Phe(4-tBu), Phe(4-guanidino), Phe[4-(2-aminoethoxy)], Phe(4-CO<sub>2</sub>H), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), or Phe(4-Me). In one embodiment, Xaa10 is Phe[4-(2-aminoethoxy)] or Phe(4-OMe). In one embodiment, Xaa10 is Phe[4-(2-aminoethoxy)]. In certain embodiments, the peptide is cyclized via Abu at Xaa4 and Cys at Xaa9. In certain embodiments, the peptide is a peptide inhibitor of Formula III, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain embodiments, a peptide of Formula IIId comprises a Glu, (D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln bound to Abu. In certain embodiments, it is (D)Arg or (D)Phe.

**[00447]** In certain embodiments, the present invention includes a peptide, optionally 8 to 20, 8 to 16 or 8 to 12 amino acids, optionally cyclized, comprising or consisting of a core sequence of Formula IIIe:



**[00448]** wherein Xaa5, Xaa6 and Xaa8 are any amino acid residue. In certain embodiments, the peptide is cyclized via Abu at Xaa4 and Cys at Xaa9. In certain embodiments, the peptide is a peptide inhibitor of Formula III, and wherein in certain embodiments, X1, X2 and X3 are absent. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R. In certain

embodiments, a peptide of Formula IIIb comprises a Glu, (D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln bound to Abu. In certain embodiments, it is (D)Arg or (D)Phe.

[00449] In one embodiment, Xaa5 and Xaa8 is Gln. In one embodiment, Xaa6 is Thr. In certain embodiments, the peptide is cyclized via Abu at Xaa4 and Cys at Xaa9.

[00450] In particular embodiments of a peptide inhibitor of Formula III, the peptide inhibitor has a structure shown in any of Tables 5A-5C or comprises an amino acid sequence set forth in Tables 5A-5C.

[00451] In certain aspects, the present invention provides a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Vf):

X1-X2-X3-Abu-X5-X6-X7-X8-Cys-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20  
(Vf),

wherein:

X1 is absent;

X2 is absent or X2 is D-Asp, E, R, D-Arg, F, D-Phe, 2-Nal, T, L, D-Gln, or D-Asn;

X3 is D-Arg;

X5 is N, Q, Cit, Lys, or a Lys conjugate (e.g., Lys(IVA), Lys(biotin), Lys(octanyl), Lys(Palm), Lys(PEG), Lys(PEG8), Lys(PEG11-Palm), Lys(Ac));

X6 is T, S or V;

X7 is W, 1-Nal, or 2-Nal;

X8 is Q, Cit, N, Aib or Lys(Ac);

X10 is Phe[4-(2-aminoethoxy)], Phe[4-(2-acetylaminoethoxy)] or Phe(4-CONH<sub>2</sub>);

X11 is 2-Nal;

X12 is 4-amino-4-carboxy-tetrahydropyran, Aib,  $\alpha$ MeLeu,  $\alpha$ MeLys, or an  $\alpha$ MeLys conjugate (e.g.,  $\alpha$ MeLys(Ac),  $\alpha$ MeLys(PEG4-Palm),  $\alpha$ MeLys(PEG4-Lauryl),  $\alpha$ MeLys(PEG4IsoGluPalm),  $\alpha$ MeLys(PEG4IsoGluLauryl),  $\alpha$ MeLys(IVA),  $\alpha$ MeLys(biotin), or  $\alpha$ MeLys(octanyl));

X13 is Q, E, Cit or a Lys conjugate (e.g., Lys(Ac), Lys(PEG4-isoGlu-Palm), Lys(PEG4-octanyl), Lys(PEG4-Palm), Lys(biotin), Lys(octanyl), Lys(Palm), Pys(PEG8), or Lys(PEG11-Palm));

X14 is N, Cit, Q, L, G, S, Aib, F, 2-Nap, N-Me-Ala, R, W, nLeu, Tic or a Lys conjugate (e.g., Lys(Ac));

X15 is N, Cit, Q,  $\beta$ Ala, Lys(Ac) or Aib; and

X16, X17, X18, X19 and X20 are absent.

In particular embodiments, X2 is D-Asp, E, R, D-Arg, F, D-Phe, 2-Nal, T, L, D-Gln, or D-Asn

[00452] In certain aspects, the present invention provides a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Vh):

X1-X2-X3-Abu-X5-X6-X7-X8-Cys-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20  
(Vh),

wherein:

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any D-amino acid or absent;

X4 is Cys, hCys, Pen, hPen, Abu, Ser, hSer or chemical moiety capable of forming a bond with X9;

X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn, Gln, Arg, Ser, Glu or Thr;

X6 is Thr, Ser, Asp, Ile or any amino acid;

X7 is Trp, 6-Chloro-Trp, 1-Nap or 2-Nap;

X8 is Glu, Gln, Asn, Lys(Ac), Cit, Cav, Lys(N- $\epsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), or Lys(N- $\epsilon$ -Palmitoyl);

X9 is Cys, hCys, Pen, hPen Abu, or any amino acid or chemical moiety capable of forming a bond with X4;

X10 is 2-Nal, a Phe analog, Tyr, or a Tyr analog;

X11 is 1-Nal, 2-Nal, Phe(3,4-dimethoxy), 5-HydroxyTrp, Phe(3,4-Cl2), Trp or Tyr(3-tBu);

X12 is Aib, 4-amino-4-carboxy-tetrahydropyran, any alpha-methylamino acid, alpha-ethyl-amino acid, Achc, Acvc, Acbc, Acpc, 4-amino-4-carboxy-piperidine, 3-Pal, Agp,  $\square$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -MeLeu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, Cav, Cha, Cit, Cpa, D-Asn, Glu, His, hLeu, hArg, Lys, Leu, Octgly, Orn, piperidine, Arg, Ser, Thr or THP;

X13 is Lys(Ac), Gln, Cit, Glu, or any amino acid;

X14 is Asn, Gln, Lys(Ac), Cit, Cav, Lys(N- $\epsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), Lys(N- $\epsilon$ -Palmitoyl), Asp, or any amino acid;

X15 is  $\beta$ -Ala, Asn, Gly, Gln, Ala, Ser, Aib, Asp or Cit;

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent.

[00453] In certain embodiments of any of the peptide inhibitors described herein, including but not limited to those of Formula (If) and (Ih), the peptide inhibitor is cyclized via a bond, e.g., a thioether bond, between X4 and X9. In certain embodiments, the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

[00454] In certain embodiments, X1, X2 and X3 are absent. In certain embodiments, X1 and X2 are absent. In certain embodiments, X1 is a D-amino acid or absent. In certain embodiments, X2 is a D-amino acid or absent.

[00455] In certain embodiments, X5 is Ala,  $\alpha$ -MeOrn,  $\alpha$ -MeSer, Cit, Dap, Dab, Dap(Ac), Gly, Lys, Asn, N-MeGln, N-MeArg, Orn, Gln, Arg, Ser, or Thr;

[00456] In certain embodiments, X5 is N, X6 is T, X7 is W, or X8 is Q. In certain embodiments, X5 is N, X6 is T, X7 is W, and X8 is Q.

[00457] In certain embodiments, X5 is Q, X6 is T, X7 is W, or X8 is Q. In certain embodiments, X5 is Q, X6 is T, X7 is W, and X8 is Q.

[00458] In certain embodiments, X5 is N, X6 is T, X7 is W, and X8 is Cit.

[00459] In certain embodiments, X10 is Phe[4-(2-aminoethoxy)].

[00460] In certain embodiments, X12 is 4-amino-4-carboxy-tetrahydropyran, Aib,  $\alpha$ MeLeu, or  $\alpha$ MeLys. In certain embodiments, X12 is 4-amino-4-carboxy-tetrahydropyran.

[00461] In certain embodiments, X13 is E or Lys(Ac). In certain embodiments, X13 is Lys(Ac).

[00462] In certain embodiments, X14 is Asn, Gln, Lys(Ac), Cit, Cav, Lys(N- $\epsilon$ -(N- $\alpha$ -Palmitoyl-L- $\gamma$ -glutamyl)), Lys(N- $\epsilon$ -Palmitoyl), or any amino acid;

[00463] In certain embodiments, X15 is  $\beta$ -Ala, Asn, Gly, Gln, Ala, Ser, Aib, or Cit.

[00464] In certain embodiments, X14 is N.

[00465] In certain embodiments, X15 is N.

[00466] In certain embodiments, X16 is a D-amino acid or absent. In certain embodiments, X17 is a D-amino acid or absent. In certain embodiments, X18 is a D-amino acid or absent. In certain embodiments, X19 is a D-amino acid or absent. In certain embodiments, X20 is a D-amino acid or absent.

[00467] In certain embodiments, X2 is absent; X3 is absent; X5 is Q, X6 is T, X7 is W, and X8 is Q; X10 is Phe[4-(2-aminoethoxy)]; X12 is 4-amino-4-carboxy-tetrahydropyran, Aib,  $\alpha$ MeLeu, or  $\alpha$ MeLys; X13 is E or Lys(Ac); X14 is N; and X15 is N. In certain embodiments, X12 is 4-amino-4-carboxy-tetrahydropyran and X13 is Lys(Ac).

[00468] In certain embodiments, any of the amino acids of the peptide inhibitor are connected by a linker moiety, e.g., a PEG.

[00469] In certain embodiments, the N-terminus of the peptide inhibitor comprises an Ac group.

[00470] In certain embodiments, the C-terminus of the peptide inhibitor comprises an NH<sub>2</sub> group.


[00471] In certain embodiments, the present invention includes a peptide comprising or consisting of an amino acid sequence shown in any of the Table 4s or Table 5s, or a peptide inhibitor comprising or consisting of a structure shown in any of the Table 4s or Table 5s (or a pharmaceutically acceptable salt thereof). In particular embodiments, the peptide does not include the conjugated moieties but does include the Abu residue. In particular embodiments, the peptide or inhibitor comprises a thioether bond between the two Abu and Cys residues, or between the two outermost amino acids within the brackets following the term “cyclo”, which indicated the presence of a cyclic structure. In particular embodiments, the inhibitor is an acetate salt. The peptide sequence of illustrative inhibitors is shown in Tables 4 and 5 from N-term to C-term, with conjugated moieties, and N-terminal Ac and/or C-terminal NH<sub>2</sub> groups indicated. The cyclic structure is indicated by “Cyclo” as illustrated in Table 5, indicating the presence of a thioether bond between the bracketed Abu at X4 and Cys at X9.

Table 4. Illustrative Thioether Peptide Inhibitors

Biotin-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
--

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
Ac-E-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-[(D)Asp]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-R-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-[(D)Arg]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-F-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-[(D)Phe]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-[2-Nal]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-T-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-L-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-[(D)Gln]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-[(D)Asn]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(PEG4-Alexa488)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Alexa488]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Alexa647]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Alexa-647]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Alexa647]-[PEG12]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>
[Alexa488]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>

Table 5A. Illustrative Thioether Peptide Inhibitors



Ac-Cyclo-[[Abu]-XXWXC]-[Phe(4-OMe)]-[2-Nal]-XXXX-NH<sub>2</sub>

Sequence
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNE-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNF-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNKK-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNWW-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNT-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNG-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNPK-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNPG-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNEP-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGK-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNPT-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGF-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGW-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGQ-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGG-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNKKK-NH <sub>2</sub>

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNEEE-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNFFF-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNTTT-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGG-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGF-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGE-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGQ-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGT-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGGR-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGGF-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGGE-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGGG-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGGGGT-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNRRRR-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNGFFFF-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNEEEE-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNQQQQQ-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENNTTTTT-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-GGG-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-RRR-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-FFF-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>

Ac-EEE-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-QQQ-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-TTT-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-RG-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-FG-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-EG-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-QG-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-TG-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Palm)]-NN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(PEG11-Palm)]-NN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(isoGlu-Palm)]-NN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ahx-Palm)]-NN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(isoGlu-Ahx-Palm)]-NN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(isoGlu-Ahx-Palm)]-NN-NH <sub>2</sub>
[Palm]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Palm-isoGlu]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Palm-PEG11]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Palm-Ahx]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
[Palm-Ahx-isoGlu]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-Lys[Palm]-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-Lys[Peg11-Palm]-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-Lys[isoGlu-Palm]-NH <sub>2</sub>

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-Lys[Ahx-Palm]-NH <sub>2</sub>
Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-Lys[isoGlu-Ahx-Palm]-NH <sub>2</sub>

Table 5B. Illustrative Thioether Peptides

Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]]-END-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDD-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-END-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[Tetrahydropyran-A]-END-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[Tetrahydropyran-A]-EDN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-ETWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-END-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[Tetrahydropyran-A]-EDN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-ETWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>
Ac-[D-Arg]- Cyclo-[Abu-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[Tetrahydropyran-A]-ENN-OH

Illustrative Peptide Inhibitors Containing Cyclic Amides

[00472] In certain embodiments, the present invention includes a peptide inhibitor of an interleukin-23 receptor, wherein the peptide inhibitor has the structure of Formula IV:




[00473] or a pharmaceutically acceptable salt or solvate thereof,

[00474] wherein  $R^1$  is a bond, hydrogen, an C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing;

[00475]  $R^2$  is a bond, OH or  $NH_2$ ; and

[00476] X is an amino acid sequence of 8 to 20 amino acids, comprising or consisting of the sequence of Formula IVa:



[00477] wherein

X1 is absent or any amino acid;

X2 is absent or any amino acid;

X3 is absent or any amino acid;

X4 is Dap, Dab, Glu, Asp, (D)-Asp or (D)-Dab;

X5 is Gln, Ala, Cys, Cit, Asp, Dab, Dap, Glu, Phe, Gly, His, hCys, Lys, Leu, Met, Asn, N-Me-Ala, N-M-Asn, N-Me-Lys, N-Me-Gln, N-Me-Arg, Orn, Pro, Pen, Gln, Arg, Ser, Thr, or Val;

X6 is Thr, Asp, Glu, Phe, Asn, Pro, Arg, Ser, or Thr;

X7 is Trp, Glu, Gly, Ile, Asn, Pro, Arg, Thr or OctGly;

X8 is Gln, Glu, Phe, Lys, Asn, , Pro, Arg, Thr, or Trp;

X9 is Dap, Dab, Glu, Asp, (D)-Asp or (D)-Dab;

X10 is Tyr(OMe)Phe(4-OMe), 1-Nal, 2-Nal, Aic, a-MePhe, Bip, (D)Cys, Cha, DMT, (D)Tyr), Glu, Phe, His, hPhe(3,4-dimethoxy), hTyr, N-Me-Tyr, Trp, Phe(4-CONH<sub>2</sub>), Phe(4-phenoxy), Thr, Tic, Tyr, Tyr(3-tBu), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5-F<sub>2</sub>), Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>), Bip, Cha, 4-pyridylalanine,  $\beta$ hTyr, OctGly, Phe(4-N<sub>3</sub>), Phe(4-Br) or Phe[4-(2-aminoethoxy)];

X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe(3,5-F<sub>2</sub>), Phe(4-CONH<sub>2</sub>), Phe(4-F), 4-phenylcyclohexylalanine, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, BIP, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Trp, Tyr, Phe(4-OMe), Phe(4-Me), Trp(2,5,7-tri-tertButyl), Phe(4-OAllyl), Tyr(3-tBu), Phe(4-tBu), Phe(4-guanidino), Tyr(Bzl), or OctGly;

X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr Tle, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeSer,  $\alpha$ -MeVal,;

X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, Lys, Lys(Ac), Leu, Asn, Ogl, Pro, Gln, Arg, Ser,  $\beta$ -spiro-pip, Thr, Tba, Tlc, Val or Tyr;

X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr;

X15 is  $\beta$ -ala, Asn, Gly, (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser;

X16 is absent, Gly, Ala, Asp, Ser, Pro, Asn or Thr;

X17 is absent, Glu, Ser, Gly or Gln;

X18 is absent or any amino acid;

X19 is absent or any amino acid; and

X20 is absent or any amino acid.

[00478] In certain embodiments of IVa: X12 is  $\alpha$ -MeLys,  $\alpha$ -MeOrn,  $\alpha$ -MeLeu, Aib, (D)Ala, (D)Asn, (D)Leu, (D)Asp, (D)Phe, (D)Thr, 3-Pal, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeu, hArg, hLeu, Ile, Lys, Leu, Asn, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Arg, Ser, Thr or Tle; X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Thr, (D)Phe, Ala, Aib,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hAla,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, Glu, Phe, hLeuLys, Leu, Asn, Ogl, Pro, Gln, Arg, Ser,  $\beta$ -spiro-pip, Thr, Tba, Tlc, Val or Tyr; X14 is Asn, Glu, Phe, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic or Tyr; and X15 is Gly, (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Ala, AEA, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Gln, Arg or Ser.

[00479] In particular embodiments of a peptide inhibitor of Formula (IV): X5 is Cys, Cit, Asp, Dab, Dap, Gly, His, hCys, Lys, Met, Asn, N-Me-Ala, N-Me-Asn, N-Me-Lys, N-Me-Gln, N-Me-Arg, Orn, Pro, Pen, Gln, Val; X6 is Glu, Arg, Ser; X7 is Trp, Glu, Gly, Ile, Asn, Pro, Arg, Thr or OctGly; X8 is Phe, Asn, Pro, Arg, Thr, Trp; X10 is Phe(4-OMe), 1-Nal, 2-Nal, Aic,  $\alpha$ -MePhe, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, His, hPhe(3,4-dimethoxy), hTyr, N-Me-Tyr, Trp, Phe(4-CONH<sub>2</sub>), Phe-(4-phenoxy), Thr, Tic, Tyr(3-tBu), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5-F<sub>2</sub>), PheCH<sub>2</sub>CO<sub>2</sub>H, Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>), Bip, Cha, 4-PyridylAlanine,  $\beta$ hTyr, OctgGly, Tyr(4-N<sub>3</sub>), Phe(4-Br), Phe[4-(2-aminoethoxy)]; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe(3,5-F<sub>2</sub>), Phe(4-CONH<sub>2</sub>), Phe(4-F), 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe, Nal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, BIP, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Tyr, Phe(4-OMe), Phe(4-Me), Tyr(2,5,7-tri-tert-Butyl), Phe(4-OAllyl), Phe(3-tBu), Phe(4-tBu), Phe(4-guanidino), Tyr(Bzl), OctGly; X12 is  $\alpha$ -Me-Lys, D-Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Tyr, Aib,  $\alpha$ -MeLeu,  $\alpha$ -MeOrn, Aib,  $\beta$ -Ala,  $\beta$ hAla,  $\beta$ hArg,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Glu, hArg, Ile, Lys, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Ser, Thr, Tle, 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys(Ac), ,  $\alpha$ -MeSer,  $\alpha$ -MeVal; X13 is Lys, Lys(Ac), (D)Asn, (D)Leu, (D)Phe, (D)Thr, Ala,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, hLeu, Asn, Ogl, Pro, Gln, Ser, Thr, Tba, Tle; X14 is Glu, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic; X15 is (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Aea, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Asn, Arg or  $\beta$ -Ala; X16 is Gly, Ser, Pro, Asn, Thr; or X17 is Glu, Ser, Gly, Gln.

[00480] In particular embodiments of a peptide inhibitor of Formula (IV): X5 is Cys, Cit, Asp, Dab, Dap, Gly, His, hCys, Lys, Met, Asn, N-Me-Ala, N-Me-Asn, N-Me-Lys, N-Me-Gln, N-Me-Arg, Orn, Pro, Pen, Gln, Val; X6 is Glu, Arg, Ser; X7 is Trp, Glu, Gly, Ile, Asn, Pro, Arg, Thr or OctGly; X8 is Phe, Asn, Pro, Arg, Thr, Trp; X10 is Phe(4-OMe), 1-Nal, 2-Nal, Aic,  $\alpha$ -MePhe, Bip, (D)Cys, Cha, DMT, (D)Tyr, Glu, His, hPhe(3,4-dimethoxy), hTyr, N-Me-Tyr, Trp, Phe(4-CONH<sub>2</sub>), Phe-(4-phenoxy), Thr, Tic, Tyr(3-tBu), Phe(4-tBu), Phe(4-CN), Phe(4-Br), Phe(4-NH<sub>2</sub>), Phe(4-F), Phe(3,5-F<sub>2</sub>), PheCH<sub>2</sub>CO<sub>2</sub>H, Phe(penta-F), Phe(3,4-Cl<sub>2</sub>), Phe(4-CF<sub>3</sub>), Bip, Cha, 4-PyridylAlanine,  $\beta$ hTyr, OctgGly, Tyr(4-N<sub>3</sub>), Phe(4-Br), Phe[4-(2-aminoethoxy)]; X11 is 2-Nal, 1-Nal, 2,4-dimethylPhe, Bip, Phe(3,4-Cl<sub>2</sub>), Phe(3,5-F<sub>2</sub>), Phe(4-CONH<sub>2</sub>), Phe(4-F), 4-phenylcyclohexyl, Phe(4-CF<sub>3</sub>),  $\alpha$ -MePhe, Nal,  $\beta$ hPhe,  $\beta$ hTyr,  $\beta$ hTrp, BIP, Nva(5-phenyl), Phe, His, hPhe, Tic, Tqa, Tyr, Phe(4-OMe), Phe(4-Me), Tyr(2,5,7-tri-tert-Butyl), Phe(4-OAllyl), Phe(3-tBu), Phe(4-tBu), Phe(4-guanidino), Tyr(Bzl), OctGly; X12 is  $\alpha$ -Me-Lys, D-Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Tyr, Aib,  $\alpha$ -MeLeu,  $\alpha$ -MeOrn, Aib,  $\beta$ -Ala,  $\beta$ hAla,  $\beta$ hArg,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Glu, hArg, Ile, Lys, N-MeLeu, N-MeArg, Ogl, Orn, Pro, Gln, Ser, Thr, Tle; X13 is Lys(Ac), (D)Asn, (D)Leu, (D)Phe, (D)Thr, Ala,  $\alpha$ -MeLeu, Aib,  $\beta$ -Ala,  $\beta$ -Glu,  $\beta$ hLeu,  $\beta$ hVal,  $\beta$ -spiro-pip, Cha, Chg, Asp, Dab, Dap,  $\alpha$ -DiethylGly, hLeu, Asn, Ogl, Pro, Gln, Ser, Thr, Tba, Tle; X14 is Glu, Gly, His, Lys, Leu, Met, Asn, Pro, Gln, Arg, Ser, Thr, Tic; X15 is (D)Ala, (D)Asn, (D)Asp, (D)Leu, (D)Phe, (D)Thr, Aea, Asp, Glu, Phe, Gly, Lys, Leu, Pro, Arg; X16 is Gly, Ser, Pro, Asn, Thr; or X17 is Glu, Ser, Gly, Gln.

[00481] In certain embodiments, the peptide inhibitor is cyclized. In particular embodiments, the peptide is cyclized through an intramolecular bond between X4 and X9. In particular embodiments, the intramolecular bond is an amide bond.

[00482] In certain embodiments, the peptide inhibitor is linear or not cyclized.

[00483] In particular embodiments of a peptide inhibitor of Formula IV, one or more, two or more, or all three of X1, X2, and X3 are absent.

[00484] In certain embodiments, X3 is Glu, (D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln. In certain embodiments, X3 is (D)Arg or (D)Phe.

[00485] In particular embodiments of a peptide inhibitor of Formula IV, one or more, two or more, or all three of X17, X19 and X20 are absent.

[00486] In particular embodiments of a peptide inhibitor of Formula IV, X4 is Dap, Dab, or (D)Dab, and X9 is Glu, (D)Asp, or Asp. In particular embodiments of a peptide inhibitor of Formula I, X4 is Glu, (D)Asp or Asp, and X9 is Dab, Dap or (D)Dab.

[00487] In particular embodiments of a peptide inhibitor of Formula IV, X18 is (D)-Lys. In certain embodiments, X17 is absent and X18 is (D)-Lys.

[00488] In particular embodiments of a peptide inhibitor of Formula IV, the peptide inhibitor includes one or more, two or more, three or more, or all four of the following features: X5 is Gln; X6 is Thr; X7 is Trp; and X8 is Gln.

[00489] In particular embodiments of a peptide inhibitor of Formula IV, the peptide inhibitor includes one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following features: X10 is Tyr, Phe[4-(2-aminoethoxy)], Phe(4-CONH<sub>2</sub>) or Phe(4-OMe); X11 is 2-Nal or Trp; X12 is 4-amino-4-carboxy-tetrahydropyran, Achc Acpc, Acbc, Acvc, Aib,  $\alpha$ -DiethylGly,  $\alpha$ -MeLys,  $\alpha$ -MeLys(Ac),  $\alpha$ -Me-Leu,  $\alpha$ -MeOrn,  $\alpha$ -MeSer,  $\alpha$ -MeVal, or Arg; X13 is Glu or Lys(Ac); X14 is Asn; X15 is Gly, Asn, or  $\beta$ -Ala; and X16 is AEA. In particular embodiments of a peptide inhibitor of Formula IV, the peptide inhibitor includes one or more, two or more, three or more, four or more, five or more, six or more, or seven of the following features: X10 is Tyr; X11 is Trp; X12 is Arg; X13 is Glu; X14 is Asn; X15 is Gly; and X16 is AEA.

[00490] In particular embodiments of a peptide inhibitor of Formula IV, the peptide inhibitor includes one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more ten or more or all of the following features: X5 is Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr; X11 is Trp; X12 is Arg; X13 is Glu or Lys(Ac); X14 is Asn; X15 is Gly; and X16 is AEA. In particular embodiments of a peptide inhibitor of Formula IV, the peptide inhibitor includes one or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more ten or more or all of the following

features: X5 is Gln; X6 is Thr; X7 is Trp; X8 is Gln; X10 is Tyr; X11 is Trp; X12 is Arg; X13 is Glu; X14 is Asn; X15 is Gly; and X16 is AEA.

[00491] In certain embodiments of a peptide inhibitor of Formula IV, the peptide is cyclized via X4 and X9; X5, X6, X7 and X8 are Gln, Thr, Trp, and Gln, respectively; and X10, X11, X12, X13, X14, X15, and X16 are Tyr, Trp, Arg, Glu, Asn, Gly, and AEA, respectively.

[00492] In certain embodiments, the present invention includes a peptide of 8 to 20 amino acids, optionally cyclized, comprising or consisting of having a core sequence comprising:



[00493] wherein Xaa4 and Xaa9 are each independently selected from Dap, Dab, Glu, Asp, (D)-Asp and (D)-Dab, wherein Xaa4 and Xaa9 are capable of forming an intramolecular bond, e.g., a cyclic amide; and Xaa5, Xaa6 and Xaa8 are any amino acid residue, wherein the peptide inhibits binding of IL-23 to IL-23R. In particular embodiments, the peptide inhibitor is a peptide inhibitor of Formula IV. In particular embodiments, the peptide inhibits the binding of IL-23 to IL-23R.

[00494] In certain embodiments, of a peptide inhibitor of Formula IV, the peptide inhibitor has a structure shown in Table 7 or comprises an amino acid sequence set forth in Table 7.

[00495] Certain illustrative peptide inhibitors of the present invention are also shown in any of Formulas (Va), (Vb), (Vc), (Vd), (Ve), (Vf), (Vg) and (Vh), and in Tables 2-5, which provide the amino acid sequence of selected peptide inhibitors. These peptide inhibitors are acetate salts.

#### Optional Characteristics of Peptide Inhibitors

[00496] Any of the peptide inhibitors of the present invention may be further defined, e.g., as described below. It is understood that each of the further defining features described herein may be applied to any peptide inhibitors where the amino acids designated at particular positions allow the presence of the further defining feature.

[00497] In certain embodiments of any of the peptide inhibitors described herein, the peptide inhibitor is cyclized.

[00498] In certain embodiments of any of the peptide inhibitors described herein, the peptide inhibitor or monomer subunit thereof is linear or not cyclized. In certain embodiments where the peptide is linear or not cyclized, X4 and X9 can be any amino acid.

[00499] In certain embodiments, the peptide inhibitor is cyclized, e.g., through X4 and X9.

[00500] In various embodiments, R<sup>1</sup> is a bond, hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, or a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing, e.g., acetyl. It is understood that the R<sup>1</sup> may replace or be present in addition to the typical amine group located at the amino terminus of a peptide. It is further understood that R<sup>1</sup> may be absent. In certain embodiments, the peptide inhibitor comprises an N-terminus selected from hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, or a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing, e.g., acetyl. In particular embodiments of any of the peptide inhibitors described herein, R<sup>1</sup> or the N-terminal moiety is hydrogen. In certain embodiments, R<sup>1</sup> is a bond, e.g., a covalent bond.

[00501] In certain embodiments of any of the peptide inhibitors having any of the various Formulas set forth herein, R<sup>1</sup> or the N-terminal moiety is selected from methyl, acetyl, formyl, benzoyl, trifluoroacetyl, isovaleryl, isobutyryl, octanyl, and the conjugated amides of lauric acid, hexadecanoic acid, and  $\gamma$ -Glu-hexadecanoic acid. In one embodiment, R<sup>1</sup> or the N-terminal moiety is pGlu. In certain embodiments, R<sup>1</sup> is hydrogen. In particular embodiments, R<sup>1</sup> is acetyl, whereby the peptide inhibitor is acylated at its N-terminus, e.g., to cap or protect an N-terminal amino acid residue, e.g., an N-terminal Pen or Abu residue.

[00502] In certain embodiments of any of the peptide inhibitors described herein, R<sup>1</sup> or the N-terminal moiety is an acid. In certain embodiments, R<sup>1</sup> or the N-terminal moiety is an acid selected from acetic acid, formic acid, benzoic acid, trifluoroacetic acid, isovaleric acid, isobutyric acid, octanoic acid, lauric acid, hexadecanoic acid, 4-Biphenylacetic acid, 4-fluorophenylacetic acid, gallic acid, pyroglutamic acid, cyclopentanepropionic acid, glycolic acid, oxalic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, palmitic acid, benzoic acid, 3-(4-hydroxybenzoyl) benzoic acid, cinnamic acid, mandelic acid, 4-methylbicyclo(2.2.2)-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl

sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, an alkylsulfonic acid and an arylsulfonic acid.

**[00503]** In particular embodiments, R<sup>1</sup> or the N-terminal moiety is an alkylsulfonic acid selected from methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, and 2-hydroxyethanesulfonic acid.

**[00504]** In particular embodiments, R<sup>1</sup> or the N-terminal moiety is an arylsulfonic acid selected from benzenesulfonic acid, 4- chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, and camphorsulfonic acid.

**[00505]** In some embodiments, wherein a peptide of the present invention comprises a conjugation to an acidic compound such as, e.g., isovaleric acid, isobutyric acid, valeric acid, and the like, the presence of such a conjugation is referenced in the acid form. So, for example, but not to be limited in any way, instead of indicating a conjugation of isovaleric acid to a peptide by referencing isovaleroyl (e.g., isovaleroyl-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH<sub>2</sub>, in some embodiments, the present application references such a conjugation as isovaleric acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH<sub>2</sub>. Reference to the conjugation in its acid form is intended to encompass the form present in the peptide inhibitor.

**[00506]** In certain embodiments, the peptide inhibitor comprises a C-terminus (e.g., R<sup>2</sup> or the C-terminal moiety) selected from a bond, OH or NH<sub>2</sub>. In certain embodiments, R<sup>2</sup> is a bond. In various embodiments of any of the peptide inhibitors having any of the various Formulas set forth herein, R<sup>2</sup> or the C-terminal moiety is OH or NH<sub>2</sub>. It is understood that the R<sup>2</sup> or the C-terminal moiety may replace or be present in addition to the carboxyl group typically located at the carboxy terminus of a peptide. It is further understood that R<sup>2</sup> may be absent.

**[00507]** In particular embodiments of any of the peptide inhibitors having any of the various Formulae set forth herein, X comprises or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 20 amino acid residues, 8 to 20 amino acid residues, 9 to 20 amino acid residues, 7 to 18

amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues.

**[00508]** In certain embodiments of any of the Formulae set forth herein, X either or both does not comprise or does not consist of an amino acid sequence set forth in US Patent Application Publication No. US2013/0029907. In certain embodiments of any of the Formulae set forth herein, X either or both does not comprise or does not consist of an amino acid sequence set forth in US Patent Application Publication No. US2013/0172272.

**[00509]** In certain embodiments of any of the peptide inhibitors described herein, the peptide inhibitor, or each monomer subunit thereof, comprises or consists of at least 3, at least 4 at least 5, at least 6, or at least 7 amino acid residues carboxy terminal of the X9 amino acid residue. In particular embodiments of any of the peptide inhibitors described herein, the peptide inhibitor comprises 3 to 11, 3 to 10, 3 to 9, 3 to 8, 3 to 7, 3 to 6, 3 to 5, 3 to 4, 3, 4, 5, 6, 7, 8, 9, 10, or 11 amino acid residues carboxy terminal of the X9 amino acid residue.

**[00510]** In certain embodiments of any of the peptide inhibitors described herein, the peptide inhibitor, or each monomer subunit thereof, comprises or consists of 4 amino acid residues between X4 and X9. In one embodiment, both X4 and X9 are cysteines.

**[00511]** In certain embodiments, a peptide inhibitor of any of the Formulae described herein comprises the amino acid residues or moieties indicated as X4-X15. In particular embodiments, the peptide inhibitor does not include X1-X3 or X16-X20. In certain embodiments, the peptide inhibitors include an N-terminal extension of one to three amino acid residues corresponding to any of X1-X3. In particular embodiments, any one or more of X1, X2 and X3, when present, are a D-amino acid. In certain embodiments, the peptide inhibitors include an C-terminal extension of one to five amino acid residues corresponding to any of X16-X20. In particular embodiments, any one or more of X16, X17, X18, X19 and X20, when present, are a D-amino acid. Illustrative amino acid residues that may be present in the N-terminal and/or C-terminal extensions are shown in Tables 3 and 5. These tables each show a first peptide inhibitor, with derivates thereof comprising N-terminal extensions, C-terminal extensions, and/or conjugated moieties. The present invention includes derivatives of any of the peptide inhibitors described herein comprising one or more such N-terminal extension, C-terminal extension, and/or

conjugated moiety. In certain embodiments, any of the amino acid residues shown in the extended positions in Tables 3 and 5 may be present in any combination in a peptide inhibitor of the present invention. In particular embodiments, the N-terminal and/or C-terminal extensions are associated with an increased half-life, e.g., upon administration to a subject.

**[00512]** In certain embodiments of any of the peptide inhibitors described herein, the peptide inhibitor, or each monomer subunit thereof, comprises the amino acid sequence motif, W-X-X-Y-W, e.g., at positions X7-X11. In certain embodiments, the peptide inhibitor, or each monomer subunit thereof, comprises the amino acid sequence motif, C-X-X-W-X-C-Y-W, e.g., at positions X4-X11. In certain embodiments, the peptide inhibitor, or each monomer subunit thereof, comprises the amino acid sequence motif, Pen-X-X-W-X-Pen-Y-W, e.g., at positions X4-X11. In certain embodiments of any of the peptide inhibitors described herein, the peptide inhibitor, or both monomer subunit thereof, does not comprise the amino acid sequence motif, W-X-X-Y-W, e.g., at positions X7-X11, where X is any amino acid.

**[00513]** In certain embodiments of any of the Formula or peptide inhibitors described herein, the peptide inhibitor comprises one or more amino acid residues N-terminal to X4. In particular embodiments, X3 is present. In certain embodiments, X3 is Glu, (D)Glu, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, or (D)Gln. In certain embodiments, X3 is (D)Arg or (D)Phe.

**[00514]** In particular embodiments of any of the Formula or peptide inhibitors described herein, the peptide inhibitor comprises an amino acid at X2. In particular embodiments, X2 is Glu, (D)Asp, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln, or (D)Asn. In certain embodiments, X2 and X3 are present. In particular embodiments, X2 is Glu, (D)Asp, Arg, (D)Arg, Phe, (D)Phe, 2-Nal, Thr, Leu, (D)Gln, or (D)As, and X3 is (D)Arg.

**[00515]** In certain embodiments, a peptide inhibitor of the present invention, or one or both monomer subunits thereof, comprises, optionally at its C-terminus, one of the following amino acid sequences:

ENG;

ENN;

[4-amino-4-carboxy-tetrahydropyran]-ENN;

[Lys(Ac)]-NN;  
[ $\alpha$ -MeLys]-ENG;  
[ $\alpha$ -MeLys]- [Lys(Ac)]-NN;  
[ $\alpha$ -MeLeu]- [Lys(Ac)]-NN  
[ $\alpha$ -MeLeu]-ENG;  
[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG;  
[ $\alpha$ -MeLeu]-ENG;  
Aib-[Lys(Ac)]-NG;  
Aib-[Lys(Ac)]-NN;  
NG-[AEA]-[(D)-Lys];  
[Dapa]-NG-[AEA]-[(D)-Lys];  
[Orn]-NG-[AEA]-[(D)-Lys];  
[ $\alpha$ -MeLys]-ENN;  
[4-amino-4-carboxy-tetrahydropyran]- [Lys(Ac)]-NN;  
[Achc]-[Lys(Ac)]-NN; or  
[Acpc]-[Lys(Ac)]-NN.

**[00516]** In particular embodiments, one of these amino acid sequences constitutes the terminal C-terminal amino acids of the peptide. In particular embodiment, these amino acid sequences correspond to X13-X15 or X12-X15 or X14-X16 or X13-X17.

**[00517]** In certain embodiments, a peptide inhibitor of the present invention, or one or both monomer subunits thereof, comprises, optionally at its C-terminus, one of the following amino acid sequences:

WQCY-[2-Nal]-[ $\alpha$ -MeLys];  
WQC-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys];

WQC-[Phe(4-OMe)]-[2-Nal]-[Aib];

WQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys];

W-Xaa8-C-Phe[4-(2-aminoethoxy)]-[2-Nal];

W-Xaa8-C-Phe[4-(2-aminoethoxy)]-[1-Nal];

W-Xaa8-C-Phe[4-(2-aminoethoxy)]; or

**[00518]** W-Xaa8-C-[Phe(4-OCH<sub>3</sub>)]. In particular embodiments, one of these amino acid sequences constitutes the terminal C-terminal amino acids of the peptide. In particular embodiment, these amino acid sequences correspond to X7 to X12 or X7 to X11 or X7 to X10.

**[00519]** In certain embodiments of any of the peptide inhibitors described herein, including both peptide monomer inhibitors and monomer subunits of peptide dimer inhibitors, the peptide monomer inhibitor or monomer subunit is cyclized via a peptide bond between its N-terminal amino acid residue and its C-terminal amino acid residue. In particular embodiments, the peptide inhibitor (or monomer subunit thereof) comprises both an intramolecular bond between X4 and X9 and a peptide bond between its N-terminal amino acid residue and its C-terminal amino acid residue. In certain embodiments, the intramolecular bond is any of those described herein, e.g., a disulfide bond or a thioether bond.

**[00520]** In certain embodiments, the present invention includes a peptide inhibitor that comprises a core consensus sequence selected from one of the following (shown in N-terminal to C-terminal direction):

X1-X2-X3-Pen-X5-X6-W-X8-Pen-X10-X11-X12-X13-X14-X15;

Pen-X5-X6-W-Q-Pen;

Pen-X5-X6-W-X8-Pen;

Pen-X5-X6-W-X8-Pen-[Phe(4-CONH<sub>2</sub>)]; and

Pen-X5-X6-W-X8-Pen-[Phe[4-(2-aminoethoxy)]],

wherein the Pen residues arejoined by an intramolecular bond, e.g., disulphide bond. X1, X2, X3, X5, X6, X8, X10, X11, X12, X13, X14, and X15 may be any amino acid. In some embodiment X5 is Arg, Asn, Gln, Dap, Orn; X6 is Thr or Ser; and X8 is Gln, Val, Phe, Glu, Lys. In particular embodiments, X1, X2, X3, X5, X6, X8, X10, X11, X12, X13, X14, and X15 are defined as described in any of the various Formulas and peptide inhibitors described herein.

[00521]In certain embodiments, the present invention includes a peptide inhibitor that comprises a core consensus sequence selected from one of the following (shown in N-terminal to C-terminal direction):

X1-X2-X3-Abu-X5-X6-W-X8-C-X9-X10-X11-X12-X13-X14-X15;

Abu-X5-X6-W-Q-C;

Abu-X5-X6-W-X8-C;

Abu-X5-X6-W-X8-C-[Phe(4-CONH2)]; and

Abu-X5-X6-W-X8-C-[Phe[4-(2-aminoethoxy)]],

where Abu and C are linked through a intra moleculer thioether bond. X1, X2, X3, X5, X6, X8, X10, X11, X12, X13, X14, and X15 may be any amino acid. In some embodiment X5 is Arg, Asn, Gln, Dap, Orn; X6 is Thr or Ser; and X8 is Gln, Val, Phe, Glu, Lys. In particular embodiments, X1, X2, X3, X5, X6, X8, X10, X11, X12, X13, X14, and X15 are defined as described in any of the various Formulas and peptide inhibitors described herein.

[00522]In certain embodiments, any of the peptide inhibitors described herein may be further cyclized via a peptide bond between its N-terminal amino acid residue and its C-terminal amino acid residue. In particular embodiments, the peptide inhibitor comprises a peptide bond between X3 or X4 and any one of X9, X10, X11, X12, X13, X14, X15, X16, X17, X18, X19 or X20. In particular embodiments, peptide inhibitors of the present invention comprise a peptide bond between their N-terminal and C-terminal amino acid residues, and they also comprise an intramolecular bond between X4 and X9. In certain embodiments, the intramolecular bond is a disulfide bond, a thioether bond, a lactam bond or any of the other bonds described herein.

### Peptide Dimers

**[00523]** In certain embodiments, the present invention includes dimers of the monomer peptide inhibitors described herein, including dimers of any of the monomer peptide inhibitors described herein or in the accompanying tables, figures or sequences listing. These dimers fall within the scope of the general term “peptide inhibitors” as used herein. Illustrative dimers of the present invention are also shown in the accompanying tables, which indicate the dimerized monomer subunits in brackets followed by the linker. Unless otherwise indicated, the subunits are linked via their C-termini. The term “dimer,” as in a peptide dimer, refers to compounds in which two peptide monomer subunits are linked. A peptide dimer inhibitor of the present invention may comprise two identical monomer subunits, resulting in a homodimer, or two non-identical monomer subunits, resulting in a heterodimer. A cysteine dimer comprises two peptide monomer subunits linked through a disulfide bond between a cysteine residue in one monomer subunit and a cysteine residue in the other monomer subunit.

**[00524]** In some embodiments, the peptide inhibitors of the present invention may be active in a dimer conformation, in particular when free cysteine residues are present in the peptide. In certain embodiments, this occurs either as a synthesized dimer or, in particular, when a free cysteine monomer peptide is present and under oxidizing conditions, dimerizes. In some embodiments, the dimer is a homodimer. In other embodiments, the dimer is a heterodimer.

**[00525]** In certain embodiments, a peptide dimer inhibitor of the present invention is a peptide dimer comprising two peptide inhibitors of the invention, including but not limited to a homodimer or heterodimer comprising any of the peptide sequences shown herein, e.g., in Tables 3A-3H, 4A, 4B, 5A-5C, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15.

**[00526]** Certain amino acid sequences listed in Tables 3A-3H, 4A, 4B, 5A-5C, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 are shown using one letter codes for amino acids. Where only the monomer peptide inhibitor sequences are shown; however it is understood that, in certain embodiments, these monomer peptide inhibitors, i.e., monomer subunits, are dimerized to form peptide dimer inhibitors, in accordance with the present teaching and as shown generally, e.g., in Tables 3A-3H, 4A, 4B, 5A-5C, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15.

[00527] In certain embodiments, monomer subunits of the present invention may be dimerized by a suitable linking moiety, e.g., a disulphide bridge between two cysteine residues, one in each peptide monomer subunit, or by another suitable linker moiety, including but not limited to those defined herein. Some of the monomer subunits are shown having C- and N-termini that both comprise free amine. Thus, to produce a peptide dimer inhibitor, the monomer subunit may be modified to eliminate either the C- or N-terminal free amine, thereby permitting dimerization at the remaining free amine. Further, in some instances, a terminal end of one or more monomer subunits is acylated with an acylating organic compound selected from the group consisting of: Trifluoropentyl, Acetyl, Octonyl, Butyl, Pentyl, Hexyl, Palmityl, Trifluoromethyl butyric, cyclopentane carboxylic, cyclopropylacetic, 4-fluorobenzoic, 4-fluorophenyl acetic, 3-Phenylpropionic, tetrahedro-2H-pyran-4carboxylic, succinic acid, and glutaric acid. In some instances, monomer subunits comprise both a free carboxy terminal and a free amino terminal, whereby a user may selectively modify the subunit to achieve dimerization at a desired terminus. One having skill in the art therefore, will appreciate that the monomer subunits of the instant invention may be selectively modified to achieve a single, specific amine for a desired dimerization.







[00528] It is further understood that the C-terminal residues of the monomer subunits disclosed herein are optionally amides. Further, it is understood that, in certain embodiments, dimerization at the C-terminus is facilitated by using a suitable amino acid with a side chain having amine functionality, as is generally understood in the art. Regarding the N-terminal residues, it is generally understood that dimerization may be achieved through the free amine of the terminal residue, or may be achieved by using a suitable amino acid side chain having a free amine, as is generally understood in the art.








[00529] The linker moieties connecting monomer subunits may include any structure, length, and/or size that is compatible with the teachings herein. In at least one embodiment, a linker moiety is selected from the non-limiting group consisting of cysteine, lysine, DIG, PEG4, PEG4-biotin, PEG13, PEG25, PEG1K, PEG2K, PEG3.4K, PEG4K, PEG5K, IDA, ADA, Boc-IDA, Glutaric acid, Isophthalic acid, 1,3-phenylenediacetic acid, 1,4-phenylenediacetic acid, 1,2-phenylenediacetic acid, Triazine, Boc-Triazine, IDA-biotin, PEG4-Biotin, AADA, suitable aliphatics, aromatics, heteroaromatics, and polyethylene glycol based linkers having a molecular





weight from approximately 400Da to approximately 40,000Da. Non-limiting examples of suitable linker moieties are provided in Table 2A.

Table 2A. Illustrative Linker Moieties

Abbrivation	Discription	Structure
DIG	DIGlycolic acid,	
PEG4	Bifunctional PEG linker with 4 PolyEthylene Glycol units	
PEG13	Bifunctional PEG linker with 13 PolyEthylene Glycol units	
PEG25	Bifunctional PEG linker with 25 PolyEthylene Glycol units	
PEG1K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 1000Da	
PEG2K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 2000Da	
PEG3.4K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 3400Da	
PEG5K	Bifunctional PEG linker with PolyEthylene Glycol Mol wt of 5000Da	
DIG	DIGlycolic acid	

$\beta$ -Ala-IDA	$\beta$ -Ala-Iminodiacetic acid	
Boc- $\beta$ - Ala-IDA	Boc- $\beta$ -Ala-Iminodiacetic acid	
Ac- $\beta$ -Ala - IDA	Ac- $\beta$ -Ala-Iminodiacetic acid	
IDA- $\beta$ -Ala- Palm	Palmityl- $\beta$ -Ala-Iminodiacetic acid	
GTA	Glutaric acid	
PMA	Pemilic acid	


AZA	Azelaic acid	
DDA	Dodecanedioic acid	
IPA	Isophthalic acid	
1,3-PDA	1,3- Phenylenediacetic acid	
1,4-PDA	1,4- Phenylenediacetic acid	
1,2-PDA	1,2 - Phenylenediacetic acid	
Triazine	Amino propyl Triazine di-acid	

Boc-Triazine	Boc-Triazine di-acid	
ADA	Amino diacetic acid (which may also referred to as Iminodiacetic acid)	
AADA	n-Acetyl amino acetic acid (which may also referred to as N-acetyl Iminodiacetic acid)	
PEG4-Biotin	PEG4-Biotin (Product number 10199, QuantaBioDesign)	
IDA-Biotin	N-Biotin-β-Ala-Iminodiacetic acid	

Lys	Lysine	<chem>NC(CCO)C(=O)O</chem>
-----	--------	----------------------------

[00530] In some embodiments, a peptide dimer inhibitor is dimerized via a linker moiety. In some embodiments, a peptide dimer inhibitor is dimerized via an intermolecular disulfide bond formed between two cysteine residues, one in each monomer subunit. In some embodiments, a peptide dimer inhibitor is dimerized via both a linker moiety and an intermolecular disulfide bond formed between two cysteine residues. In some embodiments, the intramolecular bond is a thioether, lactam, triazole, selenoether, diselenide or olefin, instead of the disulfide bond.

[00531] An illustrative diagram of one embodiment of a dimer is shown below:



[00532] One having skill in the art will appreciate that the linker (e.g., C- and N-terminal linker) moieties disclosed herein are non-limiting examples of suitable, and that the present invention may include any suitable linker moiety. Thus, some embodiments of the present invention comprises a homo- or heterodimer peptide inhibitor comprised of two monomer subunits selected from the peptides shown in any of tables herein or comprising or consisting of a sequence presented in any of tables herein, wherein the C- or N-termini of the respective monomer subunits (or internal amino acid residues) are linked by any suitable linker moiety to provide a dimer peptide inhibitor having IL-23R inhibitory activity. In certain embodiments, a linker binds to the N- or C-terminus of one monomer subunit and an internal amino acid residue of the other monomer subunit making up the dimer. In certain embodiments, a linker binds to an internal amino acid residue of one monomer subunit and an internal amino acid residue of the other monomer subunit making up the dimer. In further embodiments, a linker binds to the N- or C-terminus of both subunits.

[00533] In particular embodiments, a peptide inhibitor of the present invention comprise two or more polypeptide sequences of monomer peptide inhibitors described herein.


[00534] In one embodiment, a peptide dimer inhibitor of the present invention comprises two peptide monomer subunits connected via one or more linker moieties, wherein each peptide monomer subunit comprises or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 20 amino acid residues, 8 to 20 amino acid residues, 9 to 20 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues and comprises the sequence of Formula Ia, as described herein.

[00535] In particular embodiments, one or both of the monomer subunits comprise the sequence of any one of Formula Formula X, Formula I, II, III, IV or V as described herein.

[00536] In certain embodiments, a peptide dimer inhibitor comprises two peptide monomer subunits connected via one or more linker moieties, wherein each peptide monomer subunit is 8-20 amino acids in length and comprises a sequence of any one of the formulas describd herein, e.g., Formula X, Formula I, II, III, IV or V. In certain embodiments, a peptide dimer inhibitor

comprises two peptide monomer subunits connected via one or more linker moieties, wherein each peptide monomer subunit is 8-20 amino acids in length and comprises a sequence of any one of Formula X, Formula I, II, III, IV or V.

[00537] In certain embodiments, a peptide dimer inhibitor has the structure of Formula VI:



[00538] or a pharmaceutically acceptable salt or solvate thereof,

[00539] wherein each  $R^1$  is independently absent, a bond (e.g., a covalent bond), or  $R^1$  is selected from hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing;

[00540] each  $R^2$  is independently absent, a bond (e.g., a covalent bond), or selected from OH or  $NH_2$ ;

[00541]  $L$  is a linker moiety; and

[00542] each  $X$  is an independently selected peptide monomer subunit comprising or consisting of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 20 amino acid residues, 8 to 20 amino acid residues, 9 to 20 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues amino acids in length, each comprising or consisting of the sequence of Formula Ia, as described herein. In particular embodiments, each peptide monomer subunit comprises or consists of a sequence of Formula Ix, Ia, Ib, Ic, Id, Ie, If, Ig, Ih, Ii, Ij, Ik, Il, Im, In, Io, Ip, Iq, Iq', Is, It, IIa, IIb, IIc, IIId, IIIa, IIIb, IIIc, IIId, IIIe, IVa, IVb, or Va-Vh as described herein.

[00543] In certain embodiments, one or both peptide monomer subunit of a peptide dimer inhibitor is cyclized, e.g., via an intramolecular bond between  $X_4$  and  $X_9$ . In certain embodiments wherein both peptide monomer subunits are cyclized, the intramolecular bond may be the same or different between the two peptide monomer subunits. In certain embodiments, one

or both intramolecular bond is a disulfide bond, a thioether bond, a lactam bond, a selenoether, diselenide, or an olefin bond.

[00544] In one embodiment, X4 and X9 of the one or both cyclized peptide monomer subunit is independently selected from Cys, Pen, hCys, D-Pen, D-Cys and D-hCys, and the intramolecular bond is a disulfide bond.

[00545] In one embodiment, X4 and X9 of the one or both cyclized peptide monomer subunit is independently selected from Glu, Asp, Lys, Orn, Dap, Dab, D-Dap, D-Dab, D-Asp, D-Glu and D-Lys, and the intramolecular bond is a lactam bond.

[00546] In one embodiment, X4 and X9 of the one or both cyclized peptide monomer subunit are each independently selected from  $\beta$ -azido-Ala-OH, propargylglycine, and the peptide dimer inhibitor is cyclized through a triazole ring. In one embodiment, X4 and X9 of the one or both cyclized peptide monomer subunit are each independently selected from 2-allylglycine, 2-(3'-butenyl)glycine, 2-(4'-pentenyl)glycine, 2-(5'-hexenyl)glycine, and the peptide dimer inhibitor is cyclized via a ring closing methasis to give the corresponding olefins / 'stapled peptides'.

[00547] In one embodiment, X4 of one or both cyclized peptide monomer subunit is 2-chloromethylbenzoic acid, mercapto-propanoic acid, mercapto-butyric acid, 2-chloro-acetic acid, 3-chloro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, or hSer(Cl), X9 of one or both cyclized peptide monomer subunit is hSer(Cl), Cys, Pen, hCys, D-Pen, D-Cys or D-hCys, and the intramolecular bond is a thioether bond.

[00548] In one embodiment, X4 of one or both cyclized peptide monomer subunit is 2-chloromethylbenzoic acid, 2-chloro-acetic acid, 3-chloro-propanoic acid, 4-chloro-butyric acid, 3-chloro-isobutyric acid, hSer(Cl), or Sec, X9 of one or both cyclized peptide monomer subunit is hSer(Cl) or Sec, and the intramolecular bond is a selenoether bond.

[00549] In certain embodiments, one or both intramolecular bond is a diselenide bond.

[00550] In certain embodiments, one or both peptide monomer subunits is linear or not cyclized.

[00551] In particular embodiments, of the peptide dimer inhibitors, each X7 and each X11 are both W. In certain embodiments, each X7 and each X11 are both W, each X10 is Y, and each X4 and X9 are both C. In certain embodiments, each X7 and each X11 are both W, each X10 is Y, and each X4 and X9 are amino acids capable of forming an intramolecular bond that is a thioether bond, a lactam bond, a triazole, a selenoether, a diselenide bond, or an olefin bond.

[00552] In certain embodiments of the peptide dimer inhibitors, one or both peptide monomer subunit has a structure shown herein, e.g., in Tables 3A-3I, or comprises an amino acid sequence shown herein, e.g., as set forth in Tables 3A-3I, or wherein the peptide dimer inhibitor has a structure shown herein, e.g., in Table 3F, or comprises an amino acid sequence shown herein, e.g., as set forth in Table 3F.

[00553] In particular embodiments, each R<sup>1</sup> is independently a bond (e.g., a covalent bond), or selected from hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing. In particular embodimetns, the N-terminus of each subunit includes a moiety selected from hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing.

[00554] In certain embodiments of any of the peptide inhibitors having any of the various Formulae set forth herein, each R<sup>1</sup> (or N-terminal moiety) is selected from methyl, acetyl, formyl, benzoyl, trifluoroacetyl, isovaleryl, isobutyryl, octanyl, and the conjugated amides of lauric acid, hexadecanoic acid, and  $\gamma$ -Glu-hexadecanoic acid.

[00555] In particular embodiments, each R<sup>2</sup> (or C-terminal moiety) is independently a bond (e.g., a covalent bond), or selected from OH or NH<sub>2</sub>.

[00556] In particular embodiments of any of the peptide inhibitors having any of the various Formulae set forth herein, each X comprises or consists of 7 to 35 amino acid residues, 8 to 35 amino acid residues, 9 to 35 amino acid residues, 10 to 35 amino acid residues, 7 to 25 amino acid residues, 8 to 25 amino acid residues, 9 to 25 amino acid residues, 10 to 25 amino acid residues, 7 to 18 amino acid residues, 8 to 18 amino acid residues, 9 to 18 amino acid residues, or 10 to 18 amino acid residues.

[00557] In particular embodiments, one or both X comprises or consists of the sequence of any one of the formulas described herein. In certain embodiments of any of the peptide inhibitors, including dimers, or Formulae set forth herein, an X does not comprise or consist of an amino acid sequence set forth in US Patent Application Publication No. US2013/0029907. In certain embodiments of any of the peptide inhibitors, including dimers, or formulas set forth herein, an X does not comprise or consist of an amino acid sequence set forth in US Patent Application Publication No. US2013/0172272.

[00558] In particular embodiments of peptide inhibitors of the present invention (both monomers and dimers) comprising Cys at position X4 and Cys at position X9, the Cys at position X4 and the Cys at position X9 are linked by a disulphide bridge.

[00559] In particular embodiments of peptide inhibitors of the present invention, each X7 and each X11 are not both W.

[00560] In particular embodiments of peptide inhibitors of the present invention, each X7 and each X11 are both W.

[00561] In particular embodiments of peptide inhibitors of the present invention, each X7 and each X11 are both W, X10 is Y, and X4 and X9 are both C.

[00562] In certain embodiments, at least two cysteine residues of the peptide dimer inhibitor are linked by a disulphide bridge, either intramolecular or intermolecular.

[00563] In particular embodiments of either or both monomer subunit (e.g., Ix, Ia-It where permissible) present in a peptide dimer inhibitor, X4 and X9 are both Cys.

[00564] In particular embodiments of either or both monomer subunit (e.g., Ix, Ia-It where permissible) present in a peptide dimer inhibitor, X7 and X11 are both W.

[00565] In particular embodiments of either or both monomer subunit (e.g., Ia-It where permissible) present in a peptide dimer inhibitor, X7 and X11 are both W, X10 is Y, and X4 and X9 are both Cys.

[00566] In particular embodiments of either or both monomer subunit (e.g., Ia-It where permissible) present in a peptide dimer inhibitor, X15 is Gly or Ser.

[00567] In particular embodiments of either or both monomer subunit (e.g., Ia-It where permissible) present in a peptide dimer inhibitor, X16 is AEA or AEP.

[00568] In particular embodiments of either or both monomer subunit (e.g., Ia-It where permissible) present in a peptide dimer inhibitor, X10 is Tyr or Phe, or an analog of Tyr or Phe.

[00569] In particular embodiments of either or both monomer subunit (e.g., Ia-It where permissible) present in a peptide dimer inhibitor, X11 is Trp.

[00570] In particular embodiments of any of the peptide dimer inhibitors described herein, either or both R<sup>1</sup> is hydrogen.

[00571] In particular embodiments of peptide dimer inhibitors of the present invention, the linker moiety (L) is any of the linkers described herein or shown in Table 2A or 2B. In certain embodiments, L is a lysine linker, a diethylene glycol linker, an iminodiacetic acid (IDA) linker, a  $\beta$ -Ala-iminodiacetic acid ( $\beta$ -Ala-IDA) linker, or a PEG linker.

[00572] In various embodiments of any of the peptide dimer inhibitors, each of the peptide monomer subunits is attached to a linker moiety via its N-terminus, C-terminus, or an internal amino acid residue.

[00573] In certain embodiments of any of the peptide dimer inhibitors, the N-terminus of each peptide monomer subunit is connected by a linker moiety.

[00574] In certain embodiments of any of the peptide dimer inhibitors, the C-terminus of each peptide monomer subunit is connected by a linker moiety.

[00575] In certain embodiments of any of the peptide dimer inhibitors, each peptide monomer subunit is connected by a linker moiety attached to an internal amino acid.

[00576] In certain embodiments of peptide dimer inhibitors, the linker moiety is a diethylene glycol linker, an iminodiacetic acid (IDA) linker, a  $\beta$ -Ala-iminodiacetic acid ( $\beta$ -Ala-IDA) linker, or a PEG linker.

[00577] In certain embodiments of the peptide dimer inhibitors, one or both peptide monomer subunit has a structure shown in any of the tables in the Examples or comprises an amino acid sequence set forth in any of the tables in the Examples.

[00578] In certain embodiments of any of the peptide inhibitors, including dimers, or Formulae set forth herein, an X does not comprise or consist of an amino acid sequence set forth in US Patent Application Publication No. US2013/0029907. In certain embodiments of any of the peptide inhibitors, including dimers, or Formulas set forth herein, an X does not comprise or consist of an amino acid sequence set forth in US Patent Application Publication No. US2013/0172272.

[00579] In particular embodiments of peptide inhibitors of the present invention, each X7 and each X11 are both W, X10 is Y, and X4 and X9 are both Pen.

In certain embodiments, at least two cysteine residues of the peptide dimer inhibitor are linked by a disulphide bridge, either intramolecular or intermolecular.

#### Peptide Inhibitor Conjugates and Biopolymers

[00580] In certain embodiments, peptide inhibitors of the present invention, including both monomers and dimers, comprise one or more conjugated chemical substituents, such as lipophilic substituents and polymeric moieties, which may be referred to herein as half-life extension moieties. Without wishing to be bound by any particular theory, it is believed that the lipophilic substituent binds to albumin in the bloodstream, thereby shielding the peptide inhibitor from enzymatic degradation, and thus enhancing its half-life. In addition, it is believed that polymeric moieties enhance half-life and reduce clearance in the bloodstream.


[00581] In additional embodiments, any of the peptide inhibitors, e.g. peptides of Formulas (Va)-(Vh), further comprise a linker moiety attached to an amino acid residue present in the inhibitor, e.g., a linker moiety may be bound to a side chain of any amino acid of the peptide inhibitor, to

the N-terminal amino acid of the peptide inhibitor, or to the C-terminal amino acid of the peptide inhibitor.

[00582] In additional embodiments, any of the peptide inhibitors e.g. peptides of Formulas (Va)-(Vh), further comprise half-life extension moiety attached to an amino acid residue present in the inhibitor, e.g., a half-life extension moiety may be bound to a side chain of any amino acid of the peptide inhibitor, to the N-terminal amino acid of the peptide inhibitor, or to the C-terminal amino acid of the peptide inhibitor.

[00583] In additional embodiments, any of the peptide inhibitors e.g. peptides of Formulas (Va)-(Vh), further comprise half-life extension moiety attached to a linker moiety that is attached to an amino acid residue present in the inhibitor, e.g., a half-life extension moiety may be bound to a linker moiety that is bound to a side chain of any amino acid of the peptide inhibitor, to the N-terminal amino acid of the peptide inhibitor, or to the C-terminal amino acid of the peptide inhibitor.










[00584] In particular embodiments, an IL23R analogue comprises a half-life extension moiety having the structure shown below, wherein  $n=0$  to 24 or  $n=14$  to 24:



[00585] In certain embodiments, a IL23R analogue of the present invention comprises a half-life extension moiety shown in Table 7.

Table 7. Illustrative Half-Life Extension Moieties

#	Half-Life Extension Moieties
C1	 <p>C12 (Lauric acid)</p>


C2		C14 (Mysteric acid)
C3		C16 (Palm or Palmitic acid)
C4		C18 (Stearic acid)
C5		C20
C6		C12 diacid
C7		C14 diacid
C8		C16 diacid
C9		C18 diacid
C10		C20 diacid

[00586] In certain embodiments, a half-life extension moiety is bound directly to a peptide inhibitor, while in other embodiments, a half-life extension moiety is bound to the peptide inhibitor via a linker moiety, e.g., any of those depicted in Tables 6 or 8.

[00587] Table 8. Illustrative Linker Moieties

#	Linker Moiety
L1	<p>IsoGlu</p>
L2	<p>Dapa</p>
L3	<p>Ahx</p>
L4	<p>Lipidic based linkers:</p> <p><math>n=1</math> to 24</p>
L5	<p>PEG1</p>
L6	<p>PEG2</p>

L7	<p>PEG11 (40 atoms) also known as PEG12</p>
L8	<p>n=1 to 25</p> <p>PEG based linkers</p>
L9	<p>OEG</p>
L10	<p>IsoGlu-Ahx</p>
L11	<p>IsoGlu-OEG-OEG</p>
L12	<p>IsoGlu-PEG5</p>
L13	<p>n=1-25</p> <p>IsoGlu-PEGn</p>



[00588] In particular embodiments, a peptide inhibitor of the present invention comprises any of the linker moieties shown in Table 8 and any of the half-life extension moieties shown in Table 7, including any of the following combinations shown in Table 9a.

Table 9a. Illustrative Combinations of Linkers and Half-Life Extension Moieties in Peptide Inhibitors

Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety
L1	C1	L1	C2	L1	C3
L2	C1	L2	C2	L2	C3
L3	C1	L3	C2	L3	C3
L4	C1	L4	C2	L4	C3
L5	C1	L5	C2	L5	C3
L6	C1	L6	C2	L6	C3
L7	C1	L7	C2	L7	C3
L8	C1	L8	C2	L8	C3
L9	C1	L9	C2	L9	C3
L10	C1	L10	C2	L10	C3
L11	C1	L11	C2	L11	C3
L12	C1	L12	C2	L12	C3
L13	C1	L13	C2	L13	C3
L14	C1	L14	C2	L14	C3
L15	C1	L15	C2	L15	C3

Linker	Half-Life Extension	Linker	Half-Life Extension	Linker	Half-Life Extension

	Moiety		Moiety		Moiety
L1	C4	L1	C5	L1	C6
L2	C4	L2	C5	L2	C6
L3	C4	L3	C5	L3	C6
L4	C4	L4	C5	L4	C6
L5	C4	L5	C5	L5	C6
L6	C4	L6	C5	L6	C6
L7	C4	L7	C5	L7	C6
L8	C4	L8	C5	L8	C6
L9	C4	L9	C5	L9	C6
L10	C4	L10	C5	L10	C6
L11	C4	L11	C5	L11	C6
L12	C4	L12	C5	L12	C6
L13	C4	L13	C5	L13	C6
L14	C4	L14	C5	L14	C6
L15	C4	L15	C5	L15	C6

Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety
L1	C7	L1	C8	L1	C9
L2	C7	L2	C8	L2	C9
L3	C7	L3	C8	L3	C9
L4	C7	L4	C8	L4	C9
L5	C7	L5	C8	L5	C9
L6	C7	L6	C8	L6	C9
L7	C7	L7	C8	L7	C9
L8	C7	L8	C8	L8	C9
L9	C7	L9	C8	L9	C9
L10	C7	L10	C8	L10	C9
L11	C7	L11	C8	L11	C9
L12	C7	L12	C8	L12	C9
L13	C7	L13	C8	L13	C9
L14	C7	L14	C8	L14	C9
L15	C7	L15	C8	L15	C9


Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety
L1	C10	L6	C10	L11	C10
L2	C10	L7	C10	L12	C10
L3	C10	L8	C10	L13	C10
L4	C10	L9	C10	L14	C10
L5	C10	L10	C10	L15	C10

[00589] In some embodiments there may be multiple linkers present between the peptide the conjugated moiety, e.g., half-life extension moiety, e.g., as depicted in Table 9b.


Table 9b. Illustrative Combinations of Linkers and Half-Life Extension Moieties in Peptide Inhibitors

Linker	Half-Life Extension Moiety	Linker	Half-Life Extension Moiety
L1-L2	C10	L1-L2	C8
L2-L5-L3	C10	L2-L5-L3	C8
L3-L8	C10	L3-L8	C8
L1-L2-L3	C10	L1-L2-L3	C8
L5-L3-L3-L3	C10	L5-L3-L3-L3	C8


[00590] Illustrative examples of peptide inhibitors of the present invention, including those having a conjugates linker and/or half-life extension moiety are shown below. All amino acids are L amino acids unless otherwise stated. The present invention also includes salt forms of any of these peptide inhibitors, including, but not limited to, acetate salts thereof.




Example 1: cyclo[[Abu]-QTCWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>




Example 1a: Ac-[(D)-Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>




Example 2: Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>




Example 3: cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(Linker-Half-Life Extension Moiety)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>




Example 3a: Ac-[(D)-Arg]cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(Linker-Half-Life Extension Moiety)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>




Example 4: cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Linker-Half-Life Extension Moiety)]-NN-NH<sub>2</sub>




Example 4a: Ac-[(D)-Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Linker-Half-Life Extension Moiety)]-NN-NH<sub>2</sub>




Example 5: cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-[Lys(Linker-Half-Life Extension Moiety)]-NH<sub>2</sub>




Example 5a: Ac[(D)-Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-[Lys(Linker-Half-Life Extension Moiety)]-NH<sub>2</sub>




Example 6: [Half-Life Extension Moiety-Linker]-[cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>




Example 6a: [Half-Life Extension Moiety-Linker]-[(D)-Arg]-[cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>




Example 7: [Half-Life Extension Moiety-Linker]-[Pen]-NTWQ-[Pen]-[Phe[4-(aminoethoxy)]-2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>



Example 8: Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-[Lys(Linker-Half-Life Extension Moity)]-NH<sub>2</sub>



Example 9: Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(aminoethoxy)-(Linker-Half-Life Extension Moity)-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>



Example 10: Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Linker-Half-Life Extension Moiety )]-NN-NH<sub>2</sub>.

[00591] In certain embodiments, the half-life of a peptide inhibitor of the invention that includes a conjugated chemical substituent, i.e., a half-life extension moiety, is at least 100%, at least 120%, at least 150%, at least 200%, at least 250%, at least 300%, at least 400%, or at least 500% of the half-life of the same peptide inhibitor but without the conjugated chemical substituent. In certain embodiments, the lipophilic substituents and/or polypermic moieties enhance the permeability of the peptide inhibitor through the epithelium and/or its retention in the lamina propria. In certain embodiments, the permeability through the epithelium and/or the retention in the lamina propria of a peptide inhibitor of the invention that includes a conjugated chemical substituent is at 100%, at least 120%, at least 150%, at least 200%, at least 250%, at least 300%, at least 400%, or at least 500% of the half-life of the same peptide inhibitor but without the conjugated chemical substituent.

[00592] In one embodiment, a side chain of one or more amino acid residues (e.g., Lys residues) in a peptide inhibitor of the invention is conjugated (e.g., covalently attached) to a lipophilic substituent. The lipophilic substituent may be covalently bonded to an atom in the amino acid side chain, or alternatively may be conjugated to the amino acid side chain via one or more spacers. The spacer, when present, may provide spacing between the peptide analogue and the lipophilic substituent. In particular embodiments, the peptide inhibitor comprises any of the conjugated moieties shown in Tables 2-5.


[00593] In certain embodiments, the lipophilic substituent may comprise a hydrocarbon chain having from 4 to 30 C atoms, for example at least 8 or 12 C atoms, and preferably 24 C atoms or fewer, or 20 C atoms or fewer. The hydrocarbon chain may be linear or branched and may be saturated or unsaturated. In certain embodiments, the hydrocarbon chain is substituted with a moiety which forms part of the attachment to the amino acid side chain or the spacer, for example an acyl group, a sulfonyl group, an N atom, an O atom or an S atom. In some embodiments, the hydrocarbon chain is substituted with an acyl group, and accordingly the hydrocarbon chain may form part of an alkanoyl group, for example palmitoyl, caproyl, lauroyl, myristoyl or stearoyl.


[00594] A lipophilic substituent may be conjugated to any amino acid side chain in a peptide inhibitor of the invention. In certain embodiment, the amino acid side chain includes a carboxy, hydroxyl, thiol, amide or amine group, for forming an ester, a sulphonyl ester, a thioester, an amide or a sulphonamide with the spacer or lipophilic substituent. For example, the lipophilic substituent may be conjugated to Asn, Asp, Glu, Gln, His, Lys, Arg, Ser, Thr, Tyr, Trp, Cys or Dbu, Dpr or Orn. In certain embodiments, the lipophilic substituent is conjugated to Lys. An amino acid shown as Lys in any of the formula provided herein may be replaced by, e.g., Dbu, Dpr or Orn where a lipophilic substituent is added.

[00595] In certain embodiments, the peptide inhibitors of the present invention may be modified, e.g., to enhance stability, increase permeability, or enhance drug like characteristics, through conjugation of a chemical moiety to one or more amino acid side chain within the peptide. For example, the N(epsilon) of lysine N(epsilon), the  $\beta$ -carboxyl of aspartic, or the  $\gamma$ -carboxyl of glutamic acid may be appropriately functionalized. Thus, to produce the modified peptide, an amino acid within the peptide may be appropriately modified. Further, in some instances, the side chain is acylated with an acylating organic compound selected from the group consisting of: Trifluoropentyl, Acetyl, Octonyl, Butyl, Pentyl, Hexyl, Palmityl, Trifluoromethyl butyric, cyclopentane carboxylic, cyclopropylacetic, 4-fluorobenzoic, 4-fluorophenyl acetic, 3-Phenylpropionic, tetrahedro-2H-pyran-4carboxylic, succinic acid glutaric acid or bile acids. One having skill is the art will appreciate that a series of conjugates can be linked, e.g., for example PEG4, isoglu and combinations thereof. One having skill is the art will appreciate that an amino

acid with the peptide can be isosterically replaced, for example, Lys may be replaced for Dap, Dab,  $\alpha$ -MeLys or Orn. Examples of modified residues within a peptide are shown in Table 1B.

Table 1B. Examples of modified Lysine, Asp and Asn within the peptide





[00596] In further embodiments of the present invention, alternatively or additionally, a side-chain of one or more amino acid residues in a peptide inhibitor of the invention is conjugated to a polymeric moiety, for example, in order to increase solubility and/or half-life *in vivo* (e.g. in

plasma) and/or bioavailability. Such modifications are also known to reduce clearance (e.g. renal clearance) of therapeutic proteins and peptides.

[00597] As used herein, “Polyethylene glycol” or “PEG” is a polyether compound of general formula H-(O-CH<sub>2</sub>-CH<sub>2</sub>)<sub>n</sub>-OH. PEGs are also known as polyethylene oxides (PEOs) or polyoxyethylenes (POEs), depending on their molecular weight PEO, PEE, or POG, as used herein, refers to an oligomer or polymer of ethylene oxide. The three names are chemically synonymous, but PEG has tended to refer to oligomers and polymers with a molecular mass below 20,000 Da, PEO to polymers with a molecular mass above 20,000 Da, and POE to a polymer of any molecular mass. PEG and PEO are liquids or low-melting solids, depending on their molecular weights. Throughout this disclosure, the 3 names are used indistinguishably. PEGs are prepared by polymerization of ethylene oxide and are commercially available over a wide range of molecular weights from 300 Da to 10,000,000 Da. While PEG and PEO with different molecular weights find use in different applications, and have different physical properties (e.g. viscosity) due to chain length effects, their chemical properties are nearly identical. The polymeric moiety is preferably water-soluble (amphiphilic or hydrophilic), non-toxic, and pharmaceutically inert. Suitable polymeric moieties include polyethylene glycols (PEG), homo- or co-polymers of PEG, a monomethyl-substituted polymer of PEG (mPEG), or polyoxyethylene glycerol (POG). See, for example, Int. J. Hematology 68:1 (1998); Bioconjugate Chem. 6:150 (1995); and Crit. Rev. Therap. Drug Carrier Sys. 9:249 (1992). Also encompassed are PEGs that are prepared for purpose of half life extension, for example, mono-activated, alkoxy-terminated polyalkylene oxides (POA's) such as mono-methoxy-terminated polyethylene glycols (mPEG's); bis activated polyethylene oxides (glycols) or other PEG derivatives are also contemplated. Suitable polymers will vary substantially by weights ranging from about 200 Da to about 40,000 Da or from about 200 Da to about 60,000 Da are usually selected for the purposes of the present invention. In certain embodiments, PEGs having molecular weights from 200 to 2,000 or from 200 to 500 are used. Different forms of PEG may also be used, depending on the initiator used for the polymerization process – a common common initiator is a monofunctional methyl ether PEG, or methoxypoly(ethylene glycol), abbreviated mPEG.

[00598] Lower-molecular-weight PEGs are also available as pure oligomers, referred to as monodisperse, uniform, or discrete. These are used in certain embodiments of the present invention.

[00599] PEGs are also available with different geometries: branched PEGs have three to ten PEG chains emanating from a central core group; star PEGs have 10 to 100 PEG chains emanating from a central core group; and comb PEGs have multiple PEG chains normally grafted onto a polymer backbone. PEGs can also be linear. The numbers that are often included in the names of PEGs indicate their average molecular weights (e.g. a PEG with n = 9 would have an average molecular weight of approximately 400 daltons, and would be labeled PEG 400).

[00600] As used herein, “PEGylation” is the act of covalently coupling a PEG structure to the peptide inhibitor of the invention, which is then referred to as a “PEGylated peptide inhibitor”. In certain embodiments, the PEG of the PEGylated side chain is a PEG with a molecular weight from about 200 to about 40,000. In some embodiments, a spacer of a peptide of formula I, formula I’, or formula I’’ is PEGylated. In certain embodiments, the PEG of a PEGylated spacer is PEG3, PEG4, PEG5, PEG6, PEG7, PEG8, PEG9, PEG10, or PEG11. In certain embodiments, the PEG of a PEGylated spacer is PEG3 or PEG8.

[00601] Other suitable polymeric moieties include poly-amino acids such as poly-lysine, poly-aspartic acid and poly-glutamic acid (see for example Gombotz, et al. (1995), Bioconjugate Chem., vol. 6: 332-351; Hudecz, et al. (1992), Bioconjugate Chem., vol. 3, 49-57 and Tsukada, et al. (1984), J. Natl. Cancer Inst., vol. 73, : 721-729. The polymeric moiety may be straight-chain or branched. In some embodiments, it has a molecular weight of 500-40,000 Da, for example 500-10,000 Da, 1000-5000 Da, 10,000-20,000 Da, or 20,000-40,000 Da.

[00602] In some embodiments, a peptide inhibitor of the invention may comprise two or more such polymeric moieties, in which case the total molecular weight of all such moieties will generally fall within the ranges provided above.

[00603] In some embodiments, the polymeric moiety is coupled (by covalent linkage) to an amino, carboxyl or thiol group of an amino acid side chain. Certain examples are the thiol group

of Cys residues and the epsilon amino group of Lys residues, and the carboxyl groups of Asp and Glu residues may also be involved.

[00604] The skilled worker will be well aware of suitable techniques which can be used to perform the coupling reaction. For example, a PEG moiety bearing a methoxy group can be coupled to a Cys thiol group by a maleimido linkage using reagents commercially available from Nektar Therapeutics AL. See also WO 2008/101017, and the references cited above, for details of suitable chemistry. A maleimide-functionalised PEG may also be conjugated to the side-chain sulfhydryl group of a Cys residue.

[00605] As used herein, disulfide bond oxidation can occur within a single step or is a two step process. As used herein, for a single oxidation step, the trityl protecting group is often employed during assembly, allowing deprotection during cleavage, followed by solution oxidation. When a second disulfide bond is required, one has the option of native or selective oxidation. For selective oxidation requiring orthogonal protecting groups, Acm and Trityl is used as the protecting groups for cysteine. Cleavage results in the removal of one protecting pair of cysteine allowing oxidation of this pair. The second oxidative deprotection step of the cysteine protected Acm group is then performed. For native oxidation, the trityl protecting group is used for all cysteines, allowing for natural folding of the peptide. A skilled worker will be well aware of suitable techniques which can be used to perform the oxidation step.

[00606] Several chemical moieties, including poly(ethylene)glycol, react with functional groups present in the twenty naturally occurring amino acids, such as, for example, the epsilon amino group in lysine amino acid residues, the thiol present in cysteine amino acid residues, or other nucleophilic amino acid side chains. When multiple naturally occurring amino acids react in a peptide inhibitor, these non-specific chemical reactions result in a final peptide inhibitor that contains many isomers of peptides conjugated to one or more poly(ethylene)glycol strands at different locations within the peptide inhibitor.

[00607] One advantage of certain embodiments of the present invention includes the ability to add one or more chemical moiety (such as PEG) by incorporating one or more non-natural amino acid(s) that possess unique functional groups that react with an activated PEG by way of chemistry that is unreactive with the naturally occurring amino acids present in the peptide

inhibitor. For example, azide and alkyne groups are unreactive with all naturally occurring functional groups in a protein. Thus, a non-natural amino acid may be incorporated in one or more specific sites in a peptide inhibitor where PEG or another modification is desired without the undesirable non-specific reactions. In certain embodiments, the particular chemistry involved in the reaction results in a stable, covalent link between the PEG strand and the peptide inhibitor. In addition, such reactions may be performed in mild aqueous conditions that are not damaging to most peptides. In certain embodiments, the non-natural amino acid residue is AHA.

**[00608]** Chemical moieties attached to natural amino acids are limited in number and scope. By contrast, chemical moieties attached to non-natural amino acids can utilize a significantly greater spectrum of useful chemistries by which to attach the chemical moiety to the target molecule. Essentially any target molecule, including any protein (or portion thereof) that includes a non-natural amino acid, e.g., a non-natural amino acid containing a reactive site or side chain where a chemical moiety may attach, such as an aldehyde- or keto-derivatized amino acid, can serve as a substrate for attaching a chemical moiety.

**[00609]** Numerous chemical moieties may be joined or linked to a particular molecule through various known methods in the art. A variety of such methods are described in U.S. Patent No. 8,568,706. As an illustrative example, azide moieties may be useful in conjugating chemical moieties such as PEG or others described herein. The azide moiety serves as a reactive functional group, and is absent in most naturally occurring compounds (thus it is unreactive with the native amino acids of naturally occurring compounds). Azides also undergo a selective ligation with a limited number of reaction partners, and azides are small and can be introduced to biological samples without altering the molecular size of significantly. One reaction that allows incorporation or introduction of azides to molecules is the copper-mediated Huisgen [3+2] cycloaddition of an azide. This reaction can be used for the selective PEGylation of peptide inhibitors. (Tornoe et al., *J. Org. Chem.* 67: 3057, 2002; Rostovtsev et al., *Angew. Chem., Int. Ed.* 41: 596, 2002; and Wang et al., *J. Am. Chem. Soc.* 125: 3192, 2003, Speers et al., *J. Am. Chem. Soc.*, 2003, 125, 4686).

Illustrative Peptide Inhibitors and Peptide Dimer Inhibitors, and Methods of Making the Same

[00610] The present invention thus provides various peptide inhibitors which bind or associate with IL-23, to disrupt or block binding between IL-23 and IL-23R.

[00611] Illustrative peptide inhibitors and peptide dimer inhibitors of the present invention are shown in Tables 3A-3H, 4A, 4B, 5A-5C, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 provides the amino acid sequence of selected monomer peptide inhibitors and peptide dimer inhibitors, and indicates the linker moiety present in the peptide dimer inhibitors. According to the protocols discussed herein, a number of the peptide inhibitors and peptide dimer inhibitors shown in the accompanying tables were synthesized and cyclized. Tables E3A-E3H, E4A, E4B, E5A-E5C, E6, E7, E8, E9, E10, E11, E12, E13, E14 or E15 provide the IC<sub>50</sub> values for selected monomer peptide inhibitors and peptide dimer inhibitors in inhibiting IL-23 binding to the IL-23R, or in inhibiting IL-23 signaling as determined by measuring changes in phospho-STAT3 levels, as described in the accompanying Examples. Illustrative peptide inhibitors of the present invention are shown in Formulas (V), and in Tables 2-5, which provide the amino acid sequence of selected peptide inhibitors. These peptide inhibitors are acetate salts.

[00612] The peptide inhibitors of the present invention may be synthesized by many techniques that are known to those skilled in the art. In certain embodiments, monomer subunits are synthesized, purified, and dimerized using the techniques described in the accompanying Examples. In certain embodiments, the present invention provides a method of producing a peptide inhibitor (or monomer subunit thereof) of the present invention, comprising chemically synthesizing a peptide comprising, consisting of, or consisting essentially of a peptide having an amino acid sequence described herein, including but not limited to any of the amino acid sequences set forth in any of Formulas I, II, III, IV, V or VI or tables herein. In other embodiments, the peptide is recombinantly synthesized, instead of being chemically synthesized. In certain embodiments, the peptide inhibitor is a dimer, and the method comprises synthesizing both monomer subunits of the peptide dimer inhibitor and then dimerizing the two monomer subunits to produce the peptide dimer inhibitor. In various embodiments, dimerization is accomplished via any of the various methods described herein. In particular embodiments, methods of producing a peptide inhibitor (or monomer subunit thereof) further comprise cyclizing the peptide inhibitor (or monomer subunit thereof) after its synthesis. In particular

embodiments, cyclization is accomplished via any of the various methods described herein. In certain embodiments, the present invention provides a method of producing a peptide inhibitor (or monomer subunit thereof) of the present invention, comprising introducing an intramolecular bond, e.g., a disulfide, an amide, or a thioether bond between two amino acids residues within a peptide comprising, consisting of, or consisting essentially of a peptide having an amino acid sequence described herein, including but not limited to any of the amino acid sequences set forth in any of Formulas I, II, III, IV, V or VI, or the accompanying Examples, Tables, or Sequence Listing.

**[00613]** In related embodiments, the present invention includes polynucleotides that encode a polypeptide having a sequence set forth in any one of Formulas I, II, III, IV, V or VI, or the accompanying Examples, Tables, or Sequence Listing.

**[00614]** In addition, the present invention includes vectors, e.g., expression vectors, comprising a polynucleotide of the present invention.

#### Methods of Treatment

**[00615]** In certain embodiments, the present invention includes methods of inhibiting IL-23 binding to an IL-23R on a cell, comprising contacting the IL-23 with a peptide inhibitor of the present invention. In certain embodiments, the cell is a mammalian cell. In particular embodiments, the method is performed in vitro or in vivo. Inhibition of binding may be determined by a variety of routine experimental methods and assays known in the art.

**[00616]** In certain embodiments, the present invention includes methods of inhibiting IL-23 signaling by a cell, comprising contacting the IL-23 with a peptide inhibitor of the present invention. In certain embodiments, the cell is a mammalian cell. In particular embodiments, the method is performed in vitro or in vivo. In particular embodiments, the inhibition of IL-23 signalling may be determined by measuring changes in phospho-STAT3 levels in the cell.

**[00617]** In some embodiments, the present invention provides methods for treating a subject afflicted with a condition or indication associated with IL-21 or IL-23R (e.g., activation of the IL-23/IL-23R signaling pathway), wherein the method comprises administering to the subject a peptide inhibitor of the present invention. In one embodiment, a method is provided for treating

a subject afflicted with a condition or indication characterized by inappropriate, deregulated, or increased IL-23 or IL-23R activity or signaling, comprising administering to the individual a peptide inhibitor of the present invention in an amount sufficient to inhibit (partially or fully) binding of IL-23 to IL-23R in the subject. In particular embodiments, the inhibition of IL-23 binding to IL-23R occurs in particular organs or tissues of the subject, e.g., the stomach, small intestine, large intestine/colon, intestinal mucosa, lamina propria, Peyer's Patches, mesenteric lymph nodes, or lymphatic ducts.

**[00618]** In some embodiments, methods of the present invention comprise providing a peptide inhibitor of the present invention to a subject in need thereof. In particular embodiments, the subject in need thereof has been diagnosed with or has been determined to be at risk of developing a disease or disorder associated with IL-23/IL-23R. In particular embodiments, the subject is a mammal.

**[00619]** In certain embodiments, the disease or disorder is autoimmune inflammation and related diseases and disorders, such as multiple sclerosis, asthma, rheumatoid arthritis, inflammatory bowel diseases (IBDs), juvenile IBD, adolescent IBD, Crohn's disease, sarcoidosis, Systemic Lupus Erythematosus, ankylosing spondylitis (axial spondyloarthritis), psoriatic arthritis, or psoriasis. In particular embodiments, the disease or disorder is psoriasis (e.g., plaque psoriasis, guttate psoriasis, inverse psoriasis, pustular psoriasis, Palmo-Plantar Pustulosis, psoriasis vulgaris, or erythrodermic psoriasis), atopic dermatitis, acne ectopica, ulcerative colitis, Crohn's disease, Celiac disease (nontropical Sprue), enteropathy associated with seronegative arthropathies, microscopic colitis, collagenous colitis, eosinophilic gastroenteritis/esophagitis, colitis associated with radio- or chemo-therapy, colitis associated with disorders of innate immunity as in leukocyte adhesion deficiency-1, chronic granulomatous disease, glycogen storage disease type 1b, Hermansky-Pudlak syndrome, Chediak-Higashi syndrome, Wiskott-Aldrich Syndrome, pouchitis resulting after proctocolectomy and ileoanal anastomosis, gastrointestinal cancer, pancreatitis, insulin-dependent diabetes mellitus, mastitis, cholecystitis, cholangitis, primary biliary cirrhosis, viral-associated enteropathy, pericholangitis, chronic bronchitis, chronic sinusitis, asthma, uveitis, or graft versus host disease.

[00620] In certain related embodiments, the present invention provides a method of selectively inhibiting IL-23 or IL-23R signaling (or the binding of IL-23 to IL-23R) in a subject in need thereof, comprising providing to the subject a peptide inhibitor of the present invention. In particular embodiments, the present invention includes a method of selectively inhibiting IL-23 or IL-23R signaling (or the binding of IL-23 to IL-23R) in the GI tract of a subject in need thereof, comprising providing to the subject a peptide inhibitor of the present invention by oral administration. In particular embodiments, exposure of the administered peptide inhibitor in GI tissues (e.g., small intestine or colon) is at least 10-fold, at least 20-fold, at least 50-fold, or at least 100-fold greater than the exposure in the blood. In particular embodiments, the present invention includes a method of selectively inhibiting IL23 or IL23R signaling (or the binding of IL23 to IL23R) in the GI tract of a subject in need thereof, comprising providing to the subject a peptide inhibitor, wherein the peptide inhibitor does not block the interaction between IL-6 and IL-6R or antagonize the IL-12 signaling pathway. In a further related embodiment, the present invention includes a method of inhibiting GI inflammation and/or neutrophil infiltration to the GI, comprising providing to a subject in need thereof a peptide inhibitor of the present invention. In some embodiments, methods of the present invention comprise providing a peptide inhibitor of the present invention (i.e., a first therapeutic agent) to a subject in need thereof in combination with a second therapeutic agent. In certain embodiments, the second therapeutic agent is provided to the subject before and/or simultaneously with and/or after the peptide inhibitor is administered to the subject. In particular embodiments, the second therapeutic agent is an anti-inflammatory agent. In certain embodiments, the second therapeutic agent is a non-steroidal anti-inflammatory drug, steroid, or immune modulating agent. In another embodiment, the method comprises administering to the subject a third therapeutic agent. In certain embodiments, the second therapeutic agent is an antibody that binds IL-23 or IL-23R.

[00621] In particular embodiments, the peptide inhibitor, or the pharmaceutical composition comprising a peptide inhibitor, is suspended in a sustained-release matrix. A sustained-release matrix, as used herein, is a matrix made of materials, usually polymers, which are degradable by enzymatic or acid-base hydrolysis or by dissolution. Once inserted into the body, the matrix is acted upon by enzymes and body fluids. A sustained-release matrix desirably is chosen from biocompatible materials such as liposomes, polylactides (polylactic acid), polyglycolide (polymer of glycolic acid), polylactide co-glycolide (copolymers of lactic acid and glycolic acid)

polyanhydrides, poly(ortho)esters, polypeptides, hyaluronic acid, collagen, chondroitin sulfate, carboxylic acids, fatty acids, phospholipids, polysaccharides, nucleic acids, polyamino acids, amino acids such as phenylalanine, tyrosine, isoleucine, polynucleotides, polyvinyl propylene, polyvinylpyrrolidone and silicone. One embodiment of a biodegradable matrix is a matrix of one of either polylactide, polyglycolide, or polylactide co-glycolide (co-polymers of lactic acid and glycolic acid).

**[00622]** In certain embodiments, the present invention includes pharmaceutical compositions comprising one or more peptide inhibitors of the present invention and a pharmaceutically acceptable carrier, diluent or excipient. A pharmaceutically acceptable carrier, diluent or excipient refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Prevention of the action of microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride, and the like.

**[00623]** In certain embodiments, the compositions are administered orally, parenterally, intracisternally, intravaginally, intraperitoneally, intrarectally, topically (as by powders, ointments, drops, suppository, or transdermal patch), by inhalation (such as intranasal spray), ocularly (such as intraocularly) or buccally. The term “parenteral” as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous, intradermal and intraarticular injection and infusion. Accordingly, in certain embodiments, the compositions are formulated for delivery by any of these routes of administration.

**[00624]** In certain embodiments, pharmaceutical compositions for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders, for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), carboxymethylcellulose and suitable mixtures thereof,  $\beta$ -cyclodextrin, vegetable oils (such as olive oil), and injectable organic esters such as ethyl oleate. Proper fluidity may be

maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants. These compositions may also contain adjuvants such as preservative, wetting agents, emulsifying agents, and dispersing agents. Prolonged absorption of an injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption, such as aluminum monostearate and gelatin.

[00625] Injectable depot forms include those made by forming microencapsule matrices of the peptide inhibitor in one or more biodegradable polymers such as polylactide-polyglycolide, poly(orthoesters), poly(anhydrides), and (poly)glycols, such as PEG. Depending upon the ratio of peptide to polymer and the nature of the particular polymer employed, the rate of release of the peptide inhibitor can be controlled. Depot injectable formulations are also prepared by entrapping the peptide inhibitor in liposomes or microemulsions compatible with body tissues.

[00626] The injectable formulations may be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

[00627] Topical administration includes administration to the skin or mucosa, including surfaces of the lung and eye. Compositions for topical lung administration, including those for inhalation and intranasal, may involve solutions and suspensions in aqueous and non-aqueous formulations and can be prepared as a dry powder which may be pressurized or non-pressurized. In non-pressurized powder compositions, the active ingredient may be finely divided form may be used in admixture with a larger-sized pharmaceutically acceptable inert carrier comprising particles having a size, for example, of up to 100 micrometers in diameter. Suitable inert carriers include sugars such as lactose.

[00628] Alternatively, the composition may be pressurized and contain a compressed gas, such as nitrogen or a liquefied gas propellant. The liquefied propellant medium and indeed the total composition may be such that the active ingredient does not dissolve therein to any substantial extent. The pressurized composition may also contain a surface active agent, such as a liquid or

solid non-ionic surface active agent or may be a solid anionic surface active agent. It is preferred to use the solid anionic surface active agent in the form of a sodium salt.

[00629] A further form of topical administration is to the eye. A peptide inhibitor of the invention may be delivered in a pharmaceutically acceptable ophthalmic vehicle, such that the peptide inhibitor is maintained in contact with the ocular surface for a sufficient time period to allow the peptide inhibitor to penetrate the corneal and internal regions of the eye, as for example the anterior chamber, posterior chamber, vitreous body, aqueous humor, vitreous humor, cornea, iris/ciliary, lens, choroid/retina and sclera. The pharmaceutically acceptable ophthalmic vehicle may, for example, be an ointment, vegetable oil or an encapsulating material. Alternatively, the peptide inhibitors of the invention may be injected directly into the vitreous and aqueous humour.

[00630] Compositions for rectal or vaginal administration include suppositories which may be prepared by mixing the peptide inhibitorss of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax, which are solid at room temperature but liquid at body temperature and, therefore, melt in the rectum or vaginal cavity and release the active compound.

[00631] Peptide inhibitors of the present invention may also be administered in liposomes or other lipid-based carriers. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a peptide inhibitor of the present invention, stabilizers, preservatives, excipients, and the like. In certain embodiments, the lipids comprise phospholipids, including the phosphatidyl cholines (lecithins) and serines, both natural and synthetic. Methods to form liposomes are known in the art.

[00632] Pharmaceutical compositions to be used in the invention suitable for parenteral administration may comprise sterile aqueous solutions and/or suspensions of the peptide inhibitos made isotonic with the blood of the recipient, generally using sodium chloride, glycerin, glucose, mannitol, sorbitol, and the like.

[00633] In some aspects, the invention provides a pharmaceutical composition for oral delivery. Compositions and peptide inhibitors of the instant invention may be prepared for oral administration according to any of the methods, techniques, and/or delivery vehicles described herein. Further, one having skill in the art will appreciate that the peptide inhibitors of the instant invention may be modified or integrated into a system or delivery vehicle that is not disclosed herein, yet is well known in the art and compatible for use in oral delivery of peptides.

[00634] In certain embodiments, formulations for oral administration may comprise adjuvants (e.g. resorcinols and/or nonionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to artificially increase the permeability of the intestinal walls, and/or enzymatic inhibitors (e.g. pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) or trasylool) to inhibit enzymatic degradation. In certain embodiments, the peptide inhibitor of a solid-type dosage form for oral administration can be mixed with at least one additive, such as sucrose, lactose, cellulose, mannitol, trehalose, raffinose, maltitol, dextran, starches, agar, alginates, chitins, chitosans, pectins, gum tragacanth, gum arabic, gelatin, collagen, casein, albumin, synthetic or semisynthetic polymer, or glyceride. These dosage forms can also contain other type(s) of additives, e.g., inactive diluting agent, lubricant such as magnesium stearate, paraben, preserving agent such as sorbic acid, ascorbic acid, alpha-tocopherol, antioxidants such as cysteine, disintegrators, binders, thickeners, buffering agents, pH adjusting agents, sweetening agents, flavoring agents or perfuming agents.

[00635] In particular embodiments, oral dosage forms or unit doses compatible for use with the peptide inhibitors of the present invention may include a mixture of peptide inhibitor and nondrug components or excipients, as well as other non-reusable materials that may be considered either as an ingredient or packaging. Oral compositions may include at least one of a liquid, a solid, and a semi-solid dosage forms. In some embodiments, an oral dosage form is provided comprising an effective amount of peptide inhibitor, wherein the dosage form comprises at least one of a pill, a tablet, a capsule, a gel, a paste, a drink, a syrup, ointment, and suppository. In some instances, an oral dosage form is provided that is designed and configured to achieve delayed release of the peptide inhibitor in the subject's small intestine and/or colon.

[00636] In one embodiment, an oral pharmaceutical composition comprising a peptide inhibitor of the present invention comprises an enteric coating that is designed to delay release of the peptide inhibitor in the small intestine. In at least some embodiments, a pharmaceutical composition is provided which comprises a peptide inhibitor of the present invention and a protease inhibitor, such as aprotinin, in a delayed release pharmaceutical formulation. In some instances, pharmaceutical compositions of the instant invention comprise an enteric coat that is soluble in gastric juice at a pH of about 5.0 or higher. In at least one embodiment, a pharmaceutical composition is provided comprising an enteric coating comprising a polymer having dissociable carboxylic groups, such as derivatives of cellulose, including hydroxypropylmethyl cellulose phthalate, cellulose acetate phthalate and cellulose acetate trimellitate and similar derivatives of cellulose and other carbohydrate polymers.

[00637] In one embodiment, a pharmaceutical composition comprising a peptide inhibitor of the present invention is provided in an enteric coating, the enteric coating being designed to protect and release the pharmaceutical composition in a controlled manner within the subject's lower gastrointestinal system, and to avoid systemic side effects. In addition to enteric coatings, the peptide inhibitors of the instant invention may be encapsulated, coated, engaged or otherwise associated within any compatible oral drug delivery system or component. For example, in some embodiments a peptide inhibitor of the present invention is provided in a lipid carrier system comprising at least one of polymeric hydrogels, nanoparticles, microspheres, micelles, and other lipid systems.

[00638] To overcome peptide degradation in the small intestine, some embodiments of the present invention comprise a hydrogel polymer carrier system in which a peptide inhibitor of the present invention is contained, whereby the hydrogel polymer protects the peptide inhibitor from proteolysis in the small intestine and/or colon. The peptide inhibitors of the present invention may further be formulated for compatible use with a carrier system that is designed to increase the dissolution kinetics and enhance intestinal absorption of the peptide. These methods include the use of liposomes, micelles and nanoparticles to increase GI tract permeation of peptides.

[00639] Various bioresponsive systems may also be combined with one or more peptide inhibitor of the present invention to provide a pharmaceutical agent for oral delivery. In some

embodiments, a peptide inhibitor of the instant invention is used in combination with a bioresponsive system, such as hydrogels and mucoadhesive polymers with hydrogen bonding groups (e.g., PEG, poly(methacrylic) acid [PMAA], cellulose, Eudragit®, chitosan and alginate) to provide a therapeutic agent for oral administration. Other embodiments include a method for optimizing or prolonging drug residence time for a peptide inhibitor disclosed herein, wherein the surface of the peptide inhibitor surface is modified to comprise mucoadhesive properties through hydrogen bonds, polymers with linked mucins or/and hydrophobic interactions. These modified peptide molecules may demonstrate increase drug residence time within the subject, in accordance with a desired feature of the invention. Moreover, targeted mucoadhesive systems may specifically bind to receptors at the enterocytes and M-cell surfaces, thereby further increasing the uptake of particles containing the peptide inhibitor.

**[00640]** Other embodiments comprise a method for oral delivery of a peptide inhibitor of the present invention, wherein the peptide inhibitor is provided to a subject in combination with permeation enhancers that promote the transport of the peptides across the intestinal mucosa by increasing paracellular or transcellular permeation. Various permeation enhancers and methods for the oral delivery of therapeutic agents is described in Brayden, D.J., Mrsny, R.J., 2011. Oral peptide delivery: prioritizing the leading technologies. *Ther. Delivery* 2 (12), 1567–1573.

**[00641]** In certain embodiments, pharmaceutical compositions and formulations of the present invention comprises a peptide inhibitor of the present invention and one or more permeation enhancer. Examples of absorption enhancers may include Bile salts, fatty acids, surfactants (anionic, cationic, and nonanionic) chelators, Zonular OT, esters, cyclodextrin, dextran sulfate, azone, crown ethers, EDTA, sucrose esters, and phosphatidyl choline, for example. Although absorption enhancers are not typically carriers by themselves, they are also widely associated with other carriers to improve oral bioavailability by transporting of peptides and proteins across the intestinal mucosa. Such substances can be added to the formulation as excipients or incorporated to form non specific interactions with the intended peptide inhibitor.

**[00642]** Dietary components and/or other naturally occurring substances affirmed as enhancing tight junction permeation and as Generally Recognized As Safe (GRAS) include, e.g., asglycerides, acylcarnitines, bile salts, and medium chain fatty acids. Sodium salts of medium

chain fatty acids (MCFAS) were also suggested to be permeation enhancers. The most extensively studied MCFAS is sodium caprate, a salt of capric acid, which comprises 2-3% of the fatty acids in the milk fat fraction. To date, sodium caprate is mainly used as an excipient in a suppository formulation (Doktacillin™) for improving rectal ampicillin absorption. The permeation properties of another dietary MCFAS, sodium caprylate (8-carbon), were shown in vitro to be lower when compared to sodium caprate. Sodium caprylate and a peptidic drug were formulated in an admixture with other excipients in oil to generate an oily suspension (OS) that enhanced permeability (Tuvia, S. et al., *Pharmaceutical Research*, Vol. 31, No. 8, pp. 2010-2021 (2014)).

**[00643]**For example, in one embodiment, a permeation enhancer is combined with a peptide inhibitor, wherein the permeation enhancer comprises at least one of a medium-chain fatty acid, a long-chain fatty acid, a bile salt, an amphiphilic surfactant, and a chelating agent. In certain embodiments, medium-chain fatty acid salts promote absorption by increasing paracellular permeability of the intestinal epithelium. In one embodiment, a permeation enhancer comprising sodium N-[hydroxybenzoyl]amino] caprylate is used to form a weak noncovalent association with the peptide inhibitor of the instant invention, wherein the permeation enhancer favors membrane transport and further dissociation once reaching the blood circulation. In another embodiment, a peptide inhibitor of the present invention is conjugated to oligoarginine, thereby increasing cellular penetration of the peptide into various cell types. Further, in at least one embodiment a noncovalent bond is provided between a peptide inhibitor of the present invention and a permeation enhancer selected from the group consisting of a cyclodextrin (CD) and a dendrimers, wherein the permeation enhancer reduces peptide aggregation and increasing stability and solubility for the peptide inhibitor molecule.

**[00644]**In certain embodiments, a pharmaceutical composition or formulation comprises a peptide inhibitor of the present invention and a transient permeability enhancers (TPEs). Permeation enhancers and TPEs may be used to increase orally bioavailability of the peptide inhibitor. One example of a TPE that may be used is an oily suspension formulation that disperses a powder containing sodium caprylate and a therapeutic agent (Tuvia, S. et al., *Pharmaceutical Research*, Vol. 31, No. 8, pp. 2010-2021 (2014)).

[00645] In certain embodiments, pharmaceutical composition and formulations may include a peptide inhibitor of the present invention and one or more absorption enhancers, enzyme inhibitors, or mucoso adhesive polymers.

[00646] In particular embodiments, peptide inhibtors of the present invention are formulated in a formulation vehicle, such as, e.g., emulsions, liposomes, microsphere or nanoparticles.

[00647] Other embodiments of the invention provide a method for treating a subject with a peptide inhibitor of the present invention having an increased half-life. In one aspect, the present invention provides a peptide inhibitor having a half-life of at least several hours to one day *in vitro* or *in vivo* (e.g., when administered to a human subject) sufficient for daily (q.d.) or twice daily (b.i.d.) dosing of a therapeutically effective amount. In another embodiment, the peptide inhibitor has a half-life of three days or longer sufficient for weekly (q.w.) dosing of a therapeutically effective amount. Further, in another embodiment, the peptide inhibitor has a half-life of eight days or longer sufficient for bi-weekly (b.i.w.) or monthly dosing of a therapeutically effective amount. In another embodiment, the peptide inhibitor is derivatized or modified such that it has a longer half-life as compared to the underivatized or unmodified peptide inhibitor. In another embodiment, the peptide inhibitor contains one or more chemical modifications to increase serum half-life.

[00648] When used in at least one of the treatments or delivery systems described herein, a peptide inhibitor of the present invention may be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt form.

[00649] The total daily usage of the peptide inhibitors and compositions of the present invention can be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular subject will depend upon a variety of factors including: a) the disorder being treated and the severity of the disorder; b) activity of the specific compound employed; c) the specific composition employed, the age, body weight, general health, sex and diet of the patient; d) the time of administration, route of administration, and rate of excretion of the specific peptide inhibitor employed; e) the duration of the treatment; f) drugs used in combination or coincidental with the specific peptide inhibitor employed, and like factors well known in the medical arts.

[00650] In particular embodiments, the total daily dose of the peptide inhibitors of the invention to be administered to a human or other mammal host in single or divided doses may be in amounts, for example, from 0.0001 to 300 mg/kg body weight daily or 1 to 300 mg/kg body weight daily.

#### Non-invasive Detection of Intestinal Inflammation

[00651] The peptide inhibitors of the invention may be used for detection, assessment and diagnosis of intestinal inflammation by microPET imaging, wherein the peptide inhibitor is labeled with a chelating group or a detectable label, as part of a non-invasive diagnostic procedure. In one embodiment, a peptide inhibitor is conjugated with a bifunctional chelator. In another embodiment, a peptide inhibitor is radiolabeled. The labeled peptide inhibitor is then administered to a subject orally or rectally. In one embodiment, the labeled peptide inhibitor is included in drinking water. Following uptake of the peptide inhibitor, microPET imaging may be used to visualize inflammation throughout the subject's bowels and digestive track.

#### Identification of Peptide Inhibitors that Inhibit IL-23 Signalling

[00652] As described herein, in certain embodiments, peptide inhibitors of the present invention preferentially bind to human IL-23R and/or rat IL-23R as compared to mouse IL-23R. Mouse IL-23R contains additional amino acids as compared to human IL-23R or rat IL-23R in the region corresponding to about amino acid residue 315 to about amino acid residue 340 of the mouse IL23R protein, e.g., amino acid region NWQPWSSPFVHQTSQETGKR (see, e.g., Figure 4). In particular embodiments, the peptide inhibitors bind to a region of human IL-23R from about amino acid 230 to about amino acid residue 370.

[00653] The present invention provides a new method to identify an inhibitor (e.g., a peptide inhibitor) of IL-23R, based on identifying an agent (e.g., a peptide) that preferentially binds to human IL-23R or rat IL-23R as compared to mouse IL-23R. In certain embodiments, the method comprises: (a) determining an amount of binding of a candidate agent to a human IL-23R polypeptide or a rat IL-23R polypeptide; (b) determining an amount of binding of the candidate agent to the mouse IL-23R polypeptide; and (c) comparing the determined amount of binding to the human IL-23R polypeptide or the rat IL-23R polypeptide to the determined amount of binding to the mouse IL-23R polypeptide, wherein if the determined amount of

binding to the human IL-23R polypeptide or the rat IL-23R polypeptide is greater than the amount of binding to the mouse IL-23R polypeptide, the candidate compound is an inhibitor of IL-23R. In particular embodiments, the candidate compound is identified as an inhibitor of IL-23R if the determined amount of binding to the human IL-23R polypeptide or the rat IL-23R polypeptide is at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 100-fold, at least 200-fold, at least 500-fold, or at least 100-fold the determined amount of binding to the mouse IL-23R polypeptide. In particular embodiments, the candidate compound is a peptide. In particular embodiments, the peptide is a peptide of one of the formulas described herein. In particular embodiments, the human IL-23 polypeptide or rat IL-23R polypeptide comprises or consists of the full length human IL-23R or rat IL-23R protein, respectively. In other embodiments, the human IL-23R polypeptide is a fragment of the full length human IL-23R protein, comprising 8 or more amino acid residues within the region of human IL-23R from about amino acid residue 230 to about amino acid residue 370. In other embodiments, the rat IL-23R polypeptide is a fragment of the full length rat IL-23R protein, comprising 8 or more amino acid residues within the region of rat IL-23R from about amino acid residue 245 to about amino acid residue 385.

[00654] In another embodiment, the present invention provides a new method to identify an inhibitor (e.g., a peptide inhibitor) of IL-23R, based on identifying an agent that binds to a region of human IL-23R or rat IL-23 that is disrupted in mouse IL-23R by the presence of additional amino acids from about amino acid residues 315 to about amino acid residue 340 of the mouse IL23R protein, e.g., amino acid region NWQPWSSPFVHQTSQETGKR (see, e.g., Figure 4). In certain embodiments, the method comprises: (a) determining an amount of binding of a candidate agent to a fragment of human IL-23R polypeptide that falls within about amino acid residue 230 to about amino acid residue 370, or to a fragment of rat IL-23R polypeptide that falls within about amino acid residue 245 to about amino acid residue 385; (b) determining an amount of binding of the candidate agent to a negative control (e.g., a negative control peptide unrelated to human IL-23R or rat-IL-23R); and (c) comparing the determined amount of binding to the fragment of human IL-23R polypeptide or the fragment of rat IL-23R polypeptide to the determined amount of binding to the negative control, wherein if the determined amount of binding to the human IL-23R polypeptide fragment or the rat IL-23R polypeptide fragment is

greater than the amount of binding to the negative control, the candidate compound is an inhibitor of IL-23R. In particular embodiments, the candidate compound is identified as an inhibitor of IL-23R if the determined amount of binding to the human IL-23R polypeptide fragment or the rat IL-23R polypeptide fragment is at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, at least 20-fold, at least 30-fold, at least 40-fold, at least 50-fold, at least 100-fold, at least 200-fold, at least 500-fold, or at least 100-fold the determined amount of binding to the negative control. In particular embodiments, the candidate compound is a peptide. In particular embodiments, the peptide is a peptide of one of the formulas described herein. In particular embodiments, the fragment of human IL-23R includes at least 8, at least 12, at least 20, at least 50, or at least 100, or all amino acid residues within the region of human IL-23R from about amino acid residue 230 to about amino acid residue 370. In other embodiments, the fragment of rat IL-23R polypeptide includes at least 8, at least 12, at least 20, at least 50, or at least 100, or all amino acid residues within the region of rat IL-23R from about amino acid residue 245 to about amino acid residue 385.

[00655] Methods of determining binding of a candidate compound to an IL-23 polypeptide are known in the art and include but are not limited to in vitro and cell-based binding assays, including those described herein. For example, a labeled candidate compound may be incubated with a recombinantly produced IL-23R polypeptide or negative control bound to a solid support under conditions and for a time sufficient to allow binding, and then binding determined by measuring the amount of label associated with the bound IL-23R polypeptide.

#### Non-invasive Detection of Intestinal Inflammation

[00656] The peptide inhibitors of the invention may be used for detection, assessment and diagnosis of intestinal inflammation by microPET imaging, wherein the peptide inhibitor is labeled with a chelating group or a detectable label, as part of a non-invasive diagnostic procedure. In one embodiment, a peptide inhibitor is conjugated with a bifunctional chelator. In another embodiment, a peptide inhibitor is radiolabeled. The labeled peptide inhibitor is then administered to a subject orally or rectally. In one embodiment, the labeled peptide inhibitor is included in drinking water. Following uptake of the peptide inhibitor, microPET imaging may be used to visualize inflammation throughout the subject's bowels and digestive track.

Animal Models of IBD

[00657] The present invention includes models of animal disease, including inflammatory diseases and disorders, such as inflammatory bowel diseases, e.g., Crohn's disease and colitis. As described in the accompanying Examples, several animal models of inflammatory diseases and disorders were developed.

[00658] In one embodiment, the present invention includes a method of assessing the ability of a candidate compound to inhibit or reduce an inflammatory disease disorder, comprising:

[00659] (a) providing to a rat an amount of dextran sulfate sodium (DSS) sufficient to induce IBD;

[00660] (b) providing to the rat an amount of a candidate compound; and

[00661] (c) measuring an amount of IBD symptoms present in the rat after being provided with the DSS and the candidate compound;

[00662] wherein if the amount of IBD symptoms measured in (c) are significantly lower than the amount measured in a control rat provided with the amount of DSS and either an amount of a control compound or no peptide (e.g., vehicle control), the candidate compound inhibits or reduces the inflammatory disease or disorder.

[00663] In certain embodiments, the rat is provided with DSS for about 5 to 12 days, e.g., about 9 days. In particular embodiments, the rat is provided with DSS by providing to the rat ad lib exposure to drinking water containing DSS, e.g., about 1% to about 10% DSS, about 2% to about 5% DSS, or about 3% DSS. In particular embodiments, the rat is provided with the test compound at about 5 mg/kg to about 100 mg/kg, or about 10 mg/kg to about 50 mg/kg, or about 20 mg/kg or about 30 mg/kg. In particular embodiments, the rat is provided with test compound orally, e.g., in drinking water. In certain embodiments, the DSS assay is performed as described in the accompanying Examples.

[00664] In another embodiment, the present invention includes a method of assessing the ability of a candidate compound to inhibit or reduce an inflammatory disease disorder, comprising:

[00665](a) providing to a rat an amount of 2,4,6-Trinitrobenzenesulfonic acid (TNBS) sufficient to induce IBD;

[00666](b) providing to the rat an amount of a candidate compound; and

[00667](c) measuring an amount of IBD symptoms present in the rat after being provided with the TNBS and the candidate compound;

[00668]wherein if the amount of IBD symptoms measured in (c) are significantly lower than the amount measured in a control rat provided with the amount of TNBS and either an amount of a control compound or no peptide (e.g., vehicle control), the candidate compound inhibits or reduces the inflammatory disease or disorder.

[00669]In certain embodiments, the animals are provided with about 10mg/kg to about 200 mg/kg TNBS, e.g., about 10 mg/kg, about 20 mg/kg, about 30 mg/kg, about 40 mg/kg, about 50 mg/kg, about 60 mg/kg, about 70 mg/kg, about 80 mg/kg, about 90 mg/kg, about 100 mg/kg, about 120 mg/kg, about 150 mg/kg or about 200 mg/kg of TNBS. In certain embodiments, the TNBS is in alcohol, e.g., in 45%-50% ethanol. In particular embodiments, the TNBS is administered intrarectally. In particular embodiments, the rat is provided with the test compound at about 5 mg/kg to about 100 mg/kg, or about 10 mg/kg to about 50 mg/kg, or about 20 mg/kg or about 30 mg/kg. In particular embodiments, the rat is provided with the test compound orally, e.g., in drinking water. In certain embodiments, the TNBS assay is performed as described in the accompanying Examples.

[00670]In particular embodiments IBD symptoms are measured immediately following provision of the DSS or TNBS and candidate compound (or test compound or no compound), or later, e.g., at about 3 days, 5 days, or 9 days following initial provision of DSS or TNBS and candidate compound (or test compound or no compound). In particular embodiments, the IBD symptoms measured include one or more of percent body weight loss, stool consistency, a quantitative hemoccult score, and ratio of colon weight:colon length. In certain embodiments, the IBD symptoms are measured using a disease activity index (DAI) score and/or ratio of colon weight:colon length, wherein the DAI score consists of ratings from three parameters, including

percent body weight loss, stool consistency, and a quantitative hemoccult score, and can achieve a maximum of three units.

[00671] In certain embodiments, a neutralizing anti-IL-23p19 antibody is used as a comparator or positive control.

[00672] In certain embodiments, to assess the extent of the inflammatory response, animals are observed, e.g., daily, for clinical signs which included percent body weight loss and signs of loose stools or diarrhea. Following a time period after inoculation of with DSS or TNBS (e.g., 5 days, 6, days, or seven days), rats are sacrificed and their entire colon length and colon weight from cecum to rectum recorded. The severity of colitis may be evaluated by a pathologist blinded to the identity of treatments. In addition to the colon wall thickness, the gross colon damage may be assessed based on a 0-4 scale according to Table 19 below, and histopathological scores were determined based on below parameters (Tables 20 and 21).

[00673] In certain embodiments, IBD symptoms are measured in three groups of rats, each with at least 3 animals, e.g., six animals each, wherein the three groups include: vehicle, DSS or TNBS, and DSS or TNBS with a positive control (e.g., sulfasalazine administered at 100 mg/kg PO, QD).

## EXAMPLES

### EXAMPLE 1

#### SYNTHESIS OF PEPTIDE MONOMERS

[00674] Peptide monomers of the present invention were synthesized using the Merrifield solid phase synthesis techniques on Protein Technology's Symphony multiple channel synthesizer. The peptides were assembled using HBTU (O-Benzotriazole-N,N,N',N'-tetramethyl-uronium-hexafluoro-phosphate), Diisopropylethylamine(DIEA) coupling conditions. For some amino acid couplings PyAOP(7-Azabenzotriazol-1-yloxy)trityrrolidinophosphonium hexafluorophosphate) and DIEA conditions were used. Rink Amide MBHA resin (100-200 mesh, 0.57 mmol/g) was used for peptide with C-terminal amides and pre-loaded Wang Resin with N- $\alpha$ -Fmoc protected amino acid was used for peptide with C-terminal acids. The coupling reagents (HBTU and DIEA premixed) were prepared at 100mmol concentration. Similarly amino acids solutions were

prepared at 100 mmol concentration. Peptide inhibitors of the present invention were identified based on medical chemistry optimization and/or phage display and screened to identify those having superior binding and/or inhibitory properties.

### Assembly

[00675] The peptides were assembled using standard Symphony protocols. The peptide sequences were assembled as follows: Resin (250 mg, 0.14 mmol) in each reaction vial was washed twice with 4ml of DMF followed by treatment with 2.5ml of 20% 4-methyl piperidine (Fmoc de-protection) for 10min. The resin was then filtered and washed two times with DMF (4ml) and re-treated with N-methyl piperifine for additional 30 minute. The resin was again washed three times with DMF (4 ml) followed by addition 2.5ml of amino acid and 2.5ml of HBTU-DIEA mixture. After 45min of frequent agitations, the resin was filtered and washed three timed with DMF (4 ml each). For a typical peptide of the present invention, double couplings were performed. After completing the coupling reaction, the resin was washed three times with DMF (4 ml each) before proceeding to the next amino acid coupling.

### Ring Closing Metathesis to form Olefins

[00676] The resin (100  $\mu$ mol) was washed with 2 ml of DCM ( $3 \times 1$  min) and then with 2 ml of DCE ( $3 \times 1$  min) before being treated with a solution of 2 ml of a 6 mM solution of Grubbs' first-generation catalyst in DCE (4.94 mg ml $^{-1}$ ; 20 mol% with regard to the resin substitution). The solution was refluxed overnight (12 h) under nitrogen before being drained. The resin was washed three times with DMF (4 ml each); DCM (4 ml) before being dried and cleaved.

### Cleavage

[00677] Following completion of the peptide assembly, the peptide was cleaved from the resin by treatment with cleavage reagent, such as reagent K (82.5% trifluoroacetic acid, 5% water, 5% thioanisole, 5% phenol, 2.5% 1,2-ethanedithiol). The cleavage reagent was able to successfully cleave the peptide from the resin, as well as all remaining side chain protecting groups.

[00678] The cleaved peptides were precipitated in cold diethyl ether followed by two washings with ethyl ether. The filtrate was poured off and a second aliquot of cold ether was added, and the procedure repeated. The crude peptide was dissolved in a solution of acetonitrile:water (7:3

with 1% TFA) and filtered. The quality of linear peptide was then verified using electrospray ionization mass spectrometry (ESI-MS) (Micromass/Waters ZQ) before being purified.

#### Disulfide Bond Formation via Oxidation

[00679] The peptide containing the free thiol (for example diPen) was assembled on a Rink Amide-MBHA resin following general Fmoc-SPPS procedure. The peptide was cleaved from the resin by treatment with cleavage reagent 90% trifluoroacetic acid, 5% water, 2.5% 1,2-ethanedithiol, 2.5% tri-isopropylsilane). The cleaved peptides were precipitated in cold diethyl ether followed by two washings with ethyl ether. The filtrate was poured off and a second aliquot of cold ether was added, and the procedure repeated. The crude peptide was dissolved in a solution of acetonitrile:water (7:3 with 1% TFA) and filtered giving the wanted unoxidized peptide crude peptide

[00680] The crude, cleaved peptide with X4 and X9 possessing either Cys, Pen, hCys, (D)Pen, (D)Cys or (D)hCys, was dissolved in 20ml of water : acetonitrile. Saturated Iodine in acetic acid was then added drop wise with stirring until yellow color persisted. The solution was stirred for 15 minutes, and the reaction was monitored with analytic HPLC and LCMS. When the reaction was completed, solid ascorbic acid was added until the solution became clear. The solvent mixture was then purified by first being diluted with water and then loaded onto a reverse phase HPLC machine (Luna C18 support, 10u, 100A, Mobile phase A: water containing 0.1% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA, gradient began with 5% B, and changed to 50% B over 60 minutes at a flow rate of 15ml/min). Fractions containing pure product were then freeze-dried on a lyophilyzer.

#### Lactam Bond Formation

[00681] 100mg of crude, cleaved peptide (approx. 0.12mmol) is dissolved in 100ml of anhydrous dichloromethane. HOEt (1-Hydroxybenzotriazole hydrate) (0.24mmol, 2 equivalents) is added followed by DIEA (N,N-Diisopropylethylamine) (1.2mmol, 10equivalents) and TBTU (O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate)(0.24 mmol, 2 equivalents). The mixture is stirred overnight and followed the reaction by HPLC. When the reaction is completed, dichloromethane is evaporated and diluted with water and Acetonitrile and then loaded onto a reverse phase HPLC machine (Luna C18 support, 10u, 100A, Mobile phase A:

water containing 0.1% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA, gradient begins with 5% B, and is changed to 50% B over 60 minutes at a flow rate of 15ml/min). Fractions containing pure product are then freeze-dried on a lyophilyzer.

#### Triazole bond formation

[00682] The purified peptide containing the relevant amino acids alkyne and azide was stirred at room temperature in a phosphate / MeOH (2 : 1) at pH 7.4 (1 mg per 2 ml). CuSO<sub>4</sub> · 5 H<sub>2</sub>O (10 equiv.), and sodium ascorbate (10 equiv.) was added and the mixture was agitated in at room temperature for 36 h. MeOH was removed and the solution was acidified to pH 3 with 1%TFA water mix. The solution was then filtered before being loaded onto HPLC for peptide purification.

#### Thioether Bond Formation

[00683] The peptide containing the free thiol (eg Cys) and hSer(OTBDMS) was assembled on a Rink Amide-MBHA resin following general Fmoc-SPPS procedure. Chlorination was carried out by treating the resin with PPh<sub>3</sub> (10 equiv.) and Cl<sub>3</sub>CCN (10 equiv.) in DCM for 2 h. The peptide was cleaved from the resin by treatment with cleavage reagent 90% trifluoroacetic acid, 5% water, 2.5% 1,2-ethanedithiol, 2.5% tri-isopropylsilane). The cleaved peptides were precipitated in cold diethyl ether followed by two washings with ethyl ether. The filtrate was poured off and a second aliquot of cold ether was added, and the procedure repeated. The crude peptide was dissolved in a solution of acetonitrile:water (7:3 with 1% TFA) and filtered giving the wanted uncyclized crude peptide

[00684] The crude peptide possessing a free thiol (eg Cys, Pen, hCys, (D)Pen, (D)Cys or (D)hCys) and the alkyl halide (hSer(Cl)) at either the X4 and X9 position or X9 and X4 position was dissolved in 0.1 M TRIS buffer pH 8.5. Cyclization was allowed to take place overnight at RT. The solvent mixture was then purified by first being diluted two-fold with water and then loaded onto a reverse phase HPLC machine (Luna C18 support, 10u, 100A, Mobile phase A: water containing 0.1% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA, gradient began with 5% B, and changed to 50% B over 60 minutes at a flow rate of 15ml/min). Fractions containing pure product were then freeze-dried on a lyophilyzer.

#### Selenoether Bond Formation

[00685] Crude peptide containing the thiol protected -Selenium amino acid and the alkyl halide at X4 and X9 was dissolved in 0.1 M sodium phosphate buffer pH 5.5 containing DTT (40 equiv.). Cyclization was allowed to take place over 24 h at RT. The solution was then diluted two-fold with water, and the final cyclized peptide was purified using RP-HPLC, affording the selenoether.

#### Diselenide Bond Formation

[00686] Diselenide precursor was dissolved in a solution of 0.1 M phosphate buffer pH 6.0 and isopropanol containing DTT (40 equiv), and the reaction mixture was incubated at 37°C. After 20h, additional DTT (10 equiv) was added to the reaction. After a total of 32h, the cyclization reaction was then diluted with twofold water, and the final cyclized peptide was purified using RP-HPLC, affording the diselenide.

#### Purification

[00687] Analytical reverse-phase, high performance liquid chromatography (HPLC) was performed on a Gemini C18 column (4.6 mm x 250 mm) (Phenomenex). Semi-Preparative reverse phase HPLC was performed on a Gemini 10 µm C18 column (22 mm x 250 mm) (Phenomenex) or Jupiter 10 µm, 300 Å C18 column (21.2 mm x 250 mm) (Phenomenex). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 1 mL/min (analytical) and 15 mL/min (preparative). Separations were achieved using linear gradients of buffer B in A (Mobile phase A: water containing 0.15% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA), at a flow rate of 1 mL/min (analytical) and 15mL/min (preparative).

#### Linker Activation and Dimerization

[00688] Peptid monomer subunits were linked to form peptide dimer inhibitors as described below.

[00689] Small Scale DIG Linker Activation Procedure: 5mL of NMP was added to a glass vial containing IDA diacid (304.2 mg, 1 mmol), N-hydroxysuccinimide (NHS, 253.2 mg, 2.2 eq. 2.2mmol) and a stirring bar. The mixture was stirred at room temperature to completely dissolve the solid starting materials. N, N'-Dicyclohexylcarbodiimide (DCC, 453.9mg, 2.2 eq., 2.2

mmol) was then added to the mixture. Precipitation appeared within 10 min and the reaction mixture was further stirred at room temperature overnight. The reaction mixture was then filtered to remove the precipitated dicyclohexylurea (DCU). The activated linker was kept in a closed vial prior to use for dimerization. The nominal concentration of the activated linker was approximately 0.20 M.

[00690] For dimerization using PEG linkers, there is no pre-activation step involved. Commercially available pre-activated bi-functional PEG linkers were used.

[00691] Dimerization Procedure: 2mL of anhydrous DMF was added to a vial containing peptide monomer (0.1 mmol). The pH of the peptide was the adjusted to 8~9 with DIEA. Activated linker (IDA or PEG13, PEG 25) (0.48eq relative to monomer, 0.048 mmol) was then added to the monomer solution. The reaction mixture was stirred at room temperature for one hour. Completion of the dimerization reaction was monitored using analytical HPLC. The time for completion of dimerization reaction varied depending upon the linker. After completion of reaction, the peptide was precipitated in cold ether and centrifuged. The supernatant ether layer was discarded. The precipitation step was repeated twice. The crude dimer was then purified using reverse phase HPLC (Luna C18 support, 10u, 100A, Mobile phase A: water containing 0.1% TFA, mobile phase B: Acetonitrile (ACN) containing 0.1% TFA, gradient of 15% B and change to 45% B over 60min, flow rate 15ml/min). Fractions containing pure product were then freeze-dried on a lyophilyzer.

## EXAMPLE 2

### CHARACTERIZATION OF PEPTIDE INHIBITION OF BINDING OF INTERLEUKIN-23 TO THE INTERLEUKIN-23 RECEPTOR

[00692] Peptide optimization was performed to identify peptide inhibitors of IL-23 signalling that were active at low concentrations (e.g., IC<sub>50</sub> <10 nM) while exhibiting gastrointestinal (GI) stability. Certain peptides were tested to identify peptides that inhibit the binding of IL-23 to human IL-23R and inhibit IL-23/IL-23R functional activity, as described below. Peptides tested included peptides containing a variety of different cyclization chemistries, including, e.g., cyclic amides (side chain cyclizations), peptides containing a disulfide linkage, e.g., between two Pen residues, and peptides containing a thioether linkage. Peptide inhibitors of the present invention

include but are not limited to peptides having any of the structures depicted herein. In addition, peptide inhibitors of the present invention include those having the same amino acid sequence of the peptides or structures described herein, without being required to have the same or any N- or C-terminal “capping” groups, such as, e.g., Ac or NH<sub>2</sub>.

[00693] Assays performed to determine peptide activity are described below, and the results of these assays is provided in Tables E3A-E3H, E4A and E4B, E5A-E5C, E6, E7, and E8. Human ELISA indicates the IL23-IL23R competitive binding assay described below, Rat ELISA indicates the rat IL-23R competitive binding ELISA assay described below, and pStat3HTRF indicates the DB cells IL-23R pSTAT3 cell assay described below. The peptides depicted in Tables E3B-E3E are cyclized via a disulfide bridge formed between two cysteine residues in these peptides. The peptides depicted in Table E3F are dimerized via a linker moiety or through internal cysteine moieties, as indicated. The peptides depicted in Tables E4A and E4B are cyclized via the two Pen residues present in each of these peptides. The peptides depicted in Table E5A are cyclized via a thioether bond between the indicated amino acid residues. Table E5B provides an illustrative structure depicting thioether cyclization, which is indicated in the table by the term “Cyclo,” with the cyclic region bracketed immediately following. The monomer subunits of the peptide dimers shown in Table E5C are cyclized as indicated by the term “Cyclo” and linked to each other via the indicated linker. The peptides shown in Table E6 are cyclized via ring closing metathesis of the indicated residues. Table E7 provides two illustrative structures depicting side chain cyclizations via cyclic amides, and the peptides in this table are cyclized as indicated following the term “Cyclo.” Table E8 depicts peptides cyclized via a cysteine residue and a Pen residue.

[00694] Peptide inhibitors of the present invention include both the cyclized form of the peptides shown herein, as well as the non-cyclized forms. For certain peptides, the residue Abu is present where indicated, whereas in other embodiments related to the non-cyclized form, the Abu may be referred to as a hSer(Cl) or homoSer residue.

#### IL23-IL23R Competitive Binding ELISA

[00695] An Immulon® 4HBX plate was coated with 50 ng/well of IL23R\_huFC and incubated overnight at 4°C. The wells were washed four times with PBST, blocked with PBS

containing 3% Skim Milk for 1 hour at room temperature, and washed again four times with PBST. Serial dilutions of test peptides and IL-23 at a final concentration of 2 nM diluted in Assay Buffer (PBS containing 1% Skim Milk) were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected by incubation with 50 ng/well of goat anti-p40 polyclonal antibodies (R&D Systems #AF309) diluted in Assay Buffer for 1 hour at room temperature. The wells were again washed four times with PBST. The secondary antibodies, HRP conjugated donkey anti-goat IgG (Jackson ImmunoResearch Laboratories #705-035-147) diluted 1:5000 in Assay Buffer was then added, and incubated for 30 minutes at room temperature. The plate was finally washed as above. Signals were visualized with TMB One Component HRP Membrane Substrate, quenched with 2 M sulfuric acid and read spectrophotometrically at 450 nm. IC50 values for various test peptides determined from these data are shown in Tables E3A-E3H, E4A and 4EB, E5A-E5C, E6, E7, and E8.

#### Rat IL-23R Competitive Binding ELISA

[00696] An assay plate was coated with 300 ng/well of Rat IL-23R\_huFC and incubated overnight at 4°C. The wells were washed, blocked, and washed again. Serial dilutions of test peptides and IL-23 at a final concentration of 7 nM were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected with goat anti-p40 polyclonal antibodies, followed by an HRP conjugated donkey anti-goat IgG. Signals were visualized with TMB One Component HRP Membrane Substrate and quenched with 2 M sulfuric acid. IC50 values for various test peptides determined from these data are shown in Tables E3G, E3H, E4A, E4B, E5B, E5C and E8.

#### DB Cells IL23R pSTAT3 Cell Assay

[00697] IL-23 plays a central role in supporting and maintaining Th17 differentiation *in vivo*. This process is thought to be mediated primarily through the Signal Transducer and Activator of Transcription 3 (STAT3), with phosphorylation of STAT3 (to yield pSTAT3) leading to upregulation of RORC and pro-inflammatory IL-17. This cell assay examines the levels of pSTAT3 in IL-23R-expressing DB cells when stimulated with IL-23 in the presence of test compounds. DB cells (ATCC #CRL-2289), cultured in RPMI-1640 medium (ATCC #30-2001)

supplemented with 10% FBS and 1% Glutamine, were seeded at 5 X 10E5 cells/well in a 96 well tissue culture plate. Serial dilutions of test peptides and IL-23 at a final concentration of 0.5 nM were added to each well, and incubated for 30 minutes at 37°C in a 5% CO<sub>2</sub> humidified incubator. Changes in phospho-STAT3 levels in the cell lysates were detected using the Cisbio HTRF pSTAT3 Cellular Assay Kit, according to manufacturer's Two Plate Assay protocol. IC50 values determined from these data are shown in Tables E3E, E3G, E3H, E4A, E4B, E5B, E5C, and E8 as absolute values or within ranges. Where not shown, data was not determined.

Table E3A. Illustrative Non-cyclic Peptides and Activities

SEQ ID NO.	Sequence	ELISA IL23R / IL23 in nMoles (IC50)
1	Ac-[Aib]-[Aib]-TWQDYWLY-[Aib]-R-NH <sub>2</sub>	>100,000
2	Ac-CAMTWQDYWLYGRC-NH <sub>2</sub>	7200
3	Ac-[Aib]-[Aib]-TWQDYWLYGR-NH <sub>2</sub>	>100,000
4	Ac-AMTWQDYWLYGRK-NH <sub>2</sub>	4100
5	Ac-CAMTWQDYWLYGRCK-NH <sub>2</sub>	8500
6	Ac-KAMTWQDYWLYGR-NH <sub>2</sub>	5600
7	Ac-KCAMTWQDYWLYGRC-NH <sub>2</sub>	10600
8	Ac-AMTWaibDYWLYGR-NH <sub>2</sub>	>37,500
9	Ac-AMTWQDYWLYGR-NH <sub>2</sub>	6100
10	Cyclo-[AMTWQDYWLYGR]	Not active
11	Hy-AATWQDYWLYGR-OH	7785
12	Hy-AMAWQDYWLYGR-OH	24225
13	Hy-AMTAQDYWLYGR-OH	N/A
14	Hy-AMTWADYWLYGR-OH	6248
15	Hy-AMTWQAYWLYGR-OH	9589

Table E3B. Illustrative Peptides Containing the CXXXXC Motif with IC50 &gt;1 uM in IL23-IL23R Competitive Binding ELISA

SEQ ID NO.	Sequence
87	Hy-CSDWECYWHIFG-NH <sub>2</sub>
88	Hy-CETWECYWHSFS-NH <sub>2</sub>
89	Hy-CQSWEKYWHYYG-NH <sub>2</sub>
90	Hy-CSDWRCYWHVFG-NH <sub>2</sub>
91	Hy-CHTWVCYWHEFS-NH <sub>2</sub>
92	Hy-CTDWVCYWHEYS-NH <sub>2</sub>
93	Hy-CQTWVCYWHTYG-NH <sub>2</sub>

SEQ ID NO.	Sequence
94	Hy-CGNWECYWHVYG-NH <sub>2</sub>
95	Hy-CKDWKCYWHIYG-NH <sub>2</sub>
96	Hy-CRTWVCYWHVFG-NH <sub>2</sub>
97	Hy-CAD-[1-Nal]-VCYWHTFG-NH <sub>2</sub>
98	Hy-CAD-[2-Nal]-VCYWHTFG-NH <sub>2</sub>
99	Hy-CAD-[1-BIP]-VCYWHTFG-NH <sub>2</sub>
100	Hy-CAD-[Tic]-VCYWHTFG-NH <sub>2</sub>
101	Hy-CAD-[ $\beta$ hW]-VCYWHTFG-NH <sub>2</sub>
102	Hy-CADWVCY-[1-BIP]-HTFG-NH <sub>2</sub>
103	Hy-CADWVCY-[Tic]-HTFG-NH <sub>2</sub>
104	Hy-CADWVCY-[ $\beta$ hW]-HTFG-NH <sub>2</sub>
105	Hy-CADWVCYAHHTFG-NH <sub>2</sub>
106	Hy-ACDWVCYWHTFG-NH <sub>2</sub>
107	Hy-ACDWCCYWCCTFG-NH <sub>2</sub>
108	Hy-AADWCAYWCTFG-NH <sub>2</sub>
109	Hy-CADWCCYWCCTFG-NH <sub>2</sub>
110	Hy-CADWCCYWCCTFG-NH <sub>2</sub>
111	Hy-CADWCCYWCCTFG-NH <sub>2</sub>
112	Hy-CADWVCYWHTF-NH <sub>2</sub>
113	Hy-CADWVCYWHT-NH <sub>2</sub>
114	Hy-CADWVCYW-NH <sub>2</sub>
115	Hy-[ $\beta$ -Ala]-SCADWVCYWHTFG-OH
116	Ac-[(D)Lys]-SCADWVCYWHTFG-OH
117	Ac-[(D)Lys]-[ $\beta$ -Ala]-CADWVCYWHTFG-OH
118	Hy-[AEA]-CADWVCYWHTFG-OH
119	Ac-[(D)Lys]-CADWVCYWHTFG-OH
120	Ac-CKDWVCYWHTFG-OH
121	Ac-CADWKCYWHTFG-OH
122	Ac-CADWVCYWKTFG-OH
123	Ac-CADWVCYWHKFG-OH
124	Ac-CADWVCYWHTKG-OH
125	Ac-CADWVCYWHTF-[(D)Lys]-OH
126	Ac-CADWVCYWHTFG-NH <sub>2</sub>
127	Hy-CADWVCY-[1-Nal]-HTF-OH
128	Hy-CADWVCY-[1-Nal]-HT-[N-Me-Phe]-NH <sub>2</sub>
129	Hy-CADWVCY-[1-Nal]-H-[Sarc]-F-OH
130	Hy-CADWVCY-[1-Nal]-[N-Me-His]-TF-OH
131	Hy-CADWVCYWHTFGK-OH
132	Hy-C-[Sarc]-DWVCY-[1-Nal]-HTF-OH
133	Hy-CAD-[N-Me-Trp]-VCY-[1-Nal]-HTF-OH
134	Hy-CADW-[Sarc]-CY-[1-Nal]-HTF-OH
135	Hy-CADWVCY-[1-Nal]-HT-[(D)Phe]-OH
136	Hy-CADWVCY-[1-Nal]-HTF-[Sarc]-OH

SEQ ID NO.	Sequence
137	Ac-CATWVCYWHTFG-NH <sub>2</sub>
138	Ac-CADWECYWHTFG-NH <sub>2</sub>
139	Ac-CADWVCYWHRCGWWGC-NH <sub>2</sub>
140	Ac-CADWVCY-[1-Nal]-H-[(D)Ala]-FG-NH <sub>2</sub>
141	Ac-CADWVCY-[1-Nal]-H-[Aib]-FG-NH <sub>2</sub>
142	Ac-CADWVCY-[1-Nal]-H-[b-Ala]-FG-NH <sub>2</sub>
143	Ac-CADWVCY-[1-Nal]-FTFG-NH <sub>2</sub>
144	Ac-CADWVCY-[1-Nal]-[(D)Ala]-TFG-NH <sub>2</sub>
145	Ac-CADWVCY-[1-Nal]-H-[Aib]-[(D)Phe]-G-NH <sub>2</sub>
146	Ac-CADWVCY-[1-Nal]-HTFG-[Aib]-NH <sub>2</sub>
147	Ac-CADWVCY-[1-Nal]-[N-Me-His]-[(D)Ala]-F-[Aib]-NH <sub>2</sub>
148	Ac-CADWVCY-[1-Nal]-H-[AEP]-G-NH <sub>2</sub>
149	Ac-CADWVCY-[N-MeHis]-TFG-[AEA]-[(D)Lys]-NH <sub>2</sub>
150	Ac-CADWVCY-[Aic]-HTFG-[AEA]-[(D)Lys]-NH <sub>2</sub>
151	Ac-CADWVCY-[Bip]-HTFG-[AEA]-[(D)Lys]-NH <sub>2</sub>
152	Ac-CQTWQCYW-[N-MeArg]-ENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
153	Ac-CQTWQCYWR-[N-MeArg]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
154	Ac-CQTWQCYWR-[N-MeLys]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
155	Ac-CQTWQCYWR-[Sarc]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
156	Ac-CQTWQCYWR-[(D)Glu]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
157	Ac-CQTWQCYW-[(D)Arg]-ENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
158	Ac-CQTWQCYW-[(D)Arg]-[(D)Glu]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
159	Ac-CQTWQCYW-[N-MeGlu]-NG-[AEA]-[(D)-Lys]
160	Ac-CADWVC-NH <sub>2</sub>
161	Ac-CRDWQCYW-[N-MeArg]-KFG-[AEP]-[(D)-Lys]-NH <sub>2</sub>
162	Ac-CRDWQCYWR-[(D)Lys]-FG-[AEP]-[(D)-Lys]-NH <sub>2</sub>
163	Ac-CRDWQCYW-[(D)Arg]-KFG-[AEP]-[(D)-Lys]-NH <sub>2</sub>
164	Ac-CRDWQCYW-[(D)Arg]-[(D)Lys]-FG-[AEP]-[(D)-Lys]-NH <sub>2</sub>
165	Ac-CQTWQCYW-[N-MeArg]-ENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>

Table E3C. Illustrative Peptides Containing the CXXXXC Motif with IC50 of 500nM to 1000nM in IL23-IL23R Competitive Binding ELISA

SEQ ID NO.	Sequence
166	Hy-CTDWKCYWHEFG-NH <sub>2</sub>
167	Hy-CRTWTCYWHVYG-NH <sub>2</sub>
168	Hy-CPNWECYWHRCGWWGC-NH <sub>2</sub>
169	Hy-CADWVCYWHTFG-NH <sub>2</sub>
170	Hy-CADWMCYWHEYG-NH <sub>2</sub>
171	Hy-CTTWKCYWHQYG-NH <sub>2</sub>
172	Hy-CSNWECYWHHYG-NH <sub>2</sub>
173	Hy-CSDWVCYWHVYG-NH <sub>2</sub>
174	Hy-CDTWKCYWHRQS-NH <sub>2</sub>

175	Hy-CADWVCY-[1-Nal]-HTFG-NH <sub>2</sub>
176	Hy-CADWVCY-[2-Nal]-HTFG-NH <sub>2</sub>
177	Hy-CADWVCYWHTFG-NH <sub>2</sub>
178	Ac-CADWVCYWHTFG-[(D)Lys]-OH
179	Ac-CADWVCYWHTFGAP-[(D)Lys]-OH
180	Ac-CTDWKCYWHTFG-NH <sub>2</sub>
181	Ac-CRDWVCYWHTFG-NH <sub>2</sub>
182	Ac-CADWVCYWHEFG-NH <sub>2</sub>
183	Ac-CADWVCYWHFHQLRDA-NH <sub>2</sub>
184	Ac-CADWVCYWHEHSERVG-NH <sub>2</sub>
185	Ac-CADWVCYWHNHSEGSG-NH <sub>2</sub>
186	Ac-CADWVCYWHRSTGGQH-NH <sub>2</sub>
187	Ac-[(D)Lys]-CRDWQCY-[1-Nal]-HTH-[Sarc]-[AEP]-[(D)Arg]-NH <sub>2</sub>
188	Ac-TQFDCRTWECYWHTFG-NH <sub>2</sub>
189	Ac-GGVECNDWQCYWHTFG-NH <sub>2</sub>
190	Ac-REGTCSTWKCYWHTFG-NH <sub>2</sub>
191	Ac-DTPRCRTWECYWHTFG-NH <sub>2</sub>
192	Ac-GGGECECWECYWHTFG-NH <sub>2</sub>
193	Ac-GDHKCSSWECYWHTFG-NH <sub>2</sub>
194	Ac-GSVHCMTWECYWHTFG-NH <sub>2</sub>
195	Ac-CADWVCY-[1-Nal]-VTFG-NH <sub>2</sub>
196	Ac-CADWVCYW-[(D)His]-TFG-[AEA]-[(D)Lys]-NH <sub>2</sub>

Table E3D. Illustrative Peptides Containing the CXXXXC Motif with IC50 <500nM in IL23-IL23R Competitive Binding ELISA

SEQ ID NO.	Sequence
197	Hy-CRDWQCYWHKFG-NH <sub>2</sub>
198	Hy-CSNWVCYWHTYG-NH <sub>2</sub>
199	Ac-CADWVCYWHTFG-[(D)Lys]-OH
200	Ac-CADWVCYWHTFG-[(D)Lys]-OH
201	Ac-CADWVCYWHTFG-OH
202	Ac-CADWVCYWHTFG-[(D)Lys]-OH
203	Ac-CADWVCYWHTFG-[(D)Lys]-OH
204	Ac-CADWVCYWHTFG-[(D)Lys]-OH
205	Ac-CADWVCYWHTFG-[(D)Lys]-OH
206	Ac-CADWVCYWHTFG-[(D)Lys]-OH
207	Ac-CADWVCYWHTFGP-[(D)Lys]-OH
208	Ac-CADWVCYWHTFG-[(D)Lys]-OH
209	Ac-CADWVCYWHTFGA-[(D)Lys]-OH
210	Ac-CADWVCYWHTFGAP-[(D)Lys]-OH
211	Ac-CADWVCYWHTFGAA-[(D)Lys]-OH
212	Ac-CADWVCYWHTFGAA-[(D)Lys]-OH
213	Ac-CRDWQCYWHKFG-[(D)Lys]-OH

SEQ ID NO.	Sequence
214	Ac-CATWQCYWHEYG-NH <sub>2</sub>
215	Ac-CKTWTCYWHEFG-NH <sub>2</sub>
216	Ac-CTTWTCYWHQYG-NH <sub>2</sub>
217	Ac-CRTWECYWHEFG-NH <sub>2</sub>
218	Ac-CRTWQCYWHEYG-NH <sub>2</sub>
219	Ac-CQTWQCYWRENG-NH <sub>2</sub>
220	Ac-CRTWECYWHEYG-NH <sub>2</sub>
221	Ac-CTTWECYWHEYG-NH <sub>2</sub>
222	Ac-CRTWECYWHEQS-NH <sub>2</sub>
223	Ac-CTTWECYWHQFG-NH <sub>2</sub>
224	Ac-CTTWECYWHEFG-NH <sub>2</sub>
225	Ac-CQTWECYWHLYG-NH <sub>2</sub>
226	Ac-CEDWKCYWHKYG-NH <sub>2</sub>
227	Ac-CTDWVCYWHTFG-NH <sub>2</sub>
228	Ac-CADWVCYWHTYG-NH <sub>2</sub>
229	Ac-CADWVCYWHRHDRVK-NH <sub>2</sub>
230	Ac-CADWVCYWHTFGER-NH <sub>2</sub>
231	Ac-CADWVCYWHTHGER-NH <sub>2</sub>
232	Ac-DTPRCRTWECYWHTFG-NH <sub>2</sub>
233	Ac-CQTWVCYWRENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
234	Ac-CQTWQCYWRENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
235	Ac-CQTWQCYWRTNG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
236	Ac-CQTWQCYWRKNG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
237	Ac-CQTWQCYWRRNG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
238	Ac-CQTWQCYWR-[Dapa]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
239	Ac-CQTWQCYWR-[Orn]-NG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
240	Ac-CRTWQCYWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
241	Ac-CQTWQCYWRENG-[AEA]-[(D)Arg]-NH <sub>2</sub>
242	Ac-CQTWQCYWRENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
243	Ac-CQDWQCYWRENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
244	Ac-CQTWQCYWRENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
245	Ac-CQTWQCYWRTNG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
246	Ac-CQTWVCYWRENG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
247	Ac-CQTWQCYWRKNG-[AEA]-[(D)-Lys]-NH <sub>2</sub>
248	Ac-CQTWQCYW-[Cav]-ENG-NH <sub>2</sub>
249	Ac-CQTWQCYW-[Cpa]-ENG-NH <sub>2</sub>
250	Ac-CQTWQCYWLENG-NH <sub>2</sub>
251	Ac-CQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>
252	Ac-CQTWQCYWR-[K-Ac]-NG-NH <sub>2</sub>
253	Hy-CRTWQCYWRKFG-NH <sub>2</sub>

Table E3E. IC50 of Illustrative Peptides Containing the CXXXXC Motif with Activities

SEQ ID NO.	Sequence	ELISA IL23R / IL23 in nM	pStat3 HTRF in nMoles
169	Hy-CADWVCYWHTFG-NH <sub>2</sub>	****	****
178	Ac-CADWVCYWHTFG-[(D)Lys]-OH	****	****
210	Ac-CADWVCYWHTFGAP-[(D)Lys]-OH	****	ND
211	Ac-CADWVCYWHTFGA[Azt]-[(D)Lys]-OH	****	ND
180	Ac-CTDWKCYWHTFG-NH <sub>2</sub>	****	****
196	Ac-CADWVCYW-[(D)His]-TFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	****	****
281	DIG dimererisation through N-termina Lysine (Ac-KMTWQDYWLGR-NH <sub>2</sub> ) <sub>2</sub>	*****	*****
284	DIG dimererisation through C-terminal Lysine (Ac-AMTWQDYWLGYK-NH <sub>2</sub> ) <sub>2</sub>	*****	*****

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=1000-10,000 nM.

Table E3F. IC50 of Illustrative Peptide Dimers

SEQ ID NO.	Linker Moiety	Sequence	Human IL23R/IL23 ELISA (nM)
277	oxidized dimer through the cysteine	(Hy-FPTWEWYWCNRD-NH <sub>2</sub> ) <sub>2</sub>	*****
278	oxidized dimer through the cysteine	(Hy-ALTWEFYWLCRE-NH <sub>2</sub> ) <sub>2</sub>	>10,000
291	DIG through Lysine	(Hy-[ $\beta$ Ala]SCADWVCYWHTFG-OH) <sub>2</sub> DIG	>10,000
292	DIG through Lysine	(Ac-[(D)Lys]-SCADWVCYWHTFG-OH) <sub>2</sub> DIG	>10,000
293	DIG through Lysine	(Ac-(D)Lys-[ $\beta$ Ala]-CADWVCYWHTFG-OH) <sub>2</sub> DIG	>10,000
294	DIG through Lysine	(Hy-AEA-CADWVCYWHTFG-OH) <sub>2</sub> DIG	>10,000
295	DIG through Lysine	(Ac-[(D)Lys]-CADWVCYWHTFG-OH) <sub>2</sub> DIG	>10,000
296	DIG through Lysine	(Ac-CKDWVCYWHTFG-OH) <sub>2</sub> DIG	>10,000
297	DIG through Lysine	(Ac-CADWKCYWHTFG-OH) <sub>2</sub> DIG	>10,000
298	DIG through Lysine	(Ac-CADWVCYWKTFG-OH) <sub>2</sub> DIG	
299	DIG through Lysine	(Ac-CADWVCYWHKFG-OH) <sub>2</sub> DIG	>10,000
300	DIG through Lysine	(Ac-CADWVCYWHTKG-OH) <sub>2</sub> DIG	*****
301	DIG through Lysine	(Ac-CADWVCYWHTFDK-OH) <sub>2</sub> DIG	>10,000
302	DIG through Lysine	(Ac-CADWVCYWHTFGDK) <sub>2</sub> DIG	*****
303	DIG through Lysine	(Ac-CADWVCYWHTFG-[ $\beta$ -Ala]-[(D)Lys]-OH) <sub>2</sub> DIG	***
304	DIG through Lysine	(Ac-CADWVCYWHTFG-[AEA]-[(D)Lys]-OH) <sub>2</sub> DIG	***
305	DIG through C terminal Lysine	(Hy-CADWVCYWHTFGK-OH) <sub>2</sub> DIG	*****
306	PEG25 through Lysine	(Hy-[ $\beta$ Ala]-SCADWVCYWHTFG-OH)	

SEQ ID NO.	Linker Moiety	Sequence	Human IL23R/IL23 ELISA (nM)
		<sup>2</sup> PEG25	
307	PEG25 through Lysine	(Ac-[(D)Lys]-SCADWVCYWHTFG-OH) <sub>2</sub>	
308	PEG25 through Lysine	(Ac-(D)Lys)-[βAla]-CADWVCYWHTFG-OH) <sub>2</sub>	
309	PEG25 through Lysine	(Hy-[AEA]-CADWVCYWHTFG-OH) <sub>2</sub>	
310	PEG25 through Lysine	(Ac-[(D)Lys]-CADWVCYWHTFG-OH) <sub>2</sub>	
311	PEG25 through Lysine	(Ac-CKDWVCYWHTFG-OH) <sub>2</sub>	
312	PEG25 through Lysine	(Ac-CADWKCYWHTFG-OH) <sub>2</sub>	
313	PEG25 through Lysine	(Ac-CADWVCYWKTFG-OH) <sub>2</sub>	
314	PEG25 through Lysine	(Ac-CADWVCYWHKFG-OH) <sub>2</sub>	
315	PEG25 through Lysine	(Ac-CADWVCYWHTKG-OH) <sub>2</sub>	
316	PEG25 through Lysine	(Ac-CADWVCYWHTF-[(D)Lys]-OH) <sub>2</sub>	
317	PEG25 through Lysine	(Ac-CADWVCYWHTFG-[(D)Lys]-OH) <sub>2</sub>	
318	PEG25 through Lysine	(Ac-CADWVCYWHTFG-[bAla]-[(D)Lys]-OH) <sub>2</sub>	
319	PEG25 through Lysine	(Ac-CADWVCYWHTFG-[AEA]-[(D)Lys]-OH) <sub>2</sub>	
320	PEG25 through C-terminal Lysine	(Hy-CADWVCYWHTFGK-OH) <sub>2</sub>	

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=1000-10,000 nM.

Table E3G. IC50 of Illustrative Peptides Containing the CXXWCXXXXX-[(D)Lys] Motif

SEQ ID NO.	Sequence	Human ELISA IL23/IL23R (nM)	Rat ELISA IL23/IL23R (nM)	pStat3 HTRF (nM)
16	Ac-CQDWQCYWR-[Cha]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	113		
17	Ac-CQTWQCYWR-[Ogl]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	206		
18	Ac-CQTWQCYWK-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	32		
19	Ac-CQTWQCYWH-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	49		59
20	Ac-CQTWQCYWRLFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	51		47
21	Ac-CQTWQCYW-[hArg]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	56		
22	Ac-CQTWQCYW-[Cit]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	25		
23	Ac-CQTWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	39	62	14
24	Ac-CQTWQCYWR-[Dap]-[Tic]-G-[AEA]-[(D)Lys]-NH <sub>2</sub>	892	65	12
25	Ac-CQTWQCY-[Tic]-[Orn]-KFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
26	Ac-CQTWQCYWR-[Dab]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	37		
27	Ac-CQTWQCYW-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	79	276	37
28	Ac-CQTWQCYWHENGA-[(D)Lys]-NH <sub>2</sub>	220		
29	Ac-CRTWQCYWRENGA-[(D)Lys]-NH <sub>2</sub>	102	86	17
30	Ac-CRTWQCYWREYGA-[(D)Lys]-NH <sub>2</sub>	78	80	8
31	Ac-C-[N-MeAla]-DWVCYWHTFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	183		

SEQ ID NO.	Sequence	Human ELISA IL23/IL23R (nM)	Rat ELISA IL23/IL23R (nM)	pStat3 HTRF (nM)
32	Ac-CADWVCYWRKFG-[βAla]-[(D)Lys]-NH <sub>2</sub>	57	33(1)	13
33	Ac-CADWVCYW-[Cit]-KFG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	52		29
34	Ac-CADWVCYW-[Cit]-[Tle]-FG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	518		
35	Ac-CADWVCYW-[Cit]-[Tba]-FG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	153		
36	Ac-CADWVCYW-[Cit]-[Cha]-FG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	223		
37	Ac-CADWVCY-[1-Nal]-[Cit]-VFG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	79		22
38	Ac-CADWVCYW-[Cit]-VFG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	124		
39	Ac-CADWVCYW-[Cit]-[Chg]-FG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	>30000		
40	Ac-CADWVCYW-[Cit]-[βAla]-FG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	2584		
41	Ac-CADWVCYW-[Tle]-[Tle]-FG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	~30000		
42	Ac-CADWVCYW-[Tle]-KFG-[β-Ala]-[(D)Lys]-NH <sub>2</sub>	199		
43	Ac-CQTWQCYW-[D(Ala)-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	232		
44	Ac-CQTWQCYW-[βAla]-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	2207		
45	Ac-CQTWQCYW-[D(Leu)-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	188		
46	Ac-CQTWQCYW-[D(Phe)-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	848		
47	Ac-CQTWQCYW-[D(Asn)-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	61		
48	Ac-CQTWQCYW-[D(Thr)-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	3662		
49	Ac-CQTWQCYW-[D(Asp)-VFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	129		
50	Ac-CQTWQCYW-[Cit]-[(D)Leu]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	709		
51	Ac-CQTWQCYW-[Cit]-[(D)Phe]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	1304		
52	Ac-CQTWQCYW-[Cit]-[(D)Asn]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	269		
53	Ac-CQTWQCYW-[Cit]-[(D)Thr]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	1214		
54	Ac-CQTWQCYW-[Agp]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	241		
55	Ac-CQTWQCY-[α-MeTrp]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
56	Ac-CQTWQCY-[α-MeTrp]-[Cit]-[hLeu]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
57	Ac-CQTWQCYW-[Cit]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	73		
58	Ac-CQTWQCYW-[Agp]-[Dap]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	38		
59	Ac-CQTWQCYW-[Cit]-VF-[(D)Ala]-[AEA]-[(D)Lys]-NH <sub>2</sub>	397		
60	Ac-CQTWQCYW-[Cit]-VF-[(D)Leu]-[AEA]-[(D)Lys]-NH <sub>2</sub>	444		
61	Ac-CQTWQCYW-[Cit]-VF-[(D)Phe]-[AEA]-[(D)Lys]-NH <sub>2</sub>	784		
62	Ac-CQTWQCYW-[Cit]-VF-[(D)Asn]-[AEA]-[(D)Lys]-NH <sub>2</sub>	93		
63	Ac-CQTWQCYW-[Cit]-VF-[(D)Thr]-[AEA]-[(D)Lys]-NH <sub>2</sub>	518		
64	Ac-CQTWQCYW-[Cit]-VF-[(D)Asp]-[AEA]-[(D)Lys]-NH <sub>2</sub>	551		
65	Ac-C-[N-MeArg]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	149	192	107
66	Ac-C-[N-MeQln]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	69	85	101
67	Ac-C-[Cit]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	50	76	107
68	Ac-CADWVCYW-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	382		
69	Ac-CADWVCY-[1-Nal]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	302		
70	Ac-CADWVCY-[(D)Trp]-[Orn]-[Dap]-FG-[AEA]-	>30000		

SEQ ID NO.	Sequence	Human ELISA IL23/IL23R (nM)	Rat ELISA IL23/IL23R (nM)	pStat3 HTRF (nM)
	[(D)Lys]-NH <sub>2</sub>			
71	Ac-CADWVCY-[hPhe]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~30000		
72	Ac-CADWVCY-[Bip]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
73	Ac-CADWVCY-[Phe(3,5-F <sub>2</sub> )]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
74	Ac-CADWVCY-[Phe(CONH <sub>2</sub> )]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
75	Ac-CADWVCY-[Phe(4-CF <sub>3</sub> )]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>1000		
76	Ac-CADWVCY-[Phe(2,4-Me <sub>2</sub> )]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	1525		
77	Ac-CMTWQCYWLYGR-[AEA]-[(D)Lys]-NH <sub>2</sub>	398		
77	Hy-CMTWQCYWLYGR-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
78	Ac-CADWVCY-[βhTrp]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
79	Ac-CADWVCYW-[Orn]-[α-MeLeu]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
80	Ac-CADWVCYW-[Orn]-[β-spiral-pip]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	579		
81	Ac-CADWVCY-[4-Phenylcyclohexylalanine]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>3000		
82	Ac-CADWVCYW-[Orn]-[Aib]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	1085		
83	Ac-CADWVCYW-[Orn]-[DiethylGly]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000		
84	Ac-CADWVCY-[α-MePhe(4-F)]-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
85	Ac-CQTWQCY-[βhPhe]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
86	Ac-CQTWQCY-[β(1-Nal)]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
321	Ac-CQTWQCY-[βhTyr]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
322	Ac-CQTWQCY-[βhPhe(4-F)]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
323	Ac-CQTWQCY-[βNva(5-Phenyl)]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
324	Ac-CQTWQCY-[Phe(3,4-Cl <sub>2</sub> )]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
325	Ac-CQTWQCY-[Tqa]-RVNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000		
326	Ac-CQTWQCYWR-[βhLeu]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	224		
327	Ac-CQTWQCYWR-[Aib]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	1065		
328	Ac-CQTWQCYWR-[βhAla]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	457		
329	Ac-CQTWQCYWR-[βhVal]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	328		
330	Ac-CQTWQCYWR-[β-spiral-pip]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	405		

SEQ ID NO.	Sequence	Human ELISA IL23/IL23R (nM)	Rat ELISA IL23/IL23R (nM)	pStat3 HTRF (nM)
331	Ac-CQTWQCYWR-[βGlu]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	250		
332	Ac-CQTWQCYW-[βhLeu]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	311		
333	Ac-CQTWQCYW-[βAib]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	2903		
334	Ac-CQTWQCYW-[βhAla]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	355		
335	Ac-CQTWQCYW-[βhVal]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	501		
336	Ac-CQTWQCYW-[β-spiral-pip]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>6000		
337	Ac-CQTWQCYW-[βhArg]-VNG-[AEA]-[(D)Lys]-NH <sub>2</sub>	922		
338	Ac-MRTWQ-[MeCys]-YWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	4251		
339	Ac-ACDWVCYWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	630		
340	Ac-SRTWQSYWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	2816		
341	Ac-CDWVCYWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	664		
342	Ac-ARTWQ-[MeCys]-YWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	7571		
343	Ac-ARTWQAYWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	3194		
344	Ac-CQTWQCYW-[hLeu]-EN-[AEA]-[(D)Lys]-NH <sub>2</sub>	132		
345	Ac-CQTWQCYW-[hLeu]-ENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	222		
346	Ac-CSTWECYWRVYGG-[AEA]-[(D)Lys]-NH <sub>2</sub>	47		
347	Ac-C-[Orn]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	22	69	95
348	Ac-CQTWQCYW-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	96		
349	Ac-C-[N-MeAsn]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	148		
350	Ac-C-[N-MeLys]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	80		
351	Ac-C-[Dab]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	23	51	99
352	Ac-CQTWQCYY-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	710		
353	Ac-CSTWQCYW-[Orn]-[Dap]-YG-[AEA]-[(D)Lys]-NH <sub>2</sub>	371		
354	Ac-CSTWECYW-[Cit]-[Dap]-YG-[AEA]-[(D)Lys]-NH <sub>2</sub>	74		
355	Ac-CQTWQCFF-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	4274		
356	Ac-CPTWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	422		
347	Ac-CSTWECYW-[Orn]-[Dab]-YG-[AEA]-[(D)Lys]-NH <sub>2</sub>	338		
358	Ac-CSTWECYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	48		
359	Ac-CLTWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	134		
360	Ac-CQTWQCYF-[Orn]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	1885		
461	Ac-CNTWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	21	79	96
362	Ac-C-[Dap]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	31		100
363	Ac-C-[N-Me-Ala]-TWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	139		
364	Ac-CKTWQCYWRVFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	40		
365	Ac-CQDWQCYWR-[Cha]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	113		
366	Ac-CQTWQCYWR-[Ogl]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	206		
367	Ac-CQTWQCYWK-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	32		
368	Ac-CQTWQCYWH-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	49		59
369	Ac-CQTWQCYWRLFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	51		47
370	Ac-CQTWQCYW-[hArg]-[Dap]-FG-[AEA]-[(D)Lys]-NH <sub>2</sub>	56		

Table E3H. IC50 of Illustrative Peptides Containing the CXXWXCXXXX Motif

SEQ ID NO.	Sequence	Human IL23/IL23R ELISA (nM)	Rat IL23/IL23R ELISA (nM)	pStat3 HTRF (nM)
371	Ac-CSTWECYWRTFG-NH <sub>2</sub>	252		
372	Ac-CDSWECYWRTYG-NH <sub>2</sub>	366		
373	Ac-CSTWECYWHTYG-NH <sub>2</sub>	181	286	97
374	Ac-CKTWTCYWHTYG-NH <sub>2</sub>	381		
375	Ac-CRTWECYWHEYS-NH <sub>2</sub>	416		
376	Ac-CRTWTCYWHEYG-NH <sub>2</sub>	434		
377	Ac-CFTWQCYWHEYS-NH <sub>2</sub>	515		
378	Ac-CQTWQCYW-[3-Pal]-ENG-NH <sub>2</sub>	56	20	101
379	Ac-CQTWQCY-NH <sub>2</sub>	>30000		
380	Ac-CRTWQCY-NH <sub>2</sub>	>30000		
381	Ac-CADWVCY-NH <sub>2</sub>	>30000		
382	Ac-CADWVCYW-NH <sub>2</sub>	>30000		
383	Ac-CADWVCYWH-NH <sub>2</sub>	~30000		
384	Ac-CADWVCYWHT-NH <sub>2</sub>	4795		
385	Ac-CADWVCYWHTF-NH <sub>2</sub>	3277		
386	Ac-CMTWQCYWLWGR-NH <sub>2</sub>	613		
387	Ac-CRTWQCYWHEFG-NH <sub>2</sub>			
388	Ac-CRTWECYWHTFG-NH <sub>2</sub>			
389	Ac-CQTWQCYWHEFG-NH <sub>2</sub>			
390	Ac-CRTWQCYWQQFGGE-NH <sub>2</sub>	81		
391	Ac-CRSWQCYWLNGPD-NH <sub>2</sub>	101		
392	Ac-CRTWQCYWLKGDS-NH <sub>2</sub>	39		
393	Ac-CQTWQCYWIKRDQG-NH <sub>2</sub>	67		
394	Ac-CSTWQCYWLKHGGE-NH <sub>2</sub>	19	24	2
395	Ac-CSTWECYWSQRADQ-NH <sub>2</sub>	240		
396	Ac-CQTWECYWRTFGPS-NH <sub>2</sub>	58		
397	Ac-CRTWQCYWQEKGTD-NH <sub>2</sub>	118		
398	Ac-CQTWQCYWLDSLGD-NH <sub>2</sub>	93		
399	Ac-CRTWQCYWTKFGSEP-NH <sub>2</sub>	87		57
340	Ac-CRSWQCYWNKFGADD-NH <sub>2</sub>	142		
341	Ac-CHTWQCYWLNGDEE-NH <sub>2</sub>	323		
342	Ac-CRTWQCYWLNGNEQ-NH <sub>2</sub>	127		
343	Ac-CRTWQCYWSEFGTGE-NH <sub>2</sub>	180	778	103
344	Ac-CRTWQCYWLRLGDEG-NH <sub>2</sub>	352	483	181
345	Ac-CHTWQCYWSTLGPEA-NH <sub>2</sub>	222		
346	Ac-CSTWQCYWSKQSGGS-NH <sub>2</sub>	133	204	89
347	Ac-CHTWQCYWLNNGTSQ-NH <sub>2</sub>	113		
348	Ac-CHTWQCYWRANDGRD-NH <sub>2</sub>	210		
349	Ac-SGCRTWQCYWHEFG-NH <sub>2</sub>	390		
350	Ac-NKCRTWQCYWHEYG-NH <sub>2</sub>	112		

SEQ ID NO.	Sequence	Human IL23/IL23R ELISA (nM)	Rat IL23/IL23R ELISA (nM)	pStat3 HTRF (nM)
351	Ac-SGCRTWECYWHEYG-NH <sub>2</sub>	257		
352	Ac-DACRTWECYWHKFG-NH <sub>2</sub>	165		
353	Ac-PECRTWECYWHKFG-NH <sub>2</sub>	197		
354	Ac-QVCQTWECYWREFG-NH <sub>2</sub>	145		
355	Ac-DRCVTWECYWREFG-NH <sub>2</sub>	217		
356	Ac-ADQCRTWQCYWHEFG-NH <sub>2</sub>	228		
357	Ac-KENCRTWECYWREFG-NH <sub>2</sub>	148		
358	Ac-VQECSTWQCYWRTFG-NH <sub>2</sub>	138		
359	Ac-GEECSTWQCYWRKFG-NH <sub>2</sub>	53		24
360	Ac-DGSCRTWQCYWHQFG-NH <sub>2</sub>	240		
361	Ac-NADCHSWECYWREFG-NH <sub>2</sub>	872		
362	Ac-ERNCSTWECYWRAFG-NH <sub>2</sub>	855		
363	Ac-RVGCASTWECYWREFG-NH <sub>2</sub>	417		
364	Ac-KANCRTWQCYWRKFE-NH <sub>2</sub>	412		
365	Ac-YEDCRTWQCYWENFG-NH <sub>2</sub>	280		
366	Ac-CQTWQCYWRNFGDS-NH <sub>2</sub>			
367	Ac-CQTWQCYWRNFESG-NH <sub>2</sub>			
368	Ac-CQDWQCYWREFGPG-NH <sub>2</sub>			
369	Ac-CQDWQCYWRSFGPQ-NH <sub>2</sub>			
370	Ac-CQTWQCYWRTLGPS-NH <sub>2</sub>			
371	Ac-CRTWQCYWQNFG-NH <sub>2</sub>	235		
372	Ac-CGTWQCYWRTFGPS-NH <sub>2</sub>	76		
373	Ac-CSTWQCYWHKFGNE-NH <sub>2</sub>	182		
374	Ac-CRTWECYWRTYGPS-NH <sub>2</sub>	116		
375	Ac-CRTWQCYWWENSQM-NH <sub>2</sub>	99		
376	Ac-CQTWQCYWREFGGG-NH <sub>2</sub>	165		
377	Ac-CQTWQCYWRTHGDR-NH <sub>2</sub>	83		
378	Ac-CRDWQCYWLSRP-NH <sub>2</sub>	330		
379	Ac-CQTWQCYW-[K(Palm)]-ENG-NH <sub>2</sub>	4880		
380	Ac-CQTWQCYW-[K(PEG8)]-ENG-NH <sub>2</sub>	153		
381	Ac-CQTWQCYW-[hLeu]-EQG-NH <sub>2</sub>	128		
382	Ac-CQTWQABUC-[(D)Tyr]-W-[hLeu]-ENG-NH <sub>2</sub>	>30000		
383	Ac-CQTWQC-[(N-MeTyr)]-W-[hLeu]-ENG-NH <sub>2</sub>	>30000		
384	Ac-CQTWQC-[Tic-OH]-W-[hLeu]-ENG-NH <sub>2</sub>	>30000		
385	Ac-CQTWQCEW-[hLeu]-ENG-NH <sub>2</sub>	>30000		
386	Ac-CQTWQCTW-[hLeu]-ENG-NH <sub>2</sub>	>30000		
387	Ac-CQTWQC-[Cha]-W-[hLeu]-ENG-NH <sub>2</sub>	~6000		
388	Ac-CQTWQCYW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	22	27	5
389	Ac-CQTWQCYW-[(D)Leu]-ENG-NH <sub>2</sub>	319		
390	Ac-CQTWQCYW-[hLeu]-ENG-[(D)Lys]-NH <sub>2</sub>	121		
391	Ac-CQTWQCYW-[hLeu]-ENG-OH	317		
392	Ac-CQTWQCYW-[hLeu]-ENE-NH <sub>2</sub>	222	1002	310
393	Ac-CQTWQCYW-[hLeu]-ENR-NH <sub>2</sub>	93		

SEQ ID NO.	Sequence	Human IL23/IL23R ELISA (nM)	Rat IL23/IL23R ELISA (nM)	pStat3 HTRF (nM)
394	Ac-CQTWQCYW-[hLeu]-ENF-NH <sub>2</sub>	82	182	69
395	Ac-CQTWQCYW-[hLeu]-ENP-NH <sub>2</sub>	253	114	31
396	Ac-CQTWQCYW-[hLeu]-ENQ-NH <sub>2</sub>	347		
397	Ac-CQTWQCYW-[hLeu]-ENL-NH <sub>2</sub>	45		
398	Ac-CQTWQCYW-[hLeu]-EEG-NH <sub>2</sub>	135	53	16
399	Ac-CQTWQCYW-[hLeu]-ERG-NH <sub>2</sub>	647		
400	Ac-CQTWQCYW-[hLeu]-EPG-NH <sub>2</sub>	108	140	27
401	Ac-CQTWQCYW-[hLeu]-ELG-NH <sub>2</sub>	158		
402	Ac-CQTWQCYW-[hLeu]-ETG-NH <sub>2</sub>	818		
403	Ac-CQTWQCYW-[hLeu]-FNG-NH <sub>2</sub>	395		
404	Ac-CQTWQCYW-[hLeu]-PNG-NH <sub>2</sub>	4828		
405	Ac-CQTWQCYW-[hLeu]-NNG-NH <sub>2</sub>	89		26
406	Ac-CQTWQCYW-[hLeu]-LNG-NH <sub>2</sub>	78		
407	Ac-CQTWQCYW-[hLeu]-TNG-NH <sub>2</sub>	109		
408	Ac-CQTWQCYWFENG-NH <sub>2</sub>	185		
409	Ac-CQTWQCYWPENG-NH <sub>2</sub>	>30000		
410	Ac-CQTWQCYWQENG-NH <sub>2</sub>	173		
411	Ac-CQTWQCYWTENG-NH <sub>2</sub>	114		
412	Ac-CQTWQCYWEENG-NH <sub>2</sub>	147		
413	Ac-CQTWFCYW-[hLeu]-ENG-NH <sub>2</sub>	1412		
414	Ac-CQTWPCYW-[hLeu]-ENG-NH <sub>2</sub>	2735		
415	Ac-CQTWNQCYW-[hLeu]-ENG-NH <sub>2</sub>	1849		
416	Ac-CQTWRFCYW-[hLeu]-ENG-NH <sub>2</sub>	278		
417	Ac-CQTWTCYW-[hLeu]-ENG-NH <sub>2</sub>	114		
418	Ac-CQTWEFCYW-[hLeu]-ENG-NH <sub>2</sub>	164		
419	Ac-CQTGQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
420	Ac-CQTPQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
421	Ac-CQTNQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
422	Ac-CQTRQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
423	Ac-CQTTQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
424	Ac-CQTEQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
425	Ac-CQFWQCYW-[hLeu]-ENG-NH <sub>2</sub>	1152		
426	Ac-CQPWQCYW-[hLeu]-ENG-NH <sub>2</sub>	>10,000		
427	Ac-CQNWQCYW-[hLeu]-ENG-NH <sub>2</sub>	336		
428	Ac-CQRWQCYW-[hLeu]-ENG-NH <sub>2</sub>	469		
429	Ac-CQEWFQCYW-[hLeu]-ENG-NH <sub>2</sub>	773		
450	Ac-CFTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	205		
451	Ac-CPTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	27412		
452	Ac-CNTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	61		
453	Ac-CGTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	167		
454	Ac-CTTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	59	28	10
455	Ac-CETWQCYW-[hLeu]-ENG-NH <sub>2</sub>	101		
456	Ac-CQTWQCYW-[N-MeLeu]-ENG-NH <sub>2</sub>	>6000		

SEQ ID NO.	Sequence	Human IL23/IL23R ELISA (nM)	Rat IL23/IL23R ELISA (nM)	pStat3 HTRF (nM)
457	Ac-CQTWQCYW-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	46	64	12
458	Ac-CQTWQCYW-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	28	31	7
459	Ac-CQTWQC-[ $\alpha$ -MePhe]-W-[hLeu]-ENG-NH <sub>2</sub>	~30000		
460	Ac-CQTWQCYW-[Aib]-ENG-NH <sub>2</sub>	31	34	12
461	Ac-CQTWQC-[hTyr]-W-[hLeu]-ENG-NH <sub>2</sub>	~6000		
462	Ac-CQTWQC-[Bip]-W-[hLeu]-ENG-NH <sub>2</sub>	237		
463	Ac-CQTWQCYW-[Ogl]-ENG-NH <sub>2</sub>	66	163	76
464	Ac-CQTWQCYW-[hLeu]-[Lys(Ac)]-NG-NH <sub>2</sub>	19	32	3
465	Ac-CQTWQCYW-[hLeu]-ENGG-NH <sub>2</sub>	61	140	24
466	Ac-CQTWQCYW-[hLeu]-ENGP-NH <sub>2</sub>	97		
467	Ac-CQTWQCYW-[hLeu]-ENGE-NH <sub>2</sub>	180		
468	Ac-CQTWQCYW-[hLeu]-ENG-(D)Glu-NH <sub>2</sub>	183		
469	Ac-CQTWQCY-[ $\alpha$ -McPhe]-[hLeu]-ENG-NH <sub>2</sub>	~30000		
470	Ac-CQTWQCYW-[hLeu]-ENGP-NH <sub>2</sub>	239		
471	Ac-CQTWQCYW-[hLeu]-ENGG-NH <sub>2</sub>	362		
472	Ac-CQTWQCYW-[hLeu]-ENGL-NH <sub>2</sub>	174		
473	Ac-CQTWQCYW-[hLeu]-ENGF-NH <sub>2</sub>	131		
474	Ac-CQTWQCYW-[hLeu]-ENGE-NH <sub>2</sub>	129		
475	Ac-CQTWQCYW-[hLeu]-ENGN-NH <sub>2</sub>	66		23
476	Ac-CQTWQCYW-[hLeu]-ENGT-NH <sub>2</sub>	160		
477	Ac-CQTWQCYW-[hLeu]-ENGR-NH <sub>2</sub>	>10,000		>1000
478	Ac-PCQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	97		
479	Ac-LCQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	61	26	21
480	Ac-FCQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	56	25	16
481	Ac-ECQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>			
482	Ac-NCQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>			
483	Ac-RCQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>			
484	Ac-CQTWQCY-[2-Nal]-[hLeu]-ENG-NH <sub>2</sub>			
485	Ac-CQTWQCY-[1-Nal]-[hLeu]-ENG-NH <sub>2</sub>	18	37	6
486	Ac-CQTWQC-[2-Nal]-W-[hLeu]-ENG-NH <sub>2</sub>	48	73	11
487	Ac-CQTWQC-[1-Nal]-[2-Nal]-[hLeu]-ENG-NH <sub>2</sub>	78	125	17
488	Ac-CQTWQC-[2-Nal]-[1-Nal]-[hLeu]-ENG-NH <sub>2</sub>	117		
489	Ac-CQTWQC-[Aic]-W-[hLeu]-ENG-NH <sub>2</sub>	126		
490	Ac-CQTWQCHW-[hLeu]-ENG-NH <sub>2</sub>	~6000		
491	Ac-CQTWQCYH-[hLeu]-ENG-NH <sub>2</sub>	398		
492	Ac-CQTWQC-[Tyr(OMe)]-W-[hLeu]-ENG-NH <sub>2</sub>	~30000		
493	Ac-CQTWQCY-[Bip]-[hLeu]-ENG-NH <sub>2</sub>	42	51	11
494	Ac-CQTWQCY-[Tyr(OMe)]-[hLeu]-ENG-NH <sub>2</sub>	998		
495	Ac-CQTWQCHH-[hLeu]-ENG-NH <sub>2</sub>	148		
496	Ac-CQTWQCY-[ $\alpha$ -MeTrp]-[hLeu]-EQG-NH <sub>2</sub>	>30000		
497	Ac-CQTW-[K(PEG8)]-CYWLENG-NH <sub>2</sub>	212		
498	Ac-CQTWQCYWZ-LNG-NH <sub>2</sub>	800		
499	Ac-CQTW-[K(PEG8)]CYW-[K(PEG8)]-ENG-NH <sub>2</sub>	753		

SEQ ID NO.	Sequence	Human IL23/IL23R ELISA (nM)	Rat IL23/IL23R ELISA (nM)	pStat3 HTRF (nM)
500	Ac-CQTW-[K(Palm)]-CYWLENG-NH <sub>2</sub>	~30000		
501	Ac-CQTWQCYW-[Orn]-[K(Palm)]-NG-NH <sub>2</sub>	>6000		
502	Ac-Gly-[(D)Asn]-(D)Glu-(D)Leu-(D)Trp-(D)Tyr-(D)Cys-(D)Gln-(D)Trp-(D)Thr-(D)Gln-(D)Cys-NH <sub>2</sub>	>30000		
503	Ac-CQTWQCYW-[Orn]-[K(Peg8)]-NG-NH <sub>2</sub>	169		
504	Ac-CRTWQCYWHEFG-NH <sub>2</sub>	166		
505	Ac-CRTWECYWHTFG-NH <sub>2</sub>	333		
506	Ac-CQTWQCYWHEFG-NH <sub>2</sub>	169		
507	Ac-CQTWQCYWRNFGDS-NH <sub>2</sub>	96		
508	Ac-CQTWQCYWRNFESG-NH <sub>2</sub>	315		
509	Ac-CQDWQCYWREFGPG-NH <sub>2</sub>	82		
510	Ac-CQDWQCYWRSFGPQ-NH <sub>2</sub>	117		
511	Ac-CQTWQCYWRTLGPNSH2	66		
512	Ac-CQTWQCYW-[(D)Pro]-ENG-NH <sub>2</sub>	>30000		
513	Ac-CQTWQCYWELNG-NH <sub>2</sub>	79		
514	Ac-CQTWECYWELNG-NH <sub>2</sub>	154		
515	Ac-CQTWQCY[(1-Nal)-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	22	67	13
516	Ac-CQTWQCY-[1-Nal]-[(D)Asn]-ENG-NH <sub>2</sub>	145		98
517	Ac-CQTWQCYWLE-[K(Palm)]-G-NH <sub>2</sub>	>6000		
518	Ac-CQTWQCYWLEN-[K(Palm)]-NH <sub>2</sub>	2800		
519	Ac-CSTWECYWRTFG-NH <sub>2</sub>	252		
520	Ac-CDSWECYWRTYG-NH <sub>2</sub>	366		
521	Ac-CSTWECYWHTYG-NH <sub>2</sub>	181	286	97

Table E4A. IC50 of Illustrative examples of dimers of Peptides Containing the Ac-[Pen]-XXWX-[Pen]-XXXX Motif and analogues

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
522	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> DIG	**		*
523	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> PEG25	*		**
524	[Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -Me-Leu]-QNN-NH <sub>2</sub> ] <sub>2</sub> DIG	**		**
525	[Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -Me-Leu]-QNN-NH <sub>2</sub> ] <sub>2</sub> PEG25	*		**
526	[Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub> ] <sub>2</sub> DIG	***		***
527	[Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub> ] <sub>2</sub> PEG25	**		***

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
528	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-[D]Lys] <sub>2</sub> DIG			*
529	[Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-acetylaminooethoxy)]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-[D]Lys] <sub>2</sub> DIG			*
530	[Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-acetylaminooethoxy)]-[2-Nal]-[ $\alpha$ -MeVal]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG		**	*
531	[Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-acetylaminooethoxy)]-[2-Nal]-K-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG		***	*
532	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG		**	*
533	[Ac-[ $\alpha$ -MeLys]-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG			****
534	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG	*	**	*
535	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	*	**	*
536	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
537	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Achc]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
538	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Acvc]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	*	**	*
539	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLeu]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
540	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
541	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
542	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
543	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Acvc]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
544	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG			*
545	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> IDA			*
546	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> [IDA- $\beta$ Ala]			*

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=1000-10,000 nM.

Table E4B. IC50 of Illustrative Peptides Containing the Ac-[Pen]-XXWX-[Pen]-XXXX Motif and Analogues

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
547	Ac-[Pen]-RTWQ-[Pen]-YWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	****	****	***
548	Ac-A-[Pen]-DWV-[Pen]-YWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	>30000		
549	Ac-[[Pen]-QTWQ-[Pen]-YW-[hLeu]-ENG-NH <sub>2</sub>	****		
550	Ac-[Pen]-QTWQ-[Pen]-YW [N-MeArg]-ENG-NH <sub>2</sub>	>30000		
551	Ac-[Pen]-QTWQ-[Pen]-YW-[hLeu]-ENG-NH <sub>2</sub>	****		
552	Ac-[Pen]-QTWQ-[Pen]-YW-[N-MeArg]-ENG-NH <sub>2</sub>	>30000		
553	Ac-A-[Pen]-DWV-[Pen]-YW-[Orn]-[Dap]-FG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	>30000		
554	Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	***	****	**
555	Ac-[Pen]-QTWQ-[Pen]-YW-[(D)Asn]-ENG-NH <sub>2</sub>	*****		
556	Ac-[Pen]-QTWQ-[Pen]-Y-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	****	*
557	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	****	**
558	Ac-[Pen]-QTWQ-[Pen]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****	****	**
559	Ac-[Pen]-QTWQ-[Pen]-Y-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	***	****	**
560	Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	****	****	***
561	Ac-[Pen]-QTWQ-[Pen]-Y-[1-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	****	****	***
562	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		***
563	Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		***
564	Ac-[Pen]-QTWQ-[Pen]-[Phe-(4-OMe)]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
565	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRE (nM)
566	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[1-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	****	***
567	Ac-[Pen]-QTWQ-[Pen]-[BIP]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>10,000		
568	Ac-[Pen]-QTWQ-[Pen]-Phe(3,4-Cl <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
569	Ac-[Pen]-QTWQ-[Pen]-[Phe(3,5-F <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
570	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-NH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
571	Ac-[Pen]-QTWQ-[Pen]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>10000		
572	Ac-[Pen]-QTWQ[Pen]-[Phe(3,4-Cl <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	****		
573	Ac-[Pen]-QTWQ[Pen]-[Phe(4-CN)]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	****		
574	Ac-[Pen]-QTWQ[Pen]-[Phe(3,5-F <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	****		
575	Ac-[Pen]-QTWQ[Pen]-[Phe(4-CH <sub>2</sub> CO <sub>2</sub> H)]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>			
576	Ac-[Pen]-QTWQ[Pen]-[Phe(4-CH <sub>2</sub> COEt <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>			
577	Ac-[Pen]-QTWQ[Pen]-[Phe(penta-F)]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>			
578	Ac-[Pen]-QTWQ[Pen]-[Phe(4-CF <sub>3</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>			
579	Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>			
580	Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>			
581	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-K(ivDde)-NG-NH <sub>2</sub>	****		
582	succinic acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	*
583	glutaric acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
584	4-methylmorpholine-2,6-dione-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
585	pyroglutamic acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRE (nM)
586	isovaleric acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
587	gallic acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****		
588	octanoic acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
589	4-Biphenylacetic acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
590	4-fluorophenylacetic acid-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	****	*
591	Hy-[Pen]-ADWV-[Pen]-YWHTFG-NH <sub>2</sub>	>6000		
592	Ac-[Pen]-GTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**		*
593	Ac-[Pen]-TTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**		*
594	Ac-[Pen]-STWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**		*
595	Ac-[Pen]-[Dap]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*	***	*
596	Ac-[Pen]-[ $\alpha$ -MeOrn]-TWQ[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****		
597	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*		*
598	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	***	*
599	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	**	*
600	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**		*
601	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-ENA-NH <sub>2</sub>	*	***	*
602	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*
603	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-QNN-NH <sub>2</sub>	*	*	*
604	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>	*		*
605	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-Aib-[Lys(Ac)]-NN-NH <sub>2</sub>	*		*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRE (nM)
606	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>	*		*
607	Ac-[Pen]-Dap(Ac)TWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	**		*
608	Ac-[Pen]-[ $\alpha$ -MeOrn(Ac)]-TWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	****		
609	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NG-NH <sub>2</sub>	*		*
610	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NN-NH <sub>2</sub>	*		*
611	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	**		*
612	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENA-NH <sub>2</sub>	**		*
613	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*		*
614	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-QNN-NH <sub>2</sub>	*		*
615	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>	**		*
616	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*		*
617	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>	*		*
618	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>			*
619	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-ENA-NH <sub>2</sub>			**
620	Ac-[Pen]-TTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
621	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>			*
622	Ac-[Pen]-TTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>			*
623	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>			*
624	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>			*
625	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*


SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRE (nM)
626	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
627	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
628	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
629	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
630	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>			*
631	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-ENA-NH <sub>2</sub>			*
632	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*
633	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
634	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>			*
635	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>			*
636	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>			*
637	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
638	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>			*
639	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>		**	*
640	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
641	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			*
642	Ac-E-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
643	Ac-(D)Asp-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
644	Ac-R-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
645	Ac-(D)Arg-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*












SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRE (nM)
646	Ac-Phe-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
647	Ac-(D)Phe-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
648	Ac-[2-Nal]-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
649	Ac-T-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
650	Ac-L-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
651	Ac-(D)Gln-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			
652	Ac-[(D)Asn]-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			
653	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-[(D)Lys]-NH <sub>2</sub>			*
654	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeVal]-KNN-NH <sub>2</sub>			*
655	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-K-[Lys(Ac)]-NN-NH <sub>2</sub>			***
666	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
667	Ac-[(D)Lys]-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-NH <sub>2</sub>			*****
668	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>		***	*
669	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
670	Ac-[Pen]-QTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[Phe(3,4-OMe <sub>2</sub> )]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-NH <sub>2</sub>			**
671	Ac-[(D)Phe]-[Pen]-NTWQ[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			
672	Ac-[(D)Phe]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>			
673	Ac-[(D)Phe]-[Pen]-NTWQ[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Achc]-[Lys(Ac)]-NN-NH <sub>2</sub>			
674	Ac-[(D)Phe]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Cit]-NN-NH <sub>2</sub>			
675	Ac-[(D)Phe]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Achc]-[Cit]-NN-NH <sub>2</sub>			

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRE (nM)
676	Ac-[D(Phe)-[Pen]-NTWQ-[Pen]-[Phe(4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>			
677	Ac-[D(Phe)-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>			
678	Ac-[D(Phe)-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-[Lys(Ac)]-NN-NH <sub>2</sub>			
679	Ac-[D(Phe)-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Cit]-NN-NH <sub>2</sub>			
680	Ac-[D(Phe)-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-[Cit]-NN-NH <sub>2</sub>			
681	Ac-[D(Phe)-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-ENN-NH <sub>2</sub>			
682	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			
683	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>			
684	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Achc]-[Lys(Ac)]-NN-NH <sub>2</sub>			
685	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Acpc]-[Lys(Ac)]-NN-NH <sub>2</sub>			
686	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>			
687	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			
688	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>			
689	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-[Lys(Ac)]-NN-NH <sub>2</sub>			
670	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[Acpc]-[Lys(Ac)]-NN-NH <sub>2</sub>			
671	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>			

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=1000-10,000 nM.

Table E5A. IC50 of Illustrative Peptide Inhibitors (Thioethers)

SEQ ID NO.	Sequence/Structure	Human ELISA (nM)
672	 N-MeAla-DWVCYWHTFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000

SEQ ID NO.	Sequence/Structure	Human ELISA (nM)
673	 <p>[N-MeAla]-DWV-[Pen]-YWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	>30000
674	 <p>[N-MeAla]-DWV-[(D)Pen]-YWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	>30000
675	 <p>[N-MeAla]-DWV-[hCys]-YWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~6000
676	 <p>ADWVCYWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~3000
677	 <p>ADWV-[Pen]-YWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	>30000
678	 <p>ADWV-[(D)Pen]-YWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	>30000
679	 <p>ADWV-[hCys]-YWHTFG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~6000
680	 <p>QDWQCYWRENG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	>6000
681	 <p>QDWQCYWRENG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~30000
682	 <p>QDWQCYWRENG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~6000
683	 <p>QDWQCYWRENG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~6000
684	 <p>QDWQCYWRENG-[AEA]-[(D)Lys]-NH<sub>2</sub></p>	~30000









SEQ ID NO.	Sequence/Structure	Human ELISA (nM)
685		>6000
686		>6000
687		>6000
688		>6000
689		~30000
690		>30000
691		>30000

Table E5B. IC50 of Illustrative Peptide Inhibitors (Thioethers)

Ac-Cyclo-[[Abu]-XXWC]-[Phe(4-OMe)]-[2-Nal]-XXX-NH<sub>2</sub>

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
692	Ac-Cyclo-[[Abu]RTWQC]-YWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	***	****	***

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
693	Ac-Cyclo-[CRTWQ-[Abu]]-YWRKFG-[AEA]-[(D)Lys]- NH <sub>2</sub>	****	****	***
694	Ac-Cyclo-[[Abu]-QTWQC]-YWRENG-[AEA]-[(D)Lys]- NH <sub>2</sub>	****	****	***
695	Ac-Cyclo-[[Abu]-RTWQ-[Pen]]-YWRKFG-[AEA]-[(D)Lys]- NH <sub>2</sub>	*****		
696	Ac- Cyclo-[[Pen]-RTWQ-[Abu]]-YWRKFG-[AEA]-[(D) Lys]-NH <sub>2</sub>	****		
697	Ac-Cyclo-[(D)Cys]-RTWQ-[Abu]]-YWRKFG-[AEA]-[(D)-Lys]- NH <sub>2</sub>	****		
698	Ac-Cyclo-[[Abu]-QTWQC]-YW-[Orn]-[Dap]-NG-[AEA]-[(D) Lys]- NH <sub>2</sub>	****		
699	Ac-Cyclo-[[Abu]-QTWQC]-YW-[hLeu]-ENG- NH <sub>2</sub>	***		**
700	Ac-Cyclo-[Abu]-QTWQ-(D)Cys]]-YW-[hLeu]-ENG-NH <sub>2</sub>	*****		
701	Ac-Cyclo-[[Abu]-QTWQ-[Pen]]-YW-[hLeu]-ENG-NH <sub>2</sub>	*****		
702	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	****	*
703	Ac-Cyclo-[[Abu]-QTWQC]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	**	***	*
704	Ac-Cyclo-[[Abu]-QTWQC]-Y-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	**	*
705	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	**	*
706	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	**	***	*
707	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	***	***	*
708	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	***	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
709	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	**	**
710	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	***	**
711	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[1-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
712	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	**	*
713	Ac-Cyclo-[[Abu]-QTWQC]-YW-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
714	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[(D)Asn]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	****	***
715	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-Phenoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
716	Ac-Cyclo-[[Abu]-QTWQC]-[hPhe(3,4-dimethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****		
717	Ac-Cyclo-[[Abu]-QTWQC]-[DMT]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****		
718	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	***	*
719	Ac-Cyclo-[[Abu]-QTWQC]-Phe(3,4-Cl <sub>2</sub> )[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		***
720	Ac-Cyclo-[[Abu]-QTWQ-[Pen]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****	****	***
721	Ac-Cyclo-[[Abu]-QTWQ-[Pen]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	****	***
722	Ac-Cyclo-[[Pen]-QTWQ-[Abu]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>			
723	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Trp(2,5,7-tri-tert-Butyl)]-[ $\alpha$ -	>10,000		

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
	MeLys]-ENG-NH <sub>2</sub>			
724	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]- [Phe(4-Oallyl)]-[\alpha-MeLys]-ENG-NH <sub>2</sub>	****		
725	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Tyr(3-tBu)]-[\alpha-MeLys]-ENG-NH <sub>2</sub>	***	****	**
726	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Phe(4-tBu)]-[\alpha-MeLys]-ENG-NH <sub>2</sub>	*****		
727	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Phe(4-guanidino)]-[\alpha-MeLys]-ENG-NH <sub>2</sub>	****		
728	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Phe(Bzl)]-[\alpha-MeLys]-ENG-NH <sub>2</sub>	****		
729	Ac-Cyclo-[[Abu]-QTWQC]-[Tyr(3-tBu)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	>10,000		
780	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-tBu)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	*****		
781	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-guanidino)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	***	***	***
782	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	**	**	*
783	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-CO <sub>2</sub> H)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	****		
784	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-phenoxy)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	***	***	**
785	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-CN)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	***		***
786	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-Br)]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	***	***	***
787	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-NH <sub>2</sub> )]-W-[\alpha-MeLys]-ENG-NH <sub>2</sub>	***	***	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
788	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-Phe(4-Me)-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****		
789	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	***	**
790	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	**	*
791	Ac-Cyclo-[[Abu]-QTWQC]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	****	*
792	Ac-Cyclo-[[Abu]-QTWQC]-[Bip]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
793	Ac-Cyclo-[[Abu]-QTWQC]-Cha-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****		
794	Ac-Cyclo-[[Abu]-QTWQC]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
795	Ac-Cyclo-[[Abu]-QTWQC]-[4-Pyridylalanine]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
796	Ac-Cyclo-[[Abu]-QTWQC]-[ $\beta$ -homoTyr]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	~10000		
797	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	**	*
798	Ac-Cyclo-[[Abu]-QTWQC]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	***	
799	Ac-Cyclo-[[Abu]-QT-[2-Nal]-QC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
800	Ac-Cyclo-[[Abu]-QT-[1-Nal]-QC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
801	Ac-Cyclo-[[Abu]-QTYQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	~10000		
802	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>			

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
803	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGGE-NH <sub>2</sub>	***		
804	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGAE-NH <sub>2</sub>			
805	Ac-Cyclo-[[Abu]-STWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGGE-NH <sub>2</sub>			
806	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGGE-NH <sub>2</sub>			
807	Ac-Cyclo-[[Abu]-QTWQC]-Y-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGGE-NH <sub>2</sub>			
808	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NS-NH <sub>2</sub>	**	**	*
809	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NA-NH <sub>2</sub>	*	**	*
810	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	***	*
811	Ac-Cyclo-[[Abu]-QTWQC]-[Phe-4-N <sub>3</sub> ]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	**
812	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-QG-NH <sub>2</sub>	***	***	*
813	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-[Cit]-G-NH <sub>2</sub>			
814	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-VNG-NH <sub>2</sub>	***	***	*
815	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
816	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Orn]-[Dap]-NG-NH <sub>2</sub>	****		
817	Ac-Cyclo-[[Abu]-NTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	***	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
818	Ac-Cyclo-[[Abu]-QT-[Bip]-QC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	~10000		
819	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Cha]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***	*
820	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Chg]-[Lys(Ac)]-NG-NH <sub>2</sub>	***		
821	Ac-Cyclo-[[Abu]-QT-[Octgly]-QC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>10000		
822	Ac-Cyclo-[[Abu]-QTWQC]-[Octgly]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	~10000		
823	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Octgly]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	~10000		
824	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGE-NH <sub>2</sub>	***	***	*
825	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NAE-NH <sub>2</sub>	*	**	*
826	Ac-Cyclo-[[Abu]-STWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGE-NH <sub>2</sub>	***	***	***
827	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGE-NH <sub>2</sub>	****		
828	Ac-Cyclo-[[Abu]-QTWQC]-Y-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NGE-NH <sub>2</sub>	***		
829	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*	*	*
830	Ac-Cyclo-[[Abu]-QTQQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
831	Ac-Cyclo-[[Abu]-QTHQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
832	Ac-Cyclo-[[Abu]-QT-[hPhe]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
833	Ac-Cyclo-[[Abu]-QT-[Glu(Bzl)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
834	Ac-Cyclo-[[Abu]-QT-[Bip]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
835	Ac-Cyclo-[[Abu]-QT-[Tic]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
836	Ac-Cyclo-[[Abu]-QT-[Phe[4-(2-aminoethoxy)]]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
837	Ac-Cyclo-[[Abu]-QT-[Phe(3,4-Cl <sub>2</sub> )]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
838	Ac-Cyclo-[[Abu]-QT-[Phe(4-OMe)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
839	Ac-Cyclo-[[Abu]-QT-[Orn(Benzyl)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
840	Ac-Cyclo-[[Abu]-QT-[Orn(Benzaldehyde)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>			
841	Ac-Cyclo-[[Abu]-QTWQC]-[PheOCH <sub>2</sub> CH <sub>2</sub> NHAc]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>			
842	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>			
843	Ac-Cyclo-[[Abu]-QT-[5-hydroxyTrp]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	~3000		
844	Ac-Cyclo-[[Abu]-QT-[6-chloroTrp]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	**	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
845	Ac-Cyclo-[[Abu]-QT-[N-MeTrp]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
846	Ac-Cyclo-[[Abu]-QT-[1,2,3,4-tetrahydro-norharman]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****		
847	Ac-Cyclo-[[Abu]-QT-[Phe(4-CO2H)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
848	Ac-Cyclo-[[Abu]-QT-[Ph(4-CONH2)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
849	Ac-Cyclo-[[Abu]-QT-[Ph(4-CONH2)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
850	Ac-Cyclo-[[Abu]-QT-[Phe(3,4-OMe)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	~3000		
851	Ac-Cyclo-[[Abu]-QT-[ $\alpha$ -MePhe]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****		
852	Ac-Cyclo-[[Abu]-QT-[Phe(4-CF3)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	~3000		
853	Ac-Cyclo-[[Abu]-QT-[Phe(4-tBu)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>3000		
854	Ac-Cyclo-[[Abu]-QT-[Phe(2,4-Me2)]-QC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	****		
855	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-DNG-NH <sub>2</sub>	*	**	*
856	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-QNG-NH <sub>2</sub>	*		*
857	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Benzoic acid)]-NG-NH <sub>2</sub>	*		*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
858	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(succinic acid)]-NG-NH <sub>2</sub>	*	**	*
859	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(glutaric acid)]-NG-NH <sub>2</sub>	*		*
860	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(pyroglutamic acid)]-NG-NH <sub>2</sub>	*		*
861	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(isovaleric acid)]-NG-NH <sub>2</sub>	*	**	*
862	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Palm)]-NG-NH <sub>2</sub>	~3000		
863	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-Lys[(PEG1)]-NG-NH <sub>2</sub>			
864	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(PEG2)]-NG-NH <sub>2</sub>			
865	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(Benzoic acid)]-NG-NH <sub>2</sub>			
866	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(succinic acid)]-NG-NH <sub>2</sub>			
867	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(glutaric acid)]-NG-NH <sub>2</sub>			
868	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(pyroglutamic acid)]-NG-NH <sub>2</sub>	**		*
869	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-Dap(IVA)NG-NH <sub>2</sub>			
870	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(PEG1)]-NG-NH <sub>2</sub>			
871	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(PEG2)]-NG-NH <sub>2</sub>			

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
872	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Dap(PEG2-Ac)]-NG-NH <sub>2</sub>	**		**
873	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-[AEA]-[(D)Lys]-NH <sub>2</sub>	*		*
874	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-[(D)Lys]-NH <sub>2</sub>	*		*
875	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-[AEA]-NH <sub>2</sub>	*		*
876	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-QG-NH <sub>2</sub>	**		*
877	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-QNG-NH <sub>2</sub>	*	**	*
878	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	*		*
879	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-1-Nal[Aib]-[Lys(Ac)]-NG-NH <sub>2</sub>	***		**
880	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>	*	**	*
881	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-KNG-NH <sub>2</sub>	*		*
882	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(4-CO <sub>2</sub> H)]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
883	Ac-Cyclo-[[Abu]-[Dap]-TCQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(4-Phenoxy)]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
884	Ac-Cyclo-[[Abu]-DapTCQC]-[Phe[4-(2-aminoethoxy)]]-[Phe[4-(2-aminoethoxy)]]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
885	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
886	Ac-Cyclo-[[Abu]-DabTWQC]-[Phe[4-(2-aminoethoxy)]]-[hPhe]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>1000		
887	Ac-Cyclo-[[Abu]-DapTWQC]-[Phe[4-(2-aminoethoxy)]]-[Glu(Bzl)]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
888	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -Me-Orn]-ENG-NH <sub>2</sub>	**		*
889	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*		*
890	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -Me-Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	**		**
891	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -Me-Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	*		*
892	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*	*
893	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	**		**
894	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Orn]-ENG-NH <sub>2</sub>	***		
895	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Orn]-[Dap]-NG-NH <sub>2</sub>	****		
896	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Orn]-[Dap(Ac)]-NG-NH <sub>2</sub>	****		
897	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Orn]-[Dap]-NG-NH <sub>2</sub>	***		***
898	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Orn]-[Dap(Ac)]-NG-NH <sub>2</sub>	***		
899	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[hLeu]-ENG-NH <sub>2</sub>	**		*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
900	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*	*
901	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -Me-Leu]-ENG-NH <sub>2</sub>	*		*
902	Succinyl-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*	*
903	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-[Dap]-G-NH <sub>2</sub>	*****		
904	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-[6-amino-1,4-diazepane-2,5-dione]-NH <sub>2</sub>	***		*
905	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-Chg-[Lys(Ac)]-NG-NH <sub>2</sub>	***		
906	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	*	*	*
907	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(4-CONH <sub>2</sub> )]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****		
908	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(3,4-OMe <sub>2</sub> )]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	***	*
909	Ac-Cyclo-[[Abu]-[Dap]-TWQC]-[Phe[4-(2-aminoethoxy)]]-[Tic]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
910	Ac-Cyclo-[[Abu]-DapTWQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(3,4-Cl <sub>2</sub> )]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***		
911	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENQ-NH <sub>2</sub>	*	*	*
912	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
913	Ac-Cyclo-[[Abu]-TTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*		*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
914	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -Me-Gly(Ethyl)] Lys(Ac)]-NG-NH <sub>2</sub>	*		*
915	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*	*
916	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeSer]-[Lys(Ac)]-NG-NH <sub>2</sub>	*		*
917	Ac-Cyclo-[[Abu]-QTDapQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
918	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[Dap]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
919	Ac-Cyclo-[[Abu]-QTRQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
920	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-R-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
921	Ac-Cyclo-[[Abu]-QTDapQC]-[Phe[4-(2-aminoethoxy)]]-[Dap]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
922	Ac-Cyclo-[[Abu]-QTDQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
923	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-D-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
924	Ac-Cyclo-[[Abu]-QTDQC]-[Phe[4-(2-aminoethoxy)]]-D-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
925	Ac-(D)Lys-[Cyclo-[[Abu]-QTWQC]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	*	**	*
926	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-RNG-NH <sub>2</sub>	**		*
927	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Orn]-NG-NH <sub>2</sub>	*		*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
928	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-KNG-NH <sub>2</sub>	*		*
929	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-hRNG-NH <sub>2</sub>	*		*
930	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*	*	*
931	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Cit]-[Dap]-NG-NH <sub>2</sub>	**		*
932	Ac-Cyclo-[[Abu]-[ $\alpha$ -Me-Orn]-TWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***		**
933	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-NNG-NH <sub>2</sub>	*		*
934	Ac-Cyclo-[[Abu]-STWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-KNGGE-NH <sub>2</sub>	****		
935	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENQ-NH <sub>2</sub>	*	*	*
936	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENN-NH <sub>2</sub>	*	*	*
937	Ac-Cyclo-[[Abu]-TWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>			
938	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-Me)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*		*
939	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(3-Me)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**		*
940	Ac-Cyclo-[[Abu]-QTWQC]-[hTyr]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****		
941	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[ $\alpha$ -MeTrp]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	~3000		

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
942	Ac-Cyclo-[[Abu]-[ $\alpha$ -MeSer]-TWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***		**
943	Ac-Cyclo-[[Abu]-Q-[ $\alpha$ -MeSer]-WQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
944	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[ $\alpha$ -MePhe]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	>3000		
945	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Aib]-ENG-NH <sub>2</sub>	**		*
946	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NG-NH <sub>2</sub>	*		*
947	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-E-[Dap(Ac)]-G-NH <sub>2</sub>	**		*
948	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-E-[Dab(Ac)]-G-NH <sub>2</sub>	*	*	*
949	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-E-[Lys(Ac)]-G-NH <sub>2</sub>	**		*
950	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Aib]-ENN-NH <sub>2</sub>	**		*
951	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-ENN-NH <sub>2</sub>	*	*	*
952	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(3,4-OMe <sub>2</sub> )]-[Aib]-ENG-NH <sub>2</sub>	***		**
953	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(3,4-Cl <sub>2</sub> )]-[Aib]-ENN-NH <sub>2</sub>	***		*
954	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-[Cit]-NN-NH <sub>2</sub>	*	*	*
955	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
956	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-Me)]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	**		*
957	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(3,4-F <sub>2</sub> )]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	***		**
958	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(3-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	****		
959	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(2,4-Cl <sub>2</sub> )]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	****		
960	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(3-Me)]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	**		*
961	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-Cl)]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	**		*
962	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-F)]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	****		
963	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(2,4-Cl <sub>2</sub> , 4-OBz)]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	*****		
964	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-[(D)Lys]-NH <sub>2</sub>	***		**
965	Ac-E-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
966	Ac-(D)Glu-[Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
967	Ac-Arg-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*		*
968	Ac-[(D)Arg]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
969	Ac-F-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
970	Ac-[(D)Phe]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
971	Ac-[2-Nal]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	**	*
972	Ac-T-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
973	Ac-Leu-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
974	Ac-[(D)Gln]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*	*	*
975	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acpc]-ENN-NH <sub>2</sub>			**
976	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acbc]-ENN-NH <sub>2</sub>		*	*
977	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Achc]-ENN-NH <sub>2</sub>		*	*
978	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acvc]-ENN-NH <sub>2</sub>		*	*
979	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-piperidine]-ENN-NH <sub>2</sub>			*
980	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*	*	*
981	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NG-NH <sub>2</sub>			*
982	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>			*
983	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-QNG-NH <sub>2</sub>			*
984	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-QN-[ $\beta$ Ala]-NH <sub>2</sub>			*

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
985	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-QDG-NH <sub>2</sub>			***
986	Ac-Cyclo-[[Abu]-QTCQC]-cyclo([Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-QD)-G-NH <sub>2</sub>			****
987	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-QN-[ $\beta$ Ala]-NH <sub>2</sub>			*
988	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[1,2,3,4-tetrahydro-norharman]-[Aib]-QNG-NH <sub>2</sub>			
989	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[5-hydroxyTrp]-[Aib]-QNG-NH <sub>2</sub>			**
990	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-[Asn(isobutyl)]-G-N-NH <sub>2</sub>			***
991	Ac-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-[Asp(1,4-diaminoethane)]-G-NH <sub>2</sub>			***
992	Ac-(D)Phe-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			
993	Ac-[(D)Arg]-Cyclo-[[Abu]-QTCQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=>1000 nM.

Table E5C. IC50 of Illustrative Thioether Peptide Dimers Synthesized

SEQ ID NO.	Linker Moiety	Sequence	Human ELISA (nM)	Rat ELISA A (nM)	pStat3 HTRF (nM)

994	DIG through (D)Lys	[Ac-[(D)Lys]-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub> ] <sub>2</sub> DIG	*	***	*
995	DIG through Phe[4-(2-aminoethoxy)]	[Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-QNG-NH <sub>2</sub> ] <sub>2</sub> DIG	***		**
996	DIG through $\alpha$ -MeLys	[Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> DIG	*	**	*
997	PEG25 through $\alpha$ -MeLys	[Ac-[(D)Lys]-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> PEG25	*	**	*
998	DIG through (D)Lys	[Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OBzl)]-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> DIG			*
999	PEG25 through (D)Lys	[Ac-Cyclo-[[Abu]-QTWQC]-Y(Bzl)-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> PEG25			*
	Alexa488-PEG4 through [D-Arg]	[D-Arg]-Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>			*

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=>1000 nM.

Table E5D. Illustrative Thioether Peptides

SEQ ID NO.	Sequence	Human IL23R / IL23 ELISA (nM)
993	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>	*
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]]-END-NH <sub>2</sub>	**
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDN-NH <sub>2</sub>	***
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>	**
	Ac-[D-Arg]- Cyclo-[[Abu]-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>	**

	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDD-NH <sub>2</sub>	***
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-END-NH <sub>2</sub>	**
	Ac-[D-Arg]- Cyclo-[[Abu]-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[Tetrahydropyran-A]-END-NH <sub>2</sub>	**
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDN-NH <sub>2</sub>	***
	Ac-[D-Arg]- Cyclo-[[Abu]-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDN-NH <sub>2</sub>	***
	Ac-[D-Arg]- Cyclo-[Abu-ETWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-NH <sub>2</sub>	**
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-OH	**
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-END-OH	***
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-EDN-OH	***
	Ac-[D-Arg]- Cyclo-[[Abu]-QTWEC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-OH	**
	Ac-[D-Arg]- Cyclo-[[Abu]-ETWQC]-[Phe(4-2ae)]-[2-Nal]-[THP]-ENN-OH	**


\* =  $\leq$  1 nM; \*\* = 1 nM - 10 nM; \*\*\* = 10 nM - 100 nM

Table E6. IC50 of Peptide Inhibitors (Ring Closing Metathesis)

SEQ ID NO.	Sequence/Structure	Human ELISA (nM)
1000		~ 20000
1001		~30000
1002		*****

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=1000-10,000 nM.

Table E7. IC<sub>50</sub> of Illustrative Peptides Containing Cyclic amides (side chain cyclizations)



SEQ ID NO.	Sequence	Human ELISA (nM)
1009	Ac-Cyclo-[[Dap]-QTWQE]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000
1010	Ac-Cyclo-[EQTWQ-[Dab]]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>6000
1011	Ac-Cyclo-[EQTWQ-[Dap]]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000
1012	Ac-Cyclo-[[Dab]QTWQE]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~30000
1013	Ac-Cyclo-[[Dap]-QTWQ-[(D)Asp]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000
1014	Ac-Cyclo-[[Dap]-QTWQD]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>30000

1015	Ac-Cyclo-[[DQTWQ-[Dab]]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000
1016	Ac-Cyclo-[[Dab]QTWQD]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>6000
1017	Ac-Cyclo-[[[(D)Dab]-QTWQ-[(D)Asp]]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~6000
1018	Ac-Cyclo-[[[(D)Asp]-QTWQ-[(D)Dab]]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~1400
1019	Ac-Cyclo-[[[(D)Asp]-QTWQ-[(D)Dap]]-YWRENG-[AEA]-[(D)Lys]-NH <sub>2</sub>	~30000


Table E8. IC50 of Illustrative Peptides Containing the Ac-[Pen]-XXWXXXXXX Motif and Ac-XXXWX-[Pen]-XXXX analogues

SEQ ID NO.	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
1020	Ac-[Pen]-ADWVCYWHTFG-NH <sub>2</sub>	*****		
1021	Ac-CADWV-[Pen]-YWHTFG-NH <sub>2</sub>	*****		
1022	Ac-[(D)Pen]-ADWVCYWHTFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	****	*****	****
1023	Ac-CADWV-[(D)Pen]-YWHTFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	>30000	*****	****
1024	Ac-[Pen]-RTWQCYWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	****	****	****
1025	Ac-ACDWV-[Pen]-YWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	*****		
1026	Ac-A-[Pen]-DWVCYWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	****		
1027	Ac-A-[hCys]-DWV-[Pen]-YWRKFG-[AEA]-[(D)-Lys]-NH <sub>2</sub>	~30000		
1028	Ac-CQTWQ-[Pen]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	****	****	
1029	Ac-CQTWQ-[Pen]-YW-[(D)Asn]-ENG-NH <sub>2</sub>	*****		

\*=<10nM; \*\*=10-25 nM \*\*\* = 25-100 nM, \*\*\*\* = 100-1000 nM, \*\*\*\*\*=1000-10,000 nM.

[00698] SAR analysis of the activities of the peptide inhibitors tested indicated that the CXXXXC disulphide is associated with high activity. The two Trp residues and the Phe residue are also associated with high activity, but it is recognized that these amino acids can be readily exchanged with similar homologs (e.g., 1-Nal substituted for Trp and/or Phe substituted for Tyr). In addition, the data suggested that the presence of one or more basic residues at the C-terminus

is associated with high activity. Also, His-9 can be replaced by Arg or another homolog and maintain or improve activity. The schematic below provides one illustrative consensus sequence showing certain residues associated with high activity.



### EXAMPLE 3

#### STABILITY OF PEPTIDE INHIBITORS IN SIMULATED INTESTINAL FLUID (SIF), SIMULATED GASTRIC FLUID (SGF) AND REDOX CONDITIONS

[00699] Studies were carried out in simulated intestinal fluid (SIF) and simulated gastric fluid (SGF) to evaluate gastric stability of the peptide inhibitors of the present invention. In addition, studies were carried out to assess redox stability of the peptide inhibitors of the present invention.

[00700] SIF was prepared by adding 6.8 g of monobasic potassium phosphate and 10.0 g of pancreatin to 1.0 L of water. After dissolution, the pH was adjusted to 6.8 using NaOH. DMSO stocks (2 mM) were first prepared for the test compounds. Aliquots of the DMSO solutions were dosed into 6 individual tubes, each containing 0.5 mL of SIF, which is pre-warmed to 37°C. The final test compound concentration was 20 µM. The vials were kept in a benchtop Thermomixer® for the duration of the experiment. At each timepoint (0, 5, 10, 20, 40, 60, or 360 minutes or 24 hours), 1.0 mL of acetonitrile containing 1% formic acid was added to one vial to terminate the reaction. Samples were stored at 4°C until the end of the experiment. After the final timepoint is sampled, the tubes were mixed and then centrifuged at 3,000 rpm for 10 minutes. Aliquots of the supernatant were removed, diluted 1:1 into distilled water containing internal standard, and analyzed by LCMS/MS. Percent remaining at each timepoint was calculated based on the peak area response ratio of test to compound to internal standard. Time 0 was set to 100%, and all later timepoints were calculated relative to time 0. Half-lives were calculated by fitting to a first-order exponential decay equation using Graphpad. Stability in SIF assays is shown in Tables E9 and E10.

[00701] SGF was prepared by adding 20 mg NaCl, 32 mg porcine pepsin (MP Biochemicals, catalog 02102599), and 70 $\mu$ l HCl to 10ml water (final pH=2). Aliquots of SGF (0.5ml each) were pre-warmed at 37°C. To start the reaction, 1 $\mu$ l of peptide stock solution (10mM in DMSO) was added to 0.5ml SGF and thoroughly mixed such that the final peptide concentration was 20 $\mu$ M. The reactions were incubated at 37°C with gentle shaking. At each time point (0, 15, 30, 60 min) 50 $\mu$ l aliquots were removed and added to 200 ul acetonitrile containing 0.1% formic acid to quench the reaction. Samples are stored at 4°C until the end of the experiment and centrifuged at 10,000 rpm for 5 minutes. Aliquots of the supernatant were removed, diluted 1:1 into distilled water containing internal standard, and analyzed by LCMS/MS. Percent remaining at each timepoint was calculated based on the peak area response ratio of test to compound to internal standard. Time 0 was set to 100%, and all later timepoints were calculated relative to time 0. Half-lives were calculated by fitting to a first-order exponential decay equation using GraphPad. Stability in SGF assays is shown in Tables E9 and E10.

Table E9. Stability of Illustrative Peptides containing the Ac-[Pen]-XXWX-[Pen]-XXXX Motif and Analogues in Simulated Intestinal Fluid (SIF) and Simulated Gastric Fluid (SGF)

SEQ ID NO:	Sequence	SGF t1/2 (min)	SIF t1/2 (min)
549	Ac-[Pen]-QTWQ-[Pen]-YW-[hLeu]-ENG-NH <sub>2</sub>		*****§
1030	Ac-[Pen]-QTWQ-[Pen]-YWN-Me-RENG-NH <sub>2</sub>		****§
551	Ac-[Pen]-QTWQ-[Pen]-YW-[hLeu]-ENG-NH <sub>2</sub>		*****§
552	Ac-[Pen]-QTWQ-[Pen]-YW-[N-MeArg]-ENG-NH <sub>2</sub>		***
554	Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>		**
1028	Ac-CQTWQ-[Pen]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>		*****
555	Ac-[Pen]-QTWQ-[Pen]-YW-[(D)Asn]-ENG-NH <sub>2</sub>		**
1029	Ac-CQTWQ-[Pen]-YW-[(D)Asn]-ENG-NH <sub>2</sub>		*****
556	Ac-[Pen]-QTWQ-[Pen]-Y-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>		**
557	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	**
558	Ac-[Pen]-QTWQ-[Pen]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>		**

SEQ ID NO:	Sequence	SGF t1/2 (min)	SIF t1/2 (min)
559	Ac-[Pen]-QTWQ-[Pen]-Y-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>		**
560	Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>		**
561	Ac-[Pen]-QTWQ-[Pen]-Y-[1-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>		**
1031	Ac-[Pen]-QTWQ-[Pen]-[[Phe(4-OMe)](OMe)]-[2-Nal]-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>		*
563	Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*
1032	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>		*
565	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*
566	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[1-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>		*
1033	succinic anhydride-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	*
585	pyroglutamic acid-[Pen]-QTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	*
1034	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
601	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENA-NH <sub>2</sub>	*	*
602	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	**	***
603	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-QNN-NH <sub>2</sub>	*	*
604	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>	*	*
605	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-Aib-[Lys(Ac)]-NN-NH <sub>2</sub>	**	***
606	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>	*	*
607	Ac-[Pen]-Dap(Ac)TWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]]-[2-Nal]-[ $\alpha$ -	*	*

SEQ ID NO:	Sequence	SGF t1/2 (min)	SIF t1/2 (min)
	MeLys(Ac)]-ENG-NH <sub>2</sub>		
608	Ac-[Pen]-[ $\alpha$ -MeOrn(Ac)]-TWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	**	****
609	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*
610	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
611	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	*	**
612	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENA-NH <sub>2</sub>	*	*
613	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
614	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-QNN-NH <sub>2</sub>	*	*
615	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>	*	*
616	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
617	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>	*	*
522	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> DIG	****	*
618	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>	*	***
619	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-ENA-NH <sub>2</sub>	*****	*****
620	Ac-[Pen]-TTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
625	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	**
628	Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*	*

SEQ ID NO:	Sequence	SGF t1/2 (min)	SIF t1/2 (min)
630	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-ENN-NH <sub>2</sub>	*	***
631	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-ENA-NH <sub>2</sub>	*****	*****
632	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
633	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*	**
634	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>	*	*
636	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>	*	*
637	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
638	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NQ-NH <sub>2</sub>	*	*
639	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*	*
640	Ac-[Pen]-NTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*****	*****
641	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*****	*****
669	Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[α-MeVal]-[Lys(Ac)]-NN-NH <sub>2</sub>	**	*
534	[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[α-MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG	**	*
1035	Ac-[(D)Phe]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
676	Ac-[(D)Phe]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	**	*
682	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	**	***
683	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-Lys(Ac)]-NN-NH <sub>2</sub>	**	*

SEQ ID NO:	Sequence	SGF t1/2 (min)	SIF t1/2 (min)
684	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Achc]- [Lys(Ac)]-NN-NH <sub>2</sub>	*	*
1036	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Acvc]- [Lys(Ac)]-NN-NH <sub>2</sub>	*	*
686	Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
688	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
689	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	**
1037	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[Acvc]-[Lys(Ac)]-NN-NH <sub>2</sub>	**	*
671	Ac-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*
535	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	**	**
536	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	*	*
537	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Achc]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	**	***
539	[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLeu]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	**	**

§ the matrix used is 100 fold dilution of standard SIF concentration

\*=>360min; \*\*=180-360min; \*\*\*=120-180min; \*\*\*\*=<60-120min; \*\*\*\*\*=<60min

Table E10. Stability of Illustrative Peptides Containing Thioethers Motif and Analogues Within Simulated Intestinal Fluid (SIF) and Simulated Gastric Fluid (SGF)

SEQ ID NO:	Sequence	SIF t1/2 (min)	SGF t1/2 (min)
692	Ac-Cyclo-[[Abu]RTWQC]-YWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	*****	
694	Ac-Cyclo-[[Abu]-QTWQC]-YWRENG-[AEA]-[(D)Lys]- NH <sub>2</sub>	*****	
699	Ac-Cyclo-[[Abu]-QTWQC]-YW-[hLeu]-ENG- NH <sub>2</sub>	*****	ND
700	Ac-Cyclo-[Abu]-QTWQ-(D)Cys]-YW-[hLeu]-ENG-NH <sub>2</sub>	*** §	

SEQ ID NO:	Sequence	SIF t1/2 (min)	SGF t1/2 (min)
701	Ac-Cyclo-[[Abu]-QTWQ-[Pen]]-YW-[hLeu]-ENG-NH <sub>2</sub>	*****	
703	Ac-Cyclo-[[Abu]-QTWQC]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	*****	
704	Ac-Cyclo-[[Abu]-QTWQC]-Y-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*****	
702	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	***	****
706	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	***	****
707	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeOrn]-ENG-NH <sub>2</sub>	**	****
702	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	****
709	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	****
710	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*	****
711	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[1-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	****
712	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	****
713	Ac-Cyclo-[[Abu]-QTWQC]-YW-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	
714	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[(D)Asn]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	
715	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-Phenoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	
716	Ac-Cyclo-[[Abu]-QTWQC]-[hPhe(3,4-dimethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	
717	Ac-Cyclo-[[Abu]-QTWQC]-[DMT]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	
718	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	****
719	Ac-Cyclo-[[Abu]-QTWQC]-Phe(3,4-Cl <sub>2</sub> )[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	
720	Ac-Cyclo-[[Abu]-QTWQ-[Pen]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	***
721	Ac-Cyclo-[[Abu]-QTWQ-[Pen]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	***
782	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*	***

SEQ ID NO:	Sequence	SIF t1/2 (min)	SGF t1/2 (min)
790	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	*****
791	Ac-Cyclo-[[Abu]-QTWQC]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeOrn]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	ND
794	Ac-Cyclo-[[Abu]-QTWQC]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	ND
797	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****	*****
798	Ac-Cyclo-[[Abu]-QTWQC]-[2-Nal]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	ND
810	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Aib]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	
815	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	
820	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Chg]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	
822	Ac-Cyclo-[[Abu]-QTWQC]-[Octgly]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****	
823	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Octgly]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	****	
823	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[Octgly]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****	
829	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*	*
857	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Benzoic acid)]-NG-NH <sub>2</sub>	**	*
861	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(isovaleric acid)]-NG-NH <sub>2</sub>	*	*
876	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-QG-NH <sub>2</sub>	***	*****
877	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-QNG-NH <sub>2</sub>	***	**
878	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-ENG-NH <sub>2</sub>	*****	***
879	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-1-Nal[Aib]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	*
880	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>	****	*****
891	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -Me-Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	**
892	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-	*	*

SEQ ID NO:	Sequence	SIF t1/2 (min)	SGF t1/2 (min)
	[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>		
893	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Orn]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****	
894	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Orn]-ENG-NH <sub>2</sub>	*****	
895	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Orn]-[Dap]-NG-NH <sub>2</sub>	*****	
896	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[Orn]-[Dap(Ac)]-NG-NH <sub>2</sub>	*****	
897	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Orn]-[Dap]-NG-NH <sub>2</sub>	*****	
898	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Orn]-[Dap(Ac)]-NG-NH <sub>2</sub>	*****	
899	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[hLeu]-ENG-NH <sub>2</sub>	*****	****
900	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NG-NH <sub>2</sub>	**	****
901	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -Me-Leu]-ENG-NH <sub>2</sub>	*	**
902	Succinyl-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	**
906	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-ENG-NH <sub>2</sub>	****	****
820	Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[Chg]-[Lys(Ac)]-NG-NH <sub>2</sub>	*	**
911	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENQ-NH <sub>2</sub>	**	**
912	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	**	**
913	Ac-Cyclo-[[Abu]-TTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	**	**
914	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -Me-Gly(Ethyl)] Lys(Ac)]-NG-NH <sub>2</sub>	****	****
915	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NG-NH <sub>2</sub>	***	***
916	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeSer]-[Lys(Ac)]-NG-NH <sub>2</sub>	*****	***
925	Ac-(D)Lys-[Cyclo-[[Abu]-QTWQC]]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	****	****
1039	[Ac-[(D)Lys]-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub> ] <sub>2</sub> DIG : dimerization through (D)Lys	****	****

SEQ ID NO:	Sequence	SIF t1/2 (min)	SGF t1/2 (min)
930	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[hLeu]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*****	ND
933	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-NNG-NH <sub>2</sub>	**	**
946	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NG-NH <sub>2</sub>	****	****
955	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	***	*****
1040	[Ac-Cyclo-[[Abu]-QTWQC]-Y(Bzl)-W-[α-MeLys]-ENG-NH <sub>2</sub> ] <sub>2</sub> ; PEG25 through [α-MeLys]	**	*****
965	Ac-E-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	***
966	Ac-(D)Glu-[Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	***
967	Ac-Arg-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*****	***
1041	Ac-[(D)Arg-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	**
969	Ac-F-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	****	***
970	Ac-[(D)Phe]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	**	***
972	Ac-T-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	***
973	Ac-Leu-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	***
1042	Ac-[(D)Qln]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	***
975	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acpc]-ENN-NH <sub>2</sub>	*****	*
976	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acbc]-ENN-NH <sub>2</sub>	*****	***
1043	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acpc]-ENN-NH <sub>2</sub>	**	*
978	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acvc]-ENN-NH <sub>2</sub>	***	*
979	Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-piperidine]-ENN-NH <sub>2</sub>	*	*
972	Ac-T-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[α-MeLys]-ENN-NH <sub>2</sub>	*	*

§ the matrix used is 100 fold dilution of standard SIF concentration  
 \*=>360min; \*\*=180-360min; \*\*\*=120-180min; \*\*\*\*=<60-120min; \*\*\*\*\*=<60min

[00702] For each peptide tested, the DTT stability assay was conducted by adding 5 $\mu$ l of a 10mM peptide stock solution in DMSO to 1ml of 100mM Tris-Cl, pH 7.5 (final peptide concentration is 50 $\mu$ M). At time 0 min, 5 $\mu$ l of a freshly thawed 100mM DTT solution was added to the incubation tube containing the peptide, such that the final DTT concentration was 0.5mM. The reactions were incubated at room temperature. At different time points up to 120 minutes (20 min, 40 min, 80 min, 120 min), 50 $\mu$ l aliquots were removed, and the reaction was quenched by adding 10 $\mu$ l of 5M acetic acid. To measure disappearance of the parent peptide, the quenched samples (30 $\mu$ l) were analyzed by reverse phase HPLC and UV absorbance at 220nm. The fraction oxidized remaining was graphed versus time, and half-lives were calculated by fitting to a first-order exponential decay equation using Excel. The results of these studies are shown in Table E11. The peptides having half-life >120 min are all considered stable.

Table E11. Stability of Illustrative Peptides in DTT Assay

Sequence	DTT Stability (min)
Ac-CRTWECYWHEFG-NH <sub>2</sub>	<10
Ac-CQTWQCYW-[hLeu]-ENG-NH <sub>2</sub>	*
Ac-CADWVWCYWHFGA-[Azt]-[(D)Lys]-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-RTWQC]-YWRKFG-[AEA]-[(D)Lys]-NH <sub>2</sub>	>120
Ac-[[Pen]-QTWQ-[Pen]-YW-[hLeu]-ENG-NH <sub>2</sub>	>120
Ac-[Pen]-QTWQ-[Pen]-YW-[ $\alpha$ -MeLeu]-ENG-NH <sub>2</sub>	>120
Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>120
Ac-[Pen]-QTWQ-[Pen]-[Phe(4-Ome)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>120
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	>120
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	>120
[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG	>120

\*=10-120 min

#### EXAMPLE 4

## CROSS-REACTIVITY OF PEPTIDE INHIBITORS

[00703] The amino acids of the extracellular domain of the human IL-23R are 95%, 77% and 70% identical to the cyno IL-23R, rat IL-23R and mouse IL-23R, respectively. Interestingly, the mouse receptor contains an insertion of 21 residues that are absent in human, mouse, chimp, dog and cow receptor. These additional amino acids are located in a region where human IL-23R is thought to bind to IL-23.

[00704] To identify peptide inhibitors that cross-reacted with species other than human IL-23R, the ability of certain peptide inhibitors to inhibit human IL-23R, cyno IL-23R, rat IL-23R and mouse IL-23R by ELISA assay. In line with the observation regarding the sequence differences between human IL-23R and mouse IL-23R, the peptide antagonists tested showed a lack of or very weak inhibitory activities in the mouse IL-23R ELISA (see Table E12). In contrast, the antagonists tested to date displayed comparable potency towards the rat receptor and slightly less activity towards the cyno receptor.

[00705] Various bioassays performed to determine the potency, cross reactivity and the selectivity of IL-23R antagonists are described below.

Assays for Selectivity of specific IL-23R AntagonistsHuman IL-12R $\beta$ 1 ELISA

[00706] An assay plate was coated with 100 ng/well of human IL-12R $\beta$ 1\_huFC and incubated overnight at 4°C. The wells were washed, blocked, and washed again. Serial dilutions of test peptides and IL-23 at a final concentration of 2.5 nM were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected with goat anti-p40 polyclonal antibodies, followed by an HRP conjugated donkey anti-goat IgG. Signals were visualized with TMB One Component HRP Membrane Substrate and quenched with 2 M sulfuric acid.

Mouse IL-23R Competitive Binding ELISA

[00707] An assay plate was coated with 50 ng/well of Mouse IL-23R\_huFC and incubated overnight at 4°C. The wells were washed, blocked, and washed again. Serial dilutions of test

peptides and IL-23 at a final concentration of 4 nM were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected with goat anti-p40 polyclonal antibodies, followed by an HRP conjugated donkey anti-goat IgG. Signals were visualized with TMB One Component HRP Membrane Substrate and quenched with 2 M sulfuric acid.

Rat IL-23R Competitive Binding ELISA

[00708] An assay plate was coated with 300 ng/well of Rat IL-23R\_huFC and incubated overnight at 4°C. The wells were washed, blocked, and washed again. Serial dilutions of test peptides and IL-23 at a final concentration of 7 nM were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected with goat anti-p40 polyclonal antibodies, followed by an HRP conjugated donkey anti-goat IgG. Signals were visualized with TMB One Component HRP Membrane Substrate and quenched with 2 M sulfuric acid.

Cyno IL-23R Competitive Binding ELISA

[00709] An assay plate was coated with 50 ng/well of Cyno IL-23R\_huFC and incubated overnight at 4°C. The wells were washed, blocked, and washed again. Serial dilutions of test peptides and IL-23 at a final concentration of 2 nM were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected with goat anti-p40 polyclonal antibodies, followed by an HRP conjugated donkey anti-goat IgG. Signals were visualized with TMB One Component HRP Membrane Substrate and quenched with 2 M sulfuric acid.

Table E12. Cross-Reactivity of Illustrative Peptide Inhibitors

Cmpd. Number	Human IL-23R Activity (nM)		Rodent and Cyno IL-23R Cross Reactivity (nM)		
	ELISA huIL23R IL23	Cell Assay pSTAT3 HTRF	ELISA mouse IL23R IL23	ELISA rat IL23R IL23	ELISA cyno IL23R IL23
22	+	+	-	+	+
197	++	ND	-	++	ND
169	++	++	-	++	+

198	+++	+++	ND	+++	+++
213	+++	+++	ND	+++	ND
219	+++	+++	ND	+++	ND
230	+++	+++	ND	+++	ND

+++ indicates 0-250 nM

++ indicates 251-1000 nM

+ indicates 1001-10,000 nM

- indicates > 25,000 nM

#### EXAMPLE 5

##### NK CELL ASSAY

[00710] Natural killer (NK) cells, purified from human peripheral blood of healthy donors by negative selection (Miltenyi Biotech, Cat # 130-092-657), were cultured in complete media (RPMI 1640 containing 10% FBS, L-glutamine and penicillin-streptomycin) in the presence of IL-2 (RnD, Cat # 202-IL-010/CF) at 25 ng/mL. After 7 days, cells were centrifuged, and resuspended in complete media at 1E6 cells/mL. Recombinant IL-23 at predetermined EC50 to EC75 and IL-18 (RnD, Cat # B003-5) at 10 ng/mL were mixed with varying concentrations of peptides, and added to NK cells seeded at 1E5 cells per well. After 20 to 24 hours, IFN $\gamma$  in the supernatant was quantified using Quantikine ELISA (RnD, Cat # DIF50).

Table E13. IC50 of Illustrative Peptide Inhibitors in Primary Cell Line (NK Cell Assay)

Sequence	NK cell assay (nM)
Ac-Cyclo-[[Abu]-QTWQC]-Y-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-W-[ $\alpha$ -MeLys]-ENG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(isovaleric acid)]-NG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-QNG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NA-NH <sub>2</sub>	*

Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-(acetyl-aminoethoxy))]-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[Phe(3,4-OMe <sub>2</sub> )-[ $\alpha$ -MeLys]-[Lys(Ac)]-NG-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENQ-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NG-NH <sub>2</sub>	*
[Ac-Cyclo-[[Abu]-QTWQC]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub> ]; DIG through $\alpha$ -MeLys	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-[Cit]-NN-NH <sub>2</sub>	*
Ac-[(D)Phe]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*
Ac-T-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLys]-ENN-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acbc]-ENN-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Acpc]-ENN-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Achc]-ENN-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*
Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[ $\alpha$ -MeLeu]-QN-[ $\beta$ Ala]-NH <sub>2</sub>	*
Ac-(D)Phe-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*
Ac-[(D)Arg]-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*

\*=&lt;25nM

Table 14. IC50 of Illustrative Peptides Containing the Ac-[Pen]-XXWX-[Pen]-XXXX Motif and analogues (NK cell assay)

Sequence	NK Cell assay (nM)
Ac-[Pen]-QTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*
Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeVal]-[Lys(Ac)]-NN-NH <sub>2</sub>	**
[Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-[ $\alpha$ -MeVal]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	*
[Ac-[Pen]-QTWQ[Pen]-[Phe[4-(2-acetylaminoethoxy)]-[2-Nal]-K-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG	*
[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-OMe)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG	*
[Ac-[Pen]-QTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-NN-NH <sub>2</sub> ] <sub>2</sub> DIG	*
Ac-[(D)Phe]-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Cit]-NN-NH <sub>2</sub>	*
Ac-[(D)Phe]-[Pen]-NTWQ[Pen]-[Phe(4-OMe)]-[2-Nal]-[Achc]-ENN-NH <sub>2</sub>	*
Ac-[Pen]-NTWQ[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
[Ac-[Pen]-NTWQ-[Pen]-[Phe(4-CONH <sub>2</sub> )]-[2-Nal]-[Aib]-KNN-NH <sub>2</sub> ] <sub>2</sub> DIG	*

\*=<10nM; \*\*=10-25nM

#### EXAMPLE 6

##### BIOASSAY CHARACTERIZATION OF PEPTIDE INHIBITORS

**[00711]** The potency, cross reactivity, and selectivity of certain peptide inhibitors was determined using various bioassays developed for this purpose and described below.

##### Rat Splenocyte Assay

[00712] A new assay developed was the rat splenocyte assay. This assay examined the levels of IL-17A in activated rat splenocytes following stimulation with IL-23 in the presence of test compound.

[00713] Briefly, splenocytes freshly isolated from rat were seeded in 96-well tissue culture plates in complete medium containing IL-1 $\beta$ . Serial dilutions of test compounds were distributed to each well along with rat IL-23 at a final concentration that is between EC50 to EC80 values; plates then were incubated for 3 days at 37 °C in a 5% CO<sub>2</sub> humidified incubator. Changes in IL-17A levels in the supernatants were detected using an ELISA.

#### Rat Colitis Model: 9 Days of 3% DSS-Containing Drinking Water

[00714] There is a body of evidence in the literature supporting the pathogenic role of IL-23/IL-23R signaling in animal models of colitis. For the IL-23 ligand, this requirement has been shown in multiple models, including an IL-10<sup>-/-</sup> spontaneous colitis model, a *Helicobacter hepaticus*-driven colitis model, the anti-CD40 innate colitis model, and the chronic CD45RB<sup>high</sup> CD4 $^{+}$  T-cell transfer model. For the IL-23 receptor, the requirement for colitis development has been shown in the acute models of colitis induced by DSS or by anti-CD40, as well as the chronic CD45RB<sup>high</sup> CD4 $^{+}$  T-cell transfer model. Since certain peptide inhibitors of the present invention do not cross react with the IL-23 receptor from mouse but do recognize that from the rat, a rat model of IBD relevant to the IL-23 pathway was developed.

[00715] In this model, colitis was induced in SD rats by 9 days of *ad lib* exposure to drinking water containing 3% DSS. The disease activity index (DAI) score and ratio of colon weight:colon length were compared between three study groups (n=6 rats/group): vehicle, 3% DSS, and 3% DSS with positive control (sulfasalazine administered at 100 mg/kg PO, QD). The DAI score consisted of ratings from three parameters, including percent body weight loss, stool consistency, and a quantitative hemoccult score, and could achieve a maximum value of 3 units. DSS-exposed animals displayed significantly elevated DAI score (compared to vehicle control) from Day 4 onward, with DAI values peaking at approximately 2.5 by the end of the study (Day 9). Treatment of the DSS-exposed rats with the positive control (sulfasalazine) attenuated the disease score (compared to DSS alone) from Day 5. The differences observed in the terminal

ratio of colon weight:colon length also were significant for DSS-induced disease animals with and without sulfasalazine treatment.

#### Ex Vivo Activity and Stability

[00716] Two peptides (Compound A and Compound B) were selected for use in further biological studies (shown below). One contained a thioether linkage and the other contained a Pen-Pen disulfide bond. The activity, selectivity and *ex vivo* stability profiles of the two compounds are provided herein.


[00717] Assays for selectivity of peptide inhibitors included a human IL-12R $\beta$ 1 ELISA and measurement of the production of IL-12 in PHA activated human PBMC, which are described briefly below.

#### Human IL-12R $\beta$ 1 ELISA


[00718] An assay plate was coated with 100 ng/well of human IL-12R $\beta$ 1\_huFC and incubated overnight at 4°C. The wells were washed, blocked, and washed again. Serial dilutions of test peptides and IL-23 at a final concentration of 2.5 nM were added to each well, and incubated for 2 hours at room temperature. After the wells were washed, bound IL-23 was detected with goat anti-p40 polyclonal antibodies, followed by an HRP conjugated donkey anti-goat IgG. Signals were visualized with TMB One Component HRP Membrane Substrate and quenched with 2 M sulfuric acid. Data from these assays is provided herein.

#### Production of IFN $\gamma$ by IL-12 in PHA Activated Human PBMC

[00719] This assay examined the ability of IL-23R antagonists to neutralize production of IFN $\gamma$  proteins in IL-12-stimulated human PBMCs. IL-23R peptide inhibitors specific to the IL-23/IL-23R pathway are not expected to alter the levels of IFN $\gamma$  produced. Compound A and Compound B were tested in this assay, and a graph showing that they do not alter the levels of IFN $\gamma$  produced at most concentrations tested is provided in Figure 2.



Compound A



Compound B

### In Vivo Activity

[00720] Acute colitis was induced by feeding female Sprague Dawley rats with 3% (wt/vol) DSS dissolved in drinking water. For nine days starting at the same day as DSS, Compounds A or B was administered orally three times per day at 20 mg/kg or 30 mg/kg. Compounds A was also administered intraperitoneally three times per day at 30 mg/kg. A neutralizing anti-IL-23p19 antibody was used as a comparator, and was administered intraperitoneally at 4 mg/kg on the same day and fifth day after starting DSS. To quantify colitis with clinical activity, disease activity index (DAI) was determined daily for each animal as an average of three parameters: body weight change (scale 0-3), stool consistency (scale 0-3) and hemoccult blood (scale 0-3), as shown in Table E15. At necropsy, the entire colon was removed from the cecum to the rectum. The colon was measured for length, flushed with PBS to remove feces, weighed, and opened

longitudinally to determine macroscopic score. The visible damage of the colon was scored on a scale from 0-3, as shown in Table E16.

[00721] Table E17 shows that at Day 7, treatment with Compound A and B significantly improved DAI scores compared to vehicle treated group. FIG. 1 shows results for DAI values from Day 7. In addition, a significant reduction was also observed in the colon weight to colon length ratios, and colon macroscopic scores (FIG. 3). The reduction in inflammation observed with orally delivered peptides was similar to the effects observed from neutralizing anti-IL23p19 monoclonal antibody. Statistical analysis for significance was compared to the vehicle treated group and was determined using Student's T-test (GraphPad Prism). Differences were noted as significant \*p<0.05, \*\*p<0.01, \*\*\*p<0.001, \*\*\*\*p<0.0001.

Table E15. Scoring of the Disease Activity Index

Score	Percent Body Weight Change	Stool Consistency	Hemoccult Score
0	None	Normal	Normal
1	1 to 7	Semi solid	Guaiac+
2	8 to 15	Loose	Bleeding+
3	> 15	Diarrhea	Bleeding++

Table E16. Scoring of Gross Morphologic Damage of the Colon

Score	Gross morphology
0	Normal
1	Erythema
2	Erythema, slight edema, small erosions
3	Two or more bleeding ulcers, inflammation, moderate adhesions
4	Severe ulceration, stenosis with dilations, severe adhesions

Table E17. Disease activity index scores and the individual parameters scores at Day 7, colon weight to length ratios and colon macroscopic scores at Day 9.



	Day 7	Day 9 Necropsy

	Percent Body Weight Change		Stool Consistency		Hemoccult Score		DAI		Colon Weight/Length (g/cm)		Colon Macroscopic Score	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Group												
No DSS	11.00	2.08****	0	0****	0	0****	0	0****	75.51	7.03***	ND	ND
3% DSS, Vehicle	-6.39	1.11	2.00	0.58	1.50	0.50	1.72	0.45	124.36	17.11	1.00	1.00
Anti- IL23p19 mAb	-0.05	1.92****	1.00	0.58*	0.50	0.5*	0.67	0.43**	99.96	16.19*	0.00	0**
Compound A, PO	3.18	2.09****	1.17	0.90	0.50	0.5*	0.56	0.46**	98.38	6.91*	0.00	0**
Compound B, PO	0.13	1.24****	0.83	0.69*	0.67	0.47*	0.61	0.3***	97.36	9.32*	0.00	0**
Compound A, IP	-0.50	1.88***	1.17	0.69	0.83	0.69	0.83	0.54*	104.32	12.45	0.33	0.47

### EXAMPLE 7

#### IN VITRO ASSAYS AND SURFACE PLASMON RESONANCE (SPR) ANALYSIS

[00722] In vitro assays and SPR were performed to further characterize an illustrative compound, Compound C:



Compound C.

[00723] Assays described in previous examples were performed to demonstrate that Compound C is a potent, selective and competitive inhibitor of IL-23R, showing potent inhibition of IL-23-dependent upregulation of phosphor-STAT3 (pSTAT3) in human DB cells and IFN $\gamma$  production in human peripheral blood natural killer (PB NK) cells. In addition, Compound C was selective, showing little inhibition in a cell free ELISA for human IL6R, or in IL-12-dependent production of IFN $\gamma$  in PBMC. Data is shown below in Table E18A. Compound C also cross-reacted with cynomolgus IL-23R (IC<sub>50</sub> 7 nM) and rat IL-23R (IC<sub>50</sub> 17 nM), and inhibited IL-23-dependent IL-17A production in rat splenocytes (IC<sub>50</sub> 130 nM) (data not shown).

Table E18A. In vitro Characterization of Compound C

	IC50				KD	
	pSTAT3 /DB Cell Assay	IFN $\gamma$ /PB NK Primary Cell Assay	IL-6/IL-6R ELISA	IFN $\gamma$ /IL-12 PBMC Cell Assay	IL-23R Surface	IL- 12R $\beta$ 1 Surface
Compound C	4 nM	27nM	>100 uM	>100 uM	2.4nM	None

[00724] Compound C exposure was also restricted to the GI following oral administration to rats does PO at 20 mg/kg, with AUC values of 355 ug.h/g for small intestine mucosa; 77 ug.h/g for colon mucosa; and 0.3 ug.h/mL for plasma, with a 40% recovery in feces.

[00725] Compound C was also stable in a variety of GI fluids and reducing environment, having a SIF half-life of > 24 h; a SGF half-life of > 24 h; a human intestinal fluid half-life of > 24 h, and a half-life of > 2 h in a DTT assay.

[00726] SPR experiments were carried out using a Biacore 2000 instrument and T100 optical biosensors equipped with Biacore CM4 and Xantec HC1500m sensor chips. Recombinant human IL-23R\_huFC (RnD), or recombinant human IL-12R $\beta$ 1\_huFC (RnD) or a mixture of the two receptor subunits were captured on an anti-human IgG surface. Recombinant human IL-23

(Humanzyme) or Compound C was used as the analyte. SPR sensorgrams were fitted to a one to one interaction model, giving rise to a rough estimate of the association rate constant ( $k_{on}$ ), dissociation rate constant ( $k_{off}$ ) and dissociation constant ( $K_D$ ) of the complexes, as shown in Table E11. The data show that Compound C does not bind to IL-12R $\beta$ 1, and binds to IL-23R and the mixed surface of IL-12R $\beta$ 1 and IL-23R with similar potency, at 2.42nM and 2.56nM, respectively. This affinity for IL-23R is comparable to that from IL-23. In contrast, the affinity of IL-23 to the mixed surface is approximately 14x faster than that from Compound C.

Table E18B. Binding characteristics of IL-23 and Compound C for IL-12R $\beta$ 1, IL-23R or mixed IL-12R $\beta$ 1 and IL-23R as determined by SPR.

Surface	IL-23			Compound C		
	$k_a$ (M-1 sec-1)	$k_d$ (sec-1)	$K_D$ (nM)	$k_a$ (M-1 sec-1)	$k_d$ (sec-1)	$K_D$ (nM)
IL-12R $\beta$ 1_huFC	5.01E+05	4.38E-04	0.87	does not bind up to 16.7 uM		
IL-23R_huFC	7.82E+05	0.00132	1.69	1.37E+07	0.033	2.42
IL-12R $\beta$ 1_huFC/IL-23R_huFC	6.31E+05	1.15E-04	0.18	1.59E+07	0.041	2.56

## EXAMPLE 8

### EFFICACY OF IL-23R ANTAGONISTS IN TNBS INDUCED COLITIS IN RAT

[00727] To further evaluate the efficacy of IL-23R antagonists in an animal model of disease, acute colitis was induced by providing 7-week-old female Sprague-Dawley rats with 60 mg/kg 2,4,6-Trinitrobenzenesulfonic acid (TNBS) in 45%-50% ethanol (TNBS/ethanol) administered intrarectally at Day 0. Compound C (described in Example 7) was administered orally three times a day at 20 mg/kg or 6.7 mg/kg and was provided in drinking water at 0.6 mg/mL or 0.2 mg/mL, respectively, for 8 days starting approximately 24 hours (Day -1) prior TNBS inoculation. A neutralizing anti-IL-23p19 antibody was used as a comparator, and was administered intraperitoneally at 4 mg/kg on Day -1 and again on Day 3. All animals received orally PBS (pH 7.4) vehicle which was used to formulate Compound C. The study design is shown in Figure 5.

[00728] To assess the extent of the inflammatory response, animals were observed daily for clinical signs which included percent body weight loss and signs of loose stools or diarrhea. Six days after inoculation of TNBS, rats were sacrificed and the entire colon length and colon weight

from cecum to rectum from each animal were recorded. The severity of colitis was evaluated by a pathologist blinded to the identity of treatments. In addition to the colon wall thickness, the gross colon damage was scored on a 0-4 scale according to Table E19 below, and histopathological scores were determined based on below parameters (Tables E20 and E21).

Table E19. Definitions for colon macroscopic scores

Score	Colon Gross Morphology
0	Normal
1	Erythema
2	Erythema, slight edema, small erosions
3	Two or more bleeding ulcers, inflammation, moderate adhesions
4	Severe ulceration, stenosis with dilations, severe adhesions

Table E20. Definitions for histopathology

Parameter	Definition
Inflammation	Extent and severity of inflammatory cells infiltration, localized and/or diffuse involving full thickness of the colon section (transmural). Inflammatory cells include polymorpho-nuclear leukocytes (neutrophils), mononuclear cells (macrophages + lymphocytes), fibroplasia and neovascularization.
Mucosal Necrosis	Necrosis in the mucosa with loss of surface epithelium, hemorrhage and cellular debris; measured as the length of the lesion on the total length of the colon section to determine % area affected
Gland Loss	% crypt epithelial degeneration with or without superficial mucosal erosion
Colon Thickness	the average thickness of the colon measured transmurally (full thickness) from the mucosal surface to the serosa

Table E21. Scoring criteria

Score	Inflammation
0	Normal tissue, no inflammation
0.5	Very minimal localized infiltrates in the superficial mucosa affecting <2% of the colon section
1	Minimal degree of multifocal infiltrates in the mucosa affecting approximately 2 - 10% of the colon section
2	Mild degree of multifocal infiltrates in the mucosa, submucosa, outer muscle band, and serosa affecting approximately 11 - 25% of the colon section
3	Moderate degree of multifocal infiltrates in the mucosa submucosa, outer muscle band and serosa affecting approximately 26 - 50% of the colon section
4	Marked degree of multifocal to diffuse infiltrates in the mucosa submucosa, outer muscle band and serosa affecting approximately 51- 75% of the colon

	section
5	Sever degree of multifocal to diffuse infiltrates in the mucosa submucosa, outer muscle band and serosa affecting approximately >75% of the colon section

<b>Score</b>	<b>Mucosal Necrosis</b>
0	No Necrosis
0.5	Very minimal and localized region affecting <2% of the total colon section
1	Minimal focal to multifocal regions affecting 2 - 10% of the total colon section
2	Mild focal to multifocal regions affecting 11 - 25% of the total colon section
3	Moderate focal to multifocal regions affecting 26 - 50% of the total colon section
4	Marked focal to multifocal regions affecting 51 - 75% of the total colon section
5	Severe focal to multifocal regions affecting >75% of the total colon section

<b>Score</b>	<b>Gland Loss</b>
0	No loss, normal crypt epithelium and mucosa
0.5	Very minimal loss not exceeding 1 - 2 regions of mucosa/gland affected
1	Minimal, 1-10% regions of mucosa/gland affected
2	Mild, 11-25% regions of mucosa/gland affected
3	Moderate, 26-50% regions of mucosa/gland affected
4	Marked, 51-75% regions of mucosa/gland affected
5	Severe, >75% regions of mucosa/gland affected

<b>Score</b>	<b>Colon Thickness</b>
0	Normal = <350 microns or less
0.5	Very Minimal = 351-400 microns
1	Minimal = 400- 500 microns
2	Mild = 501 - 600 microns
3	Moderate = 601 - 700 microns
4	Marked = 701 - 800 microns
5	Severe = >801 microns

[00729] Compared to the sham group, rats challenged with TNBS suffered acute weight loss, displayed increased incidence of loose stools, and increased colon weight to length ratio. These data were confirmed by the macroscopic examination of colon which revealed mild colonic injury characterized by erythema, edema and small erosions. Treatment with Compound C attenuated these changes as compared to the TNBS colitis group. At the high dose, Compound C was significantly effective in reducing the colon weight to length ratio, diminishing the thickness of the colon walls, and more importantly, improving the colon gross pathology scores to normal

in 70% of the animals. Statistical significances were observed at the low dose in all above indications except colon wall thickness although a trend was evident. The reduction in inflammation observed with orally delivered Compound C was similar to the effect observed from the neutralizing anti-IL-23p19 monoclonal antibody (Figure 6).

[00730] Histological examination of H&E stained distal colons show that the majority of the lesions observed from the vehicle group are transmural, characterized by necrosis with inflammatory cells transversing the entire thickness of the colon, presence of necrotic tissue debris on the lumen surface, and mucosa devoid of crypts. The animals treated with Compound C generally showed localized lesions limited in the mucosa and submucosa regions, with colon tissues showed potential signs of healing at sites of necrosis. Specifically, the animals treated with 160 mg/kg/d Compound C showed a significant reduction in inflammation, mucosal necrosis and colon wall thickness leading to a significant reduction in the overall histological score, comparable to that from the anti-IL-23p19 antibody control (FIG. 7).

[00731] Concentration analysis of samples collected 1 hour post the last PO dose show that the plasma concentrations of Compound C detected from all animals are  $\leq 2X$  below the IC75 of the compounds as determined in the rat splenocyte/IL-17A cell based assay or the rat IL-23R ELISA, suggesting that the efficacies observed from oral treatment are most likely due to its local activity at the colon (see FIG. 8). Collectively, these data highlights the protective effect of an IL-23R antagonist in the development of TNBS colitis.

[00732] These studies demonstrate that peptides of the present invention are potent, selective and orally efficacious IL-23R peptide antagonists that are promising therapeutics for the treatment of IBD and other disorders. As shown herein, the present invention provides peptides that are: potent blockers of IL-23/IL-23R signaling in a human cell line and in human primary cells; selective for IL-23R, and do not inhibit binding to IL-6R or signaling through IL-12R; cross-reactive towards rat and cynomolgus but not mouse homologs, enabling in vivo studies in these species; resistant to proteolytic and reducing environments of the GI, resulting in high drug levels in the intestinal tissues and limited drug concentrations in the circulation, offering potential safety advantages over systemically delivered therapeutics; and effective and

comparable to an anti-IL23p19 monoclonal antibody in attenuating colitis in a TNBS-induced rat colitis model, most like through GI-restricted activities.

## EXAMPLE 9


### *IN VITRO CHARACTERIZATION OF AN ILLUSTRATIVE IL-23R ANTAGONISTS*

[00733] To evaluate the properties of efficacy the IL-23R peptide antagonist of SEQ ID NOS: 980 (Peptide 980), 993 (Peptide 993), and 1185 (Peptide 1185) experiments to determine potency, selectivity, and stability of the peptides were performed as described above. The IC<sub>50</sub> values of the peptides as measured by quantitative ELISA for IL-23/IL-23R competitive binding assays (performed as described above in Example 2) for human (Hu), cynomolgus monkey (Cyno), and rat (Rat) IL-23 and IL-23R binding are shown in Table E22. The potencies of the peptides were also evaluated by IL-23R activity assays as described above. The IC<sub>50</sub> values of the peptides as determined by the reduction of phospho-STAT3 (pSTAT3) levels in human DB cells exposed to IL-23 (Hu DB Cell (pSTAT3); performed as described above in Example 2); by the reduction of IFN $\gamma$  produced by human NK cells exposed to IL-23 (Hu NK cell; performed as described above in Example 5); and by the reduction of IL-17A produced by activated splenocytes exposed to IL-23 (Rat(spleen); performed as described above in Example 6). Selectivity was evaluated by measuring the IC<sub>50</sub> of the peptides for inhibiting human IL-23/IL-12R beta1 interactions (see Example 4) or human IL-6/IL-6R interactions (see Example 6). The stabilities of the peptides were determined by measuring the half-life of the peptides exposed to simulated intestinal fluid (SIF), simulated gastric fluid (SGF), or human intestinal fluid (SIF).

Table E22. Properties of illustrative IL-23R antagonists Peptide


Peptide	ELISA (nM)			Cell assay (nM)			Selectivity (ELISA, nM)		Stability (t <sub>1/2</sub> , hr)		
	Hu	Cyno	Rat	Hu DB Cell (pSTAT3)	Hu NK cell	Rat (Spleen)	IL-23/IL-12R $\beta$ 1	IL-6/IL-6R	SIF	SGF	HIF
Peptide 993	2.0	2.0	2.0	0.6	2.2	18.3	>100,000	>100,000	12	6	24
Peptide 980	1	1.3	3	0.6	2.8	11.5	>100,000	>100,000	11	10	
Peptide 1185	2	3	3	2	3.9	35	>100,000	>100,000	33	12	24

The structure of Peptide 993 is shown below:




Ac-[(D)-Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub> (SEQ ID NO: 993)

The structure of Peptide 1185 is shown below:



Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub> (SEQ ID NO: 1185)

The structure of Peptide 980 is shown below:



Ac-Cyclo-[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub> (SEQ ID NO: 980)

**[00734]** The results summarized in Table E22 indicate that the peptides potently and selectively inhibit IL-23/IL-23R binding as compared to IL-23/IL-12R beta1 or IL-6/IL-6R interactions. This inhibition was confirmed with cell culture assays that measure the IL-23R-dependent such

as STAT3 phosphorylation, IFN $\gamma$  production, and IL-17A production. The peptides were also determined to have a high degree of stability when exposed to simulated intestinal fluid, simulated gastric fluid, or human intestinal fluid.

[00735] Data from the Human DB Cell (pSTAT3) experiment were used for Schild analysis (see FIG. 17). For Schild analysis, concentrations of 0.3nM, 1nM, 3nM, 10nM, 30nM, 100nM of Peptide 993 were tested. The Schild slope was determined to be 1.068, indicating that Peptide 993 behaves as a simple competitive antagonist. Schild analysis was also performed on a peptide with the structure of SEQ ID NO: 1169, which is highly similar to the structures of Peptide 1185. The Schild slope was determined to be 0.91, indicating that peptides with similar structure, including Peptide 1185, are likely behave as a simple competitive antagonist. Schild analysis was also performed on the peptide of SEQ ID NO: 1211, which has a structure similar to Peptide 980. The Schild slope was determined to be 0.76. However, when the slope was fixed to 1, the  $R^2$  value was 0.975. These data suggest that peptides with similar structure, including Peptide 1185, could behave as a simple competitive antagonist.

#### EXAMPLE 10

##### *IN VIVO* PHARMACOKINETICS OF ILLUSTRATIVE IL-23R ANTAGONISTS

[00736] Pharmacokinetic properties of example peptides were measured *in vivo*. Sprague Dawley rats were administered Peptide 993 at 10 mg/kg P.O.

[00737] A single oral dose of Peptide 993 was administered to normal female Sprague-Dawley rats (N=3 rats per time point) either with or without a drinking water dose that was provided *ad libitum* (see FIG. 11). Following the oral dose, the exposure of Peptide 993 was determined in the plasma at 0.25, 0.5, 1, 3, 6, 8, and 24 hours post-dose. The levels of Peptide 993 were also determined in small intestine, colon, small intestine mucosa, colon mucosa, small intestine mucus, Peyer's patch and Mesenteric Lymph Node (MLN) at 1, 3, 6, 8 and 24 hours post-dose. Urine and feces were collected at 6 and 24 hours to determine the excretion of Peptide 993. Plasma, feces, and tissue samples were stored at  $-80 \pm 10$  °C prior to analysis. For plasma, 50  $\mu$ L of the sample was used to extract compound by means of protein precipitation using a quench solution (MeOH:ACN w/ 0.1% formic acid, 50:50 volume) with internal standard. For feces, samples were homogenized with 0.1% Formic acid in water (water:feces ratio of 4:1) prior to

extraction. 50  $\mu$ L of fecal homogenate was used to extract compound by means of protein precipitation using a quench solution (MeOH:ACN w/ 0.1% formic acid, 50:50 volume) with internal standard. For tissues such as colon or small intestine, samples were homogenized with 0.1% Formic acid in water (water:tissue ratio of 3:1) prior to extraction. For tissue such as Peyer's Patch and Mesenteric Lymph Nodes, samples were homogenized with 0.1% Formic acid in water (water:tissue ratio of 20:1) prior to extraction. 50  $\mu$ L of tissue homogenate was used to extract compound by means of protein precipitation using a quench solution (MeOH:ACN w/ 0.1% formic acid, 50:50 volume) with internal standard. The precipitated protein was removed by filter plate and the collected supernatant was dried and reconstituted. Processed samples were analyzed on an AB/MDS Sciex API 4000 mass spectrometer. Positive ions were monitored in the multiple reaction-monitoring (MRM) mode. Quantitation was by peak area ratio.

[00738] No detectable levels of Peptide 993 were observed in rat plasma between 0 and 24 hours following administration (see FIG. 11A). In contrast, detectable levels of Peptide 993 were present in the Peyer's Patch and small intestine for at least six hours (see FIGS. 11B and 11C), and for at least 8 hours in the colon post-dose administration (see FIG 11D). Levels greater than 5% of the total administered dose of Peptide 993 were detected in the rat feces at 24 hours following administration, further indicating that Peptide 993 has a high degree of oral stability. Taken together, these results demonstrate that Peptide 993 is an orally stable GI restricted peptide, that demonstrates a high GI content and limited systemic distribution following oral administration.

[00739] Sprague Dawley rats were administered Peptide 1185 at 10 mg/kg P.O. A single oral dose of Peptide 1185 was administered to normal female Sprague-Dawley rats (N=3 rats per time point). Following the oral dose, the exposure of Peptide 1185 was determined in the plasma in samples taken up to 8 hours post-dose. Urine and feces were collected to hours to determine the excretion of Peptide 1185 (FIG 18).

[00740] Sprague Dawley rats were administered Peptide 980 at 10 mg/kg P.O. A single oral dose of Peptide 980 was administered to normal female Sprague-Dawley rats (N=3 rats per time point). Following the oral dose, the exposure of Peptide 980 was determined in the plasma in

samples taken up to 8 hours post-dose. Urine and feces were collected to hours to determine the excretion of Peptide 980 (FIG 19).

### EXAMPLE 11

#### SAFETY PROFILE OF ILLUSTRATIVE IL-23R ANTAGONISTS

[00741] Safety profiles of illustrative peptide inhibitors were characterized. Peptide 993 and Peptide 1185 were evaluated in a safety panel examining binding to a panel of 44 targets. The targets included G protein coupled receptors (GPCRs), transporters, *e.g.*, dopamine transporter (DAT), and ion channels. For all targets, these peptides displayed no activity, as defined by a change (inhibitory or stimulatory) in the target's activity of greater than 25%. The targets tested in the safety panel are listed in Table E24. For each target, Peptide 993 and Peptide 1185 was determined to be inactive at concentrations of up to 10  $\mu$ M. The safety profile of Peptide 980 was evaluated by testing compounds selected from Table E24. For all compounds tested, Peptide 980 showed only moderate activity in the acetylcholinesterase assay (33%), and otherwise did not display any activity, as defined by a change in the target's activity of greater than 25%.

Table E24. Peptide 993 and Peptide 1185 are Inactive in a Safety Panel

Panel of 44 targets		
A <sub>2A</sub>	M <sub>3</sub>	5-HT <sub>3</sub>
alpha <sub>1A</sub>	delta <sub>2</sub> (DOP)	Ca <sup>2+</sup> channel (L, dihydropyridine site)
alpha <sub>2A</sub>	kappa (KOP)	hERG (membrane preparation)
beta <sub>1</sub>	mu (MOP)	K <sub>V</sub> channel
beta <sub>2</sub>	5-HT <sub>1A</sub>	Na <sup>+</sup> channel (site 2)
CB <sub>1</sub>	5-HT <sub>1B</sub>	AR
CB <sub>2</sub>	5-HT <sub>2A</sub>	GR
CCK <sub>1</sub> (CCK <sub>A</sub> )	5-HT <sub>2B</sub>	Lck kinase
D <sub>1</sub>	V <sub>1a</sub>	COX <sub>1</sub>
D <sub>2S</sub>	dopamine transporter (DAT)	COX <sub>2</sub>
ET <sub>A</sub>	norepinephrine transporter (NET)	acetylcholinesterase

Panel of 44 targets		
H <sub>1</sub>	5-HT transporter (SERT)	MAO-A
H <sub>2</sub>	BZD (central)	PDE3A
M <sub>1</sub>	NMDA	PDE4D <sub>2</sub>
M <sub>2</sub>	N neuronal α4β2	

## EXAMPLE 12

### EFFECTS OF AN ILLUSTRATIVE PEPTIDE INHIBITOR IN A RAT MODEL OF ACUTE COLITIS

[00742] To evaluate the efficacy of the illustrative peptide inhibitor Peptide 1185 in an animal model of disease, acute colitis was induced by providing 7-week-old female Sprague-Dawley rats with 64 mg/kg TNBS in 50% ethanol administered intrarectally at Day 0. Peptide 1185 was administered orally 37mg/kg/day (combined PO and in drinking water), PO BID, day -1 to day 7. A sham group that was not exposed to TNBS and a TNBS treated group that received treatment with vehicle were used as control groups. For comparators, neutralizing anti-IL-23p19 antibody was administered intraperitoneally at 4 mg/kg on Day -1 and again on Day 3, and prednisolone was administered at 10 mg/day P.O. All animals received orally water as the vehicle which was used to formulate Peptide 993.

[00743] As described above, animals were observed daily for clinical signs which included percent body weight loss and signs of loose stools or diarrhea. Six days after inoculation of TNBS, rats were sacrificed and the entire colon length and colon weight from each animal were recorded. The severity of colitis was evaluated by a pathologist. In addition to the colon wall thickness, the gross colon damage was scored on a 0-5 scale according to Table E29, and histopathological scores were determined based on the parameters listed in Table E30.

[00744] Treatment with Peptide 1185 significantly reduced some of the disease parameters that were observed in the TNBS rat model of acute colitis. While rats in the sham group continued to gain weight over the course of the study, rats exposed to TNBS and treated with vehicle experienced weight loss. Oral treatment with prednisolone or systemic treatment with anti-IL-23p19 prevented the weight loss in TNBS exposed rats. Treatment with orally administered

Peptide 1185 did not significantly prevent the weight loss in TNBS challenged rats (see FIG. 13). A significant reduction was also observed in the colon weight to colon length ratios following treatment with prednisolone or anti-IL-23p19 as compared to treatment with vehicle. Oral administration of Peptide 1185 resulted in similar reductions of colon weight to colon length ratios and colon macroscopic scores in TNBS exposed rats. Higher colon macroscopic scores indicated a higher degree of colon pathology. The colon macroscopic score was determined by adding the scores for adhesion, strictures, ulcer, and colon wall thickness, all of which were significantly reduced by treatment with prednisolone, anti-IL-23p19, or Peptide 1185, as compared to vehicle treated controls. These data demonstrate that oral administration of Peptide 1185 has comparable efficacy to systemic administration of anti-IL-23p19 monoclonal antibody.

[00745] The pathological features of tissue sections from colon taken from rats in the sham, vehicle, anti-IL-23p19, and Peptide 1185 groups were examined. Mucosal inflammation, transmural inflammation, gland loss, and erosion were scored according to the criteria listed in Table E29. For all of these features, treatment with anti-IL-23p19 or Prednisolone reduced the histopathology scores associated with TNBS exposure. Treatment with Peptide 1185 did not significantly reduce the histopathology scores.

### EXAMPLE 13

#### EFFECTS OF AN ILLUSTRATIVE PEPTIDE INHIBITOR IN A RAT MODEL OF ACUTE COLITIS

[00746] To evaluate the efficacy of the illustrative IL-23R peptide inhibitor Peptide 993 in an animal model of disease, acute colitis was induced by providing 7-week-old female Sprague-Dawley rats with 64 mg/kg TNBS in 50% ethanol administered intrarectally at Day 0. Peptide 993 was administered orally two times a day at 10 mg/kg, for a total of 42 mg/kg per day, for 8 days starting approximately 24 hours (Day -1) prior TNBS inoculation. A sham group that was not exposed to TNBS and a TNBS treated group that received treatment with vehicle were used as control groups. All animals received orally water as the vehicle which was used to formulate Peptide 993.

[00747] As described above, animals were observed daily for clinical signs which included percent body weight loss and signs of loose stools or diarrhea. Six days after inoculation of

TNBS, rats were sacrificed and the entire colon length and colon weight from each animal were recorded. The severity of colitis was evaluated by a pathologist. In addition to the colon wall thickness, the gross colon damage was scored on a 0-5 scale according to Table E29, and histopathological scores were determined based on the parameters listed in Table E30.

[00748] Treatment with Peptide 993 significantly reduced all disease parameters that were observed in the TNBS rat model of acute colitis. While rats in the sham group continued to gain weight over the course of the study, rats exposed to TNBS and treated with vehicle experienced weight loss. Treatment with orally administered Peptide 993 also prevented the weight loss in TNBS challenged rats (see FIG. 12). In addition, a significant reduction was also observed in the colon weight to colon length ratios following treatment with oral administration of Peptide 993. Higher colon macroscopic scores indicated a higher degree of colon pathology and was significantly reduced by treatment with Peptide 993, as compared to vehicle treated controls. Oral administration of Peptide 993 has comparable efficacy to systemic administration of anti-IL-23p19 monoclonal antibody, which served as a positive control. Histopathological scores were significantly reduced in colons from Peptide 993 treated rats as compared to vehicle group.

#### EXAMPLE 14

##### EFFECTS OF AN ILLUSTRATIVE PEPTIDE INHIBITOR IN A RAT MODEL OF ACUTE COLITIS

[00749] To evaluate the efficacy of the illustrative IL-23R peptide inhibitor Peptide 980 in an animal model of disease, acute colitis was induced by providing 7-week-old female Sprague-Dawley rats with 64 mg/kg TNBS in 50% ethanol administered intrarectally at Day 0. Peptide 980 was administered orally 37mg/kg/day (combined PO and in drinking water), PO BID, day -1 to day 7. A sham group that was not exposed to TNBS and a TNBS treated group that received treatment with vehicle were used as control groups. All animals received orally PBS as the vehicle which was used to formulate Peptide 980.

[00750] As described above, animals were observed daily for clinical signs which included percent body weight loss and signs of loose stools or diarrhea. Six days after inoculation of TNBS, rats were sacrificed and the entire colon length and colon weight from each animal were recorded. The severity of colitis was evaluated by a pathologist. In addition to the colon wall thickness, the gross colon damage was scored on a 0-5 scale according to Table E29, and

histopathological scores were determined based on the parameters listed in Table E30. Treatment with Peptide 980 significantly reduced all disease parameters that were observed in the TNBS rat model of acute colitis.

[00751] Treatment with orally administered Peptide 980 prevented the weight loss in TNBS challenged rats (see FIG. 14). In addition, a significant reduction was also observed in the colon weight to colon length ratios following treatment with oral administration of Peptide 980. Higher colon macroscopic score indicates a higher degree of colon pathology, and was significantly reduced by treatment with Peptide 980, as compared to vehicle treated controls (see FIG. 14).

[00752] Pathological features of tissue sections from colon taken from rats in the sham, vehicle, and Peptide 980 treated groups were examined. Mucosal inflammation, transmural inflammation, gland loss, and erosion (see FIG. 14D) were scored according to the criteria listed in Table E30. The sum of the histopathology score was significantly reduced by Peptide 980 treatment as compared to vehicle.

#### EXAMPLE 15

##### LEVELS OF BIOMARKERS FOLLOWING TREATMENT WITH PEPTIDE INHIBITORS IN A RAT MODEL OF ACUTE COLITIS

[00753] Levels of inflammatory markers were examined in the colons. The distal colon tissue samples, designated for protein expression analysis, were flash frozen after collection. For protein extraction, the samples were thawed, weighed and homogenized in the extraction buffer (PBS pH 7.2 supplemented with Protease Inhibitors, 3x volume:weight). The homogenates were centrifuged at 13krpm at 4°C for 15 minutes, a total of two times to remove the debris. The supernatants were saved in multiple aliquots in -80°C and subsequently used for protein expression analysis on ELISA. The total protein in each sample was quantified using BCA assay. MPO, IL-1 $\beta$ , IL-6, IL-17A and IL-22 protein expression in the distal colon samples were analyzed using commercially available rat ELISA kits.

[00754] Treatment with Peptide 993 reduced levels of inflammatory markers present in the colon. The disease defining (MPO, IL-6 and IL-1 $\beta$ ) and IL-23 directed biomarkers (IL-22, and IL-

17A) were reduced by treatment with Peptide 993 as compared to vehicle treated controls (see FIG. 15). These data demonstrate that administration of Peptide 993 in amounts that can reduce pathology *in vivo* also decrease levels of biomarkers present in the colon that are associated with IL-23R activity. Treatment with Peptide 980 reduced levels MPO and IL-22 as compared to vehicle treated controls (see FIG. 16). Treatment with Peptide 1185 at the dose tested did not significantly reduce levels of MPO, IL-22, or IL-17A.

Table E29. Colon Macroscopic Score

Colonic Score Parameters: the values for each animal was summed to obtain the colonic score (maximum value = 12).
<b>Adhesions:</b> none = 0 minimal = 1 involving several bowel loops = 2
<b>Strictures:</b> none = 0 minimal = 1 mild = 2 severe, proximal dilatation = 3
<b>Ulcers:</b> none focal hyperemia, no ulcers = 1 ulcer without significant inflammation (hyperemia and bowel wall thickening = 2 ulceration of 1- $\leq$ 3 cm = 3 ulceration 3- < 6 cm = 4 ulceration $\geq$ 6 cm = 5
<b>Wall thickness:</b> Wall thickness: less than 1 mm = 0 1- 3 mm = 1 $>$ 3 mm = 2

Table E30a. Histopathology - Mucosal/Submucosal Inflammation Score

Mucosal/Submucosal Inflammation Score
The extent of macrophage, lymphocyte, neutrophil and other inflammatory infiltrates was assigned severity scores according to the following criteria:
0=Normal

1=Minimal, larger focal area with MNIC and neutrophils or minimal diffuse, no separation of glands, may be mostly in areas of submucosal edema or mesentery
2=Mild, diffuse mild, or multifocal affecting 11-25% of mucosa with minor focal or multifocal gland separation, no separation in most areas
3=Moderate, 26-50% of mucosa affected with minimal to mild focal or multifocal separation of glands by inflammatory cell infiltrate, milder in remaining areas of mucosa with some areas having no gland separation by inflammation
4=Marked, 51-75% of mucosa affected with mild to moderate separation of glands by inflammatory cell infiltrate, minimal to mild in remaining areas of mucosa but all glands have some separation by infiltrate
5=Severe, 76-100% of mucosa affected with moderate to marked areas of gland separation by inflammatory cell infiltrate, mild to moderate in remaining areas of mucosa

Table E30b.

<b>Mucosal Thickness Score</b>
Mucosal thickness was scored in the colon in a non-tangential area of the section that best represented the overall mucosal thickness. This parameter is indicative of gland elongation and mucosal hyperplasia. A hyperplasia score was determined as follows.
- =No mucosa present
0=Normal
1= Minimal, 5-10% thicker than control mucosa
2=Mild, 11-25% thicker than control mucosa
3=Moderate, 26-50% thicker than control mucosa
4=Marked, 51-75% thicker than control mucosa

Table E30c.

<b>Transmural Inflammation</b>
Presence of inflammatory cell infiltrates within the tunica muscularis mucosa and increased fibroblasts/fibrocytes with perpendicularly arranged blood vessels (granulation tissue), possibly extending to the serosa.
0=Normal
1=Minimal, 5-10% infiltration
2=Mild, 11-25% infiltration
3=Moderate, 26-50% infiltration

4=Marked, 51-75% infiltration
5=Severe, infiltrates reach the serosa and mesentery

Table E30d.

Gland Loss Score
Crypt epithelial and remaining gland epithelial loss was scored based on the approximate percent
of the mucosa that was affected as follows:
0=None
1=Minimal, 1-10% of the mucosa affected
2=Mild, 11-25% of the mucosa affected
3=Moderate, 26-50% of the mucosa affected
4=Marked, 51-75% of the mucosa affected
5=Severe, 76-100% of the mucosa affected

Table E30e.

Erosion Score
The loss of surface epithelium was scored based on the approximate percent of the mucosa that was affected as follows.
0=None
1=Minimal, 1-10% of the mucosa affected
2=Mild, 11-25% of the mucosa affected
3=Moderate, 26-50% of the mucosa affected
4=Marked, 51-75% of the mucosa affected
5=Severe, 76-100% of the mucosa affected

Table E30f.

Histopathology Sum
A sum of inflammation, gland loss, erosion, and transmural inflammation scores was calculated.

## EXAMPLE 16

### CHARACTERIZATION OF ADDITIONAL PEPTIDE INHIBITION OF BINDING OF INTERLEUKIN-23 TO THE INTERLEUKIN-23 RECEPTOR

[00755] Peptide optimization was performed to identify additional peptide inhibitors of IL-23 signalling that were active at low concentrations (e.g., IC<sub>50</sub> <10 nM) while exhibiting gastrointestinal (GI) stability. Certain peptides were tested to identify peptides that inhibit the binding of IL-23 to human IL-23R and inhibit IL-23/IL-23R functional activity, as described below. Peptides tested included peptides containing a variety of different cyclization chemistries, including, e.g., peptides containing a disulfide linkage, e.g., between two Pen residues, and peptides containing a thioether linkage. Peptide inhibitors of the present invention include but are not limited to peptides having any of the structures depicted herein. In addition, peptide inhibitors of the present invention include those having the same amino acid sequence of the peptides or structures described herein, without being required to have the same or any N- or C-terminal “capping” groups, such as, e.g., Ac or NH<sub>2</sub>.

[00756] Assays performed to determine peptide activity were performed as described in Example 2 above. Human ELISA indicates the IL23-IL23R competitive binding assay, Rat ELISA indicates the rat IL-23R competitive binding ELISA assay, and pStat3HTRF indicates the DB cells IL-23R pSTAT3 cell assay. The peptides depicted in Table E31 are cyclized via a disulfide bridge formed between two residues in these peptides. The peptides depicted in Table E32 are cyclized via a thioether bond between the indicated amino acid residues. Table E32 provides an illustrative structure depicting thioether cyclization, which may also be indicated in the table by the term “Cyclo,” with the cyclic region bracketed immediately following.

Table E31. Illustrative Peptides Containing the Ac-[Pen]-XXWX-[Pen]-XXXX Motif and Analogues

SEQ ID NO:	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
1115	[Palm]-[isoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			**
1116	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(PEG4-isoGlu-Palm)]-NN-NH <sub>2</sub>			*
1117	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1118	[Octanyl]-[IsoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1119	[Octanyl]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*


1120	[Palm]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
1121	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Octanyl)]-NN-NH <sub>2</sub>		**	*
1122	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Palm)]-NN-NH <sub>2</sub>			*
1123	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]NN-NH <sub>2</sub>			***
1124	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			**
1125	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Palm)-[Lys(Ac)]-NN-NH <sub>2</sub>			*
1126	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Lauryl)]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
1127	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			***
1128	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			***
1129	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(PEG4-IsoGlu-Palm)]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1130	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-a-Me-K(PEG4-IsoGlu-Lauryl)]-[Lys(Ac)]-NN-NH <sub>2</sub>			*
1131	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(IVA)]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1132	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(Biotin)]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	**	*
1133	Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH <sub>2</sub> )-[2-Nal]-[ $\alpha$ -MeLys(Octanyl)]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1134	Ac-[Pen]-[Lys(IVA)]-TWO-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1135	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(IVA)]-N-NH <sub>2</sub>			*
1136	Ac-[Pen]-[Lys(Biotin)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1137	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Biotin)]-N-NH <sub>2</sub>			**
1138	Ac-[Pen]-[Lys(Octanyl)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			**
1139	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(octanyl)]-N-NH <sub>2</sub>			**
1140	Ac-[Pen]-[Lys(Palm)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			>1000

1141	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Lys(Palm)]-N-NH <sub>2</sub>			>1000
1142	Ac-[Pen]-[Lys(PEG8)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			**
1143	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(PEG8)]-N-NH <sub>2</sub>			**
1144	Ac-[Pen]-K(Peg11-Palm)TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			**
1145	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Peg11-palm)]-N-NH <sub>2</sub>			**
1146	Ac-[Pen]-[Cit]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		*	*
1147	Ac-[Pen]-[Lys(Ac)]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		**	*
1148	Ac-[Pen]-NT-[Phe(3,4-OCH <sub>3</sub> ) <sub>2</sub> ]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			~1000
1149	Ac-[Pen]-NT-[Phe(2,4-CH <sub>3</sub> ) <sub>2</sub> ]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			***
1150	Ac-[Pen]-NT-[Phe(3-CH <sub>3</sub> )]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			***
1151	Ac-[Pen]-NT-[Phe(4-CH <sub>3</sub> )]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>			***
1152	Ac[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*	*	*
1153	Ac[(D)Tyr]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	*	*	*
1154	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QN-NH <sub>2</sub>			*
1155	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-N-NH <sub>2</sub>			*
1156	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Lys(Ac)]-NH <sub>2</sub>			*
1157	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QQ-NH <sub>2</sub>		**	*
1158	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Q-[βAla]-NH <sub>2</sub>		**	*
1159	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Cit]-NH <sub>2</sub>			*
1160	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-NNH <sub>2</sub>		**	*
1161	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-Q-NH <sub>2</sub>			*
1162	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-[Lys(Ac)]-NH <sub>2</sub>			**

1163	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-[Cit]-NH <sub>2</sub>			**
1164	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QN- [ $\beta$ Ala]-NH <sub>2</sub>	*	*	*
1165	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-E-[Cit]- Q-NH <sub>2</sub>			*
1166	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- CitNCitNH <sub>2</sub>		*	*
1167	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Cit]- Q-[Cit]-NH <sub>2</sub>			*
1168	Ac-[Pen]-[Cit]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>		*	*
1169	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			
1170	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QNN- NH <sub>2</sub>		*	*
1171	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-ENQ- NH <sub>2</sub>		**	*
1172	Ac-[Pen]-GPWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			>1000
1173	Ac-[Pen]-PGWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			~1000
1174	Ac-[Pen]-NTWN-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			**
1175	Ac-[Pen]-NSWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			*
1176	Ac-[Pen]-N-[Aib]-WQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			>1000
1177	Ac-[Pen]-NTW-[Aib]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]N-[Aib]-NH <sub>2</sub>			**
1178	Ac-[Pen]-QTW-[Lys(Ac)]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]- [Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>		*	*
1179	Ac-[Pen]-[Lys(Ac)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]- [Aib]-[Lys(Ac)]NNNH <sub>2</sub>			*
1180	Ac-[Pen]-QVWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			*
1181	Ac-[Pen]-NT-[2-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			**
1182	Ac-[Pen]-NT-[1-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]- [Lys(Ac)]-NN-NH <sub>2</sub>			**
1183	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]- [Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*
1184	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]- [Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*

1185	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*
1186	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*	*	*
1187	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLys]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*	*	*
1188	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*	*	*
1189	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-LN-NH <sub>2</sub>			*
1190	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-GN-NH <sub>2</sub>			*
1191	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-SN-NH <sub>2</sub>		**	*
1192	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Aib]-N-NH <sub>2</sub>			*
1193	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-FN-NH <sub>2</sub>			*
1194	Ac-[Pen]-NTW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*
1195	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Tic]-[ $\beta$ Ala]-NH <sub>2</sub>			***
1196	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[nLeu]-[ $\beta$ Ala]-NH <sub>2</sub>			*
1197	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-G-[ $\beta$ Ala]-NH <sub>2</sub>			*
1198	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-R-[ $\beta$ Ala]-NH <sub>2</sub>			*
1199	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-W-[ $\beta$ Ala]-NH <sub>2</sub>			**
1200	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-S-[ $\beta$ Ala]-NH <sub>2</sub>			*
1201	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-L-[ $\beta$ Ala]-NH <sub>2</sub>			*
1202	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[AIB]-[ $\beta$ Ala]-NH <sub>2</sub>			*
1203	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[N-MeAla]-[ $\beta$ Ala]-NH <sub>2</sub>			***
1204	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[2-Nap]-[ $\beta$ Ala]-NH <sub>2</sub>		**	*
1205	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-F-[ $\beta$ Ala]-NH <sub>2</sub>		**	*
1206	Ac-[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]NN-NH <sub>2</sub>			*

Table E32. IC50 of Illustrative Peptide Inhibitors (Thioethers)



Ac-Cyclo-[[Abu]-XXWXC]-[Phe(4-OMe)]-[2-Nal]-XXX-NH<sub>2</sub>

SEQ ID NO:	Sequence	Human ELISA (nM)	Rat ELISA (nM)	pStat3 HTRF (nM)
1207	Biotin-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	TBC	*	*
1208	Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	TBC	*	*
1209	Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	TBC	*	*
1210	Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*	*	*
1211	Ac-E-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			
1212	Ac-[(D)Asp]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			
1213	Ac-R-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*	*	*
1214	Ac-[(D)Arg]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	TBC	*	*
1215	Ac-F-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			*
1216	Ac-[(D)Phe]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			
1217	Ac-[2-Nal]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			*
1218	Ac-T-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			*
1219	Ac-L-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			*
1220	Ac-[(D)Gln]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	TBC	*	*
1221	Ac-[(D)Asn]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>		*	*

1222	Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(PEG4-Alexa488)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>			*
1223	[Alexa488]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*	*	*
1224	[Alexa647]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	*		TBC
1225	[Alexa-647]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*		
1226	[Alexa647]-[PEG12]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*		
1227	[Alexa488]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*		*

\* < 10 nM; \*\* ≥ 10 and < 100 nM; \*\*\* ≥ 100 and ≤ 1,000 nM

## EXAMPLE 17

### STABILITY OF ADDITIONAL PEPTIDE INHIBITORS IN SIMULATED INTESTINAL FLUID (SIF), SIMULATED GASTRIC FLUID (SGF) AND REDOX CONDITIONS

[00757] Studies were carried out in simulated intestinal fluid (SIF) and simulated gastric fluid (SGF) to evaluate gastric stability of additional peptide inhibitors of the present invention. In addition, studies were carried out to assess redox stability of the additional peptide inhibitors of the present invention.

Table E33. Thioethers and Dipens

SEQ ID NO:	Sequence	SIF t1/2 (min)	SGF t1/2 (min)
1228	Biotin-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	>90	>180
1229	Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180	>180
1230	Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180	<180
1231	Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180, >180	>180 >180

1232	[Octanyl]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180	>180
1233	[Palm]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180	>10
1234	Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(PEG4-Alexa488)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	> 180	>90
1235	[Alexa488]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	> 180	> 180
1236	[Alexa647]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	>180	>180
1237	Ac[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	>180	>180
1238	Ac-[(D)Tyr]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	>180	>180
1239	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QN-[βAla]-NH <sub>2</sub>	Stable	>180
1240	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[α-MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180 >180	>180, >180
1241	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180, >180	>180, >180
1242	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[α-MeLeu]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	Stable	>180
1243	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-N-[βAla]-NH <sub>2</sub>	>180	>180
1244	Ac-[Pen]-NTW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	>180	>180
1245	Ac-[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-AMINO-4-CARBOXY-TETRAHYDROPYRAN]-[Lys(Ac)]NNNH2	> 180	> 180
1246	[Alexa488]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	> 180	> 180

>90 = less than or equal to 180 min and greater than 90 min; >45 min = less than or equal to 90 min and greater than 45 min; >10 = less than or equal to 45 min and greater than 10 min; <10 = less than 10 min.

## EXAMPLE 18

### NK CELL ASSAY

[00758] Natural killer (NK) cells, purified from human peripheral blood of healthy donors by negative selection (Miltenyi Biotech, Cat # 130-092-657), were cultured in complete media (RPMI 1640 containing 10% FBS, L-glutamine and penicillin-streptomycin) in the presence of

IL-2 (RnD, Cat # 202-IL-010/CF) at 25 ng/mL. After 7 days, cells were centrifuged, and resuspended in complete media at 1E6 cells/mL. Recombinant IL-23 at predetermined EC50 to EC75 and IL-18 (RnD, Cat # B003-5) at 10 ng/mL were mixed with varying concentrations of peptides, and added to NK cells seeded at 1E5 cells per well. After 20 to 24 hours, IFN $\gamma$  in the supernatant was quantified using Quantikine ELISA (RnD, Cat # DIF50). Results are shown in Table E34. Multiple results shown for a single peptide are from separate assays.

Table E34. Primary Cell Assay (Thioethers and Dipens)

SEQ ID NO:	Sequence	NK cell Assay (nM)
1247	Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*, *, *
1248	Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*, **, **
1249	[Alexa647]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH <sub>2</sub>	**, *
1250	Ac[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	**
1251	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QN-[ $\beta$ Ala]-NH <sub>2</sub>	**
1252	Ac-[Pen]-QTW-[Lys(Ac)]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	**
1253	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[ $\alpha$ -MeLeu]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
1254	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH <sub>2</sub>	*
1255	Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH <sub>2</sub>	*
1256	Ac-[Pen]-NTW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH <sub>2</sub>	*

[00759] All of the above U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, are incorporated herein by reference, in their entirety.

[00761] From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

**CLAIMS**

What is claimed is:

1. A peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Xa):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Xa)  
wherein

X1 is any amino acid or absent;

X2 is any amino acid or absent;

X3 is any amino acid or absent;

X4 is any amino acid or chemical moiety capable of forming a bond with X9;

X5 is any amino acid;

X6 is any amino acid;

X7 is any amino acid;

X8 is any amino acid;

X9 is any amino acid or chemical moiety capable of forming a bond with X4;

X10 is any amino acid;

X11 is any amino acid;

X12 is any amino acid;

X13 is any amino acid;

X14 is any amino acid;

X15 is any amino acid,

X16 is any amino acid or absent;

X17 is any amino acid or absent;

X18 is any amino acid or absent;

X19 is any amino acid or absent; and

X20 is any amino acid or absent,

wherein the peptide inhibitor is cyclized via a bond between X4 and X9, and

wherein the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

2. The peptide inhibitor of claim 1, wherein the bond between X4 and X9, is a disulfide bond, a thioether bond, a lactam bond, a triazole ring, a selenoether bond, a diselenide bond, or an olefin bond.
3. The peptide inhibitor of claim 1, wherein X4 is Cys and X9 is Cys, and the bond is a disulfide bond.
4. The peptide inhibitor of claim 1, wherein X4 is Pen and X9 is Pen, and the bond is a disulfide bond.
5. The peptide inhibitor of claim 1, wherein the peptide inhibitor comprises an amino acid sequence of any one of SEQ ID NOS: 365-370, 857-1029.
6. The peptide inhibitor of claim 1, wherein the peptide inhibitor comprises an amino acid sequence set forth in any of Formulas (Va), (Vb), (Vc), (Vd), (Ve), (Vf), (Vg) and (Vh).
7. The peptide inhibitor of claim 1, wherein the peptide inhibitor comprises any of the following amino acid sequences:

[Palm]-[isoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NNNH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-isoGlu-Palm)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(Ac)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Octanyl]-[IsoGlu]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Octanyl]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Palm]-[PEG4]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Octanyl)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(PEG4-Palm)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Palm)-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-Lauryl)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGlu-Palm)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)-(PEG4-IsoGLu-Lauryl)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-IsoGlu-Palm)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(PEG4-IsoGlu-Lauryl)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(IV A)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(Biotin)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-QTWQ-[Pen]-Phe(4-CONH<sub>2</sub>)-[2-Nal]-[ $\alpha$ -MeLys(Octanyl)]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-[Lys(IV A)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(IV A)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Biotin)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Biotin)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Octanyl)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(octanyl)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Palm)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Lys(Palm)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Lys(PEG8)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(PEG8)]-N-NH<sub>2</sub>;

Ac-[Pen]-K(Peg11-Palm)TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Peg11-palm)]-N-NH<sub>2</sub>;

Ac-[Pen]-[Cit]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-[Lys(Ac)]-TW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(3,4-OCH<sub>3</sub>)<sub>2</sub>]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(2,4-CH<sub>3</sub>)<sub>2</sub>]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(3-CH<sub>3</sub>)]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NT-[Phe(4-CH<sub>3</sub>)]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac[(D)Tyr]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-N-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Lys(Ac)]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-QQ-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-Q-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-N-[Cit]-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-NNH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-Q-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Cit]-[Lys(Ac)]-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Lys(Ac)]-[Cit]-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QN-[βAla]-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-E-[Cit]-Q-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Cit]-N-[Cit]-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Cit]-Q-[Cit]-NH<sub>2</sub>;  
Ac-[Pen]-[Cit]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-QNN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-ENQ-NH<sub>2</sub>;  
Ac-[Pen]-GPWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-PGWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWN-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NSWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-N-[Aib]-WQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTW-[Aib]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]N-[Aib]-NH<sub>2</sub>;  
Ac-[Pen]-QTW-[Lys(Ac)]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-[Lys(Ac)]-TWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]NNNH<sub>2</sub>;  
Ac-[Pen]-QVWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NT-[2-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NT-[1-Nal]-Q-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[α-MeLeu]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[α-MeLys]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[α-MeLeu]-[Lys(Ac)]-N-[βAla]-NH<sub>2</sub>;  
Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[α-MeLys]-[Lys(Ac)]-N-[βAla]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-N-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-LN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-GN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-SN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Aib]-N-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-FN-NH<sub>2</sub>;

Ac-[Pen]-NTW-[Cit]-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[Tic]-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[nLeu]-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-G-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-R-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-W-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-S-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-L-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[AIB]-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[N-MeAla]-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-[2-Nap]-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[Aib]-[Lys(Ac)]-F-[ $\beta$ Ala]-NH<sub>2</sub>;

Ac-[(D)Arg]-[Pen]-NTWQ-[Pen]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]NNNH<sub>2</sub>;

Biotin-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

Ac-E-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Asp]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-R-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-F-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Phe]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[2-Nal]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-T-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-L-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Gln]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-[(D)Asn]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

Ac-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)-(PEG4-Alexa488)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

[Alexa488]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

[Alexa647]-[PEG4]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-ENN-NH<sub>2</sub>;

[Alexa-647]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>;

[Alexa647]-[PEG12]-[(D)Arg]-cyclo[[Abu]-QTCWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>; and

[Alexa488]-[PEG4]-[(D)Arg]-cyclo[[Abu]-QTCWQC]-[Phe[4-(2-aminoethoxy)]-[2-Nal]-[4-amino-4-carboxy-tetrahydropyran]-[Lys(Ac)]-NN-NH<sub>2</sub>,

wherein the peptide inhibitor is cyclized via a disulfide bond between two Pen residues or by a thioether bond between Abu and a Cys residue, and wherein the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

8. The peptide inhibitor of any one of claims 1-7, further comprising one or more half-life extension moiety and/or one or more linker moiety conjugated to the peptide inhibitor.

9. The peptide inhibitor of claim 8, wherein the half-life extension moiety is conjugated to the peptide inhibitor via one or more linker moieties.

10. The peptide inhibitor of any one of claims 1-9, wherein the peptide inhibitor comprises the structure of Formula I:



or a pharmaceutically acceptable salt or solvate thereof, wherein

$R^1$  is a bond, hydrogen, a C1-C6 alkyl, a C6-C12 aryl, a C6-C12 aryl, a C1-C6 alkyl, a C1-C20 alkanoyl, and including PEGylated versions alone or as spacers of any of the foregoing; X is the amino acid sequence; and

$R^2$  is OH or NH<sub>2</sub>.

11. A peptide dimer inhibitor of an interleukin-23 receptor, wherein the peptide dimer inhibitor comprises two peptide monomer subunits connected via one or more linker moieties, wherein each peptide monomer subunit has a sequence or structure set forth in any one of claims 1-10.

12. The peptide dimer inhibitor of claim 11, wherein one or both peptide monomer subunit is cyclized via an intramolecular bond between X4 and X9.

13. The peptide dimer inhibitor of claim 12, wherein one or both intramolecular bond is a disulfide bond, a thioether bond, a lactam bond, a selenoether, diselenide, or an olefin bond.
14. The peptide dimer inhibitor of any one of claims 11-13, wherein the linker moiety is a diethylene glycol linker, an iminodiacetic acid (IDA) linker, a  $\beta$ -Ala-iminodiacetic acid ( $\beta$ -Ala-IDA) linker, or a PEG linker.
15. The peptide dimer inhibitor of any one of claims 11-14, wherein the N-terminus of each peptide monomer subunit is connected by the linker moiety or wherein the C-terminus of each peptide monomer subunit is connected by the linker moiety.
16. The peptide inhibitor of any one of claims 1-10 or the peptide dimer of any of claims 11-15 further comprising a conjugated chemical substituent.
17. The peptide inhibitor or the peptide dimer of claim 16, wherein the conjugated chemical substituent is a lipophilic substituent or a polymeric moiety.
18. The peptide inhibitor or the peptide dimer of claim 16, wherein the conjugated chemical substituent is Ac, Palm, gamaGlu-Palm, isoGlu-Palm, PEG2-Ac, PEG4-isoGlu-Palm, (PEG)<sub>5</sub>-Palm, succinic acid, glutaric acid, pyroglutaric acid, benzoic acid, IVA, octanoic acid, 1,4-diaminobutane, isobutyl, or biotin.
19. The peptide inhibitor or the peptide dimer of claim 16, wherein the conjugated chemical substituent is a polyethylene glycol with a molecular mass of 400 Da to 40,000 Da.
20. A polynucleotide comprising a sequence encoding the peptide inhibitor of any one of claims 1-10 or one or both peptide monomer subunit of the peptide dimer inhibitor of any one of claims 11-15.
21. A vector comprising the polynucleotide of claim 20.

22. A pharmaceutical composition comprising the peptide inhibitor or the peptide dimer inhibitor of any one of claims 1-19, and a pharmaceutically acceptable carrier, excipient, or diluent.
23. The pharmaceutical composition of claim 22, further comprising an enteric coating.
24. The pharmaceutical composition of claim 23, wherein the enteric coating protects and releases the pharmaceutical composition within a subject's lower gastrointestinal system.
25. A method for treating an Inflammatory Bowel Disease (IBD), ulcerative colitis, Crohn's disease, Celiac disease (*nontropical Sprue*), enteropathy associated with seronegative arthropathies, microscopic colitis, collagenous colitis, eosinophilic gastroenteritis, colitis associated with radio- or chemo-therapy, colitis associated with disorders of innate immunity as in leukocyte adhesion deficiency-1, chronic granulomatous disease, glycogen storage disease type 1b, Hermansky-Pudlak syndrome, Chediak-Higashi syndrome, and Wiskott-Aldrich Syndrome, pouchitis resulting after proctocolectomy and ileoanal anastomosis, gastrointestinal cancer, pancreatitis, insulin-dependent diabetes mellitus, mastitis, cholecystitis, cholangitis, pericholangitis, chronic bronchitis, chronic sinusitis, asthma, psoriasis, psoriatic arthritis, or graft versus host disease in a subject, comprising providing to the subject an effective amount of the peptide inhibitor or peptide dimer inhibitor of any one of claims 1-19, or the pharmaceutical composition of any one of claims 22-24.
26. The method of claim 25, wherein the pharmaceutical composition is provided to the subject by an oral, parenteral, intravenous, peritoneal, intradermal, subcutaneous, intramuscular, intrathecal, inhalation, vaporization, nebulization, sublingual, buccal, parenteral, rectal, intraocular, inhalation, topically, vaginal, or topical route of administration.
27. The method of claim 25 for treating Inflammatory Bowel Disease (IBD), ulcerative colitis, Crohn's disease, wherein the pharmaceutical composition is provided to the subject orally.

28. The method of claim 25 for treating psoriasis, wherein the pharmaceutical composition is provided to the subject orally, topically, parenterally, intravenously, subcutaneously, peritonealy, or intravenously.

29. The method of any one claims 25-28, wherein the peptide inhibitor or the peptide dimer inhibitor inhibits binding of an interleukin-23 (IL-23) to the interleukin-23 receptor (IL-23R).

1/27




FIG. 1

2/27

### IL-12 Stimulation of IFN $\gamma$ Release From Human PBMCs




FIG. 2

3/27




FIG. 3A

0	Normal
1	Edema
2	Edema, slight edema, small erosions
3	Two or more bleeding ulcers, inflammation, moderate adhesions
4	Severe ulceration, stenosis with dilations, severe adhesions

FIG. 3B

4/27

II Potential Region Peptide Binds

IL23R_human	1	10	20	30
IL23R_mouse				
IL23R_rat				
IL23R_chimp				
IL23R_dog				
IL23R_cow				
IL23R_human	130	140	150	160
IL23R_mouse				
IL23R_rat				
IL23R_chimp				
IL23R_dog				
IL23R_cow				
IL23R_human	260	270	280	290
IL23R_mouse				
IL23R_rat				
IL23R_chimp				
IL23R_dog				
IL23R_cow				
IL23R_human	390	400	410	420
IL23R_mouse				
IL23R_rat				
IL23R_chimp				
IL23R_dog				
IL23R_cow				
IL23R_human	520	530	540	
IL23R_mouse				
IL23R_rat				
IL23R_chimp				
IL23R_dog				
IL23R_cow				
IL23R_human	650	660	670	
IL23R_mouse				
IL23R_rat				
IL23R_chimp				
IL23R_dog				
IL23R_cow				

FIG. 4

5/27

40	50	60	70	80
HGGITNINCSGHIWVEPATIFKMGMNISIYCQAAIKNCQPRKLHFYK				
HGGITSINCSGDMWVEPGEIFQMGGMNVSIYCQEALKNCRPRNLYFYK				
HGGIATINCSGNMWVEPGEIFQMGGMNVSVYCQEALKNCRPRNLHFYK				
HGGITNINCSGHIWVEPATIFKMGMNISIYCQAAIKNCQPRKLHFYK				
HGGITNINCSGHIWVEPATIFKMGMNISIYCQAAIKNCQPRKLYFYK				
HGGITNINCSGHIWVEPATIFKMGMNISIYCQAAIKNCQPSKLYFYK				
170	180	190	200	210
KLTYYIDTKYVVHVKSLETEEEEQQYLTSSYINISTDSLQGGKKYLVWV				
KPTYYIDTKYIVHVKSLETEEEEQQYLAQSYVKISTDSLQGGRKYLVWV				
KPTYYIDTKYTVHVKSLETEEQQQYLASNYVNISTDSLQGGRKYLVWV				
KLTYMDTKYVVHVKSLETEEEEQQYLTSSYINISTDSLQGGKKYLVWV				
KPTYYIDTKYVVHVKSLEKEEEEQQYLSSSYINISTDSLQGGKKYLVWV				
RPTYIDTKYVVHVKSLETEEEEQEYLTSSYINISTDSLQKGKKYLVWV				
300	310	320	330	340
QQSEFYLEPNIKYVFQVRCQETGKR-----YW				
QQSEFYLEPDSKYVFQVRCQETGKRKNWQPWSSPFVHQTSEQETGKRNW				
QQSEFYLEPNSKYVFQVRCQGTGKR-----NW				
QQSEFYLEPNIKYVFQVRCQETGKR-----YW				
QQSEFYLEPNTKYIFQVRCQETGKR-----YW				
QQSEFYLQPNAMYVFQVRCQETGKK-----YW				
430	440	450	460	
KRRILLLIPKWLYYEDIPNMKNNSVVKMLQENSELMNNNSSEQVLYVD				
KRKVLLMIPKWLYYEDIPNMENSNVAKLLQEKSVFENDNASEQALYVD				
KRKVLLMIPKWLYYEDIPNMENSNIAKLLQEKSVFENENASEQALYVD				
KRRILLLIPKWLHEDIPNMENSKVVEILQKQSEFMNNNSSEQVLYAD				
KRRILLLIPKWLYYEDIPNMENSKVVKILQERNEFMNNNSSEQVLYVD				
550	560	570	580	590
PVDSLDSGNPRLQKHPNFAFSVSSVNSLSNTIFLGELSLILNQGEC				
TDD-----HFARLKTYPNFQFSASSMALLNKTLLDELCVLNQGEF				
TGD-----HFARLKTYPNFTFSASSMTSLSKTLILDELSLVLNQGEF				
PVDSLDSGNPRLQKHPNFAFSVSSVNSLSNTIFLGELSLILNQGEC				
LADSSDLGGNARFKKYPNFAFSVSSMNSLSNTLFLEEELSLILNQGER				
PADSLNLGNARFKKYPDFAFSVSSTNSLSNTIFLEEELNLILNQGEC				
680	690	700	710	720

FIG. 4 (Continued)

6/27

90	100	110	120
NGIKE RF QITRINKTTARLWYKNFLEPHASMYCTAECPKHFQE TL			
NGFKE EFDITRINRTTARIWYKGFSEPHAYMHCTAECPGHFQE TL			
NGFKE RF HITRINRTTARVWYKGFSEPHASMYCTAECPGRFQE TL			
NGIKE RF QITRINKTTARLWYKNFLEPHASMYCTAECPKHFQE TL			
NGIKE RF QITRINKTTARLWYSNFLEPHASMYCTAECPGYFQE TL			
NGIKE RF HITRINKTTARLWYNFVEPQAFVYCTAECSRYFPE TL			
220	230	240	250
QAAANALGMEESKQLQIHLDDIVIPSAAVISR AETINATVPKTI IY			
QAVNSLGMENSQQLHVHLDDIVIPPSAIIISR AETTNDTVPKTI VY			
QAVNALGMENSQQLQVHLDDIVIPSPSIIISR AETTANVPKTI IY			
QAAANALGMEESKQLQIYLDDIVIPPSAIIISR AETINATVPKTI IY			
QAAANALGMEKSKQLQINLEDDIVIPPSAIIISR AEDINTTIPKTI IH			
QASNV LGMEKSKQLQIHLDDIVIPPSAIIISR AEDINTTVSKTVI IH			
340	350	360	370
380			
QPWSSLFFHKTPETVPQVTSKAFQHDTWNSGLTVASISTGHLTSD			
QPWSSPFVHQTSQTVSQVTAKS-SHEPQKMEMLSATIFRGHPASG			
QPWSSPFVHQTPQTAQSQVTSKP-PHEPQKIEMLTATIFKGHSTSD			
QPWSSPFHKTPETVPQVTSKAFQHDTWNSGLTVASISTGHLTSD			
QPWSSPFHKTAETVPQVTVKSFQHDTQNSELLIASIFKGHLTSD			
QPWSSPFHKTPETVPQVTMKSFQHDTQNSGLLIASIFKKHLTSD			
470	480	490	500
510			
PMITEIKEIFIPEHKPTDYKKE-NTGPLETRDYPQNSLF DNTT VV			
PVLTEISEIISPLEHKPTDYKEERLTGLET RD CPLGMLSTS SSS VV			
PILTEISEIISPLEHKPTDYKKDRLLTGFLET RD CPLTTLSTS SSS VV			
PMITEIKEIFIPEHKPTDYKKE-NTGPLETRDYLQNSLF DNTT VV			
PVITEIEIFLPPEEHKPTDYKAEKNTGSLET RD CQLQNSLLTNTT VV			
PVITEIEIILPPEE-KPMGYKKENNNTGCLERKESLEKSLLTDTT VV			
600	610	620	630
640			
SSPDIQNSVEEEETMLLENDSPSETIPEQTLLPDEFVSCLGIVNE			
NSLDIKNSRQEETTSIVLQSDSPSETI PAQTLLSDEFVSCLAIGNE			
NSLDI QNSRQEETSMILQNDSPSETI P VQTLLPDEFVSCLAIGSE			
SSPDIQNSVEEEETMLLENDSPSETIPEQTLLPDEFVSCLGIVNE			
SPLDMQNSVEGETMLLENAASPNETIPEQTLLPDEFVSCLGIMNE			
SPPDMQNSIEGETAMLLEDALLNETIPEQTLLPDEFVSCLGSMNK			
730	740	750	760

FIG. 4 (Continued)

7/27

Group ID	Female SD Rat ID	TWS at Day 0	ID	Formulation	Route	Dose (mg/kg)	Con. (mg/ml)	Dose Vol. (ml/kg)	Test Article		API in Drinking Water (mg/ml)	Termination at Day 6
									frequency	1 hr post morning dose		
Group-1	1-8	Sham	None	PBS	PO	NA	NA	5	TD, Day 1 to Day 6	None	None	
Group-2	11-20	60 mg/kg	None	PBS	PO	NA	NA	5	TD, Day 1 to Day 6	None	None	
Group-3	21-30	60 mg/kg	Mouse anti-H-23p19	PBS	IP	4	2	2	Day-1 and Day 3, AND PBS PO TD at 5ml/kg	None	None	
Group-4	31-40	60 mg/kg	Compound C	PBS	PO	20	4	5	TD, Day 1 to Day 6	0.6	1 hr post morning dose	
Group-5	41-50	60 mg/kg	Compound C	PBS	PO	6.7	1.3	5	TD, Day 1 to Day 6	0.2	1 hr post morning dose	

FIG. 5

8/27




FIG. 6A




FIG. 6B

9/27




FIG. 6C







FIG. 6D

FIG. 7



11/27



12/27





FIG. 9A

FIG. 9C

FIG. 9B


13/27



X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20

R = H, CH<sub>3</sub>

n1 = 0-6  
n2 = 0-6



X9 = Any amino acid capable of forming bond with X9 with chemistries shown for Y

FIG. 10

14/27




FIG. 11A

FIG. 11B

FIG. 11C

15/27




FIG. 11E




FIG. 11D




FIG. 12B




FIG. 12D




FIG. 12A




FIG. 12C

17/27




FIG. 13B




FIG. 13A




FIG. 13C




FIG. 14B




FIG. 14D




FIG. 14C

FIG. 14A

FIG. 14B

19/27




FIG. 15B




FIG. 15A




FIG. 15C

20/27

FIG. 15E





FIG. 15D



21/27

FIG. 16B





FIG. 16A



22/27




FIG. 17B




FIG. 17A

23/27





FIG. 17D

FIG. 17C

24/27




FIG. 18A

25/27




FIG. 18B

26/27




FIG. 19A

27/27




FIG. 19B

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



## (10) International Publication Number

WO 2017/011820 A3

(43) International Publication Date  
19 January 2017 (19.01.2017)

(51) International Patent Classification:  
*A61K 38/20* (2006.01) *C07K 14/54* (2006.01)  
*C07K 7/08* (2006.01)

Road, Salisbury, Queensland 4107 (AU). **PATEL, Dinesh V.**; 45109 Cougar Circle, Fremont, California 94539 (US). **LIU, David**; 201 Ferne Avenue, Palo Alto, California 94306 (US). **BHANDARI, Ashok**; 2629 Lotus Street, Pleasanton, California 94588 (US).

(21) International Application Number:  
PCT/US2016/042680

(74) Agents: **LAHERTY, Carol** et al.; Cooley LLP, 1299 Pennsylvania Avenue, NW, Suite 700, Washington, District of Columbia 20004 (US).

(22) International Filing Date:  
15 July 2016 (15.07.2016)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant: **PROTAGONIST THERAPEUTICS, INC.**


[US/US]; 521 Cottonwood Drive, Suite A, Milpitas, California 95035 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,

(72) Inventors: **BOURNE, Gregory**; 29 Moolingal Street, Jindalee, Queensland 4074 (AU). **CHENG, XiaoLi**; 263 Lassen Avenue, Mountain View, California 94043 (US). **FREDERICK, Brian Troy**; 8470 Glen Arbor Road, Ben Lomand, California 98005 (US). **ZHANG, Jie**; 101 Ness

[Continued on next page]

(54) Title: PEPTIDE INHIBITORS OF INTERLEUKIN-23 RECEPTOR AND THEIR USE TO TREAT INFLAMMATORY DISEASES



(57) Abstract: The present invention provides novel peptide inhibitors of the interleukin-23 receptor, and related compositions and methods of using these peptide inhibitors to treat or prevent a variety of diseases and disorders, including inflammatory bowel diseases.

FIG. 13A



DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

**Published:**

- with international search report (Art. 21(3))

**(88) Date of publication of the international search report:**

23 February 2017

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US16/42680

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(8) - A61K 38/20; C07K 7/08, 14/54 (2016.01)

CPC - A61K 38/20; C07K 7/08, 14/54

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): A61K 38/20; C07K 7/00, 7/08, 14/54 (2016.01)

CPC: A61K 38/20; C07K 7/00, 7/08, 14/54

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PatSeer (US, EP, WO, JP, DE, GB, CN, FR, KR, ES, AU, IN, CA, INPADOC Data); EBSCO Discovery; PubMed; Google; Google Scholar; Google Patents; The Lens; ENA; NCBI Blast; interleukin-23, receptor, cyclized, amino acid

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2013/0172272 A1 (GALLAGHER, G et al.) July 4, 2013; abstract; paragraphs [0027], [0114], [0135]	1-9
A	US 2003/0166514 A1 (JONES, T et al.) September 4, 2003; paragraph [0035]	1-9
A	US 2010/0190710 A1 (CHEMTOB, S et al.) July 29, 2010; abstract; paragraphs [0155], [0209], [0233]	1-9

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance  
 "E" earlier application or patent but published on or after the international filing date  
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
 "O" document referring to an oral disclosure, use, exhibition or other means  
 "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art  
 "&" document member of the same patent family

Date of the actual completion of the international search

08 November 2016 (08.11.2016)

Date of mailing of the international search report

13 JAN 2017

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents  
 P.O. Box 1450, Alexandria, Virginia 22313-1450  
 Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300  
 PCT OSP: 571-272-7774

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US16/42680

**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
  
2.  Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3.  Claims Nos.: 10-29 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

\*\*\*-Please See Supplemental Page-\*\*\*-

1.  As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3.  As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4.  No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-9

**Remark on Protest**

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT  
Information on patent family members

International application No.  
PCT/US16/42680

-\*\*\*Continued from Box III-\*\*\*-

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Groups I+, Claims 1-9 and SEQ ID NO: 365 are directed toward a peptide inhibitor of an interleukin-23 receptor.

The peptide inhibitor will be searched to the extent that it encompasses SEQ ID NO: 365 (first exemplary peptide inhibitor). Applicant is invited to elect additional peptide(s) with specified SEQ ID NO: for each, or specified substitution(s) at specified site(s) of a SEQ ID NO:, such that the peptide sequence is fully specified (i.e. no optional or variable residues), to be searched. Additional peptide sequence(s) will be searched upon the payment of additional fees. It is believed that claims 1 (in-part), 2 (in-part), 3 (in-part), 4 (in-part) 5 (in-part), 8 (in-part) and 9 (in-part) encompass this first named invention and thus these claims will be searched without fee to the extent that they encompass SEQ ID NO: 365 (peptide inhibitor). Applicants must specify the claims that encompass any additionally elected peptide inhibitor sequence(s). Applicants must further indicate, if applicable, the claims which encompass the first named invention, if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched/examined. An exemplary election would be a peptide inhibitor encompassing SEQ ID NO: 366 (first exemplary elected peptide inhibitor).

No technical features are shared between the peptide inhibitor sequences of Groups I+ and, accordingly, these groups lack unity a priori.

Groups I+ share the technical features including: a peptide inhibitor of an interleukin-23 receptor, or a pharmaceutically acceptable salt or solvate thereof, wherein the peptide inhibitor comprises an amino acid sequence of Formula (Xa):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Xa) wherein X1 is any amino acid or absent; X2 is any amino acid or absent; X3 is any amino acid or absent; X4 is any amino acid or chemical moiety capable of forming a bond with X9; X5 is any amino acid; X6 is any amino acid; X7 is any amino acid; X8 is any amino acid; X9 is any amino acid or chemical moiety capable of forming a bond with X4; X10 is any amino acid; X11 is any amino acid; X12 is any amino acid; X13 is any amino acid; X14 is any amino acid; X15 is any amino acid, X16 is any amino acid or absent; X17 is any amino acid or absent; X18 is any amino acid or absent; X19 is any amino acid or absent; and X20 is any amino acid or absent, wherein the peptide inhibitor is cyclized via a bond between X4 and X9, and wherein the peptide inhibitor inhibits the binding of an interleukin-23 (IL-23) to an IL-23 receptor.

However, these shared technical features are previously disclosed by US 2013/0172272 A1 to Gallagher et al. (hereinafter 'Gallagher').

Gallagher discloses a peptide inhibitor of an interleukin-23 receptor (a peptide inhibitor of an interleukin-23 receptor; abstract) wherein the peptide inhibitor comprises an amino acid sequence of Formula (Xa):

X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Xa) wherein X1 is any amino acid or absent; X2 is any amino acid or absent; X3 is any amino acid or absent; X4 is any amino acid or chemical moiety capable of forming a bond with X9; X5 is any amino acid; X6 is any amino acid; X7 is any amino acid; X8 is any amino acid; X9 is any amino acid or chemical moiety capable of forming a bond with X4; X10 is any amino acid; X11 is any amino acid; X12 is any amino acid; X13 is any amino acid; X14 is any amino acid; X15 is any amino acid, X16 is any amino acid or absent; X17 is any amino acid or absent; X18 is any amino acid or absent; X19 is any amino acid or absent; and X20 is any amino acid or absent (wherein the peptide is 12 amino acids in length and comprises cysteine residues or cysteine and trp residues that form a cyclic peptide (wherein the peptide inhibitor comprises an amino acid sequence of Formula (Xa): X1-X2-X3-X4-X5-X6-X7-X8-X9-X10-X11-X12-X13-X14-X15-X16-X17-X18-X19-X20 (Xa) wherein X1 is any amino acid or absent; X2 is any amino acid or absent; X3 is any amino acid or absent; X4 is any amino acid or chemical moiety capable of forming a bond with X9; X5 is any amino acid; X6 is any amino acid; X7 is any amino acid; X8 is any amino acid; X9 is any amino acid or chemical moiety capable of forming a bond with X4; X10 is any amino acid; X11 is any amino acid; X12 is any amino acid; X13 is any amino acid; X14 is any amino acid; X15 is any amino acid, X16 is any amino acid or absent; X17 is any amino acid or absent; X18 is any amino acid or absent; X19 is any amino acid or absent; and X20 is any amino acid or absent); paragraphs [0027], [0114], wherein the peptide inhibitor is cyclized (wherein the peptide inhibitor is cyclized; paragraphs [0114], [0135]) via a bond between a cysteine residue and a tryptophan (via a bond between a cysteine residue and a tryptophan; paragraph [0114]). Gallagher further discloses wherein the peptide comprises a cysteine residue at X4 of the peptide (wherein the peptide comprises the sequence CWQDYWC (a cysteine residue at X4 of the peptide); Table 19, SEQ ID NO: 133), as well as wherein the peptide comprises a tryptophan residue at both X8 and X9 (as well as wherein the peptide comprises a tryptophan residue at both X8 and X9; paragraph [0048]) as a part of a WXXWWW motif (as a part of a WXXWWW motif; paragraph [0048]).

Gallagher does not disclose wherein the peptide inhibitor is cyclized via a bond between X4 and X9.

It would have been obvious to a person of ordinary skill in the art at the time of the invention was made to have modified the disclosure of Gallagher to have provided examples of peptides comprising cyclization between a cysteine residue positioned adjacent the WXXWWW motif, as disclosed by Gallagher, and either the tryptophan residue at X8 or X9, as disclosed by Gallagher, in order to assess their capacity to inhibit IL-23 binding to IL-23R, as disclosed by Gallagher, as well as the stability and active half-life of the peptide, based on the disclosure of Gallagher, in order to identify a form of the peptide with sufficient binding activity, as well as improved active half-life.

Since none of the special technical features of the Groups I+ inventions is found in more than one of the inventions, and since all of the shared technical features are previously disclosed by the Gallagher reference, unity of invention is lacking.

## 摘要

本发明提供白细胞介素-23 受体的新型肽抑制剂和相关的组合物以及使用这些肽抑制剂来治疗或预防各种疾病和病症，包括炎症性肠病的方法。