

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199710939 B2
(10) Patent No. 717603

(54) Title
Process for the preparation of 1-aryl-1-cyanocyclobutane derivatives

(51)⁷ International Patent Classification(s)
C07C 253/30

(21) Application No: **199710939**

(22) Application Date: **1996.11.28**

(87) WIPO No: **WO97/20810**

(30) Priority Data

(31) Number	(32) Date	(33) Country
9524681	1995.12.02	GB

(43) Publication Date : **1997.06.27**

(43) Publication Journal Date : **1997.08.21**

(44) Accepted Journal Date : **2000.03.30**

(71) Applicant(s)
Knoll Aktiengesellschaft

(72) Inventor(s)
Stephen John Barker; Sharon Michelle Clark

(74) Agent/Attorney
**WATERMARK PATENT and TRADEMARK ATTORNEYS, Locked Bag 5, HAWTHORN
VIC 3122**

(56) Related Art
KHIM-FARM ZH. 1980, 14(2) PAGES 40-45.

OPI DATE 27/06/97 APPLN. ID 10939/97
AOJP DATE 21/08/97 PCT NUMBER PCT/EP96/05246



AU9710939

INT

<p>(51) International Patent Classification⁶ : C07C 253/30</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/20810 (43) International Publication Date: 12 June 1997 (12.06.97)</p>
<p>(21) International Application Number: PCT/EP96/05246 (22) International Filing Date: 28 November 1996 (28.11.96) (30) Priority Data: 9524681.4 2 December 1995 (02.12.95) GB (71) Applicant (for all designated States except US): KNOLL AKTIENGESELLSCHAFT [DE/DE]; Knollstrasse, D-67061 Ludwigshafen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): BARKER, Stephen, John [GB/GB]; D107 Main Road, Beeston, Nottingham, Nottinghamshire NG9 1AD (GB). CLARK, Sharon, Michelle [GB/GB]; E50 Pennyfoot Street, Nottingham, Nottinghamshire NG1 1GF (GB). (74) Agents: MILLER, Thomas, Kerr et al.; BASF Aktiengesellschaft, D-67056 Ludwigshafen (DE).</p>		<p>(81) Designated States: AU, BG, BR, CA, CN, CZ, GE, HU, IL, JP, KR, LV, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TR, UA, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: PROCESS FOR THE PREPARATION OF 1-ARYL-1-CYANOCYCLOBUTANE DERIVATIVES</p>		
<p>(57) Abstract</p> <p>A process for the preparation of arylcyclobutyl nitrile derivatives which involves reaction of a solution of a 1,3-dihalopropane and a cyanobenzyl derivative in a substantially dimethyl sulphoxide-free solvent with a suspension of a base in a substantially dimethyl sulphoxide-free solvent at a temperature of at least 35 °C.</p>		

PROCESS FOR THE PREPARATION OF 1-ARYL-1-CYANOCYCLOBUTANE DERIVATIVES

The present invention relates to an improved process for the preparation of arylcyclobutyl cyanides. 1-(4-Chlorophenyl)cyclobutyl cyanide is an intermediate
5 useful for the preparation of sibutramine, N-1-[1-(4-chlorophenyl)cyclobutyl]-3-methylbutyl-N,N-dimethylamine. Sibutramine is useful in the treatment of depression, Parkinson's disease, obesity, Non Insulin Dependent Diabetes Mellitus (NIDDM) and epilepsy.

10 The reaction of phenylacetonitrile with a 1,3-dihalopropane in aqueous sodium hydroxide using benzyltriethylammonium chloride as a catalyst to give 1-phenylcyclobutyl cyanide is reported in Roc. Chem. 40, 1647, (1966). However, the yield is low (26%) and the amount of monoalkylated uncyclised product formed is significant (20%).

15

1-(4-Chlorophenyl)cyclobutyl cyanide was prepared by reacting 4-chlorophenylacetonitrile with 1,3-dibromobutane in a mixture of dimethyl sulphoxide and ether at 25-35°C using sodium hydride as the base (J.Org. Chem. 36 (9), 1308, 1971). It is also disclosed that the process is effective if the mineral oil is removed
20 from the sodium hydride by washing with toluene and then adding a slurry of sodium hydride in toluene to the dimethyl sulphoxide. Similar preparations are also described in US4,235,926, US3,526,656, US4,348,409, US5,405,866 and J.Organomet. Chem. 448, 1-2, p9-14(1993). The yields quoted vary between 43% and 78%.

25

GB2098602A discloses a process for the preparation of 1-(4-chlorophenyl)cyclobutyl cyanide comprising the reaction of 4-chlorophenylacetonitrile with a 1,3-dibromopropane in the presence of sodium hydride (dispersed in mineral oil). The reaction is described as being carried out in dry dimethyl
30 sulphoxide under nitrogen with stirring initially at room temperature, then at a temperature in the range 30 to 35°C for 2 hours. This preparation is also reported in EP 191542 and GB 2127819.

The presence of dimethyl sulphoxide in the aqueous waste from these processes renders the waste ineligible for discharge to the chemical effluent drain of chemical production plants. The waste therefore has to be specially disposed of. This leads to high production costs and adverse environmental effects (more
5 resources and energy are required to enable safe disposal of the aqueous waste). It is therefore desirable to find a process which does not require dimethyl sulphoxide.


Initially the reaction was attempted using toluene as the solvent. However, this course of action results in a new problem in that it leads to generation of a
10 significant, delayed exotherm during addition to the reaction mixture of the acrylonitrile. Such a process is not considered safe. The problem of exotherm generation does not arise when dimethyl sulphoxide is replaced with other water-miscible solvents, such as tetrahydrofuran. However, there is a significant loss of yield which can only be improved by partial distillation of the tetrahydrofuran and
15 addition of a water-immiscible solvent, such as toluene, prior to extraction. Such a procedure has the disadvantages of requiring extra processing (increasing the cost) and of creating a tetrahydrofuran/toluene waste stream, both of which render it unsatisfactory. A similar process is described in WO93/13073 (page 180, Example N10) for producing 1-(4-trifluoromethoxyphenyl)cyclobutyl cyanide. In this process
20 two water-miscible solvents, tetrahydrofuran and dimethylformamide, are used during the reaction, with the water-immiscible solvent ether being used for extraction of the product, to obtain a 61% yield. Again this has the disadvantage of requiring extra processing and producing a tetrahydrofuran/ether waste stream.

25 WO95/00489 describes a process for producing 1-(2-pyridyl)cyclopropyl cyanide. This reaction was carried out in toluene using a 50% aqueous sodium hydroxide solution as the base. The base was added to a stirred mixture of 2-(2-pyridyl)acetonitrile, 1-bromo-2-chloroethane, benzyltriethylammonium chloride and toluene at 25°C. The mixture was then heated at 70-75°C for 2 hours. The product
30 was extracted into ether and isolated in good yield (~85%). A disadvantage of this process is the presence of water in the initial reaction. This can lead to a rather high level of impurity formation. However, addition of an equivalent amount of a solid


base at 25°C (without the presence of water) would result in a significant delayed exotherm making such a reaction unsafe. Furthermore, it is well known that cyclobutyl rings, as described by the present invention, are considerably less facile to make than the cyclopropyl rings described in the above reference. Therefore it would not be expected that the above procedure would produce as good a yield of cyclobutyl material by using 1-bromo-2-chloropropane instead of 1-bromo-2-chloroethane. Additionally problems can arise from emulsion formation when water is present initially. This may lead to lower yields.

Surprisingly, we have found a process for the preparation of arylcyclobutyl cyanides whereby dimethyl sulphoxide can be excluded, delayed exotherms and mixed solvent waste streams avoided, and impurity formation kept to a minimum whilst still giving the desired product in good yield.


The present invention provides a process for the preparation of compounds of formula I



in which R_1 represents halo and R_2 represents hydrogen or halo; including the reaction of a 1,3-dihalopropane, a compound of formula II



in which R_1 and R_2 are as defined above, and a suspension of a base which is potassium hydroxide or sodium hydroxide in a solvent in which no more than 5% of dimethyl sulphoxide is present and wherein there is a phase transfer catalyst present in the suspension of the base at a temperature of at least 35°C.




A preferred process according to the present invention provides a process for the preparation of compounds of formula I in which R_1 represents chloro and R_2 represents hydrogen or chloro including the reaction of a 1,3-dihalopropane, a compound of formula II in which R_1 represents chloro and R_2 represents hydrogen or chloro, respectively, and a suspension of a base in a solvent in which no more than 5% of dimethyl sulphoxide is present at a temperature of at least 35°C.

The preferred processes of the present invention provide a) a process for the preparation of 1-(4-chlorophenyl)cyclobutyl cyanide including the reaction of a 1,3-dihalopropane, 4-chlorophenylacetonitrile and a suspension of a base in a solvent in which no more than 5% of dimethyl sulphoxide is present at a temperature of at least 35°C; and b) a process for the preparation of 1-(3,4-dichlorophenyl)cyclobutyl cyanide comprising the reaction of a 1,3-dihalopropane, 3,4-dichlorophenylacetonitrile and a suspension of a base in a solvent in which no more than 5% of dimethyl sulphoxide is present at a temperature of at least 35°C.

A more preferred process of the present invention provides a process for the preparation of 1-(4-chlorophenyl)cyclobutyl cyanide including the reaction of a 1,3-dihalopropane, 4-chlorophenylacetonitrile and a suspension of a base in a solvent in which no more than 5% of dimethyl sulphoxide is present at a temperature of at least 35°C.

Preferably the process includes the addition of a solution of a 1,3-dihalopropane and a compound of formula II in a solvent in which no more than 5% of dimethyl sulphoxide is present to a suspension of a base in a solvent in which no more than 5% of dimethyl sulphoxide is present at a temperature of at least 35°C.


The term "solvent" defines a water-immiscible liquid which is capable of keeping the 1,3-dihalopropane and the 4-chlorophenylacetonitrile in solution at the reaction temperature. The use of a water-immiscible liquid is advantageous as the work-up procedure is simplified and therefore process costs are reduced.



Suitably no more than 5% of dimethyl sulphoxide is present in the solvent, preferably no more than 2%, and most preferably there is a complete absence of dimethyl sulphoxide.

5 Suitably, solvent in which no more than 5% of dimethyl sulphoxide is present is a water immiscible organic liquid, preferably the liquid is non-polar. More preferably, solvent in which no more than 5% of dimethyl sulphoxide is present is a hydrocarbon such as toluene or petroleum ether. Most preferably the solvent in which no more than 5% of dimethyl sulphoxide is present is toluene.

10 Suitably, the base is potassium hydroxide or sodium hydroxide. Preferably the amount of base present is at least 2 molar equivalents relative to the amount of the compound of formula II present. More preferably, the amount is in the range of 3.8 to 4.7 molar equivalents relative to the amount of the compound of formula II present.




5 The suspension of base is preferably maintained by agitation such as stirring, shaking, or bubbling an inert gas, such as nitrogen, through the solvent, but any other means of maintaining a suspension may also be used. Preferably it is a stirred suspension.

10 Preferably the reaction is carried out under an inert atmosphere such as nitrogen.

Suitably the phase transfer catalyst is a quaternary salt or a crown ether.

- 15 Preferably the catalyst is selected from one of the following: butylpyridinium bromide, tetrabutylammonium bisulphate, benzyltriethylammonium bromide, benzyltriethylammonium chloride, benzyltrimethylammonium chloride, benzyltrimethylammonium fluoride, hexadecyltriethylammonium bromide, hexadecyltriethylphosphonium bromide, hexadecyltrimethylammonium bromide,
- 20 hexadecyltrimethylammonium chloride, dibutyltrimethylammonium chloride, decyltriethylammonium bromide, hexadecyltributylphosphonium bromide, heptylpyridinium bromide, hexadecyltributylphosphonium chloride, hexyltriethylammonium bromide, dodecylpyridinium bromide, dodecyltriethylammonium bromide, methyltrinonylammonium chloride,
- 25 methyltriphenylammonium bromide, tetrabutylammonium bromide or bisulphate, tetrabutylammonium chloride, tetrabutylammonium cyanide, tetrabutylammonium fluoride, tetrabutylammonium iodide, tetrabutylammonium hydroxide, tetrabutylphosphonium chloride, tricaprylmethylammonium chloride, tetraethylammonium chloride, tetramethylammonium bromide,
- 30 trioctylethylphosphonium bromide, trioctylmethylammonium chloride, trioctylpropylammonium chloride, tetrapropylammonium bromide, tetraphenylarsonium chloride, tetraphenylphosphonium bromide,



tetraphenylphosphonium chloride, benzyltrimethylammonium hydroxide, 18-crown-6, dibenzo-18-crown-6, dicyclohexyl-18-crown-6 or mixtures thereof. More preferably the phase transfer catalyst is a quaternary ammonium salt or a crown ether. Most preferably, the phase-transfer catalyst is tetra-n-butylammonium bromide, tetra-n-
5 butylammonium hydrogen sulphate, or tetra-n-butylammonium iodide.

Preferably the amount of phase-transfer catalyst present is in the range of 0.01 to 0.2 molar equivalents relative to the amount of the compound of formula II present. More preferably the amount is in the range of 0.05 to 0.15 molar
10 equivalents relative to the amount of the compound of formula II present.

Preferably, the temperature is in the range 35-80°C, more preferably in the range 35.1-69°C, most preferably in the range 40-60°C.

15 Preferably, water is added to aid stirring when the addition is 60-85% complete, more preferably 75% complete. Suitably the volume of water added is in the range of 0 to 5.0 parts by weight relative to the weight of the compound of formula II present. Preferably the volume of water added is in the range of 0 to 1.0 parts of weight relative to the weight of the compound of formula II present. More
20 preferably the volume of water added is on the range of 0.7 to 0.9 parts by weight relative to the weight of the compound of formula II present.

Preferably, the reaction is quenched by the addition of water.

25 Preferably, the reaction is carried out at atmospheric pressure.

Suitably the 1,3-dihalopropane is 1,3-dibromopropane, 1,3-dichloropropane or 1-bromo-3-chloropropane. Preferably the 1,3-dihalopropane is 1,3-dibromopropane.

30

Suitably the amount of 1,3-dihalopropane used is in the range of 0.8 to 1.5 molar equivalents relative to the amount of the compound of formula II present.

Preferably the amount of 1,3-dihalopropane used is in the range of 0.9 to 1.2 molar equivalents relative to the amount of the compound of formula II present. Most preferably the amount of 1,3-dihalopropane used is in the range 1.0 to 1.05 molar equivalents relative to the amount of the compound of formula II present.

5

When compounds of formula I are prepared by the process which comprises the present invention, there can be observed a significant financial saving both in terms of raw materials and disposal of the aqueous waste when compared to the known process carried out in dimethyl sulphoxide. Also, there is additional benefit to the environment because it obviates the need to dispose of waste dimethyl sulphoxide. In addition certain impurities which arise from the oxidising nature of dimethyl sulphoxide are eliminated leading to simplified work-up procedures and a purer product.

15 A further advantage of the present invention is that it may avoid the need for isolation of 1-(4-chlorophenyl)cyclobutyl cyanide when it is desired to obtain sibutramine. Instead it may be possible to use the toluene solution of 1-(4-chlorophenyl)cyclobutanecarbonitrile immediately in the reaction described in GB2098602A, incorporated herein by reference.

20

In one embodiment of the present invention, the process comprises the addition of a solution of 1,3-dibromopropane and 4-chlorophenylacetonitrile in toluene to a stirred suspension of powdered potassium hydroxide with tetra-n-butylammonium bromide in toluene at a temperature in the range 35-80°C, preferably in the range 35.1-69°C, more preferably in the range 40-60°C. Water is added after 60-85% completion of the addition. The reaction is quenched by the addition of water.

25 The invention is illustrated by the following Examples which are given by way of example only. The final product of each of these Examples was characterised by one or more of the following procedures: gas-liquid chromatography; high

performance liquid chromatography; elemental analysis; nuclear magnetic resonance spectroscopy and infrared spectroscopy.

5

Examples

The following procedure was carried out under the conditions listed in Table 1 to obtain compounds of formula I.

10 75% of a mixture (m) of 4-chlorophenylacetonitrile (II) (x g) and 1,3-dibromopropane (y g) in toluene (z ml) was added to a stirred mixture of a base (a g) and a catalyst (b g) in toluene (c ml) over 1.5 hours at a temperature of d°C. Water (e ml) was added, keeping the temperature at d°C. The remaining 25% of the mixture (m) was then added over 30 minutes at d°C, and the mixture stirred for 2.5
15 hours at f°C. The reaction was then quenched by addition of water (254 ml) over 15 minutes at d°C, and the mixture stirred for 20 minutes.

The organic phase was separated and stirred with water (354 ml) and caustic soda (76 g) at f°C for 15 minutes, then allowed to settle. The organic layer was
20 separated, then stirred at f°C with water (300 ml) and concentrated hydrochloric acid (20 ml), the aqueous layer having a pH of 3 or less. The organic layer was separated, then stirred with water (300 ml) at f°C for 15 minutes and the organic layer separated. This was repeated until the aqueous layer had a pH between 6 and
8.

25

The solvent was removed in vacuo at 90°C, and the residual oil was distilled at 1.33 to 2.66 mbar under high vacuum to yield in the appropriate fractions a compound of formula I, yield g%.

Table 1

Examples 1-10

Mass of compound II (x g)	Mass of dibromopropane (y g)	Catalyst	Mass of catalyst (b g)	Mass of base (powdered potassium hydroxide)(a g)	Temperature (d°C) and (f°C)	Toluene charge (z ml) (c ml)		Water charged @ 75% of addition (e ml)	Yield (g%)
75	105	TBAB	11.8	132	40	66	360	60	53.2
75	105	TBAB	11.8	132	40	66	360	60	57.3
75	105	TBAB	11.8	132	40	66	360	60	61.4
75	105	TBAB	11.8	132	40	66	360	0	56.5
75	105	TBAB	11.8	132	60	66	360	60	57.5
75	105	TBAB	11.8	132	60	66	360	60	63.2
75	105	TBAHS	12.4	132	40	66	360	0	57.8
75	105	TBAHS	12.4	132	40	66	360	0	62.6
75	105	TBAHS	12.4	132	60	66	360	60	61.3
75	105	TBAHS	12.4	132	60	66	360	60	57.9

5 TBAB means tetra-n-butylammonium bromide; TBAHS means tetra-n-butylammonium hydrogen sulphate. Temperatures d°C and f°C are the same for Examples 1-10.

The same procedure was carried out for Comparative Examples 11-15, except that tetrahydrofuran was used in place of toluene. The temperature (d°C) for the initial reaction was the temperature used in the original process with dimethylsulphoxide, viz room temperature (20-25°C), rising to 30-35°C (f°C) for the
5 final stir (for 1.5 hours not 2.5 hours) and separation steps. The only exception was Example 15 where the separation steps were carried out at 20-25°C (d°C) instead of 30-35°C (f°C). The other conditions and yields for these Examples are listed in Table 2. It can be seen that the yields are inferior to the results obtained using toluene at a temperature of at least 35°C.

10

The same procedure was carried out for Comparative Examples 16-17, except that a mixture of dimethyl sulphoxide and toluene was used in place of toluene. The temperature (d°C) for the initial reaction was the temperature used in the original process with dimethylsulphoxide, viz room temperature (20-25°C), rising
15 to 30-35°C (f°C) for the final stir (for 1.5 hours not 2.5 hours) and separation steps. The other conditions and yields for these Examples are listed in Table 3. It can be seen that the yields are comparable to the results obtained using toluene at a temperature of at least 35°C. However, as mentioned previously there is the disadvantage of waste dimethyl sulphoxide.

Table 2

Comparative Examples 11-15

Mass of compound II (x g)	Mass of dibromopropane (y g)	Catalyst	Mass of catalyst (b g)	Mass of potassium hydroxide (a g)	Temperature (d°C, f°C)	Tetrahydrofuran charge (z+c) (including all washes) (ml)	Water charged @ 75% of addition (e ml)	Yield (g%)
75	105	TBAB	11.8	132	20-25, 30-35	640	60	29.4
75	105	TBAB	11.8	132	20-25, 30-35	640	0	36.4
75	105	TBAB	11.8	132	20-25, 30-35	790	0	20.3
75	105	TBAB	11.8	132	20-25, 30-35	640	0	44.1
75	105	TBAB	11.8	132	20-25, 30-35	840	0	35.3

5 TBAB means tetra-n-butylammonium bromide; TBAHS means tetra-n-butylammonium hydrogen sulphate.

Table 3

Comparative Examples 16-17

Mass of compound II (x g)	Mass of dibromopropane (y g)	Catalyst	Mass of catalyst (b g)	Mass of potassium hydroxide (a g)	Temperature (d°C, f°C)	DMSO/toluene charge (z+c) (including all washes) (ml)	Water charged @ 75% of addition (e ml)	Yield (g%)
150	209.6	TBAB	22.4	264	20-25, 30-35	120/1080	120	62
75	104.8	TBAB	11.2	132	20-25, 30-35	60/345	60	55


10 TBAB means tetra-n-butylammonium bromide; TBAHS means tetra-n-butylammonium hydrogen sulphate; DMSO means dimethyl sulphoxide.

Example 18


75% of a mixture (m) of 1-(3,4-dichlorophenyl)acetonitrile (92.1 g) and 1,3-dibromopropane (105 g) in toluene (66 ml) is added to a stirred mixture of powdered potassium hydroxide (132 g) and tetra-n-butylammonium bromide (11.8 g) in toluene (360 ml) over 1.5 hours at a temperature of 60°C. Water (60 ml) is added, keeping the temperature at 60°C. The remaining 25% of the mixture (m) is then added over 30 minutes at 60°C, and the mixture stirred for 2.5 hours at 60°C. The reaction is then quenched by addition of water (254 ml) over 15 minutes at 60°C, and the mixture stirred for 20 minutes. The organic phase is separated and stirred with water (354 ml) and caustic soda (76 g) at 60°C for 15 minutes, then allowed to settle. The organic layer is separated, then stirred at 60°C with water (300 ml) and concentrated hydrochloric acid (20 ml), the aqueous layer having a pH of 3 or less. The organic layer is separated, then stirred with water (300 ml) at 60°C for 15 minutes and the organic layer separated. This is repeated until the aqueous layer has a pH between 6 and 8. The solvent is removed in vacuo at 90°C, and the residual oil is distilled at 1.33 to 2.66 mbar under high vacuum to yield in the appropriate fractions of 1-(3,4-dichlorophenyl)cyclobutyl cyanide.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A process for the preparation of compounds of formula I as defined by formula I



in which R_1 represents halo and R_2 represents hydrogen or halo; including the reaction of a 1,3-dihalopropane, a compound of formula II



in which R_1 and R_2 are as defined above, and a suspension of a base which is potassium hydroxide or sodium hydroxide in a solvent in which no more than 5% of dimethyl sulphoxide is present, wherein there is a phase-transfer catalyst present in the suspension of the base, at a temperature of at least 35°C.

2. A process as claimed in claim 1 including the addition of a solution of a 1,3-dihalopropane and a compound of formula II in a solvent in which no more than 5% of dimethyl sulphoxide is present to a suspension of a base in a solvent in which no more than 5% of dimethyl sulphoxide is present at a temperature of at least 35°C.

3. A process as claimed in either claim 1 or claim 2 in which the phase-transfer catalyst is tetra-n-butylammonium bromide, tetra-n-butylammonium hydrogen sulphate, tetra-n-butylammonium iodide or a crown ether.



4. A process as claimed in any one of the preceding claims in which the solvent is toluene.
5. A process as claimed in any one of the preceding claims in which the temperature is in the range 35-80°C.
6. A process as claimed in any one of the preceding claims in which the compound of formula II is 4-chlorophenylacetonitrile.
7. A process as claimed in any one of the preceding claims in which the compound of formula II is 3,4-dichlorophenylacetonitrile.
8. A process as claimed in any one of the preceding claims in which water is added after 60-85% of the addition is complete.
9. A process as claimed in any one of the preceding claims in which the 1,3 dihalopropane is 1,3-dibromopropane.
10. A process according to claim 1 and as substantially herein described with reference to the examples.

DATED this 29th day of December 1999

KNOLL AKTIENGESELLSCHAFT

WATERMARK PATENT AND TRADEMARK ATTORNEYS
290 BURWOOD ROAD
HAWTHORN VICTORIA 3122
AUSTRALIA

LCG:CLR:VRH P2492AU00.DOC

