



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2491349 C 2011/04/26

(11)(21) **2 491 349**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2003/06/30
(87) Date publication PCT/PCT Publication Date: 2004/01/22
(45) Date de délivrance/Issue Date: 2011/04/26
(85) Entrée phase nationale/National Entry: 2004/12/30
(86) N° demande PCT/PCT Application No.: EP 2003/006888
(87) N° publication PCT/PCT Publication No.: 2004/006666
(30) Priorité/Priority: 2002/07/10 (DE102 31 295.8)

(51) Cl.Int./Int.Cl. *A01N 43/32*(2006.01),
A01N 43/653(2006.01), *A01P 3/00*(2006.01)


(72) Inventeurs/Inventors:
AMMERMANN, EBERHARD, DE;
STIERL, REINHARD, DE;
SCHOEFL, ULRICH, DE;
SCHELBERGER, KLAUS, DE;
SCHERER, MARIA, DE;
HENNINGSEN, MICHAEL, DE;
GOLD, RANDALL EVEN, DE

(73) Propriétaire/Owner:
BASF AKTIENGESELLSCHAFT, DE

(74) Agent: ROBIC

(54) Titre : MELANGES FONGICIDES A BASE DE DITHIANONE

(54) Title: FUNGICIDAL MIXTURES BASED ON DITHIANONE



(57) Abrégé/Abstract:

The invention relates to fungicidal mixtures containing, in a synergistically effective amount: A) the compound of formula (I) and; B) at least one azole derivative II selected from the group consisting of compounds (II-1) to (II-7). The invention also relates to methods for controlling harmful fungi by using mixtures of compounds (I) and (II), and to the use of compounds (I) and (II) for producing mixtures of this type.

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro(43) Internationales Veröffentlichungsdatum
22. Januar 2004 (22.01.2004)

PCT

(10) Internationale Veröffentlichungsnummer
WO 2004/006666 A3

(51) Internationale Patentklassifikation⁷: A01N 43/32 // (A01N 43/32, 43:653)

(21) Internationales Aktenzeichen: PCT/EP2003/006888

(22) Internationales Anmelde datum: 30. Juni 2003 (30.06.2003)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität: 102 31 295.8 10. Juli 2002 (10.07.2002) DE

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): BASF AKTIENGESELLSCHAFT [DE/DE]; 67056 Ludwigshafen (DE).

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): AMMERMANN, Eberhard [DE/DE]; Von-Gagern-Str. 2, 64646 Heppenheim (DE). STIERL, Reinhard [DE/DE]; Jahnstr. 8, 67251 Freinsheim (DE). SCHÖFL, Ulrich [DE/DE]; Luftschifftring 22c, 68782 Brühl (DE). SCHELBERGER, Klaus [AT/DE]; Traminerweg 2, 67161 Gönnheim (DE). SCHERER, Maria [DE/DE]; Hermann-Jürgens-Str. 30, 76829 Godramstein (DE). HENNINGSEN, Michael [DE/DE]; Tulpenstr. 7, 67227 Frankenthal (DE). GOLD, Randall, Even [US/DE]; Birkenweg 3, 67283 Obrigheim (DE).

(74) Gemeinsamer Vertreter: BASF AKTIENGESELLSCHAFT; 67056 Ludwigshafen (DE).


(81) Bestimmungsstaaten (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE,

[Fortsetzung auf der nächsten Seite]

(54) Title: FUNGICIDAL MIXTURES BASED ON DITHIANON

(54) Bezeichnung: FUNGIZIDE MISCHUNGEN AUF BASIS VON DITHIANON

WO 2004/006666 A3



(57) Abstract: The invention relates to fungicidal mixtures containing, in a synergistically effective amount: A) the compound of formula (I) and; B) at least one azole derivative II selected from the group consisting of compounds (II-1) to (II-7). The invention also relates to methods for controlling harmful fungi by using mixtures of compounds (I) and (II), and to the use of compounds (I) and (II) for producing mixtures of this type.

[Fortsetzung auf der nächsten Seite]



GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

Veröffentlicht:

- mit internationalem Recherchenbericht
- vor Ablauf der für Änderungen der Ansprüche geltenden Frist; Veröffentlichung wird wiederholt, falls Änderungen eintreffen

(84) Bestimmungsstaaten (regional): ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(88) Veröffentlichungsdatum des internationalen Recherchenberichts:

1. Juli 2004


Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

(57) Zusammenfassung: Fungizide Mischungen, enthaltend A) die Verbindung der Formel (I) und B) mindestens ein Azolderivat II ausgewählt aus der Gruppe der Verbindungen (II-1) bis (II-7) in einer synergistisch wirksamen Menge, Verfahren zur Bekämpfung von Schadpilzen mit Mischungen der Verbindungen (I) und (II) und die Verwendung der Verbindungen (I) und (II) zur Herstellung derartiger Mischungen.

FUNGICIDAL MIXTURES BASED ON DITHIANON

The present invention as broadly disclosed relates to fungicidal mixtures, comprising:


A) the compound of the formula I




I

and


10 B) at least one azole derivative II selected from the group of the compounds II-1 to II-7




II-1




II-2



II-3




II-4




II-5

2



II-6




II-7

10 in a synergistically effective amount.

The present invention as more particularly claimed concerns a fungicidal mixture, comprising:


A) the compound of the formula I:



I

and

20 B) an azole derivative II of formula II-2,



II-2

in a synergistically effective amount.

Moreover, the invention relates to a fungicidal composition comprising a solid or liquid carrier and the mixture as described above.

2a

Moreover, the invention relates to methods for controlling harmful fungi using mixtures of the compounds I and II and to the use of the compounds I and II for preparing such mixtures.

More particularly, the invention thus relates to a method for controlling harmful fungi, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with the fungicidal mixture as described above.

In a preferred embodiment, the method comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with from 5 to 2000 g/ha of the compound I.

In another preferred embodiment, the method comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with from 5 to 500 g/ha of the compound II.

The compound of the formula I (common name: dithianon) and processes for its preparation are described in GB-A 857 383.


The compounds of the formulae II-1 to II-7, their preparation and their action against harmful fungi are likewise known from the literature:

Compound No.	common name	Literature
II-1	metconazole	EP-A 267 778
II-2	epoxiconazole	EP-A 094 564
II-3	fluquinconazole	Pesticide Manual, 12th Ed., p.449 (2000)
II-4	tebuconazole	EP-A 040 345
II-5	tetraconazole	EP-A 234 242
II-6	difenoconazole	EP-A 065 485
II-7	prothioconazole	WO-A 96/16048

2b

It is an object of the present invention to provide mixtures which have improved activity against harmful fungi combined with a reduced total amount of active compounds applied (synergistic mixtures), with a view to reducing the application rates and broadening the activity spectrum of the known compounds.

We have found that this object is achieved by the mixtures defined at the outset. Moreover, we have found that applying the compounds I and II simultaneously, i.e. together or separately, or applying the compounds I and II in succession provides better



control of harmful fungi than is possible with the individual compounds alone.

Usually, mixtures of the compound I with one azole derivative II 5 are used. However, in certain cases mixtures of the compound I with two or more azole derivatives II may be advantageous.

Particular preference is given to the compounds II-1, II-2 and II-3. Especially preferred are mixtures comprising the compound 10 II-1. In another embodiment of the mixtures according to the invention, preference is given to the compound of the formula II-3.

Owing to their basic character, the compounds II-1 to II-7 are 15 capable of forming salts or adducts with inorganic or organic acids or with metal ions.

Examples of inorganic acids are hydrohalic acids, such as hydrogen fluoride, hydrogen chloride, hydrogen bromide and 20 hydrogen iodide, sulfuric acid, phosphoric acid, carbonic acid and nitric acid.

Suitable organic acids are, for example, formic acid, and alkanoic acids, such as acetic acid, trifluoroacetic acid, 25 trichloroacetic acid and propionic acid, and also glycolic acid, thiocyanic acid, lactic acid, succinic acid, citric acid, benzoic acid, cinnamic acid, oxalic acid, alkylsulfonic acids (sulfonic acids having straight-chain or branched alkyl radicals with 1 to 20 carbon atoms), arylsulfonic acids or aryldisulfonic acids 30 (aromatic radicals, such as phenyl and naphthyl, which carry one or two sulfo groups), alkylphosphonic acids (phosphonic acids having straight-chain or branched alkyl radicals with 1 to 20 carbon atoms), arylphosphonic acids or aryldiphosphonic acids (aromatic radicals, such as phenyl and naphthyl, which carry one 35 or two phosphonic acid radicals), it being possible for the alkyl or aryl radicals to carry further substituents, for example p-toluenesulfonic acid, salicylic acid, p-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, etc.

40 Suitable metal ions are, in particular, the ions of the elements of the second main group, in particular calcium and magnesium, of the third and fourth main group, in particular aluminum, tin and lead, and of the first to eighth transition group, in particular chromium, manganese, iron, cobalt, nickel, copper, zinc and 45 others. Particular preference is given to the metal ions of the elements of the transition groups of the fourth period. The

metals can be present in the various valences which they can assume.

When preparing the mixtures, it is preferred to employ the pure 5 active compounds I and II, with which further active compounds against harmful fungi or other pests, such as insects, arachnids or nematodes, or else herbicidal or growth-regulating active compounds or fertilizers can be admixed as required.

10 The mixtures of the compounds I and II, or the simultaneous joint or separate use of the compounds I and II, have outstanding action against a wide range of phytopathogenic fungi, in particular from the classes of the *Ascomycetes*, *Deuteromycetes*, *Oomycetes* and *Basidiomycetes*. Some of them act systemically and 15 are therefore also suitable for use as foliar- and soil-acting fungicides.

They are especially important for controlling fungi in a variety of crop plants, such as vegetable species (for example cucumbers, 20 beans and cucurbits), fruit species, grapevine, but also barley, grass, oats, coffee, corn, rye, soya, wheat, ornamentals, sugarcane, and a variety of seeds.

They are particularly suitable for controlling the following 25 phytopathogenic fungi: *Erysiphe graminis* (powdery mildew) in cereals, *Erysiphe cichoracearum* and *Sphaerotheca fuliginea* in cucurbits, *Podosphaera leucotricha* in apples, *Uncinula necator* in grapevines, *Puccinia* species in cereals, *Rhizoctonia* species in cotton, rice and lawns, *Ustilago* species in cereals and 30 sugarcane, *Venturia inaequalis* (scab) in apples, *Helminthosporium* species in cereals, *Septoria nodorum* in wheat, *Botrytis cinerea* (gray mold) in strawberries, vegetables, ornamentals and grapevines, *Cercospora arachidicola* in groundnuts, *Pseudocercospora herpotrichoides* in wheat and barley, 35 *Pseudoperonospora* species in cucurbits and hops, *Plasmopara viticola* in grapevines, *Alternaria* species in vegetables and fruit and *Fusarium* and *Verticillium* species.

Furthermore, they can be used in the protection of materials (for 40 example the protection of wood), for example against *Paecilomyces variotii*.

The compounds I and II can be applied simultaneously, that is either together or separately, or in succession, the sequence, in 45 the case of separate application, generally not having any effect on the control results.

The compounds I and II are usually applied in a weight ratio of from 100:1 to 1:10, preferably from 10:1 to 1:1, in particular from 5:1 to 1:1.

5 Correspondingly, the application rates of the compound I are usually from 5 to 2 000 g/ha, preferably from 10 to 1 000 g/ha, in particular from 50 to 750 g/ha.

Depending on the nature of the desired effect, the application 10 rates of the mixtures according to the invention are, for the compounds II, from 5 g/ha to 500 g/ha, preferably from 50 to 500 g/ha, in particular from 50 to 200 g/ha.

For seed treatment, the application rates of the mixture are 15 generally from 0.001 to 1 g/kg of seed, preferably from 0.01 to 0.5 g/kg, in particular from 0.01 to 0.1 g/kg.

If phytopathogenic harmful fungi are to be controlled, the separate or joint application of the compounds I and II or of the 20 mixtures of the compounds I and II is effected by spraying or dusting the seeds, the plants or the soils before or after sowing, or before or after plant emergence.

The following are examples of formulations:

25

1. Products for dilution with water

A) Water-soluble concentrates (SL)

30 10 parts by weight of the active compounds are dissolved in water or in a water-soluble solvent. As an alternative, wetters or other auxiliaries are added. The active compound dissolves upon dilution with water.

B) Dispersible concentrates (DC)

35 20 parts by weight of the active compounds are dissolved in cyclohexanone with addition of a dispersant, for example polyvinylpyrrolidone. Dilution with water gives a dispersion.

C) Emulsifiable concentrates (EC)

40 15 parts by weight of the active compounds are dissolved in xylene with addition of calcium dodecylbenzenesulfonate and castor oil ethoxylate (in each case 5% strength). Dilution with water gives an emulsion.

45 D) Emulsions (EW, EO)

40 parts by weight of the active compounds are dissolved in xylene with addition of calcium dodecylbenzenesulfonate and

castor oil ethoxylate (in each case 5% strength). This mixture is introduced into water by means of an emulsifying machine (Ultraturrax*) and made into a homogeneous emulsion. Dilution with water gives an emulsion.

E) Suspensions (SC, OD)

In an agitated ball mill, 20 parts by weight of the active compounds are comminuted with addition of dispersants, wetters and water or an organic solvent to give a fine active compound suspension. Dilution with water gives a stable suspension of the active compound.

F) Water-dispersible granules and water-soluble granules (WG, SG)

10 50 parts by weight of the active compounds are ground finely with addition of dispersants and wetters and prepared as water-dispersible or water-soluble granules by means of technical appliances (for example extrusion, spray tower, fluidized bed). Dilution with water gives a stable dispersion or solution of the active compound.

G) Water-dispersible powders and water-soluble powders (WP, SP)

75 parts by weight of the active compounds are ground in a rotor-stator mill with addition of dispersants, wetters and silica gel. Dilution with water gives a stable dispersion or solution of the active compound.

2. Products to be applied undiluted

H) Dustable powders (DP)

20 5 parts by weight of the active compounds are ground finely and mixed intimately with 95% of finely divided kaolin. This gives a dustable product.

I) Granules (GR, FG, GG, MG)

0.5 part by weight of the active compounds is ground finely and combined with 95.5% of carriers. Current methods are extrusion, spray-drying or the fluidized bed. This gives granules to be applied undiluted.


J) ULV solutions (UL)

10 parts by weight of the active compounds are dissolved in an organic solvent, for example xylene. This gives a product to be applied undiluted.

* trademark

6a

The active compounds can be used as such, in the form of their formulations or the use forms prepared therefrom, for example in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dustable products, materials for spreading, or granules, by means of



spraying, atomizing, dusting, spreading or pouring. The use forms depend entirely on the intended purposes; they are intended to ensure in each case the finest possible distribution of the active compounds according to the invention.

5 Aqueous use forms can be prepared from emulsion concentrates, pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances, as such or dissolved in an oil or solvent, can be homogenized in water by means of a wetter, tackifier, dispersant or emulsifier. However, it is also possible to prepare 10 concentrates composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

15 The active compound concentrations in the ready-to-use preparations can be varied within relatively wide ranges. In general, they are from 0.0001 to 10%, preferably from 0.01 to 1%.

20 The active compounds may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply formulations comprising over 95% by weight of active compound, or even to apply the active compound without additives.

25 Oils of various types, wetters, adjuvants, herbicides, fungicides, other pesticides, or bactericides may be added to the active compounds, even, if appropriate, not until immediately prior to use (tank mix). These agents can be admixed with the compositions according to the invention typically in a weight ratio of from 1:10 to 10:1.

The fungicidal activity of the compounds and the mixtures can be demonstrated by the following experiments:

30 The active compounds were prepared separately or jointly as a stock solution comprising 0.25% by weight of active compound in acetone or DMSO. 1% by weight of the emulsifier Uniperol® EL (wetting agent having emulsifying and dispersing action based on 35 ethoxylated alkylphenols) was added to this solution, and the mixture was diluted with water to the desired concentration.

Use Example 1 - Activity against gray mold on bell pepper leaves caused by *Botrytis cinerea*

40 Bell pepper seedlings of the cultivar "Neusiedler Ideal Elite" were, after 4-5 leaves were well developed, sprayed to runoff point with an aqueous suspension having the concentration of active compound stated below. The next day, the treated plants 45 were inoculated with a spore suspension of *Botrytis cinerea* which contained 1.7×10^6 spores/ml in a 2% strength aqueous biomalt solution. The test plants were then placed in a climatized

chamber at 22-24°C and high atmospheric humidity. After 5 days, the extent of the fungal infection on the leaves could be determined visually in %.

5 Evaluation was carried out by determining the infected leaf areas in percent. These percentages are converted into efficacies.

The efficacy (E) is calculated as follows using Abbot's formula:

10

$$E = (1 - \alpha/\beta) \cdot 100$$

α corresponds to the fungal infection of the treated plants in % and

15 β corresponds to the fungal infection of the untreated (control) plants in %

An efficacy of 0 means that the infection level of the treated plants corresponds to that of the untreated control plants; an 20 efficacy of 100 means that the treated plants were not infected.

The expected efficacies of the active compound mixtures are determined using Colby's formula [S.R. Colby, Weeds 15, 20-22 (1967)] and compared with the observed efficacies.

25

Colby's formula:

$$E = x + y - x \cdot y / 100$$

30 E expected efficacy, expressed in % of the untreated control, when using the mixture of the active compounds A and B at the concentrations a and b

35 x efficacy, expressed in % of the untreated control, when using active compound A at a concentration of a

y efficacy, expressed in % of the untreated control, when using active compound B at a concentration of b

40

45

Table A - Individual active compounds

Example	Active compounds	Concentration of active compound in the spray liquor	Efficacy in % of the untreated control
1	Control (untreated)	(99% infection)	0
2	I (dithianon)	16 4 1 0.25	0 0 0 0
3	II-1 (metconazole)	1 0.25	49 0
4	II-2 (epoxiconazole)	1 0.25	9 9
5	II-4 (tebuconazole)	4 1 0.25	0 0 0

Table B - Combinations according to the invention

Example	Active compound mixture Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
6	I + II-1 4 + 0.25 ppm 16 : 1	19	0
7	I + II-1 4 + 1 ppm 4 : 1	59	49
8	I + II-1 1 + 0.25 ppm 4 : 1	39	0
9	I + II-1 0.25 + 0.25 ppm 1 : 1	29	0
10	I + II-1 0.25 + 1 ppm 1 : 4	59	49
11	I + II-2 4 + 0.25 ppm 16 : 1	39	9
12	I + II-2 4 + 1 ppm 4 : 1	49	9

Example	Active compound mixture Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
5 13	I + II-2 1 + 0.25 ppm 4 : 1	39	9
10 14	I + II-2 1 + 1 ppm 1 : 1	59	9
15 15	I + II-2 0.25 + 0.25 ppm 1 : 1	29	9
15 16	I + II-2 0.25 + 1 ppm 1 : 4	44	9
17	I + II-4 4 + 0.25 ppm 16 : 1	29	0
20 18	I + II-4 16 + 4 ppm 4 : 1	59	0
19	I + II-4 4 + 1 ppm 4 : 1	39	0
25 20	I + II-4 1 + 1 ppm 1 : 1	29	0
21	I + II-4 0.25 + 0.25 ppm 1 : 1	19	0
30 22	I + II-2 1 + 4 ppm 1 : 4	49	0

*) efficacy calculated using Colby's formula

35 Use Example 2 - Activity against early blight of tomato caused by *Alternaria solani*

Leaves of potted plants of the cultivar "Große Fleischtomate St. Pierre" were sprayed to runoff point with an aqueous suspension

40 having the concentration of active compounds stated below. The next day, the leaves were infected with an aqueous zoospore suspension of *Alternaria solani* in 2% biomalt solution having a density of 0.17×10^6 spores/ml. The plants were then placed in a water-vapor-saturated chamber at temperatures of between 20 and
45 22°C. After 5 days, the infection on the leaves of the untreated,

11

but infected control plants had developed to such an extent that the infection could be determined visually in %.

Table C - Individual active compounds

Example	Active compound	Concentration of active compound in the spray liquor [ppm]	Efficacy in % of the untreated control
23	Control (untreated)	(81% infection)	0
24	I (dithianon)	4 1 0.25	0 0 0
25	II-1 (metconazole)	1 0.25	63 2
26	II-2 (epoxiconazole)	1 0.25	75 63
27	II-4 (tebuconazole)	1 0.25	63 0

Tabelle D - Combinations according to the invention

Example	Active compound mixture Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
28	I + II-1 4 + 0.25 ppm 16 : 1	63	2
29	I + II-1 4 + 1 ppm 4 : 1	75	63
30	I + II-1 1 + 0.25 ppm 4 : 1	26	2
31	I + II-1 0.25 + 0.25 ppm 1 : 1	63	2
32	I + II-1 0.25 + 1 ppm 1 : 4	75	63
33	I + III-2 4 + 0.25 ppm 16 : 1	75	63
34	I + III-2 1 + 0.25 ppm 4 : 1	75	63

Example	Active compound mixture Concentration Mixing ratio	Observed efficacy	Calculated efficacy*)
5 35	I + II-2 0.25 + 0.25 ppm 1 : 1	82	63
10 36	I + II-2 0.25 + 1 ppm 1 : 4	88	75
15 37	I + II-4 4 + 0.25 ppm 16 : 1	75	0
38	I + II-4 1 + 0.25 ppm 4 : 1	26	0
20 39	I + II-4 1 + 1 ppm 1 : 1	82	63
40	I + II-4 0.25 + 0.25 ppm 1 : 1	26	0
41	I + II-4 0.25 + 1 ppm 1 : 4	75	63

25 *) efficacy calculated using Colby's formula

The test results show that, for all mixing ratios, the observed efficacy is higher than the efficacy predicted using Colby's formula.

30

35


40

45

WHAT IS CLAIMED IS:


1. A fungicidal mixture, comprising:

A) the compound of the formula I:



and

B) an azole derivative II of formula II-2,




in a synergistically effective amount.






2. The fungicidal mixture as claimed in claim 1, further comprising at least one

10 azole derivative selected from the group of the compounds II-1, II-3, II-4, II-5, II-6 and

II-7:




14




3. The fungicidal mixture as claimed in claim 1 or 2, wherein the weight ratio of the compound of formula I to the azole derivative of formula II-2 is from 100:1 to 1:10.
4. A fungicidal composition comprising a solid or liquid carrier and the mixture as claimed in any one of claims 1 to 3.


5. A method for controlling harmful fungi, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with the fungicidal mixture as claimed in any one of claims 1 to 3.
6. The method as claimed in claim 5, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with from 5 to 2000 g/ha of the compound of formula I.
7. The method as claimed in claim 5, which comprises treating the harmful fungi, their habitat, or the plants, seeds, soils, areas, materials or spaces to be kept free from them with from 5 to 500 g/ha of the azole derivative of formula II-2.
- 10 8. The method of any one of claims 5 to 7 wherein the harmful fungi is *Botritis cinerea*.
9. Use of the compound of the formula I and the azole derivative of formula II-2 as set forth in claim 1 for preparing a mixture as claimed in claim 1.




I




II-1




II-2




II-3




II-4



II-5



II-6



II-7