



(86) Date de dépôt PCT/PCT Filing Date: 1994/10/21
 (87) Date publication PCT/PCT Publication Date: 1995/05/04
 (45) Date de délivrance/Issue Date: 2002/10/01
 (85) Entrée phase nationale/National Entry: 1996/04/18
 (86) N° demande PCT/PCT Application No.: US 1994/012059
 (87) N° publication PCT/PCT Publication No.: 1995/011669
 (30) Priorité/Priority: 1993/10/25 (142,743) US

(51) Cl.Int.⁶/Int.Cl.⁶ A61K 31/565, A61K 9/10
 (72) Inventeurs/Inventors:
 Guy, Yaacov J., IL;
 Friedman, Doron, IL
 (73) Propriétaire/Owner:
 Pharmos Corporation, US
 (74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : SUSPENSION D'ETABONATE DE LOTEPREDNOL
 (54) Title: SUSPENSION OF LOTEPREDNOL ETABONATE

(57) **Abrégé/Abstract:**

The invention provides novel compositions of matter for delivering water-insoluble steroid drugs suitable for therapeutic use. The invention also provides stable aqueous suspensions of water-insoluble steroid drugs of particle sizes of $\leq 15\mu\text{m}$ which remain in such a state so as to allow for immediate suspension, when desired, even after extended periods of settling.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 9/10	A1	(11) International Publication Number: WO 95/11669 (43) International Publication Date: 4 May 1995 (04.05.95)
<p>(21) International Application Number: PCT/US94/12059</p> <p>(22) International Filing Date: 21 October 1994 (21.10.94)</p> <p>(30) Priority Data: 142,743 25 October 1993 (25.10.93) US</p> <p>(71) Applicant (<i>for all designated States except US</i>): PHARMOS CORPORATION [US/US]; 101 East 52nd Street, New York, NY 10022 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (<i>for US only</i>): GUY, Yaacov, J. [US/IL]; 26 Madar Street, 76552 Rehovot (IL). FRIEDMAN, Doron [IL/IL]; 33 Alon Street, 72910 Carmei Yosef (IL).</p> <p>(74) Agents: FANUCCI, Allan, A. et al.; Pennie & Edmonds, 1155 Avenue of the Americas, New York, NY 10036 (US).</p>	<p>(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).</p> <p style="text-align: right;">2174550</p> <p>Published <i>With international search report.</i></p>	
(54) Title: SUSPENSION OF LOTEPREDNOL ETABONATE		
<p>(57) Abstract</p> <p>The invention provides novel compositions of matter for delivering water-insoluble steroid drugs suitable for therapeutic use. The invention also provides stable aqueous suspensions of water-insoluble steroid drugs of particle sizes of $\leq 15\mu\text{m}$ which remain in such a state so as to allow for immediate suspension, when desired, even after extended periods of settling.</p>		

SUSPENSION OF LOTEPREDNOL ETABONATE

Field of Invention

5 The invention relates to aqueous suspensions for treatment of ophthalmic and otolaryngological inflammations.

Background of the Invention

10 Numerous drugs are prepared in the form of suspensions for ophthalmic, oral, otic, nasal respiratory topical, and parenteral applications. Formulation of pharmaceutical dosages of water-insoluble drugs as suspensions is frequently hampered by the subsequent
15 formation of cakes resulting from aggregation of the suspended material. Polymeric compounds (e.g. polyvinyl pyrrolidone (PVP), polyvinyl alcohol (PVA), dextrin) are commonly used to stabilize such suspensions. An
20 alternative approach to the preparation of such drugs is to enhance the solubility of the drugs within the formulation by vehicles including emulsions, liposomes, and cyclodextrins. However, certain drugs, in their
25 therapeutic concentrations, are not sufficiently stabilized or solubilized by these methods for the above-mentioned applications.

30 Topical steroids such as corticosteroids are commonly used for anti-inflammatory therapy of the eye, especially for treating inflammatory conditions of the palpebral or bulbar conjunctiva, cornea and anterior
35 segment of the globe. Common therapeutic applications for steroids include allergic conjunctivitis, acne rosacea, superficial punctate keratitis and iritis cyclitis. Steroids also are used to ameliorate inflammation associated with corneal injury due to
 chemical or thermal burns, or penetration of foreign
 bodies. Such conditions may result from surgery, injury,

allergy or infection to the eye and can cause severe discomfort.

5 Despite their therapeutic advantages, topical ocular use of corticosteroids is associated with a number of complications, including posterior subcapsular cataract formation, elevation of intraocular pressure, secondary ocular infection, retardation of corneal wound healing, uveitis, mydriasis, transient ocular discomfort and ptosis. Numerous systemic complications also may arise from the topical ocular application of corticosteroids. 10 These complications include adrenal insufficiency, Cushing's syndrome, peptic ulceration, osteoporosis, hypertension, muscle weakness or atrophy, inhibition of growth, diabetes, activation of infection, mood changes and delayed wound healing.

15 Topical steroids for treating ocular inflammations can be based on soft drugs. Soft drugs, as is known in the art, are designed to provide maximal therapeutic effect and minimal side effects. By one approach, synthesis of a "soft drug" can be achieved by 20 structurally modifying a known inactive metabolite of a known active drug to produce an active metabolite that undergoes a predictable one-step transformation in-vivo back to the parent, (see, U.S. patents 4,996,335 and 4,710,495 for soft steroids) inactive metabolite. "Soft 25 drugs" therefore are biologically active chemical components characterized by predictable in vivo metabolism to non-toxic derivatives after they provide their therapeutic effect.

30 Pharmaceutical compositions of water-insoluble drugs such as corticosteroids in aqueous suspensions for ocular and other uses must satisfy constraints imposed by physiological compatibilities such as pH, osmolality, and particle size of the suspended steroids. Furthermore, these compositions must meet requirements for 35

preservative efficiency and ease of suspension over extended periods of time.

Therapeutic suspensions of corticosteroids typically employ polymeric compounds such as polyvinyl pyrrolidone ("PVP") and polyvinyl alcohol ("PVA") as suspending agents in concentrations ranging from 0.1 to 10% (U.S. Pat. 2,861,920). Combinations of polymeric compounds such as PVP, PVA, sodium carboxymethylcellulose ("CMC"), and dextrin, with surface-active agents such as Polysorbate* 80, Polysorbate 20, and tyloxapol also have been used to stabilize corticosteroid suspensions intended for ophthalmic, nasal, and otic uses.

The amounts of polymeric compounds and surface active agents must be determined to provide stability to suspensions of corticosteroids. Excessive amounts of polymeric compounds may hamper the antimicrobial effects of preservatives added to the suspension. Also, pharmaceutical ocular and nasal dosages of these suspensions either must be buffered or have an appropriate pH with no buffering capacity. These suspensions also should be isotonic.

Loteprednol etabonate ("LE") is a known soft corticosteroid based on the known inactive metabolite prednisolone acetate of the active drug prednisolone. See U.S. patents 4,996,335 and 4,710,495.

LE is an analog of prednisolone that does not have a 20-keto group attached to the 17 β -position. Instead, the 17- β position is occupied with a metabolically-labile ester function. In biological systems, LE is hydrolysed to the inactive carboxylic acid metabolite (PJ-91) that does not bind to glucocorticoid receptors. LE also provides superior safety by reducing the risk of steroid induced cataracts and elevation of intra-ocular pressure. The lability of LE to enzymes located in the blood and/or liver also reduces the likelihood of systemic side effects. LE therefore provides therapeutic advantages

* trademark

over other corticosteroids by providing efficacy similar to its parent compound, namely, prednisolone acetate, with fewer deleterious systemic side effects. Soft steroids have the potential advantage of treating inflammation without inducing elevation of intraocular pressure. In addition, soft steroids can provide the added benefit of a lower tendency to induce cataracts which may result from interaction of corticosteroids with the ocular lens proteins.

Formulation of stable aqueous suspensions of LE for ocular applications and other uses, however, has been hampered by agglomeration of the steroid particles. Unexpectedly, common tonicity agents such as aqueous solutions containing 0.9% NaCl, 0.1% EDTA, or phosphate buffer, even in concentrations as low as 1mM, can not be employed to provide stable aqueous suspensions of corticosteroids such as LE.

A need therefore exists for aqueous suspensions of corticosteroids such as LE which can be formulated without agglomeration. A further need exists for aqueous suspensions which have therapeutically effective amounts of corticosteroids such as LE but which avoid the problems associated with the steroid suspensions of the prior art.

Summary of the Invention

The invention provides novel compositions of matter containing water-insoluble drugs suitable for therapeutic use. The invention provides stable aqueous suspensions of water-insoluble drugs of mean particle sizes of $\leq 15\mu\text{m}$ which remain in such a state so as to allow for immediate suspension, when desired, even after extended periods of settling.

More particularly, the invention is directed to aqueous suspensions of soft corticosteroids such as loteprednol etabonate suitable for therapeutic use in the

eye, ear, or nose. The aqueous suspensions of LE are surprisingly stable and can remain in a state suitable for immediate suspension when desired, even after extended 5 periods of settling. The suspensions of the invention, moreover, do not cause discomfort upon application.

The aqueous suspensions of the invention comprise component (A) of a therapeutic quantity of a "soft" steroid such as LE present as particles preferably having a mean 10 diameter of less than about fifteen microns, component (B) of a suspending agent of a nonionic polymer in an aqueous medium, and component (C) of a nonionic surface active agent. Advantageous molar ratios of (A):(B):(C) can vary from about 1:0.01:0.05 to 1:20:1.

15 The steroid of component (A) preferably is loteprednol etabonate and is added to obtain a final concentration in the suspension of about 0.2-2%, preferably about 0.5-1% (w/w). The suspending agent may be any nonionic polymer which is soluble in an aqueous medium, and can be present in an amount 20 of about 0.2 to 2% by weight, and preferably about 0.4 to 1% by weight. The molar ratio of component (A) to component (B) typically is in the range of about 1:0.01 to about 1:20, preferably about 1:0.5 to about 1:10 and more preferably about 1:0.1 to 1:3.

25 The nonionic surfactant of component (C) of the composition may be any one of a wide variety of nonionic alkylene oxide condensates of an organic compounds which contain one or more hydroxyl groups. This component (C) is advantageously present in an amount of between about 0.05 and 30 1% by weight of the composition. The molar ratio of component (A) to component (C) typically is in the range of about 1:0.05 to about 1:1.

The compositions generally include component (D) of a nonionic tonicity agent for producing isotonicity, and, if 35 necessary, component (E) of one or more preservatives.

It is essential that these components (A)-(D) be nonionic insofar as possible since it has now been discovered that the presence of ions is the major cause of caking. Thus, the preferred tonicity agents would be nonionic diols such as glycerol or mannitol rather than
5 the commonly used sodium chloride. The nonionic tonicity agent is preferably present in an amount of about 1.05 to 5.75% by weight and more preferably about 1.5 to 4%.

Accepted preservatives such as benzalkonium chloride and disodium edetate (EDTA) may be included in the
10 suspensions of the invention in concentrations sufficient for effective antibacterial action, preferably about 0.0001 to 0.025%, based on the weight of the suspension.

Having briefly summarized the invention, the invention will now be described in detail by reference to
15 the following specification and non-limiting examples. Unless otherwise specified, all percentages are by weight.

Detailed Description of the Invention

20 Therapeutic suspensions of LE for ophthalmic or otolaryngological uses are made by aseptic preparation. Purity levels of all materials employed in the suspensions of the invention exceed 98%. The suspensions of the invention are prepared by thoroughly mixing the
25 drug (component (A)), suspending agent (component (B)), and surface active agent (component (C)). Optionally, tonicity agents (component (D)) and preservatives (component (E)) may be included.

Drugs of component (A), preferably soft steroids and
30 most preferably LE, can be employed. Also, other steroids such as beclomethasone, betamethasone, fluocinolone, fluorometholone, exednisolone, may be employed. The suspensions of component (A) of the invention have a particle size of about 0.1-30 μm ,
35 preferably about 1-20 μm , most preferably about 2-10 μm

994/12059 95/11669
be
se
e

in mean diameter. LE in this size range is commercially available from suppliers such as the Sipsy Co., (Avrillé, France).

5 The nonionic polymer of component (B) can be any nonionic water-soluble polymer. Typical compounds such as PVP, PVA, dextrin or cyclodextrin can be used in a concentration of about 0.2-2%, and preferably 0.4 to 1%.

10 Component (C) is a surface-active agent that is acceptable for ophthalmic or otolaryngological uses. Preferably, this surfactant is non-ionic. Generally, the nonionic surfactant is a nonionic alkylene oxide condensate of an organic compound which contains one or more hydroxyl groups. For example, ethoxylated and/or propoxylated alcohol or ester compounds or mixtures thereof are commonly available and are well known to 15 those skilled in the art. Useful surface active agents include but are not limited to POLYSORBATE 80, tyloxapol, TWEEN* 80 (ICI America Inc., Wilmington, Delaware), PLURONIC* F-68 (from BASF, Ludwigshafen, Germany) and the POLOXAMER* surfactants can also be used. The tyloxapol and TWEEN surfactants are preferred because they are FDA 20 approved for human use. The concentration in which the surface active agent may be used is only limited by neutralization of the bacteriocidal effects on the accompanying preservatives, or by concentrations which 25 may cause irritation. Advantageously, the concentration of component (C) is about 0.05 to 1% based on the weight of the suspension, and preferably about 0.1 to 0.6%.

30 The tonicity agents of component (D) can be nonionic diols including mannitol and preferably glycerol, in sufficient amounts to achieve isotonicity. The nonionic tonicity agent is present in an amount of about 1.05 to 5.75% by weight, and preferably about 1.5 to 4%.

35 The nonionic polymeric compounds of component (B), and the surface active agents of component (C) have good solubility in water, have sufficient number of hydroxyl

* trademark

groups to interact with the steroid, and have mild effects on the viscosity of the suspension. Final viscosity should not exceed 80-centipoise.

5 In a preferred aspect, stable aqueous suspensions of LE are provided by preparing aqueous suspensions of LE in concentrations of about 0.5-1% with about 0.6% PVP, about 2-2.8% glycerol, preferably about 2.2-2.6% glycerol, most preferably about 2.4% glycerol, and about 0.05-1% tyloxapol.

10 The suspensions of the invention also may include additional therapeutic drugs such as drugs for treating glaucoma, anti-inflammatory drugs, antibiotic drugs, anti-cancer drugs, anti-fungal drugs and anti-viral drugs. Examples of anti-glaucoma drugs include but are not limited to timolol-base*, betaxalol*, athenolol*,
15 levobanolol*, epinenephrin*, dipivalyl*, oxonolol*, acetazolamide*-base and methazolamide*. Examples of anti-inflammatory drugs include but are not limited to non-steroids such as piroxicam*, indomethacin*, naproxen*, phenylbutazone*, ibuprofen* and diclofenac.* Additional
20 therapeutic materials which may be employed include but are not limited to tobramycin*, gentamycin* or other antibiotics.

Health regulations in various countries generally require that ophthalmic preparations shall include a
25 preservative. Many well known preservatives that have been used in ophthalmic preparations of the prior art, however, cannot be used in the preparations of the invention, since those preservatives may no longer be considered safe for ocular use, or may interact with the
30 surfactant employed in the suspension to form a complex that reduces the bacteriocidic activity of the preservative.

The preservatives of component (E) employed in the suspensions of the invention therefore are chosen to not
35 interact with the surface active agent to an extent that

* trademark

the preservatives are prevented from protecting the suspension from microbiological contamination. In a preferred embodiment benzalkonium chloride may be employed as a safe preservative, most preferably benzalkonium chloride with EDTA. Disodium edetate has also been found to be effective in reducing microbial growth in the present formulations. Other possible preservatives include but are not limited to benzyl alcohol, methyl parabens, propyl parabens, thimerosal, chlorbutanol and benzethonium chlorides. Preferably, a preservative (or combination of preservatives) that will impart standard antimicrobial activity to the suspension and protect against oxidation of components (A)-(D) is employed. These preservatives are generally used in an amount of about 0.0001 to 0.025% by weight and preferably 0.001 to 0.015%.

Stable aqueous suspensions of the invention can be produced over a broad range of pH values. A pH of about 4.5 - 7.4 especially is useful for preparing the stable LE suspensions of the invention.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments therefore are to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. In the following examples, all parts and percentages are by weight unless otherwise indicated,

Examples 1-37

Each of Examples 1-37 are prepared by dissolving the suspending agent (Component B) in water by gentle mechanical mixing. Subsequently, the surfactant (Component C), the tonicity agent(s) and the preservatives (Components (D) and (E), respectively) are added in that order. The solution is then sterilized by

filtration or autoclaving. LE, presterilized by irradiation, is added aseptically to the solution, and the dispersion is then mixed at 12,000 rpm for one minute. The amounts of these components are shown in Table 1.

5

SIZE DETERMINATION

The size distributions of the LE particles in the samples of Table 1 are measured with a Coulter^R LS 130 instrument. An acceptable average particle size for ophthalmic suspensions is $\leq 15 \mu\text{m}$. The results appear in Table 2.

10

15

20

25

30

35

Example Number	SAMPLE COMPOSITION (%w/w)												
	LE	Tween 80	Tyloxapol	Poloxamer-188	HPMC ¹	PVA	PVP	dextrin	Osmolarity Agent	EDTA ⁴	BKA ⁵		
1	0.5	--	0.2	--	--	0.6	--	--	--	--	--	--	
2	0.5	0.2	--	--	--	--	0.2	10 mM PBS ²	--	--	--	--	
3	0.5	--	0.4	--	0.2	--	0.4	--	--	--	--	--	
4	0.5	--	0.2	--	--	0.2	--	10 mM PBS	--	--	--	--	
5	0.5	0.6	--	--	0.4	--	--	100mM PBS	--	--	--	--	
6	0.5	0.4	--	--	--	1.4	--	5mM PBS	--	--	0.001	--	
7	1	--	0.2	--	--	--	1	--	--	--	--	--	
8	1	--	0.6	--	--	--	1.4	--	--	--	--	--	
9	1	0.6	--	--	--	--	1.4	--	--	--	--	--	
10	0.5	--	0.4	--	--	1	--	0.9% saline ³	--	--	0.001	--	
11	0.5	0.4	--	--	--	1	--	0.9% saline	--	--	0.001	--	
12	0.5	0.6	--	--	--	--	2	2.4% glycerol	0.01	0.01	0.01	--	
13	0.5	--	0.3	--	--	--	1.5	2.4% glycerol	0.01	0.01	0.01	--	
14	0.5	--	0.3	--	--	--	0.6	2.4% glycerol	0.01	0.01	0.015	--	
15	0.5	0.4	--	--	--	1.4	--	2.4% glycerol	--	--	0.001	--	
16	1	--	0.2	--	--	--	1	2.4% glycerol	--	--	0.01	--	
17	0.5	--	0.6	--	--	1.4	--	2.4% glycerol	--	--	0.01	--	
18	0.5	--	--	0.6	--	--	2	2.4% glycerol	0.01	0.01	0.004	--	
19	0.5	0.4	--	--	--	--	--	2.4% glycerol	0.01	0.01	0.004	--	
20	0.5	--	0.4	--	--	--	--	2.4% glycerol	0.01	0.01	0.01	--	
21	0.5	--	0.3	--	--	--	1	2.4% glycerol	0.01	0.01	0.01	--	
22	0.5	0.6	--	--	--	1.4	--	2.4% glycerol	--	--	--	--	

SAMPLE COMPOSITION (%w/w)

Example Number	LE	Tween 80	Tyloxapol	Poloxamer-188	HPMC ¹	PVA	PVP	dextrin	Osmolarity Agent	EDTA ⁴	BKA ⁵
23	0.5	0.6	--	--	--	1.4	--	--	2.4% glycerol	--	0.004
24	0.5	0.6	--	--	--	--	--	2	2.4% glycerol	0.01	0.01
25	0.5	--	0.3	--	--	--	0.6	--	2.4% glycerol	0.01	0.01
26	0.5	--	0.3	--	--	--	0.6	0.5	2.4% glycerol	0.01	0.01
27	1	--	0.3	--	--	--	0.6	0.5	2.4% glycerol	0.01	0.015
28	1	--	0.3	--	--	--	0.6	--	2.4% glycerol	0.01	0.015
29	1	--	0.1	--	--	--	0.4	--	2.4% glycerol	0.01	0.01
30	0.5	--	0.2	--	--	--	0.6	--	2.4% glycerol	0.01	0.01
31	1	--	0.2	--	--	--	0.6	--	2.4% glycerol	0.01	0.01
32	1	--	0.2	--	--	--	0.8	--	2.4% glycerol	0.01	0.015
33	0.5	--	0.3	--	--	--	1.5	--	2.4% glycerol	0.01	0.015
34	1	--	0.4	--	--	--	0.4	--	2.4% glycerol	0.01	0.01
35	0.5	--	0.3	--	--	--	0.6	0.3	2.4% glycerol	0.01	0.01
36	0.5	--	0.1	--	--	--	0.4	0.3	2.4% glycerol	0.01	0.01
37	0.5	--	0.3	--	--	--	0.6	--	2.4% glycerol	0.01	0.015

- 1 hydroxypropylmethyl cellulose
- 2 phosphate buffered physiological saline
- 3 sodium chloride
- 4 ethylene diamine tetraacetic acid
- 5 benzalkonium chloride

TABLE 2

Example Number	Particle Size(s) (μm) and Fraction of Total Population			
	Population A	A%	Population B	B%
1	3.906+/-2.677	86.62	53.67+/-13.13	13.38
2	112.7+/-13.27	100	--	--
3	3.526+/-1.706	100	--	--
4	111.4+/-18.59	100	--	--
5	23.52+/-20.58	100	--	--
6	32.83+/-2.563	48.74	94.06+/-40.57	51.26
7	4.596+/-2.698	92.43	57.91+/-18.14	7.57
8	3.805+/-2.417	93.14	62.38+/-20.38	6.86
9	6.591+/-3.566	100	--	--
10	3.828+/-2.693	17.52	96.28+/-38.13	82.48
11	3.888+/-2.69	10.95	110.1+/-58.02	85.98
12	3.559+/-1.469	5.62	82.84+/-13.08	94.38
13	2.932+/-2.32	3.52	100.1+/-24,56	96.48
14	88.52+/-30.19	100	--	--
15	3.652+/-2.692	100	--	--
16	3.851+/-2.401	100	--	--
17	3.969+/-2.572	100	--	--
18	4.926+/-2.955	92.29	41.59+/-7.125	7.71
19	4.429+/-2,732	100	--	--
20	3.980+/-2.566	100	--	--
21	3.633+/-2.457	100	--	--
22	4.716+/-2.762	100	--	--
23	4.789+/-2.823	100	--	--
24	4.528+/-2.552	100	--	--
25	5.261+/-2.990	100	--	--
26	5.262+/-3.013	100	--	--
27	5.204+/-2.985	100	--	--
28	4.918+/-2.832	100	--	--
29	4.126+/-2.390	100	--	--
30	12.45+/-10.91	100	--	--
31	3.976+/-2.245	100	--	--
32	3.789+/-1.609	100	--	--

TABLE 2 (cont'd)

Example Number	Particle Size(s) (μm) and Fraction of Total Population			
	Population A	A%	Population B	B%
33	3.821+/-2.181	46.77	107.3+/-14.74	53.23
34	3.813+/-2.305	100	--	--
35	3.385+/-1.506	78.44	25.16+/-1.421	21.56
36	3.737+/-2.044	100	--	--
37	3.965+/-2.229	100	--	--

1. In the Coulter particle size analysis two distinct populations of particles were sometimes discerned. In these cases the two populations are denoted as populations A and B. If only a single population was detected it is denoted population A.

EVALUATION OF SUSPENDABILITY OVER TIME

Samples containing particles with desirable size distributions (average of 2-10 μm) are tested for stability using accelerated stability tests as well as "real time" studies.

Accelerated stability studies are performed by subjecting the samples to a centrifugal force of 5000xG for two minutes. The suspendability of the settled material is tested by measuring the number of seconds of wrist shaking required to eliminate visible residue attached to the container. Since existing marketed products require as much as sixty seconds of wrist shaking to suspend the entire amount of settled residue, ten seconds is determined to be an acceptable amount of time to suspend the residue. The results are shown in Table 3.

TABLE 3
RESUSPENSION OF LE SUSPENSIONS WHICH HAVE UNDERGONE
ACCELERATED AND NATURAL¹ SETTLING

Example Number	Accelerated Stability (time to resuspend) ²	Suspension of naturally settled material ¹	
		Initial Value (# inversions)	Months Tested ³
15	15	--	10(I)
16	5	--	10(I)
17	5	--	10(I)
18	5	--	9
19	5	--	9
20	--	67	9
21	--	46	9
22	--	83	9
23	--	37	9
24	--	--	6(I)
25	5	27	8
26	5	22	6(I)
27	5	35	6(I)
28	5	35	8(I)
29	--	49	7
30	5	25	7

TABLE 3 (cont'd)

Example Number	Accelerated Stability (time to resuspend) ¹	Suspension of naturally settled material ¹	
		Initial Value (# inversions)	Months Tested ³
31	5	43	7
32	--	74	7
33	--	136	3(I)
34	--	40	7
35	--	18	7
36	--	48	7
37	--	46	8

¹ Refers to settling, at room temperature, on an open shelf

² Number of seconds of wrist shaking to suspend material that was settled by application of 5000xG for 2 minutes.

³ During the test period, samples were periodically examined to verify the retention of the initial values "I" indicates instability for the noted period, i.e., agglomeration.

The results shown in Table 3 show samples which do not form agglomerates during the longest period of observation. Acceptable samples require ≤ 100 gentle inversions following the indicated period of settling.

5 The stability of suspensions intended for multiple doses is supported by the addition of preservatives which prevent potential microbiological growth. The indicated preparations are prepared under aseptic conditions and aliquots of each material are exposed to the indicated
10 microbiological organisms for four weeks and evaluated for growth as described in the U.S. Pharmacopeia. The results, shown in Table 4, indicate whether the preservative was effective (+) or ineffective (-) according to U.S.P. requirements.

15

UNIDOSE SUSPENSIONS
WITHOUT PRESERVATIVES

20 Compositions with satisfactory particle sizes and stabilities for unidose suspensions without preservatives appear in Table 5. These compositions are satisfactory for ophthalmic or otolaryngological uses when prepared under aseptic conditions and packaged in containers for
25 single doses.

30

35

35

TABLE 4

Example	Challenge Microorganism					
	Staph. aureus	P. aerug.	Candida albicans	Asper. niger	E. coli	
23	+	-	-	-	ND	
24	+	-	-	-	ND	
25	+	-	-	+	ND	
26	+	+	+	+	ND	
27	+	+	-	+	+	
28	+	+	+	+	+	
29	+	+	+	+	ND	
30	+	+	+	+	+	
31	+	+	+	+	+	
32	+	+	+	+	+	

ND: denotes not done; (+) denotes challenge withstood; (-) denotes unacceptable microbe growth
The test was performed according to U.S.P. specifications.

TABLE 5
COMPOSITIONS OF EXEMPLARY LE FORMULATIONS FOR UNIDOSE APPLICATION

Ex. No.	LE	Tween 80	Tyloxapol	Poloxamer-188	PVA	PVP	dextrin	glycerol	Purified Water
1	1	0.6	--	--	--	1.4	--	2.4	Remainder
2	0.5	--	--	0.6	--	2	--	2.4	Remainder
3	0.5	0.4	--	--	--	--	1.6	2.4	Remainder
4	0.5	--	0.4	--	--	--	2.4	2.4	Remainder
5	0.5	--	0.3	--	--	1	--	2.4	Remainder
6	0.5	--	0.6	--	0.8	--	--	2.4	Remainder
7	0.5	0.6	--	--	1.4	0.8	--	2.4	Remainder
8	0.5	0.6	--	--	--	--	2	2.4	Remainder
9	0.5	0.6	--	--	--	--	2.4	2.4	Remainder
10	0.5	--	0.3	--	--	0.6	--	2.4	Remainder
11	0.5	--	0.3	--	--	0.6	0.5	2.4	Remainder
12	1	--	0.3	--	--	0.6	0.5	2.4	Remainder
13	1	--	0.3	--	--	0.6	--	2.4	Remainder
14	1	--	0.1	--	--	0.4	--	2.4	Remainder
15	0.5	--	0.2	--	--	0.6	--	2.4	Remainder
16	1	--	0.2	--	--	0.6	--	2.4	Remainder
17	1	--	0.4	--	--	0.6	--	2.4	Remainder
18	1	--	0.2	--	--	0.8	--	2.4	Remainder
19	1	--	0.4	--	--	0.4	--	2.4	Remainder
20	0.5	--	0.4	--	--	0.4	--	2.4	Remainder
21	0.5	--	0.3	--	--	0.6	0.3	2.4	Remainder
22	0.5	--	0.1	--	--	0.4	0.3	2.4	Remainder

Example 38

The soft steroid loteprednol etabonate was formulated as an aqueous ophthalmic suspension containing polyvinyl pyrrolidone (0.6%), glycerine (2.4%), tyloxapol (0.3%), edetate disodium (0.0005%) and benzalkonium chloride (0.001%). Loteprednol etabonate (0.5%) was incorporated into this vehicle for use in clinical studies. During these studies, the formulation was evaluated on a total of 446 patients, 220 of which had giant papillary conjunctivitis ("GPC"), 145 of which had seasonal allergic conjunctivitis ("SAC") and 81 had acute anterior uveitis.

Loteprednol etabonate in this formulation was readily suspendable throughout extended periods of storage (i.e., greater than 18 months) as well as throughout the clinical treatment. The preparation was well tolerated in all patients and was significantly more effective than the vehicle itself, which was used as a placebo, with regard to the reduction of signs and symptoms of ocular inflammation.

The vehicle was administered as a placebo to 219 GPC patients and 143 SAC patients. In SAC, treatment was initiated prophylactically, and therefore it was not possible to quantitate accurately the placebo effect. The GPC patients were enrolled in the study after the appearance of signs or symptoms. A significant number of GPC patients experienced clinically meaningful relief of signs and symptoms with the application of the vehicle alone. While the use of a demulcent solution applied four times per day should have some benefit in the treatment of GPC, the extent to which this occurred was higher than expected. Specifically, the size of the papillae was reduced in 50% of the patients, itching was reduced in 78% of the patients, and contact lens comfort

was increased in 71% of the patients. This shows that the vehicle itself is useful for such treatments.

In 76 to 94% of the GPC patients, loteprednol etabonate treatment resulted in clinically meaningful improvement in the same areas. These results are
5 statistically significant ($p < 0.001$) compared to the administration of the vehicle alone. Thus, it is preferred to administrate the vehicle containing the loteprednol etabonate for optimum results.

10

15

20

25

30

35

THE CLAIMS

We claim:

1. A composition for ophthalmic or otolaryngological anti-inflammatory use comprising:

- 5 (A) a therapeutic quantity of a corticosteroid;
 (B) a nonionic polymer in an aqueous medium; and
 (C) a nonionic surface active agent in an amount sufficient to retain the corticosteroid and nonionic polymer in solution.

10 2. The composition of claim 1 wherein the molar ratio of (A):(B):(C) is about 1:20:0.05 to about 1:0.05:1.

15 3. The composition of claim 1 wherein the corticosteroid is selected from the group consisting of soft steroids having anti-inflammatory activity and is present in an amount of about 0.2 and 2% by weight.

4. The composition of claim 1 wherein the corticosteroid is loteprednol etabonate and is present in an amount of about 0.5 and 1% by weight.

20 5. The composition of claim 1 wherein said nonionic polymer is a water soluble polymer selected from the group consisting of polyvinylpyrrolidone, polyvinyl alcohol, dextrin and cyclodextrin and is present in an amount of about 0.2 to 2% by weight.

25 6. The composition of claim 1 wherein said nonionic polymer is polyvinylpyrrolidone and is present in an amount of about 0.4 to 1% by weight.

30 7. The composition of claim 1 wherein said nonionic surface active agent is tyloxapol* and is present in an amount of about 0.1 to 0.6% by weight.

8. The composition of claim 1 which further comprises a nonionic tonicity agent in an amount sufficient to achieve isotonicity.

35 9. The composition of claim 8 wherein the nonionic tonicity agent is present in an amount of about 1.05 to

* trademark

5.75% by weight and the nonionic surfactant is present in an amount of about 0.05 to 1% by weight.

10. The composition of claim 8 wherein said non-ionic tonicity agent is a nonionic diol and is present in an amount of about 1.5 to 4% by weight.

11. The composition of claim 10 wherein said diol is glycerol or mannitol.

12. The composition of claim 1 further including at least one preservative for preventing microbial formation in said composition, the preservative being present in an amount of about 0.0001 to 0.025% by weight.

13. The composition of claim 12 wherein said preservative is selected from the group consisting of benzalkonium chloride, disodium edetate and mixtures thereof.

14. The composition of claim 12 wherein said preservative is a mixture of benzalkonium chloride and disodium edetate.

15. The composition of claim 1 further comprising an additional therapeutic drug in admixture with said corticosteroid, wherein said additional therapeutic drug is selected from the group consisting of betaxalol, athenolol, livobanolol, epinenephrin, dipivalyl, oxonolol, acetazolamide-base, methazolamide, tobramycin, gentamycin, piroxicam, indomethacin, naproxen*, phenylbutazone, ibuprofen*, and diclofenac-acid.

16. The composition of claim 1 wherein said corticosteroid has a particle size less than about fifteen microns.

17. A composition for ophthalmic or otolaryngological anti-inflammatory use comprising a nonionic polymer in an aqueous medium, a nonionic tonicity agent in an amount effective to produce isotonicity, and a nonionic surface active agent in an amount sufficient to retain the polymer and tonicity agent in the aqueous medium.

* trademark

18. The composition of claim 17 wherein the nonionic polymer is present in an amount of about 0.2 to 2% by weight; the nonionic tonicity agent is present in an amount of about 1.05 to 5.75% by weight; and the nonionic surface active agent is present in an amount of about 0.05 to 1% by weight.

19. The composition of claim 17 further comprising a preservative of benzalkonium chloride, disodium edetate, and mixtures thereof in an amount of about 0.0001 to 0.025% by weight.

20. The composition of claim 17 wherein the nonionic polymer is polyvinyl pyrrolidone and is present in an amount of about 0.4 to 1% by weight, the nonionic tonicity agent is a diol and is present in an amount of about 2 to 2.8% by weight, and the nonionic surface active agent is tyloxapol and is present in an amount of about 0.1 to 0.6% by weight.

21. The composition of claim 20 wherein the nonionic tonicity agent is glycerol or mannitol and further comprising a therapeutic quantity of a corticosteroid have a particle size of 0.1 μ m to 30 μ m in diameter, wherein the molar ratio of corticosteroid to nonionic polymer to nonionic surface active agent is about 1:20:0.05 to about 1:0.05:1.

25

30

35